BIOL 101- ENVIRONMENTAL SCIENCE



Erin Rempala
San Diego City College

LibreTexts™

BIOL 101- Environmental Science

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 10/23/2026

TABLE OF CONTENTS

Licensing

Licensing

1: Introduction to Environmental Science

1.1: The Process of Science

1.2: What is Science?

1.3: Empirical and Applied Science

1.4: Human Impact on the Planet

1.5: Environmental Science

1.6: Practicing with Data- Wildlife Corridors

2: Humans and the Environment

2.1: Planetary Boundaries- Earth's Safe Operating Space

2.2: Environmental Ethics and Worldviews

2.3: Environmental Justice and Equity

2.4: Sustainability and Sustainable Development

2.5: Systems Thinking and Modeling Sustainability

2.6: Historical Foundations of Environmental Policy

3: Ecology

3.1: Population Ecology

3.1.1: Ecological Hierarchy

3.1.2: Population Distribution

3.1.3: Population Size

3.1.4: Population Growth and Regulation

3.1.5: Life History

3.2: Community Ecology

3.2.1: Biotic Interactions

3.2.1.1: Symbiosis

3.2.1.2: Trophic Interactions

3.2.1.2.1: Predation

3.2.1.2.2: Herbivory

3.2.1.2.3: Competition

3.2.2: Community Structure and Dynamics

3.2.3: Data Dive- Beaver Impacts on Wetlands

3.3: Ecosystem Ecology

3.3.1: Matter

3.3.2: Energy

3.3.3: Biogeochemical Cycles

3.3.4: Energy Flow in Food Webs
3.3.5: Ecosystem Types and Dynamics
3.3.5.1: Climate and Biomes
3.3.5.2: Terrestrial Biomes
3.3.5.3: Aquatic Biomes
3.3.6: Data Dive- Biome Carbon Storage

4: Evolution
4.1: Evolution and Natural Selection
4.2: Other Mechanisms of Evolution
4.3: Evidence for Evolution

5: Biodiversity
5.1: The Value of Biodiversity
5.2: The Importance of Biodiversity to Human Life
5.3: Levels of Biodiversity
5.4: The Biodiversity Crisis
5.5: Threats to Biodiversity
5.6: Extinction
5.7: Measuring Biodiversity Loss
5.8: Preserving Biodiversity
5.9: Policies Protecting Biodiversity
5.10: Data Dive- Threats to Biodiversity

6: Human Population Growth
6.1: History of Human Population Growth
6.2: The Rate of Human Population Growth
6.3: Factors Affecting Population Growth
6.4: Looking Ahead
6.5: Data Dive- World Population Densities

7: The Energy System
7.1: Fossil Fuels
7.1.1: Types of Fossil Fuels and Formation
7.1.2: Mining, Processing, and Generating Electricity
7.1.3: Fossil Fuel Consumption
7.1.4: Consequences of Fossil Fuels

7.2: Nuclear Energy
7.2.1: Radioactive Isotopes
7.2.2: Generating Electricity with Nuclear Energy
7.2.3: Nuclear Energy Consumption
7.2.4: Pros and Cons of Nuclear Energy

7.2.5: The Promise of Fusion
7.2.6: Data Dive- Global Nuclear Power Generation

7.3: Renewable Energy and Conservation
7.3.1: Renewable Energy History and Consumption
7.3.2: Wind Energy
7.3.3: Solar Energy
7.3.4: Hydropower
7.3.5: Geothermal Energy
7.3.6: Biofuels (Biomass Energy)
7.3.7: Energy Efficiency and Conservation
7.3.8: Data Dive- Global Renewable Energy Generation

8: Atmospheric Pollution
8.1: Earth's Atmosphere
8.2: Air Pollution
8.3: Acid Deposition
8.4: Ozone Depletion

9: Global Climate Change
9.1: Weather, Climate, and Climate Change
9.2: Earth's Energy Balance and Radiation
9.3: The Greenhouse Effect
9.4: Evidence and Causes of Climate Change
9.5: Impacts of Climate Change
9.6: Ocean Acidification- The Other Carbon Problem
9.7: Mitigating and Adapting to Climate Change
9.8: Data Dive- Flooding in Venice Italy

10: The Water System
10.1: The Hydrologic Cycle and Freshwater Distribution
10.2: Water Usage
10.3: Water Scarcity and Sustainable Water Management
10.4: Case Study - The Salton Sea
10.5: Water Pollution
10.6: Water Treatment- Protecting Drinking Water and Treating Wastewater
10.7: Preventing Water Pollution and Restoring Watersheds

11: Waste, Materials, and the Circular Economy
11.1: Waste Generation
11.2: Disposal
11.3: Environmental Impacts of Solid Waste
11.4: From Waste Management to a Circular Economy
11.5: Circular Economy and Stewardship

12: Environmental Health and Toxicology

12.1: Types of Environmental Hazards

12.2: Epidemiology

12.3: Infectious Diseases

12.4: Environmental Toxicology

12.5: Case Study - The Love Canal Disaster

12.6: Environmental Hazard Reduction

13: The Agricultural System

13.1: Food Security

13.2: Industrial Agriculture

13.3: Soils

13.4: Soil Degradation

13.5: Selective Breeding and Genetic Engineering

13.6: Sustainable and Regenerative Agriculture

Index

Glossary

Detailed Licensing

This page titled BIOL 101- Environmental Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

Licensing

A detailed breakdown of this resource's licensing can be found in ***Back Matter/Detailed Licensing***.

This page titled Licensing is shared under a license and was authored, remixed, and/or curated by LibreTexts.

Licensing

A detailed breakdown of this resource's licensing can be found in ***Back Matter/Detailed Licensing***.

This page titled Licensing is shared under a license and was authored, remixed, and/or curated by Erin Rempala.

1: INTRODUCTION TO ENVIRONMENTAL SCIENCE

The objective of this unit is to not only get you acquainted with what this book will cover but to also introduce the rigors and structure of science and how mathematics plays a crucial role in helping scientists interpret results in the most unbiased way possible. Science and mathematics are not just for scientists. A basic understanding of these subjects is crucial in our complex modern world. Without a solid understanding of how science and mathematics play a role in our everyday lives, we cannot respond appropriately in everyday situations, know how to vote on scientific laws, or hold public representatives responsible for their decisions on scientific policy.



Figure $1.a$ People doing science. Image by Yaroslav Shuraev (CC)

Attributions

Modified by Erin Rempala

Thumbnail image - "Women doing science". Image by Yaroslav Shuraev (CC)

This page titled 1: Introduction to Environmental Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- Unit 1: Introduction by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.
- 2.2: The Scientific Method by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.1: The Process of Science

Science as a Process

Many people think of science as a set of steps called the scientific method. In reality, science is not a rigid method. Science is an iterative process, meaning scientists often repeat steps, revise ideas, and ask new questions as they learn more.

Scientific investigations do not always follow the same path. One scientist might start with an observation in nature, while another might begin by analyzing existing data or reading research studies. New evidence can lead scientists to change a hypothesis, collect more data, or rethink an explanation. As a result, scientific knowledge grows and improves over time.

Environmental science is especially iterative because it combines ideas from many fields, including biology, chemistry, geology, economics, and social science. Environmental problems are often complex and involve interactions between natural and human systems. Scientists use evidence from many sources to better understand these interactions and to develop solutions.

Although scientific investigations vary, they often begin with observations and questions. Scientists then develop explanations, collect evidence, evaluate results, and share their findings with others. The process continues as new discoveries lead to new questions. In this way, science is not about finding final answers. It is an ongoing process of learning about the natural world.

Making Observations

An **observation** is something a scientist notices. Observations can come from direct experiences in nature or from information reported by other scientists.

A direct observation is something you personally see, hear, measure, or record. For example, you might notice more birds in your yard after you plant a native tree.

An indirect observation comes from information collected by others. For example, you might read a scientific study showing that birds are more common in neighborhoods with native trees.

The following example begins with a direct observation.

During the Vietnam War (figure $1.1.a$), people in North Vietnam reported an increase in birth defects. These reports led scientists to ask whether something in the environment could be causing the problem. Later, some Vietnam veterans who returned to the United States developed higher rates of certain cancers and other diseases. Scientists wondered whether these health problems could be connected to a common cause.

Observations often lead to new scientific questions.




Figure $1.1.a$: A map of Vietnam 1954-1975. Image from Bureau of Public Affairs U.S. Government Printing Office(opens in new window) (public domain).

Asking Questions

Observations often lead to **questions**. Scientists ask: What explains the pattern that was observed? A single observation can lead to many different questions. As reports of birth defects increased in Vietnam and health problems appeared in some Vietnam veterans, scientists began searching for possible causes. One question focused on Agent Orange, a herbicide used during the war to remove forest vegetation and improve visibility for military operations (figure $1.1.b - c$). Could exposure to Agent Orange (figure $1.1.d$). be linked to the birth defects and diseases that were being observed? This question became the starting point for further scientific investigation.



Figure $1.1.b$: Agent Orange drums in Vietnam. Image by U.S. Government(opens in new window) (public domain).



Figure 1.1.*c*: A healthy mangrove forest (top), and another forest after application of Agent Orange. Image by unknown author(opens in new window) (public domain).




Figure 1.1.*d*: The chemical structure of TCDD (2,3,7,8-tetrachlorodibenzo-p-dioxin), which is produced when synthesizing the chemicals in Agent Orange. It contaminates Agent Orange at low but harmful concentrations. Image by Emeldir(opens in new window) (public domain).

Forming Hypotheses

A **hypothesis** is a possible answer to a scientific question that can be tested with evidence. Hypotheses are not "educated guesses" They predict what will happen and provide a scientific reason for that prediction. They also have a particular format and must be a statement.

Consider the following examples:

- Simple hypothesis: Agent Orange affects rates of birth defects and disease.
- Better hypothesis: Agent Orange increases rates of birth defects and disease.
- Stronger hypothesis: Agent Orange increases rates of birth defects and disease because these health problems have been reported more often in people exposed to the herbicide.

Scientists generally prefer hypotheses that are both supported by evidence and as simple as possible. This idea is known as Occam's Razor. Occam's Razor states that when two or more hypotheses explain the same observations equally well, scientists usually prefer the simpler explanation. A simpler explanation is often easier to test and requires fewer assumptions.

To be useful, a hypothesis should make a clear prediction that can be tested through observation or experimentation. A common format is:

If [proposed cause], then [predicted outcome], because [scientific reasoning].

For example:

- If exposure to Agent Orange causes developmental health problems, then mice exposed to TCDD, a contaminant found in Agent Orange, during development will have a higher rate of birth defects than unexposed control mice because TCDD may interfere with normal development.

A well-written hypothesis helps guide the design of an experiment and identifies the evidence that would support or refute the proposed explanation.

A hypothesis must be **testable**. Scientists must be able to collect observations or data that could support or refute the hypothesis. For example, the hypothesis "bears prefer blue berries because they like the color blue" is difficult to test because scientists cannot directly measure what a bear is thinking.

Scientists test hypotheses by collecting evidence through observations, experiments, or other investigations. The goal is not to *prove* a hypothesis is true. Science is not based on "proof", but instead on observable evidence. Therefore, scientists look for evidence that supports or refutes a hypothesis.

If the results do not support the predictions, the hypothesis may need to be revised or rejected. If the results support the prediction, the hypothesis is supported--and probably will undergo further testing such as leading to a new round of questions and hypotheses. However, even a well-supported hypothesis can be challenged by new evidence in the future.

For this reason, scientific knowledge is always open to revision. As new evidence becomes available, explanations can be improved, refined, or replaced by better ones.

Experiment

There are many ways to do science, and not all scientific studies involve experiments. Some scientists conduct experiments, while others make observations in nature, analyze existing data, or use computer models.

When scientists design a study, they identify the **variables**. A variable is any factor that can change.

The **independent variable** is the factor that scientists think may be causing a change. In an experimental study, the scientist changes or manipulates the independent variable.

The **dependent variable** is the factor that is measured. It is the response to the independent variable.

Other factors can also affect the dependent variable. These are called **confounding variables**. Scientists try to keep confounding variables the same across groups so they do not affect the results.

For example, in a study of Agent Orange, exposure to Agent Orange is the independent variable because it is the factor being investigated. The dependent variable could be the rate of birth defects or disease. Other factors, such as age, diet, exercise, and family history, can also affect health. These are potential confounding variables. Scientists try to control these factors so they do not influence the results.

In an experimental study, scientists manipulate the independent variable and measure the response. The condition being tested is called the **treatment**.

For example, scientists could expose developing mice to TCDD, a contaminant found in Agent Orange, and then measure the frequency of birth defects. In this study, TCDD exposure is the independent variable, and the frequency of birth defects is the dependent variable (figure $1.1.e$).

Most experiments include both a treatment group and a **control group**. The treatment group receives the treatment being tested. The control group does not receive the treatment.

The control group provides a **baseline for comparison**. It shows what happens under normal conditions when the treatment is absent. Scientists compare the treatment group and the control group to determine whether the treatment affected the dependent variable.

In this example, the treatment group would include mice exposed to TCDD. The control group would include mice that were not

exposed to TCDD but were otherwise treated the same way. If birth defects occur more often in the treatment group than in the control group, the results support the hypothesis that TCDD affects development.

A well-designed experiment changes only one independent variable at a time. This helps scientists determine whether changes in the dependent variable are caused by the independent variable rather than by confounding variables. In the Agent Orange study, scientists would need to control potential confounding variables such as the age, genetics, diet, housing conditions, and overall health of the mice. If these factors differ between the treatment group and the control group, they could influence the frequency of birth defects and make it difficult to determine whether TCDD exposure caused the observed effects. By keeping confounding variables as similar as possible between groups, scientists can be more confident that any differences in outcomes are due to the independent variable rather than another factor.

To summarize:

- Independent variable = the factor being tested (TCDD exposure)
- Dependent variable = the factor being measured (birth defects in mice)
- Confounding variables = other factors that could also affect the dependent variable (diet, genetics, age, etc.)
- Control group = mice not exposed to TCDD
- Treatment group = mice exposed to TCDD



Figure 1.1.*e*: Laboratory mice. In a proper scientific study, the treatment would be applied to multiple mice. Another group of mice would not receive the treatment (the control group). Image by Aaron Logan(opens in new window) (CC-BY(opens in new window)).

If an experiment is not possible or would be unethical, scientists may conduct an **observational study**. In an observational study, scientists do not manipulate the independent variable. Instead, they collect data and compare groups that differ naturally in their level of exposure to a possible cause.

For example, scientists could compare the health of Vietnam veterans who experienced different levels of exposure to Agent Orange. Because researchers cannot control who was exposed, they must observe and analyze patterns in existing populations.

Whether a study is experimental or observational, scientists need enough data to distinguish real patterns from random variation. For this reason, scientific studies include many **replicates**. Replicates are multiple individuals, samples, or observations within a study. Using many replicates helps scientists determine whether a pattern is caused by the independent variable rather than by natural differences among individuals.

Scientists also expect studies to be **repeatable**. A study is repeatable if other researchers can follow the same methods and obtain similar results. Repeatability helps scientists evaluate whether the findings are reliable.

In addition, important scientific ideas are tested by multiple studies. When many independent studies reach similar conclusions, scientists gain greater confidence in the results. This process helps build scientific knowledge over time.

Results

After scientists design a study, they collect data and analyze the results. Analyzing data helps scientists identify patterns and determine whether the evidence supports the hypothesis. Statistics are often used to summarize data, compare groups, and determine whether observed differences are likely due to the independent variable rather than random chance.

In the Agent Orange example, scientists conducted experiments using mice. Some developing mice were exposed to high levels of 2,4,5-T, a component of Agent Orange, or TCDD, a contaminant found in Agent Orange. Other mice served as the control group and were not exposed.

The results showed that mice exposed to 2,4,5-T or TCDD developed cleft palate birth defects more often than mice in the control group (figure $1.1.f$). Scientists also found that mouse embryos exposed to TCDD were more likely to die before birth than embryos in the control group.

These results support the hypothesis that exposure to chemicals found in Agent Orange can interfere with normal development. However, as with all scientific studies, the results do not prove the hypothesis is correct. Scientists gain confidence in an explanation when multiple studies produce similar results and the evidence continues to support the hypothesis.




Figure $1.1.f$: Cleft lip and palate, a birth defect in which these structures are split. Image by James Heilman, MD(opens in new window) (CC-BY-SA(opens in new window)).

An observational study of Korean veterans who served in the Vietnam War found that veterans who reported higher levels of Agent Orange exposure also had higher rates of several health problems. These included some types of cancer, diseases of the cardiovascular and nervous systems, skin disorders, and psychological disorders.

To understand these results, it is helpful to understand different types of relationships between variables. A **positive correlation** occurs when two variables change in the same direction. As one variable increases, the other also increases. A **negative correlation** occurs when two variables change in opposite directions. As one variable increases, the other decreases. **No correlation** occurs when there is no consistent relationship between two variables. Changes in one variable do not predict changes in the other.

Some relationships are more complex and do not follow a straight-line pattern. These are called **non-linear relationships**. In a non-linear relationship, the variables are related, but the pattern may curve, level off, or change direction. For example, plant growth may increase as fertilizer is added, but only up to a certain point. Beyond that point, adding more fertilizer may have little effect or even reduce plant growth.

The study of Korean veterans found a positive correlation between Agent Orange exposure and disease. Veterans who reported higher levels of exposure also reported higher rates of health problems.

However, a correlation does not prove that one variable caused the other. Correlation is not the same as causation. Observational

studies can identify patterns, but they cannot always rule out other explanations. To establish a cause-and-effect relationship, scientists usually need additional evidence from experiments, multiple studies, or other independent lines of evidence.

Understanding the difference between correlation and causation is an important part of scientific thinking. Positive, negative, and non-linear correlations can show that variables are related, but additional evidence is needed to determine whether one variable causes changes in another.

In the case of Agent Orange, scientists did not rely on a single study. Observational studies of veterans were combined with experimental studies using mice and other lines of evidence. Together, these studies provided stronger evidence that chemicals found in Agent Orange were linked to harmful health effects.

Conclusion

After analyzing the data, scientists determine whether the evidence supports their hypothesis. In the Agent Orange example, mice exposed to TCDD and 2,4,5-T developed cleft palate birth defects more often than mice in the control group. This result matched the prediction made by the hypothesis. Observational studies also found that veterans exposed to Agent Orange had higher rates of several diseases.

Together, these findings support the hypothesis that chemicals found in Agent Orange can increase the risk of birth defects and disease. However, scientists do not consider a hypothesis "proven". New evidence can always lead to improved explanations or new questions.

Science is an ongoing process (figure $1.1.h$). Each study adds to our understanding, but no single study provides a final answer. Even when a hypothesis is strongly supported, scientists continue to test it in different ways and under different conditions. For example, researchers continue to study Agent Orange today to better understand its long-term effects on human health and the environment.

As new evidence is collected, scientific explanations become stronger, more detailed, and more reliable.




Figure $1.1.h$: The Process of Science. Erin Rempala (made with Biorender.com)

Scientific findings can influence decision making. Evidence about the health effects of Agent Orange led the United States to provide disability compensation and health benefits to many veterans who were exposed and later developed certain diseases. Scientists also discovered that TCDD, a contaminant found in Agent Orange, could harm human health and the environment. As a result, 2,4,5-T, one of the herbicides used to make Agent Orange, was banned in the United States. Although the other component, 2,4-D, is still used in some herbicides today, modern products do not contain the same combination of chemicals found in Agent Orange. The United States has also worked with Vietnam to clean up sites that remain contaminated with TCDD. These actions show how scientific evidence can influence public policy and environmental decision making.

Building on the Work of Others

Major scientific discoveries sometimes change the way people understand the world. However, even these breakthroughs rarely appear suddenly. Most scientific advances build on ideas, observations, and evidence collected by earlier scientists.

A good example is evolutionary theory. Long before Charles Darwin published his ideas, several scientists had suggested that species change over time. Charles Darwin's grandfather, Erasmus Darwin, proposed that living things were connected through common ancestry and could change over generations. Later, Jean-Baptiste Lamarck argued that species evolve in response to their environment, although scientists no longer accept his proposed mechanism for how evolution occurs.

Darwin built on these earlier ideas as well as evidence gathered by geologists, naturalists, and explorers. For example, Charles Lyell's work showed that Earth changes slowly over long periods of time, and Thomas Malthus wrote about how populations can grow faster than their resources. These ideas helped Darwin develop his explanation of evolution by natural selection.

In the 1850s, Alfred Russel Wallace independently reached many of the same conclusions about evolution by natural selection while studying plants and animals in Southeast Asia. When Wallace shared his ideas with Darwin, both scientists realized they had arrived at a similar explanation for how species change over time. Their findings were jointly presented in 1858. Wallace's discovery encouraged Darwin to publish his own work more quickly. Darwin later published *On the Origin of Species*, which presented extensive evidence for evolution by natural selection and helped establish evolutionary theory as a foundation of modern biology (figure 1.1.*g*).

Most scientific progress, however, happens gradually. Scientists build on the work of those who came before them. Each study adds new evidence, improves existing ideas, or answers a small part of a larger question. Over time, these contributions expand scientific knowledge.

Advances in science are also driven by new tools and technologies. For example, the invention of the light microscope allowed scientists to observe cells for the first time. Later, the electron microscope revealed structures that were too small to be seen with earlier tools. New technologies often lead to new discoveries and new questions.

Science advances through the combined efforts of many scientists working across generations. New discoveries, new technologies, and new ideas all contribute to a growing understanding of the natural world.



Figure 1.1.*h*: Understanding the endangered mountain yellow-legged frog through the lens of evolution and genetics has aided its conservation. Image by Wikimedia(opens in new window) [commons.wikimedia.org] [Iasaac Chellman] (CC-BY(opens in new window)).

References

Institute of Medicine (US) Committee to Review the Health Effects in Vietnam Veterans of Exposure to Herbicides. Veterans and Agent Orange: Health Effects of Herbicides Used in Vietnam(opens in new window). Washington (DC): National Academies Press

(US); 1994. 2, History of the Controversy Over the Use of Herbicides.

Neubert, D., Dillmann, I. Embryotoxic effects in mice treated with 2,4,5-trichlorophenoxyacetic acid and 2,3,7,8-tetrachlorodibenzo-p-dioxin(opens in new window). *Naunyn-Schmiedeberg's Arch. Pharmacol.* **272,** 243–264 (1972).

Stellman, J. M., & Stellman, S. D. (2018). Agent Orange During the Vietnam War: The Lingering Issue of Its Civilian and Military Health Impact(opens in new window). *American journal of public health*, *108*(6), 726–728.

Yi, S. W., Ohrr, H., Hong, J. S., & Yi, J. J. (2013). Agent Orange exposure and prevalence of self-reported diseases in Korean Vietnam veterans(opens in new window). *Journal of preventive medicine and public health = Yebang Uihakhoe chi*, *46*(5), 213–225.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- The Process of Science from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Scientific Methods from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))

This page titled 1.1: The Process of Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 2.2: The Scientific Method by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.2: What is Science?

Like other natural sciences, environmental science gathers knowledge about the natural world. Science is more than just a body of knowledge, science provides a means to evaluate and create new knowledge. The methods of science include careful observation, record keeping, logical and mathematical reasoning, experimentation, and submitting conclusions to the scrutiny of others. Science also requires considerable imagination and creativity; a well-designed experiment is commonly described as elegant or beautiful. Science has considerable practical implications and some science is dedicated to practical applications, such as the prevention of disease (figure $1.2.a$). Other science proceeds largely motivated by curiosity. Whatever its goal, there is no doubt that science has transformed human existence and will continue to do so.




Figure $1.2.a$: Biologists may choose to study *Escherichia coli* (*E. coli*), a bacterium that is a normal resident of our digestive tracts but which is also sometimes responsible for disease outbreaks. In this micrograph, the bacterium is visualized using a scanning electron microscope and digital colorization. (credit: Eric Erbe; digital colorization by Christopher Pooley, USDA-ARS)

There are areas of knowledge, however, to which the methods of science cannot be applied. These include such things as morality, aesthetics, or spirituality. Science cannot investigate these areas because they are outside the realm of material phenomena, the phenomena of matter and energy, and cannot be observed and measured.

Evidence, Measurements, and Observations

Scientists use objective evidence over subjective evidence, to reach sound and logical conclusions. An **objective** observation is without personal bias and the same by all individuals. **Bias** refers to favoring one thing over another, and it can lead to inaccurate results. Humans are biased by nature, so they cannot be completely objective; the goal is to be as unbiased as possible. A **subjective** observation is based on a person's feelings and beliefs and is unique to that individual (figure $1.2.b$).



Figure 1.2.*b*: This is Grand Canyon of the Yellowstone in Yellowstone National Park. An objective statement about this would be, “The picture is of a waterfall.” A subjective statement would be, “The picture is beautiful.”

Another way scientists avoid bias is by using quantitative over qualitative measurements whenever possible. A **quantitative measurement** is expressed with a specific numerical value. **Qualitative observations** are general or relative descriptions. For example, describing a rock as red or heavy is a qualitative observation. Determining a rock’s color by measuring wavelengths of reflected light or its density by measuring the proportions of minerals it contains is quantitative. Numerical values are more precise than general descriptions, and they can be analyzed using statistical calculations. This is why quantitative measurements are much more useful to scientists than qualitative observations.

Science is a Social Process

Scientists do not work alone. Scientists need to be aware of research that has already been published and is in the works. Scientists also share their ideas with peers seeking guidance and feedback (figure 1.2.*c*). Research papers and data submitted for publication are rigorously reviewed by qualified peers, scientists who are experts in the same field. The scientific peer review process aims to weed out misinformation, invalid research results, and wild speculation. Thus, it is slow, cautious, and conservative.

Science is a work in progress. Hypotheses and theories are often confused. Hypotheses are tentative predictive explanations and are more specific and less general than a scientific theory. Scientific theories are the foundation of scientific knowledge. A **scientific theory** is a broad overarching explanation for a set of observations or phenomena. When a theory is new, it may have little supporting evidence and the scientific community may be skeptical of its acceptance. Scientists tend to favor the theory that can explain the most data. Throughout history, many theories have been discarded after realizing that evidence does not support them, e.g. the geocentric model, miasma, and alchemy. The best theories are supported by a wealth of observations and have stood up to rigorous hypothesis-testing by numerous scientists.Those theories that are supported by an overwhelming amount of evidence from many independent researchers are the ones we are most familiar with, for example, Cell Theory, Plate Tectonics, Climate Change and Theory of Evolution by Natural Selection.

In addition, in many scientific disciplines (less so in biology) there are **scientific laws**, often expressed in mathematical formulas, which describe how elements of nature will behave under certain specific conditions, but they do not offer explanations for why they occur.

Watch this 5-minute TED-Ed video highlighting the difference between a scientific law and theory.

Laws and Theories

TED-Ed(opens in new window) [www.youtube.com]



Figure 1.2.c: Scientists share information by publishing and attending conferences. The conference shown here focuses on peanut and mycotoxin (harmful chemicals produced by fungi) research. Image by Sharon Dowdy(opens in new window) (CC-BY-NC(opens in new window)).

Characteristics of Scientists

There is nothing mysterious or even particularly unusual about the things that scientists do. There are many ways to work on scientific problems. They all require common sense. Beyond that, they all display certain features that are especially - but not uniquely - characteristic of science.

- Skepticism — Good scientists use highly-critical standards in the judging of evidence. They approach data, claims, and theories (ideally, even their own!) with healthy doses of skepticism.
- Tolerance of uncertainty — Scientists often work for years - sometimes for an entire career - trying to understand one scientific problem. This often involves finding facts that, for a time, fail to fit into any coherent pattern and that even may support mutually contradictory explanations. Sometimes, as one listens to scientists vigorously defending their views, their confidence seems absolute. But deep in their hearts, they know that their views are based on probabilities and that a new piece of evidence may turn up at any time and force a major shift in their views.
- Although they certainly have no monopoly on hard work, their willingness to work long hours and years pursuing a problem is the mark of all good scientists. For science is hard work.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0

Modified by Melissa Ha from the following sources:

- What is Science?(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))
- The Process of Science from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Scientific Methods from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))

This page titled 1.2: What is Science? is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 2.1: What is Science? by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.
- 2.2: The Scientific Method by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.3: Empirical and Applied Science

Is it valuable to pursue science for the sake of simply gaining knowledge, or does scientific knowledge only have worth if we can apply it to solving a specific problem or bettering our lives? This question focuses on the differences between two types of science: basic science and applied science.

Empirical science, sometimes called "basic" or “pure” science seeks to expand knowledge regardless of the short-term application of that knowledge. It is not focused on solving a problem or developing a product or a service of immediate public or commercial value. The immediate goal of empircal science is knowledge for knowledge’s sake, though this does not mean that in the end it may not result in an application. Questions such as, "How have plants evolved to attract pollinators?" and "What factors determine which species will co-occur with each other?" fall under the scope of basic science (figure $1.3.a$).



Figure $1.3.a$: Many different plant species are co-occurring (growing together) at the Brisbane Botanic Gardens. Exploring which plants co-occur naturally falls under the scope of basic science. Image by Brisbane City Council(opens in new window) (CC-BY(opens in new window)).

In contrast, **applied science** focuses on solving real-world problems. Examples include increasing crop yields, reducing pollution, treating diseases, conserving endangered species, and developing renewable energy technologies. In applied science, the problem is often identified first, and scientists then use research and evidence to find solutions.

Empirical science focuses on understanding how the natural world works. Scientists ask questions, collect evidence, and develop explanations, even when there is no immediate practical application. Because the goals of empirical science are often centered on understanding rather than solving a specific problem, some people mistakenly view it as less useful than applied science.

However, the history of science shows that discoveries made through empirical science often lead to important applications. Many technologies and medical advances would not exist without decades of research aimed simply at understanding nature. For this reason, applied science often depends on knowledge gained through empirical science. At the same time, real-world problems can inspire new empirical research. The two approaches work together and support one another.

One example comes from the discovery of DNA structure. Scientists first sought to understand how genetic information is stored and passed from one generation to the next. This research was an example of empirical science because its goal was to explain a natural process. As scientists learned more about DNA and DNA replication, they developed new tools and technologies. These advances are now used to diagnose genetic diseases, identify individuals from biological evidence collected at crime scenes, and determine biological relationships among family members. These are examples of applied science.

This example shows how empirical science and applied science are connected. Scientific discoveries made to increase knowledge can eventually lead to practical solutions that benefit society. Likewise, solving real-world problems often requires a strong understanding of the underlying science.

Another example of the link between basic and applied research is the Human Genome Project, a study in which each human chromosome was analyzed and mapped to determine the precise sequence of the DNA code and the exact location of each gene. (The gene is the basic unit of heredity; an individual's complete collection of genes is his or her genome.) Other organisms have also been studied as part of this project to gain a better understanding of human chromosomes. The Human Genome Project (figure $1.3.b$) relied on basic research carried out with non-human organisms and, later, with the human genome. An important end goal eventually became using the data for applied research seeking cures for genetic diseases.

The discovery of penicillin, the first antibiotic, also originated from empirical scientific research. The mold *Penicillium* (figure $1.3.c$) contaminated petri dishes of bacteria and unexpectedly inhibited their growth. Read more about The Real Story behind Penicillin(opens in new window).



Figure $1.3.c$: The mold *Penicillium* growing on a petri dish. Image by Crulina 98(opens in new window) (CC-BY-SA(opens in new window)).

Ecological modeling is an example of how empirical science and applied science work together in environmental science and systems science. An ecological model is a simplified representation of a natural system. Many ecological models use mathematical equations and computer programs to predict how ecosystems may respond to different conditions or management decisions.

Scientists build ecological models using data collected through empirical research. For example, researchers may study how fast trees grow, how wildlife populations change over time, or how ecosystems respond to drought. This information helps scientists develop and improve models of natural systems.

Once a model has been developed, it can be used to address real-world problems. For example, a forest manager might use an ecological model to predict how different patterns of tree harvesting could affect forest health, biodiversity, and timber production. Environmental scientists might use models to predict the effects of climate change, habitat loss, or pollution on ecosystems.

Because ecological models rely on data and understanding gained through empirical science, they show how scientific knowledge can be used to make informed decisions. At the same time, real-world environmental challenges often lead scientists to collect new data and improve their models. In this way, empirical science and applied science continuously support one another.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from The Process of Science from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 1.3: Empirical and Applied Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 2.4: Basic and Applied Science by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.4: Human Impact on the Planet

Human Impact on the Planet: A Historical Perspective

Earth formed about 4.5 billion years ago. During that time, life has changed dramatically. More than 99.9% of all species that have ever lived are now extinct. Compared to Earth's long history, humans are a very recent species. Modern humans (*Homo sapiens*) have existed for only about 300,000 years.

Although humans have been on Earth for a relatively short time, our impact has been enormous. Humans use a large share of Earth's biological productivity—the energy captured by plants through photosynthesis. Large areas of land have been converted to farms, cities, roads, and other human uses. Cropland continues to expand, especially to produce food for a growing population and to support livestock production.

Today, species are disappearing much faster than they would under natural conditions. Scientists estimate that current extinction rates are tens to hundreds of times higher than the background extinction rate, and some estimates are even higher. Human activities such as habitat destruction, pollution, overexploitation, climate change, and the spread of invasive species are the primary causes of this increase.

People have modified their environment for thousands of years, but the scale and speed of environmental change have increased dramatically since the Industrial Revolution. Human activities now affect ecosystems, climate, water resources, and biodiversity on a global scale.

Several factors help explain why human impacts have become so large.

- Technological advances have greatly increased the amount of food, goods, and energy that people can produce. Most of these advances have been powered by fossil fuels such as coal, oil, and natural gas. While these technologies have improved many aspects of life, they have also increased resource use, waste production, and pollution. Humans now move more soil, rock, and other materials each year than natural processes such as rivers, wind, and erosion.

- The human population has grown rapidly. At the same time, consumption of energy, food, water, and materials has increased in many parts of the world. Environmental impacts depend not only on the number of people but also on how much each person consumes. Resource use is distributed unevenly, with some populations consuming far more resources than others. Together, population growth and rising consumption place increasing pressure on Earth's limited resources.

- Humans have released thousands of synthetic chemicals into the environment. Many of these chemicals do not occur naturally and can persist in air, water, soil, and living organisms for long periods of time. Some have been linked to environmental and health problems, while the long-term effects of many others remain poorly understood. As a result, chemical pollution has become a growing global concern.

Understanding how human activities affect Earth systems is one of the central goals of environmental science. By studying these impacts, scientists can help society make informed decisions about sustainability and the future of life on our planet.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha and Rachel Schleiger from Environment and Sustainability from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 1.4: Human Impact on the Planet is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 22.2: History of Sustainability by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.
- 2.2: The Scientific Method by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.5: Environmental Science

What is Environmental Science?

Environmental science is an interdisciplinary field that examines how living and nonliving components of the environment interact, with a strong focus on human impacts. It studies the conditions and surroundings in which organisms and communities exist, and the complex relationships that connect them within the natural world.

Environmental science includes disciplines in the physical sciences (e.g. geology, soil science, physical geography, chemistry, and atmospheric sciences), life sciences (e.g. ecology, conservation biology, restoration biology, and population biology), social sciences (e.g. human geography, economics, law, political science, and anthropology), and humanities (philosophy, ethics). As such, environmental scientists are a diverse bunch. However, they all come together for the focus of studying and identifying past, current, and future environmental issues while also exploring solutions to those environmental issues.




Figure $1.5.a$: Venn diagram illustrating how environmental science is interdisciplinary between humanities, natural sciences (which includes physical and life sciences) and social sciences. Image by Rachel Schleiger (CC-BY-NC).

Why Study Environmental Science?

Meeting the needs of a global population of over eight billion requires a fair, ethical, and sustainable use of Earth's systems. This depends on understanding how human actions affect the environment, as well as the science behind interactions between people and other species. Today, human driven environmental changes—from climate change to habitat loss—are happening faster and on a larger scale than ever before. To address these challenges, we must better understand how Earth's interconnected systems work, how human systems can be redesigned to be more sustainable, and how our choices scale up to collective impacts. Solving these complex problems will require informed, evidence-based decisions and active participation in creating long-term, sustainable solutions.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and

Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha and Rachel Schleiger Environmental Science (Ha and Schleiger) from The Earth, Humans, & the Environment, Environment and Sustainability, and Environmental Ethics from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 1.5: Environmental Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 1: Environmental Science Preface by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.
- 2.2: The Scientific Method by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

1.6: Practicing with Data- Wildlife Corridors

Overview

In 1999 the Sonoran Desert Conservation Plan was created to help wildlife populations and habitats. As a result of this plan, biologists were able to identify landscapes with wildlife barriers and get funding to install wildlife corridors. On State Route (SR) 77 two wildlife corridors were installed, one overpass and one underpass. These corridors were the first of their kind to be built in the Sonoran Desert so it was difficult to estimate what level of success they would have. As such, a monitoring plan was put in place to document which species used the corridors and how often they used it as time passed since their installation. The figure below illustrates the usage of the wildlife corridors 2+ years after their installation.

How to read a graph:

- Step 1: Read title.
- Step 2: Read axes.
- Step 3: Identify trend.
- Step 4: Explain trend.
- Step 5: Read any questions that follow and select the correct answer based on the graph.




Figure $1.6.a$: Total wildlife crossings after installation for overpass and underpass on SR 77. Graph by Erin Rempala modified from Rachel Schleiger (CC-BY-NC) modified from data in Santa Catalina - Tortolita Mountain Corridor Progress Report

Questions

1. What is the independent (explanatory) variable and the dependent (response) variable?
2. What question(s) are the authors trying to answer with this graph?
3. Does this graph show a positive or negative relationship between the variables? Support your answer by referring to trends observed in the graph.
4. Do you think like the authors are satisfied with the results in the graph? Why?
5. How can we use the results of this graph to inform future wildlife corridor installations?
6. What information/patterns is not clear from this graph?

Raw Data For Above Graph(s)

Table 1.6.*a*: Raw data for total wildlife crossings after installation for overpass and underpass on SR 77. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Santa Catalina - Tortolita Mountain Corridor Progress Report

Raw data for wildlife crossings graph

Month	Overpass	Underpass
April	0	0
June	150	150
August	249	250
October	296	294
December	600	500
February	750	502
April	1000	600
June	1200	605
August	1800	750
October	1550	950
December	1700	1050
February	2000	1500
April	2400	1850
June	2500	2000
August	2700	2250
October	3000	2500
December	3200	2750

Attribution

Modified by Erin Rempala (CC-BY-NC)

Rachel Schleiger (CC-BY-NC)

This page titled 1.6: Practicing with Data- Wildlife Corridors is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 5.5: Data Dive- Wildlife Corridors by Melissa Ha and Rachel Schleiger has no license indicated.

2: HUMANS AND THE ENVIRONMENT

How can humans live well within Earth's ecological limits?

Environmental science is an interdisciplinary field that uses science to understand environmental systems and solve environmental problems. This chapter introduces five key concepts—Planetary Boundaries, environmental ethics, environmental justice, sustainability, and environmental policy—that help explain how humans interact with the environment and respond to environmental challenges. Together, these concepts provide a foundation for understanding issues such as climate change, biodiversity loss, and use of ecosystem goods and services.

2.1: Planetary Boundaries- Earth's Safe Operating Space

2.2: Environmental Ethics and Worldviews

2.3: Environmental Justice and Equity

2.4: Sustainability and Sustainable Development

2.5: Systems Thinking and Modeling Sustainability

2.6: Historical Foundations of Environmental Policy

This page titled 2: Humans and the Environment is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 22: Sustainability by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

2.1: Planetary Boundaries- Earth's Safe Operating Space

The Anthropocene

Human activities now influence nearly every part of Earth's environment. Scientists have documented changes to the climate, biodiversity, land, water, and chemical composition of the planet. These changes are so widespread that some scientists use the term **Anthropocene** to describe the current period of Earth's history. The Anthropocene refers to this new era in which human activities have become a major force shaping Earth's systems.

The Planetary Boundaries Framework

The Planetary Boundaries Framework was proposed in 2009 by Johan Rockström and an international team of Earth system scientists (figure $2.1.a$). The framework identifies Earth-system processes that help maintain a stable environment for human societies. Together, these boundaries can be thought of as environmental guardrails that define a "safe operating space for humanity". Crossing a boundary does not necessarily cause immediate disaster, but it increases the risk of large-scale environmental changes that could affect ecosystems, human health, food production, and economic stability.

The framework includes **nine planetary boundaries**:

1. Climate Change
2. Biosphere Integrity (Biodiversity)
3. Land-System Change
4. Freshwater Change (Blue and Green Water)
5. Biogeochemical Flows (Nitrogen and Phosphorus Cycles)
6. Ocean Acidification
7. Atmospheric Aerosol Loading
8. Stratospheric Ozone Depletion
9. Novel Entities (synthetic chemicals, plastics, and other human-made substances)




Figure $2.1.a$: The 2025 update to the Planetary Boundaries Framework. The green zone in the center indicates the "safe operating space". As risk increases outside of that zone the color changes from yellowish-orange to reddish orange. Licensed under CC BY-NC-ND 3.0. Credit: "Azote for Stockholm Resilience Centre, based on analysis in Sakschewski and Caesar et al. 2025".

These boundaries are interconnected. Changes in one Earth system often affect others. For example, deforestation can contribute to climate change, reduce biodiversity, alter freshwater systems, and affect nutrient cycling. Because of these connections,

environmental problems are often best understood using **systems thinking**, which examines how different parts of a system influence one another.

Scientists regularly evaluate the status of the planetary boundaries. Current evidence suggests that seven of the nine boundaries have been exceeded. This does not mean societal collapse is inevitable, but it does indicate that human activities are increasing the risk of environmental disruptions that could affect both natural ecosystems and human societies.

The Planetary Boundaries framework provides a useful organizing model for environmental science because it connects many of the environmental challenges discussed throughout this course. Throughout the semester, you will revisit this framework as you explore topics such as climate change, biodiversity, water resources, and pollution.

For more information

For the most up-to-date information on the status of the nine planetary boundaries visit Planetary Health Check.

Attribution

Stockholm Resilience Centre. (n.d.). *Planetary boundaries*. Stockholm Resilience Centre. Retrieved July 21, 2026, from Planetary Boundaries

Planetary Health Check. (n.d.). *Planetary Health Check*. Retrieved July 21, 2026, from Planetary Health Check

This page titled 2.1: Planetary Boundaries- Earth's Safe Operating Space is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

2.2: Environmental Ethics and Worldviews

Environmental Ethics and Worldviews

Ethics are the principles people use to decide what is right and wrong. Environmental decisions often involve ethical questions because different people value nature in different ways. For example, some people may support clearing part of a rainforest to grow food because it benefits humans. Others may oppose it because it harms wildlife, ecosystems, and future generations.

An individual's **environmental ethic** reflects how they think about the relationship between humans and the environment. Some people value nature mainly for its **instrumental value**, meaning the benefits it provides to people, such as food, water, and recreation. Others believe nature also has **intrinsic value**, meaning it has worth regardless of its usefulness to humans. These beliefs help shape a person's **worldview**, or general way of understanding the world.

Environmental worldviews differ in who or what receives **moral consideration**. At the most narrow level, people may focus primarily on themselves (**egocentrism**). Moral concern can expand to include other people, future generations, and all of humanity (**anthropocentrism**). When moral concern extends beyond humans, it is called **moral extensionism**.

As moral concern expands, different environmental ethics emerge. **Biocentrism** extends moral concern to all living organisms. **Ecocentrism** extends moral concern to ecosystems and ecological processes. Some people extend moral concern to the entire biosphere, including all living things and the systems that support life on Earth (Figure $2.2.a$).




Figure $2.2.a$: Continuum of worldviews from self-centered to biosphere-centered perspectives. Image by Erin Rempala (CC-BY-NC 4.0).

These different ethical perspectives help explain why people may disagree about environmental issues. For example, a person may support protecting wetlands because they provide clean water and reduce flooding (an anthropocentric justification). Another person may argue that wetlands should also be protected because the organisms living there that have their own moral value (a biocentric perspective).

Environmental ethics can also be connected to sustainability and the Planetary Boundaries framework. People may support protecting Earth's life-support systems for different reasons. Some may focus on protecting human well-being and future generations, while others may also extend moral concern to other species, ecosystems, or the biosphere as a whole. Understanding environmental ethics helps explain why societies make different choices about conservation, sustainability, and environmental policy.

A Simplified Framework

Environmental ethicists have developed many theories and frameworks for evaluating human relationships with nature, including the land ethic, deep ecology, ecofeminism, social ecology, Indigenous philosophies, animal ethics, and others. This section uses a simplified framework based on moral consideration and worldviews to introduce some of ways people think about environmental issues. These categories are useful for understanding broad patterns in environmental decision-making, but individual beliefs often combine ideas from multiple ethical perspectives.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Erin Rempala from Hourdequin. M. 20215. *Environmental Ethics: From Theory to Practice*. Bloomsbury Academic.

Modified by Rachel Schleiger from Sustainability: A Comprehensive Foundation(opens in new window) by Openstax (licensed under CC-BY(opens in new window))

This page titled 2.2: Environmental Ethics and Worldviews is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 22.3: Environmental Ethics by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

2.3: Environmental Justice and Equity

Environmental Equity

Environmental equity means that no group should bear an unfair burden of environmental harms or receive fewer environmental benefits. Differences in resource consumption, pollution, and exposure to environmental hazards raise important questions about fairness and sustainability.

Environmental Racism and Environmental Justice

Environmental problems do not affect all people equally. During the 1980s, communities across the United States began organizing against the placement of hazardous waste facilities and other polluting industries near their homes. In 1987, civil rights activist and author, Dr. Benjamin Chavis introduced the term **environmental racism** to describe situations in which people of color and other marginalized communities face a disproportionate share of environmental hazards, such as pollution, toxic waste, and harmful industrial activities (figure $2.3.a$). Environmental racism can result from a combination of historical patterns of housing discrimination, economic inequality, political power, and land-use decisions.



Figure $2.3.a$: MeetDrBen; cropped by Beyond My Ken (talk) 03:33, 14 February 2014 (UTC), CC BY-SA 3.0 <https://creativecommons.org/licenses/by-sa/3.0>, via Wikimedia Commons

These concerns helped inspire the **environmental justice** movement (figure $2.3.b$). Environmental justice is the principle that all people should receive equal protection from environmental and health hazards and have meaningful opportunities to participate in environmental decision-making. Environmental justice recognizes that environmental burdens, such as air pollution, contaminated water, and hazardous waste, have often fallen disproportionately on low-income communities, communities of color, and Tribal nations.One of the most influential leaders of this movement is Dr. Robert Bullard, who is often called the "father of environmental justice" because of his research and advocacy documenting environmental racism and promoting environmental equity throughout the United States.



Figure 2.3.b: Dr. Robert D. Bullard, Distinguised Professor and former Dean of the Barbara Jordan-Mickey Leland School of Public Affairs at Texas Southern University, October 2012. Image by David Brenner in Flickr(opens in new window) (CC-BY-SA-2.0).

One well-known example is the **Flint Water Crisis** in Michigan. In 2014, Flint changed its drinking water source to reduce costs, but the new water was not properly treated and caused lead from aging pipes to contaminate the drinking water supply. The crisis exposed 100,000 residents, including many children, to elevated levels of lead, a toxic metal than can cause brain damage, learning disabilities and other long-term health problems. Because Flint is a low-income, majority-Black community, many researchers and community advocates view the crisis as an example of environmental injustice. The case illustrates how environmental decisions can have unequal impacts on different communities and why fairness is an important consideration in environmental policy.

Indigenous Peoples and Environmental Justice

Indigenous peoples have lived on and cared for their lands for thousands of years and continue to play important roles in environmental stewardship today. However, colonization often displaced Indigenous communities from their ancestral lands and limited their ability to participate in decisions affecting natural resources. As a result, many environmental justice issues involve Indigenous rights, land access, cultural preservation, and self-determination.

Environmental decisions can affect Indigenous communities in different ways. For example, some Indigenous groups, such as the Gwich'in of Alaska, oppose oil drilling near the Arctic National Wildlife Refuge because they believe it threatens important cultural traditions, wildlife, and ecosystems (figure 2.3.b). At the same time, some Iñupiat communities, have supported oil development because of potential economic benefits. These differences highlight that Indigenous peoples are not a single group with one perspective. Environmental justice requires respecting Indigenous sovereignty and ensuring that Tribal nations can meaningfully participate in decisions that affect their lands, resources, and futures.



Figure 2.3.*b*: Bernadette Demietieff speaking in support of protect the Arctic to protect Gwich'in land from oil drilling, Alaska, c. 2019. Image by: Deb Haaland(opens in new window) in Wikimedia Commons(opens in new window) (CC-BY-SA-4.0(opens in new window)).

Just Transition

A **just transition** is the idea that society should move toward cleaner, more sustainable ways of producing energy, food, and other goods while also protecting workers and communities. Supporters argue that environmental solutions should improve both environmental and social conditions rather than create new inequalities. For example, communities that depend on fossil fuel industries may need job training, education, and economic support as society shifts toward renewable energy. A just transition seeks to create a future that is both environmentally sustainable and socially fair by ensuring that the benefits and responsibilities of environmental change are shared more equitably.

A Simplified Framework

Environmental justice is a complex field that includes many perspectives and approaches. This section introduces several key ideas that help explain how environmental benefits and burdens are distributed among people and communities, but it is not a complete overview of the field.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Erin Rempala from:

- Haudenosaunee Confederacy. (n.d.). *Values of the Haudenosaunee Confederacy*. Retrieved July 20, 2026, from Haudenosaunee Confederacy.
- Climate Justice Alliance. (n.d.). *Just transition*. Retrieved July 20, 2026, from Climate Justice Alliance. (n.d.). Just transition. Retrieved July 20, 2026, from Climate Justice Alliance [climatejusticealliance.org]
- Wikipedia contributors. (2026, July 20). *Flint water crisis*. In *Wikipedia*. Retrieved July 20, 2026, from Flint Water Crisis [en.wikipedia.org]
- Wikipedia contributors. (n.d.). *Haudenosaunee*. In *Wikipedia*. Retrieved July 20, 2026, from Wikipedia: Haudenosaunee

Modified by Melissa Ha and Rachel Schleiger from Environmental Justice and Indigenous Struggles and Environment and Sustainability from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 2.3: Environmental Justice and Equity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 22.3.1: Environmental Justice and Indigenous Struggles by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

2.4: Sustainability and Sustainable Development

Sustainability and Sustainable Development

Sustainability is the idea that humans should meet their needs without limiting the ability of future generations to meet their own needs. Although the term is relatively modern, many Indigenous cultures have long emphasized responsibilities to future generations and long-term relationships with the natural world.

In 1987, the United Nations Brundtland Commission defined **sustainable development** as "development that meets the needs of the present without compromising the ability of future generations to meet their own needs." This definition highlights the connection between environmental health, social well-being, and economic prosperity.



Figure 2.4.a: UN Sustainable Development Goals. Image by United Nations Sustainable Development Goals (CC-BY-SA4.0). The content of this publication has not been approved by the United Nations and does not reflect the views of the United Nations or its officials or Member States.

Sustainability recognizes that environmental problems are often linked to social and economic systems. Solutions that focus only on one part of the system may create new problems elsewhere. As a result, sustainability encourages people to consider environmental, social, and economic impacts together. Some scholars argue that sustainability should focus not only on maintaining systems but also on restoring degraded ecosystems and improving environmental conditions. These ideas are reflected in concepts such as restoration, regeneration, and regenerative sustainability.

Today, sustainability provides a framework for addressing environmental challenges while considering the needs of people, their communities, future generations, and the ecosystems that support life on Earth.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Erin Rempala from:

- Sustainable Development Report website based on Sustainable Development Report(opens in new window) [doi.org]

Modified by Rachel Schleiger from Sustainability: A Comprehensive Foundation by Openstax (licensed under CC-BY)

This page titled 2.4: Sustainability and Sustainable Development is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 22.1: Introduction to Sustainability by Melissa Ha and Rachel Schleiger has no license indicated.

2.5: Systems Thinking and Modeling Sustainability

Systems Thinking and Models

A **system** is a group of interacting and interdependent components that work together as a larger whole. Because the parts of a system are connected, changes in one component can influence the behavior of the entire system. Systems are often **nested** within larger systems, meaning that one system may be a component of a larger system.

For example, in the human body, red blood cells are part of blood, blood is part of the circulatory system, and the circulatory system is part of the body. Changes in one part of the system can influence the larger systems that depend on it.

Systems thinking is an approach to understanding complexity by focusing on relationships and interactions among components rather than examining each part in isolation. We will use systems thinking throughout this course to explore environmental challenges and evaluate potential solutions.

A **model** is a simplified representation of an object, process, or system. People use models to better understand systems, their components, and the relationships among those components. Although models simplify reality, they can help us understand complex environmental systems and guide decision-making related to environmental science and sustainability.

Modeling Sustainability

One common representation of sustainability is the **Triple Bottom Line**, which represents environmental, social, and economic goals as three equally important parts of a healthy system (figure $2.5.a$). Originally developed for business, this model encourages decision-makers to consider environmental and social outcomes in addition to economic growth.

However, some critics argue that it suggests that environmental, social, and economic goals are separate and equally important, rather than recognizing that human societies and economies are ultimately nested within and dependent upon the biosphere.




Figure $2.5.a$: The Triple Bottom Line. Image by Erin Rempala (CC-BY-SA4.0).

Ecological Economist, Herman Daly, created the **Nested Dependencies Model** (figure \(\PageIndex{b}\)). In this model, the economy exists within society, and society exists within the biosphere. This arrangement emphasizes that human societies and economies depend on Earth's ecosystems for energy, materials, waste processing, and life-support services. The model highlights

that economic activity must operate within ecological limits.

This idea connects closely to the Planetary Boundaries framework, which identifies environmental limits that help maintain a stable Earth system.




Figure \(\PageIndex{b}\): The Nested Dependencies Model. Image by Erin Rempala redrawn from Daly and Farley (CC-BY-SA4.0).

Renegade economist Kate Raworth expanded this idea in her **Doughnut Model** (figure $2.5.c$). The doughnut's outer ring represents Earth's ecological limits (planetary boundaries), while the inner ring represents the social foundation needed for human well-being, such as food, water, education, and healthcare. The space between these boundaries is called the **safe and just space for humanity**, where people's needs can be met without exceeding Earth's ecological limits.

Together, these models provide different ways of thinking about the relationship between humans and the natural world.




Figure $2.5.c$: The Doughnut from Doughnut Economics. The safe and just space for humanity is within the doughnut (green) between the social foundation and the ecological ceiling. Arrows inside the social foundation mean that basic needs are not being met. Arrows outside the ecological ceiling mean that we have exceeded the planetary boundaries. Image by By DoughnutEconomics - Own work, CC BY-SA 4.0, wikimedia [commons.wikimedia.org]

A Simplified Framework

The Triple Bottom Line, Nested Dependencies, and the Doughnut models each provide a different way of thinking about relationships among the environment, society, and the economy. Although these models simplify complex real-world systems, they can help us evaluate different approaches to environmental decision-making.

Which model best explains the relationship between the economy, society, and the environment?

Attribution:

- Daly, H. E., & Farley, J. (2011). *Ecological economics: Principles and applications* (2nd ed.). Island Press.
- Raworth, K., & Doughnut Economics Action Lab. (n.d.). *Doughnut Economics Action Lab*. Retrieved July 20, 2026, from Doughnut Economics [doughnuteconomics.org]

This page titled 2.5: Systems Thinking and Modeling Sustainability is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

2.6: Historical Foundations of Environmental Policy

Historical Foundations of Environmental Policy

Environmental policies are one way societies respond to environmental challenges. In the United States, environmental policy developed as European and later U.S. settlers occupied Indigenous lands and established new political and economic systems in North America. Indigenous peoples had already cared for these lands for thousands of years using diverse knowledge systems and stewardship practices. However, many settler societies increasingly viewed land, forests, water, minerals, and wildlife primarily as resources that could support agricultural expansion, industrialization, and economic growth. Some environmental historians argue that these economic and political systems shaped both environmental problems and the policies created to address them.

During the late 1800s and early 1900s, concerns about the rapid use of forests, wildlife, and other natural resources contributed to the rise of conservation and preservation efforts. Leaders such as Theodore Roosevelt and Gifford Pinchot promoted the careful management of natural resources so they would remain available for future generations. At the same time, thinkers such as Aldo Leopold emphasized that humans are members of ecological communities and have ethical responsibilities to care for the land (**The Land Ethic**). These ideas helped establish the principle that governments and citizens share responsibility for protecting the environment. Historians also note that some conservation policies displaced Indigenous communities from lands that later became parks, forests, and protected areas.

By the mid-twentieth century, environmental concerns shifted from resource depletion to pollution and public health. In 1962, Rachel Carson's *Silent Spring* showed how pesticides such as DDT could harm wildlife, ecosystems, and human health. Growing scientific evidence led to major environmental reforms, including the creation of the **Environmental Protection Agency (EPA)** in 1970 and the passage of landmark laws such as the **Clean Air Act** and **Clean Water Act**. These policies marked a growing commitment to preventing environmental harm before serious damage occurred.

Beginning in the 1970s, environmental justice advocates drew attention to the unequal distribution of environmental risks. Research and community activism showed that low-income communities, communities of color, and Tribal nations often experienced higher exposure to pollution and environmental hazards. As a result, environmental policy expanded beyond conservation and pollution control to include concerns about fairness, public participation, and equal access to healthy environments.

A Simplified Framework

The history of environmental policy in the United States is complex and includes many people, ideas, laws, social movements, and historical events.

This section briefly introduces major ideas and events that have shaped environmental policy in the United States. While these themes are useful for understanding broad patterns in environmental policy, they do not capture every perspective, event, or debate that has shaped environmental decision-making in the United States.

Today, environmental policy addresses a wide range of challenges, including climate change, biodiversity loss, water scarcity, environmental justice, and sustainable resource management. Environmental laws and regulations continue to evolve as new scientific evidence emerges and as societies debate how best to balance environmental protection, economic development, and social well-being. Understanding the history of environmental policy helps explain why environmental decisions often involve scientific, ethical, economic, and political considerations.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

This page titled 2.6: Historical Foundations of Environmental Policy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

3: ECOLOGY

Introduction

Ecology is the study of how living things interact with each other and their environment. In this chapter, you will explore how energy and matter move through ecosystems, and how populations of organisms grow, interact, and respond to changes in their environment. You will learn how factors like food, space, predators, and climate affect where organisms live and how many can survive. We will also connect these ideas to real-world issues, such as habitat loss, pollution, and conservation, including local ecosystems you will visit in this course. By the end of the chapter, you will be able to use data, graphs, and models to better understand how ecosystems function and how human actions impact the natural world.

3.1: Population Ecology

3.1.1: Ecological Hierarchy

3.1.2: Population Distribution

3.1.3: Population Size

3.1.4: Population Growth and Regulation

3.1.5: Life History

3.2: Community Ecology

3.2.1: Biotic Interactions

3.2.1.1: Symbiosis

3.2.1.2: Trophic Interactions

3.2.1.2.1: Predation

3.2.1.2.2: Herbivory

3.2.1.2.3: Competition

3.2.2: Community Structure and Dynamics

3.2.3: Data Dive- Beaver Impacts on Wetlands

3.3: Ecosystem Ecology

3.3.1: Matter

3.3.2: Energy

3.3.3: Biogeochemical Cycles

3.3.4: Energy Flow in Food Webs

3.3.5: Ecosystem Types and Dynamics

3.3.5.1: Climate and Biomes

3.3.5.2: Terrestrial Biomes

3.3.5.3: Aquatic Biomes

3.3.6: Data Dive- Biome Carbon Storage

Attributions

Modified by Erin Rempala.

Thumbnail western toad wikimedia.org by Thompsma - Own work, (CC BY 3.0).

This page titled 3: Ecology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

3.1: POPULATION ECOLOGY

Population Ecology

Populations are interacting and potentially interbreeding groups of individuals from the same species living in a common area. The study of **population ecology** focuses on population size and the factors that regulate population growth.

Wildlife Corridors

Roadkill is a familiar reminder of one of the unintended consequences of our extensive road networks. Roads, especially freeways and highways, not only increase wildlife mortality but also divide habitats into smaller, isolated patches. This **habitat fragmentation** can make it more difficult for animals to find food, shelter, and mates. Over time, fragmented habitats can reduce biodiversity, limit gene flow between populations, and increase the risk of local extinctions.

To address this problem, ecologists and conservation planners have developed **wildlife crossings**, such as overpasses and underpasses, and **wildlife corridors**, which help reconnect fragmented habitats and allow animals to move more safely across landscapes.



Figure $3.1.a$: A newly constructed wildlife overpass over a Singapore highway. Image by Benjamin P. Y-H. Lee(opens in new window) (licensed under CC-BY-4.0(opens in new window))

3.1.1: Ecological Hierarchy

3.1.2: Population Distribution

3.1.3: Population Size

3.1.4: Population Growth and Regulation

3.1.5: Life History

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and

Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Rachel Schleiger and Melissa Ha (CC-BY-NC(opens in new window))

This page titled 3.1: Population Ecology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 5: Populations by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.1.1: Ecological Hierarchy

Ecological Hierarchy

Ecological hierarchy describes the upper levels of biological organization, with each level nested within larger systems. Individual living organisms form the base of this hierarchy. Example organisms include humans (*Homo sapiens sapiens*), black bread mold fungi (*Rhizopus stolonifer*), and African lions (*Panthera leo*). When individuals of the same species live in the same area and can reproduce with one another, they form a **population**. Two or more interacting populations make up a **community**. When communities interact with each other and with nonliving environmental factors, called **abiotic factors**, they form an **ecosystem**. All ecosystems on Earth together make up the **biosphere**, which includes all regions of the planet where life exists (figure $3.1.1.a$).

This hierarchy illustrates an important principle of systems thinking: systems are often nested within larger systems. Individuals are part of populations, populations are part of communities, communities are part of ecosystems, and ecosystems are part of the biosphere. Because these systems are interconnected, changes at one level can affect other levels. For example, a decline in a population may alter community interactions, affect ecosystem processes, and influence the biodiversity of larger regions. Understanding these connections helps environmental scientists study complex environmental problems and evaluate potential solutions.




Figure $3.1.1.a$. Biological Organization from individuals to biosphere. Edited by Erin Rempala from Mikala14, (CC BY-SA 3.0), via Wikimedia Commons. (CC BY-SA 4.0)

This page titled 3.1.1: Ecological Hierarchy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

3.1.2: Population Distribution

The **distribution pattern** of a population describes the arrangement of individuals within a habitat at a particular point in time, and broad categories of patterns are used to describe them. Individuals in a population may be distributed in three common patterns:**random**, **clumped**, or **uniform** (figure $3.1.2.a$).




Figure $3.1.2.a$: Species may have a random, clumped, or uniform distribution. Plants such as (a) dandelions with wind-dispersed seeds tend to be randomly distributed (represented by dots in no particular pattern). Animals such as (b) elephants that travel in groups exhibit a clumped distribution (represented by dots clustered together). Territorial birds such as (c) penguins tend to have a uniform distribution (represented by dots evenly spaced from each other). (credit a: modification of work by Rosendahl; credit b: modification of work by Rebecca Wood; credit c: modification of work by Ben Tubby)

A **random distribution** occurs when individuals are arranged without a clear pattern. An example is dandelions and other plants with wind-dispersed seeds that grow wherever conditions are suitable.

A **clumped distribution** occurs when individuals are grouped together in patches. This often happens when resources are unevenly distributed or when organisms benefit from living near one another. For example, wolves often live and hunt in packs.

A **uniform distribution** occurs when individuals are evenly spaced. This pattern often results from competition for resources or territorial behavior. For example, saguaro cacti are spread out to reduce competition for water, and some animals maintain territories that keep individuals separated.

The distribution pattern of a population can provide important clues about how individuals interact with one another and with their environment.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Population Demographics and Dynamics from *Environmental Biology* by Matthew R. Fisher (CC-BY(opens in new window))

This page titled 3.1.2: Population Distribution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 5.1: Population Dispersion by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.1.3: Population Size

Population Size, Density, and Sampling

Populations are not fixed. Their size and composition can change over time in response to factors such as seasonal weather changes, natural disasters, disease, predation, competition, and the availability of resources. The study of populations is called **demography**.

Two important characteristics of a population are **population size** and **population density**. **Population size** is the total number of individuals in a population. **Population density** is the number of individuals living in a given area. For example, two populations may contain the same number of individuals but have different densities if one population is spread over a larger area than the other (figure $3.1.3.a$).




Figure $3.1.3.a$: Both populations shown contain seven plants, so their populations size is the same. However, the population on the left has a higher population density (2 plants per square) than the population on the right (density = 0.5 plants per square). Image by Erin Rempala modified from Melissa (public domain).

Population size and density can influence how organisms interact with one another. In dense populations, individuals may compete more strongly for food, water, space, or mates. Diseases may also spread more easily when individuals live close together. Population size can also affect genetic diversity. Small populations often have less genetic variation, which can make it more difficult for them to adapt to environmental change.

Because it is usually impossible to count every individual in a population, scientists often estimate population size by studying a sample of the population. The method used depends on the type of organism being studied.

Estimating Population Size

For plants and other organisms that do not move, scientists often use a **quadrat**, a square frame placed at random locations within a habitat. By counting the organisms inside multiple quadrats, scientists can estimate the size and density of the population (Figure $3.1.3.c$).



Figure 3.1.3.c: High school students using a quadrat in an intertidal zone. Image by kqedquest (CC-BY-NC)

For mobile organisms such as small mammals, scientists often use **mark-and-recapture** methods. First, a sample of animals is captured, marked, and released. Later, a second sample is collected and scientists record how many marked individuals are recaptured. This information can be used to estimate the size of the entire population. Although the estimate is not exact, it provides a useful way to study populations that would otherwise be difficult to count.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Population Demographics and Dynamics from *Environmental Biology* by Matthew R. Fisher (CC-BY)

This page titled 3.1.3: Population Size is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 5.2: Population Size by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.1.4: Population Growth and Regulation

Population Growth and Population Regulation

Populations change over time. Some populations increase in size, while others decrease. Ecologists study these patterns to understand how populations respond to environmental conditions and predict how they might change in the future.

The Population Growth Rate (*r*)

A population's **growth rate** depends on the balance between births, deaths, immigration and emigration. If more individuals are born than die (or immigrate), the population grows. If more individuals die than are born, or emigrate, the population shrinks. For example, if a caribou population increases from 200 individuals to 300 individuals over two years, the population has experienced positive growth. Scientists use population growth rates to compare populations and understand how quickly they are changing (figure $3.1.4.a$).



Figure $3.1.4.a$: How could we calculate the growth rate of this caribou population per year if there were 200 individuals in 2016 and 300 individuals in 2018? Image by K. Joly(opens in new window) [K. Joly] (public domain).

Doubling Time

The **doubling time** is the amount of time it takes for a population to double in size. Populations with high growth rates have shorter doubling times than populations with low growth rates. For example, a rapidly growing population of bacteria may double in a matter of hours, while a population of large mammals may take many years to double in size.

Exponential Growth

When resources are abundant and environmental conditions are favorable, populations can experience **exponential growth** for a time. In exponential growth, the population grows faster and faster over time because there are more individuals reproducing each generation. Exponential growth produces a **J-shaped curve** when population size is plotted over time. Bacteria growing in a laboratory flask with plenty of nutrients often show exponential growth for a short period. Although exponential growth can occur temporarily, it rarely continues for long in nature because resources are limited.

Logistic Growth and Carrying Capacity

In real ecosystems, resources such as food, water, space, and shelter eventually become limited. As resources become scarce, population growth slows. This pattern is called **logistic growth** and produces an **S-shaped curve** ({3{.}1{.}4{.}\eatSpaces{b}\endSpaces}\)). Logistic growth includes three stages:

1. Rapid growth when resources are abundant.
2. Slower growth as resources become limited.
3. Stabilization near the **carrying capacity**.




Figure 3.1.4.*b*: These graphs show exponential and logistic growth, with population size on the y-axis and time on the x-axis. When resources are unlimited, populations exhibit (a) exponential growth, shown in a J-shaped curve. When resources are limited, populations exhibit (b) logistic growth. In logistic growth, population expansion decreases as resources become scarce, and it levels off when the carrying capacity of the environment is reached. The logistic growth curve is S-shaped.

The **carrying capacity (K)** is the largest population size that an environment can support over time.

Many natural populations fluctuate around their carrying capacity rather than remaining at a constant size. For example, deer populations, fish populations, and seal populations often increase and decrease slightly from year to year as environmental conditions change (figure 3.1.4.*c*).




Figure 3.1.4.*c*: A natural population of seals shows logistic growth. The number of seals, ranging from 1500 to 8500, is on the y-axis, and the year, ranging from 1975-2000, is on the x-axis. Orange dots represent the observed population size each year, and the blue line represent the trend. From 1975 to 1983, the blue line curves upwards, representing an increase in both population size and population growth rate. From 1983 to 1992, the curve flattens, representing that population size is still increasing, but population growth rate is decreasing. After 1992, population fluctuates near the carrying capacity (dotted, horizontal line).

Factors That Affect Population Size

Population growth is influenced by many factors. Some factors become stronger as population density increases. These are called **density-dependent factors** and include:

- Competition for food, water, and space
- Disease
- Parasitism
- Predation

For example, diseases often spread more rapidly in crowded populations, and competition for food becomes more intense as population size increases.

Other factors affect populations regardless of their density. These are called **density-independent factors** and include:

- Droughts
- Floods
- Hurricanes (figure 3.1.4.*d*)
- Wildfires
- Extreme temperatures
- Pollution

For example, a wildfire can reduce a deer population whether there are many deer or very few deer in the area.



Figure 3.1.4.*d*: Severe weather, like Hurricane Florence of 2018, can reduce population size regardless of density. Image by NASA Goddard Space Flight Center(opens in new window) (CC-BY(opens in new window))

Population Regulation as a System

Population growth is another example of systems thinking. Birth rates, death rates, resources, competition, predators, disease, and environmental conditions all interact to influence population size. Changes in one part of the system can affect many other parts. Because populations are connected to communities and ecosystems, changes in population size can influence biodiversity, food webs, and ecosystem processes. Understanding these relationships helps scientists predict environmental change and develop effective conservation and management strategies.

A Simplified Framework

Population ecology is a large field of study that includes many mathematical models and methods. This section introduces a few key concepts, including population growth models, carrying capacity, and population regulation. These ideas provide a useful foundation for understanding how populations change over time, but they do not represent all of the factors that influence population dynamics in nature.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Population Growth and Regulation from *Environmental Biology* by Matthew R. Fisher (CC-BY(opens in new window))
- Principles of Population Growth and The Human Population from *Biology* by John W. Kimball (CC-BY(opens in new window))
- Population Dynamics and Regulation from *General Biology* by OpenStax (CC-BY(opens in new window))

This page titled 3.1.4: Population Growth and Regulation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 5.3: Population Growth and Regulation by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.1.5: Life History

Life Histories and Survivorship

Not all species live, reproduce, and die in the same way. Some species produce a small number of offspring and invest a great deal of energy in caring for them. Others produce many offspring, but provide little or no parental care. Ecologists study these patterns, known as **life histories**, to better understand how populations change over time.

Survivorship Curves

One way ecologists study life histories is by using a **survivorship curve**, which shows how many individuals survive to different ages. There are three common patterns (figure $3.1.5.a$).

- **Type I:** Most individuals survive to old age, and mortality increases later in life. Humans, elephants, and many large mammals show this pattern.
- **Type II:** Individuals have a fairly constant chance of dying throughout their lives. Many bird species show this pattern.
- **Type III:** Many individuals die at a young age, but those that survive tend to live much longer. Many fish, insects, marine invertebrates, and plants show this pattern.

Survivorship curves help scientists compare different life histories and understand how species allocate energy to growth, reproduction, and survival.




Figure $3.1.5.a$: Survivorship curves show the chance of survival in a population according to age. The number of individuals surviving (on a log scale) is on the y-axis, and time (representing relative age) is on the x-axis. Humans and most mammals have a Type I survivorship curve, because death primarily occurs in the older years. The Type I survivorship curve is represented by a convex curve, which rapidly declines at the end. Birds have a Type II survivorship curve, as death at any age is equally probable. The Type II survivorship curve is represented by a straight line, going from the upper left to lower right of the graph. Trees have a Type III survivorship curve because very few survive the younger years, but after a certain age, individuals are much more likely to survive. The Type III survivorship curve is represented by a concave curve, which initially declines rapidly and then flattens.

Reproductive Strategies

Species also differ in how they reproduce. Some species produce only a few offspring, while others produce many. These differences can be viewed as a continuum between two broad reproductive strategies.

Species on one end of the continuum (**K-selected**) produce relatively few, large offspring and often provide parental care. These organisms tend to mature slowly, live longer, and invest more energy in each offspring. Examples include humans, elephants, and many large mammals.

Species on the other end of the continuum (**r-selected**) produce many small offspring and provide little or no parental care. These organisms often mature quickly, reproduce early, and rely on large numbers of offspring to ensure that some survive. Examples include jellyfish, many insects, and dandelions (f igure $3.1.5.b$).



(a) K-selected species



(b) r-selected species

Figure $3.1.5.b$: (a) Elephants are considered K-selected species as they live long, mature late, and provide long-term parental care to few offspring. Oak trees produce many offspring that do not receive parental care, but are considered K-selected species based on longevity and late maturation. (b) Dandelions and jellyfish are both considered r-selected species as they mature early, have short lifespans, and produce many offspring that receive no parental care.

A Continuum, Not Categories

Although scientists once classified species as either K-selected or r-selected, modern ecology recognizes that most species fall somewhere between these two extremes. For example, some organisms may produce many offspring but also invest significant energy in their survival. As a result, ecologists now focus on specific life-history traits rather than placing species into strict categories.

Systems Thinking and Life Histories

Life histories are influenced by many interacting factors, including resource availability, predation, competition, disease, climate, and environmental change. Because these factors are connected, life histories provide another example of systems thinking. Changes in environmental conditions can affect survival and reproduction, which in turn influence population growth and ecosystem dynamics.

A Simplified Framework

Life-history theory is a large and active area of ecological research. This section introduces a few basic concepts, including survivorship curves and reproductive strategies. These ideas help explain how populations survive and reproduce, but they do not capture the full complexity of life-history variation found in nature.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Population Dynamics and Regulation from *General Biology* by OpenStax (CC-BY(opens in new window))
- Population Demographics and Dynamics from *Environmental Biology* by Matthew R. Fisher (CC-BY(opens in new window))
- Principles of Population Growth from *Biology* by John W. Kimball (CC-BY(opens in new window))

This page titled 3.1.5: Life History is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 5.4: Life History by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2: COMMUNITY ECOLOGY

Community Ecology

Populations typically do not live in isolation from other species. Populations that live and interact within a given area form a **community**. The organisms that make up a community share a **habitat**, the physical environment in which they live. A community includes all of the **biotic** (living) components of an area, such as plants, animals, fungi, bacteria, and other organisms. Scientists study ecology at the community level to understand how species interact with one another and how these interactions influence the distribution and abundance of organisms.

Keystone Species

Until the late 1800s, American bison (*Bison bison*) numbered in the tens of millions across the Great Plains (figure $3.2.a$). By 1890, fewer than 1,000 remained due to overhunting, habitat loss, and U.S. government policies that contributed to the displacement of Indigenous peoples and the destruction of the resources they depended upon. Conservation efforts eventually helped bison populations recover, reaching roughly half a million individuals by the early 2000s.

As bison were reintroduced to parts of the Great Plains, ecologists observed major changes in the ecosystem. Bison grazing, wallowing, trampling, and movement across the landscape helped maintain plant diversity and created habitat conditions that benefited many other species. Because bison physically modify their environment and influence the availability of resources for other organisms, they are considered an **ecosystem engineer**. Bison are also considered a **keystone species** because their presence has a disproportionately large effect on the structure and function of Great Plains ecosystems. This example demonstrates how changes to a single species can affect an entire ecological community and highlights the importance of understanding community interactions when conserving and restoring ecosystems.



Figure $3.2.a$**:** American bison with starlings on its back. Image by NPS Photo(opens in new window) [NPS Photo] [NPS photos] (Public Domain)

3.2.1: Biotic Interactions
3.2.1.1: Symbiosis
3.2.1.2: Trophic Interactions
3.2.1.2.1: Predation
3.2.1.2.2: Herbivory
3.2.1.2.3: Competition
3.2.2: Community Structure and Dynamics
3.2.3: Data Dive- Beaver Impacts on Wetlands

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains content adapted from 6: Communities originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Rachel Schleiger and Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 3.2: Community Ecology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 6: Communities by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2.1: BIOTIC INTERACTIONS

Biotic Interactions

Biotic interactions are relationships among living organisms. These interactions can occur between members of the same species (**intraspecific interactions**) or between members of different species (**interspecific interactions**).



Figure 3.2.1.a: A variety of biotic interactions occur among the species at The Seal Beach National Wildlife Refuge (top). For example, the Light-Footed Ridgway Rail (bottom) benefits from the cordgrass in which it nests. However, predators such as red foxes harm this endangered bird. The refuge manages this harmful interaction by installing fencing to exclude the red fox, which historically did not occur in the area. Top image(opens in new window) and bottom image(opens in new window) by US Fish & Wildlife Service (public domain).

Some interactions benefit one organism while harming another. These are called **antagonistic interactions**. Examples include **predation**, in which one organism eats another, and **competition**, in which organisms compete for the same limited resources such as food, water, space, or mates.

Other interactions benefit at least one species without harming the other. These are called **facilitative interactions**. In **commensalism**, one species benefits while the other is neither helped nor harmed. In **mutualism**, both species benefit from the interaction.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))



- Biotic interactions

3.2.1.1: Symbiosis

3.2.1.2: Trophic Interactions

3.2.1.2.1: Predation

3.2.1.2.2: Herbivory

3.2.1.2.3: Competition



This page titled 3.2.1: Biotic Interactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1: Biotic Interactions by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.
- 5: Populations by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2.1.1: Symbiosis

Symbiosis

Some interactions are brief, such as a hawk catching a mouse. Others involve species living in close association for long periods of time. These long-term relationships are called **symbiosis** ("living together"). In a symbiotic relationship, at least one species benefits. Depending on the interaction, the other species may be harmed (**parasitism**), unaffected (**commensalism**), or benefited (**mutualism**). Some scientists use the term symbiosis for any close relationship between species, while others reserve it for relationships in which the organisms physically live on, in, or very close to one another.

Parasitism occurs when one organism (the **parasite**) benefits while another organism (the **host**) is harmed (figure $3.2.1.1.a$). Parasites obtain food, nutrients, or shelter from their hosts, often weakening them in the process. Unlike predators, parasites usually do not kill their hosts quickly because they depend on them for survival. Some parasites live on the outside of their host, such as fleas and ticks, while others live inside the host, such as tapeworms and roundworms. Parasitism is common in many groups of organisms, including animals, plants, fungi, bacteria, and viruses. Understanding parasitic relationships helps ecologists explain how species influence one another and how populations and communities are regulated over time.



Figure $3.2.1.1.a$: Instead of conducting its own photosynthesis, the parasitic plant dodder siphons nutrients from its host plant. Image by Scot Nelson (public domain).

For example, lichens are a mutualistic symbiosis between a fungus and a photosynthetic alga or bacterium ({3{.}2{.}1{.}1{.}\eatSpaces{b} \endSpaces}\)). The photosynthetic partner provides food, while the fungus provides protection and helps retain water. Another example is the relationship between remoras and sharks ({3{.}2{.}1{.}1{.}\eatSpaces{c} \endSpaces}\)). Remoras gain food and transportation, while the shark is generally unaffected, making this a commensal relationship.



Figure 3.2.1.1.*b*: Lichen is a fungus that has symbiotic photosynthetic algae living in close association. Image modification of work by Cory Zanker.




Figure 3.2.1.1.*c*: A remora attached to the top of a shark in an aquarium. This is an example of a symbiosis because the remora lives in close association with the shark. Image by Brian Snelson(opens in new window) (CC-BY(opens in new window))

There are many examples of symbiosis in nature. **Corals** contain photosynthetic microorganisms that provide food through photosynthesis in exchange for shelter. **Mycorrhizae** are partnerships between fungi and plant roots in which the fungus helps the plant absorb water and nutrients while receiving sugars from the plant. Humans also have symbiotic relationships with many microorganisms that live on and inside our bodies. For example, bacteria in the human gut help digest food and produce certain vitamins.

Understanding these interactions helps ecologists explain how species influence one another and how communities are structured and maintained over time.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from 6.1.4 Symbiosis and 6.1.1.3 Parasitism originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Symbiosis from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Community Ecology from *Environmental Biology* by Matthew R. Fisher's (licensed under CC-BY(opens in new window))

This page titled 3.2.1.1: Symbiosis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 6.1.4: Symbiosis by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2.1.2: TROPHIC INTERACTIONS

Trophic interactions occur when one organism feeds on another. The three main types of trophic interactions are **predation** (figure 3.2.1.2.a), **herbivory**, or **parasitism**. During these interactions, one species benefits by gaining food at the expense of the other, which either dies or loses nutrients, tissues, or organs (such as leaves). Trophic interactions involve the flow of energy, and the trophic interactions in a community can be represented by **food chains** and **food webs**.



Figure 3.2.1.2.a: A hawk (*Buteo jamaicensis*) eats a vole (*Microtus californicus*), exemplifying predation, a common trophic interaction. Image by Jrockley(opens in new window) (public domain).

- Trophic Interactions

3.2.1.2.1: Predation

3.2.1.2.2: Herbivory

3.2.1.2.3: Competition

Attribution

Melissa Ha (CC-BY-NC(opens in new window))

This page titled 3.2.1.2: Trophic Interactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1.1: Trophic Interactions by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2.1.2.1: Predation

Perhaps the classical example of a biotic interaction is the predator-prey relationship. **Predation** occurs when one species (the **predator**) kills and eats multiple **prey** over its lifetime.

Predator-Prey Dynamics

Population sizes of predators and prey in a community are not constant over time, and they may vary in cycles that appear to be related. For example, 100 years of trapping data from North America shows how population dynamics of the lynx (predator) and the snowshoe hare (prey) cycle (figure $3.2.1.2.1.a$-b).



Figure $3.2.1.2.1.a$: The Canada lynx is an example of a predator. Image by Michael Zahra (CC-BY-SA).



Figure $3.2.1.2.1.b$: The snowshoe hare is the prey of the Canada lynx. Image by the National Park Service (public domain).

This cycling of predator and prey population sizes has a period of approximately ten years, with the predator population lagging one to two years behind the prey population (figure $3.2.1.2.1.a$-c). An apparent explanation for this pattern is that as the hare numbers increase, there is more food available for the lynx, allowing the lynx population to increase as well. When the lynx

population grows to a threshold level, however, they kill so many hares that hare numbers begin to decline, followed by a decline in the lynx population because of scarcity of food. When the lynx population is low, the hare population size begins to increase due, in part, to low predation pressure, starting the cycle anew.




Figure 3.2.1.2.1.*c*: The cycling of Canada lynx and snowshoe hare populations in Northern Ontario is an example of predator-prey dynamics. The x-axis shows time in years, and the y-axis shows population size in thousands of animals. The lynx population is represented in blue (shorter peaks), and the hare population is represented in red (taller peaks). Both populations increase and decrease in 10-year cycles, with changes in the lynx population lagging a few years behind that of hare.

The dynamic of closely matched population cycling occurs because the snowshoe hare is the primary food source for the lynx. When predators have generalists, feeding on a variety of prey species, and prey are consumed by a variety of predators species, their population dynamics are less likely to cycle together.

Adaptations of Predators

Predators have evolved a variety of adaptations to catch and consume prey, and the specific adaptations depend on the predator. For example, raptors, such as owls and hawks, have hooked beaks that help with tearing flesh and talons for grabbing prey. Mammal predators often have sharp teeth and claws. Some predators can run quickly to chase prey, and others "sit and wait", lunging forward when prey pass them. Some predators, like rattlesnakes and tarantulas, subdue their prey by injecting them with venom (figure 3.2.1.2.1.*d*). Predators often have large eyes located forward (like a wolf) rather than eyes spaced far apart on the sides of the head (like a sheep). Forward-facing eyes allow for depth perception, which is key to tracking prey. In contrast, peripheral vision is expanded when eyes are located to the sides, and this helps prey identify threats.



Figure 3.2.1.2.1.d: This Asian birdeater tarantula uses its fangs to inject its prey with poisonous venom. Image by Matt Reinbold (CC-BY-SA).

Defensive Traits in Prey

Prey evolve mechanical, chemical, physical, or behavioral defenses against predators. Some prey have armor (such as a turtle shell or the bony plates protecting armadillos), a **mechanical defense** that reduces predation by discouraging physical contact. Many animals produce or obtain **chemical defenses** from plants and store them to prevent predation. Other species use their body shape and coloration as **camouflage** to avoid being detected by predators, an example of a **physical defense**. The tropical walking stick is an insect with the coloration and body shape of a twig, which makes it very hard to see when it is stationary against a background of real twigs (figure 3.2.1.2.1.e-a). In another example, the chameleon can change its color to match its surroundings (figure 3.2.1.2.1.e-b).



(a) (b)

Figure 3.2.1.2.1.e: (a) The tropical walking stick and (b) the chameleon use their body shape and/or coloration to prevent detection by predators. (credit a: modification of work by Linda Tanner; credit b: modification of work by Frank Vassen)

Some species use coloration as a way of warning predators that they are distasteful or poisonous. For example, the monarch butterfly caterpillar sequesters poisons from its food (plants and milkweeds) to make itself poisonous or distasteful to potential predators. The caterpillar is bright yellow and black to advertise its toxicity. The caterpillar is also able to pass the sequestered toxins on to the adult monarch, which is also dramatically colored black and red as a warning to potential predators. Fire-bellied toads produce toxins that make them distasteful to their potential predators (figure 3.2.1.2.1.f). They have bright red or orange coloration on their bellies, which they display to a potential predator to advertise their poisonous nature and discourage an attack. Warning coloration only works if a predator uses eyesight to locate prey and can learn—a naïve predator must experience the negative consequences of eating one before it will avoid other similarly colored individuals.



Figure 3.2.1.2.1.f: The fire-bellied toad has bright coloration on its belly that serves to warn potential predators that it is toxic. (credit: modification of work by Roberto Verzo)

While some predators learn to avoid eating certain potential prey because of their coloration, other species have evolved mechanisms to mimic this coloration to avoid being eaten, even though they themselves may not be unpleasant to eat or contain toxic chemicals. In some cases of **mimicry**, a harmless species imitates the warning coloration of a harmful species. Assuming they share the same predators, this coloration then protects the harmless ones. Many insect species mimic the coloration of wasps, which are stinging, venomous insects, thereby discouraging predation (figure 3.2.1.2.1.g). In other cases of mimicry, multiple species share the same warning coloration, but all of them actually have defenses. The commonness of the signal improves the compliance of all the potential predators.



Figure 3.2.1.2.1.g: One form of mimicry is when a harmless species mimics the coloration of a harmful species, as is seen with the (a) wasp (*Polistes* sp.) and the (b) hoverfly (*Syrphus* sp.). (credit: modification of work by Tom Ings)

Running from predators, hiding, and playing dead are examples of **behavioral defenses**. Some prey also exhibit behaviors that threaten predators. For example, snapping turtles stretch their legs to appear larger and snap aggressively at predators.

Attributions

Modifed by Erin Rempala.

Modified by Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 3.2.1.2.1: Predation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1.1.1: Predation by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.2.1.2.2: Herbivory

When an animal feed on parts of another organism (usually a plant), **herbivory** occurs. For example, the koala eats the eucalyptus leaves (figure $3.2.1.2.2.a$) or a cicada feeds a plant's sap (figure $3.2.1.2.2.b$). Like predators, herbivores may feed on multiple individuals over their lives; however, herbivores do not necessarily kill the plants that they eat.



Figure $3.2.1.2.2.a$: Koalas are herbivores that feed exclusively on eucalyptus leaves. They are thus considered specialists because the specialize on just one type of food. Image by Dr. Umesh Behari Mathur (CC-BY).



Figure $3.2.1.2.2.b$: This floury baker cicada feeds on plant sap by piercing the tree with needle-like mouth parts. Image by Toby Hudson (CC-BY-SA).

Plant species have evolved numerous defenses that reduce herbivory. Thorns are an example of a mechanical defense that discourages large herbivores from feeding on a plant (Figure $3.2.1.2.2.c$-a). Many plant species produce secondary plant compounds that serve no function for the plant except that they are toxic to animals and discourage consumption, acting as a chemical defense. For example, the foxglove produces several compounds, including digitalis, that are extremely toxic when eaten (Figure $3.2.1.2.2.c$-b). (Biomedical scientists have repurposed the chemical produced by foxglove as a heart medication, which has saved lives for many decades.)



(a) (b)

Figure 3.2.1.2.2.*c*: The (a) honey locust tree uses thorns, a mechanical defense, against herbivores, while the (b) foxglove uses a chemical defense: toxins produces by the plant can cause nausea, vomiting, hallucinations, convulsions, or death when consumed. (credit a: modification of work by Huw Williams; credit b: modification of work by Philip Jägenstedt)

Attributions

Modified by Erin Rempala from Environmental Science (Ha and Schleiger) by Melissa Ha and Rachel Schleiger (CC BY-NC 4.0(opens in new window))

Modified by Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 3.2.1.2.2: Herbivory is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1.1.2: Herbivory by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.2.1.2.3: Competition

Resources are often limited within a habitat and multiple species may compete to obtain them. **Competition** occurs when organisms use the same resources, and one or both organisms is harmed. Competing species are often occupy the same trophic level and compete for food, but species on different trophic levels could still compete for space, water, etc. **Intraspecific competition** occurs within a species. For example, most penguins defend territories from other individuals of the same species because they compete for suitable habitat and other resources (figure $3.2.1.2.3.a$). **Interspecific competition** occurs between different species. For example, the **invasive** vine, kudzu, competes with trees in the southeastern United States for light (figure $3.2.1.2.3.b$). Kudzu is considered invasive in this region because it occurs outside of its historical range (Asia and Australia) and causes ecological harm.



Figure $3.2.1.2.3.a$: Adélie penguins at Cape Adare in Ross Sea, Antarctica. These penguins defend small territories in which they nest. Image by Brocken Inaglory (CC-BY-SA).



Figure $3.2.1.2.3.b$: Kudzu is a vine that covers trees, shrubs, and other structures. It can fully block its competitors' access to light, eventually killing them. Image by Katie Ashdown (CC-BY).

Ecologists have come to understand that all species have an **ecological niche**: the unique set of resources used by a species, which includes its interactions with other species. The **competitive exclusion principle** states that two species cannot occupy

the exact same niche in a habitat. In other words, different species cannot coexist in a community if they are competing for all the same resources. Competition harms one or both competitors because it wastes energy. In an experimental test of competitive exclusion, two species of the freshwater microbe *Paramecium* were cultured separately and together. When cultured separately, both species reproduced, and the number of cells (individuals) increased. However, when the two species were grown together, one species (*P. aurelia*) grew, and the the other species (*P. caudatum*) was eliminated. Because both species occupied the same ecological niche, they could not coexist. *Paramecium aurelia* was the superior competitor in this case (figure 3.2.1.2.3.c). Competitive exclusion may be avoided if a population evolves to make use of a different resource, a different area of the habitat, or feeds during a different time of day. This is called **resource partitioning**. The two species are then said to occupy different microniches. These species coexist by minimizing direct competition.




Figure 3.2.1.2.3.c: Graphs a, b, and c all plot number of cells versus time in days. When *Paramecium aurelia* and *P. caudatum* were grown individually in the laboratory (a and b), the two species both reproduced and reached a relatively high cell density. When the two species were placed together in the same test tube (habitat, c), *P. aurelia* (top, green) reached nearly the same cell density as it exhibited when grown alone, but *P. caudatum* (bottom, purple) hardly grew at all, and eventually its population drops to zero. *Paramecium aurelia* outcompeted *P. caudatum* for food, leading to the latter's eventual extinction.

Attributions

Modified by Erin Rempala from Environmental Science (Ha and Schleiger) by Melissa Ha and Rachel Schleiger (CC BY-NC 4.0(opens in new window))

Modified by Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 3.2.1.2.3: Competition is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1.2: Competition by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.2.2: Community Structure and Dynamics

Communities are complex systems that can be characterized by **community structure** (the number and size of populations and their interactions) and **community dynamics** (how the members and their interactions change over time). Understanding community structure and dynamics allows us to minimize impacts on ecosystems and manage ecological communities we benefit from.

Keystone Species

A **keystone species**, like the American bison discussed earlier in this chapter, has a much larger impact on its ecosystem than would be expected based on its abundance. Because many other organisms depend on it, changes in a keystone species can affect the structure, function, and biodiversity of an entire community. Another example is the sea star *Pisaster ochraceus*, a keystone species in rocky intertidal communities along the Pacific Coast (figure $3.2.2.a$). When sea stars are removed, mussel populations often grow rapidly and outcompete other species, reducing biodiversity. Keystone species demonstrate how changes to a single species can ripple through an ecosystem and affect many other populations. Understanding these relationships helps ecologists protect biodiversity and restore healthy ecosystems.



Figure $3.2.2.a$: Sea stars are keystone species of the intertidal zone.

Community Dynamics and Succession

Community dynamics are the changes in the species that make up a community over time. Communities are influenced by disturbances such as wildfires, floods, volcanic eruptions, droughts, storms, invasive species, and human activities. Because ecosystems are complex systems with many interacting parts, communities are constantly changing rather than remaining completely stable. Following a disturbance, a community may recover to a state similar to its previous condition, develop into a different community, or fluctuate among several possible states.

Ecological succession describes changes in community composition over time following the creation of new habitat or a disturbance. Ecologists once viewed succession as a predictable process that always led to a stable **climax community**. Today, scientists recognize that succession is often influenced by environmental conditions, species interactions, chance events, and future disturbances. As a result, communities may follow different pathways and may not reach a single permanent endpoint.

Primary Succession

Primary succession occurs when life begins to colonize newly exposed surfaces that lack soil, such as lava flows, new volcanic islands, or land exposed by retreating glaciers. The first organisms to arrive are called **pioneer species**, which may include

lichens, mosses, and hardy plants (figure 3.2.2.*b*). These organisms help break down rock and contribute organic matter, gradually creating soil that allows other species to become established. Over time, ecological communities often become more diverse and complex.



Figure 3.2.2.*b*: During primary succession in lava on Maui, Hawaii, succulent plants are the pioneer species. (credit: Forest and Kim Starr)

Secondary Succession

Secondary succession occurs after a disturbance removes part of a community but leaves the soil intact (figure 3.2.2.*c*). Common examples include areas affected by wildfire, storms, logging, or abandoned agriculture. Because soil, seeds, roots, and microorganisms remain, recovery is usually much faster than during primary succession. Early colonizing species such as grasses and wildflowers are often followed by shrubs and trees. However, the exact path of succession depends on factors such as climate, soil conditions, species interactions, and future disturbances.




Figure 3.2.2.*c*: Example of secondary succession leading, in this case, to a mature oak and hickory forest. Modified by Erin Rempala from original. Original not credited in Ha and Schleiger's Environmental Science

Succession provides an example of **systems thinking**. Changes in one part of an ecosystem can influence many other components, including species populations, nutrient cycles, soil conditions, and habitat availability. By studying succession, ecologists can better understand ecosystem recovery, restoration, and long-term environmental change.

A Simplified Framework

Ecological succession is a useful model for understanding how communities change over time, but real ecosystems are often more complex than the classic succession model suggests. Environmental conditions, species interactions, disturbances, and chance events can all influence the course of succession. As a result, ecosystems may follow multiple pathways rather than progressing toward a single, stable endpoint.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Community Ecology from *Environmental Biology* by Matthew R. Fisher's (licensed under CC-BY(opens in new window))

This page titled 3.2.2: Community Structure and Dynamics is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.2: Community Structure and Dynamics by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.2.3: Data Dive- Beaver Impacts on Wetlands

Overview

An ecosystem engineer is any animal that creates, significantly modifies, maintains or destroys a habitat. These organisms impact both the abiotic and biotic components of a habitat and can thus completely change landscapes. One 2017 study sought to understand if beavers (classic ecosystem engineer) could even be used as a tool for habitat restoration of wetlands. Specifically, this study focused on quantifying if wetland biodiversity improved as a result of reintroducing beavers on a landscape degraded by agriculture. Figure 2.4.6a below displays some of the results in this study:




Figure 3.2.3.a: Average number of species observed in sample plots 1-2 years and 10-12 years after beavers were introduced. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Law A, Graywood MJ, Jones KC, Ramsay P, and Willby NJ 2017.

Questions

1. What is the independent (explanatory) variable and the dependent (response) variable?
2. What question(s) are the authors trying to answer with this graph?
3. What trend(s) can be observed in this graph between the 1-2 and 10-12 timetables? Support your answer by referring to appropriate patterns in the graph.
4. Do you think like the authors are satisfied with the results in the graph? Why?
5. How can the results of this graph to inform future reintroduction of beavers where wetland restoration is needed?
6. What information/patterns is not clear from this graph?

Raw Data From Above Graph(s)

Table 3.2.3.a**:** Raw data for average number of species observed in sample plots 1-2 years and 10-12 years after beavers were introduced. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Law A, Graywood MJ, Jones KC, Ramsay P, and Willby NJ 2017.

Raw data for average number of species graph

Number Of Plots	Average Number Of Species After 1-2 Years	Average Number Of Species After 10-12 Years
0	3	3

Number Of Plots	Average Number Of Species After 1-2 Years	Average Number Of Species After 10-12 Years
20	10	28
40	11	35
60	12	38
80	13	39
100	13.5	40
130	14	41

Attribution

Rachel Schleiger (CC-BY-NC)

This page titled 3.2.3: Data Dive- Beaver Impacts on Wetlands is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 7.6: Data Dive- Beaver Impacts on Wetlands by Melissa Ha and Rachel Schleiger has no license indicated.

3.3: ECOSYSTEM ECOLOGY

Introduction

In semi-urban areas, beavers (*Castor canadensis*) are considered a nuisance as their dams block drainage pipes and cause flooding. However, in wild areas beavers are one of the most important species. Beaver dams are known for altering the flow of water. However, these dams not only slow down the movement of water, but they spread and store water in a way that is much more efficient than human-made dams. In addition, they slow and spread sediment and nutrients as they move through watersheds. What this does is create a mosaic of habitats across a landscape, both aquatic and terrestrial. More habitats lead to more species, both for plants and animals. In this way, beavers act as **ecosystem engineers**, to create ecosystems, one stick at a time.



Figure $3.3.a$ American beaver. Image by NeexPix(opens in new window) (Public domain)

3.3.1: Matter

3.3.2: Energy

3.3.3: Biogeochemical Cycles

3.3.4: Energy Flow in Food Webs

3.3.5: Ecosystem Types and Dynamics

3.3.5.1: Climate and Biomes

3.3.5.2: Terrestrial Biomes

3.3.5.3: Aquatic Biomes

3.3.6: Data Dive- Biome Carbon Storage

Attributions

Modified by Erin Rempala.

Rachel Schleiger (CC-BY-NC(opens in new window))

This page titled 3.3: Ecosystem Ecology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 7: Ecosystems by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.3.1: Matter

Matter, Atoms, and the Chemistry of the Environment

Everything in the environment is made of matter. **Matter** is anything that has mass and takes up space. The air we breathe, the water we drink, the soil beneath our feet, and all living organisms are made of matter. Matter moves through ecosystems as organisms interact with the atmosphere, water, and land. These movements occur through **biogeochemical cycles** such as the water, carbon, nitrogen, and phosphorus cycles.

Matter is composed of substances called elements. An **element** is a substance that cannot be broken down into simpler substances through ordinary chemical reactions. Examples of elements important to environmental science include carbon (C), hydrogen (H), oxygen (O), nitrogen (N), phosphorus (P), and sulfur (S) (figure $3.3.1.a$).




Figure $3.3.1.a$: The periodic table of elements organizes chemical elements based on their properties. Each element is represented by a one- or two-letter symbol. Additionally, the atomic number, name, and atomic mass are listed for each element. Click here for an accessible periodic table of elements. Image by OpenStax (CC-BY)

Atoms

The smallest unit of an element is an **atom**. Atoms are the building blocks of all matter. Each atom contains three main types of particles:

- **Protons**, which have a positive charge (+1)
- **Neutrons**, which have no charge
- **Electrons**, which have a negative charge (-1)

Protons and neutrons are found in the **nucleus**, or center, of the atom, while electrons move around the nucleus (figure $3.3.1.b$).

The number of protons in an atom determines which element it is. For example, every carbon atom contains six protons, while every oxygen atom contains eight protons.




Figure 3.3.1.*b*: Atoms are comprised of protons (red with a plus sign, +) and neutrons (yellow) located within the nucleus, and electrons (blue with a minus sign, –)surrounding the nucleus. The nucleus of an atom should not be confused with the nucleus of a cell.

Molecules and Compounds

Atoms can combine to form molecules. A **molecule** is made of two or more atoms bonded together. For example:

- Oxygen gas (O_2) contains two oxygen atoms.
- Water (H_2O) contains two hydrogen atoms and one oxygen atom.
- Carbon dioxide (CO_2) contains one carbon atom and two oxygen atoms.

When atoms from different elements combine, they form a compound. Many compounds are essential for life and environmental processes.

Chemical Bonds

Atoms are held together by **chemical bonds**. The type of bond affects how substances behave in nature.

Covalent Bonds

A **covalent bond** forms when atoms share electrons (figure 3.3.1.*c*). Covalent bonds are strong and common in living organisms.

Examples include:

- Water (H_2O)
- Carbon dioxide (CO_2)
- Methane (CH_4)
- The molecules that make up living organisms

Most biological molecules are held together by covalent bonds.




Figure 3.3.1.*c*: A covalent bond forming H_2 (right) where two hydrogen atoms share the two electrons. By Jacek FH - Own work, CC BY-SA 3.0, wikimedia.org [commons.wikimedia.org] [commons.wikimedia.org] [commons.wikimedia.org]

Ionic Bonds

An ionic bond forms when one atom transfers electrons to another atom. This process creates charged particles called ions (figure 3.3.1.*d*.




Figure 3.3.1.*d*: Sodium and fluorine atoms undergoing a reaction to form sodium ions and fluoride ions. Sodium loses its outer electron, and this electron enters the fluorine atom. The oppositely charged ions – typically a great many of them – are then attracted to each other to form solid sodium fluoride.
By Wdcf - Own work, CC BY-SA 3.0, [wikimedia.org [commons.wikimedia.org]]

For example, sodium (Na⃝) and chloride (Cl⃝) combine to form sodium chloride (table salt). Ions are important in environmental systems because they dissolve easily in water and influence water quality, nutrient availability, and pH.

Hydrogen Bond Attractive Force

Hydrogen bond attractive forces are weaker attractions that form between certain molecules (Figure 3.3.1.*d*). Although weaker than covalent or ionic bonds, hydrogen bond attractive forces play an important role in environmental systems. Hydrogen bond attractive forces allow water molecules to stick together, giving water many unique properties that support life.

Water: A Special Molecule

Water is one of the most important substances on Earth. A water molecule contains one oxygen atom and two hydrogen atoms joined by covalent bonds. Because oxygen attracts electrons more strongly than hydrogen, water molecules have slightly positive and slightly negative ends. This unequal distribution of charge makes water a polar molecule.

Water's polarity allows it to:

- Dissolve many substances
- Transport nutrients through ecosystems
- Support chemical reactions in cells
- Moderate temperature in organisms and environments

These properties make water essential for life and for ecosystem processes.




Figure $3.3.1.d$: Hydrogen bonds form between slightly positive and slightly negative charges of certain molecules, such as water. Each water molecule consists of an oxygen atom (O) connected to two hydrogen atoms (H) by covalent bonds (solid lines). Dashed lines represent hydrogen bonds between water molecules.

Acids, Bases, and pH

Many environmental issues involve acids and bases.

An acid is a substance that releases hydrogen ions (H^+) in water.

A base is a substance that removes hydrogen ions from water or releases hydroxide ions (OH^-).

The concentration of hydrogen ions is measured using the pH scale (figure $3.3.1.e$).

- pH 7 is neutral.
- Values below 7 are acidic.
- Values above 7 are basic (alkaline).

Examples:

- Lemon juice has a pH of about 2.
- Pure water has a pH of 7.
- Seawater has a pH of about 8.

Even small pH changes can affect organisms and ecosystems. For example, increasing carbon dioxide in the atmosphere can lower ocean pH, a process called ocean acidification that threatens corals and shell-forming organisms.




Figure 3.3.1.*e*: The pH Scale. By PHscalenolang.png: myself, PatríciaRderivative work: Chris828 (talk) - PHscalenolang.png, CC BY-SA 2.5, [wikimedia.org [commons.wikimedia.org]]

Carbon: The Foundation of Life

Carbon is one of the most important elements in environmental science. Carbon atoms can form four covalent bonds, allowing them to build a wide variety of molecules. Methane, the main component of fossil gas, is the simplest organic molecule. Methane consists of one central carbon atom covalently bonded to four hydrogen atoms.

Because of this ability, carbon serves as the foundation for most molecules found in living organisms. Life on Earth is often described as **carbon-based**.

Carbon moves continuously through the atmosphere, oceans, soil, and living organisms in the **carbon cycle**. Understanding carbon chemistry helps explain climate change, ecosystem productivity, and the role of fossil fuels in modern society.




Figure 3.3.1.*f*: The simplest carbon molecule is methane (CH_4). By DynaBlast, CC BY-SA 2.5, [wikimedia.org [commons.wikimedia.org]]

Biological Macromolecules

Living organisms use carbon to build large molecules called biological macromolecules.

Carbohydrates

Carbohydrates provide energy for cells and help build structures such as plant cell walls. Examples include glucose, starch, and cellulose.

Lipids

Lipids store energy for long periods and form important parts of cell membranes. Fats and oils are common examples.

Proteins

Proteins perform many functions in organisms. They help build tissues, transport materials, and speed up chemical reactions.

Nucleic Acids

Nucleic acids store and transmit genetic information. The two main nucleic acids are DNA and RNA.

Together, these four groups of molecules form the foundation of all living organisms.




Figure $3.3.1.g$: The double-helix model shows DNA as two parallel strands of intertwining molecules.Chemical structure of DNA; hydrogen bonds shown as dotted lines. By Madprime (talk · contribs) - Own work. This vector image was created with Inkscape by v ., CC BY-SA 3.0, [wikimedia.org] [commons.wikimedia.org]]

Isotopes and Radioactivity

Most atoms of an element contain the same number of protons but may differ in the number of neutrons. These different forms are called **isotopes** (figure $3.3.1.h$)**.**

Some isotopes are unstable and release energy as they decay. This process is called **radioactivity**.

Scientists use **radioactive isotopes** to study groundwater movement, the age of rocks and fossils, and changes in past climates.




Figure 3.3.1.h: The three naturally occurring isotopes of hydrogen. The fact that each nuclide has 1 proton makes them all isotopes of hydrogen: the identity of the isotope is given by the number of protons and neutrons. From left to right, the isotopes are protium (^{1}H) with 0 neutrons, deuterium (^{2}H) with 1 neutron, and tritium (^{3}H) with 2 neutrons. By Dirk Hünniger;Derivative work in english - Balajijagadesh - This file was derived from: Hydrogen Deuterium Tritium Nuclei Schmatic-de.svg, CC BY-SA 3.0, [wikimedia.org [commons.wikimedia.org]]

Nuclear Energy: Fission and Fusion

Chemical reactions involve changes in electron arrangements. **Nuclear reactions** involve changes in the nucleus of an atom and release much larger amounts of energy.

Nuclear Fission

Nuclear fission occurs when a large atomic nucleus splits into smaller nuclei. Modern nuclear power plants generate electricity using fission.

Nuclear Fusion

Nuclear fusion occurs when small atomic nuclei combine to form a larger nucleus. Fusion powers the Sun and all other stars. Scientists are studying fusion because it could provide large amounts of energy with fewer long-lived radioactive wastes than fission. However, commercial fusion power remains under development.

Why Environmental Scientists Study Chemistry

Many environmental challenges involve chemical processes. Understanding atoms, molecules, chemical bonds, pH, and energy transformations helps scientists explain:

- Climate change
- Ocean acidification
- Air pollution
- Water quality
- Nutrient cycling
- Renewable and nonrenewable energy systems
- Ecosystem health and sustainability

Chemistry provides the foundation for understanding how matter moves through environmental systems and how human actions affect the Earth.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Matter from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 3.3.1: Matter is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 7.2: Matter by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.3.2: Energy

What Is Energy?

Energy is the ability to cause change or do work. It exists in many forms, including light, heat, chemical, electrical, and mechanical energy. Energy can be transferred between organisms and their environment, and it can change from one form to another. For example, plants convert light energy from the Sun into chemical energy through photosynthesis. Animals convert the chemical energy stored in food into movement and body heat. Similarly, solar panels transform sunlight into **electrical energy**.

Energy can be classified as either **kinetic energy** or **potential energy**. Kinetic energy is the energy of motion. Flowing water, wind, and moving animals all have kinetic energy. Potential energy is stored energy that can be released later. Water behind a dam, a stretched rubber band, and food molecules are all examples of potential energy (figure $3.3.2.a$).

One important form of potential energy is **chemical energy**, which is stored in the bonds between atoms. During photosynthesis, plants capture **solar energy** and store it as chemical energy in sugars. Organisms release this stored energy when they break down food molecules through **cellular respiration**. The chemical energy obtained from food powers nearly all biological activities, including growth, movement, reproduction, and maintaining life processes.

Understanding how energy is stored, transformed, and transferred helps explain how organisms survive and how energy flows through ecosystems.



Figure $3.3.2.a$. Still water has potential energy; moving water, such as in a waterfall or a rapidly flowing river, has kinetic energy. (credit “dam”: modification of work by “Pascal”/Flickr; credit “waterfall”: modification of work by Frank Gualtieri)

Energy Flow in Ecosystems

Every living organism needs energy to survive. Energy is required for growth, movement, repair, reproduction, and many other life processes. Cells use energy to build molecules, transport materials, remove wastes, and maintain the conditions needed for life. The movement and use of energy in living systems is called **bioenergetics**. All of the chemical reactions that occur within an organism are collectively called **metabolism**. Some metabolic reactions release energy, while others require energy.

Almost all life on Earth depends on energy from the Sun (figure $3.3.2.b$). Plants, algae, and some bacteria capture sunlight through **photosynthesis** and store that energy as chemical energy in sugars and other organic molecules. Animals obtain this energy by eating plants or by eating other animals. Decomposers, such as fungi and bacteria, obtain energy by breaking down dead organisms and organic waste, returning nutrients to the environment.

As energy moves from the Sun to producers, consumers, and decomposers, it flows through ecosystems in a single direction. Energy enters ecosystems as sunlight, is captured by producers through photosynthesis, and is transferred to consumers and decomposers through feeding and decomposition. During each energy transfer, organisms use some of the energy to power their life processes, and a portion is released as heat.

Unlike matter, which is recycled through biogeochemical cycles, energy is not recycled. Because energy is continually lost as heat, ecosystems require a constant input of energy from the Sun. This one-way flow of energy is a fundamental characteristic of all ecosystems and helps explain why the Sun is the ultimate source of energy for nearly all life on Earth.




Figure 3.3.2.*b*: Ultimately, most life forms get their energy from the sun. Plants use photosynthesis to capture sunlight, and herbivores eat the plants to obtain energy. Carnivores eat the herbivores, and eventual decomposition of plant and animal material contributes to the nutrient pool.

Thermodynamics and Energy Flow

The flow of energy through ecosystems follows the laws of thermodynamics, which describe how energy is transferred and transformed. As sunlight moves through producers, consumers, and decomposers, energy changes form but is never created or destroyed. At the same time, some energy is lost as heat during every transfer. These patterns help explain why ecosystems depend on a continuous input of solar energy.

Thermodynamics is the study of energy transformations. In environmental science, the first two laws of thermodynamics are especially important because they explain how energy moves through organisms and ecosystems.

Shown are some examples of energy transferred and transformed from one system to another and from one form to another (figure 3.3.2.*c*). The food we consume provides our cells with the energy required to carry out bodily functions, just as light energy provides plants with the means to create the chemical energy they need.




Figure 3.3.2.c: Shown are some examples of energy transferred and transformed from one system to another and from one form to another. The food we consume provides our cells with the energy required to carry out bodily functions, just as light energy provides plants with the means to create the chemical energy they need. (credit “ice cream”: modification of work by D. Sharon Pruitt; credit “kids”: modification of work by Max from Providence; credit “leaf”: modification of work by Cory Zanker)

The First Law of Thermodynamics

The **first law of thermodynamics** states that energy cannot be created or destroyed. Instead, energy changes form or moves from one place to another. For example, during photosynthesis, plants convert sunlight into chemical energy stored in sugars. When animals eat plants, that chemical energy is transferred to their cells. Because energy is conserved, ecosystems do not create energy; they transform and transfer it.

The Second Law of Thermodynamics

The **second law of thermodynamics** states that every energy transfer becomes less useful because some energy is lost as heat (**entropy**).

For example:

- A light bulb produces light, but it also releases heat.
- A car engine moves a vehicle, but much of the energy in fuel becomes heat.
- Living organisms use energy to grow, move, and reproduce, but they also release heat to their surroundings.

Because some energy is lost as heat during every transfer, less energy is available at each trophic level of a food web. This is why ecosystems require a constant input of solar energy and why there are generally fewer organisms at higher trophic levels.

Together, the first and second laws of thermodynamics explain how energy flows through ecosystems: energy enters primarily as sunlight, is transformed into chemical energy by producers, passes through food webs, and leaves the ecosystem as heat.

Connections to Environmental Science

Energy drives all environmental processes. The Sun provides the energy that powers photosynthesis, weather patterns, ocean currents, and most ecosystems on Earth. Understanding how energy is transformed and transferred helps scientists explain many environmental challenges.

For example:

- Climate Change is occurring because human activities have altered the flow of energy in Earth's atmosphere.

- Food webs depend on the transfer of energy from producers to consumers and decomposers.
- Renewable energy technologies such as solar and wind power capture energy that originates from the Sun.
- Agriculture depends on photosynthesis, which converts solar energy into food.
- Ecosystem productivity is limited by the amount of energy available to organisms.
- Energy losses at each trophic level help explain why food webs typically contain fewer organisms at higher levels.

The laws of thermodynamics also influence sustainability. Because energy is lost as heat during every transfer, no energy system is 100% efficient. Environmental scientists use this principle to evaluate transportation systems, food production, buildings, and energy technologies. Understanding how energy flows through natural and human systems helps us make informed decisions about resource use and environmental stewardship.

Real-World Examples

Throughout this course, you will apply energy concepts to environmental issues and field experiences, including:

- Energy use and resource recovery at the Miramar Landfill.
- Energy requirements of wastewater treatment at the Point Loma Wastewater Treatment Plant.
- Food webs and ecosystem productivity at Kendall-Frost Marsh Reserve.
- Renewable energy and climate solutions explored through multiple lab activities.

Understanding energy flow provides a foundation for studying ecosystems, climate change, sustainability, and the relationship between human activities and the environment.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content **2.2: Energy** originally authored by by Matthew R. Fisher and is licensed CC BY 4.0.

Essentials of Environmental Science(opens in new window) by Kamala Doršner is licensed under CC BY 4.0. Modified from the original by Matthew R. Fisher.

This page titled 3.3.2: Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

- 2.2: Energy by Matthew R. Fisher is licensed CC BY 4.0. Original source: https://openoregon.pressbooks.pub/envirobiology.

3.3.3: Biogeochemical Cycles

Biogeochemical Cycles: Recycling Matter on Earth

Energy Flows, Matter Cycles

Energy and matter move through ecosystems in different ways.

- Energy enters most ecosystems as sunlight.
- Energy flows through food webs and is eventually released as heat.
- Matter is recycled and reused through biogeochemical cycles.

Photosynthesis, cellular respiration, decomposition, and nutrient cycling connect the flow of energy and the movement of matter. Understanding both energy flow and matter cycling is essential for understanding how ecosystems function.

Matter Cycles Through Earth Systems

Every living organism needs matter to survive. Matter provides the atoms needed to build cells, tissues, and organisms. Unlike energy, which flows through ecosystems and is eventually lost as heat, matter is continuously recycled and reused.

Biogeochemical cycles describe how matter moves through living organisms and the environment. The term biogeochemical combines three parts:

- Bio = living organisms
- Geo = Earth, including soil, rocks, water, and the atmosphere
- Chemical = elements and compounds

Biogeochemical cycles move essential elements through the atmosphere, water, soil, rocks, and living things.

The **law of conservation of mass** states that matter cannot be created or destroyed. As a result, the same atoms are continuously recycled through Earth's systems. An atom of carbon in a tree today may once have been part of a dinosaur, a shell on the ocean floor, or a molecule of carbon dioxide in the atmosphere.

Why Are Biogeochemical Cycles Important?

Biogeochemical cycles provide organisms with the nutrients they need to survive. They also help regulate many environmental processes that support life on Earth.

These cycles:

- Support plant growth
- Maintain ecosystem function
- Regulate Earth's climate
- Help maintain air and water quality
- Support food production

Major Biogeochemical Cycles

This chapter focuses on four important nutrient cycles.

Carbon Cycle

Carbon is the foundation of life and is found in living organisms, fossil fuels, the atmosphere, oceans, and rocks (figure $3.3.3.a$).




Figure $3.3.3.a$: Carbon dioxide in the atmosphere is converted to organic carbon through photosynthesis by terrestrial organisms (like trees) and marine organisms (like algae). Respiration by terrestrial organisms (like trees and deer) and marine organisms (like algae and fish) release carbon dioxide back into the atmosphere. Additionally, microbes that decompose dead organisms release carbon dioxide through respiration. Weathering of terrestrial rocks also brings carbon into the soil. Carbon in the soil enters the water through leaching and runoff. It can accumulate into ocean sediments and reenter land through uplifting. Long-term storage of organic carbon occurs when matter from living organisms is buried deep underground and becomes fossilized. Volcanic activity and, more recently, human emissions stored carbon back into the carbon cycle. Modified from John M. Evans and Howard Perlman, USGS using tree and deer (both public domain).

Carbon moves through ecosystems as living things use and recycle organic matter. During cellular respiration, plants, animals, and many microorganisms break down food molecules. This process uses oxygen and releases energy, water, and carbon dioxide into the atmosphere. When organisms die, decomposers such as bacteria and fungi break down their remains, returning carbon to the environment as carbon dioxide or methane (figure $3.3.3.b$). Photosynthesis and respiration work together in the carbon cycle. Photosynthesis removes carbon dioxide from the atmosphere and stores carbon in sugars, while respiration releases carbon dioxide back into the atmosphere. Carbon dioxide is also released by volcanoes, hydrothermal vents, and the burning of wood and fossil fuels. The oceans store large amounts of carbon and exchange carbon with the atmosphere, helping control carbon dioxide levels on Earth.



Figure 3.3.3.*b*: Decomposers will break down the organic compounds in this fallen tree at Cliffs of the Neuse State Park in Wayne County, North Carolina, releasing carbon dioxide into the atmosphere. Decomposition ensures that carbon dioxide will be available in the atmosphere for photosynthetic organisms, which then provide carbon for consumers. Image by Gerry Dincher (CC-BY-SA).

Nitrogen Cycle

Nitrogen is needed to build proteins and DNA, but most organisms cannot use atmospheric nitrogen directly (figure 3.3.3.*c*).




Figure 3.3.3.*c*: In the nitrogen cycle, nitrogen-fixing bacteria in the soil or legume root nodules convert nitrogen gas (N_2) from the atmosphere to ammonium (NH_4^+). Nitrification occurs when bacteria convert ammonium to nitrites (NO_2^-) and then to nitrates (NO_3^-). Nitrates re-enter the atmosphere as nitrogen gas through denitrification by bacteria. Plants assimilate ammonium and nitrates, producing organic nitrogen, which is available to consumers. Decomposers, including aerobic and anaerobic bacteria and fungi, break down organic nitrogen and release ammonium through ammonification. (credit: "Nitrogen cycle"(opens in new window) by Johann Dréo & Raeky is licensed under CC BY-SA 3.0)

The nitrogen cycle moves nitrogen through the atmosphere, soil, and living things. Most nitrogen in the atmosphere is in the form of nitrogen gas, which plants cannot use. Nitrogen-fixing bacteria in soil and in the roots of some plants convert nitrogen gas into ammonium (figure 3.3.3.*d*).



Figure 3.3.3.d: Nitrogen-fixing bacteria live in the spherical nodules of this soybean root. Image by United Soybean Board (CC-BY).

Other bacteria change ammonium into nitrites and then nitrates, a process called nitrification. Plants take up ammonium and nitrates from the soil and use them to build proteins and other organic molecules. Animals obtain nitrogen by eating plants or other animals. When organisms produce waste or die, decomposers such as bacteria and fungi break down the organic matter and release ammonium back into the soil through ammonification. Finally, denitrifying bacteria convert nitrates back into nitrogen gas, returning nitrogen to the atmosphere and completing the cycle.

The nitrogen cycle is important in agriculture because crops need nitrogen to grow, but excess fertilizer can pollute waterways and disrupt ecosystems (figure 3.3.3.e).



Figure 3.3.3.e: Fertilizer containing nitrogen is conventionally applied at large scales in agriculture. Image by Bob Nichols, USDA Natural Resources Conservation Service (public domain).

Phosphorus Cycle

Phosphorus is an important part of DNA, cell membranes, and ATP. Unlike carbon and nitrogen, phosphorus has no major atmospheric phase. The phosphorus cycle moves phosphorus through rocks, soil, water, and living things (figure 3.3.3.f). Most phosphorus is stored in rocks and sediments as phosphate. Weathering slowly breaks down rocks and releases phosphate into soil and water, where plants can absorb it. Animals obtain phosphorus by eating plants or other animals. Some phosphate is carried

by runoff, rivers, and groundwater into lakes and oceans, where it becomes part of aquatic food webs. Over time, phosphate can settle on the ocean floor and form sediments. Geological processes such as uplift can eventually bring these sediments back to land, allowing the cycle to continue.




Figure $3.3.3.f$: In nature, phosphorus exists as the phosphate ion (PO_4^{3-}). Phosphate enters the atmosphere from volcanic aerosols, which precipitate to Earth. Weathering of rocks also releases phosphate into the soil and water, where it becomes available to terrestrial food webs. Some of the phosphate from terrestrial food webs dissolves in streams and lakes, and the remainder enters the soil. Phosphate enters the ocean via surface runoff, groundwater flow, and river flow, where it becomes dissolved in ocean water or enters marine food webs. Some phosphate falls to the ocean floor where it becomes sediment. If uplifting occurs, this sediment can return to land. (credit: modification of work by John M. Evans and Howard Perlman, USGS)

The phosphorus cycle is important in agriculture because crops need phosphorus to grow and develop roots, but excess phosphorus from fertilizers can wash into waterways and contribute to water pollution and harmful algal blooms.

Eutrophication and Dead Zones

Eutrophication ("well nourished") occurs when too much nitrogen and phosphorus enter rivers, lakes, or oceans, often from fertilizer runoff or sewage. These extra nutrients cause algae to grow rapidly, creating algal blooms. Thick algal blooms block sunlight from reaching aquatic plants, causing many plants to die. As the dead algae and plants decompose, bacteria use large amounts of dissolved oxygen in the water. This reduces oxygen levels in the water. Low oxygen levels can kill fish, shellfish, and other aquatic organisms. In severe cases, large areas of water are called **dead zones**, where most aquatic life cannot survive. Dead zones have formed in many places around the world, including the Gulf of Mexico and the Chesapeake Bay, where nutrient runoff from agriculture and other human activities has increased nutrient pollution (figure $3.3.3.g$).




Figure 3.3.3.g: Dead zones occur when phosphorus and nitrogen from fertilizers cause excessive growth of microorganisms, which depletes oxygen and kills fauna. This map shows dead zones around the world in 2008. Worldwide, large dead zones are found in coastal areas of high population density. Image By Robert Simmon & Jesse Allen - NASA Earth Observatory, Public Domain,

Case Study: The Chesapeake Bay 3.3.3.1

The Chesapeake Bay is one of the largest and most important estuaries in the United States, but it has been affected by pollution for many years. In the 1970s, scientists identified dead zones, that harmed fish, oysters, clams, worms, and other aquatic organisms (figure 3.3.3.h). A major cause of this problem is nutrient pollution from excess nitrogen and phosphorus entering the bay. These nutrients come from fertilizers used on farms, lawns, and gardens, as well as runoff carried by more than 150 rivers and streams that flow into the bay. Because many different sources contribute to the problem, improving the health of the Chesapeake Bay requires cooperation among farmers, businesses, governments, and local communities.



(a) (b)

Figure 3.3.3.h: This (a) satellite image shows the Chesapeake Bay, an ecosystem affected by phosphate and nitrate runoff. A (b) member of the Army Corps of Engineers holds a clump of oysters being used as a part of the oyster restoration effort in the bay. (credit a: modification of work by NASA/MODIS; credit b: modification of work by U.S. Army)

The Sulfur Cycle

Sulfur is needed for some proteins and cycles among the atmosphere, water, soil, and organisms (figure 3.3.3.*i*). The sulfur cycle moves sulfur through the atmosphere, water, soil, and living things. Sulfur enters the environment from natural sources such as volcanic eruptions and from human activities such as burning fossil fuels. In the atmosphere, sulfur compounds can return to Earth in precipitation or as particles that settle on land and water. Plants absorb sulfur from the soil and use it to build proteins and other important molecules. Animals obtain sulfur by eating plants or other animals. When organisms die, decomposers return sulfur to the environment. Microorganisms also play an important role by changing sulfur between different chemical forms. Some sulfur is carried into oceans, where it can become part of marine ecosystems or be stored in sediments for long periods of time.




Figure 3.3.3.*i*: The sulfur cycle. Sulfur dioxide (SO_2) from the atmosphere is dissolved in precipitation as weak sulfuric acid or falls directly to Earth as fallout. This releases sulfates (SO_4^{2-}) into the soil and water. Soil sulfates can be carried as runoff into the water. Marine sulfate can form pyrite, and this can break down to release soil sulfates. Organisms in terrestrial and marine ecosystems assimilate sulfate, adding sulfur to organic molecules, such as proteins (not shown). Decomposition of these organisms returns sulfates to the soil. Microorganisms can convert sulfates to hydrogen sulfide (H_2S) and vice versa. Decomposition, volcanic eruptions, and human activities (including burning fossil fuels) can release hydrogen sulfide (H_2S) or sulfur dioxide into the atmosphere. (credit: modification of work by John M. Evans and Howard Perlman, USGS)

The sulfur cycle is closely linked to air and water quality because sulfur dioxide released from burning fossil fuels can contribute to acid rain, which can damage ecosystems, buildings, and aquatic habitats.

A Systems Perspective

Biogeochemical cycles move essential elements such as carbon, nitrogen, phosphorus, and sulfur through the atmosphere, water, soil, and living things. Each cycle includes **reservoirs**, where nutrients are stored, and flows (or **fluxes**), which move nutrients between reservoirs. Because Earth's systems are interconnected, changes in one cycle often affect others.

For example, increasing carbon dioxide in the atmosphere can contribute to climate change, which can alter nutrient cycling in ecosystems. Similarly, fertilizer use can affect both the nitrogen and phosphorus cycles, increasing nutrient pollution in waterways and reducing water quality and biodiversity. Understanding these connections helps scientists study how changes in one part of Earth can influence many others.

Connections to Environmental Science

Biogeochemical cycles provide a foundation for understanding many environmental issues explored throughout this course, including:

- Climate change

- Ocean acidification
- Water quality
- Agriculture and food production
- Biodiversity loss
- Ecosystem restoration
- Sustainability

By studying how nutrients move through ecosystems, scientists can predict the effects of human activities and develop solutions that support healthy and resilient ecosystems.

Human Impacts and Planetary Boundaries

Human activities such as fossil fuel combustion, agriculture, mining, and urban development have changed the way nutrients move through Earth's systems. When these changes become large enough, they can disrupt ecosystem processes and push Earth beyond conditions that have supported human societies for thousands of years.

Several Planetary Boundaries are closely linked to biogeochemical cycles:

- **The Climate Change boundary** is connected to the carbon cycle. Human activities release carbon dioxide from fossil fuels and deforestation faster than natural processes can remove it.
- **The Ocean Acidification boundary** is also linked to the carbon cycle. As more carbon dioxide dissolves into seawater, ocean pH decreases, making it harder for corals, shellfish, and other marine organisms to build and maintain calcium carbonate structures.
- **The Biogeochemical Flows boundary** focuses on human disruption of the nitrogen and phosphorus cycles. Excess fertilizer use increases the movement of these nutrients through ecosystems, contributing to eutrophication, harmful algal blooms, and dead zones.
- **The Land-System Change boundary** is connected to multiple biogeochemical cycles. Converting forests, grasslands, and wetlands to agriculture or urban areas changes how carbon, nitrogen, phosphorus, and water move through ecosystems. These changes can reduce carbon storage, increase nutrient runoff, and alter ecosystem processes.
- Changes in nutrient cycling and land use can also affect the **Biosphere Integrity boundary** by degrading habitats, reducing biodiversity, and disrupting ecosystem function

Together, these examples show that Earth's systems are deeply connected. Understanding biogeochemical cycles helps us recognize how human activities influence the environment and why sustainable management of natural systems is important for maintaining a safe and resilient planet.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Biogeochemical Cycles, Energy, and Energy Enters Ecosystems Through Photosynthesis from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Carbon Cycle and Nitrogen Cycle from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Energy Flow through Ecosystems from *General Biology* by OpenStax (licensed under CC-BY)
- Soil and Sustainability(opens in new window) and Biogeochemical Cycles and the Flow of Energy in the Earth System from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (licensed under CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Cycling of Matter from *AP Environmental Science* by University of California College Prep, University of California (licensed under CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Nutrient Cycles from *Life Sciences Grade 10* by Siyavula (CC-BY(opens in new window))

This page titled 3.3.3: Biogeochemical Cycles is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 7.3: Biogeochemical Cycles by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.3.4: Energy Flow in Food Webs

Trophic interactions in a community can be represented by diagrams called **food chains** and **food webs**. Energy flows through a community as a result of trophic interactions.

Energy Flow

Cells run on the chemical energy found mainly in carbohydrate molecules, and the majority of these molecules are produced by one process: **photosynthesis**. Through photosynthesis, certain organisms convert solar energy (sunlight) into chemical energy, which is then used to build carbohydrate molecules (figure $3.3.4.a$). The energy that is harnessed from photosynthesis enters the communities continuously and is transferred from one organism to another. Therefore, directly or indirectly, the process of photosynthesis provides most of the energy required by living things on Earth.




Figure $3.3.4.a$: Photosynthesis uses solar energy, carbon dioxide, and water to release oxygen and to produce energy-storing sugar molecules. Image and caption by OpenStax(opens in new window) (CC-BY(opens in new window)). Access for free at openstax.org(opens in new window).

Organisms that conduct photosynthesis (such as plants, algae, and some bacteria), and organisms that synthesize sugars through other means are called **producers**. Without these organisms, energy would not be available to other living organisms, and life would not be possible. **Consumers**, like animals, fungi, and various microorganisms depend on producers, either directly or indirectly. For example, a deer obtains energy by eating plants. A wolf eating a deer obtains energy that originally came from the plants eaten by that deer (figure $3.3.4.b$). Using this reasoning, all food eaten by humans can be traced back to producers that carry out photosynthesis.



Figure 3.3.4.*b*: The energy stored in carbohydrate molecules from photosynthesis passes through the food chain. The predator that eats these deer is getting energy that originated in the photosynthetic vegetation that the deer consumed. (credit: Steve VanRiper, U.S. Fish and Wildlife Service)

Consumers can be classified based on whether they eat animal or plant material (figure 3.3.4.*c*). Consumers that feed exclusively on animals are called **carnivores**. Lions, tigers, snakes, sharks, sea stars, spiders, and ladybugs are all carnivores. **Herbivores** are consumers that feed exclusively on plant material, and examples include deer, koalas, some bird species, crickets, and caterpillars. Consumers that eat both plant and animal material are considered **omnivores**. Humans, bears, chickens, cockroaches, and crayfish are examples of omnivores.



Figure 3.3.4.*c*: Carnivores like the lion (top left) eat primarily meat. The ladybug (lower left) is also a carnivore that consumes small insects called aphids. Herbivores, like the mule deer (middle) eat primarily plant material. Omnivores like the bear (top right) and crayfish (bottom right) eat both plant and animal based food. Lion by Kevin Pluck; ladybug by Jon Sullivan; mule deer by Bill Ebbesen; bear by Dave Menke; crayfish by Jon Sullivan. All from OpenStax(opens in new window) (CC-BY(opens in new window)).

Dead producers and consumers are eaten by **detritivores** (which ingest dead tissues) and **decomposers** (which further break down these tissues into simple molecules by secreting digestive enzymes). Invertebrate animals, such as vultures, worms and millipedes are examples of detritivores, whereas fungi and certain bacteria are examples decomposers.

Food Chains

A **food chain** is a linear sequence of organisms through which nutrients and energy pass as one organism eats another. Each organism in a food chain occupies a specific **trophic level** (energy level), its position in the food chain. The first trophic level (TL) in the food chain is the producers. The **primary consumers** (the herbivores that eat producers) are the second trophic level. Next are higher-level consumers. Higher-level consumers include **secondary consumers** (third TL), which

are usually carnivores that eat the primary consumers, and **tertiary consumers** (fourth TL), which are carnivores that eat other carnivores. In the Lake Ontario food chain, shown in figure 3.3.4.*d*, the Chinook salmon is the apex consumer at the top of this food chain. Some communities have additional trophic levels (quaternary consumers, fifth order consumers, etc.). Finally, detritivores and decomposers break down dead and decaying organisms from any trophic level. There is a single path through a food chain.




Figure 3.3.4.*d*: These are the trophic levels of a food chain in Lake Ontario at the United States–Canada border. Energy and nutrients flow from photosynthetic green algae (producers) at the base to the primary consumers, which are mollusks, or snails. The secondary consumers are small fish called slimy sculpin. The tertiary and apex consumer is Chinook salmon. Detritivores and decomposers are not shown. (credit: modification of work by National Oceanic and Atmospheric Administration/NOAA)

One major factor that limits the number of steps in a food chain is energy. Only about 10% of the energy in one trophic level is transferred to the next trophic level. This is because much energy is lost as heat during transfers between trophic levels or to decomposers due to the second law of thermodynamics. Thus, after four to six trophic energy transfers, the amount of energy remaining in the food chain may not be great enough to support viable populations at higher trophic levels

Certain environmental toxins can become more concentrated as they move up the food chain, with the highest concentrations occurring in the top consumers, a process called **biomagnification**. Essentially, a top consumer ingests all the toxins that had previously accumulated in the bodies of the organisms at the lower trophic levels. This explains why frequently eating certain fish, like tuna or swordfish, increases your exposure to mercury, a toxic heavy metal.

Food Webs

While food chains are simple and easy to analyze, there is a one problem when using food chains to describe most communities. Even when all organisms are grouped into appropriate trophic levels, some of these organisms can feed at more than one trophic level. In addition, species feed on and are eaten by more than one species. In other words, the linear model of trophic interactions, the food chain, is a hypothetical and overly simplistic representation of community structure. A holistic model—which includes all the interactions between different species and their complex interconnected relationships with each other and with the environment—is a more accurate and descriptive model. A **food web** is a concept that accounts for the multiple trophic interactions between each species (figure 3.3.4.*e* and f).




Figure 3.3.4.*e*: This food web shows the interactions among organisms across trophic levels. Arrows point from an organism that is consumed to the organism that consumes it and represent energy transfer. The producers (plants) capture energy from the sun. The level above the producers shows the primary consumers that eat the producers. Some examples are squirrels, mice, seed-eating birds, and beetles. Spiders and centipedes eat beetles. Robins eat beetles, spiders, and centipedes, and toads eat beetles and centipedes. Foxes eat squirrels and mice; owls eat squirrels, mice, seed-eating birds, and robins; and snakes eat mice, seed-eating birds, robins, centipedes, and toads. Some consumers are in between trophic levels because they eat a combination of primary, secondary, and/or tertiary consumers. All the producers and consumers eventually become nourishment for the decomposers (mushrooms, mold, and bacteria) and detritivores (earthworms) in the soil, which are depicted at the bottom of the illustration. (credit "fox": modification of work by Kevin Bacher, NPS; credit "owl": modification of work by John and Karen Hollingsworth, USFWS; credit "snake": modification of work by Steve Jurvetson; credit "robin": modification of work by Alan Vernon; credit "frog": modification of work by Alessandro Catenazzi; credit "spider": modification of work by "Sanba38"/Wikimedia Commons; credit "centipede": modification of work by "Bauerph"/Wikimedia Commons; credit "squirrel": modification of work by Dawn Huczek; credit "mouse": modification of work by NIGMS, NIH; credit "sparrow": modification of work by David Friel; credit "beetle": modification of work by Scott Bauer, USDA Agricultural Research Service; credit "mushrooms": modification of work by Chris Wee; credit "mold": modification of work by Dr. Lucille Georg, CDC; credit "earthworm": modification of work by Rob Hille; credit "bacteria": modification of work by Don Stalons, CDC)




Figure 3.3.4.*f*: In this food web, each organism may have multiple food sources or be eaten by multiple species. Phytoplankton are the primary producers. They are consumed by sand lances, krill, and zooplankton. Sand lances are consumed by puffins, kittiwake, and cephalopods. Krill are consumed by the sand lance, cephalopods, auklets, and salmon. Zooplankton are consumed by the sand lance, krill, auklets, and salmon. Puffins are eaten by rats and gulls. Kittiwakes are eaten by rats, foxes, and gulls. Cephalopods are consumed by puffins and gulls. Auklets are consumed by gulls and foxes. Salmon are consumed by cephalopods and gulls. Rats are consumed by foxes and gulls. Gulls are consumed by foxes. Detritivores and decomposers are not shown. Image by Mariana Ruiz Villarreal (LadyofHats) for CK-12 Foundation (CC-BY-NC(opens in new window)).

Key Concept: Gross vs. Net Primary Productivity

- **Gross Primary Productivity (GPP):** The total energy captured by producers through photosynthesis.
- **Net Primary Productivity (NPP):** The energy remaining after producers use some for their own life processes.
- **Why it matters:** NPP determines how much energy is available to consumers and influences ecosystem productivity.

Energy Transfer, Food Webs, and Food Production

Only a small fraction of the energy available at one trophic level becomes biomass that can be used by the next trophic level. Much of the energy is lost as heat during metabolism, used for movement and other life processes, or released as waste. As a result, ecosystems contain less available energy at each higher trophic level. Ecologists estimate that, on average, only about 10% of the energy at one trophic level is transferred to the next, although the exact amount varies among ecosystems. This pattern is known as the **10% rule** and helps explain why food webs typically contain fewer organisms at higher trophic levels (figure $3.3.4.g$).




Figure $3.3.4.g$: Energy Pyramid highlighting 1% energy from the sun is captured by photosynthesizers. Thereafter, 10% energy transfer between each level. Image credit: Erin Rempala (CC-BY-NC-SA 4.0).

The 10% rule also helps explain why producing animal-based foods often requires more resources than producing plant-based foods. Livestock consume large amounts of plant biomass, but only a portion of that energy is converted into animal tissue. Most of the energy is used for growth, movement, body maintenance, and heat production. Because energy is lost at each trophic transfer, producing meat generally requires more land, water, and feed than producing an equivalent amount of plant-based food.

Understanding energy transfer helps scientists evaluate the environmental impacts of food production and identify strategies that support more sustainable food systems.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Energy, Energy Enters Ecosystems Through Photosynthesis, and Energy Flow Through Ecosystems from *Environmental*

Biology by Matthew R. Fisher (licensed under CC-BY(opens in new window))

- Ecology of Ecosystems, Energy Flow through Ecosystems, and Digestive Systems(opens in new window) from *General Biology/Biology 2e* by OpenStax (licensed under CC-BY(opens in new window)). Access for free at openstax.org(opens in new window).
- Ecosystem Productivity from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))

This page titled 3.3.4: Energy Flow in Food Webs is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 6.1.1.4: Food Chains and Food Webs by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.3.5: ECOSYSTEM TYPES AND DYNAMICS

An **ecosystem** is a community of organisms (**biotic components**) and their **abiotic** (non-living) environment. Ecosystems can be small, such as the tide pools found near the rocky shores of many oceans, or large, such as those found in the tropical rainforest of the Amazon in Brazil (figure $3.3.5.a$).



(a) (b)

Figure $3.3.5.a$: A (a) tidal pool ecosystem in Matinicus Island, Maine, is a small ecosystem, while the (b) Amazon rainforest in Brazil is a large ecosystem. (credit a: modification of work by Jim Kuhn; credit b: modification of work by Ivan Mlinaric)

Ecosystem Categories

There are three broad categories of ecosystems based on their general environment: freshwater, marine, and terrestrial. Within these three categories are individual ecosystem types based on the environmental habitat and organisms present.

Freshwater ecosystems are the least common, occurring on only 1.8 percent of Earth's surface. These systems comprise lakes, rivers, streams, and springs; they are quite diverse and support a variety of animals, plants, fungi, protists and prokaryotes.

Marine ecosystems are the most common, comprising 75 percent of Earth's surface and consisting of three basic types: shallow ocean, deep ocean water, and deep ocean bottom. Shallow ocean ecosystems include extremely biodiverse coral reef ecosystems. Small photosynthetic organisms suspended in ocean waters, collectively known as **phytoplankton,** perform 40 percent of all photosynthesis on Earth. Deep ocean bottom ecosystems contain a wide variety of marine organisms. These ecosystems are so deep that light is unable to reach them. Freshwater and marine ecosystems are found in **aquatic biomes**, which are discussed in the Biomes chapter.

Terrestrial ecosystems, also known for their diversity, are grouped into large categories called biomes. A **biome** is a large-scale community of organisms, primarily defined on land by the dominant plant types that exist in geographic regions of the planet with similar climatic conditions. Examples of biomes include tropical rainforests, savannas, deserts, grasslands, temperate forests, and tundras. Grouping these ecosystems into just a few biome categories obscures the great diversity of the individual ecosystems within them. For example, the saguaro cacti (*Carnegiea gigantean*) and other plant life in the Sonoran Desert, in the United States, are relatively diverse compared with the desolate rocky desert of Boa Vista, an island off the coast of Western Africa (figure $3.3.5.b$).



(a) (b)

Figure 3.3.5.*b*: Desert ecosystems, like all ecosystems, can vary greatly. The desert in (a) Saguaro National Park, Arizona, has abundant plant life, while the rocky desert of (b) Boa Vista island, Cape Verde, Africa, is devoid of plant life. (credit a: modification of work by Jay Galvin; credit b: modification of work by Ingo Wölbern)

Ecosystem Dynamics

Ecosystems are complex with many interacting parts. They are routinely exposed to various **disturbances**, or changes in the environment that affect their compositions: yearly variations in rainfall and temperature and the slower processes of plant growth, which may take several years. Many of these disturbances are a result of natural processes. For example, when lightning causes a forest fire and destroys part of a forest ecosystem, the ground is eventually populated by grasses, then by bushes and shrubs, and later by mature trees, restoring the forest to its former state. The impact of environmental disturbances caused by human activities is as important as the changes caused by natural processes. Human agricultural practices, air pollution, acid rain, global deforestation, overfishing, eutrophication, oil spills, and illegal dumping on land and into the ocean are all issues of concern to conservationists.

Equilibrium is the steady state of an ecosystem where all organisms are in balance with their environment and with each other. In ecology, two parameters are used to measure changes in ecosystems: resistance and resilience. The ability of an ecosystem to remain at equilibrium in spite of disturbances is called **resistance**. The speed at which an ecosystem recovers equilibrium after being disturbed, called its **resilience**. Ecosystem resistance and resilience are especially important when considering human impact. The nature of an ecosystem may change to such a degree that it can lose its resilience entirely. This process can lead to the complete destruction or irreversible altering of the ecosystem.

Foundation Species

Foundation species are considered the “base” or “bedrock” of an ecosystem, having the greatest influence on its overall structure. They are often primary producers, and they are typically an abundant organism. For example, kelp, a species of brown algae, is a foundation species that forms the basis of the kelp forests off the coast of California.

Foundation species may physically modify the environment to produce and maintain habitats that benefit the other organisms that use them. Examples include the kelp described above or tree species found in a forest. The photosynthetic corals of the coral reef also provide structure by physically modifying the environment (figure 3.3.5.*c*). The calcium carbonate deposits of living and dead coral make up most of the reef structure, which protects many other species from waves and ocean currents.



Figure 3.3.5.c: Coral is the foundation species of coral reef ecosystems. (credit: Jim E. Maragos, USFWS)

3.3.5.1: Climate and Biomes

3.3.5.2: Terrestrial Biomes

3.3.5.3: Aquatic Biomes

Attribution

Modified by Melissa Ha from the following sources:

- Community Ecology from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Ecology of Ecosystems from *General Biology* by OpenStax (licensed under CC-BY(opens in new window))

This page titled 3.3.5: Ecosystem Types and Dynamics is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 7.1: Ecosystem Types and Dynamics by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.3.5.1: Climate and Biomes

Climate and Biomes

Biomes are large-scale environments that are distinguished by characteristic climate and vegetation (figure 3.3.5.1.a). Biomes are also characterized by the animals and other organisms there, which are influenced by vegetation and climate patterns. The Earth's biomes are categorized into two major groups: terrestrial and aquatic. **Terrestrial biomes** are based on land, while **aquatic biomes** include both ocean and freshwater biomes. Altitude and latitude, which affect temperature and precipitation determine the distribution of biomes.




Figure 3.3.5.1.a: Average annual temperature and annual precipitation are two climatic factors that determine the distribution of biomes. This graph shows annual precipitation in centimeters on the y-axis and average annual temperature in degrees Celsius on the x-axis. Tropical rainforests have the highest temperatures and most precipitation. Tropical seasonal forests and savannas have high temperatures (20-30°C) with precipitation from approximately 50 to 270 centimeters. Subtropical deserts have similarly high temperatures but dry conditions. Temperate grasslands and cold deserts range in average annual temperature from below 0 to 22°C and have less than 50 centimeters of precipitation per year. Woodlands/shrublands (such as the chaparral) have the same temperature range as cold deserts but can receive a little over 100 cm of precipitation per year. Temperate forests have intermediate temperatures (approximately 16-22°C), but temperate rainforests receive more precipitation than temperate seasonal forests. Boreal forests receive a little less precipitation than temperate forests, but are colder (approximately 1-10°C). The tundra has the coldest temperatures and little precipitation. Deserts have a range of temperatures but low precipitation. Image by Navarras (public domain).

Low latitudes (near the equator) have high temperatures, while high latitudes (near the poles) have low temperatures. This is because the sun hits the equator more directly. Sunlight hits the poles at an angle, reducing the intensity of light (and heat energy) per unit of area. Temperature also decreases with altitude. At high altitudes, the atmosphere is thinner and traps less heat energy from the sun. Because temperatures decline with altitude as well as latitude, similar biomes exist on mountains even when they are at low latitudes. As a rule of thumb, a climb of 1000 feet (about 300 m) is equivalent in changed flora and fauna to a trip northward of some 600 miles (966 km).

Where precipitation is moderately abundant — 40 inches (about 1 m) or more per year — and distributed fairly evenly throughout the year, the major determinant is temperature. It is not simply a matter of average temperature, but includes such limiting factors as whether it ever freezes or length of the growing season. Biomes are thus characterized not only by average temperature and precipitation but also their seasonality.

Not only does latitude influence temperature, but it also affects precipitation. For example, deserts tend to occur at latitudes of around 30° and at the poles, both north and south, driven by circulation and prevailing wind patterns in the atmosphere. The engine that drives circulation in the atmosphere and oceans is solar energy, which is determined by the average position of the sun over the Earth's surface. Direct light provides uneven heating depending on latitude and angle of incidence, with high solar

energy in the tropics, and little or no energy at the poles. Atmospheric circulation and geographic location are the primary causal agents of deserts. At approximately 30° north and south of the equator, sinking air produces **trade wind deserts** like the Sahara and the Outback of Australia (figure 3.3.5.1.*b*).




Figure 3.3.5.1.*b*: The generalized circulation of the atmosphere. Solar energy falling on the equatorial belt heats the air and causes it to rise. The rising air cools and its contained moisture falls back on the tropics as rain. The drier air then continues to spread toward the north and south where it sinks back at about 30 degrees north and south latitudes. This sinking drier air creates belts of predominant high pressure along which desert conditions prevail. These belts of predominantly high pressure have air that descends along these belts and flows either north to become the westerlies or south to become the trade winds. Note the arrows indicating general directions of winds in the latitude zones. The trade winds are predominant in the tropics and the westerlies in the mid-latitudes.

The MinuteEarth video below discusses the global climate patterns which lead to deserts.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Rain Shadow Deserts in San Diego County

San Diego County provides a clear example of the **rain shadow effect** (figure 3.3.5.1.*c*). Moist air from the Pacific Ocean moves inland and rises over the Cuyamaca and Laguna Mountains. As the air rises, it cools and releases much of its moisture as rain on the western side of the mountains. After crossing the mountains, the air descends, warms, and becomes much drier. As a result, eastern San Diego County receives far less precipitation than coastal and mountain areas. This rain shadow creates the desert conditions found in Anza-Borrego Desert State Park, where annual rainfall is much lower than along the coast. The rain shadow effect helps explain the dramatic differences in climate, vegetation, and biodiversity found across San Diego County, from coastal sage scrub and chaparral near the coast to desert ecosystems farther inland. For example, the coast may receive 10–15 inches (25–38 cm) of rain annually, while some desert areas east of the mountains receive less than 5 inches (13 cm) per year.




Figure 3.3.5.1.*c*: The rain shadow effect. Warm, moist air is carried up a mountain by prevailing winds. The rising air cools and condenses, resulting in precipitation as it moves up the mountain. The dry air advances down the other side of the mountain, resulting in a rain shadow (a dry region). Image by domdomegg (CC-BY).

The rain shadow effect demonstrates how Earth's systems interact. Mountains influence atmospheric circulation and precipitation patterns, which in turn affect water availability, ecosystem distribution, biodiversity, wildfire risk, and human land use across San Diego County.

Attributions

Modified by Erin Rempala.

Modified by Melissa Ha from the following sources:

- Biomes from *Biology* by John W. Kimball (licensed under CC-BY)
- Prelude to Deserts and The Origin of Deserts from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC BY-NC-SA)
- Community Ecology and Terrestrial Biomes from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)
- Terrestrial Biomes from *Biology 2e* by OpenStax (CC-BY). Access for free at openstax.org.

This page titled 3.3.5.1: Climate and Biomes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 8.1: Climate and Biomes by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

3.3.5.2: Terrestrial Biomes

Major Terrestrial Biomes

Because each biome is defined by climate, the same biome can occur in geographically distinct areas with similar climates (figure $3.3.5.2.a$). Scientists commonly recognize eight major terrestrial biomes: tropical rainforests, savannas, deserts, chaparral, temperate grasslands, temperate forests, taiga (boreal forests), and Arctic tundra. This classification simplifies the wide variety of ecosystems found on Earth, and many areas contain characteristics of multiple biomes or transition gradually between them.




Figure $3.3.5.2.a$: Each of the world's eight major biomes is distinguished by characteristic temperatures and amount of precipitation. Image By OpenStax - https://cnx.org/contents/z4fCt9js@6/...fig-ch44_03_01, CC BY-SA 4.0

Tropical Rainforests

Tropical rainforests are warm, wet ecosystems that contain more species than any other terrestrial biome (figure $3.3.5.2.a$). This rich biodiversity is threatened by deforestation and land conversion for agriculture. Many rainforest organisms produce unique chemicals that may be useful in developing new medicines, earning rainforests the nickname "nature's pharmacy." Temperatures remain warm year-round, typically between 20°C and 34°C (68–93°F), and abundant rainfall supports continuous plant growth. Most trees are evergreen and form several layers of vegetation, including a dense canopy that shades the forest floor. These layers create a variety of habitats that support the rainforest's exceptional diversity of plants, animals, fungi, and other organisms.



Figure $3.3.5.2.a$: Image: Moist broadleaf forest in Mudumalai National Park. By Kmathanraj - Own work, CC BY-SA 4.0.

Tropical Soils

While productivity is high in tropical rainforests, the soils themselves tend to be of very poor quality. Because of the high rainfall, nutrients are quickly washed out of the topsoil unless they are incorporated in the forest plants. As plant and animal debris falls to the ground, it is quickly decomposed because of the warmth and moisture there. Thus minerals are found mainly in the forest plants, not in the soil. When the plants are removed and cultivation attempted, the soils quickly lose fertility.

Savannas

Savannas have mostly **perennial, herbaceous** plants (non-woody plants that persist for more than one year) with scattered trees, and they are located in Africa, South America, and northern Australia. Savannas are usually hot, tropical areas with temperatures averaging from 24 °C to 29 °C (75 °F to 84 °F) and an annual rainfall of 10–40 cm (3.9–15.7 in). Savannas have an extensive dry season; for this reason, forest trees do not grow as well as they do in the tropical rainforest. As a result, grasses and **forbs** (herbaceous plants other than grasses) are dominant (figure $3.3.5.2.d - e$). Fire is an important source of disturbance in this biome and plants have evolved well-developed root systems that allow them to quickly resprout after a fire. Depending on the region, large grazing animals like giraffes, zebras, and gazelles and their predators (cheetahs, lions, etc.) are found in savannas.



Figure 3.3.5.2.d: Savannas, like this one in Taita Hills Wildlife Sanctuary in Kenya, are dominated by grasses. (credit: Christopher T. Cooper)



Figure 3.3.5.2.e: Although savannas are dominated by grasses, small woodlands, such as this one in Mount Archer National Park in Queensland, Australia, may dot the landscape. (credit: "Ethel Aardvark"/Wikimedia Commons)

Deserts

Deserts are dry regions where evaporation exceeds precipitation and annual rainfall is usually less than 25 cm (10 in). Some deserts are hot, while others experience cold winters. Because water is scarce and rainfall is unpredictable, desert plants and animals have adaptations that help them conserve water and survive extreme conditions.

Many deserts form where dry air descends from the atmosphere or where mountains block moist air, creating a rain shadow. Deserts can be hot, such as the Sahara and Sonoran deserts, or cold, such as the Gobi and Great Basin deserts. Because water is scarce and rainfall is unpredictable, deserts generally support fewer species than many other biomes. However, desert plants and animals have special adaptations that help them survive. Many plants have deep roots, small leaves, or water-storing tissues, while many animals avoid the daytime heat by being active at night (figure 3.3.5.2.f). Despite their harsh conditions, deserts support unique species that are well adapted to these environments (figure 3.3.5.2.g).



Figure 3.3.5.2.f: Blair Valley- Anza Borrego Desert State Park. Many desert plants have tiny leaves to reduce water loss. Ocotillo, teddy bear cholla, and yucca. Image By Wasquewhat - Own work, CC BY-SA 4.0.




Figure 3.3.5.2.g: Desert Bighorn Sheep at Palm Canyon oasis, Anza-Borrego Desert State Park. Image by By Greg Bulla. Original uploader was Gb1 at en.wikipedia - Transferred from en.wikipedia, CC BY-SA 2.5,

Many of the animals in the desert (mammals, lizards and snakes, insects, and even some birds) are adapted for burrowing to escape the scorching heat of the desert sun. Many of them limit their forays for food to the night.

The net primary productivity of the desert is low. High productivity can sometimes be achieved with **irrigation**, but these gains are often only temporary because it leads to the buildup of **salts** and **minerals**. Large areas of formerly unproductive desert in the United States, Israel, and Egypt have been converted into fertile fields through irrigation. However, even the best irrigation water contains dissolved salts. Because the rainfall in deserts is so low, any water that does not immediately run off remains near the surface and is largely lost by evaporation. The salts it carries are left near the top of the soil. Their accumulation may make the soil so alkaline (basic) and so salty that it is prohibitive to agriculture (figure 3.3.5.2.g). In the U. S., the situation is especially severe in the Great Basin because water flowing down from the mountains—bearing its load of dissolved salts—cannot flow on to the ocean but simply flows out onto the valley floors and evaporates.



Figure 3.3.5.2.*g*: Salt formations at the Badwater Basin in Death Valley, California. Minerals accumulate when water pools in this low-lying area, making conditions impossible for agriculture. Some plants, like pickleweed and saltgrass, are adapted to grow around the edges of the salt flats. Image by "Mike" Michael L. Baird (CC-BY).

Chaparral (Mediterranean Scrub)

The **Mediterranean scrub**, is also called **chaparral** or scrub forest and is found only in California, along the Mediterranean Sea, along the coast of Chile, South Africa, and the southern coast of Australia (figure 3.3.5.2.*h*). The annual rainfall in this biome ranges from 65 cm to 75 cm (25.6–29.5 in) and the majority of the rain falls in the winter. Summers are very dry and many chaparral plants are dormant during the summertime.



Figure 3.3.5.2.*h*: The chaparral is dominated by shrubs. (credit: Miguel Vieira)

The Mediterranean scrub vegetation is dominated by shrubs and is adapted to periodic fires, with some plants producing seeds that germinate only after a hot fire. The ashes left behind after a fire are rich in nutrients like nitrogen and fertilize the soil, promoting plant regrowth. Fire is a natural part of the maintenance of this biome. The trees in the chaparral are mostly oaks, both deciduous and evergreen. Scrub oaks and shrubs like manzanita and the California lilac (not a relative of the eastern lilac) form dense, evergreen thickets. All of these plants are adapted to drought by such mechanisms as waxy, waterproof coatings on their leaves.

Many plants that thrive in one chaparral region are successfully cultivated in other chaparral regions. For example, vineyards, olives, and figs from their native Mediterranean and eucalyptus trees from Australia thrive in California. However, blue gum eucalyptus from Australia disrupts California ecosystems by changing the hydrology and outcompeting native plants.

Temperate Grasslands

Temperate grasslands are found throughout central North America, where they are also known as **prairies** (figure 3.3.5.2.*i*), and in Eurasia, where they are known as **steppes.** Temperate grasslands have pronounced annual fluctuations in temperature with hot summers and cold winters. The annual temperature variation produces specific growing seasons for plants. Plant growth is possible when temperatures are warm enough to sustain plant growth, which occurs in the spring, summer, and fall.



Figure 3.3.5.2.*h*: The American bison (Bison bison), more commonly called the buffalo, is a grazing mammal that once populated American prairies in huge numbers. (credit: Jack Dykinga, USDA ARS)

Temperate grasslands have few trees except for those found growing along rivers or streams. The dominant vegetation tends to consist of grasses. The treeless condition is maintained by low precipitation, frequent fires, and grazing. Fires, which are a natural disturbance in temperate grasslands, can be ignited by lightning strikes. It also appears that the lightning-caused fire regime in North American grasslands was enhanced by intentional burning by humans. When fire is suppressed in temperate grasslands, the vegetation eventually converts to scrub and dense forests. Often, the restoration or management of temperate grasslands requires the use of controlled burns to suppress the growth of trees and maintain the grasses.

Annual precipitation ranges from 25.4 cm to 88.9 cm (10–35 in) and is concentrated in the summer and spring. In the plains of North America, the annual rainfall is sufficiently low (~50 cm, 20") that little or no rainfall percolates down to the water table. Calcium and other minerals are not carried below the reach of plant roots and so remain available for use. This keeps the pH and general fertility high. The grasses in undisturbed prairie are perennial. Their extensive root systems help prevent soil erosion, and the return of the season's above-ground growth to the topsoil returns minerals and provides humus to it. These advantages are diminished when annual grasses such as wheat and corn are grown instead and removed in the harvest.The self-restoring fertility of the soils of the plains states accounts for this region being the "breadbasket" of the nation (and other countries as well).

Temperate Forests

Temperate forests are the most common biome in eastern North America, Western Europe, Eastern Asia, Chile, and New Zealand (figure 3.3.5.2.*i*). This biome is found throughout mid-latitude regions. Temperatures range between –30°C and 30°C (–22°F to 86°F) and drop to below freezing on an annual basis. These temperatures mean that temperate forests have defined growing seasons during the spring, summer, and early fall.



Figure 3.3.5.2.*i*: Deciduous trees are the dominant plant in the temperate forest. (credit: Oliver Herold)

Temperate forests may be deciduous, mixed (with deciduous and evergreen trees), or coniferous (with evergreen conifers). Examples of temperate deciduous trees are beech, maple, oak, and hickory. The deciduous trees of temperate forests drop their leaves in the winter; thus, little photosynthesis occurs during the dormant winter period. (In other biomes, such as chaparral and tropical seasonal forests, trees or shrubs may drop their leaves during the summer or dry season.) Each spring, new leaves appear as temperature increases. Because of the dormant period, the net primary productivity of temperate forests is less than that of tropical rainforests. In addition, temperate forests show far less diversity of tree species than tropical rainforest biomes. Large stands dominated by a single species are common.

The deciduous trees of the temperate forests leaf out and shade much of the ground. However, more sunlight reaches the ground in this biome than in tropical rainforests because trees in temperate forests do not grow as tall as the trees in tropical rainforests. The soils of the temperate forests are rich in inorganic and organic nutrients compared to tropical rainforests. This is because of the thick layer of leaf litter on forest floors and reduced leaching of nutrients by rainfall. As this leaf litter decays, nutrients are returned to the soil. The leaf litter also protects soil from erosion, insulates the ground, and provides habitats for invertebrates and their predators. Deer, raccoons, and salamanders are characteristic inhabitants of temperate forests.

Precipitation is relatively constant throughout the year and ranges between 75 cm and 150 cm (29.5–59 in). Enough water falls on the soil so that much of it soaks deep into the ground and carries minerals with it. Such soils tend to be acidic and of low and (if unattended) diminishing fertility. Only by regular fertilization and liming (to restore calcium and raise pH) can productive agriculture be carried out in them. In the U.S., the soils east of the Appalachian Mountains tend to be of this sort.

Taiga (Boreal Forest)

The **taiga**, also known as **boreal forest** in North America, is found roughly between 50° and 60° north latitude across most of Canada, Alaska, Russia, and northern Europe ((figure 3.3.5.2.*i*). Boreal forests are also found above a certain elevation (and below high elevations where trees cannot grow) in mountain ranges throughout the Northern Hemisphere. This biome has cold, dry winters and short, cool, wet summers. The annual precipitation is from 40 cm to 100 cm (15.7–39 in) and usually takes the form of snow. Although precipitation is relatively low, the taiga is dotted with lakes, bogs, and marshes because the cold temperatures limit evaporation.



Figure 3.3.5.2.j: The boreal forest (taiga) has low lying plants and conifer trees. (credit: L.B. Brubaker, NOAA)

The long and cold winters in the boreal forest have led to the predominance of cold-tolerant cone-bearing plants. These are evergreen coniferous trees like pines, spruce, and fir, which retain their needle-shaped leaves year-round. Evergreen trees can photosynthesize earlier in the spring than deciduous trees because less energy from the sun is required to warm a needle-like leaf than a broad leaf. In the taiga, evergreen trees grow faster than deciduous trees. In addition, soils in boreal forest regions tend to be acidic with little available nitrogen. Leaves are a nitrogen-rich structure, and deciduous trees must produce a new set of these nitrogen-rich structures each year. Therefore, coniferous trees that retain nitrogen-rich needles in a nitrogen-limiting environment may have had a competitive advantage over the broad-leafed deciduous trees.

In North America, the moose is such a typical member that it has led to the name: "spruce-moose" biome. Deer, bears, and wolves are also found in the taiga. Before the long, snowy winter sets in, many of the mammals hibernate, and many of the birds migrate south.

The net primary productivity of taiga is lower than that of temperate forests and tropical rainforests. The aboveground **biomass** (mass of living organisms) of taiga is high because these slow-growing tree species are long-lived and accumulate standing biomass over time. Species diversity is less than that seen in temperate forests and tropical rainforests. Taiga lacks the layered forest structure seen in tropical rainforests or, to a lesser degree, temperate forests. The structure of the taiga is often only a layer of trees and the understory of low-lying plants and lichen. When conifer needles are dropped, they decompose more slowly than broad leaves; therefore, fewer nutrients are returned to the soil to fuel plant growth.

At the coldest parts of the taiga, the soil remains frozen year after year, forming a layer called **permafrost**. When the permafrost melts in unusually warm years, the trees are no longer supported by this hard surface and lean in different directions, forming a "drunken forest". At extreme latitudes, the trees of the taiga become stunted by the harshness of the subarctic climate. Where they finally disappear (above the "tree line"), the Arctic tundra begins.

Arctic Tundra

The **Arctic tundra** lies north of the subarctic boreal forests and is located throughout the Arctic regions of the Northern Hemisphere. Tundra also exists at elevations above the tree line on mountains. The average winter temperature is –34°C (–29.2°F), and the average summer temperature is 3°C–12°C (37°F –52°F). Plants in the Arctic tundra have a short growing season of approximately 50–60 days. However, during this time, there are almost 24 hours of daylight and plant growth is rapid. The annual precipitation of the Arctic tundra is low (15–25 cm or 6–10 in) with little annual variation in precipitation. And, as in the taiga, there is little evaporation because of the cold temperatures.

There is little species diversity, low net primary productivity, and low above-ground biomass. Like some of the taiga, the soils of the Arctic tundra may remain in a perennially frozen state, forming permafrost. The permafrost makes it impossible for roots to penetrate far into the soil and slows the decay of organic matter, which inhibits the release of nutrients from organic matter. The surface layer of soil above the permafrost melts in the brief summer, water is released and provides for a burst of productivity

while temperatures and long days permit it. Because of these harsh growing conditions, the Arctic tundra is particularly sensitive to disturbances (disruptions that decrease biomass, such as heavy machinery passing through and squishing some of the plants).

Plants in the Arctic tundra are generally low to the ground and include low shrubs, sphagnum moss, grasses, lichens, and small flowering plants (figure $3.3.5.2.k$). Caribou feed on this growth as do vast numbers of insects. Swarms of migrating birds, especially waterfowl, invade the tundra in the summer to raise their young, feeding them on a large variety of aquatic invertebrates and vertebrates. As the brief arctic summer draws to a close, the birds fly south, and all but a few of the permanent residents, in one way or another, prepare themselves to spend the winter in a dormant state.



Figure $3.3.5.2.k$: Low-growing plants such lichen and grasses are common in tundra. Credit: Nunavut tundra(opens in new window) by Flickr: My Nunavut is licensed under CC-BY.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Terrestrial Biomes from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)
- Biomes and Soil from *Biology* by John W. Kimball (licensed under CC-BY)
- Terrestrial Biomes from *Biology 2e* by OpenStax (CC-BY). Access for free at openstax.org.

This page titled 3.3.5.2: Terrestrial Biomes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 8.2: Terrestrial Biomes by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

3.3.5.3: Aquatic Biomes

Abiotic Factors Influencing Aquatic Biomes

Like terrestrial biomes, **aquatic biomes** are influenced by a series of abiotic factors. The aquatic medium—water— has different physical and chemical properties than air. Even if the water in a pond or other body of water is perfectly clear (there are no suspended particles), water still absorbs light. As one descends into a deep body of water, there will eventually be a depth which the sunlight cannot reach. While there are some abiotic and biotic factors in a terrestrial ecosystem that might obscure light (like fog, dust, or insect swarms), usually these are not permanent features of the environment. The importance of light in aquatic biomes is central to the communities of organisms found in both freshwater and marine ecosystems. In freshwater systems, temperature stratification due to differences in density is perhaps the most critical abiotic factor and is related to the energy aspects of light. The thermal properties of water (rates of heating and cooling) are significant to the function of marine systems and have major impacts on global climate and weather patterns. Marine systems are also influenced by large-scale physical water movements, such as currents; these are less important in most freshwater lakes.

Marine Biomes

The ocean is the largest **marine biome**. It is a continuous body of salt water that is relatively uniform in chemical composition; it is a weak solution of mineral salts and decayed biological matter. Within the ocean, coral reefs are a second kind of marine biome. Estuaries, coastal areas where salt water and fresh water mix, form a third unique marine biome.

Ocean

The physical diversity of the ocean is a significant influence on plants, animals, and other organisms. The ocean is categorized by several areas (Figure $3.3.5.3.a$). Each area has a distinct group of species adapted to the biotic and abiotic conditions particular to it. The **intertidal zone**, which is the zone between high and low tide, is the oceanic region that is closest to land. Generally, most people think of this portion of the ocean as a sandy beach. In some cases, the intertidal zone is indeed a sandy beach, but it can also be rocky or muddy. Organisms are exposed to air and sunlight at low tide and are underwater most of the time, especially during high tide. Therefore, living things that thrive in the intertidal zone are adapted to being dry for long periods of time. The shore of the intertidal zone is also repeatedly struck by waves, and the organisms found there are adapted to withstand damage from the pounding action of the waves (figure $3.3.5.3.b$). The exoskeletons of shoreline crustaceans (such as the shore crab, *Carcinus maenas*) are tough and protect them from desiccation (drying out) and wave damage. Another consequence of the pounding waves is that few algae and plants establish themselves in the constantly moving rocks, sand, or mud.




Figure 3.3.5.3.*a*: The ocean is divided into different zones based on distance from the shoreline and water depth. The intertidal zone is the closest to shore followed by the neretic and oceanic zone. The photic zone is 0-200 meters deep. The aphotic zone is 200-4,000 meters deep. The abyssal zone is 4,000-10,000 meters deep. These three zones form the pelagic realm, and the benthic realm is below the pelagic realm, along the continental shelf.



Figure 3.3.5.3.*b*: Sea urchins, mussel shells, and starfish are often found in the intertidal zone, shown here in Kachemak Bay, Alaska. (credit: NOAA)

The **neritic zone** extends from the intertidal zone to depths of about 200 m (or 650 ft) at the edge of the continental shelf. Because light can penetrate this depth, photosynthesis can occur. The water here contains silt and is well-oxygenated, low in pressure, and stable in temperature. Phytoplankton and floating *Sargassum* (a type of free-floating marine seaweed) provide a habitat for some sea life found in the neritic zone. Zooplankton, protists, small fishes, and shrimp are found in the neritic zone and are the base of the food chain for most of the world's fisheries.

Beyond the neritic zone is the open ocean area known as the **oceanic zone**. Within the oceanic zone there is thermal stratification where warm and cold waters mix because of ocean currents. Abundant plankton serve as the base of the food chain for larger animals such as whales and dolphins. Nutrients are scarce and this is a relatively less productive part of the marine biome. When photosynthetic organisms and the protists and animals that feed on them die, their bodies fall to the bottom of the ocean where they remain.

All of the ocean's open water is referred to as the **pelagic realm**. The pelagic realm is divided into the photic, aphotic, and abyssal zones from top to bottom based on how far light reaches into the water. The **photic zone**, which is the portion of the ocean that

light can penetrate (approximately 200 m or 650 ft). At depths greater than 200 m, light cannot penetrate; thus, this is referred to as the **aphotic zone**. The majority of organisms in the aphotic zone include sea cucumbers (phylum Echinodermata) and other organisms that survive on the nutrients contained in the dead bodies of organisms in the photic zone.

The deepest part of the ocean is the **abyssal zone**, which is at depths of 4000 m or greater. Both the aphotic and abyssal zones lack sufficient light for photosynthesis, and together, they constitute most of the ocean. The deepest part of the ocean, the Challenger Deep (in the Mariana Trench, located in the western Pacific Ocean), is about 11,000 m (about 6.8 mi) deep. To give some perspective on the depth of this trench, the ocean is, on average, 4267 m deep. These zones are relevant to freshwater lakes as well. The abyssal zone is very cold and has very high pressure, high oxygen content, and low nutrient content. There are a variety of invertebrates and fishes found in this zone, but the abyssal zone does not have plants because of the lack of light. Cracks in the Earth's crust called **hydrothermal vents** are found primarily in the abyssal zone (figure $3.3.5.3.c$). Around these vents, bacteria that utilize the hydrogen sulfide and other minerals emitted as an energy source serve as the base of the food chain found in the abyssal zone.




Figure $3.3.5.3.c$: A hydrothermal vent. Image by NOAA (public domain).

The **benthic realm**, extends along the ocean bottom from the shoreline to the deepest parts of the ocean floor. It is comprised of sand, silt, and dead organisms. This is a nutrient-rich portion of the ocean because of the dead organisms that fall from the upper layers of the ocean. Because of this high level of nutrients, a diversity of sponges, sea anemones, marine worms, sea stars, fishes, and bacteria exist.

Coral Reefs

Coral reefs are characterized by high biodiversity and the structures created by invertebrates that live in warm, shallow waters within the photic zone of the ocean. They are mostly found within 30 degrees north and south of the equator. The Great Barrier Reef is a well-known reef system located several miles off the northeastern coast of Australia. The coral organisms are colonies of saltwater polyps that secrete a calcium carbonate skeleton. These calcium-rich skeletons slowly accumulate, forming the underwater reef (figure $3.3.5.3.d$).




Figure 3.3.5.3.*d*: Coral reefs are formed by the calcium carbonate skeletons of coral organisms, which are marine invertebrates. (credit: Terry Hughes)

Corals found in shallower waters (at a depth of approximately 60 m or about 200 ft) have a mutualistic relationship with photosynthetic unicellular algae called dinoflagellates. The relationship provides corals with the majority of the nutrition and the energy they require. The waters in which these corals live are nutritionally poor, and, without this mutualism, it would not be possible for large corals to grow. Some corals living in deeper and colder water do not have a mutualistic relationship with algae; these corals attain energy and nutrients using stinging cells on their tentacles to capture prey. It is estimated that more than 4,000 fish species inhabit coral reefs. These fishes can feed on coral, other invertebrates, or the seaweed and algae that are associated with the coral.

Estuaries: Where the Ocean Meets Fresh Water

Estuaries are biomes that occur where a source of fresh water, such as a river, meets the ocean. Therefore, both fresh water and salt water are found in the same vicinity; mixing results in a diluted (brackish) saltwater. Estuaries form protected areas where many of the young offspring of crustaceans, mollusks, and fish begin their lives. Salinity is a very important factor that influences the organisms and the adaptations of the organisms found in estuaries. The salinity of estuaries varies and is based on the rate of flow of its freshwater sources. Once or twice a day, high tides bring salt water into the estuary. Low tides occurring at the same frequency reverse the current of salt water.

The short-term and rapid variation in salinity due to the mixing of fresh water and salt water is a difficult physiological challenge for the plants and animals that inhabit estuaries. Many estuarine plant species are **halophytes**, plants that can tolerate salty conditions. Halophytic plants are adapted to deal with the salinity resulting from saltwater on their roots or from sea spray. In some halophytes, filters in the roots remove the salt from the water that the plant absorbs. Other plants are able to pump oxygen into their roots. Animals, such as mussels and clams, have developed behavioral adaptations that expend a lot of energy to function in this rapidly changing environment. When these animals are exposed to low salinity, they stop feeding, close their shells, and stop using oxygen. When high tide returns to the estuary, the salinity and oxygen content of the water increases, and these animals open their shells, begin feeding, and return to using oxygen.

Estuarine ecosystems include mangrove forests, seagrass beds, salt marshes, mudflats, and oyster reefs. These habitats provide nursery grounds for fish, improve water quality, store carbon, and protect coastlines from storms.

Freshwater Biomes

Freshwater biomes include lakes and ponds (standing water) as well as rivers and streams (flowing water). They also include wetlands, which will be discussed later. Humans rely on freshwater biomes to provide aquatic resources for drinking water, crop irrigation, sanitation, and industry. These various roles and human benefits are referred to as ecosystem services. Lakes and ponds are found in terrestrial landscapes and are, therefore, connected with abiotic and biotic factors influencing these terrestrial

biomes.

Lakes and Ponds

Lakes and ponds can range in area from a few square meters to thousands of square kilometers. Temperature is an important abiotic factor affecting living things found in lakes and ponds. In the summer, thermal stratification of lakes and ponds occurs when the upper layer of water is warmed by the sun and does not mix with deeper, cooler water. Like oceans, lakes and ponds have photoic zones through which light can penetrate and aphotic zones without light. **Phytoplankton** (small photosynthetic organisms such as algae and photosynthetic bacteria that float in the water) are found here and carry out photosynthesis, providing the base of the food web of lakes and ponds. **Zooplankton** (very small animals that float in the water), such as rotifers and small crustaceans, consume these phytoplankton (figure $3.3.5.3.e$). At the bottom of lakes and ponds, bacteria in the aphotic zone break down dead organisms that sink to the bottom.




Figure $3.3.5.3.e$: A rotifer is an example of the zooplankton found in lakes and ponds. Bob Blaylock by Bob Blaylock at English Wikipedia (CC-BY-SA).

Rivers and Streams

Rivers and streams are continuously moving bodies of water that carry large amounts of water from the source, or headwater, to a lake or ocean. The largest rivers include the Nile River in Africa, the Amazon River in South America (Figure $3.3.5.3.f$), and the Mississippi River in North America. Abiotic features of rivers and streams vary along the length of the river or stream. Streams begin at a point of origin referred to as source water. The source water is usually cold, low in nutrients, and clear. The channel (the width of the river or stream) is narrower than at any other place along the length of the river or stream. Because of this, the current is often faster here than at any other point of the river or stream.



Figure 3.3.5.3.f: The Amazon River is the world's largest river. Image by Jason Hollinger (CC-BY)

The fast-moving water results in minimal silt accumulation at the bottom of the river or stream, therefore the water is clear. Photosynthesis here is mostly attributed to algae that are growing on rocks; the swift current inhibits the growth of phytoplankton. An additional input of energy can come from leaves or other organic material that falls into the river or stream from trees and other plants that border the water. When the leaves decompose, the organic material and nutrients in the leaves are returned to the water. Plants and animals have adapted to this fast-moving water. For instance, leeches have elongated bodies and suckers on both ends. These suckers attach to the substrate, keeping the leech anchored in place. Freshwater trout species are an important predator in these fast-moving rivers and streams.

As the river or stream flows away from the source, the width of the channel gradually widens and the current slows. This slow-moving water, caused by the gradient decrease and the volume increase as tributaries unite, has more sedimentation. Phytoplankton can also be suspended in slow-moving water. Therefore, the water will not be as clear as it is near the source. The water is also warmer. Worms and insects can be found burrowing into the mud. The higher order predator vertebrates include waterfowl, frogs, and fishes.

Wetlands

Wetlands are environments in which the soil is either permanently or periodically saturated with water. Wetlands are different from lakes because wetlands are shallow bodies of water that may periodically dry out. Emergent vegetation consists of wetland plants that are rooted in the soil but have portions of leaves, stems, and flowers extending above the water's surface. There are several types of wetlands including marshes, swamps, bogs, mudflats, and salt marshes (figure 3.3.5.3.g).



Figure 3.3.5.3.g: Located in southern Florida, Everglades National Park is vast array of wetland environments, including sawgrass marshes, cypress swamps, and estuarine mangrove forests. Here, a Great Egret walks among cypress trees. (credit: NPS)

Attribution

Modified by Erin Rempala.

Modified by Melissa Ha from Aquatic Biomes from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 3.3.5.3: Aquatic Biomes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 8.3: Aquatic Biomes by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

3.3.6: Data Dive- Biome Carbon Storage

Overview

Carbon sequestration is the process by which atmospheric carbon dioxide is taken up by trees, grasses, and other plants through photosynthesis and stored as carbon in biomass (trunks, branches, foliage, and roots). Carbon storage can also occur in soil formed from organic matter found in decomposing litter. Terrestrial carbon sequestration has been a recent topic of interest because of its implications for use in climate change mitigation. The carbon storage in terrestrial biomes helps to offset sources of carbon dioxide to the atmosphere, such as deforestation, wildfires, and fossil fuel emissions. As such, carbon storage in each biome can be altered by both natural and human caused events. The figure below indicates the estimated storage above (plants) and below (soil) ground by terrestrial biomes:




Figure 3.3.6.*a*: Carbon storage (in metric tons of CO2 per hectare) above (plant) and below (soil) ground in various biomes. Graph by Erin Rempala modified from Rachel Schleiger (CC-BY-NC) modified from data in Islam KK, Sato N 2012.

Questions

1. What kind of graph is this?
2. What is the independent (explanatory) variable and the dependent (response) variable?
3. Which biome has the greatest storage in plants? What about soil?
4. What are the top three terrestrial biomes in terms of their carbon storage?
5. How can the results of this graph to inform future conservation efforts?
6. What do the results of this graph make you curious about?

Raw Data From Above Graph(s)

Table 3.3.6.*a*: Raw data of carbon storage (in metric tons of CO2 per hectare) above (plant) and below (soil) ground in various biomes. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Islam KK, Sato N 2012.

Raw data of carbon storage

Biome	Plant Carbon	Soil Carbon	Total Carbon
Tropical Forests	442	450	892

Biome	Plant Carbon	Soil Carbon	Total Carbon
Temperate Forests	208	352	561
Taiga Forests	236	1260	1496
Tundra	23	467	490
Tropical Savannas	108	430	538
Temperate Grasslands	26	865	892
Deserts/Semi-Desert Lands	6	154	160

Attribution

Rachel Schleiger (CC-BY-NC)

This page titled 3.3.6: Data Dive- Biome Carbon Storage is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 8.4: Data Dive- Biome Carbon Storage by Melissa Ha and Rachel Schleiger has no license indicated.

4: EVOLUTION

Introduction

Life on Earth is not static, it is always changing.

Evolution is the process by which populations of organisms change over generations as inherited traits become more or less common in a population.This is the cause of the wide variety of life we see today!

How did the wide variety of life on Earth develop? In this chapter, you will learn how scientists discovered and tested the theory of evolution. You will investigate the mechanisms that drive evolutionary change, the evidence that supports these ideas, and the ways organisms adapt to their environments. Evolution provides the framework that helps us understand both the diversity and interconnectedness of life.

4.1: Evolution and Natural Selection

4.2: Other Mechanisms of Evolution

4.3: Evidence for Evolution

Thumbnail Darwin's finches By John Gould (14.Sep.1804 - 3.Feb.1881) - From *Voyage of the Beagle*. Public Domain.

This page titled 4: Evolution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 11: Evolution and Its Processes by OpenStax is licensed CC BY 4.0.

4.1: Evolution and Natural Selection

How do populations adapt to environmental change over time?

Life on Earth has changed over billions of years. Scientists estimate that millions of species have lived on Earth, and many are now extinct. The great diversity of life that exists today developed through the process of evolution.

Evolution helps explain why organisms are adapted to their environments and why species continue to change as environmental conditions change. Understanding evolution is important in environmental science because it helps us understand biodiversity, conservation, response of species to climate change impacts, and ecosystem resilience. Evolution can be defined as measurable genetic change in a population over generations.

Early Ideas About Evolution

Long before scientists understood evolution, people tried to explain the diversity of life on Earth. For many centuries, many believed that species never changed after they were created.

During the eighteenth century, the naturalist Georges-Louis Leclerc, Comte de Buffon suggested that species could change over time. Around the same period, the Scottish geologist James Hutton proposed that Earth changes gradually through slow processes that continue today. Later, Charles Lyell expanded these ideas and argued that Earth was much older than previously believed. A French scientist named Jean-Baptiste Lamarck proposed an early explanation for evolution. He suggested that traits acquired during an organism's lifetime could be passed to offspring. Although this explanation was later shown to be incorrect, Lamarck's work encouraged scientists to consider that species might change through time.

Together, these ideas helped establish the foundation for the theory of evolution. Scientific understanding changes as new evidence becomes available. Evolutionary theory is an example of how science develops through observation, testing, and revision.

Darwin, Wallace, and Natural Selection

In the mid-nineteenth century, Charles Darwin and Alfred Russel Wallace independently developed the theory of evolution by natural selection (Figure $4.1.a$).



(a) (b)

Figure 4.1.*a*: (a) Charles Darwin and (b) Alfred Wallace wrote scientific papers on natural selection that were presented together before the Linnean Society in 1858.

Darwin spent five years traveling aboard the HMS Beagle. During this voyage, he observed plants, animals, fossils, and geological features from around the world. Wallace made similar observations during expeditions in South America and Southeast Asia.

One of Darwin's most important observations came from the Galápagos Islands. He noticed that finches on different islands were similar but had different beak shapes. Some species had large beaks for cracking hard seeds, while others had smaller beaks better suited for other food sources (Figure 4.1.*b*).

These observations led Darwin to propose that species change over time through **natural selection**. Natural selection is the main mechanism by which evolution occurs. Natural selection occurs when organisms with beneficial inherited traits survive and reproduce more often than others, causing populations to change over time (evolve).




Figure 4.1.*b*: Darwin observed that beak shape varies among finch species. He postulated that the beak of an ancestral species had adapted over time to equip the finches to acquire different food sources. This illustration shows the beak shapes for four species of ground finch: 1. Geospiza magnirostris (the large ground finch), 2. G. fortis (the medium ground finch), 3. G. parvula (the small tree finch), and 4. Certhidea olivacea (the green-warbler finch).

How Natural Selection Works

Natural selection occurs when the following conditions are present:

- **Variation** exists – Individuals within a population possess different inherited traits.
- **Inheritance** occurs – Some of these trait differences can be passed from parents to offspring.
- **Biotic Potential** exists – All populations have the ability to grow rapidly because organisms produce more offspring than can survive.
- **Environmental Limiting Factors** occur – Environmental conditions such as limited food, water, shelter, space, predators, disease, and extreme weather prevent populations from growing indefinitely.
- **Selection** occurs – Environmental conditions favor some inherited traits over others. Traits that improve survival and reproduction are considered advantageous in that environment.
- **Differential Reproductive Success** occurs – Individuals with advantageous inherited traits are more likely to survive and reproduce. As a result, they tend to leave more offspring than individuals with less advantageous traits.
- **Time** – Over multiple generations, advantageous traits become more common within the population.

Variation and Adaptation

Natural selection depends on variation within a population. **Variation** refers to inherited genetic differences among individuals. For example, some birds may have larger beaks, while others have smaller beaks. These differences can affect how well individuals survive and reproduce in their environment.

An **adaptation** is an inherited trait that increases an organism's ability to survive and reproduce in a particular environment. Examples of adaptations include:

- Opposable thumbs that allow primates to grasp objects and use tools
- Thick fur in cold environments
- Deep roots in desert plants
- Camouflage in animals
- Salt tolerance in marsh plants

Adaptations arise over many generations through natural selection. Traits that increase survival and reproduction become more common in a population, while less favorable traits become less common. As these changes accumulate, the genetic makeup of the population changes over time. This process is known as evolution by natural selection.

The term adaptation can also describe the evolutionary process through which populations become better suited to their environment over generations. This process is sometimes called **adaptive evolution**.

Whether a trait is beneficial depends on the environment. For example, thick fur helps animals survive in cold climates but can be a disadvantage in hot desert environments. As environmental conditions change, different traits may be favored by natural selection.

Time (Generations)

A common misconception is that evolution always takes a very long time. While some evolutionary changes occur over thousands or millions of years, evolution can happen much more quickly in species with short generation times. For example, it is difficult to observe evolutionary change in elephants or humans because they reproduce slowly and have relatively few generations over a human lifetime. In contrast, organisms such as fruit flies, bacteria, and dandelions reproduce rapidly and can produce many generations in a short period of time. As a result, evolutionary change can sometimes be observed within years, months, or even days. Natural selection requires time because favorable inherited traits must be passed from one generation to the next. The more generations that occur, the greater the opportunity for populations to evolve.

Examples of rapid evolution include:

- Antibiotic resistance in bacteria
- Pesticide resistance in insects
- Herbicide resistance in weeds
- Changes in flowering times as climates warm

These examples show that evolution is an ongoing process that can occur on timescales relevant to environmental management and conservation.

4.1.1 Case Study: Galápagos Finches

One of the strongest demonstrations of natural selection comes from research conducted by Peter and Rosemary Grant on Galápagos finches. The Grants studied finch populations on Daphne Major, a small island in the Galápagos.

During a severe drought in 1977, many plants produced fewer seeds. Small, soft seeds became especially scarce. Finches with larger beaks were better able to crack the remaining hard seeds and were more likely to survive. After the drought, the average beak size in the population increased. Because beak size is inherited, the next generation also tended to have larger beaks (figure $4.1.c$).

This study provided direct evidence that environmental changes can produce natural selection and drive evolutionary change.The finch study shows that environmental conditions influence which traits are favored in a population. Similar processes occur today as species respond to climate change and habitat alteration.




Figure $4.1.c$**:** A drought on the Galápagos island of Daphne Major in 1977 reduced the number of small seeds available to finches, causing many of the small-beaked finches to die. This caused an increase in the finches' average beak size between 1976 and 1978.

Remember: VISTA

Use the acronym **VISTA** to remember the key components of natural selection:

V = Variation

I = Inheritance

S = Selection

T = Time (measured in generations)

A = Adaptation

Why Genetic Diversity Matters

Genetic diversity increases the variety of inherited traits within a population. Populations with high genetic diversity are often better able to adapt to changing environmental conditions. Populations with low genetic diversity may be more vulnerable to disease, habitat loss, climate change, and other environmental challenges.

Maintaining genetic diversity is an important conservation goal because it can increase a population's ability to adapt to future environmental change.

Patterns of Evolution

Divergent Evolution

Divergent evolution occurs when populations that share a common ancestor become increasingly different over time (figure $4.1.d$).

The Galápagos finches provide a classic example. A single ancestral finch population gave rise to several species with different beak shapes and feeding strategies.

As populations adapt to different environments, natural selection may favor different traits, causing populations to become more distinct over time.




Figure $4.1.d$**:** Flowering plants evolved from a common ancestor. Notice that the (a) dense blazing star and (b) purple coneflower vary in appearance, yet both share a similar basic morphology. (credit a, b: modification of work by Cory Zanker)

Convergent Evolution

Convergent evolution occurs when unrelated organisms independently evolve similar traits because they experience similar environmental pressures. These similarities are called **analogous traits** or **analogous structures** because they perform similar functions but evolved independently.

Examples include:

- Sharks and dolphins both have streamlined bodies for swimming.
- Bats and insects both evolved wings for flight.
- Cacti in North America and euphorbs in Africa independently evolved similar water-storing forms and modified leaves

(spines) in desert environments.

Although these organisms are not closely related, similar environmental conditions favored similar adaptations.

Environmental Science Connection

Environmental conditions can shape the evolution of species in predictable ways. Divergent evolution increases biodiversity by producing new forms and species. Convergent evolution demonstrates how similar environmental challenges can lead to similar adaptations, even in unrelated organisms.

Evolution in Changing Environments

Evolution continues to occur today.

Climate Change

Many species are changing their behavior and geographic ranges in response to increasing temperatures.

Examples include:

- Earlier flowering in plants
- Changes in migration timing
- Movement toward cooler habitats

Antibiotic Resistance

Bacteria reproduce quickly and often contain variation in resistance to antibiotics.

When antibiotics are used, antibiotic-resistant bacteria are more likely to survive and reproduce. Over time, resistant populations become more common.

Pesticide Resistance

Insects exposed to pesticides may contain individuals that naturally survive treatment. These survivors reproduce and pass resistance traits to offspring.

As a result, pesticide resistance can spread through populations within a relatively short period.

Conservation Biology

Small populations often have less genetic diversity. Maintaining biodiversity helps populations adapt to future environmental challenges such as climate change, habitat fragmentation, emerging diseases, and pollution. Understanding evolution helps scientists develop conservation strategies that protect biodiversity and improve ecosystem resilience.

Case Study: The Peppered Moth

Patterns of Natural Selection

Natural selection can affect populations in different ways.

Directional Selection

Natural selection can affect populations in different ways. One common pattern is **directional selection**.

Directional selection occurs when environmental conditions favor one extreme form of a trait.

During the Industrial Revolution in England, soot from factories darkened many tree trunks. Before pollution increased, most peppered moths were light-colored and blended into the tree bark. As trees became darker, dark-colored moths were better camouflaged and were less likely to be eaten by birds. As a result, dark-colored moths survived and reproduced more often than light-colored moths. Over time, dark-colored moths became more common in the population.

Later, air pollution controls reduced soot deposits on trees. As tree trunks became lighter again, light-colored moths gained a survival advantage and increased in frequency (figure $4.1.e$).

The peppered moth provides evidence that environmental change can drive natural selection and cause populations to evolve.

Directional selection can occur whenever environmental conditions change and favor one trait over another.

Examples include:

- Antibiotic resistance in bacteria
- Pesticide resistance in insects
- Herbicide resistance in weeds
- Climate-related shifts in plant flowering times

Environmental Science Connection

Human activities can alter environmental conditions and influence the traits favored by natural selection. Understanding evolution helps scientists predict how populations may respond to environmental challenges such as pollution, habitat loss, invasive species, and climate change.




Figure $4.1.e$**:** As the Industrial Revolution caused trees to darken from soot, darker colored peppered moths were better camouflaged than the lighter colored ones, which caused there to be more of the darker colored moths in the population.

Summary

Evolution by natural selection occurs when five conditions are present: variation, inheritance, selection, time, and adaptation (VISTA).

Individuals within a population differ in their inherited traits. Environmental conditions favor some traits over others, causing individuals with advantageous traits to survive and reproduce more successfully. Over many generations, these traits become

more common in the population.

Natural selection helps explain how populations respond to environmental challenges such as drought, habitat loss, pollution, invasive species, and climate change.

Understanding evolution allows environmental scientists to explain biodiversity, predict how populations may respond to environmental change, and develop effective conservation strategies.

Attributions

Modified by Erin Rempala.

- Samantha Fowler (Clayton State University), Rebecca Roush (Sandhills Community College), James Wise (Hampton University). Original content by OpenStax (CC BY 4.0; Access for free at https://cnx.org/contents/b3c1e1d2-83...4-e119a8aafbdd).

This page titled 4.1: Evolution and Natural Selection is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by OpenStax.

- 11.1: Discovering How Populations Change by OpenStax is licensed CC BY 4.0.

4.2: Other Mechanisms of Evolution

Other Sources of Evolutionary Change

Natural selection is the primary mechanism discussed in this chapter, but populations can also change through other evolutionary processes.

Mutation

A **mutation** is a change in DNA. Mutations create new genetic variation within populations and are the ultimate source of new inherited traits.

Most mutations have little or no effect on an organism. Some are harmful, while others may be beneficial under certain environmental conditions.

Without mutation, new genetic variation would not arise, and natural selection would have less variation to act upon.

Genetic Drift

Genetic drift is a random change in the genetic makeup of a population caused by chance events (figure $4.2.a$. Genetic drift has the greatest effect in small populations. For example, a wildfire, flood, or disease outbreak may randomly eliminate individuals regardless of their traits. As a result, some genetic variation may be lost simply by chance.

Small populations often have lower genetic diversity and may be more vulnerable to environmental change. This is one reason conservation biologists are concerned about endangered species with small populations.




Figure $4.2.a$**:** A chance event or catastrophe can reduce the genetic variability within a population.

Gene Flow

Gene flow occurs when individuals move into or out of a population and bring their genes with them (figure $4.2.b$). For example, seeds may be dispersed by wind, animals may migrate between habitats, or pollen may be carried long distances. Gene flow can increase genetic diversity and introduce new traits into a population.




Figure 4.2.*b*: Gene flow can occur when an individual travels from one geographic location to another and joins a different population of the species. In the example shown here, the brown allele is introduced into the green population.

Summary

Natural selection is not the only process that causes populations to change over time. Mutation introduces new genetic variation, genetic drift changes populations through chance events, and gene flow moves genetic variation between populations.

Together with natural selection, these processes influence how populations respond to environmental change.

Attributions

Modified by Erin Rempala.

- Samantha Fowler (Clayton State University), Rebecca Roush (Sandhills Community College), James Wise (Hampton University). Original content by OpenStax (CC BY 4.0; Access for free at https://cnx.org/contents/b3c1e1d2-83...4-e119a8aafbdd).

This page titled 4.2: Other Mechanisms of Evolution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by OpenStax.

- 11.2: Mechanisms of Evolution by OpenStax is licensed CC BY 4.0.

4.3: Evidence for Evolution

How do we know that evolution has occurred?

The evidence for evolution is compelling and extensive. Looking at every level of organization in living systems, biologists see the signature of past and present evolution. Darwin dedicated a large portion of his book, *On the Origin of Species*, identifying patterns in nature that were consistent with evolution and since Darwin our understanding has become clearer and broader.

Evidence from Fossils

Fossils provide evidence that organisms living today are different from those that lived in the past. Scientists use fossils to reconstruct Earth's biological history and observe how species have changed over time. Some fossils show intermediate forms between ancestral and modern organisms.

For example, the fossil record of horses documents gradual changes in body size, teeth, and leg structure as environments shifted from forests to grasslands (figure $4.3.a$). Fossils provide strong evidence that populations change over time and adapt to changing environmental conditions.




Figure $4.3.a$**:** This illustration shows an artist's renderings of these species derived from fossils of the evolutionary history of the horse and its ancestors. The species depicted are only four from a very diverse lineage that contains many branches, dead ends, and adaptive radiations. One of the trends, depicted here is the evolutionary tracking of a drying climate and increase in prairie versus forest habitat reflected in forms that are more adapted to grazing and predator escape through running. Przewalski's horse is one of a few living species of horse.

Evidence from Comparative Anatomy

Many organisms share similar anatomical structures because they inherited them from a common ancestor. For example, the bones in the forelimbs of humans, whales, bats, and dogs have the same basic arrangement even though they perform different functions (Figure $4.3.b$).

These shared structures are called **homologous structures** and provide evidence of common ancestry. Some organisms also possess **vestigial structures**, which are inherited features that have lost most or all of their original function.

Examples include:

- Pelvic bones in whales
- Small leg bones in some snakes
- Wings of flightless birds

These structures suggest that modern species evolved from ancestral species with different characteristics.




Figure 4.3.2**:** The similar construction of these appendages indicates that these organisms share a common ancestor.

Evidence from Biogeography

The geographic distribution of species (**biogeography**) provides evidence for evolution. Island species are often closely related to species on nearby continents but have evolved unique adaptations after becoming isolated (figure 4.3.*c*).

Examples include:

- Galápagos finches
- Hawaiian honeycreepers
- Australian marsupials
- Proteacea plant family

These patterns suggest that populations change over time as they adapt to different environments.




Figure 4.3.4**:** The Proteacea family of plants evolved before the supercontinent Gondwana broke up. Today, members of this plant family are found throughout the southern hemisphere (shown in red). (credit “Proteacea flower”: modification of work by “dorofofoto”/Flickr)

Evidence from DNA

All living organisms use DNA to store genetic information. When scientists compare DNA sequences, closely related species have more similar DNA than distantly related species. These patterns support the idea that modern species share common ancestors and have diversified through evolution.

Attributions

Modified by Erin Rempala.

- Samantha Fowler (Clayton State University), Rebecca Roush (Sandhills Community College), James Wise (Hampton University). Original content by OpenStax (CC BY 4.0; Access for free at https://cnx.org/contents/b3c1e1d2-83...4-e119a8aafbdd).

This page titled 4.3: Evidence for Evolution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by OpenStax.

- 11.3: Evidence of Evolution by OpenStax is licensed CC BY 4.0.

5: BIODIVERSITY

Biodiversity

Biodiversity refers to the variety of life on Earth. Scientists study biodiversity at three levels:

- **Genetic diversity** – variation within a species
- **Species diversity** – the number and abundance of species in an area
- **Ecosystem diversity** – the variety of habitats, ecosystems, and ecological processes

Biodiversity supports healthy ecosystems and provides benefits that people depend on every day (**ecosystem goods and services**). These benefits include food production, pollination, clean water, climate regulation, medicines, recreation, and cultural values.

High biodiversity often increases ecosystem resilience, helping ecosystems resist and recover from disturbances such as drought, wildfire, disease, and climate change.

Why This Matters

Biodiversity supports many essential **ecosystem services**, including food production, pollination, water purification, climate regulation, nutrient cycling, and cultural and recreational opportunities. Because species are connected through complex ecological relationships, the loss of biodiversity can disrupt ecosystem processes and reduce the ability of ecosystems to provide these benefits over time.

As biodiversity declines, ecosystems may become less resilient and less able to provide these services.



Figure 5.*a*: Volunteers carrying a sea turtle. Image by Aleson Padiha Pexels (Public domain)

5.1: The Value of Biodiversity

5.2: The Importance of Biodiversity to Human Life

5.3: Levels of Biodiversity

5.4: The Biodiversity Crisis
5.5: Threats to Biodiversity
5.6: Extinction
5.7: Measuring Biodiversity Loss
5.8: Preserving Biodiversity
5.9: Policies Protecting Biodiversity
5.10: Data Dive- Threats to Biodiversity

Attributions

Modified by Erin Rempala.

Rachel Schleiger (CC-BY-NC(opens in new window))

Thumbnail image - "Environmental protection"(opens in new window) is in the Public Domain

This page titled 5: Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- Unit 3: Conservation by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

5.1: The Value of Biodiversity

Why Protect Biodiversity?

Many products and services that support human well-being originate from biodiversity.

For example, many medicines were originally discovered in plants, animals, fungi, and microorganisms. Scientists continue to study species for compounds that may help treat disease.

A recent example comes from **yerba santa** (*Eriodictyon californicum*), a plant native to California (Figure $5.1.a$). Indigenous peoples of California have used yerba santa for generations because of its medicinal properties. Scientists discovered a compound in yerba santa that helps protect nerve cells in laboratory studies and may help researchers develop future treatments for Alzheimer's disease

This example highlights an important principle of conservation biology: biodiversity contains biological resources and scientific knowledge that may benefit society in ways we do not yet fully understand. Once a species becomes extinct, those potential benefits may be lost forever.



Figure $5.1.a$ Yerba santa (*Eriodictyon californicum*) in bloom. Image by Breck22 (Public domain)

Environmental Science Connection

Protecting biodiversity helps maintain ecosystem services, supports human health, and increases the ability of ecosystems to respond to environmental change.

Attribution

Modified by Erin Rempala.

Modified by Rachel Schleiger and Melissa Ha from Importance of Biodiversity from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 5.1: The Value of Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 9: The Value of Biodiversity by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

5.2: The Importance of Biodiversity to Human Life

Why Biodiversity Matters

Biodiversity has value both because it benefits people and because many believe species and ecosystems deserve protection simply because they exist. Humans depend on healthy ecosystems for food, clean water, medicine, climate regulation, and many other ecosystem services. Although extinction is a natural process, human activities are causing species to disappear much faster than normal. As biodiversity declines, ecosystems may become less resilient and less able to support human well-being.

Human Health

Many medicines come from plants, animals, fungi, and microorganisms. Scientists continue to discover new compounds that may help treat diseases. Protecting biodiversity preserves opportunities for future medical discoveries (figure $5.2.a$).



Figure

:

Catharanthus roseus, the Madagascar periwinkle, has various medicinal properties. Among other uses, it is a source of vincristine, a drug used in the treatment of lymphomas. (credit: Forest and Kim Starr)

Food Security

Biodiversity helps maintain a stable food supply. Genetic diversity in crops allows breeders to develop varieties that resist diseases, pests, and changing environmental conditions. A lack of diversity can make food systems more vulnerable to crop failures. To help protect crop diversity, seeds from around the world are stored in facilities such as the Svalbard Global Seed Vault in Norway (figure $5.2.b$).



Figure 5.2.b: The Svalbard Global Seed Vault on Spitsbergen island in Norway. (credit: Mari Tefre, Svalbard Global Seed Vault)

Ecosystem Goods and Services

Healthy ecosystems provide both **ecosystem goods** (products) and **ecosystem services** (benefits provided by natural processes) that people depend on every day.

Ecosystem goods include:

- Food crops
- Fish and seafood
- Lumber and paper products
- Medicines
- Fresh water

Ecosystem services include:

- Pollination of crops
- Soil formation and nutrient cycling
- Water purification
- Climate regulation
- Natural pest control

For example, many fruits, vegetables, and nuts depend on pollinators such as bees and other insects. Healthy soils support crop growth by recycling nutrients, while wetlands help filter pollutants from water supplies.

Wild Food Sources

Many people depend on wild fish and other organisms as important sources of food and materials. However, **overharvesting** (taking too much from nature) can reduce populations and threaten biodiversity. Examples include overfishing, hunting wildlife for bushmeat, excessive logging of forests, and collecting plants faster than they can regrow. When resources are removed faster than they can be replaced, populations may decline or even disappear.

Human Well-Being

Natural areas provide recreational, cultural, educational, and psychological benefits. Spending time in nature can reduce stress, improve mental health, increase physical activity, and strengthen people's connection to the natural world. Parks, forests, beaches, wetlands, and other natural areas also provide opportunities for outdoor recreation, scientific research, environmental education, and cultural traditions.

Summary

Protecting biodiversity helps maintain ecosystem services, supports human health and food security, and increases the resilience of ecosystems to environmental change.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Openstax and is licensed CC BY-NC-SA 4.0.

This page titled 5.2: The Importance of Biodiversity to Human Life is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

5.3: Levels of Biodiversity

Levels of Biodiversity

Scientists often describe biodiversity at three levels:

- Genetic Diversity
- Species Diversity
- Ecosystem Diversity

Genetic Diversity

Genetic diversity is the variation in inherited traits within a species. High genetic diversity increases a population's ability to adapt to environmental change, resist disease, and survive other challenges. In contrast, populations with low genetic diversity are often more vulnerable to extinction.

Why Genetic Diversity Matters

Genetic diversity provides the raw material for evolution and adaptation. For example, crop breeders rely on genetic diversity to develop plant varieties that can resist pests, diseases, and changing climate conditions.

The Tasmanian devil illustrates the importance of genetic diversity. Because Tasmanian devils have very low genetic diversity, they are especially vulnerable to a contagious cancer called Devil Facial Tumor Disease, which has caused major population declines (figure $5.3.a$).

Conserving genetic diversity helps maintain biodiversity and increases the resilience of populations in a changing environment.



Figure $5.3.a$: The Tasmanian devil in a defensive stance, at Tasmanian Devil Conservation Park, Tasman Peninsula. Image by Wayne McLean (CC-BY-SA).

Species Diversity

Species diversity refers to the variety of species in a particular area and the relative abundance of each species. Ecosystems such as tropical rainforests and coral reefs often have very high species diversity.

Species diversity includes two components:

- **Species richness** – the number of different species in an area.
- **Species evenness** – how evenly individuals are distributed among those species.

For example, two communities may each contain three species (the same species richness). However, if one community is dominated by a single species while the other has similar numbers of individuals in each species, the second community has greater species evenness and higher overall species diversity (figure $5.3.b$).




Figure $5.3.b$: Two hypothetical tree communities have the same species richness, but community #2 (bottom) has a greater species evenness. Both communities have nine trees and three tree species. In community #1, one species is dominant, represented by six individuals. There are two individuals of a conifer species, and only one individual of the final species. In community #2, there are three individuals from each species. Images compiled by Melissa Ha from Alone tree George Hodan, Old Tree Silhouette, and Tree (all public domain).

Why Species Diversity Matters

Species perform different roles within ecosystems, helping to maintain ecosystem function and stability. Ecosystems with high species diversity often have more ecological interactions and are generally more resilient to environmental change and disturbance. Conserving species diversity also helps maintain important ecosystem services such as food production, pollination, water purification, and climate regulation.

How Many Species Live on Earth?

Scientists have identified and described about 1.5 million species, but many more remain unknown. Current estimates suggest that Earth may contain between 8 and 11 million species, though the exact number is uncertain.

Many undiscovered species live in biodiversity-rich habitats such as tropical rainforests, coral reefs, and deep-ocean ecosystems. Because habitat loss and other human activities are causing rapid biodiversity decline, some species may become extinct before they are even discovered.

Scientists cannot protect species they do not know exist. Identifying and studying Earth's biodiversity helps researchers understand ecosystems, discover new medicines, improve conservation efforts, and protect ecosystem services that support human well-being. The fact that millions of species may still be undiscovered highlights how much remains unknown about Earth's biodiversity and why conservation is important.

Ecosystem Diversity

Ecosystem diversity is the variety of ecosystems within a region or across Earth. Examples include forests, wetlands, grasslands, deserts, estuaries, chaparral, and coral reefs (figure $5.3.c$). Different ecosystems, such as coral reefs and prairies, support different communities of organisms and contribute to Earth's biodiversity.

Each ecosystem supports different species and ecological processes. When ecosystems are degraded or destroyed, biodiversity declines and important ecosystem functions may be lost. For example, North American prairies once covered much of the central United States and Canada. Today, most have been converted to agriculture and urban development, reducing biodiversity and many of the ecological processes that once maintained productive soils.



Figure 5.3.c: The variety of ecosystems on Earth—from the coral reef (top) to prairie (bottom)—enables a great diversity of species to exist. Top image by Wise Hok Wai Lum (CC-BY-SA). Bottom image modified from Jim Minnerath, USFWS.

Ecosystem Goods and Services

Healthy ecosystems provide both **ecosystem goods** and **ecosystem services** that support human well-being.

Ecosystem goods are products obtained from nature, including:

- Food
- Fish and seafood
- Timber and paper products
- Fresh water
- Medicines

Ecosystem services are benefits provided by natural processes, including:

- Pollination of crops
- Soil formation and nutrient cycling
- Water purification

- Climate regulation
- Flood control
- Natural pest control

These goods and services support food production, economic activity, human health, and environmental sustainability.

Ecosystem Services and the Economy

Many ecosystem services have tremendous value, but they are often not included in traditional economic accounting. As a result, activities that damage ecosystems may appear profitable even when they cause the loss of valuable services such as pollination, water purification, flood protection, and climate regulation.

Economists describe this as a **market failure**, where the environmental costs and benefits of human activities are not fully reflected in prices. If the value of ecosystem services were included in economic decisions, conserving ecosystems would often provide greater long-term benefits than destroying them for short-term profits (figure $5.3.c$).




Figure $5.3.c$: Ecosystem services such as climate regulation, fisheries, and coral reefs provide enormous economic benefits to society, worth billions to trillions of dollars each year. Data from The Economics of Ecosystems and Biodiversity Synthesis Report. Image by Future Challenges (CC-BY-SA).

Why It Matters

Food production depends on many ecosystem services. Healthy soils recycle nutrients, pollinators help crops reproduce, and natural predators help control agricultural pests. In California, native pollinators can provide important pollination services when natural habitat is maintained near farms.

Loss of biodiversity can reduce these ecosystem services, making food production more difficult and expensive.

Protecting biodiversity helps maintain the ecosystem goods and services that support human societies, including food security, clean water, climate regulation, and long-term sustainability.

Why All Three Types of Biodiversity Matter

Conservation efforts seek to protect genetic, species, and ecosystem diversity because each contributes to the resilience and long-term sustainability of natural systems.

For example:

- Genetic diversity helps populations adapt to a changing environment.
- Species diversity supports ecosystem function.
- Ecosystem diversity provides a variety of habitats.

Together, these three levels of biodiversity help maintain healthy ecosystems and support human well-being.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from:

- 9.1: Ecosystem Diversity by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0(opens in new window).
- **9.2: Species Diversity**(opens in new window) by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0(opens in new window).
- 9.3: Genetic Diversity by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0(opens in new window).

Modified by Melissa Ha from the following sources:

- Biological from *AP Environmental Science* by University of California College Prep, University of California (licensed under CC-BY(opens in new window)). Download for free at CNX.
- Importance of Biodiversity from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 5.3: Levels of Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 9.3: Genetic Diversity by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

5.4: The Biodiversity Crisis

Biodiversity Is Unevenly Distributed

Biodiversity is not evenly distributed across Earth (Fig a). In general, biodiversity is highest near the equator and decreases toward the poles. Tropical regions such as rainforests contain large numbers of species and many endemic species, which are species found nowhere else on Earth. These areas often become priorities for conservation because they contain unique forms of life.

Biodiversity Hotspots

Some regions contain exceptionally high biodiversity and are under significant threat from human activities. These regions are called **biodiversity hotspots**. To qualify as a hotspot, an area must contain at least 1,500 **endemic** plant species (species found nowhere else on Earth) and have 30% or less of its original natural habitat remaining.

Biodiversity hotspots are a high priority for conservation because protecting a relatively small area can help preserve a large number of unique species. Many hotspots contain plants and animals that exist nowhere else on the planet.

Conservation International has identified 36 biodiversity hotspots worldwide (figure $5.4.a$). Together, these areas cover only about 2.5% of Earth's land surface, yet they contain more than half of the world's endemic plant species and nearly 43% of endemic birds, mammals, reptiles, and amphibians.

Despite covering a small fraction of Earth's land area, biodiversity hotspots provide an estimated 35% of the ecosystem services that support human well-being, including food production, water purification, pollination, climate regulation, and other benefits that people depend on every day.




Figure $5.4.a$ Map of the world's biodiversity hot spots, all of which are heavily threatened by habitat loss and degradation. By Conservation International - Own work, CC BY-SA 4.0.

Biodiversity Loss

Although extinction is a natural process, human activities have greatly increased extinction rates.
The major causes of biodiversity loss are:

1. Habitat destruction and fragmentation
2. Invasive species
3. Overharvesting of natural resources
4. Pollution
5. Climate change

Together, these pressures are causing species to disappear at rates far above natural background levels of extinction.

Case Study: Lake Victoria Cichlids

Lake Victoria in East Africa once contained hundreds of species of cichlid fish, each adapted to a different habitat or food source (figure 5.4.*b*). Many of these species declined rapidly after the introduction of the Nile perch, a large predatory fish. Habitat degradation and increased fishing pressure also contributed to biodiversity loss. Scientists estimate that many species disappeared before they were even fully studied. This example illustrates how human activities can rapidly reduce biodiversity.




Figure 5.4.*b*: Cichlids of the African rift lake system. By original figure: Joana I. Meier, David A. Marques, Salome Mwaiko, Catherine E. W Laurent Excoffier & Ole SeehausenEditing by Dennis Pietras, Buffalo, NY USA - originally figure 1 of "Ancient hybridization fuels rapid c fish adaptive radiations" doi:10.1038/ncomms14363 http://www.nature.com/articles/ncomms14363Minor editing and conversion to jp Dennis Pietras, CC BY-SA 4.0, https://commons.wikimedia.org/w/inde...curid=56290753

Conservation of Biodiversity

Conservation biologists work to protect biodiversity by:

- Protecting habitats
- Restoring ecosystems
- Managing threatened species
- Creating protected areas
- Supporting sustainable resource use

Conservation efforts help preserve biodiversity and maintain the ecosystem services that support human societies.

Summary

Biodiversity refers to the variety of life on Earth and includes genetic, species, and ecosystem diversity. Biodiversity provides important ecosystem services and supports human well-being. However, biodiversity is declining because of habitat destruction, invasive species, pollution, overharvesting, and climate change. Protecting biodiversity helps maintain ecosystem resilience, supports conservation goals, and preserves the ecosystem goods and services that humans need for survival.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from Openstax and is licensed CC BY-NC-SA 4.0.

This page titled 5.4: The Biodiversity Crisis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

5.5: Threats to Biodiversity

Major Threats to Biodiversity

Human activities are causing biodiversity to decline much faster than the natural background rate of extinction. As biodiversity decreases, ecosystems may become less resilient and less able to provide the goods and services that humans depend on.

The major threats to biodiversity are summarized in the acronym HIPPCO:

- Habitat loss and fragmentation
- Invasive species
- Pollution
- (Human) Population Growth
- Climate Change
- Overharvesting

Habitat Loss

Habitat loss is the leading cause of biodiversity decline. When forests, wetlands, grasslands, rivers, and other habitats are destroyed or fragmented, many species lose the resources they need to survive. Human activities such as agriculture, urban development, logging, mining, and road construction can reduce or eliminate habitat.

Habitat loss is the leading cause of biodiversity decline. When forests, wetlands, grasslands, rivers, and other habitats are destroyed or fragmented, many species lose the resources they need to survive. Human activities such as agriculture, urban development, logging, mining, and road construction can reduce or eliminate habitat. In Southern California, more than 90% of historic coastal wetlands have been lost to development. Remaining wetlands, such as the Tijuana River National Estuarine Research Reserve and the Kendall-Frost Marsh Reserve, provide critical habitat for fish, birds, and other wildlife and illustrate the importance of habitat conservation.



Figure $5.5.a$:] Tijuana Estuary. By Osbomb - [www.flickr.com/photos/osbornb/267928456/], CC BY 2.0, [commons.wikimedia.org/w/index.php?curid=15357086]

Consumer Choices and Habitat Conservation

Many everyday products, such as wood flooring, furniture, and building materials, can contribute to habitat loss if the wood comes from unsustainable or illegal logging. Illegal harvesting of tropical timber destroys forests, threatens biodiversity, and increases the risk of species extinction. Consumers can help reduce these impacts by choosing wood products certified by the **Forest Stewardship Council (FSC)** or by purchasing sustainably harvested domestic wood. Making informed purchasing decisions can help protect forests and the species that depend on them.



Figure 5.5.*b*: Forest Stewardship Council logo. By The logo may be obtained from Forest Stewardship Council., Fair use, [//en.wikipedia.org/w/index.php?curid=34259396]

Invasive Species

An **invasive species** is a non-native organism that spreads rapidly and harms native species or ecosystems. Invasive species may compete with native organisms for resources, prey on native species, introduce diseases, or alter habitats. For example, the accidental introduction of the brown tree snake (*Boiga irregularis*) to Guam after World War II caused severe declines in native wildlife (figure 5.5.*b*). The snake preyed on birds, lizards, and other small animals, leading to the local extinction of many native species and dramatically altering the island's ecosystem. Today, the brown tree snake remains one of the world's most damaging invasive species.



Figure 5.5.*b*: The brown tree snake, Boiga irregularis, is an exotic species that has caused numerous extinctions on the island of Guam since its accidental introduction in 1950. (credit: NPS)

Pollution

Pollution can harm ecosystems and reduce biodiversity (figure 5.5.*c*). Common examples include:

- Plastic pollution in oceans
- Pesticides that harm pollinators
- Excess nutrients that create algal blooms
- Toxic chemicals that accumulate in food webs

Pollution can directly kill organisms, reduce reproduction and survival, or degrade habitat quality. For example, excess fertilizer runoff can trigger algal blooms that remove oxygen from the water, making it difficult for fish and other aquatic organisms to survive. Plastic pollution can injure or kill wildlife through entanglement or ingestion, while pesticides can unintentionally harm beneficial insects such as bees and butterflies. Because pollution affects both individual species and entire ecosystems, it is a major contributor to biodiversity loss worldwide.

In Southern California, urban runoff can carry fertilizers, pesticides, oil, trash, and other pollutants into rivers, wetlands, and coastal ecosystems, affecting biodiversity and ecosystem health.




Figure $5.5.c$: Some sources of water and air pollution and their major effects on biodiversity. Sources of pollution include industry, car emissions, power plants, pesticides, fertilizers, and drugs. Some major consequences of air pollution include respiratory problems, acid precipitation, plant damage, ozone depletion, and climate change. Water pollution can bioaccumulate and can lead to eutrophication, death by plastic, reduced photosynthesis, and hormone disruption. Image by Leafcutter Media (CC-BY-SA).

Human Population Growth

Human population growth and increasing resource consumption drive many of the threats to biodiversity. As the human population grows, more land, water, energy, food, and natural resources are needed to support human activities. Forests may be cleared for agriculture and housing, more resources are harvested from ecosystems, pollutants are released into the environment, and greenhouse gas emissions increase. As a result, habitat loss, overharvesting, pollution, invasive species, and climate change are all closely linked to the growing demand for resources. Sustainable resource use and conservation efforts can help reduce these pressures while supporting both human well-being and biodiversity.

Climate Change

Human-caused (**anthropogenic**) climate change is altering temperatures, precipitation patterns, sea levels, and ecosystem conditions worldwide. As environmental conditions change rapidly, some species may be unable to adapt or move to suitable habitat quickly enough (figure $5.5.d$). Climate change can also increase the effects of other threats such as habitat loss, invasive species, and disease.



Figure 5.5.d: Climate change is causing some species to shift their ranges. In parts of the Arctic, grizzly bears and polar bears now overlap in areas where they were once separated, leading to rare hybrid offspring called "grolar bears."By CNX OpenStax - By The logo may be obtained from Forest Stewardship Council., Fair use,[cnx.org], CC BY 4.0, [wikimedia.org]

Overharvesting

Overharvesting occurs when organisms are removed from the environment faster than populations can recover.

Examples include:

- Overfishing of commercial fish species
- Hunting wildlife for bushmeat
- Excessive logging of forests]
- Collecting medicinal plants faster than they can regrow

Overharvesting can reduce populations, disrupt ecosystems, and increase the risk of extinction.

Summary

Most biodiversity loss today is caused by human activities. The major threats can be summarized using the acronym **HIPPCO**: habitat loss, invasive species, population growth, pollution, climate change, and overharvesting. Understanding these threats helps scientists, policymakers, and communities develop effective conservation strategies. By protecting habitats, reducing overharvesting, controlling invasive species, reducing pollution, addressing climate change, and using resources sustainably, people can help conserve biodiversity and support healthy ecosystems.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Openstax and is licensed CC BY-NC-SA 4.0.

This page titled 5.5: Threats to Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

5.6: Extinction

Extinction and Biodiversity Loss

Biodiversity changes over time through two natural processes:

- **Speciation** – the formation of new species
- **Extinction** – the loss of a species

When new species form faster than species go extinct, biodiversity increases. When extinction occurs faster than speciation, biodiversity declines.

Throughout Earth's history, biodiversity has increased and decreased in response to environmental change. Scientists have identified five major **mass extinctions**, periods when a large percentage of Earth's species disappeared in a relatively short period of geological time.

The best-known mass extinction occurred about 66 million years ago when an asteroid impact contributed to the extinction of the non-avian dinosaurs and many other species.

(Figure $5.6.a$).




Figure $5.6.a$: The fossil record shows five major mass extinctions, times when biodiversity declined rapidly because many species went extinct.

The Sixth Mass Extinction

Many scientists believe Earth is currently experiencing a **sixth mass extinction**. Unlike previous mass extinctions that were caused by natural events, the current biodiversity crisis is primarily driven by human activities. This has led some scientists to describe the present as the **Anthropocene**, a time when human activities have become the dominant influence on Earth's natural systems

Major causes of modern extinction include:

- Habitat loss
- Invasive species
- Pollution
- Climate change

- Overharvesting

According to the 2019 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) Global Assessment, about one million plant and animal species are at risk of extinction within the coming decades if current trends continue. Human activities such as habitat loss, pollution, climate change, overexploitation of resources, and invasive species are the main drivers of this biodiversity crisis. Recent UN- and IPBES-supported assessments indicate that global biodiversity loss continues and that extinction risks have increased in many regions, highlighting the need for rapid conservation and sustainability actions.

Why Extinction Matters

Extinction is a natural process, but today's extinction rates are much higher than natural background levels. **Background extinction** is the normal rate of species loss that occurs naturally over time, even without major human impacts or catastrophic events.
The loss of species can reduce ecosystem resilience and weaken ecosystem services that people depend on, including:

- Food production
- Water purification
- Pollination
- Climate regulation

Extirpation

Extinction occurs when a species disappears from Earth entirely. **Extirpation** occurs when a species disappears from a particular area but still survives elsewhere. Although extirpation does not eliminate a species globally, it can still affect ecosystem function. Conservation efforts often focus on preventing local extirpations and restoring species to habitats where they once occurred.

Yellowstone Wolves: A Case Study

The effects of extirpation can be seen in Yellowstone National Park. Wolves were eliminated from the park during the early twentieth century but survived elsewhere in North America. After wolves were reintroduced in the 1990s, scientists observed changes in elk populations, vegetation, and other parts of the ecosystem. This example shows how the loss or return of a single species can influence an entire ecosystem. The Yellowstone wolf example is often used to illustrate a **top-down effect**, where predators influence organisms at lower trophic levels.

Ecologists generally describe ecosystem interactions as:

- **Top-down effects** – Predators influence herbivores and other organisms lower in the food web. For example, wolves can affect elk populations, which in turn can influence vegetation.
- **Bottom-up effects** – The availability of resources such as sunlight, water, nutrients, or food influences populations at higher trophic levels.

In most ecosystems, both top-down and bottom-up factors interact to shape biodiversity, community structure, and ecosystem function.

Environmental Science Connection

Understanding extinction, extirpation, and species interactions helps scientists identify conservation priorities and develop strategies to protect biodiversity in a rapidly changing world.

References

UN Report: Nature's Dangerous Decline 'Unprecedented'; Species Extinction Rates 'Accelerating'. 2019. United Nations. Accessed 2020-08-01.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- The Biodiversity Crisis from *General Biology* by OpenStax (licensed under CC-BY(opens in new window))
- Preserving Biodiversity and Importance of Biodiversity from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Biodiversity, Species Loss, and Ecosystem Function by Tom Theis and Jonathan Tomkin, Editors. Download for free at CNX. (licensed under CC-BY)

This page titled 5.6: Extinction is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 10.1: Extinction by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

5.7: Measuring Biodiversity Loss

The IUCN Red List

The **International Union for Conservation of Nature (IUCN)** maintains the **Red List of Threatened Species**, one of the world's most important tools for tracking biodiversity and extinction risk.

The Red List evaluates species and places them into categories based on their risk of extinction:

- **Least Concern** – low risk of extinction
- **Near Threatened** – may become threatened in the future
- **Vulnerable** – facing a high risk of extinction
- **Endangered** – facing a very high risk of extinction
- **Critically Endangered** – facing an extremely high risk of extinction
- **Extinct in the Wild** – survives only in captivity or managed populations
- **Extinct** – no living individuals remain

Scientists use the Red List to identify species at risk and guide conservation efforts.

Examples

- The **emperor penguin** is considered **Near Threatened** because of climate change and habitat loss (figure $5.7.a$.
- The **blue whale** is classified as **Endangered** (figure $5.7.b$.
- Many amphibians, corals, and mammals are considered threatened because of habitat loss, climate change, pollution, invasive species, and overharvesting.



Figure $5.7.a$: The Emperor Penguin is an example of near threatened species. Image by Christopher Michel (CC-BY).



Figure 5.7.c: The blue whale (*Balaenoptera musculus*) is categorized as endangered according to the IUCN Red List. Image by Mike Baird (CC-BY).

The Red List helps scientists and policymakers identify conservation priorities and monitor changes in biodiversity over time. Species that move into higher-risk categories may require additional protection to prevent extinction.

References

The IUCN Red List of Threatened Species. 2020. IUCN. Accessed 2020-07-29.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Biodiversity, Species Loss, and Ecosystem Function and The Industrialization of Nature: A Modern History (1500 to the present) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors. Download for free at CNX. (licensed under CC-BY)

This page titled 5.7: Measuring Biodiversity Loss is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 10.2: Measures of Biodiversity Loss by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

5.8: Preserving Biodiversity

Conservation Biology

Conservation biology is the scientific study of how to protect biodiversity, including genes, species, ecosystems, and the ecological processes that support life on Earth. Conservation biologists study the causes of biodiversity loss and develop solutions that help protect species, restore habitats, and maintain healthy ecosystems. Because biodiversity loss is connected to environmental, social, and economic issues, conservation biology is an interdisciplinary field.

Protecting Habitats

One of the most effective ways to conserve biodiversity is to protect habitats. National parks, wildlife refuges, marine protected areas, and nature reserves help preserve ecosystems and provide habitat for thousands of species.

Local examples include:

- Kendall-Frost Marsh Reserve
- Torrey Pines State Natural Reserve
- Cabrillo National Monument
- Anza-Borrego Desert State Park

Protecting habitats not only benefits wildlife but also helps maintain ecosystem services such as clean water, pollination, carbon storage, and recreation.

The South La Jolla State Marine Reserve (SMR) and South La Jolla State Marine Conservation Area (SMCA) are adjoining Marine Protected Areas (MPAs) that together protect more than 5 square miles of kelp forest and other marine habitats near La Jolla Cove (figure $5.8.a$). Fishing and the collection of marine organisms are prohibited in SMCA, allowing ecosystems to function with minimal human disturbance. These MPAs helps protect kelp forests, rocky reefs, fish populations, marine mammals, and other species while supporting scientific research, education, and recreation.

Interactive Element

Interactive web map of South La Jolla SMCA and South La Jolla SMR and other Marine Protected Areas along the coast of Southern California.

Restoring Ecosystems

In some cases, damaged ecosystems can be restored.

Ecosystem restoration may include:

- Removing invasive species
- Replanting native vegetation
- Restoring wetlands
- Reintroducing native wildlife

In some cases, damaged ecosystems can be restored by removing invasive species, improving habitat quality, and reestablishing natural ecological processes. In San Diego County, the **San Diego River Park Foundation** works to restore habitats along the San Diego River and its estuary near Ocean Beach. Restoration efforts include removing invasive plants, planting native vegetation, improving wildlife habitat, and increasing public access to natural areas. These projects help support biodiversity, improve water quality, and strengthen ecosystem resilience.

Sustainable Resource Use

Conservation does not mean eliminating human use of natural resources. Instead, sustainability seeks to meet human needs while maintaining healthy ecosystems for future generations.

Examples include:

- Sustainable agriculture
- Sustainable forestry
- Sustainable fisheries
- Water conservation
- Renewable energy

The Role of Individuals

Individuals can help protect biodiversity by:

- Conserving water and energy
- Reducing waste
- Supporting sustainable products
- Planting native species
- Participating in habitat restoration projects
- Supporting conservation organizations

Summary

Most threats to biodiversity are caused by human activities, but human actions can also help protect biodiversity. Conserving habitats, restoring ecosystems, using resources sustainably, and reducing environmental impacts are essential for maintaining biodiversity and supporting long-term sustainability. Protecting biodiversity helps preserve ecosystem goods and services, maintain ecosystem resilience, and support the natural systems that human societies depend upon.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Openstax and is licensed CC BY-NC-SA 4.0.

This page titled 5.8: Preserving Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

5.9: Policies Protecting Biodiversity

Laws and Policies that Protect Biodiversity

Conservation is supported by local, state, national, and international laws that help protect species and habitats.

Endangered Species Protection

In the United States, the **Endangered Species Act (ESA)** protects species that are at risk of extinction. Once a species is listed as threatened or endangered, government agencies develop plans to help protect and recover populations.

California also has its own **California Endangered Species Act (CESA)**, which protects threatened and endangered species within the state.

A local example of endangered species protection is the San Diego fairy shrimp (*Branchinecta sandiegonensis*), a federally endangered species that depends on seasonal wetlands called vernal pools found in parts of San Diego County (figure $5.9.a$.








Figure 5.9.a: (Top) San Diego fairy shrimp. By USFWS Pacific Southwest, Photo by Joanna Gilkeson/USFWS. - Flickr, CC BY 2.0, [commons.wikimedia.org/w/index.php?curid=61531108][. Middle image: A vernal pool complex between mima mounds in San Diego County. By Pacific Southwest Region USFWS from Sacramento, US - A network of vernal pools flow between mima mounds, Public Domain, [commons.wikimedia.org/w/index.php?curid=68988382]. Bottom Image: Public Affairs Specialist for the U.S. Fish and Wildlife Service, Joanna Gilkeson, takes photos of federally endangered San Diego fairy shrimp in a vernal pool in Otay Mesa, California. Vernal pools are shallow seasonal ponds, and vernal pool species are uniquely adapted to changes in water availability. Image by USFWS [Maideline Sanchez] (public domain).

Marine Protection

The **Marine Mammal Protection Act** protects whales, dolphins, seals, sea lions, and other marine mammals from hunting, harassment, and other harmful activities.

California also protects marine biodiversity through a network of **Marine Protected Areas (MPAs)**. For example, the **South La Jolla State Marine Reserve** protects marine habitats near La Jolla Cove and helps conserve kelp forests, fish populations, and other marine life.

International Conservation

Some species are threatened by international trade. The **Convention on International Trade in Endangered Species (CITES)** is an international agreement that regulates or prohibits trade in certain plants and animals to reduce the risk of extinction.

Summary

Laws and policies are important conservation tools, but they work best when combined with habitat protection, ecosystem restoration, sustainable resource use, and public support for conservation efforts.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Preserving Biodiversity from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Threatened and Endangered Species by California Department of Fish and Wildlife (public domain)
- Marine Mammal Protection by NOAA Fisheries (public domain)

This page titled 5.9: Policies Protecting Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 11.1: Policies by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

5.10: Data Dive- Threats to Biodiversity

Overview

Starting in 1998, the World Wildlife Federation (WWF) began publishing the Living Planet Report. This report analyzes the health of the plant relative to the impact humanity has on it, and is updated every two years. This report continually urges humanity, especially world leaders, to work together to build a more sustainable, resilient, and healthy future for people and nature. In the 2018 update, there was an overview of the major threats to biodiversity for several animal families of concern. See the graph of their results below:




Figure $5.10.a$: Percent of biodiversity threats to various animal groups. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Living Planet Report 2018 Aiming Higher.

Questions

1. What is the independent (explanatory) variable and the dependent (response) variable?
2. What animal category above is most affected by habitat destruction? What about exploitation?
3. What category above has the largest impact (on average) on all species?
4. What is different for the fish data patterns compared to the other animal categories? Why?
5. How can the results of this graph to inform future conservation efforts?
6. Based on what you have learned in the chapter, list two other categories of life that you think should be added to the graph. Provide at least one reason for each as to why you think it should be added.

Raw Data From Above Graph(s)

Table $5.10.a$**:** Raw data for percent of biodiversity threats to various animal groups. Graph by Rachel Schleiger (CC-BY-NC) modified from data in Living Planet Report 2018 Aiming Higher.

Raw data for percent of biodiversity threats to animal groups.

Animal Group	Habitat Degradation	Exploitation	Invasive Species And Disease	Pollution	Climate Change
Birds	49	18	10	11	12
Reptiles/Amphibians	47	23	12	11	7

Animal Group	Habitat Degradation	Exploitation	Invasive Species And Disease	Pollution	Climate Change
Mammals	45	38	9	5	3
Fish	28	55	5	4	8

Attribution

Rachel Schleiger (CC-BY-NC)

This page titled 5.10: Data Dive- Threats to Biodiversity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 10.8: Data Dive- Threats to Biodiversity by Melissa Ha and Rachel Schleiger has no license indicated.

6: HUMAN POPULATION GROWTH

Introduction

Why has our human species (*Homo sapiens sapiens*) been so successful? We have had the unique ability to not just occupy, but thrive, in diverse environments, even some of the most extreme. Through our social cooperation (despite large groups with no kin), language, art, and ingenuity (especially with tools and fire), our species became unstoppable. Modern humans have permanent settlements on all but one continent (Antarctica) and have exponentially increased our population to over 8 billion people today. We have impacted every part of the world. However, this transformation has consequences for all life on the planet (including our own species). If we are to continue to thrive we will need to change our current systems to more closely mimic nature and become more sustainable.




Figure $6.a$ Map demonstrating human spread across the globe. Image by Maulucioni(opens in new window) (licensed under CC-BY 3.0(opens in new window))

Human Population Growth

Concepts from population ecology can be applied to human populations. **Demography** is the study of how human populations grow, shrink, and change in terms of age structure, sex ratios, and geographic distribution. Demographers also compare population trends among different countries and regions. The human population has increased dramatically over the past few centuries and continues to grow (figure $6.b$).

Environmental scientists are interested not only in the number of people on Earth but also in how much energy, land, water, and other resources people use. Population growth can increase pressure on natural systems, but environmental impacts also depend on patterns of consumption and resource use. A population may experience overpopulation when its combined demand for resources and ecosystem services exceeds the environment's ability to sustainably provide those resources and absorb associated wastes. The consequences can include habitat loss, resource depletion, pollution, and declines in ecosystem health, all of which can affect human well-being.

Unlike other species, humans can modify environmental conditions through technology, agriculture, trade, and the use of external energy sources. As a result, human carrying capacity is not fixed and can change over time, making it difficult to determine a single carrying-capacity value for Earth. Instead of relying solely on carrying capacity, environmental scientists often use the concept of an **ecological footprint** to assess human impacts on the environment.

An ecological footprint estimates the amount of biologically productive land and water required to provide the resources a person or population consumes and to absorb the wastes it generates. Scientists compare ecological footprints with Earth's

available **biocapacity**, the capacity of ecosystems to regenerate resources and absorb wastes. When a population's ecological footprint exceeds available biocapacity, **ecological overshoot** occurs, indicating that resource use is not sustainable in the long term. Understanding both population dynamics and patterns of resource consumption is therefore essential for evaluating environmental sustainability.




Figure 6.b: All the people in the world (2018). Each country's size represents the size of its population in 2018. Each square is 500,000 people. Image: Max Roser for OurWorldinData.org (CC-BY)

Attributions

Modified by Rachel Schleiger and Melissa Ha from the following sources:

- Human Demography(opens in new window) from *Introduction to Environmental Science,* 2nd edition by Caralyn Zehnder et al. (licensed under CC-BY-NC-SA(opens in new window)).
- The Human Population from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

6.1: History of Human Population Growth

6.2: The Rate of Human Population Growth

6.3: Factors Affecting Population Growth

6.4: Looking Ahead

6.5: Data Dive- World Population Densities

This page titled 6: Human Population Growth is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 12: The Human Population by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

6.1: History of Human Population Growth

The human population is growing rapidly. For most of human history, there were fewer than 1 billion people on the planet. During the time of the Agricultural Revolution, 10,000 B.C., there were only 5-10 million people on Earth - which is basically the population of New York City today. In 1800, during the Industrial Revolution, there were approximately 1 billion people on Earth. Continued agricultural expansion and extraction of fossil fuels and minerals led to rapid global economic growth and, in turn, population growth in the 19th century. We've added over 6 billion people to the human population in just a little over 200 years (figure $6.1.a$). As of May 2026, the global human population is around 8.2 billion people.




Figure $6.1.a$: The size of human population over the past 10,000 years. Published online at OurWorldInData.org. (CC-BY(opens in new window))

Although global population size continues to increase, the rate of human population growth has decreased. This means that the population size is not increasing as quickly as it did in the past (figure $6.1.b$).




Figure 6.1.*b*: Human population growth. Since 1700, human population size (shaded in teal) has increased. It reached 1 billion in 1803, 2 billion in 1928, 2.5 billion in 1950, 5 billion in 1987, and 8 billion in 2022. It is projected to reach more than 10 billion in 2100. The population growth rate (pink line) was only 0.04% on average between 10,000 BCE and 1700. The population growth rate peaked in 1968 at 2.1%, and since then, it has slowed to 0.9% in 2023 and is projected to be -0.1% in 2100. Image by by Max Roser and Veronika Samborska OurWorldInData.org (CC-BY).

The fundamental cause of the acceleration of growth rate for humans in the past 200 years has been the reduced death rate due to changes in public health and sanitation. Clean drinking water and proper disposal sewage has drastically improved health in developed nations. Also, medical innovations such as the use of antibiotics and vaccines have decreased the ability of infectious disease to limit human population growth. In the past, diseases such as the bubonic plaque of the fourteenth century killed between 30 and 60 percent of Europe's population and reduced the overall world population by as many as one hundred million people. Naturally, infectious disease continues to have an impact on human population growth, especially in poorer nations. For example, life expectancy in sub-Saharan Africa, which was increasing from 1950 to 1990, began to decline after 1985 largely as a result of HIV/AIDS mortality. According to a 2016 study by Marcus et al., The reduction in life expectancy caused by HIV/AIDS was estimated to be 8 years for 2016.

Human technology and particularly our harnessing of the energy contained in fossil fuels have caused unprecedented changes to Earth's environment, altering ecosystems to the point where some may be in danger of collapse. Changes on a global scale including depletion of the ozone layer, desertification and topsoil loss, and global climate change are caused by human activities.

Reference

Marcus, J. L., Chao, C. R., Leyden, W. A., Xu, L., Quesenberry, C. P., Jr, Klein, D. B., Towner, W. J., Horberg, M. A., & Silverberg, M. J. (2016). Narrowing the Gap in Life Expectancy Between HIV-Infected and HIV-Uninfected Individuals With Access to Care(opens in new window). *Journal of acquired immune deficiency syndromes (1999)*, *73*(1), 39–46.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- The Human Population from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Human Demography(opens in new window) from *Introduction to Environmental Science,* 2nd edition by Caralyn Zehnder et al. (licensed under CC-BY-NC-SA(opens in new window)).
- Human Population Growth(opens in new window) from *General Biology* by OpenStax (licensed under CC-BY(opens in new

window))
- The Industrialization of Nature: A Modern History (1500 to the present)(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors. Download for free at CNX(opens in new window). (licensed under CC-BY(opens in new window))


This page titled 6.1: History of Human Population Growth is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 12.1: History of Human Population Growth by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

6.2: The Rate of Human Population Growth

Demographic Transition

Recall from the Populations chapter that the **population growth rate** (r) equals the birth rate minus the death rate. Slowly declining birth rates following an earlier sharp decline in death rates are today characteristic in many low-income countries. The shift observed from high birth and death rates to low birth as well as death rates in countries such as the US, Japan, and much of Europe is called the **demographic transition model.** Figure $6.2.a$






Figure $6.2.a$. Top: Demographic transition model. Image By Max Roser Our World In Data(opens in new window) (CC BY 4.0(opens in new window)). Bottom: Demographic Transition in 5 countries. Image by Max Roser Our World in Data(opens in new window) (CC BY SA(opens in new window)).

Prior to World War II, advances in public health were largely limited to high-income countries. But since then, many more countries have enjoyed improvements in public health - always with dramatic effect on death rates. For example, in 1945, the death rate in Sri Lanka (then called Ceylon) was 0.022 (2.2%). In 1946, a large-scale program to control mosquitos, which transmit malaria, was started. By eliminating the mosquito, the incidence of malaria dropped sharply. After 9 years, the death rate dropped to 0.010 (1%), and by 2012 was 0.006 (0.6%). However, a compensating decline in birth rates has come more slowly; the birth rate was 0.018 (1.8% in 2012). With birth rates higher than death rates, the population was increasing at an annual rate of 0.012 (1.2%) per year, with a doubling time of 57.5 years (t = 0.69/0.012).

While the Demographic Transition Model describes a general pattern of population change observed in high income countries, it does not fully account for the historical and structural challenges faced by many low-income countries. Factors such as colonial legacies, limited economic resources, political instability, and global inequality can alter or slow the transition, meaning that not all countries will follow the same pathway taken by the United States, Europe, or Japan.

Predicting Future Population Size

Doubling time is based on a constant population growth rate, but this rate can change depending on a number of factors, including total fertility rate (TFR) and the age structure of the population.

Total Fertility Rate

The **total fertility rate** (TFR) is the average number of children that each woman will have during her lifetime. The TFR is an average because, of course, some women will have more, some fewer, and some no children at all. Theoretically, when the TFR = 2, each pair of parents just replaces itself. Actually it takes a TFR of 2.1 or 2.2 to replace each generation - this number is called the **replacement fertility rate -** because some children will die before they grow up to have their own two children. In countries with low life expectancies, the replacement rate is even higher (2.2–3). Figure $6.2.b$ compares the total fertility rate in different countries.




Figure $6.2.b$: Total fertility rate by country in 2020. Most countries in North and South America have a total fertility rate of 1.7-2.5 children per woman, but fertility rates are lower (for example, Canada and Chile) or higher (for example, Guatamala and Bolivia) in a few countries. Most of Europe and northern and Eastern Asia have low fertility rates (generally 1.4-1.6 children per woman). Many Middle Eastern countries and those in southern and southeastern Asia have slightly higher fertility rates (generally 1.7-2.5 children per woman). The fertility rate in Australia is 1.7-1.9 children per woman. High fertility rates are common in African countries. For example, the fertility rate in Niger is 7.0-7.9. Image modified from Korakys(opens in new window) (CC-BY-SA(opens in new window)).

Age Structure


The **age structure** of a population, the proportion of a population in different age classes, is an important factor in population dynamics. The relationship between TFR and the population growth rate (*r*) depends on age structure. For example, if at one


period a population has an unusually large number of children, they will — as they pass through their childbearing years — increase the *r* of the population even if their TFR goes no higher than 2. Most people have children between the ages of 15 and 49. So if a population has a large number of young people just entering their reproductive years, the rate of growth of that population is sure to rise.

Models that incorporate age structure allow better prediction of population growth, plus the ability to associate this growth with the level of economic development in a region. Countries with rapid growth associated with high birth rates have a pyramidal shape in their age structure diagrams, showing a preponderance of younger individuals, many of whom are of reproductive age (figure $6.2.c$).

The age structure of a population also reflects the recent pattern of mortality. In countries where injuries, starvation, and disease, etc. take a heavy toll throughout life, the age structure diagram has a broad base. In countries where almost everyone survives until old age, the bases is narrower.

Rapid growth is most often observed in underdeveloped countries where individuals do not live to old age because of less-than-optimal living conditions, and there is a high birth rate. Age structures of areas with slow growth, including developed countries such as the United States, still have a pyramidal structure, but with many fewer young and reproductive-aged individuals and a greater proportion of older individuals compared to underdeveloped countries. Other developed countries, such as Italy, have zero population growth. The age structure of these populations is more of an inverted pyramid with a greater percentage of middle-aged and older individuals.The actual growth rates in different countries are shown in figure $6.2.d$, with the highest rates tending to be in the less economically developed countries of Africa and Asia.


Afghanistan 2020
Male surplus
Males
Female surplus
Females


Italy, 1st of January 2023
Source: Italian National Institute of Statistics
Chart by: Kaj Tallungs
Males
Male surplus
Females
Female surplus
Age
100+
90
80
70
60
50
40
30
20
10
0
600 000
400 000
200 000
Population
200 000
400 000
600 000



Ecuador 2020
Male surplus
Males
Female surplus
Females

Figure 6.2.c: Typical age structure diagrams are shown. Top: Afghanistan's (2020) population growth model is expansive. The rapid growth diagrams has a broad base indicating a high pre-reproductive population indicating a high reproductive rate. Afghanistan (expansive). by Sdgedfegw - Own work (CC BY-SA 4.0) Middle: Italy's (2023) population growth is constrictive. Constrictive populations have fewer prereproductive and reproductive aged people than older people indicating that the population is declining. Italy (Constrictive). By Kaj Tallungs - Own work (CC BY-SA 4.0) Ecuador (stationary). Bottom: Ecuador's (2020) population model is stationary. In a stationary model, the number of pre-reproductive balances out with the reproductive individuals indicating that the population may level off. There are more older people, there is a long life expectancy, but low birth rate. By Sdgedfegw- own work (CC BY-SA 4.0).




Figure 6.2.d: The percent growth rate of populations in 2020 in different countries is shown. The population growth rate is negative (meaning the population size is declining) in Cuba, Portugal, many countries in Eastern Europe, and Japan. Mexico, much of Central America, Ecuador, Peru, Bolivia, Paraguay, and Haiti have a population growth rate of 1-2%. Most other countries in North and South America have a population growth rate of 0-1%. Western and Northern Europe and most Asian countries have a population growth rate of 0-1%. In the Middle East, Southeast Asia, Australia, northern and southern Africa, Central African Republic, and South Sudan, the population growth rate is generally 1-2%. Examples of countries with a population growth rate above 2% include the remaining African countries, Syria, Yemen, Oman, Iraq, Afghanistan, and Tajikistan. Image modified from Max Roser, Hannah Ritchie and Esteban Ortiz-Ospina (2013) - "World Population Growth". Published online at OurWorldInData.org (CC-BY).

The U.S. Baby Boom

The TFR in the United States declined from more than 4 late in the nineteenth century to less than replacement in the early 1930s. However, when the small numbers of children born in the depression years reached adulthood, they went on a childbearing spree that produced the baby-boom generation (figure 6.2.e). In 1957, more children were born in the United States than ever before (or since).




Figure 6.2.e: Age structure diagrams of the United States population in 1970 and 1985. The x-axis represents the percent of the population. The y-axis represents age groups in increments of five from 0-5 years all the way to 85-90 years. Blue bars to the left represent males, and red bars to the right represent females. Bars taper near the top of the diagram, showing that those in the oldest age categories represent a small percentage of the population. The 1970 age structure diagram bulges around the 10-15 age group, and this represents the baby boom (green oval). In 1985, this bulge has moved to the 25-30 age group. Image: Libretexts: Biology.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- The Human Population from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- The Human Population from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 6.2: The Rate of Human Population Growth is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 12.2: The Rate of Human Population Growth by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

6.3: Factors Affecting Population Growth

Introduction

Families in low- to middle-income countries are often larger but generally use fewer resources per person than families in high income nations. However, as more people seek a “Western” standard of living, average resource use increases. Raising global living standards reduces Earth’s ability to support the human population. As a result, lowering humanity’s overall environmental impact requires both reducing resource use per person and decreasing average family size.

Efforts to reduce family size depend on understanding the factors that influence family decisions and regional population trends. Economic and cultural conditions strongly shape these decisions, and different combinations of these forces can encourage either larger or smaller families.

Although population growth is often discussed at a global level, the factors that influence it vary widely across regions. Understanding these local influences is essential for addressing population growth worldwide.

Economic Factors

Economic factors strongly influence family size and population growth. In low-income countries, rural farming families often need many children to provide labor for planting, harvesting, and selling crops. Small families may not supply enough workers to sustain the farm.

In high-income countries, families tend to be smaller because raising children is expensive. High costs for food, clothing, transportation, education, and college make having many children financially difficult. Technology and education can also reduce the need for large families. Improved tools, irrigation, and farming methods increase productivity, allowing fewer people to work the same land.

Other economic factors matter as well. Limited access to medical care and birth control can increase family size, especially where child mortality is high. In countries without social security systems, parents rely on their children for support in old age, making larger families more desirable. In contrast, national retirement programs and high taxes in high income countries reduce the economic incentive for large families.

Cultural & Religious Factors

Cultural and religious factors influence family size and, in turn, population growth around the world. Religious beliefs can shape attitudes toward family planning, as some religions encourage larger families to maintain traditions or strengthen religious communities. For example, Orthodox Judaism and Roman Catholicism both place value on having many children, and Roman Catholic teaching discourages the use of artificial birth control. Individuals who strongly follow these beliefs may choose larger families even when economic conditions would otherwise encourage smaller ones. Countries with large religious populations, such as Israel and Brazil, tend to have higher birth rates as a result of these influences. Some religions, such as Buddhism, do not explicitly promote large family size.

Factors related to women also play an important role in determining family size. These include access to education, employment opportunities, age at marriage, and social acceptance of birth control. Cultural attitudes toward women often shape how these factors develop within a society.

Global data show that higher levels of education and workforce participation among women, later marriage, and greater acceptance of birth control are generally associated with smaller families. Expanding educational and career opportunities for women is therefore linked to lower population growth rates and improved living standards worldwide.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and

Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

This page titled 6.3: Factors Affecting Population Growth is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 1.8: Population Growth is licensed CC BY 2.0. Original source: https://cnx.org/contents/ApdCwdkA.

6.4: Looking Ahead

Population Projections

A consequence of human population growth is that is takes less time to add a particular number of humans to our Earth. Figure $6.4.a$ shows that 123 years were necessary to add 1 billion humans in 1930, but it only took 24 years to add two billion people between 1975 and 1999.

The human growth rate is slowing (figure $6.4.a$)--we are no longer growing exponentially(opens in new window), and our population is predicted to level off around 2100. Of the current world value for *r* (1.075% in 2019) remains unchanged, the global population would grow from its current 8.3 billion (February 2026) to 9.7 billion by 2050. Some experts predict that the global population will continue to grow until 2100, reaching a peak of almost 11 billion.





Figure $6.4.a$: The time between the addition of each billion human beings to Earth decreases over time. By 1800, there were about a billion people on Earth. It took 130 years, until 1930, for the number to reach two million. Thirty years later, in 1960, the number reached three billion, and 15 years after that, in 1975, the number reached four billion. The population reached five billion in 1987, and six billion in 1999, each twelve years apart. Currently, the world population exceeds seven billion. The population is projected to reach 8 billion in 2028, and 9 billion in 2054. (credit: modification of work by Ryan T. Cragun)Figure $6.4.b$ Population growth rate 1950 to 2100. Our World in Data (CC BY)


Long-term Consequences

Humans are not unique in their ability to alter their environment. For example, beaver dams alter the stream environment where they are built. Humans, however, have the ability to alter their environment to increase our **carrying capacity**, sometimes to the detriment of other species. Earth's human population and their use of resources are growing rapidly, to the extent that some demographers say that humans have already exceeded our carrying capacity. This would mean that the Earth cannot indefinitely support our current population size. Long-term exponential growth carries with it the potential risks of famine, disease, and large-scale death, as well as social consequences of crowding such as increased crime.

The impact of humans on our environment depends not only on population size but also on our consumption patterns. The **ecological footprint** is a measure of the area of land needed to support an individual or country. This includes agricultural lands used to provide food, other lands used to provide goods (such as forests to provide timber) and fuels (such as oil fields to provide fossil fuels), and the land needed to dispose of wastes. (figure $6.4.b$). Many factors affect the size of an ecological footprint from economic development in a country to societal consumption patterns, resource conservation, and individual product choice. Reducing our ecological footprints can increase our carrying capacity and decrease environmental impact. So maybe we should update the question of, "What is the Earth's carrying capacity for humans," to, "What is the carrying capacity of

people on Earth at which we can provide a fair and decent life (satisfying needs) for all humans while causing the least impact on the environment and other species?


World
25G
20G
15G
10G
5G
0
global hectares
1961 1963 1965 1967 1969 1971 1973 1975 1977 1979 1981 1983 1985 1987 1989 1991 1993 1995 1997 1999 2001 2003 2005 2007 2009 2011 2013 2015 2017 2019 2021 2023 2024
Ecological Footprint
Biocapacity
Ecological Deficit
Ecological Reserve
York University, FoDaFo, Global Footprint Network, 2025 National Footprint and Biocapacity Accounts Note: last three years are estimates


United States of America
7
6
5
4
3
2
1
0
Number of Earths
1961 1963 1965 1967 1969 1971 1973 1975 1977 1979 1981 1983 1985 1987 1989 1991 1993 1995 1997 1999 2001 2003 2005 2007 2009 2011 2013 2015 2017 2019 2021 2023 2024
Ecological Footprint
Biocapacity
Ecological Deficit
Ecological Reserve
York University, FoDaFo, Global Footprint Network, 2025 National Footprint and Biocapacity Accounts Note: last three years are estimates



Ecological Footprint of Countries 2022
Carbon
Fishing Grounds
Cropland
Built-up Land
Forest Products
Grazing Land
12.5
10
7.5
5
2.5
0
Number of Earths

Figure 6.4.*b*: Top: World ecological footprint. The green line indicates our biocapacity (global hectares of biologically productive area). The rest line indicates our ecological footprint. The area under the green line indicates our ecological reserve (biologically productive area available), and the red area between the red and green lines shows our ecological deficit. 2021-2024 are estimates. Middle: The ecological footprint of the United States. Rather than showing available hectares in this graph, the Y-axis has been changed to number of earths. Since we have 1 earth available the biocapacity (green line) is horizontal at 1. The ecological footprint (red line) and ecological deficit (area between green line and red line) indicate that we have ranged from needing 3-6 Earth's to support our average way of life in the United States. Bottom: Comparison of Ecological Footprint for various countries (2022). Dark blue indicates carbon footprint, light blue is fishing grounds, orange is cropland, red is built-up land, green is forest products, and yellow is grazing land. Images: Footprint Data Foundation, York University Ecological Footprint Initiative, and Global Footprint Network: National Footprint and Biocapacity Accounts, 2023 edition. Available online at https://data.footprintnetwork.org (CC BY 4.0)

Interactive Element

Are you wondering about the size of your personal ecological footprint? You can estimate it here.(opens in new window)

Earth Overshoot Day

Another way to look at resource usage based on human population growth and consumption patterns is Earth Overshoot Day (figure 6.4.*c*). The graphs below shows how human overconsumption patterns have continued to worsen over time, and countries living in an ecological deficit or reserve (Figure 6.4.*d*)




Figure 6.4.*c*:Earth Overshoot Day: Humans have progressively used more of Earth's resources than the planet can regenerate. Earth's ability to regenerate consumed resources now lasts only about 57% as long as in 1971. Chart showing progression over time of "Earth Overshoot Day", which is the first date on which the Earth cannot regenerate ecological resources used in each year Data source: Past Earth Overshoot Days. *OvershootDay.org*. Earth Overshoot Day (2025). Archived from the original on 20 June 2025. Source advises that during its annual date list updates, all previous years' values may have changed due to more comprehensive historical data and updated accounting methods. Accordingly, any updates to this chart should include the most recently published data values for all years. Image By RCraig09 - Own work, CC BY-SA 4.0, Wikimedia(opens in new window) [commons.wikimedia.org]




Figure 6.4.*d*:Nations living within their ecological means (shaded green) or in ecological overshoot (shaded red) in 2022. By Global Footprint Network, Open Data Platform - Global Footprint Network, Open Data Platform, CC BY-SA 4.0, footprintnetwork(opens in new window) [data.footprintnetwork.org]

Reducing Population Growth (Degrowth) and leveling off (Zero Population Growth)

Reducing the population growth rate (**degrowth**) will limit the consequences of overpopulation; however, regulation of the human population is a complex issue that presents ethical considerations. Childbirth can be influenced by culture, religion, societal and family pressures as well as individual choice.

All else being equal, reductions in total fertility rate (TFR) will slow human population growth. TFR is negatively correlated with improved standard of living, access to health care, and gender equality. As countries develop economically, individuals become increasingly dependent on urban jobs rather than the family farm for income and provisions. While a large family would be needed to tend the family farm, supporting many children in an urban area is expensive. As a result, economic development is associated with people choosing to have fewer children.

Access to healthcare decreases childhood mortality and provides the option of birth control and reproductive care. While decreased infant and childhood mortality alone would increase the population growth rate, people tend to have fewer children when they can be confident that all of their children are likely to survive; thus, as childhood mortality declines, total fertility rate also declines, lowering the population growth rate.

The education of women and girls is also associated with a decreased total fertility rate. Furthermore, family planning education programs have had highly positive effects in some countries on limiting population growth rates and increasing standards of living. When women have knowledge of family planning, power to make decisions about their family size, and a variety of career options, fertility rate declines. In regions that have made recent strides in educating girls and women, families more often choose to have a small number of children so that they can afford to pay for their education regardless of gender. For example Kerala, India has a Total Fertility Rate (TFR) of 1.46 as of 2025 due to high female literacy, universal education, and a strong public healthcare system, including family planning resources.

Extreme efforts to reduce population growth led to the 1979 One-Child Policy in China, which imposed fines on urban couples who had more than one child (figure 6.4.*e*). Additionally, only a couple's first child was granted educational and health care benefits. The effectiveness of the policy in limiting overall population growth is controversial, as was the policy itself. Tragically, this policy combined with the desire of many couples to have a male heir led to female infanticide. Furthermore, enforcement of the policy led to human rights abuses including forced sterilizations and forced abortions. Over the years, some restrictions associated with the One-Child Policy were lifted. For example, in 2013, an adult who was an only child was allowed to have two children. The policy ended in 2016, and as of May 2021, each couple is now permitted to have up to three children.



Figure 6.4.4: A poster explaining and promoting China's One-Child Policy. Image by Arian Zwegers (CC-BY).

Zero population growth (ZPG) occurs when the net birth rate (BR) and immigration (I) equals the net death rate (DR) and emigration (E) such that the net rate of population growth is zero (BR + I) = (DR - E) . Logically, this would occur when total fertility rate equals replacement fertility rate, but as described in the previous section, this depends on the age structure of a population.

Gapminder

Use Gapminder tools to explore demographic data. You can change the variables on each axis and compare various countries to overall world trends.

References

Population Growth (Annual %)(opens in new window). 2019 World Bank Group. Accessed 29 August 2020.

Growing at a slower pace, world population is expected to reach 9.7 billion in 2050 and could peak at nearly 11 billion around 2100(opens in new window). 2019. United Nations Department of Economic and Social Affairs. Accessed 29 August 2020.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- The Human Population from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- The Human Population from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Human Population Growth(opens in new window) from *General Biology* by OpenStax (licensed under CC-BY(opens in new window))

This page titled 6.4: Looking Ahead is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 12.3: Looking Ahead by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

6.5: Data Dive- World Population Densities

Overview

Our World in Data (OWID) is a non-profit scientific online publication associated with University of Oxford. OWID's focus is on providing visual data for global problems. Their 2025 map of human population density can be observed below.




Figure 6.5.a: World density map. Illustrating population density for the year 2025 in number of people per km2 of land area. . Density is indicated with darker or lighter shading using the following categories in persons per square kilometer (0-2, 2-5, 5-10, 10-20, 20-50, 50-100, 100-200, 200-500, 500-1000, 1000+, and no data).Graph by Ourworldindata.org/population_growth (CC BY).

Questions

1. The data on the above graph is overlaid onto a world map. What are benefits of showing the data in this way? What are the challenges of displaying the data this way?
2. What question(s) are the authors trying to answer with this graph?
3. Which areas of the globe have the highest density levels (darkest density shading)?
4. Are there any areas of the globe that surprise you (either high or low) with their density? Why?
5. How could policy makers/world leaders/health organizations use this information? Provide at least 3 ideas.
6. What is missing from this map? Hint: It's very large!

Attribution

Modified by Erin Rempala (CC-BY-NC)

Rachel Schleiger (CC-BY-NC)

This page titled 6.5: Data Dive- World Population Densities is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 12.4: Data Dive- World Population Densities by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7: THE ENERGY SYSTEM

Energy

This chapter explores how humans obtain and use energy to power modern society. Since the Industrial Revolution, fossil fuels such as coal, oil, and natural gas have been the primary energy sources for transportation, electricity generation, industry, and heating. Although these fuels have supported economic growth and technological development, they are finite resources and can cause significant air, water, and land pollution. Their extraction and combustion also release greenhouse gases, primarily carbon dioxide, which are the main drivers of modern global climate change.

When evaluating energy sources, scientists often consider their **life cycle impacts**. A **life cycle assessment** examines the environmental effects associated with all stages of an energy technology, including the extraction of raw materials, manufacturing, construction, operation, maintenance, and disposal. For example, although solar panels and wind turbines produce no greenhouse gases while generating electricity, emissions are associated with their manufacture, transportation, installation, and eventual recycling or disposal. Considering life cycle impacts provides a more complete picture of the environmental costs and benefits of different energy sources.

> **Life Cycle Assessment (LCA):** A method used to evaluate the environmental impacts of a product or technology throughout its entire life, from raw material extraction and manufacturing to operation, maintenance, and disposal.

Nuclear fission provides large amounts of electricity with relatively low greenhouse gas emissions over its life cycle. However, concerns remain about radioactive waste, reactor accidents, and the high costs associated with building and decommissioning nuclear power plants. Scientists are also investigating nuclear fusion, the process that powers the Sun. If successfully developed as a commercial energy source, fusion could provide abundant energy with less long-lived radioactive waste than fission, although it remains in the experimental stage.

Renewable energy sources, including solar, wind, hydropower, geothermal, and biomass, are naturally replenished and generally have lower life cycle greenhouse gas emissions and environmental impacts than fossil fuels. Over the past decade, renewable energy technologies have become increasingly cost-competitive, leading to rapid growth in their adoption worldwide. As societies work to reduce greenhouse gas emissions and address climate change, renewable energy is expected to play an increasingly important role in creating a more sustainable energy future.

This chapter examines the advantages, limitations, and environmental impacts of both renewable and nonrenewable energy resources, as well as the challenges and opportunities involved in transitioning to a more sustainable energy system.

7.1: Fossil Fuels

7.1.1: Types of Fossil Fuels and Formation

7.1.2: Mining, Processing, and Generating Electricity

7.1.3: Fossil Fuel Consumption

7.1.4: Consequences of Fossil Fuels

7.2: Nuclear Energy

7.2.1: Radioactive Isotopes

7.2.2: Generating Electricity with Nuclear Energy

7.2.3: Nuclear Energy Consumption

7.2.4: Pros and Cons of Nuclear Energy

7.2.5: The Promise of Fusion

7.2.6: Data Dive- Global Nuclear Power Generation

7.3: Renewable Energy and Conservation

7.3.1: Renewable Energy History and Consumption

7.3.2: Wind Energy

7.3.3: Solar Energy

7.3.4: Hydropower

7.3.5: Geothermal Energy

7.3.6: Biofuels (Biomass Energy)

7.3.7: Energy Efficiency and Conservation

7.3.8: Data Dive- Global Renewable Energy Generation

thumbnail © User:Colin / Wikimedia Commons / CC BY-SA 3.0

This page titled 7: The Energy System is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

7.1: FOSSIL FUELS

Chapter Hook

Imagine stepping into a Roman bathhouse nearly 2,000 years ago. The warm water and heated rooms that made these public spaces so popular depended on a dark rock pulled from the ground: coal. Long before modern power plants and electricity grids, people around the world had already discovered coal's usefulness. From ancient China and Greece to the Roman Empire and the Aztec civilization, coal helped provide heat, fuel crafts, and support early industries. For thousands of years, this energy-rich rock has fueled human development and transformed societies. Yet coal's history also highlights a challenge that continues today: it is a finite source of energy. As global demand for energy grows, understanding coal's past can help us think critically about the future of energy.



Figure 7.1.*a*: Fossil charcoal in weathered coal. Image by James St. John(opens in new window) in Wikimedia Commons(opens in new window) (CC-BY2.0(opens in new window))

Attribution

Modified by Erin Rempala (CC-BY-NC-SA 4.0 (opens in new window)).

Modified by Rachel Schleiger (CC-BY-NC(opens in new window)).

This page titled 7.1: Fossil Fuels is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 16: Fossil Fuels by Melissa Ha and Rachel Schleiger is licensed CC BY 4.0.

7.1.1: Types of Fossil Fuels and Formation

Fossil Fuels

Fossil fuels are nonrenewable energy sources formed from the remains of plants, algae, and microorganisms that lived millions of years ago. The energy stored in fossil fuels originally came from sunlight. Through photosynthesis, ancient organisms captured solar energy and converted it into chemical energy stored in organic molecules. For this reason, fossil fuels are sometimes described as **fossilized solar energy**.

As discussed in earlier sections on matter and energy flow, organic molecules store chemical energy within their bonds. When fossil fuels are burned, these bonds are broken and rearranged into more stable forms, releasing the stored energy as heat. This energy can then be used to generate electricity, power transportation, heat buildings, and support industrial processes.

Fossil fuels are considered **nonrenewable** because their formation requires millions of years under specific geological conditions. The rate at which humans consume fossil fuels is far greater than the rate at which they can be naturally replaced. In addition, environmental conditions millions of years ago supported exceptionally high rates of organic matter accumulation, meaning similar fossil fuel reserves may not form again even over geologic timescales.

Fossil fuels consist primarily of **hydrocarbons**, molecules made mostly of carbon and hydrogen atoms. Smaller amounts of oxygen, sulfur, nitrogen, and other elements are also present. The size and structure of these hydrocarbon molecules differ among the three major fossil fuels: coal, petroleum (oil), and natural gas.

Because coal contains relatively large and complex hydrocarbon molecules, it exists as a solid at room temperature. Oil contains smaller molecules and is a liquid, while natural gas consists mainly of very small hydrocarbons and is therefore gaseous.

The three major fossil fuels formed from different types of ancient organisms:

- **Coal** formed primarily from land plants that accumulated in ancient swamps and wetlands.
- **Petroleum (oil)** formed from the remains of microscopic marine organisms buried in ocean sediments.
- **Natural gas** formed alongside petroleum and consists primarily of methane (CH_4), the simplest hydrocarbon.

Together, coal, oil, and natural gas provide most of the world's commercial energy, but their extraction and use have significant environmental consequences, including habitat disruption, air pollution, and climate change.

Coal

Coal is a black or dark brown sedimentary rock that forms from the accumulated remains of ancient plants. Most coal deposits originated in vast swamp forests that existed hundreds of millions of years ago, although some of the oldest deposits may have formed from large accumulations of algae before land plants became widespread.

Coal formation began when dead plant material accumulated in waterlogged, oxygen-poor environments where decomposition was slow. Instead of completely decaying, the plant material gradually formed **peat**, a soft, organic-rich material.

Over millions of years, layers of sediment buried the peat. As pressure and temperature increased, water and volatile compounds were squeezed out, concentrating the carbon content. Continued burial transformed the peat into coal (figure $7.1.1.a$). The pressure required for this process came from overlying sediment layers as well as tectonic forces associated with continental movement and mountain-building events.

The formation of coal demonstrates how solar energy captured by ancient plants can remain stored underground for hundreds of millions of years before being released through human use.

Every lump of coal contains energy originally captured from sunlight by plants that lived long before humans evolved. When coal is burned, that ancient solar energy is released in a matter of seconds. This contrast between millions of years of energy storage and rapid consumption highlights one of the central challenges of modern energy use: societies are depleting finite energy reserves far faster than natural processes can replace them.




Figure 7.1.1.*a*: Coal was formed when large plants in swamps died 300 million years ago (before the dinosaurs). Over millions of years, this vegetation was buried under water and dirt (100 million years ago). Eventually, heat and pressure turned the dead plants into coal, which is found under layers of rock and dirt. Image from How coal is formed over millions of years. (opens in new window) [U.S. Energy Information Administration] (public domain).

Coal Types

Not all coal is the same. As buried plant material experiences increasing heat and pressure over millions of years, it is transformed into progressively higher-quality forms of coal (Figure 7.1.1.*b*):

Peat → Lignite → Subbituminous Coal → Bituminous Coal → Anthracite

As coal matures, its carbon content increases while moisture and other compounds are driven off. Higher-grade coals therefore contain more stored chemical energy and release more energy when burned.

- **Lignite (brown coal)** is the lowest-grade coal and has the lowest energy content.
- **Subbituminous coal** has a higher carbon content and energy value than lignite.
- **Bituminous coal** is the most abundant coal used for electricity generation and industrial purposes.
- **Anthracite** is the highest-grade coal, with the highest carbon content and energy output (figure 7.1.1.*b*



Figure 7.1.1.*b*: Anthracitic coal, the highest grade of coal.

Petroleum (Oil) and Natural Gas

Like coal, petroleum (oil) and natural gas are fossil fuels formed from ancient organic matter. However, unlike coal, which formed primarily from land plants, oil and natural gas originated mainly from microscopic marine organisms, especially **phytoplankton** and other algae that lived in ancient oceans.

When these organisms died, their remains settled to the seafloor and were buried beneath layers of sediment. Oxygen-poor conditions slowed decomposition, allowing organic matter to accumulate. Over millions of years, increasing heat and pressure transformed this organic material into petroleum and natural gas.

The fossil fuels are generated within a **source rock**, a rock layer rich in buried organic matter. Because oil and natural gas are less dense than surrounding rock and fluids, they slowly migrate upward. If they encounter a porous rock layer where they can accumulate, that rock becomes a **reservoir**.

A **trap** forms when an impermeable rock layer blocks further upward movement of oil and natural gas. Traps concentrate fossil fuels into deposits that can be economically extracted and are known as **conventional oil and natural gas reservoirs** (figure 7.1.1.*c*.

Petroleum is a liquid mixture of hydrocarbons that can be refined into fuels such as gasoline, diesel, and jet fuel. **Natural gas**, also called **fossil gas**, consists primarily of methane (CH_4), the simplest hydrocarbon, and is commonly used for heating, electricity generation, and industrial processes.




Figure 7.1.1.*c*: Petroleum (oil) and natural (methane) gas were formed from marine microorganisms. (The image text says tiny marine plants, but they were primarily phytoplankton- tiny photosynthetic floating algae and bacteria rather than true plants.) These were covered by layers of silt and sand 300-400 million years ago. Over millions of years, the remains were buried deeper and deeper. The enormous heat and pressure turned the remains into oil and natural gas. Now, oil and natural gas deposits are found underground and can be extracted via drilling through the layers of sand, silt, and rock. Image from National Energy Education Development Project(opens in new window) [U.S. Energy Information Administration] (public domain).

Unconventional Oil and Natural Gas

Not all fossil fuels occur in easily accessible reservoirs (figure 7.1.1.*d*. Fossil fuels that remain dispersed within less permeable rock formations are known as **unconventional resources**. These resources are generally more difficult and expensive to extract and often have greater environmental impacts than conventional oil and natural gas production.

Major unconventional fossil fuel resources include:

- **Oil shale**, a sedimentary rock containing **kerogen**, an organic material that can be converted into liquid fuels through heating.
- **Tight oil and tight natural gas**, which are trapped within dense rock formations and are often extracted using hydraulic fracturing (**fracking**).
- **Tar sands (oil sands)**, deposits containing **bitumen**, a thick, highly viscous form of petroleum that must be mined or heated before processing into usable fuels (figure 7.1.1.*d*. .
- **Coal-bed methane**, natural gas trapped within coal deposits that formed during coal formation.

Because unconventional fossil fuel resources require more intensive extraction and processing, they often consume more energy and water and can result in greater environmental disturbance than conventional fossil fuel production.




Figure 7.1.1.*d*: Conventional oil and natural gas deposits are trapped beneath impervious rock (gray). Conventional natural gas may be associated with oil or nonassociated. Coalbed methane and tight gas found in shale and sandstone are examples of unconventional fossil fuels. Image from EIA.gov(opens in new window) [USGS] (public domain)



Figure 7.1.1.*e*: Tar sandstone from the Miocene Monterrey Formation of California.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Challenges and Impacts of Energy Use and Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Fossil Fuels(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))
- Shale Gas 101(opens in new window). Office of Fossil Energy. U.S. Department of Energy. Accessed 01-12-2021. (public

domain)
- The Process of Unconventional Natural Gas Production(opens in new window). 2021. United States Environmental Protection Agency. Accessed 01-12-2021. (public domain)

This page titled 7.1.1: Types of Fossil Fuels and Formation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 16.1: Types of Fossil Fuels and Formation by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

7.1.2: Mining, Processing, and Generating Electricity

Mining, Extraction, and Energy Production

Before fossil fuels can be used, they must be extracted from the Earth. The extraction method depends on the type and location of the fossil fuel. Coal is mined from rock layers, whereas oil and natural gas are obtained through drilling. Once extracted, fossil fuels are transported and used to generate electricity, power vehicles, heat buildings, and manufacture a variety of products.

Coal

Coal Mining

Coal is extracted using two general methods: **surface mining** and **subsurface mining**.

Surface mining is used when coal deposits are located near the Earth's surface. Large machines remove soil and rock, known as **overburden**, to expose coal seams (figure $7.1.2.a$).

Two common forms of surface mining are:

- **Strip mining**, in which overburden is removed in successive strips to expose and extract coal.
- **Mountaintop removal**, in which explosives are used to remove the top of a mountain and expose the entire coal seam at once.Much of the removed overburden is deposited in nearby valleys, creating valley fills that can bury streams and alter watershed drainage patterns (figure $7.1.2.b$).

Surface mining is generally less expensive than underground mining but can dramatically alter landscapes, destroy habitat, and increase erosion.



Figure $7.1.2.a$: Garzweiler surface mine, Germany. Image by Raimond Spekking / CC BY-SA 4.0 (via Wikimedia Commons)




Figure 7.1.2.*b*: Left: Mountaintop removal uses explosives to loosen overburden at the top of a mountain, which is then displaced into a nearby valley (valley fill). The underlying coal seam is then extracted. A sediment pond collects soil that erodes from the valley fill. Image by EPA(opens in new window) (public domain). Right: A mountaintop removal site. Image by JW Randolph(opens in new window) (public domain)

Subsurface mining (deep mining) is used when coal deposits are buried deeper underground. Miners access coal through tunnels and shafts that may extend hundreds or thousands of meters below the surface.

Although subsurface mining disturbs less land at the surface, it presents safety risks for miners and can cause land subsidence when underground tunnels collapse (figure 7.1.2.*c*).




Figure 7.1.2.*c*: Subsurface mining involves digging tunnels to access coal deposits that are deep underground. Image by EIA(opens in new window) [National Energy Education Development Project] (public domain)

Coal Processing and Transportation

After mining, coal is cleaned and processed to remove rock, sulfur, and other impurities. This improves its heating value and reduces some emissions during combustion.

Coal is then transported to power plants, primarily by rail but also by truck, barge, or ship (figure 7.1.2.*e*). Because transportation can be costly, many coal-fired power plants are located near major coal-producing regions.



Figure 7.1.2.*e*: A freight train loaded with coal briquettes, formed from compressed coal dust, in Morwell, Victoria, Australia. Image and caption (modified) by CSIRO (opens in new window)(CC-BY(opens in new window)).

Generating Electricity from Coal

Most coal is used to generate electricity. At a coal-fired power plant, coal is pulverized into a fine powder and burned in a furnace (figure 7.1.2.*f*). The heat converts water into high-pressure steam, which spins a turbine connected to an electrical generator.

This process converts:

Chemical Energy → Thermal Energy → Mechanical Energy → Electrical Energy

After passing through the turbine, the steam is cooled, condensed back into water, and reused. Cooling water is commonly withdrawn from nearby rivers, lakes, or oceans.

The cooling water is generally returned to its source at a higher temperature than when it was withdrawn, a form of thermal pollution that can affect aquatic ecosystems.

As of 2026, there is only 1 coal power plant in California, the Argus Cogeneration Plant that is capable of generating 63 MW. It is located northeast of Los Angeles.




Figure 7.1.2.*f*: Left: Coal power plant in Helper, Utah. Right: Diagram of a typical steam-cycle coal power plant (proceeding from left to right). Coal enters the boiler (furnace), where it is burned. This boils water, producing steam. The steam turns a turbine, powering a generator. The resultant electricity is passed through a transformer (which changes the voltage) and is then sent through transmission lines. The steam cools and condenses back to liquid water in the condenser. This is facilitator by cooling water from a nearby river. Right image by Tennessee Valley Authority(opens in new window) (public domain).

This video shows how heat energy can be used to generate electricity.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Oil and Natural Gas

Conventional Oil and Natural Gas Extraction

Conventional oil and natural gas are extracted by drilling wells into underground reservoirs where fossil fuels have accumulated beneath a trap.

In many conventional reservoirs, natural gas accumulates above oil because it is less dense. However, some reservoirs contain mainly oil or mainly natural gas.

When a new well is drilled, natural underground pressure may initially force oil and gas to the surface. As pressure declines, water or gas may be injected into the reservoir to increase recovery. Additional techniques, such as steam or carbon dioxide injection, can further improve extraction from older wells.

Oil and natural gas are produced from both:

- **Onshore wells** located on land (figure $7.1.2.g$)
- **Offshore wells** located beneath the ocean floor (figure $7.1.2.h$)

Modern offshore drilling platforms can operate in water more than a mile deep.




Figure $7.1.2.g$: Injection wells transfer water, carbon dioxide, or other substances to an oil deposit increase pressure or change the oil's properties, facilitating extraction. On the right is the production well, through which extracted oil flows. This is facilitated by an injection well for enhanced recovery. On the left is an injection well for disposal through with waste water (produced water) is stored underground. Several confining formations trap substances underground. Closer to the top is the base of underground sources of drinking water, meaning that all drinking water is extracted from above this point. Image by Government Accountability Office(opens in new window) (public domain).



Figure 7.1.2.h: A platform for offshore oil drilling. Image by Pixabay.com(opens in new window) [Pixabay] (Pixabay license(opens in new window)).

Hydraulic Fracturing (Fracking)

Some fossil fuels occur in dense rock formations that do not allow oil or natural gas to flow easily. These resources can be extracted using **hydraulic fracturing**, commonly called **fracking** (figure 7.1.2.i).

Fracking involves several steps:

1. A well is drilled vertically and then horizontally through the oil- or gas-bearing rock layer.
2. Steel casing and cement are installed to help isolate the well from surrounding rock layers and groundwater.
3. Small perforating charges create openings in the casing.
4. A mixture of water, sand, and chemical additives is injected into the well at very high pressure.
5. The pressure fractures the rock, creating pathways for oil and natural gas to move through the formation.
6. Sand, called **proppant**, helps keep the fractures open after the pressure is released.
7. Oil and natural gas flow through the fractures and into the well, where they are brought to the surface.

Fracking has greatly increased oil and natural gas production, particularly from shale formations. However, the process requires large amounts of water and generates wastewater that must be managed carefully. Environmental concerns include habitat disturbance, methane leakage, groundwater contamination from improperly constructed wells or wastewater handling, and induced earthquakes associated primarily with deep wastewater injection.




Figure 7.1.2.*i*: Two diagrams of fracking. Water is mixed with sand and chemicals and then injected into shale trapping tight oil and natural gas or tight sand. This flushes fossil fuels from the fissures that were previously created by explosives. Top: This section shows four underground layers. From top to bottom, they are the shallow aquifer, aquiclude (impermeable layer), deep aquifer, and another aquiclude. Fracking fluid is injected underground through a well surrounded by casing into the gas-bearing formation. Methane (red arrows) might escape from hydraulic fractures in this formation. Additionally, when the fractures intersect with a pre-existing fault, induced seismicity (earthquakes) is possible. Aboveground, fracking fluid is stored in wastewater ponds. Blue arrows and question marks indicate places where toxic wastewater may escape and contaminate groundwater such as from storage containers, wastewater ponds, casing, or faults. Bottom: Proper treatment and disposal of the wastewater is needed to limit environmental impact. 1: Water is acquired. 2: Chemicals are mixed. 3: Fracking fluid is injected into the well. Here, natural gas flows from fissures into the well. 4: Fracking results in flowback and produced water (wastewater). 5: Wastewater undergoes treatment and disposal. Bottom image by USGS(opens in new window) (public domain).

Unconventional Oil Resources

Other unconventional fossil fuel resources require additional processing before use.

- **Oil sands** (also called **tar sands**) contain bitumen, a dense and highly viscous form of petroleum that must be heated or processed before it can be used.
- **Oil shale** contains kerogen, an organic material that can be combusted directly for energy or heated to convert kerogen into liquid and gaseous hydrocarbons.
  (figure 7.1.2.*j*).
- **Coal-bed methane** is natural gas trapped within coal deposits.

Because these resources require more intensive extraction and processing, they generally have larger environmental footprints than conventional oil and natural gas.



Figure 7.1.2.j: Underground mining of oil shale in Estonia.

Refining Crude Oil

The oil that comes from the ground, called **crude oil**, is a mixture of many different hydrocarbons. Before it can be used, it must be separated into useful products at a refinery.

The most important refining process is **fractional distillation** (figure 7.1.2.k). Crude oil is heated, and its components separate according to their boiling points.

The heaviest hydrocarbons remain liquid and collect near the bottom of the distillation column, while lighter hydrocarbons rise higher before condensing.

This process produces products such as:

- Gasoline
- Diesel fuel
- Jet fuel
- Heating oil
- Lubricants
- Asphalt

Additional chemical processing can further modify these products to meet consumer demand (figure 7.1.2.l.




Figure 7.1.2.*k*: The process of fractional distillation involves heating crude oil and allowing the components to cool. As they move up the distillation column, they condense at different levels based on their boiling points. The heaviest compounds have the highest boiling points and condense at the bottom of the column while lighter compounds (low boiling point) condense at the top. From lowest to highest boiling point the petrochemicals produced are residual fuel oil (1050 °F), heavy gas oil (650-1050 °F), diesel and heating oil (450-650 °F), kerosene and jet fuel (350-450 °F), naphtha (used to make gasoline, solvents, cleaning solutions, etc.; 185-350 °F), gasoline blending components (85-185 °F), and butane and lighter products (< 85 °F). Image by U.S. Energy Information Administration(opens in new window) (public domain).

The video below explains the process of fractional distillation. The labeled distillation column at 3:00 shows heated crude oil (400 °C) separating into various petrochemicals. From bottom to top, they are bitumen (> 350 °C), diesel (250-350 °C), kerosene (160-250 °C), naphtha (70-160 °C), petrol (20-70 °C), and gas (< 20 °C).

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online




Figure 7.1.2.*l*: Main products (measured in gallons) made from a barrel of crude oil in 2019. These include residual fuel oil (0.9 gallons), hydrocarbon gas (1.5), other products (6.0), jet fuel (4.4), distillate (12.5), and gasoline (19.4). Note: A 42-gallon (U.S.) barrel of crude oil yields about 45 gallons of petroleum products because of refinery processing gain. The sum of the product amounts in the image may not equal 45 because of independent rounding. Image by EIA(opens in new window) (public domain).

Transporting Oil and Natural Gas

After extraction and refining, oil and natural gas are transported through extensive pipeline networks (figure 7.1.2.*m*). Crude oil is transported to refineries, while refined fuels are distributed to storage facilities, airports, industries, and gas stations.

Natural gas is transported primarily through high-pressure pipelines that connect production areas with homes, businesses, and power plants.

The United States has an extensive network of interstate and intrastate natural gas pipelines.



Figure 7.1.2.*m*: Construction of the controversial Dakota Access Pipeline, which stretches from North Dakota to Illinois. The pictured portion of the pipeline is in Central Iowa. You can learn more about the 2016 protests against this pipeline here.(opens in new window) Image by Dakota Access Pipe Line(opens in new window) (CC-BY(opens in new window)).

Generating Electricity from Natural Gas (Fossil Gas, or Methane Gas)

Natural gas (Fossil gas) is one of the most widely used fuels for electricity generation.

In a natural gas power plant, methane and compressed air are mixed in the combustion chamber and burned. to produce hot gases that spin a turbine connected to a generator (figure $7.1.2.n$). Some power plants also capture waste heat to produce additional steam and electricity, increasing efficiency.

Compared with coal, natural gas generally produces:

- Less carbon dioxide per unit of electricity
- Lower emissions of sulfur dioxide and particulate matter
- Greater flexibility for meeting changes in electricity demand

However, methane leakage during production and transportation can reduce some of these climate advantages because methane is a powerful greenhouse gas.




Figure $7.1.2.n$: This combustion chamber burns either natural gas or oil. Fuel flows through a natural gas line or from oil storage into the combustion chamber. Air passes through the air intake and is compressed in the compressor. Natural gas and compressed air are mixed with compressed air in the combustion chamber and burned. High-pressure combustion gases spin the turbine, which drives the generator. The resultant electric current is then passed through a transformer, which alters voltage. Image and caption (modified) by Tennessee Valley Authority(opens in new window) (public domain).

Although fossil fuels have fueled industrial development and modern societies, their extraction, transportation, and combustion can affect ecosystems and human health. Mining can alter landscapes and habitats, drilling can contaminate water and cause spills, and burning fossil fuels releases carbon dioxide and other pollutants that contribute to climate change and air pollution. These human health and environmental impacts must be weighed against the energy benefits provided by fossil fuels.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Fossil Fuels(opens in new window) and Mining(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))

- Chapter 4: Non-Renewable Energy(opens in new window) from *Introduction to Environmental Science: 2nd Edition* (2018) Biological Sciences Open Textbooks by Zehnder, Caralyn; Manoylov, Kalina; Mutiti, Samuel; Mutiti, Christine; VandeVoort, Allison; and Bennett, Donna (licensed under CC-BY-NC-SA(opens in new window)).
- Fossil Energy Study Guide: Natural Gas(opens in new window). 2014. U.S. Department of Energy. Accessed 01-12-2021.
- Fossil Energy Study Guide: Oil(opens in new window). 2013. U.S. Department of Energy. Accessed 01-12-2021.

This page titled 7.1.2: Mining, Processing, and Generating Electricity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 16.2: Mining, Processing, and Generating Electricity by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

7.1.3: Fossil Fuel Consumption


Fossil Fuel Consumption

Fossil fuels remain the dominant source of energy worldwide. Petroleum, natural gas, and coal together provide most of the energy used for transportation, electricity generation, manufacturing, and heating. Although renewable energy sources such as solar and wind power are growing rapidly, global energy systems continue to rely heavily on fossil fuels.

Fossil Fuels in the United States

In the United States, natural gas is currently the largest source of electricity generation, followed by nuclear energy, renewable energy sources, and coal (figure ({7{.}1{.}3{.}\eatSpaces{a} \endSpaces}\)) Petroleum contributes little to electricity generation because it is used primarily as a transportation fuel.

Coal once dominated electricity production in the United States and played a major role in industrial development. The United States continues to possess large coal reserves, making coal an important domestic energy resource even though coal consumption has declined significantly in recent decades.






Figure 7.1.3.a**: Top:** Contribution of Each Energy Source to Electricity Generation in the U.S. in 2025. U.S. Energy Information Administration(opens in new window) (public domain). Bottom: United States energy consumption by source and sector (2024). Image by US Energy Information Administration (public domain).

Global Energy Use

Globally, fossil fuels continue to provide most of the world's energy. Petroleum remains the largest single energy source, while coal and natural gas continue to play major roles in meeting global energy demand. Renewable energy sources such as wind, solar, hydroelectricity, and geothermal energy have expanded rapidly in recent decades but still provide a smaller share of total global energy consumption than fossil fuels (figure ({7{.}1{.}3{.}\eatSpaces{b} \endSpaces}\)).

The growing use of fossil fuels is closely tied to the Industrial Revolution. Abundant coal resources were one of several factors that contributed to Britain's industrial development during the eighteenth and nineteenth centuries. Coal provided a concentrated source of energy for steam engines, manufacturing, and transportation, helping transform economies and societies around the world.

Interactive Element

Use the interactive below to explore how global energy consumption has changed since 1800 and compare trends among different energy sources.

Interactive Content

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure 7.1.3.*b*: The bottom interactive graph shows the percentage that each energy source contributes to global energy consumption over time since 1800. Global consumption of most energy sources has increased over time. Images by Our Wold in Data (CC-BY).

Fossil Fuel Reserves and Energy Security

Because fossil fuels are nonrenewable, scientists, economists, and policy-makers pay close attention to the size and distribution of remaining **reserves**. A reserve is a known deposit of a fossil fuel that can be extracted economically using current technology and market conditions. Reserve estimates are not fixed and may change over time as new deposits are discovered, extraction technologies improve, prices fluctuate, and patterns of consumption shift. For this reason, reserve estimates should be viewed as snapshots based on current knowledge and conditions rather than exact predictions of when a fuel source will be exhausted (figure 7.1.3.*c*).

The geographic distribution of fossil fuel reserves is highly uneven. Large oil and natural gas reserves are concentrated in a relatively small number of countries, particularly in the Middle East, while other nations possess few domestic reserves and depend heavily on imports. This unequal distribution has important economic and political consequences because access to energy is essential for transportation, electricity generation, food production, manufacturing, and other societal needs.

These concerns are closely tied to **energy security**, which refers to the reliable and affordable availability of energy needed to support society. Countries that rely heavily on imported oil or natural gas may be vulnerable to supply disruptions caused by geopolitical conflicts, trade disputes, natural disasters, or fluctuations in global energy markets. As a result, many nations seek to enhance energy security by diversifying their energy sources, improving energy efficiency, and expanding renewable energy production.




Figure 7.1.3.c: Top: Global proved reserves of oil (top, 2017) and natural gas (bottom, 2020) by country. The area of each blue circle is scaled to reported reserve size. The maps show that fossil fuel reserves are unevenly distributed across the world, with large concentrations of oil and natural gas in the Middle East and substantial natural gas reserves in Russia. Because reserve estimates change over time as new deposits are discovered, extraction technologies change, and economic conditions shift, these maps should be interpreted as snapshots rather than fixed quantities. Source: U.S. Energy Information Administration (EIA), public domain.

Peak Oil

In the 1950s, geologist M. King Hubbert proposed that oil production in a region would eventually reach a maximum rate, known as **peak oil**, before beginning to decline. Hubbert correctly predicted that oil production in the United States would peak around 1970. However, advances in technology, particularly hydraulic fracturing and horizontal drilling, allowed previously inaccessible oil and natural gas resources to be extracted and led to a resurgence in U.S. production.

As a result, the timing of global peak oil production remains a topic of ongoing debate. Some researchers argue that production will eventually decline as reserves are depleted, while others emphasize that technological innovation and changing economic conditions can continue to expand accessible supplies.

Fossil Fuels and the Global Economy

Although the United States continues to import petroleum, domestic oil and natural gas production increased substantially during the early twenty-first century due to advances in drilling and hydraulic fracturing technologies. Consequently, U.S. dependence on imported petroleum is lower than it was during the late twentieth century.

The global oil market is strongly influenced by the Organization of Petroleum Exporting Countries (OPEC), a group of major oil-producing nations that coordinate production targets in an effort to influence oil prices and market stability. Because many OPEC member countries possess large oil reserves, the organization continues to play an important role in global energy markets.

The economic benefits of fossil fuel production are distributed unevenly among nations and communities. In some cases, fossil fuel wealth has stimulated economic growth and development. In others, it has been associated with political conflict, economic inequality, and environmental degradation. These challenges highlight the complex relationship between energy production, economic development, and environmental sustainability.

Looking Ahead

Fossil fuels have powered industrialization, economic growth, and modern society for more than two centuries. However, their finite nature, uneven distribution, and environmental impacts raise important questions about the future of global energy systems. Understanding how fossil fuels are used, where they are found, and how long they may remain economically recoverable provides a foundation for evaluating alternative energy sources and future energy policies.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Challenges and Impacts of Energy Use and Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Hannah Ritchie (2017) - "Fossil Fuels(opens in new window)". Published online at OurWorldInData.org. Accessed 01-16-2021. (licensed under CC-BY(opens in new window))

This page titled 7.1.3: Fossil Fuel Consumption is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 16.3: Fossil Fuel Consumption by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.
- 16.4: Consequences of Fossil Fuels by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

7.1.4: Consequences of Fossil Fuels

History & Benefits of Using Fossil Fuels.

For most of history, people got energy from simple sources like their own strength, animals, and burning wood. Then the Industrial Revolution changed everything by introducing fossil fuels, which powered huge advances in technology, society, and the economy. (figure $7.1.4.a$). The world is currently dependent on fossil fuels, and existing infrastructure and technologies facilitate their continued use.

An advantage of using coal for electricity is that it plentiful and inexpensive, especially in the United States, which has larger coal reserves than any other country. Furthermore, coal mining is a source of jobs and tax income. Coal's economic advantage is dwindling, however, as technologies associated with renewable sources of energy, such as solar and wind, become more efficient and inexpensive. The U.S. Energy Information Administration compared the levelized cost of electricity (LCOE) for technologies that will begin use in 2023, and the cost of coal-generated electricity exceeded that of many renewable sources (figure $7.1.4.b$). The LCOE accounts for the building and operation costs of power plants, solar panels, wind turbines, etc.




Figure $7.1.4.b$: The **levelized cost of electricity** (**LCOE**) for various energy sources. Of these, hydroelectric, geothermal, natural gas, onshore/offshore wind, solar, and biomass are considered renewable. Nuclear and coal are nonrenewable energy sources. Values are in 2018 dollars per **megawatt hour** ($/MWh). As a reference, the average U.S. household, uses about 0.909 MWh of electricity per month. These values are for the newest technologies that will being use in 2023. Hydroelectric energy is the least expensive, at about $39/MWh. Hydroelectric energy is the least expensive, at about $39/MWh. Next is geothermal ($41/MWh), natural gas ($41/MWh), onshore wind ($56/MWh), solar ($60/MWh), nuclear ($78/MWh), biomass ($92/MWh), coal ($99/MWh), and offshore wind ($130/MWh). Graph by Melissa Ha (CC-BY-NC(opens in new window)) using data U.S. Energy Information Administration, Annual Energy Outlook 2019 (opens in new window)(public domain).

Oil and natural gas continue to meet global energy needs. Despite expansions in renewable energy use, no alternative energy source is currently sufficient to replace oil and natural gas. (A combination of different renewable sources could be possible in the future.) While the United States does rely on imported oil, it continues to produce some oil and natural gas (mostly through fracking), bolstering U.S. energy independence. Local and state economics in regions rich in oil and natural reserves depend on continued extraction of these fossil fuels.

While all fossil fuels harm cause some degree of environmental harm, natural gas is a preferred fossil fuel for electricity generation when considering its environmental impacts. When burned, coal emits nearly double the carbon dioxide that natural gas does. Additionally, much less nitrogen oxides and sulfur dioxide (both air pollutants) are emitted from burning natural gas. It also does not produce ash as coal does (see below).

Health and Environmental Impacts

Fossil fuel usage is the main cause of Global Climate Change (See Global Climate Change(opens in new window) for more details). The negative impacts of fossil fuel use begin with the extraction of the resource. Fossils fuels are often located far from where they are utilized so they need to be transported by pipeline, tankers, rail or trucks. These all present the potential for accidents, leakage, and spills. Additional negative impacts are associated with processing, electricity generation, and disposal of the waste generated.

Coal Mining and Usage

Surface mining of coal disrupts local ecosystems above coal deposits as overburden is removed to access them (figure $7.1.4.c$). In mountaintop removal, a large volume of overburden is dumped over nearby habitats, causing further destruction (figure $7.1.4.d$). Mountaintop removal has affected large areas of the Appalachian Mountains in West Virginia and Kentucky. Habitat loss from coal mining decreases biodiversity, resulting in a loss of ecosystem services. Both surface and subsurface mining expose rocks that can contain contaminants, such as heavy metals or sulfates, which them leach into streams or other bodies of water. This not only harms aquatic life, but it also disrupts nutrient cycling. One of the largest environmental impacts of subsurface mining may be the methane gas that must be vented out of mines to make the mines a safe place to work. Methane gas is a potent greenhouse gas and contributes to Global Climate Change. Finally, the process of mining ultimately compacts soil. This combined with the loss of trees, which slow the flow of run-off and promote infiltration, increases the risk of flooding.



Figure $7.1.4.c$: Peak Downs and Saraji Coal Mine in Queensland, Australia. To access the coal, vegetation was removed, destroying habitat for native organisms. Image by Lock the Gate Alliance(opens in new window) (CC-BY(opens in new window)).



Figure 7.1.4.d: Mountaintop Removal Coal Mining in Pike County, Kentucky Photograph shows mountaintop coal removal mining in Pike County, Kentucky. Source: By https://www.flickr.com/people/15648670@N00 iLoveMountains.org - Mountaintop Removal Mine Site above Route 23 in Pike County, Kentucky, CC BY 2.0, https://commons.wikimedia.org/w/inde...curid=33452045

Coal miners face health hazards such as explosions, mine collapse, and exposure to toxic fumes. **Black lung disease** is a respiratory condition characterized by coughing and shortness of breath that occurs in miners exposed to too much coal dust. Residents near mines also risk exposure to coal dust and underground toxins following explosions. As a result of exposure to toxins, birth defects and other health problems are common in residents near mines.

Coal is considered the "dirtiest" source of energy because its combustion results in the most air pollution. Coal power plants emits a variety of air pollutants including sulfur dioxide, nitrogen oxide, and heavy metals. Sulfur dioxide and nitrogen oxide are sources of acid rain (acid deposition), smog, and health issues. Heavy metals cause neurological and developmental problems in humans and other animals. Burning of coal emits particulate matter and higher amounts of carbon dioxide per unit of energy than the use of oil or natural gas. Carbon dioxide is the most frequently emitted greenhouse gas and causes climate change. In 2018, electricity generation was responsible 27% of greenhouse gas emissions in the United States, and much of this was emitted from coal power plants. The transportation of coal usually relies on fossil fuels, releasing further pollution.

Ash (including fly ash and bottom ash) is a residue created when coal is burned at power plants. In the past, fly ash was released into the air through the smokestack, where it would contribute to particulate matter air pollution. Laws now require that much of the fly ash now must be be captured by pollution control devices, like scrubbers. In the United States, fly ash is generally stored at coal power plants or placed in landfills. Ash from storage or landfills can spill or leach into groundwater, resulting in water pollution.

Conventional Extraction Oil and Natural Gas

Exploring and drilling for oil degrades land and ocean habitats. On land, extensive infrastructure such as road networks, transport pipelines and housing for workers are needed to support a full-scale drilling operation. These can pollute soil and water, fragment habitats, and disturb wildlife. Extraction of oil and natural gas is also hazardous for workers, who have a high incidence of cancer and heart disease.

Human-caused oil spills in rivers and oceans harm ecosystems. From an economic perspective, oil spills disrupt the fishing industry and tourism. Oil spills at sea are generally much more damaging than those on land, since they can spread for hundreds of nautical miles in a thin oil slick which can cover beaches with a thin coating of oil. This can kill sea birds, mammals, shellfish and other organisms it coats. Oil spills on land are more readily containable if a makeshift earth dam can be rapidly bulldozed around the spill site before most of the oil escapes, and land animals can avoid the oil more easily.

Oil spills can result from supertanker accidents such as the Exxon Valdez in 1989, which spilled 10 million gallons of oil into the rich ecosystem of coastal Alaska and killed massive numbers of animals. The largest marine oil spill began in April 2010 when a

natural gas explosion at an oil well 65 km offshore of Louisiana occurred on Deepwater Horizon Oil Rig. It killed 11 employees and flowed for 3 months in 2010, releasing an estimated 200 million gallons of oil (figure 7.1.4.*d*). Wildlife, ecosystems, and people's livelihood were adversely affected. A lot of money and huge amounts of energy were expended on immediate clean-up efforts. The long-term impacts are still not known. The National Commission on the Deepwater Horizon Oil Spill and Offshore Drilling was set up to study what went wrong. The worst oil spill ever occurred during the Persian Gulf war of 1991, when Iraq deliberately dumped approximately 200 million gallons of oil in offshore Kuwait and set more than 700 oil well fires that released enormous clouds of smoke and acid rain for over nine months.



Figure 7.1.4.*d*: Platform supply vessels battle the blazing remnants of the off shore oil rig Deepwater Horizon. A Coast Guard MH-65C dolphin rescue helicopter and crew document the fire aboard the mobile offshore drilling unit Deepwater Horizon, while searching for survivors. Multiple Coast Guard helicopters, planes and cutters responded to rescue the Deepwater Horizon's 126 person crew. Image and caption (modified) from US Coast Guard(opens in new window) (public domain).

During an oil spill on water, oil floats to the surface because it is less dense than water, and the lightest hydrocarbons evaporate, decreasing the size of the spill but polluting the air. Then, bacteria begin to decompose the remaining oil, in a process that can take many years. After several months only about 15% of the original volume may remain, but it is in thick asphalt lumps, a form that is particularly harmful to birds, fish, and shellfish. Cleanup operations can include a variety of components, but each has its procs and cons. Skimmer ships that vacuum oil from the water surface, but these are effective only for small spills. Controlled burning works only in early stages before the light, ignitable part evaporates, but this also pollutes the air. **Dispersants** are detergents that break up oil to accelerate its decomposition, but some dispersants may be toxic to the ecosystem. **Bioremediation** refers to adding microorganisms that specialize in quickly decomposing oil, but this can disrupt the natural ecosystem.

Unconventional Extraction of Oil and Natural Gas

Fracking causes more environmental damage than conventional extraction. The considerable use of water (figure 7.1.4.*e*) may affect the availability of water for other uses in some regions, and this can affect aquatic habitats. In fact, fracking consumes more water than the use of nuclear energy, coal, or conventional oil and natural gas. If mismanaged, hydraulic fracturing fluid can be released by spills, leaks, or various other exposure pathways that contaminate land and groundwater (figure 7.1.4.*f*). Fracking fluid flowback – the fluid pumped out of the well and separated from oil and gas – not only contains the chemical additives used in the drilling process but also contains heavy metals, radioactive materials (which release radiation), volatile organic compounds, benzene (a carcinogen), toluene, ethylbenze, xylene, and other toxic air pollutants. Volatile organic compounds (VOCs) can react with the atmosphere to form ground-level ozone, which is associated with respiratory disease. Toulene can cause dizziness, confusion, headaches, and miscarriages. Ethylbenzene is a possible carcinogen that also causes dizziness, eye irritation, and hearing loss. Xylene also causes dizziness and headaches and furthermore can be fatal at high concentrations. In some cases, this contaminated water is sent to water treatment plants that are not equipped to deal with some of these classes of contamination. Finally, injecting wastewater for disposal can even induce earthquakes.




Figure 7.1.4.e: Fracking influences the water cycle. Water acquisition is the withdrawal of groundwater or surface water to make hydraulic fracturing fluids. Chemical mixing involves combining a base fluid, sand, and additives at the well site to create hydraulic fracturing fluids. During well injection, hydraulic fracturing fluids move through the oil and gas production well and in the targeted rock formation. Produced water handling refers to the onsite collection and handling of water that returns to the surface after hydraulic fracturing and the transportation of that water for disposal or reuse. Finally, the disposal and reuse of wastewater occurs. If this is not done properly, wastewater can pollute surrounding areas. Image and caption (modified) by EPA(opens in new window) (public domain).




Figure 7.1.4.f: The toxic fluid injected during fracking can escape, contaminating groundwater. For example, it can leak through a hole in the casing, move through faulty cement, or move through a fault in the rock that confines the deposit. Image by Government Accountability Office(opens in new window) (public domain).

Other unconventional sources of fossil fuels can also harm the environment. Surface mining of tar sands or oil shales requires the removal of all vegetation and leaves pollutants behind, causing habitat loss (figure 7.1.4.g).



Figure 7.1.4.*g*: An open pit mine in the tar sands oil fields of Alberta, Canada. Conifer trees, which previously provided habitat to native animals, have been cleared to allow for mining. Image by Howl Arts Collective(opens in new window) (CC-BY(opens in new window)).

Transportation, Refineries, and Combustion

Natural gas is released into the atmosphere from coal mines, oil and gas wells, and natural gas storage tanks, pipelines, and processing plants. These leaks are the source of about 25% of total U.S. methane emissions, which translates to three percent of total U.S. greenhouse gas emissions, contributing to climate change. When natural gas is produced but cannot be captured and transported economically, it is "flared," or burned at well sites, which converts it to carbon dioxide. This is considered to be safer and better than releasing methane into the atmosphere because carbon dioxide is a less potent greenhouse gas than methane. However, when natural gas with high concentrations of the toxic gas hydrogen sulfide is flared, it produces carbon dioxide, carbon monoxide, sulfur dioxide, nitrogen oxides, and many other compounds (see Air Pollution for more details).

Leaks also happen when we use petrochemicals on land. For example, gasoline sometimes drips onto the ground when people are filling their gas tanks, when motor oil gets thrown away after an oil change, or when fuel escapes from a leaky storage tank. When it rains, the spilled petrochemicals get washed into the gutter and eventually flow to rivers and into the ocean. Another way that oil sometimes gets into water is when fuel is leaked from motorboats and jet skis. When a leak in a storage tank or pipeline occurs, petrochemicals can also get into the ground, and the ground must be cleaned up. To prevent leaks from underground storage tanks, all buried tanks are supposed to be replaced by tanks with a double lining.

Oil refining emits a variety of toxins and is the single largest source of benzene (figure 7.1.4.*g*). As a result, residents living near oil refineries have a high incidence of cancer, asthma, and birth defects. When petrochemicals such as gasoline or diesel are burned, they release a variety of air pollutants, including carbon dioxide (a cause of climate change), sulfur dioxide, nitrous oxides, volatile organic compounds (VOCs), particulate matter, and lead (see Air Pollution for more details). The transport of oil by ship or trunk also requires energy in the form of fossil fuels, generating more pollutants. Compared to oil and coal, burning natural gas releases the fewest pollutants and greenhouse gases.



Figure 7.1.4.*g*: Pollutants are emitted from the Pine Bend oil refinery in Rosemount, Minnesota. Image by Tony Webster(opens in new window) (CC-BY(opens in new window)).

Solutions

The main solution to the pollution and climate destabilizing effects of fossil fuels is to stop using fossil fuels as quickly as possible and switch to renewables for all our energy needs (see Renewable Energy and Conservation for more information).

Reclamation can help reduce the habitat damage caused by mining or fossil-fuel extraction by restoring the land once operations end. This process may include reshaping disturbed ground, covering it with topsoil to shield organisms from heavy metals, radioactive materials, and other toxins, and neutralizing acidic runoff created when sulfates leach from exposed rock. Afterward, vegetation is replanted and disrupted water flow is partially restored.

However, reclamation cannot fully recreate the complex landscapes, stream networks, mature forests, or ecological systems that existed before mining. Even so, it does make it easier for native species to return than if the land were left untreated.

So-called **"clean coal"** technologies have existed since the 1980s and can reduce some air pollution from burning coal, but they are not truly *clean*. Burning coal still creates major environmental and health risks (figure 7.1.4.*i*). Some technologies remove toxins like mercury before combustion, while others capture pollutants such as sulfur dioxide or coal ash during burning. For example, **smokestack scrubbers** filter out sulfur dioxide, nitrogen oxides, particulate matter, and mercury before the exhaust is released. In effect, these systems turn what would have been air pollution into solid waste that still must be safely disposed of. Today, the term **"clean coal"** usually refers to **carbon capture and storage (CCS)**(figure 7.1.4.*h*). In CCS, emissions from burning fossil fuels are passed through a solvent that traps carbon dioxide, which is then separated out. In theory, some CO_2 could be reused in industrial processes, and the rest stored deep underground. While CCS can reduce some emissions and pollutants, it does not eliminate fossil-fuel air pollution. It also requires 25-40% more energy, lowering the efficiency of power plants. CCS facilities are expensive to build and maintain, and retrofitting older coal plants would further raise electricity costs. The technology also carries risks, including leakage, groundwater contamination, and increased seismic activity.

By the middle of this century, countries would need to put 6 billion tons of carbon dioxide underground each year, according to the United Nations. This plan would require much more land and infrastructure, hundreds of thousands of miles of new pipelines, and many new tankers to move the gas. It would also need about 2,000 places underground to store the CO_2, meaning a new storage site would have to be built every 4 days for the next 25 years. This is estimated to cost tens of trillions of dollars. Many of the 12 existing CCS sites hold less CO2 than estimated, some have been plagued by broken pipes, leaky rocks, and other problems.

MIT civil and environmental engineering professor Charles Harvey and Kurt House, CEO of KoBold Metals, argue that **carbon capture and storage (CCS)** is a poor investment because it mainly enables oil and gas companies to continue producing fossil

fuels rather than reducing them. They note that CCS projects have repeatedly failed, cost taxpayers billions, and cut far fewer emissions than renewable energy. In their view, funding CCS slows the transition to clean energy, and those resources would be better spent on proven cheaper, solutions like wind and solar power..

Interactive Element

For more on the problems with carbon capture and why it is not the solution to the climate crisis see this excellent 6/25/26 interactive article from ProPublica: Why Carbon Capture Can't Conceivably Solve the Climate Change.

.




Figure 7.1.4.*h*: Burning fossil fuels releases carbon dioxide into the atmosphere, but there are also opportunities to sequester (store) carbon dioxide, offsetting some fo this damage. Some of the carbon dioxide injected during tertiary recovery (enhanced oil recovery) remains underground. Additionally, carbon dioxide may be captured from smoke released by power plants and other industries and stored under ground. Image by USGS(opens in new window) (public domain).



Figure 7.1.4.*i*: A protest against clean coal at the United Nations Climate Change Conference (COP19) in 2013. Image by 350.org(opens in new window) (CC-BY-NC-SA(opens in new window)).

Because fossil fuels are limited, their reserves will eventually run out, and the world will have to rely on other energy sources. In practice, we won't extract the very last fossil fuels because the cost will become too high to make it worthwhile. The market is already shifting toward renewables, which are cheaper and far less harmful to people and the environment. Alongside expanding renewable energy, **energy conservation** — using energy more efficiently and avoiding waste — is essential, since the cleanest and safest energy is the energy we never have to use.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0 and:

- Gidden, M.J., Joshi, S., Armitage, J.J. *et al.* A prudent planetary limit for geologic carbon storage. *Nature* **645**, 124–132 (2025). Nature(opens in new window) [doi.org]
- Grossman, D. (2022, May 19). *Everything you need to know about clean coal: Is carbon capture and storage technology a pipe dream or panacea? Popular Mechanics.*
- Harvey, C., & House, K. (2022, August 16). *Every dollar spent on this climate technology is a waste. The New York Times. NYTimes(opens in new window) [www.nytimes.com]*
- MIT Climate Portal. (n.d.). *How much carbon dioxide would we have to remove from the air to counteract climate change?* Massachusetts Institute of Technology. MIT.edu(opens in new window) [climate.mit.edu]
- Ritchie, H., & Roser, M. (n.d.). *Fossil fuels.* Our World in Data. Our World in Data(opens in new window) [ourworldindata.org]
- Worth, K., & Waldron, L. (2026, June 25). *Why carbon capture and storage won't fix our climate crisis.* ProPublica. [projects.propublica.org/why-carbon-capture-cant-solve-climate-change/]

Modified by Melissa Ha from the following sources:

- Challenges and Impacts of Energy Use, Non-Renewable Energy Sources, and Water Pollution from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Fossil Fuels(opens in new window) from *An Introduction to Geology* by Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))
- Chapter 4: Non-Renewable Energy(opens in new window) from *Introduction to Environmental Science: 2nd Edition* (2018) Biological Sciences Open Textbooks by Zehnder, Caralyn; Manoylov, Kalina; Mutiti, Samuel; Mutiti, Christine; VandeVoort, Allison; and Bennett, Donna (licensed under CC-BY-NC-SA(opens in new window)).

This page titled 7.1.4: Consequences of Fossil Fuels is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 16.4: Consequences of Fossil Fuels by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

7.2: NUCLEAR ENERGY

Case Study: The Legacy of Chernobyl

It looks like a scene from a post-apocalyptic novel: a city slowly deteriorating after decades without human inhabitants. Schools, homes, and businesses stand abandoned, with personal belongings still scattered across floors and tables, suggesting the hurried evacuation of thousands of residents. Yet Chernobyl is not fiction. It is the site of the world's worst nuclear power plant accident.

On April 26, 1986, Reactor 4 at the Chernobyl Nuclear Power Plant, then part of the Soviet Union, exploded during a safety test. The accident released large amounts of radioactive material into the atmosphere, contaminating surrounding ecosystems and exposing millions of people across Europe to radioactive fallout. More than 100,000 residents were evacuated from the region, and a restricted exclusion zone remains around the site today.

The Chernobyl disaster directly killed 31 people in the days and months following the accident. However, radiation exposure affected millions of people across Europe, and scientists continue to study the long-term health impacts, including increased risks of certain cancers.

The Chernobyl disaster continues to shape public perceptions of nuclear energy. Supporters of nuclear power point to its ability to generate large amounts of electricity with very low greenhouse gas emissions, while critics emphasize the risks of accidents, radioactive waste, and long-term environmental contamination. As you read this section consider the following question: *How should societies weigh the benefits and risks of different energy sources when meeting their energy needs?*



Figure 7.2.a: Chernobyl radiation map. Image Sting(opens in new window) (vectorisation), MTruch(opens in new window) (English translation), Makeemlighter(opens in new window) (English translation) in Wikimedia Commons(opens in new window) (CC-BY-SA2.5(opens in new window))

Nuclear Power

Nuclear power is energy produced from **nuclear fission**, a process in which the nuclei of heavy elements such as uranium-235 split apart and release large amounts of energy. In nuclear power plants, the heat generated by fission reactions is used to produce steam, which drives turbines to generate electricity. Because nuclear power plants emit little carbon dioxide during operation, nuclear energy is often considered an alternative to fossil fuels for electricity generation.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha and Rachel Schleiger from Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

7.2.1: Radioactive Isotopes

7.2.2: Generating Electricity with Nuclear Energy

7.2.3: Nuclear Energy Consumption

7.2.4: Pros and Cons of Nuclear Energy

7.2.5: The Promise of Fusion

7.2.6: Data Dive- Global Nuclear Power Generation

This page titled 7.2: Nuclear Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.2.1: Radioactive Isotopes

Radioactive Isotopes and Half-Life

Atoms are made of protons and neutrons in a central nucleus surrounded by electrons. Atoms of the same element always have the same number of protons, but they may differ in the number of neutrons. These different forms of an element are called **isotopes**.

For example, all carbon atoms contain six protons. Carbon-12 has six neutrons, while carbon-14 has eight neutrons. Because both have six protons, they are both carbon, but they are different isotopes of carbon.

Some isotopes are unstable because their nuclei contain excess energy. These unstable isotopes, called **radioactive isotopes** or **radioisotopes**, spontaneously emit radiation as they decay into more stable forms (figure $7.2.1.a$). **Radiation** is the emission of energy or particles from unstable atoms as they undergo radioactive decay. Radiation may be released as particles or electromagnetic energy. Depending on the type and amount of exposure, radiation can damage living cells and tissues.

During **radioactive decay**, one element may be transformed into another. For example, carbon-14 gradually decays into nitrogen-14 while releasing radiation (figure $7.2.1.b$). Radioactive decay occurs at a predictable rate that is unique to each radioisotope.




Figure $7.2.1.a$: Isotopes of hydrogen. All of these atoms have one proton (pink circle labeled "p^+"), but protium has no neutrons, deuterium has one neutron (orange circle labeled "n"), and tritium has two neutrons. The proton and neutron(s) are located in the center of the atom (the nucleus). An electron (blue circle labeled "e^-") rotates around each atom nucleus. Image from National Isotope Development Center [National Isotope Development Center](public domain).





Figure 7.2.1.*b*: A **radioactive isotope** of carbon (carbon-14) has six protons and eight neutrons. It decays to a stable isotope of nitrogen (nitrogen-14), which has seven protons and seven neutrons. Radioactive decay releases radiation. (The particular type of radiation that occurs in this example is called beta minus decay, β-.) Carbon-14 decays at a predictable rate, with half of it decaying every 5730 years. Because carbon is abundant in organisms, this predictable decay rate is commonly used for dating fossils. Image from CDC (public domain).

Types of Radiation

Radioactive isotopes release different types of radiation as they decay(figure 7.2.1.*a*). The three most common forms are **alpha (α) particles**, **beta (β) particles**, and **gamma (γ) rays**. These types of radiation differ in their ability to penetrate matter and their potential effects on living organisms.

Alpha particles consist of two protons and two neutrons. They are relatively large and can be stopped by a sheet of paper or even the outer layer of human skin. **Beta particles** are high-energy electrons that can penetrate farther than alpha particles but are usually blocked by metal or other dense materials. **Gamma rays** are high-energy electromagnetic radiation that can penetrate much more deeply and require thick layers of lead, concrete, or steel for shielding. Some radioactive materials also emit **neutrons**, which can be blocked by water or other hydrogen-rich materials.

The health risks of radiation depend on the type of radiation, the amount of exposure, and whether radioactive materials enter the body. Understanding how different types of radiation behave is important for protecting people and ecosystems from radioactive contamination.



Figure

: Different types of radiation vary in their ability to penetrate matter. Alpha particles are stopped by paper, beta particles are stopped by metal, and gamma rays require dense shielding such as lead or concrete. Water is effective at reducing neutron radiation.

Image and caption (modified) from Flowers, Theopold, and Langley/OpenStax (CC BY-SA 4.0; OpenStax).

Half-Life

The **half-life** of a radioactive isotope is the time required for half of the original amount of the isotope to decay (figure $7.2.1.d$). For example, the half-life of carbon-14 is 5,730 years, while the half-life of uranium-238 is about 4.5 billion years.

After one half-life, 50% of the original radioactive isotope remains (figure $7.2.1.e$).. After two half-lives, 25% remains. After three half-lives, only 12.5% remains. Because radioactive decay occurs at a predictable rate, scientists can estimate how long radioactive materials remain in the environment.

Understanding half-life is important when evaluating the environmental impacts of nuclear energy. Some radioactive materials produced during nuclear power generation remain hazardous for thousands of years and require careful long-term storage.




Figure 7.2.1.*d*: After each half life, 50% of a radioactive isotope decays. After one half-life, 50% (1/2; 0.5) of a radioactive isotope remains. After two half-lives, only 25% of the original radioactive isotope remains. After three half-lives, it is 12.5%; four half-lives = 6.25%; five half-lives = 3.125%. Image by Openstax [Fraknoi, Morrison, and Wolff] (CC-BY). Access for free at openstax.org.




Figure 7.2.1.*e*: During radioactive decay, an unstable isotope can release radiation and transform into a different element. In this example, uranium-238 emits an alpha particle and becomes thorium-234. Image by OpenStax (CC-BY). Download for free at openstaxcollege.org.

Nuclear Fission and Energy Production

Natural radioactive decay releases energy, but much larger amounts of energy can be produced through **nuclear fission**. During nuclear fission, the nucleus of a large atom splits into smaller nuclei, releasing energy and additional neutrons (figure 7.2.1.*f*).

Most nuclear power plants use **uranium-235** as fuel. When a uranium-235 nucleus absorbs a neutron, it becomes unstable and splits into smaller nuclei. This process releases a large amount of heat energy along with additional neutrons. These neutrons can trigger fission in nearby uranium atoms, creating a self-sustaining **chain reaction**.

The large amount of heat produced by nuclear fission can be used to generate electricity. In a nuclear power plant, the heat is used to boil water and produce steam. The steam spins turbines connected to generators, converting the energy released by fission into electrical energy.

Because nuclear power plants emit very little carbon dioxide during operation, nuclear energy is often considered a low-

carbon source of electricity. However, concerns remain about radioactive waste, nuclear accidents, and the long-term storage of radioactive materials. These topics are discussed in later sections.




Figure 7.2.1.*f*: Nuclear fission and fusion are physical processes that produce energy from atoms. Nuclear fission involves splitting the nucleus of an atom, but nuclear fusion involves combining smaller nuclei into a larger one. Image by US DOE [Sarah Harmann](public domain).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Matter from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 7.2.1: Radioactive Isotopes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.2.2: Generating Electricity with Nuclear Energy

The Nuclear Fuel Cycle

Uranium used in nuclear power plants must be mined, processed, and enriched before it can be used as fuel. The **nuclear fuel cycle** describes the stages through which nuclear fuel passes from extraction to disposal (figure $7.2.2.a$).

The process begins with **uranium mining**, in which uranium-containing rock is extracted from the ground. The ore is then processed to concentrate the uranium into a material called **yellowcake**. Next, the uranium is chemically processed and **enriched** to increase the proportion of uranium-235, the isotope most commonly used in nuclear reactors. The enriched uranium is manufactured into small ceramic pellets that serve as nuclear fuel.

After the fuel has been used in a nuclear reactor, it becomes **spent fuel**. Although it can no longer efficiently sustain nuclear reactions, it remains radioactive and must be stored, reprocessed, or disposed of safely. Managing radioactive waste is one of the major challenges associated with nuclear energy.




Figure $7.2.2.a$: The nuclear fuel cycle begins with uranium mining and processing, followed by fuel fabrication and use in a nuclear reactor. After use, spent fuel may be stored, reprocessed, or disposed of as radioactive waste. Image by U.S. NRC (public domain)

Nuclear Reactors

The processed uranium fuel is formed into small ceramic pellets and sealed inside metal tubes called **fuel rods** (figure $7.2.2.b$). Thousands of fuel rods are bundled together to form the **reactor core**, where nuclear fission occurs. A single uranium fuel pellet contains approximately the same amount of usable energy as a ton of coal.



Figure 7.2.2.*b*: An assembly of fuel rods containing pellets of nuclear fuel. Image by Alternative Energies and Atomic Energy Commission, France (public domain)

Inside the reactor core, uranium-235 nuclei absorb neutrons and split in a process called **nuclear fission**. Each fission reaction releases energy and additional neutrons, which can trigger further fission reactions. This series of self-sustaining reactions is called a **chain reaction** (figure 7.2.2.*c*).




Figure 7.2.2.*c*: A nuclear chain reaction begins when a uranium-235 atom absorbs a neutron and undergoes fission, releasing energy and additional neutrons that can trigger more fission reactions. Image and caption (modified) by Fastfission (public domain).

To prevent the chain reaction from occurring too rapidly, reactors use **control rods** made of neutron-absorbing materials such as boron. Inserting control rods into the reactor core absorbs more neutrons and slows the reaction. Withdrawing the rods allows more fission reactions to occur and increases heat production. By carefully adjusting the position of the control rods, operators can regulate the amount of energy produced.

Generating Electricity from Nuclear Energy

A **nuclear reactor** contains the reactor core and the equipment needed to convert the energy released during fission into electricity (figure 7.2.2.*d*). The reactor core is surrounded by water, which serves two important functions. First, it transfers heat away from the fuel rods. Second, it slows down neutrons, making them more likely to cause additional fission reactions.

The heat produced by nuclear fission converts water into steam. The steam turns a turbine connected to a generator, producing electricity. After passing through the turbine, the steam is cooled, condensed back into liquid water, and reused. In this way, nuclear power plants operate similarly to coal- or natural gas-fired power plants; the primary difference is that heat is produced by nuclear fission rather than by burning fuel.




Figure 7.2.2.*d*: In a nuclear power plant, heat from nuclear fission is used to produce steam, which turns a turbine connected to a generator. Afterward, the steam is cooled and recycled. Image and caption (modified) by EPA (public domain).

Types of Nuclear Reactors

Most commercial nuclear power plants use either **pressurized water reactors (PWRs)** or **boiling water reactors (BWRs)**. Both designs use heat from nuclear fission to generate steam and electricity, but they differ in how water is circulated and heated within the system.

In a pressurized water reactor, water surrounding the reactor core is kept under high pressure so that it does not boil (figure 7.2.2.*e*). Instead, it transfers heat to a separate water system, where steam is produced to drive a turbine. Keeping the reactor water separate from the steam-producing water reduces the spread of radioactive materials beyond the reactor core.




Figure 7.2.2.*e*: There are three separate streams of water in a pressurized water reactor. The first stream is associated with the reactor core in the reactor pressure vessel. The second stream is associated with the steam generator. The third stream is from an external source and is used to condense the steam. The reactor pressure vessel contains control rods and fuel rods (not labeled). The pressure tank regulates water pressure. Hot water from the reactor moves to the steam generator, where it heats water in the second stream to produce steam. Steam from the steam generator moves through the steam line and turns a turbine, powering the electric generator. Steam cools in the condenser, and the cooled water is then pumped back into the steam generator. The reactor pressure vessel, pressure tank, and steam generator are all in a containment structure. Water from an external source is used to cool steam in the condenser. The water from the external source cools in the cone-shaped cooling tower. Image byDOE [Office of Nuclear Energy] (public domain).

Boiling Water Reactor

The reactor Boiling water reactors operate somewhat differently because water in contact with the reactor core is allowed to boil directly, producing steam that drives the turbine (figure 7.2.2.*f*). Despite these design differences, both reactor types generate electricity from the heat released during nuclear fission.




Figure 7.2.2.*f*: A boiling water reactor has two streams of water. In the first stream, the same water that bathes the reactor core in the reactor pressure vessel and turns to steam and turns a turbine. The third stream is from an external source and is used to condense the steam. The reactor pressure vessel contains control rods and fuel rods (not labeled). It is within a larger containment structure. When water is heated by the reactor, it evaporates to steam. This moves through the steam line and turns a turbine, powering the electric generator. Steam cools in the condenser, and the cooled water is then pumped back into the reactor pressure vessel. Water from an external source is used to cool steam in the condenser. The water from the external source cools in the cone-shaped cooling tower. Image by Office of Nuclear Energy (DOE) [Office of Nuclear Energy] (public domain).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Non-renewable Energy Sources from *AP Environmental Science* by University of California College Prep (CC-BY). Download for free at CNX.
- Stages of the Nuclear Fuel Cycle. 2020. U.S. NRC. Accessed 01-16-2021 (public domain).
- Nuclear explained: The nuclear fuel cycle. 2020. U.S. Energy Information Administration. Accessed 01-16-2021 (public domain).

This page titled 7.2.2: Generating Electricity with Nuclear Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.2.3: Nuclear Energy Consumption

Nuclear Energy Consumption

Nuclear energy became an important source of electricity within a few decades of the development of nuclear technology during World War II. The world's first commercial nuclear power station began operating at Calder Hall in England in 1956, demonstrating that nuclear fission could be used to generate electricity on a large scale. Over the following decades, countries around the world invested in nuclear power, and the number of operating reactors increased steadily (figure $7.2.3.a$).

Growth in nuclear energy slowed after several high-profile accidents raised concerns about reactor safety. The 1979 accident at Three Mile Island in Pennsylvania and the 1986 Chernobyl disaster in Ukraine led many countries to reconsider plans for expanding nuclear power. Following the 2011 accident at the Fukushima Daiichi Nuclear Power Station in Japan, several nations reduced their reliance on nuclear energy, and Japan temporarily shut down all of its reactors while safety reviews were conducted.

Despite these challenges, nuclear energy remains an important source of electricity in many countries. Advocates emphasize that nuclear power produces large amounts of electricity with very low greenhouse gas emissions during operation, while critics point to concerns about accidents, radioactive waste, and high construction costs. As a result, nuclear energy continues to be the subject of debate as societies seek to reduce carbon emissions while meeting growing energy demands.



Figure $7.2.3.a$: Arkansas Nuclear One, a power plant in Russellville, Arkansas. Image by Edibobb(opens in new window) (CC-BY(opens in new window)).

Nuclear energy provides a substantial share of the world's electricity. In 2019, nuclear power generated about 10% of global electricity. Although nuclear generation expanded rapidly during the second half of the twentieth century, growth slowed after accidents at Three Mile Island, Chernobyl, and Fukushima raised concerns about reactor safety.

Interactive Element

Use the interactive graph below to investigate how nuclear electricity generation has changed over time. Compare the United States, the world, and one other country using the country-selection menu on the right. As you examine the data, look for similarities, differences, and any major increases or decreases in nuclear power generation. Press the play button below the graph to view changes over time.

Interactive Content

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure 7.2.3.*b*: Global nuclear power generation in TWh over time. Nuclear power production initially increased but remained relative flat in the early 2000s. The dip in 2011 is due to the temporary shutdown of nuclear reactors in Japan following the Fukushima Daiichi Nuclear Power Station meltdown. Image from Hannah Ritchie & Pablo Rosado, Our World in Data (CC-BY).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 7.2.3: Nuclear Energy Consumption is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.2.4: Pros and Cons of Nuclear Energy

Weighing the Benefits and Risks of Nuclear Energy

Nuclear energy presents a complex sustainability challenge. On one hand, nuclear power plants generate large amounts of electricity while emitting very little carbon dioxide during operation, making nuclear energy a potential tool for reducing greenhouse gas emissions and addressing climate change. On the other hand, nuclear power produces radioactive wastes that must be managed for thousands of years, carries the risk of serious accidents, and raises concerns about nuclear weapons proliferation. As societies seek reliable and low-carbon sources of energy, scientists, policymakers, and citizens must carefully evaluate these benefits and risks when considering the role of nuclear energy in a sustainable future.

Advantages of Nuclear Energy

Compared to fossil fuels, nuclear power generates electricity with very low greenhouse gas emissions and little air pollution (figure $7.2.4.a$). Because nuclear power plants do not burn fuel during operation, they can produce large amounts of electricity without releasing significant amounts of carbon dioxide. As a result, some scientists and policymakers view nuclear energy as a potential tool for reducing greenhouse gas emissions and slowing climate change.

Nuclear energy is also a highly concentrated source of energy. Small amounts of uranium can produce large amounts of electricity, and known uranium reserves could supply nuclear reactors for many decades. In addition, nuclear power plants are among the most reliable sources of electricity. Their **capacity** is the maximum amount of electricity they can generate when operating at full power, and their **capacity factor** measures how often they operate near that maximum output. In 2020, nuclear power plants in the United States had an average capacity factor of 92.5%, meaning they produced electricity at nearly full capacity for most of the year (figure $7.2.4.b$).

Because nuclear power provides reliable electricity regardless of weather conditions and emits relatively little greenhouse gas over its lifecycle, it is often considered a low-carbon alternative to fossil fuels.




Figure 7.2.4.*a*:

Comparison of the health impacts and lifecycle greenhouse gas emissions of major energy sources. Death rates are measured as deaths per terawatt-hour (TWh) of electricity produced and include deaths associated with accidents and air pollution. Greenhouse gas emissions are measured as carbon dioxide-equivalent emissions over the entire life cycle of each energy source, including fuel extraction, construction, operation, and decommissioning. Compared to fossil fuels, nuclear energy is associated with relatively low death rates and very low greenhouse gas emissions, although other environmental and social concerns remain important considerations. Image by Hannah Ritchie and Max Roser/Our World in Data (CC-BY).




Figure 7.2.4.*b*: Capacity factor measures how often a power plant operates at or near its maximum power output. In 2020, nuclear power had the highest capacity factor (92.5%), meaning nuclear plants generated electricity at nearly full capacity for most of the year. Geothermal (74.3%) and natural gas (56.6%) also had relatively high capacity factors, while hydropower (41.5%), coal (40.2%), wind (35.4%), and solar (24.9%) operated at lower average capacities. Image from the Office of Nuclear Energy, U.S. Department of Energy (public domain).

Negative Impacts of Nuclear Energy

Despite its advantages, nuclear energy has several environmental, economic, and social drawbacks. One concern is the environmental impact of the nuclear fuel cycle. Mining uranium can disturb habitats, increase erosion, and release pollutants

from underground rock formations. Nuclear power plants also require large amounts of water for cooling. Water released back into rivers, lakes, or oceans is often warmer than the surrounding water, causing **thermal pollution**, which can alter aquatic ecosystems and stress organisms adapted to cooler temperatures.

Nuclear power plants are expensive to build and maintain, requiring substantial investments in specialized materials, infrastructure, and safety systems. In addition, technologies used to enrich uranium for nuclear fuel can potentially be adapted to produce materials for nuclear weapons, raising concerns about **nuclear proliferation** (figure 7.2.4.c).

Perhaps the greatest concerns associated with nuclear energy are **radioactive waste** and the possibility of **nuclear accidents**. Although major accidents are rare, they can have long-lasting environmental, economic, and social consequences. Furthermore, spent nuclear fuel remains radioactive for thousands of years, and no permanent disposal solution has been widely implemented. As a result, the safe management of radioactive waste remains one of the greatest challenges facing the nuclear energy industry.

The issues of radioactive waste storage and nuclear accidents are discussed in more detail below.




Figure 7.2.4.c: Pie charts showing the relative proportions of uranium-238 (blue) and uranium-235 (red) at different levels of enrichment. Nuclear weapons require more highly enriched uranium than is needed for nuclear fuel. Natural uranium (NU) consists of more than 99.2% U-238 and 0.72% or less of U-235. Low-enriched uranium (LEU) consists of less than 20% U-235, with reactor-grade uranium typically 2-5% U-235. High-enriched uranium (HEU) consists of 20-85% U-235 with weapons grade at least 85% U-235. Image and caption (modified) by Fastfission (public domain).

Nuclear Waste

One of the greatest environmental challenges associated with nuclear energy is the management of **radioactive waste**. Because many radioactive materials have long half-lives, they can remain hazardous for thousands of years. Nuclear waste is commonly classified as **high-level radioactive waste**, **low-level radioactive waste**, and **uranium mill tailings**.

High-Level Radioactive Waste

High-level radioactive waste (HLRW) consists primarily of **spent nuclear fuel** removed from nuclear reactors. Although the fuel can no longer efficiently sustain fission reactions, it remains highly radioactive and continues to generate heat.

Spent fuel is initially stored in deep pools of water, which cool the fuel and shield workers from radiation. After sufficient cooling, the fuel may be transferred to **dry casks**, heavily shielded steel and concrete containers designed for long-term interim storage (figure 7.2.4.d). In the United States, most spent nuclear fuel remains stored at or near nuclear power plants because a permanent national repository has not yet been established.




Figure 7.2.4.*d*: Dry casks seal spent fuel rods in a canister, which is surrounded by a larger storage cask. Bundles of used fuel assemblies are inside. Image by NRC (public domain).

One proposed solution for long-term storage was **Yucca Mountain** in Nevada, a deep geological repository designed to isolate radioactive waste underground for thousands of years. Although construction of the facility began, political opposition and concerns about transportation and long-term safety have prevented its use (figure 7.2.4.*e*).



Figure 7.2.4.*e*: Yucca Mountain is the proposed site for long-term storage of high-level radioactive waste in the U.S., but it is not in use due to political controversy. Image by Public Domain.

Some countries, including France and Japan, **reprocess** spent nuclear fuel to recover usable materials and reduce the volume of waste requiring disposal. Reprocessing can extract additional energy from nuclear fuel and reduce long-term storage needs. However, it is expensive and raises concerns about the potential diversion of nuclear materials for weapons production. The United States currently does not commercially reprocess spent nuclear fuel.

Low-Level Radioactive Waste

Low-level radioactive waste (LLRW) includes items contaminated with small amounts of radioactive material, such as protective clothing, tools, filters, and laboratory equipment. Although less hazardous than spent fuel, these materials must still be handled carefully.

Low-level radioactive waste is typically stored, treated, and disposed of in specialized facilities designed to prevent the release of radioactive materials into the environment.

Uranium Mill Tailings

Uranium mill tailings are the waste materials left behind after uranium ore is processed. Although the uranium has been removed, the remaining material may still contain radioactive elements and can release **radon**, a radioactive gas.

Tailings represent the largest category of nuclear waste by volume. They are typically stored in engineered containment areas called **impoundments**, which are designed to limit the spread of radioactive materials into surrounding ecosystems. Although less radioactive than spent fuel, their large volume and long-term persistence present ongoing environmental challenges.

Key Takeaway

Unlike many other forms of waste, radioactive waste can remain hazardous for thousands of years. Finding safe and secure methods for storing and managing this waste remains one of the most significant challenges associated with nuclear energy.

Nuclear Accidents and Reactor Safety

Modern nuclear power plants are designed with multiple safety systems intended to prevent accidents and contain radioactive materials. Despite these precautions, accidents can occur because of equipment failures, human error, design flaws, natural disasters, or unexpected events. Although major nuclear accidents are rare, they can have significant environmental, economic, and social consequences. Three accidents in particular, Three Mile Island, Chernobyl, and Fukushima Daiichi, have shaped public perceptions of nuclear energy.

The Three Mile Island accident occurred in Pennsylvania in 1979 when a combination of equipment malfunctions and operator errors caused a partial meltdown of a reactor core. Small amounts of radioactive gases were released, but no deaths were directly attributed to the accident. Long-term studies have found little evidence that radiation exposure from the accident increased cancer rates in the surrounding population.

The Chernobyl disaster occurred in 1986 at a nuclear power plant in what is now Ukraine. During a safety test, a flawed reactor design combined with operator mistakes led to a power surge, explosions, and a fire that released large amounts of radioactive material into the atmosphere. Thirty-one workers and emergency responders died in the days and months following the accident, and scientists estimate that thousands of additional cancer deaths may eventually result from radiation exposure. More than 100,000 people were permanently evacuated from the surrounding area, and an exclusion zone remains around the site today.

The Fukushima Daiichi nuclear disaster occurred in Japan in 2011 after a powerful earthquake and tsunami disabled reactor cooling systems. Several reactor cores overheated and partially melted, releasing radioactive materials into the environment (figure $7.2.4.f$). Although no deaths were caused by acute radiation exposure, more than 100,000 people were evacuated, and the disaster caused widespread social and economic disruption. Cleanup and decommissioning efforts are expected to continue for decades.

These accidents illustrate both the potential consequences of nuclear power and the importance of reactor design, safety systems, emergency preparedness, and effective regulation. They remain central to ongoing debates about the future role of nuclear energy in a sustainable energy system.




Figure 7.2.4.f: Left: Diagram (approximate) of the Fukushima I Nuclear Power plant accidents. (1) Unit 1: Explosion, roof blown off on 12 March. (2) Unit 2: Explosion on 15 March; contaminated water in underground trench, possible leak from suppression chamber. (3) Unit 3: Explosion, most of concrete building destroyed on 14 March, possible plutonium leak. (4) Fire on 15 March; water level in spent fuel pools partly restored. (5) Multiple trenches: probable source of contaminated water, partly underground, leaked stopped on 6 April. Image (modified) by Sodacan (CC-BY). Bottom: The Fukushima I Nuclear Power Plant after the 2011 Tōhoku earthquake and tsunami. Reactor 1 to 4 from right to left. Image by Digital Globe (CC-BY-SA 3.0).

Interactive Element

This three-minute *NPR: All Things Considered* segment, What Recovery Looks Like In Japan Almost A Decade After Fukushima Nuclear Disaster(opens in new window), provides on update on evacuees from the Fukushima nuclear disaster.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Non-Renewable Energy Sources from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Nuclear Energy(opens in new window), Environmental Challenges in Energy, Carbon Dioxide, Air, Water and Land Use(opens in new window), and Systems of Waste Management(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors. Textbook content produced by Tom Theis and Jonathan (CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Non-renewable Energy Sources(opens in new window) from *AP Environmental Science* by University of California College Prep (CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Stages of the Nuclear Fuel Cycle(opens in new window) and Low-Level Waste(opens in new window). 2020. U.S. NRC. Accessed 01-16-2021 (public domain).
- What is Generation Capacity?(opens in new window) 2020. Mike Mueller. Office of Nuclear Energy. U.S. Department of Energy. Accessed 01-16-2021 (public domain)

This page titled 7.2.4: Pros and Cons of Nuclear Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.2.5: The Promise of Fusion

Fusion Power

The Sun produces energy through nuclear fusion, a process that joins small atomic nuclei together to form larger nuclei. On Earth, scientists are working to recreate this process by heating hydrogen isotopes to extremely high temperatures, creating a superheated state of matter called plasma. When atomic nuclei collide with enough energy, they can fuse and release large amounts of energy. A common fusion reaction combines deuterium and tritium, two forms of hydrogen, to produce helium and energy (figure $7.2.5.a$). Although fusion power plants do not yet generate electricity commercially, researchers continue to study this technology because of its potential to provide a low-carbon energy source.





Figure $7.2.5.a$: Deuterium-Tritium nuclear reaction. Deuterium and Tritium are isotopes of Hydrogen. Credit © ITER Organization(opens in new window), ITER.org(opens in new window) [www.iter.org]

Although no commercial fusion power plants are operating today, fusion energy could offer several important benefits if scientists and engineers succeed in producing electricity from this process. Fusion reactions are controlled by powerful magnetic fields, so if the power supply is interrupted, the reaction stops almost immediately. As a result, fusion reactors cannot experience the type of runaway meltdown associated with some nuclear fission reactors. Fusion is also expected to produce far less long-lived radioactive waste than current nuclear technologies. Because its fuel can be derived from hydrogen isotopes found in water, fusion has the potential to provide a reliable, low-carbon source of energy with a strong safety profile.

Scientists and engineers are actively working to make fusion energy a viable source of electricity. In San Diego, the DIII-D National Fusion Facility has been studying fusion technology for many years. One of the most ambitious international projects is ITER, a large experimental fusion reactor being built in southern France through collaboration among many nations. ITER is designed to test technologies needed for future fusion power plants. These systems typically use a doughnut-shaped chamber called a tokamak, where strong magnetic fields hold and control superheated plasma so that fusion reactions can occur (figure $7.2.5.b$).






Figure 7.2.5.*b*). Top: A cut-away look inside the ITER vaccum vessel. The copper colored "D" rings are electromagnets. Middle: A look inside the JET (Joint European Torus) TOKAMAK with the plasma highlighted in red during operation on the left, and without fusion on the right. Bottom: A visualization of the plasma moving around a TOKAMAK *Visualization courtesy of Jamison Daniel, Oak Ridge Leadership Computing Facility*. Image credits: © ITER Organization(opens in new window), ITER.org(opens in new window) [www.iter.org]

This page titled 7.2.5: The Promise of Fusion is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

7.2.6: Data Dive- Global Nuclear Power Generation

Figure 7.2.6.a: Global nuclear power generation. Graph by Our World in Data (CC-BY)

Questions

1. What countries generate the most energy via nuclear power?
2. Is there anything that surprises you about the patterns observed on this graph?
3. Based on what you learned in this chapter about nuclear power, do you think the trends you see in this 2019 map continue for the next 50 years? Explain.

Raw Data For Above Graph(s)

Raw data table link for Our World In Data graph above. (CC-BY)

Attribution

Rachel Schleiger (CC-BY-NC)

This page titled 7.2.6: Data Dive- Global Nuclear Power Generation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

7.3: RENEWABLE ENERGY AND CONSERVATION

Hidden Sources of Renewable Energy

When most people think of renewable energy, they picture wind turbines spinning on a hillside or solar panels covering rooftops. Few would expect a city's drinking water system to generate electricity. Yet in Portland, Oregon, hydroelectric turbines installed inside municipal water pipes produce power around the clock as water flows downhill to homes and businesses. Because the system uses existing infrastructure, it generates renewable electricity without building a dam or disrupting river ecosystems.

Portland's project highlights an important idea: renewable energy can come from many sources and can be produced in unexpected places. As societies work to reduce greenhouse gas emissions and dependence on fossil fuels, scientists and engineers continue to develop new ways to harness Earth's renewable energy flows. Could the energy systems of the future be built into the infrastructure we already use every day?



Figure 7.3.a: Water turbine example. Image by Pixabay(opens in new window) (Public Domain)

7.3.1: Renewable Energy History and Consumption

7.3.2: Wind Energy

7.3.3: Solar Energy

7.3.4: Hydropower

7.3.5: Geothermal Energy

7.3.6: Biofuels (Biomass Energy)

7.3.7: Energy Efficiency and Conservation

7.3.8: Data Dive- Global Renewable Energy Generation

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Rachel Schleiger (CC-BY-NC(opens in new window)).

This page titled 7.3: Renewable Energy and Conservation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18: Renewable Energy by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.1: Renewable Energy History and Consumption

Renewable Energy: History and Consumption

Renewable energy comes from sources that are naturally replenished on human timescales. Major renewable energy sources include **solar energy**, **wind energy**, **hydropower**, **geothermal energy**, **biomass energy**, and **tidal energy**.

Although these energy sources appear very different, most can be traced to just three fundamental sources of energy. The sun provides energy for solar power and drives many other renewable energy sources. Solar heating of Earth's surface creates the temperature differences that generate wind, drives the water cycle that powers rivers and hydropower systems, and fuels photosynthesis, which produces biomass. **Geothermal energy** comes from heat within Earth's interior, and **tidal energy** comes from the gravitational pull of the Moon and Sun on Earth's oceans (figure $7.3.1.a$).



Figure $7.3.1.a$: A tidal current energy generator at Race Rocks in British Columbia, Canada. Image by Garry Fletcher(opens in new window) (CC-BY-SA(opens in new window)).

The Sun is also the ultimate source of energy stored in **fossil fuels** because coal, oil, and natural gas formed from ancient organisms that captured solar energy through photosynthesis. However, fossil fuels require millions of years to form and are therefore considered **nonrenewable**.

Most renewable energy sources produce little or no greenhouse gas emissions during operation and generally cause less air pollution than fossil fuels. However, renewable energy technologies can still affect the environment. For example, hydropower dams can alter river ecosystems and fish migration, large solar and wind facilities can require substantial land area, and biomass combustion releases air pollutants and carbon dioxide. As a result, renewable energy sources are often considered more sustainable than fossil fuels, but they are not entirely impact free.

Interest in renewable energy increased during the energy crises of the 1970s, when oil shortages and rising fuel prices highlighted the risks of depending heavily on fossil fuels. During the following decades, countries expanded their use of different energy sources based on local conditions. For example, Iceland developed geothermal energy, Scandinavian countries expanded hydropower and wind energy, and many regions invested in solar energy as costs declined.

In recent decades, concerns about climate change, air pollution, and energy security have accelerated the growth of renewable energy worldwide. Improvements in technology and falling costs have made solar and wind power among the fastest-growing sources of electricity. Together with hydropower, geothermal energy, biomass, and emerging technologies such as tidal energy, renewable energy is playing an increasingly important role in the global transition to lower-carbon energy systems.

Use the interactive map below to explore the global transition to renewable energy (figure $7.3.1.b$). Compare the electricity mixes

of the United States and one additional country of interest. What energy sources dominate electricity production in each location? How have the contributions of fossil fuels, nuclear power, and renewable energy changed over time? Press the play button to view shifts in electricity generation through the years.

Interactive Content

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Interactive Element




Figure 7.3.1.*b*: Per capita electricity generation by source interactive graph. By Hannah Ritchie and Pablo Rosado Data source: Ember (2026); Energy Institute - Statistical Review of World Energy (2025); Population based on various sources (2024). OurWorldinData.org/electricity-mix |

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Renewable Energy(opens in new window) and Challenges and Impacts of Energy

Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 7.3.1: Renewable Energy History and Consumption is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.1: Renewable Energy History and Consumption by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

7.3.2: Wind Energy

Wind Energy

Wind energy is generated from the movement of air, which is ultimately driven by uneven heating of Earth's surface by the Sun. As warm air rises and cool air sinks, differences in air pressure create wind. Wind turbines capture the kinetic energy of moving air and convert it into electricity (figure $7.3.2.a$).

The blades of a wind turbine function similarly to airplane wings. As wind flows across the blades, differences in air pressure cause the rotor to spin. The rotor is connected to a generator, either directly or through a gearbox that increases the rotational speed. As the rotor turns, magnets spin within coils of wire, generating electricity through electromagnetic induction. This is the same basic process used in coal-, natural gas-, hydroelectric-, and nuclear-powered generators.

Wind energy has become one of the fastest-growing sources of renewable electricity worldwide. In the United States, wind accounted for approximately 26% of new electricity-generating capacity added in 2025. Improvements in turbine design and declining costs have also made wind energy one of the least expensive sources of new electricity generation. In 2026, the levelized cost of electricity from onshore wind was estimated at approximately $26 to $54 per megawatt-hour globally and about $30 to $40 per megawatt-hour in the United States.

> **Note:** Comparisons of energy costs often use the **levelized cost of energy (LCOE)**, which estimates the average cost of generating electricity over a power plant's lifetime. While LCOE is useful for comparing technologies, it does not account for factors such as energy storage, transmission upgrades, grid reliability, or the variability of wind and solar resources. As a result, the overall cost of integrating renewable energy into the electrical grid may differ from its reported LCOE.




Figure $7.3.2.a$: (1) As the wind blows over the blades of a wind turbine, it causes the blades to lift and rotate. (2) The rotating blades turn a shaft that is connected to a generator. (3) The generator creates electricity as it turns. Image and caption (modified) from EPA(opens in new window) (public domain).

Interactive Element

Click here for an interactive animation illustrating how wind turbines work(opens in new window).

Types of Wind Turbines

Wind turbines vary in size, design, and location, but all convert the kinetic energy of moving air into electricity.

Onshore Wind Turbines

Onshore wind turbines are installed on land and are the most common type of wind energy technology (figure $7.3.2.b$). They are often grouped into large **wind farms** in areas with strong and consistent winds, such as grasslands, mountain passes, and agricultural regions. Onshore wind energy is currently one of the least expensive sources of new electricity generation in many parts of the world.



Figure $7.3.2.b$: Left: The Nordex(opens in new window) N50 wind turbine. Right: visitor centre of Lamma Winds(opens in new window) in Hong Kong(opens in new window), China. By Patrickmak (opens in new window)- Own work, CC BY-SA 3.0, wikimedia.org(opens in new window) [commons.wikimedia.org]. Right image by Joshua Winchell/USFWS (public domain).

Onshore Wind Turbines

Onshore wind turbines are installed on land and are the most common type of wind energy technology (figure $7.3.2.c$). They are often grouped into large **wind farms** in areas with strong and consistent winds, such as grasslands, mountain passes, and agricultural regions. Onshore wind energy is currently one of the least expensive sources of new electricity generation in many parts of the world.






Figure 7.3.2.*c*: Top image: Thorntonbank Wind Farm(opens in new window), using REpower(opens in new window) 5M 5 MW turbines in the North Sea(opens in new window) off the coast of Belgium(opens in new window). By © Hans Hillewaert, CC BY-SA 4.0(opens in new window), wikimedia.org(opens in new window) [commons.wikimedia.org]. Bottom image: CGP(opens in new window) Grey(opens in new window) by CGP Grey(opens in new window) (CC-BY(opens in new window)).

Unconventional Wind Turbines

Some wind turbines use alternative designs that differ from the familiar three-bladed horizontal-axis turbine. Examples include **vertical-axis wind turbines**, **counter-rotating turbines**, and small turbines integrated into buildings, rooftops, or light poles (figure 7.3.2.*d*). These designs are often used for small-scale or specialized applications and may be better suited to urban environments where wind direction changes frequently. However, they generally produce less electricity than large utility-scale wind turbines.








Figure 7.3.2.d: Top: Counterrotating turbine animation by Wikideas1(opens in new window) - Own work, CC BY-SA 4.0(opens in new window). Bottom left: Vertical-axis offshore wind turbine(opens in new window) animation by Wikideas1(opens in new window) - Own work, CC BY-SA 4.0(opens in new window). Bottom right: a light pole turbine (opens in new window)animation by Wikideas1(opens in new window) - Own work, CC BY-SA 4.0(opens in new window).

Advantages of Wind Energy

Wind energy is one of the lowest-cost sources of new electricity generation, and the expansion of wind power creates jobs in manufacturing, construction, installation, and maintenance (figure 7.3.2.e). Because wind turbines generate electricity without burning fuel, they produce little air pollution and emit very little greenhouse gas over their lifetimes compared to fossil fuel power plants.

Unlike thermoelectric power plants that burn coal, natural gas, or use nuclear fission, wind turbines require little water to operate. This reduces water consumption and avoids the thermal pollution associated with power plant cooling systems.

Wind turbines also have a relatively small physical footprint. Although wind farms may occupy large areas of land, most of the space between turbines can still be used for agriculture or grazing. This can provide an additional source of income for farmers and ranchers while allowing working landscapes to remain intact.

Offshore wind farms may offer additional advantages because winds over oceans and large lakes are often stronger and more consistent than winds over land. As a result, offshore turbines can generate more electricity under favorable conditions. In some locations, offshore wind development may also reduce conflicts with agriculture and other land uses.




Figure 7.3.2.e: Samsø, Denmark, transitioned from dependence on fossil fuels to a renewable energy system based on wind, solar, and biomass. The island's experience illustrates how a combination of renewable energy sources can provide electricity and heat while reducing greenhouse gas emissions and increasing energy independence. Images by GDS Infographics(opens in new window) (CC-BY(opens in new window)) and National Geospatial-Intelligence Agency(opens in new window) (public domain).

Disadvantages of Wind Energy

Wind energy has several challenges. Most importantly, wind turbines can generate electricity only when sufficient wind is available. Because wind speeds change over time, wind power is **intermittent**, meaning that electricity production can fluctuate from hour to hour and season to season. These variations can be reduced by connecting wind farms across large geographic areas, storing excess energy in batteries or other storage systems, and combining wind power with other energy sources. However, these solutions require additional infrastructure and can increase costs.

Some people also view wind turbines as visually intrusive, particularly in scenic landscapes, and others object to the noise produced by rotating blades. Although modern turbines are generally quieter than earlier designs, concerns about noise and aesthetics can influence public support for wind energy projects.

Wind turbines can affect wildlife, particularly birds and bats that collide with turbine blades. However, wind energy accounts for only a small fraction of human-caused bird mortality. Buildings, vehicles, power lines, domestic cats, and habitat loss kill far more birds each year than wind turbines. Careful site selection, turbine design, and operational practices can further reduce wildlife impacts. When the broader environmental effects of fuel extraction, air pollution, and climate change are considered, wind energy generally has a smaller overall ecological impact than fossil fuel energy sources.

Like all forms of energy production, wind power requires materials, land, and infrastructure. Constructing turbines, roads, transmission lines, and concrete foundations consumes energy and raw materials and can alter local habitats. Although these impacts are generally smaller than those associated with fossil fuel extraction and combustion, they should still be considered when evaluating the sustainability of wind energy.

No energy source is without environmental impacts. Wind energy reduces greenhouse gas emissions and air pollution compared to fossil fuels, but it can affect wildlife, landscapes, and land use. Evaluating sustainability requires considering both the benefits and drawbacks of each energy source and comparing their overall impacts across environmental, social, and economic

dimensions.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Renewable Energy(opens in new window) and Challenges and Impacts of Energy Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- The Power of Wind(opens in new window). Office of Energy Efficiency and Renewable Energy. U.S. Department of Energy. Accessed 01-17-2021 (public domain).

This page titled 7.3.2: Wind Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.2: Wind Energy by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.3: Solar Energy

Solar Energy

Solar energy is energy from sunlight that can be used to generate electricity, provide heating, and illuminate buildings. It is by far the most abundant renewable energy source available to humanity. Every hour, Earth receives enough solar energy from the Sun to exceed the amount of energy used by all people worldwide in an entire year. Although practical limitations such as energy storage, transmission, and land use prevent all of this energy from being captured, solar energy has enormous potential to meet society's energy needs.

As solar technologies have improved and costs have declined, solar energy has become one of the fastest-growing energy sources worldwide. In the United States, solar power accounted for approximately 45% of all new electricity-generating capacity added in 2025, more than any other energy source.

Solar energy can be used in several ways. **Passive solar technologies** use building design to capture, store, and distribute solar energy without mechanical equipment. **Active solar technologies** use pumps, fans, or other devices to collect and transfer solar energy. Solar energy may also be converted directly into electricity using **photovoltaic (PV) systems** or captured as heat using **solar thermal systems**. Together, these technologies allow solar energy to be used for lighting, heating, water heating, and electricity generation at scales ranging from individual homes to large utility-scale power plants.

The following sections examine the major types of solar energy technologies, their benefits, and their limitations.

Passive Solar

Solar technologies can be classified as either **passive** or **active**. **Passive solar technologies** use building design to capture, store, and distribute solar energy without the use of pumps, fans, or other mechanical devices. Because they rely on natural processes, passive solar systems are often simple, inexpensive, and energy efficient.

One of the simplest forms of passive solar energy is the use of natural daylight. Windows, skylights, and other building features can provide illumination during the day, reducing the need for electric lighting (figure $7.3.3.a$). Buildings can also be designed to maximize passive solar heating by placing windows where they receive the most winter sunlight and using materials such as brick, stone, or concrete to absorb heat during the day and release it slowly at night.

Another passive solar technology is the **solar tube** (or tubular skylight), which channels sunlight from the roof into interior spaces through a highly reflective tube. Solar tubes can illuminate rooms that receive little direct sunlight and are often installed similarly to conventional light fixtures (figure $7.3.3.b$).

Passive solar design can reduce energy consumption for lighting, heating, and cooling while requiring little maintenance and producing no emissions during operation. As a result, passive solar strategies are often incorporated into sustainable building design.




Figure 7.3.3.*a*: Passive solar technologies use sunlight without mechanical equipment. Examples include windows and skylights that provide natural lighting and solar tubes that channel sunlight from rooftops into interior spaces. Careful building design can also use passive solar heating to reduce energy needs. Left image by U.S. Department of Energy(opens in new window) (public domain), and right image by Energy Star(opens in new window) (public domain).

Passive solar heating uses building design to capture and store solar energy. South-facing windows allow sunlight to enter during winter, while materials such as concrete, brick, or tile absorb and slowly release heat. Roof overhangs and other shading features help keep buildings cooler during summer (figure 7.3.3.*b*).




Figure 7.3.3.*b*: This building design uses the sun for heating during the winter and is thus a passive solar technology. The roof overhang serves as a control, blocking the higher summer sun while allowing the lower winter sunlight to pass through the window (aperture). Heat energy from the sun is then distributed in the building. Thermal mass (such as a tile or stone floor) absorbs and later releases heat energy. Image by U.S. Department of Energy(opens in new window) (public domain).

Solar Water Heaters

Solar water heaters use sunlight to heat water for household or commercial use. In their simplest form, they consist of dark-colored tubes or collectors that absorb solar energy and transfer the heat to water flowing through the system (figure 7.3.3.*c*). Some designs use natural convection to circulate water and are considered **passive solar technologies**. More commonly, solar water heaters use pumps and controls to move water or a heat-transfer fluid between the solar collector and a storage tank, making them **active solar technologies**.

Because heating water accounts for a substantial portion of household energy use, solar water heaters can reduce electricity or natural gas consumption while producing little environmental impact during operation.



Figure 7.3.3.*c*: A solar water heater captures the Sun's energy to heat water for household use. Depending on the design, water may circulate naturally (passive solar) or be moved by pumps (active solar). Solar water heating is one of the most direct and efficient uses of solar energy. Image by Pixabay(opens in new window) [Vijayanarasimha] (public domain).

Why aren't solar water heaters more common in the United States?
Although solar water heaters are an efficient use of solar energy, they must compete with relatively inexpensive natural gas, electric water heaters, and modern heat-pump water heaters. In recent decades, falling costs of photovoltaic (PV) panels have also shifted investment toward solar electricity generation rather than solar water heating.

Active Solar Technologies

Photovoltaics (PV)

Active Solar Technologies: Photovoltaics (PV)

Active solar technologies use mechanical or electrical components to collect, convert, or distribute solar energy. One of the most common active solar technologies is the **photovoltaic (PV) system**, which converts sunlight directly into electricity using solar panels (figure$7.3.3.d$).

Solar panels are made up of many **photovoltaic cells** (figure$7.3.3.e$). These cells contain **semiconductors**, materials that conduct electricity only under certain conditions. When sunlight strikes a PV cell, energy from the light excites electrons within the semiconductor. The movement of these electrons creates an electric current that can be used to power homes, businesses, and other electrical devices.

PV systems can be installed on rooftops, integrated into buildings, or arranged in large solar farms that generate electricity for the electrical grid. Because they produce electricity without burning fuel, photovoltaic systems generate very little air pollution and emit very little greenhouse gas over their lifetimes compared with fossil fuel power plants.



Figure $7.3.3.d$: Rooftop solar installation on Douglas Hall at the University of Illinois at Chicago has no effect on land resources, while producing electricity with zero emissions. Source: Office of Sustainability, UIC




Figure 7.3.3.*e*: Diagram of a **photovoltaic cell** consisting of two layers of semiconductors. The top (n-type) semiconductor has extra electrons and the bottom (p-type) semiconductor has extra spots (holes) for electrons. A junction divides these two layers. Light frees the electrons and allows them to move through a wire from the n-type to fill "electron holes" in the p-type semiconductor. The movement of electrons through the wire results in an electrical current. Image by U.S. Energy Information Administration(opens in new window) (public domain).

This video explains how the photovoltaic cells within solar panels generate electricity.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Solar Thermal Energy

Another example of an **active solar technology** is **solar thermal energy**, which uses sunlight to produce heat rather than directly generating electricity with photovoltaic cells. In many solar thermal power plants, large arrays of mirrors focus sunlight onto a central receiver or a series of pipes containing a heat-transfer fluid (figure7.3.3.*f*). The concentrated solar energy produces high temperatures that are used to heat water and generate steam. The steam then turns a turbine connected to a generator, producing electricity.

Like coal-, natural gas-, and nuclear-powered plants, solar thermal power plants generate electricity by using steam to spin a turbine. The difference is that the heat comes from concentrated sunlight rather than from combustion or nuclear fission. Because sunlight is only available during the day, some solar thermal facilities store heat in materials such as molten salt, allowing electricity generation to continue after sunset.

This approach is often called **concentrated solar power (CSP)** and is most effective in regions that receive abundant direct sunlight, such as deserts of the southwestern United States, northern Africa, and parts of Australia.




Figure 7.3.3.f: In solar thermal technology, (1) mirrors or reflectors concentrate the sun's rays to heat a special kind of liquid, which absorbs heat. (2) The heat from this liquid boils water to create steam, facilitated by the heat exchanger. (3) Steam spins a turbine that is connected to a generator, which creates electricity. (4) The steam cools and condenses back to water, which is recycled, reheated, and converted into steam again. Image and caption (modified) from EPA(opens in new window) (public domain).

Advantages and Disadvantages of Solar Energy

Solar energy is abundant, widely available, and produces very little air pollution or greenhouse gas emissions during operation. Although manufacturing solar panels requires energy and materials, the lifecycle greenhouse gas emissions associated with solar power are much lower than those of fossil fuels. The rapid growth of the solar industry has also created jobs in manufacturing, installation, maintenance, and research.

Solar energy can be deployed at a variety of scales, from small rooftop systems on homes and businesses to large utility-scale solar farms. Rooftop solar panels can generate electricity close to where it is used, reducing the need for additional transmission infrastructure. Solar technologies can also be integrated into buildings, parking structures, and other developed areas.

Like wind energy, however, solar energy is **intermittent**. Solar panels generate electricity only when sunlight is available, and electricity production varies with weather, season, and time of day. Energy storage systems, such as batteries, can help address this challenge, but they add cost and require additional materials. In addition, some regions receive less direct sunlight and are therefore less suitable for large-scale solar energy production.

The environmental impacts of solar energy depend partly on where solar facilities are located. Installing solar panels on rooftops, parking lots, and other already-developed areas generally minimizes habitat disturbance. In contrast, large solar farms may require substantial land areas and can affect desert ecosystems, wildlife habitat, or agricultural land if poorly sited.

Solar energy technologies also require raw materials, manufacturing, and disposal. The production of photovoltaic cells involves chemicals, metals, and energy-intensive processes that can generate waste and pollution if not properly managed. Some solar thermal facilities require water for cleaning mirrors and cooling equipment, which can be a concern in arid regions where water is limited.

Like all energy sources, solar power involves trade-offs. Compared to fossil fuels, solar energy generally produces less air pollution, fewer greenhouse gas emissions, and fewer impacts on human health. However, solar technologies still require land, materials, water, and infrastructure. Evaluating the sustainability of solar energy requires considering both its benefits and its environmental and social costs.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Renewable Energy(opens in new window) and Challenges and Impacts of Energy Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 7.3.3: Solar Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 19.3: Solar Energy by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.4: Hydropower

Hydropower

Hydropower (hydroelectric energy) is electricity generated from the movement of water. Like wind energy, hydropower is ultimately powered by the Sun. Solar energy drives the water cycle, causing water to evaporate from Earth's surface, condense in the atmosphere, and return as precipitation. Water stored at higher elevations then flows downhill under the force of gravity.

As moving water passes through a dam, river, or other hydropower system, it spins a turbine (figure $7.3.4.a$). The rotating turbine turns a generator, which converts the kinetic energy of moving water into electricity.

Hydropower generates electricity using the same basic principle as many other energy technologies, including wind turbines, coal- and natural-gas-fired power plants, nuclear power plants, and concentrated solar thermal facilities. In each case, a turbine spins a generator to produce electricity. The primary difference is the source of energy that turns the turbine.




Figure $7.3.4.a$: Top: A water turbine. Moving water turns the blades of the turbine. The wicket gate controls the water flow. The moving water causes magnets in the rotor to spin and move electrons within the loops of wire in the stator, generating an electric current. Image by U.S. Army Corps of Engineers (public domain). Bottom: A propeller-type runner rated 28,000 hp (21 MW). Wtshymanski - own work by Wtshymanski http://en.wikipedia.org/wiki/Image:H...bineRunner.png (public domain).

Types of Hydropower

There are several ways to generate electricity from moving water, but all hydropower technologies rely on the same basic principle: flowing water spins a turbine connected to a generator.

The most familiar form of hydropower uses **dams**. A dam blocks a river, creating a reservoir that stores water at a higher elevation. When electricity is needed, water is released through a large pipe called a **penstock**, where its movement spins a turbine and generates electricity (figure $7.3.4.b$). Because the flow of water can be controlled, dam-based hydropower can provide electricity on demand and help stabilize electrical grids.

Another approach is **run-of-the-river hydropower**, in which turbines are placed directly in a river or in a channel that diverts part of the river's flow (figure $7.3.4.c$). Because these systems do not require large reservoirs, they generally have smaller environmental impacts than conventional dams. However, electricity production depends on natural river flow and may vary seasonally.

Other forms of hydropower harness energy from oceans. **Tidal energy** captures energy from the rise and fall of ocean tides or from tidal currents, while **wave energy** uses the movement of ocean waves to drive generators (figure $7.3.4.d$). Unlike most forms of hydropower, which are indirectly powered by solar energy through the water cycle, tidal energy is driven primarily by the gravitational attraction of the Moon and Sun.

Although these technologies differ in design and energy source, they all convert the kinetic energy of moving water into electricity.




Figure 7.3.4.*b*: A section of a hydroelectric dam. Water in the reservoir passes through the intake into the penstock, which is a narrow channel. The moving water turns a turbine. This powers the generator in the powerhouse. The water continues down the river. Meanwhile electricity is distributed through long distance power lines. Image by Tennessee Valley Authority(opens in new window) (public domain).




Figure 7.3.4.*c*: A microhydropower system. Although there are several ways to harness the moving water to produce energy, these run-of-the-river systems, which do not require large storage reservoirs, are often used for microhydropower projects. For run-of-the-river hydro projects, a portion of a river's water is diverted to a channel, pipeline, or pressurized pipeline (penstock) that delivers it to a waterwheel or turbine. In this image, water from the river moves through the intake then to the canal, forebay, penstock, and powerhouse, which contains the turbine. The moving water rotates the wheel or turbine, which spins a shaft. The motion of the shaft can be used for mechanical processes, such as pumping water, or it can be used to produce electricity. In this image, electrical lines connect the powerhouse to a nearby house.




Figure 7.3.4.*d*: One way to harness wave energy is with an absorber. Abosorbers extract energy from the rise and fall of the waves with a buoy. The buoy consists of the float, which is above the water and a heavy plate. The narrow spar connects the two. A cable connects the heavy plate of each buoy to an undersea substation, and another cable runs from the substation to the shore. Image and caption (modified) by OpenEI (opens in new window)(public domain).

Advantages and Disadvantages of Hydropower

Hydropower is one of the oldest and most widely used sources of renewable electricity. Because hydropower plants generate electricity without burning fuel, they produce little air pollution and low greenhouse gas emissions during operation. Reservoirs created by dams can also provide additional benefits, including **flood control**, **water storage**, **irrigation**, and **recreational opportunities** such as boating and fishing (figure 7.3.4.*e*.

Hydropower is generally a reliable source of electricity because water can be stored in reservoirs and released when energy is needed. However, electricity generation depends on water availability. During droughts, reservoir levels may decline, reducing the amount of electricity that can be produced. Hydropower is therefore most effective in regions with abundant rainfall, snowmelt, or mountainous terrain.

Despite these benefits, hydropower can have significant environmental impacts. Dams alter river ecosystems by changing water flow, depth, temperature, sediment transport, and water chemistry. These changes can affect aquatic habitats both upstream and downstream of a dam.

One of the most widely recognized impacts is the disruption of fish migration. Many fish species, including salmon, migrate upstream to reproduce. Dams can block these migrations and reduce access to spawning habitat. To address this problem, some dams include **fish ladders**, which provide routes for fish to move around barriers (figure 7.3.4.*f*. However, fish traveling downstream may still be injured or killed as they pass through turbines.

Reservoirs can also flood forests, wetlands, farmland, communities, and culturally significant sites. In some cases, dam construction has displaced human populations and altered local economies. When previously dry land is flooded, submerged vegetation decomposes under low-oxygen conditions, releasing **methane**, a potent greenhouse gas. As a result, although hydropower produces relatively low greenhouse gas emissions during operation, reservoirs can contribute to climate change, particularly in warm regions where decomposition rates are high.

Not all hydropower projects have the same impacts. Run-of-the-river systems typically cause less habitat flooding and ecosystem disruption than large dams because they use the natural flow of a river rather than storing large volumes of water in a reservoir.

Like all energy sources, hydropower involves trade-offs. It can provide reliable renewable electricity and valuable water-management benefits, but it can also alter ecosystems, affect biodiversity, and transform landscapes. Evaluating the sustainability of hydropower requires weighing these benefits and costs in the context of local environmental and societal needs.



Figure 7.3.4.f: The Grand Coulee Dam on the Columbia River in Washington state created Lake Roosevelt, which provides outdoor recreation such as fishing and swimming. However, it compromised salmon populations and flooded sacred sites for people of the Spokane Tribe. Image by U.S. Bureau of Reclamation (public domain).




Figure 7.3.4.f: Left: The fish ladder at John Day Lock and Dam along the Columbia River in Oregon allows adult fish to migrate upstream of the dam on their own. Right: Diagram of a fish ladder, which allows the passage of salmon over a dam. Salmon swim upstream to spawn (reproduce) and are able to swim up the steps of the ladder. The flow from the ladder attracts migrating salmon. At the turning pool, they rotate and then proceed up the next flight of the ladder until they are over the dam. Left image and caption (modified) from U.S. Army Corps of Engineers, and right image by NOAA Fisheries (public domain).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Renewable Energy(opens in new window) and Challenges and Impacts of Energy Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 7.3.4: Hydropower is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.5: Hydropower by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.5: Geothermal Energy

Geothermal Energy

Geothermal energy originates from heat within the Earth. This heat comes from two primary sources: thermal energy remaining from the Earth's formation and the continuous radioactive decay of isotopes such as uranium, thorium, and potassium within Earth's crust and mantle.

Geothermal Power Plants

Geothermal power plants use heat from Earth's interior to generate electricity (figure $7.3.5.a - c$). In some locations, naturally occurring hot water or steam can be brought to the surface through wells. In others, water is pumped underground, where it is heated by hot rocks and then returned to the surface. The resulting steam, or heat transferred to another fluid with a lower boiling point, is used to spin a turbine connected to a generator.

Like coal-, natural gas-, nuclear-, and solar thermal-powered plants, geothermal power plants generate electricity by using heat to produce a high-pressure fluid that turns a turbine. The difference is that the heat source is Earth's internal energy rather than combustion, nuclear fission, or concentrated sunlight.



Figure $7.3.5.a$: Steam is released from a geothermal power plant. Image by Open Access Government(opens in new window) (CC-BY(opens in new window)).




Figure 7.3.5.*b*: In a geothermal power plant, geothermal fluid is injected underground. (Geothermal water refers to the hot water/steam used in this process.) It enhances the permeability of the rocks and is heated underground. It exits to the surface through production wells, where it is used to generate electricity in the geothermal power plant. Water vapor is released from the cooling facility of the power plant. To recycle the geothermal fluid, it is returned to the reservoir and injected back into the ground. Image from Office of Energy Efficiency and Renewable Energy(opens in new window) [Office of Energy Efficiency and Renewable Energy] (public domain).




Figure 7.3.5.*c*: (1) In a geothermal power plant, hot water is pumped from deep underground through a well under high pressure. (2) When the water reaches the surface, the pressure is dropped, which causes the water to turn into steam. (3) The steam spins a turbine, which is connected to a generator that produces electricity. (4) The steam cools off in a cooling tower and condenses back to water. (5) The cooled water is pumped back into the Earth to begin the process again. Image and caption (modified) from EPA(opens in new window) (public domain).

Geothermal Heat Pumps

Geothermal heat pumps (**ground-source heat pumps**) rely on cool temperatures underground to cool or heat homes (figure 7.3.5.*d*). They are sometimes considered a second type of geothermal energy, but they are also a means of energy conservation. Geothermal heat pumps use a heat-exchange system that runs in the subsurface about 20 feet (5 meters) below the surface, which is consistently cool (around 55°F, or 12.5° C). Fluid is pumped underground and then along ducts in the home. This cools the house during the summer, acting as a heat sink. During a cold winter, it warms the house to 55° F (acting as a heat source), and traditional heating systems do the rest. This greatly reduces the energy consumption required to generate heat from gas, steam, hot water, and conventional electric air-conditioning systems.




Figure 7.3.5.*d*: A ground source heat pump (geothermal heat pump) in heating mode (during the winter; left) and in cooling mode (during the summer; right). There are four steps for heating mode (left). (1) Circulation: The above-ground heat pump moves water or another fluid through a series of buried pipes or ground loops. (2) Heat absorption: As the fluid passes through the ground loop, it absorbs heat from the warmer soil, rock, or ground water around it. (3) Heat exchange and use: The heated fluid returns to the building where it used for space or water heating. The system uses a heat exchanger to transfer heat into the building's existing air handling, distribution, and ventilation system. (4) Recirculation: Once the fluid transfers its heat to the building, it returns at a lower temperature to the ground loop to be heated again. This process is repeated, moving heat from one point to another. The four steps for cooling mode (right) are similar: (1) heat exchange and absorption, (2) circulation, (3) heat discharge, and (4) recirculation. Images and caption (modified) from EPA(opens in new window) (public domain).

This video explains the construction and mechanism of geothermal heat pumps.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

A Note about Types of Heat Pumps

Geothermal heat pumps transfer heat to and from the ground, where temperatures remain relatively constant year-round, making them highly efficient but more expensive to install due to the need for underground piping. As a result, these are typically installed during new construction rather than retrofitted. In contrast, air-source heat pumps (ASHPs) are not geothermal, and instead function by exchanging heat with the outside air. This is generally easier and cheaper to access. In California, air-source heat pumps are more widely used because the state's generally mild climate allows them to operate efficiently without the higher upfront cost and installation complexity of geothermal systems. You've likely seen them on homes and apartment buildings, since they are a common energy-efficient upgrade for both new construction and retrofits, including electrification (converting a home to run entirely on electricity) (figure $7.3.5.e$). One key advantage is that a single electric heat pump can replace both a traditional air conditioner and a gas furnace, providing an efficient all-in-one heating and cooling system.



Figure $7.3.5.e$ Electric (air-source) heat pump on a balcony. Modified from image by Wikideas1(opens in new window) - Own work, CC0(opens in new window), wikimedia(opens in new window) [commons.wikimedia.org]

Advantages and Disadvantages of Geothermal Energy

Geothermal energy has several advantages. Unlike solar and wind energy, which depend on weather conditions, geothermal energy is available continuously because heat is constantly flowing from Earth's interior. As a result, geothermal power plants can provide reliable **baseload electricity**, operating day and night throughout the year. Geothermal energy can also be used directly for heating and cooling buildings through **geothermal heat pumps**, which take advantage of the relatively stable temperatures found underground. Because shallow subsurface temperatures remain fairly constant year-round, geothermal heat pumps can be used in many regions.

Geothermal power plants produce very little air pollution and low greenhouse gas emissions compared to fossil fuel power plants. Most geothermal facilities recycle water and steam by injecting them back underground after use, helping to maintain the geothermal reservoir and reduce water consumption.

Despite these advantages, geothermal electricity generation is not practical everywhere. Conventional geothermal power plants require access to hot rocks, hot water, or steam located relatively close to Earth's surface. These conditions are most common near volcanic regions, hot springs, geysers, and tectonic plate boundaries (figure $7.3.5.f$). Although newer technologies such as **enhanced geothermal systems (EGS)** may expand the geographic range of geothermal energy in the future, suitable locations remain limited compared with solar and wind energy.

Geothermal facilities can also be expensive to construct because drilling deep wells and installing specialized equipment require

substantial upfront investment. In some cases, geothermal drilling and fluid injection have triggered small earthquakes, a phenomenon known as **induced seismicity**. Geothermal fluids may also contain naturally occurring substances such as hydrogen sulfide, dissolved minerals, and trace metals that must be carefully managed to prevent environmental contamination.

Like all energy sources, geothermal energy involves trade-offs. It provides reliable, low-carbon energy with relatively small land requirements, but its availability is geographically constrained and its development can affect local geology and water resources.



Figure 7.3.5.f: Locations of the world's volcanoes. They are concentrated along the west coasts of North American and South America as well as just east of Asia and Australia. These regions, which wrap around the Pacific Ocean, form what is known as the Ring of Fire. Image by Salazar, S.S., Muñoz, Y. & Ospino, A. Analysis of geothermal energy as an alternative source for electricity in Colombia(opens in new window). *Geotherm Energy* **5,** 27 (2017). (CC-BY(opens in new window))

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Renewable Energy(opens in new window) and Challenges and Impacts of Energy Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window)

This page titled 7.3.5: Geothermal Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.4: Geothermal Energy by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.6: Biofuels (Biomass Energy)

Biofuels (Biomass Energy)

Biofuels, also called **biomass energy**, are fuels derived from recently living organisms such as plants, algae, animal waste, and organic municipal waste. Because the energy stored in biomass was originally captured from sunlight through photosynthesis, biomass is an indirect form of solar energy.

Biomass can be burned directly or converted into liquid or gaseous fuels. The heat released during combustion can be used for cooking, heating buildings, generating electricity, or powering vehicles. When biomass is used to generate electricity, the heat from combustion produces steam that turns a turbine connected to a generator, similar to many other forms of electricity generation.

Common forms of biomass energy include:

- Wood and charcoal
- Agricultural residues such as straw or corn stalks
- Municipal solid waste
- Landfill gas (biogas)
- Bioethanol
- Biodiesel

Carbon Neutrality and Biofuels

- Biofuels are often described as **carbon neutral** because the carbon dioxide released during combustion was recently removed from the atmosphere by photosynthesis (figure $7.3.6.a$). In contrast, fossil fuels release carbon that has been stored underground for millions of years.

  However, carbon neutrality is not always guaranteed. The overall climate impacts of biofuels depend on how the biomass is grown, processed, transported, and used. For example, clearing forests to grow biofuel crops can release large amounts of stored carbon and reduce biodiversity. As a result, scientists often evaluate biofuels using a life cycle assessment, which considers environmental impacts from production through disposal.




Figure 7.3.6.a: Carbon cycling in fossil fuels and biofuels. Fossil fuel combustion releases carbon that has been stored underground for millions of years, whereas biofuel combustion releases carbon that was recently absorbed from the atmosphere through photosynthesis. Image modified
from those created by Melissa Ha from NETL/DOE (public domain) and publicdomainvectors.org (public domain).

Advantages of Biomass Energy

Biomass energy offers several potential benefits.

- Biomass can be produced locally, improving energy security.
- Many biofuels are derived from waste materials that would otherwise decompose or be discarded.
- Biomass can provide energy when solar and wind resources are unavailable.
- Some forms of biomass can reduce methane emissions by capturing gases that would otherwise be released from landfills or manure storage systems.

Because biomass can often be stored and used when needed, it is generally more reliable than intermittent renewable energy sources such as solar and wind.

Disadvantages of Biomass Energy

Despite these benefits, biomass energy also has significant environmental and social challenges.

Unlike most renewable energy sources, **burning biomass releases air pollutants**, including particulate matter, carbon monoxide, and nitrogen oxides. Indoor air pollution from cooking and heating with wood or charcoal remains a major health concern in many low-income regions.

Biomass production also requires land, water, and nutrients. Growing crops for biofuel production can compete with food production and may contribute to higher food prices. In some regions, forests and natural habitats have been cleared to make room for biofuel plantations, reducing biodiversity and releasing carbon stored in ecosystems.

Like all energy sources, biomass involves trade-offs. Biomass can help reduce dependence on fossil fuels, but its sustainability depends heavily on where the biomass comes from and how it is managed. In general, biomass derived from waste materials is considered more sustainable than biomass grown specifically for energy production.

Examples of Biomass Energy

Wood and Charcoal

Wood and charcoal have been used for cooking and heating for thousands of years (figure $7.3.6.b$). When harvested sustainably from managed forests or waste wood, they can provide a renewable source of energy. However, unsustainable harvesting can contribute to deforestation, habitat loss, and soil degradation.



Figure $7.3.6.b$: Charcoal production in Uganda. (A) Trees cut down for charcoal. (B) Pieces of firewood. (C) Wood logs read for charcoal burning. (D) Charcoal packed and ready for market. Image and caption (modified) from Bamwesigye et al. Sustainability 2020, 12(20), 8337. (CC-BY)

Municipal Solid Waste

Municipal solid waste (MSW) can be burned to generate electricity or heat. Waste-to-energy facilities reduce the volume of waste sent to landfills while producing useful energy. However, they can also generate air pollution and ash that must be carefully managed.

Landfill Gas (Biogas)

As organic waste decomposes in landfills, sewage treatment plants, and manure storage systems, microorganisms produce methane-rich **biogas**. Capturing and burning this gas generates electricity while reducing methane emissions, a potent greenhouse gas that would otherwise enter the atmosphere.

Bioethanol and Biodiesel

Bioethanol is typically produced by fermenting sugars or starches from crops such as sugarcane and corn. **Biodiesel** is commonly produced from vegetable oils, animal fats, or used cooking oil. These fuels can replace some petroleum-based fuels in vehicles, but their environmental benefits depend on how feedstocks are produced and processed.

Local examples of biogas production and usage include Miramar Landfill and Point Loma Wastewater Treatment Plant's biodigesters (figure 7.3.6.*c*).




Figure 7.3.6.*c*: The collection and processing of landfill gas to produce methane for multiple uses. First, landfill gas is collected from the landfill gas well. A system of vertical and horizontal piping buried in an municipal solid waste landfill. The landfill gas is then processed and treated for use (blower/flare/treatment). Potential end uses include industrial/institutional uses, arts and crafts, pipeline gas, and vehicle fuel. Image and caption (modified) from Landfill Methane Outreach Program/EPA (public domain).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Renewable Energy(opens in new window) and Challenges and Impacts of Energy Use from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 7.3.6: Biofuels (Biomass Energy) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.6: Biofuels (Biomass Energy) by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.7: Energy Efficiency and Conservation

Energy Efficiency and Conservation

Energy conservation refers to reducing energy use through behavioral changes, improved efficiency, and the elimination of unnecessary energy waste. **Energy efficiency** means using less energy to perform the same task or achieve the same result, thereby reducing energy consumption, costs, and environmental impacts. Examples include energy-efficient appliances, LED lighting, electric vehicles, improved insulation, and smart thermostats. Building design can also conserve energy through features such as large windows for natural lighting, passive solar heating, natural ventilation, green roofs, and strategic shading that reduce the need for artificial heating, cooling, and lighting (figure $7.3.7.a$). Because the cleanest and least expensive energy is often the energy that is never used, energy conservation and efficiency are considered among the most effective strategies for reducing environmental impacts and greenhouse gas emissions.



Figure $7.3.7.a$): The Solar Settlement with the Sun Ship in the background: two PlusEnergy projects in Freiburg, Germany.The houses contain 2-3 floors and were built with ecological building materials via wooden post-and-beam construction from regional forests and prefabricated individual modules, PVC-free, and environmentally friendly insulation materials. The houses are designed for high energy efficiency, with south-facing orientation to maximize solar energy use. They follow passive house standards, including strong insulation and high-performance glazing, resulting in very low heating needs and low annual energy costs. Each home uses a ventilation system with heat recovery and is connected to a local heating network. The roofs are equipped with photovoltaic panels, creating a large solar power system for the entire site. Andrewglaser at English Wikipedia. CC BY-SA 3.0

Advances in energy efficiency have helped reduce per-capita energy consumption in the United States. As a result, Americans used less energy per person in 2024 than in 1968, even though modern lifestyles rely heavily on computers, electronics, and other technologies that consume energy (figure $7.3.7.b$).

Explore the interactive graph below showing primary energy consumption per person. Compare the United States, the global average, and one other country by selecting countries from the menu on the right. How do energy-use patterns differ among these regions? What economic, geographic, technological, or social factors might contribute to these differences?

Interactive Element

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure 7.3.7.*b*: Energy use per person interactive (measured in kilowatt-hours per person. **Data source:** U.S. Energy Information Administration (2026); Energy Institute - Statistical Review of World Energy(2025); Population based on various sources (2024) – Our World in Data(opens in new window) [OurWorldinData.org] | CC BY(opens in new window)

Personal Energy Conservation

Personal energy conservation can be achieved through both simple daily habits and larger investments. Low- or no-cost actions include turning off lights and electronics when not in use, unplugging devices that draw standby power, lowering water-heater temperatures, driving efficiently, and using passive solar heating and lighting, such as opening south-facing blinds during winter days. Other strategies require an upfront investment but can pay for themselves over time through lower energy bills. An **energy audit** can help identify where a home is losing energy and which improvements will provide the greatest benefits. Common upgrades include sealing air leaks, adding insulation, installing Energy Star–certified appliances, upgrading to energy-efficient windows, using smart thermostats, switching to LED lighting, and replacing older heating and cooling systems with air-source heat pumps (figure 7.3.7.$c-e$). Together, these measures can substantially reduce energy consumption, save money, and lower greenhouse gas emissions.



Figure 7.3.7.*c*: Energy Star appliances use energy efficiently. Image by MoneyBlogNewz (CC-BY)

This video demonstrates how to conduct an energy audit and explains how the results can be used to reduce energy consumption, lower utility bills, and improve home comfort.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Melissa Ha (CC-BY-NC) and Home Energy Audits. U.S. Department of Energy. Accessed 01-18-2021. (public domain)

This page titled 7.3.7: Energy Efficiency and Conservation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.7: Energy Conservation by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

7.3.8: Data Dive- Global Renewable Energy Generation

Exploring Global Renewable Electricity Generation

Our World in Data (OWID) is a scientific online publication that uses research and data to help explain global challenges such as poverty, disease, hunger, climate change, and sustainability. The organization is known for its data visualizations, including interactive graphs and maps that allow users to explore trends over time. The interactive graph below shows how electricity generation from different renewable energy sources has changed around the world.

Interactive Element

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure 7.3.8.a: Global renewable energy generation. Graph by Our World in Data (CC-BY).

Questions

1. Which renewable energy source has provided the most electricity globally for the longest period of time? What evidence from the graph supports your answer?
2. Press the play button and observe changes from 1965 to the present. Describe how the global use of solar energy changes over time. During which decades does solar electricity generation increase most rapidly?
3. Select the United States and compare its renewable electricity mix to the global average. Which renewable energy sources contribute a larger share in the United States, and which contribute a smaller share?
4. Compare two countries of your choice. How do their renewable energy portfolios differ? What geographic, economic, or political factors might explain these differences?
5. Based on the trends shown in the graph, which renewable energy source do you predict will experience the greatest growth globally over the next 50 years? Explain your reasoning using evidence from the graph.

Raw Data For Above Graph(s)

Raw data table link for Our World In Data graph above. (CC-BY)

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Rachel Schleiger (CC-BY-NC)

This page titled 7.3.8: Data Dive- Global Renewable Energy Generation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 18.8: Data Dive- Global Renewable Energy Generation by Melissa Ha and Rachel Schleiger has no license indicated.

8: ATMOSPHERIC POLLUTION

Introduction

During the COVID-19 pandemic, many people in the United States became accustomed to wearing face masks to reduce the spread of disease. For most Americans, however, masks are not normally worn in daily life except during events such as wildfire smoke episodes, severe allergy seasons, or outbreaks of respiratory illnesses. In contrast, residents of some highly polluted cities around the world have long used masks as a regular form of protection from poor air quality. In cities that experience frequent smog and high levels of airborne pollutants, people may wear masks during their daily commute or while working outdoors to reduce exposure. Although air quality has improved in many regions because of environmental regulations and pollution-control efforts, air pollution remains a significant challenge in many parts of the world.



Figure 8.a: Mask garbage. Image by Pixabay(opens in new window) (Public Domain).

Air pollution is one of the most widespread environmental challenges facing the world today. Human activities such as transportation, fossil fuel-based energy production, industry, and agriculture release pollutants into the atmosphere that can affect human health, ecosystems, and Earth's climate. In this chapter, you will explore the sources and impacts of air pollution, examine how pollutants move through the atmosphere, and evaluate strategies that can improve air quality and protect environmental and public health.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Rachel Schleiger (CC-BY-NC(opens in new window)).

This page titled 8: Atmospheric Pollution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20: Pollution by Melissa Ha and Rachel Schleiger has no license indicated.

8.1: Earth’s Atmosphere

The Atmosphere

Solar Energy and Earth Systems

Nearly all life and environmental processes on Earth depend on energy from the Sun. The Sun is a medium-sized star that produces energy through nuclear fusion, a process that converts hydrogen into helium in its core. This energy travels through space as **solar radiation**, also called solar energy or insolation.

Although Earth receives only a tiny fraction of the Sun's total energy output, that energy powers many of the planet's natural systems. Solar energy drives weather patterns, ocean currents, and the water cycle. It also provides the energy needed for photosynthesis, which supports nearly all food webs on Earth. Without a continuous supply of solar energy, Earth would be too cold to support most forms of life.

The balance between incoming solar energy and outgoing heat helps maintain Earth's average surface temperature of about 15°C (59°F). Understanding how solar energy moves through the Earth system is essential for understanding weather, climate, ecosystems, and environmental change.

Composition of Earth's Atmosphere

Earth's atmosphere is a mixture of gases that surrounds and protects the planet. Dry air is composed of approximately 78% nitrogen (N_2), 21% oxygen (O_2), and about 1% argon (Ar). Several other gases, including carbon dioxide (CO_2), methane (CH_4), and ozone (O_3), are present in much smaller amounts but play important roles in Earth's climate and environmental systems. These are called **trace gases** (figure $8.1.a$).

Water vapor (H_2O) is also found in the atmosphere, but its concentration varies with location, temperature, and weather conditions. Humid regions contain more water vapor than dry regions, and warm air can hold more water vapor than cold air.

The atmosphere is not a fixed mixture of gases. Instead, gases are constantly exchanged among the atmosphere, biosphere, hydrosphere, and geosphere. For example, plants and animals use oxygen and release carbon dioxide during cellular respiration, while plants use carbon dioxide and release oxygen during photosynthesis. Water evaporates from oceans, lakes, and soil and enters the atmosphere as water vapor. Volcanoes can also release gases such as carbon dioxide and sulfur dioxide (SO_2). These exchanges help regulate Earth's atmosphere and support life on our planet.




Figure 8.1.a: Pie chart showing the gases that make up Earth's atmosphere (CC BY-SA 3.0).

Properties of the Atmosphere

Water Vapor

Water vapor (H_2O) is the gaseous form of water and the most variable component of Earth's atmosphere. Its concentration ranges from nearly 4% in warm, humid regions to almost zero in cold polar regions. Water vapor moves through the water cycle by changing between liquid, solid, and gaseous forms through processes such as evaporation, condensation, melting, and freezing.

Although it makes up only a small portion of the atmosphere, water vapor is Earth's most abundant greenhouse gas. It also stores and transfers energy during phase changes, helping drive weather systems.

Air Density and Air Pressure

Gravity pulls air molecules toward Earth's surface, making the air denser near the ground and less dense at higher elevations. Air pressure is the force exerted by the weight of the air above a location. Both air density and air pressure decrease with altitude.

Air Temperature

Air temperature measures the average motion of air molecules. Faster-moving molecules have higher temperatures, while slower-moving molecules have lower temperatures. In the lowest layer of the atmosphere, temperature generally decreases with altitude because Earth's surface absorbs solar energy and warms the air above it.

Temperature Scales

Temperature is commonly measured using Fahrenheit (°F), Celsius (°C), and Kelvin (K). Scientists typically use Celsius and Kelvin because these scales are used internationally and in scientific research. Water freezes at 0°C and boils at 100°C.

K = °C + 273.15

where **K** is temperature in kelvins and **°C** is temperature in degrees Celsius.

Case Study: Vog in Hawai'i (Volcanic Air Pollution)

Most air pollution is caused by human activities, but natural processes can also affect air quality. One example is Kīlauea, an active volcano on Hawai'i's Big Island. Kīlauea releases large amounts of sulfur dioxide (SO_2) into the atmosphere (figure $8.1.b$). When sulfur dioxide reacts with oxygen, water vapor, and sunlight, it forms a hazy mixture of gases and tiny particles known as volcanic smog, or "**vog**."

The fine particles and gases in vog can irritate the eyes, nose, throat, and lungs, especially in people with asthma or other respiratory conditions. Wind patterns can carry vog far from the volcano, affecting air quality across portions of the Hawaiian Islands. During periods of increased volcanic activity, public health officials may advise residents to limit outdoor activities and take other precautions to reduce exposure.

Vog demonstrates that air pollution can come from both natural and human sources. Although volcanic emissions are natural, they can still affect human health, ecosystems, and atmospheric chemistry. Understanding both natural and human sources of air pollution helps scientists better assess air quality and environmental risks.



Figure $8.1.b$:Water vapor and other gases escaping from the Kīlauea volcano at the Halema'uma'u vent (CC BY-SA 4.0).

Layers of the Atmosphere

Earth's atmosphere is divided into several layers based on how temperature changes with altitude. Each layer has unique characteristics that influence weather, climate, and life on Earth.

The **troposphere** is the lowest layer of the atmosphere and contains most of the atmosphere's mass, water vapor, and clouds. Nearly all weather occurs in this layer. Temperature generally decreases with altitude (figure $8.1.c$).

Above the troposphere is the **stratosphere**, which contains the ozone layer. Ozone absorbs harmful ultraviolet (UV) radiation from the Sun, causing temperatures to increase with altitude.




Figure 8.1.*c*:The vertical temperature structure of Earth's atmosphere (Public Domain)(opens in new window) in kilometers (left axis) and miles (right axis). The layers are labeled as "spheres" and the boundaries between the layers are labeled as "pauses". For example the troposphere is the lowest layer and the tropopause is the boundary between the troposphere and the stratosphere. The yellow line shows temperature in °C, and the cloud indicates where most of the weather occurs.

Above the stratosphere are the **mesosphere** and **thermosphere** (figure 8.1.*d*). These upper layers contain very little air and experience large temperature changes. The thermosphere is the region where auroras occur.




Figure 8.1.*d*:This figure illustrates the relative heights of different phenomena in the various vertical layers of the atmosphere. The Kármán Line at 62 miles (100 km) designates the boundary between Earth's atmosphere and outer space. Image by NOAA Satellites(opens in new window) (Public Domain)(opens in new window).

Environmental Science Connection

The atmosphere plays a critical role in supporting life on Earth. Atmospheric gases such as water vapor and carbon dioxide help regulate Earth's temperature by trapping heat, while the ozone layer protects organisms from harmful ultraviolet (UV) radiation. Changes in air density, pressure, and temperature influence weather patterns, climate, air quality, and the distribution of ecosystems. Natural processes, such as volcanic eruptions, can also alter atmospheric conditions and affect environmental and human health. Together, these atmospheric processes shape both natural systems and human societies.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from 1.2 Overview of Earth's Atmosphere and 1.6 Layers of the Atmosphere in *Meteorology and Climate Science* originally authored by Libretexts and is licensed CC BY-NC-SA 4.0.

This page titled 8.1: Earth's Atmosphere is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

8.2: Air Pollution

Air Pollution

Air pollution occurs when harmful gases, particles, or biological materials are present in the atmosphere at concentrations that can negatively affect human health, ecosystems, or materials. Air pollutants may be released directly into the atmosphere as **primary pollutants** or formed through chemical reactions in the atmosphere as **secondary pollutants**.

Air pollution is commonly categorized as **outdoor air pollution** or **indoor air pollution**. Outdoor air pollution originates from sources such as vehicles, power plants, industry, agriculture, and wildfires. Indoor air pollution comes from sources within buildings, including combustion of fuels, tobacco smoke, building materials, household products, mold, and pollen.

Air pollution contributes to respiratory and cardiovascular disease, damages ecosystems, reduces visibility, and can affect regional and global climate.

This video summarizes the causes, consequences, and solutions to air pollution.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Criteria Air Pollutants

Although the U.S. Environmental Protection Agency (EPA) regulates hundreds of air pollutants under the **Clean Air Act**, six particularly common and harmful pollutants are classified as **criteria air pollutants** and are monitored through national air quality standards. These pollutants are **ground-level ozone (O_3), particulate matter (PM), carbon monoxide (CO), nitrogen oxides (NO_x), sulfur dioxide (SO_2)**, and **lead (Pb)**. They can harm human health, damage ecosystems, reduce visibility, and contribute to environmental problems such as smog and acid deposition.

1. Ground-Level Ozone (O_3)

Ground-level ozone is a **secondary pollutant**, meaning it is not emitted directly into the atmosphere. Instead, it forms when **nitrogen oxides (NO_x)** and **volatile organic compounds (VOCs)** react in the presence of sunlight. Major sources of these pollutants include motor vehicles, power plants, industrial facilities, gasoline vapors, and chemical solvents.

Ground-level ozone can irritate the respiratory system, worsen asthma, reduce lung function, and damage vegetation. It should not be confused with the **stratospheric ozone layer**, which protects Earth from harmful ultraviolet radiation.

2. Particulate Matter (PM)

Particulate matter consists of tiny solid particles and liquid droplets suspended in the air. Sources include fossil fuel combustion, biomass burning, industrial processes, wildfires, and wind-blown dust.

Particle size influences health effects:

- **PM_{10}** particles are small enough to enter the respiratory system.

- **$PM_{2.5}$** particles are even smaller and can penetrate deep into the lungs and enter the bloodstream.

Exposure to particulate matter is associated with respiratory disease, cardiovascular disease, and premature death.

3. Carbon Monoxide (CO)

Carbon monoxide is a colorless, odorless gas produced by incomplete combustion of fuels. Major sources include motor vehicles, generators, furnaces, and wildfires.

Carbon monoxide reduces the blood's ability to transport oxygen. At high concentrations, it can cause headaches, dizziness, unconsciousness, and death.

4. Nitrogen Oxides (NO_x)

Nitrogen oxides are produced primarily by vehicles, power plants, and industrial combustion processes. Certain soil bacteria also release nitrogen oxides naturally.

Nitrogen oxides contribute to:

- Ground-level ozone formation
- Fine particulate pollution
- Acid deposition

They can also irritate and damage the respiratory system.

5. Sulfur Dioxide (SO_2)

Sulfur dioxide is released mainly through the combustion of sulfur-containing fossil fuels, particularly coal. Volcanoes are an important natural source.

Sulfur dioxide can impair lung function and contributes to the formation of **sulfuric acid**, a major component of acid deposition.

6. Lead (Pb)

Lead is a toxic metal that occurs naturally in the environment and is also released by human activities such as mining, metal processing, and the use of some aviation fuels. Historically, leaded gasoline was a major source of lead pollution, but regulations have dramatically reduced airborne lead concentrations in many countries.

Lead exposure can damage the nervous system, particularly in children, and may impair learning, behavior, and development.

Sources of Air Pollution

Air pollutants originate from a variety of anthropogenic and natural sources.

Stationary Sources

Stationary sources (point sources) remain in one location and include:

- Power plants
- Factories
- Refineries
- Dry-cleaning facilities

These facilities can release particulate matter, sulfur dioxide, nitrogen oxides, volatile organic compounds, and other pollutants.

Mobile Sources

Mobile sources include transportation systems that move under their own power, such as:

- Cars
- Trucks
- Buses
- Aircraft
- Trains
- Boats

Mobile sources are major contributors to carbon monoxide, nitrogen oxides, particulate matter, and volatile organic compounds.

Agricultural Sources

Agricultural activities release pollutants from livestock operations, fertilizer use, and pesticide application.

Examples include:

- Ammonia from manure
- Dust from fields
- Pesticide drift
- Emissions associated with fertilizer use

Natural Sources

Not all air pollution is caused directly by human activities. Although some of these such as wildfires and dust storms can be indirectly attributed to humans (figure $8.2.a$). So-called natural sources include:

- Wildfires
- Volcanic eruptions
- Dust storms
- Sea spray
- Plant emissions



Figure $8.2.a$: Smoke from the Camp fire in November of 2018 covers much of Northern California and extends over the Pacific Ocean. Image by NASA Earth Observatory/Lauren Dauphin(opens in new window) (public domain).

Indoor Air Pollution

People spend approximately 90% of their time indoors, and indoor air can sometimes be more polluted than outdoor air.

Common indoor air pollutants include:

- Environmental tobacco smoke
- Combustion products from cooking and heating
- Mold and mildew
- Dust and allergens
- Household chemicals
- Building materials and furnishings

Poor indoor air quality can contribute to headaches, irritation of the eyes and respiratory system, allergies, and other health problems.

Sick Building Syndrome

Sick building syndrome refers to situations in which building occupants experience health symptoms associated with spending time in a particular building. Contributing factors may include poor ventilation, indoor pollutants, and biological contaminants. Improved ventilation and reduction of pollutant sources often alleviate symptoms.

Secondhand Smoke

Secondhand smoke is the combination of smoke released from burning tobacco products and smoke exhaled by smokers. It contains thousands of chemicals, including hundreds known to be harmful and many known to cause cancer. Because there is no safe level of exposure, public health agencies recommend eliminating smoking in indoor environments.

Air Quality Index (AQI)

The **Air Quality Index (AQI)** is a tool used to communicate daily air quality conditions and associated health risks. The AQI combines measurements of major pollutants into a color-coded scale that ranges from good air quality to hazardous conditions.

Sensitive populations, including children, older adults, and individuals with asthma or other respiratory diseases, are often advised to limit outdoor activities when AQI levels are elevated.

> The Air Quality Index (AQI) and how to use it. AirNow.gov

Addressing Air Pollution

Reducing air pollution generally requires a combination of regulations, economic incentives, technological innovations, and individual actions.

Government Regulation

The **Clean Air Act of 1970** has significantly reduced levels of major air pollutants in the United States. Under this law, the EPA establishes air quality standards and regulates emissions from major sources.

Economic Incentives

Governments may use **green taxes**, **subsidies**, and **cap-and-trade systems** to encourage pollution reduction.

Examples include:

- Taxes on highly polluting activities
- Incentives for renewable energy
- Carbon trading programs that place limits on emissions

Technological Solutions

Many technologies reduce air pollution by preventing emissions or removing pollutants before they enter the atmosphere.

Examples include:

- Catalytic converters on internal gas combustion vehicles
- Air filters and ventilation systems
- Smokestack scrubbers on power plants
- Electrostatic precipitators that remove particulate matter
- Electric vehicles powered by low-emission electricity sources

Catalytic converters reduce harmful vehicle emissions by converting carbon monoxide and nitrogen oxides into less harmful substances. **Smokestack scrubbers** remove sulfur dioxide from industrial emissions, and **electrostatic precipitators** capture particulate matter before it is released. **Electrostatic precipitators** remove particulate matter from industrial emissions by electrically charging particles and collecting them on oppositely charged plates (figure $8.2.b$). **Electric vehicles** (EVs) can reduce air pollution, particularly when charged using electricity generated from renewable energy sources.

Like many environmental problems, reducing air pollution requires multiple approaches working together. Regulations, technological innovation, cleaner energy sources, and individual choices all contribute to cleaner air and healthier communities.




Figure $8.2.b$: An electrostatic precipitator. Blue arrows represent the flow of air past metal rods, which negatively charge the pollutants. These are then drawn to positive electrodes (red, flat metal plates), removing the pollutants form the air. Image by Evan Mason(opens in new window) (CC-BY-SA(opens in new window)).

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored

by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Atmospheric Pollution from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Basic Information about NO_2(opens in new window). Environmental Protection Agency. Accessed 11-04-2021 (public domain).

This page titled 8.2: Air Pollution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20.2: Air Pollution by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

8.3: Acid Deposition

Acid Deposition

Acid deposition refers to the transfer of acidic compounds from the atmosphere to Earth's surface. It includes both **wet deposition** and **dry deposition**. **Wet deposition** occurs when acidic compounds are incorporated into rain, snow, sleet, fog, or hail. **Dry deposition** occurs when acidic gases and particles settle directly onto surfaces such as soil, vegetation, buildings, or bodies of water.

The primary causes of acid deposition are emissions of **sulfur dioxide (SO_2)** and **nitrogen oxides (NO_X)**, which are released mainly through the combustion of fossil fuels in power plants, vehicles, and industrial facilities (figure $8.3.a$). Volcanoes and decaying organic matter also release small amounts of these compounds naturally. In the atmosphere, sulfur dioxide and nitrogen oxides react with water vapor, oxygen, and other chemicals to form **sulfuric acid (H_2SO_4)** and **nitric acid (HNO_3)**. Because these acids form in the atmosphere rather than being emitted directly, they are considered **secondary pollutants**.

Prevailing winds can transport sulfur dioxide and nitrogen oxides hundreds or even thousands of miles before they are converted into acids and deposited, allowing one region's emissions to affect distant ecosystems.




Figure $8.3.a$: Sulfur dioxide and nitrogen oxides released from natural and human sources can react in the atmosphere to form acidic compounds. These compounds return to Earth through wet deposition (rain, snow, fog, and other precipitation) or dry deposition (particles and gases settling on surfaces).

Acid Deposition and pH

The acidity of a solution is measured using the **pH scale** (figure $8.3.b$). The pH scale ranges from acidic to basic (alkaline) conditions and is logarithmic, meaning each whole-number change represents a tenfold change in hydrogen ion concentration.

- Lower pH values indicate greater acidity.
- Higher pH values indicate greater alkalinity.
- Pure water has a pH of 7.0.

Natural rainwater is slightly acidic because carbon dioxide in the atmosphere dissolves into water droplets, forming weak **carbonic acid**. As a result, normal rain typically has a pH of approximately **5.6**.

When sulfuric and nitric acids are present, precipitation becomes more acidic than normal. During the peak of acid deposition in the eastern United States, precipitation commonly had pH values below 5 and sometimes near 4.




Figure 8.3.b: The pH scale is used to measure acidity and alkalinity. Because the scale is logarithmic, each one-unit change in pH represents a tenfold change in hydrogen ion concentration. Source: [Epa.gov], public domain.

Effects of Acid Deposition

Acid deposition affects ecosystems, infrastructure, air quality, and human health.

Aquatic Ecosystems


The ecological effects of acid deposition are most evident in lakes, streams, and wetlands. Most freshwater ecosystems function best within a relatively narrow pH range. When acid deposition lowers the pH of lakes and streams, organisms may experience stress, reduced growth, reproductive failure, or death.

Some soils can neutralize acids through a process known as **buffering**, but soils with low buffering capacity are especially vulnerable. In these ecosystems, acidification can release aluminum from soils into lakes and streams. Dissolved aluminum is highly toxic to many aquatic organisms, particularly fish and aquatic insects.

Forests and Soils

Acid deposition can damage forests directly and indirectly. It may weaken trees by damaging leaves and needles and by removing important nutrients such as calcium and magnesium from soils. These stresses can make trees more vulnerable to drought, disease, insect outbreaks, and extreme temperatures.

High-elevation forests are often particularly sensitive because they are frequently exposed to acidic clouds and fog.






Figure 8.3.*c*: Top: pH of the precipitation in the US from 1985-1987. Acid deposition was most severe in the eastern United States during the 1970s and 1980s. Bottom: pH of US precipitation 2015-2017. Reductions in sulfur dioxide and nitrogen oxide emissions have increased precipitation pH in many regions, although precipitation remains somewhat more acidic than natural rain in some areas. Images by National Atmospheric Deposition Program(opens in new window) (public domain).

Buildings and Cultural Resources

Acid deposition accelerates the weathering of stone, metal, concrete, and painted surfaces. Buildings, bridges, monuments, statues, and historic structures made from limestone and marble are especially vulnerable because acidic compounds react with these materials and gradually dissolve them (figure 8.3.*d*).



Figure 8.3.d: Acid deposition can damage buildings, monuments, and sculptures by accelerating the weathering of stone and metal surfaces.

Visibility and Human Health

Sulfur dioxide and nitrogen oxides contribute to the formation of tiny sulfate and nitrate particles in the atmosphere. These particles scatter sunlight and contribute to **haze** and reduced visibility.

Fine sulfate and nitrate particles can also be inhaled deep into the lungs and may enter the bloodstream. Exposure is associated with respiratory and cardiovascular diseases, including asthma, bronchitis, heart disease, and premature mortality. Therefore, the pollutants responsible for acid deposition affect human health even before they are deposited from the atmosphere.

Reducing Acid Deposition

Acid deposition in North America and Europe has decreased substantially since the 1970s because regulations such as the **Clean Air Act** reduced emissions of sulfur dioxide and nitrogen oxides from power plants, industries, and vehicles. Technologies such as **smokestack scrubbers**, cleaner fuels, catalytic converters, and renewable energy have played an important role in lowering emissions and improving air quality.

In recent decades, some administrations and policymakers have proposed weakening or revising portions of environmental regulations, including certain Clean Air Act rules, while others have sought to strengthen them. These debates often focus on balancing economic costs with environmental and public health benefits.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Acid Rain from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 8.3: Acid Deposition is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20.2.2: Acid Deposition by Melissa Ha and Rachel Schleiger is licensed CC BY 4.0.

8.4: Ozone Depletion

Ozone Depletion

Ozone depletion is the thinning of the stratospheric ozone layer, a region of the upper atmosphere that absorbs most of the Sun's harmful ultraviolet (UV) radiation. The ozone layer acts as Earth's natural sunscreen, helping protect humans, animals, and plants from excessive UV exposure. During the late twentieth century, scientists discovered that certain human-produced chemicals, particularly chlorofluorocarbons (**CFCs**), were damaging ozone molecules and causing substantial reductions in ozone concentrations, especially over Antarctica.

It is important to distinguish ozone depletion from climate change. Although both are global environmental issues caused in part by human activities, they involve different atmospheric processes. Ozone depletion refers to the loss of ozone in the stratosphere and results in increased exposure to ultraviolet radiation. **Climate change** refers primarily to the warming of Earth's climate system caused by greenhouse gases such as carbon dioxide, methane, and nitrous oxide. Some chemicals, such as CFCs, contribute to both problems, but ozone depletion and climate change have different causes, consequences, and solutions.

The Process of Ozone Depletion

The ozone depletion process begins when **chlorofluorocarbons (CFCs)** and other **ozone-depleting substances (ODSs)** are released into the atmosphere.

Because these compounds are extremely stable, they do not break down quickly or dissolve readily in rain. Instead, they can persist in the atmosphere for many years and gradually rise into the **stratosphere**, the atmospheric layer that contains most of Earth's ozone (figure $8.4.a$). CFCs were widely used as refrigerants, aerosol propellants, solvents, and foam-blowing agents.




Figure $8.4.b$:

Distribution of ozone in Earth's atmosphere. Most atmospheric ozone is found in the stratosphere, where it forms the ozone layer that absorbs harmful ultraviolet (UV) radiation from the Sun. Ozone depletion refers to the thinning of this protective stratospheric ozone layer. In contrast,

tropospheric (ground-level) ozone

is a secondary air pollutant that forms from chemical reactions involving nitrogen oxides (NO_x) and volatile organic compounds (VOCs) in the presence of sunlight. Unlike stratospheric ozone, ground-level ozone contributes to smog and can harm human health, vegetation, and ecosystem. Image rastrized and improved by RedAndr - Own work using: Earth Observing System (EOS) Science Plan. Chapter 7. Ozone and Stratospheric Chemistry, Public Domain, [https://commons.wikimedia.org/w/inde...curid=2269127]

Ozone (O_3) in the stratosphere is continuously produced and destroyed through natural chemical reactions (figure $8.4.c$). Under normal conditions, the rate of ozone production is approximately equal to the rate of ozone destruction, so the total amount of ozone remains relatively stable. This balance can be compared to water flowing through a stream: although individual water molecules are constantly moving, the water level remains nearly constant.

The problem arises when ozone-depleting substances reach the stratosphere. There, **ultraviolet (UV) radiation** from the Sun breaks apart CFC molecules, releasing chlorine atoms. These chlorine atoms catalyze reactions that destroy ozone molecules. Because a single chlorine atom can destroy many ozone molecules before being removed from the atmosphere, ozone is destroyed faster than it can be naturally replaced. As a result, the concentration of stratospheric ozone decreases, leading to **ozone depletion**.




Figure 8.4.*c*: Ozone concentration and ultraviolet (UV) absorption at different altitudes. The stratospheric ozone layer absorbs all UVC radiation and most UVB radiation, preventing much of the Sun's harmful ultraviolet energy from reaching Earth's surface. In contrast, most UVA radiation is not absorbed and reaches the ground. Ozone depletion reduces this protective filtering effect, increasing UV exposure at Earth's surface. Image modified from NASA - [1], Public Domain, [https://commons.wikimedia.org/w/inde...urid=12761806]

Ozone (O_3) is continuously produced and destroyed in the stratosphere through a series of natural chemical reactions (figure 8.4.*d*). Under natural conditions, the rate of ozone production is approximately equal to the rate of ozone destruction, so the overall amount of ozone remains relatively stable. This balance is similar to the water level in a stream: although individual water molecules are constantly moving, the depth of the stream remains nearly unchanged. Likewise, ozone molecules are constantly being formed and broken apart, but the total amount of ozone in the ozone layer remains fairly constant. This natural balance persisted for thousands of years. However, during the twentieth century, increasing concentrations of ozone-depleting substances (ODSs) disrupted the cycle. ODSs such as **Chlorofluorocarbons (CFCs)** are human-made chemicals once commonly used in refrigeration, air conditioning, aerosol sprays, and industrial solvents. Chlorine and bromine released from these compounds accelerate ozone destruction, causing ozone to be removed faster than it can be naturally replenished. As a result, concentrations of stratospheric ozone decline, leading to ozone depletion.




Figure 8.4.*d*: The natural ozone cycle in the stratosphere. (1) Ultraviolet (UV) radiation from the Sun breaks apart oxygen molecules ($\mathbf{O_3}$), producing individual oxygen atoms. (2) These oxygen atoms rapidly combine with oxygen molecules to form ozone ($\mathbf{O_3}$). Ozone molecules also absorb incoming UV radiation and break apart, regenerating oxygen molecules and oxygen atoms. (3) Through this continuous cycle of ozone formation and destruction, much of the Sun's harmful UV radiation is absorbed and converted into heat, warming the stratosphere. Under natural conditions, ozone production and destruction remain in balance, maintaining a relatively stable ozone layer that protects life on Earth from excessive UV exposure. Image modified from Ozone_cycle.jpg: created by NASAderivative work: Smartse (talk) - Ozone_cycle.jpg, Public Domain, [https://commons.wikimedia.org/w/inde...curid=9096489]

Policies to Reduce Ozone Destruction

One success story in reducing pollutants that harm the atmosphere concerns ozone-destroying chemicals. In 1973, scientists calculated that CFCs could reach the stratosphere and break apart. This would release chlorine atoms, which would then destroy ozone. Based only on their calculations, the United States and most Scandinavian countries banned CFCs in spray cans in 1978. More confirmation that CFCs break down ozone was needed before more was done to reduce production of ozone-destroying chemicals. In 1985, members of the British Antarctic Survey reported that a 50% reduction in the ozone layer had been found over Antarctica in the previous three springs.

Two years after the British Antarctic Survey report, the “Montreal Protocol on Substances that Deplete the Ozone Layer” was ratified by nations all over the world. The **Montreal Protocol** controls the production and consumption of 96 chemicals that damage the ozone layer. CFCs have been mostly phased out since 1995, although they were used in low-income nations until 2010. Some of the less hazardous substances will not be phased out until 2030. The Protocol also requires that wealthier nations donate money to develop technologies that will replace these chemicals.

Because CFCs take many years to reach the stratosphere and can survive there a long time before they break down, the ozone hole did not immediately disappear after CFC emissions were reduced; however, it has been shrinking (figure 8.4.*e*).




Figure 8.4.e: Ozone levels stabilized in the 1990s following the Montreal Protocol, and have started to recover. They are projected to reach pre-1980 levels before 2075. Image and caption By NASA - [https://svs.gsfc.nasa.gov/30602, Public Domain, https://commons.wikimedia.org/w/inde...urid=61499809]

Watch the short video to learn how international action helped reduce ozone-depleting substances and contributed to the recovery of the ozone layer. As you watch, consider why the Montreal Protocol is often cited as one of the most successful international environmental agreements.

Health and Environmental Effects of Ozone Layer Depletion

There are three types of ultraviolet (UV) radiation emitted by the Sun: **UVA**, **UVB**, and **UVC**. Under normal conditions, the stratospheric ozone layer absorbs all UVC radiation and most UVB radiation, while most UVA radiation reaches Earth's surface. When stratospheric ozone levels decline, more **UVB radiation** reaches the ground. The amount of UVB emitted by the Sun does not change; rather, ozone depletion reduces the atmosphere's ability to filter out this harmful radiation. During periods of severe ozone depletion over Antarctica, UVB levels at the surface can increase substantially.

Increased exposure to UVB radiation can have significant impacts on human health. Studies have shown that UVB can damage DNA and increase the risk of skin cancer, including melanoma and non-melanoma skin cancers. UVB exposure is also associated with cataracts, a clouding of the eye's lens that can impair vision. Because some UVB reaches Earth's surface even when ozone levels are normal, protecting the skin and eyes from excessive sun exposure is always important. Ozone depletion increases these risks by allowing more UVB radiation to reach the surface.

UVB radiation can also affect other organisms and ecosystems. Because UVB damages cells and genetic material, it can reduce growth, reproduction, and survival in many species. **Microorganisms**, such as bacteria and phytoplankton, are particularly vulnerable because they consist of only one or a few cells and have limited protection from UV radiation. Increased UVB can also harm plants, reduce crop yields, and disrupt aquatic food webs by damaging phytoplankton, which form the base of many marine and freshwater ecosystems.

Thus, although ozone depletion occurs high in the stratosphere, its effects can extend throughout ecosystems and directly affect human health.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from Ozone Depletion from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in

new window))

This page titled 8.4: Ozone Depletion is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20.2.1: Ozone Depletion by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9: GLOBAL CLIMATE CHANGE

Chapter Hook: Rising Seas from Venice to San Diego

The iconic Grand Canal of Venice, Italy offers a visible reminder that Earth's climate is changing. For centuries, stone steps leading into the canal marked the boundary between land and water. Because the canal is directly connected to the surrounding sea, these steps also record changes in sea level. Today, water levels are significantly higher than when many of these structures were built, and flooding events have become increasingly common (figure $9.a$).

As global temperatures rise, melting glaciers and ice sheets, combined with the expansion of warming seawater, are causing sea levels around the world to rise.

Venice is not alone in facing this challenge. Coastal communities on every continent are preparing for the impacts of rising seas, including more frequent flooding, erosion, damage to infrastructure, and the loss of coastal habitats. Scientists project that sea level will continue to rise throughout this century, threatening many low-lying cities and coastal ecosystems.

Closer to home, San Diego faces many of the same risks. Depending on future greenhouse gas emissions and the rate of ice-sheet melting, sea level along the San Diego coastline could rise by approximately 3.6 to 7 feet (1.1 to 2.1 meters) by 2100.



Figure $9.a$: Flooding along the Grand Canal in Venice, Italy. Rising sea levels and more frequent high-tide flooding have increased challenges for this historic coastal city. Image by [WorldIslandInfo.com] in Wikimedia Commons (CC-BY2.0).




Figure $9.b$: Screenshot of mid-range sea-level rise estimate for San Diego by 2100. Sea level rise predictions for San Diego are expected to be from 3.6 feet (1.09 m) - 7 feet (2.13 m). Using the NOAA sea level rise and viewer, 1.8 m mid-range estimate for San Diego has been mapped onto the coastline (figure $9.b$). Image source: National Oceanic and Atmospheric Administration Sea level Rise Viewer. [https://coast.noaa.gov/slr/#/layer/s...tion/NOS_Minor]

Venice and San Diego are separated by more than 6,000 miles and exist in very different cultural and environmental settings. Yet both communities illustrate the same reality: climate change is not a distant problem affecting only the future. Its effects are already being measured today. Rising sea level is just one of many consequences of global climate change that scientists, policymakers, and communities must confront in the coming decades.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0 as well as from the following sources:

- City of San Diego. (n.d.). *Sea level rise*. Climate Resilient SD. San Diego.gov [www.sandiego.gov]
- National Oceanic and Atmospheric Administration. (n.d.). *Sea Level Rise Viewer*. NOAA Office for Coastal Management. NOAA.gov [coast.noaa.gov]

Modified by Melissa Ha and Rachel Schleiger from the following sources:

- Climate and the Effects of Global Climate Change from *General Biology* by OpenStax (licensed under CC-BY(opens in new window))
- Climate Change from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

9.1: Weather, Climate, and Climate Change

9.2: Earth's Energy Balance and Radiation

9.3: The Greenhouse Effect

9.4: Evidence and Causes of Climate Change

9.5: Impacts of Climate Change

9.6: Ocean Acidification- The Other Carbon Problem

9.7: Mitigating and Adapting to Climate Change

9.8: Data Dive- Flooding in Venice Italy

This page titled 9: Global Climate Change is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 21: Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY 4.0.

9.1: Weather, Climate, and Climate Change

Weather, Climate, and Climate Change

The terms **weather** and **climate** are often confused, but they describe different aspects of Earth's atmosphere.

Weather refers to atmospheric conditions at a specific place and time. Weather includes factors such as:

- **Air temperature**: how hot or cold the air is
- **Air pressure**: the force exerted by the atmosphere
- **Humidity**: the amount of water vapor in the air
- **Cloud cover**: the amount and type of clouds present
- **Precipitation**: rain, snow, sleet, or hail
- **Wind**: the movement of air caused by differences in air pressure
- **Visibility**: how far one can see through the atmosphere

Weather can change from hour to hour or day to day.

In contrast, **climate** refers to the long-term patterns and average conditions of weather in a region, typically measured over decades. Climate includes not only average temperature and precipitation but also the frequency of extreme events such as heat waves, droughts, storms, and cold spells.

A useful way to think about the difference is that **weather influences what you wear today, while climate influences the type of clothing you own**. For example, a resident of Hawai'i is unlikely to need a heavy winter coat, whereas someone living in Alaska probably does.

Climate Change and Global Warming

Climate change refers to long-term changes in Earth's climate, including shifts in temperature, precipitation patterns, wind patterns, and the frequency of extreme weather events. These changes occur over decades or longer.

Global warming refers specifically to the recent increase in Earth's average surface temperature. The primary cause of modern global warming is the accumulation of greenhouse gases released by human activities, particularly the burning of fossil fuels, deforestation, and some agricultural practices. For this reason, modern climate change is often referred to as **anthropogenic climate change**.

Global warming is one aspect of climate change, but climate change also includes related impacts such as rising sea levels, changes in precipitation patterns, stronger heat waves, and shifts in ecosystems.

A common misconception is that a single cold day or unusually cool week disproves climate change. Weather events occur over short time periods and naturally vary from day to day. Climate scientists instead examine long-term trends measured over many years or decades. Just as one hot day does not indicate climate change, one cold day does not disprove it.

This page titled 9.1: Weather, Climate, and Climate Change is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

9.2: Earth's Energy Balance and Radiation

Earth's Energy Balance and Radiation

To understand climate change, it is first necessary to understand how energy moves through the Earth system. Nearly all of Earth's energy comes from the Sun in the form of **electromagnetic radiation**. Some of this incoming solar energy is reflected back into space, while the rest is absorbed by the atmosphere, oceans, land, and living organisms. The absorbed energy warms Earth and is eventually re-emitted as infrared radiation. Earth's climate depends on the balance between incoming solar radiation and outgoing infrared radiation. Changes to this energy balance, particularly those caused by increasing greenhouse gas concentrations, are responsible for modern climate change.

Radiation

Radiation can be thought of in two ways: electromagnetic waves or as photons. For the purpose of atmospheric science, we will generally consider radiation as a wave rather than a photon particle. **Electromagnetic radiation** is a type of energy produced by electric and magnetic fields, taking a variety of names depending on the wavelength. For example, you've probably heard of radio waves and x-rays, whereby the primary difference is the wavelength of the waves in each. The next image shows the relationship between wavelength, frequency, and temperature. We will go into further detail later, but colder objects radiate at lower frequencies and longer wavelengths, and warmer objects radiate at higher frequency and shorter wavelengths. This relationship holds true across all scales, from atomic nuclei to planets, and across temperatures from near absolute zero to 10s of millions of Kelvins.

Electromagnetic Radiation

Electromagnetic radiation is energy that travels through space as waves. Different forms of electromagnetic radiation differ in their **wavelength** and **frequency**. Examples include radio waves, microwaves, infrared radiation, visible light, ultraviolet (UV) radiation, X-rays, and gamma rays (figure $9.2.a$).

For climate science, the most important types of electromagnetic radiation are:

- **Visible light**, which carries much of the Sun's incoming energy to Earth.
- **Infrared radiation**, which is emitted by Earth's surface and atmosphere as heat.

In general, hotter objects emit radiation at shorter wavelengths and higher frequencies, while cooler objects emit radiation at longer wavelengths and lower frequencies.




Figure $9.2.a$: The electromagnetic spectrum, showing the range of wavelengths and frequencies of electromagnetic radiation. (CC BY-SA 3.0).

The Sun and Earth Emit Different Types of Radiation

Although both the Sun and Earth emit electromagnetic radiation, they do so at different wavelengths because they have very different temperatures. The Sun's surface temperature is about 5,800 K, causing it to emit most of its energy as **shortwave radiation**, primarily visible light and smaller amounts of ultraviolet and infrared radiation. Earth is much cooler, with an average surface temperature of about 288 K (15°C). As a result, Earth emits most of its energy as **longwave infrared radiation**. This difference between incoming solar radiation and outgoing terrestrial radiation is fundamental to understanding climate change.

Earth's Energy Budget

Earth's temperature depends on a balance between incoming solar radiation and outgoing infrared radiation (figure $9.2.b$).

- Some incoming sunlight is reflected back into space by clouds, ice, snow, and other bright surfaces.
- The remaining solar energy is absorbed by the atmosphere, oceans, land, and living organisms.
- Absorbed energy warms Earth's surface.
- The warmed Earth emits infrared radiation back toward space.

When the amount of incoming energy equals the amount of outgoing energy, Earth's average temperature remains relatively stable.

This balance is known as **Earth's energy budget**.




Figure 9.2.b: Earth's energy balance and imbalance, showing where the excess energy goes: Outgoing radiation is decreasing owing to increasing greenhouse gases in the atmosphere, leading to Earth's energy imbalance of about 460 TW. The percentage going into each domain of the climate system is also indicated. Image by Kevin E Trenberth, (CC BY-SA 4.0) [https://creativecommons.org/licenses/by-sa/4.0], via Wikimedia Commons

Why Radiation Matters for Climate Change

Greenhouse gases such as carbon dioxide (CO_2), methane (CH_4), and water vapor absorb some of the infrared radiation emitted by Earth and re-radiate part of that energy back toward the surface. This process, known as the **greenhouse effect**, slows the loss of heat to space and keeps Earth warm enough to support life.

Human activities have increased the concentration of greenhouse gases in the atmosphere. As a result, more infrared radiation is trapped, altering Earth's energy budget and causing global temperatures to rise. The next section examines the greenhouse effect in greater detail.

This page was heavily revised and rewritten by Erin Rempala and is licensed CC BY-NC-SA 4.0.

This page titled 9.2: Earth's Energy Balance and Radiation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

9.3: The Greenhouse Effect

The Greenhouse Effect

Earth's temperature depends on a balance between incoming solar radiation and outgoing infrared radiation. Some of the Sun's energy is reflected back into space, while the rest is absorbed by Earth's atmosphere, oceans, land, and living organisms. The absorbed energy warms Earth's surface, which then emits infrared radiation back toward space. Earth's climate is largely determined by the balance between incoming and outgoing energy (figure $9.3.a$).




Figure $9.3.a$: Energy entering and leaving the Earth system. The balance between incoming solar radiation and outgoing infrared radiation influences Earth's average temperature. Modified from image by Efbrazil - Own work, CC BY-SA 4.0(opens in new window)

Many factors can influence Earth's energy balance, including:

- Changes in the greenhouse effect
- Variations in the amount of solar energy reaching Earth
- Changes in Earth's reflectivity (albedo)

The greenhouse effect is particularly important because it regulates the amount of heat retained by the atmosphere and is the primary driver of modern climate change.

Natural Greenhouse Effect

The **greenhouse effect** is a natural process that warms Earth's surface and atmosphere. Sunlight passes through the atmosphere and is absorbed by Earth's land and oceans. The warmed surface then emits energy back toward space as **infrared radiation** (heat).

Certain gases in the atmosphere absorb some of this infrared radiation and re-radiate it in all directions, including back toward Earth's surface. This slows the loss of heat to space and keeps the planet warmer than it would otherwise be.

Without the natural greenhouse effect, Earth's average surface temperature would be approximately **–18°C**, far too cold for most life as we know it. Instead, the average temperature is about **15°C**, making Earth habitable.

The greenhouse effect is often compared to the warming that occurs inside a greenhouse or a parked car on a sunny day. In both cases, energy enters more easily than it escapes, resulting in warmer temperatures.

Greenhouse Gases

The gases responsible for the greenhouse effect are known as **greenhouse gases (GHGs)**. These gases are largely transparent to visible light but absorb infrared radiation because of their molecular structure.

Important greenhouse gases include:

- **Water vapor (H_2O)**
- **Carbon dioxide (CO_2)**
- **Methane (CH_4)**
- **Nitrous oxide (N_2O)**
- **Ozone (O_3)**

These gases act like a blanket surrounding Earth, helping retain heat and maintaining temperatures suitable for life.

Enhanced Greenhouse Effect

Although the natural greenhouse effect is essential for life, human activities have strengthened it. Since the Industrial Revolution, the combustion of fossil fuels, deforestation, agriculture, and industrial activities have increased atmospheric concentrations of greenhouse gases. The resulting **enhanced greenhouse effect** causes more infrared radiation to be retained within the Earth system, increasing global temperatures (figure $9.3.b$).




Figure $9.3.b$: The enhanced greenhouse effect. Step 1: Some solar radiation reaches the Earth's surface, and some is reflected back into space. Step 2: The rest of the sun's energy is absorbed by the land and the oceans, heating the Earth. Step 3: Heat radiates from Earth towards space. Step 4: Some of the heat is trapped by greenhouse gases in the atmosphere, warming the Earth. Step 5: Human activities such as burning fossil fuels, agriculture, and land clearing have increased the concentrations of greenhouse gases in the atmosphere. Step 6: This is trapping extra heat, causing the Earth's temperature to rise.

Scientists have examined many possible causes of recent climate change, including changes in solar activity, volcanic eruptions, and natural climate cycles. Although these factors influence climate, they do not explain the rapid warming observed since the mid-twentieth century.

Climate models that include only natural influences fail to reproduce the observed warming trend. In contrast, models that include both natural influences and human-caused increases in greenhouse gases closely match observed temperatures (figure $9.3.c$).




Figure 9.3.b: Observed global temperatures are best explained by climate models that include both natural factors and human-caused increases in greenhouse gases. Models that include only natural factors do not reproduce the warming observed since the mid-twentieth century.
Image relabeled from Efbrazil(opens in new window) (CC-BY-SA(opens in new window)).

As a result, the scientific evidence indicates that modern global warming is primarily caused by human activities that increase greenhouse gas concentrations and strengthen the greenhouse effect.

Attribution.

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

This page titled 9.3: The Greenhouse Effect is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 21.1: The Greenhouse Effect and Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9.4: Evidence and Causes of Climate Change

What is Global Warming and Global Climate Change?

Global warming refers to the long-term increase in Earth's average surface temperature resulting primarily from human activities. **Climate change** is a broader term that includes global warming as well as changes in precipitation patterns, storms, droughts, sea level, and other aspects of Earth's climate system.

Climate has changed naturally throughout Earth's history. However, the rapid warming observed since the Industrial Revolution cannot be explained by natural causes alone. Scientists have concluded that human activities, particularly the combustion of fossil fuels and land-use changes such as deforestation, are the primary drivers of modern climate change.

Greenhouse Gases

The primary human-caused greenhouse gases are:

- **Carbon dioxide (CO_2)**
- **Methane (CH_4)**
- **Nitrous oxide (N_2O)**

These gases absorb infrared radiation emitted by Earth, strengthening the greenhouse effect and warming the atmosphere.

Carbon Dioxide

Carbon dioxide (CO_2) is the most important greenhouse gas driving modern climate change. Although it is naturally released through respiration, decomposition, volcanic eruptions, and ocean-atmosphere exchange, human activities, especially the burning of fossil fuels, have dramatically increased atmospheric CO_2 concentrations.

Atmospheric CO_2 concentrations have increased from approximately 280 ppm before the Industrial Revolution to more than 430 ppm today (figure $9.4.a$. This increase closely corresponds with the large-scale combustion of fossil fuels that began during the Industrial Revolution. Ice core records show that current concentrations are higher than at any point during at least the last 800,000 years (figure $9.4.b - c$.




Figure 9.4.*a*: The Keeling Curve: a daily record of global atmospheric carbon dioxide concentration started in 1958 and maintained by Scripps Institution of Oceanography at UCSD. Monthly mean carbon dioxide concentration in parts per million (ppm) measured at Mauna Loa Observatory, Hawaii. The carbon dioxide data on Mauna Loa constitute the longest record of direct measurements of CO2 in the atmosphere. The fluctuating black line represents the monthly mean values, centered on the middle of each month. Carbon dioxide concentrations dip every summer due to increased photosynthesis. . Image: Scripps Institution of Oceanography at UC San Diego (CC-BY-4.0); data from: C. D. Keeling, S. C. Piper, R. B. Bacastow, M. Wahlen, T. P. Whorf, M. Heimann, and H. A. Meijer, Exchanges of atmospheric CO2 and 13CO2 with the terrestrial biosphere and oceans from 1978 to 2000. I. Global aspects, SIO Reference Series, No. 01-06, Scripps Institution of Oceanography, San Diego, 88 pages, 2001. http://escholarship.org/uc/item/09v319r9




Figure 9.4.*b*: The EastGRIP ice core, which was drilled from the Greenland Ice Sheet. You can read more here. Image by Helle Astrid Kjær (CC-BY).




Figure 9.4.c: This graph, based on the comparison of atmospheric samples contained in ice cores and more recent direct measurements, provides evidence that atmospheric CO2 has increased since the Industrial Revolution. On the x-axis are the years before today (0 = 1950). It begins8400,000 years prior to 1950. On the y-axis is the carbon dioxide level in parts per million (ppm). Carbon dioxide levels have fluctuated over the years, but they never exceeded 300 parts per million until 1950. In 2026, carbon dioxide levels reached 431 ppm. Credit: Scripps Institution of Oceanography at UC San Diego (CC-BY-4.0) Ice core data: Lüthi, D., M. Le Floch, B. Bereiter, T. Blunier, J.-M. Barnola, U. Siegenthaler, D. Raynaud, J. Jouzel, H. Fischer, K. Kawamura, and T.F. Stocker. 2008. High-resolution carbon dioxide concentration record 650,000-800,000 years before present. Nature, Vol. 453, pp. 379-382, 15 May 2008.




Figure 9.4.d: Graph of change in temperature in degrees Celsius (top, blue line) and carbon dioxide concentration in parts per million by volume (bottom, green line) measured from the Vostok, Antarctica ice core. These have been associated for over 400,000 years. As carbon dioxide concentration increased, so did temperature. As carbon dioxide concentration decreased, so did temperature. These data were collected in 1999. Since then, carbon dioxide concentrations have increased to 409.8 ppm (2019 average). Image and caption (modified) by NOAA/Autopilot (CC-BY-SA)

Methane

Although carbon dioxide is the most abundant human-produced greenhouse gas, **methane (CH_4)** is far more effective at trapping heat on a molecule-for-molecule basis. Over a 20-year period, a single methane molecule can trap more than 80 times as much heat as a carbon dioxide molecule, making methane a powerful contributor to short-term climate warming. While methane persists in the atmosphere for a shorter time than carbon dioxide, its strong warming effect means that even relatively small increases in methane concentrations can have significant impacts on Earth's climate.

Human activities are responsible for much of the increase in atmospheric methane since the Industrial Revolution. Major sources include the extraction, processing, and transport of fossil fuels; agricultural practices, particularly livestock production where cattle release methane during digestion; the decomposition of organic waste in landfills; and microbial processes associated with wastewater treatment. As these activities have expanded over time, atmospheric methane concentrations have risen dramatically and are now more than twice their pre-industrial levels. This increase strengthens the greenhouse effect and contributes to the warming observed across the globe.

Nitrous Oxide

Nitrous oxide (N_2O) is released from agricultural soils, fertilizer use, fossil fuel combustion, and industrial processes. Molecule-for-molecule, N_2O traps approximately **265 times more heat than CO_2** over a 100-year period.

Evidence for Human-Caused Climate Change

Scientists study both **direct measurements** and **proxy records** to understand how Earth's climate has changed over time. Direct measurements, such as atmospheric carbon dioxide concentrations, temperature records, and satellite observations, provide detailed information about recent climate conditions. Because these records only extend back a relatively short period, scientists also rely on proxy data, which are indirect indicators of past climate. Proxies such as ice cores, tree rings, ocean sediments, and glacier records preserve evidence of environmental conditions that existed hundreds to hundreds of thousands of years ago. By comparing patterns from multiple independent sources of direct and proxy evidence, scientists can reconstruct past climates, evaluate natural climate variability, and place modern climate change in a long-term context.

Several lines of evidence from both direct observations and proxy records support the conclusion that recent climate change is primarily driven by human activities.

Atmospheric Carbon Dioxide Is Increasing

The **Keeling Curve** shows that atmospheric CO_2 concentrations have risen continuously since measurements began in 1958 (figure $9.4.a$.

This increase closely corresponds with the large-scale combustion of fossil fuels that began during the Industrial Revolution.

Modern CO_2 Levels Are Unprecedented

Ice core records demonstrate that present-day CO_2 concentrations exceed levels observed during the last 800,000 years (figure $9.4.c$.

Temperature and Carbon Dioxide Are Closely Linked

Ice core records from Antarctica reveal a strong relationship between atmospheric CO_2 concentrations and global temperature over hundreds of thousands of years (figure $9.4.d$)

Natural Factors Alone Cannot Explain Recent Warming

Climate models that include only natural influences such as volcanic activity and solar variability fail to reproduce the warming observed since the mid-twentieth century. Models that include both natural and human influences closely match observed temperatures.

Natural Influences on Climate

Natural processes continue to affect Earth's climate, including:

- Variations in solar activity
- Volcanic eruptions
- Long-term orbital cycles (Milankovitch cycles)

These factors explain many past climate changes. However, they cannot explain the rapid warming observed during recent decades.

Albedo and Climate Feedback

Earth reflects approximately **30%** of incoming sunlight back into space. This reflectivity is called **albedo**. Bright surfaces such as clouds, snow, and ice have high albedo and reflect more solar energy. Dark surfaces such as forests and oceans absorb more sunlight.

Some climate processes amplify warming through **reinforcing feedback loops**. One example involves the loss of Arctic sea ice. As warming causes sea ice to melt, Earth's albedo decreases because darker ocean water absorbs more sunlight than reflective ice. The additional absorption of solar energy causes further warming, which leads to even more ice melt. This is an example of a **reinforcing (positive) feedback loop**, in which an initial change amplifies itself (figure $9.4.e$.




Figure $9.4.e$: An example of a reinforcing (positive) feedback in the ice-albedo system. Image created by Erin Rempala using Microsoft Copilot.

Scientific Consensus on Climate Change

The **Intergovernmental Panel on Climate Change (IPCC)** is an international scientific body established by the United Nations to assess climate change research and provide policymakers with evidence-based information. Drawing on thousands of peer-reviewed studies, the IPCC concluded in its Sixth Assessment Report (2023) that:

- Human influence has **unequivocally** warmed the atmosphere, ocean, and land. In this context, *unequivocally* means that the evidence is so strong and consistent that there is no reasonable scientific doubt that human activities are the primary cause of recent global warming.
- Earth's average surface temperature is approximately 1.1°C warmer than it was before the Industrial Revolution.
- Climate change is already affecting the atmosphere, oceans, ice sheets, ecosystems, and human societies.
- Extreme weather events such as heat waves, heavy precipitation, droughts, and some storms are becoming more frequent or intense in many regions.

- Limiting future warming requires rapid and substantial reductions in greenhouse gas emissions.

The scientific consensus on anthropogenic climate change is supported by multiple independent lines of evidence, including atmospheric measurements, ice cores, temperature records, satellite observations, and climate models.

Why Warming Will Continue

Many greenhouse gases remain in the atmosphere for decades to centuries. Even if emissions were dramatically reduced today, some future warming would still occur because the oceans absorb and store large amounts of heat and release it slowly over time. As a result, warming and many of its impacts will continue for decades even after greenhouse gas concentrations stabilize.

Key Takeaways

Multiple independent lines of evidence show that Earth's climate is warming and that human activities are the primary cause. Atmospheric concentrations of carbon dioxide, methane, and other greenhouse gases have increased dramatically since the Industrial Revolution, strengthening the greenhouse effect. Ice core records, direct atmospheric measurements, temperature records, and climate models all support the conclusion that modern climate change is primarily driven by human activities.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Other sources:

- Intergovernmental Panel on Climate Change. (n.d.). *Sixth Assessment Report (AR6) cycle*. IPCC 6th assessment report(opens in new window) [www.ipcc.ch] Retrieved 6 May 2026.
- Scripps Institution of Oceanography. (n.d.). *Keeling Curve*. University of California San Diego. Keeling Curve (opens in new window) [keelingcurve.ucsd.edu]
- National Aeronautics and Space Administration. (n.d.). *Climate change: Vital signs of the planet*. *NASA(opens in new window) [science.nasa.gov]*
- Stanford University. (n.d.). *Methane and climate change*. Stanford Doerr School of Sustainability. Stanford(opens in new window) [sustainability.stanford.edu]
- U.S. Environmental Protection Agency. (n.d.). EPA.gov [www.epa.gov]

Modified by Melissa Ha from the following sources:

- Climate and the Effects of Global Climate Change from *General Biology* by OpenStax (licensed under CC-BY(opens in new window))
- Climate Change from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Carbon Cycle from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))

This page titled 9.4: Evidence and Causes of Climate Change is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 21.1: The Greenhouse Effect and Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9.5: Impacts of Climate Change

Impact of Climate Change

Climate change is already affecting Earth's atmosphere, oceans, ice sheets, ecosystems, and human societies. Rising greenhouse gas concentrations have increased global temperatures by approximately 1.1°C above pre-industrial levels and are driving a wide range of environmental changes expected to intensify in the coming decades.

Effects of increasing greenhouse gases include:

- increase Earth's average temperature (figure $9.5.h$)
- destabilize the patterns and amounts of precipitation,
- increase severity of hurricanes and storms,
- reduce ice and snow cover, and permafrost,
- raise sea level (figure $9.5.i$) due to thermal expansion and increase in water from melting land ice and glaciers,
- increase the acidity of the oceans (ocean acidification),
- alter phenology (e.g. timing of flowering, breeding, migration) of species




Figure $9.5.h$: Temperature increases have been most pronounced in northern latitudes and over land masses. The colors represent the temperature difference between the 2011-2020 average and the 1951-1980 baseline, with warmer colors (yellow, orange, red) representing increases, and cool colors (green, blue) representing decreases. The image uses longer term averages of at least a decade to smooth out climate variability due to factors such as El Niño. Grey areas in the image have insufficient data for rendering. Image and caption (modified) from NASA's Scientific Visualization Studio/Eric Fisk (public domain).

Explore the interactive climate monitor below to compare current global temperatures with historical averages. Which regions are warming the most relative to their historical conditions? How do the patterns compare with the warming trends shown in figure ($9.5.h$)?

Interactive Element

Reuters Climate Monitor: How today's temperatures compare to the historic average (Updates daily)

Reuters.com(opens in new window) [www.reuters.com]




Figure 9.5.4: Sea height variation (mm) over time. Sea height has increased about 3.3 millimeters per year on average since 1993. Data is from satellite sea level observations by the NASA Goddard Space Flight Center. Image by NASA(opens in new window) (public domain).

Future Climate Risks

The IPCC projects that continued greenhouse gas emissions will cause additional warming and increase climate-related risks. Each additional increment of warming increases the likelihood of extreme weather, ecosystem disruption, biodiversity loss, and economic damages. Some impacts, including sea-level rise, will continue for centuries even if emissions are substantially reduced. However, rapid reductions in greenhouse gas emissions can still limit many future impacts.

Changes in Precipitation and Extreme Weather

Climate change is altering patterns of precipitation around the world. Some regions are experiencing more frequent droughts and water shortages, while others are experiencing heavier rainfall and flooding. A warmer atmosphere can hold more water vapor, increasing the potential for intense precipitation events.

Climate change can also increase the severity of some extreme weather events. Heat waves have become more frequent and intense in many regions, and warmer ocean temperatures can contribute to stronger tropical cyclones and hurricanes.

Food and Water Resources

Climate change affects agriculture by altering temperature, precipitation, and water availability. Heat stress, drought, extreme weather, and changing pest distributions can reduce crop yields in many regions.

Changes in snowpack and rainfall patterns also affect freshwater supplies. Communities that depend on seasonal snowmelt, including much of the western United States, may face increasing challenges in managing water resources.

Ice, Snow, and Permafrost

Arctic sea ice, glaciers, snow cover, and permafrost have all declined as temperatures have increased. Permafrost, ground that remains frozen for at least two consecutive years, is particularly vulnerable because thawing can release stored greenhouse gases and damage ecosystems and infrastructure. Continued warming is expected to further reduce ice and snow cover, contributing to sea-level rise and climate feedback loops.

Climate Feedback loops

Some impacts of climate change can amplify additional warming. For example, melting ice exposes darker surfaces that absorb

more sunlight, while thawing permafrost can release stored carbon dioxide and methane into the atmosphere. These processes create **reinforcing feedback loops** that reinforce climate change.

Sea Level Rise

As discussed in the chapter introduction, sea-level rise poses significant challenges for coastal communities, including San Diego. Higher sea levels increase the risk of beach erosion, coastal flooding, saltwater intrusion, and damage to infrastructure and ecosystems.

Sea level rises for two primary reasons: warming ocean water expands (**thermal expansion**) and land-based ice, including glaciers and ice sheets, melts and adds water to the ocean. Since 1870, global sea level has risen by approximately 9.8 inches and is projected to continue rising throughout this century.

The amount of sea-level rise experienced at a specific location depends not only on global sea-level rise but also on local factors such as land subsidence or uplift, ocean currents, and regional geology. As a result, some coastal regions will experience greater impacts than others.

Ocean Acidification

Ocean acidification is the gradual decrease in ocean pH caused by the absorption of atmospheric carbon dioxide, posing a growing threat to corals, shellfish, and other marine organisms. Ocean acidification is discussed in greater detail in the next section.

Phenological Effects

Climate change is disrupting **phenology**, the timing of seasonal biological events such as flowering, migration, and reproduction. Species respond to warming at different rates, creating mismatches between organisms that depend on one another. For example, plants may flower before pollinators emerge, or migratory birds may arrive after peak food availability.

Human Health

Climate change affects human health in many ways. More frequent heat waves increase heat-related illness and deaths. Wildfires can reduce air quality and worsen respiratory diseases such as asthma. Flooding and extreme weather events can cause injuries, contamination of drinking water, and damage to infrastructure. Climate change can also affect mental health through displacement, disasters, and economic disruption.

Rising temperatures are expanding the ranges of mosquitoes and other disease vectors, increasing the risk of diseases such as dengue fever, West Nile virus, Zika, and malaria in new regions. Climate change can also affect wildlife diseases, contributing to amphibian declines and expanding outbreaks of forest pests such as bark beetles.

Key Takeaways

Climate change is already affecting Earth's atmosphere, oceans, ice sheets, ecosystems, and human societies. Rising temperatures, changing precipitation patterns, sea-level rise, ecosystem disruption, and human health impacts demonstrate that climate change is a present-day challenge rather than a future possibility. While some future changes are unavoidable, the magnitude of future impacts will depend largely on how quickly greenhouse gas emissions are reduced.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

This page titled 9.5: Impacts of Climate Change is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 21.1: The Greenhouse Effect and Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9.6: Ocean Acidification- The Other Carbon Problem

Ocean Acidification

When discussing climate change, most people think about rising temperatures. However, increasing atmospheric carbon dioxide (CO^2) affects Earth in another important way: it changes the chemistry of the oceans.

The world's oceans absorb approximately 25–30% of human-caused carbon dioxide emissions. This slows the rate of atmospheric warming, but it comes with an environmental cost. As carbon dioxide dissolves in seawater, it reacts with water molecules to form **carbonic acid** (H_2CO_3), increasing the concentration of hydrogen ions (H^+) and lowering ocean pH. This process is known as **ocean acidification**.

What Is Ocean Acidification?

Ocean acidification is the gradual decrease in the pH of seawater caused primarily by the absorption of atmospheric carbon dioxide.

Since the beginning of the Industrial Revolution, average ocean pH has decreased by approximately 0.1 pH units, representing about a 30% increase in acidity. Scientists project that ocean acidity will continue to increase if atmospheric CO_2 concentrations continue to rise.

Because the pH scale is logarithmic, even seemingly small declines in pH represent substantial changes in ocean chemistry.




Figure $9.6.a$ A pteropod shell is shown dissolving over time in seawater with a lower pH. When carbon dioxide is absorbed by the ocean from the atmosphere, the chemistry of the seawater is changed. Image credit: NOAA (Public Domain).

Why Ocean Acidification Matters

Many marine organisms use **calcium carbonate** ($CaCO_3$) to build shells, skeletons, and other structures. Examples include:

- Corals
- Clams
- Oysters
- Mussels
- Pteropods ("sea butterflies")

- Some plankton

As ocean acidity increases, carbonate ions become less available. This makes it more difficult for organisms to build and maintain shells and skeletons. In severe cases, existing shells may begin to dissolve.

Effects on Coral Reefs

Coral reefs are among the ecosystems most vulnerable to ocean acidification. Corals build their skeletons from calcium carbonate. As carbonate availability declines, coral growth slows. If atmospheric CO_2 concentrations continue to increase, coral calcification rates may decrease substantially, making reefs more vulnerable to erosion, disease, bleaching, and storm damage. Because coral reefs provide habitat for thousands of species, impacts on corals can affect entire marine ecosystems.

Effects on Food Webs

Ocean acidification also affects species at the base of marine food webs. Small planktonic organisms can experience reduced growth, reproduction, and survival. Because many fish, marine mammals, and seabirds depend on these organisms as food sources, disruptions can cascade through entire ecosystems. Ocean acidification therefore has the potential to influence fisheries, biodiversity, and food security.

Ocean Acidification and Climate Change

Ocean acidification and climate change share the same primary cause: increasing atmospheric carbon dioxide.

However, they are different environmental problems:

Climate Change	Ocean Acidification
Caused by greenhouse gases trapping heat in the atmosphere	Caused by carbon dioxide dissolving in seawater
Raises global temperatures	Lowers ocean pH
Alters weather, precipitation, and sea level	Alters marine chemistry
Affects both terrestrial and marine ecosystems	Primarily affects marine ecosystems

Because both problems result from carbon dioxide emissions, reducing fossil fuel use can help address both climate change and ocean acidification.

Connection to California and San Diego

Ocean acidification is particularly important along the California coast. Seasonal upwelling brings cold, carbon-rich water to the surface, making coastal waters naturally more acidic than many other regions. Additional carbon dioxide from human activities further increases acidity, posing challenges for marine ecosystems and shellfish industries.

Researchers are monitoring ocean acidification along the Southern California coastline because of its potential impacts on kelp forests, fisheries, marine biodiversity, and coastal food webs.

Key Takeaways

- Ocean acidification is caused by the absorption of atmospheric carbon dioxide by seawater.
- Since the Industrial Revolution, ocean acidity has increased by approximately 30%.
- Acidification reduces the availability of calcium carbonate needed by many marine organisms.
- Corals, shellfish, plankton, and other calcifying organisms are particularly vulnerable.
- Ocean acidification is distinct from global warming but is caused by the same increase in atmospheric carbon dioxide.
- Reducing carbon dioxide emissions helps slow both climate change and ocean acidification

This page titled 9.6: Ocean Acidification- The Other Carbon Problem is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 21.1: The Greenhouse Effect and Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9.7: Mitigating and Adapting to Climate Change

Climate Change Affects Everyone and Action is Urgently Needed

Human societies have developed within the relatively stable climate of the **Holocene**, and our water resources, agriculture, infrastructure, ecosystems, and public health are all closely tied to that stability. As the climate warms, these systems will experience increasing stress and disruption.

Because carbon dioxide persists in the atmosphere for roughly a century, continued warming is unavoidable in the coming decades. Higher atmospheric temperatures increase the likelihood of more severe and widespread impacts across Earth's systems. The climate conditions we have relied on historically are no longer a reliable guide for future planning.

Reducing future climate risks requires rapid reductions in greenhouse gas emissions and proactive preparation for the changes already underway. Decisions made today will strongly influence environmental and societal conditions for future generations.

Meeting this challenge requires **collective action**. Governments and industries must implement large-scale systemic changes, while individuals can reduce emissions through choices at home, in transportation, and at work. Many of these actions also provide economic and health benefits. Engagement at local and state levels can further support clean energy, efficiency, and climate-resilience initiatives.

Adaptation Limits

- Many adaptation options become less effective as warming increases, and more systems will hit adaptation limits.
- Maladaptation can occur without flexible, inclusive, long-term planning.
- Losses and damages will rise as warming grows.

Carbon Budgets & Net Zero

- Limiting warming requires net-zero CO_2.
- Whether the world stays below 1.5°C or 2°C depends heavily on emissions cuts this decade.
- Existing fossil fuel infrastructure alone would exceed the remaining 1.5°C carbon budget without additional abatement.

Mitigation Pathways

- All pathways that limit warming to 1.5°C or 2°C require immediate, deep emissions reductions in all sectors.
- Net-zero CO_2 is reached in the early 2050s for 1.5°C pathways and around the 2070s for 2°C pathways.

Overshoot

- If warming exceeds 1.5°C, temperatures could be reduced again only through sustained net-negative CO_2 emissions.
- Overshoot increases irreversible impacts, risks, and sustainability concerns due to greater reliance on carbon removal.

- ## Summary of Near-Term Climate Action Needed (from IPCC AR6)

- **Urgency of near-term integrated action** — Climate change poses a major risk to human well-being and planetary health, and the window to secure a livable future is rapidly closing. Actions taken this decade will shape conditions for thousands of years.
- **Benefits of early action** — Rapid, sustained mitigation and faster adaptation now would reduce future losses and damages and deliver major co-benefits (especially for air quality and health). Delays lock in high-emissions systems and raise long-term costs.
- **Mitigation and Adaptation Across Systems** — Deep emissions cuts require rapid, far-reaching transitions across all sectors, using a wide range of feasible, effective, and low-cost mitigation and adaptation options already available.
- **Sustainable development links** — Accelerated and equitable climate action supports sustainable development, with synergies outweighing trade-offs when implemented in the right contexts.
- **Equity and inclusion** — Prioritizing equity, climate justice, and just transitions strengthens both mitigation and adaptation. Supporting highly vulnerable groups and integrating adaptation into social protection improves resilience.
- **Governance and policy** — Effective climate action depends on strong political commitment, coordinated governance, clear goals, and broad access to finance and technology. Scaled-up regulatory and economic tools can drive deep emissions cuts.
- **Finance, technology, and cooperation** — Achieving climate goals requires a many-fold increase in adaptation and mitigation finance. Global capital exists, but barriers prevent it from flowing to climate action. Technology innovation and international cooperation are essential.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

This page titled 9.7: Mitigating and Adapting to Climate Change is shared under a license and was authored, remixed, and/or curated by LibreTexts.

- 21.1: The Greenhouse Effect and Climate Change by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

9.8: Data Dive- Flooding in Venice Italy

Overview

Historically, Venice Italy has a history of flooding that is caused from a multitude of both natural and human caused phenomena. Currently, these issues are being exacerbated by sea level changes as a result of climate change. As such, the City of Venice has been carefully tracking frequency of flooding through the years. The graph below represents the frequency of water level was above 110cm per 10-year period:




Figure 9.8.*a*: Frequency of flooding exceeding 110cm in Venice Italy. Graph by Rachel Schleiger (CC-BY-NC) modified from data published by the City of Venice.

Questions

1. What is the independent (explanatory) variable and the dependent (response) variable?
2. What question(s) are the authors trying to answer with this graph?
3. What are the results of this graph illustrating about flooding in Venice?
4. How can other coastal cities around the world learn from what patterns are emerging in this graph?
5. Provide a brief reflection for how these results make you feel and what they make you think about.

Raw Data For Above Graph(s)

Table 9.8.*a*: Raw data for the frequency of flooding exceeding 110cm in Venice Italy. Graph by Rachel Schleiger (CC-BY-NC) modified from data published by the City of Venice.

Raw data for the frequency of flooding exceeding 110cm in Venice Italy.

Ten Year Interval	Frequency
1870-79	4
1880-89	2
1890-99	3
1900-09	2

Ten Year Interval	Frequency
1910-19	3
1920-29	3
1930-39	8
1940-49	5
1950-59	13
1960-69	31
1970-79	31
1980-89	26
1990-99	44
2000-09	52
2010-19	69

Attribution

Rachel Schleiger (CC-BY-NC)

This page titled 9.8: Data Dive- Flooding in Venice Italy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 21.2: Data Dive- Flooding in Venice Italy by Melissa Ha and Rachel Schleiger has no license indicated.

10: THE WATER SYSTEM

From Ancient Aqueducts to Modern Water Scarcity

During the fifteenth century, the Mexica (Aztecs) constructed the Chapultepec Aqueduct to bring freshwater from springs at Chapultepec to their capital city of Tenochtitlan, located where the historic center of Mexico City stands today. Early sections of the aqueduct were damaged by flooding in the mid-1400s and were later rebuilt and expanded using more durable materials. During the Spanish conquest of Tenochtitlan in 1521, the system was damaged, but a new aqueduct was constructed during the colonial period along much of the original route. Today, the remains of the Chapultepec Aqueduct serve as a reminder of the long history of engineering solutions developed to provide water for growing cities.

The Chapultepec Aqueduct illustrates a challenge that societies have faced for centuries: securing reliable freshwater supplies for growing populations. As cities expand, local water sources often become insufficient, requiring increasingly complex infrastructure to transport water from distant locations.

This challenge remains highly relevant in Southern California today. More than 20 million people depend on water imported through large aqueduct systems that transport water from the Colorado River, the Sacramento-San Joaquin Delta, and the Eastern Sierra Nevada. Prolonged droughts, climate change, overuse of river systems, declining groundwater supplies, and growing demand have placed increasing pressure on these water sources.

Just as the residents of Tenochtitlan relied on engineered systems to supply freshwater to their city, modern Southern Californians depend on extensive water infrastructure to support homes, agriculture, industry, and ecosystems. Understanding where water comes from, how it moves through the water cycle, and how human activities affect freshwater availability is essential for addressing one of Southern California's most important environmental challenges: ensuring a sustainable water supply for future generations.



Figure 10.a: Remains of the Chapultepec Aqueduct in Mexico City. The aqueduct was originally constructed by the Aztecs to transport freshwater to Tenochtitlan and illustrates the long history of engineering solutions to urban water supply challenges.
Image by Diana du Vent CC-BY-SA.

10.1: The Hydrologic Cycle and Freshwater Distribution

10.2: Water Usage
10.3: Water Scarcity and Sustainable Water Management
10.4: Case Study - The Salton Sea
10.5: Water Pollution
10.6: Water Treatment- Protecting Drinking Water and Treating Wastewater
10.7: Preventing Water Pollution and Restoring Watersheds

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

This page titled 10: The Water System is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 13: Water Resources by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

10.1: The Hydrologic Cycle and Freshwater Distribution

The Hydrologic Cycle and Freshwater Distribution

Freshwater is essential for life on Earth. It supports ecosystems, regulates climate, sustains agriculture, and provides drinking water for human communities. Although Earth is often called the "Water Planet," only a small fraction of the planet's water is available as freshwater. Understanding how water moves through the environment and where it is stored provides the foundation for understanding water scarcity, water quality, and sustainable water management.

The Hydrologic Cycle

The **hydrologic cycle** (water cycle) describes the continuous movement of water among the atmosphere, oceans, land surface, groundwater, glaciers, and living organisms (figure $10.1.a$). The water cycle is driven by two primary forces: **solar energy**, which powers evaporation, and **gravity**, which moves water downhill and returns precipitation to the Earth's surface. Like other environmental systems, the water cycle contains both **stocks** (reservoirs where water is stored) and **flows** (processes that move water between reservoirs).

Water enters the atmosphere through several pathways:

- **Evaporation**: liquid water changes into water vapor
- **Sublimation**: ice changes directly into water vapor
- **Transpiration**: plants release water vapor through small openings in their leaves

Ecologists often combine evaporation and transpiration into a single process called **evapotranspiration**.

As water vapor rises into the atmosphere, it cools and undergoes **condensation**, forming clouds. When water droplets or ice crystals become large enough, they fall as **precipitation**, including rain, snow, sleet, or hail. Some precipitation infiltrates into the ground to replenish groundwater, while the remainder may flow across the land surface as **runoff**, eventually entering streams, lakes, and oceans.




Figure \(\PageIndex{a}\): The hydrologic cycle. Water continuously moves among the atmosphere, land, groundwater, surface waters, and oceans through processes such as evaporation, condensation, precipitation, infiltration, and runoff. Screenshot from interactive map by United States Geological Survey (public domain); [https://water.usgs.gov/vizlab/water-cycle/index.html]

Interactive Element

Explore the Water Cycle

Have you ever wondered where the water in your drinking glass was before it reached your home? Explore the USGS Water Cycle Interactive to follow water as it moves through the atmosphere, rivers, oceans, glaciers, groundwater, and living organisms. The interactive visualization shows how Earth's water is constantly recycled and highlights the many pathways water can take through the hydrologic cycle.

Try It!

- Click on different parts of the water cycle to learn how water moves between reservoirs.
- Compare the amount of water stored in oceans, glaciers, groundwater, rivers, lakes, and the atmosphere.
- Identify examples of **stocks** (water reservoirs) and **flows** (processes that move water).
- Trace a possible journey of a water molecule from the ocean to a cloud, into a river, and eventually into an aquifer.

Think About It: If water is constantly moving through the hydrologic cycle, why can freshwater still become scarce in some regions? How might climate change, urbanization, or groundwater pumping affect the movement and storage of water in Earth's systems?

Water Reservoirs and Residence Time

Water is the only common substance that naturally occurs on Earth as a solid, liquid, and gas. The **hydrosphere** includes all water on Earth, whether it is stored in oceans, glaciers, groundwater, lakes, rivers, soils, the atmosphere, or living organisms.

Water is not distributed evenly among these reservoirs. Some reservoirs, such as the atmosphere, contain relatively small quantities of water but exchange water rapidly. Others, such as glaciers and oceans, contain enormous quantities of water that may remain stored for thousands of years.

The average amount of time a water molecule remains in a reservoir is called its **residence time**. Water may remain in the atmosphere for only a few days, while water stored in glaciers, groundwater, and oceans can remain there for centuries to millennia (figure $10.1.b$).




Figure $10.1.b$: Average residence time of water in major reservoirs of the hydrologic cycle. Image from OpenStax(opens in new window) (CC-BY(opens in new window)).

Where Is Earth's Freshwater?

Although water covers approximately 71% of Earth's surface, most of it is not available for human use. About **97.5% of Earth's water is saltwater** found in oceans and seas. Only about **2.5% of Earth's water is freshwater**.

Most freshwater is stored in:

- Glaciers and ice caps (~69%)
- Groundwater (~30%)
- Surface waters and other reservoirs (~1%)

As a result, only a very small fraction of Earth's water is readily available in rivers, lakes, wetlands, and shallow groundwater systems.

If all of Earth's water were represented by one gallon, the amount of readily accessible freshwater would be only a few tablespoons.




Figure $10.1.c$: Distribution of Earth's water among major reservoirs.
Image from USGS/Igor Shiklomanov(opens in new window) [USGS] (public domain).

Because so much freshwater is locked in glaciers and polar ice caps, **groundwater** represents the largest reservoir of readily accessible freshwater. Surface waters such as rivers and lakes account for only a tiny portion of Earth's freshwater supply, yet they support many ecosystems and provide drinking water for much of the world's population.



Figure 10.1.*d*: Mountain Glacier in Argentina. Mountain glaciers store a large percentage of Earth's freshwater. Image by Luca Galuzzi [www.galuzzi.it].

Although freshwater is unevenly distributed among Earth's reservoirs, its availability also depends on where and how often precipitation occurs. Global climate patterns strongly influence which regions receive abundant freshwater and which experience chronic water scarcity.

Why Freshwater Availability Varies Around the World

Freshwater supplies are replenished primarily through precipitation, but precipitation is not evenly distributed across the planet.

Global patterns of temperature and atmospheric circulation strongly influence where rain and snow fall. Near the equator, warm air rises, cools, and releases moisture as frequent precipitation. In contrast, descending air around 30° north and south latitude creates many of the world's deserts, including the Sahara Desert and deserts of the southwestern United States (figure 10.1.*e*).




Figure 10.1.*e*: Annual average precipitation by country in 2025. Image by OurWorldinData.org/water-use-stress (CC-BY(opens in new window)).

Local geography also affects precipitation. Mountains create **rain shadows** by forcing moist air to rise and cool on one side of the mountain range. As the air descends on the opposite side, it becomes warmer and drier, producing arid conditions (figure 10.1.*f*).




Figure 10.1.*f*: Rainshadow effect. Image by Meg Stewart (CC BY-SA 2.0) <https://creativecommons.org/licenses/by-sa/2.0>, via Wikimedia Commons [https://en.wikipedia.org/wiki/Rain_s...ow_effect.jpg]

The distribution of precipitation strongly influences freshwater availability, which in turn affects ecosystems, agriculture, and human settlement patterns. Climate change is altering precipitation patterns in many regions. Some areas are expected to experience more frequent droughts, while others may see an increase in intense rainfall events and flooding. These changes can affect freshwater availability for both ecosystems and human communities.

In the United States, the 100th Meridian roughly separates the wetter eastern portion of the country from the drier western region. Agriculture west of this boundary often depends on irrigation supplied by rivers, reservoirs, and groundwater (figure 10.1.*g*).

Southern California lies west of the 100th Meridian and therefore relies heavily on imported water, groundwater, conservation, and water-recycling projects to meet water demand.




Figure 10.1.*g*: Precipitation across the United States. Image: U.S. Geological Survey, public domain.

Watersheds, Rivers, and Lakes

Water that falls on land eventually flows downhill toward rivers, lakes, wetlands, and oceans. The geographic area drained by a river and its tributaries is called a **watershed** or **drainage basin**. Watersheds connect landscapes through the movement of

water, sediment, nutrients, and pollutants. Because water flows downhill through watersheds, activities occurring in one part of a watershed can influence water quality, habitat conditions, and flood risk many kilometers downstream.

The Tijuana River Watershed is a local example of how watersheds connect ecosystems and communities across large geographic areas (figure $10.1.h$). This binational watershed spans approximately 1,700 square miles in both Mexico and the United States before draining into the Pacific Ocean through the Tijuana River Estuary, one of Southern California's largest remaining coastal wetlands.

Because the watershed crosses an international boundary, environmental challenges often require cooperation between multiple jurisdictions. During storms, runoff can carry trash, sediment, wastewater, and other pollutants from upstream areas through the Tijuana River and into the estuary. After major rain events, large amounts of debris may accumulate in the estuary, degrading water quality, harming wildlife habitat, and affecting nearby coastal communities. These events demonstrate an important principle of watershed science: water, pollutants, and sediments do not recognize political boundaries. Effective watershed management therefore requires collaboration among local, regional, national, and international partners.




Figure $10.1.h$: The Tijuana River Watershed. Image by Geobica (CC BY-SA 4.0); Wikipedia

Likewise, the Mississippi-Atchafalaya River Basin drains about 40% of the continental United States and transports water, nutrients, and sediment from much of North America into the Gulf of Mexico (figure $10.1.i$).

Much of the watershed is used for agriculture, and fertilizers applied to farms throughout the Midwest can be transported by runoff into streams and rivers. Excess nitrogen and phosphorus carried downstream stimulate rapid growth of algae in coastal waters. When these algae die and decompose, oxygen levels decline, creating a large **hypoxic zone** (low-oxygen "dead zone") in the Gulf of Mexico that can stress or kill fish, shellfish, and other aquatic organisms.

The Mississippi River Basin illustrates how activities occurring thousands of kilometers upstream can affect ecosystems far downstream. The connection between nutrient pollution, algal growth, and oxygen depletion is known as **eutrophication**, a major water-quality issue that will be explored in more detail in the Water Pollution section of this chapter.




Figure $10.1.i$: The Mississippi-Atchafalaya River Basin. Image by US EPA, public domain

Together, the Tijuana River Watershed and Mississippi River Basin demonstrate that local actions can have regional and even international consequences for water quality and ecosystem health.

Rivers and lakes provide critical habitat for freshwater organisms and serve as important sources of water for agriculture, industry, and human communities. Many water-storage reservoirs are artificial lakes created by dams. Because rivers often cross political boundaries, access to freshwater can also become a source of conflict. Examples of rivers that have experienced disputes over water allocation include the Colorado, Nile, Euphrates, Ganges, and Jordan Rivers.

Groundwater Resources

Although rivers and lakes are highly visible components of the water cycle, **groundwater** is actually the largest reservoir of usable freshwater on Earth. An **aquifer** is a body of permeable rock, sediment, or soil that stores and transmits groundwater. Aquifers supply water to ecosystems, agriculture, municipalities, and private wells. The **water table** marks the upper boundary of the **saturated zone**, where all pore spaces are filled with water (figure $10.1.j$).



How ground water occurs in rocks.

Figure $10.1.j$: The unsaturated zone, saturated zone, and water table within an aquifer. Image by USGS (opens in new window)(public domain)

Two characteristics largely determine an aquifer's ability to store and transmit water:

- **Porosity**: the percentage of open space within a material
- **Permeability**: the degree to which pore spaces are connected

A material with high porosity can store large amounts of water, while high permeability allows water to move through the material more easily.

Groundwater Recharge and Aquifer Storage

Groundwater is replenished through **recharge**, when precipitation and surface water infiltrate into the soil and percolate downward into aquifers. Recharge commonly occurs in wetlands, floodplains, stream channels, and other areas where water can readily soak into the ground. Groundwater eventually returns to the surface through **discharge**, including springs, wetlands, streams, and lakes. As a result, groundwater and surface-water systems are closely connected, and changes to one often affect the other.

Some communities also use **aquifer storage and recovery (ASR)** systems (figure $10.1.k$). These projects intentionally recharge aquifers by injecting or infiltrating water during wet periods and storing it underground for future use. Because underground storage is protected from evaporation, ASR can increase water security in regions with seasonal rainfall or periodic droughts.




Figure $10.1.k$: Groundwater recharge and discharge processes. Image by USGS (opens in new window)(public domain).

The Hydrologic Cycle and Water Sustainability

The hydrologic cycle connects the atmosphere, oceans, rivers, lakes, groundwater, glaciers, ecosystems, and human communities through the continuous movement of water. Although water is constantly recycled, freshwater availability varies across regions because of differences in climate, geology, topography, and land use.

Understanding how water moves through the hydrologic cycle and where it is stored provides the foundation for understanding water scarcity, groundwater depletion, water-quality challenges, and sustainable water management. Although water is continually recycled through Earth's systems, freshwater supplies can become stressed when withdrawals exceed replenishment or when climate and land-use changes alter natural hydrologic processes.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

- Water Cycle and Fresh Water Supply, Water Supply Problems and Solutions, and Biogeochemical Cycles from *Environmental*

Biology by Matthew R. Fisher (licensed under CC-BY(opens in new window))

- The Water Cycle(opens in new window), Water Use and Distribution(opens in new window), and Groundwater(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))
- Biogeochemical Cycles and the Flow of Energy in the Earth System(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (licensed under CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Biogeochemical Cycles(opens in new window) from *Biology 2e* by OpenStax (CC-BY(opens in new window)). Access for free at openstax.org(opens in new window).

This page titled 10.1: The Hydrologic Cycle and Freshwater Distribution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 13.1: Fresh Water Supply and the Water Cycle by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

10.2: Water Usage

Water Usage

Freshwater supply is one of the most important ecosystem services. In 2014, global water consumption was 3999 km^3 per year (over 1,000 trillion gallons!). According to the United Nations, approximately half of the world experiences economic or physical water scarcity each year. The greatest use of this water is for irrigation in agriculture, but significant quantities of water are also extracted for public and municipal use, as well as industrial applications and power generation (figure $10.2.a$). Increasingly, AI data servers require a tremendous amount of water, energy and carbon emissions. According to the Environmental and Energy Studies Institute(opens in new window), in 2025, a large data center can use up to 5 million gallons of water each day (MGD), and a 100-word AI prompt needs the equivalent of one bottle of water (519 ml) for cooling.




Figure $10.2.a$: Global water usage by sector 2024. Municipal refers to domestic use, such as by local businesses and households. Data from UNESCO. Image by Erin Rempala (CC-BY-NC(opens in new window)).

Humans require only about 1 gallon per day to survive, but a typical person in the United States uses approximately 82-100 gallons per day, which includes cooking, washing dishes and clothes, flushing the toilet, and bathing (2015 data, EPA). Additionally, we rely on the food, energy, and mineral resources, which all require water to produce. For example, approximately 21 gallons are needed to produce a salad with tomato lettuce and cucumber, 52 gallons for one egg, and 674 gallons for a 6-ounce steak. 20-50 gallons are needed to produce one kilowatt-hour (kWh) of electricity from a traditional power plant (about 1 kWh is needed to heat an oven for 30 minutes), one pair of blue jeans requires about 1000 gallons, and one ton of steel consumes about 63,000 gallons of water.

> Water Footprint: You can learn more about your water footprint and the embedded water footprint of the resources you use in the following water footprint calculator(opens in new window)

The **water demand** of an area is a function of the population and other uses of water. In the United States, 281 billion gallons of water were withdrawn per day in 2015, of which 82 billion gallons are fresh groundwater (figure $10.2.b$). In 2015, the state of California accounted for 9% of national water withdrawal (figure $10.2.c$). In 2025, agriculture is approximately 40% of California's annual water use,10% is used by communities and the remaining 50% returns to the environment. San Diego has been improving its per capita water usage since 1991. To learn more go to the San Diego Water Authority Water-Use Efficiency Fact Sheet(opens in new window) for 2025.




Figure 10.2.*b*: Average daily freshwater withdrawals (in billion gallons per day, left y-axis) over time (x-axis) in the United States. Light blue bars (left) represent groundwater withdrawals, medium blue (middle) represents withdrawals from surface water, and dark blue bars (right) represent total withdrawals in billions of gallons per day. In 1950, total withdrawals were around 170 billion gallons per day. This gradually increased until 1980 at just over 350 billion gallons per day. In 1985, total withdrawals dropped to around 325 billion gallons and stayed relatively constant until decreasing in 2010. In 2015, total withdrawals were around 260 billion gallons. Most of the changes in withdrawals overtime were due to changes in surface water withdrawals rather than groundwater withdrawals. Surface water withdrawals have generally been 2-3 times greater than groundwater withdrawals. The pink line represents the United States population size in millions (right y-axis), which has increased from roughly 170 million in 1950 to about 325 million in 2015. Image by USGS(opens in new window) (public domain).




Figure 10.2.c: Total water withdrawals by state in 2015. The lightest blue (Z) indicates 0-2,000 million gallons per day, and the darkest blue (V) indicates 20,001-28,800 million gallons per day. California and Texas consume the most water, and Alaska, Hawaii, the Dakotas, Oklahoma, and some New England states consume the least. Image modified from USGS (public domain).California water use varies dramatically across regions and between wet and dry years. From https://www.ppic.org/publication/wat...in-california/ (SOURCE: Department of Water Resources.) NOTE: Data for 2017 were not available.

Attributions

Modified by Erin Rempala (CC BY-NC-SA 4.0(opens in new window))

Modified by Melissa Ha from the following sources

- Water Use and Distribution(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC BY-NC-SA(opens in new window))
- Water Cycle and Fresh Water Supply from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Biogeochemical Cycles and the Flow of Energy in the Earth System(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (licensed under CC-BY(opens in new

window)). Download for free at CNX(opens in new window).

This page titled 10.2: Water Usage is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 13.2: Water Usage by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

10.3: Water Scarcity and Sustainable Water Management

Water Supply Problems: Resource Depletion

Freshwater supplies can be depleted when water is withdrawn faster than natural processes can replace it. One common example is **groundwater mining**, which occurs when groundwater is pumped from an aquifer faster than it is replenished. Pumping from a water well creates a localized lowering of the water table called a **cone of depression**. When many wells withdraw groundwater over a long period, the regional water table can decline substantially. As groundwater levels fall, wells may need to be drilled deeper, increasing costs and sometimes accessing more saline water.

Surface water can also be depleted through overuse. Many rivers, lakes, and reservoirs experience declining water levels when withdrawals exceed replenishment. Some major rivers, including the Colorado River in the United States and the Yellow River in China, have periodically failed to reach their usual endpoints because so much water is diverted for human use. Lake systems can also shrink due to excessive water withdrawals, as illustrated by the dramatic decline of the Aral Sea. In addition, glaciers that supply freshwater to downstream communities are shrinking because of accelerated melting associated with climate warming.

Groundwater depletion creates several additional environmental challenges. Along coastlines, excessive pumping of freshwater aquifers can lead to **saltwater intrusion**, a process in which seawater moves into freshwater aquifers and contaminates water supplies. Lowering the water table can also alter groundwater flow patterns. In some cases, pollutants that would normally move away from a well may instead be drawn toward it.

A declining water table can also cause the land surface to sink. **Subsidence** is the gradual sinking of land over a broad area, whereas **sinkholes** are sudden collapses that occur over a smaller area. Both can damage infrastructure, alter drainage patterns, and increase the costs of managing water resources (figure $10.3.a$).




Figure $10.3.a$. Formation of a Cone of Depression around a Pumping Water Well. Image: By Tara Gross - https://www.usgs.gov/media/images/co...level-lowering, Public Domain, [https://commons.wikimedia.org/w/inde...rid=146710900]

The Global Water Crisis

The **water crisis** refers to the growing challenge of providing sufficient quantities of clean, accessible water for people and ecosystems. Water shortages, also called **water scarcity**, can result from either a lack of available water or limited access to safe water supplies.

Two forms of water scarcity are commonly recognized:

- **Physical water scarcity:** insufficient water is available to meet demand.
- **Economic water scarcity:** water is available, but infrastructure, governance, or financial limitations prevent access to it.

Water stress is generally greatest in regions with low precipitation, high population densities, or both. Arid regions of North Africa and the Middle East experience some of the highest levels of water stress, while rapidly growing populations in parts of Asia have increased demand on already limited freshwater supplies.

Climate change is expected to worsen water scarcity in many regions. Changes in precipitation patterns may increase drought frequency in some areas, while shrinking mountain glaciers may reduce streamflow during dry seasons. Rising sea levels can also increase saltwater intrusion into coastal aquifers, further reducing freshwater availability (figure $10.3.b$).




Figure $10.3.b$. Map of global water stress (a symptom of water scarcity) in 2019. Water stress is the ratio of water use relative to water availability and is therefore a demand-driven scarcity. Image by Genetics4good, GFDL [http://www.gnu.org/copyleft/fdl.html], via Wikimedia Commons.

Water scarcity can affect food production, human health, economic development, and political stability. Because many rivers and aquifers cross political boundaries, competition for freshwater can contribute to regional tensions. Areas that have experienced disputes related to water use include the Nile River Basin in Africa, the Euphrates and Tigris Rivers in the Middle East, the Ganges River system in South Asia, and the Aral Sea Basin in Central Asia.

Moving Toward Sustainable Water Use

Addressing water scarcity requires a combination of technologies, policies, and conservation practices that reduce demand and protect freshwater systems.

Dams and Reservoirs

Dams create reservoirs that store water during wet periods and make it available during dry periods (figure $10.3.c$). Reservoirs can provide drinking water, generate hydroelectric power, reduce flooding, and support recreation.

However, reservoirs can have environmental costs. Water stored in reservoirs may be lost through evaporation, especially in arid climates. Dams also alter river ecosystems, change fish migration routes, and modify downstream sediment transport and river flow patterns.



Figure $10.3.c$. Parker Dam on Lake Havasu at the start of the Colorado River Aqueduct. Image credits Erin Rempala (CC BY-NC-SA 4.0).

Aqueducts and Water Transfers

Aqueducts transport water from regions where water is relatively abundant to regions where demand is high. These systems support agriculture and urban development in areas with limited local water supplies. California has several major water-transfer systems, including the Colorado River Aqueduct, the State Water Project, the Los Angeles Aqueduct, and the San Diego Aqueduct.

Water-transfer projects can be controversial because diverting water from one watershed may reduce water availability and alter ecosystems in another. Before water diversions began, the Owens Valley was a productive agricultural region where farms and ranches relied on the Owens River for irrigation and livestock production. In the early twentieth century, the City of Los Angeles purchased land and water rights throughout the valley and diverted much of the river's flow through the Los Angeles Aqueduct to meet the city's growing water demand.

These diversions contributed to the desiccation of Owens Lake and lowered water levels in Mono Lake, causing ecological damage and reducing water availability for local communities. After years of legal challenges and environmental advocacy, conservation measures and water-management agreements helped stabilize Mono Lake and partially restore its ecosystem. Owens Lake, however, remains largely dry and requires ongoing dust-control efforts because winds can carry fine sediments from the exposed lakebed into surrounding communities.

Another major example is the Colorado River Aqueduct, which transports water from the Colorado River to Southern California (figure $10.3.d$). Completed in 1941, the aqueduct carries water approximately 252 miles (406 km) across deserts and over mountain ranges to supply communities in the Los Angeles metropolitan area and other parts of Southern California. This engineering project helped support the rapid growth of cities in one of the driest regions of the United States.

However, the Colorado River is shared by seven U.S. states, multiple Tribal Nations, and Mexico, making allocation of its water highly contentious. Water rights along the river were largely divided during the twentieth century using estimates of river flow that are now known to have been unusually high. Today, increasing demand, prolonged drought, and climate change have reduced river flows while agricultural, urban, tribal, recreational, and ecological water needs continue to grow. As water levels in major reservoirs such as Lake Mead and Lake Powell decline, disagreements have intensified over how limited Colorado River water should be distributed among competing users and how much water should remain in the river to support ecosystems.

The Owens Valley and Colorado River examples illustrate how transferring water between watersheds can produce both benefits and costs. Aqueducts have enabled the growth of major cities and agricultural regions in arid environments, but they have also transformed ecosystems, affected rural communities, and generated conflicts over water rights and environmental protection. Understanding these tradeoffs is an important part of sustainable water management.





Figure 10.3.*d*. Top: The California Aqueduct California Aqueduct in southern California. Bottom: Gene Pumping Station off Lake Havasu at the start of the Colorado River Aqueduct. Image credits Erin Rempala (CC BY-NC-SA 4.0).

Interactive Element

Interactive Element: Follow the Journey of Colorado River Water

Ever wondered how water reaches Southern California's cities and suburbs from hundreds of miles away? Explore the Metropolitan Water District's interactive Colorado River Aqueduct Map to trace the 252-mile journey of water from Lake Havasu across deserts, through tunnels, and over mountain ranges to communities throughout Southern California.

Try It!

- Locate Parker Dam and the Gene Pumping Plant, where water enters the aqueduct.
- Follow the aqueduct's route across the desert to Southern California.
- Identify the pumping plants, reservoirs, and tunnels that help move water uphill and across rugged terrain.
- Consider the energy, infrastructure, and cooperation required to deliver water to millions of people in one of the driest regions of North America.

As you explore, think about this question: What are the environmental benefits and tradeoffs of transporting water hundreds of miles to support cities and agriculture in arid regions?

Desalination

Desalination removes dissolved salts from seawater or saline groundwater to produce freshwater. Because it creates a new source of freshwater, desalination can help meet water demand in coastal regions.

Despite its benefits, desalination is energy intensive and expensive compared with most conventional water sources. The process also produces concentrated brine that must be disposed of carefully to minimize environmental impacts. Desalination facilities are particularly common in arid regions of the Middle East, where freshwater is scarce and energy resources are readily available.

Water Conservation

Water conservation reduces water use while maintaining essential human activities and ecosystem functions. Conservation measures can be implemented at household, municipal, and agricultural scales.

Examples of household conservation include:

- Installing high-efficiency appliances
- Using low-flow toilets and showerheads
- Repairing leaks
- Landscaping with drought-tolerant native plants
- Reducing unnecessary water use

Agriculture offers some of the greatest opportunities for conservation because irrigation accounts for most freshwater withdrawals worldwide. Effective strategies include:

- Growing crops suited to local climate conditions
- Using efficient irrigation systems, such as drip irrigation
- Practicing no-till agriculture to reduce evaporation from soil
- Reusing treated wastewater for irrigation

Interactive Element

Explore California Native Plants with Calscape

Looking for a beautiful, water-wise landscape that supports local biodiversity? The **California Native Plant Society's Calscape Plant Finder** can help you discover native plants that are naturally adapted to your local climate, soils, and wildlife. Whether you want to create habitat for hummingbirds, butterflies, native bees, or other pollinators, Calscape makes it easy to find species that match your landscaping goals.

Many California native plants are highly drought tolerant and require little to no supplemental irrigation once established, making them an excellent choice for water conservation. Native plant landscapes can also improve soil health, reduce the need for fertilizers and pesticides, and provide food and shelter for local wildlife.

In many parts of California, native plants can also be incorporated into **Firewise native plant landscaping** practices. Firewise landscapes are designed to reduce wildfire risk around homes by creating defensible space, using fire-resistant plant communities, and minimizing the accumulation of highly flammable vegetation. When properly selected and maintained, native plants can help create landscapes that are both ecologically beneficial and better adapted to California's fire-prone environment.

Try It! Visit **Calscape.org**. Search for native plants that:

- Require little or no summer irrigation

- Attract birds, butterflies, or native pollinators
- Bloom during different seasons of the year
- Can be used in a Firewise landscape

How might replacing a portion of a conventional lawn with native plants conserve water while supporting local ecosystems and increasing resilience to drought and wildfire?

Rainwater Harvesting

Rainwater harvesting captures and stores rainfall for later use before it becomes runoff. Water reuse systems can also recycle treated wastewater for irrigation, industrial processes, and groundwater recharge. These approaches reduce demand on rivers, reservoirs, and aquifers while increasing the resilience of local water supplies.

Greywater Reuse

Along with rainwater harvesting, wastewater recycling, and efficient irrigation, **greywater reuse** can help communities reduce freshwater withdrawals and increase the resilience of local water supplies. Greywater is household wastewater from showers, bathtubs, bathroom sinks, and laundry that has not come into contact with toilet waste. Because it typically contains fewer contaminants than sewage, greywater can often be reused for irrigation and other nonpotable applications. Greywater reuse can be particularly valuable in arid regions facing increasing water scarcity due to population growth, drought, and climate change.

Together, rainwater harvesting, greywater reuse, wastewater recycling, improved infrastructure, and sustainable management of freshwater ecosystems can help communities meet water needs while protecting aquatic systems for future generations.

Groundwater Recharge

Groundwater recharge is the process by which water moves from the land surface into the ground and replenishes aquifers. Recharge occurs naturally when precipitation infiltrates the soil and percolates downward through porous rock and sediment. In many regions, however, urbanization has reduced natural recharge because roads, parking lots, and buildings prevent water from soaking into the ground.

Communities can enhance groundwater recharge through a variety of strategies. These include protecting open spaces and wetlands, using **permeable pavement** that allows water to infiltrate the soil, installing **bioswales** and **rain gardens** to capture and filter stormwater, directing runoff into infiltration basins, and recharging aquifers with highly treated recycled water (figure $10.3.e$). These green infrastructure approaches increase infiltration, improve water quality, and reduce flooding while helping replenish groundwater supplies. Artificial recharge projects can help offset groundwater depletion, reduce the impacts of drought, and store water underground for future use.



Figure 10.3.e: Runoff from the vicinity flows into an adjacent bioswale. Image: Erica Fischer, CC BY 2.0 <https://creativecommons.org/licenses/by/2.0>, via Wikimedia Commons [https://commons.wikimedia.org/wiki/F..._Bioswale.jpg]

Groundwater recharge provides several environmental benefits. Replenished aquifers can support drinking water supplies, maintain streamflow during dry periods, reduce land subsidence, and help prevent saltwater intrusion in coastal regions. Because groundwater is naturally protected from evaporation, aquifers can serve as important long-term water-storage systems in arid climates.

However, recharge projects must be carefully managed to prevent contamination of groundwater resources. Water used for recharge must meet strict quality standards, and land uses in recharge areas must be monitored to reduce the risk of pollutants entering aquifers.

Case Study: Building Water Resilience in San Diego County

Like many communities in the arid southwestern United States, San Diego County relies on a diverse portfolio of water sources to improve long-term water security and reduce dependence on imported water. Historically, San Diego County has relied heavily on imported water from the Colorado River and Northern California, making it vulnerable to drought, infrastructure disruptions, and changes in water availability elsewhere in the western United States. To increase local water reliability in the face of drought, climate change, and growing demand, the region has invested in several major water-supply projects that illustrate many of the water-management strategies discussed in this chapter, including desalination, potable water reuse, groundwater recharge, conservation, and diversification of water supplies.

The Claude "Bud" Lewis Carlsbad Desalination Plant, which began operation in 2015, is the largest seawater desalination facility in the United States. The plant can produce up to 50 million gallons per day (MGD), equivalent to approximately 190 million liters per day, making it one of the largest locally controlled water sources in San Diego County. By converting seawater into freshwater, the facility provides a drought-resistant water supply, although desalination requires substantial energy and produces concentrated brine (salty water left over from the process) that must be carefully managed.

The Pure Water San Diego program is designed to produce drinking water from highly treated recycled wastewater. Using advanced purification technologies such as microfiltration, reverse osmosis, and ultraviolet disinfection, the project will provide nearly half of the City of San Diego's drinking water supply by 2035. By reusing water that would otherwise be discharged to the ocean, Pure Water San Diego increases local water independence while reducing wastewater discharges. **Potable water reuse** is often considered one of the most sustainable approaches to water management because it treats wastewater as a valuable water source rather than a waste product.

In East County, the East County Advanced Water Purification (East County AWP) Program is developing a similar potable water reuse system to improve regional resilience. The project will purify treated wastewater to drinking-water standards and use it to replenish local reservoirs. In addition to providing a reliable local water source, the project is expected to reduce reliance on imported water and increase drought preparedness for communities in eastern San Diego County.

Groundwater recharge is also playing an increasingly important role in California water management. In San Diego County, projects that restore wetlands, floodplains, and other natural infiltration areas help increase groundwater recharge, while highly treated recycled water can be used to replenish reservoirs and aquifers. These approaches help offset groundwater depletion and provide additional water storage that is protected from evaporation.

Together, desalination, potable water reuse, groundwater recharge, conservation, and imported water supplies illustrate how San Diego County is diversifying its water portfolio to build a more resilient and sustainable water system in an increasingly variable climate.

Despite these investments, imported water from the Colorado River and State Water Project continues to supply a substantial portion of San Diego County's water demand. The region's water strategy therefore combines imported water, desalination, potable reuse, groundwater recharge, local storage, and conservation measures to improve resilience to drought and climate change.

San Diego County's experiences demonstrate that no single solution can address water scarcity. Instead, communities often combine conservation, water recycling, desalination, imported water, and ecosystem protection to create resilient water systems capable of adapting to drought, population growth, and climate change.

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from **7.2: Water Supply Problems and Solutions** by Matthew R. Fisher originally licensed CC BY 4.0. Original source: https://openoregon.pressbooks.pub/envirobiology (CC BY-NC-SA 4.0).

And additional sources cited:

- Claude "Bud" Lewis Carlsbad Desalination Plant
- Pure Water San Diego
- East County Advanced Water Purification
- Metropolitan Water District of Southern California

Modified by Matthew R. Fisher from Essentials of Environmental Science(opens in new window) by Kamala Doršner is licensed under CC BY 4.0.

This page titled 10.3: Water Scarcity and Sustainable Water Management is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

- 7.2: Water Supply Problems and Solutions by Matthew R. Fisher is licensed CC BY 4.0. Original source: https://openoregon.pressbooks.pub/envirobiology.

10.4: Case Study - The Salton Sea

Case Study: The Salton Sea

History

The Salton Sea is California's largest lake (35 miles long by 15 miles wide), located in the Sonoran Desert of southeastern California within the Imperial and Coachella Valleys (figure $10.4.a$). Unlike most lakes, the Salton Sea has no natural outlet. Water leaves the lake primarily through evaporation, causing dissolved salts and other substances to accumulate over time.

Although the modern Salton Sea formed in 1905, the basin has a long geologic history of periodically filling with water. For thousands of years, flooding from the Colorado River occasionally created a large freshwater lake known as **Lake Cahuilla**, which at times covered much of the present-day Imperial and Coachella Valleys. Lake Cahuilla eventually dried when the Colorado River changed course, leaving behind a dry lakebed known as the **Salton Basin**.

The modern Salton Sea was created accidentally in 1905 when flooding on the Colorado River breached irrigation works associated with the Imperial Canal and diverted the river's flow into the low-lying Salton Basin for nearly two years. By the time engineers stopped the flow in 1907, a large inland lake had once again formed in the desert.

Salton Sea map showing project locations. Contact DWR if you need more information about this map.

Figure $10.4.a$: Salton Sea Ecological Restoration Project Map. Image credit: California Department of Water Resources, public domain.

For much of the twentieth century, the Salton Sea was sustained by agricultural runoff from surrounding farms. Water from the Colorado River, delivered through irrigation canals, flowed onto agricultural fields and eventually drained into the lake. These inflows helped maintain water levels and created habitat for millions of migratory birds traveling along the Pacific Flyway. The Salton Sea also supported recreational activities including boating, fishing, and tourism, making it a popular destination during the 1950s and 1960s.

The Shrinking Salton Sea

Beginning in the late twentieth century, water conservation measures and water-transfer agreements reduced the amount of agricultural runoff entering the lake. As inflows declined, evaporation exceeded water inputs and the lake began to shrink. Falling water levels exposed large areas of previously submerged lakebed known as **playa** (figure $10.4.b$.



Figure $10.4.b$: Playa along Bombay Beach, Salton Sea. Image by Erin Rempala (CC BY-NC-SA 4.0)

Ecological Impacts

As the Salton Sea contracts, its salinity continues to increase. Today, the lake is considerably saltier than the Pacific Ocean. Rising salinity has reduced fish populations and altered the food webs that support migratory birds. Large fish die-offs and harmful algal blooms have become increasingly common, further degrading water quality and wildlife habitat.

The decline of the Salton Sea also threatens one of California's most important remaining wetland ecosystems. As wetlands throughout the western United States have been drained or developed, the Salton Sea has become a critical stopover and wintering habitat for migratory birds along the Pacific Flyway. Continued habitat loss could negatively affect numerous bird species that depend on the lake during migration (figure $10.4.c$).



Figure $10.4.c$: A burrowing owl at the Sonny Bono Salton Sea National Wildlife Refuge. Image: Erin Rempala (CC BY-NC-SA 4.0).

Human Health and Environmental Justice

The shrinking shoreline has also created significant human health concerns. Winds can lift fine sediments from the exposed playa and transport dust into nearby communities. These dust particles may contain salts, pesticides, and other contaminants that accumulated in lake sediments over decades. Exposure to airborne dust has been linked to respiratory illnesses, including asthma.

Many communities surrounding the Salton Sea are economically disadvantaged and have been disproportionately affected by declining environmental conditions and poor air quality. As a result, the Salton Sea is often cited as an example of an environmental justice issue, where environmental burdens are not distributed equally among populations.

Restoration Challenges and Lessons Learned

Recognizing these concerns, California agencies have initiated restoration projects aimed at reducing dust emissions, creating wetland habitat, and improving ecological conditions around the lake. However, restoration remains challenging because of competing demands for Colorado River water, prolonged drought, climate change, and the large scale of environmental degradation.

The Salton Sea illustrates how water-management decisions can have unintended ecological and social consequences. A water system originally developed to support agriculture created a valuable aquatic ecosystem, but changes in water use have led to declining water levels, increasing salinity, habitat loss, and public health concerns. The Salton Sea serves as a powerful example of how altering the movement of water across a landscape can transform both ecosystems and human communities for generations.

Think About It: Water-management decisions often involve tradeoffs among agriculture, urban water supplies, ecosystem health, and public well-being. Based on the strategies discussed in this chapter, what combination of conservation, potable water reuse, groundwater recharge, habitat restoration, and policy changes might help address the challenges facing the Salton Sea while balancing the needs of people and wildlife?

Sources:

California Department of Water Resources: Salton Sea Unit

The Salton Sea Authority

This page titled 10.4: Case Study - The Salton Sea is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala via source content that was edited to the style and standards of the LibreTexts platform.

- 7.5: Case Study - The Aral Sea - Going, Going, Gone by Matthew R. Fisher is licensed CC BY 4.0. Original source: https://openoregon.pressbooks.pub/envirobiology.

10.5: Water Pollution

Water Pollution

Water pollution occurs when harmful substances such as bacteria or contaminants enter rivers, lakes, wetlands, oceans, or groundwater and degrade ecosystems, reduce water quality, or threaten human health. Because water moves continuously through the hydrologic cycle, pollutants can be transported across watersheds, accumulate in sediments, enter food webs, and affect communities far from their original source.

This section examines the major sources of water pollution, the different types of pollutants that affect aquatic ecosystems, and how nutrient pollution can trigger eutrophication, hypoxia, and aquatic dead zones.

Sources of Water Pollution

Water can become contaminated through both natural processes and human activities. Agricultural operations, urban development, industry, mining, landfills, wastewater treatment systems, and atmospheric deposition can all contribute pollutants to water bodies.

As water moves through the hydrologic cycle, it dissolves and transports contaminants. The environmental impact of a pollutant depends on its concentration, toxicity, persistence, and residence time within the ecosystem.

Point Source Pollution

Point source pollution originates from a single, identifiable source that can be located and monitored. Because the source of contamination is known, point-source pollution is often easier to regulate than pollution that comes from many dispersed locations.

Common examples of point-source pollution include wastewater discharge pipes, industrial outfalls, chemical spills, landfill leachate, concentrated animal feeding operations (**CAFOs**), and combined sewer overflow systems (**CSOs**). **Leachate** is contaminated liquid that forms when rainwater filters through a landfill and dissolves chemicals from decomposing waste. If not properly collected and treated, leachate can contaminate nearby soil, surface water, and groundwater.

Concentrated animal feeding operations (CAFOs) house large numbers of livestock in relatively small areas and generate substantial quantities of manure and wastewater. If these wastes escape storage ponds or are improperly applied to agricultural fields, nutrients, pathogens, and organic matter can enter nearby waterways. In California's Imperial Valley, a major agricultural region east of San Diego County, runoff from livestock operations and agricultural activities contributes nutrients and other pollutants to rivers, drainage canals, and eventually the Salton Sea. Nutrient pollution from these sources can encourage algal blooms, reduce dissolved oxygen levels, and degrade aquatic habitats (figure $10.5.a$).



Figure $10.5.a$: Concentrated animal feeding operations (CAFOs) are potential point sources of water contamination. Image by U.S. Environmental Protection Agency(opens in new window) (public domain).

Another example of point-source pollution is a **combined sewer overflow (CSO)** system (figure $10.5.b$). In some cities, stormwater runoff and sewage are transported through the same set of pipes. During periods of heavy rainfall, the volume of water may exceed the capacity of the treatment system, causing untreated sewage and stormwater to be released directly into rivers, lakes, or coastal waters. These discharges can introduce pathogens, nutrients, and oxygen-demanding wastes into aquatic ecosystems.

Because the source of contamination can be identified, environmental regulations often focus on monitoring and limiting pollutant discharges from point sources. However, accidental spills, infrastructure failures, and extreme weather events can still result in significant water-quality problems.

Combined sewer overflows are more common in older cities such as Chicago, New York, and parts of the Great Lakes region, where stormwater and sewage are transported through the same pipe network. San Diego generally relies on separate stormwater and wastewater systems, although storm events can still contribute to water-quality problems through runoff and infrastructure failures.




Figure $10.5.b$: Combined sewer overflow systems (CSO) can release untreated sewage during major storm events. Sewage from domestic, commercial, and industrial sources enter the system through the down spout and storm water enters through the storm drain. During dry weather (and small storms), all sewage is handled by the publicly owned treatment works (POTW). A dam prevents raw sewage from flowing into a water body through the outfall pipe. During wet weather (large storms), the relief structure allows some of the combined sewage and storm water to be discharged untreated to an adjacent water body. Image by U.S. Environmental Protection Agency (opens in new window) (public domain).

Because the source can be identified, point-source pollution is often easier to monitor and regulate.

Nonpoint Source Pollution

Nonpoint source pollution comes from many dispersed sources across a landscape. It is currently the leading cause of water pollution in the United States.

Common examples include:

- Fertilizer runoff from farms and lawns
- Urban stormwater runoff
- Oil and grease from roads and parking lots
- Sediment from erosion
- Animal waste
- Pesticides and herbicides

Rainfall and irrigation water transport these pollutants through watersheds into rivers, lakes, wetlands, estuaries, and groundwater.

The Tijuana River Watershed provides a Southern California example of nonpoint source pollution. During storms, runoff can transport trash, sediment, wastewater, fertilizers, and other pollutants through the watershed into the Tijuana River Estuary near Imperial Beach, CA. Because these contaminants originate from many locations across the watershed, addressing the problem requires coordinated management throughout the entire binational watershed rather than focusing on a single pollution source (figure $10.5.c$).




Figure 10.5.c: Point source contamination can be traced to specific points of discharge. Air pollution is an example of nonpoint source contamination because contaminants are throughout the air and landscape, ultimately entering the water from multiple locations. Eroded soil and sediment are another source of nonpoint source pollution. Image by USGS(opens in new window) (public domain).

Major Categories of Water Pollutants

Although thousands of contaminants can affect water quality, most pollutants fall into three broad categories: **chemical pollutants**, **biological pollutants**, and **physical pollutants**. Understanding these categories helps scientists identify pollution sources, assess environmental impacts, and develop appropriate strategies to protect and restore water quality.

Chemical Pollutants

Chemical pollution arises from agriculture, industry, mining, transportation, urban runoff, and waste disposal.

Concentrations of water pollutants are often extremely small and are commonly measured in **parts per million (ppm)** or **parts per billion (ppb)**. One part per million is equivalent to one drop of a substance in about 50 liters of water, while one part per billion is one thousand times smaller. To provide context, rainwater typically contains about 4 ppm of dissolved solids, river water averages about 120 ppm, and seawater contains roughly 35,000 ppm of dissolved solids. These measurements help scientists assess water quality and determine whether pollutant concentrations may pose risks to ecosystems or human health.

Organic Pollutants

Organic pollutants include herbicides, pesticides, industrial solvents, petroleum products, pharmaceuticals, and synthetic hormones. Many of these substances can persist in the environment long after their release. Some are classified as **persistent organic pollutants (POPs)** because they break down very slowly, accumulate in the tissues of living organisms (**bioaccumulation**), and become increasingly concentrated as they move through food webs through a process known as **biomagnification** (covered in toxicology section of the book).

Examples of **persistent organic pollutants (POPs)** include **DDT** (dichlorodiphenyltrichloroethane), a synthetic insecticide that contributed to eggshell thinning in many bird species; **polychlorinated biphenyls (PCBs)**, industrial chemicals once used in electrical equipment that persist in sediments and aquatic food webs; and **dioxins**, toxic byproducts of industrial processes that can interfere with hormone function, reproduction, and development. Like many POPs, these chemicals resist decomposition, bioaccumulate in organisms, and biomagnify through food webs.

Case Study: Love Canal

One of the most famous examples of organic chemical contamination occurred at Love Canal in Niagara Falls, New York (figure $10.5.d$).



Figure $10.5.d$: Love Canal, Niagra Falls, NY. Image by U.S. Environmental Protection Agency (public domain).

Between 1942 and 1952, industrial chemical wastes were buried in an abandoned canal. The site was later developed in residential neighborhood and school district. Residents eventually experienced elevated rates of illness, including cancers, chemical residues appeared in basements and soils.

Community activism led by then 27- year old housewife, Lois Gibbs, helped bring national attention to the crisis and contri to the establishment of the **Comprehensive Environmental Response, Compensation and Liability Act (CERCLA** commonly known as the **Superfund Program** (figure $10.5.f$). Today, the EPA uses Superfund authority to identify and clea some of the nation's most hazardous waste sites.

Inorganic Pollutants

Inorganic pollutants include nutrients such as nitrate and phosphate, heavy metals, salts, radioactive materials, and substances that alter water acidity. These pollutants can originate from natural geologic sources, but human activities such as agriculture, mining, industry, energy production, and urban development have greatly increased their concentrations in many aquatic ecosystems.

Nutrient Pollution

Nitrogen and phosphorus are essential nutrients required for plant growth. However, excessive inputs of these nutrients from fertilizers, sewage, and animal waste can degrade water quality. When excess nitrogen and phosphorus enter waterways, they can stimulate rapid growth of algae and aquatic plants, a process known as **eutrophication**. Because nutrient pollution plays a central role in algal blooms, oxygen depletion, and hypoxic "dead zones," it is discussed in greater detail later in this chapter.

Heavy Metals

Some inorganic pollutants are **heavy metals**, including arsenic, mercury, lead, cadmium, and chromium. Unlike many organic pollutants, heavy metals do not degrade over time. As a result, they can accumulate in sediments and living organisms. Many heavy metals also bioaccumulate in individual organisms and biomagnify through food webs, exposing top predators, including

humans, to potentially harmful concentrations. This will be covered in more detail in the toxicology chapter.

Case Study: The Bangladesh Arsenic Crisis

One of the most significant examples of heavy-metal contamination occurred in Bangladesh. During the 1970s, millions of shallow groundwater wells were installed to reduce waterborne diseases caused by contaminated surface water. Although the wells successfully reduced exposure to pathogens, many tapped naturally occurring arsenic-rich groundwater. As a result, millions of people were chronically exposed to arsenic in their drinking water, making the Bangladesh arsenic crisis one of the largest mass poisonings in human history.

The Bangladesh arsenic crisis illustrates how efforts to solve one environmental health problem can sometimes create unintended consequences when geologic and water-quality conditions are not fully understood.

Case Study: Minamata Disease

Another well-known example of heavy-metal pollution occurred in Minamata, Japan. Beginning in the 1930s, industrial wastewater containing **methylmercury**, a highly toxic form of mercury, was discharged into Minamata Bay. Mercury entered the aquatic food web and bioaccumulated in fish and shellfish that were regularly consumed by local residents.

Over time, thousands of people developed severe neurological disorders, including numbness, loss of coordination, hearing and vision impairment, developmental abnormalities in children, and death. The disease became known as **Minamata disease** and is one of the most famous examples of environmental contamination affecting human health.

The Minamata disaster demonstrated how pollutants released into the environment can move through ecosystems and eventually affect people, even when the source of contamination is not immediately obvious. Later in the Environmental Health chapter, we will explore the related concepts of **bioaccumulation** (the buildup of contaminants within organisms) and **biomagnification** (the increase in contaminant concentrations at higher levels of a food web), both of which contributed to the severity of mercury exposure in Minamata.

Salt Contamination

Salt is another common inorganic pollutant. Elevated salinity in freshwater systems can result from natural mineral deposits, agricultural irrigation practices, road-salt application, or **saltwater intrusion** into coastal aquifers (figure $10.5.e$). Increased salinity can harm freshwater organisms, alter ecosystem processes, reduce crop productivity, and degrade drinking-water supplies.




Figure 10.5.e: Top: Saltwater Intrusion. Image by USGS (public domain).

pH and Water Quality

The acidity or alkalinity of water also influences water quality. **pH** measures the concentration of hydrogen ions in a solution and indicates whether water is acidic, neutral, or basic. Most freshwater ecosystems function within a relatively narrow pH range. When water becomes too acidic or too alkaline, aquatic organisms may experience reduced growth, impaired reproduction, or increased susceptibility to pollutants. Acid deposition and ocean acidification, two major causes of pH changes in aquatic ecosystems, are discussed later in this text.

Hard Water

Hard water contains relatively high concentrations of dissolved calcium and magnesium. These minerals can reduce the effectiveness of soaps and lead to scale buildup on pipes, appliances, and other surfaces. Although hard water is often considered a nuisance for household and industrial uses, it generally does not pose a threat to human health.

San Diego County is known for having relatively hard water, largely because much of the region's water originates from the Colorado River and local watersheds that flow through mineral-rich rocks and sediments. As water moves across the landscape, it dissolves calcium and magnesium, increasing water hardness. Evaporation from reservoirs and other surface-water storage systems can further concentrate these dissolved minerals. Many households in the region use water-softening systems to reduce mineral buildup and improve the performance of plumbing fixtures and appliances.

Biological Pollutants

Biological pollutants include disease-causing bacteria, viruses, protozoans, and parasites that contaminate water supplies. These organisms, collectively known as **pathogens**, commonly enter waterways through untreated sewage, leaking septic systems, animal waste, livestock operations, and combined sewer overflows.

Pathogen contamination remains one of the leading causes of waterborne disease worldwide. Illnesses such as cholera, dysentery, hepatitis A, and giardiasis are spread through contaminated water, particularly where sanitation and wastewater-treatment infrastructure are inadequate.

Even in developed countries, heavy rainfall can overwhelm sewer systems and increase pathogen contamination in rivers, lakes, and coastal waters. Water treatment technologies such as filtration, chlorination, ozonation, and ultraviolet disinfection help reduce health risks by removing or inactivating disease-causing organisms.

Physical Sources of Pollution

Physical pollutants include trash, sediment, and thermal pollution (figure $10.5.h$). Trash was discussed extensively in the Solid Waste Management chapter.



Figure $10.5.f$: Garbage floating in water, Circular Quay, Sydney, Australia. Image by MrActiniuM, CC BY-SA 4.0 <https://creativecommons.org/licenses/by-sa/4.0>, via Wikimedia Commons.

Excess sediment enters bodies of water when activities such as agriculture, construction, mining, and deforestation increase erosion. Sediments can carry toxins, nutrients, and other contaminants into waterways while also increasing **turbidity**, a measure of water cloudiness. High turbidity reduces the amount of sunlight that penetrates the water, limiting photosynthesis by aquatic plants and algae and disrupting aquatic ecosystems.

Thermal pollution occurs when water temperatures rise above their natural range. Many power plants, including coal, natural gas, and nuclear facilities, use water for cooling and then release it back into rivers, lakes, or coastal waters at elevated temperatures. Warmer water holds less dissolved oxygen than cooler water, making it more difficult for aquatic organisms to obtain the oxygen they need for cellular respiration (figure $10.5.g$). Elevated temperatures can also alter species distributions, disrupt reproduction, and increase physiological stress. Dams can contribute to thermal pollution by changing natural water-flow patterns and water temperatures downstream.




Figure $10.5.g$: Water temperature affects dissolved oxygen concentrations in bodies of water. From January to December 2017, water temperature (blue) increases in summer months and then decreases. Dissolved oxygen levels (orange) dips in the summer when temperature increases. Image and caption (modified) from USGS(opens in new window) (public domain).

Locally, following major storms, large amounts of trash and sediment can accumulate in the Tijuana River Estuary, demonstrating how physical pollutants can be transported through watersheds and affect ecosystems far downstream. This example highlights an important concept in watershed science: activities occurring far upstream can influence water quality and habitat conditions many kilometers away.

Eutrophication, Biochemical Oxygen Demand, and Hypoxia

As discussed in the previous section on watersheds, nutrient runoff from Midwestern agricultural fields enters streams and rivers throughout the **Mississippi-Atchafalaya River Basin** and is ultimately transported to the Gulf of Mexico. This nutrient pollution contributes to one of the world's largest seasonal hypoxic zones, demonstrating how activities on land can affect ecosystems far downstream.

Eutrophication

Aquatic plants and algae require a variety of nutrients to grow, but **nitrogen (N)** and **phosphorus (P)** are often in relatively short supply. Because these nutrients frequently limit plant growth, they are known as **limiting nutrients**. This is why nitrogen and phosphorus are major ingredients in agricultural fertilizers.

When excessive amounts of nitrogen and phosphorus enter waterways from fertilizers, sewage, animal waste, and urban runoff, they can stimulate rapid growth of algae and aquatic plants. This process is known as **cultural eutrophication**, a form of water pollution caused by human activities.

Excessive algal growth can reduce water quality in several ways. Dense mats of algae may block sunlight needed by aquatic plants, clog fish gills, and alter aquatic habitats. Some species of algae and cyanobacteria (blue-green algae) also produce toxins that can harm wildlife, pets, livestock, and humans. Rapid population explosions of these organisms are known as **harmful algal blooms (HABs)** or **toxic algal blooms** (figure $10.5.h$).



Figure \(\PageIndex{h}\): A cyanobacterial bloom in Lake Erie caused by excess nutrient pollution. Image by NASA(opens in new window) (public domain).

Although the Gulf of Mexico dead zone is one of the best-known examples of nutrient pollution, similar processes can occur at smaller scales in California. Nutrient-rich runoff from agriculture and livestock operations in the Imperial Valley contributes to water-quality challenges in the Salton Sea, including harmful algal blooms and declining ecosystem health.

Biochemical Oxygen Demand (BOD)

Eventually, the large quantities of algae and aquatic plants produced during eutrophication die and become organic matter. As bacteria and other decomposers break down this material, they consume oxygen during cellular respiration.

The amount of dissolved oxygen required by microorganisms to decompose organic matter is called **biochemical oxygen demand (BOD)**. High BOD indicates that large amounts of organic material are being broken down, leaving less oxygen available for fish and other aquatic organisms.

Common sources of **oxygen-demanding waste** include untreated sewage, animal manure, food-processing waste, decaying vegetation, and algal blooms. As microorganisms decompose this organic matter, they consume dissolved oxygen from the water. When oxygen consumption exceeds the rate at which oxygen is replenished, dissolved oxygen concentrations decline, placing stress on fish, aquatic invertebrates, and other organisms that rely on oxygen for survival.

Fish species differ in their tolerance to low oxygen conditions. Sensitive species such as trout require relatively high dissolved oxygen concentrations, whereas more tolerant species such as carp and catfish can survive in degraded waters with lower oxygen level (figure $10.5.i$).



Figure

: top image: Rainbow trout thrive in waters with low biological oxygen demand and high dissolved oxygen concentrations. Bottom: Carp are more tolerant of waters with high biological oxygen demand and lower oxygen levels. Top image by Engbretson Eric/U.S. Fish and Wildlife Service

(opens in new window)
(public domain). Bottom image by
X posid(opens in new window)
(public domain).

Hypoxia and Dead Zones

When dissolved oxygen concentrations fall below approximately 2 ppm, aquatic systems enter a condition known as **hypoxia**. Organisms that cannot escape these low-oxygen waters may die from lack of oxygen, creating a biological **dead zone** (figure $10.5.j$).

Approximately half of the lakes in North America, Europe, and Asia are affected by some degree of eutrophication. One of the best-known examples is the large seasonal dead zone that develops in the Gulf of Mexico each summer. Nutrient-rich runoff from agricultural fields throughout the Mississippi River Basin fuels algal growth in coastal waters. When the algae die and decompose, oxygen levels decline dramatically, creating hypoxic conditions over thousands of square kilometers.

Eutrophication and hypoxia are particularly difficult to address because they are primarily driven by **non-point source pollution**,

which originates from many dispersed locations rather than a single identifiable source. Reducing nutrient pollution often requires coordinated watershed-scale management involving agriculture, wastewater treatment, stormwater management, and land-use planning.




Figure 10.5.j: The process of eutrophication. Excess nutrients stimulate algal growth, increasing organic matter available for decomposition. As bacteria break down this material, oxygen levels decline, potentially producing hypoxic conditions and aquatic dead zones. Image credit Libretexts.

Emerging Contaminants

Scientists are increasingly concerned about **emerging contaminants**, which are substances that were not traditionally monitored in water supplies. These include pharmaceuticals, personal-care products, microplastics, and **per-** and **polyfluoroalkyl substances (PFAS)**, often called "forever chemicals" because they persist in the environment for long periods.

These contaminants have been detected in rivers, lakes, groundwater, oceans, and drinking-water sources worldwide. Although many occur at low concentrations, researchers continue to investigate their long-term effects on ecosystems and human health. Unlike many naturally occurring substances, emerging contaminants are largely the result of modern industrial and consumer activities. They are sometimes referred to as **novel entities** because they represent new chemicals, materials, and engineered substances that did not previously exist in Earth's systems.

The accumulation of novel entities has become a global environmental concern. In the Planetary Boundaries framework, the Novel Entities boundary includes synthetic chemicals, plastics, pesticides, pharmaceuticals, and other human-made substances that can disrupt ecological and Earth-system processes. Scientists have concluded that humanity has likely exceeded this planetary boundary because the production and release of novel entities now outpaces our ability to fully assess, monitor, and manage their environmental impacts. The growing presence of emerging contaminants in water supplies illustrates how local pollution issues can contribute to global environmental change.

Local Connection: PFAS in Southern California

PFAS contamination has been detected in groundwater and drinking-water sources throughout California, including parts of Southern California. State monitoring programs have identified PFAS in multiple groundwater basins and public water systems, leading to increased testing, treatment efforts, and regulatory oversight. Because PFAS resist degradation and can persist in water, soils, wildlife, and human tissues for many years, they have become a major focus of water-quality management. Their persistence highlights the challenge of managing contaminants that remain in the environment long after they are released and illustrates why synthetic chemicals are included within the **Novel Entities** planetary boundary. More broadly, PFAS exemplify a growing challenge associated with novel entities: many synthetic chemicals are intentionally designed to be durable and resistant to breakdown. While these properties can make products more effective, they can also allow contaminants to persist in ecosystems and drinking-water supplies long after their release.

Water Pollution and Sustainability

Water pollution illustrates how activities on land can affect ecosystems and communities throughout a watershed. Pollutants released from agriculture, industry, transportation, urban development, and waste disposal can travel through rivers, groundwater, wetlands, estuaries, and oceans, often impacting locations far from their original source.

Protecting water quality requires reducing pollution at its source, improving wastewater treatment, protecting wetlands and watersheds, and adopting land-use practices that reduce contaminant runoff. Because water connects ecosystems and human communities across landscapes, effective solutions often require cooperation among individuals, industries, governments, and entire watersheds.

Think About It: Treating Symptoms or Solving Problems?

Many water-pollution problems are addressed using **end-of-pipe solutions**, which remove contaminants after they have already entered the environment. Examples include wastewater treatment plants, trash booms in rivers, drinking-water filtration systems, and beach cleanups that remove litter from shorelines.

Other approaches attempt to reduce pollution before it enters waterways. Examples include reducing fertilizer use, redesigning products and packaging, improving agricultural practices, protecting wetlands, and managing entire watersheds to reduce runoff.

Both approaches can improve water quality, but they operate at different points within a system.

Reflect: Imagine volunteers collect thousands of pounds of trash during a beach cleanup. What environmental benefits result from that effort? What problems might still remain after the cleanup is finished? Compare the strengths and limitations of removing pollution after it enters the environment with strategies that aim to prevent pollution from being created in the first place. Which approaches seem most effective, and why?

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Sources:

California State Water Resources Control Board. *Per- and Polyfluoroalkyl Substances (PFAS).* [https://www.waterboards.ca.gov/pfas/]

U.S. Environmental Protection Agency. *PFAS Explained.* [https://www.epa.gov/pfas]

Smith, A. H., Lingas, E. O., & Rahman, M. (2000). Contamination of drinking-water by arsenic in Bangladesh: A public health emergency. *Bulletin of the World Health Organization, 78*(9), 1093-1103.

Modified by Melissa Ha from the following sources:

- Water Contamination(opens in new window) from *An Introduction to Geology* by Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))
- Water Pollution from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- The Industrialization of Nature: A Modern History (1500 to the present)(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors. Download for free at CNX(opens in new window). (licensed under CC-BY(opens in new window))

This page titled 10.5: Water Pollution is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20.1.1: Water Pollutants and Their Sources by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

10.6: Water Treatment- Protecting Drinking Water and Treating Wastewater

Water Treatment: Protecting Drinking Water and Treating Wastewater

The most deadly form of water pollution is contamination by disease-causing microorganisms. According to the World Health Organization, unsafe drinking water, inadequate sanitation, and poor hygiene contribute to hundreds of thousands of deaths each year, primarily in low-income countries. Preventing these illnesses requires both access to safe drinking water and effective treatment of wastewater.

Although the terms are sometimes used interchangeably, **drinking-water treatment** and **wastewater treatment** serve different purposes. Drinking-water treatment removes contaminants from rivers, reservoirs, lakes, and groundwater to produce water that is safe to consume. In contrast, wastewater treatment removes contaminants from sewage after water has been used in homes, businesses, and industries. Both systems are essential for protecting public health and maintaining healthy aquatic ecosystems.

Drinking Water Treatment

Before water reaches homes and businesses, it usually undergoes several treatment steps designed to remove sediments, contaminants, and pathogens.

Most drinking-water treatment plants use a combination of:

1. **Coagulation and flocculation**, which cause small particles suspended in water to clump together.
2. **Sedimentation**, which allows these larger particles to settle out of the water (figure \(\PageIndex{a}\)).
3. **Filtration**, which removes remaining particles by passing water through sand, gravel, activated carbon, or membrane filters.
4. **Disinfection**, which uses chlorine, ozone, or ultraviolet (UV) light to kill disease-causing microorganisms.
5. **Distribution**, which delivers treated water through pipes to homes and businesses.

These processes significantly reduce the risk of waterborne diseases and improve water quality before the water is consumed.



Figure 10.6.a: Top: The Metropolitan Water District of Southern California's Robert B. Diemer Water Treatment Plant in Yorba Linda, California. Bottom: Sedimentation basins at the plant. Image credits: Erin Rempala (CC BY-NC-SA 4.0).

Wastewater Collection Systems

After water is used, it becomes **wastewater** and must be treated before it is returned to the environment or reused.

Most urban households are connected to sewer systems that transport wastewater to centralized treatment facilities. In the United States, roughly 80% of households are connected to municipal sewer systems.

Some older cities use **combined sewer systems**, in which sewage and stormwater runoff are transported through the same pipes. During heavy rainfall, these systems can become overwhelmed and release untreated sewage into nearby rivers, lakes, and coastal waters. Modern communities typically use **separate sewer systems**, which transport sewage and stormwater through different pipe networks. This reduces the likelihood that untreated sewage will enter aquatic ecosystems during storms.

Although major investments have improved wastewater infrastructure, sewer overflows still occur and remain an important source of water pollution.

Wastewater Treatment

Wastewater treatment removes contaminants from sewage before treated water is discharged into the environment or reused. A typical wastewater treatment plant includes several stages (figure $10.6.a$).

Primary Treatment

During **primary treatment**, wastewater is held in settling tanks where gravity separates materials based on density. Heavier solid particles sink to the bottom and form **sludge**, while lighter materials such as oils, grease, and floating debris rise to the surface and form **scum**. Both sludge and scum are removed for further processing. Primary treatment removes much of the suspended material in wastewater but only a portion of its biological oxygen demand (BOD).

Secondary Treatment

During **secondary treatment**, oxygen is added to wastewater to support the growth of **aerobic bacteria**, which consume and decompose dissolved organic matter. This stage removes much of the remaining organic pollution and can reduce biological oxygen demand by up to 90% when combined with primary treatment.

Tertiary Treatment

Although primary and secondary treatment remove most organic material, they do not remove all nutrients. **Tertiary treatment** provides additional purification by removing nitrogen, phosphorus, and other dissolved substances that can contribute to eutrophication and water-quality problems downstream. Some treatment facilities also use advanced filtration technologies to remove additional contaminants.

Disinfection

Before treated wastewater is discharged or reused, it undergoes **disinfection**. Chlorine, ozone, ultraviolet (UV) light, or other treatments are used to kill disease-causing microorganisms that may remain in the water.

Sludge Treatment and Resource Recovery

The sludge removed during wastewater treatment is often processed in large tanks called **digesters**. In these tanks, **anaerobic microorganisms** break down organic matter in the absence of oxygen through a process known as **anaerobic digestion**. This process reduces the volume of waste and produces **methane**, a renewable energy source that can be captured and used to generate heat or electricity.

Discharge and Reuse

After treatment, the water may be released into rivers, lakes, wetlands, or coastal waters. Increasingly, treated wastewater is viewed as a valuable resource and may be reused for irrigation, habitat restoration, groundwater recharge, or advanced water-purification programs that produce drinking water.




Figure 10.6.*a*: Major steps in wastewater treatment.Image by Leonard G. (CC-BY-SA).

Resource Recovery, Water Recycling, and Circular Water Systems

Modern wastewater treatment plants are increasingly designed not only to remove pollutants but also to recover valuable resources. Rather than viewing wastewater as a waste product that must be disposed of, many communities now recognize it as a potential source of water, nutrients, and renewable energy.

One of the most important resource-recovery strategies is **water recycling**, in which highly treated wastewater is reused instead of being discharged to the environment. Recycled water can be used for landscape irrigation, industrial processes, habitat restoration, groundwater recharge, and, after advanced purification, even drinking-water supplies. By reusing water, communities can reduce demand on rivers, reservoirs, aquifers, and imported water sources while increasing resilience to drought.

Wastewater treatment can also recover nutrients and energy. Nitrogen and phosphorus removed during treatment can potentially be reused, reducing reliance on synthetic fertilizers. In addition, methane produced during sludge digestion can be captured and used as a renewable energy source, helping offset the energy demands of treatment facilities.

Example 10.6.1

A local example is the Point Loma Wastewater Treatment Plant in San Diego. The facility captures methane generated during sludge digestion and uses it to help power plant operations. Excess biogas is upgraded to renewable natural gas, which can be supplied to the energy grid. Point Loma illustrates how modern wastewater-treatment facilities can simultaneously protect water quality and recover valuable resources.

Example 10.6.2

San Diego is also expanding water recycling through programs such as Pure Water San Diego and the East County Advanced Water Purification Program. These projects use advanced purification technologies, including microfiltration, reverse osmosis, and ultraviolet disinfection, to transform treated wastewater into a reliable local water source. Rather than discharging treated wastewater to the ocean, these programs recycle water to supplement local drinking-water supplies, reduce dependence on imported water, and improve long-term drought resilience.

These local examples of water recycling here in San Diego reflect a broader shift toward **circular water management**, in which water, nutrients, and energy are recovered and reused rather than discarded. As populations grow and climate change places increasing pressure on freshwater supplies, resource recovery and water recycling are becoming increasingly important components of sustainable water management.

Septic Tank System

Not all homes are connected to centralized wastewater-treatment facilities. In many rural areas, wastewater is treated using an individual **septic system** (figure $10.6.b$). A typical septic system includes a sewer line that carries wastewater from the home to a **septic tank**, where solids settle out and are broken down by microorganisms. The liquid portion of the wastewater then flows to a **drain field**, a network of perforated pipes buried in the soil that allows partially treated water to slowly infiltrate into the ground.

Within the septic tank, solids accumulate as **sludge** and are decomposed by **anaerobic microorganisms**, which break down organic matter in the absence of oxygen. As wastewater moves through the drain field, soil particles and naturally occurring microorganisms help remove additional contaminants before the water reaches groundwater. Together, these processes provide a simple and effective method of wastewater treatment for homes that are not connected to municipal sewer systems.

Proper installation and maintenance are essential for septic systems to function effectively. When septic systems fail because of poor maintenance, damaged components, or unsuitable soil conditions, untreated wastewater can contaminate nearby groundwater and surface waters with pathogens, nutrients, and other pollutants.




Figure $10.6.b$: A septic tank system treats wastewater from individual homes. Wastewater flows from the house into a septic tank, where solids settle to the bottom as sludge and are broken down by anaerobic bacteria. The partially treated liquid then flows into the drain field, where it slowly infiltrates into the soil and receives additional treatment by soil microorganisms before entering groundwater. Image by United States Geological Survey (public domain).

Addressing Water Treatment Globally

Many communities around the world still lack access to safe drinking water and adequate sanitation infrastructure. The World Health Organization estimates that investments in clean water and sanitation yield substantial benefits through improved health, increased productivity, reduced healthcare costs, and fewer deaths from waterborne disease.

In regions where conventional treatment facilities are not feasible, lower-cost alternatives can significantly improve water quality. Examples include household filtration systems, chlorination, solar disinfection, and constructed wetlands.

Constructed wetlands are engineered ecosystems that use wetland plants, soils, and microorganisms to remove pollutants from wastewater. They are often less expensive to build and maintain than conventional treatment plants while also providing wildlife habitat and other ecosystem benefits.

Example 10.6.3

One notable example is the Arcata Marsh and Wildlife Sanctuary in Northern California, which integrates wastewater treatment with wetland habitat restoration.

Think About It

For much of the twentieth century, wastewater treatment focused on removing pollutants before treated water was discharged into the environment. Today, many treatment facilities are shifting toward resource recovery and water recycling.

Reflect: Should wastewater be viewed primarily as a waste product to dispose of or as a resource to recover? What environmental, economic, and social benefits might result from treating wastewater as part of a circular water system rather than a linear one?

Attributions

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Additional sources:

Arcata Marsh & Wildlife Sanctuary. Accessed 2-3-2026.

Modified by Melissa Ha from the following sources:

- Water Treatment from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Sewage Treatment from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Former National Compliance Initiative: Keeping Raw Sewage and Contaminated Stormwater Out of Our Nation's Waters. 2020. EPA. Accessed 01-18-2021. (public domain)

This page titled 10.6: Water Treatment- Protecting Drinking Water and Treating Wastewater is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 20.1.2: Water Treatment by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

10.7: Preventing Water Pollution and Restoring Watersheds

Preventing Water Pollution and Restoring Watersheds

As discussed earlier in this chapter, water moves through connected watersheds. Pollutants released anywhere within a watershed can be transported downstream through rivers, groundwater, wetlands, estuaries, and coastal ecosystems. As a result, managing water quality requires more than treating pollution at a single location. It requires managing the entire watershed.

Environmental Regulations

One of the most effective tools for improving water quality is environmental regulation. During the early twentieth century, rapid industrialization in the United States led to widespread contamination of rivers, lakes, and coastal waters. Industries often discharged untreated wastes directly into waterways, and many rivers became heavily polluted. One of the most famous examples was the Cuyahoga River in Ohio, which caught fire multiple times because oil and industrial wastes floating on the river's surface ignited (figure $10.7.a$).

The 1969 Cuyahoga River fire helped draw national attention to water pollution and contributed to the passage of the **Clean Water Act (CWA)** in 1972. The primary goal of the Clean Water Act is to restore and maintain the chemical, physical, and biological integrity of the nation's waters. The Act regulates point-source pollution by establishing permits and discharge standards for wastewater treatment plants, industries, and other identifiable pollution sources.

Since its passage, water quality in many rivers, lakes, and estuaries has improved substantially. However, non-point source pollution, which originates from many dispersed sources across a watershed, remains one of the most difficult water-quality challenges to address.



Figure $10.7.a$: Cuyahoga River on fire in 1969. Image from USEPA Environmental-Protection-Agency (public domain).

Remediation: Cleaning Up Contamination

Despite efforts to prevent pollution, contaminants sometimes enter the environment and require cleanup. **Remediation** is the process of removing pollutants or reducing their harmful effects.

Bioremediation

Bioremediation uses living organisms such as bacteria, fungi, and plants to break down, remove, or neutralize contaminants. Because it relies on natural biological processes, bioremediation is often less disruptive and less expensive than many mechanical cleanup methods.

Microorganisms are widely used in wastewater treatment plants to decompose organic wastes, but they can also be used to treat contaminated soils and groundwater. Certain microorganisms can transform toxic chemicals into less harmful forms, while others can degrade petroleum products, pesticides, and industrial pollutants.

A specialized form of bioremediation known as **phytoremediation** uses plants to absorb contaminants from soil and water. Over time, these plants may accumulate nutrients, metals, or other pollutants, helping restore contaminated sites.

One of the most common applications of bioremediation is the cleanup of oil spills. Naturally occurring bacteria can use petroleum hydrocarbons as a food source, breaking them into simpler compounds. Following large spills such as the Exxon Valdez oil spill in Alaska, cleanup efforts often include stimulating the growth of oil-degrading microorganisms by adding nutrients to the affected area (figure $10.7.b$).



(a) (b)

Figure $10.7.b$: (a) Cleaning up oil after the Exxon Valdez spill in Alaska (1989), the workers hosed oil from beaches and then used a floating boom to corral the oil, which was finally skimmed from the water surface. Some species of bacteria are able to solubilize and degrade the oil. (b) One of the most catastrophic consequences of oil spills is the damage to fauna. (credit a: modification of work by NOAA; credit b: modification of work by GOLUBENKOV, NGO: Saving Taman)

Bioremediation is not a quick solution, but it can significantly accelerate the natural recovery of contaminated environments.

Chemical and Physical Remediation

Some pollutants cannot be effectively removed through biological processes alone. **Chemical remediation** uses chemical reactions to neutralize contaminants or convert them into less harmful forms. For example, limestone barriers are often used to treat acid mine drainage by increasing pH and causing dissolved metals to precipitate out of solution.

Physical remediation removes contamination through methods such as filtration, excavation, containment, or **pump-and-treat** systems. In a pump-and-treat system, contaminated groundwater is pumped to the surface, treated, and then discharged or returned to the aquifer. Although physical remediation can be effective, it is often expensive and energy intensive.

Watershed Management: Preventing Pollution at Its Source

While remediation can reduce environmental damage, preventing pollution before it enters waterways is often the most effective long-term strategy.

Watershed management recognizes that water quality is influenced by activities throughout an entire drainage basin. Agricultural practices, urban development, transportation systems, wastewater infrastructure, habitat restoration, and

conservation efforts can all affect the movement of water, sediment, nutrients, and pollutants. Rather than focusing on individual pollution sources, watershed management seeks to improve the health and resilience of an entire interconnected system.

Watershed management focuses on improving land-use practices throughout an entire watershed to reduce runoff, erosion, and pollutant transport. Because non-point source pollution originates from many dispersed locations, watershed-scale approaches are often more effective than trying to regulate each source individually.

As discussed earlier in this chapter, activities occurring upstream can strongly influence water quality downstream. The impacts of pollution in the Tijuana River Watershed and nutrient runoff throughout the Mississippi-Atchafalaya River Basin illustrate how actions in one location can affect ecosystems and communities many kilometers away.

Riparian Buffers

One of the most effective methods of watershed protection is maintaining or restoring **riparian zones**, the vegetated areas adjacent to rivers, streams, lakes, and wetlands (figure $10.7.c$).

Riparian vegetation improves water quality in several ways. Plant roots stabilize soil and reduce erosion, preventing sediment from entering waterways. Vegetation absorbs excess nutrients such as nitrogen and phosphorus before they reach rivers and lakes, reducing the risk of eutrophication. Trees provide shade that helps keep water cool, increasing dissolved oxygen levels and improving habitat for aquatic organisms. Riparian vegetation also slows runoff and promotes infiltration, allowing pollutants to be filtered by soils while helping recharge groundwater supplies.

Because they improve water quality, reduce flooding, stabilize stream banks, and provide wildlife habitat, riparian buffers are often considered one of the most cost-effective forms of pollution prevention.






Figure 10.7.c: Top: Riparian areas reduce water pollution by absorbing nutrients, keeping water cool, limiting erosion, and promoting infiltration. Image by United States Bureau of Land Management, (public domain). [https://www.blm.gov]. Bottom: Riparian buffers can be designed to balance environmental protection and human land use. For example, some buffer systems combine natural vegetation closest to the stream with carefully managed trees or shrubs farther from the water. These layered buffers help reduce pollution while allowing limited agricultural or economic uses of the landscape. Image and caption by USDA National Agroforestry Center (public domain).

Wetland Restoration

Wetlands are often called nature's kidneys because they filter sediments, nutrients, and other pollutants while supporting biodiversity and improving ecosystem resilience.

Wetlands are among the most effective natural water-treatment systems on Earth. As water moves slowly through wetlands, sediments settle out and plants and microorganisms absorb nutrients and other pollutants. Wetlands can improve water quality, reduce flooding, provide wildlife habitat, and promote groundwater recharge. Restoring degraded wetlands can therefore address several environmental problems at once while strengthening watershed resilience.

Example 10.7.1

Location Connection: Kendall-Frost Marsh and ReWild Mission Bay

One local example of wetland restoration is Kendall-Frost Marsh Reserve in Mission Bay, one of the last remaining coastal salt marshes in San Diego County. Coastal wetlands such as Kendall-Frost Marsh provide habitat for fish, birds, invertebrates, and other wildlife while also improving water quality by trapping sediments and absorbing excess nutrients before they reach coastal waters.

Restoration efforts at Kendall-Frost Marsh are part of a broader collaboration known as ReWild Mission Bay, a long-term project that seeks to transform portions of Mission Bay from heavily altered landscapes into more natural wetlands and coastal habitats. The project aims to restore tidal wetlands, improve wildlife habitat, increase climate resilience, and enhance ecosystem services such as water filtration, flood protection, and carbon storage.

As sea levels rise and urban development continues to pressure coastal ecosystems, wetland restoration projects such as Kendall-Frost Marsh and ReWild Mission Bay demonstrate how communities can work to recover ecological functions that have been lost over time. These projects illustrate an important principle of watershed management: protecting water quality often involves restoring natural systems that can filter pollutants, store floodwaters, support biodiversity, and increase ecosystem resilience.

Green Infrastructure

Many communities now use **green infrastructure** to manage stormwater and reduce pollution. Unlike conventional stormwater systems that rapidly move water into drains and pipes, green infrastructure works with natural processes to slow, filter, and absorb runoff. Green infrastructure is increasingly viewed as a form of **natural infrastructure** because it uses ecological processes to provide services that might otherwise require expensive engineered systems.

Examples include rain gardens, bioswales, permeable pavement, urban wetlands, and green roofs. These systems reduce runoff, remove pollutants, promote infiltration, and help recharge groundwater. They also provide additional benefits such as reducing urban flooding, supporting biodiversity, and improving community resilience to climate change.

Many of the stormwater-management and groundwater-recharge strategies discussed earlier in this chapter also function as pollution-prevention tools by keeping contaminants out of waterways in the first place.

Healthy watersheds are not maintained only through regulations and treatment plants. They also depend on ecological restoration, community involvement, and long-term stewardship.

Example 10.7.1

Local Connection: Rewilding the San Diego River

The San Diego River Park Foundation promotes a watershed-scale vision known as "River Park from the Mountains to the Sea." The long-term goal is to reconnect, restore, and protect the San Diego River corridor from its headwaters near Julian and the Cuyamaca Mountains to its mouth at Ocean Beach, where the river enters the Pacific Ocean. Rather than viewing the river as a series of disconnected parks, channels, and open spaces, this approach recognizes the river as a single interconnected watershed whose ecological health depends on conditions throughout the entire system.

Restoration efforts along the San Diego River include removing invasive species, planting native vegetation, improving wildlife habitat, restoring riparian areas, managing erosion, and increasing public access to the river corridor (Figure $10.7.d$). These projects help improve water quality by reducing sediment and pollutant runoff, stabilizing stream banks, promoting groundwater recharge, and providing habitat for birds, fish, insects, and other wildlife.

The San Diego River illustrates an important principle of watershed management: protecting water quality requires more than managing the river itself. Land-use decisions throughout the watershed influence the movement of water, nutrients, sediments, and pollutants. By restoring natural ecosystem functions along the river corridor, rewilding efforts help support both ecological health and the communities that depend on the watershed.

Similar watershed-scale approaches are being used throughout Southern California, including efforts to improve water quality, restore habitat, and reduce pollution in the Tijuana River Watershed. These projects demonstrate that effective watershed management often requires cooperation among communities, agencies, and governments across an entire drainage basin.



Figure 10.7.*d*:Orignal Map of San Diego River Improvement Project. Restoration and conservation efforts along the San Diego River aim to reconnect habitat, improve water quality, support biodiversity, and create a continuous river park from the mountains near Julian to the Pacific Ocean at Ocean Beach. Rewilding projects demonstrate how watershed restoration can benefit both people and ecosystems. (Copyright; author By Wimmer Yamada & Caughey, Public Domain, [https://commons.wikimedia.org/w/inde...rid=151734059]

From Pollution Control to Water Stewardship

Environmental regulations, water treatment, remediation, watershed restoration, and green infrastructure all help protect water quality. However, these approaches operate at different points within a system. Some focus on removing pollution after it has entered the environment, whereas others prevent pollution from occurring in the first place.

Increasingly, sustainable water management combines both approaches. Effective water stewardship requires thinking at the scale of entire watersheds and recognizing that rivers, groundwater, wetlands, estuaries, ecosystems, and human communities are closely interconnected. Protecting water quality therefore depends not only on cleaning up pollution, but also on preventing contamination, restoring ecosystems, and managing watersheds as integrated systems.

Think About It

Throughout this chapter, you explored water pollution, wastewater treatment, water recycling, groundwater recharge, riparian restoration, bioswales, and watershed management.

Reflect: Which strategies are primarily designed to clean up pollution after it occurs, and which are intended to prevent pollution before it enters waterways? If your community could invest in only one approach, would you prioritize pollution cleanup, wastewater treatment, ecosystem restoration, green infrastructure, or pollution prevention? What evidence from this chapter supports your decision?

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Sources:

- Kendall-Frost Marsh Reserve
- ReWild Mission Bay
- San Diego River Park Foundation

Modified by Melissa Ha from the following sources:

- Water Contamination from *An Introduction to Geology* by Johnson et al. (licensed under CC-BY-NC-SA(opens in new window))

- Water Treatment and Bioremediation from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 10.7: Preventing Water Pollution and Restoring Watersheds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 20.1.3: Mitigating Water Pollution by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

11: WASTE, MATERIALS, AND THE CIRCULAR ECONOMY

Where is "away"?

A package of "flushable" wipes seems like a simple convenience. After all, if a product is labeled *flushable*, it should safely disappear after it is flushed down a toilet, right? We throw things "away," but where exactly is *away*?

In reality, many wastewater professionals, plumbers, and utility managers argue that so-called flushable wipes do not break down quickly enough in water and can accumulate in pipes, sewer systems, and wastewater-treatment facilities. Materials that leave our homes do not simply disappear. Instead, they enter waste-management systems where they must be collected, transported, processed, and sometimes removed by hand.

Around the world, flushable wipes have contributed to costly pipe blockages, sewer overflows, and large masses of congealed waste known as *fatbergs*. Although manufacturers may label products as flushable, there are few universal standards that determine what qualifies for that claim. As a result, a product designed for convenience can create environmental, economic, and infrastructure challenges long after it leaves a consumer's hands.

The story of flushable wipes highlights a broader question explored throughout this chapter: *What happens to the things we throw away, flush, recycle, compost, or discard once they leave our homes?* Understanding the answer requires looking beyond the trash can or toilet and examining how modern societies manage materials, generate waste, and design systems that can either discard resources or keep them circulating in a more sustainable, circular economy.



Figure 11.*a*: Wipe isle at the store. Image by 維基小霸王(opens in new window) in Wikimedia Commons(opens in new window) (CC-BY-SA4.0(opens in new window))

Solid Waste and a Circular Economy

Every day, individuals, businesses, and communities generate large quantities of **solid waste**, including food scraps, paper, plastics, metals, electronics, and construction materials. As populations grow and consumption increases, managing this waste has become an important environmental challenge. Improperly managed waste can contaminate soil and water, harm wildlife, release greenhouse gases, and contribute to broader environmental problems.

For much of modern history, societies have followed a **linear economy** in which materials are extracted, manufactured into products, used, and then discarded. Increasingly, environmental scientists, engineers, and policymakers are promoting a **circular economy**, which seeks to reduce waste by designing products to be reused, repaired, remanufactured, recycled, or composted (figure 11.*b*). Rather than viewing discarded materials as useless trash, a circular-economy approach treats many waste streams

as valuable resources that can remain in use for as long as possible.

Understanding where solid waste comes from, how it is managed, and how waste can be reduced is an important part of building more sustainable communities. In this chapter, we will examine the environmental impacts of solid waste, explore strategies such as source reduction, reuse, recycling, composting, and waste-to-energy systems, and evaluate how circular-economy approaches can help reduce the amount of material sent to landfills.




Figure 11.*b*: Linear versus circular economy. By Original: PennineFoxhound Vector: ClonewayxCatherine Weetman, CC BY-SA 4.0 <https://creativecommons.org/licenses/by-sa/4.0>, prostřednictvím Wikimedia Commons - Own work based on: Linear versus circular.jpg by PennineFoxhound, CC BY-SA 4.0, [//commons.wikimedia.org/w/index.php?curid=10823214 0]

Guiding question for this chapter

How can communities meet human needs while generating less waste and keeping materials circulating within the economy rather than disposing of them in landfills or the environment?

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Rachel Schleiger (CC-BY-NC(opens in new window)).

This page titled 11: Waste, Materials, and the Circular Economy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

- 19: Solid Waste Management by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

11.1: Waste Generation

Waste in Natural and Human Systems

In natural ecosystems, there is no such thing as "waste." Matter and nutrients continuously cycle through biological and geological processes. Dead organisms are decomposed by fungi, bacteria, and detritivores, and the materials they contain become nutrients that support new life. In this sense, ecosystems operate as largely circular systems in which materials are continually reused.

Human societies often function differently. Many modern economies follow a **linear model** in which materials are extracted, manufactured into products, used, and then discarded (figure $11.1.a$). This "take-make-waste" approach generates large quantities of **solid waste**, the discarded materials that humans consider no longer useful or valuable.

Many products are designed to be durable, resistant to decomposition, or difficult to repair or recycle. As a result, materials that could potentially be reused often accumulate in landfills, waterways, and the environment. However, what is considered waste in one context may be a valuable resource in another. Increasingly, scientists, engineers, and policymakers are exploring ways to reduce waste and keep materials in circulation through reuse, repair, recycling, composting, and circular-economy strategies.



Figure $11.1.a$: Trash is a uniquely human concept. Image by Piqsels (opens in new window)(public domain).

The quantity and composition of waste provide insights into how societies use materials and energy. Examining where waste comes from and what it is made of can help identify opportunities to reduce waste and move toward more circular systems.

> **Think About It**
>
> A fallen tree in a forest is eventually decomposed and recycled into the ecosystem. Why do many human-made materials accumulate as waste instead of being reused in the same way?

Types of Solid Waste

Solid waste includes many different materials with varying environmental impacts.

Some materials are **biodegradable**, meaning they can be broken down by microorganisms into simpler substances. Food scraps, paper, yard trimmings, and untreated wood are examples of biodegradable wastes (figure $11.1.b$).

Other materials are **nondegradable** or break down very slowly. Plastics, synthetic fabrics, glass, and many manufactured products may persist in the environment for decades or centuries.

Particular attention is often given to **hazardous waste**, which includes materials that are toxic, carcinogenic (cause cancer), mutagenic (cause DNA mutations), teratogenic (cause birth defects), highly flammable, corrosive, reactive, or explosive. Because these substances can pose risks to human health and ecosystems, they often require specialized handling, recycling, and disposal procedures.

Common examples include batteries, fluorescent bulbs, solvents, paints, pesticides, and discarded electronic devices.

Electronic waste, or **e-waste**, is one of the fastest-growing waste streams globally. Although electronics often contain valuable materials such as copper, gold, silver, and rare earth elements, they may also contain hazardous substances that require specialized recycling and disposal methods.



Figure 11.1.*b*: Disposable utensils made from wood are biodegradable and can be broken down by microorganisms, whereas conventional plastic utensils may persist in the environment for many years. Image by [pixabay.com/photos/cutlery-utensils-kitchenware-knife-5507916/](opens in new window) [BRRT] (Pixabay license).

Where Does Waste Come From?

When many people think about waste, they think about trash cans, recycling bins, and landfills. However, household garbage represents only a small portion of total waste generation.

Globally, agriculture, industry, construction, mining, and energy production generate far more waste than households. These waste streams are collectively known as **non-municipal solid waste** and include materials such as mining residues, construction debris, industrial byproducts, wastewater-treatment sludge, and agricultural wastes.

The waste generated by homes, businesses, schools, and local governments is called **municipal solid waste (MSW)**. Although MSW receives considerable attention because people interact with it daily, it accounts for only a small fraction of total waste generation (figure 11.1.*c*).




Figure 11.1.*c*: Global waste generation by category. Agriculture and industry generate far more waste than municipal sources. Image from What a Waste 2.0 : A Global Snapshot of Solid Waste Management to 2050 (CC-BY)

Municipal Solid Waste

The world generates more than 2 billion metric tons (2 trillion kg) of municipal solid waste each year, and waste generation is projected to continue increasing as populations grow, urbanization expands, and consumption rises. In the United States, municipal solid waste generation averages about 4.9 pounds (2.2 kg) per person per day, among the highest rates in the world.

Municipal solid waste includes a diverse mix of materials, including paper and cardboard, food waste, plastics, yard trimmings, metals, glass, wood, and textiles. Although the composition of waste varies among communities, studies consistently show that a large fraction of municipal waste could potentially be reduced, reused, repaired, recycled, or composted rather than being disposed of in landfills.

This observation highlights an important challenge in waste management: many materials currently treated as waste still retain value and could remain in use through more circular approaches to production and consumption.

What Happens to Municipal Solid Waste?

The composition of municipal solid waste provides insight into what communities throw away, but it does not reveal what happens to those materials after disposal. Some materials are recycled, composted, or used to recover energy, while others are landfilled or incinerated.

Comparing the composition of municipal solid waste with the materials that are recovered through recycling and composting reveals important patterns. Some materials, such as paper and cardboard, are recycled at relatively high rates because collection systems and markets for recycled paper are well established. Metals are also commonly recycled because they retain significant economic value. In contrast, plastics make up a large fraction of the waste stream but are recycled at much lower rates. Food waste and yard trimmings can be composted, yet substantial quantities are still disposed of in landfills.

The two pie charts in figure (11.1.*d*) compare the composition of municipal solid waste generated in the United States with the materials recovered through recycling and composting. Together, these figures illustrate an important principle of waste management: materials that enter the waste stream do not all have the same likelihood of being recovered. Understanding why some materials are successfully recycled while others are commonly discarded can help identify opportunities to reduce waste and move toward a more circular economy.




Figure 11.1.*d*: Top: Composition of municipal solid waste generated in the United States before recycling. Bottom: Composition of materials recovered through recycling and composting in the United States. Images from the U.S. Environmental Protection Agency; (public domain).

Think About It

Compare the pie charts in Figure (11.1.*d*). Which materials appear to be recycled or composted at high rates relative to the amount generated? Which materials are underrepresented in the recycling stream? What factors might explain these differences?

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Systems of Waste Management from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (CC-BY). Download for free at CNX.

- Solid Waste form *AP Environmental Science* by University of California College Prep (CC-BY). Download for free at CNX.
- Kaza, Silpa; Yao, Lisa C.; Bhada-Tata, Perinaz; Van Woerden, Frank. 2018. What a Waste 2.0 : A Global Snapshot of Solid Waste Management to 2050. Urban Development;. Washington, DC: World Bank. © World Bank. (CC-BY)

This page titled 11.1: Waste Generation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 19.1: Waste Generation by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

11.2: Disposal

Waste Disposal: Where Does Our Trash Go?

After efforts to reduce, reuse, repair, recycle, and compost materials, some waste still requires disposal. Historically, societies have used a variety of methods to manage unwanted materials. Today, the most common disposal methods are **open dumping, sanitary landfilling, and incineration** (figure 11.2.a).

Each approach has environmental tradeoffs. Modern waste-management systems attempt to reduce pollution, protect human health, and minimize impacts on air, water, soil, wildlife, and climate.

Interactive Element

Where Does Our Trash Go?

When you throw something away, where does it actually end up? Explore the interactive figure below to see how municipal solid waste is managed around the world. Compare how different countries handle waste through landfilling, recycling, composting, incineration, and open dumping, and look for patterns related to income, infrastructure, and environmental policy.

Try It!

- Select different countries and compare how they manage municipal solid waste.
- Which countries rely most heavily on landfills? Which divert larger amounts of waste through recycling or composting?
- How does waste management differ between high-income and low-income countries?
- Look for regions where open dumping remains common. What environmental and public-health challenges might result?

Think About It: If a community generates the same amount of waste but manages it differently, how might the environmental impacts change? Which waste-management practices appear most consistent with the goal of moving toward a circular economy?

Interactive Content

View this Interactive Content online

Figure 11.2.a How solid municipal waste is managed. **Data source:** Anshassi and Townsend (2025). OurWorldinData.org/waste-management | (CC-BY-4.0)

Open Dumps

An **open dump** is the simplest form of waste disposal, in which trash is deposited in a designated area with little or no environmental protection (figure 11.2.b). Open dumps remain common in some parts of the world but are illegal in many countries

because of their impacts on human health and the environment.

Open dumps can attract rodents, insects, and other organisms that spread disease. Rainwater flowing through waste creates **leachate**, a contaminated liquid that may carry heavy metals, pesticides, persistent organic pollutants, and other hazardous substances into soil, surface water, and groundwater.

Because waste is exposed to the atmosphere, open dumps may also release smoke from fires, produce unpleasant odors, and contribute to air pollution.



Figure $11.2.b$: An open dump in Vietnam. Image by Julien Belli (CC-BY).

Sanitary Landfills

Today, the most common method of municipal waste disposal in the United States is the **sanitary landfill**.

Unlike open dumps, sanitary landfills are engineered facilities designed to isolate waste from the surrounding environment (figure $11.2.c - d$).

Waste is compacted and buried in layers, while liners, caps, and collection systems help reduce contamination of soil and groundwater.

Modern sanitary landfills are engineered to minimize environmental impacts. They use **impermeable liners** to reduce the movement of leachate into surrounding soils and groundwater, **leachate collection and treatment systems** to capture contaminated liquids, and **groundwater-monitoring wells** to detect potential leaks. Waste is routinely covered with soil or other materials to reduce odors, litter, and access by animals, while **landfill gas collection systems** capture methane and other gases produced during decomposition. These design features make sanitary landfills much safer than open dumps, although they still require long-term monitoring and maintenance.

As organic waste decomposes in oxygen-poor conditions, microorganisms produce **landfill gas**, which consists largely of methane and carbon dioxide. Methane is a potent greenhouse gas, but many modern landfills capture it and use it to generate electricity or heat.



Figure 11.2.*c*: A sanitary landfill in 1972. Image by Bill Shrout/EPA (public domain).




Figure 11.2.*d*: Components of a sanitary landfill. Image by EPA/National Energy Education Development Project (public domain).

Example 11.2.1

Local Connection: Miramar Landfill

The Miramar Landfill in San Diego captures methane generated during decomposition and uses it to produce renewable energy. This system reduces greenhouse-gas emissions while generating electricity from materials that would otherwise be wasted.

Challenges of Landfills

Although sanitary landfills are much safer than open dumps, they still have environmental and practical limitations. Landfills require large areas of land and must be monitored for decades to ensure that leachate does not contaminate surrounding soil and groundwater. As organic waste decomposes, methane can escape into the atmosphere, contributing to climate change if it is not captured and used as an energy source. In addition, many materials buried in landfills retain value and could potentially be reused, repaired, recycled, or composted instead of being discarded. As populations grow and waste generation increases, many communities are also facing a shortage of available landfill space, prompting interest in alternative waste-management

strategies that keep materials in circulation rather than burying them.

As a result, many communities are seeking to reduce reliance on landfills through recycling, composting, and circular-economy strategies.

Incineration and Waste-to-Energy

Incineration uses high temperatures to burn waste. Modern waste-to-energy facilities use the heat generated from combustion to produce steam, which can then generate electricity (figure $11.2.e - f$).



Figure $11.2.e$: Wheelabrator Technologies' Waste-to-Energy plant in Saugus, Massachusetts has been in service since 1975. Image and caption (modified) from Fletcher6 (CC-BY).




Figure $11.2.f$: A schematic diagram of a waste incineration facility (waste-to energy system). A grabber collects the waste and transfers it into the incinerator. Heat from combusting the trash is used to produce steam and generate electricity. Emissions (flue gases) pass through a scrubber and particulate removal system to limit pollution before the gases are released through a chimney. Ash remains after incineration. Image by Kaza, Silpa; Bhada-Tata, Perinaz. 2018. Decision Maker's Guides for Solid Waste Management Technologies. Urban Development Series Knowledge Papers;. World Bank, Washington, DC. © World Bank. (CC-BY)

One advantage of **incineration** is that it dramatically reduces the volume of waste entering landfills. In some cases, incineration can reduce waste volume by as much as 85%. It can also be used to generate electricity through **waste-to-energy** systems.

However, incineration has significant drawbacks. Even with modern pollution-control technologies, incinerators may release air pollutants such as particulate matter, nitrogen oxides, sulfur dioxide, heavy metals, dioxins, and greenhouse gases. In addition, combustion produces ash that may contain toxic substances and requires careful management and disposal.

Two common waste-to-energy approaches are **mass-burn incineration** and **refuse-derived fuel (RDF)** systems. In mass-burn incineration, mixed municipal waste is burned directly to generate heat and electricity. In contrast, RDF systems remove recyclable and noncombustible materials before processing the remaining combustible waste into fuel pellets. Although RDF systems recover more recyclable materials than mass-burn facilities, both approaches ultimately destroy potentially useful materials through combustion rather than keeping them in circulation through reuse, repair, recycling, or composting.

Comparing Disposal Methods

Each disposal method presents environmental tradeoffs.

Method	Advantages	Disadvantages
Open Dumps	Low cost, simple	Disease, leachate, air pollution, habitat degradation
Sanitary Landfills	Better environmental protection, methane recovery possible	Land use, methane emissions, long-term monitoring
Incineration	Reduces waste volume, generates electricity	Air pollution, toxic ash, loss of recoverable materials

Because all disposal methods have environmental costs, many sustainability experts argue that the best solution is to avoid creating waste in the first place.

From Disposal to Circular Systems

Traditional waste management focuses on what happens after materials are discarded. Increasingly, communities are shifting toward a **circular economy**, which seeks to keep materials in use through reduction, reuse, repair, recycling, and composting rather than disposing of them in landfills or incinerators.

In this framework, disposal is viewed as a last resort rather than the primary solution to waste generation.

> **Think About It**
>
> A landfill equipped with methane recovery can generate renewable energy, while a waste-to-energy incinerator can produce electricity from trash.
>
> **Reflect:** Are these examples of sustainable waste management, or do they primarily address problems after waste has already been created? How do they compare with strategies that prevent waste from being generated in the first place?

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Systems of Waste Management from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (CC-BY). Download for free at CNX.
- Solid Waste from *AP Environmental Science* by University of California College Prep (CC-BY). Download for free at CNX.
- Biomass explained: Landfill gas and biogas. 2020. EIA. Accessed 01-18-2021 (public domain).

This page titled 11.2: Disposal is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 19.2: Waste Disposal by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

11.3: Environmental Impacts of Solid Waste

Environmental Impacts of Solid Waste

Solid waste affects ecosystems, wildlife, human health, and climate at local, regional, and global scales. While improperly managed waste can contaminate land, water, and air, even properly managed waste streams often require energy, land, and long-term environmental oversight. Understanding these impacts can help communities make more informed decisions about waste reduction, recycling, composting, and circular-economy strategies.

Land and Water Pollution

Improperly managed waste can contaminate soil, surface water, and groundwater. Open dumps and poorly managed disposal sites often generate **leachate**, a contaminated liquid produced when water passes through waste and dissolves pollutants. Leachate can contain heavy metals, nutrients, hazardous chemicals, and other contaminants that may harm ecosystems and drinking-water supplies.

Litter and illegal dumping can also degrade habitats, reduce aesthetic value, and introduce pollutants into waterways. Because water transports contaminants across landscapes, waste discarded in one location may ultimately affect rivers, wetlands, estuaries, and coastal ecosystems far away.

Climate Impacts of Solid Waste

Solid waste can also contribute to climate change. As organic materials such as food scraps, paper, and yard trimmings decompose in landfills, microorganisms produce methane, a greenhouse gas that traps significantly more heat than carbon dioxide over short time scales. Although many landfills capture landfill gas, some methane still escapes into the atmosphere.

Producing, transporting, and disposing of products also requires energy and materials. As a result, reducing waste and extending product lifetimes can lower greenhouse-gas emissions throughout the economy.

Plastic Pollution and Marine Debris

Historically, some coastal communities disposed of waste by dumping it into the ocean. In this practice, barges transported garbage offshore and released it directly into marine waters. Because ocean dumping created severe environmental problems, it has been banned in the United States. However, large amounts of waste still reach the ocean through littering, illegal dumping, stormwater runoff, rivers, and wind-blown debris (figure $11.3.a$).



Figure 11.3.*a*: Left: A garbage barge leaves Manhattan. Right: Marine debris at the beach. Left image and caption by EPA (public domain), and right image and caption by NOAA (public domain).

Once in the environment, plastics can persist for decades or even centuries. Rather than completely biodegrading, most plastics gradually fragment into smaller pieces through exposure to sunlight, physical abrasion, and weathering. These fragments, known as **microplastics** when they are smaller than 5 millimeters, are now found throughout the world's oceans, freshwater systems, soils, and even the atmosphere.

Garbage Patches

Ocean currents transport floating debris across vast distances. In some regions, circular ocean-current systems known as **gyres** concentrate floating plastics and other debris into areas commonly called **garbage patches** (figure 11.3.*b*).

The most famous example is the Great Pacific Garbage Patch located between California and Hawaii. Despite their name, garbage patches are not floating islands of trash. Instead, they consist of large concentrations of plastic fragments and microplastics distributed throughout the water column. Much of this material is too small to be seen from the surface and may extend several meters below the water.




Figure 11.3.*b*: Locations of major garbage patches in the Pacific Ocean. Image by Boyan Slat (public domain).

Impacts on Wildlife

Plastic pollution harms wildlife in numerous ways. Discarded fishing gear, six-pack rings, ropes, and nets can entangle marine organisms, restricting movement, causing injury, or leading to death. Sea turtles, seabirds, fish, marine mammals, and many other species are vulnerable to entanglement.

Wildlife may also mistake plastic debris for food. Ingested plastic can block digestive systems, damage internal tissues, reduce feeding, and create a false sense of fullness. Plastics can also transport chemicals and contaminants through ecosystems.

One well-known example involves seabirds that feed plastic fragments to their chicks. In some cases, large quantities of plastic accumulate in the digestive system, contributing to starvation or other health problems (figure 11.3.*c*).



Figure 11.3.*c*: Much of this dead albatross's digestive system was filled with plastic debris. Image by Forest and Kim Starr (CC-BY).

Microplastics and Emerging Concerns

Scientists are increasingly concerned about **microplastics**, which originate from the breakdown of larger plastic items as well as products such as synthetic clothing fibers, tire wear particles, and microbeads formerly used in personal-care products.

Microplastics have been detected in marine organisms, freshwater ecosystems, agricultural soils, drinking water, and even human tissues.

Although research is ongoing, the widespread presence of microplastics highlights the challenge of managing durable materials that persist long after disposal. Like PFAS and other synthetic contaminants discussed previously, plastic pollution illustrates how modern materials can continue affecting ecosystems long after they are discarded.

Edible six-pack rings offer a solution to the problem of plastic six-pack rings harming wildlife. You can read more about them here.

Connections to Planetary Boundaries

Plastic pollution is also connected to the Novel Entities planetary boundary. Novel entities are human-made materials and chemicals, such as plastics, pesticides, PFAS, and many industrial compounds, that are released into the environment faster than scientists and regulators can fully understand or manage their impacts. The persistence of plastic waste and microplastics in ecosystems around the world highlights how products designed for short-term use can create environmental consequences that last for decades or even centuries.

Think About It

A plastic bottle discarded on a city street may eventually be transported by wind and stormwater runoff into a creek, carried through a river system, and ultimately deposited in the ocean.

Reflect: At what stage in this journey would it be easiest and most effective to prevent environmental harm? What actions by individuals, businesses, manufacturers, and governments could keep plastic from reaching natural ecosystems in the first place?

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Melissa Ha (CC-BY-NC) and Solid Waste from *AP Environmental Science* by University of California College Prep (CC-BY). Download for free at CNX.

This page titled 11.3: Environmental Impacts of Solid Waste is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 19.3: Solid Waste and Marine Life by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

11.4: From Waste Management to a Circular Economy

The **waste management hierarchy** lists processes to address waste in order of preference with the goals of limiting waste, minimizing environmental impact, and maximizing additional benefits (such as electricity generation or production of recycled goods). It is symbolized as an upside triangle, meaning that the largest volume of waste should be handled using the first process (most preferred), and the smallest volume should be handled using the last process (least preferred). These processes in order of preference are reduction, recycling, energy recovery (such as incineration), treatment, and disposal (figure $11.4.a$). Individuals can exercise the first two processes by applying the four R's (see below). Unfortunately, the U.S. and the world overall, handles a large volume of waste using the least preferred method (disposal). As previously mentioned, about half of municipal solid waste (MSW) in the United States goes to a sanitary landfill for disposal.

In contrast, only about 32% of municipal solid waste (MSW) in the United States is recycled or composted, and approximately 12% is combusted for energy recovery (most recent national estimates from 2018–2021). Waste stream percentages vary widely by region. For example, San Francisco, California diverts around 80% of its waste through recycling and composting programs, while Houston, Texas diverts roughly 18% (2026).




Figure $11.4.a$**: Left**: The waste management hierarchy lists processes for handling waste from most to least preferable. They are source reduction, recycling, energy recovery, treatment, and disposal or other releases. **Right:** The Wasted Food Scale shows the most preferred to least preferred ways to handle wasted food in order to reduce environmental impacts. The best option is source reduction, which means to prevent wasted food. Next preferred is donating food to feed hungry people. Extra food can be donated to food banks, shelters, and soup kitchens. The third level of preference is to feed animals, where food scraps are used as animal feed. After that, composting food scraps will return nutrients to soil, or industrial anaerobic digestion turns the resulting digested materials and biosolids into into energy or fuel. The least preferred option is landfill or anaerobic digestion where disposal is the end result. Incineration of any kind should be avoided. Images by EPA (public domain).

Source reduction (waste minimization) refers to the strategies that minimize the amount of generated waste and/or reduce the toxicity of the resultant waste when designing or manufacturing products or services. Waste can be reduced by reusing materials, using less hazardous substitute materials, or by modifying components of design and processing. Source reduction in manufacturing not only saves resources, but it also reduces costs for the manufacturer and indirectly for the consumer. For example, minimal packaging reduces material use, increases distribution efficiency, and reduces fuel consumption and resulting air emissions. Similarly, building materials can be designed to reduce the overall mass of material needed for a given structure.

Recycling refers to recovery and reprocessing of useful materials. Numerous examples of successful recycling and reuse efforts are encountered every day. Many construction materials can be reused, including concrete, asphalt materials, masonry, and reinforcing steel. "Green" plant-based wastes are often recovered and immediately reused for mulch or fertilizer applications. Many industries also recover various byproducts for reuse. In some cases, the recycled materials are used as input materials and are heavily processed into end products. Common examples include the use of scrap paper for new paper manufacturing, or the processing of old aluminum cans into new aluminum products. In other cases, reclaimed materials undergo little or no processing prior to their reuse. Some common examples include the use of tree waste as wood chips, or the use of brick and other fixtures into new structural construction. In any case, the success of recycling depends on effective collection and processing of recyclables, markets for reuse, and public acceptance and promotion of recycled products and applications utilizing recycled materials.

Energy Recovery is a process that captures usable energy from food waste. This happens after higher options on the Wasted Food Scale (like feeding people or animals) have been used. Energy recovery can include processes such as using a **biodigester** to break down organic waste with bacteria. During this process, the bacteria produce **methane gas**, which can be collected and used as a source of energy, such as for electricity. This helps reduce the amount of waste sent to landfills while also creating a useful energy resource. It could also include burning the waste for energy recovery, although this is in the EPA's "avoid" category on the food waste reduction scale (figure $11.4.a$) (right).

Treatment refers to using physical, chemical or biological processes to change the volume, toxicity, or environmental impact of the waste, or make it safer to handle and dispose of. These processes include breaking down organic waste with microorganisms (e.g. biodigestion), using chemicals to neutralize harmful substances, or physically altering waste to make it smaller or more stable. Treatment helps prepare waste for safer final disposal when higher options are not possible.

Disposal is the final step in the waste hierarchy and includes **landfilling, burning (incineration), or release to the environment**. In a landfill, waste is buried and may break down slowly, sometimes producing harmful gases like methane. Incineration burns waste to reduce its volume, but it can create **air pollution** when solid waste is released into the atmosphere as gases and particles. These methods can have significant environmental impacts, so **disposal is the least preferred option** and should only be used when no better choices are available.

The Six Rs of Sustainability

The six **R's** (rethink, refuse, reduce, reuse, repair, and recycle) are strategies that allow individuals to limit the volume and negative impacts of the waste they generate. They resemble the waste management hierarchy, but focus at the individual scale rather than the scale of a company or waste management system. They are listed in order of most to least environmentally beneficial, and they are all better alternatives than discarding trash into the landfill. Recycling is last because it requires energy to reprocess waste.

To **rethink** can include asking if you really need something, or rethinking a system and changing how it works. For example, this could include thinking about your consumption patterns before you buy, or changing our energy system to focus on small-scale community power driven by solar and battery storage rather than a large centralized grid focused on non-renewable energy. **Refuse** means to decline products or processes that harm the environment if you do not need them. To **reduce** is to choose products or processes with lower **ecological footprint** (the area of land required to produce them). Choosing products without packaging or wrapped in paper is an example of both reducing your impact and refusing to support the oil industry's plastics. **Reuse** refers to using a product multiple times or finding an alternative use for it. For example, reuse a bag at the grocery store. Secondhand clothing and furniture has already been produced so buying used will dramatically decrease your carbon, water, and energy footprint. A plastic yogurt or pudding container could be repurposed for storage or gardening. **Repair** means to fix things that you have. Can you patch a hole in your jeans using visible or invisible mending (figure $11.4.b$)? Can you take something to a tailor to get it taken in so that it fits better rather than getting rid of it? As noted above, **recycling** means to return used items to be reprocessed (figure $11.4.b$). Commonly recycled materials include cardboard, glass, cans, and certain plastics. However, approximately 90% of plastics are not recycled, so if given the choice, try to reduce use of single use plastics and refuse them.



Figure 11.4.*b*: **Top:** Pair of jeans visibly mended by weaving and darning using mint green embroidery thread. By Lisenka92 - Own work, CC BY-SA 4.0, wikimedia. **Bottom:** Recycling, or returning materials to be reprocessed, is the last of the six R's. If items cannot be refused, reduced, reused, or repaired, recycling is a better alternative than discarding into the landfill. Image by Laura Jameson (Pexels) Pexels(opens in new window) [www.pexels.com]

Resin identification codes are the triangular symbols on recyclable plastics that indicate their composition (figure 11.4.*c*). A variety of materials make up plastics, and they are not all recycled the same way. Furthermore, your local recycling service may recycle some types of plastics but not others. Plastics with codes 1 and 2 are the only ones that are commonly recycled. For the codes, check with your local recycling service. Local businesses may house collection bins for plastics that are not commonly recycled. For example, grocery stores commonly have receptacles for recycling plastic bags, which fall under code 4.




Figure 11.4.*c*): The resin identification coding system explains the chemical composition of various plastics. Some are commonly recycled, and others are not. The text in the image is, "1: PET Polyethylene terephthalate, 2: HDPE High-density polyethylene, 3: PVC Polyvinyl chloride, 4: LDPE Low-density polyethylene, 5: PP Polypropylene, 6: PS Polystyrene, 7: OTHER Other plastics, including acrylic, acrylonitrile butadiene styrene, fiberglass, nylon, polycarbonate and polylactic acid. Image by Filtre/OpenClipart Library(opens in new window) (public domain).

Interactive Element

Which items can be recycled is based on where you live and the waste management service available there. The Environmental Protection Agency(opens in new window) provides general recycling guidelines. You local waste collector may provide an interactive website to help you determine which items to recycle. For example Recology, which serves Northern California, has a searchable website called What Bin(opens in new window).

Composting

Compost is organic material that can be added to soil to help plants grow (figure $11.4.d$). Food scraps and yard waste together currently make up more than 30% of what we throw away and could be composted instead. Making compost keeps these materials out of landfills where they take up space and release methane, a potent greenhouse gas that contributes to climate change. Furthermore, compost enriches soil, helping retain moisture and suppress plant diseases and pests. It reduces the need for chemical fertilizers and encourages the production of beneficial bacteria and fungi. Composting can be done at an industrial scale, such as with yard waste collected from homes in a city. It can also be done at home.



Figure $11.4.d$): Compost is a nutrient-rich substance formed from decomposition of yard and food waste that enriches garden soil. Image by EPA(opens in new window) (public domain).

All composting requires three basic ingredients: browns, greens, and water. "Browns" are materials such as dead leaves, branches, and twigs. "Greens" are materials such as grass clippings, vegetable waste, fruit scraps, and coffee grounds. The brown materials provide carbon for your compost, the green materials provide nitrogen, and the water provides moisture to help break down the organic matter. A compost pile should have an equal amount of browns to greens. Layers of these organic materials with different particle sizes should be alternated.

Backyard Composting

Backyard composting requires a compost bin or pile to be placed in dry, shady spot near a water source (figure $11.4.e$). Brown and green materials are added as they are collected, and larger pieces are chopped or shredded before they are added to the pile. Dry materials are moistened as they are added. Once the compost pile is established, grass clippings and green waste are mixed into the pile when added. When fruit and vegetable wastes are added, they are buried under 10 inches of compost material. Some homeowners cover the top of the compost with a tarp to keep it moist. When the material at the bottom is dark and rich in color, the compost is ready to use. This usually takes anywhere between two months to two years.



Figure 11.4.e: This compost bin contains food scraps and yard waste. Image by Ben_Kerckx/Pixabay(opens in new window) (Pixabay license(opens in new window))

Indoor Composting

For those who do not have the space for an outdoor compost pile, materials can be composted indoors using a special type of bin, which can be purchased at a local hardware store, gardening supplies store, or made at home. Care must be taken to properly manage the compost pile and keep track of what is added such that it does not attract pests or rodents or smell bad. The compost should be ready in two to five weeks.

Interactive Element

Not all food scraps can go into compost. If you want to start a compost pile and are not sure which food scraps are suitable, see the lists on the Environmental Protection Agency(opens in new window) website.

Biodigesters

Biodigesters are systems that use microorganisms (like bacteria) to break down organic waste, such as food scraps or sewage, in an environment without oxygen. As the waste is digested, the bacteria produce **biogas**, mainly methane, which can be collected and used to generate heat or electricity. Biodigesters also create a nutrient-rich material that can be used as fertilizer. For example, EDCO in La Mesa, CA uses anaerobic digestion to process organic waste and recover energy, while the Point Loma Wastewater Treatment Plant in San Diego uses similar processes to treat sewage and capture biogas for energy use to power the plant and sell back to SDG&E to power homes in Point Loma. These systems help reduce waste, lower greenhouse gas emissions, and turn organic material into useful resources.

Attribution

Modified by Melissa Ha from the following sources:

- Systems of Waste Management (opens in new window)from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Composting at Home(opens in new window). Environmental Protection Agency. Accessed 01-18-2021 (public domain).

This page titled 11.4: From Waste Management to a Circular Economy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 19.4: Waste Reduction by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

11.5: Circular Economy and Stewardship

Circular Economy and Stewardship

For much of modern history, societies have followed a **linear economy** in which materials are extracted, manufactured into products, used, and then discarded. This "take-make-waste" approach has contributed to resource depletion, pollution, greenhouse-gas emissions, and growing quantities of waste requiring disposal.

A **circular economy** seeks to redesign this system by keeping products, materials, and nutrients in use for as long as possible (figure $11.5.a$). Rather than viewing discarded items as worthless trash, a circular economy treats them as resources that can be reused, repaired, refurbished, remanufactured, recycled, composted, or recovered. The goal is to minimize waste generation while reducing demand for new raw materials and energy. **Dematerialization** is the process of providing the same function or value while using fewer physical materials. Examples include digital media, product lightweighting, and sharing systems that reduce the need for individual ownership, such as a car share program.




Figure $11.5.a$: In a circular economy, items that would be wastefully discarded can instead be sent back to an earlier point in the construction process. Image by Geissdoerfer, M., Pieroni, M.P., Pigosso, D.C. and Soufani, K., (CC BY 4.0) <https://creativecommons.org/licenses/by/4.0>, via Wikimedia Commons; [commons.wikimedia.org/wiki/File:The_Circular_Economy_concept.png]

The waste hierarchy and the Six R's introduced earlier in this chapter provide practical pathways toward a more circular economy. Strategies such as refusing unnecessary products, reducing consumption, repairing damaged items, reusing materials, recycling, composting, and recovering energy from organic wastes all help keep materials circulating through society rather than being sent to landfills or incinerators.

Circular systems can be found at many scales. Individuals may carry reusable water bottles, purchase secondhand clothing, or repair electronics. Communities may expand composting programs and repair cafés. Businesses may redesign products to last longer, use recycled materials, or offer take-back programs that recover materials at the end of a product's life. Together, these efforts reduce waste while conserving energy, materials, and ecological communities.

Ultimately, moving toward a circular economy requires a shift in how society views waste. In natural ecosystems, materials continuously cycle through biological and geological processes rather than being discarded. A circular economy seeks to mimic these cycles by designing products and systems that keep materials in use through durability, repair, reuse, recycling, composting, and resource recovery. In this vision, the concept of waste largely disappears as materials circulate through society in much the same way that nutrients cycle through ecosystems. Although achieving complete circularity may not be possible, moving toward more circular systems can reduce pollution, conserve materials and energy, decrease dependence on continual resource extraction, and help create a more sustainable and resilient future.

Example 11.5.1

Local Connection: Zero Waste in San Diego

Many communities are beginning to adopt circular-economy principles. In California, state laws support recycling, composting, and the diversion of organic waste from landfills. Initiatives such as residential composting programs, food-waste recovery efforts, and organic-waste processing facilities help keep materials circulating while reducing methane emissions from landfills.

Programs such as EDCO's anaerobic digestion facility, organic-waste composting operations, and regional recycling systems demonstrate how communities can recover resources from materials that were once considered waste.

Think About It

Throughout this chapter, you explored waste generation, disposal, recycling, composting, biodigesters, and the environmental impacts of waste.

Reflect: Which requires a bigger change in society: becoming better at managing waste after it is created, or redesigning products and systems so less waste is created in the first place? What examples from your own life illustrate the difference between a linear economy and a circular economy?

This page titled 11.5: Circular Economy and Stewardship is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.

12: ENVIRONMENTAL HEALTH AND TOXICOLOGY

Introduction

In 1854, a cholera outbreak in England began killing people within a day of symptoms appearing. Most people did not understand how the disease spread, but one physician, Dr. John Snow, suspected it was linked to contaminated water. Over about a month, he gathered evidence showing that nearly all the deaths were connected to a single well. Later, scientists discovered that the water had been contaminated by a bacterium called *Vibrio cholerae*, named after the disease it causes. Cholera still exists today. The World Health Organization estimates that there are between 1.3 and 4 million cases each year, with 21,000 to 143,000 deaths. In some areas, regular testing and prevention help stop outbreaks before people get sick. In other places, however, action is often taken only after an epidemic has already begun.




Figure 12.a: Original map made by John Snow in 1854. Cholera cases are highlighted in black, showing the clusters of cholera cases (indicated by stacked rectangles) in the London epidemic of 1854. The map was created in order to better understand the pattern of cholera spread in the 1854 Broad Street cholera outbreak, which Snow would use as an example of how cholera spread via the fecal-oral route through water systems as opposed to the miasma (poisonous vapor in air) theory of disease spread. The contaminated pump is located at the intersection of Broad Street and Cambridge Street (now Lexington Street), running into Little Windmill Street. The map marks an important part of the development of epidemiology as a field, and of disease mapping as a whole. Map of the book "On the Mode of Communication of Cholera" by John Snow, originally published in 1854 by C.F. Cheffins, Lith, Southhampton Buildings, London, England. Image by Wikimedia Commons (Public Domain).

12.1: Types of Environmental Hazards

12.2: Epidemiology

12.3: Infectious Diseases

12.4: Environmental Toxicology

12.5: Case Study - The Love Canal Disaster

12.6: Environmental Hazard Reduction

Attribution

Modified by Erin Rempala.

Modified by Rachel Schleiger (CC-BY-NC(opens in new window)).

This page titled 12: Environmental Health and Toxicology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15: Environmental Health by Melissa Ha and Rachel Schleiger is licensed CC BY 4.0.

12.1: Types of Environmental Hazards

Environmental health is the study of how both natural environments and human-made surroundings, along with human behaviors, affect people's health. This field aims to prevent disease, death, and disability by reducing contact with environmental hazards and encouraging healthier behaviors. **Environmental hazards** are factors in the environment that can harm human health and well-being (table $12.1.a$).

Table $12.1.a$: Typical Environmental Health Issues: Determinants and Health Consequences.

Causes (Determinants)	Possible Adverse Health and Safety Consequences
Inadequate water (quantity and quality), sanitation and solid waste disposal, improper hygiene (handwashing)	Diarrhea and vector-related diseases (for example, malaria, schistosomiasis, and dengue)
Improper water resource management, including poor drainage	Vector-related diseases
Crowded housing and poor ventilation of smoke	Acute and chronic respiratory disease, including lung cancer from coal and tobacco inhalation
Exposures to vehicular and industrial air pollution	Respiratory diseases, some cancers, and loss of IQ in children
Population movement and encroachment and construction, which affect feeding and breeding grounds of vectors, such as mosquitoes	Vector-related diseases May also spread other infectious diseases (for example, HIV/AIDS, Ebola)
Exposure to naturally occurring toxic substances	Poising from substances such as arsenic, manganese, and fluorides
Natural resources degradation (for example, landslides, poor drainage, erosion)	Injury and death from landslides and flooding
Climate change, partly from combustion of fossil fuel and release of greenhouse gases in transportation, industry, and poor energy conservation in housing, fuel, commerce, and industry	Injury/death from extreme heat/cold, floods, and fires Indirect effects spread of vector-borne diseases, aggravation of respiratory diseases, population dislocation, water pollution from sea level rise, etc.
Ozone depletion from industrial and commercial activity	Skin cancer, cataracts Indirect effect: compromised food production

Table based on Lvovsky/World Bank (CC-BY)

Traditional versus Modern Environmental Hazards

Environmental hazards are often described as traditional or modern. Traditional hazards are linked to poverty and include risks such as unsafe water, poor sanitation, and indoor air pollution. Modern hazards arise from technological and industrial development, such as chemical exposure, urban air pollution, and climate change. While traditional hazards remain more common in low-income settings, both types now occur worldwide. Many regions, including many low- and Middle-Income Countries (LMICs), face a mix of these risks, sometimes called a "double burden".

As of 2025, about 2.1 billion people worldwide—roughly one in four—still depend on polluting fuels such as kerosene, coal, and

biomass (including wood, animal dung, and crop waste) for cooking and heating. These fuels are often burned in open fires or inefficient stoves with poor ventilation, producing harmful household air pollution (figure 12.1.a). In 2021, household air pollution was responsible for an estimated 2.9 million deaths each year, including more than 309,000 deaths among children under five. Women and children—who are typically responsible for cooking and collecting fuel—experience the greatest health impacts from these exposures (figure 12.1.b).

Expanding access to cleaner fuels and technologies is essential to reduce these risks. Cleaner options include electricity, solar energy, biogas, liquefied petroleum gas (LPG), natural gas, alcohol fuels, and improved biomass stoves that meet World Health Organization emission standards. However, at current rates, only about 78% of the global population is expected to have access to clean cooking by 2030, leaving nearly 1.8 billion people without safe cooking solutions.



Figure 12.1.a: A traditional wood-fired 3-stone stove in Guatemala, which causes indoor air pollution. Image from gringologue - Flickr.com [www.flickr.com] [www.flickr.com] [www.flickr.com] (CC-BY 2.0).

Interactive Element

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure 12.1.b: Death rates from indoor air pollution. The estimated number of age-standardized deaths from all causes attributed to household air pollution from solid fuels, per 100,000 people.Source IHME, Global Burden of Disease (2025) – with major processing by Our World in Data. Published online at OurWorldInData.org.

The contribution of modern environmental risks to the disease burden in most low-income countries is similar to – and in quite a few countries, greater than – that in rich countries (figure 12.1.a). Urban air pollution, for example, is highest in parts of China, India, and some cities in Asia and Latin America. Low-income people increasingly experience a "**double burden**" of traditional and modern environmental health risks. In rich countries, they experience twice the burden of illness and death from all causes and 10 times greater disease burden from environmental risks.

Biological, Chemical, and Physical Environmental Hazards

Environmental hazards can also be classified into three interrelated categories (biological, chemical, and physical) based on

the properties of their causes. These categories are not mutually exclusive with traditional versus modern hazards. For example, indoor air pollution is both a traditional and chemical hazard. Different hazards can interact and exacerbate one another. For example, a flood is primarily a physical hazard, but it can lead to the spread of waterborne disease (a biological hazard). Similarly, air pollution (a chemical hazard) can damage respiratory tissue, making the body more vulnerable to a respiratory infection (a biological hazard). Infectious diseases (biological hazards) can also weaken the immune system, making an individual more vulnerable to chemical hazards.

Biological Hazards

For most of human history, biological hazards were the most significant factor in health. **Biological hazards** are infectious (communicable) diseases caused by **pathogens** (disease-causing organisms or infectious particles) such as bacteria, fungi, parasitic worms, protozoa, viruses, and prions. **Bacteria** are single-celled organisms with small, simple cells. Examples of bacterial diseases include tuberculosis, cholera, bacterial pneumonia, and dysentery. **Fungi** may have one or multiple cells and have a more complex cell type than bacteria. Fungal diseases include minor infections like candidiasis (yeast infection) or athlete's foot, but they can also causes severe respiratory infections (histoplasmosis, coccidioidomycosis, etc.) particularly in individuals with compromised immune systems. **Parasitic worms** are animals from several phyla (groups) that siphon nutrients from their hosts (figure $12.1.c$). Examples include tapeworms, commonly acquired through consuming undercooked meat, and blood flukes (*Schistosoma*). Like fungi, **protozoa** have larger, more complex cells than bacteria, but they are single celled and lack the rigid cell wall that surrounds fungal cells. Malaria (figure $12.1.c$), African trypanosomiasis (sleeping sickness), and giardiasis are caused by protozoa. **Viruses** are infectious particles with genetic information surrounded by a protein coat, but they are not technically considered organisms in part because they do not consist fo cells. COVID-19, influenza, measles, the common cold, ebola viral disease (Ebola hemorrhagic fever), and human immunodeficiency virus (HIV)/acquired immune deficiency syndrome (AIDS) are all caused by viruses. **Prions** (proteinaceous infectious particles) are even simpler than viruses because they lack genetic material and only contain protein.




Figure 12.1.c: Top: Dracunculus medinensis (Guinea worm), dracunculiasis. Dracunculus medinensis, or Guinea worm, is a nematode that infects humans and domestic animals through contaminated water.The stick has been removed and larvae have been squeezed from the flattened exterior portion of the worm. Image by Otis Historical Archives of "National Museum of Health & Medicine" (OTIS Archive 1), (CC BY 2.0). Bottom: Red blood cells infected with the protozoan Plasmodium, which causes malaria, under the microscope. Image by Kim-Sung Lee, Janet Cox-Singh, Balbir Singh (CC-BY).

While the proportion of deaths caused by in infectious diseases has overall decreased (with a higher proportion of deaths caused by noncommunicable diseases such as cancer and cardiovascular disease), infectious diseases still caused about one in five deaths in 2017. These deaths occurred at the highest rates in low-income countries and many were in children. Malnutrition, unclean water, poor sanitary conditions and lack of proper medical care all play roles in transmission and high death rates from infectious diseases Compounding the problems of infectious diseases are factors such as antibiotic-resistant pathogens, pesticide-resistant disease vectors, and overpopulation.

Chemical Hazards

Chemical hazards are toxic substances, which cause damage to living organisms. Air pollutants (such as secondhand smoke or carbon monoxide), heavy metals, and pesticides are a few examples. We can be exposed to these contaminants from a variety of residential, commercial, and industrial sources. Sometimes harmful environmental contaminants occur biologically, such as those from mold or a toxic algae bloom. Toxins can be classified based on their origin, purpose, chemical structure and properties, or

effects. Table 12.1.*b* describes a few categories of toxins based on their effects and provides examples. A few of these examples are discussed more specifically below.

Table 12.1.*b*: *Classification of Environmental Contaminants*

Contaminant	Definition
Carcinogen	An agent which may produce cancer (uncontrolled cell growth), either by itself or in conjunction with another substance. Examples include formaldehyde, asbestos, radon, vinyl chloride, and tobacco.
Teratogen	A substance which can cause physical defects in a developing embryo. Examples include alcohol and cigarette smoke.
Mutagen	A material that induces genetic changes (mutations) in the DNA. Examples include radioactive substances (such as radon and nuclear fuel and waste) and nitrous acid. Some forms of radiation (see Physical Hazards) are also mutagens.
Neurotoxin	A substance that can cause an adverse effect on the chemistry, structure or function of the nervous system. Examples include lead and mercury.
Endocrine disruptor	A chemical that may interfere with the body's endocrine (hormonal) system and produce adverse developmental, reproductive, neurological, and immune effects in both humans and wildlife. A wide range of substances, both natural and man-made, are thought to cause endocrine disruption, including pharmaceuticals, dioxin and dioxin-like compounds, arsenic, polychlorinated biphenyls (PCBs), DDT and other pesticides, per- and polyfluoroalkyl substances (PFAS), pthalates, and plasticizers such as bisphenol A (BPA).

Formaldehyde

Formaldehyde is a colorless, flammable gas or liquid that has a pungent, suffocating odor. It is a **volatile organic compound**, which is a compound containing carbon and hydrogen that easily becomes a vapor or gas. It is also naturally produced in small, harmless amounts in the human body. The primary way we can be exposed to formaldehyde is by breathing air containing it. Formaldehyde is released into the air by industries using or manufacturing formaldehyde, wood products (such as particle-board, plywood, and furniture), automobile exhaust, cigarette smoke, paints and varnishes, and carpets and permanent press fabrics. Nail polish and commercially applied floor finish emit formaldehyde (figure 12.1.*d*).



Figure 12.1.*d*: Nail products are known to contain toxic chemicals, such as dibutyl phthalate (DBP), toluene, and formaldehyde.

In general, indoor environments consistently have higher concentrations than outdoor environments because many building materials, consumer products, and fabrics emit formaldehyde. Levels of formaldehyde measured in indoor air range from 0.02–4 parts per million (ppm). Formaldehyde levels in outdoor air range from 0.001 to 0.02 ppm in urban areas.

Environmental Working Group

For more information on safety in cosmetics and household cleaning products, tap water, sunscreen, etc. check out the Environmental Working Group (EWG)'s Skin Deep database, Guide to Healthy Cleaning, and others.

Heavy Metals

Heavy metals are chemical elements of high density that form a special type of bond (called metallic bonds, in which electrons are shared but in a less constricted way than in covalent bonds). Arsenic, mercury, lead, and cadmium are examples of heavy metals.

Arsenic (As) is a naturally occurring element that is normally present throughout our environment in water, soil, dust, air, and food. Levels of arsenic can regionally vary due to farming and industrial activity as well as natural geological processes. The arsenic from farming and smelting tends to bind strongly to soil and is expected to remain near the surface of the land for hundreds of years as a long-term source of exposure. Wood that has been treated with chromated copper arsenate (CCA) is commonly found in decks and railings in existing homes and outdoor structures such as playground equipment. Some underground aquifers are located in rock or soil that has naturally high arsenic content.

Most arsenic gets into the body through ingestion of food or water. Arsenic in drinking water is a problem in many countries around the world, including Bangladesh, Chile, China, Vietnam, Taiwan, India, and the United States. Arsenic may also be found in foods, including rice and some fish, where it is present due to uptake from soil and water. It can also enter the body by breathing dust containing arsenic.

Arsenic poisoning causes a variety of symptoms and serious health conditions (figure $12.1.e$). Researchers are finding that arsenic, even at low levels, can interfere with the body's endocrine system. Arsenic is also a known human carcinogen associated with skin, lung, bladder, kidney, and liver cancer.



Figure $12.1.e$: Patchy areas of dark skin pigmentation (arsenical hyperkaratosis) on the palms of the hands is a symptom of arsenic poisoning. Image and caption (modified) from Agency for Toxic Substances and Disease Registry/CDC (public domain).

Mercury (Hg) is a naturally occurring metal, a useful chemical in some products, and a potential health risk. Mercury exists in several forms; the types people are usually exposed to are methylmercury and elemental mercury. Elemental mercury at room temperature is a shiny, silver-white liquid which can produce a harmful odorless vapor. Methylmercury, an organic compound, can build up in the bodies of long-living, predatory fish (Biomagnification). Although fish and shellfish have many nutritional benefits, consuming large quantities of fish increases a person's exposure to mercury. Pregnant women who eat fish high in mercury on a regular basis run the risk of permanently damaging their developing fetuses. Children born to these mothers may exhibit motor difficulties, sensory problems and cognitive deficits. The United States Environmental Protection Agency thus recommends that pregnant women and young children should not consume any swordfish, shark, king mackerel, or tilefish because of their high

mercury content. These individuals are advised to eat fish low in mercury such as salmon, shrimp, pollock, and catfish (figure 12.1.f). To keep mercury out of the fish we eat and the air we breathe, it's important to take mercury-containing products to a hazardous waste facility for disposal. Common products sold today that contain small amounts of mercury include fluorescent lights and button-cell batteries (figure 12.1.g).



Figure 12.1.f: Fish classified based on mercury levels. The best choices (black sea bass, catfish, herring, trout, and many others) have the lowest mercury levels, and two or three servings of these choices can be safely consumed each week. The good choices (carp, halibut, yellowfin tuna, etc.) have moderate levels, and it is safe to eat one serving per week. The choices to avoid, such as shark and swordfish, have the highest mercury levels and should be avoided. A table listing all fish in each category is available here. Image by EPA and FDA (public domain).Seafood WatchFor regular updates on seafood recommendations based on the science (population sizes, fishing practices, mercury levels, etc) go to Monterey Bay Aquarium's Seafood watch.



Figure 12.1.g: Button-cell batteries found in small devices like watches and hearing aids contain mercury and must be discarded at the proper hazardous waste facility. Image by Lead holder (CC-BY-SA).

Lead (Pb) is a metal that occurs naturally in the rocks and soil of the Earth's crust. It is also released from mining, manufacturing, and the combustion (burning) fossil fuels such as coal, oil, gasoline, and natural gas. Lead has no distinctive taste or smell. Lead is used to produce batteries, pipes, roofing, scientific electronic equipment, military tracking systems, medical devices, and products to shield X-rays and nuclear radiation. It is used in ceramic glazes and crystal glassware. Because of health concerns,

lead and lead compounds were banned from house paint in 1978; from solder used on water pipes in 1986; from gasoline in 1995; from solder used on food cans in 1996; and from tin-coated foil on wine bottles in 1996. The U.S. Food and Drug Administration has set a limit on the amount of lead that can be used in ceramics.

Lead and lead compounds are listed as “reasonably anticipated to be a human carcinogen”. It can affect almost every organ and system in your body. It can be equally harmful if breathed or swallowed. The part of the body most sensitive to lead exposure is the central nervous system, especially in children, who are more vulnerable to lead poisoning than adults. A child who swallows large amounts of lead can develop brain damage that can cause convulsions and death; the child can also develop blood anemia, kidney damage, colic, and muscle weakness. Repeated low levels of exposure to lead can alter a child’s normal mental and physical growth and result in learning or behavioral problems. Exposure to high levels of lead for pregnant women can cause miscarriage, premature births, and smaller babies. Repeated or chronic exposure can cause lead to accumulate in your body, leading to lead poisoning.

The video below explains how Flint, Michigan's water supply was polluted with lead in 2014 and the problematic government response that followed. This video was made in 2016.

Flint now has access to a safe water supply, and the city is working to compensate residents for the harm caused. However, no amount of money can truly make up for the loss or suffering experienced by families whose loved ones were injured or died. (Click here to read a 2020 update from Politico.)

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Asbestos

Asbestos is a mineral fiber that occurs in rock and soil. Because of its fiber strength and heat resistance asbestos has been used in a variety of building construction materials for insulation and as a fire retardant. Asbestos has also been used in a wide range of manufactured goods, mostly in building materials (roofing shingles, ceiling and floor tiles, paper products, and asbestos cement products), friction products (automobile clutch, brake, and transmission parts), heat-resistant fabrics, packaging, gaskets, and coatings. Exposure to asbestos is associated with cancers (lung cancer and mesothelioma) and another lung disease called asbestosis. In the United States, certain uses of asbestos, including in corrugated paper, flooring, and building insulation, are banned under the Toxic Substances Control Act and Clean Air Act (figure $12.1.h$).

Asbestos has been banned in more than 60–70 countries worldwide as of 2025. In the United States, the EPA attempted to ban most asbestos products in 1989, but the rule was largely overturned by a court decision in 1991. While the U.S. has since implemented additional restrictions—including a 2024 rule banning certain ongoing uses—it still does not have a complete ban on asbestos.



Figure 12.1.*h*: An asbestos pipe wrap. While this use of asbestos is banned in the United States, other uses are still permitted. Image by EPA (public domain).

Per- and polyfluoroalkyl substances (PFAS)

Per- and polyfluoroalkyl substances (PFAS) are a group of manufactured organic chemicals used in a variety of industries (figure 12.1.*i*). They can be found in food packaging, stain- and water-repellent fabrics, nonstick products (such as Teflon), polishes, waxes, paints, cleaning products, and fire-fighting foams.




Figure 12.1.*i*: The chemical structure of perfluorooctanoic acid (PFOA), a PFAS. This molecule has eight carbon atoms (black) connected in a chain. Most of them are bound to fluorine atoms (green). The carbon at the end of the chain forms two covalent bonds with an oxygen and is single bounded to another oxygen atom, which is also attached to a hydrogen. Image by National Institute of Environmental Health Sciences (public domain).

Studies indicate that some PFAS can cause reproductive and developmental, liver and kidney, and immunological effects in laboratory animals. More limited findings associate some PFAS with low infant birth weights, effects on the immune system, cancer, and thyroid hormone disruption in humans.

Eight major chemical manufacturers in the U.S. phased out the use certain PFAS (called perfluorooctanoic acid, PFOA, and perfluorooctane sulfonate, PFOS) and related chemicals in their products and as emissions from their facilities. However, these PFAS can still be imported and other PFAS are still manufactured in the U.S.

Polychlorinated Biphenyls (PCBs)

Polychlorinated biphenyls (PCBs) are a group of manufactured organic chemicals. They belong to a broad family of chemicals known as chlorinated hydrocarbons, which consisting of carbon, hydrogen and chlorine atoms (figure $12.1.j$). The number of chlorine atoms and their location in a PCB molecule determine many of its physical and chemical properties. PCBs have no known taste or smell, and range in consistency from an oil to a waxy solid.




Figure $12.1.j$: A variety of polychlorinated biphenyls (PCBs). They each have two rings of six carbon atoms each attached together. Additionally, each has multiple chlorine atoms attached to the rings, but the exact number and placement of chlorine atoms varies. Image by Leyo/M. Van den Berg et al. (public domain).

Polychlorinated biphenyls have been shown to cause cancer, cause birth defects, and affect the immune, reproductive, nervous, and endocrine systems in animals. Studies in humans support evidence for potential carcinogenic and non-carcinogenic effects of PCBs. The different health effects of PCBs may be interrelated. Alterations in one system may have significant implications for the other systems of the body.

The manufacture of PCBs in the U.S. began in 1929 until it was banned in 1979 under the Toxic Substances Control Act. Due to their non-flammability, chemical stability, high boiling point and electrical insulating properties, PCBs were used in hundreds of industrial and commercial applications including in electrical equipment, coolants paints, plastics, rubber products, pigments, and dyes.

Bisphenol A (BPA)

Bisphenol A (BPA) is a chemical synthesized in large quantities for use primarily in the production of polycarbonate plastics and epoxy resins. Polycarbonate plastics have many applications including use in some food and drink packaging such as water and infant bottles, impact-resistant safety equipment, and medical devices (figure $12.1.k$). Epoxy resins are used as lacquers to coat metal products such as food cans, bottle tops, and water supply pipes. Some dental sealants and composites may also contribute to BPA exposure. The primary source of exposure to BPA for most people is through the diet. Bisphenol A can leach into food from the protective internal epoxy resin coatings of canned foods and from consumer products such as polycarbonate tableware, food storage containers, water bottles, and baby bottles. The degree to which BPA leaches from polycarbonate bottles into liquid may depend more on the temperature of the liquid or bottle, than the age of the container. It has also be found in breast milk.



Figure 12.1.*k*: A water bottle is labeled as BPA free. Image by Hteink.min (CC-BY-SA).

Some animal studies suggest that infants and children may be the most vulnerable to the effects of BPA. It disrupts signaling by estrogen, a naturally produced hormone, and the U.S. National Toxicology Program (NTP) documented concerns about its effects on the behavior, brain, and prostate in young children and developing fetuses.

The following personal choices can reduce exposure to BPA:

- Avoid microwaving polycarbonate plastic food containers. Polycarbonate is strong and durable, but over time it may break down from over use at high temperatures.
- Note recycle codes on the bottom of plastic containers. Some, but not all, plastics that are marked with recycle codes 3 or 7 may be made with BPA.
- Reduce your use of canned foods.
- When possible, opt for glass, porcelain or stainless steel containers, particularly for hot food or liquids.

While BPA is not banned in the U.S., the Food and Drug Adminsitration banned its use in baby bottles and sippy cups in 2012 and its use in the coating of infant formula containers in 2013. However, similar compounds such as bisphenol S (BPS) are now used as replacements.

Phthalates

Phthalates are a group of synthetic chemicals used to soften and increase the flexibility of plastic and vinyl. Polyvinyl chloride (PVC) is made softer and more flexible by the addition of phthalates. Phthalates are used in hundreds of consumer products. Phthalates are used in cosmetics and personal care products, including perfume, hair spray, soap, shampoo, nail polish, and skin moisturizers (figure 12.1.*l*). They are used in consumer products such as flexible plastic and vinyl toys, shower curtains, wallpaper, vinyl miniblinds, food packaging, and plastic wrap. Exposure to low levels of phthalates may come from eating food packaged in plastic that contains phthalates or breathing dust in rooms with vinyl miniblinds, wallpaper, or recently installed flooring that contain phthalates. We can be exposed to phthalates by drinking water that contains phthalates.



Figure $12.1.l$: Phthalates are often used in shampoos and other personal care products, but some brands produce phthalate-free products. Image by inf per Open Food Facts (CC-BY-SA).

Phthalates are suspected to be endocrine disrupters. Some types of phthalates have affected the reproductive system of laboratory animals. In 2017, The U.S. Consumer Product Safety Commission (CPSC) has banned several phthalates from being used at concentrations greater than 0.1% in toys and products designed to be used by children three years old or younger.

Radon

Radon is a radioactive gas that is naturally-occurring, colorless, and odorless (figure $12.1.m$). It comes from the natural decay of uranium or thorium found in nearly all soils. It typically moves up through the ground and into the home through cracks in floors, walls and foundations. It can also be released from building materials or from well water. Radon breaks down quickly, giving off radioactive particles. Long-term exposure to these particles can lead to lung cancer. Radon is the leading cause of lung cancer among nonsmokers, according to the U.S. Environmental Protection Agency, and the second leading cause behind smoking. To reduce the risk of radon exposure, the Department of Urban and Housing Development recommends testing your home for radon, avoiding smoking to reduce the risk of lung cancer, and ensuring proper ventilation in your home.




Figure $12.1.m$: This information graphic from the Centers for Disease Control explains how people can be exposed to radon from underground and its health risks. It reads "Protect your family from radon. Radon is a gas that you can't see, smell, or taste— but it can be dangerous. It's the second leading cause of lung cancer in the U.S. Radon is in the ground naturally. But sometimes it gets into homes through cracks in the floors or walls." Image by CDC (public domain).

Dichlorodiphenyltrichloroethane (DDT)

Dichlorodiphenyltrichloroethane (DDT) was the first of a long line of chlorinated hydrocarbon insecticides (figure $12.1.n$). These compounds are chains of carbon and hydrogen with chlorine atoms replacing some of the hydrogen atoms. Introduced during World War II, DDT, along with penicillin and the sulfa drugs, was responsible for the fact that this was the first war in history where trauma killed more people - combatants and noncombatants alike - than infectious disease.

DDT is effective against many crop pests as well as vectors of human diseases such as the mosquitoes that spread malaria and yellow fever and fleas, which transmit the plague. Prior to the introduction of DDT, the number of cases of malaria in Ceylon (now Sri Lanka) was more than a million a year. By 1963 the disease had been practically eliminated from the island. However, growing concern about the hazards of DDT led to its abandonment there in the mid-1960s, and soon thereafter malaria became common once again. Because it remains in the environment and is resistant to breakdown, DDT was especially effective against malarial mosquitoes. One or two sprays a year on the walls of homes kept them free of mosquitoes. It was also inexpensive, further adding to its appeal, but DDT has several serious drawbacks.






Figure $12.1.n$: Top: DDT was a commonly-used pesticide. The bag of DDT on the right says, "Insect spray containing 50% DDT. Destroys many common insects. Use on potatoes, peas, corn, fruits, and ornamentals." Top image by Leyo (public domain), and middle image by Xanthis (public domain).

DDT accumulates in fatty tissues (bioaccumulation) and becomes more concentrated at higher levels of the food chain (biomagnification), making it especially harmful to top predators, such as bald eagles. It was discovered that DDT caused the eggshells of birds to become fragile and break, making reproduction impossible.These effects were documented in the 1960s bestseller *Silent Spring* by marine biologist Rachel Carson (figure $12.1.p0$. Through this work, she brought widespread attention to the dangers of DDT and other synthetic pesticides. Her efforts helped strengthen the U.S. environmental movement and contributed to the growing public concern that ultimately led to the creation of the U.S. Environmental Protection Agency (EPA) in 1970.

As a result, the bald eagle was listed as an endangered species under U.S. law. After DDT was banned in the United States in 1972, affected bird populations made noticeable recoveries, including the iconic bald eagle.





Figure 12.1.*o*: DDT pesticide residue in fish and other prey poisoned Bald Eagles, causing egg shell thinning that resulted in widespread nesting failures. Image by Ron Holmes/USFWS (public domain).Figure 12.1.*p*: Rachel Carson in 1943. By U.S. Fish and Wildlife Service - This image originates from the National Digital Library of the United States Fish and Wildlife Service at this page http://training.fws.gov/history/carson/carson.html, Public Domain, https://commons.wikimedia.org/w/inde...p?curid=277288.

Physical Hazards

Physical hazards are additional forces that can imperil humans. Physical hazards may arise naturally such as natural disasters (earthquakes, wildfires, landslides, etc.) or extreme weather (figure 12.1.*q*). Others may arise from human structures or activities (traffic accident, building collapse, injury from mechanical equipment, strain on the body from repeated movements, etc.) Some physical hazards, such as explosions or radiation, can arise from natural or human sources.



Figure 12.1.*q*: This section of E. Grace Street, Richmond, VA, collapsed during tropical storm Gaston. Gaston dropped twelve inches of rain in the area. Image by Liz Roll/FEMA Photo Library (public domain).

Radiation is energy given off by matter in the form of rays or high-speed particles, and some types of radiation present a physical hazard. Some familiar forms of radiation are infrared radiation (heat), visible light, ultraviolet (UV) light, radio waves, and microwaves. We are exposed to radiation every day from natural sources. For example, the sun exposes us to UV radiation. We are also exposed to radiation from human-made sources like medical X-rays and smoke detectors. We're even exposed to low levels of radiation on cross-country flights, from watching television, and even from some construction materials. Some types of radioactive materials are more dangerous than others. Specifically, **ionizing radiation**, like X rays and gamma rays (one of the forms of radiation emitted from nuclear fuel and waste), have enough energy to break molecular bonds and displace (or remove) electrons from atoms.

Attribution

Modified by Erin Rempala.

Modified by Melissa Ha from the following sources:

- *Environmental Biology* by Mathew R. Fisher (licensed under CC-BY(opens in new window))
  - The Impacts of Environmental Conditions
  - Environmental Health
  - Environmental Toxicology
  - Pests and Pesticides
  - Energy Flow through Ecosystems
- Insecticides from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Impact on Human Health from *AP Environmental Science* by University of California College Prep (licensed under CC-BY)
- Environmental Health at a glance. 2003. World Bank. Washington, DC. (licensed under CC-BY)
- Asbestos. 2018. Environment Protection Agency. Accessed 01-07-2021. (public domain)
- Per- and Polyfluoroalkyl Substances (PFAS). 2018. Environment Protection Agency. Accessed 01-07-2021. (public domain)
- Polychlorinated Biphenyls (PCBs). 2020. Environment Protection Agency. Accessed 01-08-2021. (public domain)
- What are DALYs? National Institute of Mental Health. Accessed 01-06-2021. (public domain)
- Radiation Basics. 2020. U.S. NRC. Accessed 01-07-2021 (public domain)
- Phthalates. 2017. National Biomonitoring Program. Centers for Disease Control. Accessed 01-08-2021. (public domain)

This page titled 12.1: Types of Environmental Hazards is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15.1: Types of Environmental Hazards by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

12.2: Epidemiology

The field of **epidemiology** studies the geographical distribution and timing of infectious disease occurrences and how they are transmitted and maintained in nature, with the goal of recognizing and controlling outbreaks. The science of epidemiology includes **etiology** (the study of the causes of disease) and investigation of disease transmission (mechanisms by which a disease is spread). **Epidemiologists** are thus scientists who study the causes and patterns of human disease, which involves examining statistics to identify health threats and recommending strategies to reduce these threats.

The work of epidemiologist Alice Wang is highlighted in the video below. While many of the examples on this page focus on infectious disease (which result from biological hazards), epidemiologists can study noncommunicable diseases too (such as poisoning or obesity). The text at the end of the video says, "Are You CDC? For more information bout employment opportunities with CDC, please visit: jobs.cdc.gov".

YouTube Video


This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

History of Epidemiology

The studies of 19th century physicians and researchers such as John Snow, Florence Nightingale, Ignaz Semmelweis, Joseph Lister, Robert Koch, Louis Pasteur, and others sowed the seeds of modern epidemiology.

John Snow (figure $12.2.a$) was a British physician known as the father of epidemiology for determining the source of the 1854 Broad Street cholera epidemic in London. Based on observations he had made during an earlier cholera outbreak (1848–1849), Snow proposed that cholera was spread through a fecal-oral route of transmission and that a microbe was the infectious agent. He investigated the 1854 cholera epidemic in two ways. First, suspecting that contaminated water was the source of the epidemic, Snow identified the source of water for those infected. He found a high frequency of cholera cases among individuals who obtained their water from the River Thames downstream from London. This water contained the refuse and sewage from London and settlements upstream. He also noted that brewery workers did not contract cholera and on investigation found the owners provided the workers with beer to drink and stated that they likely did not drink water. Second, he also painstakingly mapped the incidence of cholera and found a high frequency among those individuals using a particular water pump located on Broad Street. In response to Snow's advice, local officials removed the pump's handle, resulting in the containment of the Broad Street cholera epidemic. John Snow's own account of his work has additional links and information.



(a) (b)


Figure 12.2.*a*: (a) John Snow (1813–1858), British physician and father of epidemiology. (b) Snow's detailed mapping of cholera incidence led to the discovery of the contaminated water pump on Broad street (red square) responsible for the 1854 cholera epidemic. (credit a: modification of work by "Rsabbatini"/Wikimedia Commons)

Snow's work represents an early epidemiological study and it resulted in the first known public health response to an epidemic. Snow's meticulous case-tracking methods are now common practice in studying disease outbreaks and in associating new diseases with their causes. His work further shed light on unsanitary sewage practices and the effects of waste dumping in the Thames. Additionally, his work supported the germ theory of disease, which argued disease could be transmitted through contaminated items, including water contaminated with fecal matter.

Florence Nightingale's work is another example of an early epidemiological study. In 1854, Nightingale was part of a contingent of nurses dispatched by the British military to care for wounded soldiers during the Crimean War. Nightingale kept meticulous records regarding the causes of illness and death during the war. Her recordkeeping was a fundamental task of what would later become the science of epidemiology. Her analysis of the data she collected was published in 1858. In this book, she presented monthly frequency data on causes of death in a wedge chart histogram (figure 12.2.*b*). This graphical presentation of data, unusual at the time, powerfully illustrated that the vast majority of casualties during the war occurred not due to wounds sustained in action but to what Nightingale deemed preventable infectious diseases. Often these diseases occurred because of poor sanitation and lack of access to hospital facilities. Nightingale's findings led to many reforms in the British military's system of medical care. Learn more about Nightingale's wedge chart here.



(a)



(b)

Figure 12.2.*b*: (a) Florence Nightingale reported on the data she collected as a nurse in the Crimean War. (b) Nightingale's diagram shows the number of fatalities in soldiers by month of the conflict from various causes. The total number dead in a particular month is equal to the area of the wedge for that month. The colored sections of the wedge represent different causes of death: wounds (pink), preventable infectious diseases (gray), and all other causes (brown).

Joseph Lister provided early epidemiological evidence leading to good public health practices in clinics and hospitals. Most physicians did not wash their hands between patient visits or clean and sterilize their surgical tools. Lister, however, discovered the disinfecting properties of carbolic acid, also known as phenol. He introduced several disinfection protocols that dramatically lowered post-surgical infection rates. He demanded that surgeons who worked for him use a 5% carbolic acid solution to clean their surgical tools between patients, and even went so far as to spray the solution onto bandages and over the surgical site during operations (figure 12.2.*c*). He also took precautions not to introduce sources of infection from his skin or clothing by removing his coat, rolling up his sleeves, and washing his hands in a dilute solution of carbolic acid before and during the surgery.




Figure 12.2.*c*: Joseph Lister initiated the use of a carbolic acid (phenol) during surgeries. This illustration of a surgery shows a pressurized canister of carbolic acid being sprayed over the surgical site.

Analyzing Disease in a Population

Epidemiological analyses are always carried out with reference to a population, which is the group of individuals that are at risk for the disease or condition. The population can be defined geographically, but if only a portion of the individuals in that

area are susceptible, additional criteria may be required. Susceptible individuals may be defined by particular behaviors, such as intravenous drug use, owning particular pets, or membership in an institution, such as a college. Being able to define the population is important because most measures of interest in epidemiology are made with reference to the size of the population.

The state of being diseased is called **morbidity**. Morbidity in a population can be expressed in a few different ways. Morbidity, or **total morbidity,** is expressed in numbers of individuals without reference to the size of the population. The **morbidity rate** can be expressed as the number of diseased individuals out of a standard number of individuals in the population, such as 100,000, or as a percent of the population.

There are two aspects of morbidity that are relevant to an epidemiologist: a disease's prevalence and its incidence. **Prevalence** is the number, or proportion, of individuals with a particular illness in a given population at a point in time. For example, the Centers for Disease Control and Prevention (CDC) estimated that about 1.8 million people in the United States lived with an active human immunodeficiency virus (HIV) infection in 2023. Expressed as a proportion, or rate, this is a prevalence of 534 infected persons per 100,000 in the population. On the other hand, **incidence** is the number or proportion of new cases in a period of time. For the same year, the CDC estimates that there were 58,512 newly diagnosed cases of HIV infection, which is an incidence of 17.3 new cases per 100,000 in the population. The relationship between incidence and prevalence can be seen in figure $12.2.d$. For a chronic disease like HIV infection, prevalence will generally be higher than incidence because it represents the cumulative number of new cases over many years minus the number of cases that are no longer active (because the patient died or was cured).

Interactive Element

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this Interactive Content online

Figure $12.2.d$: New Cases, Deaths, and people living with HIV/AIDS, world (1980-present). By Max Roser(opens in new window) and Hannah Ritchie(opens in new window) **Data source:** IHME, Global Burden of Disease (2025) – Learn more about this data. OurWorldinData.org(opens in new window) [OurWorldinData.org] | CC BY(opens in new window)

In addition to morbidity rates, the incidence and prevalence of **mortality** (death) may also be reported. A mortality rate can be expressed as the percentage of the population that has died from a disease or as the number of deaths per 100,000 persons (or other suitable standard number).

Patterns of Incidence

Diseases that are seen only occasionally, and usually without geographic concentration, are called **sporadic diseases**. Examples of sporadic diseases include tetanus, rabies, and plague. In the United States, *Clostridium tetani*, the bacterium that causes tetanus, is ubiquitous in the soil environment, but incidences of infection occur only rarely and in scattered locations because most individuals are vaccinated, clean wounds appropriately, or are only rarely in a situation that would cause infection. Likewise in the United States there are a few scattered cases of plague each year, usually contracted from rodents in rural areas in the western states.

Diseases that are constantly present (often at a low level) in a population within a particular geographic region are called **endemic diseases**. For example, malaria is endemic to some regions of Brazil, but is not endemic to the United States.

Diseases for which a larger than expected number of cases occurs in a short time within a geographic region are called **epidemic diseases**. Influenza is a good example of a commonly epidemic disease. Incidence patterns of influenza tend to rise each winter in the northern hemisphere. These seasonal increases are expected, so it would not be accurate to say that influenza is epidemic every winter; however, some winters have an usually large number of seasonal influenza cases in particular regions, and such situations would qualify as epidemics (figures $12.2.e - f$).




Figure 12.2.*e*: The 2007–2008 influenza season in the United States saw much higher than normal numbers of visits to emergency departments for influenza-like symptoms as compared to the previous and the following years. The national baseline is near 2.5%. All years have a small peak in January and a larger peak February to April. 2007-2008 had the largest peak in February to April. 2008 – 2009 had an additional peak from May to September. (credit: modification of work by Centers for Disease Control and Prevention)




Figure 12.2.*f*: The seasonal epidemic threshold (blue curve) is set by the CDC-based data from the previous five years. The seasonal baseline fluctuates to a high in winter. The epidemic threshold is just a half percent higher than the baseline. The actual mortality fluctuates above and below both of these lines. When actual mortality rates exceed this threshold, a disease is considered to be epidemic. As this graph shows, pneumonia- and influenza-related mortality saw pronounced epidemics during the winters of 2003–2004, 2005, and 2008. (credit: modification of work by Centers for Disease Control and Prevention)

An epidemic that occurs on a worldwide scale is called a **pandemic disease**. For example, HIV/AIDS (acquired immunodeficiency syndrome) and coronavirus disease 19 (COVID-19) are pandemic diseases, and novel influenza virus strains often become pandemic.

Etiology

When studying an epidemic, an epidemiologist's first task is to determinate the cause of the disease, called the etiologic agent or causative agent. Connecting a disease to a specific pathogen can be challenging because of the extra effort typically required to demonstrate direct causation as opposed to a simple association. It is not enough to observe an association between a disease and a suspected pathogen; controlled experiments are needed to eliminate other possible causes. In addition, pathogens are

typically difficult to detect when there is no immediate clue as to what is causing the outbreak. Signs and symptoms of disease are also commonly nonspecific, meaning that many different agents can give rise to the same set of signs and symptoms. This complicates diagnosis even when a causative agent is familiar to scientists.

Robert Koch was the first scientist to specifically demonstrate the causative agent of a disease (anthrax) in the late 1800s. Koch developed four criteria, now known as Koch's postulates, which had to be met in order to positively link a disease with a pathogenic microbe (figure $12.2.g$). Between 1876 and 1905, many common diseases were linked with their etiologic agents, including cholera, diphtheria, gonorrhea, meningitis, plague, syphilis, tetanus, and tuberculosis. Today, we use the molecular Koch's postulates, a variation of Koch's original postulates. (You can read more about both here.)




Figure $12.2.g$: The steps for confirming that a pathogen is the cause of a particular disease using Koch's postulates. (1) Only diseased organisms should contain the suspected causative agent (potential pathogen). (2) The causative agent from diseased organisms must be cultured (grown in the lab). (3) The cultured agent must then cause the same disease when a healthy, susceptible individual is infected with it (inoculated). (4) The same causative agent should be isolated and cultured from the individual in the previous step. Image and caption (modified) from OpenStax (CC-BY(opens in new window)).

How Diseases Spread

Understanding how infectious pathogens spread is critical to preventing infectious disease. For pathogens to persist over long periods of time they require **reservoirs** where they normally reside. Reservoirs can be living organisms (rats, bats, racoons, livestock, etc.) or nonliving sites. Nonliving reservoirs can include soil and water in the environment. These may naturally harbor the organism because it may grow in that environment. These environments may also become contaminated with pathogens in human feces, pathogens shed by intermediate hosts, or pathogens contained in the remains of intermediate hosts.

An individual capable of transmitting a pathogen without displaying symptoms is referred to as a **carrier**. A **passive carrier** is contaminated with the pathogen and can mechanically transmit it to another host; however, a passive carrier is not infected. By contrast, an **active carrier** is an infected individual who can transmit the disease to others. An active carrier may or may not exhibit signs or symptoms of infection.

Regardless of the reservoir, transmission must occur for an infection to spread. **Contact transmission** includes **direct contact** or **indirect contact**. Person-to-person transmission is a form of direct contact transmission. Here the agent is transmitted by physical contact between two individuals (figure $12.2.h$) through actions such as touching, kissing, sexual intercourse, or droplet sprays. Indirect contact transmission involves inanimate objects called **fomites** that become contaminated by pathogens from an infected individual or reservoir (figure $12.2.i$). For example, an individual with the common cold may sneeze, causing droplets to land on a fomite such as a tablecloth or carpet, or the individual may wipe her nose and then transfer mucus to a fomite such as a doorknob or towel.



Figure 12.2.*h*: Direct contact transmission of pathogens can occur through physical contact. Many pathogens require contact with a mucous membrane to enter the body, but the host may transfer the pathogen from another point of contact (for example, hand) to a mucous membrane (for example, mouth or eye). (credit left: modification of work by Lisa Doehnert)



Figure 12.2.*i*: Fomites are nonliving objects that facilitate the indirect transmission of pathogens. Contaminated doorknobs, towels, and syringes are all common examples of fomites. (credit left: modification of work by Kate Ter Haar; credit middle: modification of work by Vernon Swanepoel; credit right: modification of work by "Zaldylmg"/Flickr)

Vehicle transmission refers to the transmission of pathogens through vehicles such as water, food, and air. Water contamination through poor sanitation methods leads to waterborne transmission of disease. **Waterborne** disease remains a serious problem in many regions throughout the world. The World Health Organization (WHO) estimates that contaminated drinking water is responsible for more than 500,000 deaths each year. Similarly, food contaminated through poor handling or storage can lead to **foodborne** transmission of disease. Dust and fine particles known as **aerosols**, which can float in the air, can carry pathogens and facilitate the **airborne** transmission of disease.

Diseases can also be transmitted by a mechanical or biological **vector**, an animal (typically an arthropod) that carries the disease from one host to another. Ticks, fleas, and mosquitos are examples of vectors (figure 12.2.*j*). **Mechanical transmission** is facilitated by a **mechanical vector**, an animal that carries a pathogen from one host to another without being infected itself. **Biological transmission** occurs when the pathogen reproduces within a **biological vector** that transmits the pathogen from one host to another (figure 12.2.*k*).




Figure 12.2.*j*: The *Anopheles* mosquito is the **vector** for *Plasmodium*, which causes malaria. Image by Alan R Walker(opens in new window) (CC-BY-SA(opens in new window)).




Figure 12.2.*k*: (a) A mechanical vector carries a pathogen on its body from one host to another, not as an infection. Step 1: fly picks up pathogen from fecal matter and carries it on its body. 2: Fly transfers pathogen to food. 3: Person eats contaminated food and gets sick. (b) A biological vector carries a pathogen from one host to another after becoming infected itself. Step 1: Infected mosquito bites uninfected person. 2: Infections spreads through body and into red blood cells. 3: Second mosquito bites infected person. Mosquito may now transmit infection to another person.

Ecosystem degradation can increase disease spread through several mechanisms. First, it can promote vector populations. For example, deforestation, dams, and urbanization increase the amount of standing water, increasing habitat for vectors, such as mosquitoes. Additionally, climate change may expand the range of disease vectors that are limited to tropical or subtropical regions. Second, disruptions in the water cycle can create conditions that favor pathogens. For example, fertilizer runoff from agriculture increases nutrient levels, making bodies of water more suitable for the bacterium that causes cholera. Increased flooding as a result of changes in land use and climate change also facilitates the spread of waterborne disease.

A third way that ecosystem degradation can increase disease spread is through decreasing biodiversity. For example, Lyme disease is caused by the bacterium *Borrelia* and is transmitted to humans from small mammals (the reservoirs) by ticks (figures $12.2.l - m$). Some reservoirs (squirrels and opossums) remove ticks, reducing infection, but field mice do not. Habitat fragmentation has increased field mice relative to squirrels and opossums and thus increased the reservoir for Lyme disease.




Figure 12.2.*l*: Drawing of the bacterium *Borrelia burgdorferi*, which causes Lyme disease. Image by Lamiot(opens in new window) (CC-BY-SA(opens in new window)).




Figure 12.2.*m*: Life cycle of the bacterium *Borrelia burgdorferi*, which causes Lyme disease. Ticks can acquire this bacterium when the feed on the initial host and then transmit the disease to humans. Step 1: Adult females drop off host to lay eggs. 2: Eggs hatch into six-legged larvae. 3: Larvae attach to and feed on first host (various small mammals) and may acquire *B. burgdorferi*. 4: Larvae molt into nymphs after leaving first host. 5: Nymphs attach to and feed on second host and may acquire *B. burgdorferi*. 6: Nymphs molt into adults after leaving second host. 7: Adults attach to the third host (such as deer) for feeding and mating. 8: An arrow branches from step 4 indicating that infected nymphs feed on humans, transmitting *B. burgdorferi*. 9: Infected adults feed on dogs and sometimes humans, transmitting *B. burgdorferi*. Image from Tick-Borne Disease Working Group 2018 Report to Congress(opens in new window).

Individuals suspected or known to have been exposed to certain contagious pathogens may be **quarantined**, or isolated to prevent transmission of the disease to others. Hospitals and other health-care facilities generally set up special wards to isolate patients with particularly hazardous diseases such as tuberculosis or COVID-19 (figure 12.2.*n*). Depending on the setting, these wards may be equipped with special air-handling methods, and personnel may implement special protocols to limit the risk of transmission, such as personal protective equipment or the use of chemical disinfectant sprays upon entry and exit of medical personnel.



(a) (b)

Figure 12.2.n: (a) The Aeromedical Biological Containment System (ABCS) is a module designed by the CDC and Department of Defense specifically for transporting highly contagious patients by air. (b) An isolation ward for Ebola patients in Lagos, Nigeria. (credit a: modification of work by Centers for Disease Control and Prevention; credit b: modification of work by CDC Global)

Attribution

Modified by Melissa Ha from Disease and Epidemiology from *Microbiology* by OpenStax (licensed under CC-BY(opens in new window)). Access for free at openstax.org(opens in new window).

This page titled 12.2: Epidemiology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15.2: Epidemiology by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

12.3: Infectious Diseases

Worldwide, infectious diseases (biological hazards) accounted for three of the 10 leading causes of death in 2020. In children under five leading causes of death include acute respiratory infections (from indoor air pollution); diarrheal diseases (mostly from poor water, sanitation, and hygiene); and other infectious diseases such as malaria. Children are especially susceptible to environmental factors that put them at risk of developing illness early in life. **Malnutrition** (the condition that occurs when body does not get enough nutrients) is an important contributor to child mortality—malnutrition and environmental infections are inextricably linked.

The World Health Organization (WHO) recently concluded that about 50% of the consequences of malnutrition are in fact caused by inadequate water and sanitation provision and poor hygienic practices. For example, intestinal worms, which thrive in poor sanitary conditions, infect close to 90 percent of children in the low-income countries and, depending on the severity of the infection may lead to malnutrition, anemia, or stunted growth. About 6 million people are blind from trachoma, a disease caused by the lack of clean water combined with poor hygiene practices.

An **emerging infectious disease** is either new to the human population or has shown an increase in prevalence in the previous twenty years. Examples include coronavirus disease 19 (COVID-19), Ebola virus disease, West Nile virus, Zika virus, sudden acute respiratory syndrome (SARS), H1N1 influenza; swine and avian influenza (swine, bird flu), human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDs), and a variety of other viral, bacterial, and protozoal diseases (table $12.3.a$). Whether the disease is new or conditions have changed to cause an increase in frequency, its status as emerging implies the need to apply resources to understand and control its growing impact. Emerging diseases may change their frequency gradually over time, or they may experience sudden widespread growth.

Table $12.3.a$***:*** *Some Emerging and Reemerging Infectious Diseases*

Disease	Pathogen	Year Discovered	Affected Regions	Transmission
AIDS	HIV	1981	Worldwide	Contact with infected body fluids
Chikungunya fever	Chikungunya virus	1952	Africa, Asia, India; spreading to Europe and the Americas	Mosquito-borne
Ebola virus disease	Ebola virus	1976	Central and Western Africa	Contact with infected body fluids
H1N1 Influenza (swine flu)	H1N1 virus	2009	Worldwide	Droplet transmission
Lyme disease	*Borrelia burgdorferi* bacterium	1981	Northern hemisphere	From mammal reservoirs to humans by tick vectors
West Nile virus disease	West Nile virus	1937	Africa, Australia, Canada to Venezuela, Europe, Middle East, Western Asia	Mosquito-borne
COVID-19	SARS-CoV-2(Severe Acute Respiratory Syndrome Coronavirus 2)	2019	Worldwide	infectious particles are breathed in or come into contact with the eyes, nose, or mouth

Emerging diseases are found in all countries, low-, middle- and and high-income nations. Some nations are better equipped to deal with them. National and international public health agencies watch for epidemics in low-income countries because those countries rarely have the health-care infrastructure and expertise to deal with large outbreaks effectively. In addition to the altruistic goal of saving lives and assisting nations lacking in resources, the global nature of transportation means that an outbreak anywhere can spread quickly to every corner of the planet. Managing an epidemic in one location—its source—is far easier than fighting it on many fronts.

In 2015, the WHO set priorities on several emerging diseases that had a high probability of causing epidemics and that were poorly understood (and thus urgently required research and development efforts).

A **reemerging infectious disease** is a disease that is increasing in frequency after a previous period of decline. Its reemergence may be a result of changing conditions or old prevention regimes that are no longer working. Examples of such diseases are drug-resistant forms of tuberculosis, bacterial pneumonia, and malaria. Drug-resistant strains of the bacteria causing gonorrhea and syphilis are also becoming more widespread, raising concerns of untreatable infections.

A variety of environmental factors may contribute to re-emergence of a particular disease, including temperature, moisture, human food or animal feed sources, etc. Disease re-emergence may be caused by the coincidence of several of these environmental and/or social factors to allow optimal conditions for transmission of the disease.

It seems likely that a wide variety of infectious diseases have affected human populations for thousands of years emerging when the environmental, host, and agent conditions were favorable. Expanding human populations have increased the potential for transmission of infectious disease as a result of close human proximity and increased likelihood for humans to be in “the wrong place at the right time” for disease to occur (eg, natural disasters or political conflicts). Global travel increases the potential for a carrier of disease to transmit infection thousands of miles away in just a few hours, as evidenced by WHO precautions concerning international travel and health.

Selected Infectious Diseases

COVID-19

Coronavirus disease 19 (**COVID-19**) is a respiratory illness caused by a virus called novel species of coronavirus (SARS-CoV-2). Symptoms include fever or chills, cough, shortness or breath or difficulty breathing, fatigue, headache, and sore throat. The WHO declared coronavirus disease 19 (COVID-19) a public health emergency of international concern on January 30 and a pandemic on March 11, 2020. A seemingly limited cluster of cases of pneumonia linked to a sea food market in Wuhan, China has become one of the worst pandemics in human history with a staggering number of more than 90.2 million infections in 177 countries and 1.9 million deaths globally as of January 10, 2021 (figure 12.3.a). This illustrates how locally emerging pathogens have the capacity to spread rapidly and cross borders and become an imminent public health threat to the entire world. You can view the the latest COVID-19 data here.




Figure 12.3.a: New confirmed COVID-19 cases per million people from March 1, 2020 to January 10, 2021 in selected countries. Image by Max Roser, Hannah Ritchie, Esteban Ortiz-Ospina and Joe Hasell (2020) - "Coronavirus Pandemic (COVID-19)". Published online at OurWorldInData.org. (CC-BY).

Studies indicate that SARS-CoV-2 might originate from bat coronaviruses and infects humans directly or through other animals as intermediaries. It primarily spreads between humans through the respiratory tract by involving droplets, aerosols, respiratory

secretions, or saliva. It is most often spread by direct contact transmission, but it airborne transmission has been documented in certain situations, such as in an enclosed space (figure 12.3.*b*). Researchers continue to investigate airborne transmission of the virus. This coronavirus is infectious during its incubation period, which is reported to be 3-7 days, at most 14 days, when no symptoms are shown in patients. Those infected may experience mild to moderate disease (80%), severe disease (15%), and critical illness (5%) with an overall case fatality rate of 0.5–2.8% with much higher rates (3.7–14.8%) in octogenarians. The severe and critical illness categories (about 20% of all infections) have overwhelmed health systems worldwide (figure 12.3.*c*). The Centers for Disease Control and Prevention's COVID-19 page has the latest information about preventing transmission, vaccines, symptoms, and testing.




Figure 12.3.*b*: Origin of SARS-CoV-2 and potential transmission routes to humans. SARS-CoV-2 transmitted to humans from bats directly or by intermediate hosts such as rats, pangolins, snakes, and rats. The virus is spread among humans by different routes including droplets, aerosols, direct contact, and other potential routes such as urine, stool, and fecal swabs. Human organs with a protein called the ACE2 receptor (ACE2+) are targets for viral infection. Disease spread can be limited through quarantining when sick, social distancing, avoiding large gatherings (not shown), wearing masks (not shown), and vaccinations (not shown). Image and caption (modified) from Tizaoui, K., Zidi, I., Lee, K. H., Ghayda, R. A., Hong, S. H., Li, H., Smith, L., Koyanagi, A., Jacob, L., Kronbichler, A., & Shin, J. I. (2020). Update of the current knowledge on genetics, evolution, immunopathogenesis, and transmission for coronavirus disease 19 (COVID-19). *International journal of biological sciences*, *16*(15), 2906–2923. https://doi.org/10.7150/ijbs.48812 (CC-BY).



Figure 12.3.c: Lieutenant (junior grade) Natasha McClinton, a surgical nurse, prepares a patient for a procedure in the intensive care unit aboard the U.S. hospital ship USNS Comfort. The ship cares for critical and non-critical patients without regard to their COVID-19 status. In April of 2020, Comfort worked with Javits New York Medical Station as an integrated system to relieve the New York City medical system as a response to the COVID-19 pandemic. Image and caption (modified) from US Navy Mass Communication Specialist 2nd Class Sara Eshleman (public domain).

Ebola Virus Disease

Ebola virus disease, previously known as Ebola hemorrhagic fever, is a rare and deadly disease caused by infection with one of the Ebola virus strains (figure 12.3.d). Ebola can cause disease in humans and nonhuman primates. Health experts had been aware of the Ebola virus since the 1970s, but outbreak at scale that had not yet been seen occurred in western Africa through 2014–2016. Previous human epidemics had been small, isolated, and contained. Indeed, the gorilla and chimpanzee populations of western Africa had suffered far worse from Ebola than the human population. The pattern of small isolated human epidemics changed in 2014. Its high transmission rate, coupled with cultural practices for treatment of the dead and perhaps its emergence in an urban setting, caused the disease to spread rapidly, and thousands of people died. The international public health community responded with a large emergency effort to treat patients and contain the epidemic.




Figure 12.3.d: The Ebola virus. Image by CDC (public domain).

The 2014-2016 Ebola epidemic was the largest in history (with 28,616 cases and 11,310 deaths), affecting multiple countries in West Africa. There were a small number of cases reported in Nigeria and Mali and a single case reported in Senegal; however, these cases were contained, with no further spread in these countries. Even with the support of international agencies, the

systems in western Africa struggled to identify and care for the sick and control spread. Engaging local leaders in prevention programs and messaging, along with careful policy implementation at the national and global level, helped to eventually contain the spread of the virus and put an end to this outbreak. The U.S. Food and Drug Administration approved a vaccine for Ebola late in 2019.

HIV/AIDS

Human immunodeficiency virus (**HIV**) is a virus that attacks the body's immune system. If HIV is not treated, it can lead to **acquired immunodeficiency syndrome** (**AIDS**). Those with AIDS have weakened immune systems and are thus more susceptible to cancers and other infections. First identified in 1981, the **HIV/AIDS** epidemic has spread with ferocious speed. In 2019, 38 million people were living HIV globally. Over 95 percent of people living with HIV are in low- and middle- income countries. There were 1.7 million new infections in 2019, and more than half of new infections are among young people below age 25. In 2019 alone, 690,000 million people died from AIDS and related illnesses. In 2016, HIV/AIDS is was the second leading cause of death in Africa. At its peak, the epidemic cut life expectancy by more than 10 years in several nations (figure $12.3.e$).




Figure $12.3.e$: Life expectancy in six countries in Sub-Saharan Africa dropped dramatically in the late 1990s and early 2000s. As the efforts to control the pandemic reduced the number of new infections (incidence) and death rates from HIV, life expectancy has again increased. Image by Max Roser and Hannah Ritchie (2018) - "HIV / AIDS". Published online at OurWorldInData.org. (CC-BY).

Malaria

Approximately 40% of the world's people—mostly those living in the world's poorest countries—are at risk from malaria. **Malaria** is an infectious disease spread by mosquitoes but caused by a single-celled protozoan parasite called *Plasmodium*. There were 229 million cases of malaria and 409,000 deaths in 2019 with most cases and deaths found in Sub-Saharan Africa (**figure** $12.3.f$ **).** However, Asia, Latin America, the Middle East, and parts of Europe are also affected. Pregnant women are especially at high risk of malaria. Non-immune pregnant women risk both acute and severe clinical disease, resulting in fetal loss in up to 60% of such women and maternal deaths in more than 10%, including a 50% mortality rate for those with severe disease. Semi-immune pregnant women with malaria infection risk severe anemia (lack of sufficient red blood cells to deliver oxygen) and impaired fetal growth, even if they show no signs of acute clinical disease. An estimated 10,000 women and 200,000 infants die annually as a result of malaria infection during pregnancy.




Figure 12.3.*f*: Deaths caused by malaria from 1990 to 2017. Deaths peaked above 800,000 around 2004 and were just above 600,000 in 2017. The largest share of deaths occurred in Sub-Saharan Africa. Image by Max Roser and Hannah Ritchie (2013) - "Malaria". Published online at OurWorldInData.org. (CC-BY).

Tuberculosis

The WHO estimates that 10.4 million new cases and 1.5 million deaths occur from **tuberculosis** (**TB**) each year. One-third of TB cases remain unknown to the health care system. For those accessing treatment, however, prevalence and mortality have declined significantly, and millions of lives have been saved.

Tuberculosis is caused by the bacterium *Mycobacterium tuberculosis*, which is transmitted between humans through the respiratory route and most commonly affects the lungs but can damage any tissue. When an infected individual coughs, exhales, speaks, etc., the bacterium is released in droplets that can remain suspended in the air for several hours. Only a minority (approximately 10 percent) of individuals infected with *M. tuberculosis* progress to active TB disease, while the remainder may maintain a latent infection that serves as a reservoir. TB has special challenges, including (a) a substantial number of patients with active disease are asymptomatic, capable of transmitting infection without knowing it; (b) patients must maintain compliance with treatment for six to nine months; and (c) the pathogen persists in many infected individuals in a latent state for many years but can be reactivated over a lifetime to cause disease and become transmissible.

This video by the CDC explains the basics of TB.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Disease Eradication

Many diseases have been eradicated or are nearly eradicated worldwide (Table 12.3.*b*). Strategies to eradicate disease infectious diseases include vaccination, improved hygiene and sanitation, preventative medication, and public health education. Some of these strategies are discussed in Environmental Health Reduction, and vaccination is discussed below.

Table 12.3.*b*: *Infectious diseases that have been eradicated and could be eradicated in the future.*

Disease	Burden of disease	Cause	Ways to eradicate	Fatality
Smallpox	Declared eradicated in 1980	Variola virus	Eradicated using vaccination	30%
Rinderpest	Declared eradicated in 2011	Rinderpest virus	Sanitary measures and vaccination	100%
Poliomielitis	116 cases in 2017	Poliovirus	Vaccination	For paralytic polio 2-5% in children and increases to 15-30% in adults
Guinea worm disease	30 cases in 2017	Parasitic worm *Dracunculus medinensis*	Hygiene, water decontamination and health education	Not fatal but debilitating
Measles	173,457 reported cases to WHO in 2017	Measles morbillivirus	Vaccination	15%
Mumps	560,622 reported cases to WHO in 2017	Mumps orthorubulavirus	Vaccination	0.01% for mumps-caused encephalitis
Rubella	6,789 reported cases to WHO in 2017	Rubella virus	Vaccination	Not reported
Lymphatic filariasis	No estimate available. In 2014, 68 million people were infected and 790 million people where at risk of infection	Roundworms: *W. bancrofti, B. malayi, B. timori*	Preventive chemotherapy	Not fatal but highly debilitating
Cysticercosis	2.56–8.30 million cases estimated by the WHO	Tapeworms: *T. solium, T. saginata, T. asiatica*	Sanitation and health education. Vaccination of pigs	Varies between countries <1-30%

Table modified from Max Roser, Sophie Ochmann, Hannah Behrens, Hannah Ritchie and Bernadeta Dadonaite (2014) - "Eradication of Diseases". Published online at OurWorldInData.org. (CC-BY).

Vaccination

Vaccines function in inducing a specific immune response that can immediately counter a pathogen if exposed. Vaccines lead to the production of specialized cells and antibodies, which can efficiently and effectively target pathogens. There are several types of vaccines. Some consists of dead or inactivated pathogens or parts of them. The Pfizer-BioNTech and Moderna COVID-19 vaccines, approved by the FDA in December of 2020 consist of a piece of the the virus's genetic material (called mRNA). When injected, the body uses this genetic information to synthesize the spike protein of the virus, which stimulates an immune responses without causing infection.

Development and approval of a vaccine does not automatically eradicate a disease (figure 12.3.*g*), for the vaccine must be distributed to those who need it. Access is particularly challenging in low-income countries. In the case of tuberculosis, which remains widespread, a vaccine has been available for 100 years, but it is more effective in protecting severe forms of the disease in children than lung infections in adults. A new, more effective vaccine is likely needed to eradicate tuberculosis.




Figure $12.3.g$: Global deaths in 2017 that were caused by diseases with existing vaccines. Tuberculosis was responsible for 1.18 million deaths followed by measles (95,290), whooping cough (91,804), hepatitis B (85,590), Hib meningitis (75,703), tetanus (38,134), meningococcal meningitis (29,967), yellow fever (4,786), and diphtheria (3,624). Image by Samantha Vanderslott and Bernadeta Dadonaite (2013) "Vaccination". Published online at OurWorldInData.org. (CC-BY).

New vaccines must be rigorously tested before they are approved. Not only are vaccines safe, but we would face spikes in deadly infectious diseases without them. The antivax movement has unfortunately spread false information questioning the safety of vaccines, but you can educate yourself with reliable information about the development, testing, and safety of vaccines here.

Antibiotic Resistance

Antibiotics and similar drugs, together called antimicrobial agents, have been used for the last 70 years to treat patients who have infectious diseases. Technically, **antibiotics** are substances that are produced by certain microorganisms (such as the fungus, *Penicillium*, or the bacterium *Streptomyces*) that inhibit the growth of other microorganisms. However, any drugs used to treat bacterial infections are commonly referred to as antibiotics. Neither of these definitions includes substances used to treat viral infections, which are called antivirals. Since the 1940s, the use of antiobitics has greatly reduced illness and death from infectious diseases. However, these drugs have been used so widely and for so long that the infectious organisms the antibiotics are designed to kill have adapted to them, making the drugs less effective.

Overuse of antibiotics have allowed pathogens to evolve **antibiotic resistance**, which occurs when bacteria change in a way that reduces the effectiveness of drugs, chemicals, or other agents designed to cure or prevent infections (figure $12.3.h$). Overuse takes many forms, such as the inclusion of antibiotics in household products or prescribing antibiotics to treat infections without first determining whether the infection is bacterial or viral (in which case, the antibiotic would not be effective). Similar to the evolution of pesticide resistance, widespread use of antibiotics favors individuals that happen to have gene versions conferring antibiotic resistance. These individuals have the best chance of surviving and reproducing, so much so, that an antibiotic-resistant strain evolves. The initial antibiotic is not longer effective against this strain, and it must be treated with alternative medications. For example, tuberculosis (TB) was nearly eliminated in most parts of the world, but drug-resistant strains have now reversed that trend.

A similar example of drug resistance has evolved in *Plasmodium*, the protozoa that cause malaria. The insecticide dichlorodiphenyltrichloroethane (DDT) was widely used to control malaria-carrying mosquito populations in tropical regions. However, after many years the mosquitoes developed a natural resistance to DDT and again spread the disease widely. Anti-malarial medicines were also over-prescribed, which allowed *Plasmodium* to become drug-resistant.




Figure 12.3.*h*: Antibiotic resistance is of increasing concern to medical professionals. (1) Initially, a variety of microbes (germs) are present, and a few are drug resistant. (2) When antibiotics are used, they kill both disease-causing microbes and beneficial ones. However, drug-resistant microbes remain. (3) The drug-resistant microbes multiple and can eventually produce a drug-resistant strain. (4) Drug-resistant bacteria can even pass their resistance genes to other bacteria, including those of different species.

New forms of antibiotic resistance can cross international boundaries and spread between continents with ease. Many forms of resistance spread with remarkable speed. Each year in the United States, at least 2 million people acquire serious infections with bacteria that are resistant to one or more of the antibiotics designed to treat those infections. At least 23,000 people die each year in the US as a direct result of these antibiotic-resistant infections. Many more die from other conditions that were complicated by an antibiotic-resistant infection. The use of antibiotics is the single most important factor leading to antibiotic resistance around the world. Antibiotics are among the most commonly prescribed drugs used in human medicine, but up to 50% of all the antibiotics prescribed for people are not needed or are not optimally effective as prescribed.

The video below shows an experiment done at Harvard Medical school, where they show bacteria adapting very quickly to apparently deadly conditions.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

During recent years, there has been growing concern over methicillin-resistant *Staphylococcus aureus* (**MRSA**)**,** a bacterium that is resistant to many antibiotics. In the community, most MRSA infections are skin infections. In medical facilities, MRSA causes life-threatening bloodstream infections, pneumonia and surgical site infections.

Attribution

Modified by Erin Rempala.

Modified by Melissa Ha from the following sources:

- Environmental Health and Sustainable Agriculture from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)
- Environmental Health and Child Survival: Epidemiology, Economics, Experiences. Environment and Development. 2008.

World Bank. Washington, DC. CC-BY.
- Global Public Health from *Microbiology* by OpenStax (CC-BY). Access for free at openstax.org.
- Patel, V., D. Chisholm., T. Dua, R. Laxminarayan, and M. E. Medina-Mora, editors. 2015. *Mental, Neurological, and Substance Use Disorders*. Disease Control Priorities, third edition, volume 4. Washington, DC: World Bank. doi:10.1596/978-1-4648-0426-7. (CC-BY).
- Balkhair A. A. 2020. COVID-19 Pandemic: A New Chapter in the History of Infectious Diseases. *Oman medical journal*, *35*(2), e123. https://doi.org/10.5001/omj.2020.41 (CC-BY-NC).
- Tizaoui, K., Zidi, I., Lee, K. H., Ghayda, R. A., Hong, S. H., Li, H., Smith, L., Koyanagi, A., Jacob, L., Kronbichler, A., & Shin, J. I. (2020). Update of the current knowledge on genetics, evolution, immunopathogenesis, and transmission for coronavirus disease 19 (COVID-19). *International journal of biological sciences*, *16*(15), 2906–2923. https://doi.org/10.7150/ijbs.48812 (CC-BY).
- Ebola (Ebola Virus Disease). 2020. Centers for Disease Control and Prevention. Accessed 10 January 2021. (public domain)
- 2014-2016 Ebola Outbreak in West Africa. 2019. Centers for Disease Control and Prevention. Accessed 10 January 2021. (public domain)
- About HIV. 2020. Centers for Disease Control and Prevention. Accessed 10 January 2021. (public domain)
- HIV/AIDS at a glance. 2003. World Bank. Washington, DC. CC-BY.

This page titled 12.3: Infectious Diseases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15.3: Infectious Diseases by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

12.4: Environmental Toxicology

Environmental Toxicology

Environmental toxicology is the study of how natural and synthetic chemicals affect living organisms and ecosystems. A chemical that causes harm to an organism is called a **toxin**, and its ability to cause harm is known as its **toxicity**. Toxicologists study the movement of toxins through the environment, how they enter organisms, and the health effects they can cause.

Humans and other organisms are exposed to chemicals through the air they breathe, the water they drink, the food they eat, and the products they use. Understanding how toxins move through organisms and ecosystems helps scientists identify environmental hazards and reduce risks to human and ecological health.

Routes of Exposure to Chemicals

For a chemical to cause harm, it must first enter the body. Once absorbed, it may travel through the bloodstream and reach target organs such as the lungs, liver, kidneys, brain, or reproductive organs.

The three primary **routes of exposure** are (figure $12.4.a$):

Inhalation

Inhalation occurs when gases, vapors, smoke, dust, or aerosols are breathed into the lungs. Because the lungs are specialized for gas exchange, many inhaled chemicals can quickly enter the bloodstream.

Ingestion

Ingestion occurs when contaminated food, water, soil, or other materials are swallowed. Chemicals absorbed through the digestive system may enter the bloodstream and be transported throughout the body.

Skin and Eye Contact

Some chemicals can be absorbed directly through the skin or eyes. Others cause localized irritation or tissue damage. While healthy skin provides an effective barrier against many chemicals, some substances can penetrate intact skin and enter the bloodstream.




Figure 12.4.a: The routes of exposure through which toxins can enter the body (inhalation, ingestion, and skin or eye contact) are boxed in pink. Image by BruceBlaus (opens in new window)(CC-BY-SA(opens in new window)).

What Determines Toxicity?

Not all chemicals are equally hazardous. Several factors influence the degree of harm a chemical may cause:

- **Dose:** the amount of a substance that enters the body.
- Duration of exposure: how long exposure lasts.
- **Potency:** the amount needed to cause harm.
- **Persistence:** how long the chemical remains in the environment.
- **Solubility:** whether the chemical dissolves in water or fats.
- Individual characteristics: age, genetics, health status, and life stage.

Fat-soluble chemicals are often of particular concern because they can accumulate in body tissues and remain there for long periods. Persistent chemicals also present greater risks because they may remain in ecosystems for years or even decades.

Persistent Organic Pollutants (POPs)

Persistent organic pollutants (POPs) are synthetic organic chemicals that resist environmental breakdown, accumulate in living organisms, and can cause harmful effects to humans and wildlife. Examples include the pesticide **DDT**, industrial chemicals called **polychlorinated biphenyls (PCBs)**, some **per- and polyfluoroalkyl substances (PFAS)**, and dioxins such as **TCDD**.

POPs share three defining characteristics:

1. **Persistent:** They remain in soils, sediments, water, and organisms for years or decades.
2. **Bioaccumulative:** They build up in organisms over time.
3. **Toxic:** They can damage living organisms even at low concentrations.

Because of these properties, many POPs have been banned or restricted through international agreements such as the Stockholm Convention on Persistent Organic Pollutants. However, many remain in the environment because they degrade very slowly.

Long-Distance Transport: The Grasshopper Effect

POPs can travel far from their original source through a process known as the **grasshopper effect**. A POP may evaporate into the atmosphere in a warm region, travel with air currents, and then condense and settle in a cooler location. Repeated cycles of evaporation and deposition allow POPs to "hop" thousands of kilometers across the globe.

As a result, POPs are commonly detected in remote regions such as the Arctic, far from the areas where they were originally released.

Bioaccumulation and Biomagnification

Some toxins become increasingly concentrated within organisms and food webs.

Bioaccumulation

Bioaccumulation is the gradual buildup of a chemical within an organism over its lifetime. This occurs when a substance enters the body faster than it can be eliminated.

Many POPs bioaccumulate because they dissolve readily in fats and are stored in fatty tissues. However, some water-soluble substances can also bioaccumulate. For example, lead can accumulate in bones and teeth, while mercury may accumulate in the brain and kidneys.

Biomagnification

Biomagnification is the increase in toxin concentration at each successive trophic level in a food chain (figure $12.4.b$). When contaminated organisms are eaten, toxins are transferred to predators. As predators consume many prey items throughout their lifetimes, toxin concentrations can become progressively higher.

As a result, top predators such as tuna, swordfish, eagles, seals, whales, polar bears, and humans may contain the highest concentrations of certain toxins.




Figure $12.4.b$: Biomagnification causes toxin concentrations to increase at successively higher trophic levels. Organisms at the top of food chains often contain much higher concentrations of contaminants than organisms lower in the food web. Image credit: Patricia Van Hoof, NOAA, GLERL

One of the most famous examples involved the insecticide **DDT**. Small amounts accumulated in aquatic organisms and became increasingly concentrated in fish-eating birds such as bald eagles, pelicans, and ospreys, where they caused eggshell thinning and reproductive failure.

Health Effects of Toxins

The effects of a toxin depend on the chemical itself as well as the characteristics of the exposed individual.

Children are often more vulnerable to toxins because their organs and nervous systems are still developing. Older adults may also be more susceptible because aging can reduce the body's ability to process and eliminate contaminants. Genetics, overall health, and exposure to other chemicals can also influence toxicity.

Many toxins have been linked to:

- cancer
- neurological damage
- developmental abnormalities
- reproductive problems
- immune system suppression
- endocrine disruption

Endocrine Disruptors

Some environmental contaminants can interfere with hormones and are known as **endocrine disruptors**. Because hormones regulate growth, development, metabolism, and reproduction, endocrine disruptors may affect organisms even at very low concentrations.

Acute and Chronic Effects

An **acute effect** occurs rapidly after exposure, usually following a relatively high dose. A **chronic effect** develops after repeated or long-term exposure to lower doses and may take months or years to become apparent. For example, carbon monoxide poisoning is primarily an acute effect, whereas lead exposure and many cancers result from chronic exposure.

Toxicological Interactions

Organisms are often exposed to multiple chemicals simultaneously. These mixtures may interact in different ways. An **additive effect** occurs when the combined effect equals the sum of the individual effects. A **synergistic effect** occurs when the combined effect is greater than expected. For example, smoking and asbestos exposure together greatly increase the risk of lung cancer. An **antagonistic effect** occurs when one chemical reduces the effect of another. For example, ethanol can be used to treat methanol poisoning because it slows the conversion of methanol into more toxic compounds.




Figure 12.4.*f*: Chemical interactions may be additive, synergistic, or antagonistic. Image modified from Tae Jin Cho et al., MDPI, Basel, Switzerland(opens in new window) (CC-BY).

Dose, Risk, and the Precautionary Principle

Toxicologists often say: "The dose makes the poison."

In general, larger doses are more likely to produce harmful effects than smaller doses. Scientists use **dose-response relationships** to determine how organisms respond to increasing amounts of a chemical and to estimate safe exposure levels.

However, uncertainty often exists, especially for chronic exposures and new chemicals whose long-term effects are not fully understood. For this reason, environmental health decisions often apply the **precautionary principle**, which states that when there is a plausible risk of harm, protective action should be taken even if cause-and-effect relationships have not been fully

established scientifically.

In simple terms, the precautionary principle can be summarized as:

Better safe than sorry.

The precautionary principle influences environmental regulations around the world. For example, some countries require manufacturers to demonstrate that a chemical is safe before it can be marketed, while others permit use until evidence of harm accumulates. By emphasizing prevention rather than reaction, the precautionary principle seeks to protect both human health and ecological systems from avoidable harm.

Attribution

This page was heavily revised and rewritten by Erin Rempala and contains adapted content from content originally authored by Melissa Ha and Rachel Schleiger and is licensed CC BY-NC-SA 4.0.

Modified by Melissa Ha from the following sources:

- Persistent Organic Pollutants: Backyards to Borders. 2004. World Bank. (licensed under CC-BY)
- *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)
  - Pests and Pesticides
  - Environmental Toxicology
  - The Impacts of Environmental Conditions
  - Environmental Health
  - Energy Flow through Ecosystems
- Biomagnification, Insecticides, and the LD50 test from *Biology* by John W. Kimball (licensed under CC-BY)

This page titled 12.4: Environmental Toxicology is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15.4: Environmental Toxicology by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

12.5: Case Study - The Love Canal Disaster

One of the most famous and important examples of groundwater pollution in the U.S. is the **Love Canal tragedy** in Niagara Falls, New York. It is important because the pollution disaster at Love Canal, along with similar pollution calamities at that time (Times Beach, Missouri and Valley of Drums, Kentucky), helped to create **Superfund**, a federal program instituted in 1980 and designed to identify and clean up the worst of the hazardous chemical waste sites in the U.S.



Figure 12.5.1. Love Canal. Source: US Environmental Protection Agency

Love Canal is a neighborhood in Niagara Falls named after a large ditch (approximately 15 m wide, 3–12 m deep, and 1600 m long) that was dug in the 1890s for hydroelectric power. The ditch was abandoned before it actually generated any power and went mostly unused for decades, except for swimming by local residents. In the 1920s Niagara Falls began dumping urban waste into Love Canal, and in the 1940s the U.S. Army dumped waste from World War II there, including waste from the frantic effort to build a nuclear bomb. Hooker Chemical purchased the land in 1942 and lined it with clay. Then, the company put into Love Canal an estimated 21,000 tons of hazardous chemical waste, including the carcinogens benzene, dioxin, and PCBs in large metal barrels and covered them with more clay. In 1953, Hooker sold the land to the Niagara Falls school board for $1, and included a clause in the sales contract that both described the land use (filled with chemical waste) and absolved them from any future damage claims from the buried waste. The school board promptly built a public school on the site and sold the surrounding land for a housing project that built 200 or so homes along the canal banks and another 1,000 in the neighborhood (Figure 12.5.1). During construction, the canal's clay cap and walls were breached, damaging some of the metal barrels.

Eventually, the chemical waste seeped into people's basements, and the metal barrels worked their way to the surface. Trees and gardens began to die; bicycle tires and the rubber soles of children's shoes disintegrated in noxious puddles. From the 1950s to the late 1970s, residents repeatedly complained of strange odors and substances that surfaced in their yards. City officials investigated the area, but did not act to solve the problem. Local residents allegedly experienced major health problems including high rates of miscarriages, birth defects, and chromosome damage, but studies by the New York State Health Department disputed that. Finally, in 1978 President Carter declared a state of emergency at Love Canal, making it the first human-caused environmental problem to be designated that way. The Love Canal incident became a symbol of improperly stored chemical waste. Clean up of Love Canal, which was funded by Superfund and completely finished in 2004, involved removing contaminated soil, installing drainage pipes to capture contaminated groundwater for treatment, and covering it with clay and plastic. In 1995, Occidental Chemical (the modern name for Hooker Chemical) paid $102 million to Superfund for cleanup and $27 million to Federal Emergency Management Association for the relocation of more than 1,000 families. New York State paid $98 million to EPA and the US government paid $8 million for pollution by the Army. The total clean up cost was estimated to be $275 million.

The Love Canal tragedy helped to create Superfund, which has analyzed tens of thousands of hazardous waste sites in the U.S. and cleaned up hundreds of the worst ones. Nevertheless, over 1,000 major hazardous waste sites with a significant risk to human health or the environment are still in the process of being cleaned.

Contributors and Attributions

- Essentials of Environmental Science(opens in new window) by Kamala Doršner is licensed under CC BY 4.0(opens in new window). Modified from the original by Matthew R. Fisher.

This page titled 12.5: Case Study - The Love Canal Disaster is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Matthew R. Fisher (OpenOregon) via source content that was edited to the style and standards of the LibreTexts platform.

- 6.5: Case Study - The Love Canal Disaster by Matthew R. Fisher is licensed CC BY 4.0. Original source: https://openoregon.pressbooks.pub/envirobiology.

12.6: Environmental Hazard Reduction

Public Health Organizations

A large number of international programs and agencies are involved in efforts to promote global public health. Among their goals are developing infrastructure in health care, public sanitation, and public health capacity; monitoring infectious disease occurrences around the world; coordinating communications between national public health agencies in various countries; and coordinating international responses to major health crises. In large part, these international efforts are necessary because disease-causing microorganisms know no national boundaries.

International public health issues are coordinated by the **World Health Organization** (WHO), an agency of the United Nations. Of its roughly $4 billion budget for 2015–16, about $1 billion was funded by member states and the remaining $3 billion by voluntary contributions. In addition to monitoring and reporting on infectious disease, WHO also develops and implements strategies for their control and prevention. WHO has had a number of successful international public health campaigns. For example, its vaccination program against smallpox, begun in the mid-1960s, resulted in the global eradication of the disease by 1980. WHO continues to be involved in infectious disease control, primarily in low-income countries, with programs targeting malaria, HIV/AIDS, and tuberculosis, among others. It also runs programs to reduce illness and mortality that occur as a result of violence, accidents, lifestyle-associated illnesses such as diabetes, and poor health-care infrastructure.

WHO maintains a global alert and response system that coordinates information from member nations. In the event of a public health emergency or epidemic, it provides logistical support and coordinates international response to the emergency. The United States contributes to this effort through the **Centers for Disease Control and Prevention** (CDC), an agency of the Department of Health and Human Services (figure 12.6.*a*). The CDC carries out international monitoring and public health efforts, mainly in the service of protecting US public health in an increasingly connected world. Similarly, the European Union maintains a Health Security Committee that monitors disease outbreaks within its member countries and internationally, coordinating with WHO.



Figure 12.6.*a*: The Centers for Disease Control and Prevention is charged with protecting the public from disease and injury. Image by CDC (public domain).

One way that the CDC carries out this mission is by overseeing the National Notifiable Disease Surveillance System (NNDSS) in cooperation with regional, state, and territorial public health departments. The NNDSS monitors diseases considered to be of public health importance on a national scale. Such diseases are called notifiable diseases or reportable diseases because all cases must be reported to the CDC. A physician treating a patient with a notifiable disease is legally required to submit a report on the case. Notifiable diseases include HIV infection, measles, West Nile virus infections, and many others. Some states have their own lists of notifiable diseases that include diseases beyond those on the CDC's list.

Notifiable diseases are tracked by epidemiological studies and the data is used to inform health-care providers and the public about possible risks. The CDC publishes the *Morbidity and Mortality Weekly Report* (*MMWR*), which provides physicians and health-care workers with updates on public health issues and the latest data pertaining to notifiable diseases. Table 12.6.*a* is an example of the kind of data contained in the *MMWR*.

***Table** 12.6.a: Incidence of Four Notifiable Diseases in the United States, Week Ending January 2, 2016*

Disease	Current Week (Jan 2, 2016)	Median of Previous 52 Weeks	Maximum of Previous 52 Weeks	Cumulative Cases 2015
Campylobacteriosis	406	869	1,385	46,618
Chlamydia trachomatis infection	11,024	28,562	31,089	1,425,303
Giardiasis	115	230	335	11,870
Gonorrhea	3,207	7,155	8,283	369,926

The current *Morbidity and Mortality Weekly Report* is available online.

Strategies for Reducing Environmental Hazards

Providing access to clean water is an important strategy for reducing environmental hazards. Because lack of clean water is responsible for diarrheal and other infectious diseases, particular in low-income countries, access to clean water limits disease spread (a biological hazard). Furthermore, clean water reduces exposure to toxins (chemical hazards). This can be achieved through digging wells and establishing water treatment facilities. Improving sanitation and hygiene (for example, by supplying latrines, [figure 12.6.*b*] and handwashing stations) further reduces the spread of infectious disease.



Figure 12.6.*b*: When adequate plumbing is not available, latrines, such as this pit latrine, prevent sewage from polluting water supplies. Image by SuSanA Secretariat(opens in new window) (CC-BY(opens in new window)).

In Sub-Saharan Africa and South Asia, the use of pipe filters helped eradicate Guinea worm disease, which is caused by a parasitic worm. People are infected by this disease when they drink water contaminated by small animals called copepods that house the larvae. Once in the body, the larvae mature in the abdomen, and adult worms eventually (and painfully) exit through the skin. If an infected person enters a body of water at this stage, adult worms release more larvae into the water, continuing their life cycle. Pipe filters are straw-like structures that contain openings small enough to allow the passage of water but not copepods, preventing infection.

Another strategy to limit exposure to biological hazards is to reduce exposure to disease vectors. Depending on the specific vector and disease this could involve removing standing water (which facilitates mosquito reproduction), application of pesticides,

or use of netting (figure $12.6.c$). Biological control is also slowly emerging in vector control in public health and in areas that for a long time mainly focused on chemical vector control of the *Anopheles* mosquito (the vector of malaria) and the black fly (the vector of river blindness, caused by a parasitic worm). The release of sterile males has been used to control the tsetse fly, the vector of African sleeping sickness. Vaccinating animals that harbor diseases (reservoirs) can also limit the chance of infection.



Figure $12.6.c$: A net excludes mosquitos, which are disease vectors. The *Anopheles* mosquito is only active at night, and thus sleeping inside of a net can limit exposure to malaria. Image by Presidents Mosquito Initiative a component of the U.S. Government's Global Health Initiative(opens in new window) (public domain).

Because air pollution exposes individuals to toxins, limiting air pollution is another means of reducing environmental hazards. This can involve energy conservation, using clean energy, such as solar or wind (rather than burning fossil fuels), enforcing air pollution standards on industry, and implementing pollution-reducing technologies, at an industry or household scale. A main source of indoor air pollution is cooking using a fire indoors. Solar ovens provide a pollution-free alternative for people who do not have access to electricity or gas for cooking (figure $12.6.d$).



Figure $12.6.d$: Solar ovens are a pollution-free option for cooking food if electricity or gas is not available. Image by SUN OVEN(opens in new window) (public domian).

The public have an important role in environmental hazard reduction. Firstly, the public must engage in behaviors to prevent disease spread and minimize contact with toxins, and public health education is critical to this. In the case of Guinea worm disease, individuals learned how to use pipe filters and shy they must avoid submerging wounds from the worms in

bodies of water to limit disease spread. In the case of COVID-19, individuals are learning to reduce disease transmission by employing frequent handwashing, mask wearing, and social distancing (figure 12.6.e).




Figure 12.6.e: The California Department of Public Health website includes guidelines for limiting disease spread. The text reads "Stay Up To Date on COVID-19. Wash Your Hands. Wear a Mask. Keep Your Distance. Stay Home if You're Sick." Image from the CDPH website(opens in new window) (public domain).

The public can also support policies that reduce exposure to environmental hazards. For example, many states have laws that reduce exposure to secondhand smoke. The Toxic Substances Control Act (TSCA) bans or regulates harmful chemicals, such as asbestos and heavy metals (figure 12.6.f)..



Figure 12.6.f: Asbestos removal from a building in the United Kingdom (UK). While asbestos is fully banned in the UK, only certain uses of asbestos are banned in the United States under the Toxic Substances Control Act . Image by Fevs101(opens in new window) (CC-BY-SA(opens in new window)).

Many strategies for reducing environmental health hazards directly align with the United Nations Sustainable Development Goals, particularly goals 3 (Good Health and Well-Being) and 6 (Clean Water and Sanitation).

Attribution

Modified by Melissa Ha from the following sources:

- Disease and Epidemiology from *Microbiology* by OpenStax (licensed under CC-BY(opens in new window))
- Sustainable Agriculture from *Environmental Biology* by Mathew R. Fisher (licensed under CC-BY(opens in new window))

This page titled 12.6: Environmental Hazard Reduction is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 15.5: Environmental Hazard Reduction by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

13: THE AGRICULTURAL SYSTEM

Introduction

Steep mountain landscapes aren't an ideal place for farming. Yet, for thousands of years farmers in mountainous regions all over the world have grown crops, even on some of the steepest slopes. To accomplish this effectively, farmers needed to carve gigantic steps across mountainous contours, forming terraces. These terraces create enchanting ripple like patterns across the landscape and make one almost forget that these farms are meant to help feed local and global communities.



Figure 13.a: Terraced rice fields. Image by PxHere(opens in new window) (Public Domain).

Since the transition to agriculture 10,000 years ago, human communities have struggled against the reality that soil suffers nutrient depletion through constant plowing and harvesting (mostly nitrogen loss). The specter of a significant die-off in human population owing to stagnant crop yields was averted in the 1970s by the **Green Revolution**, which, through the engineering of new crop varieties, large-scale irrigation projects, and the massive application of petroleum-based fertilizers to supplement nitrogen, increased staple crop production with such success that the numbers suffering malnutrition actually declined worldwide in the last two decades of the 20th century, from 1.9 to 1.4 billion, even as the world's population increased at 100 times background rates. The prospects for expanding those gains in the new century are nevertheless threatened by the success of industrial agriculture itself. Soil depletion, declining water resources, and the diminishing returns of fertilizer technology—all the products of a half-century of industrial agriculture—have seen increases in crop yields level off.

Thumbnail image - Women farmers planting a rice field in West Sumatra.

Attribution

Modified by Melissa Ha and Rachel Schleiger from The Industrialization of Nature: A Modern History (1500 to the present)(opens in new window) from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors. Download for free at CNX(opens in new window). (licensed under CC-BY(opens in new window))

13.1: Food Security

13.2: Industrial Agriculture

13.3: Soils

13.4: Soil Degradation

13.5: Selective Breeding and Genetic Engineering
13.6: Sustainable and Regenerative Agriculture

This page titled 13: The Agricultural System is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 14: Agriculture by Melissa Ha and Rachel Schleiger is licensed CC BY-NC 4.0.

13.1: Food Security

Progress continues in the fight against hunger, yet an unacceptably large number of people lack the food they need for an active and healthy life. In 2019, 690 million people in the world (8.9%) were undernourished (lacked sufficient calories; see The State of Food Security...), and 8.2% of the global population lived in poverty (see Goal 1...). Poverty is internationally defined as living on less than $1.90 per day. Poverty—not food availability—is the major driver of food insecurity. Improvements in agricultural productivity are necessary to increase rural household incomes and access to available food but are insufficient to ensure food security. Evidence indicates that poverty reduction and food security do not necessarily move in tandem. The main problem is lack of economic (social and physical) access to food at national and household levels and inadequate nutrition. Food security not only requires an adequate supply of food but also entails availability, access, and utilization by all—people of all ages, gender, ethnicity, religion, and socioeconomic levels.

From Agriculture to Food Security

Agriculture and food security are inextricably linked. The agricultural sector in each country is dependent on the available natural resources, as well as the politics that govern those resources. **Staple food crops** are the main source of dietary energy in the human diet and include things such as rice, wheat, sweet potatoes, maize (corn), and cassava (figure $13.1.a$).



Figure $13.1.a$: Cassava tubers are a staple food crop in tropical regions. Image by Renatosjoao (CC-BY-SA).

Food Security

An individual must have access to food in ample quantity and of sufficient nutritional quality at all times to be **food secure**. Those who never have sufficient quality food are **chronically food insecure**. Food security is determined by availability, access, and utilization.

Food availability refers to whether enough food is produced globally to feed the world's population. In fact, enough food is produced globally, but it is not accessible to everyone who needs it. **Food access** refers to the ability to obtain food in ample quantity and of nutritional quality. In the United States, residents of **food deserts** face limited access to nutritious foods, particularly fresh produce (figure $13.1.b$). This problem is compounded when residents do not own a vehicle or have access to public transportation to obtain foods from surrounding areas or lack the funds to do so.




Figure 13.1.*b*: Food deserts in the contiguous United States are shaded in bright green. Food deserts are distributed over the United States. Examples of large food deserts are in southeastern Oregon and northeastern Arizona. Image by Economic Research Services, USDA (public domain).

Food utilization essentially translates the food available to a household into nutritional security for its members. One aspect of utilization is analyzed in terms of distribution according to need. Nutritional standards exist for the actual nutritional needs, which differ by gender, age, and life phase (for example, pregnancy), but these "needs" are often socially constructed based on culture. For example, in South Asia evidence shows that women eat after everyone else has eaten and are less likely than men in the same household to consume preferred foods such as meats and fish.

Food insecurity can result in **undernourishment** and **malnutrition** (a lack of essential nutrients; figure 13.1.*c*). The economic costs of undernourishment and malnutrition are substantial, potentially costing individuals 10% of their lifetime earnings and costing nations 2-3% of gross domestic product (GDP) in the worst-affected countries (Alderman 2005). This in part due to the reduced ability of under- and malnourished individuals to work. Additionally, without proper nutrition, children can develop disabilities that make it more difficult for them to join to workforce as adults. The economic costs reported here do not even account for the healthcare costs associated with under- and malnutrition.



Figure 13.1.*c*: A goiter is an enlarged thyroid gland that can result from deficiency of iodine, an essential nutrient. Iodized salt, commonly used in the United States, is one strategy to prevent iodine deficiencies. Image by a goiter is an enlarged thyroid glad from iodine deficiency. [Almazi] (CC-BY). Access for free at openstax.org.

Obesity

Obesity is a chronic disease caused by a mix of factors, including genetics, neurobiology, eating habits, access to healthy food, economic influences, and the surrounding environment. In recent decades, it has increased worldwide as many countries have gained more reliable food supplies, improved living conditions, and experienced changes in diet, physical activity, and behavior due to globalization and large-scale food production. Obesity is not just a cosmetic issue—it is a serious health condition that raises the risk of diseases such as type 2 diabetes, heart disease, and certain cancers.

Although obesity has become a significant global health challenge, it is preventable and reversible. Since the 1970s, obesity rates have rapidly increased, resulting in a global obesity epidemic. Thirty-nine percent of the world's adults, or 1.9 billion people, are overweight in 2016. Of these, 650 million are obese (Obesity and Overweight). Obesity was linked to 4.7 million premature deaths in 2017 (about 8% of all deaths that year; OurWorldinData.org [Hannah]). The economic cost of obesity has been estimated at $2 trillion (Tremmel et al.).

Initially centered in developed countries, the obesity epidemic now affects countries of all economic statuses. In low-income countries, this results in a triple burden of undernourishment, malnutrition, and obesity. There is significant variation by region; some have very high rates of undernourishment and low rates of obesity, while in other regions the opposite is true (figure $13.1.d$).




Figure $13.1.d$: Obesity (red bars on the right) and undernourishment (pink bars on the left) by region. The share of the population (%) that is undernourished is approximately 24% in Sub-Saharan Africa while approximately 7% of the population in this region is obese. For South Asia 16% of the population is undernourished, and 4% is obese. These percentages 11% and 5% for East Asia and the Pacific, 5% and 15% for Latin America and the Caribbean, 7% and 17% for Europe and Central Asia, and 9% and 20% for the Middle East and North Africa, respectively.

Obesity accounts for a growing level and share of worldwide noncommunicable diseases, including diabetes, heart disease, and certain cancers that can reduce quality of life and increase public health costs of already under-resourced low-income countries. Driven primarily by increasing availability of processed, affordable, and effectively marketed food, the global food system is falling short with rising obesity and related poor health outcomes. Due to established health implications and rapid increase in prevalence, obesity is now a recognized major global health challenge.

Supplemental Reading

The United Nations explains global food supply and strategies for addressing world hunger here using a variety of informational graphics and videos.

References

Alderman, H. 2005. Linkages between Poverty Reduction Strategies and Child Nutrition: An Asian Perspective. *Economic and Political Weekly, 40*(46), 4837-4842. Accessed 2021-01-03.

Hannah Ritchie. 2017. Obesity. Our World in Data. Accessed 2021-01-03.

Goal 1: End poverty in all its forms everywhere. Sustainable Development. Department of Economic and Social Affairs. United Nations. Accessed 2021-01-03.

Obesity and Overweight. 2020. World Health Organization. Accessed 2021-01-03.

The State of Food Security and Nutrition in the World in 2020. Food and Agriculture Organization of the United Nations. Accessed 2021-01-03.

Tremmel, M., Gerdtham, U. G., Nilsson, P. M., & Saha, S. 2017. Economic Burden of Obesity: A Systematic Literature Review. *International journal of environmental research and public health*, *14*(4), 435. Accessed 2021-01-03.

Attribution

Modified by Melissa Ha from Food Security from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY)

This page titled 13.1: Food Security is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 14.1: Food Security by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

13.2: Industrial Agriculture

Also known as conventional agriculture, **industrial agriculture** is a method of farming that involves use of synthetic fertilizer, synthetic pesticides, and machinery. Industrial agriculture is dependent on large investments in mechanized equipment powered mostly by fossil fuels. In the case of livestock, most production comes from systems where animals are highly concentrated and confined.

Synthetic Pesticides

Pests are organisms that occur where they are not wanted or that cause damage to crops, humans or other animals (figure $13.2.a$). Thus, the term "pest" is a highly subjective term. A **pesticide** is a term for any substance intended for preventing, destroying, repelling, or mitigating any pest. Though often misunderstood to refer only to insecticides, the term pesticide also applies to **herbicides** (weed killers), fungicides, and various other substances used to control pests. Chemical pesticides can be effective, fast acting, adaptable to all crops and situations. When first applied, pesticides can result in impressive production gains of crops. However, despite these initial gains, excessive use of pesticides can be ecologically unsound (see Disadvantages of Industrial Agriculture). By their very nature, most pesticides create some risk of harm—pesticides can cause harm to humans, animals, and/or the environment because they are designed to kill or otherwise adversely affect living things. At the same time, pesticides are useful to society because they can kill potential disease-causing organisms and control insects, weeds, worms, and fungi.



Figure $13.2.a$: The cotton boll weevil is considered a major pest because of the damage it does to cotton plants. Image by ARS USDA (public domain).

Monoculture Farming

Industrial agriculture employs **monoculture** farming, which involves growing only one species of crop in a large area (figure $13.2.b$). The plants in a monoculture are evenly spaced and have the same planting, irrigation, fertilizer, harvesting, etc. requirements, and this uniformity results in efficient use of farming machinery. Often, the crop is a **high yield variety** (high yielding variety; HYV), which is produced through selectively breeding individual plants with desirable characteristics. Compared to traditional crops, HYVs can produce more per unit area of land. Monoculture farming compromises genetic and species diversity in agriculture, however, promoting the spread of pests and risking that all the plants for hundreds of acres could be susceptible to the same disease (see Genetic Diversity).



Figure 13.2.*b*: Monocultures like this potato farm (left) and cotton farm (right) make it easier to use farming equipment; however, this reduces genetic diversity and consumes fossil fuels. Left image by NightThree (CC-BY) and right image by Kimberly Vardeman (CC-BY).

Advantages of Industrial Agriculture

Industrial agriculture has delivered tremendous gains in productivity and efficiency. Food production worldwide has overall risen since the 1940s (figure 13.2.*c*); the World Bank estimates that between 70 percent and 90 percent of the recent increases in food production are the result of industrial agriculture rather than greater acreage under cultivation (figure 13.2.*d*). United States consumers have come to expect abundant and inexpensive food. Additionally, the use of synthetic pesticides ensures that produce is relatively free of blemishes (figure 13.2.*e*).




Figure 13.2.*c*: Corn yields in metric tons (tonnes; 1000 kilograms) per hectare over time. While yields have fluctuated from year to year, they have overall increased since the 1940s. This improving agricultural efficiency is partly due to industrial agricultural practices. Image by OurWorldinData.org [Hannah Ritchie and Max Roser] (CC-BY).




Figure 13.2.*d*: The arable (farmable) land required to produce a fixed amount of a crop has decreased over time due to improved agricultural efficiency. Image by OurWorldinData.org [Hannah Ritchie and Max Roser] (CC-BY).



Figure 13.2.*e*: Many of us carefully inspect produce at the grocery store and select those without blemishes. This visually appealing produce results from synthetic pesticides use. Image by Amanda Mills (public domain).

Industrial agriculture expands where crops can be cultivated. Applying synthetic fertilizers to poor soils can make them fertile. Additionally, irrigation systems promote high agricultural productivity in regions and during seasons that would otherwise only support the most drought-tolerant species. For example, about 25% of the food consumed in the United States is produced in California's Central Valley, which receives little rain in the summer months (figure 13.2.*f*).



Figure 13.2.*f*: The four regions of California's Central Valley from north to south are the Sacramento Valley, Delta and Eastside Streams, San Joaquin Basin, and Tulare Basin. Image by USGS (public domain).

Disadvantages of Industrial Agriculture

Economically, the U.S. agricultural sector includes a history of increasingly large federal expenditures. Also observed is a widening disparity among the income of farmers and the escalating concentration of **agribusiness**—industries involved with manufacture, processing, and distribution of farm products—into fewer and fewer hands. Market competition is limited and farmers have little control over prices of their goods, and they continue to receive a smaller and smaller portion of consumer dollars spent on agricultural products.

Economic pressures have led to a tremendous loss to the number of farms, particularly small farms, and farmers during the past

few decades. More than 155,000 farms were lost from 1987 to 1997. Economically, it is very difficult for potential farmers to enter the business today because of the high cost of doing business. Farming equipment is expensive, and the industrial agricultural system favors large corporate farms, which can more easily invest in this equipment. While this equipment promotes efficiency, it also reduces the number of jobs in agriculture (figure 13.2.*g*).




Figure 13.2.*g*: The number of people working in agriculture in selected countries over time (from 1801-2015). Generally, the number of agricultural workers has declined as farming practices increased in efficiency and as farming machinery completed tasks that were once manual. This is both an advantage because people are free to pursue other careers and a disadvantage because of the reduction in jobs. In Japan, the number of agricultural workers peaked above 16 million around 1900 and again around 1960. In the United States, it peaked to nearly 12 million around 1900. France followed a similar pattern, peaking around 1925 at approximately 9 million workers. There were over 6 million agricultural workers in South Korea in 1950, and this has overall declined (although there was a spike around 1980). In Spain the number of workers was over 5 million in 1950 and has since decreased. In the United Kingdom, the number of agricultural workers peaked around 1960 at about 3.5 million. The other countries shown (Netherlands, Finland, Sweden, and Belgium) have consistently had under 1 million agricultural workers. Image byOurWorldinData [Max Roser] (CC-BY).

The mechanical farming equipment required for industrial agriculture relies on fossil fuels. Burning fossil fuels releases air pollution, including greenhouse gases, which cause climate change. Additionally, the synthetic fertilizers and pesticides used in industrial agriculture are produced from fossil fuels. Furthermore, agriculture areas rarely contain as much biomass (organic matter that comprises living things) as biodiverse, intact ecosystems. Biomass is an important carbon sink because its organic molecules store carbon that could otherwise be in the atmosphere as carbon dioxide or methane, both greenhouse gases. Soils with a substantial organic component are also carbon sinks. Industrial agricultural practices that deplete these soils (see below) contribute to climate change, but alternative practices can promote this benefit of healthy soils (see Sustainable Agriculture).

While synthetic fertilizers provide crops with the necessary nutrients for high productivity, excess nutrients from fertilizers can enter bodies of water through runoff. This can lead to a bloom of algae or photosynthetic bacteria in lakes, rivers, and bays in a process called **eutrophication**. These photosynthetic microorganisms sometimes produce toxins that kill aquatic animals and can even harm the humans who consume these animals. Furthermore, they block aquatic plants from accessing light. Eventual decomposition of algal blooms requires oxygen, which results in **hypoxia** (low levels of dissolved oxygen), further harming aquatic species. Additionally, industrial agriculture pollutes water with pesticides and sediments. Pesticides are not only widespread in surface waters, but pesticides from every chemical class have been detected in groundwater.

Industrial agricultural practices often deplete soil quality. For example, wind and water erosion of exposed topsoil removes particles and nutrients from the soil (which also contributes to eutrophication and sediment pollution). Tilling (mixing the soil) and overgrazing of livestock exacerbates erosion. When livestock remove too much vegetation from an area, plant roots no longer anchor the soil in place, and erosion ensues. Farming equipment compacts the soil. This limits infiltration of water into the soil (decreases soil permeability), makes it more difficult for roots to penetrate the soil, and hinders gas exchange in roots. In extreme cases, soil has eroded and lost its ability to retain water that once arable (farmable) land becomes a desert-like (**desertification**; see Soil Degradation; figure 13.2.*h*).



Figure 13.2.*h*: Overgrazing depleted the soil in the Rukwa Plains of southwestern Tanzania. Image by Lichinga (CC-BY-SA).

Irrigation can lead to **salinization** (increased salinity). While the water itself eventually evaporates, is transpired, or drains from the soil, the minerals dissolved in the water may remain in the soil. Over time, these minerals can accumulate to levels that are toxic for most plants. As discussed in Water Usage, irrigation for agriculture accounts for 69% of global water usage. Withdrawal of sufficient water for irrigation contributes to water shortages and diverts water from ecosystems (see Aqueducts).

Repeated pesticide application exerts selective pressure on insects, fungi, and other crop pests to evolve pesticide resistance. By chance, some individuals in a pest population may carry gene versions that confer pesticide resistance. When these populations are exposed to pesticides, individuals with these gene versions are the most likely to survive and reproduce. They then pass these genes to their offspring, and pesticide resistance becomes more and more common in the population over time (figure 13.2.*i*). At this point, pesticides may need to be applied at greater concentrations or more frequently to achieve the same effect. If pesticide that was initially used is no longer effective at all, the farmer would be forced to find a different pesticide. Over 400 insects and mite pests and more than 70 fungal pathogens have become resistant to one or more pesticides.




Figure 13.2.*i*: Pesticide application can exert selection for the evolution of pesticide resistance. In this hypothetical example, the purple beetle has a gene version that makes it resistant to pesticides (upper left). Pesticides are applied, killing most of the susceptible individuals (white beetles; upper right). In a later generation, purple, resistant beetles are more common because they had a better chance of surviving and reproducing (lower left). Upon another pesticide application, an even greater proportion of the population is resistant. Image modified from Delldot (CC-BY-SA).

The human health and environmental costs of pesticide use have been unevenly distributed. Despite the fact that the lion's share of chemical pesticides are applied in developed countries, 99 percent of all pesticide poisoning cases occur in low-income countries where regulatory, health and education systems are weakest. Farmers in low-income countries may overuse pesticides or not take proper safety precautions because they might not understand the risks and fear smaller harvests (or the instructions to the pesticide are in English, which they may not read). Making matters worse, low-income countries seldom have strong regulatory systems for dangerous chemicals; pesticides banned or restricted in high income countries are used widely in low-income countries. Farmers' perceptions of appropriate pesticide use may also vary by setting and culture. Prolonged exposure to pesticides has been associated with several chronic and acute health effects like non-Hodgkin's lymphoma, leukemia, as well as cardiopulmonary disorders, neurological and hematological symptoms, and skin diseases.

Pesticides have also placed stresses on pollinators and other beneficial insect species. They can lead to the destruction of natural enemies, which would regulate pest populations in an intact ecosystem. For example, once apple growers began controlling pests with the pesticide dichlorodiphenyltrichloroethane (DDT), which has since been banned, they quickly found their orchards being attacked by scale insects and mites. The reason: DDT had killed off their natural enemies. More generally, pollution from industrial agriculture along with habitat loss due to converting healthy ecosystems into agricultural fields is a major threat to biodiversity. As mentioned previously, monoculture farming limits biodiversity in crops.

With respect to animal agriculture, keeping livestock are kept in close quarters increases the risk of disease spread among them. For these reason, food animals are commonly given antibiotics, either preventatively or to treat existing infections. As discussed in the Infectious Diseases section, the overuse of antibiotics (whether in humans or farm animals) increases selection for the evolution of antibiotic-resistant strains of bacteria or other disease-causing organisms. Infections by antibiotic-resistant strains are more difficult to treat.

As with many industrial practices, potential health hazards are often tied to farming practices. Under research and investigation currently is the sub-therapeutic use of antibiotics in animal production, and pesticide and nitrate contamination of water and food. Farm worker health is also a consideration in all farming practices.

Attributions

Modified by Melissa Ha from the following sources:

- Conventional Agriculture and Pests and Pesticides from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Insecticides from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Sustainable Agriculture: Definitions and Terms by Mary V. Gold, Alternative Farming Systems Information Center from National Agricultural Library, USDA (public domain)

This page titled 13.2: Industrial Agriculture is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 14.2: Industrial Agriculture by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

13.3: Soils

Soil is the outer loose layer that covers the surface of Earth. Soil quality is a major determinant, along with climate, of plant distribution and growth. Soil quality depends not only on the chemical composition of the soil, but also climate, topography, and organisms living in the soil. In agriculture, the history of the soil, such as the cultivating practices and previous crops, modify the characteristics and fertility of that soil.

Importance of Soil

Soil is important to our society primarily because it provides the foundation of agriculture and forestry. Soil plays a key role in plant growth. Beneficial aspects to plants include providing physical support, water, heat, nutrients, and oxygen. Mineral nutrients from the soil can dissolve in water and then become available to plants. Through their roots, plants absorb water and minerals (e.g., nitrates, phosphates, potassium, copper, zinc). With these and carbon dioxide acquired during photosynthesis, plants produce carbohydrates, proteins, lipids, nucleic acids, and the vitamins, on which consumers depend.

Soil plays a role in nearly all biogeochemical cycles on the Earth's surface. Global cycling of key elements such as carbon (C), nitrogen (N), phosphorous (P), and sulfur (S) all pass through soil. In the hydrologic (water) cycle, soil helps to mediate **infiltration** (percolating) from the surface into the groundwater. Microorganisms living in soil can also be important components of biogeochemical cycles through the action of decomposition and other processes such as nitrogen fixation.

Several elements are considered essential for plant growth. Carbon (C), hydrogen (H), and oxygen (O) are required in large quantities, but they are not absorbed as mineral nutrients from the soil. Plants obtain carbon from carbon dioxide in the atmosphere and hydrogen from the water absorbed by the roots. Oxygen atoms come from carbon dioxide acquired during and gaseous oxygen in the atmosphere (acquire through aerobic cellular respiration) as well as water. Of the mineral nutrients absorbed from the soil, **macronutrients**, including nitrogen (N), potassium (K), calcium (Ca), magnesium (Mg), phosphorus (P), sulfur (S), and silicon (Si) are needed by plants in significant quantities. **Micronutrients** are essential elements that are needed only in small quantities, but can still be limiting to plant growth since these nutrients are not so abundant in nature. Micronutrients include chlorine (Cl), iron (Fe), boron (B), manganese (Mn), sodium (Na), zinc (Zn), copper (Cu), nickel (Ni), and molybdenum (Mo). There are some other elements that tend to aid plant growth but are not absolutely essential.

Although many aspects of soil are beneficial to plants, excessively high levels of trace metals (either naturally occurring or added by humans) or applied herbicides can be toxic to some plants (figure $13.3.a$).



Figure $13.3.a$: These native plants are affected by herbicides that were used to kill nearby weeds. Image by Matt Lavin (CC-BY-SA).

Micronutrients and macronutrients are desirable in particular concentrations and can be detrimental to plant growth when

concentrations in soil solution are either too low (limiting) or too high (toxicity). Mineral nutrients are useful to plants only if they are in an extractable form in soil solutions, such as a dissolved ion rather than in solid mineral. Many nutrients move through the soil and into the root system as a result of concentration gradients, moving by diffusion from high to low concentrations. However, some nutrients are selectively absorbed by the root membranes, enabling concentrations to become higher inside the plant than in the soil.

An important factor affecting soil fertility is soil **pH** (the negative log of the hydrogen ion concentration). Soil pH is a measure of the acidity or alkalinity of the soil solution. On the pH scale (0 to 14) a value of seven represents a neutral solution; a value less than seven represents an **acidic** solution and a value greater than seven represents an **alkaline** solution (figure $13.3.b$). Soil pH affects the health of microorganisms in the soil and controls the availability of nutrients in the soil solution. Strongly acidic soils (less than 5.5) hinder the growth of bacteria that decompose organic matter in the soil. This results in a buildup of organic matter that has yet to be decomposed, which leaves important nutrients such as nitrogen in forms that are unusable by plants. Soil pH also affects the solubility of nutrient-bearing minerals. This is important because the nutrients must be dissolved in solution for plants to assimilate them through their roots. Most minerals are more soluble in slightly acidic soils than in neutral or slightly alkaline soils. Strongly acid soils (pH four to five), though, can result in high concentrations of aluminum, iron and manganese in the soil solution, which may inhibit the growth of some plants.




Figure $13.3.b$: The pH scale measures the amount of hydrogen ions (H^+) in a substance. This shows the pH of common household solutions. Those with a pH greater than 7 are basic (alkaline), and those with a pH less than 7 are acidic. The solutions from high to low pH are bleach (13), soapy water (12), ammonia solution (11), milk of magnesia (10), baking soda (9), seawater (8), distilled water (7), urine (6), black coffee (5), tomato juice (4), orange juice (3), lemon juice (2), and gastric acid (1). Note that pH values are approximate. Image and caption (modified) from Edward Stevens [OpenStax](CC-BY). Access for free at openstax.org.

Several factors determine soil pH. Organic material in soil decreases pH to an extent, but it also acts as a **buffer**, limiting changes in pH. Climate is also important, with high amounts of rainfall increasing leaching and lowering pH. Some types of parent material, such as those high in silicon, decrease pH, while others, such as limestone increase pH.

Soil Composition

Soil consists of organic matter (about 5%), inorganic mineral matter (40-45% of soil volume), water (about 25%) and air (about 25%). The amount of each of the four major components of soil depends on the amount of vegetation, soil compaction, and water present in the soil.

The organic material consists of dead organisms in various stages of decomposition. It is dark-colored because it contains **humus**, partially decayed matter containing organic acids. Humus enriches the soil with nutrients, gives the soil a loose texture that holds water, and allows air to diffuse through it. Oxygen is important for plant roots and many inhabitants of the soil. The organic component of soil serves as a cementing agent, returns nutrients to the plant, allows soil to store moisture, makes soil tillable for farming, and provides energy for soil microorganisms. Most soil microorganisms—bacteria, algae, or fungi—are dormant in dry soil, but become active once moisture is available.

The inorganic material of soil consists of rock, slowly broken down into smaller particles that vary in size. Soil particles that are 100 µm to 2 mm in diameter are **sand**. (A **micrometer**, µm, 10^{-6} m, or a millionth of a meter.) Soil particles between 2 and 100 µm are called **silt**, and even smaller particles, less than 2 µm in diameter, are called **clay**.

Soil should ideally contain 50 percent solid material and 50 percent pore space (figure 13.3.*c*). **Pore space** refers to the gaps in between soil particles. The larger the soil particles, the larger the pore spaces. Water can quickly pass through large pore spaces, so soils high in sand drain easily. Smaller soil particles have more surface area relative to volume and produce narrow pore spaces. Water clings to these surfaces, and soils high in clay thus retain water. (Clay is also negatively charged, which attracts water.) About one-half of the pore space should contain water, and the other half should contain air.




Figure 13.3.*c*: Soil particles and the pore spaces in between them. The black shading represents water, which fills the pore spaces in the lower portion of the soil and clings to narrow portions of pore space in the top portion of the soil. Image from "Forest physiography; physiography of the United States and principles of soils in relation to forestry" (public domain).

Soil texture is based on percentages of sand, silt, and clay (figure 13.3.*d*). Soils that have a high percentage of one particle size are named after that particle (a clay soil has a high percentage of clay). Other soils have a mixture of two particle sizes and very little of the third size. For example, silty clay has roughly 50% clay and 50% silt while sandy clay has 50-60% sand and 35-50% clay. Some soils have no dominant particle size and contain a mixture of sand, silt, and humus. These soils are called **loams**, and they are optimal for agriculture. A medium loam has roughly 40% sand, 40% silt, and 20% clay. Larger particles (sand) facilitate drainage, and small particles (clay) facilitate water retention, so loam soils both have good drainage and can remain moist. Soils that deviate slightly from a medium loam include loamy sand, sandy loam, sandy clay loam, clay loam, silty clay loam, and silty loam.




Figure 13.3.*d*: A soil texture triangle is used to determine the soil texture based on the percentages of sand, silt, and clay. The left axis represents the percentage of clay. The right axis represents the percentage of silt. The bottom axis represents the percentage of sand. Most soils that contain more than 40% are classified as clays. Exceptions are silty clays, which contain 40-60% clay and 40-60% silt. Another exception are sandy clays, which contain 35-50% clay and 50-70% sand. Clay loams and silty clay loams contain 25-40% clay. Clay loams contain 20-60% silt, and silty clay loams contain 60-75% silt. Sandy clay loams contain 20-35% clay and 50-80% sand. Medium loams contain 5-25% clay, 20-50% silt, and 30-55% sand. Soils with more than 50% silt and no more than 25% clay are either silty loams or silts. Silts have 80-100% silt and no more than 15% clay. Sand have 90-100% sand. Loamy sands have 75-90% sand and no more than 15% clay. A soil with a composition that does not fit into the already described categories would be a sandy loam. To determine the soil texture, first find the percentage of clay and trace a line horizontally to the right. At the same time, find the percentage of silt and trace a line diagonally (down and to the left, parallel to the clay axis). The two lines will meet at the correct soil type. For example, traced lines from 20% clay and 40% silt intersect at the medium loam category. To confirm, trace a diagonal line (down and to the right, parallel to the sit axis) from this point to the sand axis. This line cross the axis at 40%. In summary, a line can be traced from each axis that is parallel to axis that is counterclockwise of it. Image by Richard Wheeler [Richard Wheeler] (CC-BY-SA).

Organic Versus Mineral Soils

Soils can be divided into two groups based on how they form. **Organic soils** are those that are formed from sedimentation and often contain more than 30% organic matter. They form when organic matter, such as leaf litter, is deposited more quickly than it can be decomposed (figure 13.3.*e*). **Mineral soils** are formed from the weathering of rocks, typically contain no more than 30% organic matter, and are primarily composed of inorganic material. **Weathering** occurs when biological, physical, and chemical processes, such as erosion, leaching, or high temperatures, break down rocks.



Figure $13.3.e$): Decomposition of organic material occurs very slowly in this acidic bog. The organic material accumulates, which is characteristic of organic soils. Image by William L. Farr (CC-BY-SA).

Soil Horizons

Soil distribution is not uniform because its formation results in the production of layers; together, the vertical section of a soil is called the **soil profile**. Within the soil profile, soil scientists define zones called horizons. A **horizon** is a soil layer with distinct physical and chemical properties that differ from those of other layers.

The soil profile has four distinct layers: 1) O horizon; 2) A horizon; 3) B horizon and 4) C horizon (figure $13.3.f$-g). Upper horizons (labeled as the A and O horizons) are richer in organic material and so are important in plant growth, while deeper layers (such as the B and C horizons) retain more of the original features of the bedrock below. Some soils may have additional layers (like the E horizon, figure $13.3.f$), or lack one of these layers. The thickness of the layers is also variable, and depends on the factors that influence soil formation. In general, immature soils may have O, A, and C horizons, whereas mature soils may display all of these, plus additional layers.




Figure $13.3.f$): Soil Profile. Photograph shows a soil profile from South Dakota revealing three horizons. The A horizon (topsoil) is dark brown and extend about 1 foot. Below it is the white E (eluviated) horizon, which is about 6 inches thick. The B horizon (subsoil, labeled Bt) is the medium brown and the lowest visible layer The yellow arrows symbolize translocation of fine clays to the B horizon. The scale is in feet. Source: University of Idaho and modified by D. Grimley.




Figure $13.3.g$): This image shows the different horizons, or layers, in soil. Vegetation grows at the surface. The top, deep brown layer is the O (organic) horizon. Below this is the gray A horizon (topsoil, or surface). Next is the light brown B horizon (subsoil). The C horizon (substratum, or soil base) is tan and includes larger pieces of rocks. The gray bedrock is the deepest layer. The E (eluviated) horizon that is sometimes found between the A and B horizons is absent here. This work(opens in new window) by Wilsonbiggs is licensed under CC BY-SA 4.0

O horizon

The very top of the **O horizon** consists of partially decayed organic debris like leaves. This horizon is usually dark in color because of humus.

A horizon

The **A horizon** (topsoil) consists of a mixture of organic material with inorganic products of weathering, and it is therefore the beginning of true mineral soil. In this area, rainwater percolates through the soil and carries materials from the surface. The A horizon may be only 5 cm (2 in.), or it may over a meter. For instance, river deltas like the Mississippi River delta have deep layers of topsoil. Microbial processes occur in the top soil, and this horizon supports plant growth. Many organisms, such as earthworms and insects live among the plant roots in this horizon.

B horizon

The **B horizon** (subsoil) consists of small particles that have moved downward, resulting in a dense layer in the soil. In some soils, the B horizon contains nodules or a layer of calcium carbonate. The subsoil is usually lighter in color than topsoil and often contains an accumulation of minerals.

C horizon

The **C horizon** (soil base), includes the **parent material**, the organic and inorganic substances from which soils form. Weathering parent material represents the first steps in the chemical breakdown of rock into soil. Often the weathered parent material is underlain by the parent material itself, although in some places it has been carried from another location by wind, water, or glaciers. Beneath the C horizon lies bedrock. The chemical nature of the parent material, whether granite, limestone, or sandstone, for example, has a great influence on the fertility of the soil derived from it.

Factors Affecting Soil Formation and Composition

The fundamental factors that affect soil genesis can be categorized into five elements: climate, organisms, topography, parent material, and time. One could say that the relief, climate, and organisms dictate the local soil environment and act together to cause weathering and mixing of the soil parent material over time.

Climate

The role of climate in soil development includes aspects of temperature and precipitation. Soils in very cold areas with permafrost conditions (such as the arctic tundra) tend to be shallow and weakly developed due to the short growing season. In warm, tropical climates, soils tend to be thicker (but lacking in organic matter), with extensive leaching and mineral alteration. In such climates, organic matter decomposition and chemical weathering occur at an accelerated rate. The presence of moisture and nutrients from weathering will also promote biological activity: a key component of a quality soil. See the Biomes chapter for more details about the effect of climate on soils.

Ancient soils, sometimes buried and preserved in the subsurface, are referred to as **paleosols** (figure $13.3.h$) and reflect past climatic and environmental conditions.




Figure $13.3.h$: Modern versus Buried Soil Profiles. A buried soil profile, or paleosol (above geologist 's head), represents soil development during the last interglacial period. A modern soil profile (Alfisol) occurs near the land surface. Source: D. Grimley.

Organisms

The presence of living organisms in the soil (**soil biota**) greatly affects soil formation and structure. A diversity of animals are found in the soil such as nematodes, spiders, insects, centipedes, millipedes, pillbugs, slugs, and earthworms (figure $13.3.i$). The soil also contains microorganisms like bacteria, archaea, fungi, and "protists". Animals and microorganisms can produce pores and crevices, and plant roots can penetrate into crevices to produce more fragmentation. Additionally, leaves and other material that fall from plants decompose and contribute to soil composition. Microorganisms not only decompose organic matter, but contribute to other processes in nutrient cycles, such as nitrogen fixation.




Figure 13.3.*i*: Illustration of animals commonly found in the soil. Image by Arne Hendriks (CC-BY)

Parent Material

Mineral soils form directly from the weathering of bedrock, the solid rock that lies beneath the soil, and therefore, they have a similar composition to the original rock. Other soils form in materials that came from elsewhere, such as sand and glacial drift. Materials located in the depth of the soil are relatively unchanged compared with the deposited material. Sediments in rivers may have different characteristics, depending on whether the stream moves quickly or slowly. A fast-moving river could have sediments of rocks and sand, whereas a slow-moving river could have fine-textured material, such as clay.

The type of parent material may also affect the rapidity of soil development. Parent materials that are highly weatherable (such as volcanic ash) will transform more quickly into highly developed soils, whereas parent materials that are quartz-rich, for example, will take longer to develop. Parent materials also provide nutrients to plants and can affect soil internal drainage.

Topography

Regional surface features (familiarly called “the lay of the land”) can have a major influence on the characteristics and fertility of a soil. Topography affects water runoff, which strips away parent material and affects plant growth. Soils on steep slopes are more prone to erosion and may be thinner than soils that are on relatively level ground. **Infiltration**, the percolating of water through the soil, is limited in steep soils.

The local topography can have important microclimatic effects. In the northern hemisphere, south-facing slopes are exposed to more direct sunlight angles and are thus warmer and drier than north-facing slopes. The cooler, moister north-facing slopes have a more dynamic plant community and thicker soils because extensive root systems stabilize the soil and reduce erosion (figure 13.3.*j*).



Figure 13.3.j: A north-facing slope (left) and south-facing slope (right) in the Mediterranean climate (chaparral) of the Santa Monica Mountains, California. Vegetation on the north-facing slope is denser due to cool, moist conditions relative to the south-facing slope. Image by Noah Elhardt (CC-BY).

Time

Time is an important factor in soil formation because soils develop over long periods. Soil formation is a dynamic process. Materials are deposited over time, decompose, and transform into other materials that can be used by living organisms or deposited onto the surface of the soil.

In general, soil profiles tend to become thicker (deeper), more developed, and more altered over time. However, the rate of change is greater for soils in youthful stages of development. The degree of soil alteration and deepening slows with time and at some point, after tens or hundreds of thousands of years, may approach an equilibrium condition where erosion and deepening (removals and additions) become balanced. **Young soils** (< 10,000 years old) are strongly influenced by parent material and typically develop horizons and character rapidly. Over time, as weathering processes deepen, mix, and alter the soil, the parent material becomes less recognizable as chemical, physical, and biological processes take effect. **Moderate age soils** (roughly 10,000 to 500,000 years old) are slowing in profile development and deepening, and may begin to approach equilibrium conditions. **Old soils** (>500,000 years old) have generally reached their limit as far as soil horizonation and physical structure, but may continue to alter chemically or mineralogically.

Soil development is not always continual. Geologic events such as landslides, glacier advance, or the rising of shorelines can rapidly bury soils. Erosion in rivers and shorelines can cause removal or truncation of soils, and wind or flooding slowly deposit sediment that adds to the soil. Animals can mix the soil and sometimes cause **soil regression**, a reversal or "bump in the road" for the normal path of development, and this increases development over time.

Soil Taxonomy

Soils are classified into one of 12 soil orders based on soil horizons, how they form, and their chemical compositions. For example, Mollisols (figure 13.3.f), which are found in temperate grasslands, have a thick topsoil rich in organic content. Aridisols, on the other hand, are dry soils that contain calcium carbonate and are found in deserts. Each soil order is further divided into suborders. See USDA's The Twelve Orders of Soil Taxonomy and The Twelve Soil Orders from the University of Idaho for more details.

Attributions

Modified by Melissa Ha from the following sources:

- The Soil from *General Biology* by OpenStax (CC-BY(opens in new window))
- Soil Profiles and Processes and Soil-Plant Interactions from *Environmental Biology* by Matthew R. Fisher (licensed under CC-

BY(opens in new window))
- Soil from *Biology* by John W. Kimball (licensed under CC-BY)

This page titled 13.3: Soils is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 7.4: Soils by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

13.4: Soil Degradation

Once fertile topsoil is lost, it is not easily replaced. **Soil degradation** refers to deterioration in the quality of the soil and the resultant reduction in its capacity to produce. Soils are degraded primarily by erosion, compaction, and salinization. Such processes often arise from poor soil management during agricultural activities. In extreme cases, soil degradation can lead to **desertification** (conversion of land to desert-like conditions) of croplands and rangelands in semi-arid regions. See the Sustainable Agriculture section for strategies to preserve soil quality (**soil conservation**).

Erosion

Erosion is the biggest cause of soil degradation. Soil productivity is reduced as a result of losses of nutrients, water holding capacity, and organic matter. **Water holding capacity** is a measure of of the soils ability to retain water. The two agents of erosion are wind and water, which act to remove the finer particles from the soil. **Wind erosion** occurs mostly in flat, dry areas and moist, sandy areas along bodies of water. Wind not only removes soil, but also dries and degrades the soil structure. **Water erosion** is the most prevalent type of erosion. It occurs when raindrops splash on the ground and when water moves down slope as a thin film, small streams, or a large stream.

Some amount of soil erosion is a natural process along sloping areas and/or in areas with soft or materials that do not stick together and are susceptible to movement by water, wind, or gravity. For instance, soil material can be mobilized in strong windstorms, along the banks of rivers, in landslides, or by wave action along coastlines. However, human activities such as construction, logging, and off-road vehicle use promote erosion by removing the natural vegetation cover protecting the soil. Agricultural practices such as overgrazing and leaving plowed fields bare for extended periods contribute to farmland erosion. Each year, an estimated two billion metric tons of soil are eroded from farmlands in the United States alone. The soil transported by the erosion processes can also create problems elsewhere (for example, by clogging waterways and filling ditches and low-lying land areas). The areas most vulnerable to soil erosion include locations with thin organic (A and O) horizons and hilly terrains (figure $13.4.a$).




Figure $13.4.a$: Global map of water erosion vulnerability. There are four levels of vulnerability: very high (red), high (orange), moderate (yellow), and low (light green). Examples of regions with very high vulnerability to erosion by water include the west coast of South America, parts of the eastern United States, regions around the Mediterranean Sea, a strip across western and central Africa, Korean and Japan, Northern Australia, and New Zealand. Examples of regions with low vulnerability include northern South America, the Democratic Republic of Congo, northern India and Nepal, and western Russia. Other regions are labeled as dry (gray), cold (light blue), depositional (bright green), or ice/glacier (white). For example, northern regions are cold and ice/glacier. Much of the western United States, Chile and Argentina, northern Africa, the Middle East, Kazakhstan, Mongolia, and much of Australia are considered dry. The largest depositional area (region where sediments accumulate) is northeastern Europe. Image by USDA-NRCS, Soil Science Division, World Soil Resources (public domain).

Compaction

In modern agricultural practices, heavy machinery is used to prepare the seedbed, for planting, to control weeds, and to harvest the crop. The use of heavy equipment has many advantages in saving time and labor, but can cause compaction of soil and disruption of the natural soil biota. Much compaction is reversible and some is unavoidable with modern practices; however, serious compaction issues can occur when the equipment is used excessively during times when the soil has a high water content. The problem with soil compaction is that increased soil density limits root penetration depth and may inhibit proper plant growth.

Salinization

When considerable quantities of salt accumulate in the soil in a process known as **salinization**, many plants are unable to grow properly or even survive. This is especially a problem in irrigated farmland. Groundwater used for irrigation contains small amounts of dissolved salts. Irrigation water that is not absorbed into the soil evaporates, leaving the salts behind. This process repeats itself and eventually severe salinization of the soil occurs. A related problem is water logging of the soil. When cropland is irrigated with excessive amounts of water in order to leach salts that have accumulated in the soil, the excess water is sometimes unable to drain away properly. In this case it accumulates underground and causes a rise in the subsurface water table. If the saline water rises to the level of the plant roots, plant growth is inhibited.

Desertification

Land that was previously suited for growing crops may be turned into desert by climate change and the activities of humans, such as poor farming practices, livestock overgrazing, and overuse of available water. This process, called **desertification**, is a serious problem worldwide. Plants and soil types that are non-arid (not dry) specifically help water soak into the ground (infiltration) and water retention. When desertification begins, it leads to a reduction in vegetation and degraded soil quality, and this further increases the aridity and spreads the desert via a **positive feedback loop** (meaning that the processes feed on themselves promoting an increasing spiral).

Figure $13.4.b$ shows areas of the world and their vulnerability to desertification. Note the red and orange areas in the western and midwestern United States. The Dust Bowl of the 1930s is a classic example of human-caused desertification (figure $13.4.c$). Poor cultivation and grazing practices -- coupled with severe drought conditions -- led to severe wind erosion of soil in a region of the Great Plains that became known as the "Dust Bowl". Wind stripped large areas of farmlands of topsoil, and formed clouds of dust that traveled as far as the eastern United States.

Sometimes there is a conflict between what is known to prevent desertification and what an individual farmer feels is needed to do to make a living. Mitigating the desertification process includes both societal steps and individual education on alternatives.




Figure 13.4.*b*: World map showing desertification vulnerability. There are four levels of vulnerability: very high (red), high (orange), moderate (yellow), and low (light green). Examples of regions with a very high vulnerability to desertification include the northwestern United States, a strip through western and central Africa, parts of the middle East, and parts of Australia. Examples of regions with low vulnerability include parts of Canada, parts of western Europe, and parts of Angola and the Democratic Republic of Congo. Other regions are labeled as dry (gray), cold (light blue), humid/not vulnerable (bright green), or ice/glacier (white). For example, northern regions are cold and ice/glacier. Some of southwestern United States, Chile and Argentina, northern Africa, the Middle East, Kazakhstan, Mongolia, and much of Australia are considered dry. Examples of humid/not vulnerable areas include the eastern United States, northern and eastern South America, western Europe, Central Africa, and East Asia. Image by USDA (public domain).



Figure 13.4.*c*: A cloud of dust in Kansas in 1935, part of the Dust Bowl. Image by NOAA Photo Library (public domain).

Attribution

Modified by Melissa Ha from the following sources:

- The Great Basin and the Basin and the Range(opens in new window) from *An Introduction to Geology* by Chris Johnson et al. (licensed under CC BY-NC-SA(opens in new window))
- Soils from *AP Environmental Science* by University of California College Prep, University of California (licensed under CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Soil and Sustainability from *Sustainability: A Comprehensive Foundation* by Tom Theis and Jonathan Tomkin, Editors (licensed under CC-BY). Download for free at CNX(opens in new window).

This page titled 13.4: Soil Degradation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 7.5: Soil Degradation by Melissa Ha and Rachel Schleiger is licensed CC BY-NC-SA 4.0.

13.5: Selective Breeding and Genetic Engineering

The development of a new crop variety is an example of **agricultural biotechnology**, a range of tools that include both traditional breeding techniques and more modern lab-based methods. Traditional methods date back thousands of years, whereas biotechnology uses the tools of genetic engineering developed over the last few decades.

Selective Breeding (Artificial Selection)

Nearly all the fruits and vegetables found in your local market would not occur naturally. In fact, they exist only because of human intervention that began thousands of years ago. Humans created the vast majority of crop species by using traditional breeding practices on naturally-occurring, wild plants. These practices rely upon **selective breeding** (artificial selection), human-facilitated reproduction of individuals with desirable traits. For example, high yield varieties were produced through selective breeding. Traditional breeding practices, although low-tech and simple to perform, have the practical outcome of modifying an organism's genetic information, thus producing new traits.

Selective breeding is limited, however, by the life cycle of the plant and the genetic variants that are naturally present. For example, even the fastest flowering corn variety has a generation time of 60 days (the time required for a seed to germinate, produce a mature plant, get pollinated, and ultimately produce more seeds) in perfect conditions. Each generation provides an opportunity to selectively breed individual plants and generate seeds that are slightly closer to the desired outcome (for example, producing bigger, juicier kernels). Furthermore, if no individuals happen to possess gene variants that result in bigger, juicier kernels, it is not possible to artificially select this trait. Finally, traditional breeding shuffles all of the genes between the two individuals being bred, which can number into the tens of thousands (maize, for example, has 32,000 genes). When mixing such a large number of genes, the results can be unpredictable.

An interesting example is maize (corn). Biologists have discovered that maize was developed from a wild plant called teosinte. Through traditional breeding practices, humans living thousands of years ago in what is now Southern Mexico began selecting for desirable traits until they were able to transform the plant into what is now known as maize (figure $13.5.a$). In doing so, they permanently (and unknowingly) altered its genetic instructions.




Figure $13.5.a$: A wild grass called teosinte was genetically modified through selective breeding to produce what is now known as maize (corn). This process of transformation started thousands of years ago by indigenous people of what is now Mexico. Image by Nicolle Rager FullerNational Science Foundation [] (public domain).

This history of genetic modification is common to nearly all crop species. For example, cabbage, broccoli, Brussel sprouts, cauliflower, and kale were all developed from a single species of wild mustard plant (figure $13.5.b$). Wild nightshade was the source of tomatoes, eggplant, tobacco, and potatoes, the latter developed by humans 7,000 – 10,000 years ago in South America.



Figure 13.5.*b*: *Brassica oleracea* is a plant in the mustard family and is known as wild cabbage. From it were developed many familiar crops, such as cauliflower, broccoli, Brussel sprouts, and of course, cabbage. Image by Kulac (CC-BY-SA).

Genetic Engineering

Genetic engineering is the process of directly altering an organism's DNA to produce the desired crops more rapidly than selective breeding. Because genes can be obtained from other species or even synthesized in the lab, scientists are not limited by existing genetic variation within a crop species (or closely related species with which they can be crossed). This broadens the possible traits that can be added to crops. Modern genetic engineering is more precise than selective breeding in the sense that biologists can modify just a single gene. Also, genetic engineering can introduce a gene between two distantly-related species, such as inserting a bacterial gene into a plant (figure 13.5.*c*).




Figure $13.5.c$: Both selective (traditional) breeding and modern genetic engineering produce genetic modifications. Genetic engineering allows for fewer and more precise genetic modifications. The traditional plant breeding process introduces a number of genes into the plant. These genes may include the gene responsible for the desired characteristic, as well as genes responsible for unwanted characteristics. In traditional breeding, the donor variety DNA strand (A) and recipient variety DNA strand (C) are combined to produce the new variety DNA strand. The donor DNA strand contains a portion of an organism's entire genome, including the desired gene (B). In the new variety DNA strand, many genes are transferred with the desired gene. Genetic engineering enables the introduction into the plant of the specific gene or genes responsible for the characteristics of interest. By narrowing the introduction to one of a few identified genes, scientists can introduce the desired characteristic without also introducing genes responsible for unwanted characteristics. In genetic engineering, the desired gene (B) is copied from the donor organism's genome (A). The desired gene from the donor organism DNA strand and the recipient variety DNA strand (C) are combined to produce the new variety DNA strand. Only the desired gene is transferred to a location in the recipient genome. Modified from Michael J. ErmarthFDA [] (public domain).

Genetically modified organisms (**GMOs**) are those that have had their DNA altered through genetic engineering. Genetically modified crops are sometimes called genetically engineered (GE) crops. **Transgenic organisms** are a type of genetically modified organism that contains genes from a different species. Because they contain unique combinations of genes and are not restricted to the laboratory, transgenic plants and other GMOs are closely monitored by government agencies to ensure that they are fit for human consumption and do not endanger other plant and animal life. Because these foreign genes (transgenes) can spread to other species in the environment, particularly in the pollen and seeds of plants, extensive testing is required to ensure ecological stability.

How to Genetically Modify Plant Cells

DNA can be inserted into plant cells through various techniques. For example, a **gene gun** propels DNA bound to gold particles into plant cells. (DNA is negatively charge and clings to positively charged gold.) A more traditional approach employs the plant pathogen ***Agrobacterium tumefaciens*** (figure $13.5.d$). Ordinarily, this bacterium causes crown gall disease in plants by inserting a circular piece of DNA, called the **Ti plasmid**, into plant cells. This DNA incorporates into plant chromosomes, giving them genes to produce the gall (figure $13.5.e$), which provides a home for the bacterial pathogen.




Figure 13.5.*d*: An *Agrobacterium tumefaciens* cell (A) inserts the Ti plasmid (C), which contains T-DNA (a) into a plant cell (D). The T-DNA eventually enters the plant nucleus (G), where the plant DNA is stored. Image by Chandres (CC-BY-SA).



Figure 13.5.*e*: A crown gall caused by *Agrobacterium tumefaciens*. This bacterium inserts DNA into plant cells, which causes them to produce the crown gall. This process can be adjusted by scientists to insert genes that code for desirable traits into crop species. Image by Scott Nelson (public domain).

Scientists alters the process by which *Agrobacterium* infects and genetically alter plant cells to produce genetically modified plants with agriculturally beneficial traits as follows (figure 13.5.*f*):

1. T-DNA, which codes for the crown gall is removed from the Ti plasmid, and genes for desired traits are added.
2. The modified plasmid is then added back to *Agrobacterium*.
3. *Agrobacterium* infects undifferentiated plant cells (stem cells that can develop into any part of the plant; figure 13.5.*g*).
4. The modified plant cells are given hormones to produce the entire plant.




Figure 13.5.*f*: The process of producing a transgenic plant using *Agrobacterium*. The Ti plasmid was engineered to include the gene of interest and marker gene, which will later indicate which cells are transgenic. A rectangle represents *Agrobacterium*. It contains regular bacterial DNA and the Ti plasmid, represented by a circle. The T-DNA (which provides instructors for producing a crown gall) is removed, and a transgene (gene of interest) and maker gene are inserted instead. The gene of interest is colored green, and the marker gene (for resistance to the antibiotic kanamycin) is colored blue. Next, plant cells are infected with *Agrobacterium*. The bacterial cell inserts the Ti plasmid into a polyhedral plant cell. The transgene is transferred to the plant cell nucleus, which contains regular plant DNA. Next, the plant cells with the gene of interest are selected and allowed them to divide. This is done by culturing the plant cells on kanamycin media in a petri dish. Calluses, clusters of undifferentiated plant cells, are produced. Finally, hormones are applied to induce shoot and root growth. An arrow points from the petri dish to a potted transgenic plant. Image by Melissa Ha (CC-BY).




Figure 13.5.*g*: Individual plant cells are first genetically engineered with *Agrobacterium*. They then grow into **calluses** (blobs of undifferentiated plant cells) and are given hormones to induce root and shoot development. Because all the cells in the plant that eventually grows descended from a single genetically engineered cell, the entire plant is transgenic. Image by Igge (CC-BY-SA).

Examples of Genetically Modified Crops

Many genetically modified crops have been approved in the U.S. and produce our foods. The first genetically modified organism approved by the U.S. Food and Drug Administration (FDA) in 1994 was **Flavr Savr™ tomatoes**, which have a longer shelf life (delayed rotting) because a gene responsible for breaking down cells in inhibited. Flavr Savr tomatoes are genetically modified (because their DNA has been altered) but not trasgenic (because they do not contain genes from another species). The Flavr Savr tomato did not successfully stay in the market because of problems maintaining and shipping the crop. **Golden rice** produces β-

carotene, a precursor to vitamin A (figure 13.5.h; β-carotene is also in high concentrations in carrots, sweet potatoes, and cantaloupe, giving them their orange color.) **Roundup Ready®** corn, cotton, and soybeans are resistant to this common herbicide, making it easier to uniformly spray it in a field to kill the weeds without harming the crops (figure 13.5.i).



Figure 13.5.h: Golden rice has an orange-yellow color because it contains up to 35 µg β-carotene (a precursor to vitamin A) per gram of rice, which could prevent millions of cases of blindness worldwide. Image by International Rice Research Institute (CC-BY).



Figure 13.5.i: Roundup is a common herbicide. Roundup Ready® plants are genetically modified to resist roundup, meaning the herbicide does not kill them. This allows farmers to uniformly spray Roundup, killing weeds without harming the crops. Image by Mike Mozart (CC-BY).

Crops have also been engineered to produce insecticides. *Bacillus thuringiensis* (Bt) is a bacterium that produces protein crystals that are toxic to many insect species that feed on plants. Insects that have eaten Bt toxin stop feeding on the plants within a few hours. After the toxin is activated in the intestines of the insects, death occurs within a couple of days. The gene to produce **Bt toxin** has been added to many crops including corn (figure 13.5.j), potatoes, and cotton, providing plants with defense against insects.




Figure 13.5.*j*: Genetic engineering to produce Bt corn. The gene from *Bacillus thuringiensis* that produces Bt toxin is removed and inserted into the corn plant. The Bt corn now produces the insecticide (Bt toxin) and can defend itself against pests. Image by FDA (public domain).

Genetically modified foods are widespread in the United States. For example, 94% of soy crops were genetically modified for herbicide resistance in 2020. Likewise, 8% of cotton and 10% of corn crops were modified for herbicide resistance in addition to the 83% of cotton and 79% of corn crops that were genetically modified in multiple ways.

Genetically modified animals have recently entered the market as well. **AquaAdvantage® salmon** are modified to grow more rapidly and were approved in November of 2015. However, as of March 2021, they have still not been sold due to legal challenges. In 2020, the FDA approved **GalSafe™ pigs** for medicine and food production. These pigs lack a molecule on the outside of their cells that cause allergies in some people.

Advantages of Genetically Modified Crops

Advances in biotechnology may provide consumers with foods that are nutritionally-enriched, longer-lasting, or that contain lower levels of certain naturally occurring toxins present in some food plants. For example, researchers are using biotechnology to try to reduce saturated fats in cooking oils and reduce allergens in foods. Whether these benefits will reach the people who need them most remains to be seen. While cultivating golden rice could address vitamin A deficiency in millions of people, it has not historically been accessible to these people because it is patented and expensive. Similarly, genetically modified seeds could increase the income of impoverished farmers if they were available at low or no cost, but this is not always the case.

Rainbow and SunUp papayas are a success story of how genetically modified crops can benefit small farmers and the economy in general. In the early 1990s, an emerging disease was destroying Hawaii's production of papaya and threatening to decimate the $11-million industry (figure 13.5.*k*). Fortunately, a man named Dennis Gonsalves (figure 13.5.*l*), who was raised on a sugar plantation and then became a plant physiologist at Cornell University, would develop papaya plants genetically engineered to resist the deadly virus. By the end of the decade, the Hawaiian papaya industry and the livelihoods of many farmers were saved thanks to the free distribution of Dr. Gonsalves's seeds.




Figure 13.5.k: The symptoms of papaya ringspot virus are shown on the tree (a) and fruit (b). Image by APS (public domain).



Figure 13.5.l: Dennis Gonsalves genetically engineered papayas to resist the ringspot virus. Image by ARS USDA (public domain).

The effect of genetically modified crops on the environment depends on the specific genetic modification and which agricultural practices it promotes. For example, Bt crops produce their own insecticides such that external application of these chemicals is unnecessary, reducing the negative impacts of industrial agriculture. Ongoing research is exploring whether crops can be engineered to fix nitrogen in the atmosphere (as some bacteria do) rather than relying on ammonium, nitrites, and nitrates in the soil. If these crops were successfully engineered, they could reduce synthetic fertilizer application and minimize nutrient runoff that leads to eutrophication.

Genetically modified crops may have the potential to conserve natural resources, enable animals to more effectively use nutrients present in feed, and help meet the increasing world food and land demands. In practice, however, countries that use genetically modified crops compared to those that do not only enjoy a slight (or nonexistent) increase in yield.

Disadvantages of Genetically Modified Crops

Social Concerns

Intellectual property rights are one of the important factors in the current debate on genetically modified crops. Genetically

modified crops can be patented by agribusinesses, which can lead to them controlling and potentially exploiting agricultural markets. Some accuse companies, such as Monsanto, of allegedly controlling seed production and pricing, much to the detriment of farmers (figure $13.5.m$).



Figure $13.5.m$: Protesters in Washington D.C. oppose genetically modified organisms and Monsanto specifically. Image cropped from Sarah Stierch (CC-BY).

Environmental Concerns

Genetically modified crops present several environmental concerns. Monoculture farming already reduces biodiversity, and cultivating genetically modified crops, for which individual plants are quite similar genetically, exacerbates this. The use of Roundup Ready® crops naturally encourages widespread herbicide use, which could unintentionally kill nearby native plants. This practice would also increase herbicide residues on produce. While Bt crops are beneficial in the sense that they do not require external insecticide application, but Bt toxin is spread in their pollen. An early study found that Bt corn pollen may be harmful to monarch caterpillars (figure $13.5.n$), but only at concentrations that are seldom reached in nature. Follow-up studies found that most of Bt corn grown did not harm monarchs; however, the one strain of Bt corn did was consequently removed from the market.



Figure 13.5.*n*: An early study suggested that pollen containing Bt toxin can harm beneficial and native insects, like this monarch caterpillar. However, without Bt crops, farmers are more likely to spray insecticides, circulating more harmful chemicals than pollen from Bt crops does. Image by Judy Gallagher (CC-BY).

Through interbreeding, or hybridization, genetically modified crops might share their transgenes with wild relatives. This could affect the genetics of those wild relatives and have unforeseen consequences on their populations and could even have implications for the larger ecosystem. For example, if a gene engineered to confer herbicide resistance were to pass from a genetically modified crop to a wild relative, it might transform the wild species into a **superweed** – a species that could not be controlled by herbicide. Its rampant growth could then displace other wild species and the wildlife that depends on it, thus inflecting ecological harm.

Not only could escaped genes alter weedy species, but they could also enter populations of native species. This could make some native species better competitors than they were previously, disrupting ecosystem dynamics. (They could potentially outcompete other native species with which they would otherwise coexist.)

While there is evidence of genetic transfer between genetically modified crops and wild relatives, there is not yet evidence of ecological harm from that transfer. Clearly, continued monitoring, especially for newly-developed crops, is warranted.

The escape of genetically modified animals has potential to disrupt ecosystems as well. For example, if AquaAdvantage salmon were to escape into natural ecosystem, as farmed fish often do, they could outcompete native salmon, including endangered species. Their genetic modification, which facilitates rapid growth, could result in a competitive advantage.

Health Concerns

In addition to environmental risks, some people are concerned about potential health risks of genetically modified crops because they feel that genetic modification alters the intrinsic properties, or essence, of an organism. As discussed above, however, it is known that both traditional breeding practices and modern genetic engineering produce permanent genetic changes. Furthermore, selective breeding actually has a larger and more unpredictable impact on a species's genetics because of its comparably crude nature.

To address these concerns (and others), the US National Academies of Sciences, Engineering, and Medicine (NASEM) published a comprehensive, 500-page report in 2016 that summarized the current scientific knowledge regarding genetically modified crops. The report, titled *Genetically Engineered Crops: Experiences and Prospects*, reviewed more than 900 research articles, in addition to public comments and expert testimony. NASEM's GE Crop Report found "no substantiated evidence of a difference in risks to human health between current commercially available genetically engineered (GE) crops and conventionally bred crops, nor did it find conclusive cause-and-effect evidence of environmental problems from the GE crops." Additionally, the UN's Food and Agriculture Organization has concluded that risks to human and animal health from the use of GMOs are negligible. The scientific consensus on genetically modified crops is quite clear: they are safe for human consumption.

The potential of genetically modified crops to be allergenic is one of the potential adverse health effects, and it should continue to be studied, especially because some scientific evidence indicates that animals fed genetically modified crops have been harmed. NASEM's GE Crop Report concluded that when developing new crops, it is the product that should be studied for potential health and environmental risks, not the process that achieved that product. What this means is, because both traditional breeding practices and modern genetic engineering produce new traits through genetic modification, they both present potential risks. Thus, for the safety of the environment and human health, both should be adequately studied.

Are Genetically Modified Crops the Solution We Need?

Significant resources, both financial and intellectual, have been allocated to answering the question: are genetically modified crops safe for human consumption? After many hundreds of scientific studies, the answer is yes. But a significant question still remains: are they necessary? Certainly, such as in instances like Hawaii's papaya, which were threatened with eradication due to an aggressive disease, genetic engineering was a quick and effective solution that would have been extremely difficult, if not impossible, to solve using traditional breeding practices.

However, in many cases, the early promises of genetically engineered crops – that they would improve nutritional quality of foods, confer disease resistance, and provide unparalleled advances in crop yields – have largely failed to come to fruition. NASEM's GE Crop Report states that while genetically modified crops have resulted in the reduction of agricultural loss from pests, reduced pesticide use, and reduced rates of injury from insecticides for farm workers, they have not increased the rate at which crop yields are advancing when compared to non-GE crops. Additionally, while there are some notable exceptions like golden rice or virus-resistant papayas, very few genetically engineered crops have been produced to increase nutritional capacity or to prevent plant disease that can devastate a farmer's income and reduce food security. The vast majority of genetically modified crops are developed for only two purposes: to introduce herbicide resistance or pest resistance. Genetically modified crops are concentrated in developed countries, and their availability in low-income countries, where they are perhaps most needed, is limited (figure 13.5.*o*).




Figure 13.5.*o*: Global area of genetically modified crops in millions of hectares 1996–2015. Overall, the hectares of genetically modified crops has increased, and in 2011, low-income countries surpassed industrial countries. By 2015, out of the 28 countries that grew biotech crops, 20 were developing, and only eight were developed countries. Latin American, African, and Asian farmers together grew 97.1 million hectares (54%) of the global 179.7 million biotech hectares, whereas industrial countries only planted 83 million hectares or 46%. Image and caption (modified) from Taheri, F., Azadi, H., & D'Haese, M. (2017). A World without Hunger: Organic or GM Crops? *Sustainability*, *9*(4), 580. doi:10.3390/su9040580. (CC-BY)

Suggested Supplementary Reading

NASEM. 2016. Genetically Engineered Crops: Experiences and Prospects.

Attributions

Modified by Melissa Ha from the following sources:

- Biotechnology and Genetic Engineering from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Biotechnology in Medicine and Agriculture from *Concepts in Biology* by OpenStax (licensed under CC-BY(opens in new window))

This page titled 13.5: Selective Breeding and Genetic Engineering is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 14.3: Selective Breeding and Genetic Engineering by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

13.6: Sustainable and Regenerative Agriculture

Sustainable agriculture is often described as farming that can continue over time without harming the environment or using up natural resources. However, some scientists and farmers argue that no form of farming is truly “sustainable” in the strict sense. Any time we farm, we remove plants, nutrients, and energy from the land. This means we are always causing some level of change or loss in the ecosystem.

Because of this, a more accurate term is **regenerative agriculture**. Instead of trying to simply maintain the current state of the land, regenerative agriculture focuses on improving and rebuilding soil, water systems, and ecosystems over time. The goal is not just to avoid harm, but to actively restore natural processes and increase biodiversity.

Regenerative farming often works with natural systems. For example, farmers may:

- add compost or cover crops to rebuild soil nutrients,
- reduce tilling to protect soil structure,
- use diverse crops to support insects and wildlife,
- and rely on natural predator–prey relationships to manage pests.

These practices aim to replace what is taken from the land and improve its health over time, rather than simply minimizing damage.

Earlier definitions of sustainable agriculture, such as those in the 1977 and 1990 U.S. Farm Bills, focused on meeting human needs while protecting environmental quality, using resources efficiently, keeping farms economically viable, and supporting quality of life. These goals are still important today. However, regenerative agriculture builds on them by recognizing that farming systems should not just “do less harm,” but should actively repair and strengthen the natural systems they depend on.

In this way, regenerative agriculture offers a more realistic and forward-looking approach: it acknowledges that farming always has impacts, while also creating opportunities to restore ecosystems and improve long-term resilience.

Promoting biodiversity is a key part of regenerative agriculture because diverse systems are healthier and more resilient. When many different species are present, important ecosystem processes—like nutrient cycling, soil formation, and natural pest control—work more effectively. Instead of just trying to maintain current conditions, regenerative agriculture aims to restore and strengthen these natural systems over time.

One way this happens is by moving away from monocultures (growing only one crop) and toward **polycultures**, where multiple types of plants are grown together. Polycultures reduce the spread of pests and diseases and help keep soils rich in nutrients. Different plants use and return nutrients in different ways, which helps rebuild soil rather than simply using it up.

Regenerative agriculture also emphasizes **genetic diversity** in crops, which helps farms adapt to changing conditions and remain resilient. Seed banks play a key role by storing a wide variety of seeds at low temperatures to keep them viable for long periods. By protecting this diversity, farmers have more options for crops that can adapt to changing climate, disease, and natural disasters. Globally, they act as an insurance policy for the world's food supply. [1, 2, 3, 4, 5] (figure $13.6.a$).

Overall, regenerative agriculture focuses on increasing biodiversity and using natural processes not just to sustain farming systems, but to improve them for the future.



Figure 13.6.*a*: The Svalbard Global Seed Vault in Norway. Image by Miksu (CC-BY-SA).

Integrated Pest Management

Integrated Pest Management (IPM) refers to a mix of farmer-driven, ecologically-based pest control practices that seek to reduce reliance on synthetic chemical pesticides. It uses several methods simultaneously to control pest populations. The steps of integrated pest management are to (1) identify true pests, (2) set thresholds and monitor (figure 13.6.*b*), and (3) develop an action plan. As many insects, microbes, and other organisms found in an agricultural area have a neutral or beneficial effect, it is not necessary to remove them. True pests are those that are causing economic harm, and they are managed (kept below economically damaging levels) rather than eradicated.



Figure 13.6.*b*: A retired Navy commander and landowner inspect an insect trap as part of an Integrated Pest Management (IPM) conservation practice. Image by NRCS (public domain).

An IPM Action Plan draws on four types of control: cultural, mechanical, biological, and chemical. These methods are listed in order of least to most environmental impact and are thus applied in this order. For example, cultural control is attempted first. If that is not effective, mechanical control is added to the plan, and so on. Chemical control is used as a last resort and is de-emphasized in an IPM plan. When pesticides must be used, they are selected and applied in a way that minimizes adverse effects on beneficial organisms, humans, and the environment. It is commonly understood that applying an IPM approach does not necessarily mean eliminating pesticide use, although this is often the case because pesticides are often over-used for a variety

of reasons.

An Alternative to Spraying: Bollworm Control in Shandong

Farmers in Shandong (China) have been using innovative methods to control bollworm infestation in cotton when this insect became resistant to most pesticides. Among the control measures implemented were:

1. The use of pest resistant cultivars and interplanting of cotton with wheat or maize.
2. Use of lamps and poplar twigs to trap and kill adults to lessen the number of adults.
3. If pesticides were used, they were applied on parts of the cotton plant's stem rather than by spraying the whole field (to protect natural enemies of the bollworm).

These and some additional biological control tools have been effective in controlling insect populations and insect resistance, protecting surroundings and lowering costs.

Cultural Control

Cultural control refers to minimizing the conditions that allow pests to thrive and spread. Examples include alternating which crops are planted each year (**crop rotation**), planting multiple types of crops near each other (**intercropping**; figure $13.6.c$), selecting pest-resistant varieties, and planting pest-free rootstock (underground plant parts). Crop rotation prevents pests that are specialized to a particular type of crop from continuing year after year because their host plants are only available in certain years. Similarly, intercropping spatially limits the spread of pests. **Strip cropping** is a type of intercropping that involves growing different types of crops in alternating rows. Pests may infest one (or several) rows of their host species, but they would have to move past multiple rows of non-hosts to access additional host plants. Additionally, cultural control can involve optimizing irrigation and fertilizer application to promote plant defense and limit disease spread. Cultural control methods can be very effective and cost-efficient and present little to no risk to people or the environment.



Figure $13.6.c$: *Momordica charantia* (bitter melon) intercropped with rice in the Philippines (cultural control). Bitter melon plants emerge from holes in plastic, which is used for mechanical control of weeds. Image by Judgefloro (CC-BY-SA).

Mechanical Control

Mechanical control refers to physically removing pests or excluding them with barriers (figure $13.6.d$). Mechanical controls that remove pests include sticky insect traps, mole traps, and removing weeds by hand. Netting to exclude birds, deer fencing, and weed cloth, plastic weed barriers (figure $13.6.c$) or mulch are additional examples of mechanical control.



Figure 13.6.*d*: This solar insect trap is an example of mechanical control. It emits ultraviolet light to attract insects and is recharged using small solar panels. Image by MGK Solar Trap (CC-BY-SA).

Biological Control

Biological control is the use of organisms to reduce pest populations (also see Invasive Species). One biological control strategy involves releasing natural enemies, such as predators, parasites, or parasitoids of the pest organisms (parasitoids are similar to parasites, but they consistently kill their hosts). Home gardeners can also rely on natural enemies by purchasing preying mantises, ladybugs (ladybird beetles), or lacewings for release (figure 13.6.*e*). Successful examples include ladybird beetles to depress aphid populations, parasitoid wasps to control whiteflies, and fungi, such as *Trichoderma*, to suppress fungal-caused plant diseases.



Figure 13.6.*e*: Lacewings are natural predators of garden and crop pests. Image by artsehnPixabay [] (Pixabay license).

If not used carefully, chemical control can have unintended negative consequences for biological control. In 1887, the cottony cushion scale (native to Australia) was devastating the citrus groves of California. A U.S. entomologist went to Australia to find a natural enemy and came back with the **vedalia beetle**, a species of ladybird beetle. Released in California, the beetle quickly brought the scale under control, at least until 1946. In that year, the pest made a dramatic comeback. This coincided with the first use of the now-banned pesticide dichlorodiphenyltrichloroethane (DDT) in the groves. Not only did DDT kill the target pest insects, but it killed the vedalia beetle as well. Only by altering spray procedures and reintroducing the beetle was the scale insect again controlled.

Another strategy for biological control involves releasing sterile males, which compete with fertile males for mates, ultimately decreasing pest population size. This technique was first applied against the screwworm fly, a serious pest of cattle (figure $13.6.f$). The female flies lay their eggs in sores or other open wounds on the animals. After hatching, the larvae eat the tissues of their host. As they do so, they expose a still larger area to egg laying, often finally killing the host.



Figure $13.6.f$: Thought to be eradicated in the United States by 1966, the screwworm fly, a cattle pest, reemerged in deer at Key Deer National Refuge in Florida in 2016. Sterile males were again released, and the pest was eradicated from Florida in 2017. Image by Judy Gallagher (CC-BY).

Prior to its eradication from the southeastern United States, the screwworm was causing huge annual livestock losses. The sterile male technique involves releasing factory-reared and sterilized flies into the natural population. Sterilization is done by exposing the factory flies to just enough gamma radiation to make them sterile but not enough to reduce their general vigor.

Starting in early 1958, up to 50 million sterilized flies were released each week from aircraft flying over Florida and parts of the adjoining states. Each time a fertile female in the natural population mated with a sterile male, the female layed sterile eggs. Since the females mate only once, her reproductive career was at an end. By early 1959, the pest was totally eliminated east of the Mississippi River. The problem in southwestern states was more challenging because the fly winters in Mexico and could move across the border with each new season. Even so, by expanding the program to include Mexico as well, the screwworm fly was finally eliminated from Mexico in 1991.

The sterile male technique also successfully controlled the Mediterranean fruit fly ("medfly"), a destructive fruit fly of citrus, peaches, pears, and apples in California.

Chemical Control

Chemical control refers to the use of pesticides. If chemical control is needed, IPM favors highly targeted chemicals, such as pheromones to disrupt pest mating. **Pheromones** are chemical signals released by animals to communicate with other members of their species. Humans and many insect species alike release pheromones that function in attracting mates. Releasing the pheremones of insect pests can confuse males seeking mates and ultimately hinder them from reproducing (figure $13.6.g$). This "male confusion" has been successful against the pink bollworm that infests cotton and reduced the need for conventional chemical insecticides by 90%. Pheromones have also been successful against pests that attack tomatoes, grapes, and peaches. If targeted chemicals are not effective, IPM may employ conventional pesticides, ideally only applying them to the spots that they are needed and at the lowest effective concentration. Broadcast spraying of non-specific pesticides is a last resort.



Figure $13.6.g$: This pheromone trap is used to control the Asiatic rice borer. The caterpillars of this moth species infest rice stems. Image by Mehdi (CC-BY-SA).

Practices to Maintain Soil Fertility

A variety of practices can maintain soil quality. Many of these strategies have additional benefits such as regulating pests, limiting climate change, and preventing water pollution. These methods enrich the soil with nutrients, ensure proper water holding capacity (ability of the soil to retain water), and limit soil-degrading processes such as erosion and compaction.

Crop Rotation

As previously mentioned, **crop rotations** are planned sequences of different crops over time on the same field (figure $13.6.h$). Rotating crops provides productivity benefits by improving soil nutrient levels and breaking crop pest cycles. Farmers may also choose to rotate crops in order to reduce their production risk through diversification or to manage scarce resources, such as labor, during planting and harvesting. This strategy reduces the pesticide costs by naturally breaking the cycle of weeds, insects and diseases. Also, grass and legumes in a rotation protect water quality by preventing excess nutrients or chemicals from entering water supplies.



Figure 13.6.*h*: Crop rotation can be used at a large agricultural scale (left) or in a smaller garden (right). In each situation, which crop species are planted in each location changes year to year. Left image by USDA (public domain) and right image by Sten Porse (CC-BY-SA).

Intercropping

Intercropping means growing two or more crops in close proximity to each other during part or all of their life cycles to promote soil improvement, biodiversity, and pest management. Incorporating intercropping principles into an agricultural operation increases diversity and interaction between plants, arthropods, mammals, birds and microorganisms resulting in a more stable crop-ecosystem and a more efficient use of space, water, sunlight, and nutrients (figure 13.6.*i*). This collaborative type of crop management mimics nature and is subject to fewer pest outbreaks, improved nutrient cycling and crop nutrient uptake, and increased water infiltration and moisture retention. Soil quality, water quality and wildlife habitat all benefit.



Figure 13.6.*i*: Intercropping of beans and maize (corn). Beans have root nodules that house nitrogen-fixing bacteria, enriching the soil. Image by AnnaJB (CC-BY-SA).

A common example of strip cropping (a type of intercropping; see above) involves alternating a row crop such as corn with a ground-covering crop such as alfalfa. The ground-covering crop helps reduce water runoff and traps soil eroded from the row crop. If this ground-covering crop is a member of the legume family such as alfalfa or soybeans and is associated with nitrogen-fixing bacteria, then alternating the strips from one planting to the next can also help maintain topsoil fertility.

Cover Crops

Cover crops are those that are planted in the off-season to avoid leaving the soil bare. They can prevent soil and wind erosion, improve soil's physical and biological properties, supply nutrients, suppress weeds, improve the availability of soil water, and break pest cycles along with various other benefits. Cover crops are often members of the legume family and help enrich the soil with usable nitrogen. A single species or mix of cover crop species may be planted (figure 13.6.*j*).



Figure 13.6.*j*: A roller (crimper) on a tractor being used to roll down a rye and hairy vetch cover crop in April in the United States. While hairy vetch can limit soil erosion and add nitrogen to the soil, it is also an invasive species that must be carefully managed. Image and caption (modified) from Mr. 1032 (CC-BY-SA).

Agroforestry

Agroforestry is the process of planting rows of trees interspersed with a cash crop (figure $13.6.k - l$). Besides helping to prevent wind and water erosion of the soil, the trees provide shade which helps promote soil moisture retention. Decaying tree litter also provides some nutrients for the interplanted crops. The trees themselves may provide a cash crop. For example, fruit or nut trees may be planted with a grain crop. You can learn more about agroforestry using this interactive site by the United States Department of Agriculture and Forest Service.



Figure $13.6.k$: Through agroforestry, high-value non-timber crops (food, medicinal plants, woody florals, and crafts) can be cultivated under the protection of a forest canopy that has been managed to provide a favorable crop environment. Image and caption (modified) from USDA/ UFS USDA [USDA] [USDA] (public domain).

Contour and Terrace Farming

Contour farming involves plowing and planting crop rows along the natural contours of gently sloping land (figure $13.6.l$). The lines of crop rows perpendicular to the slope help to slow water runoff, inhibit the formation of channels of water, and limit soil erosion (and the resultant loss of nutrients). **Terracing** is a common technique used to control water erosion on more steeply sloped hills and mountains (figure $13.6.l$). Broad, level terraces are constructed along the contours of the slopes, and these act as dams trapping water for crops, reducing runoff, and limiting erosion.



Figure 13.6.*l*: Contour farming (left) and terracing (right). In contour farming, rows of crops wave perpendicular to the slopes of the land. This example (left) also employs agroforestry. Terrace farming involves making step-like structures (terraces) perpendicular to the incline of a slope to reduce erosion. Left image by Pixture2016 (CC-BY-SA). Right image by USDA (public domain).

Minimal Tillage and No-till Agriculture

In modern agricultural practices, heavy machinery is used to prepare the seedbed for planting, to control weeds, and to harvest the crop. The use of heavy equipment has many advantages in saving time and labor, but can cause compaction of soil and disruption of the natural soil organisms. The problem with soil compaction is that increased soil density limits root penetration depth and may inhibit proper plant growth. Another aspect of soil tillage (mixing the soil) is that it may lead to more rapid decomposition of organic matter due to greater soil aeration. Over large areas of farmland, this has the unintended consequence of releasing more carbon and nitrous oxides (greenhouse gases) into the atmosphere, thereby contributing to climate change.

One of the easiest ways to prevent these problems is to minimize the amount of **tillage**, or turning over of the soil. In **minimal tillage** (conservation tillage) or **no-till agriculture**, the land is disturbed as little as possible by leaving crop residue in the fields (figure 13.6.*m*). Special seed drills inject new seeds and fertilizer into the unplowed soil. Tillage of fields does help to break up clods that were previously compacted, so best practices may vary at sites with different soil textures and composition. With proper planning, minimal tillage and no-till agriculture can simultaneously limit soil erosion and compaction, protect soil organisms, reduce costs (if performed correctly), and promote water infiltration. Furthermore, carbon can actually become sequestered into the soil with these methods, thus mitigating climate change. Minimal or no-till agriculture have proved a major success in Latin

America and are being used in South Asia and Africa. However, a drawback of this method is that the crop residue can serve as a good habitat for insect pests and plant diseases.



Figure $13.6.m$: No-till agriculture is an important tool to prevent loss of soil moisture. Image by USDA (public domain).

Windbreaks

Creating **windbreaks** by planting tall trees along the perimeter of farm fields can help control the effects of wind erosion of soil (figure $13.6.n$). Windbreaks reduce wind speed at ground level, an important factor in wind erosion. They also help trap snow in the winter months, leaving soil less exposed. As a side benefit, windbreaks also provide a habitat for birds and animals. One drawback is that windbreaks can be costly to farmers because they reduce the amount of available cropland.



Figure $13.6.n$: Windbreaks are rows of trees and shrubs that reduce wind speed. They improve crop yields, reduce soil erosion, improve water-efficiency, protect livestock, and conserve energy. Image and caption (modified) from USDA [USDA/UFS] (public domain).

Organic Agriculture

Organic agriculture is often incorporated into regenerative farming practices (figure $13.6.o$). To be certified as organic, farms must avoid using synthetic pesticides, synthetic fertilizers, and genetically modified organisms (figure $13.6.p$). Organic meat,

poultry, eggs, and dairy products come from animals that are given no antibiotics or growth hormones. Pests may instead be controlled by natural enemies or naturally produced substances such as neem oil or diatomaceous earth. Some organic alternatives to synthetic pesticides have relatively low environmental impact (such as salt spray), but there are many naturally produced compounds that are still toxic to people or cause ecological harm when widely used. Interestingly, external application of Bt toxin is approved for organic farming, but use of genetically engineered Bt crops is not. The latter results in lower concentrations of Bt toxin in the environment because it is locally produced directly by the plants themselves. In summary, although many practices of organic agriculture benefit the environment and meet sustainability goals, some organic farms are not regenerative and some regenerative farms are not organic.



Figure 13.6.*o*: United States Department of Agriculture (USDA) Conservationist Shemekia Mosley (right) works with Nuri Icgoren (left) who operates Urban Sprout Farms, a biodynamic, certified organic urban farm Lakewood Heights, Georgia. Image and caption (modified) by Preston KeresUSDA [] (public domain).



Figure 13.6.*p*: The USDA certified organic label. Modified from image by nikoretro (CC-BY-SA).

Consumer Choices that Support a More Sustainable Agricultural System

Even if you're not a farmer or lawmaker, you have the power to promote more sustainable agricultural practices as a consumer. As discussed in Food Chains and Food Webs, it generally requires more land and more energy to produce meat compared to plant-based foods due to the inefficient transfer of energy from one trophic level to the next. For this reason, plant-based diets tend to be more sustainable, but this depends on the types of foods consumed and how they were produced.

Local food is not only fresher, but it requires fewer food miles (figure 13.6.*q*). In some ways, **organic food** causes less environmental degradation, but note that organic does not necessarily mean sustainable. Because it limits machinery, synthetic pesticides, and synthetic fertilizers, it has a smaller **carbon footprint** (meaning it minimizes contribution to climate change). The

Clean 15 is a list of produce that is low in pesticide residues. If you can't afford all organic foods, these are the best items to purchase non-organic. Examples include avocados, sweet corn, and pineapple. The Dirty Dozen lists produce that has the most pesticide residues. If you can only afford a few organic items, these are best to purchase organic. Examples include strawberries, spinach, and nectarines. Finally, some foods, such as beef, have higher **carbon footprint** and **water footprint** than others.



Figure $13.6.q$: Farmers' markets are one option for purchasing locally grown foods. Image by NatalieMaynor (CC-BY).

The Future of the Sustainable Agriculture Concept

Many in the agricultural community have adopted the sense of urgency and direction pointed to by the sustainable agriculture concept (figure $13.6.r$). Sustainability has become an integral component of many government, commercial, and non-profit agriculture research efforts, and it is beginning to be woven into agricultural policy. Increasing numbers of farmers and ranchers have embarked on their own paths to sustainability, incorporating integrated and innovative approaches into their own enterprises.




Figure $13.6.r$: Sustainable agriculture combines many approaches. (1) If chemical are used, they are applied only where needed. (2) Through the Conservation Reserve Program, the most ecologically important patches are left intact. (3) Terraces on slopes reduce soil erosion. (4) Crops are scouted for pests as part of monitoring, a component of integrated pest management. (5) Cover crops prevent erosion and enrich the soil during off-seasons. (6) Different types of crops are included are planted in close proximity (intercropping) or in different years (crop rotation). (7) Land is either not tilled at all or tilled minimally (conservation tillage). (8) Through precision nutrient management, soil is sampled at different locations, and fertilizer is only applied where needed. (9) Heavy equipment is used only as needed and in such a way that conserves fuel. (10) Irrigation is precisely timed to conserve water. (11) Water storage ponds help recharge groundwater. Image by the United Soybean Board (CC-BY).

Attributions

Modified by Erin Rempala.

Modified by Melissa Ha from the following sources:

- Sustainable Agriculture from *Environmental Biology* by Matthew R. Fisher (licensed under CC-BY(opens in new window))
- Soils from *AP Environmental Science* by University of California College Prep, University of California (licensed under CC-BY(opens in new window)). Download for free at CNX(opens in new window).
- Sustainability Definitions. USDA. Accessed 01-05-2021. (public domain)
- Biological Control from *Biology* by John W. Kimball (licensed under CC-BY(opens in new window))
- Integrated Pest Management (IPM) Principles. EPA. Accessed 01-05-2021 (public domain)
- Cover Crops and Soil Health. Natural Resources Conservation Service, USDA. Accessed 01-06-2021. (public domain)

This page titled 13.6: Sustainable and Regenerative Agriculture is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Melissa Ha and Rachel Schleiger (ASCCC Open Educational Resources Initiative).

- 14.4: Sustainable Agriculture by Melissa Ha and Rachel Schleiger is licensed CC BY-SA 4.0.

INDEX

A

abiotic factors
3.1.1: Ecological Hierarchy

acid rain
8.3: Acid Deposition

adaptive radiation
5.4: The Biodiversity Crisis
5.6: Extinction

age structure
6: Human Population Growth

air pressure
8.1: Earth's Atmosphere
9.1: Weather, Climate, and Climate Change

Air Temperature
9.1: Weather, Climate, and Climate Change

Aposematic Coloration
3.2.1.2.3: Competition

aquatic biomes
3.3.5.3: Aquatic Biomes

atmosphere
8.1: Earth's Atmosphere

atmospheric science
8.1: Earth's Atmosphere
9.2: Earth's Energy Balance and Radiation

Author tag:OpenStax
5.4: The Biodiversity Crisis

B

Batesian mimicry
3.2.1.2.3: Competition

biodiversity
3: Ecology
3.1.1: Ecological Hierarchy
5.4: The Biodiversity Crisis
5.6: Extinction

biodiversity hotspot
5.4: The Biodiversity Crisis
5.6: Extinction

bioenergetics
3.3.2: Energy

biological agents
12.1: Types of Environmental Hazards

biological organization
3.1.1: Ecological Hierarchy

biome
3.3.5.1: Climate and Biomes

biosphere
3.1.1: Ecological Hierarchy

biotic interactions
3: Ecology

bush meat
5.5: Threats to Biodiversity

C

camouflage
3.2.1.2.3: Competition

carbon cycle
9.3: The Greenhouse Effect

carrying capacity
3.1.4: Population Growth and Regulation

Centers for Disease Control and Prevention
12.2: Epidemiology

chemical diversity
5.4: The Biodiversity Crisis
5.6: Extinction

chytridiomycosis
5.5: Threats to Biodiversity

clean energy
7.2.5: The Promise of Fusion

climate
3: Ecology
9.1: Weather, Climate, and Climate Change

climate change
9.1: Weather, Climate, and Climate Change
9.3: The Greenhouse Effect

climax community
3.2.1.2.3: Competition

clouds
9.1: Weather, Climate, and Climate Change

coal
7: The Energy System

commensalism
3.2.1.1: Symbiosis
3.2.1.2.3: Competition

communities
3.1.1: Ecological Hierarchy

community
3.2.1: Biotic Interactions

community dynamics
3: Ecology

community ecology
3.2.1: Biotic Interactions

competitive exclusion principle
3.2.1: Biotic Interactions
3.2.1.2.3: Competition

conservation
3: Ecology

Crop rotations
13.6: Sustainable and Regenerative Agriculture

D

DDT
12.1: Types of Environmental Hazards

density
8.1: Earth's Atmosphere

DNA barcoding
5.8: Preserving Biodiversity

E

Earth's atmosphere
9.3: The Greenhouse Effect

ecological footprint
1.4: Human Impact on the Planet

ecological hierarchy
3.1.1: Ecological Hierarchy

ecosystem diversity
5.4: The Biodiversity Crisis
5.6: Extinction

ecosystem structure
3: Ecology

ecosystems
3.1.1: Ecological Hierarchy

electric fields
9.2: Earth's Energy Balance and Radiation

electromagnetic radiation
9.2: Earth's Energy Balance and Radiation

electromagnetic waves
9.2: Earth's Energy Balance and Radiation

Emsleyan/Mertensian mimicry
3.2.1.2.3: Competition

Endemic diseases
12.2: Epidemiology

endemic species
5.4: The Biodiversity Crisis
5.6: Extinction

energy
9.2: Earth's Energy Balance and Radiation

energy balance
9.3: The Greenhouse Effect

energy conservation
7: The Energy System

energy efficiency
7: The Energy System

energy generation
7.2.5: The Promise of Fusion

environmental disturbance
3.2.1.2.3: Competition

environmental impacts
7: The Energy System

environmental toxicology
12.4: Environmental Toxicology

Epidemic diseases
12.2: Epidemiology

epidemiology
12.2: Epidemiology

exotic species
5.5: Threats to Biodiversity

exponential growth
3.1.4: Population Growth and Regulation

extinction
5.4: The Biodiversity Crisis
5.6: Extinction

extinction rate
5.4: The Biodiversity Crisis
5.6: Extinction

extinction risks
3: Ecology

extraction methods
7: The Energy System

F

Food Security
13.1: Food Security

fossil
4.3: Evidence for Evolution

fossil fuels
7: The Energy System

foundation species
3.2.1.2.3: Competition
frequency
9.2: Earth's Energy Balance and Radiation
fusion reactors
7.2.5: The Promise of Fusion

G

gases
8.1: Earth's Atmosphere
genetic diversity
5.4: The Biodiversity Crisis
5.6: Extinction
genetic drift
4.2: Other Mechanisms of Evolution
genetic engineering
13.5: Selective Breeding and Genetic Engineering
global warming
9.3: The Greenhouse Effect
greenhouse effect
9.3: The Greenhouse Effect
greenhouse gas
8.1: Earth's Atmosphere
greenhouse gas emissions
7: The Energy System
greenhouse gases
9.3: The Greenhouse Effect

H

Herbivory
3.2.1: Biotic Interactions
heterogeneity
5.4: The Biodiversity Crisis
5.6: Extinction
history
7: The Energy System
host
3.2.1.2.3: Competition
human activities
9.3: The Greenhouse Effect
human population
6: Human Population Growth
humidity
9.1: Weather, Climate, and Climate Change
hydrogen isotopes
7.2.5: The Promise of Fusion

I

incidence
12.2: Epidemiology
infrared radiation
9.3: The Greenhouse Effect
integrated pest management
13.6: Sustainable and Regenerative Agriculture
intercropping
13.6: Sustainable and Regenerative Agriculture
international collaboration
7.2.5: The Promise of Fusion
island biogeography
3.2.1.2.3: Competition
ITER
7.2.5: The Promise of Fusion

K

keystone species
3.2.1.2.3: Competition
Koch's postulates
12.2: Epidemiology

L

logistic growth
3.1.4: Population Growth and Regulation

M

magnetic confinement
7.2.5: The Promise of Fusion
magnetic fields
9.2: Earth's Energy Balance and Radiation
megafauna
5.4: The Biodiversity Crisis
5.6: Extinction
Mertensian mimicry
3.2.1.2.3: Competition
meteorology
8.1: Earth's Atmosphere
mimicry
3.2.1: Biotic Interactions
minimal tillage
13.6: Sustainable and Regenerative Agriculture
morbidity
12.2: Epidemiology
morbidity rates
12.2: Epidemiology
mortality
12.2: Epidemiology
mortality rates
12.2: Epidemiology
Müllerian Mimicry
3.2.1.2.3: Competition
mutualism
3.2.1.1: Symbiosis
3.2.1.2.3: Competition

N

natural selection
4.2: Other Mechanisms of Evolution
nuclear fission
7.2.5: The Promise of Fusion
nuclear fusion
7.2.5: The Promise of Fusion

O

organic farming
13.6: Sustainable and Regenerative Agriculture
organisms
3.1.1: Ecological Hierarchy
overpopulation
6.4: Looking Ahead
ozone depletion
8.4: Ozone Depletion

P

paleosols
13.3: Soils
pandemic diseases
12.2: Epidemiology
parasite
3.2.1.2.3: Competition
parasitism
3.2.1.1: Symbiosis
photons
9.2: Earth's Energy Balance and Radiation
pioneer species
3.2.1.2.3: Competition
population ecology
3: Ecology
population growth
3.1.4: Population Growth and Regulation
populations
3.1.1: Ecological Hierarchy
precautionary principle
1.4: Human Impact on the Planet
precipitation
9.1: Weather, Climate, and Climate Change
predator
3.2.1: Biotic Interactions
preservation
5.8: Preserving Biodiversity
prevalence
12.2: Epidemiology
primary succession
3.2.1.2.3: Competition
pseudogenes
3.2.1.1: Symbiosis

R

radiation
9.2: Earth's Energy Balance and Radiation
relative species abundance
3.2.1.2.3: Competition
Remediation
10.5: Water Pollution
renewable energy
7: The Energy System

S

secondary plant compound
5.2: The Importance of Biodiversity to Human Life
secondary succession
3.2.1.2.3: Competition
Soil processes
13.3: Soils
Soil Profiles
13.3: Soils
solar energy
8.1: Earth's Atmosphere
9.3: The Greenhouse Effect
species richness
3.2.1.2.3: Competition
Sporadic diseases
12.2: Epidemiology
sun
8.1: Earth's Atmosphere
sustainable agriculture
13.6: Sustainable and Regenerative Agriculture
sustainable energy
7: The Energy System

symbiosis
3.2.1: Biotic Interactions
3.2.1.1: Symbiosis
3.2.1.2.3: Competition

T

temperature
8.1: Earth's Atmosphere
terrestrial biomes
3: Ecology
3.3.5.2: Terrestrial Biomes
Total Fertility Rate
6.4: Looking Ahead
Tragedy of the Commons
5.5: Threats to Biodiversity

W

wavelength
9.2: Earth's Energy Balance and Radiation
weather
9.1: Weather, Climate, and Climate Change
weather events
9.1: Weather, Climate, and Climate Change
wildlife
3.1.1: Ecological Hierarchy
wildlife corridors
3: Ecology
wind
9.1: Weather, Climate, and Climate Change

This page titled Index is shared under a license and was authored, remixed, and/or curated by Erin Rempala.

GLOSSARY

S

Sample Word 1 | Sample Definition 1

This page titled Glossary is shared under a license and was authored, remixed, and/or curated by Erin Rempala.

DETAILED LICENSING

Overview

Title: BIOL 101- Environmental Science

Webpages: 136

Applicable Restrictions: Noncommercial

All licenses found:

- CC BY-NC-SA 4.0: 91.9% (125 pages)
- Undeclared: 7.4% (10 pages)
- Other: 0.7% (1 page)

By Page

- BIOL 101- Environmental Science - *CC BY-NC-SA 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Other*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Introduction to Environmental Science - *CC BY-NC-SA 4.0*
    - 1.1: The Process of Science - *CC BY-NC-SA 4.0*
    - 1.2: What is Science? - *CC BY-NC-SA 4.0*
    - 1.3: Empirical and Applied Science - *CC BY-NC-SA 4.0*
    - 1.4: Human Impact on the Planet - *CC BY-NC-SA 4.0*
    - 1.5: Environmental Science - *CC BY-NC-SA 4.0*
    - 1.6: Practicing with Data- Wildlife Corridors - *CC BY-NC-SA 4.0*
  - 2: Humans and the Environment - *CC BY-NC-SA 4.0*
    - 2.1: Planetary Boundaries- Earth's Safe Operating Space - *CC BY-NC-SA 4.0*
    - 2.2: Environmental Ethics and Worldviews - *CC BY-NC-SA 4.0*
    - 2.3: Environmental Justice and Equity - *CC BY-NC-SA 4.0*
    - 2.4: Sustainability and Sustainable Development - *CC BY-NC-SA 4.0*
    - 2.5: Systems Thinking and Modeling Sustainability - *CC BY-NC-SA 4.0*
    - 2.6: Historical Foundations of Environmental Policy - *CC BY-NC-SA 4.0*
  - 3: Ecology - *CC BY-NC-SA 4.0*
    - 3.1: Population Ecology - *CC BY-NC-SA 4.0*
      - 3.1.1: Ecological Hierarchy - *CC BY-NC-SA 4.0*
      - 3.1.2: Population Distribution - *CC BY-NC-SA 4.0*
      - 3.1.3: Population Size - *CC BY-NC-SA 4.0*
      - 3.1.4: Population Growth and Regulation - *CC BY-NC-SA 4.0*
      - 3.1.5: Life History - *CC BY-NC-SA 4.0*
    - 3.2: Community Ecology - *CC BY-NC-SA 4.0*
      - 3.2.1: Biotic Interactions - *CC BY-NC-SA 4.0*
        - 3.2.1.1: Symbiosis - *CC BY-NC-SA 4.0*
        - 3.2.1.2: Trophic Interactions - *CC BY-NC-SA 4.0*
          - 3.2.1.2.1: Predation - *CC BY-NC-SA 4.0*
          - 3.2.1.2.2: Herbivory - *CC BY-NC-SA 4.0*
          - 3.2.1.2.3: Competition - *CC BY-NC-SA 4.0*
      - 3.2.2: Community Structure and Dynamics - *CC BY-NC-SA 4.0*
      - 3.2.3: Data Dive- Beaver Impacts on Wetlands - *CC BY-NC-SA 4.0*
    - 3.3: Ecosystem Ecology - *CC BY-NC-SA 4.0*
      - 3.3.1: Matter - *CC BY-NC-SA 4.0*
      - 3.3.2: Energy - *CC BY-NC-SA 4.0*
      - 3.3.3: Biogeochemical Cycles - *CC BY-NC-SA 4.0*
      - 3.3.4: Energy Flow in Food Webs - *CC BY-NC-SA 4.0*
      - 3.3.5: Ecosystem Types and Dynamics - *CC BY-NC-SA 4.0*
        - 3.3.5.1: Climate and Biomes - *CC BY-NC-SA 4.0*
        - 3.3.5.2: Terrestrial Biomes - *CC BY-NC-SA 4.0*
        - 3.3.5.3: Aquatic Biomes - *CC BY-NC-SA 4.0*
      - 3.3.6: Data Dive- Biome Carbon Storage - *CC BY-NC-SA 4.0*

- 4: Evolution - *CC BY-NC-SA 4.0*
  - 4.1: Evolution and Natural Selection - *CC BY-NC-SA 4.0*
  - 4.2: Other Mechanisms of Evolution - *CC BY-NC-SA 4.0*
  - 4.3: Evidence for Evolution - *CC BY-NC-SA 4.0*
- 5: Biodiversity - *CC BY-NC-SA 4.0*
  - 5.1: The Value of Biodiversity - *CC BY-NC-SA 4.0*
  - 5.2: The Importance of Biodiversity to Human Life - *CC BY-NC-SA 4.0*
  - 5.3: Levels of Biodiversity - *CC BY-NC-SA 4.0*
  - 5.4: The Biodiversity Crisis - *CC BY-NC-SA 4.0*
  - 5.5: Threats to Biodiversity - *CC BY-NC-SA 4.0*
  - 5.6: Extinction - *CC BY-NC-SA 4.0*
  - 5.7: Measuring Biodiversity Loss - *CC BY-NC-SA 4.0*
  - 5.8: Preserving Biodiversity - *CC BY-NC-SA 4.0*
  - 5.9: Policies Protecting Biodiversity - *CC BY-NC-SA 4.0*
  - 5.10: Data Dive- Threats to Biodiversity - *CC BY-NC-SA 4.0*
- 6: Human Population Growth - *CC BY-NC-SA 4.0*
  - 6.1: History of Human Population Growth - *CC BY-NC-SA 4.0*
  - 6.2: The Rate of Human Population Growth - *CC BY-NC-SA 4.0*
  - 6.3: Factors Affecting Population Growth - *CC BY-NC-SA 4.0*
  - 6.4: Looking Ahead - *CC BY-NC-SA 4.0*
  - 6.5: Data Dive- World Population Densities - *CC BY-NC-SA 4.0*
- 7: The Energy System - *CC BY-NC-SA 4.0*
  - 7.1: Fossil Fuels - *CC BY-NC-SA 4.0*
    - 7.1.1: Types of Fossil Fuels and Formation - *CC BY-NC-SA 4.0*
    - 7.1.2: Mining, Processing, and Generating Electricity - *CC BY-NC-SA 4.0*
    - 7.1.3: Fossil Fuel Consumption - *CC BY-NC-SA 4.0*
    - 7.1.4: Consequences of Fossil Fuels - *CC BY-NC-SA 4.0*
  - 7.2: Nuclear Energy - *CC BY-NC-SA 4.0*
    - 7.2.1: Radioactive Isotopes - *CC BY-NC-SA 4.0*
    - 7.2.2: Generating Electricity with Nuclear Energy - *CC BY-NC-SA 4.0*
    - 7.2.3: Nuclear Energy Consumption - *CC BY-NC-SA 4.0*
    - 7.2.4: Pros and Cons of Nuclear Energy - *CC BY-NC-SA 4.0*
    - 7.2.5: The Promise of Fusion - *CC BY-NC-SA 4.0*
    - 7.2.6: Data Dive- Global Nuclear Power Generation - *CC BY-NC-SA 4.0*
  - 7.3: Renewable Energy and Conservation - *CC BY-NC-SA 4.0*
    - 7.3.1: Renewable Energy History and Consumption - *CC BY-NC-SA 4.0*
    - 7.3.2: Wind Energy - *CC BY-NC-SA 4.0*
    - 7.3.3: Solar Energy - *CC BY-NC-SA 4.0*
    - 7.3.4: Hydropower - *CC BY-NC-SA 4.0*
    - 7.3.5: Geothermal Energy - *CC BY-NC-SA 4.0*
    - 7.3.6: Biofuels (Biomass Energy) - *CC BY-NC-SA 4.0*
    - 7.3.7: Energy Efficiency and Conservation - *CC BY-NC-SA 4.0*
    - 7.3.8: Data Dive- Global Renewable Energy Generation - *CC BY-NC-SA 4.0*
- 8: Atmospheric Pollution - *CC BY-NC-SA 4.0*
  - 8.1: Earth's Atmosphere - *CC BY-NC-SA 4.0*
  - 8.2: Air Pollution - *CC BY-NC-SA 4.0*
  - 8.3: Acid Deposition - *CC BY-NC-SA 4.0*
  - 8.4: Ozone Depletion - *CC BY-NC-SA 4.0*
- 9: Global Climate Change - *CC BY-NC-SA 4.0*
  - 9.1: Weather, Climate, and Climate Change - *CC BY-NC-SA 4.0*
  - 9.2: Earth's Energy Balance and Radiation - *CC BY-NC-SA 4.0*
  - 9.3: The Greenhouse Effect - *CC BY-NC-SA 4.0*
  - 9.4: Evidence and Causes of Climate Change - *CC BY-NC-SA 4.0*
  - 9.5: Impacts of Climate Change - *CC BY-NC-SA 4.0*
  - 9.6: Ocean Acidification- The Other Carbon Problem - *CC BY-NC-SA 4.0*
  - 9.7: Mitigating and Adapting to Climate Change - *Undeclared*
  - 9.8: Data Dive- Flooding in Venice Italy - *CC BY-NC-SA 4.0*
- 10: The Water System - *CC BY-NC-SA 4.0*
  - 10.1: The Hydrologic Cycle and Freshwater Distribution - *CC BY-NC-SA 4.0*
  - 10.2: Water Usage - *CC BY-NC-SA 4.0*
  - 10.3: Water Scarcity and Sustainable Water Management - *CC BY-NC-SA 4.0*
  - 10.4: Case Study - The Salton Sea - *CC BY-NC-SA 4.0*
  - 10.5: Water Pollution - *CC BY-NC-SA 4.0*
  - 10.6: Water Treatment- Protecting Drinking Water and Treating Wastewater - *CC BY-NC-SA 4.0*
  - 10.7: Preventing Water Pollution and Restoring Watersheds - *CC BY-NC-SA 4.0*
- 11: Waste, Materials, and the Circular Economy - *CC BY-NC-SA 4.0*
  - 11.1: Waste Generation - *CC BY-NC-SA 4.0*
  - 11.2: Disposal - *CC BY-NC-SA 4.0*
  - 11.3: Environmental Impacts of Solid Waste - *CC BY-NC-SA 4.0*
  - 11.4: From Waste Management to a Circular Economy - *CC BY-NC-SA 4.0*
  - 11.5: Circular Economy and Stewardship - *CC BY-NC-SA 4.0*
- 12: Environmental Health and Toxicology - *CC BY-*

NC-SA 4.0
- 12.1: Types of Environmental Hazards - *CC BY-NC-SA 4.0*
- 12.2: Epidemiology - *CC BY-NC-SA 4.0*
- 12.3: Infectious Diseases - *CC BY-NC-SA 4.0*
- 12.4: Environmental Toxicology - *CC BY-NC-SA 4.0*
- 12.5: Case Study - The Love Canal Disaster - *CC BY-NC-SA 4.0*
- 12.6: Environmental Hazard Reduction - *CC BY-NC-SA 4.0*

◦ 13: The Agricultural System - *CC BY-NC-SA 4.0*
- 13.1: Food Security - *CC BY-NC-SA 4.0*
- 13.2: Industrial Agriculture - *CC BY-NC-SA 4.0*
- 13.3: Soils - *CC BY-NC-SA 4.0*
- 13.4: Soil Degradation - *CC BY-NC-SA 4.0*
- 13.5: Selective Breeding and Genetic Engineering - *CC BY-NC-SA 4.0*
- 13.6: Sustainable and Regenerative Agriculture - *CC BY-NC-SA 4.0*

◦ Back Matter - *Undeclared*
- Index - *Undeclared*
- Glossary - *Undeclared*
- Detailed Licensing - *Undeclared*

This page titled BIOL 101- Environmental Science is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Erin Rempala.