# BIOLOGY FOR MAJORS I



Lumen Learning

LibreTexts™

# Book- Biology for Majors I (Lumen)

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.




LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 05/03/2026

# TABLE OF CONTENTS


Licensing

## 1: Course Contents

- 1.1: About This Course
- 1.2: Course Contents at a Glance
- 1.3: Learning Outcomes

## 2: Faculty Resources

- 2.1: Faculty Resources Overview
- 2.2: Pacing
- 2.3: PDF
- 2.4: PowerPoints
- 2.5: Assignments
- 2.6: Assignments with Samples
- 2.7: Question Banks
- 2.8: I Need Help

## 3: Module 1- Introduction to Biology

- 3.1: Why It Matters- Introduction to Biology
- 3.2: Introduction to Characteristics of Life
- 3.3: Properties of Life
- 3.4: Introduction to Prokaryotes and Eukaryotes
- 3.5: Comparing Prokaryotic and Eukaryotic Cells
- 3.6: Levels of Organization of Living Things
- 3.7: Introduction to Taxonomy
- 3.8: The Diversity of Life
- 3.9: Phylogenetic Trees
- 3.10: Taxonomy
- 3.11: Introduction to the Study of Biology
- 3.12: The Branches of Biology
- 3.13: Scientific Inquiry
- 3.14: Experiments and Hypotheses
- 3.15: Interpreting Results
- 3.16: Putting It Together- Introduction to Biology

## 4: Module 2- Chemistry of Life

- 4.1: Why It Matters- The Chemistry of Life
- 4.2: Introduction to Atoms and Elements
- 4.3: Atoms
- 4.4: The Periodic Table of Elements
- 4.5: Electrons
- 4.6: Introduction to Atomic Bonds
- 4.7: Ionic Bonds
- 4.8: Covalent Bonds

- 4.9: Hydrogen Bonds
- 4.10: Introduction to Functional Groups
- 4.11: Functional Groups
- 4.12: Introduction to The pH Scale
- 4.13: Acids and Bases
- 4.14: Buffers
- 4.15: Introduction to Chemical Reactions
- 4.16: Chemical Reactions and Molecules
- 4.17: Anabolic and Catabolic Pathways
- 4.18: Enzymes
- 4.19: Introduction to Thermodynamics
- 4.20: Open and Closed Systems
- 4.21: The First Law of Thermodynamics
- 4.22: The Second Law of Thermodynamics
- 4.23: Putting It Together- Chemistry of Life
- 4.24: Assignment- Biological Astronaut
- 4.25: Assignment- Interacting with Attributes of Molecules


## 5: Module 3- Important Biological Macromolecules


- 5.1: Why It Matters- Important Biological Macromolecules
- 5.2: Introduction to Proteins
- 5.3: Amino Acids
- 5.4: Protein Structure
- 5.5: Function of Proteins
- 5.6: Introduction to Lipids
- 5.7: Lipids
- 5.8: Introduction to Carbohydrates
- 5.9: Structure and Function of Carbohydrates
- 5.10: Introduction to Nucleic Acids
- 5.11: Structure of Nucleic Acids
- 5.12: DNA and RNA
- 5.13: Introduction to Comparing Biological Macromolecules
- 5.14: Different Types of Biological Macromolecules
- 5.15: Putting It Together- Important Biological Macromolecules
- 5.16: Assignment- Nutritionist for a Day


## 6: Module 4- Cellular Structure


- 6.1: Why It Matters- Cellular Structure
- 6.2: Introduction to Microscopes
- 6.3: Microscopy
- 6.4: Introduction to Organelles
- 6.5: Cytoplasm
- 6.6: Nucleus
- 6.7: Ribosomes, Mitochondria, and Peroxisomes
- 6.8: Endomembrane System
- 6.9: Vesicles
- 6.10: Unique Features of Animal and Plant Cells
- 6.11: Introduction to Cytoskeleton
- 6.12: Microfilaments

- 6.13: Intermediate Filaments
- 6.14: Microtubules
- 6.15: Introduction to Cell Surface
- 6.16: Cell Membranes
- 6.17: Cell Junctions
- 6.18: The Cell Wall
- 6.19: Putting It Together- Cellular Structure
- 6.20: Assignment- Cell Builder

## 7: Module 5- Cell Membranes

- 7.1: Why It Matters- Cell Membranes
- 7.2: Introduction to Structure of the Membrane
- 7.3: Structure of the Cell Membrane
- 7.4: Phospholipids
- 7.5: Introduction to Kinds of Transport
- 7.6: Passive Transport
- 7.7: Active Transport
- 7.8: Membranes and Transport
- 7.9: Introduction to Endocytosis and Exocytosis
- 7.10: Endocytosis
- 7.11: Exocytosis
- 7.12: Putting It Together- Cell Membranes
- 7.13: Assignment- Membranes Alive!

## 8: Module 6- Metabolic Pathways

- 8.1: Why It Matters- Metabolic Pathways
- 8.2: Introduction to Redox Reactions
- 8.3: Electrons and Energy
- 8.4: ATP in Living Systems
- 8.5: Introduction to Photosynthesis
- 8.6: An Overview of Photosynthesis
- 8.7: Spectrums of Light
- 8.8: Light-Dependent Reactions
- 8.9: Light-Independent Reactions
- 8.10: Photosynthesis
- 8.11: Introduction to Cellular Respiration
- 8.12: Glycolysis
- 8.13: Pyruvate Oxidation
- 8.14: Citric Acid Cycle
- 8.15: Electron Transport Chain
- 8.16: Cellular Respiration
- 8.17: Introduction to Fermentation
- 8.18: Metabolism without Oxygen
- 8.19: Types of Fermentation
- 8.20: Putting It Together- Metabolic Pathways
- 8.21: Assignment- Observing Energy Transactions

## 9: Module 7- Cell Communication


- 9.1: Why It Matters- Cell Communication
- 9.2: Introduction to Signaling Molecules and Cellular Receptors
- 9.3: Types of Signals
- 9.4: Signaling Molecules
- 9.5: Signaling Receptors
- 9.6: Introduction to Propagation of the Signal
- 9.7: Binding Initiates a Signaling Pathway
- 9.8: Methods of Intracellular Signaling
- 9.9: Introduction to Response to the Signal
- 9.10: Expression and Growth
- 9.11: Cell Death
- 9.12: Introduction to Signaling in Single-Celled Organisms
- 9.13: Signaling in Yeast
- 9.14: Signaling in Bacteria
- 9.15: Putting It Together- Cell Communication


## 10: Module 8- Cell Division


- 10.1: Why It Matters- Cell Division
- 10.2: Introduction to Chromosomes and DNA Packaging
- 10.3: DNA and Chromosomes
- 10.4: Chromosome Structure
- 10.5: Introduction to the Cell Cycle
- 10.6: Interphase
- 10.7: Mitosis
- 10.8: Cytokinesis
- 10.9: The Complete Cycle
- 10.10: Introduction to Cell Cycle Checkpoints
- 10.11: Control of the Cell Cycle
- 10.12: Cancer and the Cell Cycle
- 10.13: Introduction to Meiosis
- 10.14: Stages of Meiosis
- 10.15: Meiosis I
- 10.16: Meiosis II
- 10.17: Meiosis
- 10.18: Introduction to Genetic Diversity
- 10.19: Sexual Reproduction
- 10.20: Genetic Variation in Meiosis
- 10.21: Chromosomal Structural Rearrangements
- 10.22: Mitosis, Meiosis, and Sexual Reproduction
- 10.23: Introduction to Errors in Chromosome Number
- 10.24: Karyotypes
- 10.25: Common Disorders
- 10.26: Putting It Together- Cell Division
- 10.27: Assignment- Mitosis and Meiosis Worksheets

## 11: Module 9- DNA Structure and Replication


- 11.1: Why It Matters- DNA Structure and Replication
- 11.2: Introduction to Storing Genetic Information
- 11.3: Structure of DNA
- 11.4: Genetic Information
- 11.5: DNA Packaging in Eukaryotes and Prokaryotes
- 11.6: Introduction to DNA Base Pairs and Replication
- 11.7: The History of DNA
- 11.8: Basics of DNA Replication
- 11.9: Major Enzymes
- 11.10: “Proofreading” DNA
- 11.11: Telomeres
- 11.12: Introduction to DNA Mutations
- 11.13: What is a Mutation?
- 11.14: Mutations in Somatic Cells and in Gametes
- 11.15: Major Types of Mutations
- 11.16: Putting It Together- DNA Structure and Replication
- 11.17: Assignment- Which Has More DNA?


## 12: Module 10- DNA Transcription and Translation


- 12.1: Why It Matters- DNA Transcription and Translation
- 12.2: Introduction to Transcription
- 12.3: Steps of Genetic Transcription
- 12.4: RNA Polymerase
- 12.5: pre-RNA and mRNA
- 12.6: RNA Post-Translational Modification
- 12.7: Introduction to Translation
- 12.8: Steps of Translation
- 12.9: Ribosomes
- 12.10: tRNA
- 12.11: Codons
- 12.12: Protein Modifications
- 12.13: Introduction to Prokaryotic Transcription and Translation
- 12.14: Prokaryotic Transcription
- 12.15: Prokaryotic Translation
- 12.16: Introduction to the Central Dogma
- 12.17: The Central Dogma
- 12.18: Putting It Together- DNA Transcription and Translation


## 13: Module 11- Gene Expression


- 13.1: Why It Matters- Gene Expression
- 13.2: Introduction to Regulation of Gene Expression
- 13.3: Expression of Genes
- 13.4: Prokaryotic and Eukaryotic Gene Regulation
- 13.5: Introduction to Prokaryotic Gene Regulation
- 13.6: Gene Regulation in Prokaryotes
- 13.7: Repressors
- 13.8: Activators and Inducers

- 13.9: Prokaryotic Gene Regulation at Work
- 13.10: Introduction to Eukaryotic Gene Regulation
- 13.11: Eukaryotic Epigenetic Gene Regulation
- 13.12: Eukaryotic Transcription Gene Regulation
- 13.13: Post-Transcriptional Control of Gene Expression
- 13.14: Putting It Together- Gene Expression
- 13.15: Assignment- How Mutations Work


## 14: Module 12- Trait Inheritance


- 14.1: Why It Matters- Trait Inheritance
- 14.2: Introduction to the Father of Genetics
- 14.3: Mendel's Experiments and Heredity
- 14.4: Characteristics and Traits
- 14.5: Laws of Inheritance
- 14.6: Heredity
- 14.7: Introduction to Beyond Dominance and Recessiveness
- 14.8: Non-Mendelian Inheritance
- 14.9: Non-Mendelian Punnett Squares
- 14.10: Multiple Alleles
- 14.11: Introduction to Heredity and Disease
- 14.12: Penetrance and Expressivity
- 14.13: Pedigrees and Disease
- 14.14: Pedigree Flowchart
- 14.15: Genetic Disorder and Pedigrees
- 14.16: Introduction to Genetics and the Environment
- 14.17: Epistasis
- 14.18: Polygenic Inheritance and Environmental Effects
- 14.19: Effect of the Environment
- 14.20: Pleiotropy and Human Disorders
- 14.21: Putting It Together- Trait Inheritance
- 14.22: Assignment- Dragon Genetics


## 15: Module 13- Theory of Evolution


- 15.1: Why It Matters- Theory of Evolution
- 15.2: Introduction to Charles Darwin
- 15.3: Darwin and Descent with Modification
- 15.4: Darwin and the Theory of Evolution
- 15.5: Natural Selection
- 15.6: Introduction to Evidence for Evolution
- 15.7: Physical Evidence
- 15.8: Biological Evidence
- 15.9: Misconceptions of Evolution
- 15.10: Introduction to Mutations and Evolution
- 15.11: Population Genetics
- 15.12: Selective and Environmental Pressures
- 15.13: Introduction to Phylogenetic Trees
- 15.14: Scientific Classification
- 15.15: Structure of Phylogenetic Trees
- 15.16: Limitations of Phylogenetic Trees

- 15.17: The Taxonomic Classification System
- 15.18: Putting It Together- Theory of Evolution

## 16: Module 14- Modern Biology

- 16.1: Why It Matters- Modern Biology
- 16.2: Introduction to Key Technologies
- 16.3: Manipulating Genetic Material
- 16.4: DNA Sequencing
- 16.5: Cloning
- 16.6: Genetic Engineering
- 16.7: Introduction to Biotechnology Applications
- 16.8: Medicinal Biotechnology
- 16.9: Agricultural Biotechnology
- 16.10: Biotechnology at the Cutting Edge
- 16.11: Introduction to Risks and Benefits of Genomic Science
- 16.12: Genetic Information Used for Identification
- 16.13: Medical Uses of Genetic Information
- 16.14: Genomics in Agriculture
- 16.15: Putting It Together- Modern Biology
- 16.16: Assignment- When Jabberjays Attack!

## Index

## Glossary

## Detailed Licensing

# Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

# CHAPTER OVERVIEW

## 1: Course Contents

1.1: About This Course
1.2: Course Contents at a Glance
1.3: Learning Outcomes

---

1: Course Contents is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 1.1: About This Course

This gateway biology course provides a strong foundation in the principles of biology for students majoring in Natural Science, medical and healthcare fields. It is the second of a two-course sequence. Primary topics such as the history of life, body systems, and ecology are covered in this course.

This course was developed by Lumen Learning, with contributing work from Shelli Carter. The course is based on the OpenStax textbook Biology, supplemented with relevant materials from Khan Academy and videos from multiple sources. Original practice activities were authored by Shelli Carter and Lumen Learning in the development of this course.

## What's New?

We believe in making continuous improvements to our courses in order to enhance and facilitate student learning. This newest version of the course includes a vast number of data-driven improvements to assessment questions and text content in order to better illustrate, clarify, and evaluate concepts.

## About Lumen

Lumen Learning's mission is to make great learning opportunities available to all students, regardless of socioeconomic background.

We do this by using open educational resources (OER) to create well-designed and low-cost course materials that replace expensive textbooks. Because learning is about more than affordability and access, we also apply learning science insights and efficacy research to develop learning activities that are engineered to improve subject mastery, course completion and retention.

If you'd like to connect with us to learn more about adopting this course, please Contact Us.

You can also make an appointment for OER Office Hours to connect virtually with a live Lumen expert about any question you may have.

## Contributors and Attributions

CC licensed content, Original

- About This Course. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

1.1: About This Course is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 1.2: Course Contents at a Glance



The following list shows a summary of the topics covered in this course. To see all of the course pages, visit the Table of Contents.

## Module 1: Introduction to Biology

- The Characteristics of Life
- Prokaryotes and Eukaryotes
- Taxonomy
- The Study of Biology

## Module 2: Chemistry of Life

- Atoms and Elements
- Atomic Bonds
- Functional Groups
- The pH Scale
- Chemical Reactions
- Thermodynamics

## Module 3: Important Biological Macromolecules

- Proteins
- Lipids
- Carbohydrates
- Nucleic Acids
- Comparing Biological Macromolecules

## Module 4: Cellular Structure

- Microscopy
- Organelles
- The Cytoskeleton
- The Cell Surface

## Module 5: Cell Membranes

- Structure of the Membrane
- Kinds of Transport
- Endocytosis and Exocytosis

## Module 6: Metabolic Pathways

- Redox Reactions
- Photosynthesis
- Cellular Respiration
- Fermentation

## Module 7: Cell Communication

- Signaling Molecules and Cellular Receptors
- Propagation of the Signal

- Response to the Signal
- Signaling in Single-Celled Organisms

## Module 8: Cell Division

- Chromosomes and DNA Packaging
- The Cell Cycle
- Cell Cycle Checkpoints
- Meiosis
- Genetic Diversity
- Errors in Chromosome Number

## Module 9: DNA Structure and Replication

- Storing Genetic Information
- DNA Base Pairs and Replication
- DNA Mutations

## Module 10: DNA Transcription and Translation

- Transcription
- Translation
- Prokaryotic Transcription and Translation
- The Central Dogma

## Module 11: Gene Expression

- Regulation of Gene Expression
- Prokaryotic Gene Regulation
- Eukaryotic Gene Regulation

## Module 12: Trait Inheritance

- The Father of Genetics
- Beyond Dominance and Recessiveness
- Heredity and Disease
- Genetics and the Environment

## Module 13: Theory of Evolution

- Charles Darwin
- Evidence for Evolution
- Mutations and Evolution
- Phylogenetic Trees

## Module 14: Modern Biology

- Key Technologies
- Biotechnology Applications
- Risks and Benefits of Genomic Science

## Contributors and Attributions

CC licensed content, Original

- Course Contents at a Glance. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Binoculars Icon. **Authored by**: Musmellow. **Provided by**: Noun Project. **Located at**: https://thenounproject.com/search/?q=binoculars&i=1234056. **License**: *CC BY: Attribution*

1.2: Course Contents at a Glance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 1.3: Learning Outcomes



The content, assignments, and assessments for Biology for Majors I are aligned to the following learning outcomes. Click here for a more detailed list of outcomes.

## Module 1: Introduction to Biology

### Define biology and apply its principles

- List the defining characteristics of biological life
- Identify the different kinds of cells that make up different kinds of organisms
- Describe classification and organizational tools biologists use, including modern taxonomy
- Describe biology as a science and identify the key components of scientific inquiry

## Module 2: Chemistry of Life

### Identify the principles of chemistry that are integral to biology

- Identify the building blocks of matter
- Classify different types of atomic bonds
- Categorize molecules according to their functional group
- Demonstrate familiarity with the pH scale
- Understand basic chemical reactions
- Demonstrate familiarity with the first and second laws of thermodynamics

## Module 3: Important Biological Macromolecules

### Identify and describe the main features of the four main classes of important biological macromolecules

- Describe the structure and function of proteins
- Illustrate different types of lipids and relate their structure to their role in biological systems
- Summarize the roles carbohydrates play in biological systems
- Discuss nucleic acids and the role they play in DNA and RNA
- Discuss macromolecules and the differences between the four classes

## Module 4: Cellular Structure

### Identify and explain a variety of cellular components

- Understand why and how the light microscope and electron microscope are used in biology
- Identify membrane-bound organelles found in eukaryotic cells
- Demonstrate familiarity with various components of the cytoskeleton, including monomeric units
- Demonstrate familiarity with various cell surface specializations

## Module 5: Cell Membranes

### Describe and explain the structure and function of membranes

- Describe the structure and function of membranes, especially the phospholipid bilayer
- Distinguish between passive and active transport; explain how substances are directly transported across a membrane
- Describe the primary mechanisms by which cells import and export macromolecules

## Module 6: Metabolic Pathways

### Explain the metabolic pathways involved in the capture and release of energy in cells

- Understand the role movement of electrons plays in energy exchanges in cells
- Identify the basic components and steps of photosynthesis
- Identify the reactants and products of cellular respiration and where these reactions occur in a cell
- Illustrate the basic components and steps of fermentation

## Module 7: Cell Communication

### Identify the main components of a signal transduction pathway

- Differentiate between different types of signals
- Describe how a cell propagates a signal
- Describe how a cell responds to a signal
- Describe how a cell responds to a signal

## Module 8: Cell Division

### Describe and explain the various stages of cell division

- Understand chromosome structure and organization in eukaryotic cells
- Identify the stages of the cell cycle, by picture and by description of major milestones
- Identify and explain the important checkpoints that a cell passes through during the cell cycle
- Identify the stages of meiosis by picture and by description of major milestones; explain why meiosis involves two rounds of nuclear division
- Describe and explain a range of mechanisms for generating genetic diversity
- Examine karyotypes and identify the effects of significant changes in chromosome number

## Module 9: DNA Structure and Replication

### Relate DNA structure to the process of DNA replication

- Explain how DNA stores genetic information
- Explain the role of complementary base pairing in the precise replication process of DNA
- Recognize the impact of DNA mutations

## Module 10: DNA Transcription and Translation

### Describe the conversion of DNA to RNA to proteins

- Outline the process of genetic transcription
- Summarize the process of genetic translation
- Outline the process of prokaryotic transcription and translation
- Identify the central dogma of life

## Module 11: Gene Expression

### Explain the regulation of gene expression

- Define the term regulation as it applies to genes
- Discuss different components of prokaryotic gene regulation
- Discuss different components and types of epigenetic gene regulation

## Module 12: Trait Inheritance

### Complete monohybrid and dihybrid crosses and family pedigrees, and explain the inheritance of various traits

- Identify the impact of Gregor Mendel on the field of genetics and apply Mendel's two laws of genetics
- Explain complications to the phenotypic expression of genotype, including mutations

- Explain the conventions of a family pedigree and predict whether a disease will be passed through a family in one of three modes
- Discuss the role environment plays on phenotypes

## Module 13: Theory of Evolution

### Explain the theory of evolution, which documents the change in the genetic makeup of a biological population over time

- Describe the work of Charles Darwin in the Galapagos Islands, especially his discovery of natural selection in finch populations
- Describe how the theory of evolution by natural selection is supported by evidence
- Recognize that mutations are the basis of microevolution; and that adaptations enhance the survival and reproduction of individuals in a population
- Read and analyze a phylogenetic tree that documents evolutionary relationships

## Module 14: Modern Biology

### Describe and discuss techniques used in modern biology

- List key technologies enabling modern uses of biology
- Identify societal uses of biotechnology
- Discuss the risks and benefits involved in applications of genetic and genomic science

## Contributors and Attributions

CC licensed content, Original

- Learning Outcomes. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Magnify. **Authored by**: Eucalyp. **Provided by**: Lumen Learning. **Located at**: https://thenounproject.com/term/magnify/1276779/. **License**: *CC BY: Attribution*

---

1.3: Learning Outcomes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW


## 2: Faculty Resources


2.1: Faculty Resources Overview
2.2: Pacing
2.3: PDF
2.4: PowerPoints
2.5: Assignments
2.6: Assignments with Samples
2.7: Question Banks
2.8: I Need Help


---

2: Faculty Resources is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.1: Faculty Resources Overview



We've seen overwhelming demand for high quality, openly-licensed course materials, including supplemental resources to enrich teaching and learning and to make life easier for instructors. To support this need, we've developed and curated faculty resources to use with this course.

## Free and Open Supplemental Materials

On the following pages, you will find supplemental resources that are freely available to use with the interactive learning materials for this course. Since these resources are openly licensed, you may use them as is or adapt them to your needs.

## Additional Faculty Resources

Additional supplemental resources, learning tools, and support services are available to faculty who adopt Waymaker, OHM, or Candela courses with paid support from Lumen Learning. For many courses, these include some combination of summative assessments, answer keys, solutions manuals, or other materials shared only with authorized instructors in order to protect academic integrity.

Click here to learn more about additional instructor tools and resources available to faculty who adopt Lumen-supported courseware. Information about pricing and payment options is available on this page. Lumen's low-cost support fees replace the cost of expensive textbooks and may be paid by students or by the institution directly.

## Continuously Improving Learning Materials

Are you interested in collaborating with us to make these course materials better? We use learning data to identify where content improvements are needed, and then we invite faculty and subject matter experts to work with us developing continuous improvements aimed at increasing learning.

Learn more from this blog post, or sign up here to join our continuous improvement mailing list and stay up to date about upcoming OER hackathons and other continuous improvement activities.

## Contributors and Attributions

CC licensed content, Original

- Faculty Resources Overview. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Books Icon. **Authored by**: Ilham Fitrotul Hayat. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/books/2792694/. **License**: *CC BY: Attribution*

2.1: Faculty Resources Overview is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.2: Pacing



The Biology for Majors I textbook contains fourteen modules—roughly one chapter per week for a 16-week semester. If you need to modify the pace and cover the material more quickly, the following chapters work well together:

- **Module 2: Chemistry of Life** and **Module 3: Important Biological Macromolecules**
- **Module 4: Cellular Structure** and **Module 5: Cell Membranes**
- **Module 9: DNA Structure and Replication** and **Module 10: DNA Transcription and Translation**
- **Module 11: Gene Expression** and **Module 12: Trait Inheritance**
- **Module 13: Theory of Evolution** and **Module 14: Modern Biology**

We recommend NOT doubling up **Module 6: Metabolic Pathways**, because it is long and especially challenging. Students will need more time for mastery and completion of assignments. In a longer semester, it can be taught over two weeks instead of one.

## Contributors and Attributions

CC licensed content, Original

- Pacing. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Stopwatch. **Authored by**: Adrien Coquet, FR. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/stopwatch/1094680/. **License**: *CC BY: Attribution*

2.2: Pacing is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.3: PDF



Lumen makes significant investments to ensure our digital courseware is accessible, allowing students to learn using superior interactivity, multimedia, and a variety of accommodations for individuals with varying abilities.

PDFs offer an inferior learning experience compared to the richness and interactivity in our digital courseware. A PDF version of the textbook is available as a print alternative. The PDF does not include interactive content such as simulations, videos, and quizzes and is not vetted for accessibility. For these reasons, **we do not recommend using the textbook in the PDF form**. The offline version should be used as a print backup rather than as the primary textbook.

- Download the course PDF (58.2 MB)

To share these files with your students, copy and paste the text and download link above into a page or announcement in your learning management system (Blackboard, Canvas, etc.).

## Contributors and Attributions

CC licensed content, Original

- PDF. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Kindle Oasis. **Authored by**: Richard Slater. **Located at**: thenounproject.com/term/kindle-oasis/611350/. **License**: *CC BY: Attribution*

2.3: PDF is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.4: PowerPoints



A full set of PowerPoint decks is provided for download below. All decks are tightly aligned to the modules in this course. Since they are openly licensed, you are welcome to retain, reuse, revise, remix, and redistribute as desired.

These PowerPoint files are accessible. If you do revise them, make sure to follow these guidelines for creating accessible PowerPoints.

Use the following link to download all PowerPoint decks in a single .zip file (40 MB), or download each individual deck below:

- Module 1: Introduction to Biology
- Module 2: Chemistry of Life
- Module 3: Important Biological Macromolecules
- Module 4: Cellular Structure
- Module 5: Cell Membranes
- Module 6: Metabolic Pathways
- Module 7: Cell Communication
- Module 8: Cell Division
- Module 9: DNA Structure and Replication
- Module 10: DNA Transcription and Translation
- Module 11: Gene Expression
- Module 12: Trait Inheritance
- Module 13: Theory of Evolution
- Module 14: Modern Biology
-

## Contributors and Attributions

CC licensed content, Original

- PowerPoints. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Projector Screen. **Authored by**: Denis Shumaylov. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/projector-screen/1211212/. **License**: *CC BY: Attribution*

2.4: PowerPoints is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.5: Assignments



The assignments in this course are openly licensed, and are available as-is, or can be modified to suit your students' needs. Sample completed assignments are available to faculty who adopt Waymaker, OHM, or Candela courses with paid support from Lumen Learning. This approach helps us protect the academic integrity of these materials by ensuring they are shared only with authorized and institution-affiliated faculty and staff.

This course provides 10 prompts and grading rubrics for written assignments, designed to align with course outcomes. You can view them by clicking on the assignment links in course modules.

If you import this course into your learning management (Blackboard, Canvas, etc.), the assignments will automatically be loaded into the assignment tool. You can view them below or throughout the course.

| Assignment Files | Module Alignment |
|---|---|
| Biological Astronaut (download the .docx, .rtf) | After both Module 1: Introduction to Biology and Module 2: Chemistry of Life |
| Nutritionist for a Day (download the .docx, .rtf) | After Module 3: Important Biological Macromolecules |
| Cell Builder (download the .docx, .rtf) | After Module 4: Cellular Structure |
| Membranes Alive! (download the .docx, .rtf) | After Module 5: Cell Membranes |
| Observing Energy Transactions (download the .docx, .rtf) | After Module 6: Metabolic Pathways |
| Mitosis and Meiosis Internet Quests (download the .docx, .rtf) | After Module 7: Cell Division |
| Which Has More DNA? (download the .docx, .rtf) | After Module 8: DNA Structure and Replication |
| How Mutations Work (download the .docx, .rtf) | After both Module 9: DNA Transcription, Module 10: Translation, and Module 11: Gene Expression |
| Dragon Genetics (download the .docx, .rtf) | Module 12: After Trait Inheritance |
| When Jabberjays Attack! (download the .docx, .rtf) | After both Module 13: Theory of Evolution and Module 14: Modern Biology |

## Contributors and Attributions

CC licensed content, Original

- Assignments. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Pencil Cup. **Authored by**: IconfactoryTeam. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/pencil-cup/628840/. **License**: *CC BY: Attribution*

2.5: Assignments is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.6: Assignments with Samples



The assignments in this course are openly licensed, and are available as-is, or can be modified to suit your students' needs.

This course provides 10 prompts and grading rubrics for written assignments, designed to align with course outcomes.

If you import this course into your learning management (Blackboard, Canvas, etc.), the assignments will automatically be loaded into the assignment tool. You can view them below or throughout the course.

| Assignment Files | Module Alignment |
|---|---|
| Biological Astronaut (download the .docx, .rtf) | After both Module 1: Introduction to Biology and Module 2: Chemistry of Life |
| Nutritionist for a Day (download the .docx, .rtf)<br>Example completed assignment to distribute to students | After Module 3: Important Biological Macromolecules |
| Cell Builder (download the .docx, .rtf)<br>Here is an example of what students might create | After Module 4: Cellular Structure |
| Membranes Alive! (download the .docx, .rtf)<br>Example completed assignment to distribute to students | After Module 5: Cell Membranes |
| Observing Energy Transactions (download the .docx, .rtf)<br>Here is an example of what students might create | After Module 6: Metabolic Pathways |
| Mitosis and Meiosis Internet Quests (download the .docx, .rtf)<br>See this example of what students might submit | After Module 7: Cell Division |
| Which Has More DNA? (download the .docx, .rtf)<br>See the examples contained here | After Module 8: DNA Structure and Replication |
| How Mutations Work (download the .docx, .rtf)<br>This document shows an example submission<br>Please note that it only includes answers to a few questions and a single snapshot to give you an idea of how to complete this assignment. | After both Module 9: DNA Transcription, Module 10: Translation, and Module 11: Gene Expression |
| Dragon Genetics (download the .docx, .rtf)<br>Here is a worked example showing a screenshot and required text explanation | Module 12: After Trait Inheritance |
| When Jabberjays Attack! (download the .docx, .rtf)<br>Hints document to distribute to students | After both Module 13: Theory of Evolution and Module 14: Modern Biology |

## Contributors and Attributions

CC licensed content, Original

- Assignments. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Pencil Cup. **Authored by**: IconfactoryTeam. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/pencil-cup/628840/. **License**: *CC BY: Attribution*

2.6: Assignments with Samples is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.7: Question Banks



*Note: It is your responsibility to handle question banks and answer keys **securely** and appropriately to prevent them from being widely available and searchable via the Internet.*

The machine-graded assessments in this course are organized by module and are aligned to specific learning outcomes. Waymaker assess students in three locations: Show What You Know (ungraded pretests at the beginning of each module), Self-Checks (short, ungraded practice tests), and end-of-module quizzes (graded).

Faculty can edit and delete any of the quiz questions and also add new ones. To learn more about editing the assessments in Waymaker, visit the Waymaker Faculty User Guide AFTER the course package has been imported to your learning management system (Blackboard, Canvas, etc.).

## Optional Download

The Show What You Know and Self-Check questions are also available as a separate file that can be downloaded and imported into the assessment tool in your LMS. This format and extra step are needed ONLY if you want to make your own customized quiz, midterm, or final exam that combines quiz questions from multiple modules.

There are 14 question banks with a total of 620 multiple choice questions.

We provide quiz banks as QTI files, a standard format that allows you to import questions, build, and customize quizzes in most learning management systems (Canvas, Blackboard, etc.). **These files cannot be opened outside of an LMS.**

According to Lumen company policy, we do not provide quizzes in written (e.g. MS Word) form, in order to preserve the academic integrity of our quiz banks on behalf of everyone using our materials. We adopted this policy after observing how easy it is for quiz banks in written formats to find their way onto websites that promote academic cheating.

Click the name of your LMS to prompt the download of the file.

- Canvas
- Blackboard
- D2L
- Moodle

Once you've downloaded the file, visit the Quiz Imports page and click through to the directions for your LMS.

## Contributors and Attributions

CC licensed content, Original

- Quiz Banks. **Authored by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Usability Testing. **Authored by**: Maxim Basinski. **Provided by**: Noun Project . **Located at**: thenounproject.com/term/usability-testing/1290736/. **License**: *CC BY: Attribution*

2.7: Question Banks is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 2.8: I Need Help



Need more information about this course? Have questions about faculty resources? Can’t find what you’re looking for? Experiencing technical difficulties?

We’re here to help! Take advantage of the following Lumen customer-support resources:

- Check out one of Lumen’s Faculty User Guides here.
- Submit a support ticket here and tell us what you need.
- Talk and screen-share with a live human during Lumen’s OER office hours. See available times here.

## Contributors and Attributions

CC licensed content, Original

- I Need Help. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Help. **Authored by**: Michael Finney. **Provided by**: Noun Project. **Located at**: thenounproject.com/term/help/1112130/. **License**: *CC BY: Attribution*

2.8: I Need Help is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 3: Module 1- Introduction to Biology


3.1: Why It Matters- Introduction to Biology
3.2: Introduction to Characteristics of Life
3.3: Properties of Life
3.4: Introduction to Prokaryotes and Eukaryotes
3.5: Comparing Prokaryotic and Eukaryotic Cells
3.6: Levels of Organization of Living Things
3.7: Introduction to Taxonomy
3.8: The Diversity of Life
3.9: Phylogenetic Trees
3.10: Taxonomy
3.11: Introduction to the Study of Biology
3.12: The Branches of Biology
3.13: Scientific Inquiry
3.14: Experiments and Hypotheses
3.15: Interpreting Results
3.16: Putting It Together- Introduction to Biology



3: Module 1- Introduction to Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.1: Why It Matters- Introduction to Biology

## Why define biology and apply its principles?



Figure 1. The pangolin (also known as the scaly anteater) is a unique animal. It walks on just its hind legs and uses its front claws to tear open termite mounds.

One night while she was scrolling through her social media feeds, Cristina saw that her brother had linked to an article about some of the world's weirdest animals. As she paged through the article, Cristina became increasingly interested in the different features these animals had: some were eyeless, some were colorless, and others had even stranger features.

Before Cristina could dig deeper into these animals, she got a message from her cousin Samuel. He'd sent a link to an article about genetically modified foods and the dangers they inherently contain. Cristina was only halfway through reading the first paragraph of the article when Samuel sent her another article: this one lauding the paleo diet and its benefits. Cristina started to read the article, but before she got too far, she remembered that she had a paper due the next day. She made a mental note to come back to the articles from her cousin, and she bookmarked the animals article.

Though Cristina might not realize it, she's just been presented with three different biological questions. How did these animals develop such unique characteristics? Are GMOs dangerous? Are extreme diets (like the paleo diet) beneficial?

Cristina still has to come to her own conclusions and make her own choices, but having an understanding of biology will help her make the best choices she can.

### Learning Outcomes

- List the defining characteristics of biological life
- Identify the different kinds of cells that make up different kinds of organisms
- Describe classification and organizational tools biologists use, including modern taxonomy
- Describe biology as a science and identify the key components of scientific inquiry

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Introduction to Biology. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Scaly Anteater. **Authored by**: David Brossard. **Located at**: https://flic.kr/p/dqrpWk. **License**: *CC BY-SA: Attribution-ShareAlike*

3.1: Why It Matters- Introduction to Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.2: Introduction to Characteristics of Life

## List the defining characteristics of biological life

Biology is the science that studies life, but what exactly is life? This may sound like a silly question with an obvious response, but it is not always easy to define life. For example, a branch of biology called virology studies viruses, which exhibit some of the characteristics of living entities but lack others. It turns out that although viruses can attack living organisms, cause diseases, and even reproduce, they do not meet the criteria that biologists use to define life. Consequently, virologists are not biologists, strictly speaking. Similarly, some biologists study the early molecular evolution that gave rise to life; since the events that preceded life are not biological events, these scientists are also excluded from biology in the strict sense of the term.

From its earliest beginnings, biology has wrestled with these questions: What are the shared properties that make something "alive"? And once we know something is alive, how do we find meaningful levels of organization in its structure?

## What You'll Learn to Do

- List the properties of life
- Order the levels of organization of living things

### Learning Activities

The learning activities for this section include the following:

- Properties of Life
- Levels of Organization of Living Things
- Self Check: Characteristics of Life

## Contributors and Attributions

CC licensed content, Original

- Introduction to Characteristics of Life. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

3.2: Introduction to Characteristics of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.3: Properties of Life

All living organisms share several key characteristics or functions: order, sensitivity or response to the environment, reproduction, growth and development, regulation, homeostasis, and energy processing. When viewed together, these characteristics serve to define life.

## Order



Figure 1. A toad represents a highly organized structure consisting of cells, tissues, organs, and organ systems.

Organisms are highly organized, coordinated structures that consist of one or more cells. Even very simple, single-celled organisms are remarkably complex: inside each cell, atoms make up molecules; these in turn make up cell organelles and other cellular inclusions.

In multicellular organisms (Figure 1), similar cells form tissues. Tissues, in turn, collaborate to create organs (body structures with a distinct function). Organs work together to form organ systems.

## Sensitivity or Response to Stimuli



Figure 2. The leaves of this sensitive plant (*Mimosa pudica*) will instantly droop and fold when touched. After a few minutes, the plant returns to normal. (credit: Alex Lomas)

Organisms respond to diverse stimuli. For example, plants can bend toward a source of light, climb on fences and walls, or respond to touch (Figure 2). Even tiny bacteria can move toward or away from chemicals (a process called *chemotaxis*) or light (*phototaxis*). Movement toward a stimulus is considered a positive response, while movement away from a stimulus is considered a negative response.

Watch this video to see how plants respond to a stimulus—from opening to light, to wrapping a tendril around a branch, to capturing prey.

## Reproduction

Single-celled organisms reproduce by first duplicating their DNA, and then dividing it equally as the cell prepares to divide to form two new cells. Multicellular organisms often produce specialized reproductive germline cells that will form new individuals. When reproduction occurs, genes containing DNA are passed along to an organism's offspring. These genes ensure that the offspring will belong to the same species and will have similar characteristics, such as size and shape.

## Growth and Development



Figure 3. Although no two look alike, these puppies have inherited genes from both parents and share many of the same characteristics.

Organisms grow and develop following specific instructions coded for by their genes. These genes provide instructions that will direct cellular growth and development, ensuring that a species' young (Figure 3) will grow up to exhibit many of the same characteristics as its parents.

## Regulation

Even the smallest organisms are complex and require multiple regulatory mechanisms to coordinate internal functions, respond to stimuli, and cope with environmental stresses. Two examples of internal functions regulated in an organism are nutrient transport and blood flow. Organs (groups of tissues working together) perform specific functions, such as carrying oxygen throughout the body, removing wastes, delivering nutrients to every cell, and cooling the body.

## Homeostasis



Figure 4. Polar bears (*Ursus maritimus*) and other mammals living in ice-covered regions maintain their body temperature by generating heat and reducing heat loss through thick fur and a dense layer of fat under their skin. (credit: "longhorndave"/Flickr)

In order to function properly, cells need to have appropriate conditions such as proper temperature, pH, and appropriate concentration of diverse chemicals. These conditions may, however, change from one moment to the next. Organisms are able to maintain internal conditions within a narrow range almost constantly, despite environmental changes, through **homeostasis** (literally, "steady state")—the ability of an organism to maintain constant internal conditions. For example, an organism needs to regulate body temperature through a process known as thermoregulation. Organisms that live in cold climates, such as the polar bear (Figure 4), have body structures that help them withstand low temperatures and conserve body heat. Structures that aid in this type of insulation include fur, feathers, blubber, and fat. In hot climates, organisms have methods (such as perspiration in humans or panting in dogs) that help them to shed excess body heat.

## Energy Processing

All organisms use a source of energy for their metabolic activities. Some organisms capture energy from the sun and convert it into chemical energy in food (photosynthesis); others use chemical energy in molecules they take in as food (cellular respiration).



Figure 5. The California condor (*Gymnogyps californianus*) uses chemical energy derived from food to power flight. California condors are an endangered species; this bird has a wing tag that helps biologists identify the individual.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- fresh litter. **Authored by**: Magalie LAbbe. **Located at**: https://flic.kr/p/9yeYXd. **License**: *CC BY-NC: Attribution-NonCommercial*

Public domain content

- Bufo viridis/European Green Toad. **Authored by**: Ivengo(RUS). **Located at**: https://commons.wikimedia.org/wiki/File:Bufo_viridis.jpg. **License**: *Public Domain: No Known Copyright*
- Young California condor (Gymnogyps californianus) ready for flight. **Provided by**: US Fish and Wildlife Service. **Located at**: https://en.Wikipedia.org/wiki/File:California-condor.jpg. **License**: *Public Domain: No Known Copyright*

---

3.3: Properties of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.4: Introduction to Prokaryotes and Eukaryotes

## Identify the different kinds of cells that make up different kinds of organisms

There are two types of cells: prokaryotic and eukaryotic. In this section, we'll explore the similarities and differences between these two types.

## What You'll Learn to Do

- Identify features common to all cells
- Contrast the composition and size of prokaryotic and eukaryotic cells

### Learning Activities

The learning activities for this section include the following:

- Comparing Prokaryotic and Eukaryotic Cells
- Self Check: Prokaryotes and Eukaryotes

## Contributors and Attributions

CC licensed content, Original

- Introduction to Prokaryotes and Eukaryotes. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

3.4: Introduction to Prokaryotes and Eukaryotes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.5: Comparing Prokaryotic and Eukaryotic Cells

Cells fall into one of two broad categories: prokaryotic and eukaryotic. The single-celled organisms of the domains Bacteria and Archaea are classified as prokaryotes (*pro* = before; *karyon*– = nucleus). Animal cells, plant cells, fungi, and protists are eukaryotes (*eu* = true).

## Components of Prokaryotic Cells

All cells share four common components: (1) a plasma membrane, an outer covering that separates the cell's interior from its surrounding environment; (2) cytoplasm, consisting of a jelly-like region within the cell in which other cellular components are found; (3) DNA, the genetic material of the cell; and (4) ribosomes, particles that synthesize proteins. However, prokaryotes differ from eukaryotic cells in several ways.





Figure 1. This figure shows the generalized structure of a prokaryotic cell.

A **prokaryotic cell** is a simple, single-celled (unicellular) organism that lacks a nucleus, or any other membrane-bound organelle. We will shortly come to see that this is significantly different in eukaryotes. Prokaryotic DNA is found in the central part of the cell: a darkened region called the nucleoid (Figure 1).

Unlike Archaea and eukaryotes, bacteria have a cell wall made of peptidoglycan, comprised of sugars and amino acids, and many have a polysaccharide capsule (Figure 1). The cell wall acts as an extra layer of protection, helps the cell maintain its shape, and prevents dehydration. The capsule enables the cell to attach to surfaces in its environment. Some prokaryotes have flagella, pili, or fimbriae. Flagella are used for locomotion, while most pili are used to exchange genetic material during a type of reproduction called conjugation.

## Eukaryotic Cells

In nature, the relationship between form and function is apparent at all levels, including the level of the cell, and this will become clear as we explore eukaryotic cells. The principle "form follows function" is found in many contexts. It means that, in general, one can deduce the function of a structure by looking at its form, because the two are matched. For example, birds and fish have streamlined bodies that allow them to move quickly through the medium in which they live, be it air or water.

A **eukaryotic cell** is a cell that has a membrane-bound nucleus and other membrane-bound compartments or sacs, called **organelles**, which have specialized functions. The word eukaryotic means "true kernel" or "true nucleus," alluding to the presence of the membrane-bound nucleus in these cells. The word "organelle" means "little organ," and, as we learned earlier, organelles have specialized cellular functions, just as the organs of your body have specialized functions.

## Cell Size

At 0.1–5.0 µm in diameter, prokaryotic cells are significantly smaller than eukaryotic cells, which have diameters ranging from 10–100 µm (Figure 2). The small size of prokaryotes allows ions and organic molecules that enter them to quickly spread to other parts of the cell. Similarly, any wastes produced within a prokaryotic cell can quickly move out. However, larger eukaryotic cells have evolved different structural adaptations to enhance cellular transport. Indeed, the large size of these cells would not be possible without these adaptations. In general, cell size is limited because volume increases much more quickly than does cell surface area. As a cell becomes larger, it becomes more and more difficult for the cell to acquire sufficient materials to support the processes inside the cell, because the relative size of the surface area across which materials must be transported declines.





Figure 2. This figure shows the relative sizes of different kinds of cells and cellular components. An adult human is shown for comparison.

## Learning Objectives

Prokaryotes are single-celled organisms of the domains Bacteria and Archaea. All prokaryotes have plasma membranes, cytoplasm, ribosomes, a cell wall, DNA, and lack membrane-bound organelles. Many also have polysaccharide capsules. Prokaryotic cells range in diameter from 0.1–5.0 µm.

Like a prokaryotic cell, a eukaryotic cell has a plasma membrane, cytoplasm, and ribosomes, but a eukaryotic cell is typically larger than a prokaryotic cell, has a true nucleus (meaning its DNA is surrounded by a membrane), and has other membrane-bound organelles that allow for compartmentalization of functions. Eukaryotic cells tend to be 10 to 100 times the size of prokaryotic cells.

## Contributors and Attributions

CC licensed content, Shared previously

- Concepts of Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/b3c1e1d2-839c-42b0-a314-e119a8aafbdd@8.10. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/b3c1e1d2-839...9a8aafbdd@8.10

3.5: Comparing Prokaryotic and Eukaryotic Cells is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.6: Levels of Organization of Living Things

Living things are highly organized and structured, following a hierarchy that can be examined on a scale from small to large. The **atom** is the smallest and most fundamental unit of matter. It consists of a nucleus surrounded by electrons. Atoms form molecules. A **molecule** is a chemical structure consisting of at least two atoms held together by one or more chemical bonds. Many molecules that are biologically important are **macromolecules**, large molecules that are typically formed by polymerization (a polymer is a large molecule that is made by combining smaller units called monomers, which are simpler than macromolecules). An example of a macromolecule is deoxyribonucleic acid (DNA) (Figure 1), which contains the instructions for the structure and functioning of all living organisms.




Figure 1. All molecules, including this DNA molecule, are composed of atoms. (credit: “brian0918″/Wikimedia Commons)

Some cells contain aggregates of macromolecules surrounded by membranes; these are called **organelles**. Organelles are small structures that exist within cells. Examples of organelles include mitochondria and chloroplasts, which carry out indispensable functions: mitochondria produce energy to power the cell, while chloroplasts enable green plants to utilize the energy in sunlight to make sugars. All living things are made of cells; the **cell** itself is the smallest fundamental unit of structure and function in living organisms. (This requirement is why viruses are not considered living: they are not made of cells. To make new viruses, they have to invade and hijack the reproductive mechanism of a living cell; only then can they obtain the materials they need to reproduce.) Some organisms consist of a single cell and others are multicellular. Cells are classified as prokaryotic or eukaryotic. **Prokaryotes** are single-celled or colonial organisms that do not have membrane-bound nuclei or organelles; in contrast, the cells of **eukaryotes** do have membrane-bound organelles and a membrane-bound nucleus.

In larger organisms, cells combine to make **tissues**, which are groups of similar cells carrying out similar or related functions. **Organs** are collections of tissues grouped together performing a common function. Organs are present not only in animals but also in plants. An **organ system** is a higher level of organization that consists of functionally related organs. Mammals have many organ systems. For instance, the circulatory system transports blood through the body and to and from the lungs; it includes organs such as the heart and blood vessels. **Organisms** are individual living entities. For example, each tree in a forest is an organism. Single-celled prokaryotes and single-celled eukaryotes are also considered organisms and are typically referred to as microorganisms.

All the individuals of a species living within a specific area are collectively called a **population**. For example, a forest may include many pine trees. All of these pine trees represent the population of pine trees in this forest. Different populations may live in the same specific area. For example, the forest with the pine trees includes populations of flowering plants and also insects and microbial populations. A **community** is the sum of populations inhabiting a particular area. For instance, all of the trees, flowers, insects, and other populations in a forest form the forest’s community. Keep in mind that the community level only consists of living organisms. The forest itself is an ecosystem; this is the first level that contains non-living aspects of a given area that impact the living things in that environment. An **ecosystem** consists of all the living things in a particular area together with the abiotic, non-living parts of that environment such as nitrogen in the soil or rain water. At the highest level of organization (Figure 2), the **biosphere** is the collection of all ecosystems, and it represents the zones of life on earth. It includes land, water, and even the atmosphere to a certain extent.

## Practice Question

From a single organelle to the entire biosphere, living organisms are parts of a highly structured hierarchy.





Figure 2. The biological levels of organization of living things are shown. From a single organelle to the entire biosphere, living organisms are parts of a highly structured hierarchy. (credit “organelles”: modification of work by Umberto Salvagnin; credit “cells”: modification of work by Bruce Wetzel, Harry Schaefer/ National Cancer Institute; credit “tissues”: modification of work by Kilbad; Fama Clamosa; Mikael Häggström; credit “organs”: modification of work by Mariana Ruiz Villareal; credit “organisms”: modification of work by “Crystal”/Flickr; credit “ecosystems”: modification of work by US Fish and Wildlife Service Headquarters; credit “biosphere”: modification of work by NASA)

Which of the following statements is false?

a. Tissues exist within organs, which exist within organ systems.
b. Communities exist within populations, which exist within ecosystems.
c. Organelles exist within cells, which exist within tissues.
d. Communities exist within ecosystems, which exist in the biosphere.

[reveal-answer q=”750531″]**Show Answer**[/reveal-answer]
[hidden-answer a=”750531″]Statement b is false: populations exist within communities.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

3.6: Levels of Organization of Living Things is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.7: Introduction to Taxonomy

## Describe classification and organizational tools biologists use, including modern taxonomy



Figure 1. Our planet

Viewed from space, Earth offers no clues about the diversity of life forms that reside there. The first forms of life on Earth are thought to have been microorganisms that existed for billions of years in the ocean before plants and animals appeared. The mammals, birds, and flowers so familiar to us are all relatively recent, originating 130 to 200 million years ago. Humans have inhabited this planet for only the last 2.5 million years, and only in the last 200,000 years have humans started looking like we do today.

When faced with the remarkable diversity of life, how do we organize the different kinds of organisms so that we can better understand them? As new organisms are discovered every day, biologists continue to seek answers to these and other questions. In this outcome, we will discuss *taxonomy*, which both demonstrates the vast diversity of life and tries to organize these organisms in a way we can understand.

## What You'll Learn to Do

- Explain the "diversity of life"
- Explain the purpose of phylogenetic trees
- Explain how relationships are indicated by the binomial naming system

### Learning Activities

The learning activities for this section include the following:

- The Diversity of Life
- Phylogenetic Trees
- Taxonomy
- Self Check: Taxonomy

## Contributors and Attributions

CC licensed content, Original

- Introduction to Taxonomy. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

Public domain content

- EPIC Earth Image. **Provided by**: NASA. **Located at**: http://www.nasa.gov/image-feature/nasa-captures-epic-earth-image. **License**: *Public Domain: No Known Copyright*

3.7: Introduction to Taxonomy is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.8: The Diversity of Life



Figure 1. Life on earth is incredibly diverse.

Biological diversity is the variety of life on earth. This includes all the different plants, animals, and microorganisms; the genes they contain; and the ecosystems they form on land and in water. Biological diversity is constantly changing. It is increased by new genetic variation and reduced by extinction and habitat degradation.

## What Is Biodiversity?

Biodiversity refers to the variety of life and its processes, including the variety of living organisms, the genetic differences among them, and the communities and ecosystems in which they occur. Scientists have identified about 1.9 million species alive today. They are divided into the six kingdoms of life shown in Figure 2. Scientists are still discovering new species. Thus, they do not know for sure how many species really exist today. Most estimates range from 5 to 30 million species.





Figure 2. Click for a larger image. Known life on earth


## Cogs and Wheels

To save every cog and wheel is the first precaution of intelligent tinkering.

—Aldo Leopold, *Round River: from the Journals of Aldo Leopold*, 1953

Leopold—often considered the father of modern ecology—would have likely found the term *biodiversity* an appropriate description of his "cogs and wheels," even though idea did not become a vital component of biology until nearly 40 years after his death in 1948.

Literally, the word *biodiversity* means the many different kinds (*diversity*) of life (*bio-*), or the number of species in a particular area.




Biologists, however, are always alert to levels of organization, and have identified three unique measures of life's variation:

- The most precise and specific measure of biodiversity is **genetic diversity or genetic variation** within a species. This measure of diversity looks at differences among individuals within a population, or at difference across different populations of the same species.
- The level just broader is **species diversity**, which best fits the literal translation of *biodiversity*: the number of different species in a particular ecosystem or on Earth. This type of diversity simply looks at an area and reports what can be found there.
- At the broadest most encompassing level, we have **ecosystem diversity**. As Leopold clearly understood, the "cogs and wheels" include not only life but also the land, sea, and air that support life. In ecosystem diversity, biologists look at the many types of functional units formed by living communities interacting with their environments.

Although all three levels of diversity are important, the term biodiversity usually refers to ***species diversity!***

**Learning Objectives**

Watch this discussion about biodiversity:

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for "Biodiversity from 'the Wild Classroom'" here (link opens in new window).

Biodiversity provides us with all of our food. It also provides for many medicines and industrial products, and it has great potential for developing new and improved products for the future. Perhaps most importantly, biological diversity provides and maintains a wide array of ecological "services." These include provision of clean air and water, soil, food and shelter. The quality—and the

continuation— of our life and our economy is dependent on these “services.”

Try It



Figure 2. The short-beaked echidna is endemic to Australia. This animal—along with the platypus and three other species of echidnas—is one of the five surviving species of egg-laying mammals.

The long isolation of Australia over much of the last 50 million years and its northward movement have led to the evolution of a distinct biota. Significant features of Australia’s biological diversity include:

- A high percentage of endemic species (that is, they occur nowhere else):
  - over 80% of flowering plants
  - over 80% of land mammals
  - 88% of reptiles
  - 45% of birds
  - 92% of frogs
- Wildlife groups of great richness. Australia has an exceptional diversity of lizards in the arid zone, many ground orchids, and a total invertebrate fauna estimated at 200,000 species with more than 4,000 different species of ants alone. Marsupials and monotremes collectively account for about 56% of native terrestrial mammals in Australia.
- Wildlife of major evolutionary importance. For example, Australia has 12 of the 19 known families of primitive flowering plants, two of which occur nowhere else. Some species, such as the Queensland lungfish and peripatus, have remained relatively unchanged for hundreds of millions of years.

## Contributors and Attributions

CC licensed content, Original

- Revision and adaptation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-NC: Attribution-NonCommercial*

CC licensed content, Shared previously

- Biodiversity. **Provided by**: CK-12. **Located at**: http://www.ck12.org/biology/Biodiversity/lesson/Biodiversity-BIO/?referrer=featured_content. **License**: *CC BY-NC: Attribution-NonCommercial*
- Conserving Australia's biological diversity. **Provided by**: Australian Government, Department of the Environment and Energy. **Located at**: www.environment.gov.au/sustainability/education/publications/conserving-australias-biological-diversity-teachers-notes. **License**: *CC BY: Attribution*
- Long-beaked Echidna. **Authored by**: Jaganath. **Located at**: https://commons.wikimedia.org/wiki/File:Long-beakedEchidna.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Biodiversity Diagram. **Provided by**: Alberta Biodiversity Monitoring Institute (ABMI). **Located at**: www.abmi.ca/home/biodiversity/what-is-biodiversity.html. **License**: *CC BY: Attribution*

All rights reserved content

- Biodiversity - from The Wild Classroom. **Authored by**: Rob & Jonas Filmmaking Tips. **Located at**: https://youtu.be/vGxJArebKoc. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

3.8: The Diversity of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.9: Phylogenetic Trees

In scientific terms, the evolutionary history and relationship of an organism or group of organisms is called phylogeny. **Phylogeny** describes the relationships of one organism to others—such as which organisms it is thought to have evolved from, which species it is most closely related to, and so forth. Phylogenetic relationships provide information on shared ancestry but not necessarily on how organisms are similar or different.

Scientists use a tool called a phylogenetic tree to show the evolutionary pathways and connections among organisms. A **phylogenetic tree** is a diagram used to reflect evolutionary relationships among organisms or groups of organisms. Scientists consider phylogenetic trees to be a hypothesis of the evolutionary past since one cannot go back to confirm the proposed relationships. In other words, a "tree of life" can be constructed to illustrate when different organisms evolved and to show the relationships among different organisms (Figure 1).




Figure 1. Both of these phylogenetic trees shows the relationship of the three domains of life—Bacteria, Archaea, and Eukarya—but the (a) rooted tree attempts to identify when various species diverged from a common ancestor while the (b) unrooted tree does not. (credit a: modification of work by Eric Gaba)

A phylogenetic tree can be read like a map of evolutionary history. Many phylogenetic trees have a single lineage at the base representing a common ancestor. Scientists call such trees rooted, which means there is a single ancestral lineage (typically drawn from the bottom or left) to which all organisms represented in the diagram relate. Notice in the rooted phylogenetic tree that the three domains—Bacteria, Archaea, and Eukarya—diverge from a single point and branch off. The small branch that plants and animals (including humans) occupy in this diagram shows how recent and minuscule these groups are compared with other organisms. Unrooted trees don't show a common ancestor but do show relationships among species.

## Carl Woese and the Phylogenetic Tree

In the past, biologists grouped living organisms into five kingdoms: animals, plants, fungi, protists, and bacteria. The organizational scheme was based mainly on physical features, as opposed to physiology, biochemistry, or molecular biology, all of which are used by modern systematics. The pioneering work of American microbiologist Carl Woese in the early 1970s has shown, however, that life on Earth has evolved along three lineages, now called domains—Bacteria, Archaea, and Eukarya. The first two are prokaryotic groups of microbes that lack membrane-enclosed nuclei and organelles. The third domain contains the eukaryotes and includes unicellular microorganisms together with the four original kingdoms (excluding bacteria). Woese defined Archaea as a new domain, and this resulted in a new taxonomic tree (Figure 1a). Many organisms belonging to the Archaea domain live under extreme conditions and are called extremophiles. To construct his tree, Woese used genetic relationships rather than similarities based on morphology (shape).

Woese's tree was constructed from comparative sequencing of the genes that are universally distributed, present in every organism, and conserved (meaning that these genes have remained essentially unchanged throughout evolution). Woese's approach was revolutionary because comparisons of physical features are insufficient to differentiate between the prokaryotes that appear fairly similar in spite of their tremendous biochemical diversity and genetic variability (Figure 2). The comparison of homologous DNA and RNA sequences provided Woese with a sensitive device that revealed the extensive variability of prokaryotes, and which justified the separation of the prokaryotes into two domains: bacteria and archaea.




(a)



(b)



(c)



(d)

Figure 2. These organisms represent different domains. The (a) bacteria in this micrograph belong to Domain Bacteria, while the (b) extremophiles (not visible) living in this hot vent belong to Domain Archaea. Both the (c) sunflower and (d) lion are part of Domain Eukarya. (credit a: modification of work by Drew March; credit b: modification of work by Steve Jurvetson; credit c: modification of work by Michael Arrighi; credit d: modification of work by Leszek Leszcynski)

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

3.9: Phylogenetic Trees is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.10: Taxonomy

**Taxonomy** (which literally means “arrangement law”) is the science of classifying organisms to construct internationally shared classification systems with each organism placed into more and more inclusive groupings. Think about how a grocery store is organized. One large space is divided into departments, such as produce, dairy, and meats. Then each department further divides into aisles, then each aisle into categories and brands, and then finally a single product. This organization from larger to smaller, more specific categories is called a hierarchical system.

The taxonomic classification system (also called the Linnaean system after its inventor, Carl Linnaeus, a Swedish botanist, zoologist, and physician) uses a hierarchical model. Moving from the point of origin, the groups become more specific, until one branch ends as a single species. For example, after the common beginning of all life, scientists divide organisms into three large categories called a domain: Bacteria, Archaea, and Eukarya. Within each domain is a second category called a **kingdom**. After kingdoms, the subsequent categories of increasing specificity are: **phylum**, **class**, **order**, **family**, **genus**, and **species** (Figure 1).





Figure 1. The taxonomic classification system uses a hierarchical model to organize living organisms into increasingly specific categories. The common dog, Canis lupus familiaris, is a subspecies of Canis lupus, which also includes the wolf and dingo. (credit “dog”: modification of work by Janneke Vreugdenhil)

The kingdom Animalia stems from the Eukarya domain. For the common dog, the classification levels would be as shown in Figure 1. Therefore, the full name of an organism technically has eight terms. For the dog, it is: Eukarya, Animalia, Chordata, Mammalia, Carnivora, Canidae, *Canis*, and *lupus*. Notice that each name is capitalized except for species, and the genus and species names are italicized. Scientists generally refer to an organism only by its genus and species, which is its two-word scientific name, in what is called **binomial nomenclature**. Each species has a unique binomial nomenclature to allow for proper identification.

Therefore, the scientific name of the dog is *Canis lupus*.

It is important that the correct formatting (capitalization and italics) is used when calling an organism by its specific binomial.

The name at each level is also called a **taxon**. In other words, dogs are in order Carnivora. Carnivora is the name of the taxon at the order level; Canidae is the taxon at the family level, and so forth. Organisms also have a common name that people typically use, in this case, dog. Note that the dog is additionally a subspecies: the "*familiaris*" in *Canis lupus familiaris.* Subspecies are members of the same species that are capable of mating and reproducing viable offspring, but they are considered separate subspecies due to geographic or behavioral isolation or other factors.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

3.10: Taxonomy is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.11: Introduction to the Study of Biology

## Describe biology as a science and identify the key components of scientific inquiry

The scope of biology is vast. Biologists may study anything from the microscopic (Figure 1a) or submicroscopic view of a cell to ecosystems (Figure 1b) and the whole living planet.





Figure 1. (a) Formerly called blue-green algae, these cyanobacteria, shown here at 300× magnification under a light microscope, are some of Earth's oldest life forms. (b) These stromatolites along the shores of Lake Thetis in Western Australia are ancient structures formed by the layering of cyanobacteria in shallow waters.

Listening to the daily news, you will quickly realize how many aspects of biology are discussed every day. You may hear about *E. coli (Escherichia coli)* outbreaks in spinach or listeria contamination in ice cream. Or you may hear about efforts toward finding a cure for Ebola, Alzheimer's disease, and cancer. On a global scale, many researchers are committed to finding ways to protect the planet, solve environmental issues, and reduce the effects of climate change. All of these diverse endeavors are related to different facets of the discipline of biology.

Regardless of their particular focus of study, all biologists use the same methodology as they seek new discoveries: scientific inquiry.

## What You'll Learn to Do

- Identify the main branches of biology
- Describe "scientific inquiry" and identify its scope of coverage
- Form a hypothesis and use it to design a scientific experiment
- Analyze simple data and graphed results

### Learning Activities

The learning activities for this section include the following:

- The Branches of Biology
- Scientific Inquiry
- Experiments and Hypotheses
- Interpreting Results
- Self Check: The Study of Biology

## Contributors and Attributions

CC licensed content, Original

- Revision, adaptation, and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Stromatolites at Lake Thetis. **Authored by**: Ruth Ellison. **Located at**: https://flic.kr/p/ezabK. **License**: *CC BY-NC: Attribution-NonCommercial*

Public domain content

- Modification of Cyanobacteria; scale-bar data from Matt Russell. **Provided by**: NASA. **Located at**: https://commons.wikimedia.org/wiki/File:Cyanobacteria_guerrero_negro.jpg. **License**: *Public Domain: No Known Copyright*

---

3.11: Introduction to the Study of Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.12: The Branches of Biology

The scope of biology is broad and therefore contains many branches and sub-disciplines. Biologists may pursue one of those sub-disciplines and work in a more focused field. For instance, molecular biology and biochemistry study biological processes at the molecular and chemical level, including interactions among molecules such as DNA, RNA, and proteins, as well as the way they are regulated. Microbiology, the study of microorganisms, is the study of the structure and function of single-celled organisms. It is quite a broad branch itself, and depending on the subject of study, there are also microbial physiologists, ecologists, and geneticists, among others.

## Forensic Science



Figure 1. This forensic scientist works in a DNA extraction room at the U.S. Army Criminal Investigation Laboratory at Fort Gillem, GA. (credit: United States Army CID Command Public Affairs)

Forensic science is the application of science to answer questions related to the law. Biologists as well as chemists and biochemists can be forensic scientists. Forensic scientists provide scientific evidence for use in courts, and their job involves examining trace materials associated with crimes. Interest in forensic science has increased in the last few years, possibly because of popular television shows that feature forensic scientists on the job. Also, the development of molecular techniques and the establishment of DNA databases have expanded the types of work that forensic scientists can do.

Their job activities are primarily related to crimes against people such as murder, rape, and assault. Their work involves analyzing samples such as hair, blood, and other body fluids and also processing DNA (Figure 1) found in many different environments and materials.

Forensic scientists also analyze other biological evidence left at crime scenes, such as insect larvae or pollen grains. Students who want to pursue careers in forensic science will most likely be required to take chemistry and biology courses as well as some intensive math courses.

Another field of biological study, neurobiology, studies the biology of the nervous system, and although it is considered a branch of biology, it is also recognized as an interdisciplinary field of study known as neuroscience. Because of its interdisciplinary nature, this sub-discipline studies different functions of the nervous system using molecular, cellular, developmental, medical, and computational approaches.



Figure 2. Researchers work on excavating dinosaur fossils at a site in Castellón, Spain. (credit: Mario Modesto)

Paleontology, another branch of biology, uses fossils to study life's history (Figure 2). Zoology and botany are the study of animals and plants, respectively. Biologists can also specialize as biotechnologists, ecologists, or physiologists, to name just a few areas. This is just a small sample of the many fields that biologists can pursue.

Biology is the culmination of the achievements of the natural sciences from their inception to today. Excitingly, it is the cradle of emerging sciences, such as the biology of brain activity, genetic engineering of custom organisms, and the biology of evolution that uses the laboratory tools of molecular biology to retrace the earliest stages of life on earth. A scan of news headlines—whether reporting on immunizations, a newly discovered species, sports doping, or a genetically modified food—demonstrates the way biology is active in and important to our everyday world.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

3.12: The Branches of Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.13: Scientific Inquiry

Science deals with testable knowledge about physical phenomena in the universe. The goal of science is to understand how the universe works. Biology focuses on understanding living things. To gain knowledge about nature and physical phenomena, scientists use a particular approach called “scientific inquiry.”

Scientific inquiry is the best approach we have to understanding the natural world and predicting natural phenomena. Evidence for this claim can be found in the successes of science-based technologies. Take medicine, for example. Prior to the 1700s, most medical practices were based on folk traditions or on ideas promoted by religious leaders. Some of these prescientific remedies worked, but the process for discovering new treatments was a slow and haphazard system of trial and error. Ineffective treatments were often accepted simply because there was no clear procedure for evaluating them. Today, with science-based medicine and public health practices, we have gained unprecedented control over threats to our health. According to the Centers for Disease Control, the average life expectancy in the United States has increased by more than 30 years since 1900.

Scientific inquiry has not displaced faith, intuition, and dreams. These traditions and ways of knowing have emotional value and provide moral guidance to many people. But hunches, feelings, deep convictions, old traditions, or dreams cannot be accepted directly as scientifically valid. Instead, science limits itself to ideas that can be tested through verifiable observations. Supernatural claims that events are caused by ghosts, devils, God, or other spiritual entities cannot be tested in this way.

## Practice Question

Your friend sees this image of a circle of mushrooms and excitedly tells you it was caused by fairies dancing in a circle on the grass the night before. Can your friend’s explanation be studied using the process of science?



Figure 1. A fairy ring

[reveal-answer q=”665464″]**Show Answer**[/reveal-answer]
[hidden-answer a=”665464″]In theory, you might try to observe the fairies. But fairies are magical or supernatural beings. We have never observed them using any verifiable method, so scientists agree that they cannot be studied using scientific tools. Instead, science has an explanation supported by strong evidence: “fairy rings” result when a single colony of fungus spreads out into good habitat over a period of many years. The core area is clear of mushrooms because the soil nutrients have been partly depleted there. This idea can be evaluated with repeated observations over time using chemical soil tests and other verifiable measurements. [/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Scientific Inquiry. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5c2680020ca6017c03488572e0f8. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

3.13: Scientific Inquiry is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.14: Experiments and Hypotheses

Now we'll focus on the methods of scientific inquiry. Science often involves making observations and developing hypotheses. Experiments and further observations are often used to test the hypotheses.

A scientific experiment is a carefully organized procedure in which the scientist intervenes in a system to change something, then observes the result of the change. Scientific inquiry often involves doing experiments, though not always. For example, a scientist studying the mating behaviors of ladybugs might begin with detailed observations of ladybugs mating in their natural habitats. While this research may not be experimental, it is scientific: it involves careful and verifiable observation of the natural world. The same scientist might then treat some of the ladybugs with a hormone hypothesized to trigger mating and observe whether these ladybugs mated sooner or more often than untreated ones. This would qualify as an experiment because the scientist is now making a change in the system and observing the effects.

## Forming a Hypothesis

When conducting scientific experiments, researchers develop hypotheses to guide experimental design. A hypothesis is a suggested explanation that is both testable and falsifiable. You must be able to test your hypothesis, and it must be possible to prove your hypothesis true or false.

For example, Michael observes that maple trees lose their leaves in the fall. He might then propose a possible explanation for this observation: "cold weather causes maple trees to lose their leaves in the fall." This statement is testable. He could grow maple trees in a warm enclosed environment such as a greenhouse and see if their leaves still dropped in the fall. The hypothesis is also falsifiable. If the leaves still dropped in the warm environment, then clearly temperature was not the main factor in causing maple leaves to drop in autumn.

In the Try It below, you can practice recognizing scientific hypotheses. As you consider each statement, try to think as a scientist would: can I test this hypothesis with observations or experiments? Is the statement falsifiable? If the answer to either of these questions is "no," the statement is not a valid scientific hypothesis.

### Practice Questions

Determine whether each following statement is a scientific hypothesis.

Air pollution from automobile exhaust can trigger symptoms in people with asthma.

a. No. This statement is not testable or falsifiable.
b. No. This statement is not testable.
c. No. This statement is not falsifiable.
d. Yes. This statement is testable and falsifiable.

[reveal-answer q="429550″]**Show Answer**[/reveal-answer]
[hidden-answer a="429550″]d: Yes. This statement is testable and falsifiable. This could be tested with a number of different kinds of observations and experiments, and it is possible to gather evidence that indicates that air pollution is not linked with asthma.

[/hidden-answer]

Natural disasters, such as tornadoes, are punishments for bad thoughts and behaviors.

a. No. This statement is not testable or falsifiable.
b. No. This statement is not testable.
c. No. This statement is not falsifiable.
d. Yes. This statement is testable and falsifiable.

[reveal-answer q="74245″]Show Answer[/reveal-answer]
[hidden-answer a="74245″]

a: No. This statement is not testable or falsifiable. "Bad thoughts and behaviors" are excessively vague and subjective variables that would be impossible to measure or agree upon in a reliable way. The statement might be "falsifiable" if you came up with

a counterexample: a “wicked” place that was not punished by a natural disaster. But some would question whether the people in that place were really wicked, and others would continue to predict that a natural disaster was bound to strike that place at some point. There is no reason to suspect that people’s immoral behavior affects the weather unless you bring up the intervention of a supernatural being, making this idea even harder to test.

[/hidden-answer]

## Testing a Vaccine

Let’s examine the scientific process by discussing an actual scientific experiment conducted by researchers at the University of Washington. These researchers investigated whether a vaccine may reduce the incidence of the human papillomavirus (HPV). The experimental process and results were published in an article titled, “A controlled trial of a human papillomavirus type 16 vaccine.”

Preliminary observations made by the researchers who conducted the HPV experiment are listed below:

- Human papillomavirus (HPV) is the most common sexually transmitted virus in the United States.
- There are about 40 different types of HPV. A significant number of people that have HPV are unaware of it because many of these viruses cause no symptoms.
- Some types of HPV can cause cervical cancer.
- About 4,000 women a year die of cervical cancer in the United States.

### Practice Question

Researchers have developed a potential vaccine against HPV and want to test it. What is the first testable hypothesis that the researchers should study?

a. HPV causes cervical cancer.
b. People should not have unprotected sex with many partners.
c. People who get the vaccine will not get HPV.
d. The HPV vaccine will protect people against cancer.

[reveal-answer q=”20917″]**Show Answer**[/reveal-answer]
[hidden-answer a=”20917″]Hypothesis A is not the best choice because this information is already known from previous studies. Hypothesis B is not testable because scientific hypotheses are not value statements; they do not include judgments like “should,” “better than,” etc. Scientific evidence certainly might support this value judgment, but a hypothesis would take a different form: “Having unprotected sex with many partners increases a person’s risk for cervical cancer.” Before the researchers can test if the vaccine protects against cancer (hypothesis D), they want to test if it protects against the virus. This statement will make an excellent hypothesis for the next study. The researchers should first test hypothesis C—whether or not the new vaccine can prevent HPV.[/hidden-answer]

## Experimental Design

You’ve successfully identified a hypothesis for the University of Washington’s study on HPV: People who get the HPV vaccine will not get HPV.

The next step is to design an experiment that will test this hypothesis. There are several important factors to consider when designing a scientific experiment. First, scientific experiments must have an experimental group. This is the group that receives the experimental treatment necessary to address the hypothesis.

The experimental group receives the vaccine, but how can we know if the vaccine made a difference? Many things may change HPV infection rates in a group of people over time. To clearly show that the vaccine was effective in helping the experimental group, we need to include in our study an otherwise similar control group that does not get the treatment. We can then compare the two groups and determine if the vaccine made a difference. The control group shows us what happens in the absence of the factor under study.

However, the control group cannot get “nothing.” Instead, the control group often receives a placebo. A *placebo* is a procedure that has no expected therapeutic effect—such as giving a person a sugar pill or a shot containing only plain saline solution with no drug.

Scientific studies have shown that the “placebo effect” can alter experimental results because when individuals are told that they are or are not being treated, this knowledge can alter their actions or their emotions, which can then alter the results of the experiment.

Moreover, if the doctor knows which group a patient is in, this can also influence the results of the experiment. Without saying so directly, the doctor may show—through body language or other subtle cues—his or her views about whether the patient is likely to get well. These errors can then alter the patient’s experience and change the results of the experiment. Therefore, many clinical studies are “double blind.” In these studies, neither the doctor nor the patient knows which group the patient is in until all experimental results have been collected.

Both placebo treatments and double-blind procedures are designed to prevent bias. Bias is any systematic error that makes a particular experimental outcome more or less likely. Errors can happen in any experiment: people make mistakes in measurement, instruments fail, computer glitches can alter data. But most such errors are random and don’t favor one outcome over another. Patients’ belief in a treatment can make it more likely to appear to “work.” Placebos and double-blind procedures are used to level the playing field so that both groups of study subjects are treated equally and share similar beliefs about their treatment.

### Practice Questions

The scientists who are researching the effectiveness of the HPV vaccine will test their hypothesis by separating 2,392 young women into two groups: the control group and the experimental group. Answer the following questions about these two groups.

1. Which of the following groups most likely represents the control group?
    a. This group is given a placebo.
    b. This group is deliberately infected with HPV.
    c. This group is given nothing.
    d. This group is given the HPV vaccine.
2. Which of the following groups most likely represents the experimental group?
    a. This group is given a placebo.
    b. This group is deliberately infected with HPV.
    c. This group is given nothing.
    d. This group is given the HPV vaccine.

[reveal-answer q=”918962″]**Show Answers**[/reveal-answer]
[hidden-answer a=”918962″]

1. a: This group is given a placebo. A placebo will be a shot, just like the HPV vaccine, but it will have no active ingredient. It may change peoples’ thinking or behavior to have such a shot given to them, but it will not stimulate the immune systems of the subjects in the same way as predicted for the vaccine itself.
2. d: This group is given the HPV vaccine. The experimental group will receive the HPV vaccine and researchers will then be able to see if it works, when compared to the control group.

[/hidden-answer]

## Experimental Variables

A variable is a characteristic of a subject (in this case, of a person in the study) that can vary over time or among individuals. Sometimes a variable takes the form of a category, such as male or female; often a variable can be measured precisely, such as body height. Ideally, only one variable is different between the control group and the experimental group in a scientific experiment. Otherwise, the researchers will not be able to determine which variable caused any differences seen in the results. For example, imagine that the people in the control group were, on average, much more sexually active than the people in the experimental group. If, at the end of the experiment, the control group had a higher rate of HPV infection, could you confidently determine why? Maybe the experimental subjects were protected by the vaccine, but maybe they were protected by their low level of sexual contact.

To avoid this situation, experimenters make sure that their subject groups are as similar as possible in all variables except for the variable that is being tested in the experiment. This variable, or factor, will be deliberately changed in the experimental group. The one variable that is different between the two groups is called the independent variable. An independent variable is known or hypothesized to cause some outcome. Imagine an educational researcher investigating the effectiveness of a new teaching strategy in a classroom. The experimental group receives the new teaching strategy, while the control group receives the traditional strategy. It is the teaching strategy that is the independent variable in this scenario. In an experiment, the independent variable is the variable that the scientist deliberately changes or imposes on the subjects.

Dependent variables are known or hypothesized consequences; they are the effects that result from changes or differences in an independent variable. In an experiment, the dependent variables are those that the scientist measures before, during, and particularly at the end of the experiment to see if they have changed as expected. The dependent variable must be stated so that it is clear how it will be observed or measured. Rather than comparing "learning" among students (which is a vague and difficult to measure concept), an educational researcher might choose to compare test scores, which are very specific and easy to measure.

In any real-world example, many, many variables MIGHT affect the outcome of an experiment, yet only one or a few independent variables can be tested. Other variables must be kept as similar as possible between the study groups and are called *control variables*. For our educational research example, if the control group consisted only of people between the ages of 18 and 20 and the experimental group contained people between the ages of 30 and 35, we would not know if it was the teaching strategy or the students' ages that played a larger role in the results. To avoid this problem, a good study will be set up so that each group contains students with a similar age profile. In a well-designed educational research study, student age will be a controlled variable, along with other possibly important factors like gender, past educational achievement, and pre-existing knowledge of the subject area.

## Practice Questions

What is the independent variable in this experiment?

a. Sex (all of the subjects will be female)
b. Presence or absence of the HPV vaccine
c. Age
d. Presence or absence of HPV (the virus)

[reveal-answer q="68680"]Show Answer[/reveal-answer]
[hidden-answer a="68680"]Answer b. Presence or absence of the HPV vaccine. This is the variable that is different between the control and the experimental groups. All the subjects in this study are female, so this variable is the same in all groups. In a well-designed study, the two groups will be of similar age. The presence or absence of the virus is what the researchers will measure at the *end* of the experiment. Ideally the two groups will both be HPV-free at the start of the experiment.

[/hidden-answer]

List three control variables other than age.

[practice-area rows="3"][/practice-area]
[reveal-answer q="903121"]Show Answer[/reveal-answer]
[hidden-answer a="903121"]Some possible control variables would be: general health of the women, sexual activity, lifestyle, diet, socioeconomic status, etc.

[/hidden-answer]

What is the dependent variable in this experiment?

a. Sex (male or female)
b. Rates of HPV infection
c. Age (years)

[reveal-answer q="907103"]Show Answer[/reveal-answer]
[hidden-answer a="907103"]Answer b. Rates of HPV infection. The researchers will measure how many individuals got infected with HPV after a given period of time.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Original

- Revision and adaptation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

CC licensed content, Shared previously

- Scientific Inquiry. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5c2680020ca6017c03488572e0f8. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

---

3.14: Experiments and Hypotheses is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.15: Interpreting Results

## Gathering Data

After the experiment is completed, the data gathered are carefully interpreted. This involves the measurement of the dependent variable. In the case of our HPV experiment, remember, the dependent variable is the rate of HPV infection.

**Practice Question**

The researchers found that, of the 1,200 women in the control group, nine were infected with HPV at the end of the study. Of the 1,200 women in the experimental group, zero were infected with HPV. Does this result support the original hypothesis, that the HPV vaccine would reduce HPV infection?

[practice-area rows="2"][/practice-area]
[reveal-answer q="318356"]**Show Answer**[/reveal-answer]
[hidden-answer a="318356"]The results demonstrate that fewer individuals who got the vaccine were infected with HPV when compared to the number of infected individuals who did not get the vaccine. This supports the hypothesis that the new vaccine protects against HPV.[/hidden-answer]

## Significance

Although the HPV study suggests that the vaccine protects against infection by HPV, is the finding significant? In science, as in life, things can happen for many different reasons. A convincing study will rule out "luck" (random chance) as an explanation for the results. Strong results are said to be significant: very unlikely to occur by chance or random events.

Whether the outcome is significant often depends on the size of study; the larger the number of individuals enrolled, the more convincing the results are likely to be. For example, imagine only 10 women were enrolled in the study. In the control group, 2 in 5 of the women became infected. In the experimental group, 0 in 5 were infected. At first you might think this proves the vaccine's effectiveness, but it is NOT a convincing or significant result. Why not? Random events could easily explain the difference between the groups. For example, perhaps none of the five women in the experimental group were sexually active over the study period. They therefore stood no chance of acquiring HPV. The vaccine might appear to work, but a skeptical reader could account for the results by proposing many other scenarios.

However, imagine if the same study were done with 10,000 women, and the infection rates were 2,000 of 5,000 in the control group and zero of 5,000 in the experimental group. Random events would be spread out among a very large group of people in this study; on average, the two big groups should have similar sexual behavior and other factors influencing infection rates. If there is a big difference at the end of the study, it is very unlikely that *this* result occurred by random chance.

Statistical analyses did support the significance of the HPV vaccine result.

**Practice Question**

Researchers reported significant results from the HPV vaccine experiment. If the results had NOT been significant, what would this mean?

[practice-area rows="2"][/practice-area]
[reveal-answer q="185159"]**Show Answer**[/reveal-answer]
[hidden-answer a="185159"]If the results from this experiment were not significant, it would mean that the number of women who did not get HPV after being vaccinated was the same number that would be seen just from random chance.[/hidden-answer]

After the results are interpreted and conclusions are drawn, researchers often return to their work and begin asking further questions. In this way, scientific inquiry is a powerful tool for exploration.

Practice Question

What would be a good next question the HPV vaccine researchers may want to test? (*More than one answer is correct.*)

a. At what age is this vaccine most effective (what age group should be vaccinated)?
b. Does the HPV vaccine protect males as well as females?
c. Does protecting against HPV protect against cancer?
d. Should young girls be forced to get the vaccine against their parents' wishes?

[reveal-answer q="689643"]**Show Answers**[/reveal-answer]
[hidden-answer a="689643"]


Questions a, b, anc c are the best questions. These questions are currently being studied.

Although question d is important and has generated a lot of political heat, it is not a scientific question. Scientists can predict what will happen if girls are or are not vaccinated, but cannot decide whether disease risk or parental authority is more important.

[/hidden-answer]

## Graphing Data

Watch this ten-minute video about simple graphing:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=76

You can view the transcript for "A Beginner's Guide to Graphing Data" here (link opens in new window).

## Contributors and Attributions

CC licensed content, Original

- Revision and adaptation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

CC licensed content, Shared previously

- Scientific Inquiry. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5c2680020ca6017c03488572e0f8. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

All rights reserved content

- A Beginner's Guide to Graphing Data. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/9BkbYeTC6Mo. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

3.15: Interpreting Results is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 3.16: Putting It Together- Introduction to Biology

Biology is the study of life. As we've learned, this field covers a broad scope of subjects. As you progress through this course, you'll gain the knowledge you need to make informed decisions. Let's think back to the articles Cristina encountered at the beginning of this module: how could a knowledge of biological principle help with her understanding of each article?

## Think about It

Cristina read an article about some of the world's weirdest animals. What biological principle would help her understand why these animals have the features they do?

[practice-area rows="4"][/practice-area]
[reveal-answer q="775039"]See Our Thoughts[/reveal-answer]
[hidden-answer a="775039"]By learning about evolution and natural selection, Cristina could begin to see the different evolutionary reasons behind the extreme features some animals have. For example, the aye-aye, a mammal native to Madagascar, has evolved to have an extra long middle finger, which it can use to dig grubs out of trees. We'll learn more about how these types of traits are selected for in the module "Theory of Evolution."

[/hidden-answer]

The next article Cristina read talked about GMOs (genetically modified organisms) and the risks they have. What biological principle would help her understand the potential risks of genetically modified foods?

[practice-area rows="4"][/practice-area]
[reveal-answer q="781328"]See Our Thoughts[/reveal-answer]
[hidden-answer a="781328"]In order to understand potential risks of GMOs, Cristina will need to first understand the science behind these foods. How are GMOs created? We'll learn more about this in the module "Modern Biology."

[/hidden-answer]

Cristina then looked at an article about the paleo diet. What biological principle would help her understand the pros and cons of an extreme diet?

[practice-area rows="4"][/practice-area]
[reveal-answer q="781326"]See Our Thoughts[/reveal-answer]
[hidden-answer a="781326"]In order to survive, humans require specific nutrients. While we won't get into too much depth about nutrition in this course, we will learn about different biological macromolecules. These macromolecules include categories you may recognize, like proteins, lipids (more commonly called fats), and carbohydrates. In that module, we'll learn about the roles these molecules play in our bodies—we'll learn the essential functions they perform. With this knowledge, Cristina could better decide what she should or shouldn't remove from her diet.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: Introduction to Biology. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

3.16: Putting It Together- Introduction to Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 4: Module 2- Chemistry of Life


4.1: Why It Matters- The Chemistry of Life
4.2: Introduction to Atoms and Elements
4.3: Atoms
4.4: The Periodic Table of Elements
4.5: Electrons
4.6: Introduction to Atomic Bonds
4.7: Ionic Bonds
4.8: Covalent Bonds
4.9: Hydrogen Bonds
4.10: Introduction to Functional Groups
4.11: Functional Groups
4.12: Introduction to The pH Scale
4.13: Acids and Bases
4.14: Buffers
4.15: Introduction to Chemical Reactions
4.16: Chemical Reactions and Molecules
4.17: Anabolic and Catabolic Pathways
4.18: Enzymes
4.19: Introduction to Thermodynamics
4.20: Open and Closed Systems
4.21: The First Law of Thermodynamics
4.22: The Second Law of Thermodynamics
4.23: Putting It Together- Chemistry of Life
4.24: Assignment- Biological Astronaut
4.25: Assignment- Interacting with Attributes of Molecules



4: Module 2- Chemistry of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.1: Why It Matters- The Chemistry of Life

## Why identify the principles of chemistry that are integral to biology?

You may have heard the phrase “carbon-based life” when people discuss life on earth. Carbon is an element—one of the basic substances everything is made up of—but just what it mean to say life is “carbon-based”?

In order to understand this concept, we’ll need to understand more about elements, which means learning about chemistry. While chemistry is a separate field, its principles form the basis of biology. As you learned in the last module, all living things demonstrate hierarchical organization. You cannot truly understand the higher levels of organization (such as organisms or ecosystems) without understanding their component parts (like cells, molecules, and atoms).

Professionals who use this chemistry in their daily work include nutritionists, healthcare workers (especially when prescribing and administering medications), geneticists, and pharmacologists.

**Try It**

Nutritionists are often responsible for planning healthy menus and meal plans, for example in schools or daycares. They may advise individuals on changes in diet in order to achieve a particular health goal. Why do you think they might need to know about chemistry?

### Learning Outcomes

- Identify the building blocks of matter
- Classify different types of atomic bonds
- Categorize molecules according to their functional group
- Demonstrate familiarity with the pH scale
- Understand basic chemical reactions
- Demonstrate familiarity with the first and second laws of thermodynamics

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: The Chemistry of Life. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

4.1: Why It Matters- The Chemistry of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.2: Introduction to Atoms and Elements

## Identify the building blocks of matter

At its most fundamental level, life is made up of matter. **Matter** is any substance that occupies space and has mass. **Elements** are unique forms of matter with specific chemical and physical properties that cannot be broken down into smaller substances by ordinary chemical reactions. There are 118 elements, but only 92 occur naturally. The remaining elements are synthesized in laboratories and are unstable.

Each element is designated by its chemical symbol, which is a single capital letter or, when the first letter is already "taken" by another element, a combination of two letters. Some elements follow the English term for the element, such as C for carbon and Ca for calcium. Other elements' chemical symbols derive from their Latin names; for example, the symbol for sodium is Na, referring to *natrium*, the Latin word for sodium.

The four elements common to all living organisms are oxygen (O), carbon (C), hydrogen (H), and nitrogen (N). In the non-living world, elements are found in different proportions, and some elements common to living organisms are relatively rare on the earth as a whole, as shown in Table 1. For example, the atmosphere is rich in nitrogen and oxygen but contains little carbon and hydrogen, while the earth's crust, although it contains oxygen and a small amount of hydrogen, has little nitrogen and carbon. In spite of their differences in abundance, all elements and the chemical reactions between them obey the same chemical and physical laws regardless of whether they are a part of the living or non-living world.

Table 1. Approximate Percentage of Elements in Living Organisms (Humans) Compared to the Non-living World

| Element | Life (Humans) | Atmosphere | Earth's Crust |
|---|---|---|---|
| Oxygen (O) | 65% | 21% | 46% |
| Carbon (C) | 18% | trace | trace |
| Hydrogen (H) | 10% | trace | 0.1% |
| Nitrogen (N) | 3% | 78% | trace |

## What You'll Learn to Do

- Draw a diagram of an atom, according to current scientific understanding
- Understand the periodic table of elements and how to use it to understand elements
- Describe the behavior and location of electrons, and how these factors influence bond formation between atoms

### Learning Activities

The learning activities for this section include the following:

- Atoms
- The Periodic Table of Elements
- Electrons
- Self Check: Atoms and Elements

## Contributors and Attributions

CC licensed content, Original

- Introduction to Atoms and Elements. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.2: Introduction to Atoms and Elements is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.3: Atoms

## Structure of the Atom

To understand how elements come together, we must first discuss the smallest component or building block of an element, the atom. An **atom** is the smallest unit of matter that retains all of the chemical properties of an element. For example, one gold atom has all of the properties of gold in that it is a solid metal at room temperature. A gold coin is simply a very large number of gold atoms molded into the shape of a coin and containing small amounts of other elements known as impurities. Gold atoms cannot be broken down into anything smaller while still retaining the properties of gold.

An atom is composed of two regions: the **nucleus**, which is in the center of the atom and contains protons and neutrons, and the outermost region of the atom which holds its electrons in orbit around the nucleus, as illustrated in Figure 1. Atoms contain protons, electrons, and neutrons, among other subatomic particles. The only exception is hydrogen (H), which is made of one proton and one electron with no neutrons.





Figure 1. Elements, such as helium, depicted here, are made up of atoms. Atoms are made up of protons and neutrons located within the nucleus, with electrons in orbitals surrounding the nucleus.

Protons and neutrons have approximately the same mass, about $1.67 \times 10^{-24}$ grams. Scientists arbitrarily define this amount of mass as one atomic mass unit (amu) or one Dalton, as shown in Table 1. Although similar in mass, protons and neutrons differ in their electric charge. A **proton** is positively charged whereas a **neutron** is uncharged. Therefore, the number of neutrons in an atom contributes significantly to its mass, but not to its charge. **Electrons** are much smaller in mass than protons, weighing only $9.11 \times 10^{-28}$ grams, or about 1/1800 of an atomic mass unit. Hence, they do not contribute much to an element's overall atomic mass. Therefore, when considering atomic mass, it is customary to ignore the mass of any electrons and calculate the atom's mass based on the number of protons and neutrons alone. Although not significant contributors to mass, electrons do contribute greatly to the atom's charge, as each electron has a negative charge equal to the positive charge of a proton. In uncharged, neutral atoms, the number of electrons orbiting the nucleus is equal to the number of protons inside the nucleus. In these atoms, the positive and negative charges cancel each other out, leading to an atom with no net charge.

Accounting for the sizes of protons, neutrons, and electrons, most of the volume of an atom—greater than 99 percent—is, in fact, empty space. With all this empty space, one might ask why so-called solid objects do not just pass through one another. The reason they do not is that the electrons that surround all atoms are negatively charged and negative charges repel each other.

Table 1. Protons, Neutrons, and Electrons

| | Charge | Mass (amu) | Location |
|---|---|---|---|
| Proton | +1 | 1 | nucleus |
| Neutron | 0 | 1 | nucleus |
| Electron | –1 | 0 | orbitals |

## Atomic Number and Mass

Atoms of each element contain a characteristic number of protons and electrons. The number of protons determines an element's **atomic number** and is used to distinguish one element from another. The number of neutrons is variable, resulting in isotopes,

which are different forms of the same atom that vary only in the number of neutrons they possess. Together, the number of protons and the number of neutrons determine an element's mass number, as illustrated in Figure 2. Note that the small contribution of mass from electrons is disregarded in calculating the mass number. This approximation of mass can be used to easily calculate how many neutrons an element has by simply subtracting the number of protons from the mass number. Since an element's isotopes will have slightly different mass numbers, scientists also determine the **atomic mass**, which is the calculated mean of the mass number for its naturally occurring isotopes. Often, the resulting number contains a fraction. For example, the atomic mass of chlorine (Cl) is 35.45 because chlorine is composed of several isotopes, some (the majority) with atomic mass 35 (17 protons and 18 neutrons) and some with atomic mass 37 (17 protons and 20 neutrons).

### Practice Question

Carbon has an atomic number of six, and two stable isotopes with mass numbers of twelve and thirteen, respectively. Its atomic mass is 12.11.




Figure 2. Carbon-12 and carbon-13

How many neutrons do carbon-12 and carbon-13 have, respectively?

[practice-area rows="2"][/practice-area]
[reveal-answer q="912178"]**Show Answer**[/reveal-answer]
[hidden-answer a="912178"]Carbon-12 has six neutrons. Carbon-13 has seven neutrons.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.3: Atoms is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.4: The Periodic Table of Elements

The different elements are organized and displayed in the periodic table. Devised by Russian chemist Dmitri Mendeleev (1834–1907) in 1869, the table groups elements that, although unique, share certain chemical properties with other elements. The properties of elements are responsible for their physical state at room temperature: they may be gases, solids, or liquids. Elements also have specific chemical reactivity, the ability to combine and to chemically bond with each other.

In the periodic table, shown in Figure 1, the elements are organized and displayed according to their atomic number and are arranged in a series of rows and columns based on shared chemical and physical properties. In addition to providing the atomic number for each element, the periodic table also displays the element's atomic mass. Looking at carbon, for example, its symbol (C) and name appear, as well as its atomic number of six (in the upper left-hand corner) and its atomic mass of 12.11.





Figure 1. The periodic table shows the atomic mass and atomic number of each element. The atomic number appears above the symbol for the element and the approximate atomic mass appears below it.

The periodic table groups elements according to chemical properties. The differences in chemical reactivity between the elements are based on the number and spatial distribution of an atom's electrons. Atoms that chemically react and bond to each other form molecules. Molecules are simply two or more atoms chemically bonded together. Logically, when two atoms chemically bond to form a molecule, their electrons, which form the outermost region of each atom, come together first as the atoms form a chemical bond.

Watch this video for a more in-depth introduction to the periodic table:



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=88




## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- [redacted]**thored by**: CrashCourse. **Located at**: https://youtu.be/0RRVV4Diomg. [redacted]ard YouTube License

---

4.4: The Periodic Table of Elements is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.5: Electrons

## Electron Shells and the Bohr Model





Figure 1. The Bohr model.

It should be stressed that there is a connection between the number of protons in an element, the atomic number that distinguishes one element from another, and the number of electrons it has. In all electrically neutral atoms, the number of electrons is the same as the number of protons. Thus, each element, at least when electrically neutral, has a characteristic number of electrons equal to its atomic number.

An early model of the atom was developed in 1913 by Danish scientist Niels Bohr (1885–1962). In this model, electrons exist within principal shells. An electron normally exists in the lowest energy shell available, which is the one closest to the nucleus. Energy from a photon of light can bump it up to a higher energy shell, but this situation is unstable, and the electron quickly decays back to the ground state. In the process, a photon of light is released.

The Bohr model shows the atom as a central nucleus containing protons and neutrons, with the electrons in circular **orbitals** at specific distances from the nucleus, as illustrated in Figure 1. These orbits form electron shells or energy levels, which are a way of visualizing the number of electrons in the outermost shells. These energy levels are designated by a number and the symbol “n.” For example, 1n represents the first energy level located closest to the nucleus.

Electrons fill orbitals in a consistent order: they first fill the orbitals closest to the nucleus, then they continue to fill orbitals of increasing energy further from the nucleus. If there are multiple orbitals of equal energy, they will be filled with one electron in each energy level before a second electron is added. The electrons of the outermost energy level determine the energetic stability of the atom and its tendency to form chemical bonds with other atoms to form molecules.

Under standard conditions, atoms fill the inner shells first, often resulting in a variable number of electrons in the outermost shell. The innermost shell has a maximum of two electrons but the next two electron shells can each have a maximum of eight electrons. This is known as the **octet rule**, which states, with the exception of the innermost shell, that atoms are more stable energetically when they have eight electrons in their **valence shell**, the outermost electron shell. Examples of some neutral atoms and their electron configurations are shown in Figure 2. Notice that in this figure, helium has a complete outer electron shell, with two electrons filling its first and only shell. Similarly, neon has a complete outer 2n shell containing eight electrons. In contrast, chlorine and sodium have seven and one in their outer shells, respectively, but theoretically they would be more energetically stable if they followed the octet rule and had eight.

Practice Question





Figure 2. Bohr diagrams for hydrogen, helium, lithium, carbon, fluorine, neon, sodium, silicon, chlorine, and argon.

Bohr diagrams indicate how many electrons fill each principal shell. Group 18 elements (helium, neon, and argon are shown in Figure 2) have a full outer, or valence, shell. A full valence shell is the most stable electron configuration. Elements in other groups have partially filled valence shells and gain or lose electrons to achieve a stable electron configuration.

An atom may give, take, or share electrons with another atom to achieve a full valence shell, the most stable electron configuration. Looking at this figure, how many electrons do elements in group 1 need to lose in order to achieve a stable electron configuration? How many electrons do elements in groups 14 and 17 need to gain to achieve a stable configuration?

[practice-area rows="2"][/practice-area]
[reveal-answer q="766153"]**Show Answer**[/reveal-answer]
[hidden-answer a="766153"]Elements in group 1 need to lose one electron to achieve a stable electron configuration. Elements in groups 14 and 17 need to gain four and one electrons, respectively, to achieve a stable configuration.[/hidden-answer]

Understanding that the organization of the periodic table is based on the total number of protons (and electrons) helps us know how electrons are distributed among the outer shell. The periodic table is arranged in columns and rows based on the number of electrons and where these electrons are located. Take a closer look at the some of the elements in the periodic table's far right column in Figure 3.





Figure 3. The periodic table shows the atomic mass and atomic number of each element. The atomic number appears above the symbol for the element and the approximate atomic mass appears below it.

The group 18 atoms helium (He), neon (Ne), and argon (Ar) all have filled outer electron shells, making it unnecessary for them to share electrons with other atoms to attain stability; they are highly stable as single atoms. Their non-reactivity has resulted in their being named the **inert gases** (or **noble gases**). Compare this to the group 1 elements in the left-hand column. These elements, including hydrogen (H), lithium (Li), and sodium (Na), all have one electron in their outermost shells. That means that they can achieve a stable configuration and a filled outer shell by donating or sharing one electron with another atom or a molecule such as water. Hydrogen will donate or share its electron to achieve this configuration, while lithium and sodium will donate their electron to become stable. As a result of losing a negatively charged electron, they become positively charged **ions**. Group 17 elements, including fluorine and chlorine, have seven electrons in their outmost shells, so they tend to fill this shell with an electron from other atoms or molecules, making them negatively charged ions. Group 14 elements, of which carbon is the most important to living systems, have four electrons in their outer shell allowing them to make several covalent bonds (discussed below) with other atoms. Thus, the columns of the periodic table represent the potential shared state of these elements' outer electron shells that is responsible for their similar chemical characteristics.

## Electron Orbitals

Although useful to explain the reactivity and chemical bonding of certain elements, the Bohr model of the atom does not accurately reflect how electrons are spatially distributed surrounding the nucleus. They do not circle the nucleus like the earth orbits the sun, but are found in **electron orbitals**. These relatively complex shapes result from the fact that electrons behave not just like particles, but also like waves. Mathematical equations from quantum mechanics known as wave functions can predict within a certain level of probability where an electron might be at any given time. The area where an electron is most likely to be found is called its orbital.





Figure 4. Click for a larger image. The *s* subshells are shaped like spheres. Both the 1n and 2n principal shells have an *s* orbital, but the size of the sphere is larger in the 2n orbital. Each sphere is a single orbital. *p* subshells are made up of three dumbbell-shaped orbitals. Principal shell 2n has a *p* subshell, but shell 1 does not.

Recall that the Bohr model depicts an atom's electron shell configuration. Within each electron shell are subshells, and each subshell has a specified number of orbitals containing electrons. While it is impossible to calculate exactly where an electron is located, scientists know that it is most probably located within its orbital path. Subshells are designated by the letters *s*, *p*, *d*, and *f*. The *s* subshell is spherical in shape and has one orbital. Principal shell 1n has only a single *s* orbital, which can hold two electrons. Principal shell 2n has one *s* and one *p* subshell, and can hold a total of eight electrons. The *p* subshell has three dumbbell-shaped orbitals, as illustrated in Figure 4. Subshells *d* and *f* have more complex shapes and contain five and seven orbitals, respectively. These are not shown in the illustration. Principal shell 3n has *s*, *p*, and *d* subshells and can hold 18 electrons. Principal shell 4n has *s*, *p*, *d* and *f* orbitals and can hold 32 electrons. Moving away from the nucleus, the number of electrons and orbitals found in the energy levels increases. Progressing from one atom to the next in the periodic table, the electron structure can be worked out by fitting an extra electron into the next available orbital.

The closest orbital to the nucleus, called the 1*s* orbital, can hold up to two electrons. This orbital is equivalent to the innermost electron shell of the Bohr model of the atom. It is called the 1*s* orbital because it is spherical around the nucleus. The 1*s* orbital is the closest orbital to the nucleus, and it is always filled first, before any other orbital can be filled. Hydrogen has one electron; therefore, it has only one spot within the 1*s* orbital occupied. This is designated as $1s^1$, where the superscripted 1 refers to the one electron within the 1*s* orbital. Helium has two electrons; therefore, it can completely fill the 1*s* orbital with its two electrons. This is designated as $1s^2$, referring to the two electrons of helium in the 1*s* orbital. On the periodic table Figure 4, hydrogen and helium are the only two elements in the first row (period); this is because they only have electrons in their first shell, the 1*s* orbital. Hydrogen and helium are the only two elements that have the 1*s* and no other electron orbitals in the electrically neutral state.

The second electron shell may contain eight electrons. This shell contains another spherical *s* orbital and three "dumbbell" shaped *p* orbitals, each of which can hold two electrons, as shown in Figure 4. After the 1*s* orbital is filled, the second electron shell is filled, first filling its 2*s* orbital and then its three *p* orbitals. When filling the *p* orbitals, each takes a single electron; once each *p* orbital has an electron, a second may be added. Lithium (Li) contains three electrons that occupy the first and second shells. Two electrons fill the 1*s* orbital, and the third electron then fills the 2*s* orbital. Its **electron configuration** is $1s^2 2s^1$. Neon (Ne), on the other hand, has a total of ten electrons: two are in its innermost 1*s* orbital and eight fill its second shell (two each in the 2*s* and three *p* orbitals); thus, it is an inert gas and energetically stable as a single atom that will rarely form a chemical bond with other atoms. Larger elements have additional orbitals, making up the third electron shell. While the concepts of electron shells and orbitals are closely related, orbitals provide a more accurate depiction of the electron configuration of an atom because the orbital model specifies the different shapes and special orientations of all the places that electrons may occupy.


Watch this visual animation to see the spatial arrangement of the *p* and *s* orbitals. Note that this video has no audio.

A link to an interactive elements can be found at the bottom of this page.

You can view the audio description text for "Atomic Orbitals – Electron Configuration of Scandium" here (opens in new window).

Learning Objectives

This video gives another overview of the electron:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=90

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Atomic orbitals - electron configuration of Scandium (Z=21). **Authored by**: Manuel Moreirea Baptista. **Located at**: https://youtu.be/sMt5Dcex0kg. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License
- The Electron: Crash Course Chemistry #5. **Authored by**: CrashCourse. **Located at**: https://youtu.be/rcKilE9CdaA. **License**: *Public Domain: No Known Copyright*. **License Terms**: Standard YouTube License

4.5: Electrons is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.6: Introduction to Atomic Bonds

## Classify different types of atomic bonds

When atoms bond together, they create elements. The different types of bonds (ionic and covalent, polar and non-polar) have an impact on the elements they create. Understanding the types of bonds that create things can help us understand those things themselves.

## What You'll Learn to Do

- Describe the characteristics of ionic bonds and identify common ions
- Describe the characteristics of covalent bonds and differentiate between polar and non-polar bonds
- Model a Hydrogen bond and identify its unique qualities

### Learning Activities

The learning activities for this section include the following:

- Ionic Bonds
- Covalent Bonds
- Hydrogen Bonds
- Self Check: Atomic Bonds

## Contributors and Attributions

CC licensed content, Original

- Introduction to Atomic Bonds. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

4.6: Introduction to Atomic Bonds is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.7: Ionic Bonds

Some atoms are more stable when they gain or lose an electron (or possibly two) and form ions. This fills their outermost electron shell and makes them energetically more stable. Because the number of electrons does not equal the number of protons, each ion has a net charge. **Cations** are positive ions that are formed by losing electrons. Negative ions are formed by gaining electrons and are called anions. **Anions** are designated by their elemental name being altered to end in "-ide": the anion of chlorine is called chloride, and the anion of sulfur is called sulfide, for example.

This movement of electrons from one element to another is referred to as **electron transfer**. As Figure 1 illustrates, sodium (Na) only has one electron in its outer electron shell. It takes less energy for sodium to donate that one electron than it does to accept seven more electrons to fill the outer shell. If sodium loses an electron, it now has 11 protons, 11 neutrons, and only 10 electrons, leaving it with an overall charge of +1. It is now referred to as a sodium ion. Chlorine (Cl) in its lowest energy state (called the ground state) has seven electrons in its outer shell. Again, it is more energy-efficient for chlorine to gain one electron than to lose seven. Therefore, it tends to gain an electron to create an ion with 17 protons, 17 neutrons, and 18 electrons, giving it a net negative (–1) charge. It is now referred to as a chloride ion. In this example, sodium will donate its one electron to empty its shell, and chlorine will accept that electron to fill its shell. Both ions now satisfy the octet rule and have complete outermost shells. Because the number of electrons is no longer equal to the number of protons, each is now an ion and has a +1 (sodium cation) or –1 (chloride anion) charge. Note that these transactions can normally only take place simultaneously: in order for a sodium atom to lose an electron, it must be in the presence of a suitable recipient like a chlorine atom.






Figure 1. In the formation of an ionic compound, metals lose electrons and nonmetals gain electrons to achieve an octet. Ionic bonds are formed between ions with opposite charges. For instance, positively charged sodium ions and negatively charged chloride ions bond together to make crystals of sodium chloride, or table salt, creating a crystalline molecule with zero net charge.

**Ionic bonds** are formed between ions with opposite charges. For instance, positively charged sodium ions and negatively charged chloride ions bond together to make crystals of sodium chloride, or table salt, creating a crystalline molecule with zero net charge.

Certain salts are referred to in physiology as **electrolytes** (including sodium, potassium, and calcium), ions necessary for nerve impulse conduction, muscle contractions and water balance. Many sports drinks and dietary supplements provide these ions to replace those lost from the body via sweating during exercise.

## Video Review

This video shows how ionic compounds form from anions and cations.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=94

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Ionic Bonding. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/hiyTfhjeF_U. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

4.7: Ionic Bonds is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.8: Covalent Bonds

Another way the octet rule can be satisfied is by the sharing of electrons between atoms to form **covalent bonds**. These bonds much more common than ionic bonds in the molecules of living organisms. Covalent bonds are commonly found in carbon-based organic molecules, such as our DNA and proteins. Covalent bonds are also found in inorganic molecules like $H_2O$, $CO_2$, and $O_2$. One, two, or three pairs of electrons may be shared, making single, double, and triple bonds, respectively. The more covalent bonds between two atoms, the stronger their connection. Thus, triple bonds are the strongest.

The strength of different levels of covalent bonding is one of the main reasons living organisms have a difficult time in acquiring nitrogen for use in constructing their molecules, even though molecular nitrogen, $N_2$, is the most abundant gas in the atmosphere. Molecular nitrogen consists of two nitrogen atoms triple bonded to each other and, as with all molecules, the sharing of these three pairs of electrons between the two nitrogen atoms allows for the filling of their outer electron shells, making the molecule more stable than the individual nitrogen atoms. This strong triple bond makes it difficult for living systems to break apart this nitrogen in order to use it as constituents of proteins and DNA.

The formation of water molecules provides an example of covalent bonding. The hydrogen and oxygen atoms that combine to form water molecules are bound together by covalent bonds. The electron from the hydrogen splits its time between the incomplete outer shell of the hydrogen atoms and the incomplete outer shell of the oxygen atoms. To completely fill the outer shell of oxygen, which has six electrons in its outer shell but which would be more stable with eight, two electrons (one from each hydrogen atom) are needed: hence the well-known formula $H_2O$. The electrons are shared between the two elements to fill the outer shell of each, making both elements more stable.

View this short video to see an animation of ionic and covalent bonding.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript of “Ionic and Covalent Bonding Animation” here (opens in new window).

## Polar Covalent Bonds

There are two types of covalent bonds: polar and nonpolar. In a **polar covalent bond**, shown in Figure 1, the electrons are unequally shared by the atoms and are attracted more to one nucleus than the other. Because of the unequal distribution of electrons between the atoms of different elements, a slightly positive ($\delta+$) or slightly negative ($\delta-$) charge develops. This partial charge is an important property of water and accounts for many of its characteristics.

Water is a polar molecule, with the hydrogen atoms acquiring a partial positive charge and the oxygen a partial negative charge. This occurs because the nucleus of the oxygen atom is more attractive to the electrons of the hydrogen atoms than the hydrogen nucleus is to the oxygen’s electrons. Thus oxygen has a higher **electronegativity** than hydrogen and the shared electrons spend more time in the vicinity of the oxygen nucleus than they do near the nucleus of the hydrogen atoms, giving the atoms of oxygen and hydrogen slightly negative and positive charges, respectively. Another way of stating this is that the probability of finding a shared electron near an oxygen nucleus is more likely than finding it near a hydrogen nucleus. Either way, the atom’s relative electronegativity contributes to the development of partial charges whenever one element is significantly more electronegative than the other, and the charges generated by these polar bonds may then be used for the formation of hydrogen bonds based on the attraction of opposite partial charges. (Hydrogen bonds, which are discussed in detail below, are weak bonds between slightly positively charged hydrogen atoms to slightly negatively charged atoms in other molecules.) Since macromolecules often have atoms within them that differ in electronegativity, polar bonds are often present in organic molecules.

## Nonpolar Covalent Bonds


| | Bond type | Molecular shape | Molecular type |
|---|---|---|---|
| **Water** | δ– O H δ+ <br> Polar covalent | δ+ H H δ+ <br> O <br> δ– <br> Bent | Polar |
| **Methane** | C H <br> Nonpolar covalent | H <br> H C H <br> H <br> Tetrahedral | Nonpolar |
| **Carbon dioxide** | δ– O = C δ+ <br> Polar covalent | O = C = O <br> Linear | Nonpolar |

Figure 1. Whether a molecule is polar or nonpolar depends both on bond type and molecular shape. Both water and carbon dioxide have polar covalent bonds, but carbon dioxide is linear, so the partial charges on the molecule cancel each other out.

**Nonpolar covalent bonds** form between two atoms of the same element or between different elements that share electrons equally. For example, molecular oxygen ($O_2$) is nonpolar because the electrons will be equally distributed between the two oxygen atoms.

Another example of a nonpolar covalent bond is methane ($CH_4$), also shown in Figure 1. Carbon has four electrons in its outermost shell and needs four more to fill it. It gets these four from four hydrogen atoms, each atom providing one, making a stable outer shell of eight electrons. Carbon and hydrogen do not have the same electronegativity but are similar; thus, nonpolar bonds form. The hydrogen atoms each need one electron for their outermost shell, which is filled when it contains two electrons. These elements share the electrons equally among the carbons and the hydrogen atoms, creating a nonpolar covalent molecule.

Watch this video for another explanation of covalent bonds and how they form:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=96

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Ionic and covalent bonding animation. **Authored by**: Edu2000.org. **Located at**: https://youtu.be/QqjcCvzWwww. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License
- Covalent Bonding. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/Mo4Vfqt5v2A. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License


---

4.8: Covalent Bonds is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.9: Hydrogen Bonds

Ionic and covalent bonds between elements require energy to break. Iconic bonds are not as strong as covalent, which determines their behavior in biological systems. However, not all bonds are ionic or covalent bonds. Weaker bonds can also form between molecules. Two weak bonds that occur frequently are hydrogen bonds and van der Waals interactions. Without these two types of bonds, life as we know it would not exist. Hydrogen bonds provide many of the critical, life-sustaining properties of water and also stabilize the structures of proteins and DNA, the building block of cells.

When polar covalent bonds containing hydrogen form, the hydrogen in that bond has a slightly positive charge because hydrogen's electron is pulled more strongly toward the other element and away from the hydrogen. Because the hydrogen is slightly positive, it will be attracted to neighboring negative charges. When this happens, a weak interaction occurs between the $\delta+$ of the hydrogen from one molecule and the $\delta-$ charge on the more electronegative atoms of another molecule, usually oxygen or nitrogen, or within the same molecule. This interaction is called a **hydrogen bond**. This type of bond is common and occurs regularly between water molecules. Individual hydrogen bonds are weak and easily broken; however, they occur in very large numbers in water and in organic polymers, creating a major force in combination. Hydrogen bonds are also responsible for zipping together the DNA double helix.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=98

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Water: A Polar Molecule. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/iOOvX0jmhJ4. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

4.9: Hydrogen Bonds is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.10: Introduction to Functional Groups

## Categorize molecules according to their functional group

In this outcome, we'll learn about functional groups and the effects they have on the molecules they are bonded to.

## What You'll Learn to Do

- Identify the attributes of molecules with hydroxyl groups
- Identify the attributes of molecules with carboxyl groups
- Identify the attributes of molecules with amino groups
- Identify the attributes of molecules with phosphate groups
- Identify the attributes of molecules with methyl groups
- Identify the attributes of molecules with carbonyl groups
- Identify the attributes of molecules with sulfhydryl groups

### Learning Activities

The learning activities for this section include the following:

- Functional Groups
- Self Check: Functional Groups

## Contributors and Attributions


CC licensed content, Original

- Introduction to Functional Groups. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*


4.10: Introduction to Functional Groups is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.11: Functional Groups

Functional groups are groups of atoms that occur within organic molecules and confer specific chemical properties to those molecules. When functional groups are shown, the organic molecule is sometimes denoted as “R.” For example, ethanol is typically drawn like this:



In order to condense the structure and focus on the hydroxyl group (the oxygen and hydrogen bound to the second carbon), everything besides the hydroxyl group would replaced with an R, as follows:



**Note:** R doesn’t always stand for the same organic molecule. It can stand in for an infinite variety of molecules.

Functional groups are found along the “carbon backbone” of macromolecules which is formed by chains and/or rings of carbon atoms with the occasional substitution of an element such as nitrogen or oxygen. Molecules with other elements in their carbon backbone are substituted hydrocarbons. Each of the four types of macromolecules—proteins, lipids, carbohydrates, and nucleic acids—has its own characteristic set of functional groups that contributes greatly to its differing chemical properties and its function in living organisms.








## Properties of Functional Groups

A functional group can participate in specific chemical reactions. Some of the important functional groups in biological molecules include: hydroxyl, methyl, carbonyl, carboxyl, amino, phosphate, and sulfhydryl groups. These groups play an important role in the formation of molecules like DNA, proteins, carbohydrates, and lipids.

## Classifying Functional Groups


Functional groups are usually classified as hydrophobic or hydrophilic depending on their charge or polarity. An example of a hydrophobic group is the non-polar methane molecule. Among the hydrophilic functional groups is the carboxyl group found in amino acids, some amino acid side chains, and the fatty acid heads that form triglycerides and phospholipids. This carboxyl group ionizes to release hydrogen ions ($H^+$) from the —COOH group resulting in the negatively charged —$COO^-$ group; this contributes to the hydrophilic nature of whatever molecule it is found on. Other functional groups, such as the carbonyl group, have a partially negatively charged oxygen atom that may form hydrogen bonds with water molecules, again making the molecule more hydrophilic.

Table 1. Important Functional Groups in Biology

| Functional Group | Structure | Properties | General Features |
|---|---|---|---|
| Hydroxyl |  | Polar<br>Hydrophilic | Characterized by presence of H and O<br>Simple structure |
| Sulfhydryl |  | Polar | Characterized by presence of S<br>Simple branched structure |
| Methyl |  | Nonpolar | Characterized by presence of H and C<br>Simple structure |
| Carbonyl |  | Polar | Characterized by central C and O<br>Bound to 2 organic side groups<br>Double bond to oxygen increases the polarity |
| Carboxyl |  | Charged, ionized to release $H^+$. Since carboxyl groups can release $H^+$ ions into a solution, they are considered acidic. | Characterized by central C bound to O and OH<br>Acidic |
| Amino |  | Charged, accepts $H^+$ to form $NH_3^+$. Since amino groups can remove $H^+$ from solution, they are considered basic. | Characterized by presence of N<br>Branched structure |
| Phosphate |  | Charged, ionizes to release $H^+$. Since phosphate groups can release $H^+$ ions into solution, they are considered acidic.<br>Acidic | Characterized by presence of P<br>Complex structure |


## Practice Questions

Fructose is a common sugar that you've probably come into contact with in your life. What functional groups can be found in a fructose molecule?




[practice-area rows="2"][/practice-area]
[reveal-answer q="860203"]Show Answer[/reveal-answer]

[hidden-answer a="860203"]Fructose has hydroxyl and carbonyl groups. The hydroxyl groups are circled in red, and the carbonyl group is circled in purple:




[/hidden-answer]

Leucine is an amino acid that plays an important role in muscle development. What functional groups can be found in a leucine molecule?



[practice-area rows="2"][/practice-area]
[reveal-answer q="516947"]Show Answer[/reveal-answer]
[hidden-answer a="516947"]Leucine has carboxyl, amino, and methyl groups. The carboxyl group is circled in blue, the amnio group is circled in green, and the methyl group is circled in purple.



[/hidden-answer]

## Hydrogen Bonds between Functional Groups

Hydrogen bonds between functional groups (within the same molecule or between different molecules) are important to the function of many macromolecules and help them to fold properly into and maintain the appropriate shape for functioning.

Hydrogen bonds are also involved in various recognition processes, such as DNA complementary base pairing and the binding of an enzyme to its substrate, as illustrated in Figure 1.





Figure 1. Hydrogen bonds connect two strands of DNA together to create the double-helix structure.

## Learning Objectives

The unique properties of carbon make it a central part of biological molecules. Carbon binds to oxygen, hydrogen, and nitrogen covalently to form the many molecules important for cellular function. Carbon has four electrons in its outermost shell and can form four bonds. Carbon and hydrogen can form hydrocarbon chains or rings. Functional groups are groups of atoms that confer specific properties to hydrocarbon (or substituted hydrocarbon) chains or rings that define their overall chemical characteristics and function.

## Contributors and Attributions

CC licensed content, Original

- Revision and Adaptation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-SA: Attribution-ShareAlike*

CC licensed content, Shared previously

- Functional Groups. **Provided by**: Boundless. **Located at**: https://courses.lumenlearning.com/boundless-biology/. **License**: *CC BY-SA: Attribution-ShareAlike*

4.11: Functional Groups is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.12: Introduction to The pH Scale

## Demonstrate familiarity with the pH scale

In this outcome we'll learn about the pH scale and the importance of acids and bases.

## What You'll Learn to Do

- Identify the characteristics of acids
- Identify the characteristics of bases
- Define "buffers" and discuss the role they play in human biology

### Learning Activities

The learning activities for this section include the following:

- Acids and Bases
- Buffers
- Self Check: The pH Scale

## Contributors and Attributions


CC licensed content, Original

- Introduction to The pH Scale. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

4.12: Introduction to The pH Scale is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.13: Acids and Bases

The pH of a solution indicates its acidity or alkalinity.



**Litmus** or pH paper is filter paper that has been treated with a natural water-soluble dye so it can be used as a pH indicator, to test how much acid (acidity) or base (alkalinity) exists in a solution. You might have even used some to test whether the water in a swimming pool is properly treated. In both cases, the pH test measures the concentration of hydrogen ions in a given solution.

Hydrogen ions are spontaneously generated in pure water by the dissociation (ionization) of a small percentage of water molecules into equal numbers of hydrogen ($H^+$) ions and hydroxide ($OH^–$) ions. While the hydroxide ions are kept in solution by their hydrogen bonding with other water molecules, the hydrogen ions, consisting of naked protons, are immediately attracted to un-ionized water molecules, forming hydronium ions ($H_3O^+$). Still, by convention, scientists refer to hydrogen ions and their concentration as if they were free in this state in liquid water.

The concentration of hydrogen ions dissociating from pure water is $1 \times 10^{-7}$ moles $H^+$ ions per liter of water. Moles (mol) are a way to express the amount of a substance (which can be atoms, molecules, ions, etc), with one mole being equal to 6.02 x 1023 particles of the substance. Therefore, 1 mole of water is equal to $6.02 \times 10^{23}$ water molecules. The pH is calculated as the negative of the base 10 logarithm of this concentration. The log10 of $1 \times 10^{-7}$ is –7.0, and the negative of this number (indicated by the "p" of "pH") yields a pH of 7.0, which is also known as neutral pH. The pH inside of human cells and blood are examples of two areas of the body where near-neutral pH is maintained.

Non-neutral pH readings result from dissolving acids or bases in water. Using the negative logarithm to generate positive integers, high concentrations of hydrogen ions yield a low pH number, whereas low levels of hydrogen ions result in a high pH. An **acid** is a substance that increases the concentration of hydrogen ions ($H^+$) in a solution, usually by having one of its hydrogen atoms dissociate. A **base** provides either hydroxide ions ($OH^–$) or other negatively charged ions that combine with hydrogen ions, reducing their concentration in the solution and thereby raising the pH. In cases where the base releases hydroxide ions, these ions bind to free hydrogen ions, generating new water molecules.





Figure 1. The pH scale measures the concentration of hydrogen ions ($H^+$) in a solution.

The stronger the acid, the more readily it donates $H^+$. For example, hydrochloric acid (HCl) completely dissociates into hydrogen and chloride ions and is highly acidic, whereas the acids in tomato juice or vinegar do not completely dissociate and are considered weak acids.

Conversely, strong bases are those substances that readily donate $OH^-$ or take up hydrogen ions. Sodium hydroxide (NaOH) and many household cleaners are highly alkaline and give up $OH^-$ rapidly when placed in water, thereby raising the pH. An example of a weak basic solution is seawater, which has a pH near 8.0, close enough to neutral pH that marine organisms adapted to this saline environment are able to thrive in it.

The **pH scale** is, as previously mentioned, an inverse logarithm and ranges from 0 to 14 (Figure 1). Anything below 7.0 (ranging from 0.0 to 6.9) is acidic, and anything above 7.0 (from 7.1 to 14.0) is alkaline. Extremes in pH in either direction from 7.0 are usually considered inhospitable to life. The pH inside cells (6.8) and the pH in the blood (7.4) are both very close to neutral.

However, the environment in the stomach is highly acidic, with a pH of 1 to 2. So how do the cells of the stomach survive in such an acidic environment? How do they homeostatically maintain the near neutral pH inside them? The answer is that they cannot do it and are constantly dying. New stomach cells are constantly produced to replace dead ones, which are digested by the stomach acids. It is estimated that the lining of the human stomach is completely replaced every seven to ten days.

Watch this video for a straightforward explanation of pH and its logarithmic scale.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for “MediCastExtra pH Scale” here (opens in new window).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- PH Scale. **Authored by**: Edward Stevens. **Located at**: https://commons.wikimedia.org/wiki/File:PH_Scale.svg. **License**: *CC BY: Attribution*

All rights reserved content

- MedicCastExtra - pH Scale. **Authored by**: MedicCast TV EMS Weekly Tips. **Located at**: https://youtu.be/u837KYKyr9c. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

4.13: Acids and Bases is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.14: Buffers

So how can organisms whose bodies require a near-neutral pH ingest acidic and basic substances (a human drinking orange juice, for example) and survive? **Buffers** are the key. Buffers readily absorb excess $H^+$ or $OH^-$, keeping the pH of the body carefully maintained in the narrow range required for survival. Maintaining a constant blood pH is critical to a person's well-being. The buffer maintaining the pH of human blood involves carbonic acid ($H_2CO_3$), bicarbonate ion ($HCO_3^-$), and carbon dioxide ($CO_2$). When bicarbonate ions combine with free hydrogen ions and become carbonic acid, hydrogen ions are removed, moderating pH changes. Similarly, as shown in Figure 1, excess carbonic acid can be converted to carbon dioxide gas and exhaled through the lungs. This prevents too many free hydrogen ions from building up in the blood and dangerously reducing the blood's pH. Likewise, if too much $OH^-$ is introduced into the system, carbonic acid will combine with it to create bicarbonate, lowering the pH. Without this buffer system, the body's pH would fluctuate enough to put survival in jeopardy.





Figure 1. This diagram shows the body's buffering of blood pH levels. The blue arrows show the process of raising pH as more $CO_2$ is made. The purple arrows indicate the reverse process: the lowering of pH as more bicarbonate is created.

Other examples of buffers are antacids used to combat excess stomach acid. Many of these over-the-counter medications work in the same way as blood buffers, usually with at least one ion capable of absorbing hydrogen and moderating pH, bringing relief to those that suffer "heartburn" after eating. The unique properties of water that contribute to this capacity to balance pH—as well as water's other characteristics—are essential to sustaining life on Earth.

## In Summary: Buffers

The pH scale is a measure of acidity/alkalinity and provides information about how substances tend to act in aqueous solutions. All living processes occur in an ideal pH range. Buffers act together to keep the pH within a certain range, based off the release or absorption of hydrogen ions.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.14: Buffers is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.15: Introduction to Chemical Reactions

## Understand basic chemical reactions

In this outcome, we will discuss chemical bonds and reactions.

## What You'll Learn to Do

- Identify the components of simple chemical reactions
- Differentiate between catabolic and anabolic reactions
- Identify enzymes and their role in chemical reactions

### Learning Activities

The learning activities for this section include the following:

- Chemical Reactions and Molecules
- Anabolic and Catabolic Pathways
- Enzymes
- Self Check: Chemical Reactions

## Contributors and Attributions

CC licensed content, Original

- Introduction to Chemical Reactions. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

4.15: Introduction to Chemical Reactions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.16: Chemical Reactions and Molecules

All elements are most stable when their outermost shell is filled with electrons according to the octet rule. This is because it is energetically favorable for atoms to be in that configuration and it makes them stable. However, since not all elements have enough electrons to fill their outermost shells, atoms form **chemical bonds** with other atoms thereby obtaining the electrons they need to attain a stable electron configuration. When two or more atoms chemically bond with each other, the resultant chemical structure is a molecule. The familiar water molecule, $H_2O$, consists of two hydrogen atoms and one oxygen atom; these bond together to form water, as illustrated in Figure 1. Atoms can form molecules by donating, accepting, or sharing electrons to fill their outer shells.





Figure 1. Two or more atoms may bond with each other to form a molecule. When two hydrogens and an oxygen share electrons via covalent bonds, a water molecule is formed.

**Chemical reactions** occur when two or more atoms bond together to form molecules or when bonded atoms are broken apart. The substances used in the beginning of a chemical reaction are called the **reactants** (usually found on the left side of a chemical equation), and the substances found at the end of the reaction are known as the **products** (usually found on the right side of a chemical equation). An arrow is typically drawn between the reactants and products to indicate the direction of the chemical reaction; this direction is not always a "one-way street." For the creation of the water molecule shown above, the chemical equation would be:

$$2H + O \rightarrow H_2O$$

An example of a simple chemical reaction is the breaking down of hydrogen peroxide molecules, each of which consists of two hydrogen atoms bonded to two oxygen atoms ($H_2O_2$). The reactant hydrogen peroxide is broken down into water, containing one oxygen atom bound to two hydrogen atoms ($H_2O$), and oxygen, which consists of two bonded oxygen atoms ($O_2$). In the equation below, the reaction includes two hydrogen peroxide molecules and two water molecules. This is an example of a **balanced chemical equation**, wherein the number of atoms of each element is the same on each side of the equation. According to the law of conservation of matter, the number of atoms before and after a chemical reaction should be equal, such that no atoms are, under normal circumstances, created or destroyed.

$$H_2O_2 \text{ (hydrogen peroxide)} \rightarrow H_2O\text{(water)} + O_2 \text{ (oxygen)}$$

Even though all of the reactants and products of this reaction are molecules (each atom remains bonded to at least one other atom), in this reaction only hydrogen peroxide and water are representative of a subclass of molecules known as **compounds**: they contain atoms of more than one type of element. Molecular oxygen, on the other hand, as shown in Figure 2, consists of two doubly bonded oxygen atoms and is not classified as a compound but as an element.




Figure 2. The oxygen atoms in an $O_2$ molecule are joined by a double bond.

Some chemical reactions, such as the one shown above, can proceed in one direction until the reactants are all used up. The equations that describe these reactions contain a unidirectional arrow and are **irreversible**. **Reversible reactions** are those that can go in either direction. In reversible reactions, reactants are turned into products, but when the concentration of product goes beyond a certain threshold (characteristic of the particular reaction), some of these products will be converted back into reactants; at this point, the designations of products and reactants are reversed. This back and forth continues until a certain relative balance between reactants and products occurs—a state called **equilibrium**. These situations of reversible reactions are often denoted by a chemical equation with a double headed arrow pointing towards both the reactants and products.

For example, in human blood, excess hydrogen ions ($H^+$) bind to bicarbonate ions ($HCO_3^-$) forming an equilibrium state with carbonic acid ($H_2CO_3$). If carbonic acid were added to this system, some of it would be converted to bicarbonate and hydrogen ions.

$$HCO_3^- + H^+ \leftrightarrow H_2CO_3$$

In biological reactions, however, equilibrium is rarely obtained because the concentrations of the reactants or products or both are constantly changing, often with a product of one reaction being a reactant for another. To return to the example of excess hydrogen ions in the blood, the formation of carbonic acid will be the major direction of the reaction. However, the carbonic acid can also leave the body as carbon dioxide gas (via exhalation) instead of being converted back to bicarbonate ion, thus driving the reaction to the right by the chemical law known as **law of mass action**. These reactions are important for maintaining the homeostasis of our blood.

$$CO_3 \leftrightarrow CO_2 + H_2O$$

## In Summary: Chemical Reactions and Molecules

The outer electron shell dictates how readily and what type of chemical bonds a particular atom will form. The formation of compounds is often visually outlined in chemical equations which show the reactants participating in chemical reactions to form products.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.16: Chemical Reactions and Molecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.17: Anabolic and Catabolic Pathways

Anabolic pathways require an input of energy to synthesize complex molecules from simpler ones. Synthesizing sugar from $CO_2$ is one example. Other examples are the synthesis of large proteins from amino acid building blocks, and the synthesis of new DNA strands from nucleic acid building blocks. These biosynthetic processes are critical to the life of the cell, take place constantly, and demand energy provided by ATP and other high-energy molecules like NADH (nicotinamide adenine dinucleotide) and NADPH (Figure 1).

ATP is an important molecule for cells to have in sufficient supply at all times. The breakdown of sugars illustrates how a single molecule of glucose can store enough energy to make a great deal of ATP, 36 to 38 molecules. This is a catabolic pathway. Catabolic pathways involve the degradation (or breakdown) of complex molecules into simpler ones. Molecular energy stored in the bonds of complex molecules is released in catabolic pathways and harvested in such a way that it can be used to produce ATP. Other energy-storing molecules, such as fats, are also broken down through similar catabolic reactions to release energy and make ATP (Figure 1).





Figure 1. Anabolic pathways are those that require energy to synthesize larger molecules. Catabolic pathways are those that generate energy by breaking down larger molecules. Both types of pathways are required for maintaining the cell's energy balance.

It is important to know that the chemical reactions of metabolic pathways don't take place spontaneously. Each reaction step is facilitated, or catalyzed, by a protein called an enzyme. Enzymes are important for catalyzing all types of biological reactions—those that require energy as well as those that release energy.

**Remember:** Anabolic pathways assemble large molecules form smaller ones. Catabolic pathways break large molecules into small pieces.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.17: Anabolic and Catabolic Pathways is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.18: Enzymes

Enzymes are proteins that have the ability to bind substrate in their active site and then chemically modify the bound substrate, converting it to a different molecule — the *product* of the reaction. *Substrates* bind to enzymes just like ligands bind to proteins. However, when substrates bind to enzymes, they undergo an enzyme-induced chemical change, and are converted to products.





Figure 1. Compare the protein-ligand interaction to the enzyme-substrate interaction. Notice that both binding proteins and enzymes have binding sites for their ligands (L) and substrates (S), respectively. This area of the enzyme is called the active site because it also contains amino acids that are important for the conversion of substrate to product.

The substrate binds to the enzyme by interacting with amino acids in the *binding site*. The binding site on enzymes is often referred to as the *active site* because it contains amino acids that both bind the substrate and aid in its conversion to product.

You can often recognize that a protein is an enzyme by its name. Many enzyme names end with *–ase*. For example, the enzyme lactase is used to break down the sugar lactose, found in mammalian milk. Other enzymes are known by a common name, such as pepsin, which is an enzyme that aids in the digestion of proteins in your stomach by breaking the peptide bonds in the proteins.

Enzymes are catalysts, meaning that they make a reaction go faster, but the enzymes themselves are not altered by the overall reaction. Examine this image to see how enzymes work.





Figure 2. Simplified enzymatic reaction. The substrate reversibly binds to the active site of the enzyme, forming the enzyme-substrate (ES) complex. The bound substrate is converted to product by catalytic groups in the active site, forming the enzyme-product complex (EP). The bound products are released, returning the enzyme to its unbound form, ready to catalyze another round of converting substrate to product.

The amino acids in the active site of enzymes play two roles, and sometimes those roles overlap. Some of the amino acids in the active site are responsible for binding of the substrate and others are responsible for facilitating the chemical reaction. Enzymes are generally quite specific for their substrates. Although lactase and pepsin both catalyze the same type of reaction, breaking a bond using water (hydrolysis: “hydro” means “water” and “lysis” means “to break”), lactase only functions when lactose is its substrate and pepsin can only break peptide bonds.

## Practice Question

Two substrates—lactose and a short protein—are shown on the left. Two enzymes are shown on the right, labeled A and B. Which of the two enzymes is lactase?




[practice-area rows="2"][/practice-area]
[reveal-answer q="551250"]**Show Answer**[/reveal-answer]
[hidden-answer a="551250"]Enzyme A. The active site on this enzyme is the same shape as lactose. The hexagonal shape on the lactose is a poor fit for the rectangular shaped active site on enzyme B.[/hidden-answer]

## How Enzymes Work





Figure 3. Diagram of a catalytic reaction showing difference in activation energy in uncatalysed and catalysed reaction. The enzyme reduces the energy barrier required to activate the substrate, allowing more substrates to become activated, which increases the rate of product formation. Note that the energy difference between the substrate and the product is not changed by the enzyme.

In all chemical reactions, there is an initial input of energy that is required before the reaction can occur. If this initial energy requirement (called the activation energy or energy barrier) is small, then the reaction will happen quickly and easily. If the activation energy is large, then the reaction will take longer to occur. Enzymes function to reduce the activation energy required for a chemical reaction to occur.

First, the enzyme binds to the substrate and slightly distorts its shape. The change in shape activates the substrate molecule and decreases the total activation energy required for the substrate to be turned into product. As the number of activated substrate molecules increases, so does the conversion of substrate to product. An analogy for this effect is a ski hill, with skiers at the bottom of one side of the hill representing substrates, skiers on the top of the hill representing activated substrates, and the products being the number of skiers that ski down the other side. If the height of the hill is lowered (due to the presence of the enzyme), then more skiers can make it to the top, increasing the number that ski down to become products.

### Practice Questions

Fill in the blank: When an enzyme catalyzes a reaction, _____.

a. it raises the activation energy of the reaction.
b. it is used once and discarded.
c. it becomes a product.
d. it acts as a reactant.
e. it lowers the activation energy of the reaction.

[reveal-answer q="101995"]Show Answer[/reveal-answer]
[hidden-answer a="101995"]Answer e. When an enzyme catalyzes a reaction, it lowers the activation energy of the reaction; lowering the activation energy of a reaction makes it more likely to occur.

[/hidden-answer]

What will happen to the rate at which a chemical reaction proceeds if the activation energy is increased?

a. The reaction will happen faster (at a higher rate).
b. The reaction will happen slower (at a lower rate).
c. The reaction rate will not change.

[reveal-answer q="42089″]**Show Answers**[/reveal-answer]
[hidden-answer a="42089″]Answer b. If the activation energy goes up, then more energy is required for the reaction to occur, so it will happen less often.[/hidden-answer]

## Learning Objectives

Enzymes are proteins that speed up reactions by reducing the activation energy. Each enzyme typically binds only one substrate. Enzymes are not consumed during a reaction; instead they are available to bind new substrates and catalyze the same reaction repeatedly.

## Contributors and Attributions

CC licensed content, Shared previously

- How Enzymes Work. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5d6380020ca600f6190aedc2f27b. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*
- Induced fit diagram. **Authored by**: TimVickers; vectorized by Fvasconcellos. **Located at**: https://en.Wikipedia.org/wiki/File:Induced_fit_diagram.svg. **License**: *CC0: No Rights Reserved*
- Activation2 updated. **Authored by**: Jerry Crimson Mann, vectorized by Tutmosis, corrected by Fvasconcellos. **Located at**: https://commons.wikimedia.org/wiki/File:Activation2_updated.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

4.18: Enzymes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.19: Introduction to Thermodynamics

## Demonstrate familiarity with the first and second laws of thermodynamics

There are four laws of thermodynamics; however, for this course only the first two are relevant:

1. Energy may be transferred from place to place or transformed into different forms, but it cannot be created or destroyed.
2. In any given system, the entropy (the amount of energy) will either increase or stay the same.

In this outcome, we will learn exactly how these laws are important to understanding biology.

## What You'll Learn to Do

- Distinguish between an open and a closed system
- Understand how the first law of thermodynamics applies to biological systems
- Understand how the second law of thermodynamics applies to biological systems

### Learning Activities

The learning activities for this section include the following:

- Open and Closed Systems
- The First Law of Thermodynamics
- The Second Law of Thermodynamics
- Self Check: Thermodynamics

## Contributors and Attributions

CC licensed content, Original

- Introduction to Thermodynamics. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

4.19: Introduction to Thermodynamics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.20: Open and Closed Systems

Thermodynamics refers to the study of energy and energy transfer involving physical matter. The matter and its environment relevant to a particular case of energy transfer are classified as a system, and everything outside of that system is called the surroundings. For instance, when heating a pot of water on the stove, the system includes the stove, the pot, and the water. Energy is transferred within the system (between the stove, pot, and water). There are two types of systems: open and closed. An open system is one in which energy can be transferred between the system and its surroundings. The stovetop system is open because heat can be lost into the air. A closed system is one that cannot transfer energy to its surroundings.

Biological organisms are open systems. Energy is exchanged between them and their surroundings, as they consume energy-storing molecules and release energy to the environment by doing work. Like all things in the physical world, energy is subject to the laws of physics. The laws of thermodynamics govern the transfer of energy in and among all systems in the universe.

In studying energy, scientists use the term “system” to refer to the matter and its environment involved in energy transfers. Everything outside of the system is called the surroundings. Single cells are biological systems. Systems can be thought of as having a certain amount of order. It takes energy to make a system more ordered. The more ordered a system is, the lower its entropy. Entropy is a measure of the disorder of a system. As a system becomes more disordered, the lower its energy and the higher its entropy become.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.20: Open and Closed Systems is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.21: The First Law of Thermodynamics

The first law of thermodynamics deals with the total amount of energy in the universe. It states that this total amount of energy is constant. In other words, there has always been, and always will be, exactly the same amount of energy in the universe. Energy exists in many different forms. According to the first law of thermodynamics, energy may be transferred from place to place or transformed into different forms, but it cannot be created or destroyed. The transfers and transformations of energy take place around us all the time. Light bulbs transform electrical energy into light energy. Gas stoves transform chemical energy from natural gas into heat energy. Plants perform one of the most biologically useful energy transformations on earth: that of converting the energy of sunlight into the chemical energy stored within organic molecules. Some examples of energy transformations are shown in Figure 1.





Figure 1. Shown are two examples of energy being transferred from one system to another and transformed from one form to another. Humans can convert the chemical energy in food, like this ice cream cone, into kinetic energy (the energy of movement to ride a bicycle). Plants can convert electromagnetic radiation (light energy) from the sun into chemical energy. (credit “ice cream”: modification of work by D. Sharon Pruitt; credit “kids on bikes”: modification of work by Michelle Riggen-Ransom; credit “leaf”: modification of work by Cory Zanker)

The challenge for all living organisms is to obtain energy from their surroundings in forms that they can transfer or transform into usable energy to do work. Living cells have evolved to meet this challenge very well. Chemical energy stored within organic molecules such as sugars and fats is transformed through a series of cellular chemical reactions into energy within molecules of ATP. Energy in ATP molecules is easily accessible to do work. Examples of the types of work that cells need to do include building complex molecules, transporting materials, powering the beating motion of cilia or flagella, contracting muscle fibers to create movement, and reproduction.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.21: The First Law of Thermodynamics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.22: The Second Law of Thermodynamics

A living cell's primary tasks of obtaining, transforming, and using energy to do work may seem simple. However, the second law of thermodynamics explains why these tasks are harder than they appear. None of the energy transfers we've discussed, along with all energy transfers and transformations in the universe, is completely efficient. In every energy transfer, some amount of energy is lost in a form that is unusable. In most cases, this form is heat energy. Thermodynamically, heat energy is defined as the energy transferred from one system to another that is not doing work. For example, when an airplane flies through the air, some of the energy of the flying plane is lost as heat energy due to friction with the surrounding air. This friction actually heats the air by temporarily increasing the speed of air molecules. Likewise, some energy is lost as heat energy during cellular metabolic reactions. This is good for warm-blooded creatures like us, because heat energy helps to maintain our body temperature. Strictly speaking, no energy transfer is completely efficient, because some energy is lost in an unusable form.





Figure 1. Entropy is a measure of randomness or disorder in a system. Gases have higher entropy than liquids, and liquids have higher entropy than solids.

An important concept in physical systems is that of order and disorder (also known as randomness). The more energy that is lost by a system to its surroundings, the less ordered and more random the system is. Scientists refer to the measure of randomness or disorder within a system as entropy. High entropy means high disorder and low energy (Figure 1). To better understand entropy, think of a student's bedroom. If no energy or work were put into it, the room would quickly become messy. It would exist in a very disordered state, one of high entropy. Energy must be put into the system, in the form of the student doing work and putting everything away, in order to bring the room back to a state of cleanliness and order. This state is one of low entropy. Similarly, a car or house must be constantly maintained with work in order to keep it in an ordered state. Left alone, the entropy of the house or car gradually increases through rust and degradation. Molecules and chemical reactions have varying amounts of entropy as well. For example, as chemical reactions reach a state of equilibrium, entropy increases, and as molecules at a high concentration in one place diffuse and spread out, entropy also increases.

## Try It Yourself

Set up a simple experiment to understand how energy is transferred and how a change in entropy results.

1. Take a block of ice. This is water in solid form, so it has a high structural order. This means that the molecules cannot move very much and are in a fixed position. The temperature of the ice is 0°C. As a result, the entropy of the system is low.
2. Allow the ice to melt at room temperature. What is the state of molecules in the liquid water now? How did the energy transfer take place? Is the entropy of the system higher or lower? Why?
3. Heat the water to its boiling point. What happens to the entropy of the system when the water is heated?

All physical systems can be thought of in this way: Living things are highly ordered, requiring constant energy input to be maintained in a state of low entropy. As living systems take in energy-storing molecules and transform them through chemical reactions, they lose some amount of usable energy in the process, because no reaction is completely efficient. They also produce waste and by-products that aren't useful energy sources. This process increases the entropy of the system's surroundings. Since all energy transfers result in the loss of some usable energy, the second law of thermodynamics states that every energy transfer or

transformation increases the entropy of the universe. Even though living things are highly ordered and maintain a state of low entropy, the entropy of the universe in total is constantly increasing due to the loss of usable energy with each energy transfer that occurs. Essentially, living things are in a continuous uphill battle against this constant increase in universal entropy.

## Contributors and Attributions


CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

4.22: The Second Law of Thermodynamics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.23: Putting It Together- Chemistry of Life

As we've just learned, chemistry is essential to life: we are all made of compounds and molecules. Think back to the beginning of this module, where we briefly discussed the term *carbon-basedlife*. In the video below, we'll learn about why carbon is so essential to life:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=126

## Occupation Spotlight: Nutritionists

Let's look back to our earlier occupation spotlight on the nutritionist. You should now be able see how an understanding of chemistry is essential to this job: a nutritionist needs to understand how the body builds complex molecules from the food a person ingests. A nutritionist also should understand how energy is used and moved about in the body. All this requires a basic understanding of elements, their structures, and how they interact.

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

All rights reserved content

- That's Why Carbon Is A Tramp: Crash Course Biology #1. **Authored by**: CrashCourse. **Located at**: https://youtu.be/QnQe0xW_JY4. **Project**: Crash Course Biology. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

4.23: Putting It Together- Chemistry of Life is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.24: Assignment- Biological Astronaut

Imagine you are an exploratory astronaut looking for life throughout the universe. One day you encounter a planet that has no carbon present on its surface. However, your instruments register movement and a variety of other signs that make you think life exists on the surface.

- **Part 1**: Before taking a potentially dangerous trip to the surface, you must outline a theoretical framework in which another element can serve as a backbone for macromolecules. (Hint: look for an element on the periodic table that would act similarly to carbon.) Begin by describing this new backbone, including how compounds and macromolecules would form. Detail at least 2 chemical reactions forming macromolecules with this backbone. You may wish add supporting diagrams (created or obtained). Be sure to include references as appropriate.
- **Part 2**: Your theoretical framework is deemed strong enough to justify a trip landside. Once there, you are authorized to collect a simple "organism" for experimental use. Collect your specimen(s) and then design a full experiment that will test at least two characteristics that define biological life on Earth. Be sure to include all the relevant parts of an experiment and describe how you would analyze and present the data, results and conclusions.

Take a look at this breakdown for hints on completing this assignment.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references

## Rubric

Outcomes: Define biology and apply its principles. Identify the principles of chemistry that are integral to biology.

Rubric for Biological Astronaut Assignment

| Criteria | Ratings | | | Max Points |
|---|---|---|---|---|
| Define atoms and elements through the identification of an element capable of replacing carbon as the backbone atom in macromolecules. Illustrations are encouraged here. | Submission identified element that can replace carbon as a backbone element. Illustrations and descriptions included that directly compare the two elements.<br>5.0 pts | Submission identifies element that can replace carbon as a backbone element. Description directly compares two elements.<br>4.0 pts | No or wrong element chosen or justification provided is inaccurate/insufficient for identifying replacement backbone element.<br>0.0 pts | 5 pts |
| Discuss the role electrons play in enabling chosen element to replace carbon. | Electrons are discussed in terms of atomic bonding behaviors. Direct comparison is provided between carbon and new element. Illustrations are included to show similarities in bonding activity.<br>5.0 pts | Electrons are discussed in terms of atomic bonding behaviors. Direct comparison is provided between carbon and new element.<br>4.0 pts | Electrons are not discussed or are discussed inaccurately in terms of bonding behaviors.<br>0.0 pts | 5 pts |
| Identify the components of two macromolecule formation reactions and the macromolecules formed. Illustrations/diagrams are encouraged here. | Two macromolecules discussed with the new backbone element; all components in the formation of these molecules are correctly identified. Illustrations are included. There is a direct comparison between new macromolecules and traditional macromolecules.<br>5.0 pts | Two macromolecules discussed with the new backbone element; all components in the formation of these molecules are correctly identified. Illustrations may be included.<br>4.0 pts | Less than two formation reactions are included or all the components are not properly identified.<br>0.0 pts | 5 pts |
| Identify at least 4 altered functional groups formed without the use of carbon. | At least 4 functional groups identified that would form without carbon. Descriptions are detailed and well diagrammed images/figures included to illustrate these groups.<br>5.0 pts | At least 4 new functional groups identified that would form without carbon. Descriptions are adequate or basic illustrations included.<br>4.0 pts | Less than 4 functional groups identified/discussed or new functional groups are inaccurate.<br>0.0 pts | 5 pts |
| Demonstrate the critical thinking required to conduct a scientific experiment by designing a full experiment to test if an organism is ‘alive’. This experiment contains all the required | All components of an experiment are included and clearly discussed. At least 3 standardized variables included. Hypothesis is clear and testable. Potential compounding factors or areas of uncertainty identified.<br>5.0 pts | All components of an experiment are included. At least 3 standardized variables included. Hypothesis is clear and testable.<br>4.0 pts | Experiment does not include all the required components of an experiment or components are inaccurately identified/labeled. Hypothesis is not testable.<br>0.0 pts | 5 pts |

| Criteria | Ratings | | | Max Points |
|---|---|---|---|---|
| components of an experiment. | | | | |
| The designed experiment tests at least one characteristic that defines biological life. | Experiment tests at least one characteristic of biological life and the way this is tested is clearly explained.<br>5.0 pts | Experiment tests at least one characteristic of biological life.<br>4.0 pts | Experiment does not test characteristic of biological life.<br>0.0 pts | 5 pts |
| Possible results from the experiment are discussed, and how it would be concluded this organism is living or not. Organism is also identified as analogous to either prokaryotic or eukaryotic organism. | Potential results are discussed and the discussed results logically follow from outlined experiment. Nuances of possible conclusions discussed. Organism identified by cell type.<br>5.0 pts | Potential results are discussed and the discussed results logically follow from outlined experiment. Organism identified by cell type.<br>4.0 pts | Potential results are not discussed or the discussed results do not logically follow from outlined experiment. Organism not identified by cell type.<br>0.0 pts | 5 pts |
| | | | Total points: | 35 |

## Contributors and Attributions

CC licensed content, Original

- Performance Assessment: Biological Astronaut. **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

4.24: Assignment- Biological Astronaut is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 4.25: Assignment- Interacting with Attributes of Molecules

*Open Pedagogy Assignments are assignments in which students use their agency and creativity to create knowledge artifacts that can support their own learning, their classmates' learning, and the learning of students around the world. (See this peer-reviewed article for more details.) The assignment on this page is aligned to a learning outcome of Biology for Majors I and we've identified the module where the reading appears. The assignment can be created with basic web and computing tools, a cell phone camera or any video recording device, Google or Word documents, and your learning management system.*

## Learning Objectives

Categorize molecules according to their functional group

- Identify the attributes of molecules with hydroxyl groups
- Identify the attributes of molecules with carboxyl groups
- Identify the attributes of molecules with amino groups
- Identify the attributes of molecules with phosphate groups
- Identify the attributes of molecules with methyl groups
- Identify the attributes of molecules with carbonyl groups
- Identify the attributes of molecules with sulfhydryl groups

One topic that is important to learn about the chemistry of life is functional groups and the effects they have one the molecules they are bonded to. For this assignment, you are going to create an interactive, or a short learning activity, about a functional group. The purpose of the interactive is to provide yourself and others the opportunity to better understand concepts from the content in order to make it more approachable and interesting. The best interactives may be eligible to be incorporated into the course for future students!


**Instructions:**

STEP 1: Research functional groups. Use the content in the Chemistry of Life module for starters, but also review current research and journal articles related to the topic. You are required to include at least one reference to an outside source in the interactive. Any and all content you use inside of the interactive needs to be written in your own words and appropriately cited.


STEP 2: Write out your plan for the interactive. It works well to storyboard it first on a document so you know what you'll want to include on each slide. Here's an example:

- Slide 1: Introduction paragraph on the topic. Introductory example.
- Slide 2: Multiple-choice question about the example, with feedback.
- Slide 3: Introductory text. Picture.
- Slide 4: Matching activity.
- Slide 5: Conclusion.

STEP 3: Go to H5P.org and create a free account. Click "Create New Content."



Read through the tutorial on creating a course presentation. Select the type of content you want to create, and get started.




STEP 4: Create your interactive! Read through this brief tutorial on adding attributions or use the guides within H5P to help you create the type of content you need.

STEP 5: Copy and paste the URL of your interactive to share it with your instructor. Include a short description (about 100 words) of what you learned from the assignment. You may include how you made the interactive and the additional resources you found.

---

**A Note To Teachers:** This assignment can be readily adapted to align with any learning outcome or topic in the course. Identify an area that your students struggle with most and point your students there.

Additonally, you may want to offer some scaffold for the assignment by asking students to submit their plan from Step 3. This will offer you the cha

The first time your students complete this assignment, choose the best ones and ask students for permission to include them in future sections. In the first term, students will create interactives, and with their permission, you can post links to them in your course in order to show examples for the next term. The idea is to have students generate content that other students can learn from in this assignment.

## About the tools:

You will make these interactives using H5P, using the course presentation tool. The presentation should have a minimum of 4 slides and a maximum of 10. The course presentation feature in H5P works like this:

- H5P Course presentation

The other types of H5P tools you'll likely to include within the H5P presentation are:

- H5P Image Sequencing
- H5P Drag the words (this is best for matching)
- H5P Fill in the blanks
- H5P Multiple choice

The goal is not to simply duplicate the multiple-choice practice questions that you already see in course, but to create an activity that provides a deeper review or an opportunity to elaborate upon important concepts. You could also pull in current research or other engaging articles to expound upon a topic.

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: William Carr and Lumen Learning. **Provided by**: William Carr, Associate Professor of Biology at Medgar Evars College. **License**: *CC BY: Attribution*

---

4.25: Assignment- Interacting with Attributes of Molecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 5: Module 3- Important Biological Macromolecules


5.1: Why It Matters- Important Biological Macromolecules
5.2: Introduction to Proteins
5.3: Amino Acids
5.4: Protein Structure
5.5: Function of Proteins
5.6: Introduction to Lipids
5.7: Lipids
5.8: Introduction to Carbohydrates
5.9: Structure and Function of Carbohydrates
5.10: Introduction to Nucleic Acids
5.11: Structure of Nucleic Acids
5.12: DNA and RNA
5.13: Introduction to Comparing Biological Macromolecules
5.14: Different Types of Biological Macromolecules
5.15: Putting It Together- Important Biological Macromolecules
5.16: Assignment- Nutritionist for a Day



5: Module 3- Important Biological Macromolecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.1: Why It Matters- Important Biological Macromolecules

## Why identify and describe the main features of the four main classes of important biological macromolecules?

Fad diets: we've all heard about them and maybe followed one or two in our lives. These diets have strict rules, and often have restrictions on eating a certain thing like fats or carbs (carbohydrates).

There are several things to critically consider about this type of diet. First off, is it even possible for a person to cut all carbs out of his or her diet? More importantly, is it actually healthy to remove an entire class of molecules from the diet? Fats aren't really important right? Certainly cholesterol is bad—right?

Before you decide to swear off carbs or fats, you should know that these types of foods are named after the kind of molecules that build them. Then you should learn what they actually do in cells.

Biological macromolecules are large molecules, necessary for life, that are built from smaller organic molecules. There are four major classes of biological macromolecules: carbohydrates, lipids, proteins, and nucleic acids (found in DNA and RNA). We'll discuss each class and how they compare to each other.

### Learning Outcomes

- Describe the structure and function proteins
- Illustrate different types of lipids and relate their structure to their role in biological systems
- Summarize the roles carbohydrates play in biological systems
- Discuss nucleic acids and the role they play in DNA and RNA
- Discuss macromolecules and the differences between the four classes

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Important Biological Macromolecules. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.1: Why It Matters- Important Biological Macromolecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.2: Introduction to Proteins

## Describe the structure and function proteins

Proteins are polymers of amino acids. Each amino acid contains a central carbon, a hydrogen, a carboxyl group, an amino group, and a variable R group. The R group specifies which class of amino acids it belongs to: electrically charged hydrophilic side chains, polar but uncharged side chains, nonpolar hydrophobic side chains, and special cases.

Proteins have different “layers” of structure: primary, secondary, tertiary, quaternary.

Proteins have a variety of function in cells. Major functions include acting as enzymes, receptors, transport molecules, regulatory proteins for gene expression, and so on. Enzymes are biological catalysts that speed up a chemical reaction without being permanently altered. They have “active sites” where the substrate/reactant binds, and they can be either activated or inhibited (competitive and/or noncompetitive inhibitors).

## What You’ll Learn to Do

- Demonstrate familiarity with monomeric units of proteins: amino acids
- Define the different layers of protein structure
- Identify several major functions of proteins

### Learning Activities

The learning activities for this section include the following:

- Amino Acids
- Protein Structure
- Function of Proteins
- Self Check: Proteins

## Contributors and Attributions

CC licensed content, Original

- Introduction to Proteins. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

5.2: Introduction to Proteins is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.3: Amino Acids

**Proteins** are one of the most abundant organic molecules in living systems and have the most diverse range of functions of all macromolecules. Proteins may be structural, regulatory, contractile, or protective; they may serve in transport, storage, or membranes; or they may be toxins or enzymes. Each cell in a living system may contain thousands of proteins, each with a unique function. Their structures, like their functions, vary greatly. They are all, however, polymers of **amino acids**, arranged in a linear sequence.





Figure 1. Amino acids have a central asymmetric carbon to which an amino group, a carboxyl group, a hydrogen atom, and a side chain (R group) are attached.

**Amino acids** are the monomers that make up proteins. Each amino acid has the same fundamental structure, which consists of a central carbon atom, also known as the alpha ($\alpha$) carbon, bonded to an amino group (NH2), a carboxyl group (COOH), and to a hydrogen atom. Every amino acid also has another atom or group of atoms bonded to the central atom known as the R group (Figure 1).

The name “amino acid” is derived from the fact that they contain both amino group and carboxyl-acid-group in their basic structure. As mentioned, there are 20 amino acids present in proteins. Ten of these are considered essential amino acids in humans because the human body cannot produce them and they are obtained from the diet.

For each amino acid, the R group (or side chain) is different (Figure 2).





Figure 2. There are 20 common amino acids commonly found in proteins, each with a different R group (variant group) that determines its chemical nature.

Which categories of amino acid would you expect to find on the surface of a soluble protein, and which would you expect to find in the interior? What distribution of amino acids would you expect to find in a protein embedded in a lipid bilayer?

[practice-area rows="2"][/practice-area]
[reveal-answer q="539127"]**Show Answer**[/reveal-answer]
[hidden-answer a="539127"]Polar and charged amino acid residues (the remainder after peptide bond formation) are more likely to be found on the surface of soluble proteins where they can interact with water, and nonpolar (e.g., amino acid side chains) are more likely to be found in the interior where they are sequestered from water. In membrane proteins, nonpolar and hydrophobic amino acid side chains associate with the hydrophobic tails of phospholipids, while polar and charged amino acid side chains interact with the polar head groups or with the aqueous solution. However, there are exceptions. Sometimes, positively and negatively charged amino acid side chains interact with one another in the interior of a protein, and polar or charged amino acid side chains that interact with a ligand can be found in the ligand binding pocket.[/hidden-answer]

The chemical nature of the side chain determines the nature of the amino acid (that is, whether it is acidic, basic, polar, or nonpolar). For example, the amino acid glycine has a hydrogen atom as the R group. Amino acids such as valine, methionine, and alanine are nonpolar or hydrophobic in nature, while amino acids such as serine, threonine, and cysteine are polar and have hydrophilic side chains. The side chains of lysine and arginine are positively charged, and therefore these amino acids are also known as basic amino acids. Proline has an R group that is linked to the amino group, forming a ring-like structure. Proline is an exception to the standard structure of an animo acid since its amino group is not separate from the side chain (Figure 2).

Amino acids are represented by a single upper case letter or a three-letter abbreviation. For example, valine is known by the letter V or the three-letter symbol val. Just as some fatty acids are essential to a diet, some amino acids are necessary as well. They are known as essential amino acids, and in humans they include isoleucine, leucine, and cysteine. Essential amino acids refer to those necessary for construction of proteins in the body, although not produced by the body; which amino acids are essential varies from organism to organism.





Figure 3. Peptide bond formation is a dehydration synthesis reaction. The carboxyl group of one amino acid is linked to the amino group of the incoming amino acid. In the process, a molecule of water is released.

The sequence and the number of amino acids ultimately determine the protein's shape, size, and function. Each amino acid is attached to another amino acid by a covalent bond, known as a peptide bond, which is formed by a dehydration reaction. The carboxyl group of one amino acid and the amino group of the incoming amino acid combine, releasing a molecule of water. The resulting bond is the peptide bond (Figure 3).

The products formed by such linkages are called peptides. As more amino acids join to this growing chain, the resulting chain is known as a polypeptide. Each polypeptide has a free amino group at one end. This end is called the N terminal, or the amino terminal, and the other end has a free carboxyl group, also known as the C or carboxyl terminal. While the terms polypeptide and protein are sometimes used interchangeably, a polypeptide is technically a polymer of amino acids, whereas the term protein is used for a polypeptide or polypeptides that have combined together, often have bound non-peptide prosthetic groups, have a distinct shape, and have a unique function. After protein synthesis (translation), most proteins are modified. These are known as post-translational modifications. They may undergo cleavage, phosphorylation, or may require the addition of other chemical groups. Only after these modifications is the protein completely functional.

## The Evolutionary Significance of Cytochrome c

Cytochrome c is an important component of the electron transport chain, a part of cellular respiration, and it is normally found in the cellular organelle, the mitochondrion. This protein has a heme prosthetic group, and the central ion of the heme gets alternately reduced and oxidized during electron transfer. Because this essential protein's role in producing cellular energy is crucial, it has changed very little over millions of years. Protein sequencing has shown that there is a considerable amount of cytochrome c amino acid sequence homology among different species; in other words, evolutionary kinship can be assessed by measuring the similarities or differences among various species' DNA or protein sequences.

Scientists have determined that human cytochrome c contains 104 amino acids. For each cytochrome c molecule from different organisms that has been sequenced to date, 37 of these amino acids appear in the same position in all samples of cytochrome c. This indicates that there may have been a common ancestor. On comparing the human and chimpanzee protein sequences, no sequence difference was found. When human and rhesus monkey sequences were compared, the single difference found was in one amino acid. In another comparison, human to yeast sequencing shows a difference in the 44th position.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.3: Amino Acids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.4: Protein Structure

As discussed earlier, the shape of a protein is critical to its function. For example, an enzyme can bind to a specific substrate at a site known as the active site. If this active site is altered because of local changes or changes in overall protein structure, the enzyme may be unable to bind to the substrate. To understand how the protein gets its final shape or conformation, we need to understand the four levels of protein structure: primary, secondary, tertiary, and quaternary.

## Primary Structure

The unique sequence of amino acids in a polypeptide chain is its primary structure. For example, the pancreatic hormone insulin has two polypeptide chains, A and B, and they are linked together by disulfide bonds. The N terminal amino acid of the A chain is glycine, whereas the C terminal amino acid is asparagine (Figure 1). The sequences of amino acids in the A and B chains are unique to insulin.





Figure 1. Bovine serum insulin is a protein hormone made of two peptide chains, A (21 amino acids long) and B (30 amino acids long). In each chain, primary structure is indicated by three-letter abbreviations that represent the names of the amino acids in the order they are present. The amino acid cysteine (cys) has a sulfhydryl (SH) group as a side chain. Two sulfhydryl groups can react in the presence of oxygen to form a disulfide (S-S) bond. Two disulfide bonds connect the A and B chains together, and a third helps the A chain fold into the correct shape. Note that all disulfide bonds are the same length, but are drawn different sizes for clarity.





Figure 2. The beta chain of hemoglobin is 147 residues in length, yet a single amino acid substitution leads to sickle cell anemia. In normal hemoglobin, the amino acid at position seven is glutamate. In sickle cell hemoglobin, this glutamate is replaced by a valine.

The unique sequence for every protein is ultimately determined by the gene encoding the protein. A change in nucleotide sequence of the gene's coding region may lead to a different amino acid being added to the growing polypeptide chain, causing a change in

protein structure and function. In sickle cell anemia, the hemoglobin $\beta$ chain (a small portion of which is shown in Figure 2) has a single amino acid substitution, causing a change in protein structure and function.

Specifically, the amino acid glutamic acid is substituted by valine in the $\beta$ chain. What is most remarkable to consider is that a hemoglobin molecule is made up of two alpha chains and two beta chains that each consist of about 150 amino acids. The molecule, therefore, has about 600 amino acids. The structural difference between a normal hemoglobin molecule and a sickle cell molecule—which dramatically decreases life expectancy—is a single amino acid of the 600. What is even more remarkable is that those 600 amino acids are encoded by three nucleotides each, and the mutation is caused by a single base change (point mutation), 1 in 1800 bases.





Figure 3. In this blood smear, visualized at 535x magnification using bright field microscopy, sickle cells are crescent shaped, while normal cells are disc-shaped. (credit: modification of work by Ed Uthman; scale-bar data from Matt Russell)

Because of this change of one amino acid in the chain, hemoglobin molecules form long fibers that distort the biconcave, or disc-shaped, red blood cells and assume a crescent or “sickle” shape, which clogs arteries (Figure 3). This can lead to myriad serious health problems such as breathlessness, dizziness, headaches, and abdominal pain for those affected by this disease.

## Secondary Structure

The local folding of the polypeptide in some regions gives rise to the secondary structure of the protein. The most common are the $\alpha$-helix and $\beta$-pleated sheet structures (Figure 4). Both structures are the $\alpha$-helix structure—the helix held in shape by hydrogen bonds. The hydrogen bonds form between the oxygen atom in the carbonyl group in one amino acid and another amino acid that is four amino acids farther along the chain.





Figure 4. The α-helix and β-pleated sheet are secondary structures of proteins that form because of hydrogen bonding between carbonyl and amino groups in the peptide backbone. Certain amino acids have a propensity to form an α-helix, while others have a propensity to form a β-pleated sheet.

Every helical turn in an alpha helix has 3.6 amino acid residues. The R groups (the variant groups) of the polypeptide protrude out from the $\alpha$-helix chain. In the $\beta$-pleated sheet, the "pleats" are formed by hydrogen bonding between atoms on the backbone of the polypeptide chain. The R groups are attached to the carbons and extend above and below the folds of the pleat. The pleated segments align parallel or antiparallel to each other, and hydrogen bonds form between the partially positive nitrogen atom in the amino group and the partially negative oxygen atom in the carbonyl group of the peptide backbone. The $\alpha$-helix and $\beta$-pleated sheet structures are found in most globular and fibrous proteins and they play an important structural role.

## Tertiary Structure

The unique three-dimensional structure of a polypeptide is its tertiary structure (Figure 5). This structure is in part due to chemical interactions at work on the polypeptide chain. Primarily, the interactions among R groups creates the complex three-dimensional tertiary structure of a protein. The nature of the R groups found in the amino acids involved can counteract the formation of the hydrogen bonds described for standard secondary structures. For example, R groups with like charges are repelled by each other and those with unlike charges are attracted to each other (ionic bonds). When protein folding takes place, the hydrophobic R groups of nonpolar amino acids lay in the interior of the protein, whereas the hydrophilic R groups lay on the outside. The former types of interactions are also known as hydrophobic interactions. Interaction between cysteine side chains forms disulfide linkages in the presence of oxygen, the only covalent bond forming during protein folding.





Figure 5. The tertiary structure of proteins is determined by a variety of chemical interactions. These include hydrophobic interactions, ionic bonding, hydrogen bonding and disulfide linkages.

All of these interactions, weak and strong, determine the final three-dimensional shape of the protein. When a protein loses its three-dimensional shape, it may no longer be functional.

## Quaternary Structure

In nature, some proteins are formed from several polypeptides, also known as subunits, and the interaction of these subunits forms the quaternary structure. Weak interactions between the subunits help to stabilize the overall structure. For example, insulin (a globular protein) has a combination of hydrogen bonds and disulfide bonds that cause it to be mostly clumped into a ball shape. Insulin starts out as a single polypeptide and loses some internal sequences in the presence of post-translational modification after the formation of the disulfide linkages that hold the remaining chains together. Silk (a fibrous protein), however, has a $\beta$-pleated sheet structure that is the result of hydrogen bonding between different chains.

The four levels of protein structure (primary, secondary, tertiary, and quaternary) are illustrated in Figure 6.





Figure 6. The four levels of protein structure can be observed in these illustrations. (credit: modification of work by National Human Genome Research Institute)

## Denaturation and Protein Folding

Each protein has its own unique sequence and shape that are held together by chemical interactions. If the protein is subject to changes in temperature, pH, or exposure to chemicals, the protein structure may change, losing its shape without losing its primary sequence in what is known as denaturation. Denaturation is often reversible because the primary structure of the polypeptide is conserved in the process if the denaturing agent is removed, allowing the protein to resume its function. Sometimes denaturation is irreversible, leading to loss of function. One example of irreversible protein denaturation is when an egg is fried. The albumin protein in the liquid egg white is denatured when placed in a hot pan. Not all proteins are denatured at high temperatures; for instance, bacteria that survive in hot springs have proteins that function at temperatures close to boiling. The stomach is also very acidic, has a low pH, and denatures proteins as part of the digestion process; however, the digestive enzymes of the stomach retain their activity under these conditions.

Protein folding is critical to its function. It was originally thought that the proteins themselves were responsible for the folding process. Only recently was it found that often they receive assistance in the folding process from protein helpers known as chaperones (or chaperonins) that associate with the target protein during the folding process. They act by preventing aggregation of polypeptides that make up the complete protein structure, and they disassociate from the protein once the target protein is folded.

For an additional perspective on proteins, view this animation called “Biomolecules: The Proteins.”

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.4: Protein Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 5.5: Function of Proteins

The primary types and functions of proteins are listed in Table 1.

Table 1. Protein Types and Functions

| Type | Examples | Functions |
|---|---|---|
| Digestive Enzymes | Amylase, lipase, pepsin, trypsin | Help in digestion of food by catabolizing nutrients into monomeric units |
| Transport | Hemoglobin, albumin | Carry substances in the blood or lymph throughout the body |
| Structural | Actin, tubulin, keratin | Construct different structures, like the cytoskeleton |
| Hormones | Insulin, thyroxine | Coordinate the activity of different body systems |
| Defense | Immunoglobulins | Protect the body from foreign pathogens |
| Contractile | Actin, myosin | Effect muscle contraction |
| Storage | Legume storage proteins, egg white (albumin) | Provide nourishment in early development of the embryo and the seedling |

Two special and common types of proteins are enzymes and hormones. **Enzymes**, which are produced by living cells, are catalysts in biochemical reactions (like digestion) and are usually complex or conjugated proteins. Each enzyme is specific for the substrate (a reactant that binds to an enzyme) it acts on. The enzyme may help in breakdown, rearrangement, or synthesis reactions. Enzymes that break down their substrates are called catabolic enzymes, enzymes that build more complex molecules from their substrates are called anabolic enzymes, and enzymes that affect the rate of reaction are called catalytic enzymes. It should be noted that all enzymes increase the rate of reaction and, therefore, are considered to be organic catalysts. An example of an enzyme is salivary amylase, which hydrolyzes its substrate amylose, a component of starch.

**Hormones** are chemical-signaling molecules, usually small proteins or steroids, secreted by endocrine cells that act to control or regulate specific physiological processes, including growth, development, metabolism, and reproduction. For example, insulin is a protein hormone that helps to regulate the blood glucose level.

Proteins have different shapes and molecular weights; some proteins are globular in shape whereas others are fibrous in nature. For example, hemoglobin is a globular protein, but collagen, found in our skin, is a fibrous protein. Protein shape is critical to its function, and this shape is maintained by many different types of chemical bonds. Changes in temperature, pH, and exposure to chemicals may lead to permanent changes in the shape of the protein, leading to loss of function, known as denaturation. All proteins are made up of different arrangements of the same 20 types of amino acids.

### In Summary: Function of Proteins

Proteins are a class of macromolecules that perform a diverse range of functions for the cell. They help in metabolism by providing structural support and by acting as enzymes, carriers, or hormones. The building blocks of proteins (monomers) are amino acids. Each amino acid has a central carbon that is linked to an amino group, a carboxyl group, a hydrogen atom, and an R group or side chain. There are 20 commonly occurring amino acids, each of which differs in the R group. Each amino acid is linked to its neighbors by a peptide bond. A long chain of amino acids is known as a polypeptide.

Proteins are organized at four levels: primary, secondary, tertiary, and (optional) quaternary. The primary structure is the unique sequence of amino acids. The local folding of the polypeptide to form structures such as the $\alpha$ helix and $\beta$-pleated sheet constitutes the secondary structure. The overall three-dimensional structure is the tertiary structure. When two or more polypeptides combine to form the complete protein structure, the configuration is known as the quaternary structure of a protein. Protein shape and

function are intricately linked; any change in shape caused by changes in temperature or pH may lead to protein denaturation and a loss in function.

## Contributors and Attributions

CC licensed content, Original

- Revision, adaptation, and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

5.5: Function of Proteins is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.6: Introduction to Lipids

## Illustrate different types of lipids and relate their structure to their role in biological systems

In this outcome, we will discuss lipids, or fats, and the role they plan in our bodies.

## What You'll Learn to Do

- Distinguish between the different kinds of lipids
- Identify several major functions of lipids

### Learning Activities

The learning activities for this section include the following:

- Lipids
- Self Check: Lipids

## Contributors and Attributions

CC licensed content, Original

- Introduction to Lipids. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

5.6: Introduction to Lipids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.7: Lipids



Figure 1. Hydrophobic lipids in the fur of aquatic mammals, such as this river otter, protect them from the elements. (credit: Ken Bosma)

Lipids include a diverse group of compounds that are largely nonpolar in nature. This is because they are hydrocarbons that include mostly nonpolar carbon–carbon or carbon–hydrogen bonds. Non-polar molecules are hydrophobic (“water fearing”), or insoluble in water. Lipids perform many different functions in a cell. Cells store energy for long-term use in the form of fats. Lipids also provide insulation from the environment for plants and animals (Figure 1). For example, they help keep aquatic birds and mammals dry when forming a protective layer over fur or feathers because of their water-repellant hydrophobic nature. Lipids are also the building blocks of many hormones and are an important constituent of all cellular membranes. Lipids include fats, oils, waxes, phospholipids, and steroids.

## Fats and Oils

A fat molecule consists of two main components—glycerol and fatty acids. Glycerol is an organic compound (alcohol) with three carbons, five hydrogens, and three hydroxyl (OH) groups. Fatty acids have a long chain of hydrocarbons to which a carboxyl group is attached, hence the name “fatty acid.” The number of carbons in the fatty acid may range from 4 to 36; most common are those containing 12–18 carbons. In a fat molecule, the fatty acids are attached to each of the three carbons of the glycerol molecule with an ester bond through an oxygen atom (Figure 2).





Figure 2. Triacylglycerol is formed by the joining of three fatty acids to a glycerol backbone in a dehydration reaction. Three molecules of water are released in the process.

During this ester bond formation, three water molecules are released. The three fatty acids in the triacylglycerol may be similar or dissimilar. Fats are also called **triacylglycerols** or **triglycerides** because of their chemical structure. Some fatty acids have common names that specify their origin. For example, palmitic acid, a **saturated fatty acid**, is derived from the palm tree. Arachidic acid is derived from *Arachis hypogea*, the scientific name for groundnuts or peanuts.

Fatty acids may be saturated or unsaturated. In a fatty acid chain, if there are only single bonds between neighboring carbons in the hydrocarbon chain, the fatty acid is said to be saturated. Saturated fatty acids are saturated with hydrogen; in other words, the number of hydrogen atoms attached to the carbon skeleton is maximized. Stearic acid is an example of a saturated fatty acid (Figure 3)




Figure 3. Stearic acid is a common saturated fatty acid.

When the hydrocarbon chain contains a double bond, the fatty acid is said to be unsaturated. Oleic acid is an example of an unsaturated fatty acid (Figure 4).





Figure 4. Oleic acid is a common unsaturated fatty acid.

Most unsaturated fats are liquid at room temperature and are called oils. If there is one double bond in the molecule, then it is known as a monounsaturated fat (e.g., olive oil), and if there is more than one double bond, then it is known as a polyunsaturated fat (e.g., canola oil).

When a fatty acid has no double bonds, it is known as a saturated fatty acid because no more hydrogen may be added to the carbon atoms of the chain. A fat may contain similar or different fatty acids attached to glycerol. Long straight fatty acids with single bonds tend to get packed tightly and are solid at room temperature. Animal fats with stearic acid and palmitic acid (common in meat) and the fat with butyric acid (common in butter) are examples of saturated fats. Mammals store fats in specialized cells called adipocytes, where globules of fat occupy most of the cell's volume. In plants, fat or oil is stored in many seeds and is used as a source of energy during seedling development. Unsaturated fats or oils are usually of plant origin and contain *cis* unsaturated fatty acids. *Cis* and *trans* indicate the configuration of the molecule around the double bond. If hydrogens are present in the same plane, it is referred to as a cis fat; if the hydrogen atoms are on two different planes, it is referred to as a trans fat. The *cis* double bond causes a bend or a "kink" that prevents the fatty acids from packing tightly, keeping them liquid at room temperature (Figure 5). Olive oil, corn oil, canola oil, and cod liver oil are examples of unsaturated fats. Unsaturated fats help to lower blood cholesterol levels whereas saturated fats contribute to plaque formation in the arteries.


**Saturated fatty acid**

Stearic acid



**Unsaturated fatty acids**

*Cis* oleic acid



*Trans* oleic acid




Figure 5. Saturated fatty acids have hydrocarbon chains connected by single bonds only. Unsaturated fatty acids have one or more double bonds. Each double bond may be in a cis or trans configuration. In the cis configuration, both hydrogens are on the same side of the hydrocarbon chain. In the trans configuration, the hydrogens are on opposite sides. A cis double bond causes a kink in the chain.

## Trans Fats

In the food industry, oils are artificially hydrogenated to make them semi-solid and of a consistency desirable for many processed food products. Simply speaking, hydrogen gas is bubbled through oils to solidify them. During this hydrogenation process, double bonds of the *cis*– conformation in the hydrocarbon chain may be converted to double bonds in the *trans*– conformation.

Margarine, some types of peanut butter, and shortening are examples of artificially hydrogenated *trans* fats. Recent studies have shown that an increase in *trans* fats in the human diet may lead to an increase in levels of low-density lipoproteins (LDL), or “bad” cholesterol, which in turn may lead to plaque deposition in the arteries, resulting in heart disease. Many fast food restaurants have recently banned the use of *trans* fats, and food labels are required to display the *trans* fat content.

## Omega Fatty Acids





Figure 6. Alpha-linolenic acid is an example of an omega-3 fatty acid. It has three cis double bonds and, as a result, a curved shape. For clarity, the carbons are not shown. Each singly bonded carbon has two hydrogens associated with it, also not shown.

Essential fatty acids are fatty acids required but not synthesized by the human body. Consequently, they have to be supplemented through ingestion via the diet. Omega-3 fatty acids (like that shown in Figure 6) fall into this category and are one of only two known for humans (the other being omega-6 fatty acid). These are polyunsaturated fatty acids and are called omega-3 because the third carbon from the end of the hydrocarbon chain is connected to its neighboring carbon by a double bond.

The farthest carbon away from the carboxyl group is numbered as the omega ($\omega$) carbon, and if the double bond is between the third and fourth carbon from that end, it is known as an omega-3 fatty acid. Nutritionally important because the body does not make them, omega-3 fatty acids include alpha-linoleic acid (ALA), eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA), all of which are polyunsaturated. Salmon, trout, and tuna are good sources of omega-3 fatty acids. Research indicates that omega-3 fatty acids reduce the risk of sudden death from heart attacks, reduce triglycerides in the blood, lower blood pressure, and prevent thrombosis by inhibiting blood clotting. They also reduce inflammation, and may help reduce the risk of some cancers in animals.

Like carbohydrates, fats have received a lot of bad publicity. It is true that eating an excess of fried foods and other “fatty” foods leads to weight gain. However, fats do have important functions. Many vitamins are fat soluble, and fats serve as a long-term storage form of fatty acids: a source of energy. They also provide insulation for the body. Therefore, “healthy” fats in moderate amounts should be consumed on a regular basis.

## Waxes



Figure 7. Waxy coverings on some leaves are made of lipids. (credit: Roger Griffith)

**Wax** covers the feathers of some aquatic birds and the leaf surfaces of some plants. Because of the hydrophobic nature of waxes, they prevent water from sticking on the surface (Figure 7). Waxes are made up of long fatty acid chains esterified to long-chain alcohols.

## Phospholipids

**Phospholipids** are major constituents of the plasma membrane, the outermost layer of animal cells. Like fats, they are composed of fatty acid chains attached to a glycerol or sphingosine backbone. Instead of three fatty acids attached as in triglycerides, however, there are two fatty acids forming diacylglycerol, and the third carbon of the glycerol backbone is occupied by a modified phosphate group (Figure 8).





Figure 8. A phospholipid is a molecule with two fatty acids and a modified phosphate group attached to a glycerol backbone. The phosphate may be modified by the addition of charged or polar chemical groups. Two chemical groups that may modify the phosphate, choline and serine, are shown here. Both choline and serine attach to the phosphate group at the position labeled R.

A phosphate group alone attached to a diaglycerol does not qualify as a phospholipid; it is phosphatidate (diacylglycerol 3-phosphate), the precursor of phospholipids. The phosphate group is modified by an alcohol. Phosphatidylcholine and phosphatidylserine are two important phospholipids that are found in plasma membranes.A phospholipid is an amphipathic molecule, meaning it has a hydrophobic and a hydrophilic part. The fatty acid chains are hydrophobic and cannot interact with water, whereas the phosphate-containing group is hydrophilic and interacts with water (Figure 9).






Figure 9. The phospholipid bilayer is the major component of all cellular membranes. The hydrophilic head groups of the phospholipids face the aqueous solution. The hydrophobic tails are sequestered in the middle of the bilayer.

The head is the hydrophilic part, and the tail contains the hydrophobic fatty acids. In a membrane, a bilayer of phospholipids forms the matrix of the structure, the fatty acid tails of phospholipids face inside, away from water, whereas the phosphate group faces the outside, aqueous side (Figure 9).


Phospholipids are responsible for the dynamic nature of the plasma membrane. If a drop of phospholipids is placed in water, it spontaneously forms a structure known as a micelle, where the hydrophilic phosphate heads face the outside and the fatty acids face the interior of this structure.

## Steroids

Unlike the phospholipids and fats discussed earlier, steroids have a fused ring structure. Although they do not resemble the other lipids, they are grouped with them because they are also hydrophobic and insoluble in water. All steroids have four linked carbon rings and several of them, like cholesterol, have a short tail (Figure 10). Many steroids also have the –OH functional group, which puts them in the alcohol classification (sterols).




Cholesterol




Cortisol

Figure 10. Steroids such as cholesterol and cortisol are composed of four fused hydrocarbon rings.

Cholesterol is the most common steroid. Cholesterol is mainly synthesized in the liver and is the precursor to many steroid hormones such as testosterone and estradiol, which are secreted by the gonads and endocrine glands. It is also the precursor to Vitamin D. Cholesterol is also the precursor of bile salts, which help in the emulsification of fats and their subsequent absorption by cells. Although cholesterol is often spoken of in negative terms by lay people, it is necessary for proper functioning of the body. It is a component of the plasma membrane of animal cells and is found within the phospholipid bilayer. Being the outermost structure in animal cells, the plasma membrane is responsible for the transport of materials and cellular recognition and it is involved in cell-to-cell communication.

For an additional perspective on lipids, explore this interactive animation.

## Learning Objectives

Lipids are a class of macromolecules that are nonpolar and hydrophobic in nature. Major types include fats and oils, waxes, phospholipids, and steroids. Fats are a stored form of energy and are also known as triacylglycerols or triglycerides. Fats are made up of fatty acids and either glycerol or sphingosine. Fatty acids may be unsaturated or saturated, depending on the presence or absence of double bonds in the hydrocarbon chain. If only single bonds are present, they are known as saturated fatty acids. Unsaturated fatty acids may have one or more double bonds in the hydrocarbon chain. Phospholipids make up the matrix of membranes. They have a glycerol or sphingosine backbone to which two fatty acid chains and a phosphate-containing group are attached. Steroids are another class of lipids. Their basic structure has four fused carbon rings. Cholesterol is a type of steroid and is an important constituent of the plasma membrane, where it helps to maintain the fluid nature of the membrane. It is also the precursor of steroid hormones such as testosterone.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.7: Lipids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.8: Introduction to Carbohydrates

## Summarize the roles carbohydrates play in biological systems

Are carbohydrates good for you? People who wish to lose weight are often told that carbohydrates are bad for them and should be avoided. Some diets completely forbid carbohydrate consumption, claiming that a low-carbohydrate diet helps people to lose weight faster. However, carbohydrates have been an important part of the human diet for thousands of years; artifacts from ancient civilizations show the presence of wheat, rice, and corn in our ancestors' storage areas.

Carbohydrates should be supplemented with proteins, vitamins, and fats to be parts of a well-balanced diet. Calorie-wise, a gram of carbohydrate provides 4.3 Kcal. For comparison, fats provide 9 Kcal/g, a less desirable ratio. Carbohydrates contain soluble and insoluble elements; the insoluble part is known as fiber, which is mostly cellulose. Fiber has many uses; it promotes regular bowel movement by adding bulk, and it regulates the rate of consumption of blood glucose. Fiber also helps to remove excess cholesterol from the body. In addition, a meal containing whole grains and vegetables gives a feeling of fullness. As an immediate source of energy, glucose is broken down during the process of cellular respiration, which produces ATP, the energy currency of the cell. Without the consumption of carbohydrates, the availability of "instant energy" would be reduced.

Eliminating carbohydrates from the diet is not the best way to lose weight. A low-calorie diet that is rich in whole grains, fruits, vegetables, and lean meat, together with plenty of exercise and plenty of water, is the more sensible way to lose weight.

## What You'll Learn to Do

- Distinguish between monosaccharides, disaccharides, and polysaccharides
- Identify several major functions of carbohydrates

### Learning Activities

The learning activities for this section include the following:

- Structure and Function of Carbohydrates
- Self Check: Carbohydrates

## Contributors and Attributions

CC licensed content, Original

- Introduction to Carbohydrates. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.8: Introduction to Carbohydrates is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.9: Structure and Function of Carbohydrates

Most people are familiar with carbohydrates, one type of macromolecule, especially when it comes to what we eat. To lose weight, some individuals adhere to “low-carb” diets. Athletes, in contrast, often “carb-load” before important competitions to ensure that they have enough energy to compete at a high level. Carbohydrates are, in fact, an essential part of our diet; grains, fruits, and vegetables are all natural sources of carbohydrates. Carbohydrates provide energy to the body, particularly through glucose, a simple sugar that is a component of starch and an ingredient in many staple foods. Carbohydrates also have other important functions in humans, animals, and plants.

**Carbohydrates** can be represented by the stoichiometric formula $(CH_2O)n$, where n is the number of carbons in the molecule. In other words, the ratio of carbon to hydrogen to oxygen is 1:2:1 in carbohydrate molecules. This formula also explains the origin of the term “carbohydrate”: the components are carbon (“carbo”) and the components of water (hence, “hydrate”). Carbohydrates are classified into three subtypes: monosaccharides, disaccharides, and polysaccharides.

## Monosaccharides

**Monosaccharides** (*mono*– = “one”; *sacchar*– = “sweet”) are simple sugars, the most common of which is glucose. In monosaccharides, the number of carbons usually ranges from three to seven. Most monosaccharide names end with the suffix –*ose*. If the sugar has an aldehyde group (the functional group with the structure R-CHO), it is known as an aldose, and if it has a ketone group (the functional group with the structure RC(=O)R′), it is known as a ketose. Depending on the number of carbons in the sugar, they also may be known as trioses (three carbons), pentoses (five carbons), and or hexoses (six carbons). See Figure 1 for an illustration of the monosaccharides.





Figure 1. Monosaccharides are classified based on the position of their carbonyl group and the number of carbons in the backbone. Aldoses have a carbonyl group (indicated in green) at the end of the carbon chain, and ketoses have a carbonyl group in the middle of the carbon chain. Trioses, pentoses, and hexoses have three, five, and six carbon backbones, respectively.

The chemical formula for glucose is $C_6H_{12}O_6$. In humans, glucose is an important source of energy. During cellular respiration, energy is released from glucose, and that energy is used to help make adenosine triphosphate (ATP). Plants synthesize glucose using carbon dioxide and water, and glucose in turn is used for energy requirements for the plant. Excess glucose is often stored as starch that is catabolized (the breakdown of larger molecules by cells) by humans and other animals that feed on plants.

Galactose and fructose are other common monosaccharides — galactose is found in milk sugars and fructose is found in fruit sugars. Although glucose, galactose, and fructose all have the same chemical formula ($C_6H_{12}O_6$), they differ structurally and chemically (and are known as isomers) because of the different arrangement of functional groups around the asymmetric carbon; all of these monosaccharides have more than one asymmetric carbon (Figure 2).





Figure 2. Glucose, galactose, and fructose are all hexoses. They are structural isomers, meaning they have the same chemical formula ($C_6H_{12}O_6$) but a different arrangement of atoms.

What kind of sugars are these, aldose or ketose?

[practice-area rows="2"][/practice-area]
[reveal-answer q="972235"]**Show Answer**[/reveal-answer]
[hidden-answer a="972235"]Glucose and galactose are aldoses. Fructose is a ketose.[/hidden-answer]

Monosaccharides can exist as a linear chain or as ring-shaped molecules; in aqueous solutions they are usually found in ring forms (Figure 3). Glucose in a ring form can have two different arrangements of the hydroxyl group (−OH) around the anomeric carbon (carbon 1 that becomes asymmetric in the process of ring formation). If the hydroxyl group is below carbon number 1 in the sugar, it is said to be in the alpha ($\alpha$) position, and if it is above the plane, it is said to be in the beta ($\beta$) position.





Figure 3. Five and six carbon monosaccharides exist in equilibrium between linear and ring forms. When the ring forms, the side chain it closes on is locked into an α or β position. Fructose and ribose also form rings, although they form five-membered rings as opposed to the six-membered ring of glucose.

## Disaccharides

**Disaccharides** (*di–* = "two") form when two monosaccharides undergo a dehydration reaction (also known as a condensation reaction or dehydration synthesis). During this process, the hydroxyl group of one monosaccharide combines with the hydrogen of another monosaccharide, releasing a molecule of water and forming a covalent bond. A covalent bond formed between a carbohydrate molecule and another molecule (in this case, between two monosaccharides) is known as a **glycosidic bond** (Figure 4). Glycosidic bonds (also called glycosidic linkages) can be of the alpha or the beta type.





Figure 4. Sucrose is formed when a monomer of glucose and a monomer of fructose are joined in a dehydration reaction to form a glycosidic bond. In the process, a water molecule is lost. By convention, the carbon atoms in a monosaccharide are numbered from the terminal carbon closest to the carbonyl group. In sucrose, a glycosidic linkage is formed between carbon 1 in glucose and carbon 2 in fructose.

Common disaccharides include lactose, maltose, and sucrose (Figure 5). Lactose is a disaccharide consisting of the monomers glucose and galactose. It is found naturally in milk. Maltose, or malt sugar, is a disaccharide formed by a dehydration reaction between two glucose molecules. The most common disaccharide is sucrose, or table sugar, which is composed of the monomers glucose and fructose.





Figure 5. Common disaccharides include maltose (grain sugar), lactose (milk sugar), and sucrose (table sugar).

## Polysaccharides

A long chain of monosaccharides linked by glycosidic bonds is known as a **polysaccharide** (*poly–* = "many"). The chain may be branched or unbranched, and it may contain different types of monosaccharides. The molecular weight may be 100,000 daltons or more depending on the number of monomers joined. Starch, glycogen, cellulose, and chitin are primary examples of polysaccharides.

Starch is the stored form of sugars in plants and is made up of a mixture of amylose and amylopectin (both polymers of glucose). Plants are able to synthesize glucose, and the excess glucose, beyond the plant's immediate energy needs, is stored as starch in different plant parts, including roots and seeds. The starch in the seeds provides food for the embryo as it germinates and can also act as a source of food for humans and animals. The starch that is consumed by humans is broken down by enzymes, such as salivary amylases, into smaller molecules, such as maltose and glucose. The cells can then absorb the glucose.

Starch is made up of glucose monomers that are joined by $\alpha$ 1-4 or $\alpha$ 1-6 glycosidic bonds. The numbers 1-4 and 1-6 refer to the carbon number of the two residues that have joined to form the bond. As illustrated in Figure 6, amylose is starch formed by unbranched chains of glucose monomers (only $\alpha$ 1-4 linkages), whereas amylopectin is a branched polysaccharide ($\alpha$ 1-6 linkages at the branch points).





Figure 6. Amylose and amylopectin are two different forms of starch. Amylose is composed of unbranched chains of glucose monomers connected by α 1,4 glycosidic linkages. Amylopectin is composed of branched chains of glucose monomers connected by α 1,4 and α 1,6 glycosidic linkages. Because of the way the subunits are joined, the glucose chains have a helical structure. Glycogen (not shown) is similar in structure to amylopectin but more highly branched.

**Glycogen** is the storage form of glucose in humans and other vertebrates and is made up of monomers of glucose. Glycogen is the animal equivalent of starch and is a highly branched molecule usually stored in liver and muscle cells. Whenever blood glucose levels decrease, glycogen is broken down to release glucose in a process known as glycogenolysis.

**Cellulose** is the most abundant natural biopolymer. The cell wall of plants is mostly made of cellulose; this provides structural support to the cell. Wood and paper are mostly cellulosic in nature. Cellulose is made up of glucose monomers that are linked by $\beta$ 1-4 glycosidic bonds (Figure 7).





Figure 7. In cellulose, glucose monomers are linked in unbranched chains by β 1-4 glycosidic linkages. Because of the way the glucose subunits are joined, every glucose monomer is flipped relative to the next one resulting in a linear, fibrous structure.

As shown in Figure 7, every other glucose monomer in cellulose is flipped over, and the monomers are packed tightly as extended long chains. This gives cellulose its rigidity and high tensile strength—which is so important to plant cells. While the *β* 1-4 linkage cannot be broken down by human digestive enzymes, herbivores such as cows, koalas, buffalos, and horses are able, with the help of the specialized flora in their stomach, to digest plant material that is rich in cellulose and use it as a food source. In these animals, certain species of bacteria and protists reside in the rumen (part of the digestive system of herbivores) and secrete the enzyme cellulase. The appendix of grazing animals also contains bacteria that digest cellulose, giving it an important role in the digestive systems of ruminants. Cellulases can break down cellulose into glucose monomers that can be used as an energy source by the animal. Termites are also able to break down cellulose because of the presence of other organisms in their bodies that secrete cellulases.



Figure 8. Insects have a hard outer exoskeleton made of chitin, a type of polysaccharide.

Carbohydrates serve various functions in different animals. Arthropods (insects, crustaceans, and others) have an outer skeleton, called the exoskeleton, which protects their internal body parts (as seen in the bee in Figure 8).

This exoskeleton is made of the biological macromolecule chitin, which is a polysaccharide-containing nitrogen. It is made of repeating units of N-acetyl-*β*-d-glucosamine, a modified sugar. Chitin is also a major component of fungal cell walls; fungi are neither animals nor plants and form a kingdom of their own in the domain Eukarya.

## Learning Objectives

Carbohydrates are a group of macromolecules that are a vital energy source for the cell and provide structural support to plant cells, fungi, and all of the arthropods that include lobsters, crabs, shrimp, insects, and spiders. Carbohydrates are classified as monosaccharides, disaccharides, and polysaccharides depending on the number of monomers in the molecule. Monosaccharides are linked by glycosidic bonds that are formed as a result of dehydration reactions, forming disaccharides and polysaccharides with the elimination of a water molecule for each bond formed. Glucose, galactose, and fructose are common monosaccharides, whereas common disaccharides include lactose, maltose, and sucrose. Starch and glycogen, examples of polysaccharides, are the storage forms of glucose in plants and animals, respectively. The long polysaccharide

chains may be branched or unbranched. Cellulose is an example of an unbranched polysaccharide, whereas amylopectin, a constituent of starch, is a highly branched molecule. Storage of glucose, in the form of polymers like starch of glycogen, makes it slightly less accessible for metabolism; however, this prevents it from leaking out of the cell or creating a high osmotic pressure that could cause excessive water uptake by the cell.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

Public domain content

- Bee On Flowers. **Authored by**: Larisa Koshkina. **Located at**: http://www.publicdomainpictures.net/view-image.php?image=22204. **License**: *Public Domain: No Known Copyright*

5.9: Structure and Function of Carbohydrates is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.10: Introduction to Nucleic Acids

## Discuss nucleic acids and the role they play in DNA and RNA

DNA is the set of instructions for our cells. Our DNA determines who and what we are.

## What You'll Learn to Do

- Describe the basic structure of nucleic acids
- Compare and contrast the structure of DNA and RNA

### Learning Activities

The learning activities for this section include the following:

- Structure of Nucleic Acids
- DNA and RNA
- Self Check: Nucleic Acids

## Contributors and Attributions

CC licensed content, Original

- Introduction to Nucleic Acids. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

5.10: Introduction to Nucleic Acids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.11: Structure of Nucleic Acids

**Nucleic acids** are the most important macromolecules for the continuity of life. They carry the genetic blueprint of a cell and carry instructions for the functioning of the cell.

The two main types of nucleic acids are **deoxyribonucleic acid (DNA)** and **ribonucleic acid (RNA)**. DNA is the genetic material found in all living organisms, ranging from single-celled bacteria to multicellular mammals. It is found in the nucleus of eukaryotes and in the organelles, chloroplasts, and mitochondria. In prokaryotes, the DNA is not enclosed in a membranous envelope.

The entire genetic content of a cell is known as its genome, and the study of genomes is genomics. In eukaryotic cells but not in prokaryotes, DNA forms a complex with histone proteins to form chromatin, the substance of eukaryotic chromosomes. A chromosome may contain tens of thousands of genes. Many genes contain the information to make protein products; other genes code for RNA products. DNA controls all of the cellular activities by turning the genes “on” or “off.”

The other type of nucleic acid, RNA, is mostly involved in protein synthesis. The DNA molecules never leave the nucleus but instead use an intermediary to communicate with the rest of the cell. This intermediary is the **messenger RNA (mRNA)**. Other types of RNA—like rRNA, tRNA, and microRNA—are involved in protein synthesis and its regulation.

DNA and RNA are made up of monomers known as nucleotides. The nucleotides combine with each other to form a polynucleotide, DNA or RNA. Each nucleotide is made up of three components: a nitrogenous base, a pentose (five-carbon) sugar, and a phosphate group (Figure 1). Each nitrogenous base in a nucleotide is attached to a sugar molecule, which is attached to one or more phosphate groups.





Figure 1. A nucleotide is made up of three components: a nitrogenous base, a pentose sugar, and one or more phosphate groups. Carbon residues in the pentose are numbered 1′ through 5′ (the prime distinguishes these residues from those in the base, which are numbered without using a prime notation). The base is attached to the 1′ position of the ribose, and the phosphate is attached to the 5′ position. When a polynucleotide is formed, the 5′ phosphate of the incoming nucleotide attaches to the 3′ hydroxyl group at the end of the growing chain. Two types of pentose are found in nucleotides, deoxyribose (found in DNA) and ribose (found in RNA). Deoxyribose is similar in structure to ribose, but it has an H instead of an OH at the 2′ position. Bases can be divided into two categories: purines and pyrimidines. Purines have a double ring structure, and pyrimidines have a single ring.

The nitrogenous bases, important components of nucleotides, are organic molecules and are so named because they contain carbon and nitrogen. They are bases because they contain an amino group that has the potential of binding an extra hydrogen, and thus, decreases the hydrogen ion concentration in its environment, making it more basic. Each nucleotide in DNA contains one of four possible nitrogenous bases: adenine (A), guanine (G) cytosine (C), and thymine (T). RNA nucleotides also contain one of four possible bases: adenine, guanine, cytosine, and uracil (U) rather than thymine.

Adenine and guanine are classified as **purines**. The primary structure of a purine is two carbon-nitrogen rings. Cytosine, thymine, and uracil are classified as **pyrimidines** which have a single carbon-nitrogen ring as their primary structure (Figure 1). Each of these basic carbon-nitrogen rings has different functional groups attached to it. In molecular biology shorthand, the nitrogenous bases are simply known by their symbols A, T, G, C, and U. DNA contains A, T, G, and C whereas RNA contains A, U, G, and C.

The pentose sugar in DNA is deoxyribose, and in RNA, the sugar is ribose (Figure 1). The difference between the sugars is the presence of the hydroxyl group on the second carbon of the ribose and hydrogen on the second carbon of the deoxyribose. The carbon atoms of the sugar molecule are numbered as 1′, 2′, 3′, 4′, and 5′ (1′ is read as “one prime”). The phosphate residue is attached to the hydroxyl group of the 5′ carbon of one sugar and the hydroxyl group of the 3′ carbon of the sugar of the next nucleotide, which forms a 5′–3′ **phosphodiester** linkage. The phosphodiester linkage is not formed by simple dehydration reaction like the other linkages connecting monomers in macromolecules: its formation involves the removal of two phosphate groups. A polynucleotide may have thousands of such phosphodiester linkages.




Figure 2. DNA is an antiparallel double helix. The phosphate backbone (the curvy lines) is on the outside, and the bases are on the inside. Each base interacts with a base from the opposing strand. (credit: Jerome Walker/Dennis Myts)

DNA has a double-helix structure (Figure 2). The sugar and phosphate lie on the outside of the helix, forming the backbone of the DNA. The nitrogenous bases are stacked in the interior, like the steps of a staircase, in pairs; the pairs are bound to each other by hydrogen bonds. Every base pair in the double helix is separated from the next base pair by 0.34 nm.

The two strands of the helix run in opposite directions, meaning that the 5′ carbon end of one strand will face the 3′ carbon end of its matching strand. (This is referred to as antiparallel orientation and is important to DNA replication and in many nucleic acid interactions.)

Only certain types of base pairing are allowed. For example, a certain purine can only pair with a certain pyrimidine. This means A can pair with T, and G can pair with C, as shown in Figure 3. This is known as the base complementary rule. In other words, the DNA strands are complementary to each other. If the sequence of one strand is AATTGGCC, the complementary strand would have the sequence TTAACCGG. During DNA replication, each strand is copied, resulting in a daughter DNA double helix containing one parental DNA strand and a newly synthesized strand.

Practice Question





Figure 3. In a double stranded DNA molecule, the two strands run antiparallel to one another so that one strand runs 5′ to 3′ and the other 3′ to 5′. The phosphate backbone is located on the outside, and the bases are in the middle. Adenine forms hydrogen bonds (or base pairs) with thymine, and guanine base pairs with cytosine.

A mutation occurs, and cytosine is replaced with adenine. What impact do you think this will have on the DNA structure?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”263813″]**Show Answer**[/reveal-answer]
[hidden-answer a=”263813″]Adenine is larger than cytosine and will not be able to base pair properly with the guanine on the opposing strand. This will cause the DNA to bulge. DNA repair enzymes may recognize the bulge and replace the incorrect nucleotide.[/hidden-answer]

## RNA

Ribonucleic acid, or RNA, is mainly involved in the process of protein synthesis under the direction of DNA. RNA is usually single-stranded and is made of ribonucleotides that are linked by phosphodiester bonds. A ribonucleotide in the RNA chain contains ribose (the pentose sugar), one of the four nitrogenous bases (A, U, G, and C), and the phosphate group.

There are four major types of RNA: messenger RNA (mRNA), ribosomal RNA (rRNA), transfer RNA (tRNA), and microRNA (miRNA). The first, mRNA, carries the message from DNA, which controls all of the cellular activities in a cell. If a cell requires a certain protein to be synthesized, the gene for this product is turned "on" and the messenger RNA is synthesized in the nucleus. The RNA base sequence is complementary to the coding sequence of the DNA from which it has been copied. However, in RNA, the base T is absent and U is present instead. If the DNA strand has a sequence AATTGCGC, the sequence of the complementary RNA is UUAACGCG. In the cytoplasm, the mRNA interacts with ribosomes and other cellular machinery (Figure 4).





Figure 4. A ribosome has two parts: a large subunit and a small subunit. The mRNA sits in between the two subunits. A tRNA molecule recognizes a codon on the mRNA, binds to it by complementary base pairing, and adds the correct amino acid to the growing peptide chain.

The mRNA is read in sets of three bases known as codons. Each codon codes for a single amino acid. In this way, the mRNA is read and the protein product is made. Ribosomal RNA (rRNA) is a major constituent of ribosomes on which the mRNA binds. The rRNA ensures the proper alignment of the mRNA and the ribosomes; the rRNA of the ribosome also has an enzymatic activity (peptidyl transferase) and catalyzes the formation of the peptide bonds between two aligned amino acids. Transfer RNA (tRNA) is one of the smallest of the four types of RNA, usually 70–90 nucleotides long. It carries the correct amino acid to the site of protein synthesis. It is the base pairing between the tRNA and mRNA that allows for the correct amino acid to be inserted in the polypeptide chain. microRNAs are the smallest RNA molecules and their role involves the regulation of gene expression by interfering with the expression of certain mRNA messages.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.11: Structure of Nucleic Acids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.12: DNA and RNA

While DNA and RNA are similar, they have very distinct differences. Table 1 summarizes features of DNA and RNA.

Table 1. Features of DNA and RNA

| | DNA | RNA |
|---|---|---|
| Function | Carries genetic information | Involved in protein synthesis |
| Location | Remains in the nucleus | Leaves the nucleus |
| Structure | DNA is double-stranded “ladder”: sugar-phosphate backbone, with base rungs. | Usually single-stranded |
| Sugar | Deoxyribose | Ribose |
| Pyrimidines | Cytosine, thymine | Cytosine, uracil |
| Purines | Adenine, guanine | Adenine, guanine |

One other difference bears mention. There is only one type of DNA. DNA is the heritable information that is passed along to each generation of cells; its strands can be “unzipped” with small amount of energy when DNA needs to replicate, and DNA is transcribed into RNA. There are mutliple types of RNA: Messenger RNA is a temporary molecule that transports the information necessary to make a protein from the nucleus (where the DNA remains) to the cytoplasm, where the ribosomes are. Other kinds of RNA include ribosomal RNA (rRNA), transfer RNA (tRNA), small nuclear RNA (snRNA), and microRNA.

Even though the RNA is single stranded, most RNA types show extensive intramolecular base pairing between complementary sequences, creating a predictable three-dimensional structure essential for their function.

As you have learned, information flow in an organism takes place from DNA to RNA to protein. DNA dictates the structure of mRNA in a process known as transcription, and RNA dictates the structure of protein in a process known as translation. This is known as the Central Dogma of Life, which holds true for all organisms; however, exceptions to the rule occur in connection with viral infections.

## In Summary: DNA and RNA

Nucleic acids are molecules made up of nucleotides that direct cellular activities such as cell division and protein synthesis. Each nucleotide is made up of a pentose sugar, a nitrogenous base, and a phosphate group. There are two types of nucleic acids: DNA and RNA. DNA carries the genetic blueprint of the cell and is passed on from parents to offspring (in the form of chromosomes). It has a double-helical structure with the two strands running in opposite directions, connected by hydrogen bonds, and complementary to each other. RNA is single-stranded and is made of a pentose sugar (ribose), a nitrogenous base, and a phosphate group. RNA is involved in protein synthesis and its regulation. Messenger RNA (mRNA) is copied from the DNA, is exported from the nucleus to the cytoplasm, and contains information for the construction of proteins. Ribosomal RNA (rRNA) is a part of the ribosmes at the site of protein synthesis, whereas transfer RNA (tRNA) carries the amino acid to the site of protein synthesis. microRNA regulates the use of mRNA for protein synthesis.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

5.12: DNA and RNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.13: Introduction to Comparing Biological Macromolecules

## Discuss biological macromolecules and the differences between the four classes

As we've learned, there are four major classes of biological macromolecules:

- Proteins (polymers of amino acids)
- Carbohydrates (polymers of sugars)
- Lipids (polymers of lipid monomers)
- Nucleic acids (DNA and RNA; polymers of nucleotides)

Let's take a closer look at the differences between the difference classes.

## What You'll Learn to Do

- Define the term "macromolecule"
- Distinguish between the 4 classes of macromolecules

### Learning Activities

The learning activities for this section include the following:

- Different Types of Biological Macromolecules
- Self Check: Carbohydrates

## Contributors and Attributions

CC licensed content, Original


- Introduction to Comparing Biological Macromolecules. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

5.13: Introduction to Comparing Biological Macromolecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.14: Different Types of Biological Macromolecules

Now that we've discussed the four major classes of biological macromolecules (carbohydrates, lipids, proteins, and nucleic acids), let's talk about macromolecules as a whole. Each is an important cell component and performs a wide array of functions. Combined, these molecules make up the majority of a cell's dry mass (recall that water makes up the majority of its complete mass). Biological macromolecules are organic, meaning they contain carbon. In addition, they may contain hydrogen, oxygen, nitrogen, and additional minor elements.

## You are What You Eat




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=160

## Comparing the Biological Macromolecules

| Macromolecule | Basic Formula, key features | Monomer | Examples | Uses |
|---|---|---|---|---|
| Proteins | | Amino acids | Enzymes, some hormones | Storage; Signals; Structural; Contractile; Defensive; Enzyme; Transport; Receptors |
| Lipids | C:H:O<br>Greater than 2:1 H:O (carboxyl group) | Fatty acid and glycerol | Butter, oil, cholesterol, beeswax | Energy storage; Protection; Chemical messengers; Repel water |
| Carbohydrates | C:H:O<br>1:2:1 | Monosaccharides | Glucose, Fructose, Starch, Glycogen, Cellulose | Energy storage; Structure |
| Nucleic Acids | CHONP<br>pentose, nitrogenous base, phosphate | Nucleotides | DNA, RNA | Genetic information |

## Dehydration Synthesis

Most macromolecules are made from single subunits, or building blocks, called **monomers**. The monomers combine with each other using covalent bonds to form larger molecules known as **polymers**. In doing so, monomers release water molecules as byproducts. This type of reaction is known as **dehydration synthesis**, which means "to put together while losing water."





Figure 1. In the dehydration synthesis reaction depicted above, two molecules of glucose are linked together to form the disaccharide maltose. In the process, a water molecule is formed.

In a dehydration synthesis reaction (Figure 1), the hydrogen of one monomer combines with the hydroxyl group of another monomer, releasing a molecule of water. At the same time, the monomers share electrons and form covalent bonds. As additional monomers join, this chain of repeating monomers forms a polymer. Different types of monomers can combine in many configurations, giving rise to a diverse group of macromolecules. Even one kind of monomer can combine in a variety of ways to form several different polymers: for example, glucose monomers are the constituents of starch, glycogen, and cellulose.

## Hydrolysis

Polymers are broken down into monomers in a process known as hydrolysis, which means “to split water,” a reaction in which a water molecule is used during the breakdown (Figure 2). During these reactions, the polymer is broken into two components: one part gains a hydrogen atom (H+) and the other gains a hydroxyl molecule (OH–) from a split water molecule.





Figure 2. In the hydrolysis reaction shown here, the disaccharide maltose is broken down to form two glucose monomers with the addition of a water molecule. Note that this reaction is the reverse of the synthesis reaction shown in Figure 1.

Dehydration and **hydrolysis reactions** are catalyzed, or “sped up,” by specific enzymes; dehydration reactions involve the formation of new bonds, requiring energy, while hydrolysis reactions break bonds and release energy. These reactions are similar for most macromolecules, but each monomer and polymer reaction is specific for its class. For example, in our bodies, food is hydrolyzed, or broken down, into smaller molecules by catalytic enzymes in the digestive system. This allows for easy absorption of nutrients by cells in the intestine. Each macromolecule is broken down by a specific enzyme. For instance, carbohydrates are broken down by amylase, sucrase, lactase, or maltase. Proteins are broken down by the enzymes pepsin and peptidase, and by hydrochloric acid. Lipids are broken down by lipases. Breakdown of these macromolecules provides energy for cellular activities.

Visit this site to see visual representations of dehydration synthesis and hydrolysis.

### Learning Objectives

Proteins, carbohydrates, nucleic acids, and lipids are the four major classes of biological macromolecules—large molecules necessary for life that are built from smaller organic molecules. Macromolecules are made up of single units known as monomers that are joined by covalent bonds to form larger polymers. The polymer is more than the sum of its parts: it acquires new characteristics, and leads to an osmotic pressure that is much lower than that formed by its ingredients; this is an important advantage in the maintenance of cellular osmotic conditions. A monomer joins with another monomer with the release of a water molecule, leading to the formation of a covalent bond. These types of reactions are known as dehydration or condensation reactions. When polymers are broken down into smaller units (monomers), a molecule of water is used for each bond broken by these reactions; such reactions are known as hydrolysis reactions. Dehydration and hydrolysis reactions are similar for all macromolecules, but each monomer and polymer reaction is specific to its class. Dehydration reactions typically require an investment of energy for new bond formation, while hydrolysis reactions typically release energy by breaking bonds.

## Contributors and Attributions

CC licensed content, Original

- Revision, adaptation, and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Biological Molecules - You Are What You Eat: Crash Course Biology #3. **Authored by**: Crash Course. **Located at**: https://youtu.be/H8WJ2KENlK0. **Project**: Crash Course Biology. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

5.14: Different Types of Biological Macromolecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.15: Putting It Together- Important Biological Macromolecules

Now that we've learned about the different macromolecules our bodies need and use, let's return to our the questions we asked at the beginning of the modules about healthy diets:

## Think about It

- Is it even possible for a person to cut all carbs out of his or her diet?
- Is it actually healthy to remove an entire class of molecules from the diet?
- Fats and cholesterol are strictly bad—right?

[practice-area rows="4"][/practice-area]
[reveal-answer q="193414"]See Our Thoughts[/reveal-answer]
[hidden-answer a="193414"]

It is, in fact, impossible to have a no carb or no fat diet. These molecules are in all cells and cells makes up what we eat. More importantly though, each of these biological macromolecules has a very important role to play. If you cut too much fat from your diet, for example, it is possible for your fat stores dip low enough that your hair will fall out!

Even the much maligned cholesterol is a requirement for a healthy body and lifestyle: without sufficient cholesterol, your body doesn't make enough sex hormones (estrogen or testosterone depending on if you're male or female). The trick is to make healthy choices overall without too much of any biological macromolecules—after all, there can certainly be too much of a good thing.

[/hidden-answer]

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

5.15: Putting It Together- Important Biological Macromolecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 5.16: Assignment- Nutritionist for a Day

You have been hired as a high school nutritionist and your first big task is to carry out a “Healthy Lifestyles” nutrition campaign. You want the students you work with to understand the relationship between the four categories of biological macromolecules and the food they eat.

Prepare a brochure or flyer that clearly and accurately educates your students about lipids, carbohydrates, proteins, and nucleic acids. Your flyer should include the following information about each type of macromolecule:

- structural makeup
- overall function in the body
- types of food where each biomolecule can be found

Your flyer should also include an overall theme that emphasizes the importance of a healthy diet.

Take a look at the hints in this worked example for help completing this assessment.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references
- Assignment includes a food source for each biomolecule illustrated.
- Assignment emphasizes an overall healthy diet.

## Rubric

| Nutritionist for a Day |
| --- |
| Outcome: Identify and describe the main features of the four main classes of important biomolecules |

| Nutritionist for a Day | | | | |
|---|---|---|---|---|
| Criteria | Ratings | | | Pts |
| The role of carbohydrates in the body is identified and their structural makeup summarized. | Carbohydrates and their role in the body identified; structural makeup discussed/illustrated in detail in connection to role in the body.<br>5.0 pts | Carbohydrates and their basic role in the body identified; structural makeup summarized/illustrated.<br>4.0 pts | Carbohydrates are not discussed sufficiently.<br>0.0 pts | 5 pts |
| The role of lipids in the body is identified and the structural makeup of each type is summarized. | Lipids and their role in the body identified; structural makeup discussed/illustrated in detail in connection to role in the body.<br>5.0 pts | Lipids and their basic role in the body identified; structural makeup summarized/illustrated.<br>4.0 pts | Lipids are not discussed sufficiently.<br>0.0 pts | 5 pts |
| The role of proteins in the body is identified and their structural makeup summarized. | Proteins and their role in the body identified; structural makeup discussed/illustrated in detail in connection to role in the body.<br>5.0 pts | Proteins and their basic role in the body identified; structural makeup summarized/illustrated.<br>4.0 pts | Proteins are not discussed sufficiently.<br>0.0 pts | 5 pts |
| The role of nucleic acids in the body is identified and the structural makeup of DNA and RNA summarized. | Nucleic acids and their role in the body identified; structural makeup discussed/illustrated in detail in connection to role in the body.<br>5.0 pts | Nucleic acids and their basic role in the body identified; structural makeup summarized/illustrated.<br>4.0 pts | Nucleic acids are not discussed sufficiently.<br>0.0 pts | 5 pts |
| | | | | Total points: 20 |

## Contributors and Attributions

CC licensed content, Original

- Performance Assessment: Nutritionist for a Day. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

5.16: Assignment- Nutritionist for a Day is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 6: Module 4- Cellular Structure


6.1: Why It Matters- Cellular Structure
6.2: Introduction to Microscopes
6.3: Microscopy
6.4: Introduction to Organelles
6.5: Cytoplasm
6.6: Nucleus
6.7: Ribosomes, Mitochondria, and Peroxisomes
6.8: Endomembrane System
6.9: Vesicles
6.10: Unique Features of Animal and Plant Cells
6.11: Introduction to Cytoskeleton
6.12: Microfilaments
6.13: Intermediate Filaments
6.14: Microtubules
6.15: Introduction to Cell Surface
6.16: Cell Membranes
6.17: Cell Junctions
6.18: The Cell Wall
6.19: Putting It Together- Cellular Structure
6.20: Assignment- Cell Builder



6: Module 4- Cellular Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.1: Why It Matters- Cellular Structure

## Why identify and explain a variety of cellular components?

The cell is the basic structural, functional, and biological unit of all known living organisms: all living things are made up of cells (or a single cell, in some cases). However, it's important to remember that cells are simply the smallest unit that can act by themselves. Cells are composed of many different subunits that work in harmony. Each of these different units—membranes, organelles, filaments, etc.—performs a unique function that facilitates life.

**Try It**

When a specific organelle performs incorrectly, it can result in various diseases. For example, the following diseases are linked directly to specific cellular components:

- **Pompe Disease:** characterized by excess accumulation of glycogen in muscle cells
- **Leigh Disease:** progressive disorder of lesions (dead or dying cells) in the brain
- **Emery-Dreifuss muscular dystrophy:** wasting and weakness in muscles of the shoulders, upper arms, and calf muscles

What cellular components could each of these be connected to? Understanding how cellular components work can help you understand health issues related to cellular function.

### Learning Outcomes

- Understand why and how the light microscope and electron microscope are used in biology
- Identify membrane-bound organelles found in eukaryotic cells
- Demonstrate familiarity with various components of the cytoskeleton, including monomeric units
- Demonstrate familiarity with various cell surface specializations

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

6.1: Why It Matters- Cellular Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.2: Introduction to Microscopes

## Understand why and how the light microscope and electron microscope are used in biology

A cell is the smallest unit of a living thing. A living thing, whether made of one cell (like bacteria) or many cells (like a human), is called an organism. Thus, cells are the basic building blocks of all organisms.

Several cells of one kind that interconnect with each other and perform a shared function form tissues, several tissues combine to form an organ (your stomach, heart, or brain), and several organs make up an organ system (such as the digestive system, circulatory system, or nervous system). Several systems that function together form an organism (like a human being). Here, we will examine the structure and function of cells.

There are many types of cells, all grouped into one of two broad categories: prokaryotic and eukaryotic. For example, both animal and plant cells are classified as eukaryotic cells, whereas bacterial cells are classified as prokaryotic. Regardless of the type of cell, be it a component in a multicelled eukaryotic animal's nervous system or a single-celled prokaryotic lifeform, biologists will use microscopes to study them.

## What You'll Learn to Do

- Identify the general applications of light microscopes in biology
- Discuss the advantages and disadvantages of electron microscopes for biological studies

### Learning Activities

The learning activities for this section include the following:

- Microscopy
- Self Check: Microscopes

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.2: Introduction to Microscopes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.3: Microscopy

Cells vary in size. With few exceptions, individual cells cannot be seen with the naked eye, so scientists use microscopes (*micro* = "small"; *scope* = "to look at") to study them. A **microscope** is an instrument that magnifies an object. Most photographs of cells are taken with a microscope, and these images can also be called micrographs.


The optics of a microscope's lenses change the orientation of the image that the user sees. A specimen that is right-side up and facing right on the microscope slide will appear upside-down and facing left when viewed through a microscope, and vice versa. Similarly, if the slide is moved left while looking through the microscope, it will appear to move right, and if moved down, it will seem to move up. This occurs because microscopes use two sets of lenses to magnify the image. Because of the manner by which light travels through the lenses, this system of two lenses produces an inverted image (binocular, or dissecting microscopes, work in a similar manner, but include an additional magnification system that makes the final image appear to be upright).

## Light Microscopes

To give you a sense of cell size, a typical human red blood cell is about eight millionths of a meter or eight micrometers (abbreviated as eight µm) in diameter; the head of a pin of is about two thousandths of a meter (two mm) in diameter. That means about 250 red blood cells could fit on the head of a pin.

Most student microscopes are classified as **light microscopes** (Figure 1a). Visible light passes and is bent through the lens system to enable the user to see the specimen. Light microscopes are advantageous for viewing living organisms, but since individual cells are generally transparent, their components are not distinguishable unless they are colored with special stains. Staining, however, usually kills the cells.

Light microscopes commonly used in the undergraduate college laboratory magnify up to approximately 400 times. Two parameters that are important in microscopy are magnification and resolving power. Magnification is the process of enlarging an object in appearance. Resolving power is the ability of a microscope to distinguish two adjacent structures as separate: the higher the resolution, the better the clarity and detail of the image. When oil immersion lenses are used for the study of small objects, magnification is usually increased to 1,000 times. In order to gain a better understanding of cellular structure and function, scientists typically use electron microscopes.



(a) (b)

Figure 1. (a) Most light microscopes used in a college biology lab can magnify cells up to approximately 400 times and have a resolution of about 200 nanometers. (b) Electron microscopes provide a much higher magnification, 100,000x, and a have a resolution of 50 picometers. (credit a: modification of work by "GcG"/Wikimedia Commons; credit b: modification of work by Evan Bench)

## Electron Microscopes

In contrast to light microscopes, **electron microscopes** (EM) (Figure 1b) use a beam of electrons instead of a beam of light. Not only does this allow for higher magnification (up to 5 million times with some specialized EMs) and, thus, more detail (Figure 2), it also provides higher resolving power. The method used to prepare the specimen for viewing with an electron microscope kills the specimen. Electrons have short wavelengths (shorter than photons) that move best in a vacuum, so living cells cannot be viewed with most electron microscopes.

In a scanning electron microscope, a beam of electrons moves back and forth across a cell's surface, creating details of cell surface characteristics. In a transmission electron microscope, the electron beam penetrates the cell and provides details of a cell's internal structures. As you might imagine, electron microscopes are significantly more bulky and expensive than light microscopes.





Figure 2. (a) These *Salmonella* bacteria appear as tiny purple dots when viewed with a light microscope. (b) This scanning electron microscope micro graph shows *Salmonella* bacteria (in red) invading human cells (yellow). Even though subfigure (b) shows a different *Salmonella* specimen than subfigure (a), you can still observe the comparative increase in magnification and detail. (credit a: modification of work by CDC/Armed Forces Institute of Pathology, Charles N. Farmer, Rocky Mountain Laboratories; credit b: modification of work by NIAID, NIH; scale-bar data from Matt Russell)

For another perspective on cell size, try the HowBig interactive at this site.

## Cell Theory

The microscopes we use today are far more complex than those used in the 1600s by Antony van Leeuwenhoek, a Dutch shopkeeper who had great skill in crafting lenses. Despite the limitations of his now-ancient lenses, van Leeuwenhoek observed the movements of protista (a type of single-celled organism) and sperm, which he collectively termed "animalcules."

In a 1665 publication called *Micrographia*, experimental scientist Robert Hooke coined the term "cell" for the box-like structures he observed when viewing cork tissue through a lens. In the 1670s, van Leeuwenhoek discovered bacteria and protozoa. Later advances in lenses, microscope construction, and staining techniques enabled other scientists to see some components inside cells.

By the late 1830s, botanist Matthias Schleiden and zoologist Theodor Schwann were studying tissues and proposed the **unified cell theory**, which states that all living things are composed of one or more cells, the cell is the basic unit of life, and new cells arise from existing cells. Rudolf Virchow later made important contributions to this theory.





Figure 3. These uterine cervix cells, viewed through a light microscope, were obtained from a Pap smear. Normal cells are on the left. The cells on the right are infected with human papillomavirus (HPV). Notice that the infected cells are larger; also, two of these cells each have two nuclei instead of one, the normal number. (credit: modification of work by Ed Uthman, MD; scale-bar data from Matt Russell)

Have you ever heard of a medical test called a Pap smear (shown in Figure 3)? In this test, a doctor takes a small sample of cells from the uterine cervix of a patient and sends it to a medical lab where a cytotechnologist stains the cells and examines them for any changes that could indicate cervical cancer or a microbial infection.

Cytotechnologists (*cyto* = "cell") are professionals who study cells via microscopic examinations and other laboratory tests. They are trained to determine which cellular changes are within normal limits and which are abnormal. Their focus is not limited to cervical cells; they study cellular specimens that come from all organs. When they notice abnormalities, they consult a pathologist, who is a medical doctor who can make a clinical diagnosis.

Cytotechnologists play a vital role in saving people's lives. When abnormalities are discovered early, a patient's treatment can begin sooner, which usually increases the chances of a successful outcome.

## Learning Objectives

A cell is the smallest unit of life. Most cells are so tiny that they cannot be seen with the naked eye. Therefore, scientists use microscopes to study cells. Electron microscopes provide higher magnification, higher resolution, and more detail than light microscopes. The unified cell theory states that all organisms are composed of one or more cells, the cell is the basic unit of life, and new cells arise from existing cells.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.3: Microscopy is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.4: Introduction to Organelles

## Identify membrane-bound organelles found in eukaryotic cells

Have you ever heard the phrase “form follows function?” It’s a philosophy practiced in many industries. In architecture, this means that buildings should be constructed to support the activities that will be carried out inside them. For example, a skyscraper should be built with several elevator banks; a hospital should be built so that its emergency room is easily accessible.

Our natural world originated the principle of form following function, especially in cell biology, and this will become clear as we explore eukaryotic cells. Unlike prokaryotic cells, **eukaryotic cells** have: (1) a membrane-bound nucleus; (2) numerous membrane-bound **organelles**—such as the endoplasmic reticulum, Golgi apparatus, chloroplasts, mitochondria, and others; and (3) several, rod-shaped chromosomes. Because a eukaryotic cell’s nucleus is surrounded by a membrane, it is often said to have a “true nucleus.” The word “organelle” means “little organ,” and, as already mentioned, organelles have specialized cellular functions, just as the organs of your body have specialized functions.





Figure 1. These figures show the major organelles and other cell components of (a) a typical animal cell and (b) a typical eukaryotic plant cell. The plant cell has a cell wall, chloroplasts, plastids, and a central vacuole—structures not found in animal cells. Plant cells do not have lysosomes or centrosomes.

## What You’ll Learn to Do

- Describe the basic composition of cytoplasm
- Describe the structure and function of the nucleus and nuclear membrane
- Describe the structure and function of ribosomes
- Describe the structure and function of mitochondria
- Describe the structure and function of peroxisomes
- Describe the structure, function, and components of the endomembrane system
- Describe the structure and functions of vesicles
- Identify key organelles present only in animal cells, including centrosomes and lysosomes

- Identify key organelles present only in plant cells, including chloroplasts and vaculoles

### Learning Activities

The learning activities for this section include the following:

- Cytoplasm
- Nucleus
- Ribosomes, Mitochondria, and Peroxisomes
- Endomembrane System
- Vesicles
- Unique Features of Animal and Plant Cells
- Self Check: Organelles

## Contributors and Attributions

CC licensed content, Original

- Revision, adaptation, and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

6.4: Introduction to Organelles is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.5: Cytoplasm

Before we begin looking at individual organelles, we do need to briefly address the matrix in which they sit: the **cytoplasm**. The part of the cell referred to as cytoplasm is slightly different in eukaryotes and prokaryotes. In eukaryotic cells, which have a nucleus, the cytoplasm is everything between the plasma membrane and the nuclear envelope. In prokaryotes, which lack a nucleus, cytoplasm simply means everything found inside the plasma membrane.

One major component of the cytoplasm in both prokaryotes and eukaryotes is the gel-like **cytosol**, a water-based solution that contains ions, small molecules, and macromolecules. In eukaryotes, the cytoplasm also includes membrane-bound organelles, which are suspended in the cytosol. The cytoskeleton, a network of fibers that supports the cell and gives it shape, is also part of the cytoplasm and helps to organize cellular components.

Even though the cytosol is mostly water, it has a semi-solid, Jello-like consistency because of the many proteins suspended in it. The cytosol contains a rich broth of macromolecules and smaller organic molecules, including glucose and other simple sugars, polysaccharides, amino acids, nucleic acids, and fatty acids. Ions of sodium, potassium, calcium, and other elements are also found in the cytosol. Many metabolic reactions, including protein synthesis, take place in this part of the cell.

## Contributors and Attributions

CC licensed content, Shared previously

- Plasma membrane and cytoplasm. **Provided by**: Khan Academy . **Located at**: https://www.khanacademy.org/science/biology/structure-of-a-cell/prokaryotic-and-eukaryotic-cells/a/plasma-membrane-and-cytoplasm. **Project**: Biology. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

6.5: Cytoplasm is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.6: Nucleus





Figure 1. The nucleus stores chromatin (DNA plus proteins) in a gel-like substance called the nucleoplasm. The nucleolus is a condensed region of chromatin where ribosome synthesis occurs. The boundary of the nucleus is called the nuclear envelope. It consists of two phospholipid bilayers: an outer membrane and an inner membrane. The nuclear membrane is continuous with the endoplasmic reticulum. Nuclear pores allow substances to enter and exit the nucleus.

Typically, the nucleus is the most prominent organelle in a cell. The **nucleus** (plural = *nuclei*) houses the cell's DNA and directs the synthesis of ribosomes and proteins. Let's look at it in more detail (Figure 1).

## Nuclear Envelope

The **nuclear envelope** is a double-membrane structure that constitutes the outermost portion of the nucleus (Figure 1). Both the inner and outer membranes of the nuclear envelope are phospholipid bilayers.

The nuclear envelope is punctuated with pores that control the passage of ions, molecules, and RNA between the nucleoplasm and cytoplasm. The **nucleoplasm** is the semi-solid fluid inside the nucleus, where we find the chromatin and the nucleolus.

## Chromatin and Chromosomes

To understand chromatin, it is helpful to first consider chromosomes. **Chromosomes** are structures within the nucleus that are made up of DNA, the hereditary material. In prokaryotes, DNA is organized into a single circular chromosome. In eukaryotes, chromosomes are linear structures. Every eukaryotic species has a specific number of chromosomes in the nuclei of its body's cells. For example, in humans, the chromosome number is 46, while in fruit flies, it is eight. Chromosomes are only visible and distinguishable from one another when the cell is getting ready to divide. When the cell is in the growth and maintenance phases of its life cycle, proteins are attached to chromosomes, and they resemble an unwound, jumbled bunch of threads. These unwound protein-chromosome complexes are called **chromatin** (Figure 2); chromatin describes the material that makes up the chromosomes both when condensed and decondensed. We will focus on chromatin and chromosomes in greater detail later.





Figure 2. (a) This image shows various levels of the organization of chromatin (DNA and protein). (b) This image shows paired chromosomes. (credit b: modification of work by NIH; scale-bar data from Matt Russell)

## Nucleolus

We already know that the nucleus directs the synthesis of ribosomes, but how does it do this? Some chromosomes have sections of DNA that encode ribosomal RNA. A darkly staining area within the nucleus called the nucleolus (plural = nucleoli) aggregates the ribosomal RNA with associated proteins to assemble the ribosomal subunits that are then transported out through the pores in the nuclear envelope to the cytoplasm.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

6.6: Nucleus is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.7: Ribosomes, Mitochondria, and Peroxisomes

## Ribosomes





Figure 1. Ribosomes are made up of a large subunit (top) and a small subunit (bottom). During protein synthesis, ribosomes assemble amino acids into proteins.

**Ribosomes** are the cellular structures responsible for protein synthesis. When viewed through an electron microscope, ribosomes appear either as clusters (polyribosomes) or single, tiny dots that float freely in the cytoplasm. They may be attached to the cytoplasmic side of the plasma membrane or the cytoplasmic side of the endoplasmic reticulum and the outer membrane of the nuclear envelope. Electron microscopy has shown us that ribosomes, which are large complexes of protein and RNA, consist of two subunits, aptly called large and small (Figure 1). Ribosomes receive their "orders" for protein synthesis from the nucleus where the DNA is transcribed into messenger RNA (mRNA). The mRNA travels to the ribosomes, which translate the code provided by the sequence of the nitrogenous bases in the mRNA into a specific order of amino acids in a protein. Amino acids are the building blocks of proteins.

Because proteins synthesis is an essential function of all cells, ribosomes are found in practically every cell. Ribosomes are particularly abundant in cells that synthesize large amounts of protein. For example, the pancreas is responsible for creating several digestive enzymes and the cells that produce these enzymes contain many ribosomes. Thus, we see another example of form following function.

## Mitochondria

**Mitochondria** (singular = *mitochondrion*) are often called the "powerhouses" or "energy factories" of a cell because they are responsible for making adenosine triphosphate (ATP), the cell's main energy-carrying molecule. ATP represents the short-term stored energy of the cell. Cellular respiration is the process of making ATP using the chemical energy found in glucose and other nutrients. In mitochondria, this process uses oxygen and produces carbon dioxide as a waste product. In fact, the carbon dioxide that you exhale with every breath comes from the cellular reactions that produce carbon dioxide as a byproduct.

In keeping with our theme of form following function, it is important to point out that muscle cells have a very high concentration of mitochondria that produce ATP. Your muscle cells need a lot of energy to keep your body moving. When your cells don't get enough oxygen, they do not make a lot of ATP. Instead, the small amount of ATP they make in the absence of oxygen is accompanied by the production of lactic acid.





Figure 2. This electron micrograph shows a mitochondrion as viewed with a transmission electron microscope. This organelle has an outer membrane and an inner membrane. The inner membrane contains folds, called cristae, which increase its surface area. The space between the two membranes is called the intermembrane space, and the space inside the inner membrane is called the mitochondrial matrix. ATP synthesis takes place on the inner membrane. (credit: modification of work by Matthew Britton; scale-bar data from Matt Russell)

Mitochondria are oval-shaped, double membrane organelles (Figure 2) that have their own DNA and ribosomes (we'll talk about these later!). Each membrane is a phospholipid bilayer embedded with proteins. The inner layer has folds called cristae. The area surrounded by the folds is called the mitochondrial matrix. The cristae and the matrix have different roles in cellular respiration.

## Peroxisomes

Peroxisomes are small, round organelles enclosed by single membranes. They carry out oxidation reactions that break down fatty acids and amino acids. They also detoxify many poisons that may enter the body. Many of these oxidation reactions release hydrogen peroxide, $H_2O_2$, which would be damaging to cells; however, when these reactions are confined to peroxisomes, enzymes safely break down the $H_2O_2$ into oxygen and water. For example, alcohol is detoxified by peroxisomes in liver cells. Glyoxysomes, which are specialized peroxisomes in plants, are responsible for converting stored fats into sugars.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.7: Ribosomes, Mitochondria, and Peroxisomes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.8: Endomembrane System





Figure 1. Membrane and secretory proteins are synthesized in the rough endoplasmic reticulum (RER). The RER's membrane also sometimes modifies proteins. (credit: modification of work by Magnus Manske)

The endomembrane system (*endo* = "within") is a group of membranes and organelles (Figure 1) in eukaryotic cells that works together to modify, package, and transport lipids and proteins. It includes the nuclear envelope, lysosomes, vesicles, and the endoplasmic reticulum and Golgi apparatus, which we will cover shortly. Although not technically *within* the cell, the plasma membrane is included in the endomembrane system because, as you will see, it interacts with the other endomembranous organelles. The endomembrane system does not include the membranes of either mitochondria or chloroplasts.

Figure 1 illustrates the connections of the endomembrane system as a (green) integral membrane protein in the ER is modified by attachment of a (purple) carbohydrate. Vesicles with the integral protein bud from the ER and fuse with the *cis* face of the Golgi apparatus. As the protein passes along the Golgi's cisternae, it is further modified by the addition of more carbohydrates. After its synthesis is complete, it exits as integral membrane protein of the vesicle that bud from the Golgi's *trans* face and when the vesicle fuses with the cell membrane the protein becomes integral portion of that cell membrane.

## Endoplasmic Reticulum

The endoplasmic reticulum (ER) (Figure 1) is a series of interconnected membranous sacs and tubules that collectively modifies proteins and synthesizes lipids. However, these two functions are performed in separate areas of the ER: the rough ER and the smooth ER, respectively.

The hollow portion of the ER tubules is called the lumen or cisternal space. The membrane of the ER, which is a phospholipid bilayer embedded with proteins, is continuous with the nuclear envelope.

### Rough ER

The rough endoplasmic reticulum (RER) is so named because the ribosomes attached to its cytoplasmic surface give it a studded appearance when viewed through an electron microscope (Figure 2).





Figure 2. This transmission electron micrograph shows the rough endoplasmic reticulum and other organelles in a pancreatic cell. (credit: modification of work by Louisa Howard)

Ribosomes transfer their newly synthesized proteins into the lumen of the RER where they undergo structural modifications, such as folding or the acquisition of side chains. These modified proteins will be incorporated into cellular membranes—the membrane of the ER or those of other organelles—or secreted from the cell (such as protein hormones or enzymes). The RER also makes phospholipids for cellular membranes.

If the phospholipids or modified proteins are not destined to stay in the RER, they will reach their destinations via transport vesicles that bud from the RER's membrane (Figure 1).

Since the RER is engaged in modifying proteins (such as enzymes, for example) that will be secreted from the cell, you would be correct in assuming that the RER is abundant in cells that secrete proteins. This is the case with cells of the liver, for example.

### Smooth ER

The smooth endoplasmic reticulum (SER) is continuous with the RER but has few or no ribosomes on its cytoplasmic surface. Functions of the SER include synthesis of carbohydrates, lipids, and steroid hormones; detoxification of medications and poisons; and storage of calcium ions.

In muscle cells, a specialized SER called the sarcoplasmic reticulum is responsible for storage of the calcium ions that are needed to trigger the coordinated contractions of the muscle cells.

**Try It**

Heart disease is the leading cause of death in the United States. This is primarily due to our sedentary lifestyle and our high trans-fat diets.

Heart failure is just one of many disabling heart conditions. Heart failure does not mean that the heart has stopped working. Rather, it means that the heart can't pump with sufficient force to transport oxygenated blood to all the vital organs. Left untreated, heart failure can lead to kidney failure and failure of other organs.

The wall of the heart is composed of cardiac muscle tissue. Heart failure occurs when the endoplasmic reticula of cardiac muscle cells do not function properly. As a result, an insufficient number of calcium ions are available to trigger a sufficient contractile force.

Cardiologists (*cardi* = "heart"; *ologist* = "one who studies") are doctors who specialize in treating heart diseases, including heart failure. Cardiologists can make a diagnosis of heart failure via physical examination, results from an electrocardiogram (ECG, a test that measures the electrical activity of the heart), a chest X-ray to see whether the heart is enlarged, and other tests. If heart failure is diagnosed, the cardiologist will typically prescribe appropriate medications and recommend a reduction in table salt intake and a supervised exercise program.

## Golgi Apparatus





Figure 3. The Golgi apparatus in this white blood cell is visible as a stack of semicircular, flattened rings in the lower portion of the image. Several vesicles can be seen near the Golgi apparatus. (credit: modification of work by Louisa Howard)

We have already mentioned that vesicles can bud from the ER and transport their contents elsewhere, but where do the vesicles go? Before reaching their final destination, the lipids or proteins within the transport vesicles still need to be sorted, packaged, and tagged so that they wind up in the right place. Sorting, tagging, packaging, and distribution of lipids and proteins takes place in the Golgi apparatus (also called the Golgi body), a series of flattened membranes (Figure 3).

The receiving side of the Golgi apparatus is called the *cis* face. The opposite side is called the *trans* face. The transport vesicles that formed from the ER travel to the *cis* face, fuse with it, and empty their contents into the lumen of the Golgi apparatus. As the proteins and lipids travel through the Golgi, they undergo further modifications that allow them to be sorted. The most frequent modification is the addition of short chains of sugar molecules. These newly modified proteins and lipids are then tagged with phosphate groups or other small molecules so that they can be routed to their proper destinations.

Finally, the modified and tagged proteins are packaged into secretory vesicles that bud from the *trans* face of the Golgi. While some of these vesicles deposit their contents into other parts of the cell where they will be used, other secretory vesicles fuse with the plasma membrane and release their contents outside the cell.

In another example of form following function, cells that engage in a great deal of secretory activity (such as cells of the salivary glands that secrete digestive enzymes or cells of the immune system that secrete antibodies) have an abundance of Golgi.

In plant cells, the Golgi apparatus has the additional role of synthesizing polysaccharides, some of which are incorporated into the cell wall and some of which are used in other parts of the cell.

### Try It

Many diseases arise from genetic mutations that prevent the synthesis of critical proteins. One such disease is Lowe disease (also called oculocerebrorenal syndrome, because it affects the eyes, brain, and kidneys). In Lowe disease, there is a deficiency in an enzyme localized to the Golgi apparatus. Children with Lowe disease are born with cataracts, typically develop kidney disease after the first year of life, and may have impaired mental abilities.

Lowe disease is a genetic disease caused by a mutation on the X chromosome. The X chromosome is one of the two human sex chromosome, as these chromosomes determine a person's sex. Females possess two X chromosomes while males possess one X and one Y chromosome. In females, the genes on only one of the two X chromosomes are expressed. Therefore, females who carry the Lowe disease gene on one of their X chromosomes have a 50/50 chance of having the disease. However, males only have one X chromosome and the genes on this chromosome are always expressed. Therefore, males will always have Lowe disease if their X chromosome carries the Lowe disease gene. The location of the mutated gene, as well as the locations of many other mutations that cause genetic diseases, has now been identified. Through prenatal testing, a woman can find out if the fetus she is carrying may be afflicted with one of several genetic diseases.

Geneticists analyze the results of prenatal genetic tests and may counsel pregnant women on available options. They may also conduct genetic research that leads to new drugs or foods, or perform DNA analyses that are used in forensic investigations.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.8: Endomembrane System is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.9: Vesicles

**Vesicles**, like vacuoles (which we will address soon), are membrane-bound sacs that function in storage and transport. Other than the fact that vacuoles are somewhat larger than vesicles, there is a very subtle distinction between them: the membranes of vesicles can fuse with either the plasma membrane or other membrane systems within the cell. The membrane of a vacuole does not fuse with the membranes of other cellular components.

A vesicle consists of fluid enclosed by a lipid bilayer. Vesicles form naturally during the processes of secretion (exocytosis), uptake (phagocytosis) and transport of materials within the cytoplasm. Alternatively, they may be prepared artificially, in which case they are called liposomes. Vesicles can fuse with the plasma membrane to release their contents outside the cell. Vesicles can also fuse with other organelles within the cell.

Vesicles perform a variety of functions. Because they are separated from the cytosol, the inside of a vesicle can be different from the cytosolic environment. For this reason, vesicles are a basic tool used by the cell for organizing cellular substances. Vesicles are involved in metabolism, transport, buoyancy control, and enzyme storage. They can also act as chemical reaction chambers.

## Contributors and Attributions

CC licensed content, Original

- Revision and adaptation. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Vesicles and Vacuoles. **Provided by**: Boundless. **Located at**: https://courses.lumenlearning.com/boundless-biology/. **License**: *CC BY-SA: Attribution-ShareAlike*

6.9: Vesicles is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.10: Unique Features of Animal and Plant Cells

At this point, you know that each eukaryotic cell has a plasma membrane, cytoplasm, a nucleus, ribosomes, mitochondria, peroxisomes, and in some, vacuoles, but there are some striking differences between animal and plant cells. While both animal and plant cells have microtubule organizing centers (MTOCs), animal cells also have centrioles associated with the MTOC: a complex called the centrosome. Animal cells each have a centrosome and lysosomes, whereas plant cells do not. Plant cells have a cell wall, chloroplasts and other specialized plastids, and a large central vacuole, whereas animal cells do not.

## Properties of Animal Cells





Figure 1. The centrosome consists of two centrioles that lie at right angles to each other. Each centriole is a cylinder made up of nine triplets of microtubules. Nontubulin proteins (indicated by the green lines) hold the microtubule triplets together.

### Centrosome

The centrosome is a microtubule-organizing center found near the nuclei of animal cells. It contains a pair of centrioles, two structures that lie perpendicular to each other (Figure 1). Each centriole is a cylinder of nine triplets of microtubules.

The centrosome (the organelle where all microtubules originate) replicates itself before a cell divides, and the centrioles appear to have some role in pulling the duplicated chromosomes to opposite ends of the dividing cell. However, the exact function of the centrioles in cell division isn't clear, because cells that have had the centrosome removed can still divide, and plant cells, which lack centrosomes, are capable of cell division.

### Lysosomes





Figure 2. A macrophage has engulfed (phagocytized) a potentially pathogenic bacterium and then fuses with a lysosomes within the cell to destroy the pathogen. Other organelles are present in the cell but for simplicity are not shown.

In addition to their role as the digestive component and organelle-recycling facility of animal cells, lysosomes are considered to be parts of the endomembrane system.

Lysosomes also use their hydrolytic enzymes to destroy pathogens (disease-causing organisms) that might enter the cell. A good example of this occurs in a group of white blood cells called macrophages, which are part of your body's immune system. In a process known as phagocytosis or endocytosis, a section of the plasma membrane of the macrophage invaginates (folds in) and engulfs a pathogen. The invaginated section, with the pathogen inside, then pinches itself off from the plasma membrane and becomes a vesicle. The vesicle fuses with a lysosome. The lysosome's hydrolytic enzymes then destroy the pathogen (Figure 2).

## Chloroplasts





Figure 3. The chloroplast has an outer membrane, an inner membrane, and membrane structures called thylakoids that are stacked into grana. The space inside the thylakoid membranes is called the thylakoid space. The light harvesting reactions take place in the thylakoid membranes, and the synthesis of sugar takes place in the fluid inside the inner membrane, which is called the stroma. Chloroplasts also have their own genome, which is contained on a single circular chromosome.

Like the mitochondria, chloroplasts have their own DNA and ribosomes (we'll talk about these later!), but chloroplasts have an entirely different function. Chloroplasts are plant cell organelles that carry out photosynthesis. Photosynthesis is the series of reactions that use carbon dioxide, water, and light energy to make glucose and oxygen. This is a major difference between plants and animals; plants (autotrophs) are able to make their own food, like sugars, while animals (heterotrophs) must ingest their food.

Like mitochondria, chloroplasts have outer and inner membranes, but within the space enclosed by a chloroplast's inner membrane is a set of interconnected and stacked fluid-filled membrane sacs called thylakoids (Figure 3). Each stack of thylakoids is called a granum (plural = grana). The fluid enclosed by the inner membrane that surrounds the grana is called the stroma.

The chloroplasts contain a green pigment called chlorophyll, which captures the light energy that drives the reactions of photosynthesis. Like plant cells, photosynthetic protists also have chloroplasts. Some bacteria perform photosynthesis, but their chlorophyll is not relegated to an organelle.

**Try It**

Click through this activity to learn more about chloroplasts and how they work.

A link to an interactive elements can be found at the bottom of this page.

**Try It**

We have mentioned that both mitochondria and chloroplasts contain DNA and ribosomes. Have you wondered why? Strong evidence points to endosymbiosis as the explanation.

Symbiosis is a relationship in which organisms from two separate species depend on each other for their survival. Endosymbiosis (*endo–* = "within") is a mutually beneficial relationship in which one organism lives inside the other. Endosymbiotic relationships abound in nature. We have already mentioned that microbes that produce vitamin K live inside the human gut. This relationship is beneficial for us because we are unable to synthesize vitamin K. It is also beneficial for the microbes because they are protected from other organisms and from drying out, and they receive abundant food from the environment of the large intestine.

Scientists have long noticed that bacteria, mitochondria, and chloroplasts are similar in size. We also know that bacteria have DNA and ribosomes, just as mitochondria and chloroplasts do. Scientists believe that host cells and bacteria formed an endosymbiotic relationship when the host cells ingested both aerobic and autotrophic bacteria (cyanobacteria) but did not destroy them. Through many millions of years of evolution, these ingested bacteria became more specialized in their functions, with the aerobic bacteria becoming mitochondria and the autotrophic bacteria becoming chloroplasts.

## Vacuoles

**Vacuoles** are membrane-bound sacs that function in storage and transport. The membrane of a vacuole does not fuse with the membranes of other cellular components. Additionally, some agents such as enzymes within plant vacuoles break down macromolecules.

If you look at Figure 4b, you will see that plant cells each have a large central vacuole that occupies most of the area of the cell. The **central vacuole** plays a key role in regulating the cell's concentration of water in changing environmental conditions. Have you ever noticed that if you forget to water a plant for a few days, it wilts? That's because as the water concentration in the soil becomes lower than the water concentration in the plant, water moves out of the central vacuoles and cytoplasm. As the central vacuole shrinks, it leaves the cell wall unsupported. This loss of support to the cell walls of plant cells results in the wilted appearance of the plant.

The central vacuole also supports the expansion of the cell. When the central vacuole holds more water, the cell gets larger without having to invest a lot of energy in synthesizing new cytoplasm. You can rescue wilted celery in your refrigerator using this process. Simply cut the end off the stalks and place them in a cup of water. Soon the celery will be stiff and crunchy again.





Figure 4. These figures show the major organelles and other cell components of (a) a typical animal cell and (b) a typical eukaryotic plant cell. The plant cell has a cell wall, chloroplasts, plastids, and a central vacuole—structures not found in animal cells. Plant cells do not have lysosomes or centrosomes.

## Contributors and Attributions

CC licensed content, Original

- Interacting with Plant Organelles. **Authored by**: Ting Chen and Lumen Learning. **Project**: This interactive was originally authored by TIng Chen as a student in Dr. William Carr's Biology for Majors I course at Medgar Evers College in fall 2019, revised by Lumen Learning in Spring 2020.. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.10: Unique Features of Animal and Plant Cells is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.11: Introduction to Cytoskeleton

## Demonstrate familiarity with various components of the cytoskeleton, including monomeric units

If you were to remove all the organelles from a cell, would the plasma membrane and the cytoplasm be the only components left? No. Within the cytoplasm, there would still be ions and organic molecules, plus a network of protein fibers that help maintain the shape of the cell, secure some organelles in specific positions, allow cytoplasm and vesicles to move within the cell, and enable cells within multicellular organisms to move. Collectively, this network of protein fibers is known as the **cytoskeleton**. There are three types of fibers within the cytoskeleton: microfilaments, intermediate filaments, and microtubules (Figure 1). In this outcome, we will examine each.





Figure 1. Microfilaments thicken the cortex around the inner edge of a cell; like rubber bands, they resist tension. Microtubules are found in the interior of the cell where they maintain cell shape by resisting compressive forces. Intermediate filaments are found throughout the cell and hold organelles in place.

## What You'll Learn to Do

- Describe the structure and function of microfilaments
- Describe the structure and function of intermediate filaments
- Describe the structure and function of microtubules

### Learning Activities

The learning activities for this section include the following:

- Reading: Microfilaments
- Reading: Intermediate Filaments
- Reading: Microtubules
- Self Check: Cytoskeleton

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.11: Introduction to Cytoskeleton is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.12: Microfilaments





Figure 1. Microfilaments are made of two intertwined strands of actin.

Of the three types of protein fibers in the cytoskeleton, **microfilaments** are the narrowest. They function in cellular movement, have a diameter of about 7 nm, and are made of two intertwined strands of a globular protein called actin (Figure 1). For this reason, microfilaments are also known as actin filaments.

Actin is powered by ATP to assemble its filamentous form, which serves as a track for the movement of a motor protein called myosin. This enables actin to engage in cellular events requiring motion, such as cell division in animal cells and cytoplasmic streaming, which is the circular movement of the cell cytoplasm in plant cells. Actin and myosin are plentiful in muscle cells. When your actin and myosin filaments slide past each other, your muscles contract.

Microfilaments also provide some rigidity and shape to the cell. They can depolymerize (disassemble) and reform quickly, thus enabling a cell to change its shape and move. White blood cells (your body's infection-fighting cells) make good use of this ability. They can move to the site of an infection and phagocytize the pathogen.

Look below see an example of a white blood cell in action. Watch this short time-lapse video of the cell capturing two bacteria. It engulfs one and then moves on to the other. Note that this video has no audio.

A link to an interactive elements can be found at the bottom of this page.

You can view the audio description text for "White Blood Cell Chases Bacteria" here (link opens in new window).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- white blood cell chases bacteria. **Authored by**: Brad Cameron. **Located at**: https://youtu.be/AUih856vaQY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

6.12: Microfilaments is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.13: Intermediate Filaments

Intermediate filaments are made of several strands of fibrous proteins that are wound together (Figure 1). These elements of the cytoskeleton get their name from the fact that their diameter, 8 to 10 nm, is between those of microfilaments and microtubules.



Figure 1. Intermediate filaments consist of several intertwined strands of fibrous proteins.

Intermediate filaments have no role in cell movement. Their function is purely structural. They bear tension, thus maintaining the shape of the cell, and anchor the nucleus and other organelles in place. Figure 2 shows how intermediate filaments create a supportive scaffolding inside the cell.

The intermediate filaments are the most diverse group of cytoskeletal elements. Several types of fibrous proteins are found in the intermediate filaments. You are probably most familiar with keratin, the fibrous protein that strengthens your hair, nails, and the epidermis of the skin.





Figure 2. Microfilaments thicken the cortex around the inner edge of a cell; like rubber bands, they resist tension. Microtubules are found in the interior of the cell where they maintain cell shape by resisting compressive forces. Intermediate filaments are found throughout the cell and hold organelles in place.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.13: Intermediate Filaments is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.14: Microtubules

As their name implies, microtubules are small hollow tubes. The walls of the microtubule are made of polymerized dimers of *α*-tubulin and *β*-tubulin, two globular proteins (Figure 1). With a diameter of about 25 nm, microtubules are the widest components of the cytoskeleton. They help the cell resist compression, provide a track along which vesicles move through the cell, and pull replicated chromosomes to opposite ends of a dividing cell. Like microfilaments, microtubules can dissolve and reform quickly.





Figure 1. Microtubules are hollow. Their walls consist of 13 polymerized dimers of α-tubulin and β-tubulin (right image). The left image shows the molecular structure of the tube.

Microtubules are also the structural elements of flagella, cilia, and centrioles (the latter are the two perpendicular bodies of the centrosome). In fact, in animal cells, the centrosome is the microtubule-organizing center. In eukaryotic cells, flagella and cilia are quite different structurally from their counterparts in prokaryotes, as discussed below.

## Flagella and Cilia





Figure 2. This transmission electron micrograph of two flagella shows the 9 + 2 array of microtubules (credit: modification of work by Dartmouth Electron Microscope Facility, Dartmouth College; scale-bar data from Matt Russell)

To refresh your memory, flagella (singular = *flagellum*) are long, hair-like structures that extend from the plasma membrane and are used to move an entire cell (for example, sperm, *Euglena*). When present, the cell has just one flagellum or a few flagella. When cilia (singular = *cilium*) are present, however, many of them extend along the entire surface of the plasma membrane. They are short, hair-like structures that are used to move entire cells (such as paramecia) or substances along the outer surface of the cell (for example, the cilia of cells lining the Fallopian tubes that move the ovum toward the uterus, or cilia lining the cells of the respiratory tract that trap particulate matter and move it toward your nostrils.)

Despite their differences in length and number, flagella and cilia share a common structural arrangement of microtubules called a "9 + 2 array." This is an appropriate name because a single flagellum or cilium is made of a ring of nine microtubule doublets, surrounding a single microtubule doublet in the center (Figure 2).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

6.14: Microtubules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.15: Introduction to Cell Surface

## Demonstrate familiarity with various cell surface specializations

In this outcome, we'll learn about the cell surface including the cell membrane, cell junctions, and the cell wall.

## What You'll Learn to Do

- Explain the structure and function of cell membranes
- Describe cell junctions found in plant cells (plasmodesmata) and animal cells (tight junctions, desmosomes, gap junctions)
- Describe the structure and function of the cell wall

### Learning Activities

The learning activities for this section include the following:

- Cell Membranes
- Cell Junctions
- The Cell Wall
- Self Check: Cell Surface

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

6.15: Introduction to Cell Surface is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.16: Cell Membranes

Both prokaryotic and eukaryotic cells have a **plasma membrane** (Figure 1), a phospholipid bilayer with embedded proteins, that separates the internal contents of the cell from its surrounding environment. A phospholipid is a lipid molecule with two fatty acid chains and a phosphate-containing group. The plasma membrane controls the passage of organic molecules, ions, water, and oxygen into and out of the cell. Wastes (such as carbon dioxide and ammonia) also leave the cell by passing through the plasma membrane. We will cover the plasma membrane in more detail in a later unit but here is an overview of this cell surface structure.





Figure 1. The eukaryotic plasma membrane is a phospholipid bilayer with proteins and cholesterol embedded in it.

The plasma membranes of cells that specialize in absorption are folded into fingerlike projections called *microvilli* (singular = *microvillus*); (Figure 2). Such cells are typically found lining the small intestine, the organ that absorbs nutrients from digested food. This is an excellent example of form following function. People with celiac disease have an immune response to gluten, which is a protein found in wheat, barley, and rye. The immune response damages microvilli, and thus, afflicted individuals cannot absorb nutrients. This leads to malnutrition, cramping, and diarrhea. Patients suffering from celiac disease must follow a gluten-free diet.





Figure 2. Microvilli, shown here as they appear on cells lining the small intestine, increase the surface area available for absorption. These microvilli are only found on the area of the plasma membrane that faces the cavity from which substances will be absorbed. (credit “micrograph”: modification of work by Louisa Howard)

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.16: Cell Membranes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.17: Cell Junctions

## Extracellular Matrix of Animal Cells

Most animal cells release materials into the extracellular space. The primary components of these materials are proteins, and the most abundant protein is collagen. Collagen fibers are interwoven with carbohydrate-containing protein molecules called proteoglycans. Collectively, these materials are called the **extracellular matrix** (Figure 1). Not only does the extracellular matrix hold the cells together to form a tissue, but it also allows the cells within the tissue to communicate with each other. How can this happen?





Figure 1. The extracellular matrix consists of a network of proteins and carbohydrates.

Cells have protein receptors on the extracellular surfaces of their plasma membranes. When a molecule within the matrix binds to the receptor, it changes the molecular structure of the receptor. The receptor, in turn, changes the conformation of the microfilaments positioned just inside the plasma membrane. These conformational changes induce chemical signals inside the cell that reach the nucleus and turn “on” or “off” the transcription of specific sections of DNA, which affects the production of associated proteins, thus changing the activities within the cell.

Blood clotting provides an example of the role of the extracellular matrix in cell communication. When the cells lining a blood vessel are damaged, they display a protein receptor called tissue factor. When tissue factor binds with another factor in the extracellular matrix, it causes platelets to adhere to the wall of the damaged blood vessel, stimulates the adjacent smooth muscle cells in the blood vessel to contract (thus constricting the blood vessel), and initiates a series of steps that stimulate the platelets to produce clotting factors.

## Intercellular Junctions

Cells can also communicate with each other via direct contact, referred to as intercellular junctions. There are some differences in the ways that plant and animal cells do this. Plasmodesmata are junctions between plant cells, whereas animal cell contacts include tight junctions, gap junctions, and desmosomes.

### Plasmodesmata

In general, long stretches of the plasma membranes of neighboring plant cells cannot touch one another because they are separated by the cell wall that surrounds each cell. How then, can a plant transfer water and other soil nutrients from its roots, through its stems, and to its leaves? Such transport uses the vascular tissues (xylem and phloem) primarily. There also exist structural modifications called plasmodesmata (singular = plasmodesma), numerous channels that pass between cell walls of adjacent plant cells, connect their cytoplasm, and enable materials to be transported from cell to cell, and thus throughout the plant (Figure 2).





Figure 2. A plasmodesma is a channel between the cell walls of two adjacent plant cells. Plasmodesmata allow materials to pass from the cytoplasm of one plant cell to the cytoplasm of an adjacent cell.

## Tight Junctions

A **tight junction** is a watertight seal between two adjacent animal cells (Figure 3). The cells are held tightly against each other by proteins (predominantly two proteins called claudins and occludins).





Figure 3. Tight junctions form watertight connections between adjacent animal cells. Proteins create tight junction adherence. (credit: modification of work by Mariana Ruiz Villareal)

This tight adherence prevents materials from leaking between the cells; tight junctions are typically found in epithelial tissues that line internal organs and cavities, and comprise most of the skin. For example, the tight junctions of the epithelial cells lining your urinary bladder prevent urine from leaking out into the extracellular space.

## Desmosomes

Also found only in animal cells are **desmosomes**, which act like spot welds between adjacent epithelial cells (Figure 4). Short proteins called cadherins in the plasma membrane connect to intermediate filaments to create desmosomes. The cadherins join two adjacent cells together and maintain the cells in a sheet-like formation in organs and tissues that stretch, like the skin, heart, and muscles.





Figure 4. A desmosome forms a very strong spot weld between cells. It is created by the linkage of cadherins and intermediate filaments. (credit: modification of work by Mariana Ruiz Villareal)

## Gap Junctions

**Gap junctions** in animal cells are like plasmodesmata in plant cells in that they are channels between adjacent cells that allow for the transport of ions, nutrients, and other substances that enable cells to communicate (Figure 5). Structurally, however, gap junctions and plasmodesmata differ.





Figure 5. A gap junction is a protein-lined pore that allows water and small molecules to pass between adjacent animal cells. (credit: modification of work by Mariana Ruiz Villareal)

Gap junctions develop when a set of six proteins (called connexins) in the plasma membrane arrange themselves in an elongated donut-like configuration called a connexon. When the pores (“doughnut holes”) of connexons in adjacent animal cells align, a channel between the two cells forms. Gap junctions are particularly important in cardiac muscle: The electrical signal for the muscle to contract is passed efficiently through gap junctions, allowing the heart muscle cells to contract in tandem.

To conduct a virtual microscopy lab and review the parts of a cell, work through the steps of this interactive assignment.

### Learning Objectives

Animal cells communicate via their extracellular matrices and are connected to each other via tight junctions, desmosomes, and gap junctions. Plant cells are connected and communicate with each other via plasmodesmata.

When protein receptors on the surface of the plasma membrane of an animal cell bind to a substance in the extracellular matrix, a chain of reactions begins that changes activities taking place within the cell. Plasmodesmata are channels between adjacent plant cells, while gap junctions are channels between adjacent animal cells. However, their structures are quite different. A tight junction is a watertight seal between two adjacent cells, while a desmosome acts like a spot weld.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

6.17: Cell Junctions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.18: The Cell Wall

Plant cells have a cell wall, chloroplasts and other specialized plastids, and a large central vacuole, whereas animal cells do not.





Figure 1. These figures show the major organelles and other cell components of (a) a typical animal cell and (b) a typical eukaryotic plant cell. The plant cell has a cell wall, chloroplasts, plastids, and a central vacuole—structures not found in animal cells. Plant cells do not have lysosomes or centrosomes.

If you examine Figure 1b, the diagram of a plant cell, you will see a structure external to the plasma membrane called the cell wall. The **cell wall** is a rigid covering that protects the cell, provides structural support, and gives shape to the cell. Fungal and protistan cells also have cell walls, as do some prokaryotic cells. While the chief component of prokaryotic cell walls is peptidoglycan, the major organic molecule in the plant cell wall is cellulose (Figure 2), a polysaccharide made up of glucose units. Have you ever noticed that when you bite into a raw vegetable, like celery, it crunches? That's because you are tearing the rigid cell walls of the celery cells with your teeth.




Figure 2. Cellulose is a long chain of β-glucose molecules connected by a 1-4 linkage. The dashed lines at each end of the figure indicate a series of many more glucose units. The size of the page makes it impossible to portray an entire cellulose molecule.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.18: The Cell Wall is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 6.19: Putting It Together- Cellular Structure

In this module, we've learned about several different cell components. Table 1 provides a summary of all the components we covered, as well as their functions and locations.

Table 1. Components of Prokaryotic and Eukaryotic Cells

| Cell Component | Function | Present in Prokaryotes? | Present in Animal Cells? | Present in Plant Cells? |
|---|---|---|---|---|
| Cytoplasm | Provides turgor pressure to plant cells as fluid inside the central vacuole; site of many metabolic reactions; medium in which organelles are found | Yes | Yes | Yes |
| Nucleus | Cell organelle that houses DNA and directs synthesis of ribosomes and proteins | No | Yes | Yes |
| Nucleolus | Darkened area within the nucleus where ribosomal subunits are synthesized. | No | Yes | Yes |
| Ribosomes | Protein synthesis | Yes | Yes | Yes |
| Mitochondria | ATP production/cellular respiration | No | Yes | Yes |
| Peroxisomes | Oxidizes and thus breaks down fatty acids and amino acids, and detoxifies poisons | No | Yes | Yes |
| Endoplasmic reticulum | Modifies proteins and synthesizes lipids | No | Yes | Yes |
| Golgi apparatus | Modifies, sorts, tags, packages, and distributes lipids and proteins | No | Yes | Yes |
| Vesicles and vacuoles | Storage and transport; digestive function in plant cells | No | Yes | Yes |
| Centrosome | Unspecified role in cell division in animal cells; source of microtubules in animal cells | No | Yes | No |
| Lysosomes | Digestion of macromolecules; recycling of worn-out organelles | No | Yes | No |
| Chloroplasts | Photosynthesis | No | No | Yes |
| Cytoskeleton | Maintains cell’s shape, secures organelles in specific positions, allows cytoplasm and vesicles | Yes | Yes | Yes |

Table 1. Components of Prokaryotic and Eukaryotic Cells

| Cell Component | Function | Present in Prokaryotes? | Present in Animal Cells? | Present in Plant Cells? |
|---|---|---|---|---|
| | to move within cell, and enables unicellular organisms to move independently | | | |
| Flagella | Cellular locomotion | Some | Some | No, except for some plant sperm cells. |
| Cilia | Cellular locomotion, movement of particles along extracellular surface of plasma membrane, and filtration | Some | Some | No |
| Plasma membrane | Separates cell from external environment; controls passage of organic molecules, ions, water, oxygen, and wastes into and out of cell | Yes | Yes | Yes |
| Cell wall | Protection, structural support and maintenance of cell shape | Yes, primarily peptidoglycan | No | Yes, primarily cellulose |

## INHERITED DISEASES

Let's look back to our earlier disease list:

- **Pompe Disease:** characterized by excess accumulation of glycogen in muscle cells
- **Leigh Disease:** progressive disorder of lesions (dead or dying cells) in the brain
- **Emery-Dreifuss muscular dystrophy:** wasting and weakness in muscles of the shoulders, upper arms, and calf muscles

The following organelles are what cause the each disorder. As you explore the different websites that highlight each disease, think about how the function of these organelles directly relates to the disease symptoms.

- Pompe disease: lysosomes
- Leigh's Disease: mitochondria
- Emery-Dreifuss muscular dystrophy: nuclear envelope

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

6.19: Putting It Together- Cellular Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 6.20: Assignment- Cell Builder

Create a model of a eukaryotic cell using any material of your choice. In your model be sure to include all the organelles appropriate to your cell (either plant or animal). Once complete, take multiple photographs of your model from all angles. Include these images in a document that also contains the following in table format:

- A detailed key/legend that matches the model;
- Each organelle or part with its basic function;
- A disease or disorder that is associated with the malfunction of each cellular component
- How this organelle is visualized microscopically
- References

Note for the disease information, you can list a disease in either animals or plants, regardless of what type of cell you are modeling. In other words, its okay to discuss a “human” disease even if you are making a plant model, provided the organelle is present in both types of cells.

Some suggestions for 3D models include Legos, a decorated cake with candy toppings, or standard Styrofoam base with appropriate pieces attached. You can also draw or illustrate a model. Here is an example of what you might make.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references
- Assignment includes photographs or images of created model from all angles.
- Assignment includes a key documenting how each organelle is represented in the model.
- Assignment includes a completed table such as the one illustrated in the example document.
- Assignment includes a disease caused by malfunction of each identified component in the model.
- Assignment includes at least 7 organelles in the model and table.

## Rubric

| Cell Builder |
| --- |
| Outcome: Identify and explain a variety of cellular components |
| |

| Cell Builder | | | | |
|---|---|---|---|---|
| Criteria | Ratings | | | Pts |
| At least 7 membrane-bound organelles found in eukaryotic cells are contained in the model and their functions identified in the table. | Organelles and functions are outlined in detail.<br>5.0 pts | Organelles and functions are outlined in minimal but accurate terms.<br>4.0 pts | Organelles and functions are mismatched in the table or insufficiently discussed.<br>0.0 pts | 5 pts |
| The structure and function of cell membranes is included in the model and table. | Cell membranes are illustrated in detail in model and discussed in detail.<br>5.0 pts | Cell membranes are included and explained in the table/model in minimal but accurate terms.<br>4.0 pts | Cell membranes are inaccurately discussed.<br>0.0 pts | 5 pts |
| At least 2 components of the cytoskeleton are identified by form and function in the model and table. | Components of the cytoskeleton and functions are outlined in detail.<br>5.0 pts | Components of the cytoskeleton and functions are outlined in minimal but accurate terms.<br>4.0 pts | Components of the cytoskeleton are mismatched in the table or insufficiently discussed.<br>0.0 pts | 5 pts |
| At least one cell surface specialization is included in the model and table. | Cell surface specialization(s) is/are illustrated in detail in model and discussed in detail.<br>5.0 pts | Cell surface specialization(s) is/are included and explained in the table/model in minimal but accurate terms.<br>4.0 pts | Cell surface specialization(s) is/are inaccurately discussed.<br>0.0 pts | 5 pts |
| | | | | Total points: 20 |

## Contributors and Attributions

CC licensed content, Original

- Performance Assessments: Cell Builder. **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

6.20: Assignment- Cell Builder is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 7: Module 5- Cell Membranes


7.1: Why It Matters- Cell Membranes
7.2: Introduction to Structure of the Membrane
7.3: Structure of the Cell Membrane
7.4: Phospholipids
7.5: Introduction to Kinds of Transport
7.6: Passive Transport
7.7: Active Transport
7.8: Membranes and Transport
7.9: Introduction to Endocytosis and Exocytosis
7.10: Endocytosis
7.11: Exocytosis
7.12: Putting It Together- Cell Membranes
7.13: Assignment- Membranes Alive!



7: Module 5- Cell Membranes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.1: Why It Matters- Cell Membranes

## Why describe and explain the structure and function of membranes?

Cystic fibrosis (CF) is a genetic disorder that primarily affects the lungs, as well as the pancreas, liver, and intestine. CF is characterized by abnormal transport of chloride and sodium across body tissues, leading to thick, viscous secretions. The hallmark symptoms of cystic fibrosis are salty tasting skin, poor growth, poor weight gain despite a normal food intake, accumulation of thick, sticky mucus, frequent chest infections, and coughing or shortness of breath. Symptoms often appear in infancy and childhood, such as bowel obstruction in newborn babies.

The most serious symptoms of CF are difficulty breathing and frequent lung infections. Often, lung transplantation is ultimately necessary as CF worsens. Other symptoms, including sinus infections, poor growth, and infertility affect other parts of the body.

Cystic fibrosis results from the malfunctioning of a single membrane transporter. How could this error in membrane transport result in such a disease?

### Learning Outcomes

- Describe the structure and function of membranes, especially the phospholipid bilayer
- Distinguish between passive and active transport; explain how substances are directly transported across a membrane
- Describe the primary mechanisms by which cells import and export macromolecules
- Identify the main components of a signal transduction pathway

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Cystic Fibrosis. **Provided by**: Boundless. **Located at**: https://courses.lumenlearning.com/boundless-biology/. **License**: *CC BY-SA: Attribution-ShareAlike*

7.1: Why It Matters- Cell Membranes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.2: Introduction to Structure of the Membrane

## Describe the structure and function of membranes, especially the phospholipid bilayer

In this outcome, we'll learn about the structure of membranes.

## What You'll Learn to Do

- Describe the structure of cell membranes
- Identify components of the cell membrane, including phospholipids, cholesterol, proteins, and carbohydrates
- Explain why hydrophilic substances cannot pass through the interior of the cell membrane

### Learning Activities

The learning activities for this section include the following:

- Structure of the Cell Membrane
- Phospholipids
- Self Check: Structure of the Membrane

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

7.2: Introduction to Structure of the Membrane is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.3: Structure of the Cell Membrane

A cell's plasma membrane defines the boundary of the cell and determines the nature of its contact with the environment. Cells exclude some substances, take in others, and excrete still others, all in controlled quantities. Plasma membranes enclose the borders of cells, but rather than being a static bag, they are dynamic and constantly in flux. The plasma membrane must be sufficiently flexible to allow certain cells, such as red blood cells and white blood cells, to change shape as they pass through narrow capillaries. These are the more obvious functions of a plasma membrane. In addition, the surface of the plasma membrane carries markers that allow cells to recognize one another, which is vital as tissues and organs form during early development, and which later plays a role in the "self" versus "non-self" distinction of the immune response.

The plasma membrane also carries receptors, which are attachment sites for specific substances that interact with the cell. Each receptor is structured to bind with a specific substance. For example, surface receptors of the membrane create changes in the interior, such as changes in enzymes of metabolic pathways. These metabolic pathways might be vital for providing the cell with energy, making specific substances for the cell, or breaking down cellular waste or toxins for disposal. Receptors on the plasma membrane's exterior surface interact with hormones or neurotransmitters, and allow their messages to be transmitted into the cell. Some recognition sites are used by viruses as attachment points. Although they are highly specific, pathogens like viruses may evolve to exploit receptors to gain entry to a cell by mimicking the specific substance that the receptor is meant to bind. This specificity helps to explain why human immunodeficiency virus (HIV) or any of the five types of hepatitis viruses invade only specific cells.

## Fluid Mosaic Model

In 1972, S. J. Singer and Garth L. Nicolson proposed a new model of the plasma membrane that, compared to earlier understanding, better explained both microscopic observations and the function of the plasma membrane. This was called the **fluid mosaic model**. The model has evolved somewhat over time, but still best accounts for the structure and functions of the plasma membrane as we now understand them. The fluid mosaic model describes the structure of the plasma membrane as a mosaic of components—including phospholipids, cholesterol, proteins, and carbohydrates—in which the components are able to flow and change position, while maintaining the basic integrity of the membrane. Both phospholipid molecules and embedded proteins are able to diffuse rapidly and laterally in the membrane (Figure 1). The fluidity of the plasma membrane is necessary for the activities of certain enzymes and transport molecules within the membrane. Plasma membranes range from 5–10 nm thick. As a comparison, human red blood cells, visible via light microscopy, are approximately 8 µm thick, or approximately 1,000 times thicker than a plasma membrane.





Figure 1. The fluid mosaic model of the plasma membrane structure describes the plasma membrane as a fluid combination of phospholipids, cholesterol, proteins, and carbohydrates.

The plasma membrane is made up primarily of a bilayer of phospholipids with embedded proteins, carbohydrates, glycolipids, and glycoproteins, and, in animal cells, cholesterol. The amount of cholesterol in animal plasma membranes regulates the fluidity of the

membrane and changes based on the temperature of the cell's environment. In other words, cholesterol acts as antifreeze in the cell membrane and is more abundant in animals that live in cold climates.

The main fabric of the membrane is composed of two layers of phospholipid molecules, and the polar ends of these molecules (which look like a collection of balls in an artist's rendition of the model) (Figure 1) are in contact with aqueous fluid both inside and outside the cell. Thus, both surfaces of the plasma membrane are hydrophilic ("water-loving"). In contrast, the interior of the membrane, between its two surfaces, is a hydrophobic ("water-hating") or nonpolar region because of the fatty acid tails. This region has no attraction for water or other polar molecules (we will discuss this further in the next page).

Proteins make up the second major chemical component of plasma membranes. Integral proteins are embedded in the plasma membrane and may span all or part of the membrane. Integral proteins may serve as channels or pumps to move materials into or out of the cell. Peripheral proteins are found on the exterior or interior surfaces of membranes, attached either to integral proteins or to phospholipid molecules. Both integral and peripheral proteins may serve as enzymes, as structural attachments for the fibers of the cytoskeleton, or as part of the cell's recognition sites.

Carbohydrates are the third major component of plasma membranes. They are always found on the exterior surface of cells and are bound either to proteins (forming glycoproteins) or to lipids (forming glycolipids). These carbohydrate chains may consist of 2–60 monosaccharide units and may be either straight or branched. Along with peripheral proteins, carbohydrates form specialized sites on the cell surface that allow cells to recognize each other.

## How Viruses Infect Specific Organs





Figure 2. HIV docks at and binds to the CD4 receptor, a glycoprotein on the surface of T cells, before entering, or infecting, the cell. (credit: modification of work by US National Institutes of Health/National Institute of Allergy and Infectious Diseases)

Specific glycoprotein molecules exposed on the surface of the cell membranes of host cells are exploited by many viruses to infect specific organs. For example, HIV is able to penetrate the plasma membranes of specific kinds of white blood cells called T-helper cells and monocytes, as well as some cells of the central nervous system. The hepatitis virus attacks only liver cells.

These viruses are able to invade these cells, because the cells have binding sites on their surfaces that the viruses have exploited with equally specific glycoproteins in their coats. (Figure 2). The cell is tricked by the mimicry of the virus coat molecules, and the virus is able to enter the cell. Other recognition sites on the virus's surface interact with the human immune system, prompting the body to produce antibodies. Antibodies are made in response to the antigens (or proteins associated with invasive pathogens). These same sites serve as places for antibodies to attach, and either destroy or inhibit the activity of the virus. Unfortunately, these sites on HIV are encoded by genes that change quickly, making the production of an effective vaccine against the virus very difficult. The virus population within an infected individual quickly evolves through mutation into different populations, or variants, distinguished by differences in these recognition sites. This rapid change of viral surface markers decreases the effectiveness of the person's immune system in attacking the virus, because the antibodies will not recognize the new variations of the surface patterns.

## Learning Objectives

The modern understanding of the plasma membrane is referred to as the fluid mosaic model. The plasma membrane is composed of a bilayer of phospholipids, with their hydrophobic, fatty acid tails in contact with each other. The landscape of the membrane is studded with proteins, some of which span the membrane. Some of these proteins serve to transport materials into or out of the cell. Carbohydrates are attached to some of the proteins and lipids on the outward-facing surface of the

membrane. These form complexes that function to identify the cell to other cells. The fluid nature of the membrane owes itself to the configuration of the fatty acid tails, the presence of cholesterol embedded in the membrane (in animal cells), and the mosaic nature of the proteins and protein-carbohydrate complexes, which are not firmly fixed in place. Plasma membranes enclose the borders of cells, but rather than being a static bag, they are dynamic and constantly in flux.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

7.3: Structure of the Cell Membrane is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 7.4: Phospholipids

As we just learned, the main fabric of the membrane is composed of two layers of phospholipid molecules. The hydrophilic or "water-loving" areas of these molecules (which looks like a collection of balls in an artist's rendition of the model) (Figure 1) are in contact with the aqueous fluid both inside and outside the cell. Thus, both surfaces of the plasma membrane are hydrophilic. In contrast, the interior of the membrane, between its two surfaces, is a hydrophobic or nonpolar region because of the fatty acid tails. This region has no attraction for water or other polar molecules (we will discuss this further in the next page).





Figure 1. The fluid mosaic model of the plasma membrane structure describes the plasma membrane as a fluid combination of phospholipids, cholesterol, proteins, and carbohydrates.

Hydrophobic, or water-hating molecules, tend to be non-polar. They interact with other non-polar molecules in chemical reactions, but generally do not interact with polar molecules. When placed in water, hydrophobic molecules tend to form a ball or cluster. The hydrophilic regions of the phospholipids tend to form hydrogen bonds with water and other polar molecules on both the exterior and interior of the cell. Thus, the membrane surfaces that face the interior and exterior of the cell are hydrophilic. In contrast, the interior of the cell membrane is hydrophobic and will not interact with water. Therefore, phospholipids form an excellent two-layer cell membrane that separates fluid within the cell from the fluid outside of the cell.





Figure 2. This phospholipid molecule is composed of a hydrophilic head and two hydrophobic tails. The hydrophilic head group consists of a phosphate-containing group attached to a glycerol molecule. The hydrophobic tails, each containing either a saturated or an unsaturated fatty acid, are long hydrocarbon chains.

A phospholipid molecule (Figure 2) consists of a three-carbon glycerol backbone with two fatty acid molecules attached to carbons 1 and 2, and a phosphate-containing group attached to the third carbon.

This arrangement gives the overall molecule an area described as its head (the phosphate-containing group), which has a polar character or negative charge, and an area called the tail (the fatty acids), which has no charge. The head can form hydrogen bonds, but the tail cannot. A molecule with this arrangement of a positively or negatively charged area and an uncharged, or non-polar, area is referred to as amphiphilic or “dual-loving.”

This characteristic is vital to the structure of a plasma membrane because, in water, phospholipids tend to become arranged with their hydrophobic tails facing each other and their hydrophilic heads facing out. In this way, they form a lipid bilayer—a barrier composed of a double layer of phospholipids that separates the water and other materials on one side of the barrier from the water and other materials on the other side. In fact, phospholipids heated in an aqueous solution tend to spontaneously form small spheres or droplets (called micelles or liposomes), with their hydrophilic heads forming the exterior and their hydrophobic tails on the inside (Figure 3).





Figure 3. In an aqueous solution, phospholipids tend to arrange themselves with their polar heads facing outward and their hydrophobic tails facing inward. (credit: modification of work by Mariana Ruiz Villareal)

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

7.4: Phospholipids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.5: Introduction to Kinds of Transport

## Explain how substances are directly transported across a membrane

In this outcome we'll learn about passive and active transport; this includes diffusion and osmosis.

## What You'll Learn to Do

- Define and describe passive transport
- Define and describe active transport

### Learning Activities

The learning activities for this section include the following:

- Passive Transport
- Active Transport
- Membranes and Transport
- Self Check: Kinds of Transport

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

7.5: Introduction to Kinds of Transport is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.6: Passive Transport

Plasma membranes must allow certain substances to enter and leave a cell, and prevent some harmful materials from entering and some essential materials from leaving. In other words, plasma membranes are **selectively permeable**—they allow some substances to pass through, but not others. If they were to lose this selectivity, the cell would no longer be able to sustain itself, and it would be destroyed. Some cells require larger amounts of specific substances than do other cells; they must have a way of obtaining these materials from extracellular fluids. This may happen passively, as certain materials move back and forth, or the cell may have special mechanisms that facilitate transport. Some materials are so important to a cell that it spends some of its energy, hydrolyzing adenosine triphosphate (ATP), to obtain these materials. Red blood cells use some of their energy doing just that. All cells spend the majority of their energy to maintain an imbalance of sodium and potassium ions between the interior and exterior of the cell.

The most direct forms of membrane transport are passive. **Passive transport** is a naturally occurring phenomenon and does not require the cell to exert any of its energy to accomplish the movement. In passive transport, substances move from an area of higher concentration to an area of lower concentration. A physical space in which there is a range of concentrations of a single substance is said to have a **concentration gradient**.

## Selective Permeability

Plasma membranes are asymmetric: the interior of the membrane is not identical to the exterior of the membrane. In fact, there is a considerable difference between the array of phospholipids and proteins between the two leaflets that form a membrane. On the interior of the membrane, some proteins serve to anchor the membrane to fibers of the cytoskeleton. There are peripheral proteins on the exterior of the membrane that bind elements of the extracellular matrix. Carbohydrates, attached to lipids or proteins, are also found on the exterior surface of the plasma membrane. These carbohydrate complexes help the cell bind substances that the cell needs in the extracellular fluid. This adds considerably to the selective nature of plasma membranes (Figure 1).





Figure 1. The eukaryotic plasma membrane is a phospholipid bilayer with proteins and cholesterol embedded in it.

Recall that plasma membranes are amphiphilic: they have hydrophilic and hydrophobic regions. This characteristic helps the movement of some materials through the membrane and hinders the movement of others. Lipid-soluble material with a low molecular weight can easily slip through the hydrophobic lipid core of the membrane. Substances such as the fat-soluble vitamins A, D, E, and K readily pass through the plasma membranes in the digestive tract and other tissues. Fat-soluble drugs and hormones also gain easy entry into cells and are readily transported into the body's tissues and organs. Molecules of oxygen and carbon dioxide have no charge and so pass through membranes by simple diffusion.

Polar substances present problems for the membrane. While some polar molecules connect easily with the outside of a cell, they cannot readily pass through the lipid core of the plasma membrane. Additionally, while small ions could easily slip through the spaces in the mosaic of the membrane, their charge prevents them from doing so. Ions such as sodium, potassium, calcium, and

chloride must have special means of penetrating plasma membranes. Simple sugars and amino acids also need help with transport across plasma membranes, achieved by various transmembrane proteins (channels).

## Diffusion



Figure 2. Dispersion

**Diffusion** is a passive process of transport (see Figure 2). A single substance tends to move from an area of high concentration to an area of low concentration until the concentration is equal across a space. You are familiar with diffusion of substances through the air.

For example, think about someone opening a bottle of ammonia in a room filled with people. The ammonia gas is at its highest concentration in the bottle; its lowest concentration is at the edges of the room. The ammonia vapor will diffuse, or spread away, from the bottle, and gradually, more and more people will smell the ammonia as it spreads. Materials move within the cell's cytosol by diffusion, and certain materials move through the plasma membrane by diffusion (Figure 3). Diffusion expends no energy. On the contrary, concentration gradients are a form of potential energy, dissipated as the gradient is eliminated.





Figure 3. Diffusion through a permeable membrane moves a substance from an area of high concentration (extracellular fluid, in this case) down its concentration gradient (into the cytoplasm). (credit: modification of work by Mariana Ruiz Villareal)

Each separate substance in a medium, such as the extracellular fluid, has its own concentration gradient, independent of the concentration gradients of other materials. In addition, each substance will diffuse according to that gradient. Within a system, there will be different rates of diffusion of the different substances in the medium.

### Factors That Affect Diffusion

Molecules move constantly in a random manner, at a rate that depends on their mass, their environment, and the amount of thermal energy they possess, which in turn is a function of temperature. This movement accounts for the diffusion of molecules through whatever medium in which they are localized. A substance will tend to move into any space available to it until it is evenly distributed throughout it. After a substance has diffused completely through a space, removing its concentration gradient, molecules will still move around in the space, but there will be no *net* movement of the number of molecules from one area to another. This lack of a concentration gradient in which there is no net movement of a substance is known as dynamic equilibrium. While diffusion will go forward in the presence of a concentration gradient of a substance, several factors affect the rate of diffusion.

- Extent of the concentration gradient: The greater the difference in concentration, the more rapid the diffusion. The closer the distribution of the material gets to equilibrium, the slower the rate of diffusion becomes.
- Mass of the molecules diffusing: Heavier molecules move more slowly; therefore, they diffuse more slowly. The reverse is true for lighter molecules.
- Temperature: Higher temperatures increase the energy and therefore the movement of the molecules, increasing the rate of diffusion. Lower temperatures decrease the energy of the molecules, thus decreasing the rate of diffusion.
- Solvent density: As the density of a solvent increases, the rate of diffusion decreases. The molecules slow down because they have a more difficult time getting through the denser medium. If the medium is less dense, diffusion increases. Because cells primarily use diffusion to move materials within the cytoplasm, any increase in the cytoplasm's density will inhibit the movement of the materials. An example of this is a person experiencing dehydration. As the body's cells lose water, the rate of diffusion decreases in the cytoplasm, and the cells' functions deteriorate. Neurons tend to be very sensitive to this effect. Dehydration frequently leads to unconsciousness and possibly coma because of the decrease in diffusion rate within the cells.
- Solubility: As discussed earlier, nonpolar or lipid-soluble materials pass through plasma membranes more easily than polar materials, allowing a faster rate of diffusion.
- Surface area and thickness of the plasma membrane: Increased surface area increases the rate of diffusion, whereas a thicker membrane reduces it.
- Distance travelled: The greater the distance that a substance must travel, the slower the rate of diffusion. This places an upper limitation on cell size. A large, spherical cell will die because nutrients or waste cannot reach or leave the center of the cell, respectively. Therefore, cells must either be small in size, as in the case of many prokaryotes, or be flattened, as with many single-celled eukaryotes.

A variation of diffusion is the process of filtration. In filtration, material moves according to its concentration gradient through a membrane; sometimes the rate of diffusion is enhanced by pressure, causing the substances to filter more rapidly. This occurs in the kidney, where blood pressure forces large amounts of water and accompanying dissolved substances, or solutes, out of the blood and into the renal tubules. The rate of diffusion in this instance is almost totally dependent on pressure. One of the effects of high blood pressure is the appearance of protein in the urine, which is "squeezed through" by the abnormally high pressure.

## Facilitated Transport

In facilitated transport, also called facilitated diffusion, materials diffuse across the plasma membrane with the help of membrane proteins. A concentration gradient exists that would allow these materials to diffuse into the cell without expending cellular energy. However, these materials are ions or polar molecules that are repelled by the hydrophobic parts of the cell membrane. Facilitated transport proteins shield these materials from the repulsive force of the membrane, allowing them to diffuse into the cell.

The material being transported is first attached to protein or glycoprotein receptors on the exterior surface of the plasma membrane. This allows the material that is needed by the cell to be removed from the extracellular fluid. The substances are then passed to specific integral proteins that facilitate their passage. Some of these integral proteins are collections of beta pleated sheets that form a pore or channel through the phospholipid bilayer. Others are carrier proteins which bind with the substance and aid its diffusion through the membrane.

## Channels





Figure 4. Facilitated transport moves substances down their concentration gradients. They may cross the plasma membrane with the aid of channel proteins. (credit: modification of work by Mariana Ruiz Villareal)

The integral proteins involved in facilitated transport are collectively referred to as **transport proteins**, and they function as either channels for the material or carriers. In both cases, they are transmembrane proteins. Channels are specific for the substance that is being transported. **Channel proteins** have hydrophilic domains exposed to the intracellular and extracellular fluids; they additionally have a hydrophilic channel through their core that provides a hydrated opening through the membrane layers (Figure 4). Passage through the channel allows polar compounds to avoid the nonpolar central layer of the plasma membrane that would otherwise slow or prevent their entry into the cell. **Aquaporins** are channel proteins that allow water to pass through the membrane at a very high rate.

Channel proteins are either open at all times or they are “gated,” which controls the opening of the channel. The attachment of a particular ion to the channel protein may control the opening, or other mechanisms or substances may be involved. In some tissues, sodium and chloride ions pass freely through open channels, whereas in other tissues a gate must be opened to allow passage. An example of this occurs in the kidney, where both forms of channels are found in different parts of the renal tubules. Cells involved in the transmission of electrical impulses, such as nerve and muscle cells, have gated channels for sodium, potassium, and calcium in their membranes. Opening and closing of these channels changes the relative concentrations on opposing sides of the membrane of these ions, resulting in the facilitation of electrical transmission along membranes (in the case of nerve cells) or in muscle contraction (in the case of muscle cells).

## Carrier Proteins

Another type of protein embedded in the plasma membrane is a **carrier protein**. This aptly named protein binds a substance and, in doing so, triggers a change of its own shape, moving the bound molecule from the outside of the cell to its interior (Figure 5); depending on the gradient, the material may move in the opposite direction. Carrier proteins are typically specific for a single substance. This selectivity adds to the overall selectivity of the plasma membrane. The exact mechanism for the change of shape is poorly understood. Proteins can change shape when their hydrogen bonds are affected, but this may not fully explain this mechanism. Each carrier protein is specific to one substance, and there are a finite number of these proteins in any membrane. This can cause problems in transporting enough of the material for the cell to function properly. When all of the proteins are bound to their ligands, they are saturated and the rate of transport is at its maximum. Increasing the concentration gradient at this point will not result in an increased rate of transport.





Figure 5. Some substances are able to move down their concentration gradient across the plasma membrane with the aid of carrier proteins. Carrier proteins change shape as they move molecules across the membrane. (credit: modification of work by Mariana Ruiz Villareal)

An example of this process occurs in the kidney. Glucose, water, salts, ions, and amino acids needed by the body are filtered in one part of the kidney. This filtrate, which includes glucose, is then reabsorbed in another part of the kidney. Because there are only a finite number of carrier proteins for glucose, if more glucose is present than the proteins can handle, the excess is not transported and it is excreted from the body in the urine. In a diabetic individual, this is described as “spilling glucose into the urine.” A different group of carrier proteins called glucose transport proteins, or GLUTs, are involved in transporting glucose and other hexose sugars through plasma membranes within the body.

Channel and carrier proteins transport material at different rates. Channel proteins transport much more quickly than do carrier proteins. Channel proteins facilitate diffusion at a rate of tens of millions of molecules per second, whereas carrier proteins work at a rate of a thousand to a million molecules per second.

## Osmosis

Osmosis is the movement of water through a semipermeable membrane according to the concentration gradient of water across the membrane, which is inversely proportional to the concentration of solutes. While diffusion transports material across membranes and within cells, osmosis transports *only water* across a membrane and the membrane limits the diffusion of solutes in the water. Not surprisingly, the aquaporins that facilitate water movement play a large role in osmosis, most prominently in red blood cells and the membranes of kidney tubules.

### Mechanism





Figure 6. In osmosis, water always moves from an area of higher water concentration to one of lower concentration. In the diagram shown, the solute cannot pass through the selectively permeable membrane, but the water can.

Osmosis is a special case of diffusion. Water, like other substances, moves from an area of high concentration to one of low concentration. An obvious question is what makes water move at all? Imagine a beaker with a semipermeable membrane separating the two sides or halves (Figure 6). On both sides of the membrane the water level is the same, but there are different concentrations of a dissolved substance, or **solute**, that cannot cross the membrane (otherwise the concentrations on each side would be balanced by the solute crossing the membrane). If the volume of the solution on both sides of the membrane is the same, but the concentrations of solute are different, then there are different amounts of water, the solvent, on either side of the membrane.

To illustrate this, imagine two full glasses of water. One has a single teaspoon of sugar in it, whereas the second one contains one-quarter cup of sugar. If the total volume of the solutions in both cups is the same, which cup contains more water? Because the large amount of sugar in the second cup takes up much more space than the teaspoon of sugar in the first cup, the first cup has more water in it.

Returning to the beaker example, recall that it has a mixture of solutes on either side of the membrane. A principle of diffusion is that the molecules move around and will spread evenly throughout the medium if they can. However, only the material capable of getting through the membrane will diffuse through it. In this example, the solute cannot diffuse through the membrane, but the water can. Water has a concentration gradient in this system. Thus, water will diffuse down its concentration gradient, crossing the membrane to the side where it is less concentrated. This diffusion of water through the membrane—osmosis—will continue until the concentration gradient of water goes to zero or until the hydrostatic pressure of the water balances the osmotic pressure. Osmosis proceeds constantly in living systems.

## Tonicity

**Tonicity** describes how an extracellular solution can change the volume of a cell by affecting osmosis. A solution's tonicity often directly correlates with the osmolarity of the solution. **Osmolarity** describes the total solute concentration of the solution. A solution with low osmolarity has a greater number of water molecules relative to the number of solute particles; a solution with high osmolarity has fewer water molecules with respect to solute particles. In a situation in which solutions of two different osmolarities are separated by a membrane permeable to water, though not to the solute, water will move from the side of the membrane with lower osmolarity (and more water) to the side with higher osmolarity (and less water). This effect makes sense if you remember that the solute cannot move across the membrane, and thus the only component in the system that can move—the water—moves along its own concentration gradient. An important distinction that concerns living systems is that osmolarity measures the number of particles (which may be molecules) in a solution. Therefore, a solution that is cloudy with cells may have a lower osmolarity than a solution that is clear, if the second solution contains more dissolved molecules than there are cells.

### Hypotonic Solutions

Three terms—hypotonic, isotonic, and hypertonic—are used to relate the osmolarity of a cell to the osmolarity of the extracellular fluid that contains the cells. In a hypotonic situation, the extracellular fluid has lower osmolarity than the fluid inside the cell, and water enters the cell. (In living systems, the point of reference is always the cytoplasm, so the prefix *hypo–* means that the extracellular fluid has a lower concentration of solutes, or a lower osmolarity, than the cell cytoplasm.) It also means that the extracellular fluid has a higher concentration of water in the solution than does the cell. In this situation, water will follow its concentration gradient and enter the cell.

### Hypertonic Solutions

As for a hypertonic solution, the prefix *hyper–* refers to the extracellular fluid having a higher osmolarity than the cell's cytoplasm; therefore, the fluid contains less water than the cell does. Because the cell has a relatively higher concentration of water, water will leave the cell.

### Isotonic Solutions

In an isotonic solution, the extracellular fluid has the same osmolarity as the cell. If the osmolarity of the cell matches that of the extracellular fluid, there will be no net movement of water into or out of the cell, although water will still move in and out. Blood cells and plant cells (Figure 7) in hypertonic, isotonic, and hypotonic solutions take on characteristic appearances.

Practice Question





Figure 7. Osmotic pressure changes the shape of red blood cells in hypertonic, isotonic, and hypotonic solutions. The turgor pressure within a plant cell depends on the tonicity of the solution that it is bathed in. (credit: modification of work by Mariana Ruiz Villareal)

A doctor injects a patient with what the doctor thinks is an isotonic saline solution. The patient dies, and an autopsy reveals that many red blood cells have been destroyed. Do you think the solution the doctor injected was really isotonic?

[practice-area rows="2"][/practice-area]
[reveal-answer q="619831"]**Show Answer**[/reveal-answer]
[hidden-answer a="619831"]No, it must have been hypotonic as a hypotonic solution would cause water to enter the cells, thereby making them burst.[/hidden-answer]

## Tonicity in Living Systems

In a hypotonic environment, water enters a cell, and the cell swells. In an isotonic condition, the relative concentrations of solute and solvent are equal on both sides of the membrane. There is no net water movement; therefore, there is no change in the size of the cell. In a hypertonic solution, water leaves a cell and the cell shrinks. If either the hypo- or hyper- condition goes to excess, the cell's functions become compromised, and the cell may be destroyed.

A red blood cell will burst, or lyse, when it swells beyond the plasma membrane's capability to expand. Remember, the membrane resembles a mosaic, with discrete spaces between the molecules composing it. If the cell swells, and the spaces between the lipids and proteins become too large, the cell will break apart.

In contrast, when excessive amounts of water leave a red blood cell, the cell shrinks, or crenates. This has the effect of concentrating the solutes left in the cell, making the cytosol denser and interfering with diffusion within the cell. The cell's ability to function will be compromised and may also result in the death of the cell.

Various living things have ways of controlling the effects of osmosis—a mechanism called osmoregulation. Some organisms, such as plants, fungi, bacteria, and some protists, have cell walls that surround the plasma membrane and prevent cell lysis in a hypotonic solution. The plasma membrane can only expand to the limit of the cell wall, so the cell will not lyse. In fact, the cytoplasm in plants is always slightly hypertonic to the cellular environment, and water will always enter a cell if water is available. This inflow of water produces turgor pressure, which stiffens the cell walls of the plant (Figure 7). In nonwoody plants, turgor pressure supports the plant. Conversly, if the plant is not watered, the extracellular fluid will become hypertonic, causing water to leave the cell. In this condition, the cell does not shrink because the cell wall is not flexible. However, the cell membrane detaches from the wall and constricts the cytoplasm. This is called plasmolysis. Plants lose turgor pressure in this condition and wilt (Figure 8).



Figure 8. Without adequate water, the plant on the left has lost turgor pressure, visible in its wilting; the turgor pressure is restored by watering it (right). (credit: Victor M. Vicente Selvas)





Figure 9. A paramecium's contractile vacuole, here visualized using bright field light microscopy at 480× magnification, continuously pumps water out of the organism's body to keep it from bursting in a hypotonic medium. (credit: modification of work by NIH; scale-bar data from Matt Russell)


Tonicity is a concern for all living things. For example, paramecia and amoebas, which are protists that lack cell walls, have contractile vacuoles. This vesicle collects excess water from the cell and pumps it out, keeping the cell from lysing as it takes on water from its environment (Figure 9).

Many marine invertebrates have internal salt levels matched to their environments, making them isotonic with the water in which they live. Fish, however, must spend approximately five percent of their metabolic energy maintaining osmotic homeostasis. Freshwater fish live in an environment that is hypotonic to their cells. These fish actively take in salt through their gills and excrete diluted urine to rid themselves of excess water. Saltwater fish live in the reverse environment, which is hypertonic to their cells, and they secrete salt through their gills and excrete highly concentrated urine.

In vertebrates, the kidneys regulate the amount of water in the body. Osmoreceptors are specialized cells in the brain that monitor the concentration of solutes in the blood. If the levels of solutes increase beyond a certain range, a hormone is released that retards water loss through the kidney and dilutes the blood to safer levels. Animals also have high concentrations of albumin, which is produced by the liver, in their blood. This protein is too large to pass easily through plasma membranes and is a major factor in controlling the osmotic pressures applied to tissues.

## Learning Objectives

Watch this review of osmosis and diffusion




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=220

### In Summary: Passive Transport

No energy is required. The “driving force” is a difference in the concentration of a substance on one side of the membrane compared that on the other side.

- Simple diffusion ($O_2$, $CO_2$, $H_20$. Water and nonpolar molecules).
  - Osmosis is a special kind of simple diffusion for water only.
- Familiarize yourself with the terms *hypotonic*, *isotonic*, and *hypertonic*, and be able to indicate what either a plant or an animal cell will look like if placed in a particular kind of solution.
  - Example: a plant cell has a cell wall and is full and happy when placed in water (a hypotonic solution). An animal cell does not have a cell wall and will swell and burst if placed in water. This is why a patient should never receive an IV injection of water: it will cause their red blood cells to burst.
- Facilitated diffusion (sugars, ions, amino acids, etc. Charged or polar molecules).
  - Carrier proteins
  - Channel proteins (such as ion channels or aquaporin)

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Diffusion and Osmosis. **Authored by**: Sal Khan. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/test-prep/mcat/cells/transport-across-a-cell-membrane/v/diffusion-and-osmosis. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

Public domain content

- Dispersion. **Authored by**: CTho. **Located at**: https://commons.wikimedia.org/wiki/File:Dispersion.gif. **License**: *Public Domain: No Known Copyright*

---

7.6: Passive Transport is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.7: Active Transport

**Active transport** mechanisms require the use of the cell's energy, usually in the form of adenosine triphosphate (ATP). If a substance must move into the cell against its concentration gradient—that is, if the concentration of the substance inside the cell is greater than its concentration in the extracellular fluid (and vice versa)—the cell must use energy to move the substance. Some active transport mechanisms move small-molecular weight materials, such as ions, through the membrane. Other mechanisms transport much larger molecules.

## Electrochemical Gradient

We have discussed simple concentration gradients—differential concentrations of a substance across a space or a membrane—but in living systems, gradients are more complex. Because ions move into and out of cells and because cells contain proteins that do not move across the membrane and are mostly negatively charged, there is also an electrical gradient, a difference of charge, across the plasma membrane. The interior of living cells is electrically negative with respect to the extracellular fluid in which they are bathed, and at the same time, cells have higher concentrations of potassium ($K^+$) and lower concentrations of sodium ($Na^+$) than does the extracellular fluid. So in a living cell, the concentration gradient of $Na^+$ tends to drive it into the cell, and the electrical gradient of $Na^+$ (a positive ion) also tends to drive it inward to the negatively charged interior. The situation is more complex, however, for other elements such as potassium. The electrical gradient of $K^+$, a positive ion, also tends to drive it into the cell, but the concentration gradient of $K^+$ tends to drive $K^+$ *out* of the cell (Figure 1). The combined gradient of concentration and electrical charge that affects an ion is called its **electrochemical gradient**.

Practice Question





Figure 1. Electrochemical gradients arise from the combined effects of concentration gradients and electrical gradients. (credit: "Synaptitude"/Wikimedia Commons)

Injection of a potassium solution into a person's blood is lethal; this is used in capital punishment and euthanasia. Why do you think a potassium solution injection is lethal?

[practice-area rows="2"][/practice-area]
[reveal-answer q="900539"]**Show Answer**[/reveal-answer]
[hidden-answer a="900539"]Cells typically have a high concentration of potassium in the cytoplasm and are bathed in a high concentration of sodium. Injection of potassium dissipates this electrochemical gradient. In heart muscle, the sodium/potassium potential is responsible for transmitting the signal that causes the muscle to contract. When this potential is dissipated, the signal

can’t be transmitted, and the heart stops beating. Potassium injections are also used to stop the heart from beating during surgery. [/hidden-answer]

### Moving Against a Gradient

To move substances against a concentration or electrochemical gradient, the cell must use energy. This energy is harvested from ATP generated through the cell’s metabolism. Active transport mechanisms, collectively called **pumps**, work against electrochemical gradients. Small substances constantly pass through plasma membranes. Active transport maintains concentrations of ions and other substances needed by living cells in the face of these passive movements. Much of a cell’s supply of metabolic energy may be spent maintaining these processes. (Most of a red blood cell’s metabolic energy is used to maintain the imbalance between exterior and interior sodium and potassium levels required by the cell.) Because active transport mechanisms depend on a cell’s metabolism for energy, they are sensitive to many metabolic poisons that interfere with the supply of ATP.

Two mechanisms exist for the transport of small-molecular weight material and small molecules. **Primary active transport** moves ions across a membrane and creates a difference in charge across that membrane, which is directly dependent on ATP. **Secondary active transport** describes the movement of material that is due to the electrochemical gradient established by primary active transport that does not directly require ATP.

### Carrier Proteins for Active Transport

An important membrane adaption for active transport is the presence of specific carrier proteins or pumps to facilitate movement: there are three types of these proteins or **transporters** (Figure 2). A **uniporter** carries one specific ion or molecule. A **symporter** carries two different ions or molecules, both in the same direction. An **antiporter** also carries two different ions or molecules, but in different directions. All of these transporters can also transport small, uncharged organic molecules like glucose. These three types of carrier proteins are also found in facilitated diffusion, but they do not require ATP to work in that process. Some examples of pumps for active transport are $Na^{+}$–$K^{+}$ ATPase, which carries sodium and potassium ions, and $H^{+}$–$K^{+}$ ATPase, which carries hydrogen and potassium ions. Both of these are antiporter carrier proteins. Two other carrier proteins are $Ca2^{+}$ ATPase and $H^{+}$ ATPase, which carry only calcium and only hydrogen ions, respectively. Both are pumps.





Figure 2. A uniporter carries one molecule or ion. A symporter carries two different molecules or ions, both in the same direction. An antiporter also carries two different molecules or ions, but in different directions. (credit: modification of work by “Lupask”/Wikimedia Commons)

## Primary Active Transport

The primary active transport that functions with the active transport of sodium and potassium allows secondary active transport to occur. The second transport method is still considered active because it depends on the use of energy as does primary transport (Figure 3).





Figure 3. Primary active transport moves ions across a membrane, creating an electrochemical gradient (electrogenic transport). (credit: modification of work by Mariana Ruiz Villareal)

One of the most important pumps in animals cells is the sodium-potassium pump ($Na^+$-$K^+$ ATPase), which maintains the electrochemical gradient (and the correct concentrations of $Na^+$ and $K^+$) in living cells. The sodium-potassium pump moves $K^+$ into the cell while moving Na+ out at the same time, at a ratio of three $Na^+$ for every two $K^+$ ions moved in. The $Na^+$-$K^+$ ATPase exists in two forms, depending on its orientation to the interior or exterior of the cell and its affinity for either sodium or potassium ions. The process consists of the following six steps.

1. With the enzyme oriented towards the interior of the cell, the carrier has a high affinity for sodium ions. Three ions bind to the protein.
2. ATP is hydrolyzed by the protein carrier and a low-energy phosphate group attaches to it.
3. As a result, the carrier changes shape and re-orients itself towards the exterior of the membrane. The protein's affinity for sodium decreases and the three sodium ions leave the carrier.
4. The shape change increases the carrier's affinity for potassium ions, and two such ions attach to the protein. Subsequently, the low-energy phosphate group detaches from the carrier.
5. With the phosphate group removed and potassium ions attached, the carrier protein repositions itself towards the interior of the cell.
6. The carrier protein, in its new configuration, has a decreased affinity for potassium, and the two ions are released into the cytoplasm. The protein now has a higher affinity for sodium ions, and the process starts again.

Several things have happened as a result of this process. At this point, there are more sodium ions outside of the cell than inside and more potassium ions inside than out. For every three ions of sodium that move out, two ions of potassium move in. This results in the interior being slightly more negative relative to the exterior. This difference in charge is important in creating the conditions necessary for the secondary process. The sodium-potassium pump is, therefore, an electrogenic pump (a pump that creates a charge imbalance), creating an electrical imbalance across the membrane and contributing to the membrane potential.

## Secondary Active Transport (Co-transport)

Secondary active transport brings sodium ions, and possibly other compounds, into the cell. As sodium ion concentrations build outside of the plasma membrane because of the action of the primary active transport process, an electrochemical gradient is created. If a channel protein exists and is open, the sodium ions will be pulled through the membrane. This movement is used to transport other substances that can attach themselves to the transport protein through the membrane (Figure 4). Many amino acids, as well as glucose, enter a cell this way. This secondary process is also used to store high-energy hydrogen ions in the mitochondria of plant and animal cells for the production of ATP. The potential energy that accumulates in the stored hydrogen ions is translated into kinetic energy as the ions surge through the channel protein ATP synthase, and that energy is used to convert ADP into ATP.

Practice Question

An electrochemical gradient, created by primary active transport, can move other substances against their concentration gradients, a process called co-transport or secondary active transport.





Figure 4. (credit: modification of work by Mariana Ruiz Villareal)

[practice-area rows="2"][/practice-area]
[reveal-answer q="483973"]**Show Answer**[/reveal-answer]
[hidden-answer a="483973"]A decrease in pH means an increase in positively charged $H^+$ ions, and an increase in the electrical gradient across the membrane. The transport of amino acids into the cell will increase.[/hidden-answer]

### In Summary: Active Transport

Energy is required.

- Primary active transport (ATP is the "driving force").
- Secondary active transport (the energy is provided by an electrochemical gradient).

## Contributors and Attributions


CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

7.7: Active Transport is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.8: Membranes and Transport

Now that we've learned about active and passive transport separately, let's review both topics together. In addition to reviewing the ways transport works across membranes, this video will discuss the reasons cells must be selectively permeable.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=224

## Contributors and Attributions

All rights reserved content

- In Da Club - Membranes & Transport: Crash Course Biology #5. **Authored by**: CrashCourse. **Located at**: https://youtu.be/dPKvHrD1eS4. **Project**: Crash Course Biology. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

7.8: Membranes and Transport is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.9: Introduction to Endocytosis and Exocytosis

## Describe the primary mechanisms by which cells import and export macromolecules

In addition to moving small ions and molecules through the membrane, cells also need to remove and take in larger molecules and particles. Some cells are even capable of engulfing entire unicellular microorganisms. You might have correctly hypothesized that the uptake and release of large particles by the cell requires energy. A large particle, however, cannot pass through the membrane, even with energy supplied by the cell.

There are two primary mechanisms that transport these large particles: endocytosis and exocytosis.

## What You'll Learn to Do

- Describe endocytosis and identify different varieties of import, including phagocytosis, pinocytosis, and receptor-mediated endocytosis
- Identify the steps of exocytosis

### Learning Activities

The learning activities for this section include the following:

- Endocytosis
- Exocytosis
- Self Check: Endocytosis and Exocytosis

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

7.9: Introduction to Endocytosis and Exocytosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.10: Endocytosis

**Endocytosis** is a type of active transport that moves particles, such as large molecules, parts of cells, and even whole cells, into a cell. There are different variations of endocytosis, but all share a common characteristic: the plasma membrane of the cell invaginates, forming a pocket around the target particle. The pocket pinches off, resulting in the particle being contained in a newly created intracellular vesicle formed from the plasma membrane.

## Phagocytosis





Figure 1. In phagocytosis, the cell membrane surrounds the particle and engulfs it. (credit: Mariana Ruiz Villareal)

Phagocytosis (the condition of “cell eating”) is the process by which large particles, such as cells or relatively large particles, are taken in by a cell. For example, when microorganisms invade the human body, a type of white blood cell called a neutrophil will remove the invaders through this process, surrounding and engulfing the microorganism, which is then destroyed by the neutrophil (Figure 1).

In preparation for phagocytosis, a portion of the inward-facing surface of the plasma membrane becomes coated with a protein called *clathrin*, which stabilizes this section of the membrane. The coated portion of the membrane then extends from the body of the cell and surrounds the particle, eventually enclosing it. Once the vesicle containing the particle is enclosed within the cell, the clathrin disengages from the membrane and the vesicle merges with a lysosome for the breakdown of the material in the newly formed compartment (endosome). When accessible nutrients from the degradation of the vesicular contents have been extracted, the newly formed endosome merges with the plasma membrane and releases its contents into the extracellular fluid. The endosomal membrane again becomes part of the plasma membrane.

## Pinocytosis





Figure 2. In pinocytosis, the cell membrane invaginates, surrounds a small volume of fluid, and pinches off. (credit: Mariana Ruiz Villareal)

A variation of endocytosis is called pinocytosis. This literally means “cell drinking” and was named at a time when the assumption was that the cell was purposefully taking in extracellular fluid. In reality, this is a process that takes in molecules, including water, which the cell needs from the extracellular fluid. Pinocytosis results in a much smaller vesicle than does phagocytosis, and the vesicle does not need to merge with a lysosome (Figure 2).

A variation of pinocytosis is called potocytosis. This process uses a coating protein, called caveolin, on the cytoplasmic side of the plasma membrane, which performs a similar function to clathrin. The cavities in the plasma membrane that form the vacuoles have membrane receptors and lipid rafts in addition to caveolin.

The vacuoles or vesicles formed in caveolae (singular caveola) are smaller than those in pinocytosis. Potocytosis is used to bring small molecules into the cell and to transport these molecules through the cell for their release on the other side of the cell, a process called transcytosis.

## Receptor-Mediated Endocytosis





Figure 3. In receptor-mediated endocytosis, uptake of substances by the cell is targeted to a single type of substance that binds to the receptor on the external surface of the cell membrane. (credit: modification of work by Mariana Ruiz Villareal)

A targeted variation of endocytosis employs receptor proteins in the plasma membrane that have a specific binding affinity for certain substances (Figure 3).

In receptor-mediated endocytosis, as in phagocytosis, clathrin is attached to the cytoplasmic side of the plasma membrane. If uptake of a compound is dependent on receptor-mediated endocytosis and the process is ineffective, the material will not be removed from the tissue fluids or blood. Instead, it will stay in those fluids and increase in concentration.

Some human diseases are caused by the failure of receptor-mediated endocytosis. For example, the form of cholesterol termed low-density lipoprotein or LDL (also referred to as “bad” cholesterol) is removed from the blood by receptor-mediated endocytosis. In the human genetic disease familial hypercholesterolemia, the LDL receptors are defective or missing entirely. People with this condition have life-threatening levels of cholesterol in their blood, because their cells cannot clear LDL particles from their blood.

Although receptor-mediated endocytosis is designed to bring specific substances that are normally found in the extracellular fluid into the cell, other substances may gain entry into the cell at the same site. Flu viruses, diphtheria, and cholera toxin all have sites that cross-react with normal receptor-binding sites and gain entry into cells.

### Learning Objectives

Cells perform three main types of endocytosis. Phagocytosis is the process by which cells ingest large particles, including other cells, by enclosing the particles in an extension of the cell membrane and budding off a new vesicle. During pinocytosis, cells take in molecules such as water from the extracellular fluid. Finally, receptor-mediated endocytosis is a targeted version of endocytosis where receptor proteins in the plasma membrane ensure only specific, targeted substances are brought into the cell.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

7.10: Endocytosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.11: Exocytosis

The reverse process of moving material into a cell is the process of exocytosis. Exocytosis is the opposite of the processes discussed in the last section in that its purpose is to expel material from the cell into the extracellular fluid. Waste material is enveloped in a membrane and fuses with the interior of the plasma membrane. This fusion opens the membranous envelope on the exterior of the cell, and the waste material is expelled into the extracellular space (Figure 1). Other examples of cells releasing molecules via exocytosis include the secretion of proteins of the extracellular matrix and secretion of neurotransmitters into the synaptic cleft by synaptic vesicles.





Figure 1. In exocytosis, vesicles containing substances fuse with the plasma membrane. The contents are then released to the exterior of the cell. (credit: modification of work by Mariana Ruiz Villareal)

A summary of the cellular transport methods discussed is contained in Table 1, which also includes the energy requirements and materials transported by each.

Table 1. Methods of Transport, Energy Requirements, and Types of Material Transported

| Transport Method | Active/Passive | Material Transported |
|---|---|---|
| Diffusion | Passive | Small-molecular weight material |
| Osmosis | Passive | Water |
| Facilitated transport/diffusion | Passive | Sodium, potassium, calcium, glucose |
| Primary active transport | Active | Sodium, potassium, calcium |
| Secondary active transport | Active | Amino acids, lactose |
| Phagocytosis | Active | Large macromolecules, whole cells, or cellular structures |
| Pinocytosis and potocytosis | Active | Small molecules (liquids/water) |
| Receptor-mediated endocytosis | Active | Large quantities of macromolecules |
| Exocytosis | Active | Waste materials, proteins for the extracellular matrix, neurotransmitters |

## In Summary: Exocytosis

Exocytosis in many ways is the reverse process from endocytosis. Here cells expel material through the fusion of vesicles with the plasma membrane and subsequent dumping of their content into the extracellular fluid.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

7.11: Exocytosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.12: Putting It Together- Cell Membranes

Let's return to our discussion of cystic fibrosis. Cystic fibrosis (CF) is caused by a defect in a single transmembrane protein: cystic fibrosis transmembrane conductance regulator (CFTR), as seen in Figure 1. This regulator is a chloride ion channel that crosses through the plasma membrane. This channel is specifically active in epithelial tissues where it normally helps produce thin mucus via the movement of water. When mutated, the channel does not work properly and mucus becomes thick and viscous. This in turn leads directly to many of the symptoms of CF: thick, sticky mucus, frequent chest infections, and coughing or shortness of breath.





Figure 1. Panel 1 shows a properly working CFTR protein. Panel 2 shows a malfunctioning CFTR protein.

## Treatment

Cystic fibrosis is a difficult disease to treat. As we mentioned at the beginning of the module, patients with CF often suffer lung infections and sometimes require lung transplants. In addition to this, many CF patients are on one or more antibiotics at all times —even when healthy—to suppress infection. Several mechanical techniques are used to dislodge sputum and encourage its expectoration. In the hospital setting, chest physiotherapy is utilized. As lung disease worsens, mechanical breathing support may become necessary. Bi-lateral lung transplantation often becomes necessary for individuals with cystic fibrosis as lung function and exercise tolerance declines.

Gene therapy has been explored as a potential cure for cystic fibrosis. Ideally, gene therapy attempts to place a normal copy of the CFTR gene into affected cells. Transferring the normal CFTR gene into the affected epithelium cells would result in the production of functional CFTR in all target cells, without adverse reactions or an inflammation response. Studies have shown that to prevent the lung manifestations of cystic fibrosis, only 5–10 percent the normal amount of CFTR gene expression is needed.

Finally, a number of small molecules that aim at compensating various mutations of the CFTR gene are under development. About 10 percent of CF cases result from a premature stop codon in the DNA, leading to early termination of protein synthesis and truncated proteins. One approach to combating a faulty receptor is to develop drugs that get the ribosome to overcome this premature stop codon and synthesize a full-length CFTR protein.

**Try It**

- Cystic Fibrosis Foundation
- Cystic Fibrosis on the Mayo Clinic

## Contributors and Attributions

CC licensed content, Original

- Revision and Modification. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-SA: Attribution-ShareAlike*

CC licensed content, Shared previously

- Cystic Fibrosis. **Provided by**: Boundless. **Located at**: https://courses.lumenlearning.com/boundless-ap/. **License**: *CC BY-SA: Attribution-ShareAlike*
- CFTR Protein Panels. **Authored by**: Lbudd14. **Located at**: https://commons.wikimedia.org/wiki/File:CFTR_Protein_Panels.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

---

7.12: Putting It Together- Cell Membranes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 7.13: Assignment- Membranes Alive!

For this assessment you will conduct a kitchen science experiment in which you study the movement of materials into and out of cells. Individual cells are generally too small to see; however, a chicken egg is actually a single cell. Please note that on the whole this lab will take you 3 days to complete. However, there is less than 1 hour of 'hands-on' work on the first and last days.

Use the Membrane Lab procedure for your experimental set up.

When complete, write a lab report in MLA format that includes the following sections.

- Introduction
- Hypotheses
- Materials and Methods
- Results
- Discussion
- Conclusion
- References

Use this example to give you an idea of how your experiment may look in progress, and to help with completing your lab report.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references
- Assignment is formatted throughout using the MLA style.
- Assignment includes all the identified components of a lab report (hypothesis, materials and methods, results, discussion, conclusion). Images and tables are included where required and as appropriate.

## Rubric

| Membranes Alive! |
| --- |
| Outcome: Describe and explain the structure and function of membranes |
| |

| Membranes Alive! | | | | |
|---|---|---|---|---|
| Criteria | Ratings | | | Pts |
| Introduction and conclusion sections discuss how a membrane is selectively permeable and how the experimental results demonstrate that. | Introduction and conclusion sections address selectively permeable nature of membranes and experimental results in detail.<br>5.0 pts | Introduction and conclusion sections address selectively permeable nature of membranes and experimental results.<br>4.0 pts | Introduction and/or conclusion sections lack sufficient detail or do not connect to the experiment and results.<br>0.0 pts | 5 pts |
| Discussion and conclusion section address the movement of the tested substances across the membrane. | Movement of tested substances addressed correctly and in detail in both discussion and conclusion sections. Illustrations may be included.<br>5.0 pts | Movement of tested substances addressed correctly in both discussion and conclusion sections.<br>4.0 pts | Movement of tested substances not addressed correctly in both discussion and conclusion sections.<br>0.0 pts | 5 pts |
| Proper terminology is used to demonstrate an understanding of transport mechanisms and how substances move across membranes in different types of solutions. Both direction and relative speed of transport is discussed. | Terminology is used accurately and speed/direction are discussed correctly and in detail. Illustrations may be included.<br>5.0 pts | Terminology is used accurately and speed/direction are discussed correctly. Illustrations may be included.<br>4.0 pts | Terminology is used incorrectly; direction and/or speed are not correctly discussed.<br>0.0 pts | 5 pts |
| | | | | Total points: 15 |

## Contributors and Attributions

CC licensed content, Original

- Performance Assessment: Membranes Alive!. **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

7.13: Assignment- Membranes Alive! is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 8: Module 6- Metabolic Pathways


8.1: Why It Matters- Metabolic Pathways
8.2: Introduction to Redox Reactions
8.3: Electrons and Energy
8.4: ATP in Living Systems
8.5: Introduction to Photosynthesis
8.6: An Overview of Photosynthesis
8.7: Spectrums of Light
8.8: Light-Dependent Reactions
8.9: Light-Independent Reactions
8.10: Photosynthesis
8.11: Introduction to Cellular Respiration
8.12: Glycolysis
8.13: Pyruvate Oxidation
8.14: Citric Acid Cycle
8.15: Electron Transport Chain
8.16: Cellular Respiration
8.17: Introduction to Fermentation
8.18: Metabolism without Oxygen
8.19: Types of Fermentation
8.20: Putting It Together- Metabolic Pathways
8.21: Assignment- Observing Energy Transactions



8: Module 6- Metabolic Pathways is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.1: Why It Matters- Metabolic Pathways

## Why explain the metabolic pathways involved in the capture and release of energy in cells?

Every time you move—or even breathe—you're using energy. All living things are continually using energy; thus, they need a way to create or obtain new energy. Two of these ways are photosynthesis and cellular respiration.

Plants (and other autotrophs) undergo photosynthesis to create energy. Humans (and other heterotrophs) on the other hand must consume something that has energy (like plants or other animals)—we take this energy and convert it into a form our body can use. This process is known as cellular respiration.

Watch this 5 minute video for an overview of why even small changes in the global climate have the potential for big impacts on our daily lives through our food sources.



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=238

- What role does farming play in giving us energy to use every day?
- How do plants get energy to grow, and how do we then get our energy from them?

Ok, let's see where we get all the energy to stay awake during biology class!

### Learning Outcomes

- Understand the role movement of electrons plays in energy exchanges in cells
- Identify the basic components and steps of photosynthesis
- Identify the reactants and products of cellular respiration and where these reactions occur in a cell
- Illustrate the basic components and steps of fermentation

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Metabolic Pathways. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

All rights reserved content

- Withering Crops -- Changing Planet. **Authored by**: National Science Foundation . **Located at**: https://youtu.be/cBqr43QbfBE. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

8.1: Why It Matters- Metabolic Pathways is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.2: Introduction to Redox Reactions

## Understand the role movement of electrons plays in energy exchanges in cells

Energy production within a cell involves many coordinated chemical pathways. Most of these pathways are combinations of oxidation and reduction reactions. Oxidation and reduction occur in tandem. An oxidation reaction strips an electron from an atom in a compound, and the addition of this electron to another compound is a reduction reaction. Because oxidation and reduction usually occur together, these pairs of reactions are called oxidation reduction reactions, or **redox reactions**.

## What You'll Learn to Do

- Relate the movement of electrons to oxidation-reduction (redox) reactions
- Describe how cells store and transfer free energy using ATP

### Learning Activities

The learning activities for this section include the following:

- Electrons and Energy
- ATP in Living Systems
- Self Check: Redox Reactions

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.2: Introduction to Redox Reactions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 8.3: Electrons and Energy

Let's imagine that you are a cell. You've just been given a big, juicy glucose molecule, and you'd like to convert some of the energy in this glucose molecule into a more usable form, one that you can use to power your metabolic reactions. How can you go about this? What's the best way for you to squeeze as much energy as possible out of that glucose molecule, and to capture this energy in a handy form?

Fortunately for us, our cells—and those of other living organisms—are excellent at harvesting energy from glucose and other organic molecules, such as fats and amino acids. Here, we'll go through a quick overview of how cells break down fuels, then look at the electron transfer reactions (redox reactions) that are key to this process.

### Overview of Fuel Breakdown Pathways

The reactions that allow energy to be extracted from molecules such as glucose, fats, and amino acids are called **catabolic reactions**, meaning that they involve breaking a larger molecule into smaller pieces. For example, when glucose is broken down in the presence of oxygen, it's converted into six carbon dioxide molecules and six water molecules. The overall reaction for this process can be written as:



This reaction, as written, is simply a combustion reaction, similar to what takes place when you burn a piece of wood in a fireplace or gasoline in an engine. Does this mean that glucose is continually combusting inside of your cells? Thankfully, not quite! The combustion reaction describes the overall process that takes place, but inside of a cell, this process is broken down into many smaller steps. Energy contained in the bonds of glucose is released in small bursts, and some of it can be captured in the form of adenosine triphosphate (ATP), a small molecule that is used to power reactions in the cell. Much of the energy from glucose is still lost as heat, but enough is captured to keep the metabolism of the cell running.

As a glucose molecule is gradually broken down, some of the breakdowns steps release energy that is captured directly as ATP. In these steps, a phosphate group is transferred from a pathway intermediate straight to ADP, a process known as **substrate-level phosphorylation**. Many more steps, however, produce ATP in an indirect way. In these steps, electrons from glucose are transferred to small molecules known as electron carriers. The electron carriers take the electrons to a group of proteins in the inner membrane of the mitochondrion, called the electron transport chain. As electrons move through the electron transport chain, they go from a higher to a lower energy level and are ultimately passed to oxygen (forming water). Energy released in the electron transport chain is captured as a proton gradient, which powers production of ATP by a membrane protein called ATP synthase. This process is known as **oxidative phosphorylation**. A simplified diagram of oxidative and substrate-level phosphorylation is shown below.





Image modified from "Etc4" by Fvasconcellos (public domain).

When organic fuels like glucose are broken down using an electron transport chain that ends with oxygen, the breakdown process is known as **aerobic respiration** (aerobic = oxygen-requiring). Most eukaryotic cells, as well as many bacteria and other prokaryotes, can carry out aerobic respiration. Some prokaryotes have pathways similar to aerobic respiration, but with a different inorganic molecule, such as sulfur, substituted for oxygen. These pathways are not oxygen-dependent, so the breakdown process is called **anaerobic respiration** (anaerobic = non-oxygen-requiring). Officially, both processes are examples of **cellular respiration**, the breakdown of down organic fuels using an electron transport chain. However, cellular respiration is commonly used as a synonym for aerobic respiration, and we'll use it that way here[1].

### Redox Reactions


Cellular respiration involves many reactions in which electrons are passed from one molecule to another. Reactions involving electron transfers are known as **oxidation-reduction reactions** (or **redox reactions**), and they play a central role in the metabolism of a cell. In a redox reaction, one of the reacting molecules loses electrons and is said to be **oxidized**, while another reacting molecule gains electrons (the ones lost by the first molecule) and is said to be **reduced**. You can remember what oxidation and reduction mean with the handy mnemonic "LEO goes GER": *L*ose *E*lectrons, *O*xidized; *G*ain *E*lectrons,*R*educed. The formation of magnesium chloride is one simple example of a redox reaction:



In this reaction, the magnesium atom loses two electrons, so it is oxidized. These two electrons are accepted by chlorine, which is reduced. The atom or molecule that donates electrons (in this case, magnesium) is called the **reducing agent**, because its donation of electrons allows another molecule to become reduced. The atom or molecule that accepts the electrons (in this case, chlorine) is known as the **oxidizing agent**, because its acceptance of electrons allows the other molecule to become oxidized.

### Redox Reactions with Carbon-containing Molecules

When a reaction involves the formation of ions, as in the example with magnesium and chlorine above, it's relatively easy to see that electrons are being transferred. Not all redox reactions involve the complete transfer of electrons, though, and this is particularly true of reactions important in cellular metabolism. Instead, some redox reactions simply change the amount of electron density on a particular atom by altering how it shares electrons in covalent bonds. As an example, let's consider the combustion of butane:



Butane: [latex]2\text{C}_4\text{H}_{10}+13\text{O}_2\to8\text{CO}_2+10\text{H}_2\text{O}[/latex]

What's the electron-sharing situation at the start of the reaction? In butane, the carbon atoms are all bonded to other carbons and hydrogens. In $\text{C}-\text{C}$ bonds, electrons are shared equally, and in $\text{C}-\text{H}$ bonds, the $\text{C}$ atom has a very slight negative charge (since it's a bit more electronegative than hydrogen). Similarly, when oxygens are bonded to one another in $\text{O}_2$, start subscript, 2, end subscript, electrons are shared very equally. After the reaction, however, the electron-sharing picture looks quite different. Oxygen is much more electronegative than carbon, so the in the $\text{C}=\text{O}$ bonds of carbon dioxide, oxygen will "hog" the bond electrons. In the $\text{O}-\text{H}$ bonds of water, oxygen will similarly pull electrons away from the hydrogen atoms. Thus, relative to its state before the reaction, carbon has lost electron density (because oxygen is now hogging its electrons), while oxygen has gained electron density (because it can now hog electrons shared with other elements). It's thus reasonable to say that carbon was oxidized during this reaction, while oxygen was reduced. (Hydrogen arguably loses a little electron density too, though its electrons were being hogged to some degree in either case.) Biologists often refer to whole molecules, rather than individual atoms, as being reduced or oxidized; thus, we can say that butane—the source of the carbons—is oxidized, while molecular oxygen—the source of the oxygen atoms—is reduced.


It's important to understand that oxidation and reduction reactions are fundamentally about the transfer of electrons. In the context of biology, however, you may find it helpful to use the gain or loss of H and O atoms as a proxy for the transfer of electrons. As a general rule of thumb, if a carbon-containing molecule gains H atoms or loses O atoms during a reaction, it's likely been reduced (gained electrons). Conversely, if it loses H atoms or gains O atoms, it's probably been oxidized (lost electrons). For example, let's go back to the reaction for glucose breakdown, $\text{C}_6\text{H}_{12}\text{O}_6+6\text{O}_2\to6\text{CO}_2+6\text{H}_2\text{O}$. In glucose, carbon is associated with H atoms, while in carbon dioxide, no Hs are present. Thus, we would predict that glucose is oxidized in this reaction.

We can confirm this if we look at the actual electron shifts involved, as in the video below:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=242

## Energy in Redox Reactions




Click on the image for a larger view. Image based on similar diagram by Ryan Gutierrez.

Like other chemical reactions, redox reactions involve a free energy change. Reactions that move the system from a higher to a lower energy state are spontaneous and release energy, while those that do the opposite require an input of energy. In redox reactions, energy is released when an electron loses potential energy as a result of the transfer. Electrons have more potential energy when they are associated with less electronegative atoms (such as C or H), and less potential energy when they are associated with a more electronegative atom (such as O). Thus, a redox reaction that moves electrons or electron density from a less to a more electronegative atom will be spontaneous and release energy. For instance, the combustion of butane (above) releases energy because there is a net shift of electron density away from carbon and hydrogen and onto oxygen. If you've heard it said that molecules like glucose have "high-energy" electrons, this is a reference to the relatively high potential energy of the electrons in their $\text{C}-\text{C}$ and $\text{C}-\text{H}$ bonds.

Quite a bit of energy can be released when electrons in $\text{C}-\text{C}$ and $\text{C}-\text{H}$ bonds are shifted to oxygen. In a cell, however, it's not a great idea to release all that energy at once in a combustion reaction. Instead, cells harvest energy from glucose in a controlled fashion, capturing as much of it as possible in the form of ATP. This is accomplished by oxidizing glucose in a gradual, rather than an explosive, sort of way. There are two important ways in which this oxidation is gradual:


- Rather than pulling all the electrons off of glucose at the same time, cellular respiration strips them away in pairs. The redox reactions that remove electron pairs from glucose transfer them to small molecules called electron carriers.
- The electron carriers deposit their electrons in the electron transport chain, a series of proteins and organic molecules in the inner mitochondrial membrane. Electrons are passed from one component to the next in a series of energy-releasing steps, allowing energy to be captured in the form of an electrochemical gradient.

We'll look at both redox carriers and the electron transport chain in more detail below.

The removal of an electron from a molecule, oxidizing it, results in a decrease in potential energy in the oxidized compound. The electron (sometimes as part of a hydrogen atom), does not remain unbonded, however, in the cytoplasm of a cell. Rather, the electron is shifted to a second compound, reducing the second compound. The shift of an electron from one compound to another removes some potential energy from the first compound (the oxidized compound) and increases the potential energy of the second compound (the reduced compound). The transfer of electrons between molecules is important because most of the energy stored in atoms and used to fuel cell functions is in the form of high-energy electrons. The transfer of energy in the form of electrons allows the cell to transfer and use energy in an incremental fashion—in small packages rather than in a single, destructive burst. This module focuses on the extraction of energy from food; you will see that as you track the path of the transfers, you are tracking the path of electrons moving through metabolic pathways.

## Electron Carriers

Electron carriers, sometimes called electron shuttles, are small organic molecules that readily cycle between oxidized and reduced forms and are used to transport electrons during metabolic reactions. There are two electron carriers that play particularly important roles during cellular respiration: **$\text{NAD}^+$** (nicotinamide adenine dinucleotide, shown below) and **FAD** (flavin adenine dinucleotide). Both $\text{NAD}^+$ and FAD can serve as oxidizing agents, accepting a pair of electrons, along with one or more protons, to switch to their reduced forms. $\text{NAD}^+$ accepts two electrons and one $\text{H}^+$ to become NADH, while FAD accepts two electrons and two $\text{H}^+$ to become $\text{FADH}_2$. $\text{NAD}^+$ is the primary electron carrier used during cellular respiration, with FAD participating in just one (or two sometimes two) reactions.




The oxidized form of the electron carrier ($NAD^+$) is shown on the left and the reduced form (NADH) is shown on the right. The nitrogenous base in NADH has one more hydrogen ion and two more electrons than in $NAD^+$.

As shown in the image above, $NAD^+$ is a small organic molecule whose structure includes the RNA nucleotide adenine. (FAD is a similar type of molecule, although its functional groups are different.) Both molecules are B vitamin derivatives, with $NAD^+$ produced from niacin and FAD produced from riboflavin. $NAD^+$ and FAD are **coenzymes**, organic molecules that serve as helpers during enzyme-catalyzed reactions, and they receive electrons and protons as part of these reactions. Specifically, both $NAD^+$ and FAD serve as cofactors for enzymes called **dehydrogenases**, which remove one or more hydrogen atoms from their substrates.

## Electron Transport Chain

In their reduced forms, NADH and $FADH_2$ carry electrons to the electron transport chain in the inner mitochondrial membrane. They deposit their electrons at or near the beginning of the transport chain, and the electrons are then passed along from one protein or organic molecule to the next in a predictable series of steps. Importantly, the movement of electrons through the transport chain is energetically "downhill," such that energy is released at each step. In redox terms, this means that each member of the electron transport chain is more electronegative (electron-hungry) that the one before it, and less electronegative than the one after[2]. $NAD^+$, which deposits its electrons at the beginning of the chain as NADH, is the least electronegative, while oxygen, which receives the electrons at the end of the chain (along with $H^+$) to form water, is the most electronegative. As electrons trickle "downhill" through the transport chain, they release energy, and some of this energy is captured in the form of an electrochemical gradient and used to make ATP.

1. Reece, J. B., Urry, L. A., Cain, M. L., Wasserman, S. A., Minorsky, P. V., and Jackson, R. B. (2011). Cellular respiration and fermentation. In *Campbell biology* (10th ed., pp. 162–184). San Francisco, CA: Pearson. ↵
2. *Ibid*. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Introduction to cellular respiration and redox. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/cellular-respiration-and-fermentation/intro-to-cellular-respiration/a/intro-to-cellular-respiration-and-redox. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*
- Oxidation and reduction in cellular respiration. **Authored by**: Sal Khan. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/cellular-respiration-and-fermentation/intro-to-cellular-respiration/v/oxidation-and-reduction-in-cellular-respiration. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

8.3: Electrons and Energy is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.4: ATP in Living Systems

A living cell cannot store significant amounts of free energy. Excess free energy would result in an increase of heat in the cell, which would result in excessive thermal motion that could damage and then destroy the cell. Rather, a cell must be able to handle that energy in a way that enables the cell to store energy safely and release it for use only as needed. Living cells accomplish this by using the compound adenosine triphosphate (ATP). ATP is often called the “energy currency” of the cell, and, like currency, this versatile compound can be used to fill any energy need of the cell. How? It functions similarly to a rechargeable battery.

When ATP is broken down, usually by the removal of its terminal phosphate group, energy is released. The energy is used to do work by the cell, usually by the released phosphate binding to another molecule, activating it. For example, in the mechanical work of muscle contraction, ATP supplies the energy to move the contractile muscle proteins. Recall the active transport work of the sodium-potassium pump in cell membranes. ATP alters the structure of the integral protein that functions as the pump, changing its affinity for sodium and potassium. In this way, the cell performs work, pumping ions against their electrochemical gradients.

## ATP Structure and Function





Figure 1. ATP (adenosine triphosphate) has three phosphate groups that can be removed by hydrolysis to form ADP (adenosine diphosphate) or AMP (adenosine monophosphate).The negative charges on the phosphate group naturally repel each other, requiring energy to bond them together and releasing energy when these bonds are broken.

At the heart of ATP is a molecule of adenosine monophosphate (AMP), which is composed of an adenine molecule bonded to a ribose molecule and to a single phosphate group (Figure 1). Ribose is a five-carbon sugar found in RNA, and AMP is one of the nucleotides in RNA. The addition of a second phosphate group to this core molecule results in the formation of adenosine diphosphate (ADP); the addition of a third phosphate group forms adenosine triphosphate (ATP).

The addition of a phosphate group to a molecule requires energy. Phosphate groups are negatively charged and thus repel one another when they are arranged in series, as they are in ADP and ATP. This repulsion makes the ADP and ATP molecules inherently unstable. The release of one or two phosphate groups from ATP, a process called **dephosphorylation**, releases energy.

### Energy from ATP

Hydrolysis is the process of breaking complex macromolecules apart. During hydrolysis, water is split, or lysed, and the resulting hydrogen atom ($H^+$) and a hydroxyl group ($OH^-$) are added to the larger molecule. The hydrolysis of ATP produces ADP, together with an inorganic phosphate ion ($P_i$), and the release of free energy. To carry out life processes, ATP is continuously broken down into ADP, and like a rechargeable battery, ADP is continuously regenerated into ATP by the reattachment of a third phosphate group. Water, which was broken down into its hydrogen atom and hydroxyl group during ATP hydrolysis, is regenerated when a third phosphate is added to the ADP molecule, reforming ATP.

Obviously, energy must be infused into the system to regenerate ATP. Where does this energy come from? In nearly every living thing on earth, the energy comes from the metabolism of glucose. In this way, ATP is a direct link between the limited set of exergonic pathways of glucose catabolism and the multitude of endergonic pathways that power living cells.

## Phosphorylation

Recall that, in some chemical reactions, enzymes may bind to several substrates that react with each other on the enzyme, forming an intermediate complex. An intermediate complex is a temporary structure, and it allows one of the substrates (such as ATP) and reactants to more readily react with each other; in reactions involving ATP, ATP is one of the substrates and ADP is a product. During an endergonic chemical reaction, ATP forms an intermediate complex with the substrate and enzyme in the reaction. This

intermediate complex allows the ATP to transfer its third phosphate group, with its energy, to the substrate, a process called phosphorylation. **Phosphorylation** refers to the addition of the phosphate (~P). This is illustrated by the following generic reaction:

$$A + \text{enzyme} + ATP \rightarrow [A - \text{enzyme} - \sim P] \rightarrow B + \text{enzyme} + ADP + \text{phosphate ion}$$

When the intermediate complex breaks apart, the energy is used to modify the substrate and convert it into a product of the reaction. The ADP molecule and a free phosphate ion are released into the medium and are available for recycling through cell metabolism.

## Substrate Phosphorylation

ATP is generated through two mechanisms during the breakdown of glucose. A few ATP molecules are generated (that is, regenerated from ADP) as a direct result of the chemical reactions that occur in the catabolic pathways. A phosphate group is removed from an intermediate reactant in the pathway, and the free energy of the reaction is used to add the third phosphate to an available ADP molecule, producing ATP (Figure 2). This very direct method of phosphorylation is called **substrate-level phosphorylation**.





Figure 2. In phosphorylation reactions, the gamma phosphate of ATP is attached to a protein.

## Oxidative Phosphorylation

Most of the ATP generated during glucose catabolism, however, is derived from a much more complex process, chemiosmosis, which takes place in mitochondria (Figure 3) within a eukaryotic cell or the plasma membrane of a prokaryotic cell.





Figure 3. The mitochondria (Credit: modification of work by Mariana Ruiz Villareal)

**Chemiosmosis**, a process of ATP production in cellular metabolism, is used to generate 90 percent of the ATP made during glucose catabolism and is also the method used in the light reactions of photosynthesis to harness the energy of sunlight. The production of ATP using the process of chemiosmosis is called **oxidative phosphorylation** because of the involvement of oxygen in the process.

**Try It**

What happens when the critical reactions of cellular respiration do not proceed correctly? Mitochondrial diseases are genetic disorders of metabolism. Mitochondrial disorders can arise from mutations in nuclear or mitochondrial DNA, and they result in the production of less energy than is normal in body cells. In type 2 diabetes, for instance, the oxidation efficiency of NADH is reduced, impacting oxidative phosphorylation but not the other steps of respiration. Symptoms of mitochondrial diseases can include muscle weakness, lack of coordination, stroke-like episodes, and loss of vision and hearing. Most affected people are

diagnosed in childhood, although there are some adult-onset diseases. Identifying and treating mitochondrial disorders is a specialized medical field. The educational preparation for this profession requires a college education, followed by medical school with a specialization in medical genetics. Medical geneticists can be board certified by the American Board of Medical Genetics and go on to become associated with professional organizations devoted to the study of mitochondrial diseases, such as the Mitochondrial Medicine Society and the Society for Inherited Metabolic Disease.

## Learning Objectives

ATP functions as the energy currency for cells. It allows the cell to store energy briefly and transport it within the cell to support endergonic chemical reactions. The structure of ATP is that of an RNA nucleotide with three phosphates attached. As ATP is used for energy, a phosphate group or two are detached, and either ADP or AMP is produced. Energy derived from glucose catabolism is used to convert ADP into ATP. When ATP is used in a reaction, the third phosphate is temporarily attached to a substrate in a process called phosphorylation. The two processes of ATP regeneration that are used in conjunction with glucose catabolism are substrate-level phosphorylation and oxidative phosphorylation through the process of chemiosmosis.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.4: ATP in Living Systems is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.5: Introduction to Photosynthesis

## Identify the basic components and steps of photosynthesis

The processes in all organisms—from bacteria to humans—require energy. To get this energy, many organisms access stored energy by eating, that is, by ingesting other organisms. But where does the stored energy in food originate? All of this energy can be traced back to photosynthesis.

Photosynthesis is essential to all life on earth; both plants and animals depend on it. It is the only biological process that can capture energy that originates in outer space (sunlight) and convert it into chemical compounds (carbohydrates) that every organism uses to power its metabolism. In brief, the energy of sunlight is captured and used to energize electrons, which are then stored in the covalent bonds of sugar molecules. How long lasting and stable are those covalent bonds? The energy extracted today by the burning of coal and petroleum products represents sunlight energy captured and stored by photosynthesis around 300 million years ago.

Plants, algae, and a group of bacteria called cyanobacteria are the only organisms capable of performing photosynthesis (Figure 1). Because they use light to manufacture their own food, they are called **photoautotrophs** (literally, “self-feeders using light”). Other organisms, such as animals, fungi, and most other bacteria, are termed **heterotrophs** (“other feeders”), because they must rely on the sugars produced by photosynthetic organisms for their energy needs. A third very interesting group of bacteria synthesize sugars, not by using sunlight’s energy, but by extracting energy from inorganic chemical compounds; hence, they are referred to as **chemoautotrophs**.





Figure 2. Photoautotrophs including (a) plants, (b) algae, and (c) cyanobacteria synthesize their organic compounds via photosynthesis using sunlight as an energy source. Cyanobacteria and planktonic algae can grow over enormous areas in water, at times completely covering the surface. In a (d) deep sea vent, chemoautotrophs, such as these (e) thermophilic bacteria, capture energy from inorganic compounds to produce organic compounds. The ecosystem surrounding the vents has a diverse array of animals, such as tubeworms, crustaceans, and octopuses that derive energy from the bacteria. (credit a: modification of work by Steve Hillebrand, U.S. Fish and Wildlife Service; credit b: modification of work by “eutrophication&hypoxia”/Flickr; credit c: modification of work by NASA; credit d: University of Washington, NOAA; credit e: modification of work by Mark Amend, West Coast and Polar Regions Undersea Research Center, UAF, NOAA)



Figure 2. The energy stored in carbohydrate molecules from photosynthesis passes through the food chain. The predator that eats these deer receives a portion of the energy that originated in the photosynthetic vegetation that the deer consumed. (credit: modification of work by Steve VanRiper, U.S. Fish and Wildlife Service)

The importance of photosynthesis is not just that it can capture sunlight's energy. A lizard sunning itself on a cold day can use the sun's energy to warm up. Photosynthesis is vital because it evolved as a way to store the energy in solar radiation (the "photo" part) as high-energy electrons in the carbon-carbon bonds of carbohydrate molecules (the "synthesis" part). Those carbohydrates are the energy source that heterotrophs use to power the synthesis of ATP via respiration. Therefore, photosynthesis powers 99 percent of Earth's ecosystems. When a top predator, such as a wolf, preys on a deer (Figure 2), the wolf is at the end of an energy path that went from nuclear reactions on the surface of the sun, to light, to photosynthesis, to vegetation, to deer, and finally to wolf.

## What You'll Learn to Do

- Identify the reactants and products of photosynthesis
- Describe the visible and electromagnetic spectrums of light as they applies to photosynthesis
- Describe the light-dependent reactions that take place during photosynthesis
- Identify the light-independent reactions in photosynthesis

### Learning Activities

The learning activities for this section include the following:

- An Overview of Photosynthesis
- Spectrums of Light
- Light-Dependent Reactions
- Light-Independent Reactions
- Photosynthesis
- Self Check: Photosynthesis

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

8.5: Introduction to Photosynthesis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.6: An Overview of Photosynthesis

Photosynthesis is a multi-step process that requires sunlight, carbon dioxide (which is low in energy), and water as substrates (Figure 1). After the process is complete, it releases oxygen and produces glyceraldehyde-3-phosphate (GA3P), simple carbohydrate molecules (which are high in energy) that can subsequently be converted into glucose, sucrose, or any of dozens of other sugar molecules. These sugar molecules contain energy and the energized carbon that all living things need to survive.





Figure 1. Photosynthesis uses solar energy, carbon dioxide, and water to produce energy-storing carbohydrates. Oxygen is generated as a waste product of photosynthesis.

The following is the chemical equation for photosynthesis (Figure 2):

| Photosynthesis Equation | | | | | | | |
|---|---|---|---|---|---|---|---|
| Carbon dioxide | + | Water | SUNLIGHT → | Sugar | + | Oxygen | |
| $6CO_2$ | | $6H_2O$ | | $C_6H_{12}O_6$ | | $6O_2$ | |

Figure 2. The basic equation for photosynthesis is deceptively simple. In reality, the process takes place in many steps involving intermediate reactants and products. Glucose, the primary energy source in cells, is made from two three-carbon GA3Ps.

Although the equation looks simple, the many steps that take place during photosynthesis are actually quite complex. Before learning the details of how photoautotrophs turn sunlight into food, it is important to become familiar with the structures involved.

In plants, photosynthesis generally takes place in leaves, which consist of several layers of cells. The process of photosynthesis occurs in a middle layer called the **mesophyll**. The gas exchange of carbon dioxide and oxygen occurs through small, regulated openings called **stomata** (singular: stoma), which also play roles in the regulation of gas exchange and water balance. The stomata

are typically located on the underside of the leaf, which helps to minimize water loss. Each stoma is flanked by guard cells that regulate the opening and closing of the stomata by swelling or shrinking in response to osmotic changes.

In all autotrophic eukaryotes, photosynthesis takes place inside an organelle called a **chloroplast**. For plants, chloroplast-containing cells exist in the mesophyll. Chloroplasts have a double membrane envelope (composed of an outer membrane and an inner membrane). Within the chloroplast are stacked, disc-shaped structures called **thylakoids**. Embedded in the thylakoid membrane is chlorophyll, a **pigment** (molecule that absorbs light) responsible for the initial interaction between light and plant material, and numerous proteins that make up the electron transport chain. The thylakoid membrane encloses an internal space called the **thylakoid lumen**. As shown in Figure 3, a stack of thylakoids is called a **granum**, and the liquid-filled space surrounding the granum is called **stroma** or "bed" (not to be confused with stoma or "mouth," an opening on the leaf epidermis).

Practice Question





Figure 3. Photosynthesis takes place in chloroplasts, which have an outer membrane and an inner membrane. Stacks of thylakoids called grana form a third membrane layer.

On a hot, dry day, plants close their stomata to conserve water. What impact will this have on photosynthesis?

[practice-area rows="2"][/practice-area]
[reveal-answer q="173275"]**Show Answer**[/reveal-answer]
[hidden-answer a="173275"]Levels of carbon dioxide (a necessary photosynthetic substrate) will immediately fall. As a result, the rate of photosynthesis will be inhibited.[/hidden-answer]

## Two Parts of Photosynthesis

Photosynthesis takes place in two sequential stages: the light-dependent reactions and the light independent-reactions. In the **light-dependent reactions**, energy from sunlight is absorbed by chlorophyll and that energy is converted into stored chemical energy. In the **light-independent reactions**, the chemical energy harvested during the light-dependent reactions drive the assembly of sugar molecules from carbon dioxide. Therefore, although the light-independent reactions do not use light as a reactant, they require the products of the light-dependent reactions to function. In addition, several enzymes of the light-independent reactions are activated by light. The light-dependent reactions utilize certain molecules to temporarily store the energy: These are referred to as energy carriers. The energy carriers that move energy from light-dependent reactions to light-independent reactions can be thought of as "full" because they are rich in energy. After the energy is released, the "empty" energy carriers return to the light-dependent reaction to obtain more energy. Figure 4 illustrates the components inside the chloroplast where the light-dependent and light-independent reactions take place.





Figure 4. Photosynthesis takes place in two stages: light dependent reactions and the Calvin cycle. Light-dependent reactions, which take place in the thylakoid membrane, use light energy to make ATP and NADPH. The Calvin cycle, which takes place in the stroma, uses energy derived from these compounds to make GA3P from $CO_2$.

Click the link to learn more about photosynthesis.

## Try It




Figure 5. Foods that humans consume originate from photosynthesis. (credit: Associação Brasileira de Supermercados)

Major grocery stores in the United States are organized into departments, such as dairy, meats, produce, bread, cereals, and so forth. Each aisle (Figure 5) contains hundreds, if not thousands, of different products for customers to buy and consume.

Although there is a large variety, each item links back to photosynthesis. Meats and dairy link because the animals were fed plant-based foods. The breads, cereals, and pastas come largely from starchy grains, which are the seeds of photosynthesis-dependent plants. What about desserts and drinks? All of these products contain sugar—sucrose is a plant product, a disaccharide, a carbohydrate molecule, which is built directly from photosynthesis. Moreover, many items are less obviously derived from plants: for instance, paper goods are generally plant products, and many plastics (abundant as products and packaging) can be derived from algae or from oil, the fossilized remains of photosynthetic organisms. Virtually every spice and flavoring in the spice aisle was produced by a plant as a leaf, root, bark, flower, fruit, or stem. Ultimately, photosynthesis connects to every meal and every food a person consumes.

## Learning Objectives

The process of photosynthesis transformed life on Earth. By harnessing energy from the sun, photosynthesis evolved to allow living things access to enormous amounts of energy. Because of photosynthesis, living things gained access to sufficient energy that allowed them to build new structures and achieve the biodiversity evident today.

Only certain organisms, called photoautotrophs, can perform photosynthesis; they require the presence of chlorophyll, a specialized pigment that absorbs certain portions of the visible spectrum and can capture energy from sunlight. Photosynthesis uses carbon dioxide and water to assemble carbohydrate molecules and release oxygen as a waste product into the atmosphere. Eukaryotic autotrophs, such as plants and algae, have organelles called chloroplasts in which photosynthesis takes place, and starch accumulates. In prokaryotes, such as cyanobacteria, the process is less localized and occurs within folded membranes, extensions of the plasma membrane, and in the cytoplasm.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.6: An Overview of Photosynthesis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.7: Spectrums of Light

How can light be used to make food? When a person turns on a lamp, electrical energy becomes light energy. Like all other forms of kinetic energy, light can travel, change form, and be harnessed to do work. In the case of photosynthesis, light energy is converted into chemical energy, which photoautotrophs use to build carbohydrate molecules. However, autotrophs only use a few specific components of sunlight.

## What Is Light Energy?

The sun emits an enormous amount of electromagnetic radiation (solar energy). Humans can see only a fraction of this energy, which portion is therefore referred to as “visible light.” The manner in which solar energy travels is described as waves. Scientists can determine the amount of energy of a wave by measuring its wavelength, the distance between consecutive points of a wave. A single wave is measured from two consecutive points, such as from crest to crest or from trough to trough (Figure 1).





Figure 1. The wavelength of a single wave is the distance between two consecutive points of similar position (two crests or two troughs) along the wave.

Visible light constitutes only one of many types of electromagnetic radiation emitted from the sun and other stars. Scientists differentiate the various types of radiant energy from the sun within the electromagnetic spectrum. The electromagnetic spectrum is the range of all possible frequencies of radiation (Figure 2). The difference between wavelengths relates to the amount of energy carried by them.





Figure 2. The sun emits energy in the form of electromagnetic radiation. This radiation exists at different wavelengths, each of which has its own characteristic energy. All electromagnetic radiation, including visible light, is characterized by its wavelength.

Each type of electromagnetic radiation travels at a particular wavelength. The longer the wavelength (or the more stretched out it appears in the diagram), the less energy is carried. Short, tight waves carry the most energy. This may seem illogical, but think of it in terms of a piece of moving a heavy rope. It takes little effort by a person to move a rope in long, wide waves. To make a rope move in short, tight waves, a person would need to apply significantly more energy.

The electromagnetic spectrum (Figure 2) shows several types of electromagnetic radiation originating from the sun, including X-rays and ultraviolet (UV) rays. The higher-energy waves can penetrate tissues and damage cells and DNA, explaining why both X-rays and UV rays can be harmful to living organisms.

## Absorption of Light

Light energy initiates the process of photosynthesis when pigments absorb the light. Organic pigments, whether in the human retina or the chloroplast thylakoid, have a narrow range of energy levels that they can absorb. Energy levels lower than those represented by red light are insufficient to raise an orbital electron to a populatable, excited (quantum) state. Energy levels higher than those in blue light will physically tear the molecules apart, called bleaching. So retinal pigments can only “see” (absorb) 700 nm to 400 nm light, which is therefore called visible light. For the same reasons, plants pigment molecules absorb only light in the wavelength range of 700 nm to 400 nm; plant physiologists refer to this range for plants as photosynthetically active radiation.

The visible light seen by humans as white light actually exists in a rainbow of colors. Certain objects, such as a prism or a drop of water, disperse white light to reveal the colors to the human eye. The visible light portion of the electromagnetic spectrum shows the rainbow of colors, with violet and blue having shorter wavelengths, and therefore higher energy. At the other end of the spectrum toward red, the wavelengths are longer and have lower energy (Figure 3).





Figure 3. The colors of visible light do not carry the same amount of energy. Violet has the shortest wavelength and therefore carries the most energy, whereas red has the longest wavelength and carries the least amount of energy. (credit: modification of work by NASA)

## Understanding Pigments

Different kinds of pigments exist, and each has evolved to absorb only certain wavelengths (colors) of visible light. Pigments reflect or transmit the wavelengths they cannot absorb, making them appear in the corresponding color.

Chlorophylls and carotenoids are the two major classes of photosynthetic pigments found in plants and algae; each class has multiple types of pigment molecules. There are five major chlorophylls: *a*, *b*, *c* and *d* and a related molecule found in prokaryotes called bacteriochlorophyll. **Chlorophyll *a*** and **chlorophyll *b*** are found in higher plant chloroplasts and will be the focus of the following discussion.

With dozens of different forms, carotenoids are a much larger group of pigments. The carotenoids found in fruit—such as the red of tomato (lycopene), the yellow of corn seeds (zeaxanthin), or the orange of an orange peel (β-carotene)—are used as advertisements to attract seed dispersers. In photosynthesis,
**carotenoids** function as photosynthetic pigments that are very efficient molecules for the disposal of excess energy. When a leaf is exposed to full sun, the light-dependent reactions are required to process an enormous amount of energy; if that energy is not handled properly, it can do significant damage. Therefore, many carotenoids reside in the thylakoid membrane, absorb excess energy, and safely dissipate that energy as heat.

Each type of pigment can be identified by the specific pattern of wavelengths it absorbs from visible light, which is the **absorption spectrum**. The graph in Figure 4 shows the absorption spectra for chlorophyll *a*, chlorophyll *b*, and a type of carotenoid pigment called β-carotene (which absorbs blue and green light). Notice how each pigment has a distinct set of peaks and troughs, revealing a highly specific pattern of absorption. Chlorophyll *a* absorbs wavelengths from either end of the visible spectrum (blue and red), but not green. Because green is reflected or transmitted, chlorophyll appears green. Carotenoids absorb in the short-wavelength blue region, and reflect the longer yellow, red, and orange wavelengths.





Figure 4. (a) Chlorophyll a, (b) chlorophyll b, and (c) β-carotene are hydrophobic organic pigments found in the thylakoid membrane. Chlorophyll a and b, which are identical except for the part indicated in the red box, are responsible for the green color of leaves. β-carotene is responsible for the orange color in carrots. Each pigment has (d) a unique absorbance spectrum.




Figure 5. Plants that commonly grow in the shade have adapted to low levels of light by changing the relative concentrations of their chlorophyll pigments. (credit: Jason Hollinger)

Many photosynthetic organisms have a mixture of pigments; using them, the organism can absorb energy from a wider range of wavelengths. Not all photosynthetic organisms have full access to sunlight. Some organisms grow underwater where light intensity and quality decrease and change with depth. Other organisms grow in competition for light. Plants on the rainforest floor must be able to absorb any bit of light that comes through, because the taller trees absorb most of the sunlight and scatter the remaining solar radiation (Figure 5).

When studying a photosynthetic organism, scientists can determine the types of pigments present by generating absorption spectra. An instrument called a **spectrophotometer** can differentiate which wavelengths of light a substance can absorb. Spectrophotometers measure transmitted light and compute from it the absorption. By extracting pigments from leaves and placing these samples into a spectrophotometer, scientists can identify which wavelengths of light an organism can absorb. Additional methods for the identification of plant pigments include various types of chromatography that separate the pigments by their relative affinities to solid and mobile phases.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

8.7: Spectrums of Light is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.8: Light-Dependent Reactions

The overall function of light-dependent reactions is to convert solar energy into chemical energy in the form of NADPH and ATP. This chemical energy supports the light-independent reactions and fuels the assembly of sugar molecules. The light-dependent reactions are depicted in Figure 1. Protein complexes and pigment molecules work together to produce NADPH and ATP.





Figure 1. A photosystem consists of a light-harvesting complex and a reaction center. Pigments in the light-harvesting complex pass light energy to two special chlorophyll a molecules in the reaction center. The light excites an electron from the chlorophyll a pair, which passes to the primary electron acceptor. The excited electron must then be replaced. In (a) photosystem II, the electron comes from the splitting of water, which releases oxygen as a waste product. In (b) photosystem I, the electron comes from the chloroplast electron transport chain discussed below.

The actual step that converts light energy into chemical energy takes place in a multiprotein complex called a **photosystem**, two types of which are found embedded in the thylakoid membrane, **photosystem II** (PSII) and **photosystem I** (PSI) (Figure 2). The two complexes differ on the basis of what they oxidize (that is, the source of the low-energy electron supply) and what they reduce (the place to which they deliver their energized electrons).

Both photosystems have the same basic structure; a number of antenna proteins to which the chlorophyll molecules are bound surround the reaction center where the photochemistry takes place. Each photosystem is serviced by the light-harvesting complex, which passes energy from sunlight to the reaction center; it consists of multiple antenna proteins that contain a mixture of 300–400 chlorophyll *a* and *b* molecules as well as other pigments like carotenoids. The absorption of a single photon or distinct quantity or "packet" of light by any of the chlorophylls pushes that molecule into an excited state. In short, the light energy has now been captured by biological molecules but is not stored in any useful form yet. The energy is transferred from chlorophyll to chlorophyll until eventually (after about a millionth of a second), it is delivered to the reaction center. Up to this point, only energy has been transferred between molecules, not electrons.

Practice Question





Figure 2. The photosystem II (PSII) reaction center and the photosystem I (PSI).

In the photosystem II (PSII) reaction center, energy from sunlight is used to extract electrons from water. The electrons travel through the chloroplast electron transport chain to photosystem I (PSI), which reduces $NADP^+$ to NADPH. The electron transport chain moves protons across the thylakoid membrane into the lumen. At the same time, splitting of water adds protons to the lumen, and reduction of NADPH removes protons from the stroma. The net result is a low pH in the thylakoid lumen, and a high pH in the stroma. ATP synthase uses this electrochemical gradient to make ATP. What is the initial source of electrons for the chloroplast electron transport chain?

a. water
b. oxygen
c. carbon dioxide
d. NADPH

[reveal-answer q=”829552″]Show Answer[/reveal-answer]
[hidden-answer a=”829552″]Water is the initial source.[/hidden-answer]

The reaction center contains a pair of chlorophyll *a* molecules with a special property. Those two chlorophylls can undergo oxidation upon excitation; they can actually give up an electron in a process called a **photoact**. It is at this step in the reaction center, that light energy is converted into an excited electron. All of the subsequent steps involve getting that electron onto the energy carrier NADPH for delivery to the Calvin cycle where the electron is deposited onto carbon for long-term storage in the form of a carbohydrate. PSII and PSI are two major components of the photosynthetic **electron transport chain**, which also includes the **cytochrome complex**. The cytochrome complex, an enzyme composed of two protein complexes, transfers the electrons from the carrier molecule plastoquinone (Pq) to the protein plastocyanin (Pc), thus enabling both the transfer of protons across the thylakoid membrane and the transfer of electrons from PSII to PSI.

The reaction center of PSII (called **P680**) delivers its high-energy electrons, one at the time, to the **primary electron acceptor**, and through the electron transport chain (Pq to cytochrome complex to plastocyanine) to PSI. P680's missing electron is replaced by extracting a low-energy electron from water; thus, water is split and PSII is re-reduced after every photoact. Splitting one $H_2O$ molecule releases two electrons, two hydrogen atoms, and one atom of oxygen. Splitting two molecules is required to form one molecule of diatomic $O_2$ gas. About 10 percent of the oxygen is used by mitochondria in the leaf to support oxidative phosphorylation. The remainder escapes to the atmosphere where it is used by aerobic organisms to support respiration.

As electrons move through the proteins that reside between PSII and PSI, they lose energy. That energy is used to move hydrogen atoms from the stromal side of the membrane to the thylakoid lumen. Those hydrogen atoms, plus the ones produced by splitting water, accumulate in the thylakoid lumen and will be used synthesize ATP in a later step. Because the electrons have lost energy prior to their arrival at PSI, they must be re-energized by PSI, hence, another photon is absorbed by the PSI antenna. That energy is relayed to the PSI reaction center (called **P700**). P700 is oxidized and sends a high-energy electron to $NADP^+$ to form NADPH. Thus, PSII captures the energy to create proton gradients to make ATP, and PSI captures the energy to reduce $NADP^+$ into NADPH. The two photosystems work in concert, in part, to guarantee that the production of NADPH will roughly equal the production of ATP. Other mechanisms exist to fine tune that ratio to exactly match the chloroplast's constantly changing energy needs.

## Generating an Energy Carrier: ATP

As in the intermembrane space of the mitochondria during cellular respiration, the buildup of hydrogen ions inside the thylakoid lumen creates a concentration gradient. The passive diffusion of hydrogen ions from high concentration (in the thylakoid lumen) to low concentration (in the stroma) is harnessed to create ATP, just as in the electron transport chain of cellular respiration. The ions build up energy because of diffusion and because they all have the same electrical charge, repelling each other.

To release this energy, hydrogen ions will rush through any opening, similar to water jetting through a hole in a dam. In the thylakoid, that opening is a passage through a specialized protein channel called the ATP synthase. The energy released by the hydrogen ion stream allows ATP synthase to attach a third phosphate group to ADP, which forms a molecule of ATP (Figure 2). The flow of hydrogen ions through ATP synthase is called chemiosmosis because the ions move from an area of high to an area of low concentration through a semi-permeable structure.

Visit this site and click through the animation to view the process of photosynthesis within a leaf.

### Learning Objectives

The pigments of the first part of photosynthesis, the light-dependent reactions, absorb energy from sunlight. A photon strikes the antenna pigments of photosystem II to initiate photosynthesis. The energy travels to the reaction center that contains chlorophyll *a* to the electron transport chain, which pumps hydrogen ions into the thylakoid interior. This action builds up a high concentration of ions. The ions flow through ATP synthase via chemiosmosis to form molecules of ATP, which are used for the formation of sugar molecules in the second stage of photosynthesis. Photosystem I absorbs a second photon, which results in the formation of an NADPH molecule, another energy and reducing power carrier for the light-independent reactions.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.8: Light-Dependent Reactions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.9: Light-Independent Reactions

After the energy from the sun is converted into chemical energy and temporarily stored in ATP and NADPH molecules, the cell has the fuel needed to build carbohydrate molecules for long-term energy storage. The products of the light-dependent reactions, ATP and NADPH, have lifespans in the range of millionths of seconds, whereas the products of the light-independent reactions (carbohydrates and other forms of reduced carbon) can survive for hundreds of millions of years. The carbohydrate molecules made will have a backbone of carbon atoms. Where does the carbon come from? It comes from carbon dioxide, the gas that is a waste product of respiration in microbes, fungi, plants, and animals.

In plants, carbon dioxide ($CO_2$) enters the leaves through stomata, where it diffuses over short distances through intercellular spaces until it reaches the mesophyll cells. Once in the mesophyll cells, $CO_2$ diffuses into the stroma of the chloroplast—the site of light-independent reactions of photosynthesis. These reactions actually have several names associated with them. Another term, the **Calvin cycle**, is named for the man who discovered it, and because these reactions function as a cycle. Others call it the Calvin-Benson cycle to include the name of another scientist involved in its discovery. The most outdated name is dark reactions, because light is not directly required (Figure 1). However, the term dark reaction can be misleading because it implies incorrectly that the reaction only occurs at night or is independent of light, which is why most scientists and instructors no longer use it.





Figure 1. Light reactions harness energy from the sun to produce chemical bonds, ATP, and NADPH. These energy-carrying molecules are made in the stroma where carbon fixation takes place.

The light-independent reactions of the Calvin cycle can be organized into three basic stages: fixation, reduction, and regeneration.

## Stage 1: Fixation

In the stroma, in addition to $CO_2$, two other components are present to initiate the light-independent reactions: an enzyme called ribulose bisphosphate carboxylase (RuBisCO), and three molecules of ribulose bisphosphate (RuBP), as shown in Figure 2. RuBP has five atoms of carbon, flanked by two phosphates.

**Practice Question**





Figure 2. The Calvin cycle has three stages.

In stage 1, the enzyme RuBisCO incorporates carbon dioxide into an organic molecule, 3-PGA. In stage 2, the organic molecule is reduced using electrons supplied by NADPH. In stage 3, RuBP, the molecule that starts the cycle, is regenerated so that the cycle can continue. Only one carbon dioxide molecule is incorporated at a time, so the cycle must be completed three times to produce a single three-carbon GA3P molecule, and six times to produce a six-carbon glucose molecule.

Which of the following statements is true?

a. In photosynthesis, oxygen, carbon dioxide, ATP, and NADPH are reactants. GA3P and water are products.
b. In photosynthesis, chlorophyll, water, and carbon dioxide are reactants. GA3P and oxygen are products.
c. In photosynthesis, water, carbon dioxide, ATP, and NADPH are reactants. RuBP and oxygen are products.
d. In photosynthesis, water and carbon dioxide are reactants. GA3P and oxygen are products.

[reveal-answer q="542141"]**Show Answer**[/reveal-answer]
[hidden-answer a="542141"]Answer d is true.[/hidden-answer]

RuBisCO catalyzes a reaction between $CO_2$ and RuBP. For each $CO_2$ molecule that reacts with one RuBP, two molecules of another compound (3-PGA) form. PGA has three carbons and one phosphate. Each turn of the cycle involves only one RuBP and one carbon dioxide and forms two molecules of 3-PGA. The number of carbon atoms remains the same, as the atoms move to form new bonds during the reactions (3 atoms from $3CO_2$ + 15 atoms from 3RuBP = 18 atoms in 3 atoms of 3-PGA). This process is called **carbon fixation**, because $CO_2$ is "fixed" from an inorganic form into organic molecules.

## Stage 2: Reduction

ATP and NADPH are used to convert the six molecules of 3-PGA into six molecules of a chemical called glyceraldehyde 3-phosphate (G3P). That is a reduction reaction because it involves the gain of electrons by 3-PGA. Recall that a **reduction** is the gain of an electron by an atom or molecule. Six molecules of both ATP and NADPH are used. For ATP, energy is released with the loss of the terminal phosphate atom, converting it into ADP; for NADPH, both energy and a hydrogen atom are lost, converting it into $NADP^+$. Both of these molecules return to the nearby light-dependent reactions to be reused and reenergized.

## Stage 3: Regeneration

Interestingly, at this point, only one of the G3P molecules leaves the Calvin cycle and is sent to the cytoplasm to contribute to the formation of other compounds needed by the plant. Because the G3P exported from the chloroplast has three carbon atoms, it takes three “turns” of the Calvin cycle to fix enough net carbon to export one G3P. But each turn makes two G3Ps, thus three turns make six G3Ps. One is exported while the remaining five G3P molecules remain in the cycle and are used to regenerate RuBP, which enables the system to prepare for more $CO_2$ to be fixed. Three more molecules of ATP are used in these regeneration reactions.

This link leads to an animation of the Calvin cycle. Click stage 1, stage 2, and then stage 3 to see G3P and ATP regenerate to form RuBP.

Try It



Figure 3. The harsh conditions of the desert have led plants like these cacti to evolve variations of the light-independent reactions of photosynthesis. These variations increase the efficiency of water usage, helping to conserve water and energy. (credit: Piotr Wojtkowski)

During the evolution of photosynthesis, a major shift occurred from the bacterial type of photosynthesis that involves only one photosystem and is typically anoxygenic (does not generate oxygen) into modern oxygenic (does generate oxygen) photosynthesis, employing two photosystems. This modern oxygenic photosynthesis is used by many organisms—from giant tropical leaves in the rainforest to tiny cyanobacterial cells—and the process and components of this photosynthesis remain largely the same. Photosystems absorb light and use electron transport chains to convert energy into the chemical energy of ATP and NADH. The subsequent light-independent reactions then assemble carbohydrate molecules with this energy.

Photosynthesis in desert plants has evolved adaptations that conserve water. In the harsh dry heat, every drop of water must be used to survive. Because stomata must open to allow for the uptake of $CO_2$, water escapes from the leaf during active photosynthesis. Desert plants have evolved processes to conserve water and deal with harsh conditions. A more efficient use of $CO_2$ allows plants to adapt to living with less water. Some plants such as cacti (Figure 3) can prepare materials for photosynthesis during the night by a temporary carbon fixation/storage process, because opening the stomata at this time conserves water due to cooler temperatures. In addition, cacti have evolved the ability to carry out low levels of photosynthesis without opening stomata at all, a mechanism to face extremely dry periods.

## Contributors and Attributions


CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.9: Light-Independent Reactions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.10: Photosynthesis

Now that we've learned about the different pieces of photosynthesis, let's put it all together. This video walks you through the process of photosynthesis as a whole:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=256

## Contributors and Attributions

All rights reserved content

- Photosynthesis: Crash Course Biology #8. **Authored by**: CrashCourse. **Located at**: https://youtu.be/sQK3Yr4Sc_k. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

8.10: Photosynthesis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.11: Introduction to Cellular Respiration

## Identify the reactants and products of cellular respiration and where these reactions occur in a cell

Now that we've learned how autotrophs like plants convert sunlight to sugars, let's take a look at how all eukaryotes—which includes humans!—make use of those sugars.

In the process of photosynthesis, plants and other photosynthetic producers create glucose, which stores energy in its chemical bonds. Then, both plants and consumers, such as animals, undergo a series of metabolic pathways—collectively called cellular respiration. Cellular respiration extracts the energy from the bonds in glucose and converts it into a form that all living things can use.

## What You'll Learn to Do

- Describe the process of glycolysis and identify its reactants and products
- Describe the process of pyruvate oxidation and identify its reactants and products
- Describe the process of the citric acid cycle (Krebs cycle) and identify its reactants and products
- Describe the respiratory chain (electron transport chain) and its role in cellular respiration

### Learning Activities

The learning activities for this section include the following:

- Glycolysis
- Pyruvate Oxidation
- Citric Acid Cycle
- Electron Transport Chain
- ATP & Respiration
- Self Check: Cellular Respiration

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Modification of Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.11: Introduction to Cellular Respiration is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.12: Glycolysis

**Glycolysis** is the first step in the breakdown of glucose to extract energy for cellular metabolism. Nearly all living organisms carry out glycolysis as part of their metabolism. The process does not use oxygen and is therefore **anaerobic** (processes that use oxygen are called aerobic). Glycolysis takes place in the cytoplasm of both prokaryotic and eukaryotic cells. Glucose enters heterotrophic cells in two ways.

1. Through secondary active transport in which the transport takes place against the glucose concentration gradient.
2. Through a group of integral proteins called GLUT proteins, also known as glucose transporter proteins. These transporters assist in the facilitated diffusion of glucose.

Glycolysis begins with the six carbon ring-shaped structure of a single glucose molecule and ends with two molecules of a three-carbon sugar called **pyruvate** (Figure 1).





Figure 1. Reactants and products of glycolysis.

Glycolysis consists of ten steps divided into two distinct halves. The first half of the glycolysis is also known as the **energy-requiring steps**. This pathway traps the glucose molecule in the cell and uses energy to modify it so that the six-carbon sugar molecule can be split evenly into the two three-carbon molecules. The second half of glycolysis (also known as the **energy-releasing steps**) extracts energy from the molecules and stores it in the form of ATP and NADH, the reduced form of NAD.

## First Half of Glycolysis (Energy-Requiring Steps)





Figure 2. The first half of glycolysis uses two ATP molecules in the phosphorylation of glucose, which is then split into two three-carbon molecules.

Step 1. The first step in glycolysis is catalyzed by hexokinase, an enzyme with broad specificity that catalyzes the phosphorylation of six-carbon sugars. Hexokinase phosphorylates glucose using ATP as the source of the phosphate, producing glucose-6-phosphate, a more reactive form of glucose. This reaction prevents the phosphorylated glucose molecule from continuing to interact with the GLUT proteins, and it can no longer leave the cell because the negatively charged phosphate will not allow it to cross the hydrophobic interior of the plasma membrane.

Step 2. In the second step of glycolysis, an isomerase converts glucose-6-phosphate into one of its isomers, fructose-6-phosphate. An **isomerase** is an enzyme that catalyzes the conversion of a molecule into one of its isomers. This change from phosphoglucose

to phosphofructose allows the eventual split of the sugar into two three-carbon molecules.

Step 3. The third step is the phosphorylation of fructose-6-phosphate, catalyzed by the enzyme phosphofructokinase. A second ATP molecule donates a high-energy phosphate to fructose-6-phosphate, producing fructose-1,6-bisphosphate. In this pathway, phosphofructokinase is a rate-limiting enzyme. It is active when the concentration of ADP is high; it is less active when ADP levels are low and the concentration of ATP is high. Thus, if there is "sufficient" ATP in the system, the pathway slows down. This is a type of end product inhibition, since ATP is the end product of glucose catabolism.

Step 4. The newly added high-energy phosphates further destabilize fructose-1,6-bisphosphate. The fourth step in glycolysis employs an enzyme, aldolase, to cleave 1,6-bisphosphate into two three-carbon isomers: dihydroxyacetone-phosphate and glyceraldehyde-3-phosphate.

Step 5. In the fifth step, an isomerase transforms the dihydroxyacetone-phosphate into its isomer, glyceraldehyde-3-phosphate. Thus, the pathway will continue with two molecules of a single isomer. At this point in the pathway, there is a net investment of energy from two ATP molecules in the breakdown of one glucose molecule.

## Second Half of Glycolysis (Energy-Releasing Steps)

So far, glycolysis has cost the cell two ATP molecules and produced two small, three-carbon sugar molecules. Both of these molecules will proceed through the second half of the pathway, and sufficient energy will be extracted to pay back the two ATP molecules used as an initial investment and produce a profit for the cell of two additional ATP molecules and two even higher-energy NADH molecules.





Figure 3. The second half of glycolysis involves phosphorylation without ATP investment (step 6) and produces two NADH and four ATP molecules per glucose.

Step 6. The sixth step in glycolysis (Figure 3) oxidizes the sugar (glyceraldehyde-3-phosphate), extracting high-energy electrons, which are picked up by the electron carrier $NAD^+$, producing NADH. The sugar is then phosphorylated by the addition of a second phosphate group, producing 1,3-bisphosphoglycerate. Note that the second phosphate group does not require another ATP molecule.

Here again is a potential limiting factor for this pathway. The continuation of the reaction depends upon the availability of the oxidized form of the electron carrier, $NAD^+$. Thus, NADH must be continuously oxidized back into $NAD^+$ in order to keep this step going. If $NAD^+$ is not available, the second half of glycolysis slows down or stops. If oxygen is available in the system, the NADH will be oxidized readily, though indirectly, and the high-energy electrons from the hydrogen released in this process will be used to produce ATP. In an environment without oxygen, an alternate pathway (fermentation) can provide the oxidation of NADH to $NAD^+$.

Step 7. In the seventh step, catalyzed by phosphoglycerate kinase (an enzyme named for the reverse reaction), 1,3-bisphosphoglycerate donates a high-energy phosphate to ADP, forming one molecule of ATP. (This is an example of substrate-level phosphorylation.) A carbonyl group on the 1,3-bisphosphoglycerate is oxidized to a carboxyl group, and 3-phosphoglycerate is formed.

Step 8. In the eighth step, the remaining phosphate group in 3-phosphoglycerate moves from the third carbon to the second carbon, producing 2-phosphoglycerate (an isomer of 3-phosphoglycerate). The enzyme catalyzing this step is a mutase (a type of isomerase).

Step 9. Enolase catalyzes the ninth step. This enzyme causes 2-phosphoglycerate to lose water from its structure; this is a dehydration reaction, resulting in the formation of a double bond that increases the potential energy in the remaining phosphate bond and produces phosphoenolpyruvate (PEP).

Step 10. The last step in glycolysis is catalyzed by the enzyme pyruvate kinase (the enzyme in this case is named for the reverse reaction of pyruvate's conversion into PEP) and results in the production of a second ATP molecule by substrate-level phosphorylation and the compound pyruvic acid (or its salt form, pyruvate). Many enzymes in enzymatic pathways are named for the reverse reactions, since the enzyme can catalyze both forward and reverse reactions.

## Outcomes of Glycolysis

Glycolysis starts with glucose and ends with two pyruvate molecules, a total of four ATP molecules and two molecules of NADH. Two ATP molecules were used in the first half of the pathway to prepare the six-carbon ring for cleavage, so the cell has a net gain of two ATP molecules and two NADH molecules for its use.

If the cell cannot catabolize the pyruvate molecules further, it will harvest only two ATP molecules from one molecule of glucose. Mature mammalian red blood cells are not capable of **aerobic respiration**—the process in which organisms convert energy in the presence of oxygen—and glycolysis is their sole source of ATP. If glycolysis is interrupted, these cells lose their ability to maintain their sodium-potassium pumps, and eventually, they die.

The last step in glycolysis will not occur if pyruvate kinase, the enzyme that catalyzes the formation of pyruvate, is not available in sufficient quantities. In this situation, the entire glycolysis pathway will proceed, but only two ATP molecules will be made in the second half. Thus, pyruvate kinase is a rate-limiting enzyme for glycolysis.

### Learning Objectives

Glycolysis is the first pathway used in the breakdown of glucose to extract energy. It was probably one of the earliest metabolic pathways to evolve and is used by nearly all of the organisms on earth. Glycolysis consists of two parts: The first part prepares the six-carbon ring of glucose for cleavage into two three-carbon sugars. ATP is invested in the process during this half to energize the separation. The second half of glycolysis extracts ATP and high-energy electrons from hydrogen atoms and attaches them to $NAD^+$. Two ATP molecules are invested in the first half and four ATP molecules are formed by substrate phosphorylation during the second half. This produces a net gain of two ATP and two NADH molecules for the cell.

Figure 4 shows the entire process of glycolysis in one image:





Figure 4. Glycolysis

## Contributors and Attributions

CC licensed content, Original

- Modification and revision of Biology. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Modification of Glycolysis metabolic pathway 3 annotated. **Authored by**: Thomas Shafee. **Located at**: https://commons.wikimedia.org/wiki/File:Glycolysis_metabolic_pathway_3_annotated.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

8.12: Glycolysis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.13: Pyruvate Oxidation

If oxygen is available, aerobic respiration will go forward. In eukaryotic cells, the pyruvate molecules produced at the end of glycolysis are transported into mitochondria, which are the sites of cellular respiration. There, pyruvate will be transformed into an acetyl group that will be picked up and activated by a carrier compound called coenzyme A (CoA). The resulting compound is called acetyl CoA. CoA is made from vitamin B5, pantothenic acid. Acetyl CoA can be used in a variety of ways by the cell, but its major function is to deliver the acetyl group derived from pyruvate to the next stage of the pathway in glucose catabolism.

## Breakdown of Pyruvate

In order for pyruvate (which is the product of glycolysis) to enter the Citric Acid Cycle (the next pathway in cellular respiration), it must undergo several changes. The conversion is a three-step process (Figure 1).





Figure 1. Upon entering the mitochondrial matrix, a multi-enzyme complex converts pyruvate into acetyl CoA. In the process, carbon dioxide is released and one molecule of NADH is formed.

**Step 1. A carboxyl group is removed from pyruvate, releasing a molecule of carbon dioxide into the surrounding medium.** The result of this step is a two-carbon hydroxyethyl group bound to the enzyme (pyruvate dehydrogenase). This is the first of the six carbons from the original glucose molecule to be removed. This step proceeds twice (remember: there are *two* pyruvate molecules produced at the end of glycolysis) for every molecule of glucose metabolized; thus, two of the six carbons will have been removed at the end of both steps.

**Step 2. $NAD^+$ is reduced to NADH.** The hydroxyethyl group is oxidized to an acetyl group, and the electrons are picked up by $NAD^+$, forming NADH. The high-energy electrons from NADH will be used later to generate ATP.

**Step 3. An acetyl group is transferred to conenzyme A, resulting in acetyl CoA.** The enzyme-bound acetyl group is transferred to CoA, producing a molecule of acetyl CoA.

Note that during the second stage of glucose metabolism, whenever a carbon atom is removed, it is bound to two oxygen atoms, producing carbon dioxide, one of the major end products of cellular respiration.

## Acetyl CoA to $CO_2$

In the presence of oxygen, acetyl CoA delivers its acetyl group to a four-carbon molecule, oxaloacetate, to form citrate, a six-carbon molecule with three carboxyl groups; this pathway will harvest the remainder of the extractable energy from what began as a glucose molecule. This single pathway is called by different names, but we will primarily call it the **Citric Acid Cycle**.

### Learning Objectives

In the presence of oxygen, pyruvate is transformed into an acetyl group attached to a carrier molecule of coenzyme A. The resulting acetyl CoA can enter several pathways, but most often, the acetyl group is delivered to the citric acid cycle for further catabolism. During the conversion of pyruvate into the acetyl group, a molecule of carbon dioxide and two high-energy electrons are removed. The carbon dioxide accounts for two (conversion of two pyruvate molecules) of the six carbons of the original glucose molecule. The electrons are picked up by $NAD^+$, and the NADH carries the electrons to a later pathway for ATP production. At this point, the glucose molecule that originally entered cellular respiration has been completely oxidized.

> Chemical potential energy stored within the glucose molecule has been transferred to electron carriers or has been used to synthesize a few ATPs.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.13: Pyruvate Oxidation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.14: Citric Acid Cycle

Like the conversion of pyruvate to acetyl CoA, the citric acid cycle takes place in the matrix of mitochondria.This single pathway is called by different names: the citric acid cycle (for the first intermediate formed—citric acid, or citrate—when acetate joins to the oxaloacetate), the TCA cycle (since citric acid or citrate and isocitrate are tricarboxylic acids), and the Krebs cycle, after Hans Krebs, who first identified the steps in the pathway in the 1930s in pigeon flight muscles.

Almost all of the enzymes of the citric acid cycle are soluble, with the single exception of the enzyme succinate dehydrogenase, which is embedded in the inner membrane of the mitochondrion. Unlike glycolysis, the citric acid cycle is a closed loop: The last part of the pathway regenerates the compound used in the first step. The eight steps of the cycle are a series of redox, dehydration, hydration, and decarboxylation reactions that produce two carbon dioxide molecules, one GTP/ATP, and reduced forms of NADH and $FADH_2$ (Figure 1). This is considered an aerobic pathway because the NADH and $FADH_2$ produced must transfer their electrons to the next pathway in the system, which will use oxygen. If this transfer does not occur, the oxidation steps of the citric acid cycle also do not occur. Note that the citric acid cycle produces very little ATP directly and does not directly consume oxygen.





Figure 1. In the citric acid cycle, the acetyl group from acetyl CoA is attached to a four-carbon oxaloacetate molecule to form a six-carbon citrate molecule. Through a series of steps, citrate is oxidized, releasing two carbon dioxide molecules for each acetyl group fed into the cycle. In the process, three NAD+ molecules are reduced to NADH, one FAD molecule is reduced to FADH2, and one ATP or GTP (depending on the cell type) is produced (by substrate-level phosphorylation). Because the final product of the citric acid cycle is also the first reactant, the cycle runs continuously in the presence of sufficient reactants. (credit: modification of work by “Yikrazuul”/Wikimedia Commons)

## Steps in the Citric Acid Cycle

Step 1. Prior to the start of the first step, pyruvate oxidation must occur. Then, the first step of the cycle begins: This is a condensation step, combining the two-carbon acetyl group with a four-carbon oxaloacetate molecule to form a six-carbon molecule of **citrate**. CoA is bound to a sulfhydryl group (-SH) and diffuses away to eventually combine with another acetyl group. This step is irreversible because it is highly exergonic. The rate of this reaction is controlled by negative feedback and the amount of ATP available. If ATP levels increase, the rate of this reaction decreases. If ATP is in short supply, the rate increases.

Step 2. In step two, citrate loses one water molecule and gains another as citrate is converted into its isomer, **isocitrate**.

Step 3. In step three, isocitrate is oxidized, producing a five-carbon molecule, **α-ketoglutarate**, together with a molecule of $CO_2$ and two electrons, which reduce $NAD^+$ to NADH. This step is also regulated by negative feedback from ATP and NADH, and a positive effect of ADP.

Steps 3 and 4. Steps three and four are both oxidation and decarboxylation steps, which release electrons that reduce $NAD^+$ to NADH and release carboxyl groups that form $CO_2$ molecules. α-Ketoglutarate is the product of step three, and a **succinyl** group is the product of step four. CoA binds the succinyl group to form succinyl CoA. The enzyme that catalyzes step four is regulated by feedback inhibition of ATP, succinyl CoA, and NADH.

Step 5. In step five, a phosphate group is substituted for coenzyme A, and a high-energy bond is formed. This energy is used in substrate-level phosphorylation (during the conversion of the succinyl group to succinate) to form either guanine triphosphate (GTP) or ATP. There are two forms of the enzyme, called isoenzymes, for this step, depending upon the type of animal tissue in which they are found. One form is found in tissues that use large amounts of ATP, such as heart and skeletal muscle. This form produces ATP. The second form of the enzyme is found in tissues that have a high number of anabolic pathways, such as liver. This form produces GTP. GTP is energetically equivalent to ATP; however, its use is more restricted. In particular, protein synthesis primarily uses GTP.

Step 6. Step six is a dehydration process that converts **succinate** into **fumarate**. Two hydrogen atoms are transferred to FAD, producing $FADH_2$. The energy contained in the electrons of these atoms is insufficient to reduce $NAD^+$ but adequate to reduce FAD. Unlike NADH, this carrier remains attached to the enzyme and transfers the electrons to the electron transport chain directly. This process is made possible by the localization of the enzyme catalyzing this step inside the inner membrane of the mitochondrion.

Step 7. Water is added to fumarate during step seven, and **malate** is produced. The last step in the citric acid cycle regenerates **oxaloacetate** by oxidizing malate. Another molecule of NADH is produced in the process.

*You can click through each step of the citric acid cycle here.*

## Products of the Citric Acid Cycle

Two carbon atoms come into the citric acid cycle from each acetyl group, representing four out of the six carbons of one glucose molecule. Two carbon dioxide molecules are released on each turn of the cycle; however, these do not necessarily contain the most recently added carbon atoms. The two acetyl carbon atoms will eventually be released on later turns of the cycle; thus, all six carbon atoms from the original glucose molecule are eventually incorporated into carbon dioxide. Each turn of the cycle forms three NADH molecules and one $FADH_2$ molecule. These carriers will connect with the last portion of aerobic respiration to produce ATP molecules. One GTP or ATP is also made in each cycle. Several of the intermediate compounds in the citric acid cycle can be used in synthesizing non-essential amino acids; therefore, the cycle is amphibolic (both catabolic and anabolic).

### Learning Objectives

The citric acid cycle is a series of redox and decarboxylation reactions that remove high-energy electrons and carbon dioxide. The electrons temporarily stored in molecules of NADH and $FADH_2$ are used to generate ATP in a subsequent pathway. One molecule of either GTP or ATP is produced by substrate-level phosphorylation on each turn of the cycle. There is no comparison of the cyclic pathway with a linear one.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

8.14: Citric Acid Cycle is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.15: Electron Transport Chain

You have just read about two pathways in cellular respiration—glycolysis and the citric acid cycle—that generate ATP. However, most of the ATP generated during the aerobic catabolism of glucose is not generated directly from these pathways. Rather, it is derived from a process that begins with moving electrons through a series of electron transporters that undergo redox reactions: **the electron transport chain**. This causes hydrogen ions to accumulate within the matrix space. Therefore, a concentration gradient forms in which hydrogen ions diffuse out of the matrix space by passing through ATP synthase. The current of hydrogen ions powers the catalytic action of ATP synthase, which phosphorylates ADP, producing ATP.





Figure 1. The electron transport chain is a series of electron transporters embedded in the inner mitochondrial membrane that shuttles electrons from NADH and $FADH_2$ to molecular oxygen. In the process, protons are pumped from the mitochondrial matrix to the intermembrane space, and oxygen is reduced to form water.

The electron transport chain (Figure 1) is the last component of aerobic respiration and is the only part of glucose metabolism that uses atmospheric oxygen. Oxygen continuously diffuses into plants; in animals, it enters the body through the respiratory system. Electron transport is a series of redox reactions that resemble a relay race or bucket brigade in that electrons are passed rapidly from one component to the next, to the endpoint of the chain where the electrons reduce molecular oxygen, producing water. There are four complexes composed of proteins, labeled I through IV in Figure 1, and the aggregation of these four complexes, together with associated mobile, accessory electron carriers, is called the electron transport chain. The electron transport chain is present in multiple copies in the inner mitochondrial membrane of eukaryotes and the plasma membrane of prokaryotes. Note, however, that the electron transport chain of prokaryotes may not require oxygen as some live in anaerobic conditions. The common feature of all electron transport chains is the presence of a proton pump to create a proton gradient across a membrane.

## Complex I

To start, two electrons are carried to the first complex aboard NADH. This complex, labeled I, is composed of flavin mononucleotide (FMN) and an iron-sulfur (Fe-S)-containing protein. FMN, which is derived from vitamin $B_2$, also called riboflavin, is one of several prosthetic groups or co-factors in the electron transport chain. A **prosthetic group** is a non-protein molecule required for the activity of a protein. Prosthetic groups are organic or inorganic, non-peptide molecules bound to a protein that facilitate its function; prosthetic groups include co-enzymes, which are the prosthetic groups of enzymes. The enzyme in complex I is NADH dehydrogenase and is a very large protein, containing 45 amino acid chains. Complex I can pump four hydrogen ions across the membrane from the matrix into the intermembrane space, and it is in this way that the hydrogen ion gradient is established and maintained between the two compartments separated by the inner mitochondrial membrane.

## Q and Complex II

Complex II directly receives $FADH_2$, which does not pass through complex I. The compound connecting the first and second complexes to the third is **ubiquinone** (Q). The Q molecule is lipid soluble and freely moves through the hydrophobic core of the membrane. Once it is reduced, ($QH_2$), ubiquinone delivers its electrons to the next complex in the electron transport chain. Q receives the electrons derived from NADH from complex I and the electrons derived from $FADH_2$ from complex II, including

succinate dehydrogenase. This enzyme and $FADH_2$ form a small complex that delivers electrons directly to the electron transport chain, bypassing the first complex. Since these electrons bypass and thus do not energize the proton pump in the first complex, fewer ATP molecules are made from the $FADH_2$ electrons. The number of ATP molecules ultimately obtained is directly proportional to the number of protons pumped across the inner mitochondrial membrane.

## Complex III

The third complex is composed of cytochrome b, another Fe-S protein, Rieske center (2Fe-2S center), and cytochrome c proteins; this complex is also called cytochrome oxidoreductase. Cytochrome proteins have a prosthetic group of heme. The heme molecule is similar to the heme in hemoglobin, but it carries electrons, not oxygen. As a result, the iron ion at its core is reduced and oxidized as it passes the electrons, fluctuating between different oxidation states: $Fe^{++}$ (reduced) and $Fe^{+++}$ (oxidized). The heme molecules in the cytochromes have slightly different characteristics due to the effects of the different proteins binding them, giving slightly different characteristics to each complex. Complex III pumps protons through the membrane and passes its electrons to cytochrome c for transport to the fourth complex of proteins and enzymes (cytochrome c is the acceptor of electrons from Q; however, whereas Q carries pairs of electrons, cytochrome c can accept only one at a time).

## Complex IV

The fourth complex is composed of cytochrome proteins c, a, and $a_3$. This complex contains two heme groups (one in each of the two cytochromes, a, and $a_3$) and three copper ions (a pair of $Cu_A$ and one $Cu_B$ in cytochrome $a_3$). The cytochromes hold an oxygen molecule very tightly between the iron and copper ions until the oxygen is completely reduced. The reduced oxygen then picks up two hydrogen ions from the surrounding medium to make water ($H_2O$). The removal of the hydrogen ions from the system contributes to the ion gradient used in the process of chemiosmosis.

## Chemiosmosis

In chemiosmosis, the free energy from the series of redox reactions just described is used to pump hydrogen ions (protons) across the membrane. The uneven distribution of $H^+$ ions across the membrane establishes both concentration and electrical gradients (thus, an electrochemical gradient), owing to the hydrogen ions' positive charge and their aggregation on one side of the membrane.

If the membrane were open to diffusion by the hydrogen ions, the ions would tend to diffuse back across into the matrix, driven by their electrochemical gradient. Recall that many ions cannot diffuse through the nonpolar regions of phospholipid membranes without the aid of ion channels. Similarly, hydrogen ions in the matrix space can only pass through the inner mitochondrial membrane through an integral membrane protein called ATP synthase (Figure 2). This complex protein acts as a tiny generator, turned by the force of the hydrogen ions diffusing through it, down their electrochemical gradient. The turning of parts of this molecular machine facilitates the addition of a phosphate to ADP, forming ATP, using the potential energy of the hydrogen ion gradient.

**Practice Question**





Figure 2. ATP synthase is a complex, molecular machine that uses a proton (H+) gradient to form ATP from ADP and inorganic phosphate (Pi). (Credit: modification of work by Klaus Hoffmeier)

Dinitrophenol (DNP) is an uncoupler that makes the inner mitochondrial membrane leaky to protons. It was used until 1938 as a weight-loss drug. What effect would you expect DNP to have on the change in pH across the inner mitochondrial membrane? Why do you think this might be an effective weight-loss drug?

[practice-area rows="2"][/practice-area]
[reveal-answer q="346232"]**Show Answer**[/reveal-answer]
[hidden-answer a="346232"]After DNP poisoning, the electron transport chain can no longer form a proton gradient, and ATP synthase can no longer make ATP. DNP is an effective diet drug because it uncouples ATP synthesis; in other words, after taking it, a person obtains less energy out of the food he or she eats. Interestingly, one of the worst side effects of this drug is hyperthermia, or overheating of the body. Since ATP cannot be formed, the energy from electron transport is lost as heat.[/hidden-answer]

Chemiosmosis (Figure 3) is used to generate 90 percent of the ATP made during aerobic glucose catabolism; it is also the method used in the light reactions of photosynthesis to harness the energy of sunlight in the process of photophosphorylation. Recall that the production of ATP using the process of chemiosmosis in mitochondria is called oxidative phosphorylation. The overall result of these reactions is the production of ATP from the energy of the electrons removed from hydrogen atoms. These atoms were originally part of a glucose molecule. At the end of the pathway, the electrons are used to reduce an oxygen molecule to oxygen ions. The extra electrons on the oxygen attract hydrogen ions (protons) from the surrounding medium, and water is formed.

Practice Question





Figure 3. In oxidative phosphorylation, the pH gradient formed by the electron transport chain is used by ATP synthase to form ATP.

Cyanide inhibits cytochrome c oxidase, a component of the electron transport chain. If cyanide poisoning occurs, would you expect the pH of the intermembrane space to increase or decrease? What effect would cyanide have on ATP synthesis?

[practice-area rows="2"][/practice-area]
[reveal-answer q="26135"]**Show Answer**[/reveal-answer]
[hidden-answer a="26135"]After cyanide poisoning, the electron transport chain can no longer pump electrons into the intermembrane space. The pH of the intermembrane space would increase, the pH gradient would decrease, and ATP synthesis would stop.[/hidden-answer]

## ATP Yield

The number of ATP molecules generated from the catabolism of glucose varies. For example, the number of hydrogen ions that the electron transport chain complexes can pump through the membrane varies between species. Another source of variance stems from the shuttle of electrons across the membranes of the mitochondria. (The NADH generated from glycolysis cannot easily enter mitochondria.) Thus, electrons are picked up on the inside of mitochondria by either $NAD^+$ or $FAD^+$. As you have learned earlier, these $FAD^+$ molecules can transport fewer ions; consequently, fewer ATP molecules are generated when $FAD^+$ acts as a carrier. $NAD^+$ is used as the electron transporter in the liver and $FAD^+$ acts in the brain.

Another factor that affects the yield of ATP molecules generated from glucose is the fact that intermediate compounds in these pathways are used for other purposes. Glucose catabolism connects with the pathways that build or break down all other biochemical compounds in cells, and the result is somewhat messier than the ideal situations described thus far. For example, sugars other than glucose are fed into the glycolytic pathway for energy extraction. Moreover, the five-carbon sugars that form nucleic acids are made from intermediates in glycolysis. Certain nonessential amino acids can be made from intermediates of both glycolysis and the citric acid cycle. Lipids, such as cholesterol and triglycerides, are also made from intermediates in these pathways, and both amino acids and triglycerides are broken down for energy through these pathways. Overall, in living systems, these pathways of glucose catabolism extract about 34 percent of the energy contained in glucose.

### Learning Objectives

The electron transport chain is the portion of aerobic respiration that uses free oxygen as the final electron acceptor of the electrons removed from the intermediate compounds in glucose catabolism. The electron transport chain is composed of four large, multiprotein complexes embedded in the inner mitochondrial membrane and two small diffusible electron carriers shuttling electrons between them. The electrons are passed through a series of redox reactions, with a small amount of free

energy used at three points to transport hydrogen ions across a membrane. This process contributes to the gradient used in chemiosmosis. The electrons passing through the electron transport chain gradually lose energy, High-energy electrons donated to the chain by either NADH or $FADH_2$ complete the chain, as low-energy electrons reduce oxygen molecules and form water. The level of free energy of the electrons drops from about 60 kcal/mol in NADH or 45 kcal/mol in $FADH_2$ to about 0 kcal/mol in water. The end products of the electron transport chain are water and ATP. A number of intermediate compounds of the citric acid cycle can be diverted into the anabolism of other biochemical molecules, such as nonessential amino acids, sugars, and lipids. These same molecules can serve as energy sources for the glucose pathways.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.15: Electron Transport Chain is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.16: Cellular Respiration

Cellular respiration is a process that all living things use to convert glucose into energy. Autotrophs (like plants) produce glucose during photosynthesis. Heterotrophs (like humans) ingest other living things to obtain glucose. While the process can seem complex, this page takes you through the key elements of each part of cellular respiration.

Cellular respiration is a collection of three unique metabolic pathways: glycolysis, the citric acid cycle, and the electron transport chain. Glycolysis is an anaerobic process, while the other two pathways are aerobic. In order to move from glycolysis to the citric acid cycle, pyruvate molecules (the output of glycolysis) must be oxidized in a process called pyruvate oxidation.

## Glycolysis

Glycolysis is the first pathway in cellular respiration. This pathway is anaerobic and takes place in the cytoplasm of the cell. This pathway breaks down 1 glucose molecule and produces 2 pyruvate molecules. There are two halves of glycolysis, with five steps in each half. The first half is known as the “energy requiring” steps. This half splits glucose, and uses up 2 ATP. If the concentration of pyruvate kinase is high enough, the second half of glycolysis can proceed. In the second half, the “energy releasing: steps, 4 molecules of ATP and 2 NADH are released. Glycolysis has a **net gain** of 2 ATP molecules and 2 NADH.

Some cells (e.g., mature mammalian red blood cells) cannot undergo aerobic respiration, so glycolysis is their **only** source of ATP. However, most cells undergo pyruvate oxidation and continue to the other pathways of cellular respiration.

## Pyruvate Oxidation

In eukaryotes, pyruvate oxidation takes place in the mitochondria. Pyruvate oxidation can only happen if oxygen is available. In this process, the pyruvate created by glycolysis is oxidized. In this oxidation process, a carboxyl group is removed from pyruvate, creating acetyl groups, which compound with coenzyme A (CoA) to form acetyl CoA. This process also releases $CO_2$.

## Citric Acid Cycle

The citric acid cycle (also known as the Krebs cycle) is the second pathway in cellular respiration, and it also takes place in the mitochondria. The rate of the cycle is controlled by ATP concentration. When there is more ATP available, the rate slows down; when there is less ATP the rate increases. This pathway is a closed loop: the final step produces the compound needed for the first step.

The citric acid cycle is considered an aerobic pathway because the NADH and $FADH_2$ it produces act as temporary electron storage compounds, transferring their electrons to the next pathway (electron transport chain), which uses atmospheric oxygen. Each turn of the citric acid cycle provides a **net gain** of $CO_2$, 1 GTP or ATP, and 3 NADH and 1 $FADH_2$.

## Electron Transport Chain

Most ATP from glucose is generated in the electron transport chain. It is the only part of cellular respiration that directly consumes oxygen; however, in some prokaryotes, this is an anaerobic pathway. In eukaryotes, this pathway takes place in the inner mitochondrial membrane. In prokaryotes it occurs in the plasma membrane.

The electron transport chain is made up of 4 proteins along the membrane and a proton pump. A cofactor shuttles electrons between proteins I–III. If NAD is depleted, skip I: $FADH_2$ starts on II. In chemiosmosis, a proton pump takes hydrogens from inside mitochondria to the outside; this spins the “motor” and the phosphate groups attach to that. The movement changes from ADP to ATP, creating 90% of ATP obtained from aerobic glucose catabolism.

## Let’s Practice

Now that you’ve reviewed cellular respiration, this practice activity will help you see how well you know cellular respiration:

A link to an interactive elements can be found at the bottom of this page.

Click here for a text-only version of the activity.

## Contributors and Attributions

CC licensed content, Original

- Cellular Respiration. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Understanding Cellular Respiration. **Provided by**: Lumen Learning. **Located at**: www.oppia.org/explore/LG5n93fp89oh. **License**: *CC BY-SA: Attribution-ShareAlike*

---

8.16: Cellular Respiration is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.17: Introduction to Fermentation

## Illustrate the basic components and steps of fermentation.

The final metabolic pathway we'll discuss is fermentation. This is an anaerobic process (it occurs without oxygen).

## What You'll Learn to Do

- Describe fermentation as anaerobic energy production.
- Identify the process, products, and reactants of lactic acid fermentation.
- Identify the process, products, and reactants of alcohol fermentation.

### Learning Activities

The learning activities for this section include the following:

- Metabolism without Oxygen
- Types of Fermentation
- Self Check: Fermentation

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

8.17: Introduction to Fermentation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.18: Metabolism without Oxygen




Figure 1. The green color seen in these coastal waters is from an eruption of hydrogen sulfide-producing bacteria. These anaerobic, sulfate-reducing bacteria release hydrogen sulfide gas as they decompose algae in the water. (credit: modification of work by NASA/Jeff Schmaltz, MODIS Land Rapid Response Team at NASA GSFC, Visible Earth Catalog of NASA images)

In aerobic respiration, the final electron acceptor is an oxygen molecule, $O_2$. If aerobic respiration occurs, then ATP will be produced using the energy of high-energy electrons carried by NADH or $FADH_2$ to the electron transport chain. If aerobic respiration does not occur, NADH must be reoxidized to $NAD^+$ for reuse as an electron carrier for the glycolytic pathway to continue. How is this done? Some living systems use an organic molecule as the final electron acceptor. Processes that use an organic molecule to regenerate $NAD^+$ from NADH are collectively referred to as **fermentation**. In contrast, some living systems use an inorganic molecule as a final electron acceptor. Both methods are called **anaerobic cellular respiration** in which organisms convert energy for their use in the absence of oxygen.

## Anaerobic Cellular Respiration

Certain prokaryotes, including some species of bacteria and Archaea, use anaerobic respiration. For example, the group of Archaea called methanogens reduces carbon dioxide to methane to oxidize NADH. These microorganisms are found in soil and in the digestive tracts of ruminants, such as cows and sheep. Similarly, sulfate-reducing bacteria and Archaea, most of which are anaerobic (Figure 1), reduce sulfate to hydrogen sulfide to regenerate $NAD^+$ from NADH.

Visit this site to see anaerobic cellular respiration in action.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

8.18: Metabolism without Oxygen is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.19: Types of Fermentation

## Lactic Acid Fermentation

The fermentation method used by animals and certain bacteria, like those in yogurt, is **lactic acid fermentation** (Figure 1). This type of fermentation is used routinely in mammalian red blood cells and in skeletal muscle that has an insufficient oxygen supply to allow aerobic respiration to continue (that is, in muscles used to the point of fatigue). In muscles, lactic acid accumulation must be removed by the blood circulation and the lactate brought to the liver for further metabolism. The chemical reactions of lactic acid fermentation are the following:




The enzyme used in this reaction is lactate dehydrogenase (LDH). The reaction can proceed in either direction, but the reaction from left to right is inhibited by acidic conditions. Such lactic acid accumulation was once believed to cause muscle stiffness, fatigue, and soreness, although more recent research disputes this hypothesis. Once the lactic acid has been removed from the muscle and circulated to the liver, it can be reconverted into pyruvic acid and further catabolized for energy.

Practice Question





Figure 1. Lactic acid fermentation is common in muscle cells that have run out of oxygen.

Tremetol, a metabolic poison found in the white snake root plant, prevents the metabolism of lactate. When cows eat this plant, it is concentrated in the milk they produce. Humans who consume the milk become ill. Symptoms of this disease, which include vomiting, abdominal pain, and tremors, become worse after exercise. Why do you think this is the case?

[practice-area rows="2"][/practice-area]
[reveal-answer q="453242"]**Show Answer**[/reveal-answer]
[hidden-answer a="453242"]The illness is caused by lactate accumulation. Lactate levels rise after exercise, making the symptoms worse. Milk sickness is rare today, but was common in the Midwestern United States in the early 1800s.[/hidden-answer]



Figure 2. Fermentation of grape juice into wine produces $CO_2$ as a byproduct. Fermentation tanks have valves so that the pressure inside the tanks created by the carbon dioxide produced can be released.

Another familiar fermentation process is **alcohol fermentation** (Figure 3) that produces ethanol, an alcohol (because of this, this kind of fermentation is also sometimes known as **ethanol fermentation**). There are two main reactions in alcohol fermentation.

The first reaction is catalyzed by pyruvate decarboxylase, a cytoplasmic enzyme, with a coenzyme of thiamine pyrophosphate (TPP, derived from vitamin B1 and also called thiamine). A carboxyl group is removed from pyruvic acid, releasing carbon dioxide as a gas. The loss of carbon dioxide reduces the size of the molecule by one carbon, making acetaldehyde. The second reaction is catalyzed by alcohol dehydrogenase to oxidize NADH to $NAD^+$ and reduce acetaldehyde to ethanol. The fermentation of pyruvic acid by yeast produces the ethanol found in alcoholic beverages. Ethanol tolerance of yeast is variable, ranging from about 5 percent to 21 percent, depending on the yeast strain and environmental conditions.





Figure 3. Diagram of alcohol fermentation

## Other Types of Fermentation

Other fermentation methods occur in bacteria. Many prokaryotes are facultatively anaerobic. This means that they can switch between aerobic respiration and fermentation, depending on the availability of oxygen. Certain prokaryotes, like *Clostridia*, are obligate anaerobes. Obligate anaerobes live and grow in the absence of molecular oxygen. Oxygen is a poison to these microorganisms and kills them on exposure.

It should be noted that all forms of fermentation, except lactic acid fermentation, produce gas. The production of particular types of gas is used as an indicator of the fermentation of specific carbohydrates, which plays a role in the laboratory identification of the bacteria. Various methods of fermentation are used by assorted organisms to ensure an adequate supply of $NAD^+$ for the sixth step in glycolysis. Without these pathways, that step would not occur and no ATP would be harvested from the breakdown of glucose.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Ethanol fermentation. **Authored by**: Davidcarmack. **Located at**: https://en.Wikipedia.org/wiki/File:Ethanol_fermentation-1.svg. **License**: *CC BY-SA: Attribution-ShareAlike*

---

8.19: Types of Fermentation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.20: Putting It Together- Metabolic Pathways

Whether the organism is a bacterium, plant, or animal, all living things access energy by breaking down carbohydrate molecules. But if plants make carbohydrate molecules, why would they need to break them down, especially when it has been shown that the gas organisms release as a "waste product" ($CO_2$) acts as a substrate for the formation of more food in photosynthesis? Remember, living things need energy to perform life functions. In addition, an organism can either make its own food or eat another organism—either way, the food still needs to be converted to a form cells can actually use. Finally, in that process of conversion, called cellular respiration, organisms release needed energy and produce "waste" in the form of $CO_2$ gas.






Figure 1. Photosynthesis consumes carbon dioxide and produces oxygen. Aerobic respiration consumes oxygen and produces carbon dioxide. These two processes play an important role in the carbon cycle. (credit: modification of work by Stuart Bassil)

In nature, there is no such thing as waste. Every single atom of matter and energy is conserved, recycling over and over infinitely. Substances change form or move from one type of molecule to another, but their constituent atoms never disappear (Figure 1).

While you may be tempted to call $CO_2$ a waste product, you should remember that oxygen is a "waste product" of photosynthesis: $CO_2$ and oxygen are byproducts of reactions that move on to other reactions. Photosynthesis absorbs light energy to build carbohydrates in chloroplasts, and aerobic cellular respiration releases energy by using oxygen to metabolize carbohydrates in the cytoplasm and mitochondria. Both processes use electron transport chains to capture the energy necessary to drive other reactions. These two powerhouse processes, photosynthesis and cellular respiration, function in biological, cyclical harmony to allow organisms to access life-sustaining energy that originates millions of miles away in a burning star humans call the sun.

## Biofuels

Obviously its important for providing energy for living organisms to power themselves. But is that the only power that photosynthesis provides? What about biofuels? Watch this 14 minute video for an amazing discussion of a proposed biofuel source that doesn't use arable land, doesn't take away food crops, and utilizes wastewater from cities.



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=276

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Jonathan Trent: Energy from floating algae pods. **Authored by**: TED. **Located at**: https://youtu.be/X-HE4Hfa-OY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

8.20: Putting It Together- Metabolic Pathways is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 8.21: Assignment- Observing Energy Transactions

For this assessment you will conduct two kitchen science experiments that each focus on energy processes in living organisms. You will complete the experiments separately but then discuss how the two processes are related.

- **Part 1: Photosynthesis.** Use the Photosynthesis lab procedure to conduct your experiment. Your results (table, graph, photographs) should be compiled as "Part 1 data." Note this experiment will take you around 2 hours to complete.
- **Part 2: Cellular respiration.** Use the Cell Respiration lab procedure to conduct your experiment. Your results (table, graph, photographs) should be compiled as "Part 2 data." Note this experiment will take you around 1 hour to complete.
- **Part 3: Comparison.** After you have compiled both sets of data, answer all the questions in the Part 3 comparison.

Submit all three parts as a single document. Be sure to add clear headings separating the three parts. Here is an example of what you might see or how you may photograph your experiments.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references
- Assignment includes table, bar graph, and photographs associated with both Part 1 and 2 experiments.

## Rubric

| Observing Energy Transactions | | | | |
|---|---|---|---|---|
| Outcome: Explain the metabolic pathways involved in the capture and release of energy in cells | | | | |
| Criteria | Ratings | | | Pts |
| Demonstrate an understanding of photosynthesis through the completion of Part 1 experiment and Part 3 questions. | Part 1 experiment completed and well documented and Part 3 questions answered accurately in detail. 5.0 pts | Part 1 experiment completed and minimally documented and Part 3 questions answered accurately but with minimal detail. 4.0 pts | Part 1 experiment not completed fully and/or Part 3 questions answered inaccurately 0.0 pts | 5 pts |
| A basic understanding of the steps of cellular respiration is illustrated through the completion of Part 2 experiment and Part 3 questions. | Part 2 experiment completed and well documented and Part 3 questions answered accurately in detail. 5.0 pts | Part 2 experiment completed and minimally documented and Part 3 questions answered accurately but with minimal detail. 4.0 pts | Part 2 experiment not completed fully and/or Part 3 questions answered inaccurately 0.0 pts | 5 pts |
| | | | | Total points: 10 |

## Contributors and Attributions

CC licensed content, Original

- Performance Assessment: Observing Energy Transactions. **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

8.21: Assignment- Observing Energy Transactions is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 9: Module 7- Cell Communication


9.1: Why It Matters- Cell Communication
9.2: Introduction to Signaling Molecules and Cellular Receptors
9.3: Types of Signals
9.4: Signaling Molecules
9.5: Signaling Receptors
9.6: Introduction to Propagation of the Signal
9.7: Binding Initiates a Signaling Pathway
9.8: Methods of Intracellular Signaling
9.9: Introduction to Response to the Signal
9.10: Expression and Growth
9.11: Cell Death
9.12: Introduction to Signaling in Single-Celled Organisms
9.13: Signaling in Yeast
9.14: Signaling in Bacteria
9.15: Putting It Together- Cell Communication


---


9: Module 7- Cell Communication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.1: Why It Matters- Cell Communication

## Why identify the main components of a signal transduction pathway?

 Imagine what life would be like if you and the people around you could not communicate. You would not be able to express your wishes to others, nor could you ask questions to find out more about your environment. Social organization is dependent on communication between the individuals that comprise that society; without communication, society would fall apart.

As with people, it is vital for individual cells to be able to interact with their environment. This is true whether a cell is growing by itself in a pond or is one of many cells that form a larger organism. In order to properly respond to external stimuli, cells have developed complex mechanisms of communication that can receive a message, transfer the information across the plasma membrane, and then produce changes within the cell in response to the message.

In multicellular organisms, cells send and receive chemical messages constantly to coordinate the actions of distant organs, tissues, and cells. The ability to send messages quickly and efficiently enables cells to coordinate and fine-tune their functions.

While the necessity for cellular communication in larger organisms seems obvious, even single-celled organisms communicate with each other. Yeast cells signal each other to aid mating. Some forms of bacteria coordinate their actions in order to form large complexes called biofilms or to organize the production of toxins to remove competing organisms. The ability of cells to communicate through chemical signals originated in single cells and was essential for the evolution of multicellular organisms. The efficient and error-free function of communication systems is vital for all life as we know it.

### Learning Outcomes

- Differentiate between different types of signals
- Describe how a cell propagates a signal
- Describe how a cell responds to a signal
- Discuss the process of signaling in single-celled organisms

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Communication. **Authored by**: Lorenzo Stella. **Located at**: https://thenounproject.com/term/communication/27324/. **License**: *CC BY: Attribution*

9.1: Why It Matters- Cell Communication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.2: Introduction to Signaling Molecules and Cellular Receptors

## Differentiate between different types of signals

There are two kinds of communication in the world of living cells. Communication between cells is called **intercellular signaling**, and communication within a cell is called **intracellular signaling**. An easy way to remember the distinction is by understanding the Latin origin of the prefixes: *inter–* means “between” (for example, intersecting lines are those that cross each other) and *intra–* means “inside” (like intravenous).

Chemical signals are released by **signaling cells** in the form of small, usually volatile or soluble molecules called ligands. A **ligand** is a molecule that binds another specific molecule, in some cases, delivering a signal in the process. Ligands can thus be thought of as signaling molecules. Ligands interact with proteins in **target cells**, which are cells that are affected by chemical signals; these proteins are also called **receptors**. Ligands and receptors exist in several varieties; however, a specific ligand will have a specific receptor that typically binds only that ligand.

## What You’ll Learn to Do

- Differentiate between different types of signals used by multicellular organisms
- Identify different types of signaling molecules
- Identify types of receptors, their molecular composition, and the differences among them

### Learning Activities

The learning activities for this section include the following:

- Types of Signals
- Signaling Molecules
- Signaling Receptors
- Self Check: Signaling Molecules and Cellular Receptors

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.2: Introduction to Signaling Molecules and Cellular Receptors is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.3: Types of Signals

There are four categories of chemical signaling found in multicellular organisms: paracrine signaling, endocrine signaling, autocrine signaling, and direct signaling across gap junctions (Figure 1). The main difference between the different categories of signaling is the distance that the signal travels through the organism to reach the target cell. Not all cells are affected by the same signals.





Figure 1. In chemical signaling, a cell may target itself (autocrine signaling), a cell connected by gap junctions, a nearby cell (paracrine signaling), or a distant cell (endocrine signaling). Paracrine signaling acts on nearby cells, endocrine signaling uses the circulatory system to transport ligands, and autocrine signaling acts on the signaling cell. Signaling via gap junctions involves signaling molecules moving directly between adjacent cells.

## Paracrine Signaling





Figure 2. The distance between the presynaptic cell and the postsynaptic cell—called the synaptic gap—is very small and allows for rapid diffusion of the neurotransmitter. Enzymes in the synaptic cleft degrade some types of neurotransmitters to terminate the signal.

Signals that act locally between cells that are close together are called **paracrine signals**. Paracrine signals move by diffusion through the extracellular matrix. These types of signals usually elicit quick responses that last only a short amount of time. In order to keep the response localized, paracrine ligand molecules are normally quickly degraded by enzymes or removed by neighboring cells. Removing the signals will reestablish the concentration gradient for the signal, allowing them to quickly diffuse through the intracellular space if released again.

One example of paracrine signaling is the transfer of signals across synapses between nerve cells. A nerve cell consists of a cell body, several short, branched extensions called dendrites that receive stimuli, and a long extension called an axon, which transmits signals to other nerve cells or muscle cells. The junction between nerve cells where signal transmission occurs is called a synapse. A **synaptic signal** is a chemical signal that travels between nerve cells. Signals within the nerve cells are propagated by fast-moving electrical impulses. When these impulses reach the end of the axon, the signal continues on to a dendrite of the next cell by the release of chemical ligands called **neurotransmitters** by the presynaptic cell (the cell emitting the signal). The neurotransmitters are transported across the very small distances between nerve cells, which are called **chemical synapses** (Figure 2). The small distance between nerve cells allows the signal to travel quickly; this enables an immediate response, such as, Take your hand off the stove!

When the neurotransmitter binds the receptor on the surface of the postsynaptic cell, the electrochemical potential of the target cell changes, and the next electrical impulse is launched. The neurotransmitters that are released into the chemical synapse are degraded quickly or get reabsorbed by the presynaptic cell so that the recipient nerve cell can recover quickly and be prepared to respond rapidly to the next synaptic signal.

## Endocrine Signaling

Signals from distant cells are called **endocrine signals**, and they originate from **endocrine cells**. (In the body, many endocrine cells are located in endocrine glands, such as the thyroid gland, the hypothalamus, and the pituitary gland.) These types of signals usually produce a slower response but have a longer-lasting effect. The ligands released in endocrine signaling are called hormones, signaling molecules that are produced in one part of the body but affect other body regions some distance away.

Hormones travel the large distances between endocrine cells and their target cells via the bloodstream, which is a relatively slow way to move throughout the body. Because of their form of transport, hormones get diluted and are present in low concentrations when they act on their target cells. This is different from paracrine signaling, in which local concentrations of ligands can be very high.

## Autocrine Signaling

**Autocrine signals** are produced by signaling cells that can also bind to the ligand that is released. This means the signaling cell and the target cell can be the same or a similar cell (the prefix *auto-* means self, a reminder that the signaling cell sends a signal to itself). This type of signaling often occurs during the early development of an organism to ensure that cells develop into the correct tissues and take on the proper function. Autocrine signaling also regulates pain sensation and inflammatory responses. Further, if a cell is infected with a virus, the cell can signal itself to undergo programmed cell death, killing the virus in the process. In some cases, neighboring cells of the same type are also influenced by the released ligand. In embryological development, this process of stimulating a group of neighboring cells may help to direct the differentiation of identical cells into the same cell type, thus ensuring the proper developmental outcome.

## Direct Signaling Across Gap Junctions

Gap junctions in animals and plasmodesmata in plants are connections between the plasma membranes of neighboring cells. These water-filled channels allow small signaling molecules, called **intracellular mediators**, to diffuse between the two cells. Small molecules, such as calcium ions ($Ca^{2+}$), are able to move between cells, but large molecules like proteins and DNA cannot fit through the channels. The specificity of the channels ensures that the cells remain independent but can quickly and easily transmit signals. The transfer of signaling molecules communicates the current state of the cell that is directly next to the target cell; this allows a group of cells to coordinate their response to a signal that only one of them may have received. In plants, plasmodesmata are ubiquitous, making the entire plant into a giant, communication network.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.3: Types of Signals is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.4: Signaling Molecules

Produced by signaling cells and the subsequent binding to receptors in target cells, ligands act as chemical signals that travel to the target cells to coordinate responses. The types of molecules that serve as ligands are incredibly varied and range from small proteins to small ions like calcium ($Ca^{2+}$).

## Small Hydrophobic Ligands





Figure 1. Steroid hormones have similar chemical structures to their precursor, cholesterol. Because these molecules are small and hydrophobic, they can diffuse directly across the plasma membrane into the cell, where they interact with internal receptors.

Small hydrophobic ligands can directly diffuse through the plasma membrane and interact with internal receptors. Important members of this class of ligands are the steroid hormones. Steroids are lipids that have a hydrocarbon skeleton with four fused rings; different steroids have different functional groups attached to the carbon skeleton. Steroid hormones include the female sex hormone, estradiol, which is a type of estrogen; the male sex hormone, testosterone; and cholesterol, which is an important structural component of biological membranes and a precursor of steroid hormones (Figure 1). Other hydrophobic hormones include thyroid hormones and vitamin D. In order to be soluble in blood, hydrophobic ligands must bind to carrier proteins while they are being transported through the bloodstream.

## Water-Soluble Ligands

Water-soluble ligands are polar and therefore cannot pass through the plasma membrane unaided; sometimes, they are too large to pass through the membrane at all. Instead, most water-soluble ligands bind to the extracellular domain of cell-surface receptors. This group of ligands is quite diverse and includes small molecules, peptides, and proteins.

## Other Ligands

Nitric oxide (NO) is a gas that also acts as a ligand. It is able to diffuse directly across the plasma membrane, and one of its roles is to interact with receptors in smooth muscle and induce relaxation of the tissue. NO has a very short half-life and therefore only functions over short distances. Nitroglycerin, a treatment for heart disease, acts by triggering the release of NO, which causes blood vessels to dilate (expand), thus restoring blood flow to the heart. NO has become better known recently because the pathway that it affects is targeted by prescription medications for erectile dysfunction, such as Viagra (erection involves dilated blood vessels).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.4: Signaling Molecules is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.5: Signaling Receptors

Receptors are protein molecules in the target cell or on its surface that bind ligand. There are two types of receptors, internal receptors and cell-surface receptors.

## Internal Receptors





Figure 1. Hydrophobic signaling molecules typically diffuse across the plasma membrane and interact with intracellular receptors in the cytoplasm. Many intracellular receptors are transcription factors that interact with DNA in the nucleus and regulate gene expression.

**Internal receptors**, also known as intracellular or cytoplasmic receptors, are found in the cytoplasm of the cell and respond to hydrophobic ligand molecules that are able to travel across the plasma membrane. Once inside the cell, many of these molecules bind to proteins that act as regulators of mRNA synthesis (transcription) to mediate gene expression. Gene expression is the cellular process of transforming the information in a cell's DNA into a sequence of amino acids, which ultimately forms a protein. When the ligand binds to the internal receptor, a conformational change is triggered that exposes a DNA-binding site on the protein. The ligand-receptor complex moves into the nucleus, then binds to specific regulatory regions of the chromosomal DNA and promotes the initiation of transcription (Figure 1). Transcription is the process of copying the information in a cell's DNA into a special form of RNA called messenger RNA (mRNA); the cell uses information in the mRNA (which moves out into the cytoplasm and associates with ribosomes) to link specific amino acids in the correct order, producing a protein. Internal receptors can directly influence gene expression without having to pass the signal on to other receptors or messengers.

## Cell-Surface Receptors

**Cell-surface receptors**, also known as transmembrane receptors, are cell surface, membrane-anchored (integral) proteins that bind to external ligand molecules. This type of receptor spans the plasma membrane and performs signal transduction, in which an extracellular signal is converted into an intercellular signal. Ligands that interact with cell-surface receptors do not have to enter the cell that they affect. Cell-surface receptors are also called cell-specific proteins or markers because they are specific to individual cell types.

Each cell-surface receptor has three main components: an external ligand-binding domain, a hydrophobic membrane-spanning region, and an intracellular domain inside the cell. The ligand-binding domain is also called the **extracellular domain**. The size and extent of each of these domains vary widely, depending on the type of receptor.

Because cell-surface receptor proteins are fundamental to normal cell functioning, it should come as no surprise that a malfunction in any one of these proteins could have severe consequences. Errors in the protein structures of certain receptor molecules have been shown to play a role in hypertension (high blood pressure), asthma, heart disease, and cancer.

**Try It**

Unlike living cells, many viruses do not have a plasma membrane or any of the structures necessary to sustain life. Some viruses are simply composed of an inert protein shell containing DNA or RNA. To reproduce, viruses must invade a living cell, which serves as a host, and then take over the host's cellular apparatus. But how does a virus recognize its host?

Viruses often bind to cell-surface receptors on the host cell. For example, the virus that causes human influenza (flu) binds specifically to receptors on membranes of cells of the respiratory system. Chemical differences in the cell-surface receptors among hosts mean that a virus that infects a specific species (for example, humans) cannot infect another species (for example, chickens).

However, viruses have very small amounts of DNA or RNA compared to humans, and, as a result, viral reproduction can occur rapidly. Viral reproduction invariably produces errors that can lead to changes in newly produced viruses; these changes mean that the viral proteins that interact with cell-surface receptors may evolve in such a way that they can bind to receptors in a new host. Such changes happen randomly and quite often in the reproductive cycle of a virus, but the changes only matter if a virus with new binding properties comes into contact with a suitable host. In the case of influenza, this situation can occur in settings where animals and people are in close contact, such as poultry and swine farms. Once a virus jumps to a new host, it can spread quickly. Scientists watch newly appearing viruses (called emerging viruses) closely in the hope that such monitoring can reduce the likelihood of global viral epidemics.

Cell-surface receptors are involved in most of the signaling in multicellular organisms. There are three general categories of cell-surface receptors: ion channel-linked receptors, G-protein-linked receptors, and enzyme-linked receptors.





Figure 2. Gated ion channels form a pore through the plasma membrane that opens when the signaling molecule binds. The open pore then allows ions to flow into or out of the cell.

**Ion channel-linked receptors** bind a ligand and open a channel through the membrane that allows specific ions to pass through. To form a channel, this type of cell-surface receptor has an extensive membrane-spanning region. In order to interact with the phospholipid fatty acid tails that form the center of the plasma membrane, many of the amino acids in the membrane-spanning region are hydrophobic in nature. Conversely, the amino acids that line the inside of the channel are hydrophilic to allow for the passage of water or ions. When a ligand binds to the extracellular region of the channel, there is a conformational change in the proteins structure that allows ions such as sodium, calcium, magnesium, and hydrogen to pass through (Figure 2).

**G-protein-linked receptors** bind a ligand and activate a membrane protein called a G-protein. The activated G-protein then interacts with either an ion channel or an enzyme in the membrane (Figure 3). All G-protein-linked receptors have seven transmembrane domains, but each receptor has its own specific extracellular domain and G-protein-binding site.

Cell signaling using G-protein-linked receptors occurs as a cyclic series of events. Before the ligand binds, the inactive G-protein can bind to a newly revealed site on the receptor specific for its binding. Once the G-protein binds to the receptor, the resultant shape change activates the G-protein, which releases GDP and picks up GTP. The subunits of the G-protein then split into the $\alpha$ subunit and the $\beta\gamma$ subunit. One or both of these G-protein fragments may be able to activate other proteins as a result. After awhile, the GTP on the active $\alpha$ subunit of the G-protein is hydrolyzed to GDP and the $\beta\gamma$ subunit is deactivated. The subunits reassociate to form the inactive G-protein and the cycle begins anew.





Figure 3. Heterotrimeric G proteins have three subunits: *α*, *β*, and *γ*. When a signaling molecule binds to a G-protein-coupled receptor in the plasma membrane, a GDP molecule associated with the *α* subunit is exchanged for GTP. The *β* and *γ* subunits dissociate from the *α* subunit, and a cellular response is triggered either by the *α* subunit or the dissociated *βγ* pair. Hydrolysis of GTP to GDP terminates the signal.

G-protein-linked receptors have been extensively studied and much has been learned about their roles in maintaining health. Bacteria that are pathogenic to humans can release poisons that interrupt specific G-protein-linked receptor function, leading to illnesses such as pertussis, botulism, and cholera.





Figure 4. Transmitted primarily through contaminated drinking water, cholera is a major cause of death in the developing world and in areas where natural disasters interrupt the availability of clean water. (credit: New York City Sanitary Commission)

In cholera (Figure 4), for example, the water-borne bacterium *Vibrio cholerae* produces a toxin, choleragen, that binds to cells lining the small intestine. The toxin then enters these intestinal cells, where it modifies a G-protein that controls the opening of a chloride channel and causes it to remain continuously active, resulting in large losses of fluids from the body and potentially fatal dehydration as a result.

Modern sanitation eliminates the threat of cholera outbreaks, such as the one that swept through New York City in 1866. This poster from that era shows how, at that time, the way that the disease was transmitted was not understood.

**Enzyme-linked receptors** are cell-surface receptors with intracellular domains that are associated with an enzyme. In some cases, the intracellular domain of the receptor itself is an enzyme. Other enzyme-linked receptors have a small intracellular domain that interacts directly with an enzyme. The enzyme-linked receptors normally have large extracellular and intracellular domains, but the membrane-spanning region consists of a single alpha-helical region of the peptide strand. When a ligand binds to the extracellular domain, a signal is transferred through the membrane, activating the enzyme. Activation of the enzyme sets off a chain of events within the cell that eventually leads to a response. One example of this type of enzyme-linked receptor is the tyrosine kinase receptor (Figure 5). A kinase is an enzyme that transfers phosphate groups from ATP to another protein. The tyrosine kinase receptor transfers phosphate groups to tyrosine molecules (tyrosine residues). First, signaling molecules bind to the extracellular domain of two nearby tyrosine kinase receptors. The two neighboring receptors then bond together, or dimerize. Phosphates are then added to tyrosine residues on the intracellular domain of the receptors (phosphorylation). The phosphorylated residues can then transmit the signal to the next messenger within the cytoplasm.

Practice Question





Figure 5. A receptor tyrosine kinase is an enzyme-linked receptor with a single transmembrane region, and extracellular and intracellular domains. Binding of a signaling molecule to the extracellular domain causes the receptor to dimerize. Tyrosine residues on the intracellular domain are then autophosphorylated, triggering a downstream cellular response. The signal is terminated by a phosphatase that removes the phosphates from the phosphotyrosine residues.

HER2 is a receptor tyrosine kinase. In 30 percent of human breast cancers, HER2 is permanently activated, resulting in unregulated cell division. Lapatinib, a drug used to treat breast cancer, inhibits HER2 receptor tyrosine kinase autophosphorylation (the process by which the receptor adds phosphates onto itself), thus reducing tumor growth by 50 percent. Besides autophosphorylation, which of the following steps would be inhibited by Lapatinib?

a. Signaling molecule binding, dimerization, and the downstream cellular response
b. Dimerization, and the downstream cellular response
c. The downstream cellular response
d. Phosphatase activity, dimerization, and the downsteam cellular response

[reveal-answer q=”222996″]**Show Answer**[/reveal-answer]
[hidden-answer a=”222996″]Answer c. The downstream cellular response would be inhibited.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.5: Signaling Receptors is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.6: Introduction to Propagation of the Signal

## Describe how a cell propagates a signal

Once a ligand binds to a receptor, the signal is transmitted through the membrane and into the cytoplasm. Continuation of a signal in this manner is called **signal transduction**. Signal transduction only occurs with cell-surface receptors because internal receptors are able to interact directly with DNA in the nucleus to initiate protein synthesis.

When a ligand binds to its receptor, conformational changes occur that affect the receptor's intracellular domain. Conformational changes of the extracellular domain upon ligand binding can propagate through the membrane region of the receptor and lead to activation of the intracellular domain or its associated proteins. In some cases, binding of the ligand causes **dimerization** of the receptor, which means that two receptors bind to each other to form a stable complex called a dimer. A **dimer** is a chemical compound formed when two molecules (often identical) join together. The binding of the receptors in this manner enables their intracellular domains to come into close contact and activate each other.

## What You'll Learn to Do

- Explain how the binding of a ligand initiates signal transduction throughout a cell
- Recognize the role of phosphorylation in the transmission of intracellular signals
- Evaluate the role of second messengers in signal transmission

### Learning Activities

The learning activities for this section include the following:

- Binding Initiates a Signaling Pathway
- Methods of Intracellular Signaling
- Self Check: Propagation of the Signal

## Contributors and Attributions

CC licensed content, Original

- Introduction to Propagation of the Signal. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.6: Introduction to Propagation of the Signal is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.7: Binding Initiates a Signaling Pathway

After the ligand binds to the cell-surface receptor, the activation of the receptor's intracellular components sets off a chain of events that is called a signaling pathway or a signaling cascade. This process is sometimes called signal transduction. In a **signaling pathway**, second messengers, enzymes, and activated proteins interact with specific proteins, which are in turn activated in a chain reaction that eventually leads to a change in the cell's environment (Figure 1). The events in the cascade occur in a series, much like a current flows in a river. Interactions that occur before a certain point are defined as upstream events, and events after that point are called downstream events.

## Practice Question





Figure 1. The epidermal growth factor (EGF) receptor (EGFR) is a receptor tyrosine kinase involved in the regulation of cell growth, wound healing, and tissue repair. When EGF binds to the EGFR, a cascade of downstream events causes the cell to grow and divide. If EGFR is activated at inappropriate times, uncontrolled cell growth (cancer) may occur.

In certain cancers, the GTPase activity of the RAS G-protein is inhibited. This means that the RAS protein can no longer hydrolyze GTP into GDP. What effect would this have on downstream cellular events?

[practice-area rows="2"][/practice-area]
[reveal-answer q="265793"]**Show Answer**[/reveal-answer]
[hidden-answer a="265793"]ERK would become permanently activated, resulting in cell proliferation, migration, adhesion, and the growth of new blood vessels. Apoptosis would be inhibited.[/hidden-answer]

Signaling pathways can get very complicated very quickly because most cellular proteins can affect different downstream events, depending on the conditions within the cell. A single pathway can branch off toward different endpoints based on the interplay between two or more signaling pathways, and the same ligands are often used to initiate different signals in different cell types. This variation in response is due to differences in protein expression in different cell types. Another complicating element is **signal integration** of the pathways, in which signals from two or more different cell-surface receptors merge to activate the same response in the cell. This process can ensure that multiple external requirements are met before a cell commits to a specific response.

The effects of extracellular signals can also be amplified by enzymatic cascades. At the initiation of the signal, a single ligand binds to a single receptor. However, activation of a receptor-linked enzyme can activate many copies of a component of the signaling

cascade, which amplifies the signal.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

9.7: Binding Initiates a Signaling Pathway is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.8: Methods of Intracellular Signaling

The induction of a signaling pathway depends on the modification of a cellular component by an enzyme. There are numerous enzymatic modifications that can occur, and they are recognized in turn by the next component downstream. The following are some of the more common events in intracellular signaling.

Observe an animation of cell signaling at this site.

## Phosphorylation





Figure 1. In protein phosphorylation, a phosphate group ($PO_4^{-3}$) is added to residues of the amino acids serine, threonine, and tyrosine.

One of the most common chemical modifications that occurs in signaling pathways is the addition of a phosphate group ($PO_4^{-3}$) to a molecule such as a protein in a process called phosphorylation. The phosphate can be added to a nucleotide such as GMP to form GDP or GTP. Phosphates are also often added to serine, threonine, and tyrosine residues of proteins, where they replace the hydroxyl group of the amino acid (Figure 1). The transfer of the phosphate is catalyzed by an enzyme called a **kinase**. Various kinases are named for the substrate they phosphorylate. Phosphorylation of serine and threonine residues often activates enzymes. Phosphorylation of tyrosine residues can either affect the activity of an enzyme or create a binding site that interacts with downstream components in the signaling cascade. Phosphorylation may activate or inactivate enzymes, and the reversal of phosphorylation, dephosphorylation by a phosphatase, will reverse the effect.

## Second Messengers

**Second messengers** are small molecules that propagate a signal after it has been initiated by the binding of the signaling molecule to the receptor. These molecules help to spread a signal through the cytoplasm by altering the behavior of certain cellular proteins.

Calcium ion is a widely used second messenger. The free concentration of calcium ions ($Ca^{2+}$) within a cell is very low because ion pumps in the plasma membrane continuously use adenosine-5′-triphosphate (ATP) to remove it. For signaling purposes, $Ca^{2+}$ is stored in cytoplasmic vesicles, such as the endoplasmic reticulum, or accessed from outside the cell. When signaling occurs, ligand-gated calcium ion channels allow the higher levels of $Ca^{2+}$ that are present outside the cell (or in intracellular storage compartments) to flow into the cytoplasm, which raises the concentration of cytoplasmic $Ca^{2+}$. The response to the increase in $Ca^{2+}$ varies, depending on the cell type involved. For example, in the β-cells of the pancreas, $Ca^{2+}$ signaling leads to the release of insulin, and in muscle cells, an increase in $Ca^{2+}$ leads to muscle contractions.

Another second messenger utilized in many different cell types is **cyclic AMP (cAMP)**. Cyclic AMP is synthesized by the enzyme adenylyl cyclase from ATP (Figure 2). The main role of cAMP in cells is to bind to and activate an enzyme called cAMP-dependent kinase (A-kinase). A-kinase regulates many vital metabolic pathways: It phosphorylates serine and threonine residues of its target

proteins, activating them in the process. A-kinase is found in many different types of cells, and the target proteins in each kind of cell are different. Differences give rise to the variation of the responses to cAMP in different cells.





Figure 2. This diagram shows the mechanism for the formation of cyclic AMP (cAMP). cAMP serves as a second messenger to activate or inactivate proteins within the cell. Termination of the signal occurs when an enzyme called phosphodiesterase converts cAMP into AMP.

Present in small concentrations in the plasma membrane, **inositol phospholipids** are lipids that can also be converted into second messengers. Because these molecules are membrane components, they are located near membrane-bound receptors and can easily interact with them. Phosphatidylinositol (PI) is the main phospholipid that plays a role in cellular signaling. Enzymes known as kinases phosphorylate PI to form PI-phosphate (PIP) and PI-bisphosphate ($PIP_2$).

The enzyme phospholipase C cleaves $PIP_2$ to form **diacylglycerol (DAG) and inositol triphosphate ($IP_3$)** (Figure 3). These products of the cleavage of $PIP_2$ serve as second messengers. Diacylglycerol (DAG) remains in the plasma membrane and activates protein kinase C (PKC), which then phosphorylates serine and threonine residues in its target proteins. $IP_3$ diffuses into the cytoplasm and binds to ligand-gated calcium channels in the endoplasmic reticulum to release $Ca^{2+}$ that continues the signal cascade.





Figure 3. The enzyme phospholipase C breaks down $PIP_2$ into $IP_3$ and DAG, both of which serve as second messengers.

## Learning Objectives

Ligand binding to the receptor allows for signal transduction through the cell. The chain of events that conveys the signal through the cell is called a signaling pathway or cascade. Signaling pathways are often very complex because of the interplay between different proteins. A major component of cell signaling cascades is the phosphorylation of molecules by enzymes known as kinases. Phosphorylation adds a phosphate group to serine, threonine, and tyrosine residues in a protein, changing their shapes, and activating or inactivating the protein. Small molecules like nucleotides can also be phosphorylated. Second messengers are small, non-protein molecules that are used to transmit a signal within a cell. Some examples of second messengers are calcium ions ($Ca^{2+}$), cyclic AMP (cAMP), diacylglycerol (DAG), and inositol triphosphate ($IP_3$).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.8: Methods of Intracellular Signaling is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.9: Introduction to Response to the Signal

## Describe how a cell responds to a signal

Inside the cell, ligands bind to their internal receptors, allowing them to directly affect the cell's DNA and protein-producing machinery. Using signal transduction pathways, receptors in the plasma membrane produce a variety of effects on the cell. The results of signaling pathways are extremely varied and depend on the type of cell involved as well as the external and internal conditions. A small sampling of responses is described in this outcome.

## What You'll Learn to Do

- Describe how signaling pathways direct protein expression, cellular metabolism, and cell growth
- Recognize the role of apoptosis in the development and maintenance of a healthy organism

### Learning Activities

The learning activities for this section include the following:

- Expression and Growth
- Cell Death
- Self Check: Response to the Signal

## Contributors and Attributions

CC licensed content, Original

- Introduction to Response to the Signal. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

9.9: Introduction to Response to the Signal is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.10: Expression and Growth

## Gene Expression

Some signal transduction pathways regulate the transcription of RNA. Others regulate the translation of proteins from mRNA. An example of a protein that regulates translation in the nucleus is the MAP kinase ERK. ERK is activated in a phosphorylation cascade when epidermal growth factor (EGF) binds the EGF receptor. Upon phosphorylation, ERK enters the nucleus and activates a protein kinase that, in turn, regulates protein translation (Figure 1).





Figure 1. ERK is a MAP kinase that activates translation when it is phosphorylated. ERK phosphorylates MNK1, which in turn phosphorylates eIF-4E, an elongation initiation factor that, with other initiation factors, is associated with mRNA. When eIF-4E becomes phosphorylated, the mRNA unfolds, allowing protein synthesis in the nucleus to begin.

The second kind of protein with which PKC can interact is a protein that acts as an inhibitor. An **inhibitor** is a molecule that binds to a protein and prevents it from functioning or reduces its function. In this case, the inhibitor is a protein called Iκ-B, which binds to the regulatory protein NF-κB. (The symbol κ represents the Greek letter kappa.) When Iκ-B is bound to NF-κB, the complex cannot enter the nucleus of the cell, but when Iκ-B is phosphorylated by PKC, it can no longer bind NF-κB, and NF-κB (a transcription factor) can enter the nucleus and initiate RNA transcription. In this case, the effect of phosphorylation is to inactivate an inhibitor and thereby activate the process of transcription.

## Increase in Cellular Metabolism

The result of another signaling pathway affects muscle cells. The activation of β-adrenergic receptors in muscle cells by adrenaline leads to an increase in cyclic AMP (cAMP) inside the cell. Also known as epinephrine, adrenaline is a hormone (produced by the adrenal gland attached to the kidney) that readies the body for short-term emergencies. Cyclic AMP activates PKA (protein kinase A), which in turn phosphorylates two enzymes. The first enzyme promotes the degradation of glycogen by activating intermediate glycogen phosphorylase kinase (GPK) that in turn activates glycogen phosphorylase (GP) that catabolizes glycogen into glucose. (Note that your body converts excess glucose to glycogen for short-term storage. When energy is needed, glycogen is quickly reconverted to glucose.) Phosphorylation of the second enzyme, glycogen synthase (GS), inhibits its ability to form glycogen from glucose. In this manner, a muscle cell obtains a ready pool of glucose by activating its formation via glycogen degradation and by inhibiting the use of glucose to form glycogen, thus preventing a futile cycle of glycogen degradation and synthesis. The glucose is then available for use by the muscle cell in response to a sudden surge of adrenaline—the “fight or flight” reflex.

## Cell Growth

Cell signaling pathways also play a major role in cell division. Cells do not normally divide unless they are stimulated by signals from other cells. The ligands that promote cell growth are called **growth factors**. Most growth factors bind to cell-surface receptors that are linked to tyrosine kinases. These cell-surface receptors are called receptor tyrosine kinases (RTKs). Activation of RTKs initiates a signaling pathway that includes a G-protein called RAS, which activates the MAP kinase pathway described earlier. The enzyme MAP kinase then stimulates the expression of proteins that interact with other cellular components to initiate cell division.

## Try It

Cancer biologists study the molecular origins of cancer with the goal of developing new prevention methods and treatment strategies that will inhibit the growth of tumors without harming the normal cells of the body. As mentioned earlier, signaling pathways control cell growth. These signaling pathways are controlled by signaling proteins, which are, in turn, expressed by genes. Mutations in these genes can result in malfunctioning signaling proteins. This prevents the cell from regulating its cell cycle, triggering unrestricted cell division and cancer. The genes that regulate the signaling proteins are one type of oncogene, which is a gene that has the potential to cause cancer. The gene encoding RAS is an oncogene that was originally discovered when mutations in the RAS protein were linked to cancer. Further studies have indicated that 30 percent of cancer cells have a mutation in the RAS gene that leads to uncontrolled growth. If left unchecked, uncontrolled cell division can lead tumor formation and metastasis, the growth of cancer cells in new locations in the body.

Cancer biologists have been able to identify many other oncogenes that contribute to the development of cancer. For example, HER2 is a cell-surface receptor that is present in excessive amounts in 20 percent of human breast cancers. Cancer biologists realized that gene duplication led to HER2 overexpression in 25 percent of breast cancer patients and developed a drug called Herceptin (trastuzumab). Herceptin is a monoclonal antibody that targets HER2 for removal by the immune system. Herceptin therapy helps to control signaling through HER2. The use of Herceptin in combination with chemotherapy has helped to increase the overall survival rate of patients with metastatic breast cancer.

More information on cancer biology research can be found at the National Cancer Institute website.

## Learning Objectives

Many pathways influence the cell by initiating gene expression, and the methods utilized are quite numerous. Some pathways activate enzymes that interact with DNA transcription factors. Others modify proteins and induce them to change their location in the cell. Depending on the status of the organism, cells can respond by storing energy as glycogen or fat, or making it available in the form of glucose. A signal transduction pathway allows muscle cells to respond to immediate requirements for energy in the form of glucose. Cell growth is almost always stimulated by external signals called growth factors. Uncontrolled cell growth leads to cancer, and mutations in the genes encoding protein components of signaling pathways are often found in tumor cells.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.10: Expression and Growth is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.11: Cell Death




Figure 1. The histological section of a foot of a 15-day-old mouse embryo, visualized using light microscopy, reveals areas of tissue between the toes, which apoptosis will eliminate before the mouse reaches its full gestational age at 27 days. (credit: modification of work by Michal Mañas)

When a cell is damaged, superfluous, or potentially dangerous to an organism, a cell can initiate a mechanism to trigger programmed cell death, or **apoptosis**. Apoptosis allows a cell to die in a controlled manner that prevents the release of potentially damaging molecules from inside the cell. There are many internal checkpoints that monitor a cell's health; if abnormalities are observed, a cell can spontaneously initiate the process of apoptosis. However, in some cases, such as a viral infection or uncontrolled cell division due to cancer, the cell's normal checks and balances fail. External signaling can also initiate apoptosis. For example, most normal animal cells have receptors that interact with the extracellular matrix, a network of glycoproteins that provides structural support for cells in an organism. The binding of cellular receptors to the extracellular matrix initiates a signaling cascade within the cell. However, if the cell moves away from the extracellular matrix, the signaling ceases, and the cell undergoes apoptosis. This system keeps cells from traveling through the body and proliferating out of control, as happens with tumor cells that metastasize.

Another example of external signaling that leads to apoptosis occurs in T-cell development. T-cells are immune cells that bind to foreign macromolecules and particles, and target them for destruction by the immune system. Normally, T-cells do not target "self" proteins (those of their own organism), a process that can lead to autoimmune diseases. In order to develop the ability to discriminate between self and non-self, immature T-cells undergo screening to determine whether they bind to so-called self proteins. If the T-cell receptor binds to self proteins, the cell initiates apoptosis to remove the potentially dangerous cell.

Apoptosis is also essential for normal embryological development. In vertebrates, for example, early stages of development include the formation of web-like tissue between individual fingers and toes (Figure 1). During the course of normal development, these unneeded cells must be eliminated, enabling fully separated fingers and toes to form. A cell signaling mechanism triggers apoptosis, which destroys the cells between the developing digits.

## Termination of the Signal Cascade

The aberrant signaling often seen in tumor cells is proof that the termination of a signal at the appropriate time can be just as important as the initiation of a signal. One method of stopping a specific signal is to degrade the ligand or remove it so that it can no longer access its receptor. One reason that hydrophobic hormones like estrogen and testosterone trigger long-lasting events is because they bind carrier proteins. These proteins allow the insoluble molecules to be soluble in blood, but they also protect the hormones from degradation by circulating enzymes.

Inside the cell, many different enzymes reverse the cellular modifications that result from signaling cascades. For example, **phosphatases** are enzymes that remove the phosphate group attached to proteins by kinases in a process called dephosphorylation. Cyclic AMP (cAMP) is degraded into AMP by **phosphodiesterase**, and the release of calcium stores is reversed by the $Ca^{2+}$ pumps that are located in the external and internal membranes of the cell.

## Learning Objectives

Programmed cell death, or apoptosis, is important for removing damaged or unnecessary cells. The use of cellular signaling to organize the dismantling of a cell ensures that harmful molecules from the cytoplasm are not released into the spaces between cells, as they are in uncontrolled death, necrosis. Apoptosis also ensures the efficient recycling of the components of the dead cell. Termination of the cellular signaling cascade is very important so that the response to a signal is appropriate in both timing and intensity. Degradation of signaling molecules and dephosphorylation of phosphorylated intermediates of the pathway by phosphatases are two ways to terminate signals within the cell.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.11: Cell Death is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.12: Introduction to Signaling in Single-Celled Organisms

## Discuss the process of signaling in single-cell organisms

Within-cell signaling allows bacteria to respond to environmental cues, such as nutrient levels, some single-celled organisms also release molecules to signal to each other.

## What You'll Learn to Do

- Describe how single-celled yeasts use cell signaling to communicate with one another
- Relate the role of quorum sensing to the ability of some bacteria to form biofilms

### Learning Activities

The learning activities for this section include the following:

- Signaling in Yeast
- Signaling in Bacteria
- Self Check: Signaling in Single-Celled Organisms

## Contributors and Attributions

CC licensed content, Original

- Introduction to Signaling in Single-Celled Organisms. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC0: No Rights Reserved*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.12: Introduction to Signaling in Single-Celled Organisms is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.13: Signaling in Yeast





Figure 1. Micrograph of budding *Saccharomyces cerevisiae* yeast cells.

Yeasts are eukaryotes (fungi), and the components and processes found in yeast signals are similar to those of cell-surface receptor signals in multicellular organisms. Budding yeasts are able to participate in a process that is similar to sexual reproduction that entails two haploid cells (cells with one-half the normal number of chromosomes) combining to form a diploid cell (a cell with two sets of each chromosome, which is what normal body cells contain). Figure 1, shows a micrograph of this process. The budding *Saccharomyces cerevisiae* yeast cells are visualized using differential interference contrast microscopy, a light microscopy technique that enhances the contrast of the sample.

In order to find another haploid yeast cell that is prepared to mate, budding yeasts secrete a signaling molecule called **mating factor**. When mating factor binds to cell-surface receptors in other yeast cells that are nearby, they stop their normal growth cycles and initiate a cell signaling cascade that includes protein kinases and GTP-binding proteins that are similar to G-proteins.

## Cellular Communication in Yeasts

The first life on our planet consisted of single-celled prokaryotic organisms that had limited interaction with each other. While some external signaling occurs between different species of single-celled organisms, the majority of signaling within bacteria and yeasts concerns only other members of the same species. The evolution of cellular communication is an absolute necessity for the development of multicellular organisms, and this innovation is thought to have required approximately 2.5 billion years to appear in early life forms.

Yeasts are single-celled eukaryotes, and therefore have a nucleus and organelles characteristic of more complex life forms. Comparisons of the genomes of yeasts, nematode worms, fruit flies, and humans illustrate the evolution of increasingly complex signaling systems that allow for the efficient inner workings that keep humans and other complex life forms functioning correctly.

Kinases are a major component of cellular communication, and studies of these enzymes illustrate the evolutionary connectivity of different species. Yeasts have 130 types of kinases. More complex organisms such as nematode worms and fruit flies have 454 and 239 kinases, respectively. Of the 130 kinase types in yeast, 97 belong to the 55 subfamilies of kinases that are found in other eukaryotic organisms. The only obvious deficiency seen in yeasts is the complete absence of tyrosine kinases. It is hypothesized that phosphorylation of tyrosine residues is needed to control the more sophisticated functions of development, differentiation, and cellular communication used in multicellular organisms.

Because yeasts contain many of the same classes of signaling proteins as humans, these organisms are ideal for studying signaling cascades. Yeasts multiply quickly and are much simpler organisms than humans or other multicellular animals. Therefore, the signaling cascades are also simpler and easier to study, although they contain similar counterparts to human signaling.[1]

## Practice Question

What characteristics make yeasts a good model for learning about signaling in humans?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”285048″]Show Answer[/reveal-answer]

[hidden-answer a="285048"]Yeasts are eukaryotes and have many of the same systems that humans do; however, they are single-celled, so they are easy to grow, grow rapidly, have a short generation time, and are much simpler than humans.[/hidden-answer]

1. G. Manning, G.D. Plowman, T. Hunter, S. Sudarsanam, "Evolution of Protein Kinase Signaling from Yeast to Man," *Trends in Biochemical Sciences* 27, no. 10 (2002): 514–520. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

9.13: Signaling in Yeast is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.14: Signaling in Bacteria

Signaling in bacteria enables bacteria to monitor extracellular conditions, ensure that there are sufficient amounts of nutrients, and ensure that hazardous situations are avoided. There are circumstances, however, when bacteria communicate with each other.

The first evidence of bacterial communication was observed in a bacterium that has a symbiotic relationship with Hawaiian bobtail squid. When the population density of the bacteria reaches a certain level, specific gene expression is initiated, and the bacteria produce bioluminescent proteins that emit light. Because the number of cells present in the environment (cell density) is the determining factor for signaling, bacterial signaling was named **quorum sensing**. In politics and business, a quorum is the minimum number of members required to be present to vote on an issue.

Quorum sensing uses autoinducers as signaling molecules. **Autoinducers** are signaling molecules secreted by bacteria to communicate with other bacteria of the same kind. The secreted autoinducers can be small, hydrophobic molecules such as acyl-homoserine lactone, (AHL) or larger peptide-based molecules; each type of molecule has a different mode of action. When AHL enters target bacteria, it binds to transcription factors, which then switch gene expression on or off (see Figure 1). The peptide autoinducers stimulate more complicated signaling pathways that include bacterial kinases. The changes in bacteria following exposure to autoinducers can be quite extensive. The pathogenic bacterium *Pseudomonas aeruginosa* has 616 different genes that respond to autoinducers.

## Practice Question





Figure 1. Autoinducers are small molecules or proteins produced by bacteria that regulate gene expression.

Which of the following statements about quorum sensing is false?


a. Autoinducer must bind to receptor to turn on transcription of genes responsible for the production of more autoinducer.
b. The receptor stays in the bacterial cell, but the autoinducer diffuses out.
c. Autoinducer can only act on a different cell: it cannot act on the cell in which it is made.
d. Autoinducer turns on genes that enable the bacteria to form a biofilm.

[reveal-answer q="147760"]Show Answer[/reveal-answer]
[hidden-answer a="147760"]Answer c is false.[/hidden-answer]

Some species of bacteria that use quorum sensing form biofilms, complex colonies of bacteria (often containing several species) that exchange chemical signals to coordinate the release of toxins that will attack the host. Bacterial biofilms (Figure 2) can sometimes be found on medical equipment; when biofilms invade implants such as hip or knee replacements or heart pacemakers, they can cause life-threatening infections.

Cell-cell communication enables *Staphylococcus aureus* bacteria (Figure 2a) to work together to form a biofilm inside a hospital patient's catheter, seen here via scanning electron microscopy. *S. aureus* is the main cause of hospital-acquired infections. Hawaiian bobtail squid (Figure 2b) have a symbiotic relationship with the bioluminescent bacteria *Vibrio fischeri*. The luminescence makes it difficult to see the squid from below because it effectively eliminates its shadow. In return for camouflage, the squid provides food for the bacteria. Free-living *V. fischeri* do not produce luciferase, the enzyme responsible for luminescence, but *V. fischeri* living in a symbiotic relationship with the squid do. Quorum sensing determines whether the bacteria should produce the luciferase enzyme.

## Practice Question




(a) (b)

Figure 2. (a) Staphylococcus aureus bacteria. (b) Hawaiian bobtail squid. (credit a: modifications of work by CDC/Janice Carr; credit b: modifications of work by Cliff1066/Flickr)

What advantage might biofilm production confer on the *S. aureus* inside the catheter?

[practice-area rows="2"][/practice-area]
[reveal-answer q="10146"]Show Answer[/reveal-answer]
[hidden-answer a="10146"]*S. aureus* produces a biofilm because the higher cell density in the biofilm permits the formation of a dense surface that helps protect the bacteria from antibiotics.[/hidden-answer]

Research on the details of quorum sensing has led to advances in growing bacteria for industrial purposes. Recent discoveries suggest that it may be possible to exploit bacterial signaling pathways to control bacterial growth; this process could replace or supplement antibiotics that are no longer effective in certain situations.

Watch geneticist Bonnie Bassler discuss her discovery of quorum sensing in biofilm bacteria in squid.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for "How Bacteria 'Talk' " here (link opens in new window).

Watch this collection of interview clips with biofilm researchers in "What Are Bacterial Biofilms?"

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for "What are Bacterial Biofilms? A Six Minute Montage" here (link opens in new window).

## Practice Questions

Quorum sensing is triggered to begin when _____.

a. treatment with antibiotics occurs
b. bacteria release growth hormones
c. bacterial protein expression is switched on
d. a sufficient number of bacteria are present

[reveal-answer q="578321"]Show Answer[/reveal-answer]
[hidden-answer a="578321"]Answer d. Quorum sensing is triggered to begin when a sufficient number of bacteria are present.

[/hidden-answer]

Why is signaling in multicellular organisms more complicated than signaling in single-celled organisms?

[practice-area rows="2"][/practice-area]
[reveal-answer q="43987"]Show Answer[/reveal-answer]
[hidden-answer a="43987"]Multicellular organisms must coordinate many different events in different cell types that may be very distant from each other. Single-celled organisms are only concerned with their immediate environment and the presence of other cells in the area.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content


- How bacteria talk - Bonnie Bassler. **Authored by**: Bonnie Bassler. **Provided by**: TED-Ed. **Located at**: https://youtu.be/KXWurAmtf78. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License
- What Are Bacterial Biofilms? A Six Minute Montage. **Authored by**: biofilm. **Located at**: https://youtu.be/lpI4WCM_9pM. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

9.14: Signaling in Bacteria is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 9.15: Putting It Together- Cell Communication

Now that we've learned about cell communication, let's the process of a single message as the cell passes it along:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=310

We just watched how epinephrine is used as a sample messenger to trigger the release of glucose from cells in the liver. This is an example of phosphorylation, one of the two methods of intracellular signaling. The other method—second messengers—can be seen in $Ca^{2+}$ signaling in muscle cells, which leads to muscle contractions.

These are just two examples of the many, many body functions that rely on cell communication. As we discussed in opening this module, imagine again what would happen if we could not communicate in society. Now, with your new appreciation of cell communication, imagine what would happen if even a single aspect of that process broke down.

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: Cell Communication. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

All rights reserved content

- Signal Transduction Pathways. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/qOVkedxDqQo. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

9.15: Putting It Together- Cell Communication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 10: Module 8- Cell Division


10.1: Why It Matters- Cell Division
10.2: Introduction to Chromosomes and DNA Packaging
10.3: DNA and Chromosomes
10.4: Chromosome Structure
10.5: Introduction to the Cell Cycle
10.6: Interphase
10.7: Mitosis
10.8: Cytokinesis
10.9: The Complete Cycle
10.10: Introduction to Cell Cycle Checkpoints
10.11: Control of the Cell Cycle
10.12: Cancer and the Cell Cycle
10.13: Introduction to Meiosis
10.14: Stages of Meiosis
10.15: Meiosis I
10.16: Meiosis II
10.17: Meiosis
10.18: Introduction to Genetic Diversity
10.19: Sexual Reproduction
10.20: Genetic Variation in Meiosis
10.21: Chromosomal Structural Rearrangements
10.22: Mitosis, Meiosis, and Sexual Reproduction
10.23: Introduction to Errors in Chromosome Number
10.24: Karyotypes
10.25: Common Disorders
10.26: Putting It Together- Cell Division
10.27: Assignment- Mitosis and Meiosis Worksheets



10: Module 8- Cell Division is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.1: Why It Matters- Cell Division

## Why describe and explain the various stages of cell division?

Cell division is key to life: from the moment we are first conceived, we are continually changing and growing. In order for our bodies to grow and develop, they must produce new cells—and allow for the death of old cells. Cell division is also an essential component of injury repair. If our cells couldn't divide and create new cells, our bodies could never produce new skin cells to heal road rash, or grow a fingernail back. However, when cell division goes awry, dramatic results may occur. Without sufficient cellular oversight, repeated rounds of unregulated cell division can lead to a minor condition like psoriasis or a life-threatening disease like cancer. Cell division takes occurs by a strict cycle, with multiple stages and checkpoints to ensure things *don't* go awry.

Perhaps most importantly, without cell division, no species would be able to reproduce—life would simply end (or would have ended a long time ago). Every human, as well as every sexually reproducing organism, begins life as a fertilized egg (embryo) or zygote. Trillions of cell divisions subsequently occur in a controlled manner to produce a complex, multicellular human. In other words, that original single cell is the ancestor of every other cell in the body. Single-celled organisms use cell division as their method of reproduction.





Figure 1. A sea urchin begins life as a single cell that (a) divides to form two cells, visible by scanning electron microscopy. After four rounds of cell division, (b) there are 16 cells, as seen in this SEM image. After many rounds of cell division, the individual develops into a complex, multicellular organism, as seen in this (c) mature sea urchin. (credit a: modification of work by Evelyn Spiegel, Louisa Howard; credit b: modification of work by Evelyn Spiegel, Louisa Howard; credit c: modification of work by Marco Busdraghi; scale-bar data from Matt Russell)

### Learning Outcomes

- Understand chromosome structure and organization in eukaryotic cells
- Identify the stages of the cell cycle, by picture and by description of major milestones
- Identify and explain the important checkpoints that a cell passes through during the cell cycle
- Identify the stages of meiosis by picture and by description of major milestones; explain why meiosis involves two rounds of nuclear division
- Describe and explain a range of mechanisms for generating genetic diversity
- Examine karyotypes and identify the effects of significant changes in chromosome number

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Cell Division. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*
- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.1: Why It Matters- Cell Division is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.2: Introduction to Chromosomes and DNA Packaging

## Understand chromosome structure and organization in eukaryotic cells

In this outcome, we will discuss autosomes and sex chromosomes and the differences between the two as well as how cells package DNA.

## What You'll Learn to Do

- Differentiate between two kinds of chromosomes: autosomes and sex chromosomes
- Understand how DNA is protected and compacted inside cells

### Learning Activities

The learning activities for this section include the following:

- DNA and Chromosomes
- Chromosome Structure
- Self Check: Types of Chromosomes

## Contributors and Attributions

CC licensed content, Original

- Introduction to Chromosomes and DNA Packaging. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*


10.2: Introduction to Chromosomes and DNA Packaging is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.3: DNA and Chromosomes

When a cell divides in two, one of its main jobs is to make sure that each of the two new cells gets a full, perfect copy of genetic material. Mistakes during copying, or unequal division of the genetic material between cells, can lead to cells that are unhealthy or nonfunctional (and even to diseases such as cancer). But what exactly is this genetic material, and how does it behave over the course of a cell division?

## DNA and Genomes

**DNA (deoxyribonucleic acid)** is the genetic material of living organisms. In humans, DNA is found in almost all the cells of the body and provides the instructions they need to grow, function, and respond to their environment. When a cell of the body divides, it will pass on a copy of its DNA to each of its daughter cells. DNA is also passed on at the at the level of organisms, with the DNA in sperm and egg cells combining to form a new organism that has genetic material from both its parents. Physically speaking, DNA is a long string of paired chemical units (nucleotides) that come in four different types, and it carries information organized into units called genes. Genes typically provide instructions for making proteins, which give cells and organisms their functional characteristics.




In eukaryotes such as plants and animals, the great majority of DNA is found in the nucleus and is called **nuclear DNA**. Mitochondria, organelles that harvest energy for the cell, contain their own **mitochondrial DNA**, and chloroplasts, organelles that carry out photosynthesis in plant cells, also have **chloroplast DNA**. The amounts of DNA found in mitochondria and chloroplasts are much smaller than the amount found in the nucleus. In bacteria and other prokaryotes, most of the DNA is found in a central region of the cell called the **nucleoid**, which functions similarly to a nucleus but is not surrounded by a membrane.

A cell's set of DNA is called its **genome**. Since all of the cells in an organism (with a few exceptions) contain the same DNA, you can also say that an organism has its own genome, and since the members of a species typically have similar genomes, you can also describe the genome of a species. In general, when people refer to the human genome, or any other eukaryotic genome, they mean the set of DNA found in the nucleus (that is, the nuclear genome). Mitochondria and chloroplasts are considered to have their own separate genomes.

## Chromosomes

Each species has its own characteristic number of chromosomes. Humans, for instance, have 46 chromosomes in a typical body cell, while dogs have 78. Like many species of animals and plants, humans are **diploid (2*n*)**, meaning that most of their chromosomes come in matched sets known as **homologous pairs**. Thus, the 46 chromosomes of a human cell are organized into 23 pairs, and the two members of each pair are said to be **homologues** of one another (with the slight exception of the X and Y chromosomes; see below).

Human sperm and eggs, which have only one homologous chromosome from each pair, are said to be **haploid (1*n*)**. When a sperm and egg fuse, their genetic material combines to form one complete, diploid set of chromosomes. So, for each homologous pair of chromosomes in your genome, one of the homologues comes from your mom and the other from your dad.




Image modified from "Karyotype," by the National Institutes of Health (public domain).

The two chromosomes in a homologous pair are generally very similar to one another. They're the same size and shape, and have the same pattern of light and dark bands, as you can see in the human **karyotype** (image of the chromosomes) shown above. Bands appear when the chromosomes are stained with a dye, and the dark bands mark more compacted DNA (usually, with fewer genes), while the light bands mark less compacted DNA (usually, with more genes). Most importantly, the two homologues in a pair carry the same type of genetic information. For instance, there is a gene found near the bottom of chromosome 15 that affects eye color[1]. A person might have the blue version, or **allele**, of this gene on one homologue, but the brown version on the other. Both homologues have the same type of gene in the same place, but they can (and often do!) have different versions of genes.

In humans, the X and Y chromosomes determine a person's biological sex, with XX for female and XY for male. While the two X chromosomes in a woman's cells are genuinely homologous, the X and Y chromosomes of a man's cells are not. They differ in size and shape, with the X being much larger than the Y, and contain different mostly different genes (although they do have small regions of similarity). The X and Y chromosomes are known as **sex chromosomes**, while the other 44 human chromosomes are called **autosomes**.

1. The Tech Museum of Innovation. (2013). Don't it make your brown eyes blue? In *Stanford at the tech: understanding genetics*. Retrieved from http://genetics.thetech.org/original_news/news39. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Chromosomes. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/cellular-molecular-biology/intro-to-cell-division/a/dna-and-chromosomes-article. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

10.3: DNA and Chromosomes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.4: Chromosome Structure

The continuity of life from one cell to another has its foundation in the reproduction of cells by way of the cell cycle. The **cell cycle** is an orderly sequence of events that describes the stages of a cell's life from the division of a single parent cell to the production of two new daughter cells. The mechanisms involved in the cell cycle are highly regulated.

## Eukaryotic Chromosomal Structure and Compaction

If the DNA from all 46 chromosomes in a human cell nucleus was laid out end to end, it would measure approximately two meters; however, its diameter would be only 2 nm. Considering that the size of a typical human cell is about 10 µm (100,000 cells lined up to equal one meter), DNA must be tightly packaged to fit in the cell's nucleus. At the same time, it must also be readily accessible for the genes to be expressed. During some stages of the cell cycle, the long strands of DNA are condensed into compact chromosomes. There are a number of ways that chromosomes are compacted.

In the first level of compaction, short stretches of the DNA double helix wrap around a core of eight histone proteins at regular intervals along the entire length of the chromosome (Figure 1). The DNA-histone complex is called chromatin. The beadlike, histone DNA complex is called a nucleosome, and DNA connecting the nucleosomes is called linker DNA. A DNA molecule in this form is about seven times shorter than the double helix without the histones, and the beads are about 10 nm in diameter, in contrast with the 2-nm diameter of a DNA double helix. The next level of compaction occurs as the nucleosomes and the linker DNA between them are coiled into a 30-nm chromatin fiber. This coiling further shortens the chromosome so that it is now about 50 times shorter than the extended form. In the third level of packing, a variety of fibrous proteins is used to pack the chromatin. These fibrous proteins also ensure that each chromosome in a non-dividing cell occupies a particular area of the nucleus that does not overlap with that of any other chromosome.






Figure 1. Double-stranded DNA wraps around histone proteins to form nucleosomes that have the appearance of "beads on a string." The nucleosomes are coiled into a 30-nm chromatin fiber. When a cell undergoes mitosis, the chromosomes condense even further.

DNA replicates in the S phase of interphase. After replication, the chromosomes are composed of two linked sister chromatids. When fully compact, the pairs of identically packed chromosomes are bound to each other by cohesin proteins. The connection between the sister chromatids is closest in a region called the centromere. The conjoined sister chromatids, with a diameter of about 1 µm, are visible under a light microscope. The centromeric region is highly condensed and thus will appear as a constricted area.

This animation illustrates the different levels of chromosome packing:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=320

## Learning Objectives

DNA in eukaryotes is highly structured and organized in all stages of an organisms life. Diploid organisms contain a pair of each chromosome; humans have 23 pairs for a total number of 46 chromosomes. Pairs of chromosomes, also known as homologous chromosomes, contain the same genes though there may be differences between the version of gene on each member of the pair. DNA is normally tightly packed into the nucleus of a eukaryotic cell, through protein-DNA complexes that form the characteristic condensed 'chromosome' shape. DNA compacts even further in preparation for cell division.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- How DNA is Packaged (Advanced). **Authored by**: DNA Learning Center. **Located at**: https://youtu.be/gbSIBhFwQ4s. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

10.4: Chromosome Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.5: Introduction to the Cell Cycle

## Identify the stages of the cell cycle, by picture and by description of major milestones

The **cell cycle** is an ordered series of events involving cell growth and cell division that produces two new daughter cells. Cells on the path to cell division proceed through a series of precisely timed and carefully regulated stages of growth, DNA replication, and division that produces two identical (clone) cells. The cell cycle has two major phases: interphase and the mitotic phase (Figure 1). During **interphase**, the cell grows and DNA is replicated. During the **mitotic phase**, the replicated DNA and cytoplasmic contents are separated, and the cell divides.





Figure 1. The cell cycle

## What You'll Learn to Do

- Identify the characteristics and sub-phases of interphase
- Identify the characteristics and stages of mitosis
- Identify the characteristics of cytokinesis

### Learning Activities

The learning activities for this section include the following:

- Interphase
- Mitosis
- Cytokinesis
- The Complete Cycle
- Self Check: The Cell Cycle

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

10.5: Introduction to the Cell Cycle is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.6: Interphase

During interphase, the cell undergoes normal growth processes while also preparing for cell division. In order for a cell to move from interphase into the mitotic phase, many internal and external conditions must be met. The three stages of interphase are called $G_1$, S, and $G_2$.

## $G_1$ Phase (First Gap)

The first stage of interphase is called the **$G_1$ phase** (first gap) because, from a microscopic aspect, little change is visible. However, during the $G_1$ stage, the cell is quite active at the biochemical level. The cell is accumulating the building blocks of chromosomal DNA and the associated proteins as well as accumulating sufficient energy reserves to complete the task of replicating each chromosome in the nucleus.

## S Phase (Synthesis of DNA)

Throughout interphase, nuclear DNA remains in a semi-condensed chromatin configuration. In the **S phase**, DNA replication can proceed through the mechanisms that result in the formation of identical pairs of DNA molecules—sister chromatids—that are firmly attached to the centromeric region. The centrosome is duplicated during the S phase. The two centrosomes will give rise to the **mitotic spindle**, the apparatus that orchestrates the movement of chromosomes during mitosis. At the center of each animal cell, the centrosomes of animal cells are associated with a pair of rod-like objects, the **centrioles**, which are at right angles to each other. Centrioles help organize cell division. Centrioles are not present in the centrosomes of other eukaryotic species, such as plants and most fungi.

## $G_2$ Phase (Second Gap)

In the **$G_2$ phase**, the cell replenishes its energy stores and synthesizes proteins necessary for chromosome manipulation. Some cell organelles are duplicated, and the cytoskeleton is dismantled to provide resources for the mitotic phase. There may be additional cell growth during $G_2$. The final preparations for the mitotic phase must be completed before the cell is able to enter the first stage of mitosis.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.6: Interphase is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.7: Mitosis

The mitotic phase (also known as M phase) is a multistep process during which the duplicated chromosomes are aligned, separated, and move into two new, identical daughter cells. The first portion of the mitotic phase is called **karyokinesis**, or nuclear division. The second portion of the mitotic phase, called cytokinesis, is the physical separation of the cytoplasmic components into the two daughter cells.

## Karyokinesis (Mitosis)

Karyokinesis, also known as mitosis, is divided into a series of phases—prophase, prometaphase, metaphase, anaphase, and telophase—that result in the division of the cell (Figure 1).





Figure 1. Stages of mitosis

During **prophase**, the "first phase," the nuclear envelope starts to dissociate into small vesicles, and the membranous organelles (such as the Golgi complex or Golgi apparatus, and endoplasmic reticulum), fragment and disperse toward the periphery of the cell. The nucleolus disappears (disperses). The centrosomes begin to move to opposite poles of the cell. Microtubules that will form the mitotic spindle extend between the centrosomes, pushing them farther apart as the microtubule fibers lengthen. The sister chromatids begin to coil more tightly with the aid of **condensin** proteins and become visible under a light microscope.





Figure 2. During prometaphase, mitotic spindle microtubules from opposite poles attach to each sister chromatid at the kinetochore. In anaphase, the connection between the sister chromatids breaks down, and the microtubules pull the chromosomes toward opposite poles.

During **prometaphase**, the "first change phase," many processes that were begun in prophase continue to advance. The remnants of the nuclear envelope fragment. The mitotic spindle continues to develop as more microtubules assemble and stretch across the length of the former nuclear area. Chromosomes become more condensed and discrete. Each sister chromatid develops a protein

structure called a kinetochore in the centromeric region (Figure 2). The proteins of the kinetochore attract and bind mitotic spindle microtubules. As the spindle microtubules extend from the centrosomes, some of these microtubules come into contact with and firmly bind to the kinetochores. Once a mitotic fiber attaches to a chromosome, the chromosome will be oriented until the kinetochores of sister chromatids face the opposite poles. Eventually, all the sister chromatids will be attached via their kinetochores to microtubules from opposing poles. Spindle microtubules that do not engage the chromosomes are called polar microtubules. These microtubules overlap each other midway between the two poles and contribute to cell elongation. Astral microtubules are located near the poles, aid in spindle orientation, and are required for the regulation of mitosis.

During **metaphase**, the "change phase," all the chromosomes are aligned in a plane called the **metaphase plate**, or the equatorial plane, midway between the two poles of the cell. The sister chromatids are still tightly attached to each other by cohesin proteins. At this time, the chromosomes are maximally condensed.

During **anaphase**, the "upward phase," the cohesin proteins degrade, and the sister chromatids separate at the centromere. Each chromatid, now called a chromosome, is pulled rapidly toward the centrosome to which its microtubule is attached. The cell becomes visibly elongated (oval shaped) as the polar microtubules slide against each other at the metaphase plate where they overlap.

During **telophase**, the "distance phase," the chromosomes reach the opposite poles and begin to decondense (unravel), relaxing into a chromatin configuration. The mitotic spindles are depolymerized into tubulin monomers that will be used to assemble cytoskeletal components for each daughter cell. Nuclear envelopes form around the chromosomes, and nucleosomes appear within the nuclear area.

The activity below will walk you through mitosis—providing you with the chance to review the different steps of the process and how they work together.

A link to an interactive elements can be found at the bottom of this page.

Click here for a text-only version of the activity.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Mitosis diagram. **Authored by**: Marek Kultys. **Located at**: https://commons.wikimedia.org/wiki/File:Mitosis_diagram.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*
- Mitosis. **Provided by**: Lumen Learning. **Located at**: www.oppia.org/explore/q4NBr1JG7Qrt. **License**: *CC BY-SA: Attribution-ShareAlike*

Public domain content

- Modification of Mitosis cells sequence. **Authored by**: LadyofHats. **Located at**: https://commons.wikimedia.org/wiki/File:Mitosis_cells_sequence.svg. **License**: *Public Domain: No Known Copyright*

---

10.7: Mitosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.8: Cytokinesis

**Cytokinesis**, or “cell motion,” is the second main stage of the mitotic phase during which cell division is completed via the physical separation of the cytoplasmic components into two daughter cells. Division is not complete until the cell components have been apportioned and completely separated into the two daughter cells. Although the stages of mitosis are similar for most eukaryotes, the process of cytokinesis is quite different for eukaryotes that have cell walls, such as plant cells.

In cells such as animal cells that lack cell walls, cytokinesis follows the onset of anaphase. A contractile ring composed of actin filaments forms just inside the plasma membrane at the former metaphase plate. The actin filaments pull the equator of the cell inward, forming a fissure. This fissure, or “crack,” is called the **cleavage furrow**. The furrow deepens as the actin ring contracts, and eventually the membrane is cleaved in two (Figure 1).





Figure 1. During cytokinesis in animal cells, a ring of actin filaments forms at the metaphase plate. The ring contracts, forming a cleavage furrow, which divides the cell in two. In plant cells, Golgi vesicles coalesce at the former metaphase plate, forming a phragmoplast. A cell plate formed by the fusion of the vesicles of the phragmoplast grows from the center toward the cell walls, and the membranes of the vesicles fuse to form a plasma membrane that divides the cell in two.

In plant cells, a new cell wall must form between the daughter cells. During interphase, the Golgi apparatus accumulates enzymes, structural proteins, and glucose molecules prior to breaking into vesicles and dispersing throughout the dividing cell. During telophase, these Golgi vesicles are transported on microtubules to form a phragmoplast (a vesicular structure) at the metaphase plate. There, the vesicles fuse and coalesce from the center toward the cell walls; this structure is called a **cell plate**. As more vesicles fuse, the cell plate enlarges until it merges with the cell walls at the periphery of the cell. Enzymes use the glucose that has accumulated between the membrane layers to build a new cell wall. The Golgi membranes become parts of the plasma membrane on either side of the new cell wall (Figure 1).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.8: Cytokinesis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.9: The Complete Cycle

Remember, mitosis is the process of cell division, but it's just a portion of the full cell cycle. Figure 1 shows approximately how long a cell spends in each stage of the cell cycle:





Figure 1. The cell cycle consists of interphase and the mitotic phase. During interphase, the cell grows and the nuclear DNA is duplicated. Interphase is followed by the mitotic phase. During the mitotic phase, the duplicated chromosomes are segregated and distributed into daughter nuclei. The cytoplasm is usually divided as well, resulting in two daughter cells.

As you can see, cells spend most of their time in interphase.

## Video Review: The Cell Cycle

This video reviews all the steps of mitosis; seeing it all together is a great review at this stage.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=330

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Mitosis: Splitting Up Is Complicated: Crash Course Biology #12. **Authored by**: CrashCourse. **Located at**: https://youtu.be/L0k-enzoeOM. **Project**: Crash Course Biology. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

10.9: The Complete Cycle is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.10: Introduction to Cell Cycle Checkpoints

## Identify and explain the important checkpoints that a cell passes through during the cell cycle

As we just learned, the cell cycle is a fairly complicated process. In order to make sure everything goes right, there are checkpoints in the cycle. Let's learn about these and how they help control the cell cycle.

## What You'll Learn to Do

- Identify important checkpoints in cell division
- Explain how errors in cell division are related to cancer

### Learning Activities

The learning activities for this section include the following:

- Control of the Cell Cycle
- Cancer and the Cell Cycle
- Self Check: Cell Cycle Checkpoints

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

10.10: Introduction to Cell Cycle Checkpoints is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.11: Control of the Cell Cycle

The length of the cell cycle is highly variable, even within the cells of a single organism. In humans, the frequency of cell turnover ranges from a few hours in early embryonic development, to an average of two to five days for epithelial cells, and to an entire human lifetime spent in $G_0$ by specialized cells, such as cortical neurons or cardiac muscle cells. There is also variation in the time that a cell spends in each phase of the cell cycle. When fast-dividing mammalian cells are grown in culture (outside the body under optimal growing conditions), the length of the cycle is about 24 hours. In rapidly dividing human cells with a 24-hour cell cycle, the $G_1$ phase lasts approximately nine hours, the S phase lasts 10 hours, the $G_2$ phase lasts about four and one-half hours, and the M phase lasts approximately one-half hour. In early embryos of fruit flies, the cell cycle is completed in about eight minutes. The timing of events in the cell cycle is controlled by mechanisms that are both internal and external to the cell.

## Regulation of the Cell Cycle by External Events

Both the initiation and inhibition of cell division are triggered by events external to the cell when it is about to begin the replication process. An event may be as simple as the death of a nearby cell or as sweeping as the release of growth-promoting hormones, such as human growth hormone (HGH). A lack of HGH can inhibit cell division, resulting in dwarfism, whereas too much HGH can result in gigantism. Crowding of cells can also inhibit cell division. Another factor that can initiate cell division is the size of the cell; as a cell grows, it becomes inefficient due to its decreasing surface-to-volume ratio. The solution to this problem is to divide.

Whatever the source of the message, the cell receives the signal, and a series of events within the cell allows it to proceed into interphase. Moving forward from this initiation point, every parameter required during each cell cycle phase must be met or the cycle cannot progress.

## Regulation at Internal Checkpoints

It is essential that the daughter cells produced be exact duplicates of the parent cell. Mistakes in the duplication or distribution of the chromosomes lead to mutations that may be passed forward to every new cell produced from an abnormal cell. To prevent a compromised cell from continuing to divide, there are internal control mechanisms that operate at three main cell cycle checkpoints. A checkpoint is one of several points in the eukaryotic cell cycle at which the progression of a cell to the next stage in the cycle can be halted until conditions are favorable. These checkpoints occur near the end of $G_1$, at the $G_2$/M transition, and during metaphase (Figure 1).





Figure 1. The cell cycle is controlled at three checkpoints. The integrity of the DNA is assessed at the $G_1$ checkpoint. Proper chromosome duplication is assessed at the $G_2$ checkpoint. Attachment of each kinetochore to a spindle fiber is assessed at the M checkpoint.

### $G_1$ Checkpoint

The $G_1$ checkpoint determines whether all conditions are favorable for cell division to proceed. The $G_1$ checkpoint, also called the restriction point (in yeast), is a point at which the cell irreversibly commits to the cell division process. External influences, such as growth factors, play a large role in carrying the cell past the $G_1$ checkpoint. In addition to adequate reserves and cell size, there is a check for genomic DNA damage at the $G_1$ checkpoint. A cell that does not meet all the requirements will not be allowed to progress into the S phase. The cell can halt the cycle and attempt to remedy the problematic condition, or the cell can advance into $G_0$ and await further signals when conditions improve.

### $G_2$ Checkpoint

The $G_2$ checkpoint bars entry into the mitotic phase if certain conditions are not met. As at the $G_1$ checkpoint, cell size and protein reserves are assessed. However, the most important role of the $G_2$ checkpoint is to ensure that all of the chromosomes have been replicated and that the replicated DNA is not damaged. If the checkpoint mechanisms detect problems with the DNA, the cell cycle is halted, and the cell attempts to either complete DNA replication or repair the damaged DNA.

### M Checkpoint

The M checkpoint occurs near the end of the metaphase stage of karyokinesis. The M checkpoint is also known as the spindle checkpoint, because it determines whether all the sister chromatids are correctly attached to the spindle microtubules. Because the separation of the sister chromatids during anaphase is an irreversible step, the cycle will not proceed until the kinetochores of each pair of sister chromatids are firmly anchored to at least two spindle fibers arising from opposite poles of the cell.

Watch what occurs at the $G_1$, $G_2$, and M checkpoints by downloading this animation of the cell cycle.

## Regulator Molecules of the Cell Cycle

In addition to the internally controlled checkpoints, there are two groups of intracellular molecules that regulate the cell cycle. These regulatory molecules either promote progress of the cell to the next phase (positive regulation) or halt the cycle (negative regulation). Regulator molecules may act individually, or they can influence the activity or production of other regulatory proteins. Therefore, the failure of a single regulator may have almost no effect on the cell cycle, especially if more than one mechanism

controls the same event. Conversely, the effect of a deficient or non-functioning regulator can be wide-ranging and possibly fatal to the cell if multiple processes are affected.

## Positive Regulation of the Cell Cycle

Two groups of proteins, called cyclins and cyclin-dependent kinases (Cdks), are responsible for the progress of the cell through the various checkpoints. The levels of the four cyclin proteins fluctuate throughout the cell cycle in a predictable pattern (Figure 2). Increases in the concentration of cyclin proteins are triggered by both external and internal signals. After the cell moves to the next stage of the cell cycle, the cyclins that were active in the previous stage are degraded.





Figure 2. The concentrations of cyclin proteins change throughout the cell cycle. There is a direct correlation between cyclin accumulation and the three major cell cycle checkpoints. Also note the sharp decline of cyclin levels following each checkpoint (the transition between phases of the cell cycle), as cyclin is degraded by cytoplasmic enzymes. (credit: modification of work by "WikiMiMa"/Wikimedia Commons)

Cyclins regulate the cell cycle only when they are tightly bound to Cdks. To be fully active, the Cdk/cyclin complex must also be phosphorylated in specific locations. Like all kinases, Cdks are enzymes (kinases) that phosphorylate other proteins. Phosphorylation activates the protein by changing its shape. The proteins phosphorylated by Cdks are involved in advancing the cell to the next phase (Figure 3). The levels of Cdk proteins are relatively stable throughout the cell cycle; however, the concentrations of cyclin fluctuate and determine when Cdk/cyclin complexes form. The different cyclins and Cdks bind at specific points in the cell cycle and thus regulate different checkpoints.



Figure 3. Cyclin-dependent kinases (Cdks) are protein kinases that, when fully activated, can phosphorylate and thus activate other proteins that advance the cell cycle past a checkpoint. To become fully activated, a Cdk must bind to a cyclin protein and then be phosphorylated by another kinase.

Since the cyclic fluctuations of cyclin levels are based on the timing of the cell cycle and not on specific events, regulation of the cell cycle usually occurs by either the Cdk molecules alone or the Cdk/cyclin complexes. Without a specific concentration of fully activated cyclin/Cdk complexes, the cell cycle cannot proceed through the checkpoints.

Although the cyclins are the main regulatory molecules that determine the forward momentum of the cell cycle, there are several other mechanisms that fine-tune the progress of the cycle with negative, rather than positive, effects. These mechanisms essentially block the progression of the cell cycle until problematic conditions are resolved. Molecules that prevent the full activation of Cdks are called Cdk inhibitors. Many of these inhibitor molecules directly or indirectly monitor a particular cell cycle event. The block placed on Cdks by inhibitor molecules will not be removed until the specific event that the inhibitor monitors is completed.

### Negative Regulation of the Cell Cycle

The second group of cell cycle regulatory molecules are negative regulators. Negative regulators halt the cell cycle. Remember that in positive regulation, active molecules cause the cycle to progress.

The best understood negative regulatory molecules are retinoblastoma protein (Rb), p53, and p21. Retinoblastoma proteins are a group of tumor-suppressor proteins common in many cells. The 53 and 21 designations refer to the functional molecular masses of the proteins (p) in kilodaltons. Much of what is known about cell cycle regulation comes from research conducted with cells that have lost regulatory control. All three of these regulatory proteins were discovered to be damaged or non-functional in cells that had begun to replicate uncontrollably (became cancerous). In each case, the main cause of the unchecked progress through the cell cycle was a faulty copy of the regulatory protein.

Rb, p53, and p21 act primarily at the $G_1$ checkpoint. p53 is a multi-functional protein that has a major impact on the commitment of a cell to division because it acts when there is damaged DNA in cells that are undergoing the preparatory processes during $G_1$. If damaged DNA is detected, p53 halts the cell cycle and recruits enzymes to repair the DNA. If the DNA cannot be repaired, p53 can trigger apoptosis, or cell suicide, to prevent the duplication of damaged chromosomes. As p53 levels rise, the production of p21 is triggered. p21 enforces the halt in the cycle dictated by p53 by binding to and inhibiting the activity of the Cdk/cyclin complexes. As a cell is exposed to more stress, higher levels of p53 and p21 accumulate, making it less likely that the cell will move into the S phase.

Rb exerts its regulatory influence on other positive regulator proteins. Chiefly, Rb monitors cell size. In the active, dephosphorylated state, Rb binds to proteins called transcription factors, most commonly, E2F (Figure 4). Transcription factors "turn on" specific genes, allowing the production of proteins encoded by that gene. When Rb is bound to E2F, production of proteins necessary for the $G_1$/S transition is blocked. As the cell increases in size, Rb is slowly phosphorylated until it becomes inactivated. Rb releases E2F, which can now turn on the gene that produces the transition protein, and this particular block is removed. For the cell to move past each of the checkpoints, all positive regulators must be "turned on," and all negative regulators must be "turned off."

Practice Question





Figure 4. Rb halts the cell cycle and releases its hold in response to cell growth.

Rb and other proteins that negatively regulate the cell cycle are sometimes called tumor suppressors. Why do you think the name tumor suppressor might be appropriate for these proteins?

[practice-area rows="2"][/practice-area]
[reveal-answer q="518676"]**Show Answer**[/reveal-answer]
[hidden-answer a="518676"]Rb and other negative regulatory proteins control cell division and therefore prevent the formation of tumors. Mutations that prevent these proteins from carrying out their function can result in cancer.[/hidden-answer]

## Learning Objectives

Each step of the cell cycle is monitored by internal controls called checkpoints. There are three major checkpoints in the cell cycle: one near the end of $G_1$, a second at the $G_2$/M transition, and the third during metaphase. Positive regulator molecules allow the cell cycle to advance to the next stage. Negative regulator molecules monitor cellular conditions and can halt the cycle until specific requirements are met.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.11: Control of the Cell Cycle is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.12: Cancer and the Cell Cycle

Cancer comprises many different diseases caused by a common mechanism: uncontrolled cell growth. Despite the redundancy and overlapping levels of cell cycle control, errors do occur. One of the critical processes monitored by the cell cycle checkpoint surveillance mechanism is the proper replication of DNA during the S phase. Even when all of the cell cycle controls are fully functional, a small percentage of replication errors (mutations) will be passed on to the daughter cells. If changes to the DNA nucleotide sequence occur within a coding portion of a gene and are not corrected, a gene mutation results. All cancers start when a gene mutation gives rise to a faulty protein that plays a key role in cell reproduction. The change in the cell that results from the malformed protein may be minor: perhaps a slight delay in the binding of Cdk to cyclin or an Rb protein that detaches from its target DNA while still phosphorylated. Even minor mistakes, however, may allow subsequent mistakes to occur more readily. Over and over, small uncorrected errors are passed from the parent cell to the daughter cells and amplified as each generation produces more non-functional proteins from uncorrected DNA damage. Eventually, the pace of the cell cycle speeds up as the effectiveness of the control and repair mechanisms decreases. Uncontrolled growth of the mutated cells outpaces the growth of normal cells in the area, and a tumor (*~oma*) can result.

## Proto-oncogenes

The genes that code for the positive cell cycle regulators are called **proto-oncogenes**. Proto-oncogenes are normal genes that, when mutated in certain ways, become **oncogenes**, genes that cause a cell to become cancerous. Consider what might happen to the cell cycle in a cell with a recently acquired oncogene. In most instances, the alteration of the DNA sequence will result in a less functional (or non-functional) protein. The result is detrimental to the cell and will likely prevent the cell from completing the cell cycle; however, the organism is not harmed because the mutation will not be carried forward. If a cell cannot reproduce, the mutation is not propagated and the damage is minimal. Occasionally, however, a gene mutation causes a change that increases the activity of a positive regulator. For example, a mutation that allows Cdk to be activated without being partnered with cyclin could push the cell cycle past a checkpoint before all of the required conditions are met. If the resulting daughter cells are too damaged to undergo further cell divisions, the mutation would not be propagated and no harm would come to the organism. However, if the atypical daughter cells are able to undergo further cell divisions, subsequent generations of cells will probably accumulate even more mutations, some possibly in additional genes that regulate the cell cycle.

The Cdk gene in the above example is only one of many genes that are considered proto-oncogenes. In addition to the cell cycle regulatory proteins, any protein that influences the cycle can be altered in such a way as to override cell cycle checkpoints. An oncogene is any gene that, when altered, leads to an increase in the rate of cell cycle progression.

## Tumor Suppressor Genes

Like proto-oncogenes, many of the negative cell cycle regulatory proteins were discovered in cells that had become cancerous. **Tumor suppressor genes** are segments of DNA that code for negative regulator proteins, the type of regulators that, when activated, can prevent the cell from undergoing uncontrolled division. The collective function of the best-understood tumor suppressor gene proteins, Rb, p53, and p21, is to put up a roadblock to cell cycle progression until certain events are completed. A cell that carries a mutated form of a negative regulator might not be able to halt the cell cycle if there is a problem. Tumor suppressors are similar to brakes in a vehicle: malfunctioning brakes can contribute to a car crash.

Mutated p53 genes have been identified in more than one-half of all human tumor cells. This discovery is not surprising in light of the multiple roles that the p53 protein plays at the G1 checkpoint. A cell with a faulty p53 may fail to detect errors present in the genomic DNA (Figure 1). Even if a partially functional p53 does identify the mutations, it may no longer be able to signal the necessary DNA repair enzymes. Either way, damaged DNA will remain uncorrected. At this point, a functional p53 will deem the cell unsalvageable and trigger programmed cell death (apoptosis). The damaged version of p53 found in cancer cells, however, cannot trigger apoptosis.






Figure 1. The role of normal p53 is to monitor DNA and the supply of oxygen (hypoxia is a condition of reduced oxygen supply). If damage is detected, p53 triggers repair mechanisms. If repairs are unsuccessful, p53 signals apoptosis. A cell with an abnormal p53 protein cannot repair damaged DNA and thus cannot signal apoptosis. Cells with abnormal p53 can become cancerous. (credit: modification of work by Thierry Soussi)

The loss of p53 function has other repercussions for the cell cycle. Mutated p53 might lose its ability to trigger p21 production. Without adequate levels of p21, there is no effective block on Cdk activation. Essentially, without a fully functional p53, the $G_1$ checkpoint is severely compromised and the cell proceeds directly from $G_1$ to S regardless of internal and external conditions. At the completion of this shortened cell cycle, two daughter cells are produced that have inherited the mutated p53 gene. Given the non-optimal conditions under which the parent cell reproduced, it is likely that the daughter cells will have acquired other mutations in addition to the faulty tumor suppressor gene. Cells such as these daughter cells quickly accumulate both oncogenes and non-functional tumor suppressor genes. Again, the result is tumor growth.

## Learning Objectives

This video reviews the ways that cancer is a by-product of broken DNA replication:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=336

Cancer is the result of unchecked cell division caused by a breakdown of the mechanisms that regulate the cell cycle. The loss of control begins with a change in the DNA sequence of a gene that codes for one of the regulatory molecules. Faulty instructions lead to a protein that does not function as it should. Any disruption of the monitoring system can allow other mistakes to be passed on to the daughter cells. Each successive cell division will give rise to daughter cells with even more accumulated damage. Eventually, all checkpoints become nonfunctional, and rapidly reproducing cells crowd out normal cells, resulting in a tumor or leukemia (blood cancer).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Cell cycle regulation, cancer, and stem cells. **Authored by**: Sal Khan. **Provided by**: Khan Academy. **Located at**: http://www.khanacademy.org/video?v=RZhL7LDPk8w. **Project**: Biology. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

---

10.12: Cancer and the Cell Cycle is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.13: Introduction to Meiosis

## Identify the stages of meiosis by picture and by description of major milestones; explain why meiosis involves two rounds of nuclear division

Just what is the difference between mitosis and meiosis? We know that mitosis produces autosomes; meiosis produces sex chromosomes. Let's learn just how similar and different these processes are.

## What You'll Learn to Do

- Identify the stages of meiosis
- Describe the steps of meiosis I
- Describe the steps of meiosis II

### Learning Activities

The learning activities for this section include the following:

- Stages of Meiosis
- Meiosis I
- Meiosis II
- Meiosis
- Self Check: Meiosis

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

10.13: Introduction to Meiosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.14: Stages of Meiosis

The ability to reproduce *in kind* is a basic characteristic of all living things. *In kind* means that the offspring of any organism closely resemble their parent or parents. Hippopotamuses give birth to hippopotamus calves, Joshua trees produce seeds from which Joshua tree seedlings emerge, and adult flamingos lay eggs that hatch into flamingo chicks. *In kind* does not generally mean *exactly the same*.



(a) (b) (c)

Figure 1. Each of us, like these other large multicellular organisms, begins life as a fertilized egg. After trillions of cell divisions, each of us develops into a complex, multicellular organism. (credit a: modification of work by Frank Wouters; credit b: modification of work by Ken Cole, USGS; credit c: modification of work by Martin Pettitt)

As you have learned, mitosis is the part of a cell reproduction cycle that results in identical daughter nuclei that are also genetically identical to the original parent nucleus. In mitosis, both the parent and the daughter nuclei are at the same ploidy level—diploid for most plants and animals. While many unicellular organisms and a few multicellular organisms can produce genetically identical clones of themselves through mitosis, many single-celled organisms and most multicellular organisms reproduce regularly using another method: **meiosis**. Sexual reproduction, specifically meiosis and fertilization, introduces variation into offspring that may account for the evolutionary success of sexual reproduction. The vast majority of eukaryotic organisms, both multicellular and unicellular, can or must employ some form of meiosis and fertilization to reproduce.

Meiosis employs many of the same mechanisms as mitosis. However, the starting nucleus is always diploid and the nuclei that result at the end of a meiotic cell division are haploid. To achieve this reduction in chromosome number, meiosis consists of one round of chromosome duplication and two rounds of nuclear division.

Because the events that occur during each of the division stages are analogous to the events of mitosis, the same stage names are assigned. However, because there are two rounds of division, the major process and the stages are designated with a “I” or a “II.” Thus, **meiosis I** is the first round of meiotic division and consists of prophase I, prometaphase I, and so on. **Meiosis II**, in which the second round of meiotic division takes place, includes prophase II, prometaphase II, and so on.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.14: Stages of Meiosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.15: Meiosis I

Meiosis is preceded by an interphase consisting of the $G_1$, S, and $G_2$ phases, which are nearly identical to the phases preceding mitosis. The $G_1$ phase, which is also called the first gap phase, is the first phase of the interphase and is focused on cell growth. The S phase is the second phase of interphase, during which the DNA of the chromosomes is replicated. Finally, the $G_2$ phase, also called the second gap phase, is the third and final phase of interphase; in this phase, the cell undergoes the final preparations for meiosis.

During DNA duplication in the S phase, each chromosome is replicated to produce two identical copies, called sister chromatids, that are held together at the centromere by **cohesin** proteins. Cohesin holds the chromatids together until anaphase II. The centrosomes, which are the structures that organize the microtubules of the meiotic spindle, also replicate. This prepares the cell to enter prophase I, the first meiotic phase.

## Prophase I

Early in prophase I, before the chromosomes can be seen clearly microscopically, the homologous chromosomes are attached at their tips to the nuclear envelope by proteins. As the nuclear envelope begins to break down, the proteins associated with homologous chromosomes bring the pair close to each other. (Recall that, in mitosis, homologous chromosomes do not pair together. In mitosis, homologous chromosomes line up end-to-end so that when they divide, each daughter cell receives a sister chromatid from both members of the homologous pair.) The **synaptonemal complex**, a lattice of proteins between the homologous chromosomes, first forms at specific locations and then spreads to cover the entire length of the chromosomes. The tight pairing of the homologous chromosomes is called **synapsis**. In synapsis, the genes on the chromatids of the homologous chromosomes are aligned precisely with each other. The synaptonemal complex supports the exchange of chromosomal segments between non-sister homologous chromatids, a process called crossing over. Crossing over can be observed visually after the exchange as **chiasmata** (singular = *chiasma*) (Figure 1).





Figure 1. Early in prophase I, homologous chromosomes come together to form a synapse. The chromosomes are bound tightly together and in perfect alignment by a protein lattice called a synaptonemal complex and by cohesin proteins at the centromere.

In species such as humans, even though the X and Y sex chromosomes are not homologous (most of their genes differ), they have a small region of homology that allows the X and Y chromosomes to pair up during prophase I. A partial synaptonemal complex develops only between the regions of homology.

Located at intervals along the synaptonemal complex are large protein assemblies called **recombination nodules**. These assemblies mark the points of later chiasmata and mediate the multistep process of **crossover**—or genetic recombination—between the non-sister chromatids. Near the recombination nodule on each chromatid, the double-stranded DNA is cleaved, the cut ends are modified, and a new connection is made between the non-sister chromatids. As prophase I progresses, the synaptonemal complex begins to break down and the chromosomes begin to condense. When the synaptonemal complex is gone, the homologous chromosomes remain attached to each other at the centromere and at chiasmata. The chiasmata remain until anaphase I. The number of chiasmata varies according to the species and the length of the chromosome. There must be at least one chiasma per chromosome for proper separation of homologous chromosomes during meiosis I, but there may be as many as 25. Following crossover, the synaptonemal complex breaks down and the cohesin connection between homologous pairs is also removed. At the

end of prophase I, the pairs are held together only at the chiasmata (Figure 2) and are called **tetrads** because the four sister chromatids of each pair of homologous chromosomes are now visible.





Figure 2. Crossover occurs between non-sister chromatids of homologous chromosomes. The result is an exchange of genetic material between homologous chromosomes.

The crossover events are the first source of genetic variation in the nuclei produced by meiosis. A single crossover event between homologous non-sister chromatids leads to a reciprocal exchange of equivalent DNA between a maternal chromosome and a paternal chromosome. Now, when that sister chromatid is moved into a gamete cell it will carry some DNA from one parent of the individual and some DNA from the other parent. The sister recombinant chromatid has a combination of maternal and paternal genes that did not exist before the crossover. Multiple crossovers in an arm of the chromosome have the same effect, exchanging segments of DNA to create recombinant chromosomes.

## Prometaphase I

The key event in prometaphase I is the attachment of the spindle fiber microtubules to the kinetochore proteins at the centromeres. Kinetochore proteins are multiprotein complexes that bind the centromeres of a chromosome to the microtubules of the mitotic spindle. Microtubules grow from centrosomes placed at opposite poles of the cell. The microtubules move toward the middle of the

cell and attach to one of the two fused homologous chromosomes. The microtubules attach at each chromosomes' kinetochores. With each member of the homologous pair attached to opposite poles of the cell, in the next phase, the microtubules can pull the homologous pair apart. A spindle fiber that has attached to a kinetochore is called a kinetochore microtubule. At the end of prometaphase I, each tetrad is attached to microtubules from both poles, with one homologous chromosome facing each pole. The homologous chromosomes are still held together at chiasmata. In addition, the nuclear membrane has broken down entirely.

## Metaphase I

During metaphase I, the homologous chromosomes are arranged in the center of the cell with the kinetochores facing opposite poles. The homologous pairs orient themselves randomly at the equator. For example, if the two homologous members of chromosome 1 are labeled a and b, then the chromosomes could line up a-b, or b-a. This is important in determining the genes carried by a gamete, as each will only receive one of the two homologous chromosomes. Recall that homologous chromosomes are not identical. They contain slight differences in their genetic information, causing each gamete to have a unique genetic makeup.

This randomness is the physical basis for the creation of the second form of genetic variation in offspring. Consider that the homologous chromosomes of a sexually reproducing organism are originally inherited as two separate sets, one from each parent. Using humans as an example, one set of 23 chromosomes is present in the egg donated by the mother. The father provides the other set of 23 chromosomes in the sperm that fertilizes the egg. Every cell of the multicellular offspring has copies of the original two sets of homologous chromosomes. In prophase I of meiosis, the homologous chromosomes form the tetrads. In metaphase I, these pairs line up at the midway point between the two poles of the cell to form the metaphase plate. Because there is an equal chance that a microtubule fiber will encounter a maternally or paternally inherited chromosome, the arrangement of the tetrads at the metaphase plate is random. Any maternally inherited chromosome may face either pole. Any paternally inherited chromosome may also face either pole. The orientation of each tetrad is independent of the orientation of the other 22 tetrads.

This event—the random (or independent) assortment of homologous chromosomes at the metaphase plate—is the second mechanism that introduces variation into the gametes or spores. In each cell that undergoes meiosis, the arrangement of the tetrads is different. The number of variations is dependent on the number of chromosomes making up a set. There are two possibilities for orientation at the metaphase plate; the possible number of alignments therefore equals $2^n$, where $n$ is the number of chromosomes per set. Humans have 23 chromosome pairs, which results in over eight million ($2^{23}$) possible genetically-distinct gametes. This number does not include the variability that was previously created in the sister chromatids by crossover. Given these two mechanisms, it is highly unlikely that any two haploid cells resulting from meiosis will have the same genetic composition (Figure 3).





Figure 3. Random, independent assortment during metaphase I can be demonstrated by considering a cell with a set of two chromosomes ($n$ = 2). In this case, there are two possible arrangements at the equatorial plane in metaphase I. The total possible number of different gametes is $2n$, where $n$ equals the number of chromosomes in a set. In this example, there are four possible genetic combinations for the gametes. With $n$ = 23 in human cells, there are over 8 million possible combinations of paternal and maternal chromosomes.

To summarize the genetic consequences of meiosis I, the maternal and paternal genes are recombined by crossover events that occur between each homologous pair during prophase I. In addition, the random assortment of tetrads on the metaphase plate produces a unique combination of maternal and paternal chromosomes that will make their way into the gametes.

## Anaphase I

In anaphase I, the microtubules pull the linked chromosomes apart. The sister chromatids remain tightly bound together a the centromere. The chiasmata are broken in anaphase I as the microtubules attached to the fused kinetochores pull the homologous chromosomes apart (Figure 4).





Figure 4. The process of chromosome alignment differs between meiosis I and meiosis II. In prometaphase I, microtubules attach to the fused kinetochores of homologous chromosomes, and the homologous chromosomes are arranged at the midpoint of the cell in metaphase I. In anaphase I, the homologous chromosomes are separated. In prometaphase II, microtubules attach to the kinetochores of sister chromatids, and the sister chromatids are arranged at the midpoint of the cells in metaphase II. In anaphase II, the sister chromatids are separated.

## Telophase I and Cytokinesis

In telophase, the separated chromosomes arrive at opposite poles. The remainder of the typical telophase events may or may not occur, depending on the species. In some organisms, the chromosomes decondense and nuclear envelopes form around the chromatids in telophase I. In other organisms, cytokinesis—the physical separation of the cytoplasmic components into two daughter cells—occurs without reformation of the nuclei. In nearly all species of animals and some fungi, cytokinesis separates the cell contents via a cleavage furrow (constriction of the actin ring that leads to cytoplasmic division). In plants, a cell plate is formed during cell cytokinesis by Golgi vesicles fusing at the metaphase plate. This cell plate will ultimately lead to the formation of cell walls that separate the two daughter cells.

Two haploid cells are the end result of the first meiotic division. The cells are haploid because at each pole, there is just one of each pair of the homologous chromosomes. Therefore, only one full set of the chromosomes is present. This is why the cells are considered haploid—there is only one chromosome set, even though each homolog still consists of two sister chromatids. Recall that sister chromatids are merely duplicates of one of the two homologous chromosomes (except for changes that occurred during crossing over). In meiosis II, these two sister chromatids will separate, creating four haploid daughter cells.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

10.15: Meiosis I is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.16: Meiosis II

In some species, cells enter a brief interphase, or **interkinesis**, before entering meiosis II. Interkinesis lacks an S phase, so chromosomes are not duplicated. The two cells produced in meiosis I go through the events of meiosis II in synchrony. During meiosis II, the sister chromatids within the two daughter cells separate, forming four new haploid gametes. The mechanics of meiosis II is similar to mitosis, except that each dividing cell has only one set of homologous chromosomes. Therefore, each cell has half the number of sister chromatids to separate out as a diploid cell undergoing mitosis.

## Prophase II

If the chromosomes decondensed in telophase I, they condense again. If nuclear envelopes were formed, they fragment into vesicles. The centrosomes that were duplicated during interkinesis move away from each other toward opposite poles, and new spindles are formed.

## Prometaphase II

The nuclear envelopes are completely broken down, and the spindle is fully formed. Each sister chromatid forms an individual kinetochore that attaches to microtubules from opposite poles.

## Metaphase II

The sister chromatids are maximally condensed and aligned at the equator of the cell.

## Anaphase II

The sister chromatids are pulled apart by the kinetochore microtubules and move toward opposite poles. Non-kinetochore microtubules elongate the cell.





Figure 1. The process of chromosome alignment differs between meiosis I and meiosis II. In prometaphase I, microtubules attach to the fused kinetochores of homologous chromosomes, and the homologous chromosomes are arranged at the midpoint of the cell in metaphase I. In anaphase I, the homologous chromosomes are separated. In prometaphase II, microtubules attach to the kinetochores of sister chromatids, and the sister chromatids are arranged at the midpoint of the cells in metaphase II. In anaphase II, the sister chromatids are separated.

## Telophase II and Cytokinesis

The chromosomes arrive at opposite poles and begin to decondense. Nuclear envelopes form around the chromosomes. Cytokinesis separates the two cells into four unique haploid cells. At this point, the newly formed nuclei are both haploid. The cells produced are genetically unique because of the random assortment of paternal and maternal homologs and because of the recombining of maternal and paternal segments of chromosomes (with their sets of genes) that occurs during crossover. The entire process of meiosis is outlined in Figure 2.





Figure 2. An animal cell with a diploid number of four ($2n = 4$) proceeds through the stages of meiosis to form four haploid daughter cells.

Review the process of meiosis, observing how chromosomes align and migrate, at Meiosis: An Interactive Animation.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.16: Meiosis II is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.17: Meiosis

This video walks you through meiosis I and meiosis II:




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=346

## Contributors and Attributions

All rights reserved content

- Meiosis: Where the Sex Starts - Crash Course Biology #13. **Authored by**: CrashCourse. **Located at**: https://youtu.be/qCLmR9-YY7o. **Project**: Crash Course Biology. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

10.17: Meiosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.18: Introduction to Genetic Diversity

## Describe and explain a range of mechanisms for generating genetic diversity

Now that we know how meiosis works, let’s see how it and its involved processes contribute to genetic diversity.

## What You’ll Learn to Do

- Understand how sexual reproduction leads to different sexual life cycles
- Understand how meiosis contributes to genetic diversity
- Understand the impact of translocation between non-homologous chromosomes during meiosis or mitosis
- Understand how mitosis, meiosis, and random fertilization all result in genetically unique individuals

### Learning Activities

The learning activities for this section include the following:

- Sexual Reproduction
- Genetic Variation in Meiosis
- Chromosomal Structural Rearrangements
- Mitosis, Meiosis, and Sexual Reproduction
- Self Check: Genetic Diversity

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

10.18: Introduction to Genetic Diversity is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.19: Sexual Reproduction

Sexual reproduction was an early evolutionary innovation after the appearance of eukaryotic cells. It appears to have been very successful because most eukaryotes are able to reproduce sexually, and in many animals, it is the only mode of reproduction. And yet, scientists recognize some real disadvantages to sexual reproduction. On the surface, creating offspring that are genetic clones of the parent appears to be a better system. If the parent organism is successfully occupying a habitat, offspring with the same traits would be similarly successful. There is also the obvious benefit to an organism that can produce offspring whenever circumstances are favorable by asexual budding, fragmentation, or asexual eggs. These methods of reproduction do not require another organism of the opposite sex. Indeed, some organisms that lead a solitary lifestyle have retained the ability to reproduce asexually. In addition, in asexual populations, every individual is capable of reproduction. In sexual populations, the males are not producing the offspring themselves, so in theory an asexual population could grow twice as fast.

However, multicellular organisms that exclusively depend on asexual reproduction are exceedingly rare. Why is sexuality (and meiosis) so common? This is one of the important unanswered questions in biology and has been the focus of much research beginning in the latter half of the twentieth century. There are several possible explanations, one of which is that the variation that sexual reproduction creates among offspring is very important to the survival and reproduction of the population. Thus, on average, a sexually reproducing population will leave more descendants than an otherwise similar asexually reproducing population. The only source of variation in asexual organisms is mutation. This is the ultimate source of variation in sexual organisms, but in addition, those different mutations are continually reshuffled from one generation to the next when different parents combine their unique genomes and the genes are mixed into different combinations by crossovers during prophase I and random assortment at metaphase I.

**Try It**

It is not in dispute that sexual reproduction provides evolutionary advantages to organisms that employ this mechanism to produce offspring. But why, even in the face of fairly stable conditions, does sexual reproduction persist when it is more difficult and costly for individual organisms? Variation is the outcome of sexual reproduction, but why are ongoing variations necessary? Enter the Red Queen hypothesis, first proposed by Leigh Van Valen in 1973. The concept was named in reference to the Red Queen's race in Lewis Carroll's book, *Through the Looking-Glass*.

All species co-evolve with other organisms; for example predators evolve with their prey, and parasites evolve with their hosts. Each tiny advantage gained by favorable variation gives a species an edge over close competitors, predators, parasites, or even prey. The only method that will allow a co-evolving species to maintain its own share of the resources is to also continually improve its fitness. As one species gains an advantage, this increases selection on the other species; they must also develop an advantage or they will be outcompeted. No single species progresses too far ahead because genetic variation among the progeny of sexual reproduction provides all species with a mechanism to improve rapidly. Species that cannot keep up become extinct. The Red Queen's catchphrase was, "It takes all the running you can do to stay in the same place." This is an apt description of co-evolution between competing species.

## Life Cycles of Sexually Reproducing Organisms

Fertilization and meiosis alternate in sexual **life cycles**. What happens between these two events depends on the organism. The process of meiosis reduces the chromosome number by half. Fertilization, the joining of two haploid gametes, restores the diploid condition. There are three main categories of life cycles in multicellular organisms: **diploid-dominant**, in which the multicellular diploid stage is the most obvious life stage, such as with most animals including humans; **haploid-dominant**, in which the multicellular haploid stage is the most obvious life stage, such as with all fungi and some algae; and **alternation of generations**, in which the two stages are apparent to different degrees depending on the group, as with plants and some algae.

### Diploid-Dominant Life Cycle

Nearly all animals employ a diploid-dominant life-cycle strategy in which the only haploid cells produced by the organism are the gametes. Early in the development of the embryo, specialized diploid cells, called **germ cells**, are produced within the gonads, such as the testes and ovaries. Germ cells are capable of mitosis to perpetuate the cell line and meiosis to produce gametes. Once the

haploid gametes are formed, they lose the ability to divide again. There is no multicellular haploid life stage. Fertilization occurs with the fusion of two gametes, usually from different individuals, restoring the diploid state (Figure 1).





Figure 1. In animals, sexually reproducing adults form haploid gametes from diploid germ cells. Fusion of the gametes gives rise to a fertilized egg cell, or zygote. The zygote will undergo multiple rounds of mitosis to produce a multicellular offspring. The germ cells are generated early in the development of the zygote.

## Haploid-Dominant Life Cycle

Most fungi and algae employ a life-cycle type in which the “body” of the organism—the ecologically important part of the life cycle—is haploid. The haploid cells that make up the tissues of the dominant multicellular stage are formed by mitosis. During sexual reproduction, specialized haploid cells from two individuals, designated the (+) and (−) mating types, join to form a diploid zygote. The zygote immediately undergoes meiosis to form four haploid cells called spores. Although haploid like the “parents,” these spores contain a new genetic combination from two parents. The spores can remain dormant for various time periods. Eventually, when conditions are conducive, the spores form multicellular haploid structures by many rounds of mitosis (Example 1).

**Practice Question**





Figure 2. Fungi, such as black bread mold (Rhizopus nigricans), have haploid-dominant life cycles. The haploid multicellular stage produces specialized haploid cells by mitosis that fuse to form a diploid zygote. The zygote undergoes meiosis to produce haploid spores. Each spore gives rise to a multicellular haploid organism by mitosis. (credit “zygomycota” micrograph: modification of work by “Fanaberka”/Wikimedia Commons)

If a mutation occurs so that a fungus is no longer able to produce a minus mating type, will it still be able to reproduce?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”139510″]**Show Answer**[/reveal-answer]
[hidden-answer a=”139510″]Most likely yes as the fungus can likely reproduce asexually.[/hidden-answer]

## Alternation of Generations

The third life-cycle type, employed by some algae and all plants, is a blend of the haploid-dominant and diploid-dominant extremes. Species with alternation of generations have both haploid and diploid multicellular organisms as part of their life cycle. The haploid multicellular plants are called **gametophytes**, because they produce gametes from specialized cells. Meiosis is not directly involved in the production of gametes in this case, because the organism that produces the gametes is already a haploid. Fertilization between the gametes forms a diploid zygote. The zygote will undergo many rounds of mitosis and give rise to a diploid multicellular plant called a **sporophyte**. Specialized cells of the sporophyte will undergo meiosis and produce haploid spores. The spores will subsequently develop into the gametophytes (Figure 3).





Figure 3. Plants have a life cycle that alternates between a multicellular haploid organism and a multicellular diploid organism. In some plants, such as ferns, both the haploid and diploid plant stages are free-living. The diploid plant is called a sporophyte because it produces haploid spores by meiosis. The spores develop into multicellular, haploid plants called gametophytes because they produce gametes. The gametes of two individuals will fuse to form a diploid zygote that becomes the sporophyte. (credit "fern": modification of work by Cory Zanker; credit "sporangia": modification of work by "Obsidian Soul"/Wikimedia Commons; credit "gametophyte and sporophyte": modification of work by "Vlmastra"/Wikimedia Commons)

Although all plants utilize some version of the alternation of generations, the relative size of the sporophyte and the gametophyte and the relationship between them vary greatly. In plants such as moss, the gametophyte organism is the free-living plant, and the sporophyte is physically dependent on the gametophyte. In other plants, such as ferns, both the gametophyte and sporophyte plants are free-living; however, the sporophyte is much larger. In seed plants, such as magnolia trees and daisies, the gametophyte is composed of only a few cells and, in the case of the female gametophyte, is completely retained within the sporophyte.

Sexual reproduction takes many forms in multicellular organisms. However, at some point in each type of life cycle, meiosis produces haploid cells that will fuse with the haploid cell of another organism. The mechanisms of variation—crossover, random assortment of homologous chromosomes, and random fertilization—are present in all versions of sexual reproduction. The fact that nearly every multicellular organism on Earth employs sexual reproduction is strong evidence for the benefits of producing offspring with unique gene combinations, though there are other possible benefits as well.

## Learning Objectives

Nearly all eukaryotes undergo sexual reproduction. The variation introduced into the reproductive cells by meiosis appears to be one of the advantages of sexual reproduction that has made it so successful. Meiosis and fertilization alternate in sexual life cycles. The process of meiosis produces unique reproductive cells called gametes, which have half the number of chromosomes as the parent cell. Fertilization, the fusion of haploid gametes from two individuals, restores the diploid condition. Thus, sexually reproducing organisms alternate between haploid and diploid stages. However, the ways in which reproductive cells are produced and the timing between meiosis and fertilization vary greatly. There are three main categories of life cycles: diploid-dominant, demonstrated by most animals; haploid-dominant, demonstrated by all fungi and some algae; and the alternation of generations, demonstrated by plants and some algae.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---


10.19: Sexual Reproduction is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.20: Genetic Variation in Meiosis

The gametes produced in meiosis aren't genetically identical to the starting cell, and they also aren't identical to one another. As an example, consider the meiosis II diagram above, which shows the end products of meiosis for a simple cell with a diploid number of $2n = 4$ chromosomes. The four gametes produced at the end of meiosis II are all slightly different, each with a unique combination of the genetic material present in the starting cell.

As it turns out, there are many more potential gamete types than just the four shown in the diagram, even for a simple cell with with only four chromosomes. This diversity of possible gametes reflects two factors: crossing over and the random orientation of homologue pairs during metaphase of meiosis I.

- **Crossing over.** The points where homologues cross over and exchange genetic material are chosen more or less at random, and they will be different in each cell that goes through meiosis. If meiosis happens many times, as it does in human ovaries and testes, crossovers will happen at many different points. This repetition produces a wide variety of recombinant chromosomes, chromosomes where fragments of DNA have been exchanged between homologues.
- **Random orientation of homologue pairs.** The random orientation of homologue pairs during metaphase of meiosis I is another important source of gamete diversity.




What exactly does *random orientation* mean here? Well, a homologous pair consists of one homologue from your dad and one from your mom, and you have 23 pairs of homologous chromosomes all together, counting the X and Y as homologous for this purpose. During meiosis I, the homologous pairs will separate to form two equal groups, but it's not usually the case that all the paternal—dad—chromosomes will go into one group and all the maternal—mom—chromosomes into the other.

Instead, each pair of homologues will effectively flip a coin to decide which chromosome goes into which group. In a cell with just two pairs of homologous chromosomes, like the one at right, random metaphase orientation allows for $2^2 = 4$ different types of possible gametes. In a human cell, the same mechanism allows for $2^{23} = 8{,}388{,}608$ different types of possible gametes[1]. And that's not even considering crossovers!

Given those kinds of numbers, it's very unlikely that any two sperm or egg cells made by a person will be the same. It's even more unlikely that you and your sister or brother will be genetically identical, unless you happen to be identical twins, thanks to the process of fertilization (in which a unique egg from Mom combines with a unique sperm from Dad, making a zygote whose genotype is well beyond one-in-a-trillion!)[2].

Meiosis and fertilization create genetic variation by making new combinations of gene variants (alleles). In some cases, these new combinations may make an organism more or less fit (able to survive and reproduce), thus providing the raw material for natural selection. Genetic variation is important in allowing a population to adapt via natural selection and thus survive in the long term.

---

1. Reece, J. B., L. A. Urry, M. L. Cain, S. A. Wasserman, P. V. Minorksy, and R. B. Jackson. "Genetic Variation Produced in Sexual Life Cycles Contributes to Evolution." In *Campbell Biology*, 263–65. 10th ed. San Francisco, CA: Pearson, 2011. ↵
2. Ibid. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Meiosis. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/cellular-molecular-biology/meiosis/a/phases-of-meiosis. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

---

10.20: Genetic Variation in Meiosis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.21: Chromosomal Structural Rearrangements

Cytologists have characterized numerous structural rearrangements in chromosomes, but chromosome inversions and translocations are the most common. Both are identified during meiosis by the adaptive pairing of rearranged chromosomes with their former homologs to maintain appropriate gene alignment. If the genes carried on two homologs are not oriented correctly, a recombination event could result in the loss of genes from one chromosome and the gain of genes on the other. This would produce aneuploid gametes.

## Chromosome Inversions

A **chromosome inversion** is the detachment, 180° rotation, and reinsertion of part of a chromosome. Inversions may occur in nature as a result of mechanical shear, or from the action of transposable elements (special DNA sequences capable of facilitating the rearrangement of chromosome segments with the help of enzymes that cut and paste DNA sequences). Unless they disrupt a gene sequence, inversions only change the orientation of genes and are likely to have more mild effects than aneuploid errors. However, altered gene orientation can result in functional changes because regulators of gene expression could be moved out of position with respect to their targets, causing aberrant levels of gene products.

An inversion can be **pericentric** and include the centromere, or **paracentric** and occur outside of the centromere (Figure 1). A pericentric inversion that is asymmetric about the centromere can change the relative lengths of the chromosome arms, making these inversions easily identifiable.





Figure 1. Pericentric inversions include the centromere, and paracentric inversions do not. A pericentric inversion can change the relative lengths of the chromosome arms; a paracentric inversion cannot.

When one homologous chromosome undergoes an inversion but the other does not, the individual is described as an inversion heterozygote. To maintain point-for-point synapsis during meiosis, one homolog must form a loop, and the other homolog must mold around it. Although this topology can ensure that the genes are correctly aligned, it also forces the homologs to stretch and can be associated with regions of imprecise synapsis (Figure 2).





Figure 2. When one chromosome undergoes an inversion but the other does not, one chromosome must form an inverted loop to retain point-for-point interaction during synapsis. This inversion pairing is essential to maintaining gene alignment during meiosis and to allow for recombination.

### The Chromosome 18 Inversion

Not all structural rearrangements of chromosomes produce nonviable, impaired, or infertile individuals. In rare instances, such a change can result in the evolution of a new species. In fact, a pericentric inversion in chromosome 18 appears to have contributed to the evolution of humans. This inversion is not present in our closest genetic relatives, the chimpanzees. Humans and chimpanzees differ cytogenetically by pericentric inversions on several chromosomes and by the fusion of two separate chromosomes in chimpanzees that correspond to chromosome two in humans.

The pericentric chromosome 18 inversion is believed to have occurred in early humans following their divergence from a common ancestor with chimpanzees approximately five million years ago. Researchers characterizing this inversion have suggested that approximately 19,000 nucleotide bases were duplicated on 18p, and the duplicated region inverted and reinserted on chromosome 18 of an ancestral human.

A comparison of human and chimpanzee genes in the region of this inversion indicates that two genes—*ROCK1* and *USP14*—that are adjacent on chimpanzee chromosome 17 (which corresponds to human chromosome 18) are more distantly positioned on human chromosome 18. This suggests that one of the inversion breakpoints occurred between these two genes. Interestingly, humans and chimpanzees express *USP14* at distinct levels in specific cell types, including cortical cells and fibroblasts. Perhaps the chromosome 18 inversion in an ancestral human repositioned specific genes and reset their expression levels in a useful way. Because both *ROCK1* and *USP14* encode cellular enzymes, a change in their expression could alter cellular function. It is not known how this inversion contributed to hominid evolution, but it appears to be a significant factor in the divergence of humans from other primates.

## Translocations

A **translocation** occurs when a segment of a chromosome dissociates and reattaches to a different, nonhomologous chromosome. Translocations can be benign or have devastating effects depending on how the positions of genes are altered with respect to regulatory sequences. Notably, specific translocations have been associated with several cancers and with schizophrenia. Reciprocal translocations result from the exchange of chromosome segments between two nonhomologous chromosomes such that there is no gain or loss of genetic information (Figure 3).





Figure 3. A reciprocal translocation occurs when a segment of DNA is transferred from one chromosome to another, nonhomologous chromosome. (credit: modification of work by National Human Genome Research/USA)

## Learning Objectives

The overall structure of chromosomes can be changed, in both minor and dramatic ways. Chromosome inversions involve the detachment, flipping, and reinsertion of a portion of a chromosome. Pericentric inversions involve the centromere while paracentric do not. Chromosome translocations involve the insertion of a piece of DNA from one chromosome into a non-homologous chromosome. Either of these types of structural changes may cause effects that range from minor to drastic.

## Contributors and Attributions


CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.21: Chromosomal Structural Rearrangements is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.22: Mitosis, Meiosis, and Sexual Reproduction

As you now know, genetic variation is very important. Genetic variation is introduced in multiple ways, including changes in mitosis, crossing over and random orientation in meiosis, and random fertilization. The video below offers you a nice overview of how each contributes to genetic diversity.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=356

## Contributors and Attributions

CC licensed content, Shared previously

- Mitosis, Meiosis and Sexual Reproduction. **Authored by**: Khan Academy. **Located at**: https://youtu.be/kaSIjIzAtYA. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

10.22: Mitosis, Meiosis, and Sexual Reproduction is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.23: Introduction to Errors in Chromosome Number

## Examine karyotypes and identify the effects of significant changes in chromosome number

We previously learned how errors in mitosis can potentially lead to cancer. What could errors in meiosis result in? In this outcome, we'll learn what happens when errors occur in chromosome number.

## What You'll Learn to Do

- Identify a karyotype and describe its uses in biology
- Identify common errors that can create an abnormal karyotype
- Identify syndromes that result from a significant change in chromosome number

### Learning Activities

The learning activities for this section include the following:

- Karyotypes
- Common Disorders
- Self Check: Errors in Chromosome Number

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

10.23: Introduction to Errors in Chromosome Number is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.24: Karyotypes

The isolation and microscopic observation of chromosomes forms the basis of cytogenetics and is the primary method by which clinicians detect chromosomal abnormalities in humans. A **karyotype** is the number and appearance of chromosomes, and includes their length, banding pattern, and centromere position. To obtain a view of an individual's karyotype, cytologists photograph the chromosomes and then cut and paste each chromosome into a chart, or **karyogram**, also known as an ideogram (Figure 1).





Figure 1. This karyotype is of a female human. Notice that homologous chromosomes are the same size, and have the same centromere positions and banding patterns. A human male would have an XY chromosome pair instead of the XX pair shown. (credit: Andreas Blozer et al)

In a given species, chromosomes can be identified by their number, size, centromere position, and banding pattern. In a human karyotype, **autosomes** or "body chromosomes" (all of the non–sex chromosomes) are generally organized in approximate order of size from largest (chromosome 1) to smallest (chromosome 22). The X and Y chromosomes are not autosomes. However, chromosome 21 is actually shorter than chromosome 22. This was discovered after the naming of Down syndrome as trisomy 21, reflecting how this disease results from possessing one extra chromosome 21 (three total). Not wanting to change the name of this important disease, chromosome 21 retained its numbering, despite describing the shortest set of chromosomes. The chromosome "arms" projecting from either end of the centromere may be designated as short or long, depending on their relative lengths. The short arm is abbreviated *p* (for "petite"), whereas the long arm is abbreviated *q* (because it follows "p" alphabetically). Each arm is further subdivided and denoted by a number. Using this naming system, locations on chromosomes can be described consistently in the scientific literature.

## Geneticists Use Karyograms to Identify Chromosomal Aberrations

Although Mendel is referred to as the "father of modern genetics," he performed his experiments with none of the tools that the geneticists of today routinely employ. One such powerful cytological technique is karyotyping, a method in which traits characterized by chromosomal abnormalities can be identified from a single cell. To observe an individual's karyotype, a person's cells (like white blood cells) are first collected from a blood sample or other tissue. In the laboratory, the isolated cells are stimulated to begin actively dividing. A chemical called colchicine is then applied to cells to arrest condensed chromosomes in metaphase. Cells are then made to swell using a hypotonic solution so the chromosomes spread apart. Finally, the sample is preserved in a fixative and applied to a slide.

The geneticist then stains chromosomes with one of several dyes to better visualize the distinct and reproducible banding patterns of each chromosome pair. Following staining, the chromosomes are viewed using bright-field microscopy. A common stain choice is the Giemsa stain. Giemsa staining results in approximately 400–800 bands (of tightly coiled DNA and condensed proteins) arranged along all of the 23 chromosome pairs; an experienced geneticist can identify each band. In addition to the banding patterns, chromosomes are further identified on the basis of size and centromere location. To obtain the classic depiction of the karyotype in which homologous pairs of chromosomes are aligned in numerical order from longest to shortest, the geneticist obtains a digital image, identifies each chromosome, and manually arranges the chromosomes into this pattern (Figure 1).

At its most basic, the karyogram may reveal genetic abnormalities in which an individual has too many or too few chromosomes per cell. Examples of this are Down Syndrome, which is identified by a third copy of chromosome 21, and Turner Syndrome, which is characterized by the presence of only one X chromosome in women instead of the normal two. Geneticists can also identify large deletions or insertions of DNA. For instance, Jacobsen Syndrome—which involves distinctive facial features as well as heart and bleeding defects—is identified by a deletion on chromosome 11. Finally, the karyotype can pinpoint translocations,

which occur when a segment of genetic material breaks from one chromosome and reattaches to another chromosome. Translocations are implicated in certain cancers, including chronic myelogenous leukemia.

During Mendel's lifetime, inheritance was an abstract concept that could only be inferred by performing crosses and observing the traits expressed by offspring. By observing a karyogram, today's geneticists can actually visualize the chromosomal composition of an individual to confirm or predict genetic abnormalities in offspring, even before birth.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

10.24: Karyotypes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.25: Common Disorders

Of all of the chromosomal disorders, abnormalities in chromosome number are the most obviously identifiable from a karyogram. Disorders of chromosome number include the duplication or loss of entire chromosomes, as well as changes in the number of complete sets of chromosomes. They are caused by nondisjunction, which occurs when pairs of homologous chromosomes or sister chromatids fail to separate during meiosis. Misaligned or incomplete synapsis, or a dysfunction of the spindle apparatus that facilitates chromosome migration, can cause nondisjunction. The risk of nondisjunction occurring increases with the age of the parents.

Nondisjunction can occur during either meiosis I or II, with differing results (Figure 1). If homologous chromosomes fail to separate during meiosis I, the result is two gametes that lack that particular chromosome and two gametes with two copies of the chromosome. If sister chromatids fail to separate during meiosis II, the result is one gamete that lacks that chromosome, two normal gametes with one copy of the chromosome, and one gamete with two copies of the chromosome.

Practice Question





Figure 1. Nondisjunction occurs when homologous chromosomes or sister chromatids fail to separate during meiosis, resulting in an abnormal chromosome number. Nondisjunction may occur during meiosis I or meiosis II.

Which of the following statements about nondisjunction is true?

a. Nondisjunction only results in gametes with n+1 or n–1 chromosomes.
b. Nondisjunction occurring during meiosis II results in 50 percent normal gametes.
c. Nondisjunction during meiosis I results in 50 percent normal gametes.
d. Nondisjunction always results in four different kinds of gametes.

[reveal-answer q="235653"]**Show Answer**[/reveal-answer]
[hidden-answer a="235653"]Answer b is true.[/hidden-answer]

## Aneuploidy





Figure 2. The incidence of having a fetus with trisomy 21 increases dramatically with maternal age.

An individual with the appropriate number of chromosomes for their species is called **euploid**; in humans, euploidy corresponds to 22 pairs of autosomes and one pair of sex chromosomes. An individual with an error in chromosome number is described as **aneuploid**, a term that includes **monosomy** (loss of one chromosome) or **trisomy** (gain of an extraneous chromosome). Monosomic human zygotes missing any one copy of an autosome invariably fail to develop to birth because they lack essential genes. This underscores the importance of "gene dosage" in humans. Most autosomal trisomies also fail to develop to birth; however, duplications of some of the smaller chromosomes (13, 15, 18, 21, or 22) can result in offspring that survive for several weeks to many years. Trisomic individuals suffer from a different type of genetic imbalance: an excess in gene dose. Individuals with an extra chromosome may synthesize an abundance of the gene products encoded by that chromosome. This extra dose (150 percent) of specific genes can lead to a number of functional challenges and often precludes development. The most common trisomy among viable births is that of chromosome 21, which corresponds to Down Syndrome. Individuals with this inherited disorder are characterized by short stature and stunted digits, facial distinctions that include a broad skull and large tongue, and significant developmental delays. The incidence of Down syndrome is correlated with maternal age; older women are more likely to become pregnant with fetuses carrying the trisomy 21 genotype (Figure 2).

## Polyploidy



Figure 3. As with many polyploid plants, this triploid orange daylily (*Hemerocallis fulva*) is particularly large and robust, and grows flowers with triple the number of petals of its diploid counterparts. (credit: Steve Karg)

An individual with more than the correct number of chromosome sets (two for diploid species) is called **polyploid**. For instance, fertilization of an abnormal diploid egg with a normal haploid sperm would yield a triploid zygote. Polyploid animals are extremely rare, with only a few examples among the flatworms, crustaceans, amphibians, fish, and lizards. Polyploid animals are sterile

because meiosis cannot proceed normally and instead produces mostly aneuploid daughter cells that cannot yield viable zygotes. Rarely, polyploid animals can reproduce asexually by haplodiploidy, in which an unfertilized egg divides mitotically to produce offspring. In contrast, polyploidy is very common in the plant kingdom, and polyploid plants tend to be larger and more robust than euploids of their species (Figure 3).

## Sex Chromosome Nondisjunction in Humans

Humans display dramatic deleterious effects with autosomal trisomies and monosomies. Therefore, it may seem counterintuitive that human females and males can function normally, despite carrying different numbers of the X chromosome. Rather than a gain or loss of autosomes, variations in the number of sex chromosomes are associated with relatively mild effects. In part, this occurs because of a molecular process called **X inactivation**. Early in development, when female mammalian embryos consist of just a few thousand cells (relative to trillions in the newborn), one X chromosome in each cell inactivates by tightly condensing into a quiescent (dormant) structure called a Barr body. The chance that an X chromosome (maternally or paternally derived) is inactivated in each cell is random, but once the inactivation occurs, all cells derived from that one will have the same inactive X chromosome or Barr body. By this process, females compensate for their double genetic dose of X chromosome.



Figure 4. In cats, the gene for coat color is located on the X chromosome. In the embryonic development of female cats, one of the two X chromosomes is randomly inactivated in each cell, resulting in a tortoiseshell pattern if the cat has two different alleles for coat color. Male cats, having only one X chromosome, never exhibit a tortoiseshell coat color. (credit: Michael Bodega)

In so-called “tortoiseshell” cats, embryonic X inactivation is observed as color variegation (Figure 4). Females that are heterozygous for an X-linked coat color gene will express one of two different coat colors over different regions of their body, corresponding to whichever X chromosome is inactivated in the embryonic cell progenitor of that region.

An individual carrying an abnormal number of X chromosomes will inactivate all but one X chromosome in each of her cells. However, even inactivated X chromosomes continue to express a few genes, and X chromosomes must reactivate for the proper maturation of female ovaries. As a result, X-chromosomal abnormalities are typically associated with mild mental and physical defects, as well as sterility. If the X chromosome is absent altogether, the individual will not develop in utero.

Several errors in sex chromosome number have been characterized. Individuals with three X chromosomes, called triplo-X, are phenotypically female but express developmental delays and reduced fertility. The XXY genotype, corresponding to one type of Klinefelter syndrome, corresponds to phenotypically male individuals with small testes, enlarged breasts, and reduced body hair. More complex types of Klinefelter syndrome exist in which the individual has as many as five X chromosomes. In all types, every X chromosome except one undergoes inactivation to compensate for the excess genetic dosage. This can be seen as several Barr bodies in each cell nucleus. Turner syndrome, characterized as an X0 genotype (i.e., only a single sex chromosome), corresponds to a phenotypically female individual with short stature, webbed skin in the neck region, hearing and cardiac impairments, and sterility.

## Duplications and Deletions

In addition to the loss or gain of an entire chromosome, a chromosomal segment may be duplicated or lost. Duplications and deletions often produce offspring that survive but exhibit physical and mental abnormalities. Duplicated chromosomal segments may fuse to existing chromosomes or may be free in the nucleus. Cri-du-chat (from the French for “cry of the cat”) is a syndrome associated with nervous system abnormalities and identifiable physical features that result from a deletion of most of 5p (the small arm of chromosome 5) (Figure 5). Infants with this genotype emit a characteristic high-pitched cry on which the disorder’s name is based.




Figure 5. This individual with cri-du-chat syndrome is shown at two, four, nine, and 12 years of age. (credit: Paola Cerruti Mainardi)

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

10.25: Common Disorders is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.26: Putting It Together- Cell Division

Mitosis and meiosis are both forms of division of the nucleus in eukaryotic cells. They share some similarities, but also exhibit distinct differences that lead to very different outcomes (Figure 1 and Table 1). Mitosis is a single nuclear division that results in two nuclei that are usually partitioned into two new cells. The nuclei resulting from a mitotic division are genetically identical to the original nucleus. They have the same number of sets of chromosomes, one set in the case of haploid cells and two sets in the case of diploid cells. In most plants and all animal species, it is typically diploid cells that undergo mitosis to form new diploid cells. In contrast, meiosis consists of two nuclear divisions resulting in four nuclei that are usually partitioned into four new cells. The nuclei resulting from meiosis are not genetically identical and they contain one chromosome set only. This is half the number of chromosome sets in the original cell, which is diploid.

The main differences between mitosis and meiosis occur in meiosis I, which is a very different nuclear division than mitosis. In meiosis I, the homologous chromosome pairs become associated with each other, are bound together with the synaptonemal complex, develop chiasmata and undergo crossover between sister chromatids, and line up along the metaphase plate in tetrads with kinetochore fibers from opposite spindle poles attached to each kinetochore of a homolog in a tetrad. All of these events occur only in meiosis I.





Figure 1. Meiosis and mitosis are both preceded by one round of DNA replication; however, meiosis includes two nuclear divisions. The four daughter cells resulting from meiosis are haploid and genetically distinct. The daughter cells resulting from mitosis are diploid and identical to the parent cell.

Table 1. Meiosis v. Mitosis

| | Meiosis | Mitosis |
|---|---|---|
| DNA Synthesis | Occurs in S phase of Interphase | Occurs in S phase of Interphase |
| Synapsis of homolgous chromosomes | During prophase I | Does not occur in mitosis |
| Crossover | During prophase I | Does not occur in mitosis |
| Homologous chromosomes line up at metaphase plate | During metaphase I | Does not occur in mitosis |
| Sister chromatids line up at metaphase plate | During metaphase II | During metaphase |
| Outcome: Number and genetic composition of daughter cells | Four haploid cells at the end of meiosis II | Two diploid cells at the end of mitosis |

## The Mystery of the Evolution of Meiosis

Some characteristics of organisms are so widespread and fundamental that it is sometimes difficult to remember that they evolved like other simpler traits. Meiosis is such an extraordinarily complex series of cellular events that biologists have had trouble hypothesizing and testing how it may have evolved. Although meiosis is inextricably entwined with sexual reproduction and its advantages and disadvantages, it is important to separate the questions of the evolution of meiosis and the evolution of sex, because early meiosis may have been advantageous for different reasons than it is now. Thinking outside the box and imagining what the early benefits from meiosis might have been is one approach to uncovering how it may have evolved.

Meiosis and mitosis share obvious cellular processes and it makes sense that meiosis evolved from mitosis. The difficulty lies in the clear differences between meiosis I and mitosis. Adam Wilkins and Robin Holliday[1] summarized the unique events that needed to occur for the evolution of meiosis from mitosis. These steps are homologous chromosome pairing, crossover exchanges, sister chromatids remaining attached during anaphase, and suppression of DNA replication in interphase. They argue that the first step is the hardest and most important, and that understanding how it evolved would make the evolutionary process clearer. They suggest genetic experiments that might shed light on the evolution of synapsis.

There are other approaches to understanding the evolution of meiosis in progress. Different forms of meiosis exist in single-celled protists. Some appear to be simpler or more “primitive” forms of meiosis. Comparing the meiotic divisions of different protists may shed light on the evolution of meiosis. Marilee Ramesh and colleagues[2] compared the genes involved in meiosis in protists to understand when and where meiosis might have evolved. Although research is still ongoing, recent scholarship into meiosis in protists suggests that some aspects of meiosis may have evolved later than others. This kind of genetic comparison can tell us what aspects of meiosis are the oldest and what cellular processes they may have borrowed from in earlier cells.

1. Adam S. Wilkins and Robin Holliday, “The Evolution of Meiosis from Mitosis,” *Genetics* 181 (2009): 3–12. ↵
2. Marilee A. Ramesh, Shehre-Banoo Malik and John M. Logsdon, Jr, “A Phylogenetic Inventory of Meiotic Genes: Evidence for Sex in *Giardia* and an Early Eukaryotic Origin of Meiosis,” *Current Biology* 15 (2005):185–91. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

10.26: Putting It Together- Cell Division is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 10.27: Assignment- Mitosis and Meiosis Worksheets

Use the two documents linked below to complete an internet hands-on activity involving mitosis and meiosis. During these activities you will demonstrate your understanding of cell division by identifying and drawing various stages of these events as well as answering questions about each.

## Download Worksheets

You may chose to print these sheets, write your answers out and then rescan the documents to create your single submission file. Or you may chose to type your answers into the Word document and submit that file. Do keep in mind that you must complete the drawing activities as well so in some fashion you have to get those images into your final file.

- Mitosis Worksheet
  - PDF
  - docx
- Meiosis Worksheet
  - PDF
  - docx

See this example for one way you may insert your images.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references
- Assignment includes answers to all questions and required images

## Rubric

| Mitosis and Meiosis Internet Quests | | | | |
|---|---|---|---|---|
| Outcome: Describe and explain the various stages of cell division | | | | |
| Criteria | Ratings | | | Pts |
| Identify the stages of the cell cycle, by picture and description of major milestones. | More than 22 questions answered correctly and all included images are correct/accurate.<br>5.0 pts | 20–22 mitosis questions answered correctly and 7 correct images included<br>4.0 pts | Less than 20 mitosis questions answered correctly and/or less than 7 correct images included<br>0.0 pts | 5 pts |
| Identify the stages of meiosis by picture and by description of major milestones; describe why meiosis involves two rounds of nuclear division. | More than 13 meiosis questions answered correctly.<br>5.0 pts | 12–13 meiosis questions answered correctly<br>4.0 pts | Less than 12 meiosis questions answered correctly<br>0.0 pts | 5 pts |
| | | | | Total points: 10 |

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Mitosis--Internet Lesson. **Provided by**: Biologycorner. **Located at**: https://www.biologycorner.com/worksheets/mitosis.html. **License**: *CC BY-NC: Attribution-NonCommercial*
- Meiosis--Internet Lesson. **Provided by**: Biologycorner. **Located at**: https://www.biologycorner.com//worksheets/meiosis_internet.html. **License**: *CC BY-NC: Attribution-NonCommercial*

---

10.27: Assignment- Mitosis and Meiosis Worksheets is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 11: Module 9- DNA Structure and Replication


11.1: Why It Matters- DNA Structure and Replication
11.2: Introduction to Storing Genetic Information
11.3: Structure of DNA
11.4: Genetic Information
11.5: DNA Packaging in Eukaryotes and Prokaryotes
11.6: Introduction to DNA Base Pairs and Replication
11.7: The History of DNA
11.8: Basics of DNA Replication
11.9: Major Enzymes
11.10: “Proofreading” DNA
11.11: Telomeres
11.12: Introduction to DNA Mutations
11.13: What is a Mutation?
11.14: Mutations in Somatic Cells and in Gametes
11.15: Major Types of Mutations
11.16: Putting It Together- DNA Structure and Replication
11.17: Assignment- Which Has More DNA?



11: Module 9- DNA Structure and Replication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.1: Why It Matters- DNA Structure and Replication

## Why relate DNA structure to the process of DNA replication?

You're one of a kind. It's not just your eyes, smile, and personality. Your health, risk for disease, and the ways you respond to medicines are also unique. Medicines that work well for some people may not help you at all. In fact, they might even cause problems. Wouldn't it be nice if treatments and preventive care could be designed just for you?

The careful matching of your biology to your medical care is known as personalized medicine. It's already being used by health care providers nationwide.



The story of personalized medicine begins with the unique DNA you inherited from your parents. DNA is responsible for all the physical traits that make you function as a human organism. It plays a vital role in life on this planet. DNA stores genetic information.

Genes are stretches of DNA that serve as a sort of instruction manual telling your body how to make the proteins and perform the other tasks that your body needs. The same genes often differ slightly between people. Bases may be switched, missing, or added here and there. Most of these variations have no effect on your health. But some can create unusual proteins that might boost your risk for certain diseases. Some variants can affect how well a medicine works in your body. Or they might cause a medicine to have different side effects in you than in someone else.

It's becoming more common for doctors to test for gene variants before prescribing certain drugs. "If doctors know your genes, they can predict drug response and incorporate this information into the medical decisions they make," says Dr. Rochelle Long, a pharmacogenomics expert at NIH. "By screening to know who shouldn't get certain drugs, we can prevent life-threatening side effects," Long says.

Even one of the oldest and most common drugs, aspirin, can have varying effects based on your genes. Millions of people take a daily aspirin to lower their risk for heart attack and stroke. Aspirin helps by preventing blood clots that could clog arteries. But aspirin doesn't reduce heart disease risk in everyone.

**Try It**

Here are just a few treatments that benefit from personalization:

- Blood clot treatments
- Colorectal cancer treatments
- Breast cancer treatments

So what does this mean for you? Is personalized medicine something you should look into? Let's learn more about DNA and genetics to see if we can answer this question.

### Learning Outcomes

- Explain how DNA stores genetic information
- Explain the role of complementary base pairing in the precise replication process of DNA

- Recognize the impact of DNA mutations

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- DNA. **Authored by**: Aneeque Ahmed. **Provided by**: The Noun Project. **Located at**: thenounproject.com/search/?q=DNA&i=783433. **License**: *CC BY: Attribution*

Public domain content

- Personalized Medicine: Matching Treatments to Your Genes. **Authored by**: Vicki Contie, Dana Steinberg, Carol Torgan, and Katherine Wendelsdorf. **Provided by**: NIH. **Located at**: https://newsinhealth.nih.gov/issue/dec2013/feature1. **License**: *Public Domain: No Known Copyright*

---

11.1: Why It Matters- DNA Structure and Replication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.2: Introduction to Storing Genetic Information

## Explain how DNA stores genetic information

In this outcome, you will learn to describe the double helix structure of DNA: its sugar-phosphate backbone ladder with nitrogenous base “rungs” of ladder.

## What You’ll Learn to Do

- Diagram the structure of DNA
- Relate the structure of DNA to the storage of genetic information
- Explain how DNA packaging in both eukaryotic and prokaryotic cells protects genetic information

### Learning Activities

The learning activities for this section include the following:

- Structure of DNA
- Genetic Information
- DNA Packaging in Eukaryotes and Prokaryotes
- Self Check: Storing Genetic Information

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

11.2: Introduction to Storing Genetic Information is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.3: Structure of DNA

The building blocks of DNA are nucleotides. The important components of each nucleotide are a nitrogenous base, deoxyribose (5-carbon sugar), and a phosphate group (see Figure 1). Each nucleotide is named depending on its nitrogenous base. The nitrogenous base can be a purine, such as adenine (A) and guanine (G), or a pyrimidine, such as cytosine (C) and thymine (T). Uracil (U) is also a pyrimidine (as seen in Figure 1), but it only occurs in RNA, which we will talk more about later.






Figure 1. Each nucleotide is made up of a sugar, a phosphate group, and a nitrogenous base. The sugar is deoxyribose in DNA and ribose in RNA.

Purines have a double ring structure with a six-membered ring fused to a five-membered ring. Pyrimidines are smaller in size; they have a single six-membered ring structure. The carbon atoms of the five-carbon sugar are numbered 1′, 2′, 3′, 4′, and 5′ (1′ is read as “one prime”). The nucleotides combine with each other by covalent bonds known as phosphodiester bonds or linkages. The phosphate residue is attached to the hydroxyl group of the 5′ carbon of one sugar of one nucleotide and the hydroxyl group of the 3′ carbon of the sugar of the next nucleotide, thereby forming a 5′-3′ phosphodiester bond.

In the 1950s, Francis Crick and James Watson worked together to determine the structure of DNA at the University of Cambridge, England. Other scientists like Linus Pauling and Maurice Wilkins were also actively exploring this field. Pauling had discovered the secondary structure of proteins using X-ray crystallography. In Wilkins’ lab, researcher Rosalind Franklin was using X-ray diffraction methods to understand the structure of DNA. Watson and Crick were able to piece together the puzzle of the DNA molecule on the basis of Franklin’s data because Crick had also studied X-ray diffraction (Figure 2). In 1962, James Watson, Francis Crick, and Maurice Wilkins were awarded the Nobel Prize in Medicine. Unfortunately, by then Franklin had died, and Nobel prizes are not awarded posthumously.



(a) (b)

Figure 2. The work of pioneering scientists (a) James Watson, Francis Crick, and Maclyn McCarty led to our present day understanding of DNA. Scientist Rosalind Franklin discovered (b) the X-ray diffraction pattern of DNA, which helped to elucidate its double helix structure. (credit a: modification of work by Marjorie McCarty, Public Library of Science)

Watson and Crick proposed that DNA is made up of two strands that are twisted around each other to form a right-handed helix. Base pairing takes place between a purine and pyrimidine; namely, A pairs with T and G pairs with C. Adenine and thymine are complementary base pairs, and cytosine and guanine are also complementary base pairs. The base pairs are stabilized by hydrogen bonds; adenine and thymine form two hydrogen bonds and cytosine and guanine form three hydrogen bonds. The two strands are anti-parallel in nature; that is, the 3′ end of one strand faces the 5′ end of the other strand. The sugar and phosphate of the nucleotides form the backbone of the structure, whereas the nitrogenous bases are stacked inside. Each base pair is separated from the other base pair by a distance of 0.34 nm, and each turn of the helix measures 3.4 nm. Therefore, ten base pairs are present per turn of the helix. The diameter of the DNA double helix is 2 nm, and it is uniform throughout. Only the pairing between a purine and pyrimidine can explain the uniform diameter. The twisting of the two strands around each other results in the formation of uniformly spaced major and minor grooves (Figure 3).





Figure 3. DNA has (a) a double helix structure and (b) phosphodiester bonds. The (c) major and minor grooves are binding sites for DNA binding proteins during processes such as transcription (the copying of RNA from DNA) and replication.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStaxCNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

11.3: Structure of DNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.4: Genetic Information

The genetic information of an organism is stored in DNA molecules. How can one kind of molecule contain all the instructions for making complicated living beings like ourselves? What component or feature of DNA can contain this information? It has to come from the nitrogen bases, because, as you already know, the backbone of all DNA molecules is the same. But there are only four bases found in DNA: G, A, C, and T. The sequence of these four bases can provide all the instructions needed to build any living organism. It might be hard to imagine that 4 different “letters” can communicate so much information. But think about the English language, which can represent a huge amount of information using just 26 letters. Even more profound is the binary code used to write computer programs. This code contains only ones and zeros, and think of all the things your computer can do. The DNA alphabet can encode very complex instructions using just four letters, though the messages end up being really long. For example, the *E. coli* bacterium carries its genetic instructions in a DNA molecule that contains more than five million nucleotides. The human genome (all the DNA of an organism) consists of around three billion **nucleotides** divided up between 23 paired DNA molecules, or **chromosomes**.

The information stored in the order of bases is organized into **genes**: each gene contains information for making a functional product. The genetic information is first copied to another **nucleic acid polymer**, **RNA** (ribonucleic acid), preserving the order of the nucleotide bases. Genes that contain instructions for making proteins are converted to messenger RNA (mRNA). Some specialized genes contain instructions for making functional RNA molecules that don’t make **proteins**. These RNA molecules function by affecting cellular processes directly; for example some of these RNA molecules regulate the expression of mRNA. Other genes produce RNA molecules that are required for **protein synthesis**, **transfer RNA** (**tRNA**), and **ribosomal RNA** (**rRNA**).

In order for DNA to function effectively at storing information, two key processes are required. First, information stored in the DNA molecule must be copied, with minimal errors, every time a cell divides. This ensures that both daughter cells inherit the complete set of genetic information from the parent cell. Second, the information stored in the DNA molecule must be **translated**, or expressed. In order for the stored information to be useful, cells must be able to access the instructions for making specific proteins, so the correct proteins are made in the right place at the right time.





Figure 1. DNA’s double helix. Graphic modified from “DNA chemical structure,” by Madeleine Price Ball, CC-BY-SA-2.0

Both copying and reading the information stored in DNA relies on base pairing between two **nucleic acid** polymer strands. Recall that DNA structure is a double helix (see Figure 1).

The sugar deoxyribose with the **phosphate group** forms the scaffold or backbone of the molecule (highlighted in yellow in Figure 1). Bases point inward. Complementary bases form hydrogen bonds with each other within the double helix. See how the bigger bases (**purines**) pair with the smaller ones (**pyrimidines**). This keeps the width of the double helix constant. More specifically, A pairs with T and C pairs with G. As we discuss the function of DNA in subsequent sections, keep in mind that there is a chemical reason for specific pairing of bases.

To illustrate the connection between information in DNA and an observable characteristic of an organism, let’s consider a gene that provides the instructions for building the hormone insulin. Insulin is responsible for regulating blood sugar levels. The insulin gene contains instructions for assembling the protein insulin from individual amino acids. Changing the sequence of nucleotides in the DNA molecule can change the amino acids in the final protein, leading to protein malfunction. If insulin does not function

correctly, it might be unable to bind to another protein (insulin receptor). On the organismal level of organization, this molecular event (change of DNA sequence) can lead to a disease state—in this case, diabetes.

## Practice Questions

The order of nucleotides in a gene (in DNA) is the key to how information is stored. For example, consider these two words: stable and tables. Both words are built from the same letters (subunits), but the different order of these subunits results in very different meanings. In DNA, the information is stored in units of 3 letters. Use the following key to decode the encrypted message. This should help you to see how information can be stored in the linear order of nucleotides in DNA.

| | | | |
|---|---|---|---|
| ABC = a | DEF = d | GHI = e | JKL = f |
| MNO = h | PQR = i | STU = m | VWX = n |
| YZA = o | BCD = r | EFG = s | HIJ = t |
| KLM = w | NOP = j | QRS = p | TUV = y |

**Encrypted Message:** HIJMNOPQREFG – PQREFG – MNOYZAKLM – DEFVWXABC – EFGHIJYZABCDGHIEFG – PQRVWXJKLYZABCDSTUABCHIJPQRYZAVWX

[practice-area rows="2"][/practice-area]
[reveal-answer q="236947"]**Show Answer**[/reveal-answer]
[hidden-answer a="236947"]This is how DNA stores information.

[/hidden-answer]

Where in the DNA is information stored?

a. The shape of the DNA
b. The sugar-phosphate backbone
c. The sequence of bases
d. The presence of two strands.

[reveal-answer q="767717"]Show Answer[/reveal-answer]
[hidden-answer a="767717"]Answer c. The sequence of the bases codes for the instructions for protein synthesis. The shape is DNA is not related to information storage. The sugar-phosphate backbone only acts as a scaffold. The presence of two strands is important for replication, but their information content is equivalent, as they are complementary to each other.

[/hidden-answer]

Which statement is correct?

a. The sequence of DNA bases is arranged into chromosomes, most of which contain the instructions to build an amino acid.
b. The sequence of DNA strands is arranged into chromosomes, most of which contain the instructions to build a protein.
c. The sequence of DNA bases is arranged into genes, most of which contain the instructions to build a protein.
d. The sequence of DNA phosphates is arranged into genes, most of which contain the instructions to build a cell.

[reveal-answer q="363484"]Show Answer[/reveal-answer]
[hidden-answer a="363484"]Answer c. **The sequence of DNA bases is arranged into genes, most of which contain the instructions to build a protein.** DNA stores information in the sequence of its bases. The information is grouped into genes. Protein is what is mainly coded.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Information Content of DNA. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5f7d80020ca600a650db8880db33. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

11.4: Genetic Information is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.5: DNA Packaging in Eukaryotes and Prokaryotes

When comparing prokaryotic cells to eukaryotic cells, prokaryotes are much simpler than eukaryotes in many of their features (Figure 1). Most prokaryotes contain a single, circular chromosome that is found in an area of the cytoplasm called the nucleoid.

## Practice Question





Figure 1. A eukaryote contains a well-defined nucleus, whereas in prokaryotes, the chromosome lies in the cytoplasm in an area called the nucleoid.

In eukaryotic cells, DNA and RNA synthesis occur in a separate compartment from protein synthesis. In prokaryotic cells, both processes occur together. What advantages might there be to separating the processes?

[practice-area rows="2"][/practice-area]
[reveal-answer q="306617"]**Show Answer**[/reveal-answer]
[hidden-answer a="306617"]What advantages might there be to having them occur together? Compartmentalization enables a eukaryotic cell to divide processes into discrete steps so it can build more complex protein and RNA products. But there is an advantage to having a single compartment as well: RNA and protein synthesis occurs much more quickly in a prokaryotic cell. [/hidden-answer]

The size of the genome in one of the most well-studied prokaryotes, *E.coli*, is 4.6 million base pairs (approximately 1.1 mm, if cut and stretched out). So how does this fit inside a small bacterial cell? The DNA is twisted by what is known as supercoiling. Supercoiling means that DNA is either under-wound (less than one turn of the helix per 10 base pairs) or over-wound (more than 1 turn per 10 base pairs) from its normal relaxed state. Some proteins are known to be involved in the supercoiling; other proteins and enzymes such as DNA gyrase help in maintaining the supercoiled structure.

Eukaryotes, whose chromosomes each consist of a linear DNA molecule, employ a different type of packing strategy to fit their DNA inside the nucleus (Figure 2). At the most basic level, DNA is wrapped around proteins known as histones to form structures called nucleosomes. The histones are evolutionarily conserved proteins that are rich in basic amino acids and form an octamer. The DNA (which is negatively charged because of the phosphate groups) is wrapped tightly around the histone core. This nucleosome is linked to the next one with the help of a linker DNA. This is also known as the "beads on a string" structure. This is further compacted into a 30 nm fiber, which is the diameter of the structure. At the metaphase stage, the chromosomes are at their most compact, are approximately 700 nm in width, and are found in association with scaffold proteins.

In interphase, eukaryotic chromosomes have two distinct regions that can be distinguished by staining. The tightly packaged region is known as heterochromatin, and the less dense region is known as euchromatin. Heterochromatin usually contains genes that are not expressed, and is found in the regions of the centromere and telomeres. The euchromatin usually contains genes that are transcribed, with DNA packaged around nucleosomes but not further compacted.





Figure 2. These figures illustrate the compaction of the eukaryotic chromosome.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

11.5: DNA Packaging in Eukaryotes and Prokaryotes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.6: Introduction to DNA Base Pairs and Replication

## Explain the role of complementary base pairing in the precise replication process of DNA

In this outcome, we'll learn more about the precise structure of DNA and how it replicates.

## What You'll Learn to Do

- Understand the historical basis of our understanding of DNA
- Outline the basic steps in DNA replication
- Identify the major enzymes that play a role in DNA replication
- Identify the key proofreading processes in DNA replication
- Understand the basic role of telomeres in protecting DNA from replication errors

### Learning Activities

The learning activities for this section include the following:

- The History of DNA
- Basics of DNA Replication
- Major Enzymes
- "Proofreading" DNA
- Telomeres
- Self Check: DNA Base Pairs and Replication

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

11.6: Introduction to DNA Base Pairs and Replication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.7: The History of DNA



Figure 1. Friedrich Miescher (1844–1895) discovered nucleic acids.

Modern understandings of DNA have evolved from the discovery of nucleic acid to the development of the double-helix model. In the 1860s, Friedrich Miescher (Figure 1), a physician by profession, was the first person to isolate phosphate-rich chemicals from white blood cells or leukocytes. He named these chemicals (which would eventually be known as RNA and DNA) nuclein because they were isolated from the nuclei of the cells.

To see Miescher conduct an experiment step-by-step, click through this review of how he discovered the key role of DNA and proteins in the nucleus.

A half century later, British bacteriologist Frederick Griffith was perhaps the first person to show that hereditary information could be transferred from one cell to another "horizontally," rather than by descent. In 1928, he reported the first demonstration of bacterial **transformation**, a process in which external DNA is taken up by a cell, thereby changing morphology and physiology. He was working with *Streptococcus pneumoniae*, the bacterium that causes pneumonia. Griffith worked with two strains, rough (R) and smooth (S). The R strain is non-pathogenic (does not cause disease) and is called rough because its outer surface is a cell wall and lacks a capsule; as a result, the cell surface appears uneven under the microscope. The S strain is pathogenic (disease-causing) and has a capsule outside its cell wall. As a result, it has a smooth appearance under the microscope. Griffith injected the live R strain into mice and they survived. In another experiment, when he injected mice with the heat-killed S strain, they also survived. In a third set of experiments, a mixture of live R strain and heat-killed S strain were injected into mice, and—to his surprise—the mice died. Upon isolating the live bacteria from the dead mouse, only the S strain of bacteria was recovered. When this isolated S strain was injected into fresh mice, the mice died. Griffith concluded that something had passed from the heat-killed S strain into the live R strain and transformed it into the pathogenic S strain, and he called this the transforming principle (Figure 2). These experiments are now famously known as Griffith's transformation experiments.



Mouse injected with heat-killed virulant S strain lives.



Mouse injected with both heat-killed S strain and live non-virulant R strain dies.

Figure 2. Two strains of *S. pneumoniae* were used in Griffith's transformation experiments. The R strain is non-pathogenic. The S strain is pathogenic and causes death. When Griffith injected a mouse with the heat-killed S strain and a live R strain, the mouse died. The S strain was recovered from the dead mouse. Thus, Griffith concluded that something had passed from the heat-killed S strain to the R strain, transforming the R strain into S strain in the process. (credit "living mouse": modification of work by NIH; credit "dead mouse": modification of work by Sarah Marriage)

Scientists Oswald Avery, Colin MacLeod, and Maclyn McCarty (1944) were interested in exploring this transforming principle further. They isolated the S strain from the dead mice and isolated the proteins and nucleic acids, namely RNA and DNA, as these were possible candidates for the molecule of heredity. They conducted a systematic elimination study. They used enzymes that specifically degraded each component and then used each mixture separately to transform the R strain. They found that when DNA was degraded, the resulting mixture was no longer able to transform the bacteria, whereas all of the other combinations were able to transform the bacteria. This led them to conclude that DNA was the transforming principle.

## Forensic Scientists and DNA Analysis

DNA evidence was used for the first time to solve an immigration case. The story started with a teenage boy returning to London from Ghana to be with his mother. Immigration authorities at the airport were suspicious of him, thinking that he was traveling on a forged passport. After much persuasion, he was allowed to go live with his mother, but the immigration authorities did not drop the case against him. All types of evidence, including photographs, were provided to the authorities, but deportation proceedings were started nevertheless. Around the same time, Dr. Alec Jeffreys of Leicester University in the United Kingdom had invented a technique known as DNA fingerprinting. The immigration authorities approached Dr. Jeffreys for help. He took DNA samples from the mother and three of her children, plus an unrelated mother, and compared the samples with the boy's DNA. Because the biological father was not in the picture, DNA from the three children was compared with the boy's DNA. He found a match in the boy's DNA for both the mother and his three siblings. He concluded that the boy was indeed the mother's son.

Forensic scientists analyze many items, including documents, handwriting, firearms, and biological samples. They analyze the DNA content of hair, semen, saliva, and blood, and compare it with a database of DNA profiles of known criminals. Analysis includes DNA isolation, sequencing, and sequence analysis; most forensic DNA analysis involves polymerase chain reaction (PCR) amplification of short tandem repeat (STR) loci and electrophoresis to determine the length of the PCR-amplified fragment. Only mitochondrial DNA is sequenced for forensics. Forensic scientists are expected to appear at court hearings to present their findings. They are usually employed in crime labs of city and state government agencies. Geneticists experimenting with DNA techniques also work for scientific and research organizations, pharmaceutical industries, and college and university labs. Students wishing to pursue a career as a forensic scientist should have at least a bachelor's degree in chemistry, biology, or physics, and preferably some experience working in a laboratory.

Experiments conducted by Martha Chase and Alfred Hershey in 1952 provided confirmatory evidence that DNA was the genetic material and not proteins. Chase and Hershey were studying a bacteriophage, which is a virus that infects bacteria. Viruses typically have a simple structure: a protein coat, called the capsid, and a nucleic acid core that contains the genetic material, either DNA or RNA. The bacteriophage infects the host bacterial cell by attaching to its surface, and then it injects its nucleic acids inside the cell. The phage DNA makes multiple copies of itself using the host machinery, and eventually the host cell bursts, releasing a large number of bacteriophages. Hershey and Chase labeled one batch of phage with radioactive sulfur, $^{35}S$, to label the protein coat. Another batch of phage were labeled with radioactive phosphorus, $^{32}P$. Because phosphorous is found in DNA, but not protein, the DNA and not the protein would be tagged with radioactive phosphorus.

Each batch of phage was allowed to infect the cells separately. After infection, the phage bacterial suspension was put in a blender, which caused the phage coat to be detached from the host cell. The phage and bacterial suspension was spun down in a centrifuge. The heavier bacterial cells settled down and formed a pellet, whereas the lighter phage particles stayed in the supernatant. In the tube that contained phage labeled with $^{35}S$, the supernatant contained the radioactively labeled phage, whereas no radioactivity was detected in the pellet. In the tube that contained the phage labeled with $^{32}P$, the radioactivity was detected in the pellet that contained the heavier bacterial cells, and no radioactivity was detected in the supernatant. Hershey and Chase concluded that it was the phage DNA that was injected into the cell and carried information to produce more phage particles, thus providing evidence that DNA was the genetic material and not proteins (Figure 3).





Figure 3. In Hershey and Chase's experiments, bacteria were infected with phage radiolabeled with either $^{35}S$, which labels protein, or $^{32}P$, which labels DNA. Only $^{32}P$ entered the bacterial cells, indicating that DNA is the genetic material.

Around this same time, Austrian biochemist Erwin Chargaff examined the content of DNA in different species and found that the amounts of adenine, thymine, guanine, and cytosine were not found in equal quantities, and that it varied from species to species, but not between individuals of the same species. He found that the amount of adenine equals the amount of thymine, and the amount of cytosine equals the amount of guanine, or A = T and G = C. This is also known as Chargaff's rules. This finding proved immensely useful when Watson and Crick were getting ready to propose their DNA double helix model.

## Practice Question

The experiments by Hershey and Chase helped confirm that DNA was the hereditary material on the basis of the finding that:

a. radioactive phage were found in the pellet
b. radioactive cells were found in the supernatant
c. radioactive sulfur was found inside the cell
d. radioactive phosphorus was found in the cell

[reveal-answer q="549343"]Show Answer[/reveal-answer]
[hidden-answer a="549343"]d. radioactive phosphorus was found in the cell[/hidden-answer]

## In Summary: The History of DNA

DNA was first isolated from white blood cells by Friedrich Miescher, who called it nuclein because it was isolated from nuclei. Frederick Griffith's experiments with strains of *Streptococcus pneumoniae* provided the first hint that DNA may be the transforming principle. Avery, MacLeod, and McCarty proved that DNA is required for the transformation of bacteria. Later experiments by Hershey and Chase using bacteriophage T2 proved that DNA is the genetic material. Chargaff found that the ratio of A = T and C = G, and that the percentage content of A, T, G, and C is different for different species.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

11.7: The History of DNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.8: Basics of DNA Replication





Figure 1. The three suggested models of DNA replication. Grey indicates the original DNA strands, and blue indicates newly synthesized DNA.

The elucidation of the structure of the double helix provided a hint as to how DNA divides and makes copies of itself. This model suggests that the two strands of the double helix separate during replication, and each strand serves as a template from which the new complementary strand is copied. What was not clear was how the replication took place. There were three models suggested: conservative, semi-conservative, and dispersive (see Figure 1).

In conservative replication, the parental DNA remains together, and the newly formed daughter strands are together. The semi-conservative method suggests that each of the two parental DNA strands act as a template for new DNA to be synthesized; after replication, each double-stranded DNA includes one parental or “old” strand and one “new” strand. In the dispersive model, both copies of DNA have double-stranded segments of parental DNA and newly synthesized DNA interspersed.

Meselson and Stahl were interested in understanding how DNA replicates. They grew *E. coli* for several generations in a medium containing a “heavy” isotope of nitrogen ($^{15}N$) that gets incorporated into nitrogenous bases, and eventually into the DNA (Figure 2).





Figure 2. Meselson and Stahl experimented with E. coli grown first in heavy nitrogen ($^{15}N$) then in $^{14}N$. DNA grown in $^{15}N$ (red band) is heavier than DNA grown in $^{14}N$ (orange band), and sediments to a lower level in cesium chloride solution in an ultracentrifuge. When DNA grown in $^{15}N$ is switched to media containing $^{14}N$, after one round of cell division the DNA sediments halfway between the $^{15}N$ and $^{14}N$ levels, indicating that it now contains fifty percent $^{14}N$. In subsequent cell divisions, an increasing amount of DNA contains $^{14}N$ only. This data supports the semi-conservative replication model. (credit: modification of work by Mariana Ruiz Villareal)

The *E. coli* culture was then shifted into medium containing $^{14}N$ and allowed to grow for one generation. The cells were harvested and the DNA was isolated. The DNA was centrifuged at high speeds in an ultracentrifuge. Some cells were allowed to grow for one more life cycle in $^{14}N$ and spun again. During the density gradient centrifugation, the DNA is loaded into a gradient (typically a salt such as cesium chloride or sucrose) and spun at high speeds of 50,000 to 60,000 rpm. Under these circumstances, the DNA will form a band according to its density in the gradient. DNA grown in $^{15}N$ will band at a higher density position than that grown in $^{14}N$. Meselson and Stahl noted that after one generation of growth in $^{14}N$ after they had been shifted from $^{15}N$, the single band observed was intermediate in position in between DNA of cells grown exclusively in $^{15}N$ and $^{14}N$. This suggested either a semi-conservative or dispersive mode of replication. The DNA harvested from cells grown for two generations in $^{14}N$ formed two bands: one DNA band was at the intermediate position between $^{15}N$ and $^{14}N$, and the other corresponded to the band of $^{14}N$ DNA. These results could only be explained if DNA replicates in a semi-conservative manner. Therefore, the other two modes were ruled out.

During DNA replication, each of the two strands that make up the double helix serves as a template from which new strands are copied. The new strand will be complementary to the parental or “old” strand. When two daughter DNA copies are formed, they have the same sequence and are divided equally into the two daughter cells.

Click through this tutorial on DNA replication.

## In Summary: Basics of DNA Replication

The model for DNA replication suggests that the two strands of the double helix separate during replication, and each strand serves as a template from which the new complementary strand is copied. In conservative replication, the parental DNA is conserved, and the daughter DNA is newly synthesized. The semi-conservative method suggests that each of the two parental DNA strands acts as template for new DNA to be synthesized; after replication, each double-stranded DNA includes one parental or “old” strand and

one “new” strand. The dispersive mode suggested that the two copies of the DNA would have segments of parental DNA and newly synthesized DNA. Experimental evidence showed DNA replication is semi-conservative.

## Contributors and Attributions


CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

11.8: Basics of DNA Replication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 11.9: Major Enzymes

The process of **DNA replication** is catalyzed by a type of enzyme called **DNA polymerase** (*poly* meaning many, *mer* meaning pieces, and *–ase* meaning enzyme; so an enzyme that attaches many pieces of DNA). Observe Figure 1: the double helix of the original DNA molecule separates (blue) and new strands are made to match the separated strands. The result will be two DNA molecules, each containing an old and a new strand. Therefore, DNA replication is called semiconservative. The term *semiconservative* refers to the fact that half of the original molecule (one of the two strands in the double helix) is “conserved” in the new molecule. The original strand is referred to as the *template strand* because it provides the information, or template, for the newly synthesized strand.



3′ synthesis shown, no enzymes in diagram. ” width=”1024″ height=”508″> Figure 1. By Madprime(Wikipedia) (DNA replication split horizontal) CC BY-SA 2.0





Figure 2. Primer and Template

**DNA replication** relies on the double-stranded nature of the molecule. One double stranded DNA molecule, when replicated, will become two double-stranded molecules, each containing one original strand and one newly synthesized strand. You remember that the two strands of DNA run antiparallel: one from the 5′ to the 3′, and the other from the 3′ to the 5′. The synthesis of the new DNA strand can only happen in one direction: from the 5′ to the 3′ end. In other words, the new bases are always added to the 3′ end of the newly synthesized DNA strand. So if the new nucleotide is always added to the 3′ end of an existing nucleotide, where does the *first* nucleotide come from? In fact, **DNA polymerase** needs an “anchor” to start adding nucleotides: a short sequence of DNA or RNA that is complementary to the template strand will work to provide a free 3′ end. This sequence is called a *primer* (Figure 2).

How does **DNA polymerase** know in what order to add nucleotides? Specific base pairing in DNA is the key to copying the DNA: if you know the sequence of one strand, you can use base pairing rules to build the other strand. Bases form pairs (base pairs) in a very specific way. Figure 3 shows how *A* (adenine) pairs with *T* (thymine) and *G* (guanine) pairs with *C* (cytosine). It is important to remember that this binding is specific: *T* pairs with *A*, but not with *C*. The molecular recognition occurs because of the ability of bases to form specific hydrogen bonds: atoms align just right to make hydrogen bonds possible. Also note that a larger base (purine, *A* or *G*) always pairs with a smaller base (pyrimidine, *C* or *T*).





Figure 3. DNA chemical structure. Modification of DNA chemical structure by Madeleine Price Ball; CC-BY-SA-2.0

### Practice Questions

True/False: DNA replication requires an enzyme.

[practice-area rows=”1″][/practice-area]
[reveal-answer q=”527189″]Show Answer[/reveal-answer]
[hidden-answer a=”527189″]True. Most biological reactions rely on the enzyme to speed up the reaction. In the case of DNA replication, this enzyme is DNA polymerase.

[/hidden-answer]

What are the building blocks on DNA?

a. Deoxyribonucleotides
b. Fatty acids
c. Ribonucleotides
d. Amino acids

[reveal-answer q=”93495″]Show Answer[/reveal-answer]
[hidden-answer a=”93495″]Answer a. DNA is a double helix made up of two long chains of deoxyribonucleotides.

[/hidden-answer]

True/False: DNA replication requires energy.

[practice-area rows="1"][/practice-area]
[reveal-answer q="62103"]Show Answer[/reveal-answer]
[hidden-answer a="62103"]True. Making large molecules from small subunits (anabolism) requires energy. What supplies the energy? The building blocks themselves serve as a source of energy. As they get incorporated into the DNA polymer, two phosphate groups are broken off to release energy, some of which is used for making the polymer. Deoxyribonucleotides differ from nucleotides like ATP only by one missing oxygen atom.

[/hidden-answer]

We have the building blocks, a source of energy, and a catalyst. What's missing? We need instruction about the order of nucleotides in the new polymer. Which molecule provides these instructions?

a. Protein
b. DNA
c. Carbohydrate
d. Lipid

[reveal-answer q="494506"]Show Answer[/reveal-answer]
[hidden-answer a="494506"]Answer b. We refer to this DNA as a template. The original information stored in the order of bases will direct the synthesis of the new DNA via base pairing.

[/hidden-answer]

There is one more thing required by the DNA polymerase. It cannot just start making a DNA copy of the template strand; it needs a short piece of DNA or RNA with a free hydroxyl group in the right place to attach the nucleotides to. (Remember that synthesis always occurs in one direction—new building blocks are added to the 3′ end.) This component starts the process by giving DNA polymerase something to bind to. What might you call this short piece of nucleic acid?

a. A solvent
b. A primer
c. A converter
d. A sealant

[reveal-answer q="529681"]Show Answer[/reveal-answer]
[hidden-answer a="529681"]Answer b. A primer is used to start this process by giving DNA polymerase something to bind the new nucleotide to.[/hidden-answer]

Now that you understand the basics of **DNA replication**, we can add a bit of complexity. The two strands of DNA have to be temporarily separated from each other; this job is done by a special enzyme, *helicase*, that helps unwind and separate the DNA helices (Figure 4). Another issue is that the **DNA polymerase** only works in one direction along the strand (5′ to 3′), but the double-stranded DNA has two strands oriented in opposite directions. This problem is solved by synthesizing the two strands slightly differently: one new strand grows continuously, the other in bits and pieces. Short fragments of RNA are used as primers for the **DNA polymerase**.





Figure 4. By Mariana Ruiz (DNA replication) Public Domain

## Practice Questions

Which of these separates the two complementary strands of DNA?

a. DNA polymerase
b. helicase
c. RNA primer
d. single-strand binding protein

[reveal-answer q="197431"]Show Answer[/reveal-answer]
[hidden-answer a="197431"]Answer b. Helicase breaks the hydrogen bonds holding together the two strands of DNA.

[/hidden-answer]

Which of these attaches complementary bases to the template strand?

a. DNA polymerase
b. helicase
c. RNA primer
d. single-strand binding protein

[reveal-answer q="381621"]Show Answer[/reveal-answer]
[hidden-answer a="381621"]Answer a. DNA polymerase builds the new strand of DNA.

[/hidden-answer]

Which of these is later replaced with DNA bases?

a. DNA polymerase
b. helicase
c. RNA primer
d. single-strand binding protein

[reveal-answer q="2143"]Show Answers[/reveal-answer]
[hidden-answer a="2143"]Answer c. the RNA primer is replaced with DNA nucleotides.[/hidden-answer]

## In Summary: Major Enzymes

Replication in eukaryotes starts at multiple origins of replication. A primer is required to initiate synthesis, which is then extended by DNA polymerase as it adds nucleotides one by one to the growing chain. The leading strand is synthesized continuously, whereas the lagging strand is synthesized in short stretches called Okazaki fragments. The RNA primers are replaced with DNA nucleotides; the DNA remains one continuous strand by linking the DNA fragments with DNA ligase. Below is a summary table of the major enzymes addressed in this reading, listed in rough order of activity during replication.

| Important Enzymes in DNA Replication | |
|---|---|
| Enzyme | Function |
| Topoisomerase | Relaxes the super-coiled DNA |
| DNA helicase | Unwinds the double helix at the replication fork |
| Primase | Provides the starting point for DNA polymerase to begin synthesis of the new strand |
| DNA polymerase | Synthesizes the new DNA strand; also proofreads and corrects some errors |
| DNA ligase | Re-joins the two DNA strands into a double helix and joins Okazaki fragments of the lagging strand |

## Contributors and Attributions

CC licensed content, Original

- Revision, adaptation, and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

CC licensed content, Shared previously

- DNA Replication. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5f7e80020ca6000225da6a4220c9. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

11.9: Major Enzymes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.10: “Proofreading” DNA

DNA replication is a highly accurate process, but mistakes can occasionally occur, such as a DNA polymerase inserting a wrong base. Uncorrected mistakes may sometimes lead to serious consequences, such as cancer. Repair mechanisms correct the mistakes. In rare cases, mistakes are not corrected, leading to mutations; in other cases, repair enzymes are themselves mutated or defective.

Most of the mistakes during DNA replication are promptly corrected by DNA polymerase by proofreading the base that has been just added (Figure 1). In **proofreading**, the DNA pol reads the newly added base before adding the next one, so a correction can be made. The polymerase checks whether the newly added base has paired correctly with the base in the template strand. If it is the right base, the next nucleotide is added. If an incorrect base has been added, the enzyme makes a cut at the phosphodiester bond and releases the wrong nucleotide. This is performed by the exonuclease action of DNA pol III. Once the incorrect nucleotide has been removed, a new one will be added again.





Figure 1. Proofreading by DNA polymerase corrects errors during replication.

Some errors are not corrected during replication, but are instead corrected after replication is completed; this type of repair is known as **mismatch repair** (Figure 2). The enzymes recognize the incorrectly added nucleotide and excise it; this is then replaced by the correct base. If this remains uncorrected, it may lead to more permanent damage. How do mismatch repair enzymes recognize which of the two bases is the incorrect one? In *E. coli*, after replication, the nitrogenous base adenine acquires a methyl group; the parental DNA strand will have methyl groups, whereas the newly synthesized strand lacks them. Thus, DNA polymerase is able to remove the wrongly incorporated bases from the newly synthesized, non-methylated strand. In eukaryotes, the mechanism is not very well understood, but it is believed to involve recognition of unsealed nicks in the new strand, as well as a short-term continuing association of some of the replication proteins with the new daughter strand after replication has completed.





Figure 2. In mismatch repair, the incorrectly added base is detected after replication. The mismatch repair proteins detect this base and remove it from the newly synthesized strand by nuclease action. The gap is now filled with the correctly paired base.

In another type of repair mechanism, **nucleotide excision repair**, enzymes replace incorrect bases by making a cut on both the 3′ and 5′ ends of the incorrect base (Figure 3).





Figure 3. Nucleotide excision repairs thymine dimers. When exposed to UV, thymines lying adjacent to each other can form thymine dimers. In normal cells, they are excised and replaced.

The segment of DNA is removed and replaced with the correctly paired nucleotides by the action of DNA pol. Once the bases are filled in, the remaining gap is sealed with a phosphodiester linkage catalyzed by DNA ligase. This repair mechanism is often employed when UV exposure causes the formation of pyrimidine dimers.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

11.10: "Proofreading" DNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.11: Telomeres

As you've learned, the enzyme DNA pol can add nucleotides only in the 5′ to 3′ direction. In the leading strand, synthesis continues until the end of the chromosome is reached. On the lagging strand, DNA is synthesized in short stretches, each of which is initiated by a separate primer. When the replication fork reaches the end of the linear chromosome, there is no place for a primer to be made for the DNA fragment to be copied at the end of the chromosome. These ends thus remain unpaired, and over time these ends may get progressively shorter as cells continue to divide.

The ends of the linear chromosomes are known as **telomeres**, which have repetitive sequences that code for no particular gene. In a way, these telomeres protect the genes from getting deleted as cells continue to divide. In humans, a six base pair sequence, TTAGGG, is repeated 100 to 1000 times. The discovery of the enzyme telomerase (Figure 1) helped in the understanding of how chromosome ends are maintained. The **telomerase** enzyme contains a catalytic part and a built-in RNA template. It attaches to the end of the chromosome, and complementary bases to the RNA template are added on the 3′ end of the DNA strand. Once the 3′ end of the lagging strand template is sufficiently elongated, DNA polymerase can add the nucleotides complementary to the ends of the chromosomes. Thus, the ends of the chromosomes are replicated.





Figure 1. The ends of linear chromosomes are maintained by the action of the telomerase enzyme.



Figure 2. Elizabeth Blackburn, 2009 Nobel Laureate, is the scientist who discovered how telomerase works. (credit: US Embassy Sweden)

Telomerase is typically active in germ cells and adult stem cells. It is not active in adult somatic cells. For her discovery of telomerase and its action, Elizabeth Blackburn (Figure 2) received the Nobel Prize for Medicine and Physiology in 2009.

## Telomerase and Aging

Cells that undergo cell division continue to have their telomeres shortened because most somatic cells do not make telomerase. This essentially means that telomere shortening is associated with aging. With the advent of modern medicine, preventative health care, and healthier lifestyles, the human life span has increased, and there is an increasing demand for people to look younger and have a better quality of life as they grow older.

In 2010, scientists found that telomerase can reverse some age-related conditions in mice. This may have potential in regenerative medicine.[1] Telomerase-deficient mice were used in these studies; these mice have tissue atrophy, stem cell depletion, organ system failure, and impaired tissue injury responses. Telomerase reactivation in these mice caused extension of telomeres, reduced DNA damage, reversed neurodegeneration, and improved the function of the testes, spleen, and intestines. Thus, telomere reactivation may have potential for treating age-related diseases in humans.

Cancer is characterized by uncontrolled cell division of abnormal cells. The cells accumulate mutations, proliferate uncontrollably, and can migrate to different parts of the body through a process called metastasis. Scientists have observed that cancerous cells have considerably shortened telomeres and that telomerase is active in these cells. Interestingly, only after the telomeres were shortened in the cancer cells did the telomerase become active. If the action of telomerase in these cells can be inhibited by drugs during cancer therapy, then the cancerous cells could potentially be stopped from further division.

### Learning Objectives

The ends of the chromosomes pose a problem during DNA replication as polymerase is unable to extend them without a primer. Telomerase, an enzyme with a built-in RNA template, extends the ends by copying the RNA template and extending one end of the chromosome. DNA polymerase can then extend the DNA using the primer. In this way, the ends of the chromosomes are protected. This is important as evidence indicates telomere length may play a role in regulating cell division and the process of aging.

1. Jaskelioff et al., “Telomerase reactivation reverses tissue degeneration in aged telomerase-deficient mice,” *Nature* 469 (2011): 102–7. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

11.11: Telomeres is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.12: Introduction to DNA Mutations

## Recognize the impact of DNA mutations

In this outcome, we'll learn just what mutations are and how they're often connected to our DNA.

## What You'll Learn to Do

- Understand what a mutation is and how one generally occurs
- Understand the impact of mutations in somatic cells versus gametes
- Identify the major types of DNA mutations

### Learning Activities

The learning activities for this section include the following:

- What is a Mutation?
- Mutations in Somatic Cells and in Gametes
- Major Types of Mutations
- Self Check: DNA Mutations

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

11.12: Introduction to DNA Mutations is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.


## 11.13: What is a Mutation?

Over a lifetime, our DNA can undergo changes or **mutations** in the sequence of bases: A, C, G and T. This results in changes in the proteins that are made. This can be a bad or a good thing.

A mutation is a change that occurs in our DNA sequence, either due to mistakes when the DNA is copied or as the result of environmental factors such as UV light and cigarette smoke. Mutations can occur during DNA replication if errors are made and not corrected in time. Mutations can also occur as the result of exposure to environmental factors such as smoking, sunlight and radiation. Often cells can recognize any potentially mutation-causing damage and repair it before it becomes a fixed mutation.

Mutations contribute to genetic variation within species. Mutations can also be inherited, particularly if they have a positive effect. For example, the disorder sickle cell anaemia is caused by a mutation in the gene that instructs the building of a protein called hemoglobin. This causes the red blood cells to become an abnormal, rigid, sickle shape. However, in African populations, having this mutation also protects against malaria.

However, mutation can also disrupt normal gene activity and cause diseases, like cancer. Cancer is the most common human genetic disease; it is caused by mutations occurring in a number of growth-controlling genes. Sometimes faulty, cancer-causing genes can exist from birth, increasing a person's chance of getting cancer.




An illustration to show an example of a DNA mutation. Image credit: Genome Research Limited

### Contributors and Attributions

CC licensed content, Shared previously

- What is a mutation?. **Authored by**: Genome Research Limited. **Located at**: http://www.yourgenome.org/facts/what-is-a-mutation. **Project**: YourGenome. **License**: *CC BY-SA: Attribution-ShareAlike*

11.13: What is a Mutation? is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.14: Mutations in Somatic Cells and in Gametes

Let's begin with a question: *What is a gene mutation and how do mutations occur?*

A gene mutation is a permanent alteration in the DNA sequence that makes up a gene, such that the sequence differs from what is found in most people. Mutations range in size; they can affect anywhere from a single DNA building block (base pair) to a large segment of a chromosome that includes multiple genes.

Gene mutations can be classified in two major ways:

- **Hereditary mutations** are inherited from a parent and are present throughout a person's life in virtually every cell in the body. These mutations are also called germline mutations because they are present in the parent's egg or sperm cells, which are also called germ cells. When an egg and a sperm cell unite, the resulting fertilized egg cell receives DNA from both parents. If this DNA has a mutation, the child that grows from the fertilized egg will have the mutation in each of his or her cells.
- **Acquired (or somatic) mutations** occur at some time during a person's life and are present only in certain cells, not in every cell in the body. These changes can be caused by environmental factors such as ultraviolet radiation from the sun, or can occur if a mistake is made as DNA copies itself during cell division. Acquired mutations in somatic cells (cells other than sperm and egg cells) cannot be passed on to the next generation.

Genetic changes that are described as de novo (new) mutations can be either hereditary or somatic. In some cases, the mutation occurs in a person's egg or sperm cell but is not present in any of the person's other cells. In other cases, the mutation occurs in the fertilized egg shortly after the egg and sperm cells unite. (It is often impossible to tell exactly when a de novo mutation happened.) As the fertilized egg divides, each resulting cell in the growing embryo will have the mutation. De novo mutations may explain genetic disorders in which an affected child has a mutation in every cell in the body but the parents do not, and there is no family history of the disorder.

Somatic mutations that happen in a single cell early in embryonic development can lead to a situation called mosaicism. These genetic changes are not present in a parent's egg or sperm cells, or in the fertilized egg, but happen a bit later when the embryo includes several cells. As all the cells divide during growth and development, cells that arise from the cell with the altered gene will have the mutation, while other cells will not. Depending on the mutation and how many cells are affected, mosaicism may or may not cause health problems.

Most disease-causing gene mutations are uncommon in the general population. However, other genetic changes occur more frequently. Genetic alterations that occur in more than 1 percent of the population are called polymorphisms. They are common enough to be considered a normal variation in the DNA. Polymorphisms are responsible for many of the normal differences between people such as eye color, hair color, and blood type. Although many polymorphisms have no negative effects on a person's health, some of these variations may influence the risk of developing certain disorders.

## Contributors and Attributions

Public domain content

- What is a gene mutation and how do mutations occur?. **Provided by**: NIH. **Located at**: https://ghr.nlm.nih.gov/primer/mutationsanddisorders/genemutation. **Project**: US National Library of Medicine. **License**: *Public Domain: No Known Copyright*

11.14: Mutations in Somatic Cells and in Gametes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.15: Major Types of Mutations



Figure 1. Xeroderma pigmentosa is a condition in which thymine dimerization from exposure to UV is not repaired. Exposure to sunlight results in skin lesions. (credit: James Halpern et al.)

A well-studied example of a mutation is seen in people suffering from xeroderma pigmentosa (Figure 1). Affected individuals have skin that is highly sensitive to UV rays from the sun.


When individuals are exposed to UV, pyrimidine dimers, especially those of thymine, are formed; people with xeroderma pigmentosa are not able to repair the damage. These are not repaired because of a defect in the nucleotide excision repair enzymes, whereas in normal individuals, the thymine dimers are excised and the defect is corrected. The thymine dimers distort the structure of the DNA double helix, and this may cause problems during DNA replication. People with xeroderma pigmentosa may have a higher risk of contracting skin cancer than those who don't have the condition.

Errors during DNA replication are not the only reason why mutations arise in DNA. **Mutations**, variations in the nucleotide sequence of a genome, can also occur because of damage to DNA. Such mutations may be of two types: induced or spontaneous. **Induced mutations** are those that result from an exposure to chemicals, UV rays, x-rays, or some other environmental agent. **Spontaneous mutations** occur without any exposure to any environmental agent; they are a result of natural reactions taking place within the body.

Mutations may have a wide range of effects. Some mutations are not expressed; these are known as **silent mutations**. **Point mutations** are those mutations that affect a single base pair. The most common nucleotide mutations are substitutions, in which one base is replaced by another. These can be of two types, either transitions or transversions. **Transition substitution** refers to a purine or pyrimidine being replaced by a base of the same kind; for example, a purine such as adenine may be replaced by the purine guanine. Transversion substitution refers to a purine being replaced by a pyrimidine, or vice versa; for example, cytosine, a pyrimidine, is replaced by adenine, a purine. Mutations can also be the result of the addition of a base, known as an insertion, or the removal of a base, also known as deletion. Sometimes a piece of DNA from one chromosome may get moved to another chromosome or to another region of the same chromosome; this is also known as translocation.

Mutations in repair genes have been known to cause cancer. Many mutated repair genes have been implicated in certain forms of pancreatic cancer, colon cancer, and colorectal cancer. Mutations can affect either somatic cells or germ cells. If many mutations accumulate in a somatic cell, they may lead to problems such as the uncontrolled cell division observed in cancer. If a mutation takes place in germ cells, the mutation will be passed on to the next generation, as in the case of hemophilia and xeroderma pigmentosa.

## Causes of Genetic Mutations



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=398

## Learning Objectives

DNA polymerase can make mistakes while adding nucleotides. Most mistakes are corrected, but if they are not, they may result in a mutation defined as a permanent change in the DNA sequence. Mutations can be of many types, such as **substitution**, **deletion**, **insertion**, and **translocation**. Mutations in repair genes may lead to serious consequences such as cancer. Mutations can be induced or may occur spontaneously.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- The causes of genetic mutations. **Authored by**: Ross Firestone. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/video/the-causes-of-genetic-mutations. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

---

11.15: Major Types of Mutations is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.16: Putting It Together- DNA Structure and Replication

Let's expand just a bit on how the process of personalized medicine works. Within any given genome there will be small differences in the DNA sequence. All humans, for example, are 99.99% identical and only differ by 0.01%. These small differences account for many features in an individual, including how he or she responds to disease treatments.

## Personalized Medicine in Practice

Let's return to our examples from the beginning of the module:

- **Blood clot treatments:** Screening the genome of a patient prior to prescribing Warafin for blood clots allows clinicians to determine if the drug will even work and if so, to pinpoint a specific dosage.
- **Colorectal cancer treatments:** In the case of colorectal cancer, the KRAS protein can be examined to determine if cetuximab will be effective. This is an important step as this drug is ineffective in about 40% of patients.
- **Breast cancer treatments:** Anti-breast cancer treatments can be assessed prior to trial-and-error in a patient based on the DNA sequence of key receptors in the body.

So, should you have your genes tested? Unfortunately, the decision about whether to get a particular genetic test can be complicated.



Getting a genetic test usually isn't difficult. Doctors generally take a sample of body fluid or tissue, such as blood, saliva or skin, and send it to a lab. Most genetic tests used today analyze just one or a few genes, often to help diagnose disease. Newborns, for example, are routinely screened for several genetic disorders by taking a few drops of blood from their heels. When life-threatening conditions are caught early, infants can be treated right away to prevent problems.

Genetic tests are now available for about 2,500 diseases, and that number keeps growing. Your doctor might advise you to get tested for specific genetic diseases if they tend to run in your family or if you have certain symptoms.

"While there are many genetic tests, they vary as to how well they predict risk," says Dr. Lawrence Brody, a genetic testing expert at NIH.

For some diseases, such as sickle cell anemia or cystic fibrosis, inheriting 2 copies of abnormal genes means a person will get that disease. But for other diseases and conditions, the picture is more complex. For type 2 diabetes, testing positive for some specific gene variants may help predict risk, but no better than other factors—such as obesity, high blood pressure and having a close relative with the disease.

The latest approach to personalized medicine is to get your whole genome sequenced. That's still expensive, but the cost has dropped dramatically over the past decade and will likely continue to fall. Since your genome essentially stays the same over time, this information might one day become part of your medical record, so doctors could consult it as needed.

You can start to get a sense of your genetic risks by putting together your family's health history. A free online tool called My Family Health Portrait from the U.S. Surgeon General can help you and your doctor spot early warning signs of conditions that run

in your family.

But personalized medicine isn't just about genes. You can learn a lot about your health risks by taking a close look at your current health and habits. Smoking, a poor diet, and lack of exercise can raise your risks for life-threatening health problems, such as heart disease and cancer. Talk to your health care provider about the steps you can take to understand and reduce your unique health risks.

## Learn More

Visit this site to learn more about how FDA regulations come into play.

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: DNA Structure and Replication. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- adult, africa, african, black, care, cheerful, clinic, doctor,. **Authored by**: Ilmicrofono Oggiono . **Provided by**: www.audio-luci-store.it. **Located at**: https://flic.kr/p/qtWa9C. **License**: *CC BY: Attribution*

Public domain content

- Personalized Medicine: Matching Treatments to Your Genes. **Authored by**: Vicki Contie, Dana Steinberg, Carol Torgan, and Katherine Wendelsdorf. **Provided by**: NIH. **Located at**: https://newsinhealth.nih.gov/issue/dec2013/feature1. **License**: *Public Domain: No Known Copyright*

---

11.16: Putting It Together- DNA Structure and Replication is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 11.17: Assignment- Which Has More DNA?

**Question:** Would a kiwifruit and strawberry that are approximately the same size also have approximately the same amount of DNA?

**Background:** You have learned that DNA is carried on chromosomes. All mammals are diploid; however, not all plants are diploid. The common strawberry is octoploid (8*n*) and the cultivated kiwi is hexaploid (6*n*). Research the total number of chromosomes in the cells of each of these fruits and think about how this might correspond to the amount of DNA in these fruits' cell nuclei.

**Hypothesis:** Hypothesize whether you would be able to detect a difference in DNA quantity from similarly sized strawberries and kiwis. Which fruit do you think would yield more DNA?

Test your hypothesis. Perform the DNA extraction experiment.

**Record your observations:** Because you are not quantitatively measuring DNA volume, you can record for each trial whether the two fruits produced the same or different amounts of DNA as observed by eye. If one or the other fruit produced noticeably more DNA, record this as well. Determine whether your observations are consistent with several pieces of each fruit.

**Analyze your data:** Did you notice an obvious difference in the amount of DNA produced by each fruit? Were your results reproducible?

**Draw a conclusion:** Given what you know about the number of chromosomes in each fruit, can you conclude that chromosome number necessarily correlates to DNA amount? Can you identify any drawbacks to this procedure? If you had access to a laboratory, how could you standardize your comparison and make it more quantitative?

To complete this assessment, address all of the points above in a lab write up. Be sure to include images of your experiment in progress. See the examples contained here.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references.
- Assignment addresses the required points noted above (hypothesis, observations/data, analysis of data, conclusion).

## Rubric

| Which Has More DNA? | | | | |
|---|---|---|---|---|
| Outcome: Relate DNA structure to the process of DNA replication | | | | |
| Criteria | Ratings | | | Pts |
| Hypothesis addresses how the DNA stores genetic information and the role of complementary base pairing in DNA storage/replication. | Hypothesis is testable and clearly addresses storage and replication of genetic information.<br>5.0 pts | Hypothesis is testable and addresses genetic information storage and replication with some explanation required.<br>4.0 pts | Hypothesis is not testable or does not address storage of genetic information and replication of DNA bases.<br>0.0 pts | 5 pts |
| Conclusion sections addresses how DNA is used to store genetic information. | Conclusion addresses storage of genetic information in detail and references experimental findings directly and in detail.<br>5.0 pts | Conclusion addresses storage of genetic information at high level and mentions experimental findings.<br>4.0 pts | Conclusion does not address or incorrectly addresses storage of genetic information.<br>0.0 pts | 5 pts |
| | | | | Total points: 10 |

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

11.17: Assignment- Which Has More DNA? is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 12: Module 10- DNA Transcription and Translation


12.1: Why It Matters- DNA Transcription and Translation
12.2: Introduction to Transcription
12.3: Steps of Genetic Transcription
12.4: RNA Polymerase
12.5: pre-RNA and mRNA
12.6: RNA Post-Translational Modification
12.7: Introduction to Translation
12.8: Steps of Translation
12.9: Ribosomes
12.10: tRNA
12.11: Codons
12.12: Protein Modifications
12.13: Introduction to Prokaryotic Transcription and Translation
12.14: Prokaryotic Transcription
12.15: Prokaryotic Translation
12.16: Introduction to the Central Dogma
12.17: The Central Dogma
12.18: Putting It Together- DNA Transcription and Translation



12: Module 10- DNA Transcription and Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.1: Why It Matters- DNA Transcription and Translation

## Why describe the conversion of DNA to RNA to proteins?




Figure 1. Before buildings are created, plans are drawn up. The plans direct the construction process.

By now you're familiar with genes and DNA sequences and how changes in the DNA can have a big impact. But the question still remains . . . how? How does DNA cause anything to happen?

Our bodies contain trillions of cells that need to be constructed in very precise ways. Much like when a construction company creates blueprints before they actually begin a project, your body needs some kind of plan to accomplish this.

DNA is acts as a blueprint. With this plan in every cell, your body is able to convert DNA into action molecules, which are proteins, by way of an intermediary, RNA. This idea is so central to biology that it is often called the central dogma of biology: DNA is transcribed to RNA which is translated to protein. This multi-step process is one of the reasons for the diversity we see in the world around us.

### Learning Outcomes

- Outline the process of eukaryotic transcription
- Summarize the process of translation
- Outline the process of prokaryotic transcription and translation
- Identify the central dogma of life

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: DNA Transcription and Translation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

Public domain content

- Joy Oil gas station blueprints. **Provided by**: Joy Oil Co Ltd. **Located at**: https://commons.wikimedia.org/wiki/File:Joy_Oil_gas_station_blueprints.jpg. **License**: *Public Domain: No Known Copyright*

12.1: Why It Matters- DNA Transcription and Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.2: Introduction to Transcription

## Outline the process of eukaryotic transcription

DNA is copied into RNA in a process called genetic transcription. To *transcribe* means to “put down something in writing.” The information in DNA is transcribed—or rewritten—into a smaller version (RNA) that can be used by the cell.

## What You’ll Learn to Do

- Understand the basic steps in the transcription of DNA into RNA
- Describe the role of RNA polymerase
- Understand the difference between pre-RNA and mRNA
- Describe RNA post-translational modification and its purpose

### Learning Activities

The learning activities for this section include the following:

- Steps of Genetic Transcription
- RNA Polymerase
- pre-RNA and mRNA
- RNA Post-Translational Modification
- Self Check: Transcription

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Transcription of DNA to RNA. **Provided by**: CK-12. **Located at**: www.ck12.org/biology/Transcription-of-DNA-to-RNA/lesson/Transcription-of-DNA-to-RNA-BIO/. **License**: *CC BY-NC: Attribution-NonCommercial*

12.2: Introduction to Transcription is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.3: Steps of Genetic Transcription

Transcription takes place in the nucleus. It uses DNA as a template to make an RNA (mRNA) molecule. During transcription, a strand of mRNA is made that is complementary to a strand of DNA. Figure 1 shows how this occurs.





Figure 1. Overview of Transcription. Transcription uses the sequence of bases in a strand of DNA to make a complementary strand of mRNA. Triplets are groups of three successive nucleotide bases in DNA. Codons are complementary groups of bases in mRNA.

You can also watch this more detailed video about transcription.

Transcription takes place in three steps: initiation, elongation, and termination. The steps are illustrated in Figure 2.





Figure 2. Transcription occurs in the three steps—initiation, elongation, and termination—all shown here.

## Step 1: Initiation

**Initiation** is the beginning of transcription. It occurs when the enzyme RNA polymerase binds to a region of a gene called the promoter. This signals the DNA to unwind so the enzyme can ‘‘read’’ the bases in one of the DNA strands. The enzyme is now ready to make a strand of mRNA with a complementary sequence of bases.


## Step 2: Elongation

**Elongation** is the addition of nucleotides to the mRNA strand. RNA polymerase reads the unwound DNA strand and builds the mRNA molecule, using complementary base pairs. There is a brief time during this process when the newly formed RNA is bound to the unwound DNA. During this process, an adenine (A) in the DNA binds to an uracil (U) in the RNA.

## Step 3: Termination

**Termination** is the ending of transcription, and occurs when RNA polymerase crosses a stop (termination) sequence in the gene. The mRNA strand is complete, and it detaches from DNA.

This video provides a review of these steps. You can stop watching the video at 5:35. (After this point, it discusses translation, which we'll discuss in the next outcome.)




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=410

## Contributors and Attributions

CC licensed content, Shared previously

- Transcription of DNA to RNA. **Provided by**: CK-12. **Located at**: www.ck12.org/biology/Transcription-of-DNA-to-RNA/lesson/Transcription-of-DNA-to-RNA-BIO/. **License**: *CC BY-NC: Attribution-NonCommercial*

All rights reserved content

- Transcription and Translation. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/h3b9ArupXZg. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

12.3: Steps of Genetic Transcription is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.4: RNA Polymerase

This section will expand upon the specific role of RNA polymerases during transcription. Read on to learn the role of RNA polymerases at each stage of transcription.

## Initiation of Transcription

Unlike the prokaryotic polymerase that can bind to a DNA template on its own, eukaryotes require several other proteins, called transcription factors, to first bind to the promoter region and then help recruit the appropriate polymerase.

### The Three Eukaryotic RNA Polymerases

The features of eukaryotic mRNA synthesis are markedly more complex those of prokaryotes. Instead of a single polymerase comprising five subunits, the eukaryotes have three polymerases that are each made up of 10 subunits or more. Each eukaryotic polymerase also requires a distinct set of transcription factors to bring it to the DNA template.

RNA polymerase I is located in the nucleolus, a specialized nuclear substructure in which ribosomal RNA (rRNA) is transcribed, processed, and assembled into ribosomes (Table 1). The rRNA molecules are considered structural RNAs because they have a cellular role but are not translated into protein. The rRNAs are components of the ribosome and are essential to the process of translation. RNA polymerase I synthesizes all of the rRNAs except for the 5S rRNA molecule. The “S” designation applies to “Svedberg” units, a nonadditive value that characterizes the speed at which a particle sediments during centrifugation.

Table 1. Locations, Products, and Sensitivities of the Three Eukaryotic RNA Polymerases

| RNA Polymerase | Cellular Compartment | Product of Transcription | α-Amanitin Sensitivity |
|---|---|---|---|
| I | Nucleolus | All rRNAs except 5S rRNA | Insensitive |
| II | Nucleus | All protein-coding nuclear pre-mRNAs | Extremely sensitive |
| III | Nucleus | 5S rRNA, tRNAs, and small nuclear RNAs | Moderately sensitive |

RNA polymerase II is located in the nucleus and synthesizes all protein-coding nuclear pre-mRNAs. Eukaryotic pre-mRNAs undergo extensive processing after transcription but before translation (Figure 1). For clarity, this module’s discussion of transcription and translation in eukaryotes will use the term “mRNAs” to describe only the mature, processed molecules that are ready to be translated. RNA polymerase II is responsible for transcribing the overwhelming majority of eukaryotic genes.





Figure 1. Eukaryotic mRNA contains introns that must be spliced out. A 5′ cap and 3′ poly-A tail are also added.

RNA polymerase III is also located in the nucleus. This polymerase transcribes a variety of structural RNAs that includes the 5S pre-rRNA, transfer pre-RNAs (pre-tRNAs), and **small nuclear** pre-**RNAs**. The tRNAs have a critical role in translation; they serve as the adaptor molecules between the mRNA template and the growing polypeptide chain. Small nuclear RNAs have a variety of functions, including “splicing” pre-mRNAs and regulating transcription factors.

A scientist characterizing a new gene can determine which polymerase transcribes it by testing whether the gene is expressed in the presence of a particular mushroom poison, α-amanitin (Table 1). Interestingly, α-amanitin produced by *Amanita phalloides*, the

Death Cap mushroom, affects the three polymerases very differently. RNA polymerase I is completely insensitive to α-amanitin, meaning that the polymerase can transcribe DNA in vitro in the presence of this poison. In contrast, RNA polymerase II is extremely sensitive to α-amanitin, and RNA polymerase III is moderately sensitive. Knowing the transcribing polymerase can clue a researcher into the general function of the gene being studied. Because RNA polymerase II transcribes the vast majority of genes, we will focus on this polymerase in our subsequent discussions about eukaryotic transcription factors and promoters.

## Structure of an RNA Polymerase II Promoter

Eukaryotic promoters are much larger and more complex than prokaryotic promoters, but both have a TATA box. For example, in the mouse thymidine kinase gene, the TATA box is located at approximately -30 relative to the initiation (+1) site (Figure 2). For this gene, the exact TATA box sequence is TATAAAA, as read in the 5′ to 3′ direction on the nontemplate strand. The thermostability of A–T bonds is low and this helps the DNA template to locally unwind in preparation for transcription.





Figure 2. A generalized promoter of a gene transcribed by RNA polymerase II is shown. Transcription factors recognize the promoter. RNA polymerase II then binds and forms the transcription initiation complex.

The mouse genome includes one gene and two pseudogenes for cytoplasmic thymidine kinase. Pseudogenes are genes that have lost their protein-coding ability or are no longer expressed by the cell. These pseudogenes are copied from mRNA and incorporated into the chromosome. For example, the mouse thymidine kinase promoter also has a conserved **CAAT box** (GGCCAATCT) at approximately -80. This sequence is essential and is involved in binding transcription factors. Further upstream of the TATA box, eukaryotic promoters may also contain one or more **GC-rich boxes** (GGCG) or **octamer boxes** (ATTTGCAT). These elements bind cellular factors that increase the efficiency of transcription initiation and are often identified in more “active” genes that are constantly being expressed by the cell.

## Transcription Factors for RNA Polymerase II

The complexity of eukaryotic transcription does not end with the polymerases and promoters. An army of basal transcription factors, enhancers, and silencers also help to regulate the frequency with which pre-mRNA is synthesized from a gene. Enhancers and silencers affect the efficiency of transcription but are not necessary for transcription to proceed. Basal transcription factors are crucial in the formation of a **preinitiation complex** on the DNA template that subsequently recruits RNA polymerase II for transcription initiation.

The names of the basal transcription factors begin with “TFII” (this is the transcription factor for RNA polymerase II) and are specified with the letters A–J. The transcription factors systematically fall into place on the DNA template, with each one further stabilizing the preinitiation complex and contributing to the recruitment of RNA polymerase II.

The processes of bringing RNA polymerases I and III to the DNA template involve slightly less complex collections of transcription factors, but the general theme is the same. Eukaryotic transcription is a tightly regulated process that requires a variety of proteins to interact with each other and with the DNA strand. Although the process of transcription in eukaryotes involves a greater metabolic investment than in prokaryotes, it ensures that the cell transcribes precisely the pre-mRNAs that it needs for protein synthesis.

### The Evolution of Promoters

The evolution of genes may be a familiar concept. Mutations can occur in genes during DNA replication, and the result may or may not be beneficial to the cell. By altering an enzyme, structural protein, or some other factor, the process of mutation can transform functions or physical features. However, eukaryotic promoters and other gene regulatory sequences may evolve as well. For instance, consider a gene that, over many generations, becomes more valuable to the cell. Maybe the gene encodes a structural protein that the cell needs to synthesize in abundance for a certain function. If this is the case, it would be beneficial to the cell for that gene's promoter to recruit transcription factors more efficiently and increase gene expression.

Scientists examining the evolution of promoter sequences have reported varying results. In part, this is because it is difficult to infer exactly where a eukaryotic promoter begins and ends. Some promoters occur within genes; others are located very far upstream, or even downstream, of the genes they are regulating. However, when researchers limited their examination to human core promoter sequences that were defined experimentally as sequences that bind the preinitiation complex, they found that promoters evolve even faster than protein-coding genes.

It is still unclear how promoter evolution might correspond to the evolution of humans or other higher organisms. However, the evolution of a promoter to effectively make more or less of a given gene product is an intriguing alternative to the evolution of the genes themselves.[1]

### Promoter Structures for RNA Polymerases I and III

In eukaryotes, the conserved promoter elements differ for genes transcribed by RNA polymerases I, II, and III. RNA polymerase I transcribes genes that have two GC-rich promoter sequences in the –45 to +20 region. These sequences alone are sufficient for transcription initiation to occur, but promoters with additional sequences in the region from –180 to –105 upstream of the initiation site will further enhance initiation. Genes that are transcribed by RNA polymerase III have upstream promoters or promoters that occur within the genes themselves.

## Elongation and Termination

Following the formation of the preinitiation complex, the polymerase is released from the other transcription factors, and elongation is allowed to proceed as it does in prokaryotes with the polymerase synthesizing pre-mRNA in the 5′ to 3′ direction. As discussed previously, RNA polymerase II transcribes the major share of eukaryotic genes, so this section will focus on how this polymerase accomplishes elongation and termination.

Although the enzymatic process of elongation is essentially the same in eukaryotes and prokaryotes, the DNA template is more complex. When eukaryotic cells are not dividing, their genes exist as a diffuse mass of DNA and proteins called chromatin. The DNA is tightly packaged around charged histone proteins at repeated intervals. These DNA–histone complexes, collectively called nucleosomes, are regularly spaced and include 146 nucleotides of DNA wound around eight histones like thread around a spool.

For polynucleotide synthesis to occur, the transcription machinery needs to move histones out of the way every time it encounters a nucleosome. This is accomplished by a special protein complex called **FACT**, which stands for "facilitates chromatin transcription." This complex pulls histones away from the DNA template as the polymerase moves along it. Once the pre-mRNA is synthesized, the FACT complex replaces the histones to recreate the nucleosomes.

The termination of transcription is different for the different polymerases. Unlike in prokaryotes, elongation by RNA polymerase II in eukaryotes takes place 1,000–2,000 nucleotides beyond the end of the gene being transcribed. This pre-mRNA tail is subsequently removed by cleavage during mRNA processing. On the other hand, RNA polymerases I and III require termination signals. Genes transcribed by RNA polymerase I contain a specific 18-nucleotide sequence that is recognized by a termination protein. The process of termination in RNA polymerase III involves an mRNA hairpin similar to rho-independent termination of transcription in prokaryotes.

1. H Liang et al., "Fast evolution of core promoters in primate genomes," *Molecular Biology and Evolution* 25 (2008): 1239–44. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

12.4: RNA Polymerase is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.5: pre-RNA and mRNA

After transcription, eukaryotic pre-mRNAs must undergo several processing steps before they can be translated. Eukaryotic (and prokaryotic) tRNAs and rRNAs also undergo processing before they can function as components in the protein synthesis machinery.

## mRNA Processing

The eukaryotic pre-mRNA undergoes extensive processing before it is ready to be translated. The additional steps involved in eukaryotic mRNA maturation create a molecule with a much longer half-life than a prokaryotic mRNA. Eukaryotic mRNAs last for several hours, whereas the typical *E. coli* mRNA lasts no more than five seconds.

Pre-mRNAs are first coated in RNA-stabilizing proteins; these protect the pre-mRNA from degradation while it is processed and exported out of the nucleus. The three most important steps of pre-mRNA processing are the addition of stabilizing and signaling factors at the 5′ and 3′ ends of the molecule, and the removal of intervening sequences that do not specify the appropriate amino acids. In rare cases, the mRNA transcript can be "edited" after it is transcribed.

### 5′ Capping

While the pre-mRNA is still being synthesized, a **7-methylguanosine cap** is added to the 5′ end of the growing transcript by a phosphate linkage. This moiety (functional group) protects the nascent mRNA from degradation. In addition, factors involved in protein synthesis recognize the cap to help initiate translation by ribosomes.

### 3′ Poly-A Tail

Once elongation is complete, the pre-mRNA is cleaved by an endonuclease between an AAUAAA consensus sequence and a GU-rich sequence, leaving the AAUAAA sequence on the pre-mRNA. An enzyme called poly-A polymerase then adds a string of approximately 200 A residues, called the **poly-A tail**. This modification further protects the pre-mRNA from degradation and signals the export of the cellular factors that the transcript needs to the cytoplasm.

### Pre-mRNA Splicing

Eukaryotic genes are composed of **exons**, which correspond to protein-coding sequences (*ex*-on signifies that they are *ex*pressed), and *int*ervening sequences called **introns** (*int*ron denotes their *int*ervening role), which may be involved in gene regulation but are removed from the pre-mRNA during processing. Intron sequences in mRNA do not encode functional proteins.

The discovery of introns came as a surprise to researchers in the 1970s who expected that pre-mRNAs would specify protein sequences without further processing, as they had observed in prokaryotes. The genes of higher eukaryotes very often contain one or more introns. These regions may correspond to regulatory sequences; however, the biological significance of having many introns or having very long introns in a gene is unclear. It is possible that introns slow down gene expression because it takes longer to transcribe pre-mRNAs with lots of introns. Alternatively, introns may be nonfunctional sequence remnants left over from the fusion of ancient genes throughout evolution. This is supported by the fact that separate exons often encode separate protein subunits or domains. For the most part, the sequences of introns can be mutated without ultimately affecting the protein product.

All of a pre-mRNA's introns must be completely and precisely removed before protein synthesis. If the process errs by even a single nucleotide, the reading frame of the rejoined exons would shift, and the resulting protein would be dysfunctional. The process of removing introns and reconnecting exons is called **splicing** (Figure 1). Introns are removed and degraded while the pre-mRNA is still in the nucleus. Splicing occurs by a sequence-specific mechanism that ensures introns will be removed and exons rejoined with the accuracy and precision of a single nucleotide. The splicing of pre-mRNAs is conducted by complexes of proteins and RNA molecules called spliceosomes.

Practice Question





Figure 1. Pre-mRNA splicing involves the precise removal of introns from the primary RNA transcript. The splicing process is catalyzed by protein complexes called spliceosomes that are composed of proteins and RNA molecules called snRNAs. Spliceosomes recognize sequences at the 5′ and 3′ end of the intron.

Errors in splicing are implicated in cancers and other human diseases. What kinds of mutations might lead to splicing errors?

[practice-area rows="2"][/practice-area]
[reveal-answer q="454729"]Show Answer[/reveal-answer]
[hidden-answer a="454729"]Think of different possible outcomes if splicing errors occur. Mutations in the spliceosome recognition sequence at each end of the intron, or in the proteins and RNAs that make up the spliceosome, may impair splicing. Mutations may also add new spliceosome recognition sites. Splicing errors could lead to introns being retained in spliced RNA, exons being excised, or changes in the location of the splice site.[/hidden-answer]

Note that more than 70 individual introns can be present, and each has to undergo the process of splicing—in addition to 5′ capping and the addition of a poly-A tail—just to generate a single, translatable mRNA molecule.

See how introns are removed during RNA splicing at this website.

Try It





Figure 2. *Trypanosoma brucei* is the causative agent of sleeping sickness in humans. The mRNAs of this pathogen must be modified by the addition of nucleotides before protein synthesis can occur. (credit: modification of work by Torsten Ochsenreiter)

The trypanosomes are a group of protozoa that include the pathogen *Trypanosoma brucei*, which causes sleeping sickness in humans (Figure 2). Trypanosomes, and virtually all other eukaryotes, have organelles called mitochondria that supply the cell with chemical energy. Mitochondria are organelles that express their own DNA and are believed to be the remnants of a symbiotic relationship between a eukaryote and an engulfed prokaryote. The mitochondrial DNA of trypanosomes exhibit an interesting exception to The Central Dogma: their pre-mRNAs do not have the correct information to specify a functional protein. Usually, this is because the mRNA is missing several U nucleotides. The cell performs an additional RNA processing step called RNA editing to remedy this.

Other genes in the mitochondrial genome encode 40- to 80-nucleotide guide RNAs. One or more of these molecules interacts by complementary base pairing with some of the nucleotides in the pre-mRNA transcript. However, the guide RNA has more A nucleotides than the pre-mRNA has U nucleotides to bind with. In these regions, the guide RNA loops out. The 3′ ends of guide RNAs have a long poly-U tail, and these U bases are inserted in regions of the pre-mRNA transcript at which the guide RNAs are

looped. This process is entirely mediated by RNA molecules. That is, guide RNAs—rather than proteins—serve as the catalysts in RNA editing.

RNA editing is not just a phenomenon of trypanosomes. In the mitochondria of some plants, almost all pre-mRNAs are edited. RNA editing has also been identified in mammals such as rats, rabbits, and even humans. What could be the evolutionary reason for this additional step in pre-mRNA processing? One possibility is that the mitochondria, being remnants of ancient prokaryotes, have an equally ancient RNA-based method for regulating gene expression. In support of this hypothesis, edits made to pre-mRNAs differ depending on cellular conditions. Although speculative, the process of RNA editing may be a holdover from a primordial time when RNA molecules, instead of proteins, were responsible for catalyzing reactions.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.5: pre-RNA and mRNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.6: RNA Post-Translational Modification

**Learning Objectives**

- Gene expression can be regulated at various stages after an RNA transcript has been produced.
- Some transcripts can undergo **alternative splicing**. This regulated process makes different mRNAs and proteins from the same initial RNA transcript.
- Some mRNAs are targeted by small regulatory RNAs, including **miRNAs**, which can cause mRNA degradation or block translation.
- A protein's activity may be regulated after translation by mechanisms such as proteolysis ("snipping out" of pieces) and addition of chemical groups.

The genes that a eukaryotic cell turns "on" largely determine its identity and properties. For instance, a photoreceptor cell in your eye can detect light because it expresses genes for light-sensitive proteins, as well as as genes for neurotransmitters that allow signals to be relayed to the brain.

In eukaryotic cells like photoreceptors, gene expression is often controlled primarily at the level of transcription. However, that doesn't mean transcription is the last chance for regulation. Later stages of gene expression can also be regulated, including the following:

- RNA processing, such as splicing, capping, and addition of a poly-A tail
- Messenger RNA (mRNA) translation and lifetime in the cytosol
- Protein modifications, such as addition of chemical groups or removal of amino acids

In the sections below, we'll discuss some common types of gene regulation that occur after an RNA transcript has been made.

## Regulation of RNA processing

When a eukaryotic gene is transcribed in the nucleus, the primary transcript (freshly made RNA molecule) isn't yet considered a messenger RNA. Instead, it's an "immature" molecule called a pre-mRNA.

The pre-mRNA has to go through some modifications to become a mature mRNA molecule that can leave the nucleus and be translated. These include splicing, capping, and addition of a poly-A tail, all of which can potentially be regulated – sped up, slowed down, or altered to result in a different product.

### Alternative splicing

Most pre-mRNA molecules have sections that are removed from the molecule, called **introns**, and sections that are linked or together to make the final mRNA, called **exons**. This process is called **splicing**.

In the process of **alternative splicing**, different portions of an mRNA can be selected for use as exons. This allows either of two (or more) mRNA molecules to be made from one pre-mRNA.





Figure 1. Image modified from "Eukaryotic Post-transcriptional Gene Regulation," by OpenStax College, Biology (CC BY 3.0).

Alternative splicing is not a random process. Instead, it's typically controlled by regulatory proteins. The proteins bind to specific sites on the pre-mRNA and "tell" the splicing factors which exons should be used. Different cell types may express different regulatory proteins, so different exon combinations can be used in each cell type, leading to the production of different proteins.

## Small regulatory RNA

Once an mRNA has left the nucleus, it may or may not be translated many times to make proteins. Two key determinants of how much protein is made from an mRNA are its “lifespan” (how long it floats around in the cytosol) and how readily the translation machinery, such as the ribosome, can attach to it.

A recently discovered class of regulators, called small regulatory RNAs, can control mRNA lifespan and translation. Let’s see how this works.

### microRNAs

**microRNAs (miRNAs)** were among the first small regulatory RNAs to be discovered. A miRNA is first transcribed as a long RNA molecule, which forms base pairs with itself and folds over to make a hairpin. Next, the hairpin is chopped up by enzymes, releasing a small double-stranded fragment of about 20 nucleotides. One of the strands in this fragment is the mature miRNA, which binds to a specific protein to make an RNA-protein complex.





Figure 2. Image modifed from “miRNA biogenesis,” by Narayanese (CC BY-SA 3.0). The modified image is licensed under a CC BY-SA 3.0 license.

The miRNA directs the protein complex to “matching” mRNA molecules (ones that form base pairs with the miRNA). When the RNA-protein complex binds:

- If the miRNA and its target match perfectly, an enzyme in the RNA-protein complex will typically chop the mRNA in half, leading to its breakdown.
- If the miRNA and its target have some mismatches, the RNA-protein complex may instead bind to the mRNA and keep it from being translated.

These are not the only ways that miRNAs inhibit expression of their targets, and scientists are still investigating their many modes of action.[1]

---

1. Carthew, R. W. and Sontheimer, E. J. (2009). Origins and mechanisms of miRNAs and siRNAs. *Cell*, *136*(4), 642–655. [1]http://dx.doi.org/10.1016/j.cell.2009.01.035. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Regulation after transcription. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/gene-regulation/gene-regulation-in-eukaryotes/a/regulation-after-transcription. **License**: *CC BY-NC-SA: Attribution-NonCommercial-*

*ShareAlike*

12.6: RNA Post-Translational Modification is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.7: Introduction to Translation

## Summarize the process of translation

RNA undergoes genetic translation, a process that produces proteins.

## What You'll Learn to Do

- Outline the basic steps of translation
- Describe the composition and role of ribosomes in translation, focusing on rRNA
- Understand the role of tRNA in translation
- Describe a codon and how they are used in translation
- Describe how proteins may be modified after translation

### Learning Activities

The learning activities for this section include the following:

- Steps of Translation
- Ribosomes
- tRNA
- Codons
- Protein Modifications
- Self Check: Translation

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

12.7: Introduction to Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.8: Steps of Translation

As with mRNA synthesis, protein synthesis can be divided into three phases: initiation, elongation, and termination. The process of translation is similar in prokaryotes and eukaryotes. Here we'll explore how translation occurs in *E. coli*, a representative prokaryote, and specify any differences between prokaryotic and eukaryotic translation.

## Initiation of Translation

Protein synthesis begins with the formation of an initiation complex. In *E. coli*, this complex involves the small 30S ribosome, the mRNA template, three initiation factors (IFs; IF-1, IF-2, and IF-3), and a special **initiator tRNA**, called $\text{tRNA}^{\text{Met}}_{\text{f}}$. The initiator tRNA interacts with the start codon AUG (or rarely, GUG), links to a formylated methionine called fMet, and can also bind IF-2. Formylated methionine is inserted by $\text{fMet} - \text{tRNA}^{\text{ Met}}_{\text{ f}}$ at the beginning of every polypeptide chain synthesized by *E. coli*, but it is usually clipped off after translation is complete. When an in-frame AUG is encountered during translation elongation, a non-formylated methionine is inserted by a regular Met-$\text{tRNA}^{\text{Met}}$.

In *E. coli* mRNA, a sequence upstream of the first AUG codon, called the **Shine-Dalgarno sequence** (AGGAGG), interacts with the rRNA molecules that compose the ribosome. This interaction anchors the 30S ribosomal subunit at the correct location on the mRNA template. Guanosine triphosphate (GTP), which is a purine nucleotide triphosphate, acts as an energy source during translation—both at the start of elongation and during the ribosome's translocation.

In eukaryotes, a similar initiation complex forms, comprising mRNA, the 40S small ribosomal subunit, IFs, and nucleoside triphosphates (GTP and ATP). The charged initiator tRNA, called Met-tRNAi, does not bind fMet in eukaryotes, but is distinct from other Met-tRNAs in that it can bind IFs.

Instead of depositing at the Shine-Dalgarno sequence, the eukaryotic initiation complex recognizes the 7-methylguanosine cap at the 5′ end of the mRNA. A cap-binding protein (CBP) and several other IFs assist the movement of the ribosome to the 5′ cap. Once at the cap, the initiation complex tracks along the mRNA in the 5′ to 3′ direction, searching for the AUG start codon. Many eukaryotic mRNAs are translated from the first AUG, but this is not always the case. According to **Kozak's rules**, the nucleotides around the AUG indicate whether it is the correct start codon. Kozak's rules state that the following consensus sequence must appear around the AUG of vertebrate genes: 5′-gccRccAUGG-3′. The R (for purine) indicates a site that can be either A or G, but cannot be C or U. Essentially, the closer the sequence is to this consensus, the higher the efficiency of translation.

Once the appropriate AUG is identified, the other proteins and CBP dissociate, and the 60S subunit binds to the complex of Met-tRNAi, mRNA, and the 40S subunit. This step completes the initiation of translation in eukaryotes.

## Translation, Elongation, and Termination

In prokaryotes and eukaryotes, the basics of elongation are the same, so we will review elongation from the perspective of *E. coli*. The 50S ribosomal subunit of *E. coli* consists of three compartments: the A (aminoacyl) site binds incoming charged aminoacyl tRNAs. The P (peptidyl) site binds charged tRNAs carrying amino acids that have formed peptide bonds with the growing polypeptide chain but have not yet dissociated from their corresponding tRNA. The E (exit) site releases dissociated tRNAs so that they can be recharged with free amino acids. There is one exception to this assembly line of tRNAs: in *E. coli*, $\text{fMet-tRNA}^{\text{ Met}}_{\text{ f}}$ is capable of entering the P site directly without first entering the A site. Similarly, the eukaryotic Met-tRNAi, with help from other proteins of the initiation complex, binds directly to the P site (Figure 1). In both cases, this creates an initiation complex with a free A site ready to accept the tRNA corresponding to the first codon after the AUG.





Figure 1. Ribosome mRNA translation

During translation elongation, the mRNA template provides specificity. As the ribosome moves along the mRNA, each mRNA codon comes into register, and specific binding with the corresponding charged tRNA anticodon is ensured. If mRNA were not present in the elongation complex, the ribosome would bind tRNAs nonspecifically.

Elongation proceeds with charged tRNAs entering the A site and then shifting to the P site followed by the E site with each single-codon "step" of the ribosome. Ribosomal steps are induced by conformational changes that advance the ribosome by three bases in the 3′ direction. The energy for each step of the ribosome is donated by an elongation factor that hydrolyzes GTP. Peptide bonds form between the amino group of the amino acid attached to the A-site tRNA and the carboxyl group of the amino acid attached to the P-site tRNA. The formation of each peptide bond is catalyzed by **peptidyl transferase**, an RNA-based enzyme that is integrated into the 50S ribosomal subunit. The energy for each peptide bond formation is derived from GTP hydrolysis, which is catalyzed by a separate elongation factor. The amino acid bound to the P-site tRNA is also linked to the growing polypeptide chain. As the ribosome steps across the mRNA, the former P-site tRNA enters the E site, detaches from the amino acid, and is expelled (Figure 2). Amazingly, the *E. coli* translation apparatus takes only 0.05 seconds to add each amino acid, meaning that a 200-amino acid protein can be translated in just 10 seconds.





Figure 2. Translation begins when an initiator tRNA anticodon recognizes a codon on mRNA. The large ribosomal subunit joins the small subunit, and a second tRNA is recruited. As the mRNA moves relative to the ribosome, the polypeptide chain is formed. Entry of a release factor into the A site terminates translation and the components dissociate.

## Practice Questions

Many antibiotics inhibit bacterial protein synthesis. For example, tetracycline blocks the A site on the bacterial ribosome, and chloramphenicol blocks peptidyl transfer. What specific effect would you expect each of these antibiotics to have on protein synthesis?

Tetracycline would directly affect:

a. tRNA binding to the ribosome
b. ribosome assembly
c. growth of the protein chain

[reveal-answer q=”10129″]**Show Answer**[/reveal-answer]
[hidden-answer a=”10129″]Answer a. Tetracycline would directly affect tRNA binding to the ribosome.

[/hidden-answer]

Chloramphenicol would directly affect

a. tRNA binding to the ribosome
b. ribosome assembly
c. growth of the protein chain

[reveal-answer q="10029"]**Show Answer**[/reveal-answer]
[hidden-answer a="10029"]Answer c. Chloramphenicol would directly affect growth of the protein chain.[/hidden-answer]

Termination of translation occurs when a nonsense codon (UAA, UAG, or UGA) is encountered. Upon aligning with the A site, these nonsense codons are recognized by release factors in prokaryotes and eukaryotes that instruct peptidyl transferase to add a water molecule to the carboxyl end of the P-site amino acid. This reaction forces the P-site amino acid to detach from its tRNA, and the newly made protein is released. The small and large ribosomal subunits dissociate from the mRNA and from each other; they are recruited almost immediately into another translation initiation complex. After many ribosomes have completed translation, the mRNA is degraded so the nucleotides can be reused in another transcription reaction.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.8: Steps of Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.9: Ribosomes





Figure 1. A peptide bond links the carboxyl end of one amino acid with the amino end of another, expelling one water molecule. For simplicity in this image, only the functional groups involved in the peptide bond are shown. The R and R′ designations refer to the rest of each amino acid structure.

The synthesis of proteins consumes more of a cell's energy than any other metabolic process. In turn, proteins account for more mass than any other component of living organisms (with the exception of water), and proteins perform virtually every function of a cell. The process of translation, or protein synthesis, involves the decoding of an mRNA message into a polypeptide product. Amino acids are covalently strung together by interlinking peptide bonds in lengths ranging from approximately 50 amino acid residues to more than 1,000. Each individual amino acid has an amino group ($NH_2$) and a carboxyl (COOH) group. Polypeptides are formed when the amino group of one amino acid forms an amide (i.e., peptide) bond with the carboxyl group of another amino acid (Figure 1). This reaction is catalyzed by ribosomes and generates one water molecule.

## Protein Synthesis Machinery

In addition to the mRNA template, many molecules and macromolecules contribute to the process of translation. The composition of each component may vary across species; for instance, ribosomes may consist of different numbers of rRNAs and polypeptides depending on the organism. However, the general structures and functions of the protein synthesis machinery are comparable from bacteria to human cells. Translation requires the input of an mRNA template, ribosomes, tRNAs, and various enzymatic factors.

Click through the steps of this PBS interactive to see protein synthesis in action.

### Polysomes

Even before an mRNA is translated, a cell must invest energy to build each of its ribosomes. In *E. coli*, there are between 10,000 and 70,000 ribosomes present in each cell at any given time. A ribosome is a complex macromolecule composed of structural and catalytic rRNAs, and many distinct polypeptides. In eukaryotes, the nucleolus is completely specialized for the synthesis and assembly of rRNAs.

Ribosomes exist in the cytoplasm in prokaryotes and in the cytoplasm and rough endoplasmic reticulum in eukaryotes. Mitochondria and chloroplasts also have their own ribosomes in the matrix and stroma, which look more similar to prokaryotic ribosomes (and have similar drug sensitivities) than the ribosomes just outside their outer membranes in the cytoplasm. Ribosomes dissociate into large and small subunits when they are not synthesizing proteins and reassociate during the initiation of translation. In *E. coli*, the small subunit is described as 30S, and the large subunit is 50S, for a total of 70S (recall that Svedberg units are not additive). Mammalian ribosomes have a small 40S subunit and a large 60S subunit, for a total of 80S. The small subunit is responsible for binding the mRNA template, whereas the large subunit sequentially binds tRNAs. Each mRNA molecule is simultaneously translated by many ribosomes, all synthesizing protein in the same direction: reading the mRNA from 5′ to 3′ and synthesizing the polypeptide from the N terminus to the C terminus. The complete mRNA/poly-ribosome structure is called a **polysome**.

### tRNAs

The tRNAs are structural RNA molecules that were transcribed from genes by RNA polymerase III. Depending on the species, 40 to 60 types of tRNAs exist in the cytoplasm. Serving as adaptors, specific tRNAs bind to sequences on the mRNA template and add

the corresponding amino acid to the polypeptide chain. Therefore, tRNAs are the molecules that actually “translate” the language of RNA into the language of proteins.

Of the 64 possible mRNA codons—or triplet combinations of A, U, G, and C—three specify the termination of protein synthesis and 61 specify the addition of amino acids to the polypeptide chain. Of these 61, one codon (AUG) also encodes the initiation of translation. Each tRNA anticodon can base pair with one of the mRNA codons and add an amino acid or terminate translation, according to the genetic code. For instance, if the sequence CUA occurred on an mRNA template in the proper reading frame, it would bind a tRNA expressing the complementary sequence, GAU, which would be linked to the amino acid leucine.

As the adaptor molecules of translation, it is surprising that tRNAs can fit so much specificity into such a small package. Consider that tRNAs need to interact with three factors:

1. They must be recognized by the correct aminoacyl synthetase.
2. They must be recognized by ribosomes.
3. They must bind to the correct sequence in mRNA.

### Aminoacyl tRNA Synthetases

The process of pre-tRNA synthesis by RNA polymerase III only creates the RNA portion of the adaptor molecule. The corresponding amino acid must be added later, once the tRNA is processed and exported to the cytoplasm. Through the process of tRNA “charging,” each tRNA molecule is linked to its correct amino acid by a group of enzymes called aminoacyl tRNA synthetases. At least one type of **aminoacyl tRNA synthetase** exists for each of the 20 amino acids; the exact number of aminoacyl tRNA synthetases varies by species. These enzymes first bind and hydrolyze ATP to catalyze a high-energy bond between an amino acid and adenosine monophosphate (AMP); a pyrophosphate molecule is expelled in this reaction. The activated amino acid is then transferred to the tRNA, and AMP is released.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.9: Ribosomes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.10: tRNA

## Processing of tRNAs and rRNAs





Figure 1. This is a space-filling model of a tRNA molecule that adds the amino acid phenylalanine to a growing polypeptide chain. The anticodon AAG binds the Codon UUC on the mRNA. The amino acid phenylalanine is attached to the other end of the tRNA.

The tRNAs and rRNAs are structural molecules that have roles in protein synthesis; however, these RNAs are not themselves translated. Pre-rRNAs are transcribed, processed, and assembled into ribosomes in the nucleolus. Pre-tRNAs are transcribed and processed in the nucleus and then released into the cytoplasm where they are linked to free amino acids for protein synthesis.

Most of the tRNAs and rRNAs in eukaryotes and prokaryotes are first transcribed as a long precursor molecule that spans multiple rRNAs or tRNAs. Enzymes then cleave the precursors into subunits corresponding to each structural RNA. Some of the bases of pre-rRNAs are methylated; that is, a $–CH_3$ moiety (methyl functional group) is added for stability. Pre-tRNA molecules also undergo methylation. As with pre-mRNAs, subunit excision occurs in eukaryotic pre-RNAs destined to become tRNAs or rRNAs.

Mature rRNAs make up approximately 50 percent of each ribosome. Some of a ribosome's RNA molecules are purely structural, whereas others have catalytic or binding activities. Mature tRNAs take on a three-dimensional structure through intramolecular hydrogen bonding to position the amino acid binding site at one end and the **anticodon** at the other end (Figure 1).

The anticodon is a three-nucleotide sequence in a tRNA that interacts with an mRNA codon through complementary base pairing.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.10: tRNA is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.11: Codons

Given the different numbers of “letters” in the mRNA and protein “alphabets,” scientists theorized that combinations of nucleotides corresponded to single amino acids. Nucleotide doublets would not be sufficient to specify every amino acid because there are only 16 possible two-nucleotide combinations (42). In contrast, there are 64 possible nucleotide triplets (43), which is far more than the number of amino acids. Scientists theorized that amino acids were encoded by nucleotide triplets and that the genetic code was **degenerate**. In other words, a given amino acid could be encoded by more than one nucleotide triplet. This was later confirmed experimentally; Francis Crick and Sydney Brenner used the chemical mutagen proflavin to insert one, two, or three nucleotides into the gene of a virus. When one or two nucleotides were inserted, protein synthesis was completely abolished. When three nucleotides were inserted, the protein was synthesized and functional. This demonstrated that three nucleotides specify each amino acid. These nucleotide triplets are called **codons**. The insertion of one or two nucleotides completely changed the triplet reading frame, thereby altering the message for every subsequent amino acid (Figure 1). Though insertion of three nucleotides caused an extra amino acid to be inserted during translation, the integrity of the rest of the protein was maintained.





Figure 1. The deletion of two nucleotides shifts the reading frame of an mRNA and changes the entire protein message, creating a nonfunctional protein or terminating protein synthesis altogether.

Scientists painstakingly solved the genetic code by translating synthetic mRNAs in vitro and sequencing the proteins they specified (Figure 2).





Figure 2. This figure shows the genetic code for translating each nucleotide triplet in mRNA into an amino acid or a termination signal in a nascent protein. (credit: modification of work by NIH)

In addition to instructing the addition of a specific amino acid to a polypeptide chain, three of the 64 codons terminate protein synthesis and release the polypeptide from the translation machinery. These triplets are called nonsense codons, or stop codons. Another codon, AUG, also has a special function. In addition to specifying the amino acid methionine, it also serves as the start codon to initiate translation. The reading frame for translation is set by the AUG start codon near the 5′ end of the mRNA.

The genetic code is universal. With a few exceptions, virtually all species use the same genetic code for protein synthesis. Conservation of codons means that a purified mRNA encoding the globin protein in horses could be transferred to a tulip cell, and the tulip would synthesize horse globin. That there is only one genetic code is powerful evidence that all of life on Earth shares a common origin, especially considering that there are about 1084 possible combinations of 20 amino acids and 64 triplet codons.

Transcribe a gene and translate it to protein using complementary pairing and the genetic code at this site.

Degeneracy is believed to be a cellular mechanism to reduce the negative impact of random mutations. Codons that specify the same amino acid typically only differ by one nucleotide. In addition, amino acids with chemically similar side chains are encoded by similar codons. This nuance of the genetic code ensures that a single-nucleotide substitution mutation might either specify the same amino acid but have no effect or specify a similar amino acid, preventing the protein from being rendered completely nonfunctional.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

12.11: Codons is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.12: Protein Modifications

During and after translation, individual amino acids may be chemically modified, signal sequences may be appended, and the new protein “folds” into a distinct three-dimensional structure as a result of intramolecular interactions. A **signal sequence** is a short tail of amino acids that directs a protein to a specific cellular compartment. These sequences at the amino end or the carboxyl end of the protein can be thought of as the protein’s “train ticket” to its ultimate destination. Other cellular factors recognize each signal sequence and help transport the protein from the cytoplasm to its correct compartment. For instance, a specific sequence at the amino terminus will direct a protein to the mitochondria or chloroplasts (in plants). Once the protein reaches its cellular destination, the signal sequence is usually clipped off.

Many proteins fold spontaneously, but some proteins require helper molecules, called chaperones, to prevent them from aggregating during the complicated process of folding. Even if a protein is properly specified by its corresponding mRNA, it could take on a completely dysfunctional shape if abnormal temperature or pH conditions prevent it from folding correctly.

## Chemical Modifications, Protein Activity, and Longevity

Proteins can be chemically modified with the addition of groups including methyl, phosphate, acetyl, and ubiquitin groups. The addition or removal of these groups from proteins regulates their activity or the length of time they exist in the cell. Sometimes these modifications can regulate where a protein is found in the cell—for example, in the nucleus, the cytoplasm, or attached to the plasma membrane.





Figure 1. Proteins with ubiquitin tags are marked for degradation within the proteasome.

Chemical modifications occur in response to external stimuli such as stress, the lack of nutrients, heat, or ultraviolet light exposure. These changes can alter epigenetic accessibility, transcription, mRNA stability, or translation—all resulting in changes in expression of various genes. This is an efficient way for the cell to rapidly change the levels of specific proteins in response to the environment. Because proteins are involved in every stage of gene regulation, the phosphorylation of a protein (depending on the protein that is modified) can alter accessibility to the chromosome, can alter translation (by altering transcription factor binding or function), can change nuclear shuttling (by influencing modifications to the nuclear pore complex), can alter RNA stability (by binding or not binding to the RNA to regulate its stability), can modify translation (increase or decrease), or can change post-translational modifications (add or remove phosphates or other chemical modifications).

The addition of an ubiquitin group to a protein marks that protein for degradation. Ubiquitin acts like a flag indicating that the protein lifespan is complete. These proteins are moved to the **proteasome**, an organelle that functions to remove proteins, to be degraded (Figure 1). One way to control gene expression, therefore, is to alter the longevity of the protein.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.12: Protein Modifications is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.13: Introduction to Prokaryotic Transcription and Translation

## Outline the process of prokaryotic transcription and translation

The prokaryotes, which include bacteria and archaea, are mostly single-celled organisms that, by definition, lack membrane-bound nuclei and other organelles. A bacterial chromosome is a covalently closed circle that, unlike eukaryotic chromosomes, is not organized around histone proteins. The central region of the cell in which prokaryotic DNA resides is called the nucleoid. In addition, prokaryotes often have abundant **plasmids**, which are shorter circular DNA molecules that may only contain one or a few genes. Plasmids can be transferred independently of the bacterial chromosome during cell division and often carry traits such as antibiotic resistance. Because of these unique features, transcription and gene regulation is somewhat different between prokaryotic cells and eukaryotic ones.

## What You'll Learn to Do

- Understand the basic steps in the transcription of DNA into RNA in prokaryotic cells
- Understand the basics of prokaryotic translation and how it differs from eukaryotic translation

### Learning Activities

The learning activities for this section include the following:

- Prokaryotic Transcription
- Prokaryotic Translation
- Self Check: Prokaryotic Transcription and Translation

## Contributors and Attributions

CC licensed content, Original

- Introduction to Prokaryotic Transcription and Translation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.13: Introduction to Prokaryotic Transcription and Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.14: Prokaryotic Transcription

Prokaryotes do not have membrane-enclosed nuclei. Therefore, the processes of transcription, translation, and mRNA degradation can all occur simultaneously. The intracellular level of a bacterial protein can quickly be amplified by multiple transcription and translation events occurring concurrently on the same DNA template. Prokaryotic transcription often covers more than one gene and produces polycistronic mRNAs that specify more than one protein.

Our discussion here will exemplify transcription by describing this process in *Escherichia coli*, a well-studied bacterial species. Although some differences exist between transcription in *E. coli* and transcription in archaea, an understanding of *E. coli* transcription can be applied to virtually all bacterial species.

## Prokaryotic RNA Polymerase

Prokaryotes use the same RNA polymerase to transcribe all of their genes. In *E. coli*, the polymerase is composed of five polypeptide subunits, two of which are identical. Four of these subunits, denoted $\alpha$, $\alpha$, $\beta$, and $\beta'$ comprise the polymerase **core enzyme**. These subunits assemble every time a gene is transcribed, and they disassemble once transcription is complete. Each subunit has a unique role; the two $\alpha$-subunits are necessary to assemble the polymerase on the DNA; the $\beta$-subunit binds to the ribonucleoside triphosphate that will become part of the nascent "recently born" mRNA molecule; and the $\beta'$ binds the DNA template strand. The fifth subunit, $\sigma$, is involved only in transcription initiation. It confers transcriptional specificity such that the polymerase begins to synthesize mRNA from an appropriate initiation site. Without $\sigma$, the core enzyme would transcribe from random sites and would produce mRNA molecules that specified protein gibberish. The polymerase comprised of all five subunits is called the **holoenzyme** (a holoenzyme is a biochemically active compound comprised of an enzyme and its coenzyme).

## Prokaryotic Promoters





Figure 1. The σ subunit of prokaryotic RNA polymerase recognizes consensus sequences found in the promoter region upstream of the transcription start sight. The σ subunit dissociates from the polymerase after transcription has been initiated.

A **promoter** is a DNA sequence onto which the transcription machinery binds and initiates transcription. In most cases, promoters exist upstream of the genes they regulate. The specific sequence of a promoter is very important because it determines whether the corresponding gene is transcribed all the time, some of the time, or infrequently. Although promoters vary among prokaryotic genomes, a few elements are conserved. At the -10 and -35 regions upstream of the initiation site, there are two promoter **consensus** sequences, or regions that are similar across all promoters and across various bacterial species (Figure 1).

The -10 consensus sequence, called the -10 region, is TATAAT. The -35 sequence, TTGACA, is recognized and bound by $\sigma$. Once this interaction is made, the subunits of the core enzyme bind to the site. The A–T-rich -10 region facilitates unwinding of the DNA template, and several phosphodiester bonds are made. The transcription initiation phase ends with the production of abortive transcripts, which are polymers of approximately 10 nucleotides that are made and released.

View this MolecularMovies animation to see the first part of transcription and the base sequence repetition of the TATA box.

## Elongation and Termination in Prokaryotes

The transcription elongation phase begins with the release of the $\sigma$ subunit from the polymerase. The dissociation of $\sigma$ allows the core enzyme to proceed along the DNA template, synthesizing mRNA in the 5′ to 3′ direction at a rate of approximately 40 nucleotides per second. As elongation proceeds, the DNA is continuously unwound ahead of the core enzyme and rewound behind it (Figure 2). The base pairing between DNA and RNA is not stable enough to maintain the stability of the mRNA synthesis

components. Instead, the RNA polymerase acts as a stable linker between the DNA template and the nascent RNA strands to ensure that elongation is not interrupted prematurely.





Figure 2. Click for a larger image. During elongation, the prokaryotic RNA polymerase tracks along the DNA template, synthesizes mRNA in the 5′ to 3′ direction, and unwinds and rewinds the DNA as it is read.

## Prokaryotic Termination Signals

Once a gene is transcribed, the prokaryotic polymerase needs to be instructed to dissociate from the DNA template and liberate the newly made mRNA. Depending on the gene being transcribed, there are two kinds of termination signals. One is protein-based and the other is RNA-based. **Rho-dependent termination** is controlled by the rho protein, which tracks along behind the polymerase on the growing mRNA chain. Near the end of the gene, the polymerase encounters a run of G nucleotides on the DNA template and it stalls. As a result, the rho protein collides with the polymerase. The interaction with rho releases the mRNA from the transcription bubble.

**Rho-independent termination** is controlled by specific sequences in the DNA template strand. As the polymerase nears the end of the gene being transcribed, it encounters a region rich in C–G nucleotides. The mRNA folds back on itself, and the complementary C–G nucleotides bind together. The result is a stable **hairpin** that causes the polymerase to stall as soon as it begins to transcribe a region rich in A–T nucleotides. The complementary U–A region of the mRNA transcript forms only a weak interaction with the template DNA. This, coupled with the stalled polymerase, induces enough instability for the core enzyme to break away and liberate the new mRNA transcript.

Upon termination, the process of transcription is complete. By the time termination occurs, the prokaryotic transcript would already have been used to begin synthesis of numerous copies of the encoded protein because these processes can occur concurrently. The unification of transcription, translation, and even mRNA degradation is possible because all of these processes occur in the same 5′ to 3′ direction, and because there is no membranous compartmentalization in the prokaryotic cell (Figure 3). In contrast, the presence of a nucleus in eukaryotic cells precludes simultaneous transcription and translation.






Figure 3. Multiple polymerases can transcribe a single bacterial gene while numerous ribosomes concurrently translate the mRNA transcripts into polypeptides. In this way, a specific protein can rapidly reach a high concentration in the bacterial cell.

Visit this BioStudio animation to see the process of prokaryotic transcription.



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=432

## Practice Questions

Which subunit of the *E. coli* polymerase confers specificity to transcription?

a. $\alpha$
b. $\beta$
c. $\beta'$
d. $\sigma$

[reveal-answer q="861545"]Show Answer[/reveal-answer]
[hidden-answer a="861545"]The $\sigma$ subunit of the *E. coli* polymerase confers specificity to transcription.

[/hidden-answer]

The -10 and -35 regions of prokaryotic promoters are called consensus sequences because _____.

a. they are identical in all bacterial species
b. they are similar in all bacterial species
c. they exist in all organisms
d. they have the same function in all organisms

[reveal-answer q="827251"]Show Answer[/reveal-answer]
[hidden-answer a="827251"]The -10 and -35 regions of prokaryotic promoters are called consensus sequences because they are similar in all bacterial species.[/hidden-answer]

## Learning Objectives

In prokaryotes, mRNA synthesis is initiated at a promoter sequence on the DNA template comprising two consensus sequences that recruit RNA polymerase. The prokaryotic polymerase consists of a core enzyme of four protein subunits and a $\sigma$ protein that assists only with initiation. Elongation synthesizes mRNA in the 5′ to 3′ direction at a rate of 40 nucleotides per second. Termination liberates the mRNA and occurs either by rho protein interaction or by the formation of an mRNA hairpin.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.14: Prokaryotic Transcription is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.15: Prokaryotic Translation

Translation is similar in prokaryotes and eukaryotes. Here we will explore how translation occurs in *E. coli*, a representative prokaryote, and specify any differences between bacterial and eukaryotic translation.

## Initiation

The **initiation of protein synthesis** begins with the formation of an initiation complex. In ***E. coli***, this complex involves the small 30S ribosome, the mRNA template, three initiation factors that help the ribosome assemble correctly, guanosine triphosphate (GTP) that acts as an energy source, and a special initiator tRNA carrying ***N*-formyl-methionine** ($fMet\text{-}tRNAf^{Met}$) (Figure 1). The initiator tRNA interacts with the start codon AUG of the mRNA and carries a formylated methionine (fMet). Because of its involvement in initiation, fMet is inserted at the beginning (N terminus) of every polypeptide chain synthesized by *E. coli*. In *E. coli* mRNA, a leader sequence upstream of the first AUG codon, called the **Shine-Dalgarno sequence** (also known as the ribosomal binding site AGGAGG), interacts through complementary base pairing with the rRNA molecules that compose the ribosome. This interaction anchors the 30S ribosomal subunit at the correct location on the mRNA template. At this point, the 50S ribosomal subunit then binds to the initiation complex, forming an intact ribosome.

In eukaryotes, initiation complex formation is similar, with the following differences:

- The initiator tRNA is a different specialized tRNA carrying methionine, called Met-tRNAi
- Instead of binding to the mRNA at the Shine-Dalgarno sequence, the eukaryotic initiation complex recognizes the 5′ cap of the eukaryotic mRNA, then tracks along the mRNA in the 5′ to 3′ direction until the AUG start codon is recognized. At this point, the 60S subunit binds to the complex of Met-tRNAi, mRNA, and the 40S subunit.





Figure 1. Translation in bacteria begins with the formation of the initiation complex, which includes the small ribosomal subunit, the mRNA, the initiator tRNA carrying N-formyl-methionine, and initiation factors. Then the 50S subunit binds, forming an intact ribosome.

## Elongation

In prokaryotes and eukaryotes, the basics of **elongation of translation** are the same. In ***E. coli***, the binding of the 50S ribosomal subunit to produce the intact ribosome forms three functionally important ribosomal sites: The **A (aminoacyl) site** binds incoming charged aminoacyl tRNAs. The **P (peptidyl) site** binds charged tRNAs carrying amino acids that have formed peptide bonds with the growing polypeptide chain but have not yet dissociated from their corresponding tRNA. The **E (exit) site** releases dissociated tRNAs so that they can be recharged with free amino acids. There is one notable exception to this assembly line of tRNAs: During initiation complex formation, bacterial fMet−tRNA$^{fMet}$ or eukaryotic Met-tRNAi enters the P site directly without first entering the A site, providing a free A site ready to accept the tRNA corresponding to the first codon after the AUG.

Elongation proceeds with single-codon movements of the ribosome each called a translocation event. During each translocation event, the charged tRNAs enter at the A site, then shift to the P site, and then finally to the E site for removal. Ribosomal movements, or steps, are induced by conformational changes that advance the ribosome by three bases in the 3′ direction. Peptide bonds form between the amino group of the amino acid attached to the A-site tRNA and the carboxyl group of the amino acid attached to the P-site tRNA. The formation of each peptide bond is catalyzed by **peptidyl transferase**, an RNA-based ribozyme that is integrated into the 50S ribosomal subunit. The amino acid bound to the P-site tRNA is also linked to the growing polypeptide chain. As the ribosome steps across the mRNA, the former P-site tRNA enters the E site, detaches from the amino acid, and is expelled. Several of the steps during elongation, including binding of a charged aminoacyl tRNA to the A site and translocation, require energy derived from GTP hydrolysis, which is catalyzed by specific elongation factors. Amazingly, the *E. coli* translation apparatus takes only 0.05 seconds to add each amino acid, meaning that a 200 amino-acid protein can be translated in just 10 seconds.

## Termination

The **termination of translation** occurs when a **nonsense codon** (UAA, UAG, or UGA) is encountered for which there is no complementary tRNA. On aligning with the A site, these nonsense codons are recognized by release factors in prokaryotes and eukaryotes that result in the P-site amino acid detaching from its tRNA, releasing the newly made polypeptide. The small and large ribosomal subunits dissociate from the mRNA and from each other; they are recruited almost immediately into another translation init iation complex.

In summary, there are several key features that distinguish prokaryotic gene expression from that seen in eukaryotes. These are illustrated in Figure 2 and listed in Table 1.





Figure 2. (a) In prokaryotes, the processes of transcription and translation occur simultaneously in the cytoplasm, allowing for a rapid cellular response to an environmental cue. (b) In eukaryotes, transcription is localized to the nucleus and translation is localized to the cytoplasm, separating these processes and necessitating RNA processing for stability.

| Table 1. Comparison of Translation in Bacteria Versus Eukaryotes | | |
|---|---|---|
| Property | Bacteria | Eukaryotes |
| Ribosomes | 70S<br>• 30S (small subunit) with 16S rNA subunit<br>• 50S (large subunit) with 5S and 23S rRNA subunits | 80S<br>• 40S (small subunit) with 18S rRNA subunit<br>• 60S (large subunit) with 5S, 5.8S, and 28S rRNA subunits |
| Amino acid carried by initiator tRNA | fMet | Met |
| Shine-Dalgarno sequence in mRNA | Present | Absent |
| Simultaneous transcription and translation | Yes | No |

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

12.15: Prokaryotic Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.16: Introduction to the Central Dogma

## Identify the central dogma of life

As you have learned, information flow in an organism takes place from DNA to RNA to protein. DNA dictates the structure of mRNA in a process known as transcription, and RNA dictates the structure of protein in a process known as translation. This is known as the **Central Dogma of Life**, which holds true for all organisms.

### Learning Activities

The learning activities for this section include the following:

- The Central Dogma
- Self Check: Central Dogma

## Contributors and Attributions

CC licensed content, Original

- Introduction to the Central Dogma. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

12.16: Introduction to the Central Dogma is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.17: The Central Dogma

As you have learned, information flow in an organism takes place from DNA to RNA to protein:

- DNA is transcribed to RNA via complementary base pairing rules (but with U instead of T in the transcript)
- The RNA transcript, specifically mRNA, is then translated to an amino acid polypeptide
- Final folding and modifications of the polypeptide lead to functional proteins that actually do things in cells

This is known as the **Central Dogma of Life**, which holds true for all organisms.





Figure 1. Click for a larger image. Instructions on DNA are transcribed onto messenger RNA. Ribosomes are able to read the genetic information inscribed on a strand of messenger RNA and use this information to string amino acids together into a protein.

A link to an interactive elements can be found at the bottom of this page.

Click here for a text-only version of the activity.

## Contributors and Attributions

CC licensed content, Original

- The Central Dogma. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously


- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- The Central Dogma of Life. **Provided by**: Lumen Learning. **Located at**: www.oppia.org/explore/exjRkfVQADh2. **License**: *CC BY-SA: Attribution-ShareAlike*

12.17: The Central Dogma is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 12.18: Putting It Together- DNA Transcription and Translation

As we've seen, transcription and translation are essential processes that enable life. These processes, also called the central dogma, are how our bodies—and all living organisms on earth—are built.

This video compares transcription and translation to something you may be a little more familiar with: creating a Hot Pocket.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=440

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: DNA Transcription and Translation. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

All rights reserved content

- DNA, Hot Pockets, and The Longest Word Ever: Crash Course Biology #11. **Authored by**: CrashCourse. **Located at**: https://youtu.be/itsb2SqR-R0. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

12.18: Putting It Together- DNA Transcription and Translation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 13: Module 11- Gene Expression


13.1: Why It Matters- Gene Expression
13.2: Introduction to Regulation of Gene Expression
13.3: Expression of Genes
13.4: Prokaryotic and Eukaryotic Gene Regulation
13.5: Introduction to Prokaryotic Gene Regulation
13.6: Gene Regulation in Prokaryotes
13.7: Repressors
13.8: Activators and Inducers
13.9: Prokaryotic Gene Regulation at Work
13.10: Introduction to Eukaryotic Gene Regulation
13.11: Eukaryotic Epigenetic Gene Regulation
13.12: Eukaryotic Transcription Gene Regulation
13.13: Post-Transcriptional Control of Gene Expression
13.14: Putting It Together- Gene Expression
13.15: Assignment- How Mutations Work



13: Module 11- Gene Expression is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.1: Why It Matters- Gene Expression

## Why explain the regulation of gene expression?

Cancer is one of the top ten causes of death in the United States. It is not a single disease but includes many different diseases. In cancer cells, mutations modify cell-cycle control, and cells don't stop growing as they normally would. Mutations can also alter the growth rate or the progression of the cell through the cell cycle.

Thus, cancer can be described as a disease of altered gene expression. Changes at every level of eukaryotic gene expression can be detected in some form of cancer at some point in time. In order to understand how changes to gene expression can cause cancer, it is critical to understand how each stage of gene regulation works in normal cells. By understanding the mechanisms of control in normal, non-diseased cells, it will be easier for scientists to understand what goes wrong in disease states including complex ones like cancer.

### Learning Outcomes

- Explain how DNA stores genetic information
- Explain the role of complementary base pairing in the precise replication process of DNA
- Recognize the impact of DNA mutations

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Gene Expression. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.1: Why It Matters- Gene Expression is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.2: Introduction to Regulation of Gene Expression

## Define the term regulation as it applies to genes

For a cell to function properly, necessary proteins must be synthesized at the proper time. All cells control or regulate the synthesis of proteins from information encoded in their DNA. The process of turning on a gene to produce RNA and protein is called **gene expression**. Whether in a simple unicellular organism or a complex multi-cellular organism, each cell controls when and how its genes are expressed. For this to occur, there must be a mechanism to control when a gene is expressed to make RNA and protein, how much of the protein is made, and when it is time to stop making that protein because it is no longer needed.

The regulation of gene expression conserves energy and space. It would require a significant amount of energy for an organism to express every gene at all times, so it is more energy efficient to turn on the genes only when they are required. In addition, only expressing a subset of genes in each cell saves space because DNA must be unwound from its tightly coiled structure to transcribe and translate the DNA. Cells would have to be enormous if every protein were expressed in every cell all the time.

The control of gene expression is extremely complex. Malfunctions in this process are detrimental to the cell and can lead to the development of many diseases, including cancer.

## What You'll Learn to Do

- Discuss why every cell does not express all of its genes
- Compare prokaryotic and eukaryotic gene regulation

### Learning Activities

The learning activities for this section include the following:

- Expression of Genes
- Prokaryotic and Eukaryotic Gene Regulation
- Self Check: Regulation of Gene Expression

## Contributors and Attributions

CC licensed content, Original

- Outcome: Regulation of Gene Expression. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.2: Introduction to Regulation of Gene Expression is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.3: Expression of Genes

For a cell to function properly, necessary proteins must be synthesized at the proper time. All cells control or regulate the synthesis of proteins from information encoded in their DNA. The process of turning on a gene to produce RNA and protein is called **gene expression**. Whether in a simple unicellular organism or a complex multi-cellular organism, each cell controls when and how its genes are expressed. For this to occur, there must be a mechanism to control when a gene is expressed to make RNA and protein, how much of the protein is made, and when it is time to stop making that protein because it is no longer needed.

The regulation of gene expression conserves energy and space. It would require a significant amount of energy for an organism to express every gene at all times, so it is more energy efficient to turn on the genes only when they are required. In addition, only expressing a subset of genes in each cell saves space because DNA must be unwound from its tightly coiled structure to transcribe and translate the DNA. Cells would have to be enormous if every protein were expressed in every cell all the time.

The control of gene expression is extremely complex. Malfunctions in this process are detrimental to the cell and can lead to the development of many diseases, including cancer.

## Gene regulation makes cells different

**Gene regulation** is how a cell controls which genes, out of the many genes in its genome, are "turned on" (expressed). Thanks to gene regulation, each cell type in your body has a different set of active genes—despite the fact that almost all the cells of your body contain the exact same DNA. These different patterns of gene expression cause your various cell types to have different sets of proteins, making each cell type uniquely specialized to do its job.

For example, one of the jobs of the liver is to remove toxic substances like alcohol from the bloodstream. To do this, liver cells express genes encoding subunits (pieces) of an enzyme called alcohol dehydrogenase. This enzyme breaks alcohol down into a non-toxic molecule. The neurons in a person's brain don't remove toxins from the body, so they keep these genes unexpressed, or "turned off." Similarly, the cells of the liver don't send signals using neurotransmitters, so they keep neurotransmitter genes turned off (Figure 1).





Figure 1. Different cells have different genes "turned on."

There are many other genes that are expressed differently between liver cells and neurons (or any two cell types in a multicellular organism like yourself).

## How do cells “decide” which genes to turn on?

Now there’s a tricky question! Many factors that can affect which genes a cell expresses. Different cell types express different sets of genes, as we saw above. However, two different cells of the same type may also have different gene expression patterns depending on their environment and internal state.

Broadly speaking, we can say that a cell’s gene expression pattern is determined by information from both inside and outside the cell.

- Examples of information from *inside* the cell: the proteins it inherited from its mother cell, whether its DNA is damaged, and how much ATP it has.
- Examples of information from *outside* the cell: chemical signals from other cells, mechanical signals from the extracellular matrix, and nutrient levels.

How do these cues help a cell “decide” what genes to express? Cells don’t make decisions in the sense that you or I would. Instead, they have molecular pathways that convert information—such as the binding of a chemical signal to its receptor—into a change in gene expression.

As an example, let’s consider how cells respond to growth factors. A growth factor is a chemical signal from a neighboring cell that instructs a target cell to grow and divide. We could say that the cell “notices” the growth factor and “decides” to divide, but how do these processes actually occur?





Figure 2. Growth factor prompting cell division

- The cell detects the growth factor through physical binding of the growth factor to a receptor protein on the cell surface.
- Binding of the growth factor causes the receptor to change shape, triggering a series of chemical events in the cell that activate proteins called transcription factors.
- The transcription factors bind to certain sequences of DNA in the nucleus and cause transcription of cell division-related genes.
- The products of these genes are various types of proteins that make the cell divide (drive cell growth and/or push the cell forward in the cell cycle).

This is just one example of how a cell can convert a source of information into a change in gene expression. There are many others, and understanding the logic of gene regulation is an area of ongoing research in biology today.

Growth factor signaling is complex and involves the activation of a variety of targets, including both transcription factors and non-transcription factor proteins.

Learning Objectives

- Gene regulation is the process of controlling which genes in a cell's DNA are expressed (used to make a functional product such as a protein).
- Different cells in a multicellular organism may express very different sets of genes, even though they contain the same DNA.
- The set of genes expressed in a cell determines the set of proteins and functional RNAs it contains, giving it its unique properties.
- In eukaryotes like humans, gene expression involves many steps, and gene regulation can occur at any of these steps. However, many genes are regulated primarily at the level of transcription.

[reveal-answer q="915423"]Show References[/reveal-answer]
[hidden-answer a="915423"]

Alcohol dehydrogenase. (2016, January 6). Retrieved April 26, 2016 from Wikipedia: https://en.Wikipedia.org/wiki/Alcohol_dehydrogenase.

Cooper, G. M. (2000). Regulation of transcription in eukaryotes. In *The cell: A molecular approach*. Sunderland, MA: Sinauer Associates. Retrieved from www.ncbi.nlm.nih.gov/books/NBK9904/.

Kimball, John W. (2014, April 19). The human and chimpanzee genomes. In *Kimball's biology pages*. Retrieved from http://www.biology-pages.info/H/HominoidClade.html.

OpenStax College, Biology. (2016, March 23). Eukaryotic transcription gene regulation. In _OpenStax CNX. Retrieved from http://cnx.org/contents/GFy_h8cu@10....cription-Gene-.

OpenStax College, Biology. (2016, March 23). Regulation of gene expression. In _OpenStax CNX. Retrieved from http://cnx.org/contents/GFy_h8cu@10....ene-Expression

Phillips, T. (2008). Regulation of transcription and gene expression in eukaryotes. *Nature Education*, *1*(1), 199. Retrieved from http://www.nature.com/scitable/topic...ession-in-1086.

Purves, W. K., Sadava, D. E., Orians, G. H., and Heller, H.C. (2003). Transcriptional regulation of gene expression. In *Life: The science of biology* (7th ed., pp. 290-296). Sunderland, MA: Sinauer Associates.

Reece, J. B., Urry, L. A., Cain, M. L., Wasserman, S. A., Minorsky, P. V., and Jackson, R. B. (2011). Eukaryotic gene expression is regulated at many stages. In *Campbell Biology* (10th ed., pp. 365-373). San Francisco, CA: Pearson.

[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Overview: Eukaryotic gene regulation. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/gene-regulation/gene-regulation-in-eukaryotes/a/overview-of-eukaryotic-gene-regulation. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

13.3: Expression of Genes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.4: Prokaryotic and Eukaryotic Gene Regulation

To understand how gene expression is regulated, we must first understand how a gene codes for a functional protein in a cell. The process occurs in both prokaryotic and eukaryotic cells, just in slightly different manners.

Prokaryotic organisms are single-celled organisms that lack a cell nucleus, and their DNA therefore floats freely in the cell cytoplasm. To synthesize a protein, the processes of transcription and translation occur almost simultaneously. When the resulting protein is no longer needed, transcription stops. As a result, the primary method to control what type of protein and how much of each protein is expressed in a prokaryotic cell is the regulation of DNA transcription. All of the subsequent steps occur automatically. When more protein is required, more transcription occurs. Therefore, in prokaryotic cells, the control of gene expression is mostly at the transcriptional level.

Eukaryotic cells, in contrast, have intracellular organelles that add to their complexity. In eukaryotic cells, the DNA is contained inside the cell's nucleus and there it is transcribed into RNA. The newly synthesized RNA is then transported out of the nucleus into the cytoplasm, where ribosomes translate the RNA into protein. The processes of transcription and translation are physically separated by the nuclear membrane; transcription occurs only within the nucleus, and translation occurs only outside the nucleus in the cytoplasm. The regulation of gene expression can occur at all stages of the process (Figure 1). Regulation may occur when the DNA is uncoiled and loosened from nucleosomes to bind transcription factors (**epigenetic** level), when the RNA is transcribed (transcriptional level), when the RNA is processed and exported to the cytoplasm after it is transcribed (**post-transcriptional** level), when the RNA is translated into protein (translational level), or after the protein has been made (**post-translational** level).





Figure 1. Prokaryotic transcription and translation occur simultaneously in the cytoplasm, and regulation occurs at the transcriptional level. Eukaryotic gene expression is regulated during transcription and RNA processing, which take place in the nucleus, and during protein translation, which takes place in the cytoplasm. Further regulation may occur through post-translational modifications of proteins.

The differences in the regulation of gene expression between prokaryotes and eukaryotes are summarized in Table 1. The regulation of gene expression is discussed in detail in subsequent modules.

| Table 1. Differences in the Regulation of Gene Expression of Prokaryotic and Eukaryotic Organisms | |
|---|---|
| Prokaryotic organisms | Eukaryotic organisms |
| Lack nucleus | Contain nucleus |
| DNA is found in the cytoplasm | DNA is confined to the nuclear compartment |
| RNA transcription and protein formation occur almost simultaneously | RNA transcription occurs prior to protein formation, and it takes place in the nucleus. Translation of RNA to protein occurs in the cytoplasm. |
| Gene expression is regulated primarily at the transcriptional level | Gene expression is regulated at many levels (epigenetic, transcriptional, nuclear shuttling, post-transcriptional, translational, and post-translational) |

## Evolution of Gene Regulation

Prokaryotic cells can only regulate gene expression by controlling the amount of transcription. As eukaryotic cells evolved, the complexity of the control of gene expression increased. For example, with the evolution of eukaryotic cells came compartmentalization of important cellular components and cellular processes. A nuclear region that contains the DNA was formed. Transcription and translation were physically separated into two different cellular compartments. It therefore became possible to control gene expression by regulating transcription in the nucleus, and also by controlling the RNA levels and protein translation present outside the nucleus.

Some cellular processes arose from the need of the organism to defend itself. Cellular processes such as gene silencing developed to protect the cell from viral or parasitic infections. If the cell could quickly shut off gene expression for a short period of time, it would be able to survive an infection when other organisms could not. Therefore, the organism evolved a new process that helped it survive, and it was able to pass this new development to offspring.

## Practice Questions

Control of gene expression in eukaryotic cells occurs at which level(s)?

a. only the transcriptional level
b. epigenetic and transcriptional levels
c. epigenetic, transcriptional, and translational levels
d. epigenetic, transcriptional, post-transcriptional, translational, and post-translational levels

[reveal-answer q=”386227″]Show Answer[/reveal-answer]
[hidden-answer a=”386227″]Answer d. Control of gene expression in eukaryotic cells occurs at epigenetic, transcriptional, post-transcriptional, translational, and post-translational levels.

[/hidden-answer]

Post-translational control refers to the:

a. regulation of gene expression after transcription
b. regulation of gene expression after translation
c. control of epigenetic activation
d. period between transcription and translation

[reveal-answer q=”960802″]Show Answer[/reveal-answer]
[hidden-answer a=”960802″]Answer b. Post-translational control refers to the regulation of gene expression after translation[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

13.4: Prokaryotic and Eukaryotic Gene Regulation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.5: Introduction to Prokaryotic Gene Regulation

## Discuss different components of prokaryotic gene regulation

The DNA of prokaryotes is organized into a circular chromosome supercoiled in the nucleoid region of the cell cytoplasm. Proteins that are needed for a specific function are encoded together in blocks called **operons**. For example, all of the genes needed to use lactose as an energy source are coded next to each other in the lactose (or lac) operon.

In prokaryotic cells, there are three types of regulatory molecules that can affect the expression of operons: repressors, activators, and inducers. **Repressors** are proteins that suppress transcription of a gene in response to an external stimulus, whereas **activators** are proteins that increase the transcription of a gene in response to an external stimulus. Finally, **inducers** are small molecules that either activate or repress transcription depending on the needs of the cell and the availability of substrate.

## What You'll Learn to Do

- Understand the basic steps in gene regulation in prokaryotic cells
- Explain the roles of repressors in negative gene regulation
- Explain the role of activators and inducers in positive gene regulation

### Learning Activities

The learning activities for this section include the following:

- Gene Regulation in Prokaryotes
- Repressors
- Activators and Inducers
- Self Check: Prokaryotic Gene Regulation

## Contributors and Attributions

CC licensed content, Original

- Outcome: Prokaryotic Gene Regulation. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.5: Introduction to Prokaryotic Gene Regulation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.6: Gene Regulation in Prokaryotes

In bacteria and archaea, structural proteins with related functions—such as the genes that encode the enzymes that catalyze the many steps in a single biochemical pathway—are usually encoded together within the genome in a block called an **operon** and are transcribed together under the control of a single **promoter.** This forms a polycistronic transcript (Figure 1). The promoter then has simultaneous control over the regulation of the transcription of these structural genes because they will either all be needed at the same time, or none will be needed.





Figure 1. In prokaryotes, structural genes of related function are often organized together on the genome and transcribed together under the control of a single promoter. The operon's regulatory region includes both the promoter and the operator. If a repressor binds to the operator, then the structural genes will not be transcribed. Alternatively, activators may bind to the regulatory region, enhancing transcription.

French scientists François **Jacob** (1920–2013) and Jacques **Monod** at the Pasteur Institute were the first to show the organization of bacterial genes into operons, through their studies on the ***lac* operon** of *E. coli*. They found that in *E. coli*, all of the structural genes that encode enzymes needed to use lactose as an energy source lie next to each other in the lactose (or *lac*) operon under the control of a single promoter, the *lac* promoter. For this work, they won the Nobel Prize in Physiology or Medicine in 1965.

Although eukaryotic genes are not organized into operons, prokaryotic operons are excellent models for learning about gene regulation generally. There are some gene clusters in eukaryotes that function similar to operons. Many of the principles can be applied to eukaryotic systems and contribute to our understanding of changes in gene expression in eukaryotes that can result pathological changes such as cancer.

Each operon includes DNA sequences that influence its own transcription; these are located in a region called the regulatory region. The regulatory region includes the promoter and the region surrounding the promoter, to which **transcription factors**, proteins encoded by regulatory genes, can bind. Transcription factors influence the binding of **RNA polymerase** to the promoter and allow its progression to transcribe structural genes. A **repressor** is a transcription factor that suppresses transcription of a gene in response to an external stimulus by binding to a DNA sequence within the regulatory region called the **operator**, which is located between the RNA polymerase binding site of the promoter and the transcriptional start site of the first structural gene. Repressor binding physically blocks RNA polymerase from transcribing structural genes. Conversely, an **activator** is a transcription factor that increases the transcription of a gene in response to an external stimulus by facilitating RNA polymerase binding to the promoter. An **inducer**, a third type of regulatory molecule, is a small molecule that either activates or represses transcription by interacting with a repressor or an activator.

In prokaryotes, there are examples of operons whose gene products are required rather consistently and whose expression, therefore, is unregulated. Such operons are **constitutively expressed**, meaning they are transcribed and translated continuously to provide the cell with constant intermediate levels of the protein products. Such genes encode enzymes involved in housekeeping functions required for cellular maintenance, including DNA replication, repair, and expression, as well as enzymes involved in core

metabolism. In contrast, there are other prokaryotic operons that are expressed only when needed and are regulated by repressors, activators, and inducers.

## Practice Questions

What are the parts in the DNA sequence of an operon?

[practice-area rows="2"][/practice-area]
[reveal-answer q="665976"]Show Answer[/reveal-answer]
[hidden-answer a="665976"]An operon is composed of a promoter, an operator, and the structural genes. They must occur in that order.

[/hidden-answer]

What types of regulatory molecules are there?

[practice-area rows="2"][/practice-area]
[reveal-answer q="665979"]Show Answer[/reveal-answer]
[hidden-answer a="665979"]There are three types of regulatory molecules: repressors, activators, and inducers.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.6: Gene Regulation in Prokaryotes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.7: Repressors

## Tryptophan Operon: A Repressor Operon

Bacteria such as *E. coli* need amino acids to survive. **Tryptophan** is one such amino acid that *E. coli* can ingest from the environment. *E. coli* can also synthesize tryptophan using enzymes that are encoded by five genes. These five genes are next to each other in what is called the **tryptophan (*trp*) operon** (Figure 1). If tryptophan is present in the environment, then *E. coli* does not need to synthesize it and the switch controlling the activation of the genes in the *trp* operon is switched off. However, when tryptophan availability is low, the switch controlling the operon is turned on, transcription is initiated, the genes are expressed, and tryptophan is synthesized.





Figure 1. The five genes that are needed to synthesize tryptophan in *E. coli* are located next to each other in the *trp* operon. When tryptophan is plentiful, two tryptophan molecules bind the repressor protein at the operator sequence. This physically blocks the RNA polymerase from transcribing the tryptophan genes. When tryptophan is absent, the repressor protein does not bind to the operator and the genes are transcribed.

A DNA sequence that codes for proteins is referred to as the coding region. The five coding regions for the tryptophan biosynthesis enzymes are arranged sequentially on the chromosome in the operon. Just before the coding region is the **transcriptional start site**. This is the region of DNA to which RNA polymerase binds to initiate transcription. The promoter sequence is upstream of the transcriptional start site; each operon has a sequence within or near the promoter to which proteins (activators or repressors) can bind and regulate transcription.

A DNA sequence called the operator sequence is encoded between the promoter region and the first *trp* coding gene. This **operator** contains the DNA code to which the repressor protein can bind. When tryptophan is present in the cell, two tryptophan molecules bind to the *trp* repressor, which changes shape to bind to the *trp* operator. Binding of the tryptophan–repressor complex at the operator physically prevents the RNA polymerase from binding, and transcribing the downstream genes.

When tryptophan is not present in the cell, the repressor by itself does not bind to the operator; therefore, the operon is active and tryptophan is synthesized. Because the repressor protein actively binds to the operator to keep the genes turned off, the *trp* operon is negatively regulated and the proteins that bind to the operator to silence *trp* expression are **negative regulators**.

Watch this video to learn more about the *trp* operon.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for “operon Trp” here (opens in new window).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

13.7: Repressors is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.8: Activators and Inducers

## Catabolite Activator Protein (CAP): An Activator Regulator

Just as the *trp* operon is negatively regulated by tryptophan molecules, there are proteins that bind to the operator sequences that act as a **positive regulator** to turn genes on and activate them. For example, when glucose is scarce, *E. coli* bacteria can turn to other sugar sources for fuel. To do this, new genes to process these alternate sugars must be transcribed. When glucose levels drop, cyclic AMP (cAMP) begins to accumulate in the cell. The cAMP molecule is a signaling molecule that is involved in glucose and energy metabolism in *E. coli*. When glucose levels decline in the cell, accumulating cAMP binds to the positive regulator **catabolite activator protein (CAP)**, a protein that binds to the promoters of operons that control the processing of alternative sugars. When cAMP binds to CAP, the complex binds to the promoter region of the genes that are needed to use the alternate sugar sources (Figure 1). In these operons, a CAP binding site is located upstream of the RNA polymerase binding site in the promoter. This increases the binding ability of RNA polymerase to the promoter region and the transcription of the genes.






Figure 1. When glucose levels fall, *E. coli* may use other sugars for fuel but must transcribe new genes to do so. As glucose supplies become limited, cAMP levels increase. This cAMP binds to the CAP protein, a positive regulator that binds to an operator region upstream of the genes required to use other sugar sources.

## Lactose Operon: An Inducer Operon

The third type of gene regulation in prokaryotic cells occurs through **inducible operons**, which have proteins that bind to activate or repress transcription depending on the local environment and the needs of the cell. The *lac* operon is a typical inducible operon. As mentioned previously, *E. coli* is able to use other sugars as energy sources when glucose concentrations are low. To do so, the cAMP–CAP protein complex serves as a positive regulator to induce transcription. One such sugar source is lactose. The ***lac* operon** encodes the genes necessary to acquire and process the lactose from the local environment. CAP binds to the operator sequence upstream of the promoter that initiates transcription of the *lac* operon. However, for the *lac* operon to be activated, two conditions must be met. First, the level of glucose must be very low or non-existent. Second, lactose must be present. Only when glucose is absent and lactose is present will the *lac* operon be transcribed. This makes sense for the cell, because it would be energetically wasteful to create the proteins to process lactose if glucose was plentiful or lactose was not available.

**Practice Question**

Transcription of the *lac* operon is carefully regulated so that its expression only occurs when glucose is limited and lactose is present to serve as an alternative fuel source.




In *E. coli*, the *trp* operon is on by default, while the *lac* operon is off. Why do you think this is the case?

[practice-area rows="2"][/practice-area]
[reveal-answer q="703127"]Show Answer[/reveal-answer]
[hidden-answer a="703127"]Tryptophan is an amino acid essential for making proteins, so the cell always needs to have some on hand. However, if plenty of tryptophan is present, it is wasteful to make more, and the expression of the *trp* receptor is repressed. Lactose, a sugar found in milk, is not always available. It makes no sense to make the enzymes necessary to digest an energy source that is not available, so the lac operon is only turned on when lactose is present.[/hidden-answer]

If glucose is absent, then CAP can bind to the operator sequence to activate transcription. If lactose is absent, then the repressor binds to the operator to prevent transcription. If either of these requirements is met, then transcription remains off. Only when both conditions are satisfied is the *lac* operon transcribed (Table 1).

Table 1. Signals that Induce or Repress Transcription of the *lac* Operon

| Glucose | CAP binds | Lactose | Repressor binds | Transcription |
|---|---|---|---|---|
| + | – | – | + | No |
| + | – | + | – | Some |
| – | + | – | + | No |
| – | + | + | – | Yes |

Watch an animated tutorial about the workings of *lac* operon here.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for “Operon LAC” here (link opens in new window).

## Practice Questions

If glucose is absent, but so is lactose, the lac operon will be _____.

1. activated
2. repressed
3. activated, but only partially
4. mutated

[reveal-answer q=”88859″]Show Answer[/reveal-answer]
[hidden-answer a=”88859″]Answer b. If glucose is absent, but so is lactose, the lac operon will be repressed.

[/hidden-answer]

Describe how transcription in prokaryotic cells can be altered by external stimulation such as excess lactose in the environment.

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”742266″]Show Answer[/reveal-answer]
[hidden-answer a=”742266″]Environmental stimuli can increase or induce transcription in prokaryotic cells. In this example, lactose in the environment will induce the transcription of the *lac* operon, but only if glucose is not available in the environment.

[/hidden-answer]

What is the difference between a repressible and an inducible operon?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”353005″]Show Answer[/reveal-answer]
[hidden-answer a=”353005″]A repressible operon uses a protein bound to the promoter region of a gene to keep the gene repressed or silent. This repressor must be actively removed in order to transcribe the gene. An inducible operon is either activated or repressed depending on the needs of the cell and what is available in the local environment.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.8: Activators and Inducers is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.9: Prokaryotic Gene Regulation at Work

As we've just learned, there are three types of regulatory molecules that can affect the expression of operons: repressors, activators, and inducers.

- **Repressors** are proteins that suppress transcription of a gene in response to an external stimulus. In other words, a repressor keeps a gene "off."
- **Activators** are proteins that increase the transcription of a gene in response to an external stimulus. In other words, an activator turns a gene "on."
- **Inducers** are small molecules that either activate or repress transcription depending on the needs of the cell and the availability of substrate. Inducers basically help speed up or slow down "on" or "off" by binding to a repressor or activator. In other words: they don't work alone.

In the interactive below, we will focus on the differences between activators and repressors:

A link to an interactive elements can be found at the bottom of this page.

Click here for a text-only version of the activity.

## Contributors and Attributions

CC licensed content, Original

- Prokaryotic Gene Regulation at Work. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Activators and Repressors. **Provided by**: Lumen Learning. **Located at**: www.oppia.org/explore/5HfYP2FK5YCD. **License**: *CC BY-SA: Attribution-ShareAlike*

13.9: Prokaryotic Gene Regulation at Work is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.10: Introduction to Eukaryotic Gene Regulation

## Discuss different components and types of epigenetic gene regulation

Eukaryotic gene expression is more complex than prokaryotic gene expression because the processes of transcription and translation are physically separated. Unlike prokaryotic cells, eukaryotic cells can regulate gene expression at many different levels. Eukaryotic gene expression begins with control of access to the DNA. This form of regulation, called epigenetic regulation, occurs even before transcription is initiated.

## What You'll Learn to Do

- Explain the process of epigenetic regulation
- Discuss the role of transcription factors in gene regulation
- Understand RNA splicing and explain its role in regulating gene expression
- Describe the importance of RNA stability in gene regulation

### Learning Activities

The learning activities for this section include the following:

- Eukaryotic Epigenetic Gene Regulation
- Eukaryotic Transcription Gene Regulation
- Post-Translational Control of Gene Expression
- Self Check: Eukaryotic Gene Regulation

## Contributors and Attributions

CC licensed content, Original

- Outcome: Eukaryotic Gene Regulation. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.10: Introduction to Eukaryotic Gene Regulation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.11: Eukaryotic Epigenetic Gene Regulation

The human genome encodes over 20,000 genes; each of the 23 pairs of human chromosomes encodes thousands of genes. The DNA in the nucleus is precisely wound, folded, and compacted into chromosomes so that it will fit into the nucleus. It is also organized so that specific segments can be accessed as needed by a specific cell type.

The first level of organization, or packing, is the winding of DNA strands around histone proteins. Histones package and order DNA into structural units called nucleosome complexes, which can control the access of proteins to the DNA regions (Figure 1a). Under the electron microscope, this winding of DNA around histone proteins to form nucleosomes looks like small beads on a string (Figure 1b). These beads (histone proteins) can move along the string (DNA) and change the structure of the molecule.





Figure 1. DNA is folded around histone proteins to create (a) nucleosome complexes. These nucleosomes control the access of proteins to the underlying DNA. When viewed through an electron microscope (b), the nucleosomes look like beads on a string. (credit “micrograph”: modification of work by Chris Woodcock)

If DNA encoding a specific gene is to be transcribed into RNA, the nucleosomes surrounding that region of DNA can slide down the DNA to open that specific chromosomal region and allow for the transcriptional machinery (RNA polymerase) to initiate transcription (Figure 2). Nucleosomes can move to open the chromosome structure to expose a segment of DNA, but do so in a very controlled manner.

## Practice Question





Figure 2. Nucleosomes can slide along DNA. When nucleosomes are spaced closely together (top), transcription factors cannot bind and gene expression is turned off. When the nucleosomes are spaced far apart (bottom), the DNA is exposed. Transcription factors can bind, allowing gene expression to occur. Modifications to the histones and DNA affect nucleosome spacing.

In females, one of the two X chromosomes is inactivated during embryonic development because of epigenetic changes to the chromatin. What impact do you think these changes would have on nucleosome packing?

[practice-area rows="2"][/practice-area]
[reveal-answer q="670204"]Show Answer[/reveal-answer]
[hidden-answer a="670204"]The nucleosomes would pack more tightly together.[/hidden-answer]

How the histone proteins move is dependent on signals found on both the histone proteins and on the DNA. These signals are tags added to histone proteins and DNA that tell the histones if a chromosomal region should be open or closed (Figure 3 depicts modifications to histone proteins and DNA). These tags are not permanent, but may be added or removed as needed. They are chemical modifications (phosphate, methyl, or acetyl groups) that are attached to specific amino acids in the protein or to the nucleotides of the DNA. The tags do not alter the DNA base sequence, but they do alter how tightly wound the DNA is around the histone proteins. DNA is a negatively charged molecule; therefore, changes in the charge of the histone will change how tightly wound the DNA molecule will be. When unmodified, the histone proteins have a large positive charge; by adding chemical modifications like acetyl groups, the charge becomes less positive.

The DNA molecule itself can also be modified. This occurs within very specific regions called CpG islands. These are stretches with a high frequency of cytosine and guanine dinucleotide DNA pairs (CG) found in the promoter regions of genes. When this configuration exists, the cytosine member of the pair can be methylated (a methyl group is added). This modification changes how the DNA interacts with proteins, including the histone proteins that control access to the region. Highly methylated (hypermethylated) DNA regions with deacetylated histones are tightly coiled and transcriptionally inactive.





Figure 3. Histone proteins and DNA nucleotides can be modified chemically. Modifications affect nucleosome spacing and gene expression. (credit: modification of work by NIH)

This type of gene regulation is called epigenetic regulation. Epigenetic means “around genetics.” The changes that occur to the histone proteins and DNA do not alter the nucleotide sequence and are not permanent. Instead, these changes are temporary (although they often persist through multiple rounds of cell division) and alter the chromosomal structure (open or closed) as needed. A gene can be turned on or off depending upon the location and modifications to the histone proteins and DNA. If a gene is to be transcribed, the histone proteins and DNA are modified surrounding the chromosomal region encoding that gene. This opens the chromosomal region to allow access for RNA polymerase and other proteins, called transcription factors, to bind to the promoter region, located just upstream of the gene, and initiate transcription. If a gene is to remain turned off, or silenced, the histone proteins and DNA have different modifications that signal a closed chromosomal configuration. In this closed configuration, the RNA polymerase and transcription factors do not have access to the DNA and transcription cannot occur (Figure 2).

View this video that describes how epigenetic regulation controls gene expression.

A link to an interactive elements can be found at the bottom of this page.

## In Summary: Eukaryotic Epigenetic Gene Regulation

In eukaryotic cells, the first stage of gene expression control occurs at the epigenetic level. Epigenetic mechanisms control access to the chromosomal region to allow genes to be turned on or off. These mechanisms control how DNA is packed into the nucleus by regulating how tightly the DNA is wound around histone proteins. The addition or removal of chemical modifications (or flags) to histone proteins or DNA signals to the cell to open or close a chromosomal region. Therefore, eukaryotic cells can control whether a gene is expressed by controlling accessibility to transcription factors and the binding of RNA polymerase to initiate transcription.

## Contributors and Attributions

CC licensed content, Shared previously


- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.11: Eukaryotic Epigenetic Gene Regulation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.12: Eukaryotic Transcription Gene Regulation

Like prokaryotic cells, the transcription of genes in eukaryotes requires the actions of an RNA polymerase to bind to a sequence upstream of a gene to initiate transcription. However, unlike prokaryotic cells, the eukaryotic RNA polymerase requires other proteins, or transcription factors, to facilitate transcription initiation. Transcription factors are proteins that bind to the promoter sequence and other regulatory sequences to control the transcription of the target gene. RNA polymerase by itself cannot initiate transcription in eukaryotic cells. Transcription factors must bind to the promoter region first and recruit RNA polymerase to the site for transcription to be established.

View the process of transcription—the making of RNA from a DNA template:



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=466

## Promoter and the Transcription Machinery

Genes are organized to make the control of gene expression easier. The promoter region is immediately upstream of the coding sequence. This region can be short (only a few nucleotides in length) or quite long (hundreds of nucleotides long). The longer the promoter, the more available space for proteins to bind. This also adds more control to the transcription process. The length of the promoter is gene-specific and can differ dramatically between genes. Consequently, the level of control of gene expression can also differ quite dramatically between genes. The purpose of the promoter is to bind transcription factors that control the initiation of transcription.





Figure 1. An enhancer is a DNA sequence that promotes transcription. Each enhancer is made up of short DNA sequences called distal control elements. Activators bound to the distal control elements interact with mediator proteins and transcription factors. Two different genes may have the same promoter but different distal control elements, enabling differential gene expression.

Within the promoter region, just upstream of the transcriptional start site, resides the TATA box. This box is simply a repeat of thymine and adenine dinucleotides (literally, TATA repeats). RNA polymerase binds to the transcription initiation complex, allowing transcription to occur. To initiate transcription, a transcription factor (TFIID) is the first to bind to the TATA box. Binding of TFIID recruits other transcription factors, including TFIIB, TFIIE, TFIIF, and TFIIH to the TATA box. Once this complex is assembled, RNA polymerase can bind to its upstream sequence. When bound along with the transcription factors, RNA polymerase is phosphorylated. This releases part of the protein from the DNA to activate the transcription initiation complex and places RNA polymerase in the correct orientation to begin transcription; DNA-bending protein brings the enhancer, which can be quite a distance from the gene, in contact with transcription factors and mediator proteins (Figure 1).

In addition to the general transcription factors, other transcription factors can bind to the promoter to regulate gene transcription. These transcription factors bind to the promoters of a specific set of genes. They are not general transcription factors that bind to every promoter complex, but are recruited to a specific sequence on the promoter of a specific gene. There are hundreds of transcription factors in a cell that each bind specifically to a particular DNA sequence motif. When transcription factors bind to the promoter just upstream of the encoded gene, it is referred to as a *cis*-acting element, because it is on the same chromosome just next to the gene. The region that a particular transcription factor binds to is called the transcription factor binding site. Transcription factors respond to environmental stimuli that cause the proteins to find their binding sites and initiate transcription of the gene that is needed.

## Enhancers and Transcription

In some eukaryotic genes, there are regions that help increase or enhance transcription. These regions, called enhancers, are not necessarily close to the genes they enhance. They can be located upstream of a gene, within the coding region of the gene, downstream of a gene, or may be thousands of nucleotides away.

Enhancer regions are binding sequences, or sites, for transcription factors. When a DNA-bending protein binds, the shape of the DNA changes (Figure 1). This shape change allows for the interaction of the activators bound to the enhancers with the transcription factors bound to the promoter region and the RNA polymerase. Whereas DNA is generally depicted as a straight line in two dimensions, it is actually a three-dimensional object. Therefore, a nucleotide sequence thousands of nucleotides away can fold over and interact with a specific promoter.

## Turning Genes Off: Transcriptional Repressors

Like prokaryotic cells, eukaryotic cells also have mechanisms to prevent transcription. Transcriptional repressors can bind to promoter or enhancer regions and block transcription. Like the transcriptional activators, repressors respond to external stimuli to prevent the binding of activating transcription factors.

### Learning Objectives

To start transcription, general transcription factors, such as TFIID, TFIIH, and others, must first bind to the TATA box and recruit RNA polymerase to that location. The binding of additional regulatory transcription factors to *cis*-acting elements will either increase or prevent transcription. In addition to promoter sequences, enhancer regions help augment transcription. Enhancers can be upstream, downstream, within a gene itself, or on other chromosomes. Transcription factors bind to enhancer regions to increase or prevent transcription.

### Practice Questions

The binding of _____ is required for transcription to start.

a. a protein
b. DNA polymerase
c. RNA polymerase
d. a transcription factor

[reveal-answer q="670222"]Show Answer[/reveal-answer]
[hidden-answer a="670222"]Answer c. The binding of RNA polymerase is required for transcription to start.

[/hidden-answer]

What will result from the binding of a transcription factor to an enhancer region?

a. decreased transcription of an adjacent gene
b. increased transcription of a distant gene
c. alteration of the translation of an adjacent gene
d. initiation of the recruitment of RNA polymerase

[reveal-answer q="829037"]Show Answer[/reveal-answer]
[hidden-answer a="829037"]Answer b. Increased transcription of a distant gene will result from the binding of a transcription factor to an enhancer region.

[/hidden-answer]

A mutation within the promoter region can alter transcription of a gene. Describe how this can happen.

[practice-area rows="2"][/practice-area]
[reveal-answer q="332179"]Show Answer[/reveal-answer]
[hidden-answer a="332179"]A mutation in the promoter region can change the binding site for a transcription factor that normally binds to increase transcription. The mutation could either decrease the ability of the transcription factor to bind, thereby decreasing transcription, or it can increase the ability of the transcription factor to bind, thus increasing transcription.

[/hidden-answer]

What could happen if a cell had too much of an activating transcription factor present?

[practice-area rows="2"][/practice-area]
[reveal-answer q="162780"]Show Answer[/reveal-answer]
[hidden-answer a="162780"]If too much of an activating transcription factor were present, then transcription would be increased in the cell. This could lead to dramatic alterations in cell function. [/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

13.12: Eukaryotic Transcription Gene Regulation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.13: Post-Transcriptional Control of Gene Expression

RNA is transcribed, but must be processed into a mature form before translation can begin. This processing after an RNA molecule has been transcribed, but before it is translated into a protein, is called post-transcriptional modification. As with the epigenetic and transcriptional stages of processing, this post-transcriptional step can also be regulated to control gene expression in the cell. If the RNA is not processed, shuttled, or translated, then no protein will be synthesized.

## RNA splicing, the first stage of post-transcriptional regulation

In eukaryotic cells, the RNA transcript often contains regions, called introns, that are removed prior to translation. The regions of RNA that code for protein are called exons (Figure 1). After an RNA molecule has been transcribed, but prior to its departure from the nucleus to be translated, the RNA is processed and the introns are removed by splicing.





Figure 1. Pre-mRNA can be alternatively spliced to create different proteins.

Try It






Figure 2. There are five basic modes of alternative splicing.

In the 1970s, genes were first observed that exhibited alternative RNA splicing. Alternative RNA splicing is a mechanism that allows different protein products to be produced from one gene when different combinations of introns, and sometimes exons, are removed from the transcript (Figure 2). This alternative splicing can be haphazard, but more often it is controlled and acts as a mechanism of gene regulation, with the frequency of different splicing alternatives controlled by the cell as a way to control the production of different protein products in different cells or at different stages of development. Alternative splicing is now understood to be a common mechanism of gene regulation in eukaryotes; according to one estimate, 70 percent of genes in humans are expressed as multiple proteins through alternative splicing.

How could alternative splicing evolve? Introns have a beginning and ending recognition sequence; it is easy to imagine the failure of the splicing mechanism to identify the end of an intron and instead find the end of the next intron, thus removing two introns and the intervening exon. In fact, there are mechanisms in place to prevent such intron skipping, but mutations are likely to lead to their failure. Such “mistakes” would more than likely produce a nonfunctional protein. Indeed, the cause of many genetic diseases is alternative splicing rather than mutations in a sequence. However, alternative splicing would create a protein variant without the loss of the original protein, opening up possibilities for adaptation of the new variant to new functions. Gene duplication has played

an important role in the evolution of new functions in a similar way by providing genes that may evolve without eliminating the original, functional protein.

Visualize how mRNA splicing happens by watching the process in action in this video:



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=468

## Control of RNA Stability

Before the mRNA leaves the nucleus, it is given two protective "caps" that prevent the end of the strand from degrading during its journey. The **5′ cap**, which is placed on the 5′ end of the mRNA, is usually composed of a methylated guanosine triphosphate molecule (GTP). The **poly-A tail**, which is attached to the 3′ end, is usually composed of a series of adenine nucleotides. Once the RNA is transported to the cytoplasm, the length of time that the RNA resides there can be controlled. Each RNA molecule has a defined lifespan and decays at a specific rate. This rate of decay can influence how much protein is in the cell. If the decay rate is increased, the RNA will not exist in the cytoplasm as long, shortening the time for translation to occur. Conversely, if the rate of decay is decreased, the RNA molecule will reside in the cytoplasm longer and more protein can be translated. This rate of decay is referred to as the RNA stability. If the RNA is stable, it will be detected for longer periods of time in the cytoplasm.

Binding of proteins to the RNA can influence its stability. Proteins, called **RNA-binding proteins**, or RBPs, can bind to the regions of the RNA just upstream or downstream of the protein-coding region. These regions in the RNA that are not translated into protein are called the **untranslated regions**, or UTRs. They are not introns (those have been removed in the nucleus). Rather, these are regions that regulate mRNA localization, stability, and protein translation. The region just before the protein-coding region is called the **5′ UTR**, whereas the region after the coding region is called the **3′ UTR** (Figure 3). The binding of RBPs to these regions can increase or decrease the stability of an RNA molecule, depending on the specific RBP that binds.





Figure 3. The protein-coding region of mRNA is flanked by 5′ and 3′ untranslated regions (UTRs). The presence of RNA-binding proteins at the 5′ or 3′ UTR influences the stability of the RNA molecule.

## RNA Stability and microRNAs

In addition to RBPs that bind to and control (increase or decrease) RNA stability, other elements called microRNAs can bind to the RNA molecule. These **microRNAs**, or miRNAs, are short RNA molecules that are only 21–24 nucleotides in length. The miRNAs are made in the nucleus as longer pre-miRNAs. These pre-miRNAs are chopped into mature miRNAs by a protein called **dicer**. Like transcription factors and RBPs, mature miRNAs recognize a specific sequence and bind to the RNA; however, miRNAs also associate with a ribonucleoprotein complex called the **RNA-induced silencing complex (RISC)**. RISC binds along with the miRNA to degrade the target mRNA. Together, miRNAs and the RISC complex rapidly destroy the RNA molecule.

## Learning Objectives

Post-transcriptional control can occur at any stage after transcription, including RNA splicing, nuclear shuttling, and RNA stability. Once RNA is transcribed, it must be processed to create a mature RNA that is ready to be translated. This involves the removal of introns that do not code for protein. Spliceosomes bind to the signals that mark the exon/intron border to remove the introns and ligate the exons together. Once this occurs, the RNA is mature and can be translated. RNA is created and spliced in the nucleus, but needs to be transported to the cytoplasm to be translated. RNA is transported to the cytoplasm through the nuclear pore complex. Once the RNA is in the cytoplasm, the length of time it resides there before being degraded, called RNA stability, can also be altered to control the overall amount of protein that is synthesized. The RNA stability can be increased, leading to longer residency time in the cytoplasm, or decreased, leading to shortened time and less protein synthesis. RNA stability is controlled by RNA-binding proteins (RPBs) and microRNAs (miRNAs). These RPBs and miRNAs bind to the 5′ UTR or the 3′ UTR of the RNA to increase or decrease RNA stability. Depending on the RBP, the stability can be increased or decreased significantly; however, miRNAs always decrease stability and promote decay.

## Practice Questions

Which of the following are involved in post-transcriptional control?

a. control of RNA splicing
b. control of RNA shuttling
c. control of RNA stability
d. all of the above

[reveal-answer q=”681081″]Show Answer[/reveal-answer]
[hidden-answer a=”681081″]Answer d. All of the above (control of RNA splicing, RNA shuttling, and RNA stability) are involved in post-transcriptional control.

[/hidden-answer]

Binding of an RNA binding protein will _____ the stability of the RNA molecule.

a. increase
b. decrease
c. neither increase nor decrease
d. either increase or decrease

[reveal-answer q=”464261″]Show Answer[/reveal-answer]
[hidden-answer a=”464261″]Answer d. Binding of an RNA binding protein will either increase or decrease the stability of the RNA molecule.

[/hidden-answer]

Describe how RBPs can prevent miRNAs from degrading an RNA molecule.

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”16680″]Show Answer[/reveal-answer]
[hidden-answer a=”16680″]RNA binding proteins (RBP) bind to the RNA and can either increase or decrease the stability of the RNA. If they increase the stability of the RNA molecule, the RNA will remain intact in the cell for a longer period of time than normal. Since both RBPs and miRNAs bind to the RNA molecule, RBP can potentially bind first to the RNA and prevent the binding of the miRNA that will degrade it.

[/hidden-answer]

How can external stimuli alter post-transcriptional control of gene expression?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”598728″]Show Answer[/reveal-answer]

[hidden-answer a=”598728″]External stimuli can modify RNA-binding proteins (i.e., through phosphorylation of proteins) to alter their activity.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.13: Post-Transcriptional Control of Gene Expression is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.14: Putting It Together- Gene Expression

Let's return to our earlier look at cancer: One example of a gene modification that alters the growth rate is increased phosphorylation of cyclin B, a protein that controls the progression of a cell through the cell cycle and serves as a cell-cycle checkpoint protein.

For cells to move through each phase of the cell cycle, the cell must pass through checkpoints. This ensures that the cell has properly completed the step and has not encountered any mutation that will alter its function. Many proteins, including cyclin B, control these checkpoints. The phosphorylation of cyclin B, a post-translational event, alters its function. As a result, cells can progress through the cell cycle unimpeded, even if mutations exist in the cell and its growth should be terminated.

## Cancer Treatment

Cancer therapies are designed to target actively dividing cells. Besides cancer cells, what other cells actively divide in humans? One of the most physically obvious side effects of cancer treatments is hair loss. This is because the living cells in the hair root continually divide to make hair grow longer. These cells therefore are often impacted by broad scale cancer treatments like chemotherapy drugs and radiation localized to the head. Shortly after cancer treatments, a patient's blood count may also drop. This is due to the rapid division of blood cells throughout a person's life. Unlike hair cells though, blood cells divide fairly often and rapidly. Hence, the blood count often returns to normal much faster than hair regrows.

**Try It**

Scientists are using what is known about the regulation of gene expression in disease states, including cancer, to develop new ways to treat and prevent disease development. Many scientists are designing drugs on the basis of the gene expression patterns within individual tumors. This idea, that therapy and medicines can be tailored to an individual, has given rise to the field of personalized medicine. With an increased understanding of gene regulation and gene function, medicines can be designed to specifically target diseased cells without harming healthy cells. Some new medicines, called targeted therapies, have exploited the overexpression of a specific protein or the mutation of a gene to develop a new medication to treat disease. One such example is the use of anti-EGF receptor medications to treat the subset of breast cancer tumors that have very high levels of the EGF protein. Undoubtedly, more targeted therapies will be developed as scientists learn more about how gene expression changes can cause cancer.

As scientists learn more about cell division and the unique ways it malfunctions in cancer cells, they are able to develop targeted therapies. These drugs are still chemotherapies, but they are often focused on a particular feature of different types of cancer cells, making them less likely to target non-cancerous dividing cells. This reduces global side effects. Unfortunately, our understanding of cancer is still incomplete. Therefore, every day cancer researchers and clinicians work to manage and treat these horrible diseases.

Read more at www.cancer.org

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: Gene Expression. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

13.14: Putting It Together- Gene Expression is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 13.15: Assignment- How Mutations Work

For this hands on activity you should download the "Mutations: Changing the Genetic Code" simulation from MOLO.

Complete all 7 pages of the simulation, answering all the questions and taking the snapshots as directed. For each snapshot you should explain what your image is showing. When you have completed the simulation, click on the button at the bottom of the screen that says **Create a report for printing and submission**. Choose to **Print without login** and enter your name in the appropriate field. Instead of school, please enter the term. When the next screen pops up, hit the print button. Instead of printing though, you can choose to save the file as a PDF. Then return here to submit your PDF document.

Note there are no optional questions for you; all questions in the simulation are required even if marked "optional."

This document shows you an example submission; please note that it only includes answers to a few questions and a single snapshot to give you an idea of how to complete this assignment.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment includes 6 snapshots.
- Assignment includes answer to all 13 questions.

## Rubric

| How Mutations Work | | | | |
|---|---|---|---|---|
| Outcome: Describe the conversion of DNA to RNA to proteins. Explain the regulation of gene expression. | | | | |
| Criteria | Ratings | | | Pts |
| Outline the process of genetic transcription. | Answered all 3 questions correctly and included snapshots.<br>5.0 pts | Answered at least 2 of 3 questions correctly and included snapshots.<br>4.0 pts | Did not answer questions 1–3 correctly and/or did not include all snapshots.<br>0.0 pts | 5 pts |
| Summarize the process of genetic translation. | Answered 9–10 questions correctly and included snapshots.<br>5.0 pts | Answered 8 questions correctly and included snapshots.<br>4.0 pts | Answered less than 8 questions correctly and/or did not include all snapshots.<br>0.0 pts | 5 pts |
| | | | | Total points: 10 |

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Mutations: Changing the Genetic Code (a 7-page activity). **Authored by**: Concord Consortium. **Located at**: http://molo.concord.org/database/activities/102.html. **License**: *CC BY-NC: Attribution-NonCommercial*

13.15: Assignment- How Mutations Work is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 14: Module 12- Trait Inheritance


14.1: Why It Matters- Trait Inheritance
14.2: Introduction to the Father of Genetics
14.3: Mendel’s Experiments and Heredity
14.4: Characteristics and Traits
14.5: Laws of Inheritance
14.6: Heredity
14.7: Introduction to Beyond Dominance and Recessiveness
14.8: Non-Mendelian Inheritance
14.9: Non-Mendelian Punnett Squares
14.10: Multiple Alleles
14.11: Introduction to Heredity and Disease
14.12: Penetrance and Expressivity
14.13: Pedigrees and Disease
14.14: Pedigree Flowchart
14.15: Genetic Disorder and Pedigrees
14.16: Introduction to Genetics and the Environment
14.17: Epistasis
14.18: Polygenic Inheritance and Environmental Effects
14.19: Effect of the Environment
14.20: Pleiotropy and Human Disorders
14.21: Putting It Together- Trait Inheritance
14.22: Assignment- Dragon Genetics



14: Module 12- Trait Inheritance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.1: Why It Matters- Trait Inheritance

## Why complete monohybrid and dihybrid crosses and family pedigrees, and explain the inheritance of various traits?

Take a moment to pause and think about yourself and your family. How much do you look like your parents? What about the other members of your family: aunts, uncles, cousins, full siblings, half-siblings? Why do some family members look nearly identical to each other, while other family members seem as if they don't share two traits? The answer to this boils down to *how* traits are inherited.

Parents pass on traits to their children. Traits like blood type, cleft chin, dimples, and widow's peaks are all inherited in a fairly straight-forward, simple fashion. However, the inheritance of other traits is much more complex and harder to understand: these traits include height, skin color, and eye color. These more complex paths of inheritance are what cause some siblings to look so different from each other—despite having the same two parents.

Unfortunately, genetics also plays a role in the inheritance of some diseases such as Huntingdon's, sickle cell anemia, and Tay-Sachs disease. A degree in genetics can be used in careers ranging from a forensic examiner, a genetic counselor, a medical geneticist, a statistical geneticist, to a clinical technician.

**Try It**

Genetic counselors use their understanding of the rules of heredity to help gauge the risks of genetic diseases. They may advise a couple of their risks for passing on certain genetic disorders, order genetic testing, and then explain the chances that some genetic diseases will show up in any offspring. Genetic counselors are there to help potential parents understand risk factors and the prognosis of genetic disorders, including basic treatment and management plans in the case of certain disorders.

### Learning Outcomes

- Identify the impact of Gregor Mendel on the field of genetics and apply Mendel's two laws of genetics
- Explain complications to the phenotypic expression of genotype, including mutations
- Explain the conventions of a family pedigree and predict whether a disease will be passed through a family in one of three modes
- Discuss the role environment plays on phenotypes

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Trait Inheritance. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

14.1: Why It Matters- Trait Inheritance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.2: Introduction to the Father of Genetics

## Identify the impact of Gregor Mendel on the field of genetics and apply Mendel's two laws of genetics

Gregor Mendel is often referred to as the Father of Genetics. But just what did he do to earn this honorary title? In this outcome we'll examine the work he did and how his work still impacts genetics today.

## What You'll Learn to Do

- Describe Mendel's study of garden peas and hereditary
- Understand how the inheritance of a genotype generates a phenotype
- Apply the law of segregation
- Apply the law of independent assortment

### Learning Activities

The learning activities for this section include the following:

- Mendel's Experiments and Heredity
- Characteristics and Traits
- Laws of Inheritance
- Heredity
- Self Check: The Father of Genetics

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

14.2: Introduction to the Father of Genetics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.3: Mendel's Experiments and Heredity



Figure 1. Experimenting with thousands of garden peas, Mendel uncovered the fundamentals of genetics. (credit: modification of work by Jerry Kirkhart)

Genetics is the study of heredity. Johann Gregor Mendel set the framework for genetics long before chromosomes or genes had been identified, at a time when meiosis was not well understood. Mendel selected a simple biological system and conducted methodical, quantitative analyses using large sample sizes. Because of Mendel's work, the fundamental principles of heredity were revealed. We now know that genes, carried on chromosomes, are the basic functional units of heredity with the capability to be replicated, expressed, or mutated. Today, the postulates put forth by Mendel form the basis of classical, or Mendelian, genetics. Not all genes are transmitted from parents to offspring according to Mendelian genetics, but Mendel's experiments serve as an excellent starting point for thinking about inheritance.

## Mendel's Experiments and the Laws of Probability



Figure 2. Johann Gregor Mendel is considered the father of genetics.

Johann Gregor Mendel (1822–1884) (Figure 2) was a lifelong learner, teacher, scientist, and man of faith. As a young adult, he joined the Augustinian Abbey of St. Thomas in Brno in what is now the Czech Republic. Supported by the monastery, he taught physics, botany, and natural science courses at the secondary and university levels. In 1856, he began a decade-long research pursuit involving inheritance patterns in honeybees and plants, ultimately settling on pea plants as his primary **model system** (a system with convenient characteristics used to study a specific biological phenomenon to be applied to other systems). In 1865, Mendel presented the results of his experiments with nearly 30,000 pea plants to the local Natural History Society. He demonstrated that traits are transmitted faithfully from parents to offspring independently of other traits and in dominant and recessive patterns. In 1866, he published his work, *Experiments in Plant Hybridization*, in the proceedings of the Natural History Society of Brünn.

Mendel's work went virtually unnoticed by the scientific community that believed, incorrectly, that the process of inheritance involved a blending of parental traits that produced an intermediate physical appearance in offspring; this hypothetical process appeared to be correct because of what we know now as continuous variation. **Continuous variation** results from the action of many genes to determine a characteristic like human height. Offspring appear to be a "blend" of their parents' traits when we look at characteristics that exhibit continuous variation. The **blending theory of inheritance** asserted that the original parental traits were lost or absorbed by the blending in the offspring, but we now know that this is not the case. Mendel was the first researcher to see it. Instead of continuous characteristics, Mendel worked with traits that were inherited in distinct classes (specifically, violet

versus white flowers); this is referred to as **discontinuous variation**. Mendel's choice of these kinds of traits allowed him to see experimentally that the traits were not blended in the offspring, nor were they absorbed, but rather that they kept their distinctness and could be passed on. In 1868, Mendel became abbot of the monastery and exchanged his scientific pursuits for his pastoral duties. He was not recognized for his extraordinary scientific contributions during his lifetime. In fact, it was not until 1900 that his work was rediscovered, reproduced, and revitalized by scientists on the brink of discovering the chromosomal basis of heredity.

## Mendel's Model System

Mendel's seminal work was accomplished using the garden pea, *Pisum sativum*, to study inheritance. This species naturally self-fertilizes, such that pollen encounters ova within individual flowers. The flower petals remain sealed tightly until after pollination, preventing pollination from other plants. The result is highly inbred, or "true-breeding," pea plants. These are plants that always produce offspring that look like the parent. By experimenting with true-breeding pea plants, Mendel avoided the appearance of unexpected traits in offspring that might occur if the plants were not true breeding. The garden pea also grows to maturity within one season, meaning that several generations could be evaluated over a relatively short time. Finally, large quantities of garden peas could be cultivated simultaneously, allowing Mendel to conclude that his results did not come about simply by chance.

## Mendelian Crosses

Mendel performed **hybridizations**, which involve mating two true-breeding individuals that have different traits. In the pea, which is naturally self-pollinating, this is done by manually transferring pollen from the anther of a mature pea plant of one variety to the stigma of a separate mature pea plant of the second variety. In plants, pollen carries the male gametes (sperm) to the stigma, a sticky organ that traps pollen and allows the sperm to move down the pistil to the female gametes (ova) below. To prevent the pea plant that was receiving pollen from self-fertilizing and confounding his results, Mendel painstakingly removed all of the anthers from the plant's flowers before they had a chance to mature.

Plants used in first-generation crosses were called $P_0$, or parental generation one, plants (Figure 3). Mendel collected the seeds belonging to the $P_0$ plants that resulted from each cross and grew them the following season. These offspring were called the $\mathbf{F_1}$, or the first filial (*filial* = offspring, daughter or son), generation. Once Mendel examined the characteristics in the $F_1$ generation of plants, he allowed them to self-fertilize naturally. He then collected and grew the seeds from the $F_1$ plants to produce the $\mathbf{F_2}$, or second filial, generation. Mendel's experiments extended beyond the $F_2$ generation to the $F_3$ and $F_4$generations, and so on, but it was the ratio of characteristics in the $P_0$−$F_1$−$F_2$ generations that were the most intriguing and became the basis for Mendel's postulates.





Figure 3. In one of his experiments on inheritance patterns, Mendel crossed plants that were true-breeding for violet flower color with plants true-breeding for white flower color (the $P_0$ generation). The resulting hybrids in the $F_1$ generation all had violet flowers. In the $F_2$ generation, approximately three quarters of the plants had violet flowers, and one quarter had white flowers.

## Garden Pea Characteristics Revealed the Basics of Heredity

In his 1865 publication, Mendel reported the results of his crosses involving seven different characteristics, each with two contrasting traits. A **trait** is defined as a variation in the physical appearance of a heritable characteristic. The characteristics included plant height, seed texture, seed color, flower color, pea pod size, pea pod color, and flower position. For the characteristic of flower color, for example, the two contrasting traits were white versus violet. To fully examine each characteristic, Mendel generated large numbers of $F_1$ and $F_2$ plants, reporting results from 19,959 $F_2$ plants alone. His findings were consistent.

What results did Mendel find in his crosses for flower color? First, Mendel confirmed that he had plants that bred true for white or violet flower color. Regardless of how many generations Mendel examined, all self-crossed offspring of parents with white flowers had white flowers, and all self-crossed offspring of parents with violet flowers had violet flowers. In addition, Mendel confirmed that, other than flower color, the pea plants were physically identical.

Once these validations were complete, Mendel applied the pollen from a plant with violet flowers to the stigma of a plant with white flowers. After gathering and sowing the seeds that resulted from this cross, Mendel found that 100 percent of the $F_1$ hybrid generation had violet flowers. Conventional wisdom at that time would have predicted the hybrid flowers to be pale violet or for hybrid plants to have equal numbers of white and violet flowers. In other words, the contrasting parental traits were expected to blend in the offspring. Instead, Mendel's results demonstrated that the white flower trait in the $F_1$ generation had completely disappeared.

Importantly, Mendel did not stop his experimentation there. He allowed the $F_1$ plants to self-fertilize and found that, of $F_2$-generation plants, 705 had violet flowers and 224 had white flowers. This was a ratio of 3.15 violet flowers per one white flower, or approximately 3:1. When Mendel transferred pollen from a plant with violet flowers to the stigma of a plant with white flowers and vice versa, he obtained about the same ratio regardless of which parent, male or female, contributed which trait. This is called a **reciprocal cross**—a paired cross in which the respective traits of the male and female in one cross become the respective traits of the female and male in the other cross. For the other six characteristics Mendel examined, the $F_1$ and $F_2$ generations behaved in the same way as they had for flower color. One of the two traits would disappear completely from the $F_1$ generation only to reappear in the $F_2$ generation at a ratio of approximately 3:1 (Table 1).

Table 1. The Results of Mendel's Garden Pea Hybridizations

| Characteristic | Contrasting $P_0$ Traits | $F_1$ Offspring Traits | $F_2$ Offspring Traits | $F_2$ Trait Ratios |
|---|---|---|---|---|
| Flower color | Violet vs. white | 100 percent violet | • 705 violet<br>• 224 white | 3.15:1 |
| Flower position | Axial vs. terminal | 100 percent axial | • 651 axial<br>• 207 terminal | 3.14:1 |
| Plant height | Tall vs. dwarf | 100 percent tall | • 787 tall<br>• 277 dwarf | 2.84:1 |
| Seed texture | Round vs. wrinkled | 100 percent round | • 5,474 round<br>• 1,850 wrinkled | 2.96:1 |
| Seed color | Yellow vs. green | 100 percent yellow | • 6,022 yellow<br>• 2,001 green | 3.01:1 |
| Pea pod texture | Inflated vs. constricted | 100 percent inflated | • 882 inflated<br>• 299 constricted | 2.95:1 |
| Pea pod color | Green vs. yellow | 100 percent green | • 428 green<br>• 152 yellow | 2.82:1 |

Upon compiling his results for many thousands of plants, Mendel concluded that the characteristics could be divided into expressed and latent traits. He called these, respectively, dominant and recessive traits. **Dominant traits** are those that are inherited unchanged in a hybridization. **Recessive traits** become latent, or disappear, in the offspring of a hybridization. The recessive trait does, however, reappear in the progeny of the hybrid offspring. An example of a dominant trait is the violet-flower trait. For this same characteristic (flower color), white-colored flowers are a recessive trait. The fact that the recessive trait reappeared in the $F_2$ generation meant that the traits remained separate (not blended) in the plants of the $F_1$ generation. Mendel also proposed that plants possessed two copies of the trait for the flower-color characteristic, and that each parent transmitted one of its two copies to its offspring, where they came together. Moreover, the physical observation of a dominant trait could mean that the genetic composition of the organism included two dominant versions of the characteristic or that it included one dominant and one recessive version. Conversely, the observation of a recessive trait meant that the organism lacked any dominant versions of this characteristic.

So why did Mendel repeatedly obtain 3:1 ratios in his crosses? To understand how Mendel deduced the basic mechanisms of inheritance that lead to such ratios, we must first review the laws of probability.

## Probability Basics

Probabilities are mathematical measures of likelihood. The empirical probability of an event is calculated by dividing the number of times the event occurs by the total number of opportunities for the event to occur. It is also possible to calculate theoretical probabilities by dividing the number of times that an event is expected to occur by the number of times that it could occur. Empirical probabilities come from observations, like those of Mendel. Theoretical probabilities come from knowing how the events are produced and assuming that the probabilities of individual outcomes are equal. A probability of one for some event indicates that it is guaranteed to occur, whereas a probability of zero indicates that it is guaranteed not to occur. An example of a genetic event is a round seed produced by a pea plant. In his experiment, Mendel demonstrated that the probability of the event "round seed" occurring was one in the $F_1$ offspring of true-breeding parents, one of which has round seeds and one of which has wrinkled seeds. When the $F_1$ plants were subsequently self-crossed, the probability of any given $F_2$ offspring having round seeds was now three out of four. In other words, in a large population of $F_2$ offspring chosen at random, 75 percent were expected to have round seeds, whereas 25 percent were expected to have wrinkled seeds. Using large numbers of crosses, Mendel was able to calculate probabilities and use these to predict the outcomes of other crosses.

### The Product Rule and Sum Rule

Mendel demonstrated that the pea-plant characteristics he studied were transmitted as discrete units from parent to offspring. As will be discussed, Mendel also determined that different characteristics, like seed color and seed texture, were transmitted independently of one another and could be considered in separate probability analyses. For instance, performing a cross between a plant with green, wrinkled seeds and a plant with yellow, round seeds still produced offspring that had a 3:1 ratio of green:yellow seeds (ignoring seed texture) and a 3:1 ratio of round:wrinkled seeds (ignoring seed color). The characteristics of color and texture did not influence each other.

The **product** rule of probability can be applied to this phenomenon of the independent transmission of characteristics. The product rule states that the probability of two independent events occurring together can be calculated by multiplying the individual probabilities of each event occurring alone. To demonstrate the product rule, imagine that you are rolling a six-sided die (D) and flipping a penny (P) at the same time. The die may roll any number from 1–6 (D#), whereas the penny may turn up heads (PH) or tails (PT). The outcome of rolling the die has no effect on the outcome of flipping the penny and vice versa. There are 12 possible outcomes of this action (Table 2), and each event is expected to occur with equal probability.

Table 2. Twelve Equally Likely Outcomes of Rolling a Die and Flipping a Penny

| Rolling Die | Flipping Penny |
|---|---|
| D1 | PH |
| D1 | PT |
| D2 | PH |
| D2 | PT |
| D3 | PH |
| D3 | PT |
| D4 | PH |
| D4 | PT |
| D5 | PH |
| D5 | PT |
| D6 | PH |
| D6 | PT |

Of the 12 possible outcomes, the die has a 2/12 (or 1/6) probability of rolling a two, and the penny has a 6/12 (or 1/2) probability of coming up heads. By the product rule, the probability that you will obtain the combined outcome 2 and heads is: (D2) x (PH) = (1/6) x (1/2) or 1/12 (Table 3). Notice the word "and" in the description of the probability. The "and" is a signal to apply the product rule. For example, consider how the product rule is applied to the dihybrid cross: the probability of having both dominant traits in the F2progeny is the product of the probabilities of having the dominant trait for each characteristic, as shown here:

On the other hand, the **sum rule** of probability is applied when considering two mutually exclusive outcomes that can come about by more than one pathway. The sum rule states that the probability of the occurrence of one event or the other event, of two mutually exclusive events, is the sum of their individual probabilities. Notice the word "or" in the description of the probability. The "or" indicates that you should apply the sum rule. In this case, let's imagine you are flipping a penny (P) and a quarter (Q). What is the probability of one coin coming up heads and one coin coming up tails? This outcome can be achieved by two cases: the penny may be heads (PH) and the quarter may be tails (QT), or the quarter may be heads (QH) and the penny may be tails (PT). Either case fulfills the outcome. By the sum rule, we calculate the probability of obtaining one head and one tail as [(PH) × (QT)] + [(QH) × (PT)] = [(1/2) × (1/2)] + [(1/2) × (1/2)] = 1/2 (Table 3). You should also notice that we used the product rule to calculate the probability of PH and QT, and also the probability of PT and QH, before we summed them. Again, the sum rule can be applied to show the probability of having just one dominant trait in the F2 generation of a dihybrid cross:

Table 3. The Product Rule and Sum Rule

| Product Rule | Sum Rule |
|---|---|
| For independent events A and B, the probability (P) of them both occurring (A *and* B) is (PA × PB) | For mutually exclusive events A and B, the probability (P) that at least one occurs (A *or* B) is (PA + PB) |

To use probability laws in practice, it is necessary to work with large sample sizes because small sample sizes are prone to deviations caused by chance. The large quantities of pea plants that Mendel examined allowed him calculate the probabilities of the traits appearing in his $F_2$ generation. As you will learn, this discovery meant that when parental traits were known, the offspring's traits could be predicted accurately even before fertilization.

## Learning Objectives

Working with garden pea plants, Mendel found that crosses between parents that differed by one trait produced $F_1$ offspring that all expressed the traits of one parent. Observable traits are referred to as dominant, and non-expressed traits are described as recessive. When the offspring in Mendel's experiment were self-crossed, the $F_2$ offspring exhibited the dominant trait or the recessive trait in a 3:1 ratio, confirming that the recessive trait had been transmitted faithfully from the original $P_0$ parent. Reciprocal crosses generated identical $F_1$ and $F_2$ offspring ratios. By examining large sample sizes, Mendel showed that his crosses behaved reproducibly according to the laws of probability, and that the traits were inherited as independent events.

Two rules in probability can be used to find the expected proportions of offspring of different traits from different crosses. To find the probability of two or more independent events occurring together, apply the product rule and multiply the probabilities of the individual events. The use of the word "and" suggests the appropriate application of the product rule. To find the probability of two or more events occurring in combination, apply the sum rule and add their individual probabilities together. The use of the word "or" suggests the appropriate application of the sum rule.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

14.3: Mendel's Experiments and Heredity is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.4: Characteristics and Traits

The seven characteristics that Mendel evaluated in his pea plants were each expressed as one of two versions, or traits. The physical expression of characteristics is accomplished through the expression of genes carried on chromosomes. The genetic makeup of peas consists of two similar or homologous copies of each chromosome, one from each parent. Each pair of homologous chromosomes has the same linear order of genes. In other words, peas are diploid organisms in that they have two copies of each chromosome. The same is true for many other plants and for virtually all animals. Diploid organisms utilize meiosis to produce haploid gametes, which contain one copy of each homologous chromosome that unite at fertilization to create a diploid zygote.

For cases in which a single gene controls a single characteristic, a diploid organism has two genetic copies that may or may not encode the same version of that characteristic. Gene variants that arise by mutation and exist at the same relative locations on homologous chromosomes are called alleles. Mendel examined the inheritance of genes with just two allele forms, but it is common to encounter more than two alleles for any given gene in a natural population.

## Phenotypes and Genotypes

Two alleles for a given gene in a diploid organism are expressed and interact to produce physical characteristics. The observable traits expressed by an organism are referred to as its **phenotype**. An organism's underlying genetic makeup, consisting of both physically visible and non-expressed alleles, is called its **genotype**. Mendel's hybridization experiments demonstrate the difference between phenotype and genotype. When true-breeding plants in which one parent had yellow pods and one had green pods were cross-fertilized, all of the $F_1$ hybrid offspring had yellow pods. That is, the hybrid offspring were phenotypically identical to the true-breeding parent with yellow pods. However, we know that the allele donated by the parent with green pods was not simply lost because it reappeared in some of the $F_2$ offspring. Therefore, the $F_1$ plants must have been genotypically different from the parent with yellow pods.

The $P_0$ plants that Mendel used in his experiments were each homozygous for the trait he was studying. Diploid organisms that are **homozygous** at a given gene, or locus, have two identical alleles for that gene on their homologous chromosomes. Mendel's parental pea plants always bred true because both of the gametes produced carried the same trait. When $P_0$ plants with contrasting traits were cross-fertilized, all of the offspring were heterozygous for the contrasting trait, meaning that their genotype reflected that they had different alleles for the gene being examined.

### Dominant and Recessive Alleles

Our discussion of homozygous and heterozygous organisms brings us to why the $F_1$ heterozygous offspring were identical to one of the parents, rather than expressing both alleles. In all seven pea-plant characteristics, one of the two contrasting alleles was dominant, and the other was recessive. Mendel called the dominant allele the expressed unit factor; the recessive allele was referred to as the latent unit factor. We now know that these so-called unit factors are actually genes on homologous chromosome pairs. For a gene that is expressed in a dominant and recessive pattern, homozygous dominant and heterozygous organisms will look identical (that is, they will have different genotypes but the same phenotype). The recessive allele will only be observed in homozygous recessive individuals (Table 1).

| Table 1. Human Inheritance in Dominant and Recessive Patterns | |
|---|---|
| Dominant Traits | Recessive Traits |
| Achondroplasia | Albinism |
| Brachydactyly | Cystic fibrosis |
| Huntington's disease | Duchenne muscular dystrophy |
| Marfan syndrome | Galactosemia |
| Neurofibromatosis | Phenylketonuria |
| Widow's peak | Sickle-cell anemia |
| Wooly hair | Tay-Sachs disease |

Several conventions exist for referring to genes and alleles. For the purposes of this chapter, we will abbreviate genes using the first letter of the gene's corresponding dominant trait. For example, violet is the dominant trait for a pea plant's flower color, so the flower-color gene would be abbreviated as *V* (note that it is customary to italicize gene designations). Furthermore, we will use uppercase and lowercase letters to represent dominant and recessive alleles, respectively. Therefore, we would refer to the genotype of a homozygous dominant pea plant with violet flowers as *VV*, a homozygous recessive pea plant with white flowers as *vv*, and a heterozygous pea plant with violet flowers as *Vv*.

## Punnett Square Approach for a Monohybrid Cross

When fertilization occurs between two true-breeding parents that differ in only one characteristic, the process is called a **monohybrid** cross, and the resulting offspring are monohybrids. Mendel performed seven monohybrid crosses involving contrasting traits for each characteristic. On the basis of his results in $F_1$ and $F_2$ generations, Mendel postulated that each parent in the monohybrid cross contributed one of two paired unit factors to each offspring, and every possible combination of unit factors was equally likely.

To demonstrate a monohybrid cross, consider the case of true-breeding pea plants with yellow versus green pea seeds. The dominant seed color is yellow; therefore, the parental genotypes were *YY* for the plants with yellow seeds and *yy* for the plants with green seeds, respectively. A **Punnett square**, devised by the British geneticist Reginald Punnett, can be drawn that applies the rules of probability to predict the possible outcomes of a genetic cross or mating and their expected frequencies. To prepare a Punnett square, all possible combinations of the parental alleles are listed along the top (for one parent) and side (for the other parent) of a grid, representing their meiotic segregation into haploid gametes. Then the combinations of egg and sperm are made in the boxes in the table to show which alleles are combining. Each box then represents the diploid genotype of a zygote, or fertilized egg, that could result from this mating. Because each possibility is equally likely, genotypic ratios can be determined from a Punnett square. If the pattern of inheritance (dominant or recessive) is known, the phenotypic ratios can be inferred as well. For a monohybrid cross of two true-breeding parents, each parent contributes one type of allele. In this case, only one genotype is possible. All offspring are *Yy* and have yellow seeds (Figure 1).





Figure 1. In the $P_0$ generation, pea plants that are true-breeding for the dominant yellow phenotype are crossed with plants with the recessive green phenotype. This cross produces $F_1$ heterozygotes with a yellow phenotype. Punnett square analysis can be used to predict the genotypes of the $F_2$ generation.

A self-cross of one of the *Yy* heterozygous offspring can be represented in a 2 × 2 Punnett square because each parent can donate one of two different alleles. Therefore, the offspring can potentially have one of four allele combinations: *YY*, *Yy*, *yY*, or *yy* (Figure 1). Notice that there are two ways to obtain the *Yy* genotype: a *Y* from the egg and a *y* from the sperm, or a *y* from the egg and a *Y* from the sperm. Both of these possibilities must be counted. Recall that Mendel's pea-plant characteristics behaved in the same way in reciprocal crosses. Therefore, the two possible heterozygous combinations produce offspring that are genotypically and phenotypically identical despite their dominant and recessive alleles deriving from different parents. They are grouped together. Because fertilization is a random event, we expect each combination to be equally likely and for the offspring to exhibit a ratio of *YY*:*Yy*:*yy* genotypes of 1:2:1 (Figure 1). Furthermore, because the *YY* and *Yy* offspring have yellow seeds and are phenotypically identical, applying the sum rule of probability, we expect the offspring to exhibit a phenotypic ratio of 3 yellow:1 green. Indeed, working with large sample sizes, Mendel observed approximately this ratio in every $F_2$ generation resulting from crosses for individual traits.

Mendel validated these results by performing an $F_3$ cross in which he self-crossed the dominant- and recessive-expressing $F_2$ plants. When he self-crossed the plants expressing green seeds, all of the offspring had green seeds, confirming that all green seeds had homozygous genotypes of *yy*. When he self-crossed the $F_2$ plants expressing yellow seeds, he found that one-third of the plants bred true, and two-thirds of the plants segregated at a 3:1 ratio of yellow:green seeds. In this case, the true-breeding plants had homozygous (*YY*) genotypes, whereas the segregating plants corresponded to the heterozygous (*Yy*) genotype. When these plants self-fertilized, the outcome was just like the $F_1$ self-fertilizing cross.

## Test Cross Distinguishes the Dominant Phenotype

Beyond predicting the offspring of a cross between known homozygous or heterozygous parents, Mendel also developed a way to determine whether an organism that expressed a dominant trait was a heterozygote or a homozygote. Called the test cross, this technique is still used by plant and animal breeders. In a test cross, the dominant-expressing organism is crossed with an organism that is homozygous recessive for the same characteristic. If the dominant-expressing organism is a homozygote, then all $F_1$ offspring will be heterozygotes expressing the dominant trait (Figure 2). Alternatively, if the dominant expressing organism is a

heterozygote, the $F_1$ offspring will exhibit a 1:1 ratio of heterozygotes and recessive homozygotes (Figure 2). The test cross further validates Mendel's postulate that pairs of unit factors segregate equally.

Practice Question





Figure 2. A test cross can be performed to determine whether an organism expressing a dominant trait is a homozygote or a heterozygote.

In pea plants, round peas (*R*) are dominant to wrinkled peas (*r*). You do a test cross between a pea plant with wrinkled peas (genotype *rr*) and a plant of unknown genotype that has round peas. You end up with three plants, all which have round peas. From this data, can you tell if the round pea parent plant is homozygous dominant or heterozygous? If the round pea parent plant is heterozygous, what is the probability that a random sample of 3 progeny peas will all be round?

[practice-area rows="2"][/practice-area]
[reveal-answer q="837873"]**Show Answer**[/reveal-answer]
[hidden-answer a="837873"]You cannot be sure if the plant is homozygous or heterozygous as the data set is too small: by random chance, all three plants might have acquired only the dominant gene even if the recessive one is present. If the round pea parent is heterozygous, there is a one-eighth probability that a random sample of three progeny peas will all be round.[/hidden-answer]

Many human diseases are genetically inherited. A healthy person in a family in which some members suffer from a recessive genetic disorder may want to know if he or she has the disease-causing gene and what risk exists of passing the disorder on to his or her offspring. Of course, doing a test cross in humans is unethical and impractical. Instead, geneticists use pedigree analysis to study the inheritance pattern of human genetic diseases (Figure 3).

Practice Question




Figure 3. Pedigree Analysis for Alkaptonuria

Alkaptonuria is a recessive genetic disorder in which two amino acids, phenylalanine and tyrosine, are not properly metabolized. Affected individuals may have darkened skin and brown urine, and may suffer joint damage and other complications. In this pedigree, individuals with the disorder are indicated in blue and have the genotype aa. Unaffected individuals are indicated in yellow and have the genotype AA or Aa. Note that it is often possible to determine a person's genotype from the genotype of their offspring. For example, if neither parent has the disorder but their child does, they must be heterozygous. Two individuals on the pedigree have an unaffected phenotype but unknown genotype. Because they do not have the disorder, they must have at least one normal allele, so their genotype gets the "A?" designation.

What are the genotypes of the individuals labeled 1, 2 and 3?

[practice-area rows="2"][/practice-area]
[reveal-answer q="567889"]**Show Answer**[/reveal-answer]
[hidden-answer a="567889"]Individual 1 has the genotype *aa*. Individual 2 has the genotype *Aa*. Individual 3 has the genotype *Aa*.[/hidden-answer]

## Learning Objectives

When true-breeding or homozygous individuals that differ for a certain trait are crossed, all of the offspring will be heterozygotes for that trait. If the traits are inherited as dominant and recessive, the $F_1$ offspring will all exhibit the same phenotype as the parent homozygous for the dominant trait. If these heterozygous offspring are self-crossed, the resulting $F_2$ offspring will be equally likely to inherit gametes carrying the dominant or recessive trait, giving rise to offspring of which one quarter are homozygous dominant, half are heterozygous, and one quarter are homozygous recessive. Because homozygous dominant and heterozygous individuals are phenotypically identical, the observed traits in the $F_2$ offspring will exhibit a ratio of three dominant to one recessive.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

14.4: Characteristics and Traits is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.5: Laws of Inheritance

Mendel generalized the results of his pea-plant experiments into four postulates, some of which are sometimes called “laws,” that describe the basis of dominant and recessive inheritance in diploid organisms. As you will learn, more complex extensions of Mendelism exist that do not exhibit the same $F_2$ phenotypic ratios (3:1). Nevertheless, these laws summarize the basics of classical genetics.

## Pairs of Unit Factors, or Genes

Mendel proposed first that paired unit factors of heredity were transmitted faithfully from generation to generation by the dissociation and reassociation of paired factors during gametogenesis and fertilization, respectively. After he crossed peas with contrasting traits and found that the recessive trait resurfaced in the $F_2$ generation, Mendel deduced that hereditary factors must be inherited as discrete units. This finding contradicted the belief at that time that parental traits were blended in the offspring.

## Alleles Can Be Dominant or Recessive



Figure 1. The child in the photo expresses albinism, a recessive trait.

Mendel’s **law of dominance** states that in a heterozygote, one trait will conceal the presence of another trait for the same characteristic. Rather than both alleles contributing to a phenotype, the dominant allele will be expressed exclusively. The recessive allele will remain “latent” but will be transmitted to offspring by the same manner in which the dominant allele is transmitted. The recessive trait will only be expressed by offspring that have two copies of this allele (Figure 1), and these offspring will breed true when self-crossed.

Since Mendel’s experiments with pea plants, other researchers have found that the law of dominance does not always hold true. Instead, several different patterns of inheritance have been found to exist, which we will explore this later in the module.

## Equal Segregation of Alleles

Observing that true-breeding pea plants with contrasting traits gave rise to $F_1$ generations that all expressed the dominant trait and $F_2$ generations that expressed the dominant and recessive traits in a 3:1 ratio, Mendel proposed the **law of segregation**. This law states that paired unit factors (genes) must segregate equally into gametes such that offspring have an equal likelihood of inheriting either factor. For the $F_2$ generation of a monohybrid cross, the following three possible combinations of genotypes could result: homozygous dominant, heterozygous, or homozygous recessive. Because heterozygotes could arise from two different pathways (receiving one dominant and one recessive allele from either parent), and because heterozygotes and homozygous dominant individuals are phenotypically identical, the law supports Mendel’s observed 3:1 phenotypic ratio. The equal segregation of alleles is the reason we can apply the Punnett square to accurately predict the offspring of parents with known genotypes. The physical basis of Mendel’s law of segregation is the first division of meiosis, in which the homologous chromosomes with their different versions of each gene are segregated into daughter nuclei. The role of the meiotic segregation of chromosomes in sexual reproduction was not understood by the scientific community during Mendel’s lifetime.

## Independent Assortment

Mendel’s **law of independent assortment** states that genes do not influence each other with regard to the sorting of alleles into gametes, and every possible combination of alleles for every gene is equally likely to occur. The independent assortment of genes

can be illustrated by the **dihybrid** cross, a cross between two true-breeding parents that express different traits for two characteristics. Consider the characteristics of seed color and seed texture for two pea plants, one that has green, wrinkled seeds (*yyrr*) and another that has yellow, round seeds (*YYRR*). Because each parent is homozygous, the law of segregation indicates that the gametes for the green/wrinkled plant all are *yr*, and the gametes for the yellow/round plant are all *YR*. Therefore, the $F_1$ generation of offspring all are*YyRr* (Figure 2).





Figure 2. This dihybrid cross of pea plants involves the genes for seed color and texture.

**Practice Question**

In pea plants, purple flowers (P) are dominant to white flowers (p) and yellow peas (Y) are dominant to green peas (y). What are the possible genotypes and phenotypes for a cross between PpYY and ppYy pea plants? How many squares do you need to do a Punnett square analysis of this cross?

[practice-area rows="2"][/practice-area]
[reveal-answer q="733053"]**Show Answer**[/reveal-answer]
[hidden-answer a="733053"]The possible genotypes are PpYY, PpYy, ppYY, and ppYy. The former two genotypes would result in plants with purple flowers and yellow peas, while the latter two genotypes would result in plants with white flowers with yellow peas, for a 1:1 ratio of each phenotype. You only need a 2 × 2 Punnett square (four squares total) to do this analysis because two of the alleles are homozygous.[/hidden-answer]

For the $F_2$ generation, the law of segregation requires that each gamete receive either an *R* allele or an *r* allele along with either a *Y* allele or a *y* allele. The law of independent assortment states that a gamete into which an *r* allele sorted would be equally likely to contain either a *Y* allele or a *y* allele. Thus, there are four equally likely gametes that can be formed when the *YyRr* heterozygote is self-crossed, as follows: *YR*, *Yr*, *yR*, and *yr*. Arranging these gametes along the top and left of a 4 × 4 Punnett square (Figure 2) gives us 16 equally likely genotypic combinations. From these genotypes, we infer a phenotypic ratio of 9 round/yellow:3 round/green:3 wrinkled/yellow:1 wrinkled/green (Figure 2). These are the offspring ratios we would expect, assuming we performed the crosses with a large enough sample size.

Because of independent assortment and dominance, the 9:3:3:1 dihybrid phenotypic ratio can be collapsed into two 3:1 ratios, characteristic of any monohybrid cross that follows a dominant and recessive pattern. Ignoring seed color and considering only

seed texture in the above dihybrid cross, we would expect that three quarters of the $F_2$ generation offspring would be round, and one quarter would be wrinkled. Similarly, isolating only seed color, we would assume that three quarters of the $F_2$ offspring would be yellow and one quarter would be green. The sorting of alleles for texture and color are independent events, so we can apply the product rule. Therefore, the proportion of round and yellow $F_2$ offspring is expected to be (3/4) × (3/4) = 9/16, and the proportion of wrinkled and green offspring is expected to be (1/4) × (1/4) = 1/16. These proportions are identical to those obtained using a Punnett square. Round, green and wrinkled, yellow offspring can also be calculated using the product rule, as each of these genotypes includes one dominant and one recessive phenotype. Therefore, the proportion of each is calculated as (3/4) × (1/4) = 3/16.

The law of independent assortment also indicates that a cross between yellow, wrinkled (*YYrr*) and green, round (*yyRR*) parents would yield the same $F_1$ and $F_2$ offspring as in the *YYRR* x *yyrr* cross.

The physical basis for the law of independent assortment also lies in meiosis I, in which the different homologous pairs line up in random orientations. Each gamete can contain any combination of paternal and maternal chromosomes (and therefore the genes on them) because the orientation of tetrads on the metaphase plane is random.

In pea plants, purple flowers (P) are dominant to white flowers (p) and yellow peas (Y) are dominant to green peas (y). What are the possible genotypes and phenotypes for a cross between PpYY and ppYy pea plants? How many squares do you need to do a Punnett square analysis of this cross? The possible genotypes are PpYY, PpYy, ppYY, and ppYy. The former two genotypes would result in plants with purple flowers and yellow peas, while the latter two genotypes would result in plants with white flowers with yellow peas, for a 1:1 ratio of each phenotype. You only need a 2 × 2 Punnett square (four squares total) to do this analysis because two of the alleles are homozygous.

## Forked-Line Method

When more than two genes are being considered, the Punnett-square method becomes unwieldy. For instance, examining a cross involving four genes would require a 16 × 16 grid containing 256 boxes. It would be extremely cumbersome to manually enter each genotype. For more complex crosses, the forked-line and probability methods are preferred.

To prepare a forked-line diagram for a cross between $F_1$ heterozygotes resulting from a cross between *AABBCC* and *aabbcc* parents, we first create rows equal to the number of genes being considered, and then segregate the alleles in each row on forked lines according to the probabilities for individual monohybrid crosses (Figure 3). We then multiply the values along each forked path to obtain the $F_2$ offspring probabilities. Note that this process is a diagrammatic version of the product rule. The values along each forked pathway can be multiplied because each gene assorts independently. For a trihybrid cross, the $F_2$ phenotypic ratio is 27:9:9:9:3:3:3:1.





Figure 3. The forked-line method can be used to analyze a trihybrid cross. Here, the probability for color in the $F_2$ generation occupies the top row (3 yellow:1 green). The probability for shape occupies the second row (3 round:1 wrinked), and the probability for height occupies the third row (3 tall:1 dwarf). The probability for each possible combination of traits is calculated by multiplying the probability for each individual trait. Thus, the probability of $F_2$ offspring having yellow, round, and tall traits is 3 × 3 × 3, or 27.

### Probability Method

While the forked-line method is a diagrammatic approach to keeping track of probabilities in a cross, the probability method gives the proportions of offspring expected to exhibit each phenotype (or genotype) without the added visual assistance. Both methods make use of the product rule and consider the alleles for each gene separately. Earlier, we examined the phenotypic proportions for a trihybrid cross using the forked-line method; now we will use the probability method to examine the genotypic proportions for a cross with even more genes.

For a trihybrid cross, writing out the forked-line method is tedious, albeit not as tedious as using the Punnett-square method. To fully demonstrate the power of the probability method, however, we can consider specific genetic calculations. For instance, for a tetrahybrid cross between individuals that are heterozygotes for all four genes, and in which all four genes are sorting independently and in a dominant and recessive pattern, what proportion of the offspring will be expected to be homozygous recessive for all four alleles? Rather than writing out every possible genotype, we can use the probability method. We know that for each gene, the fraction of homozygous recessive offspring will be 1/4. Therefore, multiplying this fraction for each of the four genes, (1/4) × (1/4) × (1/4) × (1/4), we determine that 1/256 of the offspring will be quadruply homozygous recessive.

For the same tetrahybrid cross, what is the expected proportion of offspring that have the dominant phenotype at all four loci? We can answer this question using phenotypic proportions, but let's do it the hard way—using genotypic proportions. The question asks for the proportion of offspring that are 1) homozygous dominant at *A* or heterozygous at *A*, and 2) homozygous at *B* or heterozygous at *B*, and so on. Noting the "or" and "and" in each circumstance makes clear where to apply the sum and product rules. The probability of a homozygous dominant at *A* is 1/4 and the probability of a heterozygote at *A* is 1/2. The probability of the homozygote or the heterozygote is 1/4 + 1/2 = 3/4 using the sum rule. The same probability can be obtained in the same way for each of the other genes, so that the probability of a dominant phenotype at *A* and *B* and *C* and *D* is, using the product rule, equal to 3/4 × 3/4 × 3/4 × 3/4, or 27/64. If you are ever unsure about how to combine probabilities, returning to the forked-line method should make it clear.

### Rules for Multihybrid Fertilization

Predicting the genotypes and phenotypes of offspring from given crosses is the best way to test your knowledge of Mendelian genetics. Given a multihybrid cross that obeys independent assortment and follows a dominant and recessive pattern, several generalized rules exist; you can use these rules to check your results as you work through genetics calculations (Table 1). To apply these rules, first you must determine *n*, the number of heterozygous gene pairs (the number of genes segregating two alleles each). For example, a cross between *AaBb* and *AaBb* heterozygotes has an *n* of 2. In contrast, a cross between *AABb* and *AABb* has an *n* of 1 because *A* is not heterozygous.

Table 1. General Rules for Multihybrid Crosses

| General Rule | Number of Heterozygous Gene Pairs |
|---|---|
| Number of different $F_1$ gametes | $2^n$ |
| Number of different $F_2$ genotypes | $3^n$ |
| Given dominant and recessive inheritance, the number of different $F_2$ phenotypes | $2^n$ |

## Linked Genes Violate the Law of Independent Assortment

Although all of Mendel's pea characteristics behaved according to the law of independent assortment, we now know that some allele combinations are not inherited independently of each other. Genes that are located on separate non-homologous chromosomes will always sort independently. However, each chromosome contains hundreds or thousands of genes, organized linearly on chromosomes like beads on a string. The segregation of alleles into gametes can be influenced by linkage, in which genes that are located physically close to each other on the same chromosome are more likely to be inherited as a pair. However, because of the process of recombination, or "crossover," it is possible for two genes on the same chromosome to behave independently, or as if they are not linked. To understand this, let's consider the biological basis of gene linkage and recombination.

Homologous chromosomes possess the same genes in the same linear order. The alleles may differ on homologous chromosome pairs, but the genes to which they correspond do not. In preparation for the first division of meiosis, homologous chromosomes replicate and synapse. Like genes on the homologs align with each other. At this stage, segments of homologous chromosomes exchange linear segments of genetic material (Figure 4). This process is called recombination, or crossover, and it is a common genetic process. Because the genes are aligned during recombination, the gene order is not altered. Instead, the result of recombination is that maternal and paternal alleles are combined onto the same chromosome. Across a given chromosome, several recombination events may occur, causing extensive shuffling of alleles.





Figure 4. The process of crossover, or recombination, occurs when two homologous chromosomes align during meiosis and exchange a segment of genetic material. Here, the alleles for gene C were exchanged. The result is two recombinant and two non-recombinant chromosomes.

When two genes are located in close proximity on the same chromosome, they are considered linked, and their alleles tend to be transmitted through meiosis together. To exemplify this, imagine a dihybrid cross involving flower color and plant height in which the genes are next to each other on the chromosome. If one homologous chromosome has alleles for tall plants and red flowers, and the other chromosome has genes for short plants and yellow flowers, then when the gametes are formed, the tall and red alleles will go together into a gamete and the short and yellow alleles will go into other gametes. These are called the parental genotypes because they have been inherited intact from the parents of the individual producing gametes. But unlike if the genes were on different chromosomes, there will be no gametes with tall and yellow alleles and no gametes with short and red alleles. If you create the Punnett square with these gametes, you will see that the classical Mendelian prediction of a 9:3:3:1 outcome of a dihybrid cross would not apply. As the distance between two genes increases, the probability of one or more crossovers between them increases, and the genes behave more like they are on separate chromosomes. Geneticists have used the proportion of recombinant gametes (the ones not like the parents) as a measure of how far apart genes are on a chromosome. Using this information, they have constructed elaborate maps of genes on chromosomes for well-studied organisms, including humans.

Mendel's seminal publication makes no mention of linkage, and many researchers have questioned whether he encountered linkage but chose not to publish those crosses out of concern that they would invalidate his independent assortment postulate. The garden pea has seven chromosomes, and some have suggested that his choice of seven characteristics was not a coincidence. However, even if the genes he examined were not located on separate chromosomes, it is possible that he simply did not observe linkage because of the extensive shuffling effects of recombination.

## Testing the Hypothesis of Independent Assortment

To better appreciate the amount of labor and ingenuity that went into Mendel's experiments, proceed through one of Mendel's dihybrid crosses.

**Question:** What will be the offspring of a dihybrid cross?

**Background:** Consider that pea plants mature in one growing season, and you have access to a large garden in which you can cultivate thousands of pea plants. There are several true-breeding plants with the following pairs of traits: tall plants with inflated pods, and dwarf plants with constricted pods. Before the plants have matured, you remove the pollen-producing organs from the tall/inflated plants in your crosses to prevent self-fertilization. Upon plant maturation, the plants are manually crossed by transferring pollen from the dwarf/constricted plants to the stigmata of the tall/inflated plants.

**Hypothesis:** Both trait pairs will sort independently according to Mendelian laws. When the true-breeding parents are crossed, all of the $F_1$ offspring are tall and have inflated pods, which indicates that the tall and inflated traits are dominant over the dwarf and constricted traits, respectively. A self-cross of the $F_1$ heterozygotes results in 2,000 $F_2$ progeny.

**Test the hypothesis:** Because each trait pair sorts independently, the ratios of tall:dwarf and inflated:constricted are each expected to be 3:1. The tall/dwarf trait pair is called *T/t*, and the inflated/constricted trait pair is designated *I/i*. Each member of the $F_1$ generation therefore has a genotype of *TtIi*. Construct a grid analogous to Figure 5, in which you cross two *TtIi* individuals. Each individual can donate four combinations of two traits: *TI*, *Ti*, *tI*, or *ti*, meaning that there are 16 possibilities of offspring genotypes. Because the *T* and *I* alleles are dominant, any individual having one or two of those alleles will express the tall or inflated phenotypes, respectively, regardless if they also have a *t* or *i* allele. Only individuals that are *tt* or *ii* will express the dwarf and constricted alleles, respectively. As shown in Figure 5, you predict that you will observe the following offspring proportions: tall/inflated : tall/constricted : dwarf/inflated : dwarf/constricted in a 9:3:3:1 ratio. Notice from the grid that when considering the tall/dwarf and inflated/constricted trait pairs in isolation, they are each inherited in 3:1 ratios.



Figure 5. This figure shows all possible combinations of offspring resulting from a dihybrid cross of pea plants that are heterozygous for the tall/dwarf and inflated/constricted alleles.

**Test the hypothesis:** You cross the dwarf and tall plants and then self-cross the offspring. For best results, this is repeated with hundreds or even thousands of pea plants. What special precautions should be taken in the crosses and in growing the plants?

**Analyze your data:** You observe the following plant phenotypes in the $F_2$ generation: 2706 tall/inflated, 930 tall/constricted, 888 dwarf/inflated, and 300 dwarf/constricted. Reduce these findings to a ratio and determine if they are consistent with Mendelian laws.

**Form a conclusion:** Were the results close to the expected 9:3:3:1 phenotypic ratio? Do the results support the prediction? What might be observed if far fewer plants were used, given that alleles segregate randomly into gametes? Try to imagine growing that many pea plants, and consider the potential for experimental error. For instance, what would happen if it was extremely windy one day?

## Learning Objectives

Mendel postulated that genes (characteristics) are inherited as pairs of alleles (traits) that behave in a dominant and recessive pattern. Alleles segregate into gametes such that each gamete is equally likely to receive either one of the two alleles present in a diploid individual. In addition, genes are assorted into gametes independently of one another. That is, alleles are generally not more likely to segregate into a gamete with a particular allele of another gene. A dihybrid cross demonstrates independent assortment when the genes in question are on different chromosomes or distant from each other on the same chromosome. For crosses involving more than two genes, use the forked line or probability methods to predict offspring genotypes and phenotypes rather than a Punnett square.

Although chromosomes sort independently into gametes during meiosis, Mendel's law of independent assortment refers to genes, not chromosomes, and a single chromosome may carry more than 1,000 genes. When genes are located in close proximity on the same chromosome, their alleles tend to be inherited together. This results in offspring ratios that violate Mendel's law of independent assortment. However, recombination serves to exchange genetic material on homologous chromosomes such that maternal and paternal alleles may be recombined on the same chromosome. This is why alleles on a given chromosome are not always inherited together. Recombination is a random event occurring anywhere on a chromosome. Therefore, genes that are far apart on the same chromosome are likely to still assort independently because of recombination events that occurred in the intervening chromosomal space.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

14.5: Laws of Inheritance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.6: Heredity

Watch this video for a nice summary of Mendel's contribution to the field of genetics and how a genotype leads to a phenotype.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=486

## Contributors and Attributions

All rights reserved content

- Heredity: Crash Course Biology #9. **Authored by**: CrashCourse. **Located at**: https://youtu.be/CBezq1fFUEA. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

14.6: Heredity is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.7: Introduction to Beyond Dominance and Recessiveness

## Explain complications to the phenotypic expression of genotype, including mutations

Mendel's experiments with pea plants suggested the following:

1. Two "units" or alleles exist for every gene.
2. Alleles maintain their integrity in each generation (no blending).
3. In the presence of the dominant allele, the recessive allele is hidden and makes no contribution to the phenotype.

Therefore, recessive alleles can be "carried" and not expressed by individuals. Such heterozygous individuals are sometimes referred to as "carriers." Further genetic studies in other plants and animals have shown that much more complexity exists, but that the fundamental principles of Mendelian genetics still hold true. In the sections to follow, we consider some of the extensions of Mendelism. If Mendel had chosen an experimental system that exhibited these genetic complexities, it's possible that he would not have understood what his results meant.

## What You'll Learn to Do

- Explain how a trait with incomplete dominance will appear in a population
- Explain how a trait with codominant inheritance will appear in a population
- Explain how a trait with sex-linkage will appear in a population
- Explain how mutli-allele inheritance will impact a trait within in a population
- Describe the impacts of penetrance and expressivity on a trait's expression in a population

### Learning Activities

The learning activities for this section include the following:

- Variations of Dominance
- Sex-Linked Traits
- Multiple Alleles
- Penetrance and Expressivity
- Self Check: Beyond Dominance and Recessiveness

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

14.7: Introduction to Beyond Dominance and Recessiveness is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.8: Non-Mendelian Inheritance

## Incomplete Dominance



Figure 1. These pink flowers of a heterozygote snapdragon result from incomplete dominance. (credit: “storebukkebruse”/Flickr)

Mendel’s results, that traits are inherited as dominant and recessive pairs, contradicted the view at that time that offspring exhibited a blend of their parents’ traits. However, the heterozygote phenotype occasionally does appear to be intermediate between the two parents. For example, in the snapdragon, *Antirrhinum majus* (Figure 1), a cross between a homozygous parent with white flowers ($C^WC^W$) and a homozygous parent with red flowers ($C^RC^R$) will produce offspring with pink flowers ($C^RC^W$). (Note that different genotypic abbreviations are used for Mendelian extensions to distinguish these patterns from simple dominance and recessiveness.) This pattern of inheritance is described as **incomplete dominance**, denoting the expression of two contrasting alleles such that the individual displays an intermediate phenotype. The allele for red flowers is incompletely dominant over the allele for white flowers. However, the results of a heterozygote self-cross can still be predicted, just as with Mendelian dominant and recessive crosses. In this case, the genotypic ratio would be 1 $C^RC^R$:2 $C^RC^W$:1 $C^WC^W$, and the phenotypic ratio would be 1:2:1 for red:pink:white.

Incomplete dominance can be seen in several types of flowers, including pink tulips, carnations and roses—any pink flowers in these are due to the mixing of red and white alleles. Incomplete dominance can also be observed in some animals, such as rabbits. When a long-furred Angora breeds with a short-furred Rex, the offspring have medium-length fur. Tail length in dogs is similarly impacted by genes that display incomplete dominance patterns.

## Codominant Inheritance



Figure 2. Red Roan Horse

A variation on incomplete dominance is **codominance**, in which both alleles for the same characteristic are simultaneously expressed in the heterozygote. An example of codominance is the MN blood groups of humans. The M and N alleles are expressed in the form of an M or N antigen present on the surface of red blood cells. Homozygotes ($L^ML^M$ and $L^NL^N$) express either the M or the N allele, and heterozygotes ($L^ML^N$) express both alleles equally. In a self-cross between heterozygotes expressing a codominant

trait, the three possible offspring genotypes are phenotypically distinct. However, the 1:2:1 genotypic ratio characteristic of a Mendelian monohybrid cross still applies.

Codominance can also be seen in human blood types: the AB blood type is a result of both the $I^A$ allele and the $I^B$ allele being codominant. The roan coat color in horses is also an example of codominance. A “red” roan results from the mating of a chestnut parent and a white parent (Figure 2). We know this is codominance because individual hairs are either chestnut or they are white, leading to the red roan overall appearance.

**Practice Question**

So what’s the difference between incomplete dominance and codominant inheritance? While they are very similar, the key difference is this: in incomplete dominance, the two traits are blended together, whereas in codominance, both traits are expressed.

We’ve already discussed incomplete dominance in flowers (Figure 1). What do you think a flower would look like if the red and white phenotypes were codominant instead?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”528012″]Show Answer[/reveal-answer]
[hidden-answer a=”528012″]The flower would have both red and white petals, like this Rhododendron:



Figure 3. Codominance is shown in this rhododendron.

[/hidden-answer]

## Sex-Linked Traits

In humans, as well as in many other animals and some plants, the sex of the individual is determined by sex chromosomes. The sex chromosomes are one pair of non-homologous chromosomes. Until now, we have only considered inheritance patterns among non-sex chromosomes, or **autosomes**. In addition to 22 homologous pairs of autosomes, human females have a homologous pair of X chromosomes, whereas human males have an XY chromosome pair. Although the Y chromosome contains a small region of similarity to the X chromosome so that they can pair during meiosis, the Y chromosome is much shorter and contains many fewer genes. When a gene being examined is present on the X chromosome, but not on the Y chromosome, it is said to be **X-linked**.



Figure 4. In Drosophila, the gene for eye color is located on the X chromosome. Clockwise from top left are brown, cinnabar, sepia, vermilion, white, and red. Red eye color is wild-type and is dominant to white eye color.

Eye color in *Drosophila* was one of the first X-linked traits to be identified. Thomas Hunt Morgan mapped this trait to the X chromosome in 1910. Like humans, *Drosophila* males have an XY chromosome pair, and females are XX. In flies, the wild-type eye color is red ($X^W$) and it is dominant to white eye color ($X^w$) (Figure 4). Because of the location of the eye-color gene, reciprocal crosses do not produce the same offspring ratios. Males are said to be **hemizygous**, because they have only one allele for any X-linked characteristic. Hemizygosity makes the descriptions of dominance and recessiveness irrelevant for XY males. *Drosophila* males lack a second allele copy on the Y chromosome; that is, their genotype can only be $X^WY$ or $X^wY$. In contrast, females have two allele copies of this gene and can be $X^WX^W$, $X^WX^w$, or $X^wX^w$.

In an X-linked cross, the genotypes of $F_1$ and $F_2$ offspring depend on whether the recessive trait was expressed by the male or the female in the $P_0$ generation. With regard to *Drosophila* eye color, when the $P_0$ male expresses the white-eye phenotype and the female is homozygous red-eyed, all members of the $F_1$ generation exhibit red eyes. The $F_1$ females are heterozygous ($X^WX^w$), and the males are all $X^WY$, having received their X chromosome from the homozygous dominant $P_0$ female and their Y chromosome from the $P_0$ male. A subsequent cross between the $X^WX^w$ female and the $X^WY$ male would produce only red-eyed females (with $X^WX^W$ or $X^WX^w$ genotypes) and both red- and white-eyed males (with $X^WY$ or $X^wY$ genotypes). Now, consider a cross between a homozygous white-eyed female and a male with red eyes (Figure 5). The $F_1$ generation would exhibit only heterozygous red-eyed females ($X^WX^w$) and only white-eyed males ($X^wY$). Half of the $F_2$ females would be red-eyed ($X^WX^w$) and half would be white-eyed ($X^wX^w$). Similarly, half of the $F_2$ males would be red-eyed ($X^WY$) and half would be white-eyed ($X^wY$).

Practice Question





Figure 5. Punnett square analysis is used to determine the ratio of offspring from a cross between a red-eyed male fruit fly and a white-eyed female fruit fly.

What ratio of offspring would result from a cross between a white-eyed male and a female that is heterozygous for red eye color?

[practice-area rows="2"][/practice-area]
[reveal-answer q="722618"]**Show Answer**[/reveal-answer]
[hidden-answer a="722618"]Half of the female offspring would be heterozygous ($X^WX^w$) with red eyes, and half would be homozygous recessive ($X^wX^w$) with white eyes. Half of the male offspring would be hemizygous dominant ($X^WY$) with red yes, and half would be hemizygous recessive ($X^wY$) with white eyes.[/hidden-answer]

Discoveries in fruit fly genetics can be applied to human genetics. When a female parent is homozygous for a recessive X-linked trait, she will pass the trait on to 100 percent of her offspring. Her male offspring are, therefore, destined to express the trait, as they will inherit their father's Y chromosome. In humans, the alleles for certain conditions (some forms of color blindness, hemophilia, and muscular dystrophy) are X-linked. Females who are heterozygous for these diseases are said to be carriers and may not exhibit any phenotypic effects. These females will pass the disease to half of their sons and will pass carrier status to half of their daughters; therefore, recessive X-linked traits appear more frequently in males than females.

In some groups of organisms with sex chromosomes, the gender with the non-homologous sex chromosomes is the female rather than the male. This is the case for all birds. In this case, sex-linked traits will be more likely to appear in the female, in which they are hemizygous.

## Contributors and Attributions

CC licensed content, Original

- Modification and original content. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Co-dominance Rhododendron. **Authored by**: darwin cruz. **Located at**: https://commons.wikimedia.org/wiki/File:Co-dominance_Rhododendron.jpg. **License**: *CC BY: Attribution*
- Red roan Quarter Horse. **Authored by**: Betty Wills. **Provided by**: Wikimedia Commons. **Located at**: https://commons.wikimedia.org/wiki/File:Red_roan_Quarter_Horse.jpg. **License**: *CC BY-SA: Attribution-ShareAlike*

---

14.8: Non-Mendelian Inheritance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.9: Non-Mendelian Punnett Squares

This practice activity will help you remember the difference between types of non-Mendelian inheritance and remember just how they work.

A link to an interactive elements can be found at the bottom of this page.

Click here for a text-only version of the activity.

## Video Review

Watch this video for a summary of the three “special” cases of non-Mendelian inheritance you just practiced.

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

CC licensed content, Original

- Non-Mendelian Punnett Squares. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Non-Mendelian Trait Inheritance. **Provided by**: Lumen Learning. **Located at**: www.oppia.org/explore/huVa7Jm_04aL. **License**: *CC BY-SA: Attribution-ShareAlike*

All rights reserved content

- Incomplete Dominance, Codominance, and Sex-Linked. **Authored by**: MissKmetBioClass. **Located at**: https://youtu.be/fQvER3MyI2c. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

14.9: Non-Mendelian Punnett Squares is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.10: Multiple Alleles

Mendel implied that only two alleles, one dominant and one recessive, could exist for a given gene. We now know that this is an oversimplification. Although individual humans (and all diploid organisms) can only have two alleles for a given gene, multiple alleles may exist at the population level such that many combinations of two alleles are observed. Note that when many alleles exist for the same gene, the convention is to denote the most common phenotype or genotype among wild animals as the **wild type** (often abbreviated "+"); this is considered the standard or norm. All other phenotypes or genotypes are considered **variants** of this standard, meaning that they deviate from the wild type. The variant may be recessive or dominant to the wild-type allele.

An example of multiple alleles is coat color in rabbits (Figure 1). Here, four alleles exist for the *c* gene. The wild-type version, $C^+C^+$, is expressed as brown fur. The chinchilla phenotype, $c^{ch}c^{ch}$, is expressed as black-tipped white fur. The Himalayan phenotype, $c^hc^h$, has black fur on the extremities and white fur elsewhere. Finally, the albino, or "colorless" phenotype, *cc*, is expressed as white fur. In cases of multiple alleles, dominance hierarchies can exist. In this case, the wild-type allele is dominant over all the others, chinchilla is incompletely dominant over Himalayan and albino, and Himalayan is dominant over albino. This hierarchy, or allelic series, was revealed by observing the phenotypes of each possible heterozygote offspring.





Figure 1. Four different alleles exist for the rabbit coat color (*C*) gene.



Figure 2. As seen in comparing the wild-type *Drosophila* (left) and the *Antennapedia* mutant (right), the Antennapedia mutant has legs on its head in place of antennae.

The complete dominance of a wild-type phenotype over all other mutants often occurs as an effect of "dosage" of a specific gene product, such that the wild-type allele supplies the correct amount of gene product whereas the mutant alleles cannot. For the allelic

series in rabbits, the wild-type allele may supply a given dosage of fur pigment, whereas the mutants supply a lesser dosage or none at all. Interestingly, the Himalayan phenotype is the result of an allele that produces a temperature-sensitive gene product that only produces pigment in the cooler extremities of the rabbit's body.

Alternatively, one mutant allele can be dominant over all other phenotypes, including the wild type. This may occur when the mutant allele somehow interferes with the genetic message so that even a heterozygote with one wild-type allele copy expresses the mutant phenotype. One way in which the mutant allele can interfere is by enhancing the function of the wild-type gene product or changing its distribution in the body.

One example of this is the *Antennapedia* mutation in *Drosophila* (Figure 2). In this case, the mutant allele expands the distribution of the gene product, and as a result, the *Antennapedia* heterozygote develops legs on its head where its antennae should be.

## Multiple Alleles Confer Drug Resistance in the Malaria Parasite

Malaria is a parasitic disease in humans that is transmitted by infected female mosquitoes, including *Anopheles gambiae* (Figure 3a), and is characterized by cyclic high fevers, chills, flu-like symptoms, and severe anemia. *Plasmodium falciparum* and *P. vivax* are the most common causative agents of malaria, and *P. falciparum* is the most deadly (Figure 3b). When promptly and correctly treated, *P. falciparum*malaria has a mortality rate of 0.1 percent. However, in some parts of the world, the parasite has evolved resistance to commonly used malaria treatments, so the most effective malarial treatments can vary by geographic region.






Figure 3. The (a) Anopheles gambiae, or African malaria mosquito, acts as a vector in the transmission to humans of the malaria-causing parasite (b) Plasmodium falciparum, here visualized using false-color transmission electron microscopy. (credit a: James D. Gathany; credit b: Ute Frevert; false color by Margaret Shear; scale-bar data from Matt Russell)

In Southeast Asia, Africa, and South America, *P. falciparum* has developed resistance to the anti-malarial drugs chloroquine, mefloquine, and sulfadoxine-pyrimethamine. *P. falciparum*, which is haploid during the life stage in which it is infectious to humans, has evolved multiple drug-resistant mutant alleles of the *dhps* gene. Varying degrees of sulfadoxine resistance are associated with each of these alleles. Being haploid, *P. falciparum* needs only one drug-resistant allele to express this trait.

In Southeast Asia, different sulfadoxine-resistant alleles of the *dhps* gene are localized to different geographic regions. This is a common evolutionary phenomenon that occurs because drug-resistant mutants arise in a population and interbreed with other *P. falciparum* isolates in close proximity. Sulfadoxine-resistant parasites cause considerable human hardship in regions where this drug is widely used as an over-the-counter malaria remedy. As is common with pathogens that multiply to large numbers within an infection cycle, *P. falciparum* evolves relatively rapidly (over a decade or so) in response to the selective pressure of commonly used anti-malarial drugs. For this reason, scientists must constantly work to develop new drugs or drug combinations to combat the worldwide malaria burden.[1]

## Multiple Alleles (ABO Blood Types) and Punnett Squares




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=494

---

1. Sumiti Vinayak, et al., "Origin and Evolution of Sulfadoxine Resistant *Plasmodium falciparum*," *Public Library of Science Pathogens* 6, no. 3 (2010): e1000830, doi:10.1371/journal.ppat.1000830. ↵

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Multiple Alleles (ABO Blood Types) and Punnett Squares. **Authored by**: Amoeba Sisters. **Located at**: https://youtu.be/9O5JQqlngFY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

14.10: Multiple Alleles is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.11: Introduction to Heredity and Disease

## Explain the conventions of a family pedigree and predict whether a disease will be passed through a family in one of three modes

Over the years, we've seen that some diseases are hereditary. In this outcome, we'll learn just how some diseases can be passed through a family line.

## What You'll Learn to Do

- Understand why geneticists use pedigrees to track human diseases
- Understand how a pedigree analysis is used to identify the inheritance pattern of a genetic disorder

### Learning Activities

The learning activities for this section include the following:

- Pedigrees and Disease
- Pedigree Flowchart
- Genetic Disorder and Pedigrees
- Self Check: Heredity and Disease

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

14.11: Introduction to Heredity and Disease is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

## 14.12: Penetrance and Expressivity

Penetrance refers to the probability of a gene or trait being expressed. In some cases, despite the presence of a dominant allele, a phenotype may not be present. One example of this is polydactyly in humans (extra fingers and/or toes). A dominant allele produces polydactyly in humans but not all humans with the allele display the extra digits. “Complete” penetrance means the gene or genes for a trait are expressed in all the population who have the genes. “Incomplete” or ‘reduced’ penetrance means the genetic trait is expressed in only part of the population. The penetrance of expression may also change in different age groups of a population. Reduced penetrance probably results from a combination of genetic, environmental, and lifestyle factors, many of which are unknown. This phenomenon can make it challenging for genetics professionals to interpret a person’s family medical history and predict the risk of passing a genetic condition to future generations.





Illustration modeled after similar image by Steven M. Carr, Penetrance versus expressivity.

Expressivity on the other hand refers to variation in phenotypic expression when an allele is penetrant. Back to the polydactyly example, an extra digit may occur on one or more appendages. The digit can be full size or just a stub. Hence, this allele has reduced penetrance as well as variable expressivity. Variable expressivity refers to the range of signs and symptoms that can occur in different people with the same genetic condition. As with reduced penetrance, variable expressivity is probably caused by a combination of genetic, environmental, and lifestyle factors, most of which have not been identified. If a genetic condition has highly variable signs and symptoms, it may be challenging to diagnose.





Illustration modeled after similar image by Steven M. Carr, Penetrance versus expressivity.
For more information about reduced penetrance and variable expressivity take a look at the interactive tutorial on penetrance the PHG Foundation offers. The tutorial explains the differences between reduced penetrance and variable expressivity.

### Variable Expressivity and Incomplete Penetrance

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for “Variable Expressivity and Incomplete Penetrance” here (link opens in new window).

### Contributors and Attributions

All rights reserved content

- Variable Expressivity and Incomplete Penetrance. **Authored by**: Learning Simply. **Located at**: https://youtu.be/1XW-j3VQGRY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

Public domain content

- What are reduced penetrance and variable expressivity?. **Provided by**: National Institutes of Health. **Located at**: http://ghr.nlm.nih.gov/handbook/inheritance/penetranceexpressivity. **Project**: Genetics Home Reference. **License**: *Public Domain: No Known Copyright*

14.12: Penetrance and Expressivity is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.13: Pedigrees and Disease

Health care professionals have known for a long time that common diseases—like heart disease, cancer, and diabetes—and even rare diseases—like hemophilia, cystic fibrosis, and sickle cell anemia—can run in families. If one generation of a family has high blood pressure, it is not unusual for the next generation to have similarly high blood pressure. Therefore, family history can be a powerful screening tool and has often been referred to as the best "genetic test."

Both common diseases and rare diseases can run in families. Therefore, family history can be a powerful screening tool. Family history should be updated on each visit and patients should be made aware of its significance to their health.

## Importance of Family History

Family history holds important information about an individual's past and future life. Family history can be used as a diagnostic tool and help guide decisions about genetic testing for the patient and at-risk family members. If a family is affected by a disease, an accurate family history will be important to establish a pattern of transmission. In addition, a family history can even help to exclude genetic diseases, particularly for common diseases in which lifestyle and environment play strong roles. Lastly, a family history can identify potential health problems that an individual may be at increased risk for in the future. Early identification of increased risk can allow the individual and health professional to take steps to reduce risk by implementing lifestyle changes and increasing disease surveillance.

While many of the well-known genetic disorders are of childhood onset, many complex, adult-onset conditions can also run in families. For example, about 5 to 10 percent of all breast cancers are hereditary. These cancers may be caused by mutations in particular genes, such as BRCA1 or BRCA2. An individual may be at high risk of hereditary breast cancer and genetic testing should be considered if her family history includes more than one first-degree (mother, sister, or daughter) or second-degree relative (aunt, grandmother, or niece) with breast or ovarian cancer, particularly if the diagnosis of breast or ovarian cancer in those relatives was at a young age (50 or younger).

Another example of an adult-onset disease that can be inherited is Alzheimer's disease. Although most of Alzheimer's disease cases are sporadic, a small number are hereditary. Hereditary Alzheimer's disease is an extremely aggressive form of the disease and typically manifests before the age of 65. Three genes that cause early-onset Alzheimer's disease have been identified.

Notwithstanding the importance of family history to help define occurrence of a genetic disorder within a family, it should be noted that some genetic diseases are caused by spontaneous mutations, such as for single gene disorders like Duchenne muscular dystrophy and hemophilia A, as well as for most cases of Down syndrome, chromosomal deletion syndromes, and other chromosomal disorders. Therefore, a genetic disorder cannot be ruled out in the absence of a family history.


## How to Take a Family Medical History

A basic family history should include three generations. To begin taking a family history, start by asking the patient about his/her health history and then ask about siblings and parents. Questions should include:


1. General information such as names and birthdates
2. Family's origin or racial/ethnic background
3. Health status
4. Age at death and cause of death of each family member
5. Pregnancy outcomes of the patient and genetically-related relatives It may be easier to list all the members of the nuclear family first and then go back and ask about the health status of each one. After you have taken the family history of the patient's closest relatives, go back one generation at a time and ask about aunts, uncles, grandparents, and first cousins.

## Pedigrees

One way to record a family history is by drawing a family tree called a "pedigree." A pedigree represents family members and relationships using standardized symbols (see below). As patients relate information to you about their family history, a pedigree can be drawn much quicker than recording the information in writing and allows patterns of disease to emerge as the pedigree is drawn. Since the family history is continually changing, the pedigree can be easily updated on future visits. Patients should be encouraged to record information and update their family history regularly.




The sample pedigree below contains information such as age or date of birth (and, for all deceased family members, age at death and cause of death), major medical problems with age of onset, birth defects, learning problems and mental retardation, and vision loss/hearing loss at a young age. For family members with known medical problems, ask whether they smoke, what their diet and exercise habits are if known, and if they are overweight.




[reveal-answer q="724528"]Show References[/reveal-answer]
[hidden-answer a="724528"]

American Society for Human Genetics. (2004). *Your Family History*. www.ashg.org/genetics/ashg/educ/007.shtml.

Bennett RL. *The Practical Guide to the Genetic Family History.* New York: Wiley-Liss, Inc. 1999.

Centers for Disease Control and Prevention. Office of Genomics and Disease Prevention. *Using Family History to Promote Health*. www.cdc.gov/genomics/public/famhistMain.htm.

Genetic Alliance. (2004). *Taking a Family History.* www.geneticalliance.org/ws_di...asp?filter=fhh.

*March of Dimes–Genetics and Your Practice*. http://www.marchofdimes.com/gyponline/index.bm2.

*My Family Health Portrait*. familyhistory.genome.gov.

U.S. Department of Health and Human Services. (2004) *U.S. Surgeon General's Family History Initiative.* http://www.hhs.gov/familyhistory/.

[/hidden-answer]

## Contributors and Attributions

CC licensed content, Shared previously

- Chapter 3: Pedigree and Family History Taking. **Authored by**: Genetic Alliance. **Provided by**: National Center for Biotechnology Information. **Located at**: www.ncbi.nlm.nih.gov/books/NBK132175/. **Project**: Understanding Genetics: A New England Guide for Patients and Health Professionals. **License**: *CC BY: Attribution*

---

14.13: Pedigrees and Disease is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.14: Pedigree Flowchart

Watch this video to understand the basics of conducting a pedigree analysis of a human genetic disorder. At the end, you should feel comfortable with the basic steps used to conduct a pedigree investigation.

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

All rights reserved content

- Pedigree Flowchart. **Authored by**: Biologybyme's channel. **Located at**: https://youtu.be/z0cwhooDZ0M. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

14.14: Pedigree Flowchart is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.15: Genetic Disorder and Pedigrees

The following video provides a summary of all you have just learned about pedigrees, including differences in family inheritance patterns based on autosomal dominant, autosomal recessive, or sex-linked inheritance of a particular characteristic.

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

All rights reserved content

- Video Lecture: 9-3 Genetic Disorder and Pedigrees. **Authored by**: jlambbio. **Located at**: https://youtu.be/-Su7OAqU72U. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

14.15: Genetic Disorder and Pedigrees is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.16: Introduction to Genetics and the Environment

## Discuss the role environment plays on phenotypes

In recent years, scientists have begun to research how our environment can impact our phenotypes. In this outcome, we'll learn about a few different ways our environment can impact us.

## What You'll Learn to Do

- Explain how epistasis impacts trait expression
- Describe polygenic inheritance and how to recognize it
- Describe continuous variation and how to recognize it
- Identify gene-environment interaction and how this impacts trait expression
- Explain pleiotropy and its impact on traits in a population

### Learning Activities

The learning activities for this section include the following:

- Epistasis
- Polygenic Inheritance and Environmental Effects
- Effect of the Environment
- Pleiotropy and Human Disorders
- Self Check: Genetics and the Environment

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

14.16: Introduction to Genetics and the Environment is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.17: Epistasis

Mendel's studies in pea plants implied that the sum of an individual's phenotype was controlled by genes (or as he called them, unit factors), such that every characteristic was distinctly and completely controlled by a single gene. In fact, single observable characteristics are almost always under the influence of multiple genes (each with two or more alleles) acting in unison. For example, at least eight genes contribute to eye color in humans.

Eye color in humans is determined by multiple genes. Use the Eye Color Calculator to predict the eye color of children from parental eye color.

In some cases, several genes can contribute to aspects of a common phenotype without their gene products ever directly interacting. In the case of organ development, for instance, genes may be expressed sequentially, with each gene adding to the complexity and specificity of the organ. Genes may function in complementary or synergistic fashions, such that two or more genes need to be expressed simultaneously to affect a phenotype. Genes may also oppose each other, with one gene modifying the expression of another.

In epistasis, the interaction between genes is antagonistic, such that one gene masks or interferes with the expression of another. "Epistasis" is a word composed of Greek roots that mean "standing upon." The alleles that are being masked or silenced are said to be hypostatic to the epistatic alleles that are doing the masking. Often the biochemical basis of epistasis is a gene pathway in which the expression of one gene is dependent on the function of a gene that precedes or follows it in the pathway.

An example of epistasis is pigmentation in mice. The wild-type coat color, agouti (*AA*), is dominant to solid-colored fur (*aa*). However, a separate gene (*C*) is necessary for pigment production. A mouse with a recessive *c* allele at this locus is unable to produce pigment and is albino regardless of the allele present at locus *A* (Figure 1). Therefore, the genotypes *AAcc*, *Aacc*, and *aacc* all produce the same albino phenotype. A cross between heterozygotes for both genes (*AaCc* x *AaCc*) would generate offspring with a phenotypic ratio of 9 agouti:3 solid color:4 albino (Figure 1). In this case, the *C* gene is epistatic to the *A* gene.



Figure 1. In mice, the mottled agouti coat color (*A*) is dominant to a solid coloration, such as black or gray. A gene at a separate locus (*C*) is responsible for pigment production. The recessive *c* allele does not produce pigment, and a mouse with the homozygous recessive *cc* enotype is albino regardless of the allele present at the *A* locus. Thus, the *C* gene is epistatic to the *A* gene.

Epistasis can also occur when a dominant allele masks expression at a separate gene. Fruit color in summer squash is expressed in this way. Homozygous recessive expression of the *W* gene (*ww*) coupled with homozygous dominant or heterozygous expression of the *Y* gene (*YY* or *Yy*) generates yellow fruit, and the *wwyy* genotype produces green fruit. However, if a dominant copy of the *W* gene is present in the homozygous or heterozygous form, the summer squash will produce white fruit regardless of the *Y* alleles. A cross between white heterozygotes for both genes (*WwYy* × *WwYy*) would produce offspring with a phenotypic ratio of 12 white:3 yellow:1 green.

Finally, epistasis can be reciprocal such that either gene, when present in the dominant (or recessive) form, expresses the same phenotype. In the shepherd's purse plant (*Capsella bursa-pastoris*), the characteristic of seed shape is controlled by two genes in a dominant epistatic relationship. When the genes *A* and *B* are both homozygous recessive (*aabb*), the seeds are ovoid. If the dominant allele for either of these genes is present, the result is triangular seeds. That is, every possible genotype other than *aabb* results in triangular seeds, and a cross between heterozygotes for both genes (*AaBb* x *AaBb*) would yield offspring with a phenotypic ratio of 15 triangular:1 ovoid.

As you work through genetics problems, keep in mind that any single characteristic that results in a phenotypic ratio that totals 16 is typical of a two-gene interaction. Recall the phenotypic inheritance pattern for Mendel's dihybrid cross, which considered two

non-interacting genes—9:3:3:1. Similarly, we would expect interacting gene pairs to also exhibit ratios expressed as 16 parts. Note that we are assuming the interacting genes are not linked; they are still assorting independently into gametes.

For an excellent review of Mendel's experiments and to perform your own crosses and identify patterns of inheritance, visit the Mendel's Peas web lab.

Whether or not they are sorting independently, genes may interact at the level of gene products such that the expression of an allele for one gene masks or modifies the expression of an allele for a different gene. This is called epistasis.

## In Summary: Epistasis

Epistasis is a form on non-Mendelian inheritance in which one gene is capable of interfering with expression of another. This is often found associated with gene pathways where the expression of one gene is directly dependent on the presence or absence of another gene product within the pathway.

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Epistasis!. **Authored by**: Science Club!. **Located at**: https://youtu.be/ctjw-ZijciM. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

14.17: Epistasis is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.18: Polygenic Inheritance and Environmental Effects

## How is Height Inherited?

Many heritable human characteristics don't seem to follow Mendelian rules in their inheritance patterns. For example, consider human height. Unlike a simple Mendelian characteristic, human height displays:

- **Continuous variation.** Unlike Mendel's pea plants, humans don't come in two clear-cut "tall" and "short" varieties. In fact, they don't even come in four heights, or eight, or sixteen. Instead, it's possible to get humans of many different heights, and height can vary in increments of inches or fractions of inches. As an example, consider the bell curve-shaped graph in Figure 1, which shows the heights of a group of male high school seniors.
- **A complex inheritance pattern.** If you've paid attention to the heights of your friends and family, you may have noticed that many different patterns of inheritance are possible. Tall parents can have a short child, short parents can have a tall child, and two parents of different heights may or may not have a child of intermediate height. In addition, siblings with the same two parents may have a range of heights, ones that don't fall into clear, distinct categories. Simple models involving one or two genes can't accurately predict all of these inheritance patterns.





Figure 1. Heights of male high school seniors. Image modified from "Continuous variation: Quantitative traits," by J. W. Kimball (CC BY 3.0).


How, then, is height inherited? Height and other similar features are controlled not just by one gene, but rather, by multiple (often many) genes that each make a small contribution to the overall outcome. This inheritance pattern is called **polygenic inheritance** (*poly–* = many). For instance, a recent study found over 400 genes linked to variation in height[1]. When there are large numbers of genes involved, it becomes hard to distinguish the effect of each individual gene, and even harder to see that gene variants (alleles) are inherited according to Mendelian rules. In a further complication, height doesn't just depend on genetics: it also depends a lot on environmental factors, such as a child's overall health and the type of nutrition he or she receives while growing up.

In this article, we'll look in more detail at how complex human traits such as height are inherited, as well as how factors like genetic background and environment can influence the **phenotype** (observable features) produced by a particular **genotype** (set of gene variants, or alleles).

## Polygenic Inheritance

Some human characteristics, such as height, eye color, and hair color, don't come in just a few distinct forms. Instead, they vary in small gradations, forming a spectrum or continuum of possible phenotypes. Features like these are called quantitative characters, and they're typically controlled by multiple genes (often, many genes), each of which contributes to the overall phenotype. For example, although there are two major eye color genes, there are at least 14 additional genes that play roles in determining a person's exact eye color[2].

Looking at a real example of a human polygenic trait would get complicated, largely because we'd have to keep track of tens, or even hundreds, of different allele pairs. However, we can use an example involving the color of wheat kernels to see how Mendelian inheritance of multiple genes (plus a little incomplete dominance of alleles) can produce a broad spectrum of phenotypes[3]. In this example, there are three genes that make reddish pigment in wheat kernels, which we'll call *A*, *B*, and *C*. Each comes in two alleles, one of which makes a unit of pigment (the capital-letter allele) and one of which does not make any pigment (the lowercase allele). Thus, the *aa* genotype would contribute zero units of pigment, the *Aa* genotype would contribute one unit, and the *AA* genotype would contribute two—basically, a form of incomplete dominance.




Now, let's imagine that two plants heterozygous for all three genes (*AaBbCc*) were crossed to one another (or, equivalently, allowed to self-fertilize). Each of the parent plants would have three units of pigment, or pinkish kernels. Their offspring, however, could display seven different categories of phenotypes, ranging from zero units of pigment (*aabbcc*) and pure white kernels to six units of pigment (*AABBCC*) and dark red kernels, with the intermediate phenotypes being most common.

This example illustrates how we can get a spectrum of slightly different phenotypes (something approaching continuous variation) with just three genes whose alleles display incomplete dominance. It's not hard to imagine that, as we increased the number of genes involved, we'd be able to get even finer variations in color, or in another trait such as height. Real polygenic traits aren't usually quite this clean and simple. (For instance, genes may make unequal contributions to the phenotype, alleles may or may not display incomplete dominance, and there may be non-additive interactions between genes.) However, the basic idea—that multiple genes obeying Mendelian rules can produce a spectrum of finely differing phenotypes—holds true for human traits such as skin and eye color.

### PRactice Questions

We've learned about polygenic inheritance and continuous variation. Just what is the difference between these two types of inheritance?

[practice-area rows="4"][/practice-area]
[reveal-answer q="646463"]Show Answer[/reveal-answer]

[hidden-answer a=”646463″]*Polygenic traits* are traits that rely on multiple genes. *Continuous variation* describes traits whose phenotypes occur on a continuum, rather than having a limited number of possible phenotypes. Traits with continuous variation are often also polygenic traits, but not always, and not all polygenic traits have continuous variation.[/hidden-answer]


1. Wood, A. R., Esko, T., Yang, J., Vedantam, S., Pers, T. H., Gustafsson, S., ... Frayling, T. M. (2014). Defining the role of common variation in the genomic and biological architecture of adult human height. *Nature Genetics*, *46*, 1173–1186. http://dx.doi.org/10.1038/ng.3097. ↵
2. White, D. and Rabago-Smith, M. (2011). Genotype-phenotype associations and human eye color. *Journal of Human Genetics*, *56*, 5–7. http://dx.doi.org/10.1038/jhg.2010.126. ↵
3. Kimball, J. W. (2011, March 8). Continuous variation: Quantitative traits. Retrieved from users.rcn.com/jkimball.ma.ultranet/BiologyPages/Q/QTL.html. ↵


## Contributors and Attributions


CC licensed content, Shared previously

- Polygenic inheritance and environmental effects. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/classical-genetics/variations-on-mendelian-genetics/a/polygenic-inheritance-and-environmental-effects. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

14.18: Polygenic Inheritance and Environmental Effects is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.19: Effect of the Environment

Characteristics that are influenced by environmental as well as genetic factors are called *multifactorial*. The idea of “nature versus nurture” — in other words, the relative influence of genetics versus environmental factors — has been and still is debated. Just looking at the genes of a given organism will not determine how that organism will develop and act. Even identical twins will show different characteristics, depending on the environment in which they live. Everyone is a product of their environment as well as their genetics.



Figure 1. Taking a newborn blood sample for PKU testing. By Staff Sgt Eric T. Sheler, U.S. Air Force (Phenylketonuria testing) Public Domain

Even when influenced by the environment, phenotypes have a normal range of expression. For instance, human height varies based on nutrition and genetics, but not many people are shorter than 4½ feet or taller than 7 feet. The range of phenotypic possibilities is called the *norm of reaction*. Hydrangeas, for example, may be blue, pink, or purple, but they are never naturally orange. Hydrangeas are blue in acidic soil with available aluminum, and they are pink in alkaline soil without available aluminum.

Some human characteristics have a narrow norm of reaction, such as blood type. Others have a wide norm of reaction, such as the number of blood cells in humans, which varies depending on factors that include physical fitness, presence or history of infections, and even the altitude at which a person lives.

The environment also affects human genes. Serotonin, a neurotransmitter that acts inside brain cells, lowers anxiety and depression during traumatic times. Mutations in the serotonin transporter gene may cause a reduced ability to cope with stress. That does not mean that the person is always depressed, but if the environment produces stress, the person may become depressed more easily than a person with unmutated serotonin transporter genes.

You already learned about PKU, a pleiotropic disorder caused by defects in a single gene coding for an enzyme that converts the amino acid phenylalanine to tyrosine. Newborns are tested for this defect very early in life (Figure 1), so that if the results are positive, they can be given a diet limiting phenylalanine ingestion. That way, the toxic buildup is prevented and the children can develop normally. PKU is an example in which environmental factors can modify gene expression.

## Practice Question

Two identical twins (female) live in different parts of the country. One is very committed to a healthy lifestyle: not smoking, exercising regularly, eating a diet rich in fresh produce, and avoiding red meats and processed foods. The other is not as careful: she smokes, is overweight, and often eats fast and processed foods. They are aware that several women in their family have had breast cancer, and decide to consult a doctor about their odds of developing the disease. Which of the following statements by the doctor sounds most correct?

1. As identical twins, you are genetically the same, so your chances of developing breast cancer are identical.
2. The twin with the healthy lifestyle should not be terribly concerned, while the one with the unhealthy lifestyle is at a higher risk.
3. Breast cancer has a genetic component, and the twins have identical genes, so they have the same genetic risk. However, environmental factors such as smoking, obesity, and consumption of red meat have been shown to increase the risk of cancer. While both twins should monitor themselves closely, the twin who smokes and is overweight may want to consider a healthier lifestyle to decrease her risk of breast cancer.

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”41921″]Show Answer[/reveal-answer]

[hidden-answer a=”41921″]Option one is wrong; while it has been shown that certain genes may predispose people to cancer, there are many associations between environmental effects and cancer. Option two is also wrong; familial cancers have a genetic component which may or may not be balanced by a healthy lifestyle. Option three is the most correct answer; lifestyle choices are important, but genetic influences are to be taken seriously, especially if there is a family pattern associated with them.[/hidden-answer]

## In Summary: Effect of the Environment

While genes and genetic causes play a large role in health and phenotypes, the environment also plays an important role. Understanding this can enable the treatment of some disorders, such as the case with PKU in which limiting the intake of phenylalanine can prevent toxic build up of this amino acid. Often the norm of reaction is set by genetic factors but ultimately determined by environmental exposures.

## Contributors and Attributions

CC licensed content, Shared previously

- Application Spotlight: Effect of the Environment. **Provided by**: Open Learning Initiative. **Located at**: https://oli.cmu.edu/jcourse/workbook/activity/page?context=434a5f4180020ca600915a031e68a874. **Project**: Introduction to Biology (Open + Free). **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

14.19: Effect of the Environment is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.20: Pleiotropy and Human Disorders


Based on Mendel's experiments, you might imagine that all genes control a single characteristic, are present in two copies, and affect some harmless aspect of an organism's appearance (such as color, height, or shape). Although those predictions are accurate in many cases, there are also some important exceptions. For instance, how can we explain observations like the following?

- The genetic disorder Marfan syndrome is caused by a mutation in one gene, yet it affects many aspects of growth and development, including height, vision, and heart function.

To understand observations like these, we need to look more deeply at what genes are. Rather than abstract "heritable factors," genes are stretches of DNA found on chromosomes, and most of them encode (specify the sequence of) proteins that do a certain job in the cell or body. In this article, we'll look in more detail at genes affecting multiple characteristics (pleiotropy).

## Pleiotropy

When we discussed Mendel's experiments with purple-flowered and white-flowered plants, we didn't mention any other phenotypes associated with the two flower colors. However, Mendel noticed that the flower colors were always correlated with two other features: the color of the seed coat (covering of the seed) and the color of the axils (junctions where the leaves met the main stem)[1]. In plants with white flowers, the seed coats and axils were colorless, while in plants with purple flowers, the seed coats were brown-gray and the axils were reddish. Thus, rather than affecting just one characteristic, the flower color gene actually affected three.




Based on similar diagram by Ingrid Lobo

Genes like this, which affect multiple, seemingly unrelated aspects of an organism's phenotype, are said to be **pleiotropic** (*pleio–* = many, *–tropic* = effects)[2]. We now know that Mendel's flower color gene encodes a regulator protein that activates pigment biosynthesis, and that it works in several different parts of the pea plant (flowers, seed coat, and leaf axils). Thus, the seemingly unrelated phenotypes can all be traced back to a defect in a single gene with several jobs.

Alleles of pleiotropic genes are transmitted in the same way as alleles of genes that affect single traits. In these cases, the difference is that the phenotype contains multiple elements. These elements are specified as a package, and there would be both a dominant and recessive version of this package of traits.

## Pleiotropy in Human Genetic Disorders

Genes affected in human genetic disorders are often pleiotropic. For example, people with the hereditary disorder Marfan syndrome may have a constellation of seemingly unrelated symptoms[3]:

- Unusually tall height
- Thin fingers and toes
- Dislocation of the lens of the eye
- Heart problems (in which the aorta, the large blood vessel carrying blood away from the heart, bulges or ruptures).

These symptoms don't appear directly related to one another, but as it turns out, they can all be traced back to the mutation of a single gene. This gene encodes a protein that assembles into chains, making elastic fibrils that give strength and flexibility to the body's connective tissues[4]. Disease-causing mutations in the Marfan syndrome reduce the amount of functional protein produced,

resulting in fewer fibrils. The eye and the aorta normally contain many fibrils that help maintain structure, explaining why these two organs are strongly affected in Marfan syndrome[5]. In addition, the fibrils serve as “storage shelves” for growth factors. When there are fewer of them in Marfan syndrome, the growth factors cannot be shelved and thus cause excess growth (leading to the characteristic tall, thin Marfan build)[6].


1. Lobo, I. (2008). Pleiotropy: One gene can affect multiple traits. *Nature Education*, *1*(1), 10. Retrieved from http://www.nature.com/scitable/topicpage/pleiotropy-one-gene-can-affect-multiple-traits-569. ↵
2. *Ibid*. ↵
3. Marfan syndrome. (2012). In *Genetics home reference*. Retrieved from http://ghr.nlm.nih.gov/condition/marfan-syndrome. ↵
4. FBN1. (2015). In *Genetics home reference*. Retrieved from http://ghr.nlm.nih.gov/gene/FBN1. ↵
5. Marfan syndrome. (2015, November 3). Retrieved November 21, 2015 from Wikipedia: https://en.Wikipedia.org/wiki/Marfan_syndrome. ↵
6. FBN1. (2015). In *Genetics home reference*. Retrieved from http://ghr.nlm.nih.gov/gene/FBN1. ↵


## Contributors and Attributions


CC licensed content, Shared previously

- Pleiotropy, lethal alleles, and sex linkage. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/classical-genetics/variations-on-mendelian-genetics/a/pleiotropy-lethal-alleles-and-sex-linkage. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

14.20: Pleiotropy and Human Disorders is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.21: Putting It Together- Trait Inheritance

Now that you understand more about genetic inheritance, you can appreciate better the complications associated with predicting traits in children. Unfortunately, these complications apply to inherited disease just as much as they apply to predicting a child's height or eye color.

There are over 6,000 monogenetic disorders and thousands more polygenetic disorders. All of this is in addition to the complications brought about by environmental influences and random genetic mutations or damage. Despite this complexity, we continually learn new things about the genetic root of human diseases. This knowledge can be used not only to predict occurrence, as a genetic counselor does, but also to develop treatments, cures, or preventatives as clinicians and researchers do.

One organization that puts this modern understanding of genetics to use is the Dor Yeshorim, an organization that offers genetic screening to members of the Jewish community worldwide. Its objective is to minimize, and eventually eliminate, the incidence of genetic disorders common to Jewish people, such as Tay–Sachs disease.





Figure 1. Tay-Sachs is an autosomal recessive disease. This means that children can only inherit the Tay-Sachs if both of their parents are carriers for the disease.

In both the Ashkenazi and Sephardi Jewish communities, there is an increased rate of a number of genetic disorders such as Tay–Sachs disease, an autosomal recessive disorder that goes unnoticed in carriers, but is fatal within the first few years of life in almost all homozygotes.

Dor Yeshorim screens only for recessive traits that give rise to lethal or severely debilitating disorders, providing preventative, rather than diagnostic services. They do not screen for disorders arising from dominant gene mutations, as these cannot be prevented by informed mate selection. Orthodox Judaism generally opposes selective abortion, and although preimplantation genetic diagnosis (PGD) is often approved by Halakha, it is a difficult and costly process. By avoiding marriages between carriers of the diseases, the incidence of the disorders decreases without having to resort to these methods.

Dor Yeshorim advocates anonymous testing. Individuals are tested during large sessions in Jewish schools and processed anonymously with only a PIN linking the sample with the candidate. When two members of the system contemplate marriage, they contact the organization and enter both their PINs. When both carry a gene for the same disorder, the risk of affected offspring is 25 percent, and it is considered advisable to discontinue the plans. In the context of shidduchim (a system of matchmaking in Jewish communities), the "carriership check" is often run within the first three dates, to avoid disappointments and heartbreak.

According to the Dor Yeshorim website, they are "credited with singlehandedly eradicating Tay Sachs from the Jewish community and effectively closing the Tay Sachs ward at Kingsbrook Medical center forever."[1]

1. "Our History", *Dor Yeshorim*, https://doryeshorim.org/history-achievements/. ↵

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: Trait Inheritance. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Modification of Dor Yeshorim. **Provided by**: Wikipedia. **Located at**: https://en.Wikipedia.org/wiki/Dor_Yeshorim. **License**: *CC BY-SA: Attribution-ShareAlike*

Public domain content

- Autorecessive. **Provided by**: National Institutes of Health. **Located at**: https://commons.wikimedia.org/wiki/File:Autorecessive.jpg. **License**: *Public Domain: No Known Copyright*

---

14.21: Putting It Together- Trait Inheritance is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 14.22: Assignment- Dragon Genetics

Begin this assessment by downloading the Modern Genetics simulator (links to an external site) from Concord.org.

Run the simulation and complete the exercises in it. While doing so, gather answers to these questions. Take screenshots as directed; save each screenshot as its own document. You will then complete the quiz entitled “C10: Dragon Genetics” to submit your answers and upload your documents. (Download the quiz file here.)

Please note the following:

- The questions in the document are broken down by parts in the simulator; this is to help you as you move through the exercise. They are not separated this way in the quiz but they will follow the same order.
- When performing your pedigree crosses, if you obtain offspring of a single sex, simply repeat the exact same parental cross to get more offspring. Do this until you have at least one of each sex in that generation.

Here is a worked example showing a screenshot and required text explanation.

## Contributors and Attributions

CC licensed content, Original

- Performance Assessment: Dragon Genetics. **Authored by**: Shelli Carter. **Provided by**: Columbia Basin College. **Located at**: https://www.columbiabasin.edu/. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Modern Genetics simulator. **Provided by**: Concord Consortium. **Located at**: http://concord.org/stem-resources/modern-genetics. **License**: *Other*. **License Terms**: GNU LGPL

14.22: Assignment- Dragon Genetics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 15: Module 13- Theory of Evolution


15.1: Why It Matters- Theory of Evolution
15.2: Introduction to Charles Darwin
15.3: Darwin and Descent with Modification
15.4: Darwin and the Theory of Evolution
15.5: Natural Selection
15.6: Introduction to Evidence for Evolution
15.7: Physical Evidence
15.8: Biological Evidence
15.9: Misconceptions of Evolution
15.10: Introduction to Mutations and Evolution
15.11: Population Genetics
15.12: Selective and Environmental Pressures
15.13: Introduction to Phylogenetic Trees
15.14: Scientific Classification
15.15: Structure of Phylogenetic Trees
15.16: Limitations of Phylogenetic Trees
15.17: The Taxonomic Classification System
15.18: Putting It Together- Theory of Evolution



15: Module 13- Theory of Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.1: Why It Matters- Theory of Evolution

## Why explain the theory of evolution, which documents the change in the genetic makeup of a biological population over time?

Have you ever wondered why doctors say you should get a new flu shot every year? It's just the same old flu virus right? How much could it change from year to year?

Unfortunately, it's not just the same old flu virus—and it often changes a lot from year to year. When the flu vaccine does its work, killing the current iteration of the flu, it typically does a good job and saves its host from illness. However, as it wipes out the majority of flu viruses, the only ones that are left are those that are resistant to the current virus, which means the overall make up of the population of flu viruses has changed: the population has evolved to become resistant to one anti-viral treatment, so another must be found.

Of course, while the theory of evolution can be applied to viruses, it is more often discussed in the context of living things: bacteria, plants, animals, and even humans. The theory of evolution began as a revolutionary idea, but it remains central to the study of biology. Evolution is the unifying concept in biology. This theory documents the change in the genetic makeup of a biological population over time. Evolution helps us understand the development of antibiotic resistance in bacteria and other parasitic organisms. The following are just a few of the antibiotic resistant "bugs" plaguing humans.

- Antibiotic-Resistant Mycobacterium tuberculosis (TB)
- Methicillin-Resistant Staphylococcus aureus (MRSA)
- Vancomycin-Resistant Enterococci (VRE)
- Multidrug-Resistant Neisseria gonorrhoeae (Gonorrhea)

**Try It**

You can also check out the NIH's website about antimicrobial resistance for more information.

While evolution is easiest to see in bacteria due to their short life cycles, every living population experiences evolution of one kind or another. Let's see just how the world we live in has guided the evolutionary process.

### Learning Outcomes

- Describe the work of Charles Darwin in the Galapagos Islands, especially his discovery of natural selection in finch populations
- Describe how the theory of evolution by natural selection is supported by evidence
- Recognize that mutations are the basis of microevolution; and that adaptations enhance the survival and reproduction of individuals in a population
- Read and analyze a phylogenetic tree that documents evolutionary relationships

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

15.1: Why It Matters- Theory of Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.2: Introduction to Charles Darwin

## Describe the work of Charles Darwin in the Galapagos Islands, especially his discovery of natural selection in finch populations

Charles Darwin is best known for his discovery of natural selection. In this outcome we'll learn more about his work and how it helped develop the theory of evolution.

## What You'll Learn to Do

- Outline the work of Charles Darwin as a naturalist aboard the HMS Beagle
- Summarize the prior work and new evidence Darwin used to develop the idea of "descent with modification"
- Describe how Darwin's work developed to the theory of evolution
- Define the concept of natural selection

### Learning Activities

The learning activities for this section include the following:

- Darwin and Descent with Modification
- Darwin and the Theory of Evolution
- Natural Selection
- Self Check: Charles Darwin

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

15.2: Introduction to Charles Darwin is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.3: Darwin and Descent with Modification





Figure 1. Darwin observed that beak shape varies among finch species. He postulated that the beak of an ancestral species had adapted over time to equip the finches to acquire different food sources.

In the mid-nineteenth century, the actual mechanism for evolution was independently conceived of and described by two naturalists: Charles Darwin and Alfred Russel Wallace. Importantly, each naturalist spent time exploring the natural world on expeditions to the tropics. From 1831 to 1836, Darwin traveled around the world on *H.M.S. Beagle*, including stops in South America, Australia, and the southern tip of Africa. Wallace traveled to Brazil to collect insects in the Amazon rainforest from 1848 to 1852 and to the Malay Archipelago from 1854 to 1862. Darwin's journey, like Wallace's later journeys to the Malay Archipelago, included stops at several island chains, the last being the Galápagos Islands west of Ecuador. On these islands, Darwin observed species of organisms on different islands that were clearly similar, yet had distinct differences. For example, the ground finches inhabiting the Galápagos Islands comprised several species with a unique beak shape (Figure 1).

The species on the islands had a graded series of beak sizes and shapes with very small differences between the most similar. He observed that these finches closely resembled another finch species on the mainland of South America. Darwin imagined that the island species might be species modified from one of the original mainland species. Upon further study, he realized that the varied beaks of each finch helped the birds acquire a specific type of food. For example, seed-eating finches had stronger, thicker beaks for breaking seeds, and insect-eating finches had spear-like beaks for stabbing their prey.


Wallace and Darwin both observed similar patterns in other organisms and they independently developed the same explanation for how and why such changes could take place. Darwin called this mechanism natural selection. **Natural selection**, also known as "survival of the fittest," is the more prolific reproduction of individuals with favorable traits that survive environmental change because of those traits; this leads to evolutionary change.

For example, a population of giant tortoises found in the Galapagos Archipelago was observed by Darwin to have longer necks than those that lived on other islands with dry lowlands. These tortoises were "selected" because they could reach more leaves and access more food than those with short necks. In times of drought when fewer leaves would be available, those that could reach more leaves had a better chance to eat and survive than those that couldn't reach the food source. Consequently, long-necked tortoises would be more likely to be reproductively successful and pass the long-necked trait to their offspring. Over time, only long-necked tortoises would be present in the population.

Natural selection, Darwin argued, was an inevitable outcome of three principles that operated in nature. First, most characteristics of organisms are inherited, or passed from parent to offspring. Although no one, including Darwin and Wallace, knew how this happened at the time, it was a common understanding. Second, more offspring are produced than are able to survive, so resources for survival and reproduction are limited. The capacity for reproduction in all organisms outstrips the availability of resources to support their numbers. Thus, there is competition for those resources in each generation. Both Darwin and Wallace's understanding of this principle came from reading an essay by the economist Thomas Malthus who discussed this principle in relation to human populations. Third, offspring vary among each other in regard to their characteristics and those variations are inherited. Darwin and Wallace reasoned that offspring with inherited characteristics which allow them to best compete for limited resources will survive and have more offspring than those individuals with variations that are less able to compete. Because characteristics are inherited,

these traits will be better represented in the next generation. This will lead to change in populations over generations in a process that Darwin called descent with modification. Ultimately, natural selection leads to greater adaptation of the population to its local environment; it is the only mechanism known for adaptive evolution.

Papers by Darwin and Wallace (Figure 2) presenting the idea of natural selection were read together in 1858 before the Linnean Society in London. The following year Darwin's book, *On the Origin of Species*, was published. His book outlined in considerable detail his arguments for gradual changes and adaptive survival by natural selection.



(a) (b)

Figure 2. Both (a) Charles Darwin and (b) Alfred Wallace wrote scientific papers on natural selection that were presented together before the Linnean Society in 1858.

Demonstrations of evolution by natural selection are time consuming and difficult to obtain. One of the best examples has been demonstrated in the very birds that helped to inspire Darwin's theory: the Galápagos finches. Peter and Rosemary Grant and their colleagues have studied Galápagos finch populations every year since 1976 and have provided important demonstrations of natural selection. The Grants found changes from one generation to the next in the distribution of beak shapes with the medium ground finch on the Galápagos island of Daphne Major. The birds have inherited variation in the bill shape with some birds having wide deep bills and others having thinner bills. During a period in which rainfall was higher than normal because of an El Niño, the large hard seeds that large-billed birds ate were reduced in number; however, there was an abundance of the small soft seeds which the small-billed birds ate. Therefore, survival and reproduction were much better in the following years for the small-billed birds. In the years following this El Niño, the Grants measured beak sizes in the population and found that the average bill size was smaller. Since bill size is an inherited trait, parents with smaller bills had more offspring and the size of bills had evolved to be smaller. As conditions improved in 1987 and larger seeds became more available, the trend toward smaller average bill size ceased.

## In Summary: Darwin and Descent with Modification

While Charles Darwin is generally called "the father of evolution," the basic idea for this concept was actually developed by both Darwin and Alfred Russel Wallace. Both scientists based their hypotheses on observations of diversity among natural populations. Darwin's work in particular focused on animals of the Galapagos islands, especially finches. Over time, the idea that species changed from natural selection pressures through "descent with modification" gave rise to the idea of evolution. Data accumulated over time, for example the long study of the Galapagos finches by the Grant research team, has supported this idea and moved it into the realm of a supported theory of biology.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

---

15.3: Darwin and Descent with Modification is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.4: Darwin and the Theory of Evolution

Natural selection can only take place if there is **variation**, or differences, among individuals in a population. Importantly, these differences must have some genetic basis; otherwise, the selection will not lead to change in the next generation. This is critical because variation among individuals can be caused by non-genetic reasons such as an individual being taller because of better nutrition rather than different genes.

Genetic diversity in a population comes from two main mechanisms: mutation and sexual reproduction. Mutation, a change in DNA, is the ultimate source of new alleles, or new genetic variation in any population. The genetic changes caused by mutation can have one of three outcomes on the phenotype. A mutation affects the phenotype of the organism in a way that gives it reduced fitness—lower likelihood of survival or fewer offspring. A mutation may produce a phenotype with a beneficial effect on fitness. And, many mutations will also have no effect on the fitness of the phenotype; these are called neutral mutations. Mutations may also have a whole range of effect sizes on the fitness of the organism that expresses them in their phenotype, from a small effect to a great effect. Sexual reproduction also leads to genetic diversity: when two parents reproduce, unique combinations of alleles assemble to produce the unique genotypes and thus phenotypes in each of the offspring.

A heritable trait that helps the survival and reproduction of an organism in its present environment is called an **adaptation**. Scientists describe groups of organisms becoming adapted to their environment when a change in the range of genetic variation occurs over time that increases or maintains the “fit” of the population to its environment. The webbed feet of platypuses are an adaptation for swimming. The snow leopards’ thick fur is an adaptation for living in the cold. The cheetahs’ fast speed is an adaptation for catching prey.

Whether or not a trait is favorable depends on the environmental conditions at the time. The same traits are not always selected because environmental conditions can change. For example, consider a species of plant that grew in a moist climate and did not need to conserve water. Large leaves were selected because they allowed the plant to obtain more energy from the sun. Large leaves require more water to maintain than small leaves, and the moist environment provided favorable conditions to support large leaves. After thousands of years, the climate changed, and the area no longer had excess water. The direction of natural selection shifted so that plants with small leaves were selected because those populations were able to conserve water to survive the new environmental conditions.

The evolution of species has resulted in enormous variation in form and function. Sometimes, evolution gives rise to groups of organisms that become tremendously different from each other. When two species evolve in diverse directions from a common point, it is called divergent evolution. Such **divergent evolution** can be seen in the forms of the reproductive organs of flowering plants which share the same basic anatomies; however, they can look very different as a result of selection in different physical environments and adaptation to different kinds of pollinators (Figure 1).




Figure 1. Flowering plants evolved from a common ancestor. Notice that the (a) dense blazing star (*Liatrus spicata*) and the (b) purple coneflower (*Echinacea purpurea*) vary in appearance, yet both share a similar basic morphology. (credit a: modification of work by Drew Avery; credit b: modification of work by Cory Zanker)

In other cases, similar phenotypes evolve independently in distantly related species. For example, flight has evolved in both bats and insects, and they both have structures we refer to as wings, which are adaptations to flight. However, the wings of bats and insects have evolved from very different original structures. This phenomenon is called **convergent evolution**, where similar traits evolve independently in species that do not share a recent common ancestry. The two species came to the same function, flying, but did so separately from each other.

These physical changes occur over enormous spans of time and help explain how evolution occurs. Natural selection acts on individual organisms, which in turn can shape an entire species. Although natural selection may work in a single generation on an individual, it can take thousands or even millions of years for the genotype of an entire species to evolve. It is over these large time spans that life on earth has changed and continues to change.

## In Summary: Darwin and the Theory of Evolution

Natural selection, the driving force behind evolution, can only work if variation exists among organisms. Variation arises ultimately from genetic mutations. Diversity is further encouraged through sexual reproduction. As environments change, selective pressures shift and favor different adaptations. In this way, given thousands or millions of years, species evolve.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.4: Darwin and the Theory of Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.5: Natural Selection

This video reviews some what we've learned so far this outcome, as well as going into more depth about natural selection and how it works.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=530

## Contributors and Attributions

All rights reserved content

- Natural Selection - Crash Course Biology #14. **Authored by**: CrashCourse. **Located at**: https://youtu.be/aTftyFboC_M. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

15.5: Natural Selection is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.6: Introduction to Evidence for Evolution

## Describe how the theory of evolution by natural selection is supported by evidence

The evidence for evolution is compelling and extensive. Looking at every level of organization in living systems, biologists see the signature of past and present evolution. Darwin dedicated a large portion of his book, *On theOrigin of Species*, to identifying patterns in nature that were consistent with evolution, and since Darwin, our understanding has become clearer and broader.

## What You'll Learn to Do

- Outline physical evidence that supports the theory of evolution
- Outline biological evidence that supports the theory of evolution
- Refute common misconceptions about evolution

### Learning Activities

The learning activities for this section include the following:

- Physical Evidence
- Biological Evidence
- Misconceptions of Evolution
- Self Check: Evidence for Evolution

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

15.6: Introduction to Evidence for Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.7: Physical Evidence

## Fossils

Fossils provide solid evidence that organisms from the past are not the same as those found today, and fossils show a progression of evolution. Scientists determine the age of fossils and categorize them from all over the world to determine when the organisms lived relative to each other. The resulting fossil record tells the story of the past and shows the evolution of form over millions of years (Figure 1a). For example, scientists have recovered highly detailed records showing the evolution of humans and horses (Figure 1b).




(a) (b)

Figure 1. In this (a) display, fossil hominids are arranged from oldest (bottom) to newest (top). As hominids evolved, the shape of the skull changed. An artist's rendition of (b) extinct species of the genus Equus reveals that these ancient species resembled the modern horse (Equus ferus) but varied in size.

## Anatomy and Embryology





Figure 2. The similar construction of these appendages indicates that these organisms share a common ancestor.

Another type of evidence for evolution is the presence of structures in organisms that share the same basic form. For example, the bones in the appendages of a human, dog, bird, and whale all share the same overall construction (Figure 2) resulting from their origin in the appendages of a common ancestor. Over time, evolution led to changes in the shapes and sizes of these bones in different species, but they have maintained the same overall layout. Scientists call these synonymous parts homologous structures.

Some structures exist in organisms that have no apparent function at all, and appear to be residual parts from a past common ancestor. These unused structures without function are called vestigial structures. Some examples of vestigial structures are wings on flightless birds, leaves on some cacti, and hind leg bones in whales.

Visit this interactive site to guess which bones structures are homologous and which are analogous, and see examples of evolutionary adaptations to illustrate these concepts.

Another evidence of evolution is the convergence of form in organisms that share similar environments. For example, species of unrelated animals, such as the arctic fox and ptarmigan, living in the arctic region have been selected for seasonal white phenotypes during winter to blend with the snow and ice (Figure 3). These similarities occur not because of common ancestry, but because of similar selection pressures—the benefits of not being seen by predators.



(a) (b)

Figure 3. The white winter coat of the (a) arctic fox and the (b) ptarmigan's plumage are adaptations to their environments. (credit a: modification of work by Keith Morehouse)

Embryology, the study of the development of the anatomy of an organism to its adult form, also provides evidence of relatedness between now widely divergent groups of organisms. Mutational tweaking in the embryo can have such magnified consequences in the adult that embryo formation tends to be conserved. As a result, structures that are absent in some groups often appear in their embryonic forms and disappear by the time the adult or juvenile form is reached. For example, all vertebrate embryos, including humans, exhibit gill slits and tails at some point in their early development. These disappear in the adults of terrestrial groups but are maintained in adult forms of aquatic groups such as fish and some amphibians. Great ape embryos, including humans, have a tail structure during their development that is lost by the time of birth.

## Learning Objectives

Since Darwin developed his ideas on descent with modification and the pressures of natural selection, a variety of evidence has been gathered supporting the theory of evolution. Fossil evidence shows the changes in lineages over millions of years, such as in hominids and horses. Studying anatomy allows scientists to identify homologous structures across diverse groups of related organisms, such as leg bones. Vestigial structures also offer clues to common ancestors. Using embryology, scientists can identify common ancestors through structures present only during development and not in the adult form.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.7: Physical Evidence is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.8: Biological Evidence

## Biogeography

The geographic distribution of organisms on the planet follows patterns that are best explained by evolution in conjunction with the movement of tectonic plates over geological time. Broad groups that evolved before the breakup of the supercontinent Pangaea (about 200 million years ago) are distributed worldwide. Groups that evolved since the breakup appear uniquely in regions of the planet, such as the unique flora and fauna of northern continents that formed from the supercontinent Laurasia and of the southern continents that formed from the supercontinent Gondwana. The presence of members of the plant family Proteaceae in Australia, southern Africa, and South America is best due to their appearance prior to the southern supercontinent Gondwana breaking up.

The great diversification of marsupials in Australia and the absence of other mammals reflect Australia's long isolation. Australia has an abundance of endemic species—species found nowhere else—which is typical of islands whose isolation by expanses of water prevents species migration. Over time, these species diverge evolutionarily into new species that look very different from their ancestors that may exist on the mainland. The marsupials of Australia, the finches on the Galápagos, and many species on the Hawaiian Islands are all unique to their one point of origin, yet they display distant relationships to ancestral species on mainlands.

## Molecular Biology

Like anatomical structures, the structures of the molecules of life reflect descent with modification. Evidence of a common ancestor for all of life is reflected in the universality of DNA as the genetic material and in the near universality of the genetic code and the machinery of DNA replication and expression. Fundamental divisions in life between the three domains are reflected in major structural differences in otherwise conservative structures such as the components of ribosomes and the structures of membranes. In general, the relatedness of groups of organisms is reflected in the similarity of their DNA sequences—exactly the pattern that would be expected from descent and diversification from a common ancestor.

DNA sequences have also shed light on some of the mechanisms of evolution. For example, it is clear that the evolution of new functions for proteins commonly occurs after gene duplication events that allow the free modification of one copy by mutation, selection, or drift (changes in a population's gene pool resulting from chance), while the other copy continues to produce a functional protein.

**Learning Objectives**

Biogeography offers further clues about evolutionary relationships. The presence of related organisms across continents indicates when these organisms may have evolved. For example, some flora and fauna of the northern continents are similar across these landmasses but distinct from that of the southern continents. Islands such as Australia and the Galapagos chain often have unique species that evolved after these landmasses separated from the mainland. Finally, molecular biology provides data supporting the theory of evolution. In particular, the universality of DNA and near universality of the genetic code for proteins shows that all life once shared a common ancestor. DNA also provides clues into how evolution may have happened. Gene duplications allow one copy to undergo mutational events without harming an organism, as one copy continues to produce functional protein.

## Evolution—It's a Thing

This video defines evolution and discusses several varieties of evidence that support the Theory of Evolution:



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=536

## Contributors and Attributions

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Evolution: It's a Thing - Crash Course Biology #20. **Authored by**: CrashCourse. **Located at**: https://youtu.be/P3GagfbA2vo. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

15.8: Biological Evidence is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.9: Misconceptions of Evolution

Although the theory of evolution generated some controversy when it was first proposed, it was almost universally accepted by biologists, particularly younger biologists, within 20 years after publication of *On theOrigin of Species*. Nevertheless, the theory of evolution is a difficult concept and misconceptions about how it works abound.

This site addresses some of the main misconceptions associated with the theory of evolution.

## Evolution Is Just a Theory

Critics of the theory of evolution dismiss its importance by purposefully confounding the everyday usage of the word "theory" with the way scientists use the word. In science, a "theory" is understood to be a body of thoroughly tested and verified explanations for a set of observations of the natural world. Scientists have a theory of the atom, a theory of gravity, and the theory of relativity, each of which describes understood facts about the world. In the same way, the theory of evolution describes facts about the living world. As such, a theory in science has survived significant efforts to discredit it by scientists. In contrast, a "theory" in common vernacular is a word meaning a guess or suggested explanation; this meaning is more akin to the scientific concept of "hypothesis." When critics of evolution say evolution is "just a theory," they are implying that there is little evidence supporting it and that it is still in the process of being rigorously tested. This is a mischaracterization.

## Individuals Evolve

Evolution is the change in genetic composition of a population over time, specifically over generations, resulting from differential reproduction of individuals with certain alleles. Individuals do change over their lifetime, obviously, but this is called development and involves changes programmed by the set of genes the individual acquired at birth in coordination with the individual's environment. When thinking about the evolution of a characteristic, it is probably best to think about the change of the average value of the characteristic in the population over time. For example, when natural selection leads to bill-size change in medium-ground finches in the Galápagos, this does not mean that individual bills on the finches are changing. If one measures the average bill size among all individuals in the population at one time and then measures the average bill size in the population several years later, this average value will be different as a result of evolution. Although some individuals may survive from the first time to the second, they will still have the same bill size; however, there will be many new individuals that contribute to the shift in average bill size.

## Organisms Evolve on Purpose

Statements such as "organisms evolve in response to a change in an environment" are quite common, but such statements can lead to two types of misunderstandings. First, the statement must not be understood to mean that individual organisms evolve. The statement is shorthand for "a population evolves in response to a changing environment." However, a second misunderstanding may arise by interpreting the statement to mean that the evolution is somehow intentional. A changed environment results in some individuals in the population, those with particular phenotypes, benefiting and therefore producing proportionately more offspring than other phenotypes. This results in change in the population if the characteristics are genetically determined.

It is also important to understand that the variation that natural selection works on is already in a population and does not arise in response to an environmental change. For example, applying antibiotics to a population of bacteria will, over time, select a population of bacteria that are resistant to antibiotics. The resistance, which is caused by a gene, did not arise by mutation because of the application of the antibiotic. The gene for resistance was already present in the gene pool of the bacteria, likely at a low frequency. The antibiotic, which kills the bacterial cells without the resistance gene, strongly selects individuals that are resistant, since these would be the only ones that survived and divided. Experiments have demonstrated that mutations for antibiotic resistance do not arise as a result of antibiotic.

In a larger sense, evolution is not goal directed. Species do not become "better" over time; they simply track their changing environment with adaptations that maximize their reproduction in a particular environment at a particular time. Evolution has no goal of making faster, bigger, more complex, or even smarter species, despite the commonness of this kind of language in popular discourse. What characteristics evolve in a species are a function of the variation present and the environment, both of which are

constantly changing in a non-directional way. What trait is fit in one environment at one time may well be fatal at some point in the future. This holds equally well for a species of insect as it does the human species.

## Evolution Explains the Origin of Life

It is a common misunderstanding that evolution includes an explanation of life's origins. Conversely, some of the theory's critics believe that it cannot explain the origin of life. The theory does not try to explain the origin of life. The theory of evolution explains how populations change over time and how life diversifies the origin of species. It does not shed light on the beginnings of life including the origins of the first cells, which is how life is defined. The mechanisms of the origin of life on Earth are a particularly difficult problem because it occurred a very long time ago, and presumably it just occurred once. Importantly, biologists believe that the presence of life on Earth precludes the possibility that the events that led to life on Earth can be repeated because the intermediate stages would immediately become food for existing living things.

However, once a mechanism of inheritance was in place in the form of a molecule like DNA either within a cell or pre-cell, these entities would be subject to the principle of natural selection. More effective reproducers would increase in frequency at the expense of inefficient reproducers. So while evolution does not explain the origin of life, it may have something to say about some of the processes operating once pre-living entities acquired certain properties.

### Learning Objectives

Many misconceptions exist about the theory of evolution—including some perpetuated by critics of the theory. First, evolution as a scientific theory means that it has years of observation and accumulated data supporting it. It is not "just a theory" as a person may say in common vernacular.

Another misconception is that individuals evolve, though in fact it is populations that evolve over time. Individuals simply carry mutations. Furthermore, these mutations neither arise on purpose nor do they arise in response to an environmental pressure. Instead, mutations in DNA happen spontaneously and are already present in individuals of a population when a selective pressure occurs. Once the environment begins to favor a particular trait, then those individuals already carrying that mutation will have a selective advantage and are likely to survive better and outproduce others without the adaptation.

Finally, the theory of evolution does not in fact address the origins of life on this planet. Scientists believe that we cannot, in fact, repeat the circumstances that led to life on Earth because at this time life already exists. The presence of life has so dramatically changed the environment that the origins cannot be totally produced for study.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.9: Misconceptions of Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.10: Introduction to Mutations and Evolution

## Recognize that mutations are the basis of microevolution; and that adaptations enhance the survival and reproduction of individuals in a population

We've already learned about DNA and mutations. In this outcome, we'll understand how these mutations impact evolution and the change of species.

## What You'll Learn to Do

- Understand the connection between genetics and evolution
- Understand how environmental changes and selective pressures impact the spread of mutations, contributing to the process of evolution

### Learning Activities

The learning activities for this section include the following:

- Population Genetics
- Selective and Environmental Pressures
- Self Check: Mutations and Evolution

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

15.10: Introduction to Mutations and Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.11: Population Genetics

## Darwin Meets Mendel—Not Literally

When Darwin came up with his theories of evolution and natural selection, he knew that the processes he was describing depended on heritable variation in populations. That is, they relied on differences in the features of the organisms in a population and on the ability of these different features to be passed on to offspring.

Darwin did not, however, know how traits were inherited. Like other scientists of his time, he thought that traits were passed on via blending inheritance. In this model, parents' traits are supposed to permanently blend in their offspring. The blending model was disproven by Austrian monk Gregor Mendel, who found that traits are specified by non-blending heritable units called genes.

Although Mendel published his work on genetics just a few years after Darwin published his ideas on evolution, Darwin probably never read Mendel's work. Today, we can combine Darwin's and Mendel's ideas to arrive at a clearer understanding of what evolution is and how it takes place.

## Microevolution and Population Genetics

**Microevolution**, or evolution on a small scale, is defined as a change in the frequency of gene variants, alleles, in a population over generations. The field of biology that studies allele frequencies in populations and how they change over time is called **population genetics**.

Microevolution is sometimes contrasted with **macroevolution**, evolution that involves large changes, such as formation of new groups or species, and happens over long time periods. However, most biologists view microevolution and macroevolution as the same process happening on different timescales. Microevolution adds up gradually, over long periods of time to produce macroevolutionary changes.

Let's look at three concepts that are core to the definition of microevolution: populations, alleles, and allele frequency.

### Populations

A **population** is a group of organisms of the same species that are found in the same area and can interbreed. A population is the smallest unit that can evolve—in other words, an individual can't evolve.

### Alleles

An **allele** is a version of a **gene**, a heritable unit that controls a particular feature of an organism.

For instance, Mendel studied a gene that controls flower color in pea plants. This gene comes in a white allele, *w*, and a purple allele, *W*. Each pea plant has two gene copies, which may be the same or different alleles. When the alleles are different, one—the **dominant** allele, *W*—may hide the other—the **recessive allele**, *w*. A plant's set of alleles, called its **genotype**, determines its **phenotype**, or observable features, in this case flower color.




### Allele Frequency

**Allele frequency** refers to how frequently a particular allele appears in a population. For instance, if all the alleles in a population of pea plants were purple alleles, *W*, the allele frequency of *W* would be 100%, or 1.0. However, if half the alleles were *W* and half were *w*, each allele would have an allele frequency of 50%, or 0.5.

In general, we can define allele frequency as


$$\text{Frequency of allele } A=\frac{\text{Number of copies of allele } A\text{ in population}}{\text{Total number of } A\text{/}a\text{ gene copies in population}}$$

Sometimes there are more than two alleles in a population (e.g., there might be *A*, *a*, and $A_i$ alleles of a gene). In that case, you would want to add up *all* of the different alleles to get your denominator.

It's also possible to calculate **genotype frequencies**—the fraction of individuals with a given genotype—and **phenotype frequencies**—the fraction of individuals with a given phenotype. Keep in mind, though, that these are different concepts from allele frequency. We'll see an example of this difference next.

## Learning Objectives

This video talks about population genetics, which helps to explain the evolution of populations over time.




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=542

## Contributors and Attributions

CC licensed content, Shared previously

- Allele frequency and the gene pool. **Provided by**: Khan Academy. **Located at**: https://www.khanacademy.org/science/biology/her/heredity-and-genetics/a/allele-frequency-the-gene-pool. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

All rights reserved content

- Population Genetics: When Darwin Met Mendel - Crash Course Biology #18. **Authored by**: CrashCourse. **Located at**: https://youtu.be/WhFKPaRnTdQ. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

15.11: Population Genetics is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.12: Selective and Environmental Pressures

Natural selection only acts on the population's heritable traits: selecting for beneficial alleles and thus increasing their frequency in the population, while selecting against deleterious alleles and thereby decreasing their frequency—a process known as **adaptive evolution**. Natural selection does not act on individual alleles, however, but on entire organisms. An individual may carry a very beneficial genotype with a resulting phenotype that, for example, increases the ability to reproduce (fecundity), but if that same individual also carries an allele that results in a fatal childhood disease, that fecundity phenotype will not be passed on to the next generation because the individual will not live to reach reproductive age. Natural selection acts at the level of the individual; it selects for individuals with greater contributions to the gene pool of the next generation, known as an organism's **evolutionary (Darwinian) fitness**.

Fitness is often quantifiable and is measured by scientists in the field. However, it is not the absolute fitness of an individual that counts, but rather how it compares to the other organisms in the population. This concept, called **relative fitness**, allows researchers to determine which individuals are contributing additional offspring to the next generation, and thus, how the population might evolve.

There are several ways selection can affect population variation: stabilizing selection, directional selection, diversifying selection, frequency-dependent selection, and sexual selection. As natural selection influences the allele frequencies in a population, individuals can either become more or less genetically similar and the phenotypes displayed can become more similar or more disparate.

## Stabilizing Selection

If natural selection favors an average phenotype, selecting against extreme variation, the population will undergo **stabilizing selection** (Figure 1a). In a population of mice that live in the woods, for example, natural selection is likely to favor individuals that best blend in with the forest floor and are less likely to be spotted by predators. Assuming the ground is a fairly consistent shade of brown, those mice whose fur is most closely matched to that color will be most likely to survive and reproduce, passing on their genes for their brown coat. Mice that carry alleles that make them a bit lighter or a bit darker will stand out against the ground and be more likely to fall victim to predation. As a result of this selection, the population's genetic variance will decrease.

## Directional Selection

When the environment changes, populations will often undergo **directional selection** (Figure 1b), which selects for phenotypes at one end of the spectrum of existing variation. A classic example of this type of selection is the evolution of the peppered moth in eighteenth- and nineteenth-century England. Prior to the Industrial Revolution, the moths were predominately light in color, which allowed them to blend in with the light-colored trees and lichens in their environment. But as soot began spewing from factories, the trees became darkened, and the light-colored moths became easier for predatory birds to spot. Over time, the frequency of the melanic form of the moth increased because they had a higher survival rate in habitats affected by air pollution because their darker coloration blended with the sooty trees. Similarly, the hypothetical mouse population may evolve to take on a different coloration if something were to cause the forest floor where they live to change color. The result of this type of selection is a shift in the population's genetic variance toward the new, fit phenotype.

In science, sometimes things are believed to be true, and then new information comes to light that changes our understanding. The story of the peppered moth is an example: the facts behind the selection toward darker moths have recently been called into question. Read this article to learn more.

## Diversifying Selection

Sometimes two or more distinct phenotypes can each have their advantages and be selected for by natural selection, while the intermediate phenotypes are, on average, less fit. Known as diversifying selection (Figure 1c), this is seen in many populations of animals that have multiple male forms. Large, dominant alpha males obtain mates by brute force, while small males can sneak in for furtive copulations with the females in an alpha male's territory. In this case, both the alpha males and the "sneaking" males will be selected for, but medium-sized males, which can't overtake the alpha males and are too big to sneak copulations, are selected against. Diversifying selection can also occur when environmental changes favor individuals on either end of the phenotypic spectrum. Imagine a population of mice living at the beach where there is light-colored sand interspersed with patches of tall grass.

In this scenario, light-colored mice that blend in with the sand would be favored, as well as dark-colored mice that can hide in the grass. Medium-colored mice, on the other hand, would not blend in with either the grass or the sand, and would thus be more likely to be eaten by predators. The result of this type of selection is increased genetic variance as the population becomes more diverse.

Practice Question





Figure 1. Different types of natural selection can impact the distribution of phenotypes within a population. In (a) stabilizing selection, an average phenotype is favored. In (b) directional selection, a change in the environment shifts the spectrum of phenotypes observed. In (c) diversifying selection, two or more extreme phenotypes are selected for, while the average phenotype is selected against.

In recent years, factories have become cleaner, and less soot is released into the environment. What impact do you think this has had on the distribution of moth color in the population?

[practice-area rows="2"][/practice-area]
[reveal-answer q="173318"]**Show Answer**[/reveal-answer]
[hidden-answer a="173318"]Moths have shifted to a lighter color.[/hidden-answer]

## Frequency-dependent Selection



Figure 2. A yellow-throated side-blotched lizard is smaller than either the blue-throated or orange-throated males and appears a bit like the females of the species, allowing it to sneak copulations. (credit: “tinyfroglet”/Flickr)

Another type of selection, called **frequency-dependent selection**, favors phenotypes that are either common (positive frequency-dependent selection) or rare (negative frequency-dependent selection). An interesting example of this type of selection is seen in a unique group of lizards of the Pacific Northwest. Male common side-blotched lizards come in three throat-color patterns: orange, blue, and yellow. Each of these forms has a different reproductive strategy: orange males are the strongest and can fight other males for access to their females; blue males are medium-sized and form strong pair bonds with their mates; and yellow males (Figure 2) are the smallest, and look a bit like females, which allows them to sneak copulations. Like a game of rock-paper-scissors, orange beats blue, blue beats yellow, and yellow beats orange in the competition for females. That is, the big, strong orange males can fight off the blue males to mate with the blue’s pair-bonded females, the blue males are successful at guarding their mates against yellow sneaker males, and the yellow males can sneak copulations from the potential mates of the large, polygynous orange males.

In this scenario, orange males will be favored by natural selection when the population is dominated by blue males, blue males will thrive when the population is mostly yellow males, and yellow males will be selected for when orange males are the most populous. As a result, populations of side-blotched lizards cycle in the distribution of these phenotypes—in one generation, orange might be predominant, and then yellow males will begin to rise in frequency. Once yellow males make up a majority of the population, blue males will be selected for. Finally, when blue males become common, orange males will once again be favored.

Negative frequency-dependent selection serves to increase the population’s genetic variance by selecting for rare phenotypes, whereas positive frequency-dependent selection usually decreases genetic variance by selecting for common phenotypes.

## Sexual Selection

Males and females of certain species are often quite different from one another in ways beyond the reproductive organs. Males are often larger, for example, and display many elaborate colors and adornments, like the peacock’s tail, while females tend to be smaller and duller in decoration. Such differences are known as **sexual dimorphisms** (Figure 3), which arise from the fact that in many populations, particularly animal populations, there is more variance in the reproductive success of the males than there is of the females. That is, some males—often the bigger, stronger, or more decorated males—get the vast majority of the total matings, while others receive none. This can occur because the males are better at fighting off other males, or because females will choose to mate with the bigger or more decorated males. In either case, this variation in reproductive success generates a strong selection pressure among males to get those matings, resulting in the evolution of bigger body size and elaborate ornaments to get the females’ attention. Females, on the other hand, tend to get a handful of selected matings; therefore, they are more likely to select more desirable males.

Sexual dimorphism varies widely among species, of course, and some species are even sex-role reversed. In such cases, females tend to have a greater variance in their reproductive success than males and are correspondingly selected for the bigger body size and elaborate traits usually characteristic of males.



(a) (b) (c)

Figure 3. Sexual dimorphism is observed in (a) peacocks and peahens, (b) Argiope appensa spiders (the female spider is the large one), and in (c) wood ducks. (credit “spiders”: modification of work by “Sanba38″/Wikimedia Commons; credit “duck”: modification of work by Kevin Cole)

The selection pressures on males and females to obtain matings is known as sexual selection; it can result in the development of secondary sexual characteristics that do not benefit the individual’s likelihood of survival but help to maximize its reproductive success. Sexual selection can be so strong that it selects for traits that are actually detrimental to the individual’s survival. Think, once again, about the peacock’s tail. While it is beautiful and the male with the largest, most colorful tail is more likely to win the female, it is not the most practical appendage. In addition to being more visible to predators, it makes the males slower in their attempted escapes. There is some evidence that this risk, in fact, is why females like the big tails in the first place. The speculation is that large tails carry risk, and only the best males survive that risk: the bigger the tail, the more fit the male. This idea is known as the **handicap principle**.

The **good genes hypothesis** states that males develop these impressive ornaments to show off their efficient metabolism or their ability to fight disease. Females then choose males with the most impressive traits because it signals their genetic superiority, which they will then pass on to their offspring. Though it might be argued that females should not be picky because it will likely reduce their number of offspring, if better males father more fit offspring, it may be beneficial. Fewer, healthier offspring may increase the chances of survival more than many, weaker offspring.

In both the handicap principle and the good genes hypothesis, the trait is said to be an **honest signal** of the males’ quality, thus giving females a way to find the fittest mates— males that will pass the best genes to their offspring.

## No Perfect Organism

Natural selection is a driving force in evolution and can generate populations that are better adapted to survive and successfully reproduce in their environments. But natural selection cannot produce the perfect organism. Natural selection can only select on existing variation in the population; it does not create anything from scratch. Thus, it is limited by a population’s existing genetic variance and whatever new alleles arise through mutation and gene flow.

Natural selection is also limited because it works at the level of individuals, not alleles, and some alleles are linked due to their physical proximity in the genome, making them more likely to be passed on together (linkage disequilibrium). Any given individual may carry some beneficial alleles and some unfavorable alleles. It is the net effect of these alleles, or the organism’s fitness, upon which natural selection can act. As a result, good alleles can be lost if they are carried by individuals that also have several overwhelmingly bad alleles; likewise, bad alleles can be kept if they are carried by individuals that have enough good alleles to result in an overall fitness benefit.

Furthermore, natural selection can be constrained by the relationships between different polymorphisms. One morph may confer a higher fitness than another, but may not increase in frequency due to the fact that going from the less beneficial to the more beneficial trait would require going through a less beneficial phenotype. Think back to the mice that live at the beach. Some are light-colored and blend in with the sand, while others are dark and blend in with the patches of grass. The dark-colored mice may be, overall, more fit than the light-colored mice, and at first glance, one might expect the light-colored mice be selected for a darker coloration. But remember that the intermediate phenotype, a medium-colored coat, is very bad for the mice—they cannot blend in with either the sand or the grass and are more likely to be eaten by predators. As a result, the light-colored mice would not be

selected for a dark coloration because those individuals that began moving in that direction (began being selected for a darker coat) would be less fit than those that stayed light.

Finally, it is important to understand that not all evolution is adaptive. While natural selection selects the fittest individuals and often results in a more fit population overall, other forces of evolution, including genetic drift and gene flow, often do the opposite: introducing deleterious alleles to the population's gene pool. Evolution has no purpose—it is not changing a population into a preconceived ideal. It is simply the sum of the various forces described in this chapter and how they influence the genetic and phenotypic variance of a population.

### Learning Objectives

Because natural selection acts to increase the frequency of beneficial alleles and traits while decreasing the frequency of deleterious qualities, it is adaptive evolution. Natural selection acts at the level of the individual, selecting for those that have a higher overall fitness compared to the rest of the population. If the fit phenotypes are those that are similar, natural selection will result in stabilizing selection, and an overall decrease in the population's variation. Directional selection works to shift a population's variance toward a new, fit phenotype, as environmental conditions change. In contrast, diversifying selection results in increased genetic variance by selecting for two or more distinct phenotypes.

Other types of selection include frequency-dependent selection, in which individuals with either common (positive frequency-dependent selection) or rare (negative frequency-dependent selection) phenotypes are selected for. Finally, sexual selection results from the fact that one sex has more variance in the reproductive success than the other. As a result, males and females experience different selective pressures, which can often lead to the evolution of phenotypic differences, or sexual dimorphisms, between the two.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.12: Selective and Environmental Pressures is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.13: Introduction to Phylogenetic Trees

## Read and analyze a phylogenetic tree that documents evolutionary relationships

In scientific terms, the evolutionary history and relationship of an organism or group of organisms is called phylogeny. **Phylogeny** describes the relationships of an organism, such as from which organisms it is thought to have evolved, to which species it is most closely related, and so forth. Phylogenetic relationships provide information on shared ancestry but not necessarily on how organisms are similar or different.

## What You'll Learn to Do

- Identify how and why scientists classify the organisms on earth
- Differentiate between types of phylogenetic trees and what their structure tells us
- Identify some limitations of phylogenetic trees
- Relate the taxonomic classification system and binomial nomenclature

### Learning Activities

The learning activities for this section include the following:

- Scientific Classification
- Structure of a Phylogenetic Tree
- Limitation of Phylogenetic Trees
- The Taxonomic Classification System
- Self Check: Phylogenetic Trees

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.13: Introduction to Phylogenetic Trees is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.14: Scientific Classification



Figure 1. Only a few of the more than one million known species of insects are represented in this beetle collection. Beetles are a major subgroup of insects. They make up about 40 percent of all insect species and about 25 percent of all known species of organisms.

Why do biologists classify organisms? The major reason is to make sense of the incredible diversity of life on Earth. Scientists have identified millions of different species of organisms. Among animals, the most diverse group of organisms is the insects. More than one million different species of insects have already been described. An estimated nine million insect species have yet to be identified. A tiny fraction of insect species is shown in the beetle collection in Figure 1.

As diverse as insects are, there may be even more species of bacteria, another major group of organisms. Clearly, there is a need to organize the tremendous diversity of life. Classification allows scientists to organize and better understand the basic similarities and differences among organisms. This knowledge is necessary to understand the present diversity and the past evolutionary history of life on Earth.

## Phylogenetic Trees

Scientists use a tool called a phylogenetic tree to show the evolutionary pathways and connections among organisms. A **phylogenetic tree** is a diagram used to reflect evolutionary relationships among organisms or groups of organisms. Scientists consider phylogenetic trees to be a hypothesis of the evolutionary past since one cannot go back to confirm the proposed relationships. In other words, a “tree of life” can be constructed to illustrate when different organisms evolved and to show the relationships among different organisms (Figure 1).

Each group of organisms went through its own evolutionary journey, called its phylogeny. Each organism shares relatedness with others, and based on morphologic and genetic evidence, scientists attempt to map the evolutionary pathways of all life on Earth. Many scientists build phylogenetic trees to illustrate evolutionary relationships.

## Contributors and Attributions

CC licensed content, Shared previously

- 14.1: Form and Function. **Provided by**: CK-12. **Located at**: www.ck12.org/book/CK-12-Biology-I-Honors-CA-DTI3/section/14.1/. **License**: *CC BY-NC: Attribution-NonCommercial*
- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.14: Scientific Classification is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.15: Structure of Phylogenetic Trees

A phylogenetic tree can be read like a map of evolutionary history. Many phylogenetic trees have a single lineage at the base representing a common ancestor. Scientists call such trees rooted, which means there is a single ancestral lineage (typically drawn from the bottom or left) to which all organisms represented in the diagram relate. Notice in the rooted phylogenetic tree that the three domains—Bacteria, Archaea, and Eukarya—diverge from a single point and branch off. The small branch that plants and animals (including humans) occupy in this diagram shows how recent and miniscule these groups are compared with other organisms. Unrooted trees don't show a common ancestor but do show relationships among species.



Figure 1. Both of these phylogenetic trees shows the relationship of the three domains of life—Bacteria, Archaea, and Eukarya—but the (a) rooted tree attempts to identify when various species diverged from a common ancestor while the (b) unrooted tree does not. (credit a: modification of work by Eric Gaba)

In a rooted tree, the branching indicates evolutionary relationships (Figure 2). The point where a split occurs, called a **branch point**, represents where a single lineage evolved into a distinct new one. A lineage that evolved early from the root and remains unbranched is called **basal taxon**. When two lineages stem from the same branch point, they are called **sister taxa**. A branch with more than two lineages is called a **polytomy** and serves to illustrate where scientists have not definitively determined all of the relationships. It is important to note that although sister taxa and polytomy do share an ancestor, it does not mean that the groups of organisms split or evolved from each other. Organisms in two taxa may have split apart at a specific branch point, but neither taxa gave rise to the other.





Figure 2. The root of a phylogenetic tree indicates that an ancestral lineage gave rise to all organisms on the tree. A branch point indicates where two lineages diverged. A lineage that evolved early and remains unbranched is a basal taxon. When two lineages stem from the same branch point, they are sister taxa. A branch with more than two lineages is a polytomy.

The diagrams above can serve as a pathway to understanding evolutionary history. The pathway can be traced from the origin of life to any individual species by navigating through the evolutionary branches between the two points. Also, by starting with a single species and tracing back towards the "trunk" of the tree, one can discover that species' ancestors, as well as where lineages share a common ancestry. In addition, the tree can be used to study entire groups of organisms.

Another point to mention on phylogenetic tree structure is that rotation at branch points does not change the information. For example, if a branch point was rotated and the taxon order changed, this would not alter the information because the evolution of each taxon from the branch point was independent of the other.

Many disciplines within the study of biology contribute to understanding how past and present life evolved over time; these disciplines together contribute to building, updating, and maintaining the “tree of life.” Information is used to organize and classify organisms based on evolutionary relationships in a scientific field called systematics. Data may be collected from fossils, from studying the structure of body parts or molecules used by an organism, and by DNA analysis. By combining data from many sources, scientists can put together the phylogeny of an organism; since phylogenetic trees are hypotheses, they will continue to change as new types of life are discovered and new information is learned.

## Video Review

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Phylogeny and Phylogentic Trees. **Authored by**: Complex Life. **Located at**: https://youtu.be/iyAOkzdO3vw. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

15.15: Structure of Phylogenetic Trees is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.16: Limitations of Phylogenetic Trees

It may be easy to assume that more closely related organisms look more alike, and while this is often the case, it is not always true. If two closely related lineages evolved under significantly varied surroundings or after the evolution of a major new adaptation, it is possible for the two groups to appear more different than other groups that are not as closely related. For example, the phylogenetic tree in Figure 1 shows that lizards and rabbits both have amniotic eggs, whereas frogs do not; yet lizards and frogs appear more similar than lizards and rabbits.





Figure 1. This ladder-like phylogenetic tree of vertebrates is rooted by an organism that lacked a vertebral column. At each branch point, organisms with different characters are placed in different groups based on the characteristics they share.

Another aspect of phylogenetic trees is that, unless otherwise indicated, the branches do not account for length of time, only the evolutionary order. In other words, the length of a branch does not typically mean more time passed, nor does a short branch mean less time passed— unless specified on the diagram. For example, in Figure 1, the tree does not indicate how much time passed between the evolution of amniotic eggs and hair. What the tree does show is the order in which things took place. Again using Figure 1, the tree shows that the oldest trait is the vertebral column, followed by hinged jaws, and so forth. Remember that any phylogenetic tree is a part of the greater whole, and like a real tree, it does not grow in only one direction after a new branch develops.

So, for the organisms in Figure 1, just because a vertebral column evolved does not mean that invertebrate evolution ceased, it only means that a new branch formed. Also, groups that are not closely related, but evolve under similar conditions, may appear more phenotypically similar to each other than to a close relative.

Head to this website to see interactive exercises that allow you to explore the evolutionary relationships among species.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.16: Limitations of Phylogenetic Trees is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.17: The Taxonomic Classification System

**Taxonomy** (which literally means “arrangement law”) is the science of classifying organisms to construct internationally shared classification systems with each organism placed into more and more inclusive groupings. Think about how a grocery store is organized. One large space is divided into departments, such as produce, dairy, and meats. Then each department further divides into aisles, then each aisle into categories and brands, and then finally a single product. This organization from larger to smaller, more specific categories is called a hierarchical system.

The taxonomic classification system (also called the Linnaean system after its inventor, Carl Linnaeus, a Swedish botanist, zoologist, and physician) uses a hierarchical model. Moving from the point of origin, the groups become more specific, until one branch ends as a single species. For example, after the common beginning of all life, scientists divide organisms into three large categories called a domain: Bacteria, Archaea, and Eukarya. Within each domain is a second category called a **kingdom**. After kingdoms, the subsequent categories of increasing specificity are: **phylum**, **class**, **order**, **family**, **genus**, and **species** (Figure 1).





Figure 1. The taxonomic classification system uses a hierarchical model to organize living organisms into increasingly specific categories. The common dog, *Canis lupus familiaris*, is a subspecies of *Canis lupus*, which also includes the wolf and dingo. (credit “dog”: modification of work by Janneke Vreugdenhil)

The kingdom Animalia stems from the Eukarya domain. For the common dog, the classification levels would be as shown in Figure 1. Therefore, the full name of an organism technically has eight terms. For the dog, it is: Eukarya, Animalia, Chordata, Mammalia, Carnivora, Canidae, *Canis*, and *lupus*. Notice that each name is capitalized except for species, and the genus and species names are italicized. Scientists generally refer to an organism only by its genus and species, which is its two-word scientific name, in what is called **binomial nomenclature**. Therefore, the scientific name of the dog is *Canis lupus*. The name at each level is also called a **taxon**. In other words, dogs are in order Carnivora. Carnivora is the name of the taxon at the order level; Canidae is the taxon at the family level, and so forth. Organisms also have a common name that people typically use, in this case, dog. Note that the dog is additionally a subspecies: the "*familiaris*" in *Canis lupus familiaris*. Subspecies are members of the same species that are capable of mating and reproducing viable offspring, but they are considered separate subspecies due to geographic or behavioral isolation or other factors.

Figure 2 shows how the levels move toward specificity with other organisms. Notice how the dog shares a domain with the widest diversity of organisms, including plants and butterflies. At each sublevel, the organisms become more similar because they are more closely related. Historically, scientists classified organisms using characteristics, but as DNA technology developed, more precise phylogenies have been determined.

## Practice Question





Figure 2. At each sublevel in the taxonomic classification system, organisms become more similar. Dogs and wolves are the same species because they can breed and produce viable offspring, but they are different enough to be classified as different subspecies. (credit “plant”: modification of work by “berduchwal”/Flickr; credit “insect”: modification of work by Jon Sullivan; credit “fish”: modification of work by Christian Mehlführer; credit “rabbit”: modification of work by Aidan Wojtas; credit “cat”: modification of work by Jonathan Lidbeck; credit “fox”: modification of work by Kevin Bacher, NPS; credit “jackal”: modification of work by Thomas A. Hermann, NBII, USGS; credit “wolf”: modification of work by Robert Dewar; credit “dog”: modification of work by “digital_image_fan”/Flickr)

At what levels are cats and dogs considered to be part of the same group?

[practice-area rows=”2″][/practice-area]

[reveal-answer q=”854290″]**Show Answer**[/reveal-answer]

[hidden-answer a=”854290″]Cats and dogs are part of the same group at five levels: both are in the domain Eukarya, the kingdom Animalia, the phylum Chordata, the class Mammalia, and the order Carnivora.[/hidden-answer]

Visit this website to classify three organisms—bear, orchid, and sea cucumber—from kingdom to species. To launch the game, under Classifying Life, click the picture of the bear or the Launch Interactive button.

Recent genetic analysis and other advancements have found that some earlier phylogenetic classifications do not align with the evolutionary past; therefore, changes and updates must be made as new discoveries occur. Recall that phylogenetic trees are hypotheses and are modified as data becomes available. In addition, classification historically has focused on grouping organisms mainly by shared characteristics and does not necessarily illustrate how the various groups relate to each other from an evolutionary perspective. For example, despite the fact that a hippopotamus resembles a pig more than a whale, the hippopotamus may be the closest living relative of the whale.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

15.17: The Taxonomic Classification System is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 15.18: Putting It Together- Theory of Evolution

Let's return to our discussion of evolving bacteria from the beginning of the module. The World Health Organization (WHO) suggests that based on current evidence, gonorrhea may soon be untreatable; there are no new treatments or vaccinations in development for this bacteria. Within the United States this remain the second most common sexually transmitted disease. Almost a third of the US cases are drug resistant. MRSA remains a growing problem in our healthcare system. Staph bacteria are the most common source of health-care related infections in the United States. Of MRSA infections, nearly 14% result in death.

## Think about It

Based on this lesson you now understand that evolution is an inevitable process. Is there any hope of stopping antibiotic resistance from developing?

[practice-area rows="4"][/practice-area]
[reveal-answer q="880848"]See Our Thoughts[/reveal-answer]
[hidden-answer a="880848"]Yes and no. The average person can take a number of steps to minimize selective pressures on parasites. Without these pressures, mutations are less likely to spread in the population.

[/hidden-answer]

Is there anything you can do?

[practice-area rows="4"][/practice-area]
[reveal-answer q="558353"]See Our Thoughts[/reveal-answer]
[hidden-answer a="558353"]

- Take the full course of prescribed antibiotics
- Don't save or share leftover antibiotics
- Don't ask for antibiotics if your doctor does not recommend them
- Practice good hygiene and get recommended vaccinations
- Minimize your use of antibacterial products

[/hidden-answer]

To learn more about the topic, you can visit the following websites:

- WHO—Antimicrobial Resistance
- CDC—Antibiotic Resistance Threats
- CDC—Mission Critical: Preventing Antibiotic Resistance
- FDA—Antibacterial Soap

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

15.18: Putting It Together- Theory of Evolution is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# CHAPTER OVERVIEW

## 16: Module 14- Modern Biology


16.1: Why It Matters- Modern Biology
16.2: Introduction to Key Technologies
16.3: Manipulating Genetic Material
16.4: DNA Sequencing
16.5: Cloning
16.6: Genetic Engineering
16.7: Introduction to Biotechnology Applications
16.8: Medicinal Biotechnology
16.9: Agricultural Biotechnology
16.10: Biotechnology at the Cutting Edge
16.11: Introduction to Risks and Benefits of Genomic Science
16.12: Genetic Information Used for Identification
16.13: Medical Uses of Genetic Information
16.14: Genomics in Agriculture
16.15: Putting It Together- Modern Biology
16.16: Assignment- When Jabberjays Attack!


16: Module 14- Modern Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.1: Why It Matters- Modern Biology

## Why describe and discuss techniques used in modern biology?

Biology has impacted nearly every facet of human life. The entire field of medicine is based on biological discovery and exploration; without knowledge of how the human body functions, we would have no idea how to fix our bodies when something goes wrong.

Over the decades, new techniques have been developed to study biology. Scientists recently mapped the complete human genome. New technologies were developed to help speed this process up. Forensic science has also been vastly influenced by biological discovery. New techniques enable law enforcement to gather DNA evidence at crime scenes, which helps ensure they identify the right suspect. DNA based information can be applied to many such situations. Consider the following scenario.

On Memorial Day 1984, the remains of a soldier killed during the Vietnam War were laid to rest in the Tomb of the Unknown at Arlington National Cemetery. A decade later, however, questions began to arise about the identity of this unknown soldier. In particular, people began to wonder if, due to scientific improvement and developments, the soldier could be identified.

### Learning Outcomes

- List key technologies enabling modern uses of biology
- Identify societal uses of biotechnology
- Discuss the risks and benefits involved in applications of genetic and genomic science

## Contributors and Attributions

CC licensed content, Original

- Why It Matters: Modern Biology. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

16.1: Why It Matters- Modern Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.2: Introduction to Key Technologies

## List key technologies enabling modern uses of biology

Biotechnology is the use of biological agents for technological advancement. Biotechnology was used for breeding livestock and crops long before the scientific basis of these techniques was understood. Since the discovery of the structure of DNA in 1953, the field of biotechnology has grown rapidly through both academic research and private companies. The primary applications of this technology are in medicine (production of vaccines and antibiotics) and agriculture (genetic modification of crops, such as to increase nutrient content). Biotechnology also has many industrial applications, such as fermentation, the treatment of oil spills, and the production of biofuels (Figure 1).





Figure 1. Antibiotics are chemicals produced by fungi, bacteria, and other organisms that have antimicrobial properties. The first antibiotic discovered was penicillin. Antibiotics are now commercially produced and tested for their potential to inhibit bacterial growth. (credit “advertisement”: modification of work by NIH; credit “test plate”: modification of work by Don Stalons/CDC; scale-bar data from Matt Russell)

In this outcome, we will learn about some modern technologies used in biology today.

## What You’ll Learn to Do

- List basic techniques to manipulation genetic information (DNA and RNA)
- Recognize technologies used for molecular, cellular, and reproductive cloning
- Understand the basics of genetic engineering

### Learning Activities

The learning activities for this section include the following:

- Manipulating Genetic Material
- Cloning
- Genetic Engineering
- Self Check: Key Technologies

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.2: Introduction to Key Technologies is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.3: Manipulating Genetic Material

To understand the basic techniques used to work with nucleic acids, remember that nucleic acids are macromolecules made of nucleotides (a sugar, a phosphate, and a nitrogenous base) linked by phosphodiester bonds. The phosphate groups on these molecules each have a net negative charge. An entire set of DNA molecules in the nucleus is called the genome. DNA has two complementary strands linked by hydrogen bonds between the paired bases. The two strands can be separated by exposure to high temperatures (DNA denaturation) and can be reannealed by cooling. The DNA can be replicated by the DNA polymerase enzyme. Unlike DNA, which is located in the nucleus of eukaryotic cells, RNA molecules leave the nucleus. The most common type of RNA that is analyzed is the messenger RNA (mRNA) because it represents the protein-coding genes that are actively expressed. However, RNA molecules present some other challenges to analysis, as they are often less stable than DNA.

## DNA and RNA Extraction

To study or manipulate nucleic acids, the DNA or RNA must first be isolated or extracted from the cells. Various techniques are used to extract different types of DNA (Figure 1). Most nucleic acid extraction techniques involve steps to break open the cell and use enzymatic reactions to destroy all macromolecules that are not desired (such as degradation of unwanted molecules and separation from the DNA sample). Cells are broken using a **lysis buffer** (a solution which is mostly a detergent); lysis means “to split.” These enzymes break apart lipid molecules in the cell membranes and nuclear membranes. Macromolecules are inactivated using enzymes such as **proteases** that break down proteins, and **ribonucleases** (RNAses) that break down RNA. The DNA is then precipitated using alcohol. Human genomic DNA is usually visible as a gelatinous, white mass. The DNA samples can be stored frozen at –80°C for several years.





Figure 1. This diagram shows the basic method used for extraction of DNA.

RNA analysis is performed to study gene expression patterns in cells. RNA is naturally very unstable because RNAses are commonly present in nature and very difficult to inactivate. Similar to DNA, RNA extraction involves the use of various buffers and enzymes to inactivate macromolecules and preserve the RNA.




Figure 2. Shown are DNA fragments from seven samples run on a gel, stained with a fluorescent dye, and viewed under UV light. (credit: James Jacob, Tompkins Cortland Community College)

## Gel Electrophoresis

Because nucleic acids are negatively charged ions at neutral or basic pH in an aqueous environment, they can be mobilized by an electric field. **Gel electrophoresis** is a technique used to separate molecules on the basis of size, using this charge. The nucleic acids can be separated as whole chromosomes or fragments. The nucleic acids are loaded into a slot near the negative electrode of a semisolid, porous gel matrix and pushed toward the positive electrode at the opposite end of the gel. Smaller molecules move through the pores in the gel faster than larger molecules; this difference in the rate of migration separates the fragments on the basis of size. There are molecular weight standard samples that can be run alongside the molecules to provide a size comparison. Nucleic acids in a gel matrix can be observed using various fluorescent or colored dyes. Distinct nucleic acid fragments appear as bands at specific distances from the top of the gel (the negative electrode end) on the basis of their size (Figure 2). A mixture of genomic DNA fragments of varying sizes appear as a long smear, whereas uncut genomic DNA is usually too large to run through the gel and forms a single large band at the top of the gel.

### Amplification of Nucleic Acid Fragments by Polymerase Chain Reaction

Although genomic DNA is visible to the naked eye when it is extracted in bulk, DNA analysis often requires focusing on one or more specific regions of the genome. **Polymerase chain reaction (PCR)** is a technique used to amplify specific regions of DNA for further analysis (Figure 3). PCR is used for many purposes in laboratories, such as the cloning of gene fragments to analyze genetic diseases, identification of contaminant foreign DNA in a sample, and the amplification of DNA for sequencing. More practical applications include the determination of paternity and detection of genetic diseases.





Figure 3. Polymerase chain reaction, or PCR, is used to amplify a specific sequence of DNA. Primers—short pieces of DNA complementary to each end of the target sequence—are combined with genomic DNA, Taq polymerase, and deoxynucleotides. Taq polymerase is a DNA polymerase isolated from the thermostable bacterium *Thermus aquaticus* that is able to withstand the high temperatures used in PCR. *Thermus aquaticus* grows naturally in the Lower Geyser Basin of Yellowstone National Park. Reverse transcriptase PCR (RT-PCR) is similar to PCR, but cDNA is made from an RNA template before PCR begins.

DNA fragments can also be amplified from an RNA template in a process called reverse **transcriptase PCR (RT-PCR)**. The first step is to recreate the original DNA template strand (called cDNA) by applying DNA nucleotides to the mRNA. This process is called reverse transcription. This requires the presence of an enzyme called reverse transcriptase. After the cDNA is made, regular PCR can be used to amplify it.

Deepen your understanding of the polymerase chain reaction by clicking through this interactive exercise.

## Hybridization, Southern Blotting, and Northern Blotting

Nucleic acid samples, such as fragmented genomic DNA and RNA extracts, can be probed for the presence of certain sequences. Short DNA fragments called **probes** are designed and labeled with radioactive or fluorescent dyes to aid detection. Gel electrophoresis separates the nucleic acid fragments according to their size. The fragments in the gel are then transferred onto a nylon membrane in a procedure called **blotting** (Figure 4). The nucleic acid fragments that are bound to the surface of the membrane can then be probed with specific radioactively or fluorescently labeled probe sequences. When DNA is transferred to a

nylon membrane, the technique is called **Southern blotting**, and when RNA is transferred to a nylon membrane, it is called **northern blotting**. Southern blots are used to detect the presence of certain DNA sequences in a given genome, and northern blots are used to detect gene expression. Note that "Southern" blotting is capitalized but no other type of blotting; Southern blotting is named after the scientist who pioneered this technique, Edwin Southern. The other types of blotting were named in reference to the original Southern blot.





Figure 4. Southern blotting is used to find a particular sequence in a sample of DNA. DNA fragments are separated on a gel, transferred to a nylon membrane, and incubated with a DNA probe complementary to the sequence of interest. Northern blotting is similar to Southern blotting, but RNA is run on the gel instead of DNA. In western blotting, proteins are run on a gel and detected using antibodies.

## Contributors and Attributions

CC licensed content, Original

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.3: Manipulating Genetic Material is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.4: DNA Sequencing

Until the 1990s, the sequencing of DNA (reading the sequence of DNA) was a relatively expensive and long process. Using radiolabeled nucleotides also compounded the problem through safety concerns. With currently available technology and automated machines, the process is cheap, safer, and can be completed in a matter of hours. Fred Sanger developed the sequencing method used for the human genome sequencing project, which is widely used today (Figure 1).





Figure 1. In Frederick Sanger's dideoxy chain termination method, dye-labeled dideoxynucleotides are used to generate DNA fragments that terminate at different points. The DNA is separated by capillary electrophoresis on the basis of size, and from the order of fragments formed, the DNA sequence can be read. The DNA sequence readout is shown on an electropherogram that is generated by a laser scanner.

Visit this site to watch a video explaining the DNA sequence reading technique that resulted from Sanger's work.

The method is known as the dideoxy chain termination method. The sequencing method is based on the use of chain terminators, the dideoxynucleotides (ddNTPs). The dideoxynucleotides, or ddNTPSs, differ from the deoxynucleotides by the lack of a free 3′ OH group on the five-carbon sugar. If a ddNTP is added to a growing a DNA strand, the chain is not extended any further because the free 3′ OH group needed to add another nucleotide is not available. By using a predetermined ratio of deoxyribonucleotides to dideoxynucleotides, it is possible to generate DNA fragments of different sizes.




Figure 2. DNA can be separated on the basis of size using gel electrophoresis. (credit: James Jacob, Tompkins Cortland Community College)

The DNA sample to be sequenced is denatured or separated into two strands by heating it to high temperatures. The DNA is divided into four tubes in which a primer, DNA polymerase, and all four nucleotides (A, T, G, and C) are added. In addition to each of the four tubes, limited quantities of one of the four dideoxynucleotides are added to each tube respectively. The tubes are labeled as A, T, G, and C according to the ddNTP added. For detection purposes, each of the four dideoxynucleotides carries a different fluorescent label. Chain elongation continues until a fluorescent dideoxy nucleotide is incorporated, after which no further

elongation takes place. After the reaction is over, electrophoresis is performed. Even a difference in length of a single base can be detected. The sequence is read from a laser scanner. For his work on DNA sequencing, Sanger received a Nobel Prize in chemistry in 1980.

As already mentioned, gel **electrophoresis** (Figure 2) is a technique used to separate DNA fragments of different sizes. Usually the gel is made of a chemical called agarose. Agarose powder is added to a buffer and heated. After cooling, the gel solution is poured into a casting tray. Once the gel has solidified, the DNA is loaded on the gel and electric current is applied. The DNA has a net negative charge and moves from the negative electrode toward the positive electrode. The electric current is applied for sufficient time to let the DNA separate according to size; the smallest fragments will be farthest from the well (where the DNA was loaded), and the heavier molecular weight fragments will be closest to the well. Once the DNA is separated, the gel is stained with a DNA-specific dye for viewing it.

Sanger's genome sequencing has led to a race to sequence human genomes at a rapid speed and low cost, often referred to as the $1000 in one day sequence. Learn more by watching the Sequencing at Speed animation here.

## Neanderthal Genome: How Are We Related?

The first draft sequence of the Neanderthal genome was published by Richard E. Green et al. in 2010.[1] Neanderthals are the closest ancestors of present-day humans. They were known to have lived in Europe and Western Asia before they disappeared from fossil records approximately 30,000 years ago. Green's team studied almost 40,000-year-old fossil remains that were selected from sites across the world. Extremely sophisticated means of sample preparation and DNA sequencing were employed because of the fragile nature of the bones and heavy microbial contamination. In their study, the scientists were able to sequence some four billion base pairs. The Neanderthal sequence was compared with that of present-day humans from across the world. After comparing the sequences, the researchers found that the Neanderthal genome had 2 to 3 percent greater similarity to people living outside Africa than to people in Africa. While current theories have suggested that all present-day humans can be traced to a small ancestral population in Africa, the data from the Neanderthal genome may contradict this view. Green and his colleagues also discovered DNA segments among people in Europe and Asia that are more similar to Neanderthal sequences than to other contemporary human sequences. Another interesting observation was that Neanderthals are as closely related to people from Papua New Guinea as to those from China or France. This is surprising because Neanderthal fossil remains have been located only in Europe and West Asia. Most likely, genetic exchange took place between Neanderthals and modern humans as modern humans emerged out of Africa, before the divergence of Europeans, East Asians, and Papua New Guineans.

Several genes seem to have undergone changes from Neanderthals during the evolution of present-day humans. These genes are involved in cranial structure, metabolism, skin morphology, and cognitive development. One of the genes that is of particular interest is *RUNX2*, which is different in modern day humans and Neanderthals. This gene is responsible for the prominent frontal bone, bell-shaped rib cage, and dental differences seen in Neanderthals. It is speculated that an evolutionary change in *RUNX2* was important in the origin of modern-day humans, and this affected the cranium and the upper body.

Watch Svante Pääbo's talk explaining the Neanderthal genome research at the 2011 annual TED (Technology, Entertainment, Design) conference.



A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=566

1. Richard E. Green et al., “A Draft Sequence of the Neandertal Genome,” *Science* 328 (2010): 710–22. ↵

## Contributors and Attributions

CC licensed content, Original

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Svante Paabo: DNA clues to our inner neanderthal. **Authored by**: TED. **Located at**: https://youtu.be/kU0ei9ApmsY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

16.4: DNA Sequencing is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.5: Cloning

## Molecular Cloning

In general, the word “cloning” means the creation of a perfect replica; however, in biology, the re-creation of a whole organism is referred to as “reproductive cloning.” Long before attempts were made to clone an entire organism, researchers learned how to reproduce desired regions or fragments of the genome, a process that is referred to as molecular cloning.

Cloning small fragments of the genome allows for the manipulation and study of specific genes (and their protein products), or noncoding regions in isolation. A plasmid (also called a vector) is a small circular DNA molecule that replicates independently of the chromosomal DNA of microorganisms such as *E. coli*. In cloning, the plasmid molecules can be used to provide a “folder” in which to insert a desired DNA fragment. Plasmids are usually introduced into a bacterial host for proliferation. In the bacterial context, the fragment of DNA from the human genome (or the genome of another organism that is being studied) is referred to as **foreign DNA**, or a transgene, to differentiate it from the DNA of the bacterium, which is called the **host DNA**.

Plasmids occur naturally in bacterial populations (such as *Escherichia coli*) and have genes that can contribute favorable traits to the organism, such as **antibiotic resistance** (the ability to be unaffected by antibiotics). Plasmids have been repurposed and engineered as vectors for molecular cloning and the large-scale production of important reagents, such as insulin and human growth hormone. An important feature of plasmid vectors is the ease with which a foreign DNA fragment can be introduced via the **multiple cloning site (MCS)**. The MCS is a short DNA sequence containing multiple sites that can be cut with different commonly available restriction endonucleases. **Restriction endonucleases** recognize specific DNA sequences and cut them in a predictable manner; they are naturally produced by bacteria as a defense mechanism against foreign DNA. Many restriction endonucleases make staggered cuts in the two strands of DNA, such that the cut ends have a 2- or 4-base single-stranded overhang. Because these overhangs are capable of annealing with complementary overhangs, these are called “sticky ends.” Addition of an enzyme called DNA ligase permanently joins the DNA fragments via phosphodiester bonds. In this way, any DNA fragment generated by restriction endonuclease cleavage can be spliced between the two ends of a plasmid DNA that has been cut with the same restriction endonuclease (Figure 1).

### Recombinant DNA Molecules

Plasmids with foreign DNA inserted into them are called **recombinant DNA** molecules because they are created artificially and do not occur in nature. They are also called chimeric molecules because the origin of different parts of the molecules can be traced back to different species of biological organisms or even to chemical synthesis. Proteins that are expressed from recombinant DNA molecules are called **recombinant proteins**. Not all recombinant plasmids are capable of expressing genes. The recombinant DNA may need to be moved into a different vector (or host) that is better designed for gene expression. Plasmids may also be engineered to express proteins only when stimulated by certain environmental factors, so that scientists can control the expression of the recombinant proteins.

View an animation of recombination in cloning from the DNA Learning Center.

## Cellular Cloning

Unicellular organisms, such as bacteria and yeast, naturally produce clones of themselves when they replicate asexually by binary fission; this is known as **cellular cloning**. The nuclear DNA duplicates by the process of mitosis, which creates an exact replica of the genetic material.

Practice Question





Figure 1. This diagram shows the steps involved in molecular cloning.

You are working in a molecular biology lab and, unbeknownst to you, your lab partner left the foreign genomic DNA that you are planning to clone on the lab bench overnight instead of storing it in the freezer. As a result, it was degraded by nucleases, but still used in the experiment. The plasmid, on the other hand, is fine. What results would you expect from your molecular cloning experiment?

a. There will be no colonies on the bacterial plate.
b. There will be blue colonies only.
c. There will be blue and white colonies.
d. The will be white colonies only.

[reveal-answer q="511969"]**Show Answer**[/reveal-answer]
[hidden-answer a="511969"]Answer b. The experiment would result in blue colonies only.[/hidden-answer]

## Reproductive Cloning

**Reproductive cloning** is a method used to make a clone or an identical copy of an entire multicellular organism. Most multicellular organisms undergo reproduction by sexual means, which involves genetic hybridization of two individuals (parents), making it impossible for generation of an identical copy or a clone of either parent. Recent advances in biotechnology have made it possible to artificially induce asexual reproduction of mammals in the laboratory.

Parthenogenesis, or “virgin birth,” occurs when an embryo grows and develops without the fertilization of the egg occurring; this is a form of asexual reproduction. An example of parthenogenesis occurs in species in which the female lays an egg and if the egg is fertilized, it is a diploid egg and the individual develops into a female; if the egg is not fertilized, it remains a haploid egg and develops into a male. The unfertilized egg is called a parthenogenic, or virgin, egg. Some insects and reptiles lay parthenogenic eggs that can develop into adults.

Sexual reproduction requires two cells; when the haploid egg and sperm cells fuse, a diploid zygote results. The zygote nucleus contains the genetic information to produce a new individual. However, early embryonic development requires the cytoplasmic material contained in the egg cell. This idea forms the basis for reproductive cloning. Therefore, if the haploid nucleus of an egg cell is replaced with a diploid nucleus from the cell of any individual of the same species (called a donor), it will become a zygote that is genetically identical to the donor. Somatic cell nuclear transfer is the technique of transferring a diploid nucleus into an enucleated egg. It can be used for either therapeutic cloning or reproductive cloning.

The first cloned animal was Dolly, a sheep who was born in 1996. The success rate of reproductive cloning at the time was very low. Dolly lived for seven years and died of respiratory complications (Figure 2). There is speculation that because the cell DNA belongs to an older individual, the age of the DNA may affect the life expectancy of a cloned individual. Since Dolly, several animals such as horses, bulls, and goats have been successfully cloned, although these individuals often exhibit facial, limb, and cardiac abnormalities. There have been attempts at producing cloned human embryos as sources of embryonic stem cells, sometimes referred to as cloning for therapeutic purposes. Therapeutic cloning produces stem cells to attempt to remedy detrimental diseases or defects (unlike reproductive cloning, which aims to reproduce an organism). Still, therapeutic cloning efforts have met with resistance because of bioethical considerations.

Practice Question





Figure 2. Dolly the sheep was the first mammal to be cloned. To create Dolly, the nucleus was removed from a donor egg cell. The nucleus from a second sheep was then introduced into the cell, which was allowed to divide to the blastocyst stage before being implanted in a surrogate mother. (credit: modification of work by “Squidonius”/Wikimedia Commons)

Do you think Dolly was a Finn-Dorset or a Scottish Blackface sheep?

[practice-area rows=”2″][/practice-area]
[reveal-answer q=”985505″]**Show Answer**[/reveal-answer]

[hidden-answer a=”985505″]Dolly was a Finn-Dorset sheep because even though the original cell came from a Scottish blackface sheep and the surrogate mother was a Scottish blackface, the DNA came from a Finn-Dorset.[/hidden-answer]

## Contributors and Attributions

CC licensed content, Original

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.5: Cloning is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.6: Genetic Engineering

**Genetic engineering** is the alteration of an organism's genotype using recombinant DNA technology to modify an organism's DNA to achieve desirable traits. The addition of foreign DNA in the form of recombinant DNA vectors generated by molecular cloning is the most common method of genetic engineering. The organism that receives the recombinant DNA is called a **genetically modified organism (GMO)**. If the foreign DNA that is introduced comes from a different species, the host organism is called **transgenic**. Bacteria, plants, and animals have been genetically modified since the early 1970s for academic, medical, agricultural, and industrial purposes. In the US, GMOs such as Roundup-ready soybeans and borer-resistant corn are part of many common processed foods.

## Gene Targeting

Although classical methods of studying the function of genes began with a given phenotype and determined the genetic basis of that phenotype, modern techniques allow researchers to start at the DNA sequence level and ask: "What does this gene or DNA element do?" This technique, called reverse genetics, has resulted in reversing the classic genetic methodology. This method would be similar to damaging a body part to determine its function. An insect that loses a wing cannot fly, which means that the function of the wing is flight. The classical genetic method would compare insects that cannot fly with insects that can fly, and observe that the non-flying insects have lost wings. Similarly, mutating or deleting genes provides researchers with clues about gene function. The methods used to disable gene function are collectively called gene targeting. **Gene targeting** is the use of recombinant DNA vectors to alter the expression of a particular gene, either by introducing mutations in a gene, or by eliminating the expression of a certain gene by deleting a part or all of the gene sequence from the genome of an organism.

## Contributors and Attributions

CC licensed content, Original

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.6: Genetic Engineering is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.7: Introduction to Biotechnology Applications

## Identify societal uses of biotechnology

In this outcome we will discuss the ways biotechnology is used in our society.

## What You'll Learn to Do

- List at least 3 medicinal uses of biotechnology advances
- List at least 3 agricultural uses of biotechnology advances

### Learning Activities

The learning activities for this section include the following:

- Medicinal Biotechnology
- Agricultural Biotechnology
- Biotechnology at the Cutting Edge
- Self Check: Biotechnology Applications

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

16.7: Introduction to Biotechnology Applications is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.8: Medicinal Biotechnology

It is easy to see how biotechnology can be used for medicinal purposes. Knowledge of the genetic makeup of our species, the genetic basis of heritable diseases, and the invention of technology to manipulate and fix mutant genes provides methods to treat the disease.

## Genetic Diagnosis and Gene Therapy





Figure 1. Gene therapy using an adenovirus vector can be used to cure certain genetic diseases in which a person has a defective gene. (credit: NIH)

The process of testing for suspected genetic defects before administering treatment is called **genetic diagnosis** by **genetic testing**. Depending on the inheritance patterns of a disease-causing gene, family members are advised to undergo genetic testing. For example, women diagnosed with breast cancer are usually advised to have a biopsy so that the medical team can determine the genetic basis of cancer development. Treatment plans are based on the findings of genetic tests that determine the type of cancer. If the cancer is caused by inherited gene mutations, other female relatives are also advised to undergo genetic testing and periodic screening for breast cancer. Genetic testing is also offered for fetuses (or embryos with *in vitro* fertilization) to determine the presence or absence of disease-causing genes in families with specific debilitating diseases.

**Gene therapy** is a genetic engineering technique used to cure disease. In its simplest form, it involves the introduction of a good gene at a random location in the genome to aid the cure of a disease that is caused by a mutated gene. The good gene is usually introduced into diseased cells as part of a vector transmitted by a virus that can infect the host cell and deliver the foreign DNA (Figure 1). More advanced forms of gene therapy try to correct the mutation at the original site in the genome, such as is the case with treatment of severe combined immunodeficiency (SCID).

Here's a website to help you learn more information on SCID and its gene therapy trials.

## Production of Vaccines, Antibiotics, and Hormones

Traditional vaccination strategies use weakened or inactive forms of microorganisms to mount the initial immune response. Modern techniques use the genes of microorganisms cloned into vectors to mass produce the desired antigen. The antigen is then introduced into the body to stimulate the primary immune response and trigger immune memory. Genes cloned from the influenza virus have been used to combat the constantly changing strains of this virus.

Antibiotics are a biotechnological product. They are naturally produced by microorganisms, such as fungi, to attain an advantage over bacterial populations. Antibiotics are produced on a large scale by cultivating and manipulating fungal cells.

Recombinant DNA technology was used to produce large-scale quantities of human insulin in *E. coli* as early as 1978. Previously, it was only possible to treat diabetes with pig insulin, which caused allergic reactions in humans because of differences in the gene product. In addition, human growth hormone (HGH) is used to treat growth disorders in children. The HGH gene was cloned from a cDNA library and inserted into *E. coli* cells by cloning it into a bacterial vector.

## Learning Objectives

Transgenic organisms possess DNA from a different species, usually generated by molecular cloning techniques. Vaccines, antibiotics, and hormones are examples of products obtained by recombinant DNA technology. Transgenic plants are usually created to improve characteristics of crop plants.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.8: Medicinal Biotechnology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.9: Agricultural Biotechnology

Biotechnology in agriculture can enhance resistance to disease, pest, and environmental stress, and improve both crop yield and quality.

## Transgenic Animals

Although several recombinant proteins used in medicine are successfully produced in bacteria, some proteins require a eukaryotic animal host for proper processing. For this reason, the desired genes are cloned and expressed in animals, such as sheep, goats, chickens, and mice. Animals that have been modified to express recombinant DNA are called transgenic animals. Several human proteins are expressed in the milk of transgenic sheep and goats, and some are expressed in the eggs of chickens. Mice have been used extensively for expressing and studying the effects of recombinant genes and mutations.

## Transgenic Plants



Figure 1. Corn, a major agricultural crop used to create products for a variety of industries, is often modified through plant biotechnology. (credit: Keith Weller, USDA)

Manipulating the DNA of plants (i.e., creating GMOs) has helped to create desirable traits, such as disease resistance, herbicide and pesticide resistance, better nutritional value, and better shelf-life (Figure 1). Plants are the most important source of food for the human population. Farmers developed ways to select for plant varieties with desirable traits long before modern-day biotechnology practices were established.

Plants that have received recombinant DNA from other species are called transgenic plants. Because they are not natural, transgenic plants and other GMOs are closely monitored by government agencies to ensure that they are fit for human consumption and do not endanger other plant and animal life. Because foreign genes can spread to other species in the environment, extensive testing is required to ensure ecological stability. Staples like corn, potatoes, and tomatoes were the first crop plants to be genetically engineered.

### Transformation of Plants Using *Agrobacterium tumefaciens*

Gene transfer occurs naturally between species in microbial populations. Many viruses that cause human diseases, such as cancer, act by incorporating their DNA into the human genome. In plants, tumors caused by the bacterium *Agrobacterium tumefaciens* occur by transfer of DNA from the bacterium to the plant. Although the tumors do not kill the plants, they make the plants stunted and more susceptible to harsh environmental conditions. Many plants, such as walnuts, grapes, nut trees, and beets, are affected by *A. tumefaciens*. The artificial introduction of DNA into plant cells is more challenging than in animal cells because of the thick plant cell wall.

Researchers used the natural transfer of DNA from *Agrobacterium* to a plant host to introduce DNA fragments of their choice into plant hosts. In nature, the disease-causing *A. tumefaciens* have a set of plasmids, called the Ti plasmids (tumor-inducing plasmids), that contain genes for the production of tumors in plants. DNA from the Ti plasmid integrates into the infected plant cell's genome. Researchers manipulate the Ti plasmids to remove the tumor-causing genes and insert the desired DNA fragment for transfer into the plant genome. The Ti plasmids carry antibiotic resistance genes to aid selection and can be propagated in *E. coli* cells as well.

### Organic Insecticide *Bacillus thuringiensis*

*Bacillus thuringiensis* (Bt) is a bacterium that produces protein crystals during sporulation that are toxic to many insect species that affect plants. Bt toxin has to be ingested by insects for the toxin to be activated. Insects that have eaten Bt toxin stop feeding on the plants within a few hours. After the toxin is activated in the intestines of the insects, death occurs within a couple of days. Modern biotechnology has allowed plants to encode their own crystal Bt toxin that acts against insects. The crystal toxin genes have been cloned from Bt and introduced into plants. Bt toxin has been found to be safe for the environment, non-toxic to humans and other mammals, and is approved for use by organic farmers as a natural insecticide.

### Flavr Savr Tomato

The first GM crop to be introduced into the market was the Flavr Savr Tomato produced in 1994. Antisense RNA technology was used to slow down the process of softening and rotting caused by fungal infections, which led to increased shelf life of the GM tomatoes. Additional genetic modification improved the flavor of this tomato. The Flavr Savr tomato did not successfully stay in the market because of problems maintaining and shipping the crop. However, since that time numerous crop plants have been developed and approved for sale and consumption. Corn, soybeans, and cotton in particular have been widely adopted by U.S. farmers.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

16.9: Agricultural Biotechnology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.10: Biotechnology at the Cutting Edge

This video provides an introduction to some of the newest breakthroughs in biotechnology: biofuel, health, and agriculture.

A link to an interactive elements can be found at the bottom of this page.

## Contributors and Attributions

All rights reserved content

- Biotechnology at the Cutting Edge. **Authored by**: National Museum of American History. **Located at**: https://youtu.be/ebSWG1QmxnY. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

16.10: Biotechnology at the Cutting Edge is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.11: Introduction to Risks and Benefits of Genomic Science

## Discuss the risks and benefits involved in applications of genetic and genomic science

Scientific developments can be used in a lot of different ways that impact our lives. In this outcome, we'll learn about the risks and benefits of genomic science.

## What You'll Learn to Do

- Outline how genetic information is used in personal identification
- Discuss medical uses of genetic information and the potential benefits and risks of this
- Outline the potential benefits and risks associated with agricultural uses of biotechnology

### Learning Activities

The learning activities for this section include the following:

- Genetic Information Used for Identification
- Medical Uses of Genetic Information
- Genomics in Agriculture
- Self Check: Risks and Benefits of Genomic Science

## Contributors and Attributions

CC licensed content, Original

- **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

16.11: Introduction to Risks and Benefits of Genomic Science is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.12: Genetic Information Used for Identification

## DNA as a Forensic Tool

Information and clues obtained from DNA samples found at crime scenes have been used as evidence in court cases, and genetic markers have been used in forensic analysis. Genomic analysis has also become useful in this field. In 2001, the first use of genomics in forensics was published. It was a collaborative attempt between academic research institutions and the FBI to solve the mysterious cases of anthrax communicated via the US Postal Service. Using microbial genomics, researchers determined that a specific strain of anthrax was used in all the mailings.

## Mitochondrial Genomics

Mitochondria are intracellular organelles that contain their own DNA. Mitochondrial DNA mutates at a rapid rate and is often used to study evolutionary relationships. Another feature that makes studying the mitochondrial genome interesting is that the mitochondrial DNA in most multicellular organisms is passed on only from the mother during the process of fertilization. For this reason, mitochondrial genomics is often used to trace genealogy.

## DNA Fingerprinting


**DNA fingerprinting** (also called DNA profiling, DNA testing, or DNA typing) is a forensic technique used to identify individuals by characteristics of their DNA. A DNA profile is a small set of DNA variations that is very likely to be different in all unrelated individuals, thereby being as unique to individuals as are fingerprints (hence the name for the technique).

Although 99.9% of human DNA sequences are the same in every person, enough of the DNA is different that it is possible to distinguish one individual from another, unless they are monozygotic (“identical”) twins. DNA fingerprinting uses repetitive sequences that are highly variable, called variable number tandem repeats (VNTRs). Modern law enforcement in particular uses short tandem repeats (STRs). STR loci are very similar between closely related individuals, but are so variable that unrelated individuals are extremely unlikely to have the same STRs. The combination of STRs used by law enforcement enable identification though because even closely related individuals will not share all the same STR loci.

The modern process of DNA fingerprinting was developed in 1984 by Sir Alec Jeffreys, while he was working in the Department of Genetics at the University of Leicester. DNA fingerprinting can be used to identify a person or to place a person at a crime scene and to help clarify paternity. DNA fingerprinting has also been widely used in the study of animal and floral populations and has revolutionized the fields of zoology, botany, and agriculture.

### Learning Objectives

Watch this video on the process of DNA fingerprinting and DNA profiling




A YouTube element has been excluded from this version of the text. You can view it online here: pb.libretexts.org/biom1/?p=582

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8
- Modification of DNA profiling. **Provided by**: Wikipedia. **Located at**: https://en.Wikipedia.org/wiki/DNA_profiling. **License**: *CC BY-SA: Attribution-ShareAlike*

All rights reserved content

- DNA Fingerprinting. **Authored by**: Bozeman Science. **Located at**: https://youtu.be/DbR9xMXuK7c. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

16.12: Genetic Information Used for Identification is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.13: Medical Uses of Genetic Information

Watch this video and consider whether you would be interested in knowing details about your own personal disease risk or susceptibility.

A link to an interactive elements can be found at the bottom of this page.

## Predicting Disease Risk at the Individual Level

Predicting the risk of disease involves screening currently healthy individuals by genome analysis at the individual level. Intervention with lifestyle changes and drugs can be recommended before disease onset. However, this approach is most applicable when the problem resides within a single gene defect. Such defects only account for approximately 5 percent of diseases in developed countries. Most of the common diseases, such as heart disease, are multi-factored or polygenic, which is a phenotypic characteristic that involves two or more genes, and also involve environmental factors such as diet. In April 2010, scientists at Stanford University published the genome analysis of a healthy individual (Stephen Quake, a scientist at Stanford University, who had his genome sequenced); the analysis predicted his propensity to acquire various diseases. A risk assessment was performed to analyze Quake's percentage of risk for 55 different medical conditions. A rare genetic mutation was found, which showed him to be at risk for sudden heart attack. He was also predicted to have a 23 percent risk of developing prostate cancer and a 1.4 percent risk of developing Alzheimer's. The scientists used databases and several publications to analyze the genomic data. Even though genomic sequencing is becoming more affordable and analytical tools are becoming more reliable, ethical issues surrounding genomic analysis at a population level remain to be addressed.

Debate remains over what to do with individual level data as well, such as the data from the genomic analysis of Quake's DNA. As a result of the study it was recommended that Quake start a regiment of preventative statins; the long-term effects of this study or treatment remain unknown at this stage.





Figure 1. *PCA3* is a gene that is expressed in prostate epithelial cells and overexpressed in cancerous cells. A high concentration of *PCA3* in urine is indicative of prostate cancer. The *PCA3* test is considered to be a better indicator of cancer than the more well know PSA test, which measures the level of PSA (prostate-specific antigen) in the blood.

For example, in 2011, the United States Preventative Services Task Force recommended against using the PSA test to screen healthy men for prostate cancer. Their recommendation is based on evidence that screening does not reduce the risk of death from prostate cancer. Prostate cancer often develops very slowly and does not cause problems, while the cancer treatment can have severe side effects. The *PCA3* (Figure 1) test is considered to be more accurate, but screening may still result in men who would not have been harmed by the cancer itself suffering side effects from treatment.

What do you think? Should all healthy men be screened for prostate cancer using the *PCA3* or PSA test? Should people in general be screened to find out if they have a genetic risk for cancer or other diseases?

There are no right or wrong answers to these questions. While it is true that prostate cancer treatment itself can be harmful, many men would rather be aware that they have cancer so they can monitor the disease and begin treatment if it progresses. And while genetic screening may be useful, it is expensive and may cause needless worry. People with certain risk factors may never develop the disease, and preventative treatments may do more harm than good.

## Pharmacogenomics and Toxicogenomics

Pharmacogenomics, also called toxicogenomics, involves evaluating the effectiveness and safety of drugs on the basis of information from an individual's genomic sequence. Genomic responses to drugs can be studied using experimental animals (such as laboratory rats or mice) or live cells in the laboratory before embarking on studies with humans. Studying changes in gene expression could provide information about the transcription profile in the presence of the drug, which can be used as an early indicator of the potential for toxic effects. For example, genes involved in cellular growth and controlled cell death, when disturbed, could lead to the growth of cancerous cells. Genome-wide studies can also help to find new genes involved in drug toxicity. Personal genome sequence information can be used to prescribe medications that will be most effective and least toxic on the basis of the individual patient's genotype. The gene signatures may not be completely accurate, but can be tested further before pathologic symptoms arise.

## Microbial Genomics

### Metagenomics





Figure 2. Metagenomics involves isolating DNA from multiple species within an environmental niche.

Traditionally, microbiology has been taught with the view that microorganisms are best studied under pure culture conditions, which involves isolating a single type of cell and culturing it in the laboratory. Because microorganisms can go through several generations in a matter of hours, their gene expression profiles adapt to the new laboratory environment very quickly. In addition, the vast majority of bacterial species resist being cultured in isolation. Most microorganisms do not live as isolated entities, but in microbial communities known as biofilms. For all of these reasons, pure culture is not always the best way to study microorganisms. Metagenomics is the study of the collective genomes of multiple species that grow and interact in an environmental niche. Metagenomics can be used to identify new species more rapidly and to analyze the effect of pollutants on the environment (Figure 2).

### Multitude of Benefits

Knowledge of the genomics of microorganisms is being used to find better ways to harness biofuels from algae and cyanobacteria. The primary sources of fuel today are coal, oil, wood, and other plant products, such as ethanol. Although plants are renewable resources, there is still a need to find more alternative renewable sources of energy to meet our population's energy demands. The microbial world is one of the largest resources for genes that encode new enzymes and produce new organic compounds, and it remains largely untapped. Microorganisms are used to create products, such as enzymes that are used in research, antibiotics, and other anti-microbial mechanisms. Microbial genomics is helping to develop diagnostic tools, improved vaccines, new disease treatments, and advanced environmental cleanup techniques.

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- Personalized Medicine: Do You Want to Know?. **Authored by**: World Science Festival. **Located at**: https://youtu.be/OON70d8gTnc. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

---

16.13: Medical Uses of Genetic Information is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.14: Genomics in Agriculture

Genomics can reduce the trials and failures involved in scientific research to a certain extent, which could improve the quality and quantity of crop yields in agriculture. Linking traits to genes or gene signatures helps to improve crop breeding to generate hybrids with the most desirable qualities. Scientists use genomic data to identify desirable traits, and then transfer those traits to a different organism. Scientists are discovering how genomics can improve the quality and quantity of agricultural production. For example, scientists could use desirable traits to create a useful product or enhance an existing product, such as making a drought-sensitive crop more tolerant of the dry season.

## GMO Controversies: Science versus Public Fear

Watch Borut Bohanec, the Chair of the Department of Agronomy at the University of Ljubljana (in Slovenia), as he discusses the fears and potential benefits surrounding GMOs.

A link to an interactive elements can be found at the bottom of this page.

You can view the transcript for “GMO Controversies – Science vs. Public Fear” here (opens in new window).

## Contributors and Attributions

CC licensed content, Shared previously

- Biology. **Provided by**: OpenStax CNX. **Located at**: http://cnx.org/contents/185cbf87-c72e-48f5-b51e-f14f21b5eabd@10.8. **License**: *CC BY: Attribution*. **License Terms**: Download for free at http://cnx.org/contents/185cbf87-c72...f21b5eabd@10.8

All rights reserved content

- GMO controversies - science vs. public fear: Borut Bohanec at TEDxLjubljana. **Authored by**: TEDx Talks. **Located at**: https://youtu.be/mz4_TwdaYeI. **License**: *All Rights Reserved*. **License Terms**: Standard YouTube License

16.14: Genomics in Agriculture is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.15: Putting It Together- Modern Biology



Figure 1. A photo of Michael Blassie in his Air Force Academy Cadet Uniform.

Let's return to our discussion of the unknown solider we discussed at the beginning of the module. First Lieutenant Michael Joseph Blassie, 24, was shot down over South Vietnam in 1972 and presumed dead.

In the early 1990s, Blassie's family received word that his remains might be buried in the Tomb of the Unknown. They petitioned the Department of Defense to open the site and conduct DNA testing, a technology that had been unavailable when the remains were first brought to and buried in the Tomb of the Unknown. In 1998, the Tomb of the Unknown was opened and the remains of the Vietnam Unknown—previously identified as X-26—were removed.

Forensic anthropologists took the aged and damaged samples of bone for mitochondrial DNA (mtDNA) testing. Because mtDNA is passed along the maternal line, scientists compared the Unknown Soldier's DNA against two samples submitted by First Lieutenant Blassie's mother and sister and found a match.

On July 11, 1998, 1st Lt. Michael Blassie was buried with full military honors in Jefferson National Cemetery, Missouri, near his hometown, in the same cemetery as his father.

## Contributors and Attributions

CC licensed content, Original

- Putting It Together: Modern Biology. **Authored by**: Shelli Carter and Lumen Learning. **Provided by**: Lumen Learning. **License**: *CC BY: Attribution*

Public domain content

- Michael Blassie. **Provided by**: US Airforce. **Located at**: https://en.Wikipedia.org/wiki/File:MichaelBlassie.jpg. **License**: *Public Domain: No Known Copyright*
- Michael Blassie unknown no more. **Provided by**: National Institutes of Health. **Located at**: https://www.nlm.nih.gov/exhibition/visibleproofs/galleries/cases/blassie.html. **Project**: Visible Proofs. **License**: *Public Domain: No Known Copyright*

16.15: Putting It Together- Modern Biology is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# 16.16: Assignment- When Jabberjays Attack!

Read this New York Times article by author James Gorman which will introduce you to the idea of DIY biology. After reading it proceed with the activities below.

- **Part 1:** Using your understanding of DNA technology and evolution, summarize changes that would need to take place in order to create organisms such as the Jabberjay and mockingjay.
- **Part 2:** Conduct further research into the idea of DIY biology and genetic engineering. Briefly summarize your findings and provide a specific bioengineering example. Cite references using MLA format.
- **Part 3:** State your opinion of the DIYbio movement and whether you believe government regulation or publishing restrictions should be put into place. Why or why not? Provide specific evidence and a logical argument to support your opinion. Simply stating you 'feel' or 'believe' something is insufficient justification.

Your essay should be prepared and cited using MLA format. Submit your final work as a single document.

Use this document to find some helpful hints for completing this assignment.

Basic Requirements (the assignment will not be accepted or assessed unless the follow criteria have been met):

- Assignment has been proofread and does not contain any major spelling or grammatical errors
- Assignment includes appropriate references.
- Assignment is formatted throughout using MLA style.

## Rubric

| When Jabberjays Attack! |
| --- |
| Outcome: Explain the theory of evolution, which documents the change in the genetic makeup of a biological population over time. Describe and discuss techniques used in modern biology. |

| When Jabberjays Attack! | | | | |
|---|---|---|---|---|
| Criteria | Ratings | | | Pts |
| Jabberjay creation discussed referencing specific DNA technologies possibly used for their engineering. | DNA technologies identified and discussed in detail that could have created jabberjays.<br>5.0 pts | DNA technologies identified that could have created jabberjays.<br>4.0 pts | Incorrect DNA technologies discussed or not sufficiently justified in discussion.<br>0.0 pts | 5 pts |
| Jabberjay and mockingjay evolutionary relationships discussed in order to highlight the way that adaptations enhance the survival and reproduction of individuals in a population. | Evolutionary relationship of jabberjays/mockingjays discussed correctly and in detail.<br>5.0 pts | Evolutionary relationship of jabberjays/mockingjays discussed correctly.<br>4.0 pts | Jabberjay/mockingjay relationship not logical or insufficiently discussed.<br>0.0 pts | 5 pts |
| Discuss mockingjay survival as an example of natural selection. | Natural selection discussion accurate and detailed.<br>5.0 pts | Natural selection discussion accurate.<br>4.0 pts | Natural selection discussion incorrect or illogical.<br>0.0 pts | 5 pts |
| DIY biology and specific bioengineering example discussed in order to highlight societal use of biotechnology | DIY biology and specific example used to highlight societal uses of biotechnology and potential scenarios provided illustrating pluses/minuses<br>5.0 pts | DIY biology and specific example used to highlight societal uses of biotechnology<br>4.0 pts | DIY biology and specific example not used to highlight societal uses of biotechnology<br>0.0 pts | 5 pts |
| Paper includes personal opinion on gene engineering, supported by evidence beyond simple opinion, that discusses possible risks and benefits of the application genomic and genetic science. | Personal opinion well supported by evidence, example, and/or scenarios.<br>5.0 pts | Personal opinion supported by at least one full scenario or example.<br>4.0 pts | Personal opinion not fully supported.<br>0.0 pts | 5 pts |
| | | | | Total points: 25 |

## Contributors and Attributions

CC licensed content, Shared previously

- 'Hunger Games' Science: Investigating Genetically Engineered Organisms. **Authored by**: Jeff Gerlach. **Located at**: https://www.oercommons.org/authoring/4514-hunger-games-science-investigating-genetically-eng/view. **License**: *CC BY: Attribution*

16.16: Assignment- When Jabberjays Attack! is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

# Index

# Glossary

**Sample Word 1** | Sample Definition 1

# Detailed Licensing

## Overview

**Title:** Biology for Majors I (Lumen)

**Webpages:** 296

**All licenses found:**

- Undeclared: 100% (296 pages)

## By Page


- Biology for Majors I (Lumen) - *Undeclared*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Course Contents - *Undeclared*
    - 1.1: About This Course - *Undeclared*
    - 1.2: Course Contents at a Glance - *Undeclared*
    - 1.3: Learning Outcomes - *Undeclared*
  - 2: Faculty Resources - *Undeclared*
    - 2.1: Faculty Resources Overview - *Undeclared*
    - 2.2: Pacing - *Undeclared*
    - 2.3: PDF - *Undeclared*
    - 2.4: PowerPoints - *Undeclared*
    - 2.5: Assignments - *Undeclared*
    - 2.6: Assignments with Samples - *Undeclared*
    - 2.7: Question Banks - *Undeclared*
    - 2.8: I Need Help - *Undeclared*
  - 3: Module 1- Introduction to Biology - *Undeclared*
    - 3.1: Why It Matters- Introduction to Biology - *Undeclared*
    - 3.2: Introduction to Characteristics of Life - *Undeclared*
    - 3.3: Properties of Life - *Undeclared*
    - 3.4: Introduction to Prokaryotes and Eukaryotes - *Undeclared*
    - 3.5: Comparing Prokaryotic and Eukaryotic Cells - *Undeclared*
    - 3.6: Levels of Organization of Living Things - *Undeclared*
    - 3.7: Introduction to Taxonomy - *Undeclared*
    - 3.8: The Diversity of Life - *Undeclared*
    - 3.9: Phylogenetic Trees - *Undeclared*
    - 3.10: Taxonomy - *Undeclared*
    - 3.11: Introduction to the Study of Biology - *Undeclared*
    - 3.12: The Branches of Biology - *Undeclared*
    - 3.13: Scientific Inquiry - *Undeclared*
    - 3.14: Experiments and Hypotheses - *Undeclared*
    - 3.15: Interpreting Results - *Undeclared*
    - 3.16: Putting It Together- Introduction to Biology - *Undeclared*
  - 4: Module 2- Chemistry of Life - *Undeclared*
    - 4.1: Why It Matters- The Chemistry of Life - *Undeclared*
    - 4.2: Introduction to Atoms and Elements - *Undeclared*
    - 4.3: Atoms - *Undeclared*
    - 4.4: The Periodic Table of Elements - *Undeclared*
    - 4.5: Electrons - *Undeclared*
    - 4.6: Introduction to Atomic Bonds - *Undeclared*
    - 4.7: Ionic Bonds - *Undeclared*
    - 4.8: Covalent Bonds - *Undeclared*
    - 4.9: Hydrogen Bonds - *Undeclared*
    - 4.10: Introduction to Functional Groups - *Undeclared*
    - 4.11: Functional Groups - *Undeclared*
    - 4.12: Introduction to The pH Scale - *Undeclared*
    - 4.13: Acids and Bases - *Undeclared*
    - 4.14: Buffers - *Undeclared*
    - 4.15: Introduction to Chemical Reactions - *Undeclared*
    - 4.16: Chemical Reactions and Molecules - *Undeclared*
    - 4.17: Anabolic and Catabolic Pathways - *Undeclared*
    - 4.18: Enzymes - *Undeclared*
    - 4.19: Introduction to Thermodynamics - *Undeclared*
    - 4.20: Open and Closed Systems - *Undeclared*
    - 4.21: The First Law of Thermodynamics - *Undeclared*
    - 4.22: The Second Law of Thermodynamics - *Undeclared*
    - 4.23: Putting It Together- Chemistry of Life - *Undeclared*
    - 4.24: Assignment- Biological Astronaut - *Undeclared*

- 4.25: Assignment- Interacting with Attributes of Molecules - *Undeclared*

- 5: Module 3- Important Biological Macromolecules - *Undeclared*
  - 5.1: Why It Matters- Important Biological Macromolecules - *Undeclared*
  - 5.2: Introduction to Proteins - *Undeclared*
  - 5.3: Amino Acids - *Undeclared*
  - 5.4: Protein Structure - *Undeclared*
  - 5.5: Function of Proteins - *Undeclared*
  - 5.6: Introduction to Lipids - *Undeclared*
  - 5.7: Lipids - *Undeclared*
  - 5.8: Introduction to Carbohydrates - *Undeclared*
  - 5.9: Structure and Function of Carbohydrates - *Undeclared*
  - 5.10: Introduction to Nucleic Acids - *Undeclared*
  - 5.11: Structure of Nucleic Acids - *Undeclared*
  - 5.12: DNA and RNA - *Undeclared*
  - 5.13: Introduction to Comparing Biological Macromolecules - *Undeclared*
  - 5.14: Different Types of Biological Macromolecules - *Undeclared*
  - 5.15: Putting It Together- Important Biological Macromolecules - *Undeclared*
  - 5.16: Assignment- Nutritionist for a Day - *Undeclared*
- 6: Module 4- Cellular Structure - *Undeclared*
  - 6.1: Why It Matters- Cellular Structure - *Undeclared*
  - 6.2: Introduction to Microscopes - *Undeclared*
  - 6.3: Microscopy - *Undeclared*
  - 6.4: Introduction to Organelles - *Undeclared*
  - 6.5: Cytoplasm - *Undeclared*
  - 6.6: Nucleus - *Undeclared*
  - 6.7: Ribosomes, Mitochondria, and Peroxisomes - *Undeclared*
  - 6.8: Endomembrane System - *Undeclared*
  - 6.9: Vesicles - *Undeclared*
  - 6.10: Unique Features of Animal and Plant Cells - *Undeclared*
  - 6.11: Introduction to Cytoskeleton - *Undeclared*
  - 6.12: Microfilaments - *Undeclared*
  - 6.13: Intermediate Filaments - *Undeclared*
  - 6.14: Microtubules - *Undeclared*
  - 6.15: Introduction to Cell Surface - *Undeclared*
  - 6.16: Cell Membranes - *Undeclared*
  - 6.17: Cell Junctions - *Undeclared*
  - 6.18: The Cell Wall - *Undeclared*
  - 6.19: Putting It Together- Cellular Structure - *Undeclared*
  - 6.20: Assignment- Cell Builder - *Undeclared*
- 7: Module 5- Cell Membranes - *Undeclared*
  - 7.1: Why It Matters- Cell Membranes - *Undeclared*
  - 7.2: Introduction to Structure of the Membrane - *Undeclared*
  - 7.3: Structure of the Cell Membrane - *Undeclared*
  - 7.4: Phospholipids - *Undeclared*
  - 7.5: Introduction to Kinds of Transport - *Undeclared*
  - 7.6: Passive Transport - *Undeclared*
  - 7.7: Active Transport - *Undeclared*
  - 7.8: Membranes and Transport - *Undeclared*
  - 7.9: Introduction to Endocytosis and Exocytosis - *Undeclared*
  - 7.10: Endocytosis - *Undeclared*
  - 7.11: Exocytosis - *Undeclared*
  - 7.12: Putting It Together- Cell Membranes - *Undeclared*
  - 7.13: Assignment- Membranes Alive! - *Undeclared*
- 8: Module 6- Metabolic Pathways - *Undeclared*
  - 8.1: Why It Matters- Metabolic Pathways - *Undeclared*
  - 8.2: Introduction to Redox Reactions - *Undeclared*
  - 8.3: Electrons and Energy - *Undeclared*
  - 8.4: ATP in Living Systems - *Undeclared*
  - 8.5: Introduction to Photosynthesis - *Undeclared*
  - 8.6: An Overview of Photosynthesis - *Undeclared*
  - 8.7: Spectrums of Light - *Undeclared*
  - 8.8: Light-Dependent Reactions - *Undeclared*
  - 8.9: Light-Independent Reactions - *Undeclared*
  - 8.10: Photosynthesis - *Undeclared*
  - 8.11: Introduction to Cellular Respiration - *Undeclared*
  - 8.12: Glycolysis - *Undeclared*
  - 8.13: Pyruvate Oxidation - *Undeclared*
  - 8.14: Citric Acid Cycle - *Undeclared*
  - 8.15: Electron Transport Chain - *Undeclared*
  - 8.16: Cellular Respiration - *Undeclared*
  - 8.17: Introduction to Fermentation - *Undeclared*
  - 8.18: Metabolism without Oxygen - *Undeclared*
  - 8.19: Types of Fermentation - *Undeclared*
  - 8.20: Putting It Together- Metabolic Pathways - *Undeclared*
  - 8.21: Assignment- Observing Energy Transactions - *Undeclared*
- 9: Module 7- Cell Communication - *Undeclared*
  - 9.1: Why It Matters- Cell Communication - *Undeclared*
  - 9.2: Introduction to Signaling Molecules and Cellular Receptors - *Undeclared*

- 9.3: Types of Signals - *Undeclared*
- 9.4: Signaling Molecules - *Undeclared*
- 9.5: Signaling Receptors - *Undeclared*
- 9.6: Introduction to Propagation of the Signal - *Undeclared*
- 9.7: Binding Initiates a Signaling Pathway - *Undeclared*
- 9.8: Methods of Intracellular Signaling - *Undeclared*
- 9.9: Introduction to Response to the Signal - *Undeclared*
- 9.10: Expression and Growth - *Undeclared*
- 9.11: Cell Death - *Undeclared*
- 9.12: Introduction to Signaling in Single-Celled Organisms - *Undeclared*
- 9.13: Signaling in Yeast - *Undeclared*
- 9.14: Signaling in Bacteria - *Undeclared*
- 9.15: Putting It Together- Cell Communication - *Undeclared*

10: Module 8- Cell Division - *Undeclared*

- 10.1: Why It Matters- Cell Division - *Undeclared*
- 10.2: Introduction to Chromosomes and DNA Packaging - *Undeclared*
- 10.3: DNA and Chromosomes - *Undeclared*
- 10.4: Chromosome Structure - *Undeclared*
- 10.5: Introduction to the Cell Cycle - *Undeclared*
- 10.6: Interphase - *Undeclared*
- 10.7: Mitosis - *Undeclared*
- 10.8: Cytokinesis - *Undeclared*
- 10.9: The Complete Cycle - *Undeclared*
- 10.10: Introduction to Cell Cycle Checkpoints - *Undeclared*
- 10.11: Control of the Cell Cycle - *Undeclared*
- 10.12: Cancer and the Cell Cycle - *Undeclared*
- 10.13: Introduction to Meiosis - *Undeclared*
- 10.14: Stages of Meiosis - *Undeclared*
- 10.15: Meiosis I - *Undeclared*
- 10.16: Meiosis II - *Undeclared*
- 10.17: Meiosis - *Undeclared*
- 10.18: Introduction to Genetic Diversity - *Undeclared*
- 10.19: Sexual Reproduction - *Undeclared*
- 10.20: Genetic Variation in Meiosis - *Undeclared*
- 10.21: Chromosomal Structural Rearrangements - *Undeclared*
- 10.22: Mitosis, Meiosis, and Sexual Reproduction - *Undeclared*
- 10.23: Introduction to Errors in Chromosome Number - *Undeclared*
- 10.24: Karyotypes - *Undeclared*
- 10.25: Common Disorders - *Undeclared*
- 10.26: Putting It Together- Cell Division - *Undeclared*
- 10.27: Assignment- Mitosis and Meiosis Worksheets - *Undeclared*

11: Module 9- DNA Structure and Replication - *Undeclared*

- 11.1: Why It Matters- DNA Structure and Replication - *Undeclared*
- 11.2: Introduction to Storing Genetic Information - *Undeclared*
- 11.3: Structure of DNA - *Undeclared*
- 11.4: Genetic Information - *Undeclared*
- 11.5: DNA Packaging in Eukaryotes and Prokaryotes - *Undeclared*
- 11.6: Introduction to DNA Base Pairs and Replication - *Undeclared*
- 11.7: The History of DNA - *Undeclared*
- 11.8: Basics of DNA Replication - *Undeclared*
- 11.9: Major Enzymes - *Undeclared*
- 11.10: “Proofreading” DNA - *Undeclared*
- 11.11: Telomeres - *Undeclared*
- 11.12: Introduction to DNA Mutations - *Undeclared*
- 11.13: What is a Mutation? - *Undeclared*
- 11.14: Mutations in Somatic Cells and in Gametes - *Undeclared*
- 11.15: Major Types of Mutations - *Undeclared*
- 11.16: Putting It Together- DNA Structure and Replication - *Undeclared*
- 11.17: Assignment- Which Has More DNA? - *Undeclared*

12: Module 10- DNA Transcription and Translation - *Undeclared*

- 12.1: Why It Matters- DNA Transcription and Translation - *Undeclared*
- 12.2: Introduction to Transcription - *Undeclared*
- 12.3: Steps of Genetic Transcription - *Undeclared*
- 12.4: RNA Polymerase - *Undeclared*
- 12.5: pre-RNA and mRNA - *Undeclared*
- 12.6: RNA Post-Translational Modification - *Undeclared*
- 12.7: Introduction to Translation - *Undeclared*
- 12.8: Steps of Translation - *Undeclared*
- 12.9: Ribosomes - *Undeclared*
- 12.10: tRNA - *Undeclared*
- 12.11: Codons - *Undeclared*
- 12.12: Protein Modifications - *Undeclared*
- 12.13: Introduction to Prokaryotic Transcription and Translation - *Undeclared*
- 12.14: Prokaryotic Transcription - *Undeclared*

- 12.15: Prokaryotic Translation - *Undeclared*
- 12.16: Introduction to the Central Dogma - *Undeclared*
- 12.17: The Central Dogma - *Undeclared*
- 12.18: Putting It Together- DNA Transcription and Translation - *Undeclared*

13: Module 11- Gene Expression - *Undeclared*

- 13.1: Why It Matters- Gene Expression - *Undeclared*
- 13.2: Introduction to Regulation of Gene Expression - *Undeclared*
- 13.3: Expression of Genes - *Undeclared*
- 13.4: Prokaryotic and Eukaryotic Gene Regulation - *Undeclared*
- 13.5: Introduction to Prokaryotic Gene Regulation - *Undeclared*
- 13.6: Gene Regulation in Prokaryotes - *Undeclared*
- 13.7: Repressors - *Undeclared*
- 13.8: Activators and Inducers - *Undeclared*
- 13.9: Prokaryotic Gene Regulation at Work - *Undeclared*
- 13.10: Introduction to Eukaryotic Gene Regulation - *Undeclared*
- 13.11: Eukaryotic Epigenetic Gene Regulation - *Undeclared*
- 13.12: Eukaryotic Transcription Gene Regulation - *Undeclared*
- 13.13: Post-Transcriptional Control of Gene Expression - *Undeclared*
- 13.14: Putting It Together- Gene Expression - *Undeclared*
- 13.15: Assignment- How Mutations Work - *Undeclared*

14: Module 12- Trait Inheritance - *Undeclared*

- 14.1: Why It Matters- Trait Inheritance - *Undeclared*
- 14.2: Introduction to the Father of Genetics - *Undeclared*
- 14.3: Mendel's Experiments and Heredity - *Undeclared*
- 14.4: Characteristics and Traits - *Undeclared*
- 14.5: Laws of Inheritance - *Undeclared*
- 14.6: Heredity - *Undeclared*
- 14.7: Introduction to Beyond Dominance and Recessiveness - *Undeclared*
- 14.8: Non-Mendelian Inheritance - *Undeclared*
- 14.9: Non-Mendelian Punnett Squares - *Undeclared*
- 14.10: Multiple Alleles - *Undeclared*
- 14.11: Introduction to Heredity and Disease - *Undeclared*
- 14.12: Penetrance and Expressivity - *Undeclared*
- 14.13: Pedigrees and Disease - *Undeclared*
- 14.14: Pedigree Flowchart - *Undeclared*
- 14.15: Genetic Disorder and Pedigrees - *Undeclared*
- 14.16: Introduction to Genetics and the Environment - *Undeclared*
- 14.17: Epistasis - *Undeclared*
- 14.18: Polygenic Inheritance and Environmental Effects - *Undeclared*
- 14.19: Effect of the Environment - *Undeclared*
- 14.20: Pleiotropy and Human Disorders - *Undeclared*
- 14.21: Putting It Together- Trait Inheritance - *Undeclared*
- 14.22: Assignment- Dragon Genetics - *Undeclared*

15: Module 13- Theory of Evolution - *Undeclared*

- 15.1: Why It Matters- Theory of Evolution - *Undeclared*
- 15.2: Introduction to Charles Darwin - *Undeclared*
- 15.3: Darwin and Descent with Modification - *Undeclared*
- 15.4: Darwin and the Theory of Evolution - *Undeclared*
- 15.5: Natural Selection - *Undeclared*
- 15.6: Introduction to Evidence for Evolution - *Undeclared*
- 15.7: Physical Evidence - *Undeclared*
- 15.8: Biological Evidence - *Undeclared*
- 15.9: Misconceptions of Evolution - *Undeclared*
- 15.10: Introduction to Mutations and Evolution - *Undeclared*
- 15.11: Population Genetics - *Undeclared*
- 15.12: Selective and Environmental Pressures - *Undeclared*
- 15.13: Introduction to Phylogenetic Trees - *Undeclared*
- 15.14: Scientific Classification - *Undeclared*
- 15.15: Structure of Phylogenetic Trees - *Undeclared*
- 15.16: Limitations of Phylogenetic Trees - *Undeclared*
- 15.17: The Taxonomic Classification System - *Undeclared*
- 15.18: Putting It Together- Theory of Evolution - *Undeclared*

16: Module 14- Modern Biology - *Undeclared*

- 16.1: Why It Matters- Modern Biology - *Undeclared*
- 16.2: Introduction to Key Technologies - *Undeclared*
- 16.3: Manipulating Genetic Material - *Undeclared*
- 16.4: DNA Sequencing - *Undeclared*
- 16.5: Cloning - *Undeclared*
- 16.6: Genetic Engineering - *Undeclared*

- 16.7: Introduction to Biotechnology Applications - *Undeclared*
- 16.8: Medicinal Biotechnology - *Undeclared*
- 16.9: Agricultural Biotechnology - *Undeclared*
- 16.10: Biotechnology at the Cutting Edge - *Undeclared*
- 16.11: Introduction to Risks and Benefits of Genomic Science - *Undeclared*
- 16.12: Genetic Information Used for Identification - *Undeclared*
- 16.13: Medical Uses of Genetic Information - *Undeclared*
- 16.14: Genomics in Agriculture - *Undeclared*
- 16.15: Putting It Together- Modern Biology - *Undeclared*
- 16.16: Assignment- When Jabberjays Attack! - *Undeclared*

- Back Matter - *Undeclared*
  - Index - *Undeclared*
  - Glossary - *Undeclared*
  - Detailed Licensing - *Undeclared*