CHE 125: GOB INTRODUCTORY CHEMISTRY



Erin O'Connell
UW-Milwaukee/Washington County &
Waukesha

LibreTexts™

CHE 125: GOB Introductory Chemistry

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide and accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 06/07/2025

TABLE OF CONTENTS

Licensing

1: Chemistry as a Science

- 1.1: The Scope of Chemistry
- 1.2: Chemicals Compose Ordinary Things
- 1.3: Hypothesis, Theories, and Laws
- 1.4: The Scientific Method - How Chemists Think

2: Measurements and Density

- 2.1: Taking Measurements
- 2.2: Scientific Notation - Writing Large and Small Numbers
- 2.3: Significant Figures - Writing Numbers to Reflect Precision
- 2.4: Significant Figures in Calculations
- 2.5: The Basic Units of Measurement
- 2.6: Problem Solving and Unit Conversions
- 2.7: Solving Multistep Conversion Problems
- 2.8: Units Raised to a Power
- 2.9: Density
- 2.10: Dosage Calculations

3: Matter

- 3.1: In Your Room
- 3.2: What is Matter?
- 3.3: Classifying Matter According to Its State- Solid, Liquid, and Gas
- 3.4: Classifying Matter According to Its Composition
- 3.5: Differences in Matter- Physical and Chemical Properties
- 3.6: Changes in Matter - Physical and Chemical Changes
- 3.7: Conservation of Mass - There is No New Matter
- 3.E: Chemistry, Matter, and Measurement (Exercises)

4: Atoms, Elements, and the Periodic Table

- 4.1: Prelude to Elements, Atoms, and the Periodic Table
- 4.2: The Elements
- 4.3: Atomic Theory
- 4.4: Atomic Theory
- 4.5: Defining Isotopes
- 4.6: Atomic Masses
- 4.7: Arrangements of Electrons
- 4.8: Orbital Shape
- 4.9: The Periodic Table
- 4.10: Ions - Losing and Gaining Electrons
- 4.E: Elements, Atoms, and the Periodic Table (Exercises)

5: Compounds- Ionic and Covalent

- 5.1: Sugar and Salt
- 5.2: Compounds Display Constant Composition
- 5.3: Chemical Formulas- How to Represent Compounds
- 5.4: A Molecular View of Elements and Compounds
- 5.5: Writing Formulas for Ionic Compounds
- 5.6: Nomenclature- Naming Compounds
- 5.7: Naming Ionic Compounds
- 5.8: Naming Molecular Compounds
- 5.9: Naming Acids
- 5.10: Nomenclature Summary
- 5.11: Formula Mass- The Mass of a Molecule or Formula Unit
- 5.12: Moles, Molecular Formulas, and Calculating Empirical Formulas for Compounds

6: Chemical Bonding

- 6.1: Representing Valence Electrons with Dots
- 6.2: Lewis Structures of Ionic Compounds- Electrons Transferred
- 6.3: Covalent Lewis Structures- Electrons Shared
- 6.4: Writing Lewis Structures for Covalent Compounds
- 6.5: Resonance - Equivalent Lewis Structures for the Same Molecule
- 6.6: Predicting the Shapes of Molecules
- 6.7: Electronegativity and Polarity - Why Oil and Water Don’t Mix

7: Chemical Reactions

- 7.1: Prelude to Introduction to Chemical Reactions
- 7.2: Laws of Conservation : A Review
- 7.3: Evidence of a Chemical Reaction
- 7.4: Chemical Equations
- 7.5: How to Write Balanced Chemical Equations
- 7.6: Aqueous Solutions and Solubility - Compounds Dissolved in Water
- 7.7: Precipitation Reactions
- 7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
- 7.9: Acid–Base and Gas Evolution Reactions
- 7.10: Oxidation-Reduction Reactions
- 7.11: Classifying Chemical Reactions
- 7.12: The Activity Series- Predicting Spontaneous Redox Reactions

8: Quantities in Chemical Reactions

- 8.1: Prelude to Quantities in Chemical Reactions
- 8.2: The Mole
- 8.3: Atomic and Molar Masses
- 8.4: Mole-Mass Conversions
- 8.5: Mole-Mole Relationships in Chemical Reactions
- 8.6: Mole-Mass and Mass-Mass Problems
- 8.E: Quantities in Chemical Reactions (Exercise)
- 8.S: Quantities in Chemical Reactions (Summary)

9: Solutions

- 9.1: Prelude to Solutions
- 9.2: Solutions
- 9.3: Concentration
- 9.4: The Dissolution Process
- 9.5: Properties of Solutions
- 9.E: Solutions (Exercises)
- 9.S: Solutions (Summary)

10: Equilibrium

- 10.1: The Rate of a Chemical Reaction
- 10.2: The Idea of Dynamic Chemical Equilibrium
- 10.3: The Equilibrium Constant - A Measure of How Far a Reaction Goes
- 10.4: Disturbing a Reaction at Equilibrium- Le Châtelier's Principle
- 10.5: The Effect of a Concentration Change on Equilibrium
- 10.6: The Effect of a Volume Change on Equilibrium
- 10.7: The Effect of Temperature Changes on Equilibrium
- 10.8: The Path of a Reaction and the Effect of a Catalyst

11: Energy and Chemical Processes

- 11.1: Prelude to Energy and Chemical Processes
- 11.2: Energy and Its Units
- 11.3: Heat
- 11.4: Phase Changes
- 11.5: Bond Energies and Chemical Reactions
- 11.6: The Energy of Biochemical Reactions
- 11.E: Energy and Chemical Processes (Exercises)
- 11.S: Energy and Chemical Processes (Summary)

12: Solids, Liquids, and Gases

- 12.1: A Molecular Comparison of Gases, Liquids, and Solids
- 12.2: Kinetic-Molecular Theory
- 12.3: The Ideal Gas Equation
- 12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases
- 12.5: Dalton's Law (Law of Partial Pressures)
- 12.6: Evaporation and Condensation
- 12.7: Intermolecular forces

13: Acids and Bases

- 13.1: Prelude to Acids and Bases
- 13.2: Arrhenius Definition of Acids and Bases
- 13.3: Brønsted-Lowry Definition of Acids and Bases
- 13.4: Water - Both an Acid and a Base
- 13.5: The Strengths of Acids and Bases
- 13.6: Buffers
- 13.E: Acids and Bases (Exercises)
- 13.S: Acids and Bases (Summary)

14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons

- 14.1: Organic Chemistry
- 14.2: Structures and Names of Alkanes
- 14.3: Alkenes- Structures and Names
- 14.4: Alkynes
- 14.5: Branched-Chain Alkanes
- 14.6: Condensed Structural and Line-Angle Formulas
- 14.7: IUPAC Nomenclature
- 14.8: Physical Properties of Alkanes
- 14.9: Physical Properties of Alkenes
- 14.10: Chemical Properties of Alkanes
- 14.11: Chemical Properties of Alkenes
- 14.12: Halogenated Hydrocarbons
- 14.13: Cycloalkanes
- 14.14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons
- 14.15: Organic Reactions
- 14.E: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Exercises)
- 14.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary)
- 14.S: Unsaturated and Aromatic Hydrocarbons (Summary)

Index

Index

Glossary

Detailed Licensing

Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

CHAPTER OVERVIEW

1: Chemistry as a Science

1.1: The Scope of Chemistry
1.2: Chemicals Compose Ordinary Things
1.3: Hypothesis, Theories, and Laws
1.4: The Scientific Method - How Chemists Think

1: Chemistry as a Science is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

1.1: The Scope of Chemistry

Learning Objectives

- To recognize the breadth, depth, and scope of chemistry.
- Define chemistry in relation to other sciences.
- Identify the main disciplines of chemistry.

Chemistry is the study of matter—what it consists of, what its properties are, and how it changes. Matter is anything that has mass and takes up space—that is, anything that is physically real. Some things are easily identified as matter—the screen on which you are reading this book, for example. Others are not so obvious. Because we move so easily through air, we sometimes forget that it, too, is matter. Because of this, chemistry is a science that has its fingers in just about everything. Being able to describe the ingredients in a cake and how they change when the cake is baked, for example, is chemistry!

Chemistry is one branch of science. Science is the process by which we learn about the natural universe by observing, testing, and then generating models that explain our observations. Because the physical universe is so vast, there are many different branches of science (Figure 1.1.1). Thus, chemistry is the study of matter, biology is the study of living things, and geology is the study of rocks and the earth. Mathematics is the language of science, and we will use it to communicate some of the ideas of chemistry.




Figure 1.1.1: The Relationships between Some of the Major Branches of Science. Chemistry lies more or less in the middle, which emphasizes its importance to many branches of science.

Although we divide science into different fields, there is much overlap among them. For example, some biologists and chemists work in both fields so much that their work is called biochemistry. Similarly, geology and chemistry overlap in the field called geochemistry. Figure 1.1.1 shows how many of the individual fields of science are related. At some level, all of these fields depend on matter because they all involve "stuff"; because of this, chemistry has been called the "central science", linking them all together.

> *There are many other fields of science, in addition to the ones (biology, medicine, etc.) listed here.*

Example 1.1.1: Science Fields

Which fields of study are branches of science? Explain.

a. sculpture
b. astronomy

Solution

a. Sculpture is not considered a science because it is not a study of some aspect of the natural universe.
b. Astronomy is the study of stars and planets, which are part of the natural universe. Astronomy is therefore a field of science.

? Exercise 1.1.1

Which fields of study are branches of science?

a. physiology (the study of the function of an animal's or a plant's body)
b. geophysics
c. agriculture
d. politics

Answer a:

yes

Answer b:

yes

Answer c:

yes

Answer d:

no

Areas of Chemistry

The study of modern chemistry has many branches, but can generally be broken down into five main disciplines, or areas of study:

- **Physical chemistry:** Physical chemistry is the study of macroscopic properties, atomic properties, and phenomena in chemical systems. A physical chemist may study such things as the rates of chemical reactions, the energy transfers that occur in reactions, or the physical structure of materials at the molecular level.
- **Organic chemistry:** Organic chemistry is the study of chemicals containing carbon. Carbon is one of the most abundant elements on Earth and is capable of forming a tremendously vast number of chemicals (over twenty million so far). Most of the chemicals found in all living organisms are based on carbon.
- **Inorganic chemistry:** Inorganic chemistry is the study of chemicals that, in general, are not primarily based on carbon. Inorganic chemicals are commonly found in rocks and minerals. One current important area of inorganic chemistry deals with the design and properties of materials involved in energy and information technology.
- **Analytical chemistry:** Analytical chemistry is the study of the composition of matter. It focuses on separating, identifying, and quantifying chemicals in samples of matter. An analytical chemist may use complex instruments to analyze an unknown material in order to determine its various components.
- **Biochemistry:** Biochemistry is the study of chemical processes that occur in living things. Research may cover anything from basic cellular processes up to understanding disease states so that better treatments can be developed.



Figure 1.1.2: (left) Measurement of trace metals using atomic spectroscopy. (right) Measurement of hormone concentrations.

In practice, chemical research is often not limited to just one of the five major disciplines. A particular chemist may use biochemistry to isolate a particular chemical found in the human body such as hemoglobin, the oxygen carrying component of red

blood cells. He or she may then proceed to analyze the hemoglobin using methods that would pertain to the areas of physical or analytical chemistry. Many chemists specialize in areas that are combinations of the main disciplines, such as bioinorganic chemistry or physical organic chemistry.

History of Chemistry

The **history** of chemistry is an interesting and challenging one. Very early chemists were often motivated mainly by the achievement of a specific goal or product. Making perfume or soaps did not need a lot of **theory**, just a good recipe and careful attention to detail. There was no standard way of naming materials (and no periodic table that we could all agree on). It is often difficult to figure out exactly what a particular person was using. However, the science developed over the centuries by trial and error.

Major progress was made toward putting chemistry on a solid foundation when Robert Boyle (1637-1691) began his **research** in chemistry (Figure 1.1.3). He developed the basic ideas about the behavior of gases. He could then describe gases mathematically. Boyle also helped form the idea that small particles could combine to form molecules. Many years later, John Dalton used these ideas to develop the atomic theory.



Figure 1.1.3: Robert Boyle.

The field of chemistry began to develop rapidly in the 1700's. Joseph Priestley (1733-1804) isolated and characterized several gases: oxygen, carbon monoxide, and nitrous oxide. It was later discovered that nitrous oxide ("laughing gas") worked as an anesthetic. This gas was used for that purpose for the first time in 1844 during a tooth extraction. Other gases discovered during that time were chlorine, by C.W. Scheele (1742-1786) and nitrogen, by Antoine Lavoisier (1743-1794). Lavoisier has been considered by many scholars to be the "father of chemistry". Among other accomplishments, he discovered the role of oxygen in combustion and definitively formulated the law of conservation of matter.

Chemists continued to discover new compounds in the 1800's. The science also began to develop a more theoretical foundation. John Dalton (1766-1844) put forth his atomic theory in 1807. This idea allowed scientists to think about chemistry in a much more systematic way. Amadeo Avogadro (1776-1856) laid the groundwork for a more quantitative approach to chemistry by calculating the number of particles in a given amount of a gas. A lot of effort was put forth in studying chemical reactions. These efforts led to new materials being produced. Following the invention of the battery by Alessandro Volta (1745-1827), the field of electrochemistry (both theoretical and applications) developed through major contributions by Humphry Davy (1778-1829) and Michael Faraday (1791-1867). Other areas of the discipline also progressed rapidly.

It would take a large book to cover developments in chemistry during the twentieth century and up to today. One major area of expansion was in the area of the chemistry of living processes. Research in photosynthesis in plants, the discovery and characterization of enzymes as biochemical catalysts, elucidation of the structures of biomolecules such as insulin and DNA—these efforts gave rise to an explosion of information in the field of biochemistry.

The practical aspects of chemistry were not ignored. The work of Volta, Davy, and Faraday eventually led to the development of batteries that provided a source of electricity to power a number of devices (Figure 1.1.4).




Figure 1.1.4: Battery developed by Volta. (CC BY-SA 3.0; (left) GuidoB and (right) Kkkdc).

Charles Goodyear (1800-1860) discovered the process of vulcanization, allowing a stable rubber product to be produced for the tires of all the vehicles we have today. Louis Pasteur (1822-1895) pioneered the use of heat sterilization to eliminate unwanted microorganisms in wine and milk. Alfred Nobel (1833-1896) invented dynamite (Figure 1.1.5). After his death, the fortune he made from this product was used to fund the Nobel Prizes in science and the humanities. J.W. Hyatt (1837-1920) developed the first plastic. Leo Baekeland (1863-1944) developed the first synthetic resin, widely used for inexpensive and sturdy dinnerware.



Figure 1.1.5: Dynamite explosion in Panama, Central America (1908).

Today, chemistry continues to be essential to the development of new materials and technologies, from semiconductors for electronics to powerful new medicines, and beyond.

Summary

- Chemistry is the study of matter and the changes it undergoes and considers both *macroscopic* and *microscopic* information.
- Matter is anything that has mass and occupies space.
- The five main disciplines of chemistry are physical chemistry, organic chemistry, inorganic chemistry, analytical chemistry and biochemistry.
- Many civilizations contributed to the growth of chemistry. A lot of early chemical research focused on practical uses. Basic chemistry theories were developed during the nineteenth century. New materials and batteries are a few of the products of modern chemistry.

1.1: The Scope of Chemistry is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **1.7: The Scope of Chemistry** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.2: Chemicals Compose Ordinary Things

Chemistry is the branch of science dealing with the structure, composition, properties, and the reactive characteristics of matter. Matter is anything that has mass and occupies space. Thus, chemistry is the study of literally everything around us – the liquids that we drink, the gasses we breathe, the composition of everything from the plastic case on your phone to the earth beneath your feet. Moreover, chemistry is the study of the transformation of matter. Crude oil is transformed into more useful petroleum products such as gasoline and kerosene by the process of refining. Some of these products are further transformed into plastics. Crude metal ores are transformed into metals, that can then be fashioned into everything from foil to automobiles. Potential drugs are identified from natural sources, isolated and then prepared in the laboratory. Their structures are systematically modified to produce the pharmaceuticals that have led to vast advances in modern medicine. Chemistry is at the center of all of these processes and chemists are the people that study the nature of matter and learn to design, predict and control these chemical transformations. Within the branches of chemistry you will find several apparent subdivisions. Inorganic chemistry, historically, focused on minerals and metals found in the earth, while organic chemistry dealt with carbon-containing compounds that were first identified in living things. Biochemistry is an outgrowth of the application of organic chemistry to biology and relates to the chemical basis for living things. In the later chapters of this text we will explore organic and biochemistry in a bit more detail and you will notice examples of organic compounds scattered throughout the text. Today, the lines between the various fields have blurred significantly and a contemporary chemist is expected to have a broad background in all of these areas.

In this chapter we will discuss some of the properties of matter, how chemists measure those properties and we will introduce some of the vocabulary that is used throughout chemistry and the other physical sciences.

Let's begin with matter. **Matter** is defined as any substance that has mass. It's important to distinguish here between weight and mass. **Weight** is the result of the pull of gravity on an object. On the Moon, an object will weigh less than the same object on Earth because the pull of gravity is less on the Moon. The **mass** of an object, however, is an inherent property of that object and does not change, regardless of location, gravitational pull, or whatever. It is a property that is solely dependent on the quantity of matter within the object.

Contemporary theories suggests that matter is composed of **atoms**. Atoms themselves are constructed and characterized from the subatomic particles called neutrons, protons and electrons that the atom possesses. Atoms will be studied closer in chapter 4 of this text. There are at least 116 different types of atoms recognized by scientists today, each type having a different number of protons in its nucleus. These different types of atoms are called **elements**.

As different elements were discovered and named, abbreviations of their names were developed to allow for a convenient chemical shorthand. The abbreviation for an element is called its chemical symbol. A chemical symbol consists of one or two letters, and the relationship between the symbol and the name of the element is generally apparent. Thus helium has the chemical symbol He, nitrogen is N, and lithium is Li. Sometimes the symbol is less apparent but is decipherable; magnesium is Mg, strontium is Sr, and manganese is Mn. Symbols for elements that have been known since ancient times, however, are often based on Latin or Greek names and appear somewhat obscure from their modern English names. For example, copper is Cu (from cuprum), silver is Ag (from argentum), gold is Au (from aurum), and iron is (Fe from ferrum). Throughout your study of chemistry, you will routinely use chemical symbols and it is important that you begin the process of learning the names and chemical symbols for the common elements. By the time you complete this course, you should be able to name and identify the symbols for common elements. Table 1.2.1 contains a starter list of common elements that you should begin learning now, as they will be the ones you are expected to know for this course.

Table 1.2.1: Names and Chemical Symbols for Common Elements

Element	Chemical Symbol	Element	Chemical Symbol
Aluminum	Al	Gold	Au
Beryllium	Be	Argon	Ar
Boron	B	Potassium	K
Carbon	C	Calcium	Ca
Fluorine	F	Zinc	Zn
Helium	He	Sulfur	S

Element	Chemical Symbol	Element	Chemical Symbol
Hydrogen	H	Phosphorus	P
Lithium	Li	Chlorine	Cl
Magnesium	Mg	Iodine	I
Manganese	Mn	Arsenic	As
Mercury	Hg	Barium	Ba
Neon	Ne	Bromine	Br
Nitrogen	N	Iron	Fe
Oxygen	O	Copper	Cu
Silicon	Si	Lead	Pb
Sodium	Na	Silver	Ag
Tin	Sn	Chromium	Cr

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Paul R. Young, Professor of Chemistry, University of Illinois at Chicago, Wiki: AskTheNerd; PRY@askthenerd.com - pyoung@uic.edu; ChemistryOnline.com
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

1.2: Chemicals Compose Ordinary Things is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

1.3: Hypothesis, Theories, and Laws

Learning Objectives

- Describe the difference between hypothesis and theory as scientific terms.
- Describe the difference between a theory and scientific law.

Although many have taken science classes throughout the course of their studies, people often have incorrect or misleading ideas about some of the most important and basic principles in science. Most students have heard of hypotheses, theories, and laws, but what do these terms really mean? Prior to reading this section, consider what you have learned about these terms before. What do these terms mean to you? What do you read that contradicts or supports what you thought?

What is a Fact?

A fact is a basic statement established by experiment or observation. All facts are true under the specific conditions of the observation.

What is a Hypothesis?

One of the most common terms used in science classes is a "hypothesis". The word can have many different definitions, depending on the context in which it is being used:

- An educated guess: a scientific hypothesis provides a suggested solution based on evidence.
- Prediction: if you have ever carried out a science experiment, you probably made this type of hypothesis when you predicted the outcome of your experiment.
- Tentative or proposed explanation: hypotheses can be suggestions about why something is observed. In order for it to be scientific, however, a scientist must be able to test the explanation to see if it works and if it is able to correctly predict what will happen in a situation. For example, "if my hypothesis is correct, we should see ___ result when we perform ___ test."

> *A hypothesis is very tentative; it can be easily changed.*

What is a Theory?

The *United States National Academy of Sciences* describes what a theory is as follows:

> "Some scientific explanations are so well established that no new evidence is likely to alter them. The explanation becomes a scientific theory. In everyday language a theory means a hunch or speculation. Not so in science. In science, the word **theory** refers to a comprehensive explanation of an important feature of nature supported by facts gathered over time. Theories also allow scientists to make predictions about as yet unobserved phenomena."
>
> "A scientific theory is a well-substantiated explanation of some aspect of the natural world, based on a body of facts that have been repeatedly confirmed through observation and experimentation. Such fact-supported theories are not "guesses" but reliable accounts of the real world. The theory of biological evolution is more than "just a theory." It is as factual an explanation of the universe as the atomic theory of matter (stating that everything is made of atoms) or the germ theory of disease (which states that many diseases are caused by germs). Our understanding of gravity is still a work in progress. But the phenomenon of gravity, like evolution, is an accepted fact.

Note some key features of theories that are important to understand from this description:

- Theories are explanations of natural phenomena. They aren't predictions (although we may use theories to make predictions). They are explanations as to why we observe something.
- Theories aren't likely to change. They have a large amount of support and are able to satisfactorily explain numerous observations. Theories can, indeed, be facts. Theories can change, but it is a long and difficult process. In order for a theory to change, there must be many observations or pieces of evidence that the theory cannot explain.
- Theories are not guesses. The phrase "just a theory" has no room in science. To be a scientific theory carries a lot of weight; it is not just one person's idea about something

> *Theories aren't likely to change.*

What is a Law?

Scientific laws are similar to scientific theories in that they are principles that can be used to predict the behavior of the natural world. Both scientific laws and scientific theories are typically well-supported by observations and/or experimental evidence. Usually scientific laws refer to rules for how nature will behave under certain conditions, frequently written as an equation. Scientific theories are more overarching explanations of how nature works and why it exhibits certain characteristics. As a comparison, theories explain why we observe what we do and laws describe what happens.

For example, around the year 1800, Jacques Charles and other scientists were working with gases to, among other reasons, improve the design of the hot air balloon. These scientists found, after many, many tests, that certain patterns existed in the observations on gas behavior. If the temperature of the gas is increased, the volume of the gas increased. This is known as a natural law. A law is a relationship that exists between variables in a group of data. Laws describe the patterns we see in large amounts of data, but do not describe why the patterns exist.

What is a Belief?

A **belief** is a statement that is not scientifically provable. Beliefs may or may not be incorrect; they just are outside the realm of science to explore.

Laws vs. Theories

A common misconception is that scientific theories are rudimentary ideas that will eventually graduate into scientific laws when enough data and evidence has accumulated. A theory does not change into a scientific law with the accumulation of new or better evidence. Remember, **theories are explanations** and **laws are patterns** we see in large amounts of data, frequently written as an equation. A theory will always remain a theory; a law will always remain a law.



Video **1.3.1**: What's the difference between a scientific law and theory?

Summary

- A hypothesis is a tentative explanation that can be tested by further investigation.
- A theory is a well-supported explanation of observations.
- A scientific law is a statement that summarizes the relationship between variables.
- An experiment is a controlled method of testing a hypothesis.

Contributions & Attributions

-
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

1.3: Hypothesis, Theories, and Laws is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **1.6: Hypothesis, Theories, and Laws** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.4: The Scientific Method - How Chemists Think

Learning Objectives

- Identify the components of the scientific method.

Scientists search for answers to questions and solutions to problems by using a procedure called the scientific method. This procedure consists of making observations, formulating hypotheses, and designing experiments; which leads to additional observations, hypotheses, and experiments in repeated cycles (Figure 1.4.1).




Figure 1.4.1: The Steps in the Scientific Method.

Step 1: Make observations

Observations can be qualitative or quantitative. **Qualitative observations** describe properties or occurrences in ways that do not rely on numbers. Examples of qualitative observations include the following: "the outside air temperature is cooler during the winter season," "table salt is a crystalline solid," "sulfur crystals are yellow," and "dissolving a penny in dilute nitric acid forms a blue solution and a brown gas." **Quantitative observations** are measurements, which by definition consist of both a number and a unit. Examples of quantitative observations include the following: "the melting point of crystalline sulfur is 115.21° Celsius," and "35.9 grams of table salt—the chemical name of which is sodium chloride—dissolve in 100 grams of water at 20° Celsius." For the question of the dinosaurs' extinction, the initial observation was quantitative: iridium concentrations in sediments dating to 66 million years ago were 20–160 times higher than normal.

Step 2: Formulate a hypothesis

After deciding to learn more about an observation or a set of observations, scientists generally begin an investigation by forming a hypothesis, a tentative explanation for the observation(s). The hypothesis may not be correct, but it puts the scientist's understanding of the system being studied into a form that can be tested. For example, the observation that we experience alternating periods of light and darkness corresponding to observed movements of the sun, moon, clouds, and shadows is consistent with either one of two hypotheses:

a. Earth rotates on its axis every 24 hours, alternately exposing one side to the sun.
b. The sun revolves around Earth every 24 hours.

Suitable experiments can be designed to choose between these two alternatives. For the disappearance of the dinosaurs, the hypothesis was that the impact of a large extraterrestrial object caused their extinction. Unfortunately (or perhaps fortunately), this hypothesis does not lend itself to direct testing by any obvious experiment, but scientists can collect additional data that either support or refute it.

Step 3: Design and perform experiments

After a hypothesis has been formed, scientists conduct experiments to test its validity. Experiments are systematic observations or measurements, preferably made under controlled conditions—that is—under conditions in which a single variable changes.

Step 4: Accept or modify the hypothesis

A properly designed and executed experiment enables a scientist to determine whether or not the original hypothesis is valid. If the hypothesis is valid, the scientist can proceed to step 5. In other cases, experiments often demonstrate that the hypothesis is incorrect or that it must be modified and requires further experimentation.

Step 5: Development into a law and/or theory

More experimental data are then collected and analyzed, at which point a scientist may begin to think that the results are sufficiently reproducible (i.e., dependable) to merit being summarized in a law, a verbal or mathematical description of a phenomenon that allows for general predictions. A law simply states what happens; it does not address the question of why.

One example of a law, the law of definite proportions, which was discovered by the French scientist Joseph Proust (1754–1826), states that a chemical substance always contains the same proportions of elements by mass. Thus, sodium chloride (table salt) always contains the same proportion by mass of sodium to chlorine, in this case 39.34% sodium and 60.66% chlorine by mass, and sucrose (table sugar) is always 42.11% carbon, 6.48% hydrogen, and 51.41% oxygen by mass.

Whereas a law states only what happens, a theory attempts to explain why nature behaves as it does. Laws are unlikely to change greatly over time unless a major experimental error is discovered. In contrast, a theory, by definition, is incomplete and imperfect, evolving with time to explain new facts as they are discovered.

Because scientists can enter the cycle shown in Figure 1.4.1 at any point, the actual application of the scientific method to different topics can take many different forms. For example, a scientist may start with a hypothesis formed by reading about work done by others in the field, rather than by making direct observations.

✔ Example 1.4.1

Classify each statement as a law, a theory, an experiment, a hypothesis, an observation.

a. Ice always floats on liquid water.
b. Birds evolved from dinosaurs.
c. Hot air is less dense than cold air, probably because the components of hot air are moving more rapidly.
d. When 10 g of ice were added to 100 mL of water at 25°C, the temperature of the water decreased to 15.5°C after the ice melted.
e. The ingredients of Ivory soap were analyzed to see whether it really is 99.44% pure, as advertised.

Solution

a. This is a general statement of a relationship between the properties of liquid and solid water, so it is a law.
b. This is a possible explanation for the origin of birds, so it is a hypothesis.
c. This is a statement that tries to explain the relationship between the temperature and the density of air based on fundamental principles, so it is a theory.
d. The temperature is measured before and after a change is made in a system, so these are observations.
e. This is an analysis designed to test a hypothesis (in this case, the manufacturer's claim of purity), so it is an experiment.

? Exercise 1.4.1

Classify each statement as a law, a theory, an experiment, a hypothesis, a qualitative observation, or a quantitative observation.

a. Measured amounts of acid were added to a Rolaids tablet to see whether it really "consumes 47 times its weight in excess stomach acid."
b. Heat always flows from hot objects to cooler ones, not in the opposite direction.
c. The universe was formed by a massive explosion that propelled matter into a vacuum.
d. Michael Jordan is the greatest pure shooter to ever play professional basketball.
e. Limestone is relatively insoluble in water, but dissolves readily in dilute acid with the evolution of a gas.

Answer a

experiment

Answer b

law

Answer c

theory

Answer d

hypothesis

Answer e

observation

Summary

The scientific method is a method of investigation involving experimentation and observation to acquire new knowledge, solve problems, and answer questions. The key steps in the scientific method include the following:

- Step 1: Make observations.
- Step 2: Formulate a hypothesis.
- Step 3: Test the hypothesis through experimentation.
- Step 4: Accept or modify the hypothesis.
- Step 5: Develop into a law and/or a theory.

Contributions & Attributions

-
- Wikipedia

1.4: The Scientific Method - How Chemists Think is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **1.3: The Scientific Method - How Chemists Think** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

2: Measurements and Density

2.1: Taking Measurements
2.2: Scientific Notation - Writing Large and Small Numbers
2.3: Significant Figures - Writing Numbers to Reflect Precision
2.4: Significant Figures in Calculations
2.5: The Basic Units of Measurement
2.6: Problem Solving and Unit Conversions
2.7: Solving Multistep Conversion Problems
2.8: Units Raised to a Power
2.9: Density
2.10: Dosage Calculations

2: Measurements and Density is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

2.1: Taking Measurements

Learning Objectives

- Express quantities properly, using a number and a unit.

A coffee maker's instructions tell you to fill the coffee pot with 4 cups of water and to use 3 scoops of coffee. When you follow these instructions, you are measuring. When you visit a doctor's office, a nurse checks your temperature, height, weight, and perhaps blood pressure (Figure 2.1.1); the nurse is also measuring.



Figure 2.1.1: Measuring Blood Pressure. A nurse or a doctor measuring a patient's blood pressure is taking a measurement. (GFDL; Pia von Lützau).

Chemists measure the properties of matter and express these measurements as quantities. A quantity is an amount of something and consists of a number and a unit. The number tells us how many (or how much), and the unit tells us what the scale of measurement is. For example, when a distance is reported as "5 kilometers," we know that the quantity has been expressed in units of kilometers and that the number of kilometers is 5. If you ask a friend how far they walk from home to school, and the friend answers "12" without specifying a unit, you do not know whether your friend walks 12 kilometers, 12 miles, 12 furlongs, or 12 yards. *Both a number and a unit must be included to express a quantity properly.*

To understand chemistry, we need a clear understanding of the units chemists work with and the rules they follow for expressing numbers. The next two sections examine the rules for expressing numbers.

Example 2.1.1

Identify the number and the unit in each quantity.

a. one dozen eggs
b. 2.54 centimeters
c. a box of pencils
d. 88 meters per second

Solution

a. The number is one, and the unit is a dozen eggs.
b. The number is 2.54, and the unit is centimeter.
c. The number 1 is implied because the quantity is only *a* box. The unit is box of pencils.
d. The number is 88, and the unit is meters per second. Note that in this case the unit is actually a combination of two units: meters and seconds.

Key Take Away

- Identify a quantity properly with a number and a unit.

2.1: Taking Measurements is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.1: Taking Measurements** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

2.2: Scientific Notation - Writing Large and Small Numbers

Learning Objectives

- Express a large number or a small number in scientific notation.
- Carry out arithmetical operations and express the final answer in scientific notation

Chemists often work with numbers that are exceedingly large or small. For example, entering the mass in grams of a hydrogen atom into a calculator would require a display with at least 24 decimal places. A system called **scientific notation** avoids much of the tedium and awkwardness of manipulating numbers with large or small magnitudes. In scientific notation, these numbers are expressed in the form

$$N \times 10^n$$

where N is greater than or equal to 1 and less than 10 ($1 \le N < 10$), and n is a positive or negative integer ($10^0 = 1$). The number 10 is called the base because it is this number that is raised to the power n. Although a base number may have values other than 10, the base number in scientific notation is always 10.

A simple way to convert numbers to scientific notation is to move the decimal point as many places to the left or right as needed to give a number from 1 to 10 (N). The magnitude of n is then determined as follows:

- If the decimal point is moved to the left n places, n is positive.
- If the decimal point is moved to the right n places, n is negative.

Another way to remember this is to recognize that as the number N decreases in magnitude, the exponent increases and vice versa. The application of this rule is illustrated in Example 2.2.1.

Example 2.2.1: Expressing Numbers in Scientific Notation

Convert each number to scientific notation.

a. 637.8
b. 0.0479
c. 7.86
d. 12,378
e. 0.00032
f. 61.06700
g. 2002.080
h. 0.01020

Solution

Solutions to Example 2.2.1

	Explanation	Answer
a	To convert 637.8 to a number from 1 to 10, we move the decimal point two places to the left: 637.8 Because the decimal point was moved two places to the left, n = 2.	6.378×10^2
b	To convert 0.0479 to a number from 1 to 10, we move the decimal point two places to the right: 0.0479 Because the decimal point was moved two places to the right, n = −2.	4.79×10^{-2}
c	This is usually expressed simply as 7.86. (Recall that $10^0 = 1$.)	7.86×10^0
d	Because the decimal point was moved four places to the left, n = 4.	1.2378×10^4

	Explanation	Answer
e	Because the decimal point was moved four places to the right, n = −4.	3.2×10^{-4}
f	Because the decimal point was moved one place to the left, n = 1.	6.106700×10^{1}
g	Because the decimal point was moved three places to the left, n = 3.	2.002080×10^{3}
h	Because the decimal point was moved two places to the right, n = -2.	1.020×10^{-2}

Addition and Subtraction

Before numbers expressed in scientific notation can be added or subtracted, they must be converted to a form in which all the exponents have the same value. The appropriate operation is then carried out on the values of N. Example $2.2.2$ illustrates how to do this.

Example 2.2.2: Expressing Sums and Differences in Scientific Notation

Carry out the appropriate operation and then express the answer in scientific notation.

a. $(1.36 \times 10^{2}) + (4.73 \times 10^{3})$
b. $(6.923 \times 10^{-3}) - (8.756 \times 10^{-4})$

Solution

Solutions to Example 2.2.2.

	Explanation	Answer
a	Both exponents must have the same value, so these numbers are converted to either $(1.36 \times 10^{2}) + (47.3 \times 10^{2}) =$ $(1.36 + 47.3) \times 10^{2} = 48.66 \times 10^{2}$ or $(0.136 \times 10^{3}) + (4.73 \times 10^{3}) =$ $(0.136 + 4.73) \times 10^{3}) = 4.87 \times 10^{3}$. Choosing either alternative gives the same answer, reported to two decimal places. In converting 48.66 × 10^{2} to scientific notation, n has become more positive by 1 because the value of N has decreased.	4.8
b	Converting the exponents to the same value gives either $(6.923 \times 10^{-3}) - (0.8756 \times 10^{-3}) =$ $(6.923 - 0.8756) \times 10^{-3}$ or $(69.23 \times 10^{-4}) - (8.756 \times 10^{-4}) =$ $(69.23 - 8.756) \times 10^{-4} = 60.474 \times 10^{-4}$. In converting 60.474 × 10^{-4} to scientific notation, n has become more positive by 1 because the value of N has decreased.	6.0

Multiplication and Division

When multiplying numbers expressed in scientific notation, we multiply the values of N and add together the values of n. Conversely, when dividing, we divide N in the dividend (the number being divided) by N in the divisor (the number by which we are dividing) and then

subtract n in the divisor from n in the dividend. In contrast to addition and subtraction, the exponents do not have to be the same in multiplication and division. Examples of problems involving multiplication and division are shown in Example 2.2.3.

Example 2.2.3: Expressing Products and Quotients in Scientific Notation

Perform the appropriate operation and express your answer in scientific notation.

a. $(6.022 \times 10^{23})(6.42 \times 10^{-2})$

b. $\dfrac{1.67 \times 10^{-24}}{9.12 \times 10^{-28}}$

c. $\dfrac{(6.63 \times 10^{-34})(6.0 \times 10)}{8.52 \times 10^{-2}}$

Solution

Solution to Example 2.2.3

	Explanation	
a	In multiplication, we add the exponents: $(6.022 \times 10^{23})(6.42 \times 10^{-2}) = (6.022)(6.42) \times 10^{[23+(-2)]} = 38.7 \times 10^{21}$ In converting 38.7×10^{21} to scientific notation, n has become more positive by 1 because the value of N has decreased.	3.87×10^{22}
b	In division, we subtract the exponents: $\frac{1.67 \times 10^{-24}}{9.12 \times 10^{-28}} = \frac{1.67}{9.12} \times 10^{[-24-(-28)]} = 0.183 \times 10^{4}$ In converting 0.183×10^{4} to scientific notation, n has become more negative by 1 because the value of N has increased.	1.83×10^{3}
c	This problem has both multiplication and division: $\frac{(6.63 \times 10^{-34})(6.0 \times 10)}{(8.52 \times 10^{-2})} = \frac{39.78}{8.52} \times 10^{[-34+1-(-2)]}$	4.7×10^{-31}

2.2: Scientific Notation - Writing Large and Small Numbers is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.2: Scientific Notation - Writing Large and Small Numbers** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.3: Significant Figures - Writing Numbers to Reflect Precision

Learning Objectives

- Identify the number of significant figures in a reported value.

The **significant figures** in a measurement consist of all the certain digits in that measurement plus one uncertain or estimated digit. In the ruler illustration below, the bottom ruler gave a length with 2 significant figures, while the top ruler gave a length with 3 significant figures. In a correctly reported measurement, the final digit is significant but not certain. Insignificant digits are not reported. With either ruler, it would not be possible to report the length at $2.553\ \text{cm}$ as there is no possible way that the thousandths digit could be estimated. The 3 is not significant and would not be reported.




Figure 2.3.1: Measurement with two different rulers.
Ruler A's measurement can be rounded to 2.55, with 2 certain digits, while Ruler B's measurement of 2.5 has 1 certain digit

Measurement Uncertainty


Some error or **uncertainty** always exists in any measurement. The amount of uncertainty depends both upon the skill of the measurer and upon the quality of the measuring tool. While some balances are capable of measuring masses only to the nearest $0.1\ \text{g}$, other highly sensitive balances are capable of measuring to the nearest $0.001\ \text{g}$ or even better. Many measuring tools such as rulers and graduated cylinders have small lines which need to be carefully read in order to make a measurement. Figure 2.3.1 shows two rulers making the same measurement of an object (indicated by the blue arrow).

With either ruler, it is clear that the length of the object is between 2 and 3 cm. The bottom ruler contains no millimeter markings. With that ruler, the tenths digit can be estimated and the length may be reported as $2.5\ \text{cm}$. However, another person may judge that the measurement is $2.4\ \text{cm}$ or perhaps $2.6\ \text{cm}$. While the 2 is known for certain, the value of the tenths digit is uncertain.

The top ruler contains marks for tenths of a centimeter (millimeters). Now the same object may be measured as $2.55\ \text{cm}$. The measurer is capable of estimating the hundredths digit because he can be certain that the tenths digit is a 5. Again, another measurer may report the length to be $2.54\ \text{cm}$ or $2.56\ \text{cm}$. In this case, there are two certain digits (the 2 and the 5), with the hundredths digit being uncertain. Clearly, the top ruler is a superior ruler for measuring lengths as precisely as possible.


Example 2.3.1: Reporting Measurements to the Proper Number of Significant Figures

Use each diagram to report a measurement to the proper number of significant figures.



a.

b.



Ruler measuring a rectangle in units of centimeters, with the rectangle's edge between 1.2 and 1.3 cm marks


Solutions

Solutions to Example 2.3.1

	Explanation	Answer
a.	The arrow is between 4.0 and 5.0, so the measurement is at least 4.0. The arrow is between the third and fourth small tick marks, so it's at least 0.3. We will have to estimate the last place. It looks like about one-third of the way across the space, so let us estimate the hundredths place as 3. The symbol psi stands for "pounds per square inch" and is a unit of pressure, like air in a tire. The measurement is reported to three significant figures.	4.33 psi
b.	The rectangle is at least 1.0 cm wide but certainly not 2.0 cm wide, so the first significant digit is 1. The rectangle's width is past the second tick mark but not the third; if each tick mark represents 0.1, then the rectangle is at least 0.2 in the next significant digit. We have to estimate the next place because there are no markings to guide us. It appears to be about halfway between 0.2 and 0.3, so we will estimate the next place to be a 5. Thus, the measured width of the rectangle is 1.25 cm. The measurement is reported to three significant figures.	1.25 cm

? Exercise 2.3.1

What would be the reported width of this rectangle?



Answer

1.25 cm

When you look at a reported measurement, it is necessary to be able to count the number of significant figures. The table below details the rules for determining the number of significant figures in a reported measurement. For the examples in the table, assume that the quantities are correctly reported values of a measured quantity.

Table 2.3.1: Significant Figure Rules

Rule	Examples
1. All nonzero digits in a measurement are significant.	• 237 has three significant figures. • 1.897 has four significant figures.
2. Zeros that appear between other nonzero digits (middle zeros) are always significant.	• 39,004 has five significant figures. • 5.02 has three significant figures.
3. Zeros that appear in front of all of the nonzero digits are called leading zeros. Leading zeros are never significant.	• 0.008 has one significant figure. • 0.000416 has three significant figures.
4. Zeros that appear after all nonzero digits are called trailing zeros. A number with trailing zeros that lacks a decimal point may or may not be significant. **Use scientific notation to indicate the appropriate number of significant figures.**	• 1400 is ambiguous. ◦ 1.4×10^3 has two significant figures. ◦ 1.40×10^3 three significant figures. ◦ 1.400×10^3 has four significant figures.
5. Trailing zeros in a number with a decimal point are significant. This is true whether the zeros occur before or after the decimal point.	• 620.0 has four significant figures. • 19.000 has five significant figures.

Exact Numbers

Integers obtained either by counting objects or from definitions are exact numbers, which are considered to have infinitely many significant figures. If we have counted four objects, for example, then the number 4 has an infinite number of significant figures (i.e., it represents 4.000…). Similarly, 1 foot (ft) is defined to contain 12 inches (in), so the number 12 in the following equation has infinitely many significant figures:

Example 2.3.2

Give the number of significant figures in each. Identify the rule for each.

a. 5.87
b. 0.031
c. 52.90
d. 00.2001
e. 500
f. 6 atoms

Solution

Solution to Example 2.3.2

	Explanation	Answer
a	All three numbers are significant (rule 1).	5.87, three significant figures
b	The leading zeros are not significant (rule 3). The 3 and the 1 are significant (rule 1).	0.031, two significant figures
c	The 5, the 2 and the 9 are significant (rule 1). The trailing zero is also significant (rule 5).	52.90, four significant figures

	Explanation	Answer
d	The leading zeros are not significant (rule 3). The 2 and the 1 are significant (rule 1) and the middle zeros are also significant (rule 2).	00.2001, four significant figures
e	The number is ambiguous. It could have one, two or three significant figures.	500, ambiguous
f	The 6 is a counting number. A counting number is an exact number.	6, infinite

? Exercise 2.3.2

Give the number of significant figures in each.

a. 36.7 m
b. 0.006606 s
c. 2,002 kg
d. 306,490,000 people
e. 3,800 g

Answer a

three significant figures

Answer b

four significant figures

Answer c

four significant figures

Answer d

infinite (exact number)

Answer e

Ambiguous, could be two, three or four significant figures.

Accuracy and Precision


Measurements may be accurate, meaning that the measured value is the same as the true value; they may be precise, meaning that multiple measurements give nearly identical values (i.e., reproducible results); they may be both accurate and precise; or they may be neither accurate nor precise. The goal of scientists is to obtain measured values that are both accurate and precise. The video below demonstrates the concepts of accuracy and precision.



Video 2.3.1: Difference between precision and accuracy.

Example 2.3.3

The following archery targets show marks that represent the results of four sets of measurements.



Which target shows

a. a precise, but inaccurate set of measurements?
b. a set of measurements that is both precise and accurate?
c. a set of measurements that is neither precise nor accurate?

Solution

a. Set **(a)** is precise, but inaccurate.
b. Set **(c)** is both precise and accurate.
c. Set **(d)** is neither precise nor accurate.

Summary

Uncertainty exists in all measurements. The degree of uncertainty is affected in part by the quality of the measuring tool. Significant figures give an indication of the certainty of a measurement. Rules allow decisions to be made about how many digits to use in any given situation.

2.3: Significant Figures - Writing Numbers to Reflect Precision is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.3: Significant Figures - Writing Numbers to Reflect Precision** by Henry Agnew, Marisa Alviar-Agnew, Sridhar Budhi is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.4: Significant Figures in Calculations

Learning Objectives

- Use significant figures correctly in arithmetical operations.

Rounding

Before dealing with the specifics of the rules for determining the significant figures in a calculated result, we need to be able to round numbers correctly. To **round** a number, first decide how many significant figures the number should have. Once you know that, round to that many digits, starting from the left. If the number immediately to the right of the last significant digit is less than 5, it is dropped and the value of the last significant digit remains the same. If the number immediately to the right of the last significant digit is greater than or equal to 5, the last significant digit is increased by 1.

Consider the measurement $207.518\ \text{m}$ Right now, the measurement contains six significant figures. How would we successively round it to fewer and fewer significant figures? Follow the process as outlined in Table $2.4.1$.

Table 2.4.1: Rounding examples

Number of Significant Figures	Rounded Value	Reasoning
6	207.518	All digits are significant
5	207.52	8 rounds the 1 up to 2
4	207.5	2 is dropped
3	208	5 rounds the 7 up to 8
2	210	8 is replaced by a 0 and rounds the 0 up to 1
1	200	1 is replaced by a 0

Notice that the more rounding that is done, the less reliable the figure is. An approximate value may be sufficient for some purposes, but scientific work requires a much higher level of detail.

It is important to be aware of significant figures when you are mathematically manipulating numbers. For example, dividing 125 by 307 on a calculator gives 0.4071661238… to an infinite number of digits. But do the digits in this answer have any practical meaning, especially when you are starting with numbers that have only three significant figures each? When performing mathematical operations, there are two rules for limiting the number of significant figures in an answer—one rule is for addition and subtraction, and one rule is for multiplication and division.

> *In operations involving significant figures, the answer is reported in such a way that it reflects the reliability of the **least precise** operation. An answer is no more precise than the least precise number used to get the answer.*

Multiplication and Division

For multiplication or division, the rule is to count the number of significant figures in each number being multiplied or divided and then limit the significant figures in the answer to the lowest count. An example is as follows:

$$\underbrace{38.65}_{\text{4 sig figs}} \times \underbrace{105.93}_{\text{5 sig figs}} = \underbrace{4{,}094.1945}_{\text{reduce to 4 sig figs}}$$

The final answer, limited to four significant figures, is 4,094. The first digit dropped is 1, so we do not round up.

Scientific notation provides a way of communicating significant figures without ambiguity. You simply include all the significant figures in the leading number. For example, the number 450 has two significant figures and would be written in scientific notation

as 4.5×10^2, whereas 450.0 has four significant figures and would be written as 4.500×10^2. In scientific notation, all significant figures are listed explicitly.

Example 2.4.1

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 23.096×90.300
b. 125×9.000

Solution

a

Explanation	Answer
The calculator answer is 2,085.5688, but we need to round it to five significant figures. Because the first digit to be dropped (in the tenths place) is greater than 5, we round up to 2,085.6.	2.0856×10^3

b

Explanation	Answer
The calculator gives 1,125 as the answer, but we limit it to three significant figures.	1.13×10^3

Addition and Subtraction

How are significant figures handled in calculations? It depends on what type of calculation is being performed. If the calculation is an addition or a subtraction, the rule is as follows: limit the reported answer to the rightmost column that all numbers have significant figures in common. For example, if you were to add 1.2 and 4.71, we note that the first number stops its significant figures in the tenths column, while the second number stops its significant figures in the hundredths column. We therefore limit our answer to the tenths column.

$$\begin{array}{r} 1.2 \\ \underline{4.41} \\ 5.61 \end{array}$$

↑ limit final answer to the tenths column: 5.6

We drop the last digit—the 1—because it is not significant to the final answer.

The dropping of positions in sums and differences brings up the topic of rounding. Although there are several conventions, in this text we will adopt the following rule: the final answer should be rounded up if the first dropped digit is 5 or greater, and rounded down if the first dropped digit is less than 5.

$$\begin{array}{r} 77.2 \\ \underline{10.46} \\ 87.66 \end{array}$$

↑ limit final answer to the tenths column and round up: 87.7

Example 2.4.2

a. 13.77 + 908.226
b. 1,027 + 611 + 363.06

Solution

a

Explanation	Answer
The calculator answer is 921.996, but because 13.77 has its farthest-right significant figure in the hundredths place, we need to round the final answer to the hundredths position. Because the first digit to be dropped (in the thousandths place) is greater than 5, we round up to 922.00	$922.00 = 9.2200 \times 10^2$

b

Explanation	Answer
The calculator gives 2,001.06 as the answer, but because 611 and 1027 has its farthest-right significant figure in the ones place, the final answer must be limited to the ones position.	$2,001.06 = 2.001 \times 10^3$

Exercise 2.4.2

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 217 ÷ 903
b. 13.77 + 908.226 + 515
c. 255.0 − 99
d. 0.00666 × 321

Answer a:

$0.240 = 2.40 \times 10^{-1}$

Answer b:

$1,437 = 1.437 \times 10^3$

Answer c:

$156 = 1.56 \times 10^2$

Answer d:

$2.14 = 2.14 \times 10^0$

Remember that calculators do not understand significant figures. *You* are the one who must apply the rules of significant figures to a result from your calculator.

Calculations Involving Multiplication/Division and Addition/Subtraction

In practice, chemists generally work with a calculator and carry all digits forward through subsequent calculations. When working on paper, however, we often want to minimize the number of digits we have to write out. Because successive rounding can compound inaccuracies, intermediate rounding needs to be handled correctly. When working on paper, always round an intermediate result so as to retain at least one more digit than can be justified and carry this number into the next step in the calculation. The final answer is then rounded to the correct number of significant figures at the very end.



Video 2.4.1: Significant figures in mixed operations (https://www.youtube.com/watch?v=yBntMndXQWA).



Video 2.4.2: https://www.youtube.com/watch?v=__csP0NtlGI

In the worked examples in this text, we will often show the results of intermediate steps in a calculation. In doing so, we will show the results to only the correct number of significant figures allowed for that step, in effect treating each step as a separate calculation. This procedure is intended to reinforce the rules for determining the number of significant figures, but in some cases it may give a final answer that differs in the last digit from that obtained using a calculator, where all digits are carried through to the last step.

Example 2.4.3

a. 2(1.008 g) + 15.99 g
b. 137.3 s + 2(35.45 s)
c. $\frac{118.7g}{2} - 35.5g$

Solution

a.

Explanation	Answer

Explanation	Answer
2(1.008 g) + 15.99 g = Perform multiplication first. 2 (1.008 g *4 sig figs*) = **2.016 g** *4 sig figs* The number with the least number of significant figures is 1.008 g; *the number 2 is an exact number and therefore has an infinite number of significant figures*. **Then, perform the addition.** **2.016 g** *thousandths place* + 15.99 g *hundredths place (least precise)* = 18.006 g **Round the final answer.** Round the final answer to the hundredths place since 15.99 has its farthest right significant figure in the hundredths place (least precise).	18.01 g (rounding up)

b.

Explanation	Answer
137.3 s + 2(35.45 s) = **Perform multiplication first.** 2(35.45 s *4 sig figs*) = **70.90 s** *4 sig figs* The number with the least number of significant figures is 35.45; *the number 2 is an exact number and therefore has an infinite number of significant figures.* **Then, perform the addition.** 137.3 s *tenths place (least precise)* + **70.90 s** *hundredths place* = 208.20 s **Round the final answer.** Round the final answer to the tenths place based on 137.3 s.	208.2 s

c.

Explanation	Answer
$\frac{118.7g}{2} - 35.5g =$ **Perform division first.** $\frac{118.7g}{2}$ *4 sig figs* = **59.35 g** *4 sig figs* The number with the least number of significant figures is 118.7 g; *the number 2 is an exact number and therefore has an infinite number of significant figures.* **Perform subtraction next.** **59.35 g** *hundredths place* − 35.5 g *tenths place (least precise)* = 23.85 g **Round the final answer.** Round the final answer to the tenths place based on 35.5 g.	23.9 g (rounding up)

Exercise 2.4.3

Complete the calculations and report your answers using the correct number of significant figures.

a. 5(1.008s) - 10.66 s
b. 99.0 cm+ 2(5.56 cm)

Answer a

-5.62 s

Answer b

110.2 cm

Summary

- Rounding
  - If the number to be dropped is greater than or equal to 5, increase the number to its left by 1 (e.g. 2.9699 rounded to three significant figures is 2.97).
  - If the number to be dropped is less than 5, there is no change (e.g. 4.00443 rounded to four significant figures is 4.004).
- The rule in multiplication and division is that the final answer should have the same number of significant figures as there are in the number with the fewest significant figures.
- The rule in addition and subtraction is that the answer is given the same number of decimal places as the term with the fewest decimal places.

2.4: Significant Figures in Calculations is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.4: Significant Figures in Calculations** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.5: The Basic Units of Measurement

Learning Objectives

- State the different measurement systems used in chemistry.
- Describe how prefixes are used in the metric system and identify how the prefixes milli-, centi-, and kilo- compare to the base unit.

How long is a yard? It depends on whom you ask and when you asked the question. Today we have a standard definition of the yard, which you can see marked on every football field. If you move the ball ten yards, you get a first down and it does not matter whether you are playing in Los Angeles, Dallas, or Green Bay. But at one time that yard was arbitrarily defined as the distance from the tip of the king's nose to the end of his outstretched hand. Of course, the problem there is simple: new king, new distance (and then you have to re-mark all of those football fields).




Figure 2.5.1: Meter standard (left) and Kilogram standard (right).

SI Base Units

All measurements depend on the use of units that are well known and understood. The **English system** of measurement units (inches, feet, ounces, etc.) are not used in science because of the difficulty in converting from one unit to another. The **metric system** is used because all metric units are based on multiples of 10, making conversions very simple. The metric system was originally established in France in 1795. **The International System of Units** is a system of measurement based on the metric system. The acronym **SI** is commonly used to refer to this system and stands for the French term, *Le Système International d'Unités*. The SI was adopted by international agreement in 1960 and is composed of seven base units in Table 2.5.1.

Table 2.5.1: SI Base Units of Measurement

Quantity	SI Base Unit	Symbol
Length	meter	m
Mass	kilogram	kg
Temperature	kelvin	K
Time	second	s
Amount of a Substance	mole	mol
Electric Current	ampere	A
Luminous Intensity	candela	cd

The first units are frequently encountered in chemistry. All other measurement quantities, such as volume, force, and energy, can be derived from these seven base units.

Unfortunately, the Metric System is Not Ubiquitous

The map below shows the adoption of the SI units in countries around the world. The United States has legally adopted the metric system for measurements, but does not use it in everyday practice. Great Britain and much of Canada use a combination of metric and imperial units.




Figure 2.5.1: Areas of world using metric system (in green). Only a few countries are slow or resistant to adoption including the United States.

Prefix Multipliers

Conversions between metric system units are straightforward because the system is based on powers of ten. For example, meters, centimeters, and millimeters are all metric units of length. There are 10 millimeters in 1 centimeter and 100 centimeters in 1 meter. **Metric prefixes** are used to distinguish between units of different size. These prefixes all derive from either Latin or Greek terms. For example, *mega* comes from the Greek word $\mu\varepsilon\gamma\alpha\varsigma$, meaning "great". Table $2.5.2$ lists the most common metric prefixes and their relationship to the central unit that has no prefix. Length is used as an example to demonstrate the relative size of each prefixed unit.

Table 2.5.2: SI Prefixes

Prefix	Unit Abbreviation	Meaning	Example
giga	G	1,000,000,000	1 gigameter $(\mathrm{Gm}) = 10^9$ m
mega	M	1,000,000	1 megameter $(\mathrm{Mm}) = 10^6$ m
kilo	k	1,000	1 kilometer $(\mathrm{km}) = 1,000$ m
hecto	h	100	1 hectometer $(\mathrm{hm}) = 100$ m
deka	da	10	1 dekameter $(\mathrm{dam}) = 10$ m
		1	1 meter (m)
deci	d	1/10	1 decimeter $(\mathrm{dm}) = 0.1$ m
centi	c	1/100	1 centimeter $(\mathrm{cm}) = 0.01$ m
milli	m	1/1,000	1 millimeter $(\mathrm{mm}) = 0.001$ m
micro	μ	1/1,000,000	1 micrometer $(\mu\mathrm{m}) = 10^{-6}$ m
nano	n	1/1,000,000,000	1 nanometer $(\mathrm{nm}) = 10^{-9}$ m
pico	p	1/1,000,000,000,000	1 picometer $(\mathrm{pm}) = 10^{-12}$ m

There are a couple of odd little practices with the use of metric abbreviations. Most abbreviations are lowercase. We use "m" for meter and not "M". However, when it comes to volume, the base unit "liter" is abbreviated as "L" and not "l". So we would write 3.5 milliliters as $3.5\ \mathrm{mL}$.

As a practical matter, whenever possible you should express the units in a small and manageable number. If you are measuring the weight of a material that weighs 6.5 kg, this is easier than saying it weighs 6500 g or 0.65 dag. All three are correct, but the kg

units in this case make for a small and easily managed number. However, if a specific problem needs grams instead of kilograms, go with the grams for consistency.

Example 2.5.1: Unit Abbreviations

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kiloliter
b. microsecond
c. decimeter
d. nanogram

Solutions

Solutions to Example 2.5.1

	Explanation	Answer
a	The prefix kilo means “1,000 ×,” so 1 kL equals 1,000 L.	kL
b	The prefix micro implies 1/1,000,000th of a unit, so 1 µs equals 0.000001 s.	µs
c	The prefix deci means 1/10th, so 1 dm equals 0.1 m.	dm
d	The prefix nano means 1/1000000000, so a nanogram is equal to 0.000000001 g.	ng

Exercise 2.5.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kilometer
b. milligram
c. nanosecond
d. centiliter

Answer a:
km

Answer b:
mg

Answer c:
ns

Answer d:
cL

Summary

- Metric prefixes derive from Latin or Greek terms. The prefixes are used to make the units manageable.
- The SI system is based on multiples of ten. There are seven basic units in the SI system. Five of these units are commonly used in chemistry.

2.5: The Basic Units of Measurement is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.5: The Basic Units of Measurement** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.6: Problem Solving and Unit Conversions

Learning Objectives

- To convert a value reported in one unit to a corresponding value in a different unit using conversion factors.

During your studies of chemistry (and physics also), you will note that mathematical equations are used in many different applications. Many of these equations have a number of different variables with which you will need to work. You should also note that these equations will often require you to use measurements with their units. Algebra skills become very important here!

Converting Between Units with Conversion Factors

A **conversion factor** is a factor used to convert one unit of measurement into another. A simple conversion factor can convert meters into centimeters, or a more complex one can convert miles per hour into meters per second. Since most calculations require measurements to be in certain units, you will find many uses for conversion factors. Always remember that a conversion factor has to represent a fact; this fact can either be simple or more complex. For instance, you already know that 12 eggs equal 1 dozen. A more complex fact is that the speed of light is 1.86×10^5 miles/sec. Either one of these can be used as a conversion factor depending on what type of calculation you are working with (Table $2.6.1$).

Table $2.6.1$: Conversion Factors from SI units to English Units

English Units	Metric Units	Quantity
1 ounce (oz)	28.35 grams (g)	*mass
1 fluid once (oz)	29.6 mL	volume
2.205 pounds (lb)	1 kilogram (kg)	*mass
1 inch (in)	2.54 centimeters (cm)	length
0.6214 miles (mi)	1 kilometer (km)	length
1 quarter (qt)	0.95 liters (L)	volume

*Pounds and ounces are technically units of force, not mass, but this fact is often ignored by the non-scientific community.

Of course, there are other ratios which are not listed in Table $2.6.1$. They may include:

- Ratios embedded in the text of the problem (using words such as *per* or *in each*, or using symbols such as / or %).
- Conversions in the metric system, as covered earlier in this chapter.
- Common knowledge ratios (such as 60 seconds = 1 minute).

If you learned the SI units and prefixes described, then you know that 1 cm is 1/100th of a meter.

$$1 \text{ cm} = \frac{1}{100} \text{ m} = 10^{-2} \text{m}$$

or

$$100 \text{ cm} = 1 \text{ m}$$

Suppose we divide both sides of the equation by 1m (both the number *and* the unit):

$$\frac{100 \text{ cm}}{1 \text{ m}} = \frac{1 \text{ m}}{1 \text{ m}}$$

As long as we perform the same operation on both sides of the equals sign, the expression remains an equality. Look at the right side of the equation; it now has the same quantity in the numerator (the top) as it has in the denominator (the bottom). Any fraction that has the same quantity in the numerator and the denominator has a value of 1:

$$\frac{100 \text{ cm}}{1 \text{ m}} = \frac{1000 \text{ mm}}{1 \text{ m}} = \frac{1 \times 10^6 \mu\text{m}}{1 \text{ m}} = 1$$

We know that 100 cm *is* 1 m, so we have the same quantity on the top and the bottom of our fraction, although it is expressed in different units.

Performing Dimensional Analysis

Dimensional analysis is amongst the most valuable tools that physical scientists use. Simply put, it is the conversion between an amount in one unit to the corresponding amount in a desired unit using various conversion factors. This is valuable because certain measurements are more accurate or easier to find than others. The use of units in a calculation to ensure that we obtain the final proper units is called *dimensional analysis*.

Here is a simple example. How many centimeters are there in 3.55 m? Perhaps you can determine the answer in your head. If there are 100 cm in every meter, then 3.55 m equals 355 cm. To solve the problem more formally with a conversion factor, we first write the quantity we are given, 3.55 m. Then we multiply this quantity by a conversion factor, which is the same as multiplying it by 1. We can write 1 as $\frac{100\text{ cm}}{1\text{ m}}$ and multiply:

$$3.55\text{ m} \times \frac{100\text{ cm}}{1\text{ m}}$$

The 3.55 m can be thought of as a fraction with a 1 in the denominator. Because m, the abbreviation for meters, occurs in both the numerator *and* the denominator of our expression, they cancel out:

$$\frac{3.55\ \cancel{\text{m}}}{1} \times \frac{100\text{ cm}}{1\ \cancel{\text{m}}}$$

The final step is to perform the calculation that remains once the units have been canceled:

$$\frac{3.55}{1} \times \frac{100\text{ cm}}{1} = 355\text{ cm}$$

In the final answer, we omit the 1 in the denominator. Thus, by a more formal procedure, we find that 3.55 m equals 355 cm. A generalized description of this process is as follows:

quantity (in old units) × conversion factor = quantity (in new units)

You may be wondering why we use a seemingly complicated procedure for a straightforward conversion. In later studies, the conversion problems you encounter *will not always be so simple*. If you master the technique of applying conversion factors, you will be able to solve a large variety of problems.

In the previous example, we used the fraction $\frac{100\text{ cm}}{1\text{ m}}$ as a conversion factor. Does the conversion factor $\frac{1\text{ m}}{100\text{ cm}}$ also equal 1? Yes, it does; it has the same quantity in the numerator as in the denominator (except that they are expressed in different units). Why did we not use *that* conversion factor? If we had used the second conversion factor, the original unit would not have canceled, and the result would have been meaningless. Here is what we would have gotten:

$$3.55\text{ m} \times \frac{1\text{ m}}{100\text{ cm}} = 0.0355\frac{\text{m}^2}{\text{cm}}$$

For the answer to be meaningful, we have to *construct the conversion factor in a form that causes the original unit to cancel out*. Figure 2.6.1 shows a concept map for constructing a proper conversion.




Figure 2.6.1: A Concept Map for Conversions. This is how you construct a conversion factor to convert from one unit to another. Converting meters to centimeters: the unit you are converting is on the bottom of the fraction, and the unit to convert to is on top

General Steps in Performing Dimensional Analysis

1. Identify the "**given**" information in the problem. Look for a number with units to start this problem with.
2. What is the problem asking you to "**find**"? In other words, what unit will your answer have?
3. Use **ratios** and conversion factors to cancel out the units that aren't part of your answer, and leave you with units that are part of your answer.
4. When your units cancel out correctly, you are ready to do the **math**. You are multiplying fractions, so you multiply the top numbers and divide by the bottom numbers in the fractions.



Significant Figures in Conversions

How do conversion factors affect the determination of significant figures?

- Numbers in conversion factors based on prefix changes, such as kilograms to grams, are *not* considered in the determination of significant figures in a calculation because the numbers in such conversion factors are exact.
- Exact numbers are defined or counted numbers, not measured numbers, and can be considered as having an infinite number of significant figures. (In other words, 1 kg is exactly 1,000 g, by the definition of kilo-.)
- Counted numbers are also exact. If there are 16 students in a classroom, the number 16 is exact.
- In contrast, conversion factors that come from measurements (such as density, as we will see shortly) or that are approximations have a limited number of significant figures and should be considered in determining the significant figures of the final answer.

Example 2.6.1

Steps for Problem Solving for Example 2.6.1 and 2.6.2

	Example 2.6.1	**Example 2.6.2**
Steps for Problem Solving	The average volume of blood in an adult male is 4.7 L. What is this volume in milliliters?	A hummingbird can flap its wings once in 18 ms. How many seconds are in 18 ms?
Identify the "given" information and what the problem is asking you to "find."	Given: 4.7 L Find: mL	Given: 18 ms Find: s
List other known quantities.	$1\,mL = 10^{-3}L$	$1\,ms = 10^{-3}s$
Prepare a concept map and use the proper conversion factor.	 $\frac{1mL}{10^{-3}L}$	 $\frac{10^{-3}s}{1ms}$
Cancel units and calculate.	$4.7\,\cancel{L} \times \frac{1\text{ mL}}{10^{-3}\ \cancel{L}} = 4,700\text{ mL}$ or $4.7\,\cancel{L} \times \frac{1,000\text{ mL}}{1\ \cancel{L}} = 4,700\text{ mL}$ or 4.7 x 10^3 2SF, not ambiguous	$18\ \cancel{ms} \times \frac{10^{-3}\text{ s}}{1\ \cancel{ms}} = 0.018\text{ s}$ or $18\ \cancel{ms} \times \frac{1\text{ s}}{1,000\ \cancel{ms}} = 0.018\text{ s}$
Think about your result.	The amount in mL should be 1000 times larger than the given amount in L.	The amount in s should be 1/1000 the given amount in ms.

Exercise 2.6.1

Perform each conversion.

a. 101,000 ns to seconds
b. 32.08 kg to grams

c. 1.53 grams to cg

Answer a:

$1.01000x10^{-4}s$

Answer b:

$3.208x10^{4}g$

Answer c:

$1.53x10^{2}cg$

Summary

- Conversion factors are used to convert one unit of measurement into another.
- Dimensional analysis (unit conversions) involves the use of conversion factors that will cancel unwanted units and produce the appropriate units.

2.6: Problem Solving and Unit Conversions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.6: Problem Solving and Unit Conversions** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.


2.7: Solving Multistep Conversion Problems

Multiple Conversions

Sometimes you will have to perform more than one conversion to obtain the desired unit. For example, suppose you want to convert 54.7 km into millimeters. We will set up a series of conversion factors so that each conversion factor produces the next unit in the sequence. We first convert the given amount in km to the base unit which is meters. We know that 1,000 m =1 km.

Then we convert meters to mm, remembering that $1\ \text{mm} = 10^{-3}\ \text{m}$.

Concept Map



Calculation

$$54.7\ \cancel{\text{km}} \times \frac{1,000\ \cancel{\text{m}}}{1\ \cancel{\text{km}}} \times \frac{1\ \cancel{\text{mm}}}{10^{-3}\cancel{\text{m}}} = 54,700,000\ \text{mm} \tag{2.7.1}$$


$$= 5.47 \times 10^{7}\ \text{mm} \tag{2.7.2}$$

> *In each step, the previous unit is canceled and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.*

Example 2.7.1: Unit Conversion

Convert 58.2 ms to megaseconds in one multistep calculation.

SOLUTION

Steps for Problem Solving	Unit Conversion
Identify the "given"information and what the problem is asking you to "find."	Given: 58.2 ms Find: Ms
List other known quantities	$1ms = 10^{-3}s$ $1Ms = 10^{6}s$
Prepare a concept map	
Calculate	$58.2\ \cancel{\text{ms}} \times \frac{10^{-3}\ \cancel{s}}{1\ \cancel{\text{ms}}} \times \frac{1\ \text{Ms}}{1,000,000\ \cancel{s}} = 0.0000000582\ \text{Ms}$ $= 5.82 \times 10^{-8}\ \text{Ms}$ Neither conversion factor affects the number of significant figures in the final answer.

Example 2.7.2: Unit Conversion

How many seconds are in a day?

Solution

Steps for Problem Solving	Unit Conversion
Identify the "given"information and what the problem is asking you to "find."	Given: 1 day Find: s
List other known quantities	1 day = 24 hours 1 hour = 60 minutes 1 minute = 60 seconds
Prepare a concept map	
Calculate	$1\text{ d} \times \frac{24\text{ hr}}{1\text{ d}} \times \frac{60\text{ min}}{1\text{ hr}} \times \frac{60\text{ s}}{1\text{ min}} = 86,400\text{ s}$

Exercise 2.7.1

Perform each conversion in one multistep calculation.

a. 43.007 ng to kg
b. 1005 in to ft
c. 12 mi to km

Answer a:

$4.3007x10^{-11}kg$

Answer b:

83.75 ft

Answer c:

19 km

Career Focus: Pharmacist

A pharmacist dispenses drugs that have been prescribed by a doctor. Although that may sound straightforward, pharmacists in the United States must hold a doctorate in pharmacy and be licensed by the state in which they work. Most pharmacy programs require four years of education in a specialty pharmacy school. Pharmacists must know a lot of chemistry and biology so they can understand the effects that drugs (which are chemicals, after all) have on the body. Pharmacists can advise physicians on the selection, dosage, interactions, and side effects of drugs. They can also advise patients on the proper use of their medications, including when and how to take specific drugs properly. Pharmacists can be found in drugstores, hospitals, and other medical facilities. Curiously, an outdated name for pharmacist is *chemist*, which was used when pharmacists formerly did a lot of drug preparation, or *compounding*. In modern times, pharmacists rarely compound their own drugs, but their knowledge of the sciences, including chemistry, helps them provide valuable services in support of everyone's health.



A woman consulting with a pharmacist. This image was released by the National Cancer Institute, an agency part of the National Institutes of Health. (Public Domain; Rhoda Baer (Photographer) via NIH).

Summary

In multistep conversion problems, the previous unit is canceled for each step and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.

Contributuors

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

2.7: Solving Multistep Conversion Problems is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

2.8: Units Raised to a Power


Learning Objectives

- To convert a value reported in one unit raised to a power of 10, to a corresponding value in a different unit raised to the same power of 10, using conversion factors.

Conversion factors for area and volume can also be produced by the dimensional analysis method. Just remember that if a quantity is raised to a power of 10, both the number and the unit must be raised to the same power of 10. For example, to convert 1500 cm^2 to m^2, we need to start with the relationship between centimeter and meter. We know that 1 cm = 10^{-2} m or 100 cm =1 m, but since we are given the quantity in 1500 cm^2, then we have to use the relationship:

$$1\,cm^2 = (10^{-2}\,m)^2 = 10^{-4}\,m^2$$

CONCEPT MAP



To convert centimeters squared to meters squared, use the conversion factor 0.01 meters per 1 centimeter, squared overall

CALCULATION

$$1500\ \cancel{\text{cm}^2} \times \left(\frac{10^{-2}\ \text{m}}{1\ \cancel{\text{cm}}}\right)^2 = 0.15\ \text{m}^2$$

or

$$1500\ \cancel{\text{cm}^2} \times \left(\frac{1\ \text{m}}{100\ \cancel{\text{cm}}}\right)^2 = 0.15\ \text{m}^2$$

or

$$1500\ \cancel{\text{cm}^2} \times \frac{1\ \text{m}^2}{10,000\ \cancel{\text{cm}^2}} = 0.15\ \text{m}^2$$


Example 2.8.1: Volume of a Sphere

What is the volume of a sphere (radius 4.30 inches) in cubic cm (cm^3)?

Solution

Solution for Example 2.8.1

Steps for Problem Solving	What is the volume of a sphere (radius 4.30 inches) in cubic cm (cm^3)?
Identify the "given" information and what the problem is asking you to "find."	Given: radius = 4.30 in Find: cm^3 (volume)
Determine other known quantities.	Volume of a sphere: V = $\frac{4}{3} \times \pi \times r^3$ = $\frac{4}{3} \times 3.1416 \times (4.3\underline{0}in)^3$ = $33\underline{3}.04in^3$

Steps for Problem Solving	What is the volume of a sphere (radius 4.30 inches) in cubic cm (cm^3)?
Prepare a concept map.	 $\left(\frac{2.54\ cm}{1\ in}\right)^3$ To convert inches cubed to centimeters cubed, use conversion factor 2.54 centimeters per 1 inch, cubed overall
Calculate.	$333.04\ \cancel{in^3}\left(\frac{2.54cm}{1\ \cancel{in}}\right)^3 = 5.46 \times 10^3 cm^3$
Think about your result.	A centimeter is a smaller unit than an inch, so the answer in cubic centimeters is larger than the given value in cubic inches.

? Exercise 2.8.1

Lake Tahoe has a surface area of 191 square miles. What is the area in square km (km^2)?

Answer

495 km^2

Contributions & Attributions

2.8: Units Raised to a Power is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.8: Units Raised to a Power** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.9: Density

Learning Objectives

- Define density.
- Use density as a conversion factor.

Density (ρ) is a physical property found by dividing the mass of an object by its volume. Regardless of the sample size, density is always constant. For example, the density of a pure sample of tungsten is always 19.25 grams per cubic centimeter. This means that whether you have one gram or one kilogram of the sample, the density will never vary. The equation, as we already know, is as follows:

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

or just

$$\rho = \frac{m}{V} \tag{2.9.1}$$

Based on this equation, it's clear that density can, and does, vary from element to element and substance to substance due to differences in the relationship of mass and volume. Pure water, for example, has a density of 0.998 g/cm^3 at 25° C. The average densities of some common substances are in Table 2.9.1. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will "float."

Table 2.9.1: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Density can be measured for all substances—solids, liquids and gases. For solids and liquids, density is often reported using the units of g/cm^3. Densities of gases, which are significantly lower than the densities of solids and liquids, are often given using units of g/L.

Example 2.9.1: Ethyl Alcohol

Calculate the density of a 30.2 mL sample of ethyl alcohol with a mass of 23.71002 g

Solution

This is a direct application of Equation 2.9.1:

$$\rho = \frac{23.71002\,g}{30.2\,mL} = 0.785\,g/mL$$

Exercise 2.9.1

a. Find the density (in kg/L) of a sample that has a volume of 36.5 L and a mass of 10.0 kg.
b. If you have a 2.130 mL sample of acetic acid with mass 0.002234 kg, what is the density in kg/L?

Answer a

$0.274\ kg/L$

Answer b

$1.049\ kg/L$

Density as a Conversion Factor

Conversion factors can also be constructed for converting between different kinds of units. For example, density can be used to convert between the mass and the volume of a substance. Consider mercury, which is a liquid at room temperature and has a density of 13.6 g/mL. The density tells us that 13.6 g of mercury have a volume of 1 mL. We can write that relationship as follows:

13.6 g mercury = 1 mL mercury

This relationship can be used to construct two conversion factors:

$$\frac{13.6\ g}{1\ mL} = 1$$

and

$$\frac{1\ mL}{13.6\ g} = 1$$

Which one do we use? It depends, as usual, on the units we need to cancel and introduce. For example, suppose we want to know the mass of 2.0 mL of mercury. We would use the conversion factor that has milliliters on the bottom (so that the milliliter unit cancels) and grams on top, so that our final answer has a unit of mass:

$$2.0\ \cancel{\text{mL}} \times \frac{13.6\ \text{g}}{1\ \cancel{\text{mL}}} = 27.2\ \text{g} = 27\ \text{g}$$

In the last step, we limit our final answer to two significant figures because the volume quantity has only two significant figures; the 1 in the volume unit is considered an exact number, so it does not affect the number of significant figures. The other conversion factor would be useful if we were given a mass and asked to find volume, as the following example illustrates.


> *Density can be used as a conversion factor between mass and volume.*

Example 2.9.2: Mercury Thermometer Steps for Problem Solving

A mercury thermometer for measuring a patient's temperature contains 0.750 g of mercury. What is the volume of this mass of mercury?

Solution

Solution to Example 2.9.2

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 0.750 g Find: mL
List other known quantities.	13.6 g/mL (density of mercury)
Prepare a concept map.	
Calculate.	$0.750\ \cancel{\text{g}} \times \frac{1\ \text{mL}}{13.6\ \cancel{\text{g}}} = 0.055147\ldots\ \text{mL} \approx 0.0551\ \text{mL}$ We have limited the final answer to three significant figures.

Exercise 2.9.2

What is the volume of 100.0 g of air if its density is 1.3 g/L?

Answer

$77\ L$

Summary

- Density is defined as the mass of an object divided by its volume.
- Density can be used as a conversion factor between mass and volume.

Contributions & Attributions

2.9: Density is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **2.9: Density** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.10: Dosage Calculations

Learning Objectives

- Calculate drug dosages using conversion factors.

To Your Health: Dosages

A medicine can be more harmful than helpful if it is not taken in the proper dosage. A dosage (or dose) is the specific amount of a medicine that is known to be therapeutic for an ailment in a patient of a certain size. Dosages of the active ingredient in medications are usually described by units of mass, typically grams or milligrams, and generally are equated with a number of capsules or teaspoonfuls to be swallowed or injected. The amount of the active ingredient in a medicine is carefully controlled so that the proper number of pills or spoonfuls contains the proper dose.

Most drugs must be taken in just the right amount. If too little is taken, the desired effects will not occur (or will not occur fast enough for comfort); if too much is taken, there may be potential side effects that are worse than the original ailment. Some drugs are available in multiple dosages. For example, tablets of the medication levothyroxine sodium, a synthetic thyroid hormone for those suffering from decreased thyroid gland function, are available in 11 different doses, ranging from 25 micrograms (µg) to 300 µg. It is a doctor's responsibility to prescribe the correct dosage for a patient, and it is a pharmacist's responsibility to provide the patient with the correct medicine at the dosage prescribed. Thus, proper quantities—which are expressed using numbers and their associated units—are crucial for keeping us healthy.

Effects are dose-dependent

Chemicals are the most common things for which doses are measured, but there are others, such as radiation exposure. For humans, most doses of micronutrients and medications are measured in milligrams (mg), but some are measured in micrograms because of their potency. Nonmedicinal poisons span the measurement scale; some poisons are so dangerous that a single microgram of it could be deadly, whereas other substances take much more. For example, even water is toxic when consumed in large enough quantities.

Dosage (the size of each dose) determines the strength and duration of the health benefits of nutrients, and also of the therapeutic effects of medical treatments. Dosage also determines the severity of adverse effects of treatments and toxins.

Duration of exposure, that is, the period of time over which the dose was received (all at once or gradually) also determines its effects (the body may build tolerance to gradual exposure to a drug, while a large immediate dose could be deadly).

The route by which a dose is exposed to, may affect the outcome, because some medications have different effects depending on whether they are inhaled, ingested, taken transdermally, injected, or inserted.

The dosage, route, concentration, and division over time may all be critical considerations in the administering of drugs, or in responding to exposure to a toxin. In nutrition, the route is usually a given, as nutrients are generally eaten; while dosage and the frequency of ingestion of nutrients are very important variables in preventing disease and promoting overall health.

Calculation of dose

Calculating drug dosages for humans based on the doses used in animal studies can be based on weight (e.g., mg/kg) or surface area (e.g., mg/m2) based on weight2/3.[1]

Drug dosage calculations

Drug dosage calculation is required if the physician's order is different from what is available.

Example 2.10.1

a. A physician ordered 100 mg of Demerol. Demerol is available as 50 mg per tablet. How many tablets should the nurse administer?
b. The doctor's order is 1.2 g of Folic Acid. Folic Acid is available as 800 mg per tablet. How many tablets should be taken?

Solution

a. We start with the given, 100 mg. We want to change the unit from *mg* to *tablets*. There are 50 mg in 1 tablet (Remember that *per* tablet means *one* tablet. From this relationship, we can construct the conversion factor. We use the conversion factor that will cancel out the original unit, *mg*, and introduce the unit we are converting to, which is *tablet*.

$$100 \text{ mg} \times \frac{1 \text{ tablet}}{50 \text{ mg}} = 2 \text{ tablets}$$

Hence, the nurse should administer 2 tablets.

b. We start with the given, 1.2 g and we want to change *grams* to number of *tablets*. First, we convert *1.2 g* to *mg* and then convert *mg to tablets*. We need a conversion factor for each step.

$$1.2 \text{ g} \times \frac{1{,}000 \text{ mg}}{1 \text{ g}} \times \frac{1 \text{ tablet}}{800 \text{ mg}} = 1.5 \text{ tablets}$$

Hence, 1.5 tablets should be taken.

Exercise 2.10.1

Calculate each of the following.

a. The physician ordered 20 mg of Valium. Valium is available as 10 mg per tablet. How many tablets should the nurse administer?
b. The doctor's order is 1 g of Calcium. What is on hand is Calcium as 500 mg per tablet. How many tablets should be taken?

Answer a

Start with 20 mg of Valium. For the conversion factor, we know that 10 mg Valium = 1 tablet

$$20 \text{ mg} \times \frac{1 \text{ tablet}}{10 \text{ mg}} = 2 \text{ tablets}$$

Answer b

Start with 1 g (1000 mg) of Calcium. For the conversion factor, we know that 500 mg Calcium = 1 tablet

$$1000 \text{ mg} \times \frac{1 \text{ tablet}}{500 \text{ mg}} = 2 \text{ tablets}$$

Drug Dosage Calculation based on Body Weight. Many drugs (especially in children) are dosed according to body weight (mg/kg). These calculations are carried in 3-step conversions. The first step is to convert the body weight from *pounds (lbs)* to *kg*. The second step is to convert *kg* to *mg* (the total *mg dose* calculated based on body weight). And, finally, the *mg dose* is converted to the number of *tablets*. (as in Example 1.8.1).

Example 2.10.2

a. Demerol is ordered 1.5mg/kg for a patient that is 220 lbs. Demerol is available as 50 mg per tablet. How many tablets should the nurse administer?
b. A doctor prescribes amoxicillin 30mg/kg to a child weighing 73.5 lbs. Amoxicillin is available as 500 mg tablets. How many tablets should the nurse administer?

Solution

a. We start with the given, 220 *lbs*. We want to change the unit from *lbs* to *kg*, and then, from *kg* to *total dose (mg)* and then *the mg dose to tablets*. The first conversion factor will cancel out the original unit, *lbs*, and introduce the unit we are converting to, which is *kg*. The second conversion factor will cancel out *kg*, and introduce the unit of the *dose* (usually *mg*) and then *mg* to *tablet*.

$$220 \text{ lbs} \times \frac{1 \text{ kg}}{2.2 \text{ lbs}} \times \frac{1.5 \text{ mg}}{1 \text{ kg}} \times \frac{1 \text{ tablet}}{50 \text{ mg}} = 3 \text{ tablets}$$

Hence, the nurse administers 3 tablets.

b. Start with the given, 73.5 *lbs*. We want to change the unit from *lbs* to *kg*, and then, from *kg* to total dose (*mg*) and then *mg* to *tablets*. The first conversion factor will cancel out the original unit, *lbs*, and convert to *kg*. The second conversion factor will cancel out *kg*, and convert to the total *mg dose* and the final conversion will cancel *mg* to introduce the final unit, *tablet*.

$$73.5 \text{ lbs} \times \frac{1 \text{ kg}}{2.2 \text{ lbs}} \times \frac{30 \text{ mg}}{1 \text{ kg}} \times \frac{1 \text{ tablet}}{500 \text{ mg}} = 2 \text{ tablets}$$

Hence, the nurse administers 2 tablets.

? Exercise 2.10.2

Calculate each of the following.

a. Vancomycin is ordered 15mg/kg for a patient that is 110 lbs. Vancomycin is available as 250 mg per capsule. How many capsules should the nurse administer? (ans. 3 capsules)
b. A doctor prescribes ampicillin 40mg/kg to a patient weighing 55 lbs. Ampicillin is available as 500 mg tablets. How many tablets should the nurse administer? (ans. 2 tablets)

Answer a

$$110 \text{ lbs} \times \frac{1 \text{ kg}}{2.2 \text{ lbs}} \times \frac{15 \text{ mg}}{1 \text{ kg}} \times \frac{1 \text{ tablet}}{250 \text{ mg}} = 3 \text{ tablets}$$

Answer b

$$55 \text{ lbs} \times \frac{1 \text{ kg}}{2.2 \text{ lbs}} \times \frac{40 \text{ mg}}{1 \text{ kg}} \times \frac{1 \text{ tablet}}{500 \text{ mg}} = 2 \text{ tablets}$$

References

1. The Use of Body Surface Area as a Criterion of Drug Dosage in Cancer Chemotherapy(opens in new window) [cancerres.aacrjournals.org] "The Use of Body Surface Area as a Criterion of Drug Dosage in Cancer Chemotherapy" D Pinkel. Cancer Research 1958

Contributors and Attributions

- Wikipedia
- Leticia Colmenares (UH-WCC)

2.10: Dosage Calculations is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **1.8: Dosage Calculations** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

3: Matter

3.1: In Your Room
3.2: What is Matter?
3.3: Classifying Matter According to Its State- Solid, Liquid, and Gas
3.4: Classifying Matter According to Its Composition
3.5: Differences in Matter- Physical and Chemical Properties
3.6: Changes in Matter - Physical and Chemical Changes
3.7: Conservation of Mass - There is No New Matter
3.E: Chemistry, Matter, and Measurement (Exercises)

3: Matter is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

3.1: In Your Room

Matter is any substance that has mass and takes up space (volume); this includes atoms and anything made up of these, but not other energy phenomena or waves such as light or sound. While this simple definition is easily applied, the way people view matter is often broken down into two characteristic length scales: the macroscopic and the microscopic.



Figure 3.1.1: A typical American university and college dormitory room in 2002. (CC BY-SA 3.0; Raul654).

The *macroscopic scale* is the length scale on which objects or phenomena are large enough to be visible almost practically with the naked eye, without magnifying optical instruments. Everything that one can see, touch and handle in the dorm room of Figure 3.1.1 is in the macroscopic scale and to describe each of these objects only few macroscopic properties are required. However, each of these items can be decomposed into smaller microscopic scale properties.

The *microscopic scale* is the scale of objects and events smaller than those that can easily be seen by the naked eye, requiring a lens or microscope to see them clearly. All the everyday objects that we can bump into, touch or squeeze are ultimately composed of atoms. This ordinary atomic matter is in turn made up of interacting subatomic particles—usually a nucleus of protons and neutrons, and a cloud of orbiting electrons. Because of this, a large number of variables are needed to describe such a system which complicates the characterization.

Matter vs. Mass

Matter should not be confused with mass, as the two are not the same in modern physics. Matter is itself a physical substance of which systems may be composed, while mass is not a substance but rather a quantitative property of matter and other substances or systems.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Wikipedia

3.1: In Your Room is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

3.2: What is Matter?

Learning Objectives

- Define matter and explain how it is composed of building blocks known as "atoms".

We are all familiar with matter. For most common objects that we deal with every day, it is fairly simple to demonstrate that they have mass and take up space. You might be able to imagine, however, the difficulty for people several hundred years ago to demonstrate that air has mass and volume. Air (and all other gases) are invisible to the eye, have very small masses compared to equal amounts of solids and liquids, and are quite easy to compress (change volume). Without sensitive equipment, it would have been difficult to convince people that gases are matter. Today, we can measure the mass of a small balloon when it is deflated and then blow it up, tie it off, and measure its mass again to detect the additional mass due to the air inside. The mass of air, under room conditions, that occupies a one quart jar is approximately 0.0002 pounds. This small amount of mass would have been difficult to measure in times before balances were designed to accurately measure very small masses. Later, scientists were able to compress gases into such a small volume that the gases turned into liquids, which made it clear that gases are matter.



Figure 3.2.1: Everything from an ant, to a truck, to the earth and even the entire galaxy is composed of matter. Images used with permission from Wikipedia (CC_SA-BY-3.0; credit High Contrast)

Even though the universe consists of "things" as wildly different as ants and galaxies, the matter that makes up all of these "things" is composed of a very limited number of building blocks. These building blocks are known as **atoms** (the smallest unit of matter), and so far, scientists have discovered or created a grand total of 118 different types of atoms. Scientists have given a name to each different type of atom. A substance that is composed of only one type of atom is called an element. At this point, what should amaze you is that all forms of matter in our universe are made with only 118 different building blocks. In some ways, it's sort of like cooking a gourmet, five-course meal using only three ingredients! How is it possible? To answer that question, you have to understand the ways in which different elements are put together to form matter.

The most important method that nature uses to organize atoms into matter is the formation of **molecules**. Molecules are groups of two or more atoms that have been bonded together. There are millions of different ways to bond atoms together, which means that there are millions of different possible molecules. Each of these molecules has its own set of chemical properties, and it's these

properties with which chemists are most concerned. You will learn a lot more about atoms and molecules, including how they were discovered, in a later part of the textbook.

Summary

All matter has mass and occupies space. All physical objects are made of matter. Matter itself is composed of tiny building blocks known as "atoms". There are only 118 different types of atoms known to man. Frequently, atoms are bonded together to form "molecules".

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Wikipedia
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

3.2: What is Matter? is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

3.3: Classifying Matter According to Its State- Solid, Liquid, and Gas

Learning Objectives

- To describe the solid, liquid and gas phases.

Water can take many forms. At low temperatures (below $0^{o}C$), it is a solid. When at "normal" temperatures (between $0^{o}C$ and $100^{o}C$), it is a liquid. While at temperatures above $100^{o}C$, water is a gas (steam). The state that water is in depends upon the temperature. Each state has its own unique set of physical properties. Matter typically exists in one of three states: **solid**, **liquid**, or **gas**.




Figure 3.3.1: Matter is usually classified into three classical states, with plasma sometimes added as a fourth state. From left to right: quartz (solid), water (liquid), nitrogen dioxide (gas).

The state that a given substance exhibits is also a physical property. Some substances exist as gases at room temperature (oxygen and carbon dioxide), while others, like water and mercury metal, exist as liquids. Most metals exist as solids at room temperature. All substances can exist in any of these three states. Figure 3.3.2 shows the differences among solids, liquids, and gases at the molecular level. A solid has definite volume and shape, a liquid has a definite volume but no definite shape, and a gas has neither a definite volume nor shape.




Figure 3.3.2: A Representation of the Solid, Liquid, and Gas States. (a) Solid O_2 has a fixed volume and shape, and the molecules are packed tightly together. (b) Liquid O_2 conforms to the shape of its container but has a fixed volume; it contains relatively densely packed molecules. (c) Gaseous O_2 fills its container completely—regardless of the container's size or shape—and consists of widely separated molecules.

Plasma: A Fourth State of Matter

Technically speaking, a fourth state of matter called plasma exists, but it does not naturally occur on earth, so we will omit it from our study here.



A plasma globe operating in a darkened room. (CC BY-SA 3.0; Chocolateoak).

Solids

In the solid state, the individual particles of a substance are in fixed positions with respect to each other because there is not enough thermal energy to overcome the intermolecular interactions between the particles. As a result, solids have a definite shape and volume. Most solids are hard, but some (like waxes) are relatively soft. Many solids composed of ions can also be quite brittle.

Solids are defined by the following characteristics:

- Definite shape (rigid)
- Definite volume
- Particles vibrate around fixed axes

If we were to cool liquid mercury to its freezing point of $-39^{\circ}\mathrm{C}$, and under the right pressure conditions, we would notice all of the liquid particles would go into the solid state. Mercury can be solidified when its temperature is brought to its freezing point. However, when returned to room temperature conditions, mercury does not exist in solid state for long, and returns back to its more common liquid form.

Solids usually have their constituent particles arranged in a regular, three-dimensional array of alternating positive and negative ions called a **crystal**. The effect of this regular arrangement of particles is sometimes visible macroscopically, as shown in Figure 3.3.3. Some solids, especially those composed of large molecules, cannot easily organize their particles in such regular crystals and exist as amorphous (literally, "without form") solids. Glass is one example of an amorphous solid.




Figure 3.3.3: (left) The periodic crystalline lattice structure of quartz SiO_2 in two-dimensions. (right) The random network structure of glassy SiO_2 in two-dimensions. Note that, as in the crystal, each Silicon atom is bonded to 4 oxygen atoms, where the fourth oxygen atom is obscured from view in this plane. Images used with permission (public domain).

Liquids

If the particles of a substance have enough energy to partially overcome intermolecular interactions, then the particles can move about each other while remaining in contact. This describes the liquid state. In a liquid, the particles are still in close contact, so liquids have a definite volume. However, because the particles can move about each other rather freely, a liquid has no definite shape and takes a shape dictated by its container.

Liquids have the following characteristics:

- No definite shape (takes the shape of its container).
- Has definite volume.
- Particles are free to move over each other, but are still attracted to each other.

A familiar liquid is mercury metal. Mercury is an anomaly. It is the only metal we know of that is liquid at room temperature. Mercury also has an ability to stick to itself (surface tension)—a property that all liquids exhibit. Mercury has a relatively high surface tension, which makes it very unique. Here you see mercury in its common liquid form.



***Video* 3.3.1** *: Mercury boiling to become a gas.*

If we heat liquid mercury to its boiling point of $357^{\circ}\mathrm{C}$ under the right pressure conditions, we would notice all particles in the liquid state go into the gas state.

Gases

If the particles of a substance have enough energy to completely overcome intermolecular interactions, then the particles can separate from each other and move about randomly in space. This describes the gas state, which we will consider in more detail elsewhere. Like liquids, gases have no definite shape, but unlike solids and liquids, gases have no definite volume either. The change from solid to liquid usually does not significantly change the volume of a substance. However, the change from a liquid to a gas significantly increases the volume of a substance, by a factor of 1,000 or more. Gases have the following characteristics:

- No definite shape (takes the shape of its container)
- No definite volume
- Particles move in random motion with little or no attraction to each other
- Highly compressible

Table 3.3.1: Characteristics of the Three States of Matter

Characteristics	Solids	Liquids	Gases
shape	definite	indefinite	indefinite
volume	definite	definite	indefinite
relative intermolecular interaction strength	strong	moderate	weak
relative particle positions	in contact and fixed in place	in contact but not fixed	not in contact, random positions

Example 3.3.1

What state or states of matter does each statement, describe?

a. This state has a definite volume, but no definite shape.
b. This state has no definite volume.
c. This state allows the individual particles to move about while remaining in contact.

Solution

a. This statement describes the liquid state.
b. This statement describes the gas state.
c. This statement describes the liquid state.

Exercise 3.3.1

What state or states of matter does each statement describe?

a. This state has individual particles in a fixed position with regard to each other.
b. This state has individual particles far apart from each other in space.
c. This state has a definite shape.

Answer a:

solid

Answer b:

gas

Answer c:

solid

Summary

- Three states of matter exist—solid, liquid, and gas.
- Solids have a definite shape and volume.
- Liquids have a definite volume, but take the shape of the container.
- Gases have no definite shape or volume.

3.3: Classifying Matter According to Its State- Solid, Liquid, and Gas is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **3.3: Classifying Matter According to Its State—Solid, Liquid, and Gas** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.4: Classifying Matter According to Its Composition

Learning Objectives

- Explain the difference between a pure substance and a mixture.
- Explain the difference between an element and a compound.
- Explain the difference between a homogeneous mixture and a heterogeneous mixture.

One useful way of organizing our understanding of matter is to think of a hierarchy that extends down from the most general and complex to the simplest and most fundamental (Figure 3.4.1). Matter can be classified into two broad categories: pure substances and mixtures. A **pure substance** is a form of matter that has a constant composition (meaning that it is the same everywhere) and properties that are constant throughout the sample (meaning that there is only one set of properties such as melting point, color, boiling point, etc. throughout the matter). A material composed of two or more substances is a **mixture**. Elements and compounds are both examples of pure substances. A substance that cannot be broken down into chemically simpler components is an **element**. Aluminum, which is used in soda cans, is an element. A substance that can be broken down into chemically simpler components (because it has more than one element) is a **compound**. For example, water is a compound composed of the elements hydrogen and oxygen. Today, there are about 118 elements in the known universe. In contrast, scientists have identified tens of millions of different compounds to date.




Figure 3.4.1: Relationships between the Types of Matter and the Methods Used to Separate Mixtures

Ordinary table salt is called sodium chloride. It is considered a **substance** because it has a uniform and definite composition. All samples of sodium chloride are chemically identical. Water is also a pure substance. Salt easily dissolves in water, but salt water cannot be classified as a substance because its composition can vary. You may dissolve a small amount of salt or a large amount into a given amount of water. A mixture is a physical blend of two or more components, each of which retains its own identity and properties in the **mixture**. Only the form of the salt is changed when it is dissolved into water. It retains its composition and properties.

A **homogeneous** mixture is a mixture in which the composition is uniform throughout the mixture. The salt water described above is homogeneous because the dissolved salt is evenly distributed throughout the entire salt water sample. Often it is easy to confuse a homogeneous mixture with a pure substance because they are both uniform. The difference is that the composition of the substance is always the same. The amount of salt in the salt water can vary from one sample to another. All solutions are considered homogeneous because the dissolved material is present in the same amount throughout the solution.

A **heterogeneous mixture** is a mixture in which the composition is not uniform throughout the mixture. Vegetable soup is a heterogeneous mixture. Any given spoonful of soup will contain varying amounts of the different vegetables and other components of the soup.

Phase

A phase is any part of a sample that has a uniform composition and properties. By definition, a pure substance or a homogeneous mixture consists of a single phase. A heterogeneous mixture consists of two or more phases. When oil and water are combined, they do not mix evenly, but instead form two separate layers. Each of the layers is called a phase.

Example 3.4.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. filtered tea
b. freshly squeezed orange juice
c. a compact disc
d. aluminum oxide, a white powder that contains a 2:3 ratio of aluminum and oxygen atoms
e. selenium

Given: a chemical substance

Asked for: its classification

Strategy:

A. Decide whether a substance is chemically pure. If it is pure, the substance is either an element or a compound. If a substance can be separated into its elements, it is a compound.
B. If a substance is not chemically pure, it is either a heterogeneous mixture or a homogeneous mixture. If its composition is uniform throughout, it is a homogeneous mixture.

Solution

a. **A)** Tea is a solution of compounds in water, so it is not chemically pure. It is usually separated from tea leaves by filtration.
B) Because the composition of the solution is uniform throughout, it is ***a homogeneous mixture***.
b. **A)** Orange juice contains particles of solid (pulp) as well as liquid; it is not chemically pure.
B) Because its composition is not uniform throughout, orange juice is ***a heterogeneous mixture***.
c. **A)** A compact disc is a solid material that contains more than one element, with regions of different compositions visible along its edge. Hence, a compact disc is not chemically pure.
B) The regions of different composition indicate that a compact disc is ***a heterogeneous mixture.***
d. **A)** Aluminum oxide is a single, chemically ***pure compound***.
e. **A)** Selenium is one of the known ***elements***.

Exercise 3.4.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. white wine
b. mercury
c. ranch-style salad dressing
d. table sugar (sucrose)

Answer a:
homogeneous mixture (solution)

Answer b:
element

Answer c:
heterogeneous mixture

Answer d:
compound

Example 3.4.2

How would a chemist categorize each example of matter?

a. saltwater
b. soil
c. water
d. oxygen

Solution

a. Saltwater acts as if it were a single substance even though it contains two substances—salt and water. Saltwater is a homogeneous mixture, or a solution.
b. Soil is composed of small pieces of a variety of materials, so it is a heterogeneous mixture.
c. Water is a substance. More specifically, because water is composed of hydrogen and oxygen, it is a compound.
d. Oxygen, a substance, is an element.

Exercise 3.4.2

How would a chemist categorize each example of matter?

a. coffee
b. hydrogen
c. an egg

Answer a:
a homogeneous mixture (solution), assuming it is filtered coffee

Answer b:
element

Answer c:
heterogeneous mixture

Summary

Matter can be classified into two broad categories: pure substances and mixtures. A pure substance is a form of matter that has a constant composition and properties that are constant throughout the sample. Mixtures are physical combinations of two or more elements and/or compounds. Mixtures can be classified as homogeneous or heterogeneous. Elements and compounds are both examples of pure substances. Compounds are substances that are made up of more than one type of atom. Elements are the simplest substances made up of only one type of atom.

Vocabulary

- Element: a substance that is made up of only one type of atom.
- Compound:a substance that is made up of more than one type of atom bonded together.
- Mixture: a combination of two or more elements or compounds which have not reacted to bond together; each part in the mixture retains its own properties.

Contributions & Attributions

-
- Stephen Lower, Professor Emeritus (Simon Fraser U.) Chem1 Virtual Textbook

---

3.4: Classifying Matter According to Its Composition is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **3.4: Classifying Matter According to Its Composition** by Henry Agnew, Marisa Alviar-Agnew, Stephen Lower is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.5: Differences in Matter- Physical and Chemical Properties

Learning Objectives

To separate physical from chemical properties.

All matter has physical and chemical properties. Physical properties are characteristics that scientists can measure without changing the composition of the sample under study, such as mass, color, and volume (the amount of space occupied by a sample). Chemical properties describe the characteristic ability of a substance to react to form new substances; they include its flammability and susceptibility to corrosion. All samples of a pure substance have the same chemical and physical properties. For example, pure copper is always a reddish-brown solid (a physical property) and always dissolves in dilute nitric acid to produce a blue solution and a brown gas (a chemical property).

Physical Property

A *physical property* is a characteristic of a substance that can be observed or measured without changing the identity of the substance. Silver is a shiny metal that conducts electricity very well. It can be molded into thin sheets, a property called malleability. Salt is dull and brittle and conducts electricity when it has been dissolved into water, which it does quite easily. Physical properties of matter include color, hardness, malleability, solubility, electrical conductivity, density, melting point, and boiling point.

For the elements, color does not vary much from one element to the next. The vast majority of elements are colorless, silver, or gray. Some elements do have distinctive colors: sulfur and chlorine are yellow, copper is (of course) copper-colored, and elemental bromine is red. However, density can be a very useful parameter for identifying an element. Of the materials that exist as solids at room temperature, iodine has a very low density compared to zinc, chromium, and tin. Gold has a very high density, as does platinum. Pure water, for example, has a density of 0.998 g/cm^3 at 25°C. The average densities of some common substances are in Table 3.5.1. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will “float.”

Table 3.5.1: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Hardness helps determine how an element (especially a metal) might be used. Many elements are fairly soft (silver and gold, for example) while others (such as titanium, tungsten, and chromium) are much harder. Carbon is an interesting example of hardness. In graphite, (the "lead" found in pencils) the carbon is very soft, while the carbon in a diamond is roughly seven times as hard.



Figure 3.5.1: Pencil (left) and Diamond ring (right). Both are a form of carbon, but exhibit very different physical properties.

Melting and boiling points are somewhat unique identifiers, especially of compounds. In addition to giving some idea as to the identity of the compound, important information can be obtained about the purity of the material.

Chemical Properties

Chemical properties of matter describe its potential to undergo some chemical change or reaction by virtue of its composition. The elements, electrons, and bonds that are present give the matter potential for chemical change. It is quite difficult to define a chemical property without using the word "change". Eventually, after studying chemistry for some time, you should be able to look at the formula of a compound and state some chemical property. For example, hydrogen has the potential to ignite and explode given the right conditions—this is a chemical property. Metals in general have the chemical property of reacting with an acid. Zinc reacts with hydrochloric acid to produce hydrogen gas—this is a chemical property.



Figure 3.5.2: Heavy rust on the links of a chain near the Golden Gate Bridge in San Francisco; it was continuously exposed to moisture and salt spray, causing surface breakdown, cracking, and flaking of the metal. (CC BY-SA 3.0; Marlith).

A chemical property of iron is its capability of combining with oxygen to form iron oxide, the chemical name of rust (Figure 3.5.2). The more general term for rusting and other similar processes is corrosion. Other terms that are commonly used in descriptions of chemical changes are burn, rot, explode, decompose, and ferment. Chemical properties are very useful in identifying substances. However, unlike physical properties, chemical properties can only be observed as the substance is in the process of being changed into a different substance.

Table 3.5.2: Contrasting Physical and Chemical Properties

Physical Properties	Chemical Properties
Gallium metal melts at 30 ^{o}C.	Iron metal rusts.
Mercury is a very dense liquid.	A green banana turns yellow when it ripens.
Gold is shiny.	A dry piece of paper burns.

Example 3.5.1

Which of the following is a chemical property of iron?

a. Iron corrodes in moist air.
b. Density = 7.874 g/cm^3
c. Iron is soft when pure.
d. Iron melts at 1808 K.

Solution

"Iron corrodes in moist air" is the only chemical property of iron from the list.

Exercise 3.5.1A

Which of the following is a physical property of matter?

a. corrosiveness
b. pH (acidity)
c. density
d. flammability

Answer

c

Exercise 3.5.1B

Which of the following is a chemical property?

a. flammability
b. melting point
c. boiling point
d. density

Answer

a

Summary

A physical property is a characteristic of a substance that can be observed or measured without changing the identity of the substance. Physical properties include color, density, hardness, and melting and boiling points. A chemical property describes the ability of a substance to undergo a specific chemical change. To identify a chemical property, we look for a chemical change. A chemical change always produces one or more types of matter that differ from the matter present before the change. The formation of rust is a chemical change because rust is a different kind of matter than the iron, oxygen, and water present before the rust formed.

3.5: Differences in Matter- Physical and Chemical Properties is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **3.5: Differences in Matter- Physical and Chemical Properties** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.6: Changes in Matter - Physical and Chemical Changes

Learning Objectives

- Label a change as chemical or physical.
- List evidence that can indicate a chemical change occurred.

Change is happening all around us all of the time. Just as chemists have classified elements and compounds, they have also classified types of changes. Changes are classified as either physical or chemical changes. Chemists learn a lot about the nature of matter by studying the changes that matter can undergo. Chemists make a distinction between two different types of changes that they study—physical changes and chemical changes.

Physical Change

Physical changes are changes in which no bonds are broken or formed. This means that the same types of compounds or elements that were there at the beginning of the change are there at the end of the change. Because the ending materials are the same as the beginning materials, the properties (such as color, boiling point, etc.) will also be the same. Physical changes involve moving molecules around, but not changing them. Some types of physical changes include:

- Changes of state (changes from a solid to a liquid or a gas and vice versa).
- Separation of a mixture.
- Physical deformation (cutting, denting, stretching).
- Making solutions (special kinds of mixtures).

As an ice cube melts, its shape changes as it acquires the ability to flow. However, its composition does not change. Melting is an example of a physical change. A physical change is a change to a sample of matter in which some properties of the material change, but the identity of the matter does not. When liquid water is heated, it changes to water vapor. However, even though the physical properties have changed, the molecules are exactly the same as before. We still have each water molecule containing two hydrogen atoms and one oxygen atom covalently bonded. When you have a jar containing a mixture of pennies and nickels and you sort the mixture so that you have one pile of pennies and another pile of nickels, you have not altered the identity of the pennies or the nickels—you've merely separated them into two groups. This would be an example of a physical change. Similarly, if you have a piece of paper, you don't change it into something other than a piece of paper by ripping it up. What was paper before you started tearing is still paper when you are done. Again, this is an example of a physical change.



Figure 3.6.1: Ice melting is a physical change. When liquid water (H_2O) freezes into a solid state (ice), it appears changed; however, this change is only physical, as the composition of the constituent molecules is the same: 11.19% hydrogen and 88.81% oxygen by mass. (Public Domain; Moussa).

Physical changes can further be classified as reversible or irreversible. The melted ice cube may be refrozen, so melting is a reversible physical change. Physical changes that involve a change of state are all reversible. Other changes of state include **vaporization** (liquid to gas), **freezing** (liquid to solid), and **condensation** (gas to liquid). Dissolving is also a reversible physical change. When salt is dissolved into water, the salt is said to have entered the aqueous state. The salt may be regained by boiling off the water, leaving the salt behind.

Chemical Change

Chemical changes occur when bonds are broken and/or formed between molecules or atoms. This means that one substance with a certain set of properties (such as melting point, color, taste, etc) is turned into a different substance with different properties. Chemical changes are frequently harder to reverse than physical changes.

One good example of a chemical change is burning a candle. The act of burning paper actually results in the formation of new chemicals (carbon dioxide and water) from the burning of the wax. Another example of a chemical change is what occurs when natural gas is burned in your furnace. This time, on the left there is a molecule of methane, $\mathrm{CH_4}$, and two molecules of oxygen, $\mathrm{O_2}$; on the right are two molecules of water, $\mathrm{H_2O}$, and one molecule of carbon dioxide, $\mathrm{CO_2}$. In this case, not only has the appearance changed, but the structure of the molecules has also changed. The new substances do not have the same chemical properties as the original ones. Therefore, this is a chemical change.



Figure 3.6.2: Burning of wax to generate water and carbon dioxide is a chemical reaction. (CC-SA-BY-3.0; Andrikkos)

We can't actually see molecules breaking and forming bonds, although that's what defines chemical changes. We have to make other observations to indicate that a chemical change has happened. Some of the evidence for chemical change will involve the energy changes that occur in chemical changes, but some evidence involves the fact that new substances with different properties are formed in a chemical change.

Observations that help to indicate chemical change include:

- Temperature changes (either the temperature increases or decreases).
- Light given off.
- Unexpected color changes (a substance with a different color is made, rather than just mixing the original colors together).
- Bubbles are formed (but the substance is not boiling—you made a substance that is a gas at the temperature of the beginning materials, instead of a liquid).
- Different smell or taste (do not taste your chemistry experiments, though!).
- A solid forms if two clear liquids are mixed (look for *floaties*—technically called a precipitate).

Example 3.6.1

Label each of the following changes as a physical or chemical change. Give evidence to support your answer.

a. Boiling water.
b. A nail rusting.
c. A green solution and colorless solution are mixed. The resulting mixture is a solution with a pale green color.
d. Two colorless solutions are mixed. The resulting mixture has a yellow precipitate.

Solution

a. Physical: boiling and melting are physical changes. When water boils, no bonds are broken or formed. The change could be written: $\mathrm{H_2O}\,(l) \rightarrow \mathrm{H_2O}\,(g)$
b. Chemical: The dark grey nail changes color to form an orange flaky substance (the rust); this must be a chemical change. Color changes indicate chemical change. The following reaction occurs: $\mathrm{Fe} + \mathrm{O_2} \rightarrow \mathrm{Fe_2O_3}$

c. Physical: because none of the properties changed, this is a physical change. The green mixture is still green and the colorless solution is still colorless. They have just been spread together. No color *change* occurred or other evidence of chemical change.
d. Chemical: the formation of a precipitate and the color change from colorless to yellow indicate a chemical change.

? Exercise 3.6.1

Label each of the following changes as a physical or chemical change.

a. A mirror is broken.
b. An iron nail corroded in moist air
c. Copper metal is melted.
d. A catalytic converter changes nitrogen dioxide to nitrogen gas and oxygen gas.

Answer a:

physical change

Answer b:

chemical change

Answer c:

physical change

Answer d:

chemical change

Separating Mixtures Through Physical Changes

Homogeneous mixtures (solutions) can be separated into their component substances by physical processes that rely on differences in some physical property, such as differences in their boiling points. Two of these separation methods are distillation and crystallization. Distillation makes use of differences in volatility, a measure of how easily a substance is converted to a gas at a given temperature. A simple distillation apparatus for separating a mixture of substances, at least one of which is a liquid. The most volatile component boils first and is condensed back to a liquid in the water-cooled condenser, from which it flows into the receiving flask. If a solution of salt and water is distilled, for example, the more volatile component, pure water, collects in the receiving flask, while the salt remains in the distillation flask.




Figure 3.6.3: *The Distillation of a Solution of Table Salt in Water. The solution of salt in water is heated in the distilling flask until it boils. The resulting vapor is enriched in the more volatile component (water), which condenses to a liquid in the cold condenser and is then collected in the receiving flask.*

Parts of a distillation setup: Bunsen burner, salt water in distilling flask, condenser with cool water in and warm water out, pure water in receiving flask

Mixtures of two or more liquids with different boiling points can be separated with a more complex distillation apparatus. One example is the refining of crude petroleum into a range of useful products: aviation fuel, gasoline, kerosene, diesel fuel, and lubricating oil (in the approximate order of decreasing volatility). Another example is the distillation of alcoholic spirits such as brandy or whiskey. This relatively simple procedure caused more than a few headaches for federal authorities in the 1920s during the era of Prohibition, when illegal stills proliferated in remote regions of the United States.

Another example for using physical properties to separate mixtures is filtration (Figure 3.6.4). Filtration is any mechanical, physical or biological operation that separates solids from fluids (liquids or gases) by adding a medium through which only the fluid can pass. The fluid that passes through is called the filtrate. There are many different methods of filtration; all aim to attain the separation of substances. Separation is achieved by some form of interaction between the substance or objects to be removed and the filter. The substance that is to pass through the filter must be a fluid, i.e. a liquid or gas. Methods of filtration vary depending on the location of the targeted material, i.e. whether it is dissolved in the fluid phase or suspended as a solid.




Figure 3.6.4: *Filtration for the separation of solids from a hot solution. (CC BY-SA 4.0; Suman6395).*

Summary

- Chemists make a distinction between two different types of changes that they study—physical changes and chemical changes.
- Physical changes are changes that do not alter the identity of a substance.
- Chemical changes are changes that occur when one substance is turned into another substance.
- Chemical changes are frequently harder to reverse than physical changes. Observations that indicate a chemical change has occurred include color change, temperature change, light given off, formation of bubbles, formation of a precipitate, etc.

Contributions & Attributions

-
- Boundless (www.boundless.com)

3.6: Changes in Matter - Physical and Chemical Changes is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **3.6: Changes in Matter - Physical and Chemical Changes** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.7: Conservation of Mass - There is No New Matter

It may seem as though burning destroys matter, but the same amount, or mass, of matter still exists after a campfire as before. Look at Figure 3.7.1 below. It shows that when wood burns, it combines with oxygen and changes not only to ashes, but also to carbon dioxide and water vapor. The gases float off into the air, leaving behind just the ashes. Suppose you had measured the mass of the wood before it burned and the mass of the ashes after it burned. Also suppose you had been able to measure the oxygen used by the fire and the gases produced by the fire. What would you find? The total mass of matter after the fire would be the same as the total mass of matter before the fire.




Figure 3.7.1: Burning is a chemical process. The flames are caused as a result of a fuel undergoing combustion (burning). (CC BY-SA 2.5; Einar Helland Berger for fire and Walter Siegmund for ash).

Law of Conservation of Mass

The law of conservation of mass was created in 1789 by a French chemist, Antoine Lavoisier. The **law of conservation of mass** states that matter cannot be created or destroyed in a chemical reaction. For example, when wood burns, the mass of the soot, ashes, and gases equals the original mass of the charcoal and the oxygen when it first reacted. So the mass of the product equals the mass of the reactant. A reactant is the chemical reaction of two or more elements to make a new substance, and a product is the substance that is formed as the result of a chemical reaction (Video 3.7.1). Matter and its corresponding mass may not be able to be created or destroyed, but can change forms to other substances like liquids, gases, and solids.



Video 3.7.1: This is a nice little demonstration showing the Conservation of Mass in action.

If you witness a 300 kg tree burn to the ground, there are only ashes left after the burn, and all of them together weigh 10 kg. It may make you wonder where the other 290 kg went. The missing 290 kg was released into the atmosphere as smoke, so the only thing left that you can see is the 10 kg of ash. If you know the law of conservation of mass, then you know that the other 290 kg has to go somewhere, because it has to equal the mass of the tree before it burnt down.

Example 3.7.1

If heating 10.0 grams of calcium carbonate ($CaCO_3$) produces 4.4 g of carbon dioxide (CO_2) and 5.6 g of calcium oxide (CaO), show that these observations are in agreement with the law of conservation of mass.

Solution

$$\text{Mass of the reactants} = \text{Mass of the products}$$

$$10.0\ \text{g of } \ce{CaCO3} = 4.4\ \text{g of } \ce{CO2} + 5.6\ \text{g of CaO}$$

$$10.0\ \text{g of reactant} = 10.0\ \text{g of products}$$

Because the mass of the reactant is equal to the mass of the products, the observations are in agreement with the law of conservation of mass.

? Exercise 3.7.1

Potassium hydroxide ($\ce{KOH}$) readily reacts with carbon dioxide ($\ce{CO2}$) to produce potassium carbonate ($\ce{K2CO3}$) and water ($\ce{H2O}$). How many grams of potassium carbonate are produced if 224.4 g of $\ce{KOH}$ reacts with 88.0 g of $\ce{CO2}$? The reaction also produces 36.0 g of water.

Answer

276.4 g of potassium carbonate

The Law is also applicable to both chemical and physical changes. For example, if you have an ice cube that melts into a liquid and you heat that liquid up, it becomes a gas. It will appear to have disappeared, but is still there.

Summary

- Burning and other changes in matter do not destroy matter.
- The mass of matter is always the same before and after the changes occur.
- The law of conservation of mass states that matter cannot be created or destroyed.

Contributions & Attributions

- Binod Shrestha (University of Lorraine)

3.7: Conservation of Mass - There is No New Matter is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **3.7: Conservation of Mass - There is No New Matter** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.E: Chemistry, Matter, and Measurement (Exercises)

1. Does each statement refer to a chemical property or a physical property?
    a. Balsa is a very light wood.
    b. If held in a flame, magnesium metal burns in air.
    c. Mercury has a density of 13.6 g/mL.
    d. Human blood is red.
2. Does each statement refer to a chemical property or a physical property?
    a. The elements sodium and chlorine can combine to make table salt.
    b. The metal tungsten does not melt until its temperature exceeds 3,000°C.
    c. The ingestion of ethyl alcohol can lead to disorientation and confusion.
    d. The boiling point of isopropyl alcohol, which is used to sterilize cuts and scrapes, is lower than the boiling point of water.
3. Define *element*. How does it differ from a compound?
4. Define *compound*. How does it differ from an element?
5. Give two examples of a heterogeneous mixture.
6. Give two examples of a homogeneous mixture.
7. Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.
    a. xenon, a substance that cannot be broken down into chemically simpler components
    b. blood, a substance composed of several types of cells suspended in a salty solution called plasma
    c. water, a substance composed of hydrogen and oxygen
8. Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.
    a. sugar, a substance composed of carbon, hydrogen, and oxygen
    b. hydrogen, the simplest chemical substance
    c. dirt, a combination of rocks and decaying plant matter
9. Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.
    a. air, primarily a mixture of nitrogen and oxygen
    b. ringer's lactate, a standard fluid used in medicine that contains salt, potassium, and lactate compounds all dissolved in sterile water
    c. tartaric acid, a substance composed of carbon, hydrogen, and oxygen
10. Identify each material as an element, a compound, a heterogeneous mixture, or a solution.
    a. equal portions of salt and sand placed in a beaker and shaken up
    b. a combination of beeswax dissolved in liquid hexane
    c. hydrogen peroxide, a substance composed of hydrogen and oxygen
11. What word describes each phase change?
    a. solid to liquid
    b. liquid to gas
    c. solid to gas
12. What word describes each phase change?
    a. liquid to solid
    b. gas to liquid
    c. gas to solid

Answers

1.
    a. physical property

b. chemical property
c. physical property
d. physical property

2.

a. chemical property
b. physical property
c. chemical property
d. physical property

3. An element is a substance that cannot be broken down into chemically simpler components. Compounds can be broken down into simpler substances.

4. A compound is composed of two or more elements combined in a fixed ratio. An element is the simplest chemical substance.

5. a salt and pepper mix and a bowl of cereal (answers will vary)

6. vinegar and rubbing alcohol (answers will vary)

7.

a. element
b. heterogeneous mixture
c. compound

8.

a. compound
b. element
c. heterogeneous mixture

9.

a. solution
b. solution
c. compound

10.

a. heterogeneous mixture
b. solution
c. compound

11.

a. melting or fusion
b. boiling or evaporation
c. sublimation

12.

a. freezing
b. condensation
c. deposition

3.E: Chemistry, Matter, and Measurement (Exercises) is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

4: Atoms, Elements, and the Periodic Table

4.1: Prelude to Elements, Atoms, and the Periodic Table
4.2: The Elements
4.3: Atomic Theory
4.4: Atomic Theory
4.5: Defining Isotopes
4.6: Atomic Masses
4.7: Arrangements of Electrons
4.8: Orbital Shape
4.9: The Periodic Table
4.10: Ions - Losing and Gaining Electrons
4.E: Elements, Atoms, and the Periodic Table (Exercises)

4: Atoms, Elements, and the Periodic Table is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

4.1: Prelude to Elements, Atoms, and the Periodic Table

The hardest material in the human body is tooth enamel. It has to be hard so that our teeth can serve us for a lifetime of biting and chewing; however, tough as it is, tooth enamel is susceptible to chemical attack. Acids found in some foods or made by bacteria that feed on food residues on our teeth are capable of dissolving enamel. Unprotected by enamel, a tooth will start to decay, thus developing cavities and other dental problems.

In the early 1900s, a dentist in Colorado Springs, Colorado, noted that many people who lived in the area had brown-stained teeth that, while unsightly, were surprisingly resistant to decay. After years of study, excess fluorine compounds in the drinking water were discovered to be the cause of both these effects. Research continued, and in the 1930s, the US Public Health Service found that low levels of fluorine in water would provide the benefit of resisting decay without discoloring teeth.

The protective effects of fluorine have a simple chemical explanation. Tooth enamel consists mostly of a mineral called hydroxyapatite, which is composed of calcium, phosphorus, oxygen, and hydrogen. We know now that fluorine combines with hydroxyapatite to make fluorapatite, which is more resistant to acidic decay than hydroxyapatite is. Currently about 50% of the US population drinks water that has some fluorine added (in the form of sodium fluoride, NaF) to reduce tooth decay. This intentional fluoridation, coupled with the use of fluoride-containing toothpastes and improved oral hygiene, has reduced tooth decay by as much as 60% in children. The nationwide reduction of tooth decay has been cited as an important public health advance in history. (Another important advance was the eradication of polio.)

4.1: Prelude to Elements, Atoms, and the Periodic Table is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.0: Prelude to Elements, Atoms, and the Periodic Table** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.2: The Elements

Learning Objectives

- Define a chemical element and give examples of the abundance of different elements.
- Represent a chemical element with a chemical symbol.

An element is a substance that cannot be broken down into simpler chemical substances. There are about 90 naturally occurring elements known on Earth. Using technology, scientists have been able to create nearly 30 additional elements that do not occur in nature. Today, chemistry recognizes 118 elements—some of which were created an atom at a time. Figure 4.2.1 shows some of the chemical elements.



Figure 4.2.1: Samples of Elements. Gold is a yellowish solid, iron is a silvery solid, while mercury is a silvery liquid at room temperature. © Thinkstock

Abundance

The elements vary widely in abundance. In the universe as a whole, the most common element is hydrogen (about 90% of atoms), followed by helium (most of the remaining 10%). All other elements are present in relatively minuscule amounts, as far as we can detect.

Table 4.2.1: Elemental Composition of Earth

Earth's Crust		Earth (overall)	
Element	**Percentage**	**Element**	**Percentage**
oxygen	46.1	iron	34.6
silicon	28.2	oxygen	29.5
aluminum	8.23	silicon	15.2
iron	5.53	magnesium	12.7
calcium	4.15	nickel	2.4
sodium	2.36	sulfur	1.9
magnesium	2.33	all others	3.7
potassium	2.09		
titanium	0.565		
hydrogen	0.14		
phosphorus	0.105		
all others	0.174		

Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 14–17.

On the planet Earth, however, the situation is rather different. Oxygen makes up 46.1% of the mass of Earth's crust (the relatively thin layer of rock forming Earth's surface), mostly in combination with other elements, while silicon makes up 28.2%. Hydrogen,

the most abundant element in the universe, makes up only 0.14% of Earth's crust. Table 4.2.1 lists the relative abundances of elements on Earth as a whole and in Earth's crust. Table 4.2.2 lists the relative abundances of elements in the human body. If you compare Table 4.2.1 and Table 4.2.2, you will find disparities between the percentage of each element in the human body and on Earth. Oxygen has the highest percentage in both cases, but carbon, the element with the second highest percentage in the body, is relatively rare on Earth and does not even appear as a separate entry in Table 4.2.1; carbon is part of the 0.174% representing "other" elements. How does the human body concentrate so many apparently rare elements?

Table 4.2.2: Elemental Composition of a Human Body

Element	Percentage by Mass
oxygen	61
carbon	23
hydrogen	10
nitrogen	2.6
calcium	1.4
phosphorus	1.1
sulfur	0.20
potassium	0.20
sodium	0.14
chlorine	0.12
magnesium	0.027
silicon	0.026
iron	0.006
fluorine	0.0037
zinc	0.0033
all others	0.174

Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 7–24.

The relative amounts of elements in the body have less to do with their abundances on Earth than with their availability in a form we can assimilate. We obtain oxygen from the air we breathe and the water we drink. We also obtain hydrogen from water. On the other hand, although carbon is present in the atmosphere as carbon dioxide, and about 80% of the atmosphere is nitrogen, we obtain those two elements from the food we eat, not the air we breathe.

LOOKING CLOSER: PHOSPHOROUS, THE CHEMICAL BOTTLENECK

There is an element that we need more of in our bodies than is proportionately present in Earth's crust, and *this* element is not easily accessible. Phosphorus makes up 1.1% of the human body but only 0.105% of Earth's crust. We need phosphorus for our bones and teeth, and it is a crucial component of all living cells. Unlike carbon, which can be obtained from carbon dioxide through photosynthesis, there is no phosphorus compound present in our surroundings that can serve as a convenient source. Phosphorus, then, is nature's bottleneck. Its availability limits the amount of life our planet can sustain.

Higher forms of life, such as humans, can obtain phosphorus by selecting a proper diet (plenty of protein); but lower forms of life, such as algae, must absorb it from the environment. When phosphorus-containing detergents were introduced in the 1950s, wastewater from normal household activities greatly increased the amount of phosphorus available to algae and other plant life. Lakes receiving this wastewater experienced sudden increases in growth of algae. When the algae died, concentrations of bacteria that ate the dead algae increased. Because of the large bacterial concentrations, the oxygen content of the water

dropped, causing fish to die in large numbers. This process, called *eutrophication*, is considered a negative environmental impact.

Today, many detergents are made without phosphorus so the detrimental effects of eutrophication are minimized. You may even see statements to that effect on detergent boxes. It can be sobering to realize how much impact a single element can have on life—or the ease with which human activity can affect the environment.

Names and Symbols

Each element has a name. Some of these names date from antiquity, while others are quite new. Today, the names for new elements are proposed by their discoverers but must be approved by the International Union of Pure and Applied Chemistry, an international organization that makes recommendations concerning all kinds of chemical terminology.

> *Today, new elements are usually named after famous scientists.*

The names of the elements can be cumbersome to write in full, especially when combined to form the names of compounds. Therefore, each element name is abbreviated as a one- or two-letter chemical symbol. By convention, the first letter of a chemical symbol is a capital letter, while the second letter (if there is one) is a lowercase letter. The first letter of the symbol is usually the first letter of the element's name, while the second letter is some other letter from the name. Some elements have symbols that derive from earlier, mostly Latin names, so the symbols may not contain any letters from the English name. Table 4.2.3 lists the names and symbols of some of the most familiar elements.

Table 4.2.3: Element Names and Symbols

aluminum	Al	magnesium	Mg
argon	Ar	manganese	Mn
arsenic	As	mercury	Hg*
barium	Ba	neon	Ne
bismuth	Bi	nickel	Ni
boron	B	nitrogen	N
bromine	Br	oxygen	O
calcium	Ca	phosphorus	P
carbon	C	platinum	Pt
chlorine	Cl	potassium	K*
chromium	Cr	silicon	Si
copper	Cu*	silver	Ag*
fluorine	F	sodium	Na*
gold	Au*	strontium	Sr
helium	He	sulfur	S
hydrogen	H	tin	Sn*
iron	Fe	tungsten	W†
iodine	I	uranium	U
lead	Pb*	zinc	Zn

*The symbol comes from the Latin name of element.

†The symbol for tungsten comes from its German name—*wolfram*.

lithium	Li	zirconium	Zr

*The symbol comes from the Latin name of element.

†The symbol for tungsten comes from its German name—*wolfram*.

Element names in languages other than English are often close to their Latin names. For example, gold is *oro* in Spanish and *or* in French (close to the Latin *aurum*), tin is *estaño* in Spanish (compare to *stannum*), lead is *plomo* in Spanish and *plomb* in French (compare to *plumbum*), silver is *argent* in French (compare to *argentum*), and iron is *fer* in French and *hierro* in Spanish (compare to *ferrum*). The closeness is even more apparent in pronunciation than in spelling.

Example 4.2.1

Write the chemical symbol for each element without consulting Table 4.2.3 "Element Names and Symbols".

a. bromine
b. boron
c. carbon
d. calcium
e. gold

Answer a

Br

Answer b

B

Answer c

C

Answer d

Ca

Answer e

Au

Exercise 4.2.1

Write the chemical symbol for each element without consulting Table 4.2.3.

a. manganese
b. magnesium
c. neon
d. nitrogen
e. silver

Answer a

Mn

Answer b

Mg

Answer c

Ne

Answer d

N

Answer e

Ag

Example 4.2.2

What element is represented by each chemical symbol?

a. Na
b. Hg
c. P
d. K
e. I

Answer a

sodium

Answer b

mercury

Answer c

phosphorus

Answer d

potassium

Answer e

iodine

Exercise 4.2.2

What element is represented by each chemical symbol?

a. Pb
b. Sn
c. U
d. O
e. F

Answer a

lead

Answer b

tin

Answer c

uranium

Answer d

oxygen

Answer e

fluorine

Concept Review Exercises

1. What is an element?
2. Give some examples of how the abundance of elements varies.
3. Why are chemical symbols so useful? What is the source of the letter(s) for a chemical symbol?

Answers

1. An element is the basic chemical building block of matter; it is the simplest chemical substance.
2. Elements vary from being a small percentage to more than 30% of the atoms around us.
3. Chemical symbols are useful to concisely represent the elements present in a substance. The letters usually come from the name of the element.

Key Takeaways

- All matter is composed of elements.
- Chemical elements are represented by a one- or two-letter symbol.

4.2: The Elements is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.1: The Elements** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.3: Atomic Theory

Learning Objectives

- Explain how all matter is composed of atoms.
- Describe the modern atomic theory.
- Recognize which elements exist as diatomic molecules.

Take some aluminum foil. Cut it in half. Now you have two smaller pieces of aluminum foil. Cut one of the pieces in half again. Cut one of those smaller pieces in half again. Continue cutting, making smaller and smaller pieces of aluminum foil.

It should be obvious that the pieces are still aluminum foil; they are just becoming smaller and smaller. But how far can you take this exercise, at least in theory? Can you continue cutting the aluminum foil into halves forever, making smaller and smaller pieces? Or is there some limit, some absolute smallest piece of aluminum foil? (Thought experiments like this—and the conclusions based on them—were debated as far back as the fifth century BC.)

The modern atomic theory, proposed about 1803 by the English chemist John Dalton (Figure 4.3.1), is a fundamental concept that states that all elements are composed of atoms. Previously, we defined an atom as the smallest part of an element that maintains the identity of that element. Individual atoms are extremely small; even the largest atom has an approximate diameter of only 5.4×10^{-10} m. With that size, it takes over 18 million of these atoms, lined up side by side, to equal the width of your little finger (about 1 cm).




Figure 4.3.1 John Dalton was an English scientist who enunciated the modern atomic theory.

Most elements in their pure form exist as **individual atoms**. For example, a macroscopic chunk of iron (Fe) metal is composed, microscopically, of individual atoms. Some elements, however, exist as groups of atoms called molecules. Several important elements exist as two-atom combinations and are called **diatomic molecules**. In representing a diatomic molecule, we use the symbol of the element and include the ***subscript*** "2" to indicate that two atoms of that element are joined together. The elements that exist as diatomic molecules are hydrogen (H_2), nitrogen (N_2), fluorine (F_2), oxygen (O_2), iodine (I_2), chlorine (Cl_2) and bromine (Br_2). A few other elements can exist as 3-atom molecules like ozone (O_3) and 4-atom molecules like phosphorus (P_4). The most common form of the element sulfur is composed of molecules that consist of eight atoms of sulfur; its molecular formula is S_8 (Figure 4.3.2).




Figure 4.3.2: A molecule of sulfur is composed of eight sulfur atoms and is therefore written as S_8. It can be represented as (a) a structural formula, (b) a ball-and-stick model, and (c) a space-filling model. Sulfur atoms are represented by yellow spheres.

Figure A shows eight sulfur atoms, symbolized with the letter S, that are bonded to each other to form an octagon. Figure B shows a 3-D, ball-and-stick model of the arrangement of the sulfur atoms. The shape is clearly not octagonal as it is represented in the structural formula. Figure C is a space-filling model that shows each sulfur atom is partially embedded into the sulfur atom it bonds with.

It is important to note that a subscript following a symbol and a number in front of a symbol do not represent the same thing; for example, H_2 and 2H represent distinctly different species. H_2 is a molecular formula; it represents a diatomic molecule of hydrogen, consisting of two atoms of the element that are chemically bonded together. The expression 2H, on the other hand, indicates two separate hydrogen atoms that are **not** combined as a unit. The expression $2H_2$ represents two molecules of diatomic hydrogen (Figure 4.3.3).




Figure 4.3.3: The symbols H, 2H, H_2, and $2H_2$ represent very different entities.

This figure shows four diagrams. The diagram for H shows a single, white sphere and is labeled one H atom. The diagram for 2 H shows two white spheres that are not bonded together. It is labeled 2 H atoms. The diagram for H subscript 2 shows two white spheres bonded together. It is labeled one H subscript 2 molecule. The diagram for 2 H subscript 2 shows two sets of bonded, white spheres. It is labeled 2 H subscript 2 molecules.

Example 4.3.1

Write the chemical formula of the following elements:

a. oxygen
b. carbon
c. potassium

Solution

Memorizing the diatomic molecules is worthwhile for our future endeavors. A mnemonic device to help in the memorization of the diatomic elements is as follows: **H**ave **N**o **F**ear **O**f **I**ce **C**old **B**eer. (hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine).

a. Oxygen is one of the seven diatomic molecular elements. Its chemical formula is O_2.
b. Carbon is monatomic, hence its formula is C.
c. Potassium is monatomic hence its formula is K.

rcise

Write the chemical formula of the following elements:

a. hydrogen
b. nitrogen
c. neon

Answer a

Hydrogen is one of the seven diatomic molecular elements. Its chemical formula is H_2.

Answer b

Nitrogen is one of the seven diatomic molecular elements. Its chemical formula is N_2.

Answer c

Neon is a monatomic element, hence its formula is Ne.

Looking Closer: Atomic Theory

Dalton's ideas are called the *modern* atomic theory because the concept of atoms is very old. The Greek philosophers Leucippus and Democritus originally introduced atomic concepts in the fifth century BC. (The word *atom* comes from the Greek word *atomos*, which means "indivisible" or "uncuttable.") Dalton had something that the ancient Greek philosophers didn't have, however; he had experimental evidence, such as the formulas of simple chemicals and the behavior of gases. In the 150 years or so before Dalton, natural philosophy had been maturing into modern science, and the scientific method was being used to study nature. So when Dalton announced a modern atomic theory, he was proposing a fundamental theory to describe many previous observations of the natural world; he was not just participating in a philosophical discussion.

Concept Review Exercises

1. What is the modern atomic theory?
2. What are atoms?

Answers

1. The modern atomic theory states that all matter is composed of atoms.
2. Atoms are the smallest parts of an element that maintain the identity of that element.

Key Takeaways

- **Atoms** are the ultimate building blocks of all matter.
- The modern atomic theory establishes the concepts of **atoms** and how they compose matter.
- The **diatomic** elements are hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine.

4.3: Atomic Theory is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.2: Atomic Theory** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.4: Atomic Theory

Learning Objectives

- State the modern atomic theory.
- Learn how atoms are constructed.
- Describe the locations, charges, and masses of the three main subatomic particles.
- Determine the number of protons and electrons in an atom.
- Define atomic mass unit (amu).

The smallest piece of an element that maintains the identity of that element is called an **atom**. Individual atoms are extremely small. It would take about fifty million atoms in a row to make a line that is 1 cm long. The period at the end of a printed sentence has several million atoms in it. Atoms are so small that it is difficult to believe that all matter is made from atoms-but it is.

The concept that atoms play a fundamental role in chemistry is formalized by the modern atomic theory, first stated by John Dalton, an English scientist, in 1808. It consists of three parts:

1. All matter is composed of atoms.
2. Atoms of the same element are the same; atoms of different elements are different.
3. Atoms combine in whole-number ratios to form compounds.

These concepts form the basis of chemistry. Although the word *atom* comes from a Greek word that means "indivisible," we understand now that atoms themselves are composed of smaller parts called *subatomic particles*. The first part to be discovered was the electron, a tiny subatomic particle with a negative charge. It is often represented as e^-, with the right superscript showing the negative charge. Later, two larger particles were discovered. The proton, a subatomic particle with a positive charge. is a more massive (but still tiny) subatomic particle with a positive charge, represented as p. The neutron is a subatomic particle with about the same mass as a proton but no charge. It is represented as either n. We now know that all atoms of all elements are composed of electrons, protons, and (with one exception) neutrons. Table 4.4.1 summarizes the properties of these three subatomic particles.

Even though electrons, protons, and neutrons are all types of subatomic particles, they are not all the same size. When you compare the masses of electrons, protons, and neutrons, what you find is that electrons have an extremely small mass, compared to either protons or neutrons. On the other hand, the masses of protons and neutrons are fairly similar, although technically, the mass of a neutron is slightly larger than the mass of a proton. Because protons and neutrons are so much more massive than electrons, almost all of the mass of any atom comes from the nucleus, which contains all of the neutrons and protons.

Table 4.4.1: Properties of the Three Subatomic Particles

Particle	Symbol	Mass (amu)	Relative Mass (proton = 1)	Relative Charge	Location
proton	p	1	1	+1	inside the nucleus
electron	e^-	5.45×10^{-4}	0.00055	−1	outside nucleus
neutron	n	1	1	0	inside the nucleus

The third column shows the masses of the three subatomic particles in "atomic mass units." An **atomic mass unit (amu)** is defined as one-twelfth the mass of a carbon-12 atom. Atomic mass units (amu) are useful, because, as you can see, the mass of a proton and the mass of a neutron are almost exactly 11 in this unit system.

How are these particles arranged in atoms? They are not arranged at random. Experiments by Ernest Rutherford in England in the 1910s pointed to a nuclear model with atoms that has the protons and neutrons in a central nucleus with the electrons in orbit about the nucleus. The relatively massive protons and neutrons are collected in the center of an atom, in a region called the nucleus of the atom (plural *nuclei*). The electrons are outside the nucleus and spend their time orbiting in space about the nucleus. (Figure 4.4.1).




Figure 4.4.1: The Structure of the Atom.Atoms have protons and neutrons in the center, making the nucleus, while the electrons orbit the nucleus.

The modern atomic theory states that atoms of one element are the same, while atoms of different elements are different. What makes atoms of different elements different? The fundamental characteristic that all atoms of the same element share is the *number of protons*. All atoms of hydrogen have one and only one proton in the nucleus; all atoms of iron have 26 protons in the nucleus. This number of protons is so important to the identity of an atom that it is called the atomic number. The number of protons in an atom is the atomic number of the element (Z). Thus, hydrogen has an atomic number of 1, while iron has an atomic number of 26. Each element has its own characteristic atomic number.

Atoms of the same element can have different numbers of neutrons, however. Atoms of the same element (i.e., atoms with the same number of protons) with different numbers of neutrons are called **isotopes**. The sum of the number of protons and neutrons in the nucleus is called the mass number of the isotope. We will discuss isotopes and their symbols more in section 4.5 of this chapter.

Neutral atoms have the same number of electrons as they have protons, so their overall charge is zero. Negative and positive charges of equal magnitude cancel each other out. This means that the negative charge on an electron perfectly balances the positive charge on the proton. In other words, a neutral atom must have exactly one electron for every proton. If a neutral atom has 1 proton, it must have 1 electron. If a neutral atom has 2 protons, it must have 2 electrons. If a neutral atom has 10 protons, it must have 10 electrons. You get the idea. In order to be neutral, an atom must have the same number of electrons and protons. If the number of electrons and protons are not equal, the atom is called an **ion**. Ions and ionic compounds will be discussed in detail in section 4.10 and chapter 5.

Example 4.4.1:

a. The most common carbon atoms have six protons and six neutrons in their nuclei. What are the atomic number and the mass number of these carbon atoms?
b. An isotope of uranium has an atomic number of 92 and a mass number of 235. What are the number of protons and neutrons in the nucleus of this atom?

Solution

a. If a carbon atom has six protons in its nucleus, its atomic number is 6. If it also has six neutrons in the nucleus, then the mass number is 6 + 6, or 12.

b. If the atomic number of uranium is 92, then that is the number of protons in the nucleus. Because the mass number is 235, then the number of neutrons in the nucleus is 235 − 92, or 143.

Exercise 4.4.1

The number of protons in the nucleus of a tin atom is 50, while the number of neutrons in the nucleus is 68. What are the atomic number and the mass number of this isotope?

Answer

Atomic number = 50, mass number = 118

When referring to an atom, we simply use the element's name: the term *sodium* refers to the element as well as an atom of sodium. But it can be unwieldy to use the name of elements all the time, and therefore scientists use the chemical symbols discussed in section 1.2 and 4.2.The elements are grouped together in a special chart called the periodic table of all the elements. A simple periodic table is shown in Figure 4.4.2, while one may view a more extensive periodic table. The elements on the periodic table are listed in order of ascending atomic number. We will learn more about how the periodic table is organized as we move through the semester. One immediate use of the periodic table helps us identify metals and nonmetals. Nonmetals are in the upper right corner of the periodic table, on one side of the heavy line splitting the right-hand part of the chart. All other elements are metals.




Figure 4.4.1: A Simple Periodic Table

Exercise 4.4.2

How many protons are in $^{23}_{11}\text{Na}$

Answer

11 protons

Key Takeaways

- Chemistry is based on the modern atomic theory, which states that all matter is composed of atoms.
- Atoms themselves are composed of protons, neutrons, and electrons.
- Each element has its own atomic number, which is equal to the number of protons in its nucleus.
- Isotopes of an element contain different numbers of neutrons.
- Elements are represented by an atomic symbol.
- The periodic table is a chart that organizes all the elements.

4.4: Atomic Theory is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

4.5: Defining Isotopes

Learning Objectives

- Define and differentiate between the atomic number and the mass number of an element.
- Explain how isotopes differ from one another.

Now that we know how atoms are generally constructed, what do atoms of any particular element look like? How many protons, neutrons, and electrons are in a specific kind of atom? First, if an atom is electrically neutral overall, then the number of protons equals the number of electrons. Because these particles have the same but opposite charges, equal numbers cancel out, producing a neutral atom.

Atomic Number

In the 1910s, experiments with x-rays led to this useful conclusion: the magnitude of the positive charge in the nucleus of every atom of a particular element is the same. In other words, all atoms of the same element have the same number of protons. Furthermore, different elements have a different number of protons in their nuclei, so the number of protons in the nucleus of an atom is characteristic of a particular element. This discovery was so important to our understanding of atoms that the number of protons in the nucleus of an atom is called the atomic number (Z).

For example, hydrogen has the atomic number 1; all hydrogen atoms have 1 proton in their nuclei. Helium has the atomic number 2; all helium atoms have 2 protons in their nuclei. There is no such thing as a hydrogen atom with 2 protons in its nucleus; a nucleus with 2 protons would be a helium atom. The atomic number *defines* an element. Table 4.5.1 lists some common elements and their atomic numbers. Based on its **atomic number**, you can determine the **number of protons in the nucleus** of an atom. The largest atoms have over 100 protons in their nuclei.

Table 4.5.1: Some Common Elements and Their Atomic Numbers

Element	Atomic Number	Element	Atomic Nmbers
aluminum (Al)	13	magnesium (Mg)	12
americium (Am)	95	manganese (Mn)	25
argon (Ar)	18	mercury (Hg)	80
barium (Ba)	56	neon (Ne)	10
beryllium (Be)	4	nickel (Ni)	28
bromine (Br)	35	nitrogen (N)	7
calcium (Ca)	20	oxygen (O)	8
carbon (C)	6	phosphorus (P)	15
chlorine (Cl)	17	platinum (Pt)	78
chromium (Cr)	24	potassium (K)	19
cesium (Cs)	55	radon (Rn)	86
fluorine (F)	9	silver (Ag)	47
gallium (Ga)	31	sodium (Na)	11
gold (Au)	79	strontium (Sr)	38
helium (He)	2	sulfur (S)	16
hydrogen (H)	1	titanium (Ti)	22

Element	Atomic Number	Element	Atomic Nmbers
iron (Fe)	26	tungsten (W)	74
iodine (I)	53	uranium (U)	92
lead (Pb)	82	zinc (Zn)	30
lithium (Li)	3	zirconium (Zr)	40

Example 4.5.1

What is the number of protons in the nucleus of each element?

a. aluminum
b. iron
c. carbon

Answer a

According to Table 2.4.1, aluminum has an atomic number of 13. Therefore, every aluminum atom has 13 protons in its nucleus.

Answer b

Iron has an atomic number of 26. Therefore, every iron atom has 26 protons in its nucleus.

Answer c

Carbon has an atomic number of 6. Therefore, every carbon atom has 6 protons in its nucleus.

Exercise 4.5.1

What is the number of protons in the nucleus of each element? Use Table 2.4.1.

a. sodium
b. oxygen
c. chlorine

Answer a

Sodium has 11 protons in its nucleus.

Answer b

Oxygen has 8 protons in its nucleus.

Answer c

Chlorine has 17 protons in its nucleus

How many electrons are in an atom? Previously we said that for an electrically neutral atom, the **number of electrons equals the number of protons**, so the total opposite charges cancel. Thus, the **atomic number** of an element also gives the **number of electrons** in an atom of that element. (Later we will find that some elements may gain or lose electrons from their atoms, so those atoms will no longer be electrically neutral. Thus we will need a way to differentiate the number of electrons for those elements.)

Example 4.5.2

How many electrons are present in the atoms of each element?

a. sulfur
b. tungsten

c. argon

Answer a

The atomic number of sulfur is 16. Therefore, in a neutral atom of sulfur, there are 16 electrons.

Answer b

The atomic number of tungsten is 74. Therefore, in a neutral atom of tungsten, there are 74 electrons.

Answer c

The atomic number of argon is 18. Therefore, in a neutral atom of argon, there are 18 electrons.

? Exercise 4.5.2

How many electrons are present in the atoms of each element?

a. magnesium
b. potassium
c. iodine

Answer a

Mg has 12 electrons.

Answer b

K has 19 electrons.

Answer c

I has 53 electrons.

Isotopes

How many neutrons are in atoms of a particular element? At first it was thought that the number of neutrons in a nucleus was also characteristic of an element. However, it was found that atoms of the same element can have *different* numbers of neutrons. Atoms of the same element (i.e., same atomic number, Z) that have different numbers of neutrons are called isotopes. For example, 99% of the carbon atoms on Earth have 6 neutrons and 6 protons in their nuclei; about 1% of the carbon atoms have 7 neutrons in their nuclei. Naturally occurring carbon on Earth, therefore, is actually a mixture of isotopes, albeit a mixture that is 99% carbon with 6 neutrons in each nucleus.

An important series of isotopes is found with hydrogen atoms. Most hydrogen atoms have a nucleus with only a single proton. About 1 in 10,000 hydrogen nuclei, however, also has a neutron; this particular isotope is called *deuterium*. An extremely rare hydrogen isotope, *tritium*, has 1 proton and 2 neutrons in its nucleus. Figure 4.5.1 compares the three isotopes of hydrogen.




Figure 4.5.1: Isotopes of Hydrogen. Most hydrogen atoms have only a proton in the nucleus (a). A small amount of hydrogen exists as the isotope deuterium, which has one proton and one neutron in its nucleus (b). A tiny amount of the hydrogen isotope tritium, with one proton and two neutrons in its nucleus, also exists on Earth (c). The nuclei and electrons are proportionately much smaller than depicted here.

The discovery of isotopes required a minor change in Dalton's atomic theory. Dalton thought that all atoms of the same element were exactly the same.

Most elements exist as mixtures of isotopes. In fact, there are currently over 3,500 isotopes known for all the elements. When scientists discuss individual isotopes, they need an efficient way to specify the number of neutrons in any particular nucleus. The *mass number* (A) of an atom is the sum of the numbers of protons and neutrons in the nucleus. Given the mass number for a nucleus (and knowing the atomic number of that particular atom), you can determine the number of neutrons by subtracting the atomic number from the mass number.

A simple way of indicating the mass number of a particular isotope is to list it as a superscript on the left side of an element's symbol. Atomic numbers are often listed as a subscript on the left side of an element's symbol. Thus, we might see

$$\begin{matrix}\text{mass number}\longrightarrow\\ \text{atomic number}\longrightarrow\end{matrix}{}^{56}_{26}\text{Fe} \tag{4.5.1}$$

which indicates a particular isotope of iron. The 26 is the atomic number (which is the same for all iron atoms), while the 56 is the mass number of the isotope. To determine the number of neutrons in this isotope, we subtract 26 from 56: 56 − 26 = 30, so there are 30 neutrons in this atom.

Example 4.5.3

How many protons and neutrons are in each atom?

a. $^{35}_{17}\text{Cl}$
b. $^{127}_{53}\text{I}$

Answer a

In $^{35}_{17}\text{Cl}$ there are 17 protons, and 35 − 17 = 18 neutrons in each nucleus.

Answer b

In $^{127}_{53}\text{I}$ there are 53 protons, and 127 − 53 = 74 neutrons in each nucleus.

Exercise 4.5.3

How many protons and neutrons are in each atom?

a. $^{197}_{79}\text{Au}$
b. $^{23}_{11}\text{Na}$

Answer a

In $^{197}_{79}\text{Au}$ there are 79 protons, and 197 − 79 = 118 neutrons in each nucleus.

Answer b

In $^{23}_{11}\text{Na}$ there are 11 protons, and 23 − 11 = 12 neutrons in each nucleus.

It is not absolutely necessary to indicate the atomic number as a subscript because each element has its own unique atomic number. Many isotopes are indicated with a superscript only, such as ^{13}C or ^{235}U. You may also see isotopes represented in print as, for example, carbon-13 or uranium-235.

Summary

The atom consists of discrete particles that govern its chemical and physical behavior. Each atom of an element contains the same number of protons, which is the **atomic number (Z)**. Neutral atoms have the same number of electrons and protons. Atoms of an element that contain different numbers of neutrons are called **isotopes**. Each isotope of a given element has the same atomic number but a different **mass number (A)**, which is the **sum of the numbers of protons and neutrons**.

Almost all of the mass of an atom is from the total protons and neutrons contained within a tiny (and therefore very dense) nucleus. The majority of the volume of an atom is the surrounding space in which the electrons reside. A representation of a carbon-12 atom is shown below in Figure 4.5.2.




Figure 4.5.2: Formalism used for identifying specific nuclide (any particular kind of nucleus)

The atomic symbol, X, is the abbreviation used to represent an atom in chemical formulas. The mass number, A, is the number of protons and neutrons in the atom, which is to the left of the X. The atomic number, Z, is the number of protons in the atom, which is to the left of the X and below the A.

Concept Review Exercises

1. Why is the atomic number so important to the identity of an atom?
2. What is the relationship between the number of protons and the number of electrons in an atom?
3. How do isotopes of an element differ from each other?
4. What is the mass number of an element?

Answers

1. The atomic number defines the identity of an element. It describes the number of protons in the nucleus.
2. In an electrically neutral atom, the number of protons equals the number of electrons.
3. Isotopes of an element have the same number of protons but have different numbers of neutrons in their nuclei.
4. The mass number is the sum of the numbers of protons and neutrons in the nucleus of an atom.

Key Takeaways

- Each element is identified by its atomic number. The atomic number provides the element's location on the periodic table
- The isotopes of an element have different masses and are identified by their mass numbers.

4.5: Defining Isotopes is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.4: Nuclei of Atoms** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.6: Atomic Masses

Learning Objectives

- Define atomic mass and atomic mass unit.
- Calculate atomic mass.

Even though atoms are very tiny pieces of matter, they have mass. Their masses are so small, however, that chemists often use a unit other than grams to express them—the atomic mass unit.

Atomic Mass Unit

The atomic mass unit (abbreviated u, although amu is also used) is defined as 1/12 of the mass of a ^{12}C atom:

$$1 \text{ u} = \frac{1}{12} \text{ the mass of } {}^{12}\text{C atom} \tag{4.6.1}$$

It is equal to 1.661×10^{-24} g.

Masses of other atoms are expressed with respect to the atomic mass unit. For example, the mass of an atom of ^{1}H is 1.008 u, the mass of an atom of ^{16}O is 15.995 u, and the mass of an atom of ^{32}S is 31.97 u. Note, however, that these masses are for particular isotopes of each element. Because most elements exist in nature as a mixture of isotopes, any sample of an element will actually be a mixture of atoms having slightly different masses (because neutrons have a significant effect on an atom's mass). How, then, do we describe the mass of a given element? By calculating an average of an element's atomic masses, weighted by the natural abundance of each isotope, we obtain a weighted average mass called the atomic mass (also commonly referred to as the *atomic weight*) of an element.

Atomic Mass is the Weighted Average Mass of Isotopes

As stated above, most elements occur naturally as a mixture of two or more isotopes. Listed below (Table 4.6.1) are the naturally occurring isotopes of selected elements along with the percent natural abundance of each.

Table 4.6.1: Atomic Masses and Percent Abundances of Some Natural Isotopes

Element	Isotope (Symbol)	Percent Natural Abundance	Atomic Mass (amu)	Average Atomic Mass (amu)
Hydrogen	$^{1}_{1}H$	99.985	1.0078	1.0079
	$^{2}_{1}H$	0.015	2.0141	
	$^{3}_{1}H$	negligible	3.0160	
Carbon	$^{12}_{6}C$	98.89	12.000	12.011
	$^{13}_{6}C$	1.11	13.003	
	$^{14}_{6}C$	trace	14.003	
Oxygen	$^{16}_{8}O$	99.759	15.995	15.999
	$^{17}_{8}O$	0.037	16.995	
	$^{18}_{8}O$	0.204	17.999	
Chlorine	$^{35}_{17}Cl$	75.77	34.969	35.453
	$^{38}_{17}Cl$	24.23	36.966	
Copper	$^{63}_{29}Cu$	69.17	62.930	63.546
	$^{65}_{29}Cu$	30.83	64.928	

For some elements, one particular isotope is much more abundant than any other isotopes. For example, naturally occurring hydrogen is nearly all hydrogen-1, and naturally occurring oxygen is nearly all oxygen-16. For many other elements, however, more than one isotope may exist in substantial quantities. Chlorine (atomic number 17) is yellowish-green toxic gas. About three quarters of all chlorine atoms have 18 neutrons, giving those atoms a mass number of 35. About one quarter of all chlorine atoms have 20 neutrons, giving those atoms a mass number of 37. Were you to simply calculate the arithmetic average of the precise atomic masses, you would get approximately 36.

$$\frac{34.969\,u + 36.966\,u}{2} = 35.968\,u$$

As you can see, the average atomic mass given in the last column of the table above (35.453) is significantly lower. Why? The reason is that we need to take into account the natural abundance percentages of each isotope in order to calculate what is called the **weighted average**. The atomic mass *of an element is the weighted average of the atomic masses of the naturally occurring isotopes of that element.*

$$0.7577\,(34.969\,u) + 0.2423\,(36.966\,u) = 35.453\,u$$

The weighted average is determined by multiplying the percent of natural abundance by the actual mass of the isotope. This is repeated until there is a term for each isotope. For chlorine, there are only two naturally occurring isotopes so there are only two terms.

Atomic mass = (%1)(mass 1) + (%2)(mass 2) + ⋯

Another example: oxygen exists as a mixture that is 99.759% ^{16}O, 0.037% ^{17}O and 0.204% ^{18}O. The atomic mass of oxygen (use percent natural abundance data from Table 2.5.1) would be calculated as follows:

Atomic mass = (%1)(mass 1) + (%2)(mass 2) + (%3)(mass 3)

$$0.99759\,(15.995u) + 0.00037\,(16.995u) + 0.00204\,(17.999u) = 15.999u$$

To confirm your answer, compare the calculated value to the weighted mass displayed on the periodic table.

Example 4.6.1

Calculate the atomic mass of oxygen. Oxygen exists as a mixture of 3 isotopes. Their respective masses and natural abundance are shown below.

- ^{16}O: 15.995 u (99.759%)
- ^{17}O: 16.995 u (0.037%)
- ^{18}O: 17.999 u (0.204%)

Solution

Multiply the isotope abundance by the actual mass of the isotope, and then sum up the products.

$$0.99759\,(15.995\,u) + 0.00037\,(16.995\,u) + 0.00204\,(17.999\,u) = 15.999\,u$$

rcise

Calculate the atomic mass of copper. Copper exists as a mixture of 2 isotopes. Their respective masses and natural abundance are shown below.


- ^{63}Cu: 62.930 u (69.17%)
- ^{65}Cu: 64.928 u (30.83%)

Answer

63.546 u

The **atomic mass** of each element is found under the element symbol in the **periodic table.** Examples are shown below. The atomic mass of tin (Sn) is 118.71 u while the atomic mass of carbon (C) is 12.011 u. On the other hand, the **atomic number (Z)** of

each element is found **above** the atomic symbol.



Atomic mass indicated on entries of the Periodic Table. (public Domain; Pubchem)

The **periodic table** is found in this link:

https://pubchem.ncbi.nlm.nih.gov/periodic-table/png/Periodic_Table_of_Elements_w_Atomic_Mass_PubChem.png

Example 4.6.2: Mass of Carbon

What is the average mass of a carbon atom in grams? The atomic mass is found in the Periodic Table. Please use two decimal places.

Solution

This is a simple one-step conversion, similar to conversions we did in Chapter 1. We use the fact that 1 u = 1.661×10^{-24} g:

$$12.01 \ \cancel{\text{u}} \times \frac{1.661 \times 10^{-24} \text{ g}}{1 \ \cancel{\text{u}}} = 1.995 \times 10^{-23} \text{ g}$$

This is an extremely small mass, which illustrates just how small individual atoms are.

Exercise 4.6.2: Mass of Tin

What is the average mass of a tin atom in grams? The average atomic mass is found in the Periodic Table. Please use two decimal places.

Answer

$$118.71 \ \cancel{\text{u}} \times \frac{1.661 \times 10^{-24} \text{ g}}{1 \ \cancel{\text{u}}} = 1.972 \times 10^{-22} \text{ g}$$

Concept Review Exercises

1. Define atomic mass. Why is it considered a weighted average?
2. What is an atomic mass unit?

Answers

1. The atomic mass is an average of an element's atomic masses, weighted by the natural abundance of each isotope of that element. It is a weighted average because different isotopes have different masses.
2. An atomic mass unit is 1/12th of the mass of a ^{12}C atom.

Key Takeaway

- Atoms have a mass that is based largely on the number of protons and neutrons in their nucleus.
- The atomic mass of each element in the Periodic Table is the weighted average of the mass of all its isotopes.

4.6: Atomic Masses is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.5: Atomic Masses** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.7: Arrangements of Electrons

Learning Objectives

- Describe how electrons are grouped within atoms.

Although we have discussed the general arrangement of subatomic particles in atoms, we have said little about how electrons occupy the space about the nucleus. Do they move around the nucleus at random, or do they exist in some ordered arrangement?

The modern theory of electron behavior is called quantum mechanics. It makes the following statements about electrons in atoms:

- Electrons in atoms can have only certain specific energies. We say that the energies of the electrons are quantized.
- Electrons are organized according to their energies into sets called **shells** (labeled by the principle quantum number, ***n***). Generally the higher the energy of a shell, the farther it is (on average) from the nucleus. Shells do not have specific, fixed distances from the nucleus, but an electron in a higher-energy shell will spend more time farther from the nucleus than does an electron in a lower-energy shell.
- Shells are further divided into subsets of electrons called **subshells**. The first shell has only one subshell, the second shell has two subshells, the third shell has three subshells, and so on. The subshells of each shell are labeled, in order, with the letters *s*, *p*, *d*, and *f*. Thus, the first shell has only a single *s* subshell (called **1*s***), the second shell has **2*s*** and **2*p*** subshells, the third shell has **3*s***, **3*p***, and **3*d*** and so forth.

Table 2.6.1: Shells and Subshells

Shell	Number of Subshells	Names of Subshells
1	1	*1s*
2	2	*2s* and *2p*
3	3	*3s*, *3p* and *3d*
4	4	*4s*, *4p*, *4d* and *4f*

- Different subshells hold a different maximum number of electrons. **Any *s* subshell can hold up to 2 electrons; *p*, 6; *d*, 10; and *f*, 14.**

Table 2.6.2 Number of Electrons

Subshell	Maximum Number of Electrons
s	2
p	6
d	10
f	14

It is the arrangement of electrons into shells and subshells that most concerns us here, so we will focus on that.

We use numbers to indicate which shell an electron is in. As shown in Table **2.6.1**, the first shell, closest to the nucleus and with the lowest-energy electrons, is shell 1. This first shell has only one subshell, which is labeled 1*s* and can hold a maximum of 2 electrons. We combine the shell and subshell labels when referring to the organization of electrons about a nucleus and use a superscript to indicate how many electrons are in a subshell. Thus, because a hydrogen atom has its single electron in the *s* subshell of the first shell, we use $\mathbf{1}s^{\mathbf{1}}$ to describe the electronic structure of hydrogen. This structure is called an electron configuration. Electron configurations are shorthand descriptions of the arrangements of electrons in atoms. The electron configuration of a hydrogen atom is spoken out loud as “one-ess-one.”

Helium atoms have 2 electrons. Both electrons fit into the 1*s* subshell because *s* subshells can hold up to 2 electrons; therefore, the electron configuration for helium atoms is $\mathbf{1}s^{\mathbf{2}}$ (spoken as “one-ess-two”).

The 1*s* subshell cannot hold 3 electrons (because an *s* subshell can hold a maximum of 2 electrons), so the electron configuration for a lithium atom cannot be $1s^3$. Two of the lithium electrons can fit into the 1*s* subshell, but the third electron must go into the

second shell. The second shell has two subshells, *s* and *p*, which fill with electrons in that order. The 2*s* subshell holds a maximum of 2 electrons, and the 2*p* subshell holds a maximum of 6 electrons. Because lithium's final electron goes into the 2*s* subshell, we write the electron configuration of a lithium atom as $\mathbf{1}s^{\mathbf{2}}\mathbf{2}s^{\mathbf{1}}$**.** The shell diagram for a lithium atom is shown below. The shell closest to the nucleus (first shell) has 2 dots representing the 2 electrons in **1*s***, while the outermost shell (**2*s***) has 1 electron.




Figure 2.6.1 Shell diagram of lithium (Li) atom.

The next largest atom, beryllium, has 4 electrons, so its electron configuration is $1s^2 2s^2$. Now that the 2*s* subshell is filled, electrons in larger atoms start filling the 2*p* subshell. Thus, the electron configurations for the next six atoms are as follows:

- B: $1s^2 2s^2 2p^1$
- C: $1s^2 2s^2 2p^2$
- N: $1s^2 2s^2 2p^3$
- O: $1s^2 2s^2 2p^4$
- F: $1s^2 2s^2 2p^5$
- Ne: $1s^2 2s^2 2p^6$

With neon, the 2*p* subshell is completely filled. Because the second shell has only two subshells, atoms with more electrons now must begin the third shell. The third shell has three subshells, labeled *s*, *p*, and *d*. The *d* subshell can hold a maximum of 10 electrons. The first two subshells of the third shell are filled in order—for example, the electron configuration of aluminum, with 13 electrons, is $1s^2 2s^2 2p^6 3s^2 3p^1$. However, a curious thing happens after the 3*p* subshell is filled: the 4*s* subshell begins to fill before the 3*d* subshell does. In fact, the exact ordering of subshells becomes more complicated at this point (after argon, with its 18 electrons), so we will not consider the electron configurations of larger atoms.

A fourth subshell, the *f* subshell, is needed to complete the electron configurations for all elements. An *f* subshell can hold up to 14 electrons.

Electron filling always starts with **1*s***, the subshell closest to the nucleus. Next is **2*s*, 2*p*, 3*s*, 3*p*, 4*s*, 3*d*, 4*p***, etc., shown in the electron shell filling order diagram in Figure **2.6.2**. Follow each red arrow in order from top to bottom. The subshells you reach along each arrow give the ordering of filling of subshells in larger atoms.




Figure 2.6.2 *The order of electron filling in an atom.*

Example 4.7.1: Electronic Configuration of Phosphorus Atoms

What is the electron configuration of a neutral phosphorus atom?

Solution

A neutral phosphorus atom has 15 electrons. Two electrons can go into the 1*s* subshell, 2 can go into the 2*s* subshell, and 6 can go into the 2*p* subshell. That leaves 5 electrons. Of those 5 electrons, 2 can go into the 3*s* subshell, and the remaining 3

electrons can go into the 3*p* subshell. Thus, the electron configuration of neutral phosphorus atoms is $1s^2 2s^2 2p^6 3s^2 3p^3$.

Exercise 4.7.1: Electronic Configuration of Chlorine Atoms

What is the electron configuration of a neutral chlorine atom?

Answer

A neutral chlorine atom has 17 electrons. Two electrons can go into the 1*s* subshell, 2 can go into the 2*s* subshell, and 6 can go into the 2*p* subshell. That leaves 7 electrons. Of those 7 electrons, 2 can go into the 3*s* subshell, and the remaining 5 electrons can go into the 3*p* subshell. Thus, the electron configuration of neutral chlorine atoms is $1s^2 2s^2 2p^6 3s^2 3p^5$.

Valence Electrons

In the study of chemical reactivity, we will find that the electrons in the outermost principal energy level are very important and so they are given a special name. **Valence electrons** are the electrons in the highest occupied principal energy level of an atom. In the second period elements, the two electrons in the $1s$ sublevel are called **inner-shell electrons** and are not involved directly in the element's reactivity or in the formation of compounds. Lithium has a single electron in the second principal energy level and so we say that lithium has one valence electron. Beryllium has two valence electrons. How many valence electrons does boron have? You must recognize that the second principal energy level consists of both the $2s$ and the $2p$ sublevels and so the answer is three. In fact, the number of valence electrons goes up by one for each step across a period until the last element is reached. Neon, with its configuration ending in $2s^2 2p^6$, has eight valence electrons.

A chemical reaction results from electron removal, electron addition, or electron sharing of the valence electrons of the different atoms. The path a specific element will take depends on where the electrons are in the atom and how many there are. Thus, it is convenient to separate electrons into two groups. Valence shell electrons (or, more simply, the *valence electrons*) are the electrons in the highest-numbered shell, or valence shell, while core electrons are the electrons in lower-numbered shells. We can see from the electron configuration of a carbon atom—$1s^2 2s^2 2p^2$—that it has 4 valence electrons ($2s^2 2p^2$) and 2 core electrons ($1s^2$). You will see in the next chapters that the chemical properties of elements are determined by the number of valence electrons.

Example 4.7.2: Counting Valence Electrons in Phosphorus Atoms

From the electron configuration of neutral phosphorus atoms in Example 4.7.1, how many valence electrons and how many core electrons does a neutral phosphorus atom have?

Solution

The highest-numbered shell is the third shell, which has 2 electrons in the 3*s* subshell and 3 electrons in the 3*p* subshell. That gives a total of 5 electrons, so neutral phosphorus atoms have **5 valence electrons**. The 10 remaining electrons, from the first and second shells, are core electrons.

Exercise 4.7.2: Counting Valence Electrons in Chlorine Atoms

From the electron configuration of neutral chlorine atoms (Exercise 4.7.1), how many valence electrons and how many core electrons does a neutral chlorine atom have?

Answer

The highest-numbered shell is the third shell, which has 2 electrons in the 3*s* subshell and 5 electrons in the 3*p* subshell. That gives a total of 7 electrons, so neutral chlorine atoms have **7 valence electrons**. The 10 remaining electrons, from the first and second shells, are core electrons.

Exercise 4.7.2 Counting Valence Electrons in Sodium Atoms


What is the electron configuration of a neutral atom of sodium (Na)? How many core electrons are there? How many valence electrons? How do you draw the shell diagram of sodium atom?

Answer

Sodium (Na) has atomic number 11, hence, 11 electrons. The electron configuration is: $1s^2 2s^2 2p^6 3s^1$. This means the **first shell (1s) has 2 electrons**. The **second shell** (2s and 2p) has a **total of 8 electrons**. And, the **third (last) shell has 1 electron**.

The first and second shells comprise the core (inner) electrons = 2 + 8 = **10 electrons**. The outermost **(valence) has 1 electron**.

The **shell diagram** of the Na atom is shown here. The shell nearest the nucleus (first shell) has 2 electrons, the second shell has 8 electrons and the last (outermost) shell has 1 electron (**2.8.1**)



Concept Review Exercises

1. How are electrons organized in atoms?
2. What information does an electron configuration convey?
3. What is the difference between core electrons and valence electrons?

Answers

1. Electrons are organized into shells and subshells around nuclei.
2. The electron configuration states the arrangement of electrons in shells and subshells.
3. Valence electrons are in the highest-numbered shell; all other electrons are core electrons.

Key Takeaway



- Electrons are organized into shells and subshells about the nucleus of an atom.
- The valence electrons determine the reactivity of an atom.

Exercises

2.6: Arrangements of Electrons

1. What is the maximum number of electrons that can fit in an *s* subshell? Does it matter what shell the *s* subshell is in?
2. What is the maximum number of electrons that can fit in a *p* subshell? Does it matter what shell the *p* subshell is in?
3. What is the maximum number of electrons that can fit in a *d* subshell? Does it matter what shell the *d* subshell is in?
4. What is the maximum number of electrons that can fit in an *f* subshell? Does it matter what shell the *f* subshell is in?
5. What is the electron configuration of a carbon atom?
6. What is the electron configuration of a sulfur atom?
7. What is the valence shell electron configuration of a calcium atom?
8. What is the valence shell electron configuration of a selenium atom?
9. What atom has the electron configuration $1s^2 2s^2 2p^5$?
10. What atom has the electron configuration $1s^2 2s^2 2p^6 3s^2 3p^3$?
11. Draw a representation of the electronic structure of an oxygen atom.
12. Draw a representation of the electronic structure of a phosphorus atom.
13. A potassium atom has ____ core electrons and ____ valence electrons.
14. A silicon atom has ____ core electrons and ____ valence electrons.

Answers

1. 2; no
2. 6; no
3. 10; no
4. 14; no
5. $1s^2 2s^2 2p^2$
6. $1s^2 2s^2 2p^6 3s^2 3p^4$
7. $4s^2$
8. $4s^2 4s^4$
9. fluorine
10. phosphorus
11. 
12. 
13. 18; 1
14. 10; 4

4.7: Arrangements of Electrons is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

4.8: Orbital Shape

The shell diagram example shown in the previous section, also known as a Bohr model, is a useful way to begin understanding how electrons fill orbitals. However, it is not quite this simple. Think back to section 4.4 on atomic theory, and how it discussed the electrons being in a cloud. Figure 4.4.1 showed no defined orbitals circling the nucleus in rings, but rather an overall area around the nucleus where the electrons were located. In an electron cloud model the electrons still exist within orbitals, but the shapes of the orbitals are more like a probability map of the location of electrons.

The reason for this is that particles are always in motion. What this means, is that the electrons do not sit still. Instead, they move around throughout the electron cloud within the various orbitals. This can occur because of the negative charge the electrons carry. Like charges repel one another, and opposite charges attract. You can think of this like magnets. If you bring two north ends of a magnet near one another, they will push each other away (repel), however if you bring a north and a south end of a magnet together they will attract and stick to each other. Electrons will repel one another because of their like charges. Since they are always in motion, they continually will change paths to keep from interacting too closely with other electrons. At the same time, the electrons stay within the electron cloud of the atom because the positive charge of the protons in the nucleus attract the electrons.

This isn't to say electrons cannot leave an electron cloud. We have already discussed previously that ions form when an atom does not have equal numbers of electrons and protons. We will discuss this idea more later. For now, the focus is on understanding basic electron behavior.

At this point you might be wondering about how orbitals are shaped if not in rings around the nucleus? The answer to this is that the shape of an orbital is dependent upon how many electrons can be held within the orbital. We already know that s-orbitals hold two electrons. The shape of this orbital is a sphere. The p-orbital (which holds a maximum of 6 electrons) is a peanut or dumbbell shape, and the d-orbital (holding a maximum of 10 electrons) is a cross peanut or cross dumbbell shape. Figure 4.8.1 shows the breakdown for basic orbital shapes, and the number of suborbitals within each orbital type. Each suborbital can hold a maximum of two electrons. You are only required to know s, p, and d orbital shapes for this class, but it is important to know other orbital shapes exist. Figure 4.8.1 below shows up through the f-orbital.




Figure 4.8.1: Orbital shapes and their orientation along various axes.

What you will notice is that for any orbital that holds more than two electrons there are various points in space in which those electrons may be held. Electrons arrange themselves into orbitals based on the amount of energy exists within the atom. The more electrons that exist within an atom, the more orbitals that exist. To fit multiple of each type of orbitals, the orbital space expands to overlap other orbitals. Since the electrons cannot interact, any electron in higher energy orbitals is pushed farther out from the nucleus. In the end, electrons closer to the nucleus are held more tightly by the positive charge of the protons. Meanwhile, electrons further from the nucleus have less attraction to the positively charged protons at the center of the atom. It is the electrons farthest from the nucleus that we call **valence electrons**. It is the valence electrons that interact with other atoms causing the formation of ionic and covalent compounds.

Note

You may find the video below from Khan Academy helpful.

Video 4.8.1 walks through using the periodic table to write out ground state electron configurations for orbitals, instead of using the chart shown in section 4.7.



4.8: Orbital Shape is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

4.9: The Periodic Table

Learning Objectives

- Explain how elements are organized into the periodic table.
- Describe how some characteristics of elements relate to their positions on the periodic table.

In the 19th century, many previously unknown elements were discovered, and scientists noted that certain sets of elements had similar chemical properties. For example, chlorine, bromine, and iodine react with other elements (such as sodium) to make similar compounds. Likewise, lithium, sodium, and potassium react with other elements (such as oxygen) to make similar compounds. Why is this so?

In 1864, Julius Lothar Meyer, a German chemist, organized the elements by atomic mass and grouped them according to their chemical properties. Later that decade, Dmitri Mendeleev, a Russian chemist, organized all the known elements according to similar properties. He left gaps in his table for what he thought were undiscovered elements, and he made some bold predictions regarding the properties of those undiscovered elements. When elements were later discovered whose properties closely matched Mendeleev's predictions, his version of the table gained favor in the scientific community. Because certain properties of the elements repeat on a regular basis throughout the table (that is, they are periodic), it became known as the periodic table.

> *Mendeleev had to list some elements out of the order of their atomic masses to group them with other elements that had similar properties.*

The periodic table is one of the cornerstones of chemistry because it organizes all the known elements on the basis of their chemical properties. A modern version is shown in Figure 4.9.1. Most periodic tables provide additional data (such as atomic mass) in a box that contains each element's symbol. The elements are listed in order of atomic number.




Figure 4.9.1: Modern Periodic Table. (Public Domain; PubChem modified Leticia Colmenares).

Features of the Periodic Table

Elements that have similar chemical properties are grouped in columns called **groups** (or families). As well as being numbered, some of these groups have names—for example, *alkali metals* (the first column of elements), *alkaline earth metals* (the second column of elements), *halogens* (the next-to-last column of elements), and *noble gases* (the last column of elements).

Each row of elements on the periodic table is called a **period**. Periods have different lengths; the first period has only 2 elements (hydrogen and helium), while the second and third periods have 8 elements each. The fourth and fifth periods have 18 elements each, and later periods are so long that a segment from each is removed and placed beneath the main body of the table.

Metals, Nonmetals and Metalloids

Certain elemental properties become apparent in a survey of the periodic table as a whole. Every element can be classified as either a metal, a nonmetal, or a semimetal, as shown in Figure 4.9.2. A metal is a substance that is shiny, typically (but not always) silvery in color, and an excellent conductor of electricity and heat. Metals are also malleable (they can be beaten into thin sheets) and ductile (they can be drawn into thin wires). A nonmetal is typically dull and a poor conductor of electricity and heat. Solid nonmetals are also very brittle. As shown in Figure 4.9.2, **metals** occupy the left three-fourths of the periodic table, while **nonmetals** (except for hydrogen) are clustered in the upper right-hand corner of the periodic table. The elements with properties intermediate between those of metals and nonmetals are called **semimetals** (or **metalloids**). Elements adjacent to the **bold zigzag** line in the right-hand portion of the periodic table have semimetal properties.




Figure 4.9.2: Types of Elements. Elements are either metals, nonmetals, or semimetals. Each group is located in a different part of the periodic table.

Example 4.9.1

Based on its position in the periodic table, do you expect selenium (Se) to be a metal, a nonmetal, or a semimetal?

Solution

The atomic number of selenium is 34, which places it in period 4 and group 16. In Figure 2.7.1, selenium lies above and to the right of the diagonal line marking the boundary between metals and nonmetals, so it should be a nonmetal. Note, however, that because selenium is close to the metal-nonmetal dividing line, it would not be surprising if selenium were similar to a semimetal in some of its properties.

rcise

Based on its location in the periodic table, do you expect indium (In) to be a nonmetal, a metal, or a semimetal?

Answer

metal

Representative, Transition and Inner-transition

Another way to categorize the elements of the periodic table is shown in Figure 4.9.3. The first two columns on the left and the last six columns on the right are called the **main group** or **representative** elements. The ten-column block between these columns

contains the **transition metals**. The two rows beneath the main body of the periodic table contain the inner transition metals. The elements in these two rows are also referred to as, respectively, the **lanthanide metals** and the **actinide metals**.




Figure 4.9.3: Special Names for Sections of the Periodic Table. Some sections of the periodic table have special names. The elements lithium, sodium, potassium, rubidium, cesium, and francium are collectively known as alkali metals.

Alkali metals are the first column. Alkaline earth metals are the second. Halogens are the second to last column. Noble gases are the last column.

To Your Health: Transition Metals in the Body

Most of the elemental composition of the human body consists of main group elements. The first element appearing on the list that is not a main group element is iron, at 0.006 percentage by mass. Because iron has relatively massive atoms, it would appear even lower on a list organized in terms of percent by *atoms* rather than percent by mass.

Iron is a transition metal. Transition metals have interesting chemical properties, partially because some of their electrons are in *d* subshells. The chemistry of iron makes it a key component in the proper functioning of red blood cells.

Red blood cells are cells that transport oxygen from the lungs to cells of the body and then transport carbon dioxide from the cells to the lungs. Without red blood cells, animal respiration as we know it would not exist. The critical part of the red blood cell is a protein called *hemoglobin*. Hemoglobin combines with oxygen and carbon dioxide, transporting these gases from one location to another in the body. Hemoglobin is a relatively large molecule, with a mass of about 65,000 u.

The crucial atom in the hemoglobin protein is iron. Each hemoglobin molecule has four iron atoms, which act as binding sites for oxygen. It is the presence of this particular transition metal in your red blood cells that allows you to use the oxygen you inhale.

Other transition metals have important functions in the body, despite being present in low amounts. Zinc is needed for the body's immune system to function properly, as well as for protein synthesis and tissue and cell growth. Copper is also needed for several proteins to function properly in the body. Manganese is needed for the body to metabolize oxygen properly. Cobalt is a necessary component of vitamin B-12, a vital nutrient. These last three metals are not listed explicitly in Table 2.1.2, so they are present in the body in very small quantities. However, even these small quantities are required for the body to function properly.

As previously noted, the periodic table is arranged so that elements with similar chemical behaviors are in the same group. Chemists often make general statements about the properties of the elements in a group using descriptive names with historical origins. For example, the elements of Group 1 are known as the alkali metals, Group 2 are the alkaline earth metals, Group 17 are the halogens, and Group 18 are the noble gases.

Group 1: The Alkali Metals

The alkali metals are lithium, sodium, potassium, rubidium, cesium, and francium. Hydrogen is unique in that it is generally placed in Group 1, but it is not a metal. The compounds of the alkali metals are common in nature and daily life. One example is table salt (sodium chloride); lithium compounds are used in greases, in batteries, and as drugs to treat patients who exhibit manic-depressive, or bipolar, behavior. Although lithium, rubidium, and cesium are relatively rare in nature, and francium is so unstable and highly radioactive that it exists in only trace amounts, sodium and potassium are the seventh and eighth most abundant elements in Earth's crust, respectively.

Group 2: The Alkaline Earth Metals

The alkaline earth metals are beryllium, magnesium, calcium, strontium, barium, and radium. Beryllium, strontium, and barium are rare, and radium is unstable and highly radioactive. In contrast, calcium and magnesium are the fifth and sixth most abundant elements on Earth, respectively; they are found in huge deposits of limestone and other minerals.

Group 17: The Halogens

The halogens are fluorine, chlorine, bromine, iodine, and astatine. The name halogen is derived from the Greek words for "salt forming," which reflects that all the halogens react readily with metals to form compounds, such as sodium chloride and calcium chloride (used in some areas as road salt).

Compounds that contain the fluoride ion are added to toothpaste and the water supply to prevent dental cavities. Fluorine is also found in Teflon coatings on kitchen utensils. Although chlorofluorocarbon propellants and refrigerants are believed to lead to the depletion of Earth's ozone layer and contain both fluorine and chlorine, the latter is responsible for the adverse effect on the ozone layer. Bromine and iodine are less abundant than chlorine, and astatine is so radioactive that it exists in only negligible amounts in nature.

Group 18: The Noble Gases

The noble gases are helium, neon, argon, krypton, xenon, and radon. Because the noble gases are composed of only single atoms, they are called monatomic. At room temperature and pressure, they are **unreactive gases**. Because of their lack of reactivity, for many years they were called **inert gases** or rare gases. However, the first chemical compounds containing the noble gases were prepared in 1962. Although the noble gases are relatively minor constituents of the atmosphere, natural gas contains substantial amounts of helium. Because of its low reactivity, argon is often used as an unreactive (inert) atmosphere for welding and in light bulbs. The red light emitted by neon in a gas discharge tube is used in neon lights.

To Your Health: Radon

Radon is an invisible, odorless noble gas that is slowly released from the ground, particularly from rocks and soils whose uranium content is high. Because it is a noble gas, radon is not chemically reactive. Unfortunately, it is radioactive, and increased exposure to it has been correlated with an increased lung cancer risk.

Because radon comes from the ground, we cannot avoid it entirely. Moreover, because it is denser than air, radon tends to accumulate in basements, which if improperly ventilated can be hazardous to a building's inhabitants. Fortunately, specialized ventilation minimizes the amount of radon that might collect. Special fan-and-vent systems are available that draw air from below the basement floor, before it can enter the living space, and vent it above the roof of a house.

After smoking, radon is thought to be the second-biggest *preventable* cause of lung cancer in the United States. The American Cancer Society estimates that 10% of all lung cancers are related to radon exposure. There is uncertainty regarding what levels of exposure cause cancer, as well as what the exact causal agent might be (either radon or one of its breakdown products, many of which are also radioactive and, unlike radon, not gases). The US Environmental Protection Agency recommends testing every floor below the third floor for radon levels to guard against long-term health effects.

Why do elements in a given group have similar properties?

The periodic table is organized on the basis of similarities in elemental properties, but what explains these similarities? It turns out that the shape of the periodic table reflects the filling of subshells with electrons, as shown in Figure 4.9.4. Starting with the first period and going from left to right, the table reproduces the order of filling of the electron subshells in atoms. Furthermore, elements in the same *group* share the same valence shell electron configuration. For example, all elements in the first column have a single *s* electron in their valence shells, so their electron configurations can be described as ns^1 (where *n* represents the shell number). This last observation is crucial. Chemistry is largely the result of interactions between the valence electrons of different atoms. Thus, atoms that have **the same valence shell electron configuration** will have **similar chemistry.**




Figure 4.9.4 The Shape of the Periodic Table. The shape of the periodic table reflects the order in which electron shells and subshells fill with electrons.

Example 4.9.1

Using the variable *n* to represent the number of the valence electron shell, write the valence shell electron configuration for each group.

a. the alkaline earth metals
b. the column of elements headed by carbon

Answer a

The alkaline earth metals are in the second column of the periodic table. This column corresponds to the *s* subshell being filled with 2 electrons. Therefore, the valence shell electron configuration is ns^2.

Answer b

The electron configuration of carbon is $1s^2 2s^2 2p^2$. Its valence shell electron configuration is $2s^2 2p^2$. Every element in the same column should have a similar valence shell electron configuration, which we can represent as $ns^2 np^2$.

Exercise 4.9.1

Using the variable *n* to represent the number of the valence electron shell, write the valence shell electron configuration for each group.

a. the halogens
b. the column of elements headed by oxygen

Answer a

The halogens are in the 17th column (or Group 7A) of the periodic table. This column corresponds to the *p* subshell being filled with 5 electrons. Therefore, the valence shell electron configuration is $ns^2 np^5$.

Answer b

The column headed by O is the 16th column (or Group 6A). This column corresponds to the *p* subshell being filled with 4 electrons. Therefore, the valence shell electron configuration is $ns^2 np^4$.

Valence Electrons and Group Number

The number of valence electrons of an element can be determined by the periodic table group (vertical column) in which the element is categorized. With the exception of groups 3–12 (the transition metals), the units digit of the group number identifies how many valence electrons are associated with a neutral atom of an element listed under that particular column.

Table 4.9.1. The Group number and the number of valence electrons.

Periodic table group	Valence electrons
Group 1 (I) (alkali metals)	1

Group 2 (II) (alkaline earth metals)	2
Groups 3-12 (transition metals)	2*
Group 13 (III) (boron group)	3
Group 14 (IV) (carbon group)	4
Group 15 (V) (pnictogens)	5
Group 16 (VI) (chalcogens)	6
Group 17 (VII) (halogens)	7
Group 18 (VIII or 0) (noble gases)	8**

** The general method for counting valence electrons is generally not useful for transition metals.*

*** Except for helium, which has only two valence electrons.*

Atomic Radius

The periodic table is useful for understanding atomic properties that show periodic trends. One such property is the atomic radius (Figure 4.9.5). The atomic radius is defined as one-half the distance between the nuclei of identical atoms that are bonded together. The units for atomic radii are picometers, equal to 10^{-12} meters. As an example, the internuclear distance between the two hydrogen atoms in an H_2 molecule is measured to be 74 pm. Therefore, the atomic radius of a hydrogen atom is $\frac{74}{2} = 37$ pm.

As mentioned earlier, the higher the shell number, the farther from the nucleus the electrons in that shell are likely to be. In other words, the size of an atom is generally determined by the number of the valence electron shell. Therefore, as we go down a column on the periodic table, the atomic radius increases. As we go *across* a period on the periodic table, however, electrons are being added to the *same* valence shell; meanwhile, more protons are being added to the nucleus, so the positive charge of the nucleus is increasing. The increasing positive charge attracts the electrons more strongly, pulling them closer to the nucleus. Consequently, as we go across a period, from left to right, the atomic radius decreases. These trends are seen clearly in Figure 4.9.5




Figure 4.9.5 Trends on the Periodic Table. Atomic radii of the representative elements measured in picometers. The relative sizes of the atoms show several trends with regard to the structure of the periodic table. Atoms become larger going down a group and going from right to left across a period.

Example 4.9.2

Using the periodic table (rather than Figure 4.9.5), which atom is larger?

a. N or Bi
b. Mg or Cl

Answer a

Bi is below N in Group 5A in the periodic table and has electrons in higher-numbered shells, hence we expect that Bi atoms are larger than N atoms.

Answer b

Both Mg and Cl are in period 3 of the periodic table, but Cl lies farther to the right. Therefore we expect Mg atoms to be larger than Cl atoms.

Exercise 4.9.2

Using the periodic table (rather than Figure 4.9.5), which atom is larger?

a. Li or F
b. Na or K

Answer a

Li and F are on the same period, but F lies farther to the right. Therefore, we expect Li to be larger than F atoms.

Answer b

K lies below Na in Group 1A, hence has more electron shells, making it larger than Na.

Career Focus: Clinical Chemist

Clinical chemistry is the area of chemistry concerned with the analysis of body fluids to determine the health status of the human body. Clinical chemists measure a variety of substances, ranging from simple elements such as sodium and potassium to complex molecules such as proteins and enzymes, in blood, urine, and other body fluids. The absence or presence, or abnormally low or high amounts, of a substance can be a sign of some disease or an indication of health. Many clinical chemists use sophisticated equipment and complex chemical reactions in their work, so they not only need to understand basic chemistry, but also be familiar with special instrumentation and how to interpret test results.

Concept Review Exercises

1. How are the elements organized into the periodic table?
2. Looking at the periodic table, where do the following elements appear?
   a. the metals
   b. the nonmetals
   c. the halogens
   d. the transition metals
   e. the noble gases
3. Describe the trends in atomic radii as related to an element’s position on the periodic table.

Answers

1. Elements are organized in order of increasing atomic number.
2. a. the left three-quarters of the periodic table (to the left of the zigzag band)
   b. the right quarter of the periodic table (to the right of the zigzag band)
   c. the next-to-last column of the periodic table
   d. the middle section of the periodic table

e. the last column of the periodic table

3. As you go across the periodic table, atomic radii decrease; as you go down the periodic table, atomic radii increase.

Key Takeaways

- The chemical elements are arranged in a chart called the periodic table.
- Some characteristics of the elements are related to their position on the periodic table.
- The number of valence electrons of an element can be determined by the group (vertical column) number in the Periodic Table. Elements with the same number of valence electrons have similar chemical properties.

4.9: The Periodic Table is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **2.7: The Periodic Table** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.10: Ions - Losing and Gaining Electrons

Learning Objectives

- Define the two types of ions.

Atoms tend to lose, gain, or share some valance electrons, making bonds to acquire the electron configuration of the nearest noble gas, i.e., usually eight valence electrons for the main group elements, called an octet. Most atoms do not have eight electrons in their valence electron shell. Some atoms have only a few electrons in their outer shell, while others lack only one or two electrons to have an octet. In cases where an atom has three or fewer valence electrons, the atom may lose those valence electrons quite easily until what remains is a lower shell that contains an octet. Atoms that lose electrons acquire a positive charge because they are left with fewer negatively charged electrons to balance the positive charges of the protons in the nucleus. Positively charged ions are called **cations**. Most metals become cations when they make ionic compounds.

Cations

A neutral sodium atom is likely to achieve an octet in its outermost shell by losing its one valence electron.

$$\mathrm{Na} \rightarrow \mathrm{Na^+} + \mathrm{e^-}$$

The cation produced in this way, $\mathrm{Na^+}$, is called the sodium ion to distinguish it from the element. The outermost shell of the sodium ion is the second electron shell, which has eight electrons in it. The octet rule has been satisfied. Figure 4.10.1 is a graphical depiction of this process.




Figure 4.10.1: The Formation of a Sodium Ion. On the left, a sodium atom has 11 electrons. On the right, the sodium ion only has 10 electrons and a 1+ charge. Neutral sodium atom on left has 11 protons and 11 electrons. Sodium ion on right has 11 protons and 10 electrons, with a +1 overall charge.

Anions

Some atoms have nearly eight electrons in their valence shell and can gain additional valence electrons until they have an octet. When these atoms gain electrons, they acquire a negative charge because they now possess more electrons than protons. Negatively charged ions are called **anions**. Most nonmetals become anions when they make ionic compounds.

A neutral chlorine atom has seven electrons in its outermost shell. Only one more electron is needed to achieve an octet in chlorine's valence shell. (In table salt, this electron comes from the sodium atom.)

$$\mathrm{e^-} + \mathrm{Cl} \longrightarrow \mathrm{Cl^-}$$

In this case, the ion has the same outermost shell as the original atom, but now that shell has eight electrons in it. Once again, the octet rule has been satisfied. The resulting anion, $\mathrm{Cl^-}$, is called the chloride ion; note the slight change in the suffix (*-ide* instead of *-ine*) to create the name of this anion. Figure 4.10.2is a graphical depiction of this process.




Figure 4.10.2: The Formation of a Chlorine Ion. On the left, the chlorine atom has 17 electrons. On the right, the chloride ion has 18 electrons and has a 1− charge. Neutral chlorine atom on left has 17 protons and 17 electrons. Sodium ion on right has 17 protons and 18 electrons, with a -1 overall charge.

> *The names for positive and negative ions are pronounced CAT-eye-ons and ANN-eye-ons, respectively.*

In many cases, elements that belong to the same group (vertical column) on the periodic table form ions with the same charge because they have the same number of valence electrons. Thus, the periodic table becomes a tool for remembering the charges on many ions. For example, all ions made from alkali metals, the first column on the periodic table, have a 1+ charge. Ions made from alkaline earth metals, the second group on the periodic table, have a 2+ charge. On the other side of the periodic table, the next-to-last column, the halogens, form ions having a 1− charge. Figure 4.10.3 shows how the charge on many ions can be predicted by the location of an element on the periodic table. Note the convention of first writing the number and then the sign on a ion with multiple charges. The barium cation is written Ba^{2+}, not Ba^{+2}.




Figure 4.10.3: Predicting Ionic Charges. The charge that an atom acquires when it becomes an ion is related to the structure of the periodic table. Within a group (family) of elements, atoms form ions of a certain charge.

4.10: Ions - Losing and Gaining Electrons is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

- **4.7: Ions - Losing and Gaining Electrons** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

4.E: Elements, Atoms, and the Periodic Table (Exercises)

02.E: Exercises II

4.E: Elements, Atoms, and the Periodic Table (Exercises) is shared under a Public Domain license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

5: Compounds- Ionic and Covalent

Template:HideTOC

There are many substances that exist as two or more atoms connected together so strongly that they behave as a single particle. These multiatom combinations are called moleculesThe smallest part of a substance that has the physical and chemical properties of that substance.. A molecule is the smallest part of a substance that has the physical and chemical properties of that substance. In some respects, a molecule is similar to an atom. A molecule, however, is composed of more than one atom.

5.1: Sugar and Salt
5.2: Compounds Display Constant Composition
5.3: Chemical Formulas- How to Represent Compounds
5.4: A Molecular View of Elements and Compounds
5.5: Writing Formulas for Ionic Compounds
5.6: Nomenclature- Naming Compounds
5.7: Naming Ionic Compounds
5.8: Naming Molecular Compounds
5.9: Naming Acids
5.10: Nomenclature Summary
5.11: Formula Mass- The Mass of a Molecule or Formula Unit
5.12: Moles, Molecular Formulas, and Calculating Empirical Formulas for Compounds

Template:HideTOC

5: Compounds- Ionic and Covalent is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

5.1: Sugar and Salt

Sodium chloride also known as table salt, is an ionic compound with the chemical formula NaCl, representing a 1:1 ratio of sodium and chloride ions. It is commonly used as a condiment and food preservative. Salt can be created by adding two very reactive elements together: sodium Na (s) metal and chlorine $\mathrm{Cl_2}$ (g) gas.

$$2\,\mathrm{Na(s)} + \mathrm{Cl_2(g)} \rightarrow 2\,\mathrm{NaCl(s)} \tag{5.1.1}$$

The element sodium (Figure 5.1.1*a*) is a very reactive metal; given the opportunity, it will react with the sweat on your hands and form sodium hydroxide, which is a very corrosive substance. The element chlorine (Figure 5.1.1*b*) is a pale yellow, corrosive gas that should not be inhaled due to its poisonous nature. Bring these two hazardous substances together, however, and they react to make the ionic compound sodium chloride (Figure 5.1.1*c*), known simply as salt.




Figure 5.1.1: Sodium + Chlorine = Sodium Chloride (a) Sodium is a very reactive metal. (b) Chlorine is a pale yellow, noxious gas. (c) Together, sodium and chlorine make sodium chloride—salt—which is necessary for our survival. Source: Photo on the left courtesy of reenhorn1 and photo in the center courtesy of Benjah-bmm27. Photo on the right © Thinkstock.

Salt is necessary for life. $\mathrm{Na^+}$ ions are one of the main ions in the human body and are necessary to regulate the fluid balance in the body. $\mathrm{Cl^-}$ ions are necessary for proper nerve function and respiration. Both of these ions are supplied by salt. The taste of salt is one of the fundamental tastes; salt is probably the most ancient flavoring known, and one of the few rocks we eat. Clearly when the elemental sodium and chlorine combine (Equation 5.1.1), the resulting salt product has radically different properties (both physical and chemical). This reaction is spectacular to observed (Video 5.1.1).



***Video* 5.1.1:** *Making Table Salt using Sodium Metal and Chlorine gas*

Another compound is sugar, which is the generic name for sweet, soluble carbohydrates, many of which are used in food. Sugar has the chemical formulate $\mathrm{C_{12}H_{22}O_{11}}$ and is constructed from different elements than salt: carbon, hydrogen and oxygen. While sugar qualitatively resembles table salt (often confused in the kitchen), they two have distinctly different physical and chemical properties. There are various types of sugar derived from different sources. While sugar is made with carbon, hydrogen and oxygen, it is considerably hard to syntheses from the constituent elements than table salt (Equation 5.1.1). However, the thermal decomposition is considerably easiler and can be represented as as a dehydration of sucrose to pure carbon and water vapor in Equation 5.1.2 and demonstrated in Video 5.1.2.

$$\mathrm{C_{12}H_{22}O_{11}(s) + heat \rightarrow 12\,C(s) + 11\,H_2O(g)} \tag{5.1.2}$$



***Video* 5.1.2:** *A science experiment in the kitchen shows what happens to sugar molecules when they are heated. The experiment did not disappoint!*

As with salt, sugar has radically different properties (both physical and chemical) than its constituent elements. That is a central feature of chemical reactions as this chapter will discuss.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Wikipedia

5.1: Sugar and Salt is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

5.2: Compounds Display Constant Composition

When building a house, the starting point is a blueprint of what the house will look like. The plan states how many windows and what kind, how many doors and what style, how many rooms and what type (bedroom, kitchen, other). The blueprint shows how the different pieces will go together to make the house. As long as the blueprint is followed and exactly the same items are used, the house will be identical to its blueprint.

Compounds

A **compound** is a substance that contains two or more elements chemically combined in a fixed proportion. The elements carbon and hydrogen combine to form many different compounds. One of the simplest is called methane, in which there are always four times as many hydrogen particles as carbon particles. Methane is a pure substance because it always has the same composition. However, it is not an element because it can be broken down into simpler substances—carbon and hydrogen.

Recall that the components of a mixture can be separated from one another by physical means. This is not true for a compound. Table salt is a compound consisting of equal parts of the elements sodium and chlorine. Salt cannot be separated into its two elements by filtering, distillation, or any other physical process. Salt and other compounds can only be decomposed into their elements by a chemical process. A **chemical change** is a change that produces matter with a different composition. Many compounds can be decomposed into their elements by heating. When sugar is heated, it decomposes into carbon and water. Water is still a compound, but one which cannot be broken down into hydrogen and oxygen by heating. Instead, the passage of an electrical current through water will produce hydrogen and oxygen gases.

The properties of compounds are generally very different than the properties of the elements from which the compound is formed. Sodium is an extremely reactive soft metal that cannot be exposed to air or water. Chlorine is a deadly gas. The compound sodium chloride is a white solid which is essential for all living things (see below).




Figure 5.2.1: (A) Sodium is so reactive that it must be stored under oil. (B) Chlorine is a poisonous yellow-green gas. (C) Salt crystals, a compound of sodium and chlorine.

Summary

- A compound is a substance that contains two or more elements chemically combined in a fixed proportion.
- A chemical change is a change that produces matter with a different composition.

5.2: Compounds Display Constant Composition is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.2: Compounds Display Constant Composition** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.3: Chemical Formulas- How to Represent Compounds

Learning Objectives

- Determine the number of different atoms in a formula.
- Define chemical formula, molecular formula, and empirical formula.

A **chemical formula** is an expression that shows the elements in a compound and the relative proportions of those elements. Water is composed of hydrogen and oxygen in a 2:1 ratio. The chemical formula for water is H_2O. Sulfuric acid is one of the most widely produced chemicals in the United States and is composed of the elements hydrogen, sulfur, and oxygen. The chemical formula for sulfuric acid is H_2SO_4.

Certain groups of atoms are bonded together to form what is called a polyatomic ion that acts as a single unit. Polyatomic ions are discussed in more detail in Section 5.5. Polyatomic ions are enclosed in parenthesis followed by a subscript if more than one of the same ion exist in a chemical formula. The formula $Ca_3(PO_4)_2$ represents a compound with the following:

3 Ca atoms + 2 PO_4^{3-} ions

To count the total number of atoms for formulas with polyatomic ions enclosed in parenthesis, use the subscript as a multiplier for each atom or number of atoms.

$$Ca_3(PO_4)_2$$

3 Ca + 2 x1 P + 2 x 4 O = 3 Ca atoms + 2 P atoms + 8 O atoms

Molecular Formula

A **molecular formula** is a chemical formula of a molecular compound that shows the kinds and numbers of atoms present in a molecule of the compound. Ammonia is a compound of nitrogen and hydrogen as shown below:




Figure 5.3.1: The molecular formula for ammonia. NH3. There is one atom of nitrogen and 3 atoms of hydrogen in a molecule of ammonia.

Note from the example that there are some standard rules to follow in writing molecular formulas. The arrangements of the elements depend on the particular structure, which is not of concern at this point. The number of atoms of each kind is indicated by a subscript following the atom. If there is only one atom, no number is written. If there is more than one atom of a specific kind, the number is written as a subscript following the atom. We would not write N_3H for ammonia, because that would mean that there are three nitrogen atoms and one hydrogen atom in the molecule, which is incorrect.

Empirical Formula

An **empirical formula** is a formula that shows the elements in a compound in their lowest whole-number ratio. Glucose is an important simple sugar that cells use as their primary source of energy. Its molecular formula is $C_6H_{12}O_6$. Since each of the subscripts is divisible by 6, the empirical formula for glucose is CH_2O. When chemists analyze an unknown compound, often the first step is to determine its empirical formula.

- molecular formula: $C_6H_{12}O_6$
- empirical formula: CH_2O

There are a great many compounds whose molecular and empirical formulas are the same. If the molecular formula cannot be simplified into a smaller whole-number ratio, as in the case of H_2O or P_2O_5, then the empirical formula is also the molecular formula.

Summary

- A chemical formula is an expression that shows the elements in a compound and the relative proportions of those elements.
- If only one atom of a specific type is present, no subscript is used.
- For atoms that have two or more of a specific type of atom present, a subscript is written after the symbol for that atom.
- Polyatomic ions in chemical formulas are enclosed in parentheses followed by a subscript if more than one of the same type of polyatomic ion exist.
- Molecular formulas do not indicate how the atoms are arranged in the molecule.
- The empirical formula tells the lowest whole-number ratio of elements in a compound. The empirical formula does not show the actual number of atoms.

5.3: Chemical Formulas- How to Represent Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.3: Chemical Formulas - How to Represent Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.4: A Molecular View of Elements and Compounds

Learning Objectives

- Classify substances as atomic elements, molecular elements, molecular compounds, or ionic compounds.

Atomic Elements

Most elements exist with **individual atoms** as their basic unit. It is assumed that there is only one atom in a formula if there is no numerical subscript on the right side of an element's symbol.

Molecular Elements

There are many substances that exist as two or more atoms connected together so strongly that they behave as a single particle. These multi-atom combinations are called **molecules**. A molecule is the smallest part of a substance that has the physical and chemical properties of that substance. In some respects, a molecule is similar to an atom. A molecule, however, is composed of more than one atom.

Table 5.4.1: Elements That Exist as Diatomic Molecules

Hydrogen, H	Oxygen	Nitrogen	Fluorine	Chlorine	Bromine	Iodine

Some elements exist naturally as molecules. For example, hydrogen and oxygen exist as two-atom molecules. Other elements also exist naturally as diatomic molecules—a molecule with only two atoms (Table 5.4.1). As with any molecule, these elements are labeled with a **molecular formula**, a formal listing of what and how many atoms are in a molecule. (Sometimes only the word *formula* is used, and its meaning is inferred from the context.) For example, the molecular formula for elemental hydrogen is H_2, with H being the symbol for hydrogen and the subscript 2 implying that there are two atoms of this element in the molecule. Other diatomic elements have similar formulas: O_2, N_2, and so forth. Other elements exist as molecules—for example, sulfur normally exists as an eight-atom molecule, S_8, while phosphorus exists as a four-atom molecule, P_4 (Figure 5.4.1).




Figure 5.4.1: Molecular Art of S_8 and P_4 Molecules. If each green ball represents a sulfur atom, then the diagram on the left represents an S_8 molecule. The molecule on the right shows that one form of elemental phosphorus exists, as a four-atom molecule.

Figure 5.4.1 shows two examples of how molecules will be represented in this text. An atom is represented by a small ball or sphere, which generally indicates where the nucleus is in the molecule. A cylindrical line connecting the balls represents the connection between the atoms that make this collection of atoms a molecule. This connection is called a chemical bond.

Ionic Compounds

The elements in the periodic table are divided into specific groupings; the metals, the non-metals, the semi-metals, and so on. These groupings are largely based on physical properties and on the tendency of the various elements to bond with other elements by forming either an ionic or a covalent bond. As a general rule of thumb, compounds that involve a metal binding with either a non-metal or a semi-metal will display ionic bonding. Thus, the compound formed from sodium and chlorine will be ionic (a metal and a non-metal). The basic unit of ionic compounds is the **formula unit**.

Molecular Compounds

Compounds that are composed of only non-metals or semi-metals with non-metals will display covalent bonding and will be classified as molecular compounds. Nitrogen monoxide (NO) will be a covalently bound molecule (two non-metals) and silicon

dioxide (SiO_2) will also be a covalently bound molecule (a semi-metal and a non-metal). The basic unit of molecular compounds is the **molecule**.

Example 5.4.1

Provide the classification (i.e. atomic element, molecular element, molecular compound, or ionic compound) of each substance.

a. Fe
b. PCl_3
c. LiBr
d. P_4
e. oxygen gas

Solution

a. **Fe** (iron) is an element that is represented with no subscript, so it is an **atomic element**.
b. **PCl_3** is made up of two nonmetals, so it is a **molecular compound**.
c. **LiBr** is made up of lithium, a metal, and bromine, a nonmetal, so it is an **ionic compound**.
d. **P_4** is a substance that is made up of four atoms of the same element, so it is a **molecular element**.
e. The formula for **oxygen gas** is **O_2** so it is a molecular element.

Exercise 5.4.1

Provide the classification (i.e. atomic element, molecular element, molecular compound, or ionic compound) of each substance.

a. I_2
b. He
c. H_2O
d. Al
e. CuCl

Answer a:

molecular element

Answer b:

atomic element

Answer c:

molecular compound

Answer d:

atomic element

Answer e:

ionic compound

5.4: A Molecular View of Elements and Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.4: A Molecular View of Elements and Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.5: Writing Formulas for Ionic Compounds

Learning Objectives

- Write the correct formula for an ionic compound.
- Recognize polyatomic ions in chemical formulas.

Ionic compounds do not exist as molecules. In the solid state, ionic compounds are in crystal lattice containing many ions each of the cation and anion. An ionic formula, like $NaCl$, is an empirical formula. This formula merely indicates that sodium chloride is made of an equal number of sodium and chloride ions. Sodium sulfide, another ionic compound, has the formula Na_2S. This formula indicates that this compound is made up of twice as many sodium ions as sulfide ions. This section will teach you how to find the correct ratio of ions, so that you can write a correct formula.

If you know the name of a binary ionic compound, you can write its **chemical formula**. Start by writing the metal ion with its charge, followed by the nonmetal ion with its charge. Because the overall compound must be electrically neutral, decide how many of each ion is needed in order for the positive and negative charges to cancel each other out.

Example 5.5.1: Aluminum Nitride and Lithium Oxide

Write the formulas for aluminum nitride and lithium oxide.

Solution

Solution to Example 5.5.1

	Write the formula for aluminum nitride	Write the formula for lithium oxide
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Al^{3+} N^{3-}	Li^{+} O^{2-}
2. Use a multiplier to make the total charge of the cations and anions equal to each other.	total charge of cations = total charge of anions 1(3+) = 1(3-) +3 = -3	total charge of cations = total charge of anions 2(1+) = 1(2-) +2 = -2
3. Use the multipliers as subscript for each ion.	Al_1N_1	Li_2O_1
4. Write the final formula. Leave out all charges and all subscripts that are 1.	AlN	Li_2O

An alternative way to writing a correct formula for an ionic compound is to use the **crisscross method**. In this method, the numerical value of each of the ion charges is crossed over to become the subscript of the other ion. Signs of the charges are dropped.

Example 5.5.2: The Crisscross Method for Lead (IV) Oxide

Write the formula for lead (IV) oxide.

Solution

Solution to Example 5.5.2

Crisscross Method	Write the formula for lead (IV) oxide
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Pb^{4+} O^{2-}

Crisscross Method	Write the formula for lead (IV) oxide
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	
3. Reduce to the lowest ratio.	Pb_2O_4
4. Write the final formula. Leave out all subscripts that are 1.	PbO_2

Exercise 5.5.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the calcium ion and the oxygen ion
b. the 2+ copper ion and the sulfur ion
c. the 1+ copper ion and the sulfur ion

Answer a:

CaO

Answer b:

CuS

Answer c:

Cu_2S

> *Be aware that ionic compounds are empirical formulas and so must be written as the lowest ratio of the ions.*

Example 5.5.3: Sulfur Compound

Write the formula for sodium combined with sulfur.

Solution

Solution to Example 5.5.3

Crisscross Method	Write the formula for sodium combined with sulfur
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	$Na^+ \quad S^{2-}$
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	
3. Reduce to the lowest ratio.	This step is not necessary.
4. Write the final formula. Leave out all subscripts that are 1.	Na_2S

Exercise 5.5.3

Write the formula for each ionic compound.

a. sodium bromide
b. lithium chloride
c. magnesium oxide

Answer a:

NaBr

Answer b:

LiCl

Answer c:

MgO

Polyatomic Ions

Some ions consist of groups of atoms bonded together and have an overall electric charge. Because these ions contain more than one atom, they are called polyatomic ions. Polyatomic ions have characteristic formulas, names, and charges that should be memorized. For example, NO_3^- is the nitrate ion; it has one nitrogen atom and three oxygen atoms and an overall 1− charge. Table 5.5.1 lists the most common polyatomic ions.

Table 5.5.1: Some Polyatomic Ions

Name	Formula
ammonium ion	NH_4^+
acetate ion	$C_2H_3O_2^-$ (also written $CH_3CO_2^-$)
carbonate ion	CO_3^{2-}
chromate ion	CrO_4^{2-}
dichromate ion	$Cr_2O_7^{2-}$
hydrogen carbonate ion (bicarbonate ion)	HCO_3^-
cyanide ion	CN^-
hydroxide ion	OH^-
nitrate ion	NO_3^-
nitrite ion	NO_2^-
permanganate ion	MnO_4^-
phosphate ion	PO_4^{3-}
hydrogen phosphate ion	HPO_4^{2-}
dihydrogen phosphate ion	$H_2PO_4^-$
sulfate ion	SO_4^{2-}
hydrogen sulfate ion (bisulfate ion)	HSO_4^-
sulfite ion	SO_3^{2-}

The rule for constructing formulas for ionic compounds containing polyatomic ions is the same as for formulas containing monatomic (single-atom) ions: the positive and negative charges must balance. If more than one of a particular polyatomic ion is needed to balance the charge, the *entire formula* for the polyatomic ion must be enclosed in parentheses, and the numerical subscript is placed *outside* the parentheses. This is to show that the subscript applies to the entire polyatomic ion. An example is $Ba(NO_3)_2$.

Writing Formulas for Ionic Compounds Containing Polyatomic Ions

Writing a formula for ionic compounds containing polyatomic ions also involves the same steps as for a binary ionic compound. Write the symbol and charge of the cation followed by the symbol and charge of the anion.

Example 5.5.4: Calcium Nitrate

Write the formula for calcium nitrate.

Solution

Solution to Example 5.5.4

Crisscross Method	Write the formula for calcium nitrate
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Ca^{2+} NO_3^-
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	 The 2+ charge on Ca becomes the subscript of NO3 and the 1- charge on NO3 becomes the subscript of Ca.
3. Reduce to the lowest ratio.	$Ca_1(NO_3)_2$
4. Write the final formula. Leave out all subscripts that are 1. If there is only 1 of the polyatomic ion, leave off parentheses.	$Ca(NO_3)_2$

Example 5.5.5

Write the chemical formula for an ionic compound composed of the potassium ion and the sulfate ion.

Solution

Solution to Example 5.5.5

Explanation	Answer
Potassium ions have a charge of 1+, while sulfate ions have a charge of 2−. We will need two potassium ions to balance the charge on the sulfate ion, so the proper chemical formula is K_2SO_4.	K_2SO_4

Exercise 5.5.5

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the magnesium ion and the carbonate ion
b. the aluminum ion and the acetate ion

Answer a:

$MgCO_3$

Answer b:

$Al(CH_3COO)_3$

Recognizing Ionic Compounds

There are two ways to recognize ionic compounds.

Method 1

Compounds between metal and nonmetal elements are usually ionic. For example, $CaBr_2$ contains a metallic element (calcium, a group 2 [or 2A] metal) and a nonmetallic element (bromine, a group 17 [or 7A] nonmetal). Therefore, it is most likely an ionic compound (in fact, it *is* ionic). In contrast, the compound NO_2 contains two elements that are both nonmetals (nitrogen, from group 15 [or 5A], and oxygen, from group 16 [or 6A]. It is not an ionic compound; it belongs to the category

of covalent compounds discussed elsewhere. Also note that this combination of nitrogen and oxygen has no electric charge specified, so it is *not* the nitrite ion.

Method 2

Second, if you recognize the formula of a polyatomic ion in a compound, the compound is ionic. For example, if you see the formula $\mathrm{Ba(NO_3)_2}$, you may recognize the "$\mathrm{NO_3}$" part as the nitrate ion, $\mathrm{NO_3^-}$. (Remember that the convention for writing formulas for ionic compounds is not to include the ionic charge.) This is a clue that the other part of the formula, Ba, is actually the $\mathrm{Ba^{2+}}$ ion, with the 2+ charge balancing the overall 2− charge from the two nitrate ions. Thus, this compound is also ionic.

Example 5.5.6

Identify each compound as ionic or not ionic.

a. $\mathrm{Na_2O}$
b. $\mathrm{PCl_3}$
c. $\mathrm{NH_4Cl}$
d. $\mathrm{OF_2}$

Solution

Solution to Example 5.5.6

Explanation	Answer
a. Sodium is a metal, and oxygen is a nonmetal. Therefore, $\mathrm{Na_2O}$ is expected to be ionic via method 1.	$\mathrm{Na_2O}$, ionic
b. Both phosphorus and chlorine are nonmetals. Therefore, $\mathrm{PCl_3}$ is not ionic via method 1	$\mathrm{PCl_3}$, not ionic
c. The $\mathrm{NH_4}$ in the formula represents the ammonium ion, $\mathrm{NH_4^+}$, which indicates that this compound is ionic via method 2	$\mathrm{NH_4Cl}$, ionic
d. Both oxygen and fluorine are nonmetals. Therefore, $\mathrm{OF_2}$ is not ionic via method 1	$\mathrm{OF_2}$, not ionic

Exercise 5.5.6

Identify each compound as ionic or not ionic.

a. $\mathrm{N_2O}$
b. $\mathrm{FeCl_3}$
c. $\mathrm{(NH_4)_3PO_4}$
d. $\mathrm{SOCl_2}$

Answer a:

not ionic

Answer b:

ionic

Answer c:

ionic

Answer d:

not ionic

Summary

Formulas for ionic compounds contain the symbols and number of each atom present in a compound in the lowest whole number ratio.

5.5: Writing Formulas for Ionic Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.5: Writing Formulas for Ionic Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.6: Nomenclature- Naming Compounds

Nomenclature is the process of naming chemical compounds so that they can be easily identified as separate chemicals. The primary function of chemical nomenclature is to ensure that a spoken or written chemical name leaves no ambiguity concerning which chemical compound the name refers to—each chemical name should refer to a single substance. A less important aim is to ensure that each substance has a single name, although a limited number of alternative names is acceptable in some cases. Preferably, the name also conveys some information about the structure or chemistry of a compound. A common name will often suffice to identify a chemical compound in a particular set of circumstances. To be more generally applicable, the name should indicate at least the chemical formula. To be more specific still, the three-dimensional arrangement of the atoms may need to be specified.

Contributions & Attributions

- Wikipedia (CC-BY-SA-3.0)

5.6: Nomenclature- Naming Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.6: Nomenclature- Naming Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.7: Naming Ionic Compounds

Learning Objectives

- To use the rules for naming ionic compounds.

After learning a few more details about the names of individual ions, you will be one step away from knowing how to name ionic compounds. This section begins the formal study of nomenclature, the systematic naming of chemical compounds.

Naming Ions

The name of a monatomic cation is simply the name of the element followed by the word *ion*. Thus, Na^+ is the sodium ion, Al^{3+} is the aluminum ion, Ca^{2+} is the calcium ion, and so forth.

We have seen that some elements lose different numbers of electrons, producing ions of different charges (Figure 3.3). Iron, for example, can form two cations, each of which, when combined with the same anion, makes a different compound with unique physical and chemical properties. Thus, we need a different name for each iron ion to distinguish Fe^{2+} from Fe^{3+}. The same issue arises for other ions with more than one possible charge.

There are two ways to make this distinction. In the simpler, more modern approach, called the **Stock system**, an ion's positive charge is indicated by a roman numeral in parentheses after the element name, followed by the word *ion*. Thus, Fe^{2+} is called the iron(II) ion, while Fe^{3+} is called the iron(III) ion. This system is used only for elements that form more than one common positive ion. We do not call the Na^+ ion the sodium(I) ion because (I) is unnecessary. Sodium forms only a 1+ ion, so there is no ambiguity about the name *sodium ion*.

Table 5.7.1: The Modern and Common System of Cation Names

Element	Stem	Charge	Modern Name	Common Name
iron	ferr-	2+	iron(II) ion	ferrous ion
		3+	iron(III) ion	ferric ion
copper	cupr-	1+	copper(I) ion	cuprous ion
		2+	copper(II) ion	cupric ion
tin	stann-	2+	tin(II) ion	stannous ion
		4+	tin(IV) ion	stannic ion
lead	plumb-	2+	lead(II) ion	plumbous ion
		4+	lead(IV) ion	plumbic ion
chromium	chrom-	2+	chromium(II) ion	chromous ion
		3+	chromium(III) ion	chromic ion
gold	aur-	1+	gold(I) ion	aurous ion
		3+	gold(III) ion	auric ion

The second system, called the **common system**, is not conventional but is still prevalent and used in the health sciences. This system recognizes that many metals have two common cations. The common system uses two suffixes (*-ic* and *-ous*) that are appended to the stem of the element name. The *-ic* suffix represents the greater of the two cation charges, and the *-ous* suffix represents the lower one. In many cases, the stem of the element name comes from the Latin name of the element. Table 5.7.1 lists the elements that use the common system, along with their respective cation names.

Table 5.7.2: Some Monatomic Anions

Ion	Name

Ion	Name
F^-	fluoride ion
Cl^-	chloride ion
Br^-	bromide ion
I^-	iodide ion
O^{2-}	oxide ion
S^{2-}	sulfide ion
P^{3-}	phosphide ion
N^{3-}	nitride ion

The name of a monatomic anion consists of the stem of the element name, the suffix *-ide*, and then the word *ion*. Thus, as we have already seen, Cl^- is "chlor-" + "-ide ion," or the chloride ion. Similarly, O^{2-} is the oxide ion, Se^{2-} is the selenide ion, and so forth. Table 5.7.2 lists the names of some common monatomic ions. The polyatomic ions have their own characteristic names, as discussed earlier.

Example 5.7.1

Name each ion.

a. Ca^{2+}
b. S^{2-}
c. SO_3^{2-}
d. NH_4^+
e. Cu^+

Solution

a. the calcium ion
b. the sulfide ion
c. the sulfite ion
d. the ammonium ion
e. the copper(I) ion or the cuprous ion

Exercise 5.7.1

Name each ion.

a. Fe^{2+}
b. Fe^{3+}
c. SO_4^{2-}
d. Ba^{2+}
e. HCO_3^-

Answer a:
iron(II) ion
Answer b:
iron(III) ion
Answer c:
sulfate ion
Answer d:
barium ion

Answer e:

hydrogen carbonate ion or bicarbonate ion

Example 5.7.2

Write the formula for each ion.

a. the bromide ion
b. the phosphate ion
c. the cupric ion
d. the magnesium ion

Solution

a. Br^-
b. PO_4^{3-}
c. Cu^{2+}
d. Mg^{2+}

Exercise 5.7.2

Write the formula for each ion.

a. the fluoride ion
b. the carbonate ion
c. the stannous ion
d. the potassium ion

Answer a:

F^-

Answer b:

CO_3^{2-}

Answer c:

Sn^{2+}

Answer d:

K^+

Naming Binary Ionic Compounds with a Metal that Forms Only One Type of Cation

A **binary** ionic compound is a compound composed of a **monatomic** metal **cation** and a monatomic nonmetal **anion**. The metal cation is named first, followed by the nonmetal anion as illustrated in Figure 5.7.1 for the compound $BaCl_2$. The word *ion* is dropped from both parts.




Figure 5.7.1: Naming $BaCl_2$
Naming formula: Name of metal cation + base
name of nonmetal anion + suffix -ide. BaCl2 is named as barium chloride.

Subscripts in the formula do not affect the name.

Example 5.7.3: Naming Ionic Compounds

Name each ionic compound.

a. $CaCl_2$
b. AlF_3
c. KCl

Solution

a. Using the names of the ions, this ionic compound is named calcium chloride.
b. The name of this ionic compound is aluminum fluoride.
c. The name of this ionic compound is potassium chloride

Exercise 5.7.3

Name each ionic compound.

a. AgI
b. MgO
c. Ca_3P_2

Answer a:

silver iodide

Answer b:

magnesium oxide

Answer c:

calcium phosphide


Naming Binary Ionic Compounds with a Metal That Forms More Than One Type of Cation

If you are given a formula for an ionic compound whose cation can have more than one possible charge, you must first determine the charge on the cation before identifying its correct name. For example, consider $FeCl_2$ and $FeCl_3$. In the first compound, the iron ion has a 2+ charge because there are two Cl^- ions in the formula (1− charge on each chloride ion). In the second compound, the

iron ion has a 3+ charge, as indicated by the three Cl^- ions in the formula. These are two different compounds that need two different names. By the Stock system, the names are iron(II) chloride and iron(III) chloride (Figure 5.7.2).

Table 5.7.3: Naming the $FeCl_2$ and $FeCl_3$ Compounds in the Modern/Stock System.


Name of cation (metal) + (Roman Numeral in parenthesis) + Base name of anion (nonmetal) and *-ide*



Remove the word "ion"

iron (II) + chloride

iron (II) chloride



Remove the word "ion"

iron (III) + chloride


iron (III) chloride

If we were to use the stems and suffixes of the common system, the names would be ferrous chloride and ferric chloride, respectively (Figure 5.7.3) .

Table 5.7.4: Naming the $FeCl_2$ and $FeCl_3$ Compounds in the Old/Common System.

"Old" base name of cation (metal) and *-ic or -ous* + Base name of anion (nonmetal) and *-ide*

-ous (for ions with lower charge)




Remove the word "ion"

ferrous + chloride

ferrous chloride

-ic (for ions with higher charge)



Remove the word "ion"

ferric + chloride

ferric chloride

Example 5.7.4:

Name each ionic compound.

a. Co_2O_3
b. FeCl2

Solution

Solutions to Example 5.7.4

	Explanation	Answer
a	We know that cobalt can have more than one possible charge; we just need to determine what it is. • Oxide always has a 2− charge, so with three oxide ions, we have a total negative charge of 6−. • This means that the two cobalt ions have to contribute 6+, which for two cobalt ions means that each one is 3+. • Therefore, the proper name for this ionic compound is cobalt(III) oxide.	cobalt(III) oxide
b	Iron can also have more than one possible charge. • Chloride always has a 1− charge, so with two chloride ions, we have a total negative charge of 2−. • This means that the one iron ion must have a 2+ charge. • Therefore, the proper name for this ionic compound is iron(II) chloride.	iron(II) chloride

Exercise 5.7.4

Name each ionic compound.

a. $AuCl_3$
b. PbO_2
c. CuO

Answer a:
gold(III) chloride
Answer b:
lead(IV) oxide
Answer c:
copper(II) oxide

Naming Ionic Compounds with Polyatomic Ions

The process of naming ionic compounds with polyatomic ions is the same as naming binary ionic compounds. The cation is named first, followed by the anion. One example is the ammonium sulfate compound in Figure 5.7.6.




Figure 5.7.2: Naming Ionic Compounds with Polyatomic Ions

Example 5.7.5: Naming Ionic Compounds

Write the proper name for each ionic compound.

a. $(NH_4)_2S$
b. $AlPO_4$,
c. $Fe_3(PO_4)_2$

Solution

Solutions to Example 5.7.5

Explanation	Answer
a. The ammonium ion has a 1+ charge and the sulfide ion has a 2− charge. Two ammonium ions need to balance the charge on a single sulfide ion. The compound’s name is ammonium sulfide.	ammonium sulfide
b. The ions have the same magnitude of charge, one of each (ion) is needed to balance the charges. The name of the compound is aluminum phosphate.	aluminum phosphate
c. Neither charge is an exact multiple of the other, so we have to go to the least common multiple of 6. To get 6+, three iron(II) ions are needed, and to get 6−, two phosphate ions are needed . The compound’s name is iron(II) phosphate.	iron(II) phosphate

Exercise 5.7.5A

Write the proper name for each ionic compound.

a. $(NH_4)_3PO_4$
b. $Co(NO_2)_3$

Answer a:

ammonium phosphate

Answer b:

cobalt(III) nitrite

Figure 5.7.1 is a synopsis of how to name simple ionic compounds.




Figure 5.7.3: A Guide to Naming Simple Ionic Compounds.

Exercise 5.7.5B

Name each ionic compound.

a. $ZnBr_2$
b. Al_2O_3
c. $(NH_4)_3PO_4$
d. AuF_3
e. AgF

Answer a:

zinc bromide

Answer b:

aluminum oxide

Answer c:

ammonium phosphate

Answer d:

gold(III) fluoride or auric fluoride

Answer e:

silver fluoride

Summary

- Ionic compounds are named by stating the cation first, followed by the anion.
- Positive and negative charges must balance.
- Some anions have multiple forms and are named accordingly with the use of roman numerals in parentheses.
- Ternary compounds are composed of three or more elements.

5.7: Naming Ionic Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.7: Naming Ionic Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.8: Naming Molecular Compounds

Learning Objectives

- Determine the name of a simple molecular compound from its chemical formula.

Molecular Compounds

Molecular compounds are inorganic compounds that take the form of discrete **molecules**. Examples include such familiar substances as water (H_2O) and carbon dioxide (CO_2). These compounds are very different from ionic compounds like sodium chloride ($NaCl$). Ionic compounds are formed when metal atoms lose one or more of their electrons to nonmetal atoms. The resulting cations and anions are electrostatically attracted to each other.

So what holds the atoms of a molecule together? Rather than forming ions, the atoms of a molecule share their **electrons** in such a way that a **bond** forms between a pair of atoms. In a carbon dioxide molecule, there are two of these bonds, each occurring between the carbon atom and one of the two oxygen atoms.




Figure 5.8.1: Carbon dioxide molecules consist of a central carbon atom bonded to 2 oxygen atoms.

Larger molecules can have many, many bonds that serve to keep the molecule together. In a large sample of a given molecular compound, all of the individual molecules are identical.

Naming Binary Molecular Compounds

Recall that a molecular formula shows the number of atoms of each element that a molecule contains. A molecule of water contains two hydrogen atoms and one oxygen atom, so its formula is H_2O. A molecule of octane, which is a component of gasoline, contains 8 atoms of carbon and 18 atoms of hydrogen. The molecular formula of octane is C_8H_{18}.



Figure 5.8.2: Nitrogen dioxide (NO_2) is a reddish-brown toxic gas that is a prominent air pollutant produced by internal combustion engines.


Naming *binary* (two-element) molecular compounds is similar to naming simple ionic compounds. The first element in the formula is simply listed using the name of the element. The second element is named by taking the stem of the element name and adding the suffix *-ide*. A system of numerical prefixes is used to specify the number of atoms in a molecule. Table 5.8.1 lists these numerical prefixes.

Table 5.8.1: Numerical Prefixes for Naming Binary Covalent Compounds

Number of Atoms in Compound	Prefix on the Name of the Element
1	mono-*
2	di-
3	tri-
4	tetra-
5	penta-

Number of Atoms in Compound	Prefix on the Name of the Element
6	hexa-
7	hepta-
8	octa-
9	nona-
10	deca-

*This prefix is not used for the first element's name.



Note

- Generally, the less electronegative element is written first in the formula, though there are a few exceptions. Carbon is always first in a formula and hydrogen is after nitrogen in a formula such as NH_3. The order of common nonmetals in binary compound formulas is C, P, N, H, S, I, Br, Cl, O, F.
- The *a* or *o* at the end of a prefix is usually dropped from the name when the name of the element begins with a vowel. As an example, four oxygen atoms, is tetr**o**xide instead of tetr**ao**xide.
- The prefix is "mono" is not added to the first element's name if there is only one atom of the first element in a molecule.

Some examples of molecular compounds are listed in Table 5.8.2.

Table 5.8.2

Formula	Name
NO	nitrogen monoxide
N_2O	dinitrogen monoxide
S_2Cl_2	disulfur dichloride
Cl_2O_7	dichlorine heptoxide

Notice that the *mono-* prefix is not used with the nitrogen in the first compound, but is used with the oxygen in both of the first two examples. The S_2Cl_2 emphasizes that the formulas for molecular compounds are not reduced to their lowest ratios. The *o* of the *mono-* and the *a* of *hepta-* are dropped from the name when paired with oxide.

Exercise 5.8.1

Write the name for each compound.

a. CF_4
b. $SeCl_2$

c. SO_3

Answer a:

carbon tetrafluoride

Answer b:

selenium dichloride

Answer c:

sulfur trioxide

Simple molecular compounds with common names

For some simple covalent compounds, we use common names rather than systematic names. We have already encountered these compounds, but we list them here explicitly:

- H_2O: water
- NH_3: ammonia
- CH_4: methane
- H_2O_2: hydrogen peroxide

Methane is the simplest organic compound. Organic compounds are compounds with carbon atoms and are named by a separate nomenclature system.

Some Compounds Have Both Covalent and Ionic Bonds

If you recall the introduction of polyatomic ions, you will remember that the bonds that hold the polyatomic ions together are covalent bonds. Once the polyatomic ion is constructed with covalent bonds, it reacts with other substances as an ion. The bond between a polyatomic ion and another ion will be ionic. An example of this type of situation is in the compound sodium nitrate. Sodium nitrate is composed of a sodium ion and a nitrate ion. The nitrate ion is held together by covalent bonds and the nitrate ion is attached to the sodium ion by an ionic bond.

Summary

- A molecular compound is usually composed of two or more nonmetal elements.
- Molecular compounds are named with the first element first and then the second element by using the stem of the element name plus the suffix -ide. Numerical prefixes are used to specify the number of atoms in a molecule.

5.8: Naming Molecular Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.8: Naming Molecular Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.9: Naming Acids

A spot test for gold has been in use for decades. The sample is first treated with nitric acid. Other metals may react or dissolve in this acid, but gold will not. Then the sample is added to a mixture of nitric acid and hydrochloric acid. Gold will only dissolve in this mixture. The term "acid test" arose from the California gold rush in the late 1840's when this combination was used to test for the presence of real gold. It has since come to mean, "tested and approved" in a number of fields.

Acids

An **acid** can be defined in several ways. The most straightforward definition is that an acid is a molecular compound that contains one or more hydrogen atoms and produces hydrogen ions $\left(H^+\right)$ when dissolved in water.



Figure 5.9.1: (A) Vinegar comes in a variety of types, but all contain acetic acid. (B) Citrus fruits like grapefruit contain citric and ascorbic acids.

This is a different type of compound than the others we have seen so far. Acids are molecular, which means that in their pure state they are individual molecules and do not adopt the extended three-dimensional structures of ionic compounds like $NaCl$. However, when these molecules are dissolved in water, the chemical bond between the hydrogen atom and the rest of the molecule breaks, leaving a positively-charged hydrogen ion and an anion. This can be symbolized in a chemical equation:

$$HCl \rightarrow H^+ + Cl^-$$


Since acids produce H^+ cations upon dissolving in water, the H of an acid is written first in the formula of an inorganic acid. The remainder of the acid (other than the H) is the anion after the acid dissolves. Organic acids are also an important class of compounds, but will not be discussed here.

Naming Acids

Since all acids contain hydrogen, the name of an acid is based on the anion that goes with it. These anions can either be monatomic or polyatomic.

Naming Binary acids (in aqueous form)

A **binary acid** is an acid that consists of hydrogen and one other element. The most common binary acids contain a halogen. The acid name begins with the prefix *hydro-*. followed by the base name of the anion, followed by the suffix *-ic*.




Formula for naming acids: Hydro- and Base name of nonmetal and -ic + acid. Example: HCl is hydrochloric acid.

Naming Oxyacids

An **oxyacid** is an acid that consists of hydrogen, oxygen, and a third element. The third element is usually a nonmetal.

a. Oxyanions with ***-ite*** ending.


The name of the acid is the root of the anion followed by the suffix ***-ous***. There is no prefix.



Formula for naming oxyanions with -ite ending: Base name of oxyanion and -ous + acid. Example: H2SO3 is sulfurous acid.

b. Oxyanions with ***-ate*** ending.

The name of the acid is the root of the anion followed by the suffix ***-ic***. There is no prefix.




Formula for naming oxyanions with -ate ending: Base name of oxyanion and -ic + acid. Example: H3PO4 is phosphoric acid.

Note

The base name for sulfur containing oxyacid is ***sulfur-*** instead of just ***sulf-***. The same is true for a phosphorus containing oxyacid. The base name is ***phosphor-*** instead of simply ***phosph-***.

Writing Formulas for Acids

Like other compounds that we have studied, acids are electrically neutral. Therefore, the charge of the anion part of the formula must be exactly balanced out by the H^+ ions. Another way to think about writing the correct formula is to utilize the crisscross method, shown below for sulfuric acid.



H2SO4: H has +1 charge and SO4 has -2 charge, so there must be 2 H+ and 1 SO4 for the charges to balance out.

Formula: H_2SO_4

Figure 5.9.2: *Crisscross approach to writing formula for sulfuric acid.*


Summary

- Acids are molecular compounds that release hydrogen ions.
- A binary acid consists of hydrogen and one other element.
- Oxyacids contain hydrogen, oxygen, and one other element.
- The name of the acid is based on the anion attached to the hydrogen.

5.9: Naming Acids is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.9: Naming Acids** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.10: Nomenclature Summary



Contributions & Attributions

5.10: Nomenclature Summary is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.10: Nomenclature Summary** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.11: Formula Mass- The Mass of a Molecule or Formula Unit

Learning Objectives

- To determine the formula mass of an ionic or molecular compound.

A necessary skill for future chapters is the ability to determine the mass of the formula of an ionic compound. This quantity is called the formula mass. The formula mass is obtained by adding the masses of each individual atom in the formula of the compound. Because a proper formula is electrically neutral (with no net electrons gained or lost), the ions can be considered atoms for the purpose of calculating the formula mass.

Let us start by calculating the formula mass of sodium chloride (NaCl). This formula mass is the sum of the atomic masses of one sodium atom and one chlorine atom, which we find from the periodic table; here, we use the masses to two decimal places:

Na: 22.99 amu

Cl: +35.34 amu

Total: 58.44 amu

To two decimal places, the formula mass of NaCl is 58.44 amu.

When an ionic compound has more than one anion or cation, you must remember to use the proper multiple of the atomic mass for the element in question. For the formula mass of calcium fluoride (CaF_2), we must multiply the mass of the fluorine atom by 2 to account for the two fluorine atoms in the chemical formula:

Ca: 1 x 40.08 = 40.08 amu

F: 2 x 19.00 = +38.00 amu

Total = 78.08 amu

The formula mass of CaF_2 is 78.08 amu.

For ionic compounds with polyatomic ions, the sum must include the number and mass of each atom in the formula for the polyatomic ion. For example, potassium nitrate (KNO_3) has one potassium atom, one nitrogen atom, and three oxygen atoms:

K: 1 x 39.10 = 39.10 amu

N: 1 x 14.00 = +14.00 amu

O: 3 x 16.00 = +48.00 amu

Total = 101.10 amu

The formula mass of KNO_3 is 101.10 amu.

> Potassium nitrate is a key ingredient in gunpowder and has been used clinically as a diuretic.

When a formula contains more than one polyatomic unit in the chemical formula, as in $Ca(NO_3)_2$, do not forget to multiply the atomic mass of every atom inside of the parentheses by the subscript outside of the parentheses. This is necessary because the subscript refers to the *entire polyatomic ion*. Thus, for $Ca(NO_3)_2$, the subscript 2 implies two complete nitrate ions, so we must sum the masses of two (1 × 2) nitrogen atoms and six (3 × 2) oxygen atoms, along with the mass of a single calcium atom:

Ca: 1 x 40.08 = 40.08 amu

N: 2 x 14.00 = +28.00 amu

O: 6 x 16.00 = +96.00 amu

Total = 164.08 amu

The key to calculating the formula mass of an ionic compound is to correctly count each atom in the formula and multiply the atomic masses of its atoms accordingly.

Example 5.11.1

Use the atomic masses (rounded to two decimal places) to determine the formula mass for each ionic compound.

a. $FeCl_3$
b. $(NH_4)_3PO_4$

Solution

a.

Fe: 1 x 55.85 = 55.85 amu

Cl: 1 x 35.45 = +106.35 amu

Total = 162.20 amu

The formula mass of $FeCl_3$ is 162.2 amu.

b. When we distribute the subscript 3 through the parentheses containing the formula for the ammonium ion, we see that we have 3 nitrogen atoms and 12 hydrogen atoms. Thus, we set up the sum as follows:

N: 3 x 14.00 = 42.00 amu

H: 12 x 1.00 = +12.00 amu

P: 1 x 30.97 = +30.97 amu

O: 4 x 16.00 = +64.00 amu

Total = 148.97 amu

The formula mass for $(NH_4)_3PO_4$ is 149.0 amu.

Exercise 5.11.1

Use the atomic masses (rounded to two decimal places) to determine the formula mass for each ionic compound.

a. TiO_2
b. AgBr
c. $Au(NO_3)_3$
d. $Fe_3(PO_4)_2$

Answer

a. 79.87 amu
b. 187.77 amu
c. 383.0 amu

To Your Health: Hydrates

Some ionic compounds have water (H_2O) incorporated within their formula unit. These compounds, called *hydrates*, have a characteristic number of water units associated with each formula unit of the compound. Hydrates are solids, not liquids or solutions, despite the water they contain.

To write the chemical formula of a hydrate, write the number of water units per formula unit of compound after its chemical formula. The two chemical formulas are separated by a vertically centered dot. The hydrate of copper(II) sulfate has five water units associated with each formula unit, so it is written as $CuSO_4 \cdot 5\,H_2O$. The name of this compound is copper(II) sulfate pentahydrate, with the *penta-* prefix indicating the presence of five water units per formula unit of copper(II) sulfate.



Cast (CC BY-SA 4.0; JanSLWC via Wikipedia).

Hydrates have various uses in the health industry. Calcium sulfate hemihydrate ($CaSO_4 \cdot \frac{1}{2}H_2O$), known as **plaster of Paris**, is used to make casts for broken bones. Epsom salt ($MgSO_4 \cdot 7\,H_2O$) is used as a bathing salt and a laxative. Aluminum chloride hexahydrate is an active ingredient in antiperspirants. Table $5.11.1$ lists some useful hydrates.

Table 5.11.1: Names and Formulas of Some Widely Used Hydrates

Formula	Name	Uses
$AlCl_3$•$6H_2O$	aluminum chloride hexahydrate	antiperspirant
$CaSO_4$•½H_2O	calcium sulfate hemihydrate (plaster of Paris)	casts (for broken bones and castings)
$CaSO_4$•$2H_2O$	calcium sulfate dihydrate (gypsum)	drywall component
$CoCl_2$•$6H_2O$	cobalt(II) chloride hexahydrate	drying agent, humidity indicator
$CuSO_4$•$5H_2O$	copper(II) sulfate pentahydrate	fungicide, algicide, herbicide
$MgSO_4$•$7H_2O$	magnesium sulfate heptahydrate (Epsom salts)	laxative, bathing salt
Na_2CO_3•$10H_2O$	sodium carbonate decahydrate (washing soda)	laundry additive/cleaner

Key Takeaway

- Formula masses of ionic compounds can be determined from the masses of the atoms in their formulas.

5.11: Formula Mass- The Mass of a Molecule or Formula Unit is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.11: Formula Mass - The Mass of a Molecule or Formula Unit** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

5.12: Moles, Molecular Formulas, and Calculating Empirical Formulas for Compounds

Learning Objectives

- Define molecular formula.
- Define empirical formula.
- Identify the difference between molecular and empirical formulas.
- Determine empirical formula from percent composition of a compound.

When studying chemical formulas, they may be written in several different ways. The most common way is via a **molecular formula**. Molecular formulas tell us the number of atoms of each type of element in a molecule or compound. This formula does not necessarily tell us the order in which the atoms are connected, but will provide the total number of each type of atom. Take the molecular formula for sugar $C_6H_{12}O_6$. This is the formula for both glucose and galactose. Both sugars have the same molecular formula, but different structures. In section 5.11 we covered calculating formula mass values. The mass of any compound represents the mass of all atoms of the substance in 1 mole of a substance. We will cover more about moles later, but for now it is important to understand that moles are the way we quantify a grouping of atoms for a specific substance. A great example of this is taking a carton of eggs. We measure eggs by the dozen. It is commonly accepted that one dozen eggs is the same as 12 eggs. When calculating the mass of a substance, it is accepted that the mass of each element shown on the periodic table is the mass for one mole (or one grouping of atoms) for that substance. The units used here for mass will be grams. Hence, if you look at any element on the periodic table the mass shown is the number of grams of atoms in one mole of that substance.

Another way in which a formula may be written is called an **empirical formula**. Empirical formulas do not tell you the number of teach type of atom that is in a substance, but rather only the types of atoms involved in a substance. Often times empirical formulas are found, and then through taking measurements the molecular formula is determined. We will cover the concept of empirical formulas in this unit.

Determining Empirical Formulas

An empirical formula tells us the relative ratios of different atoms in a compound. The ratios hold true on the *molar* level as well. Thus, H_2O is composed of two atoms of hydrogen and 1 atom of oxygen. Likewise, **1.0 mole of H_2O** is composed **of 2.0 moles of hydrogen** and **1.0 mole of oxygen**. We can also work backwards from molar ratios since *if we know the molar amounts of each element in a compound we can determine the empirical formula*.

> *In a procedure called elemental analysis, an unknown compound can be analyzed in the laboratory in order to determine the percentages of each element contained within it. These percentages can be transformed into the mole ratio of the elements, which leads to the empirical formula.*

Note

Steps to determine empirical formula.

1. Assume a 100 g sample of the compound so that the given percentages can be directly converted into grams.
2. Use each element's molar mass to convert the grams of each element to moles.
3. In order to find a whole-number ratio, divide the moles of each element by whichever of the moles from step 2 is the smallest.
4. If all the moles at this point are whole numbers (or very close), the empirical formula can be written with the moles as the subscript of each element.
5. In some cases, one or more of the moles calculated in step 3 will not be whole numbers. Multiply each of the moles by the smallest whole number that will convert each into a whole number. Write the empirical formula.

Example 5.12.1

A compound of iron and oxygen is analyzed and found to contain 69.94% iron and 30.06% oxygen. Find the empirical formula of the compound.

SOLUTION

Steps for Problem Solving	**Find the empirical formula of a compound of 69.94% iron and 30.06% oxygen.**
Identify the "given"information and what the problem is asking you to "find."	Given: $\%$ of $\mathrm{Fe} = 69.94\%$ $\%$ of $\mathrm{O} = 30.06\%$ Find: Empirical formula $= \mathrm{Fe}_?\mathrm{O}_?$
Calculate	
a. Assume a 100 g sample, convert the same % values to grams.	$69.94\ \mathrm{g\ Fe}$ $30.06\ \mathrm{g\ O}$
b. Convert to moles.	$69.94\ \mathrm{g\ Fe} \times \dfrac{1\ \mathrm{mol\ Fe}}{55.85\ \mathrm{g\ Fe}} = 1.252\ \mathrm{mol\ Fe}$ $30.06\ \mathrm{g\ O} \times \dfrac{1\ \mathrm{mol\ O}}{16.00\ \mathrm{g\ O}} = 1.879\ \mathrm{mol\ O}$
c. Divide both moles by the smallest of the results.	$\mathrm{Fe}: \dfrac{1.252\ \mathrm{mol}}{1.252}$ $\mathrm{O}: \dfrac{1.879\ \mathrm{mol}}{1.252}$ The "non- whole number" empirical formula of the compound is $\mathrm{Fe}_1\mathrm{O}_{1.5}$
Multiply each of the moles by the smallest whole number that will convert each into a whole number.	Fe:O = 2 (1:1.5) = 2:3 Since the moles of O is still not a whole number, both moles can be multiplied by 2, while rounding to a whole number.
Write the empirical formula.	The empirical formula of the compound is $\mathrm{Fe}_2\mathrm{O}_3$.
Think about your result.	The subscripts are whole numbers and represent the mole ratio of the elements in the compound. The compound is the ionic compound iron (III) oxide.

Exercise 5.12.1

Mercury forms a compound with chlorine that is 73.9% mercury and 26.1% chlorine by mass. What is the empirical formula?

Answer

HgCl_2

Summary

- A process is described for the calculation of the empirical formula for a compound based on the percent composition of that compound.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

-
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

5.12: Moles, Molecular Formulas, and Calculating Empirical Formulas for Compounds is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

6: Chemical Bonding

Template:HideTOC

How do atoms make compounds? Typically they join together in such a way that they lose their identities as elements and adopt a new identity as a compound. These joins are called *chemical bonds*. But how do atoms join together? Ultimately, it all comes down to electrons. Before we discuss how electrons interact, we need to introduce a tool to simply illustrate electrons in an atom.

6.1: Representing Valence Electrons with Dots
6.2: Lewis Structures of Ionic Compounds- Electrons Transferred
6.3: Covalent Lewis Structures- Electrons Shared
6.4: Writing Lewis Structures for Covalent Compounds
6.5: Resonance - Equivalent Lewis Structures for the Same Molecule
6.6: Predicting the Shapes of Molecules
6.7: Electronegativity and Polarity - Why Oil and Water Don’t Mix

Template:HideTOC

6: Chemical Bonding is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

6.1: Representing Valence Electrons with Dots

Learning Objective

- Draw a Lewis electron dot diagram for an atom or a monatomic ion.

In almost all cases, chemical bonds are formed by interactions of valence electrons in atoms. To facilitate our understanding of how valence electrons interact, a simple way of representing those valence electrons would be useful.

A **Lewis electron dot diagram** (or electron dot diagram, or a Lewis diagram, or a Lewis structure) is a representation of the valence electrons of an atom that uses dots around the symbol of the element. The number of dots equals the number of valence electrons in the atom. These dots are arranged to the right and left and above and below the symbol, with no more than two dots on a side. (The order in which the positions are used does not matter.) For example, the Lewis electron dot diagram for hydrogen is simply

$$\mathbf{H}\cdot$$

Because the side is not important, the Lewis electron dot diagram could also be drawn as follows:

$$\dot{\mathbf{H}} \quad or \quad \cdot\mathbf{H} \quad or \quad \underset{\cdot}{\mathbf{H}}$$

The electron dot diagram for helium, with two valence electrons, is as follows:

$$\mathbf{He:}$$

By putting the two electrons together on the same side, we emphasize the fact that these two electrons are both in the 1*s* subshell; this is the common convention we will adopt, although there will be exceptions later. The next atom, lithium, has an electron configuration of $1s^2 2s^1$, so it has only one electron in its valence shell. Its electron dot diagram resembles that of hydrogen, except the symbol for lithium is used:

$$\mathbf{Li}\cdot$$

Beryllium has two valence electrons in its 2*s* shell, so its electron dot diagram is like that of helium:

$$\mathbf{Be:}$$

The next atom is boron. Its valence electron shell is $2s^2 2p^1$, so it has three valence electrons. The third electron will go on another side of the symbol:

$$\dot{\mathbf{B}}\mathbf{:}$$

Again, it does not matter on which sides of the symbol the electron dots are positioned.

For carbon, there are four valence electrons, two in the 2*s* subshell and two in the 2*p* subshell. As usual, we will draw two dots together on one side, to represent the 2*s* electrons. However, conventionally, we draw the dots for the two *p* electrons on different sides. As such, the electron dot diagram for carbon is as follows:

$$\cdot\dot{\mathbf{C}}\mathbf{:}$$

With N, which has three *p* electrons, we put a single dot on each of the three remaining sides:

$$\cdot\underset{\cdot}{\dot{\mathbf{N}}}\mathbf{:}$$

For oxygen, which has four *p* electrons, we now have to start doubling up on the dots on one other side of the symbol. When doubling up electrons, make sure that each side has no more than two electrons.

$$\cdot\underset{\cdot}{\ddot{\mathbf{O}}}\mathbf{:}$$

Fluorine and neon have seven and eight dots, respectively:

$$\mathbf{:}\underset{\cdot}{\ddot{\mathbf{F}}}\mathbf{:}$$



With the next element, sodium, the process starts over with a single electron because sodium has a single electron in its highest-numbered shell, the $n = 3$ shell. By going through the periodic table, we see that the Lewis electron dot diagrams of atoms will never have more than eight dots around the atomic symbol.

Example 6.1.1: Lewis Dot Diagrams

What is the Lewis electron dot diagram for each element?

a. aluminum
b. selenium

Solution

a. The valence electron configuration for aluminum is $3s^2 3p^1$. So it would have three dots around the symbol for aluminum, two of them paired to represent the 3*s* electrons:



2. The valence electron configuration for selenium is $4s^2 4p^4$. In the highest-numbered shell, the $n = 4$ shell, there are six electrons. Its electron dot diagram is as follows:



Exercise 6.1.1

What is the Lewis electron dot diagram for each element?

a. phosphorus
b. argon

Answer a



Answer b



Summary

- Lewis electron dot diagrams use dots to represent valence electrons around an atomic symbol.
- Lewis electron dot diagrams for ions have less (for cations) or more (for anions) dots than the corresponding atom.

6.1: Representing Valence Electrons with Dots is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.2: Representing Valence Electrons with Dots** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

6.2: Lewis Structures of Ionic Compounds- Electrons Transferred

Learning Objectives

- State the octet rule.
- Define *ionic bond*.
- Draw Lewis structures for ionic compounds.

In Section 4.7, we demonstrated that ions are formed by losing electrons to make cations, or by gaining electrons to form anions. The astute reader may have noticed something: many of the ions that form have eight electrons in their valence shell. Either atoms gain enough electrons to have eight electrons in the valence shell and become the appropriately charged anion, or they lose the electrons in their original valence shell; the *lower* shell, now the valence shell, has eight electrons in it, so the atom becomes positively charged. For whatever reason, having eight electrons in a valence shell is a particularly energetically stable arrangement of electrons. The **octet rule** explains the favorable trend of atoms having eight electrons in their valence shell. When atoms form compounds, the octet rule is not always satisfied for all atoms at all times, but it is a very good rule of thumb for understanding the kinds of bonding arrangements that atoms can make.

It is not impossible to violate the octet rule. Consider sodium: in its elemental form, it has one valence electron and is stable. It is rather reactive, however, and does not require a lot of energy to remove that electron to make the Na^+ ion. We *could* remove another electron by adding even more energy to the ion, to make the Na^{2+} ion. However, that requires much more energy than is normally available in chemical reactions, so sodium stops at a 1+ charge after losing a single electron. It turns out that the Na^+ ion has a complete octet in its new valence shell, the n = 2 shell, which satisfies the octet rule. The octet rule is a result of trends in energies and is useful in explaining why atoms form the ions that they do.

Now consider an Na atom in the presence of a Cl atom. The two atoms have these Lewis electron dot diagrams and electron configurations:

$$\mathbf{Na}\cdot \qquad \cdot\ddot{\underset{\cdot\cdot}{\mathbf{Cl}}}\mathbf{:}$$

$$[Ne]\,3s^1 \qquad [Ne]\,3s^2 3p^5$$

For the Na atom to obtain an octet, it must lose an electron; for the Cl atom to gain an octet, it must gain an electron. An electron transfers from the Na atom to the Cl atom:

$$\mathbf{Na}\cdot \curvearrowright \cdot\ddot{\underset{\cdot\cdot}{\mathbf{Cl}}}\mathbf{:}$$

resulting in two ions—the Na^+ ion and the Cl^- ion:

$$\mathbf{Na}^+ \qquad \mathbf{:}\ddot{\underset{\cdot\cdot}{\mathbf{Cl}}}\mathbf{:}^-$$

$$[Ne] \qquad [Ne]\,3s^2 3p^6$$

Both species now have complete octets, and the electron shells are energetically stable. From basic physics, we know that opposite charges attract. This is what happens to the Na^+ and Cl^- ions:

$$\mathbf{Na}^+ + \mathbf{:}\ddot{\underset{\cdot\cdot}{\mathbf{Cl}}}\mathbf{:}^- \rightarrow Na^+Cl^- \quad or \quad NaCl$$

where we have written the final formula (the formula for sodium chloride) as per the convention for ionic compounds, without listing the charges explicitly. The attraction between oppositely charged ions is called an **ionic bond,** and it is one of the main types of chemical bonds in chemistry. Ionic bonds are caused by electrons *transferring* from one atom to another.

In electron transfer, the number of electrons lost must equal the number of electrons gained. We saw this in the formation of NaCl. A similar process occurs between Mg atoms and O atoms, except in this case two electrons are transferred:



The two ions each have octets as their valence shell, and the two oppositely charged particles attract, making an ionic bond:

$$\mathbf{Mg}^{2+} + \left[:\ddot{\mathbf{O}}:\right]^{2-} \quad Mg^{2+}O^{2-} \text{ or } MgO$$

Remember, in the final formula for the ionic compound, we do not write the charges on the ions.

What about when an Na atom interacts with an O atom? The O atom needs two electrons to complete its valence octet, but the Na atom supplies only one electron:

$$\mathbf{Na}\cdot \curvearrowright \cdot\ddot{\mathbf{O}}:$$

The O atom still does not have an octet of electrons. What we need is a second Na atom to donate a second electron to the O atom:




These three ions attract each other to give an overall neutral-charged ionic compound, which we write as Na_2O. The need for the number of electrons lost being equal to the number of electrons gained explains why ionic compounds have the ratio of cations to anions that they do. This is required by the law of conservation of matter as well.

Example 6.2.1: Synthesis of Calcium Chloride from Elements

With arrows, illustrate the transfer of electrons to form calcium chloride from Ca atoms and Cl atoms.

Solution

A Ca atom has two valence electrons, while a Cl atom has seven electrons. A Cl atom needs only one more to complete its octet, while Ca atoms have two electrons to lose. Thus we need two Cl atoms to accept the two electrons from one Ca atom. The transfer process looks as follows:




The oppositely charged ions attract each other to make $CaCl_2$.

Exercise 6.2.1

With arrows, illustrate the transfer of electrons to form potassium sulfide from K atoms and S atoms.

Answer



Summary

- The tendency to form species that have eight electrons in the valence shell is called the octet rule.
- The attraction of oppositely charged ions caused by electron transfer is called an ionic bond.
- The strength of ionic bonding depends on the magnitude of the charges and the sizes of the ions.

6.2: Lewis Structures of Ionic Compounds- Electrons Transferred is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.3: Lewis Structures of Ionic Compounds- Electrons Transferred** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

6.3: Covalent Lewis Structures- Electrons Shared

Learning Objectives

- Define *covalent bond*.
- Illustrate covalent bond formation with Lewis electron dot diagrams.

Ionic bonding typically occurs when it is easy for one atom to lose one or more electrons and another atom to gain one or more electrons. However, some atoms won't give up or gain electrons easily. Yet they still participate in compound formation. How? There is another mechanism for obtaining a complete valence shell: *sharing* electrons. When electrons are shared between two atoms, they make a bond called a **covalent bond**.


Let us illustrate a covalent bond by using H atoms, with the understanding that H atoms need only two electrons to fill the 1*s* subshell. Each H atom starts with a single electron in its valence shell:

$$\mathbf{H}\cdot \quad \cdot\mathbf{H}$$

The two H atoms can share their electrons:

$$\mathbf{H:H}$$


We can use circles to show that each H atom has two electrons around the nucleus, completely filling each atom's valence shell:



Because each H atom has a filled valence shell, this bond is stable, and we have made a diatomic hydrogen molecule. (This explains why hydrogen is one of the diatomic elements.) For simplicity's sake, it is not unusual to represent the covalent bond with a dash, instead of with two dots:

H–H


Because two atoms are sharing one pair of electrons, this covalent bond is called a **single bond**. As another example, consider fluorine. F atoms have seven electrons in their valence shell:



These two atoms can do the same thing that the H atoms did; they share their unpaired electrons to make a covalent bond.



Note that each F atom has a complete octet around it now:




We can also write this using a dash to represent the shared electron pair:




There are two different types of electrons in the fluorine diatomic molecule. The **bonding electron pair** makes the covalent bond. Each F atom has three other pairs of electrons that do not participate in the bonding; they are called **lone pair electrons**. Each F atom has one bonding pair and three lone pairs of electrons.

Covalent bonds can be made between different elements as well. One example is HF. Each atom starts out with an odd number of electrons in its valence shell:



The two atoms can share their unpaired electrons to make a covalent bond:



We note that the H atom has a full valence shell with two electrons, while the F atom has a complete octet of electrons.

Example 6.3.1:


Use Lewis electron dot diagrams to illustrate the covalent bond formation in HBr.

Solution

HBr is very similar to HF, except that it has Br instead of F. The atoms are as follows:



The two atoms can share their unpaired electron:



Exercise 6.3.1

Use Lewis electron dot diagrams to illustrate the covalent bond formation in Cl_2.

Answer



When working with covalent structures, it sometimes looks like you have leftover electrons. You apply the rules you learned so far, and there are still some electrons that remain unattached. You can't just leave them there. So where do you put them?

Multiple Covalent Bonds

Some molecules are not able to satisfy the octet rule by making only single covalent bonds between the atoms. Consider the compound ethene, which has a molecular formula of C_2H_4. The carbon atoms are bonded together, with each carbon also bonded to two hydrogen atoms.

$$\text{two C atoms} = 2 \times 4 = 8 \text{ valence electrons}$$

$$\text{four H atoms} = 4 \times 1 = 4 \text{ valence electrons}$$

$$\text{total of 12 valence electrons in the molecule}$$

If the Lewis electron dot structure was drawn with a single bond between the carbon atoms and with the octet rule followed, it would look like this:




Figure 6.3.1: Incorrect dot structure of ethene. (CK12 License)

This Lewis structure is incorrect because it contains a total of 14 electrons. However, the Lewis structure can be changed by eliminating the lone pairs on the carbon atoms and having to share two pairs instead of only one pair.




Figure 6.3.2: Correct dot structure for ethene. (CK12 License)

A **double covalent bond** is a covalent bond formed by atoms that share two pairs of electrons. The double covalent bond that occurs between the two carbon atoms in ethane can also be represented by a structural formula and with a molecular model as shown in the figure below.




Figure 6.3.3: (A) The structural model for C_2H_4 consists of a double covalent bond between the two carbon atoms and single bonds to the hydrogen atoms. (B) Molecular model of C_2H_4.

A **triple covalent bond** is a covalent bond formed by atoms that share three pairs of electrons. The element nitrogen is a gas that composes the majority of Earth's atmosphere. A nitrogen atom has five valence electrons, which can be shown as one pair and three single electrons. When combining with another nitrogen atom to form a diatomic molecule, the three single electrons on each atom combine to form three shared pairs of electrons.




Figure 6.3.4: Triple bond in N_2.

Each nitrogen atom follows the octet rule with one lone pair of electrons, and six electrons that are shared between the atoms.

Summary

- Covalent bonds are formed when atoms share electrons.
- Lewis electron dot diagrams can be drawn to illustrate covalent bond formation.
- Double bonds or triple bonds between atoms may be necessary to properly illustrate the bonding in some molecules.

Contributions & Attributions

-
- Anonymous by request

6.3: Covalent Lewis Structures- Electrons Shared is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.4: Covalent Lewis Structures- Electrons Shared** by Henry Agnew, Marisa Alviar-Agnew is licensed notset.

6.4: Writing Lewis Structures for Covalent Compounds

Learning Objectives

- Draw Lewis structures for covalent compounds.

The following procedure can be used to construct Lewis electron structures for more complex molecules and ions.

How-to: Constructing Lewis electron structures

1. Determine the total number of valence electrons in the molecule or ion.

- Add together the valence electrons from each atom. (Recall that the number of valence electrons is indicated by the position of the element in the periodic table.)
- If the species is a polyatomic ion, remember to add or subtract the number of electrons necessary to give the total charge on the ion.

  For CO_3^{2-}, for example, we add two electrons to the total because of the -2 charge.

2. Arrange the atoms to show specific connections.

- When there is a central atom, it is usually the least electronegative element in the compound. Chemists usually list this central atom first in the chemical formula (as in CCl_4 and CO_3^{2-}, which both have C as the central atom), which is another clue to the compound's structure.
- Hydrogen and the halogens are almost always connected to only one other atom, so they are usually *terminal* rather than central.

3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond.

- In H_2O, for example, there is a bonding pair of electrons between oxygen and each hydrogen.

4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).

- These electrons will usually be lone pairs.

5. If any electrons are left over, place them on the central atom.

- We will explain later that some atoms are able to accommodate more than eight electrons.

6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.

- This will not change the number of electrons on the terminal atoms.

7. Final check

- Always make sure all valence electrons are accounted for and that each atom has an octet of electrons, except for hydrogen (with two electrons).
- The central atom is usually the least electronegative element in the molecule or ion; hydrogen and the halogens are usually terminal.

Now let's apply this procedure to some particular compounds, beginning with one we have already discussed.


Example 6.4.1: Water

Write the Lewis Structure for H_2O.

Solution

Solutions to Example 10.4.1

Steps for Writing Lewis Structures	Example 6.4.1
1. Determine the total number of valence electrons in the molecule or ion.	Each H atom (group 1) has 1 valence electron, and the O atom (group 16) has 6 valence electrons, for a total of 8 valence electrons.


Steps for Writing Lewis Structures	Example 6.4.1
2. Arrange the atoms to show specific connections.	H O H Because H atoms are almost always terminal, the arrangement within the molecule must be HOH.
3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond. 4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).	Placing one bonding pair of electrons between the O atom and each H atom gives H -O- H with 4 electrons left over. Each H atom has a full valence shell of 2 electrons.
5. If any electrons are left over, place them on the central atom.	Adding the remaining 4 electrons to the oxygen (as two lone pairs) gives the following structure: 
6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.	Not necessary.
7. Final check.	The Lewis structure gives oxygen an octet and each hydrogen 2 electrons.



Example 6.4.2

Write the Lewis structure for the CH_2O molecule

Solution

Solutions to Example 10.4.2


Steps for Writing Lewis Structures	Example 6.4.2
1. Determine the total number of valence electrons in the molecule or ion.	Each hydrogen atom (group 1) has 1 valence electron, carbon (group 14) has 4 valence electrons, and oxygen (group 16) has 6 valence electrons, for a total of [(2)(1) + 4 + 6] = 12 valence electrons.
2. Arrange the atoms to show specific connections.	 Because carbon is less electronegative than oxygen and hydrogen is normally terminal, C must be the central atom.
3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond.	Placing a bonding pair of electrons between each pair of bonded atoms gives the following:  6 electrons are used, and 6 are left over.

Steps for Writing Lewis Structures	Example 6.4.2
4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).	Adding all 6 remaining electrons to oxygen (as three lone pairs) gives the following:  Although oxygen now has an octet and each hydrogen has 2 electrons, carbon has only 6 electrons.
5. If any electrons are left over, place them on the central atom.	Not necessary. There are no electrons left to place on the central atom.
6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.	To give carbon an octet of electrons, we use one of the lone pairs of electrons on oxygen to form a carbon–oxygen double bond: 
7. Final check	Both the oxygen and the carbon now have an octet of electrons, so this is an acceptable Lewis electron structure. The O has two bonding pairs and two lone pairs, and C has four bonding pairs. This is the structure of formaldehyde, which is used in embalming fluid.

? Exercise 6.4.1


Write Lewis electron structures for CO_2 and SCl_2, a vile-smelling, unstable red liquid that is used in the manufacture of rubber.

Answer CO_2



Carbon dioxide

Answer SCl_2



Sulfur dichloride

The United States Supreme Court has the unenviable task of deciding what the law is. This responsibility can be a major challenge when there is no clear principle involved or where there is a new situation not encountered before. Chemistry faces the same challenge in extending basic concepts to fit a new situation. Drawing of Lewis structures for polyatomic ions uses the same approach, but tweaks the process a little to fit a somewhat different set of circumstances.

Writing Lewis Structures for Polyatomic Ions (CK-12)

Recall that a polyatomic ion is a group of atoms that are covalently bonded together and which carry an overall electrical charge. The ammonium ion, NH_4^+, is formed when a hydrogen ion $\left(H^+\right)$ attaches to the lone pair of an ammonia $\left(NH_3\right)$ molecule in a coordinate covalent bond.




Figure 6.4.3: The ammonium ion. (CK12 License)

When drawing the Lewis structure of a polyatomic ion, the charge of the ion is reflected in the number of total valence electrons in the structure. In the case of the ammonium ion:

$$1 \text{ N atom} = 5 \text{ valence electrons}$$

$$4 \text{ H atoms} = 4 \times 1 = 4 \text{ valence electrons}$$

$$\text{subtract 1 electron for the } 1+ \text{ charge of the ion}$$


$$\text{total of 8 valence electrons in the ion}$$

It is customary to put the Lewis structure of a polyatomic ion into a large set of brackets, with the charge of the ion as a superscript outside of the brackets.

Exercise 6.4.2

Draw the Lewis electron dot structure for the sulfate ion.

Answer (CK12 License)



Exceptions to the Octet Rule (BC Campus)


As important and useful as the octet rule is in chemical bonding, there are some well-known violations. This does not mean that the octet rule is useless—quite the contrary. As with many rules, there are exceptions, or violations.

There are three violations to the octet rule. Odd-electron molecules represent the first violation to the octet rule. Although they are few, some stable compounds have an odd number of electrons in their valence shells. With an odd number of electrons, at least one atom in the molecule will have to violate the octet rule. Examples of stable odd-electron molecules are NO, NO_2, and ClO_2. The Lewis electron dot diagram for NO is as follows:




Although the O atom has an octet of electrons, the N atom has only seven electrons in its valence shell. Although NO is a stable compound, it is very chemically reactive, as are most other odd-electron compounds.


Electron-deficient molecules represent the second violation to the octet rule. These stable compounds have less than eight electrons around an atom in the molecule. The most common examples are the covalent compounds of beryllium and boron. For example, beryllium can form two covalent bonds, resulting in only four electrons in its valence shell:



Boron commonly makes only three covalent bonds, resulting in only six valence electrons around the B atom. A well-known example is BF_3:



The third violation to the octet rule is found in those compounds with more than eight electrons assigned to their valence shell. These are called expanded valence shell molecules. Such compounds are formed only by central atoms in the third row of the periodic table or beyond that have empty *d* orbitals in their valence shells that can participate in covalent bonding. One such compound is PF_5. The only reasonable Lewis electron dot diagram for this compound has the P atom making five covalent bonds:



Formally, the P atom has 10 electrons in its valence shell.

Example 6.4.3: Octet Violations

Identify each violation to the octet rule by drawing a Lewis electron dot diagram.


a. ClO
b. SF_6

Solution

a. With one Cl atom and one O atom, this molecule has 6 + 7 = 13 valence electrons, so it is an odd-electron molecule. A Lewis electron dot diagram for this molecule is as follows:



b. In SF_6, the central S atom makes six covalent bonds to the six surrounding F atoms, so it is an expanded valence shell molecule. Its Lewis electron dot diagram is as follows:



Exercise 6.4.3: Xenon Difluoride

Identify the violation to the octet rule in XeF_2 by drawing a Lewis electron dot diagram.

Answer



The Xe atom has an expanded valence shell with more than eight electrons around it.

Summary

Lewis dot symbols provide a simple rationalization of why elements form compounds with the observed stoichiometries. A plot of the overall energy of a covalent bond as a function of internuclear distance is identical to a plot of an ionic pair because both result from attractive and repulsive forces between charged entities. In Lewis electron structures, we encounter **bonding pairs**, which are shared by two atoms, and **lone pairs**, which are not shared between atoms. Lewis structures for polyatomic ions follow the same rules as those for other covalent compounds. There are three violations to the octet rule: odd-electron molecules, electron-deficient molecules, and expanded valence shell molecules.

6.4: Writing Lewis Structures for Covalent Compounds is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.5: Writing Lewis Structures for Covalent Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed notset. Original source: https://www.ck12.org/c/chemistry/.

https://chem.libretexts.org/@go/page/218508

https://chem.libretexts.org/@go/page/218508

6.5: Resonance - Equivalent Lewis Structures for the Same Molecule

Learning Objectives

- Explain the following laws within the Ideal Gas Law

Resonance

There are some cases in which more than one viable Lewis structure can be drawn for a molecule. An example is the ozone (O_3) molecule in Figure $6.5.1$. There are a total of 18 electrons in the structure and so the following two structures are possible.




Figure $6.5.1$: Resonance forms of ozone. Note the use of the double-headed arrow.

The structure on the left $(6.5.1)$ can be converted to the structure on the right by a shifting of electrons without altering the positions of the atoms.

It was once thought that the structure of a molecule such as O_3 consisted of one single bond and one double bond which then shifted back and forth as shown above. However, further studies showed that the two bonds are identical. Any double covalent bond between two given atoms is typically shorter than a single covalent bond. Studies of the O_3 and other similar molecules showed that the bonds were identical in length. Interestingly, the length of the bond is in between the lengths expected for an $O-O$ single bond and a double bond.

Resonance is the use of two or more Lewis structures to represent the covalent bonding in a molecule. One of the valid structures is referred to as a resonance structure. It is now understood that the true structure of a molecule which displays resonance is that of an average or a hybrid of all the resonance structures. In the case of the O_3 molecule, each of the covalent bonds between O atoms is best thought of as being "one and a half" bonds, as opposed to either a pure single bond or a pure double bond. This "half-bond" can be shown as a dotted line in both the Lewis structure and the molecular model (Figure $6.5.2$). Because of the half bond structure, each oxygen on either end of the single bond can be thought of as charged, since the central oxygen is donating the two electrons that make the single bond. The oxygen with three lone pairs therefore has one extra electron.




Figure $6.5.2$: "Half-bond" model of ozone molecule. This is a better description of the electornic structure of ozone than either of the resonance structures in Figure $6.5.1$.

Many polyatomic ions also display resonance. In some cases, the true structure may be an average of three valid resonance structures, as in the case of the nitrate ion, NO_3^- in Figure $6.5.3$.




Figure $6.5.3$: Resonance structure of nitrate anion.

The bond lengths between the central N atom and each O atom are identical and the bonds can be approximated as being equal to one and one-third bonds.

Summary

- Resonance structures are averages of different Lewis structure possibilities.
- Bond lengths are intermediate between covalent bonds and covalent double bonds.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

-
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

6.5: Resonance - Equivalent Lewis Structures for the Same Molecule is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

6.6: Predicting the Shapes of Molecules

Learning Objective

- Determine the shape of simple molecules.

Molecules have shapes. There is an abundance of experimental evidence to that effect—from their physical properties to their chemical reactivity. Small molecules—molecules with a single central atom—have shapes that can be easily predicted. The basic idea in molecular shapes is called **valence shell electron pair repulsion (VSEPR).** VSEPR says that electron pairs, being composed of negatively charged particles, repel each other to get as far away from one another as possible. VSEPR makes a distinction between *electron group geometry*, which expresses how electron groups (bonds and nonbonding electron pairs) are arranged, and *molecular geometry*, which expresses how the atoms in a molecule are arranged. However, the two geometries are related.

There are two types of **electron groups**: any type of bond—single, double, or triple—and lone electron pairs. When applying VSEPR to simple molecules, the first thing to do is to count the number of electron groups around the central atom. Remember that a multiple bond counts as only *one* electron group.

Any molecule with only two atoms is linear. A molecule whose central atom contains only two electron groups orients those two groups as far apart from each other as possible—180° apart. When the two electron groups are 180° apart, the atoms attached to those electron groups are also 180° apart, so the overall molecular shape is linear. Examples include BeH_2 and CO_2:



Figure 6.6.1: Beryllium hydride and carbon dioxide bonding.

The two molecules, shown in the figure below in a "ball and stick" model.




Figure 6.6.2: Beryllium hydride and carbon dioxide models. (CK12 Licence)

A molecule with three electron groups orients the three groups as far apart as possible. They adopt the positions of an equilateral triangle—120° apart and in a plane. The shape of such molecules is *trigonal planar*. An example is BF_3:




Figure 6.6.3: Boron trifluoride bonding. (CK12 Licence)

Some substances have a trigonal planar electron group distribution but have atoms bonded to only two of the three electron groups. An example is GeF_2:




Figure 6.6.4: Germanium difluoride bonding.

From an electron group geometry perspective, GeF_2 has a trigonal planar shape, but its real shape is dictated by the positions of the atoms. This shape is called *bent* or *angular*.

A molecule with four electron groups about the central atom orients the four groups in the direction of a tetrahedron, as shown in Figure 6.6.1 Tetrahedral Geometry. If there are four atoms attached to these electron groups, then the molecular shape is also

tetrahedral. Methane (CH_4) is an example.



Figure 6.6.5: Tetrahedral structure of methane. (CK12 Licence)

This diagram of CH_4 illustrates the standard convention of displaying a three-dimensional molecule on a two-dimensional surface. The straight lines are in the plane of the page, the solid wedged line is coming out of the plane toward the reader, and the dashed wedged line is going out of the plane away from the reader.




Figure 6.6.6: Methane bonding. (CK12 Licence)

NH_3 is an example of a molecule whose central atom has four electron groups, but only three of them are bonded to surrounding atoms.




Figure 6.6.7: Ammonia bonding. (CK12 Licence)

Although the electron groups are oriented in the shape of a tetrahedron, from a molecular geometry perspective, the shape of NH_3 is *trigonal pyramidal*.

H_2O is an example of a molecule whose central atom has four electron groups, but only two of them are bonded to surrounding atoms.




Figure 6.6.8: Water bonding.

Although the electron groups are oriented in the shape of a tetrahedron, the shape of the molecule is *bent* or *angular*. A molecule with four electron groups about the central atom, but only one electron group bonded to another atom, is linear because there are only two atoms in the molecule.

Double or triple bonds count as a single electron group. The Lewis electron dot diagram of formaldehyde (CH_2O) is shown in Figure 6.6.9.



Figure 6.6.9: Lewis Electron Dot Diagram of Formaldehyde.

The central C atom has three electron groups around it because the double bond counts as one electron group. The three electron groups repel each other to adopt a trigonal planar shape.




Figure 6.6.10: Formaldehyde bonding.

(The lone electron pairs on the O atom are omitted for clarity.) The molecule will not be a perfect equilateral triangle because the C–O double bond is different from the two C–H bonds, but both planar and triangular describe the appropriate approximate shape of this molecule.

Table 6.6.1 summarizes the shapes of molecules based on the number of electron groups and surrounding atoms.

Table 6.6.1: Summary of Molecular Shapes

Number of Electron Groups on Central Atom	Number of Bonding Groups	Number of Lone Pairs	Electron Geometry	Molecular Shape
2	2	0	linear	linear
3	3	0	trigonal planar	trigonal planar
3	2	1	trigonal planar	bent
4	4	0	tetrahedral	tetrahedral
4	3	1	tetrahedral	trigonal pyramidal
4	2	2	tetrahedral	bent

Example 6.6.1


What is the approximate shape of each molecule?

a. PCl_3
b. NOF

Solution


The first step is to draw the Lewis structure of the molecule.

For PCl_3, the electron dot diagram is as follows:



The lone electron pairs on the Cl atoms are omitted for clarity. The P atom has four electron groups with three of them bonded to surrounding atoms, so the molecular shape is trigonal pyramidal.

The electron dot diagram for NOF is as follows:



The N atom has three electron groups on it, two of which are bonded to other atoms. The molecular shape is bent.

Exercise 6.6.1

What is the approximate molecular shape of CH_2Cl_2?

Answer

Tetrahedral

Exercise 6.6.2

Ethylene (C_2H_4) has two central atoms. Determine the geometry around each central atom and the shape of the overall molecule. (Hint: hydrogen is a terminal atom.)

Answer

Trigonal planar about both central C atoms.

Summary

The approximate shape of a molecule can be predicted from the number of electron groups and the number of surrounding atoms.

6.6: Predicting the Shapes of Molecules is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.7: Predicting the Shapes of Molecules** by Henry Agnew, Marisa Alviar-Agnew is licensed notset.

6.7: Electronegativity and Polarity - Why Oil and Water Don't Mix

Learning Objectives

- Explain how polar compounds differ from nonpolar compounds.
- Determine if a molecule is polar or nonpolar.
- Given a pair of compounds, predict which would have a higher melting or boiling point.

Bond Polarity

The ability of an atom in a molecule to attract shared electrons is called **electronegativity**. When two atoms combine, the difference between their electronegativities is an indication of the type of bond that will form. If the difference between the electronegativities of the two atoms is small, neither atom can take the shared electrons completely away from the other atom, and the bond will be covalent. If the difference between the electronegativities is large, the more electronegative atom will take the bonding electrons completely away from the other atom (electron transfer will occur), and the bond will be ionic. This is why metals (low electronegativities) bonded with nonmetals (high electronegativities) typically produce ionic compounds.

A bond may be so polar that an electron actually transfers from one atom to another, forming a true ionic bond. How do we judge the degree of polarity? Scientists have devised a scale called **electronegativity**, a scale for judging how much atoms of any element attract electrons. Electronegativity is a unitless number; the higher the number, the more an atom attracts electrons. A common scale for electronegativity is shown in Figure 6.7.1.




Figure 6.7.1: Electronegativities of the Elements. Electronegativities are used to determine the polarity of covalent bonds.

The polarity of a covalent bond can be judged by determining the *difference* of the electronegativities of the two atoms involved in the covalent bond, as summarized in the following table:

difference of the electronegativities of the two atoms involved in the covalent bond

Electronegativity Difference	Bond Type
0–0.4	pure covalent
0.5–2.0	polar covalent
>2.0	likely ionic

Nonpolar Covalent Bonds

A bond in which the electronegativity difference is less than 1.9 is considered to be mostly covalent in character. However, at this point we need to distinguish between two general types of covalent bonds. A **nonpolar covalent bond** is a covalent bond in which the bonding electrons are shared equally between the two atoms. In a nonpolar covalent bond, the distribution of electrical charge is balanced between the two atoms.




Figure 6.7.2: A nonpolar covalent bond is one in which the distribution of electron density between the two atoms is equal.

The two chlorine atoms share the pair of electrons in the single covalent bond equally, and the electron density surrounding the Cl_2 molecule is symmetrical. Also note that molecules in which the electronegativity difference is very small (<0.5) are also considered nonpolar covalent. An example would be a bond between chlorine and bromine ($\Delta EN = 3.0 - 2.8 = 0.2$).

Polar Covalent Bonds

A bond in which the electronegativity difference between the atoms is between 0.5 and 2.0 is called a polar covalent bond. A **polar covalent bond** is a covalent bond in which the atoms have an unequal attraction for electrons and so the sharing is unequal. In a polar covalent bond, sometimes simply called a polar bond, the distribution of electrons around the molecule is no longer symmetrical.




Figure 6.7.3: In the polar covalent bond of HF, the electron density is unevenly distributed. There is a higher density (red) near the fluorine atom, and a lower density (blue) near the hydrogen atom.

An easy way to illustrate the uneven electron distribution in a polar covalent bond is to use the Greek letter delta (δ).




Figure 6.7.4: Use of δ to indicate partial charge.

The atom with the greater electronegativity acquires a partial negative charge, while the atom with the lesser electronegativity acquires a partial positive charge. The delta symbol is used to indicate that the quantity of charge is less than one. A crossed arrow can also be used to indicate the direction of greater electron density.



Figure 6.7.5: Use of crossed arrow to indicate polarity.



Electronegativity differences in bonding using Pauling scale. Differences in electronegativity classify bonds as covalent, polar covalent, or ionic.

Example 6.7.1: Bond Polarity

What is the polarity of each bond?

a. C–H
b. O–H

Solution

Using Figure 6.7.1, we can calculate the difference of the electronegativities of the atoms involved in the bond.

a. For the C–H bond, the difference in the electronegativities is 2.5 − 2.1 = 0.4. Thus we predict that this bond will be nonpolar covalent.
b. For the O–H bond, the difference in electronegativities is 3.5 − 2.1 = 1.4, so we predict that this bond will be polar covalent.

Exercise 6.7.1

What is the polarity of each bond?

a. Rb–F
b. P–Cl

Answer a

likely ionic

Answer b

polar covalent

Molecular Polarity

To determine if a molecule is polar or nonpolar, it is generally useful to look at Lewis structures. **Nonpolar compounds** will be symmetric, meaning all of the sides around the central atom are identical—bonded to the same element with no unshared pairs of electrons. **Polar molecules** are asymmetric, either containing lone pairs of electrons on a central atom or having atoms with different electronegativities bonded. This works pretty well, as long as you can visualize the molecular geometry. That's the hard part. To know how the bonds are oriented in space, you have to have a strong grasp of Lewis structures and VSEPR theory. Assuming that you do, you can look at the structure of each one and decide if it is polar or not, *whether or not you know the individual atom's electronegativity*. This is because you know that all bonds between dissimilar elements are polar, and in these particular examples, it doesn't matter which direction the dipole moment vectors are pointing (out or in).

A **polar molecule** is a molecule in which one end of the molecule is slightly positive, while the other end is slightly negative. A diatomic molecule that consists of a polar covalent bond, such as HF, is a polar molecule. The two electrically charged regions on either end of the molecule are called poles, similar to a magnet having a north and a south pole. A molecule with two poles is called a **dipole** (see figure below). Hydrogen fluoride is a dipole.



Figure 6.7.6: A dipole is any molecule with a positive end and a negative end, resulting from unequal distribution of electron density throughout the molecule.

For molecules with more than two atoms, the molecular geometry must also be taken into account when determining if the molecule is polar or nonpolar. The figure below shows a comparison between carbon dioxide and water. Carbon dioxide (CO_2) is a linear molecule. The oxygen atoms are more electronegative than the carbon atom, so there are two individual dipoles pointing outward from the C atom to each O atom. However, since the dipoles are of equal strength and are oriented this way, they cancel out and the overall molecular polarity of CO_2 is zero.

Water is a bent molecule because of the two lone pairs on the central oxygen atom. The individual dipoles point from the H atoms toward the O atom. Because of the shape, the dipoles do not cancel each other out and the water molecule is polar. In the figure below, the net dipole is shown in blue and points upward.




Figure 6.7.7: The molecular geometry of a molecule affects its polarity. Each CO bond has a dipole moment, but they point in opposite directions so that the net CO2 molecule is nonpolar. In contrast, water is polar because the OH bond moments do not cancel out.

Some other molecules are shown in the figure below. Notice that a tetrahedral molecule such as CH_4 is nonpolar. However, if one of the peripheral H atoms is replaced with another atom that has a different electronegativity, the molecule becomes polar. A trigonal planar molecule (BF_3) may be nonpolar if all three peripheral atoms are the same, but a trigonal pyramidal molecule (NH_3) is polar.




Figure 6.7.8: Some examples of polar and nonpolar molecules based on molecular geometry.

To summarize, to be polar, a molecule must:

1. Contain at least one polar covalent bond.
2. Have a molecular structure such that the sum of the vectors of each bond dipole moment do not cancel.

Steps to Identify Polar Molecules

1. Draw the Lewis structure.
2. Figure out the geometry (using VSEPR theory).
3. Visualize or draw the geometry.
4. Find the net dipole moment (you don't have to actually do calculations if you can visualize it).
5. If the net dipole moment is zero, it is non-polar. Otherwise, it is polar.

Properties of Polar Molecules

Polar molecules tend to align when placed in an electric field with the positive end of the molecule oriented toward the negative plate and the negative end toward the positive plate (Figure 6.7.14). We can use an electrically charged object to attract polar molecules, but nonpolar molecules are not attracted. Also, polar solvents are better at dissolving polar substances, and nonpolar solvents are better at dissolving nonpolar substances.




Figure 6.7.9: (a) Molecules are always randomly distributed in the liquid state in the absence of an electric field. (b) When an electric field is applied, polar molecules like HF will align to the dipoles with the field direction. (OpenStax CC-BY-SA);

While molecules can be described as "polar covalent" or "ionic", it must be noted that this is often a relative term, with one molecule simply being *more polar* or *less polar* than another. However, the following properties are typical of such molecules. Polar molecules tend to:

- have higher melting points than nonpolar molecules
- have higher boiling points than nonpolar molecules
- be more soluble in water (dissolve better) than nonpolar molecules
- have lower vapor pressures than nonpolar molecules

Example 6.7.2:

Label each of the following as polar or nonpolar.

a. Water, H_2O:

b. Methanol, CH_3OH:

c. Hydrogen Cyanide, HCN: H—C≡N

d. Oxygen, O_2:

e. Propane, C_3H_8:

Solution

a. Water is polar. Any molecule with lone pairs of electrons around the central atom is polar.
b. Methanol is polar. This is not a symmetric molecule. The $-\mathrm{OH}$ side is different from the other 3 $-\mathrm{H}$ sides.
c. Hydrogen cyanide is polar. The molecule is not symmetric. The nitrogen and hydrogen have different electronegativities, creating an uneven pull on the electrons.
d. Oxygen is nonpolar. The molecule is symmetric. The two oxygen atoms pull on the electrons by exactly the same amount.
e. Propane is nonpolar, because it is symmetric, with H atoms bonded to every side around the central atoms and no unshared pairs of electrons.

? Exercise 6.7.2

Label each of the following as polar or nonpolar.

a. SO_3
b. NH_3

Answer a

nonpolar

Answer b

polar

Contributions & Attributions

- StackExchange (thomij).

6.7: Electronegativity and Polarity - Why Oil and Water Don't Mix is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.8: Electronegativity and Polarity - Why Oil and Water Do not Mix** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

7: Chemical Reactions

7.1: Prelude to Introduction to Chemical Reactions
7.2: Laws of Conservation : A Review
7.3: Evidence of a Chemical Reaction
7.4: Chemical Equations
7.5: How to Write Balanced Chemical Equations
7.6: Aqueous Solutions and Solubility - Compounds Dissolved in Water
7.7: Precipitation Reactions
7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
7.9: Acid–Base and Gas Evolution Reactions
7.10: Oxidation-Reduction Reactions
7.11: Classifying Chemical Reactions
7.12: The Activity Series- Predicting Spontaneous Redox Reactions

7: Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

7.1: Prelude to Introduction to Chemical Reactions

Although yeast has been used for thousands of years, its true nature has been known only for the last two centuries. Yeasts are single-celled fungi. About 1,000 species are recognized, but the most common species is *Saccharomyces cerevisiae*, which is used in bread making. Other species are used for the fermentation of alcoholic beverages. Some species can cause infections in humans.

Yeasts live primarily on sugars, such as glucose ($C_6H_{12}O_6$). They convert glucose into carbon dioxide (CO_2) and ethanol (C_2H_5OH) in a chemical transformation that is represented as follows:

$$C_6H_{12}O_6(s) \rightarrow 2\,CO_2(g) + 2\,C_2H_5OH(\ell)$$

Bread making depends on the production of carbon dioxide. The gas, which is produced in tiny pockets in bread dough, acts as a leavening agent: it expands during baking and makes the bread rise. Leavened bread is softer, lighter, and easier to eat and chew than unleavened bread. The other major use of yeast, fermentation, depends on the production of ethanol, which results from the same chemical transformation. Some alcoholic beverages, such as champagne, can also be carbonated using the carbon dioxide produced by the yeast.

Yeast is among the simplest life forms on Earth, yet it is absolutely necessary for at least two major food industries. Without yeast to turn dough into bread and juice into wine, these foods and food industries would not exist today.

7.1: Prelude to Introduction to Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **5.0: Prelude to Introduction to Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.2: Laws of Conservation : A Review

Learning Objectives

- Correctly define a law as it pertains to science.
- State the law of conservation of matter.

In science, a law is a general statement that explains a large number of observations. Before being accepted, a law must be verified many times under many conditions. Laws are therefore considered the highest form of scientific knowledge and are generally thought to be inviolable. Scientific laws form the core of scientific knowledge. One scientific law that provides the foundation for understanding in chemistry is the law of conservation of matter. It states that in any given system that is closed to the transfer of matter (in and out), the amount of matter in the system stays constant. A concise way of expressing this law is to say that the amount of matter in a system is *conserved*.

With the development of more precise ideas on elements, compounds and mixtures, scientists began to investigate how and why substances react. French chemist A. Lavoisier laid the foundation to the scientific investigation of matter by describing that substances react by following certain laws. These laws are called the laws of chemical combination. These eventually formed the basis of Dalton's Atomic Theory of Matter.

Law of Conservation of Mass

According to this law, during any physical or chemical change, the total mass of the products remains equal to the total mass of the reactants.

$$\overbrace{\underbrace{\mathrm{HgO(s)}}_{100\ \mathrm{g}}}^{\text{Mercuric oxide}} \rightarrow \overbrace{\underbrace{\mathrm{Hg(l)}}_{92.6\ \mathrm{g}}}^{\text{Mercury}} + \overbrace{\underbrace{\mathrm{O_2(g)}}_{7.4\ \mathrm{g}}}^{\text{Oxygen}} \tag{7.2.1}$$

Another way of stating this is, "In a chemical reaction, matter is neither created nor destroyed." The law of conservation of mass is also known as the "law of indestructibility of matter."

Example 7.2.1

If heating 10 grams of $\mathrm{CaCO_3}$ produces 4.4 g of $\mathrm{CO_2}$ and 5.6 g of CaO, show that these observations are in agreement with the law of conservation of mass.



A sample of calcium carbonate ($CaCO_3$). (Public Domain; Walkerma).

Solution

- Mass of the reactants: $10\ g$
- Mass of the products: $4.4\ g + 5.6\ g = 10\ g$.

Because the mass of the reactants is equal to the mass of the products, the observations are in agreement with the law of conservation of mass.

What does this mean for chemistry? In any chemical change, one or more initial substances change into a different substance or substances. Both the initial and final substances are composed of atoms because all matter is composed of atoms. According to the law of conservation of matter, matter is neither created nor destroyed, so we must have the same number and type of atoms after the chemical change as were present before the chemical change.

Before looking at explicit examples of the law of conservation of matter, we need to examine the method chemists use to represent chemical changes.

Exercise 7.2.1

a. What is the law of conservation of matter?
b. How does the law of conservation of matter apply to chemistry?

Answer a:

The law of conservation of matter states that in any given system that is closed to the transfer of matter, the amount of matter in the system stays constant

Answer b:

The law of conservation of matter says that in chemical reactions, the total mass of the products must equal the total mass of the reactants.

Key Takeaway

The amount of matter in a closed system is conserved.

Contributors and Attributions

- Binod Shrestha (University of Lorraine)

Exercises

1. Express the law of conservation of matter in your own words.
2. Explain why the concept of conservation of matter is considered a scientific law.

Answer

1. Matter may not be created or destroyed.

2. The concept is a scientific law because it is based on experimentation.

7.2: Laws of Conservation : A Review is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

7.3: Evidence of a Chemical Reaction

Learning Objectives

- Identify the evidence for chemical reactions.

In a chemical change, new substances are formed. In order for this to occur, the chemical bonds of the substances break, and the atoms that compose them separate and rearrange themselves into new substances with new chemical bonds. When this process occurs, we call it a chemical reaction. A **chemical reaction** is the process in which one or more substances are changed into one or more new substances.




Figure 7.3.1: (a) Copper and nitric acid undergo a chemical change to form copper nitrate and brown, gaseous nitrogen dioxide. (b) During the combustion of a match, cellulose in the match and oxygen from the air undergo a chemical change to form carbon dioxide and water vapor. (c) Cooking red meat causes a number of chemical changes, including the oxidation of iron in myoglobin that results in the familiar red-to-brown color change. (d) A banana turning brown is a chemical change as new, darker (and less tasty) substances form. (Credit b: modification of work by Jeff Turner; credit c: modification of work by Gloria Cabada-Leman; credit d: modification of work by Roberto Verzo.)

To identify a chemical reaction, we look for a **chemical change**. A chemical change always produces one or more types of matter that differ from the matter present before the change. The formation of rust is a chemical change because rust is a different kind of matter than the iron, oxygen, and water present before the rust formed. The explosion of nitroglycerin is a chemical change because the gases produced are very different kinds of matter from the original substance. Other examples of chemical changes include: reactions that are performed in a lab (such as copper reacting with nitric acid), all forms of combustion (burning), and food being cooked, digested, or rotting (Figure 7.3.1).



Video **7.3.1** *: Evidence of a Chemical Reaction*

Example 7.3.1: Evidence of a Chemical Reaction

Which of the following is a chemical reaction?

a. Freezing liquid mercury.
b. Adding yellow to blue to make green.
c. Cutting a piece of paper into two pieces.
d. Dropping a sliced orange into a vat of sodium dydroxide.
e. Filling a balloon with natural air.

Solution

A, B, C, & E involve only physical changes. A sliced orange has acid (citric acid) that can react with sodium hydroxide, so the answer is D.

Exercise 7.3.1

Which of the following is a chemical reaction?

a. Painting a wall blue.
b. A bicycle rusting.
c. Ice cream melting.
d. Scratching a key across a desk.
e. Making a sand castle.

Answer

B

Example 7.3.2: Evidence of a Chemical Reaction

Which of the following is not a chemical reaction?

a. Shattering glass with a baseball.
b. Corroding metal.
c. Fireworks exploding.
d. Lighting a match.
e. Baking a cake.

Solution

Shattering glass with a baseball results in glass broken into many pieces but no chemical change happens, so the answer is A.

Exercise 7.3.2

Which of the following is NOT a chemical reaction?

a. Frying an egg.
b. Slicing carrots.
c. A Macbook falling out of a window.
d. Creating ATP in the human body.
e. Dropping a fizzy tablet into a glass of water.

Answer

B and C

Summary

Chemical reactions can be identified via a wide range of different observable factors including change in color, energy change (temperature change or light produced), gas production, formation of precipitate and change in properties.

7.3: Evidence of a Chemical Reaction is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.2: Evidence of a Chemical Reaction** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY 4.0. Original source: https://www.ck12.org/c/chemistry/.

7.4: Chemical Equations

Learning Objectives

- Identify the reactants and products in any chemical reaction.
- Convert word equations into chemical equations.
- Use the common symbols, (s), (l), (g), (aq), and $\rightarrow$ appropriately when writing a chemical reaction.

In a chemical change, new substances are formed. In order for this to occur, the chemical bonds of the substances break, and the atoms that compose them separate and rearrange themselves into new substances with new chemical bonds. When this process occurs, we call it a chemical reaction. A **chemical reaction** is the process in which one or more substances are changed into one or more new substances.

Reactants and Products

To describe a chemical reaction, we need to indicate what substances are present at the beginning and what substances are present at the end. The substances that are present at the beginning are called **reactants** and the substances present at the end are called **products**.

Sometimes when reactants are put into a reaction vessel, a reaction will take place to produce products. Reactants are the starting materials, that is, whatever we have as our initial ingredients. The products are just that—what is produced—or the result of what happens to the reactants when we put them together in the reaction vessel. If we think about baking chocolate chip cookies, our reactants would be flour, butter, sugar, vanilla, baking soda, salt, egg, and chocolate chips. What would be the products? Cookies! The reaction vessel would be our mixing bowl.

$$\underbrace{\text{Flour} + \text{Butter} + \text{Sugar} + \text{Vanilla} + \text{Baking Soda} + \text{Eggs} + \text{Chocolate Chips}}_{\text{Ingredients} = \text{Reactants}} \rightarrow \underbrace{\text{Cookies}}_{\text{Product}}$$

Writing Chemical Equations

When sulfur dioxide is added to oxygen, sulfur trioxide is produced. Sulfur dioxide and oxygen, $\mathrm{SO_2} + \mathrm{O_2}$, are reactants and sulfur trioxide, $\mathrm{SO_3}$, is the product.

$$\underbrace{2\,\mathrm{SO_2(g)} + \mathrm{O_2(g)}}_{\text{Reactants}} \rightarrow \underbrace{2\,\mathrm{SO_3(g)}}_{\text{Products}}$$

In chemical reactions, the reactants are found before the symbol "$\rightarrow$" and the products are found after the symbol "$\rightarrow$". The general equation for a reaction is:

$$\text{Reactants} \rightarrow \text{Products}$$

There are a few special symbols that we need to know in order to "talk" in chemical shorthand. In the table below is the summary of the major symbols used in chemical equations. Table $7.4.1$ shows a listing of symbols used in chemical equations.


Table $7.4.1$: Symbols Used in Chemical Equations

Symbol	Description	Symbol	Description
$+$	used to separate multiple reactants or products	(s)	reactant or product in the solid state
$\rightarrow$	yield sign; separates reactants from products	(l)	reactant or product in the liquid state
$\rightleftharpoons$	replaces the yield sign for reversible reactions that reach equilibrium	(g)	reactant or product in the gas state

Symbol	Description	Symbol	Description
$\xrightarrow{\text{Pt}}$	formula written above the arrow is used as a catalyst in the reaction	(aq)	reactant or product in an aqueous solution (dissolved in water)
$\xrightarrow{\Delta}$	triangle indicates that the reaction is being heated		

Chemists have a choice of methods for describing a chemical reaction.

1. They could draw a picture of the chemical reaction.



2. They could write a word equation for the chemical reaction:
"Two molecules of hydrogen gas react with one molecule of oxygen gas to produce two molecules of water vapor."

3. They could write the equation in chemical shorthand.

$$2\text{H}_2\,(g) + \text{O}_2\,(g) \rightarrow 2\text{H}_2\text{O}\,(g)$$

In the symbolic equation, chemical formulas are used instead of chemical names for reactants and products, while symbols are used to indicate the phase of each substance. It should be apparent that the chemical shorthand method is the quickest and clearest method for writing chemical equations.

We could write that an aqueous solution of calcium nitrate is added to an aqueous solution of sodium hydroxide to produce solid calcium hydroxide and an aqueous solution of sodium nitrate. Or in shorthand we could write:

$$\text{Ca}(\text{NO}_3)_2\,(aq) + 2\text{NaOH}\,(aq) \rightarrow \text{Ca}(\text{OH})_2\,(s) + 2\text{NaNO}_3\,(aq)$$

How much easier is that to read? Let's try it in reverse. Look at the following reaction in shorthand and write the word equation for the reaction:

$$\text{Cu}\,(s) + \text{AgNO}_3\,(aq) \rightarrow \text{Cu}(\text{NO}_3)_2\,(aq) + \text{Ag}\,(s)$$

The word equation for this reaction might read something like "solid copper reacts with an aqueous solution of silver nitrate to produce a solution of copper (II) nitrate with solid silver."

To turn word equations into symbolic equations, we need to follow the given steps:

1. Identify the reactants and products. This will help you know which symbols go on each side of the arrow and where the $+$ signs go.
2. Write the correct formulas for all compounds. You will need to use the rules you learned in Chapter 5 (including making all ionic compounds charge balanced).
3. Write the correct formulas for all elements. Usually this is given straight off of the periodic table. However, there are seven elements that are considered diatomic, meaning that they are always found in pairs in nature. They include those elements listed in the table.

Table 7.4.1*: Diatomic Elements*

Element Name	Hydrogen	Nitrogen	Oxygen	Fluorine	Chlorine	Bromine	Iodine
Formula	H_2	N_2	O_2	F_2	Cl_2	Br_2	I_2

Example 7.4.1

Transfer the following symbolic equations into word equations or word equations into symbolic equations.

a. $\mathrm{HCl}\,(aq) + \mathrm{NaOH}\,(aq) \rightarrow \mathrm{NaCl}\,(aq) + \mathrm{H_2O}\,(l)$
b. Gaseous propane, $\mathrm{C_3H_8}$, burns in oxygen gas to produce gaseous carbon dioxide and liquid water.
c. Hydrogen fluoride gas reacts with an aqueous solution of potassium carbonate to produce an aqueous solution of potassium fluoride, liquid water, and gaseous carbon dioxide.

Solution

a. An aqueous solution of hydrochloric acid reacts with an aqueous solution of sodium hydroxide to produce an aqueous solution of sodium chloride and liquid water.

b. Reactants: propane ($\mathrm{C_3H_8}$) and oxygen ($\mathrm{O_2}$)

Product: carbon dioxide ($\mathrm{CO_2}$) and water ($\mathrm{H_2O}$)

$$\mathrm{C_3H_8}\,(g) + \mathrm{O_2}\,(g) \rightarrow \mathrm{CO_2}\,(g) + \mathrm{H_2O}\,(l)$$

c. Reactants: hydrogen fluoride and potassium carbonate

Products: potassium fluoride, water, and carbon dioxide

$$\mathrm{HF}\,(g) + \mathrm{K_2CO_3}\,(aq) \rightarrow \mathrm{KF}\,(aq) + \mathrm{H_2O}\,(l) + \mathrm{CO_2}\,(g)$$

Exercise 7.4.1

Transfer the following symbolic equations into word equations or word equations into symbolic equations.

a. Hydrogen gas reacts with nitrogen gas to produce gaseous ammonia.
b. $\mathrm{HCl}\,(aq) + \mathrm{LiOH}\,(aq) \rightarrow \mathrm{LiCl}\,(aq) + \mathrm{H_2O}\,(l)$
c. Copper metal is heated with oxygen gas to produce solid copper(II) oxide.

Answer a

$H_2(g) + N_2(g) \rightarrow NH_3(g)$

Answer b

An aqueous solution of hydrochloric acid reacts with an aqueous solution of lithium hydroxide to produce an aqueous solution of lithium chloride and liquid water.

Answer c

$Cu(s) + O_2(g) \rightarrow CuO(s)$

Summary

- A chemical reaction is the process by which one or more substances are changed into one or more new substances.
- Chemical reactions are represented by chemical equations.
- Chemical equations have reactants on the left, an arrow that is read as "yields", and the products on the right.

7.4: Chemical Equations is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.3: The Chemical Equation** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.5: How to Write Balanced Chemical Equations

Learning Objectives

- Explain the roles of subscripts and coefficients in chemical equations.
- Balance a chemical equation when given the unbalanced equation.
- Explain the role of the Law of Conservation of Mass in a chemical reaction.

Even though chemical compounds are broken up and new compounds are formed during a chemical reaction, atoms in the reactants do not disappear, nor do new atoms appear to form the products. In chemical reactions, atoms are never created or destroyed. The same atoms that were present in the reactants are present in the products—they are merely reorganized into different arrangements. In a complete chemical equation, the two sides of the equation must be present on the reactant and the product sides of the equation.

Coefficients and Subscripts

There are two types of numbers that appear in chemical equations. There are subscripts, which are part of the chemical formulas of the reactants and products; and there are coefficients that are placed in front of the formulas to indicate how many molecules of that substance is used or produced.





Figure 7.5.1: Balancing Equations. You cannot change subscripts in a chemical formula to balance a chemical equation; you can change only the coefficients. Changing subscripts changes the ratios of atoms in the molecule and the resulting chemical properties. For example, water (H_2O) and hydrogen peroxide (H_2O_2) are chemically distinct substances. H_2O_2 decomposes to H_2O and O_2 gas when it comes in contact with the metal platinum, whereas no such reaction occurs between water and platinum.

The **subscripts** are part of the formulas and once the formulas for the reactants and products are determined, the subscripts may not be changed. The **coefficients** indicate the number of each substance involved in the reaction and may be changed in order to balance the equation. The equation above indicates that one mole of solid copper is reacting with two moles of aqueous silver nitrate to produce one mole of aqueous copper (II) nitrate and two atoms of solid silver.

Balancing a Chemical Equation

Because the identities of the reactants and products are fixed, the equation cannot be balanced by changing the subscripts of the reactants or the products. To do so would change the chemical identity of the species being described, as illustrated in Figure 7.5.1.



Original molecule H_2O: if the coefficient 2 is added in front, that makes 2 water molecules; but if the subscript 2 is added to make H_2O_2, that's hydrogen peroxide.

The simplest and most generally useful method for balancing chemical equations is “inspection,” better known as trial and error. The following is an efficient approach to balancing a chemical equation using this method.

Steps in Balancing a Chemical Equation

1. Identify the most complex substance.
2. Beginning with that substance, choose an element(s) that appears in only one reactant and one product, if possible. Adjust the coefficients to obtain the same number of atoms of this element(s) on both sides.
3. Balance polyatomic ions (if present on both sides of the chemical equation) as a unit.
4. Balance the remaining atoms, usually ending with the least complex substance and using fractional coefficients if necessary. If a fractional coefficient has been used, multiply both sides of the equation by the denominator to obtain whole numbers for the coefficients.
5. Count the numbers of atoms of each kind on both sides of the equation to be sure that the chemical equation is balanced.

Example 7.5.1: Combustion of Heptane

Balance the chemical equation for the combustion of Heptane (C_7H_{16}).

$$C_7H_{16}(l) + O_2(g) \rightarrow CO_2(g) + H_2O(g)$$

Solution

Solutions to Example 7.5.1

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is the one with the largest number of different atoms, which is C_7H_{16}. We will assume initially that the final balanced chemical equation contains 1 molecule or formula unit of this substance.
2. **Adjust the coefficients.**	a. Because one molecule of n-heptane contains 7 carbon atoms, we need 7 CO_2 molecules, each of which contains 1 carbon atom, on the right side: $$C_7H_{16}(l) + O_2(g) \rightarrow \underline{7}CO_2(g) + H_2O(g)$$ • 7 carbon atoms on both reactant and product sides b. Because one molecule of n-heptane contains 16 hydrogen atoms, we need 8 H_2O molecules, each of which contains 2 hydrogen atoms, on the right side: $$C_7H_{16}(l) + O_2(g) \rightarrow 7\,CO_2(g) + \underline{8}H_2O(g)$$ • 16 hydrogen atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	There are no polyatomic ions to be considered in this reaction.
4. **Balance the remaining atoms.**	The carbon and hydrogen atoms are now balanced, but we have 22 oxygen atoms on the right side and only 2 oxygen atoms on the left. We can balance the oxygen atoms by adjusting the coefficient in front of the least complex substance, O_2, on the reactant side: $$C_7H_{16}(l) + \underline{11}O_2(g) \rightarrow 7\,CO_2(g) + 8\,H_2O(g)$$ • 22 oxygen atoms on both reactant and product sides
5. **Check your work.**	The equation is now balanced, and there are no fractional coefficients: there are 7 carbon atoms, 16 hydrogen atoms, and 22 oxygen atoms on each side. Always check to be sure that a chemical equation is balanced.

Example 7.5.2: Combustion of Isooctane

Combustion of Isooctane (C_8H_{18})

$$C_8H_{18}(l) + O_2(g) \longrightarrow CO_2(g) + H_2O(g)$$

Solution

The assumption that the final balanced chemical equation contains only one molecule or formula unit of the most complex substance is not always valid, but it is a good place to start. The combustion of any hydrocarbon with oxygen produces carbon dioxide and water.

Solutions to Example 7.4.2

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is the one with the largest number of different atoms, which is C_8H_{18}. We will assume initially that the final balanced chemical equation contains 1 molecule or formula unit of this substance.
2. **Adjust the coefficients.**	a. The first element that appears only once in the reactants is carbon: 8 carbon atoms in isooctane means that there must be 8 CO_2 molecules in the products: $$C_8H_{18}(l) + O_2(g) \longrightarrow \underline{8}CO_2(g) + H_2O(g)$$ • 8 carbon atoms on both reactant and product sides b. 18 hydrogen atoms in isooctane means that there must be 9 H_2O molecules in the products: $$C_8H_{18}(l) + O_2(g) \longrightarrow 8\,CO_2(g) + \underline{9}H_2O(g)$$ • 18 hydrogen atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	There are no polyatomic ions to be considered in this reaction.
4. **Balance the remaining atoms.**	The carbon and hydrogen atoms are now balanced, but we have 25 oxygen atoms on the right side and only 2 oxygen atoms on the left. We can balance the least complex substance, O_2, but because there are 2 oxygen atoms per O_2 molecule, we must use a fractional coefficient ($\frac{25}{2}$) to balance the oxygen atoms: $$C_8H_{18}(l) + \underline{\frac{25}{2}}O_2(g) \rightarrow 8\,CO_2(g) + 9\,H_2O(g)$$ • 25 oxygen atoms on both reactant and product sides The equation is now balanced, but we usually write equations with whole number coefficients. We can eliminate the fractional coefficient by multiplying all coefficients on both sides of the chemical equation by **2**: $$\underline{2}C_8H_{18}(l) + \underline{25}O_2(g) \longrightarrow \underline{16}CO_2(g) + \underline{18}H_2O(g)$$

Steps	Example
5. **Check your work.**	The balanced chemical equation has 16 carbon atoms, 36 hydrogen atoms, and 50 oxygen atoms on each side. Balancing equations requires some practice on your part as well as some common sense. If you find yourself using very large coefficients or if you have spent several minutes without success, go back and make sure that you have written the formulas of the reactants and products correctly.

Example 7.5.3: Precipitation of Lead (II) Chloride

Aqueous solutions of lead (II) nitrate and sodium chloride are mixed. The products of the reaction are an aqueous solution of sodium nitrate and a solid precipitate of lead (II) chloride. Write the balanced chemical equation for this reaction.

Solution

Solutions to Example 7.4.3

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is lead (II) chloride. $\mathrm{Pb(NO_3)_2(aq) + NaCl(aq) \rightarrow NaNO_3(aq) + PbCl_2(s)}$
2. **Adjust the coefficients.**	There are twice as many chloride ions in the reactants as there are in the products. Place a 2 in front of the NaCl in order to balance the chloride ions. $\mathrm{Pb(NO_3)_2(aq) + \underline{2}NaCl(aq) \rightarrow NaNO_3(aq) + PbCl_2(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on reactant side, 1 Na atom on product side • 2 Cl atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	The nitrate ions are still unbalanced. Place a 2 in front of the $NaNO_3$. The result is: $\mathrm{Pb(NO_3)_2(aq) + 2\,NaCl(aq) \rightarrow \underline{2}NaNO_3(aq) + PbCl_2(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on both reactant and product sides • 2 Cl atoms on both reactant and product sides • 2 NO_3^- atoms on both reactant and product sides
4. **Balance the remaining atoms.**	There is no need to balance the remaining atoms because they are already balanced.
5. **Check your work.**	$\mathrm{Pb(NO_3)_2(aq) + 2\,NaCl(aq) \rightarrow 2\,NaNO_3(aq) + PbCl_2(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on both reactant and product sides • 2 Cl atoms on both reactant and product sides • 2 NO_3^- atoms on both reactant and product sides

Exercise 7.5.1

Is each chemical equation balanced?

a. $2\,Hg(\ell) + O_2(g) \rightarrow Hg_2O_2(s)$
b. $C_2H_4(g) + 2\,O_2(g) \rightarrow 2\,CO_2(g) + 2\,H_2O(g)$
c. $Mg(NO_3)_2(s) + 2\,Li(s) \rightarrow Mg(s) + 2\,LiNO_3(s)$

Answer a

yes

Answer b

no

Answer c

yes

Exercise 7.5.2

Balance the following chemical equations.

a. $N_2(g) + O_2(g) \rightarrow NO_2(g)$
b. $Pb(NO_3)_2(aq) + FeCl_3(aq) \rightarrow Fe(NO_3)_3(aq) + PbCl_2(s)$
c. $C_6H_{14}(l) + O_2(g) \rightarrow CO_2(g) + H_2O(g)$

Answer a

$N_2\,(g) + 2O_2\,(g) \rightarrow 2NO_2\,(g)$

Answer b

$3Pb(NO_3)_2(aq) + 2FeCl_3(aq) \rightarrow 2Fe(NO_3)_3(aq) + 3PbCl_2(s)$

Answer c

$2C_6H_{14}(l) + 19O_2(g) \rightarrow 12CO_2(g) + 14H_2O(g)$

Summary

- To be useful, chemical equations must always be balanced. Balanced chemical equations have the same number and type of each atom on both sides of the equation.
- The coefficients in a balanced equation must be the simplest whole number ratio. Mass is always conserved in chemical reactions.

Vocabulary

- **Chemical reaction** - The process in which one or more substances are changed into one or more new substances.
- **Reactants** - The starting materials in a reaction.
- **Products** - Materials present at the end of a reaction.
- **Balanced chemical equation** - A chemical equation in which the number of each type of atom is equal on the two sides of the equation.
- **Subscripts** - Part of the chemical formulas of the reactants and products that indicate the number of atoms of the preceding element.
- **Coefficient** - A small whole number that appears in front of a formula in a balanced chemical equation.

7.5: How to Write Balanced Chemical Equations is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.4: How to Write Balanced Chemical Equations** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

7.6: Aqueous Solutions and Solubility - Compounds Dissolved in Water

Learning Objectives

- Define and give examples of electrolytes.

When some substances are dissolved in water, they undergo either a physical or a chemical change that yields ions in solution. These substances constitute an important class of compounds called electrolytes. Substances that do not yield ions when dissolved are called nonelectrolytes. If the physical or chemical process that generates the ions is essentially 100% efficient (all of the dissolved compound yields ions), then the substance is known as a strong electrolyte. If only a relatively small fraction of the dissolved substance undergoes the ion-producing process, it is called a weak electrolyte.

Substances may be identified as strong, weak, or nonelectrolytes by measuring the electrical conductance of an aqueous solution containing the substance. To conduct electricity, a substance must contain freely mobile, charged species. Most familiar is the conduction of electricity through metallic wires, in which case the mobile, charged entities are electrons. Solutions may also conduct electricity if they contain dissolved ions, with conductivity increasing as ion concentration increases. Applying a voltage to electrodes immersed in a solution permits assessment of the relative concentration of dissolved ions, either quantitatively, by measuring the electrical current flow, or qualitatively, by observing the brightness of a light bulb included in the circuit (Figure 7.6.1).




Figure 7.6.1: Solutions of nonelectrolytes, such as ethanol, do not contain dissolved ions and cannot conduct electricity. Solutions of electrolytes contain ions that permit the passage of electricity. The conductivity of an electrolyte solution is related to the strength of the electrolyte. This diagram shows three separate beakers. Each has a wire plugged into a wall outlet. In each case, the wire leads from the wall to the beaker and is split resulting in two ends. One end leads to a light bulb and continues on to a rectangle labeled with a plus sign. The other end leads to a rectangle labeled with a minus sign. The rectangles are in a solution. In the first beaker, labeled “Ethanol No Conductivity,” four pairs of linked small green spheres suspended in the solution between the rectangles. In the second beaker, labeled “K C l Strong Conductivity,” six individual green spheres, three labeled plus and three labeled minus are suspended in the solution. Each of the six spheres has an arrow extending from it pointing to the rectangle labeled with the opposite sign. In the third beaker, labeled “Acetic acid solution Weak conductivity,” two pairs of joined green spheres and two individual spheres, one labeled plus and one labeled minus are shown suspended between the two rectangles. The plus labeled sphere has an arrow pointing to the rectangle labeled minus and the minus labeled sphere has an arrow pointing to the rectangle labeled plus.

Ionic Electrolytes

Water and other polar molecules are attracted to ions, as shown in Figure 7.6.2. The electrostatic attraction between an ion and a molecule with a dipole is called an ion-dipole attraction. These attractions play an important role in the dissolution of ionic compounds in water.




Figure 7.6.2: As potassium chloride (KCl) dissolves in water, the ions are hydrated. The polar water molecules are attracted by the charges on the K^+ and Cl^- ions. Water molecules in front of and behind the ions are not shown. The diagram shows eight purple spheres labeled K superscript plus and eight green spheres labeled C l superscript minus mixed and touching near the center of the diagram. Outside of this cluster of spheres are seventeen clusters of three spheres, which include one red and two white spheres. A red sphere in one of these clusters is labeled O. A white sphere is labeled H. Two of the green C l superscript minus spheres are surrounded by three of the red and white clusters, with the red spheres closer to the green spheres than the white spheres. One of the K superscript plus purple spheres is surrounded by four of the red and white clusters. The white spheres of these clusters are closest to the purple spheres.

When ionic compounds dissolve in water, the ions in the solid separate and disperse uniformly throughout the solution because water molecules surround and solvate the ions, reducing the strong electrostatic forces between them. This process represents a physical change known as dissociation. Under most conditions, ionic compounds will dissociate nearly completely when dissolved, and so they are classified as strong electrolytes.

Let us consider what happens at the microscopic level when we add solid KCl to water. Ion-dipole forces attract the positive (hydrogen) end of the polar water molecules to the negative chloride ions at the surface of the solid, and they attract the negative (oxygen) ends to the positive potassium ions. The water molecules penetrate between individual K^+ and Cl^- ions and surround them, reducing the strong interionic forces that bind the ions together and letting them move off into solution as solvated ions, as Figure 7.6.2 shows. The reduction of the electrostatic attraction permits the independent motion of each hydrated ion in a dilute solution, resulting in an increase in the disorder of the system, as the ions change from their fixed and ordered positions in the crystal to mobile and much more disordered states in solution. This increased disorder is responsible for the dissolution of many ionic compounds, including KCl, which dissolve with absorption of heat.

In other cases, the electrostatic attractions between the ions in a crystal are so large, or the ion-dipole attractive forces between the ions and water molecules are so weak, that the increase in disorder cannot compensate for the energy required to separate the ions, and the crystal is insoluble. Such is the case for compounds such as calcium carbonate (limestone), calcium phosphate (the inorganic component of bone), and iron oxide (rust).

Solubility Rules

Some combinations of aqueous reactants result in the formation of a solid precipitate as a product. However, some combinations will not produce such a product. If solutions of sodium nitrate and ammonium chloride are mixed, no reaction occurs. One could write a molecular equation showing a double-replacement reaction, but both products, sodium chloride and ammonium nitrate, are soluble and would remain in the solution as ions. Every ion is a spectator ion and there is no net ionic equation at all. It is useful to be able to predict when a precipitate will occur in a reaction. To do so, you can use a set of guidelines called the **solubility rules** (Tables 7.6.1 and 7.6.2).

Table 7.6.1: Solubility Rules for Soluble Substances

Soluble in Water	Important Exceptions (Insoluble)
All Group IA and NH_4^+ salts	none
All nitrates, chlorates, perchlorates and acetates	none
All sulfates	$CaSO_4$, $BaSO_4$, $SrSO_4$, $PbSO_4$
All chlorides, bromides, and iodides	AgX, Hg_2X_2, PbX_2 (X= Cl, Br, or I)

Table 7.6.2: Solubility Rules for Sparingly Soluble Substances

Sparingly Soluble in Water	Important Exceptions (Soluble)
All carbonates and phosphates	Group IA and NH_4^+ salts
All hydroxides	Group IA and NH_4^+ salts; Ba^{2+}, Sr^{2+}, Ca^{2+} sparingly soluble
All sulfides	Group IA, IIA and NH_4^+ salts; MgS, CaS, BaS sparingly soluble
All oxalates	Group IA and NH_4^+ salts
Special note: The following electrolytes are of only moderate solubility in water: CH_3COOAg, Ag_2SO_4, $KClO_4$. They will precipitate only if rather concentrated solutions are used.	

As an example on how to use the solubility rules, predict if a precipitate will form when solutions of cesium bromide and lead (II) nitrate are mixed.

$$Cs^+ (aq) + Br^- (aq) + Pb^{2+} (aq) + 2NO_3^- (aq) \rightarrow ?$$

The potential precipitates from a double-replacement reaction are cesium nitrate and lead (II) bromide. According to the solubility rules table, cesium nitrate is soluble because all compounds containing the nitrate ion, as well as all compounds containing the alkali metal ions, are soluble. Most compounds containing the bromide ion are soluble, but lead (II) is an exception. Therefore, the cesium and nitrate ions are spectator ions and the lead (II) bromide is a precipitate. The balanced net ionic reaction is:

$$Pb^{2+} (aq) + 2Br^- (aq) \rightarrow PbBr_2 (s)$$

Example 7.6.1: Solubility

Classify each compound as soluble or insoluble

a. $Zn(NO_3)_2$
b. $PbBr_2$
c. $Sr_3(PO_4)_2$

Solution

a. All nitrates are soluble in water, so $Zn(NO_3)_2$ is soluble.
b. All bromides are soluble in water, except those combined with Pb^{2+}, so $PbBr_2$ is insoluble.
c. All phosphates are insoluble, so $Sr_3(PO_4)_2$ is insoluble.

Exercise 7.6.1: Solubility

Classify each compound as soluble or insoluble.

a. $Mg(OH)_2$
b. KBr
c. $Pb(NO_3)_2$

Answer a

insoluble

Answer b

soluble

Answer c

soluble

Summary

Substances that dissolve in water to yield ions are called electrolytes. Nonelectrolytes are substances that do not produce ions when dissolved in water. Solubility rules allow prediction of what products will be insoluble in water.

Contributions & Attributions

- Paul Flowers (University of North Carolina - Pembroke), Klaus Theopold (University of Delaware) and Richard Langley (Stephen F. Austin State University) with contributing authors. Textbook content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at http://cnx.org/contents/85abf193-2bd...a7ac8df6@9.110).
-

7.6: Aqueous Solutions and Solubility - Compounds Dissolved in Water is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.7: Precipitation Reactions

Learning Objectives

- To identify a precipitation reaction and predict solubility.

A precipitation reaction is a reaction that yields an insoluble product—a precipitate—when two solutions are mixed. When a colorless solution of silver nitrate is mixed with a yellow-orange solution of potassium dichromate, a reddish precipitate of silver dichromate is produced.

$$\mathrm{AgNO_3(aq) + K_2Cr_2O_7(aq) \rightarrow Ag_2Cr_2O_7(s) + KNO_3(aq)} \tag{7.7.1}$$

This unbalanced equation has the general form of an exchange reaction:

$$\mathrm{AC + BD \rightarrow \underset{insoluble}{AD} + BC} \tag{7.7.2}$$

Thus precipitation reactions are a subclass of exchange reactions that occur between ionic compounds when one of the products is insoluble. Because both components of each compound change partners, such reactions are sometimes called **double-displacement reactions**. Precipitation reactions are used to isolate metals that have been extracted from their ores, and to recover precious metals for recycling.



Video: Mixing potassium dichromate and silver nitrate together to initiate a precipitation reaction (Equation 7.7.1).

Just as important as predicting the product of a reaction is knowing when a chemical reaction will *not* occur. Simply mixing solutions of two different chemical substances does *not* guarantee that a reaction will take place. For example, if 500 mL of aqueous $NaCl$ solution is mixed with 500 mL of aqueous KBr solution, the final solution has a volume of 1.00 L and contains $\mathrm{Na^+(aq)}$, $\mathrm{Cl^-(aq)}$, $\mathrm{K^+(aq)}$, and $\mathrm{Br^-(aq)}$. As you will see in (Figure 7.7.1), none of these species reacts with any of the others. When these solutions are mixed, the only effect is to dilute each solution with the other.





Figure 7.7.1: The Effect of Mixing Aqueous KBr and NaCl Solutions. Because no net reaction occurs, the only effect is to dilute each solution with the other. (Water molecules are omitted from molecular views of the solutions for clarity.)

Predicting Precipitation Reactions

A precipitation reaction occurs when a solid precipitate forms after mixing two strong electrolyte solutions. As stated previously, if none of the species in the solution reacts then no net reaction occurred.

Predict what will happen when aqueous solutions of barium chloride and lithium sulfate are mixed.

Change the partners of the anions and cations on the reactant side to form new compounds (products):



Chemical equation of the reactants barium chloride and lithium sulfate forming the products barium sulfate and lithium chloride.

Because barium chloride and lithium sulfate are strong electrolytes, each dissociates completely in water to give a solution that contains the constituent anions and cations. Mixing the two solutions *initially* gives an aqueous solution that contains Ba^{2+}, Cl^{-}, Li^{+}, and SO_4^{2-} ions. The only possible exchange reaction is to form $LiCl$ and $BaSO_4$.

Correct the formulas of the products based on the charges of the ions.

No need to correct the formula as both compounds already have their charges balanced.

$$BaCl_2(aq) + Li_2SO_4(aq) \rightarrow BaSO_4 + LiCl$$

Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.

$$BaCl_2(aq) + Li_2SO_4(aq) \rightarrow BaSO_4(s) + LiCl(aq)$$

Table 7.5.1 from the previous section shows that $LiCl$ is soluble in water, but $BaSO_4$ is not soluble in water.

Balance the equation:

$$BaCl_2(aq) + Li_2SO_4(aq) \rightarrow BaSO_4(s) + 2\,LiCl(aq)$$

Although soluble barium salts are toxic, $BaSO_4$ is so insoluble that it can be used to diagnose stomach and intestinal problems without being absorbed into tissues. An outline of the digestive organs appears on x-rays of patients who have been given a "barium milkshake" or a "barium enema"—a suspension of very fine $BaSO_4$ particles in water.




Figure 7.7.2: An x-ray of the digestive organs of a patient who has swallowed a “barium milkshake.” A barium milkshake is a suspension of very fine $BaSO_4$ particles in water; the high atomic mass of barium makes it opaque to x-rays. (Public Domain; Glitzy queen00 via Wikipedia).

Example 7.7.1

Predict what will happen if aqueous solutions of rubidium hydroxide and cobalt(II) chloride are mixed.

Solution

Solutions to Example 7.6.1

Steps	Example
Change the partners of the anions and cations on the reactant side to form new compounds (products).	 Chemical equation of the reactants rubidium hydroxide and cobalt(II) chloride forming the products rubidium chloride and cobalt hydroxide.
Correct the formulas of the products based on the charges of the ions.	$\mathrm{RbOH(aq) + CoCl_2(aq) \rightarrow RbCl + Co(OH)_2}$
Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.	$\mathrm{RbOH(aq) + CoCl_2(aq) \rightarrow RbCl(aq) + Co(OH)_2(s)}$
Balance the equation.	Coefficients already balanced. $\mathrm{RbOH(aq) + CoCl_2(aq) \rightarrow RbCl(aq) + Co(OH)_2(s)}$


Example 7.7.2

Predict what will happen if aqueous solutions of strontium bromide and aluminum nitrate are mixed.

Solution

Solutions for Example 7.6.2

Steps	Example

Steps	Example
Change the partners of the anions and cations on the reactant side to form new compounds (products).	 Chemical equation of the reactants strontium bromide and aluminum nitrate forming the products strontium nitrate and aluminum bromide.
Correct the formulas of the products based on the charges of the ions.	$\mathrm{SrBr_2(aq) + Al(NO_3)_3(aq) \rightarrow Sr(NO_3)_2 + AlBr_3}$
Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.	$\mathrm{SrBr_2(aq) + Al(NO_3)_3(aq) \rightarrow Sr(NO_3)_2(aq) + AlBr_3(aq)}$ According to Table 7.5.1 from the previous section, both $\mathrm{AlBr_3}$ (rule 4) and $\mathrm{Sr(NO_3)_2}$ (rule 2) are soluble.
If all possible products are soluble, then no net reaction will occur.	$\mathrm{SrBr_2(aq) + Al(NO_3)_3(aq) \rightarrow}$ **NO REACTION**

Exercise 7.7.2

Using the information in Table 7.5.1 from the previous section, predict what will happen in each case involving strong electrolytes.

a. An aqueous solution of strontium hydroxide is added to an aqueous solution of iron(II) chloride.
b. Solid potassium phosphate is added to an aqueous solution of mercury(II) perchlorate.
c. Solid sodium fluoride is added to an aqueous solution of ammonium formate.
d. Aqueous solutions of calcium bromide and cesium carbonate are mixed.

Answer a

$Fe(OH)_2$ precipitate is formed.

Answer b

$Hg_3(PO_4)_2$ precipitate is formed.

Answer c

No Reaction.

Answer d

$CaCO_3$ is precipitate formed.

Summary

In a **precipitation reaction**, a subclass of exchange reactions, an insoluble material (a **precipitate**) forms when two electrolyte solutions are mixed. To predict the product of a precipitation reaction, all species initially present in the solutions are identified, as are any combinations likely to produce an insoluble salt.

Contributions & Attributions

- Modified by Joshua Halpern (Howard University)

7.7: Precipitation Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.6: Precipitation Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

A typical precipitation reaction occurs when an aqueous solution of barium chloride is mixed with one containing sodium sulfate. The **complete chemical equation** can be written to describe what happens, and such an equation is useful in making chemical calculations.

$$\underbrace{\mathrm{BaCl_2(aq) + Na_2SO_4(aq) \longrightarrow BaSO_4(s) + 2\,NaCl(aq)}}_{\text{Complete Chemical Equation}} \tag{7.8.1}$$

However, Equation 7.8.1 does not really represent the microscopic particles (that is, the ions) present in the solution. Below is the complete ionic equation:

$$\underbrace{\mathrm{Ba^{2+}(aq) + \overbrace{2\,Cl^-(aq)}^{\text{spectator}} + \overbrace{2\,Na^+(aq)}^{\text{spectator}} + SO_4^{2-}(aq) \longrightarrow BaSO_4(s) + \overbrace{2\,Na^+(aq)}^{\text{spectator}} + \overbrace{Cl^-(aq)}^{\text{spectator}}}}_{\text{Complete Ionic Equation}} \tag{7.8.2}$$

Equation 7.8.2 is rather cumbersome and includes so many different ions that it may be confusing. In any case, we are often interested in the independent behavior of ions, not the specific compound from which they came. A precipitate of $\mathrm{BaSO_4(s)}$ will form when *any* solution containing $\mathrm{Ba^{2+}(aq)}$ is mixed with *any* solution containing $\mathrm{SO_4^{2-}(aq)}$ (provided concentrations are not extremely small). This happens independently of the $\mathrm{Cl^-(aq)}$ and $\mathrm{Na^+(aq)}$ ions in Equation 7.8.2. These ions are called **spectator ions** because they do not participate in the reaction. When we want to emphasize the independent behavior of ions, a **net ionic equation** is written, omitting the spectator ions. For precipitation of $\mathrm{BaSO_4}$ the net ionic equation is

$$\underbrace{\mathrm{Ba^{2+}(aq) + SO_4^{2-}(aq) \longrightarrow BaSO_4(s)}}_{\text{Net Ionic Equation}} \tag{7.8.3}$$

Example 7.8.1

a. When a solution of $\mathrm{AgNO_3}$ is added to a solution of $\mathrm{CaCl_2}$, insoluble AgCl precipitates. Write three equations (complete chemical equation, complete ionic equation, and net ionic equation) that describe this process.

b. Write the balanced net ionic equation to describe any reaction that occurs when the solutions of $\mathrm{Na_2SO_4}$ and $\mathrm{NH_4I}$ are mixed.

Solution

Solutions to Example 7.7.1

Equation Type	Example 7.8.1*a*	Example 7.8.1*b*
Complete Chemical Equation	$\mathrm{2\,AgNO_3(aq) + CaCl_2(aq) \longrightarrow 2\,AgCl(s) + Ca(NO_3)_2(aq)}$ The proper states and formulas of all products are written and the chemical equation is balanced.	$\mathrm{Na_2SO_4(aq) + 2\,NH_4I(aq) \longrightarrow 2\,NaI(aq) + (NH_4)_2SO_4(aq)}$ Both products are aqueous so there is no net ionic equation that can be written.
Complete Ionic Equation	$\mathrm{2\,Ag^+(aq) + 2\,NO_3^-(aq) + Ca^{2+}(aq) + 2\,Cl^-(aq) \longrightarrow 2\,AgCl(s) + Ca^{2+}(aq) + 2\,NO_3^-(aq)}$ AgCl is a solid so it does not break up into ions in solution.	
Net Ionic Equation	$\mathrm{Ag^+(aq) + Cl^-(aq) \longrightarrow AgCl(s)}$ All spectator ions are removed. The chemical equation is written using the lowest common coefficients.	NaI and $\mathrm{(NH_4)_2SO_4}$ are both soluble. No net ionic equation. There is no reaction.

The occurrence or nonoccurrence of precipitates can be used to detect the presence or absence of various species in solution. A $BaCl_2$ solution, for instance, is often used as a test for the presence of $SO_4^{2-}(aq)$ ions. There are several insoluble salts of Ba, but they all dissolve in dilute acid except for $BaSO_4$. Thus, if $BaCl_2$ solution is added to an unknown solution which has previously been acidified, the occurrence of a white precipitate is proof of the presence of the SO_4^{2-} ion.



Figure 7.8.1: The three common silver halide precipitates: AgI, $AgBr$ and $AgCl$ (left to right). The silver halides precipitate out of solution, but often form suspensions before settling. (CC BY-SA 3.0; Cychr).

$AgNO_3$ solutions are often used in a similar way to test for halide ions. If $AgNO_3$ solution is added to an acidified unknown solution, a white precipitate indicates the presence of Cl^- ions, a cream-colored precipitate indicates the presence of Br^- ions, and a yellow precipitate indicates the presence of I^- ions (Figure 7.8.1). Further tests can then be made to see whether perhaps a mixture of these ions is present. When $AgNO_3$ is added to tap water, a white precipitate is almost always formed. The Cl^- ions in tap water usually come from the Cl_2 which is added to municipal water supplies to kill microorganisms.

? Exercise 7.8.1

Write balanced net ionic equations to describe any reaction that occurs when the following solutions are mixed.

a. $K_2CO_3 + SrCl_2$
b. $FeSO_4 + Ba(NO_3)_2$

Answer a

$$Sr^{2+}(aq) + CO_3^{2-}(aq) \longrightarrow SrCO_3(s)$$

Answer b

$$Ba^{2+}(aq) + SO_4^{2-}(aq) \longrightarrow Ba(SO_4)(s)$$

Precipitates are also used for quantitative analysis of solutions, that is, to determine the amount of solute or the mass of solute in a given solution. For this purpose it is often convenient to use the first of the three types of equations described above. Then the rules of stoichiometry may be applied.

Contributions & Attributions

- Ed Vitz (Kutztown University), John W. Moore (UW-Madison), Justin Shorb (Hope College), Xavier Prat-Resina (University of Minnesota Rochester), Tim Wendorff, and Adam Hahn.

7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.9: Acid–Base and Gas Evolution Reactions

Learning Objectives

- Identify when a reaction will evolve a gas.

Neutralization Reactions

Acids and bases react chemically with each other to form *salts*. A salt is a general chemical term for any ionic compound formed from an acid and a base. In reactions where the acid is a hydrogen-ion-containing compound and the base is a hydroxide-ion-containing compound, water is also a product. The general reaction is as follows:

$$\text{acid} + \text{base} \rightarrow \text{water} + \text{salt}$$

The reaction of acid and base to make water and a salt is called **neutralization**. Like any chemical equation, a neutralization chemical equation must be properly balanced. For example, the neutralization reaction between sodium hydroxide and hydrochloric acid is as follows:

$$\ce{NaOH(aq) + HCl(aq) \rightarrow NaCl(aq) + H_2O(\ell)} \tag{7.9.1}$$

with coefficients all understood to be one. The neutralization reaction between sodium hydroxide and sulfuric acid is as follows:

$$\ce{2\,NaOH(aq) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + 2\,H_2O(\ell)} \tag{7.9.2}$$

Example 7.9.1: Neutralizing Nitric Acid

Nitric acid ($HNO_3(aq)$) can be neutralized by calcium hydroxide ($Ca(OH)_2(aq)$). Write a balanced chemical equation for the reaction between these two compounds and identify the salt that it produces.

Solution

Solutions to Example 7.8.1

Steps	Explanation	Equation
Write the unbalanced equation.	This is a double displacement reaction, so the cations and anions swap to create new products.	$Ca(OH)_2(aq) + HNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + H_2O(\ell)$
Balance the equation.	Because there are two OH^- ions in the formula for $Ca(OH)_2$, we need two moles of HNO_3 to provide H^+ ions	$Ca(OH)_2(aq) + \underline{2}HNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + \underline{2}H_2O(\ell)$
Additional step: identify the salt.		The salt formed is calcium nitrate.

Exercise 7.9.1

Hydrocyanic acid ($HCN(aq)$) can be neutralized by potassium hydroxide ($KOH(aq)$). Write a balanced chemical equation for the reaction between these two compounds and identify the salt that it produces.

Answer

\[\ce{KOH (aq) + HCN(aq) → KCN (aq) + H2O(ℓ)} \nonumber \]

Gas Evolving Reactions

A gas evolution reaction is a chemical process that produces a gas, such as oxygen or carbon dioxide. In the following examples, an acid reacts with a carbonate, producing salt, carbon dioxide, and water, respectively. For example, nitric acid reacts with sodium carbonate to form sodium nitrate, carbon dioxide, and water (Table 7.9.1):

$$\ce{2\,HNO_3(aq) + Na_2CO_3(aq) \rightarrow 2\,NaNO_3(aq) + CO_2(g) + H_2O(l)}$$

Sulfuric acid reacts with calcium carbonate to form calcium sulfate, carbon dioxide, and water:

$$\ce{H_2SO_4(aq) + CaCO_3(aq) \rightarrow CaSO_4(aq) + CO_2(g) + H_2O(l)}$$

Hydrochloric acid reacts with calcium carbonate to form calcium chloride, carbon dioxide, and water:

$$\ce{2\,HCl(aq) + CaCO_3(aq) \rightarrow CaCl_2(aq) + CO_2(g) + H_2O(l)}$$

Figure 7.9.1 demonstrates this type of reaction:





Figure 7.9.1: Reaction of acids with carbonates. In this reaction setup, lime water (water + calcium hydroxide) is poured into one of the test tubes and sealed with a stopper. A small amount of hydrochloric acid is carefully poured into the remaining test tube. A small amount of sodium carbonate is added to the acid, and the tube is sealed with a rubber stopper. The two tubes are connected. As a result of the acid-carbonate reaction, carbon dioxide is produced and the lime water turns milky.

In this reaction setup, lime water, a dilute calcium hydroxide ($Ca(OH)_2$) solution, is poured into one of the test tubes and sealed with a stopper. A small amount of hydrochloric acid is carefully poured into the remaining test tube. A small amount of sodium carbonate is added to the acid, and the tube is sealed with a rubber stopper. The two tubes are connected. As a result of the acid-carbonate reaction, carbon dioxide is produced and the lime water turns milky.

Table 7.9.1: Types of Compounds That Undergo Gas-Evolution Reactions

Reactant Type	Intermediate Product	Gas Evolved	Example
sulfide	none	H_2S	$2\,HCl(aq) + K_2S \rightarrow H_2S(g) + 2\,KCl(a$
carbonates and bicarbonates	H_2CO_3	CO_2	$2\,HCl(aq) + K_2CO_2 \rightarrow H_2O(l) + CO_2$
sulfites and bisulfites	H_2SO_3	SO_2	$2\,HCl(aq) + K_2SO_2 \rightarrow H_2O(l) + SO_2($
ammonia	NH_4OH	NH_3	\(\ce{NH4Cl(aq) + KOH \rightarrow H2O (l) + NH3(g) + 2KCl (aq)}\)

The gas-evolving experiment lime water is illustrated in the following video:



***Video* 7.9.1:** *Carbon Dioxide (CO_2) & Limewater (Chemical Reaction). As the reaction proceeds, the limewater on the turns from clear to milky; this is due to the $CO_2(g)$ reacting with the aqueous calcium hydroxide to form calcium carbonate, which is only slightly soluble in water.*

When this experiment is repeated with nitric or sulfuric acid instead of HCl, it yields the same results: the clear limewater turns milky, indicating the production of carbon dioxide. Another method to chemically generate gas is the oxidation of metals in acidic solutions. This reaction will yield a metal salt and hydrogen gas.

$$2\,HCl(aq) + Zn(s) \rightarrow ZnCl_2(aq) + H_2(g)$$

Here, hydrochloric acid oxidizes zinc to produce an aqueous metal salt and hydrogen gas bubbles. Recall that oxidation refers to a loss of electrons, and reduction refers to the gain of electrons. In the above redox reaction, neutral zinc is oxidized to Zn^{2+}, and the acid, H^+, is reduced to $H_2(g)$. The oxidation of metals by strong acids is another common example of a gas evolution reaction.

Contributors & Affiliations

- Boundless (www.boundless.com)
- Wikipedia (CC-BY-SA-3.0)
- Paul Flowers (University of North Carolina - Pembroke), Klaus Theopold (University of Delaware) and Richard Langley (Stephen F. Austin State University) with contributing authors. Textbook content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at http://cnx.org/contents/85abf193-2bd...a7ac8df6@9.110).

7.9: Acid–Base and Gas Evolution Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.8: Acid–Base and Gas Evolution Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.10: Oxidation-Reduction Reactions

Learning Objectives

- Define *oxidation* and *reduction*.
- Assign oxidation numbers to atoms in simple compounds.
- Recognize a reaction as an oxidation-reduction reaction.

Consider this chemical reaction:

$$\mathrm{Mg(s) + Cl_2(g) \rightarrow MgCl_2}$$

The reactants are two electrically neutral elements; they have the same number of electrons as protons. The product, however, is ionic; it is composed of $\mathrm{Mg^{2+}}$ and $\mathrm{Cl^-}$ ions. Somehow, the individual Mg atoms lose two electrons to make the $\mathrm{Mg^{2+}}$ ion, while the Cl atoms gain an electron to become $\mathrm{Cl^-}$ ions. This reaction involves the *transfer of electrons* between atoms.

The process of losing and gaining electrons occurs simultaneously. However, mentally we can separate the two processes. **Oxidation** is defined as the loss of one or more electrons by an atom. **Reduction** is defined as the gain of one or more electrons by an atom. So oxidation and reduction always occur together; it is only mentally that we can separate them. Chemical reactions that involve the transfer of electrons are called **oxidation-reduction (or redox) reactions**.

Redox reactions require that we keep track of the electrons assigned to each atom in a chemical reaction. How do we do that? We use **oxidation numbers** to keep track of electrons in atoms. Oxidation numbers are assigned to atoms based on four rules. Oxidation numbers are not necessarily equal to the charge on the atom (although sometimes they can be); we must keep the concepts of charge and oxidation numbers separate.

Assigning Oxidation Numbers

The rules for assigning oxidation numbers to atoms are as follows:

1. Atoms in their elemental state are assigned an oxidation number of 0.
2. Atoms in monatomic (i.e., one-atom) ions are assigned an oxidation number equal to their charge. Oxidation numbers are usually written with the sign first, then the magnitude, to differentiate them from charges.
3. In compounds, fluorine is assigned a −1 oxidation number; oxygen is usually assigned a −2 oxidation number (except in peroxide compounds [where it is −1] and in binary compounds with fluorine [where it is positive]); and hydrogen is usually assigned a +1 oxidation number (except when it exists as the hydride ion [$\mathrm{H^-}$], in which case rule 2 prevails).
4. In compounds, all other atoms are assigned an oxidation number so that the sum of the oxidation numbers on all the atoms in the species equals the charge on the species (which is zero if the species is neutral).

Here are some examples for practice. In $\mathrm{H_2}$, both H atoms have an oxidation number of 0 by rule 1. In $\mathrm{MgCl_2}$, magnesium has an oxidation number of +2, while chlorine has an oxidation number of −1 by rule 2. In $\mathrm{H_2O}$, the H atoms each have an oxidation number of +1, while the O atom has an oxidation number of −2, even though hydrogen and oxygen do not exist as ions in this compound (rule 3). By contrast, by rule 3, each H atom in hydrogen peroxide ($\mathrm{H_2O_2}$) has an oxidation number of +1, while each O atom has an oxidation number of −1. We can use rule 4 to determine oxidation numbers for the atoms in $\mathrm{SO_2}$. Each O atom has an oxidation number of −2; for the sum of the oxidation numbers to equal the charge on the species (which is zero), the S atom is assigned an oxidation number of +4. Does this mean that the sulfur atom has a 4+ charge on it? No, it means only that the S atom is assigned a +4 oxidation number by our rules of apportioning electrons among the atoms in a compound.

Example 7.10.1

Assign oxidation numbers to the atoms in each substance.

a. $\mathrm{Cl_2}$
b. $\mathrm{GeO_2}$
c. $\mathrm{Ca(NO_3)_2}$

Solution

1. Cl_2 is the elemental form of chlorine. Rule 1 states each atom has an oxidation number of 0.
2. By rule 3, oxygen is normally assigned an oxidation number of −2. For the sum of the oxidation numbers to equal the charge on the species (zero), the Ge atom is assigned an oxidation number of +4.
3. $Ca(NO_3)_2$ can be separated into two parts: the Ca^{2+} ion and the NO_3^- ion. Considering these separately, the Ca^{2+} ion has an oxidation number of +2 by rule 2. Now consider the NO_3^- ion. Oxygen is assigned an oxidation number of −2, and there are three of them. According to rule 4, the sum of the oxidation numbers on all atoms must equal the charge on the species, so we have the simple algebraic equation x + 3(−2) = −1 where x is the oxidation number of the N atom and the −1 represents the charge on the species. Evaluating for x,

$$\begin{aligned} x + (-6) &= -1 \\ x &= +5 \end{aligned}$$

Thus the oxidation number on the N atom in the NO_3^- ion is +5.

Exercise 7.10.1: Phosphoric Acid

Assign oxidation numbers to the atoms in H3PO4.

Answer

H: +1; O: −2; P: +5

All redox reactions occur with a simultaneous change in the oxidation numbers of some atoms. At least two elements must change their oxidation numbers. When an oxidation number of an atom is increased in the course of a redox reaction, that atom is being *oxidized*. When an oxidation number of an atom is decreased in the course of a redox reaction, that atom is being *reduced*. Oxidation and reduction can also be defined in terms of increasing or decreasing oxidation numbers, respectively.

Example 7.10.2

Identify what is being oxidized and reduced in this redox reaction.


$$2\,Na + Br_2 \rightarrow 2\,NaBr$$

Solution

Both reactants are the elemental forms of their atoms, so the Na and Br atoms have oxidation numbers of 0. In the ionic product, the Na^+ ions have an oxidation number of +1, while the Br^- ions have an oxidation number of −1.

$$\overset{0}{2\,Na} + \overset{0}{Br_2} \rightarrow \overset{+1\ -1}{2\,NaBr}$$

Sodium is increasing its oxidation number from 0 to +1, so it is being oxidized; bromine is decreasing its oxidation number from 0 to −1, so it is being reduced:



Because oxidation numbers are changing, this is a redox reaction. The total number of electrons being lost by sodium (two, one lost from each Na atom) is gained by bromine (two, one gained for each Br atom).

Exercise 7.10.2

Identify what is being oxidized and reduced in this redox reaction.

$$C + O_2 \rightarrow CO_2$$

Answer

C is being oxidized from 0 to +4.

O is being reduced from 0 to −2.

Oxidation reactions can become quite complex, as attested by the following redox reaction:

$$6\,H^+(aq) + 2\,\underset{+7}{MnO_4^-}(aq) + 5\,\underset{-1}{H_2O_2}(l) \longrightarrow 2\,\underset{+2}{Mn^{2+}}(aq) + 5\,\underset{0}{O_2}(g) + 8\,H_2O(l)$$

To demonstrate that this is a redox reaction, the oxidation numbers of the species being oxidized and reduced are listed; can you determine what is being oxidized and what is being reduced? This is also an example of a net ionic reaction; spectator ions that do not change oxidation numbers are not displayed in the equation. Eventually, we will need to learn techniques for writing correct (i.e., balanced) redox reactions.

Food and Drink Application: Fortifying Food with Iron

Iron is an essential mineral in our diet; iron-containing compounds like the heme protein in hemoglobin could not function without it. Most biological iron has the form of the Fe^{2+} ion; iron with other oxidation numbers is almost inconsequential in human biology (although the body does contain an enzyme to reduce Fe^{3+} to Fe^{2+}, so Fe^{3+} must have some biological significance, albeit minor). To ensure that we ingest enough iron, many foods are enriched with iron. Although Fe^{2+} compounds are the most logical substances to use, some foods use "reduced iron" as an ingredient (bread and breakfast cereals are the most well-known examples). Reduced iron is simply iron metal; iron is added as a fine metallic powder. The metallic iron is oxidized to Fe^{2+} in the digestive system and then absorbed by the body, but the question remains: Why are we ingesting metallic iron? Why not just use Fe^{2+} salts as an additive?



Figure 7.10.1 Ingredients. Many prepared foods list reduced iron in their ingredients list.

Although it is difficult to establish conclusive reasons, a search of scientific and medical literature suggests a few reasons. One reason is that fine iron filings do not affect the taste of the product. The size of the iron powder (several dozen micrometers) is not noticeable when chewing iron-supplemented foods, and the tongue does not detect any changes in flavor that can be detected when using Fe^{2+} salts. Fe^{2+} compounds can affect other properties of foodstuffs during preparation and cooking, like dough pliability, yeast growth, and color. Finally, of the common iron substances that might be used, metallic iron is the least expensive. These factors appear to be among the reasons why metallic iron is the supplement of choice in some foods.

Key Takeaways

- Oxidation-reduction (redox) reactions involve the transfer of electrons from one atom to another.
- Oxidation numbers are used to keep track of electrons in atoms.
- There are rules for assigning oxidation numbers to atoms.
- Oxidation is an increase in oxidation number (loss of electrons); reduction is a decrease in oxidation number (gain of electrons).

7.10: Oxidation-Reduction Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **14.2: Oxidation-Reduction Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/beginning-chemistry.

7.11: Classifying Chemical Reactions

Learning Objectives

- Classify a chemical reaction as a synthesis, decomposition, single replacement, double replacement, or a combustion reaction.
- Predict the products of simple reactions.

The chemical reactions we have described are only a tiny sampling of the infinite number of chemical reactions possible. How do chemists cope with this overwhelming diversity? How do they predict which compounds will react with one another and what products will be formed? The key to success is to find useful ways to categorize reactions. Familiarity with a few basic types of reactions will help you to predict the products that form when certain kinds of compounds or elements come in contact.

Most chemical reactions can be classified into one or more of five basic types: acid–base reactions, exchange reactions, condensation reactions (and the reverse, cleavage reactions), and oxidation–reduction reactions. The general forms of these five kinds of reactions are summarized in Table 7.11.1, along with examples of each. It is important to note, however, that many reactions can be assigned to more than one classification, as you will see in our discussion.

Table 7.11.1: *Basic Types of Chemical Reactions*

Name of Reaction	General Form	Examples
Oxidation–Reduction (redox)	oxidant + reductant → reduced oxidant + oxidized reductant	$C_7H_{16}(l) + 11O_2(g) \rightarrow 7CO_2(g) + 8H_2O(g)$
Acid–Base	acid + base → salt	$NaOH(aq) + HNO_3(aq) \rightarrow NaNO_3(aq) + H_2O(l)$
Exchange: Single Replacement	AB + C → AC + B	$ZnCl_2(aq) + Mg(s) \rightarrow MgCl_2(aq) + Zn(s)$
Exchange: Double Replacement	AB + CD → AD + CB	$BaCl_2(aq) + Na_2SO_4(aq) \rightarrow BaSO_4(s) + 2NaCl(aq)$
Combination (Synthesis)	A + B → AB	$CO_2(g) + H_2O(l) \rightarrow H_2CO_3(aq)$ $N_2(g) + 2O_2(g) \rightarrow 2NO_2(g)$
Decomposition	AB → A + B	$CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$

The classification scheme is only for convenience; the same reaction can be classified in different ways, depending on which of its characteristics is most important. Oxidation–reduction reactions, in which there is a net transfer of electrons from one atom to another, and condensation reactions are discussed in this section. Acid–base reactions are one kind of exchange reaction—the formation of an insoluble salt, such as barium sulfate, when solutions of two soluble salts are mixed together.

Combination Reactions

A **combination reaction** is a reaction in which two or more substances combine to form a single new substance. Combination reactions can also be called synthesis reactions. The general form of a combination reaction is:

$$\mathrm{A + B \rightarrow AB}$$

One combination reaction is two elements combining to form a compound. Solid sodium metal reacts with chlorine gas to produce solid sodium chloride.

$$\mathrm{2Na\,(\mathit{s}) + Cl_2\,(\mathit{g}) \rightarrow 2NaCl\,(\mathit{s})}$$

Notice that in order to write and balance the equation correctly, it is important to remember the seven elements that exist in nature as diatomic molecules (H_2, N_2, O_2, F_2, Cl_2, Br_2, and I_2).

One type of combination reaction that occurs frequently is the reaction of an element with oxygen to form an oxide. Metals and nonmetals both react readily with oxygen under most conditions. Magnesium reacts rapidly and dramatically when ignited, combining with oxygen from the air to produce a fine powder of magnesium oxide:

$$2\mathrm{Mg}\,(s) + \mathrm{O_2}\,(g) \rightarrow 2\mathrm{MgO}\,(s)$$

Sulfur reacts with oxygen to form sulfur dioxide:

$$\mathrm{S}\,(s) + \mathrm{O_2}\,(g) \rightarrow \mathrm{SO_2}\,(g)$$

When nonmetals react with one another, the product is a molecular compound. Often, the nonmetal reactants can combine in different ratios and produce different products. Sulfur can also combine with oxygen to form sulfur trioxide:

$$2\mathrm{S}\,(s) + 3\mathrm{O_2}\,(g) \rightarrow 2\mathrm{SO_3}\,(g)$$

Transition metals are capable of adopting multiple positive charges within their ionic compounds. Therefore, most transition metals are capable of forming different products in a combination reaction. Iron reacts with oxygen to form both iron (II) oxide and iron (III) oxide:

$$2\mathrm{Fe}\,(s) + \mathrm{O_2}\,(g) \rightarrow 2\mathrm{FeO}\,(s)$$

$$4\mathrm{Fe}\,(s) + 3\mathrm{O_2}\,(g) \rightarrow 2\mathrm{Fe_2O_3}\,(s)$$

Example 7.11.1: Combustion of Solid Potassium

Potassium is a very reactive alkali metal that must be stored under oil in order to prevent it from reacting with air. Write the balanced chemical equation for the combination reaction of potassium with oxygen.

Solution

Solutions to Example 7.10.1

Steps	Example Solution
Plan the problem.	Make sure formulas of all reactants and products are correct before balancing the equation. Oxygen gas is a diatomic molecule. Potassium oxide is an ionic compound and so its formula is constructed by the crisscross method. Potassium as an ion becomes $\mathrm{K^+}$, while the oxide ion is $\mathrm{O^{2-}}$.
Solve.	The skeleton (unbalanced) equation: $$\mathrm{K}\,(s) + \mathrm{O_2}\,(g) \rightarrow \mathrm{K_2O}\,(s)$$ The equation is then easily balanced with coefficients. $$4\mathrm{K}\,(s) + \mathrm{O_2}\,(g) \rightarrow 2\mathrm{K_2O}\,(s)$$
Think about your result.	Formulas are correct and the resulting combination reaction is balanced.

Combination reactions can also take place when an element reacts with a compound to form a new compound composed of a larger number of atoms. Carbon monoxide reacts with oxygen to form carbon dioxide according to the equation:

$$2\mathrm{CO}\,(g) + \mathrm{O_2}\,(g) \rightarrow 2\mathrm{CO_2}\,(g)$$

Two compounds may also react to form a more complex compound. A very common example is the reactions of oxides with water. Calcium oxide reacts readily with water to produce an aqueous solution of calcium hydroxide:

$$\mathrm{CaO}\,(s) + \mathrm{H_2O}\,(l) \rightarrow \mathrm{Ca(OH)_2}\,(aq)$$

Sulfur trioxide gas reacts with water to form sulfuric acid. This is an unfortunately common reaction that occurs in the atmosphere in some places where oxides of sulfur are present as pollutants. The acid formed in the reaction falls to the ground as acid rain.

$$\mathrm{SO_3}\,(g) + \mathrm{H_2O}\,(l) \rightarrow \mathrm{H_2SO_4}\,(aq)$$




Figure 7.11.1: Acid rain has severe consequences on both natural and manmade objects. Acid rain degrades marble statues like the one on the left (A). The trees in the forest on the right (B) have been killed by acid rain.

? Exercise 7.11.1

a. Write the chemical equation for the synthesis of silver bromide, AgBr.
b. Predict the products for the following reaction: $\mathrm{CO_2}\,(g) + \mathrm{H_2O}\,(l)$

Answer a:

$$2\mathrm{Ag} + \mathrm{Br_2} \rightarrow 2\mathrm{AgBr}$$

Answer b:

$$\mathrm{CO_2}\,(g) + \mathrm{H_2O}\,(l) \rightarrow \mathrm{H_2CO_3}$$

Decomposition Reactions

A **decomposition reaction** is a reaction in which a compound breaks down into two or more simpler substances. The general form of a decomposition reaction is:

$$\mathrm{AB} \rightarrow \mathrm{A} + \mathrm{B}$$

Most decomposition reactions require an input of energy in the form of heat, light, or electricity.

Binary compounds are compounds composed of just two elements. The simplest kind of decomposition reaction is when a binary compound decomposes into its elements. Mercury (II) oxide, a red solid, decomposes when heated to produce mercury and oxygen gas:

$$2\mathrm{HgO}\,(s) \rightarrow 2\mathrm{Hg}\,(l) + \mathrm{O_2}\,(g)$$



Video 7.11.2: *Mercury (II) oxide is a red solid. When it is heated, it decomposes into mercury metal and oxygen gas.*

A reaction is also considered to be a decomposition reaction even when one or more of the products are still compounds. A metal carbonate decomposes into a metal oxide and carbon dioxide gas. For example, calcium carbonate decomposes into calcium oxide and carbon dioxide:

$$\mathrm{CaCO_3}\,(s) \rightarrow \mathrm{CaO}\,(s) + \mathrm{CO_2}\,(g)$$

Metal hydroxides decompose on heating to yield metal oxides and water. Sodium hydroxide decomposes to produce sodium oxide and water:

$$2\text{NaOH}\,(s) \rightarrow \text{Na}_2\text{O}\,(s) + \text{H}_2\text{O}\,(g)$$

Some unstable acids decompose to produce nonmetal oxides and water. Carbonic acid decomposes easily at room temperature into carbon dioxide and water:

$$\text{H}_2\text{CO}_3\,(aq) \rightarrow \text{CO}_2\,(g) + \text{H}_2\text{O}\,(l)$$

Example 7.11.2: Electrolysis of Water

When an electric current is passed through pure water, it decomposes into its elements. Write a balanced equation for the decomposition of water.

Solution

Solutions to Example 7.10.2

Steps	Example Solution
Plan the problem.	Water is a binary compound composed of hydrogen and oxygen. The hydrogen and oxygen gases produced in the reaction are both diatomic molecules.
Solve.	The skeleton (unbalanced) equation: $$\text{H}_2\text{O}\,(l) \overset{\text{elec}}{\rightarrow} \text{H}_2\,(g) + \text{O}_2\,(g)$$ Note the abbreviation "elec" above the arrow to indicate the passage of an electric current to initiate the reaction. Balance the equation. $$2\text{H}_2\text{O}\,(l) \overset{\text{elec}}{\rightarrow} 2\text{H}_2\,(g) + \text{O}_2\,(g)$$
Think about your result.	The products are elements and the equation is balanced.

Exercise 7.11.2

Write the chemical equation for the decomposition of:

a. Al_2O_3
b. Ag_2S

Answer a

$2\text{Al}_2\text{O}_3 \rightarrow 4\text{Al} + 3\text{O}_2$

Answer b

$\text{Ag}_2\text{S} \rightarrow 2\text{Ag} + \text{S}$

Single Replacement Reactions

A third type of reaction is the single replacement reaction, in which one element replaces a similar element in a compound. The general form of a single-replacement (also called single-displacement) reaction is:

$$\text{A} + \text{BC} \rightarrow \text{AC} + \text{B}$$

In this general reaction, element A is a metal and replaces element B, also a metal, in the compound. When the element that is doing the replacing is a nonmetal, it must replace another nonmetal in a compound, and the general equation becomes:

$$\text{Y} + \text{XZ} \rightarrow \text{XY} + \text{Z}$$

where Y is a nonmetal and replaces the nonmetal Z in the compound with X.

Metal Replacement

Magnesium is a more reactive metal than copper. When a strip of magnesium metal is placed in an aqueous solution of copper (II) nitrate, it replaces the copper. The products of the reaction are aqueous magnesium nitrate and solid copper metal.

$$\mathrm{Mg}\,(s) + \mathrm{Cu(NO_3)_2}\,(aq) \rightarrow \mathrm{Mg(NO_3)_2}\,(aq) + \mathrm{Cu}\,(s)$$

This subcategory of single-replacement reactions is called a metal replacement reaction because it is a metal that is being replaced (copper).

Hydrogen Replacement

Many metals react easily with acids and when they do so, one of the products of the reaction is hydrogen gas. Zinc reacts with hydrochloric acid to produce aqueous zinc chloride and hydrogen (figure below).

$$\mathrm{Zn}\,(s) + 2\mathrm{HCl}\,(aq) \rightarrow \mathrm{ZnCl_2}\,(aq) + \mathrm{H_2}\,(g)$$

In a hydrogen replacement reaction, the hydrogen in the acid is replaced by an active metal. Some metals are so reactive that they are capable of replacing the hydrogen in water. The products of such a reaction are the metal hydroxide and hydrogen gas. All Group 1 metals undergo this type of reaction. Sodium reacts vigorously with water to produce aqueous sodium hydroxide and hydrogen (see figure below).

$$2\mathrm{Na}\,(s) + 2\mathrm{H_2O}\,(l) \rightarrow 2\mathrm{NaOH}\,(aq) + \mathrm{H_2}\,(g)$$



Figure 7.11.2: (First image) Zinc metal reacts with hydrochloric acid to give off hydrogen gas in a single-displacement reaction. (Second image) Sodium metal reacts vigorously with water, giving off hydrogen gas. A large piece of sodium will often generate so much heat that the hydrogen will ignite.

Halogen Replacement

The element chlorine reacts with an aqueous solution of sodium bromide to produce aqueous sodium chloride and elemental bromine:

$$\mathrm{Cl_2}\,(g) + 2\mathrm{NaBr}\,(aq) \rightarrow 2\mathrm{NaCl}\,(aq) + \mathrm{Br_2}\,(l)$$

The reactivity of the halogen group (group 17) decreases from top to bottom within the group. Fluorine is the most reactive halogen, while iodine is the least. Since chlorine is above bromine, it is more reactive than bromine and can replace it in a halogen replacement reaction.

Example 7.11.3

What are the products of the reaction between solid aluminum (Al) and iron (III) oxide ($\mathrm{Fe_2O_3}$)?

Solution

Solutions to Example 7.10.3

Steps	Example Solution

Steps	Example Solution
Plan the problem.	To predict the products, we need to know that aluminum will replace iron and form aluminum oxide (the metal will replace the metal ion in the compound). Aluminum has a charge of $+3$ and oxygen has a charge of -2. The compound formed between aluminum and oxygen, therefore, will be $\mathrm{Al_2O_3}$. Since iron is replaced in the compound by aluminum, the iron will now be the single element in the products.
Solve.	The unbalanced equation will be: $$\mathrm{Al + Fe_2O_3 \rightarrow Al_2O_3 + Fe}$$ and the balanced equation will be: $$\mathrm{2Al + Fe_2O_3 \rightarrow Al_2O_3 + 2Fe}$$
Think about your result.	This is a single replacement reaction, and when balanced the coefficients accurately reflect that the iron and aluminum have the same charge in this reaction.

? Exercise 7.11.3

a. Write the chemical equation for the single replacement reaction between zinc solid and lead (II) nitrate solution to produce zinc nitrate solution and solid lead. (Note that zinc forms ions with a $+2$ charge.)
b. Predict the products for the following reaction: $\mathrm{Fe + CuSO_4}$. (In this reaction, assume iron forms ions with a $+2$ charge.)

Answer a

$$\mathrm{Zn + Pb(NO_3)_2 \rightarrow Pb + Zn(NO_3)_2}$$

Answer b

$$\mathrm{Fe + CuSO_4 \rightarrow Cu + FeSO_4}$$

Double Replacement Reactions

A **double-replacement reaction** is a reaction in which the positive and negative ions of two ionic compounds exchange places to form two new compounds. The general form of a double-replacement (also called double-displacement) reaction is:

$$\mathrm{AB + CD \rightarrow AD + BC}$$

In this reaction, A and C are positively-charged cations, while B and D are negatively-charged anions. Double-replacement reactions generally occur between substances in aqueous solution. In order for a reaction to occur, one of the products is usually a solid precipitate, a gas, or a molecular compound such as water.

Formation of a Precipitate

A precipitate forms in a double-replacement reaction when the cations from one of the reactants combine with the anions from the other reactant to form an insoluble ionic compound. When aqueous solutions of potassium iodide and lead (II) nitrate are mixed, the following reaction occurs:

$$\mathrm{2KI\,(\mathit{aq}) + Pb(NO_3)_2\,(\mathit{aq}) \rightarrow 2KNO_3\,(\mathit{aq}) + PbI_2\,(\mathit{s})} \qquad (7.11.1)$$

There are very strong attractive forces that occur between $\mathrm{Pb^{2+}}$ and $\mathrm{I^-}$ ions and the result is a brilliant yellow precipitate (Figure 7.11.3). The other product of the reaction, potassium nitrate, remains soluble.



Figure 7.11.3: Lead (II) iodide precipitates when potassium iodide is mixed with lead (II) nitrate (Equation 7.11.1). (CC BY-SA 3.0; PRHaney).

Formation of a Gas

Some double-replacement reactions produce a gaseous product which then bubbles out of the solution and escapes into the air. When solutions of sodium sulfide and hydrochloric acid are mixed, the products of the reaction are aqueous sodium chloride and hydrogen sulfide gas:

$$\mathrm{Na_2S}\,(aq) + 2\mathrm{HCl}\,(aq) \rightarrow 2\mathrm{NaCl}\,(aq) + \mathrm{H_2S}\,(g)$$

Formation of a Molecular Compound

Another kind of double-replacement reaction is one that produces a molecular compound as one of its products. Many examples in this category are reactions that produce water. When aqueous hydrochloric acid is reacted with aqueous sodium hydroxide, the products are aqueous sodium chloride and water:

$$\mathrm{HCl}\,(aq) + \mathrm{NaOH}\,(aq) \rightarrow \mathrm{NaCl}\,(aq) + \mathrm{H_2O}\,(l)$$

Example 7.11.4

Write a complete and balanced chemical equation for the double-replacement reaction $\mathrm{NaCN}\,(aq) + \mathrm{HBr}\,(aq) \rightarrow$ (hydrogen cyanide gas is formed).

Solution

Solutions to Example 7.10.4

Steps	Example Solution
Plan the problem.	The production of a gas drives the reaction.
Solve.	The cations of both reactants are $+1$ charged ions, while the anions are -1 charged ions. After exchanging partners, the balanced equation is: $$\mathrm{NaCN}\,(aq) + \mathrm{HBr}\,(aq) \rightarrow \mathrm{NaBr}\,(aq) + \mathrm{HCN}\,(g)$$
Think about your result.	This is a double replacement reaction. All formulas are correct and the equation is balanced.

Exercise 7.11.4

Write a complete and balanced chemical equation for the double-replacement reaction $(\mathrm{NH_4})_2\mathrm{SO_4}\,(aq) + \mathrm{Ba(NO_3)_2}\,(aq) \rightarrow$ (a precipitate of barium sulfate forms).

Answer a:

$$(\mathrm{NH_4})_2\mathrm{SO_4}\,(aq) + \mathrm{Ba(NO_3)_2}\,(aq) \rightarrow 2\mathrm{NH_4NO_3}\,(aq) + \mathrm{BaSO_4}\,(s)$$

Occasionally, a reaction will produce both a gas and a molecular compound. The reaction of a sodium carbonate solution with hydrochloric acid produces aqueous sodium chloride, carbon dioxide gas, and water:

$$\mathrm{Na_2CO_3}\left(aq\right) + 2\mathrm{HCl}\left(aq\right) \rightarrow 2\ \square\ \mathrm{NaCl}\left(aq\right) + \mathrm{CO_2}\left(g\right) + \mathrm{H_2O}\left(l\right)$$

7.11: Classifying Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.10: Classifying Chemical Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

7.12: The Activity Series- Predicting Spontaneous Redox Reactions

Learning Objectives

- Use the activity series to predict if a reaction will occur.

We see below two metals that can be exposed to water. The picture on the left is of sodium, which has a violent reaction when it comes in contact with water. The picture on the right is of silver, a metal so unreactive with water that it can be made into drinking vessels. Both metals have a single s electron in their outer shell, so you would predict a similar reactivity from each. However, we have a tool that allows us to make better predictions about how certain elements will react with others.



Figure 7.12.1: On the left, sodium reacts with water. On the right, silver in the form of cups does not react with water.

The Activity Series

Single-replacement reactions only occur when the element that is doing the replacing is more reactive than the element that is being replaced. Therefore, it is useful to have a list of elements in order of their relative reactivity. The **activity series** is a list of elements in decreasing order of their reactivity. Since metals replace other metals, while nonmetals replace other nonmetals, they each have a separate activity series. The table 7.12.1 below is an activity series of most common metals, and the table 7.12.2 is an activity series of the halogens.

Table 7.12.1: Activity Series of Metal Elements

Elements, from most to least reactive	Reaction Occurring
Li K Ba Sr Ca Na	React with cold water, replacing hydrogen.
Mg Al Zn Cr Fe Cd	React with steam, but not cold water, replacing hydrogen.
Co Ni Sn Pb	Do not react with water. React with acids, replacing hydrogen.
H_2	
Cu Hg Ag Pt Au	Unreactive with water or acids.

Table 7.12.2: Activity Series of Nonmetal elements

Elements, from most to least reactive
F_2
Cl_2
Br_2
I_2

For a single-replacement reaction, a given element is capable of replacing an element that is below it in the activity series. This can be used to predict if a reaction will occur. Suppose that small pieces of the metal nickel were placed into two separate aqueous solutions: one of iron (III) nitrate and one of lead (II) nitrate. Looking at the activity series, we see that nickel is below iron, but above lead. Therefore, the nickel metal will be capable of replacing the lead in a reaction, but will not be capable of replacing iron.

$$\mathrm{Ni}\,(s) + \mathrm{Pb(NO_3)_2}\,(aq) \rightarrow \mathrm{Ni(NO_3)_2}\,(aq) + \mathrm{Pb}\,(s)$$

$$\mathrm{Ni}\,(s) + \mathrm{Fe(NO_3)_3}\,(aq) \rightarrow \mathrm{NR\ (no\ reaction)}$$

In the descriptions that accompany the activity series of metals, a given metal is also capable of undergoing the reactions described below that section. For example, lithium will react with cold water, replacing hydrogen. It will also react with steam and with acids, since that requires a lower degree of reactivity.

Examples 7.12.1

Use the activity series to predict if the following reactions will occur. If not, write NR. If the reaction does occur, write the products of the reaction and balance the equation.

a. $\mathrm{Al}\,(s) + \mathrm{Zn(NO_3)_2}\,(aq) \rightarrow$
b. $\mathrm{Ag}\,(s) + \mathrm{HCl}\,(aq) \rightarrow$

Solution

Solutions to Example 7.11.1

Steps	**Example 7.12.1*A*** $\mathrm{Al}\,(s) + \mathrm{Zn(NO_3)_2}\,(aq) \rightarrow$	**Example 7.12.1*B*** $\mathrm{Ag}\,(s) + \mathrm{HCl}\,(aq) \rightarrow$
Plan the problem.	Compare the placements of aluminum and zinc on the activity series (Table 7.12.1)	Compare the placements of silver and hydrogen (Table 7.12.1)
Solve.	Since aluminum is above zinc, it is capable of replacing it and a reaction will occur. The products of the reaction will be aqueous aluminum nitrate and solid zinc. Take care to write the correct formulas for the products before balancing the equation. Aluminum adopts a $+3$ charge in an ionic compound, so the formula for aluminum nitrate is $\mathrm{Al(NO_3)_3}$. The balanced equation is: $2\mathrm{Al}\,(s) + 3\mathrm{Zn(NO_3)_2}\,(aq) \rightarrow 2\mathrm{Al(NO_3)_3}\,(aq) + 3\mathrm{Zn}\,(s)$	Since silver is below hydrogen, it is not capable of replacing hydrogen in a reaction with an acid. $\mathrm{Ag}\,(s) + \mathrm{HCl}\,(aq) \rightarrow \mathrm{NR}$

Exercise 7.12.1

Use the activity series to predict the products, if any, of each equation.

a. $\mathrm{FeCl_2} + \mathrm{Zn} \rightarrow$

b. $HNO_3 + Au \rightarrow$

Answer a

The products are $ZnCl_2$ + Fe.

Answer b

No reaction.

Summary

- Metals and halogens are ranked according to their ability to displace other metals or halogens below them in the activity series.

7.12: The Activity Series- Predicting Spontaneous Redox Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.11: The Activity Series- Predicting Spontaneous Redox Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

8: Quantities in Chemical Reactions

So far, we have talked about chemical reactions in terms of individual atoms and molecules. Although this works, most of the reactions occurring around us involve much larger amounts of chemicals. Even a tiny sample of a substance will contain millions, billions, or a hundred billion billions of atoms and molecules. How do we compare amounts of substances to each other in chemical terms when it is so difficult to count to a hundred billion billion? Actually, there are ways to do this, which we will explore in this chapter. In doing so, we will increase our understanding of stoichiometry, which is the study of the numerical relationships between the reactants and the products in a balanced chemical reaction.

8.1: Prelude to Quantities in Chemical Reactions
8.2: The Mole
8.3: Atomic and Molar Masses
8.4: Mole-Mass Conversions
8.5: Mole-Mole Relationships in Chemical Reactions
8.6: Mole-Mass and Mass-Mass Problems
8.E: Quantities in Chemical Reactions (Exercise)
8.S: Quantities in Chemical Reactions (Summary)

Template:HideTOC

8: Quantities in Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

8.1: Prelude to Quantities in Chemical Reactions

When the disengaged gasses are carefully examined, they are found to weigh **113.7 grs.**; these are of two kinds, viz. **144 cubical inches** of carbonic acid gas, weighing **100 grs.** and **380 cubical inches** of a very light gas, weighing only **13.7 grs.**…and, when the water which has passed over into the bottle [labeled] H is carefully examined, it is found to have lost **85.7 grs.** of its weight. Thus, in this experiment, **85.7 grs.** of water, joined to **28 grs.** of charcoal, have combined in such a way as to form **100 grs.** of carbonic acid, and **13.7 grs.** of a particular gas capable of being burnt. (Bold emphasis added.)

In this paragraph from the *Elements of Chemistry*, Antoine Lavoisier (1743–94) is explaining an experiment in which he was trying to demonstrate that water is not an element but instead is composed of hydrogen (the gas "capable of being burnt") and oxygen. This is a historical account of a groundbreaking experiment and illustrates the importance of amounts in chemistry. Lavoisier was pointing out that the initial total mass of water and charcoal, 85.7 g plus 28 g, equals the final total mass of carbonic acid and the particular gas, 100 g plus 13.7 g. In this way, he was illustrating the law of conservation of matter. It is another way of saying that *amounts matter*.

Amounts do matter and in a variety of circumstances. The chapter-opening essay in Chapter 1 tells the story of a nurse who mistakenly read "2–3 mg" as "23 mg" and administered the higher and potentially fatal dose of morphine to a child. Food scientists who work in test kitchens must keep track of specific amounts of ingredients as they develop new products for us to eat. Quality control technicians measure amounts of substances in manufactured products to ensure that the products meet company or government standards. Supermarkets routinely weigh meat and produce and charge consumers by the ounce or the pound.

8.1: Prelude to Quantities in Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.0: Prelude to Quantities in Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.2: The Mole

Learning Objectives

- To define the mole unit.

Figure 8.2.1 shows that we need 2 hydrogen atoms and 1 oxygen atom to make 1 water molecule. If we want to make 2 water molecules, we will need 4 hydrogen atoms and 2 oxygen atoms. If we want to make 5 molecules of water, we need 10 hydrogen atoms and 5 oxygen atoms. The ratio of atoms we will need to make any number of water molecules is the same: 2 hydrogen atoms to 1 oxygen atom.




Figure 8.2.1 Water Molecules. The ratio of hydrogen atoms to oxygen atoms used to make water molecules is always 2:1, no matter how many water molecules are being made.

One problem we have, however, is that it is extremely difficult, if not impossible, to organize atoms one at a time. As stated in the introduction, we deal with billions of atoms at a time. How can we keep track of so many atoms (and molecules) at a time? We do it by using mass rather than by counting individual atoms.

A hydrogen atom has a mass of approximately 1 u. An oxygen atom has a mass of approximately 16 u. The ratio of the mass of an oxygen atom to the mass of a hydrogen atom is therefore approximately 16:1.

If we have 2 atoms of each element, the ratio of their masses is approximately 32:2, which reduces to 16:1—the same ratio. If we have 12 atoms of each element, the ratio of their total masses is approximately (12 × 16):(12 × 1), or 192:12, which also reduces to 16:1. If we have 100 atoms of each element, the ratio of the masses is approximately 1,600:100, which again reduces to 16:1. As long as we have equal numbers of hydrogen and oxygen atoms, the ratio of the masses will always be 16:1.

The same consistency is seen when ratios of the masses of other elements are compared. For example, the ratio of the masses of silicon atoms to equal numbers of hydrogen atoms is always approximately 28:1, while the ratio of the masses of calcium atoms to equal numbers of lithium atoms is approximately 40:7.

So we have established that the masses of atoms are constant with respect to each other, as long as we have the same number of each type of atom. Consider a more macroscopic example. If a sample contains 40 g of Ca, this sample has the same number of

atoms as there are in a sample of 7 g of Li. What we need, then, is a number that represents a convenient quantity of atoms so we can relate macroscopic quantities of substances. Clearly even 12 atoms are too few because atoms themselves are so small. We need a number that represents billions and billions of atoms.

Chemists use the term **mole** to represent a large number of atoms or molecules. Just as a dozen implies 12 things, a **mole** (abbreviated as ***mol***) represents **6.022×10^{23} things**. The number 6.022×10^{23}, called Avogadro's number after the 19th-century chemist Amedeo Avogadro, is the number we use in chemistry to represent macroscopic amounts of atoms and molecules. Thus, if we have 6.022×10^{23} Na atoms, we say we have 1 mol of Na atoms. If we have 2 mol of Na atoms, we have $2 \times (6.022 \times 10^{23})$ Na atoms, or 1.2044×10^{24} Na atoms. Similarly, if we have 0.5 mol of benzene (C_6H_6) molecules, we have $0.5 \times (6.022 \times 10^{23})$ C_6H_6 molecules, or 3.011×10^{23} C_6H_6 molecules.

> A mole represents a very large number! If 1 mol of quarters were stacked in a column, it could stretch back and forth between Earth and the sun *6.8 billion* times.

Notice that we are applying the mole unit to different types of **chemical entities**. The word *mole* represents a number of things—6.022×10^{23} of them—but does not by itself specify what "they" are. The chemical entities can be ***atoms, molecules***, ***formula units*** and ***ions***. This specific information needs to be specified accurately. Most students find this confusing hence, we need to review the composition of elements, covalent and ionic compounds.

Most elements are made up of individual ***atoms,*** such as helium. However, some elements consist of ***molecules***, such as the diatomic elements, nitrogen, hydrogen, oxygen, etc. One mole of He consists of 6.022×10^{23} He ***atoms*** but one mole of nitrogen contains 6.022×10^{23} N_2 ***molecules***. The basic units of covalent (molecular) compounds are ***molecules*** as well. The molecules of "compounds" consist of different kinds of atoms while the molecules of "elements" consist of only one type of atom. For example, the molecules of ammonia (NH_3) consist of nitrogen and hydrogen atoms while N_2 molecules have N atoms only. Compounds that are ionic, like NaCl, are represented by ionic formulas. One mole of NaCl, for example, refers to 6.022×10^{23} ***formula units*** of NaCl. And, one formula unit of NaCl consists of one sodium ***ion*** and one chloride ***ion***. Figure 6.1.2 summarizes the basic units of elements, covalent and ionic compounds




Figure 8.2.2: The basic units of elements (atoms or molecules), covalent compounds (molecules) and ionic compounds (formula units of ions).

Conversion Between Moles and Atoms, Molecules and Ions

Using our unit conversion techniques learned in Chapter 1, we can use the mole relationship and the chemical formula to convert back and forth between the moles and the number of chemical entities (atoms, molecules or ions).

Because 1 N_2 molecule contains 2 N atoms, 1 mol of N_2 molecules (6.022×10^{23} molecules) has 2 mol of N atoms. Using formulas to indicate how many atoms of each element we have in a substance, we can relate the number of moles of molecules to the number of moles of atoms. For example, in 1 mol of ethanol (C_2H_6O), we can construct the following relationships (Table 8.2.1):

Table 8.2.1: Molecular Relationships

1 Molecule of C_2H_6O Has	1 Mol of C_2H_6O Has	Molecular Relationships
2 C atoms	2 mol of C atoms	$\frac{2\text{ mol C atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{2\text{ mol C atoms}}$
6 H atoms	6 mol of H atoms	$\frac{6\text{ mol H atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{6\text{ mol H atoms}}$
1 O atom	1 mol of O atoms	$\frac{1\text{ mol O atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{1\text{ mol O atoms}}$

The following example illustrates how we can use these relationships as conversion factors.

Example 8.2.1

If a sample consists of 2.5 mol of ethanol (C_2H_6O), how many moles of carbon atoms, hydrogen atoms, and oxygen atoms does it have?

Solution

Using the relationships in Table 8.2.1, we apply the appropriate conversion factor for each element:

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \underbrace{\frac{2\text{ mol C atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}}_{\text{conversion factor}} = 5.0\text{ mol C atoms}$$

Note how the unit *mol C_2H_6O molecules* cancels algebraically. Similar equations can be constructed for determining the number of H and O atoms:

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \frac{6\text{ mol H atoms}}{1\text{ mol }C_2H_6O\text{ molecules}} = 15\text{ mol H atoms}$$

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \frac{1\text{ mol O atoms}}{1\text{ mol }C_2H_6O\text{ molecules}} = 2.5\text{ mol O atoms}$$

Exercise 8.2.1

If a sample contains 6.75 mol of Na_2SO_4, how many moles of sodium atoms, sulfur atoms, and oxygen atoms does it have?

Answer

13.5 mol Na, 6.75 mol S and 27 mol O.

We can use Avogadro's number as a conversion factor, or ratio, in dimensional analysis problems. For example, if we are dealing with element X, the mole relationship is expressed as follows:

$$1\text{ mol X} = 6.022 \times 10^{23}\text{ X atoms}$$

We can convert this relationship into two possible conversion factors shown below:

$$\frac{1\text{ mol X}}{6.022 \times 10^{23}\text{ X atoms}} \text{ or } \frac{6.022 \times 10^{23}\text{ X atoms}}{1\text{ mol X}}$$

If the number of "atoms of element X" is given, we can convert it into "moles of X" by multiplying the given value with the conversion factor at the left. However, if the number of "mol of X" is given, the appropriate conversion factor to use is the one at the right.

If we are dealing with a molecular compound (such as C_4H_{10}), the mole relationship is expressed as follows:

$$1 \text{ mol } C4H10 = 6.022 \times 10^{23} \text{ } C4H10 \text{ molecules}$$

If working with ionic compounds (such as NaCl), the mole relationship is expressed as follows:


$$1 \text{ mol NaCl} = 6.022 \times 10^{23} \text{ NaCl formula units}$$

Example 8.2.2

How many formula units are present in 2.34 mol of NaCl? How many ions are in 2.34 mol?

Solution

Typically in a problem like this, we start with what we are given and apply the appropriate conversion factor. Here, we are given a quantity of 2.34 mol of NaCl, to which we can apply the definition of a **mole as a conversion factor**:



$$2.34 \text{ mol NaCl} \times \frac{6.022 \times 10^{23} \text{ NaCl units}}{1 \text{ mol NaCl}} = 1.41 \times 10^{24} \text{ NaCl units}$$

Because there are two ions per formula unit, there are

$$1.41 \times 10^{24} \text{ NaCl units} \times \frac{2 \text{ ions}}{\text{NaCl units}} = 2.82 \times 10^{24} \text{ ions}$$

in the sample.

Exercise 8.2.2

How many molecules are present in 16.02 mol of C_4H_{10}? How many atoms are in 16.02 mol?

Answer

9.647 x 10^{24}molecules, 1.351 x 10^{26} atoms.

Key Takeaway

- A mole is 6.022 × 10^{23} things.

8.2: The Mole is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.1: The Mole** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.3: Atomic and Molar Masses

Learning Objectives

- To learn how the masses of moles of atoms and molecules are expressed.

Now that we have introduced the mole and practiced using it as a conversion factor, we ask the obvious question: why is the mole *that particular* number of things? Why is it 6.022×10^{23} and not 1×10^{23} or even 1×10^{20} ?

The number in a mole, Avogadro's number, is related to the relative sizes of the **atomic mass unit** and **gram mass units**. Whereas one hydrogen atom has a mass of approximately 1 u, **1 mol of H atoms** has a mass of approximately **1 *gram***. And whereas one sodium atom has an approximate mass of 23 u, 1 mol of Na atoms has an approximate mass of 23 *grams*.

One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units. The numbers in the periodic table that we identified as the atomic masses of the atoms not only tell us the mass of one atom in u but also tell us the mass of 1 mol of atoms in grams.

> One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units.

Example 8.3.1: Moles to Mass Conversion with Elements

What is the mass of each quantity?

a. 1 mol of Al atoms
b. 2 mol of U atoms

Solution

a. One mole of Al atoms has a mass in grams that is numerically equivalent to the atomic mass of aluminum. The periodic table shows that the atomic mass (rounded to two decimal points) of Al is 26.98, so 1 mol of Al atoms has a mass of 26.98 g.
b. According to the periodic table, 1 mol of U has a mass of 238.0 g, so the mass of 2 mol is twice that, or 476.0 g.

Exercise 8.3.1: Moles to Mass Conversion with Elements

What is the mass of each quantity?

a. 1 mol of Au atoms
b. 5 mol of Br atoms

Answer a:

197.0 g

Answer b:

5 mol Br atoms x 79.90 g/mol = 399.5 g

The mole concept can be extended to masses of formula units and molecules as well. The mass of 1 mol of molecules (or formula units) in grams is numerically equivalent to the mass of one molecule (or formula unit) in atomic mass units. For example, a single molecule of O_2 has a mass of 32.00 u, and 1 mol of O_2 molecules has a mass of 32.00 g. As with atomic mass unit–based masses, to obtain the mass of 1 mol of a substance, we simply sum the masses of the individual atoms in the formula of that substance. The mass of 1 mol of a substance is referred to as its **molar mass**, whether the substance is an element, an ionic compound, or a covalent compound.

Example 8.3.2: Moles to Mass Conversion with Compounds

What is the mass of 1 mol of each substance?

1. NaCl
2. bilirubin ($C_{33}H_{36}N_4O_6$), the principal pigment present in bile (a liver secretion)

Solution

1. Summing the molar masses of the atoms in the NaCl formula unit gives

Solutions to Example 6.2.2, Summing the molar masses of the atoms in the NaCl formula unit gives

1 Na molar mass:	22.99 g
1 Cl molar mass:	35.45 g
Total:	58.44 g

The mass of 1 mol of NaCl is 58.44 g.

2. Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

Solutions to Example 6.2.2, Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

33 C molar mass:	33 × 12.01 g	396.33 g
36 H molar mass:	36 × 1.01 =	36.36 g
4 N molar mass:	4 × 14.01 =	56.04 g
6 O molar mass:	6 × 16.00 =	96.00 g
Total:		584.73 g

The mass of 1 mol of bilirubin is 584.73 g.

Exercise 8.3.2: Moles to Mass Conversion with Compounds

What is the mass of 1 mol of each substance?

a. barium sulfate ($BaSO_4$), used to take X rays of the gastrointestional tract
b. adenosine ($C_{10}H_{13}N_5O_4$), a component of cell nuclei crucial for cell division

Answer a:

233.36 g

Answer b:

267.28 g

Be careful when counting atoms. In formulas with polyatomic ions in parentheses, the subscript outside the parentheses is applied to every atom inside the parentheses. For example, the molar mass of $Ba(OH)_2$ requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

The molar mass of Ba(OH)2 requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

1 Ba molar mass:	1 × 137.3 g =	137.3 g
2 O molar mass:	2 × 16.00 g =	32.00 g
2 H molar mass:	2 × 1.01 g =	2.02 g

1 Ba molar mass:	1 × 137.3 g =	137.3 g
Total:		171.32 g

Because molar mass is defined as the mass for 1 mol of a substance, we can refer to molar mass as grams per mole (g/mol). The division sign (/) implies “per,” and “1” is implied in the denominator. Thus, the molar mass of bilirubin can be expressed as 584.73 g/mol, which is read as “five hundred eighty four point seventy three grams per mole.”

Key Takeaway

- The mass of moles of atoms and molecules is expressed in units of grams.

8.3: Atomic and Molar Masses is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.2: Atomic and Molar Masses** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.4: Mole-Mass Conversions

Learning Objectives

- To convert quantities between mass units and mole units.

A previous Example stated that the mass of 2 mol of U is twice the molar mass of uranium. Such a straightforward exercise does not require any formal mathematical treatment. Many questions concerning mass are not so straightforward, however, and require some mathematical manipulations.

The simplest type of manipulation using molar mass as a conversion factor is a mole-mass conversion (or its reverse, a mass-mole conversion). In such a conversion, we use the molar mass of a substance as a conversion factor to convert mole units into mass units (or, conversely, mass units into mole units).

We also established that 1 mol of Al has a mass of 26.98 g (Example 8.4.1). Stated mathematically,

$$1 \text{ mol Al} = 26.98 \text{ g Al}$$

We can divide both sides of this expression by either side to get one of two possible conversion factors:

$$\frac{1 \text{ mol Al}}{26.98 \text{ g Al}} \quad \text{and} \quad \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$


The first conversion factor can be used to convert from mass to moles, and the second converts from moles to mass. Both can be used to solve problems that would be hard to do "by eye."

Example 8.4.1

What is the mass of 3.987 mol of Al?

Solution

The first step in a conversion problem is to decide what conversion factor to use. Because we are starting with mole units, we want a conversion factor that will cancel the mole unit and introduce the unit for mass in the numerator. Therefore, we should use the $\frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$ conversion factor. We start with the given quantity and multiply by the conversion factor:



$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$

Note that the mol units cancel algebraically. (The quantity 3.987 mol is understood to be in the numerator of a fraction that has 1 in the unwritten denominator.) Canceling and solving gives

$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = 107.6 \text{ g Al}$$

Our final answer is expressed to four significant figures.

Exercise 8.4.1

How many moles are present in 100.0 g of Al? (Hint: you will have to use the other conversion factor we obtained for aluminum.)

Answer

$$100.0 \text{ g Al} \times \frac{1 \text{ mol Al}}{26.98 \text{ g Al}} = 3.706 \text{ mol Al}$$

Conversions like this are possible for any substance, as long as the proper atomic mass, formula mass, or molar mass is known (or can be determined) and expressed in grams per mole. Figure 8.4.1 is a chart for determining what conversion factor is needed, and Figure 8.4.2 is a flow diagram for the steps needed to perform a conversion.




Figure 8.4.1 A Simple Flowchart for Converting between Mass and Moles of a Substance. It takes one mathematical step to convert from moles to mass or from mass to moles.




Figure 8.4.2 A Flowchart Illustrating the Steps in Performing a Unit Conversion. When performing many unit conversions, the same logical steps can be taken.

Example 8.4.2

A biochemist needs 0.00655 mol of bilirubin ($C_{33}H_{36}N_4O_6$) for an experiment. How many grams of bilirubin will that be?

Solution

To convert from moles to mass, we need the molar mass of bilirubin, which we can determine from its chemical formula:

Solutions to Example 6.3.2

33 C molar mass:	33 × 12.01 g =	396.33 g
36 H molar mass:	36 × 1.01 g =	36.36 g
4 N molar mass:	4 × 14.01 g =	56.04 g
6 O molar mass:	6 × 16.00 g =	96.00 g
Total:		584.73 g

The molar mass of bilirubin is 584.73 g. Using the relationship

1 mol bilirubin = 584.73 g bilirubin

we can construct the appropriate conversion factor for determining how many grams there are in 0.00655 mol. Following the steps from Figure 8.4.2:

$$0.00655 \text{ mol bilirubin} \times \frac{584.73 \text{ g bilirubin}}{\text{mol bilirubin}} = 3.83 \text{ g bilirubin}$$

The mol bilirubin unit cancels. The biochemist needs 3.83 g of bilirubin.

Exercise 8.4.2

A chemist needs 457.8 g of $KMnO_4$ to make a solution. How many moles of $KMnO_4$ is that?

Answer

$$457.8 \text{ g } KMnO_4 \times \frac{1 \text{ mol } KMnO_4}{158.04 \text{ g } KMnO_4} = 2.897 \text{ mol } KMnO_4$$

To Your Health: Minerals

For our bodies to function properly, we need to ingest certain substances from our diets. Among our dietary needs are minerals, the noncarbon elements our body uses for a variety of functions, such developing bone or ensuring proper nerve transmission. The US Department of Agriculture has established some recommendations for the RDIs of various minerals. The accompanying table lists the RDIs for minerals, both in mass and moles, assuming a 2,000-calorie daily diet.

Table 8.4.1: Essential Minerals and their Composition in Humans

Mineral	Male (age 19–30 y)		Female (age 19–30 y)	
Ca	1,000 mg	0.025 mol	1,000 mg	0.025 mol
Cr	35 µg	6.7×10^{-7} mol	25 µg	4.8×10^{-7} mol
Cu	900 µg	1.4×10^{-5} mol	900 µg	1.4×10^{-5} mol
F	4 mg	2.1×10^{-4} mol	3 mg	1.5×10^{-4} mol
I	150 µg	1.2×10^{-6} mol	150 µg	1.2×10^{-6} mol
Fe	8 mg	1.4×10^{-4} mol	18 mg	3.2×10^{-4} mol
K	3,500 mg	9.0×10^{-2} mol	3,500 mg	9.0×10^{-2} mol
Mg	400 mg	1.6×10^{-2} mol	310 mg	1.3×10^{-2} mol
Mn	2.3 mg	4.2×10^{-5} mol	1.8 mg	3.3×10^{-5} mol
Mo	45 mg	4.7×10^{-7} mol	45 mg	4.7×10^{-7} mol
Na	2,400 mg	1.0×10^{-1} mol	2,400 mg	1.0×10^{-1} mol
P	700 mg	2.3×10^{-2} mol	700 mg	2.3×10^{-2} mol
Se	55 µg	7.0×10^{-7} mol	55 µg	7.0×10^{-7} mol
Zn	11 mg	1.7×10^{-4} mol	8 mg	1.2×10^{-4} mol

Table 8.4.1 illustrates several things. First, the needs of men and women for some minerals are different. The extreme case is for iron; women need over twice as much as men do. In all other cases where there is a different RDI, men need more than women.

Second, the amounts of the various minerals needed on a daily basis vary widely—both on a mass scale and a molar scale. The average person needs 0.1 mol of Na a day, which is about 2.5 g. On the other hand, a person needs only about 25–35 µg of Cr per day, which is under one millionth of a mole. As small as this amount is, a deficiency of chromium in the diet can lead to diabetes-like symptoms or neurological problems, especially in the extremities (hands and feet). For some minerals, the body does not require much to keep itself operating properly.

Although a properly balanced diet will provide all the necessary minerals, some people take dietary supplements. However, too much of a good thing, even minerals, is not good. Exposure to too much chromium, for example, causes a skin irritation, and certain forms of chromium are known to cause cancer (as presented in the movie *Erin Brockovich*).

Key Takeaway

- It is possible to convert between moles of material and mass of material.

8.4: Mole-Mass Conversions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.3: Mole-Mass Conversions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.5: Mole-Mole Relationships in Chemical Reactions

Learning Objectives

- To use a balanced chemical reaction to determine molar relationships between the substances.

Previously, you learned to balance chemical equations by comparing the numbers of each type of atom in the reactants and products. The coefficients in front of the chemical formulas represent the numbers of molecules or formula units (depending on the type of substance). Here, we will extend the meaning of the coefficients in a chemical equation.

Consider the simple chemical equation

$$2\,H_2 + O_2 \rightarrow 2\,H_2O$$

The convention for writing balanced chemical equations is to use the lowest whole-number ratio for the coefficients. However, the equation is balanced as long as the coefficients are in a 2:1:2 ratio. For example, this equation is also balanced if we write it as

$$4\,H_2 + 2\,O_2 \rightarrow 4\,H_2O$$

The ratio of the coefficients is 4:2:4, which reduces to 2:1:2. The equation is also balanced if we were to write it as

$$22\,H_2 + 11\,O_2 \rightarrow 22\,H_2O$$

because 22:11:22 also reduces to 2:1:2.

Suppose we want to use larger numbers. Consider the following coefficients:

$$12.044 \times 10^{23}\ H_2 + 6.022 \times 10^{23}\ O_2 \rightarrow 12.044 \times 10^{23}\ H_2O$$


These coefficients also have the ratio 2:1:2 (check it and see), so this equation is balanced. But 6.022×10^{23} is 1 mol, while 12.044×10^{23} is 2 mol (and the number is written that way to make this more obvious), so we can simplify this version of the equation by writing it as

$$2\ \text{mol}\ H_2 + 1\ \text{mol}\ O_2 \rightarrow 2\ \text{mol}\ H_2O$$

We can leave out the word *mol* and not write the 1 coefficient (as is our habit), so the final form of the equation, still balanced, is

$$2\,H_2 + O_2 \rightarrow 2\,H_2O$$

Now we interpret the coefficients as referring to molar amounts, not individual molecules. The lesson? *Balanced chemical equations are balanced not only at the molecular level but also in terms of molar amounts of reactants and products.* Thus, we can read this reaction as “two moles of hydrogen react with one mole of oxygen to produce two moles of water.”



2 molecules H_2 1 molecule O_2 2 molecules H_2O

2 moles H_2 1 mole O_2 2 moles H_2O

2 x 2.02 g=4.04 g H_2 32.0 g O_2 2 x 18.02 g=36.04 g H_2O

Figure 8.5.1: This representation of the production of water from oxygen and hydrogen show several ways to interpret the quantitative information of a chemical reaction.

By the same token, the ratios we constructed to describe molecules reaction can also be constructed in terms of moles rather than molecules. For the reaction in which hydrogen and oxygen combine to make water, for example, we can construct the following

ratios:

$$\frac{2\text{ mol }H_2}{1\text{ mol }O_2}\text{ or }\frac{1\text{ mol }O_2}{2\text{ mol }H_2}$$

$$\frac{2\text{ mol }H_2O}{1\text{ mol }O_2}\text{ or }\frac{1\text{ mol }O_2}{2\text{ mol }H_2O}$$

$$\frac{2\text{ mol }H_2}{2\text{ mol }H_2O}\text{ or }\frac{2\text{ mol }H_2O}{2\text{ mol }H_2}$$

We can use these ratios to determine what amount of a substance, in moles, will react with or produce a given number of moles of a different substance. The study of the numerical relationships between the reactants and the products in balanced chemical reactions is called ***stoichiometry*****.** The ratio of coefficients in a balanced chemical equation, used in computations relating amounts of reactants and products is called the **stoichiometric factor.**


Example 8.5.1

How many moles of oxygen react with hydrogen to produce 27.6 mol of H_2O? The balanced equation is as follows:

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

Solution

Because we are dealing with quantities of H_2O and O_2, we will use the **stoichiometric ratio** that relates those two substances. Because we are given an amount of H_2O and want to determine an amount of O_2, we will use the ratio that has H_2O in the denominator (so it cancels) and O_2 in the numerator (so it is introduced in the answer). Thus,



$$27.6\text{ mol }H_2O \times \frac{1\text{ mol }O_2}{2\text{ mol }H_2O} = 13.8\text{ mol }O_2$$

To produce 27.6 mol of H_2O, 13.8 mol of O_2 react.

Exercise 8.5.1

Using $2H_2 + O_2 \rightarrow 2H_2O$, how many moles of hydrogen react with 3.07 mol of oxygen to produce H_2O?

Answer

$$3.07\text{ mol }O_2 \times \frac{2\text{ mol }H_2}{1\text{ mol }O_2} = 6.14\text{ mol }H_2$$

Key Takeaway

- The balanced chemical reaction can be used to determine molar relationships between substances.

8.5: Mole-Mole Relationships in Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.


- **6.4: Mole-Mole Relationships in Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.6: Mole-Mass and Mass-Mass Problems

Learning Objectives

- To convert from mass or moles of one substance to mass or moles of another substance in a chemical reaction.

We have established that a balanced chemical equation is balanced in terms of moles as well as atoms or molecules. We have used balanced equations to set up ratios, now in terms of moles of materials, that we can use as conversion factors to answer stoichiometric questions, such as how many moles of substance A react with so many moles of reactant B. We can extend this technique even further. Recall that we can relate a molar amount to a mass amount using molar mass. We can use that ability to answer stoichiometry questions in terms of the masses of a particular substance, in addition to moles. We do this using the following sequence:



Collectively, these conversions are called mole-mass calculations.

As an example, consider the balanced chemical equation

$$\mathrm{Fe_2O_3} + 3\,\mathrm{SO_3} \rightarrow \mathrm{Fe_2(SO_4)_3}$$

If we have 3.59 mol of Fe_2O_3, how many grams of SO_3 can react with it? Using the mole-mass calculation sequence, we can determine the required mass of SO_3 in two steps. First, we construct the appropriate molar ratio, determined from the balanced chemical equation, to calculate the number of moles of SO_3 needed. Then using the molar mass of SO_3 as a conversion factor, we determine the mass that this number of moles of SO_3 has.

The first step resembles the exercises we did in Section 6.4. As usual, we start with the quantity we were given:


$$3.59\ \text{mol}\ \mathrm{Fe_2O_3} \times \frac{3\ \text{mol}\ \mathrm{SO_3}}{1\ \text{mol}\ \mathrm{Fe_2O_3}} = 10.77\ \text{mol}\ \mathrm{SO_3}$$

The mol Fe_2O_3 units cancel, leaving mol SO_3 unit. Now, we take this answer and convert it to grams of SO_3, using the molar mass of SO_3 as the conversion factor:

$$10.77\ \text{mol}\ \mathrm{SO_3} \times \frac{80.07\ \text{g}\ \mathrm{SO_3}}{1\ \text{mol}\ \mathrm{SO_3}} = 862.4\ \text{g}\ \mathrm{SO_3}$$

Our final answer is expressed to three significant figures. Thus, in a two-step process, we find that 862 g of SO_3 will react with 3.59 mol of Fe_2O_3. Many problems of this type can be answered in this manner.

The same two-step problem can also be worked out in a single line, rather than as two separate steps, as follows:



$$3.59 \text{ mol } Fe_2O_3 \times \underbrace{\frac{3 \text{ mol } SO_3}{1 \text{ mol } Fe_2O_3}}_{\text{converts to moles of } SO_3} \times \underbrace{\frac{80.07 \text{ g } SO_3}{1 \text{ mol } SO_3}}_{\text{converts to grams of } SO_3} = 862.4 \text{ g } SO_3$$

We get exactly the same answer when combining all the math steps together as we do when we calculate one step at a time.

Example 8.6.1

How many grams of CO_2 are produced if 2.09 mol of HCl are reacted according to this balanced chemical equation?

$$CaCO_3 + 2\,HCl \rightarrow CaCl_2 + CO_2 + H_2O$$

Solution

Our strategy will be to convert from moles of HCl to moles of CO_2 and then from moles of CO_2 to grams of CO_2. We will need the molar mass of CO_2, which is 44.01 g/mol. Performing these two conversions in a single-line gives 46.0 g of CO_2:

$$2.09 \text{ mol HCl} \times \underbrace{\frac{1 \text{ mol } CO_2}{2 \text{ mol HCl}}}_{\text{from the coefficients of the balanced equation}} \times \frac{44.01 \text{ g } CO_2}{1 \text{ mol } CO_2} = 46.0 \text{ g } CO_2$$

The molar ratio between CO_2 and HCl comes from the balanced chemical equation.

Exercise


How many grams of glucose ($C_6H_{12}O_6$) are produced if 17.3 mol of H_2O are reacted according to this balanced chemical equation?

$$6\,CO_2 + 6\,H_2O \rightarrow C_6H_{12}O_6 + 6\,O_2$$

Answer

$$17.3 \text{ mol } H_2O \times \frac{1 \text{ mol } C_6H_{12}O_6}{6 \text{ mol } H_2O} \times \frac{180.18 \text{ g } C_6H_{12}O_6}{1 \text{ mol } C_6H_{12}O_6} = 520 \text{ g } C_6H_{12}O_6$$

It is a small step from mole-mass calculations to mass-mass calculations. If we start with a known mass of one substance in a chemical reaction (instead of a known number of moles), we can calculate the corresponding masses of other substances in the reaction. The first step in this case is to convert the known mass into moles, using the substance's molar mass as the conversion factor. Then—and only then—we use the balanced chemical equation to construct a conversion factor to convert that quantity to moles of another substance, which in turn can be converted to a corresponding mass. Sequentially, the process is as follows:



This three-part process can be carried out in three discrete steps or combined into a single calculation that contains three conversion factors. The following example illustrates both techniques.

Example 8.6.2: Chlorination of Carbon

Methane can react with elemental chlorine to make carbon tetrachloride (CCl_4). The balanced chemical equation is as follows:

$$CH_4 + 4\,Cl_2 \rightarrow CCl_4 + 4\,HCl$$

How many grams of HCl are produced by the reaction of 100.0 g of CH_4?

Solution

First, let us work the problem in stepwise fashion. We begin by converting the mass of CH_4 to moles of CH_4, using the molar mass of CH_4 (16.05 g/mol) as the conversion factor:

$$100.0\text{ g } CH_4 \times \frac{1\text{ mol } CH_4}{16.05\text{ g } CH_4} = 6.231\text{ mol } CH_4$$


Note that we inverted the molar mass so that the gram units cancel, giving us an answer in moles. Next, we use the balanced chemical equation to determine the ratio of moles CCl_4 and moles HCl and convert our first result into moles of HCl:

$$6.231\text{ mol } CH_4 \times \frac{4\text{ mol HCl}}{1\text{ mol } CH_4} = 24.92\text{ mol HCl}$$

Finally, we use the molar mass of HCl (36.46 g/mol) as a conversion factor to calculate the mass of 24.92 mol of HCl:

$$24.92\text{ mol HCl} \times \frac{36.46\text{ g HCl}}{1\text{ mol HCl}} = 908.5\text{ g HCl}$$

In each step, we have limited the answer to the proper number of significant figures. If desired, we can do all three conversions on a single line:



$$100.0 \text{ g } CH_4 \times \frac{1 \text{ mol } CH_4}{16.05 \text{ g } CH_4} \times \frac{4 \text{ mol HCl}}{1 \text{ mol } CH_4} \times \frac{36.46 \text{ g HCl}}{1 \text{ mol HCl}} = 908.7 \text{ g HCl}$$

This final answer is slightly different from our first answer because only the final answer is restricted to the proper number of significant figures. In the first answer, we limited each intermediate quantity to the proper number of significant figures. As you can see, both answers are essentially the same.

Exercise 8.6.2: Oxidation of Propanal

The oxidation of propanal (CH_3CH_2CHO) to propionic acid (CH_3CH_2COOH) has the following chemical equation:

$$CH_3CH_2CHO + 2K_2Cr_2O_7 \rightarrow CH_3CH_2COOH + \text{other products}$$

How many grams of propionic acid are produced by the reaction of 135.8 g of $K_2Cr_2O_7$?

Answer

$$135.8 \text{ g } K_2Cr_2O_7 \times \frac{1 \text{ mol } K_2Cr_2O_7}{294.20 \text{ g } K_2Cr_2O_7} \times \frac{1 \text{ mol } CH_3CH_2COOH}{2 \text{ mol } K_2Cr_2O_7} \times \frac{74.09 \text{ g } CH_3CH_2COOH}{1 \text{ mol } CH_3CH_2COOH} = 17.10 \text{ g } CH_3CH_2COOH$$

To Your Health: The Synthesis of Taxol

Taxol is a powerful anticancer drug that was originally extracted from the Pacific yew tree (*Taxus brevifolia*). As you can see from the accompanying figure, taxol is a very complicated molecule, with a molecular formula of $C_{47}H_{51}NO_{14}$. Isolating taxol from its natural source presents certain challenges, mainly that the Pacific yew is a slow-growing tree, and the equivalent of six trees must be harvested to provide enough taxol to treat a single patient. Although related species of yew trees also produce taxol in small amounts, there is significant interest in synthesizing this complex molecule in the laboratory.

After a 20-year effort, two research groups announced the complete laboratory synthesis of taxol in 1994. However, each synthesis required over 30 separate chemical reactions, with an overall efficiency of less than 0.05%. To put this in perspective, to obtain a single 300 mg dose of taxol, you would have to begin with 600 g of starting material. To treat the 26,000 women who are diagnosed with ovarian cancer each year with one dose, almost 16,000 kg (over 17 tons) of starting material must be converted to taxol. Taxol is also used to treat breast cancer, with which 200,000 women in the United States are diagnosed every year. This only increases the amount of starting material needed.

Clearly, there is intense interest in increasing the overall efficiency of the taxol synthesis. An improved synthesis not only will be easier but also will produce less waste materials, which will allow more people to take advantage of this potentially life-saving drug.




Figure 8.6.1 The Structure of the Cancer Drug Taxol. Because of the complexity of the molecule, hydrogen atoms are not shown, but they are present on every atom to give the atom the correct number of covalent bonds (four bonds for each carbon atom).

Key Takeaway

- A balanced chemical equation can be used to relate masses or moles of different substances in a reaction.

8.6: Mole-Mass and Mass-Mass Problems is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.5: Mole-Mass and Mass-Mass Problems** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.E: Quantities in Chemical Reactions (Exercise)

6.1: The Mole

Concept Review Exercise

1. What is a mole?

Answer

1. A mole is 6.022×10^{23} things.

Exercises

1. How many dozens are in 1 mol? Express your answer in proper scientific notation.
2. A gross is a dozen dozen, or 144 things. How many gross are in 1 mol? Express your answer in proper scientific notation.
3. How many moles of each type of atom are in 1.0 mol of $C_6H_{12}O_6$?
4. How many moles of each type of atom are in 1.0 mol of $K_2Cr_2O_7$?
5. How many moles of each type of atom are in 2.58 mol of Na_2SO_4?
6. How many moles of each type of atom are in 0.683 mol of $C_{34}H_{32}FeN_4O_4$? (This is the formula of heme, a component of hemoglobin.)
7. How many molecules are in 16.8 mol of H_2O?
8. How many formula units are in 0.778 mol of iron(III) nitrate?
9. A sample of gold contains 7.02×10^{24} atoms. How many moles of gold is this?
10. A flask of mercury contains 3.77×10^{22} atoms. How many moles of mercury are in the flask?
11. An intravenous solution of normal saline may contain 1.72 mol of sodium chloride (NaCl). How many sodium and chlorine atoms are present in the solution?
12. A lethal dose of arsenic is 1.00×10^{21} atoms. How many moles of arsenic is this?

Answers

1. 5.018×10^{22} dozens
2. 4.18×10^{21} grosses
3. 6.0 mol of C atoms, 12.0 mol of H atoms, and 6.0 mol of O atoms
4. 2.0 mol of K atoms, 2.0 mol of Cr atoms, and 7.0 mol of O atoms
5. 5.16 mol of Na atoms, 2.58 mol of S atoms, and 10.32 mol of O atoms
6. 23.2 mol of C atoms, 21.9 mol of H atoms, 0.683 mol of Fe, 2.73 mol of N and 2.73 mol of O atoms
7. 1.012×10^{25} molecules
8. 4.69×10^{23} formula units
9. 11.7 mol
10. 0.0626 mol
11. 1.04×10^{24} Na atoms and 1.04×10^{24} Cl atoms
12. 0.00166 mol

6.2: Atomic and Molar Masses

Concept Review Exercises

1. How are molar masses of the elements determined?
2. How are molar masses of compounds determined?

Answers

1. Molar masses of the elements are the same numeric value as the masses of a single atom in atomic mass units but in units of grams instead.
2. Molar masses of compounds are calculated by adding the molar masses of their atoms.

Exercises

1. What is the molar mass of Si? What is the molar mass of U?
2. What is the molar mass of Mn? What is the molar mass of Mg?
3. What is the molar mass of $FeCl_2$? What is the molar mass of $FeCl_3$?
4. What is the molar mass of C_6H_6? What is the molar mass of $C_6H_5CH_3$?
5. What is the molar mass of $(NH_4)_2S$? What is the molar mass of $Ca(OH)_2$?
6. What is the molar mass of $(NH_4)_3PO_4$? What is the molar mass of $Sr(HCO_3)_2$?
7. Aspirin ($C_9H_8O_4$) is an analgesic (painkiller) and antipyretic (fever reducer). What is the molar mass of aspirin?
8. Ibuprofen ($C_{13}H_{18}O_2$) is an analgesic (painkiller). What is the molar mass of ibuprofen?
9. Morphine ($C_{17}H_{19}NO_3$) is a narcotic painkiller. What is the mass of 1 mol of morphine?
10. Heroin ($C_{21}H_{23}NO_5$) is a narcotic drug that is a derivative of morphine. What is the mass of 1 mol of heroin?

Answers

1. 28.09 g/mol; 238.0 g/mol
2. 54.94 g/mol; 24.31 g/mol
3. 126.75 g/mol; 162.20 g/mol
4. 78.12 g/mol; 92.15 g/mol
5. 68.16 g/mol; 74.10 g/mol
6. 149.12 g/mol; 209.64 g/mol
7. 180.17 g/mol
8. 206.31 g/mol
9. 285.37 g
10. 369.45 g

6.3: Mole-Mass Conversions

Concept Review Exercises

1. What relationship is needed to perform mole-mass conversions?
2. What information determines which conversion factor is used in a mole-mass conversion?

Answers

1. The atomic or molar mass is needed for a mole-mass conversion.
2. The unit of the initial quantity determines which conversion factor is used.

Exercises

1. What is the mass of 8.603 mol of Fe metal?
2. What is the mass of 0.552 mol of Ag metal?
3. What is the mass of 6.24×10^4 mol of Cl_2 gas?
4. What is the mass of 0.661 mol of O_2 gas?
5. What is the mass of 20.77 mol of $CaCO_3$?
6. What is the mass of 9.02×10^{-3} mol of the hormone epinephrine ($C_9H_{13}NO_3$)?
7. How many moles are present in 977.4 g of $NaHCO_3$?
8. How many moles of erythromycin ($C_{37}H_{67}NO_{13}$), a widely used antibiotic, are in 1.00×10^3 g of the substance?

9. Cortisone ($C_{21}H_{28}O_5$) is a synthetic steroid that is used as an anti-inflammatory drug. How many moles of cortisone are present in one 10.0 mg tablet?
10. Recent research suggests that the daily ingestion of 85 mg of aspirin (also known as acetylsalicylic acid, $C_9H_8O_4$) will reduce a person's risk of heart disease. How many moles of aspirin is that?

Answers

1. 480.5 g
2. 59.6 g
3. 4.42×10^6 g
4. 21.2 g
5. 2,079 g
6. 1.65 g
7. 11.63 mol
8. 1.36 mol
9. 2.77×10^{-5} mol
10. 4.7×10^{-4} mol

6.4: Mole-Mole Relationships in Chemical Reactions

Concept Review Exercise

1. How do we relate molar amounts of substances in chemical reactions?

Answer

1. Amounts of substances in chemical reactions are related by their coefficients in the balanced chemical equation.

Exercises

1. List the molar ratios you can derive from this balanced chemical equation:

$$NH_3 + 2O_2 \rightarrow HNO_3 + H_2O$$

2. List the molar ratios you can derive from this balanced chemical equation

$$2C_2H_2 + 5O_2 \rightarrow 4CO_2 + 2H_2O$$

3. Given the following balanced chemical equation,

$$6NaOH + 3Cl_2 \rightarrow NaClO_3 + 5NaCl + 3H_2O$$

how many moles of NaCl can be formed if 3.77 mol of NaOH were to react?

4. Given the following balanced chemical equation,

$$C_5H_{12} + 8O_2 \rightarrow 5CO_2 + 6H_2O$$

how many moles of H_2O can be formed if 0.0652 mol of C_5H_{12} were to react?

5. Balance the following unbalanced equation and determine how many moles of H_2O are produced when 1.65 mol of NH_3 react.

$$NH_3 + O_2 \rightarrow N_2 + H_2O$$

6. Trinitrotoluene [$C_6H_2(NO_2)_3CH_3$], also known as TNT, is formed by reacting nitric acid (HNO_3) with toluene ($C_6H_5CH_3$):

$$HNO_3 + C_6H_5CH_3 \rightarrow C_6H_2(NO_2)_3CH_3 + H_2O$$

Balance the equation and determine how many moles of TNT are produced when 4.903 mol of HNO_3 react.

7. Chemical reactions are balanced in terms of molecules and in terms of moles. Are they balanced in terms of dozens? Defend your answer.
8. Explain how a chemical reaction balanced in terms of moles satisfies the law of conservation of matter.

Answers

1. 1 mol NH_3:2 mol O_2:1 mol HNO_3:1 mol H_2O

2. 2 mol C_2H_2:5 mol O_2:4 mol CO_2:2 mol H_2O

3. 3.14 mol

4. 0.3912 mol

5. $4NH_3 + 3O_2 \rightarrow 2N_2 + 6H_2O$; 2.48 mol

6. $3HNO_3 + C_6H_5CH_3 \rightarrow C_6H_2(NO_2)_3CH_3 + 3H_2O$; 1.634 mol

7. Yes, they are still balanced.

8. A chemical reaction, balanced in terms of moles, contains the same number of atoms of each element, before and after the reaction. This means that all the atoms and its masses are conserved.

6.5: Mole-Mass and Mass-Mass Problems

Concept Review Exercises

1. What is the general sequence of conversions for a mole-mass calculation?
2. What is the general sequence of conversions for a mass-mass calculation?

Answers

1. mol first substance → mol second substance → mass second substance
2. mass first substance → mol first substance → mol second substance → mass second substance

Exercises

1. Given the following unbalanced chemical equation,

$$H_3PO_4 + NaOH \rightarrow H_2O + Na_3PO_4$$

what mass of H_2O is produced by the reaction of 2.35 mol of H_3PO_4?

2. Given the following unbalanced chemical equation,

$$C_2H_6 + Br_2 \rightarrow C_2H_4Br_2 + HBr$$

what mass of HBr is produced if 0.884 mol of C_2H_6 is reacted?

3. Certain fats are used to make soap, the first step being to react the fat with water to make glycerol (also known as glycerin) and compounds called fatty acids. One example is as follows:

$$\underset{\text{a fat}}{C_3H_5(OOC(CH_2)_{14}CH_3)_3} + 3H_2O \rightarrow \underset{\text{glycerol}}{C_3H_5(OH)_3} + 3\underset{\text{fatty acid}}{CH_3(CH_2)_{14}COOH}$$

How many moles of glycerol can be made from the reaction of 1,000.0 g of $C_3H_5(OOC(CH_2)_{14}CH_3)_3$?

4. Photosynthesis in plants leads to the general overall reaction for producing glucose ($C_6H_{12}O_6$):

$$6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$$

How many moles of glucose can be made from the reaction of 544 g of CO_2?

5. Precipitation reactions, in which a solid (called a precipitate) is a product, are commonly used to remove certain ions from solution. One such reaction is as follows:

$$Ba(NO_3)_2(aq) + Na_2SO_4(aq) \rightarrow BaSO_4(s) + 2NaNO_3(aq)$$

How many grams of Na_2SO_4 are needed to precipitate all the barium ions produced by 43.9 g of $Ba(NO_3)_2$?

6. Nitroglycerin [$C_3H_5(ONO_2)_3$] is made by reacting nitric acid (HNO_3) with glycerol [$C_3H_5(OH)_3$] according to this reaction:

$$C_3H_5(OH)_3 + 3HNO_3 \rightarrow C_3H_5(ONO_2)_3 + 3H_2O$$

If 87.4 g of HNO_3 are reacted with excess glycerol, what mass of nitroglycerin can be made?

7. Antacids are bases that neutralize acids in the digestive tract. Magnesium hydroxide [$Mg(OH)_2$] is one such antacid. It reacts with hydrochloric acid in the stomach according to the following reaction:

$$Mg(OH)_2 + 2HCl \rightarrow MgCl_2 + 2H_2O$$

How many grams of HCl can a 200 mg dose of $Mg(OH)_2$ neutralize?

8. Acid rain is caused by the reaction of nonmetal oxides with water in the atmosphere. One such reaction involves nitrogen dioxide (NO_2) and produces nitric acid (HNO_3):

$$3NO_2 + H_2O \rightarrow 2HNO_3 + NO$$

If 1.82×10^{13} g of NO_2 enter the atmosphere every year due to human activities, potentially how many grams of HNO_3 can be produced annually?

9. A simplified version of the processing of iron ore into iron metal is as follows:

$$2Fe_2O_3 + 3C \rightarrow 4Fe + 3CO_2$$

How many grams of C are needed to produce 1.00×10^9 g of Fe?

10. The *SS Hindenburg* contained about 5.33×10^5 g of H_2 gas when it burned at Lakehurst, New Jersey, in 1937. The chemical reaction is as follows:

$$2H_2 + O_2 \rightarrow 2H_2O$$

How many grams of H_2O were produced?

Answers

1. 127 g
2. 143 g
3. 1.238 mol
4. 2.06 mol
5. 23.9 g
6. 105 g
7. 0.250 g
8. 1.66×10^{13} g
9. 1.61×10^8 g
10. 4.75×10^6 g

Additional Exercises

1. If the average male has a body mass of 70 kg, of which 60% is water, how many moles of water are in an average male?
2. If the average female is 60.0 kg and contains 0.00174% iron, how many moles of iron are in an average female?
3. How many moles of each element are present in 2.67 mol of each compound?
   a. HCl
   b. H_2SO_4
   c. $Al(NO_3)_3$
   d. $Ga_2(SO_4)_3$
4. How many moles of each element are present in 0.00445 mol of each compound?
   a. HCl
   b. H_2SO_4
   c. $Al_2(CO_3)_3$
   d. $Ga_2(SO_4)_3$
5. What is the mass of one hydrogen atom in grams? What is the mass of one oxygen atom in grams? Do these masses have a 1:16 ratio, as expected?

6. What is the mass of one sodium atom in grams?
7. If 6.63×10^{-6} mol of a compound has a mass of 2.151 mg, what is the molar mass of the compound?
8. Hemoglobin (molar mass is approximately 64,000 g/mol) is the major component of red blood cells that transports oxygen and carbon dioxide in the body. How many moles are in 0.034 g of hemoglobin?

Answers

1. 2,330 mol

2. 0.0187 mol

3. a. 2.67 mol of H and 2.67 mol of Cl
   b. 5.34 mol of H, 2.67 mol of S, and 10.68 mol of O
   c. 2.67 mol of Al, 8.01 mol of N, and 24.03 mol of O
   d. 5.34 mol of Ga, 8.01 mol of S, and 32.04 mol of O

4.
   a. 0.00445 mol of H and 0.00445 mol of Cl
   b. 0.00890 mol of H, 0.00445 mol of S, and 0.0178 mol of O
   c. 0.00890 mol of Al, 0.0134 mol of C, and 0.0401 mol of O
   d. 0.00890 mol of Ga, 0.0134 mol of S, and 0.0534 mol of O

5. H = 1.68×10^{-24} g and O = 2.66×10^{-23} g; yes, they are in a 1:16 ratio.

6. 3.819×10^{-23} g

7. 324 g/mol

8. 5.3×10^{-7} mol

8.E: Quantities in Chemical Reactions (Exercise) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.E: Quantities in Chemical Reactions (Exercise)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **6.1: The Mole** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **6.2: Atomic and Molar Masses** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **6.3: Mole-Mass Conversions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **6.4: Mole-Mole Relationships in Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **6.5: Mole-Mass and Mass-Mass Problems** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.S: Quantities in Chemical Reactions (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Chemical reactions relate quantities of reactants and products. Chemists use the **mole** unit to represent 6.022×10^{23} things, whether the things are atoms of elements or molecules of compounds. This number, called **Avogadro's number**, is important because this number of atoms or molecules has the same mass in grams as one atom or molecule has in atomic mass units. **Molar masses** of substances can be determined by summing the appropriate masses from the periodic table; the final molar mass will have units of grams.

Because one mole of a substance will have a certain mass, we can use that relationship to construct conversion factors that will convert a mole amount into a mass amount, or vice versa. Such **mole-mass conversions** typically take one algebraic step.

Chemical reactions list reactants and products in molar amounts, not just molecular amounts. We can use the coefficients of a balanced chemical equation to relate moles of one substance in the reaction to moles of other substances (stoichiometry). Chemical reactions obey the Law of Conservation of Mass. To balance a chemical reaction, the coefficients in front of each compound can be adjusted until the total number of atoms of each elements is equal on both sides of the reaction arrow.

Collision Theory can be used to described the energetic aspects of chemical reactions. The reactants and products of a chemical reaction store potential energy in the form of chemical bonds and intermolecular forces. The energy difference in the bond energies of the reactants and products is called "Enthalpy". When the products are lower in potential energy than the reactants, then this excess energy is released as heat and the reaction is described as exothermic. When the products are higher in potential energy than the reactants, then energy must be added to the reaction as heat for it to occur and the reaction is described as endothermic.

The rate of a chemical reaction is influenced by its activation energy. The larger the activation energy, the slower the reaction rate. Catalysts can be added to reactions to lower the activation energy and increase the reaction rate.

Not all reactions go to completion. For some reactions, both the forward and reverse reaction can occur simultaneously. When the forward and reverse reaction rates are equal, the reaction is described as being "at equilibrium". Another way to recognize when a system is at equilibrium is when the concentration of reactants and products remain constant. Le Chatlier's principle states that the direction of an equilibrium reaction will shift to reduce the stress.

8.S: Quantities in Chemical Reactions (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **6.S: Quantities in Chemical Reactions (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

9: Solutions

Solutions are all around us. Air, for example, is a solution. If you live near a lake, a river, or an ocean, that body of water is not pure H_2O but most probably a solution. Much of what we drink—for example, soda, coffee, tea, and milk—is at least in part a solution. Solutions are a large part of everyday life. A lot of the chemistry occurring around us happens in solution. In fact, much of the chemistry that occurs in our own bodies takes place in solution, and many solutions—such as the Ringer's lactate IV solution—are important for our health. In our understanding of chemistry, we need to understand a little bit about solutions. In this chapter, you will learn about the special characteristics of solutions, how solutions are characterized, and some of their properties.

9.1: Prelude to Solutions
9.2: Solutions
9.3: Concentration
9.4: The Dissolution Process
9.5: Properties of Solutions
9.E: Solutions (Exercises)
9.S: Solutions (Summary)

Template:HideTOC

9: Solutions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

9.1: Prelude to Solutions

If you watch any of the medical dramas on television, you may have heard a doctor (actually an actor) call for an intravenous solution of “Ringer’s lactate” (or “lactated Ringer’s” or "Ri-Lac"). So what is Ringer’s lactate?

Intravenous (IV) solutions are administered for two main reasons:

1. to introduce necessary substances into the bloodstream, such as ions for proper body function, sugar and other food substances for energy, or drugs to treat a medical condition, and
2. to increase the volume of the bloodstream.

Many people with acute or long-term medical conditions have received some type of an IV solution.

One basic IV solution, called a *normal saline solution*, is simply a dilute solution of NaCl dissolved in water. Normal saline is 9.0 g of NaCl dissolved in each liter of solution. *Ringer’s lactate* is a normal saline solution that also has small amounts of potassium and calcium ions mixed in. In addition, it contains about 2.5 g of lactate ions ($C_3H_5O_3^-$) per liter of solution. The liver metabolizes lactate ions into bicarbonate (HCO_3^-) ions, which help maintain the acid-base balance of blood. Many medical problems, such as heart attacks and shock, affect the acid-base balance of blood, and the presence of lactate in the IV solution eases problems caused by this imbalance.

Physicians can select from a range of premade IV solutions, in accordance with a patient’s particular needs. Ringer’s lactate is commonly used when a patient’s blood volume must be increased quickly. Another frequently used IV solution, called D5W, is a 5% solution of dextrose (a form of sugar) in water.

9.1: Prelude to Solutions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.0: Prelude to Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.2: Solutions

Learning Objectives

- To understand what causes solutions to form.

A solution is another name for a homogeneous mixture. A *mixture* as a material composed of two or more substances. In a solution, the combination is so intimate that the different substances cannot be differentiated by sight, even with a microscope. Compare, for example, a mixture of salt and pepper and another mixture consisting of salt and water. In the first mixture, we can readily see individual grains of salt and the flecks of pepper. A mixture of salt and pepper is not a solution. However, in the second mixture, no matter how carefully we look, we cannot see two different substances. Salt dissolved in water is a solution.

The major component of a solution, called the **solvent**, is typically the same phase as the solution itself. Each minor component of a solution (and there may be more than one) is called the **solute**. In most of the solutions we will describe in this textbook, there will be no ambiguity about whether a component is the solvent or the solute. For example, in a solution of salt in water, the solute is salt, and solvent is water.

Solutions come in all phases, and the solvent and the solute do not have to be in the same phase to form a solution (such as salt and water). For example, air is a gaseous solution of about 80% nitrogen and about 20% oxygen, with some other gases present in much smaller amounts. An alloy is a solid solution consisting of a metal (like iron) with some other metals or nonmetals dissolved in it. Steel, an alloy of iron and carbon and small amounts of other metals, is an example of a solid solution. Table 9.2.1 lists some common types of solutions, with examples of each.

Table 9.2.1: Types of Solutions

Solvent Phase	Solute Phase	Example
gas	gas	air
liquid	gas	carbonated beverages
liquid	liquid	ethanol (C_2H_5OH) in H_2O (alcoholic beverages)
liquid	solid	saltwater
solid	gas	H_2 gas absorbed by Pd metal
solid	liquid	Hg(ℓ) in dental fillings
solid	solid	steel alloys

What causes a solution to form? The simple answer is that the solvent and the solute must have similar intermolecular interactions. When this is the case, the individual particles of solvent and solute can easily mix so intimately that each particle of solute is surrounded by particles of solute, forming a solution. However, if two substances have very different intermolecular interactions, large amounts of energy are required to force their individual particles to mix intimately, so a solution does not form.

This process leads to a simple rule of thumb: *like dissolves like*. Solvents that are very polar will dissolve solutes that are very polar or even ionic. Solvents that are nonpolar will dissolve nonpolar solutes. Thus water, being polar, is a good solvent for ionic compounds and polar solutes like ethanol (C_2H_5OH). However, water does not dissolve nonpolar solutes, such as many oils and greases (Figure 9.2.1).



Figure 9.2.1: A beaker holds water with blue food dye (upper liquid layer) and a much more dense perfluoroheptane (a fluorocarbon) lower liquid layer. The two fluids cannot mix and the dye cannot dissolve in fluorocarbon. A goldfish and a crab have been introduced into the water. The goldfish cannot penetrate the dense fluorocarbon. The crab floats at the liquid boundary with only parts of his legs penetrating the fluorocarbon fluid, unable to sink to the bottom of the beaker. Quarter coins rest on the bottom of the beaker. Animals were rescued from their predicament after the photo was taken. Figure used with permission from Wikipedia (Sbharris (Steven B. Harris)).

We use the word soluble to describe a solute that dissolves in a particular solvent, and the word insoluble for a solute that does not dissolve in a solvent. Thus, we say that sodium chloride is soluble in water but insoluble in hexane (C_6H_{14}). If the solute and the solvent are both liquids and soluble in any proportion, we use the word miscible, and the word immiscible if they are not.

Example 9.2.1

Water is considered a polar solvent. Which substances should dissolve in water?

1. methanol (CH_3OH)
2. sodium sulfate (Na_2SO_4)
3. octane (C_8H_{18})

Solution

Because water is polar, substances that are polar or ionic will dissolve in it.

1. Because of the OH group in methanol, we expect its molecules to be polar. Thus, we expect it to be soluble in water. As both water and methanol are liquids, the word *miscible* can be used in place of *soluble*.
2. Sodium sulfate is an ionic compound, so we expect it to be soluble in water.
3. Like other hydrocarbons, octane is nonpolar, so we expect that it would not be soluble in water.

Exercise 9.2.1

Toluene ($C_6H_5CH_3$) is widely used in industry as a nonpolar solvent. Which substances should dissolve in toluene?

a. water (H_2O)
b. sodium sulfate (Na_2SO_4)
c. octane (C_8H_{18})

Answer:

Octane only.

Concept Review Exercises

1. What causes a solution to form?
2. How does the phrase *like dissolves like* relate to solutions?

Answers

1. Solutions form because a solute and a solvent have similar intermolecular interactions.
2. It means that substances with similar intermolecular interactions will dissolve in each other.

Key Takeaway

- Solutions form because a solute and a solvent experience similar intermolecular interactions.

Exercises

1. Define *solution*.
2. Give several examples of solutions.
3. What is the difference between a solvent and a solute?
4. Can a solution have more than one solute in it? Can you give an example?
5. Does a solution have to be a liquid? Give several examples to support your answer.
6. Give at least two examples of solutions found in the human body.
7. Which substances will probably be soluble in water, a very polar solvent?
   a. sodium nitrate ($NaNO_3$)
   b. hexane (C_6H_{14})
   c. isopropyl alcohol [$(CH_3)_2CHOH$]
   d. benzene (C_6H_6)
8. Which substances will probably be soluble in toluene ($C_6H_5CH_3$), a nonpolar solvent?
   a. sodium nitrate ($NaNO_3$)
   b. hexane (C_6H_{14})
   c. isopropyl alcohol [$(CH_3)_2CHOH$]
   d. benzene (C_6H_6)
9. The solubility of alcohols in water varies with the length of carbon chain. For example, ethanol (CH_3CH_2OH) is soluble in water in any ratio, while only 0.0008 mL of heptanol ($CH_3CH_2CH_2CH_2CH_2CH_2CH_2OH$) will dissolve in 100 mL of water. Propose an explanation for this behavior.
10. Dimethyl sulfoxide [$(CH_3)_2SO$] is a polar liquid. Based on the information in Exercise 9, which do you think will be more soluble in it—ethanol or heptanol?

Answers

1. a homogeneous mixture
2. vinegar, dextrose IV, saline IV, coffee, tea, wine
3. A solvent is the majority component of a solution; a solute is the minority component of a solution.
4. yes. Coke or Pepsi has sugar, caffeine and carbon dioxide as solutes.
5. A solution does not have to be liquid; air is a gaseous solution, while some alloys are solid solutions (answers will vary).
6. Urine, plasma
7. a. probably soluble
   b. probably not soluble
   c. probably soluble
   d. probably not soluble
8. a. probably not soluble
   b. probably soluble
   c. probably not soluble
   d. probably soluble

9. Small alcohol molecules have strong polar intermolecular interactions, so they dissolve in water. In large alcohol molecules, the nonpolar end overwhelms the polar end, so they do not dissolve very well in water.

10. Ethanol is a smaller molecule. It will be more soluble in water than heptanol.

9.2: Solutions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

9.3: Concentration

Learning Objectives

- Calculate percentage concentration (m/m, v/v, m/v), ppm and ppb.
- Calculate the molarity of a solution.
- Use concentration units to calculate the amount of solute in a solution.
- Use molarity to determine quantities in chemical reactions.
- Determine the resulting concentration of a diluted solution.

To define a solution precisely, we need to state its concentration: how much solute is dissolved in a certain amount of solvent. Words such as *dilute* or *concentrated* are used to describe solutions that have a little or a lot of dissolved solute, respectively, but these are relative terms whose meanings depend on various factors.

Solubility

There is usually a limit to how much solute will dissolve in a given amount of solvent. This limit is called the **solubility** of the solute. Some solutes have a very small solubility, while other solutes are soluble in all proportions. Table 9.3.1 lists the solubilities of various solutes in water. Solubilities vary with temperature, so Table 9.3.1 includes the temperature at which the solubility was determined.

Table 9.3.1: Solubilities of Various Solutes in Water at 25°C (Except as Noted)

Substance	Solubility (g in 100 mL of H2O)
AgCl(s)	0.019
$C_6H_6(\ell)$ (benzene)	0.178
$CH_4(g)$	0.0023
$CO_2(g)$	0.150
$CaCO_3(s)$	0.058
$CaF_2(s)$	0.0016
$Ca(NO_3)_2(s)$	143.9
$C_6H_{12}O_6$ (glucose)	120.3 (at 30°C)
KBr(s)	67.8
$MgCO_3(s)$	2.20
NaCl(s)	36.0
$NaHCO_3(s)$	8.41
$C_{12}H_{22}O_{11}$ (sucrose)	204.0 (at 20°C)

If a solution contains so much solute that its solubility limit is reached, the solution is said to be **saturated**, and its concentration is known from information contained in Table 9.3.1. If a solution contains less solute than the solubility limit, it is **unsaturated**. Under special circumstances, more solute can be dissolved even after the normal solubility limit is reached; such solutions are called ***supersaturated*** and are not stable. If the solute is solid, excess solute can easily recrystallize. If the solute is a gas, it can bubble out of solution uncontrollably, like what happens when you shake a soda can and then immediately open it.

Precipitation from Supersaturated Solutions

Recrystallization of excess solute from a supersaturated solution usually gives off energy as heat. Commercial heat packs containing supersaturated sodium acetate ($NaC_2H_3O_2$) take advantage of this phenomenon. You can probably find them at your local drugstore.



Video* 9.3.1*: Watered-down sodium acetate trihydrate. Needle crystal is truly wonderful structures

Most solutions we encounter are unsaturated, so knowing the solubility of the solute does not accurately express the amount of solute in these solutions. There are several common ways of specifying the concentration of a solution.

Percent Composition

There are several ways of expressing the concentration of a solution by using a percentage. The **mass/mass percent** (% m/m) is defined as the mass of a solute divided by the mass of a solution times 100:

$$\%\ \mathrm{m/m} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 100\%$$

If you can measure the masses of the solute and the solution, determining the mass/mass percent is easy. Each mass must be expressed in the same units to determine the proper concentration.

✔ Example 9.3.1

A saline solution with a mass of 355 g has 36.5 g of NaCl dissolved in it. What is the mass/mass percent concentration of the solution?

Solution

We can substitute the quantities given in the equation for mass/mass percent:

$$\%\ \mathrm{m/m} = \frac{36.5\ \mathrm{g}}{355\ \mathrm{g}} \times 100\% = 10.3\%$$

? Exercise 9.3.1

A dextrose (also called D-glucose, $C_6H_{12}O_6$) solution with a mass of 2.00×10^2 g has 15.8 g of dextrose dissolved in it. What is the mass/mass percent concentration of the solution?

Answer

7.90%

For gases and liquids, volumes are relatively easy to measure, so the concentration of a liquid or a gas solution can be expressed as a **volume/volume percent** (% v/v): the volume of a solute divided by the volume of a solution times 100:

$$\%\ \mathrm{v/v} = \frac{\text{volume of solute}}{\text{volume of solution}} \times 100\%$$

Again, the units of the solute and the solution must be the same. A hybrid concentration unit, **mass/volume percent** (% m/v), is commonly used for intravenous (IV) fluids (Figure 9.3.1). It is defined as the mass in grams of a solute, divided by volume in milliliters of solution times 100:

$$\%\ \text{m/v} = \frac{\text{mass of solute (g)}}{\text{volume of solution (mL)}} \times 100\%$$



Figure 9.3.1: Mass/Volume Percent.The 0.9% NaCl concentration on this IV bag is mass/volume percent (left). Such solution is used for other purposes and available in bottles (right). Figures used with permission from Wikipedia

Using Percent Composition in Calculations

The **percent concentration** can be used to produce a **conversion factor** between the **amount of solute** and the **amount of solution**. As such, concentrations can be useful in a variety of stoichiometry problems as discussed in Chapter 6. In many cases, it is best to use the original definition of the concentration unit; it is that definition that provides the conversion factor.

As an example, if the given concentration is **5% v/v solution of alcohol**, this means that there are 5 mL of alcohol dissolved in every 100 mL solution.

5 mL alcohol = 100 mL solution

The two possible conversion factors are written as follows:

$$\frac{\textbf{5 mL alcohol}}{\textbf{100 mL solution}} \text{ or } \frac{\textbf{100 mL solution}}{\textbf{5 mL alcohol}}$$

Use the first conversion factor to convert from a given amount of solution to amount of solute. The second conversion factor is used to convert from a given amount of solute to amount of solution. Given any two quantities in any percent composition, the third quantity can be calculated, as the following example illustrates.

Example 9.3.2

A sample of 45.0% v/v solution of ethanol (C_2H_5OH) in water has a volume of 115 mL. What volume of ethanol solute does the sample contain?

Solution

A percentage concentration is simply the number of parts of solute per 100 parts of solution. Thus, the percent concentration of 45.0% v/v implies the following:

$$45.0\%\ \text{v/v} \rightarrow \frac{45\ \text{mL}\ C_2H_5OH}{100\ \text{mL solution}}$$

That is, there are 45 mL of C_2H_5OH for every 100 mL of solution. We can use this fraction as a **conversion factor** to determine the amount of C_2H_5OH in 115 mL of solution:

$$115\ \text{mL solution} \times \frac{45\ \text{mL}\ C_2H_5OH}{100\ \text{mL solution}} = 51.8\ \text{mL}\ C_2H_5OH$$

Exercise 9.3.2

What volume of a 12.75% m/v solution of glucose ($C_6H_{12}O_6$) in water is needed to obtain 50.0 g of $C_6H_{12}O_6$?

Answer

$$50.0 \text{ g } C_6H_12O_6 \times \frac{100 \text{ mL solution}}{12.75 \text{ g } C_6H_12O_6} = 392 \text{ mL solution}$$

Example 9.3.3

A normal saline IV solution contains 9.0 g of NaCl in every liter of solution. What is the mass/volume percent of normal saline?

Solution

We can use the definition of mass/volume percent, but first we have to express the volume in milliliter units:

$$1 \text{ L} = 1{,}000 \text{ mL}$$

Because this is an exact relationship, it does not affect the significant figures of our result.

$$\%\text{ m/v} = \frac{9.0 \text{ g NaCl}}{1,000 \text{ mL solution}} \times 100\% = 0.90\%\text{ m/v}$$

Exercise 9.3.3

The chlorine bleach that you might find in your laundry room is typically composed of 27.0 g of sodium hypochlorite (NaOCl), dissolved to make 500.0 mL of solution. What is the mass/volume percent of the bleach?

Answer

$$\%\text{ m/v} = \frac{27.0 \text{ g NaOCl}}{500.0 \text{ mL solution}} \times 100\% = 5.40\%\text{ m/v}$$

In addition to percentage units, the units for expressing the concentration of extremely dilute solutions are parts per million (ppm) and parts per billion (ppb). Both of these units are mass based and are defined as follows:

$$\text{ppm} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 1,000,000$$

$$\text{ppb} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 1,000,000,000$$

> *Similar to parts per million and parts per billion, related units include parts per thousand (ppth) and parts per trillion (ppt).*

Concentrations of *trace elements* in the body—elements that are present in extremely low concentrations but are nonetheless necessary for life—are commonly expressed in parts per million or parts per billion. Concentrations of poisons and pollutants are also described in these units. For example, cobalt is present in the body at a concentration of 21 ppb, while the State of Oregon's Department of Agriculture limits the concentration of arsenic in fertilizers to 9 ppm.

> *In aqueous solutions, 1 ppm is essentially equal to 1 mg/L, and 1 ppb is equivalent to 1 µg/L.*

Example 9.3.4

If the concentration of cobalt in a human body is 21 ppb, what mass in grams of Co is present in a body having a mass of 70.0 kg?

Solution

A concentration of 21 ppb means "21 g of solute per 1,000,000,000 g of solution." Written as a **conversion factor,** this concentration of Co is as follows:

$$21 \text{ ppb Co} \rightarrow \frac{21 \text{ g Co}}{1,000,000,000 \text{ g solution}}$$

We can use this as a conversion factor, but first we must convert 70.0 kg to gram units:

$$70.0 \text{ kg} \times \frac{1,000 \text{ g}}{1 \text{ kg}} = 7.00 \times 10^4 \text{ g}$$

Now we determine the amount of Co:

$$7.00 \times 10^4 \text{ g solution} \times \frac{21 \text{ g Co}}{1,000,000,000 \text{ g solution}} = 0.0015 \text{ g Co}$$

This is only 1.5 mg.

? Exercise 9.3.4

An 85 kg body contains 0.012 g of Ni. What is the concentration of Ni in parts per million?

Answer

0.14 ppm

Molarity

Another way of expressing concentration is to give the number of moles of solute per unit volume of solution. Such concentration units are useful for discussing chemical reactions in which a solute is a product or a reactant. Molar mass can then be used as a conversion factor to convert amounts in moles to amounts in grams.

Molarity is defined as the number of moles of a solute dissolved per liter of solution:

$$\text{molarity} = \frac{\text{number of moles of solute}}{\text{number of liters of solution}}$$


Molarity is abbreviated M (often referred to as "molar"), and the units are often abbreviated as mol/L. It is important to remember that "mol" in this expression refers to moles of solute and that "L" refers to liters of solution. For example, if you have 1.5 mol of NaCl dissolved in 0.500 L of solution, its molarity is therefore

$$\frac{1.5 \text{ mol NaCl}}{0.500 \text{ L solution}} = 3.0 \text{ M NaCl}$$

which is read as "three point oh **molar** sodium chloride." Sometimes (aq) is added when the solvent is water, as in "3.0 M NaCl(aq)."

Before a molarity concentration can be calculated, the amount of the **solute must be expressed in moles**, and the **volume of the solution** must be expressed in **liters**.

If the quantity of the solute is given in mass units, you must convert mass units to mole units before using the definition of molarity to calculate concentration. For example, what is the molar concentration of a solution of 22.4 g of HCl dissolved in 1.56 L?



Step 1: convert the **mass of solute** to **moles** using the **molar mass of HCl (36.46 g/mol):**

$$22.4\ g\cancel{HCl} \times \frac{1\ mol\ HCl}{36.46\ g\cancel{HCl}} = 0.614\ mol\ HCl$$

Step 2: use the **definition of molarity** to determine the concentration:

$$M\ =\ \frac{0.614\ mol\ HCl}{1.56 L\ solution} = 0.394\ MHCl$$

Example 9.3.5

What is the molarity of an aqueous solution of 25.0 g of NaOH in 750 mL?

Solution

Before we substitute these quantities into the definition of molarity, we must convert them to the proper units. The mass of NaOH must be converted to moles of NaOH. The molar mass of NaOH is 40.00 g/mol:

$$25.0\ \text{g NaOH} \times \frac{1\ \text{mol NaOH}}{40.00\ \text{g NaOH}} = 0.625\ \text{mol NaOH}$$

Next, we convert the volume units from milliliters to liters:

$$750\ \text{mL} \times \frac{1\ \text{L}}{1,000\ \text{mL}} = 0.750\ \text{L}$$

Now that the quantities are expressed in the proper units, we can substitute them into the definition of molarity:

$$\text{M} = \frac{0.625\ \text{mol NaOH}}{0.750\ \text{L}} = 0.833\ \text{M NaOH}$$

Exercise 9.3.5

If a 350 mL cup of coffee contains 0.150 g of caffeine ($C_8H_{10}N_4O_2$), what is the molarity of this caffeine solution?

Answer

0.00221 M

Using Molarity in Calculations

The definition of molarity can also be used to calculate a needed volume of solution, given its concentration and the number of moles desired, or the number of moles of solute (and subsequently, the mass of the solute), given its concentration and volume. As in the percent concentration, molarity can also be expressed as a **conversion factor**.


Molarity is defined as moles solute per liter solution. There is an understood 1 in the denominator of the conversion factor. For example, a **3.0 M solution of sucrose** means that there are three moles of sucrose dissolved in every **liter of solution**. Mathematically, this is stated as follows:

3.0 moles sucrose = 1 L solution

Dividing both sides of this expression by either side, we generate two possible conversion factors:

$$\frac{\textbf{3.0 mol sucrose}}{\textbf{1 L solution}} \text{ or } \frac{\textbf{1 L solution}}{\textbf{3.0 mol sucrose}}$$

The first conversion factor can be used to convert from *volume (L) of solution to moles solute*, and the *second converts from moles of solute to volume (L) of solution*.



For example, suppose we are asked how many moles of sucrose are present in 0.108 L of a 3.0 M sucrose solution. The given volume (0.108 L) is multiplied by the first conversion factor to cancel the L units, and find that 0.32 moles of sucrose are present.

$$0.108\ L\ \cancel{solution} \times \frac{3.0\ mol\ sucrose}{1 L\ \cancel{solution}} = 0.32\ mol\ sucrose$$

How many liters of 3.0 M sucrose solution are needed to obtain 4.88 mol of sucrose? In such a conversion, we multiply the given (4.88 moles sucrose) with the second conversion factor. This cancels the moles units and converts it to liters of solution.

$$4.88\ mol\ \cancel{sucrose} \times \frac{1\ L\ solution}{3.0\ mol\ \cancel{sucrose}} = 1.63\ L\ solution$$

Example 9.3.6

1. What volume of a 0.0753 M solution of dimethylamine [$(CH_3)_2NH$] is needed to obtain 0.450 mol of the compound?
2. Ethylene glycol ($C_2H_6O_2$) is mixed with water to make auto engine coolants. How many grams of $C_2H_6O_2$ are in 5.00 L of a 6.00 M aqueous solution?

Solution

1. To solve for the volume, multiply the "given" (0.450 mol of dimethylamine) with the molarity conversion factor (0.0753 M). Use the proper conversion factor to cancel the unit "mol" and get the unit volume (L) of solution:

$$0.450\ \text{mol dimethylamine} \times \frac{1\ \text{L solution}}{0.0753\ \text{mol dimethylamine}} = 5.98\ \text{L solution}$$

2. The strategy in solving this problem is to convert the given volume (5.00 L) using the 6.00 M (conversion factor) to solve for moles of ethylene glycol, which can then be converted to grams.


Step 1: Convert the given volume (5.00 L) to moles ethylene glycol.

$$5.00\ \text{L solution} \times \frac{6.00\ \text{mol}\ C_2H_6O_2}{1\ \text{L solution}} = 30.0\ \text{mol}\ C_2H_6O_2$$

Step 2: Convert 30.0 mols $C_2H_6O_2$ to grams $C_2H_6O_2$. Molar mass of $C_2H_6O_2$= 62.08 g/mol

$$30.0\ \text{mol}\ C_2H_6O_2 \times \frac{62.08\ \text{g}\ C_2H_6O_2}{1\ \text{mol}\ C_2H_6O_2} = 1,860\ \text{g}\ C_2H_6O_2$$

The same two-step problem can also be worked out in a single line, rather than as two separate steps, as follows:



$$5.00\ \text{L solution} \times \frac{6.00\ \text{mol}\ C_2H_6O_2}{1\ \text{L solution}} \times \frac{62.08\ \text{g}\ C_2H_6O_2}{1\ \text{mol}\ C_2H_6O_2} = 1,860\ \text{g}\ C_2H_6O_2$$

The final answer is rounded off to 3 significant figures. Thus, there are 1,860 g of $C_2H_6O_2$ in the specified amount of engine coolant.

Note: Dimethylamine has a “fishy” odor. In fact, organic compounds called amines cause the odor of decaying fish.

Example 9.3.6A

1. What volume of a 0.0753 M solution of dimethylamine [$(CH_3)_2NH$] is needed to obtain 0.450 mol of the compound?
2. Ethylene glycol ($C_2H_6O_2$) is mixed with water to make auto engine coolants. How many grams of $C_2H_6O_2$ are in 5.00 L of a 6.00 M aqueous solution?

Solution

This is an alternative method in case you don't want to use the conversion factor for molarity. In both parts, we will use the definition of molarity to solve for the desired quantity.

1. $0.0753\ \text{M} = \dfrac{0.450\ \text{mol}\ (\text{CH}_3)_2\text{NH}}{\text{volume of solution}}$

   To solve for the volume of solution, we multiply both sides by volume of solution and divide both sides by the molarity value to isolate the volume of solution on one side of the equation:

   $$\text{volume of solution} = \frac{0.450\ \text{mol}\ (\text{CH}_3)_2\text{NH}}{0.0753\ \text{M}} = 5.98\ \text{L}$$

   Note that because the definition of molarity is mol/L, the division of mol by M yields L, a unit of volume.

2. The molar mass of $C_2H_6O_2$ is 62.08 g/mol., so

   $$6.00\ \text{M} = \frac{\text{moles of solute}}{5.00\ \text{L}}$$

   To solve for the number of moles of solute, we multiply both sides by the volume:

   $$\text{moles of solute} = (6.00\ \text{M})(5.00\ \text{L}) = 30.0\ \text{mol}$$

   Note that because the definition of molarity is mol/L, the product M × L gives mol, a unit of amount. Now, using the molar mass of $C_3H_8O_3$, we convert mol to g:

   $$30.0\ \text{mol} \times \frac{62.08\ \text{g}}{\text{mol}} = 1,860\ \text{g}$$

   Thus, there are 1,860 g of $C_2H_6O_2$ in the specified amount of engine coolant.

? Exercise 9.3.6

a. What volume of a 0.0902 M solution of formic acid (HCOOH) is needed to obtain 0.888 mol of HCOOH?
b. Acetic acid ($HC_2H_3O_2$) is the acid in vinegar. How many grams of $HC_2H_3O_2$ are in 0.565 L of a 0.955 M solution?

Answer

a. 9.84 L

b. 32.4 g

Using Molarity in Stoichiometry Problems

Of all the ways of expressing concentration, molarity is the one most commonly used in stoichiometry problems because it is directly related to the mole unit. Consider the following chemical equation:

$$\text{HCl(aq)} + \text{NaOH(s)} \rightarrow \text{H}_2\text{O}(\ell) + \text{NaCl(aq)}$$

Suppose we want to know how many liters of aqueous HCl solution will react with a given mass of NaOH. A typical approach to answering this question is as follows:




***Figure* 9.3.2** *: Typical approach to solving Molarity problems*

In itself, each step is a straightforward conversion. It is the combination of the steps that is a powerful quantitative tool for problem solving.

Example 9.3.7

How many milliliters of a 2.75 M HCl solution are needed to react with 185 g of NaOH? The balanced chemical equation for this reaction is as follows:

$$\mathrm{HCl(aq) + NaOH(s) \rightarrow H_2O(\ell) + NaCl(aq)}$$

Solution

We will follow the flowchart to answer this question. First, we convert the mass of NaOH to moles of NaOH using its molar mass, 40.00 g/mol:

$$185\ \text{g NaOH} \times \frac{1\ \text{mol NaOH}}{40.00\ \text{g NaOH}} = 4.63\ \text{mol NaOH}$$

Using the balanced chemical equation, we see that there is a one-to-one ratio of moles of HCl to moles of NaOH. We use this to determine the number of moles of HCl needed to react with the given amount of NaOH:

$$4.63\ \text{mol NaOH} \times \frac{1\ \text{mol HCl}}{1\ \text{mol NaOH}} = 4.63\ \text{mol HCl}$$


Finally, we use the definition of molarity to determine the volume of 2.75 M HCl needed:

$$2.75\ \text{M HCl} = \frac{4.63\ \text{mol HCl}}{\text{volume of HCl solution}}$$

$$\text{volume of HCl} = \frac{4.63\ \text{mol HCl}}{2.75\ \text{M HCl}} = 1.68\ \text{L} \times \frac{1,000\ \text{mL}}{1\ \text{L}} = 1,680\ \text{mL}$$

We need 1,680 mL of 2.75 M HCl to react with the NaOH.

The same multi-step problem can also be worked out in a single line, rather than as separate steps, as follows:



$$185 \text{ g NaOH} \times \frac{1 \text{ mol NaOH}}{40.00 \text{ g NaOH}} \times \frac{1 \text{ mol HCl}}{1 \text{ mol NaOH}} \times \frac{1 \text{ L HCl solution}}{2.75 \text{ mol HCl}} \times \frac{1000 \text{ mL HCl solution}}{1 \text{ L HCl solution}} = 1,680 \text{ mL HCl solution}$$

Our final answer (rounded off to three significant figures) is 1,680 mL HCl solution.

Exercise 9.3.7

How many milliliters of a 1.04 M H_2SO_4 solution are needed to react with 98.5 g of $Ca(OH)_2$? The balanced chemical equation for the reaction is as follows:

$$H_2SO_{4(aq)} + Ca(OH)_{2(s)} \rightarrow 2H_2O_{(\ell)} + CaSO_{4(aq)}$$

Answer

1,280 mL

The general steps for performing stoichiometry problems such as this are shown in Figure 9.3.3. You may want to consult this figure when working with solutions in chemical reactions. The double arrows in Figure 9.3.3 indicate that you can start at either end of the chart and, after a series of simple conversions, determine the quantity at the other end.




Figure 9.3.3: Diagram of Steps for Using Molarity in Stoichiometry Calculations. When using molarity in stoichiometry calculations, a specific sequence of steps usually leads you to the correct answer.

Many of the fluids found in our bodies are solutions. The solutes range from simple ionic compounds to complex proteins. Table 9.3.2 lists the typical concentrations of some of these solutes.

Table 9.3.2: Approximate Concentrations of Various Solutes in Some Solutions in the Body*

Solution	Solute	Concentration (M)
blood plasma	Na^+	0.138
	K^+	0.005
	Ca^{2+}	0.004
	Mg^{2+}	0.003
	Cl^-	0.110
	HCO_3^-	0.030
stomach acid	HCl	0.10
urine	NaCl	0.15
	PO_4^{3-}	0.05
	NH_2CONH_2 (urea)	0.30

*Note: Concentrations are approximate and can vary widely.

Looking Closer: The Dose Makes the Poison

Why is it that we can drink 1 qt of water when we are thirsty and not be harmed, but if we ingest 0.5 g of arsenic, we might die? There is an old saying: *the dose makes the poison*. This means that what may be dangerous in some amounts may not be dangerous in other amounts.

Take arsenic, for example. Some studies show that arsenic deprivation limits the growth of animals such as chickens, goats, and pigs, suggesting that arsenic is actually an essential trace element in the diet. Humans are constantly exposed to tiny amounts of arsenic from the environment, so studies of completely arsenic-free humans are not available; if arsenic is an essential trace mineral in human diets, it is probably required on the order of 50 ppb or less. A toxic dose of arsenic corresponds to about 7,000 ppb and higher, which is over 140 times the trace amount that may be required by the body. Thus, arsenic is not poisonous in and of itself. Rather, it is the amount that is dangerous: the dose makes the poison.

Similarly, as much as water is needed to keep us alive, too much of it is also risky to our health. Drinking too much water too fast can lead to a condition called water intoxication, which may be fatal. The danger in water intoxication is not that water itself becomes toxic. It is that the ingestion of too much water too fast dilutes sodium ions, potassium ions, and other salts in the bloodstream to concentrations that are not high enough to support brain, muscle, and heart functions. Military personnel, endurance athletes, and even desert hikers are susceptible to water intoxication if they drink water but do not replenish the salts lost in sweat. As this example shows, even the right substances in the wrong amounts can be dangerous!

Equivalents

Concentrations of ionic solutes are occasionally expressed in units called equivalents (Eq). One equivalent equals 1 mol of positive or negative charge. Thus, 1 mol/L of Na^+(aq) is also 1 Eq/L because sodium has a 1+ charge. A 1 mol/L solution of Ca^{2+}(aq) ions has a concentration of 2 Eq/L because calcium has a 2+ charge. Dilute solutions may be expressed in milliequivalents (mEq)—for example, human blood plasma has a total concentration of about 150 mEq/L. (For more information about the ions present in blood plasma, see Chapter 3, Section 3.3.)

Dilutions

When additional water is added to an aqueous solution, the concentration of that solution decreases. This is because the number of **moles of the solute does not change**, while the **volume of the solution increases**. We can set up an equality between the moles of the solute before the dilution (1) and the moles of the solute after the dilution (2).

$$\text{mol}_1 = \text{mol}_2$$

Since the moles of solute in a solution is equal to the molarity multiplied by the liters, we can set those equal.

$$M_1 \times L_1 = M_2 \times L_2$$

Finally, because the two sides of the equation are set equal to one another, the volume can be in any units we choose, as long as that unit is the same on both sides. Our equation for calculating the molarity of a diluted solution becomes:

$$M_1 \times V_1 = M_2 \times V_2$$

Suppose that you have $100.\ \text{mL}$ of a $2.0\ \text{M}$ solution of HCl. You dilute the solution by adding enough water to make the solution volume $500.\ \text{mL}$. The new molarity can easily be calculated by using the above equation and solving for M_2.

$$M_2 = \frac{M_1 \times V_1}{V_2} = \frac{2.0\ \text{M} \times 100.\ \text{mL}}{500.\ \text{mL}} = 0.40\ \text{M HCl}$$

The solution has been diluted by one-fifth since the new volume is five times as great as the original volume. Consequently, the molarity is one-fifth of its original value.




Figure 9.3.1: Figure 9.3.4: Diluting a solution by adding more solvent. Through the dilution process, the volume increases and the concentrations of the solutes decrease. As a consequence, there is more solution in the beaker after dilution, and the color of the solution becomes fainter. The inset shows that at the particular level (picometer to nanometer scale), there are less solute particles (red triangles) per volume. The content in the central box shows the dilution law, i.e. that the product of concentration and volume stays constant upon dilution. CC BY-SA 4.0.

Dilution Equations

Any units of concentration and volume can be used, as long as both concentrations and both volumes have the same unit. For example, if we are using M (molarity), then we can express the equation as follows:

$$\text{Molarity}_{\text{initial}} \times \text{volume}_{\text{initial}} = \text{Molarity}_{\text{final}} \times \text{volume}_{\text{final}} \text{ or}$$

$$\boldsymbol{M_1V_1 = M_2V_2}$$

If we are using percent, the dilution equation is as follows:

$$\%_{\text{initial}} \times \text{volume}_{\text{initial}} = \%_{\text{final}} \times \text{volume}_{\text{final}} \text{ or}$$

$$\boldsymbol{\%_1V_1 = \%_2V_2}$$

Example 9.3.8

A 125 mL sample of 0.900 M NaCl is diluted to 1,125 mL. What is the final concentration of the diluted solution?

Solution

Because the volume units are the same, and we are looking for the molarity of the final solution, we can use (concentration × volume)$_{\text{initial}}$ = (concentration × volume)$_{\text{final}}$:

$$(0.900\ \text{M} \times 125\ \text{mL}) = (\text{concentration} \times 1{,}125\ \text{mL})$$

We solve by isolating the unknown concentration by itself on one side of the equation. Dividing by 1,125 mL gives

$$\text{concentration} = \frac{0.900\ \text{M} \times 125\ \text{mL}}{1{,}125\ \text{mL}} = 0.100\ \text{M}$$

as the final concentration.

Exercise 9.3.8

a. A nurse uses a syringe to inject 5.00 mL of 0.550 M heparin solution (heparin is an anticoagulant drug) into a 250 mL IV bag, for a final volume of 255 mL. What is the concentration of the resulting heparin solution?

b. A 0.885 M solution of KBr whose initial volume is 76.5 mL has more water added until its concentration is 0.500 M. What is the new volume of the solution?

Answer

a. 0.0108 M

b. 135.4 mL

Preparing IV Solutions

In a hospital emergency room, a physician orders an intravenous (IV) delivery of 100 mL of 0.5% KCl for a patient suffering from hypokalemia (low potassium levels). Does an aide run to a supply cabinet and take out an IV bag containing this concentration of KCl?

Not likely. It is more probable that the aide must make the proper solution from an IV bag of sterile solution and a more concentrated, sterile solution, called a *stock solution*, of KCl. The aide is expected to use a syringe to draw up some stock solution and inject it into the waiting IV bag and dilute it to the proper concentration. Thus the aide must perform a dilution calculation.



Figure 9.3.5 *Preparing IV Solution © Thinkstock. Medical personnel commonly must perform dilutions for IV solutions.*

If the stock solution is 10.0% KCl and the final volume and concentration need to be 100 mL and 0.50%, respectively, then it is an easy calculation to determine how much stock solution to use:

$$(10\%)V_1 = (0.50\%)(100\text{ mL})$$

$$V_1 = 5\text{ mL}$$

Of course, the addition of the stock solution affects the total volume of the diluted solution, but the final concentration is likely close enough even for medical purposes.

Medical and pharmaceutical personnel are constantly dealing with dosages that require concentration measurements and dilutions. It is an important responsibility: calculating the *wrong* dose can be useless, harmful, or even fatal!

Key Takeaways

- Various concentration units are used to express the amounts of solute in a solution.
- Concentration units can be used as conversion factors in stoichiometry problems.
- New concentrations can be easily calculated if a solution is diluted.

9.3: Concentration is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.2: Concentration** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.4: The Dissolution Process

Learning Objectives

- To describe the dissolution process at the molecular level

The Dissolution Process

What occurs at the molecular level to cause a solute to dissolve in a solvent? The answer depends in part on the solute, but there are some similarities common to all solutes.

Recall the rule that *like dissolves like*. This means that substances must have similar intermolecular forces to form solutions. When a soluble solute is introduced into a solvent, the particles of solute can interact with the particles of solvent. In the case of a solid or liquid solute, the interactions between the solute particles and the solvent particles are so strong that the individual solute particles separate from each other and, surrounded by solvent molecules, enter the solution. (Gaseous solutes already have their constituent particles separated, but the concept of being surrounded by solvent particles still applies.) This process is called **solvation** and is illustrated in Figure 9.4.1. When the solvent is water, the word **hydration**, rather than solvation, is used.




Figure 9.4.1: Solvation. When a solute dissolves, the individual particles of solute become surrounded by solvent particles. Eventually the particle detaches from the remaining solute, surrounded by solvent molecules in solution. Source: Photo © Thinkstock

Ionic Compounds and Covalent Compounds as Solutes

In the case of molecular solutes like glucose, the solute particles are individual molecules. However, if the solute is ionic, the individual ions separate from each other and become surrounded by solvent particles. That is, the cations and anions of an ionic solute separate when the solute dissolves. This process is referred to as **dissociation** (Figure 9.4.1).

The dissociation of soluble ionic compounds gives solutions of these compounds an interesting property: they conduct electricity. Because of this property, soluble ionic compounds are referred to as electrolytes. Many ionic compounds dissociate completely and are therefore called **strong electrolytes**. Sodium chloride is an example of a strong electrolyte. Some compounds dissolve but dissociate only partially, and solutions of such solutes may conduct electricity only weakly. These solutes are called **weak electrolytes**. Acetic acid (CH_3COOH), the compound in vinegar, is a weak electrolyte. Solutes that dissolve into individual neutral molecules without dissociation do not impart additional electrical conductivity to their solutions and are called nonelectrolytes. Table sugar ($C_{12}H_{22}O_{11}$) is an example of a nonelectrolyte.

> *The term electrolyte is used in medicine to mean any of the important ions that are dissolved in aqueous solution in the body. Important physiological electrolytes include Na^+, K^+, Ca^{2+}, Mg^{2+}, and Cl^-.*

Example 9.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

1. potassium chloride (KCl)
2. fructose ($C_6H_{12}O_6$)

3. isopropyl alcohol [$CH_3CH(OH)CH_3$]
4. magnesium hydroxide [$Mg(OH)_2$]

Solution

Each substance can be classified as an ionic solute or a nonionic solute. Ionic solutes are electrolytes, and nonionic solutes are nonelectrolytes.

1. Potassium chloride is an ionic compound; therefore, when it dissolves, its ions separate, making it an electrolyte.
2. Fructose is a sugar similar to glucose. (In fact, it has the same molecular formula as glucose.) Because it is a molecular compound, we expect it to be a nonelectrolyte.
3. Isopropyl alcohol is an organic molecule containing the alcohol functional group. The bonding in the compound is all covalent, so when isopropyl alcohol dissolves, it separates into individual molecules but not ions. Thus, it is a nonelectrolyte
4. Magnesium hydroxide is an ionic compound, so when it dissolves it dissociates. Thus, magnesium hydroxide is an electrolyte.

? Exercise 9.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

a. acetone (CH_3COCH_3)
b. iron(III) nitrate [$Fe(NO_3)_3$]
c. elemental bromine (Br_2)
d. sodium hydroxide (NaOH)

Answer

a. nonelectrolyte

b. electrolyte

c. nonelectrolyte

d. electrolyte

Electrolytes in Body Fluids

Our body fluids are solutions of electrolytes and many other things. The combination of blood and the circulatory system is the *river of life*, because it coordinates all the life functions. When the heart stops pumping in a heart attack, the life ends quickly. Getting the heart restarted as soon as one can is crucial in order to maintain life.

The primary electrolytes required in the body fluid are cations (of calcium, potassium, sodium, and magnesium) and anions (of chloride, carbonates, aminoacetates, phosphates, and iodide). These are nutritionally called **macrominerals**.

Electrolyte balance is crucial to many body functions. Here's some extreme examples of what can happen with an imbalance of electrolytes: elevated potassium levels may result in cardiac arrhythmias; decreased extracellular potassium produces paralysis; excessive extracellular sodium causes fluid retention; and decreased plasma calcium and magnesium can produce muscle spasms of the extremities.

When a patient is dehydrated, a carefully prepared (commercially available) electrolyte solution is required to maintain health and well being. In terms of child health, oral electrolyte is given when a child is dehydrated due to diarrhea. The use of oral electrolyte maintenance solutions, which is responsible for saving millions of lives worldwide over the last 25 years, is one of the most important medical advances in protecting the health of children in the century, explains Juilus G.K. Goepp, MD, assistant director of the Pediatric Emergency Department of the Children's Center at Johns Hopkins Hospital. If a parent provides an oral electrolyte maintenance solution at the very start of the illness, dehydration can be prevented. The functionality of electrolyte solutions is related to their properties, and interest in electrolyte solutions goes far beyond chemistry.

Sports drinks are designed to rehydrate the body after excessive fluid depletion. Electrolytes in particular promote normal rehydration to prevent fatigue during physical exertion. Are they a good choice for achieving the recommended fluid intake? Are

they performance and endurance enhancers like they claim? Who should drink them?

Typically, eight ounces of a sports drink provides between fifty and eighty calories and 14 to 17 grams of carbohydrate, mostly in the form of simple sugars. Sodium and potassium are the most commonly included electrolytes in sports drinks, with the levels of these in sports drinks being highly variable. The American College of Sports Medicine says a sports drink should contain 125 milligrams of sodium per 8 ounces as it is helpful in replenishing some of the sodium lost in sweat and promotes fluid uptake in the small intestine, improving hydration.

Gatorade

In the summer of 1965, the assistant football coach of the University of Florida Gators requested scientists affiliated with the university study why the withering heat of Florida caused so many heat-related illnesses in football players and provide a solution to increase athletic performance and recovery post-training or game. The discovery was that inadequate replenishment of fluids, carbohydrates, and electrolytes was the reason for the "wilting" of their football players. Based on their research, the scientists concocted a drink for the football players containing water, carbohydrates, and electrolytes and called it "Gatorade." In the next football season the Gators were nine and two and won the Orange Bowl. The Gators' success launched the sports-drink industry, which is now a multibillion-dollar industry that is still dominated by Gatorade.



University of Florida football player Chip Hinton testing Gatorade in 1965, pictured next to the leader of its team of inventors, Robert Cade.

Key Takeaway

- When a solute dissolves, its individual particles are surrounded by solvent molecules and are separated from each other.

9.4: The Dissolution Process is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.3: The Dissolution Process** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.5: Properties of Solutions

Learning Objectives

- To describe how the properties of solutions differ from those of pure solvents.

Solutions are likely to have properties similar to those of their major component—usually the solvent. However, some solution properties differ significantly from those of the solvent. Here, we will focus on liquid solutions that have a solid solute, but many of the effects we will discuss in this section are applicable to all solutions.

Colligative Properties

Solutes affect some properties of solutions that depend only on the concentration of the dissolved particles. These properties are called **colligative properties**. Four important colligative properties that we will examine here are vapor pressure depression, boiling point elevation, freezing point depression, and osmotic pressure.

Molecular compounds separate into individual molecules when they are dissolved, so for every 1 mol of molecules dissolved, we get 1 mol of particles. In contrast, **ionic** compounds separate into their constituent ions when they dissolve, so 1 mol of an ionic compound will produce more than 1 mol of dissolved particles. For example, **every mole of NaCl** that dissolves yields **1 mol of Na^+ ions** and **1 mol of Cl^- ions**, for a **total of 2 mol** of particles in solution. Thus, the effect on a solution's properties by dissolving NaCl may be twice as large as the effect of dissolving the same amount of moles of glucose ($C_6H_{12}O_6$).

Vapor Pressure Depression

All liquids evaporate. In fact, given enough volume, a liquid will turn completely into a vapor. If enough volume is not present, a liquid will evaporate only to the point where the rate of evaporation equals the rate of vapor condensing back into a liquid. The pressure of the vapor at this point is called the **vapor pressure** of the liquid.

The presence of a dissolved solid lowers the characteristic vapor pressure of a liquid so that it evaporates more slowly. (The exceptions to this statement are if the solute itself is a liquid or a gas, in which case the solute will also contribute something to the evaporation process. We will not discuss such solutions here.) This property is called vapor pressure depression and is depicted in Figure 9.5.1.




Figure 9.5.1: Vapor Pressure Depression. The presence of solute particles blocks some of the ability for liquid particles to evaporate. Thus, solutions of solid solutes typically have a lower vapor pressure than the pure solvent.

Boiling Point and Freezing Point Effects

A related property of solutions is that their boiling points are higher than the boiling point of the pure solvent. Because the presence of solute particles decreases the vapor pressure of the liquid solvent, a higher temperature is needed to reach the boiling point. This phenomenon is called boiling point elevation. **For every mole of particles dissolved in a liter of water, the boiling point of water increases by about 0.5°C.** The addition of one mole of sucrose (molecular compound) in one liter of water will raise the boiling point from 100^0C to 100.5^0C but the addition of one mole of NaCl in one liter of water will raise the boiling point by 2 x 0.5^0C = 1^0C. Furthermore, the addition of one mole of $CaCl_2$ in one liter of water will raise the boiling point by 3 x 0.5^0C = 1.5^0C.

Some people argue that putting a pinch or two of salt in water used to cook spaghetti or other pasta makes a solution that has a higher boiling point, so the pasta cooks faster. In actuality, the amount of solute is so small that the boiling point of the water is practically unchanged.

The presence of solute particles has the opposite effect on the freezing point of a solution. When a solution freezes, only the solvent particles come together to form a solid phase, and the presence of solute particles interferes with that process. Therefore, for the liquid solvent to freeze, more energy must be removed from the solution, which lowers the temperature. Thus, solutions have lower freezing points than pure solvents do. This phenomenon is called freezing point depression. **For every mole of particles in a liter of water, the freezing point decreases by about 1.9°C.**




Figure 9.5.2: Comparison of boiling and freezing points of a pure liquid (right side) with a solution (left side).

Both boiling point elevation and freezing point depression have practical uses. For example, solutions of water and ethylene glycol ($C_2H_6O_2$) are used as coolants in automobile engines because the boiling point of such a solution is greater than 100°C, the normal boiling point of water. In winter, salts like NaCl and $CaCl_2$ are sprinkled on the ground to melt ice or keep ice from forming on roads and sidewalks (Figure 9.5.2). This is because the solution made by dissolving sodium chloride or calcium chloride in water has a lower freezing point than pure water, so the formation of ice is inhibited.



Figure 9.5.2: Effect of Freezing Point Depression. The salt sprinkled on this sidewalk makes the water on the sidewalk have a lower freezing point than pure water, so it does not freeze as easily. This makes walking on the sidewalk less hazardous in winter. © Thinkstock

✔ Example 9.5.1

Which solution's freezing point deviates more from that of pure water—a 1 M solution of NaCl or a 1 M solution of $CaCl_2$?

Solution

Colligative properties depend on the number of dissolved particles, so the solution with the greater number of particles in solution will show the greatest deviation. When NaCl dissolves, it separates into two ions, Na^+ and Cl^-. But when $CaCl_2$ dissolves, it separates into three ions—one Ca^{2+} ion and two Cl^- ions. Thus, mole for mole, $CaCl_2$ will have 50% more impact on freezing point depression than NaCl.

? Exercise 9.5.1

Which solution's boiling point deviates more from that of pure water—a 1 M solution of $CaCl_2$ or a 1 M solution of $MgSO_4$?

Answer

$CaCl_2$

Example 9.5.2

Estimate the boiling point of 0.2 M $CaCl_2$ solution.

Solution

The boiling point increases 0.5^0C for every mole of solute per liter of water. For this estimation, let's assume that 1 liter of solution is roughly the same volume as 1 liter of water. A 0.2 M $CaCl_2$ solution contains 0.2 moles of $CaCl_2$ solution formula units per liter of solution. Each $CaCl_2$ unit separates into three ions.

$$0.2 \text{ mol } CaCl_2 \times \frac{3 \text{ mol ions}}{1 \text{ mol } CaCl_2} \times \frac{0.5 \text{ deg C}}{1 \text{ mol ion}} = 0.3 \text{ deg C}$$

The normal boiling point of water is 100^0C, so the boiling point of the solution is raised to 100.3^0C.

rcise

Estimate the freezing point of 0.3 M $CaCl_2$ solution.

Answer

minus 1.7^0C

Osmotic Pressure

The last colligative property of solutions we will consider is a very important one for biological systems. It involves **osmosis**, the process by which solvent molecules can pass through certain membranes but solute particles cannot. When two solutions of different concentration are present on either side of these membranes (called *semipermeable membranes*), there is a tendency for solvent molecules to move from the more dilute solution to the more concentrated solution until the concentrations of the two solutions are equal. This tendency is called **osmotic pressure**. External pressure can be exerted on a solution to counter the flow of solvent; the pressure required to halt the osmosis of a solvent is equal to the osmotic pressure of the solution.

Osmolarity (osmol) is a way of reporting the total number of particles in a solution to determine osmotic pressure. It is defined as the molarity of a solute times the number of particles a formula unit of the solute makes when it dissolves (represented by i):

$$osmol = M \times i \tag{9.5.1}$$

If more than one solute is present in a solution, the individual osmolarities are additive to get the total osmolarity of the solution. Solutions that have the same osmolarity have the same osmotic pressure. If solutions of differing osmolarities are present on opposite sides of a semipermeable membrane, solvent will transfer from the lower-osmolarity solution to the higher-osmolarity solution. Counterpressure exerted on the high-osmolarity solution will reduce or halt the solvent transfer. An even higher pressure can be exerted to force solvent from the high-osmolarity solution to the low-osmolarity solution, a process called *reverse osmosis*. Reverse osmosis is used to make potable water from saltwater where sources of fresh water are scarce.

Example 9.5.3

A 0.50 M NaCl aqueous solution and a 0.30 M $Ca(NO_3)_2$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Solution

The solvent will flow into the solution of higher osmolarity. The NaCl solute separates into two ions—Na^+ and Cl^-—when it dissolves, so its osmolarity is as follows:

osmol (NaCl) = 0.50 M × 2 = 1.0 osmol

The $Ca(NO_3)_2$ solute separates into three ions—one Ca^{2+} and two NO_3^-—when it dissolves, so its osmolarity is as follows:

osmol [$Ca(NO_3)_2$] = 0.30 M × 3 = 0.90 osmol

The osmolarity of the $Ca(NO_3)_2$ solution is lower than that of the NaCl solution, so water will transfer through the membrane from the $Ca(NO_3)_2$ solution to the NaCl solution.

Exercise 9.5.3

A 1.5 M $C_6H_{12}O_6$ aqueous solution and a 0.40 M $Al(NO_3)_3$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Answer

osmol $C_6H_{12}O_6$ = 1.5; osmol $Al(NO_3)_3$ = 1.6

The solvent flows from $C_6H_{12}O_6$ solution (lower osmolarity) to $Al(NO_3)_3$ solution (higher osmolarity).

To Your Health: Dialysis

The main function of the kidneys is to filter the blood to remove wastes and extra water, which are then expelled from the body as urine. Some diseases rob the kidneys of their ability to perform this function, causing a buildup of waste materials in the bloodstream. If a kidney transplant is not available or desirable, a procedure called dialysis can be used to remove waste materials and excess water from the blood.

In one form of dialysis, called *hemodialysis*, a patient's blood is passed though a length of tubing that travels through an *artificial kidney machine* (also called a *dialysis machine*). A section of tubing composed of a semipermeable membrane is immersed in a solution of sterile water, glucose, amino acids, and certain electrolytes. The osmotic pressure of the blood forces waste molecules and excess water through the membrane into the sterile solution. Red and white blood cells are too large to pass through the membrane, so they remain in the blood. After being cleansed in this way, the blood is returned to the body.



Figure 9.5.3: A patient undergoing hemodialysis depends on osmosis to cleanse the blood of waste products that the kidneys are incapable of removing due to disease. from Wikipedia.

Dialysis is a continuous process, as the osmosis of waste materials and excess water takes time. Typically, 5–10 lb of waste-containing fluid is removed in each dialysis session, which can last 2–8 hours and must be performed several times a week. Although some patients have been on dialysis for 30 or more years, dialysis is always a temporary solution because waste materials are constantly building up in the bloodstream. A more permanent solution is a kidney transplant.

Cell walls are semipermeable membranes, so the osmotic pressures of the body's fluids have important biological consequences. If solutions of different osmolarity exist on either side of the cells, solvent (water) may pass into or out of the cells, sometimes with disastrous results. Consider what happens if red blood cells are placed in a *hypotonic* solution, meaning a solution of lower osmolarity than the liquid inside the cells. The cells swell up as water enters them, disrupting cellular activity and eventually causing the cells to burst. This process is called *hemolysis*. If red blood cells are placed in a *hypertonic* solution, meaning one having a higher osmolarity than exists inside the cells, water leaves the cells to dilute the external solution, and the red blood cells shrivel and die. This process is called *crenation*. Only if red blood cells are placed in *isotonic* solutions that have the same osmolarity as exists inside the cells are they unaffected by negative effects of osmotic pressure. **Glucose solutions of about 0.31 M, or sodium chloride solutions of about 0.16 M, are isotonic with blood plasma.**

The concentration of an isotonic sodium chloride (NaCl) solution is only half that of an isotonic glucose ($C_6H_{12}O_6$) solution because NaCl produces two ions when a formula unit dissolves, while molecular $C_6H_{12}O_6$ produces only one particle when a formula unit dissolves. The osmolarities are therefore the same even though the concentrations of the two solutions are different.

Osmotic pressure explains why you should not drink seawater if you are abandoned in a life raft in the middle of the ocean. Its osmolarity is about three times higher than most bodily fluids. You would actually become thirstier as water from your cells was drawn out to dilute the salty ocean water you ingested. Our bodies do a better job coping with hypotonic solutions than with hypertonic ones. The excess water is collected by our kidneys and excreted.

Osmotic pressure effects are used in the food industry to make pickles from cucumbers and other vegetables and in brining meat to make corned beef. It is also a factor in the mechanism of getting water from the roots to the tops of trees!

Career Focus: Perfusionist

A perfusionist is a medical technician trained to assist during any medical procedure in which a patient's circulatory or breathing functions require support. The use of perfusionists has grown rapidly since the advent of open-heart surgery in 1953.

Most perfusionists work in operating rooms, where their main responsibility is to operate heart-lung machines. During many heart surgeries, the heart itself must be stopped. In these situations, a heart-lung machine keeps the patient alive by aerating the blood with oxygen and removing carbon dioxide. The perfusionist monitors both the machine and the status of the blood, notifying the surgeon and the anesthetist of any concerns and taking corrective action if the status of the blood becomes abnormal.

Despite the narrow parameters of their specialty, perfusionists must be highly trained. Certified perfusion education programs require a student to learn anatomy, physiology, pathology, chemistry, pharmacology, math, and physics. A college degree is usually required. Some perfusionists work with other external artificial organs, such as hemodialysis machines and artificial livers.

9.5: Properties of Solutions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.4: Properties of Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.E: Solutions (Exercises)

9.1: Solutions

Concept Review Exercises

1. What causes a solution to form?
2. How does the phrase *like dissolves like* relate to solutions?

Answers

1. Solutions form because a solute and a solvent have similar intermolecular interactions.
2. It means that substances with similar intermolecular interactions will dissolve in each other.

Exercises

1. Define *solution*.
2. Give several examples of solutions.
3. What is the difference between a solvent and a solute?
4. Can a solution have more than one solute in it? Can you give an example?
5. Does a solution have to be a liquid? Give several examples to support your answer.
6. Give at least two examples of solutions found in the human body.
7. Which substances will probably be soluble in water, a very polar solvent?
   a. sodium nitrate ($NaNO_3$)
   b. hexane (C_6H_{14})
   c. isopropyl alcohol [$(CH_3)_2CHOH$]
   d. benzene (C_6H_6)
8. Which substances will probably be soluble in toluene ($C_6H_5CH_3$), a nonpolar solvent?
   a. sodium nitrate ($NaNO_3$)
   b. hexane (C_6H_{14})
   c. isopropyl alcohol [$(CH_3)_2CHOH$]
   d. benzene (C_6H_6)
9. The solubility of alcohols in water varies with the length of carbon chain. For example, ethanol (CH_3CH_2OH) is soluble in water in any ratio, while only 0.0008 mL of heptanol ($CH_3CH_2CH_2CH_2CH_2CH_2CH_2OH$) will dissolve in 100 mL of water. Propose an explanation for this behavior.
10. Dimethyl sulfoxide [$(CH_3)_2SO$] is a polar liquid. Based on the information in Exercise 9, which do you think will be more soluble in it—ethanol or heptanol?

Answers

1. a homogeneous mixture
2. vinegar, dextrose IV, saline IV, coffee, tea, wine
3. A solvent is the majority component of a solution; a solute is the minority component of a solution.
4. yes. Coke or Pepsi has sugar, caffeine and carbon dioxide as solutes.
5. A solution does not have to be liquid; air is a gaseous solution, while some alloys are solid solutions (answers will vary).
6. Urine, plasma
7. a. probably soluble
   b. probably not soluble
   c. probably soluble
   d. probably not soluble

8.

a. probably not soluble
b. probably soluble
c. probably not soluble
d. probably soluble

9. Small alcohol molecules have strong polar intermolecular interactions, so they dissolve in water. In large alcohol molecules, the nonpolar end overwhelms the polar end, so they do not dissolve very well in water.

10. Ethanol is a smaller molecule. It will be more soluble in water than heptanol.

9.2: Concentration

Concept Review Exercises

1. What are some of the units used to express concentration?
2. Distinguish between the terms *solubility* and *concentration*.

Answers

1. % m/m, % m/v, ppm, ppb, molarity, and Eq/L (answers will vary)
2. Solubility is typically a limit to how much solute can dissolve in a given amount of solvent. Concentration is the quantitative amount of solute dissolved at any concentration in a solvent.

Exercises

1. Define *solubility*. Do all solutes have the same solubility?
2. Explain why the terms *dilute* or *concentrated* are of limited usefulness in describing the concentration of solutions.
3. If the solubility of sodium chloride (NaCl) is 30.6 g/100 mL of H_2O at a given temperature, how many grams of NaCl can be dissolved in 250.0 mL of H_2O?
4. If the solubility of glucose ($C_6H_{12}O_6$) is 120.3 g/100 mL of H_2O at a given temperature, how many grams of $C_6H_{12}O_6$ can be dissolved in 75.0 mL of H_2O?
5. How many grams of sodium bicarbonate ($NaHCO_3$) can a 25.0°C saturated solution have if 150.0 mL of H_2O is used as the solvent?
6. If 75.0 g of potassium bromide (KBr) are dissolved in 125 mL of H_2O, is the solution saturated, unsaturated, or supersaturated?
7. Calculate the mass/mass percent of a saturated solution of NaCl. Use the data from Table $9.E.1$ "Solubilities of Various Solutes in Water at 25°C (Except as Noted)", assume that masses of the solute and the solvent are additive, and use the density of H_2O (1.00 g/mL) as a conversion factor.
8. Calculate the mass/mass percent of a saturated solution of $MgCO_3$ Use the data from Table $9.E.1$ "Solubilities of Various Solutes in Water at 25°C (Except as Noted)", assume that masses of the solute and the solvent are additive, and use the density of H_2O (1.00 g/mL) as a conversion factor.
9. Only 0.203 mL of C_6H_6 will dissolve in 100.000 mL of H_2O. Assuming that the volumes are additive, find the volume/volume percent of a saturated solution of benzene in water.
10. Only 35 mL of aniline ($C_6H_5NH_2$) will dissolve in 1,000 mL of H_2O. Assuming that the volumes are additive, find the volume/volume percent of a saturated solution of aniline in water.
11. A solution of ethyl alcohol (C_2H_5OH) in water has a concentration of 20.56% v/v. What volume of C_2H_5OH is present in 255 mL of solution?
12. What mass of KCl is present in 475 mL of a 1.09% m/v aqueous solution?
13. The average human body contains 5,830 g of blood. What mass of arsenic is present in the body if the amount in blood is 0.55 ppm?

14. The Occupational Safety and Health Administration has set a limit of 200 ppm as the maximum safe exposure level for carbon monoxide (CO). If an average breath has a mass of 1.286 g, what is the maximum mass of CO that can be inhaled at that maximum safe exposure level?
15. Which concentration is greater—15 ppm or 1,500 ppb?
16. Express the concentration 7,580 ppm in parts per billion.
17. What is the molarity of 0.500 L of a potassium chromate solution containing 0.0650 mol of K_2CrO_4?
18. What is the molarity of 4.50 L of a solution containing 0.206 mol of urea [$(NH_2)_2CO$]?
19. What is the molarity of a 2.66 L aqueous solution containing 56.9 g of NaBr?
20. If 3.08 g of $Ca(OH)_2$ is dissolved in enough water to make 0.875 L of solution, what is the molarity of the $Ca(OH)_2$?
21. What mass of HCl is present in 825 mL of a 1.25 M solution?
22. What mass of isopropyl alcohol (C_3H_8O) is dissolved in 2.050 L of a 4.45 M aqueous C_3H_8O solution?
23. What volume of 0.345 M NaCl solution is needed to obtain 10.0 g of NaCl?
24. How many milliliters of a 0.0015 M cocaine hydrochloride ($C_{17}H_{22}ClNO_4$) solution is needed to obtain 0.010 g of the solute?
25. Aqueous calcium chloride reacts with aqueous silver nitrate according to the following balanced chemical equation:

$$CaCl_2(aq) + 2AgNO_3(aq) \rightarrow 2AgCl(s) + Ca(NO_3)_2(aq)$$

How many moles of AgCl(s) are made if 0.557 L of 0.235 M $CaCl_2$ react with excess $AgNO_3$? How many grams of AgCl are made?

26. Sodium bicarbonate ($NaHCO_3$) is used to react with acid spills. The reaction with sulfuric acid (H_2SO_4) is as follows:

$$2NaHCO_3(s) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(\ell) + 2CO_2(g)$$

If 27.6 mL of a 6.25 M H_2SO_4 solution were spilled, how many moles of $NaHCO_3$ would be needed to react with the acid? How many grams of $NaHCO_3$ is this?

27. The fermentation of glucose to make ethanol and carbon dioxide has the following overall chemical equation:

$$C_6H_{12}O_6(aq) \rightarrow 2C_2H_5OH(aq) + 2CO_2(g)$$

If 1.00 L of a 0.567 M solution of $C_6H_{12}O_6$ were completely fermented, what would be the resulting concentration of the C_2H_5OH solution? How many moles of CO_2 would be formed? How many grams is this? If each mole of CO_2 had a volume of 24.5 L, what volume of CO_2 is produced?

28. Aqueous sodium bisulfite gives off sulfur dioxide gas when heated:

$$2NaHSO_3(aq) \rightarrow Na_2SO_3(aq) + H_2O(\ell) + SO_2(g)$$

If 567 mL of a 1.005 M $NaHSO_3$ solution were heated until all the $NaHSO_3$ had reacted, what would be the resulting concentration of the Na_2SO_3 solution? How many moles of SO_2 would be formed? How many grams of SO_2 would be formed? If each mole of SO_2 had a volume of 25.78 L, what volume of SO_2 would be produced?

29. What is the concentration of a 1.0 M solution of $K^+(aq)$ ions in equivalents/liter?
30. What is the concentration of a 1.0 M solution of $SO_4^{2-}(aq)$ ions in equivalents/liter?
31. A solution having initial concentration of 0.445 M and initial volume of 45.0 mL is diluted to 100.0 mL. What is its final concentration?
32. A 50.0 mL sample of saltwater that is 3.0% m/v is diluted to 950 mL. What is its final mass/volume percent?

Answers

1. Solubility is the amount of a solute that can dissolve in a given amount of solute, typically 100 mL. The solubility of solutes varies widely.

2. The term *dilute* means relatively less solute and the term *concentrated* implies relatively more solute. Both are of limited usefulness because these are not accurate.

3. 76.5 g
4. 90.2 g
5. 12.6 g
6. unsaturated
7. 26.5%
8. 2.15%
9. 0.203%
10. 3.4%
11. 52.4 mL
12. 5.18 g
13. 0.00321 g
14. 2.57×10^{-4} g
15. 15 ppm
16. 7,580,000 ppb
17. 0.130 M
18. 0.0458 M
19. 0.208 M
20. 0.0475 M
21. 37.6 g
22. 548 g
23. 0.496 L
24. 20 mL
25. 0.262 mol; 37.5 g
26. 0.345 mol; 29.0 g
27. 1.13 M C_2H_5OH; 1.13 mol of CO_2; 49.7 g of CO_2; 27.7 L of CO_2
28. 0.503 M Na_2SO_3; 0.285 mol SO_2; 18.3 g SO_2; 471 L SO_2
29. 1.0 Eq/L
30. 2.0 Eq/L
31. 0.200 M
32. 0.16 % m/v

9.3: The Dissolution Process

Concept Review Exercise

1. Explain how the solvation process describes the dissolution of a solute in a solvent.

Answer

1. Each particle of the solute is surrounded by particles of the solvent, carrying the solute from its original phase.

Exercises

1. Describe what happens when an ionic solute like Na_2SO_4 dissolves in a polar solvent.
2. Describe what happens when a molecular solute like sucrose ($C_{12}H_{22}O_{11}$) dissolves in a polar solvent.
3. Classify each substance as an electrolyte or a nonelectrolyte. Each substance dissolves in H_2O to some extent.
    a. NH_4NO_3

b. CO_2
c. NH_2CONH_2
d. HCl

4. Classify each substance as an electrolyte or a nonelectrolyte. Each substance dissolves in H_2O to some extent.

a. $CH_3CH_2CH_2OH$
b. $Ca(CH_3CO_2)_2$
c. I_2
d. KOH

5. Will solutions of each solute conduct electricity when dissolved?

a. $AgNO_3$
b. $CHCl_3$
c. $BaCl_2$
d. Li_2O

6. Will solutions of each solute conduct electricity when dissolved?

a. CH_3COCH_3
b. $N(CH_3)_3$
c. $CH_3CO_2C_2H_5$
d. $FeCl_2$

Answers

1. Each ion of the ionic solute is surrounded by particles of solvent, carrying the ion from its associated crystal.

2. Each sucrose molecule is surrounded by solvent molecules (attracted to each other via intermolecular forces of attraction).

3. a. electrolyte
b. nonelectrolyte
c. nonelectrolyte
d. electrolyte

4.

a. nonelectrolyte
b. electrolyte
c. nonelectrolyte
d. electrolyte

5.

a. yes
b. no
c. yes
d. yes

6.

a. no
b. no
c. no
d. yes

9.4: Properties of Solutions

Concept Review Exercises

1. What are the colligative properties of solutions?

2. Explain how the following properties of solutions differ from those of the pure solvent: vapor pressure, boiling point, freezing point, and osmotic pressure.

Answers

1. Colligative properties are characteristics that a solution has that depend on the number, not the identity, of solute particles.
2. In solutions, the vapor pressure is lower, the boiling point is higher, the freezing point is lower, and the osmotic pressure is higher.

Exercises

1. In each pair of aqueous systems, which will have the lower vapor pressure?
   a. pure water or 1.0 M NaCl
   b. 1.0 M NaCl or 1.0 M $C_6H_{12}O_6$
   c. 1.0 M $CaCl_2$ or 1.0 M $(NH_4)_3PO_4$
2. In each pair of aqueous systems, which will have the lower vapor pressure?
   a. 0.50 M $Ca(NO_3)_2$ or 1.0 M KBr
   b. 1.5 M $C_{12}H_{22}O_{11}$ or 0.75 M $Ca(OH)_2$
   c. 0.10 M $Cu(NO_3)_2$ or pure water
3. In each pair of aqueous systems, which will have the higher boiling point?
   a. pure water or a 1.0 M NaCl
   b. 1.0 M NaCl or 1.0 M $C_6H_{12}O_6$
   c. 1.0 M $CaCl_2$ or 1.0 M $(NH_4)_3PO_4$
4. In each pair of aqueous systems, which will have the higher boiling point?
   a. 0.50 M $Ca(NO_3)_2$ or 1.0 M KBr
   b. 1.5 M $C_{12}H_{22}O_{11}$ or 0.75 M $Ca(OH)_2$
   c. 0.10 M $Cu(NO_3)_2$ or pure water
5. Estimate the boiling point of each aqueous solution. The boiling point of pure water is 100.0°C.
   a. 0.50 M NaCl
   b. 1.5 M Na_2SO_4
   c. 2.0 M $C_6H_{12}O_6$
6. Estimate the freezing point of each aqueous solution. The freezing point of pure water is 0.0°C.
   a. 0.50 M NaCl
   b. 1.5 M Na_2SO_4
   c. 2.0 M $C_6H_{12}O_6$
7. Explain why salt (NaCl) is spread on roads and sidewalks to inhibit ice formation in cold weather.
8. Salt (NaCl) and calcium chloride ($CaCl_2$) are used widely in some areas to minimize the formation of ice on sidewalks and roads. One of these ionic compounds is better, mole for mole, at inhibiting ice formation. Which is that likely to be? Why?
9. What is the osmolarity of each aqueous solution?
   a. 0.500 M NH_2CONH_2
   b. 0.500 M NaBr
   c. 0.500 M $Ca(NO_3)_2$
10. What is the osmolarity of each aqueous solution?
    a. 0.150 M KCl
    b. 0.450 M $(CH_3)_2CHOH$
    c. 0.500 M $Ca_3(PO_4)_2$
11. A 1.0 M solution of an unknown soluble salt has an osmolarity of 3.0 osmol. What can you conclude about the salt?

12. A 1.5 M NaCl solution and a 0.75 M $Al(NO_3)_3$ solution exist on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and the direction of solvent flow, if any, across the membrane.

Answers

1. a. 1.0 M NaCl
   b. 1.0 M NaCl
   c. 1.0 M $(NH_4)_3PO_4$
2.
   a. 1.0 M KBr
   b. 0.75 M $Ca(OH)_2$
   c. 0.10 M $Cu(NO_3)_2$
3.
   a. 1.0 M NaCl
   b. 1.0 M NaCl
   c. 1.0 M $(NH_4)_3PO_4$
4.
   a. 1.0 M KBr
   b. 0.75 M $Ca(OH)_2$
   c. 0.10 M $Cu(NO_3)_2$
5.
   a. 100.5°C
   b. 102.3°C
   c. 101°C
6.
   a. -1.9°C
   b. -8.6°C
   c. -3.8°C

7. NaCl lowers the freezing point of water, so it needs to be colder for the water to freeze.

8. $CaCl_2$ splits up into 3 ions while NaCl splits up into 2 ions only. $CaCl_2$ will be more effective.

9.
   a. 0.500 osmol
   b. 1.000 osmol
   c. 1.500 osmol
10.
   a. 0.300 osmol
   b. 0.450 osmol
   c. 2.50 osmol

11. It must separate into three ions when it dissolves.

12. Both NaCl and $Al(NO_3)_3$ have 3.0 osmol. There will be no net difference in the solvent flow.

9.5: Chemical Equilibrium

Concept Review Exercises

1. What is chemical equilibrium?

2. What does the equilibrium constant tell us?

Answers

1. The rate of the forward reaction equals the rate of the reverse reaction.

2. The ratio of products and reactants when the system is at equilibrium.

EXERCISES

1. If the reaction $H_2 + I_2 \rightleftharpoons 2HI$ is at equilibrium, do the concentrations of HI, H_2, and I_2 have to be equal?

2. Do the concentrations at equilibrium depend upon how the equilibrium was reached?

3. What does it mean if the K_{eq} is > 1?

4. What does it mean if the K_{eq} is < 1?

5. Does the equilibrium state depend on the starting concentrations?

6. Write an expression for the equilibrium constant K equation.

   a. $PCl_5\ (g) \rightleftarrows PCl_3\ (g) + Cl_2\ (g)$

   b. $2\ O_3\ (g) \rightleftharpoons 3\ O_2\ (g)$

7. Benzene is one of the compounds used as octane enhancers in unleaded gasoline. It is manufactured by the catalytic conversion of acetylene to benzene: $3C_2H_2(g) \longrightarrow C_6H_6(g)$. Which value of *K* would make this reaction most useful commercially? Explain your answer.

   a. $K \approx 0.01$

   b. $K \approx 1$

   c. $K \approx 10$.

8. Tell whether the reactants or the products are favored at equilibrium:

   a. $2NH_3(g) \rightleftharpoons N_2(g) + 3H_2(g)$ $K = 172$

   b. $2SO_3(g) \rightleftharpoons 2SO_2(g) + O_2(g)$ $K = 0.230$

   c. $2NO(g) + Cl_2(g) \rightleftharpoons 2NOCl(g)$ $K = 4.6\times10^4$

   d. $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$ $K = 0.050$

AnswerS

1. No, the concentrations are constant but the concentrations do not have to be equal.

2. No.

3. More products than reactants are present at equilibrium.

4. More reactants than products present at equilibrium.

5. No. The equilibrium ratio does not depend on the initial concentrations.

6.

   a. $K = \dfrac{[PCl_3][Cl_2]}{[PCl_5]}$

   b. $K = \dfrac{[O_2]^3}{[O_3]^2}$

7. The answer is c. $K \approx 10$.

Since $K = \dfrac{[C_6H_6]^2}{[C_2H_2]^3}$ (K≈10), this means that C_6H_6 predominates over C_2H_2. In such a case, the reaction would be commercially feasible if the rate to equilibrium is suitable.

8.

a. products

b. reactants

c. products

d. reactants

9.6: Le Chatelier's Principle

Concept Review Exercises

1. Define Le Chatelier’s principle.

2. List the three factors types of changes that can disturb the equilibrium of a system.

Answers

1. Le Chatelier’s principle states that a system at equilibrium is disturbed, it will respond in a way to minimize the disturbance.

2. temperature, change in amount of substance, change in pressure through change in volume

EXERCISES

1. How will each change affect the reaction?

$PCl_5(g)$ + heat $\rightleftharpoons PCl_3(g) + Cl_2(g)$

a. Addition of PCl_5
b. Addition of Cl_2
c. Removal of PCl_3
d. Increasing temperature
e. Decreasing temperature
f. Decreasing volume

2. How will each change affect the reaction?

$HNO_2(aq) \rightleftharpoons H^+(aq) + NO_2^-(aq)$

a. Removal of HNO_2
b. Addition of HCl (i.e. adding more H^+)
c. Increasing volume
d. Decreasing volume
e. Removal of NO_2^-
f. Addition of OH^- (which will react with and remove H^+)

3. How will each change affect the reaction?

$CO_2(g) + C(s) \rightleftharpoons 2CO(g)$ $\Delta H=172.5kJ$

a. Addition of CO_2
b. Removal of CO_2
c. Increasing temperature
d. Decreasing temperature
e. Increasing volume
f. Addition of CO

4. How will each change affect the reaction?

$H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ $\Delta H=-9.48kJ$

a. Addition of H_2
b. Removal of H_2
c. Increasing temperature

d. Decreasing temperature
e. Increasing volume
f. Decreasing volume

AnswerS

1.

a. shift right
b. shift left
c. shift right
d. shift right
e. shift left
f. shift left

2.

a. shift left
b. shift left
c. no effect
d. no effect
e. shift right
f. shift right

3.

a. shift right
b. shift left
c. shift right
d. shift left
e. shift right
f. shift left

4.

a. shift right
b. shift left
c. shift left
d. shift right
e. no effect
f. no effect

9.7: Osmosis and Diffusion

Concept Review Exercises

1. What are some of the features of a semipermeable membrane?

2. What do the prefixes hyper, hypo, and iso mean?

Answers

1. A semipermeable membrane allows some substances to pass through but not others.

2. hyper – higher; hypo – lower; iso - same

EXERCISES

1. Two solutions are separated by a semipermeable membrane. Solution A contains 25.0 g of NaCl in 100.0 mL of water and solution B contains 35.0 g of NaCl in 100.0 mL of water.

a. Which one has a higher concentration?
b. Which way will water molecules flow?
c. Which volume will increase?
d. Which volume will decrease?
e. What will happen to the concentration of solution A?
f. What will happen to the concentration of solution B?

2. Two solutions with different concentrations and compositions are separated by a semipermeable membrane. The left-hand solution is a .50 M solution of $MgSO_4$, while the right-hand solution contains $CaCl_2$ at a concentration of .40 M. Determine the direction of the flow of solvent, left or right.

3. Given the following situations, wherein two tanks of different solutions are separated by a semipermeable membrane, determine the direction of the flow of solvent (water).

a. Solution A contains a 0.40 M concentration of $CaCl_2$, while Solution B contains a 0.45 M concentration of KI

b. Solution A contains a 1.00 M concentration of NH_4Cl, while Solution B contains a 1.00 M concentration of CH_2O

4. Cells are placed in a solution and the cells then undergo hemolysis. What can be said about the relative concentrations of solute in the cell and the solution?

5. Describe the relative concentrations inside and outside a red blood cell when crenation occurs.

6. A saltwater fish is placed in a freshwater tank. What will happen to the fish? Describe the flow of water molecules to explain the outcome.

7. What makes up the "head" region of a phospholipid? Is it hydrophobic or hydrophilic?

8. What makes up the "tail" region of a phospholipid? Is it hydrophobic or hydrophilic?

Answers

1. Two solutions are separated by a semipermeable membrane. Solution A contains 25.0 g of NaCl in 100.0 mL of water and solution B contains 35.0 g of NaCl in 100.0 mL of water.

a. Solution B
b. A → → B
c. B
d. A
e. increase
f. decrease

2. Water (solvent) flows from left to right.

3.

a. Water flows from Solution B to Solution A.

b. Water flows from Solution B to Solution A.

4. Cells contain fluid with higher concentration than solution outside the cell.

5. Cells contain fluid with a lower concentration than the solution outside the cell.

6. Water molecules will flow from the tank water into the fish because the fish has a higher concentration of salt. If the fish absorbs too much water, it will die.

7. The "head" region is a phosphate group and it is hydrophilic.

8. The "tail" is a hydrocarbon tail and it is hydrophobic.

Contributors

-
- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)

Additional Exercises

1. Calcium nitrate reacts with sodium carbonate to precipitate solid calcium carbonate:

$$Ca(NO_3)_{2(aq)} + Na_2CO_{3(aq)} \rightarrow CaCO_{3(s)} + NaNO_{3(aq)} \tag{9.E.1}$$

   a. Balance the chemical equation.
   b. How many grams of Na_2CO_3 are needed to react with 50.0 mL of 0.450 M $Ca(NO_3)_2$?
   c. Assuming that the Na_2CO_3 has a negligible effect on the volume of the solution, find the osmolarity of the $NaNO_3$ solution remaining after the $CaCO_3$ precipitates from solution.

2. The compound HCl reacts with sodium carbonate to generate carbon dioxide gas:

$$HCl_{(aq)} + Na_2CO_{3(aq)} \rightarrow H_2O_{(\ell)} + CO_{2(g)} + NaCl_{(aq)} \tag{9.E.2}$$

   a. Balance the chemical equation.
   b. How many grams of Na_2CO_3 are needed to react with 250.0 mL of 0.755 M HCl?
   c. Assuming that the Na_2CO_3 has a negligible effect on the volume of the solution, find the osmolarity of the NaCl solution remaining after the reaction is complete.

3. Estimate the freezing point of concentrated aqueous HCl, which is usually sold as a 12 M solution. Assume complete ionization into H^+ and Cl^- ions.

4. Estimate the boiling point of concentrated aqueous H_2SO_4, which is usually sold as an 18 M solution. Assume complete ionization into H^+ and HSO_4^- ions.

5. Seawater can be approximated by a 3.0% m/m solution of NaCl in water. Determine the molarity and osmolarity of seawater. Assume a density of 1.0 g/mL.

6. Human blood can be approximated by a 0.90% m/m solution of NaCl in water. Determine the molarity and osmolarity of blood. Assume a density of 1.0 g/mL.

7. How much water must be added to 25.0 mL of a 1.00 M NaCl solution to make a resulting solution that has a concentration of 0.250 M?

8. Sports drinks like Gatorade are advertised as capable of resupplying the body with electrolytes lost by vigorous exercise. Find a label from a sports drink container and identify the electrolytes it contains. You should be able to identify several simple ionic compounds in the ingredients list.

9. Occasionally we hear a sensational news story about people stranded in a lifeboat on the ocean who had to drink their own urine to survive. While distasteful, this act was probably necessary for survival. Why not simply drink the ocean water? (Hint: See Exercise 5 and Exercise 6 above. What would happen if the two solutions in these exercises were on opposite sides of a semipermeable membrane, as we would find in our cell walls?)

Answers

1.

   a. $Ca(NO_3)_2(aq) + Na_2CO_3(aq) \rightarrow CaCO_3(s) + 2NaNO_3(aq)$
   b. 2.39 g
   c. 1.80 osmol

2.

   a. $2HCl\ (aq) + Na_2CO_3(aq) \rightarrow H_2O(\ell) + CO_2(g) + 2NaCl\ (aq)$
   b. 10.0 g
   c. 1.51 M

3. −45.6°C
4. 118°C
5. 0.513 M; 1.026 osmol
6. molarity = 0.15 M; osmolarity = 0.31 M
7. 75.0 mL
8. magnesium chloride, calcium chloride (answers may vary)
9. The osmotic pressure of seawater is too high. Drinking seawater would cause water to go from inside our cells into the more concentrated seawater, ultimately killing the cells.

9.E: Solutions (Exercises) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.1: Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **9.2: Concentration** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **9.3: The Dissolution Process** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **9.4: Properties of Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **9.5: Chemical Equilibrium** by CK-12 Foundation is licensed CK-12. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **9.6: Le Chatelier's Principle** by CK-12 Foundation is licensed CK-12.
- **8.4: Osmosis and Diffusion** by CK-12 Foundation is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

9.S: Solutions (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **solution** is a homogeneous mixture. The major component is the **solvent**, while the minor component is the **solute**. Solutions can have any phase; for example, an **alloy** is a solid solution. Solutes are **soluble** or **insoluble**, meaning they dissolve or do not dissolve in a particular solvent. The terms **miscible** and **immiscible**, instead of soluble and insoluble, are used for liquid solutes and solvents. The statement *like dissolves like* is a useful guide to predicting whether a solute will dissolve in a given solvent.

The amount of solute in a solution is represented by the **concentration** of the solution. The maximum amount of solute that will dissolve in a given amount of solvent is called the **solubility** of the solute. Such solutions are **saturated**. Solutions that have less than the maximum amount are **unsaturated**. Most solutions are unsaturated, and there are various ways of stating their concentrations. **Mass/mass percent**, **volume/volume percent**, and **mass/volume percent** indicate the percentage of the overall solution that is solute. **Parts per million (ppm)** and **parts per billion (ppb)** are used to describe very small concentrations of a solute. **Molarity**, defined as the number of moles of solute per liter of solution, is a common concentration unit in the chemistry laboratory. **Equivalents** express concentrations in terms of moles of charge on ions. When a solution is diluted, we use the fact that the amount of solute remains constant to be able to determine the volume or concentration of the final diluted solution.

Dissolving occurs by **solvation**, the process in which particles of a solvent surround the individual particles of a solute, separating them to make a solution. For water solutions, the word **hydration** is used. If the solute is molecular, it dissolves into individual molecules. If the solute is ionic, the individual ions separate from each other, forming a solution that conducts electricity. Such solutions are called **electrolytes**. If the dissociation of ions is complete, the solution is a **strong electrolyte**. If the dissociation is only partial, the solution is a **weak electrolyte**. Solutions of molecules do not conduct electricity and are called **nonelectrolytes**.

Solutions have properties that differ from those of the pure solvent. Some of these are **colligative** properties, which are due to the number of solute particles dissolved, not the chemical identity of the solute. Colligative properties include **vapor pressure depression**, **boiling point elevation**, **freezing point depression**, and **osmotic pressure**. Osmotic pressure is particularly important in biological systems. It is caused by **osmosis**, the passage of solvents through certain membranes like cell walls. The **osmolarity** of a solution is the product of a solution's molarity and the number of particles a solute separates into when it dissolves. Osmosis can be reversed by the application of pressure; this reverse osmosis is used to make fresh water from saltwater in some parts of the world. Because of osmosis, red blood cells placed in hypotonic or hypertonic solutions lose function through either hemolysis or crenation. If they are placed in isotonic solutions, however, the cells are unaffected because osmotic pressure is equal on either side of the cell membrane.

9.S: Solutions (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **9.S: Solutions (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

10: Equilibrium

10.1: The Rate of a Chemical Reaction
10.2: The Idea of Dynamic Chemical Equilibrium
10.3: The Equilibrium Constant - A Measure of How Far a Reaction Goes
10.4: Disturbing a Reaction at Equilibrium- Le Châtelier's Principle
10.5: The Effect of a Concentration Change on Equilibrium
10.6: The Effect of a Volume Change on Equilibrium
10.7: The Effect of Temperature Changes on Equilibrium
10.8: The Path of a Reaction and the Effect of a Catalyst

10: Equilibrium is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

10.1: The Rate of a Chemical Reaction

Learning Objectives

- Describe the conditions for successful collisions that cause reactions.
- Describe *rate* in terms of the conditions of successful collisions.
- Describe how changing the temperature, concentration of a reactant, or surface area of a reaction affects the rate of a reaction.
- Define a catalyst and how a catalyst affects the rate of a reaction.


We know that a chemical system can be made up of atoms (H_2, N_2, K, etc.), ions (NO_3^-, Cl^-, Na^+, etc.), or molecules (H_2O, $C_{12}H_{22}O_{11}$, etc.). We also know that in a chemical system, these particles are moving around in random motion. The **collision theory** explains why reactions occur at this particle level between these atoms, ions, and/or molecules. It also explains how it is possible to speed up or slow down reactions that are occurring.

Collision Theory

The collision theory provides us with the ability to predict what conditions are necessary for a successful reaction to take place. These conditions include:


1. The particles must collide with each other.
2. The particles must collide with sufficient energy to break the old bonds.
3. The particles must have proper orientation.

A chemical reaction involves breaking bonds in the reactants, rearranging the atoms into new groupings (the products), and forming new bonds in the products.



This collision is successful and results in reaction. (Source: Richard Parsons. CC-BY-SA)

Therefore, a collision must not only occur between reactant particles, but the collision also has to have sufficient energy to break all the reactant bonds that need to be broken in order to form the products. Some reactions need less collision energy than others. The amount of energy that reactant particles must have in order to break the old bonds for a reaction to occur is called the **activation energy**, abbreviated E_a. Another way to think of this is to look at an energy diagram, as shown in the figure. Particles must be able to get over the "bump"—the activation energy—if they are going to react. If the reactant particles collide with less than the activation energy, the particles will rebound (bounce off of each other), and no reaction will occur.



Each reaction has its own activation energy, E_a. The smaller the "bump", the less energy particles must have to react.

Reaction Rate

Chemists use reactions to generate a product for which they have a use. For the most part, the reactions that produce some desired compound are only useful if the reaction occurs at a reasonable rate. For example, using a reaction to produce brake fluid would not be useful if the reaction required 8,000 years to complete the product. Such a reaction would also not be useful if the reaction was so fast that it was explosive. For these reasons, chemists wish to be able to control reaction rates. In some cases, chemists wish to speed up reactions that are too slow or slow down reactions that are too fast. In order to gain any control over reaction rates, we must know the factors that affect reaction rates. Chemists have identified many factors that affect the rate of a reaction.

The rate, or speed, at which a reaction occurs depends on the frequency of successful collisions. Remember, a successful collision occurs when two reactants collide with enough energy and with the right orientation. That means if there is an increase in the number of collisions, an increase in the number of particles that have enough energy to react, and/or an increase in the number of particles with the correct orientation, the rate of reaction will increase.

Effect of Temperature on Rate of Reaction

The rate of reaction was discussed in terms of three factors: collision frequency, the collision energy, and the geometric orientation. Remember that the collision frequency is the number of collisions per second. The collision frequency is dependent, among other factors, on the temperature of the reaction.

When the temperature is increased, the average velocity of the particles is increased. The average kinetic energy of these particles is also increased. The result is that the particles will collide more frequently, because the particles move around faster and will encounter more reactant particles. However, this is only a minor part of the reason why the rate is increased. Just because the particles are colliding more frequently does not mean that the reaction will definitely occur.

The major effect of increasing the temperature is that more of the particles that collide will have the amount of energy needed to have an effective collision. In other words, more particles will have the necessary activation energy.

At room temperature, the hydrogen and oxygen in the atmosphere do not have sufficient energy to attain the activation energy needed to produce water:

$$\mathrm{O_2}\,(g) + \mathrm{H_2}\,(g) \rightarrow \text{No reaction}$$

At any one moment in the atmosphere, there are many collisions occurring between these two reactants. But what we find is that water is not formed from the oxygen and hydrogen molecules colliding in the atmosphere, because the activation energy barrier is just too high, and all the collisions are resulting in rebound. When we increase the temperature of the reactants or give them energy in some other way, the molecules have the necessary activation energy and are able to react to produce water:

$$\mathrm{O_2}\,(g) + \mathrm{H_2}\,(g) \rightarrow \mathrm{H_2O}\,(l)$$

There are times when the rate of a reaction needs to be slowed down. Lowering the temperature could also be used to decrease the number of collisions that would occur and lowering the temperature would also reduce the kinetic energy available for activation energy. If the particles have insufficient activation energy, the collisions will result in rebound rather than reaction. Using this idea, when the rate of a reaction needs to be lower, keeping the particles from having sufficient activation energy will definitely keep the reaction at a lower rate.

Society uses the effects of temperature on reaction rate every day. Food storage is a prime example of how the temperature effect on reaction rate is utilized by society. Consumers store food in freezers and refrigerators to slow down the processes that cause it to spoil. The decrease in temperature decreases the rate at which food will break down or be broken down by bacteria. In the early years of the 20th century, explorers were fascinated with being the first to reach the South Pole. In order to attempt such a difficult task at a time without most of the technology that we take for granted today, they devised a variety of ways of surviving. One method was to store their food in the snow to be used later during their advances to the pole. On some explorations, they buried so much food that they didn't need to use all of it, and some was left behind. Many years later, when this food was located and thawed, it was found to still be edible.

When milk, for example, is stored in the refrigerator, the molecules in the milk have less energy. This means that while molecules will still collide with other molecules, few of them will react (which means in this case "spoil") because the molecules do not have sufficient energy to overcome the activation energy barrier. The molecules do have energy and are colliding, however, and so, over time, even in the refrigerator, the milk will spoil. Eventually the higher energy molecules will gain the energy needed to react and when enough of these reactions occur, the milk becomes "soured".


However, if that same carton of milk was at room temperature, the milk would react (in other words, "spoil") much more quickly. Most of the molecules would have sufficient energy to overcome the energy barrier at room temperature, and many more collisions would occur. This allows for the milk to spoil in a fairly short amount of time. This is also the reason why most fruits and vegetables ripen in the summer when the temperature is much warmer. You may have experienced this first hand if you have ever bitten into an unripe banana—it was probably sour tasting and might even have felt like biting into a piece of wood! When a banana ripens, numerous reactions occur that produce all the compounds that we expect to taste in a banana. But this can only happen if the temperature is high enough to allow these reactions to make those products.

Effect of Concentration on Rate of Reaction

If you had an enclosed space, like a classroom, and there was one red ball and one green ball flying around the room in random motion, undergoing perfectly elastic collisions with the walls and with each other, in a given amount of time, the balls would collide with each other a certain number of times determined by probability. If you now put two red balls and one green ball in the room under the same conditions, the probability of a collision between a red ball and the green ball would exactly double. The green ball would have twice the chance of encountering a red ball in the same amount of time.

In terms of chemical reactions, a similar situation exists. Particles of two gaseous reactants or two reactants in solution have a certain probability of undergoing collisions with each other in a reaction vessel. If you double the concentration of either reactant, the probability of a collision doubles. The rate of reaction is proportional to the number of collisions per unit time. If one concentration is doubled, the number of collisions will also double. Assuming that the percent of collisions that are successful does not change, then having twice as many collisions will result in

twice as many successful collisions. The rate of reaction is proportional to the number of collisions over time; increasing the concentration of either reactant increases the number of collisions, and therefore increases the number of successful collisions and the reaction rate.




The reaction mixture on the left is less concentrated, so particles will not collide as often. The reaction will be slower. Because the reacting particles on the right image have a greater change of colliding, the reaction will go faster. CC – Tracy Poulsen

For example, the chemical test used to identify a gas as oxygen, or not, relies on the fact that increasing the concentration of a reactant increases reaction rate. The reaction we call combustion refers to a reaction in which a flammable substance reacts with oxygen. If we light a wooden splint (a thin splinter of wood) on fire and then blow the fire out, the splint will continue to glow in air for a period of time. If we insert that glowing splint into any gas that does not contain oxygen, the splint will immediately cease to glow—that is, the reaction stops. Oxygen is the only gas that will support combustion, Air is approximately 20% oxygen gas. If we take that glowing splint and insert it into pure oxygen gas, the reaction will increase its rate by a factor of five, since pure oxygen has 5 times the concentration of oxygen that is in the air. When the reaction occurring on the glowing splint increases its rate by a factor of five, the glowing splint will suddenly burst back into full flame. This test, of thrusting a glowing splint into a gas, is used to identify the gas as oxygen. Only a greater concentration of oxygen than that found in air will cause the glowing splint to burst into flame.

Effect of Surface Area on Rate of Reaction

The very first requirement for a reaction to occur between reactant particles is that the particles must collide with one another. The previous section pointed out how increasing the concentration of the reactants increases reaction rate because it increases the frequency of collisions between particles. It can be shown that the number of collisions that occur between reactant particles is also dependent on the surface area of solid reactants. Consider a reaction between reactant RED and reactant BLUE in which reactant blue is in the form of a single lump. Then compare this to the same reaction where reactant blue has been broken up into many smaller pieces.



In these figures, only the particles on the outside of the solid blue reactant have a chance to collide with the red reactant. In figure B, the same amount of solid reactant as used in A was crushed into smaller particles. This means that more particles on the outside of the reactant have an opportunity to collide with the red reactant and speeds up the reaction.

In the diagram, only the blue particles on the outside surface of the lump are available for collision with reactant red. The blue particles on the interior of the lump are protected by the blue particles on the surface. In Figure A, if you count the number of blue particles available for collision, you will find that only 20 blue particles could be struck by a particle of reactant red. In Figure A, there are a number of blue particles on the interior of the lump that cannot be struck. In Figure B, however, the lump has been broken up into smaller pieces and all the interior blue particles are now on a surface and available for collision. In Figure B, more collisions between the blue and red will occur, and therefore, the reaction in Figure B will occur at a faster rate than the same reaction in Figure A. Increasing the surface area of a reactant increases the frequency of collisions and increases the reaction rate.

Several smaller particles have more surface area than one large particle. The more surface area that is available for particles to collide, the faster the reaction will occur. You can see an example of this in everyday life if you have ever tried to start a fire in the fireplace. If you hold a match up against a large log in an attempt to start the log burning, you will find it to be an unsuccessful effort. Holding a match against a large log will not cause enough reactions to occur in order to keep the fire going by providing sufficient activation energy for further reactions. In order to start a wood fire, it is common to break a log up into many small, thin sticks called kindling. These thinner sticks of wood provide many times the surface area of a single log. The match will successfully cause enough reactions in the kindling so that sufficient heat is given off to provide activation energy for further reactions.

There have been, unfortunately, cases where serious accidents were caused by the failure to understand the relationship between surface area and reaction rate. One such example occurred in flour mills. A grain of wheat is not very flammable. It takes significant effort to get a grain of wheat to burn. If the grain of wheat, however, is pulverized and scattered through the air, only a spark is necessary to cause an explosion. When the wheat is ground to make flour, it is pulverized into a fine powder and some of the powder gets scattered around in the air. A small spark then, is sufficient to start a very rapid reaction which can destroy the entire flour mill. In a 10-year period from 1988 to 1998, there were 129 grain dust explosions in mills in the United States. Efforts are now made in flour mills to have huge fans circulate the air in the mill through filters to remove the majority of the flour dust particles.

Another example is in the operation of coal mines. Coal will of course burn, but it takes an effort to get the coal started; once it is burning, it burns slowly because only the surface particles are available to collide with oxygen particles. The interior particles of coal have to wait until the outer surface of the coal lump burns off before they can collide with oxygen. In coal mines, huge blocks of coal must be broken up before the coal can be brought out of the mine. In the process of breaking up the huge blocks of coal, drills are used to drill into the walls of coal. This drilling produces fine coal dust that mixes into the air; then a spark from a tool can cause a massive explosion in the mine. There are explosions in coal mines for other reasons, but coal dust explosions have contributed to the death of many miners. In modern coal mines, lawn sprinklers are used to spray water through the air in the mine and this reduces the coal dust in the air, and eliminates coal dust explosions.

Effect of a Catalyst on Rate of Reaction

The final factor that affects the rate of the reaction is the effect of a catalyst. A **catalyst** is a substance that speeds up the rate of the reaction, without being consumed by the reaction itself.


In the reaction of potassium chlorate breaking down to potassium chloride and oxygen, a catalyst is available to make this reaction occur much faster than it would occur by itself under room conditions. The reaction is:

$$2\mathrm{KClO_3}\,(s) \xrightarrow{\mathrm{MnO_2}(s)} 2\mathrm{KCl}\,(s) + 3\mathrm{O_2}\,(g)$$

The catalyst is manganese dioxide and its presence causes the reaction shown above to run many times faster than it occurs without the catalyst. When the reaction has reached completion, the $\mathrm{MnO_2}$ can be removed from the reaction vessel and its condition is exactly the same as it was before the reaction. This is part of the definition of a catalyst—that it is not consumed by the reaction. You should note that the catalyst is not written into the equation as a reactant or a product, but is noted above the yields arrow. This is standard notation for the use of a catalyst.

Some reactions occur very slowly without the presence of a catalyst. In other words, the activation energy for these reactions is very high. When the catalyst is added, the activation energy is lowered because the catalyst provides a new reaction pathway with lower activation energy.

In the figure on the right, the endothermic reaction shows the catalyst reaction in red with the lower activation energy, designated E_a'. The new reaction pathway has lower activation energy but has no effect on the energy of the reactants, the products, or the value of ΔH. The same is true for the exothermic reaction. The activation energy of the catalyzed reaction is lower than that of the uncatalyzed reaction. The new reaction pathway provided by the catalyst affects the energy required for reactant bonds to break, and product bonds to form.



A catalyst speeds up a reaction by lowering the activation energy, E_a. Because less energy is required to react, more particles have the necessary energy.

While many reactions in the laboratory can be increased by increasing the temperature, this is not possible for all of the reactions that occur in our bodies throughout our entire lives. In fact, the body needs to be maintained at a very specific temperature: $98.6^{\circ}\mathrm{F}$ or $37^{\circ}\mathrm{C}$. Of course there are times, such as when the body is fighting infection, when the body temperature may be increased. But generally, in a healthy person, the temperature is quite consistent. However, many of the reactions that a healthy body depends on could never occur at body temperature. The answer to this dilemma is catalysts—also referred to as enzymes. Many of these enzymes are made in human cells because human DNA carries the directions to make them. However, there are some enzymes required by the body that are not made by human cells. These catalysts must be supplied to our bodies in the food we eat and are called vitamins.

Reversible Reactions

Typically when we think of a chemical reaction, we think of the reactants getting totally used up so that none are left, and that we end up with only products. Also, we generally consider chemical reactions as one-way events. You may well have learned during earlier science classes that this is one way to distinguish chemical change from physical change—physical changes (such as the melting and freezing of ice) are easily reversed, but chemical changes cannot be reversed (pretty tough to un-fry an egg).

Throughout this chaper, we will see that this isn't always the case. We will see that many chemical reactions are, in fact, reversible under the right conditions. And because many reactions can be reversed, our idea of a reaction ending with no reactants left, only products, will need to be modified.

Here are some examples of reactions that can be reversed:

1.	Nitrogen dioxide, $\mathrm{NO_2}$, a reddish-brown gas, reacts to form colorless dinitrogen tetroxide, $\mathrm{N_2O_4}$: $2\,\mathrm{NO_2(g)} \rightarrow \mathrm{N_2O_4(g)}$ But the reaction can also go the other way—dinitrogen tetroxide also readily breaks down to form nitrogen dioxide: $\mathrm{N_2O_4(g)} \rightarrow 2\,\mathrm{NO_2(g)}$ We typically write a reaction that can go in both directions by using a double arrow (which will sometimes appear as $\leftrightarrow$ in these online notes): $2\,\mathrm{NO_2(g)} \leftrightarrow \mathrm{N_2O_4(g)}$ Because the reaction continues in both directions at the same time, we never run out of either $\mathrm{NO_2}$ or $\mathrm{N_2O_4}$. $\mathrm{NO_2}$ is continually being used up to form $\mathrm{N_2O_4}$, but at the same time $\mathrm{N_2O_4}$ is forming more $\mathrm{NO_2}$

2.

When hydrogen gas is passed over heated iron oxide, iron and steam are produced:

(1)	$\mathrm{Fe_3O_4(s) + 4\,H_2(g) \rightarrow 3\,Fe(s) + 4}$
The reverse reaction can occur when steam is passed over red-hot iron:	
(2)	$\mathrm{3\,Fe(s) + 4\,H_2O(g) \rightarrow Fe_3O_4(s) +}$
We can write these two equations together as:	
(3)	\(\ce{Fe_3O_4(s) + 4H_2(g) \leftrightarrow 3Fe(s) + 4H_2O(g)}\)

When we have a reversible reaction written in this way, we need to be able to distinguish between which way the reaction is headed. As written above in Reaction (3), we would say that in the ***forward reaction***, iron oxide and hydrogen gas, the reactants, produce the products iron and steam.

During the ***reverse reaction***, iron reacts with steam to produce the products iron oxide and hydrogen gas.

It is important to understand the terminology, and to use the terms correctly.

Does it matter which way we write our reversible reaction? It can also be written as

$$\mathrm{3\,Fe(s) + 4\,H_2O(g) \leftrightarrow Fe_3O_4(s) + 4\,H_2(g)}$$

Now iron and steam are reactants of the forward direction, and iron oxide and hydrogen gas would be the reactants of the reverse direction.

Summary

- The collision theory explains why reactions occur between atoms, ions, and molecules.
- In order for a reaction to be effective, particles must collide with enough energy, and have the correct orientation.
- With an increase in temperature, there is an increase in energy that can be converted into activation energy in a collision, and that will increase the reaction rate. A decrease in temperature would have the opposite effect.
- With an increase in temperature, there is an increase in the number of collisions.
- Increasing the concentration of a reactant increases the frequency of collisions between reactants and will, therefore, increase the reaction rate.
- Increasing the surface area of a reactant (by breaking a solid reactant into smaller particles) increases the number of particles available for collision and will increase the number of collisions between reactants per unit time.
- A catalyst is a substance that speeds up the rate of the reaction without being consumed by the reaction itself. When a catalyst is added, the activation energy is lowered because the catalyst provides a new reaction pathway with lower activation energy.

Vocabulary

- **Catalyst** - A substance that speeds up the rate of the reaction without being consumed by the reaction itself.
- **Surface area to volume ratio** - The comparison of the volume inside a solid to the area exposed on the surface.

Further Reading/Supplemental Links

- Activation Energy: http://www.mhhe.com/physsci/chemistr...sh/activa2.swf
- learner.org/resources/series61.html The ***learner.org*** website allows users to view streaming videos of the Annenberg series of chemistry videos. You are required to register before you can watch the videos, but there is no charge. The website has one video that relates to this lesson called ***Molecules in Action***.
- www.vitamins-guide.net
- en.Wikipedia.org/wiki
- Observing molecules during chemical reactions helps explain the role of catalysts. Dynamic equilibrium is also demonstrated. Molecules in Action (www.learner.org/vod/vod_window.html?pid=806)
- Surface science examines how surfaces react with each other at the molecular level. On the Surface (www.learner.org/vod/vod_window.html?pid=812)

10.1: The Rate of a Chemical Reaction is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.2: The Rate of a Chemical Reaction** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.



https://chem.libretexts.org/@go/page/218526

10.2: The Idea of Dynamic Chemical Equilibrium

Template:HideTOC

Learning Objectives

- Describe the three possibilities that exist when reactants come together.
- Describe what is occurring in a system at equilibrium.

Think for a minute about sitting down to a table to eat dinner. There are three possibilities that could happen when you eat dinner. You could (1) finish your entire dinner, (2) you could not want any of it and leave it all on your plate, or (3) you could eat some of it and leave some of it. Reactions have the same possibilities. Reactions also do not always proceed all the way from start to finish. You may have reactions that (1) go to completion so that at the end the reaction vessel contains all products and only products. Some reactions (2) may not start at all so at the end the reaction vessel contains all reactants and only reactants. And some reactions (3) may start but not go to completion, that is, the reaction might start but not go completely to products. In this last case, the reaction vessel would contain some reactants and some products. In this section, we are going to take a closer look at the third type of reaction.

Reversible Reactions and Equilibrium

Consider the hypothetical reaction:

$$\mathrm{A}+\mathrm{B}\rightarrow\mathrm{C}+\mathrm{D}. \tag{10.2.1}$$

If we looked at this reaction using what we have learned, this reaction will keep going, forming C and D until A and B run out. This is what we call an "irreversible reaction" or a "reaction that goes to completion".

Some reactions, however, are **reversible**, meaning the reaction can go backwards in which products react to form reactants, so that: $\mathrm{A}+\mathrm{B}\leftarrow\mathrm{C}+\mathrm{D}$. The direction of the arrow shows that C and D are reacting to form A and B. What if the two reactions, the forward reaction and the reverse reaction, were occurring at the same time? What would this look like? If you could peer into the reaction, you would be able to find A, B, C, and D particles. A and B would react to form C and D at the same time that C and D are reacting to form A and B.

If the forward and reverse reactions are happening at the same rate, the reaction is said to be at **equilibrium** or **dynamic equilibrium**. At this point, the concentrations of A, B, C, and D are not changing (or are constant) and we would see no difference in our reaction container, but reactions are still occurring in both directions. It is important to point out that having constant amounts of reactants and products does NOT mean that the concentration of the reactants is equal to the concentration of the products. It means they are not changing. These reactions appear to have stopped before one of the reactants has run out.

Chemists use a double-headed arrow, $\rightleftharpoons$, to show that a reaction is at equilibrium. We would write the example reaction as:

$$\mathrm{A}+\mathrm{B}\rightleftharpoons\mathrm{C}+\mathrm{D}. \tag{10.2.2}$$

The arrow indicates that both directions of the reaction are happening.

Another way to think about reversible and irreversible reactions is to compare them to two types of games of tag. Reversible reactions are in many ways like a traditional game of tag: The "it" person can become "not it" and somebody who is "not it" is tagged and becomes "it". In this way it is a reversible change. It is also like a reaction at equilibrium, because overall no change is occurring. There is always a constant number of "it" people and "not it" people in the game. Also, having constant numbers of "it" and "not it" people in our game does not mean that the number of "it" people (reactants) is equal to the number of "not it" people (products). Furthermore, this is similar to equilibrium in that this game never truly ends (unless everybody gets tired of playing). The game could go on forever. We could write this as the following reversible reaction:

$$\text{"It"}\rightleftharpoons\text{"Not it"} \tag{10.2.3}$$

Irreversible reactions (those that only go in one direction from reactants to products and cannot reach a state of equilibrium) is more like a game of sharks and minnows. In sharks and minnows almost everybody starts out as a minnow. Once tagged, they become a shark. However, the difference here is that once you are a shark you are always a shark; there is no way to go back to becoming a minnow. The game continues until everybody has been tagged and becomes a shark. This is similar to irreversible reactions in that the reactants turn into products, but can't change back. Furthermore, the reaction will proceed until the reactants have been used up and there isn't any more left. We could write the reaction as:

$$\text{Minnow} \rightarrow \text{Shark} \tag{10.2.4}$$

Here's another example of a reversible reaction - dissolving salt in a beaker of water, described by the following reaction:

$$\text{NaCl(s)} \leftrightarrow \text{NaCl(aq)} \tag{10.2.5}$$

If you keep adding more and more solid salt, eventually you'll reach the point where no more salt dissolves, and the excess sits at the bottom of the beaker. At this point we have a saturated solution. Has the dissolving reaction stopped? It would appear so, but that's not the case (wouldn't ***that*** be too easy?).

What happens in our saturated solution, which has reached the point of **equilibrium**, is that both the forward

$$\text{NaCl(s)} \rightarrow \text{NaCl(aq)} \tag{10.2.6}$$

and reverse

$$\text{NaCl(aq)} \rightarrow \text{NaCl(s)} \tag{10.2.7}$$

reactions are still going on, but at the same ***rate***. This in effect cancels out any observable, or measurable, changes in our system. At the same rate that solid NaCl produces aqueous NaCl (dissolved salt), the dissolved salt is recrystallizing to form more solid NaCl.

- Equilibrium is the state at which the rate of the forward reaction equals the rate of the reverse reaction.
- At the point of equilibrium, no more measurable or observable changes in the system can be noted.

It is important for you to understand that equilibrium means the ***rates*** of the forward and reverse reactions are equal; it does ***not*** mean that there are equal amount of reactants and products present at equilibrium.

For example, the following reaction was allowed to come to the point of equilibrium, and concentrations of all reaction participants were measured at that time:

$$\text{H}_2\text{(g)} + \text{I}_2\text{(g)} \leftrightarrow 2\,\text{HI(g)}$$

At equilibrium:	$[\text{H}_2] = 0.022\,\text{M}$ $[\text{I}_2] = 0.022\,\text{M}$ $[\text{HI}] = 0.156\,\text{M}$

For this particular reversible reaction, there is more HI at equilibrium (0.156 M) than there is of H_2 and I_2 (both at 0.022M). We say the product side of the reaction is favored.

> *Equilibrium does not mean equal amounts at equilibrium!*

Here's something to help you understand about how equilibrium works. Imagine yourself on a escalator that is going down. You start at the top (reactants) and end up at the bottom (products). But when you are partway down you start walking up the escalator as it continues going down. If you match your rate of walking up to the same rate that the escalator is going down, you make no progress and appear to be at a standstill. To an observer it would look as if you and the escalator had come to a stop, but actually both upward and downward movements continue.


Equilibrium is **dynamic** - both forward and reverse reactions continue, even though the reaction appears to have stopped. And this equilibrium does not need to occur right in the middle of two floors - you could be near the bottom, near the top, or anywhere in between when you carry out your reverse process.

In order for a reversible reaction to reach the point of equilibrium, the reaction must be carried out in a closed system - no additional reactants can be added or products removed. If, in our last example, the product HI was removed as it formed, the reaction would never reach the point of equilibrium - instead H_2 and I_2 would continue to react to produce HI until one or both of the reactants was used up.

> If reactants are constantly being added, and products removed as they form, the system would appear to be at equilibrium because to an outside observer it would appear that the reaction has stopped, but that would not be the case. This situation - with new material constantly being added as products are removed - is called a **steady state system**. A factory with an assembly line is a steady state system - new raw materials are constantly being added; finished products are removed. A campfire with wood being added to the fire is another steady state system. Be careful not to confuse steady state with equilibrium.

How do the rates of the forward and reverse reactions as the reaction heads towards equilibrium (before it reaches equilibrium?)

If we start our above reaction with H_2 and I_2, and with no HI, the two gases will react at a certain rate. But remember that the rate of a reaction slows down over time, as the reactants get used up (and lower their concentration). Eventually, however, the amount of the product HI increases, and it will begin producing H_2 and I_2. Thus the rate of the reverse reaction starts out slowly (there is no HI present), but will speed up as the concentration of HI increases. Eventually both rates will level off (not always to the same level as shown by this example, however):



Chemists have found that there is a mathematical relationship that exists between the concentration of the reactants and products once equilibrium has reached that is independent of the initial concentration of the participants. For any general reaction:

$$\mathrm{aA} + \mathrm{bB} \rightleftharpoons \mathrm{cD} + \mathrm{dD} \tag{10.2.8}$$

an equilibrium constant expression can be written as:

$$\mathrm{K_{eq}} = \frac{[\mathrm{C}]^{\mathrm{c}} \times [\mathrm{D}]^{\mathrm{d}}}{[\mathrm{A}]^{\mathrm{a}} \times [\mathrm{B}]^{\mathrm{b}}} \tag{10.2.9}$$

This mathematical relationship exists for all equilibrium systems, and produces a constant ratio called the ***equilibrium constant, K_{eq}*** .

law of mass-action.

Equation 10.2.9 is sometimes called the Law of Mass-Action.

This relationship will be very important to us for the next few units, so it is important that you understand how to set this relationship up and what it tells us about an equilibrium system.

The products of the reaction (C and D) are placed in the numerator, and their concentrations are raised to the power of the coefficients from the balanced equation. The reactants (A and B) are placed in the denominator, with their concentrations raised to the power of *their* coefficients.

Example 10.2.1

For the reaction between hydrogen and iodine gas to produce hydrogen iodide:

$$H_2(g) + I_2(g) \rightleftharpoons 2\,HI(g) \tag{10.2.10}$$

the equilibrium constant expression will be:

$$K_{eq} = \frac{[HI]^2}{[H_2] \times [I_2]} \tag{10.2.11}$$

Using the example we examined in our last section, equilibrium concentrations for each substance were measured at equilibrium and found to be:

At equilibrium:	$[H_2] = 0.022\ M$ $[I_2] = 0.022\ M$ $[HI] = 0.156\ M$

We substitute these values into our equilibrium expression and solve for K_{eq}:

$$K_{eq} = \frac{[HI]^2}{[H_2] \times [I_2]} = \frac{(0.156)^2}{(0.022)(0.022)} = 50.3 \tag{10.2.12}$$

The value of K_{eq}, which has no units, is a constant for any particular reaction, and its value does not change unless the temperature of the system is changed. It does not depend on the initial concentrations used to reach the point of equilibrium.

For example, the following data were obtained for equilibrium concentrations of H_2, I_2 and HI, and the value of K_{eq} was calculated for each trial:

Trial	$[HI]$	$[H_2]$	$[I_2]$	K_{eq}
1	0.156	0.0220	0.0220	50.3
2	0.750	0.106	0.106	50.1
3	1.00	0.820	0.0242	50.4
4	1.00	0.0242	0.820	50.4
5	1.56	0.220	0.220	50.3

Aside from accounting for slight experimental variation between trials, the value for K_{eq} is the same despite differences in equilibrium concentrations for the individual participants.

There is one other important point to make at this time.

K_{eq} relates the concentrations of products to reactants at equilibrium.

For aqueous solutions, concentration is often measured as mol · L^{-1}. For gases, concentration is often measured as partial pressure in atmospheres (atm). We have not yet gotten to our unit on gases, and therefore the concept on partial pressure. However, this does not impact our ability to understand equilibrium, since equilibrium is a ratio of concentrations. Whether the concentrations are in molarity for aqueous solutions, or atmospheres for gases.

The concentrations of both aqueous solutions and gases change during the progress of a reaction. For reactions involving a solid or a liquid, while the ***amounts*** of the solid or liquid will change during a reaction, their ***concentrations*** (much like their densities) will ***not*** change during the reaction. Instead, their values will remain constant. Because they are constant, their values **are not included in the equilibrium constant expression**.

Example 10.2.2

For example, consider the reaction showing the formation of solid calcium carbonate from solid calcium oxide and carbon dioxide gas:

$$CaO(s) + CO_2(g) \leftrightarrow CaCO_3(s) \tag{10.2.13}$$

The equilibrium constant for this reaction is (before modification)

$$K_{eq} = \frac{[CaCO_3]}{[CaO] \times [CO_2]} \tag{10.2.14}$$

But we remove those participants whose state is either a solid or a liquid, which leaves us with the following equilibrium constant expression:

$$K_{eq} = \frac{1}{[CO_2]} \tag{10.2.15}$$

Lesson Summary

- There are a few possible ways a reaction can go: It can go to completion ($\text{reactants} \rightarrow \text{products}$); it can occur but not go to completion. Instead is would reach chemical equilibrium ($\text{reactants} \rightleftharpoons \text{products}$).
- Chemical equilibrium occurs when the number of particles becoming products is equal to the number of particles becoming reactants.
- A dynamic equilibrium is a state where the rate of the forward reaction is equal to the rate of the reverse reaction.

Vocabulary

- Equilibrium: A state that occurs when the rate of the forward reaction is equal to the rate of the reverse reaction.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

-

10.2: The Idea of Dynamic Chemical Equilibrium is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

10.3: The Equilibrium Constant - A Measure of How Far a Reaction Goes

Learning Objectives

- Write equilibrium constant expressions.
- Use equilibrium constant expressions to solve for unknown concentrations.
- Use known concentrations to solve for the equilibrium constants.
- Explain what the value of K means in terms of relative concentrations of reactants and products.

In the previous section, you learned about reactions that can reach a state of equilibrium, in which the concentration of reactants and products aren't changing. If these amounts are changing, we should be able to make a relationship between the amount of product and reactant when a reaction reaches equilibrium.

The Equilibrium Constant

Equilibrium reactions are those that do not go to completion, but are in a state where the reactants are reacting to yield products and the products are reacting to produce reactants. In a reaction at equilibrium, the equilibrium concentrations of all reactants and products can be measured. The **equilibrium constant (K)** is a mathematical relationship that shows how the concentrations of the products vary with the concentration of the reactants. Sometimes, subscripts are added to the equilibrium constant symbol K, such as K_{eq}, K_{c}, K_{p}, K_{a}, K_{b}, and K_{sp}. These are all equilibrium constants and are subscripted to indicate special types of equilibrium reactions.

There are some rules about writing equilibrium constant expressions that need to be learned:

1. Concentrations of products are multiplied on the top of the expression. Concentrations of reactants are multiplied together on the bottom.
2. Coefficients in the equation become exponents in the equilibrium constant expression.
3. Solids, liquids, and solvents are assigned a value of 1, so their concentrations do not affect the value of K.

Example 10.3.1

Write the equilibrium constant expression for:

$$\text{CO}\,(g) + 3\text{H}_2\,(g) \rightleftharpoons \text{CH}_4\,(g) + \text{H}_2\text{O}\,(g)$$

Solution

$$K = \frac{[\text{CH}_4]\,[\text{H}_2\text{O}]}{[\text{CO}]\,[\text{H}_2]^3}$$

*Note that the coefficients become exponents. Also, note that the concentrations of products in the numerator are *multiplied*. The same is true of the reactants in the denominator.

Example 10.3.2

Write the equilibrium constant expression for:

$$2\text{TiCl}_3\,(s) + 2\text{HCl}\,(g) \rightleftharpoons 2\text{TiCl}_4\,(s) + \text{H}_2\,(g)$$

Solution

$$K = \frac{[\text{H}_2]}{[\text{HCl}]^2}$$

*Note that the solids have a value of 1, and multiplying or dividing by 1 does not change the value of K.

Example 10.3.2

Write the equilibrium constant expression for:

$$\mathrm{P_4}\,(s) + 6\mathrm{Cl_2}\,(g) \rightleftharpoons 4\mathrm{PCl_3}\,(s)$$

Solution

$$K = \frac{1}{[\mathrm{Cl_2}]^6}$$

*Note that the only product is a solid, which is defined to have a value of 1. That leaves just 1 on top in the numerator.

Example 10.3.3

Write the equilibrium constant expression for:

$$\mathrm{H_2O}\,(l) \rightleftharpoons \mathrm{H^+}\,(aq) + \mathrm{OH^-}\,(aq)$$

Solution

$$K = [\mathrm{H^+}]\,[\mathrm{OH^-}]$$

*Note that the water is the solvent, and thus has a value of 1. Dividing by 1 does not change the value of K.

Equilibrium Constant Expressions

The equilibrium constant value is the ratio of the concentrations of the products over the reactants. This means that we can use the value of K to predict whether there are more products or reactants at equilibrium for a given reaction. What can the value of K_{eq} tell us about a reaction?

- **If K_{eq} is very large**, the concentration of the products is much greater than the concentration of the reactants. The reaction essentially "goes to completion"; all, or most of, the reactants are used up to form the products.
- **If K_{eq} is very small**, the concentration of the reactants is much greater than the concentration of the products. The reaction does not occur to any great extent—most of the reactants remain unchanged, and there are few products produced.
- **When K_{eq} is not very large or very small** (close to a value of 1) then roughly equal amounts of reactants and products are present at equilibrium.

Here are some examples to consider:

Reaction	Chemical Equations	Equilibrium Constant
the decomposition of ozone, $\mathrm{O_3}$	$2\,\mathrm{O_3(g)} \rightleftharpoons 3\,\mathrm{O_2(g)}$	$K_{eq} = 2.0 \times 10^{57}$
	K_{eq} is very large, indicating that mostly $\mathrm{O_2}$ is present in an equilibrium system, with very little $\mathrm{O_3}$.	
production of nitrogen monoxide	$\mathrm{N_2(g) + O_2(g)} \rightleftharpoons 2\,\mathrm{NO(g)}$	$K_{eq} = 1.0 \times 10^{-25}$
	Very little NO is produced by this reaction; $\mathrm{N_2}$ and $\mathrm{O_2}$ do not react readily to produce NO (this is lucky for us—otherwise we would have little oxygen to breath in our atmosphere!).	
reaction of carbon monoxide and water	$\mathrm{CO(g) + H_2O(g)} \rightleftharpoons \mathrm{CO_2(g) + H_2(g)}$	$K_{eq} = 5.09\,(\text{at } 700\text{ K})$
	The concentrations of the reactants are very close to the concentrations of the products at equilibrium.	

If the equilibrium constant is 1 or nearly 1, it indicates that the molarities of the reactants and products are about the same. If the equilibrium constant value is a large number, like 100, or a very large number, like 1×10^{15}, it indicates that the products (numerator) are a great deal larger than the reactants. This means that at equilibrium, the great majority of the material is in the form of products and it is said that the "products are strongly favored". If the equilibrium constant is small, like 0.10, or very small,

like 1×10^{-12}, it indicates that the reactants are much larger than the products and the reactants are strongly favored. With large K values, most of the material at equilibrium is in the form of products and with small K values, most of the material at equilibrium is in the form of the reactants.

The equilibrium constant expression is an equation that we can use to solve for K or for the concentration of a reactant or product.

Example 10.3.4

Determine the value of K for the reaction

$$\mathrm{SO_2}\,(g) + \mathrm{NO_2}\,(g) \rightleftharpoons \mathrm{SO_3}\,(g) + \mathrm{NO}\,(g)$$

when the equilibrium concentrations are: $[\mathrm{SO_2}] = 1.20\ \mathrm{M}$, $[\mathrm{NO_2}] = 0.60\ \mathrm{M}$, $[\mathrm{NO}] = 1.6\ \mathrm{M}$, and $[\mathrm{SO_3}] = 2.2\ \mathrm{M}$.

Solution

Step 1: Write the equilibrium constant expression:

$$K = \frac{[\mathrm{SO_3}]\,[\mathrm{NO}]}{[\mathrm{SO_2}]\,[\mathrm{NO_2}]}$$

Step 2: Substitute in given values and solve:

$$K = \frac{(2.2)\,(1.6)}{(1.20)\,(0.60)} = 4.9$$

Example 10.3.5

Consider the following reaction:

$$\mathrm{CO}\,(g) + \mathrm{H_2O}\,(g) \rightleftharpoons \mathrm{H_2}\,(g) + \mathrm{CO_2}\,(g)$$

with $K = 1.34$. If the $[\mathrm{H_2O}] = 0.100\ \mathrm{M}$, $[\mathrm{H_2}] = 0.100\ \mathrm{M}$, and $[\mathrm{CO_2}] = 0.100\ \mathrm{M}$ at equilibrium, what is the equilibrium concentration of CO?

Solution

Step 1: Write the equilibrium constant expression:

$$K = \frac{[\mathrm{H_2}]\,[\mathrm{CO_2}]}{[\mathrm{CO}]\,[\mathrm{H_2O}]}$$

Step 2: Substitute in given values and solve:

$$1.34 = \frac{(0.100)\,(0.100)}{[\mathrm{CO}]\,(0.100)}$$

Solving for $[\mathrm{CO}]$, we get: $[\mathrm{CO}] = 0.0746\ \mathrm{M}$

Summary

- The equilibrium constant expression is a mathematical relationship that shows how the concentrations of the products vary with the concentration of the reactants.
- If the value of K is greater than 1, the products in the reaction are favored. If the value of K is less than 1, the reactants in the reaction are favored. If K is equal to 1, neither reactants nor products are favored.

Vocabulary

- **Equilibrium constant** (K) - A mathematical ratio that shows the concentrations of the products divided by the concentrations of the reactants.

Contributions & Attributions

-
- Modified by Tom Neils (Grand Rapids Community College)

10.3: The Equilibrium Constant - A Measure of How Far a Reaction Goes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.4: The Equilibrium Constant - A Measure of How Far a Reaction Goes** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

10.4: Disturbing a Reaction at Equilibrium- Le Châtelier's Principle

Learning Objectives

- State Le Chatelier's Principle.
- Describe the effect of concentration on an equilibrium system.
- Describe the effect of temperature as a stress on an equilibrium system.

When a reaction has reached equilibrium with a given set of conditions, if the conditions are not changed, the reaction will remain at equilibrium forever. The forward and reverse reactions continue at the same equal and opposite rates and the macroscopic properties remain constant.

It is possible, however, to disturb that equilibrium by changing conditions. For example, you could increase the concentration of one of the products, or decrease the concentration of one of the reactants, or change the temperature. When a change of this type is made within a reaction at equilibrium, the reaction is no longer in equilibrium. When you alter something in a reaction at equilibrium, chemists say that you put **stress** on the equilibrium. When this occurs, the reaction will no longer be in equilibrium and the reaction itself will begin changing the concentrations of reactants and products until the reaction comes to a new position of equilibrium. How a reaction will change when a stress is applied can be explained and predicted. That is the topic of this section.

Le Chatelier's Principle

Think back to our escalator example, with you walking up a downward moving escalator. With the rate of the moving stairs and your walking evenly matched, you appear to be at a standstill. But what happens if the escalator begins moving just a little faster? If you want to maintain the same position you had, at some specific point between the bottom and the top of the stairs, you'll also need to make some adjustments. Chemical systems at equilibrium tend to make these adjustments as well.

In the late 1800's, a chemist by the name of Henry-Louis Le Chatelier was studying stresses that were applied to chemical equilibria. He formulated a principle from this research and, of course, the principle is called Le Chatelier's Principle. **Le Chatelier's Principle** states that when a stress is applied to a system at equilibrium, the equilibrium will shift in a direction to partially counteract the stress and once again reach equilibrium. Le Chatelier's principle is not an explanation of what happens on the molecular level to cause the equilibrium shift, it is simply a quick way to determine which way the reaction will run in response to a stress applied to the system at equilibrium.

Le Chatelier's Principle

If a system at **equilibrium** is subjected to an external stress, the equilibrium will shift to **minimize** the effects of that stress.

Equilibrium is all about rates—the rate of the forward reaction is equal to the rate of the reverse reaction. External stresses are factors that will cause the rate of either the forward or reverse reaction to change, throwing the system out of balance. Le Chatelier's Principle allows us to predict how this will affect our system.

In our unit on Kinetics we examined factors that influenced reaction rates. Recall these factors:

1. concentration
2. pressure and volume
3. temperature
4. catalysts

We will see how changing these factors affects a system at equilibrium.

Effect of Concentration Changes on a System at Equilibrium

For instance, if a stress is applied by increasing the concentration of a reactant, the reaction will adjust in such a way that the reactants and products can get back to equilibrium. In the case of too much reactant, the reaction will use up some of the reactant to make more product. It is said in this scenario that the reaction "shifts to the products" or "shifts to the right". If the concentration of a product is increased, there is an opposite effect. The reaction will use up some of the product to make more reactant. The reaction "shifts to the reactants" or "shifts to the left".

What if some reactant or product is removed? If a stress is applied by lowering a reactant concentration, the reaction will try to replace some of the missing reactant. It uses up some of the product to make more reactant, and the reaction "shifts to the reactants". If a stress is applied by reducing the concentration of a product, the equilibrium position will shift toward the products.

10.4: Disturbing a Reaction at Equilibrium- Le Châtelier's Principle is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.7: Disturbing a Reaction at Equilibrium- Le Châtelier's Principle** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

10.5: The Effect of a Concentration Change on Equilibrium

Consider the following system under equilibrium:

$$\underbrace{\mathrm{Fe}^{3+}(\mathrm{aq})}_{\text{colorless}} + \underbrace{\mathrm{SCN}^{-}(\mathrm{aq})}_{\text{colorless}} \rightleftharpoons \underbrace{\mathrm{FeSCN}^{2+}(\mathrm{aq})}_{\text{red}}$$

If more Fe^{3+} is added to the reaction, what will happen?

According to Le Chatelier's Principle, the system will react to minimize the stress. Since Fe^{3+} is on the reactant side of this reaction, the rate of the forward reaction will increase in order to "use up" the additional reactant. This will cause the equilibrium to **shift to the right**, producing more $FeSCN^{2+}$. For this particular reaction, we will be able to see that this has happened, as the solution will become a darker red color.

There are a few different ways to state what happens here when more Fe^{3+} is added, all of which have the same meaning:

- equilibrium shifts to the right
- equilibrium shifts to the product side
- the forward reaction is favored

What changes does this cause in the concentrations of the reaction participants?

Changes within reaction participants

Fe^{3+}	Since this is what was added to cause the stress, the concentration of Fe^{3+} will increase. (A shorthand way to indicate this: $[\mathrm{Fe}]^{3+} \uparrow$ (Reminder: the square brackets represent "concentration")
$\mathrm{SCN}^{-}(\mathrm{aq})$	Equilibrium will shift to the right, which will use up the reactants. The concentration of $\mathrm{SCN}^{-}(\mathrm{aq})$ will decrease $[\mathrm{SCN}]^{-} \downarrow$ as the rate of the forward reaction increases.
FeSCN^{2+}	When the forward reaction rate increases, more products are produced, and the concentration of FeSCN^{2+} will increase. $[\mathrm{FeSCN}]^{2+} \uparrow$

How about the value of K_{eq}? Notice that the concentration of some reaction participants have increased, while others have decreased. Once equilibrium has re-established itself, the value of K_{eq} will be unchanged.

> *The value of K_{eq} does not change when changes in concentration cause a shift in equilibrium.*

What if more $FeSCN^{2+}$ is added?

Again, equilibrium will shift to use up the added substance. In this case, equilibrium will shift to favor the *reverse* reaction, since the reverse reaction will use up the additional $FeSCN^{2+}$.

- equilibrium shifts to the left
- equilibrium shifts to the reactant side
- the reverse reaction is favored

How do the concentrations of reaction participants change?

Change of concentrations of reaction participants when adding substance

Fe^{3+}	$[\mathrm{Fe}]^{3+} \uparrow$ as the reverse reaction is favored
$\mathrm{SCN}^{-}(\mathrm{aq})$	$[\mathrm{SCN}]^{-} \uparrow$ as the reverse reaction is favored
FeSCN^{2+}	$[\mathrm{FeSCN}]^{2+} \uparrow$ because this is the substance that was added

Concentration can also be changed by **removing** a substance from the reaction. This is often accomplished by adding another substance that reacts (in a side reaction) with something already in the reaction.

Let's remove SCN^- from the system (perhaps by adding some Pb^{2+} ions—the lead(II) ions will form a precipitate with SCN^-, removing them from the solution). What will happen now? Equilibrium will shift to **replace** SCN^-—the reverse reaction will be favored because that is the direction that produces more SCN^-.

- equilibrium shifts to the left
- equilibrium shifts to the reactant side
- the reverse reaction is favored

How do the concentrations of reaction participants change?

Change of concentrations of reaction participants when removing a substance

Fe^{3+}	$[Fe]^{3+}$ ↑ as the reverse reaction is favored
SCN^-	$[SCN]^-$ ↑ as the reverse reaction is favored (but also ↓ because it was removed)
$FeSCN^{2+}$	$[FeSCN]^{2+}$ ↑ because this is the substance that was added

---

10.5: The Effect of a Concentration Change on Equilibrium is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.8: The Effect of a Concentration Change on Equilibrium** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

10.6: The Effect of a Volume Change on Equilibrium

Changing the pressure or volume of a container enclosing an equilibrium system will only affect the reaction if gases are present.

You may remember from earlier chemistry classes that equal volumes of gases contain an equal number of particles and, under standard conditions of temperature and pressure (STP), one mole of gas occupies a volume of 22.4 L. This is known as the molar volume of gases. So, two moles of any gas will occupy a volume of 44.8 L and one-half mole would occupy 11.2 L.

How does changing pressure and volume affect equilibrium systems?

- If you increase the pressure of a system at equilibrium (typically by reducing the volume of the container), the stress will best be reduced by reaction that favors the side with the ***fewest*** moles of gas, since fewer moles will occupy the smallest volume.
- Conversely, if you decrease the pressure (by increasing the volume of the container), equilibrium will shift to favor the side with the ***most*** moles of gas, since more moles will occupy a greater volume.
- If both sides of the equation have the same number of moles of gas, then there will be no change in the position of equilibrium.

When considering the effect of changing volume or pressure on equilibrium systems, be sure to only count the number of moles of **gases** on each side of the equation. Solids, liquids, and aqueous solutions will not be affected by changing pressure and volume.

Example 10.6.1

Predict the effect on equilibrium when the pressure is increased for the following reaction:

$$\mathrm{N_2O_4(g) \rightleftharpoons 2\,NO_2(g)}$$

Solution

The reactant side of the equation has 1 mole of a gas; the product side has 2 moles. Increasing the pressure favors the side with the fewest moles of gas, so the equilibrium will shift to the left (the reverse reaction will be favored).

10.6: The Effect of a Volume Change on Equilibrium is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.9: The Effect of a Volume Change on Equilibrium** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

10.7: The Effect of Temperature Changes on Equilibrium

Learning Objectives

- Explain the following laws within the Ideal Gas Law

When temperature is the stress that affects a system at equilibrium, there are two important consequences:

- an increase in temperature will favor that reaction direction that ***absorbs*** heat (i.e. the endothermic reaction)
- the value of K_{eq} will change

Consider the following equilibrium system

$$\mathrm{N_2O_4(g) \leftrightarrow 2\,NO_2(g)} \tag{10.7.1}$$

with $\Delta H^{\circ} = 58 \cdot 0 \text{ kJ}$

Delta H is a measurement showing heat was gained by the reaction. We will cover heat and energy soon.

We see by the sign of ΔH° that the forward reaction is endothermic. Heat is absorbed (required as a reactant) when the reaction proceeds as

$$\mathrm{N_2O_4(g) \rightarrow 2\,NO_2(g)} \tag{10.7.2}$$

By adding more heat, equilibrium will shift to use up the additional heat, thus favoring this forward direction.

Why will K_{eq} change, when it did not change when concentration, pressure, and volume were the applied stresses?

When temperature changes cause an equilibrium to shift, one entire side of the reaction equation is favored over the other side. Mathematically, this will alter the value of K_{eq} as follows:

$$K_{eq} = \frac{[\text{products}]}{[\text{reactants}]} \tag{10.7.3}$$

if the forward reaction is favored	more products are produced; fewer reactants	K_{eq} will increase
if the reverse reaction is favored	fewer products; more reactants	K_{eq} will decrease

So in our example given above, increasing the temperature will favor the forward direction. The value of K_{eq} will increase. **Removing heat (making the system colder) will favor the exothermic reaction** - the exothermic reaction ***releases heat*** to the surroundings, thus "replacing" the heat that has been removed.

10.7: The Effect of Temperature Changes on Equilibrium is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

10.8: The Path of a Reaction and the Effect of a Catalyst

The addition of a catalyst to an equilibrium system is a final stress factor. We consider how adding a catalyst affects the following:

$$\mathrm{N_2(g) + O_2(g) \rightleftharpoons 2\,NO(g)}$$

Adding a catalyst to this, or any other equilibrium system, will ***not*** affect the position of an equilibrium. A catalyst speeds up both the forward and the reverse reactions, so there is no uneven change in reaction rates. Generally, a catalyst will help a reaction to reach the point of equilibrium *sooner*, but it will not affect the equilibrium otherwise.

10.8: The Path of a Reaction and the Effect of a Catalyst is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **15.12: The Path of a Reaction and the Effect of a Catalyst** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

CHAPTER OVERVIEW

11: Energy and Chemical Processes

Energy is a vital component of the world around us. Nearly every physical and chemical process, including all the chemical reactions discussed in previous chapters, occurs with a simultaneous energy change. In this chapter, we will explore the nature of energy and how energy and chemistry are related.

11.1: Prelude to Energy and Chemical Processes
11.2: Energy and Its Units
11.3: Heat
11.4: Phase Changes
11.5: Bond Energies and Chemical Reactions
11.6: The Energy of Biochemical Reactions
11.E: Energy and Chemical Processes (Exercises)
11.S: Energy and Chemical Processes (Summary)

Template:HideTOC

11: Energy and Chemical Processes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

11.1: Prelude to Energy and Chemical Processes

Metabolism is the collective term for the chemical reactions that occur in cells and provide energy to keep cells alive. Some of the energy from metabolism is in the form of heat, and some animals use this heat to regulate their body temperatures. Such *warm-blooded* animals are called *endotherms*. In endotherms, problems with metabolism can lead to fluctuations in body temperature. When humans get sick, for instance, our body temperatures can rise higher than normal; we develop a fever. When food is scarce (especially in winter), some endotherms go into a state of controlled decreased metabolism called *hibernation*. During hibernation, the body temperatures of these endotherms actually decrease. In hot weather or when feverish, endotherms will pant or sweat to rid their bodies of excess heat.

Table 11.1.1: Average Body Temperatures of Selected Endotherms

Endotherm	Body Temperature (°F)	Body Temperature (°C)
bird	up to 110	up to 43.5
cat	101.5	38.6
dog	102	38.9
horse	100.0	37.8
human	98.6	37.0
pig	102.5	39.2

Ectotherms, sometimes called *cold-blooded* animals, do not use the energy of metabolism to regulate body temperature. Instead, they depend on external energy sources, such as sunlight. Fish, for example, will seek out water of different temperatures to regulate body temperature. The amount of energy available is directly related to the metabolic rate of the animal. When energy is scarce, ectotherms may also hibernate.

The connection between metabolism and body temperature is a reminder that energy and chemical reactions are intimately related. A basic understanding of this relationship is especially important when those chemical reactions occur within our own bodies.

11.1: Prelude to Energy and Chemical Processes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.0: Prelude to Energy and Chemical Processes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.2: Energy and Its Units

Learning Objectives

- To define *energy* and *heat*.
- To relate calories to nutrition and exercise.

Energy is the ability to do work. You can understand what this means by thinking about yourself when you feel "energetic." You feel ready to go—to jump up and get something done. When you have a lot of energy, you can perform a lot of work. By contrast, if you do not feel energetic, you have very little desire to do much of anything. This description is not only applicable to you but also to all physical and chemical processes. The quantity of work that can be done is related to the quantity of energy available to do it.

Energy can be transferred from one object to another if the objects have different temperatures. The transfer of energy due to temperature differences is called heat. For example, if you hold an ice cube in your hand, the ice cube slowly melts as energy in the form of heat is transferred from your hand to the ice. As your hand loses energy, it starts to feel cold.

Because of their interrelationships, energy, work, and heat have the same units. The SI unit of energy, work, and heat is the joule (J). A joule is a tiny amount of energy. For example, it takes about 4 J to warm 1 mL of H_2O by 1°C. Many processes occur with energy changes in thousands of joules, so the kilojoule (kJ) is also common. Another unit of energy, used widely in the health professions and everyday life, is the calorie (cal). The calorie was initially defined as the amount of energy needed to warm 1 g of H_2O by 1°C, but in modern times, the calorie is related directly to the joule, as follows:

$$1\ cal = 4.184\ J$$

We can use this relationship to convert quantities of energy, work, or heat from one unit to another.

> *Although the joule is the proper SI unit for energy, we will use the calorie or the kilocalorie (or Calorie) in this chapter because they are widely used by health professional*

Example 11.2.1

The energy content of a single serving of bread is 70.0 Cal. What is the energy content in calories? In joules?

Solution

This is a simple conversion-factor problem. Using the relationship 1 Cal = 1,000 cal, we can answer the first question with a one-step conversion:

$$70.0\ \text{Cal} \times \frac{1,000\ \text{cal}}{1\ \text{Cal}} = 70,000\ \text{cal}$$

Then we convert calories into joules

$$70,000\ \text{cal} \times \frac{4.184\ \text{J}}{1\ \text{cal}} = 293,000\ \text{J}$$

and then kilojoules

$$293,000\ \text{J} \times \frac{1\ \text{kJ}}{1,000\ \text{J}} = 293\ \text{kJ}$$

The energy content of bread comes mostly from carbohydrates.

cise

The energy content of one cup of honey is 1,030 Cal. What is its energy content in calories and joules?

Answer

1,030,000 (1.03 x 10^6) cal; 4,309,520 (4.31 x 10^6) J

The calorie is used in nutrition to express the energy content of foods. However, because a calorie is a rather small quantity, nutritional energies are usually expressed in **kilocalories (kcal)**, also called **Calories (capitalized; Cal)**. For example, a candy bar may provide 120 Cal (nutritional calories) of energy, which is equal to 120,000 cal. Figure 11.2.1 shows an example.




Figure 11.2.1: Figure 11.2.1: Nutritional Energy. A sample nutrition facts label, with instructions from the U.S. Food and Drug Administration. from Wikipedia.

The caloric content of foods is determined by analyzing the food for protein, carbohydrate, fat, water, and “minerals” and then calculating the caloric content using the average values for each component that produces energy (**9 Cal/g for fats, 4 Cal/g for carbohydrates and proteins, and 0 Cal/g for water and minerals**). An example of this approach is shown in Table 11.2.1 for a slice of roast beef.

Table 11.2.1: Approximate Composition and Fuel Value of an 8 oz Slice of Roast Beef

Composition	**Calories**
97.5 g of water	× 0 Cal/g = 0
58.7 g of protein	× 4 Cal/g = 235
69.3 g of fat	× 9 Cal/g = 624
0 g of carbohydrates	× 4 Cal/g = 0
1.5 g of minerals	× 0 Cal/g = 0
Total mass: 227.0 g	**Total calories: about 900 Cal**

The compositions and caloric contents of some common foods are given in 11.2.2

Table 11.2.2: Approximate Compositions and Fuel Values of Some Common Foods

Food (quantity)	**Approximate Composition (%)**				**Food Value (Cal/g)**	**Calories**
	Water	**Carbohydrate**	**Protein**	**Fat**		
beer (12 oz)	92	3.6	0.3	0	0.4	150
coffee (6 oz)	100	~0	~0	~0	~0	~0
milk (1 cup)	88	4.5	3.3	3.3	0.6	150

Food (quantity)	Approximate Composition (%)				Food Value (Cal/g)	Calories
egg (1 large)	75	2	12	12	1.6	80
butter (1 tbsp)	16	~0	~0	79	7.1	100
apple (8 oz)	84	15	~0	0.5	0.6	125
bread, white (2 slices)	37	48	8	4	2.6	130
brownie (40 g)	10	55	5	30	4.8	190
hamburger (4 oz)	54	0	24	21	2.9	326
fried chicken (1 drumstick)	53	8.3	22	15	2.7	195
carrots (1 cup)	87	10	1.3	~0	0.4	70

Because the Calorie represents such a large amount of energy, a few of them go a long way. An average 73 kg (160 lb) person needs about 67 Cal/h (1600 Cal/day) to fuel the basic biochemical processes that keep that person alive. This energy is required to maintain body temperature, keep the heart beating, power the muscles used for breathing, carry out chemical reactions in cells, and send the nerve impulses that control those automatic functions. Physical activity increases the amount of energy required but not by as much as many of us hope. A moderately active individual requires about 2500−3000 Cal/day; athletes or others engaged in strenuous activity can burn 4000 Cal/day. Any excess caloric intake is stored by the body for future use, usually in the form of fat, which is the most compact way to store energy. When more energy is needed than the diet supplies, stored fuels are mobilized and oxidized. We usually exhaust the supply of stored carbohydrates before turning to fats, which accounts in part for the popularity of low-carbohydrate diets.

To Your Health: Energy Expenditures

Most health professionals agree that exercise is a valuable component of a healthy lifestyle. Exercise not only strengthens the body and develops muscle tone but also expends energy. After obtaining energy from the foods we eat, we need to expend that energy somehow, or our bodies will store it in unhealthy ways (e.g., fat). Like the energy content in food, the energy expenditures of exercise are also reported in kilocalories, usually kilocalories per hour of exercise. These expenditures vary widely, from about 440 kcal/h for walking at a speed of 4 mph to 1,870 kcal/h for mountain biking at 20 mph. Table 11.2.3 lists the energy expenditure for a variety of exercises.

Table 11.2.3: Energy Expenditure of a 180-Pound Person during Selected Exercises

Exercise	Energy Expended (kcal/h)
aerobics, low-level	325
basketball	940
bike riding, 20 mph	830
golfing, with cart	220
golfing, carrying clubs	425
jogging, 7.5 mph	950
racquetball	740
skiing, downhill	520
soccer	680
walking upstairs	1,200

Exercise	Energy Expended (kcal/h)
yoga	280

Because some forms of exercise use more energy than others, anyone considering a specific exercise regimen should consult with his or her physician first.

Summary

Energy is the ability to do work. Heat is the transfer of energy due to temperature differences. Energy and heat are expressed in units of joules.

11.2: Energy and Its Units is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.1: Energy and Its Units** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.3: Heat

Learning Objectives

- To relate heat transfer to temperature change.

Heat is a familiar manifestation of energy. When we touch a hot object, energy flows from the hot object into our fingers, and we perceive that incoming energy as the object being “hot.” Conversely, when we hold an ice cube in our palms, energy flows from our hand into the ice cube, and we perceive that loss of energy as “cold.” In both cases, the temperature of the object is different from the temperature of our hand, so we can conclude that differences in temperatures are the ultimate cause of heat transfer.

Suppose we consider the transfer of heat , represented by the letter q, from the opposite perspective—namely, what happens to a system that gains or loses heat? Generally, the system’s temperature changes. (We will address a few exceptions later.) The greater the original temperature difference, the greater the transfer of heat, and the greater the ultimate temperature change. The relationship between the amount of heat transferred and the temperature change can be written as

$$\text{heat} \propto \Delta T \tag{11.3.1}$$

where $\propto$ means “is proportional to” and ΔT is the change in temperature of the system. Any change in a variable is always defined as “the final value minus the initial value” of the variable, so ΔT is $T_{\text{final}} - T_{\text{initial}}$. In addition, the greater the mass of an object, the more heat is needed to change its temperature. We can include a variable representing mass (m) to the proportionality as follows:

$$\text{heat} \propto m\Delta T \tag{11.3.2}$$

To change this proportionality into an equality, we include a proportionality constant. The proportionality constant is called the specific heat and is commonly symbolized by c:

$$\text{q} = mc\Delta T \tag{11.3.3}$$

Every substance has a characteristic specific heat, which is reported in units of cal/g•°C or cal/g•K, depending on the units used to express ΔT. The specific heat of a substance is the amount of energy that must be transferred to or from 1 g of that substance to change its temperature by 1°. Table 11.3.1 lists the specific heats for various materials.

Table 11.3.1: Specific Heats of Selected Substances

Substance	c (cal/g•°C)
aluminum (Al)	0.215
aluminum oxide (Al_2O_3)	0.305
benzene (C_6H_6)	0.251
copper (Cu)	0.092
ethanol (C_2H_6O)	0.578
hexane (C_6H_{14})	0.394
hydrogen (H_2)	3.419
ice [$H_2O(s)$]	0.492
iron (Fe)	0.108
iron(III) oxide (Fe_2O_3)	0.156
mercury (Hg)	0.033
oxygen (O_2)	0.219
sodium chloride (NaCl)	0.207
steam [$H_2O(g)$]	0.488
water [$H_2O(\ell)$]	1.00

> *The proportionality constant c is sometimes referred to as the specific heat capacity or (incorrectly) the heat capacity.*

The *direction* of heat flow is not shown in q = *mc*Δ*T*. If energy goes into an object (endothermic), the total energy of the object increases, and the values of heat Δ*T* are positive. If energy is coming out of an object (exothermic), the total energy of the object decreases, and the values of heat and Δ*T* are negative.

Example 11.3.1

What quantity of heat is transferred when a 150.0 g block of iron metal is heated from 25.0°C to 73.3°C? What is the direction of heat flow?

Solution

We can use q = *mc*Δ*T* to determine the amount of heat, but first we need to determine Δ*T*. Because the final temperature of the iron is 73.3°C and the initial temperature is 25.0°C, Δ*T* is as follows:

$$\begin{aligned}\Delta T = T_{final} - T_{initial} \\ &= 73.3^oC - 25.0^oC \\ &= 48.3^oC\end{aligned}$$

The mass is given as 150.0 g, and Table 11.3.1 gives the specific heat of iron as 0.108 cal/g•°C. Substitute the known values into heat = *mc*Δ*T* and solve for amount of heat:

$$\text{q} = (150.0\ \text{g})\left(0.108\ \frac{\text{cal}}{\text{g}\cdot{}^\circ\text{C}}\right)(48.3^\circ\text{C}) = 782\ \text{cal}$$

Note how the gram and °C units cancel algebraically, leaving only the calorie unit, which is a unit of heat. Because the temperature of the iron increases, energy (as heat) must be flowing *into* the metal.

Exercise 11.3.1

What quantity of heat is transferred when a 295.5 g block of aluminum metal is cooled from 128.0°C to 22.5°C? What is the direction of heat flow?

Example 11.3.2

A 10.3 g sample of a reddish-brown metal gave off 71.7 cal of heat as its temperature decreased from 97.5°C to 22.0°C. What is the specific heat of the metal? Can you identify the metal from the data in Table 11.3.1?

Solution

The question gives us the heat, the final and initial temperatures, and the mass of the sample. The value of Δ*T* is as follows:

$$\Delta T = T_{\text{final}} - T_{\text{initial}} = 22.0^\circ\text{C} - 97.5^\circ\text{C} = -75.5^\circ\text{C}$$

If the sample gives off 71.7 cal, it loses energy (as heat), so the value of heat is written as a negative number, −71.7 cal. Substitute the known values into heat = *mc*Δ*T* and solve for *c*:

$$-71.7\ \text{cal} = (10.3\ \text{g})(c)(-75.5^\circ\text{C})$$

$$c = \frac{-71.7\ \text{cal}}{(10.3\ \text{g})(-75.5^\circ\text{C})}$$

$$c = 0.0923\ \text{cal/g}\cdot{}^\circ\text{C}$$

This value for specific heat is very close to that given for copper in Table 11.3.1.

Exercise 11.3.2

A 10.7 g crystal of sodium chloride (NaCl) had an initial temperature of 37.0°C. What is the final temperature of the crystal if 147 cal of heat were supplied to it?

Summary

Heat transfer is related to temperature change.

Concept Review Exercise

1. Describe the relationship between heat transfer and the temperature change of an object.

Answer

1. Heat is equal to the product of the mass, the change in temperature, and a proportionality constant called the specific heat.

Exercises

1. A pot of water is set on a hot burner of a stove. What is the direction of heat flow?
2. Some uncooked macaroni is added to a pot of boiling water. What is the direction of heat flow?
3. How much energy in calories is required to heat 150 g of H_2O from 0°C to 100°C?
4. How much energy in calories is required to heat 125 g of Fe from 25°C to 150°C?
5. If 250 cal of heat were added to 43.8 g of Al at 22.5°C, what is the final temperature of the aluminum?
6. If 195 cal of heat were added to 33.2 g of Hg at 56.2°C, what is the final temperature of the mercury?
7. A sample of copper absorbs 145 cal of energy, and its temperature rises from 37.8°C to 41.7°C. What is the mass of the copper?
8. A large, single crystal of sodium chloride absorbs 98.0 cal of heat. If its temperature rises from 22.0°C to 29.7°C, what is the mass of the NaCl crystal?
9. If 1.00 g of each substance in Table 7.3 were to absorb 100 cal of heat, which substance would experience the largest temperature change?
10. If 1.00 g of each substance in Table 7.3 were to absorb 100 cal of heat, which substance would experience the smallest temperature change?
11. Determine the heat capacity of a substance if 23.6 g of the substance gives off 199 cal of heat when its temperature changes from 37.9°C to 20.9°C.
12. What is the heat capacity of gold if a 250 g sample needs 133 cal of energy to increase its temperature from 23.0°C to 40.1°C?

Answers

1. Heat flows into the pot of water.
2. Heat flows to the macaroni.
3. 15,000 cal
4. 1,690 cal
5. 49.0°C
6. 234°C
7. 404 g
8. 61 g
9. Mercury would experience the largest temperature change.
10. hydrogen (H_2)
11. 0.496 cal/g•°C

12. 0.031 cal/g•°C

11.3: Heat is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

11.4: Phase Changes

Learning Objectives

- Determine the heat associated with a phase change.

Matter can exist in one of several different states, including a gas, liquid, or solid state. The amount of energy in molecules of matter determines the **state of matter**.

- A **gas** is a state of matter in which atoms or molecules have enough energy to move freely. The molecules come into contact with one another only when they randomly collide.
- A **liquid** is a state of matter in which atoms or molecules are constantly in contact but have enough energy to keep changing positions relative to one another.
- A **solid** is a state of matter in which atoms or molecules do not have enough energy to move. They are constantly in contact and in fixed positions relative to one another.




Figure 11.4.1: States of Matter. All three containers contain a substance with the same mass, but the substances are in different states. In the left-hand container, the substance is a gas, which has spread to fill its container. It takes both the shape and volume of the container. In the middle container, the substance is a liquid, which has spread to take the shape of its container but not the volume. In the right-hand container, the substance is a solid, which takes neither the shape nor the volume of its container.

The following are the changes of state:

Changes of State

Solid → Liquid	Melting or fusion
Liquid → Gas	Vaporization
Liquid → Solid	Freezing
Gas → Liquid	Condensation
Solid → Gas	Sublimation

- If heat is added to a substance, such as in melting, vaporization, and sublimation, the process is **endothermic**. In this instance, heat is increasing the speed of the molecules causing them move faster (examples: solid to liquid; liquid to gas; solid to gas).
- If heat is removed from a substance, such as in freezing and condensation, then the process is **exothermic**. In this instance, heat is decreasing the speed of the molecules causing them move slower (examples: liquid to solid; gas to liquid). These changes **release heat** to the surroundings.
- The amount of heat needed to change a sample from solid to liquid would be the same to reverse from liquid to solid. The only difference is the direction of heat transfer.

Example 11.4.1

Label each of the following processes as endothermic or exothermic.

a. water boiling
b. ice forming on a pond

Solution

a. endothermic - you must put a pan of water on the stove and give it heat in order to get water to boil. Because you are adding heat/energy, the reaction is endothermic.
b. exothermic - think of ice forming in your freezer instead. You put water into the freezer, which takes heat out of the water, to get it to freeze. Because heat is being pulled out of the water, it is exothermic. Heat is leaving.

cise

Label each of the following processes as endothermic or exothermic.

a. water vapor condensing
b. gold melting

Answer

a. exothermic

b. endothermic

A phase change is a physical process in which a substance goes from one phase to another. Usually the change occurs when adding or removing heat at a particular temperature, known as the melting point or the boiling point of the substance. The melting point is the temperature at which the substance goes from a solid to a liquid (or from a liquid to a solid). The boiling point is the temperature at which a substance goes from a liquid to a gas (or from a gas to a liquid). The nature of the phase change depends on the direction of the heat transfer. Heat going *into* a substance changes it from a solid to a liquid or a liquid to a gas. Removing heat *from* a substance changes a gas to a liquid or a liquid to a solid.

Two key points are worth emphasizing. First, at a substance's melting point or boiling point, two phases can exist simultaneously. Take water (H_2O) as an example. On the Celsius scale, H_2O has a melting point of 0°C and a boiling point of 100°C. At 0°C, both the solid and liquid phases of H_2O can coexist. However, if heat is added, some of the solid H_2O will melt and turn into liquid H_2O. If heat is removed, the opposite happens: some of the liquid H_2O turns into solid H_2O. A similar process can occur at 100°C: adding heat increases the amount of gaseous H_2O, while removing heat increases the amount of liquid H_2O (Figure 11.4.1).




Figure 11.4.2: Heating curve for water. As heat is added to solid water, the temperature increases until it reaches 0 °C, the melting point. At this point, the phase change, added heat goes into changing the state from a solid to liquid. Only when this phase change is complete, the temperature can increase. (CC BY 3.0 Unported; Community College Consortium for Bioscience Credentials).

Water is a good substance to use as an example because many people are already familiar with it. Other substances have melting points and boiling points as well.

Second, as shown in Figure 11.4.1, **the temperature of a substance does not change** as the substance goes **from one phase to another**. In other words, phase changes are isothermal (isothermal means "constant temperature"). Again, consider H_2O as an example. Solid water (ice) can exist at 0°C. If heat is added to ice at 0°C, some of the solid changes phase to make liquid, which is also at 0°C. Remember, the solid and liquid phases of H_2O can coexist at 0°C. Only after all of the solid has melted into liquid does the addition of heat change the temperature of the substance.

For each phase change of a substance, there is a characteristic quantity of heat needed to perform the phase change per gram (or per mole) of material. The heat of fusion (ΔH_{fus}) is the amount of heat per gram (or per mole) required for a phase change that occurs at the melting point. The heat of vaporization (ΔH_{vap}) is the amount of heat per gram (or per mole) required for a phase change that occurs at the boiling point. If you know the total number of grams or moles of material, you can use the ΔH_{fus} or the ΔH_{vap} to determine the total heat being transferred for melting or solidification using these expressions:

$$\text{heat} = n \times \Delta H_{fus} \tag{11.4.1}$$

where n is the number of moles and ΔH_{fus} is expressed in energy/mole or

$$\text{heat} = m \times \Delta H_{fus} \tag{11.4.2}$$

where m is the mass in grams and ΔH_{fus} is expressed in energy/gram.

For the boiling or condensation, use these expressions:

$$\text{heat} = n \times \Delta H_{vap} \tag{11.4.3}$$

where n is the number of moles) and ΔH_{vap} is expressed in energy/mole or

$$\text{heat} = m \times \Delta H_{vap} \tag{11.4.4}$$

where m is the mass in grams and ΔH_{vap} is expressed in energy/gram.

Remember that a phase change depends on the direction of the heat transfer. If heat transfers in, solids become liquids, and liquids become solids at the melting and boiling points, respectively. If heat transfers out, liquids solidify, and gases condense into liquids. At these points, there are no changes in temperature as reflected in the above equations.

Example 11.4.2

How much heat is necessary to melt 55.8 g of ice (solid H_2O) at 0°C? The heat of fusion of H_2O is 79.9 cal/g.

Solution

We can use the relationship between heat and the heat of fusion (Equation 11.4.1) to determine how many cal of heat are needed to melt this ice:

$$\text{heat} = \text{m} \times \Delta \text{H}_{\text{fus}}$$

$$\text{heat} = (55.8 \ \cancel{\text{g}}) \left(\frac{79.9 \text{ cal}}{\cancel{\text{g}}} \right) = 4,460 \text{ cal}$$

Exercise 11.4.2

How much heat is necessary to vaporize 685 g of H_2O at 100°C? The heat of vaporization of H_2O is 540 cal/g.

Answer

$$\text{heat} = \text{m} \times \Delta \text{H}_{\text{vap}}$$

$$\text{heat} = (685 \ \cancel{\text{g}}) \left(\frac{540 \text{ cal}}{\cancel{\text{g}}} \right) = 370,000 \text{ cal}$$

Table 11.4.1 lists the heats of fusion and vaporization for some common substances. Note the units on these quantities; when you use these values in problem solving, make sure that the other variables in your calculation are expressed in units consistent with the

units in the specific heats or the heats of fusion and vaporization.

Table 11.4.1: Heats of Fusion and Vaporization for Selected Substances

Substance	ΔH_{fus} (cal/g)	ΔH_{vap} (cal/g)
aluminum (Al)	94.0	2,602
gold (Au)	15.3	409
iron (Fe)	63.2	1,504
water (H_2O)	79.9	540
sodium chloride (NaCl)	123.5	691
ethanol (C_2H_5OH)	45.2	200.3
benzene (C_6H_6)	30.4	94.1

Sublimation

There is also a phase change where a solid goes directly to a gas:

$$\text{solid} \rightarrow \text{gas} \tag{11.4.5}$$

This phase change is called *sublimation*. Each substance has a characteristic heat of sublimation associated with this process. For example, the heat of sublimation (ΔH_{sub}) of H_2O is 620 cal/g.

We encounter sublimation in several ways. You may already be familiar with dry ice, which is simply solid carbon dioxide (CO_2). At −78.5°C (−109°F), solid carbon dioxide sublimes, changing directly from the solid phase to the gas phase:

$$CO_2(s) \xrightarrow{-78.5^\circ C} CO_2(g) \tag{11.4.6}$$

Solid carbon dioxide is called dry ice because it does not pass through the liquid phase. Instead, it does directly to the gas phase. (Carbon dioxide *can* exist as liquid but only under high pressure.) Dry ice has many practical uses, including the long-term preservation of medical samples.

Even at temperatures below 0°C, solid H_2O will slowly sublime. For example, a thin layer of snow or frost on the ground may slowly disappear as the solid H_2O sublimes, even though the outside temperature may be below the freezing point of water. Similarly, ice cubes in a freezer may get smaller over time. Although frozen, the solid water slowly sublimes, redepositing on the colder cooling elements of the freezer, which necessitates periodic defrosting (frost-free freezers minimize this redeposition). Lowering the temperature in a freezer will reduce the need to defrost as often.

Under similar circumstances, water will also sublime from frozen foods (e.g., meats or vegetables), giving them an unattractive, mottled appearance called freezer burn. It is not really a “burn,” and the food has not necessarily gone bad, although it looks unappetizing. Freezer burn can be minimized by lowering a freezer’s temperature and by wrapping foods tightly so water does not have any space to sublime into.

Key Takeaway

- There is an energy change associated with any phase change.

11.4: Phase Changes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.3: Phase Changes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.5: Bond Energies and Chemical Reactions

Learning Objectives

The Learning Objectives of this Module are to:

- Define *bond energy*.
- Calculate enthalpy change or heat of reaction.
- Determine if a chemical process is exothermic or endothermic.

What happens when you take a basketball, place it halfway up a playground slide, and then let it go? The basketball rolls down the slide. What happens if you do it again? Does the basketball roll down the slide? It should.

If you were to perform this experiment over and over again, do you think the basketball would ever roll *up* the slide? Probably not. Why not? Well, for starters, in all of our experience, the basketball has always moved to a lower position when given the opportunity. The gravitational attraction of Earth exerts a force on the basketball, and given the chance, the basketball will move down. We say that the basketball is going to a lower *gravitational potential energy*. The basketball can move up the slide, but only if someone exerts some effort (that is, work) on the basketball. A general statement, based on countless observations over centuries of study, is that all objects tend to move spontaneously to a position of minimum energy unless acted on by some other force or object.

Bond Energy

A similar statement can be made about atoms in compounds. Atoms bond together to form compounds because in doing so they attain lower energies than they possess as individual atoms. A quantity of energy, equal to the difference between the energies of the bonded atoms and the energies of the separated atoms, is released, usually as heat. That is, the bonded atoms have a lower energy than the individual atoms do. *When atoms combine to make a compound, energy is always given off, and the compound has a lower overall energy.* In making compounds, atoms act like a basketball on a playground slide; they move in the direction of decreasing energy.

We can reverse the process, just as with the basketball. If we put energy into a molecule, we can cause its bonds to break, separating a molecule into individual atoms. Bonds between certain specific elements usually have a characteristic energy, called the bond energy, that is needed to break the bond. The same amount of energy was liberated when the atoms made the chemical bond in the first place. The term *bond energy* is usually used to describe the strength of interactions between atoms that make covalent bonds. A C–C bond has an approximate bond energy of 80 kcal/mol, while a C=C has a bond energy of about 145 kcal/mol. The C=C bond is stronger than C-C (as discussed in relation to bond length in Section 4.4). For atoms in ionic compounds attracted by opposite charges, the term lattice energy is used. For now, we will deal with covalent bonds in molecules.

Although each molecule has its own characteristic bond energy, some generalizations are possible. For example, although the exact value of a C–H bond energy depends on the particular molecule, all C–H bonds have a bond energy of roughly the same value because they are all C–H bonds. It takes roughly 100 kcal of energy to break 1 mol of C–H bonds, so we speak of the bond energy of a C–H bond as being about 100 kcal/mol. Table 11.5.1 lists the approximate bond energies of various covalent bonds.

Table 11.5.1: Approximate Bond Energies

Bond	Bond Energy (kcal/mol)
C–H	100
C–O	86
C=O	190
C–N	70
C–C	85
C=C	145
C≡C	200

Bond	Bond Energy (kcal/mol)
N–H	93
H–H	105
Br-Br	46
Cl–Cl	58
O–H	110
O=O	119
H–Br	87
H–Cl	103

When a chemical reaction occurs, the atoms in the reactants rearrange their chemical bonds to make products. The new arrangement of bonds does not have the same total energy as the bonds in the reactants. Therefore, when chemical reactions occur, *there will always be an accompanying energy change*. The enthalpy change, for a given reaction can be calculated using the bond energy values from Table 11.5.1.

Enthalpy Change or Heat of Reaction, Δ*H*

Energy changes in chemical reactions are usually measured as changes in enthalpy. In this process, one adds energy to the reaction to break bonds, and extracts energy for the bonds that are formed.


$$\text{enthalpy change} \approx \sum(\text{bonds broken}) - \sum(\text{bonds formed})$$

The ≈ sign is used because we are adding together *average* bond energies (i.e., over many different molecules). Hence, this approach does not give exact values for the enthalpy change, Δ*H*.

Let's consider the reaction of 2 mols of hydrogen gas (H_2) with 1 mol of oxygen gas (O_2) to give 2 mol water:

$$2\,H_2(g) + O_2(g) \rightarrow 2\,H_2O(g)$$

or shown graphically:



with these average bond energies involved:

- H– H = 105 kcal/mol
- O=O = 119 kcal/mol
- O– H = 110 kcal/mol

In this reaction, 2 H–H bonds and 1 O=O bonds are broken, while 4 O–H bonds (two for each H_2O) are formed. The energy changes can be tabulated and calculated as follows:


2 H–H bonds and 1 O=O bonds are broken, while 4 O–H bonds (two for each H2O) are formed. The energy changes can be tabulated and calculated as follows:

Bonds Broken (kcal/mol)		Bonds Formed (kcal/mol)	
2 H–H	2 mol x 105 kcal/mol = 210 kcal	4 O–H	4 mols x 110 kcal/mol = 440 kcal
1 O=O	1 mol x 119 kcal/mol = 119 kcal		

Bonds Broken (kcal/mol)		Bonds Formed (kcal/mol)	
	Total = 329 kcal		**Total** = 440 kcal

$$\Delta H \approx \sum(\text{bonds broken}) - \sum(\text{bonds formed})$$
$$\approx 329 \text{ kcal} - 440 \text{ kcal}$$
$$\approx -111 \text{ kcal}$$

The enthalpy change (ΔH) of the reaction is approximately −111 kcal/mol. This means that bonds in the products (440 kcal) are stronger than the bonds in the reactants (329 kcal) by about 111 kcal/mol. Because the bonds in the products are stronger than those in the reactants, the reaction releases more energy than it consumes. This excess energy is **released** as heat, so the reaction is **exothermic**. Hence, we can re-write the reaction with the heat released (111 kcal) on the **product side** of the equation, as follows:



We can also re-write the reaction equation with the ΔH information (see below). Note that an **exothermic** reaction has a **negative ΔH** value.

$$2\,H_2(g) + O_2(g) \rightarrow 2\,H_2O(g) \qquad \Delta H = -111 \text{ kcal}$$

Example 11.5.1

What is the enthalpy change for this reaction? Is the reaction exothermic or endothermic?

$$H_2(g) + Br_2(g) \rightarrow 2\,HBr(g)$$

Solution

Step 1- First look at the equation and identify which bonds exist on in the reactants (bonds broken).

- one H-H bond and
- one Br-Br bond

Step 2- Do the same for the products (bonds formed)

- two H-Br bonds

Step 3- Identify the bond energies of these bonds from Table 11.5.1:

- H-H bonds: 105 kcal/mol
- Br-Br bonds: 46 kcal/mol

Step 4- Set up the table (see below) and apply the formula for enthalpy change.

Solutions to Example 7.4.1

Bonds Broken (kcal/mol)		Bonds Formed (kcal/mol)	
1 H–H	1 mol x 105 kcal/mol = 105 kcal	2 H–*Br*	2 mols x 87 kcal/mol = 174 kcal
1 Br–Br	1 mol x 46 kcal/mol = 46 kcal		
	Total = 151 kcal		**Total** = 174 kcal

$$\Delta H \approx 151 \text{ kcal} - 174 \text{ kcal}$$
$$\approx -23 \text{ kcal}$$

Step 5- Since ΔH is negative (−23 kcal), the reaction is *exothermic*.

cise

Using the bond energies given in the chart above, find the enthalpy change for the thermal decomposition of water:

$$H_2(g) + Cl_2(g) \rightarrow 2\,HCl(g)$$

Is the reaction written above exothermic or endothermic? Explain.

Answer

ΔH = −43 kcal

Since ΔH is negative (−43 kcal), the reaction is *exothermic*.

Endothermic and Exothermic Reactions

Endothermic and exothermic reactions can be thought of as having energy as either a reactant of the reaction or a product. **Endothermic** reactions require energy, so energy is a reactant. Heat flows from the surroundings to the system (reaction mixture) and the enthalpy of the system increases (ΔH is positive). As discussed in the previous section, in an **exothermic** reaction, heat is released (considered a product) and the enthalpy of the system decreases (ΔH is negative).

In the course of an **endothermic** process, the system gains heat from the surroundings and so the **temperature of the surroundings decreases (gets cold).** A chemical reaction is **exothermic** if heat is released by the system into the surroundings. Because the surroundings is gaining heat from the system, **the temperature of the surroundings increases.** See Figure 11.5.1.





Figure 11.5.1: (A) Endothermic reaction. (B) Exothermic reaction.


Endothermic reaction: surroundings get cooler and delta H is greater than 0, Exothermic reaction: surroundings get warmer and delta H is less than 0

Endothermic Reaction: When 1 mol of calcium carbonate decomposes into 1 mol of calcium oxide and 1 mol of carbon dioxide, 177.8 kJ of heat is absorbed. Because the heat is absorbed by the system, the 177.8 kJ is written as a **reactant**. The **ΔH is positive** for an endothermic reaction.



$$CaCO_3\,(s) \rightarrow CaO\,(s) + CO_2\,(g) \qquad \Delta H = +177.8\ \text{kJ}$$

Exothermic Reaction: When methane gas is combusted, heat is released, making the reaction exothermic. Specifically, the combustion of 1 mol of methane releases 890.4 kilojoules of heat energy. This information can be shown as part of the balanced equation in two ways. First, the amount of heat released can be written in the **product** side of the reaction. Another way is to write the **heat of reaction** (ΔH) information with a **negative** sign, -890.4 kJ.



$$CH_4\,(g) + 2O_2\,(g) \rightarrow CO_2\,(g) + 2H_2O\,(l) \qquad \Delta H = -890.4\ \text{kJ}$$

Example 11.5.2

Is each chemical reaction exothermic or endothermic?

a. $CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(\ell) + 213$ kcal
b. $N_2(g) + O_2(g) + 45$ kcal $\rightarrow 2NO(g)$

Solution

a. Because energy (213 kcal) is a product, energy is given off by the reaction. Therefore, this reaction is exothermic.
b. Because energy (45 kcal) is a reactant, energy is absorbed by the reaction. Therefore, this reaction is endothermic.

Exercise 11.5.2

Is each chemical reaction exothermic or endothermic?

a. $H_2(g) + F_2(g) \rightarrow 2HF\ (g) + 130$ kcal
b. $2C(s) + H_2(g) + 5.3$ kcal $\rightarrow C_2H_2(g)$

Answer

a. The energy (130 kcal) is produced, hence the reaction is exothermic

b. The energy (5.3 kcal) is supplied or absorbed to react, hence, the reaction is endothermic

Energy Diagrams

Endothermic and exothermic reactions can be visually represented by *energy-level diagrams* like the ones in **Figure 11.5.2**. In endothermic reactions, the reactants have higher bond energy (stronger bonds) than the products. **Strong bonds have lower potential energy than weak bonds**. Hence, the **energy of the reactants is lower than that of the products**. This type of reaction is represented by an "uphill" energy-level diagram shown in Figure $11.5.2A$. For an endothermic chemical reaction to proceed, the reactants must absorb energy from their environment to be converted to products.

In an exothermic reaction, the bonds in the product have higher bond energy (stronger bonds) than the reactants. In other words, the **energy of the products is lower than the energy of the reactants,** hence is energetically downhill, shown in Figure $11.5.2B$. Energy is given off as reactants are converted to products. The energy given off is usually in the form of heat (although a few reactions give off energy as light). In the course of an exothermic reaction, heat flows from the system to its surroundings, and thus, gets warm.




Figure 11.5.2A: Endothermic Reactions




Figure 11.5.2B: Exothermic Reactions

Table 11.5.2: Endothermic and Exothermic Reactions

Endothermic Reactions	Exothermic Reactions
Heat is absorbed by reactants to form products.	Heat is released.
Heat is absorbed from the surroundings; as a result, the surroundings get cold.	Heat is released by the reaction to surroundings; surroundings feel hot.
ΔH_{rxn} is positive	ΔH_{rxn} is negative
The bonds broken in the reactants are stronger than the bonds formed in the products	The bonds formed in the products are stronger than the bonds broken in the reactants
The reactants are lower in energy than the products	The products are lower in energy than the reactants
Represented by an "uphill" energy diagram	Represented by an "downhill" energy diagram

Key Takeaways

- Atoms are held together by a certain amount of energy called bond energy.
- Energy is required to break bonds. Energy is released when chemical bonds are formed because atoms become more stable.
- Chemical processes are labeled as exothermic or endothermic based on whether they give off or absorb energy, respectively.

11.5: Bond Energies and Chemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.4: Bond Energies and Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.6: The Energy of Biochemical Reactions

Learning Objectives

- To relate the concept of energy change to chemical reactions that occur in the body.

The chemistry of the human body, or any living organism, is very complex. Even so, the chemical reactions found in the human body follow the same principles of energy that other chemical reactions follow.

Where does the energy that powers our bodies come from? The details are complex, but we can look at some simple processes at the heart of cellular activity.

An important reaction that provides energy for our bodies is the oxidation of glucose ($C_6H_{12}O_6$):

$$C_6H_{12}O_6(s) + 6\,O_2(g) \rightarrow 6\,CO_2(g) + 6\,H_2O(\ell) + 670\text{ kcal} \tag{11.6.1}$$

Considering that 1 mol of $C_6H_{12}O_6(s)$ has a volume of about 115 mL, we can see that glucose is a compact source of energy.

Glucose and other sugars are examples of carbohydrates, which are one of the three main dietary components of a human diet. All carbohydrates supply approximately 4 kcal/g. You can verify that by taking the heat of reaction for glucose oxidation and dividing it by its molar mass. Proteins, the building blocks of structural tissues like muscle and skin, also supply about 4 kcal/g. Other important energy sources for the body are fats, which are largely hydrocarbon chains. Fats provide even more energy per gram, about 9 kcal/g.

Another important reaction is the conversion of adenosine triphosphate (ATP) to adenosine diphosphate (ADP), which is shown in Figure 11.6.1. Under physiological conditions, the breaking of an O–P bond and the formation of an O–P and two O–H bonds gives off about 7.5 kcal/mol of ATP. This may not seem like much energy, especially compared to the amount of energy given off when glucose reacts. It is enough energy, however, to fuel other biochemically important chemical reactions in our cells.




Figure 11.6.1: ATP to ADP. The conversion of ATP (top) to ADP (bottom) provides energy for the cells of the body.

Even complex biological reactions must obey the basic rules of chemistry.

Career Focus: Dietitian

A dietitian is a nutrition expert who communicates food-related information to the general public. In doing so, dietitians promote the general well-being among the population and help individuals recover from nutritionally related illnesses.

Our diet does not just supply us with energy. We also get vitamins, minerals, and even water from what we eat. Eating too much, too little, or not enough of the right foods can lead to a variety of problems. Dietitians are trained to make specific dietary recommendations to address particular issues relating to health. For example, a dietitian might work with a person to develop an overall diet that would help that person lose weight or control diabetes. Hospitals employ dietitians in planning menus for patients, and many dietitians work with community organizations to improve the eating habits of large groups of people.

Key Takeaway

- Energy to power the human body comes from chemical reactions.

11.6: The Energy of Biochemical Reactions is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.5: The Energy of Biochemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.E: Energy and Chemical Processes (Exercises)

7.1: Energy and Its Units

Concept Review Exercises

1. What is the relationship between energy and heat?
2. What units are used to express energy and heat?

Answers

1. Heat is the exchange of energy from one part of the universe to another. Heat and energy have the same units.
2. Joules and calories are the units of energy and heat.

Exercises

1. Define *energy*.
2. What is heat?
3. What is the relationship between a calorie and a joule? Which unit is larger?
4. What is the relationship between a calorie and a kilocalorie? Which unit is larger?
5. Express 1,265 cal in kilocalories and in joules.
6. Express 9,043.3 J in calories and in kilocalories.
7. One kilocalorie equals how many kilojoules?
8. One kilojoule equals how many kilocalories?
9. Many nutrition experts say that an average person needs 2,000 Cal per day from his or her diet. How many joules is this?
10. Baby formula typically has 20.0 Cal per ounce. How many ounces of formula should a baby drink per day if the RDI is 850 Cal?

Answers

1. Energy is the ability to do work.
2. Heat is a form of enery (thermal) that can be transferred from one object to another.
3. 1 cal = 4.184 J; the calorie is larger.
4. 1 kilocalorie(1 Cal) = 1000 cal; the kcal is larger.
5. 1.265 kcal; 5,293 J
6. 2161.4 cal; 2.1614 kcal
7. 1 kcal = 4.184 kJ
8. 1 kJ = 0.239 kcal
9. 8.4×10^6 J
10. 42.5 oz

7.2: Heat and Temperature

Concept Review Exercise

1. Describe the relationship between heat transfer and the temperature change of an object.
2. Describe what happens when two objects that have different temperatures come into contact with one another.

Answer

1. Heat is equal to the product of the mass, the change in temperature, and a proportionality constant called the specific heat.

2. The temperature of the hot object decreases and the temperature of the cold object increases as heat is transferred from the hot object to the cold object. The change in temperature of each depends on the identity and properties of each substance.

Exercises

1. The melting point of mercury is −38.84°C. Convert this value to degrees Fahrenheit and Kelvin.

2. A pot of water is set on a hot burner of a stove. What is the direction of heat flow?

3. Some uncooked macaroni is added to a pot of boiling water. What is the direction of heat flow?

4. How much energy in calories is required to heat 150 g of H_2O from 0°C to 100°C?

5. How much energy in calories is required to heat 125 g of Fe from 25°C to 150°C?

6. If 250 cal of heat were added to 43.8 g of Al at 22.5°C, what is the final temperature of the aluminum?

7. If 195 cal of heat were added to 33.2 g of Hg at 56.2°C, what is the final temperature of the mercury?

8. A sample of copper absorbs 145 cal of energy, and its temperature rises from 37.8°C to 41.7°C. What is the mass of the copper?

9. A large, single crystal of sodium chloride absorbs 98.0 cal of heat. If its temperature rises from 22.0°C to 29.7°C, what is the mass of the NaCl crystal?

10. If 1.00 g of each substance in Table 7.2.1 were to absorb 100 cal of heat, which substance would experience the largest temperature change?

11. If 1.00 g of each substance in Table 7.2.1 were to absorb 100 cal of heat, which substance would experience the smallest temperature change?

12. Determine the heat capacity of a substance if 23.6 g of the substance gives off 199 cal of heat when its temperature changes from 37.9°C to 20.9°C.

13. What is the heat capacity of gold if a 250 g sample needs 133 cal of energy to increase its temperature from 23.0°C to 40.1°C?

Answers

1. -37.91^{0}F and 234.31 K

2. Heat flows into the pot of water.

3. Heat flows to the macaroni.

4. 15,000 cal

5. 1,690 cal

6. 49.0°C

7. 234°C

8. 404 g

9. 61 g

10. Mercury would experience the largest temperature change.

11. hydrogen (H_2)

12. 0.496 cal/g•°C

13. 0.031 cal/g•°C

7.3: Phase Changes

Concept Review Exercises

1. Explain what happens when heat flows into or out of a substance at its melting point or boiling point.
2. How does the amount of heat required for a phase change relate to the mass of the substance?
3. What is the direction of heat transfer in boiling water?
4. What is the direction of heat transfer in freezing water?
5. What is the direction of heat transfer in sweating?

Answers

1. The energy goes into changing the phase, not the temperature.

2. The amount of heat is a constant per gram of substance.

3. Boiling. Heat is being added to the water to get it from the liquid state to the gas state.

4. Freezing. Heat is exiting the system in order to go from liquid to solid. Another way to look at it is to consider the opposite process of melting. Energy is consumed (endothermic) to melt ice (solid to liquid) so the opposite process (liquid to solid) must be exothermic.

5. Sweating. Heat is consumed to evaporate the moisture on your skin which lowers your temperature.

Exercises

1. How much energy is needed to melt 43.8 g of Au at its melting point of 1,064°C?
2. How much energy is given off when 563.8 g of NaCl solidifies at its freezing point of 801°C?
3. What mass of ice can be melted by 558 cal of energy?
4. How much ethanol (C_2H_5OH) in grams can freeze at its freezing point if 1,225 cal of heat are removed?
5. What is the heat of vaporization of a substance if 10,776 cal are required to vaporize 5.05 g? Express your final answer in joules per gram.
6. If 1,650 cal of heat are required to vaporize a sample that has a heat of vaporization of 137 cal/g, what is the mass of the sample?
7. What is the heat of fusion of water in calories per mole?
8. What is the heat of vaporization of benzene (C_6H_6) in calories per mole?
9. What is the heat of vaporization of gold in calories per mole?
10. What is the heat of fusion of iron in calories per mole?

Answers

1. 670 cal
2. 69,630 cal
3. 6.98 g
4. 27.10 g
5. 8,930 J/g
6. 12.0 g
7. 1,440 cal/mol
8. 7,350 cal/mol
9. 80,600 cal/mol
10. 3,530 cal/mol

7.4: Bond Energies and Chemical Reactions

Concept Review Exercises

1. What is the connection between energy and chemical bonds?
2. Why does energy change during the course of a chemical reaction?
3. Two different reactions are performed in two identical test tubes. In reaction A, the test tube becomes very warm as the reaction occurs. In reaction B, the test tube becomes cold. Which reaction is endothermic and which is exothermic? Explain.
4. Classify "burning paper" as endothermic or exothermic processes.

Answers

1. Chemical bonds have a certain energy that is dependent on the elements in the bond and the number of bonds between the atoms.
2. Energy changes because bonds rearrange to make new bonds with different energies.
3. Reaction A is exothermic because heat is leaving the system making the test tube feel hot. Reaction B is endothermic because heat is being absorbed by the system making the test tube feel cold.
4. "Burning paper" is exothermic because burning (also known as combustion) releases heat

Exercises

1. Using the data in Table 7.4.1, calculate the energy of one C–H bond (as opposed to 1 mol of C–H bonds). Recall that 1 mol = 6.022×10^{23} C–H bonds
2. Using the data in Table 7.4.1, calculate the energy of one C=C bond (as opposed to 1 mol of C=C bonds). Recall that 1 mol = 6.022×10^{23} C=C bonds
3. Is a bond-breaking process exothermic or endothermic?
4. Is a bond-making process exothermic or endothermic?
5. Is each chemical reaction exothermic or endothermic?
   a. $2SnCl_2(s) + 33\ kcal \rightarrow Sn(s) + SnCl_4(s)$
   b. $CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(\ell) + 213\ kcal$
6. Is each chemical reaction exothermic or endothermic?
   a. $C_2H_4(g) + H_2(g) \rightarrow C_2H_6(g) + 137\ kJ$
   b. $C(s, graphite) + 1.9\ kJ \rightarrow C(s, diamond)$

Answers

1. 1.661×10^{-19} cal
2. 2.408×10^{-19} cal
3. endothermic
4. exothermic
5. a. endothermic
   b. exothermic
6. a. exothermic
   b. endothermic

7.5: The Energy of Biochemical Reactions

Concept Review Exercise

1. What is the energy content per gram of proteins, carbohydrates, and fats?

Answer

1. proteins and carbohydrates: 4 kcal/g; fats: 9 kcal/g

Exercises

1. An 8 oz serving of whole milk has 8.0 g of fat, 8.0 g of protein, and 13 g of carbohydrates. Approximately how many kilocalories does it contain?
2. A serving of potato chips has 160 kcal. If the chips have 15 g of carbohydrates and 2.0 g of protein, about how many grams of fat are in a serving of potato chips?
3. The average body temperature of a person is 37°C, while the average surrounding temperature is 22°C. Is overall human metabolism exothermic or endothermic?
4. Cold-blooded animals absorb heat from the environment for part of the energy they need to survive. Is this an exothermic or an endothermic process?
5. If the reaction ATP → ADP gives off 7.5 kcal/mol, then the reverse process, ADP → ATP requires 7.5 kcal/mol to proceed. How many moles of ADP can be converted to ATP using the energy from 1 serving of potato chips (see Exercise 2)?
6. If the oxidation of glucose yields 670 kcal of energy per mole of glucose oxidized, how many servings of potato chips (see Exercise 2) are needed to provide the same amount of energy?

Answers

1. 156 kcal
2. 10.2 g
3. exothermic
4. endothermic
5. 21.3 mol
6. 4.2 servings

Additional Exercises

1. Sulfur dioxide (SO_2) is a pollutant gas that is one cause of acid rain. It is oxidized in the atmosphere to sulfur trioxide (SO_3), which then combines with water to make sulfuric acid (H_2SO_4).
   a. Write the balanced reaction for the oxidation of SO_2 to make SO_3. (The other reactant is diatomic oxygen.)
   b. When 1 mol of SO_2 reacts to make SO_3, 23.6 kcal of energy are given off. If 100 lb (1 lb = 454 g) of SO_2 were converted to SO_3, what would be the total energy change?
2. Ammonia (NH_3) is made by the direct combination of H_2 and N_2 gases according to this reaction:

$$N_2(g) + 3H_2(g) \rightarrow 2NH_3(g) + 22.0 \text{ kcal}$$

   a. Is this reaction endothermic or exothermic?
   b. What is the overall energy change if 1,500 g of N_2 are reacted to make ammonia?
3. A 5.69 g sample of iron metal was heated in boiling water to 99.8°C. Then it was dropped into a beaker containing 100.0 g of H_2O at 22.6°C. Assuming that the water gained all the heat lost by the iron, what is the final temperature of the H_2O and Fe?
4. A 5.69 g sample of copper metal was heated in boiling water to 99.8°C. Then it was dropped into a beaker containing 100.0 g of H_2O at 22.6°C. Assuming that the water gained all the heat lost by the copper, what is the final temperature of the H_2O and Cu?
5. When 1 g of steam condenses, 540 cal of energy is released. How many grams of ice can be melted with 540 cal?

6. When 1 g of water freezes, 79.9 cal of energy is released. How many grams of water can be boiled with 79.9 cal?
7. The change in energy is +65.3 kJ for each mole of calcium hydroxide [$Ca(OH)_2$] according to the following reaction:

$$Ca(OH)_2(s) \rightarrow CaO(s) + H_2O(g)$$

How many grams of $Ca(OH)_2$ could be reacted if 575 kJ of energy were available?

8. The thermite reaction gives off so much energy that the elemental iron formed as a product is typically produced in the liquid state:

$$2Al(s) + Fe_2O_3(s) \rightarrow Al_2O_3(s) + 2Fe(\ell) + 204 \text{ kcal}$$

How much heat will be given off if 250 g of Fe are to be produced?

9. A normal adult male requires 2,500 kcal per day to maintain his metabolism.
    a. Nutritionists recommend that no more than 30% of the calories in a person's diet come from fat. At 9 kcal/g, what is the maximum mass of fat an adult male should consume daily?
    b. At 4 kcal/g each, how many grams of protein and carbohydrates should an adult male consume daily?
10. A normal adult male requires 2,500 kcal per day to maintain his metabolism.
    a. At 9 kcal/g, what mass of fat would provide that many kilocalories if the diet was composed of nothing but fats?
    b. At 4 kcal/g each, what mass of protein and/or carbohydrates is needed to provide that many kilocalories?
11. The volume of the world's oceans is approximately 1.34×10^{24} cm^3.
    a. How much energy would be needed to increase the temperature of the world's oceans by 1°C? Assume that the heat capacity of the oceans is the same as pure water.
    b. If Earth receives 6.0×10^{22} J of energy per day from the sun, how many days would it take to warm the oceans by 1°C, assuming all the energy went into warming the water?
12. Does a substance that has a small specific heat require a small or large amount of energy to change temperature? Explain.
13. Some biology textbooks represent the conversion of adenosine triphosphate (ATP) to adenosine diphosphate (ADP) and phosphate ions as follows:

$$ATP \rightarrow ADP + \text{phosphate} + \text{energy}$$

What is wrong with this reaction?

14. Assuming that energy changes are additive, how much energy is required to change 15.0 g of ice at −15°C to 15.0 g of steam at 115°C? (Hint: you will have five processes to consider.)

Answers

1.
    a. $2SO_2 + O_2 \rightarrow 2SO_3$
    b. 16,700 kcal
2.
    a. exothermic
    b. 1177 kcal
3. about 23.1°C
4. about 23.0°C
5. 6.76 g
6. 0.148 g

7. 652 g

8. 457 kcal

9.

   a. 83.3 g
   b. 438 g

10.

   a. 278 g
   b. 625 g

11.

   a. 1.34×10^{24} cal
   b. 93 days

12. A substance with **smaller specific heat** requires **less energy** per unit of mass to raise its temperature,

13. A reactant is missing: H_2O is missing.

14. Total energy = 11,019 cal

11.E: Energy and Chemical Processes (Exercises) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.1: Energy and Its Units** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **7.2: Heat and Temperature** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **7.3: Phase Changes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **7.4: Bond Energies and Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **7.5: The Energy of Biochemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.S: Energy and Chemical Processes (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Energy is the ability to do work. The transfer of energy from one place to another is **heat**. Heat and energy are measured in units of **joules**, **calories**, or kilocalories (equal to 1,000 calories). The amount of heat gained or lost when the temperature of an object changes can be related to its mass and a constant called the **specific heat** of the substance.

The transfer of energy can also cause a substance to change from one phase to another. During the transition, called a **phase change**, heat is either added or lost. Despite the fact that heat is going into or coming out of a substance during a phase change, the temperature of the substance does not change until the phase change is complete; that is, phase changes are **isothermal**. Analogous to specific heat, a constant called the **heat of fusion** of a substance describes how much heat must be transferred for a substance to melt or solidify (that is, to change between solid and liquid phases), while the **heat of vaporization** describes the amount of heat transferred in a boiling or condensation process (that is, to change between liquid and gas phases).

Every chemical change is accompanied by an energy change. This is because the interaction between atoms bonding to each other has a certain **bond energy**, the energy required to break the bond (called **lattice energy** for ionic compounds), and the bond energies of the reactants will not be the same as the bond energies of the products. Reactions that give off energy are called **exothermic**, while reactions that absorb energy are called **endothermic**. Energy-level diagrams can be used to illustrate the energy changes that accompany chemical reactions.

Even complex biochemical reactions have to follow the rules of simple chemistry, including rules involving energy change. Reactions of **carbohydrates** and **proteins** provide our bodies with about 4 kcal of energy per gram, while **fats** provide about 9 kcal per gram.

11.S: Energy and Chemical Processes (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **7.S: Energy and Chemical Processes (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

12: Solids, Liquids, and Gases

12.1: A Molecular Comparison of Gases, Liquids, and Solids
12.2: Kinetic-Molecular Theory
12.3: The Ideal Gas Equation
12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases
12.5: Dalton's Law (Law of Partial Pressures)
12.6: Evaporation and Condensation
12.7: Intermolecular forces

12: Solids, Liquids, and Gases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

12.1: A Molecular Comparison of Gases, Liquids, and Solids

Skills to Develop

- To be familiar with the kinetic molecular description of liquids.

The ***physical*** properties of a substance depends upon its physical state. Water vapor, liquid water and ice all have the same ***chemical*** properties, but their ***physical*** properties are considerably different. In general ***Covalent bonds*** determine: molecular shape, bond energies, ***chemical*** properties, while ***intermolecular forces*** (non-covalent bonds) influence the ***physical*** properties of liquids and solids. The kinetic molecular theory of gases gives a reasonably accurate description of the behavior of gases. A similar model can be applied to liquids, but it must take into account the nonzero volumes of particles and the presence of strong intermolecular attractive forces.



Figure 12.1.1: *The four fundamental states of matter. Clockwise from top left, they are solid, liquid, plasma, and gas, represented by an ice sculpture, a drop of water, electrical arcing from a tesla coil, and the air around clouds, respectively. Images used with permission from Wikipedia.*

The ***state*** of a substance depends on the balance between the ***kinetic energy*** of the individual particles (molecules or atoms) and the ***intermolecular forces***. The k**inetic energy** keeps the molecules apart and moving around, and is a function of the temperature of the substance and the intermolecular forces try to draw the particles together (Figure 12.1.2). A discussed previously, gasses are very sensitive to temperatures and pressure. However, these also affect liquids and solids too. Heating and cooling can change the ***kinetic energy*** of the particles in a substance, and so, we can change the physical state of a substance by heating or cooling it. Increasing the pressure on a substance forces the molecules closer together, which ***increases*** the strength of intermolecular forces




Figure 12.1.2: *Molecular level picture of gases, liquids and solids. Image used with permission from WIkipedia (Yelod).*

Below is an overview of the general properties of the three different phases of matter.

Properties of Gases

- A collection of widely separated molecules
- The kinetic energy of the molecules is greater than any attractive forces between the molecules
- The lack of any significant attractive force between molecules allows a gas to expand to fill its container
- If attractive forces become large enough, then the gases exhibit ***non-ideal*** behavior

Properties of Liquids

- The intermolecular attractive forces are strong enough to hold molecules close together
- Liquids are more dense and less compressible than gasses
- Liquids have a definite volume, independent of the size and shape of their container
- The attractive forces are ***not*** strong enough, however, to keep neighboring molecules in a fixed position and molecules are free to move past or slide over one another

> *Thus, liquids can be poured and assume the shape of their containers*

Properties of Solids

- The intermolecular forces between neighboring molecules are strong enough to keep them locked in position
- Solids (like liquids) are not very compressible due to the lack of space between molecules
- If the molecules in a solid adopt a highly ordered packing arrangement, the structures are said to be ***crystalline***

> *Due to the strong intermolecular forces between neighboring molecules, solids are rigid*

Note

- Cooling a gas may change the state to a liquid
- Cooling a liquid may change the state to a solid
- Increasing the pressure on a gas may change the state to a liquid
- Increasing the pressure on a liquid may change the state to a solid



***Video* 12.1.1** *: Video highlighting the properties for the three states of matter. Source found at https://www.youtube.com/watch?v=s-KvoVzukHo.*

Physical Properties of Liquids

In a gas, the distance between molecules, whether monatomic or polyatomic, is very large compared with the size of the molecules; thus gases have a low density and are highly compressible. In contrast, the molecules in liquids are very close together, with essentially no empty space between them. As in gases, however, the molecules in liquids are in constant motion, and their kinetic energy (and hence their speed) depends on their temperature. We begin our discussion by examining some of the characteristic properties of liquids to see how each is consistent with a modified kinetic molecular description.

The properties of liquids can be explained using a modified version of the kinetic molecular theory of gases described previously This model explains the higher density, greater order, and lower compressibility of liquids versus gases; the thermal expansion of liquids; why they diffuse; and why they adopt the shape (but not the volume) of their containers. A kinetic molecular description of liquids must take into account both the nonzero volumes of particles and the presence of strong intermolecular attractive forces. Solids and liquids have particles that are fairly close to one another, and are thus called "**condensed phases**" to distinguish them from gases

- **Density**: The molecules of a liquid are packed relatively close together. Consequently, liquids are much denser than gases. The density of a liquid is typically about the same as the density of the solid state of the substance. Densities of liquids are therefore more commonly measured in units of grams per cubic centimeter (g/cm^3) or grams per milliliter (g/mL) than in grams per liter (g/L), the unit commonly used for gases.

- **Molecular Order:** Liquids exhibit short-range order because strong intermolecular attractive forces cause the molecules to pack together rather tightly. Because of their higher kinetic energy compared to the molecules in a solid, however, the molecules in a liquid move rapidly with respect to one another. Thus unlike the ions in the ionic solids, the molecules in liquids are not arranged in a repeating three-dimensional array. Unlike the molecules in gases, however, the arrangement of the molecules in a liquid is not completely random.
- **Compressibility**: Liquids have so little empty space between their component molecules that they cannot be readily compressed. Compression would force the atoms on adjacent molecules to occupy the same region of space.
- **Thermal Expansion**: The intermolecular forces in liquids are strong enough to keep them from expanding significantly when heated (typically only a few percent over a 100°C temperature range). Thus the volumes of liquids are somewhat fixed. Notice from Table S1 (with a shorten version in Table 12.1.1) that the density of water, for example, changes by only about 3% over a 90-degree temperature range.

***Table* 12.1.1**: *The Density of Water at Various Temperatures*

T (°C)	Density (g/cm^3)
0	0.99984
30	0.99565
60	0.98320
90	0.96535

- **Diffusion**: Molecules in liquids diffuse because they are in constant motion. A molecule in a liquid cannot move far before colliding with another molecule, however, so the mean free path in liquids is very short, and the rate of diffusion is much slower than in gases.
- **Fluidity**: Liquids can flow, adjusting to the shape of their containers, because their molecules are free to move. This freedom of motion and their close spacing allow the molecules in a liquid to move rapidly into the openings left by other molecules, in turn generating more openings, and so forth (Figure 11.1.3).




***Figure* 112.1.3**: *Why Liquids Flow. Molecules in a liquid are in constant motion. Consequently, when the flask is tilted, molecules move to the left and down due to the force of gravity, and the openings are occupied by other molecules. The result is a net flow of liquid out of the container.*

Contributors

- Mike Blaber (Florida State University)

12.1: A Molecular Comparison of Gases, Liquids, and Solids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

12.2: Kinetic-Molecular Theory

Learning Outcomes

- Define pressure and describe how gases exert pressure.
- Convert between units of gas pressure (atm, mmHg, torr, and Pa).
- Relate temperature to average kinetic energy.
- Relate mass to molecular speed.
- Describe the behavior of an ideal gas.

The **kinetic-molecular theory** *is a theory that explains the states of matter and is based on the idea that matter is composed of tiny particles that are always in motion*. The theory helps explain observable properties and behaviors of solids, liquids, and gases. However, the theory is most easily understood as it applies to gases. The theory applies specifically to a model of a gas called an ideal gas. An **ideal gas** *is an imaginary gas whose behavior perfectly fits all the assumptions of the kinetic-molecular theory*. In reality, gases are not ideal, but they are very close to being so under most everyday conditions.

The kinetic-molecular theory, as it applies to gases, has five basic assumptions.

1. **Gases consist of very large numbers of tiny spherical particles that are far apart from one another compared to their size**. The particles of a gas may be either atoms or molecules. The distance between the particles of a gas is much, much greater than the distances between the particles of a liquid or a solid. Most of the volume of a gas, therefore, is composed of the empty space between the particles. In fact, the volume of the particles themselves is considered to be insignificant compared to the volume of the empty space.
2. **Gas particles are in constant rapid motion in random directions**. The fast motion of gas particles gives them a relatively large amount of kinetic energy. Recall that kinetic energy is the energy that an object possesses because of its motion. The particles of a gas move in a straight line until they collide with another particle or with one of the walls of their container (see figure below).
3. **Collisions between gas particles and between particles and the container walls are elastic collisions**. An elastic collision is one in which there is no overall loss of kinetic energy. Kinetic energy may be transferred from one particle to another during an elastic collision, but there is no change in the total energy of the colliding particles.
4. **There are no forces of attraction or repulsion between gas particles**. Attractive forces are responsible for particles of a real gas condensing together to form a liquid. It is assumed that the particles of an ideal gas have no such attractive forces. The motion of each particle is completely independent of the motion of all other particles.
5. **The average kinetic energy of gas particles is dependent upon the temperature of the gas**. As the temperature of a gas is increased, its component particles begin to move faster, resulting in an increase in their kinetic energies. Not all particles in a given sample have the same speed, so the sample will contain particles with a range of different kinetic energies. However, the average kinetic energy of the particles in a sample is proportional to its temperature.




Figure 12.2.1: Gas particles move in a random, linear fashion according to the kinetic-molecular theory. The space between particles is very large compared to the size of the particles.

Gas Pressure

Pressure *is defined as the force per unit area on a surface.*

$$\text{Pressure} = \frac{\text{force}}{\text{area}} \tag{12.2.1}$$

When a person stands on the floor, his feet exert pressure on the surface. That pressure is related to both the mass of the person and the surface area of his feet. If the person were holding a heavy object, the pressure would increase because of a greater force. Alternatively, if the person stands on his toes, the pressure also increases because of a decrease in the surface area.

Gas molecules also exert pressure. Earth's atmosphere exerts pressure because gravity acts on the huge number of gas particles contained in the atmosphere, holding it in place. Pressure is also exerted by small samples of gas, such as the outward pressure exerted by the gas inside a balloon. **Gas pressure** *is the pressure that results from collisions of gas particles with an object*. Inside the balloon, the gas particles collide with the balloon's inner walls. It is those collisions that keep the balloon inflated. If the gas particles were to suddenly stop moving, the balloon would instantly deflate. The figure below is an illustration of gas particles exerting pressure inside a container.




Figure 12.2.2: Gas pressure results from collisions with the surface of its container.

Measuring Pressure

Atmospheric pressure *is the pressure exerted by the gas particles in Earth's atmosphere as those particles collide with objects*. A **barometer** *is an instrument used to measure atmospheric pressure*. A traditional mercury barometer consists of an evacuated tube immersed in a container of mercury. Air molecules from the atmosphere push down on the outer surface of the mercury, but, because the inside of the tube is a vacuum, there is no corresponding downward push on the mercury in the tube. As a result, the mercury rises inside the tube. The height to which the mercury rises is dependent on the external air pressure.

At sea level, a mercury column will rise a distance of 760 mm. This atmospheric pressure is reported as 760 mm Hg (millimeters of mercury). At higher altitudes, the atmospheric pressure is lower, so the column of mercury will not rise as high. For example, on the summit of Mt. Everest (at an elevation of 8848 m), the air pressure is 253 mm Hg. Atmospheric pressure is also slightly dependent on weather conditions.

A more convenient barometer, called an aneroid barometer, measures pressure by the expansion and contraction of a small spring within an evacuated metal capsule.

Units of Gas Pressure

A barometers measures gas pressure by the height of the column of mercury. One unit of gas pressure is the millimeter of mercury (mm Hg). An equivalent unit to the mm Hg is called the torr, in honor of the inventor of the barometer, Evangelista Torricelli. The **pascal** (Pa) *is the standard unit of pressure*. A pascal is a very small amount of pressure, so a more useful unit for everyday gas

pressures is the kilopascal (kPa). A kilopascal is equal to 1000 pascals. Another commonly used unit of pressure is the atmosphere (atm). Standard atmospheric pressure is called $1\ \mathrm{atm}$ of pressure and is equal to $760\ \mathrm{mm\ Hg}$ and $101.3\ \mathrm{kPa}$ The relationships between the most common units of pressure are shown below.

$$1\ \mathrm{atm} = 760\ \mathrm{mm\ Hg} = 760\ \mathrm{torr} = 101.3\ \mathrm{kPa} \tag{12.2.2}$$

Example 12.2.1

The atmospheric pressure in a mountainous location is measured to be $613\ \mathrm{mm\ Hg}$. What is this pressure in atm and in kPa?

Solution:

Step 1: List the known quantities and plan the problem.

Known

- Given: $613\ \mathrm{mm\ Hg}$
- $1\ \mathrm{atm} = 760\ \mathrm{mm\ Hg}$
- $101.3\ \mathrm{kPa} = 760\ \mathrm{mm\ Hg}$

Unknown

- Pressure $=?\ \mathrm{atm}$
- Pressure $=?\ \mathrm{kPa}$

Use conversion factors from the equivalent pressure units to convert from $\mathrm{mm\ Hg}$ to atm and from $\mathrm{mm\ Hg}$ to kPa.

Step 2: Solve.

$$613\ \mathrm{mm\ Hg} \times \frac{1\ \mathrm{atm}}{760\ \mathrm{mm\ Hg}} = 0.807\ \mathrm{atm}$$

$$613\ \mathrm{mm\ Hg} \times \frac{101.3\ \mathrm{kPa}}{760\ \mathrm{mm\ Hg}} = 81.7\ \mathrm{kPa}$$

Step 3: Think about your result.

The air pressure is about 80% of the standard atmospheric pressure at sea level. The standard pressure of $760\ \mathrm{mm\ Hg}$ can be considered to have three significant figures.

Kinetic Energy and Temperature

As stated in the kinetic-molecular theory, the temperature of a substance is related to the average kinetic energy of the particles of that substance. When a substance is heated, some of the absorbed energy is stored within the particles, while some of the energy increases the speeds at which the particles are moving. This is observed as an increase in the temperature of the substance.

Average Kinetic Energy

At any given temperature, not all of the particles in a sample of matter have the same kinetic energy. Instead, the particles display a wide range of kinetic energies. Most of the particles have a kinetic energy near the middle of the range. However, some of the particles have kinetic energies a great deal lower or a great deal higher than the average (see figure below).




Figure 12.2.3: *A distribution of molecular kinetic energies as a function of temperature. The blue curve is for a low temperature, while the red curve is for a high temperature.*

The blue curve shown (see figure above) is for a sample of matter at a relatively low temperature, while the red curve is for a sample at a relatively high temperature. In both cases, most of the particles have intermediate kinetic energies, close to the average. Notice that as temperature increases, the range of kinetic energies increases and the distribution curve "flattens out".

At a given temperature, the particles of any substance have the same average kinetic energy. At room temperature, the molecules in a sample of liquid water have the same average kinetic energy as the molecules in a sample of oxygen gas or the ions in a sample of sodium chloride.

Absolute Zero

As a sample of matter is continually cooled, the average kinetic energy of its particles decreases. Eventually, one would expect the particles to stop moving completely. **Absolute zero** *is the temperature at which the motion of particles theoretically ceases.* Absolute zero has never been attained in the laboratory, but temperatures on the order of 1×10^{-10} K have been achieved. The Kelvin temperature scale is based on this theoretical limit, so absolute zero is equal to 0 K. The Kelvin temperature of a substance is directly proportional to the average kinetic energy of the particles of the substance. For example, the particles in a sample of hydrogen gas at 200 K have twice the average kinetic energy as the particles in a hydrogen sample at 100 K.

Contributors

-
- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)

This page titled 12.2: Kinetic-Molecular Theory is shared under a not declared license and was authored, remixed, and/or curated by Allison Soult.

12.3: The Ideal Gas Equation

Learning Outcomes

- State the ideal gas law and identify the variables.
- Use the ideal gas law to solve for an unknown.
- State the combined gas law.
- Simplify the combined gas law for any values that are constant.
- Use the combined gas law to solve for an unknown value.

Individual Gas Laws

Properties of gases such as pressure (*P*), volume (*V*), temperature (*T*), and moles(*n*) are relatively easy to measure. Unlike with liquids and solids, the particles (molecules or atoms) in a gas phase sample are very far apart from one another. As a result, their behavior is much more predictable because intermolecular forces become insignificant for most samples in the gas phase even over a wide range of conditions. The presence of intermolecular forces in liquid and solid samples makes their behavior harder to predict.

Experiments with gas phase samples over time showed the relationship between pairs of variables (*P*, *V*, *T*, and *n*) and individual gas laws (equations) show the quantitative relationship between those variables. Avogadro's law tells us that at constant *P* and *T*, the volume of a gas is directly proportional to the amount of gas. Boyle's law says that volume is inversely proportional to pressure at constant *T* and *n*. Charles' indicates that volume is directly proportional to temperature at constant *P* and *n*.

The video below shows a situation where 3 variables, pressure, volume, and amount of substance (moles) are all interrelated: inside our lungs.



As you can see in the video, when the pink balloon on the bottom (the "diaphragm") is pulled down, the balloon inside expands. This expansion causes a decrease in pressure (Boyle's Law). The pressure decrease causes a pressure differential, drawing air in through the straw, an increase in the amount of air (moles). So in your lungs, volume, pressure, and amount of air are all related. But none of the current laws explain the relation between 2 variables. How can this be resolved?

The Ideal Gas Law

These three laws may all be applied at once if we write ($\propto$ means "proportional to"):

$$V \propto n \times \frac{1}{P} \times T \tag{12.3.1}$$

or, introducing a constant of proportionality *R*,

$$V = R\,\frac{nT}{P} \tag{12.3.2}$$

This is known as the ideal gas law which results from the combination of the individual gas laws. Equation 12.3.2 applies to all gases at low pressures and high temperatures and is a very good approximation under nearly all conditions. The value of *R*, the **gas**

constant, is independent of the kind of gas, the temperature, or the pressure and has a value of $\frac{0.08206\ L{\cdot}atm}{mol{\cdot}K}$.

Equation 12.3.2 is usually rearranged by multiplying both sides by *P*, so that it reads

$$PV = nRT \tag{12.3.3}$$

This is called the ideal gas equation or the ideal gas law. With the ideal gas equation we can convert from volume of a gas to amount of substance (provided that *P* and *T* are known). This is very useful since the volume, pressure, and temperature of a gas are easier to measure than mass, and because knowledge of the molar mass is unnecessary.

Note that for any gas law calculations, the temperature must be in units of Kelvin. The relationship between ^{o}C and K is K = °C + 273.15.

Example 12.3.1 : MOLES of Gas

Calculate the moles of gas in a 0.100 L sample at a temperature of 300 K and a pressure of 0.987 atm.

Solution

$$PV = nRT \tag{12.3.4}$$

$$(0.987\ atm)\,(0.100\ L) = n\left(\frac{0.08206\ L\cdot atm}{mol\cdot K}\right)(300\ K) \tag{12.3.5}$$

$$n = 0.00401\ mol \tag{12.3.6}$$

Example 12.3.2 : unit considerations

A sample of benzene (C_6H_6) was heated to 100.°C in an evacuated flask whose volume was 247.2 ml, a sample of benzene vaporized. When the benzene was condensed to a liquid, its mass was found to be 0.616 g. What was the pressure in the flask?

Solution

The problem gives values for temperature, volume, and mass of the sample. Since R has units of \(\left (\frac{L\cdot atm}{mol\cdot K} \right)), we need to have the temperature in units of Kelvin, the volume in liters, and the amount of sample in moles.

Temperature:

K = °C + 273.15

K = 100.°C + 273.15

K = 373 K

Volume: $247.2\ mL\left(\frac{10^{-3}\ L}{1\ mL}\right) = 0.2472\ L$

Moles: $0.616\ g\left(\frac{1\ mol}{78.11\ g}\right) = 7.89\times10^{-3}\ mol$

Now that all of the values are in the correct units, the value for the unknown pressure can be determined.

$$PV = nRT \tag{12.3.7}$$

$$P\,(0.2472\ L) = \left(7.89\times10^{-3}\ mol\right)\left(\frac{0.08206\ L\cdot atm}{mol\cdot K}\right)(373\ K) \tag{12.3.8}$$

$$P = .977atm \tag{12.3.9}$$

Combined Gas law

While the ideal gas law is useful in solving for a single unknown when the other values are known, the combined gas law is useful when comparing initial and final situations. The ideal gas law can be rearranged to solve for R, the gas constant.

$$R = \frac{PV}{nT}$$

Under the initial conditions, $R = \frac{P_iV_i}{n_iT_i}$ and under final conditions, $R = \frac{P_fV_f}{n_fT_f}$. Since both expressions are equal to R, they are equal to each other.

$$\frac{P_iV_i}{n_iT_i} = \frac{P_fV_f}{n_fT_f}$$

This equation is typically used when one or more of the variables is constant. As a result, that variable is canceled from the equation. For example, the equation $2x^2 = 2y$ can be simplified to $x^2 = y$ since the 2 is on both sides of the equation.

What happens to the combined gas law equation when the initial and final pressures are equal ($P_i = P_f$)? Since they are equal, P_i can replace P_f.

$$\frac{P_iV_i}{n_iT_i} = \frac{P_iV_f}{n_fT_f}$$

which simplifies to

$$\frac{V_i}{n_iT_i} = \frac{V_f}{n_fT_f}$$

If two variables are constant, the equation can be simplified even more. If temperature and volume are constant, then $T_i = T_f$ and $V_i = V_f$. Then,

$$\frac{P_iV_i}{n_iT_i} = \frac{P_fV_i}{n_fT_i}$$

simplifies to

$$\frac{P_i}{n_i} = \frac{P_f}{n_f}$$

Example 12.3.3

Imagine a 1855 L balloon initially at 30°C and 745 mmHg. The balloon rises to an altitude of 23,000 ft and that the pressure and temperature at that altitude were 312 mmHg and −30°C, respectively. To what volume would the balloon have to expand to hold the same amount of hydrogen gas at the higher altitude?

Solution:

Begin by setting up a table of the two sets of conditions (note that some values will need to be converted to different units):

Initial	Final
$P_i = 745\ \text{mmHg} = 0.980\ \text{atm}$	$P_f = 312\ \text{mmHg} = 0.411\ \text{atm}$
$T_i = 30\ ^\circ\text{C} = 303\ \text{K}$	$T_f = -30\ ^\circ\text{C} = 243\ \text{K}$
$V_i = 1855\ \text{L}$	$V_f = ?$

By eliminating the constant property (n) of the gas, the combined gas law is simplified to

$$\frac{P_iV_i}{T_i} = \frac{P_fV_f}{T_f} \tag{12.3.10}$$

By solving the equation for V_f, we get:

$$\frac{P_iV_i}{T_i} = \frac{P_fV_f}{T_f} \tag{12.3.11}$$

$$\frac{0.980\ atm \cdot 1855\ L}{303\ K} = \frac{0.411\ atm \cdot V_f}{243\ K} \tag{12.3.12}$$

$$V_f = 3.55 \times 10^3\ L \tag{12.3.13}$$

Contributors

- Ed Vitz (Kutztown University), John W. Moore (UW-Madison), Justin Shorb (Hope College), Xavier Prat-Resina (University of Minnesota Rochester), Tim Wendorff, and Adam Hahn.
- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)
-

This page titled 12.3: The Ideal Gas Equation is shared under a not declared license and was authored, remixed, and/or curated by Allison Soult.

12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases

Learning Objectives

- Use the kinetic-molecular theory's postulates to explain the gas laws

The gas laws that we have seen to this point, as well as the ideal gas equation, are empirical, that is, they have been derived from experimental observations. The mathematical forms of these laws closely describe the macroscopic behavior of most gases at pressures less than about 1 or 2 atm. Although the gas laws describe relationships that have been verified by many experiments, they do not tell us why gases follow these relationships.

The test of the KMT and its postulates is its ability to explain and describe the behavior of a gas. The various gas laws can be derived from the assumptions of the KMT, which have led chemists to believe that the assumptions of the theory accurately represent the properties of gas molecules. We will first look at the individual gas laws (Boyle's, Charles's, Amontons's, Avogadro's, and Dalton's laws) conceptually to see how the KMT explains them. Then, we will more carefully consider the relationships between molecular masses, speeds, and kinetic energies with temperature, and explain Graham's law.

The Kinetic-Molecular Theory Explains the Behavior of Gases, Part I

Recalling that gas pressure is exerted by rapidly moving gas molecules and depends directly on the number of molecules hitting a unit area of the wall per unit of time, we see that the KMT conceptually explains the behavior of a gas as follows:

- *Amontons's law.* If the temperature is increased, the average speed and kinetic energy of the gas molecules increase. If the volume is held constant, the increased speed of the gas molecules results in more frequent and more forceful collisions with the walls of the container, therefore increasing the pressure (Figure 12.4.1*a*).
- *Charles's law.* If the temperature of a gas is increased, a constant pressure may be maintained only if the volume occupied by the gas increases. This will result in greater average distances traveled by the molecules to reach the container walls, as well as increased wall surface area. These conditions will decrease both the frequency of molecule-wall collisions and the number of collisions per unit area, the combined effects of which balance the effect of increased collision forces due to the greater kinetic energy at the higher temperature.
- *Boyle's law.* If the gas volume is decreased, the container wall area decreases and the molecule-wall collision frequency increases, both of which increase the pressure exerted by the gas (Figure 12.4.1*b*).
- *Avogadro's law.* At constant pressure and temperature, the frequency and force of molecule-wall collisions are constant. Under such conditions, increasing the number of gaseous molecules will require a proportional increase in the container volume in order to yield a decrease in the number of collisions per unit area to compensate for the increased frequency of collisions (Figure 12.4.1*c*).
- *Dalton's Law.* Because of the large distances between them, the molecules of one gas in a mixture bombard the container walls with the same frequency whether other gases are present or not, and the total pressure of a gas mixture equals the sum of the (partial) pressures of the individual gases.




Figure 12.4.1: (a) When gas temperature increases, gas pressure increases due to increased force and frequency of molecular collisions. (b) When volume decreases, gas pressure increases due to increased frequency of molecular collisions. (c) When the amount of gas increases at a constant pressure, volume increases to yield a constant number of collisions per unit wall area per unit time.

The Kinetic-Molecular Theory Explains the Behavior of Gases, Part II

According to Graham's law, the molecules of a gas are in rapid motion and the molecules themselves are small. The average distance between the molecules of a gas is large compared to the size of the molecules. As a consequence, gas molecules can move past each other easily and diffuse at relatively fast rates.

The rate of effusion of a gas depends directly on the (average) speed of its molecules:

$$\text{effusion rate} \propto u_{\text{rms}} \tag{12.4.1}$$

Using this relation, and the equation relating molecular speed to mass, Graham's law may be easily derived as shown here:

$$u_{\text{rms}} = \sqrt{\frac{3RT}{m}} \tag{12.4.2}$$

$$m = \frac{3RT}{u_{\text{rms}}^2} = \frac{3RT}{\overline{u}^2} \tag{12.4.3}$$

$$\frac{\text{effusion rate A}}{\text{effusion rate B}} = \frac{u_{\text{rms A}}}{u_{\text{rms B}}} = \frac{\sqrt{\frac{3RT}{m_{\text{A}}}}}{\sqrt{\frac{3RT}{m_{\text{B}}}}} = \sqrt{\frac{m_{\text{B}}}{m_{\text{A}}}} \tag{12.4.4}$$

The ratio of the rates of effusion is thus derived to be inversely proportional to the ratio of the square roots of their masses. This is the same relation observed experimentally and expressed as Graham's law.

Summary

The kinetic molecular theory is a simple but very effective model that effectively explains ideal gas behavior. The theory assumes that gases consist of widely separated molecules of negligible volume that are in constant motion, colliding elastically with one another and the walls of their container with average velocities determined by their absolute temperatures. The individual molecules of a gas exhibit a range of velocities, the distribution of these velocities being dependent on the temperature of the gas and the mass of its molecules.

Key Equations

- $u_{\text{rms}} = \sqrt{\overline{u^2}} = \sqrt{\dfrac{u_1^2 + u_2^2 + u_3^2 + u_4^2 + \ldots}{n}}$
- $\text{KE}_{\text{avg}} = \dfrac{3}{2}RT$
- $u_{\text{rms}} = \sqrt{\dfrac{3RT}{m}}$

Summary

kinetic molecular theory

theory based on simple principles and assumptions that effectively explains ideal gas behavior

root mean square velocity (u_{rms})

measure of average velocity for a group of particles calculated as the square root of the average squared velocity

Contributors and Attributions

- Paul Flowers (University of North Carolina - Pembroke), Klaus Theopold (University of Delaware) and Richard Langley (Stephen F. Austin State University) with contributing authors. Textbook content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at http://cnx.org/contents/85abf193-2bd...a7ac8df6@9.110).

This page titled 12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

12.5: Dalton's Law (Law of Partial Pressures)

Dalton’s Law, or the Law of Partial Pressures, states that the total pressure exerted by a mixture of gases is equal to the sum of the partial pressures of the gases in the mixture.

Dalton's Law of Partial Pressures

Based on the kinetic theory of gases, gas pressure results from collisions between gas particles and the inside walls of their container. If more gas is added to a rigid container, the gas pressure increases. The identities of the two gases do not matter. An ideal gas will expand in a container to fill up the space it is in and does not have any forces of attraction between the molecules. In other words, the different molecules in a mixture of gases are so far apart that they act independently; they do not react with each other. The pressure of an ideal gas is determined by its collisions with the container, not collisions with molecules of other substances since there are no other collisions. Since the gases in a mixture of gases are in one container, the Volume (V) and Temperature (T) for the different gases are the same as well. Each gas exerts its own pressure on the system, which can be added up to find the total pressure of the mixture of gases in a container.

John Dalton, the English chemist who proposed the atomic theory, also studied mixtures of gases. He found that each gas in a mixture exerts a pressure independently of every other gas in the mixture. For example, our atmosphere is composed of about 78% nitrogen and 21% oxygen, with smaller amounts of several other gases making up the rest. Since nitrogen makes up 78% of the gas particles in a given sample of air, it exerts 78% of the pressure. If the overall atmospheric pressure is $1.00\ \text{atm}$, then the pressure of just the nitrogen in the air is $0.78\ \text{atm}$. The pressure of the oxygen in the air is $0.21\ \text{atm}$.

The **partial pressure** of a gas is the contribution that gas makes to the total pressure when the gas is part of a mixture. The partial pressure of nitrogen is represented by P_{N_2}. **Dalton's law of partial pressures** states that the total pressure of a mixture of gases is equal to the sum of all of the partial pressures of the component gases. Dalton's law can be expressed with the following equation:

$$P_{\text{total}} = P_1 + P_2 + P_3 + \cdots \tag{12.5.1}$$

The figure below shows two gases that are in separate, equal-sized containers at the same temperature and pressure. Each exerts a different pressure, P_1 and P_2, reflective of the number of particles in the container. On the right, the two gases are combined into the same container, with no volume change. The total pressure of the gas mixture is equal to the sum of the individual pressures. If $P_1 = 300\ \text{mm Hg}$ and $P_2 = 500\ \text{mm Hg}$, then $P_{\text{total}} = 800\ \text{mm Hg}$.



Figure 14.12.1: Dalton's law says that the pressure of a gas mixture is equal to the partial pressures of the combining gases. (CC BY-NC; CK-12)

Derivation

We have, from the Ideal Gas Law

$$PV = nRT \tag{2}$$

If we know the molar composition of the gas, we can write

$$n_{total} = n_a + n_b + \ldots \tag{3}$$

Again, based on the kinetic theory of gases and the ideal gas law, Dalton's law can also be applied to the number of moles so that the total number of moles equals the sum of the number of moles of the individual gases. Here, the pressure, temperature and volume are held constant in the system. The total volume of a gas can be found the same way, although this is not used as much. This yields the equation,

$$P_{total} V = n_{total} RT \tag{4}$$

We can rearrange this equation to find the total number of moles. Sometimes masses of each sample of gas are given and students are asked to find the total pressure. This can be done by converting grams to moles and using Dalton's law to find the pressure.

Example

Suppose that we had 0.010 mol of a gas in a 250-ml container at a temperature of 32°C. The pressure would be

$$P = \frac{RT}{V} n = \frac{0.0820 \text{ liter atm mol}^{-1} \text{ K}^{-1} \times 305 \text{ K}}{0.250 \text{ liter}} \times 0.010 \text{ mol} \tag{12.5.2}$$

$$= 1.00 \text{ atm} \tag{12.5.3}$$

Now suppose we filled the same container with 0.004 mol $H_2(g)$ at the same temperature. The pressure would be

$$p_{H_2} = \frac{0.0820 \text{ liter atm mol}^{-1} \text{ K}^{-1} \times 305 \text{ K}}{0.250 \text{ liter}} \times 0.004 \text{ mol} \tag{12.5.4}$$

$$= 0.40 \text{ atm} \tag{12.5.5}$$

If we put 0.006 mol N_2 in the container,

$$p_{N_2} = \frac{0.0820 \text{ liter atm mol}^{-1} \text{ K}^{-1} \times 305 \text{ K}}{0.250 \text{ liter}} \times 0.006 \text{ mol} = 0.60 \text{ atm} \tag{12.5.6}$$

Now suppose we put both the 0.004 mol H_2 and the 0.006 mol N_2 into the same flask together. What would the pressure be? Since the ideal gas law does not depend on *which* gas we have but only on the amount of *any* gas, the pressure of the (0.004 + 0.006) mol, or 0.010 mol, would be exactly what we got in our first calculation. But this is just the sum of the pressure that H_2 would exert if it occupied the container alone plus the pressure of N_2 if it were the only gas present. That is,

$$P_{total} = p_{H_2} + p_{N_2} \tag{12.5.7}$$

The figure below demonstrates the concept of partial pressure in more concrete terms, showing the pressure of each gas alone in a container and then showing the gases combined pressure once mixed.



Figure 12.5.1: Above, the figure shows how Dalton's law of partial pressure works. Alone, Hydrogen has a pressure of 0.4 atm, while Nitrogen has a pressure of 0.6 atm. When placed in the same container, the total pressure is 1 atm, with each gas contributing its partial pressure. Image credits: OpenStax College

We have just worked out an example of **Dalton's law of partial pressures** (named for John Dalton, its discoverer). This law states that *in a mixture of two or more gases, the total pressure is the sum of the partial pressures of all the components*. The **partial pressure** of a gas is the pressure that gas would exert if it occupied the container by itself. Partial pressure is represented by a lowercase letter *p*.

Mole Ratio

From the partial pressure of a certain gas and the total pressure of a certain mixture, the mole ratio, called Xi, of a gas can be found. The mole ratio describes what fraction of the mixture is a specific gas. For example, if oxygen exerts 4 atm of pressure in a mixture and the total pressure of the system is 10 atm, the mole ratio would be 4/10 or 0.4. The mole ratio applies to pressure, volume, and moles as seen by the equation below. This also means that 0.4 moles of the mixture is made up of gas i.

$$X_i = \frac{P_i}{P_{tot}} = \frac{n_i}{n_{tot}} = \frac{V_i}{V_{tot}} \tag{5}$$

The mole ratio, (X_i) is often used to determine the composition of gases in a mixture. The sum of the mole ratios of each gas in a mixture should always equal one since they represent the proportion of each gas in the mixture.

Collection of a Gas Over Water

The Law of Partial Pressures is commonly applied in looking at the pressure of a closed container of gas and water. The total pressure of this system is the pressure that the gas exerts on the liquid. The gas is made up of whatever sample of gas there is plus the evaporated water. The pressure of the gas on the liquid consists of the pressure of the evaporated water and the pressure of the gas collected. Based on Dalton's law, the pressure of the gas collected can be calculated by subtracting the pressure of the water vapor from the total pressure. This is commonly encountered when a gas is collected by displacement of water, as shown in Figure 12.5.2.



Figure 12.5.2: The total pressure exerted by a wet gas is equal to the sum of the partial pressure of the gas itself + the vapor pressure of water at that temperature. (At 20°C the vapor pressure of water is 17.3 mmHg.)

Because the gas has been bubbled through water, it contains some water molecules and is said to be "wet." The total pressure of this wet gas is the sum of the partial pressure of the gas itself and the partial pressure of the water vapor it contains. The latter partial pressure is called **the vapor pressure** of water. It depends only on the temperature of the experiment and may be obtained from a handbook or from Table 1.

TABLE 12.5.1: Vapor Pressure of Water as a Function of Temperature.

Temperature(°C)	Vapor Pressure (mmHg)	Vapor Pressure (kPa)
0	4.6	0.61
5	6.5	0.87
10	9.2	1.23
15	12.8	1.71
20	17.5	2.33
25	23.8	3.17
30	31.8	4.24
50	92.5	12.33

70	233.7	31.16
75	289.1	38.63
80	355.1	47.34
85	433.6	57.81
90	525.8	70.10
95	633.9	84.51
100	760.0	101.32

Example 12.5.1: Volume of Hydrogen

Assume 0.321 g zinc metal is allowed to react with excess hydrochloric acid (an aqueous solution of HCl gas) according to the equation

$$\mathrm{Zn}(s) + 2\mathrm{HCL}(aq) \rightarrow \mathrm{ZnCl_2}(aq) + \mathrm{H_2}(g) \tag{12.5.8}$$

The resulting hydrogen gas is collected over water at 25°C, while the barometric pressure is 745.4 mmHg. What volume of wet hydrogen will be collected?

Solution From Table 1 we find that at 25°C the vapor pressure of water is 23.8 mmHg. Accordingly

$$p_{H_2} = p_{total} - p_{H_2O} = 754 \text{ mmHg} - 23.8 \text{ mmHg} = 721.6 \text{ mmHg}.$$

This must be converted to units compatible *R*:

$$p_{\mathrm{H_2}} = 721.6 \text{ mmHg} \times \frac{1 \text{ atm}}{760 \text{ mmHg}} = 0.949 \text{ atm} \tag{12.5.9}$$

The road map for this problem is

$$m_{\mathrm{Zn}} \xrightarrow{M_{\mathrm{Zn}}} n_{\mathrm{Zn}} \xrightarrow{S(\mathrm{H_2/Zn})} n_{\mathrm{H_2}} \xrightarrow{RT/P} V_{\mathrm{H_2}} \tag{12.5.10}$$

Thus

$$V_{\mathrm{H_2}} = 0.321 \text{ g Zn} \times \frac{1 \text{ mol Zn}}{65.38 \text{ g Zn}} \times \frac{1 \text{ mol H}_2}{2 \text{ mol Zn}} \times \frac{0.0820 \text{ liter atm}}{1 \text{ K mol H}_2} \times \frac{2930.15 \text{ K}}{0.987 \text{ atm}} \tag{12.5.11}$$

$$= 0.126 \text{ liter} \tag{12.5.12}$$

Real Gases

Real gases are gases that do not behave ideally. That is, they violate one or more of the rules of the kinetic theory of gases. Real gases behave ideally when the gases are at low pressure and high temperature. Therefore at high pressures and low temperatures, Dalton's law is not applicable since the gases are more likely to react and change the pressure of the system. For example, if there are forces of attraction between the molecules, the molecules would get closer together and the pressure would be adjusted because the molecules are interacting with each other.

Summary

- The total pressure in a system is equal to the sums of the partial pressures of the gases present.

Contributors and Attributions

- Ed Vitz (Kutztown University), John W. Moore (UW-Madison), Justin Shorb (Hope College), Xavier Prat-Resina (University of Minnesota Rochester), Tim Wendorff, and Adam Hahn.
-

Problems

1. A sample of gas A evaporates over water in a closed system. What is the pressure of gas A if the total pressure is 780 torr and water vapor pressure is 1 atm?
2. There is a mixture of 4 moles of hydrogen gas, 8 moles of oxygen, 12 moles of helium, and 6 moles of nitrogen in a closed container. What is the total number of moles in this system?
3. If there is a mixture of hydrogen and oxygen gas in a container with 10 moles total and the mole ratio of hydrogen is .67 $mol_{Hydrogen}$ to 1 mol_{total}, how many moles of each gas are there?
4. 24.0 L of nitrogen gas at 2 atm and 12.0 L of oxygen gas at 2 atm are added to a 10 L container at 273 K. Find the partial pressure of nitrogen and oxygen and then find the total pressure.
5. Flourine gas is in a 5.0 L container that is 25 C and 2 atm. A certain amount of hydrogen with a partial pressure of .5 is added to the container. What is the mole ratio of hydrogen?

Solutions

1:

- Convert pressure to same units so 780 torr=1.03 atm
- Subtract water vapor pressure from total pressure to get partial pressure of gas A: P_A=1.03 atm- 1 atm=**0.03 atm**

2. The law of partial pressures also applies to the total number of moles if the other values are constant, so

4 mol Hydrogen+8 mol Oxygen+12 mol Helium+6 mol Nitrogen=**30 moles total**

3.

- 10 mol_{total} x .67 $mol_{Hydrogen}$/ 1 mol_{total}= 6.7 moles H_2 gas
- 10 mol_{total}-6.7 $mol_{Hydrogen}$=3.3 moles O gas

4.

- Find the number of moles of oxygen and nitrogen using PV=nRT which is n=PV/RT
  1. oxygen: ((1 atm)(12L))/(0.08206 atm L mol^{-1} K^{-1})(273 K)=0.536 moles oxygen
  2. nitrogen: ((1 atm)(24.0L))/(.08206 atm L mol^{-1} K^{-1})(273 K)=1.07 moles of Nitrogen
  3. add to get n_{tot}: .536 mol_{Oxygen}+1.07 $mol_{Nitrogen}$=1.61 moles total
- Use PV=nRT or P=(nRT)/V to find the total pressure
  1. Ptot=((1.61 mol_{total})(0.08206 atm L mol^{-1} K^{-1})(273 K))/(10.0 L)=3.61 atm
- $P_A/P_{tot}=n_A/N_{tot}$ can be rearranged to $P_{A=(}P_{tot)(}n_A/N_{tot)}$ to find the partial pressures
  1. P_{oxygen}=(3.61 atm_{total})(.536 mol_{Oxygen}/1.61 mol_{total})=1.20 atm_{Oxygen}
  2. $P_{nitrogen}$=3.61 atm_{total}-1.20 atm_{Oxygen}=2.41 $atm_{Nitrogen}$

5.

- total pressure is 2 $atm_{Flourine}$+.5 $atm_{Hydrogen}$=2.5 atm_{total}
- remember that $p_H/p_{tot}=n_H/n_{tot}=v_h/v_{tot}$ so the mole ratio of hydrogen to the mixture is is .5/2.5=**.2 $mol_{Hydrogen}$ to 1 mol_{total}**

References

1. Dutton, F.B. "Dalton's Law of Partial Pressures." Journal of Chemical Education Aug 1961. http://pubs.acs.org/doi/pdfplus/10.1021/ed038pA545.1
2. Blauch, David. "Gas Laws." Davidson Chemistry. 2009. http://www.chm.davidson.edu/vce/gaslaws/daltonslaw.html
3. Petrucci, Ralph H. "General Chemistry: Principles & Modern Application, 9th Edition." New Jersey: Pearson Prentice Hall. 2007.

4. Toupadakis, Andreas. "Chemistry Reader 2A." Kendall Hunt Publishing. 2009.
5. http://pubs.acs.org/doi/pdfplus/10.1021/ed038pA545.1
6. http://pubs.acs.org/doi/pdf/10.1021/ed084p469

12.5: Dalton's Law (Law of Partial Pressures) is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

12.6: Evaporation and Condensation

Learning Objectives

- Explain the following laws within the Ideal Gas Law

On the roof of the house in the picture below is a device known as a "swamp cooler". This piece of equipment traces its origin back to the ancient Egyptians who hung wet blankets across the doors of their homes. As the warm air passed through the blankets, water would evaporate and cool the air. The royalty went one step further and had servants fan wet cloths over jugs of water to get more evaporation and cooling.



Figure 12.6.1: A home with a swamp cooler attached to the roof.

The origin of the term "swamp cooler" is not known - they certainly don't work in a swamp. Best conditions for cooling include a high temperature (over $80^{o}\mathrm{F}$) and a low humidity (preferably less than 30%). These coolers work well in desert areas, but don't provide any cooling in the humid areas of the country.

Evaporation

A puddle of water left undisturbed eventually disappears. The liquid molecules escape into the gas phase, becoming water vapor. **Vaporization** is the process in which a liquid is converted to a gas. **Evaporation** is the conversion of a liquid to its vapor below the boiling temperature of the liquid. If the water is instead kept in a closed container, the water vapor molecules do not have a chance to escape into the surroundings and so the water level does not change. As some water molecules become vapor, an equal number of water vapor molecules condense back into the liquid state. **Condensation** is the change of state from a gas to a liquid.




Figure 12.6.2: Evaporation (A) and condensation (B).

In order for a liquid molecule to escape into the gas state, the molecule must have enough kinetic energy to overcome the intermolecular attractive forces in the liquid. Recall that a given liquid sample will have molecules with a wide range of kinetic energies. Liquid molecules that have this certain threshold kinetic energy escape the surface and become vapor. As a result, the liquid molecules that remain now have lower kinetic energy. As evaporation occurs, the temperature of the remaining liquid decreases. You have observed the effects of evaporative cooling. On a hot day, the water molecules in your perspiration absorb body heat and evaporate from the surface of your skin. The evaporating process leaves the remaining perspiration cooler, which in turn absorbs more heat from your body.

A given liquid will evaporate more quickly when it is heated. This is because the heating process results in a greater fraction of the liquid's molecules having the necessary kinetic energy to escape the surface of the liquid. The figure below shows the kinetic energy distribution of liquid molecules at two temperatures. The numbers of molecules that have the required kinetic energy to

evaporate are shown in the shaded area under the curve at the right. The higher temperature liquid (T_2) has more molecules that are capable of escaping into the vapor phase than the lower temperature liquid (T_1).




Figure 12.6.3: Kinetic energy distribution curves for a liquid at two temperatures T_1 and T_2. The shaded area is the molecules with enough kinetic energy to escape the liquid and become vapor.

At 29,029 feet $(8848\ \text{m})$, Mount Everest in the Himalayan range on the border between China and Nepal is the highest point on the earth. Its altitude presents many practical problems to climbers. The oxygen content of the air is much lower than at sea level, making it necessary to bring oxygen tanks along (although a few climbers have reached the peak without oxygen). One other problem is that of boiling water for cooking food. Although water boils at 100°C at sea level, the boiling point on top of Mount Everest is only about 70°C. This difference makes it very difficult to get a decent cup of tea (which definitely frustrated some of the British climbers).

Boiling

As a liquid is heated, the average kinetic energy of its particles increases. The rate of evaporation increases as more and more molecules are able to escape the liquid's surface into the vapor phase. Eventually a point is reached when the molecules all throughout the liquid have enough kinetic energy to **vaporize**. At this point the liquid begins to boil. The **boiling point** is the temperature at which the vapor pressure of a liquid is equal to the external pressure. The figure below illustrates the boiling of liquid.




Figure 12.6.4: Comparison between evaporation and boiling.

In the picture on the left, the liquid is below its boiling point, yet some of the liquid evaporates. On the right, the temperature has been increased until bubbles begin to form in the body of the liquid. When the vapor pressure inside the bubble is equal to the external atmospheric pressure, the bubbles rise to the surface of the liquid and burst. The temperature at which this process occurs is the boiling point of the liquid.

The normal boiling point is the temperature at which the vapor pressure of the liquid is equal to standard pressure. Because atmospheric pressure can change based on location, the boiling point of a liquid changes with the external pressure. The normal boiling point is a constant because it is defined relative to the standard atmospheric pressure of $760\ \text{mm Hg}$ (or $1\ \text{atm}$ or $101.3\ \text{kPa}$).




Figure 12.6.5: Influence of altitude on the boiling point of water.

Summary




Figure 12.6.1: Change of State Comparison

- The boiling point is the temperature at which the vapor pressure of a liquid is equal to the external pressure.
- As the altitude increases, the boiling point decreases.
- Evaporation is the conversion of a liquid to its vapor below the boiling temperature of the liquid.
- Condensation is the change of state from a gas to a liquid.
- As the temperature increases, the rate of evaporation increases.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

-
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

12.6: Evaporation and Condensation is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

12.7: Intermolecular forces

Learning Objective

- To describe the intermolecular forces in liquids.

The properties of liquids are intermediate between those of gases and solids but are more similar to solids. In contrast to *intra*molecular forces, such as the covalent bonds that hold atoms together in molecules and polyatomic ions, *inter*molecular forces hold molecules together in a liquid or solid. Intermolecular forces are generally much weaker than covalent bonds. For example, it requires 927 kJ to overcome the intramolecular forces and break both O–H bonds in 1 mol of water, but it takes only about 41 kJ to overcome the intermolecular attractions and convert 1 mol of liquid water to water vapor at 100°C. (Despite this seemingly low value, the intermolecular forces in liquid water are among the strongest such forces known!) Given the large difference in the strengths of intra- and intermolecular forces, changes between the solid, liquid, and gaseous states almost invariably occur for molecular substances *without breaking covalent bonds*.

Note the Pattern

The properties of liquids are intermediate between those of gases and solids but are more similar to solids.

Intermolecular forces determine bulk properties such as the melting points of solids and the boiling points of liquids. Liquids boil when the molecules have enough thermal energy to overcome the intermolecular attractive forces that hold them together, thereby forming bubbles of vapor within the liquid. Similarly, solids melt when the molecules acquire enough thermal energy to overcome the intermolecular forces that lock them into place in the solid.

Intermolecular forces are electrostatic in nature; that is, they arise from the interaction between positively and negatively charged species. Like covalent and ionic bonds, intermolecular interactions are the sum of both attractive and repulsive components. Because electrostatic interactions fall off rapidly with increasing distance between molecules, intermolecular interactions are most important for solids and liquids, where the molecules are close together. These interactions become important for gases only at very high pressures, where they are responsible for the observed deviations from the ideal gas law at high pressures.

In this section, we explicitly consider three kinds of intermolecular interactions:There are two additional types of electrostatic interaction that you are already familiar with: the ion–ion interactions that are responsible for ionic bonding and the ion–dipole interactions that occur when ionic substances dissolve in a polar substance such as water. The intermolecular forces between neutral molecules are *dipole–dipole interactions*, *London dispersion forces*, and *hydrogen bonds*. The first two are often described collectively as van der Waals forcesThe intermolecular forces known as dipole–dipole interactions and London dispersion forces..

Dipole–Dipole Interactions

Recall that polar covalent bonds behave as if the bonded atoms have localized fractional charges that are equal but opposite (i.e., the two bonded atoms generate a *dipole*). If the structure of a molecule is such that the individual bond dipoles do not cancel one another, then the molecule has a net dipole moment. Molecules with net dipole moments tend to align themselves so that the positive end of one dipole is near the negative end of another and vice versa, as shown in part (a) in Figure 12.7.1 . These arrangements are more stable than arrangements in which two positive or two negative ends are adjacent (part (c) in Figure 12.7.1). Hence dipole–dipole interactionsA kind of intermolecular interaction (force) that results between molecules with net dipole moments., such as those in part (b) in Figure 12.7.1 , are *attractive intermolecular interactions*, whereas those in part (d) in Figure 12.7.1 are *repulsive intermolecular interactions*. Because molecules in a liquid move freely and continuously, molecules always experience both attractive and repulsive dipole–dipole interactions simultaneously, as shown in Figure 12.7.2 . On average, however, the attractive interactions dominate.




Figure 12.7.1 Attractive and Repulsive Dipole–Dipole Interactions *(a and b) Molecular orientations in which the positive end of one dipole (δ^+) is near the negative end of another (δ^-) (and vice versa) produce attractive interactions. (c and d) Molecular orientations that juxtapose the positive or negative ends of the dipoles on adjacent molecules produce repulsive interactions.*




Figure 12.7.2 Both Attractive and Repulsive Dipole–Dipole Interactions Occur in a Liquid Sample with Many Molecules

Because each end of a dipole possesses only a fraction of the charge of an electron, dipole–dipole interactions are substantially weaker than the interactions between two ions, each of which has a charge of at least ±1, or between a dipole and an ion, in which one of the species has at least a full positive or negative charge. In addition, the attractive interaction between dipoles falls off much more rapidly with increasing distance than do ion–ion interactions. The attractive energy between two ions is proportional to $1/r$, where r is the distance between the ions. Doubling the distance ($r \rightarrow 2r$) decreases the attractive energy by one-half. In contrast, the energy of the interaction of two dipoles is proportional to $1/r^6$, so doubling the distance between the dipoles decreases the strength of the interaction by 2^6, or 64-fold. Thus a substance such as HCl, which is partially held together by dipole–dipole interactions, is a gas at room temperature and 1 atm pressure, whereas NaCl, which is held together by interionic interactions, is a high-melting-point solid. Within a series of compounds of similar molar mass, the strength of the intermolecular interactions increases as the dipole moment of the molecules increases, as shown in Table 12.7.1. Using what we learned previously about predicting relative bond polarities from the electronegativities of the bonded atoms, we can make educated guesses about the relative boiling points of similar molecules.

Table 12.7.1 Relationships between the Dipole Moment and the Boiling Point for Organic Compounds of Similar Molar Mass


Compound	Molar Mass (g/mol)	Dipole Moment (D)	Boiling Point (K)
C_3H_6 (cyclopropane)	42	0	240
CH_3OCH_3 (dimethyl ether)	46	1.30	248
CH_3CN (acetonitrile)	41	3.9	355

Note the Pattern

The attractive energy between two ions is proportional to $1/r$, whereas the attractive energy between two dipoles is proportional to $1/r^6$.

Example 12.7.1

Arrange ethyl methyl ether ($CH_3OCH_2CH_3$), 2-methylpropane [isobutane, $(CH_3)_2CHCH_3$], and acetone (CH_3COCH_3) in order of increasing boiling points. Their structures are as follows:



Given: compounds

Asked for: order of increasing boiling points

Strategy:

Compare the molar masses and the polarities of the compounds. Compounds with higher molar masses and that are polar will have the highest boiling points.

Solution:

The three compounds have essentially the same molar mass (58–60 g/mol), so we must look at differences in polarity to predict the strength of the intermolecular dipole–dipole interactions and thus the boiling points of the compounds. The first compound, 2-methylpropane, contains only C–H bonds, which are not very polar because C and H have similar electronegativities. It should therefore have a very small (but nonzero) dipole moment and a very low boiling point. Ethyl methyl ether has a structure similar to H_2O; it contains two polar C–O single bonds oriented at about a 109° angle to each other, in addition to relatively nonpolar C–H bonds. As a result, the C–O bond dipoles partially reinforce one another and generate a significant dipole moment that should give a moderately high boiling point. Acetone contains a polar C=O double bond oriented at about 120° to two methyl groups with nonpolar C–H bonds. The C–O bond dipole therefore corresponds to the molecular dipole, which should result in both a rather large dipole moment and a high boiling point. Thus we predict the following order of boiling points: 2-methylpropane < ethyl methyl ether < acetone. This result is in good agreement with the actual data: 2-methylpropane, boiling point = −11.7°C, and the dipole moment (μ) = 0.13 D; methyl ethyl ether, boiling point = 7.4°C and μ = 1.17 D; acetone, boiling point = 56.1°C and μ = 2.88 D.

Exercise

Arrange carbon tetrafluoride (CF_4), ethyl methyl sulfide ($CH_3SC_2H_5$), dimethyl sulfoxide [$(CH_3)_2S{=}O$], and 2-methylbutane [isopentane, $(CH_3)_2CHCH_2CH_3$] in order of decreasing boiling points.

Answer: dimethyl sulfoxide (boiling point = 189.9°C) > ethyl methyl sulfide (boiling point = 67°C) > 2-methylbutane (boiling point = 27.8°C) > carbon tetrafluoride (boiling point = −128°C)

London Dispersion Forces

Thus far we have considered only interactions between polar molecules, but other factors must be considered to explain why many nonpolar molecules, such as bromine, benzene, and hexane, are liquids at room temperature, and others, such as iodine and naphthalene, are solids. Even the noble gases can be liquefied or solidified at low temperatures, high pressures, or both (Table 12.7.2)

What kind of attractive forces can exist between nonpolar molecules or atoms? This question was answered by Fritz London (1900–1954), a German physicist who later worked in the United States. In 1930, London proposed that temporary fluctuations in the electron distributions within atoms and nonpolar molecules could result in the formation of short-lived instantaneous dipole momentsThe short-lived dipole moment in atoms and nonpolar molecules caused by the constant motion of their electrons, which results in an asymmetrical distribution of charge at any given instant., which produce attractive forces called London dispersion forcesA kind of intermolecular interaction (force) that results from temporary fluctuations in the electron distribution within atoms and nonpolar molecules. between otherwise nonpolar substances.

Table 12.7.2 Normal Melting and Boiling Points of Some Elements and Nonpolar Compounds

Substance	Molar Mass (g/mol)	Melting Point (°C)	Boiling Point (°C)
Ar	40	−189.4	−185.9
Xe	131	−111.8	−108.1
N_2	28	−210	−195.8
O_2	32	−218.8	−183.0
F_2	38	−219.7	−188.1
I_2	254	113.7	184.4
CH_4	16	−182.5	−161.5

Consider a pair of adjacent He atoms, for example. On average, the two electrons in each He atom are uniformly distributed around the nucleus. Because the electrons are in constant motion, however, their distribution in one atom is likely to be asymmetrical at any given instant, resulting in an instantaneous dipole moment. As shown in part (a) in Figure 12.7.3, the instantaneous dipole moment on one atom can interact with the electrons in an adjacent atom, pulling them toward the positive end of the instantaneous dipole or repelling them from the negative end. The net effect is that the first atom causes the temporary formation of a dipole, called an induced dipoleA short-lived dipole moment that is created in atoms and nonpolar molecules adjacent to atoms or molecules with an instantaneous dipole moment., in the second. Interactions between these temporary dipoles cause atoms to be attracted to one another. These attractive interactions are weak and fall off rapidly with increasing distance. London was able to show with quantum mechanics that the attractive energy between molecules due to temporary dipole–induced dipole interactions falls off as $1/r^6$. Doubling the distance therefore decreases the attractive energy by 2^6, or 64-fold.




Figure 12.7.3 Instantaneous Dipole Moments *The formation of an instantaneous dipole moment on one He atom (a) or an H_2 molecule (b) results in the formation of an induced dipole on an adjacent atom or molecule.*

Instantaneous dipole–induced dipole interactions between nonpolar molecules can produce intermolecular attractions just as they produce interatomic attractions in monatomic substances like Xe. This effect, illustrated for two H_2 molecules in part (b) in Figure 12.7.3 tends to become more pronounced as atomic and molecular masses increase (Table 12.7.2). For example, Xe boils at −108.1°C, whereas He boils at −269°C. The reason for this trend is that the strength of London dispersion forces is related to the ease with which the electron distribution in a given atom can be perturbed. In small atoms such as He, the two 1*s* electrons are held

close to the nucleus in a very small volume, and electron–electron repulsions are strong enough to prevent significant asymmetry in their distribution. In larger atoms such as Xe, however, the outer electrons are much less strongly attracted to the nucleus because of filled intervening shells. As a result, it is relatively easy to temporarily deform the electron distribution to generate an instantaneous or induced dipole. The ease of deformation of the electron distribution in an atom or molecule is called its polarizability. Because the electron distribution is more easily perturbed in large, heavy species than in small, light species, we say that heavier substances tend to be much more *polarizable* than lighter ones.

Note the Pattern

For similar substances, London dispersion forces get stronger with increasing molecular size.

The polarizability of a substance also determines how it interacts with ions and species that possess permanent dipoles. Thus London dispersion forces are responsible for the general trend toward higher boiling points with increased molecular mass and greater surface area in a homologous series of compounds, such as the alkanes (part (a) in Figure 12.7.4). The strengths of London dispersion forces also depend significantly on molecular shape because shape determines how much of one molecule can interact with its neighboring molecules at any given time. For example, part (b) in Figure 12.7.4 shows 2,2-dimethylpropane (neopentane) and *n*-pentane, both of which have the empirical formula C_5H_{12}. Neopentane is almost spherical, with a small surface area for intermolecular interactions, whereas *n*-pentane has an extended conformation that enables it to come into close contact with other *n*-pentane molecules. As a result, the boiling point of neopentane (9.5°C) is more than 25°C lower than the boiling point of *n*-pentane (36.1°C).




Figure 12.7.4 Mass and Surface Area Affect the Strength of London Dispersion Forces *(a) In this series of four simple alkanes, larger molecules have stronger London forces between them than smaller molecules and consequently higher boiling points. (b) Linear n-pentane molecules have a larger surface area and stronger intermolecular forces than spherical neopentane molecules. As a result, neopentane is a gas at room temperature, whereas n-pentane is a volatile liquid.*

All molecules, whether polar or nonpolar, are attracted to one another by London dispersion forces in addition to any other attractive forces that may be present. In general, however, dipole–dipole interactions in small polar molecules are significantly stronger than London dispersion forces, so the former predominate.

Example 12.7.2

Arrange *n*-butane, propane, 2-methylpropane [isobutene, $(CH_3)_2CHCH_3$], and *n*-pentane in order of increasing boiling points.

Given: compounds

Asked for: order of increasing boiling points

Strategy:

Determine the intermolecular forces in the compounds and then arrange the compounds according to the strength of those forces. The substance with the weakest forces will have the lowest boiling point.

Solution:

The four compounds are alkanes and nonpolar, so London dispersion forces are the only important intermolecular forces. These forces are generally stronger with increasing molecular mass, so propane should have the lowest boiling point and *n*-pentane should have the highest, with the two butane isomers falling in between. Of the two butane isomers, 2-methylpropane is more compact, and *n*-butane has the more extended shape. Consequently, we expect intermolecular interactions for *n*-butane to be stronger due to its larger surface area, resulting in a higher boiling point. The overall order is thus as follows, with actual boiling points in parentheses: propane (−42.1°C) < 2-methylpropane (−11.7°C) < *n*-butane (−0.5°C) < *n*-pentane (36.1°C).

Exercise

Arrange GeH_4, $SiCl_4$, SiH_4, CH_4, and $GeCl_4$ in order of decreasing boiling points.

Answer: $GeCl_4$ (87°C) > $SiCl_4$ (57.6°C) > GeH_4 (−88.5°C) > SiH_4 (−111.8°C) > CH_4 (−161°C)

Hydrogen Bonds

Molecules with hydrogen atoms bonded to electronegative atoms such as O, N, and F (and to a much lesser extent Cl and S) tend to exhibit unusually strong intermolecular interactions. These result in much higher boiling points than are observed for substances in which London dispersion forces dominate, as illustrated for the covalent hydrides of elements of groups 14–17 in Figure 12.7.5 . Methane and its heavier congeners in group 14 form a series whose boiling points increase smoothly with increasing molar mass. This is the expected trend in nonpolar molecules, for which London dispersion forces are the exclusive intermolecular forces. In contrast, the hydrides of the lightest members of groups 15–17 have boiling points that are more than 100°C greater than predicted on the basis of their molar masses. The effect is most dramatic for water: if we extend the straight line connecting the points for H_2Te and H_2Se to the line for period 2, we obtain an estimated boiling point of −130°C for water! Imagine the implications for life on Earth if water boiled at −130°C rather than 100°C.




Figure 12.7.5 The Effects of Hydrogen Bonding on Boiling Points *These plots of the boiling points of the covalent hydrides of the elements of groups 14–17 show that the boiling points of the lightest members of each series for which hydrogen bonding is possible (HF, NH_3, and H_2O) are anomalously high for compounds with such low molecular masses.*

Why do strong intermolecular forces produce such anomalously high boiling points and other unusual properties, such as high enthalpies of vaporization and high melting points? The answer lies in the highly polar nature of the bonds between hydrogen and very electronegative elements such as O, N, and F. The large difference in electronegativity results in a large partial positive charge on hydrogen and a correspondingly large partial negative charge on the O, N, or F atom. Consequently, H–O, H–N, and H–F bonds have very large bond dipoles that can interact strongly with one another. Because a hydrogen atom is so small, these dipoles can also approach one another more closely than most other dipoles. The combination of large bond dipoles and short dipole–dipole distances results in very strong dipole–dipole interactions called hydrogen bondsAn unusually strong dipole-dipole interaction (intermolecular force) that results when hydrogen is bonded to very electronegative elements, such as O, N, and F., as shown for ice in Figure 12.7.6 . A hydrogen bond is usually indicated by a dotted line between the hydrogen atom attached to O, N, or F (the *hydrogen bond donor*) and the atom that has the lone pair of electrons (the *hydrogen bond acceptor*). Because each water molecule contains two hydrogen atoms and two lone pairs, a tetrahedral arrangement maximizes the number of hydrogen bonds that can be

formed. In the structure of ice, each oxygen atom is surrounded by a distorted tetrahedron of hydrogen atoms that form bridges to the oxygen atoms of adjacent water molecules. The bridging hydrogen atoms are *not* equidistant from the two oxygen atoms they connect, however. Instead, each hydrogen atom is 101 pm from one oxygen and 174 pm from the other. In contrast, each oxygen atom is bonded to two H atoms at the shorter distance and two at the longer distance, corresponding to two O–H covalent bonds and two O·H hydrogen bonds from adjacent water molecules, respectively. The resulting open, cagelike structure of ice means that the solid is actually slightly less dense than the liquid, which explains why ice floats on water rather than sinks. Remember that the hydrogens in one water molecule point towards one of the lone pairs of tetrahedal sp^3 orbitals of the oxygen atom on another water molecule.




Figure 12.7.6 The Hydrogen-Bonded Structure of Ice *Each water molecule accepts two hydrogen bonds from two other water molecules and donates two hydrogen atoms to form hydrogen bonds with two more water molecules, producing an open, cagelike structure. The structure of liquid water is very similar, but in the liquid, the hydrogen bonds are continually broken and formed because of rapid molecular motion.*

Note the Pattern

Hydrogen bond formation requires *both* a hydrogen bond donor *and* a hydrogen bond acceptor.

Because ice is less dense than liquid water, rivers, lakes, and oceans freeze from the top down. In fact, the ice forms a protective surface layer that insulates the rest of the water, allowing fish and other organisms to survive in the lower levels of a frozen lake or sea. If ice were denser than the liquid, the ice formed at the surface in cold weather would sink as fast as it formed. Bodies of water would freeze from the bottom up, which would be lethal for most aquatic creatures. The expansion of water when freezing also explains why automobile or boat engines must be protected by “antifreeze” and why unprotected pipes in houses break if they are allowed to freeze.

Although hydrogen bonds are significantly weaker than covalent bonds, with typical dissociation energies of only 15–25 kJ/mol, they have a significant influence on the physical properties of a compound. Compounds such as HF can form only two hydrogen bonds at a time as can, on average, pure liquid NH_3. Consequently, even though their molecular masses are similar to that of water, their boiling points are significantly lower than the boiling point of water, which forms *four* hydrogen bonds at a time.

Example 12.7.3

Considering CH_3OH, C_2H_6, Xe, and $(CH_3)_3N$, which can form hydrogen bonds with themselves? Draw the hydrogen-bonded structures.

Given: compounds

Asked for: formation of hydrogen bonds and structure

Strategy:


A Identify the compounds with a hydrogen atom attached to O, N, or F. These are likely to be able to act as hydrogen bond donors.

B Of the compounds that can act as hydrogen bond donors, identify those that also contain lone pairs of electrons, which allow them to be hydrogen bond acceptors. If a substance is both a hydrogen donor and a hydrogen bond acceptor, draw a structure showing the hydrogen bonding.

Solution:

A Of the species listed, xenon (Xe), ethane (C_2H_6), and trimethylamine [$(CH_3)_3N$] do not contain a hydrogen atom attached to O, N, or F; hence they cannot act as hydrogen bond donors.


B The one compound that can act as a hydrogen bond donor, methanol (CH_3OH), contains both a hydrogen atom attached to O (making it a hydrogen bond donor) and two lone pairs of electrons on O (making it a hydrogen bond acceptor); methanol can thus form hydrogen bonds by acting as either a hydrogen bond donor or a hydrogen bond acceptor. The hydrogen-bonded structure of methanol is as follows:




Exercise

Considering CH_3CO_2H, $(CH_3)_3N$, NH_3, and CH_3F, which can form hydrogen bonds with themselves? Draw the hydrogen-bonded structures.

Answer: CH_3CO_2H and NH_3;





Example 12.7.4

Arrange C_{60} (buckminsterfullerene, which has a cage structure), NaCl, He, Ar, and N_2O in order of increasing boiling points.

Given: compounds

Asked for: order of increasing boiling points

Strategy:

Identify the intermolecular forces in each compound and then arrange the compounds according to the strength of those forces. The substance with the weakest forces will have the lowest boiling point.

Solution:

Electrostatic interactions are strongest for an ionic compound, so we expect NaCl to have the highest boiling point. To predict the relative boiling points of the other compounds, we must consider their polarity (for dipole–dipole interactions), their ability to form hydrogen bonds, and their molar mass (for London dispersion forces). Helium is nonpolar and by far the lightest, so it should have the lowest boiling point. Argon and N_2O have very similar molar masses (40 and 44 g/mol, respectively), but N_2O is polar while Ar is not. Consequently, N_2O should have a higher boiling point. A C_{60} molecule is nonpolar, but its molar mass is 720 g/mol, much greater than that of Ar or N_2O. Because the boiling points of nonpolar substances increase rapidly with molecular mass, C_{60} should boil at a higher temperature than the other nonionic substances. The predicted order is thus as follows, with actual boiling points in parentheses: He (−269°C) < Ar (−185.7°C) < N_2O (−88.5°C) < C_{60} (>280°C) < NaCl (1465°C).

Exercise

Arrange 2,4-dimethylheptane, Ne, CS_2, Cl_2, and KBr in order of decreasing boiling points.

Answer: KBr (1435°C) > 2,4-dimethylheptane (132.9°C) > CS_2 (46.6°C) > Cl_2 (−34.6°C) > Ne (−246°C)

Summary

Molecules in liquids are held to other molecules by intermolecular interactions, which are weaker than the intramolecular interactions that hold the atoms together within molecules and polyatomic ions. Transitions between the solid and liquid or the liquid and gas phases are due to changes in intermolecular interactions but do not affect intramolecular interactions. The three

major types of intermolecular interactions are dipole–dipole interactions, London dispersion forces (these two are often referred to collectively as **van der Waals forces**), and hydrogen bonds. **Dipole–dipole interactions** arise from the electrostatic interactions of the positive and negative ends of molecules with permanent dipole moments; their strength is proportional to the magnitude of the dipole moment and to $1/r^6$, where r is the distance between dipoles. **London dispersion forces** are due to the formation of **instantaneous dipole moments** in polar or nonpolar molecules as a result of short-lived fluctuations of electron charge distribution, which in turn cause the temporary formation of an **induced dipole** in adjacent molecules. Like dipole–dipole interactions, their energy falls off as $1/r^6$. Larger atoms tend to be more **polarizable** than smaller ones because their outer electrons are less tightly bound and are therefore more easily perturbed. **Hydrogen bonds** are especially strong dipole–dipole interactions between molecules that have hydrogen bonded to a highly electronegative atom, such as O, N, or F. The resulting partially positively charged H atom on one molecule (the *hydrogen bond donor*) can interact strongly with a lone pair of electrons of a partially negatively charged O, N, or F atom on adjacent molecules (the *hydrogen bond acceptor*). Because of strong O·H> hydrogen bonding between water molecules, water has an unusually high boiling point, and ice has an open, cagelike structure that is less dense than liquid water.

Key Takeaway


- Intermolecular forces are electrostatic in nature and include van der Waals forces and hydrogen bonds.

Conceptual Problems

1. What is the main difference between intramolecular interactions and intermolecular interactions? Which is typically stronger? How are changes of state affected by these different kinds of interactions?
2. Describe the three major kinds of intermolecular interactions discussed in this chapter and their major features. The hydrogen bond is actually an example of one of the other two types of interaction. Identify the kind of interaction that includes hydrogen bonds and explain why hydrogen bonds fall into this category.
3. Which are stronger—dipole–dipole interactions or London dispersion forces? Which are likely to be more important in a molecule with heavy atoms? Explain your answers.
4. Explain why hydrogen bonds are unusually strong compared to other dipole–dipole interactions. How does the strength of hydrogen bonds compare with the strength of covalent bonds?
5. Liquid water is essential for life as we know it, but based on its molecular mass, water should be a gas under standard conditions. Why is water a liquid rather than a gas under standard conditions?
6. Describe the effect of polarity, molecular mass, and hydrogen bonding on the melting point and boiling point of a substance.
7. Why are intermolecular interactions more important for liquids and solids than for gases? Under what conditions must these interactions be considered for gases?
8. Using acetic acid as an example, illustrate both attractive and repulsive intermolecular interactions. How does the boiling point of a substance depend on the magnitude of the repulsive intermolecular interactions?
9. In group 17, elemental fluorine and chlorine are gases, whereas bromine is a liquid and iodine is a solid. Why?
10. The boiling points of the anhydrous hydrogen halides are as follows: HF, 19°C; HCl, −85°C; HBr, −67°C; and HI, −34°C. Explain any trends in the data, as well as any deviations from that trend.
11. Identify the most important intermolecular interaction in each of the following.
    1. SO_2
    2. HF
    3. CO_2
    4. CCl_4
    5. CH_2Cl_2
12. Identify the most important intermolecular interaction in each of the following.
    1. LiF
    2. I_2
    3. ICl
    4. NH_3

5. NH_2Cl

13. Would you expect London dispersion forces to be more important for Xe or Ne? Why? (The atomic radius of Ne is 38 pm, whereas that of Xe is 108 pm.)
14. Arrange Kr, Cl_2, H_2, N_2, Ne, and O_2 in order of increasing polarizability. Explain your reasoning.
15. Both water and methanol have anomalously high boiling points due to hydrogen bonding, but the boiling point of water is greater than that of methanol despite its lower molecular mass. Why? Draw the structures of these two compounds, including any lone pairs, and indicate potential hydrogen bonds.
16. The structures of ethanol, ethylene glycol, and glycerin are as follows:



Arrange these compounds in order of increasing boiling point. Explain your rationale.

17. Do you expect the boiling point of H_2S to be higher or lower than that of H_2O? Justify your answer.
18. Ammonia (NH_3), methylamine (CH_3NH_2), and ethylamine ($CH_3CH_2NH_2$) are gases at room temperature, while propylamine ($CH_3CH_2CH_2NH_2$) is a liquid at room temperature. Explain these observations.
19. Why is it not advisable to freeze a sealed glass bottle that is completely filled with water? Use both macroscopic and microscopic models to explain your answer. Is a similar consideration required for a bottle containing pure ethanol? Why or why not?
20. Which compound in the following pairs will have the higher boiling point? Explain your reasoning.
    1. NH_3 or PH_3
    2. ethylene glycol ($HOCH_2CH_2OH$) or ethanol
    3. 2,2-dimethylpropanol [$CH_3C(CH_3)_2CH_2OH$] or *n*-butanol ($CH_3CH_2CH_2CH_2OH$)
21. Some recipes call for vigorous boiling, while others call for gentle simmering. What is the difference in the temperature of the cooking liquid between boiling and simmering? What is the difference in energy input?
22. Use the melting of a metal such as lead to explain the process of melting in terms of what is happening at the molecular level. As a piece of lead melts, the temperature of the metal remains constant, even though energy is being added continuously. Why?
23. How does the O–H distance in a hydrogen bond in liquid water compare with the O–H distance in the covalent O–H bond in the H_2O molecule? What effect does this have on the structure and density of ice?
24. 1. Explain why the hydrogen bonds in liquid HF are stronger than the corresponding intermolecular <math display="inline" xml:id="av_1.0-ch11_m003"><semantics><mrow><mtext>H</mtext><mo>·</mo><mo>·</mo><mo>·</mo><mtext>I</mtext></mrow></semantics></math> interactions in liquid HI.
    2. In which substance are the individual hydrogen bonds stronger: HF or H_2O? Explain your reasoning.
    3. For which substance will hydrogen bonding have the greater effect on the boiling point: HF or H_2O? Explain your reasoning.

Answers

1.
2.
3.
4.
5. Water is a liquid under standard conditions because of its unique ability to form four strong hydrogen bonds per molecule.
6.
7.
8.
9. As the atomic mass of the halogens increases, so does the number of electrons and the average distance of those electrons from the nucleus. Larger atoms with more electrons are more easily polarized than smaller atoms, and the increase in polarizability

with atomic number increases the strength of London dispersion forces. These intermolecular interactions are strong enough to favor the condensed states for bromine and iodine under normal conditions of temperature and pressure.

10.

11.

1. The V-shaped SO_2 molecule has a large dipole moment due to the polar S=O bonds, so dipole–dipole interactions will be most important.
2. The H–F bond is highly polar, and the fluorine atom has three lone pairs of electrons to act as hydrogen bond acceptors; hydrogen bonding will be most important.
3. Although the C=O bonds are polar, this linear molecule has no net dipole moment; hence, London dispersion forces are most important.
4. This is a symmetrical molecule that has no net dipole moment, and the Cl atoms are relatively polarizable; thus, London dispersion forces will dominate.
5. This molecule has a small dipole moment, as well as polarizable Cl atoms. In such a case, dipole–dipole interactions and London dispersion forces are often comparable in magnitude.

12.

13.

14.

15. Water has two polar O–H bonds with H atoms that can act as hydrogen bond donors, plus two lone pairs of electrons that can act as hydrogen bond acceptors, giving a net of *four* hydrogen bonds per H_2O molecule. Although methanol also has two lone pairs of electrons on oxygen that can act as hydrogen bond acceptors, it only has one O–H bond with an H atom that can act as a hydrogen bond donor. Consequently, methanol can only form *two* hydrogen bonds per molecule on average, versus four for water. Hydrogen bonding therefore has a much greater effect on the boiling point of water.

16.

17.

18.

19.

20.

21. Vigorous boiling causes more water molecule to escape into the vapor phase, but does not affect the temperature of the liquid. Vigorous boiling requires a higher energy input than does gentle simmering.

22.

23.

24.

Contributors

- Anonymous

Modified by Joshua Halpern, Scott Sinex and Scott Johnson

12.7: Intermolecular forces is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

13: Acids and Bases

Many of us are familiar with the group of chemicals called acids. But do you know what it takes for a compound to be an acid? Actually, there are several different definitions of acid that chemistry uses, and each definition is appropriate under different circumstances. Less familiar—but just as important to chemistry and ultimately to us—is the group of chemicals known as bases. Both acids and bases are important enough that we devote an entire chapter to them—their properties and their reactions.

13.1: Prelude to Acids and Bases
13.2: Arrhenius Definition of Acids and Bases
13.3: Brønsted-Lowry Definition of Acids and Bases
13.4: Water - Both an Acid and a Base
13.5: The Strengths of Acids and Bases
13.6: Buffers
13.E: Acids and Bases (Exercises)
13.S: Acids and Bases (Summary)

Template:HideTOC

13: Acids and Bases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

13.1: Prelude to Acids and Bases

One of the most concentrated acids in the body is stomach acid, which can be approximated as a 0.05 M hydrochloric acid solution. Special cells in the stomach wall secrete this acid, along with special enzymes, as part of the digestion process. In a laboratory, a 0.05 M solution of hydrochloric acid would dissolve some metals. How does the stomach survive the presence of such a reactive acid?

Actually, the stomach has several mechanisms for withstanding this chemical onslaught. First, the lining of the stomach is coated with a thin layer of mucus that contains some bicarbonate ions (HCO_3^-). These react with the hydrochloric acid to produce water, carbon dioxide, and harmless chloride ions. If any acid penetrates through the mucus, it can attack the surface layer of stomach cells, called the *gastric epithelium*. Cells in the gastric epithelium are being constantly shed, so damaged cells are quickly removed and replaced with healthy cells.



Figure 13.1.1: Stomach with acidic digestive juices. (CC BY-SA 4.0; BruceBlaus).

However, if the gastric epithelium is destroyed faster than it can be replaced, the acid may reach the wall of the stomach, resulting in ulcers. If an ulcer grows large enough, it can expose blood vessels in the stomach wall, causing bleeding. In extreme situations, the loss of blood through a severe ulcer can threaten a person's health.

Ulcers can also result from the presence of a certain bacterium—*Helicobacter pylori*—in the stomach. The mechanism for this ulcer formation is not the same as that for ulcers caused by stomach acid and is not completely understood. However, there are two main treatments for ulcers: (1) *antacids* to react chemically with excess hydrochloric acid in the stomach and (2) *antibiotics* to destroy the *H. pylori* bacteria in the stomach.

13.1: Prelude to Acids and Bases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.


- **10.0: Prelude to Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.2: Arrhenius Definition of Acids and Bases

Learning Objectives

- To recognize a compound as an Arrhenius acid or an Arrhenius base.
- To describe characteristics of acids and bases.
- To write equations of neutralization reactions.

One way to define a class of compounds is by describing the various characteristics its members have in common. In the case of the compounds known as acids, the common characteristics include a sour taste, the ability to change the color of the vegetable dye *litmus* to red, and the ability to dissolve certain metals and simultaneously produce hydrogen gas. For the compounds called bases, the common characteristics are a slippery texture, a bitter taste, and the ability to change the color of litmus to blue. Acids and bases also react with each other to form compounds generally known as salts.



Although we include their tastes among the common characteristics of acids and bases, we never advocate tasting an unknown chemical!

Chemists prefer, however, to have definitions for acids and bases in chemical terms. The Swedish chemist Svante Arrhenius developed the first chemical definitions of acids and bases in the late 1800s. Arrhenius defined an acid as a compound that increases the concentration of hydrogen ion (H^+) in aqueous solution. Many acids are simple compounds that release a hydrogen cation into solution when they dissolve. Similarly, Arrhenius defined a base as a compound that increases the concentration of hydroxide ion (OH^-) in aqueous solution. Many bases are ionic compounds that have the hydroxide ion as their anion, which is released when the base dissolves in water.

Table 13.2.1: Formulas and Names for Some Acids and Bases

Acids		**Bases**	
Formula	Name	Formula	Name
HCl(aq)	hydrochloric acid	NaOH(aq)	sodium hydroxide
HBr(aq)	hydrobromic acid	KOH(aq)	potassium hydroxide
HI(aq)	hydriodic acid	$Mg(OH)_2$(aq)	magnesium hydroxide
H_2S(aq)	hydrosulfuric acid	$Ca(OH)_2$(aq)	calcium hydroxide
$HC_2H_3O_2$(aq)	acetic acid	NH_3(aq)	ammonia
HNO_3(aq)	nitric acid	$NaHCO_3$ (aq)	sodium bicarbonate
HNO_2(aq)	nitrous acid	$CaCO_3$ (aq)	calcium carbonate
H_2SO_4(aq)	sulfuric acid		
H_2SO_3(aq)	sulfurous acid		
$HClO_3$(aq)	chloric acid		
$HClO_4$(aq)	perchloric acid		
$HClO_2$(aq)	chlorous acid		
H_3PO_4(aq)	phosphoric acid		

Acids		Bases	
H_3PO_3(aq)	phosphorous acid		
H_2CO_3(aq)	carbonic acid		

Many bases and their aqueous solutions are named using the normal rules of ionic compounds that were presented previously; that is, they are named as hydroxide compounds. For example, the base sodium hydroxide (NaOH) is both an ionic compound and an aqueous solution. However, aqueous solutions of acids have their own naming rules. The names of *binary acids* (compounds with hydrogen and one other element in their formula) are based on the root of the name of the other element preceded by the prefix *hydro-* and followed by the suffix *-ic acid*. Thus, an aqueous solution of HCl [designated "HCl(aq)"] is called hydrochloric acid, H_2S(aq) is called hydrosulfuric acid, and so forth. Acids composed of more than two elements (typically hydrogen and oxygen and some other element) have names based on the name of the other element, followed by the suffix *-ic acid* or *-ous acid*, depending on the number of oxygen atoms in the acid's formula. Other prefixes, like *per-* and *hypo-*, also appear in the names for some acids. Unfortunately, there is no strict rule for the number of oxygen atoms that are associated with the *-ic acid* suffix; the names of these acids are best memorized. Table 13.2.1 lists some acids and bases and their names. Note that acids have hydrogen written first, as if it were the cation, while most bases have the negative hydroxide ion, if it appears in the formula, written last.

> *The name oxygen comes from the Latin meaning "acid producer" because its discoverer, Antoine Lavoisier, thought it was the essential element in acids. Lavoisier was wrong, but it is too late to change the name now.*

Example 13.2.1

Name each substance.

a. HF(aq)
b. $Sr(OH)_2$(aq)

Solution

a. This acid has only two elements in its formula, so its name includes the *hydro-* prefix. The stem of the other element's name, fluorine, is *fluor*, and we must also include the *-ic acid* ending. Its name is hydrofluoric acid.
b. This base is named as an ionic compound between the strontium ion and the hydroxide ion: strontium hydroxide.

Exercise 13.2.1

Name each substance.

a. H_2Se(aq)
b. $Ba(OH)_2$(aq)

Answer

a. hydroselenic acid

b. barium hydroxide

Notice that one base listed in Table 13.2.1—ammonia—does not have hydroxide as part of its formula. How does this compound increase the amount of hydroxide ion in aqueous solution? Instead of dissociating into hydroxide ions, ammonia molecules react with water molecules by taking a hydrogen ion from the water molecule to produce an ammonium ion and a hydroxide ion:

$$NH_{3(aq)} + H_2O_{(\ell)} \rightarrow NH^+_{4(aq)} + OH^-_{(aq)} \tag{13.2.1}$$

Because this reaction of ammonia with water causes an increase in the concentration of hydroxide ions in solution, ammonia satisfies the Arrhenius definition of a base. Many other nitrogen-containing compounds are bases because they too react with water to produce hydroxide ions in aqueous solution.

Neutralization

As we noted previously, acids and bases react chemically with each other to form *salts*. A salt is a general chemical term for any ionic compound formed from an acid and a base. In reactions where the acid is a hydrogen ion containing compound and the base is a hydroxide ion containing compound, water is also a product. The general reaction is as follows:

$$\text{acid + base} \rightarrow \text{water + salt}$$

The reaction of acid and base to make water and a salt is called **neutralization**. Like any chemical equation, a neutralization chemical equation must be properly balanced. For example, the neutralization reaction between sodium hydroxide and hydrochloric acid is as follows:

$$NaOH(aq) + HCl_{(aq)} \rightarrow NaCl_{(aq)} + H_2O_{(\ell)} \tag{13.2.2}$$

with coefficients all understood to be one. The neutralization reaction between sodium hydroxide and sulfuric acid is as follows:

$$2NaOH_{(aq)} + H_2SO_{4(aq)} \rightarrow Na_2SO_{4(aq)} + 2H_2O_{(\ell)} \tag{13.2.3}$$

Once a neutralization reaction is properly balanced, we can use it to perform stoichiometry calculations, such as the ones we practiced earlier.

There are a number of examples of acid-base chemistry in everyday life. One example is the use of baking soda, or sodium bicarbonate in baking. $NaHCO_3$ is a base. When it reacts with an acid such as lemon juice, buttermilk, or sour cream in a batter, bubbles of carbon dioxide gas are formed from decomposition of the resulting carbonic acid, and the batter "rises." Baking powder is a combination of sodium bicarbonate, and one or more acid salts that react when the two chemicals come in contact with water in the batter.

$$HCO_3^-(aq) + H^+(aq) \rightarrow H_2CO_3(aq) \tag{13.2.4}$$

$$H_2CO_3(aq) \rightarrow CO_2(g) + H_2O(l)$$

Example 13.2.2

Nitric acid [HNO_3(aq)] can be neutralized by calcium hydroxide [$Ca(OH)_2$(aq)].


a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 16.8 g of HNO_3 is present initially. How many grams of $Ca(OH)_2$ are needed to neutralize that much HNO_3?
c. In a second reaction, 805 mL of 0.672 M $Ca(OH)_2$ is present initially. What volume of 0.432 M HNO_3 solution is necessary to neutralize the $Ca(OH)_2$ solution?

Solution

a. Because there are two OH^- ions in the formula for $Ca(OH)_2$, we need two moles of HNO_3 to provide H^+ ions. The balanced chemical equation is as follows: $Ca(OH)_2(aq) + 2HNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + 2H_2O(\ell)$

   The salt formed is calcium nitrate.

b. This calculation is much like the calculations we did in Chapter 6. First we convert the mass of HNO_3 to moles using its molar mass of 1.01 + 14.01 + 3(16.00) = 63.02 g/mol; then we use the balanced chemical equation to determine the related number of moles of $Ca(OH)_2$ needed to neutralize it; and then we convert that number of moles of $Ca(OH)_2$ to the mass of $Ca(OH)_2$ using its molar mass of 40.08 + 2(1.01) + 2(16.00) = 74.10 g/mol.



$$16.8\ \text{g HNO}_3 \times \frac{1\ \text{mol HNO}_3}{63.02\ \text{g HNO}_3} \times \frac{1\ \text{mol Ca(OH)}_2}{2\ \text{mol HNO}_3} \times \frac{74.10\ \text{g Ca(OH)}_2}{1\ \text{mol Ca(OH)}_2} = 9.88\ \text{g Ca(OH)}_2\ \text{needed}$$

c. Having concentration information allows us to employ the skills we developed in Chapter 9. We have two alternative solutions: the multi-step process and the combined one-line process (found at the bottom).

First, we use the concentration and volume data to determine the number of moles of $Ca(OH)_2$ present. Recognizing that 805 mL = 0.805 L,

$$0.672\ \text{M Ca(OH)}_2 = \frac{\text{mol Ca(OH)}_2}{0.805\ \text{L soln}}$$

$$0.672\ \text{M CaOH)}_2 \times (0.805\ \text{L soln}) = \text{mol Ca(OH)}_2 = 0.541\ \text{mol Ca(OH)}_2$$


We combine this information with the proper ratio from the balanced chemical equation to determine the number of moles of HNO_3 needed:

$$0.541\ \text{mol Ca(OH)}_2 \times \frac{2\ \text{mol HNO}_3}{1\ \text{mol Ca(OH)}_2} = 1.08\ \text{mol HNO}_3$$

Now, using the definition of molarity one more time, we determine the volume of acid solution needed:

$$0.432\ \text{M HNO}_3 = \frac{1.08\ \text{mol HNO}_3}{\text{volume of HNO}_3}$$

$$\text{volume of HNO}_3 = \frac{1.08\ \text{mol HNO}_3}{0.432\ \text{M HNO}_3} = 2.50\ \text{L} = 2.50 \times 10^3\ \text{mL HNO}_3$$



$$0.805\ \text{L Ca(OH)}_2\ \text{soln} \times \frac{0.672\ \text{mol Ca(OH)}_2}{1\ \text{L Ca(OH)}_2\ \text{soln}} \times \frac{2\ \text{mol HNO}_3}{1\ \text{mol Ca(OH)}_2} \times \frac{1\ \text{L HNO}_3\ \text{soln}}{0.432\ \text{mol HNO}_3} = 2.50\ \text{L HNO}_3\ \text{soln needed}$$

? Exercise 13.2.2

Hydrocyanic acid [HCN(aq)] can be neutralized by potassium hydroxide [KOH(aq)].

a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 37.5 g of HCN is present initially. How many grams of KOH are needed to neutralize that much HCN?
c. In a second reaction, 43.0 mL of 0.0663 M KOH is present initially. What volume of 0.107 M HCN solution is necessary to neutralize the KOH solution?

Answer

a. KOH(aq) + HCN(aq) → KCN(aq) + $H_2O(\ell)$ KCN is the salt.

b. 77.8 g

c. 0.0266 L or 26.6 mL

Hydrocyanic acid (HCN) is one exception to the acid-naming rules that specify using the prefix *hydro-* for binary acids (acids composed of hydrogen and only one other element).

Stomach Antacids

Our stomachs contain a solution of roughly 0.03 *M* HCl, which helps us digest the food we eat. The burning sensation associated with heartburn is a result of the acid of the stomach leaking through the muscular valve at the top of the stomach into the lower reaches of the esophagus. The lining of the esophagus is not protected from the corrosive effects of stomach acid the way the lining of the stomach is, and the results can be very painful. When we have heartburn, it feels better if we reduce the excess acid in the esophagus by taking an antacid. As you may have guessed, antacids are bases. One of the most common antacids is calcium carbonate, $CaCO_3$. The reaction,

$$CaCO_3(s) + 2HCl(aq) \rightleftharpoons CaCl_2(aq) + H_2O(l) + CO_2(g)$$

not only neutralizes stomach acid, it also produces $CO_2(g)$, which may result in a satisfying belch.

Milk of Magnesia is a suspension of the sparingly soluble base magnesium hydroxide, $Mg(OH)_2$. It works according to the reaction:

$$Mg(OH)_2(s) \rightleftharpoons Mg^{2+}(aq) + 2OH^-(aq)$$

The hydroxide ions generated in this equilibrium then go on to react with the hydronium ions from the stomach acid, so that :

$$H_3O^+ + OH^- \rightleftharpoons 2H_2O(l)$$

This reaction does not produce carbon dioxide, but magnesium-containing antacids can have a laxative effect. Several antacids have aluminum hydroxide, $Al(OH)_3$, as an active ingredient. The aluminum hydroxide tends to cause constipation, and some antacids use aluminum hydroxide in concert with magnesium hydroxide to balance the side effects of the two substances.

Example 13.2.3

Assume that the stomach of someone suffering from acid indigestion contains 75 mL of 0.20 M HCl. How many Tums tablets are required to neutralize 90% of the stomach acid, if each tablet contains 500 mg of $CaCO_3$? (Neutralizing all of the stomach acid is not desirable because that would completely shut down digestion.)

Solution

A. Write the balanced chemical equation for the reaction and then decide whether the reaction will go to completion.
B. Calculate the number of moles of acid present. Multiply the number of moles by the percentage to obtain the quantity of acid that must be neutralized. Using mole ratios, calculate the number of moles of base required to neutralize the acid.
C. Calculate the number of moles of base contained in one tablet by dividing the mass of base by the corresponding molar mass. Calculate the number of tablets required by dividing the moles of base by the moles contained in one tablet.

A. We first write the balanced chemical equation for the reaction:

$$2HCl(aq) + CaCO_3(s) \rightarrow CaCl_2(aq) + H_2CO_3(aq)$$

Each carbonate ion can react with 2 mol of H^+ to produce H_2CO_3, which rapidly decomposes to H_2O and CO_2. Because HCl is a strong acid and CO_3^{2-} is a weak base, the reaction will go to completion.

B. Next we need to determine the number of moles of HCl present:

$$75 \ \cancel{mL}\left(\frac{1 \ \cancel{L}}{1000 \ \cancel{mL}}\right)\left(\frac{0.20 \ mol \ HCl}{\cancel{L}}\right) = 0.015 \ mol \ HCl$$

Because we want to neutralize only 90% of the acid present, we multiply the number of moles of HCl by 0.90:

$$(0.015 \ mol \ HCl)(0.90) = 0.014 \ mol \ HCl$$

We know from the stoichiometry of the reaction that each mole of $CaCO_3$ reacts with 2 mol of HCl, so we need

$$moles \ CaCO_3 = 0.014 \ \cancel{mol \ HCl}\left(\frac{1 \ mol \ CaCO_3}{2 \ \cancel{mol \ HCl}}\right) = 0.0070 \ mol \ CaCO_3$$

C. Each Tums tablet contains

$$\left(\frac{500\ \cancel{mg\,CaCO_3}}{1\ Tums\ tablet}\right)\left(\frac{1\ \cancel{g}}{1000\ \cancel{mg\,CaCO_3}}\right)\left(\frac{1\ mol\ CaCO_3}{100.1\ \cancel{g}}\right) = 0.00500\ mol\ CaCO_3$$

Thus we need $\frac{0.0070\ \cancel{mol\ CaCO_3}}{0.00500\ \cancel{mol\ CaCO_3}} = 1.4$ Tums tablets.

cise

Assume that as a result of overeating, a person's stomach contains 300 mL of 0.25 M HCl. How many Rolaids tablets must be consumed to neutralize 95% of the acid, if each tablet contains 400 mg of $NaAl(OH)_2CO_3$? The neutralization reaction can be written as follows:

$$NaAl(OH)_2CO_3(s) + 4HCl(aq) \rightarrow AlCl_3(aq) + NaCl(aq) + CO_2(g) + 3H_2O(l)$$

Answer

6.4 tablets

Key Takeaway

- An Arrhenius acid increases the H^+ ion concentration in water, while an Arrhenius base increases the OH^- ion concentration in water.

13.2: Arrhenius Definition of Acids and Bases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.1: Arrhenius Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.3: Brønsted-Lowry Definition of Acids and Bases

Learning Objectives

1. Recognize a compound as a Brønsted-Lowry acid or a Brønsted-Lowry base.
2. Illustrate the proton transfer process that defines a Brønsted-Lowry acid-base reaction.


Ammonia (NH_3) increases the hydroxide ion concentration in aqueous solution by reacting with water rather than releasing hydroxide ions directly. In fact, the Arrhenius definitions of an acid and a base focus on hydrogen ions and hydroxide ions. Are there more fundamental definitions for acids and bases?

In 1923, the Danish scientist Johannes Brønsted and the English scientist Thomas Lowry independently proposed new definitions for acids and bases. Rather than considering both hydrogen and hydroxide ions, they focused on the hydrogen ion only. A Brønsted-Lowry acid is a compound that supplies a hydrogen ion in a reaction. A Brønsted-Lowry base, conversely, is a compound that accepts a hydrogen ion in a reaction. Thus, the Brønsted-Lowry definitions of an acid and a base focus on the movement of hydrogen ions in a reaction, rather than on the production of hydrogen ions and hydroxide ions in an aqueous solution.

Let us use the reaction of ammonia in water to demonstrate the Brønsted-Lowry definitions of an acid and a base. Ammonia and water molecules are reactants, while the ammonium ion and the hydroxide ion are products:

$$NH_3(aq) + H_2O(\ell) \rightarrow NH_4^+(aq) + OH^-(aq) \tag{13.3.1}$$

What has happened in this reaction is that the original water molecule has donated a hydrogen ion to the original ammonia molecule, which in turn has accepted the hydrogen ion. We can illustrate this as follows:



Because the water molecule donates a hydrogen ion to the ammonia, it is the Brønsted-Lowry acid, while the ammonia molecule—which accepts the hydrogen ion—is the Brønsted-Lowry base. Thus, ammonia acts as a base in both the Arrhenius sense and the Brønsted-Lowry sense.


Is an Arrhenius acid like hydrochloric acid still an acid in the Brønsted-Lowry sense? Yes, but it requires us to understand what really happens when HCl is dissolved in water. Recall that the hydrogen *atom* is a single proton surrounded by a single electron. To make the hydrogen *ion*, we remove the electron, leaving a bare proton. Do we *really* have bare protons floating around in aqueous solution? No, we do not. What really happens is that the H^+ ion attaches itself to H_2O to make H_3O^+, which is called the *hydronium ion*. For most purposes, H^+ and H_3O^+ represent the same species, but writing H_3O^+ instead of H^+ shows that we understand that there are no bare protons floating around in solution. Rather, these protons are actually attached to solvent molecules.

> A proton in aqueous solution may be surrounded by more than one water molecule, leading to formulas like $H_5O_2^+$ or $H_9O_4^+$ rather than H_3O^+. It is simpler, however, to use H_3O^+.

With this in mind, how do we define HCl as an acid in the Brønsted-Lowry sense? Consider what happens when HCl is dissolved in H_2O:

$$HCl(g) + H_2O(\ell) \rightarrow H_3O^+(aq) + Cl^-(aq) \tag{13.3.2}$$

We can depict this process using Lewis electron dot diagrams:



Now we see that a hydrogen ion is transferred from the HCl molecule to the H_2O molecule to make chloride ions and hydronium ions. As the hydrogen ion donor, **HCl acts as a Brønsted-Lowry acid**; as a hydrogen ion acceptor, **H_2O is a Brønsted-Lowry base**. So HCl is an acid not just in the Arrhenius sense but also in the Brønsted-Lowry sense. Moreover, by the Brønsted-Lowry definitions, H_2O is a base in the formation of aqueous HCl. So the Brønsted-Lowry definitions of an acid and a base classify the dissolving of HCl in water as a reaction between an acid and a base—although the Arrhenius definition would not have labeled H_2O a base in this circumstance.

> All Arrhenius acids and bases are Brønsted-Lowry acids and bases as well. However, not all Brønsted-Lowry acids and bases are Arrhenius acids and bases.

✔ Example 13.3.1

Aniline ($C_6H_5NH_2$) is slightly soluble in water. It has a nitrogen atom that can accept a hydrogen ion from a water molecule just like the nitrogen atom in ammonia does. Write the chemical equation for this reaction and identify the Brønsted-Lowry acid and base.

Solution

$C_6H_5NH_2$ and H_2O are the reactants. When $C_6H_5NH_2$ accepts a proton from H_2O, it gains an extra H and a positive charge and leaves an OH^- ion behind. The reaction is as follows:

$$C_6H_5NH_2(aq) + H_2O(\ell) \rightarrow C_6H_5NH_3^+(aq) + OH^-(aq)$$

Because $C_6H_5NH_2$ accepts a proton, it is the Brønsted-Lowry base. The H_2O molecule, because it donates a proton, is the Brønsted-Lowry acid.

? Exercise 13.3.1

Caffeine ($C_8H_{10}N_4O_2$) is a stimulant found in coffees and teas. When dissolved in water, it can accept a proton from a water molecule. Write the chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer

$C_8H_{10}N_4O_2(aq) + H_2O(\ell) \rightarrow C_8H_{11}N_4O_2^+(aq) + OH^-(aq)$

B-L base B-L acid

The Brønsted-Lowry definitions of an acid and a base can be applied to chemical reactions that occur in solvents other than water. The following example illustrates.

✔ Example 13.3.2

Sodium amide ($NaNH_2$) dissolves in methanol (CH_3OH) and separates into sodium ions and amide ions (NH_2^-). The amide ions react with methanol to make ammonia and the methoxide ion (CH_3O^-). Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Solution

The equation for the reaction is between NH_2^- and CH_3OH to make NH_3 and CH_3O^- is as follows:

$$NH_2^-(solv) + CH_3OH(\ell) \rightarrow NH_3(solv) + CH_3O^-(solv)$$

The label *(solv)* indicates that the species are dissolved in some solvent, in contrast to *(aq)*, which specifies an aqueous (H_2O) solution. In this reaction, we see that the NH_2^- ion accepts a proton from a CH_3OH molecule to make an NH_3 molecule. Thus, as the proton acceptor, NH_2^- is the Brønsted-Lowry base. As the proton donor, CH_3OH is the Brønsted-Lowry acid.

? Exercise 13.3.2

Pyridinium chloride (C_5H_5NHCl) dissolves in ethanol (C_2H_5OH) and separates into pyridinium ions ($C_5H_5NH^+$) and chloride ions. The pyridinium ion can transfer a hydrogen ion to a solvent molecule. Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer

$C_5H_5NH^+(solv) + C_2H_5OH(\ell) \rightarrow C_5H_5N(solv) + C_2H_5OH_2^+(solv)$

B-L acid B-L base

Application in Everyday Life

Many people like to put lemon juice or vinegar, both of which are acids, on cooked fish (Figure 13.3.1). It turns out that fish have volatile amines (bases) in their systems, which are neutralized by the acids to yield involatile ammonium salts. This reduces the odor of the fish, and also adds a "sour" taste that we seem to enjoy.



Figure 13.3.1: A neutralization reaction takes place between citric acid or acetic acid (proton donors) in lemons or vinegar and putrescine (proton acceptor) in the flesh of fish. (CC BY 4.0; OpenStax)

Pickling is a method used to preserve vegetables using a naturally produced acidic environment. The vegetable, such as a cucumber, is placed in a sealed jar submerged in a brine solution. The brine solution favors the growth of beneficial bacteria and suppresses the growth of harmful bacteria. The beneficial bacteria feed on starches in the cucumber and produce lactic acid as a waste product in a process called fermentation. The lactic acid eventually increases the acidity of the brine to a level that kills any harmful bacteria, which require a basic environment. Without the harmful bacteria consuming the cucumbers they are able to last much longer than if they were unprotected. A byproduct of the pickling process changes the flavor of the vegetables with the acid making them taste sour.

To Your Health: Brønsted-Lowry Acid-Base Reactions in Pharmaceuticals

There are many interesting applications of Brønsted-Lowry acid-base reactions in the pharmaceutical industry. For example, drugs often need to be water soluble for maximum effectiveness. However, many complex organic compounds are not soluble or are only slightly soluble in water. Fortunately, those drugs that contain proton-accepting nitrogen atoms (and there are a lot of them) can be reacted with dilute hydrochloric acid [HCl(aq)]. The nitrogen atoms—acting as Brønsted-Lowry bases—accept the hydrogen ions from the acid to make an ion, which is usually much more soluble in water. The modified drug molecules can then be isolated as chloride salts:

\[\mathrm{RN(sl\: aq) + H^+(aq) \rightarrow RNH^+(aq) \xrightarrow{Cl^-(aq)} RNHCl(s)} \label{Eq3} \]

where RN represents some organic compound containing nitrogen. The label *(sl aq)* means “slightly aqueous,” indicating that the compound RN is only slightly soluble. Drugs that are modified in this way are called *hydrochloride salts*. Examples include the powerful painkiller codeine, which is commonly administered as codeine hydrochloride. Acids other than hydrochloric acid are also used. Hydrobromic acid, for example, gives *hydrobromide salts*. Dextromethorphan, an ingredient in many cough medicines, is dispensed as dextromethorphan hydrobromide. The accompanying figure shows another medication (lidocaine) as a hydrochloride salt.



Figure used with permission from Wikipedia.

Conjugate Acid-Base Pairs

According to the Bronsted-Lowry theory of acids and bases, an acid is a proton donor and a base is a proton acceptor. Once an acid has given up a proton, the part that remains is called the acid's **conjugate base**. This species is a base because it can accept a proton (to re-form the acid). The conjugate base of HF (first example below) is fluoride ion, F^-.

$$\text{Acid} = H^+ + \text{Conjugate base of Acid}^-$$

$$HF \rightleftharpoons H^+ + F^-$$

$$H_2O \rightleftharpoons H^+ + OH^-$$

$$NH_4^+ \rightleftharpoons H^+ + NH_3$$

Similarly, the part of the base that remains after a base accepts a proton is called the base's **conjugate acid**. This species is an acid because it can give up a proton (and thus re-form the base). The conjugate acid of fluoride ion, F^- (first example below) is HF.

$$H^+ + \text{Base} = \text{Conjugate acid of Base}^+$$

$$H^+ + F^- \rightleftharpoons HF$$

$$H^+ + OH^- \rightleftharpoons H_2O$$

$$H^+ + H_2O \rightleftharpoons H_3O^+$$

$$H^+ + NH_3 \rightleftharpoons NH_4^+$$

To summarize, the conjugate base of HF is fluoride ion, F^-, and the conjugate acid of fluoride ion, F^-, is HF. The HF/F- pair is referred to as a conjugate acid-base pair. The difference in the formulas of a conjugate acid-base pair (example: HF and F^-) is H^+. The table below lists conjugate acid-base pairs for your reference so that you can figure out the strategy of identifying them. For any given acid or base, you should be able to give its conjugate base or conjugate acid. The formula of an acid's conjugate base is generated by **removing a proton** (H^+) from the acid formula. The formula of the base's conjugate acid is formed by **adding a proton** (H^+) to the formula of the base.

Table 13.3.1. Conjugate acid-base pairs.

Conjugate Acid	Conjugate Base
H_3O^+	H_2O
H_2O	OH^-
H_2SO_4	HSO_4^-
HSO_4^-	SO_4^{2-}
NH_4^+	NH_3

Conjugate Acid	Conjugate Base
NH_3	NH_2^-
CH_3COOH	CH_3COO^-
$CH_3NH_3^+$	CH_3NH_2

Example 13.3.3 : Conjugate Pairs

Write the formula of the conjugate base of (a) HCl and (b) HCO_3^-.

Write the formula of the conjugate acid of (c) CH_3NH_2 and (d) OH^-.

Solution:

A conjugate base is formed by removing a proton (H^+). A conjugate acid is formed by accepting a proton (H^+).

a. After HCl donates a proton, a Cl^- ion is produced, and so Cl^- is the conjugate base.
b. After hydrogen carbonate ion, HCO_3^-, donates a proton, its conjugate base, CO_3^{2-} is produced.
c. After accepting a proton (H^+), CH_3NH_2 is converted to $CH_3NH_3^+$, its conjugate acid.
d. After accepting a proton (H^+), OH^- is converted to H_2O, its conjugate acid.

Exercise : Conjugate Pairs

Write the formula of the conjugate base of (a) HNO_2 and (b) H_2CO_3.

Write the formula of the conjugate acid of (c) C_6H5NH_2 and (d) HCO_3^-.

Answer


a. NO_2^- is the conjugate base of HNO_2.

b. HCO_3^- is the conjugate base of H_2CO_3

c. $C_6H5NH_3^+$ is the conjugate acid of C_6H5NH_2.


d. H_2CO_3 is the conjugate acid of HCO_3^-

In the reaction of ammonia with water to give ammonium ions and hydroxide ions, ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The **conjugate acid–base pairs** for this reaction are NH_4^+/NH_3 and H_2O/OH^-. This means that the conjugate acid of the base NH_3 is NH_4^+ while the conjugate base of the acid NH_4^+ is NH_3. Similarly, the conjugate base of the acid H_2O is OH^-, and the conjugate acid of the base OH^- is H_2O.




This figure has three parts in two rows. In the first row, two diagrams of acid-base pairs are shown. On the left, a space filling model of H subscript 2 O is shown with a red O atom at the center and two smaller white H atoms attached in a bent shape. Above this model is the label “H subscript 2 O (acid)” in purple. An arrow points right, which is labeled “Remove H superscript plus.” To the right is another space filling model with a single red O atom to which a single smaller white H atom is attached. The label in purple above this model reads, “O H superscript negative (conjugate base).” Above both of these red and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” To the right is a space filling model with a central blue N atom to which three smaller white H atoms are attached in a triangular pyramid arrangement. A label in green above reads “N H subscript 3 (base).” An arrow labeled “Add H superscript plus” points right. To the right of the arrow is another space filling model with a blue central N atom and four smaller white H atoms in a tetrahedral arrangement. The green label above reads “N H subscript 3 superscript plus (conjugate acid).” Above both of these blue and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” The second row of the figure shows the chemical reaction, H subscript 2 O (l) is shown in purple, and is labeled below in purple as “acid,” plus N H subscript 3 (a q) in green, labeled below in green as “base,” followed by a double sided arrow arrow and O H superscript negative (a q) in purple, labeled in purple as “conjugate base,” plus N H subscript 4 superscript plus (a q)” in green, which is labeled in green as “conjugate acid.” The acid on the left side of the equation is connected to the conjugate base on the right with a purple line. Similarly, the base on the left is connected to the conjugate acid on the right side.

In the **forward** reaction, the parent acid is H_2O and and the parent base is NH_3 (shown in the illustration below). The acid H_2O loses a proton (H^+) to form its conjugate base OH^-. The base NH_3 gains a proton, to produce its conjugate acid NH_4^+. In the **reverse** reaction, the acid NH_4^+ loses a proton (H^+) to form its conjugate base NH_3. The base OH^- gains a proton, to produce its conjugate acid H_2O.



When hydrogen fluoride (HF) dissolves in water and ionizes, protons are transferred from hydrogen fluoride (parent acid) molecules to water (parent base) molecules, yielding hydronium ions (conjugate acid of water) and fluoride ions (conjugate base of HF):

$$HF + H_2O \rightleftharpoons H_3O^+ + F^-$$



https://chem.libretexts.org/@go/page/218564

Example 13.3.3

Identify the conjugate acid-base pairs in this equilibrium.


$$CH_3CO_2H + H_2O \rightleftharpoons H_3O^+ + CH_3CO_2^-$$

Solution

Similarly, in the reaction of acetic acid with water, acetic acid **donates** a proton to water, which acts as the base. In the reverse reaction, H_3O^+ is the acid that donates a proton to the acetate ion, which acts as the base.

Once again, we have two conjugate acid–base pairs:

- the parent acid and its conjugate base ($CH_3CO_2H/CH_3CO_2^-$) and
- the parent base and its conjugate acid (H_3O^+/H_2O).



Example 13.3.4

Identify the conjugate acid-base pairs in this equilibrium.

$$(CH_3)_3N + H_2O \rightleftharpoons (CH_3)_3NH^+ + OH^-$$

Solution

One pair is H_2O and OH^-, where H_2O has one more H^+ and is the conjugate acid, while OH^- has one less H^+ and is the conjugate base.

The other pair consists of $(CH_3)_3N$ and $(CH_3)_3NH^+$, where $(CH_3)_3NH^+$ is the conjugate acid (it has an additional proton) and $(CH_3)_3N$ is the conjugate base.

Exercise 13.3.4

Identify the conjugate acid-base pairs in this equilibrium.

$$NH_2^- + H_2O \rightleftharpoons NH_3 + OH^-$$

Answer:

H_2O (acid) and OH^- (base); NH_2^- (base) and NH_3 (acid)

The use of conjugate acid-base pairs allows us to make a very simple statement about relative strengths of acids and bases. *The stronger an acid, the weaker its conjugate base*, and, conversely, *the stronger a base, the weaker its conjugate acid.*

Key Takeaways

- A Brønsted-Lowry acid is a proton donor, and a Brønsted-Lowry base is a proton acceptor.
- Brønsted-Lowry acid-base reactions are essentially proton transfer reactions.

13.3: Brønsted-Lowry Definition of Acids and Bases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.2: Brønsted-Lowry Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.4: Water - Both an Acid and a Base

Learning Objectives

- To write chemical equations for water acting as an acid and as a base.

Water (H_2O) is an interesting compound in many respects. Here, we will consider its ability to behave as an acid or a base.

In some circumstances, a water molecule will accept a proton and thus act as a **Brønsted-Lowry base**. We saw an example in the dissolving of HCl in H_2O:

$$\mathrm{HCl} + \mathrm{H_2O}(\ell) \rightarrow \mathrm{H_3O^+}(\mathrm{aq}) + \mathrm{Cl^-}(\mathrm{aq}) \tag{13.4.1}$$

In other circumstances, a water molecule can donate a proton and thus act as a **Brønsted-Lowry acid**. For example, in the presence of the amide ion (see Example 4 in Section 10.2), a water molecule donates a proton, making ammonia as a product:

$$H_2O(\ell) + NH_2^-(aq) \rightarrow OH^-(aq) + NH_3(aq) \tag{13.4.2}$$

In this case, NH_2^- is a Brønsted-Lowry base (the proton acceptor).

So, depending on the circumstances, H_2O can act as either a Brønsted-Lowry acid or a Brønsted-Lowry base. Water is not the only substance that can react as an acid in some cases or a base in others, but it is certainly the most common example—and the most important one. A substance that can either donate or accept a proton, depending on the circumstances, is called an **amphiprotic** compound.

A water molecule can act as an acid or a base even in a sample of pure water. About 6 in every 100 million (6 in 10^8) water molecules undergo the following reaction:

$$H_2O(\ell) + H_2O(\ell) \rightarrow H_3O^+(aq) + OH^-(aq) \tag{13.4.3}$$

This process is called the **autoionization of water** (Figure 13.4.1) and occurs in every sample of water, whether it is pure or part of a solution. Autoionization occurs to some extent in any amphiprotic liquid. (For comparison, liquid ammonia undergoes autoionization as well, but only about 1 molecule in a million billion (1 in 10^{15}) reacts with another ammonia molecule.)

Note

It is rare to truly have pure water. Water exposed to air will usually be slightly acidic because dissolved carbon dioxide gas, or carbonic acid, decreases the pH slightly below 7. Alternatively, dissolved minerals, like calcium carbonate (limestone), can make water slightly basic.




Figure 13.4.1 Autoionization. A small fraction of water molecules—approximately 6 in 100 million—ionize spontaneously into hydronium ions and hydroxide ions. This picture necessarily overrepresents the amount of autoionization that really occurs in pure water.

Example 13.4.1

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $H_2O(\ell) + NO_2^-(aq) \rightarrow HNO_2(aq) + OH^-(aq)$
2. $HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)$

Solution

1. In this reaction, the water molecule donates a proton to the NO_2^- ion, making $OH^-(aq)$. As the proton donor, H_2O acts as a Brønsted-Lowry acid.
2. In this reaction, the water molecule accepts a proton from $HC_2H_3O_2$, becoming $H_3O^+(aq)$. As the proton acceptor, H_2O is a Brønsted-Lowry base.

Exercise 13.4.2

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $HCOOH(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + HCOO^-(aq)$
2. $H_2O(\ell) + PO_4^{3-}(aq) \rightarrow OH^-(aq) + HPO_4^{2-}(aq)$

Answer

1. H_2O acts as the proton acceptor (Brønsted-Lowry base)
2. H_2O acts as the proton donor (Brønsted-Lowry acid)

Key Takeaway

- Water molecules can act as both an acid and a base, depending on the conditions.

13.4: Water - Both an Acid and a Base is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.3: Water - Both an Acid and a Base** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.5: The Strengths of Acids and Bases

Learning Objectives

- Describe the difference between strong and weak acids and bases.
- Describe how a chemical reaction reaches chemical equilibrium.
- Define the pH scale and use it to describe acids and bases.

Acids and bases do not all demonstrate the same degree of chemical activity in solution. Different acids and bases have different strengths.

Strong and Weak Acids

Let us consider the strengths of acids first. A small number of acids ionize completely in aqueous solution. For example, when HCl dissolves in water, every molecule of HCl separates into a hydronium ion and a chloride ion:

$$\mathrm{HCl(g) + H_2O(l) \xrightarrow{\sim 100\%} H_3O^+(aq) + Cl^-(aq)} \tag{13.5.1}$$

HCl(aq) is one example of a **strong acid**, which is a compound that is essentially **100% ionized** in aqueous solution. There are very few strong acids. The important ones are listed in Table 13.5.1.

Table 13.5.1: Strong Acids and Bases (All in Aqueous Solution)

Acids	Bases
HCl	LiOH
HBr	NaOH
HI	KOH
HNO_3	$Mg(OH)_2$
H_2SO_4	$Ca(OH)_2$
$HClO_4$	

By analogy, a strong base is a compound that is essentially 100% ionized in aqueous solution. As with acids, there are only a few strong bases, which are also listed in Table 13.5.1.

If an acid is **not listed** in Table 13.5.1, it is likely a **weak acid**, which is a compound that is not 100% ionized in aqueous solution. Similarly, a weak base is a compound that is not 100% ionized in aqueous solution. For example, acetic acid ($HC_2H_3O_2$) is a weak acid. The ionization reaction for acetic acid is as follows:

$$\mathrm{HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)} \tag{13.5.2}$$

Depending on the concentration of $HC_2H_3O_2$, the ionization reaction may occur only for 1%–5% of the acetic acid molecules.

Looking Closer: Household Acids and Bases

Many household products are acids or bases. For example, the owner of a swimming pool may use muriatic acid to clean the pool. Muriatic acid is another name for hydrochloric acid [$HCl(aq)$]. Vinegar has already been mentioned as a dilute solution of acetic acid [$HC_2H_3O_2(aq)$]. In a medicine chest, one may find a bottle of vitamin C tablets; the chemical name of vitamin C is ascorbic acid ($HC_6H_7O_6$).

One of the more familiar household bases is ammonia (NH_3), which is found in numerous cleaning products. As we mentioned previously, ammonia is a base because it increases the hydroxide ion concentration by reacting with water:

$$\mathrm{NH_3(aq) + H_2O(\ell) \rightarrow NH_4^+(aq) + OH^-(aq)} \tag{13.5.3}$$

Many soaps are also slightly basic because they contain compounds that act as Brønsted-Lowry bases, accepting protons from water and forming excess hydroxide ions. This is one reason that soap solutions are slippery.



Figure 13.5.1: (left) Bottles of alkaline drain cleaners containing sodium hydroxide can dissolve greases and hair. (right) Chemical burns caused by sodium hydroxide solution photographed 44 hours after exposure. Figures used with permission from Wikipedia.

Perhaps the most dangerous household chemical is the lye-based drain cleaner. Lye is a common name for sodium hydroxide, although it is also used as a synonym for potassium hydroxide. Lye is an extremely caustic chemical that can react with grease, hair, food particles, and other substances that may build up and form a clog in a pipe. Unfortunately, lye can also attack tissues and other substances in our bodies. Thus, when we use lye-based drain cleaners, we must be very careful not to touch any of the solid drain cleaner or spill the water it was poured into. Safer, nonlye drain cleaners use peroxide compounds to react on the materials in the clog and clear the drain.

Chemical Equilibrium in Weak Acids and Bases

The behavior of **weak acids and bases** illustrates a key concept in chemistry. Does the chemical reaction describing the ionization of a weak acid or base just stop when the acid or base is done ionizing? Actually, no. Rather, the reverse process—the reformation of the molecular form of the acid or base—occurs, ultimately at the same rate as the ionization process. For example, the ionization of the weak acid $\mathrm{HC_2H_3O_2(aq)}$ is as follows:

$$\mathrm{HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)} \tag{13.5.4}$$

The reverse process also begins to occur:

$$\mathrm{H_3O^+(aq) + C_2H_3O_2^-(aq) \rightarrow HC_2H_3O_2(aq) + H_2O(\ell)} \tag{13.5.5}$$

Eventually, there is a balance between the two opposing processes, and no additional change occurs. The chemical reaction is better represented at this point with a double arrow:

$$\mathrm{HC_2H_3O_2(aq) + H_2O(\ell) \rightleftharpoons H_3O^+(aq) + C_2H_3O_2^-(aq)} \tag{13.5.6}$$

The $\rightleftharpoons$ implies that **both the forward and reverse reactions** are occurring, and their effects cancel each other out. A process at this point is considered to be at **chemical equilibrium** (or equilibrium). It is important to note that the processes do not stop. They balance out each other so that there is no further net change; that is, chemical equilibrium is a *dynamic equilibrium*.

Example 13.5.1: Partial Ionization

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HNO_2(aq)$
b. $C_5H_5N(aq)$

Solution

a. $HNO_2(aq) + H_2O(\ell) \leftrightarrows NO_2^-(aq) + H_3O^+(aq)$
b. $C_5H_5N(aq) + H_2O(\ell) \leftrightarrows C_5H_5NH^+(aq) + OH^-(aq)$

? Exercise 13.5.1

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HF_{(aq)}$
b. $AgOH_{(aq)}$
c. $CH_3NH_2(aq)$

Answer

a. $HF(aq) + H_2O(\ell) \leftrightarrows F^-(aq) + H_3O^+(aq)$

b. $AgOH(aq) \leftrightarrows Ag^+(aq) + OH^-(aq)$

c. $CH_3NH_2(aq) + H_2O(\ell) \leftrightarrows CH_3NH_3^+(aq) + OH^-(aq)$

Acid Ionization Constant, K_a

The ionization for a general weak acid, HA, can be written as follows:

$$\mathrm{HA}\,(aq) \rightleftharpoons \mathrm{H^+}\,(aq) + \mathrm{A^-}\,(aq)$$

Because the acid is weak, an equilibrium expression can be written. An **acid ionization constant** (K_a) is the equilibrium constant for the ionization of an acid.

$$K_a = \frac{\left[\mathrm{H^+}\right]\left[\mathrm{A^-}\right]}{[\mathrm{HA}]}$$

The acid ionization represents the fraction of the original acid that has been ionized in solution. Therefore, the numerical value of K_a is a reflection of the strength of the acid. Weak acids with relatively higher K_a values are stronger than acids with relatively lower K_a values. Because strong acids are essentially 100% ionized, the concentration of the acid in the denominator is nearly zero and the K_a value approaches infinity. For this reason, K_a values are generally reported for weak acids only.

The table below is a listing of acid ionization constants for several acids. Note that polyprotic acids have a distinct ionization constant for each ionization step, with each successive ionization constant being smaller than the previous one.

Table 13.5.2: Acid Ionization Constants at $25^\circ\mathrm{C}$

Name of Acid	Ionization Equation	K_a
Sulfuric acid	$\mathrm{H_2SO_4} \rightleftharpoons \mathrm{H^+} + \mathrm{HSO_4^-}$ $\mathrm{HSO_4} \rightleftharpoons \mathrm{H^+} + \mathrm{SO_4^{2-}}$	very large 1.3×10^{-2}
Hydrofluoric acid	$\mathrm{HF} \rightleftharpoons \mathrm{H^+} + \mathrm{F^-}$	7.1×10^{-4}
Nitrous acid	$\mathrm{HNO_2} \rightleftharpoons \mathrm{H^+} + \mathrm{NO_2^-}$	4.5×10^{-4}
Benzoic acid	$\mathrm{C_6H_5COOH} \rightleftharpoons \mathrm{H^+} + \mathrm{C_6H_5COO^-}$	6.5×10^{-5}
Acetic acid	$\mathrm{CH_3COOH} \rightleftharpoons \mathrm{H^+} + \mathrm{CH_3COO^-}$	1.8×10^{-5}
Carbonic acid	$\mathrm{H_2CO_3} \rightleftharpoons \mathrm{H^+} + \mathrm{HCO_3^-}$ $\mathrm{HCO_3^-} \rightleftharpoons \mathrm{H^+} + \mathrm{CO_3^{2-}}$	4.2×10^{-7} 4.8×10^{-11}

Hydrofluoric acid $HF_{(aq)}$ reacts directly with glass (very few chemicals react with glass). Hydrofluoric acid is used in glass etching.

Strong and Weak Bases and Base Ionization Constant, K_b

As with acids, bases can either be strong or weak, depending on their extent of ionization. A **strong base** is a base, which ionizes completely in an aqueous solution. The most common strong bases are soluble metal hydroxide compounds such as potassium

hydroxide. Some metal hydroxides are not as strong simply because they are not as soluble. Calcium hydroxide is only slightly soluble in water, but the portion that does dissolve also dissociates into ions.

A **weak base** is a base that ionizes only slightly in an aqueous solution. Recall that a base can be defined as a substance, which accepts a hydrogen ion from another substance. When a weak base such as ammonia is dissolved in water, it accepts an H^+ ion from water, forming the hydroxide ion and the conjugate acid of the base, the ammonium ion.

$$NH_3\,(aq) + H_2O\,(l) \rightleftharpoons NH_4^+\,(aq) + OH^-\,(aq)$$

The equilibrium greatly favors the reactants and the extent of ionization of the ammonia molecule is very small.

An equilibrium expression can be written for the reactions of weak bases with water. Because the concentration of water is extremely large and virtually constant, the water is not included in the expression. A **base ionization constant** (K_b) is the equilibrium constant for the ionization of a base. For ammonia the expression is:

$$K_b = \frac{[NH_4^+]\,[OH^-]}{[NH_3]}$$

The numerical value of K_b is a reflection of the strength of the base. Weak bases with relatively higher K_b values are stronger than bases with relatively lower K_b values. Table 13.5.3is a listing of base ionization constants for several weak bases.

Table 13.5.3: Base Ionization Constants at 25°C

Name of Base	Ionization Equation	K_b
Methylamine	$CH_3NH_2 + H_2O \rightleftharpoons CH_3NH_3^+ + OH^-$	5.6×10^{-4}
Ammonia	$NH_3 + H_2O \rightleftharpoons NH_4^+ + OH^-$	1.8×10^{-5}
Pyridine	$C_5H_5N + H_2O \rightleftharpoons C_5H_5NH^+ + OH^-$	1.7×10^{-9}
Acetate ion	$CH_3COO^- + H_2O \rightleftharpoons CH_3COOH + OH^-$	5.6×10^{-10}
Fluoride ion	$F^- + H_2O \rightleftharpoons HF + OH^-$	1.4×10^{-11}
Urea	$H_2NCONH_2 + H_2O \rightleftharpoons H_2NCONH_3^+ + OH^-$	1.5×10^{-14}

The Ion-Product of Water

As we have already seen, H_2O can act as an acid or a base. Within any given sample of water, some H_2O molecules are acting as acids, and other H_2O molecules are acting as bases. The chemical equation is as follows:

$$\underbrace{H_2O}_{\text{acid}} + \underbrace{H_2O}_{\text{base}} \rightleftharpoons H_3O^+ + OH^- \tag{13.5.7}$$




Figure 13.5.2: Autoionization of water, giving hydroxide and hydronium ions.

Similar to a weak acid, the autoionization of water is an equilibrium process, and is more properly written as follows:

$$H_2O(\ell) + H_2O(\ell) \rightleftharpoons H_3O^+(aq) + OH^-(aq) \tag{13.5.8}$$

We often use the simplified form of the reaction:

$$H_2O(l) \rightleftharpoons H^+(aq) + OH-(aq)$$

The equilibrium constant for the autoionization of water is referred to as the ion-product for water and is given the symbol Kw.

$$K_w = [H^+][OH^-]$$

The **ion-product of water (*Kw*)** is the mathematical product of the concentration of hydrogen ions and hydroxide ions. Note that H_2O is not included in the ion-product expression because it is a pure liquid. The value of Kw is very small, in accordance with a

reaction that favors the reactants. At 25°C, the experimentally determined value of K_w in pure water is 1.0×10^{-14}.

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14}$$

In a sample of pure water, the concentrations of hydrogen and hydroxide ions are equal to one another. Pure water or any other aqueous solution in which this ratio holds is said to be **neutral.** To find the molarity of each ion, the square root of Kw is taken.

$$[H^+] = [OH^-] = 1.0\times10^{-7}$$

The product of these two concentrations is 1.0×10^{-14}

$$[H^+]\times[OH^-] = (1.0 \times 10^{-7})(1.0 \times 10^{-7}) = 1.0 \times 10^{-14}$$

- For acids, the concentration of H^+ or $[H^+]$) is greater than 1.0×10^{-7} M
- For bases, the concentration of OH^- or $[OH^-]$ is greater than 1.0×10^{-7} M.

Aqueous HCl is an example of acidic solution. Hydrogen chloride (HCl) ionizes to produce H^+ and Cl^- ions upon dissolving in water. This **increases the concentration of H^+ ions** in the solution. According to **Le Chatelier's principle**, the equilibrium represented by

$$H_2O(l) \rightleftharpoons H^+(aq) + OH^-(aq)$$

$$HCl(g) \longrightarrow H^+(aq) + Cl^-(aq)$$

is **forced to the left**, towards the reactant. As a result, the **concentration of the hydroxide ion decreases**.

Now, consider KOH (aq), a **basic solution**. Solid potassium hydroxide (KOH) dissociates in water to yield potassium ions and hydroxide ions.

$$KOH(s) \rightarrow K^+(aq) + OH^-(aq)$$

The **increase in concentration of the OH^- ions** will cause a decrease in the concentration of the H^+ ions.

No matter whether the aqueous solution is an acid, a base, or neutral:and the ion-product of $[H^+][OH^-]$ remains constant.

- For acidic solutions, $[H^+]$) is greater than $[OH^-]$.
- For basic solutions, [OH−] is greater than $[H^+]$.
- For neutral solutions, $[H_3O^+] = [OH-] = 1.0\times10^{-7}$M

This means that if you know $[H^+]$ for a solution, you can calculate what $[OH^-]$) has to be for the product to equal 1.0×10^{-14}, or if you know $[OH^-]$), you can calculate $[H^+]$. This also implies that as one concentration goes up, the other must go down to compensate so that their product always equals the value of K_w.

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \tag{13.5.9}$$

Example 13.5.2

Hydrochloric acid (HCl) is a strong acid, meaning it is 100% ionized in solution. What is the $[H^+]$ and the $[OH^-]$ in a solution of 2.0×10^{-3} M HCl?

Solution

Step 1: List the known values and plan the problem.

Known

- [HCl] = 2.0×10^{-3} M
- Kw = 1.0×10^{-14}

Unknown

- $[H^+]$=?M
- $[OH^-]$=?M

Because HCl is 100% ionized, the concentration of H+ ions in solution will be equal to the original concentration of HCl. Each HCl molecule that was originally present ionizes into one H+ ion and one Cl− ion. The concentration of OH− can then be determined from the [H+] and Kw.

Step 2: Solve.

$$[H^+]=2.0\times10^{-3}\,M$$

$$Kw = [H^+][OH^-] = 1.0\times10^{-14}$$

$$[OH^-] = Kw/[H^+] = 1.0\times10^{-14}/2.0\times10^{-3} = 5.0\times10^{-12}\,M$$

Step 3: Think about your result.

The $[H^+]$ is much higher than the $[OH^-]$ because the solution is acidic. As with other equilibrium constants, the unit for Kw is customarily omitted.

cise

Sodium hydroxide (NaOH) is a strong base. What is the $[H^+]$ and the $[OH^-]$ in a 0.001 M NaOH solution at 25 °C?

Answer

$[OH^-]$ = 0.001M or 1 x 10^{-3}M; $[H^+]=1\times10^{-11}$M.

The pH Scale

One qualitative measure of the strength of an acid or a base solution is the pH scale, which is based on the concentration of the hydronium (or hydrogen) ion in aqueous solution.

$$pH = -\log[H^+]$$

or

$$pH = -\log[H_3O^+]$$

Figure 13.5.3illustrates this relationship, along with some examples of various solutions. Because hydrogen ion concentrations are generally less than one (for example $1.3 \times 10^{-3}\ M$), the log of the number will be a negative number. To make pH even easier to work with, pH is defined as the ***negative*** **log of** $[H_3O^+]$, which will give a positive value for pH.




Figure 13.5.3: The relationaship between $[H^+]$ and *pH values for several common materials.*

A neutral (neither acidic nor basic) solution has a pH of 7. A pH below 7 means that a solution is acidic, with lower values of pH corresponding to increasingly acidic solutions. A pH greater than 7 indicates a basic solution, with higher values of pH corresponding to increasingly basic solutions. Thus, given the pH of several solutions, you can state which ones are acidic, which ones are basic, and which are more acidic or basic than others. These are summarized in Table \(\PageIndex{4}.

Table 13.5.4: Acidic, Basic and Neutral pH Values

Acidic, Basic and Neutral pH Values

Classification	Relative Ion Concentrations	pH at 25 °C
acidic	$[H^+] > [OH^-]$	pH < 7
neutral	$[H^+] = [OH^-]$	pH = 7
basic	$[H^+] < [OH^-]$	pH > 7

Example 13.5.3

Find the pH, given the $[H^+]$ of the following:

a. 1×10^{-3} M
b. 2.5×10^{-11} M
c. 4.7×10^{-9} M

Solution

pH = - log $[H_3O^+]$

Substitute the known quantity into the equation and solve. Use a scientific calculator for b and c.

a. pH = - log $[1 \times 10^{-3}]$ = 3.0 (1 decimal place since 1 has 1 significant figure)
b. pH = - log $[2.5 \times 10^{-11}]$ = 10.60 (2 decimal places since 2.5 has 2 significant figures)
c. pH = - log $[4.7 \times 10^{-9}]$ = 8.33 (2 decimal places since 4.7 has 2 significant figures)

Note on significant figures:

Because the number(s) before the decimal point in the pH value relate to the power on 10, the number of digits *after* the decimal point (underlined) is what determines the number of significant figures in the final answer.

cise

Find the pH, given $[H^+]$ of the following:

a. 5.8×10^{-4} M
b. 1.0×10^{-7} M

Answer

a. 3.24

b. 7.00

Table 13.5.5 lists the pH of several common solutions. The most acidic among the listed solutions is battery acid with the lowest pH value (0.3). The most basic is 1M NaOH solution with the highest pH value of 14.0. Notice that some biological fluids (stomach acid and urine) are nowhere near neutral. You may also notice that many food products are slightly acidic. They are acidic because they contain solutions of weak acids. If the acid components of these foods were strong acids, the food would likely be inedible.

Table 13.5.5: The pH Values of Some Common Solutions

Solution	pH
battery acid	0.3
stomach acid	1–2
lemon or lime juice	2.1

Solution	pH
vinegar	2.8–3.0
Coca-Cola	3
wine	2.8–3.8
beer	4–5
coffee	5
milk	6
urine	6
pure H_2O	7
(human) blood	7.3–7.5
sea water	8
antacid (milk of magnesia)	10.5
NH_3 (1 M)	11.6
bleach	12.6
NaOH (1 M)	14.0

Example 13.5.4

Label each solution as acidic, basic, or neutral based only on the stated pH.

a. milk of magnesia, pH = 10.5
b. pure water, pH = 7
c. wine, pH = 3.0

Solution

a. With a pH greater than 7, milk of magnesia is basic. (Milk of magnesia is largely $Mg(OH)_2$.)
b. Pure water, with a pH of 7, is neutral.
c. With a pH of less than 7, wine is acidic.

cise

Identify each substance as acidic, basic, or neutral based only on the stated pH.

a. human blood with pH = 7.4
b. household ammonia with pH = 11.0
c. cherries with pH = 3.6

Answer

a. slightly basic

b. basic

c. acidic

Acid Rain

Normal rainwater has a pH between 5 and 6 due to the presence of dissolved CO_2 which forms carbonic acid:

$$H_2O(l) + CO_2(g) \longrightarrow H_2CO_3(aq) \tag{13.5.10}$$

$$H_2CO_3(aq) \rightleftharpoons H^+(aq) + HCO_3^-(aq) \tag{13.5.11}$$

Acid rain is rainwater that has a pH of less than 5, due to a variety of nonmetal oxides, including CO_2, SO_2, SO_3, NO, and NO_2 being dissolved in the water and reacting with it to form not only carbonic acid, but sulfuric acid and nitric acid. The formation and subsequent ionization of sulfuric acid are shown here:

$$H_2O(l) + SO_3(g) \longrightarrow H_2SO_4(aq) \tag{13.5.12}$$

$$H_2SO_4(aq) \longrightarrow H^+(aq) + HSO_4^-(aq) \tag{13.5.13}$$

Carbon dioxide is naturally present in the atmosphere because we and most other organisms produce it as a waste product of metabolism. Carbon dioxide is also formed when fires release carbon stored in vegetation or when we burn wood or fossil fuels. Sulfur trioxide in the atmosphere is naturally produced by volcanic activity, but it also stems from burning fossil fuels, which have traces of sulfur, and from the process of "roasting" ores of metal sulfides in metal-refining processes. Oxides of nitrogen are formed in internal combustion engines where the high temperatures make it possible for the nitrogen and oxygen in air to chemically combine.

Acid rain is a particular problem in industrial areas where the products of combustion and smelting are released into the air without being stripped of sulfur and nitrogen oxides. In North America and Europe until the 1980s, it was responsible for the destruction of forests and freshwater lakes, when the acidity of the rain actually killed trees, damaged soil, and made lakes uninhabitable for all but the most acid-tolerant species. Acid rain also corrodes statuary and building facades that are made of marble and limestone (Figure 13.5.4). Regulations limiting the amount of sulfur and nitrogen oxides that can be released into the atmosphere by industry and automobiles have reduced the severity of acid damage to both natural and manmade environments in North America and Europe. It is now a growing problem in industrial areas of China and India.



Figure 13.5.4: (a) Acid rain makes trees more susceptible to drought and insect infestation, and depletes nutrients in the soil. (b) It also is corrodes statues that are carved from marble or limestone. (credit a: modification of work by Chris M Morris; credit b: modification of work by "Eden, Janine and Jim"/Flickr)

Two photos are shown. Photograph a on the left shows the upper portion of trees against a bright blue sky. The tops of several trees at the center of the photograph have bare branches and appear to be dead. Image b shows a statue of a man that appears to from the revolutionary war era in either marble or limestone.

Key Takeaways

- Acids and bases can be strong or weak depending on the extent of ionization in solution.
- Most chemical reactions reach equilibrium at which point there is no net change.
- The ion-product of $[H^+][OH^-]$ in an aqueous solution remains constant.
- A ***p*H** value is simply the *negative* of the logarithm of the H^+ ion concentration ($-\log[H^+]$).
- The pH scale is used to succinctly communicate the acidity or basicity of a solution.
- A solution is **acidic** if *p*H < 7.
- A solution is **basic** if *p*H > 7.
- A solution is **neutral** if *p*H = 7.

13.5: The Strengths of Acids and Bases is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.4: The Strengths of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.6: Buffers

Learning Objectives

- To define *buffer* and describe how it reacts with an acid or a base.

Weak acids are relatively common, even in the foods we eat. But we occasionally come across a strong acid or base, such as stomach acid, that has a strongly acidic pH of 1–2. By definition, strong acids and bases can produce a relatively large amount of hydrogen or hydroxide ions and, as a consequence, have a marked chemical activity. In addition, very small amounts of strong acids and bases can change the pH of a solution very quickly. If 1 mL of stomach acid [which we will approximate as 0.05 M HCl(aq)] is added to the bloodstream, and if no correcting mechanism is present, the pH of the blood would go from about 7.4 to about 4.9—a pH that is not conducive to continued living. Fortunately, the body has a mechanism for minimizing such dramatic pH changes.

The mechanism involves a **buffer**, a solution that resists dramatic changes in pH. A *buffer* (or *buffered*) solution is one that resists a change in its pH when H^+ or OH^- ions are added or removed owing to some other reaction taking place in the same solution. Buffers do so by being composed of certain pairs of solutes: either a **weak acid plus its conjugate base** or a **weak base plus its conjugate acid**.

For example, a buffer can be composed of dissolved **acetic acid ($HC_2H_3O_2$, a weak acid) and sodium acetate ($NaC_2H_3O_2$)**. Sodium acetate is a salt that dissociates into sodium ions and acetate ions in solution. For as long as acetic acid and acetate ions are present in significant amounts a solution, this can resist dramatic pH changes. Another example of a buffer is a solution containing **ammonia (NH_3, a weak base) and ammonium chloride (NH_4Cl).** Ammonium acetate is also a salt that dissociates into ammonium ions and chloride ions in solution. The presence of ammonium ions with ammonia molecules satisfies the requisite condition for a buffer solution.

How Buffers Work

The essential component of a buffer system is a **conjugate acid-base pair** whose concentration is fairly high in relation to the concentrations of added H^+ or OH^- it is expected to buffer against. Let us use an acetic acid–sodium acetate buffer to demonstrate how buffers work. If a strong base—a source of $OH^-(aq)$ ions—is added to the buffer solution, those hydroxide ions will react with the acetic acid in an acid-base reaction:

$$HC_2H_3O_2(aq) + OH^-(aq) \longrightarrow H_2O(\ell) + C_2H_3O_2^-(aq) \tag{13.6.1}$$

Rather than changing the pH dramatically by making the solution basic, the added hydroxide ions react to make water, and the pH does not change much.

> Many people are aware of the concept of buffers from *buffered aspirin*, which is aspirin that also has magnesium carbonate, calcium carbonate, magnesium oxide, or some other salt. The salt acts like a base, while aspirin is itself a weak acid.

If a strong acid—a source of H^+ ions—is added to the buffer solution, the H^+ ions will react with the anion from the salt. Because $HC_2H_3O_2$ is a weak acid, it is not ionized much. This means that if lots of hydrogen ions and acetate ions (from sodium acetate) are present in the same solution, they will come together to make acetic acid:

$$H^+(aq) + C_2H_3O_2^-(aq) \longrightarrow HC_2H_3O_2(aq) \tag{13.6.2}$$

Rather than changing the pH dramatically and making the solution acidic, the added hydrogen ions react to make molecules of a weak acid. Figure 13.6.1 illustrates both actions of a buffer.




Figure 13.6.1: The Action of Buffers. Buffers can react with both strong acids (top) and strong bases (bottom) to minimize large changes in pH.

A simple buffer system might be a 0.2 *M* solution of sodium acetate; the conjugate pair here is acetic acid HAc and its conjugate base, the acetate ion Ac^-. The idea is that this conjugate pair "pool" will be available to gobble up any small ($\leq 10^{-3}$ *M*) addition of H+ or OH^- that may result from other processes going on in the solution.




Figure 13.6.2: How HAc/Ac buffer works

Buffers work well only for limited amounts of added strong acid or base. Once either solute is all reacted, the solution is no longer a buffer, and rapid changes in pH may occur. We say that a buffer has a certain capacity. Buffers that have more solute dissolved in them to start with have larger capacities, as might be expected.

Buffers made from weak bases and salts of weak bases act similarly. For example, in a buffer containing NH_3 and NH_4Cl, ammonia molecules can react with any excess hydrogen ions introduced by strong acids:

$$NH_{3(aq)} + H^+_{(aq)} \rightarrow NH^+_{4(aq)} \tag{13.6.3}$$

while the ammonium ion [NH_4^+(aq)] can react with any hydroxide ions introduced by strong bases:

$$NH^+_{4(aq)} + OH^-_{(aq)} \rightarrow NH_{3(aq)} + H_2O_{(\ell)} \tag{13.6.4}$$




Figure 13.6.3: How NH_4^+/NH_3 buffer works

Example 13.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $HCHO_2$ and $NaCHO_2$
b. HCl and NaCl
c. CH_3NH_2 and CH_3NH_3Cl
d. NH_3 and NaOH

Solution

a. Formic acid ($HCHO_2$) is a weak acid, while $NaCHO_2$ is a salt supplying —formate ion (CHO_2^-), the conjugate base of $HCHO_2$. The combination of these two solutes would make a buffer solution.
b. Hydrochloric acid (HCl) is a strong acid, not a weak acid, so the combination of these two solutes would not make a buffer solution.
c. Methylamine (CH_3NH_2) is like ammonia, a weak base. The compound CH_3NH_3Cl is a salt supplying $CH_3NH_3^+$, the conjugate acid of CH_3NH_2. The combination of these two solutes would make a buffer solution.
d. Ammonia (NH_3) is a weak base, but NaOH is a strong base. The combination of these two solutes would not make a buffer solution.

? Exercise 13.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $NaHCO_3$ and NaCl
b. H_3PO_4 and NaH_2PO_4
c. NH_3 and $(NH_4)_3PO_4$
d. NaOH and NaCl

Answer

b. H_3PO_4 (weak acid) and $H_2PO_4^-$ (conjugate base of H_3PO_4) make a buffer.

c. NH_3 (weak base) and NH_4^+ (conjugate acid of NH_3) make a buffer

✔ Food and Drink App: The Acid That Eases Pain

Although medicines are not exactly "food and drink," we do ingest them, so let's take a look at an acid that is probably the most common medicine: acetylsalicylic acid, also known as aspirin. Aspirin is well known as a pain reliever and antipyretic (fever reducer).

The structure of aspirin is shown in the accompanying figure. The acid part is circled; it is the H atom in that part that can be donated as aspirin acts as a Brønsted-Lowry acid. Because it is not given in Table 10.5.1, acetylsalicylic acid is a weak acid. However, it is still an acid, and given that some people consume relatively large amounts of aspirin daily, its acidic nature can cause problems in the stomach lining, despite the stomach's defenses against its own stomach acid.




Figure 13.6.4 The Molecular Structure of Aspirin. The circled atoms are the acid part of the molecule.

Because the acid properties of aspirin may be problematic, many aspirin brands offer a "buffered aspirin" form of the medicine. In these cases, the aspirin also contains a buffering agent-usually MgO-that regulates the acidity of the aspirin to minimize its acidic side effects.

As useful and common as aspirin is, it was formally marketed as a drug starting in 1899. The US Food and Drug Administration (FDA), the governmental agency charged with overseeing and approving drugs in the United States, wasn't formed until 1906. Some have argued that if the FDA had been formed before aspirin was introduced, aspirin may never have gotten approval due to its potential for side effects-gastrointestinal bleeding, ringing in the ears, Reye's syndrome (a liver problem), and some allergic reactions. However, recently aspirin has been touted for its effects in lessening heart attacks and strokes, so it is likely that aspirin is here to stay.

Buffer solutions are essential components of all living organisms.

- Our blood is buffered to maintain a pH of 7.4 that must remain unchanged as metabolically-generated CO_2 (carbonic acid) is added and then removed by our lungs.
- Buffers in the oceans, in natural waters such as lakes and streams, and within soils help maintain their environmental stability against acid rain and increases in atmospheric CO_2.
- Many industrial processes, such as brewing, require buffer control, as do research studies in biochemistry and physiology that involve enzymes, are active only within certain pH ranges.

The pH in living systems (Figure \(\PageIndex{1}\)) is maintained by buffer systems.

Table 7.3.2: pH in Living Systems

Compartment	pH
Gastric Acid	1
Lysosomes	4.5
Granules of Chromaffin Cells	5.5
Human Skin	5.5
Urine	6
Neutral H_2O at 37 °C	6.81
Cytosol	7.2
Cerebrospinal Fluid	7.3
Blood	7.43-7.45
Mitochondrial Matrix	7.5
Pancreas Secretions	8.1

Human blood has a buffering system to minimize extreme changes in pH. One buffer in blood is based on the presence of HCO_3^- and H_2CO_3 [H_2CO_3 is another way to write CO_2(aq)]. With this buffer present, even if some stomach acid were to find its way directly into the bloodstream, the change in the pH of blood would be minimal. Inside many of the body's cells, there is a buffering system based on phosphate ions.

Medicine: The Buffer System in Blood

The normal pH of human blood is about 7.4. The carbonate buffer system in the blood uses the following equilibrium reaction:

$$\mathrm{CO_2}(g) + 2\,\mathrm{H_2O}(l) \rightleftharpoons \mathrm{H_2CO_3}(aq) \rightleftharpoons \mathrm{HCO_3^-}(aq) + \mathrm{H_3O^+}(aq)$$

The concentration of carbonic acid, H_2CO_3 is approximately 0.0012 *M*, and the concentration of the hydrogen carbonate ion, $\mathrm{HCO_3^-}$, is around 0.024 *M*. Using the Henderson-Hasselbalch equation and the pK_a of carbonic acid at body temperature, we can calculate the pH of blood:

$$\mathrm{pH} = \mathrm{p}K_a + \log\frac{[\mathrm{base}]}{[\mathrm{acid}]} = 6.1 + \log\frac{0.024}{0.0012} = 7.4$$

The fact that the H_2CO_3 concentration is significantly lower than that of the $\mathrm{HCO_3^-}$ ion may seem unusual, but this imbalance is due to the fact that most of the by-products of our metabolism that enter our bloodstream are acidic. Therefore, there must be a larger proportion of base than acid, so that the capacity of the buffer will not be exceeded.

Lactic acid is produced in our muscles when we exercise. As the lactic acid enters the bloodstream, it is neutralized by the $\mathrm{HCO_3^-}$ ion, producing H_2CO_3. An enzyme then accelerates the breakdown of the excess carbonic acid to carbon dioxide and water, which can be eliminated by breathing. In fact, in addition to the regulating effects of the carbonate buffering system on the pH of blood, the body uses breathing to regulate blood pH. If the pH of the blood decreases too far, an increase in breathing removes CO_2 from the blood through the lungs driving the equilibrium reaction such that [H_3O^+] is lowered. If the blood is too

alkaline, a lower breath rate increases CO_2 concentration in the blood, driving the equilibrium reaction the other way, increasing $[H^+]$ and restoring an appropriate pH.

Career Focus: Blood Bank Technology Specialist

At this point in this text, you should have the idea that the chemistry of blood is fairly complex. Because of this, people who work with blood must be specially trained to work with it properly.

A blood bank technology specialist is trained to perform routine and special tests on blood samples from blood banks or transfusion centers. This specialist measures the pH of blood, types it (according to the blood's ABO+/− type, Rh factors, and other typing schemes), tests it for the presence or absence of various diseases, and uses the blood to determine if a patient has any of several medical problems, such as anemia. A blood bank technology specialist may also interview and prepare donors to give blood and may actually collect the blood donation.

Blood bank technology specialists are well trained. Typically, they require a college degree with at least a year of special training in blood biology and chemistry. In the United States, training must conform to standards established by the American Association of Blood Banks.

Key Takeaway

- A buffer is a solution that resists sudden changes in pH.

Concept Review Exercise

1. Explain how a buffer prevents large changes in pH.

Answer

1. A buffer has components that react with both strong acids and strong bases to resist sudden changes in pH.

Exercises

1. Describe a buffer. What two related chemical components are required to make a buffer?
2. Can a buffer be made by combining a strong acid with a strong base? Why or why not?
3. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. HCl and NaCl
    2. HNO_2 and $NaNO_2$
    3. NH_4NO_3 and HNO_3
    4. NH_4NO_3 and NH_3
4. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. H_3PO_4 and Na_3PO_4
    2. $NaHCO_3$ and Na_2CO_3
    3. $NaNO_3$ and $Ca(NO_3)_2$
    4. HN_3 and NH_3
5. For each combination in Exercise 3 that is a buffer, write the chemical equations for the reactions of the buffer components when a strong acid and a strong base is added.
6. For each combination in Exercise 4 that is a buffer, write the chemical equations for the reaction of the buffer components when a strong acid and a strong base is added.
7. The complete phosphate buffer system is based on four substances: H_3PO_4, $H_2PO_4^-$, HPO_4^{2-}, and PO_4^{3-}. What different buffer solutions can be made from these substances?
8. Explain why NaBr cannot be a component in either an acidic or a basic buffer.
9. Explain why $Mg(NO_3)_2$ cannot be a component in either an acidic or a basic buffer.

Answers

1. A buffer resists sudden changes in pH. It has a weak acid or base and a salt of that weak acid or base.
2. No. Combining a strong acid and a strong base will produce salt and water. Excess strong acid or strong base will not act as a buffer.

3. 1. not a buffer
   2. buffer
   3. not a buffer
   4. buffer

4. 1. not a buffer

2. buffer

3. not a buffer

4. not buffer

5. 3b: strong acid: $H^+ + NO_2^- \rightarrow HNO_2$; strong base: $OH^- + HNO_2 \rightarrow H_2O + NO_2^-$; 3d: strong acid: $H^+ + NH_3 \rightarrow NH_4^+$; strong base: $OH^- + NH_4^+ \rightarrow H_2O + NH_3$
6. 4b: strong acid: $H^+ + CO_3^{2-} \rightarrow HCO3^-$; strong base: $OH^- + HCO_3^- \rightarrow H_2O + CO_3^{2-}$;
7. Buffers can be made by combining H_3PO_4 and $H_2PO_4^-$, $H_2PO_4^-$ and HPO_4^{2-}, and HPO_4^{2-} and PO_4^{3-}.
8. NaBr splits up into two ions in solution, Na^+ and Br^-. Na^+ will not react with any added base knowing that NaOH is a strong base. Br- will not react with any added acid knowing that HBr is a strong acid. Because NaBr will not react with any added base or acid, it does not resist change in pH and is not a buffer.
9. $Mg(NO_3)_2$ includes two types of ions, Mg^{2+} and NO_3^-. $Mg(OH)_2$ is strong base and completely dissociates (100% falls apart), so Mg^{2+} will not react with any added base (0% combines with OH^-). HNO_3 is strong acid and completely dissociates (100% falls apart), so NO_3^- will not react with any added acid (0% combines with H^+). Because $Mg(NO_3)_2$ will not react with any added base or acid, it does not resist change in pH and is not a buffer.

13.6: Buffers is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.5: Buffers** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.E: Acids and Bases (Exercises)

10.1: Arrhenius Definition of Acids and Bases

Concept Review Exercises

1. Give the Arrhenius definitions of an acid and a base.
2. What is neutralization?

Answers

1. Arrhenius acid: a compound that increases the concentration of hydrogen ion (H^+) in aqueous solution; Arrhenius base: a compound that increases the concentration of hydroxide ion (OH^-) in aqueous solution.
2. the reaction of an acid and a base

Exercises

1. Give two examples of Arrhenius acids.
2. Give two examples of Arrhenius bases.
3. List the general properties of acids.
4. List the general properties of bases.
5. Name each compound. (For acids, look up the name in Table 10.1.1. For bases, use the rules for naming ionic compounds from Chapter 3.)
    a. HBr(aq)
    b. $Ca(OH)_2$(aq)
    c. HNO_3(aq)
    d. $Fe(OH)_3$(aq)
6. Name each compound.
    a. HI(aq)
    b. $Cu(OH)_2$(aq)
    c. H_3PO_4(aq)
    d. CsOH(aq)
7. Write a balanced chemical equation for the neutralization of $Ba(OH)_2$(aq) with HNO_3(aq).
8. Write a balanced chemical equation for the neutralization of H_2SO_4(aq) with $Cr(OH)_3$(aq).
9. Gastric juice, the digestive fluid produced in the stomach, contains hydrochloric acid, HCl. Milk of Magnesia, a suspension of solid $Mg(OH)_2$ in an aqueous medium, is sometimes used to neutralize excess stomach acid. Write a complete balanced equation for the neutralization reaction.
10. Identify the salt produced in each acid-base reaction below. Then, balance the equation.
    a. $2HCl + Sr(OH)_2 \rightarrow 2H_2O$ + ??
    b. KNO_3; $HNO_3 + KOH \rightarrow$?? + H_2O
    c. $HF + Ca(OH)_2$ ---> ?? + H_2O
11. How many moles of sodium hydroxide (NaOH) are needed to neutralize 0.844 mol of acetic acid ($HC_2H_3O_2$)? (Hint: begin by writing a balanced chemical equation for the process.)
12. How many moles of perchloric acid ($HClO_4$) are needed to neutralize 0.052 mol of calcium hydroxide [$Ca(OH)_2$]? (Hint: begin by writing a balanced chemical equation for the process
13. Hydrazoic acid (HN_3) can be neutralized by a base.
    a. Write the balanced chemical equation for the reaction between hydrazoic acid and calcium hydroxide.

b. How many milliliters of 0.0245 M $Ca(OH)_2$ are needed to neutralize 0.564 g of HN_3?

14. Citric acid ($H_3C_6H_5O_7$) has three hydrogen atoms that can form hydrogen ions in solution.

a. Write the balanced chemical equation for the reaction between citric acid and sodium hydroxide.

b. If an orange contains 0.0675 g of $H_3C_6H_5O_7$, how many milliliters of 0.00332 M NaOH solution are needed to neutralize the acid?

15. Magnesium hydroxide [$Mg(OH)_2$] is an ingredient in some antacids. How many grams of $Mg(OH)_2$ are needed to neutralize the acid in 158 mL of 0.106 M HCl(aq)? It might help to write the balanced chemical equation first.

16. Aluminum hydroxide [$Al(OH)_3$] is an ingredient in some antacids. How many grams of $Al(OH)_3$ are needed to neutralize the acid in 96.5 mL of 0.556 M H_2SO_4(aq)? It might help to write the balanced chemical equation first.

17. Write the balanced chemical equation for the reaction between HBr and $Ca(OH)_2$. What volume of 0.010 M HBr solution is be required to neutralize 25 mL of a 0.0100*M* $Ca(OH)_2$ solution?

18. Write the balanced chemical equation for the reaction between HNO_3 and KOH. What volume of 0.5M HNO_3 is required to neutralize 60 mL of 0.4M KOH solution?

Answers

1. HCl and HNO_3 (answers will vary)
2. NaOH and $Ca(OH)_2$ (answers will vary)
3. sour taste, react with metals, react with bases, and turn litmus red
4. bitter taste, feels slippery, react with acids and turn litmus blue
5. a. hydrobromic acid

b. calcium hydroxide

c. nitric acid

d. iron(III) hydroxide

6. a. hydroiodic acid

b. cupric hydroxide

c. phosphoric acid

d. cesium hydroxide

7. $2HNO_3(aq) + Ba(OH)_2(aq) \rightarrow Ba(NO_3)_2(aq) + 2H_2O$

8. $3H_2SO_4(aq) + 2Cr(OH)_3(aq) \rightarrow Cr_2(SO_4)_3(aq) + 6H_2O$

9. $Mg(OH)_2 + 2HCl$ --> $MgCl_2 + 2H_2O$

10. a. $SrCl_2$; $2HCl + Sr(OH)_2 \rightarrow 2H_2O + SrCl_2$

b. KNO_3; $HNO_3 + KOH \rightarrow KNO_3 + H_2O$

c. CaF_2; $2HF + Ca(OH)_2 \rightarrow CaF_2 + 2H_2O$

11. 0.844 mol

12. 0.104 mol

13. Part 1: $2HN_3(aq) + Ca(OH)_2 \rightarrow Ca(N_3)_2 + 2H_2O$

Part 2: 268 mL

14. Part 1: $H_3C_6H_5O_7(aq) + 3NaOH(aq) \rightarrow Na_3C_6H_5O_7(aq) + 3H_2O$

Part 2: 317.5 mL

15. 0.488 g

16. 2.79 g

17. $2HBr + Ca(OH)_2 \rightarrow CaBr_2 + 2H_2O$; 50 mL HBr

18. $HNO_3 + KOH \rightarrow KNO_3 + H_2O$; 48 mL HNO_3

10.2: Brønsted-Lowry Definition of Acids and Bases

Concept Review Exercise

1. Give the definitions of a Brønsted-Lowry acid and a Brønsted-Lowry base.

Answer

1. A Brønsted-Lowry acid is a proton donor, while a Brønsted-Lowry base is a proton acceptor.

Exercises

1. Label each reactant as a Brønsted-Lowry acid or a Brønsted-Lowry base.

   $HCl(aq) + NH_3(aq) \rightarrow NH_4^+(aq) + Cl^-(aq)$

2. Label each reactant as a Brønsted-Lowry acid or a Brønsted-Lowry base.

   $H_2O(\ell) + N_2H_4(aq) \rightarrow N_2H_5^+(aq) + OH^-(aq)$

3. Explain why a Brønsted-Lowry acid can be called a proton donor.
4. Explain why a Brønsted-Lowry base can be called a proton acceptor.
5. Write the chemical equation of the reaction of ammonia in water and label the Brønsted-Lowry acid and base.
6. Write the chemical equation of the reaction of methylamine (CH_3NH_2) in water and label the Brønsted-Lowry acid and base.
7. Demonstrate that the dissolution of HNO_3 in water is actually a Brønsted-Lowry acid-base reaction by describing it with a chemical equation and labeling the Brønsted-Lowry acid and base.
8. Identify the Brønsted-Lowry acid and base in the following chemical equation:

   $C_3H_7NH_2(aq) + H_3O^+(aq) \rightarrow C_3H_7NH_3^+(aq) + H_2O(\ell)$

9. Identify the Brønsted-Lowry acid and the Brønsted-Lowry base in each of the following equations

   1. $NO_2^- + H_2O \longrightarrow HNO_2 + OH^-$
   2. $HBr + H_2O \longrightarrow H_3O^+ + Br^-$
   3. $HS^- + H_2O \longrightarrow H_2S + OH^-$
   4. $H_2PO_4^- + OH^- \longrightarrow HPO_4^{2-} + H_2O$
   5. $H_2PO_4^- + HCl \longrightarrow H_3PO_4 + Cl^-$

10. Write the chemical equation for the reaction that occurs when cocaine hydrochloride ($C_{17}H_{22}ClNO_4$) dissolves in water and donates a proton to a water molecule. (When hydrochlorides dissolve in water, they separate into chloride ions and the appropriate cation.)

11. If codeine hydrobromide has the formula $C_{18}H_{22}BrNO_3$, what is the formula of the parent compound codeine?

Answers

1. HCl: Brønsted-Lowry acid; NH_3: Brønsted-Lowry base
2. H_2O: Brønsted-Lowry acid; N_2H_4: Brønsted-Lowry base
3. A Brønsted-Lowry acid gives away an H^+ ion—nominally, a proton—in an acid-base reaction.
4. A Brønsted-Lowry base accepts an H^+ ion (a proton) in an acid-base reaction.
5. $NH_3 + H_2O \rightarrow NH_4^+ + OH^-$ (here NH_3 = Brønsted-Lowry base; H_2O = Brønsted-Lowry acid)
6. $CH_3NH_2 + H_2O \rightarrow CH_3NH_3^+ + OH^-$ (here CH_3NH_2 = Brønsted-Lowry base; H_2O = Brønsted-Lowry acid)
7. $HNO_3 + H_2O \rightarrow H_3O^+ + NO_3^-$ (here HNO_3 = Brønsted-Lowry acid; H_2O = Brønsted-Lowry base)
8. $C_3H_7NH_2(aq) + H_3O^+(aq) \rightarrow C_3H_7NH_3^+(aq) + H_2O(\ell)$ (here H_3O^+ = Brønsted-Lowry acid; $C_3H_7NH_2$ = Brønsted-Lowry base)

10.3: Water - Both an Acid and a Base

Concept Review Exercises

1. Explain how water can act as an acid.
2. Explain how water can act as a base.

Answers

1. Under the right conditions, H_2O can donate a proton, making it a Brønsted-Lowry acid.
2. Under the right conditions, H_2O can accept a proton, making it a Brønsted-Lowry base.

Exercises

1. Is $H_2O(\ell)$ acting as an acid or a base?

   $H_2O(\ell) + NH_4^+(aq) \rightarrow H_3O^+(aq) + NH_3(aq)$

2. Is $H_2O(\ell)$ acting as an acid or a base?

   $CH_3^-(aq) + H_2O(\ell) \rightarrow CH_4(aq) + OH^-(aq)$

3. In the aqueous solutions of some salts, one of the ions from the salt can react with water molecules. In some $C_2H_3O_2^-$ solutions, the following reaction can occur:

   $C_2H_3O_2^-(aq) + H_2O(\ell) \rightarrow HC_2H_3O_2(aq) + OH^-(aq)$

   Is H_2O acting as an acid or a base in this reaction?

4. In the aqueous solutions of some salts, one of the ions from the salt can react with water molecules. In some NH_4^+ solutions, the following reaction can occur:

   $NH_4^+(aq) + H_2O \rightarrow NH_3(aq) + H_3O^+(aq)$

   Is H_2O acting as an acid or a base in this reaction?

5. Why is pure water considered neutral?

Answers

1. base
2. acid
3. acid
4. base
5. When water ionizes, equal amounts of H^+ (acid) and OH^- (base) are formed, so the solution is neither acidic nor basic: $H_2O(\ell) \rightarrow H^+(aq) + OH^-(aq)$

10.4: The Strengths of Acids and Bases

Concept Review Exercises

1. Explain the difference between a strong acid or base and a weak acid or base.
2. Explain what is occurring when a chemical reaction reaches equilibrium.
3. Define *pH*.

AnswerS

1. A strong acid or base is 100% ionized in aqueous solution; a weak acid or base is less than 100% ionized.
2. The overall reaction progress stops because the reverse process balances out the forward process.
3. pH is a measure of the hydrogen ion concentration.

Exercises

1. Name a strong acid and a weak acid. (Hint: use Table 10.4.1.)
2. Name a strong base and a weak base. (Hint: use Table 10.4.1.)
3. Is each compound a strong acid or a weak acid? Assume all are in aqueous solution. (Hint: use Table 10.4.1.)
    1. HF
    2. $HC_2H_3O_2$
    3. HCl
    4. $HClO_4$
4. Is each compound a strong acid or a weak acid? Assume all are in aqueous solution. (Hint: use Table 10.4.1.)
    1. H_2SO_4
    2. HSO_4^-
    3. HPO_4^{2-}
    4. HNO_3
5. Is each compound a strong base or a weak base? Assume all are in aqueous solution. (Hint: use Table 10.4.1.)
    1. NH_3
    2. NaOH
    3. $Mg(OH)_2$
    4. $Cu(OH)_2$
6. Is each compound a strong base or a weak base? Assume all are in aqueous solution. (Hint: use Table 10.4.1.)
    1. KOH
    2. H_2O
    3. $Fe(OH)_2$
    4. $Fe(OH)_3$
7. Write the chemical equation for the equilibrium process for each weak acid in Exercise 3.
8. Write the chemical equation for the equilibrium process for each weak acid in Exercise 4.
9. Write the chemical equation for the equilibrium process for each weak base in Exercise 5.
10. Write the chemical equation for the equilibrium process for each weak base in Exercise 6.
11. Which is the stronger acid—HCl(aq) or HF(aq)?
12. Which is the stronger base—KOH(aq) or $Ni(OH)_2$(aq)?
13. Consider the two acids in Exercise 11. For solutions that have the same concentration, which one would you expect to have a lower pH?
14. Consider the two bases in Exercise 12. For solutions that have the same concentration, which one would you expect to have a higher pH?
15. Consider the list of substances in Table \PageIndex3\PageIndex3.2"The pH Values of Some Common Solutions". What is the most acidic substance on the list that you have encountered recently?
16. Consider the list of substances in Table \PageIndex3\PageIndex3.2"The pH Values of Some Common Solutions". What is the most basic substance on the list that you have encountered recently?
17. Indicate whether solutions with the following pH values are acidic, basic, or neutral:
    1. pH = 9.4
    2. pH = 7.0
    3. pH = 1.2
    4. pH = 6.5

Answers

1. strong acid: HCl; weak acid: $HC_2H_3O_2$ (answers will vary)
2. strong base: NaOH; weak base: NH_3 (answers will vary)
3.
    1. weak
    2. weak
    3. strong
    4. strong

4. 1. strong
   2. weak
   3. weak
   4. strong
5. 1. weak
   2. strong
   3. strong
   4. weak
6. 1. strong
   2. weak
   3. weak
   4. weak

7. 3a: $HF(aq) \leftrightarrows H^+(aq) + F^-(aq)$; 3b: $HC_2H_3O_2(aq) \leftrightarrows H^+(aq) + C_2H_3O_2^-(aq)$

8. 4b: $HSO_4^-(aq) \leftrightarrows H^+(aq) + SO_4^{2-}(aq)$; 4c: $HPO_4^{2-}(aq) \leftrightarrows H^+(aq) + PO_4^{3-}(aq)$

9. 5a: $NH_3(aq) + H_2O \leftrightarrows NH_4^+(aq) + OH^-(aq)$; 5d: $Cu(OH)_2(aq) \leftrightarrows Cu^{2+}(aq) + 2OH^-(aq)$

10. 6b: $H_2O + H_2O \leftrightarrows H_3O^+(aq) + OH^-(aq)$; 6c: $Fe(OH)_2(aq) \leftrightarrows Fe^{2+}(aq) + 2OH^-(aq)$; 6d: $Fe(OH)3(aq) \leftrightarrows Fe^{3+}(aq) + 3OH^-(aq)$

11. HCl(aq)

12. KOH(aq)

13. HCl(aq)

14. KOH(aq)

15. (answers will vary)

16. (answers will vary)

17. 1. basic

2. neutral

3. acidic (strongly)

4. acidic (mildly)

10.5: Buffers

Concept Review Exercise

1. Explain how a buffer prevents large changes in pH.

Answer

1. A buffer has components that react with both strong acids and strong bases to resist sudden changes in pH.

Exercises

1. Describe a buffer. What two related chemical components are required to make a buffer?
2. Can a buffer be made by combining a strong acid with a strong base? Why or why not?
3. Which solute combinations can make a buffer? Assume all are aqueous solutions.
   1. HCl and NaCl
   2. HNO_2 and $NaNO_2$
   3. NH_4NO_3 and HNO_3
   4. NH_4NO_3 and NH_3
4. Which solute combinations can make a buffer? Assume all are aqueous solutions.
   1. H_3PO_4 and Na_3PO_4
   2. $NaHCO_3$ and Na_2CO_3

3. $NaNO_3$ and $Ca(NO_3)_2$
4. HN_3 and NH_3

5. For each combination in Exercise 3 that is a buffer, write the chemical equations for the reactions of the buffer components when a strong acid and a strong base is added.
6. For each combination in Exercise 4 that is a buffer, write the chemical equations for the reaction of the buffer components when a strong acid and a strong base is added.
7. The complete phosphate buffer system is based on four substances: H_3PO_4, $H_2PO_4^-$, HPO_4^{2-}, and PO_4^{3-}. What different buffer solutions can be made from these substances?
8. Explain why NaBr cannot be a component in either an acidic or a basic buffer.
9. Explain why $Mg(NO_3)_2$ cannot be a component in either an acidic or a basic buffer.

Answers

1. A buffer resists sudden changes in pH. It has a weak acid or base and a salt of that weak acid or base.
2. No. Combining a strong acid and a strong base will produce salt and water. Excess strong acid or strong base will not act as a buffer.
3. 1. not a buffer
   2. buffer
   3. not a buffer
   4. buffer
4. 1. not a buffer

   2. buffer

   3. not a buffer

   4. not buffer
5. 3b: strong acid: $H^+ + NO_2^- \rightarrow HNO_2$; strong base: $OH^- + HNO_2 \rightarrow H_2O + NO_2^-$; 3d: strong acid: $H^+ + NH_3 \rightarrow NH_4^+$; strong base: $OH^- + NH_4^+ \rightarrow H_2O + NH_3$
6. 4b: strong acid: $H^+ + CO_3^{2-} \rightarrow HCO3^-$; strong base: $OH^- + HCO_3^- \rightarrow H_2O + CO_3^{2-}$;
7. Buffers can be made by combining H_3PO_4 and $H_2PO_4^-$, $H_2PO_4^-$ and HPO_4^{2-}, and HPO_4^{2-} and PO_4^{3-}.
8. NaBr splits up into two ions in solution, Na^+ and Br^-. Na^+ will not react with any added base knowing that NaOH is a strong base. Br- will not react with any added acid knowing that HBr is a strong acid. Because NaBr will not react with any added base or acid, it does not resist change in pH and is not a buffer.
9. $Mg(NO_3)_2$ includes two types of ions, Mg^{2+} and NO_3^-. $Mg(OH)_2$ is strong base and completely dissociates (100% falls apart), so Mg^{2+} will not react with any added base (0% combines with OH^-). HNO_3 is strong acid and completely dissociates (100% falls apart), so NO_3^- will not react with any added acid (0% combines with H^+). Because $Mg(NO_3)_2$ will not react with any added base or acid, it does not resist change in pH and is not a buffer.

Additional Exercises

1. The properties of a 1.0 M HCl solution and a 1.0 M $HC_2H_3O_2$ solution are compared. Measurements show that the hydrochloric acid solution has a higher osmotic pressure than the acetic acid solution. Explain why.
2. Of a 0.50 M HNO_3 solution and a 0.50 M $HC_2H_3O_2$ solution, which should have the higher boiling point? Explain why.
3. The reaction of sulfuric acid [$H_2SO_4(aq)$] with sodium hydroxide [NaOH(aq)] can be represented by two separate steps, with only one hydrogen ion reacting in each step. Write the chemical equation for each step.
4. The reaction of aluminum hydroxide [$Al(OH)_3(aq)$] with hydrochloric acid [HCl(aq)] can be represented by three separate steps, with only one hydroxide ion reacting in each step. Write the chemical equation for each step.
5. A friend brings you a small sample of an unknown chemical. Assuming that the chemical is soluble in water, how would you determine if the chemical is an acid or a base?
6. A neutral solution has a hydrogen ion concentration of about 1×10^{-7} M. What is the concentration of the hydroxide ion in a neutral solution?
7. The Lewis definitions of an acid and a base are based on electron pairs, not protons. A Lewis acid is an electron pair acceptor, while a Lewis base is an electron pair donor. Use Lewis diagrams to show that

$H^+(aq) + OH^-(aq) \rightarrow H_2O(\ell)$

is an acid-base reaction in the Lewis sense as well as in the Arrhenius and Brønsted-Lowry senses.

8. Given the chemical reaction

$NH_3(g) + BF_3(g) \rightarrow NH_3—BF_3(s)$

show that the reaction illustrated by this equation is an acid-base reaction if we use the Lewis definitions of an acid and a base (see Exercise 7). The product contains a bond between the N and B atoms.

Answers

1. HCl is a strong acid and yields more ions in solution. $HC_2H_3O_2$ is a weak acid and undergoes partial ionization in solution.
2. HNO_3 is a strong acid while $HC_2H_3O_2$ is a weak acid. HNO_3 dissociates 100% and its solution contains more ions. The more ions the solution contains the lower is its vapor pressure; the higher temperature is required for it to boil.
3. $H_2SO_4 + NaOH \rightarrow NaHSO_4 + H_2O$; $NaHSO_4 + NaOH \rightarrow Na_2SO_4 + H_2O$
4. $Al(OH)_3 + HCl \rightarrow Al(OH)_2Cl + H_2O$; $Al(OH)_2Cl + HCl \rightarrow Al(OH)Cl_2 + H_2O$; $Al(OH)Cl_2 + HCl \rightarrow AlCl3 + H_2O$
5. One way is to add it to $NaHCO_3$; if it bubbles, it is an acid. Alternatively, add the sample to litmus and look for a characteristic color change (red for acid, blue for base).
6. In a neutral solution, $[OH^-] = [H^+] = 1.0 \times 10^{-7}$ M
7. The O atom is donating an electron pair to the H^+ ion, making the base an electron pair donor and the acid an electron pair acceptor.
8. The N atom is donating a lone pair to B in BF_3, Hence NH_3 is the Lewis base and BF_3 is the Lewis acid.

13.E: Acids and Bases (Exercises) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.E: Acids and Bases (Exercises)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **10.1: Arrhenius Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **10.2: Brønsted-Lowry Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **10.3: Water - Both an Acid and a Base** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **10.4: The Strengths of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **10.5: Buffers** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

13.S: Acids and Bases (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

The earliest chemical definition of an acid, the **Arrhenius definition**, says that an **acid** is a compound that increases the amount of hydrogen ion (H^+) in aqueous solution. An **Arrhenius base** is a compound that increases the amount of hydroxide ion (OH^-) in aqueous solution. While most bases are named as ionic hydroxide compounds, aqueous acids have a naming system unique to acids. Acids and bases react together in a characteristic chemical reaction called **neutralization**, in which the products are water and a salt. The principles of stoichiometry, along with the balanced chemical equation for a reaction between an acid and a base, can be used to determine how much of one compound will react with a given amount of the other.

A **Brønsted-Lowry acid** is any substance that donates a proton to another substance. A **Brønsted-Lowry base** is any substance that accepts a proton from another substance. The reaction of ammonia with water to make ammonium ions and hydroxide ions can be used to illustrate Brønsted-Lowry acid and base behavior.

Some compounds can either donate or accept protons, depending on the circumstances. Such compounds are called **amphiprotic**. Water is one example of an amphiprotic compound. One result of water being amphiprotic is that a water molecule can donate a proton to another water molecule to make a hydronium ion and a hydroxide ion. This process is called the **autoionization of water** and occurs in any sample of water.

Not all acids and bases are equal in chemical strength. A **strong acid** is an acid whose molecules are all dissociated into ions in aqueous solution. Hydrochloric acid is an example of a strong acid. Similarly, a **strong base** is a base whose molecules are dissociated into ions in aqueous solution. Sodium hydroxide is an example of a strong base. Any acid or base whose molecules are not all dissociated into ions in aqueous solution is a **weak acid** or a **weak base**. Solutions of weak acids and weak bases reach a **chemical equilibrium** between the un-ionized form of the compound and the dissociated ions. It is a dynamic equilibrium because acid and base molecules are constantly dissociating into ions and reassociating into neutral molecules.

The **pH** scale is a scale used to express the concentration of hydrogen ions in solution. A neutral solution, neither acidic nor basic, has a pH of 7. Acidic solutions have a pH lower than 7, while basic solutions have a pH higher than 7.

Buffers are solutions that resist dramatic changes in pH when an acid or a base is added to them. They contain a weak acid and a salt of that weak acid, or a weak base and a salt of that weak base. When a buffer is present, any strong acid reacts with the anion of the salt, forming a weak acid and minimizing the presence of hydrogen ions in solution. Any strong base reacts with the weak acid, minimizing the amount of additional hydroxide ions in solution. However, buffers only have limited **capacity**; there is a limit to the amount of strong acid or strong base any given amount of buffer will react with.

13.S: Acids and Bases (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **10.S: Acids and Bases (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons

We begin our study of organic chemistry with the alkanes, compounds containing only two elements, carbon and hydrogen, and having only single bonds. There are several other kinds of hydrocarbons, distinguished by the types of bonding between carbon atoms and by the properties that result from that bonding. We will first examine hydrocarbons with double bonds, with triple bonds, and with a special kind of bonding called aromaticity. Then we will study some compounds considered to be derived from hydrocarbons by replacing one or more hydrogen atoms with an oxygen-containing group. Finally, we focuse on organic acids and bases, after which we will be ready to look at the chemistry of life itself—biochemistry—in the remaining five chapters.

14.1: Organic Chemistry
14.2: Structures and Names of Alkanes
14.3: Alkenes- Structures and Names
14.4: Alkynes
14.5: Branched-Chain Alkanes
14.6: Condensed Structural and Line-Angle Formulas
14.7: IUPAC Nomenclature
14.8: Physical Properties of Alkanes
14.9: Physical Properties of Alkenes
14.10: Chemical Properties of Alkanes
14.11: Chemical Properties of Alkenes
14.12: Halogenated Hydrocarbons
14.13: Cycloalkanes
14.14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons
14.15: Organic Reactions
14.E: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Exercises)
14.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary)
14.S: Unsaturated and Aromatic Hydrocarbons (Summary)

Template:HideTOC

14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

14.1: Organic Chemistry

Learning Objectives

- To recognize the composition and properties typical of organic and inorganic compounds.

Scientists of the 18th and early 19th centuries studied compounds obtained from plants and animals and labeled them *organic* because they were isolated from "organized" (living) systems. Compounds isolated from nonliving systems, such as rocks and ores, the atmosphere, and the oceans, were labeled *inorganic*. For many years, scientists thought organic compounds could be made by only living organisms because they possessed a vital force found only in living systems. The vital force theory began to decline in 1828, when the German chemist Friedrich Wöhler synthesized urea from inorganic starting materials. He reacted silver cyanate ($AgOCN$) and ammonium chloride (NH_4Cl), expecting to get ammonium cyanate (NH_4OCN). What he expected is described by the following equation.

$$AgOCN + NH_4Cl \rightarrow AgCl + NH_4OCN \tag{14.1.1}$$

Instead, he found the product to be urea (NH_2CONH_2), a well-known organic material readily isolated from urine. This result led to a series of experiments in which a wide variety of organic compounds were made from inorganic starting materials. The vital force theory gradually went away as chemists learned that they could make many organic compounds in the laboratory.

Today organic chemistry is the study of the chemistry of the carbon compounds, and inorganic chemistry is the study of the chemistry of all other elements. It may seem strange that we divide chemistry into two branches—one that considers compounds of only one element and one that covers the 100-plus remaining elements. However, this division seems more reasonable when we consider that of tens of millions of compounds that have been characterized, the overwhelming majority are carbon compounds.

> The word *organic* has different meanings. Organic fertilizer, such as cow manure, is organic in the original sense; it is derived from living organisms. Organic foods generally are foods grown without synthetic pesticides or fertilizers. Organic chemistry is the chemistry of compounds of carbon.

Carbon is unique among the other elements in that its atoms can form stable covalent bonds with each other and with atoms of other elements in a multitude of variations. The resulting molecules can contain from one to millions of carbon atoms. We previously surveyed organic chemistry by dividing its compounds into families based on functional groups. We begin with the simplest members of a family and then move on to molecules that are organic in the original sense—that is, they are made by and found in living organisms. These complex molecules (all containing carbon) determine the forms and functions of living systems and are the subject of biochemistry.

Organic compounds, like inorganic compounds, obey all the natural laws. Often there is no clear distinction in the chemical or physical properties among organic and inorganic molecules. Nevertheless, it is useful to compare typical members of each class, as in Table 14.1.1.

Table 14.1.1: General Contrasting Properties and Examples of Organic and Inorganic Compounds

Organic	Hexane		Inorganic	NaCl
low melting points	−95°C		high melting points	801°C
low boiling points	69°C		high boiling points	1,413°C
low solubility in water; high solubility in nonpolar solvents	insoluble in water; soluble in gasoline		greater solubility in water; low solubility in nonpolar solvents	soluble in water; insoluble in gasoline
flammable	highly flammable		nonflammable	nonflammable
aqueous solutions do not conduct electricity	nonconductive		aqueous solutions conduct electricity	conductive in aqueous solution
exhibit covalent bonding	covalent bonds		exhibit ionic bonding	ionic bonds

Keep in mind, however, that there are exceptions to every category in this table. To further illustrate typical differences among organic and inorganic compounds, Table 14.1.1 also lists properties of the inorganic compound sodium chloride (common table salt, NaCl) and the organic compound hexane (C_6H_{14}), a solvent that is used to extract soybean oil from soybeans (among other uses). Many compounds can be classified as organic or inorganic by the presence or absence of certain typical properties, as illustrated in Table 14.1.1.

14.1: Organic Chemistry is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.1: Organic Chemistry** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.2: Structures and Names of Alkanes

Learning Objectives

- To identify and name simple (straight-chain) alkanes given formulas and write formulas for straight-chain alkanes given their names.

We begin our study of organic chemistry with the hydrocarbons, the simplest organic compounds, which are composed of carbon and hydrogen atoms only. As we noted, there are several different kinds of hydrocarbons. They are distinguished by the types of bonding between carbon atoms and the properties that result from that bonding. Hydrocarbons with only carbon-to-carbon single bonds (C–C) and existing as a continuous chain of carbon atoms also bonded to hydrogen atoms are called alkanes (or saturated hydrocarbons). *Saturated*, in this case, means that each carbon atom is bonded to four other atoms (hydrogen or carbon)—the most possible; there are no double or triple bonds in the molecules.

> The word *saturated* has the same meaning for hydrocarbons as it does for the dietary fats and oils: the molecule has no carbon-to-carbon double bonds (C=C).

We previously introduced the three simplest alkanes—methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) and they are shown again in Figure 14.2.1.




Figure 14.2.1: The Three Simplest Alkanes

The flat representations shown do not accurately portray bond angles or molecular geometry. Methane has a tetrahedral shape that chemists often portray with wedges indicating bonds coming out toward you and dashed lines indicating bonds that go back away from you. An ordinary solid line indicates a bond in the plane of the page. Recall that the VSEPR theory correctly predicts a tetrahedral shape for the methane molecule (Figure 14.2.2).




Figure 14.2.2: The Tetrahedral Methane Molecule

Methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) are the beginning of a series of compounds in which any two members in a sequence differ by one carbon atom and two hydrogen atoms—namely, a CH_2 unit. The first 10 members of this series are given in Table 14.2.1.

Table 14.2.1: The First 10 Straight-Chain Alkanes

Name	Molecular Formula (C_nH_{2n+2})	Condensed Structural Formula	Number of Possible Isomers
methane	CH_4	CH_4	—
ethane	C_2H_6	CH_3CH_3	—
propane	C_3H_8	$CH_3CH_2CH_3$	—
butane	C_4H_{10}	$CH_3CH_2CH_2CH_3$	2
pentane	C_5H_{12}	$CH_3CH_2CH_2CH_2CH_3$	3

Name	Molecular Formula (C_nH_{2n+2})	Condensed Structural Formula	Number of Possible Isomers
hexane	C_6H_{14}	$CH_3CH_2CH_2CH_2CH_2CH_3$	5
heptane	C_7H_{16}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_3$	9
octane	C_8H_{18}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	18
nonane	C_9H_{20}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	35
decane	$C_{10}H_{22}$	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	75

Consider the series in Figure 14.2.3. The sequence starts with C_3H_8, and a CH_2 unit is added in each step moving up the series. Any family of compounds in which adjacent members differ from each other by a definite factor (here a CH_2 group) is called a homologous series. The members of such a series, called *homologs*, have properties that vary in a regular and predictable manner. The principle of *homology* gives organization to organic chemistry in much the same way that the periodic table gives organization to inorganic chemistry. Instead of a bewildering array of individual carbon compounds, we can study a few members of a homologous series and from them deduce some of the properties of other compounds in the series.




Figure 14.2.3: Members of a Homologous Series. Each succeeding formula incorporates one carbon atom and two hydrogen atoms more than the previous formula.

The principle of homology allows us to write a general formula for alkanes: C_nH_{2n+2}. Using this formula, we can write a molecular formula for any alkane with a given number of carbon atoms. For example, an alkane with eight carbon atoms has the molecular formula $C_8H_{(2\times 8)+2} = C_8H_{18}$.

Key Takeaway

- Simple alkanes exist as a homologous series, in which adjacent members differ by a CH_2 unit.

14.2: Structures and Names of Alkanes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.


- **12.2: Structures and Names of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.3: Alkenes- Structures and Names

Learning Objectives

- To name alkenes given formulas and write formulas for alkenes given names.

As noted before, alkenes are hydrocarbons with carbon-to-carbon double bonds ($R_2C{=}CR_2$) and alkynes are hydrocarbons with carbon-to-carbon triple bonds (R–C≡C–R). Collectively, they are called unsaturated hydrocarbons because they have fewer hydrogen atoms than does an alkane with the same number of carbon atoms, as is indicated in the following general formulas:




Some representative alkenes—their names, structures, and physical properties—are given in Table 14.3.1.

Table 14.3.1: Physical Properties of Some Selected Alkenes

IUPAC Name	Molecular Formula	Condensed Structural Formula	Melting Point (°C)	Boiling Point (°C)
ethene	C_2H_4	$CH_2{=}CH_2$	–169	–104
propene	C_3H_6	$CH_2{=}CHCH_3$	–185	–47
1-butene	C_4H_8	$CH_2{=}CHCH_2CH_3$	–185	–6
1-pentene	C_5H_{10}	$CH_2{=}CH(CH_2)_2CH_3$	–138	30
1-hexene	C_6H_{12}	$CH_2{=}CH(CH_2)_3CH_3$	–140	63
1-heptene	C_7H_{14}	$CH_2{=}CH(CH_2)_4CH_3$	–119	94
1-octene	C_8H_{16}	$CH_2{=}CH(CH_2)_5CH_3$	–102	121

We used only condensed structural formulas in Table 14.3.1. Thus, $CH_2{=}CH_2$ stands for



The double bond is shared by the two carbons and does not involve the hydrogen atoms, although the condensed formula does not make this point obvious. Note that the molecular formula for ethene is C_2H_4, whereas that for ethane is C_2H_6.

The first two alkenes in Table 14.3.1, ethene and propene, are most often called by their common names—ethylene and propylene, respectively (Figure 14.3.1). Ethylene is a major commercial chemical. The US chemical industry produces about 25 billion kilograms of ethylene annually, more than any other synthetic organic chemical. More than half of this ethylene goes into the manufacture of polyethylene, one of the most familiar plastics. Propylene is also an important industrial chemical. It is converted to plastics, isopropyl alcohol, and a variety of other products.





Figure 14.3.1: Ethene and Propene. The ball-and-spring models of ethene/ethylene (a) and propene/propylene (b) show their respective shapes, especially bond angles.

Although there is only one alkene with the formula C_2H_4 (ethene) and only one with the formula C_3H_6 (propene), there are several alkenes with the formula C_4H_8.


Here are some basic rules for naming alkenes from the International Union of Pure and Applied Chemistry (IUPAC):


1. The longest chain of carbon atoms *containing the double bond* is considered the parent chain. It is named using the same stem as the alkane having the same number of carbon atoms but ends in *-ene* to identify it as an alkene. Thus the compound CH_2=$CHCH_3$ is *propene*.
2. If there are four or more carbon atoms in a chain, we must indicate the position of the double bond. The carbons atoms are numbered so that the first of the two that are doubly bonded is given the lower of the two possible numbers.The compound CH_3CH=$CHCH_2CH_3$, for example, has the double bond between the second and third carbon atoms. Its name is 2-pentene (not 3-pentene).
3. Substituent groups are named as with alkanes, and their position is indicated by a number. Thus, the structure below is 5-methyl-2-hexene. Note that the numbering of the parent chain is always done in such a way as to give the double bond the lowest number, even if that causes a substituent to have a higher number. *The double bond always has priority in numbering.*



Example 14.3.1

Name each compound.

a. 

b. 

Solution


a. The longest chain containing the double bond has five carbon atoms, so the compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the fourth carbon atom (rule 3), so the compound's name is 4-methyl-2-pentene.

b. The longest chain containing the double bond has five carbon atoms, so the parent compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the third carbon atom (rule 3), so the compound's name is 3-methyl-2-pentene.

Exercise 14.3.1

Name each compound.

a. $CH_3CH_2CH_2CH_2CH_2CH{=}CHCH_3$

b. 

Answer

Just as there are cycloalkanes, there are *cycloalkenes*. These compounds are named like alkenes, but with the prefix *cyclo-* attached to the beginning of the parent alkene name.

Example 14.3.2


Draw the structure for each compound.

a. 3-methyl-2-pentene
b. cyclohexene


Solution

a.

First write the parent chain of five carbon atoms: C–C–C–C–C. Then add the double bond between the second and third carbon atoms:




Now place the methyl group on the third carbon atom and add enough hydrogen atoms to give each carbon atom a total of four bonds.



b

First, consider what each of the three parts of the name means. *Cyclo* means a ring compound, *hex* means 6 carbon atoms, and *-ene* means a double bond.



Exercise 14.3.2

Draw the structure for each compound.

a. 2-ethyl-1-hexene
b. cyclopentene

Key Takeaway

- Alkenes are hydrocarbons with a carbon-to-carbon double bond.

14.3: Alkenes- Structures and Names is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **13.1: Alkenes- Structures and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.4: Alkynes

Learning Objectives

- Describe the general physical and chemical properties of alkynes.
- Name alkynes given formulas and write formulas for alkynes given names.

The simplest alkyne—a hydrocarbon with carbon-to-carbon triple bond—has the molecular formula C_2H_2 and is known by its common name—acetylene (Figure 14.4.1). Its structure is H–C≡C–H.




Figure 14.4.1: Ball-and-Spring Model of Acetylene. Acetylene (ethyne) is the simplest member of the alkyne family.

Acetylene is used in oxyacetylene torches for cutting and welding metals. The flame from such a torch can be very hot. Most acetylene, however, is converted to chemical intermediates that are used to make vinyl and acrylic plastics, fibers, resins, and a variety of other products.

Alkynes are similar to alkenes in both physical and chemical properties. For example, alkynes undergo many of the typical addition reactions of alkenes. The International Union of Pure and Applied Chemistry (IUPAC) names for alkynes parallel those of alkenes, except that the family ending is *-yne* rather than *-ene*. The IUPAC name for acetylene is ethyne. The names of other alkynes are illustrated in the following exercises.

Key Takeaway

- Alkynes are hydrocarbons with carbon-to-carbon triple bonds and properties much like those of alkenes.

14.4: Alkynes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **13.6: Alkynes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.5: Branched-Chain Alkanes


Learning Objectives

- To learn how alkane molecules can have branched chains and recognize compounds that are isomers.


We can write the structure of butane (C_4H_{10}) by stringing four carbon atoms in a row,

–C–C–C–C–

and then adding enough hydrogen atoms to give each carbon atom four bonds:



The compound butane has this structure, but there is another way to put 4 carbon atoms and 10 hydrogen atoms together. Place 3 of the carbon atoms in a row and then branch the fourth one off the middle carbon atom:



Now we add enough hydrogen atoms to give each carbon four bonds.




Figure 14.5.1).





Figure 14.5.1: Butane and Isobutane. The ball-and-stick models of these two compounds show them to be isomers; both have the molecular formula C_4H_{10}.


Notice that C_4H_{10} is depicted with a bent chain in Figure 14.5.1. The four-carbon chain may be bent in various ways because the groups can rotate freely about the C–C bonds. However, this rotation does not change the identity of the compound. It is important to realize that bending a chain does *not* change the identity of the compound; all of the following represent the same compound, butane:



The structure of isobutane shows a continuous chain of three carbon atoms only, with the fourth attached as a branch off the middle carbon atom of the continuous chain, which is different from the structures of butane (compare the two structures in Figure 14.5.1.

Unlike C_4H_{10}, the compounds methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) do not exist in isomeric forms because there is only one way to arrange the atoms in each formula so that each carbon atom has four bonds.

Next beyond C_4H_{10} in the homologous series is pentane. Each compound has the same molecular formula: C_5H_{12}. (Table 12.1.1 from the previous section has a column identifying the number of possible isomers for the first 10 straight-chain alkanes.) The compound at the far left is pentane because it has all five carbon atoms in a continuous chain. The compound in the middle is isopentane; like isobutane, it has a one CH_3 branch off the second carbon atom of the continuous chain. The compound at the far right, discovered after the other two, was named neopentane (from the Greek *neos*, meaning "new"). Although all three have the same molecular formula, they have different properties, including boiling points: pentane, 36.1°C; isopentane, 27.7°C; and neopentane, 9.5°C.



A continuous (unbranched) chain of carbon atoms is often called a *straight chain* even though the tetrahedral arrangement about each carbon gives it a zigzag shape. Straight-chain alkanes are sometimes called *normal alkanes*, and their names are given the prefix *n*-. For example, butane is called *n*-butane. We will not use that prefix here because it is not a part of the system established by the International Union of Pure and Applied Chemistry.

14.5: Branched-Chain Alkanes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.3: Branched-Chain Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.6: Condensed Structural and Line-Angle Formulas

Learning Objectives

- Write condensed structural formulas for alkanes given complete structural formulas.
- Draw line-angle formulas given structural formulas.

We use several kinds of formulas to describe organic compounds. A *molecular formula* shows only the kinds and numbers of atoms in a molecule. For example, the molecular formula C_4H_{10} tells us there are 4 carbon atoms and 10 hydrogen atoms in a molecule, but it doesn't distinguish between butane and isobutane. A structural formula shows all the carbon and hydrogen atoms and the bonds attaching them. Thus, structural formulas identify the specific isomers by showing the order of attachment of the various atoms.

Unfortunately, structural formulas are difficult to type/write and take up a lot of space. Chemists often use condensed structural formulas to alleviate these problems. The condensed formulas show hydrogen atoms right next to the carbon atoms to which they are attached, as illustrated for butane:



The ultimate condensed formula is a line-angle formula, in which carbon atoms are implied at the corners and ends of lines, and each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds. For example, we can represent pentane ($CH_3CH_2CH_2CH_2CH_3$) and isopentane [$(CH_3)_2CHCH_2CH_3$] as follows:



Pentane Isopentane

Parentheses in condensed structural formulas indicate that the enclosed grouping of atoms is attached to the adjacent carbon atom.

Key Takeaways

- Condensed chemical formulas show the hydrogen atoms (or other atoms or groups) right next to the carbon atoms to which they are attached.
- Line-angle formulas imply a carbon atom at the corners and ends of lines. Each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds.

14.6: Condensed Structural and Line-Angle Formulas is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.4: Condensed Structural and Line-Angle Formulas** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.7: IUPAC Nomenclature

Learning Objectives

- To name alkanes by the IUPAC system and write formulas for alkanes given IUPAC names



As noted in previously, the number of isomers increases rapidly as the number of carbon atoms increases. There are 3 pentanes, 5 hexanes, 9 heptanes, and 18 octanes. It would be difficult to assign unique individual names that we could remember. A systematic way of naming hydrocarbons and other organic compounds has been devised by the International Union of Pure and Applied Chemistry (IUPAC). These rules, used worldwide, are known as the IUPAC System of Nomenclature. (Some of the names we used earlier, such as isobutane, isopentane, and neopentane, do not follow these rules and are called *common names*.) A stem name (Table 14.7.1) indicates the number of carbon atoms in the longest continuous chain (LCC). Atoms or groups attached to this carbon chain, called *substituents*, are then named, with their positions indicated by numbers. For now, we will consider only those substituents called alkyl groups.

Table 14.7.1: Stems That Indicate the Number of Carbon Atoms in Organic Molecules








Stem	Number
meth-	1
eth-	2
prop-	3
but-	4
pent-	5
hex-	6
hept-	7
oct-	8
non-	9
dec-	10

An alkyl group is a group of atoms that results when one hydrogen atom is removed from an alkane. The group is named by replacing the *-ane* suffix of the parent hydrocarbon with *-yl*. For example, the $-CH_3$ group derived from methane (CH_4) results from subtracting one hydrogen atom and is called a *methyl group*. The alkyl groups we will use most frequently are listed in Table 14.7.2 Alkyl groups are not independent molecules; they are parts of molecules that we consider as a unit to name compounds systematically.

Table 14.7.2: Common Alkyl Groups

Parent Alkane		Alkyl Group		Condensed Structural Formula
methane		methyl		$CH_3–$

*There are four butyl groups, two derived from butane and two from isobutane. We will introduce the other three where appropriate.


Parent Alkane		Alkyl Group		Condensed Structural Formula
ethane		ethyl		$CH_3CH_2–$
propane		propyl		$CH_3CH_2CH_2–$
		isopropyl		$(CH_3)_2CH–$
butane		butyl*		$CH_3CH_2CH_2CH_2–$

*There are four butyl groups, two derived from butane and two from isobutane. We will introduce the other three where appropriate.

Simplified IUPAC rules for naming alkanes are as follows (demonstrated in Example 14.7.1).

1. **Name alkanes according to the LCC (longest continuous chain) of carbon atoms in the molecule (rather than the total number of carbon atoms).** This LCC, considered the parent chain, determines the base name, to which we add the suffix *-ane* to indicate that the molecule is an alkane.

2. **If the hydrocarbon is branched, number the carbon atoms of the LCC.** Numbers are assigned in the direction that gives the lowest numbers to the carbon atoms with attached substituents. Hyphens are used to separate numbers from the names of substituents; commas separate numbers from each other. (The LCC need not be written in a straight line; for example, the LCC in the following has five carbon atoms.)






3. **Place the names of the substituent groups in alphabetical order before the name of the parent compound.** If the same alkyl group appears more than once, the numbers of all the carbon atoms to which it is attached are expressed. If the same group appears more than once on the same carbon atom, the number of that carbon atom is repeated as many times as the group appears. Moreover, the number of identical groups is indicated by the Greek prefixes *di-*, *tri-*, *tetra-*, and so on. These prefixes are *not* considered in determining the alphabetical order of the substituents. For example, ethyl is listed before dimethyl; the di- is simply ignored. The last alkyl group named is prefixed to the name of the parent alkane to form one word.

When these rules are followed, every unique compound receives its own exclusive name. The rules enable us to not only name a compound from a given structure but also draw a structure from a given name. The best way to learn how to use the IUPAC system is to put it to work, not just memorize the rules. It's easier than it looks.

Example 14.7.1

Name each compound.

1. 




- 
- 

Solution

1. The LCC has five carbon atoms, and so the parent compound is pentane (rule 1). There is a methyl group (rule 2) attached to the second carbon atom of the pentane chain. The name is therefore 2-methylpentane.
2. The LCC has six carbon atoms, so the parent compound is hexane (rule 1). Methyl groups (rule 2) are attached to the second and fifth carbon atoms. The name is 2,5-dimethylhexane.
3. The LCC has eight carbon atoms, so the parent compound is octane (rule 1). There are methyl and ethyl groups (rule 2), both attached to the fourth carbon atom (counting from the *right* gives this carbon atom a lower number; rule 3). The correct name is thus 4-ethyl-4-methyloctane.

Exercise 14.7.1

Name each compound.

1. 
- 
- 

Example 14.7.2


Draw the structure for each compound.

a. 2,3-dimethylbutane
b. 4-ethyl-2-methylheptane


Solution

In drawing structures, always start with the parent chain.


a. The parent chain is butane, indicating four carbon atoms in the LCC.



Then add the groups at their proper positions. You can number the parent chain from either direction as long as you are consistent; just don't change directions before the structure is done. The name indicates two methyl (CH_3) groups, one on the second carbon atom and one on the third.




Finally, fill in all the hydrogen atoms, keeping in mind that each carbon atom must have four bonds.




- The parent chain is heptane in this case, indicating seven carbon atoms in the LCC. –C–C–C–C–C–C–C–

  Adding the groups at their proper positions gives



  Filling in all the hydrogen atoms gives the following condensed structural formulas:



  Note that the bonds (dashes) can be shown or not; sometimes they are needed for spacing.

? Exercise 14.7.2

Draw the structure for each compound.

a. 4-ethyloctane
b. 3-ethyl-2-methylpentane
c. 3,3,5-trimethylheptane

Key Takeaway

- Alkanes have both common names and systematic names, specified by IUPAC.

14.7: IUPAC Nomenclature is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.5: IUPAC Nomenclature** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.8: Physical Properties of Alkanes

Learning Objectives

- To identify the physical properties of alkanes and describe trends in these properties.

Because alkanes have relatively predictable physical properties and undergo relatively few chemical reactions other than combustion, they serve as a basis of comparison for the properties of many other organic compound families. Let's consider their physical properties first.

Table 14.8.1 describes some of the properties of some of the first 10 straight-chain alkanes. Because alkane molecules are nonpolar, they are insoluble in water, which is a polar solvent, but are soluble in nonpolar and slightly polar solvents. Consequently, alkanes themselves are commonly used as solvents for organic substances of low polarity, such as fats, oils, and waxes. Nearly all alkanes have densities less than 1.0 g/mL and are therefore less dense than water (the density of H_2O is 1.00 g/mL at 20°C). These properties explain why oil and grease do not mix with water but rather float on its surface.

Table 14.8.1: Physical Properties of Some Alkanes

Molecular Name	Formula	Melting Point (°C)	Boiling Point (°C)	Density (20°C)*	Physical State (at 20°C)
methane	CH_4	–182	–164	0.668 g/L	gas
ethane	C_2H_6	–183	–89	1.265 g/L	gas
propane	C_3H_8	–190	–42	1.867 g/L	gas
butane	C_4H_{10}	–138	–1	2.493 g/L	gas
pentane	C_5H_{12}	–130	36	0.626 g/mL	liquid
hexane	C_6H_{14}	–95	69	0.659 g/mL	liquid
octane	C_8H_{18}	–57	125	0.703 g/mL	liquid
decane	$C_{10}H_{22}$	–30	174	0.730 g/mL	liquid

*Note the change in units going from gases (grams per liter) to liquids (grams per milliliter). Gas densities are at 1 atm pressure.



Figure 14.8.1: Oil Spills. Crude oil coats the water's surface in the Gulf of Mexico after the *Deepwater Horizon* oil rig sank following an explosion. The leak was a mile below the surface, making it difficult to estimate the size of the spill. One liter of oil can create a slick 2.5 hectares (6.3 acres) in size. This and similar spills provide a reminder that hydrocarbons and water don't mix. Source: Photo courtesy of NASA Goddard / MODIS Rapid Response Team, NASA.gov, Topics, Earth Features, Oil Spill(opens in new window) [www.nasa.gov].

Looking Closer: Gas Densities and Fire Hazards

Table 14.8.1 indicates that the first four members of the alkane series are gases at ordinary temperatures. Natural gas is composed chiefly of methane, which has a density of about 0.67 g/L. The density of air is about 1.29 g/L. Because natural gas is less dense than air, it rises. When a natural-gas leak is detected and shut off in a room, the gas can be removed by opening an upper window. On the other hand, bottled gas can be either propane (density 1.88 g/L) or butanes (a mixture of butane and isobutane; density about 2.5 g/L). Both are much heavier than air (density 1.2 g/L). If bottled gas escapes into a building, it collects near the floor. This presents a much more serious fire hazard than a natural-gas leak because it is more difficult to rid the room of the heavier gas.

Also shown in Table 14.8.1 are the boiling points of the straight-chain alkanes increase with increasing molar mass. This general rule holds true for the straight-chain homologs of all organic compound families. Larger molecules have greater surface areas and consequently interact more strongly; more energy is therefore required to separate them. For a given molar mass, the boiling points of alkanes are relatively low because these nonpolar molecules have only weak dispersion forces to hold them together in the liquid state.

Looking Closer: An Alkane Basis for Properties of Other Compounds

An understanding of the physical properties of the alkanes is important in that petroleum and natural gas and the many products derived from them—gasoline, bottled gas, solvents, plastics, and more—are composed primarily of alkanes. This understanding is also vital because it is the basis for describing the properties of other organic and biological compound families. For example, large portions of the structures of lipids consist of nonpolar alkyl groups. Lipids include the dietary fats and fatlike compounds called phospholipids and sphingolipids that serve as structural components of living tissues. These compounds have both polar and nonpolar groups, enabling them to bridge the gap between water-soluble and water-insoluble phases. This characteristic is essential for the selective permeability of cell membranes.




Figure 14.8.2: Tripalmitin (a), a typical fat molecule, has long hydrocarbon chains typical of most lipids. Compare these chains to hexadecane (b), an alkane with 16 carbon atoms.

Key Takeaway

- Alkanes are nonpolar compounds that are low boiling and insoluble in water.

14.8: Physical Properties of Alkanes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.6: Physical Properties of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.9: Physical Properties of Alkenes

Learning Objectives

- To identify the physical properties of alkenes and describe trends in these properties.

The physical properties of alkenes are similar to those of the alkanes. The table at the start of the chapter shows that the boiling points of straight-chain alkenes increase with increasing molar mass, just as with alkanes. For molecules with the same number of carbon atoms and the same general shape, the boiling points usually differ only slightly, just as we would expect for substances whose molar mass differs by only 2 u (equivalent to two hydrogen atoms). Like other hydrocarbons, the alkenes are insoluble in water but soluble in organic solvents.

Looking Closer: Environmental Note

Alkenes occur widely in nature. Ripening fruits and vegetables give off ethylene, which triggers further ripening. Fruit processors artificially introduce ethylene to hasten the ripening process; exposure to as little as 0.1 mg of ethylene for 24 h can ripen 1 kg of tomatoes. Unfortunately, this process does not exactly duplicate the ripening process, and tomatoes picked green and treated this way don't taste much like vine-ripened tomatoes fresh from the garden.



The bright red color of tomatoes is due to lycopene—a polyene.

Other alkenes that occur in nature include 1-octene, a constituent of lemon oil, and octadecene ($C_{18}H_{36}$) found in fish liver. Dienes (two double bonds) and polyenes (three or more double bonds) are also common. Butadiene (CH_2=$CHCH$=CH_2) is found in coffee. Lycopene and the carotenes are isomeric polyenes ($C_{40}H_{56}$) that give the attractive red, orange, and yellow colors to watermelons, tomatoes, carrots, and other fruits and vegetables. Vitamin A, essential to good vision, is derived from a carotene. The world would be a much less colorful place without alkenes.

Key Takeaway

- The physical properties of alkenes are much like those of the alkanes: their boiling points increase with increasing molar mass, and they are insoluble in water.

14.9: Physical Properties of Alkenes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.


- **13.3: Physical Properties of Alkenes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.10: Chemical Properties of Alkanes

Learning Objectives

- To identify the main chemical properties of alkanes.

Alkane molecules are nonpolar and therefore generally do not react with ionic compounds such as most laboratory acids, bases, oxidizing agents, or reducing agents. Consider butane as an example:



Butane plus O H superscript negative sign yields no reaction. There is also no reaction of butane with H superscript positive sign and Mn O subscript 4 superscript negative sign.

Neither positive ions nor negative ions are attracted to a nonpolar molecule. In fact, the alkanes undergo so few reactions that they are sometimes called *paraffins*, from the Latin *parum affinis*, meaning "little affinity."

Two important reactions that the alkanes do undergo are combustion and halogenation. Nothing happens when alkanes are merely mixed with oxygen (O_2) at room temperature, but when a flame or spark provides the activation energy, a highly exothermic combustion reaction proceeds vigorously. For methane (CH_4), the reaction is as follows:

$$CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O + \text{heat} \tag{14.10.1}$$

If the reactants are adequately mixed and there is sufficient oxygen, the only products are carbon dioxide (CO_2), water (H_2O), and heat—heat for cooking foods, heating homes, and drying clothes. Because conditions are rarely ideal, however, other products are frequently formed. When the oxygen supply is limited, carbon monoxide (CO) is a by-product:

$$2CH_4 + 3O_2 \rightarrow 2CO + 4H_2O \tag{14.10.2}$$

This reaction is responsible for dozens of deaths each year from unventilated or improperly adjusted gas heaters. (Similar reactions with similar results occur with kerosene heaters.)

Alkanes also react with the halogens chlorine (Cl_2) and bromine (Br_2) in the presence of ultraviolet light or at high temperatures to yield chlorinated and brominated alkanes. For example, chlorine reacts with excess methane (CH_4) to give methyl chloride (CH_3Cl).

$$CH_4 + Cl_2 \rightarrow CH_3Cl + HCl \tag{14.10.3}$$

With more chlorine, a mixture of products is obtained: CH_3Cl, CH_2Cl_2, $CHCl_3$, and CCl_4. Fluorine (F_2), the lightest halogen, combines explosively with most hydrocarbons. Iodine (I_2) is relatively unreactive. Fluorinated and iodinated alkanes are produced by indirect methods.

Key Takeaway

- Alkanes react with oxygen (combustion) and with halogens (halogenation).

14.10: Chemical Properties of Alkanes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.


- **12.7: Chemical Properties of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.11: Chemical Properties of Alkenes

Learning Objectives

- To write equations for the addition reactions of alkenes with hydrogen, halogens, and water

Alkenes are valued mainly for addition reactions, in which one of the bonds in the double bond is broken. Each of the carbon atoms in the bond can then attach another atom or group while remaining joined to each other by a single bond. Perhaps the simplest addition reaction is hydrogenation—a reaction with hydrogen (H_2) in the presence of a catalyst such as nickel (Ni) or platinum (Pt).



Ethylene Hydrogen Ethane


The product is an alkane having the same carbon skeleton as the alkene.

Alkenes also readily undergo halogenation—the addition of halogens. Indeed, the reaction with bromine (Br_2) can be used to test for alkenes. Bromine solutions are brownish red. When we add a Br_2 solution to an alkene, the color of the solution disappears because the alkene reacts with the bromine:



Ethylene Bromine (brownish-red) 1,2-Dibromoethane (colorless)

Another important addition reaction is that between an alkene and water to form an alcohol. This reaction, called hydration, requires a catalyst—usually a strong acid, such as sulfuric acid (H_2SO_4):



Ethylene Water Ethanol

The hydration reaction is discussed later, where we deal with this reaction in the synthesis of alcohols.


Example 14.11.1


Write the equation for the reaction between $CH_3CH{=}CHCH_3$ and each substance.


a. H_2 (Ni catalyst)
b. Br_2
c. H_2O (H_2SO_4 catalyst)

Solution

In each reaction, the reagent adds across the double bond.

a. 

b. 

c. 

? Exercise 14.11.1

Write the equation for each reaction.

a. $CH_3CH_2CH{=}CH_2$ with H_2 (Ni catalyst)
b. $CH_3CH{=}CH_2$ with Cl_2
c. $CH_3CH_2CH{=}CHCH_2CH_3$ with H_2O (H_2SO_4 catalyst)

Key Takeaway

- Alkenes undergo addition reactions, adding such substances as hydrogen, bromine, and water across the carbon-to-carbon double bond.

14.11: Chemical Properties of Alkenes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **13.4: Chemical Properties of Alkenes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.12: Halogenated Hydrocarbons

Learning Objectives

- To name halogenated hydrocarbons given formulas and write formulas for these compounds given names.

Many organic compounds are closely related to the alkanes. As we noted previously, alkanes react with halogens to produce halogenated hydrocarbons, the simplest of which have a single halogen atom substituted for a hydrogen atom of the alkane. Even more closely related are the cycloalkanes, compounds in which the carbon atoms are joined in a ring, or cyclic fashion.

The reactions of alkanes with halogens produce halogenated hydrocarbons, compounds in which one or more hydrogen atoms of a hydrocarbon have been replaced by halogen atoms:



The replacement of only one hydrogen atom gives an alkyl halide (or haloalkane). The *common names* of alkyl halides consist of two parts: the name of the alkyl group plus the stem of the name of the halogen, with the ending *-ide*. The IUPAC system uses the name of the parent alkane with a prefix indicating the halogen substituents, preceded by number indicating the substituent's location. The prefixes are *fluoro-*, *chloro-*, *bromo-*, and *iodo-*. Thus CH_3CH_2Cl has the common name ethyl chloride and the IUPAC name chloroethane. Alkyl halides with simple alkyl groups (one to four carbon atoms) are often called by common names. Those with a larger number of carbon atoms are usually given IUPAC names.

Example 14.12.1

Give the common and IUPAC names for each compound.

a. $CH_3CH_2CH_2Br$
b. $(CH_3)_2CHCl$

Solution

a. The alkyl group ($CH_3CH_2CH_2$–) is a propyl group, and the halogen is bromine (Br). The common name is therefore propyl bromide. For the IUPAC name, the prefix for bromine (bromo) is combined with the name for a three-carbon chain (propane), preceded by a number identifying the carbon atom to which the Br atom is attached, so the IUPAC name is 1-bromopropane.
b. The alkyl group [$(CH_3)_2CH$–] has three carbon atoms, with a chlorine (Cl) atom attached to the middle carbon atom. The alkyl group is therefore isopropyl, and the common name of the compound is isopropyl chloride. For the IUPAC name, the Cl atom (prefix *chloro-*) attached to the middle (second) carbon atom of a propane chain results in 2-chloropropane.


Exercise 14.12.1


Give common and IUPAC names for each compound.

a. CH_3CH_2I
b. $CH_3CH_2CH_2CH_2F$

Example 14.12.2

Give the IUPAC name for each compound.

1. 


- 


Solution

1. The parent alkane has five carbon atoms in the longest continuous chain; it is pentane. A bromo (Br) group is attached to the second carbon atom of the chain. The IUPAC name is 2-bromopentane.
2. The parent alkane is hexane. Methyl (CH_3) and bromo (Br) groups are attached to the second and fourth carbon atoms, respectively. Listing the substituents in alphabetical order gives the name 4-bromo-2-methylhexane.

? Exercise 14.12.2

Give the IUPAC name for each compound.

a. 

- 

A wide variety of interesting and often useful compounds have one or more halogen atoms per molecule. For example, methane (CH_4) can react with chlorine (Cl_2), replacing one, two, three, or all four hydrogen atoms with Cl atoms. Several halogenated products derived from methane and ethane (CH_3CH_3) are listed in Table 14.12.1, along with some of their uses.

Table 14.12.1: Some Halogenated Hydrocarbons

Formula	Common Name	IUPAC Name	Some Important Uses
Derived from CH_4			
CH_3Cl	methyl chloride	chloromethane	refrigerant; the manufacture of silicones, methyl cellulose, and synthetic rubber
CH_2Cl_2	methylene chloride	dichloromethane	laboratory and industrial solvent
$CHCl_3$	chloroform	trichloromethane	industrial solvent
CCl_4	carbon tetrachloride	tetrachloromethane	dry-cleaning solvent and fire extinguishers (but no longer recommended for use)
$CBrF_3$	halon-1301	bromotrifluoromethane	fire extinguisher systems
CCl_3F	chlorofluorocarbon-11 (CFC-11)	trichlorofluoromethane	foaming plastics
CCl_2F_2	chlorofluorocarbon-12 (CFC-12)	dichlorodifluoromethane	refrigerant
Derived from CH_3CH_3			
CH_3CH_2Cl	ethyl chloride	chloroethane	local anesthetic
$ClCH_2CH_2Cl$	ethylene dichloride	1,2-dichloroethane	solvent for rubber
CCl_3CH_3	methylchloroform	1,1,1-trichloroethane	solvent for cleaning computer chips and molds for shaping plastics

To Your Health: Halogenated Hydrocarbons

Once widely used in consumer products, many chlorinated hydrocarbons are suspected carcinogens (cancer-causing substances) and also are known to cause severe liver damage. An example is carbon tetrachloride (CCl_4), once used as a dry-cleaning solvent

and in fire extinguishers but no longer recommended for either use. Even in small amounts, its vapor can cause serious illness if exposure is prolonged. Moreover, it reacts with water at high temperatures to form deadly phosgene ($COCl_2$) gas, which makes the use of CCl_4 in fire extinguishers particularly dangerous.

Ethyl chloride, in contrast, is used as an external local anesthetic. When sprayed on the skin, it evaporates quickly, cooling the area enough to make it insensitive to pain. It can also be used as an emergency general anesthetic.

Bromine-containing compounds are widely used in fire extinguishers and as fire retardants on clothing and other materials. Because they too are toxic and have adverse effects on the environment, scientists are engaged in designing safer substitutes for them, as for many other halogenated compounds.

To Your Health: Chlorofluorocarbons and the Ozone Layer

Alkanes substituted with both fluorine (F) and chlorine (Cl) atoms have been used as the dispersing gases in aerosol cans, as foaming agents for plastics, and as refrigerants. Two of the best known of these chlorofluorocarbons (CFCs) are listed in Table 14.12.1

Chlorofluorocarbons contribute to the greenhouse effect in the lower atmosphere. They also diffuse into the stratosphere, where they are broken down by ultraviolet (UV) radiation to release Cl atoms. These in turn break down the ozone (O_3) molecules that protect Earth from harmful UV radiation. Worldwide action has reduced the use of CFCs and related compounds. The CFCs and other Cl- or bromine (Br)-containing ozone-destroying compounds are being replaced with more benign substances. Hydrofluorocarbons (HFCs), such as CH_2FCF_3, which have no Cl or Br to form radicals, are one alternative. Another is hydrochlorofluorocarbons (HCFCs), such as $CHCl_2CF_3$. HCFC molecules break down more readily in the troposphere, and fewer ozone-destroying molecules reach the stratosphere.




Figure 14.12.1: Ozone in the upper atmosphere shields Earth's surface from UV radiation from the sun, which can cause skin cancer in humans and is also harmful to other animals and to some plants. Ozone "holes" in the upper atmosphere (the gray, pink, and purple areas at the center) are large areas of substantial ozone depletion. They occur mainly over Antarctica from late August through early October and fill in about mid-November. Ozone depletion has also been noted over the Arctic regions. The largest ozone hole ever observed occurred on 24 September 2006. Source: Image courtesy of NASA, Ozone watch (opens in new window) [ozonewatch.gsfc.nasa.gov].

Key Takeaway

- The replacement of an hydrogen atom on an alkane by a halogen atom—F, Cl, Br, or I—forms a halogenated compound.

14.12: Halogenated Hydrocarbons is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.8: Halogenated Hydrocarbons** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.13: Cycloalkanes

Learning Objectives

- To name cycloalkanes given their formulas and write formulas for these compounds given their names.

The hydrocarbons we have encountered so far have been composed of molecules with open-ended chains of carbon atoms. When a chain contains three or more carbon atoms, the atoms can join to form *ring* or *cyclic* structures. The simplest of these cyclic hydrocarbons has the formula C_3H_6. Each carbon atom has two hydrogen atoms attached (Figure 14.13.1) and is called cyclopropane.





Figure 14.13.1: Ball-and-Spring Model of Cyclopropane. The springs are bent to join the carbon atoms.

To Your Health: Cyclopropane as an Anesthetic

With its boiling point of −33°C, cyclopropane is a gas at room temperature. It is also a potent, quick-acting anesthetic with few undesirable side effects in the body. It is no longer used in surgery, however, because it forms explosive mixtures with air at nearly all concentrations.

The cycloalkanes—cyclic hydrocarbons with only single bonds—are named by adding the prefix *cyclo-* to the name of the open-chain compound having the same number of carbon atoms as there are in the ring. Thus the name for the cyclic compound C_4H_8 is cyclobutane. The carbon atoms in cyclic compounds can be represented by *line-angle formulas* that result in regular geometric figures. Keep in mind, however, that each corner of the geometric figure represents a carbon atom plus as many hydrogen atoms as needed to give each carbon atom four bonds.



cyclopropane cyclohexane

Some cyclic compounds have substituent groups attached. Example 14.13.1 interprets the name of a cycloalkane with a single substituent group.


Example 14.13.1

Draw the structure for each compound.


a. cyclopentane
b. methylcyclobutane

Solution

a. The name *cyclopentane* indicates a cyclic (cyclo) alkane with five (pent-) carbon atoms. It can be represented as a pentagon.



- The name *methylcyclobutane* indicates a cyclic alkane with four (but-) carbon atoms in the cyclic part. It can be represented as a square with a CH_3 group attached.



? Exercise 14.13.1

Draw the structure for each compound.

a. cycloheptane
b. ethylcyclohexane

The properties of cyclic hydrocarbons are generally quite similar to those of the corresponding open-chain compounds. So cycloalkanes (with the exception of cyclopropane, which has a highly strained ring) act very much like noncyclic alkanes. Cyclic structures containing five or six carbon atoms, such as cyclopentane and cyclohexane, are particularly stable. We will see later that some carbohydrates (sugars) form five- or six-membered rings in solution.

The cyclopropane ring is strained because the C–C–C angles are 60°, and the preferred (tetrahedral) bond angle is 109.5°. (This strain is readily evident when you try to build a ball-and-stick model of cyclopropane; see Figure 14.13.1) Cyclopentane and cyclohexane rings have little strain because the C–C–C angles are near the preferred angles.

Key Takeaway

- Many organic compounds have cyclic structures.

14.13: Cycloalkanes is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.9: Cycloalkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons

Hydrocarbons are the simplest organic compounds, but they have interesting physiological effects. These effects depend on the size of the hydrocarbon molecules and where on or in the body they are applied. Alkanes of low molar mass—those with from 1 to approximately 10 or so carbon atoms—are gases or light liquids that act as anesthetics. Inhaling (“sniffing”) these hydrocarbons in gasoline or aerosol propellants for their intoxicating effect is a major health problem that can lead to liver, kidney, or brain damage or to immediate death by asphyxiation by excluding oxygen.

Swallowed, liquid alkanes do little harm while in the stomach. In the lungs, however, they cause “chemical” pneumonia by dissolving fatlike molecules from cell membranes in the tiny air sacs (alveoli). The lungs become unable to expel fluids, just as in pneumonia caused by bacteria or viruses. People who swallow gasoline or other liquid alkane mixtures should not be made to vomit, as this would increase the chance of getting alkanes into the lungs. (There is no home-treatment antidote for gasoline poisoning; call a poison control center.)



Figure 14.14.1: Butane is a hydrocarbon with the formula C_4H_{10} and is a gas at room temperature and atmospheric pressure. Under excess pressures, it can be liquidized like the fuel shown in this butane lighters. (CC BY 3.0Unported; Frank Vincentz)

Liquid alkanes with approximately 5–16 carbon atoms per molecule wash away natural skin oils and cause drying and chapping of the skin, while heavier liquid alkanes (those with approximately 17 or more carbon atoms per molecule) act as emollients (skin softeners). Such alkane mixtures as mineral oil and petroleum jelly can be applied as a protective film. Water and aqueous solutions such as urine will not dissolve such a film, which explains why petroleum jelly protects a baby’s tender skin from diaper rash.

14.14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.0: Prelude to Organic Chemistry - Alkanes and Halogenated Hydrocarbons** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.15: Organic Reactions

Learning Outcomes

- Identify and describe substitution, elimination, addition, hydrolysis, and condensation reactions.
- Predict products of each reaction type.
- Predict products of hydration reactions based on Markovnikov's rule.
- Explain why condensation reactions are also called dehydration reactions.
- Define "polymer".

Organic reactions require the breaking of strong covalent bonds, which takes a considerable input of energy. In order for relatively stable organic molecules to react at a reasonable rate, they often must be modified with the use of highly reactive materials or in the presence of a catalyst. In this lesson, you will learn about several general categories of organic reactions.

Substitution Reactions

A **substitution reaction**, which is the same as a single replacement reaction in inorganic reactions, *is a reaction in which one or more atoms in a molecule are replaced with another atom or group of atoms.* Alkyl halides are formed by the substitution of a halogen atom for a hydrogen atom. When methane reacts with chlorine gas, ultraviolet light can act as a catalyst for the reaction.

$$\mathrm{CH_4\,(g) + Cl_2\,(g) \overset{\text{UV light}}{\rightarrow} CH_3Cl\,(g) + HCl\,(g)} \tag{14.15.1}$$

The reaction produces chloromethane and hydrogen chloride. When the mixture is allowed to react for longer periods of time, further substitution reactions may occur.

$$\mathrm{CH_3Cl\,(g) + Cl_2\,(g) \overset{\text{UV light}}{\rightarrow} CH_2Cl_2\,(g) + HCl\,(g)} \tag{14.15.2}$$


The product above is dichloromethane. Further substitution products trichloromethane and tetrachloromethane, commonly called carbon tetrachloride. A mixture of products occurs in the reaction, with the relative amounts dependent upon the time that the reaction is allowed to proceed. Chlorofluorocarbons are produced by reacting chloroalkanes with HF, because the fluorine atom makes a stronger bond to the carbon atom than chlorine does.

$$\mathrm{CCl_4\,(g) + HF\,(g) \overset{SbF_5}{\rightarrow} CCl_3F\,(g) + HCl\,(g)} \tag{14.15.3}$$

The fluorine atom substitutes for a chlorine atom in the reaction.

Elimination Reactions

An **elimination reaction** involves the removal of adjacent atoms from a molecule. This results in the formation of a multiple bond and the release of a small molecule, so they are called elimination reactions. They have the general form



A typical example is the conversion of ethyl chloride to ethylene:

$$CH_3CH_2Cl \rightarrow CH_2 = CH_2 + HCl \tag{14.15.4}$$

Much of the approximately 26 million tons of ethylene produced per year in the United States is used to synthesize plastics, such as polyethylene. In the above reaction, the A–B molecule eliminated is HCl, whose components are eliminated as H^+ from the carbon atom on the left and Cl^- from the carbon on the right. When an acid is produced, as occurs here, the reaction is generally carried out in the presence of a base (such as NaOH) to neutralize the acid. Other elimination reactions will produce H_2, X_2 (where X = halogen), or H_2O. These reactions are often referred to by more descriptive terms such as dehydrogenation (removing hydrogen) or dechlorination (removing chlorine).

Addition Reactions

An **addition reaction** *is a reaction in which an atom or molecule is added to an unsaturated molecule, making a single product.* An addition reaction can often be thought of as adding a molecule across the double bond of an alkene or across the triple bond of an alkyne. Knowing that "ation" means to add, the specific names of these reactions, such as hydrogenation, hydration, or chlorination, should make sense. Note that hydrogenation (adding H_2) and hydration (adding H_2O) are very different processes.

One type of addition reaction is called hydrogenation. **Hydrogenation** *is a reaction that occurs when molecular hydrogen is added to an alkene to produce an alkane* or *hydrogen is added to an alkyne to produce an alkene or alkane.* The reaction is typically performed with the use of a transition metal catalyst. For example, ethene reacts with hydrogen to form ethane.

$$\mathrm{CH_2{=}CH_2\,(g) + H_2\,(g) \overset{Pt}{\rightarrow} CH_3CH_3\,(g)} \tag{14.15.5}$$

Note that the hydrogenation reaction is also a redox reaction. Ethene is reduced, because the oxidation numbers of the carbon atoms change from -2 to -3 as a result of the reaction.


Vegetable oils consist of long carbon chains with carboxyl groups on the end; these molecules are referred to as fatty acids. The carbon chains of the fatty acids in vegetable oils are unsaturated, usually containing multiple double bonds. When hydrogen gas is blown through a sample of the oil, hydrogen atoms add across the double bonds. This conversion changes the substance from a liquid oil into a solid fat. The "hydrogenated" on a food product is an indication that oil (liquid) has been converted into fat (solid) by this process. Margarine is manufactured from unsaturated vegetable oil in this way by hydrogenating some of the double bonds making it a "partially hydrogenated vegetable oil".

Addition reactions are also useful ways to introduce a new functional group into an organic molecule. Alkyl halides can be produced from an alkene by the addition of either the elemental halogen or the hydrogen halide. A monosubstituted alkyl halide can be produced by the addition of a hydrogen halide to an alkene. Shown below is the formation of chloroethane.

$$\mathrm{CH_2{=}CH_2\,(g) + HCl\,(g) \rightarrow CH_3CH_2Cl\,(g)} \tag{14.15.6}$$

When the reactant is the diatomic halogen, the product is a disubstituted alkyl halide as in the addition of bromine to ethene.

$$\mathrm{CH_2{=}CH_2\,(g) + Br_2\,(l) \rightarrow CH_2BrCH_2Br\,(g)} \tag{14.15.7}$$




1,2-dibromoethane

The addition of bromine to an unknown organic compound can be used as a test for unsaturation in the compound. Bromine has a distinctive brownish-orange color, while most bromoalkanes are colorless. When bromine is slowly added to a solution of the compound, the orange color will fade if it undergoes an addition reaction to produce an alkyl halide. If the orange color remains, then the original compound was already saturated, and no reaction occurred.

A **hydration reaction** *is a reaction in which water is added to an alkene*. Hydration reactions can take place when the alkene and water are heated to near $100^{\circ}\mathrm{C}$ in the presence of a strong acid, which acts as a catalyst. Shown below is the hydration of ethene to produce ethanol.


$$\mathrm{CH_2{=}CH_2\,(g) + H_2O\,(l) \rightarrow CH_3CH_2OH\,(g)} \tag{14.15.8}$$

Unlike addition reactions involving H_2 or Br_2, the addition of water can have two possible products because an $-H$ and an $-OH$ are being added to the carbons in the double bond. **Markovnikov's rule** helps predict the main product in a hydration reaction. The rule states that the hydrogen atom from water will add to the carbon that originally had more hydrogen atoms. For example, look at the hydration of 1-butene.



Note that the first carbon in 1-butene started with two hydrogen atoms and the second carbon started with one hydrogen. Therefore, the hydrogen from water adds to the first carbon and the $-OH$ group adds to the second carbon. When there are equal numbers of hydrogen atoms on both carbons in a double bond then the two products will form in approximately equal amounts. For example,

the hydration of 2-pentene results in two products. In the first product, the $-OH$ group is on the third carbon and in the second product, the $-OH$ group is on the second carbon. While these two molecules will have similar properties, there will be differences.



Condensation Reactions

A **condensation reaction** *is a reaction in which two molecules combine to form a single molecule.* A small molecule, often water, is usually removed during a condensation reaction. Amino acids are important biological molecules that have an amine functional group on one end of the molecule and a carboxylic acid functional group on the other end. When two amino acids combine in a condensation reaction, a covalent bond forms between the amine nitrogen of one amino acid and the carboxyl carbon of the second amino acid. A molecule of water is then removed as a second product (see figure below).





Figure 14.15.1: Amino acids join together to form a molecule called a dipeptide. The $-OH$ from the carboxyl group of one amino acid combines with a hydrogen atom from the amine group of the other amino group to produce water (blue).

This reaction forms a molecule called a dipeptide, and the resulting carbon-nitrogen covalent bond is often called a peptide bond. When repeated numerous times, a long molecule called a protein is eventually produced.

An **esterification** *is a condensation reaction in which an ester is formed from an alcohol and a carboxylic acid.* Esterification is a subcategory of condensation reactions because a water molecule is produced in the reaction. The reaction can be catalyzed by a strong acid, usually sulfuric acid. When the carboxylic acid, butanoic acid, is heated with an excess of methanol and a few drops of sulfuric acid, the ester methyl butanoate is produced. Methyl butanoate has the scent of pineapples. The reaction is shown below with both molecular and structural formulas. Esterification reactions are reversible.

$$CH_3CH_2CH_2COOH + HOCH_3 \overset{H_2SO_4}{\rightarrow} CH_3CH_2CH_2COOCH_3 + H_2O \tag{14.15.9}$$




Saponification *describes the alkaline hydrolysis reaction of an ester*. The term saponification originally described the hydrolysis of long-chain esters called fatty acid esters to produce soap molecules, which are the salts of fatty acids. One such soap molecule is sodium stearate, formed from the hydrolysis of ethyl stearate.

$$\underset{\text{ethyl stearate}}{C_{17}H_{35}COOC_2H_5} + NaOH \rightarrow \underset{\text{sodium stearate (soap)}}{C_{17}H_{35}COO{-}Na^+} + C_2H_5OH \tag{14.15.10}$$

Hydrolysis Reactions

Hydrolysis is the reverse of condensation. "Hydro" indicates that water is involved and "lysis" means to break apart. In a hydrolysis reaction, water is added and the molecule breaks apart, usually at a $C{-}O{-}C$ linkage.



Polymerization

Polymers are very different than the other kinds of organic molecules that you have seen so far. Whereas other compounds are of relatively low molar mass, polymers are giant molecules of very high molar mass. Polymers are the primary components of all sorts of plastics and related compounds. A **polymer** *is a large molecule formed of many smaller molecules covalently bonded to one another in a repeating pattern*. The *small molecules that make up the polymer are called* **monomers**. Polymers are generally formed by either addition or condensation reactions. Teflon (see figure below) is a non-reactive, non-stick coating used on cookware as well as in containers and pipes for reactive or corrosive chemicals.




Figure 14.15.2: Polytetrafluoroethylene (also known as Teflon) is formed from the polymerization of tetrafluoroethylene.

This page titled 14.15: Organic Reactions is shared under a not declared license and was authored, remixed, and/or curated by Allison Soult.

14.E: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Exercises)

12.1: Organic Chemistry

Concept Review Exercises

1. Classify each compound as organic or inorganic.
    a. C_3H_8O
    b. $CaCl_2$
    c. $Cr(NH_3)_3Cl_3$
    d. $C_{30}H_{48}O_3N$
2. Which compound is likely organic and which is likely inorganic?
    a. a flammable compound that boils at 80°C and is insoluble in water
    b. a compound that does not burn, melts at 630°C, and is soluble in water

Answers

1.
    a. organic
    b. inorganic
    c. inorganic
    d. organic
2.
    a. organic
    b. inorganic

1. Classify each compound as organic or inorganic.
    a. C_6H_{10}
    b. $CoCl_2$
    c. $C_{12}H_{22}O_{11}$
2. Classify each compound as organic or inorganic.
    a. CH_3NH_2
    b. $NaNH_2$
    c. $Cu(NH_3)_6Cl_2$
3. Which member of each pair has a higher melting point?
    a. CH_3OH and NaOH
    b. CH_3Cl and KCl
4. Which member of each pair has a higher melting point?
    a. C_2H_6 and $CoCl_2$
    b. CH_4 and LiH

Answers

1.
    a. organic
    b. inorganic
    c. organic

3.

a. NaOH
b. KCl

12.2: Structures and Names of Alkanes

Concept Review Exercises

1. In the homologous series of alkanes, what is the molecular formula for the member just above C_8H_{18}?
2. Use the general formula for alkanes to write the molecular formula of the alkane with 12 carbon atoms.

Answers

1. C_9H_{20}
2. $C_{12}H_{26}$

Exercises

1. What compounds contain fewer carbon atoms than C_3H_8 and are its homologs?
2. What compounds contain five to eight carbon atoms and are homologs of C_4H_{10}?

Answer

1. CH_4 and C_2H_6

12.3: Branched-Chain Alkanes

Concept Review Exercises

1. In alkanes, can there be a two-carbon branch off the second carbon atom of a four-carbon chain? Explain.
2. A student is asked to write structural formulas for two different hydrocarbons having the molecular formula C_5H_{12}. She writes one formula with all five carbon atoms in a horizontal line and the other with four carbon atoms in a line, with a CH_3 group extending down from the first attached to the third carbon atom. Do these structural formulas represent different molecular formulas? Explain why or why not.

Answers

1. No; the branch would make the longest continuous chain of five carbon atoms.
2. No; both are five-carbon continuous chains.


Key Takeaway


- Alkanes with four or more carbon atoms can exist in isomeric forms.

Exercises

1. Briefly identify the important distinctions between a straight-chain alkane and a branched-chain alkane.
2. How are butane and isobutane related? How do they differ?
3. Name each compound.

a.




b. 

4. Write the structural formula for each compound.


   a. hexane
   b. octane

5. Indicate whether the structures in each set represent the same compound or isomers.

   a. $CH_3CH_2CH_2CH_3$ and



   b. $CH_3CH_2CH_2CH_2CH_3$ and



Answers

1. Straight-chain alkanes and branched-chain alkanes have different properties as well as different structures.

3.

   a. pentane
   b. heptane


5.


   a. no
   b. yes

12.4: Condensed Structural and Line-Angle Formulas


Exercises

1. Write the condensed structural formula for each structural formula.


   a. 

   b. 


https://chem.libretexts.org/@go/page/218585

c. 

2. A condensed structural formula for isohexane can be written as $(CH_3)_2CHCH_2CH_2CH_3$. Draw the line-angle formula for isohexane.
3. Draw a line-angle formula for the compound $CH_3CH_2CH(CH_3)CH_2CH_2CH_3$.
4. Give the structural formula for the compound represented by this line-angle formula:



Answers

1.
   a. CH_3CH_3
   b. $CH_3CH_2CH_3$
   c. $CH_3CH_2CH_2CH_2CH_3$
3. 

12.5: IUPAC Nomenclature

Concept Review Exercises

1. What is a CH_3 group called when it is attached to a chain of carbon atoms—a substituent or a functional group?
2. Which type of name uses numbers to locate substituents—common names or IUPAC names?

Answers


1. substituent
2. IUPAC names


Exercises

1. Briefly identify the important distinctions between an alkane and an alkyl group.
2. How many carbon atoms are present in each molecule?
   a. 2-methylbutane
   b. 3-ethylpentane
3. How many carbon atoms are present in each molecule?
   a. 2,3-dimethylbutane
   b. 3-ethyl-2-methylheptane
4. Draw the structure for each compound.
   a. 3-methylpentane
   b. 2,2,5-trimethylhexane
   c. 4-ethyl-3-methyloctane


5. Draw the structure for each compound.
   a. 2-methylpentane
   b. 4-ethyl-2-methylhexane
   c. 2,2,3,3-tetramethylbutane


6. Name each compound according to the IUPAC system.

a. 

b. 

7. Name each compound according to the IUPAC system.

a. 

b. 

8. What is a substituent? How is the location of a substituent indicated in the IUPAC system?


9. Briefly identify the important distinctions between a common name and an IUPAC name.


Answers

1. An alkane is a molecule; an alkyl group is not an independent molecule but rather a part of a molecule that we consider as a unit.


3.
   a. 6
   b. 10

5.

a. 

b. 

https://chem.libretexts.org/@go/page/218585

c. 

7.

a. 2,2,4,4-tetramethylpentane
b. 3-ethylhexane

9. Common names are widely used but not very systematic; IUPAC names identify a parent compound and name other groups as substituents.

12.6: Physical Properties of Alkanes

Concept Review Exercises

1. Without referring to a table, predict which has a higher boiling point—hexane or octane. Explain.
2. If 25 mL of hexane were added to 100 mL of water in a beaker, which of the following would you expect to happen? Explain.
   a. Hexane would dissolve in water.
   b. Hexane would not dissolve in water and would float on top.
   c. Hexane would not dissolve in water and would sink to the bottom of the container.

Answers

1. octane because of its greater molar mass
2. b; hexane is insoluble in water and less dense than water.

Exercises

1. Without referring to a table or other reference, predict which member of each pair has the higher boiling point.
   a. pentane or butane
   b. heptane or nonane
2. For which member of each pair is hexane a good solvent?
   a. pentane or water
   b. sodium chloride or soybean oil

Answer

1. a. pentane
   b. nonane

12.7: Chemical Properties of Alkanes

Concept Review Exercises

1. Why are alkanes sometimes called paraffins?
2. Which halogen reacts most readily with alkanes? Which reacts least readily?

Answers

1. Alkanes do not react with many common chemicals. They are sometimes called paraffins, from the Latin parum affinis, meaning “little affinity.”
2. most readily: F_2; least readily: I_2

Exercises

1. Why do alkanes usually not react with ionic compounds such as most laboratory acids, bases, oxidizing agents, or reducing agents?
2. Write an equation for the complete combustion of methane (CH_4), the main component of natural gas).
3. What is the most important reaction of alkanes?
4. Name some substances other than oxygen that react readily with alkanes.

Answers

1. Alkanes are nonpolar; they do not attract ions.

12.8: Halogenated Hydrocarbons

Concept Review Exercises

1. What is the IUPAC name for the HFC that has the formula CH_2FCF_3? (Hint: you must use a number to indicate the location of each substituent F atom.)
2. What is the IUPAC name for the HCFC that has the formula $CHCl_2CF_3$?

Answers

1. 1,1,1,2-tetrafluoroethane
2. 1,1,1-trifluoro-2,2-dichloroethane

Exercises

1. Write the condensed structural formula for each compound.
   a. methyl chloride
   b. chloroform
2. Write the condensed structural formula for each compound.
   a. ethyl bromide
   b. carbon tetrachloride
3. Write the condensed structural formulas for the two isomers that have the molecular formula C_3H_7Br. Give the common name and the IUPAC name of each.
4. Write the condensed structural formulas for the four isomers that have the molecular formula C_4H_9Br. Give the IUPAC name of each.
5. What is a CFC? How are CFCs involved in the destruction of the ozone layer?
6. Explain why each compound is less destructive to the ozone layer than are CFCs.
   a. fluorocarbons
   b. HCFCs


Answers

1. a. CH_3Cl
   b. $CHCl_3$
3. $CH_3CH_2CH_2Br$, propyl bromide, 1-bromopropane; $CH_3CHBrCH_3$, isopropyl bromide, 2-bromopropane
5. compounds containing Cl, F, and C; by releasing Cl atoms in the stratosphere

12.9: Cycloalkanes

Concept Review Exercises

1. What is the molecular formula of cyclooctane?
2. What is the IUPAC name for this compound?





Answers

1. C_8H_{16}
2. ethylcyclopropane


Exercises

1. Draw the structure for each compound.
   a. ethylcyclobutane
   b. propylcyclopropane
2. Draw the structure for each compound.
   a. methylcyclohexane
   b. butylcyclobutane
3. Cycloalkyl groups can be derived from cycloalkanes in the same way that alkyl groups are derived from alkanes. These groups are named as cyclopropyl, cyclobutyl, and so on. Name each cycloalkyl halide.

   a. 


   b. 


4. Halogenated cycloalkanes can be named by the IUPAC system. As with alkyl derivatives, monosubstituted derivatives need no number to indicate the position of the halogen. To name disubstituted derivatives, the carbon atoms are numbered starting at the position of one substituent (C1) and proceeding to the second substituted atom by the shortest route. Name each compound.

   a. 

   b. 

Answers

1. a. 


   b. 


3. a. cyclopentyl bromide
   b. cyclohexyl chloride


12.10: Chapter Summary

Additional Exercises

1. You find an unlabeled jar containing a solid that melts at 48°C. It ignites readily and burns readily. The substance is insoluble in water and floats on the surface. Is the substance likely to be organic or inorganic?
2. Give the molecular formulas for methylcyclopentane, 2-methylpentane, and cyclohexane. Which are isomers?
3. What is wrong with each name? (Hint: first write the structure *as if* it were correct.) Give the correct name for each compound.
   a. 2-dimethylpropane
   b. 2,3,3-trimethylbutane
   c. 2,4-diethylpentane
   d. 3,4-dimethyl-5-propylhexane
4. What is the danger in swallowing a liquid alkane?
5. Distinguish between lighter and heavier liquid alkanes in terms of their effects on the skin.
6. Following is the line formula for an alkane. Draw its structure and give its name.



7. Write equations for the complete combustion of each compound.
   a. propane (a bottled gas fuel)
   b. octane (a typical hydrocarbon in gasoline).
8. The density of a gasoline sample is 0.690 g/mL. On the basis of the complete combustion of octane, calculate the amount in grams of carbon dioxide (CO_2) and water (H_2O) formed per gallon (3.78 L) of the gasoline when used in an automobile.
9. Draw the structures for the five isomeric hexanes (C_6H_{14}). Name each by the IUPAC system.
10. Indicate whether the structures in each set represent the same compound or isomers.
    a. 
    b. 
    c. 
11. Consider the line-angle formulas shown here and answer the questions.



a. Which pair of formulas represents isomers? Draw each structure.
b. Which formula represents an alkyl halide? Name the compound and write its condensed structural formula.
c. Which formula represents a cyclic alkane? Name the compound and draw its structure.
d. What is the molecular formula of the compound represented by (i)?

Answers

1. organic

3.


a. Two numbers are needed to indicate two substituents; 2,2-dimethylpropane.
b. The lowest possible numbers were not used; 2,2,3-trimethylbutane.
c. An ethyl substituent is not possible on the second carbon atom; 3,5-dimethylheptane.
d. A propyl substituent is not possible on the fifth carbon atom; 3,4,5-trimethyloctane.

5. Lighter alkanes wash away protective skin oils; heavier alkanes form a protective layer.

7.


a. $C_3H_8 + 5O_2 \rightarrow 3CO_2 + 4H_2O$
b. $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$

9. $CH_3CH_2CH_2CH_2CH_2CH_3$; hexane




11.

a. ii and iii; $CH_3CH_2CH_2CH_2CH_2CH_2CH_3$ and



b. iv; 3-chloropentane; $CH_3CH_2CHClCH_2CH_3$

c. i; ethylcyclopentane;



d. C_7H_{14}

14.E: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Exercises) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.6: Physical Properties of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **12.7: Chemical Properties of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **12.8: Halogenated Hydrocarbons** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **12.9: Cycloalkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the summary and ask yourself how they relate to the topics in the chapter.

Organic chemistry is the chemistry of carbon compounds, and **inorganic chemistry** is the chemistry of all the other elements. Carbon atoms can form stable covalent bonds with other carbon atoms and with atoms of other elements, and this property allows the formation the tens of millions of organic compounds. **Hydrocarbons** contain only hydrogen and carbon atoms.

Hydrocarbons in which each carbon atom is bonded to four other atoms are called **alkanes** or **saturated hydrocarbons**. They have the general formula C_nH_{2n+2}. Any given alkane differs from the next one in a series by a CH_2 unit. Any family of compounds in which adjacent members differ from each other by a definite factor is called a **homologous series**.

Carbon atoms in alkanes can form straight chains or branched chains. Two or more compounds having the same molecular formula but different structural formulas are **isomers** of each other. There are no isomeric forms for the three smallest alkanes; beginning with C_4H_{10}, all other alkanes have isomeric forms.

A **structural formula** shows all the carbon and hydrogen atoms and how they are attached to one another. A **condensed structural formula** shows the hydrogen atoms right next to the carbon atoms to which they are attached. A **line-angle formula** is a formula in which carbon atoms are implied at the corners and ends of lines. Each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds.

The **IUPAC System of Nomenclature** provides rules for naming organic compounds. An **alkyl group** is a unit formed by removing one hydrogen atom from an alkane.

The physical properties of alkanes reflect the fact that alkane molecules are nonpolar. Alkanes are insoluble in water and less dense than water.

Alkanes are generally unreactive toward laboratory acids, bases, oxidizing agents, and reducing agents. They do burn (undergo **combustion reactions**).

Alkanes react with halogens by substituting one or more halogen atoms for hydrogen atoms to form **halogenated hydrocarbons**. An **alkyl halide (haloalkane)** is a compound resulting from the replacement of a hydrogen atom of an alkane with a halogen atom.

Cycloalkanes are hydrocarbons whose molecules are closed rings rather than straight or branched chains. A **cyclic hydrocarbon** is a hydrocarbon with a ring of carbon atoms

---

14.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **12.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.S: Unsaturated and Aromatic Hydrocarbons (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Any hydrocarbon containing either a double or triple bond is an **unsaturated hydrocarbon**. **Alkenes** have a carbon-to-carbon double bond. The general formula for alkenes with one double bond is C_nH_{2n}. Alkenes can be straight chain, branched chain, or cyclic. Simple alkenes often have common names, but all alkenes can be named by the system of the International Union of Pure and Applied Chemistry.

Cis-trans isomers (or geometric isomers) are characterized by molecules that differ only in their configuration around a rigid part of the structure, such as a carbon–to-carbon double bond or a ring. The molecule having two identical (or closely related) atoms or groups on the same side is the **cis isomer**; the one having the two groups on opposite sides is the **trans isomer**.

The physical properties of alkenes are quite similar to those of alkanes. Like other hydrocarbons, alkenes are insoluble in water but soluble in organic solvents.

More reactive than alkanes, alkenes undergo **addition reactions** across the double bond:

- Addition of hydrogen (**hydrogenation**):

$$CH_2{=}CH_2 + H_2 \rightarrow CH_3CH_3$$

- Addition of halogen (**halogenation**):

$$CH_2{=}CH_2 + X_2 \rightarrow XCH_2CH_2X$$

  where X = F, Cl, Br, or I.

- Addition of water (**hydration**):

$$CH_2{=}CH_2 + HOH \rightarrow HCH_2CH_2OH$$

Alkenes also undergo **addition polymerization**, molecules joining together to form long-chain molecules.

$$\ldots CH_2{=}CH_2 + CH_2{=}CH_2 + CH_2{=}CH_2 + \ldots \rightarrow \ldots CH_2CH_2{-}CH_2CH_2{-}CH_2CH_2{-}\ldots$$

The reactant units are **monomers**, and the product is a **polymer**.

Alkynes have a carbon-to-carbon triple bond. The general formula for alkynes is C_nH_{2n-2}. The properties of alkynes are quite similar to those of alkenes. They are named much like alkenes but with the ending *-yne*.

The cyclic hydrocarbon *benzene* (C_6H_6) has a ring of carbon atoms. The molecule seems to be unsaturated, but it does not undergo the typical reactions expected of alkenes. The electrons that might be fixed in three double bonds are instead *delocalized* over all six carbon atoms.

A hydrocarbon containing one or more benzene rings (or other similarly stable electron arrangements) is an **aromatic hydrocarbon**, and any related substance is an **aromatic compound**. One or more of the hydrogen atoms on a benzene ring can be replaced by other atoms. When two hydrogen atoms are replaced, the product name is based on the relative position of the replacement atoms (or atom groups). A 1,2-disubstituted benzene is designated as an *ortho* (*o*-) isomer; 1,3-, a *meta* (*m*-) isomer; and 1,4-, a *para* (*p*-) isomer. An aromatic group as a substituent is called an **aryl** group.

A **polycyclic aromatic hydrocarbon (PAH)** has fused benzene rings sharing a common side.

14.S: Unsaturated and Aromatic Hydrocarbons (Summary) is shared under a CC BY license and was authored, remixed, and/or curated by LibreTexts.

- **13.S: Unsaturated and Aromatic Hydrocarbons (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

Index

A

addition reaction
14.15: Organic Reactions
anion
4.10: Ions - Losing and Gaining Electrons
atomic theory
4.4: Atomic Theory
atoms
4.4: Atomic Theory

B

Balancing a Chemical Equation
7.5: How to Write Balanced Chemical Equations
barometer
12.2: Kinetic-Molecular Theory

C

cation
4.10: Ions - Losing and Gaining Electrons
chemical change
3.6: Changes in Matter - Physical and Chemical Changes
chemical property
3.5: Differences in Matter- Physical and Chemical Properties
12.1: A Molecular Comparison of Gases, Liquids, and Solids
Coefficients and Subscripts
7.5: How to Write Balanced Chemical Equations
collision theory
10.1: The Rate of a Chemical Reaction
complete ionic equation
7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
compound
3.4: Classifying Matter According to Its Composition
condensation reaction
14.15: Organic Reactions
condensed phase
12.1: A Molecular Comparison of Gases, Liquids, and Solids
conservation of mass
3.7: Conservation of Mass - There is No New Matter
covalent bond
12.1: A Molecular Comparison of Gases, Liquids, and Solids

D

Dalton's Law of Partial Pressure
12.5: Dalton's Law (Law of Partial Pressures)
dipole force
12.7: Intermolecular forces

E

electron
4.4: Atomic Theory
element
3.4: Classifying Matter According to Its Composition
elimination reaction
14.15: Organic Reactions

G

gas constant
12.3: The Ideal Gas Equation

H

hydrolysis
14.15: Organic Reactions
hypothesis
1.3: Hypothesis, Theories, and Laws

I

ideal gas law
12.3: The Ideal Gas Equation
intermolecular force
12.1: A Molecular Comparison of Gases, Liquids, and Solids
intermolecular forces
12.7: Intermolecular forces
ions
4.10: Ions - Losing and Gaining Electrons

K

kinetic energy
12.1: A Molecular Comparison of Gases, Liquids, and Solids
kinetic molecular theory
12.2: Kinetic-Molecular Theory
12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases

L

law
1.3: Hypothesis, Theories, and Laws
law of conservation of mass
7.2: Laws of Conservation : A Review

M

matter
3.2: What is Matter?
4.4: Atomic Theory
mixture
3.4: Classifying Matter According to Its Composition

N

net ionic equation
7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
neutron
4.4: Atomic Theory

O

oxidation
7.10: Oxidation-Reduction Reactions
oxidation numbers
7.10: Oxidation-Reduction Reactions

P

physical change
3.6: Changes in Matter - Physical and Chemical Changes
physical property
3.5: Differences in Matter- Physical and Chemical Properties
12.1: A Molecular Comparison of Gases, Liquids, and Solids
polymerization
14.15: Organic Reactions
precipitation reaction
7.7: Precipitation Reactions
proton
4.4: Atomic Theory

R

redox
7.10: Oxidation-Reduction Reactions
redox reaction
7.10: Oxidation-Reduction Reactions
redox reactions
7.10: Oxidation-Reduction Reactions
reduction
7.10: Oxidation-Reduction Reactions
resonance
6.5: Resonance - Equivalent Lewis Structures for the Same Molecule
Ringer’s lactate
9.1: Prelude to Solutions
root mean square velocity
12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases

S

Separation of Mixtures
3.6: Changes in Matter - Physical and Chemical Changes
significant figures
2.4: Significant Figures in Calculations
spectator ions
7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
Stock system
5.7: Naming Ionic Compounds
substitution reaction
14.15: Organic Reactions

T

theory
1.3: Hypothesis, Theories, and Laws

Index

A

addition reaction
- 14.15: Organic Reactions

anion
- 4.10: Ions - Losing and Gaining Electrons

atomic theory
- 4.4: Atomic Theory

atoms
- 4.4: Atomic Theory

B

Balancing a Chemical Equation
- 7.5: How to Write Balanced Chemical Equations

barometer
- 12.2: Kinetic-Molecular Theory

C

cation
- 4.10: Ions - Losing and Gaining Electrons

chemical change
- 3.6: Changes in Matter - Physical and Chemical Changes

chemical property
- 3.5: Differences in Matter- Physical and Chemical Properties
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

Coefficients and Subscripts
- 7.5: How to Write Balanced Chemical Equations

collision theory
- 10.1: The Rate of a Chemical Reaction

complete ionic equation
- 7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

compound
- 3.4: Classifying Matter According to Its Composition

condensation reaction
- 14.15: Organic Reactions

condensed phase
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

conservation of mass
- 3.7: Conservation of Mass - There is No New Matter

covalent bond
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

D

Dalton's Law of Partial Pressure
- 12.5: Dalton's Law (Law of Partial Pressures)

dipole force
- 12.7: Intermolecular forces

E

electron
- 4.4: Atomic Theory

element
- 3.4: Classifying Matter According to Its Composition

elimination reaction
- 14.15: Organic Reactions

G

gas constant
- 12.3: The Ideal Gas Equation

H

hydrolysis
- 14.15: Organic Reactions

hypothesis
- 1.3: Hypothesis, Theories, and Laws

I

ideal gas law
- 12.3: The Ideal Gas Equation

intermolecular force
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

intermolecular forces
- 12.7: Intermolecular forces

ions
- 4.10: Ions - Losing and Gaining Electrons

K

kinetic energy
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

kinetic molecular theory
- 12.2: Kinetic-Molecular Theory
- 12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases

L

law
- 1.3: Hypothesis, Theories, and Laws

law of conservation of mass
- 7.2: Laws of Conservation : A Review

M

matter
- 3.2: What is Matter?
- 4.4: Atomic Theory

mixture
- 3.4: Classifying Matter According to Its Composition

N

net ionic equation
- 7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

neutron
- 4.4: Atomic Theory

O

oxidation
- 7.10: Oxidation-Reduction Reactions

oxidation numbers
- 7.10: Oxidation-Reduction Reactions

P

physical change
- 3.6: Changes in Matter - Physical and Chemical Changes

physical property
- 3.5: Differences in Matter- Physical and Chemical Properties
- 12.1: A Molecular Comparison of Gases, Liquids, and Solids

polymerization
- 14.15: Organic Reactions

precipitation reaction
- 7.7: Precipitation Reactions

proton
- 4.4: Atomic Theory

R

redox
- 7.10: Oxidation-Reduction Reactions

redox reaction
- 7.10: Oxidation-Reduction Reactions

redox reactions
- 7.10: Oxidation-Reduction Reactions

reduction
- 7.10: Oxidation-Reduction Reactions

resonance
- 6.5: Resonance - Equivalent Lewis Structures for the Same Molecule

Ringer’s lactate
- 9.1: Prelude to Solutions

root mean square velocity
- 12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases

S

Separation of Mixtures
- 3.6: Changes in Matter - Physical and Chemical Changes

significant figures
- 2.4: Significant Figures in Calculations

spectator ions
- 7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

Stock system
- 5.7: Naming Ionic Compounds

substitution reaction
- 14.15: Organic Reactions

T

theory
- 1.3: Hypothesis, Theories, and Laws

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: CHE 125: GOB Introductory Chemistry

Webpages: 153

All licenses found:

- CC BY 4.0: 71.2% (109 pages)
- Undeclared: 11.8% (18 pages)
- CC BY-SA 4.0: 9.2% (14 pages)
- Public Domain: 7.8% (12 pages)

By Page

- CHE 125: GOB Introductory Chemistry - *CC BY 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Chemistry as a Science - *CC BY-SA 4.0*
    - 1.1: The Scope of Chemistry - *CC BY-SA 4.0*
    - 1.2: Chemicals Compose Ordinary Things - *CC BY-SA 4.0*
    - 1.3: Hypothesis, Theories, and Laws - *CC BY-SA 4.0*
    - 1.4: The Scientific Method - How Chemists Think - *CC BY-SA 4.0*
  - 2: Measurements and Density - *CC BY 4.0*
    - 2.1: Taking Measurements - *CC BY 4.0*
    - 2.2: Scientific Notation - Writing Large and Small Numbers - *CC BY 4.0*
    - 2.3: Significant Figures - Writing Numbers to Reflect Precision - *CC BY 4.0*
    - 2.4: Significant Figures in Calculations - *CC BY 4.0*
    - 2.5: The Basic Units of Measurement - *CC BY 4.0*
    - 2.6: Problem Solving and Unit Conversions - *CC BY 4.0*
    - 2.7: Solving Multistep Conversion Problems - *CC BY 4.0*
    - 2.8: Units Raised to a Power - *CC BY 4.0*
    - 2.9: Density - *CC BY 4.0*
    - 2.10: Dosage Calculations - *CC BY 4.0*
  - 3: Matter - *CC BY-SA 4.0*
    - 3.1: In Your Room - *CC BY-SA 4.0*
    - 3.2: What is Matter? - *CC BY-SA 4.0*
    - 3.3: Classifying Matter According to Its State- Solid, Liquid, and Gas - *CC BY-SA 4.0*
    - 3.4: Classifying Matter According to Its Composition - *CC BY-SA 4.0*
    - 3.5: Differences in Matter- Physical and Chemical Properties - *CC BY-SA 4.0*
    - 3.6: Changes in Matter - Physical and Chemical Changes - *CC BY-SA 4.0*
    - 3.7: Conservation of Mass - There is No New Matter - *CC BY-SA 4.0*
    - 3.E: Chemistry, Matter, and Measurement (Exercises) - *CC BY-SA 4.0*
  - 4: Atoms, Elements, and the Periodic Table - *Public Domain*
    - 4.1: Prelude to Elements, Atoms, and the Periodic Table - *Public Domain*
    - 4.2: The Elements - *Public Domain*
    - 4.3: Atomic Theory - *Public Domain*
    - 4.4: Atomic Theory - *Public Domain*
    - 4.5: Defining Isotopes - *Public Domain*
    - 4.6: Atomic Masses - *Public Domain*
    - 4.7: Arrangements of Electrons - *Public Domain*
    - 4.8: Orbital Shape - *Public Domain*
    - 4.9: The Periodic Table - *Public Domain*
    - 4.10: Ions - Losing and Gaining Electrons - *Public Domain*
    - 4.E: Elements, Atoms, and the Periodic Table (Exercises) - *Public Domain*
  - 5: Compounds- Ionic and Covalent - *CC BY 4.0*
    - 5.1: Sugar and Salt - *CC BY 4.0*
    - 5.2: Compounds Display Constant Composition - *CC BY 4.0*
    - 5.3: Chemical Formulas- How to Represent Compounds - *CC BY 4.0*
    - 5.4: A Molecular View of Elements and Compounds - *CC BY 4.0*
    - 5.5: Writing Formulas for Ionic Compounds - *CC BY 4.0*
    - 5.6: Nomenclature- Naming Compounds - *CC BY 4.0*
    - 5.7: Naming Ionic Compounds - *CC BY 4.0*
    - 5.8: Naming Molecular Compounds - *CC BY 4.0*

- 5.9: Naming Acids - *CC BY 4.0*
- 5.10: Nomenclature Summary - *CC BY 4.0*
- 5.11: Formula Mass- The Mass of a Molecule or Formula Unit - *CC BY 4.0*
- 5.12: Moles, Molecular Formulas, and Calculating Empirical Formulas for Compounds - *Undeclared*

6: Chemical Bonding - *CC BY 4.0*

- 6.1: Representing Valence Electrons with Dots - *CC BY 4.0*
- 6.2: Lewis Structures of Ionic Compounds- Electrons Transferred - *CC BY 4.0*
- 6.3: Covalent Lewis Structures- Electrons Shared - *CC BY 4.0*
- 6.4: Writing Lewis Structures for Covalent Compounds - *CC BY 4.0*
- 6.5: Resonance - Equivalent Lewis Structures for the Same Molecule - *CC BY 4.0*
- 6.6: Predicting the Shapes of Molecules - *CC BY 4.0*
- 6.7: Electronegativity and Polarity - Why Oil and Water Don’t Mix - *CC BY 4.0*

7: Chemical Reactions - *CC BY 4.0*

- 7.1: Prelude to Introduction to Chemical Reactions - *CC BY 4.0*
- 7.2: Laws of Conservation : A Review - *CC BY 4.0*
- 7.3: Evidence of a Chemical Reaction - *CC BY 4.0*
- 7.4: Chemical Equations - *CC BY 4.0*
- 7.5: How to Write Balanced Chemical Equations - *CC BY 4.0*
- 7.6: Aqueous Solutions and Solubility - Compounds Dissolved in Water - *CC BY 4.0*
- 7.7: Precipitation Reactions - *CC BY 4.0*
- 7.8: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations - *CC BY 4.0*
- 7.9: Acid–Base and Gas Evolution Reactions - *CC BY 4.0*
- 7.10: Oxidation-Reduction Reactions - *CC BY 4.0*
- 7.11: Classifying Chemical Reactions - *CC BY 4.0*
- 7.12: The Activity Series- Predicting Spontaneous Redox Reactions - *CC BY 4.0*

8: Quantities in Chemical Reactions - *CC BY 4.0*

- 8.1: Prelude to Quantities in Chemical Reactions - *CC BY 4.0*
- 8.2: The Mole - *CC BY 4.0*
- 8.3: Atomic and Molar Masses - *CC BY 4.0*
- 8.4: Mole-Mass Conversions - *CC BY 4.0*
- 8.5: Mole-Mole Relationships in Chemical Reactions - *CC BY 4.0*
- 8.6: Mole-Mass and Mass-Mass Problems - *CC BY 4.0*
- 8.E: Quantities in Chemical Reactions (Exercise) - *CC BY 4.0*
- 8.S: Quantities in Chemical Reactions (Summary) - *CC BY 4.0*

9: Solutions - *CC BY 4.0*

- 9.1: Prelude to Solutions - *CC BY 4.0*
- 9.2: Solutions - *CC BY 4.0*
- 9.3: Concentration - *CC BY 4.0*
- 9.4: The Dissolution Process - *CC BY 4.0*
- 9.5: Properties of Solutions - *CC BY 4.0*
- 9.E: Solutions (Exercises) - *CC BY 4.0*
- 9.S: Solutions (Summary) - *CC BY 4.0*

10: Equilibrium - *CC BY 4.0*

- 10.1: The Rate of a Chemical Reaction - *CC BY 4.0*
- 10.2: The Idea of Dynamic Chemical Equilibrium - *CC BY 4.0*
- 10.3: The Equilibrium Constant - A Measure of How Far a Reaction Goes - *CC BY 4.0*
- 10.4: Disturbing a Reaction at Equilibrium- Le Châtelier’s Principle - *CC BY 4.0*
- 10.5: The Effect of a Concentration Change on Equilibrium - *CC BY 4.0*
- 10.6: The Effect of a Volume Change on Equilibrium - *CC BY 4.0*
- 10.7: The Effect of Temperature Changes on Equilibrium - *CC BY 4.0*
- 10.8: The Path of a Reaction and the Effect of a Catalyst - *CC BY 4.0*

11: Energy and Chemical Processes - *CC BY 4.0*

- 11.1: Prelude to Energy and Chemical Processes - *CC BY 4.0*
- 11.2: Energy and Its Units - *CC BY 4.0*
- 11.3: Heat - *CC BY 4.0*
- 11.4: Phase Changes - *CC BY 4.0*
- 11.5: Bond Energies and Chemical Reactions - *CC BY 4.0*
- 11.6: The Energy of Biochemical Reactions - *CC BY 4.0*
- 11.E: Energy and Chemical Processes (Exercises) - *CC BY 4.0*
- 11.S: Energy and Chemical Processes (Summary) - *CC BY 4.0*

12: Solids, Liquids, and Gases - *CC BY 4.0*

- 12.1: A Molecular Comparison of Gases, Liquids, and Solids - *Undeclared*
- 12.2: Kinetic-Molecular Theory - *Undeclared*
- 12.3: The Ideal Gas Equation - *Undeclared*
- 12.4: The Kinetic-Molecular Theory Explains the Behavior of Gases - *CC BY 4.0*
- 12.5: Dalton's Law (Law of Partial Pressures) - *Undeclared*

- 12.6: Evaporation and Condensation - *Undeclared*
- 12.7: Intermolecular forces - *Undeclared*

- 13: Acids and Bases - *CC BY 4.0*
  - 13.1: Prelude to Acids and Bases - *CC BY 4.0*
  - 13.2: Arrhenius Definition of Acids and Bases - *CC BY 4.0*
  - 13.3: Brønsted-Lowry Definition of Acids and Bases - *CC BY 4.0*
  - 13.4: Water - Both an Acid and a Base - *CC BY 4.0*
  - 13.5: The Strengths of Acids and Bases - *CC BY 4.0*
  - 13.6: Buffers - *CC BY 4.0*
  - 13.E: Acids and Bases (Exercises) - *CC BY 4.0*
  - 13.S: Acids and Bases (Summary) - *CC BY 4.0*
- 14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons - *CC BY 4.0*
  - 14.1: Organic Chemistry - *CC BY 4.0*
  - 14.2: Structures and Names of Alkanes - *CC BY 4.0*
  - 14.3: Alkenes- Structures and Names - *CC BY 4.0*
  - 14.4: Alkynes - *CC BY 4.0*
  - 14.5: Branched-Chain Alkanes - *CC BY 4.0*
  - 14.6: Condensed Structural and Line-Angle Formulas - *CC BY 4.0*
  - 14.7: IUPAC Nomenclature - *CC BY 4.0*
  - 14.8: Physical Properties of Alkanes - *CC BY 4.0*
  - 14.9: Physical Properties of Alkenes - *CC BY 4.0*
  - 14.10: Chemical Properties of Alkanes - *CC BY 4.0*
  - 14.11: Chemical Properties of Alkenes - *CC BY 4.0*
  - 14.12: Halogenated Hydrocarbons - *CC BY 4.0*
  - 14.13: Cycloalkanes - *CC BY 4.0*
  - 14.14: Organic Chemistry - Alkanes and Halogenated Hydrocarbons - *CC BY 4.0*
  - 14.15: Organic Reactions - *Undeclared*
  - 14.E: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Exercises) - *CC BY 4.0*
  - 14.S: Organic Chemistry- Alkanes and Halogenated Hydrocarbons (Summary) - *CC BY 4.0*
  - 14.S: Unsaturated and Aromatic Hydrocarbons (Summary) - *CC BY 4.0*
- Back Matter - *Undeclared*
  - Index - *Undeclared*
  - Index - *Undeclared*
  - Glossary - *Undeclared*
  - Detailed Licensing - *Undeclared*