CHEM 1121: GENERAL, ORGANIC, AND BIOCHEMISTRY



LibreTexts™

CHEM 1121: General, Organic, and Biochemistry

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 01/12/2026

TABLE OF CONTENTS

Licensing

1: Measuring Matter and Energy

- 1.1: What is Chemistry
- 1.2: Classification of Matter
- 1.3: Introduction to Kinetic and Potential Energy
- 1.4: Measurements
- 1.5: Expressing Numbers- Scientific Notation
- 1.6: Expressing Numbers- Significant Figures
- 1.7: The International System of Units
- 1.8: Converting Units
- 1.9: Temperature Conversions
- 1.10: Specific Heat
- 1.11: Dosage Calculations
- 1.12: Chapter Summary

2: Atomic Structure

- 2.1: The Elements
- 2.2: Atomic Theory
- 2.3: The Structure of Atoms
- 2.4: Nuclei of Atoms
- 2.5: Arrangement of Electron (Shell Model)
- 2.6: The Periodic Table
- 2.7: Chapter Summary

3: Nuclear Chemistry

- 3.1: Radioactivity
- 3.2: Half-Life
- 3.3: Units of Radioactivity
- 3.4: Uses of Radioactive Isotopes
- 3.5: Nuclear Energy
- 3.6: 3.6 Chapter Summary
- 3.7: Ionic Bonding and Simple Ionic Compounds

4: Ionic Bonding and Simple Ionic Compounds

- 4.1: Two Types of Bonding
- 4.2: Ions
- 4.3: Formulas for Ionic Compounds
- 4.4: Ionic Nomenclature
- 4.5: Chapter Summary

5: Covalent Bonding and Simple Molecular Compounds

- 5.1: Covalent Bonds
- 5.2: Covalent Compounds- Formulas and Names

- 5.3: Multiple Covalent Bonds
- 5.4: Characteristics of Covalent Bonds
- 5.5: Characteristics of Molecules
- 5.6: Intermolecular Forces
- 5.7: Chapter Summary

6: Introduction to Chemical Reactions

- 6.1: Prelude to Introduction to Chemical Reactions
- 6.2: The Law of Conservation of Matter
- 6.3: Chemical Equations
- 6.4: Quantitative Relationships Based on Chemical Equations
- 6.5: Types of Chemical Reactions
- 6.6: Oxidation-Reduction (Redox) Reactions
- 6.7: Redox Reactions in Organic Chemistry and Biochemistry
- 6.E: Introduction to Chemical Reactions (Exercises)
- 6.S: Introduction to Chemical Reactions (Summary)

7: Quantities in Chemical Reactions

- 7.1: Formula Mass
- 7.2: The Mole
- 7.3: Atomic and Molar Masses
- 7.4: Mole-Mass Conversions
- 7.5: Mole-Mole Relationships in Chemical Reactions
- 7.6: Energetics of Biochemical Reactions
- 7.7: Factors Affecting Reaction Rates
- 7.8: Catalysis
- 7.9: Equilbria and Le Chatelier's Principle
- 7.10: Chapter Summary

8: States of Matter and the Gas Laws

- 8.1: Phase Changes
- 8.2: Solids, Liquids, and Gases
- 8.3: Gases and Pressure
- 8.4: Gas Laws
- 8.5: Chapter Summary

9: Solutions

- 9.1: Introduction to Solutions
- 9.2: Solution Concentration- Molarity
- 9.3: Other Units for Solution Concentrations
- 9.4: The Dissolution Process
- 9.5: Properties of Solutions
- 9.6: Dialysis and Tonicity
- 9.7: Chapter Summary

10: Acids and Bases

- 10.1: Arrhenius Definition of Acids and Bases
- 10.2: Brønsted-Lowry Definition of Acids and Bases

- 10.3: Water is both an Acid and a Base
- 10.4: The Strengths of Acids and Bases
- 10.5: pH and Kw
- 10.6: Buffers
- 10.7: End-of-Chapter Material

11: Organic Chemistry of Hydrocarbons

- 11.1: Introduction to Organic Chemistry
- 11.2: Organic Chemistry
- 11.3: Structures and Names of Alkanes
- 11.4: Branched-Chain Alkanes
- 11.5: Condensed Structural and Line-Angle Formulas
- 11.6: Cycloalkanes
- 11.7: IUPAC Nomenclature
- 11.8: Alkenes- Structure and Nomenclature
- 11.9: Alkenes- Cis-Tran Isomerization
- 11.10: Alkynes
- 11.11: Aromatic Compounds
- 11.12: Aromatic Compounds- Structure and Nomenclature
- 11.13: Physical Properties of Hydrocarbons
- 11.14: Chapter Summary

12: Organic Functional Groups - Structure and Nomenclature

- 12.1: Organic Compounds with Functional Groups
- 12.2: 9.2 Alcohols and Phenols- Nomenclature and Classification
- 12.3: Glycols and Glycerol
- 12.4: Ethers
- 12.5: Aldehydes and Ketones- Structure and Names
- 12.6: Organic Sulfur Compounds
- 12.7: Carboxylic Acids- Structures and Names
- 12.8: Esters- Structures and Names
- 12.9: Phosphate Esters
- 12.10: Amines
- 12.11: Amides

13: Organic Reactions

- 13.1: Organic Reactions
- 13.2: Organic Reactions (Exercises)

14: Properties of Organic Functional Groups

- 14.1: Physical Properties of Alcohols
- 14.2: Physical Properties of Aldehydes and Ketones
- 14.3: Physical Properties of Carboxylic Acids
- 14.4: Physical Properties of Esters
- 14.5: Physical Properties of Amines
- 14.6: Physical Properties of Amides
- 14.7: Ionization of Carboxylic Acids
- 14.8: Amines as Bases

- 14.9: Chirality
- 14.10: End-of-Chapter Material

15: Carbohydrates

- 15.1: Introduction to Carbohydrates
- 15.2: Classes of Monosaccharides
- 15.3: Important Hexoses
- 15.4: Cyclic Structures of Monosaccharides
- 15.5: Properties of Monosaccharides
- 15.6: Disaccharides
- 15.7: Polysaccharides
- 15.8: Diabetes
- 15.9: Chapter Summary

16: Lipids

- 16.1: Introduction to Lipids
- 16.2: Fatty Acids
- 16.3: Fats and Oils
- 16.4: Membranes and Membrane Lipids
- 16.5: Steroids
- 16.6: Chapter Summary

17: Proteins and Enzymes

- 17.1: Properties of Amino Acids
- 17.2: Reactions of Amino Acids
- 17.3: Peptides
- 17.4: Proteins
- 17.5: Enzymes
- 17.6: Enzymatic Catalysis
- 17.7: Enzymatic Activity
- 17.8: Enzyme Inhibition
- 17.9: Enzyme Cofactors and Vitamins
- 17.10: Chapter Summary

18: Nucleic Acids

- 18.1: Nucleotides
- 18.2: Nucleic Acid Structure
- 18.3: Replication and Expression of Genetic Information
- 18.4: Protein Synthesis and the Genetic Code
- 18.5: Mutations and Genetic Diseases
- 18.6: Viruses
- 18.7: Chapter Summary

19: Energy Metabolism

- 19.1: Prelude to Energy Metabolism
- 19.2: ATP- the Universal Energy Currency
- 19.3: Stage I of Catabolism
- 19.4: Overview of Stage II of Catabolism

- 19.5: Stage III of Catabolism
- 19.6: Stage II of Carbohydrate Catabolism
- 19.7: Stage II of Lipid Catabolism
- 19.8: Stage II of Protein Catabolism
- 19.9: Oxidative Phosphorylation
- 19.10: Energy Metabolism (Exercises)
- 19.11: Energy Metabolism (Summary)

Index

Glossary

Detailed Licensing

Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

CHAPTER OVERVIEW

1: Measuring Matter and Energy

The study of chemistry will open your eyes to a fascinating world. Chemical processes are continuously at work all around us. They happen as you cook and eat food, strike a match, shampoo your hair, and even read this page. Chemistry is called the central science because a knowledge of chemical principles is essential for other sciences. You might be surprised at the extent to which chemistry pervades your life.

1.1: What is Chemistry
1.2: Classification of Matter
1.3: Introduction to Kinetic and Potential Energy
1.4: Measurements
1.5: Expressing Numbers- Scientific Notation
1.6: Expressing Numbers- Significant Figures
1.7: The International System of Units
1.8: Converting Units
1.9: Temperature Conversions
1.10: Specific Heat
1.11: Dosage Calculations
1.12: Chapter Summary

1: Measuring Matter and Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

1.1: What is Chemistry

Learning Objectives

1. Define chemistry in relation to other sciences.
2. Identify the general steps in the scientific method.

Chemistry is the study of matter—what it consists of, what its properties are, and how it changes. Being able to describe the ingredients in a cake and how they change when the cake is baked is called chemistry. Matter is anything that has mass and takes up space—that is, anything that is physically real. Some things are easily identified as matter—this book, for example. Others are not so obvious. Because we move so easily through air, we sometimes forget that it, too, is matter.

Chemistry is one branch of science. Science is the process by which we learn about the natural universe by observing, testing, and then generating models that explain our observations. Because the physical universe is so vast, there are many different branches of science (Figure **1.1.1**). Thus, chemistry is the study of matter, biology is the study of living things, and geology is the study of rocks and the earth. Mathematics is the language of science, and we will use it to communicate some of the ideas of chemistry.




Figure **1.1.1**: The Relationships between Some of the Major Branches of Science. Chemistry lies more or less in the middle, which emphasizes its importance to many branches of science.

Although we divide science into different fields, there is much overlap among them. For example, some biologists and chemists work in both fields so much that their work is called biochemistry. Similarly, geology and chemistry overlap in the field called geochemistry. Figure **1.1.1** shows how many of the individual fields of science are related.


> *There are many other fields of science in addition to the ones listed here.*

Alchemy

As our understanding of the universe has changed over time, so has the practice of science. Chemistry in its modern form, based on principles that we consider valid today, was developed in the 1600s and 1700s. Before that, the study of matter was known as *alchemy* and was practiced mainly in China, Arabia, Egypt, and Europe.

Alchemy was a somewhat mystical and secretive approach to learning how to manipulate matter. Practitioners, called alchemists, thought that all matter was composed of different proportions of the four basic elements—fire, water, earth, and air—and believed that if you changed the relative proportions of these elements in a substance, you could change the substance. The long-standing attempts to “transmute” common metals into gold represented one goal of alchemy. Alchemy’s other major

goal was to synthesize the philosopher's stone, a material that could impart long life—even immortality. Alchemists used symbols to represent substances, some of which are shown in the accompanying figure. This was not done to better communicate ideas, as chemists do today, but to maintain the secrecy of alchemical knowledge, keeping others from sharing in it.



The first affinity table. Table of different relations observed in chemistry between different substances; Memoirs of the Royal Academy of Sciences, p. 202-212. Alchemists used symbols like these to represent substances.

In spite of this secrecy, in its time alchemy was respected as a serious, scholarly endeavor. Isaac Newton, the great mathematician and physicist, was also an alchemist.

Exercise 1.1.1

Which fields of study are branches of science? Explain.

a. sculpture
b. astronomy

Answer a

Sculpture is not considered a science because it is not a study of some aspect of the natural universe.

Answer b

Astronomy is the study of stars and planets, which are part of the natural universe. Astronomy is therefore a field of science.

Exercise 1.1.2

Which fields of study are branches of science?

a. physiology (the study of the function of an animal's or a plant's body)
b. geophysics
c. agriculture
d. politics

Answer

a, b and c only

How do scientists work? Generally, they follow a process called the scientific method. The scientific method is an organized procedure for learning answers to questions and making explanations for observations. To find the answer to a question (for example, “Why do birds fly toward Earth’s equator during the cold months?”), a scientist goes through the following steps, which are also illustrated in Figure **1.1.2**:




Figure **1.1.2**: The General Steps of the Scientific Method. After an observation is made or a question is identified, a hypothesis is made and experiments are designed to test the hypothesis.

The steps may not be as clear-cut in real life as described here, but most scientific work follows this general outline.

1. **Propose a hypothesis.** A scientist generates a testable idea, or hypothesis, to try to answer a question or explain an observation about how the natural universe works. Some people use the word *theory* in place of hypothesis, but the word hypothesis is the proper word in science. For scientific applications, the word theory is a general statement that describes a large set of observations and data. A theory represents the highest level of scientific understanding.
2. **Test the hypothesis.** A scientist evaluates the hypothesis by devising and carrying out experiments to test it. If the hypothesis passes the test, it may be a proper answer to the question. If the hypothesis does not pass the test, it may not be a good answer.
3. **Refine the hypothesis if necessary.** Depending on the results of experiments, a scientist may want to modify the hypothesis and then test it again. Sometimes the results show the original hypothesis to be completely wrong, in which case a scientist will have to devise a new hypothesis.

Not all scientific investigations are simple enough to be separated into these three discrete steps. But these steps represent the general method by which scientists learn about our natural universe.

1.1.3 Exercise

Define science and chemistry.

Answer

Science is a process by which we learn about the natural universe by observing, testing, and then generating models that explain our observations. Chemistry is the study of matter.

1.1.4 Exercise

Name the steps of the scientific method.

Answer

After identifying the problem or making an observation, propose a hypothesis, test the hypothesis, and refine the hypothesis, if necessary

Key Takeaways

- Chemistry is the study of matter and how it behaves.
- The scientific method is the general process by which we learn about the natural universe.

1. Based on what you know, which fields are branches of science?

    a. meteorology (the study of weather)
    b. astrophysics (the physics of planets and stars)
    c. economics (the study of money and monetary systems)
    d. astrology (the prediction of human events based on planetary and star positions)
    e. political science (the study of politics)

2. Based on what you know, which fields are a branches of science?

    a. history (the study of past events)
    b. ornithology (the study of birds)
    c. paleontology (the study of fossils)
    d. zoology (the study of animals)
    e. phrenology (using the shape of the head to determine personal characteristics)

3. Which of the following are examples of matter?

    a. a baby
    b. an idea
    c. the Empire State Building
    d. an emotion
    e. the air
    f. Alpha Centauri, the closest known star (excluding the sun) to our solar system

4. Which of the following are examples of matter?

    a. your textbook
    b. brain cells
    c. love
    d. a can of soda
    e. breakfast cereal

5. Suggest a name for the science that studies the physics of rocks and the earth.
6. Suggest a name for the study of the physics of living organisms.
7. Engineering is the practical application of scientific principles and discoveries to develop things that make our lives easier. Is medicine science or engineering? Justify your answer.
8. Based on the definition of engineering in Exercise 7, would building a bridge over a river or road be considered science or engineering? Justify your answer.
9. When someone says, “I have a theory that excess salt causes high blood pressure,” does that person really have a theory? If it is not a theory, what is it?
10. When a person says, “My hypothesis is that excess calcium in the diet causes kidney stones,” what does the person need to do to determine if the hypothesis is correct?
11. Some people argue that many scientists accept many scientific principles on faith. Using what you know about the scientific method, how might you argue against that assertion?
12. Most students take multiple English classes in school. Does the study of English use the scientific method?

Answers

1.
    a. science
    b. science
    c. not science
    d. not science
    e. not science

2.
    a. not science
    b. science
    c. science
    d. science
    e. not science

3.
    a. matter
    b. not matter
    c. matter
    d. not matter
    e. matter
    f. matter

4.
    a. matter
    b. matter
    c. not matter
    d. matter
    e. matter

5. geophysics
6. biophysics
7. Medicine is probably closer to a field of engineering than a field of science, but this may be arguable. Ask your doctor.
8. Engineering
9. In scientific terms, this person has a hypothesis.
10. Conduct experiments to determine if kidney stones contain calcium.
11. Science is based on reproducible facts, not blind belief.
12. No.

1.1: What is Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.1: What is Chemistry?** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.2: Classification of Matter

Learning Objectives

- Use physical and chemical properties, including phase, to describe matter.
- Identify a sample of matter as an element, a compound, or a mixture.

Part of understanding matter is being able to describe it. One way chemists describe matter is to assign different kinds of properties to different categories.

Physical and Chemical Properties

The properties that chemists use to describe matter fall into two general categories. **Physical properties** are characteristics that describe matter. They include characteristics such as size, shape, color, and mass. These characteristics can be observed or measured without changing the *identity* of the matter in question. **Chemical properties** are characteristics that describe how matter changes its chemical structure or composition. An example of a chemical property is flammability—a material's ability to burn—because burning (also known as combustion) changes the chemical composition of a material. The observation of chemical properties involves a *chemical change* of the matter in question, resulting in matter with a different *identity* and different physical and chemical properties.



Figure **1.2.1**: (left) Ice Melting is a physical change. When liquid water (H_2O) freezes into a solid state (ice), it appears changed; However, this change is only physical as the the composition of the constituent molecules is the same: 11.19% hydrogen and 88.81% oxygen by mass. (right) Burning of wax to generate water and carbon dioxide is a chemical reaction. (CC-SA-BY-3.0; Andrikkos)

Elements and Compounds

Any sample of matter that has the *same physical and chemical properties throughout* the sample is called a **substance**. There are two types of substances. A substance that cannot be broken down into chemically simpler components is called an **element**. Aluminum, which is used in soda cans and is represented by the symbol Al, is an element. A substance that can be broken down into chemically simpler components (because it consists of more than one element) is called a **compound**. Water is a compound composed of the elements hydrogen and oxygen and is described by the chemical formula, H_2O. Today, there are about 118 elements in the known universe. In contrast, scientists have identified tens of millions of different compounds to date.

> Sometimes the word *pure* is used to describe a substance, but this is not absolutely necessary. By definition, any single substance, element or compound is *pure*.

The smallest part of an element that maintains the identity of that element is called an **atom**. Atoms are extremely tiny; to make a line of iron atoms that is 1 inch long, you would need approximately 217 million iron atoms. The smallest part of a compound that maintains the identity of that compound is called a **molecule**. Molecules are composed of two or more atoms that are attached

together and behave as a unit. Scientists usually work with millions and millions of atoms and molecules at a time. When a scientist is working with large numbers of atoms or molecules at a time, the scientist is studying the *macroscopic viewpoint* of the universe. However, scientists can also describe chemical events on the level of individual atoms or molecules, which is referred to as the *microscopic viewpoint*. We will see examples of both macroscopic and microscopic viewpoints throughout this book (Figure **1.2.2**).




Figure **1.2.2**: How Many Particles Are Needed for a Period in a Sentence? Although we do not notice it from a macroscopic perspective, matter is composed of microscopic particles so tiny that billions of them are needed to make a speck we can see with the naked eye. The ×25 and ×400,000,000 indicate the number of times the image is magnified.

Mixtures

A material composed of two or more substances is a **mixture**. In a mixture, the individual substances maintain their chemical identities. Many mixtures are obvious combinations of two or more substances, such as a mixture of sand and water. Such mixtures are called **heterogeneous mixtures**. In some mixtures, the components are so intimately combined that they act like a single substance (even though they are not). Mixtures with a consistent or uniform composition throughout are called **homogeneous mixtures** (or solutions). For example, when sugar is dissolved in water to form a liquid solution, the individual properties of the components cannot be distinguished. Other examples or homogenous mixtures include solid solutions, like the metal alloy steel, and gaseous solutions, like air which is a mixture of mainly nitrogen and oxygen.

✔ Example 1.2.1

How would a chemist categorize each example of matter?

a. saltwater
b. soil
c. water
d. oxygen

Answer a

Saltwater acts as if it were a single substance even though it contains two substances—salt and water. Saltwater is a *homogeneous mixture*, or a solution.

Answer b

Soil is composed of small pieces of a variety of materials, so it is a ***h**eterogeneous mixture*.

Answer c

Water is a *substance*; more specifically, because water is composed of hydrogen and oxygen, it is a *compound*.

Answer d

Oxygen, a *substance*, is an *element*.

? Exercise 1.2.2

How would a chemist categorize each example of matter?

a. breakfast coffee
b. hydrogen
c. an egg

Answer a

homogeneous mixture or solution

Answer b

element

Answer c

heterogeneous mixture

Phases or Physical States of Matter

All matter can be further classified by one of three physical **states** or **phases**, solid, liquid or gas. These three descriptions each imply that the matter has certain physical properties when in these states. A solid has a definite shape and a definite volume. Liquids ordinarily have a definite volume but not a definite shape; they take the shape of their containers. Gases have neither a definite shape nor a definite volume, and they expand to fill their containers.




Figure 1.2.3: The three most common states or phases of matter are solid, liquid, and gas. (CC BY-4.0; OpenStax)

A beaker labeled solid contains a cube of red matter and says has fixed shape and volume. A beaker labeled liquid contains a brownish-red colored liquid. This beaker says takes shape of container, forms horizontal surfaces, has fixed volume. The beaker labeled gas is filled with a light brown gas. This beaker says expands to fill container.

We encounter matter in each phase every day; in fact, we regularly encounter water in all three phases: ice (solid), water (liquid), and steam (gas) (Figure 1.2.2).



Figure 1.2.5: Boiling Water. When liquid water boils to make gaseous water, it undergoes a phase change. (CC BY-SA 3.0 Unported; Markus Schweiss via Wikipedia)

We know from our experience with water that substances can change from one phase to another if the conditions are right. Typically, varying the temperature of a substance (and, less commonly, the pressure exerted on it) can cause a **phase change**, a physical process in which a substance changes from one phase to another (Figure 1.2.5). Phase changes are identified by particular names depending on what phases are involved, as summarized in Table 1.2.1.

Table 1.2.1: Phase Changes

Change	Name
solid to liquid	melting, fusion
solid to gas	sublimation
liquid to gas	boiling, evaporation
liquid to solid	solidification, freezing
gas to liquid	condensation
gas to solid	deposition

Figure **1.2.3** illustrates the relationships between the different ways matter can be classified.




Figure **1.2.6**: The Classification of Matter. Matter can be classified in a variety of ways, depending on its properties.
This table starts with a substance. If there is only one present, it can either be an element or a compound. If there is more than one present, it can either be a homogeneous mixture or a heterogeneous mixture. From these choices, the substance can be in a certain phase. It can either be solid, liquid, or gas, and each phase can go from one to another.

Concept Review Exercises

1. Explain the differences between the physical properties of matter and the chemical properties of matter.
2. What is the difference between a heterogeneous mixture and a homogeneous mixture? Give an example of each.
3. Give at least two examples of a phase change and state the phases involved in each.

Answers

1. Physical properties describe the existence of matter, and chemical properties describe how substances change into other substances.
2. A heterogeneous mixture is obviously a mixture, such as dirt; a homogeneous mixture behaves like a single substance, such as saltwater.
3. solid to liquid (melting) and liquid to gas (boiling) (answers will vary)

Key Takeaways

- Matter can be described with both physical properties and chemical properties.
- Matter can be identified as an element, a compound, or a mixture

Exercise 1.2.3

Does each statement refer to a chemical property or a physical property?

1. Balsa is a very light wood.
2. If held in a flame, magnesium metal burns in air.
3. Mercury has a density of 13.6 g/mL.
4. Human blood is red.

Answer

1. physical property
2. chemical property
3. physical property
4. physical property

Exercise 1.2.4

Does each statement refer to a chemical property or a physical property?

1. The elements sodium and chlorine can combine to make table salt.
2. The metal tungsten does not melt until its temperature exceeds 3,000°C.
3. The ingestion of ethyl alcohol can lead to disorientation and confusion.
4. The boiling point of isopropyl alcohol, which is used to sterilize cuts and scrapes, is lower than the boiling point of water

Answer

1. chemical property
2. physical property
3. chemical property
4. physical property

Exercise 1.2.5

Define *element*. How does it differ from a compound?

Answer

An element is a substance that cannot be broken down into chemically simpler components. Compounds can be broken down into simpler substances.

Exercise 1.2.6

Define *compound*. How does it differ from an element?

Answer

A compound is composed of two or more elements combined in a fixed ratio. An element is the simplest chemical substance.

Exercise 1.2.7

Give two examples of a heterogeneous mixture.

Answer

a salt and pepper mix and a bowl of cereal (answers will vary)

Exercise 1.2.8

Give two examples of a homogeneous mixture.

Answer

vinegar and rubbing alcohol (answers will vary)

Exercise 1.2.9

Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.

1. xenon, a substance that cannot be broken down into chemically simpler components
2. blood, a substance composed of several types of cells suspended in a salty solution called plasma
3. water, a substance composed of hydrogen and oxygen

Answer

1. element
2. heterogeneous mixture
3. compound

Exercise 1.2.10

Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.

1. sugar, a substance composed of carbon, hydrogen, and oxygen
2. hydrogen, the simplest chemical substance
3. dirt, a combination of rocks and decaying plant matter

Answer

1. compound
2. element
3. heterogeneous mixture

Exercise 1.2.11

Identify each substance as an element, a compound, a heterogeneous mixture, or a solution.

1. air, primarily a mixture of nitrogen and oxygen
2. ringer's lactate, a standard fluid used in medicine that contains salt, potassium, and lactate compounds all dissolved in sterile water
3. tartaric acid, a substance composed of carbon, hydrogen, and oxygen

Answer

1. heterogeneous mixture
2. solution
3. compound

Exercise 1.2.12

What word describes each phase change?

1. solid to liquid
2. liquid to gas
3. solid to gas

Answer

1. melting or fusion
2. boiling or evaporation
3. sublimation

Exercise 1.2.13

1. What word describes each phase change?
    1. liquid to solid
    2. gas to liquid
    3. gas to solid

Answer

1. freezing
2. condensation
3. deposition

1.2: Classification of Matter is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.2: The Classification of Matter** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.3: Introduction to Kinetic and Potential Energy


Learning Objectives

- Define energy and its two main types listed below
- Describe the conservation of energy

Energy is the capacity to do work. You have a sense of what work is from regular life, it's things that require effort. Energy and work have the same units.

Kinetic Energy

At the end of the seventeenth century, a quantity was introduced into mechanics to explain collisions between two perfectly elastic bodies, in which one body makes a head-on collision with an identical body at rest. The first body stops, and the second body moves off with the initial velocity of the first body. (If you have ever played billiards or croquet, or seen a model of Newton's Cradle, you have observed this type of collision.) The idea behind this quantity was related to the forces acting on a body and was referred to as "the energy of motion." Later on, during the eighteenth century, the name **kinetic energy** was given to energy of motion.



Newton's cradle in motion. One ball is set in motion and soon collides with the rest, conveying the energy through the rest of the balls and eventually to the last ball, which in turn is set in motion. Image used with permission (CC SA-BY 3.0; Dominique Toussaint).

The equation for kinetic energy is

$$KE = \frac{1}{2}mv^2 \tag{1.3.1}$$

where KE is kinetic energy, m is mass, and v is velocity. This definition should make sense: big things moving fast have the most energy, the most ability to shove other things or knock them over, etc.

Example 1.3.1: Kinetic Energy of an Object

a. What is the kinetic energy of an 80-kg athlete, running at 10 m/s?

b. The Chicxulub crater in Yucatan, one of the largest existing impact craters on Earth, is thought to have been created by an asteroid, traveling at 22 km/s and releasing 4.2 x 10^{23} J of kinetic energy upon impact. What was its mass?

c. In nuclear reactors, thermal neutrons, traveling at about 2.2 km/s, play an important role. What is the kinetic energy of such a particle?

Strategy

To answer these questions, you can use the definition of kinetic energy in Equation 1.3.1. You also have to look up the mass of a neutron.

Solution

Do not forget to convert km into m to do these calculations, although, to save space, we omitted showing these conversions.

a. $$KE = \frac{1}{2}(80\ kg)(10\ m/s)^2 = 4.0\ kJ.$$

b. $$m = \frac{2K}{v^2} = \frac{2(4.2 \times 10^{23}\ J)}{22\ km/s)^2} = 1.7 \times 10^{15}\ kg.$$

c. $$KE = \frac{1}{2}(1.68 \times 110^{-27}\ kg)(2.2\ km/s)^2 = 4.1 \times 10^{-21}\ J.$$

Significance

In this example, we used the way mass and speed are related to kinetic energy, and we encountered a very wide range of values for the kinetic energies. Different units are commonly used for such very large and very small values. The energy of the impactor in part (b) can be compared to the explosive yield of TNT and nuclear explosions, 1 megaton = 4.18 x 10^{15} J. The Chicxulub asteroid's kinetic energy was about a hundred million megatons. At the other extreme, the energy of subatomic particle is expressed in electron-volts, 1 eV = 1.6 x 10^{-19} J. The thermal neutron in part (c) has a kinetic energy of about one fortieth of an electronvolt.

Exercise 1.3.1

a. A car and a truck are each moving with the same kinetic energy. Assume that the truck has more mass than the car. Which has the greater speed?
b. A car and a truck are each moving with the same speed. Which has the greater kinetic energy?

Potential Energy

Potential energy is energy that comes from position. There are many different flavors of potential energy with two discussed below.

Gravitational Potential Energy

Gravitational potential energy is the energy that objects that are "high up" in a gravitation field (e.g., surface of earth). The equation for potential energy from gravity is

$$PE = mgh \tag{1.3.2}$$

where PE is potential energy, m is mass, g is the acceleration of gravity, and h is the height. This makes the units of energy very clear:

$$\text{mass} \times \text{distance} \times \text{acceleration}$$

or

$$\text{force} \times \text{distance}$$

which both comes to units of kg•m^2s^{-2}.

Example 1.3.2

A 15 gram ball sits on top of a 2m high refrigerator. What is the potential energy of the ball at the top of the refrigerator?

Solution

Use equation for gravitational potential energy (Equation 1.3.2) with mass =15 grams. This mass however has to be in kilograms. The conversion to grams to kilograms is:

- 1,000 grams per 1 kg
- height=2
- g=9.8

$$PE = (0.015)(9.8)(2) = 0.294\ J$$

Exercise 1.3.2A

How much, in grams, does a cart full of groceries sitting on top of a 2 m hill weigh if its PE is 0.3 J?

Answer

$$0.3 = m(9.8)(2)$$
$$m = 0.015kg$$
$$= 15\,g$$

Exercise 1.3.2B

A 200 gram weight is placed on top of a shelf with a potential energy of 5 J. How high is the weight?

Answer

$$5 = \left(\frac{200}{1000}\right)(9.8)(h)$$
$$h = 2.55\,m$$

Coulombic Potential Energy


In chemistry, the force that leads to potential energy is almost always the Coulomb force, not gravity. The PE of two charged particles at a distance can be found through the equation:

$$E = \frac{q_1 q_2}{4\pi\epsilon_o r} \tag{1.3.3}$$

- r is distance
- q_1 and q_2 are the charges
- ϵ_o is $8.85 \times 10^{-12}\,C^2/J\,m$

When the charges have the same sign, they repel and will accelerate away from each other if allowed to move; the potential energy has a positive sign. When the charges have the opposite sign, they attract each other and have negative potential energy. If they are allowed to get closer together, the potential energy will get more negative. If they are separated, r gets bigger and the potential energy approaches zero.

For charges with the same sign, E has a + sign and tends to get smaller as r increases. This can explain why like charges repel each other. Systems prefer a low PE and thus repel each other which increases the distance between them (r) and lowers the PE.



For charges with different charges, the opposite of what is stated above is true. E has a - sign which becomes even more negative as the opposite charged particles attract, or come closer together.

Example 1.3.3

Calculate the potential energy of two particles with charges of $3 \times 10^{-6}\,C$ and $3.9 \times 10^{-6}\,C$ are separated by a distance of $1\,m$.

Solution

$$E = \frac{(3 \times 10^{-6})(3.9 \times 10^{-6})}{4\pi 8.85 \times 10^{-12}}$$
$$E = 0.105\,J$$

Exercise 1.3.3

Find the distance between two particles that have a PE of 0.2 J and charges of $2.5 \times 10^{-6}\,C$ and $3.1 \times 10^{-6}\,C$.

Answer


$$0.2\,J = \frac{(2.5 \times 10^{-6})(3.1 \times 10^{-6})}{4\pi 8.85 \times 10^{-12} r})$$
$$= \frac{(8.99 \times 10^{9})(7.75 \times 10^{-11})}{r}$$
$$= \frac{0.6967}{r}$$

cross multiply and solve for r

$$r = 3.5\,m$$

Conservation of Energy

You probably learned about **conservation of energy** already in a physics class. For instance, if you have a pendulum as shown, at Position 1 the weight has some potential energy, but no kinetic energy. When you release the weight, the weight falls, moving through Position 2. At Position 2, some of the potential energy has been converted to kinetic energy. Finally, at Position 3, all the potential energy has been converted to kinetic energy. As it passes 3, the process is reversed, and kinetic energy is converted to potential energy. When the weight reaches Position 4, all the kinetic energy has been converted back to the same amount of potential energy it started with at 1. This is just one example of conservation of energy. It is a general observation that the amount of energy in the universe doesn't change, and the amount of energy in a particular system doesn't change unless there is a flow of energy in or out.



A pendulum. The red ball is raised to Position 1 and released; it swings through Positions 2 and 3 to Position 4 on the other side.

Contributors

- Emily V Eames (City College of San Francisco)
- Brittanie Harbick (UCD), Laura Suh (UCD), Amrit Paul Bains (UCD)
- Kinetic Energy by OpenStax (CC BY 4.0).

1.3: Introduction to Kinetic and Potential Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

1.4: Measurements

Learning Objectives

- Express quantities properly using a number and a unit.

A coffee maker's instructions tell you to fill the coffeepot with 4 cups of water and use 3 scoops of coffee. When you follow these instructions, you are measuring. When you visit a doctor's office, a nurse checks your temperature, height, weight, and perhaps blood pressure (Figure **1.4.1**); the nurse is also measuring.



Figure **1.4.1**: Measuring Blood Pressure. A nurse or a doctor measuring a patient's blood pressure is taking a measurement. (GNU Free Documentation License; Pia von Lützau via Wikipedia).

Chemists measure the properties of matter using a variety of devices or measuring tools, many of which are similar to those used in everyday life. Rulers are used to measure length, balances (scales) are used to measure mass (weight), and graduated cylinders or pipettes are used to measure volume. Measurements made using these devices are expressed as quantities. A **quantity** is an amount of something and consists of a **number** and a **unit**. The number tells us how many (or how much), and the unit tells us what the scale of measurement is. For example, when a distance is reported as "5.2 kilometers," we know that the quantity has been expressed in units of kilometers and that the number of kilometers is 5.2.

$$\underbrace{5.2}_{\text{number}}\ \underbrace{\text{kilometers}}_{\text{unit}}$$

If you ask a friend how far he or she walks from home to school, and the friend answers "12" without specifying a unit, you do not know whether your friend walks—for example, 12 miles, 12 kilometers, 12 furlongs, or 12 yards.

Without units, a number can be meaningless, confusing, or possibly life threatening. Suppose a doctor prescribes phenobarbital to control a patient's seizures and states a dosage of "100" without specifying units. Not only will this be confusing to the medical professional giving the dose, but the consequences can be dire: 100 mg given three times per day can be effective as an anticonvulsant, but a single dose of 100 g is more than 10 times the lethal amount.

> *Both a number and a unit must be included to express a quantity properly.*

To understand chemistry, we need a clear understanding of the units chemists work with and the rules they follow for expressing numbers. The next two sections examine the rules for expressing numbers.

Exercise 1.4.1

Identify the ***number*** and the ***unit*** in each quantity.

a. one dozen eggs
b. 2.54 centimeters
c. a box of pencils
d. 88 meters per second

Answer a

The number is one, and the unit is dozen.

Answer b

The number is 2.54, and the unit is centimeter.

Answer c

The number 1 is implied because the quantity is only *a* box. The unit is box of pencils.

Answer d

The number is 88, and the unit is meters per second. Note that in this case the unit is actually a combination of two units: meters and seconds.

? Exercise 1.4.2

Identify the ***number*** and the ***unit*** in each quantity.

a. 99 bottles of soda
b. 60 miles per hour
c. 32 fluid ounces
d. 98.6 degrees Fahrenheit

Answer a

The number is 99, and the unit is bottles of soda.

Answer b

The number is 60, and the unit is miles per hour.

Answer c

The number 32, and the unit is fluid ounces

Answer d

The number is 98.6, and the unit is degrees Fahrenheit

? Exercise 1.4.2

What are the two necessary parts of a quantity?

Answer

The two necessary parts are the number and the unit.

Key Takeaway

- Identify a quantity properly with a number and a unit.

1.4: Measurements is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.3: Measurements** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.5: Expressing Numbers- Scientific Notation

Learning Objectives

- Express a large number or a small number in scientific notation.
- Convert a number in scientific notation to standard conventional form.

The instructions for making a pot of coffee specified 3 scoops (rather than 12,000 grounds) because any measurement is expressed more efficiently with units that are appropriate in size. In science, however, we often must deal with quantities that are extremely small or incredibly large. For example, you may have 5,000,000,000,000 red blood cells in a liter of blood, and the diameter of an iron atom is 0.000000014 inches. Numbers with many zeros can be cumbersome to work with, so scientists use scientific notation.

Scientific notation is a system for expressing very large or very small numbers in a compact manner. It uses the idea that such numbers can be rewritten as a simple number multiplied by 10 raised to a certain exponent, or power.

Let us look first at very large numbers. Suppose a spacecraft is 1,500,000 miles from Mars. The number 1,500,000 can be thought of as follows:

$$1.5 \times \underbrace{1{,}000{,}000}_{\underbrace{10 \times 10 \times 10 \times 10 \times 10 \times 10}_{10^6}} = 1.5 \times 10^6$$

That is, 1,500,000 is the same as 1.5 times 1 million, and 1 million is 10 × 10 × 10 × 10 × 10 × 10, or 10^6 (which is read as "ten to the sixth power"). Therefore, 1,500,000 can be rewritten as 1.5 times 10^6, or 1.5×10^6. The distance of the spacecraft from Mars can therefore be expressed as 1.5×10^6 miles.

Recall that:

- $10^0 = 1$
- $10^1 = 10$
- $10^2 = 100$
- $10^3 = 1{,}000$
- $10^4 = 10{,}000$
- and so forth

The standard convention for expressing numbers in scientific notation is to write a single *nonzero* first digit, a decimal point, and the rest of the digits, excluding any trailing zeros (see rules for significant figures in the next section for more details on what to exclude). This number is followed by a multiplication sign and then by 10 raised to the power necessary to reproduce the original number. For example, although 1,500,000 can also be written as $15. \times 10^5$ (which would be 15. × 100,000), the convention is to have only one digit before the decimal point. How do we know to what power 10 is raised? The power is the number of places you have to move the decimal point to the *left* to place it after the first digit, so that the number being multiplied is *between 1 and 10*:

$$1{,}500{,}000 = 1.5 \times 10^6$$

Example 1.5.1: Scientific Notation

Express each number in scientific notation.

a. 67,000,000,000
b. 1,689
c. 12.6

Answer a

Moving the decimal point 10 places to the left gives 6.7×10^{10}.

Answer b

The decimal point is assumed to be at the end of the number, so moving it three places to the left gives 1.689×10^3.

Answer c

In this case, we need to move the decimal point only one place to the left, which yields 1.26×10^1.

? Exercise 1.5.1

Express each number in scientific notation.

a. 1,492
b. 102,000,000
c. 101,325

Answer a

Moving the decimal point 3 places to the left gives 1.492×10^3.

Answer b

The decimal point is assumed to be at the end of the number, so moving it 8 places to the left gives 1.02×10^8.

Answer c

Moving the decimal point 5 places to the left yields 1.01325×10^5.

To change a number in **scientific notation** to **standard form**, we reverse the process, moving the decimal point to the right. Add zeros to the end of the number being converted, if necessary, to produce a number of the proper magnitude. Lastly, we drop the number 10 and its power.

$$1.5 \times 10^6 = 1.500000 = 1{,}500{,}000$$

✓ Example 1.5.2

Express each number in standard, or conventional notation.

a. 5.27×10^4
b. 1.0008×10^6

Answer a

Moving the decimal four places to the right and adding zeros give 52,700.

Answer b

Moving the decimal six places to the right and adding zeros give 1,000,800.

✓ Exercise 1.5.2

Express each number in standard, or conventional notation.

a. 6.98×10^8
b. 1.005×10^2

Answer a

Moving the decimal point eight places to the right and adding zeros give 698,000,000.

Answer b

Moving the decimal point two places to the right gives 100.5

We can also use scientific notation to express numbers whose magnitudes are less than 1. For example, the quantity 0.006 centimeters can be expressed as follows:

$$6 \times \underbrace{\frac{1}{1{,}000}}_{\underbrace{\frac{1}{10} \times \frac{1}{10} \times \frac{1}{10}}_{10^{-3}}} = 6 \times 10^{-3}$$

That is, 0.006 centimeters is the same as 6 *divided by* one thousand, which is the same as 6 *divided* by 10 x 10 x 10 or 6 *times* 10^{-3} (which is read as "ten to the negative third power"). Therefore, 0.006 centimeters can be rewritten as 6 times 10^{-3}, or 6×10^{-3} centimeters.

Recall that:

- $10^{-1} = 1/10$
- $10^{-2} = 1/100$
- $10^{-3} = 1/1{,}000$
- $10^{-4} = 1/10{,}000$
- $10^{-5} = 1/100{,}000$
- and so forth

We use a negative number as the power to indicate the number of places we have to move the decimal point to the right to make it follow the first nonzero digit so that the number is between 1 and 10. This is illustrated as follows:

$$0.006 = 6 \times 10^{-3}$$

Note:

In writing scientific notations, the convention is to have only one digit before the decimal point.

- Numbers that are greater than one have a positive power in scientific notation. If the decimal point is moved to the left n places, the power (n) of 10 is positive. $1{,}500{,}000 = 1.5 \times 10^{6}$
- Numbers that are less than one have a negative power in scientific notation. If the decimal point is moved to the right n places, the power (n) of 10 is negative. $0.006 = 6 \times 10^{-3}$

Example 1.5.3

Express each number in scientific notation.

a. 0.000006567
b. −0.0004004
c. 0.000000000000123

Answer a

Move the decimal point six places to the right to get 6.567×10^{-6}.

Answer b

Move the decimal point four places to the right to get -4.004×10^{-4}. The negative sign on the number itself does not affect how we apply the rules of scientific notation.

Answer c

Move the decimal point 13 places to the right to get 1.23×10^{-13}.

Exercise 1.5.3

Express each number in scientific notation.

a. 0.000355
b. 0.314159
c. −0.051204

Answer a

Moving the decimal point four places to the right gives 3.55×10^{-4}.

Answer b

Moving the decimal point one place to the right gives 3.14159×10^{-1}.

Answer c

Moving the decimal point one place to the right gives -5.1204×10^{-2}.

As with numbers with positive powers of 10, when changing from **scientific** notation to **standard or conventional** format, we reverse the process.

$$6 \times 10^{-3} = .006 = 0.006$$

Note

Changing a number in scientific notation to standard form:

- If the scientific notation has a positive power, the standard number is greater than one. Example: $8 \times 10^{4} = 80{,}000$
- If the scientific notation has a negative power, then the standard number is less than one. Example: $8 \times 10^{-2} = 0.08$

Example 1.5.4

Change the number in scientific notation to standard form.

a. 6.22×10^{-2}
b. 9.9×10^{-9}

Answer a

0.0622

Answer b

0.0000000099

Exercise 1.5.4

Change the number in scientific notation to standard form.

a. 9.98×10^{-5}
b. 5.109×10^{-8}

Answer a

0.0000998

Answer b

0.00000005109

Although calculators can show 8 to 10 digits in their display windows, that is not always enough when working with very large or very small numbers. For this reason, many calculators are designed to handle scientific notation. The method for entering scientific notation differs for each calculator model, so take the time to learn how to do it properly on your calculator, *asking your instructor for assistance if necessary*. If you do not learn to enter scientific notation into your calculator properly, you will not get the correct final answer when performing a calculation.

? Exercise 1.5.1

Why it is easier to use scientific notation to express very large or very small numbers?

Answer

Scientific notation is more convenient than listing a large number of zeros.

? Exercise 1.5.2

What is the relationship between how many places a decimal point moves and the power of 10 used in changing a conventional number into scientific notation?

Answer

The number of places the decimal point moves equals the power of 10. The power of 10 is positive if the decimal point moves to the left and negative if the decimal point moves to the right.

Key Takeaway

- Large or small numbers are expressed in scientific notation, which use powers of 10.

1.5: Expressing Numbers- Scientific Notation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.4: Expressing Numbers - Scientific Notation** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.6: Expressing Numbers- Significant Figures

Learning Objectives

- Understand the importance of significant figures in measured numbers.
- Identify the number of significant figures in a reported value.
- Use significant figures correctly in arithmetical operations.

Scientists have established certain conventions for communicating the degree of **precision** of a measurement, which is dependent on the measuring device used. Imagine, for example, that you are using a meterstick to measure the width of a table. The centimeters (cm) marked on the meterstick, tell you how many *centimeters* wide the table is. Many metersticks also have markings for millimeters (mm), so we can measure the table to the nearest *millimeter*. Most metersticks do not have any smaller (or more precise) markings indicated, so you cannot report the measured width of the table any more precise than to the nearest millimeter. However, you can *estimate* one past the smallest marking, in this case the millimeter, to the next decimal place in the measurement (Figure 1.6.1).




Figure 1.6.1: Measuring an Object to the Correct Number of Digits. How many digits should be reported for the length of this object?

The concept of **significant figures** takes this limitation into account. The significant figures of a measured quantity are defined as all the digits known with *certainty* (those indicated by the markings on the measuring device) ***and*** the first uncertain, or estimated, digit (one digit past the smallest marking on the measuring device). It makes no sense to report any digits after the first uncertain one, so it is the last digit reported in a measurement. Zeros are used when needed to place the significant figures in their correct positions. Thus, zeros are sometimes counted as significant figures but are sometimes only used as placeholders.

> *"Sig figs" is a common abbreviation for significant figures.*

Consider the earlier example of measuring the width of a table with a meterstick. If the table is measured and reported as being 1,357 mm wide, the number 1,357 has four significant figures. The 1 (thousands place), the 3 (hundreds place), and the 5 (tens place) are certain; the 7 (ones place) is assumed to have been estimated. It would make no sense to report such a measurement as 1,357.0 (five Sig Figs) or 1,357.00 (six Sig Figs) because that would suggest the measuring device was able to determine the width to the nearest tenth or hundredth of a millimeter, when in fact it shows only tens of millimeters and therefore the ones place was estimated.

On the other hand, if a measurement is reported as 150 mm, the 1 (hundreds) and the 5 (tens) are known to be significant, but how do we know whether the zero is or is not significant? The measuring device could have had marks indicating every 100 mm or marks indicating every 10 mm. How can you determine if the zero is significant (the estimated digit), or if the 5 is significant and the zero a value placeholder?

The **rules** for deciding which digits in a measurement are significant are as follows:

1. All nonzero digits are significant. In 1,357 mm, all the digits are significant.
2. *Sandwiched (or embedded) zeros*, those between significant digits, are significant. Thus, 405 g has *three* significant figures.
3. *Leading zeros*, which are zeros at the beginning of a decimal number less than 1, are not significant. In 0.000458 mL, the first four digits are leading zeros and are not significant. The zeros serve only to put the digits 4, 5, and 8 in the correct decimal positions. This number has three significant figures.
4. *Trailing zeros*, which are zeros at the end of a number, are significant only if the number has a decimal point. Thus, in 1,500 m, the two trailing zeros are not significant because the number is written without a decimal point; the number has two significant

figures. However, in 1,500.00 m, all six digits are significant because the number has a decimal point.

Example 1.6.1

How many significant figures does each number have?

a. 6,798,000
b. 6,000,798
c. 6,000,798.00
d. 0.0006798

Answer a

four (by rules 1 and 4)

Answer b

seven (by rules 1 and 2)

Answer c

nine (by rules 1, 2, and 4)

Answer d

four (by rules 1 and 3)

Exercise 1.6.1

How many significant figures does each number have?

a. 2.1828
b. 0.005505
c. 55,050
d. 5
e. 500

Answer a

five

Answer b

four

Answer c

four

Answer d

one

Answer e

one

Rounding off numbers

Before dealing with the specifics of the rules for determining the significant figures in a calculated result, we need to be able to round numbers correctly. To **round** a number, first decide how many significant figures the number should have. Once you know that, round to that many digits, starting from the left. **If the number immediately to the right of the last significant digit is less than 5, it is dropped and the value of the last significant digit remains the same. If the number immediately to the right of the last significant digit is greater than or equal to 5, the last significant digit is increased by 1.**

Consider the measurement 207.518m. Right now, the measurement contains six significant figures. How would we successively round it to fewer and fewer significant figures? Follow the process as outlined in Table 1.5.1.

Table 1.5.1: Rounding examples

Number of Significant Figures	Rounded Value	Reasoning
6	207.518	All digits are significant
5	207.52	8 rounds the 1 up to 2
4	207.5	2 is dropped
3	208	5 rounds the 7 up to 8
2	210	8 is replaced by a 0 and rounds the 0 up to 1
1	200	1 is replaced by a 0

Notice that the more rounding that is done, the less reliable the figure is. An approximate value may be sufficient for some purposes, but scientific work requires a much higher level of detail.

It is important to be aware of significant figures when you are mathematically manipulating numbers. For example, dividing 125 by 307 on a calculator gives 0.4071661238… to an infinite number of digits. But do the digits in this answer have any practical meaning, especially when you are starting with numbers that have only three significant figures each? When performing mathematical operations, there are two rules for limiting the number of significant figures in an answer—one rule is for addition and subtraction, and one rule is for multiplication and division.

> *In operations involving significant figures, the answer is reported in such a way that it reflects the reliability of the **least precise** operation. An answer is no more precise than the least precise number used to get the answer.*

Combining Numbers

For **addition or subtraction**, the rule is to stack all the numbers with their decimal points aligned and then limit (round to) the answer's significant figures to the rightmost column for which all the numbers have significant figures. Consider the following:

```
    56.789
+   102.2
+ 1,300.099
= 1,459.088
          ↑ Limit to this column
```

The arrow points to the rightmost column in which all the numbers have significant figures—in this case, the tenths place. Therefore, we will limit our final answer to the tenths place. Is our final answer therefore 1,459.0? No, because when we drop digits from the end of a number, we also have to round the number. Notice that the first dropped digit, in the hundredths place, is 8. This suggests that the answer is actually closer to 1,459.1 than it is to 1,459.0, so we need to round up to **1,459.1**. The standard rules for rounding numbers are simple: If the first dropped digit is 5 or higher, round up. If the first dropped digit is lower than 5, do not round up.

For **multiplication or division**, the rule is to count the number of significant figures in each number being multiplied or divided and then limit the significant figures in the answer to the lowest count. An example is as follows:

$$\underbrace{38.65}_{\text{4 sig figs}} \times \underbrace{105.93}_{\text{5 sig figs}} = \underbrace{4{,}094.1945}_{\text{reduce to 4 sig figs}}$$

The final answer, limited to four significant figures, is **4,094**. The first digit dropped is 1, so we do not round up.

Scientific notation provides a way of communicating significant figures without ambiguity. You simply include all the significant figures in the leading number. For example, the number 4,000 has one significant figure and should be written as the number 4×10^4. The number 450 has two significant figures and would be written in scientific notation as 4.5×10^2, whereas 450.0 has four significant figures and would be written as 4.500×10^2. In scientific notation, all reported digits are significant.

Example 1.6.2

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 23.096 × 90.300
b. 125 × 9.000
c. 1,027 + 610.0 + 363.06

Answer a

The calculator answer is 2,085.5688, but we need to round it to five significant figures. Because the first digit to be dropped (in the hundredths place) is greater than 5, we round up to 2,085.6, which in scientific notation is 2.0856×10^3.

Answer b

The calculator gives 1,125 as the answer, but we limit it to three significant figures and convert into scientific notation: 1.13×10^3.

Answer c

The calculator gives 2,000.06 as the answer, but because 1,027 has its farthest-right significant figure in the ones column, our answer must be limited to the ones position: 2,000 which in scientific notation is 2.000×10^3.

Exercise 1.6.2

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 217 ÷ 903
b. 13.77 + 908.226 + 515
c. 255.0 − 99
d. 0.00666 × 321

Answer a

$0.240 = 2.40 \times 10^{-1}$

Answer b

$1437 = 1.437 \times 10^3$

Answer c

$156 = 1.56 \times 10^2$

Answer d

$2.14 = 2.14 \times 10^0$

Remember that calculators do not understand significant figures. *You* are the one who must apply the rules of significant figures to a result from your calculator.

Exercise 1.6.1

Explain why the concept of significant figures is important in scientific measurements.

Answer

Significant figures represent all the known digits of a measurement plus the first estimated one. It gives information about how precise the measuring device and measurement is.

? Exercise 1.6.2

State the rules for determining the significant figures in a measurement.

Answer

All nonzero digits are significant; zeros between nonzero digits are significant; zeros at the end of a nondecimal number or the beginning of a decimal number are not significant; zeros at the end of a decimal number are significant.

? Exercise 1.6.3

When do you round a number up, and when do you not round a number up?

Answer

Round up only if the first digit dropped is 5 or higher.

Key Takeaways

- Significant figures properly report the number of measured and estimated digits in a measurement.
- The rule in multiplication and division is that the final answer should have the same number of significant figures as there are in the value with the fewest significant figures.
- The rule in addition and subtraction is that the final answer should have the same number of decimal places as the term with the fewest decimal places.

1.6: Expressing Numbers- Significant Figures is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.5: Expressing Numbers - Significant Figures** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.7: The International System of Units

Learning Objectives

- Recognize the SI base units and explain the system of prefixes used with them.
- Define and calculate density.

People who live in the United States measure weight in pounds, height in feet and inches, and a car's speed in miles per hour. In contrast, chemistry and other branches of science use the International System of Units (also known as **SI** after *Système Internationale d'Unités*), which was established so that scientists around the world could communicate efficiently with each other. Many countries have also adopted SI units for everyday use as well. The United States is one of the few countries that has not.

Base SI Units

Base (or basic) units, are the fundamental units of SI. There are seven base units, which are listed in Table **1.7.1**, Chemistry uses five of the base units: the mole for amount, the kilogram for mass, the meter for length, the second for time, and the kelvin for temperature. The degree Celsius (°C) is also commonly used for temperature. The numerical relationship between kelvins and degrees Celsius is as follows:

$$K = °C + 273 \tag{1.7.1}$$

The United States uses the English (sometimes called Imperial) system of units for many quantities. Inches, feet, miles, gallons, pounds, and so forth, are all units connected with the English system of units. There have been many mistakes due to the improper conversion of units between the SI and English systems.

Table **1.7.1**: The Seven Base SI Units

Property	Unit	Abbreviation
length	meter	m
mass	kilogram	kg
time	second	s
amount	mole	mol
temperature	kelvin	K
electrical current	ampere	amp
luminous intensity	candela	cd

The size of each base unit is defined by international convention. For example, the *kilogram* is defined as the quantity of mass of a special metal cylinder kept in a vault in France (Figure **1.7.1**). The other base units have similar definitions and standards. The sizes of the base units are not always convenient for all measurements. For example, a meter is a rather large unit for describing the width of something as narrow as human hair. Instead of reporting the diameter of hair as 0.00012 m or as 1.2×10^{-4} m using scientific notation as discussed in section 1.4, SI also provides a series of **prefixes** that can be attached to the units, creating units that are larger or smaller by powers of 10.

Common prefixes and their multiplicative factors are listed in Table **1.7.2**. (Perhaps you have already noticed that the base unit *kilogram* is a combination of a prefix, kilo- meaning 1,000 ×, and a unit of mass, the gram.) Some prefixes create a multiple of the original unit: 1 kilogram equals 1,000 grams, and 1 megameter equals 1,000,000 meters. Other prefixes create a fraction of the original unit. Thus, 1 centimeter equals 1/100 of a meter, 1 millimeter equals 1/1,000 of a meter, 1 microgram equals 1/1,000,000 of a gram, and so forth.

Table **1.7.2**: Prefixes Used with SI Units

Prefix	Abbreviation	Multiplicative Factor	Multiplicative Factor in Scientific Notation
giga-	G	1,000,000,000 ×	10^{9} ×
mega-	M	1,000,000 ×	10^{6} ×
kilo-	k	1,000 ×	10^{3} ×
deca-	D	10 ×	10^{1} ×
deci-	d	1/10 ×	10^{-1} ×
centi-	c	1/100 ×	10^{-2} ×
milli-	m	1/1,000 ×	10^{-3} ×
micro-	μ*	1/1,000,000 ×	10^{-6} ×
nano-	n	1/1,000,000,000 ×	10^{-9} ×

*The letter μ is the Greek lowercase letter for *m* and is called “mu,” which is pronounced “myoo.”

Both SI units and prefixes have abbreviations, and the combination of a prefix abbreviation with a base unit abbreviation gives the abbreviation for the modified unit. For example, kg is the abbreviation for kilogram. We will be using these abbreviations throughout this book.



Figure **1.7.1**: The Kilogram. The standard for the kilogram is a platinum-iridium cylinder kept in a special vault in France. Source: Photo reproduced by permission of the Bureau International des Poids et Mesures, who retain full internationally protected copyright.

The Difference Between Mass and Weight

The mass of a body is a measure of its inertial property or how much matter it contains. The weight of a body is a measure of the force exerted on it by gravity or the force needed to support it. Gravity on earth gives a body a downward acceleration of about 9.8 m/s^2. In common parlance, weight is often used as a synonym for mass in weights and measures. For instance, the verb “to weigh” means “to determine the mass of” or “to have a mass of.” The incorrect use of weight in place of mass should be phased out, and the term mass used when mass is meant. The SI unit of mass is the kilogram (kg). In science and technology, the weight of a body in a particular reference frame is defined as the force that gives the body an acceleration equal to the local acceleration of free fall in that reference frame. Thus, the SI unit of the quantity weight defined in this way (force) is the newton (N).

Derived SI Units

Derived units are combinations of SI base units. Units can be multiplied and divided, just as numbers can be multiplied and divided. For example, the area of a square having a side of 2 cm is 2 cm × 2 cm, or 4 cm^2 (read as “four centimeters squared” or “four square centimeters”). Notice that we have squared a length unit, the centimeter, to get a derived unit for area, the square centimeter.




Figure 1.7.2: The relative lengths of 1 m, 1 yd, 1 cm, and 1 in. are shown (not actual size), as well as comparisons of 2.54 cm and 1 in., and of 1 m and 1.094 yd. (CC BY 4.0; OpenStax)

Volume is an important quantity that uses a derived unit. **Volume** is the amount of space that a given substance occupies and is defined geometrically as length × width × height. Each distance can be expressed using the meter unit, so volume has the derived unit m × m × m, or m^3 (read as “meters cubed” or “cubic meters”). A cubic meter is a rather large volume, so scientists typically express volumes in terms of 1/1,000 of a cubic meter. This unit has its own name—the liter (L). A liter is a little larger than 1 US quart in volume. Below are approximate equivalents for some of the units used in chemistry.

Approximate Equivalents to Some SI Units

- 1 m ≈ 39.36 in. ≈ 3.28 ft ≈ 1.09 yd
- 1 in. ≈ 2.54 cm
- 1 km ≈ 0.62 mi
- 1 kg ≈ 2.20 lb
- 1 lb ≈ 454 g
- 1 L ≈ 1.06 qt
- 1 qt ≈ 0.946 L

As shown in Figure 1.7.3, a liter is also 1,000 cm^3. By definition, there are 1,000 mL in 1 L, so 1 milliliter and 1 cubic centimeter represent the same volume.

$$1\ mL = 1\ cm^3 \tag{1.7.2}$$

Example 1.7.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kiloliter
b. microsecond
c. decimeter
d. nanogram

Answer a

The abbreviation for a kiloliter is kL. Because kilo means “1,000 ×,” 1 kL equals 1,000 L.

Answer b

The abbreviation for microsecond is µs. Micro implies 1/1,000,000th of a unit, so 1 µs equals 0.000001 s.

Answer c

The abbreviation for decimeter is dm. Deci means 1/10th, so 1 dm equals 0.1 m.

Answer d

The abbreviation for nanogram is ng and equals 0.000000001 g.

? Exercise 1.7.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kilometer
b. milligram
c. nanosecond
d. centiliter

Answer a

km (1,000 m)

Answer b

mg (0.001 g)

Answer c

ns (0.000000001 s)

Answer d

cL (0.01L)




Figure **1.7.3**: Units of Volume. (a) The relative volumes are shown for cubes of 1 m^3, 1 dm^3 (1 L), and 1 cm^3 (1 mL) (not to scale). A liter (L) is defined as a cube 1 dm (1/10th of a meter) on a side. A milliliter (mL), 1/1,000th of a liter, is equal to 1 cubic centimeter. (b) The diameter of a dime is compared relative to the edge length of a 1-cm^3 (1-mL) cube. (CC BY 4.0; OpenStax)

Energy, another important quantity in chemistry, is the ability to perform work, such as moving a box of books from one side of a room to the other side. It has a derived unit of kg•m^2/s^2. (The dot between the kg and m units implies the units are multiplied together.) Because this combination is cumbersome, this collection of units is redefined as a **joule** (J). An older unit of energy, but

likely more familiar to you, the calorie (cal), is also widely used. There are 4.184 J in 1 cal. Energy changes occur during all chemical processes and will be discussed in a later chapter.

To Your Health: Energy and Food

The food in our diet provides the energy our bodies need to function properly. The energy contained in food could be expressed in joules or calories, which are the conventional units for energy, but the food industry prefers to use the kilocalorie and refers to it as the Calorie (with a capital C). The average daily energy requirement of an adult is about 2,000–2,500 Calories, which is 2,000,000–2,500,000 calories (with a lowercase c).

If we expend the same amount of energy that our food provides, our body weight remains stable. If we ingest more Calories from food than we expend, however, our bodies store the extra energy in high-energy-density compounds, such as fat, and we gain weight. On the other hand, if we expend more energy than we ingest, we lose weight. Other factors affect our weight as well—genetic, metabolic, behavioral, environmental, cultural factors—but dietary habits are among the most important.

In 2008 the US Centers for Disease Control and Prevention issued a report stating that 73% of Americans were either overweight or obese. More alarmingly, the report also noted that 19% of children aged 6–11 and 18% of adolescents aged 12–19 were overweight—numbers that had tripled over the preceding two decades. Two major reasons for this increase are excessive calorie consumption (especially in the form of high-fat foods) and reduced physical activity. Partly because of that report, many restaurants and food companies are working to reduce the amounts of fat in foods and provide consumers with more healthy food options.

Density is defined as the mass of an object divided by its volume; it describes the amount of matter contained in a given amount of space.

$$\text{density} = \frac{\text{mass}}{\text{volume}} \tag{1.7.3}$$

Thus, the units of density are the units of mass divided by the units of volume: g/cm^3 or g/mL (for solids and liquids), g/L (for gases), kg/m^3, and so forth. For example, the density of water is about 1.00 g/cm^3, while the density of mercury is 13.6 g/mL. (Remember that 1 mL equals 1 cm^3.) Mercury is over 13 times as dense as water, meaning that it contains over 13 times the amount of matter in the same amount of space. The density of air at room temperature is about 1.3 g/L. Table 1.6.3 shows the densities of some common substances.

Table **1.7.2**: Densities of Common Substances

Solids	Liquids	Gases (at 25 °C and 1 atm)
ice (at 0 °C) 0.92 g/cm^3	water 1.0 g/cm^3	dry air 1.20 g/L
oak (wood) 0.60–0.90 g/cm^3	ethanol 0.79 g/cm^3	oxygen 1.31 g/L
iron 7.9 g/cm^3	acetone 0.79 g/cm^3	nitrogen 1.14 g/L
copper 9.0 g/cm^3	glycerin 1.26 g/cm^3	carbon dioxide 1.80 g/L
lead 11.3 g/cm^3	olive oil 0.92 g/cm^3	helium 0.16 g/L
silver 10.5 g/cm^3	gasoline 0.70–0.77 g/cm^3	neon 0.83 g/L
gold 19.3 g/cm^3	mercury 13.6 g/cm^3	radon 9.1 g/L

Example 1.7.2: Density of Bone

What is the density of a section of bone if a 25.3 cm^3 sample has a mass of 27.8 g?

Solution

Because density is defined as the mass of an object divided by its volume, we can set up the following relationship:

$$\begin{aligned} \text{density} &= \frac{mass}{volume} \\ &= \frac{27.8\ g}{25.3\ cm^3} \\ &= 1.10\ g/cm^3 \end{aligned}$$

Note that we have limited our final answer to three significant figures.

? Exercise 1.7.2: Density of Oxygen

What is the density of oxygen gas if a 15.0 L sample has a mass of 21.7 g?

Answer

1.45 g/L

Density can be used to convert between the mass and the volume of a substance. This will be discussed in the next section.

Concept Review Exercises

1. What is the difference between a base unit and a derived unit? Give two examples of each type of unit.
2. Do units follow the same mathematical rules as numbers do? Give an example to support your answer.
3. What is density?

Answers

1. Base units are the seven fundamental units of SI; derived units are constructed by making combinations of the base units; Two examples of base units: kilograms and meters (answers will vary); Two examples of derived units: grams per milliliter and joules (answers will vary).
2. yes; $\mathbf{mL} \times \frac{\mathbf{g}}{\mathbf{mL}} = \mathbf{g}$ (answers will vary)
3. Density is defined as the mass of an object divided by its volume

Key Takeaways

- Recognize the SI base units and derived units.
- Combining prefixes with base units creates new units of larger or smaller sizes.

1.7: The International System of Units is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.6: The International System of Units** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.8: Converting Units

Learning Objectives

- Convert a value reported in one unit to a corresponding value in a different unit.

The ability to convert from one unit to another is an important skill. For example, a nurse with 50 mg aspirin tablets who must administer 0.2 g of aspirin to a patient needs to know that 0.2 g equals 200 mg, so 4 tablets are needed. Fortunately, there is a simple way to convert from one unit to another.

Conversion Factors

If you learned the SI units and prefixes described, then you know that 1 cm is 1/100th of a meter.

$$1\ \text{cm} = \frac{1}{100}\ \text{m}$$

or

$$100\ \text{cm} = 1\ \text{m}$$

Suppose we divide both sides of the equation by 1 m (both the number *and* the unit):

$$\frac{100\ \text{cm}}{1\ \text{m}} = \frac{1\ \text{m}}{1\ \text{m}}$$

As long as we perform the same operation on both sides of the equals sign, the expression remains an equality. Look at the right side of the equation; it now has the same quantity in the numerator (the top) as it has in the denominator (the bottom). Any fraction that has the same quantity in the numerator and the denominator has a value of 1:

$$\text{same quantity} \left\langle \frac{100\ \text{cm}}{1\ \text{m}} = 1 \right.$$

We know that 100 cm *is* 1 m, so we have the same quantity on the top and the bottom of our fraction, although it is expressed in different units. A fraction that has equivalent quantities in the numerator and the denominator but expressed in *different units* is called a **conversion factor**.

Here is a simple example. How many centimeters are there in 3.55 m? Perhaps you can determine the answer in your head. If there are 100 cm in every meter, then 3.55 m equals 355 cm. To solve the problem more formally with a conversion factor, we first write the quantity we are given, 3.55 m. Then we multiply this quantity by a conversion factor, which is the same as multiplying it by 1. We can write 1 as $\frac{100\ \text{cm}}{1\ \text{m}}$ and multiply:

$$3.55\ \text{m} \times \frac{100\ \text{cm}}{1\ \text{m}}$$

The 3.55 m can be thought of as a fraction with a 1 in the denominator. Because m, the abbreviation for meters, occurs in both the numerator *and* the denominator of our expression, they cancel out:

$$\frac{3.55\ \cancel{\text{m}}}{1} \times \frac{100\ \text{cm}}{1\ \cancel{\text{m}}}$$

The final step is to perform the calculation that remains once the units have been canceled:

$$\frac{3.55}{1} \times \frac{100\ \text{cm}}{1} = 355\ \text{cm} \tag{1.8.1}$$

In the final answer, we omit the 1 in the denominator. Thus, by a more formal procedure, we find that 3.55 m equals 355 cm. A generalized description of this process is as follows:

$$\text{quantity (in old units)} \times \text{conversion factor} = \text{quantity (in new units)}$$

You may be wondering why we use a seemingly complicated procedure for a straightforward conversion. In later studies, the conversion problems you will encounter *will not always be so simple*. If you can master the technique of applying conversion factors, you will be able to solve a large variety of problems.

In the previous example (Equation 1.8.1), we used the fraction $\frac{100\ \text{cm}}{1\ \text{m}}$ as a conversion factor. Does the conversion factor $\frac{1\ \text{m}}{100\ \text{cm}}$ also equal 1? Yes, it does; it has the same quantity in the numerator as in the denominator (except that they are expressed in different units). Why did we not use *that* conversion factor? If we had used the second conversion factor, the original unit would not have canceled, and the result would have been meaningless. Here is what we would have gotten:

$$3.55\ \text{m} \times \frac{1\ \text{m}}{100\ \text{cm}} = 0.0355 \frac{\text{m}^2}{\text{cm}}$$

For the answer to be meaningful, we have to *construct the conversion factor in a form that causes the original unit to cancel out*. Figure 1.8.1 shows a **concept map** for constructing a proper conversion.




Figure 1.8.1: A Concept Map for Conversions. This is how you construct a conversion factor to convert from one unit to another. Meters converted to centimeters. A fraction showing 100 cm over 1 m. The unit you convert to is on top, and the unit you convert from is on bottom.

Significant Figures in Conversions

How do conversion factors affect the determination of significant figures? Numbers in conversion factors based on prefix changes, such as kilograms to grams, are *not* considered in the determination of significant figures in a calculation because the numbers in such conversion factors are exact. **Exact numbers** are *defined* or *counted* numbers, not measured numbers, and can be considered as having an infinite number of significant figures. (In other words, 1 kg is exactly 1,000 g, by the definition of kilo-.) Counted numbers are also exact. If there are 16 students in a classroom, the number 16 is exact. In contrast, conversion factors that come from measurements (such as density, as we will see shortly) or are approximations have a limited number of significant figures and should be considered in determining the significant figures of the final answer.

Example 1.8.1

a. The average volume of blood in an adult male is 4.7 L. What is this volume in milliliters?
b. A hummingbird can flap its wings once in 18 ms. How many seconds are in 18 ms?

Solution

a. We start with what we are given, 4.7 L. We want to change the unit from liters to milliliters. There are 1,000 mL in 1 L. From this relationship, we can construct two conversion factors:

$$\frac{1\ \text{L}}{1,000\ \text{mL}} \quad \text{or} \quad \frac{1,000\ \text{mL}}{1\ \text{L}}$$

We use the conversion factor that will cancel out the original unit, liters, and introduce the unit we are converting to, which is milliliters. The conversion factor that does this is the one on the right.




$$4.7 \cancel{L} \times \frac{1,000 \text{ mL}}{1 \cancel{L}} = 4,700 \text{ mL}$$

Because the numbers in the conversion factor are exact, we do not consider them when determining the number of significant figures in the final answer. Thus, we report two significant figures in the final answer.

b. We can construct two conversion factors from the relationships between milliseconds and seconds:

$$\frac{1,000 \text{ ms}}{1 \text{ s}} \text{ or } \frac{1 \text{ s}}{1,000 \text{ ms}}$$

To convert 18 ms to seconds, we choose the conversion factor that will cancel out milliseconds and introduce seconds. The conversion factor on the right is the appropriate one. We set up the conversion as follows:



$$18 \cancel{\text{ms}} \times \frac{1 \text{ s}}{1,000 \cancel{\text{ms}}} = 0.018 \text{ s}$$

The conversion factor's numerical values do not affect our determination of the number of significant figures in the final answer.

? Exercise 1.8.1

Perform each conversion.

a. 101,000 ns to seconds
b. 32.08 kg to grams

Answer a

$$101,000 \cancel{\text{ns}} \times \frac{1 \text{ s}}{1,000,000,000 \cancel{\text{ns}}} = 0.000101 \text{ s}$$

Answer b

$$32.08 \cancel{\text{kg}} \times \frac{1,000 \text{ g}}{1 \cancel{\text{kg}}} = 32,080 \text{ g}$$

Conversion Factors From Different Units

Conversion factors can also be constructed for converting between different kinds of units. For example, density can be used to convert between the mass and the volume of a substance. Consider mercury, which is a liquid at room temperature and has a density of 13.6 g/mL. The density tells us that 13.6 g of mercury have a volume of 1 mL. We can write that relationship as follows:

13.6 g mercury = 1 mL mercury

This relationship can be used to construct two conversion factors:

$$\frac{13.6\ \text{g}}{1\ \text{mL}} \text{ and } \frac{1\ \text{mL}}{13.6\ \text{g}}$$

Which one do we use? It depends, as usual, on the units we need to cancel and introduce. For example, suppose we want to know the mass of 16 mL of mercury. We would use the conversion factor that has milliliters on the bottom (so that the milliliter unit cancels) and grams on top so that our final answer has a unit of mass:

$$16\ \cancel{\text{mL}} \times \frac{13.6\ \text{g}}{1\ \cancel{\text{mL}}} = 217.6\ \text{g}$$

$$\approx 220\ \text{g}$$

In the last step, we limit our final answer to two significant figures because the volume quantity has only two significant figures; the 1 in the volume unit is considered an exact number, so it does not affect the number of significant figures. The other conversion factor would be useful if we were given a mass and asked to find volume, as the following example illustrates.

Density can be used as a conversion factor between mass and volume.

Example 1.8.2: Mercury Thermometer

A mercury thermometer for measuring a patient's temperature contains 0.750 g of mercury. What is the volume of this mass of mercury?

Solution

Because we are starting with grams, we want to use the conversion factor that has grams in the denominator. The gram unit will cancel algebraically, and milliliters will be introduced in the numerator.

$$0.750\ \cancel{\text{g}} \times \frac{1\ \text{mL}}{13.6\ \cancel{\text{g}}} = 0.055147\ldots\ \text{mL}$$

$$\approx 0.0551\ \text{mL}$$

We have limited the final answer to three significant figures.

Exercise 1.8.2

What is the volume of 100.0 g of air if its density is 1.3 g/L?

Answer

$$100.0\ \cancel{\text{g}} \times \frac{1\ \text{L}}{1.3\ \cancel{\text{g}}} = 76.92307692\ \text{L} \approx 77\text{L}$$

Because the density (1.3 g/L) has only 2 significant figures, we are rounding off the final answer to 2 significant figures.

Looking Closer: Density and the Body

The densities of many components and products of the body have a bearing on our health.

Bones. Bone density is important because bone tissue of lower-than-normal density is mechanically weaker and susceptible to breaking. The density of bone is, in part, related to the amount of calcium in one's diet; people who have a diet deficient in calcium, which is an important component of bones, tend to have weaker bones. Dietary supplements or adding dairy products to the diet seems to help strengthen bones. As a group, women experience a decrease in bone density as they age. It has been estimated that fully half of women over age 50 suffer from excessive bone loss, a condition known as osteoporosis. Exact bone densities vary within the body, but for a healthy 30-year-old female, it is about 0.95–1.05 g/cm^3. Osteoporosis is diagnosed if the bone density is below 0.6–0.7 g/cm^3.

Urine. The density of urine can be affected by a variety of medical conditions. Sufferers of diabetes produce an abnormally large volume of urine with a relatively low density. In another form of diabetes, called diabetes mellitus, there is excess glucose dissolved in the urine, so that the density of urine is abnormally high. The density of urine may also be abnormally high because of excess protein in the urine, which can be caused by congestive heart failure or certain renal (kidney) problems. Thus, a urine density test can provide clues to various kinds of health problems. The density of urine is commonly expressed as a specific gravity, which is a unitless quantity defined as

$$\frac{\text{density of some material}}{\text{density of water}}$$

Normal values for the specific gravity of urine range from 1.002 to 1.028.

Body Fat. The overall density of the body is one indicator of a person's total body fat. Fat is less dense than muscle and other tissues, so as it accumulates, the overall density of the body decreases. Measurements of a person's weight and volume provide the overall body density, which can then be correlated to the percentage of body fat. (The body's volume can be measured by immersion in a large tank of water. The amount of water displaced is equal to the volume of the body.)

Problem Solving With Multiple Conversions

Sometimes you will have to perform more than one conversion to obtain the desired unit. For example, suppose you want to convert 54.7 km into millimeters. You can either memorize the relationship between kilometers and millimeters, or you can do the conversion in two steps. Most people prefer to convert in steps.

To do a stepwise conversion, we first convert the given amount to the base unit. In this example, the base unit is meters. We know that there are 1,000 m in 1 km:

$$54.7 \cancel{\text{km}} \times \frac{1,000 \text{ m}}{1 \cancel{\text{km}}} = 54,700 \text{ m}$$


Then we take the result (54,700 m) and convert it to millimeters, remembering that there are $1,000 \text{ mm}$ for every 1 m:

$$54,700 \cancel{\text{m}} \times \frac{1,000 \text{ mm}}{1 \cancel{\text{m}}} = 54,700,000 \text{ mm}$$
$$= 5.47 \times 10^7 \text{ mm}$$

We have expressed the final answer in scientific notation.

As a shortcut, both steps in the conversion can be combined into a single, multistep expression:

Concept Map



Calculation

$$54.7 \cancel{\text{km}} \times \frac{1,000 \cancel{\text{m}}}{1 \cancel{\text{km}}} \times \frac{1,000 \text{ mm}}{1 \cancel{\text{m}}} = 54,700,000 \text{ mm}$$
$$= 5.47 \times 10^7 \text{ mm}$$

In each step, the previous unit is canceled and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.

Either method—one step at a time or all the steps together—is acceptable. If you do all the steps together, the restriction for the proper number of significant figures should be done after the last step. As long as the math is performed correctly, you should get

the same answer no matter which method you use.


Example 1.8.3

Convert 58.2 ms to megaseconds in one multistep calculation.

Solution

First, convert the given unit (ms) to the base unit—in this case, seconds—and then convert seconds to the final unit, megaseconds:

Concept Map



Calculation

$$58.2 \cancel{\text{ms}} \times \frac{1 \cancel{\text{s}}}{1{,}000 \cancel{\text{ms}}} \times \frac{1 \text{ Ms}}{1{,}000{,}000 \cancel{\text{s}}} = 0.0000000582 \text{ Ms}$$

$$= 5.82 \times 10^{-8} \text{ Ms}$$

Neither conversion factor affects the number of significant figures in the final answer.

Exercise 1.8.3

Convert 43.007 mg to kilograms in one multistep calculation.

Answer

$$43.007 \cancel{\text{mg}} \times \frac{1 \cancel{\text{g}}}{1{,}000 \cancel{\text{mg}}} \times \frac{1 \text{ kg}}{1{,}000 \cancel{\text{g}}} = 0.000043007 \text{ kg}$$

$$= 4.3007 \times 10^{-5} \text{ kg}$$

.

Neither conversion factor affects the number of significant figures in the final answer.

Career Focus: Pharmacist

A pharmacist dispenses drugs that have been prescribed by a doctor. Although that may sound straightforward, pharmacists in the United States must hold a doctorate in pharmacy and be licensed by the state in which they work. Most pharmacy programs require four years of education in a specialty pharmacy school.

Pharmacists must know a lot of chemistry and biology so they can understand the effects that drugs (which are chemicals, after all) have on the body. Pharmacists can advise physicians on the selection, dosage, interactions, and side effects of drugs. They can also advise patients on the proper use of their medications, including when and how to take specific drugs properly. Pharmacists can be found in drugstores, hospitals, and other medical facilities.

Curiously, an outdated name for pharmacist is *chemist*, which was used when pharmacists formerly did a lot of drug preparation, or *compounding*. In modern times, pharmacists rarely compound their own drugs, but their knowledge of the sciences, including chemistry, helps them provide valuable services in support of everyone's health.

Key Takeaway

- A unit can be converted to another unit of the same type with a conversion factor.

Concept Review Exercises

1. How do you determine which quantity in a conversion factor goes in the denominator of the fraction?
2. State the guidelines for determining significant figures when using a conversion factor.
3. Write a concept map (a plan) for how you would convert $\mathbf{1.0 \times 10^{12}}$ *nano*liters (nL) to *kilo*liters (kL).

Answers

1. The unit you want to cancel from the numerator goes in the denominator of the conversion factor.
2. Exact numbers that appear in many conversion factors do not affect the number of significant figures; otherwise, the normal rules of multiplication and division for significant figures apply.
3. Concept Map: Convert the given (nanoliters, nL) to liters; then convert liters to kiloliters.



1.0 times 10 to the 12th nL converts to 1000 Liters converts to 1 kiloLiter. One Liter is 1.0 times 10 to the 9th nL. One kiloLiter is 1000 Liters.

1.8: Converting Units is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **1.7: Converting Units** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.9: Temperature Conversions

Learning Objectives

- To identify the different between temperature and heat
- To recognize the different scales used to measuring temperature

The concept of temperature may seem familiar to you, but many people confuse temperature with heat. **Temperature** is a measure of how hot or cold an object is relative to another object (its thermal energy content), whereas **heat** is the flow of thermal energy between objects with different temperatures.

Three different scales are commonly used to measure temperature: Fahrenheit (expressed as °F), Celsius (°C), and Kelvin (K). Thermometers measure temperature by using materials that expand or contract when heated or cooled. Mercury or alcohol thermometers, for example, have a reservoir of liquid that expands when heated and contracts when cooled, so the liquid column lengthens or shortens as the temperature of the liquid changes.

The Fahrenheit Scale

The Fahrenheit temperature scale was developed in 1717 by the German physicist Gabriel Fahrenheit, who designated the temperature of a bath of ice melting in a solution of salt as the zero point on his scale. Such a solution was commonly used in the 18th century to carry out low-temperature reactions in the laboratory. The scale was measured in increments of 12; its upper end, designated as 96°, was based on the armpit temperature of a healthy person—in this case, Fahrenheit's wife. Later, the number of increments shown on a thermometer increased as measurements became more precise. The upper point is based on the boiling point of water, designated as 212° to maintain the original magnitude of a Fahrenheit degree, whereas the melting point of ice is designated as 32°.

The Celsius Scale

The Celsius scale was developed in 1742 by the Swedish astronomer Anders Celsius. It is based on the melting and boiling points of water under normal atmospheric conditions. The current scale is an inverted form of the original scale, which was divided into 100 increments. Because of these 100 divisions, the Celsius scale is also called the *centigrade scale*.

The Kelvin Scale

Lord Kelvin, working in Scotland, developed the Kelvin scale in 1848. His scale uses molecular energy to define the extremes of hot and cold. Absolute zero, or 0 K, corresponds to the point at which molecular energy is at a minimum. The Kelvin scale is preferred in scientific work, although the Celsius scale is also commonly used. Temperatures measured on the Kelvin scale are reported simply as K, not °K.




Figure 1.9.1: A Comparison of the Fahrenheit, Celsius, and Kelvin Temperature Scales. Because the difference between the freezing point of water and the boiling point of water is 100° on both the Celsius and Kelvin scales, the size of a degree Celsius (°C) and a kelvin (K) are precisely the same. In contrast, both a degree Celsius and a kelvin are 9/5 the size of a degree Fahrenheit (°F).

Converting between Scales

The kelvin is the same size as the Celsius degree, so measurements are easily converted from one to the other. The freezing point of water is 0°C = 273.15 K; the boiling point of water is 100°C = 373.15 K. The Kelvin and Celsius scales are related as follows:

$$T \text{ (in °C)} + 273.15 = T \text{ (in K)}$$

$$T \text{ (in K)} - 273.15 = T \text{ (in °C)}$$

Degrees on the Fahrenheit scale, however, are based on an English tradition of using 12 divisions, just as 1 ft = 12 in. The relationship between degrees Fahrenheit and degrees Celsius is as follows:where the coefficient for degrees Fahrenheit is exact. (Some calculators have a function that allows you to convert directly between °F and °C.) There is only one temperature for which the numerical value is the same on both the Fahrenheit and Celsius scales: −40°C = −40°F. The relationship between the scales are as follows:

$$°C = (5/9)*(°F-32)$$

$$°F = (9/5)*(°C)+32$$

? Exercise 1.9.1

Convert the temperature of the surface of the sun (5800 K) and the boiling points of gold (3080 K) and liquid nitrogen (77.36 K) to °C and °F.A student is ill with a temperature of 103.5°F. What is her temperature in °C and K?

1.9: Temperature Conversions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **Temperature Basics** is licensed CC BY-NC-SA 4.0.

1.10: Specific Heat

Learning Objectives

- To relate heat transfer to temperature change.
- Memorize temperature equations for Celsius, Fahrenheit and Kelvin conversions.
- Understand how body temperature can vary.

The concept of temperature may seem familiar to you, but many people confuse temperature with heat. **Temperature** is a measure of how hot or cold an object is relative to another object (its thermal energy content), whereas **heat** is the flow of thermal energy between objects with different temperatures. When we touch a hot object, energy flows from the hot object into our fingers, and we perceive that incoming energy as the object being "hot." Conversely, when we hold an ice cube in our palms, energy flows from our hand into the ice cube, and we perceive that loss of energy as "cold." In both cases, the temperature of the object is different from the temperature of our hand, so we can conclude that differences in temperatures are the ultimate cause of heat transfer.

Temperature Scales

Temperature is a measure of the average amount of energy of motion, or *kinetic energy*, a system contains. Temperatures are expressed using scales that use units called degrees. There are three scales used for reporting temperatures. Figure **1.10.1** compares the three temperature scales: Fahrenheit (expressed as °F), Celsius (°C), and Kelvin (K). Thermometers measure temperature by using materials that expand or contract when heated or cooled. Mercury or alcohol thermometers, for example, have a reservoir of liquid that expands when heated and contracts when cooled, so the liquid column lengthens or shortens as the temperature of the liquid changes.




Figure **1.10.1**: Fahrenheit, Celsius, and Kelvin Temperatures. A comparison of the three temperature scales. These thermometers have a red-colored liquid which contains alcohol and food coloring. Silver looking thermometers contain mercury, which is a neurotoxin.

In the United States, the commonly used temperature scale is the *Fahrenheit scale* (symbolized by °F and spoken as "degrees Fahrenheit"). On this scale, the freezing point of liquid water (the temperature at which liquid water turns to solid ice) is 32 °F, and the boiling point of water (the temperature at which liquid water turns to steam) is 212 °F.

Science also uses other scales to express temperature. For example, the Celsius scale (symbolized by °C and spoken as "degrees Celsius") defines 0°C as the freezing point of water and 100 °C as the boiling point of water. This scale is divided into 100

divisions between these two landmarks and extended higher and lower as well. By comparing the Fahrenheit and Celsius scales, a conversion between the two scales can be determined:

$$^{\circ}C = (^{\circ}F - 32) \times \frac{5}{9} \tag{1.10.1}$$

$$^{\circ}F = \left(^{\circ}C \times \frac{9}{5}\right) + 32 \tag{1.10.2}$$

Note that science uses the Celsius and Kelvin scales almost exclusively; virtually no practicing chemist expresses laboratory-measured temperatures with the Fahrenheit scale. (In fact, the United States is one of the few countries in the world that still uses the Fahrenheit scale on a daily basis. People driving near the borders of Canada or Mexico may pick up local radio stations on the other side of the border that express the daily weather in degrees Celsius, so do not get confused by their weather reports.)

✔ Example 1.10.1: Conversions

a. What is 98.6°F in degrees Celsius?
b. What is 25.0°C in degrees Fahrenheit?

Solution

a. Using Equation 1.10.1, we have

$$\begin{aligned} ^{\circ}C &= (98.6 - 32) \times \frac{5}{9} \\ &= 66.6 \times \frac{5}{9} \\ &= 37.0^{\circ}C \end{aligned}$$

b. Using Equation 1.10.2, we have

$$\begin{aligned} ^{\circ}F &= (25.0 \times \frac{9}{5}) + 32 \\ &= 45.0 + 32 \\ &= 77.0^{\circ}F \end{aligned}$$

For more examples of how to perform these types of problems, click on this video to see your professor in action.

? Exercise 1.10.1

a. Convert 0°F to degrees Celsius.
b. Convert 212°C to degrees Fahrenheit.

Answer a

−17.8°C

Answer b

414°F

The fundamental unit of temperature in SI is the Kelvin (K). The Kelvin temperature scale (note that the name of the scale capitalizes the word *Kelvin*, but the unit itself is lowercase) uses degrees that are the same size as the Celsius degree, but the numerical scale is shifted up by 273.15 units. That is, the conversion between the Kelvin and Celsius scales is as follows:

$$K =^{\circ} C + 273.15 \tag{1.10.3}$$

For most purposes, it is acceptable to use 273 instead of 273.15 in Equation 1.10.3.

Note that the Kelvin scale does not use the word *degrees*; a temperature of 295 K is spoken of as “two hundred ninety-five kelvin” and not “two hundred ninety-five degrees Kelvin.”

The reason that the Kelvin scale is defined this way is that there exists a minimum possible temperature called absolute zero (zero Kelvin). The Kelvin temperature scale is set so that 0 K is absolute zero, and the temperature is counted upward from there. Normal room temperature is about 295 K, as seen in the following example.

Example 1.10.2: Room Temperature

If the normal room temperature is 72.0°F, what is room temperature in degrees Celsius and kelvin?

Solution

First, we use Equation 1.10.1 to determine the temperature in degrees Celsius:

$$\begin{aligned} ^\circ C &= (72.0 - 32) \times \frac{5}{9} \\ &= 40.0 \times \frac{5}{9} \\ &= 22.2^\circ C \end{aligned}$$

Then we use Equation 1.10.3 to determine the temperature in the Kelvin scale:

$$\begin{aligned} K &= 22.2^\circ C + 273.15 \\ &= 295.4 K \end{aligned}$$

So, room temperature is about 295 K.

Health Application: Body temperature

Normal body temperature is defined as being 98.6°F (+/- 1.0°F). To determine body temperature, thermometers can be placed inside or on the surface of the body. The two best methods of obtaining body temperature are placing the thermometer either under the tongue or inside the rectum. Typically, children are capable of holding a thermometer in their mouths around the age of four (have fun before that age).

Fever is defined as body temperature being above 100°F (adults). High fever status occurs at and above 104°F. For adults, these adults should seek medical attention immediately if the fever exceeds 104°F. For children, these values are much lower.

Hyperthermia (inability of the body to regulate heat) occurs when normal body temperature is exceeded. Conditions that can cause hyperthermia are fever (infection), heat stroke, thyroid disorders, heart attack, or traumatic injury. Medications for cancer, arthritis, and thyroid patients can cause the body temperature to rise. Symptoms of hyperthermia include sweating, confusion, nausea, and dizziness.

Hypothermia (exposure to cold environments) occurs when normal body temperature dips below 95.0°F. When this occurs, the affected person(s) should seek immediate medical attention. During hypothermia, the body has problems producing heat. Medical conditions like diabetes, infection, and thyroid dysfunction can cause hypothermia. Watch this video of American Marines attempting to survive extreme environments. Symptoms of this condition involve shivering, confusion, and sluggish behavior.

Treating hyperthermia could involve hydrating a patient. If a fever is due to infection, analgesics (like Tylenol, Advil, aspirin, or Aleve) can help alleviate fever. Placing someone in a cool bath can also relieve symptoms.

To combat hypothermia, one must remove wet clothing, redress in warm materials, and participate in physical movement.

Heat transfer

Heat is a familiar manifestation of energy. When we touch a hot object, energy flows from the hot object into our fingers, and we perceive that incoming energy as the object being “hot.” Conversely, when we hold an ice cube in our palms, energy flows from

our hand into the ice cube, and we perceive that loss of energy as “cold.” In both cases, the temperature of the object is different from the temperature of our hand, so we can conclude that differences in temperatures are the ultimate cause of heat transfer.

Suppose we consider the transfer of heat from the opposite perspective—namely, what happens to a system that gains or loses heat? Generally, the system’s temperature changes. (We will address a few exceptions later.) The greater the original temperature difference, the greater the transfer of heat, and the greater the ultimate temperature change. The relationship between the amount of heat transferred and the temperature change can be written as

$$\text{heat} \propto \Delta T \tag{1.10.4}$$

where ∝ means “is proportional to” and **ΔT is the change in temperature of the system**. Any change in a variable is always defined as “the final value minus the initial value” of the variable, so ΔT is $T_{\text{final}} - T_{\text{initial}}$. In addition, the greater the mass of an object, the more heat is needed to change its temperature. We can include a variable representing mass (m) to the proportionality as follows:

$$\text{heat} \propto m\Delta T \tag{1.10.5}$$

To change this proportionality into an equality, we include a proportionality constant. The proportionality constant is called the specific heat and is commonly symbolized by c:

$$\text{heat} = mc\Delta T \tag{1.10.6}$$

Every substance has a characteristic **specific heat**, which is reported in units of cal/g•°C or cal/g•K, depending on the units used to express ΔT. The specific heat of a substance is the amount of energy that must be transferred to or from 1 g of that substance to change its temperature by 1°C. Table **1.10.1** lists the specific heats for various materials.

Table **1.10.1**: Specific Heats of Selected Substances

Substance	c (cal/g•°C)
aluminum (Al)	0.215
aluminum oxide (Al_2O_3)	0.305
benzene (C_6H_6)	0.251
copper (Cu)	0.092
ethanol (C_2H_6O)	0.578
hexane (C_6H_{14})	0.394
hydrogen (H_2)	3.419
ice [$H_2O(s)$]	0.492
iron (Fe)	0.108
iron(III) oxide (Fe_2O_3)	0.156
mercury (Hg)	0.033
oxygen (O_2)	0.219
sodium chloride (NaCl)	0.207
steam [$H_2O(g)$]	0.488
water [$H_2O(\ell)$]	1.00

> *The proportionality constant c is sometimes referred to as the specific heat capacity or (incorrectly) the heat capacity.*

The *direction* of heat flow is not shown in heat = $mc\Delta T$. If energy goes into an object, the total energy of the object increases, and the values of heat ΔT are positive. If energy is coming out of an object, the total energy of the object decreases, and the values of heat and ΔT are negative.

Example 1.10.3

What quantity of heat is transferred when a 150.0 g block of iron metal is heated from 25.0°C to 73.3°C? What is the direction of heat flow?

Solution

We can use Equation 1.10.6 to determine the amount of heat, but first we need to determine ΔT. Because the final temperature of the iron is 73.3°C and the initial temperature is 25.0°C, ΔT is as follows:

$$\begin{aligned}\Delta T &= T_{final} - T_{initial} \\ &= 73.3^{o}C - 25.0^{o}C \\ &= 48.3^{o}C\end{aligned}$$

The mass is given as 150.0 g, and Table 1.10.1 gives the specific heat of iron as 0.108 cal/g•°C. Substitute the known values into Equation 1.10.6 and solve for amount of heat:

$$\text{heat} = (150.0\ \text{g})\left(0.108\ \frac{\text{cal}}{\text{g}\cdot{}^{\circ}\text{C}}\right)(48.3^{\circ}\text{C}) = 782\ \text{cal}$$

Note how the gram and °C units cancel algebraically, leaving only the calorie unit, which is a unit of heat. Because the temperature of the iron increases, energy (as heat) must be flowing *into* the metal.

1.10.3 Exercise

What quantity of heat is transferred when a 295.5 g block of aluminum metal is cooled from 128.0°C to 22.5°C? What is the direction of heat flow?

Answer

6,700 cal

Example 1.10.4

A 10.3 g sample of a reddish-brown metal gave off 71.7 cal of heat as its temperature decreased from 97.5°C to 22.0°C. What is the specific heat of the metal? Can you identify the metal from the data in Table 1.10.1?

Solution

The question gives us the heat, the final and initial temperatures, and the mass of the sample. The value of ΔT is as follows:

$$\Delta T = T_{\text{final}} - T_{\text{initial}} = 22.0^{\circ}\text{C} - 97.5^{\circ}\text{C} = -75.5^{\circ}\text{C}$$

If the sample gives off 71.7 cal, it loses energy (as heat), so the value of heat is written as a negative number, −71.7 cal. Substitute the known values into heat = $mc\Delta T$ and solve for c:

$$-71.7\ \text{cal} = (10.3\ \text{g})(c)(-75.5^{\circ}\text{C})$$

$$c = \frac{-71.7\ \text{cal}}{(10.3\ \text{g})(-75.5^{\circ}\text{C})}$$

$$c = 0.0923\ \text{cal/g}\cdot{}^{\circ}\text{C}$$

This value for specific heat is very close to that given for copper in Table 1.10.1.

1.10.4 Exercise

A 10.7 g crystal of sodium chloride (NaCl) had an initial temperature of 37.0°C. What is the final temperature of the crystal if 147 cal of heat were supplied to it?

Answer

103.4°C

Notice that water has a very high specific heat compared to most other substances. Water is commonly used as a coolant for machinery because it is able to absorb large quantities of heat (see table above). Coastal climates are much more moderate than inland climates because of the presence of the ocean. Water in lakes or oceans absorbs heat from the air on hot days and releases it back into the air on cool days.



Figure **1.10.2**: This power plant in West Virginia, like many others, is located next to a large lake so that the water from the lake can be used as a coolant. Cool water from the lake is pumped into the plant, while warmer water is pumped out of the plant and back into the lake.

Summary

- Heat transfer is related to temperature change.
- The specific heat of a substance is the amount of energy required to raise the temperature of 1 gram of the substance by **1°C**.

1.10: Specific Heat is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **7.2: Heat and Temperature** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.11: Dosage Calculations

A medicine can be more harmful than helpful if it is not taken in the proper dosage. A dosage (or dose) is the specific amount of a medicine that is known to be therapeutic for an ailment in a patient of a certain size. Dosages of the active ingredient in medications are usually described by units of mass, typically grams or milligrams, and generally are equated with a number of capsules or teaspoonfuls to be swallowed or injected. The amount of the active ingredient in a medicine is carefully controlled so that the proper number of pills or spoonfuls contains the proper dose.

Most drugs must be taken in just the right amount. If too little is taken, the desired effects will not occur (or will not occur fast enough for comfort); if too much is taken, there may be potential side effects that are worse than the original ailment. Some drugs are available in multiple dosages. For example, tablets of the medication levothyroxine sodium, a synthetic thyroid hormone for those suffering from decreased thyroid gland function, are available in 11 different doses, ranging from 25 micrograms (µg) to 300 µg. It is a doctor's responsibility to prescribe the correct dosage for a patient, and it is a pharmacist's responsibility to provide the patient with the correct medicine at the dosage prescribed. Thus, proper quantities—which are expressed using numbers and their associated units—are crucial for keeping us healthy.

Effects are dose-dependent

Chemicals are the most common things for which doses are measured, but there are others, such as radiation exposure. For humans, most doses of micronutrients and medications are measured in milligrams (mg), but some are measured in micrograms because of their potency. Nonmedicinal poisons span the measurement scale; some poisons are so dangerous that a single microgram of it could be deadly, whereas other substances take much more. For example, even water is toxic when consumed in large enough quantities.

- Dosage (the size of each dose) determines the strength and duration of the health benefits of nutrients, and also of the therapeutic effects of medical treatments. Dosage also determines the severity of adverse effects of treatments and toxins.
- Duration of exposure, that is, the period of time over which the dose was received (all at once or gradually) also determines its effects (the body may build tolerance to gradual exposure to a drug, while a large immediate dose could be deadly).
- The route by which a dose is exposed to, may affect the outcome, because some medications have different effects depending on whether they are inhaled, ingested, taken transdermally, injected, or inserted.

The dosage, route, concentration, and division over time may all be critical considerations in the administering of drugs, or in responding to exposure to a toxin. In nutrition, the route is usually a given, as nutrients are generally eaten; while dosage and the frequency of ingestion of nutrients are very important variables in preventing disease and promoting overall health.

> *Calculating drug dosages for humans based on the doses used in animal studies can be based on weight (e.g., mg/kg) or surface area (e.g., mg/m^2) based on weight. Desirable dosages may also vary among different persons according to body weight and other factors.*

Biological agents

Biological agents (bacteria, viruses, parasites) may have different dosage units. This is because it is the ability of the organism to cause effects that is the important unit, not a specific quantity by weight, volume, or even numerical count. Often, the unit used is CFU (colony forming units), which is proportionate to the number of organisms present multiplied times the number able to reproduce on a culture medium such as a Petri dish.

Toxicology

In the realm of toxicology, several measures are commonly used to describe toxic dosages according to the degree of effect on an organism or a population, and some are specifically defined by various laws or organizational usage. These include:

- LD50 = Median lethal dose, a dose that will kill 50% of an exposed population
- NOEL = No Observed Effect Level, the highest dose known to show no effect
- NOAEL = No Observed Adverse Effect Level, the highest dose known to show no adverse effects

- PEL = Personal Exposure Limit, the highest concentration permitted under US OSHA regulations
- STEL = Short-Term Exposure Limit, the highest concentration permitted for short periods of time, in general 15–30 minutes
- TWA = Time-Weighted Average, the average amount of an agent's concentration over a specified period of time, usually 8 hours.

References

1. http://www.unc.edu/~bangel/quiz/quiz5.htm
2. http://cancerres.aacrjournals.org/cg...t/18/7/853.pdf "The Use of Body Surface Area as a Criterion of Drug Dosage in Cancer Chemotherapy" D Pinkel. Cancer Research 1958

Contributors

- Wikipedia
- Anonymous

1.11: Dosage Calculations is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

1.12: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Chemistry is the study of **matter**, which is anything that has mass and takes up space. Chemistry is one branch of **science**, which is the study of the natural universe. Like all branches of science, chemistry relies on the **scientific method**, which is a process of learning about the world around us. In the scientific method, a guess or **hypothesis** is tested through experiment and measurement.

Matter can be described in a number of ways. **Physical properties** describe characteristics of a sample that do not change the chemical identity of the material (size, shape, color, and so on), while **chemical properties** describe how a sample of matter changes its chemical composition. A **substance** is any material that has the same physical and chemical properties throughout. An **element** is a substance that cannot be broken down into chemically simpler components. The smallest chemically identifiable piece of an element is an **atom**. A substance that can be broken down into simpler chemical components is a **compound**. The smallest chemically identifiable piece of a compound is a **molecule**. Two or more substances combine physically to make a **mixture**. If the mixture is composed of discrete regions that maintain their own identity, the mixture is a **heterogeneous mixture**. If the mixture is so thoroughly mixed that the different components are evenly distributed throughout, it is a **homogeneous mixture**. Another name for a homogeneous mixture is a **solution**. Substances can also be described by their **phase**: solid, liquid, or gas.

Scientists learn about the universe by making measurements of **quantities**, which consist of **numbers** (how many) and **units** (of what). The numerical portion of a quantity can be expressed using **scientific notation**, which is based on **powers**, or exponents, of 10. Large numbers have positive powers of 10, while numbers less than 1 have negative powers of 10. The proper reporting of a measurement requires proper use of **significant figures**, which are all the known digits of a measurement plus the first estimated digit. The number of figures to report in the result of a calculation based on measured quantities depends on the numbers of significant figures in those quantities. For addition and subtraction, the number of significant figures is determined by position; for multiplication and division, it is decided by the number of significant figures in the original measured values. Nonsignificant digits are dropped from a final answer in accordance with the rules of **rounding**.

Chemistry uses SI, a system of units based on seven **basic units**. The most important ones for chemistry are the units for length, mass, amount, time, and temperature. Basic units can be combined with numerical prefixes to change the size of the units. They can also be combined with other units to make **derived units**, which are used to express other quantities such as **volume**, **density**, or **energy**. A formal conversion from one unit to another uses a **conversion factor**, which is constructed from the relationship between the two units. Numbers in conversion factors may affect the number of significant figures in a calculated quantity, depending on whether the conversion factor is **exact**. Conversion factors can be applied in separate computations, or several can be used at once in a single, longer computation.

Additional Exercises

1. A sample of urine has a density of 1.105 g/cm^3. What is the mass of 0.255 L of this urine?
2. The hardest bone in the body is tooth enamel, which has a density of 2.91 g/cm^3. What is the volume, in liters, of 75.9 g of tooth enamel?
3. Some brands of aspirin have 81 mg of aspirin in each tablet. If a person takes 8 tablets per day, how many grams of aspirin is that person consuming every day?
4. The US government has a recommended daily intake (RDI) of 5 µg of vitamin D per day. (The name *recommended daily allowance* was changed to RDI in 1997.) If milk contains 1.2 µg per 8 oz glass, how many ounces of milk are needed to supply the RDI of vitamin D?
5. The population of the United States, according to the 2000 census, was 281.4 million people.
   a. How many significant figures does this number have?
   b. What is the unit in this quantity?
   c. Express this quantity in proper scientific notation.

6. The United States produces 24,800,000,000 lb of sugar each year, and much of it is exported to other countries.
   a. How many significant figures does this number have?
   b. What is the unit in this quantity?
   c. Express this quantity in proper scientific notation.
7. Construct a conversion factor that can convert from one unit to the other in each pair of units.
   a. from millimeters to kilometers
   b. from kilograms to micrograms
   c. from centimeters to micrometers
8. Construct a conversion factor that can convert from one unit to the other in each pair of units.
   a. from kilometers to micrometers
   b. from decaliters to milliliters
   c. from megagrams to milligrams
9. What is the density of a dextrose solution if 355 mL of the solution has a mass of 406.9 g?
10. What is the density of a dental amalgam (an alloy used to fill cavities) if 1.005 kg of the material has a volume of 433 mL? Express your final answer in grams per milliliter.

For Exercises 11–16, see the accompanying table for the relationships between English and SI units.

1 m ≈ 39.36 in. ≈ 3.28 ft ≈ 1.09 yd
1 cm ≈ 2.54 in.
1 km ≈ 0.62 mi
1 kg ≈ 2.20 lb
1 lb ≈ 454 g
1 L ≈ 1.06 qt
1 qt ≈ 0.946 L

11. Approximately how many inches are in 4.76 m?
12. Approximately how many liters are in 1 gal, which is exactly 4 qt?
13. Approximately how many kilograms are in a person who weighs 170 lb?
14. The average distance between Earth and the sun is 9.3×10^7 mi. How many kilometers is that?
15. Show mathematically that 1 L equals 1 dm^3.
16. Show mathematically that 1 L equals 1,000 cm^3.

1.12: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

2: Atomic Structure

Just as a language has an alphabet from which words are built, chemistry has an alphabet from which matter is described. However, the chemical alphabet is larger than the one we use for spelling. You may have already figured out that the chemical alphabet consists of the chemical elements. Their role is central to chemistry, for they combine to form the millions and millions of known compounds.

2.1: The Elements
2.2: Atomic Theory
2.3: The Structure of Atoms
2.4: Nuclei of Atoms
2.5: Arrangement of Electron (Shell Model)
2.6: The Periodic Table
2.7: Chapter Summary

2: Atomic Structure is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

2.1: The Elements

Learning Objectives

- Define a chemical element and give examples of the abundance of different elements.
- Represent a chemical element with a chemical symbol.

An element is a substance that cannot be broken down into simpler chemical substances. There are about 90 naturally occurring elements known on Earth. Using technology, scientists have been able to create nearly 30 additional elements that do not occur in nature. Today, chemistry recognizes 118 elements—some of which were created an atom at a time. Figure **2.1.1** shows some of the chemical elements.



Figure **2.1.1**: Samples of Elements. Gold is a yellowish solid, iron is a silvery solid, while mercury is a silvery liquid at room temperature. © Thinkstock

Abundance

The elements vary widely in abundance. In the universe as a whole, the most common element is hydrogen (about 90% of atoms), followed by helium (most of the remaining 10%). All other elements are present in relatively minuscule amounts, as far as we can detect.

Table **2.1.1**: Elemental Composition of Earth

Earth's Crust		Earth (overall)	
Element	**Percentage**	**Element**	**Percentage**
oxygen	46.1	iron	34.6
silicon	28.2	oxygen	29.5
aluminum	8.23	silicon	15.2
iron	5.53	magnesium	12.7
calcium	4.15	nickel	2.4
sodium	2.36	sulfur	1.9
magnesium	2.33	all others	3.7
potassium	2.09		
titanium	0.565		
hydrogen	0.14		
phosphorus	0.105		
all others	0.174		

Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 14–17.

On the planet Earth, however, the situation is rather different. Oxygen makes up 46.1% of the mass of Earth's crust (the relatively thin layer of rock forming Earth's surface), mostly in combination with other elements, while silicon makes up 28.2%. Hydrogen, the most abundant element in the universe, makes up only 0.14% of Earth's crust. Table **2.1.1** lists the relative abundances of elements on Earth as a whole and in Earth's crust. Table **2.1.2** lists the relative abundances of elements in the human body. If you compare Table **2.1.1** and Table **2.1.2**, you will find disparities between the percentage of each element in the human body and on Earth. Oxygen has the highest percentage in both cases, but carbon, the element with the second highest percentage in the body, is relatively rare on Earth and does not even appear as a separate entry in Table **2.1.1**; carbon is part of the 0.174% representing "other" elements. How does the human body concentrate so many apparently rare elements?

Table **2.1.2**: Elemental Composition of a Human Body

Element	Percentage by Mass
oxygen	61
carbon	23
hydrogen	10
nitrogen	2.6
calcium	1.4
phosphorus	1.1
sulfur	0.20
potassium	0.20
sodium	0.14
chlorine	0.12
magnesium	0.027
silicon	0.026
iron	0.006
fluorine	0.0037
zinc	0.0033
all others	0.174

Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 7–24.

The relative amounts of elements in the body have less to do with their abundances on Earth than with their availability in a form we can assimilate. We obtain oxygen from the air we breathe and the water we drink. We also obtain hydrogen from water. On the other hand, although carbon is present in the atmosphere as carbon dioxide, and about 80% of the atmosphere is nitrogen, we obtain those two elements from the food we eat, not the air we breathe.

LOOKING CLOSER: PHOSPHOROUS, THE CHEMICAL BOTTLENECK

There is an element that we need more of in our bodies than is proportionately present in Earth's crust, and *this* element is not easily accessible. Phosphorus makes up 1.1% of the human body but only 0.105% of Earth's crust. We need phosphorus for our bones and teeth, and it is a crucial component of all living cells. Unlike carbon, which can be obtained from carbon dioxide through photosynthesis, there is no phosphorus compound present in our surroundings that can serve as a convenient source. Phosphorus, then, is nature's bottleneck. Its availability limits the amount of life our planet can sustain.

Higher forms of life, such as humans, can obtain phosphorus by selecting a proper diet (plenty of protein); but lower forms of life, such as algae, must absorb it from the environment. When phosphorus-containing detergents were introduced in the 1950s, wastewater from normal household activities greatly increased the amount of phosphorus available to algae and other plant life. Lakes receiving this wastewater experienced sudden increases in growth of algae. When the algae died, concentrations of bacteria that ate the dead algae increased. Because of the large bacterial concentrations, the oxygen content of the water dropped, causing fish to die in large numbers. This process, called *eutrophication*, is considered a negative environmental impact.

Today, many detergents are made without phosphorus so the detrimental effects of eutrophication are minimized. You may even see statements to that effect on detergent boxes. It can be sobering to realize how much impact a single element can have on life—or the ease with which human activity can affect the environment.

Names and Symbols

Each element has a name. Some of these names date from antiquity, while others are quite new. Today, the names for new elements are proposed by their discoverers but must be approved by the International Union of Pure and Applied Chemistry, an international organization that makes recommendations concerning all kinds of chemical terminology.

> *Today, new elements are usually named after famous scientists.*

The names of the elements can be cumbersome to write in full, especially when combined to form the names of compounds. Therefore, each element name is abbreviated as a one- or two-letter chemical symbol. By convention, the first letter of a chemical symbol is a capital letter, while the second letter (if there is one) is a lowercase letter. The first letter of the symbol is usually the first letter of the element's name, while the second letter is some other letter from the name. Some elements have symbols that derive from earlier, mostly Latin names, so the symbols may not contain any letters from the English name. Table **2.1.3** lists the names and symbols of some of the most familiar elements.

Table **2.1.3**: Element Names and Symbols

aluminum	Al	magnesium	Mg
argon	Ar	manganese	Mn
arsenic	As	mercury	Hg*
barium	Ba	neon	Ne
bismuth	Bi	nickel	Ni
boron	B	nitrogen	N
bromine	Br	oxygen	O
calcium	Ca	phosphorus	P
carbon	C	platinum	Pt
chlorine	Cl	potassium	K*
chromium	Cr	silicon	Si
copper	Cu*	silver	Ag*
fluorine	F	sodium	Na*
gold	Au*	strontium	Sr
helium	He	sulfur	S
hydrogen	H	tin	Sn*
iron	Fe	tungsten	W†
iodine	I	uranium	U
lead	Pb*	zinc	Zn
lithium	Li	zirconium	Zr

*The symbol comes from the Latin name of element.

†The symbol for tungsten comes from its German name—*wolfram*.

Element names in languages other than English are often close to their Latin names. For example, gold is *oro* in Spanish and *or* in French (close to the Latin *aurum*), tin is *estaño* in Spanish (compare to *stannum*), lead is *plomo* in Spanish and *plomb* in French (compare to *plumbum*), silver is *argent* in French (compare to *argentum*), and iron is *fer* in French and *hierro* in Spanish (compare to *ferrum*). The closeness is even more apparent in pronunciation than in spelling.

Example 2.1.1

Write the chemical symbol for each element without consulting Table **2.1.3** "Element Names and Symbols".

a. bromine
b. boron
c. carbon
d. calcium
e. gold

Answer a

Br

Answer b

B

Answer c

C

Answer d

Ca

Answer e

Au

Exercise 2.1.1

Write the chemical symbol for each element without consulting Table 2.1.3.

a. manganese
b. magnesium
c. neon
d. nitrogen
e. silver

Answer a

Mn

Answer b

Mg

Answer c

Ne

Answer d

N

Answer e

Ag

Example 2.1.2

What element is represented by each chemical symbol?

a. Na
b. Hg
c. P
d. K
e. I

Answer a

sodium

Answer b

mercury

Answer c

phosphorus

Answer d

potassium

Answer e

iodine

? Exercise 2.1.2

What element is represented by each chemical symbol?

a. Pb
b. Sn
c. U
d. O
e. F

Answer a

lead

Answer b

tin

Answer c

uranium

Answer d

oxygen

Answer e

fluorine

Concept Review Exercises

1. What is an element?
2. Give some examples of how the abundance of elements varies.
3. Why are chemical symbols so useful? What is the source of the letter(s) for a chemical symbol?

Answers

1. An element is the basic chemical building block of matter; it is the simplest chemical substance.
2. Elements vary from being a small percentage to more than 30% of the atoms around us.
3. Chemical symbols are useful to concisely represent the elements present in a substance. The letters usually come from the name of the element.

Key Takeaways

- All matter is composed of elements.
- Chemical elements are represented by a one- or two-letter symbol.

2.1: The Elements is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.1: The Elements** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.2: Atomic Theory

Learning Objectives

- Explain how all matter is composed of atoms.
- Describe the modern atomic theory.
- Recognize which elements exist as diatomic molecules.

Take some aluminum foil. Cut it in half. Now you have two smaller pieces of aluminum foil. Cut one of the pieces in half again. Cut one of those smaller pieces in half again. Continue cutting, making smaller and smaller pieces of aluminum foil.

It should be obvious that the pieces are still aluminum foil; they are just becoming smaller and smaller. But how far can you take this exercise, at least in theory? Can you continue cutting the aluminum foil into halves forever, making smaller and smaller pieces? Or is there some limit, some absolute smallest piece of aluminum foil? (Thought experiments like this—and the conclusions based on them—were debated as far back as the fifth century BC.)

The modern atomic theory, proposed about 1803 by the English chemist John Dalton (Figure **2.2.1**), is a fundamental concept that states that all elements are composed of atoms. Previously, we defined an atom as the smallest part of an element that maintains the identity of that element. Individual atoms are extremely small; even the largest atom has an approximate diameter of only 5.4×10^{-10} m. With that size, it takes over 18 million of these atoms, lined up side by side, to equal the width of your little finger (about 1 cm).




Figure **2.2.1** John Dalton was an English scientist who enunciated the modern atomic theory.

Most elements in their pure form exist as **individual atoms**. For example, a macroscopic chunk of iron (Fe) metal is composed, microscopically, of individual atoms. Some elements, however, exist as groups of atoms called molecules. Several important elements exist as two-atom combinations and are called **diatomic molecules**. In representing a diatomic molecule, we use the symbol of the element and include the ***subscript*** "2" to indicate that two atoms of that element are joined together. The elements that exist as diatomic molecules are hydrogen (H_2), nitrogen (N_2), fluorine (F_2), oxygen (O_2), iodine (I_2), chlorine (Cl_2) and bromine (Br_2). A few other elements can exist as 3-atom molecules like ozone (O_3) and 4-atom molecules like phosphorus (P_4). The most common form of the element sulfur is composed of molecules that consist of eight atoms of sulfur; its molecular formula is S_8 (Figure **2.2.2**).




Figure **2.2.2**: A molecule of sulfur is composed of eight sulfur atoms and is therefore written as S_8. It can be represented as (a) a structural formula, (b) a ball-and-stick model, and (c) a space-filling model. Sulfur atoms are represented by yellow spheres.

Figure A shows eight sulfur atoms, symbolized with the letter S, that are bonded to each other to form an octagon. Figure B shows a 3-D, ball-and-stick model of the arrangement of the sulfur atoms. The shape is clearly not octagonal as it is represented in the structural formula. Figure C is a space-filling model that shows each sulfur atom is partially embedded into the sulfur atom it bonds with.

It is important to note that a subscript following a symbol and a number in front of a symbol do not represent the same thing; for example, H_2 and 2H represent distinctly different species. H_2 is a molecular formula; it represents a diatomic molecule of hydrogen, consisting of two atoms of the element that are chemically bonded together. The expression 2H, on the other hand, indicates two separate hydrogen atoms that are **not** combined as a unit. The expression $2H_2$ represents two molecules of diatomic hydrogen (Figure **2.2.3**).




Figure **2.2.3**: The symbols H, 2H, H_2, and $2H_2$ represent very different entities.

This figure shows four diagrams. The diagram for H shows a single, white sphere and is labeled one H atom. The diagram for 2 H shows two white spheres that are not bonded together. It is labeled 2 H atoms. The diagram for H subscript 2 shows two white spheres bonded together. It is labeled one H subscript 2 molecule. The diagram for 2 H subscript 2 shows two sets of bonded, white spheres. It is labeled 2 H subscript 2 molecules.

✔ Example 2.2.1

Write the chemical formula of the following elements:

a. oxygen
b. carbon
c. potassium

Solution

Memorizing the diatomic molecules is worthwhile for our future endeavors. A mnemonic device to help in the memorization of the diatomic elements is as follows: **H**ave **N**o **F**ear **O**f **I**ce **C**old **B**eer. (hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine).

a. Oxygen is one of the seven diatomic molecular elements. Its chemical formula is O_2.
b. Carbon is monatomic, hence its formula is C.
c. Potassium is monatomic hence its formula is K.

2.2.1 Exercise

Write the chemical formula of the following elements:

a. hydrogen
b. nitrogen
c. neon

Answer a

Hydrogen is one of the seven diatomic molecular elements. Its chemical formula is H_2.

Answer b

Nitrogen is one of the seven diatomic molecular elements. Its chemical formula is N_2.

Answer c

Neon is a monatomic element, hence its formula is Ne.

Looking Closer: Atomic Theory

Dalton's ideas are called the *modern* atomic theory because the concept of atoms is very old. The Greek philosophers Leucippus and Democritus originally introduced atomic concepts in the fifth century BC. (The word *atom* comes from the Greek word *atomos*, which means "indivisible" or "uncuttable.") Dalton had something that the ancient Greek philosophers didn't have, however; he had experimental evidence, such as the formulas of simple chemicals and the behavior of gases. In the 150 years or so before Dalton, natural philosophy had been maturing into modern science, and the scientific method was being used to study nature. So when Dalton announced a modern atomic theory, he was proposing a fundamental theory to describe many previous observations of the natural world; he was not just participating in a philosophical discussion.

Concept Review Exercises

1. What is the modern atomic theory?
2. What are atoms?

Answers

1. The modern atomic theory states that all matter is composed of atoms.
2. Atoms are the smallest parts of an element that maintain the identity of that element.

Key Takeaways

- **Atoms** are the ultimate building blocks of all matter.
- The modern atomic theory establishes the concepts of **atoms** and how they compose matter.
- The **diatomic** elements are hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine.

2.2: Atomic Theory is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.2: Atomic Theory** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.3: The Structure of Atoms

Learning Objectives

- Describe the three main subatomic particles.
- State how the subatomic particles are arranged in atoms.

There have been several minor but important modifications to Dalton's atomic theory. For one thing, Dalton considered atoms to be indivisible. We know now that atoms not only can be divided but also are composed of three different kinds of particle, *subatomic particles*, with their own properties that are different from the chemical properties of atoms.

Subatomic Particles

The first subatomic particle was identified in 1897 and called the electron. It is an extremely tiny particle, with a mass of about 9.109×10^{-31} kg. Experiments with magnetic fields showed that the electron has a negative electrical charge. By 1920, experimental evidence indicated the existence of a second particle. A proton has the same amount of charge as an electron, but its charge is positive, not negative. Another major difference between a proton and an electron is mass. Although still incredibly small, the mass of a proton is 1.673×10^{-27} kg, which is almost 2,000 times greater than the mass of an electron. Because opposite charges attract each other (while like charges repel each other), protons attract electrons (and vice versa).

Finally, additional experiments pointed to the existence of a third particle. Evidence produced in 1932 established the existence of the neutron, a particle with about the same mass as a proton but with no electrical charge, it is *neutral*. We understand now that all atoms can be broken down into subatomic particles: protons, neutrons, and electrons. Table **2.3.1** lists some of their important characteristics and the symbols used to represent each particle.

Table **2.3.1**: Properties of the Subatomic Particles

Particle	Symbol	Mass (kg)	Relative Mass (proton = 1)	Relative Charge
proton	p^+	1.673×10^{-27}	1	+1
neutron	n^0	1.675×10^{-27}	1	0
electron	e^-	9.109×10^{-31}	0.00055	−1

The Nucleus

How are these subatomic particles arranged? Between 1909 and 1911, Ernest Rutherford, a Cambridge physicist, and his associates Hans Geiger and Ernest Marsden performed experiments that provided strong evidence concerning the internal structure of an atom. They took a very thin metal foil, such as gold or platinum, and aimed a beam of positively charged particles (called alpha particles, which are combinations of two protons and two neutrons) from a radioactive source toward the foil. Surrounding the foil was a detector—either a scintillator (a material that glows when hit by such particles) or some unexposed film (which is exposed where the particles hit it). The detector allowed the scientists to determine the distribution of the alpha particles after they interacted with the foil. Figure **2.3.1** shows a diagram of the experimental setup.




Figure **2.3.1**: The Geiger-Marsden Experimental Setup. Experiments using this setup were used to investigate the structure of atoms. In this experiment, most of the particles traveled straight through the foil, but some alpha particles were deflected off to one side. Some were even deflected back toward the source. This was unexpected. Rutherford once said, "It was almost as incredible as if you fired a 15-inch shell at a piece of tissue paper and it came back and hit you." (CC SA-BY 3.0; Kurzon).

Rutherford proposed the following model to explain these experimental results. Protons and neutrons are concentrated in a central region he called the nucleus (plural, *nuclei*) of the atom. Electrons are outside the nucleus and orbit about it because they are attracted to the positive charge in the nucleus. Most of the mass of an atom is in the nucleus, while the orbiting electrons account for an atom's size. As a result, an atom consists largely of empty space. Rutherford called his description the "planetary model" of the atom. Figure **2.3.2** shows how this model explains the experimental results.




Figure **2.3.2**: Rutherford's Metal Foil Experiments. Rutherford explained the results of the metal foil experiments by proposing that most of the mass and the positive charge of an atom are located in its nucleus, while the relatively low-mass electrons orbit about the nucleus. Most alpha particles go straight through the empty space, a few particles are deflected, and fewer still ricochet back toward the source. The nucleus is much smaller proportionately than depicted here.

> *The planetary model of the atom replaced the plum pudding model, which had electrons floating around aimlessly like plums in a "pudding" of positive charge.*

Rutherford's model is essentially the same model that we use today to describe atoms but with one important modification. The planetary model suggests that electrons occupy certain specific, circular orbits about the nucleus. We know now that this model is overly simplistic. A better description is that electrons form fuzzy clouds around nuclei. Figure **2.3.3** shows a more modern version of our understanding of atomic structure.




Figure **2.3.3**: A Modern Depiction of Atomic Structure. A more modern understanding of atoms, reflected in these representations of the electron in a hydrogen atom, is that electrons occupy regions of space about the nucleus; they are not in discrete orbits like planets around the sun. (a) The darker the color, the higher the probability that an electron will be at that point. (b) In a two-dimensional cross section of the electron in a hydrogen atom, the more crowded the dots, the higher the probability that an electron will be at that point. In both (a) and (b), the nucleus is in the center of the diagram.

Concept Review Exercises

1. What are the charges and the relative masses of the three subatomic particles?
2. Describe the structure of an atom in terms of its protons, neutrons, and electrons.

Answers

1. proton: +1, large; neutron: 0, large; electron: −1, small
2. Protons and neutrons are located in a central nucleus, while electrons orbit about the nucleus.

Key Takeaways

- Atoms are composed of three main subatomic particles: protons, neutrons, and electrons.
- Protons and neutrons are grouped together in the nucleus of an atom, while electrons orbit about the nucleus.

2.3: The Structure of Atoms is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.3: The Structure of Atoms** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.4: Nuclei of Atoms

Learning Objectives

- Define and differentiate between the atomic number and the mass number of an element.
- Explain how isotopes differ from one another.

Now that we know how atoms are generally constructed, what do atoms of any particular element look like? How many protons, neutrons, and electrons are in a specific kind of atom? First, if an atom is electrically neutral overall, then the number of protons equals the number of electrons. Because these particles have the same but opposite charges, equal numbers cancel out, producing a neutral atom.

Atomic Number

In the 1910s, experiments with x-rays led to this useful conclusion: the magnitude of the positive charge in the nucleus of every atom of a particular element is the same. In other words, all atoms of the same element have the same number of protons. Furthermore, different elements have a different number of protons in their nuclei, so the number of protons in the nucleus of an atom is characteristic of a particular element. This discovery was so important to our understanding of atoms that the number of protons in the nucleus of an atom is called the atomic number (Z).

For example, hydrogen has the atomic number 1; all hydrogen atoms have 1 proton in their nuclei. Helium has the atomic number 2; all helium atoms have 2 protons in their nuclei. There is no such thing as a hydrogen atom with 2 protons in its nucleus; a nucleus with 2 protons would be a helium atom. The atomic number *defines* an element. Table **2.4.1** lists some common elements and their atomic numbers. Based on its **atomic number**, you can determine the **number of protons in the nucleus** of an atom. The largest atoms have over 100 protons in their nuclei.

Example 2.4.1

What is the number of protons in the nucleus of each element?

a. aluminum
b. iron
c. carbon

Answer a

According to Table 2.4.1, aluminum has an atomic number of 13. Therefore, every aluminum atom has 13 protons in its nucleus.

Answer b

Iron has an atomic number of 26. Therefore, every iron atom has 26 protons in its nucleus.

Answer c

Carbon has an atomic number of 6. Therefore, every carbon atom has 6 protons in its nucleus.

Exercise 2.4.1

What is the number of protons in the nucleus of each element? Use Table 2.4.1.

a. sodium
b. oxygen
c. chlorine

Answer a

Sodium has 11 protons in its nucleus.

Answer b

Oxygen has 8 protons in its nucleus.

Answer c

Chlorine has 17 protons in its nucleus

How many electrons are in an atom? Previously we said that for an electrically neutral atom, the **number of electrons equals the number of protons**, so the total opposite charges cancel. Thus, the **atomic number** of an element also gives the **number of electrons** in an atom of that element. (Later we will find that some elements may gain or lose electrons from their atoms, so those atoms will no longer be electrically neutral. Thus we will need a way to differentiate the number of electrons for those elements.)

Example 2.4.2

How many electrons are present in the atoms of each element?

a. sulfur
b. tungsten
c. argon

Answer a

The atomic number of sulfur is 16. Therefore, in a neutral atom of sulfur, there are 16 electrons.

Answer b

The atomic number of tungsten is 74. Therefore, in a neutral atom of tungsten, there are 74 electrons.

Answer c

The atomic number of argon is 18. Therefore, in a neutral atom of argon, there are 18 electrons.

Exercise 2.4.2

How many electrons are present in the atoms of each element?

a. magnesium
b. potassium
c. iodine

Answer a

Mg has 12 electrons.

Answer b

K has 19 electrons.

Answer c

I has 53 electrons.

Table **2.4.1**: Some Common Elements and Their Atomic Numbers

Element	Atomic Number	Element	Atomic Nmbers
aluminum (Al)	13	magnesium (Mg)	12
americium (Am)	95	manganese (Mn)	25
argon (Ar)	18	mercury (Hg)	80
barium (Ba)	56	neon (Ne)	10
beryllium (Be)	4	nickel (Ni)	28
bromine (Br)	35	nitrogen (N)	7
calcium (Ca)	20	oxygen (O)	8
carbon (C)	6	phosphorus (P)	15
chlorine (Cl)	17	platinum (Pt)	78
chromium (Cr)	24	potassium (K)	19
cesium (Cs)	55	radon (Rn)	86
fluorine (F)	9	silver (Ag)	47
gallium (Ga)	31	sodium (Na)	11
gold (Au)	79	strontium (Sr)	38
helium (He)	2	sulfur (S)	16
hydrogen (H)	1	titanium (Ti)	22
iron (Fe)	26	tungsten (W)	74
iodine (I)	53	uranium (U)	92
lead (Pb)	82	zinc (Zn)	30
lithium (Li)	3	zirconium (Zr)	40

Isotopes

How many neutrons are in atoms of a particular element? At first it was thought that the number of neutrons in a nucleus was also characteristic of an element. However, it was found that atoms of the same element can have *different* numbers of neutrons. Atoms of the same element (i.e., same atomic number, Z) that have different numbers of neutrons are called isotopes. For example, 99% of the carbon atoms on Earth have 6 neutrons and 6 protons in their nuclei; about 1% of the carbon atoms have 7 neutrons in their nuclei. Naturally occurring carbon on Earth, therefore, is actually a mixture of isotopes, albeit a mixture that is 99% carbon with 6 neutrons in each nucleus.




Figure **2.4.1**: Isotopes of Hydrogen. Most hydrogen atoms have only a proton in the nucleus (a). A small amount of hydrogen exists as the isotope deuterium, which has one proton and one neutron in its nucleus (b). A tiny amount of the hydrogen isotope tritium, with one proton and two neutrons in its nucleus, also exists on Earth (c). The nuclei and electrons are proportionately much smaller than depicted here.

An important series of isotopes is found with hydrogen atoms. Most hydrogen atoms have a nucleus with only a single proton. About 1 in 10,000 hydrogen nuclei, however, also has a neutron; this particular isotope is called *deuterium*. An extremely rare hydrogen isotope, *tritium*, has 1 proton and 2 neutrons in its nucleus. Figure **2.4.1** compares the three isotopes of hydrogen.

> The discovery of isotopes required a minor change in Dalton's atomic theory. Dalton thought that all atoms of the same element were exactly the same.

Most elements exist as mixtures of isotopes. In fact, there are currently over 3,500 isotopes known for all the elements. When scientists discuss individual isotopes, they need an efficient way to specify the number of neutrons in any particular nucleus. The *mass number* (A) of an atom is the sum of the numbers of protons and neutrons in the nucleus. Given the mass number for a nucleus (and knowing the atomic number of that particular atom), you can determine the number of neutrons by subtracting the atomic number from the mass number.

A simple way of indicating the mass number of a particular isotope is to list it as a superscript on the left side of an element's symbol. Atomic numbers are often listed as a subscript on the left side of an element's symbol. Thus, we might see

$$\begin{matrix}\text{mass number}\longrightarrow 56\\ \text{atomic number}\longrightarrow 26\end{matrix}\mathrm{Fe} \tag{2.4.1}$$

which indicates a particular isotope of iron. The 26 is the atomic number (which is the same for all iron atoms), while the 56 is the mass number of the isotope. To determine the number of neutrons in this isotope, we subtract 26 from 56: 56 − 26 = 30, so there are 30 neutrons in this atom.

✔ Example 2.4.3

How many protons and neutrons are in each atom?

a. $^{35}_{17}\mathrm{Cl}$
b. $^{127}_{53}\mathrm{I}$

Answer a

In $^{35}_{17}\mathrm{Cl}$ there are 17 protons, and 35 − 17 = 18 neutrons in each nucleus.

Answer b

In $^{127}_{53}\mathrm{I}$ there are 53 protons, and 127 − 53 = 74 neutrons in each nucleus.

? Exercise 2.4.3

How many protons and neutrons are in each atom?

a. $^{197}_{79}\mathrm{Au}$
b. $^{23}_{11}\mathrm{Na}$

Answer a

In $^{197}_{79}\mathrm{Au}$ there are 79 protons, and 197 − 79 = 118 neutrons in each nucleus.

Answer b

In $^{23}_{11}\mathrm{Na}$ there are 11 protons, and 23 − 11 = 12 neutrons in each nucleus.

It is not absolutely necessary to indicate the atomic number as a subscript because each element has its own unique atomic number. Many isotopes are indicated with a superscript only, such as $^{13}\mathrm{C}$ or $^{235}\mathrm{U}$. You may also see isotopes represented in print as, for example, carbon-13 or uranium-235.

Summary

The atom consists of discrete particles that govern its chemical and physical behavior. Each atom of an element contains the same number of protons, which is the **atomic number (Z)**. Neutral atoms have the same number of electrons and protons. Atoms of an element that contain different numbers of neutrons are called **isotopes**. Each isotope of a given element has the same atomic number but a different **mass number (A)**, which is the **sum of the numbers of protons and neutrons**.

Almost all of the mass of an atom is from the total protons and neutrons contained within a tiny (and therefore very dense) nucleus. The majority of the volume of an atom is the surrounding space in which the electrons reside. A representation of a carbon-12 atom is shown below in Figure **2.4.2**.




Figure **2.4.2**: Formalism used for identifying specific nuclide (any particular kind of nucleus)

The atomic symbol, X, is the abbreviation used to represent an atom in chemical formulas. The mass number, A, is the number of protons and neutrons in the atom, which is to the left of the X. The atomic number, Z, is the number of protons in the atom, which is to the left of the X and below the A.

Concept Review Exercises

1. Why is the atomic number so important to the identity of an atom?
2. What is the relationship between the number of protons and the number of electrons in an atom?
3. How do isotopes of an element differ from each other?
4. What is the mass number of an element?

Answers

1. The atomic number defines the identity of an element. It describes the number of protons in the nucleus.
2. In an electrically neutral atom, the number of protons equals the number of electrons.
3. Isotopes of an element have the same number of protons but have different numbers of neutrons in their nuclei.
4. The mass number is the sum of the numbers of protons and neutrons in the nucleus of an atom.

Key Takeaways

- Each element is identified by its atomic number. The atomic number provides the element's location on the periodic table
- The isotopes of an element have different masses and are identified by their mass numbers.

2.4: Nuclei of Atoms is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.4: Nuclei of Atoms** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.5: Arrangement of Electron (Shell Model)

An electron shell is the outside part of an atom around the atomic nucleus. It is a group of atomic orbitals with the same value of the principal quantum number n. Electron shells have one or more electron subshells, or sublevels. The name for electron shells comes from the Bohr model, in which groups of electrons were believed to go around the nucleus at certain distances, so that their orbits formed "shells".

Introduction

What more could we want to know about the structure of an atom? We know that atoms contain positively and negatively charged particles, and that the number of these charges in each atom is different for each element. We also know that the positive charges are concentrated in a tiny nucleus, and that the electrons move around the nucleus in a space that is much, much larger than the nucleus.

However, some of the most important questions we asked in the previous Concept Development Study are still unanswered. Remember that we saw that carbon and nitrogen have very similar atomic masses. Now we can add that these elements have very similar atomic numbers, so their atoms have similar numbers of protons and electrons. But carbon and nitrogen are, in most chemical and physical ways, very different. Similarly, some elements like sodium and potassium have very different atomic numbers but have quite similar chemical and physical properties. It seems that comparing the properties of two different atoms is not very easy to understand just from comparing the numbers of protons and electrons the atoms contain.

To continue to understand the answers to these questions, we need even more detail about the structure of each type of atom. Knowledge of the electron configuration of different atoms is useful in understanding the structure of the periodic table of elements. The concept is also useful for describing the chemical bonds that hold atoms together.

Shells

An electron shell may be thought of as an orbit followed by electrons around an atom nucleus. Because each shell can contain only a fixed number of electrons, each shell is associated with a particular range of electron energy, and thus each shell must fill completely before electrons can be added to an outer shell. The electrons in the outermost shell determine the chemical properties of the atom (see Valence shell). For an explanation of why electrons exist in these shells see electron configuration.




***Figure* 2.5.1**: *A shell Diagram of lithium (left) and Sodium (right)*

The electron shells are labeled K, L, M, N, O, P, and Q; or 1, 2, 3, 4, 5, 6, and 7; going from innermost shell outwards. Electrons in outer shells have higher average energy and travel farther from the nucleus than those in inner shells. This makes them more important in determining how the atom reacts chemically and behaves as a conductor, because the pull of the atom's nucleus upon them is weaker and more easily broken. In this way, a given element's reactivity is highly dependent upon its electronic configuration.

Valence Electrons

The valence shell is the outermost shell of an atom in its uncombined state, which contains the electrons most likely to account for the nature of any reactions involving the atom and of the bonding interactions it has with other atoms. Valence electrons are electrons that are associated with an atom, and that can participate in the formation of a chemical bond; in a single covalent bond, both atoms in the bond contribute one valence electron in order to form a shared pair. The presence of valence electrons can determine the element's chemical properties and whether it may bond with other elements: For a main group element, a valence electron can only be in the outermost electron shell.

An atom with a closed shell of valence electrons tends to be chemically inert. An atom with one or two valence electrons more than a closed shell is highly reactive, because the extra valence electrons are easily removed to form a positive ion. An atom with one or two valence electrons fewer than a closed shell is also highly reactive, because of a tendency either to gain the missing valence electrons (thereby forming a negative ion), or to share valence electrons (thereby forming a covalent bond).

Like an electron in an inner shell, a valence electron has the ability to absorb or release energy in the form of a photon. An energy gain can trigger an electron to move (jump) to an outer shell; this is known as atomic excitation. Or the electron can even break free from its associated atom's valence shell; this is ionization to form a positive ion. When an electron loses energy (thereby causing a photon to be emitted), then it can move to an inner shell which is not fully occupied.

The number of valence electrons of an element can be determined by the periodic table group (vertical column) in which the element is categorized. With the exception of groups 3–12 (the transition metals), the units digit of the group number identifies how many valence electrons are associated with a neutral atom of an element listed under that particular column.

***Table* 2.5.1**

Periodic table group	Valence electrons
Group 1 (I) (alkali metals)	1
Group 2 (II) (alkaline earth metals)	2
Groups 3-12 (transition metals)	2*
Group 13 (III) (boron group)	3
Group 14 (IV) (carbon group)	4
Group 15 (V) (pnictogens)	5
Group 16 (VI) (chalcogens)	6
Group 17 (VII) (halogens)	7
Group 18 (VIII or 0) (noble gases)	8**

** The general method for counting valence electrons is generally not useful for transition metals.*

*** Except for helium, which has only two valence electrons.*

Contributors

- Connections (John Hutchinson)
- Wikipedia

2.5: Arrangement of Electron (Shell Model) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

2.6: The Periodic Table

Learning Objectives

- Explain how elements are organized into the periodic table.
- Describe how some characteristics of elements relate to their positions on the periodic table.

In the 19th century, many previously unknown elements were discovered, and scientists noted that certain sets of elements had similar chemical properties. For example, chlorine, bromine, and iodine react with other elements (such as sodium) to make similar compounds. Likewise, lithium, sodium, and potassium react with other elements (such as oxygen) to make similar compounds. Why is this so?

In 1864, Julius Lothar Meyer, a German chemist, organized the elements by atomic mass and grouped them according to their chemical properties. Later that decade, Dmitri Mendeleev, a Russian chemist, organized all the known elements according to similar properties. He left gaps in his table for what he thought were undiscovered elements, and he made some bold predictions regarding the properties of those undiscovered elements. When elements were later discovered whose properties closely matched Mendeleev's predictions, his version of the table gained favor in the scientific community. Because certain properties of the elements repeat on a regular basis throughout the table (that is, they are periodic), it became known as the periodic table.

> *Mendeleev had to list some elements out of the order of their atomic masses to group them with other elements that had similar properties.*

The periodic table is one of the cornerstones of chemistry because it organizes all the known elements on the basis of their chemical properties. A modern version is shown in Figure **2.6.1**. Most periodic tables provide additional data (such as atomic mass) in a box that contains each element's symbol. The elements are listed in order of atomic number.




Figure **2.6.1**: Modern Periodic Table. (Public Domain; PubChem modified Leticia Colmenares).

Features of the Periodic Table

Elements that have similar chemical properties are grouped in columns called **groups** (or families). As well as being numbered, some of these groups have names—for example, *alkali metals* (the first column of elements), *alkaline earth metals* (the second column of elements), *halogens* (the next-to-last column of elements), and *noble gases* (the last column of elements).

Each row of elements on the periodic table is called a **period**. Periods have different lengths; the first period has only 2 elements (hydrogen and helium), while the second and third periods have 8 elements each. The fourth and fifth periods have 18 elements each, and later periods are so long that a segment from each is removed and placed beneath the main body of the table.

Metals, Nonmetals and Metalloids

Certain elemental properties become apparent in a survey of the periodic table as a whole. Every element can be classified as either a metal, a nonmetal, or a semimetal, as shown in Figure **2.6.2**. A metal is a substance that is shiny, typically (but not always) silvery in color, and an excellent conductor of electricity and heat. Metals are also malleable (they can be beaten into thin sheets) and ductile (they can be drawn into thin wires). A nonmetal is typically dull and a poor conductor of electricity and heat. Solid nonmetals are also very brittle. As shown in Figure **2.6.2**, **metals** occupy the left three-fourths of the periodic table, while **nonmetals** (except for hydrogen) are clustered in the upper right-hand corner of the periodic table. The elements with properties intermediate between those of metals and nonmetals are called **semimetals** (or **metalloids**). Elements adjacent to the **bold zigzag** line in the right-hand portion of the periodic table have semimetal properties.




Figure **2.6.2**: Types of Elements. Elements are either metals, nonmetals, or semimetals. Each group is located in a different part of the periodic table.

Example 2.6.1

Based on its position in the periodic table, do you expect selenium (Se) to be a metal, a nonmetal, or a semimetal?

Solution

The atomic number of selenium is 34, which places it in period 4 and group 16. In Figure 2.7.1, selenium lies above and to the right of the diagonal line marking the boundary between metals and nonmetals, so it should be a nonmetal. Note, however, that because selenium is close to the metal-nonmetal dividing line, it would not be surprising if selenium were similar to a semimetal in some of its properties.

2.6.1 Exercise

Based on its location in the periodic table, do you expect indium (In) to be a nonmetal, a metal, or a semimetal?

Answer

metal

Representative, Transition and Inner-transition

Another way to categorize the elements of the periodic table is shown in Figure **2.6.3**. The first two columns on the left and the last six columns on the right are called the **main group** or **representative** elements. The ten-column block between these columns contains the **transition metals**. The two rows beneath the main body of the periodic table contain the inner transition metals. The elements in these two rows are also referred to as, respectively, the **lanthanide metals** and the **actinide metals**.




Figure **2.6.3**: Special Names for Sections of the Periodic Table. Some sections of the periodic table have special names. The elements lithium, sodium, potassium, rubidium, cesium, and francium are collectively known as alkali metals.

Alkali metals are the first column. Alkaline earth metals are the second. Halogens are the second to last column. Noble gases are the last column.

To Your Health: Transition Metals in the Body

Most of the elemental composition of the human body consists of main group elements. The first element appearing on the list that is not a main group element is iron, at 0.006 percentage by mass. Because iron has relatively massive atoms, it would appear even lower on a list organized in terms of percent by *atoms* rather than percent by mass.

Iron is a transition metal. Transition metals have interesting chemical properties, partially because some of their electrons are in *d* subshells. The chemistry of iron makes it a key component in the proper functioning of red blood cells.

Red blood cells are cells that transport oxygen from the lungs to cells of the body and then transport carbon dioxide from the cells to the lungs. Without red blood cells, animal respiration as we know it would not exist. The critical part of the red blood cell is a protein called *hemoglobin*. Hemoglobin combines with oxygen and carbon dioxide, transporting these gases from one location to another in the body. Hemoglobin is a relatively large molecule, with a mass of about 65,000 u.

The crucial atom in the hemoglobin protein is iron. Each hemoglobin molecule has four iron atoms, which act as binding sites for oxygen. It is the presence of this particular transition metal in your red blood cells that allows you to use the oxygen you inhale.

Other transition metals have important functions in the body, despite being present in low amounts. Zinc is needed for the body's immune system to function properly, as well as for protein synthesis and tissue and cell growth. Copper is also needed for several proteins to function properly in the body. Manganese is needed for the body to metabolize oxygen properly. Cobalt is a necessary component of vitamin B-12, a vital nutrient. These last three metals are not listed explicitly in Table 2.1.2, so they are present in the body in very small quantities. However, even these small quantities are required for the body to function properly.

As previously noted, the periodic table is arranged so that elements with similar chemical behaviors are in the same group. Chemists often make general statements about the properties of the elements in a group using descriptive names with historical origins. For example, the elements of Group 1 are known as the alkali metals, Group 2 are the alkaline earth metals, Group 17 are the halogens, and Group 18 are the noble gases.

Group 1: The Alkali Metals

The alkali metals are lithium, sodium, potassium, rubidium, cesium, and francium. Hydrogen is unique in that it is generally placed in Group 1, but it is not a metal. The compounds of the alkali metals are common in nature and daily life. One example is table salt (sodium chloride); lithium compounds are used in greases, in batteries, and as drugs to treat patients who exhibit manic-depressive, or bipolar, behavior. Although lithium, rubidium, and cesium are relatively rare in nature, and francium is so unstable and highly radioactive that it exists in only trace amounts, sodium and potassium are the seventh and eighth most abundant elements in Earth's crust, respectively.

Group 2: The Alkaline Earth Metals

The alkaline earth metals are beryllium, magnesium, calcium, strontium, barium, and radium. Beryllium, strontium, and barium are rare, and radium is unstable and highly radioactive. In contrast, calcium and magnesium are the fifth and sixth most abundant elements on Earth, respectively; they are found in huge deposits of limestone and other minerals.

Group 17: The Halogens

The halogens are fluorine, chlorine, bromine, iodine, and astatine. The name halogen is derived from the Greek words for "salt forming," which reflects that all the halogens react readily with metals to form compounds, such as sodium chloride and calcium chloride (used in some areas as road salt).

Compounds that contain the fluoride ion are added to toothpaste and the water supply to prevent dental cavities. Fluorine is also found in Teflon coatings on kitchen utensils. Although chlorofluorocarbon propellants and refrigerants are believed to lead to the depletion of Earth's ozone layer and contain both fluorine and chlorine, the latter is responsible for the adverse effect on the ozone layer. Bromine and iodine are less abundant than chlorine, and astatine is so radioactive that it exists in only negligible amounts in nature.

Group 18: The Noble Gases

The noble gases are helium, neon, argon, krypton, xenon, and radon. Because the noble gases are composed of only single atoms, they are called monatomic. At room temperature and pressure, they are **unreactive gases**. Because of their lack of reactivity, for many years they were called **inert gases** or rare gases. However, the first chemical compounds containing the noble gases were prepared in 1962. Although the noble gases are relatively minor constituents of the atmosphere, natural gas contains substantial amounts of helium. Because of its low reactivity, argon is often used as an unreactive (inert) atmosphere for welding and in light bulbs. The red light emitted by neon in a gas discharge tube is used in neon lights.

To Your Health: Radon

Radon is an invisible, odorless noble gas that is slowly released from the ground, particularly from rocks and soils whose uranium content is high. Because it is a noble gas, radon is not chemically reactive. Unfortunately, it is radioactive, and increased exposure to it has been correlated with an increased lung cancer risk.

Because radon comes from the ground, we cannot avoid it entirely. Moreover, because it is denser than air, radon tends to accumulate in basements, which if improperly ventilated can be hazardous to a building's inhabitants. Fortunately, specialized ventilation minimizes the amount of radon that might collect. Special fan-and-vent systems are available that draw air from below the basement floor, before it can enter the living space, and vent it above the roof of a house.

After smoking, radon is thought to be the second-biggest *preventable* cause of lung cancer in the United States. The American Cancer Society estimates that 10% of all lung cancers are related to radon exposure. There is uncertainty regarding what levels of exposure cause cancer, as well as what the exact causal agent might be (either radon or one of its breakdown products, many of which are also radioactive and, unlike radon, not gases). The US Environmental Protection Agency recommends testing every floor below the third floor for radon levels to guard against long-term health effects.

Why do elements in a given group have similar properties?

The periodic table is organized on the basis of similarities in elemental properties, but what explains these similarities? It turns out that the shape of the periodic table reflects the filling of subshells with electrons, as shown in Figure **2.6.4**. Starting with the first

period and going from left to right, the table reproduces the order of filling of the electron subshells in atoms. Furthermore, elements in the same *group* share the same valence shell electron configuration. For example, all elements in the first column have a single *s* electron in their valence shells, so their electron configurations can be described as ns^1 (where *n* represents the shell number). This last observation is crucial. Chemistry is largely the result of interactions between the valence electrons of different atoms. Thus, atoms that have **the same valence shell electron configuration** will have **similar chemistry.**




Figure **2.6.4** The Shape of the Periodic Table. The shape of the periodic table reflects the order in which electron shells and subshells fill with electrons.

Example 2.6.1

Using the variable *n* to represent the number of the valence electron shell, write the valence shell electron configuration for each group.

a. the alkaline earth metals
b. the column of elements headed by carbon

Answer a

The alkaline earth metals are in the second column of the periodic table. This column corresponds to the *s* subshell being filled with 2 electrons. Therefore, the valence shell electron configuration is ns^2.

Answer b

The electron configuration of carbon is $1s^22s^22p^2$. Its valence shell electron configuration is $2s^22p^2$. Every element in the same column should have a similar valence shell electron configuration, which we can represent as ns^2np^2.

Exercise 2.6.1

Using the variable *n* to represent the number of the valence electron shell, write the valence shell electron configuration for each group.

a. the halogens
b. the column of elements headed by oxygen

Answer a

The halogens are in the 17th column (or Group 7A) of the periodic table. This column corresponds to the *p* subshell being filled with 5 electrons. Therefore, the valence shell electron configuration is ns^2np^5.

Answer b

The column headed by O is the 16th column (or Group 6A). This column corresponds to the *p* subshell being filled with 4 electrons. Therefore, the valence shell electron configuration is ns^2np^4.

Valence Electrons and Group Number

The number of valence electrons of an element can be determined by the periodic table group (vertical column) in which the element is categorized. With the exception of groups 3–12 (the transition metals), the units digit of the group number identifies how many valence electrons are associated with a neutral atom of an element listed under that particular column.

Table **2.6.1**. The Group number and the number of valence electrons.

Periodic table group	Valence electrons
Group 1 (I) (alkali metals)	1
Group 2 (II) (alkaline earth metals)	2
Groups 3-12 (transition metals)	2*
Group 13 (III) (boron group)	3
Group 14 (IV) (carbon group)	4
Group 15 (V) (pnictogens)	5
Group 16 (VI) (chalcogens)	6
Group 17 (VII) (halogens)	7
Group 18 (VIII or 0) (noble gases)	8**

** The general method for counting valence electrons is generally not useful for transition metals.*

*** Except for helium, which has only two valence electrons.*

Atomic Radius

The periodic table is useful for understanding atomic properties that show periodic trends. One such property is the atomic radius (Figure **2.6.5**). The atomic radius is defined as one-half the distance between the nuclei of identical atoms that are bonded together. The units for atomic radii are picometers, equal to 10^{-12} meters. As an example, the internuclear distance between the two hydrogen atoms in an $\mathbf{H_2}$ molecule is measured to be $\mathbf{74\ pm}$. Therefore, the atomic radius of a hydrogen atom is $\frac{74}{2} = \mathbf{37\ pm}$.

As mentioned earlier, the higher the shell number, the farther from the nucleus the electrons in that shell are likely to be. In other words, the size of an atom is generally determined by the number of the valence electron shell. Therefore, as we go down a column on the periodic table, the atomic radius increases. As we go *across* a period on the periodic table, however, electrons are being added to the *same* valence shell; meanwhile, more protons are being added to the nucleus, so the positive charge of the nucleus is increasing. The increasing positive charge attracts the electrons more strongly, pulling them closer to the nucleus. Consequently, as we go across a period, from left to right, the atomic radius decreases. These trends are seen clearly in Figure **2.6.5**




Figure **2.6.5** Trends on the Periodic Table. Atomic radii of the representative elements measured in picometers. The relative sizes of the atoms show several trends with regard to the structure of the periodic table. Atoms become larger going down a group and going from right to left across a period.

✔ Example 2.6.2

Using the periodic table (rather than Figure **2.6.5**), which atom is larger?

a. N or Bi
b. Mg or Cl

Answer a

Bi is below N in Group 5A in the periodic table and has electrons in higher-numbered shells, hence we expect that Bi atoms are larger than N atoms.

Answer b

Both Mg and Cl are in period 3 of the periodic table, but Cl lies farther to the right. Therefore we expect Mg atoms to be larger than Cl atoms.

? Exercise 2.6.2

Using the periodic table (rather than Figure **2.6.5**), which atom is larger?

a. Li or F
b. Na or K

Answer a

Li and F are on the same period, but F lies farther to the right. Therefore, we expect Li to be larger than F atoms.

Answer b

K lies below Na in Group 1A, hence has more electron shells, making it larger than Na.

Career Focus: Clinical Chemist

Clinical chemistry is the area of chemistry concerned with the analysis of body fluids to determine the health status of the human body. Clinical chemists measure a variety of substances, ranging from simple elements such as sodium and potassium to complex molecules such as proteins and enzymes, in blood, urine, and other body fluids. The absence or presence, or abnormally low or high amounts, of a substance can be a sign of some disease or an indication of health. Many clinical chemists use sophisticated equipment and complex chemical reactions in their work, so they not only need to understand basic chemistry, but also be familiar with special instrumentation and how to interpret test results.

Concept Review Exercises

1. How are the elements organized into the periodic table?
2. Looking at the periodic table, where do the following elements appear?
    a. the metals
    b. the nonmetals
    c. the halogens
    d. the transition metals
    e. the noble gases
3. Describe the trends in atomic radii as related to an element's position on the periodic table.

Answers

1. Elements are organized in order of increasing atomic number.
2. a. the left three-quarters of the periodic table (to the left of the zigzag band)
    b. the right quarter of the periodic table (to the right of the zigzag band)
    c. the next-to-last column of the periodic table
    d. the middle section of the periodic table
    e. the last column of the periodic table
3. As you go across the periodic table, atomic radii decrease; as you go down the periodic table, atomic radii increase.

Key Takeaways

- The chemical elements are arranged in a chart called the periodic table.
- Some characteristics of the elements are related to their position on the periodic table.
- The number of valence electrons of an element can be determined by the group (vertical column) number in the Periodic Table. Elements with the same number of valence electrons have similar chemical properties.

2.6: The Periodic Table is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.7: The Periodic Table** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.7: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms and ask yourself how they relate to the topics in the chapter.

An element is a substance that cannot be broken down into simpler chemical substances. Only about 90 naturally occurring elements are known. They have varying abundances on Earth and in the body. Each element has a one- or two-letter **chemical symbol**.

The **modern atomic theory** states that the smallest piece of an element is an **atom**. Individual atoms are extremely small, on the order of 10^{-10} m across. Most elements exist in pure form as individual atoms, but some exist as **diatomic molecules**. Atoms themselves are composed of subatomic particles. The **electron** is a tiny subatomic particle with a negative charge. The **proton** has a positive charge and, while small, is much larger than the electron. The **neutron** is also much larger than an electron but has no electrical charge.

Protons, neutrons, and electrons have a specific arrangement in an atom. The protons and neutrons are found in the center of the atom, grouped together into a **nucleus**. The electrons are found in fuzzy clouds around the nucleus.

Each element has a characteristic number of protons in its nucleus. This number of protons is the **atomic number** of the element. An element may have different numbers of neutrons in the nuclei of its atoms; such atoms are referred to as **isotopes**. Two isotopes of hydrogen are deuterium, with a proton and a neutron in its nucleus, and tritium, with a proton and two neutrons in its nucleus. The sum of the numbers of protons and neutrons in a nucleus is called the **mass number** and is used to distinguish isotopes from each other.

Masses of individual atoms are measured in **atomic mass units**. An atomic mass unit is equal to 1/12th of the mass of a single carbon-12 atom. Because different isotopes of an element have different masses, the **atomic mass** of an element is a weighted average of the masses of all the element's naturally occurring isotopes.

The modern theory of electron behavior is called **quantum mechanics**. According to this theory, electrons in atoms can only have specific, or **quantized**, energies. Electrons are grouped into general regions called **shells**, and within these into more specific regions called **subshells**. There are four types of subshells, and each type can hold up to a maximum number of electrons. Chemistry typically occurs because of interactions between the electrons of the outermost shell of different atoms, called the valence shell electrons. Electrons in inner shells are called core electrons.

Elements are grouped together by similar chemical properties into a chart called the **periodic table**. Vertical columns of elements are called **groups** or **families**. Some of the groups of elements have names, like the alkali metals, the alkaline earth metals, the halogens, and the noble gases. A horizontal row of elements is called a **period**. Periods and groups have differing numbers of elements in them. The periodic table separates elements into **metals**, **nonmetals**, and **semimetals**. The periodic table is also separated into **main group elements**, **transition metals**, **lanthanide elements**, and **actinide elements**. The lanthanide and actinide elements are also referred to as **inner transition metal elements**. The shape of the periodic table reflects the sequential filling of shells and subshells in atoms.

The periodic table helps us understand trends in some of the properties of atoms. One such property is the **atomic radius** of atoms. From top to bottom of the periodic table, atoms get bigger because electrons are occupying larger and bigger shells. From left to right across the periodic table, electrons are filling the same shell but are being attracted by an increasing positive charge from the nucleus, and thus the atoms get smaller.

2.7: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

3: Nuclear Chemistry

Most chemists pay little attention to the nucleus of an atom except to consider the number of protons it contains because that determines an element's identity. However, in nuclear chemistry, the composition of the nucleus and the changes that occur there are very important. Applications of nuclear chemistry may be more widespread than you realize. Many people are aware of nuclear power plants and nuclear bombs, but nuclear chemistry also has applications ranging from smoke detectors to medicine, from the sterilization of food to the analysis of ancient artifacts. In this chapter, we will examine some of the basic concepts of nuclear chemistry and some of the nuclear reactions that are important in our everyday lives.

3.1: Radioactivity
3.2: Half-Life
3.3: Units of Radioactivity
3.4: Uses of Radioactive Isotopes
3.5: Nuclear Energy
3.6: 3.6 Chapter Summary
3.7: Ionic Bonding and Simple Ionic Compounds

3: Nuclear Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.1: Radioactivity

Learning Objectives

- To define and give examples of the major types of radioactivity.

Atoms are composed of subatomic particles—protons, neutrons, and electrons. Protons and neutrons are located in the nucleus and provide most of the mass of the atom, while electrons circle the nucleus in shells and subshells and account for an atom's size. Remember, the notation for succinctly representing an isotope of a particular atom:

$$^{12}_{6}\mathrm{C}$$

The element in this example, represented by the symbol *C*, is carbon. Its atomic number, 6, is the lower left subscript on the symbol and is the number of protons in the atom. The mass number, the superscript to the upper left of the symbol, is the sum of the number of protons and neutrons in the nucleus of this particular isotope. In this case, the mass number is 12, which means that the number of neutrons in the atom is 12 − 6 = 6 (that is, the mass number of the atom minus the number of protons in the nucleus equals the number of neutrons). Occasionally, the atomic number is omitted in this notation because the symbol of the element itself conveys its characteristic atomic number. The two isotopes of hydrogen, $^{2}\mathrm{H}$ and $^{3}\mathrm{H}$, are given their own names: deuterium (D) and tritium (T), respectively. Another way of expressing a particular isotope is to list the mass number after the element name, like carbon-12 or hydrogen-3.

Atomic theory in the 19th century presumed that nuclei had fixed compositions. But in 1896, the French scientist Henri Becquerel found that a uranium compound placed near a photographic plate made an image on the plate, even if the compound was wrapped in black cloth. He reasoned that the uranium compound was emitting some kind of radiation that passed through the cloth to expose the photographic plate. Further investigations showed that the radiation was a combination of particles and electromagnetic rays, with its ultimate source as the atomic nucleus. These emanations were ultimately called, collectively, **radioactivity**.

There are three main forms of radioactive emissions. The first is called an alpha particle, which is symbolized by the Greek letter α. An alpha particle is composed of two protons and two neutrons, and so it is the same as a helium nucleus. (We often use $^{4}_{2}\mathrm{He}$ to represent an alpha particle.) It has a 2+ charge. When a radioactive atom emits an alpha particle, the original atom's atomic number decreases by two (because of the loss of two protons), and its mass number decreases by four (because of the loss of four nuclear particles). We can represent the emission of an alpha particle with a chemical equation—for example, the alpha-particle emission of uranium-235 is as follows:

$$^{235}_{92}\mathrm{U} \rightarrow {}^{4}_{2}\mathrm{He} + {}^{231}_{90}\mathrm{Th}$$

How do we know that a product of the reaction is $^{231}_{90}\mathrm{Th}$? We use the law of conservation of matter, which says that matter cannot be created or destroyed. This means we must have the same number of protons and neutrons on both sides of the chemical equation. If our uranium nucleus loses 2 protons, there are 90 protons remaining, identifying the element as thorium. Moreover, if we lose 4 nuclear particles of the original 235, there are 231 remaining. Thus, we use subtraction to identify the isotope of the thorium atom —in this case, $^{231}_{90}\mathrm{Th}$.

Chemists often use the names *parent isotope* and *daughter isotope* to represent the original atom and the product other than the alpha particle. In the previous example, $^{235}_{92}\mathrm{U}$ is the parent isotope, and $^{231}_{90}\mathrm{Th}$ is the daughter isotope. When one element changes into another in this manner, it undergoes *radioactive decay*.

Example 3.1.1

Write the nuclear equation that represents the radioactive decay of radon-222 by alpha particle emission and identify the daughter isotope.

Solution

Radon has an atomic number of 86, so the parent isotope is represented as $^{222}_{86}\mathrm{Rn}$. We represent the alpha particle as $^{4}_{2}\mathrm{He}$ and use subtraction (222 − 4 = 218 and 86 − 2 = 84) to identify the daughter isotope as an isotope of polonium, $^{218}_{84}\mathrm{Po}$:

$$^{222}_{86}\text{Rn} \rightarrow {}^{4}_{2}\text{He} + {}^{218}_{84}\text{Po}$$

Exercise 3.1.1

Write the nuclear equation that represents the radioactive decay of polonium-209 by alpha particle emission and identify the daughter isotope.

Answer

$$^{209}_{84}\text{Po} \rightarrow {}^{4}_{2}\text{He} + {}^{205}_{82}\text{Pb}$$

The second major type of radioactive emission is called a beta particle, symbolized by the Greek letter β. A beta particle is an electron ejected from the nucleus (not from the shells of electrons about the nucleus) and has a 1− charge. We can also represent a beta particle as $^{0}_{-1}\text{e}$ or β^{-}. The net effect of beta particle emission on a nucleus is that a neutron is converted to a proton. The overall mass number stays the same, but because the number of protons increases by one, the atomic number goes up by one. Carbon-14 decays by emitting a beta particle:

$$^{14}_{6}\text{C} \rightarrow {}^{14}_{7}\text{N} + {}^{0}_{-1}\text{e}$$

Again, the sum of the atomic numbers is the same on both sides of the equation, as is the sum of the mass numbers. (Note that the electron is assigned an "atomic number" of 1−, equal to its charge.)

The third major type of radioactive emission is not a particle but rather a very energetic form of electromagnetic radiation called gamma rays, symbolized by the Greek letter γ. Gamma rays themselves do not carry an overall electrical charge, but they may knock electrons out of atoms in a sample of matter and make it electrically charged (for which gamma rays are termed *ionizing radiation*). For example, in the radioactive decay of radon-222, both alpha and gamma radiation are emitted, with the latter having an energy of 8.2×10^{-14} J per nucleus decayed:

$$^{222}_{86}\text{Rn} \rightarrow {}^{218}_{84}\text{Po} + {}^{4}_{2}\text{He} + \gamma$$

This may not seem like much energy, but if 1 mol of radon atoms were to decay, the gamma ray energy would be 49 million kJ!

Example 3.1.2

Write the nuclear equation that represents the radioactive decay of boron-12 by beta particle emission and identify the daughter isotope. A gamma ray is emitted simultaneously with the beta particle.

Solution

The parent isotope is $^{12}_{5}\text{B}$ while one of the products is an electron, $^{0}_{-1}\text{e}$. So that the mass and atomic numbers have the same value on both sides, the mass number of the daughter isotope must be 12, and its atomic number must be 6. The element having an atomic number of 6 is carbon. Thus, the complete nuclear equation is as follows:

$$^{12}_{5}\text{B} \rightarrow {}^{12}_{6}\text{C} + {}^{0}_{-1}\text{e} + \gamma$$

The daughter isotope is $^{12}_{6}\text{C}$.

Exercise 3.1.2

Write the nuclear equation that represents the radioactive decay of iodine-131 by beta particle emission and identify the daughter isotope. A gamma ray is emitted simultaneously with the beta particle.

Answer

$$^{131}_{53}\text{I} \rightarrow {}^{131}_{54}\text{Xe} + {}^{0}_{-1}\text{e} + \gamma$$

Alpha, beta, and gamma emissions have different abilities to penetrate matter. The relatively large alpha particle is easily stopped by matter (although it may impart a significant amount of energy to the matter it contacts). Beta particles penetrate slightly into matter, perhaps a few centimeters at most. Gamma rays can penetrate deeply into matter and can impart a large amount of energy into the surrounding matter.




Figure **3.1.2**: Different emissions exhibit different penetration powers. (CC BY-NC-SA 3.0; anonymous)

Table **3.1.1** summarizes the properties of the three main types of radioactive emissions.

Table **3.1.1**: The Three Main Forms of Radioactive Emissions

Characteristic	Alpha Particles	Beta Particles	Gamma Rays
symbols	α, $^{4}_{2}\text{He}$	β, $^{0}_{-1}\text{e}$	γ
identity	helium nucleus	electron	electromagnetic radiation
charge	2+	1−	none
mass number	4	0	0
penetrating power	minimal (will not penetrate skin)	short (will penetrate skin and some tissues slightly)	deep (will penetrate tissues deeply)

Occasionally, an atomic nucleus breaks apart into smaller pieces in a radioactive process called *spontaneous fission* (or fission). Typically, the daughter isotopes produced by fission are a varied mix of products, rather than a specific isotope as with alpha and beta particle emission. Often, fission produces excess neutrons that will sometimes be captured by other nuclei, possibly inducing additional radioactive events. Uranium-235 undergoes spontaneous fission to a small extent. One typical reaction is

$$^{235}_{92}\text{U} \rightarrow {}^{139}_{56}\text{Ba} + {}^{94}_{36}\text{Kr} + 2{}^{1}_{0}\text{n}$$

where $^{1}_{0}\text{n}$ is a neutron. As with any nuclear process, the sums of the atomic numbers and the mass numbers must be the same on both sides of the equation. Spontaneous fission is found only in large nuclei. The smallest nucleus that exhibits spontaneous fission is lead-208.

> *Fission is the radioactive process used in nuclear power plants and one type of nuclear bomb.*

Key Takeaway

The major types of radioactivity include alpha particles, beta particles, and gamma rays.

3.1: Radioactivity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.1: Radioactivity** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.2: Half-Life

Learning Objectives

- To define *half-life*.
- To determine the amount of radioactive substance remaining after a given number of half-lives.

Whether or not a given isotope is radioactive is a characteristic of that particular isotope. Some isotopes are stable indefinitely, while others are radioactive and decay through a characteristic form of emission. As time passes, less and less of the radioactive isotope will be present, and the level of radioactivity decreases. An interesting and useful aspect of radioactive decay is the **half-life**. The half-life of a radioactive isotope is the amount of time it takes for one-half of the radioactive isotope to decay. The half-life of a specific radioactive isotope is constant; it is unaffected by conditions and is independent of the initial amount of that isotope.

Consider the following example. Suppose we have 100.0 g of ^{3}H (tritium, a radioactive isotope of hydrogen). It has a half-life of 12.3 y. After 12.3 y, half of the sample will have decayed to ^{3}He by emitting a beta particle, so that only 50.0 g of the original ^{3}H remains. After another 12.3 y—making a total of 24.6 y—another half of the remaining ^{3}H will have decayed, leaving 25.0 g of ^{3}H. After another 12.3 y—now a total of 36.9 y—another half of the remaining ^{3}H will have decayed, leaving 12.5 g of ^{3}H. This sequence of events is illustrated in Figure **3.2.1**.




Figure **3.2.1**: Radioactive Decay. During each successive half-life, half of the initial amount will radioactively decay.

We can determine the amount of a radioactive isotope remaining after a given number half-lives by using the following expression:

$$\text{amount remaining} = \text{initial amount} \times \left(\frac{1}{2}\right)^{n} \tag{3.2.1}$$

where n is the number of half-lives. This expression works even if the number of half-lives is not a whole number.

Example 3.2.1

The half-life of ^{20}F is 11.0 s. If a sample initially contains 5.00 g of ^{20}F, how much ^{20}F remains after 44.0 s?

Solution

If we compare the time that has passed to the isotope's half-life, we note that 44.0 s is exactly 4 half-lives, so we can use Equation 3.2.1 with $n = 4$. Substituting and solving results in the following:

$$\begin{aligned}\text{amount remaining} &= 5.00\ g \times \left(\frac{1}{2}\right)^{4} \\ &= 5.00\ g \times \frac{1}{16} \\ &= 0.313\ g\end{aligned}$$

Less than one-third of a gram of ^{20}F remains.

? Exercise 3.2.2

The half-life of ^{44}Ti is 60.0 y. A sample initially contains 0.600 g of ^{44}Ti. How much ^{44}Ti remains after 180.0 y?

Answer

0.075 g.

Half-lives of isotopes range from fractions of a microsecond to billions of years. Table **3.2.1** lists the half-lives of some isotopes.

Table **3.2.1**: Half-Lives of Various Isotopes

Isotope	Half-Life
^{3}H	12.3 y
^{14}C	5,730 y
^{40}K	1.26×10^{9} y
^{51}Cr	27.70 d
^{90}Sr	29.1 y
^{131}I	8.04 d
^{222}Rn	3.823 d
^{235}U	7.04×10^{8} y
^{238}U	4.47×10^{9} y
^{241}Am	432.7 y
^{248}Bk	23.7 h
^{260}Sg	4 ms

✓ Example 3.2.2

The isotope **I**-125 is used in certain laboratory procedures and has a half-life of 59.4 days. If the initial activity of a sample of **I**-125 is **32,000 counts per minute (cpm)**, how much activity will be present in 178.2 days?

Solution

We begin by determining how many half-lives are represented by 178.2 days:

$$\frac{\textbf{178.2 days}}{\textbf{59.4 days/half-life}} = \textbf{3 half-lives}$$

Then we simply count activity:

$$\begin{aligned}\textbf{initial activity } (t_0) &= \textbf{32,000 cpm}\\ \textbf{after one half-life} &= \textbf{16,000 cpm}\\ \textbf{after two half-lives} &= \textbf{8,000 cpm}\\ \textbf{after three half-lives} &= \textbf{4,000 cpm}\end{aligned}$$

Be sure to keep in mind that the initial count is at time zero (t_0) and we subtract from that count at the first half-life. The second half-life has an activity of half the previous count (not the initial count).


Equation 3.2.1 can be used to calculate the amount of radioactivity remaining after a given time:

$$N_t = N_0 \times (0.5)^{\text{number of half-lives}}$$

where N_t = activity at time t and N_0 = initial activity at time $t = 0$.

If we have an initial activity of $32,000$ cpm, what will the activity be after four half-lives?

$$\begin{aligned} N_t &= N_0\,(0.5)^4 \\ &= (32,000)\,(0.5)\,(0.5)\,(0.5)\,(0.5) \\ &= 2000 \text{ cpm} \end{aligned}$$



Typical radioactive decay curve. (CC BY-NC-SA 3.0; LibreTexts)

The graph above illustrates a typical decay curve for $\mathrm{I-125}$. The activity decreases by one-half during each succeeding half-life.

3.2.2 Exercise

A sample of Ac-225 originally contained 80 grams and after 50 days only 2.5 grams of the original Ac-225 remain. What is the half life of Ac-225?

Answer

5 days

Looking Closer: Half-Lives of Radioactive Elements

Many people think that the half-life of a radioactive element represents the amount of time an element is radioactive. In fact, it is the time required for half—not all—of the element to decay radioactively. Occasionally, however, the daughter element is also radioactive, so its radioactivity must also be considered.

The expected working life of an ionization-type smoke detector (described in the opening essay) is about 10 years. In that time, americium-241, which has a half-life of about 432 y, loses less than 4% of its radioactivity. A half-life of 432 y may seem long to us, but it is not very long as half-lives go. Uranium-238, the most common isotope of uranium, has a half-life of about 4.5×10^9 y, while thorium-232 has a half-life of 14×10^9 y.

On the other hand, some nuclei have extremely short half-lives, presenting challenges to the scientists who study them. The longest-lived isotope of lawrencium, ^{262}Lr, has a half-life of 3.6 h, while the shortest-lived isotope of lawrencium, ^{252}Lr, has a half-life of 0.36 s. As of this writing, the largest atom ever detected has atomic number 118, mass number 293, and a half-life of 120 ns. Can you imagine how quickly an experiment must be done to determine the properties of elements that exist for so short a time?

Key Takeaways

- Natural radioactive processes are characterized by a half-life, the time it takes for half of the material to decay radioactively.
- The amount of material left over after a certain number of half-lives can be easily calculated.

3.2: Half-Life is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.2: Half-Life** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.3: Units of Radioactivity

Learning Objectives

- To express amounts of radioactivity in a variety of units.

Previously, we used mass to indicate the amount of radioactive substance present. This is only one of several units used to express amounts of radiation. Some units describe the number of radioactive events occurring per unit time, while others express the amount of a person's exposure to radiation.

Perhaps the direct way of reporting radioactivity is the number of radioactive decays per second. One decay per second is called one becquerel (Bq). Even in a small mass of radioactive material, however, there are many thousands of decays or disintegrations per second. The unit curie (Ci), now defined as 3.7×10^{10} decays per second, was originally defined as the number of decays per second in 1 g of radium. Many radioactive samples have activities that are on the order of microcuries (μCi) or more. Both the becquerel and curie can be used in place of grams to describe quantities of radioactive material. As an example, the amount of americium in an average smoke detector has an activity of 0.9 μCi.

> *The unit becquerel is named after Henri Becquerel, who discovered radioactivity in 1896. The unit curie is named after Polish scientist Marie Curie, who performed some of the initial investigations into radioactive phenomena and discovered the elements, polonium (Po) and radium (Ra) in the early 1900s.*

Example 3.3.1

A sample of radium has an activity of 16.0 mCi (millicuries). If the half-life of radium is 1,600 y, how long before the sample's activity is 1.0 mCi?

Solution

The following table shows the activity of the radium sample over multiple half-lives:

Solutions to Example 11.3.1

Time in Years	Activity
0	16.0 mCi
1,600	8.0 mCi
3,200	4.0 mCi
4,800	2.0 mCi
6,400	1.0 mCi

Over a period of 4 half-lives, the activity of the radium will be halved four times, at which point its activity will be 1.0 mCi. Thus, it takes 4 half-lives, or 4 × 1,600 y = 6,400 y, for the activity to decrease to 1.0 mCi.

Exercise 3.3.1

A sample of radon has an activity of 60,000 Bq. If the half-life of radon is 15 h, how long before the sample's activity is 3,750 Bq?

Answer

60 hrs.

Other measures of radioactivity are based on the effects it has on living tissue. Radioactivity can transfer energy to tissues in two ways: through the kinetic energy of the particles hitting the tissue and through the electromagnetic energy of the gamma rays being absorbed by the tissue. Either way, the transferred energy—like thermal energy from boiling water—can damage the tissue.

The rad (an acronym for radiation absorbed dose) is a unit equivalent to a gram of tissue absorbing 0.01 J:

$$1 \text{ rad} = 0.01 \text{ J/g}$$

Another unit of radiation absorption is the gray (Gy):

$$1 \text{ Gy} = 100 \text{ rad}$$

The rad is more common. To get an idea of the amount of energy this represents, consider that the absorption of 1 rad by 70,000 g of H_2O (approximately the same mass as a 150 lb person) would increase its temperature by only 0.002°C. This may not seem like a lot, but it is enough energy to break about 1×10^{21} molecular C–C bonds in a person's body. That amount of damage would not be desirable.

Predicting the effects of radiation is complicated by the fact that various tissues are affected differently by different types of emissions. To quantify these effects, the unit **rem** (an acronym for roentgen equivalent, man) is defined as

$$\text{rem} = \text{rad} \times \text{RBE}$$

where RBE is the *relative biological effectiveness factor* is a number greater than or equal to 1 that takes into account the type of radioactive emission and sometimes the type of tissue being exposed. For beta particles, RBE factor equals 1. For alpha particles striking most tissues, the factor is 10, but for eye tissue, the factor is 30. Most radioactive emissions that people are exposed to are on the order of a few dozen millirems (mrem) or less; a medical X ray is about 20 mrem. A sievert (Sv) is a related unit and is defined as 100 rem.

What is a person's annual exposure to radioactivity and radiation? Table **3.3.1** lists the sources and annual amounts of radiation exposure. It may surprise you to learn that fully 82% of the radioactivity and radiation exposure we receive is from natural sources—sources we cannot avoid. Fully 10% of the exposure comes from our own bodies—largely from ^{14}C and ^{40}K.

Table **3.3.1**: Average Annual Radiation Exposure (Approximate)

Source	Amount (mrem)
radon gas	200
medical sources	53
radioactive atoms in the body naturally	39
terrestrial sources	28
cosmic sources	28
consumer products	10
nuclear energy	0.05
Total	358

> *Flying from New York City to San Francisco adds 5 mrem to your overall radiation exposure because the plane flies above much of the atmosphere, which protects us from most cosmic radiation.*

The actual effects of radioactivity and radiation exposure on a person's health depend on the type of radioactivity, the length of exposure, and the tissues exposed. Table **3.3.2** lists the potential threats to health at various amounts of exposure over short periods of time (hours or days).

Table **3.3.2**: Effects of Short-Term Exposure to Radioactivity and Radiation

Exposure (rem)	Effect
1 (over a full year)	no detectable effect
~20	increased risk of some cancers
~100	damage to bone marrow and other tissues; possible internal bleeding; decrease in white blood cell count
200–300	visible “burns” on skin, nausea, vomiting, and fatigue
>300	loss of white blood cells; hair loss
~600	death

One of the simplest ways of detecting radioactivity is by using a piece of photographic film embedded in a badge or a pen. On a regular basis, the film is developed and checked for exposure. A comparison of the exposure level of the film with a set of standard exposures indicates the amount of radiation a person was exposed to.



Figure **3.3.1**: Detecting Radioactivity. A Geiger counter is a common instrument used to detect radioactivity.

Another means of detecting radioactivity is an electrical device called a Geiger counter (Figure **3.3.1**). It contains a gas-filled chamber with a thin membrane on one end that allows radiation emitted from radioactive nuclei to enter the chamber and knock electrons off atoms of gas (usually argon). The presence of electrons and positively charged ions causes a small current, which is detected by the Geiger counter and converted to a signal on a meter or, commonly, an audio circuit to produce an audible “click.”

Key Takeaway

- Radioactivity can be expressed in a variety of units, including rems, rads, and curies.

3.3: Units of Radioactivity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.3: Units of Radioactivity** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.4: Uses of Radioactive Isotopes

Learning Objectives

- To learn some applications of radioactivity.

Radioactive isotopes have a variety of applications. Generally, however, they are useful either because we can detect their radioactivity or we can use the energy they release.

Radioactive isotopes are effective tracers because their radioactivity is easy to detect. A tracer is a substance that can be used to follow the pathway of that substance through some structure. For instance, leaks in underground water pipes can be discovered by running some tritium-containing water through the pipes and then using a Geiger counter to locate any radioactive tritium subsequently present in the ground around the pipes. (Recall that tritium, ^{3}H, is a radioactive isotope of hydrogen.)

Tracers can also be used to follow the steps of a complex chemical reaction. After incorporating radioactive atoms into reactant molecules, scientists can track where the atoms go by following their radioactivity. One excellent example of this is the use of radioactive carbon-14 to determine the steps involved in the photosynthesis in plants. We know these steps because researchers followed the progress of the radioactive carbon-14 throughout the process.

Radioactive isotopes are useful for establishing the ages of various objects. The half-life of radioactive isotopes is unaffected by any environmental factors, so the isotope acts like an internal clock. For example, if a rock is analyzed and is found to contain a certain amount of uranium-235 and a certain amount of its daughter isotope, we can conclude that a certain fraction of the original uranium-235 has radioactively decayed. If half of the uranium has decayed, then the rock has an age of one half-life of uranium-235, or about 4.5×10^{9} y. Many analyses like this, using a wide variety of isotopes, have indicated that the age of Earth itself is over 4×10^{9} y. In another interesting example of radioactive dating, ^{3}H dating has been used to verify the stated vintages of some old fine wines.

Carbon-14 (half-life is 5,370 y) is particularly useful in determining the age of once-living artifacts (e.g., animal or plant matter). A tiny amount of carbon-14 is produced naturally in the upper reaches of the atmosphere, and living things incorporate some of it into their tissues, building up to a constant, although very low, level. Once a living thing dies, however, it no longer acquires carbon-14, and as time passes, the carbon-14 that was in the tissues decays. If a once-living artifact is discovered and analyzed many years after its death, with the remaining carbon-14 compared to the known constant level, an approximate age of the artifact can be determined. Using such methods, scientists determined that the age of the Shroud of Turin (made of linen, which comes from the flax plant, and purported by some to be the burial cloth of Jesus Christ; Figure **3.4.1**) is about 600–700 y, not 2,000 y as claimed by some. Scientists were also able to use radiocarbon dating to show that the age of a mummified body found in the ice of the Alps was 5,300 y.




Figure **3.4.1**: Shroud of Turin. In 1989, several groups of scientists used carbon-14 dating to demonstrate that the age of the Shroud of Turin was only 600–700 y. Many people still cling to a different notion, despite the scientific evidence. (Public Domain)

The radiation emitted by some radioactive substances can be used to kill microorganisms on a variety of foodstuffs, which extends the shelf life of these products. Produce such as tomatoes, mushrooms, sprouts, and berries are irradiated with the emissions from coba . This exposure kills a lot of the bacteria that cause spoilage, so the produce stays fresh longer. Eggs and some meat, such as beef, pork, and poultry, can also be irradiated. Contrary to the belief of some people, irradiation of food *does not* make the food itself radioactive.

Radioactive isotopes have numerous medical applications—diagnosing and treating illnesses and diseases. One example of a diagnostic application is using radioactive iodine-131 to test for thyroid activity (Figure **3.4.2**). The thyroid gland in the neck is one of the few places in the body with a significant concentration of iodine. To evaluate thyroid activity, a measured dose of iodine-131 is administered to a patient, and the next day a scanner is used to measure the amount of radioactivity in the thyroid gland. The amount of radioactive iodine that collects there is directly related to the activity of the thyroid, allowing trained physicians to diagnose both hyperthyroidism and hypothyroidism. Iodine-131 has a half-life of only 8 d, so the potential for damage due to exposure is minimal. Technetium-99 can also be used to test thyroid function. Bones, the heart, the brain, the liver, the lungs, and many other organs can be imaged in similar ways by using the appropriate radioactive isotope.




Figure **3.4.2**: Medical Diagnostics. Radioactive iodine can be used to image the thyroid gland for diagnostic purposes. (CC BY 3.0 Unported; Myohan via Wikipedia).

Very little radioactive material is needed in these diagnostic techniques because the radiation emitted is so easy to detect. However, therapeutic applications usually require much larger doses because their purpose is to preferentially kill diseased tissues. For example, if a thyroid tumor is detected, a much larger infusion (thousands of rem, as opposed to a diagnostic dose of less then 40 rem) of iodine-131 could help destroy the tumor cells. Similarly, radioactive strontium is used to not only detect but also ease the pain of bone cancers. Table **3.4.1** lists several radioactive isotopes and their medical uses.

Table **3.4.1**: Some Radioactive Isotopes That Have Medical Applications

Isotope	Use
^{32}P	cancer detection and treatment, especially in eyes and skin
^{59}Fe	anemia diagnosis
^{60}Co	gamma ray irradiation of tumors
^{99m}Tc	brain, thyroid, liver, bone marrow, lung, heart, and intestinal scanning; blood volume determination
^{131}I	diagnosis and treatment of thyroid function
^{133}Xe	lung imaging
^{198}Au	liver disease diagnosis

In addition to the direct application of radioactive isotopes to diseased tissue, the gamma ray emissions of some isotopes can be directed toward the tissue to be destroyed. Cobalt-60 is a useful isotope for this kind of procedure.

To Your Health: Positron Emission Tomography Scans

One relatively rare form of radioactivity is called *positron emission*. It is similar to beta particle emission, except that instead of emitting an electron, a nucleus emits a positively charged electron, called a *positron*. A positron is actually a piece of antimatter; therefore, when a positron encounters an electron, both particles are converted into high-energy gamma radiation.

Isotopes that emit positrons can be employed in a medical imaging technique called *positron emission tomography* (*PET*). A patient receives a compound containing a positron-emitting isotope, either intravenously or by ingestion. The radioactive compound travels throughout the body, and the patient is then pushed slowly through a ring of sensors that detect the gamma radiation given off by the annihilation of positrons and electrons. A computer connected to the sensors constructs a three-dimensional image of the interior of part or all of the patient's body, allowing doctors to see organs or tumors or regulate the function of various organs (such as the brain or the heart) to diagnose the medical condition of the patient.



Figure **3.4.3**: (left) Combined apparatus for positron emission tomography (PET) and X-ray computer tomography (CT), Siemens Biograph. (CC BY-SA 3.0 Unported; Brudersohn via Wikipedia) (right) Whole-body PET scan using 18F-FDG (Public domain; Jens Maus via Wikipedia)

Two isotopes that undergo positron emission are carbon-11 and fluorine-18, with half-lives of 20.4 and 110 min, respectively. Both isotopes can be incorporated into sugar molecules and introduced into the body. Doctors can use the intensity of gamma ray emission to find tissues that metabolize the sugar faster than other tissues; fast-metabolizing tissue is one sign of a malignant (i.e., cancerous) tumor. Researchers use similar techniques to map areas of the brain that are most active during specific tasks, such as reading or speaking.

PET is one of many diagnostic and treatment methods that physicians use to improve the quality of our lives. It is one of the many positive uses of radioactivity in society.

Key Takeaway

- Radioactivity has several practical applications, including tracers, medical applications, dating once-living objects, and the preservation of food.

3.4: Uses of Radioactive Isotopes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.4: Uses of Radioactive Isotopes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.5: Nuclear Energy

Learning Objectives

- Explain where nuclear energy comes from.
- Describe the difference between fission and fusion.

Nuclear changes occur with a simultaneous release of energy. Where does this energy come from? If we could precisely measure the masses of the reactants and the products of a nuclear reaction, we would notice that the amount of mass drops slightly in the conversion from reactants to products. Consider the following nuclear reaction, in which the molar mass of each species is indicated to four decimal places:

$$\underbrace{^{235}\text{U}}_{235.0439} \rightarrow \underbrace{^{139}\text{Ba}}_{138.9088} + \underbrace{^{94}\text{Kr}}_{93.9343} + \underbrace{2\,^{1}\text{n}}_{2\times1.0087}$$

If we compare the mass of the reactant (235.0439) to the masses of the products (sum = 234.8605), we notice a mass difference of −0.1834 g, or −0.0001834 kg. Where did this mass go?

According to Albert Einstein's theory of relativity, energy (E) and mass (m) are related by the following equation:

$$E = mc^2 \tag{3.5.1}$$

where c is the speed of light, or 3.00 × 10^8 m/s. In the course of the uranium nuclear chemical reaction, the mass difference is converted to energy, which is given off by the reaction:

$$\begin{aligned} E &= (-0.0001834\ kg)(3.00 \times 10^8\ m/s)^2 \\ &= -1.65 \times 10^{13}\ J \\ &= -1.65 \times 10^{10}\ kJ \end{aligned}$$

Hence, 16.5 billion kJ of energy are given off every time 1 mol of uranium-235 undergoes this nuclear reaction. This is an extraordinary amount of energy. Compare it to combustion reactions of hydrocarbons, which give off about 650 kJ/mol of energy for every CH_2 unit in the hydrocarbon—on the order of *hundreds* of kilojoules per mole. Nuclear reactions give off *billions* of kilojoules per mole.

If this energy could be properly harvested, it would be a significant source of energy for our society. Nuclear energy involves the controlled harvesting of energy from fission reactions. The reaction can be controlled because the fission of uranium-235 (and a few other isotopes, such as plutonium-239) can be artificially initiated by injecting a neutron into a uranium nucleus. The overall nuclear equation, with energy included as a product, is then as follows:

$$^{235}\text{U} + {}^{1}\text{n} \longrightarrow {}^{139}\text{Ba} + {}^{94}\text{Kr} + 3\,{}^{1}\text{n} + \text{energy}$$

Thus, by the careful addition of extra neutrons into a sample of uranium, we can control the fission process and obtain energy that can be used for other purposes.

The Curie Family

Artificial or induced radioactivity was first demonstrated in 1934 by Irène Joliot-Curie and Frédéric Joliot, the daughter and son-in-law of Marie Curie.

Example 3.5.1

Plutonium-239 can absorb a neutron and undergo a fission reaction to produce an atom of gold-204 and an atom of phosphorus-31. Write the balanced nuclear equation for the process and determine the number of neutrons given off as part of the reaction.

Solution

Using the data given, we can write the following initial equation:

$$^{1}_{0}\text{n} + {}^{239}_{94}\text{Pu} \rightarrow {}^{204}_{79}\text{Au} + {}^{31}_{15}\text{P} + ?{}^{1}_{0}\text{n}$$

In balanced nuclear equations, the sums of the subscripts on each sides of the equation are the same, as are the sums of the superscripts. The subscripts are already balanced: 0 + 94 = 94 and 79 + 15 = 94. The superscripts on the left equal 240 (1 + 239) but equal 235 (204 + 31) on the right. We need five more mass number units on the right. Five neutrons should be the products of the process for the mass numbers to balance. (Because the atomic number of a neutron is zero, including five neutrons on the right does not change the overall sum of the subscripts.) Thus, the balanced nuclear equation is as follows:

$$^{1}_{0}\text{n} + {}^{239}_{94}\text{Pu} \rightarrow {}^{204}_{79}\text{Au} + {}^{31}_{15}\text{P} + 5{}^{1}_{0}\text{n}$$

We predict that the overall process will give off five neutrons.

? Exercise 3.5.1

Uranium-238 can absorb a neutron and undergo a fission reaction to produce an atom of cesium-135 and an atom of rubidium-96. Write the balanced nuclear equation for the process and determine the number of neutrons given off as part of the reaction.

Answer

$$^{1}_{0}\text{n} + {}^{238}_{92}\text{U} \rightarrow {}^{135}_{55}\text{Cs} + {}^{96}_{37}\text{Rb} + 8{}^{1}_{0}\text{n}$$

We predict that the overall process will give off eight neutrons.

A nuclear reactor is an apparatus designed to carefully control the progress of a nuclear reaction and extract the resulting energy for useful purposes. Figure 3.5.1 shows a simplified diagram of a nuclear reactor. The energy from the controlled nuclear reaction converts liquid water into high-pressure steam, which is used to run turbines that generate electricity.




Figure 3.5.1: A Diagram of a Nuclear Power Plant for Generating Electricity. The two main components of the power plant are the nuclear reactor itself and the steam-driven turbine and electricity generator.

Notice that the fission of uranium produces two more free neutrons than were present to begin with. These neutrons can themselves stimulate other uranium nuclei to undergo fission, releasing yet more energy and even more neutrons, which can in turn induce even more uranium fission. A single neutron can thus begin a process that grows exponentially in a phenomenon called a chain reaction:

$$1 \rightarrow 2 \rightarrow 4 \rightarrow 8 \rightarrow 16 \rightarrow 32 \rightarrow 64 \rightarrow 128 \rightarrow 256 \rightarrow 512 \rightarrow 1{,}024 \rightarrow 2{,}048 \rightarrow 4{,}096 \rightarrow 8{,}192 \rightarrow 16{,}384 \rightarrow \ldots$$

Because energy is produced with each fission event, energy is also produced exponentially and in an uncontrolled fashion. The quick production of energy creates an explosion. This is the mechanism behind the atomic bomb.

The first controlled chain reaction was achieved on December 2, 1942, in an experiment supervised by Enrico Fermi in a laboratory underneath the football stadium at the University of Chicago.

Although fairly simple in theory, an atomic bomb is difficult to produce, in part because uranium-235, the isotope that undergoes fission, makes up only 0.7% of natural uranium; the rest is mostly uranium-238, which does not undergo fission. (Remember that the radioactive process that a nucleus undergoes is characteristic of the isotope.) To make uranium useful for nuclear reactors, the uranium in uranium-235 must be *enriched* to about 3%. Enrichment of uranium is a laborious and costly series of physical and chemical separations. To be useful in an atomic bomb, the uranium in uranium-235 must be enriched to 70% or more. At lesser concentrations, the chain reaction cannot sustain itself, so no explosion is produced.

Fusion is another nuclear process that can be used to produce energy. In this process, smaller nuclei are combined to make larger nuclei, with an accompanying release of energy. One example is the hydrogen fusion, which makes helium. While the steps of the process are complicated, the net reaction is:

$$4\,^{1}\mathrm{H} \rightarrow {}^{4}\mathrm{He} + 2.58 \times 10^{12}\ J$$

Notice that the amount of energy given off per mole of reactant is only a fraction of the amount given off by the fission of 1 mol of uranium-235. On a mass (per gram) basis, however, the hydrogen fusion emits many times more energy than fission does. In addition, the product of fission is helium gas, not a wide range of isotopes (some of which are also radioactive) produced by fission.

The practical problem is that to perform fusion, extremely high pressures and temperatures are necessary. Currently, the only known stable systems undergoing fusion are the interiors of stars. The conditions necessary for fusion can be created using an atomic bomb, but the resulting fusion is uncontrollable (and the basis for another type of bomb, a hydrogen bomb). Currently, researchers are looking for safe, controlled ways of producing useful energy using fusion.

Career Focus: Nuclear Medicine Technologist

Generally speaking, a radiological technician deals with X ray equipment and procedures. A *nuclear medicine technologist* has similar responsibilities, using compounds containing radioactive isotopes to help diagnose and treat disease.

Nuclear medicine technologists administer the substances containing the radioactive isotope and subsequently operate the apparatus that detects the radiation produced by radioactive decay. The apparatus may be as simple as a piece of photographic film or as complex as a series of computer-controlled electronic detectors. The images obtained by the technologist are interpreted by a specially trained physician.

One of the chief responsibilities of a nuclear medicine technologist is safety. Improper exposure to radioactivity can be harmful to both patient and technologist alike. Therefore, the technologist must adhere to strict safety standards to keep unnecessary exposure as low as possible. The technologist must also know how to dispose of waste materials safely and appropriately.

Key Takeaways

- Nuclear energy comes from tiny mass changes in nuclei as radioactive processes occur.
- In fission, large nuclei break apart and release energy; in fusion, small nuclei merge together and release energy.

3.5: Nuclear Energy is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.5: Nuclear Energy** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.6: 3.6 Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Some atoms have unstable nuclei that emit particles and high-energy electromagnetic radiation to form new elements that are more stable. This emission of particles and electromagnetic radiation is called **radioactivity**. There are three main types of spontaneous radioactive emission: **alpha particles**, which are equivalent to helium nuclei; **beta particles**, which are electrons; and **gamma radiation**, which is high-energy electromagnetic radiation. Another type of radioactive process is **spontaneous fission**, in which large nuclei spontaneously break apart into smaller nuclei and, often, neutrons. In all forms of radioactivity, new elements are formed from the radioactive reactants.

Radioactive isotopes decay at different rates. The rate of an isotope's decay is expressed as a **half-life**, which is the amount of time required for half of the original material to decay. The length of its half-life is a characteristic of the particular isotope and can range from less than microseconds to billions of years.

Amounts of radioactivity are measured in several different ways. A **becquerel** is equal to one radioactive decay per second. A **curie** represents 3.7×10^{10} decays per second. Other units describe the amount of energy absorbed by body tissues. One **rad** is equivalent to 0.01 joule of energy absorbed per gram of tissue. Different tissues react differently to different types of radioactivity. The **rem** unit takes into account not only the energy absorbed by the tissues, but also includes a numerical multiplication factor to account for the type of radioactivity and the type of tissue. The average annual radiation exposure of a person is less than 360 millirem, over 80% of which is from natural sources. Radioactivity can be detected using photographic film or other devices such as **Geiger counters**.

Radioactive isotopes have many useful applications. They can be used as **tracers** to follow the journey of a substance through a system, like an underground waterway or a metabolic pathway. Radioactive isotopes can be used to date objects, since the amount of parent and daughter isotopes can sometimes be measured very accurately. Radioactive emission can be used to sterilize food for a longer edible lifetime. There are also a number of diagnostic and therapeutic medical applications for radioactive isotopes.

Radioactive processes occur with simultaneous changes in energy. This **nuclear energy** can be used to generate power for human use. **Nuclear reactors** use the energy released by fission of large isotopes to generate electricity. When carefully controlled, fission can produce a **chain reaction** that facilitates the continuous production of energy. If not carefully controlled, a very quick production of energy can result, as in an **atomic bomb**. Natural uranium does not contain enough of the proper isotope of uranium to work in a nuclear reactor, so it must first be **enriched** in uranium-235. Forcing small nuclei together to make larger nuclei, a process called **fusion**, also gives off energy; however, scientists have yet to achieve a controlled fusion process.

Template:HideTOC

3.6: 3.6 Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

SECTION OVERVIEW

3.7: Ionic Bonding and Simple Ionic Compounds

There are only 118 known chemical elements but tens of millions of known chemical compounds. Compounds can be very complex combinations of atoms, but many important compounds are fairly simple. Table salt, as we have seen, consists of only two elements: sodium and chlorine. Nevertheless, the compound has properties completely different from either elemental sodium (a chemically reactive metal) or elemental chlorine (a poisonous, green gas). We will see additional examples of such differences in this chapter and Chapter 4, as we consider how atoms combine to form compounds.

3.7: Ionic Bonding and Simple Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

4: Ionic Bonding and Simple Ionic Compounds

There are only 118 known chemical elements but tens of millions of known chemical compounds. Compounds can be very complex combinations of atoms, but many important compounds are fairly simple. Table salt, as we have seen, consists of only two elements: sodium and chlorine. Nevertheless, the compound has properties completely different from either elemental sodium (a chemically reactive metal) or elemental chlorine (a poisonous, green gas). We will see additional examples of such differences in this chapter and Chapter 4, as we consider how atoms combine to form compounds.

4.1: Two Types of Bonding

4.2: Ions

4.3: Formulas for Ionic Compounds

4.4: Ionic Nomenclature

4.5: Chapter Summary

4: Ionic Bonding and Simple Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

4.1: Two Types of Bonding

Learning Objectives

- Define the octet rule.
- Describe how ionic bonds are formed.

Atoms can join together by forming a chemical bond, which is a very strong attraction between two atoms. Chemical bonds are formed when electrons in different atoms interact with each other to make an arrangement that is more stable than when the atoms are apart.

What causes atoms to make a chemical bond with other atoms, rather than remaining as individual atoms? A clue comes by considering the noble gas elements, the rightmost column of the periodic table. These elements—helium, neon, argon, krypton, xenon, and radon—do not form compounds very easily, which suggests that they are especially stable as lone atoms. What else do the noble gas elements have in common? Except for helium, they all have eight valence electrons. Chemists have concluded that atoms are especially stable if they have eight electrons in their outermost shell. This useful rule of thumb is called the **octet rule**, and it is a key to understanding why compounds form.

> *Of the noble gases, only krypton, xenon, and radon have been found to make compounds.*

There are two ways for an atom that does not have an octet of valence electrons to obtain an octet in its outer shell. One way is the transfer of electrons between two atoms until all atoms have octets. Because some atoms will lose electrons and some atoms will gain electrons, there is no overall change in the number of electrons, but individual atoms acquire a nonzero electric charge. Those that lose electrons become positively charged, and those that gain electrons become negatively charged. Charged atoms are called ions. Because opposite charges attract (while like charges repel), these oppositely charged ions attract each other, forming **ionic bonds**. The resulting compounds are called **ionic compounds** and are the primary subject of this chapter.

The second way for an atom to obtain an octet of electrons is by sharing electrons with another atom. These shared electrons simultaneously occupy the outermost shell of more than one atom. The bond made by electron sharing is called a **covalent bond**.

Despite our focus on the octet rule, we must remember that for small atoms, such as hydrogen, helium, and lithium, the first shell is, or becomes, the outermost shell and hold only two electrons. Therefore, these atoms satisfy a "**duet rule**" rather than the octet rule.

Example 4.1.1

A sodium atom has one valence electron. Do you think it is more likely for a sodium atom to lose one electron or gain seven electrons to obtain an octet?

Solution

Although either event is possible, a sodium atom is more likely to lose its single valence electron. When that happens, it becomes an ion with a net positive charge. This can be illustrated as follows:

Table explaining the solution to the example.

Sodium atom		Sodium ion	
11 protons	11+	11 protons	11+
11 electrons	11−	10 electrons	10−
	0 overall charge		+1 overall charge

? Exercise 4.1.1

A fluorine atom has seven valence electrons. Do you think it is more likely for a fluorine atom to lose seven electrons or gain one electron to obtain an octet? Write the formula of the resulting ion.

Answer

The process that involves less number of electrons is more favorable. Fluorine would gain one electron. The formula of the resulting ion is F^-.

Key Takeaways

- Atoms have a tendency to have eight electrons in their valence shell.
- The attraction of oppositely charged ions is what makes ionic bonds.

Exercises

1. What is the octet rule?

2. How are ionic bonds formed?

3. Why is an ionic compound unlikely to consist of two positively charged ions?

4. Why is an ionic compound unlikely to consist of two negatively charged ions?

5. A calcium atom has two valence electrons. Do you think it will lose two electrons or gain six electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

6. An aluminum atom has three valence electrons. Do you think it will lose three electrons or gain five electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

7. A selenium atom has six valence electrons. Do you think it will lose six electrons or gain two electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

8. An iodine atom has seven valence electrons. Do you think it will lose seven electrons or gain one electron to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

Answers

1. The octet rule is the concept that atoms tend to have eight electrons in their valence electron shell.

2. Ionic bonds are formed by the attraction between oppositely charged ions.

3. Positive charges repel each other, so an ionic compound is not likely between two positively charged ions.

4. Negative charges repel each other also. 5. Ca atom is more likely to lose two electrons. It will become Ca^{2+} ion. 6. An Al atom is more likely to lose three electrons. It will become Al^{3+} ion.

7. Selenium is more likely to gain two electrons. It will become Se^{2-} ion.

8. Iodine is more likely to gain one electron. It will become I^- ion.

4.1: Two Types of Bonding is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **3.1: Two Types of Bonding** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.2: Ions

Learning Objectives

- Define the two types of ions.
- Use Lewis diagrams to illustrate ion formation.

Most atoms do not have eight electrons in their valence electron shell. Some atoms have only a few electrons in their outer shell, while some atoms lack only one or two electrons to have an octet. In cases where an atom has three or fewer valence electrons, the atom may lose those valence electrons quite easily until what remains is a lower shell that contains an octet. Atoms that lose electrons acquire a positive charge as a result because they are left with fewer negatively charged electrons to balance the positive charges of the protons in the nucleus. Positively charged ions are called cations. Most metals become cations when they make ionic compounds.

Some atoms have nearly eight electrons in their valence shell and can gain additional valence electrons until they have an octet. When these atoms gain electrons, they acquire a negative charge because they now possess more electrons than protons. Negatively charged ions are called anions. Most nonmetals become anions when they make ionic compounds.

> The names for positive and negative ions are pronounced CAT-eye-ons (cations) and ANN-eye-ons (anions), respectively.

Electron Transfer

We can use electron configurations to illustrate the electron transfer process between sodium atoms and chlorine atoms.

$$\text{Na: } 1s^2 2s^2 2p^6 3s^1$$

As demonstrated here, a sodium atom (Na) has one valence electron in the third principal energy level. It is likely to achieve an octet in its outermost shell by losing its one valence electron. The cation produced in this way, Na^+, is called the sodium ion to distinguish it from the element. The sodium ion, Na^+, has the electron configuration with an octet of electrons from the second principal energy level. It is now the same as that of the noble gas neon. The term **isoelectronic** refers to an atom and an ion of a different atom (or two different ions) that have the same electron configuration. The sodium ion is isoelectronic with the neon atom. The equation below illustrates this process.

$$\begin{array}{ccc} \mathbf{Na} & \rightarrow & \mathbf{Na^+ + e^-} \\ \mathbf{1s^2\ 2s^2\ 2p^6\ 3s^1} & & \mathbf{1s^2\ 2s^2\ 2p^6\ (octet)} \end{array}$$

Figure **4.2.1** is a graphical depiction of this process.




Figure **4.2.1**: The Formation of a Sodium Ion. On the left, a sodium atom has 11 electrons. On the right, the sodium ion only has 10 electrons and a 1+ charge.

Now, let's consider chlorine atom, Cl: $1s^2 2s^2 2p^6 3s^2 3p^5$

Only one more electron is needed to achieve an octet in chlorine's valence shell. When a chlorine atom gains an electron, its outermost principal energy level achieves an octet. In this case, the ion has the same outermost shell as the original atom, but now that shell has eight electrons in it. Once again, the octet rule has been satisfied. The resulting anion, Cl^-, is called the chloride ion;

note the slight change in the suffix (*-ide* instead of *-ine*) to create the name of this anion. This process is illustrated below. (In table salt, this electron comes from the sodium atom.)

$$\begin{array}{lcl} \mathrm{Cl} + \mathrm{e}^- & \rightarrow & \mathrm{Cl}^- \\ 1s^2\,2s^2\,2p^6\,3s^2\,3p^5 & & 1s^2\,2s^2\,2p^6\,3s^2\,3p^6\,(\text{octet}) \end{array}$$

Figure **4.2.2** is a graphical depiction of this process.




Figure **4.2.2**: The Formation of a Chlorine Ion. On the left, the chlorine atom has 17 electrons. On the right, the chloride ion has 18 electrons and has a 1− charge.

With two oppositely charged ions, there is an electrostatic attraction between them because opposite charges attract. The resulting combination is the compound sodium chloride. Notice that there are no leftover electrons. The number of electrons lost by the sodium atom (one) equals the number of electrons gained by the chlorine atom (one), so the compound is electrically neutral. In macroscopic samples of sodium chloride, there are billions and billions of sodium and chloride ions, although there is always the same number of cations and anions.

✔ Example 4.2.1

Write the electron configuration of aluminum atom (Z=13). How many electrons must Al lose/gain to achieve octet? Write the formula of the resulting ion and its electron configuration.

Solution

The electron configuration of Al atom is $1s^2 2s^2 2p^6 3s^2 3p^1$. The second shell has octet ($2s^2 2p^6$) while the valence shell has 3 electrons ($3s^2 3p^1$). Mg can achieve octet by losing the 3 valence electrons. The resulting cation is Al^{3+} with electron configuration, $1s^2 2s^2 2p^6$.

4.2.1 Exercise

Write the electron configuration of oxygen atom (Z=8). How many electrons must O lose/gain to achieve octet? Write the formula of the resulting ion and its electron configuration.

Answer

The electron configuration of O atom is $1s^2 2s^2 2p^4$. The second shell has six electrons ($2s^2 2p^4$) and needs two electrons to achieve octet. Oxygen will gain 2 electrons. The resulting anion is O^{2-} with electron configuration, $1s^2 2s^2 2p^6$.

In many cases, elements that belong to the same group (vertical column) on the periodic table form ions with the same charge because they have the same number of valence electrons. Thus, the periodic table becomes a tool for remembering the charges on many ions. For example, all ions made from alkali metals, the first column on the periodic table, have a 1+ charge. Ions made from alkaline earth metals, the second group on the periodic table, have a 2+ charge. On the other side of the periodic table, the next-to-last column, the halogens, form ions having a 1− charge. Figure **4.2.3** shows how the charge on many ions can be predicted by the location of an element on the periodic table. Note the convention of first writing the number and then the sign on a multiply charged ion. The barium cation is written Ba^{2+}, not Ba^{+2}.




Figure **4.2.3**: Predicting Ionic Charges. The charge that an atom acquires when it becomes an ion is related to the structure of the periodic table. Within a group (family) of elements, atoms form ions of a certain charge.

Example 4.2.2

Which of these ions is *not likely* to form?

a. Mg^{+}
b. K^{+}

Solution

(a) Mg is in Group 2A and has two valence electrons. It achieves octet by losing two electrons to form Mg^{2+} cation. Losing only one electron to form Mg^{+} does not make an octet, hence, Mg^{+} is not likely to form.

4.2.2 Exercise

Which of these ions is *not likely* to form?

a. S^{3-}
b. N^{3-}

Answer

(a) S is in Group 6A and has six valence electrons. It achieves octet by gaining two electrons to form S^{2-} anion. Gaining three electrons to form S^{3-} does not make it octet, hence, S^{3-} is not likely to form.

Lewis Diagrams

Chemists use simple diagrams to show an atom's valence electrons and how they transfer. These diagrams have two advantages over the electron shell diagrams. First, they show only valence electrons. Second, instead of having a circle around the chemical symbol to represent the electron shell, they have up to eight dots around the symbol; each dot represents a valence electron. These dots are arranged to the right and left and above and below the symbol, **with no more than two dots on a side**. For example, the representation for sodium is as follows:



and the representation for chlorine is as follows:



> For the above diagrams, it does not matter what sides the dots are placed on in Lewis diagrams as long as each side has a maximum of two dots.

These diagrams are called **Lewis electron dot diagrams**, or simply Lewis diagrams, after Gilbert N. Lewis, the American chemist who introduced them. To write an element's Lewis dot symbol, place the dots representing its valence electrons, one at a time, around the element's chemical symbol. Up to four dots are placed above, below, to the left, and to the right of the symbol (in any order, as long as elements with four or fewer valence electrons have no more than one dot in each position). The next dots, for elements with more than four valence electrons, are again distributed one at a time, each paired with one of the first four. In other words, **place the dots singly on each side before pairing them**. The Lewis electron dot diagram of fluorine, for example, with seven valence electrons, is constructed as follows:





Figure 4.2.4 shows the electron configurations and Lewis diagrams of the elements lithium through neon, which is the entire second period of the periodic table. For the **main group** elements, the **number of valence electrons** is the **same as the group number** listed at the top of the periodic table.



Figure 4.2.4: Lewis Electron Dot Diagrams of the Elements Lithium through Neon

The transfer of electrons can be illustrated easily with Lewis diagrams:




In representing the final formula, the dots are omitted.

✔ Example 4.2.3

Starting with lithium and bromine atoms, use Lewis diagrams to show the formation of the ionic compound LiBr.

Solution


From the periodic table, we see that lithium is in the same column as sodium, so it will have the same valence shell electron configuration. That means that the neutral lithium atom will have the same Lewis diagram that the sodium atom has. Similarly, bromine is in the same column as chlorine, so it will have the same Lewis diagram that chlorine has. Therefore,




? Exercise 4.2.3

Starting with magnesium and oxygen atoms, use Lewis diagrams to show the formation of the ionic compound MgO.

Answer



Some ionic compounds have different numbers of cations and anions. In those cases, electron transfer occurs between more than one atom. For example, here is the formation of $MgBr_2$:



Notice that in this example there are two bromide ions (1– charge) needed for every one magnesium ion (2+ charge) in order for the overall charge of the compound to equal zero. This is called **charge balance**. The number of each type of ion is indicated in the formula by the subscript.

Most of the elements that make ionic compounds form an ion that has a characteristic charge. For example, sodium makes ionic compounds in which the sodium ion always has a 1+ charge. Chlorine makes ionic compounds in which the chloride ion always has a 1− charge. Some elements, especially transition metals, can form ions of multiple charges. Figure **4.2.5** shows the characteristic charges for some of these ions. As we saw in Figure **4.2.1**, there is a pattern to the charges on many of the main group ions, but there is no simple pattern for transition metal ions (or for the larger main group elements).




Figure **4.2.5**: Charges of the Monatomic Ions. Note that some atoms commonly form ions of variable charges.

Key Takeaways

- Ions can be positively charged or negatively charged.
- A Lewis diagram is used to show how electrons are transferred to make ions and ionic compounds.

Exercises

1. What are the two types of ions?
2. Use Lewis diagrams to illustrate the formation of an ionic compound from a potassium atom and an iodine atom.

3. When the following atoms become ions, what charges do they acquire?

a. Li
b. S
c. Ca
d. F

4. Identify each as a cation, an anion, or neither.

a. H^+
b. Cl^-
c. O_2
d. Ba^{2+}
e. CH_4
f. CS_2

5. Identify each as a cation, an anion, or neither.

a. NH_3
b. Br^-
c. H^-
d. Hg^{2+}

e. CCl_4
f. SO_3

6. Write the electron configuration for each ion.

a. Li^+
b. Mg^{2+}
c. F^-
d. S^{2-}

7. Write the electron configuration for each ion.

a. Na^+
b. Be^{2+}
c. Cl^-
d. O^{2-}

8. Draw Lewis diagrams for the ions listed in Exercise 6. Also include Lewis diagrams for the respective neutral atoms as a comparison.

9. Draw Lewis diagrams for the ions listed in Exercise 7. Also include Lewis diagrams for the respective neutral atoms as a comparison.

10. Using Lewis diagrams, show the electron transfer for the formation of LiF.

11. Using Lewis diagrams, show the electron transfer for the formation of MgO.

12. Using Lewis diagrams, show the electron transfer for the formation of Li_2O.

13. Using Lewis diagrams, show the electron transfer for the formation of CaF_2.

14. What characteristic charge do atoms in the first column of the periodic table have when they become ions?


15. What characteristic charge do atoms in the second column of the periodic table have when they become ions?

16. What characteristic charge do atoms in the third-to-last column of the periodic table have when they become ions?

17. What characteristic charge do atoms in the next-to-last column of the periodic table have when they become ions?

Answers

1. Cations (positive charged) and anions (negative charged).

2. 

3.
a. 1+
b. 2−
c. 2+
d. 1−

4.
a. cation
b. anion
c. neither
d. cation
e. neither
f. neither

5.
a. neither
b. anion

c. anion
d. cation
e. neither
f. neither

6.

a. $1s^2$
b. $1s^22s^22p^6$
c. $1s^22s^22p^6$
d. $1s^22s^22p^63s^23p^6$

7.

a. $1s^22s^22p^6$
b. $1s^2$
c. $1s^22s^22p^63s^23p^6$
d. $1s^22s^22p^6$

8.

a. $Li\bullet$, Li^+

b. $\bullet Mg\bullet$, Mg^{2+}

c. $:\ddot{F}:$, $:\ddot{F}:^-$

d. $:\ddot{S}:$, $:\ddot{S}:^{2-}$


$Na\bullet$, Na^+

$\bullet Be\bullet$, Be^{2+}

$:\ddot{Cl}:$, $:\ddot{Cl}:^-$


$:\ddot{O}:$, $:\ddot{O}:^{2-}$

9.



10.

11.



12.



13.

14. 1+

15. 2+

16. 2−

17. 1−

4.2: Ions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **3.2: Ions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.3: Formulas for Ionic Compounds

Learning Objectives

- Write the chemical formula for a simple ionic compound.
- Recognize polyatomic ions in chemical formulas.

We have already encountered some chemical formulas for simple ionic compounds. A chemical formula is a concise list of the elements in a compound and the ratios of these elements. To better understand what a chemical formula means, we must consider how an ionic compound is constructed from its ions.

Ionic compounds exist as alternating positive and negative ions in regular, three-dimensional arrays called crystals (Figure 4.3.1). As you can see, there are no individual NaCl "particles" in the array; instead, there is a continuous lattice of alternating sodium and chloride ions. However, we can use the ratio of sodium ions to chloride ions, expressed in the lowest possible whole numbers, as a way of describing the compound. In the case of sodium chloride, the ratio of sodium ions to chloride ions, expressed in lowest whole numbers, is 1:1, so we use NaCl (one Na symbol and one Cl symbol) to represent the compound. Thus, NaCl is the chemical formula for sodium chloride, which is a concise way of describing the relative number of different ions in the compound. A macroscopic sample is composed of myriads of NaCl pairs; each individual pair called a **formula unit**. Although it is convenient to think that NaCl crystals are composed of individual NaCl units, Figure 4.3.1 shows that no single ion is exclusively associated with any other single ion. Each ion is surrounded by ions of opposite charge.




Figure 4.3.1: A Sodium Chloride Crystal. A crystal contains a three-dimensional array of alternating positive and negative ions. The precise pattern depends on the compound. A crystal of sodium chloride, shown here, is a collection of alternating sodium and chlorine ions.

The formula for an ionic compound follows several conventions. First, the **cation** is written **before** the **anion**. Because most metals form cations and most nonmetals form anions, formulas typically list the metal first and then the nonmetal. Second, **charges** are **not written** in a formula. Remember that in an ionic compound, the component species are ions, not neutral atoms, even though the formula does not contain charges. Finally, the proper formula for an ionic compound always has ***a net zero charge,*** meaning the total positive charge must equal the total negative charge. To determine the proper formula of any combination of ions, determine how many of each ion is needed to balance the total positive and negative charges in the compound.

> *This rule is ultimately based on the fact that matter is, overall, electrically neutral.*

By convention, assume that there is only one atom if a subscript is not present. We do not use 1 as a subscript.

If we look at the ionic compound consisting of lithium ions and bromide ions, we see that the lithium ion has a 1+ charge and the bromide ion has a 1− charge. Only one ion of each is needed to balance these charges. The formula for lithium bromide is LiBr.

When an ionic compound is formed from magnesium and oxygen, the magnesium ion has a 2+ charge, and the oxygen atom has a 2− charge. Although both of these ions have higher charges than the ions in lithium bromide, they still balance each other in a one-to-one ratio. Therefore, the proper formula for this ionic compound is MgO.

Now consider the ionic compound formed by magnesium and chlorine. A magnesium ion has a 2+ charge, while a chlorine ion has a 1− charge:

$$Mg^{2+}Cl^{-}$$


Combining one ion of each does not completely balance the positive and negative charges. The easiest way to balance these charges is to assume the presence of *two* chloride ions for each magnesium ion:

$$Mg^{2+}\ Cl^{-}\ Cl^{-}$$


Now the positive and negative charges are balanced. We could write the chemical formula for this ionic compound as **MgClCl**, but the convention is to use a numerical subscript when there is more than one ion of a given type—**$MgCl_2$**. This chemical formula says that there are one magnesium ion and *two* chloride ions in this formula. (Do not read the "Cl_2" part of the formula as a molecule of the diatomic elemental chlorine. Chlorine does not exist as a diatomic element in this compound. Rather, it exists as two individual chloride ions.) By convention, the **lowest whole number ratio** is used in the formulas of ionic compounds. The formula **Mg_2Cl_4** has balanced charges with the ions in a 1:2 ratio, but it is not the lowest whole number ratio.

> By convention, the lowest whole-number ratio of the ions is used in ionic formulas. There are exceptions for certain ions, such as **Hg_2^{2+}**.

For compounds in which the ratio of ions is not as obvious, the subscripts in the formula can be obtained by **crossing charges**: use the absolute value of the charge on one ion as the subscript for the other ion. This method is shown schematically in Figure 3.3.2.



The image shows M with a charge of n+ and X with a charge of m-. The charge on the M becomes the subscript of X, and the charge on X becomes the subscript of M, making the final product M subscript m X subscript n.



The image asks to write the formula for the compound formed by aluminum and oxygen. It shows Al with a charge of 3+ and O with a charge of 2-. By crossing charges, the final result is Al subscript 2 O subscript 3.

Figure **4.3.2**: Crossing charges. One method for obtaining subscripts in the empirical formula is by crossing charges.

When crossing charges, it is sometimes necessary to reduce the subscripts to their simplest ratio to write the empirical formula. Consider, for example, the compound formed by Pb^{4+} and O^{2-}. Using the absolute values of the charges on the ions as subscripts gives the formula Pb_2O_4. This simplifies to its correct empirical formula **PbO_2**. The empirical formula has one Pb^{4+} ion and two O^{2-} ions.

✔ Example 4.3.1

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the sodium ion and the sulfur ion
b. the aluminum ion and the fluoride ion
c. the 3+ iron ion and the oxygen ion

Solution

a. To obtain a valence shell octet, sodium forms an ion with a 1+ charge, while the sulfur ion has a 2− charge. Two sodium 1+ ions are needed to balance the 2− charge on the sulfur ion. Rather than writing the formula as $\mathbf{NaNaS}$, we shorten it by convention to $\mathbf{Na_2S}$.

b. The aluminum ion has a 3+ charge, while the fluoride ion formed by fluorine has a 1− charge. Three fluorine 1− ions are needed to balance the 3+ charge on the aluminum ion. This combination is written as $\mathbf{AlF_3}$.

c. Iron can form two possible ions, but the ion with a 3+ charge is specified here. The oxygen atom has a 2− charge as an ion. To balance the positive and negative charges, we look to the least common multiple—6: two iron 3+ ions will give 6+, while three 2− oxygen ions will give 6−, thereby balancing the overall positive and negative charges. Thus, the formula for this ionic compound is $\mathbf{Fe_2O_3}$. Alternatively, use the crossing charges method shown in Figure 3.3.2.

? Exercise 4.3.1

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the calcium ion and the oxygen ion
b. the 2+ copper ion and the sulfur ion
c. the 1+ copper ion and the sulfur ion

Answer a:

CaO

Answer b:


CuS

Answer c:

Cu_2S

Polyatomic Ions

Some ions consist of groups of atoms ***covalently*** bonded together and have an **overall electric charge**. Because these ions contain more than one atom, they are called polyatomic ions. The Lewis structures, names and formulas of some polyatomic ions are found in Table 3.3.1.



Some Polyatomic Ions


Polyatomic ions have defined formulas, names, and charges that cannot be modified in any way. Table **4.3.2** lists the ion names and ion formulas of the most common polyatomic ions. For example, $\mathbf{NO_3^-}$ is the nitrate ion; it has one nitrogen atom and three oxygen atoms and an overall 1− charge. Figure **4.3.2** lists the most common polyatomic ions.

Table **4.3.2**: Ion Names and Ion Formulas of Common Polyatomic Ions

Ion Name	Ion Formula
ammonium ion	NH_4^{+1}
hydroxide ion	OH^{-1}
cyanide ion	CN^{-1}
carbonate ion	CO_3^{-2}
bicarbonate or hydrogen carbonate	HCO_3^-
acetate ion	$C_2H_3O_2^{-1}$ **or** $CH_3CO_2^{-1}$
nitrate ion	NO_3^{-1}
nitrite ion	NO_2^{-1}
sulfate ion	SO_4^{-2}
sulfite ion	SO_3^{-2}
phosphate ion	PO_4^{-3}
phosphite ion	PO_3^{-3}

Note that only one polyatomic ion in this Table, the ammonium ion ($NH_4^{+1)}$, is a cation. This polyatomic ion contains one nitrogen and four hydrogens that collectively bear a +1 charge. The remaining polyatomic ions are all negatively-charged and, therefore, are classified as anions. However, only two of these, the hydroxide ion and the cyanide ion, are named using the "-ide" suffix that is typically indicative of negatively-charged particles. The remaining polyatomic anions, which all contain oxygen, in combination with another non-metal, exist as part of a series in which the number of oxygens within the polyatomic unit can vary. As has been repeatedly emphasized in several sections of this text, no two chemical formulas should share a common chemical name. A single suffix, "-ide," is insufficient for distinguishing the names of the anions in a related polyatomic series. Therefore, "-ate" and "-ite" suffixes are employed, in order to denote that the corresponding polyatomic ions are part of a series. Additionally, these suffixes also indicate the relative number of oxygens that are contained within the polyatomic ions. Note that all of the polyatomic ions whose names end in "-ate" contain one more oxygen than those polyatomic anions whose names end in "-ite." Unfortunately, much like the common system for naming transition metals, these suffixes only indicate the *relative* number of oxygens that are contained within the polyatomic ions. For example, the nitrate ion, which is symbolized as NO_3^{-1}, has one more oxygen than the nitrite ion, which is symbolized as NO_2^{-1}. However, the sulfate ion is symbolized as SO_4^{-2}. While both the nitrate ion and the sulfate ion share an "-ate" suffix, the former contains three oxygens, but the latter contains four. Additionally, both the nitrate ion and the sulfite ion contain three oxygens, but these polyatomic ions do not share a common suffix. Unfortunately, the relative nature of these suffixes mandates that the ion formula/ion name combinations of the polyatomic ions must simply be memorized.

The rule for constructing formulas for ionic compounds containing polyatomic ions is the same as for formulas containing monatomic (single-atom) ions: the positive and negative charges must balance. If more than one of a particular polyatomic ion is needed to balance the charge, the *entire formula* for the polyatomic ion must be enclosed in parentheses, and the numerical subscript is placed *outside* the parentheses. This is to show that the subscript applies to the entire polyatomic ion. Two examples are shown below:



The image asks a two-part question: Write the formula for the compound formed by (a) barium and nitrate, and (b) ammonium and phosphate. For (a), Ba has a charge of 2+ and NO_3 has a charge of -1. By crossing charges, the formula is $Ba(NO_3)_2$. For (b), NH_4 has a charge of +1 and PO_4 has a charge of 3-, so by crossing charges the formula is $(NH_4)_3PO_4$.

Example 4.3.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the potassium ion and the sulfate ion
b. the calcium ion and the nitrate ion

Solution

a. Potassium ions have a charge of 1+, while sulfate ions have a charge of 2−. We will need two potassium ions to balance the charge on the sulfate ion, so the proper chemical formula is $\mathbf{K_2SO_4}$.
b. Calcium ions have a charge of 2+, while nitrate ions have a charge of 1−. We will need two nitrate ions to balance the charge on each calcium ion. The formula for nitrate must be enclosed in parentheses. Thus, we write $\mathbf{Ca(NO_3)_2}$ as the formula for this ionic compound.

Exercise 4.3.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the magnesium ion and the carbonate ion
b. the aluminum ion and the acetate ion

Answer a:

Mg^{2+} and $CO_3^{2-} = MgCO_3$

Answer b:

Al^{3+} and $C_2H_3O_2^- = Al(C_2H_3O_2)_3$

Recognizing Ionic Compounds

There are two ways to recognize ionic compounds. First, compounds between **metal and nonmetal** elements are usually **ionic**. For example, $CaBr_2$ contains a metallic element (calcium, a group 2A metal) and a nonmetallic element (bromine, a group 7A nonmetal). Therefore, it is most likely an ionic compound. (In fact, it *is* ionic.) In contrast, the compound NO_2 contains two elements that are both nonmetals (nitrogen, from group 5A, and oxygen, from group 6A). It is not an ionic compound; it belongs to the category of covalent compounds discuss elsewhere. Also note that this combination of nitrogen and oxygen has no electric charge specified, so it is *not* the nitrite ion.

Second, if you recognize the formula of a **polyatomic ion** in a compound, the compound is **ionic**. For example, if you see the formula $\mathbf{Ba(NO_3)_2}$, you may recognize the "NO_3" part as the nitrate ion, $\mathbf{NO_3^-}$. (Remember that the convention for writing formulas for ionic compounds is not to include the ionic charge.) This is a clue that the other part of the formula, **Ba**, is actually the $\mathbf{Ba^{2+}}$ ion, with the 2+ charge balancing the overall 2− charge from the two nitrate ions. Thus, this compound is also ionic.

Example 4.3.3

Identify each compound as ionic or not ionic.

a. Na_2O
b. PCl_3
c. NH_4Cl
d. OF_2

Solution

a. Sodium is a metal, and oxygen is a nonmetal; therefore, Na_2O is expected to be ionic.
b. Both phosphorus and chlorine are nonmetals. Therefore, PCl_3 is not ionic.
c. The NH_4 in the formula represents the ammonium ion, NH_4^+, which indicates that this compound is ionic.
d. Both oxygen and fluorine are nonmetals. Therefore, OF_2 is not ionic.

Exercise 4.3.3

Identify each compound as ionic or not ionic.

a. N_2O
b. $FeCl_3$
c. $(NH_4)_3PO_4$
d. $SOCl_2$

Answer a:

not ionic

Answer b:

ionic

Answer c:

ionic

Answer d:

not ionic

Looking Closer: Blood and Seawater

Science has long recognized that blood and seawater have similar compositions. After all, both liquids have ionic compounds dissolved in them. The similarity may be more than mere coincidence; many scientists think that the first forms of life on Earth arose in the oceans. A closer look, however, shows that blood and seawater are quite different. A 0.9% solution of sodium chloride approximates the salt concentration found in blood. In contrast, seawater is principally a 3% sodium chloride solution, over three times the concentration in blood. Here is a comparison of the amounts of ions in blood and seawater:

Table showing a comparison of the amounts of ions in blood and seawater.

Ion	Percent in Seawater	Percent in Blood
Na^+	2.36	0.322
Cl^-	1.94	0.366
Mg^{2+}	0.13	0.002
SO_4^{2-}	0.09	—
K^+	0.04	0.016
Ca^{2+}	0.04	0.0096
HCO_3^-	0.002	0.165
HPO_4^{2-}, $H_2PO_4^-$	—	0.01

Most ions are more abundant in seawater than they are in blood, with some important exceptions. There are far more hydrogen carbonate ions ($\mathbf{HCO_3^-}$) in blood than in seawater. This difference is significant because the hydrogen carbonate ion and some related ions have a crucial role in controlling the acid-base properties of blood. The amount of hydrogen phosphate ions—$\mathbf{HPO_4^{2-}}$ and $\mathbf{H_2PO_4^-}$—in seawater is very low, but they are present in higher amounts in blood, where they also affect acid-base properties. Another notable difference is that blood does not have significant amounts of the sulfate ion ($\mathbf{SO_4^{2-}}$), but this ion is present in seawater.

Key Takeaways

- Proper chemical formulas for ionic compounds balance the total positive charge with the total negative charge.
- Groups of atoms with an overall charge, called polyatomic ions, also exist.

EXERCISES

1. What information is contained in the formula of an ionic compound?

- Why do the chemical formulas for some ionic compounds contain subscripts, while others do not?

3. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Mg^{2+} and I^-
   b. Na^+ and O^{2-}

4. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Na^+ and Br^-
   b. Mg^{2+} and Br^-
   c. Mg^{2+} and S^{2-}

5. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. K^+ and Cl^-
   b. Mg^{2+} and Cl^-
   c. Mg^{2+} and Se^{2-}

6. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Na^+ and N^{3-}
   b. Mg^{2+} and N^{3-}
   c. Al^{3+} and S^{2-}

7. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Li^+ and N^{3-}
   b. Mg^{2+} and P^{3-}

c. Li^{+} and P^{3-}

8. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Fe^{3+} and Br^{-}
b. Fe^{2+} and Br^{-}
c. Au^{3+} and S^{2-}
d. Au^{+} and S^{2-}

9. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Cr^{3+} and O^{2-}
b. Cr^{2+} and O^{2-}
c. Pb^{2+} and Cl^{-}
d. Pb^{4+} and Cl^{-}

10. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Cr^{3+} and NO_3^{-}
b. Fe^{2+} and PO_4^{3-}
c. Ca^{2+} and CrO_4^{2-}
d. Al^{3+} and OH^{-}

11. Write the chemical formula for the ionic compound formed by each pair of ions.

a. NH_4^{+} and NO_3^{-}
b. H^{+} and $Cr_2O_7^{2-}$
c. Cu^{+} and CO_3^{2-}
d. Na^{+} and HCO_3^{-}

12. For each pair of elements, determine the charge for their ions and write the proper formula for the resulting ionic compound between them.

a. Ba and S
b. Cs and I

13. For each pair of elements, determine the charge for their ions and write the proper formula for the resulting ionic compound between them.

a. K and S
b. Sc and Br

14. Which compounds would you predict to be ionic?

a. Li_2O
b. $(NH_4)_2O$
c. CO_2
d. $FeSO_3$
e. C_6H_6
f. C_2H_6O

15. Which compounds would you predict to be ionic?

a. $Ba(OH)_2$
b. CH_2O
c. NH_2CONH_2
d. $(NH_4)_2CrO_4$
e. C_8H_{18}
f. NH_3

Answers

1. the ratio of each kind of ion in the compound

2. Sometimes more than one ion is needed to balance the charge on the other ion in an ionic compound.

3.
   a. MgI_2
   b. Na_2O

4.
   a. NaBr
   b. $MgBr_2$
   c. MgS

5.
   a. KCl
   b. $MgCl_2$
   c. MgSe

6.
   a. Na_3N
   b. Mg_3N_2
   c. Al_2S_3

7.
   a. Li_3N
   b. Mg_3P_2
   c. Li_3P

8.
   a. $FeBr_3$
   b. $FeBr_2$
   c. Au_2S_3
   d. Au_2S

9.
   a. Cr_2O_3
   b. CrO
   c. $PbCl_2$
   d. $PbCl_4$

10.
   a. $Cr(NO_3)_3$
   b. $Fe_3(PO_4)_2$
   c. $CaCrO_4$
   d. $Al(OH)_3$

11.
   a. NH_4NO_3
   b. $H_2Cr_2O_7$
   c. Cu_2CO_3
   d. $NaHCO_3$

12.

a. Ba^{2+}, S^{2-}, BaS
b. Cs^{+}, I^{-}, CsI

13.

a. K^{+}, S^{2-}, K_2S
b. Sc^{3+}, Br^{-}, $ScBr_3$

14.

a. ionic
b. ionic
c. not ionic
d. ionic
e. not ionic
f. not ionic

15.

a. ionic
b. not ionic
c. not ionic
d. ionic
e. not ionic
f. not ionic

4.3: Formulas for Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **3.3: Formulas for Ionic Compounds** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.4: Ionic Nomenclature

Learning Objectives

- To use the rules for naming ionic compounds

After learning a few more details about the names of individual ions, you will be a step away from knowing how to name ionic compounds. This section begins the formal study of nomenclature, the systematic naming of chemical compounds.

Naming Ions

The name of a monatomic cation is simply the name of the element followed by the word *ion*. Thus, Na^+ is the sodium ion, Al^{3+} is the aluminum ion, Ca^{2+} is the calcium ion, and so forth.

We have seen that some elements lose different numbers of electrons, producing ions of different charges. Iron, for example, can form two cations, each of which, when combined with the same anion, makes a different compound with unique physical and chemical properties. Thus, we need a different name for each iron ion to distinguish Fe^{2+} from Fe^{3+}. The same issue arises for other ions with more than one possible charge.

There are two ways to make this distinction. In the simpler, more modern approach, called the ***Stock system***, an ion's positive charge is indicated by a roman numeral in parentheses after the element name, followed by the word *ion*. Thus, Fe^{2+} is called the iron(II) ion, while Fe^{3+} is called the iron(III) ion. This system is used only for elements that form more than one common positive ion. We do not call the Na^+ ion the sodium(I) ion because (I) is unnecessary. Sodium forms only a 1+ ion, so there is no ambiguity about the name *sodium ion*.

Table **4.4.1**: The Common System of Cation Names

Element	Stem	Charge	Name
iron	ferr-	2+	ferrous ion
		3+	ferric ion
copper	cupr-	1+	cuprous ion
		2+	cupric ion
tin	stann-	2+	stannous ion
		4+	stannic ion
lead	plumb-	2+	plumbous ion
		4+	plumbic ion
chromium	chrom-	2+	chromous ion
		3+	chromic ion
gold	aur-	1+	aurous ion
		3+	auric ion

The second system, called the **common system**, is not conventional but is still prevalent and used in the health sciences. This system recognizes that many metals have two common cations. The common system uses two suffixes (*-ic* and *-ous*) that are appended to the stem of the element name. The *-ic* suffix represents the greater of the two cation charges, and the *-ous* suffix represents the lower one. In many cases, the stem of the element name comes from the Latin name of the element. Table **4.4.1** lists the elements that use the common system, along with their respective cation names.

The name of a monatomic anion consists of the stem of the element name, the suffix *-ide*, and then the word *ion*. Thus, as we have already seen, Cl^- is "chlor-" + "-ide ion," or the chloride ion. Similarly, O^{2-} is the oxide ion, Se^{2-} is the selenide ion, and so forth. Table **4.4.2** lists the names of some common monatomic ions.

Table **4.4.2**: Some Monatomic Anions

Ion	Name
F^-	fluoride ion
Cl^-	chloride ion
Br^-	bromide ion
I^-	iodide ion
O^{2-}	oxide ion
S^{2-}	sulfide ion
P^{3-}	phosphide ion
N^{3-}	nitride ion

The polyatomic ions have their own characteristic names, as discussed earlier.

Example 4.4.1

Name each ion.

a. Ca^{2+}
b. S^{2-}
c. SO_3^{2-}
d. NH_4^+
e. Cu^+

Answer a

the calcium ion

Answer b

the sulfide ion (from Table **4.4.2**)

Answer c

the sulfite ion

Answer d

the ammonium ion

Answer e

the copper(I) ion or the cuprous ion (copper can form cations with either a 1+ or 2+ charge, so we have to specify which charge this ion has

Exercise 4.4.1

Name each ion.

a. Fe^{2+}
b. Fe^{3+}
c. SO_4^{2-}

d. Ba^{2+}
e. HCO_3^-

Answer a

the iron (II) or ferrous ion

Answer b

the iron (III) or ferric ion

Answer c

the sulfate ion

Answer d

the barium ion

Answer e

the bicarbonate ion or hydrogen carbonate ion

Example 4.4.2

Write the formula for each ion.

a. the bromide ion
b. the phosphate ion
c. the cupric ion
d. the magnesium ion

Answer a

Br^-

Answer b

PO_4^{3-}

Answer c

Cu^{2+}

Answer d

Mg^{2+}

Exercise 4.4.2

Write the formula for each ion.

a. the fluoride ion
b. the carbonate ion
c. the ferrous ion
d. the potassium ion

Answer a

F^-

Answer b

CO_3^{2-}

Answer c


Fe^{2+}

Answer d

K^+

Naming Compounds

Now that we know how to name ions, we are ready to name ionic compounds. We do so by placing the name of the cation first, followed by the name of the anion, and dropping the word *ion* from both parts. For example, what is the name of the compound whose formula is $\mathbf{Ba(NO_3)_2}$?



The compound's name does not indicate that there are two nitrate ions for every barium ion. You must determine the relative numbers of ions by balancing the positive and negative charges.

If you are given a formula for an ionic compound whose cation can have more than one possible charge, you must first determine the charge on the cation before identifying its correct name. For example, consider $\mathbf{FeCl_2}$ and $\mathbf{FeCl_3}$. In the first compound, the iron ion has a 2+ charge because there are two $\mathbf{Cl^-}$ ions in the formula (1− charge on each chloride ion). In the second compound, the iron ion has a 3+ charge, as indicated by the three $\mathbf{Cl^-}$ ions in the formula. These are two different compounds that need two different names. By the Stock system, the names are iron(II) chloride and iron(III) chloride. If we were to use the stems and suffixes of the common system, the names would be ferrous chloride and ferric chloride, respectively.

✔ Example 4.4.3

Name each ionic compound, using both Stock and common systems if necessary.

a. $Ca_3(PO_4)_2$
b. $(NH_4)_2Cr_2O_7$
c. KCl
d. CuCl
e. SnF_2

Answer a

calcium phosphate

Answer b

ammonium dichromate (the prefix *di-* is part of the name of the anion)

Answer c

potassium chloride

Answer d

copper(I) chloride or cuprous chloride

Answer e

tin(II) fluoride or stannous fluoride

Exercise 4.4.3

Name each ionic compound, using both Stock and common systems if necessary.

a. $ZnBr_2$
b. $Fe(NO_3)_3$
c. Al_2O_3
d. CuF_2
e. AgF

Answer a

zinc bromide

Answer b

iron (III) nitrate or ferric nitrate

Answer c

aluminum oxide

Answer d

copper (II) fluoride or cupric fluoride

Answer e

silver fluoride

Figure 4.4.1 is a synopsis of how to name simple ionic compounds.




Figure 4.4.1: A Guide to Naming Simple Ionic Compounds. Follow these steps to name a simple ionic compound.

Identify the cation name and the anion name. If the cation can have more than one possible charge, either use the Stock system name of the cation and name of the anion, or use the stem of the cation name and -ic/-ous and the name of the anion. Examples of this would be $FeCl_2$, which is iron(II) chloride of ferrous chloride, $CuSO_4$, which is copper(II) sulfate or cupric sulfate, and Cr_2O_3, which is chromium(III) oxide or chromic oxide. If the cation can not have more than one possible charge, use the name of the cation and the name of the anion. Examples of this would be KBr, which is potassium bromide, $NaNO_3$, which is sodium nitrate, and $(NH_4)_2S$, which is ammonium sulfide.

KEY TAKEAWAY

- Each ionic compound has its own unique name that comes from the names of the ions.

EXERCISES

1. Briefly describe the process for naming an ionic compound.

- In what order do the names of ions appear in the names of ionic compounds?

3. Which ionic compounds can be named using two different systems? Give an example.

4. Name each ion.

   a. Ra^{2+}
   b. P^{3-}
   c. $H_2PO_4^-$
   d. Sn^{4+}

5. Name each ion.

   a. Cs^+
   b. As^{3-}
   c. HSO_4^-
   d. Sn^{2+}

6. Name the ionic compound formed by each pair of ions.

   a. Na^+ and Br^-
   b. Mg^{2+} and Br^-
   c. Mg^{2+} and S^{2-}

7. Name the ionic compound formed by each pair of ions.

   a. K^+ and Cl^-
   b. Mg^{2+} and Cl^-
   c. Mg^{2+} and Se^{2-}

8. Name the ionic compound formed by each pair of ions.

   a. Na^+ and N^{3-}
   b. Mg^{2+} and N^{3-}
   c. Al^{3+} and S^{2-}

9. Name the ionic compound formed by each pair of ions.

   a. Li^+ and N^{3-}
   b. Mg^{2+} and P^{3-}
   c. Li^+ and P^{3-}

10. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

    a. Fe^{3+} and Br^-
    b. Fe^{2+} and Br^-
    c. Au^{3+} and S^{2-}
    d. Au^+ and S^{2-}

11. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

    a. Cr^{3+} and O^{2-}
    b. Cr^{2+} and O^{2-}
    c. Pb^{2+} and Cl^-
    d. Pb^{4+} and Cl^-

12. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

    a. Cr^{3+} and NO_3^-

b. Fe^{2+} and PO_4^{3-}
c. Ca^{2+} and CrO_4^{2-}
d. Al^{3+} and OH^-

13. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

a. NH_4^+ and NO_3^-
b. K^+ and $Cr_2O_7^{2-}$
c. Cu^+ and CO_3^{2-}
d. Na^+ and HCO_3^-

14. Give two names for each compound.

a. $Al(HSO_4)_3$
b. $Mg(HSO_4)_2$

15. Give two names for each compound.

a. $Co(HCO_3)_2$
b. $LiHCO_3$

AnswerS

1. Name the cation and then the anion but don’t use numerical prefixes.
2. the cation name followed by the anion name
3. Ionic compounds in which the cation can have more than one possible charge have two naming systems. $FeCl_3$ is either iron(III) chloride or ferric chloride (answers will vary).

4.

a. the radium ion
b. the phosphide ion
c. the dihydrogen phosphate ion
d. the tin(IV) ion or the stannic ion

5.

a. the cesium ion
b. the arsenide ion
c. the hydrogen sulfate ion
d. the tin(II) ion or the stannous ion

6.

a. sodium bromide
b. magnesium bromide
c. magnesium sulfide

7.

a. potassium chloride
b. magnesium chloride
c. magnesium selenide

8.

a. sodium nitride
b. magnesium nitride
c. aluminum sulfide

9.

a. lithium nitride
b. magnesium phosphide
c. lithium phosphide

10.

a. iron(III) bromide or ferric bromide
b. iron(II) bromide or ferrous bromide
c. gold(III) sulfide or auric sulfide
d. gold(I) sulfide or aurous sulfide

11.

a. chromium(III) oxide or chromic oxide
b. chromium(II) oxide or chromous oxide
c. lead(II) chloride or plumbous chloride
d. lead(IV) chloride or plumbic chloride

12.

a. chromium(III) nitrate or chromic nitrate
b. iron(II) phosphate or ferrous phosphate
c. calcium chromate
d. aluminum hydroxide

13.

a. ammonium nitrate
b. potassium dichromate
c. copper(I) carbonate or cuprous carbonate
d. sodium hydrogen carbonate or sodium bicarbonate

14.

a. aluminum hydrogen sulfate or aluminum bisulfate
b. magnesium hydrogen sulfate or magnesium bisulfate

15.

a. cobalt hydrogen carbonate or cobalt bicarbonate
b. lithium hydrogen carbonate or lithium bicarbonate

4.4: Ionic Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **3.4: Ionic Nomenclature** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.5: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms and ask yourself how they relate to the topics in the chapter.

Atoms combine into compounds by forming **chemical bonds**. A survey of stable atoms and molecules leads to the **octet rule**, which says that stable atoms tend to have eight electrons in their outermost, or valence, shell. One way atoms obtain eight electrons in the valence shell is for some atoms to lose electrons while other atoms gain them. When this happens, the atoms take on an electrical charge. Charged atoms are called **ions**. Ions having opposite charges attract each other. This attraction is called **ionic bonding**, and the compounds formed are called **ionic compounds**.

Positively charged ions are called **cations**, while negatively charged ions are called **anions**. The formation of both cations and anions can be illustrated using electron configurations. Because elements in a column of the periodic table have the same valence shell electron configuration, atoms in the same column of the periodic table tend to form ions having the same charge. **Electron dot diagrams**, or **Lewis diagrams**, can also be used to illustrate the formation of cations and anions.

Ionic compounds are represented in writing by a **chemical formula**, which gives the lowest ratio of cations and anions present in the compound. In a formula, the symbol of the cation is written first, followed by the symbol of the anion. **Formula unit** is considered the basic unit of an ionic compound because ionic compounds do not exist as discrete units. Instead, they exist as **crystals**, three-dimensional arrays of ions, with cations surrounded by anions and anions surrounded by cations. Chemical formulas for ionic compounds are determined by balancing the positive charge from the cation(s) with the negative charge from the anion(s). A subscript to the right of the ion indicates that more than one of that ion is present in the chemical formula.

Some ions are groups of atoms bonded together and having an overall electrical charge. These are called **polyatomic ions**. Writing formulas with polyatomic ions follows the same rules as with monatomic ions, except that when more than one polyatomic ion is present in a chemical formula, the polyatomic ion is enclosed in parentheses and the subscript is outside the right parenthesis. Ionic compounds typically form between metals and nonmetals or between polyatomic ions.

Names of ionic compounds are derived from the names of the ions, with the name of the cation coming first, followed by the name of the anion. If an element can form cations of different charges, there are two alternate systems for indicating the compound's name. In the **Stock system**, a roman numeral in parentheses indicates the charge on the cation. An example is the name for $FeCl_2$, which is iron(II) chloride. In the common system, the suffixes *-ous* and *-ic* are used to stand for the lower and higher possible charge of the cation, respectively. These suffixes are attached to a stem representing the element (which frequently comes from the Latin form of the element name). An example is the common name for $FeCl_2$, which is ferrous chloride.

Template:HideTOC

4.5: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

5: Covalent Bonding and Simple Molecular Compounds

Ionic bonding results from the transfer of electrons among atoms or groups of atoms. In this chapter, we will consider another type of bonding—covalent bonding. We will examine how atoms share electrons to form these bonds, and we will begin to explore how the resulting compounds, such as cholesterol, are different from ionic compounds.

5.1: Covalent Bonds
5.2: Covalent Compounds- Formulas and Names
5.3: Multiple Covalent Bonds
5.4: Characteristics of Covalent Bonds
5.5: Characteristics of Molecules
5.6: Intermolecular Forces
5.7: Chapter Summary

5: Covalent Bonding and Simple Molecular Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

5.1: Covalent Bonds

Learning Objectives


- To describe how a covalent bond forms.
- To apply the octet rule to covalent compounds

You have already seen examples of substances that contain covalent bonds. One substance mentioned previously was water (H_2O). You can tell from its formula that it is not an ionic compound; it is not composed of a metal and a nonmetal. Consequently, its properties are different from those of ionic compounds.


Electron Sharing

Previously, we discussed ionic bonding where electrons can be transferred from one atom to another so that both atoms have an energy-stable outer electron shell. Because most filled electron shells have eight electrons in them, chemists called this tendency the octet rule. However, there is another way an atom can achieve a full valence shell: atoms can *share* electrons.

This concept can be illustrated by using two hydrogen atoms, each of which has a single electron in its valence shell. (For small atoms such as hydrogen atoms, the valence shell will be the first shell, which holds only two electrons.) We can represent the two individual hydrogen atoms as follows:



In contrast, when two hydrogen atoms get close enough together to share their electrons, they can be represented as follows:



By sharing their valence electrons, both hydrogen atoms now have two electrons in their respective valence shells. Because each valence shell is now filled, this arrangement is more stable than when the two atoms are separate. The sharing of electrons between atoms is called a covalent bond, and the two electrons that join atoms in a covalent bond are called a bonding pair of electrons. A discrete group of atoms connected by covalent bonds is called a molecule—the smallest part of a compound that retains the chemical identity of that compound.

Chemists frequently use Lewis diagrams to represent covalent bonding in molecular substances. For example, the Lewis diagrams of two separate hydrogen atoms are as follows:



The Lewis diagram of two hydrogen atoms sharing electrons looks like this:



This depiction of molecules is simplified further by using a dash to represent a covalent bond. The hydrogen molecule is then represented as follows:



Remember that the dash, also referred to as a single bond, represents a *pair* of electrons.

The bond in a hydrogen molecule, measured as the distance between the two nuclei, is about 7.4×10^{-11} m, or 74 picometers (pm; 1 pm = 1×10^{-12} m). This particular bond length represents a balance between several forces: the attractions between oppositely charged electrons and nuclei, the repulsion between two negatively charged electrons, and the repulsion between two positively charged nuclei. If the nuclei were closer together, they would repel each other more strongly; if the nuclei were farther apart, there would be less attraction between the positive and negative particles.

Fluorine is another element whose atoms bond together in pairs to form *diatomic* (two-atom) molecules. Two separate fluorine atoms have the following electron dot diagrams:




Each fluorine atom contributes one valence electron, making a single bond and giving each atom a complete valence shell, which fulfills the **octet** rule:



Two F's with two dots in between them, and two dots on the top, left/right side, and bottoms of each F.

The circles show that each fluorine atom has eight electrons around it. As with hydrogen, we can represent the fluorine molecule with a dash in place of the bonding electrons:



Two F's with a long dash in between them, and two dots on the top, left/right side, and bottoms of each F.


Each fluorine atom has six electrons, or three pairs of electrons, that are not participating in the covalent bond. Rather than being shared, they are considered to belong to a single atom. These are called nonbonding pairs (or lone pairs) of electrons.

Covalent Bonds between Different Atoms

Now that we have looked at electron sharing between atoms of the same element, let us look at covalent bond formation between atoms of different elements. Consider a molecule composed of one hydrogen atom and one fluorine atom:



Each atom needs one additional electron to complete its valence shell. By each contributing one electron, they make the following molecule:



An H and F connected by two dots, and the F has 2 dots on the top, right, and bottom. It also shows the same combination but with a long dash in between the H and F rather than two dots.

In this molecule, the hydrogen atom does not have nonbonding electrons, while the fluorine atom has six nonbonding electrons (three lone electron pairs). The circles show how the valence electron shells are filled for both atoms.

Example 5.1.1


Use Lewis diagrams to indicate the formation of the following:

a. Cl_2
b. HBr

Solution


a. When two chlorine atoms form a chlorine molecule, they share one pair of electrons. In Cl_2 molecule, each chlorine atom is surrounded by an **octet** number of electrons.

The Lewis diagram for a Cl_2 molecule is similar to the one for F_2 (shown above).



Cl with 7 dots surrounding it plus Cl with 7 dots surrounding it means they are chlorine atoms. To turn into a chlorine molecule, have two dots in between each Cl and 6 dots surrounding each Cl on the sides that aren't connected.

b. When a hydrogen atom and a bromine atom form HBr, they share one pair of electrons. In the HBr molecule, H achieves a full valence of two electrons (**duet**) while Br achieves an **octet**. The Lewis diagram for HBr is similar to that for HF shown above.



H with one dot plus Br with 7 dots turns into H connected to Br by two dots, with 6 other dots surrounding Br.

Exercise 5.1.1


Draw the Lewis diagram for each compound.

a. a molecule composed of one chlorine atom and one fluorine atom
b. a molecule composed of one hydrogen atom and one iodine atom

Answer a:




Answer b:




Covalent Bonds in Larger Molecules

The formation of a water molecule from two hydrogen atoms and an oxygen atom can be illustrated using Lewis dot symbols (shown below).



H with one dot + O with 6 dots + H with one dot turns into H connected to O with two dots connected to H with two dots. O also has two dots on top and top dots on bottom.


The structure on the right is the *Lewis electron structure*, or *Lewis structure*, for H_2O. With two bonding pairs and two lone pairs, the oxygen atom has now completed its **octet**. Moreover, by sharing a bonding pair with oxygen, each hydrogen atom now has a full valence shell of two electrons. Chemists usually indicate a bonding pair by a single line, as shown (below).



Water


Other large molecules are constructed in a similar fashion, with some atoms participating in more than one covalent bond. For example, methane (CH_4), the central carbon atom bonded to four hydrogen atoms, can be represented using either of the Lewis

structures below. Again, sharing electrons between C and H atoms results in C achieving and octet while H achieving a duet number of electrons.



How Many Covalent Bonds Are Formed?

The number of bonds that an atom can form can often be predicted from the number of electrons needed to reach an octet (eight valence electrons). In the Lewis structure, the number of bonds formed by an element in a neutral compound is the same as the number of unpaired electrons it must share with other atoms to complete its octet of electrons. For example, each atom of a **group 4A** (14) element has four electrons in its outermost shell and therefore requires four more electrons to reach an octet. These four electrons can be gained by forming **four covalent bonds**, as illustrated here for carbon in CH_4 (methane). **Group 5A** (15) elements such as nitrogen have five valence electrons in the atomic Lewis symbol: one lone pair and three unpaired electrons. To obtain an octet, these atoms form **three covalent bonds**, as in NH_3 (ammonia). Oxygen and other atoms in **group 6A** (16) obtain an octet by forming **two covalent bonds**. Fluorine and the other halogens in **group 7A** (17) have seven valence electrons and can obtain an octet by forming **one covalent bond**.



Typically, the atoms of group 4A form 4 covalent bonds; group 5A form 3 bonds; group 6A form 2 bonds; and group 7A form one bond. The number of electrons required to obtain an octet determines the number of covalent bonds an atom can form. This is summarized in the table below. In each case, the sum of the number of bonds and the number of lone pairs is 4, which is equivalent to eight (octet) electrons.

This table shows atoms and their group numbers, and how many bonds and lone pairs each has.

Atom (Group number)	Number of Bonds	Number of Lone Pairs
Carbon (Group 14 or 4A)	4	0
Nitrogen (Group 15 or 5A)	3	1
Oxygen (Group 16 or 6A)	2	2
Fluorine (Group 17 or 7A)	1	3

Because hydrogen only needs two electrons to fill its valence shell, it follows the duet rule. It is an **exception to the octet rule.** Hydrogen only needs to form one bond. This is the reason why H is always a terminal atom and never a central atom. Figure **5.1.1** shows the number of covalent bonds various atoms typically form.

The transition elements and inner transition elements also do not follow the octet rule since they have d and f electrons involved in their valence shells.





Figure **5.1.1**: How Many Covalent Bonds Are Formed? In molecules, there is a pattern to the number of covalent bonds that different atoms can form. Each block with a number indicates the number of covalent bonds formed by that atom in neutral compounds.

Example 5.1.2

Examine the Lewis structure of OF_2 below. Count the number of bonds formed by each element. Based on the element's location in the periodic table, does it correspond to the expected number of bonds shown in Table 4.1? Does the Lewis structure below follow the octet rule?




F surrounded by 6 dots, connected to O with a long dash. O surrounded by 4 dots and connects to another F with a long dash. F surrounded by 6 dots.

Solution

Yes. F (group 7A) forms one bond and O (group 6A) forms 2 bonds. Each atom is surrounded by 8 electrons. This structure satisfies the octet rule.

5.1.2 Exercise

Examine the Lewis structure of NCl_3 below. Count the number of bonds formed by each element. Based on the element's location in the periodic table, does it correspond to the expected number of bonds shown in Table 4.1? Does the Lewis structure below follow the octet rule?



Answer

Both Cl and N form the expected number of bonds. Cl (group 7A) has one bond and 3 lone pairs. The central atom N (group 5A) has 3 bonds and one lone pair. Yes, the Lewis structure of NCl_3 follows the octet rule.

Key Takeaways

- A covalent bond is formed between two atoms by sharing electrons.
- The number of bonds an element forms in a covalent compound is determined by the number of electrons it needs to reach octet.
- Hydrogen is an exception to the octet rule. H forms only one bond because it needs only two electrons.

5.1: Covalent Bonds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.1: Covalent Bonds** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.2: Covalent Compounds- Formulas and Names

Learning Objectives


- Identify covalent and ionic compounds.
- Determine the chemical formula of a simple covalent compound from its name.
- Determine the name of a simple covalent compound from its chemical formula.

COVALENT AND IONIC COMPOUNDS


What elements make covalent bonds? Covalent bonds form when two or more nonmetals combine. For example, both hydrogen and oxygen are nonmetals, and when they combine to make water, they do so by forming covalent bonds. Compounds that are composed of only non-metals or semi-metals with non-metals will display covalent bonding and will be classified as molecular compounds.

As a general rule of thumb, compounds that involve a metal binding with either a non-metal or a semi-metal will display ionic bonding. Thus, the compound formed from sodium and chlorine will be ionic (a metal and a non-metal). Nitrogen monoxide (NO) will be a covalently bound molecule (two non-metals), silicon dioxide (SiO_2) will be a covalently bound molecule (a semi-metal and a non-metal) and $MgCl_2$ will be ionic (a metal and a non-metal).

A polyatomic ion is an ion composed of two or more atoms that have a charge as a group (poly = many). The ammonium ion (see figure below) consists of one nitrogen atom and four hydrogen atoms. Together, they comprise a single ion with a 1+ charge and a formula of NH_4^+. The carbonate ion (see figure below) consists of one carbon atom and three oxygen atoms and carries an overall charge of 2−. The formula of the carbonate ion is CO_3^{2-}.



ammonium ion



carbonate ion

The atoms of a polyatomic ion are tightly bonded together and so the entire ion behaves as a single unit. Several examples are found in Table 3.3.1. Nonmetal atoms in polyatomic ions are joined by covalent bonds, but the ion as a whole participates in ionic bonding. For example, ammonium chloride (NH_4Cl) has ionic bonding between a polyatomic ion, $\mathbf{NH_4^+}$, and $\mathbf{Cl^-}$ ions, but within the ammonium ion (NH_4^+), the nitrogen and hydrogen atoms are connected by covalent bonds (shown above).

Both ionic and covalent bonding are also found in calcium carbonate. Calcium carbonate ($CaCO_3$) has ionic bonding between calcium ion $\mathbf{Ca^{2+}}$ and a polyatomic ion, $\mathbf{CO_3^{2-}}$, but within the carbonate ion (CO_3^{2-}), the carbon and oxygen atoms are connected by covalent bonds (shown above).

Characteristics of Covalent (Molecular) Compounds

Compounds that contain covalent bonds (also called molecular compounds) exhibit different physical properties than ionic compounds. Because the attraction between molecules, which are electrically neutral, is weaker than that between electrically charged ions, covalent compounds generally have much lower melting and boiling points than ionic compounds (discussed in Section 3.6). For example, water (molecular compound) boils at 100 °C while sodium chloride (ionic compound) boils at 1413 °C. In fact, many covalent compounds are liquids or gases at room temperature, and, in their solid states, they are typically much softer than ionic solids. Furthermore, whereas ionic compounds are good conductors of electricity when dissolved in water, most covalent compounds, being electrically neutral, are poor conductors of electricity in any state. The attraction between molecules (called intermolecular forces) will be discussed in more detail in Section 8.1

Example 5.2.1

Is each compound formed from ionic bonds, covalent bonds, or both?

a. Na_2O
b. Na_3PO_4
c. N_2O_4

Answer a

The elements in Na_2O are a metal and a nonmetal, which form ionic bonds.

Answer b

Because sodium is a metal and we recognize the formula for the phosphate ion, we know that this compound is ionic. However, within the polyatomic phosphate ion, the atoms are held together by covalent bonds, so this compound contains both ionic and covalent bonds.

Answer c

The elements in N_2O_4| are both nonmetals, rather than a metal and a nonmetal. Therefore, the atoms form covalent bonds.

Exercise 5.2.1

Is each compound are formed from ionic bonds, covalent bonds, or both?

a. $Ba(OH)_2$
b. F_2
c. PCl_3

Answer a:

both

Answer b:


covalent

Answer c:

covalent

MOLECULAR FORMULAS

The chemical formulas for covalent compounds are referred to as **molecular formulas** because these compounds exist as separate, discrete molecules. Typically, a molecular formula begins with the nonmetal that is closest to the lower left corner of the periodic table, except that hydrogen is almost never written first (H_2O is the prominent exception). Then the other nonmetal symbols are listed. Numerical subscripts are used if there is more than one of a particular atom. For example, we have already seen CH_4, the molecular formula for methane. Below is the molecular formula of ammonia, NH_3.



NH_3. An arrow points to the N and says, "The lack of a subscript on the N symbol indicates that there is 1 atom of nitrogen in a molecule of ammonia". An arrow points to H and says, "The subscript below the H symbol indicates that there are 3 hydrogen atoms in a molecule of ammonia".

NAMING COVALENT COMPOUNDS

Naming *binary* (two-element) covalent compounds is similar to naming simple ionic compounds. The first element in the formula is simply listed using the name of the element. The second element is named by taking the stem of the element name and adding

the suffix *-ide*. A system of numerical prefixes is used to specify the number of atoms in a molecule. Table **5.2.1** lists these numerical prefixes. Normally, no prefix is added to the first element's name if there is only one atom of the first element in a molecule. If the second element is oxygen, the trailing vowel is usually omitted from the end of a polysyllabic prefix but not a monosyllabic one (that is, we would say "monoxide" rather than "monooxide" and "trioxide" rather than "troxide").

Table **5.2.1**: Numerical Prefixes for Naming Binary Covalent Compounds

Number of Atoms in Compound	Prefix on the Name of the Element
1	mono-*
2	di-
3	tri-
4	tetra-
5	penta-
6	hexa-
7	hepta-
8	octa-
9	nona-
10	deca-

*This prefix is not used for the first element's name.

Let us practice by naming the compound whose molecular formula is CCl_4. The name begins with the name of the first element—carbon. The second element, chlor*ine*, becomes chlor*ide*, and we attach the correct numerical prefix ("tetra-") to indicate that the molecule contains four chlorine atoms. Putting these pieces together gives the name *carbon tetrachloride* for this compound.

Example 5.2.2

Write the molecular formula for each compound.

a. chlorine trifluoride
b. phosphorus pentachloride
c. sulfur dioxide
d. dinitrogen pentoxide

Solution

If there is no numerical prefix on the first element's name, we can assume that there is only one atom of that element in a molecule.

a. ClF_3
b. PCl_5
c. SO_2
d. N_2O_5 (The *di-* prefix on nitrogen indicates that two nitrogen atoms are present.)

Exercise 5.2.2

Write the molecular formula for each compound.

a. nitrogen dioxide
b. dioxygen difluoride
c. sulfur hexafluoride

d. selenium monoxide

Answer a:

a. NO_2

Answer b:

O_2F_2

Answer c:

SF_6

Answer d:

SeO

Because it is so unreactive, sulfur hexafluoride is used as a spark suppressant in electrical devices such as transformers.

Example 5.2.3

Write the name for each compound.

a. BrF_5
b. S_2F_2
c. CO

Solution

a. bromine pentafluoride
b. disulfur difluoride
c. carbon monoxide

Exercise 5.2.3

Write the name for each compound.

a. CF_4
b. $SeCl_2$
c. SO_3

Answer a:

carbon tetrafluoride

Answer b:

selenium dichloride

Answer c:

sulfur trioxide

For some simple covalent compounds, we use common names rather than systematic names. We have already encountered these compounds, but we list them here explicitly:

- H_2O: water
- NH_3: ammonia
- CH_4: methane

Methane is the simplest organic compound. Organic compounds are compounds with carbon atoms and are named by a separate nomenclature system that we will introduce in in a separate section.

Key Takeaways

- The chemical formula of a simple covalent compound can be determined from its name.
- The name of a simple covalent compound can be determined from its chemical formula.

5.2: Covalent Compounds- Formulas and Names is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.2: Covalent Compounds - Formulas and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


5.3: Multiple Covalent Bonds

Learning Objectives

- To draw Lewis structures.
- To recognize molecules that are likely to have multiple covalent bonds.

DRAWING LEWIS STRUCTURES

For very simple molecules and molecular ions, we can write the Lewis structures by merely pairing up the unpaired electrons on the constituent atoms. See these examples:



For more complicated molecules and molecular ions, it is helpful to follow the step-by-step procedure outlined here:

1. Determine the total number of valence (outer shell) electrons among all the atoms. For cations, subtract one electron for each positive charge. For anions, add one electron for each negative charge.
2. Draw a skeleton structure of the molecule or ion, arranging the atoms around a central atom. (Generally, the least electronegative element should be placed in the center.) Connect each atom to the central atom with a single bond (one electron pair).
3. Distribute the remaining electrons as lone pairs on the terminal atoms (except hydrogen), completing an octet around each atom.
4. Place all remaining electrons on the central atom.
5. Rearrange the electrons of the outer atoms to make multiple bonds with the central atom in order to obtain octets wherever possible.

Let us determine the Lewis structures of OF_2 and HCN as examples in following this procedure:

1. Determine the total number of valence (outer shell) electrons in the molecule or ion. For a molecule, we add the number of valence electrons (use the main group number) on each atom in the molecule. This is the total number of electrons that must be used in the Lewis structure.

$$O + 2\,(F) = OF_2$$

$$6e^- + (2 \times 7e^-) = 20e^-$$

$$H + C + N = HCN$$

$$1e^- + 4e^- + 5e^- = 10e^-$$

2. Draw a skeleton structure of the molecule or ion, arranging the atoms around a central atom and connecting each atom to the central atom with a single (one electron pair) bond. Note that H and F can only form one bond, and are always on the periphery rather than the central atom.

F—O—F H—C—N

3. Distribute the remaining electrons as lone pairs on the terminal atoms (except hydrogen) to complete their valence shells with an octet of electrons.


- In OF_2, six electrons are placed on each F.

- In HCN, six electrons placed on N




4. Place all remaining electrons on the central atom.

- In OF_2, 4 electrons are placed on O.
- In HCN: no electrons remain (the total valence of $10e^-$ is reached) so nothing changes.



5. Rearrange the electrons of the outer atoms to make multiple bonds with the central atom in order to obtain octets wherever possible.

- In OF_2, each atom has an octet as drawn, so nothing changes.
- In HCN, form two more C–N bonds




H dash C dash N with 6 dots around the N. 4 of the dots are pointing to the dash connecting the C and the N. This turns the formula into H dash C and three lines connecting the C and N together, with two dots on the outside of the N.

Finally, check to see if the total number of valence electrons are present in the Lewis structure. And then, inspect if the H atom has 2 electrons surrounding it and if each of the main group atoms is surrounded by 8 electrons.

MULTIPLE BONDS


In many molecules, the octet rule would not be satisfied if each pair of bonded atoms shares only two electrons. Review HCN in Step 5 above. Another example is carbon dioxide (CO_2). CO_2 has a total valence of $4e^- + (2 \times 6e^-) = 16e^-$. Following steps 1 to 4, we draw the following:




This does not give the carbon atom a complete octet; only four electrons are in its valence shell. This arrangement of shared electrons is far from satisfactory.




In this case, more than one pair of electrons must be shared between two atoms for both atoms to have an octet. A second electron pair from each oxygen atom must be shared with the central carbon atom shown by the arrows above. A lone pair from each O must be converted into a bonding pair of electrons.



In this arrangement, the carbon atom shares four electrons (two pairs) with the oxygen atom on the left and four electrons with the oxygen atom on the right. There are now eight electrons around each atom. Two pairs of electrons shared between two atoms make a double bond between the atoms, which is represented by a double dash:



Some molecules contain triple bonds (like HCN, shown above). Triple bonds are covalent bonds in which *three* pairs of electrons are shared by two atoms. Another compound that has a triple bond is acetylene (C_2H_2), whose Lewis diagram is as follows:




Example 5.3.1

Draw the Lewis diagram for each molecule.


1. **N_2**
2. **CH_2O** (The carbon atom is the central atom.) One application of CH_2O, also called formaldehyde, is the preservation of biological specimens. Aqueous solutions of CH_2O are called formalin and have a sharp, characteristic (pungent) odor.

Solution

1. The total number of electrons is 2 x 5 = 10 electrons. The bond between the two nitrogen atoms is a triple bond. The Lewis diagram for $N_2$is as follows:



2. The total number of electrons is 4 x 2(1) + 6 = 12 electrons. In CH_2O, the central atom is surrounded by two different types of atoms. The Lewis diagram that fills each atom's valence electron shell is as follows:




Exercise 5.3.1


Draw the Lewis diagram for each molecule.

a. **O_2**
b. **C_2H_4**

Answer a:





Answer b:




Example 5.3.2

Which is the correct Lewis structure for N_2H_2?

A. 

B. 

C. 

Solution


Lewis structure A is the correct answer. It has a total of (2 x $5e^-$) + (2 x $1e^-$) = $12e^-$. Each of the N atoms satisfy the octet requirement and the H atoms follow the duet rule.


Structure B is electron deficient. It has only $10e^-$ instead of 12.


Structure C has 14 (2 extra) electrons. The N atoms do not satisfy the octet.

5.3.2 Exercise

Which is the correct Lewis structure for NOCl?

A. 

B. 

C. 

Answer

Structure A violates the octet rule; N is surrounded by only $6e^-$.

Structure B violates the octet rule; Cl has $10e^-$ around it. Furthermore, there are a total of $20e^-$ instead of $18e^-$.

Structure C is the correct structure. It has a total of $6e^-$ + $5e^-$ + $7e^-$ = $18e^-$. Each atom is surrounded by 8 electrons (octet rule).

Key Takeaways

- A Lewis structure shows the bonding and nonbonding electrons around individual atoms in a molecule.
- Some molecules must have multiple covalent bonds between atoms to satisfy the octet rule.
- A double bond contains four electrons and a triple bond contains six electrons.

5.3: Multiple Covalent Bonds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.3: Drawing Lewis Structures** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.4: Characteristics of Covalent Bonds

Learning Objectives

- Recognize bond characteristics of covalent compounds: bond length and bond polarity.
- Use electronegativity values to predict bond polarity.

Bond Length

We previously stated that the covalent bond in the hydrogen molecule (H_2) has a certain length (about 7.4×10^{-11} m). Other covalent bonds also have known bond lengths, which are dependent on both the identities of the atoms in the bond and whether the bonds are single, double, or triple bonds. Table **5.4.1** lists the approximate bond lengths for some single covalent bonds. The exact bond length may vary depending on the identity of the molecule but will be close to the value given in the table.

Table **5.4.1**: Approximate Bond Lengths of Some Single Bonds

Bond	Length ($\times 10^{-12}\ m$)
H–H	74
H–C	110
H–N	100
H–O	97
H–I	161
C–C	154
C–N	147
C–O	143
N–N	145
O–O	145

Table **5.4.2** compares the lengths of single covalent bonds with those of double and triple bonds between the same atoms. Without exception, as the number of covalent bonds between two atoms increases, the bond length decreases. With more electrons between the two nuclei, the nuclei can get closer together before the internuclear repulsion is strong enough to balance the attraction.

Table **5.4.2**: Comparison of Bond Lengths for Single and Multiple Bonds

Bond	Length ($\times 10^{-12}\,m$)
C–C	154
C=C	134
C≡C	120
C–N	147
C=N	128
C≡N	116
C–O	143
C=O	120
C≡O	113
N–N	145
N=N	123
N≡N	110
O–O	145
O=O	121

Electronegativity and Bond Polarity

Although we defined covalent bonding as electron sharing, the electrons in a covalent bond are not always shared equally by the two bonded atoms. Unless the bond connects two atoms of the same element, as in H_2, there will always be one atom that attracts the electrons in the bond more strongly than the other atom does, as in HCl, shown in Figure **5.4.1**. A covalent bond that has an equal sharing of electrons (Figure **5.4.1***a*) is called a **nonpolar covalent bond**. A covalent bond that has an unequal sharing of electrons, as in Figure **5.4.1***b*, is called a **polar covalent bond**.




Figure **5.4.1** Polar versus Nonpolar Covalent Bonds. (a) The electrons in the covalent bond are equally shared by both hydrogen atoms. This is a nonpolar covalent bond. (b) The chlorine atom attracts the electrons in the bond more than the hydrogen atom does, leading to an imbalance in the electron distribution. This is a polar covalent bond.

The distribution of electron density in a polar bond is uneven. It is greater around the atom that attracts the electrons more than the other. For example, the electrons in the H–Cl bond of a hydrogen chloride molecule spend more time near the chlorine atom than near the hydrogen atom. Note that the shaded area around Cl in Figure **5.4.1***b* is much larger than it is around H.

This imbalance in electron density results in a buildup of partial negative charge (designated as δ−) on one side of the bond (Cl) and a partial positive charge (designated δ+) on the other side of the bond (H). This is seen in Figure **5.4.2***a*. The separation of charge in a polar covalent bond results in an electric dipole (two poles), represented by the arrow in Figure **5.4.2***b*. The direction of the arrow is pointed toward the δ− end while the + tail of the arrow indicates the δ+ end of the bond.




Figure **5.4.2**: (a) Unequal sharing of the bonding pair of electrons between H and Cl leads to partial positive charge on the H atom and partial negative charge on the Cl. Symbols δ+ and δ– indicate the polarity of the H–Cl bond. (b) The dipole is represented by an arrow with a cross at the tail. The cross is near the δ+ end and the arrowhead coincides with the δ–.

Any covalent bond between atoms of different elements is a polar bond, but the degree of polarity varies widely. Some bonds between different elements are only minimally polar, while others are strongly polar. Ionic bonds can be considered the ultimate in polarity, with electrons being transferred rather than shared. To judge the relative polarity of a covalent bond, chemists use **electronegativity**, which is a relative measure of how strongly an atom attracts electrons when it forms a covalent bond. There are various numerical scales for rating electronegativity. Figure **5.4.3** shows one of the most popular—the Pauling scale.





Figure **5.4.3** The electronegativity values derived by Pauling follow predictable periodic trends with the higher electronegativities toward the upper right of the periodic table. Fluorine has the highest value (4.0).

Looking Closer: Linus Pauling

Arguably the most influential chemist of the 20th century, Linus Pauling (1901–94) is the only person to have won two individual (that is, unshared) Nobel Prizes. In the 1930s, Pauling used new mathematical theories to enunciate some fundamental principles of the chemical bond. His 1939 book *The Nature of the Chemical Bond* is one of the most significant books ever published in chemistry.


By 1935, Pauling's interest turned to biological molecules, and he was awarded the 1954 Nobel Prize in Chemistry for his work on protein structure. (He was very close to discovering the double helix structure of DNA when James Watson and James Crick announced their own discovery of its structure in 1953.) He was later awarded the 1962 Nobel Peace Prize for his efforts to ban the testing of nuclear weapons.



Linus Pauling was one of the most influential chemists of the 20th century.

In his later years, Pauling became convinced that large doses of vitamin C would prevent disease, including the common cold. Most clinical research failed to show a connection, but Pauling continued to take large doses daily. He died in 1994, having spent a lifetime establishing a scientific legacy that few will ever equal.

The polarity of a covalent bond can be judged by determining the difference in the electronegativities of the two atoms making the bond. The greater the difference in electronegativities, the greater the imbalance of electron sharing in the bond. Although there are no hard and fast rules, the general rule is if the difference in electronegativities is **less than about 0.4,** the bond is considered **nonpolar;** if the difference is **greater than 0.4,** the bond is considered **polar.** If the difference in electronegativities is large enough **(**generally **greater than about 1.8),** the resulting compound is considered **ionic** rather than covalent**.** An electronegativity difference of **zero,** of course, indicates a **nonpolar covalent bond.**



Bond Type	Electronegativity Difference
pure covalent	< 0.4
polar covalent	between 0.4 and 1.8
ionic	> 1.8

Figure **5.4.4**: As the electronegativity difference increases between two atoms, the bond becomes more ionic.

Example 5.4.1

Describe the electronegativity difference between each pair of atoms and the resulting polarity (or bond type).

a. C and H
b. H and H
c. Na and Cl
d. O and H

Solution

a. Carbon has an electronegativity of 2.5, while the value for hydrogen is 2.1. The difference is 0.4, which is rather small. The C–H bond is therefore considered nonpolar.
b. Both hydrogen atoms have the same electronegativity value—2.1. The difference is zero, so the bond is nonpolar.
c. Sodium's electronegativity is 0.9, while chlorine's is 3.0. The difference is 2.1, which is rather high, and so sodium and chlorine form an ionic compound.
d. With 2.1 for hydrogen and 3.5 for oxygen, the electronegativity difference is 1.4. We would expect a very polar bond. The sharing of electrons between O and H is unequal with the electrons more strongly drawn towards O.

? Exercise 5.4.1

Describe the electronegativity (EN) difference between each pair of atoms and the resulting polarity (or bond type).

a. C and O
b. K and Br
c. N and N
d. Cs and F

Answer a:

The EN difference is 1.0 , hence polar. The sharing of electrons between C and O is unequal with the electrons more strongly drawn towards O.

Answer b:

The EN difference is greater than 1.8, hence ionic.

Answer c:

Identical atoms have zero EN difference, hence nonpolar.

Answer d:

The EN difference is greater than 1.8, hence ionic.

Bond Polarity and Molecular Polarity

If there is only one bond in the molecule, the bond polarity determines the molecular polarity. Any diatomic molecule in which the two atoms are the same element must be a **nonpolar** molecule. A diatomic molecule that consists of a polar covalent bond, such as HF, is a **polar molecule**. A **polar molecule** *is a molecule in which one end of the molecule is slightly positive, while the other end is slightly negative*. The two electrically charged regions on either end of the molecule are called poles, similar to a magnet having a north and a south pole. Hence, a *molecule with two poles is called a* **dipole**. A simplified way to depict polar molecules like HF is pictured below (see figure below).




Figure 5.4.5: A molecular dipole results from the unequal distribution of electron density throughout a molecule.

When placed between oppositely charged plates, polar molecules orient themselves so that their positive ends are closer to the negative plate and their negative ends are closer to the positive plate (see Figure 4.4.6 below).




Figure 5.4.6: Polar molecules are randomly oriented in the absence of an applied electric field (top). In an electric field, the molecules orient themselves to maximize the attraction between opposite charges (bottom).

Experimental techniques involving electric fields can be used to determine if a certain substance is composed of polar molecules and to measure the degree of polarity.

For molecules with more than two atoms, the **molecular geometry** must also be taken into account when determining if the molecule is polar or nonpolar. (Figure **5.4.7**) is a comparison between carbon dioxide and water. Carbon dioxide (CO_2) is a **linear** molecule. The oxygen atoms are more electronegative than the carbon atom, so there are two individual dipoles pointing outward from the C atom to each O atom. However, since the dipoles are of equal strength and are oriented directly opposite each other, they cancel each other out, and the overall molecular polarity of CO_2 is zero. CO_2 is a **nonpolar** molecule. In H_2O, the orientation of the two O–H bonds is **bent**: one end of the molecule has a partial positive charge, and the other end has a partial negative charge. In short, the molecule itself is **polar**. The polarity of water has an enormous impact on its physical and chemical properties. For example, the boiling point of water (100°C) is high for such a small molecule due to the fact that polar molecules attract each other strongly. On the other hand, the nonpolar carbon dioxide becomes a gas at −77°C, almost 200° lower than the temperature at which water boils. More details will be presented in the next section.



Figure **5.4.7** Physical Properties and Polarity. The physical properties of water and carbon dioxide are affected by their polarities.

Key Takeaways

- Covalent bonds between different atoms have different bond lengths.
- Covalent bonds can be polar or nonpolar, depending on the electronegativity difference between the atoms involved.

5.4: Characteristics of Covalent Bonds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.4: Characteristics of Covalent Bonds** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.5: Characteristics of Molecules

Learning Objectives

- Determine the molecular mass of a molecule.
- Predict the general shape of a simple covalent molecule.
- Predict the polarity of the molecule.
- Compare the properties of ionic and molecular compounds.

Unlike the ions in ionic compounds, which are arranged in lattices called crystals, molecules of covalent compounds exist as discrete units with a characteristic mass and a certain three-dimensional shape.

Molecular Mass

The mass of a molecule—the molecular mass (sometimes called the molecular weight)—is simply the sum of the masses of its atoms. As with formula masses, it is important that you keep track of the number of atoms of each element in the molecular formula to obtain the correct molecular mass.

Example 5.5.1

What is the molecular mass of each covalent compound?

a. H_2O
b. C_6H_6
c. NO_2
d. N_2O_5

Solution

Use the atomic masses from the Periodic Table in Section 2.7.

The molecular formula H_2O indicates that there are two hydrogen atoms and one oxygen atom in each molecule. Summing the masses of these atoms,

This table finds the mass of atoms.

2 H:	2 × 1.01 =	2.02 amu
1 O:		+ 16.00 amu
Total:		18.02 amu

The molecular mass of H_2O is **18.02** amu.

This table finds the mass of atoms.

6 C:	6 × 12.01 =	72.06 amu
6 H:	6 × 1.01 =	+ 6.06 amu
Total:		78.12 amu

The molecular mass of C_6H_6 is **78.12** amu.

This table finds the mass of atoms.

1 N:		14.01 amu
2 O:	2 × 16.00 =	+ 32.00 amu
Total:		46.01 amu

The molecular mass of NO_2 is **46.01** amu.

This table finds the mass of atoms.

2 N:	2 × 14.01 =	28.02 amu
5 O:	5 × 16.00 =	+ 80.00 amu
Total:		108.02 amu

The molecular mass of N_2O_5 is **108.02** amu.

Note that the two different nitrogen and oxygen compounds in these examples have different molecular masses.

? Exercise 5.5.1

What is the molecular mass of each covalent compound?

a. C_2H_2
b. CO
c. CO_2
d. BF_3

Answer a:

26.04 amu

Answer b:

28.01 amu

Answer c:


44.01 amu

Answer d:


67.81 amu

Molecular Shape: VSEPR Theory

Unlike ionic compounds, with their extended crystal lattices, covalent molecules are discrete units with specific three-dimensional shapes. The shape of a molecule is determined by the fact that covalent bonds, which are composed of negatively charged electrons, tend to repel one another. This concept is called the **valence shell electron pair repulsion (VSEPR) theory**. For example, the two covalent bonds in $BeCl_2$ **stay as far from each other** as possible, ending up 180° apart from each other. The result is a ***linear*** molecule:



The three covalent bonds in BF_3 repel each other to form 120° angles in a plane, in a shape called ***trigonal planar***:



The molecules $BeCl_2$ and BF_3 actually violate the octet rule; however, such exceptions are rare and will not be discussed in this text.


> Try sticking three toothpicks into a marshmallow or a gumdrop and see if you can find different positions where your "bonds" are farther apart than the planar 120° orientation.

The four covalent bonds in CCl_4 arrange themselves three dimensionally, pointing toward the corner of a tetrahedron and making bond angles of 109.5°. CCl_4 is said to have a ***tetrahedral*** shape:

Table describes how to use the number of atoms around the central atom to determine the geometry of a molecule, as well as examples.


Atoms Around Central Atom	Geometry	Example
2 AB_2	Linear	$BeCl_2$
3 AB_3	Trigonal Planar	BF_3
4 AB_4	Tetrahedral	CCl_4

In determining the shapes of molecules, it is useful to first determine the Lewis diagram for a molecule. The shapes of molecules with multiple bonds are determined by treating the multiple bonds as one bond. Thus, formaldehyde (CH_2O) has a shape similar to that of BF_3. It is ***trigonal planar***.




Molecules With Lone Pairs Around Central Atom

Molecules with lone electron pairs around the central atom have a shape based on the position of the atoms, not the electron pairs. For example, NH_3 has **one lone electron pair** and **three bonded electron pairs**. These four electron pairs repel each other and adopt a tetrahedral arrangement. However, the shape of the molecule is described in terms of the positions of the atoms, not the lone electron pairs. Thus, NH_3 is said to have a ***trigonal pyramidal*** shape, not a tetrahedral one.



Similarly, H_2O has **two lone pairs** of electrons around the central oxygen atom and **two bonded electron pairs**. Although the four electron pairs adopt a tetrahedral arrangement, the shape of the molecule is described by the positions of the atoms only. The shape of H_2O is ***bent*** with an approximate 109.5° angle.






In summary, to determine the molecular geometry:

Step 1: Draw the Lewis structure.

Step 2: Count the number of bonds (a double/triple bond counts as one) and lone pairs around the **central atom**.

Step 3: Use Table 4.5.1 to determine the molecular geometry.

Number of bonds and lone pairs	Number of bonds 0 lone pair	1 lone pair	2 lone pairs
2	 Linear		
3	 Trigonal planar	 Bent or angular	
4	 Tetrahedral	 Trigonal pyramid	 Bent or angular


Lone Pair

If a molecule has 2 bonds and 0 lone pairs, it is linear. If a molecule has 3 bonds and 0 lone pairs, it is trigonal planar. If a molecule has 3 bonds and 1 lone pair, it is bent or angular. If a molecule has 4 bonds and 0 lone pairs, it is tetrahedral. If a molecule has 4 bonds and 1 lone pair, it is trigonal pyramid. If a molecule has 4 bonds and 2 lone pairs, it is bent or angular.

Table 4.5.1: *The molecular geometry depends on the number of bonds and lone pairs around the* **central atom, A.**


Example 5.5.1

What is the geometry of the ammonium ion, NH_4^+? Its Lewis structure is shown below. How is this different from ammonia, NH_3?




Solution

In ammonium ion, the central atom N has 4 bonds and no lone pair. It is equivalent to the below in Table 4.5.1. Hence, this is *tetrahedral*.



Tetrahedral

In ammonia (NH_3), shown below, N has 3 bonds and one lone pair.




It is equivalent to the below in Table 4.5.1. Hence, the shape of this molecule is *trigonal pyramid*.



Trigonal pyramid


5.5.1 Exercise

What is the molecular shape of nitrosyl chloride, a highly corrosive, reddish-orange gas? Its Lewis structure is shown below.



Answer

Focus on the central atom, N. It has a double bond to O, count this as one bond. It also has a single bond to Cl. Thus, N has 2 bonds and one lone pair. These 3 electron pairs will spread out 120 degrees from each other. But, since the shape is defined by the arrangement of the atoms only, the shape is bent or angular. If you consult Table 4.5.1, this molecule is equivalent to the below. Hence, two bonds and one lone pair has a *bent or angular* shape.



Molecular Polarity

In general, a molecule is nonpolar if all its bonds are nonpolar. Examples are I_2, O_2, H_2, CH_4, C_2H_6 and C_3H_8.

In general, a molecule is polar if it contains polar bonds EXCEPT when the bond polarities cancel each other. As mentioned in Section 4.4, the shape of the CO_2 molecule (linear) orients the two C=O polar bonds **directly opposite** each other, thus **cancelling** each other's effect. **Carbon dioxide** (CO_2) is a ***nonpolar*** molecule.

On the other hand, water (as discussed above) is a bent molecule because of the two lone pairs on the central oxygen atom. Because of the bent shape, the dipoles do not cancel each other out and the water molecule is ***polar***. In the figure below, the individual H-O polar bonds represented by the two red arrows are not directly opposite each other. These two dipoles don't cancel each other out. In fact, the net dipole (blue arrow) points upward. There is a resultant partial positive charge at one end (between the two H atoms) and a partial negative charge on the other end (where O is located). The uneven distribution of charge or the overall dipole is shown by the blue arrow below (Figure 4.5.1). Hence, **water is *polar*** (has + and - poles) while **carbon dioxide is *nonpolar***.




Figure 4.5.1: The molecular geometry of a molecule (linear vs. bent) affects its polarity.

Similarly, in BF_3 (trigonal planar), the effect of a B-F bond is cancelled by the sum of the other two B-F bonds (see video). Hence, a trigonal planar molecule (BF_3) is nonpolar because the bond polarities cancel each other, but a trigonal pyramidal molecule (NH_3) is polar.




Figure 4.5.2: The molecular geometry of a molecule (trigonal planar vs. trigonal pyramid) affects its polarity.

Some other molecules are shown in the figure below. Notice that a tetrahedral molecule such as CCl_4 is ***nonpolar***. However, if the peripheral atoms are not of the same electronegativity, the bond polarities don't cancel and the molecule becomes ***polar***, as in CH_3Cl.




Figure 4.5.3: *The same molecular geometry but peripheral bonds are of different electronegativity. CCl_4 is nonpolar but CH_3Cl is polar.*




Figure **5.5.4**: Molecules with Polar Bonds. Individual bond dipole moments are indicated in red. Due to their different three-dimensional geometry, some molecules with polar bonds have a net dipole moment (HCl, CH_2O, NH_3, and $CHCl_3$), indicated in blue, whereas others do not because the bond dipoles cancel due to symmetry (BCl_3 and CCl_4).

Physical Properties of Molecular Compounds

The physical state and properties of a particular compound depend in large part on the type of chemical bonding it displays. **Molecular compounds**, sometimes called covalent compounds, display a wide range of physical properties due to the different types of intermolecular attractions such as different kinds of polar interactions. The melting and boiling points of molecular compounds are generally quite low compared to those of **ionic compounds**. This is because the energy required to disrupt the **intermolecular forces** (discussed in Section 8.1) between molecules is far less than the energy required to break the ionic bonds in a crystalline ionic compound. Since molecular compounds are composed of neutral molecules, their electrical conductivity is generally quite poor, whether in the solid or liquid state. Ionic compounds do not conduct electricity in the solid state because of their rigid structure, but conduct well when either molten or dissolved into a solution. The water solubility of molecular compounds is variable and depends primarily on the type of intermolecular forces involved. Substances that exhibit hydrogen bonding or dipole-dipole forces are generally water soluble, whereas those that exhibit only London dispersion forces are generally insoluble. Most, but not all, ionic compounds are quite soluble in water. The table below summarizes some of the differences between ionic and molecular compounds.

Table 4.5.2: Comparison of Ionic and Molecular Compounds

Property	Ionic Compounds	Molecular Compounds
Type of elements	Metal and nonmetal	Nonmetals only
Bonding	Ionic - transfer of electron(s) between atoms	Covalent - sharing of pair(s) of electrons between atoms
Representative unit	Formula unit	Molecule
Physical state at room temperature	Solid	Gas, liquid, or solid
Water solubility	Usually high	Variable
Melting and boiling temperatures	Generally high	Generally low
Electrical conductivity	Good when molten or in solution	Poor

In summary, covalent compounds are softer, have lower boiling and melting points, are more flammable, are less soluble in water and do not conduct electricity compared to ionic compounds. The individual melting and boiling points, solubility and other physical properties of molecular compounds depend on molecular polarity.


Example 5.5.2

Describe the shape of each molecule. Is it polar or nonpolar?

a. PCl_3
b. CO_2


Solution

a. The Lewis diagram for PCl_3 is as follows:



Focus on the central atom, P that has 3 bonds and one lone pair. The four electron pairs arrange themselves tetrahedrally, but the lone electron pair is not considered in describing the molecular shape. Like NH_3, this molecule is *pyramidal*. The 3 P-Cl bonds don't cancel each other. This is *polar.*

- The Lewis diagram for CO_2 is as follows:




Focus on the central atom, C. The multiple bonds are treated as one group, hence C has 2 bonds and zero lone pair. CO_2 has only two groups of electrons that repel each other. They will direct themselves 180° apart from each other, so CO_2 molecules are *linear*. This is highly symmetrical, with the two opposite dipoles cancelling each other. The CO_2 molecule is *nonpolar*.


Exercise 5.5.2

Describe the shape of each molecule. Is it polar or nonpolar?

a. CBr_4
b. BCl_3


Answer a:



or


The Lewis structure shows 4 groups attached to the central atom, hence *tetrahedral*. All the 4 groups are identical and the shape is symmetrical. Hence, it is *nonpolar.*

Answer b:



The Lewis diagram shows 3 groups attached to the central atom, hence *trigonal planar*. All the 3 groups are identical and shape is symmetrical, hence, it is *nonpolar.*

Key Takeaways

- A molecule has a certain mass, called the molecular mass.
- Simple molecules have geometries that can be determined from VSEPR theory.
- Polar molecules result from differences in electronegativity of the atoms in the molecule.
- Dipoles that are directly opposite one another cancel each other out.

5.5: Characteristics of Molecules is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.5: Characteristics of Molecules** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.6: Intermolecular Forces


The physical properties of melting point, boiling point, vapor pressure, evaporation, viscosity, surface tension, and solubility are related to the strength of attractive forces between molecules. These attractive forces are called Intermolecular Forces. The amount of "stick togetherness" is important in the interpretation of the various properties listed above.

Introduction

There are four types of intermolecular forces. Most of the intermolecular forces are identical to bonding between atoms in a single molecule. Intermolecular forces just extend the thinking to forces between molecules and follows the patterns already set by the bonding within molecules.


Ionic Forces

The forces holding ions together in ionic solids are electrostatic forces. Opposite charges attract each other. These are the strongest intermolecular forces. Ionic forces hold many ions in a crystal lattice structure.




Dipole Forces

Polar covalent molecules are sometimes described as "dipoles", meaning that the molecule has two "poles". One end (pole) of the molecule has a partial positive charge while the other end has a partial negative charge. The molecules will orientate themselves so that the opposite charges attract principle operates effectively. For example, hydrochloric acid (***HCl***) is a polar molecule with the partial positive charge on the hydrogen and the partial negative charge on the chlorine. A network of partial + and - charges attract molecules to each other.



Hydrogen Bonding


The hydrogen bond is really a special case of dipole forces. A hydrogen bond is the attractive force between the hydrogen attached to an electronegative atom of one molecule and an electronegative atom of a different molecule. Usually the electronegative atom is oxygen, nitrogen, or fluorine. In other words - The hydrogen on one molecule attached to O or N that is attracted to an O or N of a different molecule.



In the graphic below, the hydrogen is partially positive and attracted to the partially negative charge on the oxygen or nitrogen. Because oxygen has two lone pairs, two different hydrogen bonds can be made to each oxygen. This is a very specific bond as indicated. Some combinations that are not hydrogen bonds include: hydrogen to another hydrogen or hydrogen to a carbon.

London Dispersion or Induced Dipole or Van der Waals Forces

Forces between essentially non-polar molecules are the weakest of all intermolecular forces. "Temporary dipoles" are formed by the shifting of electron clouds within molecules. These temporary dipoles attract or repel the electron clouds of nearby non-polar molecules. The temporary dipoles may exist for only a fraction of a second but a force of attraction also exist for that fraction of time. The strength of induced dipole forces depends on how easily electron clouds can be distorted. Large atoms or molecules with many electrons far removed from the nucleus are more easily distorted.



Contributors

- Charles Ophardt, Professor Emeritus, Elmhurst College
- Virtual Chembook

5.6: Intermolecular Forces is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

5.7: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Atoms can share pairs of valence electrons to obtain a valence shell octet. This sharing of electrons is a **covalent bond**. A species formed from covalently bonded atoms is a **molecule** and is represented by a **molecular formula**, which gives the number of atoms of each type in the molecule. The two electrons shared in a covalent bond are called a **bonding pair of electrons**. The electrons that do not participate in covalent bonds are called **nonbonding pairs** (or **lone pairs**) **of electrons**. A covalent bond consisting of one pair of shared electrons is called a **single bond**.

Covalent bonds occur between nonmetal atoms. Naming simple covalent compounds follows simple rules similar to those for ionic compounds. However, for covalent compounds, numerical prefixes are used as necessary to specify the number of atoms of each element in the compound.

In some cases, more than one pair of electrons is shared to satisfy the octet rule. Two pairs of electrons are shared by two atoms to make a **double bond**. Three pairs of atoms are shared to make a **triple bond**. Single, double, and triple covalent bonds may be represented by one, two, or three dashes, respectively, between the symbols of the atoms.

The distance between two covalently bonded atoms is the **bond length**. Bond lengths depend on the types of atoms participating in the bond as well as the number of electron pairs being shared. A covalent bond can be a **polar covalent bond** if the electron sharing between the two atoms is unequal. If the sharing is equal, the bond is a **nonpolar covalent bond**. Because the strength of an atom's attraction for electrons in a bond is rated by the atom's **electronegativity**, the difference in the two atoms' electronegativities indicates how polar a covalent bond between those atoms will be.

The mass of a molecule is called its **molecular mass** and is the sum of the masses of the atoms in the molecule. The shape of a molecule can be predicted using **valence shell electron pair repulsion (VSEPR)**, which uses the fact that the negative electrons in covalent bonds repel each other as much as possible.

Organic chemistry is the chemistry of carbon compounds. Carbon forms covalent bonds with other carbon atoms and with the atoms of many other elements. The simplest organic compounds are **hydrocarbons**, which consist solely of carbon and hydrogen. Hydrocarbons containing only single bonds are called **alkanes (saturated hydrocarbons)**. Hydrocarbons containing carbon–carbon double bonds are **alkenes**, while hydrocarbons with carbon–carbon triple bonds are **alkynes**. Carbon-carbon double and triple bonds are examples of **functional groups**, atoms or bonds that impart a characteristic chemical function to the molecule. Other functional groups include the alcohol functional group (OH) and the **carboxyl functional group** (COOH). They are the characteristic functional group in organic compounds called **alcohols** and **carboxylic acids**.

5.7: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

6: Introduction to Chemical Reactions

Chemical change is a central concept in chemistry. The goal of chemists is to know how and why a substance changes in the presence of another substance or even by itself. Because there are tens of millions of known substances, there are a huge number of possible chemical reactions. In this chapter, we will find that many of these reactions can be classified into a small number of categories according to certain shared characteristics.

6.1: Prelude to Introduction to Chemical Reactions
6.2: The Law of Conservation of Matter
6.3: Chemical Equations
6.4: Quantitative Relationships Based on Chemical Equations
6.5: Types of Chemical Reactions
6.6: Oxidation-Reduction (Redox) Reactions
6.7: Redox Reactions in Organic Chemistry and Biochemistry
6.E: Introduction to Chemical Reactions (Exercises)
6.S: Introduction to Chemical Reactions (Summary)

6: Introduction to Chemical Reactions is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

6.1: Prelude to Introduction to Chemical Reactions

Although yeast has been used for thousands of years, its true nature has been known only for the last two centuries. Yeasts are single-celled fungi. About 1,000 species are recognized, but the most common species is *Saccharomyces cerevisiae*, which is used in bread making. Other species are used for the fermentation of alcoholic beverages. Some species can cause infections in humans.

Yeasts live primarily on sugars, such as glucose ($\mathrm{C_6H_{12}O_6}$). They convert glucose into carbon dioxide ($\mathrm{CO_2}$) and ethanol ($\mathrm{C_2H_5OH}$) in a chemical transformation that is represented as follows:

$$\mathrm{C_6H_{12}O_6\,(s) \longrightarrow 2\,CO_2\,(g) + 2\,C_2H_5OH(\ell)} \tag{6.1.1}$$

Bread making depends on the production of carbon dioxide. The gas, which is produced in tiny pockets in bread dough, acts as a leavening agent: it expands during baking and makes the bread rise. Leavened bread is softer, lighter, and easier to eat and chew than unleavened bread. The other major use of yeast, fermentation, depends on the production of ethanol, which results from the same chemical transformation. Some alcoholic beverages, such as champagne, can also be carbonated using the carbon dioxide produced by the yeast.

Yeast is among the simplest life forms on Earth, yet it is absolutely necessary for at least two major food industries. Without yeast to turn dough into bread and juice into wine, these foods and food industries would not exist today.

6.1: Prelude to Introduction to Chemical Reactions is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

6.2: The Law of Conservation of Matter

Learning Objectives

- Correctly define a law as it pertains to science.
- State the law of conservation of matter.

In science, a law is a general statement that explains a large number of observations. Before being accepted, a law must be verified many times under many conditions. Laws are therefore considered the highest form of scientific knowledge and are generally thought to be inviolable. Scientific laws form the core of scientific knowledge. One scientific law that provides the foundation for understanding in chemistry is the law of conservation of matter. It states that in any given system that is closed to the transfer of matter (in and out), the amount of matter in the system stays constant. A concise way of expressing this law is to say that the amount of matter in a system is *conserved*.

With the development of more precise ideas on elements, compounds and mixtures, scientists began to investigate how and why substances react. French chemist A. Lavoisier laid the foundation to the scientific investigation of matter by describing that substances react by following certain laws. These laws are called the laws of chemical combination. These eventually formed the basis of Dalton's Atomic Theory of Matter.

Law of Conservation of Mass

According to this law, during any physical or chemical change, the total mass of the products remains equal to the total mass of the reactants.

$$\underbrace{\overbrace{\mathrm{HgO\,(s)}}^{\text{Mercuric oxide}}}_{100\ \mathrm{g}} \rightarrow \underbrace{\overbrace{\mathrm{Hg\,(l)}}^{\text{Mercury}}}_{92.6\ \mathrm{g}} + \underbrace{\overbrace{\mathrm{O_2\,(g)}}^{\text{Oxygen}}}_{7.4\ \mathrm{g}}$$

Another way of stating this is, "In a chemical reaction, matter is neither created nor destroyed." The law of conservation of mass is also known as the "law of indestructibility of matter."

Example 6.2.1

If heating 10 grams of $\mathrm{CaCO_3}$ produces 4.4 g of $\mathrm{CO_2}$ and 5.6 g of CaO, show that these observations are in agreement with the law of conservation of mass.



A sample of calcium carbonate (CaCO3). (Public Domain; Walkerma).

Solution

- Mass of the reactants, $\mathrm{CaCO_3}$: $10\,g$
- Mass of the products, $\mathrm{CO_2}$ and CaO: $4.4\,g + 5.6\,g = 10\,g$.

Because the mass of the reactants = the mass of the products, the observations are in agreement with the law of conservation of mass.

What does this mean for chemistry? In any chemical change, one or more initial substances change into a different substance or substances. Both the initial and final substances are composed of atoms because all matter is composed of atoms. According to the

law of conservation of matter, matter is neither created nor destroyed, so we must have the **same number and kind of atoms** after the chemical change as were present before the chemical change.

It may seem as though burning destroys matter, but the same amount, or mass, of matter still exists after a campfire as before. Figure 5.1.1 shows that when wood burns, it combines with oxygen and changes not only to ashes, but also to carbon dioxide and water vapor. The gases float off into the air, leaving behind just the ashes. Suppose we had measured the mass of the wood before it burned and the mass of the ashes after it burned. Also suppose we had been able to measure the oxygen used by the fire and the gases produced by the fire. What would we find? The total mass of matter after the fire would be the same as the total mass of matter before the fire.




Figure 5.1.1: Burning is a chemical process. The flames are caused as a result of a fuel undergoing combustion (burning). Images used with permission (CC BY-SA 2.5; Einar Helland Berger for fire and for ash).

? Exercise 6.2.1

a. What is the law of conservation of matter?
b. How does the law of conservation of matter apply to chemistry?

Answer a:

The law of conservation of matter states that in any given system that is closed to the transfer of matter, the amount of matter in the system stays constant

Answer b:

The law of conservation of matter says that in chemical reactions, the total mass of the products must equal the total mass of the reactants.

Key Takeaway

The amount of matter in a closed system is conserved.

6.2: The Law of Conservation of Matter is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **5.1: The Law of Conservation of Matter** by Binod Shrestha is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.3: Chemical Equations

Learning Objectives

- Define chemical reaction.
- Use a balanced chemical equation to represent a chemical reaction.

Water (H_2O) is composed of hydrogen and oxygen. Suppose we imagine a process in which we take some elemental hydrogen (H_2) and elemental oxygen (O_2) and let them react to make water. The statement

"hydrogen and oxygen react to make water"

is one way to represent that process, which is called a chemical reaction. Figure **6.3.1** shows a rather dramatic example of this very reaction.




Figure **6.3.1**: The Formation of Water. Hydrogen and oxygen combine to form water. Here, the hydrogen gas in the zeppelin *SS Hindenburg* reacts with oxygen in the air to make water. Source: Photo courtesy of the US Navy. For a video of this see www.youtube.com/watch?v=CgWHbpMVQ1U.

To simplify the writing of reactions, we use formulas instead of names when we describe a reaction. We can also use symbols to represent other words in the reaction. A **plus sign** connects the initial substances (and final substances, if there is more than one), and an **arrow** (→) represents the chemical change:

$$\mathbf{H_2 + O_2 \rightarrow H_2O} \tag{6.3.1}$$

This statement is one example of a **chemical equation**, an abbreviated way of using symbols to represent a **chemical change**. The substances on the left side of the arrow are called **reactants**, and the substances on the right side of the arrow are called **products**. It is not uncommon to include a phase label with each formula—(s) for solid, (ℓ) for liquid, (g) for gas, and (aq) for a substance dissolved in water, also known as an *aqueous solution*. If we included phase labels for the reactants and products, under normal environmental conditions, the reaction would be as follows:

$$\mathbf{H_2(g) + O_2(g) \rightarrow H_2O(\ell)} \tag{6.3.2}$$

Chemical equations can also be used to describe physical changes. We will see examples of this soon.

This equation is still not complete because *it does not satisfy the* ***law of conservation of matter***. Count the **number of atoms of each element** on each side of the arrow. On the reactant side, there are two H atoms and two O atoms; on the product side, there are two H atoms and only one oxygen atom. The equation is **not balanced** because the number of oxygen atoms on each side is not the same (Figure **6.3.2**).




Figure **6.3.2**: Balanced—Yes or No?. By counting the atoms of each element, we can see that the reaction is not balanced as written.

To make this chemical equation conform to the law of conservation of matter, we must revise the amounts of the reactants and the products as necessary to get the same number of atoms of a given element on each side. Because every substance has a characteristic chemical formula, we cannot change the chemical formulas of the individual substances. For example, we cannot

change the formula for elemental oxygen to O. However, we can assume that different numbers of reactant molecules or product molecules may be involved. For instance, perhaps two water molecules are produced, not just one:

$$H_2(g) + O_2(g) \rightarrow 2H_2O(\ell) \tag{6.3.3}$$

The **2** preceding the formula for water is called a **coefficient**. It implies that two water molecules are formed. There are now two oxygen atoms on each side of the equation.

> This point is so important that we should repeat it. You **cannot** change the formula of a chemical substance to balance a chemical reaction! You **must** use the proper chemical formula of the substance.

Unfortunately, by inserting the coefficient 2 in front of the formula for water, we have also changed the number of hydrogen atoms on the product side as well. As a result, we no longer have the same number of hydrogen atoms on each side. This can be easily fixed, however, by putting a coefficient of 2 in front of the diatomic hydrogen reactant:

$$2H_2(g) + O_2(g) \rightarrow 2H_2O(\ell) \tag{6.3.4}$$

Now we have four hydrogen atoms and two oxygen atoms on each side of the equation. The **law of conservation of matter is satisfied** because we now have the same number of atoms of each element in the reactants and in the products. We say that the reaction is **balanced** (Figure **6.3.3**). The diatomic oxygen has a coefficient of 1, which typically is not written but assumed in balanced chemical equations.




Figure **6.3.3**: Balanced—Yes or No?. By counting the atoms of each element, we can see that the reaction is now balanced.

Proper chemical equations should be balanced. Writing balanced reactions is a chemist's way of acknowledging the law of conservation of matter.

✔ Example 6.3.1

Is each chemical equation balanced?

a. $2Na(s) + O_2(g) \longrightarrow 2Na_2O(s)$
b. $CH_4(g) + 2O_2(g) \longrightarrow CO_2(g) + 2H_2O(\ell)$
c. $AgNO_3(aq) + 2KCl(aq) \longrightarrow AgCl(s) + KNO_3(aq)$

Solution

a. By counting, we find two sodium atoms and two oxygen atoms in the reactants and four sodium atoms and two oxygen atoms in the products. This equation is not balanced.
b. The reactants have one carbon atom, four hydrogen atoms, and four oxygen atoms. The products have one carbon atom, four hydrogen atoms, and four oxygen atoms. This equation is balanced.
c. The reactants have one silver atom, one nitrogen atom, three oxygen atoms, two potassium atoms, and two chlorine atoms. The products have one silver atom, one chlorine atom, one potassium atom, one nitrogen atom, and three oxygen atoms. Because there are different numbers of chlorine and potassium atoms, this equation is not balanced.

? Exercise 6.3.1

Is each chemical equation balanced?

a. $2Hg(l) + O_2(g) \longrightarrow Hg_2O_2(s)$
b. $C_2H_4(g) + 2O_2(g) \longrightarrow 2CO_2(g) + 2H_2O(l)$
c. $Mg(NO_3)_2(s) + 2Li(s) \longrightarrow Mg(s) + 2LiNO_3(s)$.

Answer a:

balanced

Answer b:

O is not balanced; the 4 atoms of oxygen on the left does not balance with the 6 oxygen atoms on the right

Answer c:

balanced

How does one balance a chemical equation, starting with the correct formulas of the reactants and products? Basically, a **back-and-forth approach** is adopted, counting the number of atoms of one element on one side, checking the number of atoms of that element on the other side, and changing a coefficient if necessary. Then check another element, going back and forth from one side of the equation to another, until each element has the same number of atoms on both sides of the arrow. In many cases, it does not matter which element is balanced first and which is balanced last, as long as all elements have the same number of atoms on each side of the equation.

Below are guidelines for writing and balancing chemical equations.

Writing and Balancing Chemical Equations

1. Determine the correct chemical formulas for each reactant and product. Write the skeleton equation.
2. Count the number of atoms of each element that appears as a reactant and as a product. If a polyatomic ion is unchanged on both sides of the equation, count it as a unit.
3. Balance each element one at a time by placing coefficients in front of the formulas. No coefficient is written for a 1. It is best to begin by balancing elements that only appear in one chemical formula on each side of the equation. NEVER change the subscripts in a chemical formula - you can only balance equations by using coefficients.
4. Check each atom or polyatomic ion to be sure that they are equal on both sides of the equation.
5. Make sure that all coefficients are in the lowest possible ratio. If necessary, reduce to the lowest ratio.

For example, to balance the equation

$$\mathrm{CH_4 + Cl_2 \rightarrow CCl_4 + HCl}$$

Step 1: Write the skeleton equation with the correct formulas.

$$\mathrm{CH_4 + Cl_2 \rightarrow CCl_4 + HCl} \tag{6.3.5}$$

Step 2: Count the number of each atom or polyatomic ion on both sides of the equation.

Reactants	Products
1 C atom	1 C atom
4 H ions	1 H ions
2 Cl atom	5 Cl atoms

Step 3: We find that both sides are already balanced with one carbon atom. So we proceed to balance the hydrogen atoms. We find that the reactant side has four hydrogen atoms, so the product side must also have four hydrogen atoms. This is balanced by putting a 4 in front of the HCl:

$$\mathrm{CH_4 + Cl_2 \rightarrow CCl_4 + 4\,HCl} \tag{6.3.6}$$

Reactants	Products
1 C atom	1 C atom
4 H ions	4 H ions
2 Cl atom	8 Cl atoms

Now each side has four hydrogen atoms. The product side has a total of eight chlorine atoms (four from the $\mathrm{CCl_4}$ and four from the four molecules of HCl), so we need eight chlorine atoms as reactants. Because elemental chlorine is a diatomic molecule, we need four chlorine molecules to get a total of eight chlorine atoms. We add another 4 in front of the $\mathrm{Cl_2}$ reactant:

$$CH_4 + 4\,Cl_2 \rightarrow CCl_4 + 4\,HCl \tag{6.3.7}$$

Reactants	Products
1 C atom	1 C atom
4 H ions	4 H ions
8 Cl atom	8 Cl atoms

Step 3: Now we check: each side has one carbon atom, four hydrogen atoms, and eight chlorine atoms. The chemical equation is balanced. And, the coefficients are in the lowest possible ratio.

Example 6.3.2

Aqueous solutions of lead (II) nitrate and sodium chloride are mixed. The products of the reaction are an aqueous solution of sodium nitrate and a solid precipitate of lead (II) chloride. Write the balanced chemical equation for this reaction.

Solution

Step 1: Write the skeleton equation with the correct formulas.

$$Pb(NO_3)_2\,(aq) + NaCl\,(aq) \rightarrow NaNO_3\,(aq) + PbCl_2\,(s)$$

Step 2: Count the number of each atom or polyatomic ion on both sides of the equation.

Reactants	Products
1 Pb atom	1 Pb atom
2 NO_3^- ions	1 NO_3^- ions
1 Na atom	1 Na atom
1 Cl atom	2 Cl atoms

Step 3: Solve. The nitrate ions and the chlorine atoms are unbalanced. Start by placing a 2 in front of the $NaCl$. This increases the reactant counts to 2 Na atoms and 2 Cl atoms. Then place a 2 in front of the $NaNO_3$. The result is:

$$Pb(NO_3)_2\,(aq) + 2NaCl\,(aq) \rightarrow 2NaNO_3\,(aq) + PbCl_2\,(s)$$

Step 4: The new count for each atom and polyatomic ion becomes:

Reactants	Products
1 Pb atom	1 Pb atom
2 NO_3^- ions	2 NO_3^- ions
2 Na atom	2 Na atom
2 Cl atom	2 Cl atoms

Step 5: Think about the result.

The equation is now balanced since there are equal numbers of atoms of each element on both sides of the equation. And, the coefficients are in the lowest possible ratio.

Exercise 6.3.2

Fermentation is a biochemical process that enables yeast cells to live in the absence of oxygen. Humans have exploited it for centuries to produce wine and beer and make bread rise. In fermentation, sugars such as glucose ($C_6H_{12}O_6$) are converted to ethanol (C_2H_5OH) and carbon dioxide CO_2. Write a balanced chemical reaction for the fermentation of glucose.



Commercial use of fermentation. (a) Microbrewery vats are used to prepare beer. (b) The fermentation of glucose by yeast cells is the reaction that makes beer production possible.

Answer

$$C_6H_{12}O_6\,(s) \longrightarrow 2\,C_2H_5OH\,(l) + 2\,CO_2\,(g)$$

Key Takeaway

Chemical reactions are represented by chemical equations that list reactants and products. Proper chemical equations are balanced; the same number of each element's atoms appears on each side of the equation.

6.3: Chemical Equations is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **5.2: Chemical Equations** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.4: Quantitative Relationships Based on Chemical Equations

Learning Objectives

- To calculate the amount of one substance that will react with or be produced from a given amount of another substance.

A balanced chemical equation not only describes some of the chemical properties of substances—by showing us what substances react with what other substances to make what products—but also shows numerical relationships between the reactants and the products. The study of these numerical relationships is called stoichiometry. The stoichiometry of chemical equations revolves around the coefficients in the balanced chemical equation because these coefficients determine the molecular ratio in which reactants react and products are made.

> *The word stoichiometry is pronounced "stow-eh-key-OM-et-tree." It is of mixed Greek and English origins, meaning roughly "measure of an element."*

Looking Closer: Stoichiometry in Cooking

Let us consider a stoichiometry analogy from the kitchen. A recipe that makes 1 dozen biscuits needs 2 cups of flour, 1 egg, 4 tablespoons of shortening, 1 teaspoon of salt, 1 teaspoon of baking soda, and 1 cup of milk. If we were to write this as a chemical equation, we would write

$$\text{2 c flour + 1 egg + 4 tbsp shortening + 1 tsp salt + 1 tsp baking soda + 1 c milk} \rightarrow \text{12 biscuits}$$

(Unlike true chemical reactions, this one has all 1 coefficients written explicitly—partly because of the many different units here.) This equation gives us ratios of how much of what reactants are needed to make how much of what product. Two cups of flour, when combined with the proper amounts of the other ingredients, will yield 12 biscuits. One teaspoon of baking soda (when also combined with the right amounts of the other ingredients) will make 12 biscuits. One egg must be combined with 1 cup of milk to yield the product food. Other relationships can also be expressed.

We can use the ratios we derive from the equation for predictive purposes. For instance, if we have 4 cups of flour, how many biscuits can we make if we have enough of the other ingredients? It should be apparent that we can make a double recipe of 24 biscuits.

But how would we find this answer formally, that is, mathematically? We would set up a conversion factor, much like we did in Chapter 1. Because 2 cups of flour make 12 biscuits, we can set up an equivalency ratio:

$$\frac{\text{12 biscuits}}{\text{2 c flour}}$$

We then can use this ratio in a formal conversion of flour to biscuits:

$$\text{4 c flour} \times \frac{\text{12 biscuits}}{\text{2 c flour}} = \text{24 biscuits}$$

Similarly, by constructing similar ratios, we can determine how many biscuits we can make from any amount of ingredient. When you are doubling or halving a recipe, you are doing a type of stoichiometry. Applying these ideas to chemical reactions should not be difficult if you use recipes when you cook.



A recipe shows how much of each ingredient is needed for the proper reaction to take place. (CC BY-SA 3.0 Unported; Peng via Wikipedia)

Consider the following balanced chemical equation:

$$2\,C_2H_2 + 5\,O_2 \rightarrow 4\,CO_2 + 2\,H_2O \tag{6.4.1}$$

The coefficients on the chemical formulas give the ratios in which the reactants combine and the products form. Thus, we can make the following statements and construct the following ratios:

Table uses a statement from the balanced chemical reaction to show ratios and inverse ratios.

Statement from the Balanced Chemical Reaction	Ratio	Inverse Ratio
two C_2H_2 molecules react with five O_2 molecules	$\frac{2C_2H_2}{5O_2}$	$\frac{5O_2}{2C_2H_2}$
two C_2H_2 molecules react to make four CO_2 molecules	$\frac{2C_2H_2}{4CO_2}$	$\frac{4CO_2}{2C_2H_2}$
five O_2 molecules react to make two H_2O molecules	$\frac{5O_2}{2H_2O}$	$\frac{2H_2O}{5O_2}$
four CO_2 molecules are made at the same time as two H_2O molecules	$\frac{2H_2O}{4CO_2}$	$\frac{4CO_2}{2H_2O}$

Other relationships are possible; in fact, 12 different conversion factors can be constructed from this balanced chemical equation. In each ratio, the unit is assumed to be molecules because that is how we are interpreting the chemical equation.

Any of these fractions can be used as a conversion factor to relate an amount of one substance to an amount of another substance. For example, suppose we want to know how many CO_2 molecules are formed when 26 molecules of C_2H_2 are reacted. As usual with a conversion problem, we start with the amount we are given—$26C_2H_2$—and multiply it by a conversion factor that cancels out our original unit and introduces the unit we are converting to—in this case, CO_2. That conversion factor is $\frac{4CO_2}{2C_2H_2}$, which is composed of terms that come directly from the balanced chemical equation. Thus, we have

$$26C_2H_2 \times \frac{4CO_2}{2C_2H_2}$$

The molecules of C_2H_2 cancel, and we are left with molecules of CO_2. Multiplying through, we get

$$26C_2H_2 \times \frac{4CO_2}{2C_2H_2} = 52CO_2$$

Thus, 52 molecules of CO_2 are formed.

This application of stoichiometry is extremely powerful in its predictive ability, as long as we begin with a balanced chemical equation. Without a balanced chemical equation, the predictions made by simple stoichiometric calculations will be incorrect.

Example 6.4.1

Start with this balanced chemical equation.

$$\mathbf{KMnO_4 + 8\,HCl + 5\,FeCl_2 \longrightarrow 5\,FeCl_3 + MnCl_2 + 4\,H_2O + KCl}$$

a. Verify that the equation is indeed balanced.
b. Give 2 ratios that give the relationship between $\mathbf{HCl}$ and $\mathbf{FeCl_3}$.

Solution a

Each side has 1 K atom and 1 Mn atom. The 8 molecules of HCl yield 8 H atoms, and the 4 molecules of H_2O also yield 8 H atoms, so the H atoms are balanced. The Fe atoms are balanced, as we count 5 Fe atoms from 5 $FeCl_2$ reactants and 5 $FeCl_3$ products. As for Cl, on the reactant side, there are 8 Cl atoms from HCl and 10 Cl atoms from the 5 $FeCl_2$ formula units, for a total of 18 Cl atoms. On the product side, there are 15 Cl atoms from the 5 $FeCl_3$ formula units, 2 from the $MnCl_2$ formula unit, and 1 from the KCl formula unit. This is a total of 18 Cl atoms in the products, so the Cl atoms are balanced. All the elements are balanced, so the entire chemical equation is balanced.

Solution b

Because the balanced chemical equation tells us that 8 HCl molecules react to make 5 $FeCl_3$ formula units, we have the following 2 ratios: $\mathbf{\frac{8HCl}{5FeCl_3}}$ **and** $\mathbf{\frac{5FeCl_3}{8HCl}}$. There are a total of 42 possible ratios. Can you find the other 40 relationships?

Exercise 6.4.1

Start with this balanced chemical equation.

$$\mathbf{2KMnO_4 + 3CH_2{=}CH_2 + 4H_2O \rightarrow 2MnO_2 + 3HOCH_2CH_2OH + 2KOH}$$

a. Verify that the equation is balanced.
b. Give 2 ratios that give the relationship between $KMnO_4$ and $CH_2{=}CH_2$. (A total of 30 relationships can be constructed from this chemical equation. Can you find the other 28?)

Answer a:

Each side has 2 K atoms and 2 Mn atoms. On the reactant side, 3 $CH_2{=}CH_2$ yield 6 C atoms and on the product side, 3 $HOCH_2CH_2OH$ also yield 6 C atoms, so the C atoms are balanced. There are 20 H atoms on the reactants side: 12 H atoms from 3 $CH_2{=}CH_2$ and 8 H atoms from 4 H_2O. On the product side, there are also 20 H atoms: 18 H atoms from 3 $HOCH_2CH_2OH$ and 2 H atoms from 2 KOH. So, the H atoms are balanced. As for O, on the reactant side, there are 8 O atoms from 2 $KMnO_4$ and 4 O atoms from 4 H_2O, for a total of 12 O atoms. On the product side, there are 4 O atoms from the 2 MnO_2 formula units, 6 O atoms from 3 $HOCH_2CH_2OH$, and 2 O atoms from the 2 KOH formula units. This is a total of 12 O atoms in the products, so the O atoms are balanced. All the elements are balanced, so the entire chemical equation is balanced.

Answer b:

Because the balanced chemical equation tells us that 2 $KMnO_4$ formula units react with 3 $CH_2{=}CH_2$ molecules, we have the following 2 ratios: $\mathbf{\frac{2KMnO_4}{3CH_2{=}CH_2}}$ **and** $\mathbf{\frac{3CH_2{=}CH_2}{2KMnO_4}}$. There are a total of 30 possible ratios. Can you find the other 28 relationships?

Key Takeaway

The coefficients in a balanced chemical equation give the ratios in which molecules of substances react and are produced in a chemical reaction.

6.4: Quantitative Relationships Based on Chemical Equations is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **5.3: Quantitative Relationships Based on Chemical Equations** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.5: Types of Chemical Reactions

learning objectives

- List and define the 5 types of reactions based on the rearrangement of atoms
- Match given reactions to their respective category of reaction type
- Relate combustion reactions of hydrocarbons to climate change
- Relate decomposition reactions to production of hydrogen fuel cells.

Some Basic Types of Chemical Reactions

We are going to identify 5 basic "types" of chemical reactions because they have characteristic equations that allow us to use different techniques when we balance them. That is, in the next section when we balance a reaction, the first thing we will ask is what type of reaction it is, and then we will use the technique that is appropriate to that type reaction. There are other ways of classifying reactions that may actually tell us more about the chemistry, like are electrons transferred between atoms when a reaction occurs, which we will discuss later in this Chapter, and are called oxidation-reduction (redox) reactions, or acid base reactions (which at this level, occur when protons are transferred between between reactants).

1. **Formation Reactions (synthesis reactions)**
2. **Decomposition Reactions**
3. **Single Displacement Reactions**
4. **Double Displacement Reaction**
5. **Combustion Reactions**

Formation or Synthesis Reactions

$$H_2(g) + Cl_2(g) \text{ ® } 2HCl(g)$$

As the name implies, two or more chemicals come together and form another chemical. Here, HCl is being formed from hydrogen and chlorine.

Decomposition Reaction

$$CaCO_3(s) \text{ --> } CaCO(s) + CO_2(g)$$

As the name implies, something decomposes into something else. Here Calcium Carbonate decomposes to calcium oxide and carbon dioxide

Single Displacement Reactions

$$Fe(s) + CuSO_4(aq) \text{ -> } FeSO_4(aq) + Cu(s)$$
$$F_2(g) + BaCl_2(aq) \text{ -> } BaF_2(g) + Cl_2(g)$$

In a single displacement reaction something that is "pure" displaces something in a compound. As a rule of thumb, we say metals form cations and nonmetals from anions. So a metal can exist in two forms, as a pure metal, or as a cation (in an ionic compound). Likewise, a nonmetal can exist as something pure, or as an anion (in an ionic compound). In the first of the above two equations, the metal iron displaces the cation Copper(II) and we call that a metal single displacement reaction, while in the second, the fluorine displaces the anion chloride, and so we call that a nonmetal single displacement reaction. Note, sometimes these are called single replacement reactions.

If you note, these reactions involve a change in the charge of the metal or nonmetal, and so they involve the transfer of electrons. We will pick these up again later in this Chapter when we get to section 3.6.2 and discuss these from the perspective of electron transfer.

Metal

$$Fe(s) + AuSO_4(aq) \text{ --> } FeSO_4(aq) + Au(s)$$

Some metals prefer to be ions, and other prefer to be pure. This can be understood by comparing iron ore with gold ore. Iron (pure) spontaneously rusts and iron ore that is mined must consists of compounds like hematite (Fe_2O_3) and magnetite (Fe_3O_4), while gold, a noble metal, does not tend to form ionic compounds, and so pure gold can be found in gold ore. So the above reaction is logical, in the sense that iron as a pure metal is not stable, while gold is.




Figure **6.5.1**: Hematite ore on left (image from flickr) and gold on right (image from wikicommons).

Since the above reaction proceeds, we can correctly predict that:

$$FeSO_4(aq) + Au(s) \text{ --> No Reaction}$$

Nonmetal

$$F_2(g) + BaCl_2(aq) \text{ -> } BaF_2(g) + Cl_2(g)$$

Single Displacement Reactions are actually a subset of another class (type) of reactions called Redox or Oxidation-Reduction Reactions, where electrons are transferred between species. We will cover those in section 3.6 of this Chapter, and at the time come up with a way to predict if a single displacement reaction will occur, or if there is no reaction.

Metal-Nonmetal

A metal can also displace hydrogen from an acid or even from water (look at water as H-OH, where the anion is hydroxide). For example

$$2Na(s) + 2H_2O(l) \text{ ->} H_2(g) + 2NaOH$$

Here the pure sodium metal becomes a cation, while a hydrogen is pulled off of each water to from hydrogen gas (a diatomic), and so hydrogen is behaving similar to a cation on the reactant side, but it is not a cation, it is an acidic hydrogen atom in a covalent bond. These reactions can give off heat which can then combust the hydrogen, as seen in Video **6.5.2**.



Video 6.5.2. 1'09" YouTube uploaded by halosaqi showing alkali metals displacing hydrogen from water.

Double Displacement Reactions

$PbNO_3(aq) + NaS_2(aq) \rightarrow PbS(s) + NaNO_3(aq)$

In a double displacement reaction, the ions of two ionic compounds swap counter ions.



Video 6.5.3: 22 " YouTube uploaded by Bob Belford showing lead(II)nitrate reacting with sodium sulfide to form the black lead(II)sulfide and sodium nitrate (https://youtu.be/X6BV8Av_UnU).

We will look at these in great detail in section 3.4.

Combustion Reaction

In a combustion reaction a compound reacts with oxygen and releases energy. Carbon dioxide and water are also released if the compound contains carbon.

$CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$ + energy

Today's society uses fossil fuels as an energy source and the carbon dioxide emission is a concern to many scientists. The following youtube video was created by the EPA to describe the consequences of our economic reliance on fossil fuels. You may wish to check the link at the end of the video,
www.epa.gov/climatechange/students .



Video 6.5.4: 2'37" youtube posted by EPA (https://youtu.be/ScX29WBJI3w)

Do combustion reactions always produce carbon dioxide and water?

No there are other types of combustion reactions, like the combustion of hydrogen and magnesium.

$2H_2 + O_2$ --> $2H_2O$ + energy
$2Mg + O_2$ -->MgO + energy



Video 6.5.5: RSC video of magnesium reacting with water from Peter Wothers lecture series (https://youtu.be/gAhsvOP5yaM)



Video 6.5.6: RSC video of magnesium reacting with sand (silicon dioxide) from Peter Wothers lecture series (https://youtu.be/J60J-g4ifXg)

The hydrogen economy is considered as an alternative to fossil fuels and an area of intense current chemistry research. The concept is simple, one can use electricity generated by methods like solar, hydroelectric, wind, geothermal to split water into hydrogen and oxygen. Then you can transport the hydrogen to a new location and recombine them to produce energy where you need it, and produce no carbon dioxide. There are obvious challenges, like the storage of the hydrogen. As we shall learn later this semester, energy is the capacity to do work and transfer heat. So any energy lost as heat can not be used for work. You have probably heard of fuel cells, and they operate like a battery that consumes fuel, and can be more efficient than a combustion engine because they do not produce as much heat. In the following video the person uses a reversible fuel cell kit to demonstrate the concept of a hyrdogen economy. Simply speaking, you split water into hydrogen and oxygen when you are near a power source (solar, geothermal, wind, hydroelectric) then you recombine them where you want to use the energy, and in the process, you produce no carbon dioxide.



Video 6.5.5: 4'18" YouTube Uploaded by Andrew Copeland showing basic concepts behind hydrogen economy as an alternative to the fossil fuel economy (https://youtu.be/qZhhnBfXP7I).

Practice Problems

Exercise 6.5.1

Identify the types of reactions for each equation

a. $AlCl_3(aq) + Na_2CO_3(aq) \rightarrow Al_2(CO_3)_3(s) + NaCl(aq)$
b. $C_8H_{18} + O_2 \rightarrow CO_2 + H_2O$
c. $Li_2S + Br_2 \rightarrow 2LiBr + S$

d. H_2CO_3 --> $H_2O + CO_2$
e. $Ba(C_2H_3O_2)_2(aq) + 2HNO_3(aq)$ ® $Ba(NO_3)_2(aq) + 2HC_2H_3O_2(aq)$

Answer a

Double Displacement

Answer b

combustion

Answer c

Single Displacement (non metal)

Answer d

Decomposition

Answer e

Double Displacement (involving acid)

Exercise 6.5.21

Identify the types of reactions for each equation

a. $H_2S(g)$ ® $H_2(g) + S(s)$
b. $Pb + 2CuCl$ ® $PbCl_2 + Cu$
c. $CH_2CO(g) + 2\ O_2(g)$ ® $2\ CO_2(g) + H_2O(g)$
d. $HCl + NaOH$ --> $NaCl + H_2O$

Answer a

Decomposition

Answer b

Single Displacement (metal displacement)

Answer c

Combustion

Answer d

Double Displacement (neutralization)

Contributors and Attributions

Robert E. Belford (University of Arkansas Little Rock; Department of Chemistry). The breadth, depth and veracity of this work is the responsibility of Robert E. Belford, rebelford@ualr.edu. You should contact him if you have any concerns. This material has both original contributions, and content built upon prior contributions of the LibreTexts Community and other resources, including but not limited to:

- November Palmer & Emily Chaote (UALR)
- Ronia Kattoum (Learning Goals)

This page titled 6.5: Types of Chemical Reactions is shared under a not declared license and was authored, remixed, and/or curated by Robert Belford.

6.6: Oxidation-Reduction (Redox) Reactions

Skills to Develop

- Define oxidation and reduction in terms of a gain or loss of oxygen or hydrogen atoms.
- Identify the substances involved in oxidation and reduction in a reaction.
- Identify the oxidizing and reducing agents in a redox reaction.
- To identify a chemical reaction as an oxidation-reduction reaction.

Oxygen's Role in Reactions

Oxygen is an element that has been known for centuries. In its pure elemental form, oxygen is highly reactive, and it readily makes compounds with most other elements. It is also the most abundant element by mass in the Earth's crust. The class of reactions called oxidation and reduction were originally defined with respect to the element oxygen.

Early scientists viewed **oxidation** as a process in which a substance combines with oxygen to produce oxides. Examples of oxidation are shown below: magnesium, methane, iron and copper are being oxidized.

The heating of magnesium in air allows it to combine with oxygen from the air to form magnesium oxide.

$$2\mathrm{Mg}\,(s) + \mathrm{O_2}\,(g) \rightarrow 2\mathrm{MgO}\,(s) \tag{6.6.1}$$

The burning (combustion) of methane allows the fuel (methane) to react with oxygen (from the air), producing carbon dioxide and water (and heat).

$$\mathrm{CH_4}\,(g) + 2\mathrm{O_2}\,(g) \rightarrow \mathrm{CO_2}\,(g) + 2\mathrm{H_2O}\,(g) \tag{6.6.2}$$

The oxidation (commonly known as rusting or corrosion) of iron is shown below. Iron that is exposed to air (and water) is slowly oxidized to produce rust (an oxide of iron).


$$4\mathrm{Fe}\,(s) + 3\mathrm{O_2}\,(g) \rightarrow 2\mathrm{Fe_2O_3}\,(s) \tag{6.6.3}$$

The Statue of Liberty is a landmark every American recognizes. The Statue of Liberty is easily identified by its height, stance, and unique blue-green color. When this statue was first delivered from France, its appearance was not green. It was brown, the color of its copper “skin.” So how did the Statue of Liberty change colors? The change in appearance was a direct result of corrosion. The copper that is the primary component of the statue slowly underwent oxidation from the air. The oxidation-reduction reactions of copper metal in the environment occur in several steps. Copper metal is oxidized to copper(I) oxide Cu_2O), which is red, and then to copper(II) oxide (CuO), which is black.

$$4Cu\,(s) + O_2\,(g) \rightarrow 2Cu_2O\,(s) \tag{6.6.4}$$

$$2Cu_2O\,(s) + O_2\,(g) \rightarrow 4CuO\,(s) \tag{6.6.5}$$

Coal, which was often high in sulfur, was burned extensively in the early part of the last century. As a result, sulfur trioxide, carbon dioxide, and water all reacted with the CuO. These three compounds are responsible for the characteristic blue-green patina seen today. Fortunately, formation of the patina created a protective layer on the surface, preventing further corrosion of the copper skin.



(a) (b)

Figure 6.6.1: (a) The Statue of Liberty is covered with a copper skin, and was originally brown, as shown in this painting. (b) Exposure to the elements has resulted in the formation of the blue-green patina seen today.

Bleaches contain various compounds such as sodium hypochlorite (NaClO) that can oxidize stains by the transfer of oxygen atoms, making the molecules in the stains more water-soluble and therefore easier to rinse off. Hydrogen peroxide (H_2O_2) releases oxygen as it spontaneously decomposes. It also acts as a bleach and is used as an antiseptic that kills bacteria by oxidizing them.

We worry a lot about our smiles. Over the years teeth do discolor some, so the procedure of teeth bleaching has become more and more popular. Best done in a dentist's office, various chemical preparations containing peroxides are used to whiten teeth. Less effective, but

easier to use are "teeth-whitening" toothpastes (also containing peroxides) that promise to give you a brighter smile and improve your life in every way.

Oxidation and Reduction

The opposite of oxidation is called reduction. Since **oxidation was originally defined as the addition of oxygen**, reduction was therefore the **removal of oxygen** from a substance. Many naturally occurring metal ores are present as oxides. The pure metals can be extracted by reduction. For example, pure iron is obtained from iron (III) oxide by reacting it with carbon at high temperatures.

$$2\mathrm{Fe_2O_3}\,(s) + 3\mathrm{C}\,(s) \rightarrow 4\mathrm{Fe}\,(s) + 3\mathrm{CO_2}\,(g) \tag{6.6.6}$$

The **removal of oxygen from the Fe_2O_3 molecule means that it is being reduced** to Fe. Note that an **oxidation process is also occurring simultaneously** in this reaction; the carbon reactant is being oxidized to CO_2. This is an important concept. In looking at organic compounds, we can describe **oxidation and reduction in terms of the loss or gain of oxygen or the loss or gain of hydrogen.** The changes in number of oxygen atoms, and number of hydrogen atoms is summarized below for oxidation and reduction reactions.

Process	Change in oxygen (some reactions)	Change in hydrogen (some reactions)
Oxidation	gain	lose
Reduction	lose	gain

If we look at the **oxidation** of ethanol (CH_3CH_2OH, an alcohol) to form ethanal (CH_3CHO, an aldehyde), we see that the **number of bonds to oxygen has increased and the number of hydrogen atoms has decreased** from six to four. Either or both of these indicate that an oxidation has occurred.

For a **reduction**, we look for the **removal of oxygen or bonds to oxygen or the addition or hydrogen atoms**. In the following reaction, we see ethanoic acid (CH_3COOH, a carboxylic acid) being reduced to ethanal (CH_3CHO, an aldehyde). Note that the number of oxygen atoms is reduced from two to one. There is no change in the number of hydrogen atoms, but both changes do not need to happen for evidence of oxidation or reduction.

If we see evidence of an oxidation or reduction, we know the other must have happened as well. Oxidation and reduction must happen together. Neither can happen alone in a reaction.

Oxidation and Reduction Must Occur at the same time

Note that an oxidation process is simultaneously occurring with a reduction process. Oxidation cannot happen alone, without reduction. If reactant A is gaining oxygen (getting oxidized), reactant B must be losing the O (getting reduced). If reactant A is undergoing the removal of hydrogen (getting oxidized), reactant B must be undergoing the addition of hydrogen (getting reduced).

Redox reactions are also commonly described in terms of oxidizing and reducing agents. In the reaction below, C is being oxidized by gaining oxygen. The source of oxygen is Fe_2O_3. In other words, Fe_2O_3 is causing the C to be oxidized. Consequently, Fe_2O_3 is referred to as the oxidizing agent. Conversely, the C causes the Fe_2O_3 to lose oxygen and become reduced, so C is the reducing agent. An **oxidizing agent** ***(OA)*** *is a substance that causes oxidation by releasing oxygen*, and a **reducing agent** ***(RA)*** *is a substance that causes reduction by gaining oxygen*. Said another way, the oxidizing agent (OA) is the substance that is reduced, while the reducing agent (RA) is the substance that is oxidized. Focusing on the reaction below, **Fe_2O_3 is the substance being reduced, while C is the substance that is being oxidized. Hence,** Fe_2O_3 is the oxidizing agent (OA) and C is the reducing agent (RA).

$$2Fe_2O_3\,(s) + 3C\,(s) \rightarrow 4Fe\,(s) + 3CO_2\,(g) \tag{6.6.7}$$

Fe_2O_3 = substance reduced C = substance oxidized

Fe_2O_3 = oxidizing agent (OA) C = reducing agent (RA)

Note

The *substance oxidized* and *substance reduced* are both reactants in the reaction.

Several classes of organic compounds are related to one another by oxidation and reduction reactions. Alkanes, alkenes, and alkynes represent different levels of oxidation of a hydrocarbon. When an alkane is heated in the presence of an appropriate catalyst, it can be

oxidized to the corresponding alkene in a reaction called a dehydrogenation reaction. Two hydrogen atoms are removed in the process. The alkene can be further oxidized to an alkyne by the removal of two more hydrogen atoms.

$$\text{oxidation:} \qquad CH_3CH_3 \rightarrow CH_2{=}CH_2 \rightarrow CH{\equiv}CH$$

The reactions are reversible, so an alkyne can be reduced first to an alkene and then to an alkane. These are addition reactions.

$$\text{reduction:} \qquad CH{\equiv}CH \rightarrow CH_2{=}CH_2 \rightarrow CH_3CH_3$$

The alkane is the most reduced form of a hydrocarbon, while the alkyne is the most oxidized form.


Oxidation reactions in organic chemistry often involve the addition of oxygen to a compound (or an increase in the number of bonds to oxygen), which changes the functional group that is present. The following sequence shows how methane can be oxidized first to methanol (alcohol), then to methanal (aldehyde), then to methanoic acid (carboxylic acid), and finally to carbon dioxide. Each step in the process is either a gain of oxygen or a loss of hydrogen. Each step also represents energy, which explains why the complete combustion of alkanes to carbon dioxide is an extremely exothermic reaction. Hence, alkanes are used as fuels.

$$\underset{\text{alkane}}{CH_4} \xrightarrow{\text{gain of oxygen}} \underset{\text{alcohol}}{CH_3OH} \xrightarrow{\text{loss of hydrogen}} \underset{\text{aldehyde}}{CH_2O} \xrightarrow{\text{gain of oxygen}} \underset{\text{carboxylic acid}}{HCOOH} \xrightarrow{\text{loss of hydrogen}} \underset{\text{carbon dioxide}}{CO_2}$$

The opposite process is the reduction of a carboxylic acid to an alkane which involves the loss of oxygen and the gain of hydrogen.

When a primary alcohol is oxidized to an aldehyde in the presence of water, the reaction can be difficult to stop because the aldehyde can be further oxidized to the corresponding carboxylic acid. For example, the oxidation of ethanal produces ethanoic (acetic) acid. Ethanol-containing beverages such as wine are susceptible to such oxidation if kept for long

periods of time after having been opened and exposed to the air. Wine that has become oxidized will have an unpleasant vinegary taste due to the production of acetic acid.



Summary

- Oxidation is the gain of oxygen (O) while reduction is the loss of oxygen (O).
- Oxidation is the loss of hydrogen (H) while reduction is the gain of hydrogen (H).
- Oxidation and reduction reactions must occur together.
- The reactant oxidized is the reducing agent.
- The reactant reduced is the oxidizing agent.

To Your Health: Redox Reactions and Pacemaker Batteries

All batteries use redox reactions to supply electricity because electricity is basically a stream of electrons being transferred from one substance to another. Pacemakers—surgically implanted devices for regulating a person's heartbeat—are powered by tiny batteries, so the proper operation of a pacemaker depends on a redox reaction.

Pacemakers used to be powered by NiCad batteries, in which nickel and cadmium (hence the name of the battery) react with water according to this redox reaction:

$$Cd(s) + 2NiOOH(s) + 2H_2O(\ell) \rightarrow Cd(OH)_2(s) + 2Ni(OH)_2(s)$$

The cadmium is oxidized, while the nickel atoms in NiOOH are reduced. Except for the water, all the substances in this reaction are solids, allowing NiCad batteries to be recharged hundreds of times before they stop operating. Unfortunately, NiCad batteries are fairly heavy batteries to be carrying around in a pacemaker. Today, the lighter lithium/iodine battery is used instead. The iodine is dissolved in a solid polymer support, and the overall redox reaction is as follows:

$$2Li(s) + I_2(s) \rightarrow 2LiI\ (s)$$

Lithium is oxidized, and iodine is reduced. Although the lithium/iodine battery cannot be recharged, one of its advantages is that it lasts up to 10 years. Thus, a person with a pacemaker does not have to worry about periodic recharging; about once per decade a person requires minor surgery to replace the pacemaker/battery unit. Lithium/iodine batteries are also used to power calculators and watches.



Figure 6.6.2: A small button battery like this is used to power a watch, pacemaker, or calculator. Image used with permission from Gerhard H Wrodnigg (via Wikipedia)

Example 6.6.1

In each conversion, indicate whether oxidation or reduction is occurring.

a. $N_2 \rightarrow NH_3$
b. $CH_3CH_2OHCH_3 \rightarrow CH_3COCH_3$
c. $HCHO \rightarrow HCOOH$

SOLUTION

a. Hydrogen is being added to the original reactant molecule, so reduction is occurring.
b. Hydrogen is being removed from the original reactant molecule, so oxidation is occurring.
c. Oxygen is being added to the original reactant molecule, so oxidation is occurring.

Exercise 6.6.1

In each conversion, indicate whether oxidation or reduction is occurring.

a. $CH_4 \rightarrow CO_2 + H_2O$
b. $NO_2 \rightarrow N_2$
c. $CH_2{=}CH_2 \rightarrow CH_3CH_3$

Answer

a. oxidation
b. reduction

c. reduction

Summary

Chemical reactions in which oxygen and/or hydrogen are transferred are called oxidation-reduction, or redox, reactions. Oxidation is the gain of O or loss of H. Reduction is the loss of O or gain of H. Oxidation and reduction always occur together, even though they can be written as separate chemical equations.

Concept Review Exercises

1. Give two different definitions for oxidation and reduction.
2. Give an example of each definition of oxidation and reduction.

Answers

1. Oxidation is the addition of oxygen or the loss of hydrogen; reduction is the loss of oxygen or the addition of hydrogen.
2. $C \rightarrow CO_2$ (oxidation); $C_2H_4 + H_2 \rightarrow C_2H_6$ (reduction) (answers will vary)

Exercises

1. For the following redox reactions, identify the substance oxidized and the substance reduced.
   a. $2C_2H_2 + 5O_2 \rightarrow 4CO_2 + 2H_2O$
   b. $4Al + 3O_2 \rightarrow 2Al_2O_3$
   c. $H_2O_2 + H_2 \rightarrow 2H_2O$
   d. $2CH_3OH + 3O_2 \rightarrow 2CO_2 + 4H_2O$

2. Identify the oxidizing and reducing agent in each redox reaction below.

a. $2C_2H_2 + 5O_2 \rightarrow 4CO_2 + 2H_2O$
b. $4Al + 3O_2 \rightarrow 2Al_2O_3$
c. $H_2O_2 + H_2 \rightarrow 2H_2O$
d. $2CH_3OH + 3O_2 \rightarrow 2CO_2 + 4H_2O$

Answers

1. Look at the reactants, and examine which is the substance oxidized and the substance reduced.

a. substance oxidized:C_2H_2 since it lost H (and gained O) substance reduced: O_2 since it gained H
b. substance oxidized:Al since Al gained O substance reduced: O_2
c. substance oxidized:H_2 since it gained O substance reduced: H_2O_2 since it lost O
d. substance oxidized:CH_3OH since it lost H (and gained O) substance reduced: O_2 since it gained H

2. Identify the oxidizing and reducing agent in each reaction below.

The substance oxidized is the RA while the substance reduced is the OA.

a. RA: C_2H_2 OA: O_2
b. RA: Al OA: O_2
c. RA: H_2 OA: H_2O_2
d. RA: CH_3OH OA: O_2

6.6: Oxidation-Reduction (Redox) Reactions is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

6.7: Redox Reactions in Organic Chemistry and Biochemistry

Skills to Develop

- To identify oxidation-reduction reactions with organic compounds.

Oxidation-reduction reactions are of central importance in organic chemistry and biochemistry. The burning of fuels that provides the energy to maintain our civilization and the metabolism of foods that furnish the energy that keeps us alive both involve redox reactions.



Figure 6.7.1: The Burning of Natural Gas. The burning of natural gas is not only a combustion reaction but also a redox reaction. Similar reactions include the burning of gasoline and coal. These are also redox reactions. Image used with permission from Wikipedia.

All combustion reactions are also redox reactions. A typical combustion reaction is the burning of methane, the principal component of natural gas (Figure 6.7.1).

$$\mathrm{CH_4 + 2\,O_2 \rightarrow CO_2 + 2\,H_2O} \tag{6.7.1}$$

In respiration, the biochemical process by which the oxygen we inhale in air oxidizes foodstuffs to carbon dioxide and water, redox reactions provide energy to living cells. A typical respiratory reaction is the oxidation of glucose ($\mathrm{C_6H_{12}O_6}$), the simple sugar we encountered in the chapter-opening essay that makes up the diet of yeast:

$$\mathrm{C_6H_{12}O_6 + 6\,O_2 \rightarrow 6\,CO_2 + 6\,H_2O} \tag{6.7.2}$$

Organic chemists use a variety of redox reactions. For example, potassium dichromate ($\mathrm{K_2Cr_2O_7}$) is a common oxidizing agent that can be used to oxidize alcohols (symbolized by the general formula ROH). The product of the reaction depends on the location of the OH functional group in the alcohol molecule, the relative proportions of alcohol and the dichromate ion, and reaction conditions such as temperature. If the OH group is attached to a terminal carbon atom and the product is distilled off as it forms, the product is an aldehyde, which has a terminal *carbonyl group* (C=O) and is often written as RCHO. One example is the reaction used by the Breathalyzer to detect ethyl alcohol ($\mathrm{CH_3CH_2OH}$) in a person's breath:

$$\mathrm{3\,CH_3CH_2OH + Cr_2O_7^{2-} + 8\,H^+ \rightarrow 3\,CH_3CHO + 2\,Cr^{3+} + 7\,H_2O} \tag{6.7.3}$$

If the product acetaldehyde ($\mathrm{CH_3CHO}$) is not removed as it forms, it is further oxidized to acetic acid ($\mathrm{CH_3COOH}$). In this case, the overall reaction is as follows:

$$\mathrm{3\,CH_3CH_2OH + 2\,Cr_2O_7^{2-} + 16\,H^+ \rightarrow 3\,CH_3COOH + 4\,Cr^{3+} + 11\,H_2O} \tag{6.7.4}$$

In this reaction, the chromium atom is reduced from $Cr_2O_7^{2-}$ to Cr^{3+}, and the ethanol is oxidized to acetic acid.

When the OH group of the alcohol is bonded to an interior carbon atom, the oxidation of an alcohol will produce a ketone (the formulas of ketones are often written as RCOR, and the carbon–oxygen bond is a double bond). The simplest ketone is derived from 2-propanol ($\mathrm{CH_3CHOHCH_3}$). It is the common solvent acetone [$\mathrm{(CH_3)_2CO}$], which is used in varnishes, lacquers, rubber cement, and nail polish remover. Acetone can be formed by the following redox reaction:

$$\mathrm{3\,CH_3CHOHCH_3 + Cr_2O_7^{2-} + 8\,H^+ \rightarrow 3\,(CH_3)_2CO + 2\,Cr^{3+} + 7\,H_2O} \tag{6.7.5}$$

As we have just seen, aldehydes and ketones can be formed by the oxidation of alcohols. Conversely, aldehydes and ketones can be reduced to alcohols. Reduction of the carbonyl group is important in living organisms. For example, in anaerobic metabolism, in

which biochemical processes take place in the absence of oxygen, pyruvic acid ($CH_3COCOOH$) is reduced to lactic acid ($CH_3CHOHCOOH$) in the muscles.

$$CH_3COCOOH \rightarrow CH_3CHOHCOOH \tag{6.7.6}$$

(Pyruvic acid is both a carboxylic acid and a ketone; only the ketone group is reduced.) The buildup of lactic acid during vigorous exercise is responsible in large part for the fatigue that we experience.

In food chemistry, the substances known as antioxidants are reducing agents. Ascorbic acid (vitamin C; $C_6H_8O_6$) is thought to retard potentially damaging oxidation of living cells. In the process, it is oxidized to dehydroascorbic acid ($C_6H_6O_6$). In the stomach, ascorbic acid reduces the nitrite ion (NO_2^-) to nitric oxide (NO):

$$C_6H_8O_6 + 2H^+ + 2NO_2^- \rightarrow C_6H_6O_6 + 2H_2O + 2NO \tag{6.7.7}$$

> *If reaction in Equation 6.7.7 did not occur, nitrite ions from foods would oxidize the iron in hemoglobin, destroying its ability to carry oxygen.*

Tocopherol (vitamin E) is also an antioxidant. In the body, vitamin E is thought to act by scavenging harmful by-products of metabolism, such as the highly reactive molecular fragments called free radicals. In foods, vitamin E acts to prevent fats from being oxidized and thus becoming rancid. Vitamin C is also a good antioxidant (Figure **6.7.2**).



Figure **6.7.2**: Citrus Fruits. Citrus fruits, such as oranges, lemons, and limes, are good sources of vitamin C, which is an antioxidant. Wedges of pink grapefruit, lime, and lemon, and a half orange (clockwise from top). Image used with permission from Wikipedia.

Finally, and of greatest importance, green plants carry out the redox reaction that makes possible almost all life on Earth. They do this through a process called photosynthesis, in which carbon dioxide and water are converted to glucose ($C_6H_{12}O_6$). The synthesis of glucose requires a variety of proteins called enzymes and a green pigment called chlorophyll that converts sunlight into chemical energy (Figure **6.7.3**). The overall change that occurs is as follows:

$$6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2 \tag{6.7.8}$$

In this reaction, carbon dioxide is reduced to glucose, and water is oxidized to oxygen gas. Other reactions convert the glucose to more complex carbohydrates, plant proteins, and oils.



Figure **6.7.3**: Life on Earth. Photosynthesis is the fundamental process by which plants use sunlight to convert carbon dioxide and water into glucose and oxygen. Then plants make more complex carbohydrates. It is the ultimate source of all food on Earth, and it is a redox reaction. Image used with permission (Public Domain; Wikipedia).

Summary

Redox reactions are common in organic and biological chemistry, including the combustion of organic chemicals, respiration, and photosynthesis.

Concept Review Exercise

1. Give some biochemical examples of oxidation and reduction reactions.

Answer

1. photosynthesis and antioxidants in foods (answers will vary)

Exercises

1. A typical respiratory reaction discussed in the text is the oxidation of glucose ($C_6H_{12}O_6$):

$$C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$$

   Is this a redox reaction? If so, what are the oxidizing and reducing agents?

2. The major net reaction in photosynthesis is as follows:

$$6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$$

   Is this a redox reaction? If so, what are the oxidizing and reducing agents?

3. What would be the ultimate organic product if $CH_3CH_2CH_2OH$ were to react with a solution of $K_2Cr_2O_7$?
4. What would be the final organic product if $CH_3CH_2CHOHCH_3$ were to react with a solution of $K_2Cr_2O_7$?
5. What would be the major organic product if $CH_3CH_2CHOHCH_2CH_3$ were to react with a solution of $K_2Cr_2O_7$?
6. What alcohol is produced in the reduction of acetone [$(CH_3)_2CO$]?
7. What alcohol is produced in the reduction of propanal (CH_3CH_2CHO)?

Answers

1. yes; oxidizing agent: O_2; reducing agent: $C_6H_{12}O_6$

2. yes; OA = CO_2; RA= H_2O

3. CH_3CH_2COOH

4. $CH_3CH_2C(O)CH_3$, where the carbon is double bonded to the oxygen

5. $CH_3CH_2C(O)CH_2CH_3$, where the carbon is double bonded to the oxygen

6. $CH_3CHOHCH_3$, or isopropyl alcohol

7. $CH_3CH_2CH_2OH$, or n-propyl alcohol

6.7: Redox Reactions in Organic Chemistry and Biochemistry is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

6.E: Introduction to Chemical Reactions (Exercises)

5.1: The Law of Conservation of Matter

1. What is the law of conservation of matter?
2. How does the law of conservation of matter apply to chemistry?

ANSWERS

1. The law of conservation of matter states that in any given system that is closed to the transfer of matter, the amount of matter in the system stays constant
2. The law of conservation of matter says that in chemical reactions, the total mass of the products must equal the total mass of the reactants.

Exercises

1. Express the law of conservation of matter in your own words.

2. Explain why the concept of conservation of matter is considered a scientific law.

3. Potassium hydroxide ($\mathbf{KOH}$) readily reacts with carbon dioxide ($\mathbf{CO_2}$) to produce potassium carbonate ($\mathbf{K_2CO_3}$) and water ($\mathbf{H_2O}$). How many grams of potassium carbonate is produced if 224.4 g of $\mathbf{KOH}$ reacted with 88.0 g of $\mathbf{CO_2}$. The reaction also produced 36.0 g of water.

Answers

1. Matter may not be created or destroyed.

2. The concept is a scientific law because it is based on experimentation.

3. 276.4 g

5.2: Chemical Equations

Concept Review Exercises

a. What are the parts of a chemical equation?
b. Explain why chemical equations need to be balanced.

Answers

a. reactants and products

b. Chemical equations need to be balanced to satisfy the law of conservation of matter.

Exercises

1. Write a chemical equation to express the fact that hydrogen gas and solid iodine react to make gaseous hydrogen iodide. Make sure the equation satisfies the law of conservation of matter.
2. Write a chemical equation to express the fact that sodium metal and chlorine gas react to make solid sodium chloride. Make sure the equation satisfies the law of conservation of matter.
3. Write an equation expressing the fact that hydrogen gas and fluorine gas react to make gaseous hydrogen fluoride. Make sure the equation satisfies the law of conservation of matter.
4. Write an equation expressing the fact that solid potassium and fluorine gas react to make solid potassium fluoride. Make sure the equation satisfies the law of conservation of matter.
5. Mercury reacts with oxygen to make mercury(II) oxide. Write a balanced chemical equation that summarizes this reaction.

6. Octane (C_8H_{18}) reacts with oxygen to make carbon dioxide and water. Write a balanced chemical equation that summarizes this reaction.
7. Propyl alcohol (C_3H_7OH) reacts with oxygen to make carbon dioxide and water. Write a balanced chemical equation that summarizes this reaction.
8. Sulfuric acid (H_2SO_4) reacts with iron metal to make iron(III) sulfate and hydrogen gas. Write a balanced chemical equation that summarizes this reaction.
9. Balance each equation.
    a. $MgCl_2 + K \rightarrow KCl + Mg$
    b. $C_6H_{12}O_6 + O_2 \rightarrow CO_2 + H_2O$
    c. $NaN_3 \rightarrow Na + N_2$ (This is the reaction used to inflate airbags in cars.)
10. Balance each equation.
    a. $NH_4NO_3 \rightarrow N_2O + H_2O$
    b. $TiBr_4 + H_2O \rightarrow TiO_2 + HBr$
    c. $C_3H_5N_3O_9 \rightarrow CO_2 + N_2 + O_2 + H_2O$ (This reaction represents the decomposition of nitroglycerine.)
11. Balance each equation.
    a. $NH_3 + O_2 \rightarrow NO + H_2O$
    b. $Li + N_2 \rightarrow Li_3N$
    c. $AuCl \rightarrow Au + AuCl_3$
12. Balance each equation.
    a. $NaOH + H_3PO_4 \rightarrow Na_3PO_4 + H_2O$
    b. $N_2H_4 + Cl_2 \rightarrow N_2 + HCl$
    c. $Na_2S + H_2S \rightarrow NaSH$
13. Chromium(III) oxide reacts with carbon tetrachloride to make chromium(III) chloride and phosgene ($COCl_2$). Write the balanced chemical equation for this reaction.
14. The reaction that occurs when an Alka-Seltzer tablet is dropped into a glass of water has sodium bicarbonate reacting with citric acid ($H_3C_6H_5O_7$) to make carbon dioxide, water, and sodium citrate ($Na_3C_6H_5O_7$). Write the balanced chemical equation for this reaction.
15. When sodium hydrogen carbonate is used to extinguish a kitchen fire, it decomposes into sodium carbonate, carbon dioxide and water. Write a balanced chemical equation for this reaction.
16. Elemental bromine gas can be generated by reacting sodium bromide with elemental chlorine. The other product is sodium chloride. Write a balanced chemical equation for this reaction.

Answers

1. $H_2(g) + I_2(s) \rightarrow 2HI(g)$
2. $2Na(s) + Cl_2(g) \rightarrow 2NaCl(s)$
3. $H_2(g) + F_2(g) \rightarrow 2HF(g)$
4. $2K(s) + F_2(g) \rightarrow 2KF(s)$
5. $2Hg + O_2 \rightarrow 2HgO$
6. $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$
7. $2C_3H_7OH + 9O_2 \rightarrow 6CO_2 + 8H_2O$
8. $3H_2SO_4 + 2Fe \rightarrow Fe_2(SO_4)_3 + 3H_2$
9. a. $MgCl_2 + 2K \rightarrow 2KCl + Mg$

b. $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$
c. $2NaN_3 \rightarrow 2Na + 3N_2$

10.

a. $NH_4NO_3 \rightarrow N_2O + 2H_2O$
b. $TiBr_4 + 2H_2O \rightarrow TiO_2 + 4HBr$
c. $4C_3H_5N_3O_9 \rightarrow 12CO_2 + 6N_2 + O_2 + 10H_2O$

11.

a. $4NH_3 + 5O_2 \rightarrow 4NO + 6H_2O$
b. $6Li + N_2 \rightarrow 2Li_3N$
c. $3AuCl \rightarrow 2Au + AuCl_3$

12.

a. $3NaOH + H_3PO_4 \rightarrow Na_3PO_4 + 3H_2O$
b. $N_2H_4 + 2Cl_2 \rightarrow N_2 + 4HCl$
c. $Na_2S + H_2S \rightarrow 2NaSH$

13. $Cr_2O_3 + 3CCl_4 \rightarrow 2CrCl_3 + 3COCl_2$

14. $3NaHCO_3 + H_3C_6H_5O_7 \rightarrow 3CO_2 + 3H_2O + Na_3C_6H_5O_7$

15. $2NaHCO_3 \rightarrow Na_2CO_3 + CO_2 + H_2O$

16. $2NaBr + Cl_2 \rightarrow Br_2 + 2NaCl$

5.3: Quantitative Relationships Based on Chemical Equations

Concept Review Exercises

1. Explain how stoichiometric ratios are constructed from a chemical equation.
2. Why is it necessary for a chemical equation to be balanced before it can be used to construct conversion factors?

Answers

1. Stoichiometric ratios are made using the coefficients of the substances in the balanced chemical equation.
2. A balanced chemical equation is necessary so one can construct the proper stoichiometric ratios.

Exercises

1. Balance this equation and write every stoichiometric ratio you can from it.

$$NH_4NO_3 \rightarrow N_2O + H_2O$$

2. Balance this equation.

$$N_2 + H_2 \rightarrow NH_3$$

3. Balance this equation and write every stoichiometric ratio you can from it.

$$Fe_2O_3 + C \rightarrow Fe + CO_2$$

4. Balance this equation.

$$Fe_2O_3 + CO \rightarrow Fe + CO_2$$

5. Balance this equation and determine how many molecules of CO_2 are formed if 15 molecules of C_6H_6 are reacted.

$$C_6H_6 + O_2 \rightarrow CO_2 + H_2O$$

6. Balance this equation and determine how many formula units of $Ag_2CO_3(s)$ are produced if 20 formula units of Na_2CO_3 are reacted.

$$Na_2CO_3(aq) + AgNO_3(aq) \rightarrow NaNO_3(aq) + Ag_2CO_3(s)$$

7. Copper metal reacts with nitric acid according to this equation:

$$3Cu(s) + 8HNO_3(aq) \rightarrow 3Cu(NO_3)_2(aq) + 2NO(g) + 4H_2O(\ell)$$

   a. Verify that this equation is balanced.
   b. How many Cu atoms will react if 488 molecules of aqueous HNO_3 are reacted?

8. Gold metal reacts with a combination of nitric acid and hydrochloric acid according to this equation:

$$Au(s) + 3HNO_3(aq) + 4HCl(aq) \rightarrow HAuCl_4(aq) + 3NO_2(g) + 3H_2O(\ell)$$

   a. Verify that this equation is balanced.
   b. How many Au atoms react with 639 molecules of aqueous HNO_3?

9. Sulfur can be formed by reacting sulfur dioxide with hydrogen sulfide at high temperatures according to this equation:

$$SO_2(g) + 2H_2S(g) \rightarrow 3S(g) + 2H_2O(g)$$

   a. Verify that this equation is balanced.
   b. How many S atoms will be formed from by reacting 1,078 molecules of H_2S?

10. Nitric acid is made by reacting nitrogen dioxide with water:

$$3NO_2(g) + H_2O(\ell) \rightarrow 2HNO_3(aq) + NO(g)$$

   a. Verify that this equation is balanced.
   b. How many molecules of NO will be formed by reacting 2,268 molecules of NO_2?

Answers

1. $NH_4NO_3 \rightarrow N_2O + 2H_2O$; the stoichiometric ratios are $\frac{1NH_4NO_3}{1N_2O}$, $\frac{1NH_4NO_3}{2H_2O}$, $\frac{1N_2O}{2H_2O}$, and their reciprocals.
2. $N_2 + 3H_2 \rightarrow 2NH_3$
3. $2Fe_2O_3 + 3C \rightarrow 4Fe + 3CO_2$; the stoichiometric ratios are $\frac{2Fe_2O_3}{3C}$, $\frac{2Fe_2O_3}{4Fe}$, $\frac{2Fe_2O_3}{3CO_2}$, $\frac{3C}{4Fe}$, $\frac{3C}{3CO_2}$, $\frac{4Fe}{3CO_2}$, and their reciprocals.
4. $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$
5. $2C_6H_6 + 15O_2 \rightarrow 12CO_2 + 6H_2O$; 90 molecules
6. $Na_2CO_3(aq) + 2AgNO_3(aq) \rightarrow 2NaNO_3(aq) + Ag_2CO_3(s)$; 20 formula units
7.
   a. It is balanced.
   b. 183 atoms
8.
   a. It is balanced.
   b. 213 atoms
9.
   a. It is balanced.
   b. 1,617 atoms
10.
   a. It is balanced.
   b. 756 molecules

5.4: Some Types of Chemical Reactions

Concept Review Exercises

1. What is the difference between a combination reaction and a combustion reaction?
2. Give the distinguishing characteristic(s) of a decomposition reaction.
3. How do we recognize a combustion reaction?

Answers

1. A combination reaction produces a certain substance; a combustion reaction is a vigorous reaction, usually a combination with oxygen, that is accompanied by the production of light and/or heat.
2. In a decomposition reaction, a single substance reacts to make multiple substances as products.
3. A combustion reaction is typically a vigorous reaction accompanied by light and/or heat, usually because of reaction with oxygen.

Exercises

1. Identify each type of reaction.

    a. $C_6H_5CH_3 + 9O_2 \rightarrow 7CO_2 + 4H_2O$
    b. $2NaHCO_3 \rightarrow Na_2CO_3 + H_2O + CO_2$
    c. $C + 2H_2 \rightarrow CH_4$

2. Identify each type of reaction.

    a. $P_4O_{10} + 6H_2O \rightarrow 4H_3PO_4$
    b. $FeO + SO_3 \rightarrow FeSO_4$
    c. $CaCO_3(s) \rightarrow CO_2(g) + CaO(s)$

3. Identify each type of reaction.

    a. $2NH_4NO_3(s) \rightarrow 2N_2(g) + 4H_2O(g) + O_2(g)$
    b. $Hg(\ell) + ½O_2\ (g) \rightarrow HgO(s)$
    c. $CH_2CH_2(g) + Br_2(\ell) \rightarrow CH_2BrCH_2Br$

4. Identify each type of reaction.

    a. $Ti(s) + O_2(g) \rightarrow TiO_2(s)$
    b. $H_2SO_3(aq) \rightarrow H_2O(\ell) + SO_2(g)$
    c. $3O_2(g) \rightarrow 2O_3(g)$

Answers

1.
    a. combustion
    b. decomposition
    c. combination
2.
    a. combination
    b. combination
    c. decomposition
3.
    a. decomposition
    b. combustion (also combination)
    c. combination

4.

a. combination

b. decomposition

c. combination

5.5: Oxidation-Reduction (Redox) Reactions

Concept Review Exercises

1. Give two different definitions for oxidation and reduction.
2. Give an example of each definition of oxidation and reduction.

Answers

1. Oxidation is the loss of electrons or the addition of oxygen; reduction is the gain of electrons or the addition of hydrogen.
2. $Zn \rightarrow Zn^{2+} + 2e^-$ (oxidation); $C_2H_4 + H_2 \rightarrow C_2H_6$ (reduction) (answers will vary)

Exercises

1. Which reactions are redox reactions? For those that are redox reactions, identify the oxidizing and reducing agents.

   a. $NaOH + HCl \rightarrow H_2O + NaCl$
   b. $3Mg + 2AlCl_3 \rightarrow 2Al + 3MgCl_2$
   c. $H_2O_2 + H_2 \rightarrow 2H_2O$
   d. $KCl + AgNO_3 \rightarrow AgCl + KNO_3$

2. Which reactions are redox reactions? For those that are redox reactions, identify the oxidizing and reducing agents.

   a. $3Cu + 8HNO_3 \rightarrow 3Cu(NO_3)_2 + 2NO + 4H_2O$
   b. $2C_2H_6 + 7O_2 \rightarrow 4CO_2 + 6H_2O$
   c. $2NaHCO_3 \rightarrow Na_2CO_3 + CO_2 + H_2O$
   d. $2K + 2H_2O \rightarrow 2KOH + H_2$

3. Balance each redox reaction by writing appropriate half reactions and combining them to cancel the electrons.

   a. $Ca(s) + H^+(aq) \rightarrow Ca^{2+}(aq) + H_2(g)$
   b. $I^-(aq) + Br_2(\ell) \rightarrow Br^-(aq) + I_2(s)$

4. Balance each redox reaction by writing appropriate half reactions and combining them to cancel the electrons.

   a. $Fe(s) + Sn^{4+}(aq) \rightarrow Fe^{3+}(aq) + Sn^{2+}(aq)$
   b. $Pb(s) + Pb^{4+}(aq) \rightarrow Pb^{2+}(aq)$ (Hint: both half reactions will yield the same product.)

Answers

1.

   a. no

   b. yes; oxidizing agent: $AlCl_3$; reducing agent: Mg

   c. yes; oxidizing agent: H_2O_2; reducing agent: H_2

   d. no

2.

   a. yes; oxidizing agent: HNO_3; reducing agent: Cu
   b. yes; oxidizing agent: O_2; reducing agent: C_2H_6
   c. no

d. yes; oxidizing agent: H_2O; reducing agent: K

3.

a. Combined: $Ca + 2H^+ \rightarrow Ca^{2+} + H_2$
b. Combined: $Br_2 + 2I^- \rightarrow 2Br^- + I_2$

4.

a. $(Fe \rightarrow Fe^{3+} + 3e^-) \times 2$

$(Sn^{4+} + 2e^- \rightarrow Sn^{2+}) \times 3$

Combined: $2Fe + 3Sn^{4+} \rightarrow 2Fe^{3+} + 3Sn^{2+}$

b. $Pb \rightarrow Pb^{2+} + 2e^-$

$Pb^{4+} + 2e^- \rightarrow Pb^{2+}$

Combined: $Pb + Pb^{4+} \rightarrow 2Pb^{2+}$

5.6: Redox Reactions in Organic Chemistry and Biochemistry

Concept Review Exercise

1. Give some biochemical examples of oxidation and reduction reactions.

Answer

1. photosynthesis and antioxidants in foods (answers will vary)

Exercises

1. A typical respiratory reaction discussed in the text is the oxidation of glucose ($C_6H_{12}O_6$):

$$C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$$

Is this a redox reaction? If so, what are the oxidizing and reducing agents?

2. The major net reaction in photosynthesis is as follows:

$$6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$$

Is this a redox reaction? If so, what are the oxidizing and reducing agents?

3. What would be the ultimate organic product if $CH_3CH_2CH_2OH$ were to react with a solution of $K_2Cr_2O_7$?
4. What would be the ultimate organic product if $CH_3CH_2CH_2CH_2OH$ were to react with a solution of $K_2Cr_2O_7$?
5. What would be the final organic product if $CH_3CH_2CHOHCH_3$ were to react with a solution of $K_2Cr_2O_7$?
6. What would be the major organic product if $CH_3CH_2CHOHCH_2CH_3$ were to react with a solution of $K_2Cr_2O_7$?
7. What alcohol is produced in the reduction of acetone $[(CH_3)_2CO]$?
8. What alcohol is produced in the reduction of propanal (CH_3CH_2CHO)?

Answers

1. yes; oxidizing agent: O_2; reducing agent: $C_6H_{12}O_6$
2. yes; oxidizing agent: CO_2; reducing agent: H_2O
3. CH_3CH_2COOH
4. $CH_3CH_2CH_2COOH$
5. $CH_3CH_2C(O)CH_3$, where the carbon is double bonded to the oxygen
6. $CH_3CH_2C(O)CH_2CH_3$, carbon #3 is double bonded to the oxygen

7. $CH_3CHOHCH_3$, or isopropyl alcohol

8. $CH_3CH_2CH_2OH$

5.

Additional Exercises

1. Isooctane (C_8H_{18}) is used as a standard for comparing gasoline performance. Write a balanced chemical equation for the combustion of isooctane.

2. Heptane (C_7H_{16}), like isooctane (see Exercise 1), is also used as a standard for determining gasoline performance. Write a balanced chemical equation for the combustion of heptane.

3. What is the difference between a combination reaction and a redox reaction? Are all combination reactions also redox reactions? Are all redox reactions also combination reactions?

4. Are combustion reactions always redox reactions as well? Explain.

5. A friend argues that the equation

$$Fe^{2+} + Na \rightarrow Fe + Na^{+}$$

is balanced because each side has one iron atom and one sodium atom. Explain why your friend is incorrect.

6. Some antacids contain aluminum hydroxide [$Al(OH)_3$]. This compound reacts with excess hydrochloric acid (HCl) in the stomach to neutralize it. If the products of this reaction are water and aluminum chloride, what is the balanced chemical equation for this reaction? What is the stoichiometric ratio between the number of HCl molecules made to the number of H_2O molecules made?

7. Sulfuric acid is made in a three-step process: (1) the combustion of elemental sulfur to produce sulfur dioxide, (2) the continued reaction of sulfur dioxide with oxygen to produce sulfur trioxide, and (3) the reaction of sulfur trioxide with water to make sulfuric acid (H_2SO_4). Write balanced chemical equations for all three reactions.

8. If the products of glucose metabolism are carbon dioxide and water, what is the balanced chemical equation for the overall process? What is the stoichiometric ratio between the number of CO_2 molecules made to the number of H_2O molecules made?

9. Historically, the first true battery was the Leclanché cell, named after its discoverer, Georges Leclanché. It was based on the following reaction:

$$Zn(s) + Cu^{2+}(aq) \rightarrow Zn^{2+}(aq) + Cu(s)$$

Identify what is being oxidized, what is being reduced, and the respective reducing and oxidizing agents.

Answers

1. $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$

2. $C_7H_{16} + 11O_2 \rightarrow 7CO_2 + 8H_2O$

3. A combination reaction makes a new substance from more than one reactant; a redox reaction rearranges electrons. Most (not all) combination reactions are redox reactions. Not all redox reactions are combination reactions.

4. All combustion reactions are redox reactions. In combustion a chemical is combining with oxygen and that chemical is being oxidized. Oxygen, on the other hand, is being reduced.

5. Your friend is incorrect because the number of electrons transferring is not balanced. A balanced equation must not only have the same number of atoms of each element on each side of the equation but must also have the same charge on both sides.

6. $Al(OH)_3 + 3HCl \rightarrow 3H_2O + AlCl_3$; 1:1

7. (1) $S + O_2 \rightarrow SO_2$; (2) $2SO_2 + O_2 \rightarrow 2SO_3$; (3) $SO_3 + H_2O \rightarrow H_2SO_4$

8. $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$; 1:1

9. oxidized and reducing agent: Zn; reduced and oxidizing agent: Cu^{2+}

Additional Exercises

1. Isooctane (C_8H_{18}) is used as a standard for comparing gasoline performance. Write a balanced chemical equation for the combustion of isooctane.
2. Heptane (C_7H_{16}), like isooctane (see Exercise 1), is also used as a standard for determining gasoline performance. Write a balanced chemical equation for the combustion of heptane.
3. What is the difference between a combination reaction and a redox reaction? Are all combination reactions also redox reactions? Are all redox reactions also combination reactions?
4. Are combustion reactions always redox reactions as well? Explain.
5. A friend argues that the equation

   $$Fe^{2+} + Na \rightarrow Fe + Na^{+}$$

   is balanced because each side has one iron atom and one sodium atom. Explain why your friend is incorrect.
6. Some antacids contain aluminum hydroxide [$Al(OH)_3$]. This compound reacts with excess hydrochloric acid (HCl) in the stomach to neutralize it. If the products of this reaction are water and aluminum chloride, what is the balanced chemical equation for this reaction? What is the stoichiometric ratio between the number of HCl molecules made to the number of H_2O molecules made?
7. Sulfuric acid is made in a three-step process: (1) the combustion of elemental sulfur to produce sulfur dioxide, (2) the continued reaction of sulfur dioxide with oxygen to produce sulfur trioxide, and (3) the reaction of sulfur trioxide with water to make sulfuric acid (H_2SO_4). Write balanced chemical equations for all three reactions.
8. If the products of glucose metabolism are carbon dioxide and water, what is the balanced chemical equation for the overall process? What is the stoichiometric ratio between the number of CO_2 molecules made to the number of H_2O molecules made?
9. Historically, the first true battery was the Leclanché cell, named after its discoverer, Georges Leclanché. It was based on the following reaction:

   $$Zn(s) + Cu^{2+}(aq) \rightarrow Zn^{2+}(aq) + Cu(s)$$

   Identify what is being oxidized, what is being reduced, and the respective reducing and oxidizing agents.

Answers

1. $2C_8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O$
2. $C_7H_{16} + 11O_2 \rightarrow 7CO_2 + 8H_2O$
3. A combination reaction makes a new substance from more than one reactant; a redox reaction rearranges electrons. Most (not all) combination reactions are redox reactions. Not all redox reactions are combination reactions.
4. All combustion reactions are redox reactions. In combustion a chemical is combining with oxygen and that chemical is being oxidized. Oxygen, on the other hand, is being reduced.
5. Your friend is incorrect because the number of electrons transferring is not balanced. A balanced equation must not only have the same number of atoms of each element on each side of the equation but must also have the same charge on both sides.
6. $Al(OH)_3 + 3HCl \rightarrow 3H_2O + AlCl_3$; 1:1
7. (1) $S + O_2 \rightarrow SO_2$; (2) $2SO_2 + O_2 \rightarrow 2SO_3$; (3) $SO_3 + H_2O \rightarrow H_2SO_4$
8. $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O$; 1:1
9. oxidized and reducing agent: Zn; reduced and oxidizing agent: Cu^{2+}

6.E: Introduction to Chemical Reactions (Exercises) is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

- **5.E: Introduction to Chemical Reactions (Exercises)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

- **5.1: The Law of Conservation of Matter** by Binod Shrestha is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **5.2: Chemical Equations** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **5.3: Quantitative Relationships Based on Chemical Equations** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **5.4: Some Types of Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **5.5: Oxidation-Reduction (Redox) Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **5.6: Redox Reactions in Organic Chemistry and Biochemistry** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.S: Introduction to Chemical Reactions (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Scientific **laws** are general statements that apply to a wide variety of circumstances. One important law in chemistry is the **law of conservation of matter**, which states that in any closed system, the amount of matter stays constant.

Chemical equations are used to represent **chemical reactions**. **Reactants** change chemically into **products**. The law of conservation of matter requires that a proper chemical equation be **balanced**. **Coefficients** are used to show the relative numbers of reactant and product molecules.

In **stoichiometry**, quantities of reactants and/or products can be related to each other using the balanced chemical equation. The coefficients in a balanced chemical reaction are used to devise the proper ratios that relate the number of molecules of one substance to the number of molecules of another substance.

Chemical reactions can be classified by type. **Combination reactions** (also called **composition reactions**) make a substance from other substances. **Decomposition reactions** break one substance down into multiple substances. **Combustion reactions** combine molecular oxygen with the atoms of another reactant.

Oxidation reactions are reactions in which an atom loses an electron. **Reduction reactions** are reactions in which an atom gains an electron. These two processes always occur together, so they are collectively referred to as **oxidation-reduction** (or **redox**) **reactions**. The species being oxidized it called the **reducing agent**, while the species being reduced is the **oxidizing agent**. Alternate definitions of oxidation and reduction focus on the gain or loss of oxygen atoms, or the loss or gain of hydrogen atoms. Redox reactions are easily balanced if the overall reaction is first separated into **half reactions**, which are individually balanced.

Oxidation-reduction reactions are common in organic and biological chemistry. **Respiration**, the process by which we inhale and metabolize oxygen, is a series of redox reactions. In the absence of oxygen, redox reactions still occur in a process called **anaerobic metabolism**. **Antioxidants** such as ascorbic acid also play a part in the human diet, acting as reducing agents in various biochemical reactions. **Photosynthesis**, the process by which plants convert water and carbon dioxide to glucose, is also based on redox reactions.

6.S: Introduction to Chemical Reactions (Summary) is shared under a CC BY-SA license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

7: Quantities in Chemical Reactions

So far, we have talked about chemical reactions in terms of individual atoms and molecules. Although this works, most of the reactions occurring around us involve much larger amounts of chemicals. Even a tiny sample of a substance will contain millions, billions, or a hundred billion billions of atoms and molecules. How do we compare amounts of substances to each other in chemical terms when it is so difficult to count to a hundred billion billion? Actually, there are ways to do this, which we will explore in this chapter. In doing so, we will increase our understanding of stoichiometry, which is the study of the numerical relationships between the reactants and the products in a balanced chemical reaction.

7.1: Formula Mass
7.2: The Mole
7.3: Atomic and Molar Masses
7.4: Mole-Mass Conversions
7.5: Mole-Mole Relationships in Chemical Reactions
7.6: Energetics of Biochemical Reactions
7.7: Factors Affecting Reaction Rates
7.8: Catalysis
7.9: Equilbria and Le Chatelier's Principle
7.10: Chapter Summary

7: Quantities in Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

7.1: Formula Mass

Learning Objectives

- To determine the formula mass of an ionic compound.

One skill needed in future chapters is the ability to determine the mass of the formula of an ionic compound. This quantity is called the formula mass. The formula mass is obtained by adding the masses of each individual atom in the formula of the compound. Because a proper formula is electrically neutral (with no net electrons gained or lost), the ions can be considered atoms for the purpose of calculating the formula mass.

Let us start by calculating the formula mass of sodium chloride (NaCl). This formula mass is the sum of the atomic masses of one sodium atom and one chlorine atom, which we find from the periodic table; here, we use the masses to two decimal places:

Table shows how to calculate the formula mass of sodium chloride by using their atomic masses.

Na:	22.99 amu
Cl:	+ 35.45 amu
Total:	58.44 amu

To two decimal places, the formula mass of NaCl is 58.44 amu.

When an ionic compound has more than one anion or cation, you must remember to use the proper multiple of the atomic mass for the element in question. For the formula mass of calcium fluoride (CaF_2), we must multiply the mass of the fluorine atom by 2 to account for the two fluorine atoms in the chemical formula:

Table shows how to find the formula mass of calcium fluoride by using their atomic masses.

Ca:	1 × 40.08	40.08 amu
F:	2 × 19.00 =	+ 38.00 amu
Total:		78.08 amu

The formula mass of CaF_2 is 78.08 amu.

For ionic compounds with polyatomic ions, the sum must include the number and mass of each atom in the formula for the polyatomic ion. For example, potassium nitrate (KNO_3) has one potassium atom, one nitrogen atom, and three oxygen atoms:

Table shows how to find the formula mass of potassium nitrate by using their atomic masses.

K:	1 × 39.10	39.10 amu
N:	1 × 14.01	+ 14.01 amu
O:	3 × 16.00 =	+ 48.00 amu
Total:		101.11 amu

The formula mass of KNO_3 is 101.11 amu.

> Potassium nitrate is a key ingredient in gunpowder and has been used clinically as a diuretic.

When a formula contains more than one polyatomic unit in the chemical formula, as in $Ca(NO_3)_2$, do not forget to multiply the atomic mass of every atom inside the parentheses by the subscript outside the parentheses. This is necessary because the subscript refers to the *entire polyatomic ion*. Thus, for $Ca(NO_3)_2$, the subscript 2 implies two complete nitrate ions, so we must sum the masses of two (1 × 2) nitrogen atoms and six (3 × 2) oxygen atoms, along with the mass of a single calcium atom:

Table shows how to find the formula mass of $Ca(NO_3)_2$ by using their atomic masses.

Ca:	1 × 40.08	40.08 amu
N:	2 × 14.01 =	+ 28.02 amu
O:	6 × 16.00 =	+ 96.00 amu
Total:		164.10 amu

The key to calculating the formula mass of an ionic compound is to correctly count each atom in the formula and multiply the atomic masses of its atoms accordingly.

The Periodic Table is found in this link: Periodic Table of Elements with Atomic Mass [pubchem.ncbi.nlm.nih.gov]

Example 7.1.1

Use the atomic masses (**rounded to two decimal places**) from the **Periodic Table** to determine the formula mass for each ionic compound.

a. $FeCl_3$
b. $(NH_4)_3PO_4$

Solution

a.

Table shows how to find the formula mass of $FeCl_3$ by using their atomic masses.

Fe:		55.84 amu
Cl:	3 × 35.45 =	+ 106.35 amu
Total:		162.19 amu

The formula mass of $FeCl_3$ is 162.19 amu.

a. When we distribute the subscript 3 through the parentheses containing the formula for the ammonium ion, we see that we have 3 nitrogen atoms and 12 hydrogen atoms. Thus, we set up the sum as follows:

Table shows how to find the formula mass of $(NH_4)_3PO_4$ by using their atomic masses.

N:	3 × 14.01 =	42.03 amu
H:	12 × 1.01 =	+ 12.12 amu
P:		+ 30.97 amu
O:	4 × 16.00 =	+ 64.00 amu
Total:		149.12 amu

The formula mass for $(NH_4)_3PO_4$ is 149.12 amu.

Exercise 7.1.1

Use the atomic masses (**rounded to two decimal places**) from the **Periodic Table** to determine the formula mass for each ionic compound.

a. TiO_2
b. AgBr
c. $Au(NO_3)_3$

d. $Fe_3(PO_4)_2$

Answer a:

47.87 + 2 (16.00) = 79.87 amu

Answer b:

107.87 + 79.90 = 187.77 amu

Answer c:

196.97 + 3(14.01) + 9(16.00) = 383.00 amu

Answer d:

3(55.84) + 2(30.97) + 8(16.00) = 357.46 amu

To Your Health: Hydrates

Some ionic compounds have water (H_2O) incorporated within their formula unit. These compounds, called *hydrates*, have a characteristic number of water units associated with each formula unit of the compound. Hydrates are solids, not liquids or solutions, despite the water they contain.

To write the chemical formula of a hydrate, write the number of water units per formula unit of compound after its chemical formula. The two chemical formulas are separated by a vertically centered dot. The hydrate of copper(II) sulfate has five water units associated with each formula unit, so it is written as $CuSO_4 \cdot 5H_2O$. The name of this compound is copper(II) sulfate pentahydrate, with the *penta-* prefix indicating the presence of five water units per formula unit of copper(II) sulfate.



Magnesium sulfate heptahydrate. (Public Domain; Chemicalinterest).

Hydrates have various uses in the health industry. Calcium sulfate hemihydrate ($CaSO_4 \cdot \frac{1}{2}H_2O$), known as plaster of Paris, is used to make casts for broken bones. Epsom salt ($MgSO_4 \cdot 7H_2O$) is used as a bathing salt and a laxative. Aluminum chloride hexahydrate is an active ingredient in antiperspirants. The accompanying table lists some useful hydrates.

Table **7.1.1**: Names and Formulas of Some Widely Used Hydrates

Formula	Name	Uses
$AlCl_3 \cdot 6H_2O$	aluminum chloride hexahydrate	antiperspirant
$CaSO_4 \cdot \frac{1}{2}H_2O$	calcium sulfate hemihydrate (plaster of Paris)	casts (for broken bones and castings)
$CaSO_4 \cdot 2H_2O$	calcium sulfate dihydrate (gypsum)	drywall component
$CoCl_2 \cdot 6H_2O$	cobalt(II) chloride hexahydrate	drying agent, humidity indicator
$CuSO_4 \cdot 5H_2O$	copper(II) sulfate pentahydrate	fungicide, algicide, herbicide
$MgSO_4 \cdot 7H_2O$	magnesium sulfate heptahydrate (Epsom salts)	laxative, bathing salt
$Na_2CO_3 \cdot 10H_2O$	sodium carbonate decahydrate (washing soda)	laundry additive/cleaneKEY TAKEAWAY

KEY TAKEAWAY

- Formula masses of ionic compounds can be determined from the masses of the atoms in their formulas.

EXERCISES

1. What is the relationship between atomic mass and formula mass?

2. How are subscripts used to determine a formula mass when more than one polyatomic ion is present in a chemical formula?

3. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Na^+ and Br^-
   b. Mg^{2+} and Br^-
   c. Mg^{2+} and S^{2-}

4. What is the formula mass for the ionic compound formed by each pair of ions?

   a. K^+ and Cl^-
   b. Mg^{2+} and Cl^-
   c. Mg^{2+} and Se^{2-}

5. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Na^+ and N^{3-}
   b. Mg^{2+} and N^{3-}
   c. Al^{3+} and S^{2-}

6. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Li^+ and N^{3-}
   b. Mg^{2+} and P^{3-}
   c. Li^+ and P^{3-}

7. What is the formula mass for each compound?

   a. $FeBr_3$
   b. $FeBr_2$
   c. Au_2S_3
   d. Au_2S

8. What is the formula mass for each compound?

   a. Cr_2O_3
   b. CrO
   c. $PbCl_2$
   d. $PbCl_4$

9. What is the formula mass for each compound?

   a. $Cr(NO_3)_3$
   b. $Fe_3(PO_4)_2$
   c. $CaCrO_4$
   d. $Al(OH)_3$

10. What is the formula mass for each compound?

    a. NH_4NO_3
    b. $K_2Cr_2O_7$
    c. Cu_2CO_3
    d. $NaHCO_3$

11. What is the formula mass for each compound?

    a. $Al(HSO_4)_3$
    b. $Mg(HSO_4)_2$

12. What is the formula mass for each compound?

    a. $Co(HCO_3)_2$
    b. $LiHCO_3$

Answers

1. The formula mass is the sum of the atomic masses of the atoms in the formula.
2. The subscript is distributed throughout the parentheses to determine the total number of atoms in the formula.

3.

    a. 102.89 amu
    b. 184.11 amu
    c. 56.38 amu

4.

    a. 74.55 amu
    b. 95.21 amu
    c. 103.28 amu

5.

    a. 82.98 amu
    b. 100.95 amu
    c. 150.17 amu

6.

    a. 35.01 amu
    b. 134.87 amu
    c. 51.79 amu

7.

    a. 295.55 amu
    b. 215.65 amu
    c. 490.15 amu
    d. 426.01 amu

8.

    a. 152.00 amu
    b. 68.00 amu
    c. 278.10 amu
    d. 349.00 amu

9.

    a. 238.03 amu
    b. 357.49 amu
    c. 156.08 amu
    d. 78.01 amu

10.

    a. 80.06 amu

b. 294.20 amu
c. 187.11 amu
d. 84.01 amu

11.

a. 318.22 amu
b. 218.47 amu

12.

a. 180.97 amu
b. 67.96 amu

7.1: Formula Mass is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **3.5: Formula Mass** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.2: The Mole

Learning Objectives

- To define the mole unit.

Figure **7.2.1** shows that we need 2 hydrogen atoms and 1 oxygen atom to make 1 water molecule. If we want to make 2 water molecules, we will need 4 hydrogen atoms and 2 oxygen atoms. If we want to make 5 molecules of water, we need 10 hydrogen atoms and 5 oxygen atoms. The ratio of atoms we will need to make any number of water molecules is the same: 2 hydrogen atoms to 1 oxygen atom.




Figure **7.2.1** Water Molecules. The ratio of hydrogen atoms to oxygen atoms used to make water molecules is always 2:1, no matter how many water molecules are being made.

One problem we have, however, is that it is extremely difficult, if not impossible, to organize atoms one at a time. As stated in the introduction, we deal with billions of atoms at a time. How can we keep track of so many atoms (and molecules) at a time? We do it by using mass rather than by counting individual atoms.

A hydrogen atom has a mass of approximately 1 u. An oxygen atom has a mass of approximately 16 u. The ratio of the mass of an oxygen atom to the mass of a hydrogen atom is therefore approximately 16:1.

If we have 2 atoms of each element, the ratio of their masses is approximately 32:2, which reduces to 16:1—the same ratio. If we have 12 atoms of each element, the ratio of their total masses is approximately (12 × 16):(12 × 1), or 192:12, which also reduces to 16:1. If we have 100 atoms of each element, the ratio of the masses is approximately 1,600:100, which again reduces to 16:1. As long as we have equal numbers of hydrogen and oxygen atoms, the ratio of the masses will always be 16:1.

The same consistency is seen when ratios of the masses of other elements are compared. For example, the ratio of the masses of silicon atoms to equal numbers of hydrogen atoms is always approximately 28:1, while the ratio of the masses of calcium atoms to equal numbers of lithium atoms is approximately 40:7.

So we have established that the masses of atoms are constant with respect to each other, as long as we have the same number of each type of atom. Consider a more macroscopic example. If a sample contains 40 g of Ca, this sample has the same number of atoms as there are in a sample of 7 g of Li. What we need, then, is a number that represents a convenient quantity of atoms so we can relate macroscopic quantities of substances. Clearly even 12 atoms are too few because atoms themselves are so small. We need a number that represents billions and billions of atoms.

Chemists use the term **mole** to represent a large number of atoms or molecules. Just as a dozen implies 12 things, a **mole** (abbreviated as ***mol***) represents **6.022×10^{23} things**. The number 6.022×10^{23}, called Avogadro's number after the 19th-century chemist Amedeo Avogadro, is the number we use in chemistry to represent macroscopic amounts of atoms and molecules. Thus, if we have 6.022×10^{23} Na atoms, we say we have 1 mol of Na atoms. If we have 2 mol of Na atoms, we have $2 \times (6.022 \times 10^{23})$ Na atoms, or 1.2044×10^{24} Na atoms. Similarly, if we have 0.5 mol of benzene (C_6H_6) molecules, we have $0.5 \times (6.022 \times 10^{23})$ C_6H_6 molecules, or 3.011×10^{23} C_6H_6 molecules.

> A mole represents a very large number! If 1 mol of quarters were stacked in a column, it could stretch back and forth between Earth and the sun *6.8 billion* times.

Notice that we are applying the mole unit to different types of **chemical entities**. The word *mole* represents a number of things—6.022×10^{23} of them—but does not by itself specify what "they" are. The chemical entities can be ***atoms, molecules***, ***formula units*** and ***ions***. This specific information needs to be specified accurately. Most students find this confusing hence, we need to review the composition of elements, covalent and ionic compounds.

Most elements are made up of individual ***atoms,*** such as helium. However, some elements consist of ***molecules***, such as the diatomic elements, nitrogen, hydrogen, oxygen, etc. One mole of He consists of 6.022×10^{23} He ***atoms*** but one mole of nitrogen contains 6.022×10^{23} N_2 ***molecules***. The basic units of covalent (molecular) compounds are ***molecules*** as well. The molecules of "compounds" consist of different kinds of atoms while the molecules of "elements" consist of only one type of atom. For example, the molecules of ammonia (NH_3) consist of nitrogen and hydrogen atoms while N_2 molecules have N atoms only. Compounds that are ionic, like NaCl, are represented by ionic formulas. One mole of NaCl, for example, refers to 6.022×10^{23} ***formula units*** of NaCl. And, one formula unit of NaCl consists of one sodium ***ion*** and one chloride ***ion***. Figure 6.1.2 summarizes the basic units of elements, covalent and ionic compounds




Figure **7.2.2**: The basic units of elements (atoms or molecules), covalent compounds (molecules) and ionic compounds (formula units of ions).

Conversion Between Moles and Atoms, Molecules and Ions

Using our unit conversion techniques learned in Chapter 1, we can use the mole relationship and the chemical formula to convert back and forth between the moles and the number of chemical entities (atoms, molecules or ions).

Because 1 N_2 molecule contains 2 N atoms, 1 mol of N_2 molecules (6.022×10^{23} molecules) has 2 mol of N atoms. Using formulas to indicate how many atoms of each element we have in a substance, we can relate the number of moles of molecules to the number of moles of atoms. For example, in 1 mol of ethanol (C_2H_6O), we can construct the following relationships (Table **7.2.1**):

Table **7.2.1**: Molecular Relationships

1 Molecule of C_2H_6O Has	1 Mol of C_2H_6O Has	Molecular Relationships
2 C atoms	2 mol of C atoms	$\frac{2\ \text{mol C atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}}$ or $\frac{1\ \text{mol } C_2H_6O\ \text{molecules}}{2\ \text{mol C atoms}}$
6 H atoms	6 mol of H atoms	$\frac{6\ \text{mol H atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}}$ or $\frac{1\ \text{mol } C_2H_6O\ \text{molecules}}{6\ \text{mol H atoms}}$
1 O atom	1 mol of O atoms	$\frac{1\ \text{mol O atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}}$ or $\frac{1\ \text{mol } C_2H_6O\ \text{molecules}}{1\ \text{mol O atoms}}$

The following example illustrates how we can use these relationships as conversion factors.

Example 7.2.1

If a sample consists of 2.5 mol of ethanol (C_2H_6O), how many moles of carbon atoms, hydrogen atoms, and oxygen atoms does it have?

Solution

Using the relationships in Table **7.2.1**, we apply the appropriate conversion factor for each element:

$$2.5\ \text{mol } C_2H_6O\ \text{molecules} \times \underbrace{\frac{2\ \text{mol C atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}}}_{\text{conversion factor}} = 5.0\ \text{mol C atoms}$$

Note how the unit *mol C_2H_6O molecules* cancels algebraically. Similar equations can be constructed for determining the number of H and O atoms:

$$2.5\ \text{mol } C_2H_6O\ \text{molecules} \times \frac{6\ \text{mol H atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}} = 15\ \text{mol H atoms}$$

$$2.5\ \text{mol } C_2H_6O\ \text{molecules} \times \frac{1\ \text{mol O atoms}}{1\ \text{mol } C_2H_6O\ \text{molecules}} = 2.5\ \text{mol O atoms}$$

Exercise 7.2.1

If a sample contains 6.75 mol of Na_2SO_4, how many moles of sodium atoms, sulfur atoms, and oxygen atoms does it have?

Answer

13.5 mol Na, 6.75 mol S and 27 mol O.

We can use Avogadro's number as a conversion factor, or ratio, in dimensional analysis problems. For example, if we are dealing with element X, the mole relationship is expressed as follows:

$$1\ \text{mol X} = 6.022 \times 10^{23}\ \text{X atoms}$$

We can convert this relationship into two possible conversion factors shown below:

$$\frac{1\ \text{mol X}}{6.022 \times 10^{23}\ \text{X atoms}} \text{ or } \frac{6.022 \times 10^{23}\ \text{X atoms}}{1\ \text{mol X}}$$

If the number of "atoms of element X" is given, we can convert it into "moles of X" by multiplying the given value with the conversion factor at the left. However, if the number of "mol of X" is given, the appropriate conversion factor to use is the one at

the right.

If we are dealing with a molecular compound (such as C_4H_{10}), the mole relationship is expressed as follows:

$$1 \text{ mol } C4H10 = 6.022 \times 10^{23} \text{ } C4H10 \text{ molecules}$$

If working with ionic compounds (such as NaCl), the mole relationship is expressed as follows:


$$1 \text{ mol NaCl} = 6.022 \times 10^{23} \text{ NaCl formula units}$$

Example 7.2.2

How many formula units are present in 2.34 mol of NaCl? How many ions are in 2.34 mol?

Solution

Typically in a problem like this, we start with what we are given and apply the appropriate conversion factor. Here, we are given a quantity of 2.34 mol of NaCl, to which we can apply the definition of a **mole as a conversion factor**:



$$2.34 \text{ mol NaCl} \times \frac{6.022 \times 10^{23} \text{ NaCl units}}{1 \text{ mol NaCl}} = 1.41 \times 10^{24} \text{ NaCl units}$$

Because there are two ions per formula unit, there are

$$1.41 \times 10^{24} \text{ NaCl units} \times \frac{2 \text{ ions}}{\text{NaCl units}} = 2.82 \times 10^{24} \text{ ions}$$

in the sample.

Exercise 7.2.2

How many molecules are present in 16.02 mol of C_4H_{10}? How many atoms are in 16.02 mol?

Answer

9.647 x 10^{24}molecules, 1.351 x 10^{26} atoms.

Key Takeaway

- A mole is 6.022×10^{23} things.

7.2: The Mole is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **6.1: The Mole** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.3: Atomic and Molar Masses

Learning Objectives

- To learn how the masses of moles of atoms and molecules are expressed.

Now that we have introduced the mole and practiced using it as a conversion factor, we ask the obvious question: why is the mole *that particular* number of things? Why is it 6.022×10^{23} and not 1×10^{23} or even 1×10^{20}?

The number in a mole, Avogadro's number, is related to the relative sizes of the **atomic mass unit** and **gram mass units**. Whereas one hydrogen atom has a mass of approximately 1 u, **1 mol of H atoms** has a mass of approximately **1 *gram***. And whereas one sodium atom has an approximate mass of 23 u, 1 mol of Na atoms has an approximate mass of 23 *grams*.

One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units. The numbers in the periodic table that we identified as the atomic masses of the atoms not only tell us the mass of one atom in u but also tell us the mass of 1 mol of atoms in grams.

> One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units.

Example 7.3.1: Moles to Mass Conversion with Elements

What is the mass of each quantity?

a. 1 mol of Al atoms
b. 2 mol of U atoms

Solution

a. One mole of Al atoms has a mass in grams that is numerically equivalent to the atomic mass of aluminum. The periodic table shows that the atomic mass (rounded to two decimal points) of Al is 26.98, so 1 mol of Al atoms has a mass of 26.98 g.
b. According to the periodic table, 1 mol of U has a mass of 238.0 g, so the mass of 2 mol is twice that, or 476.0 g.

Exercise 7.3.1: Moles to Mass Conversion with Elements

What is the mass of each quantity?

a. 1 mol of Au atoms
b. 5 mol of Br atoms

Answer a

197.0 g

Answer b

5 mol Br atoms x 79.90 g/mol = 399.5 g

The mole concept can be extended to masses of formula units and molecules as well. The mass of 1 mol of molecules (or formula units) in grams is numerically equivalent to the mass of one molecule (or formula unit) in atomic mass units. For example, a single molecule of O_2 has a mass of 32.00 u, and 1 mol of O_2 molecules has a mass of 32.00 g. As with atomic mass unit–based masses, to obtain the mass of 1 mol of a substance, we simply sum the masses of the individual atoms in the formula of that substance. The mass of 1 mol of a substance is referred to as its **molar mass**, whether the substance is an element, an ionic compound, or a covalent compound.

Example 7.3.2: Moles to Mass Conversion with Compounds

What is the mass of 1 mol of each substance?

a. NaCl
b. bilirubin ($C_{33}H_{36}N_4O_6$), the principal pigment present in bile (a liver secretion)

Solution a

1. Summing the molar masses of the atoms in the NaCl formula unit gives

Solutions to Example 6.2.2, Summing the molar masses of the atoms in the NaCl formula unit gives

1 Na molar mass:	22.99 g
1 Cl molar mass:	35.45 g
Total:	58.44 g

The mass of 1 mol of NaCl is 58.44 g.

Solution b

Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

Solutions to Example 6.2.2, Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

33 C molar mass:	33 × 12.01 g	396.33 g
36 H molar mass:	36 × 1.01 =	36.36 g
4 N molar mass:	4 × 14.01 =	56.04 g
6 O molar mass:	6 × 16.00 =	96.00 g
Total:		584.73 g

The mass of 1 mol of bilirubin is 584.73 g.

Exercise 7.3.2: Moles to Mass Conversion with Compounds

What is the mass of 1 mol of each substance?

a. barium sulfate ($BaSO_4$), used to take X rays of the gastrointestional tract
b. adenosine ($C_{10}H_{13}N_5O_4$), a component of cell nuclei crucial for cell division

Answer a

233.36 g

Answer b

267.28 g

Be careful when counting atoms. In formulas with polyatomic ions in parentheses, the subscript outside the parentheses is applied to every atom inside the parentheses. For example, the molar mass of $Ba(OH)_2$ requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

The molar mass of Ba(OH)2 requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

1 Ba molar mass:	1 × 137.3 g =	137.3 g
2 O molar mass:	2 × 16.00 g =	32.00 g
2 H molar mass:	2 × 1.01 g =	2.02 g
Total:		171.32 g

Because molar mass is defined as the mass for 1 mol of a substance, we can refer to molar mass as grams per mole (g/mol). The division sign (/) implies "per," and "1" is implied in the denominator. Thus, the molar mass of bilirubin can be expressed as 584.73 g/mol, which is read as "five hundred eighty four point seventy three grams per mole."

Key Takeaway

- The mass of moles of atoms and molecules is expressed in units of grams.

7.3: Atomic and Molar Masses is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **6.2: Atomic and Molar Masses** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.4: Mole-Mass Conversions

Learning Objectives

- To convert quantities between mass units and mole units.

A previous Example stated that the mass of 2 mol of U is twice the molar mass of uranium. Such a straightforward exercise does not require any formal mathematical treatment. Many questions concerning mass are not so straightforward, however, and require some mathematical manipulations.

The simplest type of manipulation using molar mass as a conversion factor is a mole-mass conversion (or its reverse, a mass-mole conversion). In such a conversion, we use the molar mass of a substance as a conversion factor to convert mole units into mass units (or, conversely, mass units into mole units).

We also established that 1 mol of Al has a mass of 26.98 g (Example **7.4.1**). Stated mathematically,

$$1 \text{ mol Al} = 26.98 \text{ g Al}$$

We can divide both sides of this expression by either side to get one of two possible conversion factors:

$$\frac{1 \text{ mol Al}}{26.98 \text{ g Al}} \quad \text{and} \quad \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$


The first conversion factor can be used to convert from mass to moles, and the second converts from moles to mass. Both can be used to solve problems that would be hard to do “by eye.”

Example 7.4.1

What is the mass of 3.987 mol of Al?

Solution

The first step in a conversion problem is to decide what conversion factor to use. Because we are starting with mole units, we want a conversion factor that will cancel the mole unit and introduce the unit for mass in the numerator. Therefore, we should use the $\frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$ conversion factor. We start with the given quantity and multiply by the conversion factor:



$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$

Note that the mol units cancel algebraically. (The quantity 3.987 mol is understood to be in the numerator of a fraction that has 1 in the unwritten denominator.) Canceling and solving gives

$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = 107.6 \text{ g Al}$$

Our final answer is expressed to four significant figures.

Exercise 7.4.1

How many moles are present in 100.0 g of Al? (Hint: you will have to use the other conversion factor we obtained for aluminum.)

Answer

$$100.0\ \text{g Al} \times \frac{1\ \text{mol Al}}{26.98\ \text{g Al}} = 3.706\ \text{mol Al}$$

Conversions like this are possible for any substance, as long as the proper atomic mass, formula mass, or molar mass is known (or can be determined) and expressed in grams per mole. Figure **7.4.1** is a chart for determining what conversion factor is needed, and Figure **7.4.2** is a flow diagram for the steps needed to perform a conversion.




Figure **7.4.1** A Simple Flowchart for Converting between Mass and Moles of a Substance. It takes one mathematical step to convert from moles to mass or from mass to moles.




Figure **7.4.2** A Flowchart Illustrating the Steps in Performing a Unit Conversion. When performing many unit conversions, the same logical steps can be taken.

Example 7.4.2

A biochemist needs 0.00655 mol of bilirubin ($C_{33}H_{36}N_4O_6$) for an experiment. How many grams of bilirubin will that be?

Solution

To convert from moles to mass, we need the molar mass of bilirubin, which we can determine from its chemical formula:

Solutions to Example 6.3.2

33 C molar mass:	33 × 12.01 g =	396.33 g
36 H molar mass:	36 × 1.01 g =	36.36 g
4 N molar mass:	4 × 14.01 g =	56.04 g
6 O molar mass:	6 × 16.00 g =	96.00 g
Total:		584.73 g

The molar mass of bilirubin is 584.73 g. Using the relationship

1 mol bilirubin = 584.73 g bilirubin

we can construct the appropriate conversion factor for determining how many grams there are in 0.00655 mol. Following the steps from Figure **7.4.2**:

$$0.00655\ \text{mol bilirubin} \times \frac{584.73\ \text{g bilirubin}}{\text{mol bilirubin}} = 3.83\ \text{g bilirubin}$$

The mol bilirubin unit cancels. The biochemist needs 3.83 g of bilirubin.

Exercise 7.4.2

A chemist needs 457.8 g of $KMnO_4$ to make a solution. How many moles of $KMnO_4$ is that?

Answer

$$457.8\ \text{g KMnO}_4 \times \frac{1\ \text{mol KMnO}_4}{158.04\ \text{g KMnO}_4} = 2.897\ \text{mol KMnO}_4$$

To Your Health: Minerals

For our bodies to function properly, we need to ingest certain substances from our diets. Among our dietary needs are minerals, the noncarbon elements our body uses for a variety of functions, such developing bone or ensuring proper nerve transmission. The US Department of Agriculture has established some recommendations for the RDIs of various minerals. The accompanying table lists the RDIs for minerals, both in mass and moles, assuming a 2,000-calorie daily diet.

Table **7.4.1**: Essential Minerals and their Composition in Humans

Mineral	Male (age 19–30 y)		Female (age 19–30 y)	
Ca	1,000 mg	0.025 mol	1,000 mg	0.025 mol
Cr	35 µg	6.7×10^{-7} mol	25 µg	4.8×10^{-7} mol
Cu	900 µg	1.4×10^{-5} mol	900 µg	1.4×10^{-5} mol
F	4 mg	2.1×10^{-4} mol	3 mg	1.5×10^{-4} mol
I	150 µg	1.2×10^{-6} mol	150 µg	1.2×10^{-6} mol
Fe	8 mg	1.4×10^{-4} mol	18 mg	3.2×10^{-4} mol
K	3,500 mg	9.0×10^{-2} mol	3,500 mg	9.0×10^{-2} mol
Mg	400 mg	1.6×10^{-2} mol	310 mg	1.3×10^{-2} mol
Mn	2.3 mg	4.2×10^{-5} mol	1.8 mg	3.3×10^{-5} mol
Mo	45 mg	4.7×10^{-7} mol	45 mg	4.7×10^{-7} mol
Na	2,400 mg	1.0×10^{-1} mol	2,400 mg	1.0×10^{-1} mol
P	700 mg	2.3×10^{-2} mol	700 mg	2.3×10^{-2} mol
Se	55 µg	7.0×10^{-7} mol	55 µg	7.0×10^{-7} mol
Zn	11 mg	1.7×10^{-4} mol	8 mg	1.2×10^{-4} mol

Table **7.4.1** illustrates several things. First, the needs of men and women for some minerals are different. The extreme case is for iron; women need over twice as much as men do. In all other cases where there is a different RDI, men need more than women.

Second, the amounts of the various minerals needed on a daily basis vary widely—both on a mass scale and a molar scale. The average person needs 0.1 mol of Na a day, which is about 2.5 g. On the other hand, a person needs only about 25–35 µg of Cr per day, which is under one millionth of a mole. As small as this amount is, a deficiency of chromium in the diet can lead to diabetes-like symptoms or neurological problems, especially in the extremities (hands and feet). For some minerals, the body does not require much to keep itself operating properly.

Although a properly balanced diet will provide all the necessary minerals, some people take dietary supplements. However, too much of a good thing, even minerals, is not good. Exposure to too much chromium, for example, causes a skin irritation, and certain forms of chromium are known to cause cancer (as presented in the movie *Erin Brockovich*).

Key Takeaway

- It is possible to convert between moles of material and mass of material.

7.4: Mole-Mass Conversions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **6.3: Mole-Mass Conversions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.5: Mole-Mole Relationships in Chemical Reactions

Learning Objectives

- To use a balanced chemical reaction to determine molar relationships between the substances.

Previously, you learned to balance chemical equations by comparing the numbers of each type of atom in the reactants and products. The coefficients in front of the chemical formulas represent the numbers of molecules or formula units (depending on the type of substance). Here, we will extend the meaning of the coefficients in a chemical equation.

Consider the simple chemical equation

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

The convention for writing balanced chemical equations is to use the lowest whole-number ratio for the coefficients. However, the equation is balanced as long as the coefficients are in a 2:1:2 ratio. For example, this equation is also balanced if we write it as

$$4\,H_2 + 2\,O_2 \longrightarrow 4\,H_2O$$

The ratio of the coefficients is 4:2:4, which reduces to 2:1:2. The equation is also balanced if we were to write it as

$$22\,H_2 + 11\,O_2 \longrightarrow 22\,H_2O$$

because 22:11:22 also reduces to 2:1:2.

Suppose we want to use larger numbers. Consider the following coefficients:

$$12.044 \times 10^{23}\ H_2 + 6.022 \times 10^{23}\ O_2 \rightarrow 12.044 \times 10^{23}\ H_2O$$


These coefficients also have the ratio 2:1:2 (check it and see), so this equation is balanced. But 6.022×10^{23} is 1 mol, while 12.044×10^{23} is 2 mol (and the number is written that way to make this more obvious), so we can simplify this version of the equation by writing it as

$$2\ \text{mol}\ \ H_2 + 1\ \ \text{mol}\ \ O_2 \longrightarrow 2\ \text{mol}\ \ H_2O$$

We can leave out the word *mol* and not write the 1 coefficient (as is our habit), so the final form of the equation, still balanced, is

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

Now we interpret the coefficients as referring to molar amounts, not individual molecules. The lesson? *Balanced chemical equations are balanced not only at the molecular level but also in terms of molar amounts of reactants and products.* Thus, we can read this reaction as “two moles of hydrogen react with one mole of oxygen to produce two moles of water.”



2 molecules H_2 1 molecule O_2 2 molecules H_2O

2 moles H_2 1 mole O_2 2 moles H_2O

2 x 2.02 g=4.04 g H_2 32.0 g O_2 2 x 18.02 g=36.04 g H_2O

Figure 7.5.1: This representation of the production of water from oxygen and hydrogen show several ways to interpret the quantitative information of a chemical reaction.

By the same token, the ratios we constructed to describe molecules reaction can also be constructed in terms of moles rather than molecules. For the reaction in which hydrogen and oxygen combine to make water, for example, we can construct the following

ratios:

$$\frac{2\ \text{mol}\ H_2}{1\ \text{mol}\ O_2}\ \text{or}\ \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2}$$

$$\frac{2\ \text{mol}\ H_2O}{1\ \text{mol}\ O_2}\ \text{or}\ \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2O}$$

$$\frac{2\ \text{mol}\ H_2}{2\ \text{mol}\ H_2O}\ \text{or}\ \frac{2\ \text{mol}\ H_2O}{2\ \text{mol}\ H_2}$$

We can use these ratios to determine what amount of a substance, in moles, will react with or produce a given number of moles of a different substance. The study of the numerical relationships between the reactants and the products in balanced chemical reactions is called ***stoichiometry*.** The ratio of coefficients in a balanced chemical equation, used in computations relating amounts of reactants and products is called the **stoichiometric factor.**


Example 7.5.1

How many moles of oxygen react with hydrogen to produce 27.6 mol of H_2O? The balanced equation is as follows:

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

Solution

Because we are dealing with quantities of H_2O and O_2, we will use the **stoichiometric ratio** that relates those two substances. Because we are given an amount of H_2O and want to determine an amount of O_2, we will use the ratio that has H_2O in the denominator (so it cancels) and O_2 in the numerator (so it is introduced in the answer). Thus,



$$27.6\ \text{mol}\ H_2O \times \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2O} = 13.8\ \text{mol}\ O_2$$

To produce 27.6 mol of H_2O, 13.8 mol of O_2 react.

Exercise 7.5.1

Using $2H_2 + O_2 \rightarrow 2H_2O$, how many moles of hydrogen react with 3.07 mol of oxygen to produce H_2O?

Answer

$$3.07\ \text{mol}\ O_2 \times \frac{2\ \text{mol}\ H_2}{1\ \text{mol}\ O_2} = 6.14\ \text{mol}\ H_2$$

Key Takeaway

- The balanced chemical reaction can be used to determine molar relationships between substances.

7.5: Mole-Mole Relationships in Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **6.4: Mole-Mole Relationships in Chemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.6: Energetics of Biochemical Reactions

Learning Objectives

- To relate the concept of energy change to chemical reactions that occur in the body.

The chemistry of the human body, or any living organism, is very complex. Even so, the chemical reactions found in the human body follow the same principles of energy that other chemical reactions follow.

Where does the energy that powers our bodies come from? The details are complex, but we can look at some simple processes at the heart of cellular activity.

An important reaction that provides energy for our bodies is the oxidation of glucose ($C_6H_{12}O_6$):

$$C_6H_{12}O_6\,(s) + 6\,O_2\,(g) \rightarrow 6\,CO_2\,(g) + 6\,H_2O(\ell) + 670\ \text{kcal} \tag{7.6.1}$$

Considering that 1 mol of $C_6H_{12}O_6\,(s)$ has a volume of about 115 mL, we can see that glucose is a compact source of energy.

Glucose and other sugars are examples of carbohydrates, which are one of the three main dietary components of a human diet. All carbohydrates supply approximately 4 kcal/g. You can verify that by taking the heat of reaction for glucose oxidation and dividing it by its molar mass. Proteins, the building blocks of structural tissues like muscle and skin, also supply about 4 kcal/g. Other important energy sources for the body are fats, which are largely hydrocarbon chains. Fats provide even more energy per gram, about 9 kcal/g.

Another important reaction is the conversion of adenosine triphosphate (ATP) to adenosine diphosphate (ADP), which is shown in Figure **7.6.1**. Under physiological conditions, the breaking of an O–P bond and the formation of an O–P and two O–H bonds gives off about 7.5 kcal/mol of ATP. This may not seem like much energy, especially compared to the amount of energy given off when glucose reacts. It is enough energy, however, to fuel other biochemically important chemical reactions in our cells.




Figure **7.6.1**: ATP to ADP. The conversion of ATP (top) to ADP (bottom) provides energy for the cells of the body.

Even complex biological reactions must obey the basic rules of chemistry.

Career Focus: Dietitian

A dietitian is a nutrition expert who communicates food-related information to the general public. In doing so, dietitians promote the general well-being among the population and help individuals recover from nutritionally related illnesses.

Our diet does not just supply us with energy. We also get vitamins, minerals, and even water from what we eat. Eating too much, too little, or not enough of the right foods can lead to a variety of problems. Dietitians are trained to make specific dietary recommendations to address particular issues relating to health. For example, a dietitian might work with a person to develop an overall diet that would help that person lose weight or control diabetes. Hospitals employ dietitians in planning

menus for patients, and many dietitians work with community organizations to improve the eating habits of large groups of people.

Key Takeaway

- Energy to power the human body comes from chemical reactions.

7.6: Energetics of Biochemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **7.5: The Energy of Biochemical Reactions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.7: Factors Affecting Reaction Rates

Learning Objectives

- Describe the effects of chemical nature, physical state, temperature, concentration, and catalysis on reaction rates

The rates at which reactants are consumed and products are formed during chemical reactions vary greatly. We can identify five factors that affect the rates of chemical reactions: the chemical nature of the reacting substances, the state of subdivision (one large lump versus many small particles) of the reactants, the temperature of the reactants, the concentration of the reactants, and the presence of a catalyst.

The Chemical Nature of the Reacting Substances

The rate of a reaction depends on the nature of the participating substances. Reactions that appear similar may have different rates under the same conditions, depending on the identity of the reactants. For example, when small pieces of the metals iron and sodium are exposed to air, the sodium reacts completely with air overnight, whereas the iron is barely affected. The active metals calcium and sodium both react with water to form hydrogen gas and a base. Yet calcium reacts at a moderate rate, whereas sodium reacts so rapidly that the reaction is almost explosive.

The State of Subdivision of the Reactants

Except for substances in the gaseous state or in solution, reactions occur at the boundary, or interface, between two phases. Hence, the rate of a reaction between two phases depends to a great extent on the surface contact between them. A finely divided solid has more surface area available for reaction than does one large piece of the same substance. Thus a liquid will react more rapidly with a finely divided solid than with a large piece of the same solid. For example, large pieces of iron react slowly with acids; finely divided iron reacts much more rapidly (Figure **7.7.1**). Large pieces of wood smolder, smaller pieces burn rapidly, and saw dust burns explosively.



Figure **7.7.1**: (a) Iron powder reacts rapidly with dilute hydrochloric acid and produces bubbles of hydrogen gas because the powder has a large total area: 2Fe(s) + 6HCl(aq) ⟶ 2FeCl3(aq) + 3H2(g). (surfaceb) An iron nail reacts more slowly.

This figure shows two photos labeled (a) and (b). Photo (a) shows the bottom of a test tube. The test tube is filled with a dark gas, and there is a dark substance and bubbles in the bottom. Photo (b) shows a rod and bubbles in a test tube similar to photo (a), but the gas in the test tube is not as dark.



Video* 7.7.1: The reaction of cesium with water in slow motion and a discussion of how the state of reactants and particle size affect reaction rates.*

Temperature of the Reactants

Chemical reactions typically occur faster at higher temperatures. Food can spoil quickly when left on the kitchen counter. However, the lower temperature inside of a refrigerator slows that process so that the same food remains fresh for days. We use a burner or a hot plate in the laboratory to increase the speed of reactions that proceed slowly at ordinary temperatures. In many cases, an increase in temperature of only 10 °C will approximately double the rate of a reaction in a homogeneous system.

Concentrations of the Reactants

The rates of many reactions depend on the concentrations of the reactants. Rates usually increase when the concentration of one or more of the reactants increases. For example, calcium carbonate ($\mathbf{CaCO_3}$) deteriorates as a result of its reaction with the pollutant sulfur dioxide. The rate of this reaction depends on the amount of sulfur dioxide in the air (Figure **7.7.2**). As an acidic oxide, sulfur dioxide combines with water vapor in the air to produce sulfurous acid in the following reaction:

$$\mathbf{SO_{2(g)} + H_2O_{(g)} \longrightarrow H_2SO_{3(aq)}} \tag{7.7.1}$$

Calcium carbonate reacts with sulfurous acid as follows:

$$\mathbf{CaCO_{3(s)} + H_2SO_{3(aq)} \longrightarrow CaSO_{3(aq)} + CO_{2(g)} + H_2O_{(l)}} \tag{7.7.2}$$

In a polluted atmosphere where the concentration of sulfur dioxide is high, calcium carbonate deteriorates more rapidly than in less polluted air. Similarly, phosphorus burns much more rapidly in an atmosphere of pure oxygen than in air, which is only about 20% oxygen.




Figure **7.7.2**: Statues made from carbonate compounds such as limestone and marble typically weather slowly over time due to the actions of water, and thermal expansion and contraction. However, pollutants like sulfur dioxide can accelerate weathering. As the concentration of air pollutants increases, deterioration of limestone occurs more rapidly. (credit: James P Fisher III).

A photograph is shown of an angel statue. While some details of the statue, including facial features, are present, effects of weathering appear to be diminishing these features.



***Video* 7.7.2**: *Phosphorous burns rapidly in air, but it will burn even more rapidly if the concentration of oxygen in is higher.*

The Presence of a Catalyst

Hydrogen peroxide solutions foam when poured onto an open wound because substances in the exposed tissues act as catalysts, increasing the rate of hydrogen peroxide's decomposition. However, in the absence of these catalysts (for example, in the bottle in the medicine cabinet) complete decomposition can take months. A catalyst is a substance that increases the rate of a chemical reaction by lowering the activation energy without itself being consumed by the reaction. Activation energy is the minimum amount of energy required for a chemical reaction to proceed in the forward direction. A catalyst increases the reaction rate by providing an alternative pathway or mechanism for the reaction to follow (Figure **7.7.3**). Catalysis will be discussed in greater detail later in this chapter as it relates to mechanisms of reactions.




Figure **7.7.3**: The presence of a catalyst increases the rate of a reaction by lowering its activation energy.

A graph is shown with the label, "Reaction coordinate," on the x-axis. The x-axis is depicted as an arrow. The y-axis is also an arrow and is labeled, "Energy." There is a horizontal line that runs the width of the graph and appears just above the x-axis. A segment of this line is blue and is labeled, "Reactants." From the right end of this line segment, a solid black, concave down curve is shown which reaches the level just below the end of the y-axis. The curve ends at another short, blue line labeled, "Products." The "Products" line appears at a higher level than the "Reactants" line. An arrow extends from the horizontal line to the apex of the curve. The arrow is labeled, "Uncatalyzed reaction activation energy." A second, black concave down curve is shown. This curve also meets the reactants and products blue line segments, but only extends to about two-thirds the height of the initial curve. From the horizontal line is another arrow pointing to the apex of the second curve. This arrow is labeled, "Catalyzed reaction activation energy."

> *Chemical reactions occur when molecules collide with each other and undergo a chemical transformation. Before physically performing a reaction in a laboratory, scientists can use molecular modeling simulations to predict how the parameters discussed earlier will influence the rate of a reaction. Use the PhET Reactions & Rates interactive to explore how temperature, concentration, and the nature of the reactants affect reaction rates.*

Summary

The rate of a chemical reaction is affected by several parameters. Reactions involving two phases proceed more rapidly when there is greater surface area contact. If temperature or reactant concentration is increased, the rate of a given reaction generally increases as well. A catalyst can increase the rate of a reaction by providing an alternative pathway that causes the activation energy of the reaction to decrease.

Glossary

catalyst

substance that increases the rate of a reaction without itself being consumed by the reaction

This page titled 7.7: Factors Affecting Reaction Rates is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **12.3: Factors Affecting Reaction Rates** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/chemistry-2e.

7.8: Catalysis

Learning Objectives

- Explain the function of a catalyst in terms of reaction mechanisms and potential energy diagrams
- List examples of catalysis in natural and industrial processes

We have seen that the rate of many reactions can be accelerated by catalysts. A catalyst speeds up the rate of a reaction by lowering the activation energy; in addition, the catalyst is regenerated in the process. Several reactions that are thermodynamically favorable in the absence of a catalyst only occur at a reasonable rate when a catalyst is present. One such reaction is catalytic hydrogenation, the process by which hydrogen is added across an alkene C=C bond to afford the saturated alkane product. A comparison of the reaction coordinate diagrams (also known as energy diagrams) for catalyzed and uncatalyzed alkene hydrogenation is shown in Figure **7.8.1**.




Figure **7.8.1**: This graph compares the reaction coordinates for catalyzed and uncatalyzed alkene hydrogenation.

A graph is shown with the label, “Reaction coordinate,” on the x-axis and the label,“Energy,” on the y-axis. Approximately half-way up the y-axis, a short portion of a black concave down curve which has a horizontal line extended from it across the graph. The left end of this line is labeled “H subscript 2 C equals C H subscript 2 plus H subscript 2.” The black concave down curve extends upward to reach a maximum near the height of the y-axis. The peak of this curve is labeled, “Transition state.” A double sided arrow extends from the horizontal line to the peak of the curve. This arrow is labeled, “Activation energy of Uncatalyzed reation.” From the peak, the curve continues downward to a second horizontally flattened region well below the origin of the curve near the x-axis. This flattened region is shaded in blue and is labeled “H subscript 3 C dash C H subscript 3.” A double sided arrow is drawn from the lowers part of this curve at the far right of the graph to the line extending across the graph above it. This arrow is labeled, “capital delta H less than 0 : exothermic.” A second curve is drawn with the same flattened regions at the start and end of the curve. The height of this curve is about two-thirds the height of the first curve. A double sided arrow is drawn from the horizontal line that originates at the left side of the graph to the peak of this second curve. This arrow is labeled, “Activation energy of catalyzed reaction.”

Catalysts function by providing an alternate reaction mechanism that has a lower activation energy than would be found in the absence of the catalyst. In some cases, the catalyzed mechanism may include additional steps, as depicted in the reaction diagrams shown in Figure **7.8.2** This lower activation energy results in an increase in rate as described by the Arrhenius equation. Note that a catalyst decreases the activation energy for both the forward and the reverse reactions and hence *accelerates both the forward and the reverse reactions*. Consequently, the presence of a catalyst will permit a system to reach equilibrium more quickly, but it has no effect on the position of the equilibrium as reflected in the value of its equilibrium constant (see the later chapter on chemical equilibrium).





Figure **7.8.2**: This potential energy diagram shows the effect of a catalyst on the activation energy. The catalyst provides a different reaction path with a lower activation energy. As shown, the catalyzed pathway involves a two-step mechanism (note the presence of two transition states) and an intermediate species (represented by the valley between the two transitions states).

A graph is shown with the label, “Extent of reaction,” appearing in a right pointing arrow below the x-axis and the label, “Energy,” in an upward pointing arrow just left of the y-axis. Approximately one-fifth of the way up the y-axis, a very short, somewhat flattened portion of both a red and a blue curve are shown. This region is labeled “Reactants.” A red concave down curve extends upward to reach a maximum near the height of the y-axis. This curve is labeled, “Uncatalyzed pathway.” From the peak, the curve continues downward to a second horizontally flattened region at a height of about one-third the height of the y-axis. This flattened region is labeled, “Products.” A second curve is drawn in blue with the same flattened regions at the start and end of the curve. The height of this curve is about two-thirds the height of the first curve and just right of its maximum, the curve dips low, then rises back and continues a downward trend at a lower height, but similar to that of the red curve. This blue curve is labeled, “Catalyzed pathway.”

Example 7.8.1: Using Reaction Diagrams to Compare Catalyzed Reactions

The two reaction diagrams here represent the same reaction: one without a catalyst and one with a catalyst. Identify which diagram suggests the presence of a catalyst, and determine the activation energy for the catalyzed reaction:



In this figure, two graphs are shown. The x-axes are labeled, “Extent of reaction,” and the y-axes are labeled, “Energy (k J).” The y-axes are marked off from 0 to 50 in intervals of five. In a, a blue curve is shown. It begins with a horizontal segment at about 6. The curve then rises sharply near the middle to reach a maximum of about 32 and similarly falls to another horizontal segment at about 10. In b, the curve begins and ends similarly, but the maximum reached near the center of the graph is only 20.


Solution

A catalyst does not affect the energy of reactant or product, so those aspects of the diagrams can be ignored; they are, as we would expect, identical in that respect. There is, however, a noticeable difference in the transition state, which is distinctly lower in diagram (b) than it is in (a). This indicates the use of a catalyst in diagram (b). The activation energy is the difference between the energy of the starting reagents and the transition state—a maximum on the reaction coordinate diagram. The reagents are at 6 kJ and the transition state is at 20 kJ, so the activation energy can be calculated as follows:

$$E_a = 20\ \text{kJ} - 6\ \text{kJ} = 14\ \text{kJ} \tag{7.8.1}$$

? Exercise 7.8.1

Determine which of the two diagrams here (both for the same reaction) involves a catalyst, and identify the activation energy for the catalyzed reaction:



In this figure, two graphs are shown. The x-axes are labeled, “Extent of reaction,” and the y-axes are labeledc “Energy (k J).” The y-axes are marked off from 0 to 100 at intervals of 10. In a, a blue curve is shown. It begins with a horizontal segment at about 10. The curve then rises sharply near the middle to reach a maximum of about 91, then sharply falls to about 52, again rises sharply to about 73 and falls to another horizontal segment at about 5. In b, the curve begins and ends similarly, but the first peak reaches about 81, drops to about 55, then rises to about 77 before falling to the horizontal region at about 5.

Answer

Diagram (b) is a catalyzed reaction with an activation energy of about 70 kJ.

Homogeneous Catalysts

A homogeneous catalyst is present in the same phase as the reactants. It interacts with a reactant to form an intermediate substance, which then decomposes or reacts with another reactant in one or more steps to regenerate the original catalyst and form product. As an important illustration of homogeneous catalysis, consider the earth’s ozone layer. Ozone in the upper atmosphere, which protects the earth from ultraviolet radiation, is formed when oxygen molecules absorb ultraviolet light and undergo the reaction:

$$3\,O_2(g) \xrightarrow{h\nu} 2\,O_3(g) \tag{7.8.2}$$

Ozone is a relatively unstable molecule that decomposes to yield diatomic oxygen by the reverse of this equation. This decomposition reaction is consistent with the following mechanism:

$$\begin{aligned} O_3 &\longrightarrow O_2 + O \\ O + O_3 &\longrightarrow 2\,O_2 \end{aligned} \tag{7.8.3}$$

The presence of nitric oxide, NO, influences the rate of decomposition of ozone. Nitric oxide acts as a catalyst in the following mechanism:

$$\begin{aligned} NO(g) + O_3(g) &\longrightarrow NO_2(g) + O_2(g) \\ O_3(g) &\longrightarrow O_2(g) + O(g) \\ NO_2(g) + O(g) &\longrightarrow NO(g) + O_2(g) \end{aligned} \tag{7.8.4}$$


The overall chemical change for the catalyzed mechanism is the same as:

$$2\,O_3(g) \longrightarrow 3\,O_2(g) \tag{7.8.5}$$

The nitric oxide reacts and is regenerated in these reactions. It is not permanently used up; thus, it acts as a catalyst. The rate of decomposition of ozone is greater in the presence of nitric oxide because of the catalytic activity of NO. Certain compounds that contain chlorine also catalyze the decomposition of ozone.

Mario J. Molina

The 1995 Nobel Prize in Chemistry was shared by Paul J. Crutzen, Mario J. Molina (Figure 7.8.3), and F. Sherwood Rowland "for their work in atmospheric chemistry, particularly concerning the formation and decomposition of ozone." Molina, a Mexican citizen, carried out the majority of his work at the Massachusetts Institute of Technology (MIT).



(a) (b)

Figure 7.8.3: (a) Mexican chemist Mario Molina (1943 –) shared the Nobel Prize in Chemistry in 1995 for his research on (b) the Antarctic ozone hole. (credit a: courtesy of Mario Molina; credit b: modification of work by NASA)

A photograph is shown of Mario Molina. To the right of the photo, an image of Earth's southern hemisphere is shown with a central circular region in purple with a radius of about half that of the entire hemisphere. Just outside this region is a narrow royal blue band, followed by an outer thin turquoise blue band. The majority of the outermost region is green. Two small bands of yellow are present in the lower regions of the image.

In 1974, Molina and Rowland published a paper in the journal *Nature* (one of the major peer-reviewed publications in the field of science) detailing the threat of chlorofluorocarbon gases to the stability of the ozone layer in earth's upper atmosphere. The ozone layer protects earth from solar radiation by absorbing ultraviolet light. As chemical reactions deplete the amount of ozone in the upper atmosphere, a measurable "hole" forms above Antarctica, and an increase in the amount of solar ultraviolet radiation— strongly linked to the prevalence of skin cancers—reaches earth's surface. The work of Molina and Rowland was instrumental in the adoption of the Montreal Protocol, an international treaty signed in 1987 that successfully began phasing out production of chemicals linked to ozone destruction.

Molina and Rowland demonstrated that chlorine atoms from human-made chemicals can catalyze ozone destruction in a process similar to that by which NO accelerates the depletion of ozone. Chlorine atoms are generated when chlorocarbons or chlorofluorocarbons—once widely used as refrigerants and propellants—are photochemically decomposed by ultraviolet light or react with hydroxyl radicals. A sample mechanism is shown here using methyl chloride:

\[\ce{CH3Cl + OH ⟶ Cl + other\: products} \nonumber \]

Chlorine radicals break down ozone and are regenerated by the following catalytic cycle:

$$\ce{Cl + O3 ⟶ ClO + O2}$$
$$\ce{ClO + O ⟶ Cl + O2}$$
$$\textbf{overall Reaction: } \ce{O3 + O ⟶ 2\,O2}$$

A single monatomic chlorine can break down thousands of ozone molecules. Luckily, the majority of atmospheric chlorine exists as the catalytically inactive forms Cl_2 and $ClONO_2$.

Glucose-6-Phosphate Dehydrogenase Deficiency

Enzymes in the human body act as catalysts for important chemical reactions in cellular metabolism. As such, a deficiency of a particular enzyme can translate to a life-threatening disease. G6PD (glucose-6-phosphate dehydrogenase) deficiency, a genetic condition that results in a shortage of the enzyme glucose-6-phosphate dehydrogenase, is the most common enzyme deficiency in humans. This enzyme, shown in Figure 7.8.4, is the rate-limiting enzyme for the metabolic pathway that supplies NADPH to cells (FIgure 7.8.5).




Figure **7.8.4:** Glucose-6-phosphate dehydrogenase is a rate-limiting enzyme for the metabolic pathway that supplies NADPH to cells.

A colorful model of the Glucose-6-phosphate dehydrogenase structure is shown. The molecule has two distinct lobes which are filled with spiraled ribbon-like regions of yellow, lavender, blue, silver, green, and pink.

A disruption in this pathway can lead to reduced glutathione in red blood cells; once all glutathione is consumed, enzymes and other proteins such as hemoglobin are susceptible to damage. For example, hemoglobin can be metabolized to bilirubin, which leads to jaundice, a condition that can become severe. People who suffer from G6PD deficiency must avoid certain foods and medicines containing chemicals that can trigger damage their glutathione-deficient red blood cells.




Figure **7.8.5**: In the mechanism for the pentose phosphate pathway, G6PD catalyzes the reaction that regulates NAPDH, a co-enzyme that regulates glutathione, an antioxidant that protects red blood cells and other cells from oxidative damage.

A reaction mechanism is diagrammed in this figure. At the left, the name Glucose is followed by a horizontal, right pointing arrow, labeled, “Hexokinase.” Below this arrow and to the left is a yellow star shape labeled, “A T P.” A curved arrow extends from this shape to the right pointing arrow, and down to the right to a small brown oval labeled, “A D P.” To the right of the horizontal arrow is the name Glucose 6 phosphate, which is followed by another horizontal, right pointing arrow which is labeled, “G 6 P D.” A small orange rectangle below and left of this arrow is labeled “N A D P superscript plus.” A curved arrow extends from this shape to the right pointing arrow, and down to the right to a small salmon-colored rectangle labeled “N A P D H.” A curved arrow extends from this shape below and to the left, back to the orange rectangle labeled, “N A D P superscript plus.” Another curved arrow extends from a green oval labeled “G S S G” below the orange rectangle, up to the arrow curving back to the orange rectangle. This last curved arrow continues on to the lower right to a second green oval labeled, “G S H.” The end of this curved arrow is labeled, “Glutathione reductase.” To the right of the rightmost horizontal arrow appears the name 6 phosphogluconate.

Heterogeneous Catalysts

A heterogeneous catalyst is a catalyst that is present in a different phase (usually a solid) than the reactants. Such catalysts generally function by furnishing an active surface upon which a reaction can occur. Gas and liquid phase reactions catalyzed by heterogeneous catalysts occur on the surface of the catalyst rather than within the gas or liquid phase.

Heterogeneous catalysis has at least four steps:

1. Adsorption of the reactant onto the surface of the catalyst
2. Activation of the adsorbed reactant
3. Reaction of the adsorbed reactant
4. Diffusion of the product from the surface into the gas or liquid phase (desorption).

Any one of these steps may be slow and thus may serve as the rate determining step. In general, however, in the presence of the catalyst, the overall rate of the reaction is faster than it would be if the reactants were in the gas or liquid phase.

Figure 7.8.6 illustrates the steps that chemists believe to occur in the reaction of compounds containing a carbon–carbon double bond with hydrogen on a nickel catalyst. Nickel is the catalyst used in the hydrogenation of polyunsaturated fats and oils (which contain several carbon–carbon double bonds) to produce saturated fats and oils (which contain only carbon–carbon single bonds).




Figure 7.8.6: There are four steps in the catalysis of the reaction $C_2H_4 + H_2 \longrightarrow C_2H_6$ *by nickel. (a) Hydrogen is adsorbed on the surface, breaking the H–H bonds and forming Ni–H bonds. (b) Ethylene is adsorbed on the surface, breaking the π-bond and forming Ni–C bonds. (c) Atoms diffuse across the surface and form new C–H bonds when they collide. (d)* C_2H_6 *molecules escape from the nickel surface, since they are not strongly attracted to nickel.*

In this figure, four diagrams labeled a through d are shown. In each, a green square surface is shown in perspective to provide a three-dimensional appearance. In a, the label “N i surface” is placed above with a line segment extending to the green square. At the lower left and upper right, pairs of white spheres bonded tougher together appear as well as white spheres on the green surface. Black arrows are drawn from each of the white spheres above the surface to the white sphere on the green surface. In b, the white spheres are still present on the green surface. Near the center of this surface is a molecule with two central black spheres with a double bond indicated by two horizontal black rods between them. Above and below to the left and right, a total of four white spheres are connected to the black spheres with white rods. A line segment extends from this structure to the label, “Ethylene absorbed on surface breaking pi bonds.” Just above this is a nearly identical structure greyed out with three downward pointing arrows to the black and white structure to indicate downward motion. The label “Ethylene” at the top of the diagram is connected to the greyed out structure with a line segment. In c, the diagram is very similar to b except that the greyed out structure and labels are gone and one of the white spheres near the black and white structure in each pair on the green surface is greyed out. Arrows point from the greyed out white spheres to the double bond between the two black spheres. In d, only a single white sphere remains from each pair in the green surface. A curved arrow points from the middle of the green surface to a model above with two central black spheres with a single black rod indicating a single bond between them. Each of the black rods has three small white spheres bonded as indicated by white rods between the black spheres and the small white spheres. The four bonds around each black sphere are evenly distributed about the black spheres.

Other significant industrial processes that involve the use of heterogeneous catalysts include the preparation of sulfuric acid, the preparation of ammonia, the oxidation of ammonia to nitric acid, and the synthesis of methanol, CH_3OH. Heterogeneous catalysts are also used in the catalytic converters found on most gasoline-powered automobiles (Figure 7.8.7).

Automobile Catalytic Converters

Scientists developed catalytic converters to reduce the amount of toxic emissions produced by burning gasoline in internal combustion engines. Catalytic converters take advantage of all five factors that affect the speed of chemical reactions to ensure that exhaust emissions are as safe as possible.

By utilizing a carefully selected blend of catalytically active metals, it is possible to effect complete combustion of all carbon-containing compounds to carbon dioxide while also reducing the output of nitrogen oxides. This is particularly impressive when we consider that one step involves adding more oxygen to the molecule and the other involves removing the oxygen (Figure 7.8.6).




Figure 7.8.6: A catalytic converter allows for the combustion of all carbon-containing compounds to carbon dioxide, while at the same time reducing the output of nitrogen oxide and other pollutants in emissions from gasoline-burning engines.

An image is shown of a catalytic converter. At the upper left, a blue arrow pointing into a pipe that enters a larger, widened chamber is labeled, “Dirty emissions.” A small black arrow that points to the lower right is positioned along the upper left side of the widened region. This arrow is labeled, “Additional oxygen from air pump.” The image shows the converter with the upper surface removed, exposing a red-brown interior. The portion of the converter closest to the dirty emissions inlet shows small, round components in an interior layer. This layer is labeled “Three-way reduction catalyst.” The middle region shows closely packed small brown rods that are aligned parallel to the dirty emissions inlet pipe. The final quarter of the interior of the catalytic converter again shows a layer of closely packed small red brown circles. Two large light grey arrows extend from this layer to the open region at the lower right of the image to the label “Clean emissions.”

Most modern, three-way catalytic converters possess a surface impregnated with a platinum-rhodium catalyst, which catalyzes the conversion nitric oxide into dinitrogen and oxygen as well as the conversion of carbon monoxide and hydrocarbons such as octane into carbon dioxide and water vapor:

$$2\,NO_2(g) \longrightarrow N_2(g) + 2\,O_2(g)$$
$$[5pt]2\,CO(g) + O_2(g) \longrightarrow 2\,CO_2(g)$$
$$[5pt]2\,C_8H_{18}(g) + 25\,O_2(g) \longrightarrow 16\,CO_2(g) + 18\,H_2O(g)$$

In order to be as efficient as possible, most catalytic converters are preheated by an electric heater. This ensures that the metals in the catalyst are fully active even before the automobile exhaust is hot enough to maintain appropriate reaction temperatures.

Enzyme Structure and Function

The study of enzymes is an important interconnection between biology and chemistry. Enzymes are usually proteins (polypeptides) that help to control the rate of chemical reactions between biologically important compounds, particularly those that are involved in cellular metabolism. Different classes of enzymes perform a variety of functions, as shown in Table 7.8.1.

ATP hydrolysis.”" data-quail-id="131" data-mt-width="1016">

Table 7.8.1: Classes of Enzymes and Their Functions

Class	Function
oxidoreductases	redox reactions
transferases	transfer of functional groups
hydrolases	hydrolysis reactions
lyases	group elimination to form double bonds
isomerases	isomerization
ligases	bond formation with ATP hydrolysis

Enzyme molecules possess an active site, a part of the molecule with a shape that allows it to bond to a specific substrate (a reactant molecule), forming an enzyme-substrate complex as a reaction intermediate. There are two models that attempt to explain how this active site works. The most simplistic model is referred to as the lock-and-key hypothesis, which suggests that the molecular shapes of the active site and substrate are complementary, fitting together like a key in a lock. The induced fit hypothesis, on the other hand, suggests that the enzyme molecule is flexible and changes shape to accommodate a bond with the substrate. This is not to suggest that an enzyme's active site is completely malleable, however. Both the lock-and-key model and the induced fit model account for the fact that enzymes can only bind with specific substrates, since in general a particular enzyme only catalyzes a particular reaction (Figure 7.8.7).




Figure 7.8.7: (a) According to the lock-and-key model, the shape of an enzyme's active site is a perfect fit for the substrate. (b) According to the induced fit model, the active site is somewhat flexible, and can change shape in order to bond with the substrate.

A diagram is shown of two possible interactions of an enzyme and a substrate. In a, which is labeled "Lock-and-key," two diagrams are shown. The first shows a green wedge-like shape with two small depressions in the upper surface of similar size, but the depression on the left has a curved shape, and the depression on the right has a pointed shape. This green shape is labeled "Enzyme." Just above this shape are two smaller, irregular, lavender shapes each with a projection from its lower surface. The lavender shape on the left has a curved projection which matches the shape of the depression on the left in the green shape below. This projection is shaded orange and has a curved arrow extending from in to the matching depression in the green shape below. Similarly, the lavender shape on the right has a projection with a pointed tip which matches the shape of the depression on the right in the green shape below. This projection is shaded orange and has a curved arrow extending from in to the matching depression in the green shape below. Two line segments extend from the depressions in the green shape to form an inverted V shape above the depressions. Above this and between the lavender shapes is the label, "Active site is proper shape." The label "Substrates" is at the very top of the diagram with line segments extending to the two lavender shapes. To the right of this diagram is a second diagram showing the lavender shapes positioned next to each other, fit snugly into the depressions in the green shape, which is labeled "Enzyme." Above this diagram is the label, "Substrate complex formed." In b, which is labeled "Induced fit," two diagrams are shown. The first shows a green wedge-like shape with two small depressions in the upper surface of similar size, but irregular shape. This green shape is labeled "Enzyme." Just above this shape are two smaller irregular lavender shapes each with a projection from its lower surface. The lavender shape on the left has a curved projection. This projection is shaded orange and has a curved arrow extending from it to the irregular depression just below it in the green shape below. Similarly, the lavender shape on the right has a projection with a pointed tip. This projection is shaded orange and has a curved arrow extending from it to the irregular depression just below it in the green shape below. Two line segments extend from the depressions in the green shape to form an inverted V shape above the depressions. Above this and between the lavender shapes is the label, "Active site changes to fit." The label, "Substrates" is at the very top of the diagram with line segments extending to the two lavender shapes. To the right of this diagram is a second diagram showing the purple shapes positioned next to each other, fit snugly into the depressions in the green shape, which is labeled "Enzyme." Above this diagram is the label "Substrate complex formed." The projections from the lavender shapes match the depression shapes in the green shape, resulting in a proper fit.

Summary

Catalysts affect the rate of a chemical reaction by altering its mechanism to provide a lower activation energy. Catalysts can be homogenous (in the same phase as the reactants) or heterogeneous (a different phase than the reactants).

Footnotes

1. "The Nobel Prize in Chemistry 1995," Nobel Prize.org, accessed February 18, 2015, Nobel Prizes Chemistry [www.nobelprize.org].

Glossary

heterogeneous catalyst

catalyst present in a different phase from the reactants, furnishing a surface at which a reaction can occur

homogeneous catalyst

catalyst present in the same phase as the reactants

This page titled 7.8: Catalysis is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **12.8: Catalysis** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/chemistry-2e.

7.9: Equilbria and Le Chatelier's Principle

Le Châtelier's principle states that if a **dynamic equilibrium** is disturbed by changing the conditions, the position of equilibrium shifts to counteract the change to reestablish an equilibrium. If a chemical reaction is at equilibrium and experiences a change in pressure, temperature, or concentration of products or reactants, the equilibrium shifts in the opposite direction to offset the change. This page covers changes to the position of equilibrium due to such changes and discusses briefly why catalysts have no effect on the equilibrium position.

Introduction

An action that changes the temperature, pressure, or concentrations of reactants in a system at equilibrium stimulates a response that partially offsets the change while a new equilibrium condition is established (2). Hence, Le Châtelier's principle states that any change to a system at equilibrium will adjust to compensate for that change. In 1884 the French chemist and engineer Henry-Louis Le Châtelier proposed one of the central concepts of chemical equilibria, which describes what happens to a system when something briefly removes it from a state of equilibrium.

It is important to understand that Le Châtelier's principle is only a useful guide to identify what happens when the conditions are changed in a reaction in dynamic equilibrium; it does **not** give reasons for the changes at the molecular level (e.g., timescale of change and underlying reaction mechanism).

Concentration Changes


Le Châtelier's principle states that if the system is changed in a way that increases the concentration of one of the reacting species, it must favor the reaction in which that species is consumed. In other words, if there is an increase in products, the reaction quotient, Q_c, is increased, making it greater than the equilibrium constant, K_c. Consider an equilibrium established between four substances, A, B, C, and D:

$$A + 2B \rightleftharpoons C + D \tag{7.9.1}$$

Increasing a concentration

What happens if conditions are altered by increasing the concentration of A?


According to Le Châtelier, the position of equilibrium will move in such a way as to counteract the change. In this case, the equilibrium position will move so that the concentration of A decreases again by reacting it with B to form more C and D. The equilibrium moves to the right (indicated by the green arrow below).



In a practical sense, this is a useful way of converting the maximum possible amount of B into C and D; this is advantageous if, for example, B is a relatively expensive material whereas A is cheap and plentiful.

Decreasing a concentration

In the opposite case in which the concentration of A is decreased, according to Le Châtelier, the position of equilibrium will move so that the concentration of A increases again. More C and D will react to replace the A that has been removed. The position of equilibrium moves to the left.



This is essentially what happens if one of the products is removed as soon as it is formed. If, for example, C is removed in this way, the position of equilibrium would move to the right to replace it. If it is continually removed, the equilibrium position shifts further and further to the right, effectively creating a one-way, irreversible reaction.

Pressure Changes


This only applies to reactions involving gases, although not necessarily all species in the reaction need to be in the gas phase. A general homogeneous gaseous reaction is given below:

$$A(g) + 2B(g) \rightleftharpoons C(g) + D(g) \tag{7.9.2}$$

Increasing the pressure

According to Le Châtelier, if the pressure is increased, the position of equilibrium will move so that the pressure is reduced again. Pressure is caused by gas molecules hitting the sides of their container. The more molecules in the container, the higher the pressure will be. The system can reduce the pressure by reacting in such a way as to produce fewer molecules.

In this case, there are three moles on the left-hand side of the equation, but only two on the right. By forming more C and D, the system causes the pressure to reduce. Increasing the pressure on a gas reaction shifts the position of equilibrium towards the side with fewer moles of gas molecules.




Example 7.9.1: Haber Process Equilibrium

$$N_2 + 3H_2 \rightleftharpoons 2NH_3 \tag{7.9.3}$$

If this mixture is transferred from a 1.5 L flask to a 5 L flask, in which direction does a net change occur to return to equilibrium? Because the volume is increased (and therefore the pressure reduced), the shift occurs in the direction that produces more moles of gas. To restore equilibrium the shift needs to occur to the left, in the direction of the reverse reaction.

Decreasing the pressure

The equilibrium will move in such a way that the pressure increases again. It can do that by producing more gaseous molecules. In this case, the position of equilibrium will move towards the left-hand side of the reaction.



What happens if there are the same number of molecules on both sides of the equilibrium reaction?

In this case, increasing the pressure has no effect on the position of the equilibrium. Because there are equal numbers of molecules on both sides, the equilibrium cannot move in any way that will reduce the pressure again. Again, this is not a rigorous explanation of why the position of equilibrium moves in the ways described.

Summary

Three ways to change the pressure of an equilibrium mixture are: 1. Add or remove a gaseous reactant or product, 2. Add an inert gas to the constant-volume reaction mixture, or 3. Change the volume of the system.


1. Adding products makes Q_c greater than K_c. This creates a net change in the reverse direction, toward reactants. The opposite occurs when adding more reactants.
2. Adding an inert gas into a gas-phase equilibrium at constant volume does not result in a shift. This is because the addition of a non-reactive gas does not change the partial pressures of the other gases in the container. While the total pressure of the system increases, the total pressure does not have any effect on the equilibrium constant.
3. When the volume of a mixture is reduced, a net change occurs in the direction that produces fewer moles of gas. When volume is increased the change occurs in the direction that produces **more** moles of gas.

Temperature Changes

To understand how temperature changes affect equilibrium conditions, the sign of the reaction enthalpy must be known. Assume that the forward reaction is exothermic (heat is evolved):



In this reaction, 250 kJ is evolved (indicated by the negative sign) when 1 mole of A reacts completely with 2 moles of B. For reversible reactions, the enthalpy value is always given as if the reaction was one-way in the forward direction. The back reaction (the conversion of C and D into A and B) would be endothermic, absorbing the same amount of heat.




The main effect of temperature on equilibrium is in changing the value of the equilibrium constant.

Warning

It is not uncommon that textbooks and instructors to consider heat as a independent "species" in a reaction. While this is rigorously incorrect because one cannot "add or remove heat" to a reaction as with species, it serves as a convenient mechanism to predict the shift of reactions with changing temperature. For example, if heat is a "reactant" ($\Delta H > 0$), then the reaction favors the formation of products at elevated temperature. Similarly, if heat is a "product" ($\Delta H < 0$), then the reaction favors the formation of reactants. A more accurate, and hence preferred, description is discussed below.

Increasing the temperature


If the temperature is increased, then the position of equilibrium will move so that the temperature is reduced again. Suppose the system is in equilibrium at 300°C, and the temperature is increased 500°C. To cool down, it needs to absorb the extra heat added. In the case, the back reaction is that in which heat is absorbed. The position of equilibrium therefore moves to the left. The new equilibrium mixture contains more A and B, and less C and D.



If the goal is to maximize the amounts of C and D formed, increasing the temperature on a reversible reaction in which the forward reaction is exothermic is a poor approach.

Decreasing the temperature?

The equilibrium will move in such a way that the temperature increases again. Suppose the system is in equilibrium at 500°C and the temperature is reduced to 400°C. The reaction will tend to heat itself up again to return to the original temperature by favoring the exothermic reaction. The position of equilibrium will move to the right with more A and B converted into C and D at the lower temperature:



Example 7.9.2:

Consider the formation of water

$$O_2 + 2H_2 \rightleftharpoons 2H_2O \quad \Delta H = -125.7\,kJ \tag{7.9.4}$$

1. What side of the reaction is favored? Because the heat is a product of the reaction, the reactants are favored.
2. Would the conversion of O_2 and H_2 to H_2O be favored with heat as a product or as a reactant? Heat as a product would shift the reaction forward, creating H_2O. The more heat added to the reaction, the more H_2O created.

Catalysts

Adding a catalyst makes absolutely **no difference** to the position of equilibrium, and Le Châtelier's principle does not apply. This is because a catalyst speeds up the forward and back reaction to the same extent and adding a catalyst does not affect the relative rates of the two reactions, it cannot affect the position of equilibrium.

However, catalysts have some application to equilibrium systems. For a dynamic equilibrium to be set up, the rates of the forward reaction and the back reaction must be equal. This does not happen instantly and for very slow reactions, it may take years! A catalyst speeds up the rate at which a reaction reaches dynamic equilibrium.

Example 7.9.3:

You might try imagining how long it would take to establish a dynamic equilibrium if you took the visual model on the introductory page and reduced the chances of the colors changing by a factor of 1000 - from 3 in 6 to 3 in 6000 and from 1 in 6 to 1 in 6000. Starting with blue squares, by the end of the time taken for the examples on that page, you would most probably still have entirely blue squares. Eventually, though, you would end up with the same sort of patterns as before - containing 25% blue and 75% orange squares.

Summary

- Increasing the temperature of a system in dynamic equilibrium favors the endothermic reaction. The system counteracts the change by absorbing the extra heat.
- Decreasing the temperature of a system in dynamic equilibrium favors the exothermic reaction. The system counteracts the change by producing more heat.

Outside Links

- Huddle, Benjamin P. "Conceptual Questions" on LeChatelier's Principle." *J. Chem. Educ.***1998** *75* 1175.
- Thomsen, Volker B. E. " Le Châtelier's Principle in the Sciences." *J. Chem. Educ.* **2000** *77* 173.

Problems

1. Varying Concentration

What will happen to the equilibrium when more $2SO_2$ (g) is added to the following system?

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

2. Varying Pressure

What will happen to the equilibrium when the volume of the system is decreased?

$$2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)}$$

3. Varying Temperature

What will happen to the equilibrium when the temperature of the system is decreased?

$$N_{2(g)} + O_{2(g)} \rightleftharpoons 2NO_{(g)} \quad \Delta H = 180.5\ kJ$$

Solutions

1.

Adding more reactants shifts the equilibrium in the direction of the products; therefore, the equilibrium shifts to the right.

Overall, the concentration of $2SO_2$ from initial equilibrium to final equilibrium will increase because only a portion of the added amount of $2SO_2$ will be consumed.

The concentration of O_2 will decrease because as the equilibrium is reestablished, O_2 is consumed with the $2SO_2$ to create more $2SO_3$. The concentration of $2SO_3$ will be greater because none of it is lost and more is being generated.

2.

Decreasing the volume leads to an increase in pressure which will cause the equilibrium to shift towards the side with fewer moles. In this example there are 3 moles on the reactant side and 2 moles on the product side, so the new equilibrium will shift towards the products (to the right).

3.

Because ΔH is positive, the reaction is endothermic in the forward direction. Removing heat from the system forces the equilibrium to shift towards the exothermic reaction, so the reverse reaction will occur and more reactants will be produced.

References

1. Pauling, L., College Chemistry, 3rd ed., Freeman, San Francisco, CA, 1964.
2. Petrucci, R., Harwood, W., Herring, F., Madura, J., General Chemistry, 9th ed., Pearson, New Jersey, 1993.
3. http://www.jce.divched.org/Journal/I...2N08/p1190.pdf

Contributors

- Amanda Wolf (UCD)
- Jim Clark (Chemguide.co.uk)

7.9: Equilbria and Le Chatelier's Principle is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

7.10: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Chemical reactions relate quantities of reactants and products. Chemists use the **mole** unit to represent 6.022×10^{23} things, whether the things are atoms of elements or molecules of compounds. This number, called **Avogadro's number**, is important because this number of atoms or molecules has the same mass in grams as one atom or molecule has in atomic mass units. **Molar masses** of substances can be determined by summing the appropriate masses from the periodic table; the final molar mass will have units of grams.

Because one mole of a substance will have a certain mass, we can use that relationship to construct conversion factors that will convert a mole amount into a mass amount, or vice versa. Such **mole-mass conversions** typically take one algebraic step.

Chemical reactions list reactants and products in molar amounts, not just molecular amounts. We can use the coefficients of a balanced chemical equation to relate moles of one substance in the reaction to moles of other substances (stoichiometry). Chemical reactions obey the Law of Conservation of Mass. To balance a chemical reaction, the coefficients in front of each compound can be adjusted until the total number of atoms of each elements is equal on both sides of the reaction arrow.

Collision Theory can be used to described the energetic aspects of chemical reactions. The reactants and products of a chemical reaction store potential energy in the form of chemical bonds and intermolecular forces. The energy difference in the bond energies of the reactants and products is called "Enthalpy". When the products are lower in potential energy than the reactants, then this excess energy is released as heat and the reaction is described as exothermic. When the products are higher in potential energy than the reactants, then energy must be added to the reaction as heat for it to occur and the reaction is described as endothermic.

The rate of a chemical reaction is influenced by its activation energy. The larger the activation energy, the slower the reaction rate. Catalysts can be added to reactions to lower the activation energy and increase the reaction rate.

Not all reactions go to completion. For some reactions, both the forward and reverse reaction can occur simultaneously. When the forward and reverse reaction rates are equal, the reaction is described as being "at equilibrium". Another way to recognize when a system is at equilibrium is when the concentration of reactants and products remain constant. Le Chatlier's principle states that the direction of an equilibrium reaction will shift to reduce the stress.

7.10: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

8: States of Matter and the Gas Laws

Most of us are familiar with the three phases of matter: solid, liquid, and gas. Indeed, we addressed the energy changes involved in phase changes. The picture on this page shows the substance we are probably most familiar with as having those three phases: water. In everyday life, we commonly come in contact with water as a solid (ice), as a liquid, and as a gas (steam). All we have to do is change the conditions of the substance—typically temperature—and we can change the phase from solid to liquid to gas and back again. Under the proper conditions of temperature and pressure, many substances—not only water—can experience the three different phases. An understanding of the phases of matter is important for our understanding of all matter. In this chapter, we will explore the three phases of matter.

8.1: Phase Changes

8.2: Solids, Liquids, and Gases

8.3: Gases and Pressure

8.4: Gas Laws

8.5: Chapter Summary

8: States of Matter and the Gas Laws is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

8.1: Phase Changes

Learning Objectives

- Determine the heat associated with a phase change.

Matter can exist in one of several different states, including a gas, liquid, or solid state. The amount of energy in molecules of matter determines the **state of matter**.

- A **gas** is a state of matter in which atoms or molecules have enough energy to move freely. The molecules come into contact with one another only when they randomly collide.
- A **liquid** is a state of matter in which atoms or molecules are constantly in contact but have enough energy to keep changing positions relative to one another.
- A **solid** is a state of matter in which atoms or molecules do not have enough energy to move. They are constantly in contact and in fixed positions relative to one another.




Figure **8.1.1**: States of Matter. All three containers contain a substance with the same mass, but the substances are in different states. In the left-hand container, the substance is a gas, which has spread to fill its container. It takes both the shape and volume of the container. In the middle container, the substance is a liquid, which has spread to take the shape of its container but not the volume. In the right-hand container, the substance is a solid, which takes neither the shape nor the volume of its container.

The following are the changes of state:

Changes of State

Phase to Phase transition	Transition Name
Solid → Liquid	Melting or fusion
Liquid → Gas	Vaporization
Liquid → Solid	Freezing
Gas → Liquid	Condensation
Solid → Gas	Sublimation

- If heat is added to a substance, such as in melting, vaporization, and sublimation, the process is **endothermic**. In this instance, heat is increasing the speed of the molecules causing them move faster (examples: solid to liquid; liquid to gas; solid to gas).
- If heat is removed from a substance, such as in freezing and condensation, then the process is **exothermic**. In this instance, heat is decreasing the speed of the molecules causing them move slower (examples: liquid to solid; gas to liquid). These changes **release heat** to the surroundings.
- The amount of heat needed to change a sample from solid to liquid would be the same to reverse from liquid to solid. The only difference is the direction of heat transfer.

Example 8.1.1

Label each of the following processes as endothermic or exothermic.

a. water boiling
b. ice forming on a pond

Solution a

endothermic - you must put a pan of water on the stove and give it heat in order to get water to boil. Because you are adding heat/energy, the reaction is endothermic.

Solution b

exothermic - think of ice forming in your freezer instead. You put water into the freezer, which takes heat out of the water, to get it to freeze. Because heat is being pulled out of the water, it is exothermic. Heat is leaving.

8.1.1 Exercise

Label each of the following processes as endothermic or exothermic.

a. water vapor condensing
b. gold melting

Answer

a. exothermic

b. endothermic

A phase change is a physical process in which a substance goes from one phase to another. Usually the change occurs when adding or removing heat at a particular temperature, known as the melting point or the boiling point of the substance. The melting point is the temperature at which the substance goes from a solid to a liquid (or from a liquid to a solid). The boiling point is the temperature at which a substance goes from a liquid to a gas (or from a gas to a liquid). The nature of the phase change depends on the direction of the heat transfer. Heat going *into* a substance changes it from a solid to a liquid or a liquid to a gas. Removing heat *from* a substance changes a gas to a liquid or a liquid to a solid.

Two key points are worth emphasizing. First, at a substance's melting point or boiling point, two phases can exist simultaneously. Take water (H_2O) as an example. On the Celsius scale, H_2O has a melting point of 0°C and a boiling point of 100°C. At 0°C, both the solid and liquid phases of H_2O can coexist. However, if heat is added, some of the solid H_2O will melt and turn into liquid H_2O. If heat is removed, the opposite happens: some of the liquid H_2O turns into solid H_2O. A similar process can occur at 100°C: adding heat increases the amount of gaseous H_2O, while removing heat increases the amount of liquid H_2O (Figure 8.1.1).




Figure 8.1.2: Heating curve for water. As heat is added to solid water, the temperature increases until it reaches 0 °C, the melting point. At this point, the phase change, added heat goes into changing the state from a solid to liquid. Only when this phase change is complete, the temperature can increase. (CC BY 3.0 Unported; Community College Consortium for Bioscience Credentials).

> Water is a good substance to use as an example because many people are already familiar with it. Other substances have melting points and boiling points as well.

Second, as shown in Figure 8.1.1, **the temperature of a substance does not change** as the substance goes **from one phase to another**. In other words, phase changes are isothermal (isothermal means "constant temperature"). Again, consider H_2O as an example. Solid water (ice) can exist at 0°C. If heat is added to ice at 0°C, some of the solid changes phase to make liquid, which is also at 0°C. Remember, the solid and liquid phases of H_2O can coexist at 0°C. Only after all of the solid has melted into liquid does the addition of heat change the temperature of the substance.

For each phase change of a substance, there is a characteristic quantity of heat needed to perform the phase change per gram (or per mole) of material. The heat of fusion (ΔH_{fus}) is the amount of heat per gram (or per mole) required for a phase change that occurs at the melting point. The heat of vaporization (ΔH_{vap}) is the amount of heat per gram (or per mole) required for a phase change that occurs at the boiling point. If you know the total number of grams or moles of material, you can use the ΔH_{fus} or the ΔH_{vap} to determine the total heat being transferred for melting or solidification using these expressions:

$$\text{heat} = n \times \Delta H_{fus}$$

where n is the number of moles and ΔH_{fus} is expressed in energy/mole or

$$\text{heat} = m \times \Delta H_{fus}$$

where m is the mass in grams and ΔH_{fus} is expressed in energy/gram.

For the boiling or condensation, use these expressions:

$$\text{heat} = n \times \Delta H_{vap}$$

where n is the number of moles) and ΔH_{vap} is expressed in energy/mole or

$$\text{heat} = m \times \Delta H_{vap}$$

where m is the mass in grams and ΔH_{vap} is expressed in energy/gram.

Remember that a phase change depends on the direction of the heat transfer. If heat transfers in, solids become liquids, and liquids become solids at the melting and boiling points, respectively. If heat transfers out, liquids solidify, and gases condense into liquids. At these points, there are no changes in temperature as reflected in the above equations.

Example 8.1.2

How much heat is necessary to melt 55.8 g of ice (solid H_2O) at 0°C? The heat of fusion of H_2O is 79.9 cal/g.

Solution

We can use the relationship between heat and the heat of fusion (Equation **8.1.1**) to determine how many cal of heat are needed to melt this ice:

$$\text{heat} = \text{m} \times \Delta H_{\text{fus}}$$

$$\text{heat} = (55.8 \cancel{\text{g}}) \left(\frac{79.9 \text{ cal}}{\cancel{\text{g}}} \right) = 4,460 \text{ cal}$$

Exercise 8.1.2

How much heat is necessary to vaporize 685 g of H_2O at 100°C? The heat of vaporization of H_2O is 540 cal/g.

Answer

$$\text{heat} = \text{m} \times \Delta H_{\text{vap}}$$

$$\text{heat} = (685 \cancel{\text{g}}) \left(\frac{540 \text{ cal}}{\cancel{\text{g}}} \right) = 370,000 \text{ cal}$$

Table **8.1.1** lists the heats of fusion and vaporization for some common substances. Note the units on these quantities; when you use these values in problem solving, make sure that the other variables in your calculation are expressed in units consistent with the units in the specific heats or the heats of fusion and vaporization.

Table **8.1.1**: Heats of Fusion and Vaporization for Selected Substances

Substance	ΔH_{fus} (cal/g)	ΔH_{vap} (cal/g)
aluminum (Al)	94.0	2,602
gold (Au)	15.3	409
iron (Fe)	63.2	1,504
water (H_2O)	79.9	540
sodium chloride (NaCl)	123.5	691
ethanol (C_2H_5OH)	45.2	200.3
benzene (C_6H_6)	30.4	94.1

Sublimation

There is also a phase change where a solid goes directly to a gas:

\[\text{solid} \rightarrow \text{gas} \nonumber\]

This phase change is called *sublimation*. Each substance has a characteristic heat of sublimation associated with this process. For example, the heat of sublimation (ΔH_{sub}) of H_2O is 620 cal/g.

We encounter sublimation in several ways. You may already be familiar with dry ice, which is simply solid carbon dioxide (CO_2). At −78.5°C (−109°F), solid carbon dioxide sublimes, changing directly from the solid phase to the gas phase:

$$\text{CO}_2(\text{s}) \xrightarrow{-78.5^\circ\text{C}} \text{CO}_2(\text{g})$$

Solid carbon dioxide is called dry ice because it does not pass through the liquid phase. Instead, it does directly to the gas phase. (Carbon dioxide *can* exist as liquid but only under high pressure.) Dry ice has many practical uses, including the long-term preservation of medical samples.

Even at temperatures below 0°C, solid H_2O will slowly sublime. For example, a thin layer of snow or frost on the ground may slowly disappear as the solid H_2O sublimes, even though the outside temperature may be below the freezing point of water. Similarly, ice cubes in a freezer may get smaller over time. Although frozen, the solid water slowly sublimes, redepositing on the colder cooling elements of the freezer, which necessitates periodic defrosting (frost-free freezers minimize this redeposition). Lowering the temperature in a freezer will reduce the need to defrost as often.

Under similar circumstances, water will also sublime from frozen foods (e.g., meats or vegetables), giving them an unattractive, mottled appearance called freezer burn. It is not really a "burn," and the food has not necessarily gone bad, although it looks unappetizing. Freezer burn can be minimized by lowering a freezer's temperature and by wrapping foods tightly so water does not have any space to sublime into.

Key Takeaway

- There is an energy change associated with any phase change.

8.1: Phase Changes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **7.3: Phase Changes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.2: Solids, Liquids, and Gases

Learning Objectives

- To describe the solid and liquid phases.

Solids and liquids are collectively called *condensed phases* because their particles are in virtual contact. The two states share little else, however.

Solids

In the solid state, the individual particles of a substance are in fixed positions with respect to each other because there is not enough thermal energy to overcome the intermolecular interactions between the particles. As a result, solids have a definite shape and volume. Most solids are hard, but some (like waxes) are relatively soft. Many solids composed of ions can also be quite brittle.



Figure 8.2.1: Crystalline Arrangement of Quartz crystal cluster. Some large crystals look the way they do because of the regular arrangement of atoms (ions) in their crystal structure. from Wikipedia.

Solids usually have their constituent particles arranged in a regular, three-dimensional array of alternating positive and negative ions called a crystal. The effect of this regular arrangement of particles is sometimes visible macroscopically, as shown in Figure 8.2.1. Some solids, especially those composed of large molecules, cannot easily organize their particles in such regular crystals and exist as amorphous (literally, “without form”) solids. Glass is one example of an amorphous solid.

Liquids

If the particles of a substance have enough energy to partially overcome intermolecular interactions, then the particles can move about each other while remaining in contact. This describes the liquid state. In a liquid, the particles are still in close contact, so liquids have a definite volume. However, because the particles can move about each other rather freely, a liquid has no definite shape and takes a shape dictated by its container.



Figure 8.2.2: The formation of a spherical droplet of liquid water minimizes the surface area, which is the natural result of surface tension in liquids. from Wikipedia.

Gases

If the particles of a substance have enough energy to completely overcome intermolecular interactions, then the particles can separate from each other and move about randomly in space. Like liquids, gases have no definite shape, but unlike solids and liquids, gases have no definite volume either.




Figure 8.2.3: A Representation of the Solid, Liquid, and Gas States. A solid has definite volume and shape, a liquid has a definite volume but no definite shape, and a gas has neither a definite volume nor shape.

The change from solid to liquid usually does not significantly change the volume of a substance. However, the change from a liquid to a gas significantly increases the volume of a substance, by a factor of 1,000 or more. Figure 8.2.3 shows the differences among solids, liquids, and gases at the molecular level, while Table 8.2.1 lists the different characteristics of these states.

Table 8.2.1: Characteristics of the Three States of Matter

Characteristic	Solid	Liquid	Gas
shape	definite	indefinite	indefinite
volume	definite	definite	indefinite
relative intermolecular interaction strength	strong	moderate	weak
relative particle positions	in contact and fixed in place	in contact but not fixed	not in contact, random positions

Example 8.2.1

What state or states of matter does each statement, describe?

a. This state has a definite volume.
b. This state has no definite shape.
c. This state allows the individual particles to move about while remaining in contact.

Solution

a. This statement describes either the liquid state or the solid state.
b. This statement describes either the liquid state or the gas state.
c. This statement describes the liquid state.

Exercise 8.2.1

What state or states of matter does each statement describe?

a. This state has individual particles in a fixed position with regard to each other.
b. This state has individual particles far apart from each other in space.
c. This state has a definite shape.

Answer

a. solid
b. gas
c. solid

Looking Closer: Water, the Most Important Liquid

Earth is the only known body in our solar system that has liquid water existing freely on its surface. That is a good thing because life on Earth would not be possible without the presence of liquid water.

Water has several properties that make it a unique substance among substances. It is an excellent solvent; it dissolves many other substances and allows those substances to react when in solution. In fact, water is sometimes called the *universal solvent* because of this ability. Water has unusually high melting and boiling points (0°C and 100°C, respectively) for such a small molecule. The boiling points for similar-sized molecules, such as methane (BP = −162°C) and ammonia (BP = −33°C), are more than 100° lower. Though a liquid at normal temperatures, water molecules experience a relatively strong intermolecular interaction that allows them to maintain the liquid phase at higher temperatures than expected.

Unlike most substances, the solid form of water is less dense than its liquid form, which allows ice to float on water. The most energetically favorable configuration of H_2O molecules is one in which each molecule is hydrogen-bonded to four neighboring molecules. Owing to the thermal motions, this ideal is never achieved in the liquid, but when water freezes to ice, the molecules settle into exactly this kind of an arrangement in the ice crystal. This arrangement requires that the molecules be somewhat farther apart than would otherwise be the case; as a consequence, ice, in which hydrogen bonding is at its maximum, has a more open structure, and thus a lower density than water.




Figure **8.2.4**: Three-dimensional views of a typical local structure of liquid water (left) and ice (right).

Here are three-dimensional views of a typical local structure of water (left) and ice (right.) Notice the greater openness of the ice structure which is necessary to ensure the strongest degree of hydrogen bonding in a uniform, extended crystal lattice. The structure of liquid water is very similar, but in the liquid, the hydrogen bonds are continually broken and formed because of rapid molecular motion. Because ice is less dense than liquid water, rivers, lakes, and oceans freeze from the top down. In fact, the ice forms a protective surface layer that insulates the rest of the water, allowing fish and other organisms to survive in the lower levels of a frozen lake or sea. If ice were denser than the liquid, the ice formed at the surface in cold weather would sink as fast as it formed. Bodies of water would freeze from the bottom up, which would be lethal for most aquatic creatures. The expansion of water when freezing also explains why automobile or boat engines must be protected by “antifreeze” and why unprotected pipes in houses break if they are allowed to freeze.

Water also requires an unusually large amount of energy to change temperature. While 100 J of energy will change the temperature of 1 g of Fe by 230°C, this same amount of energy will change the temperature of 1 g of H_2O by only 100°C. Thus, water changes its temperature slowly as heat is added or removed. This has a major impact on weather, as storm systems like hurricanes can be impacted by the amount of heat that ocean water can store.

Water’s influence on the world around us is affected by these properties. Isn’t it fascinating that such a small molecule can have such a big impact?

Key Takeaway

- Solids and liquids are phases that have their own unique properties.

8.2: Solids, Liquids, and Gases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **8.2: Solids and Liquids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.3: Gases and Pressure

Learning Objectives

- To describe the gas phase.

The gas phase is unique among the three states of matter in that there are some simple models we can use to predict the physical behavior of all gases—independent of their identities. We cannot do this for the solid and liquid states. In fact, the development of this understanding of the behavior of gases represents the historical dividing point between alchemy and modern chemistry. Initial advances in the understanding of gas behavior were made in the mid 1600s by Robert Boyle, an English scientist who founded the Royal Society (one of the world's oldest scientific organizations).

How is it that we can model all gases independent of their chemical identity? The answer is in a group of statements called the kinetic theory of gases:

- Gases are composed of tiny particles that are separated by large distances.
- Gas particles are constantly moving, experiencing collisions with other gas particles and the walls of their container.
- The velocity of gas particles is related to the temperature of a gas.
- Gas particles do not experience any force of attraction or repulsion with each other.

Did you notice that none of these statements relates to the identity of the gas? This means that all gases should behave similarly. A gas that follows these statements perfectly is called an *ideal gas*. Most gases show slight deviations from these statements and are called *real gases*. However, the existence of real gases does not diminish the importance of the kinetic theory of gases.

One of the statements of the kinetic theory mentions collisions. As gas particles are constantly moving, they are also constantly colliding with each other and with the walls of their container. There are forces involved as gas particles bounce off the container walls (Figure **8.3.1**). The force generated by gas particles divided by the area of the container walls yields pressure. Pressure is a property we can measure for a gas, but we typically do not consider pressure for solids or liquids.




Figure **8.3.1**: Gas Pressure. Pressure is what results when gas particles rebound off the walls of their container.

The basic unit of pressure is the newton per square meter (N/m^2). This combined unit is redefined as a pascal (Pa). One pascal is not a very large amount of pressure. A more useful unit of pressure is the bar, which is 100,000 Pa (1 bar = 100,000 Pa). Other common units of pressure are the atmosphere (atm), which was originally defined as the average pressure of Earth's atmosphere at sea level; and mmHg (millimeters of mercury), which is the pressure generated by a column of mercury 1 mm high. The unit millimeters of mercury is also called a torr, named after the Italian scientist Evangelista Torricelli, who invented the barometer in the mid-1600s. A more precise definition of atmosphere, in terms of torr, is that there are exactly 760 torr in 1 atm. A bar equals 1.01325 atm. Given all the relationships between these pressure units, the ability to convert from one pressure unit to another is a useful skill.

Example 8.3.1: Converting Pressures

Write a conversion factor to determine how many atmospheres are in 1,547 mmHg.

Solution

Because 1 mmHg equals 1 torr, the given pressure is also equal to 1,547 torr. Because there are 760 torr in 1 atm, we can use this conversion factor to do the mathematical conversion:

$$1,547 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 2.04 \text{ atm}$$

Note how the torr units cancel algebraically.

? Exercise 8.3.1: Converting Pressures

Write a conversion factor to determine how many millimeters of mercury are in 9.65 atm.

Answer

$$9.65 \text{ atm} \times \frac{760 \text{ mmHg}}{1 \text{ atm}} = 7,334 \text{ mmHg}.$$

The kinetic theory also states that there is no interaction between individual gas particles. Although we know that there are, in fact, intermolecular interactions in real gases, the kinetic theory assumes that gas particles are so far apart that the individual particles don't "feel" each other. Thus, we can treat gas particles as tiny bits of matter whose identity isn't important to certain physical properties.

Key Takeaway

- The gas phase has certain general properties characteristic of that phase.

8.3: Gases and Pressure is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **8.3: Gases and Pressure** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.4: Gas Laws

Learning Objectives

- To predict the properties of gases using the gas laws.

Experience has shown that several properties of a gas can be related to each other under certain conditions. The properties are pressure (*P*), volume (*V*), temperature (*T*, in kelvins), and amount of material expressed in moles (*n*). What we find is that a sample of gas cannot have any random values for these properties. Instead, only certain values, dictated by some simple mathematical relationships, will occur.

Boyle's Law

The first simple relationship, referred to as a gas law, is between the pressure of a gas and its volume. If the amount of gas in a sample and its temperature are kept constant, then as the pressure of a gas is increased, the volume of the gas decreases proportionately. Mathematically, this is written as

$$\mathrm{P} \propto \frac{1}{\mathrm{V}}$$

where the "∝" symbol means "is proportional to." This is one form of Boyle's law, which relates the pressure of a gas to its volume.

A more useful form of Boyle's law involves a change in conditions of a gas. For a given amount of gas at a constant temperature, if we know the initial pressure and volume of a gas sample and the pressure or volume changes, we can calculate what the new volume or pressure will be. That form of Boyle's law is written

$$P_i V_i = P_f V_f \tag{8.4.1}$$

where the subscript i refers to initial conditions and the subscript f refers to final conditions.

To use 8.4.1, you need to know any three of the variables so that you can algebraically calculate the fourth variable. Also, the pressure quantities must have the same units, as must the two volume quantities. If the two similar variables don't have the same variables, one value must be converted to the other value's unit.

Example 8.4.1: Increasing Pressure in a Gas

What happens to the volume of a gas if its pressure is increased? Assume all other conditions remain the same.

Solution

If the pressure of a gas is increased, the volume decreases in response.

Exercise 8.4.1: Increasing Volume in a Gas

What happens to the pressure of a gas if its volume is increased? Assume all other conditions remain the same.

Answer

If the volume of a gas is increased, the pressure decreases.

Example 8.4.2: Gas Compression

If a sample of gas has an initial pressure of 1.56 atm and an initial volume of 7.02 L, what is the final volume if the pressure is reduced to 0.987 atm? Assume that the amount and the temperature of the gas remain constant.

Solution

The key in problems like this is to be able to identify which quantities represent which variables from the relevant equation. The way the question is worded, you should be able to tell that 1.56 atm is P_i, 7.02 L is V_i, and 0.987 atm is P_f. What we are looking for is the final volume—V_f. Therefore, substituting these values into $P_iV_i = P_fV_f$:

$$(1.56 \text{ atm})(7.02 \text{ L}) = (0.987 \text{ atm}) \times V_f$$

The expression has atmospheres on both sides of the equation, so they cancel algebraically:

$$(1.56)(7.02 \text{ L}) = (0.987) \times V_f$$

Now we divide both sides of the expression by 0.987 to isolate V_f, the quantity we are seeking:

$$\frac{(1.56)(7.02 \text{ L})}{0.987} = \mathbf{V_f}$$

Performing the multiplication and division, we get the value of V_f, which is 11.1 L. The volume increases. This should make sense because the pressure decreases, so pressure and volume are inversely related.

Exercise 8.4.2

If a sample of gas has an initial pressure of 3.66 atm and an initial volume of 11.8 L, what is the final pressure if the volume is reduced to 5.09 L? Assume that the amount and the temperature of the gas remain constant.

Answer

8.48 atm

If the units of similar quantities are not the same, one of them must be converted to the other quantity's units for the calculation to work out properly. It does not matter which quantity is converted to a different unit; the only thing that matters is that the conversion and subsequent algebra are performed properly. The following example illustrates this process.

Example 8.4.3

If a sample of gas has an initial pressure of 1.56 atm and an initial volume of 7.02 L, what is the final volume if the pressure is changed to 1,775 torr? Does the answer make sense? Assume that the amount and the temperature of the gas remain constant.

Solution

This example is similar to Example **8.4.2**, except now the final pressure is expressed in torr. For the math to work out properly, one of the pressure values must be converted to the other unit. Let us change the initial pressure to torr:

$$1.56 \text{ atm} \times \frac{760 \text{ torr}}{1 \text{ atm}} = 1,190 \text{ torr}$$

Now we can use Boyle's law:

$$(1{,}190 \text{ torr})(7.02 \text{ L}) = (1{,}775 \text{ torr}) \times V_f$$

Torr cancels algebraically from both sides of the equation, leaving

$$(1{,}190)(7.02 \text{ L}) = (1{,}775) \times V_f$$

Now we divide both sides of the equation by 1,775 to isolate V_f on one side. Solving for the final volume,

$$\mathbf{V_f} = \frac{(1,190)(7.02 \text{ L})}{1,775} = 4.71 \text{ L}$$

Because the pressure increases, it makes sense that the volume decreases.

The answer for the final volume is essentially the same if we converted the 1,775 torr to atmospheres: $1,775 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 2.336 \text{ atm}$. Using Boyle's law: (1.56 atm)(7.02 L) = (2.335 atm) × V_f; $\mathbf{V_f} = \frac{(1.56 \text{ atm})(7.02 \text{ L})}{2.336 \text{ atm}} = 4.69 \text{ L}$.

Exercise 8.4.3

If a sample of gas has an initial pressure of 375 torr and an initial volume of 7.02 L, what is the final pressure if the volume is changed to 4,577 mL? Does the answer make sense? Assume that amount and the temperature of the gas remain constant.

Answer

575 torr

To Your Health: Breathing

Breathing certainly is a major contribution to your health! Without breathing, we could not survive. Curiously, the act of breathing itself is little more than an application of Boyle's law.

The lungs are a series of ever-narrowing tubes that end in a myriad of tiny sacs called alveoli. It is in the alveoli that oxygen from the air transfers to the bloodstream and carbon dioxide from the bloodstream transfers to the lungs for exhalation. For air to move in and out of the lungs, the pressure inside the lungs must change, forcing the lungs to change volume—just as predicted by Boyle's law.

The pressure change is caused by the diaphragm, a muscle that covers the bottom of the lungs. When the diaphragm moves down, it expands the size of our lungs. When this happens, the air pressure inside our lungs decreases slightly. This causes new air to rush in, and we inhale. The pressure decrease is slight—only 3 torr, or about 0.4% of an atmosphere. We inhale only 0.5–1.0 L of air per normal breath.

Exhaling air requires that we relax the diaphragm, which pushes against the lungs and slightly decreases the volume of the lungs. This slightly increases the pressure of the air in the lungs, and air is forced out; we exhale. Only 1–2 torr of extra pressure is needed to exhale. So with every breath, our own bodies are performing an experimental test of Boyle's law.

Charles's Law

Another simple gas law relates the volume of a gas to its temperature. Experiments indicate that as the temperature of a gas sample is increased, its volume increases as long as the pressure and the amount of gas remain constant. The way to write this mathematically is

$$V \propto T$$

At this point, the concept of temperature must be clarified. Although the Kelvin scale is the preferred temperature scale, the Celsius scale is also a common temperature scale used in science. The Celsius scale is based on the melting and boiling points of water and is actually the common temperature scale used by most countries around the world (except for the United States, which still uses the Fahrenheit scale). The value of a Celsius temperature is directly related to its Kelvin value by a simple expression:

$$\text{Kelvin temperature} = \text{Celsius temperature} + 273$$

Thus, it is easy to convert from one temperature scale to another.

The Kelvin scale is sometimes referred to as the absolute scale because the zero point on the Kelvin scale is at absolute zero, the coldest possible temperature. On the other temperature scales, absolute zero is −260°C or −459°F.

The expression relating a gas volume to its temperature begs the following question: to which temperature scale is the volume of a gas related? The answer is that gas volumes are directly related to the *Kelvin temperature*. Therefore, the temperature of a gas sample should always be expressed in (or converted to) a Kelvin temperature.

Example 8.4.4: Increasing Temperature

What happens to the volume of a gas if its temperature is decreased? Assume that all other conditions remain constant.

Solution

If the temperature of a gas sample is decreased, the volume decreases as well.

Exercise 8.4.4

What happens to the temperature of a gas if its volume is increased? Assume that all other conditions remain constant.

Answer

The temperature increases.

As with Boyle's law, the relationship between volume and temperature can be expressed in terms of initial and final values of volume and temperature, as follows:

$$\frac{V_i}{T_i} = \frac{V_f}{T_f}$$

where V_i and T_i are the initial volume and temperature, and V_f and T_f are the final volume and temperature. This is Charles's law. The restriction on its use is that the pressure of the gas and the amount of gas must remain constant. (Charles's law is sometimes referred to as Gay-Lussac's law, after the scientist who promoted Charles's work.)

Example 8.4.5

A gas sample at 20°C has an initial volume of 20.0 L. What is its volume if the temperature is changed to 60°C? Does the answer make sense? Assume that the pressure and the amount of the gas remain constant.

Solution

Although the temperatures are given in degrees Celsius, we must convert them to the kelvins before we can use Charles's law. Thus,

$$20°C + 273 = 293\ K = T_i \quad 60°C + 273 = 333\ K = T_f$$

Now we can substitute these values into Charles's law, along with the initial volume of 20.0 L:

$$\frac{20.0\ L}{293\ K} = \frac{V_f}{333\ K}$$

Multiplying the 333 K to the other side of the equation, we see that our temperature units will cancel:

$$\frac{(333\ K)(20.0\ L)}{293\ K} = V_f$$

Solving for the final volume, V_f = 22.7 L. So, as the temperature is increased, the volume increases. This makes sense because volume is directly proportional to the absolute temperature (as long as the pressure and the amount of the remain constant).

Exercise 8.4.5

A gas sample at 35°C has an initial volume of 5.06 L. What is its volume if the temperature is changed to −35°C? Does the answer make sense? Assume that the pressure and the amount of the gas remain constant.

Answer

3.91 L

Combined Gas Law

Other gas laws can be constructed, but we will focus on only two more. The combined gas law brings Boyle's and Charles's laws together to relate pressure, volume, and temperature changes of a gas sample:

Double subscripts: use braces to clarify

To apply this gas law, the amount of gas should remain constant. As with the other gas laws, the temperature must be expressed in kelvins, and the units on the similar quantities should be the same. Because of the dependence on three quantities at the same time, it is difficult to tell in advance what will happen to one property of a gas sample as two other properties change. The best way to know is to work it out mathematically.

Example 8.4.6

A sample of gas has P_i = 1.50 atm, V_i = 10.5 L, and T_i = 300 K. What is the final volume if P_f = 0.750 atm and T_f = 350 K?

Solution

Using the combined gas law, substitute for five of the quantities:

$$\frac{(1.50\ \text{atm})(10.5\ \text{L})}{300\ \text{K}} = \frac{(0.750\ \text{atm})(\text{V}_\text{f})}{350\ \text{K}}$$

We algebraically rearrange this expression to isolate V_f on one side of the equation:

$$\text{V}_\text{f} = \frac{(1.50\ \text{atm})(10.5\ \text{L})(350\ \text{K})}{(300\ \text{K})(0.750\ \text{atm})} = 24.5\ \text{L}$$

Note how all the units cancel except the unit for volume.

8.4.6 Exercise

A sample of gas has P_i = 0.768 atm, V_i = 10.5 L, and T_i = 300 K. What is the final pressure if V_f = 7.85 L and T_f = 250 K?

Answer

0.856 atm

Example 8.4.7

A balloon containing a sample of gas has a temperature of 22°C and a pressure of 1.09 atm in an airport in Cleveland. The balloon has a volume of 1,070 mL. The balloon is transported by plane to Denver, where the temperature is 11°C and the pressure is 655 torr. What is the new volume of the balloon?

Solution

The first task is to convert all quantities to the proper and consistent units. The temperatures must be expressed in kelvins, and the pressure units are different so one of the quantities must be converted. Let us convert the atmospheres to torr:

$$22°\text{C} + 273 = 295\ \text{K} = T_i$$

$$11°\text{C} + 273 = 284\ \text{K} = T_f$$

$$1.09\ \text{atm} \times \frac{760\ \text{torr}}{1\ \text{atm}} = 828\ \text{torr} = \text{P}_\text{i}$$

Now we can substitute the quantities into the combined has law:

$$\frac{(828\ \text{torr})(1,070\ \text{mL})}{295\ \text{K}} = \frac{(655\ \text{torr}) \times \text{V}_\text{f}}{284\ \text{K}}$$

To solve for V_f, we multiply the 284 K in the denominator of the right side into the numerator on the left, and we divide 655 torr in the numerator of the right side into the denominator on the left:

$$\frac{(828\ \text{torr})(1,070\ \text{mL})(284\ \text{K})}{(295\ \text{K})(655\ \text{torr})} = \text{V}_\text{f}$$

Notice that torr and kelvins cancel, as they are found in both the numerator and denominator. The only unit that remains is milliliters, which is a unit of volume. So V_f = 1,300 mL. The overall change is that the volume of the balloon has increased by 230 mL.

8.4.7 Exercise

A balloon used to lift weather instruments into the atmosphere contains gas having a volume of 1,150 L on the ground, where the pressure is 0.977 atm and the temperature is 18°C. Aloft, this gas has a pressure of 6.88 torr and a temperature of −15°C. What is the new volume of the gas?

Answer

110,038 L

The Ideal Gas Law

So far, the gas laws we have used have focused on changing one or more properties of the gas, such as its volume, pressure, or temperature. There is one gas law that relates all the independent properties of a gas under any particular condition, rather than a change in conditions. This gas law is called the ideal gas law. The formula of this law is as follows:

$$PV = nRT$$

In this equation, P is pressure, V is volume, n is amount of moles, and T is temperature. R is called the ideal gas law constant and is a proportionality constant that relates the values of pressure, volume, amount, and temperature of a gas sample. The variables in this equation do not have the subscripts i and f to indicate an initial condition and a final condition. The ideal gas law relates the four independent properties of a gas under *any* conditions.

The value of R depends on what units are used to express the other quantities. If volume is expressed in liters and pressure in atmospheres, then the proper value of R is as follows:

$$\mathrm{R = 0.08205\ \frac{L \cdot atm}{mol \cdot K}}$$

This may seem like a strange unit, but that is what is required for the units to work out algebraically.

Example 8.4.8

What is the volume in liters of 1.45 mol of N_2 gas at 298 K and 3.995 atm?

Solution

Using the ideal gas law where P = 3.995 atm, n = 1.45, and T = 298,

$$\mathrm{(3.995\ atm) \times V = (1.45\ mol)\left(0.08205\ \frac{L \cdot atm}{mol \cdot K}\right)(298\ K)}$$

On the right side, the moles and kelvins cancel. Also, because atmospheres appear in the numerator on both sides of the equation, they also cancel. The only remaining unit is liters, a unit of volume. So

$$3.995 \times V = (1.45)(0.08205)(298)\ \text{L}$$

Dividing both sides of the equation by 3.995 and evaluating, we get V = 8.87 L. Note that the conditions of the gas are not changing. Rather, the ideal gas law allows us to determine what the fourth property of a gas (here, volume) *must* be if three other properties (here, amount, pressure, and temperature) are known.

8.4.8 Exercise

What is the pressure of a sample of CO_2 gas if 0.557 mol is held in a 20.0 L container at 451 K?

Answer

1.03 atm

For convenience, scientists have selected 273 K (0°C) and 1.00 atm pressure as a set of standard conditions for gases. This combination of conditions is called standard temperature and pressure (STP). Under these conditions, 1 mol of any gas has about the same volume. We can use the ideal gas law to determine the volume of 1 mol of gas at STP:

$$(1.00\ \text{atm}) \times \text{V} = (1.00\ \text{mol}) \left(0.08205\ \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (273\ \text{K})$$

This volume is 22.4 L. Because this volume is independent of the identity of a gas, the idea that 1 mol of gas has a volume of 22.4 L at STP makes a convenient conversion factor:

1 mol gas = 22.4 L (at STP)

Example 8.4.9

Cyclopropane (C_3H_6) is a gas that formerly was used as an anesthetic. How many moles of gas are there in a 100.0 L sample if the gas is at STP?

Solution

We can set up a simple, one-step conversion that relates moles and liters:

$$100.0\ \text{L}\ \text{C}_3\text{H}_6 \times \frac{1\ \text{mol}}{22.4\ \text{L}} = 4.46\ \text{mol}\ \text{C}_3\text{H}_6$$

There are almost 4.5 mol of gas in 100.0 L.

Note: Because of its flammability, cyclopropane is no longer used as an anesthetic gas.

8.4.9 Exercise

Freon is a trade name for a series of fluorine- and chlorine-containing gases that formerly were used in refrigeration systems. What volume does 8.75 mol of Freon have at STP?

Note: Many gases known as Freon are no longer used because their presence in the atmosphere destroys the ozone layer, which protects us from ultraviolet light from the sun.

Answer

196 L

Airbags

Airbags (Figure **8.4.3**) are a safety feature provided in most automobiles since the 1990s. The effective operation of an airbag requires that it be rapidly inflated with an appropriate amount (volume) of gas when the vehicle is involved in a collision. This requirement is satisfied in many automotive airbag systems through use of explosive chemical reactions, one common choice being the decomposition of sodium azide, NaN_3. When sensors in the vehicle detect a collision, an electrical current is passed through a carefully measured amount of NaN_3 to initiate its decomposition:

$$2\,\text{NaN}_3(s) \rightarrow 3\,\text{N}_2(g) + 2\,\text{Na}(s)$$

This reaction is very rapid, generating gaseous nitrogen that can deploy and fully inflate a typical airbag in a fraction of a second (~0.03–0.1 s). Among many engineering considerations, the amount of sodium azide used must be appropriate for generating enough nitrogen gas to fully inflate the air bag and ensure its proper function. For example, a small mass (~100 g) of NaN_3 will generate approximately 50 L of N_2.



Figure **8.4.3**: Airbags deploy upon impact to minimize serious injuries to passengers. (credit: Jon Seidman)

Dalton's Law of Partial Pressures

The ideal gas equation of state applies to mixtures just as to pure gases. It was in fact with a gas mixture, ordinary air, that Boyle, Gay-Lussac and Charles did their early experiments. The only new concept we need in order to deal with gas mixtures is the *partial pressure*, a concept invented by the famous English chemist John Dalton (1766-1844). Dalton reasoned that the low density and high compressibility of gases indicates that they consist mostly of empty space; from this it follows that when two or more different gases occupy the same volume, they behave entirely independently. The contribution that each component of a gaseous mixture makes to the total pressure of the gas is known as the *partial pressure* of that gas.

The definition of Dalton's Law of Partial Pressures that address this is:

The total pressure of a gas is the sum of the partial pressures of its components

which is expressed algebraically as

$$P_{total} = P_1 + P_2 + P_3 ... = \sum_i P_i$$

or, equivalently

$$P_{total} = \frac{RT}{V} \sum_i n_i$$


There is also a similar relationship based on *volume fractions*, known as *Amagat's law of partial volumes*. It is exactly analogous to Dalton's law, in that it states that the total volume of a mixture is just the sum of the partial volumes of its components. But there are two important differences: Amagat's law holds only for ideal gases which must all be at the same temperature and pressure. Dalton's law has neither of these restrictions. Although Amagat's law seems intuitively obvious, it sometimes proves useful in chemical engineering applications. We will make no use of it in this course.

Example 8.4.10

Three flasks having different volumes and containing different gases at various pressures are connected by stopcocks as shown. When the stopcocks are opened,

a. What will be the pressure in the system?
b. Which gas will be most abundant in the mixture?

Assume that the temperature is uniform and that the volume of the connecting tubes is negligible.



Solution

The trick here is to note that the total number of moles n_T and the temperature remain unchanged, so we can make use of Boyle's law PV = constant. We will work out the details for CO_2 only, denoted by subscripts *a*.

For CO_2,

$$P_aV_a = (2.13\ atm)(1.50\ L) = 3.20\ L \cdot atm$$

Adding the PV products for each separate container, we obtain

\[\sum_i P_i V_i = 6.36\; L \cdot atm = n_T RT \nonumber \]

We will call this sum P_1V_1. After the stopcocks have been opened and the gases mix, the new conditions are denoted by P_2V_2.

From Boyle's law (8.4.1,

$$P_1V_1 = P_2V_2 = 6.36\ L \cdot atm$$

$$V_2 = \sum_i V_i = 4.50\ L$$

Solving for the final pressure P_2 we obtain (6.36 L-atm)/(4.50 L) = **1.41 atm**.

For part *(b)*, note that the number of moles of each gas is $n = PV/RT$. The mole fraction of any one gas is $X_i = n_i / n_T$. For CO_2, this works out to (3.19/*RT*) / (6.36/*RT*) = 0.501. Because this exceeds 0.5, we know that this is the most abundant gas in the final mixture.

Dalton's law states that in a gas mixture (P_{total}) each gas will exert a pressure independent of the other gases (P_n) and each gas will behave as if it alone occupies the total volume. By extension, the partial pressure of each gas can be calculated by multiplying the total pressure (P_{total}) by the gas percentage (%).

$$P_{Total} = P_1 + P_2 + P_3 + P_4 + ... + P_n$$

or

$$P_n = \frac{n}{P_{Total}}$$

Table 8.4.1: Partial Pressures for the gases in air on a typical day

Gas	Partial Pressure (mm Hg)	Percentage (%)
Nitrogen, (N_2\)	P_{N_2} = 594	78
Oxygen, O_2	P_{O_2} = 160	21
Carbon Dioxide, CO_2	P_{CO_2} = 0.25	0.033
Water Vapor, H_2O	P_{H_2O} = 5.7	0.75
Other trace gases	P_{Other} = 0.05	0.22
Total air	P_{Total} = 760	1

Application of Dalton's Law: Collecting Gases over Water

A common laboratory method of collecting the gaseous product of a chemical reaction is to conduct it into an inverted tube or bottle filled with water, the opening of which is immersed in a larger container of water. This arrangement is called a *pneumatic trough*, and was widely used in the early days of chemistry. As the gas enters the bottle it displaces the water and becomes trapped in the upper part.




Figure 8.4.1:An Apparatus for Collecting Gases by the Displacement of Water

The volume of the gas can be observed by means of a calibrated scale on the bottle, but what about its pressure? The total pressure confining the gas is just that of the atmosphere transmitting its force through the water. (An exact calculation would also have to take into account the height of the water column in the inverted tube.) But liquid water itself is always in equilibrium with its vapor, so the space in the top of the tube is a mixture of two gases: the gas being collected, and gaseous H_2O. The partial pressure of H_2O is known as the vapor pressure of water and it depends on the temperature. In order to determine the quantity of gas we have collected, we must use Dalton's Law to find the partial pressure of that gas.

Example 8.4.11

Oxygen gas was collected over water as shown above. The atmospheric pressure was 754 torr, the temperature was 22°C, and the volume of the gas was 155 mL. The vapor pressure of water at 22°C is 19.8 torr. Use this information to estimate the number of moles of O_2 produced.

Solution

From Dalton's law,

$$P_{O_2} = P_{total} – P_{H_2O} = 754 – 19.8 = 734\ torr = 0.966\ atm$$

Now use the Ideal Gas Law to convert to moles

$$n = \frac{PV}{RT} = \frac{(0.966\ atm)(0.155\ L)}{(0.082\ Latmmol^{-1}K^{-1})(295\ K)} = 0.00619\ mol$$

Henry’s Law

Henry's law is one of the gas laws formulated by William Henry in 1803. It states: "At a constant temperature, the amount of a given gas that dissolves in a given type and volume of liquid is directly proportional to the partial pressure of that gas in equilibrium with that liquid." An equivalent way of stating the law is that the solubility of a gas in a liquid is directly proportional to the partial pressure of the gas above the liquid.

To explain this law, Henry derived the equation:

$$C = kP_{gas}$$

where

Henry’s Law tells us that the greater the pressure of gas above the surface of a liquid, the higher the concentration of the gas in the liquid. Also, Henry’s law tells us that gases diffuse from areas of high gas concentration to areas of low gas concentration.

Applicability of Henry's Law

- Henry's law only works if the molecules are at equilibrium.
- Henry's law does not work for gases at high pressures (e.g., $N_{2\,(g)}$ at high pressure becomes very soluble and harmful when in the blood supply).
- Henry's law does not work if there is a chemical reaction between the solute and solvent (e.g., $HCl_{(g)}$ reacts with water by a dissociation reaction to generate H_3O^+ and Cl^- ions).

Application of Henry's Law: Scuba diving

Our respiratory systems are designed to maintain the proper oxygen concentration in the blood when the partial pressure of O_2 is 0.21 atm, its normal sea-level value. Below the water surface, the pressure increases by 1 atm for each 10.3 m increase in depth; thus a scuba diver at 10.3 m experiences a total of 2 atm pressure pressing on the body. In order to prevent the lungs from collapsing, the air the diver breathes should also be at about the same pressure.




Figure **8.4.2**: Scuba Diving actively takes into account both Henry's and Dalton's Laws

But at a total pressure of 2 atm, the partial pressure of O_2 in ordinary air would be 0.42 atm; at a depth of 100 ft (about 30 m), the O_2 pressure of 0.8 atm would be far too high for health. For this reason, the air mixture in the pressurized tanks that scuba divers wear must contain a smaller fraction of O_2. This can be achieved most simply by raising the nitrogen content, but high partial pressures of N_2 can also be dangerous, resulting in a condition known as nitrogen narcosis. The preferred diluting agent for sustained deep diving is helium, which has very little tendency to dissolve in the blood even at high pressures.

Career Focus: Respiratory Therapist

Certain diseases—such as emphysema, lung cancer, and severe asthma—primarily affect the lungs. Respiratory therapists help patients with breathing-related problems. They can evaluate, help diagnose, and treat breathing disorders and even help provide emergency assistance in acute illness where breathing is compromised.

Most respiratory therapists must complete at least two years of college and earn an associate's degree, although therapists can assume more responsibility if they have a college degree. Therapists must also pass state or national certification exams. Once certified, respiratory therapists can work in hospitals, doctor's offices, nursing homes, or patient's homes. Therapists work with equipment such as oxygen tanks and respirators, may sometimes dispense medication to aid in breathing, perform tests, and educate patients in breathing exercises and other therapy.

Because respiratory therapists work directly with patients, the ability to work well with others is a must for this career. It is an important job because it deals with one of the most crucial functions of the body.

Key Takeaway

- The physical properties of gases are predictable using mathematical formulas known as gas laws.
- C is the solubility of a gas at a fixed temperature in a particular solvent (in units of M or mL gas/L)
- k is Henry's law constant (often in units of M/atm)
- P_{gas} is the partial pressure of the gas (often in units of Atm)

8.4: Gas Laws is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **8.4: Gas Laws** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

https://chem.libretexts.org/@go/page/387566

8.5: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **phase** is a certain form of matter that has the same physical properties throughout. Three phases are common: the solid, the liquid, and the gas phase. What determines the phase of a substance? Generally, the strength of the **intermolecular interactions** determines whether a substance is a solid, liquid, or gas under any particular conditions. **Covalent network bonding** is a very strong form of intermolecular interaction. Diamond is one example of a substance that has this intermolecular interaction. **Ionic interactions**, the forces of attraction due to oppositely charged ions, are also relatively strong. Covalent bonds are another type of interaction within molecules, but if the bonds are **polar covalent bonds**, then the unequal sharing of electrons can cause charge imbalances within molecules that cause interactions between molecules. These molecules are described as **polar**, and these interactions are called **dipole-dipole interactions**. A certain rather strong type of dipole-dipole interaction, involving a hydrogen atom, is called **hydrogen bonding**. On the other hand, equal sharing of electrons forms **nonpolar covalent bonds**, and the interactions between different molecules is less because the molecules are nonpolar. All substances have very weak **dispersion forces** (also called **London forces**) caused by the movement of electrons within the bonds themselves.

In the solid phase, intermolecular interactions are so strong that they hold the individual atoms or molecules in place. In many solids, the regular three-dimensional arrangement of particles makes a **crystal**. In other solids, the irregular arrangement of particles makes an **amorphous** solid. In liquids, the intermolecular interactions are strong enough to keep the particles of substance together but not in place. Thus, the particles are free to move over each other but still remain in contact.

In gases, the intermolecular interactions are weak enough that the individual particles are separated from each other in space. The **kinetic theory of gases** is a collection of statements that describe the fundamental behavior of all gases. Among other properties, gases exert a **pressure** on their container. Pressure is measured using units like **pascal**, **bar**, **atmosphere**, or **mmHg** (also called a **torr**).

There are several simple relationships between the variables used to describe a quantity of gas. These relationships are called **gas laws**. **Boyle's law** relates the pressure and volume of a gas, while **Charles's law** relates the volume and absolute temperature of a gas. The **combined gas law** relates the volume, pressure, and absolute temperature of a gas sample. All of these gas laws allow us to understand the changing conditions of a gas. The **ideal gas law** relates the pressure, volume, amount, and absolute temperature of a gas under any conditions. These four variables are related to the **ideal gas law constant**, which is the proportionality constant used to calculate the conditions of a gas. Because the conditions of a gas can change, a set of benchmark conditions called **standard temperature and pressure (STP)** is defined. Standard temperature is 0ºC, and standard pressure is 1.00 atm.

Dalton's Law of Partial Pressures describes the behavior of a mixture of gases and states that t*he total pressure of a gas is the sum of the partial pressures of its components.* **Henry's Law states** that the solubility of a gas in a liquid is directly proportional to the partial pressure of the gas above the liquid.

8.5: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

9: Solutions

Solutions are all around us. Air, for example, is a solution. If you live near a lake, a river, or an ocean, that body of water is not pure H_2O but most probably a solution. Much of what we drink—for example, soda, coffee, tea, and milk—is at least in part a solution. Solutions are a large part of everyday life. A lot of the chemistry occurring around us happens in solution. In fact, much of the chemistry that occurs in our own bodies takes place in solution, and many solutions—such as the Ringer's lactate IV solution—are important for our health. In our understanding of chemistry, we need to understand a little bit about solutions. In this chapter, you will learn about the special characteristics of solutions, how solutions are characterized, and some of their properties.

9.1: Introduction to Solutions
9.2: Solution Concentration- Molarity
9.3: Other Units for Solution Concentrations
9.4: The Dissolution Process
9.5: Properties of Solutions
9.6: Dialysis and Tonicity
9.7: Chapter Summary

9: Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

9.1: Introduction to Solutions

Learning Objectives

- To understand what causes solutions to form.

A solution is another name for a homogeneous mixture. A *mixture* as a material composed of two or more substances. In a solution, the combination is so intimate that the different substances cannot be differentiated by sight, even with a microscope. Compare, for example, a mixture of salt and pepper and another mixture consisting of salt and water. In the first mixture, we can readily see individual grains of salt and the flecks of pepper. A mixture of salt and pepper is not a solution. However, in the second mixture, no matter how carefully we look, we cannot see two different substances. Salt dissolved in water is a solution.

The major component of a solution, called the **solvent**, is typically the same phase as the solution itself. Each minor component of a solution (and there may be more than one) is called the **solute**. In most of the solutions we will describe in this textbook, there will be no ambiguity about whether a component is the solvent or the solute. For example, in a solution of salt in water, the solute is salt, and solvent is water.

Solutions come in all phases, and the solvent and the solute do not have to be in the same phase to form a solution (such as salt and water). For example, air is a gaseous solution of about 80% nitrogen and about 20% oxygen, with some other gases present in much smaller amounts. An alloy is a solid solution consisting of a metal (like iron) with some other metals or nonmetals dissolved in it. Steel, an alloy of iron and carbon and small amounts of other metals, is an example of a solid solution. Table **9.1.1** lists some common types of solutions, with examples of each.

Table **9.1.1**: Types of Solutions

Solvent Phase	Solute Phase	Example
gas	gas	air
liquid	gas	carbonated beverages
liquid	liquid	ethanol (C_2H_5OH) in H_2O (alcoholic beverages)
liquid	solid	saltwater
solid	gas	H_2 gas absorbed by Pd metal
solid	liquid	Hg(ℓ) in dental fillings
solid	solid	steel alloys

What causes a solution to form? The simple answer is that the solvent and the solute must have similar intermolecular interactions. When this is the case, the individual particles of solvent and solute can easily mix so intimately that each particle of solute is surrounded by particles of solvent, forming a solution. However, if two substances have very different intermolecular interactions, large amounts of energy are required to force their individual particles to mix intimately, so a solution does not form. Thus two alkanes like *n*-heptane, C_7H_{16}, and *n*-hexane, C_6H_{14}, are completely miscible in all proportions. The C_7H_{16} and C_6H_{14} molecules are so similar (recall Section 4.6) that there are only negligible differences in intermolecular forces.

For a similar reason, methanol, CH_3OH, is completely miscible with water. In this case both molecules are polar and can form hydrogen bonds among themselves, and so there are strong intermolecular attractions within each liquid. However, CH_3OH dipoles can align with H_2O dipoles, and CH_3OH molecules can hydrogen bond to H_2O molecules, and so the attractions among unlike molecules in the solution are similar to those among like molecules in each pure liquid.

This process leads to a simple rule of thumb: *like dissolves like*. Solvents that are very polar will dissolve solutes that are very polar or even ionic. Solvents that are nonpolar will dissolve nonpolar solutes. Thus water, being polar, is a good solvent for ionic

compounds and polar solutes like ethanol (C_2H_5OH). However, water does not dissolve nonpolar solutes, such as many oils and greases (Figure **9.1.1**).



Figure **9.1.1**: A beaker holds water with blue food dye (upper liquid layer) and a much more dense perfluoroheptane (a fluorocarbon) lower liquid layer. The two fluids cannot mix and the dye cannot dissolve in fluorocarbon. A goldfish and a crab have been introduced into the water. The goldfish cannot penetrate the dense fluorocarbon. The crab floats at the liquid boundary with only parts of his legs penetrating the fluorocarbon fluid, unable to sink to the bottom of the beaker. Quarter coins rest on the bottom of the beaker. Animals were rescued from their predicament after the photo was taken. Figure used with permission from Wikipedia (Sbharris (Steven B. Harris)).

We use the word soluble to describe a solute that dissolves in a particular solvent, and the word insoluble for a solute that does not dissolve in a solvent. Thus, we say that sodium chloride is soluble in water but insoluble in hexane (C_6H_{14}). If the solute and the solvent are both liquids and soluble in any proportion, we use the word miscible, and the word immiscible if they are not.

✔ Example 9.1.1

Water is considered a polar solvent. Which substances should dissolve in water?

1. methanol (CH_3OH)
2. sodium sulfate (Na_2SO_4)
3. octane (C_8H_{18})

Solution

Because water is polar, substances that are polar or ionic will dissolve in it.

1. Because of the OH group in methanol, we expect its molecules to be polar. Thus, we expect it to be soluble in water. As both water and methanol are liquids, the word *miscible* can be used in place of *soluble*.
2. Sodium sulfate is an ionic compound, so we expect it to be soluble in water.
3. Like other hydrocarbons, octane is nonpolar, so we expect that it would not be soluble in water.

? Exercise 9.1.1

Toluene ($C_6H_5CH_3$) is widely used in industry as a nonpolar solvent. Which substances should dissolve in toluene?

a. water (H_2O)
b. sodium sulfate (Na_2SO_4)
c. octane (C_8H_{18})

Answer

Octane only.

Example 9.1.2

Predict which of the following compounds will be most soluble in water:

a. $\underset{\text{Ethanol}}{CH_3CH_2OH}$

b. $\underset{\text{Hexanol}}{CH_3CH_2CH_2CH_2CH_2CH_2OH}$

Solution

Since ethanol contains an **OH** group, it can hydrogen bond to water. Although the same is true of hexanol, the OH group is found only at one end of a fairly large molecule. The rest of the molecule can be expected to behave much as though it were a nonpolar alkane. This substance should thus be much less soluble than the first. Experimentally we find that ethanol is completely miscible with water, while only 0.6 g hexanol dissolves in 100 g water.

9.1.2 Exercise

Would I_2 be more soluble in CCl_4 or H_2O?

Answer

I_2 is nonpolar. Of the two solvents, CCl_4 is nonpolar and H_2O is polar, so I_2 would be expected to be more soluble in CCl_4.

Key Takeaway

- Solutions form because a solute and a solvent experience similar intermolecular interactions.

9.1: Introduction to Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **9.1: Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.2: Solution Concentration- Molarity

Learning Objectives

- Use molarity to determine quantities in chemical reactions.
- Use molarity as a conversion factor in calculations.

Another way of expressing concentration is to give the number of moles of solute per unit volume of solution. Of all the quantitative measures of concentration, molarity is the one used most frequently by chemists. **Molarity** is defined as the number of moles of solute per liter of solution.

$$\text{molarity} = \frac{\text{number of moles of solute}}{\text{number of liters of solution}} \tag{9.2.1}$$

The symbol for molarity is **M** or moles/liter. Chemists also use square brackets to indicate a reference to the molarity of a substance. For example, the expression $[Ag^+]$ refers to the molarity of the silver ion in solution. Solution concentrations expressed in molarity are the easiest to perform calculations with, but the most difficult to make in the lab. Such concentration units are useful for discussing chemical reactions in which a solute is a product or a reactant. Molar mass can then be used as a conversion factor to convert amounts in moles to amounts in grams.

It is important to remember that “mol” in this expression refers to moles of solute and that “L” refers to liters of solution. For example, if you have 1.5 mol of NaCl dissolved in 0.500 L of solution, its molarity is

$$\frac{1.5 \text{ mol NaCl}}{0.500 \text{ L solution}} = 3.0 \text{ M NaCl}$$

Sometimes (aq) is added when the solvent is water, as in “3.0 M NaCl (aq).” This is read as "a 3.00 ***molar*** sodium chloride solution," meaning that there are 3.00 moles of NaOH dissolved per one liter of solution.

> *Be sure to note that molarity is calculated as the total volume of the **entire** solution, not just volume of solvent! The solute contributes to total volume.*

If the quantity of the solute is given in mass units, you must convert mass units to mole units before using the definition of molarity to calculate concentration. For example, what is the molar concentration of a solution of 22.4 g of HCl dissolved in 1.56 L?

Step 1: First, convert the mass of solute to moles using the molar mass of HCl (36.5 g/mol):

$$22.4 \, \cancel{g HCl} \times \frac{1 \, mol \, HCl}{36.5 \, \cancel{g HCl}} = 0.614 \, mol \, HCl$$

Step 2: Now we can use the definition of molarity to determine a concentration:

$$M = \frac{0.614 \, mol \, HCl}{1.56 L \, solution} = 0.394 \, M HCl$$


Before a molarity concentration can be calculated, the amount of the solute must be expressed in moles, and the volume of the solution must be expressed in liters, as demonstrated in the following example.

Example 9.2.1

A solution is prepared by dissolving **42.23 g** of NH_4Cl into enough water to make **500.0 mL** of solution. Calculate its molarity.

Solution

Solutions to Example 13.6.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: Mass $= 42.23\ \text{g}\ NH_4Cl$ Volume solution $= 500.0\ \text{mL} = 0.5000\ \text{L}$ Find: Molarity = ? M
List other known quantities.	Molar mass $NH_4Cl = 53.50\ \text{g/mol}$
Plan the problem.	1. The mass of the ammonium chloride is first converted to moles.  $\frac{1\ mol\ NH_4Cl}{53.50g\ NH_4Cl}$ 2. Then the molarity is calculated by dividing by liters. Note the given volume has been converted to liters. $\text{M} = \frac{\text{mol } NH_4Cl}{\text{L solution}}$
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $42.23\ \cancel{\text{g } NH_4Cl} \times \frac{1\ \text{mol } NH_4Cl}{53.50\ \cancel{\text{g } NH_4Cl}} = 0.7893\ \text{mol } NH_4Cl$ $\frac{0.7893\ \text{mol } NH_4Cl}{0.5000\ \text{L solution}} = 1.579\ \text{M}$
Think about your result.	The molarity is $1.579\ \text{M}$, meaning that a liter of the solution would contain $1.579\ \text{mol } NH_4Cl$. Four significant figures are appropriate.

? Exercise 9.2.1A

What is the molarity of a solution made when 66.2 g of $C_6H_{12}O_6$ are dissolved to make 235 mL of solution?

Answer

1.57 M $C_6H_{12}O_6$

? Exercise 9.2.1B

What is the concentration, in mol/L, where 137 g of NaCl has been dissolved in enough water to make 500 mL of solution?

Answer

4.69 M NaCl

Using Molarity in Calculations


Concentration can be a conversion factor between the amount of solute and the amount of solution or solvent (depending on the definition of the concentration unit). As such, concentrations can be useful in a variety of stoichiometry problems. In many cases, it is best to use the original definition of the concentration unit; it is that definition that provides the conversion factor.

A simple example of using a concentration unit as a conversion factor is one in which we use the definition of the concentration unit and rearrange; we can do the calculation again as a unit conversion, rather than as a definition.

Example 9.2.2: Determining Moles of Solute, Given the Concentration and Volume of a Solution

For example, suppose we ask how many moles of solute are present in 0.108 L of a 0.887 M NaCl solution. Because 0.887 M means 0.887 mol/L, we can use this second expression for the concentration as a conversion factor:

Solution



$$0.108 \cancel{L\ solution} \times \frac{0.887\ mol\ NaCl}{1 \cancel{L\ solution}} = 0.0958\ mol\ NaCl$$

If we used the definition approach, we get the same answer, but now we are using conversion factor skills.


Like any other conversion factor that relates two different types of units, the reciprocal of the concentration can be also used as a conversion factor.

Example 9.2.3: Determining Volume of a Solution, Given the Concentration and Moles of Solute

Using concentration as a conversion factor, how many liters of 2.35 M $CuSO_4$ are needed to obtain 4.88 mol of $CuSO_4$?

Solution

This is a one-step conversion, but the concentration must be written as the reciprocal for the units to work out:



$$4.88 \cancel{mol\ CuSO_4} \times \frac{1\ L\ solution}{2.35 \cancel{mol\ CuSO_4}} = 2.08\ L\ of\ solution$$


In a laboratory situation, a chemist must frequently prepare a given volume of solutions of a known molarity. The task is to calculate the mass of the solute that is necessary. The molarity equation can be rearranged to solve for moles, which can then be converted to grams. The following example illustrates this.

Example 9.2.4

A chemist needs to prepare **3.00 L** of a **0.250 M** solution of potassium permanganate (**$KMnO_4$**). What mass of **$KMnO_4$** does she need to make the solution?

Solution

Solutions to Example 13.6.4

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: Molarity $= 0.250\ \text{M}$ Volume $= 3.00\ \text{L}$ Find: Mass $\text{KMnO}_4 = ?\ \text{g}$
List other known quantities.	Molar mass $\text{KMnO}_4 = 158.04\ \text{g/mol}$ 0.250 mol $KMnO_4$ to 1 L of $KMnO_4$ solution
Plan the problem.	 $\frac{.250\ mol\ KMnO_4}{1\ L\ solution}$ $\frac{158.04g\ KMnO_4}{1\ mol\ KMnO_4}$
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $\text{mol KMnO}_4 = 0.250\ \text{M KMnO}_4 \times 3.00\ \text{L} = 0.750\ \text{mol KMnO}_4$ $3.00\ \cancel{L\ solution} \times \frac{0.250\ \cancel{\text{mol KMnO}_4}}{1\ \cancel{L\ solution}} \times \frac{158.04\ \text{g KMnO}_4}{1\ \cancel{\text{mol KMnO}_4}} = 119\ \text{g KMnO}_4$
Think about your result.	When $119\ \text{g}$ of potassium permanganate is dissolved into water to make $3.00\ \text{L}$ of solution, the molarity is $0.250\ \text{M}$.

Exercise 9.2.4A

Using concentration as a conversion factor, how many liters of 0.0444 M CH_2O are needed to obtain 0.0773 mol of CH_2O?

Answer

1.74 L

Exercise 9.2.4B

Answer the problems below using concentration as a conversion factor.

a. What mass of solute is present in 1.08 L of 0.0578 M H_2SO_4?
b. What volume of 1.50 M HCl solution contains 10.0 g of hydrogen chloride?

Answer a

6.12 g

Answer b

183 mL or 0.183L

How to Indicate Concentration

- Square brackets are often used to represent concentration, e.g., [NaOH] = 0.50 M.
- Use the capital letter M for molarity, not a lower case m (this is a different concentration unit called *molality*).



Watch as the Flinn Scientific Tech Staff demonstrates "How To Prepare Solutions."

It is important to note that there are many different ways you can set up and solve your chemistry equations. Some students prefer to answer multi-step calculations in one long step, while others prefer to work out each step individually. Neither method is necessarily better or worse than the other method—whichever makes the most sense to ***you*** is the one that you should use. In this text, we will typically use unit analysis (also called dimension analysis or factor analysis).

Contributors and Attributions

- Peggy Lawson (Oxbow Prairie Heights School). Funded by Saskatchewan Educational Technology Consortium.

9.2: Solution Concentration- Molarity is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **13.6: Specifying Solution Concentration- Molarity** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

9.3: Other Units for Solution Concentrations

Learning Objectives

- Define the concentration units of mass percentage, volume percentage, mass-volume percentage, parts-per-million (ppm), and parts-per-billion (ppb)
- Perform computations relating a solution's concentration and its components' volumes and/or masses using these units

In the previous section, we introduced molarity, a very useful measurement unit for evaluating the concentration of solutions. However, molarity is only one measure of concentration. In this section, we will introduce some other units of concentration that are commonly used in various applications, either for convenience or by convention.

Mass Percentage

Earlier in this chapter, we introduced percent composition as a measure of the relative amount of a given element in a compound. Percentages are also commonly used to express the composition of mixtures, including solutions. The mass percentage of a solution component is defined as the ratio of the component's mass to the solution's mass, expressed as a percentage:

$$\text{mass percentage} = \frac{\text{mass of component}}{\text{mass of solution}} \times 100\% \tag{9.3.1}$$

We are generally most interested in the mass percentages of solutes, but it is also possible to compute the mass percentage of solvent.

Mass percentage is also referred to by similar names such as *percent mass*, *percent weight*, *weight/weight percent*, and other variations on this theme. The most common symbol for mass percentage is simply the percent sign, %, although more detailed symbols are often used including %mass, %weight, and (w/w)%. Use of these more detailed symbols can prevent confusion of mass percentages with other types of percentages, such as volume percentages (to be discussed later in this section).

Mass percentages are popular concentration units for consumer products. The label of a typical liquid bleach bottle (Figure 9.3.1) cites the concentration of its active ingredient, sodium hypochlorite (NaOCl), as being 7.4%. A 100.0-g sample of bleach would therefore contain 7.4 g of NaOCl.



Figure 9.3.1: Liquid bleach is an aqueous solution of sodium hypochlorite (NaOCl). This brand has a concentration of 7.4% NaOCl by mass.

The sides of two cylindrical containers are shown. Each container's label is partially visible. The left container's label reads "Bleach." The right label contains more information about the product including the phrase, "Contains: Sodium hypochlorite 7.4 %."

Example 9.3.1: Calculation of Percent by Mass

A 5.0-g sample of spinal fluid contains 3.75 mg (0.00375 g) of glucose. What is the percent by mass of glucose in spinal fluid?

Solution

The spinal fluid sample contains roughly 4 mg of glucose in 5000 mg of fluid, so the mass fraction of glucose should be a bit less than one part in 1000, or about 0.1%. Substituting the given masses into the equation defining mass percentage yields:

$$\%\,\text{glucose} = \frac{3.75\ \text{mg glucose} \times \frac{1\ \text{g}}{1000\ \text{mg}}}{5.0\ \text{g spinal fluid}} = 0.075\%$$

The computed mass percentage agrees with our rough estimate (it's a bit less than 0.1%).

Note that while any mass unit may be used to compute a mass percentage (mg, g, kg, oz, and so on), the same unit must be used for both the solute and the solution so that the mass units cancel, yielding a dimensionless ratio. In this case, we converted the units of solute in the numerator from mg to g to match the units in the denominator. We could just as easily have converted the denominator from g to mg instead. As long as identical mass units are used for both solute and solution, the computed mass percentage will be correct.

? Exercise 9.3.1

A bottle of a tile cleanser contains 135 g of **HCl** and 775 g of water. What is the percent by mass of **HCl** in this cleanser?

Answer


14.8%

✓ Example 9.3.2: Calculations using Mass Percentage

"Concentrated" hydrochloric acid is an aqueous solution of 37.2% **HCl** that is commonly used as a laboratory reagent. The density of this solution is 1.19 g/mL. What mass of **HCl** is contained in 0.500 L of this solution?

Solution

The HCl concentration is near 40%, so a 100-g portion of this solution would contain about 40 g of HCl. Since the solution density isn't greatly different from that of water (1 g/mL), a reasonable estimate of the HCl mass in 500 g (0.5 L) of the solution is about five times greater than that in a 100 g portion, or $5 \times 40 = 200\ \text{g}$. To derive the mass of solute in a solution from its mass percentage, we need to know the corresponding mass of the solution. Using the solution density given, we can convert the solution's volume to mass, and then use the given mass percentage to calculate the solute mass. This mathematical approach is outlined in this flowchart:



A diagram of three boxes connected by a right-facing arrow in between each is shown. The box on the left contains the phrase, "Volume of solution (m L)," the middle box reads, "Mass of solution (g)," while the one on the right contains the phrase, "Mass of H C l (g)." There is a phrase under the left arrow that says, "Multiply by density (g / m L)" and under the right arrow it states, "Multiply by mass percent as ratio (g H C l / g solution)."

For proper unit cancellation, the 0.500-L volume is converted into 500 mL, and the mass percentage is expressed as a ratio, 37.2 g HCl/g solution:

$$500\ \text{mL solution}\left(\frac{1.19\ \text{g solution}}{\text{mL solution}}\right)\left(\frac{37.2\ \text{g HCl}}{100\ \text{g solution}}\right) = 221\ \text{g HCl}$$

This mass of HCl is consistent with our rough estimate of approximately 200 g.

? Exercise 9.3.2

What volume of concentrated HCl solution contains 125 g of HCl?

Answer

282 mL

Volume Percentage

Liquid volumes over a wide range of magnitudes are conveniently measured using common and relatively inexpensive laboratory equipment. The concentration of a solution formed by dissolving a liquid solute in a liquid solvent is therefore often expressed as a volume percentage, %vol or (v/v)%:

$$\text{volume percentage} = \frac{\text{volume solute}}{\text{volume solution}} \times 100\% \tag{9.3.2}$$

Example 9.3.3: Calculations using Volume Percentage

Rubbing alcohol (isopropanol) is usually sold as a 70%vol aqueous solution. If the density of isopropyl alcohol is 0.785 g/mL, how many grams of isopropyl alcohol are present in a 355 mL bottle of rubbing alcohol?

Solution

Per the definition of volume percentage, the isopropanol volume is 70% of the total solution volume. Multiplying the isopropanol volume by its density yields the requested mass:

$$\begin{aligned} &355 \text{ mL solution} \left(\frac{70 \text{ mL isopropyl alcohol}}{100 \text{ mL solution}}\right)\left(\frac{0.785 \text{ g isopropyl alcohol}}{1 \text{ mL isopropyl alcohol}}\right) \\ &= 195 \text{ g isopropyl alcohol} \end{aligned}$$

Exercise 9.3.3

Wine is approximately 12% ethanol (CH_3CH_2OH) by volume. Ethanol has a molar mass of 46.06 g/mol and a density 0.789 g/mL. How many moles of ethanol are present in a 750-mL bottle of wine?

Answer

1.5 mol ethanol

Mass-Volume Percentage

"Mixed" percentage units, derived from the mass of solute and the volume of solution, are popular for certain biochemical and medical applications. A mass-volume percent is a ratio of a solute's mass to the solution's volume expressed as a percentage. The specific units used for solute mass and solution volume may vary, depending on the solution. For example, physiological saline solution, used to prepare intravenous fluids, has a concentration of 0.9% mass/volume (m/v), indicating that the composition is 0.9 g of solute per 100 mL of solution. The concentration of glucose in blood (commonly referred to as "blood sugar") is also typically expressed in terms of a mass-volume ratio. Though not expressed explicitly as a percentage, its concentration is usually given in milligrams of glucose per deciliter (100 mL) of blood (Figure 9.3.2).



(a) (b)

Figure 9.3.2: "Mixed" mass-volume units are commonly encountered in medical settings. (a) The NaCl concentration of physiological saline is 0.9% (m/v). (b) This device measures glucose levels in a sample of blood. The normal range for glucose concentration in blood (fasting) is around 70–100 mg/dL. (credit a: modification of work by "The National Guard"/Flickr; credit b: modification of work by Biswarup Ganguly).

Two pictures are shown labeled a and b. Picture a depicts a clear, colorless solution in a plastic bag being held in a person's hand. Picture b shows a person's hand holding a detection meter with a digital readout screen while another hand holds someone's finger up to the end of the meter. The meter is pressed to the drop of blood that is at the end of the person's finger.

Parts per Million and Parts per Billion


Very low solute concentrations are often expressed using appropriately small units such as parts per million (ppm) or parts per billion (ppb). Like percentage ("part per hundred") units, ppm and ppb may be defined in terms of masses, volumes, or mixed mass-volume units. There are also ppm and ppb units defined with respect to numbers of atoms and molecules.

The mass-based definitions of ppm and ppb are given here:

$$\text{ppm} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^6 \text{ ppm} \tag{9.3.3}$$

$$\text{ppb} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^9 \text{ ppb} \tag{9.3.4}$$

Both ppm and ppb are convenient units for reporting the concentrations of pollutants and other trace contaminants in water. Concentrations of these contaminants are typically very low in treated and natural waters, and their levels cannot exceed relatively low concentration thresholds without causing adverse effects on health and wildlife. For example, the EPA has identified the maximum safe level of fluoride ion in tap water to be 4 ppm. Inline water filters are designed to reduce the concentration of fluoride and several other trace-level contaminants in tap water (Figure **9.3.3**).



(a) (b)

Figure **9.3.3**: (a) In some areas, trace-level concentrations of contaminants can render unfiltered tap water unsafe for drinking and cooking. (b) Inline water filters reduce the concentration of solutes in tap water. (credit a: modification of work by Jenn Durfey; credit b: modification of work by "vastateparkstaff"/Wikimedia commons).

Two pictures are shown labeled a and b. Picture a depicts a clear, colorless solution in a plastic bag being held in a person's hand. Picture b shows a person's hand holding a detection meter with a digital readout screen while another hand holds someone's finger up to the end of the meter. The meter is pressed to the drop of blood that is at the end of the person's finger.

Example 9.3.4: Parts per Million and Parts per Billion Concentrations

According to the EPA, when the concentration of lead in tap water reaches 15 ppb, certain remedial actions must be taken. What is this concentration in ppm? At this concentration, what mass of lead (μg) would be contained in a typical glass of water (300 mL)?

Solution

The definitions of the ppm and ppb units may be used to convert the given concentration from ppb to ppm. Comparing these two unit definitions shows that ppm is 1000 times greater than ppb (1 ppm = 10^3 ppb). Thus:

$$15 \cancel{\text{ ppb}} \times \frac{1 \text{ ppm}}{10^3 \cancel{\text{ ppb}}} = 0.015 \text{ ppm}$$

The definition of the ppb unit may be used to calculate the requested mass if the mass of the solution is provided. However, only the volume of solution (300 mL) is given, so we must use the density to derive the corresponding mass. We can assume the density of tap water to be roughly the same as that of pure water (~1.00 g/mL), since the concentrations of any dissolved substances should not be very large. Rearranging the equation defining the ppb unit and substituting the given quantities yields:

$$\text{ppb} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^9 \text{ ppb}$$

$$\text{mass solute} = \frac{\text{ppb} \times \text{mass solution}}{10^9 \text{ ppb}}$$

$$\text{mass solute} = \frac{15 \text{ ppb} \times 300 \text{ mL} \times \frac{1.00 \text{ g}}{\text{mL}}}{10^9 \text{ ppb}} = 4.5 \times 10^{-6} \text{ g}$$

Finally, convert this mass to the requested unit of micrograms:

$$4.5 \times 10^{-6} \text{ g} \times \frac{1 \ \mu\text{g}}{10^{-6} \text{ g}} = 4.5 \ \mu\text{g}$$

? Exercise 9.3.4

A 50.0-g sample of industrial wastewater was determined to contain 0.48 mg of mercury. Express the mercury concentration of the wastewater in ppm and ppb units.

Answer

9.6 ppm, 9600 ppb

Summary

In addition to molarity, a number of other solution concentration units are used in various applications. Percentage concentrations based on the solution components' masses, volumes, or both are useful for expressing relatively high concentrations, whereas lower concentrations are conveniently expressed using ppm or ppb units. These units are popular in environmental, medical, and other fields where mole-based units such as molarity are not as commonly used.

Glossary

mass percentage
ratio of solute-to-solution mass expressed as a percentage

mass-volume percent
ratio of solute mass to solution volume, expressed as a percentage

parts per billion (ppb)
ratio of solute-to-solution mass multiplied by 10^9

parts per million (ppm)
ratio of solute-to-solution mass multiplied by 10^6

volume percentage
ratio of solute-to-solution volume expressed as a percentage

This page titled 9.3: Other Units for Solution Concentrations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.4: Other Units for Solution Concentrations** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/chemistry-2e.

9.4: The Dissolution Process

http://chemwiki.ucdavis.edu/Under_Construction/GOB/09%3A_Solutions/9.3_The_Dissolution_Process

Learning Objectives

- To describe the dissolution process at the molecular level

The Dissolution Process

What occurs at the molecular level to cause a solute to dissolve in a solvent? The answer depends in part on the solute, but there are some similarities common to all solutes.

Recall the rule that *like dissolves like*. This means that substances must have similar intermolecular forces to form solutions. When a soluble solute is introduced into a solvent, the particles of solute can interact with the particles of solvent. In the case of a solid or liquid solute, the interactions between the solute particles and the solvent particles are so strong that the individual solute particles separate from each other and, surrounded by solvent molecules, enter the solution. (Gaseous solutes already have their constituent particles separated, but the concept of being surrounded by solvent particles still applies.) This process is called **solvation** and is illustrated in Figure **9.4.1**. When the solvent is water, the word **hydration**, rather than solvation, is used.




Figure **9.4.1**: Solvation. When a solute dissolves, the individual particles of solute become surrounded by solvent particles. Eventually the particle detaches from the remaining solute, surrounded by solvent molecules in solution. Source: Photo © Thinkstock

Ionic Compounds and Covalent Compounds as Solutes

In the case of molecular solutes like glucose, the solute particles are individual molecules. However, if the solute is ionic, the individual ions separate from each other and become surrounded by solvent particles. That is, the cations and anions of an ionic solute separate when the solute dissolves. This process is referred to as **dissociation** (Figure **9.4.1**).

The dissociation of soluble ionic compounds gives solutions of these compounds an interesting property: they conduct electricity. Because of this property, soluble ionic compounds are referred to as electrolytes. Many ionic compounds dissociate completely and are therefore called **strong electrolytes**. Sodium chloride is an example of a strong electrolyte. Some compounds dissolve but dissociate only partially, and solutions of such solutes may conduct electricity only weakly. These solutes are called **weak electrolytes**. Acetic acid (CH_3COOH), the compound in vinegar, is a weak electrolyte. Solutes that dissolve into individual neutral molecules without dissociation do not impart additional electrical conductivity to their solutions and are called nonelectrolytes. Table sugar ($C_{12}H_{22}O_{11}$) is an example of a nonelectrolyte.

> *The term electrolyte is used in medicine to mean any of the important ions that are dissolved in aqueous solution in the body. Important physiological electrolytes include* Na^+, K^+, Ca^{2+}, Mg^{2+}, *and* Cl^-.

Example 9.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

1. potassium chloride (KCl)
2. fructose ($C_6H_{12}O_6$)
3. isopropyl alcohol [$CH_3CH(OH)CH_3$]
4. magnesium hydroxide [$Mg(OH)_2$]

Solution

Each substance can be classified as an ionic solute or a nonionic solute. Ionic solutes are electrolytes, and nonionic solutes are nonelectrolytes.

1. Potassium chloride is an ionic compound; therefore, when it dissolves, its ions separate, making it an electrolyte.
2. Fructose is a sugar similar to glucose. (In fact, it has the same molecular formula as glucose.) Because it is a molecular compound, we expect it to be a nonelectrolyte.
3. Isopropyl alcohol is an organic molecule containing the alcohol functional group. The bonding in the compound is all covalent, so when isopropyl alcohol dissolves, it separates into individual molecules but not ions. Thus, it is a nonelectrolyte
4. Magnesium hydroxide is an ionic compound, so when it dissolves it dissociates. Thus, magnesium hydroxide is an electrolyte.

Exercise 9.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

a. acetone (CH_3COCH_3)
b. iron(III) nitrate [$Fe(NO_3)_3$]
c. elemental bromine (Br_2)
d. sodium hydroxide (NaOH)

Answer

a. nonelectrolyte

b. electrolyte

c. nonelectrolyte

d. electrolyte

Electrolytes in Body Fluids

Our body fluids are solutions of electrolytes and many other things. The combination of blood and the circulatory system is the *river of life*, because it coordinates all the life functions. When the heart stops pumping in a heart attack, the life ends quickly. Getting the heart restarted as soon as one can is crucial in order to maintain life.

The primary electrolytes required in the body fluid are cations (of calcium, potassium, sodium, and magnesium) and anions (of chloride, carbonates, aminoacetates, phosphates, and iodide). These are nutritionally called **macrominerals**.

Electrolyte balance is crucial to many body functions. Here's some extreme examples of what can happen with an imbalance of electrolytes: elevated potassium levels may result in cardiac arrhythmias; decreased extracellular potassium produces paralysis; excessive extracellular sodium causes fluid retention; and decreased plasma calcium and magnesium can produce muscle spasms of the extremities.

When a patient is dehydrated, a carefully prepared (commercially available) electrolyte solution is required to maintain health and well being. In terms of child health, oral electrolyte is given when a child is dehydrated due to diarrhea. The use of oral electrolyte maintenance solutions, which is responsible for saving millions of lives worldwide over the last 25 years, is one of the most

important medical advances in protecting the health of children in the century, explains Juilus G.K. Goepp, MD, assistant director of the Pediatric Emergency Department of the Children's Center at Johns Hopkins Hospital. If a parent provides an oral electrolyte maintenance solution at the very start of the illness, dehydration can be prevented. The functionality of electrolyte solutions is related to their properties, and interest in electrolyte solutions goes far beyond chemistry.

Sports drinks are designed to rehydrate the body after excessive fluid depletion. Electrolytes in particular promote normal rehydration to prevent fatigue during physical exertion. Are they a good choice for achieving the recommended fluid intake? Are they performance and endurance enhancers like they claim? Who should drink them?

Typically, eight ounces of a sports drink provides between fifty and eighty calories and 14 to 17 grams of carbohydrate, mostly in the form of simple sugars. Sodium and potassium are the most commonly included electrolytes in sports drinks, with the levels of these in sports drinks being highly variable. The American College of Sports Medicine says a sports drink should contain 125 milligrams of sodium per 8 ounces as it is helpful in replenishing some of the sodium lost in sweat and promotes fluid uptake in the small intestine, improving hydration.

Gatorade

In the summer of 1965, the assistant football coach of the University of Florida Gators requested scientists affiliated with the university study why the withering heat of Florida caused so many heat-related illnesses in football players and provide a solution to increase athletic performance and recovery post-training or game. The discovery was that inadequate replenishment of fluids, carbohydrates, and electrolytes was the reason for the “wilting” of their football players. Based on their research, the scientists concocted a drink for the football players containing water, carbohydrates, and electrolytes and called it “Gatorade.” In the next football season the Gators were nine and two and won the Orange Bowl. The Gators’ success launched the sports-drink industry, which is now a multibillion-dollar industry that is still dominated by Gatorade.



University of Florida football player Chip Hinton testing Gatorade in 1965, pictured next to the leader of its team of inventors, Robert Cade.

Key Takeaway

- When a solute dissolves, its individual particles are surrounded by solvent molecules and are separated from each other.

9.4: The Dissolution Process is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **9.3: The Dissolution Process** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.5: Properties of Solutions

Learning Objectives

- To describe how the properties of solutions differ from those of pure solvents.

Solutions are likely to have properties similar to those of their major component—usually the solvent. However, some solution properties differ significantly from those of the solvent. Here, we will focus on liquid solutions that have a solid solute, but many of the effects we will discuss in this section are applicable to all solutions.

Colligative Properties

Solutes affect some properties of solutions that depend only on the concentration of the dissolved particles. These properties are called **colligative properties**. Four important colligative properties that we will examine here are vapor pressure depression, boiling point elevation, freezing point depression, and osmotic pressure.

Molecular compounds separate into individual molecules when they are dissolved, so for every 1 mol of molecules dissolved, we get 1 mol of particles. In contrast, **ionic** compounds separate into their constituent ions when they dissolve, so 1 mol of an ionic compound will produce more than 1 mol of dissolved particles. For example, **every mole of NaCl** that dissolves yields **1 mol of Na^+ ions** and **1 mol of Cl^- ions**, for a **total of 2 mol** of particles in solution. Thus, the effect on a solution's properties by dissolving NaCl may be twice as large as the effect of dissolving the same amount of moles of glucose ($C_6H_{12}O_6$).

Vapor Pressure Depression

All liquids evaporate. In fact, given enough volume, a liquid will turn completely into a vapor. If enough volume is not present, a liquid will evaporate only to the point where the rate of evaporation equals the rate of vapor condensing back into a liquid. The pressure of the vapor at this point is called the **vapor pressure** of the liquid.

The presence of a dissolved solid lowers the characteristic vapor pressure of a liquid so that it evaporates more slowly. (The exceptions to this statement are if the solute itself is a liquid or a gas, in which case the solute will also contribute something to the evaporation process. We will not discuss such solutions here.) This property is called vapor pressure depression and is depicted in Figure **9.5.1**.




Figure **9.5.1**: Vapor Pressure Depression. The presence of solute particles blocks some of the ability for liquid particles to evaporate. Thus, solutions of solid solutes typically have a lower vapor pressure than the pure solvent.

Boiling Point and Freezing Point Effects

A related property of solutions is that their boiling points are higher than the boiling point of the pure solvent. Because the presence of solute particles decreases the vapor pressure of the liquid solvent, a higher temperature is needed to reach the boiling point. This phenomenon is called boiling point elevation. **For every mole of particles dissolved in a liter of water, the boiling point of water increases by about 0.5°C.** The addition of one mole of sucrose (molecular compound) in one liter of water will raise the boiling point from 100^0C to 100.5^0C but the addition of one mole of NaCl in one liter of water will raise the boiling point by 2 x 0.5^0C = 1^0C. Furthermore, the addition of one mole of $CaCl_2$ in one liter of water will raise the boiling point by 3 x 0.5^0C = 1.5^0C.

Some people argue that putting a pinch or two of salt in water used to cook spaghetti or other pasta makes a solution that has a higher boiling point, so the pasta cooks faster. In actuality, the amount of solute is so small that the boiling point of the water is practically unchanged.

The presence of solute particles has the opposite effect on the freezing point of a solution. When a solution freezes, only the solvent particles come together to form a solid phase, and the presence of solute particles interferes with that process. Therefore, for the liquid solvent to freeze, more energy must be removed from the solution, which lowers the temperature. Thus, solutions have lower freezing points than pure solvents do. This phenomenon is called freezing point depression. **For every mole of particles in a liter of water, the freezing point decreases by about 1.9°C.**




Figure **9.5.2**: Comparison of boiling and freezing points of a pure liquid (right side) with a solution (left side).

Both boiling point elevation and freezing point depression have practical uses. For example, solutions of water and ethylene glycol ($C_2H_6O_2$) are used as coolants in automobile engines because the boiling point of such a solution is greater than 100°C, the normal boiling point of water. In winter, salts like NaCl and $CaCl_2$ are sprinkled on the ground to melt ice or keep ice from forming on roads and sidewalks (Figure **9.5.2**). This is because the solution made by dissolving sodium chloride or calcium chloride in water has a lower freezing point than pure water, so the formation of ice is inhibited.



Figure **9.5.2**: Effect of Freezing Point Depression. The salt sprinkled on this sidewalk makes the water on the sidewalk have a lower freezing point than pure water, so it does not freeze as easily. This makes walking on the sidewalk less hazardous in winter. © Thinkstock

Example 9.5.1

Which solution's freezing point deviates more from that of pure water—a 1 M solution of NaCl or a 1 M solution of $CaCl_2$?

Solution

Colligative properties depend on the number of dissolved particles, so the solution with the greater number of particles in solution will show the greatest deviation. When NaCl dissolves, it separates into two ions, Na^+ and Cl^-. But when $CaCl_2$ dissolves, it separates into three ions—one Ca^{2+} ion and two Cl^- ions. Thus, mole for mole, $CaCl_2$ will have 50% more impact on freezing point depression than NaCl.

? Exercise 9.5.1

Which solution's boiling point deviates more from that of pure water—a 1 M solution of $\mathbf{CaCl_2}$ or a 1 M solution of $MgSO_4$?

Answer

$\mathbf{CaCl_2}$

✔ Example 9.5.2

Estimate the boiling point of 0.2 M $\mathbf{CaCl_2}$ solution.

Solution

The boiling point increases 0.5^0C for every mole of solute per liter of water. For this estimation, let's assume that 1 liter of solution is roughly the same volume as 1 liter of water. A 0.2 M $\mathbf{CaCl_2}$ solution contains 0.2 moles of $\mathbf{CaCl_2}$ solution formula units per liter of solution. Each $\mathbf{CaCl_2}$ unit separates into three ions.

$$0.2 \text{ mol } \mathbf{CaCl_2} \times \frac{3 \text{ mol ions}}{1 \text{ mol } \mathbf{CaCl_2}} \times \frac{0.5 \text{ deg C}}{1 \text{ mol ion}} = 0.3 \text{ deg C}$$

The normal boiling point of water is 100^0C, so the boiling point of the solution is raised to 100.3^0C.

9.5.2Exercise

Estimate the freezing point of 0.3 M $\mathbf{CaCl_2}$ solution.

Answer

minus 1.7^0C

Osmotic Pressure

The last colligative property of solutions we will consider is a very important one for biological systems. It involves **osmosis**, the process by which solvent molecules can pass through certain membranes but solute particles cannot. When two solutions of different concentration are present on either side of these membranes (called *semipermeable membranes*), there is a tendency for solvent molecules to move from the more dilute solution to the more concentrated solution until the concentrations of the two solutions are equal. This tendency is called **osmotic pressure**. External pressure can be exerted on a solution to counter the flow of solvent; the pressure required to halt the osmosis of a solvent is equal to the osmotic pressure of the solution.

Osmolarity (osmol) is a way of reporting the total number of particles in a solution to determine osmotic pressure. It is defined as the molarity of a solute times the number of particles a formula unit of the solute makes when it dissolves (represented by i):

$$osmol = M \times i \tag{9.5.1}$$

If more than one solute is present in a solution, the individual osmolarities are additive to get the total osmolarity of the solution. Solutions that have the same osmolarity have the same osmotic pressure. If solutions of differing osmolarities are present on opposite sides of a semipermeable membrane, solvent will transfer from the lower-osmolarity solution to the higher-osmolarity solution. Counterpressure exerted on the high-osmolarity solution will reduce or halt the solvent transfer. An even higher pressure can be exerted to force solvent from the high-osmolarity solution to the low-osmolarity solution, a process called *reverse osmosis*. Reverse osmosis is used to make potable water from saltwater where sources of fresh water are scarce.

✔ Example 9.5.3

A 0.50 M NaCl aqueous solution and a 0.30 M $Ca(NO_3)_2$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Solution

The solvent will flow into the solution of higher osmolarity. The NaCl solute separates into two ions—Na^+ and Cl^-—when it dissolves, so its osmolarity is as follows:

$$\text{osmol (NaCl)} = 0.50\ \text{M} \times 2 = 1.0\ \text{osmol}$$

The $Ca(NO_3)_2$ solute separates into three ions—one Ca^{2+} and two NO_3^-—when it dissolves, so its osmolarity is as follows:

$$\text{osmol } [Ca(NO_3)_2] = 0.30\ \text{M} \times 3 = 0.90\ \text{osmol}$$

The osmolarity of the $Ca(NO_3)_2$ solution is lower than that of the NaCl solution, so water will transfer through the membrane from the $Ca(NO_3)_2$ solution to the NaCl solution.

? Exercise 9.5.3

A 1.5 M $C_6H_{12}O_6$ aqueous solution and a 0.40 M $Al(NO_3)_3$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Answer

osmol $C_6H_{12}O_6$ = 1.5; osmol $Al(NO_3)_3$ = 1.6

The solvent flows from $C_6H_{12}O_6$ solution (lower osmolarity) to $\mathbf{Al(NO_3)_3}$ solution (higher osmolarity).

To Your Health: Dialysis

The main function of the kidneys is to filter the blood to remove wastes and extra water, which are then expelled from the body as urine. Some diseases rob the kidneys of their ability to perform this function, causing a buildup of waste materials in the bloodstream. If a kidney transplant is not available or desirable, a procedure called dialysis can be used to remove waste materials and excess water from the blood.

In one form of dialysis, called *hemodialysis*, a patient's blood is passed though a length of tubing that travels through an *artificial kidney machine* (also called a *dialysis machine*). A section of tubing composed of a semipermeable membrane is immersed in a solution of sterile water, glucose, amino acids, and certain electrolytes. The osmotic pressure of the blood forces waste molecules and excess water through the membrane into the sterile solution. Red and white blood cells are too large to pass through the membrane, so they remain in the blood. After being cleansed in this way, the blood is returned to the body.



Figure **9.5.3**: A patient undergoing hemodialysis depends on osmosis to cleanse the blood of waste products that the kidneys are incapable of removing due to disease. from Wikipedia.

Dialysis is a continuous process, as the osmosis of waste materials and excess water takes time. Typically, 5–10 lb of waste-containing fluid is removed in each dialysis session, which can last 2–8 hours and must be performed several times a week. Although some patients have been on dialysis for 30 or more years, dialysis is always a temporary solution because waste materials are constantly building up in the bloodstream. A more permanent solution is a kidney transplant.

Cell walls are semipermeable membranes, so the osmotic pressures of the body's fluids have important biological consequences. If solutions of different osmolarity exist on either side of the cells, solvent (water) may pass into or out of the cells, sometimes with disastrous results. Consider what happens if red blood cells are placed in a *hypotonic* solution, meaning a solution of lower

osmolarity than the liquid inside the cells. The cells swell up as water enters them, disrupting cellular activity and eventually causing the cells to burst. This process is called *hemolysis*. If red blood cells are placed in a *hypertonic* solution, meaning one having a higher osmolarity than exists inside the cells, water leaves the cells to dilute the external solution, and the red blood cells shrivel and die. This process is called *crenation*. Only if red blood cells are placed in *isotonic* solutions that have the same osmolarity as exists inside the cells are they unaffected by negative effects of osmotic pressure. **Glucose solutions of about 0.31 M, or sodium chloride solutions of about 0.16 M, are isotonic with blood plasma.**

> The concentration of an isotonic sodium chloride (NaCl) solution is only half that of an isotonic glucose ($C_6H_{12}O_6$) solution because NaCl produces two ions when a formula unit dissolves, while molecular $C_6H_{12}O_6$ produces only one particle when a formula unit dissolves. The osmolarities are therefore the same even though the concentrations of the two solutions are different.

Osmotic pressure explains why you should not drink seawater if you are abandoned in a life raft in the middle of the ocean. Its osmolarity is about three times higher than most bodily fluids. You would actually become thirstier as water from your cells was drawn out to dilute the salty ocean water you ingested. Our bodies do a better job coping with hypotonic solutions than with hypertonic ones. The excess water is collected by our kidneys and excreted.

Osmotic pressure effects are used in the food industry to make pickles from cucumbers and other vegetables and in brining meat to make corned beef. It is also a factor in the mechanism of getting water from the roots to the tops of trees!

Career Focus: Perfusionist

A perfusionist is a medical technician trained to assist during any medical procedure in which a patient's circulatory or breathing functions require support. The use of perfusionists has grown rapidly since the advent of open-heart surgery in 1953.

Most perfusionists work in operating rooms, where their main responsibility is to operate heart-lung machines. During many heart surgeries, the heart itself must be stopped. In these situations, a heart-lung machine keeps the patient alive by aerating the blood with oxygen and removing carbon dioxide. The perfusionist monitors both the machine and the status of the blood, notifying the surgeon and the anesthetist of any concerns and taking corrective action if the status of the blood becomes abnormal.

Despite the narrow parameters of their specialty, perfusionists must be highly trained. Certified perfusion education programs require a student to learn anatomy, physiology, pathology, chemistry, pharmacology, math, and physics. A college degree is usually required. Some perfusionists work with other external artificial organs, such as hemodialysis machines and artificial livers.

9.5: Properties of Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **9.4: Properties of Solutions** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.6: Dialysis and Tonicity

Dialysis is the separation of colloids from dissolved ions or molecules of small dimensions, or crystalloid, in a solution. A **colloid** is any substance that is made of particles that are of an extremely small size: larger than atoms but generally have the size of 10^{-7} cm ranging to 10^{-3} cm. A **crystalloid** is a substance that has some or all of the properties of a crystal or a substance that forms a true solution and diffuses through a membrane by dialysis. Dialysis is a process that is like osmosis. Osmosis is the process in which there is a diffusion of a solvent through a semipermeable membrane.

Introduction

In 1861, chemist Thomas Graham used the process of dialysis, a process used to separate colloidal particles from dissolved ions or molecules. Thomas Graham is also responsible for developing Graham's Law. Dialysis is possible because of the *unequal rates of diffusion* through a **semipermeable membrane** (Figure **9.6.1**). A semipermeable membrane is a membrane that lets some molecules to pass through it while not letting others. Examples of semipermeable membranes include parchment and cellophane.



Another way to think of a semipermeable membrane is to think of a net like object that traps larger objects, but lets smaller object pass through because they can pass through the holes in the net.



***Video* 9.6.1**: *Osmotic Pressure Dialysis Tubing*

When a colloidal mixture is places in a semipermeable membrane, which is then placed in an aqueous solution or pure water, dissolved ions and small molecules are allowed to pass through this membrane. This causes colloidal particles to stay in the membrane, because these particles are unable to pass through the small pores of the membrane.

Dialysis is not a quick process; the rate of dialysis depends on the speed of the unequal diffusion rates between the crystalloids and the colloids and the differences in particle size. The rate of dialysis can be changed through **heating**, or if the crystalloids are charged, then **applying an electric field**, called electrodialysis. Electrodialysis is the type of dialysis in which electrodes are placed on the sides of the membrane. In this way, positive

ions can pass through one side of this membrane while the negatively charged ions can pass through the other side of the membrane. This causes acceleration in the process of dialysis.

Hemodialysis

Hemodialysis is a method in which kidney failure is treated with the process of dialysis. In hemodialysis, blood is removed, purified through dialysis, and returned to the bloodstream. In kidney failure, there is a retention of salts and water, urea, and metabolic acids. The patient is then connected to a dialysis machine, which is also called a hemodialyzer. The blood flows through small channels made of semipermeable membranes. The dissolved substances like urea and salts pass through a sterile solution. Compounds like sugar and amino acids are added to the sterile solution. The dialysis solution is on the other side of the membranes, and the molecules flow through the membranes. The molecules diffuse from a higher concentration to a low concentration area. The concentrations of molecules needed to be removed from the blood are zero in the dialysis fluid.




***Figure* 9.6.2:** *(left) Scheme of semipermeable membrane during hemodialysis, where blood is red, dialysing fluid is blue, and the membrane is yellow. Image used with permission from Wikipedia (Freemesm). (right) Hemodialysis setup.* Blausen.com staff. "Blausen gallery 2014". Wikiversity Journal of Medicine. DOI:10.15347/wjm/2014.010. ISSN 20018762. *See a full animation of this medical topic.*

The process of hemodialysis helps many patients who have kidney failure because a person who suffers from kidney failure are at great risk, because someone who has complete kidney failure will need a kidney transplant within two weeks, or else he/she will face death. Between the time that the person finds a suitable kidney to be transplanted, the hemodialyzer comes into great help in facing the fight against death.

References

1. Petrucci, et al. *General Chemistry Principles & Modern Applications*. 9th ed. Upper Saddle River, NJ: Pearson Prentice Hall, 2007.
2. Sadava, et al. *Life: The Science of Biology*. 8th ed. New York, NY. W.H. Freeman and Company, 2007.

Contributors

- Katrina Lau (UCD)

9.6: Dialysis and Tonicity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

9.7: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **solution** is a homogeneous mixture. The major component is the **solvent**, while the minor component is the **solute**. Solutions can have any phase; for example, an **alloy** is a solid solution. Solutes are **soluble** or **insoluble**, meaning they dissolve or do not dissolve in a particular solvent. The terms **miscible** and **immiscible**, instead of soluble and insoluble, are used for liquid solutes and solvents. The statement *like dissolves like* is a useful guide to predicting whether a solute will dissolve in a given solvent.

The amount of solute in a solution is represented by the **concentration** of the solution. The maximum amount of solute that will dissolve in a given amount of solvent is called the **solubility** of the solute. Such solutions are **saturated**. Solutions that have less than the maximum amount are **unsaturated**. Most solutions are unsaturated, and there are various ways of stating their concentrations. **Mass/mass percent**, **volume/volume percent**, and **mass/volume percent** indicate the percentage of the overall solution that is solute. **Parts per million (ppm)** and **parts per billion (ppb)** are used to describe very small concentrations of a solute. **Molarity**, defined as the number of moles of solute per liter of solution, is a common concentration unit in the chemistry laboratory. **Equivalents** express concentrations in terms of moles of charge on ions. When a solution is diluted, we use the fact that the amount of solute remains constant to be able to determine the volume or concentration of the final diluted solution.

Dissolving occurs by **solvation**, the process in which particles of a solvent surround the individual particles of a solute, separating them to make a solution. For water solutions, the word **hydration** is used. If the solute is molecular, it dissolves into individual molecules. If the solute is ionic, the individual ions separate from each other, forming a solution that conducts electricity. Such solutions are called **electrolytes**. If the dissociation of ions is complete, the solution is a **strong electrolyte**. If the dissociation is only partial, the solution is a **weak electrolyte**. Solutions of molecules do not conduct electricity and are called **nonelectrolytes**.

Solutions have properties that differ from those of the pure solvent. Some of these are **colligative** properties, which are due to the number of solute particles dissolved, not the chemical identity of the solute. Colligative properties include **vapor pressure depression**, **boiling point elevation**, **freezing point depression**, and **osmotic pressure**. Osmotic pressure is particularly important in biological systems. It is caused by **osmosis**, the passage of solvents through certain membranes like cell walls. The **osmolarity** of a solution is the product of a solution's molarity and the number of particles a solute separates into when it dissolves. Osmosis can be reversed by the application of pressure; this reverse osmosis is used to make fresh water from saltwater in some parts of the world. Because of osmosis, red blood cells placed in hypotonic or hypertonic solutions lose function through either hemolysis or crenation. If they are placed in isotonic solutions, however, the cells are unaffected because osmotic pressure is equal on either side of the cell membrane.

Template:HideTOC

9.7: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

10: Acids and Bases

Many of us are familiar with the group of chemicals called acids. But do you know what it takes for a compound to be an acid? Actually, there are several different definitions of acid that chemistry uses, and each definition is appropriate under different circumstances. Less familiar—but just as important to chemistry and ultimately to us—is the group of chemicals known as bases. Both acids and bases are important enough that we devote an entire chapter to them—their properties and their reactions.

10.1: Arrhenius Definition of Acids and Bases

10.2: Brønsted-Lowry Definition of Acids and Bases

10.3: Water is both an Acid and a Base

10.4: The Strengths of Acids and Bases

10.5: pH and Kw

10.6: Buffers

10.7: End-of-Chapter Material


10: Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

10.1: Arrhenius Definition of Acids and Bases

Learning Objectives

- To recognize a compound as an Arrhenius acid or an Arrhenius base.
- To describe characteristics of acids and bases.
- To write equations of neutralization reactions.

One way to define a class of compounds is by describing the various characteristics its members have in common. In the case of the compounds known as acids, the common characteristics include a sour taste, the ability to change the color of the vegetable dye *litmus* to red, and the ability to dissolve certain metals and simultaneously produce hydrogen gas. For the compounds called bases, the common characteristics are a slippery texture, a bitter taste, and the ability to change the color of litmus to blue. Acids and bases also react with each other to form compounds generally known as salts.



Although we include their tastes among the common characteristics of acids and bases, we never advocate tasting an unknown chemical!

Chemists prefer, however, to have definitions for acids and bases in chemical terms. The Swedish chemist Svante Arrhenius developed the first chemical definitions of acids and bases in the late 1800s. Arrhenius defined an acid as a compound that increases the concentration of hydrogen ion (H^+) in aqueous solution. Many acids are simple compounds that release a hydrogen cation into solution when they dissolve. Similarly, Arrhenius defined a base as a compound that increases the concentration of hydroxide ion (OH^-) in aqueous solution. Many bases are ionic compounds that have the hydroxide ion as their anion, which is released when the base dissolves in water.

Table **10.1.1**: Formulas and Names for Some Acids and Bases

Acids		Bases	
Formula	Name	Formula	Name
HCl(aq)	hydrochloric acid	NaOH(aq)	sodium hydroxide
HBr(aq)	hydrobromic acid	KOH(aq)	potassium hydroxide
HI(aq)	hydriodic acid	$Mg(OH)_2$(aq)	magnesium hydroxide
H_2S(aq)	hydrosulfuric acid	$Ca(OH)_2$(aq)	calcium hydroxide
$HC_2H_3O_2$(aq)	acetic acid	NH_3(aq)	ammonia
HNO_3(aq)	nitric acid	$NaHCO_3$ (aq)	sodium bicarbonate
HNO_2(aq)	nitrous acid	$CaCO_3$ (aq)	calcium carbonate
H_2SO_4(aq)	sulfuric acid		
H_2SO_3(aq)	sulfurous acid		
$HClO_3$(aq)	chloric acid		
$HClO_4$(aq)	perchloric acid		
$HClO_2$(aq)	chlorous acid		
H_3PO_4(aq)	phosphoric acid		
H_3PO_3(aq)	phosphorous acid		
H_2CO_3(aq)	carbonic acid		

Many bases and their aqueous solutions are named using the normal rules of ionic compounds that were presented previously; that is, they are named as hydroxide compounds. For example, the base sodium hydroxide (NaOH) is both an ionic compound and an aqueous solution. However, aqueous solutions of acids have their own naming rules. The names of *binary acids* (compounds with hydrogen and one other element in their formula) are based on the root of the name of the other element preceded by the prefix *hydro-* and followed by the suffix *-ic acid*. Thus, an aqueous solution of HCl [designated "HCl(aq)"] is called hydrochloric acid, H_2S(aq) is called hydrosulfuric acid, and so forth. Acids composed of more than two elements (typically hydrogen and oxygen and some other element) have names based on the name of the other element, followed by the suffix *-ic acid* or *-ous acid*, depending on the number of oxygen atoms in the acid's formula. Other prefixes, like *per-* and *hypo-*, also appear in the names for some acids. Unfortunately, there is no strict rule for the number of oxygen atoms that are associated with the *-ic acid* suffix; the names of these acids are best memorized. Table **10.1.1** lists some acids and bases and their names. Note that acids have hydrogen written first, as if it were the cation, while most bases have the negative hydroxide ion, if it appears in the formula, written last.

> *The name oxygen comes from the Latin meaning "acid producer" because its discoverer, Antoine Lavoisier, thought it was the essential element in acids. Lavoisier was wrong, but it is too late to change the name now.*

Example 10.1.1

Name each substance.

a. HF(aq)
b. $Sr(OH)_2$(aq)

Solution

a. This acid has only two elements in its formula, so its name includes the *hydro-* prefix. The stem of the other element's name, fluorine, is *fluor*, and we must also include the *-ic acid* ending. Its name is hydrofluoric acid.

b. This base is named as an ionic compound between the strontium ion and the hydroxide ion: strontium hydroxide.

? Exercise 10.1.1

Name each substance.

a. $H_2Se(aq)$

b. $Ba(OH)_2(aq)$

Answer

a. hydroselenic acid

b. barium hydroxide

Notice that one base listed in Table 10.1.1—ammonia—does not have hydroxide as part of its formula. How does this compound increase the amount of hydroxide ion in aqueous solution? Instead of dissociating into hydroxide ions, ammonia molecules react with water molecules by taking a hydrogen ion from the water molecule to produce an ammonium ion and a hydroxide ion:

$$NH_{3(aq)} + H_2O_{(\ell)} \rightarrow NH^+_{4(aq)} + OH^-_{(aq)} \tag{10.1.1}$$

Because this reaction of ammonia with water causes an increase in the concentration of hydroxide ions in solution, ammonia satisfies the Arrhenius definition of a base. Many other nitrogen-containing compounds are bases because they too react with water to produce hydroxide ions in aqueous solution.

Neutralization

As we noted previously, acids and bases react chemically with each other to form *salts*. A salt is a general chemical term for any ionic compound formed from an acid and a base. In reactions where the acid is a hydrogen ion containing compound and the base is a hydroxide ion containing compound, water is also a product. The general reaction is as follows:

$$\text{acid} + \text{base} \rightarrow \text{water} + \text{salt}$$

The reaction of acid and base to make water and a salt is called **neutralization**. Like any chemical equation, a neutralization chemical equation must be properly balanced. For example, the neutralization reaction between sodium hydroxide and hydrochloric acid is as follows:

$$NaOH(aq) + HCl_{(aq)} \rightarrow NaCl_{(aq)} + H_2O_{(\ell)} \tag{10.1.2}$$

with coefficients all understood to be one. The neutralization reaction between sodium hydroxide and sulfuric acid is as follows:

$$2NaOH_{(aq)} + H_2SO_{4(aq)} \rightarrow Na_2SO_{4(aq)} + 2H_2O_{(\ell)} \tag{10.1.3}$$

Once a neutralization reaction is properly balanced, we can use it to perform stoichiometry calculations, such as the ones we practiced earlier.

There are a number of examples of acid-base chemistry in everyday life. One example is the use of baking soda, or sodium bicarbonate in baking. $NaHCO_3$ is a base. When it reacts with an acid such as lemon juice, buttermilk, or sour cream in a batter, bubbles of carbon dioxide gas are formed from decomposition of the resulting carbonic acid, and the batter "rises." Baking powder is a combination of sodium bicarbonate, and one or more acid salts that react when the two chemicals come in contact with water in the batter.

$$HCO^-_3(aq) + H^+(aq) \rightarrow H_2CO_3(aq) \tag{10.1.4}$$

$$H_2CO_3(aq) \rightarrow CO_2(g) + H_2O(l)$$

Example 10.1.2

Nitric acid [$HNO_3(aq)$] can be neutralized by calcium hydroxide [$Ca(OH)_2(aq)$].


a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 16.8 g of HNO_3 is present initially. How many grams of $Ca(OH)_2$ are needed to neutralize that much HNO_3?
c. In a second reaction, 805 mL of 0.672 M $Ca(OH)_2$ is present initially. What volume of 0.432 M HNO_3 solution is necessary to neutralize the $Ca(OH)_2$ solution?

Solution

a. Because there are two OH^- ions in the formula for $Ca(OH)_2$, we need two moles of HNO_3 to provide H^+ ions. The balanced chemical equation is as follows: $Ca(OH)_2(aq) + 2HNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + 2H_2O(\ell)$

The salt formed is calcium nitrate.

b. This calculation is much like the calculations we did in Chapter 6. First we convert the mass of HNO_3 to moles using its molar mass of 1.01 + 14.01 + 3(16.00) = 63.02 g/mol; then we use the balanced chemical equation to determine the related number of moles of $Ca(OH)_2$ needed to neutralize it; and then we convert that number of moles of $Ca(OH)_2$ to the mass of $Ca(OH)_2$ using its molar mass of 40.08 + 2(1.01) + 2(16.00) = 74.10 g/mol.



$$16.8\ \text{g HNO}_3 \times \frac{1\ \text{mol HNO}_3}{63.02\ \text{g HNO}_3} \times \frac{1\ \text{mol Ca(OH)}_2}{2\ \text{mol HNO}_3} \times \frac{74.10\ \text{g Ca(OH)}_2}{1\ \text{mol Ca(OH)}_2} = 9.88\ \text{g Ca(OH)}_2\ \text{needed}$$

c. Having concentration information allows us to employ the skills we developed in Chapter 9. We have two alternative solutions: the multi-step process and the combined one-line process (found at the bottom).

First, we use the concentration and volume data to determine the number of moles of $Ca(OH)_2$ present. Recognizing that 805 mL = 0.805 L,

$$0.672\ \text{M Ca(OH)}_2 = \frac{\text{mol Ca(OH)}_2}{0.805\ \text{L soln}}$$

$$0.672\ \text{M CaOH)}_2 \times (0.805\ \text{L soln}) = \text{mol Ca(OH)}_2 = 0.541\ \text{mol Ca(OH)}_2$$


We combine this information with the proper ratio from the balanced chemical equation to determine the number of moles of HNO_3 needed:

$$0.541\ \text{mol Ca(OH)}_2 \times \frac{2\ \text{mol HNO}_3}{1\ \text{mol Ca(OH)}_2} = 1.08\ \text{mol HNO}_3$$

Now, using the definition of molarity one more time, we determine the volume of acid solution needed:

$$0.432\ \text{M HNO}_3 = \frac{1.08\ \text{mol HNO}_3}{\text{volume of HNO}_3}$$

$$\text{volume of HNO}_3 = \frac{1.08\ \text{mol HNO}_3}{0.432\ \text{M HNO}_3} = 2.50\ \text{L} = 2.50 \times 10^3\ \text{mL HNO}_3$$



$$0.805\ \text{L Ca(OH)}_2\ \text{soln} \times \frac{0.672\ \text{mol Ca(OH)}_2}{1\ \text{L Ca(OH)}_2\ \text{soln}} \times \frac{2\ \text{mol HNO}_3}{1\ \text{mol Ca(OH)}_2} \times \frac{1\ \text{L HNO}_3\ \text{soln}}{0.432\ \text{mol HNO}_3} = 2.50\ \text{L HNO}_3\ \text{soln needed}$$

? Exercise 10.1.2

Hydrocyanic acid [HCN(aq)] can be neutralized by potassium hydroxide [KOH(aq)].

a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 37.5 g of HCN is present initially. How many grams of KOH are needed to neutralize that much HCN?
c. In a second reaction, 43.0 mL of 0.0663 M KOH is present initially. What volume of 0.107 M HCN solution is necessary to neutralize the KOH solution?

Answer

a. KOH(aq) + HCN(aq) → KCN(aq) + $H_2O(\ell)$ KCN is the salt.

b. 77.8 g

c. 0.0266 L or 26.6 mL

Hydrocyanic acid (HCN) is one exception to the acid-naming rules that specify using the prefix *hydro-* for binary acids (acids composed of hydrogen and only one other element).

Stomach Antacids

Our stomachs contain a solution of roughly 0.03 *M* HCl, which helps us digest the food we eat. The burning sensation associated with heartburn is a result of the acid of the stomach leaking through the muscular valve at the top of the stomach into the lower reaches of the esophagus. The lining of the esophagus is not protected from the corrosive effects of stomach acid the way the lining of the stomach is, and the results can be very painful. When we have heartburn, it feels better if we reduce the excess acid in the esophagus by taking an antacid. As you may have guessed, antacids are bases. One of the most common antacids is calcium carbonate, $CaCO_3$. The reaction,

$$CaCO_3(s) + 2HCl(aq) \rightleftharpoons CaCl_2(aq) + H_2O(l) + CO_2(g)$$

not only neutralizes stomach acid, it also produces $CO_2(g)$, which may result in a satisfying belch.

Milk of Magnesia is a suspension of the sparingly soluble base magnesium hydroxide, $Mg(OH)_2$. It works according to the reaction:

$$Mg(OH)_2(s) \rightleftharpoons Mg^{2+}(aq) + 2OH^-(aq)$$

The hydroxide ions generated in this equilibrium then go on to react with the hydronium ions from the stomach acid, so that :

$$H_3O^+ + OH^- \rightleftharpoons 2H_2O(l)$$

This reaction does not produce carbon dioxide, but magnesium-containing antacids can have a laxative effect. Several antacids have aluminum hydroxide, $Al(OH)_3$, as an active ingredient. The aluminum hydroxide tends to cause constipation, and some antacids use aluminum hydroxide in concert with magnesium hydroxide to balance the side effects of the two substances.

Example 10.1.3

Assume that the stomach of someone suffering from acid indigestion contains 75 mL of 0.20 M HCl. How many Tums tablets are required to neutralize 90% of the stomach acid, if each tablet contains 500 mg of $CaCO_3$? (Neutralizing all of the stomach acid is not desirable because that would completely shut down digestion.)

Solution

A. Write the balanced chemical equation for the reaction and then decide whether the reaction will go to completion.
B. Calculate the number of moles of acid present. Multiply the number of moles by the percentage to obtain the quantity of acid that must be neutralized. Using mole ratios, calculate the number of moles of base required to neutralize the acid.
C. Calculate the number of moles of base contained in one tablet by dividing the mass of base by the corresponding molar mass. Calculate the number of tablets required by dividing the moles of base by the moles contained in one tablet.

A. We first write the balanced chemical equation for the reaction:

$$2HCl(aq) + CaCO_3(s) \rightarrow CaCl_2(aq) + H_2CO_3(aq)$$

Each carbonate ion can react with 2 mol of H^+ to produce H_2CO_3, which rapidly decomposes to H_2O and CO_2. Because HCl is a strong acid and CO_3^{2-} is a weak base, the reaction will go to completion.

B. Next we need to determine the number of moles of HCl present:

$$75\ \cancel{mL}\left(\frac{1\ \cancel{L}}{1000\ \cancel{mL}}\right)\left(\frac{0.20\ mol\ HCl}{\cancel{L}}\right) = 0.015\ mol\ HCl$$

Because we want to neutralize only 90% of the acid present, we multiply the number of moles of HCl by 0.90:

$$(0.015\ mol\ HCl)(0.90) = 0.014\ mol\ HCl$$

We know from the stoichiometry of the reaction that each mole of $CaCO_3$ reacts with 2 mol of HCl, so we need

$$moles\ CaCO_3 = 0.014\ \cancel{mol\ HCl}\left(\frac{1\ mol\ CaCO_3}{2\ \cancel{mol\ HCl}}\right) = 0.0070\ mol\ CaCO_3$$

C. Each Tums tablet contains

$$\left(\frac{500\ \cancel{mg\ CaCO_3}}{1\ Tums\ tablet}\right)\left(\frac{1\ \cancel{g}}{1000\ \cancel{mg\ CaCO_3}}\right)\left(\frac{1\ mol\ CaCO_3}{100.1\ \cancel{g}}\right) = 0.00500\ mol\ CaCO_3$$

Thus we need $\dfrac{0.0070\ \cancel{mol\ CaCO_3}}{0.00500\ \cancel{mol\ CaCO_3}} = 1.4$ Tums tablets.

10.1.3 Exercise

Assume that as a result of overeating, a person's stomach contains 300 mL of 0.25 M HCl. How many Rolaids tablets must be consumed to neutralize 95% of the acid, if each tablet contains 400 mg of $NaAl(OH)_2CO_3$? The neutralization reaction can be written as follows:

$$NaAl(OH)_2CO_3(s) + 4HCl(aq) \rightarrow AlCl_3(aq) + NaCl(aq) + CO_2(g) + 3H_2O(l)$$

Answer

6.4 tablets

Key Takeaway

- An Arrhenius acid increases the H^+ ion concentration in water, while an Arrhenius base increases the OH^- ion concentration in water.

10.1: Arrhenius Definition of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.1: Arrhenius Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.2: Brønsted-Lowry Definition of Acids and Bases

Learning Objectives

1. Recognize a compound as a Brønsted-Lowry acid or a Brønsted-Lowry base.
2. Illustrate the proton transfer process that defines a Brønsted-Lowry acid-base reaction.


Ammonia (NH_3) increases the hydroxide ion concentration in aqueous solution by reacting with water rather than releasing hydroxide ions directly. In fact, the Arrhenius definitions of an acid and a base focus on hydrogen ions and hydroxide ions. Are there more fundamental definitions for acids and bases?

In 1923, the Danish scientist Johannes Brønsted and the English scientist Thomas Lowry independently proposed new definitions for acids and bases. Rather than considering both hydrogen and hydroxide ions, they focused on the hydrogen ion only. A Brønsted-Lowry acid is a compound that supplies a hydrogen ion in a reaction. A Brønsted-Lowry base, conversely, is a compound that accepts a hydrogen ion in a reaction. Thus, the Brønsted-Lowry definitions of an acid and a base focus on the movement of hydrogen ions in a reaction, rather than on the production of hydrogen ions and hydroxide ions in an aqueous solution.

Let us use the reaction of ammonia in water to demonstrate the Brønsted-Lowry definitions of an acid and a base. Ammonia and water molecules are reactants, while the ammonium ion and the hydroxide ion are products:

$$NH_3(aq) + H_2O(\ell) \rightarrow NH_4^+(aq) + OH^-(aq) \tag{10.2.1}$$

What has happened in this reaction is that the original water molecule has donated a hydrogen ion to the original ammonia molecule, which in turn has accepted the hydrogen ion. We can illustrate this as follows:



Because the water molecule donates a hydrogen ion to the ammonia, it is the Brønsted-Lowry acid, while the ammonia molecule—which accepts the hydrogen ion—is the Brønsted-Lowry base. Thus, ammonia acts as a base in both the Arrhenius sense and the Brønsted-Lowry sense.


Is an Arrhenius acid like hydrochloric acid still an acid in the Brønsted-Lowry sense? Yes, but it requires us to understand what really happens when HCl is dissolved in water. Recall that the hydrogen *atom* is a single proton surrounded by a single electron. To make the hydrogen *ion*, we remove the electron, leaving a bare proton. Do we *really* have bare protons floating around in aqueous solution? No, we do not. What really happens is that the H^+ ion attaches itself to H_2O to make H_3O^+, which is called the *hydronium ion*. For most purposes, H^+ and H_3O^+ represent the same species, but writing H_3O^+ instead of H^+ shows that we understand that there are no bare protons floating around in solution. Rather, these protons are actually attached to solvent molecules.

> A proton in aqueous solution may be surrounded by more than one water molecule, leading to formulas like $H_5O_2^+$ or $H_9O_4^+$ rather than H_3O^+. It is simpler, however, to use H_3O^+.

With this in mind, how do we define HCl as an acid in the Brønsted-Lowry sense? Consider what happens when HCl is dissolved in H_2O:

$$HCl(g) + H_2O(\ell) \rightarrow H_3O^+(aq) + Cl^-(aq) \tag{10.2.2}$$

We can depict this process using Lewis electron dot diagrams:



Now we see that a hydrogen ion is transferred from the HCl molecule to the H_2O molecule to make chloride ions and hydronium ions. As the hydrogen ion donor, **HCl acts as a Brønsted-Lowry acid**; as a hydrogen ion acceptor, **H_2O is a Brønsted-Lowry base**. So HCl is an acid not just in the Arrhenius sense but also in the Brønsted-Lowry sense. Moreover, by the Brønsted-Lowry definitions, H_2O is a base in the formation of aqueous HCl. So the Brønsted-Lowry definitions of an acid and a base classify the dissolving of HCl in water as a reaction between an acid and a base—although the Arrhenius definition would not have labeled H_2O a base in this circumstance.

All Arrhenius acids and bases are Brønsted-Lowry acids and bases as well. However, not all Brønsted-Lowry acids and bases are Arrhenius acids and bases.

Example 10.2.1

Aniline ($C_6H_5NH_2$) is slightly soluble in water. It has a nitrogen atom that can accept a hydrogen ion from a water molecule just like the nitrogen atom in ammonia does. Write the chemical equation for this reaction and identify the Brønsted-Lowry acid and base.

Solution

$C_6H_5NH_2$ and H_2O are the reactants. When $C_6H_5NH_2$ accepts a proton from H_2O, it gains an extra H and a positive charge and leaves an OH^- ion behind. The reaction is as follows:

$$C_6H_5NH_2(aq) + H_2O(\ell) \rightarrow C_6H_5NH_3^+(aq) + OH^-(aq)$$

Because $C_6H_5NH_2$ accepts a proton, it is the Brønsted-Lowry base. The H_2O molecule, because it donates a proton, is the Brønsted-Lowry acid.

Exercise 10.2.1

Caffeine ($C_8H_{10}N_4O_2$) is a stimulant found in coffees and teas. When dissolved in water, it can accept a proton from a water molecule. Write the chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer

$C_8H_{10}N_4O_2(aq) + H_2O(\ell) \rightarrow C_8H_{11}N_4O_2^+(aq) + OH^-(aq)$

B-L base B-L acid

The Brønsted-Lowry definitions of an acid and a base can be applied to chemical reactions that occur in solvents other than water. The following example illustrates.

Example 10.2.2

Sodium amide ($NaNH_2$) dissolves in methanol (CH_3OH) and separates into sodium ions and amide ions (NH_2^-). The amide ions react with methanol to make ammonia and the methoxide ion (CH_3O^-). Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Solution

The equation for the reaction is between NH_2^- and CH_3OH to make NH_3 and CH_3O^- is as follows:

$$NH_2^-(solv) + CH_3OH(\ell) \rightarrow NH_3(solv) + CH_3O^-(solv)$$

The label *(solv)* indicates that the species are dissolved in some solvent, in contrast to *(aq)*, which specifies an aqueous (H_2O) solution. In this reaction, we see that the NH_2^- ion accepts a proton from a CH_3OH molecule to make an NH_3 molecule. Thus, as the proton acceptor, NH_2^- is the Brønsted-Lowry base. As the proton donor, CH_3OH is the Brønsted-Lowry acid.

? Exercise 10.2.2

Pyridinium chloride (C_5H_5NHCl) dissolves in ethanol (C_2H_5OH) and separates into pyridinium ions ($C_5H_5NH^+$) and chloride ions. The pyridinium ion can transfer a hydrogen ion to a solvent molecule. Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer

$C_5H_5NH^+(solv) + C_2H_5OH(\ell) \rightarrow C_5H_5N(solv) + C_2H_5OH_2^+(solv)$

B-L acid B-L base

Application in Everyday Life

Many people like to put lemon juice or vinegar, both of which are acids, on cooked fish (Figure **10.2.1**). It turns out that fish have volatile amines (bases) in their systems, which are neutralized by the acids to yield involatile ammonium salts. This reduces the odor of the fish, and also adds a "sour" taste that we seem to enjoy.



$CH_3COOH + NH_2CH_2CH_2CH_2CH_2NH_2 \longrightarrow CH_3COO^- + NH_3^+CH_2CH_2CH_2CH_2NH_2$

Acetic acid + Putrescine → Acetate ion + Putrescinium ion

Figure **10.2.1**: A neutralization reaction takes place between citric acid or acetic acid (proton donors) in lemons or vinegar and putrescine (proton acceptor) in the flesh of fish. (CC BY 4.0; OpenStax)

Pickling is a method used to preserve vegetables using a naturally produced acidic environment. The vegetable, such as a cucumber, is placed in a sealed jar submerged in a brine solution. The brine solution favors the growth of beneficial bacteria and suppresses the growth of harmful bacteria. The beneficial bacteria feed on starches in the cucumber and produce lactic acid as a waste product in a process called fermentation. The lactic acid eventually increases the acidity of the brine to a level that kills any harmful bacteria, which require a basic environment. Without the harmful bacteria consuming the cucumbers they are able to last much longer than if they were unprotected. A byproduct of the pickling process changes the flavor of the vegetables with the acid making them taste sour.

To Your Health: Brønsted-Lowry Acid-Base Reactions in Pharmaceuticals

There are many interesting applications of Brønsted-Lowry acid-base reactions in the pharmaceutical industry. For example, drugs often need to be water soluble for maximum effectiveness. However, many complex organic compounds are not soluble or are only slightly soluble in water. Fortunately, those drugs that contain proton-accepting nitrogen atoms (and there are a lot of them) can be

reacted with dilute hydrochloric acid [HCl(aq)]. The nitrogen atoms—acting as Brønsted-Lowry bases—accept the hydrogen ions from the acid to make an ion, which is usually much more soluble in water. The modified drug molecules can then be isolated as chloride salts:

\[\mathrm{RN(sl\: aq) + H^+(aq) \rightarrow RNH^+(aq) \xrightarrow{Cl^-(aq)} RNHCl(s)} \label{Eq3} \]

where RN represents some organic compound containing nitrogen. The label *(sl aq)* means "slightly aqueous," indicating that the compound RN is only slightly soluble. Drugs that are modified in this way are called *hydrochloride salts*. Examples include the powerful painkiller codeine, which is commonly administered as codeine hydrochloride. Acids other than hydrochloric acid are also used. Hydrobromic acid, for example, gives *hydrobromide salts*. Dextromethorphan, an ingredient in many cough medicines, is dispensed as dextromethorphan hydrobromide. The accompanying figure shows another medication (lidocaine) as a hydrochloride salt.



Figure used with permission from Wikipedia.

Conjugate Acid-Base Pairs

According to the Bronsted-Lowry theory of acids and bases, an acid is a proton donor and a base is a proton acceptor. Once an acid has given up a proton, the part that remains is called the acid's **conjugate base**. This species is a base because it can accept a proton (to re-form the acid). The conjugate base of HF (first example below) is fluoride ion, F^-.

$$\text{Acid} = H^+ + \text{Conjugate base of Acid}^-$$

$$HF \rightleftharpoons H^+ + F^-$$

$$H_2O \rightleftharpoons H^+ + OH^-$$

$$NH_4^+ \rightleftharpoons H^+ + NH_3$$

Similarly, the part of the base that remains after a base accepts a proton is called the base's **conjugate acid**. This species is an acid because it can give up a proton (and thus re-form the base). The conjugate acid of fluoride ion, F^- (first example below) is HF.

$$H^+ + \text{Base} = \text{Conjugate acid of Base}^+$$

$$H^+ + F^- \rightleftharpoons HF$$

$$H^+ + OH^- \rightleftharpoons H_2O$$

$$H^+ + H_2O \rightleftharpoons H_3O^+$$

$$H^+ + NH_3 \rightleftharpoons NH_4^+$$

To summarize, the conjugate base of HF is fluoride ion, F^-, and the conjugate acid of fluoride ion, F^-, is HF. The HF/F- pair is referred to as a conjugate acid-base pair. The difference in the formulas of a conjugate acid-base pair (example: HF and F^-) is H^+. The table below lists conjugate acid-base pairs for your reference so that you can figure out the strategy of identifying them. For any given acid or base, you should be able to give its conjugate base or conjugate acid. The formula of an acid's conjugate base is generated by **removing a proton** (H^+) from the acid formula. The formula of the base's conjugate acid is formed by **adding a proton** (H^+) to the formula of the base.

Table **10.2.1**. Conjugate acid-base pairs.

Conjugate Acid	Conjugate Base
H_3O^+	H_2O
H_2O	OH^-
H_2SO_4	HSO_4^-
HSO_4^-	SO_4^{2-}
NH_4^+	NH_3
NH_3	NH_2^-
CH_3COOH	CH_3COO^-
$CH_3NH_3^+$	CH_3NH_2

Example 10.2.3 : Conjugate Pairs

Write the formula of the conjugate base of (a) HCl and (b) HCO_3^-.

Write the formula of the conjugate acid of (c) CH_3NH_2 and (d) OH^-.

Solution:

A conjugate base is formed by removing a proton (H^+). A conjugate acid is formed by accepting a proton (H^+).

a. After HCl donates a proton, a Cl^- ion is produced, and so Cl^- is the conjugate base.
b. After hydrogen carbonate ion, HCO_3^-, donates a proton, its conjugate base, CO_3^{2-} is produced.
c. After accepting a proton (H^+), CH_3NH_2 is converted to $CH_3NH_3^+$, its conjugate acid.
d. After accepting a proton (H^+), OH^- is converted to H_2O, its conjugate acid.

10.2.3Exercise : Conjugate Pairs

Write the formula of the conjugate base of (a) HNO_2 and (b) H_2CO_3.

Write the formula of the conjugate acid of (c) C_6H5NH_2 and (d) HCO_3^-.

Answer


a. NO_2^- is the conjugate base of HNO_2.

b. HCO_3^- is the conjugate base of H_2CO_3

c. $C_6H5NH_3^+$ is the conjugate acid of C_6H5NH_2.


d. H_2CO_3 is the conjugate acid of HCO_3^-

In the reaction of ammonia with water to give ammonium ions and hydroxide ions, ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The **conjugate acid–base pairs** for this reaction are $\boldsymbol{NH_4^+/NH_3}$ and $\boldsymbol{H_2O/OH^-}$. This means that the conjugate acid of the base NH_3 is NH_4^+ while the conjugate base of the acid NH_4^+ is NH_3. Similarly, the conjugate base of the acid H_2O is OH^-, and the conjugate acid of the base OH^- is H_2O.




This figure has three parts in two rows. In the first row, two diagrams of acid-base pairs are shown. On the left, a space filling model of H subscript 2 O is shown with a red O atom at the center and two smaller white H atoms attached in a bent shape. Above this model is the label “H subscript 2 O (acid)” in purple. An arrow points right, which is labeled “Remove H superscript plus.” To the right is another space filling model with a single red O atom to which a single smaller white H atom is attached. The label in purple above this model reads, “O H superscript negative (conjugate base).” Above both of these red and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” To the right is a space filling model with a central blue N atom to which three smaller white H atoms are attached in a triangular pyramid arrangement. A label in green above reads “N H subscript 3 (base).” An arrow labeled “Add H superscript plus” points right. To the right of the arrow is another space filling model with a blue central N atom and four smaller white H atoms in a tetrahedral arrangement. The green label above reads “N H subscript 3 superscript plus (conjugate acid).” Above both of these blue and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” The second row of the figure shows the chemical reaction, H subscript 2 O (l) is shown in purple, and is labeled below in purple as “acid,” plus N H subscript 3 (a q) in green, labeled below in green as “base,” followed by a double sided arrow arrow and O H superscript negative (a q) in purple, labeled in purple as “conjugate base,” plus N H subscript 4 superscript plus (a q)” in green, which is labeled in green as “conjugate acid.” The acid on the left side of the equation is connected to the conjugate base on the right with a purple line. Similarly, the base on the left is connected to the conjugate acid on the right side.

In the **forward** reaction, the parent acid is H_2O and and the parent base is NH_3 (shown in the illustration below). The acid H_2O loses a proton (H^+) to form its conjugate base OH^-. The base NH_3 gains a proton, to produce its conjugate acid NH_4^+. In the **reverse** reaction, the acid NH_4^+ loses a proton (H^+) to form its conjugate base NH_3. The base OH^- gains a proton, to produce its conjugate acid H_2O.



When hydrogen fluoride (HF) dissolves in water and ionizes, protons are transferred from hydrogen fluoride (parent acid) molecules to water (parent base) molecules, yielding hydronium ions (conjugate acid of water) and fluoride ions (conjugate base of HF):

$$HF + H_2O \rightleftharpoons H_3O^+ + F^-$$



https://chem.libretexts.org/@go/page/387578

Example 10.2.3

Identify the conjugate acid-base pairs in this equilibrium.


$$CH_3CO_2H + H_2O \rightleftharpoons H_3O^+ + CH_3CO_2^-$$

Solution

Similarly, in the reaction of acetic acid with water, acetic acid **donates** a proton to water, which acts as the base. In the reverse reaction, H_3O^+ is the acid that donates a proton to the acetate ion, which acts as the base.

Once again, we have two conjugate acid–base pairs:

- the parent acid and its conjugate base ($CH_3CO_2H/CH_3CO_2^-$) and
- the parent base and its conjugate acid (H_3O^+/H_2O).



Example 10.2.4

Identify the conjugate acid-base pairs in this equilibrium.

$$(CH_3)_3N + H_2O \rightleftharpoons (CH_3)_3NH^+ + OH^-$$

Solution

One pair is H_2O and OH^-, where H_2O has one more H^+ and is the conjugate acid, while OH^- has one less H^+ and is the conjugate base.

The other pair consists of $(CH_3)_3N$ and $(CH_3)_3NH^+$, where $(CH_3)_3NH^+$ is the conjugate acid (it has an additional proton) and $(CH_3)_3N$ is the conjugate base.

Exercise 10.2.4

Identify the conjugate acid-base pairs in this equilibrium.

$$NH_2^- + H_2O \rightleftharpoons NH_3 + OH^-$$

Answer:

H_2O (acid) and OH^- (base); NH_2^- (base) and NH_3 (acid)

The use of conjugate acid-base pairs allows us to make a very simple statement about relative strengths of acids and bases. *The stronger an acid, the weaker its conjugate base*, and, conversely, *the stronger a base, the weaker its conjugate acid.*

Key Takeaways

- A Brønsted-Lowry acid is a proton donor, and a Brønsted-Lowry base is a proton acceptor.
- Brønsted-Lowry acid-base reactions are essentially proton transfer reactions.

10.2: Brønsted-Lowry Definition of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.2: Brønsted-Lowry Definition of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.3: Water is both an Acid and a Base

Learning Objectives

- To write chemical equations for water acting as an acid and as a base.

Water (H_2O) is an interesting compound in many respects. Here, we will consider its ability to behave as an acid or a base.

In some circumstances, a water molecule will accept a proton and thus act as a **Brønsted-Lowry base**. We saw an example in the dissolving of HCl in H_2O:

$$\mathrm{HCl} + \mathrm{H_2O}(\ell) \rightarrow \mathrm{H_3O^+(aq)} + \mathrm{Cl^-(aq)} \tag{10.3.1}$$

In other circumstances, a water molecule can donate a proton and thus act as a **Brønsted-Lowry acid**. For example, in the presence of the amide ion (see Example 4 in Section 10.2), a water molecule donates a proton, making ammonia as a product:

$$H_2O(\ell) + NH_2^-(aq) \rightarrow OH^-(aq) + NH_3(aq) \tag{10.3.2}$$

In this case, NH_2^- is a Brønsted-Lowry base (the proton acceptor).

So, depending on the circumstances, H_2O can act as either a Brønsted-Lowry acid or a Brønsted-Lowry base. Water is not the only substance that can react as an acid in some cases or a base in others, but it is certainly the most common example—and the most important one. A substance that can either donate or accept a proton, depending on the circumstances, is called an **amphiprotic** compound.

A water molecule can act as an acid or a base even in a sample of pure water. About 6 in every 100 million (6 in 10^8) water molecules undergo the following reaction:

$$H_2O(\ell) + H_2O(\ell) \rightarrow H_3O^+(aq) + OH^-(aq) \tag{10.3.3}$$

This process is called the **autoionization of water** (Figure **10.3.1**) and occurs in every sample of water, whether it is pure or part of a solution. Autoionization occurs to some extent in any amphiprotic liquid. (For comparison, liquid ammonia undergoes autoionization as well, but only about 1 molecule in a million billion (1 in 10^{15}) reacts with another ammonia molecule.)

Note

It is rare to truly have pure water. Water exposed to air will usually be slightly acidic because dissolved carbon dioxide gas, or carbonic acid, decreases the pH slightly below 7. Alternatively, dissolved minerals, like calcium carbonate (limestone), can make water slightly basic.




Figure **10.3.1** Autoionization. A small fraction of water molecules—approximately 6 in 100 million—ionize spontaneously into hydronium ions and hydroxide ions. This picture necessarily overrepresents the amount of autoionization that really occurs in pure water.

Example 10.3.1

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $H_2O(\ell) + NO_2^-(aq) \rightarrow HNO_2(aq) + OH^-(aq)$
2. $HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)$

Solution

1. In this reaction, the water molecule donates a proton to the NO_2^- ion, making $OH^-(aq)$. As the proton donor, H_2O acts as a Brønsted-Lowry acid.
2. In this reaction, the water molecule accepts a proton from $HC_2H_3O_2$, becoming $H_3O^+(aq)$. As the proton acceptor, H_2O is a Brønsted-Lowry base.

Exercise 10.3.2

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $HCOOH(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + HCOO^-(aq)$
2. $H_2O(\ell) + PO_4^{3-}(aq) \rightarrow OH^-(aq) + HPO_4^{2-}(aq)$

Answer

1. H_2O acts as the proton acceptor (Brønsted-Lowry base)
2. H_2O acts as the proton donor (Brønsted-Lowry acid)

Key Takeaway

- Water molecules can act as both an acid and a base, depending on the conditions.

10.3: Water is both an Acid and a Base is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.3: Water - Both an Acid and a Base** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.4: The Strengths of Acids and Bases

Learning Objectives

- Describe the difference between strong and weak acids and bases.
- Describe how a chemical reaction reaches chemical equilibrium.
- Define the pH scale and use it to describe acids and bases.

Acids and bases do not all demonstrate the same degree of chemical activity in solution. Different acids and bases have different strengths.

Strong and Weak Acids

Let us consider the strengths of acids first. A small number of acids ionize completely in aqueous solution. For example, when HCl dissolves in water, every molecule of HCl separates into a hydronium ion and a chloride ion:

$$\mathrm{HCl(g) + H_2O(l) \xrightarrow{\sim 100\,\%} H_3O^+(aq) + Cl^-(aq)} \tag{10.4.1}$$

HCl(aq) is one example of a **strong acid**, which is a compound that is essentially **100% ionized** in aqueous solution. There are very few strong acids. The important ones are listed in Table **10.4.1**.

Table **10.4.1**: Strong Acids and Bases (All in Aqueous Solution)

Acids	Bases
HCl	LiOH
HBr	NaOH
HI	KOH
HNO_3	$Mg(OH)_2$
H_2SO_4	$Ca(OH)_2$
$HClO_4$	

By analogy, a strong base is a compound that is essentially 100% ionized in aqueous solution. As with acids, there are only a few strong bases, which are also listed in Table **10.4.1**.

If an acid is **not listed** in Table **10.4.1**, it is likely a **weak acid**, which is a compound that is not 100% ionized in aqueous solution. Similarly, a weak base is a compound that is not 100% ionized in aqueous solution. For example, acetic acid ($\mathrm{HC_2H_3O_2}$) is a weak acid. The ionization reaction for acetic acid is as follows:

$$\mathrm{HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)} \tag{10.4.2}$$

Depending on the concentration of $HC_2H_3O_2$, the ionization reaction may occur only for 1%–5% of the acetic acid molecules.

Looking Closer: Household Acids and Bases

Many household products are acids or bases. For example, the owner of a swimming pool may use muriatic acid to clean the pool. Muriatic acid is another name for hydrochloric acid [$\mathrm{HCl(aq)}$]. Vinegar has already been mentioned as a dilute solution of acetic acid [$\mathrm{HC_2H_3O_2(aq)}$]. In a medicine chest, one may find a bottle of vitamin C tablets; the chemical name of vitamin C is ascorbic acid ($\mathrm{HC_6H_7O_6}$).

One of the more familiar household bases is ammonia (NH_3), which is found in numerous cleaning products. As we mentioned previously, ammonia is a base because it increases the hydroxide ion concentration by reacting with water:

$$\mathrm{NH_3(aq) + H_2O(\ell) \rightarrow NH_4^+(aq) + OH^-(aq)} \tag{10.4.3}$$

Many soaps are also slightly basic because they contain compounds that act as Brønsted-Lowry bases, accepting protons from water and forming excess hydroxide ions. This is one reason that soap solutions are slippery.



Figure 10.4.1: (left) Bottles of alkaline drain cleaners containing sodium hydroxide can dissolve greases and hair. (right) Chemical burns caused by sodium hydroxide solution photographed 44 hours after exposure. Figures used with permission from Wikipedia.

Perhaps the most dangerous household chemical is the lye-based drain cleaner. Lye is a common name for sodium hydroxide, although it is also used as a synonym for potassium hydroxide. Lye is an extremely caustic chemical that can react with grease, hair, food particles, and other substances that may build up and form a clog in a pipe. Unfortunately, lye can also attack tissues and other substances in our bodies. Thus, when we use lye-based drain cleaners, we must be very careful not to touch any of the solid drain cleaner or spill the water it was poured into. Safer, nonlye drain cleaners use peroxide compounds to react on the materials in the clog and clear the drain.

Chemical Equilibrium in Weak Acids and Bases

The behavior of **weak acids and bases** illustrates a key concept in chemistry. Does the chemical reaction describing the ionization of a weak acid or base just stop when the acid or base is done ionizing? Actually, no. Rather, the reverse process—the reformation of the molecular form of the acid or base—occurs, ultimately at the same rate as the ionization process. For example, the ionization of the weak acid $\mathbf{HC_2H_3O_2\,(aq)}$ is as follows:

$$\mathbf{HC_2H_3O_2\,(aq) + H_2O(\ell) \;\rightarrow H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)} \tag{10.4.4}$$

The reverse process also begins to occur:

$$\mathbf{H_3O^+\,(aq) + C_2H_3O_2^-\,(aq) \;\rightarrow HC_2H_3O_2\,(aq) + H_2O(\ell)} \tag{10.4.5}$$

Eventually, there is a balance between the two opposing processes, and no additional change occurs. The chemical reaction is better represented at this point with a double arrow:

$$\mathbf{HC_2H_3O_2\,(aq) + H_2O(\ell) \rightleftharpoons H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)} \tag{10.4.6}$$

The $\rightleftharpoons$ implies that **both the forward and reverse reactions** are occurring, and their effects cancel each other out. A process at this point is considered to be at **chemical equilibrium** (or equilibrium). It is important to note that the processes do not stop. They balance out each other so that there is no further net change; that is, chemical equilibrium is a *dynamic equilibrium*.

Example 10.4.1: Partial Ionization

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HNO_2(aq)$
b. $C_5H_5N(aq)$

Solution

a. $HNO_2(aq) + H_2O(\ell) \leftrightarrows NO_2^-(aq) + H_3O^+(aq)$
b. $C_5H_5N(aq) + H_2O(\ell) \leftrightarrows C_5H_5NH^+(aq) + OH^-(aq)$

? Exercise 10.4.1

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HF_{(aq)}$
b. $AgOH_{(aq)}$
c. $CH_3NH_2(aq)$

Answer

a. $HF(aq) + H_2O(\ell) \leftrightarrows F^-(aq) + H_3O^+(aq)$

b. $AgOH(aq) \leftrightarrows Ag^+(aq) + OH^-(aq)$

c. $CH_3NH_2(aq) + H_2O(\ell) \leftrightarrows CH_3NH_3^+(aq) + OH^-(aq)$

Acid Ionization Constant, K_a

The ionization for a general weak acid, **HA**, can be written as follows:

$$\mathrm{HA}\,(aq) \rightleftharpoons \mathrm{H^+}\,(aq) + \mathrm{A^-}\,(aq)$$

Because the acid is weak, an equilibrium expression can be written. An **acid ionization constant** (K_a) is the equilibrium constant for the ionization of an acid.

$$K_a = \frac{[\mathrm{H^+}]\,[\mathrm{A^-}]}{[\mathrm{HA}]}$$

The acid ionization represents the fraction of the original acid that has been ionized in solution. Therefore, the numerical value of K_a is a reflection of the strength of the acid. Weak acids with relatively higher K_a values are stronger than acids with relatively lower K_a values. Because strong acids are essentially 100% ionized, the concentration of the acid in the denominator is nearly zero and the K_a value approaches infinity. For this reason, K_a values are generally reported for weak acids only.

The table below is a listing of acid ionization constants for several acids. Note that polyprotic acids have a distinct ionization constant for each ionization step, with each successive ionization constant being smaller than the previous one.

Table 10.4.2: Acid Ionization Constants at 25°C

Name of Acid	Ionization Equation	K_a
Sulfuric acid	$\mathrm{H_2SO_4 \rightleftharpoons H^+ + HSO_4^-}$ $\mathrm{HSO_4^- \rightleftharpoons H^+ + SO_4^{2-}}$	very large 1.3×10^{-2}
Hydrofluoric acid	$\mathrm{HF \rightleftharpoons H^+ + F^-}$	7.1×10^{-4}
Nitrous acid	$\mathrm{HNO_2 \rightleftharpoons H^+ + NO_2^-}$	4.5×10^{-4}
Benzoic acid	$\mathrm{C_6H_5COOH \rightleftharpoons H^+ + C_6H_5COO^-}$	6.5×10^{-5}
Acetic acid	$\mathrm{CH_3COOH \rightleftharpoons H^+ + CH_3COO^-}$	1.8×10^{-5}
Carbonic acid	$\mathrm{H_2CO_3 \rightleftharpoons H^+ + HCO_3^-}$ $\mathrm{HCO_3^- \rightleftharpoons H^+ + CO_3^{2-}}$	4.2×10^{-7} 4.8×10^{-11}

Hydrofluoric acid $HF_{(aq)}$ reacts directly with glass (very few chemicals react with glass). Hydrofluoric acid is used in glass etching.

Strong and Weak Bases and Base Ionization Constant, K_b

As with acids, bases can either be strong or weak, depending on their extent of ionization. A **strong base** is a base, which ionizes completely in an aqueous solution. The most common strong bases are soluble metal hydroxide compounds such as potassium

hydroxide. Some metal hydroxides are not as strong simply because they are not as soluble. Calcium hydroxide is only slightly soluble in water, but the portion that does dissolve also dissociates into ions.

A **weak base** is a base that ionizes only slightly in an aqueous solution. Recall that a base can be defined as a substance, which accepts a hydrogen ion from another substance. When a weak base such as ammonia is dissolved in water, it accepts an H^+ ion from water, forming the hydroxide ion and the conjugate acid of the base, the ammonium ion.

$$NH_3\,(aq) + H_2O\,(l) \rightleftharpoons NH_4^+\,(aq) + OH^-\,(aq)$$

The equilibrium greatly favors the reactants and the extent of ionization of the ammonia molecule is very small.

An equilibrium expression can be written for the reactions of weak bases with water. Because the concentration of water is extremely large and virtually constant, the water is not included in the expression. A **base ionization constant** (K_b) is the equilibrium constant for the ionization of a base. For ammonia the expression is:

$$K_b = \frac{[NH_4^+]\,[OH^-]}{[NH_3]}$$

The numerical value of K_b is a reflection of the strength of the base. Weak bases with relatively higher K_b values are stronger than bases with relatively lower K_b values. Table 10.4.3 is a listing of base ionization constants for several weak bases.

Table 10.4.3: Base Ionization Constants at 25°C

Name of Base	Ionization Equation	K_b
Methylamine	$CH_3NH_2 + H_2O \rightleftharpoons CH_3NH_3^+ + OH^-$	5.6×10^{-4}
Ammonia	$NH_3 + H_2O \rightleftharpoons NH_4^+ + OH^-$	1.8×10^{-5}
Pyridine	$C_5H_5N + H_2O \rightleftharpoons C_5H_5NH^+ + OH^-$	1.7×10^{-9}
Acetate ion	$CH_3COO^- + H_2O \rightleftharpoons CH_3COOH + OH^-$	5.6×10^{-10}
Fluoride ion	$F^- + H_2O \rightleftharpoons HF + OH^-$	1.4×10^{-11}
Urea	$H_2NCONH_2 + H_2O \rightleftharpoons H_2NCONH_3^+ + OH^-$	1.5×10^{-14}

The Ion-Product of Water

As we have already seen, H_2O can act as an acid or a base. Within any given sample of water, some H_2O molecules are acting as acids, and other H_2O molecules are acting as bases. The chemical equation is as follows:

$$\underbrace{H_2O}_{\text{acid}} + \underbrace{H_2O}_{\text{base}} \rightleftharpoons H_3O^+ + OH^- \tag{10.4.7}$$




Figure 10.4.2: Autoionization of water, giving hydroxide and hydronium ions.

Similar to a weak acid, the autoionization of water is an equilibrium process, and is more properly written as follows:

$$H_2O(\ell) + H_2O(\ell) \rightleftharpoons H_3O^+\,(aq) + OH^-\,(aq) \tag{10.4.8}$$

We often use the simplified form of the reaction:

$$H_2O\,(l) \rightleftharpoons H^+\,(aq) + OH^-\,(aq)$$

The equilibrium constant for the autoionization of water is referred to as the ion-product for water and is given the symbol Kw.

$$K_w = [H^+][OH^-]$$

The **ion-product of water (*Kw*)** is the mathematical product of the concentration of hydrogen ions and hydroxide ions. Note that H_2O is not included in the ion-product expression because it is a pure liquid. The value of Kw is very small, in accordance with a

reaction that favors the reactants. At 25°C, the experimentally determined value of K_w in pure water is 1.0×10^{-14}.

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14}$$

In a sample of pure water, the concentrations of hydrogen and hydroxide ions are equal to one another. Pure water or any other aqueous solution in which this ratio holds is said to be **neutral.** To find the molarity of each ion, the square root of Kw is taken.

$$[H^+] = [OH^-] = 1.0\times10^{-7}$$

The product of these two concentrations is 1.0×10^{-14}

$$[H^+]\times[OH^-] = (1.0 \times 10^{-7})(1.0 \times 10^{-7}) = 1.0 \times 10^{-14}$$

- For acids, the concentration of H^+ or $[H^+]$) is greater than 1.0×10^{-7} M
- For bases, the concentration of OH^- or $[OH^-]$ is greater than 1.0×10^{-7} M.

Aqueous HCl is an example of acidic solution. Hydrogen chloride (HCl) ionizes to produce H^+ and Cl^- ions upon dissolving in water. This **increases the concentration of H^+ ions** in the solution. According to **Le Chatelier's principle**, the equilibrium represented by

$$H_2O\,(l) \rightleftharpoons H^+\,(aq) + OH^-\,(aq)$$

$$HCl\,(g) \longrightarrow H^+\,(aq) + Cl^-\,(aq)$$

is **forced to the left**, towards the reactant. As a result, the **concentration of the hydroxide ion decreases**.

Now, consider KOH (aq), a **basic solution**. Solid potassium hydroxide (KOH) dissociates in water to yield potassium ions and hydroxide ions.

$$KOH(s) \rightarrow K^+(aq) + OH^-(aq)$$

The **increase in concentration of the OH^- ions** will cause a decrease in the concentration of the H^+ ions.

No matter whether the aqueous solution is an acid, a base, or neutral:and the ion-product of $[H^+][OH^-]$ remains constant.

- For acidic solutions, $[H^+]$) is greater than $[OH^-]$.
- For basic solutions, [OH−] is greater than $[H^+]$.
- For neutral solutions, $[H_3O^+] = [OH-] = 1.0\times10^{-7}$M

This means that if you know $[H^+]$ for a solution, you can calculate what $[OH^-]$) has to be for the product to equal 1.0×10^{-14}, or if you know $[OH^-]$), you can calculate $[H^+]$. This also implies that as one concentration goes up, the other must go down to compensate so that their product always equals the value of K_w.

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \tag{10.4.9}$$

Example 10.4.2

Hydrochloric acid (HCl) is a strong acid, meaning it is 100% ionized in solution. What is the $[H^+]$ and the $[OH^-]$ in a solution of 2.0×10^{-3} M HCl?

Solution

Step 1: List the known values and plan the problem.

Known

- $[HCl] = 2.0\times10^{-3}$ M
- $Kw = 1.0\times10^{-14}$

Unknown

- $[H^+]$=?M
- $[OH^-]$=?M

Because HCl is 100% ionized, the concentration of H+ ions in solution will be equal to the original concentration of HCl. Each HCl molecule that was originally present ionizes into one H+ ion and one Cl− ion. The concentration of OH− can then be determined from the [H+] and Kw.

Step 2: Solve.

$$[H^+]=2.0\times10^{-3}\,M$$

$$Kw = [H^+][OH^-] = 1.0\times10^{-14}$$

$$[OH^-] = Kw/[H^+] = 1.0\times10^{-14}/2.0\times10^{-3} = 5.0\times10^{-12}\,M$$

Step 3: Think about your result.

The $[H^+]$ is much higher than the $[OH^-]$ because the solution is acidic. As with other equilibrium constants, the unit for Kw is customarily omitted.

10.4.2 Exercise

Sodium hydroxide (NaOH) is a strong base. What is the $[H^+]$ and the $[OH^-]$ in a 0.001 M NaOH solution at 25 °C?

Answer

$[OH^-]$ = 0.001M or 1 x 10^{-3}M; $[H^+]=1\times10^{-11}$M.

The pH Scale

One qualitative measure of the strength of an acid or a base solution is the pH scale, which is based on the concentration of the hydronium (or hydrogen) ion in aqueous solution.

$$pH = -\log[H^+]$$

or

$$pH = -\log[H_3O^+]$$

Figure **10.4.3** illustrates this relationship, along with some examples of various solutions. Because hydrogen ion concentrations are generally less than one (for example $1.3 \times 10^{-3}\ M$), the log of the number will be a negative number. To make pH even easier to work with, pH is defined as the ***negative* log of** $[H_3O^+]$, which will give a positive value for pH.




Figure **10.4.3**: The relationship between $[H^+]$ and pH *values for several common materials.*

A neutral (neither acidic nor basic) solution has a pH of 7. A pH below 7 means that a solution is acidic, with lower values of pH corresponding to increasingly acidic solutions. A pH greater than 7 indicates a basic solution, with higher values of pH corresponding to increasingly basic solutions. Thus, given the pH of several solutions, you can state which ones are acidic, which ones are basic, and which are more acidic or basic than others. These are summarized in Table \(\PageIndex{4}.

Table **10.4.4**: Acidic, Basic and Neutral pH Values

Acidic, Basic and Neutral pH Values

Classification	Relative Ion Concentrations	pH at 25 °C
acidic	$[H^+] > [OH^-]$	pH < 7
neutral	$[H^+] = [OH^-]$	pH = 7
basic	$[H^+] < [OH^-]$	pH > 7

Example 10.4.3

Find the pH, given the $[H^+]$ of the following:

a. 1×10^{-3} M
b. 2.5×10^{-11} M
c. 4.7×10^{-9} M

Solution

pH = - log $[H_3O^+]$

Substitute the known quantity into the equation and solve. Use a scientific calculator for b and c.

a. pH = - log $[\underline{1} \times 10^{-3}]$ = 3.$\underline{0}$ (1 decimal place since 1 has 1 significant figure)
b. pH = - log $[\underline{2.5} \times 10^{-11}]$ = 10.$\underline{60}$ (2 decimal places since 2.5 has 2 significant figures)
c. pH = - log $[\underline{4.7} \times 10^{-9}]$ = 8.$\underline{33}$ (2 decimal places since 4.7 has 2 significant figures)

Note on significant figures:

Because the number(s) before the decimal point in the pH value relate to the power on 10, the number of digits *after* the decimal point (underlined) is what determines the number of significant figures in the final answer.

10.4.3 Exercise

Find the pH, given $[H^+]$ of the following:

a. 5.8×10^{-4} M
b. 1.0×10^{-7} M

Answer

a. 3.24

b. 7.00

Table **10.4.5** lists the pH of several common solutions. The most acidic among the listed solutions is battery acid with the lowest pH value (0.3). The most basic is 1M NaOH solution with the highest pH value of 14.0. Notice that some biological fluids (stomach acid and urine) are nowhere near neutral. You may also notice that many food products are slightly acidic. They are acidic because they contain solutions of weak acids. If the acid components of these foods were strong acids, the food would likely be inedible.

Table **10.4.5**: The pH Values of Some Common Solutions

Solution	pH
battery acid	0.3
stomach acid	1–2
lemon or lime juice	2.1
vinegar	2.8–3.0
Coca-Cola	3
wine	2.8–3.8
beer	4–5
coffee	5
milk	6
urine	6
pure H_2O	7
(human) blood	7.3–7.5
sea water	8
antacid (milk of magnesia)	10.5
NH_3 (1 M)	11.6
bleach	12.6
NaOH (1 M)	14.0

Example 10.4.4

Label each solution as acidic, basic, or neutral based only on the stated ***pH***.

a. milk of magnesia, pH = 10.5
b. pure water, pH = 7
c. wine, pH = 3.0

Solution

a. With a pH greater than 7, milk of magnesia is basic. (Milk of magnesia is largely $Mg(OH)_2$.)
b. Pure water, with a pH of 7, is neutral.
c. With a pH of less than 7, wine is acidic.

10.4.4 Exercise

Identify each substance as acidic, basic, or neutral based only on the stated ***pH***.

a. human blood with ***pH*** = 7.4
b. household ammonia with ***pH*** = 11.0
c. cherries with ***pH*** = 3.6

Answer

a. slightly basic

b. basic

c. acidic

Acid Rain

Normal rainwater has a pH between 5 and 6 due to the presence of dissolved CO_2 which forms carbonic acid:

$$\mathrm{H_2O\,(l) + CO_2\,(g) \longrightarrow H_2CO_3\,(aq)} \tag{10.4.10}$$

$$\mathrm{H_2CO_3\,(aq) \rightleftharpoons H^+\,(aq) + HCO_3^-\,(aq)} \tag{10.4.11}$$

Acid rain is rainwater that has a pH of less than 5, due to a variety of nonmetal oxides, including CO_2, SO_2, SO_3, NO, and NO_2 being dissolved in the water and reacting with it to form not only carbonic acid, but sulfuric acid and nitric acid. The formation and subsequent ionization of sulfuric acid are shown here:

$$\mathrm{H_2O\,(l) + SO_3\,(g) \longrightarrow H_2SO_4\,(aq)} \tag{10.4.12}$$

$$\mathrm{H_2SO_4\,(aq) \longrightarrow H^+\,(aq) + HSO_4^-\,(aq)} \tag{10.4.13}$$

Carbon dioxide is naturally present in the atmosphere because we and most other organisms produce it as a waste product of metabolism. Carbon dioxide is also formed when fires release carbon stored in vegetation or when we burn wood or fossil fuels. Sulfur trioxide in the atmosphere is naturally produced by volcanic activity, but it also stems from burning fossil fuels, which have traces of sulfur, and from the process of "roasting" ores of metal sulfides in metal-refining processes. Oxides of nitrogen are formed in internal combustion engines where the high temperatures make it possible for the nitrogen and oxygen in air to chemically combine.

Acid rain is a particular problem in industrial areas where the products of combustion and smelting are released into the air without being stripped of sulfur and nitrogen oxides. In North America and Europe until the 1980s, it was responsible for the destruction of forests and freshwater lakes, when the acidity of the rain actually killed trees, damaged soil, and made lakes uninhabitable for all but the most acid-tolerant species. Acid rain also corrodes statuary and building facades that are made of marble and limestone (Figure **10.4.4**). Regulations limiting the amount of sulfur and nitrogen oxides that can be released into the atmosphere by industry and automobiles have reduced the severity of acid damage to both natural and manmade environments in North America and Europe. It is now a growing problem in industrial areas of China and India.



(a) (b)

Figure **10.4.4**: (a) Acid rain makes trees more susceptible to drought and insect infestation, and depletes nutrients in the soil. (b) It also is corrodes statues that are carved from marble or limestone. (credit a: modification of work by Chris M Morris; credit b: modification of work by "Eden, Janine and Jim"/Flickr)

Two photos are shown. Photograph a on the left shows the upper portion of trees against a bright blue sky. The tops of several trees at the center of the photograph have bare branches and appear to be dead. Image b shows a statue of a man that appears to from the revolutionary war era in either marble or limestone.

Key Takeaways

- Acids and bases can be strong or weak depending on the extent of ionization in solution.
- Most chemical reactions reach equilibrium at which point there is no net change.
- The ion-product of $[H^+][OH^-]$ in an aqueous solution remains constant.
- A ***p*H** value is simply the *negative* of the logarithm of the H^+ ion concentration ($-\log[H^+]$).
- The pH scale is used to succinctly communicate the acidity or basicity of a solution.
- A solution is **acidic** if $p\mathrm{H} < 7$.
- A solution is **basic** if $p\mathrm{H} > 7$.
- A solution is **neutral** if $p\mathrm{H} = 7$.

10.4: The Strengths of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.4: The Strengths of Acids and Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.5: pH and Kw

Skills to Develop

- To define the pH scale as a measure of acidity of a solution

Because of its *amphoteric* nature (i.e., acts as both an acid or a base), water does not always remain as H_2O molecules. In fact, two water molecules react to form hydronium and hydroxide ions:

$$2\,H_2O\,(l) \rightleftharpoons H_3O^+\,(aq) + OH^-\,(aq) \tag{10.5.1}$$

This is also called the self-ionization of water. The concentration of H_3O^+ and OH^- are equal in pure water because of the 1:1 stoichiometric ratio of Equation 10.5.1. The molarity of H_3O^+ and OH^- in water are also both $1.0 \times 10^{-7}\ M$ at 25° C. Therefore, a constant of water (K_w) is created to show the equilibrium condition for the self-ionization of water. The product of the molarity of hydronium and hydroxide ion is always 1.0×10^{-14}.

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \tag{10.5.2}$$

This equations also applies to all aqueous solutions. However, K_w does change at different temperatures, which affects the pH range discussed below.

Note

H^+ and H_3O^+ is often used interchangeably to represent the hydrated proton, commonly call the hydronium ion.

pH

Because K_w is constant (1.0×10^{-14} at 25 °C, the pK_w is 14, the constant of water determines the range of the pH scale. To understand what the pK_w is, it is important to understand first what the "p" means in pOH, and pH. The Danish biochemist Søren Sørenson proposed the term pH to refer to the "potential of hydrogen ion." He defined the "p" as the negative of the logarithm, -log, of $[H^+]$. Therefore the pH is the negative logarithm of the molarity of H. The pOH is the negative logarithm of the molarity of OH^- and the pK_w is the negative logarithm of the constant of water. These definitions give the following equations:

$$pH = -\log[H^+] \tag{10.5.3}$$

$$pOH = -\log[OH^-] \tag{10.5.4}$$

$$pK_w = -\log[K_w] \tag{10.5.5}$$

A logarithm, used in the above equations, of a number is how much a power is raised to a particular base in order to produce that number. To simplify this, look at the equation: $\log_b a=x$. This correlates to $b^x=a$. A simple example of this would be $\log_{10} 100 = 2$, or $10^2 = 100$. It is assumed that the base of Logarithms is ten if it is not stated. So for the sake of pH and pOH problems it will always be ten. When x is a negative number that means you are dividing it by the power. So, if $log_{10}0.01 = -2$ which can be written $10^{-2} = 0.01$, then 10^{-2} also means $1/10^2$. The log function can be found on your scientific calculator. Now if we apply this to pH and pOH we can better understand how we calculate the values.

At room temperature,

$$K_w = 1.0 \times 10^{-14} \tag{10.5.6}$$

So

$$pK_w = -\log[1.0 \times 10^{-14}] \tag{10.5.7}$$

Using the properties of logarithms, Equation 10.5.7 can be rewritten as

$$10^{-pK_w} = 10^{-14}. \tag{10.5.8}$$

By substituting, we see that pK_w is 14. The equation also shows that each increasing unit on the scale decreases by the factor of ten on the concentration of H^+. For example, a pH of 1 has a molarity ten times more concentrated than a solution of pH 2.

Since

$$pK_w = 14 \tag{10.5.9}$$

$$pK_w = pH + pOH = 14 \tag{10.5.10}$$

Note

The pH scale is logarithmic, meaning that an increase or decrease of an integer value changes the concentration by a tenfold. For example, a pH of 3 is ten times more acidic than a pH of 4. Likewise, a pH of 3 is one hundred times more acidic than a pH of 5. Similarly a pH of 11 is ten times more basic than a pH of 10.

The Effective Range of the pH Scale

It is common that the pH scale is argued to range from 0-14 or perhaps 1-14 but neither is correct. The pH range does not have an upper nor lower bound, since as defined above, the pH is an indication of concentration of H^+. For example, at a pH of zero the hydronium ion concentration is one molar, while at pH 14 the hydroxide ion concentration is one molar. Typically the concentrations of H^+ in water in most solutions fall between a range of 1 M (pH=0) and 10^{-14} M (pH=14). Hence a range of 0 to 14 provides sensible (**but not absolute**) "bookends" for the scale (Figure **10.5.1**). One can go somewhat below zero and somewhat above 14 in water, because the concentrations of hydronium ions or hydroxide ions can exceed one molar. Figure 1 depicts the pH scale with common solutions and where they are on the scale.





Figure* 10.5.1: Solutions and the placement of them on pH scale*

- From the range 7-14, a solution is basic. The pOH should be looked in the perspective of OH^- instead lf H^+. Whenever the value of pOH is greater than 7, then it is considered basic. And therefore there are more OH^- than H^+ in the solution
- At pH 7, the substance or solution is at neutral and means that the concentration of H^+ and OH^- ion is the same.
- From the range 1-7, a solution is acidic. So, whenever the value of a pH is less than 7, it is considered acidic. There are more H^+ than OH^- in an acidic solution.

Note

The pH scale does **not** have an upper nor lower bound. Negative pH values are possible

The following PhET simulation helps to build an intuition of the pH scale.



Example 10.5.1

If the concentration of $NaOH$ in a solution is $2.5 \times 10^{-4}\ M$, what is the concentration of H_3O^+?

SOLUTION

Because

$$1.0 \times 10^{-14} = [H_3O^+][OH^-]$$

to find the concentration of H_3O^+, solve for the $[H_3O^+]$.

$$\frac{1.0 \times 10^{-14}}{[OH^-]} = [H_3O^+]$$

$$\frac{1.0 \times 10^{-14}}{2.5 \times 10^{-4}} = [H_3O^+] = 4.0 \times 10^{-11}\ M$$

Example 10.5.2

a. Find the pH of a solution of 0.002 M of HCl.
b. Find the pH of a solution of 0.00005 M NaOH.

SOLUTION

a. The equation for pH is -log [H+]

$$[H^+] = 2.0 \times 10^{-3}\ M$$

$$pH = -\log[2.0 \times 10^{-3}] = 2.70$$

b. The equation for pOH is -log [OH^-]

$$[OH^-] = 5.0 \times 10^{-5}\ M$$

$$pOH = -\log[5.0 \times 10^{-5}] = 4.30$$

$$pK_w = pH + pOH$$

and

$$pH = pK_w - pOH$$

then

$$pH = 14 - 4.30 = 9.70$$

Living Systems

Molecules that make up or are produced by living organisms usually function within a narrow pH range (near neutral) and a narrow temperature range (body temperature). Many biological solutions, such as blood, have a pH near neutral. pH influences the structure and the function of many enzymes (protein catalysts) in living systems. Many of these enzymes have narrow ranges of pH activity (Table **10.5.1**).

Table 10.5.1: pH in Living Systems

Compartment	pH
Gastric Acid	1
Lysosomes	4.5
Granules of Chromaffin Cells	5.5
Human Skin	5.5
Urine	6
Neutral H_2O at 37 °C	6.81
Cytosol	7.2
Cerebrospinal Fluid	7.3
Blood	7.43-7.45
Mitochondrial Matrix	7.5
Pancreas Secretions	8.1

Cellular pH is so important that death may occur within hours if a person becomes acidotic (having increased acidity in the blood). As one can see pH is critical to life, biochemistry, and important chemical reactions. Common examples of how pH plays a very important role in our daily lives are given below:

- Water in swimming pool is maintained by checking its pH. Acidic or basic chemicals can be added if the water becomes too acidic or too basic.
- Whenever we get a heartburn, more acid build up in the stomach and causes pain. We needs to take antacid tablets (a base) to neutralize excess acid in the stomach.
- The pH of blood is slightly basic. A fluctuation in the pH of the blood can cause in serious harm to vital organs in the body.
- Certain diseases are diagnosed only by checking the pH of blood and urine.
- Certain crops thrive better at certain pH range.
- Enzymes activate at a certain pH in our body.

Contributors

- Emmellin Tung (UCD), Sharon Tsao (UCD), Divya Singh (UCD), Patrick Gormley (Lapeer Community School District)

10.5: pH and Kw is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

10.6: Buffers

Learning Objectives

- To define *buffer* and describe how it reacts with an acid or a base.

Weak acids are relatively common, even in the foods we eat. But we occasionally come across a strong acid or base, such as stomach acid, that has a strongly acidic pH of 1–2. By definition, strong acids and bases can produce a relatively large amount of hydrogen or hydroxide ions and, as a consequence, have a marked chemical activity. In addition, very small amounts of strong acids and bases can change the pH of a solution very quickly. If 1 mL of stomach acid [which we will approximate as 0.05 M HCl(aq)] is added to the bloodstream, and if no correcting mechanism is present, the pH of the blood would go from about 7.4 to about 4.9—a pH that is not conducive to continued living. Fortunately, the body has a mechanism for minimizing such dramatic pH changes.

The mechanism involves a **buffer**, a solution that resists dramatic changes in pH. A *buffer* (or *buffered*) solution is one that resists a change in its pH when H^+ or OH^- ions are added or removed owing to some other reaction taking place in the same solution. Buffers do so by being composed of certain pairs of solutes: either a **weak acid plus its conjugate base** or a **weak base plus its conjugate acid**.

For example, a buffer can be composed of dissolved **acetic acid ($HC_2H_3O_2$, a weak acid) and sodium acetate** ($NaC_2H_3O_2$). Sodium acetate is a salt that dissociates into sodium ions and acetate ions in solution. For as long as acetic acid and acetate ions are present in significant amounts a solution, this can resist dramatic pH changes. Another example of a buffer is a solution containing **ammonia (NH_3, a weak base) and ammonium chloride** (NH_4Cl). Ammonium acetate is also a salt that dissociates into ammonium ions and chloride ions in solution. The presence of ammonium ions with ammonia molecules satisfies the requisite condition for a buffer solution.

How Buffers Work

The essential component of a buffer system is a **conjugate acid-base pair** whose concentration is fairly high in relation to the concentrations of added H^+ or OH^- it is expected to buffer against. Let us use an acetic acid–sodium acetate buffer to demonstrate how buffers work. If a strong base—a source of OH^-(aq) ions—is added to the buffer solution, those hydroxide ions will react with the acetic acid in an acid-base reaction:

$$HC_2H_3O_2\,(aq) + OH^-\,(aq) \longrightarrow H_2O(\ell) + C_2H_3O_2^-\,(aq) \tag{10.6.1}$$

Rather than changing the pH dramatically by making the solution basic, the added hydroxide ions react to make water, and the pH does not change much.

> Many people are aware of the concept of buffers from *buffered aspirin*, which is aspirin that also has magnesium carbonate, calcium carbonate, magnesium oxide, or some other salt. The salt acts like a base, while aspirin is itself a weak acid.

If a strong acid—a source of H^+ ions—is added to the buffer solution, the H^+ ions will react with the anion from the salt. Because $HC_2H_3O_2$ is a weak acid, it is not ionized much. This means that if lots of hydrogen ions and acetate ions (from sodium acetate) are present in the same solution, they will come together to make acetic acid:

$$H^+\,(aq) + C_2H_3O_2^-\,(aq) \longrightarrow HC_2H_3O_2\,(aq) \tag{10.6.2}$$

Rather than changing the pH dramatically and making the solution acidic, the added hydrogen ions react to make molecules of a weak acid. Figure 10.6.1 illustrates both actions of a buffer.




Figure **10.6.1**: The Action of Buffers. Buffers can react with both strong acids (top) and strong bases (bottom) to minimize large changes in pH.

A simple buffer system might be a 0.2 *M* solution of sodium acetate; the conjugate pair here is acetic acid HAc and its conjugate base, the acetate ion Ac^-. The idea is that this conjugate pair "pool" will be available to gobble up any small ($\leq 10^{-3}$ *M*) addition of H+ or OH^- that may result from other processes going on in the solution.




Figure **10.6.2**: How HAc/Ac buffer works

Buffers work well only for limited amounts of added strong acid or base. Once either solute is all reacted, the solution is no longer a buffer, and rapid changes in pH may occur. We say that a buffer has a certain capacity. Buffers that have more solute dissolved in them to start with have larger capacities, as might be expected.

Buffers made from weak bases and salts of weak bases act similarly. For example, in a buffer containing NH_3 and NH_4Cl, ammonia molecules can react with any excess hydrogen ions introduced by strong acids:

$$NH_{3(aq)} + H^+_{(aq)} \rightarrow NH^+_{4(aq)} \tag{10.6.3}$$

while the ammonium ion [NH_4^+(aq)] can react with any hydroxide ions introduced by strong bases:

$$NH^+_{4(aq)} + OH^-_{(aq)} \rightarrow NH_{3(aq)} + H_2O_{(\ell)} \tag{10.6.4}$$




Figure **10.6.3**: How NH_4^+/NH_3 buffer works

Example 10.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $HCHO_2$ and $NaCHO_2$
b. HCl and NaCl
c. CH_3NH_2 and CH_3NH_3Cl

d. NH_3 and NaOH

Solution

a. Formic acid ($HCHO_2$) is a weak acid, while $NaCHO_2$ is a salt supplying —formate ion (CHO_2^-), the conjugate base of $HCHO_2$. The combination of these two solutes would make a buffer solution.
b. Hydrochloric acid (HCl) is a strong acid, not a weak acid, so the combination of these two solutes would not make a buffer solution.
c. Methylamine (CH_3NH_2) is like ammonia, a weak base. The compound CH_3NH_3Cl is a salt supplying $CH_3NH_3^+$, the conjugate acid of CH_3NH_2. The combination of these two solutes would make a buffer solution.
d. Ammonia (NH_3) is a weak base, but NaOH is a strong base. The combination of these two solutes would not make a buffer solution.

? Exercise 10.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $NaHCO_3$ and NaCl
b. H_3PO_4 and NaH_2PO_4
c. NH_3 and $(NH_4)_3PO_4$
d. NaOH and NaCl

Answer

b. H_3PO_4 (weak acid) and $H_2PO_4^-$ (conjugate base of H_3PO_4) make a buffer.

c. NH_3 (weak base) and NH_4^+ (conjugate acid of NH_3) make a buffer

✓ Food and Drink App: The Acid That Eases Pain

Although medicines are not exactly "food and drink," we do ingest them, so let's take a look at an acid that is probably the most common medicine: acetylsalicylic acid, also known as aspirin. Aspirin is well known as a pain reliever and antipyretic (fever reducer).

The structure of aspirin is shown in the accompanying figure. The acid part is circled; it is the H atom in that part that can be donated as aspirin acts as a Brønsted-Lowry acid. Because it is not given in Table 10.5.1, acetylsalicylic acid is a weak acid. However, it is still an acid, and given that some people consume relatively large amounts of aspirin daily, its acidic nature can cause problems in the stomach lining, despite the stomach's defenses against its own stomach acid.




Figure 10.6.4 The Molecular Structure of Aspirin. The circled atoms are the acid part of the molecule.

Because the acid properties of aspirin may be problematic, many aspirin brands offer a "buffered aspirin" form of the medicine. In these cases, the aspirin also contains a buffering agent-usually MgO-that regulates the acidity of the aspirin to minimize its acidic side effects.

As useful and common as aspirin is, it was formally marketed as a drug starting in 1899. The US Food and Drug Administration (FDA), the governmental agency charged with overseeing and approving drugs in the United States, wasn't formed until 1906. Some have argued that if the FDA had been formed before aspirin was introduced, aspirin may never have gotten approval due to its potential for side effects-gastrointestinal bleeding, ringing in the ears, Reye's syndrome (a liver

problem), and some allergic reactions. However, recently aspirin has been touted for its effects in lessening heart attacks and strokes, so it is likely that aspirin is here to stay.

Buffer solutions are essential components of all living organisms.

- Our blood is buffered to maintain a pH of 7.4 that must remain unchanged as metabolically-generated CO_2 (carbonic acid) is added and then removed by our lungs.
- Buffers in the oceans, in natural waters such as lakes and streams, and within soils help maintain their environmental stability against acid rain and increases in atmospheric CO_2.
- Many industrial processes, such as brewing, require buffer control, as do research studies in biochemistry and physiology that involve enzymes, are active only within certain pH ranges.

The pH in living systems (Figure \(\PageIndex{1}\)) is maintained by buffer systems.

Table 7.3.2: pH in Living Systems

Compartment	pH
Gastric Acid	1
Lysosomes	4.5
Granules of Chromaffin Cells	5.5
Human Skin	5.5
Urine	6
Neutral H_2O at 37 °C	6.81
Cytosol	7.2
Cerebrospinal Fluid	7.3
Blood	7.43-7.45
Mitochondrial Matrix	7.5
Pancreas Secretions	8.1

Human blood has a buffering system to minimize extreme changes in pH. One buffer in blood is based on the presence of HCO_3^- and H_2CO_3 [H_2CO_3 is another way to write CO_2(aq)]. With this buffer present, even if some stomach acid were to find its way directly into the bloodstream, the change in the pH of blood would be minimal. Inside many of the body's cells, there is a buffering system based on phosphate ions.

Medicine: The Buffer System in Blood

The normal pH of human blood is about 7.4. The carbonate buffer system in the blood uses the following equilibrium reaction:

$$\mathbf{CO_2}(g) + \mathbf{2\,H_2O}(l) \rightleftharpoons \mathbf{H_2CO_3}(aq) \rightleftharpoons \mathbf{HCO_3^-}(aq) + \mathbf{H_3O^+}(aq)$$

The concentration of carbonic acid, H_2CO_3 is approximately 0.0012 *M*, and the concentration of the hydrogen carbonate ion, $\mathbf{HCO_3^-}$, is around 0.024 *M*. Using the Henderson-Hasselbalch equation and the pK_a of carbonic acid at body temperature, we can calculate the pH of blood:

$$\mathbf{pH} = \mathbf{p}K_\mathbf{a} + \log\frac{[\mathbf{base}]}{[\mathbf{acid}]} = 6.1 + \log\frac{0.024}{0.0012} = 7.4$$

The fact that the H_2CO_3 concentration is significantly lower than that of the $\mathbf{HCO_3^-}$ ion may seem unusual, but this imbalance is due to the fact that most of the by-products of our metabolism that enter our bloodstream are acidic. Therefore, there must be a larger proportion of base than acid, so that the capacity of the buffer will not be exceeded.

Lactic acid is produced in our muscles when we exercise. As the lactic acid enters the bloodstream, it is neutralized by the HCO_3^- ion, producing H_2CO_3. An enzyme then accelerates the breakdown of the excess carbonic acid to carbon dioxide and water, which can be eliminated by breathing. In fact, in addition to the regulating effects of the carbonate buffering system on the pH of blood, the body uses breathing to regulate blood pH. If the pH of the blood decreases too far, an increase in breathing removes CO_2 from the blood through the lungs driving the equilibrium reaction such that $[H_3O^+]$ is lowered. If the blood is too alkaline, a lower breath rate increases CO_2 concentration in the blood, driving the equilibrium reaction the other way, increasing $[H^+]$ and restoring an appropriate pH.

Career Focus: Blood Bank Technology Specialist

At this point in this text, you should have the idea that the chemistry of blood is fairly complex. Because of this, people who work with blood must be specially trained to work with it properly.

A blood bank technology specialist is trained to perform routine and special tests on blood samples from blood banks or transfusion centers. This specialist measures the pH of blood, types it (according to the blood's ABO+/− type, Rh factors, and other typing schemes), tests it for the presence or absence of various diseases, and uses the blood to determine if a patient has any of several medical problems, such as anemia. A blood bank technology specialist may also interview and prepare donors to give blood and may actually collect the blood donation.

Blood bank technology specialists are well trained. Typically, they require a college degree with at least a year of special training in blood biology and chemistry. In the United States, training must conform to standards established by the American Association of Blood Banks.

Key Takeaway

- A buffer is a solution that resists sudden changes in pH.

Concept Review Exercise

1. Explain how a buffer prevents large changes in pH.

Answer

1. A buffer has components that react with both strong acids and strong bases to resist sudden changes in pH.

Exercises

1. Describe a buffer. What two related chemical components are required to make a buffer?
2. Can a buffer be made by combining a strong acid with a strong base? Why or why not?
3. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. HCl and NaCl
    2. HNO_2 and $NaNO_2$
    3. NH_4NO_3 and HNO_3
    4. NH_4NO_3 and NH_3
4. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. H_3PO_4 and Na_3PO_4
    2. $NaHCO_3$ and Na_2CO_3
    3. $NaNO_3$ and $Ca(NO_3)_2$
    4. HN_3 and NH_3
5. For each combination in Exercise 3 that is a buffer, write the chemical equations for the reactions of the buffer components when a strong acid and a strong base is added.
6. For each combination in Exercise 4 that is a buffer, write the chemical equations for the reaction of the buffer components when a strong acid and a strong base is added.
7. The complete phosphate buffer system is based on four substances: H_3PO_4, $H_2PO_4^-$, HPO_4^{2-}, and PO_4^{3-}. What different buffer solutions can be made from these substances?

8. Explain why NaBr cannot be a component in either an acidic or a basic buffer.
9. Explain why $Mg(NO_3)_2$ cannot be a component in either an acidic or a basic buffer.

Answers

1. A buffer resists sudden changes in pH. It has a weak acid or base and a salt of that weak acid or base.
2. No. Combining a strong acid and a strong base will produce salt and water. Excess strong acid or strong base will not act as a buffer.
3. 1. not a buffer
   2. buffer
   3. not a buffer
   4. buffer

4. 1. not a buffer

2. buffer

3. not a buffer

4. not buffer

5. 3b: strong acid: $H^+ + NO_2^- \rightarrow HNO_2$; strong base: $OH^- + HNO_2 \rightarrow H_2O + NO_2^-$; 3d: strong acid: $H^+ + NH_3 \rightarrow NH_4^+$; strong base: $OH^- + NH_4^+ \rightarrow H_2O + NH_3$
6. 4b: strong acid: $H^+ + CO_3^{2-} \rightarrow HCO3^-$; strong base: $OH^- + HCO_3^- \rightarrow H_2O + CO_3^{2-}$;
7. Buffers can be made by combining H_3PO_4 and $H_2PO_4^-$, $H_2PO_4^-$ and HPO_4^{2-}, and HPO_4^{2-} and PO_4^{3-}.
8. NaBr splits up into two ions in solution, Na^+ and Br^-. Na^+ will not react with any added base knowing that NaOH is a strong base. Br- will not react with any added acid knowing that HBr is a strong acid. Because NaBr will not react with any added base or acid, it does not resist change in pH and is not a buffer.
9. $Mg(NO_3)_2$ includes two types of ions, Mg^{2+} and NO_3^-. $Mg(OH)_2$ is strong base and completely dissociates (100% falls apart), so Mg^{2+} will not react with any added base (0% combines with OH^-). HNO_3 is strong acid and completely dissociates (100% falls apart), so NO_3^- will not react with any added acid (0% combines with H^+). Because $Mg(NO_3)_2$ will not react with any added base or acid, it does not resist change in pH and is not a buffer.

10.6: Buffers is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.5: Buffers** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.7: End-of-Chapter Material

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

The earliest chemical definition of an acid, the **Arrhenius definition**, says that an **acid** is a compound that increases the amount of hydrogen ion (H^+) in aqueous solution. An **Arrhenius base** is a compound that increases the amount of hydroxide ion (OH^-) in aqueous solution. While most bases are named as ionic hydroxide compounds, aqueous acids have a naming system unique to acids. Acids and bases react together in a characteristic chemical reaction called **neutralization**, in which the products are water and a salt. The principles of stoichiometry, along with the balanced chemical equation for a reaction between an acid and a base, can be used to determine how much of one compound will react with a given amount of the other.

A **Brønsted-Lowry acid** is any substance that donates a proton to another substance. A **Brønsted-Lowry base** is any substance that accepts a proton from another substance. The reaction of ammonia with water to make ammonium ions and hydroxide ions can be used to illustrate Brønsted-Lowry acid and base behavior.

Some compounds can either donate or accept protons, depending on the circumstances. Such compounds are called **amphiprotic**. Water is one example of an amphiprotic compound. One result of water being amphiprotic is that a water molecule can donate a proton to another water molecule to make a hydronium ion and a hydroxide ion. This process is called the **autoionization of water** and occurs in any sample of water.

Not all acids and bases are equal in chemical strength. A **strong acid** is an acid whose molecules are all dissociated into ions in aqueous solution. Hydrochloric acid is an example of a strong acid. Similarly, a **strong base** is a base whose molecules are dissociated into ions in aqueous solution. Sodium hydroxide is an example of a strong base. Any acid or base whose molecules are not all dissociated into ions in aqueous solution is a **weak acid** or a **weak base**. Solutions of weak acids and weak bases reach a **chemical equilibrium** between the un-ionized form of the compound and the dissociated ions. It is a dynamic equilibrium because acid and base molecules are constantly dissociating into ions and reassociating into neutral molecules.

The **pH** scale is a scale used to express the concentration of hydrogen ions in solution. A neutral solution, neither acidic nor basic, has a pH of 7. Acidic solutions have a pH lower than 7, while basic solutions have a pH higher than 7.

Buffers are solutions that resist dramatic changes in pH when an acid or a base is added to them. They contain a weak acid and a salt of that weak acid, or a weak base and a salt of that weak base. When a buffer is present, any strong acid reacts with the anion of the salt, forming a weak acid and minimizing the presence of hydrogen ions in solution. Any strong base reacts with the weak acid, minimizing the amount of additional hydroxide ions in solution. However, buffers only have limited **capacity**; there is a limit to the amount of strong acid or strong base any given amount of buffer will react with.

10.7: End-of-Chapter Material is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **10.S: Acids and Bases (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

11: Organic Chemistry of Hydrocarbons

Hydrocarbons are organic compounds that contain **only** carbon and hydrogen. The four general classes of hydrocarbons are: alkanes, alkenes, alkynes and arenes. Aromatic compounds derive their names from the fact that many of these compounds in the early days of discovery were grouped because they were oils with fragrant odors.

11.1: Introduction to Organic Chemistry
11.2: Organic Chemistry
11.3: Structures and Names of Alkanes
11.4: Branched-Chain Alkanes
11.5: Condensed Structural and Line-Angle Formulas
11.6: Cycloalkanes
11.7: IUPAC Nomenclature
11.8: Alkenes- Structure and Nomenclature
11.9: Alkenes- Cis-Tran Isomerization
11.10: Alkynes
11.11: Aromatic Compounds
11.12: Aromatic Compounds- Structure and Nomenclature
11.13: Physical Properties of Hydrocarbons
11.14: Chapter Summary

11: Organic Chemistry of Hydrocarbons is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

11.1: Introduction to Organic Chemistry

Learning Objectives

- Define *organic chemistry*.
- Identify organic molecules as alkanes, alkenes, alkynes, alcohols, or carboxylic acids.

When methane was mentioned previously, we described it as the simplest organic compound. In this section, we introduce organic chemistry more formally. Organic chemistry is the study of the chemistry of carbon compounds. Carbon is singled out because it has a chemical diversity unrivaled by any other chemical element. Its diversity is based on the following:

- Carbon atoms bond reasonably strongly with other carbon atoms.
- Carbon atoms bond reasonably strongly with atoms of other elements.
- Carbon atoms make a large number of covalent bonds (four).

Curiously, elemental carbon is not particularly abundant. It does not even appear in the list of the most common elements in Earth's crust. Nevertheless, all living things consist of organic compounds. Most organic chemicals are covalent compounds, which is why we introduce organic chemistry here. By convention, compounds containing carbonate ions and bicarbonate ions, as well as carbon dioxide and carbon monoxide, are not considered part of organic chemistry, even though they contain carbon.

Alkanes

The simplest organic compounds are the **hydrocarbons**, compounds composed of carbon and hydrogen atoms only. Some hydrocarbons have only single bonds and appear as a chain (which can be a straight chain or can have branches) of carbon atoms also bonded to hydrogen atoms. These hydrocarbons are called **alkanes (saturated hydrocarbons)**. Each alkane has a characteristic, systematic name depending on the number of carbon atoms in the molecule. These names consist of a stem that indicates the number of carbon atoms in the chain plus the ending -ane. The stem meth- means one carbon atom, so methane is an alkane with one carbon atom. Similarly, the stem eth- means two carbon atoms; ethane is an alkane with two carbon atoms. Continuing, the stem prop- means three carbon atoms, so propane is an alkane with three carbon atoms. The stem but- means four carbon atoms; butane is an alkane with four carbon atoms. Figure 11.1.1 gives the Lewis structures, condensed structural formulas and molecular formulas of the four simplest alkanes. In the **condensed structural formula**, the covalent bonds are understood to exist between each carbon and the hydrogens associated with it, as well as between carbon atoms.




Figure 11.1.1: Formulas and Molecular Models of the Four Simplest Alkanes. The four smallest alkanes are methane, ethane, propane and butane.

Alkenes

Some hydrocarbons have one or more carbon–carbon double bonds (denoted C=C). These hydrocarbons are called alkenes. Figure 11.1.2 shows the formulas and the molecular models of the two simplest alkenes. Note that the names of alkenes have the same stem as the alkane with the same number of carbon atoms in its chain but have the ending *-ene*. Thus, ethene is an alkene with two carbon atoms per molecule, and propene is a compound with three carbon atoms and one double bond.





Figure **11.1.2**: Formulas and Molecular Models of the Two Simplest Alkenes. Ethene is commonly called ethylene, while propene is commonly called propylene.

Alkynes

Alkynes are hydrocarbons with a carbon–carbon triple bond (denoted C≡C) as part of their carbon skeleton. Below is the formula and the molecular model of the simplest alkyne and its systematic name. Its common name is acetylene. Its chemical formula is $\mathbf{C_2H_2}$.



Ethyne

The names for alkynes have the same stems as for alkanes but with the ending *-yne*.

To Your Health: Saturated and Unsaturated Fats

Hydrocarbons are not the only compounds that can have carbon–carbon double bonds. A group of compounds called fats can have them as well, and their presence or absence in the human diet is becoming increasingly correlated with health issues.

Fats are combinations of long-chain organic compounds (fatty acids) and glycerol ($C_3H_8O_3$). (For more information on fats, see Chapter 17) The long carbon chains can have either all single bonds, in which case the fat is classified as *saturated*, or one or more double bonds, in which case it is a *monounsaturated* or a *polyunsaturated* fat, respectively. Saturated fats are typically solids at room temperature; beef fat (tallow) is one example. Mono- or polyunsaturated fats are likely to be liquids at room temperature and are often called oils. Olive oil, flaxseed oil, and many fish oils are mono- or polyunsaturated fats.

Studies have linked higher amounts of saturated fats in people's diets with a greater likelihood of developing heart disease, high cholesterol, and other diet-related diseases. In contrast, increases in unsaturated fats (either mono- or polyunsaturated) have been linked to a lower incidence of certain diseases. Thus, there have been an increasing number of recommendations by government bodies and health associations to decrease the proportion of saturated fat and increase the proportion of unsaturated fat in the diet. Most of these organizations also recommend decreasing the total amount of fat in the diet.

Recently, certain fats called trans fats have been implicated in the presence of heart disease. These are fats from animal sources and are also produced when liquid oils are exposed to partial hydrogenation, an industrial process that increases their saturation. Trans fats are used in many prepared and fried foods. Because they bring with them the health risks that naturally occurring saturated fats do, there has been some effort to better quantify the presence of trans fats in food products. US law now requires that food labels list the amount of trans fat in each serving.

Since the early 1900's, the US Department of Agriculture has been providing science-based dietary guidelines for the public. The most current version is the MyPlate illustration that gives a simple, visual picture of how much of what kind of foods make up a good, balanced diet. It recommends minimizing daily intake of sugars, the "bad fats", trans and saturated fat, and sodium. "Good fats", unsaturated fats or oils, are not considered a food group but do contain essential nutrients and therefore are included as part of a healthy eating pattern. The difference as simple as the difference between a single and double carbon–carbon bond, good and bad fats, can have a significant impact on health.



MyPlate is a reminder that everything you eat and drink matters. Source: Image courtesy of the USDA, www.mypyramid.gov/downloads/MiniPoster.pdf.

Functional Groups

The carbon–carbon double and triple bonds are examples of functional groups in organic chemistry. A functional group is a specific structural arrangement of atoms or bonds that imparts a characteristic chemical reactivity to a molecule. Alkanes have no functional group. A carbon–carbon double bond is considered a functional group because carbon–carbon double bonds chemically react in specific ways that differ from reactions of alkanes (for example, under certain circumstances, alkenes react with water); a carbon–carbon triple bond also undergoes certain specific chemical reactions. In the remainder of this section, we introduce two other common functional groups.

If an OH group (also called a hydroxyl group) is substituted for a hydrogen atom in a hydrocarbon molecule, the compound is an alcohol. Alcohols are named using the parent hydrocarbon name but with the final *-e* dropped and the suffix *-ol* attached. The two simplest alcohols are methanol and ethanol. Figure **11.1.4** shows their formulas along with a molecular model of each.




Figure **11.1.4**: The Two Simplest Organic Alcohol Compounds. Alcohols have an OH functional group in the molecule.

Cholesterol, described in the chapter-opening essay, has an alcohol functional group, as its name implies.

Alcohol

Ethanol (also called ethyl alcohol) is the alcohol in alcoholic beverages. Other alcohols include methanol (or methyl alcohol), which is used as a solvent and a cleaner, and isopropyl alcohol (or rubbing alcohol), which is used as a medicinal disinfectant. Neither methanol nor isopropyl alcohol should be ingested, as they are toxic even in small quantities.

Another important family of organic compounds has a carboxyl group, in which a carbon atom is double-bonded to an oxygen atom and to an OH group. Compounds with a carboxyl functional group are called carboxylic acids, and their names end in *-oic acid*. Figure **11.1.5** shows the formulas and the molecular models of the two simplest carboxylic acids, perhaps best known by the common names formic acid and acetic acid. The carboxyl group is sometimes written in molecules as COOH.





Figure **11.1.5**: The Two Smallest Organic Acids. The two smallest carboxylic acids are formic acid (found in the stingers of ants) and acetic acid (found in vinegar).


The condensed structures of methanoic acid and ethanoic acid are **HCOOH** and **CH_3COOH**, respectively.


Many organic compounds are considerably more complex than the examples described here. Many compounds, such as cholesterol discussed in the chapter-opening essay, contain more than one functional group. The formal names can also be quite complex.

Example 11.1.1

Identify the functional group(s) in each molecule as a double bond, a triple bond, an alcohol, or a carboxyl.

a. 

b. 

c. 

d. **$CH_3CH_2CH_2CH_2OH$**

Answer a

This molecule has a double bond and a carboxyl functional group.

Answer b

This molecule has an alcohol functional group.

Answer c

This molecule has a carbon-carbon double bond and a carboxyl functional group.


Answer d


This molecule has an alcohol functional group.


Exercise 11.1.1

Identify the functional group(s) in each molecule as a double bond, a triple bond, an alcohol, or a carboxyl.

a. $CH_3C \equiv CCH_3$

b. 

c. 

d. 

Answer a:

triple bond (alkyne)

Answer b:

carboxyl group

Answer c:

alcohol group

Answer d:

double bond (alkene) and carboxyl group

Career Focus: Forensic Chemist

The main job of a forensic chemist is to identify unknown materials and their origins. Although forensic chemists are most closely associated in the public mind with crime labs, they are employed in pursuits as diverse as tracing evolutionary patterns in living organisms, identifying environmental contaminants, and determining the origin of manufactured chemicals.

In a crime lab, the forensic chemist has the job of identifying the evidence so that a crime can be solved. The unknown samples may consist of almost anything—for example, paint chips, blood, glass, cloth fibers, drugs, or human remains. The forensic chemist subjects them to a variety of chemical and instrumental tests to discover what the samples are. Sometimes these samples are extremely small, but sophisticated forensic labs have state-of-the-art equipment capable of identifying the smallest amount of unknown sample.

Another aspect of a forensic chemist's job is testifying in court. Judges and juries need to be informed about the results of forensic analyses, and it is the forensic chemist's job to explain those results. Good public-speaking skills, along with a broad background in chemistry, are necessary to be a successful forensic chemist.

Key Takeaways

- Organic chemistry is the study of the chemistry of carbon compounds.
- Organic molecules can be classified according to the types of elements and bonds in the molecules.

11.1: Introduction to Organic Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **4.6: Organic Chemistry** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.2: Organic Chemistry

Learning Objectives

- To recognize the composition and properties typical of organic and inorganic compounds.

Scientists of the 18th and early 19th centuries studied compounds obtained from plants and animals and labeled them *organic* because they were isolated from "organized" (living) systems. Compounds isolated from nonliving systems, such as rocks and ores, the atmosphere, and the oceans, were labeled *inorganic*. For many years, scientists thought organic compounds could be made by only living organisms because they possessed a vital force found only in living systems. The vital force theory began to decline in 1828, when the German chemist Friedrich Wöhler synthesized urea from inorganic starting materials. He reacted silver cyanate ($AgOCN$) and ammonium chloride (NH_4Cl), expecting to get ammonium cyanate (NH_4OCN). What he expected is described by the following equation.

$$AgOCN + NH_4Cl \rightarrow AgCl + NH_4OCN$$

However, when heated, ammonium cyanate continues to react and rearranges into urea (NH_2CONH_2), a well-known organic material readily isolated from urine.

$$NH_4OCN \longrightarrow NH_2CONH_2$$

This observation led to a series of experiments in which a wide variety of organic compounds were made from inorganic starting materials. The vital force theory gradually went away as chemists learned that they could make many organic compounds in the laboratory. Today organic chemistry is the study of the chemistry of the carbon compounds, and inorganic chemistry is the study of the chemistry of all other elements. It may seem strange that we divide chemistry into two branches—one that considers compounds of only one element and one that covers the 100-plus remaining elements. However, this division seems more reasonable when we consider that of tens of millions of compounds that have been characterized, the overwhelming majority are carbon compounds.

> The word *organic* has different meanings. Organic fertilizer, such as cow manure, is organic in the original sense; it is derived from living organisms. Organic foods generally are foods grown without synthetic pesticides or fertilizers. Organic chemistry is the chemistry of compounds of carbon.

Carbon is unique among the other elements in that its atoms can form stable covalent bonds with each other and with atoms of other elements in a multitude of variations. The resulting molecules can contain from one to millions of carbon atoms. We previously surveyed organic chemistry by dividing its compounds into families based on functional groups. We begin with the simplest members of a family and then move on to molecules that are organic in the original sense—that is, they are made by and found in living organisms. These complex molecules (all containing carbon) determine the forms and functions of living systems and are the subject of biochemistry.

Organic compounds, like inorganic compounds, obey all the natural laws. Often there is no clear distinction in the chemical or physical properties among organic and inorganic molecules. Nevertheless, it is useful to compare typical members of each class, as in Table **11.2.1**.

Table **11.2.1**: General Contrasting Properties and Examples of Organic and Inorganic Compounds

Organic	Hexane		Inorganic	NaCl
low melting points	−95°C		high melting points	801°C
low boiling points	69°C		high boiling points	1,413°C
low solubility in water; high solubility in nonpolar solvents	insoluble in water; soluble in gasoline		greater solubility in water; low solubility in nonpolar solvents	soluble in water; insoluble in gasoline
flammable	highly flammable		nonflammable	nonflammable
aqueous solutions do not conduct electricity	nonconductive		aqueous solutions conduct electricity	conductive in aqueous solution
exhibit covalent bonding	covalent bonds		exhibit ionic bonding	ionic bonds

Keep in mind, however, that there are exceptions to every category in this table. To further illustrate typical differences among organic and inorganic compounds, Table **11.2.1** also lists properties of the inorganic compound sodium chloride (common table salt, NaCl) and the organic compound hexane (C_6H_{14}), a solvent that is used to extract soybean oil from soybeans (among other uses). Many compounds can be classified as organic or inorganic by the presence or absence of certain typical properties, as illustrated in Table **11.2.1**.

11.2: Organic Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.1: Organic Chemistry** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.3: Structures and Names of Alkanes

Learning Objectives

- To identify and name simple (straight-chain) alkanes given formulas and write formulas for straight-chain alkanes given their names.

We begin our study of organic chemistry with the hydrocarbons, the simplest organic compounds, which are composed of carbon and hydrogen atoms only. As we noted, there are several different kinds of hydrocarbons. They are distinguished by the types of bonding between carbon atoms and the properties that result from that bonding. Hydrocarbons with only carbon-to-carbon single bonds (C–C) and existing as a continuous chain of carbon atoms also bonded to hydrogen atoms are called alkanes (or saturated hydrocarbons). *Saturated*, in this case, means that each carbon atom is bonded to four other atoms (hydrogen or carbon)—the most possible; there are no double or triple bonds in the molecules.

> The word *saturated* has the same meaning for hydrocarbons as it does for the dietary fats and oils: the molecule has no carbon-to-carbon double bonds (C=C).

We previously introduced the three simplest alkanes—methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) and they are shown again in Figure **11.3.1**.




Figure **11.3.1**: The Three Simplest Alkanes

The flat representations shown do not accurately portray bond angles or molecular geometry. Methane has a tetrahedral shape that chemists often portray with wedges indicating bonds coming out toward you and dashed lines indicating bonds that go back away from you. An ordinary solid line indicates a bond in the plane of the page. Recall that the VSEPR theory correctly predicts a tetrahedral shape for the methane molecule (Figure **11.3.2**).




Figure **11.3.2**: The Tetrahedral Methane Molecule

Methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) are the beginning of a series of compounds in which any two members in a sequence differ by one carbon atom and two hydrogen atoms—namely, a CH_2 unit. The first 10 members of this series are given in Table **11.3.1**.

Table **11.3.1**: The First 10 Straight-Chain Alkanes

Name	Molecular Formula (C_nH_{2n+2})	Condensed Structural Formula	Number of Possible Isomers
methane	CH_4	CH_4	—
ethane	C_2H_6	CH_3CH_3	—
propane	C_3H_8	$CH_3CH_2CH_3$	—
butane	C_4H_{10}	$CH_3CH_2CH_2CH_3$	2
pentane	C_5H_{12}	$CH_3CH_2CH_2CH_2CH_3$	3
hexane	C_6H_{14}	$CH_3CH_2CH_2CH_2CH_2CH_3$	5
heptane	C_7H_{16}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_3$	9
octane	C_8H_{18}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	18
nonane	C_9H_{20}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	35
decane	$C_{10}H_{22}$	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	75

Consider the series in Figure **11.3.3**. The sequence starts with C_3H_8, and a CH_2 unit is added in each step moving up the series. Any family of compounds in which adjacent members differ from each other by a definite factor (here a CH_2 group) is called a homologous series. The members of such a series, called *homologs*, have properties that vary in a regular and predictable manner. The principle of *homology* gives organization to organic chemistry in much the same way that the periodic table gives organization to inorganic chemistry. Instead of a bewildering array of individual carbon compounds, we can study a few members of a homologous series and from them deduce some of the properties of other compounds in the series.




Figure **11.3.3**: Members of a Homologous Series. Each succeeding formula incorporates one carbon atom and two hydrogen atoms more than the previous formula.

The principle of homology allows us to write a general formula for alkanes: C_nH_{2n+2}. Using this formula, we can write a molecular formula for any alkane with a given number of carbon atoms. For example, an alkane with eight carbon atoms has the molecular formula $C_8H_{(2\times 8)+2} = C_8H_{18}$.

Key Takeaway

- Simple alkanes exist as a homologous series, in which adjacent members differ by a CH_2 unit.

11.3: Structures and Names of Alkanes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.2: Structures and Names of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.4: Branched-Chain Alkanes


Learning Objectives

- To learn how alkane molecules can have branched chains and recognize compounds that are isomers.


We can write the structure of butane (C_4H_{10}) by stringing four carbon atoms in a row,

–C–C–C–C–

and then adding enough hydrogen atoms to give each carbon atom four bonds:



The compound butane has this structure, but there is another way to put 4 carbon atoms and 10 hydrogen atoms together. Place 3 of the carbon atoms in a row and then branch the fourth one off the middle carbon atom:



Now we add enough hydrogen atoms to give each carbon four bonds.




Figure **11.4.1**).





Figure **11.4.1**: Butane and Isobutane. The ball-and-stick models of these two compounds show them to be isomers; both have the molecular formula C_4H_{10}.


Notice that C_4H_{10} is depicted with a bent chain in Figure **11.4.1**. The four-carbon chain may be bent in various ways because the groups can rotate freely about the C–C bonds. However, this rotation does not change the identity of the compound. It is important to realize that bending a chain does *not* change the identity of the compound; all of the following represent the same compound, butane:



The structure of isobutane shows a continuous chain of three carbon atoms only, with the fourth attached as a branch off the middle carbon atom of the continuous chain, which is different from the structures of butane (compare the two structures in Figure **11.4.1**.

Unlike C_4H_{10}, the compounds methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) do not exist in isomeric forms because there is only one way to arrange the atoms in each formula so that each carbon atom has four bonds.

Next beyond C_4H_{10} in the homologous series is pentane. Each compound has the same molecular formula: C_5H_{12}. (Table 12.1.1 from the previous section has a column identifying the number of possible isomers for the first 10 straight-chain alkanes.) The compound at the far left is pentane because it has all five carbon atoms in a continuous chain. The compound in the middle is isopentane; like isobutane, it has a one CH_3 branch off the second carbon atom of the continuous chain. The compound at the far right, discovered after the other two, was named neopentane (from the Greek *neos*, meaning "new"). Although all three have the same molecular formula, they have different properties, including boiling points: pentane, 36.1°C; isopentane, 27.7°C; and neopentane, 9.5°C.



A continuous (unbranched) chain of carbon atoms is often called a *straight chain* even though the tetrahedral arrangement about each carbon gives it a zigzag shape. Straight-chain alkanes are sometimes called *normal alkanes*, and their names are given the prefix *n*-. For example, butane is called *n*-butane. We will not use that prefix here because it is not a part of the system established by the International Union of Pure and Applied Chemistry.

11.4: Branched-Chain Alkanes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.3: Branched-Chain Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.5: Condensed Structural and Line-Angle Formulas

Learning Objectives

- Write condensed structural formulas for alkanes given complete structural formulas.
- Draw line-angle formulas given structural formulas.

We use several kinds of formulas to describe organic compounds. A *molecular formula* shows only the kinds and numbers of atoms in a molecule. For example, the molecular formula C_4H_{10} tells us there are 4 carbon atoms and 10 hydrogen atoms in a molecule, but it doesn't distinguish between butane and isobutane. A structural formula shows all the carbon and hydrogen atoms and the bonds attaching them. Thus, structural formulas identify the specific isomers by showing the order of attachment of the various atoms.

Unfortunately, structural formulas are difficult to type/write and take up a lot of space. Chemists often use condensed structural formulas to alleviate these problems. The condensed formulas show hydrogen atoms right next to the carbon atoms to which they are attached, as illustrated for butane:



The ultimate condensed formula is a line-angle formula, in which carbon atoms are implied at the corners and ends of lines, and each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds. For example, we can represent pentane ($CH_3CH_2CH_2CH_2CH_3$) and isopentane [$(CH_3)_2CHCH_2CH_3$] as follows:



Parentheses in condensed structural formulas indicate that the enclosed grouping of atoms is attached to the adjacent carbon atom.

Key Takeaways

- Condensed chemical formulas show the hydrogen atoms (or other atoms or groups) right next to the carbon atoms to which they are attached.
- Line-angle formulas imply a carbon atom at the corners and ends of lines. Each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds.

11.5: Condensed Structural and Line-Angle Formulas is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.4: Condensed Structural and Line-Angle Formulas** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.6: Cycloalkanes

Learning Objectives

- To name cycloalkanes given their formulas and write formulas for these compounds given their names.

The hydrocarbons we have encountered so far have been composed of molecules with open-ended chains of carbon atoms. When a chain contains three or more carbon atoms, the atoms can join to form *ring* or *cyclic* structures. The simplest of these cyclic hydrocarbons has the formula C_3H_6. Each carbon atom has two hydrogen atoms attached (Figure **11.6.1**) and is called cyclopropane.





Figure **11.6.1**: Ball-and-Spring Model of Cyclopropane. The springs are bent to join the carbon atoms.

To Your Health: Cyclopropane as an Anesthetic

With its boiling point of −33°C, cyclopropane is a gas at room temperature. It is also a potent, quick-acting anesthetic with few undesirable side effects in the body. It is no longer used in surgery, however, because it forms explosive mixtures with air at nearly all concentrations.

The cycloalkanes—cyclic hydrocarbons with only single bonds—are named by adding the prefix *cyclo-* to the name of the open-chain compound having the same number of carbon atoms as there are in the ring. Thus the name for the cyclic compound C_4H_8 is cyclobutane. The carbon atoms in cyclic compounds can be represented by *line-angle formulas* that result in regular geometric figures. Keep in mind, however, that each corner of the geometric figure represents a carbon atom plus as many hydrogen atoms as needed to give each carbon atom four bonds.



Some cyclic compounds have substituent groups attached. Example **11.6.1** interprets the name of a cycloalkane with a single substituent group.

Example 11.6.1

Draw the structure for each compound.

a. cyclopentane
b. methylcyclobutane


Solution a

The name *cyclopentane* indicates a cyclic (cyclo) alkane with five (pent-) carbon atoms. It can be represented as a pentagon.



Solution b

The name *methylcyclobutane* indicates a cyclic alkane with four (but-) carbon atoms in the cyclic part. It can be represented as a square with a CH_3 group attached.



? Exercise 11.6.1

Draw the structure for each compound.

a. cycloheptane
b. ethylcyclohexane

The properties of cyclic hydrocarbons are generally quite similar to those of the corresponding open-chain compounds. So cycloalkanes (with the exception of cyclopropane, which has a highly strained ring) act very much like noncyclic alkanes. Cyclic structures containing five or six carbon atoms, such as cyclopentane and cyclohexane, are particularly stable. We will see later that some carbohydrates (sugars) form five- or six-membered rings in solution.

The cyclopropane ring is strained because the C–C–C angles are 60°, and the preferred (tetrahedral) bond angle is 109.5°. (This strain is readily evident when you try to build a ball-and-stick model of cyclopropane; see Figure **11.6.1**.) Cyclopentane and cyclohexane rings have little strain because the C–C–C angles are near the preferred angles.

Key Takeaway

- Many organic compounds have cyclic structures.

11.6: Cycloalkanes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.9: Cycloalkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.7: IUPAC Nomenclature

Learning Objectives

- To name alkanes by the IUPAC system and write formulas for alkanes given IUPAC names










As noted in previously, the number of isomers increases rapidly as the number of carbon atoms increases. There are 3 pentanes, 5 hexanes, 9 heptanes, and 18 octanes. It would be difficult to assign unique individual names that we could remember. A systematic way of naming hydrocarbons and other organic compounds has been devised by the International Union of Pure and Applied Chemistry (IUPAC). These rules, used worldwide, are known as the IUPAC System of Nomenclature. (Some of the names we used earlier, such as isobutane, isopentane, and neopentane, do not follow these rules and are called *common names*.) A stem name (Table **11.7.1**) indicates the number of carbon atoms in the longest continuous chain (LCC). Atoms or groups attached to this carbon chain, called *substituents*, are then named, with their positions indicated by numbers. For now, we will consider only those substituents called alkyl groups.

Table **11.7.1**: Stems That Indicate the Number of Carbon Atoms in Organic Molecules

Stem	Number
meth-	1
eth-	2
prop-	3
but-	4
pent-	5
hex-	6
hept-	7
oct-	8
non-	9
dec-	10

An alkyl group is a group of atoms that results when one hydrogen atom is removed from an alkane. The group is named by replacing the *-ane* suffix of the parent hydrocarbon with *-yl*. For example, the -CH_3 group derived from methane (CH_4) results from subtracting one hydrogen atom and is called a *methyl group*. The alkyl groups we will use most frequently are listed in Table **11.7.2**. Alkyl groups are not independent molecules; they are parts of molecules that we consider as a unit to name compounds systematically.


Table **11.7.2**: Common Alkyl Groups

Parent Alkane		Alkyl Group		Condensed Structural Formula
methane		methyl		$CH_3–$
ethane		ethyl		$CH_3CH_2–$
propane		propyl		$CH_3CH_2CH_2–$
		isopropyl		$(CH_3)_2CH–$
butane		butyl*		$CH_3CH_2CH_2CH_2–$

*There are four butyl groups, two derived from butane and two from isobutane. We will introduce the other three where appropriate.

Simplified IUPAC rules for naming alkanes are as follows (demonstrated in Example **11.7.1**).

1. **Name alkanes according to the LCC (longest continuous chain) of carbon atoms in the molecule (rather than the total number of carbon atoms).** This LCC, considered the parent chain, determines the base name, to which we add the suffix *-ane* to indicate that the molecule is an alkane.
2. **If the hydrocarbon is branched, number the carbon atoms of the LCC.** Numbers are assigned in the direction that gives the lowest numbers to the carbon atoms with attached substituents. Hyphens are used to separate numbers from the names of substituents; commas separate numbers from each other. (The LCC need not be written in a straight line; for example, the LCC in the following has five carbon atoms.)






3. **Place the names of the substituent groups in alphabetical order before the name of the parent compound.** If the same alkyl group appears more than once, the numbers of all the carbon atoms to which it is attached are expressed. If the same group appears more than once on the same carbon atom, the number of that carbon atom is repeated as many times as the group appears. Moreover, the number of identical groups is indicated by the Greek prefixes *di-*, *tri-*, *tetra-*, and so on. These prefixes are *not* considered in determining the alphabetical order of the substituents. For example, ethyl is listed before dimethyl; the di- is simply ignored. The last alkyl group named is prefixed to the name of the parent alkane to form one word.

When these rules are followed, every unique compound receives its own exclusive name. The rules enable us to not only name a compound from a given structure but also draw a structure from a given name. The best way to learn how to use the IUPAC system is to put it to work, not just memorize the rules. It's easier than it looks.

Example 11.7.1

Name each compound.




1. 
2. 
3. 

Solution

1. The LCC has five carbon atoms, and so the parent compound is pentane (rule 1). There is a methyl group (rule 2) attached to the second carbon atom of the pentane chain. The name is therefore 2-methylpentane.
2. The LCC has six carbon atoms, so the parent compound is hexne (rule 1). Methyl groups (rule 2) are attached to the second and fifth carbon atoms. The name is 2,5-dimethylhexane.
3. The LCC has eight carbon atoms, so the parent compound is octane (rule 1). There are methyl and ethyl groups (rule 2), both attached to the fourth carbon atom (counting from the *right* gives this carbon atom a lower number; rule 3). The correct name is thus 4-ethyl-4-methyloctane.

Exercise 11.7.1

Name each compound.

1. 
2. 
3. 

Example 11.7.2


Draw the structure for each compound.

a. 2,3-dimethylbutane
b. 4-ethyl-2-methylheptane


Solution a

In drawing structures, always start with the parent chain.


The parent chain is butane, indicating four carbon atoms in the LCC.



Then add the groups at their proper positions. You can number the parent chain from either direction as long as you are consistent; just don't change directions before the structure is done. The name indicates two methyl (CH_3) groups, one on the second carbon atom and one on the third.




Finally, fill in all the hydrogen atoms, keeping in mind that each carbon atom must have four bonds.




Solution b


The parent chain is heptane in this case, indicating seven carbon atoms in the LCC.



Adding the groups at their proper positions gives



Filling in all the hydrogen atoms gives the following condensed structural formulas:



Note that the bonds (dashes) can be shown or not; sometimes they are needed for spacing.

? Exercise 11.7.2

Draw the structure for each compound.

a. 4-ethyloctane
b. 3-ethyl-2-methylpentane
c. 3,3,5-trimethylheptane

Key Takeaway

- Alkanes have both common names and systematic names, specified by IUPAC.

11.7: IUPAC Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **12.5: IUPAC Nomenclature** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.8: Alkenes- Structure and Nomenclature

Learning Objectives

- To name alkenes given formulas and write formulas for alkenes given names.

As noted before, alkenes are hydrocarbons with carbon-to-carbon double bonds ($R_2C{=}CR_2$) and alkynes are hydrocarbons with carbon-to-carbon triple bonds (R–C≡C–R). Collectively, they are called unsaturated hydrocarbons because they have fewer hydrogen atoms than does an alkane with the same number of carbon atoms, as is indicated in the following general formulas:




Some representative alkenes—their names, structures, and physical properties—are given in Table **11.8.1**.

Table **11.8.1**: Physical Properties of Some Selected Alkenes

IUPAC Name	Molecular Formula	Condensed Structural Formula	Melting Point (°C)	Boiling Point (°C)
ethene	C_2H_4	$CH_2{=}CH_2$	–169	–104
propene	C_3H_6	$CH_2{=}CHCH_3$	–185	–47
1-butene	C_4H_8	$CH_2{=}CHCH_2CH_3$	–185	–6
1-pentene	C_5H_{10}	$CH_2{=}CH(CH_2)_2CH_3$	–138	30
1-hexene	C_6H_{12}	$CH_2{=}CH(CH_2)_3CH_3$	–140	63
1-heptene	C_7H_{14}	$CH_2{=}CH(CH_2)_4CH_3$	–119	94
1-octene	C_8H_{16}	$CH_2{=}CH(CH_2)_5CH_3$	–102	121

We used only condensed structural formulas in Table **11.8.1**. Thus, $CH_2{=}CH_2$ stands for



The double bond is shared by the two carbons and does not involve the hydrogen atoms, although the condensed formula does not make this point obvious. Note that the molecular formula for ethene is C_2H_4, whereas that for ethane is C_2H_6.

The first two alkenes in Table **11.8.1**, ethene and propene, are most often called by their common names—ethylene and propylene, respectively (Figure **11.8.1**). Ethylene is a major commercial chemical. The US chemical industry produces about 25 billion kilograms of ethylene annually, more than any other synthetic organic chemical. More than half of this ethylene goes into the manufacture of polyethylene, one of the most familiar plastics. Propylene is also an important industrial chemical. It is converted to plastics, isopropyl alcohol, and a variety of other products.





Figure **11.8.1**: Ethene and Propene. The ball-and-spring models of ethene/ethylene (a) and propene/propylene (b) show their respective shapes, especially bond angles.

Although there is only one alkene with the formula C_2H_4 (ethene) and only one with the formula C_3H_6 (propene), there are several alkenes with the formula C_4H_8.


Here are some basic rules for naming alkenes from the International Union of Pure and Applied Chemistry (IUPAC):


1. The longest chain of carbon atoms *containing the double bond* is considered the parent chain. It is named using the same stem as the alkane having the same number of carbon atoms but ends in *-ene* to identify it as an alkene. Thus the compound CH_2=$CHCH_3$ is *propene*.
2. If there are four or more carbon atoms in a chain, we must indicate the position of the double bond. The carbons atoms are numbered so that the first of the two that are doubly bonded is given the lower of the two possible numbers.The compound CH_3CH=$CHCH_2CH_3$, for example, has the double bond between the second and third carbon atoms. Its name is 2-pentene (not 3-pentene).
3. Substituent groups are named as with alkanes, and their position is indicated by a number. Thus, the structure below is 5-methyl-2-hexene. Note that the numbering of the parent chain is always done in such a way as to give the double bond the lowest number, even if that causes a substituent to have a higher number. *The double bond always has priority in numbering.*



Example 11.8.1

Name each compound.

a. 

b. 

Solution


a. The longest chain containing the double bond has five carbon atoms, so the compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the fourth carbon atom (rule 3), so the compound's name is 4-methyl-2-pentene.

b. The longest chain containing the double bond has five carbon atoms, so the parent compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the third carbon atom (rule 3), so the compound's name is 3-methyl-2-pentene.

Exercise 11.8.1

Name each compound.

a. $CH_3CH_2CH_2CH_2CH_2CH{=}CHCH_3$

b. 

Answer

Just as there are cycloalkanes, there are *cycloalkenes*. These compounds are named like alkenes, but with the prefix *cyclo-* attached to the beginning of the parent alkene name.

Example 11.8.2


Draw the structure for each compound.

a. 3-methyl-2-pentene
b. cyclohexene


Solution

a.

First write the parent chain of five carbon atoms: C–C–C–C–C. Then add the double bond between the second and third carbon atoms:




Now place the methyl group on the third carbon atom and add enough hydrogen atoms to give each carbon atom a total of four bonds.



b

First, consider what each of the three parts of the name means. *Cyclo* means a ring compound, *hex* means 6 carbon atoms, and *-ene* means a double bond.



Exercise 11.8.2

Draw the structure for each compound.

a. 2-ethyl-1-hexene
b. cyclopentene

Key Takeaway

- Alkenes are hydrocarbons with a carbon-to-carbon double bond.

11.8: Alkenes- Structure and Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **13.1: Alkenes- Structures and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.9: Alkenes- Cis-Tran Isomerization

Learning Objectives

- Recognize that alkenes that can exist as cis-trans isomers.
- Classify isomers as cis or trans.
- Draw structures for cis-trans isomers given their names.

There is free rotation about the carbon-to-carbon single bonds (C–C) in alkanes. In contrast, the structure of alkenes requires that the carbon atoms of a double bond and the two atoms bonded to each carbon atom all lie in a single plane, and that each doubly bonded carbon atom lies in the center of a triangle. This part of the molecule's structure is rigid; rotation about doubly bonded carbon atoms is *not* possible without rupturing the bond. Look at the two chlorinated hydrocarbons in Figure **11.9.1**.




Figure **11.9.1**: *Rotation about Bonds. In 1,2-dichloroethane (a), free rotation about the C–C bond allows the two structures to be interconverted by a twist of one end relative to the other. In 1,2-dichloroethene (b), restricted rotation about the double bond means that the relative positions of substituent groups above or below the double bond are significant.*

In 1,2-dichloroethane (part (a) of Figure **11.9.1**), there is free rotation about the C–C bond. The two models shown represent exactly the same molecule; they are *not* isomers. You can draw structural formulas that look different, but if you bear in mind the possibility of this free rotation about single bonds, you should recognize that these two structures represent the same molecule:



In 1,2-dichloroethene (Figure **11.9.1***b*), however, restricted rotation about the double bond means that the relative positions of substituent groups above or below the double bond become significant. This leads to a special kind of isomerism. The isomer in which the two chlorine (Cl) atoms lie on the same side of the molecule is called the cis isomer (Latin *cis*, meaning "on this side") and is named cis-1,2-dichloroethene. The isomer with the two Cl atoms on opposite sides of the molecule is the trans isomer (Latin *trans*, meaning "across") and is named *trans*-1,2-dichloroethene. These two compounds are cis-trans isomers (or geometric isomers), compounds that have different configurations (groups permanently in different places in space) because of the presence of a rigid structure in their molecule.

Consider the alkene with the condensed structural formula $CH_3CH{=}CHCH_3$. We could name it 2-butene, but there are actually two such compounds; the double bond results in cis-trans isomerism (Figure **11.9.2**).




Figure **11.9.2**: Ball-and-Spring Models of (a) Cis-2-Butene and (b) Trans-2-Butene. Cis-trans isomers have different physical, chemical, and physiological properties.

Cis-2-butene has both methyl groups on the same side of the molecule. *Trans*-2-butene has the methyl groups on opposite sides of the molecule. Their structural formulas are as follows:



***Figure* 11.9.3**: *Models of (left) Cis-2-Butene and (right) Trans-2-Butene.*

Note, however, that the presence of a double bond does **not** necessarily lead to cis-trans isomerism (Figure **11.9.4**). We can draw two *seemingly* different propenes:



***Figure* 11.9.4**: *Different views of the propene molecule (flip vertically). These are not isomers.*

However, these two structures are not really different from each other. If you could pick up either molecule from the page and flip it over top to bottom, you would see that the two formulas are identical. Thus there are two requirements for cis-trans isomerism:

1. Rotation must be restricted in the molecule.
2. There must be **two nonidentical groups** on **each** doubly bonded carbon atom.

In these propene structures, the second requirement for cis-trans isomerism is not fulfilled. One of the doubly bonded carbon atoms does have two different groups attached, but the rules require that *both* carbon atoms have two different groups. In general, the following statements hold true in cis-trans isomerism:

- Alkenes with a $C{=}CH_2$ unit do not exist as cis-trans isomers.
- Alkenes with a $C{=}CR_2$ unit, where the two R groups are the same, do not exist as cis-trans isomers.
- Alkenes of the type R–CH=CH–R can exist as cis and trans isomers; cis if the two R groups are on the same side of the carbon-to-carbon double bond, and trans if the two R groups are on opposite sides of the carbon-to-carbon double bond.

Advanced Note: E/Z Isomerization

If a molecule has a C=C bond with one non-hydrogen group attached to each of the carbons, cis/trans nomenclature descried above is enough to describe it. However, if you have three different groups (or four), then the cis/trans approach is insufficient to describe the different isomers, since we do not know which two of the three groups are being described. For example, if you have a C=C bond, with a methyl group and a bromine on one carbon , and an ethyl group on the other, it is neither trans nor cis, since it is not clear whether the ethyl group is trans to the bromine or the methyl. This is addressed with a more advanced E,Z Convention [E] [E] [E] discussed elsewhere.

Cis-trans isomerism also occurs in cyclic compounds. In ring structures, groups are unable to rotate about any of the ring carbon–carbon bonds. Therefore, groups can be either on the same side of the ring (cis) or on opposite sides of the ring (trans). For our purposes here, we represent all cycloalkanes as planar structures, and we indicate the positions of the groups, either above or below the plane of the ring.



Example 11.9.1

Which compounds can exist as cis-trans (geometric) isomers? Draw them.

1. CHCl=CHBr
2. CH_2=$CBrCH_3$
3. $(CH_3)_2C$=$CHCH_2CH_3$
4. CH_3CH=$CHCH_2CH_3$


Solution

All four structures have a double bond and thus meet rule 1 for cis-trans isomerism.

1. This compound meets rule 2; it has two nonidentical groups on *each* carbon atom (H and Cl on one and H and Br on the other). It exists as both cis and trans isomers:





2. This compound has two hydrogen atoms on one of its doubly bonded carbon atoms; it fails rule 2 and does not exist as cis and trans isomers.
3. This compound has two methyl (CH_3) groups on one of its doubly bonded carbon atoms. It fails rule 2 and does not exist as cis and trans isomers.
4. This compound meets rule 2; it has two nonidentical groups on *each* carbon atom and exists as both cis and trans isomers:



Exercise 11.9.1

Which compounds can exist as cis-trans isomers? Draw them.

a. CH_2=$CHCH_2CH_2CH_3$
b. CH_3CH=$CHCH_2CH_3$
c. CH_3CH_2CH=$CHCH_2CH_3$
d. 

e. 

Key Takeaway

- Cis-trans (geometric) isomerism exists when there is restricted rotation in a molecule and there are two nonidentical groups on *each* doubly bonded carbon atom.

11.9: Alkenes- Cis-Tran Isomerization is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **13.2: Cis-Trans Isomers (Geometric Isomers)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.10: Alkynes

Learning Objectives

- Describe the general physical and chemical properties of alkynes.
- Name alkynes given formulas and write formulas for alkynes given names.

The simplest alkyne—a hydrocarbon with carbon-to-carbon triple bond—has the molecular formula C_2H_2 and is known by its common name—acetylene (Figure **11.10.1**). Its structure is H–C≡C–H.




Figure **11.10.1**: Ball-and-Spring Model of Acetylene. Acetylene (ethyne) is the simplest member of the alkyne family.

Acetylene is used in oxyacetylene torches for cutting and welding metals. The flame from such a torch can be very hot. Most acetylene, however, is converted to chemical intermediates that are used to make vinyl and acrylic plastics, fibers, resins, and a variety of other products.

Alkynes are similar to alkenes in both physical and chemical properties. For example, alkynes undergo many of the typical addition reactions of alkenes. The International Union of Pure and Applied Chemistry (IUPAC) names for alkynes parallel those of alkenes, except that the family ending is *-yne* rather than *-ene*. The IUPAC name for acetylene is ethyne. The names of other alkynes are illustrated in the following exercises.

Key Takeaway

- Alkynes are hydrocarbons with carbon-to-carbon triple bonds and properties much like those of alkenes.

11.10: Alkynes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **13.6: Alkynes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.11: Aromatic Compounds

Learning Objectives


- To describe the bonding in benzene and the way typical reactions of benzene differ from those of the alkenes.

Next we consider a class of hydrocarbons with molecular formulas like those of unsaturated hydrocarbons, but which, unlike the alkenes, do not readily undergo addition reactions. These compounds comprise a distinct class, called aromatic hydrocarbons, with unique structures and properties. We start with the simplest of these compounds. Benzene (C_6H_6) is of great commercial importance, but it also has noteworthy health effects.

The formula C_6H_6 seems to indicate that benzene has a high degree of unsaturation. (Hexane, the saturated hydrocarbon with six carbon atoms has the formula C_6H_{14}—eight more hydrogen atoms than benzene.) However, despite the seeming low level of saturation, benzene is rather unreactive. It does not, for example, react readily with bromine, which, is a test for unsaturation.

Benzene is a liquid that smells like gasoline, boils at 80°C, and freezes at 5.5°C. It is the aromatic hydrocarbon produced in the largest volume. It was formerly used to decaffeinate coffee and was a significant component of many consumer products, such as paint strippers, rubber cements, and home dry-cleaning spot removers. It was removed from many product formulations in the 1950s, but others continued to use benzene in products until the 1970s when it was associated with leukemia deaths. Benzene is still important in industry as a precursor in the production of plastics (such as Styrofoam and nylon), drugs, detergents, synthetic rubber, pesticides, and dyes. It is used as a solvent for such things as cleaning and maintaining printing equipment and for adhesives such as those used to attach soles to shoes. Benzene is a natural constituent of petroleum products, but because it is a known carcinogen, its use as an additive in gasoline is now limited.

To explain the surprising properties of benzene, chemists suppose the molecule has a cyclic, hexagonal, planar structure of six carbon atoms with one hydrogen atom bonded to each. We can write a structure with alternate single and double bonds, either as a full structural formula or as a line-angle formula:



However, these structures do not explain the unique properties of benzene. Furthermore, experimental evidence indicates that all the carbon-to-carbon bonds in benzene are equivalent, and the molecule is unusually stable. Chemists often represent benzene as a hexagon with an inscribed circle:



The inner circle indicates that the valence electrons are shared equally by all six carbon atoms (that is, the electrons are *delocalized*, or spread out, over all the carbon atoms). It is understood that each corner of the hexagon is occupied by one carbon atom, and each carbon atom has one hydrogen atom attached to it. Any other atom or groups of atoms substituted for a hydrogen atom must be shown bonded to a particular corner of the hexagon. We use this modern symbolism, but many scientists still use the earlier structure with alternate double and single bonds.

To Your Health: Benzene and Us

Most of the benzene used commercially comes from petroleum. It is employed as a starting material for the production of detergents, drugs, dyes, insecticides, and plastics. Once widely used as an organic solvent, benzene is now known to have both short- and long-term toxic effects. The inhalation of large concentrations can cause nausea and even death due to respiratory or heart failure, while repeated exposure leads to a progressive disease in which the ability of the bone marrow to make new blood cells is eventually destroyed. This results in a condition called *aplastic anemia*, in which there is a decrease in the numbers of both the red and white blood cells.

Key Takeaway

- Aromatic hydrocarbons appear to be unsaturated, but they have a special type of bonding and do not undergo addition reactions.

11.11: Aromatic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **13.7: Aromatic Compounds- Benzene** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.12: Aromatic Compounds- Structure and Nomenclature

Learning Objectives

- Recognize aromatic compounds from structural formulas.
- Name aromatic compounds given formulas.
- Write formulas for aromatic compounds given their names.


Historically, benzene-like substances were called aromatic hydrocarbons because they had distinctive aromas. Today, an aromatic compound is any compound that contains a benzene ring or has certain benzene-like properties (but not necessarily a strong aroma). You can recognize the aromatic compounds in this text by the presence of one or more benzene rings in their structure. Some representative aromatic compounds and their uses are listed in Table **11.12.1**, where the benzene ring is represented as C_6H_5.


Table **11.12.1**: Some Representative Aromatic Compounds


Name	Structure	Typical Uses
aniline	$C_6H_5–NH_2$	starting material for the synthesis of dyes, drugs, resins, varnishes, perfumes; solvent; vulcanizing rubber
benzoic acid	$C_6H_5–COOH$	food preservative; starting material for the synthesis of dyes and other organic compounds; curing of tobacco
bromobenzene	$C_6H_5–Br$	starting material for the synthesis of many other aromatic compounds; solvent; motor oil additive
nitrobenzene	$C_6H_5–NO_2$	starting material for the synthesis of aniline; solvent for cellulose nitrate; in soaps and shoe polish
phenol	$C_6H_5–OH$	disinfectant; starting material for the synthesis of resins, drugs, and other organic compounds
toluene	$C_6H_5–CH_3$	solvent; gasoline octane booster; starting material for the synthesis of benzoic acid, benzaldehyde, and many other organic compounds


Example 11.12.1

Which compounds are aromatic?

1. 

2. 

3. 


4. 


Solution

1. The compound has a benzene ring (with a chlorine atom substituted for one of the hydrogen atoms); it is aromatic.
2. The compound is cyclic, but it does not have a benzene ring; it is not aromatic.
3. The compound has a benzene ring (with a propyl group substituted for one of the hydrogen atoms); it is aromatic.
4. The compound is cyclic, but it does not have a benzene ring; it is not aromatic.

? Exercise 11.12.1

Which compounds are aromatic?

1. 

2. 

3. 

In the International Union of Pure and Applied Chemistry (IUPAC) system, aromatic hydrocarbons are named as derivatives of benzene. Figure **11.12.1** shows four examples. In these structures, it is immaterial whether the single substituent is written at the top, side, or bottom of the ring: a hexagon is symmetrical, and therefore all positions are equivalent.




Figure **11.12.1**: Some Benzene Derivatives. These compounds are named in the usual way with the group that replaces a hydrogen atom named as a substituent group: Cl as chloro, Br as bromo, I as iodo, NO_2 as nitro, and CH_3CH_2 as ethyl.

Although some compounds are referred to exclusively by IUPAC names, some are more frequently denoted by common names, as is indicated in Table **11.12.1**.



When there is more than one substituent, the corners of the hexagon are no longer equivalent, so we must designate the relative positions. There are three possible disubstituted benzenes, and we can use numbers to distinguish them (Figure **11.12.2**). We start numbering at the carbon atom to which one of the groups is attached and count toward the carbon atom that bears the other substituent group by the shortest path.






Figure **11.12.2**: The Three Isomeric Dichlorobenzenes


In Figure **11.12.2**, common names are also used: the prefix *ortho* (*o*-) for 1,2-disubstitution, *meta* (*m*-) for 1,3-disubstitution, and *para* (*p*-) for 1,4-disubstitution. The substituent names are listed in alphabetical order. The first substituent is given the lowest number. When a common name is used, the carbon atom that bears the group responsible for the name is given the number 1:




Example 11.12.2

Name each compound using both the common name and the IUPAC name.

1. 


2. 

3. 

Solution


1. The benzene ring has two chlorine atoms (dichloro) in the first and second positions. The compound is *o*-dichlorobenzene or 1,2-dichlorobenzene.
2. The benzene ring has a methyl (CH_3) group. The compound is therefore named as a derivative of toluene. The bromine atom is on the fourth carbon atom, counting from the methyl group. The compound is *p*-bromotoluene or 4-bromotoluene.
3. The benzene ring has two nitro (NO_2) groups in the first and third positions. It is *m*-dinitrobenzene or 1,3-dinitrobenzene.


4. Note: The nitro (NO_2) group is a common substituent in aromatic compounds. Many nitro compounds are explosive, most notably 2,4,6-trinitrotoluene (TNT).




? Exercise 11.12.2

Name each compound using both the common name and the IUPAC name.

1. 

2. 

3. 


- Sometimes an aromatic group is found as a substituent bonded to a nonaromatic entity or to another aromatic ring. The group of atoms remaining when a hydrogen atom is removed from an aromatic compound is called an aryl group. The most common aryl group is derived from benzene (C_6H_6) by removing one hydrogen atom (C_6H_5) and is called a *phenyl* group, from *pheno*, an old name for benzene.



Phenyl group 2-Phenylheptane

Polycyclic Aromatic Hydrocarbons

Some common aromatic hydrocarbons consist of fused benzene rings—rings that share a common side. These compounds are called polycyclic aromatic hydrocarbons (PAHs).



Naphthalene Anthracene Phenanthrene

The three examples shown here are colorless, crystalline solids generally obtained from coal tar. Naphthalene has a pungent odor and is used in mothballs. Anthracene is used in the manufacture of certain dyes. Steroids, a large group of naturally occurring substances, contain the phenanthrene structure.

To Your Health: Polycyclic Aromatic Hydrocarbons and Cancer

The intense heating required for distilling coal tar results in the formation of PAHs. For many years, it has been known that workers in coal-tar refineries are susceptible to a type of skin cancer known as tar cancer. Investigations have shown that a number of PAHs






are carcinogens. One of the most active carcinogenic compounds, benzopyrene, occurs in coal tar and has also been isolated from cigarette smoke, automobile exhaust gases, and charcoal-broiled steaks. It is estimated that more than 1,000 t of benzopyrene are emitted into the air over the United States each year. Only a few milligrams of benzopyrene per kilogram of body weight are required to induce cancer in experimental animals.

Biologically Important Compounds with Benzene Rings

Substances containing the benzene ring are common in both animals and plants, although they are more abundant in the latter. Plants can synthesize the benzene ring from carbon dioxide, water, and inorganic materials. Animals cannot synthesize it, but they are dependent on certain aromatic compounds for survival and therefore must obtain them from food. Phenylalanine, tyrosine, and tryptophan (essential amino acids) and vitamins K, B_2 (riboflavin), and B_9 (folic acid) all contain the benzene ring. Many important drugs, a few of which are shown in Table **11.12.2**, also feature a benzene ring.

> So far we have studied only aromatic compounds with carbon-containing rings. However, many cyclic compounds have an element other than carbon atoms in the ring. These compounds, called *heterocyclic compounds*, are discussed later. Some of these are heterocyclic aromatic compounds.

Table **11.12.2**: Some Drugs That Contain a Benzene Ring

Name	Structure
aspirin	
acetaminophen	
ibuprofen	
amphetamine	
sulfanilamide	

Key Takeaway

- Aromatic compounds contain a benzene ring or have certain benzene-like properties; for our purposes, you can recognize aromatic compounds by the presence of one or more benzene rings in their structure.

11.12: Aromatic Compounds- Structure and Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **13.8: Structure and Nomenclature of Aromatic Compounds** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.13: Physical Properties of Hydrocarbons

Learning Objectives

- To identify the physical properties of alkanes and describe trends in these properties.

Because alkanes have relatively predictable physical properties and undergo relatively few chemical reactions other than combustion, they serve as a basis of comparison for the properties of many other organic compound families. Let's consider their physical properties first.

Table **11.13.1** describes some of the properties of some of the first 10 straight-chain alkanes. Because alkane molecules are nonpolar, they are insoluble in water, which is a polar solvent, but are soluble in nonpolar and slightly polar solvents. Consequently, alkanes themselves are commonly used as solvents for organic substances of low polarity, such as fats, oils, and waxes. Nearly all alkanes have densities less than 1.0 g/mL and are therefore less dense than water (the density of H_2O is 1.00 g/mL at 20°C). These properties explain why oil and grease do not mix with water but rather float on its surface.

Table **11.13.1**: Physical Properties of Some Alkanes

Molecular Name	Formula	Melting Point (°C)	Boiling Point (°C)	Density (20°C)*	Physical State (at 20°C)
methane	CH_4	–182	–164	0.668 g/L	gas
ethane	C_2H_6	–183	–89	1.265 g/L	gas
propane	C_3H_8	–190	–42	1.867 g/L	gas
butane	C_4H_{10}	–138	–1	2.493 g/L	gas
pentane	C_5H_{12}	–130	36	0.626 g/mL	liquid
hexane	C_6H_{14}	–95	69	0.659 g/mL	liquid
octane	C_8H_{18}	–57	125	0.703 g/mL	liquid
decane	$C_{10}H_{22}$	–30	174	0.730 g/mL	liquid

*Note the change in units going from gases (grams per liter) to liquids (grams per milliliter). Gas densities are at 1 atm pressure.



Figure **11.13.1**: Oil Spills. Crude oil coats the water's surface in the Gulf of Mexico after the *Deepwater Horizon* oil rig sank following an explosion. The leak was a mile below the surface, making it difficult to estimate the size of the spill. One liter of oil can create a slick 2.5 hectares (6.3 acres) in size. This and similar spills provide a reminder that hydrocarbons and water don't mix. Source: Photo courtesy of NASA Goddard / MODIS Rapid Response Team, NASA.gov, Topics, Earth Features, Oil Spill(opens in new window) [www.nasa.gov].

Looking Closer: Gas Densities and Fire Hazards

Table **11.13.1** indicates that the first four members of the alkane series are gases at ordinary temperatures. Natural gas is composed chiefly of methane, which has a density of about 0.67 g/L. The density of air is about 1.29 g/L. Because natural gas is less dense than air, it rises. When a natural-gas leak is detected and shut off in a room, the gas can be removed by opening an upper window. On the other hand, bottled gas can be either propane (density 1.88 g/L) or butanes (a mixture of butane and isobutane; density about 2.5 g/L). Both are much heavier than air (density 1.2 g/L). If bottled gas escapes into a building, it collects near the floor. This presents a much more serious fire hazard than a natural-gas leak because it is more difficult to rid the room of the heavier gas.

Also shown in Table **11.13.1** are the boiling points of the straight-chain alkanes increase with increasing molar mass. This general rule holds true for the straight-chain homologs of all organic compound families. Larger molecules have greater surface areas and consequently interact more strongly; more energy is therefore required to separate them. For a given molar mass, the boiling points of alkanes are relatively low because these nonpolar molecules have only weak dispersion forces to hold them together in the liquid state.

Looking Closer: An Alkane Basis for Properties of Other Compounds

An understanding of the physical properties of the alkanes is important in that petroleum and natural gas and the many products derived from them—gasoline, bottled gas, solvents, plastics, and more—are composed primarily of alkanes. This understanding is also vital because it is the basis for describing the properties of other organic and biological compound families. For example, large portions of the structures of lipids consist of nonpolar alkyl groups. Lipids include the dietary fats and fatlike compounds called phospholipids and sphingolipids that serve as structural components of living tissues. These compounds have both polar and nonpolar groups, enabling them to bridge the gap between water-soluble and water-insoluble phases. This characteristic is essential for the selective permeability of cell membranes.




Figure **11.13.2**: Tripalmitin (a), a typical fat molecule, has long hydrocarbon chains typical of most lipids. Compare these chains to hexadecane (b), an alkane with 16 carbon atoms

Key Takeaway

- Alkanes are nonpolar compounds that are low boiling and insoluble in water.

11.13: Physical Properties of Hydrocarbons is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **12.6: Physical Properties of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.14: Chapter Summary

Template:HideTOC

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the summary and ask yourself how they relate to the topics in the chapter.

Organic chemistry is the chemistry of carbon compounds, and **inorganic chemistry** is the chemistry of all the other elements. Carbon atoms can form stable covalent bonds with other carbon atoms and with atoms of other elements, and this property allows the formation the tens of millions of organic compounds. **Hydrocarbons** contain only hydrogen and carbon atoms.

Hydrocarbons in which each carbon atom is bonded to four other atoms are called **alkanes** or **saturated hydrocarbons**. They have the general formula C_nH_{2n+2}. Any given alkane differs from the next one in a series by a CH_2 unit. Any family of compounds in which adjacent members differ from each other by a definite factor is called a **homologous series**.

Carbon atoms in alkanes can form straight chains or branched chains. Two or more compounds having the same molecular formula but different structural formulas are **isomers** of each other. There are no isomeric forms for the three smallest alkanes; beginning with C_4H_{10}, all other alkanes have isomeric forms.

A **structural formula** shows all the carbon and hydrogen atoms and how they are attached to one another. A **condensed structural formula** shows the hydrogen atoms right next to the carbon atoms to which they are attached. A **line-angle formula** is a formula in which carbon atoms are implied at the corners and ends of lines. Each carbon atom is understood to be attached to enough hydrogen atoms to give each carbon atom four bonds.

The **IUPAC System of Nomenclature** provides rules for naming organic compounds. An **alkyl group** is a unit formed by removing one hydrogen atom from an alkane.

The physical properties of alkanes reflect the fact that alkane molecules are nonpolar. Alkanes are insoluble in water and less dense than water.

Alkanes are generally unreactive toward laboratory acids, bases, oxidizing agents, and reducing agents. They do burn (undergo **combustion reactions**).

Alkanes react with halogens by substituting one or more halogen atoms for hydrogen atoms to form **halogenated hydrocarbons**. An **alkyl halide (haloalkane)** is a compound resulting from the replacement of a hydrogen atom of an alkane with a halogen atom.

Cycloalkanes are hydrocarbons whose molecules are closed rings rather than straight or branched chains. A **cyclic hydrocarbon** is a hydrocarbon with a ring of carbon atoms.

11.14: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

12: Organic Functional Groups - Structure and Nomenclature

Functional groups are specific groups of atoms or bonds within molecules that are responsible for the characteristic chemical reactions of those molecules. The same functional group will undergo the same or similar chemical reaction(s) regardless of the size of the molecule it is a part of. However, its relative reactivity can be modified by other functional groups nearby. The atoms of functional groups are linked to each other and to the rest of the molecule by covalent bonds. When the group of covalently bound atoms bears a net charge, the group is referred to more properly as a polyatomic ion or a complex ion.

12.1: Organic Compounds with Functional Groups
12.2: 9.2 Alcohols and Phenols- Nomenclature and Classification
12.3: Glycols and Glycerol
12.4: Ethers
12.5: Aldehydes and Ketones- Structure and Names
12.6: Organic Sulfur Compounds
12.7: Carboxylic Acids- Structures and Names
12.8: Esters- Structures and Names
12.9: Phosphate Esters
12.10: Amines
12.11: Amides

12: Organic Functional Groups - Structure and Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.








12.1: Organic Compounds with Functional Groups

Learning Objectives

- to describe functional groups and explain why they are useful in the study of organic chemistry.

Previously, we considered several kinds of hydrocarbons. Now we examine some of the many organic compounds that contain functional groups. We first introduced the idea of the functional group, a specific structural arrangement of atoms or bonds that imparts a characteristic chemical reactivity to the molecule. If you understand the behavior of a particular functional group, you will know a great deal about the general properties of that class of compounds. In this chapter, we make a brief yet systematic study of some of organic compound families. Each family is based on a common, simple functional group that contains an oxygen atom or a nitrogen atom. Some common functional groups are listed in Table **12.1.1**.

Table **12.1.1**: Selected Organic Functional Groups

Name of Family	General Formula	Functional Group	Suffix*
alkane	RH	none	*-ane*
alkene	$R_2C{=}CR_2$		*-ene*
alkyne	RC≡CR	–C≡C–	*-yne*
alcohol	ROH	–OH	*-ol*
thiol	RSH	–SH	*-thiol*
ether	ROR	–O–	ether
aldehyde			*-al*
ketone			*-one*
carboxylic acid			*-oic acid*

*Ethers do not have a suffix in their common name; all ethers end with the word *ether*.

Summary

The functional group, a structural arrangement of atoms and/or bonds, is largely responsible for the properties of organic compound families.

12.1: Organic Compounds with Functional Groups is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.1: Organic Compounds with Functional Groups** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.2: 9.2 Alcohols and Phenols- Nomenclature and Classification


Skills to Develop

- Identify the general structure for an alcohol.
- Identify the structural feature that classifies alcohols as primary, secondary, or tertiary.
- Name alcohols with both common names and IUPAC names

An alcohol is an organic compound with a hydroxyl (OH) functional group on an aliphatic carbon atom. Because OH is the functional group of all alcohols, we often represent alcohols by the general formula ROH, where R is an alkyl group. Alcohols are common in nature. Most people are familiar with ethyl alcohol (ethanol), the active ingredient in alcoholic beverages, but this compound is only one of a family of organic compounds known as alcohols. The family also includes such familiar substances as cholesterol and the carbohydrates. Methanol (CH_3OH) and ethanol (CH_3CH_2OH) are the first two members of the homologous series of alcohols.

Nomenclature of Alcohols

Alcohols with one to four carbon atoms are frequently called by common names, in which the name of the alkyl group is followed by the word *alcohol*:




According to the International Union of Pure and Applied Chemistry (IUPAC), alcohols are named by changing the ending of the parent alkane name to *-ol*. Here are some basic IUPAC rules for naming alcohols:

1. The longest continuous chain (LCC) of carbon atoms containing the OH group is taken as the parent compound—an alkane with the same number of carbon atoms. The chain is numbered from the end nearest the OH group.
2. The number that indicates the position of the OH group is prefixed to the name of the parent hydrocarbon, and the *-e* ending of the parent alkane is replaced by the suffix *-ol*. (In cyclic alcohols, the carbon atom bearing the OH group is designated C1, but the 1 is not used in the name.) Substituents are named and numbered as in alkanes.
3. If more than one OH group appears in the same molecule (polyhydroxy alcohols), suffixes such as *-diol* and *-triol* are used. In these cases, the *-e* ending of the parent alkane is retained.
4. Figure **12.2.1** shows some examples of the application of these rules.


Rules 1 and 2




2-methylbutan-2-ol



3,5-dimethylhexan-1-ol




6-methylheptan-3-ol



2-Bromo-5-chlorocyclopentanol

Rule 3






***Figure* 12.2.1:** *IUPAC Rules for Alcohols. The names and structures of some alcohols demonstrate the use of IUPAC rules.*

ethane-1,2-diol
1,2-ethanediol
(ethylene glycol)

propane-1,2,3-triol
(glycerol)


- The longest continuous chain (LCC) of carbon atoms containing the OH group is taken as the parent compound—an alkane with the same number of carbon atoms. The chain is numbered from the end nearest the OH group.
- The number that indicates the position of the OH group is prefixed to the name of the parent hydrocarbon, and the *-e* ending of the parent alkane is replaced by the suffix *-ol*. (In cyclic alcohols, the carbon atom bearing the OH group is designated C1, but the 1 is not used in the name.) Substituents are named and numbered as in alkanes.
- If more than one OH group appears in the same molecule (polyhydroxy alcohols), suffixes such as *-diol* and *-triol* are used. In these cases, the *-e* ending of the parent alkane is retained.
-

Figure 12.2.1 shows some examples of the application of these rules.


Rules 1 and 2




2-methylbutan-2-ol



3,5-dimethylhexan-1-ol




6-methylheptan-3-ol




2-Bromo-5-chlorocyclopentanol

Rule 3



ethane-1,2-diol
1,2-ethanediol
(ethylene glycol)



propane-1,2,3-triol
(glycerol)

***Figure* 12.2.1:** *IUPAC Rules for Alcohols. The names and structures of some alcohols demonstrate the use of IUPAC rules.*

Example 9.2.2


Draw the structure for each compound.

1. 2-hexanol
2. 3-methyl-2-pentanol


SOLUTION

1. The ending *-ol* indicates an alcohol (the OH functional group), and the *hex-* stem tells us that there are six carbon atoms in the LCC. We start by drawing a chain of six carbon atoms: –C–C–C–C–C–C–.

   The 2 indicates that the OH group is attached to the second carbon atom.




Finally, we add enough hydrogen atoms to give each carbon atom four bonds.



2. The ending *-ol* indicates an OH functional group, and the *pent-* stem tells us that there are five carbon atoms in the LCC. We start by drawing a chain of five carbon atoms:

   –C–C–C–C–C–

   The numbers indicate that there is a methyl (CH_3) group on the third carbon atom and an OH group on the second carbon atom.



Exercise 9.2.2

Draw the structure for each compound.

a. 3-heptanol

b. 2-methyl-3-hexanol

Classification of Alcohols

Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.


Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.



Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.




Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.



Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.

Table 9.2.1: *Classification and Nomenclature of Some Alcohols*

Condensed Structural Formula	Class of Alcohol	Common Name	IUPAC Name
CH_3OH	—	methyl alcohol	methanol
CH_3CH_2OH	primary	ethyl alcohol	ethanol
$CH_3CH_2CH_2OH$	primary	propyl alcohol	1-propanol
$(CH_3)_2CHOH$	secondary	isopropyl alcohol	2-propanol
$CH_3CH_2CH_2CH_2OH$	primary	butyl alcohol	1-butanol
$CH_3CH_2CHOHCH_3$	secondary	*sec*-butyl alcohol	2-butanol
$(CH_3)_2(CH_3)_2CHCH_2OH$	primary	isobutyl alcohol	2-methyl-1-propanol
$(CH_3)_3COH$	tertiary	*tert*-butyl alcohol	2-methyl-2-propanol
	secondary	cyclohexyl alcohol	cyclohexanol

Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.

Some of the properties of alcohols depend on the number of carbon atoms attached to the specific carbon atom that is attached to the OH group. Alcohols can be grouped into three classes on this basis.




Phenols

Compounds in which an OH group is attached directly to an aromatic ring are designated ArOH and called phenols. Phenols differ from alcohols in that they are slightly acidic in water. They react with aqueous sodium hydroxide (NaOH) to form salts.

$$ArOH_{(aq)} + NaOH_{(aq)} \rightarrow ArONa_{(aq)} + H_2O \tag{12.2.1}$$

The parent compound, C_6H_5OH, is itself called phenol. (An old name, emphasizing its slight acidity, was *carbolic acid*.) Phenol is a white crystalline compound that has a distinctive (“hospital smell”) odor.




To Your Health: Phenols and Us

Phenols are widely used as antiseptics (substances that kill microorganisms on living tissue) and as disinfectants (substances intended to kill microorganisms on inanimate objects such as furniture or floors). The first widely used antiseptic was phenol. Joseph Lister used it for antiseptic surgery in 1867. Phenol is toxic to humans, however, and can cause severe burns when applied to the skin. In the bloodstream, it is a systemic poison—that is, one that is carried to and affects all parts of the body. Its severe side effects led to searches for safer antiseptics, a number of which have been found.



An operation in 1753, painted by Gaspare Traversi, of a surgery before antiseptics were used.


One safer phenolic antiseptic is 4-hexylresorcinol (4-hexyl-1,3-dihydroxybenzene; resorcinol is the common name for 1,3-dihydroxybenzene, and 4-hexylresorcinol has a hexyl group on the fourth carbon atom of the resorcinol ring). It is much more powerful than phenol as a germicide and has fewer undesirable side effects. Indeed, it is safe enough to be used as the active ingredient in some mouthwashes and throat lozenges.



The compound 4-hexylresorcinol is mild enough to be used as the active ingredient in antiseptic preparations for use on the skin.

Concept Review Exercises

1. Is isobutyl alcohol primary, secondary, or tertiary? Explain.



2. What is the LCC in 2-ethyl-1-hexanol? What is taken as the LCC in naming the compound? Explain.
3. How do phenols differ from alcohols in terms of structure and properties?
4. How do phenols differ in properties from aromatic hydrocarbons?

Answers

1. primary; the carbon atom bearing the OH group is attached to only one other carbon atom
2. 7 carbon atoms; the 6-atom chain includes the carbon atom bearing the OH group
3. Phenols have an OH group attached directly to an aromatic ring. Phenols are weakly acidic.
4. Phenols have an OH group and are somewhat soluble in water.


Key Takeaways


- In the IUPAC system, alcohols are named by changing the ending of the parent alkane name to *-ol*.
- Alcohols are classified according to the number of carbon atoms attached to the carbon atom that is attached to the OH group.
- Phenols are compounds in which an OH group is attached directly to an aromatic ring.
- Many phenols are used as antiseptics.

Exercises


1. Name each alcohol and classify it as primary, secondary, or tertiary.


   a. $CH_3CH_2CH_2CH_2CH_2CH_2OH$


   b. 

   c. 

2. Name each alcohol and classify it as primary, secondary, or tertiary.

   a. 

   b. 

   c. 


3. Draw the structure for each alcohol.

   a. 3-hexanol
   b. 3,3-dimethyl-2-butanol
   c. cyclobutanol

4. Draw the structure for each alcohol.


   a. cyclopentanol
   b. 4-methyl-2-hexanol
   c. 4,5-dimethyl-3-heptanol


5. Name each compound.

   a. 

   b. 

6. Name each compound.

a. 

b. 

7. Draw the structure for each compound.

a. *m*-iodophenol
b. *p*-methylphenol (*p*-cresol)

8. Draw the structure for each compound.


a. 2,4,6-trinitrophenol (picric acid)
b. 3,5-diethylphenol


Answers


1.

a. 1-hexanol; primary
b. 3-hexanol; secondary
c. 3,3-dibromo-2-methyl-2-butanol; tertiary

3.

a. 

b. 

c. 

5

a. *o*-nitrophenol
b. *p*-bromophenol

7.

a. 

b. 


12.2: 9.2 Alcohols and Phenols- Nomenclature and Classification is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

12.3: Glycols and Glycerol

Learning Objectives


- To describe the structure and uses of some common polyhydric alcohols.

Alcohols with two OH groups on adjacent carbon atoms are commonly known as glycols. The most important of these is 1,2-ethanediol (the common name is ethylene glycol), a sweet, colorless, somewhat viscous liquid.




Ethylene glycol

Another common glycol, 1,2-propanediol, is commonly called propylene glycol. Its physical properties are quite similar to those of ethylene glycol.



Propylene glycol

Commonly called glycerol or glycerin, 1,2,3-propanetriol is the most important trihydroxy alcohol. Like the two glycols, it is a sweet, syrupy liquid. Glycerol is a product of the hydrolysis of fats and oils.




Glycerol

Ethylene glycol is the main ingredient in many antifreeze mixtures for automobile radiators. The two OH groups lead to extensive intermolecular hydrogen bonding. This results in a high boiling point—198°C; thus ethylene glycol does not boil away when it is used as an antifreeze. It is also completely miscible with water. A solution of 60% ethylene glycol in water freezes at −49°C (−56°F) and thus protects an automobile radiator down to that temperature. Ethylene glycol is also used in the manufacture of polyester fiber and magnetic film used in tapes for recorders and computers.

To Your Health: Glycols and Human Health

Ethylene glycol is quite toxic. Because it is sweet, pets often lap up spills of leaked antifreeze from a garage floor or driveway. Sometimes people, especially children, drink it. As with methanol, its toxicity is due to a metabolite. Liver enzymes oxidize ethylene glycol to oxalate ion.



Ethylene glycol Oxalate ion


In the kidneys, the oxalate ion combines with the calcium (Ca^{2+}) ion, precipitating as calcium oxalate (CaC_2O_4).

$$Ca^{2+}(aq) + C_2O_4^{2-}(aq) \rightarrow CaC_2O_4(s)$$

These crystals cause renal damage and can lead to kidney failure and death.

Although propylene glycol has physical properties much like those of ethylene glycol, its physiological properties are quite different. Propylene glycol is essentially nontoxic, and it can be used as a solvent for drugs and as a moisturizing agent for foods.

Like other alcohols, propylene glycol is oxidized by liver enzymes.



Propylene glycol Pyruvate ion

In this case, however, the product is pyruvate ion, a normal intermediate in carbohydrate metabolism. Glycerol, a product of fat metabolism, is essentially nontoxic.

Summary

Glycols are alcohols with two OH groups on adjacent carbon atoms. Glycerol is the most important trihydroxy alcohol.

12.3: Glycols and Glycerol is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.6: Glycols and Glycerol** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.4: Ethers

Learning Objectives

- Describe the structural difference between an alcohol and an ether that affects physical characteristics and reactivity of each.
- Name simple ethers.
- Describe the structure and uses of some ethers.

With the general formula ROR′, an ether may be considered a derivative of water in which both hydrogen atoms are replaced by alkyl or aryl groups. It may also be considered a derivative of an alcohol (ROH) in which the hydrogen atom of the OH group is been replaced by a second alkyl or aryl group:

$$\text{HOH} \xrightarrow[\text{H atoms}]{\text{replace both}} \text{ROR}' \xleftarrow[\text{of OH group}]{\text{replace H atom}} \text{ROH}$$

Simple ethers have simple common names, formed from the names of the groups attached to oxygen atom, followed by the generic name *ether*. For example, CH_3–O–$CH_2CH_2CH_3$ is methyl propyl ether. If both groups are the same, the group name should be preceded by the prefix *di-*, as in dimethyl ether (CH_3–O–CH_3) and diethyl ether CH_3CH_2–O–CH_2CH_3.

Ether molecules have no hydrogen atom on the oxygen atom (that is, no OH group). Therefore there is no intermolecular hydrogen bonding between ether molecules, and ethers therefore have quite low boiling points for a given molar mass. Indeed, ethers have boiling points about the same as those of alkanes of comparable molar mass and much lower than those of the corresponding alcohols (Table **12.4.1**).

Table **12.4.1**: Comparison of Boiling Points of Alkanes, Alcohols, and Ethers


Condensed Structural Formula	Name	Molar Mass	Boiling Point (°C)	Intermolecular Hydrogen Bonding in Pure Liquid?
$CH_3CH_2CH_3$	propane	44	–42	no
CH_3OCH_3	dimethyl ether	46	–25	no
CH_3CH_2OH	ethyl alcohol	46	78	yes
$CH_3CH_2CH_2CH_2CH_3$	pentane	72	36	no
$CH_3CH_2OCH_2CH_3$	diethyl ether	74	35	no
$CH_3CH_2CH_2CH_2OH$	butyl alcohol	74	117	yes

Ether molecules do have an oxygen atom, however, and engage in hydrogen bonding with water molecules. Consequently, an ether has about the same solubility in water as the alcohol that is isomeric with it. For example, dimethyl ether and ethanol (both having the molecular formula C_2H_6O) are completely soluble in water, whereas diethyl ether and 1-butanol (both $C_4H_{10}O$) are barely soluble in water (8 g/100 mL of water).

Example 12.4.1

What is the common name for each ether?

a. $CH_3CH_2CH_2OCH_2CH_2CH_3$

b. 

Solution


a. The carbon groups on either side of the oxygen atom are propyl ($CH_3CH_2CH_2$) groups, so the compound is dipropyl ether.

b. The three-carbon group is attached by the middle carbon atom, so it is an isopropyl group. The one-carbon group is a methyl group. The compound is isopropyl methyl ether.

? Exercise 12.4.1


What is the common name for each ether?

a. $CH_3CH_2CH_2CH_2OCH_2CH_2CH_2CH_3$

b. 


To Your Health: Ethers as General Anesthetics

A *general anesthetic* acts on the brain to produce unconsciousness and a general insensitivity to feeling or pain. Diethyl ether ($CH_3CH_2OCH_2CH_3$) was the first general anesthetic to be used.



William Morton, a Boston dentist, introduced diethyl ether into surgical practice in 1846. This painting shows an operation in Boston in 1846 in which diethyl ether was used as an anesthetic. Inhalation of ether vapor produces unconsciousness by depressing the activity of the central nervous system. Source: Painting of William Morton by Ernest Board.

Diethyl ether is relatively safe because there is a fairly wide gap between the dose that produces an effective level of anesthesia and the lethal dose. However, because it is highly flammable and has the added disadvantage of causing nausea, it has been replaced by newer inhalant anesthetics, including the fluorine-containing compounds halothane, enflurane, and isoflurane. Unfortunately, the safety of these compounds for operating room personnel has been questioned. For example, female operating room workers exposed to halothane suffer a higher rate of miscarriages than women in the general population.



These three modern, inhalant, halogen-containing, anesthetic compounds are less flammable than diethyl ether.

Summary

To give ethers common names, simply name the groups attached to the oxygen atom, followed by the generic name *ether*. If both groups are the same, the group name should be preceded by the prefix *di-*. Ether molecules have no OH group and thus no intermolecular hydrogen bonding. Ethers therefore have quite low boiling points for a given molar mass. Ether molecules have an oxygen atom and can engage in hydrogen bonding with water molecules. An ether molecule has about the same solubility in water as the alcohol that is isomeric with it.

12.4: Ethers is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.8: Ethers** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.5: Aldehydes and Ketones- Structure and Names

Learning Objectives

- Identify the general structure for an aldehyde and a ketone.
- Use common names to name aldehydes and ketones.
- Use the IUPAC system to name aldehydes and ketones.


The next functional group we consider, the carbonyl group, has a carbon-to-oxygen double bond.




Carbonyl groups define two related families of organic compounds: the aldehydes and the ketones.

> *The carbonyl group is ubiquitous in biological compounds. It is found in carbohydrates, fats, proteins, nucleic acids, hormones, and vitamins—organic compounds critical to living systems.*

In a ketone, two carbon groups are attached to the carbonyl carbon atom. The following general formulas, in which R represents an alkyl group and Ar stands for an aryl group, represent ketones.



In an aldehyde, at least one of the attached groups must be a hydrogen atom. The following compounds are aldehydes:



In condensed formulas, we use CHO to identify an aldehyde rather than COH, which might be confused with an alcohol. This follows the general rule that in condensed structural formulas H comes after the atom it is attached to (usually C, N, or O).



An aldehyde




A ketone

The carbon-to-oxygen double bond is not shown but understood to be present. Because they contain the same functional group, aldehydes and ketones share many common properties, but they still differ enough to warrant their classification into two families.

Naming Aldehydes and Ketones

Both common and International Union of Pure and Applied Chemistry (IUPAC) names are frequently used for aldehydes and ketones, with common names predominating for the lower homologs. The common names of aldehydes are taken from the names of the acids into which the aldehydes can be converted by *oxidation*.




The stems for the common names of the first four aldehydes are as follows:

- 1 carbon atom: *form-*
- 2 carbon atoms: *acet-*
- 3 carbon atoms: *propion-*
- 4 carbon atoms: *butyr-*

Because the carbonyl group in a ketone must be attached to two carbon groups, the simplest ketone has three carbon atoms. It is widely known as *acetone*, a unique name unrelated to other common names for ketones.





Generally, the common names of ketones consist of the names of the groups attached to the carbonyl group, followed by the word *ketone*. (Note the similarity to the naming of ethers.) Another name for acetone, then, is *dimethyl ketone*. The ketone with four carbon atoms is ethyl methyl ketone.




Example 12.5.1

Classify each compound as an aldehyde or a ketone. Give the common name for each ketone.

a. 

b. 

c. 

Solution




a. This compound has the carbonyl group on an end carbon atom, so it is an aldehyde.

b. This compound has the carbonyl group on an interior carbon atom, so it is a ketone. Both alkyl groups are propyl groups. The name is therefore dipropyl ketone.

c. This compound has the carbonyl group between two alkyl groups, so it is a ketone. One alkyl group has three carbon atoms and is attached by the middle carbon atom; it is an isopropyl group. A group with one carbon atom is a methyl group. The name is therefore isopropyl methyl ketone.

Exercise 12.5.1

Classify each compound as an aldehyde or a ketone. Give the common name for each ketone.


1. 
2. 
3. 


Here are some simple IUPAC rules for naming aldehydes and ketones:


- The stem names of aldehydes and ketones are derived from those of the parent alkanes, defined by the longest continuous chain (LCC) of carbon atoms that contains the functional group.
- For an aldehyde, drop the *-e* from the alkane name and add the ending *-al*. Methanal is the IUPAC name for formaldehyde, and ethanal is the name for acetaldehyde.
- For a ketone, drop the *-e* from the alkane name and add the ending *-one*. Propanone is the IUPAC name for acetone, and butanone is the name for ethyl methyl ketone.
- To indicate the position of a substituent on an aldehyde, the carbonyl carbon atom is always considered to be C1; it is unnecessary to designate this group by number.
- To indicate the position of a substituent on a ketone, number the chain in the manner that gives the carbonyl carbon atom the lowest possible number. In cyclic ketones, it is understood that the carbonyl carbon atom is C1.

Example 12.5.2

Give the IUPAC name for each compound.

a. 

b. 

c. 

Solution


a. There are five carbon atoms in the LCC. The methyl group (CH_3) is a substituent on the second carbon atom of the chain; the aldehyde carbon atom is always C1. The name is derived from pentane. Dropping the *-e* and adding the ending *-al* gives pentanal. The methyl group on the second carbon atom makes the name 2-methylpentanal.


b. There are five carbon atoms in the LCC. The carbonyl carbon atom is C3, and there are methyl groups on C2 and C4. The IUPAC name is 2,4-dimethyl-3-pentanone.


c. There are six carbon atoms in the ring. The compound is cyclohexanone. No number is needed to indicate the position of the carbonyl group because all six carbon atoms are equivalent.

Exercise

Give the IUPAC name for each compound.

a. 

b. 

c. 


Example 12.5.3

Draw the structure for each compound.


a. 7-chlorooctanal
b. 4-methyl–3-hexanone

Solution

a. The *octan-* part of the name tells us that the LCC has eight carbon atoms. There is a chlorine (Cl) atom on the seventh carbon atom; numbering from the carbonyl group and counting the carbonyl carbon atom as C1, we place the Cl atom on the seventh carbon atom.



b. The *hexan-* part of the name tells us that the LCC has six carbon atoms. The 3 means that the carbonyl carbon atom is C3 in this chain, and the 4 tells us that there is a methyl (CH_3) group at C4:



Exercise 12.5.3

Draw the structure for each compound.

a. 5-bromo-3-iodoheptanal
b. 5-bromo-4-ethyl-2-heptanone

Summary

The common names of aldehydes are taken from the names of the corresponding carboxylic acids: formaldehyde, acetaldehyde, and so on. The common names of ketones, like those of ethers, consist of the names of the groups attached to the carbonyl group, followed by the word *ketone*. Stem names of aldehydes and ketones are derived from those of the parent alkanes, using an *-al* ending for an aldehydes and an *-one* ending for a ketone.

12.5: Aldehydes and Ketones- Structure and Names is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.9: Aldehydes and Ketones- Structure and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.6: Organic Sulfur Compounds

Learning Objectives

- Identify thiols (mercaptans) by the presence of an SH group.
- The mild oxidation of thiols gives disulfides.

Because sulfur is in the same group (6A) of the periodic table as oxygen, the two elements have some similar properties. We might expect sulfur to form organic compounds related to those of oxygen, and indeed it does. Thiols (also called mercaptans), which are sulfur analogs of alcohols, have the general formula RSH. Methanethiol (also called methyl mercaptan), has the formula CH_3SH. Ethanethiol (ethyl mercaptan) is the most common odorant for liquid propane (LP) gas.

The mild oxidation of thiols gives compounds called **disulfides**.

$$2RSH \xrightarrow{[O]} RSSR$$

The amino acids cysteine [$HSCH_2CH(NH_2)COOH$] and methionine [$CH_3SCH_2CH_2CH(NH_2)COOH$] contain sulfur atoms, as do all proteins that contain these amino acids. Disulfide linkages (–S–S–) between protein chains are extremely important in protein structure.

Thioethers, which are sulfur analogs of ethers, have the form general formula RSR′. An example is dimethylsulfide (CH_3SCH_3), which is responsible for the sometimes unpleasant odor of cooking cabbage and related vegetables. Note that methionine has a thioether functional group.

Career Focus: Paramedic

Paramedics are highly trained experts at providing emergency medical treatment. Their critical duties often include rescue work and emergency medical procedures in a wide variety of settings, sometimes under extremely harsh and difficult conditions. Like other science-based professions, their work requires knowledge, ingenuity, and complex thinking, as well as a great deal of technical skill. The recommended courses for preparation in this field include anatomy, physiology, medical terminology, and—not surprisingly—chemistry. An understanding of basic principles of organic chemistry, for example, is useful when paramedics have to deal with such traumas as burns from fuel (hydrocarbons) or solvent (alcohols, ethers, esters, and so on) fires and alcohol and drug overdoses.

To become a paramedic requires 2–4 y of training and usually includes a stint as an emergency medical technician (EMT). An EMT provides basic care, can administer certain medications and treatments, such as oxygen for respiratory problems and epinephrine (adrenalin) for allergic reactions, and has some knowledge of common medical conditions. A paramedic, in contrast, must have extensive knowledge of common medical problems and be trained to administer a wide variety of emergency drugs.

Paramedics usually work under the direction of a medical doctor with a title such as “medical director.” Some paramedics are employed by fire departments and may work from a fire engine that carries medical equipment as well as fire-fighting gear. Some work from hospital-sponsored ambulances and continue to care for their patients after reaching the hospital emergency room. Still other paramedics work for a government department responsible for emergency health care in a specific geographical area. Finally, some work for private companies that contract to provide service for a government body.

An experienced paramedic has a broad range of employment options, including training for mountain or ocean rescue, working with police department special weapons and tactics (SWAT) teams, or working in isolated settings such as on oil rigs. With their expertise at treating and stabilizing patients before quickly moving them to a hospital, paramedics often provide the first critical steps in saving an endangered life. The following quotation, inscribed on the Arlington National Cemetery headstone of Army Lieutenant R. Adams Cowley, who is often called the “father” of shock trauma medicine, serves as the motto for many paramedic units: “Next to creating a life the finest thing a man can do is save one.” —Abraham Lincoln

Summary

Thiols, thioethers, and disulfides are common in biological compounds.

12.6: Organic Sulfur Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.11: Organic Sulfur Compounds** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.7: Carboxylic Acids- Structures and Names

Learning Objectives


- Name carboxylic acids with common names.
- Name carboxylic acids according to IUPAC nomenclature.

Carboxylic acids occur widely in nature, often combined with alcohols or other functional groups, as in fats, oils, and waxes. They are components of many foods, medicines, and household products (Figure **12.7.1**). Not surprisingly, many of them are best known by common names based on Latin and Greek words that describe their source.




Figure **12.7.1**: Carboxylic Acids in the Home. Carboxylic acids occur in many common household items. (a) Vinegar contains acetic acid, (b) aspirin is acetylsalicylic acid, (c) vitamin C is ascorbic acid, (d) lemons contain citric acid, and (e) spinach contains oxalic acid. © Thinkstock

The simplest carboxylic acid, formic acid (HCOOH), was first obtained by the distillation of ants (Latin *formica*, meaning “ant”). The bites of some ants inject formic acid, and the stings of wasps and bees contain formic acid (as well as other poisonous materials).



Formic acid

The next higher homolog is acetic acid, which is made by fermenting cider and honey in the presence of oxygen. This fermentation produces vinegar, a solution containing 4%–10% acetic acid, plus a number of other compounds that add to its flavor. Acetic acid is probably the most familiar weak acid used in educational and industrial chemistry laboratories.



Acetic acid

Pure acetic acid solidifies at 16.6°C, only slightly below normal room temperature. In the poorly heated laboratories of the late 19th and early 20th centuries in northern North America and Europe, acetic acid often "froze" on the storage shelf. For that reason, pure acetic acid (sometimes called concentrated acetic acid) came to be known as *glacial acetic acid*, a name that survives to this day.

The third homolog, propionic acid (CH_3CH_2COOH), is seldom encountered in everyday life. The fourth homolog, butyric acid ($CH_3CH_2CH_2COOH$), is one of the most foul-smelling substances imaginable. It is found in rancid butter and is one of the ingredients of body odor. By recognizing extremely small amounts of this and other chemicals, bloodhounds are able to track fugitives. Models of the first four carboxylic acids are shown in Figure **12.7.2**.






Figure **12.7.2**: Ball-and-Stick Models of Carboxylic Acids. Carboxylic acids feature a carbon atom doubly bonded to an oxygen atom and also joined to an OH group. The four acids illustrated here are formic acid (a), acetic acid (b), propionic acid (c), and butyric acid (d).

The acid with the carboxyl group attached directly to a benzene ring is called benzoic acid (C_6H_5COOH).



Benzoic acid

The common names of carboxylic acids use Greek letters (α, β, γ, δ, and so forth), not numbers, to designate the position of substituent groups in acids. These letters refer to the position of the carbon atom in relation to the carboxyl carbon atom.




In the nomenclature system of the International Union of Pure and Applied Chemistry (IUPAC), the parent hydrocarbon is the one that corresponds to the longest continuous chain (LCC) containing the carboxyl group. The *-e* ending of the parent alkane is replaced by the suffix *-oic* and the word *acid*. For example, the carboxylic acid derived from pentane is pentanoic acid ($CH_3CH_2CH_2CH_2COOH$). As with aldehydes, the carboxyl carbon atom is counted first; numbers are used to indicate any substituted carbon atoms in the parent chain.

Greek letters are used with common names; numbers are used with IUPAC names.


Example 12.7.1

Give the common and IUPAC names for each compound.

1. $ClCH_2CH_2CH_2COOH$
2. 


Solution

1. The LCC contains four carbon atoms; the compound is therefore named as a substituted butyric (or butanoic) acid.



The chlorine atom is attached to the γ-carbon in the common system or C4 in the IUPAC system. The compound is γ-chlorobutyric acid or 4-chlorobutanoic acid.

2. The LCC contains four carbon atoms; the compound is therefore named as a substituted butyric (or butanoic) acid.



The bromine (Br) atom is at the α-carbon in the common system or C2 in the IUPAC system. The compound is α-bromobutyric acid or 2-bromobutanoic acid.

Exercise 12.7.1

Give the IUPAC name for each compound.

a. $ClCH_2CH_2CH_2CH_2COOH$
b. $(CH_3)_2CHCH_2CHBrCOOH$

Example 12.7.2

Write the condensed structural formula for β-chloropropionic acid.

Solution

Propionic acid has three carbon atoms: C–C–COOH. Attach a chlorine (Cl) atom to the parent chain at the beta carbon atom, the second one from the carboxyl group: Cl–C–C–COOH. Then add enough hydrogen atoms to give each carbon atom four bonds: $ClCH_2CH_2COOH$.

Exercise 12.7.2

Write the condensed structural formula for 4-bromo-5-methylhexanoic acid.

Key Takeaways

- Simple carboxylic acids are best known by common names based on Latin and Greek words that describe their source (e.g., formic acid, Latin *formica*, meaning “ant”).
- Greek letters, not numbers, designate the position of substituted acids in the common naming convention.
- IUPAC names are derived from the LCC of the parent hydrocarbon with the *-e* ending of the parent alkane replaced by the suffix *-oic* and the word *acid*.

12.7: Carboxylic Acids- Structures and Names is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.1: Carboxylic Acids - Structures and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.8: Esters- Structures and Names

Learning Objectives

- Identify the general structure for an ester.
- Use common names to name esters.
- Name esters according to the IUPAC system.

Esters have the general formula RCOOR′, where R may be a hydrogen atom, an alkyl group, or an aryl group, and R′ may be an alkyl group or an aryl group but *not* a hydrogen atom. (If it were hydrogen atom, the compound would be a carboxylic acid.) Figure 12.8.1 shows models for two common esters.




Figure 12.8.1: The Structure of Esters. Esters feature a carbon-to-oxygen double bond that is also singly bonded to a second oxygen atom, which is then joined to an alkyl or an aryl group. The esters shown here are ethyl acetate (a) and methyl butyrate (b).


Esters occur widely in nature. Unlike carboxylic acids, esters generally have pleasant odors and are often responsible for the characteristic fragrances of fruits and flowers. Once a flower or fruit has been chemically analyzed, flavor chemists can attempt to duplicate the natural odor or taste. Both natural and synthetic esters are used in perfumes and as flavoring agents.

> Fats and vegetable oils are esters of long-chain fatty acids and glycerol. Esters of phosphoric acid are of the utmost importance to life.

Names of Esters


Although esters are covalent compounds and salts are ionic, esters are named in a manner similar to that used for naming salts. The group name of the alkyl or aryl portion is given first and is followed by the name of the acid portion. In both common and International Union of Pure and Applied Chemistry (IUPAC) nomenclature, the *-ic* ending of the parent acid is replaced by the suffix *-ate* (Table 12.8.1).

Table 12.8.1: Nomenclature of Esters

Condensed Structural Formula	Common Name	IUPAC Name
$HCOOCH_3$	methyl formate	methyl methanoate
CH_3COOCH_3	methyl acetate	methyl ethanoate
$CH_3COOCH_2CH_3$	ethyl acetate	ethyl ethanoate
$CH_3CH_2COOCH_2CH_3$	ethyl propionate	ethyl propanoate
$CH_3CH_2CH_2COOCH(CH_3)_2$	isopropyl butyrate	isopropyl butanoate
	ethyl benzoate	ethyl benzoate

Example 12.8.1

Give the common and IUPAC names for each compound.

1. 
2. 


Solution

The alkyl group attached directly to the oxygen atom is a butyl group (in green).




The part of the molecule derived from the carboxylic acid (in red) has three carbon atoms. It is called propionate (common) or propanoate (IUPAC). The ester is therefore butyl propionate or butyl propanoate.

1. An alkyl group (in green) is attached directly to the oxygen atom by its middle carbon atom; it is an isopropyl group. The part derived from the acid (that is, the benzene ring and the carbonyl group, in red) is benzoate. The ester is therefore isopropyl benzoate (both the common name and the IUPAC name).



Exercise 12.8.1

Give the common and IUPAC names for each compound.


1. 

Example 12.8.2


Draw the structure for ethyl pentanoate.

Solution

Start with the portion from the acid. Draw the pentanoate (five carbon atoms) group first; keeping in mind that the last carbon atom is a part of the carboxyl group.



Then attach the ethyl group to the bond that ordinarily holds the hydrogen atom in the carboxyl group.



Exercise 12.8.2

Draw the structure for phenyl pentanoate.

Key Takeaway

- An ester has an OR group attached to the carbon atom of a carbonyl group.

12.8: Esters- Structures and Names is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **15.5: Esters - Structures and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.9: Phosphate Esters


Learning Objectives

- Describe phosphate esters.
- Understand why phosphate esters are important in living cells.


Just as carboxylic acids do, inorganic acids such as nitric acid (HNO_3), sulfuric acid (H_2SO_4), and phosphoric acid (H_3PO_4) also form esters. The esters of phosphoric acid are especially important in biochemistry. A phosphoric acid molecule can form a monoalkyl, a dialkyl, or a trialkyl ester by reaction with one, two, or three molecules of an alcohol.



Ethyl dihydrogen phosphate




Diethyl hydrogen phosphate




Triethyl phosphate


Esters of pyrophosphoric acid and triphosphoric acid are also important in biochemistry.



Phosphoric acid



Pyrophosphoric acid




Triphosphoric acid

Esters of these acids are present in every plant and animal cell. They are biochemical intermediates in the transformation of food into usable energy. The bonds between phosphate units in adenosine triphosphate (ATP) are called **phosphoanhydride bonds**. These are high-energy bonds that store energy from the metabolism of foods. Hydrolysis of ATP releases energy as it is needed for biochemical processes (for instance, for muscle contraction). Phosphate esters are also important structural constituents of phospholipids and nucleic acids.

Nitroglycerin

The explosive nitroglycerin (glyceryl trinitrate) is an ester formed from glycerol and nitric acid. It is used in medicine to relieve chest pain in heart disease.



Glycerol

Glycerol trinitrate (nitroglycerin)

12.9.1 Exercise

What compounds combine to form phosphate esters?

Answer

phosphoric acids and alcohols

Summary

Inorganic acids such as H_3PO_4 form esters. The esters of phosphoric acid are especially important in biochemistry.

12.9: Phosphate Esters is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **15.9: Esters of Phosphoric Acid** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.10: Amines


Learning Objectives

- Describe phosphate esters.
- Understand why phosphate esters are important in living cells.


Just as carboxylic acids do, inorganic acids such as nitric acid (HNO_3), sulfuric acid (H_2SO_4), and phosphoric acid (H_3PO_4) also form esters. The esters of phosphoric acid are especially important in biochemistry. A phosphoric acid molecule can form a monoalkyl, a dialkyl, or a trialkyl ester by reaction with one, two, or three molecules of an alcohol.



Ethyl dihydrogen phosphate




Diethyl hydrogen phosphate




Triethyl phosphate


Esters of pyrophosphoric acid and triphosphoric acid are also important in biochemistry.



Phosphoric acid



Pyrophosphoric acid




Triphosphoric acid

Esters of these acids are present in every plant and animal cell. They are biochemical intermediates in the transformation of food into usable energy. The bonds between phosphate units in adenosine triphosphate (ATP) are called **phosphoanhydride bonds**. These are high-energy bonds that store energy from the metabolism of foods. Hydrolysis of ATP releases energy as it is needed for biochemical processes (for instance, for muscle contraction). Phosphate esters are also important structural constituents of phospholipids and nucleic acids.

Nitroglycerin

The explosive nitroglycerin (glyceryl trinitrate) is an ester formed from glycerol and nitric acid. It is used in medicine to relieve chest pain in heart disease.



Glycerol

Glycerol trinitrate (nitroglycerin)

12.10.1 Exercise

What compounds combine to form phosphate esters?

Answer

phosphoric acids and alcohols

Summary

Inorganic acids such as H_3PO_4 form esters. The esters of phosphoric acid are especially important in biochemistry.

12.10: Amines is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **15.9: Esters of Phosphoric Acid** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

12.11: Amides

Learning Objectives


- Describe phosphate esters.
- Understand why phosphate esters are important in living cells.

Just as carboxylic acids do, inorganic acids such as nitric acid (HNO_3), sulfuric acid (H_2SO_4), and phosphoric acid (H_3PO_4) also form esters. The esters of phosphoric acid are especially important in biochemistry. A phosphoric acid molecule can form a monoalkyl, a dialkyl, or a trialkyl ester by reaction with one, two, or three molecules of an alcohol.



Ethyl dihydrogen phosphate | Diethyl hydrogen phosphate | Triethyl phosphate

Esters of pyrophosphoric acid and triphosphoric acid are also important in biochemistry.




Phosphoric acid | Pyrophosphoric acid | Triphosphoric acid

Esters of these acids are present in every plant and animal cell. They are biochemical intermediates in the transformation of food into usable energy. The bonds between phosphate units in adenosine triphosphate (ATP) are called **phosphoanhydride bonds**. These are high-energy bonds that store energy from the metabolism of foods. Hydrolysis of ATP releases energy as it is needed for biochemical processes (for instance, for muscle contraction). Phosphate esters are also important structural constituents of phospholipids and nucleic acids.

Nitroglycerin

The explosive nitroglycerin (glyceryl trinitrate) is an ester formed from glycerol and nitric acid. It is used in medicine to relieve chest pain in heart disease.



Glycerol | Glycerol trinitrate (nitroglycerin)

12.11.1 Exercise

What compounds combine to form phosphate esters?

Answer

phosphoric acids and alcohols

Summary

Inorganic acids such as H_3PO_4 form esters. The esters of phosphoric acid are especially important in biochemistry.

12.11: Amides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.9: Esters of Phosphoric Acid** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

13: Organic Reactions

Organic reactions are chemical reactions involving organic compounds. The basic organic chemistry reaction types are addition reactions, elimination reactions, substitution reactions, pericyclic reactions, rearrangement reactions, photochemical reactions and redox reactions.

13.1: Organic Reactions

13.2: Organic Reactions (Exercises)

This page titled 13: Organic Reactions is shared under a CK-12 license and was authored, remixed, and/or curated by CK-12 Foundation via source content that was edited to the style and standards of the LibreTexts platform.

13.1: Organic Reactions

Learning Outcomes

- Identify and describe substitution, elimination, addition, hydrolysis, and condensation reactions.
- Predict products of each reaction type.
- Predict products of hydration reactions based on Markovnikov's rule.
- Explain why condensation reactions are also called dehydration reactions.
- Define "polymer".

Organic reactions require the breaking of strong covalent bonds, which takes a considerable input of energy. In order for relatively stable organic molecules to react at a reasonable rate, they often must be modified with the use of highly reactive materials or in the presence of a catalyst. In this lesson, you will learn about several general categories of organic reactions.

Substitution Reactions

A **substitution reaction**, which is the same as a single replacement reaction in inorganic reactions, *is a reaction in which one or more atoms in a molecule are replaced with another atom or group of atoms.* Alkyl halides are formed by the substitution of a halogen atom for a hydrogen atom. When methane reacts with chlorine gas, ultraviolet light can act as a catalyst for the reaction.

$$\mathrm{CH_4\,(g) + Cl_2\,(g) \overset{\text{UV light}}{\rightarrow} CH_3Cl\,(g) + HCl\,(g)} \tag{13.1.1}$$

The reaction produces chloromethane and hydrogen chloride. When the mixture is allowed to react for longer periods of time, further substitution reactions may occur.

$$\mathrm{CH_3Cl\,(g) + Cl_2\,(g) \overset{\text{UV light}}{\rightarrow} CH_2Cl_2\,(g) + HCl\,(g)} \tag{13.1.2}$$


The product above is dichloromethane. Further substitution products trichloromethane and tetrachloromethane, commonly called carbon tetrachloride. A mixture of products occurs in the reaction, with the relative amounts dependent upon the time that the reaction is allowed to proceed. Chlorofluorocarbons are produced by reacting chloroalkanes with **HF**, because the fluorine atom makes a stronger bond to the carbon atom than chlorine does.

$$\mathrm{CCl_4\,(g) + HF\,(g) \overset{SbF_5}{\rightarrow} CCl_3F\,(g) + HCl\,(g)} \tag{13.1.3}$$

The fluorine atom substitutes for a chlorine atom in the reaction.

Elimination Reactions

An **elimination reaction** involves the removal of adjacent atoms from a molecule. This results in the formation of a multiple bond and the release of a small molecule, so they are called elimination reactions. They have the general form



A typical example is the conversion of ethyl chloride to ethylene:

$$\mathrm{CH_3CH_2Cl \rightarrow CH_2 = CH_2 + HCl} \tag{13.1.4}$$

Much of the approximately 26 million tons of ethylene produced per year in the United States is used to synthesize plastics, such as polyethylene. In the above reaction, the A–B molecule eliminated is HCl, whose components are eliminated as H^+ from the carbon atom on the left and Cl^- from the carbon on the right. When an acid is produced, as occurs here, the reaction is generally carried out in the presence of a base (such as NaOH) to neutralize the acid. Other elimination reactions will produce H_2, X_2 (where X = halogen), or H_2O. These reactions are often referred to by more descriptive terms such as dehydrogenation (removing hydrogen) or dechlorination (removing chlorine).

Addition Reactions

An **addition reaction** *is a reaction in which an atom or molecule is added to an unsaturated molecule, making a single product.* An addition reaction can often be thought of as adding a molecule across the double bond of an alkene or across the triple bond of an alkyne. Knowing that "ation" means to add, the specific names of these reactions, such as hydrogenation, hydration, or chlorination, should make sense. Note that hydrogenation (adding H_2) and hydration (adding H_2O) are very different processes.

One type of addition reaction is called hydrogenation. **Hydrogenation** *is a reaction that occurs when molecular hydrogen is added to an alkene to produce an alkane* or *hydrogen is added to an alkyne to produce an alkene or alkane.* The reaction is typically performed with the use of a transition metal catalyst. For example, ethene reacts with hydrogen to form ethane.

$$\mathrm{CH_2{=}CH_2\,(g) + H_2\,(g) \xrightarrow{Pt} CH_3CH_3\,(g)} \tag{13.1.5}$$

Note that the hydrogenation reaction is also a redox reaction. Ethene is reduced, because the oxidation numbers of the carbon atoms change from -2 to -3 as a result of the reaction.


Vegetable oils consist of long carbon chains with carboxyl groups on the end; these molecules are referred to as fatty acids. The carbon chains of the fatty acids in vegetable oils are unsaturated, usually containing multiple double bonds. When hydrogen gas is blown through a sample of the oil, hydrogen atoms add across the double bonds. This conversion changes the substance from a liquid oil into a solid fat. The "hydrogenated" on a food product is an indication that oil (liquid) has been converted into fat (solid) by this process. Margarine is manufactured from unsaturated vegetable oil in this way by hydrogenating some of the double bonds making it a "partially hydrogenated vegetable oil".

Addition reactions are also useful ways to introduce a new functional group into an organic molecule. Alkyl halides can be produced from an alkene by the addition of either the elemental halogen or the hydrogen halide. A monosubstituted alkyl halide can be produced by the addition of a hydrogen halide to an alkene. Shown below is the formation of chloroethane.

$$\mathrm{CH_2{=}CH_2\,(g) + HCl\,(g) \rightarrow CH_3CH_2Cl\,(g)} \tag{13.1.6}$$

When the reactant is the diatomic halogen, the product is a disubstituted alkyl halide as in the addition of bromine to ethene.

$$\mathrm{CH_2{=}CH_2\,(g) + Br_2\,(l) \rightarrow CH_2BrCH_2Br\,(g)} \tag{13.1.7}$$




1,2-dibromoethane

The addition of bromine to an unknown organic compound can be used as a test for unsaturation in the compound. Bromine has a distinctive brownish-orange color, while most bromoalkanes are colorless. When bromine is slowly added to a solution of the compound, the orange color will fade if it undergoes an addition reaction to produce an alkyl halide. If the orange color remains, then the original compound was already saturated, and no reaction occurred.


A **hydration reaction** *is a reaction in which water is added to an alkene*. Hydration reactions can take place when the alkene and water are heated to near $\mathbf{100°C}$ in the presence of a strong acid, which acts as a catalyst. Shown below is the hydration of ethene to produce ethanol.

$$\mathrm{CH_2{=}CH_2\,(g) + H_2O\,(l) \rightarrow CH_3CH_2OH\,(g)} \tag{13.1.8}$$

Unlike addition reactions involving $\mathbf{H_2}$ or $\mathbf{Br_2}$, the addition of water can have two possible products because an $\mathbf{-H}$ and an $\mathbf{-OH}$ are being added to the carbons in the double bond. **Markovnikov's rule** helps predict the main product in a hydration reaction. The rule states that the hydrogen atom from water will add to the carbon that originally had more hydrogen atoms. For example, look at the hydration of 1-butene.



Note that the first carbon in 1-butene started with two hydrogen atoms and the second carbon started with one hydrogen. Therefore, the hydrogen from water adds to the first carbon and the **–OH** group adds to the second carbon. When there are equal numbers of hydrogen atoms on both carbons in a double bond then the two products will form in approximately equal amounts. For example, the hydration of 2-pentene results in two products. In the first product, the **–OH** group is on the third carbon and in the second product, the **–OH** group is on the second carbon. While these two molecules will have similar properties, there will be differences.



Condensation Reactions

A **condensation reaction** *is a reaction in which two molecules combine to form a single molecule.* A small molecule, often water, is usually removed during a condensation reaction. Amino acids are important biological molecules that have an amine functional group on one end of the molecule and a carboxylic acid functional group on the other end. When two amino acids combine in a condensation reaction, a covalent bond forms between the amine nitrogen of one amino acid and the carboxyl carbon of the second amino acid. A molecule of water is then removed as a second product (see figure below).





Figure **13.1.1**: Amino acids join together to form a molecule called a dipeptide. The **–OH** from the carboxyl group of one amino acid combines with a hydrogen atom from the amine group of the other amino group to produce water (blue).

This reaction forms a molecule called a dipeptide, and the resulting carbon-nitrogen covalent bond is often called a peptide bond. When repeated numerous times, a long molecule called a protein is eventually produced.

An **esterification** *is a condensation reaction in which an ester is formed from an alcohol and a carboxylic acid.* Esterification is a subcategory of condensation reactions because a water molecule is produced in the reaction. The reaction can be catalyzed by a strong acid, usually sulfuric acid. When the carboxylic acid, butanoic acid, is heated with an excess of methanol and a few drops of sulfuric acid, the ester methyl butanoate is produced. Methyl butanoate has the scent of pineapples. The reaction is shown below with both molecular and structural formulas. Esterification reactions are reversible.

$$CH_3CH_2CH_2COOH + HOCH_3 \overset{H_2SO_4}{\rightarrow} CH_3CH_2CH_2COOCH_3 + H_2O \qquad (13.1.9)$$




Saponification *describes the alkaline hydrolysis reaction of an ester*. The term saponification originally described the hydrolysis of long-chain esters called fatty acid esters to produce soap molecules, which are the salts of fatty acids. One such soap molecule is sodium stearate, formed from the hydrolysis of ethyl stearate.

$$\underset{\text{ethyl stearate}}{C_{17}H_{35}COOC_2H_5} + NaOH \rightarrow \underset{\text{sodium stearate (soap)}}{C_{17}H_{35}COO^-Na^+} + C_2H_5OH \qquad (13.1.10)$$

Hydrolysis Reactions

Hydrolysis is the reverse of condensation. "Hydro" indicates that water is involved and "lysis" means to break apart. In a hydrolysis reaction, water is added and the molecule breaks apart, usually at a C–O–C linkage.



Polymerization

Polymers are very different than the other kinds of organic molecules that you have seen so far. Whereas other compounds are of relatively low molar mass, polymers are giant molecules of very high molar mass. Polymers are the primary components of all sorts of plastics and related compounds. A **polymer** *is a large molecule formed of many smaller molecules covalently bonded to one another in a repeating pattern*. The *small molecules that make up the polymer are called* **monomers**. Polymers are generally formed by either addition or condensation reactions. Teflon (see figure below) is a non-reactive, non-stick coating used on cookware as well as in containers and pipes for reactive or corrosive chemicals.




Figure **13.1.2**: Polytetrafluoroethylene (also known as Teflon) is formed from the polymerization of tetrafluoroethylene.

This page titled 13.1: Organic Reactions is shared under a CK-12 license and was authored, remixed, and/or curated by CK-12 Foundation via source content that was edited to the style and standards of the LibreTexts platform.

13.2: Organic Reactions (Exercises)

These are homework exercises to accompany Chapter 12 of the University of Kentucky's LibreText for CHE 103 - Chemistry for Allied Health. Answers are below the questions.

Questions

12.1: Organic Reactions

(click here for solutions)

Q12.1.1

Explain the difference between hydrolysis and condensation reactions.

Q12.1.2

What is being added in a hydrogenation reaction? Hydration? Bromination?

Q12.1.3

What is the product of the oxidation of a primary alcohol? A secondary alcohol?

Q12.1.4

Why is a condensation reaction also called a dehydration reaction?



Q12.1.5

Draw the structure for the primary product of each reaction..

a. $CH_3CH{=}CH_2 + Br_2 \rightarrow$
b. $CH_3CH_2CH{=}CH_2 + H_2 \rightarrow$
c. $CH_3CH{=}CHCH_3 + H_2O \rightarrow$
d. $CH_3CH_2CH{=}CH_2 + H_2O \rightarrow$
e. $CH_3CH_2CH{=}CHCH_3 + H_2O \rightarrow$



Q12.1.6

Draw the structure of the compound that would be formed from the single oxidation of each compound.

a.	
b.	
c.	




Q12.1.7

Draw the structure of the compound that would be formed from the complete oxidation of each compound.

a.	
b.	
c.	


Q12.1.8


Draw the structure of the compound that would be formed from a single reduction of each compound.


a.	
b.	
c.	


Q12.1.9

Draw the product(s) for the hydrolysis of the given reactant. The red bond in the reactant is the one that is broken.

a. 


b. 

c. 

d. 

Q12.1.10

Draw the product(s) for the condensation of the given reactants. The layout of the molecules indicates where the bond will form between them. If there are three molecules, two condensations will occur to produce one final product.



Q12.1.11

Compare the reactants and products in part d and e of 12.1.10. Are the reactants the same or different? Are the products the same or different? Why?

Answers

12.1: Organic Reactions

Q12.1.1

Hydrolysis is a molecule breaking apart after the addition of water. Condensation is two molecules coming together and producing H_2O.

Q12.1.2

hydrogenation = adding hydrogen (H_2)

hydration = adding water

bromination = adding bromine (Br_2)






Q12.1.3

A primary alcohol oxidizes to an aldehyde and then to a carboxylic acid. A secondary alcohol oxidizes to a ketone.



Q12.1.4

Because water is released during a condensation reaction.


Q12.1.5

a.	 Each bromine atom is added to one of the carbons in the double bond.
b.	$CH_3CH_2CH_2CH_3$ Each hydrogen atom is added to one of the carbons in the double bonds.
c.	 Water is added as H and OH. The carbons in the double bond each have the same number of hydrogen atoms so Markovnikov's rule does not apply. The molecule is symmetrical so adding the OH group to either carbon results in the same molecule.
d.	 Water is added as H and OH. The carbon atoms in the double bond have different number of hydrogen atoms so Markovnikov's rule does apply. The H is added to the carbon with more hydrogen atoms and the OH is added to the carbon with fewer hydrogen atoms.
e.	 and  Water is added as H and OH. The carbon atoms in the double bond have the same number of hydrogen atoms on each so Markovnikov's rule does not apply. There are two possible products because adding the OH to the carbon on the left side of the double bond results in a different molecule than adding the OH to the carbon on the right side of the double bond. This differs from c because the molecule is not symmetrical.




Q12.1.6

a.	
b.	
c.	

Q12.1.7







a.	
b.	
c.	

Q12.1.8





a.	
b.	
c.	

https://chem.libretexts.org/@go/page/387613

Q12.1.9

a.	 
b.	 
c.	 
d.	 

Q12.1.10

a.	
b.	
c.	
d.	
e.	

Q12.1.11

The reactants are the same, but the products are different because they are combined in a different order.

This page titled 13.2: Organic Reactions (Exercises) is shared under a CK-12 license and was authored, remixed, and/or curated by CK-12 Foundation via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

14: Properties of Organic Functional Groups

14.1: Physical Properties of Alcohols
14.2: Physical Properties of Aldehydes and Ketones
14.3: Physical Properties of Carboxylic Acids
14.4: Physical Properties of Esters
14.5: Physical Properties of Amines
14.6: Physical Properties of Amides
14.7: Ionization of Carboxylic Acids
14.8: Amines as Bases
14.9: Chirality
14.10: End-of-Chapter Material

14: Properties of Organic Functional Groups is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

14.1: Physical Properties of Alcohols

Learning Objectives

- Explain why the boiling points of alcohols are higher than those of ethers and alkanes of similar molar masses.
- Explain why alcohols and ethers of four or fewer carbon atoms are soluble in water while comparable alkanes are not soluble.

Alcohols can be considered derivatives of water (H_2O; also written as HOH).



Like the H–O–H bond in water, the R–O–H bond is bent, and alcohol molecules are polar. This relationship is particularly apparent in small molecules and reflected in the physical and chemical properties of alcohols with low molar mass. Replacing a hydrogen atom from an alkane with an OH group allows the molecules to associate through hydrogen bonding (Figure **14.1.1**).




Figure **14.1.1**: Intermolecular Hydrogen Bonding in Methanol. The OH groups of alcohol molecules make hydrogen bonding possible.

Recall that physical properties are determined to a large extent by the type of intermolecular forces. Table **14.1.1** lists the molar masses and the boiling points of some common compounds. The table shows that substances with similar molar masses can have quite different boiling points.

Table **14.1.1**: Comparison of Boiling Points and Molar Masses

Formula	Name	Molar Mass	Boiling Point (°C)
CH_4	methane	16	–164
HOH	water	18	100
C_2H_6	ethane	30	–89
CH_3OH	methanol	32	65
C_3H_8	propane	44	–42
CH_3CH_2OH	ethanol	46	78
C_4H_{10}	butane	58	–1
$CH_3CH_2CH_2OH$	1-propanol	60	97

Alkanes are nonpolar and are thus associated only through relatively weak dispersion forces. Alkanes with one to four carbon atoms are gases at room temperature. In contrast, even methanol (with one carbon atom) is a liquid at room temperature. Hydrogen bonding greatly increases the boiling points of alcohols compared to hydrocarbons of comparable molar mass. The boiling point is a rough measure of the amount of energy necessary to separate a liquid molecule from its nearest neighbors. If the molecules interact through hydrogen bonding, a relatively large quantity of energy must be supplied to break those intermolecular attractions. Only then can the molecule escape from the liquid into the gaseous state.

Alcohols can also engage in hydrogen bonding with water molecules (Figure **14.1.2**). Thus, whereas the hydrocarbons are insoluble in water, alcohols with one to three carbon atoms are completely soluble. As the length of the chain increases, however, the solubility of alcohols in water decreases; the molecules become more like hydrocarbons and less like water. The alcohol 1-decanol ($CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_2OH$) is essentially insoluble in water. We frequently find that the borderline of solubility in a family of organic compounds occurs at four or five carbon atoms.




Figure **14.1.2**: Hydrogen Bonding between Methanol Molecules and Water Molecules. Hydrogen bonding between the OH of methanol and water molecules accounts for the solubility of methanol in water.

Summary

Alcohols have higher boiling points than do ethers and alkanes of similar molar masses because the OH group allows alcohol molecules to engage in hydrogen bonding. Alcohols of four or fewer carbon atoms are soluble in water because the alcohol molecules engage in hydrogen bonding with water molecules; comparable alkane molecules cannot engage in hydrogen bonding.

14.1: Physical Properties of Alcohols is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **14.3: Physical Properties of Alcohols** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.2: Physical Properties of Aldehydes and Ketones

Learning Objectives

- Explain why the boiling points of aldehydes and ketones are higher than those of ethers and alkanes of similar molar masses but lower than those of comparable alcohols.
- Compare the solubilities in water of aldehydes and ketones of four or fewer carbon atoms with the solubilities of comparable alkanes and alcohols.
- Name the typical reactions take place with aldehydes and ketones.
- Describe some of the uses of common aldehydes and ketones.

The carbon-to-oxygen double bond is quite polar, more polar than a carbon-to-oxygen single bond. The electronegative oxygen atom has a much greater attraction for the bonding electron pairs than does the carbon atom. The carbon atom has a partial positive charge, and the oxygen atom has a partial negative charge:



In aldehydes and ketones, this charge separation leads to dipole-dipole interactions that are great enough to significantly affect the boiling points. Table **14.2.1** shows that the polar single bonds in ethers have little such effect, whereas hydrogen bonding between alcohol molecules is even stronger.


Table **14.2.1**: Boiling Points of Compounds Having Similar Molar Masses but Different Types of Intermolecular Forces

Compound	Family	Molar Mass	Type of Intermolecular Forces	Boiling Point (°C)
$CH_3CH_2CH_2CH_3$	alkane	58	dispersion only	–1
$CH_3OCH_2CH_3$	ether	60	weak dipole	6
CH_3CH_2CHO	aldehyde	58	strong dipole	49
$CH_3CH_2CH_2OH$	alcohol	60	hydrogen bonding	97

Formaldehyde is a gas at room temperature. Acetaldehyde boils at 20°C; in an open vessel, it boils away in a warm room. Most other common aldehydes are liquids at room temperature.

Although the lower members of the homologous series have pungent odors, many higher aldehydes have pleasant odors and are used in perfumes and artificial flavorings. As for the ketones, acetone has a pleasant odor, but most of the higher homologs have rather bland odors.


The oxygen atom of the carbonyl group engages in hydrogen bonding with a water molecule.



The solubility of aldehydes is therefore about the same as that of alcohols and ethers. Formaldehyde, acetaldehyde, and acetone are soluble in water. As the carbon chain increases in length, solubility in water decreases. The borderline of solubility occurs at about four carbon atoms per oxygen atom. All aldehydes and ketones are soluble in organic solvents and, in general, are less dense than water.

Oxidation of Aldehydes and Ketones

Aldehydes and ketones are much alike in many of their reactions, owing to the presence of the carbonyl functional group in both. They differ greatly, however, in one most important type of reaction: oxidation. Aldehydes are readily oxidized to carboxylic acids, whereas ketones resist oxidation.



The aldehydes are, in fact, among the most easily oxidized of organic compounds. They are oxidized by oxygen (O_2) in air to carboxylic acids.

$$2\,RCHO + O_2 \longrightarrow 2\,RCOOH$$

The ease of oxidation helps chemists identify aldehydes. A sufficiently mild oxidizing agent can distinguish aldehydes not only from ketones but also from alcohols. Tollens' reagent, for example, is an alkaline solution of silver (Ag^+) ion complexed with ammonia (NH_3), which keeps the Ag^+ ion in solution.

$$H_3N{-}Ag^+{-}NH_3$$

When Tollens' reagent oxidizes an aldehyde, the Ag^+ ion is reduced to free silver (Ag).

$$\underbrace{RCHO\,(aq)}_{\text{an aldehyde}} + 2\,Ag(NH_3)_2^+\,(aq) \longrightarrow RCOO^- + \underbrace{2\,Ag\,(s)}_{\text{free silver}} + 4\,NH_3\,(aq) + 2\,H_2O$$

Deposited on a clean glass surface, the silver produces a mirror (Figure 14.2.1). Ordinary ketones do not react with Tollens' reagent.



Figure 14.2.1: Aldehyde Reactions. A reaction related to the Tollens' reaction is often used to silver mirrors. These ornaments were silvered by such a reaction. Glucose, a simple sugar with an aldehyde functional group, is used as the reducing agent. Source: Photo courtesy of Krebs Glas Lauscha, commons.wikimedia.org/wiki/File:Silvering.jpg.

A person wearing gloves is holding a bunch of ornaments that she has just lifted from a basin containing a solution.

Although ketones resist oxidation by ordinary laboratory oxidizing agents, they undergo combustion, as do aldehydes.

Some Common Carbonyl Compounds

Formaldehyde has an irritating odor. Because of its reactivity, it is difficult to handle in the gaseous state. For many uses, it is therefore dissolved in water and sold as a 37% to 40% aqueous solution called *formalin*. Formaldehyde denatures proteins, rendering them insoluble in water and resistant to bacterial decay. For this reason, formalin is used in embalming solutions and in preserving biological specimens.

Aldehydes are the active components in many other familiar substances. Large quantities of formaldehyde are used to make phenol-formaldehyde resins for gluing the wood sheets in plywood and as adhesives in other building materials. Sometimes the formaldehyde escapes from the materials and causes health problems in some people. While some people seem unaffected, others experience coughing, wheezing, eye irritation, and other symptoms.

> *The odor of green leaves is due in part to a carbonyl compound, cis-3-hexenal, which with related compounds is used to impart a "green" herbal odor to shampoos and other products.*

Acetaldehyde is an extremely volatile, colorless liquid. It is a starting material for the preparation of many other organic compounds. Acetaldehyde is formed as a metabolite in the fermentation of sugars and in the detoxification of alcohol in the liver. Aldehydes are the active components of many other familiar materials (Figure **14.2.2**).





Figure **14.2.2** Some Interesting Aldehydes. (a) Benzaldehyde is an oil found in almonds; (b) cinnamaldehyde is oil of cinnamon; (c) vanillin gives vanilla its flavor; (d) *cis*-3-hexenal provides an herbal odor; and (e) *trans*-2-*cis*-6-nonadienal gives a cucumber odor.

Acetone is the simplest and most important ketone. Because it is miscible with water as well as with most organic solvents, its chief use is as an industrial solvent (for example, for paints and lacquers). It is also the chief ingredient in some brands of nail polish remover.

To Your Health: Acetone in Blood, Urine, and Breath

Acetone is formed in the human body as a by-product of lipid metabolism. Normally, acetone does not accumulate to an appreciable extent because it is oxidized to carbon dioxide and water. The normal concentration of acetone in the human body is less than 1 mg/100 mL of blood. In certain disease states, such as uncontrolled diabetes mellitus, the acetone concentration rises to higher levels. It is then excreted in the urine, where it is easily detected. In severe cases, its odor can be noted on the breath.

Ketones are also the active components of other familiar substances, some of which are noted in the accompanying figure.



Some ketones have interesting properties: (a) Butter flavoring comes from 2,3-butanedione; (b) β-ionone is responsible for the odor of violets; (c) muscone is musk oil, an ingredient in perfumes; and (d) camphor is used in some insect repellents.

Certain steroid hormones have the ketone functional group as a part of their structure. Two examples are progesterone, a hormone secreted by the ovaries that stimulates the growth of cells in the uterine wall and prepares it for attachment of a fertilized egg, and testosterone, the main male sex hormone. These and other sex hormones affect our development and our lives in fundamental ways.

Summary

The polar carbon-to-oxygen double bond causes aldehydes and ketones to have higher boiling points than those of ethers and alkanes of similar molar masses but lower than those of comparable alcohols that engage in intermolecular hydrogen bonding. Aldehydes are readily oxidized to carboxylic acids, whereas ketones resist oxidation.

14.2: Physical Properties of Aldehydes and Ketones is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **14.10: Properties of Aldehydes and Ketones** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.3: Physical Properties of Carboxylic Acids

Learning Objectives

- Compare the boiling points of carboxylic acids with alcohols of similar molar mass.
- Compare the solubilities of carboxylic acids in water with the solubilities of comparable alkanes and alcohols in water.

Many carboxylic acids are colorless liquids with disagreeable odors. The carboxylic acids with 5 to 10 carbon atoms all have "goaty" odors (explaining the odor of Limburger cheese). These acids are also produced by the action of skin bacteria on human sebum (skin oils), which accounts for the odor of poorly ventilated locker rooms. The acids with more than 10 carbon atoms are waxlike solids, and their odor diminishes with increasing molar mass and resultant decreasing volatility.

Carboxylic acids exhibit strong hydrogen bonding between molecules. They therefore have high boiling points compared to other substances of comparable molar mass.

The carboxyl group readily engages in hydrogen bonding with water molecules (Figure **14.3.1**). The acids with one to four carbon atoms are completely miscible with water. Solubility decreases as the carbon chain length increases because dipole forces become less important and dispersion forces become more predominant. Hexanoic acid [$CH_3(CH_2)_4COOH$] is barely soluble in water (about 1.0 g/100 g of water). Palmitic acid [$CH_3(CH_2)_{14}COOH$], with its large nonpolar hydrocarbon component, is essentially insoluble in water. The carboxylic acids generally are soluble in such organic solvents as ethanol, toluene, and diethyl ether.




Figure **14.3.1**: Hydrogen Bonding between an Acetic Acid Molecule and Water Molecules. Carboxylic acids of low molar mass are quite soluble in water.

Table 15.4.1 lists some physical properties for selected carboxylic acids. The first six are homologs. Notice that the boiling points increase with increasing molar mass, but the melting points show no regular pattern.

Table **14.3.1**: Physical Constants of Carboxylic Acids

Condensed Structural Formula	Name of Acid	Melting Point (°C)	Boiling Point (°C)	Solubility (g/100 g of Water)
HCOOH	formic acid	8	100	miscible
CH_3COOH	acetic acid	17	118	miscible
CH_3CH_2COOH	propionic acid	–22	141	miscible
$CH_3(CH_2)_2COOH$	butyric acid	–5	163	miscible
$CH_3(CH_2)_3COOH$	valeric acid	–35	187	5
$CH_3(CH_2)_4COOH$	caproic acid	–3	205	1.1
C_6H_5COOH	benzoic acid	122	249	0.29

Key Takeaways

- Carboxylic acids have high boiling points compared to other substances of comparable molar mass. Boiling points increase with molar mass.
- Carboxylic acids having one to four carbon atoms are completely miscible with water. Solubility decreases with molar mass.

14.3: Physical Properties of Carboxylic Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.3: Physical Properties of Carboxylic Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.4: Physical Properties of Esters

Learning Objectives

- Compare the boiling points of esters with alcohols of similar molar mass.
- Compare the solubilities of esters in water with the solubilities of comparable alkanes and alcohols in water.

Ester molecules are polar but have no hydrogen atom attached directly to an oxygen atom. They are therefore incapable of engaging in intermolecular hydrogen bonding with one another and thus have considerably lower boiling points than their isomeric carboxylic acids counterparts. Because ester molecules can engage in hydrogen bonding with water molecules, however, esters of low molar mass are somewhat soluble in water. Borderline solubility occurs in those molecules that have three to five carbon atoms. Table **14.4.1** lists the physical properties of some common esters.

Esters are common solvents. Ethyl acetate is used to extract organic solutes from aqueous solutions—for example, to remove caffeine from coffee. It also is used to remove nail polish and paint. Cellulose nitrate is dissolved in ethyl acetate and butyl acetate to form lacquers. The solvent evaporates as the lacquer "dries," leaving a thin film on the surface. High boiling esters are used as softeners (plasticizers) for brittle plastics.

Table **14.4.1**: *Physical Properties of Some Esters*

Condensed Structural Formula	Name	Molar Mass	Melting Point (°C)	Boiling Point (°C)	Aroma
$HCOOCH_3$	methyl formate	60	−99	32	
$HCOOCH_2CH_3$	ethyl formate	74	−80	54	rum
CH_3COOCH_3	methyl acetate	74	−98	57	
$CH_3COOCH_2CH_3$	ethyl acetate	88	−84	77	
$CH_3CH_2CH_2COOCH_3$	methyl butyrate	102	−85	102	apple
$CH_3CH_2CH_2COOCH_2CH_3$	ethyl butyrate	116	−101	121	pineapple
$CH_3COO(CH_2)_4CH_3$	pentyl acetate	130	−71	148	pear
$CH_3COOCH_2CH_2CH(CH_3)_2$	isopentyl acetate	130	−79	142	banana
$CH_3COOCH_2C_6H_5$	benzyl acetate	150	−51	215	jasmine
$CH_3CH_2CH_2COO(CH_2)_4CH_3$	pentyl butyrate	158	−73	185	apricot
$CH_3COO(CH_2)_7CH_3$	octyl acetate	172	−39	210	orange

Summary

Esters have polar bonds but do not engage in hydrogen bonding and are therefore intermediate in boiling points between the nonpolar alkanes and the alcohols, which engage in hydrogen bonding. Ester molecules can engage in hydrogen bonding with water, so esters of low molar mass are therefore somewhat soluble in water.

14.4: Physical Properties of Esters is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.6: Physical Properties of Esters** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.5: Physical Properties of Amines

Learning Objectives

- Explain why the boiling points of primary and secondary amines are higher than those of alkanes or ethers of similar molar mass but are lower than those of alcohols.
- Compare the boiling points of tertiary amines with alcohols, alkanes, and ethers of similar molar mass.
- Compare the solubilities in water of amines of five or fewer carbon atoms with the solubilities of comparable alkanes and alcohols in water.

Primary and secondary amines have hydrogen atoms bonded to an nitrogen atom and are therefore capable of hydrogen bonding (part (a) of Figure **14.5.1**), although not as strongly as alcohol molecules (which have hydrogen atoms bonded to an oxygen atom, which is more electronegative than nitrogen). These amines boil at higher temperatures than alkanes but at lower temperatures than alcohols of comparable molar mass. For example, compare the boiling point of methylamine (CH_3NH_2; −6°C) with those of ethane (CH_3CH_3; −89°C) and methanol (CH_3OH; 65°C). Tertiary amines have no hydrogen atom bonded to the nitrogen atom and so cannot participate in intermolecular hydrogen bonding. They have boiling points comparable to those of ethers (Table **14.5.1**).




Figure **14.5.1**: Hydrogen Bonding. (a) Amine molecules are associated through hydrogen bonding. (b) An amine molecule can form a hydrogen bond with water molecules.

Table **14.5.1**: Physical Properties of Some Amines and Comparable Oxygen-Containing Compounds

Name	Condensed Structural Formula	Class	Molar Mass	Boiling Point (°C)	Solubility at 25°C (g/100 g Water)
butylamine	$CH_3CH_2CH_2CH_2NH_2$	1°	73	78	miscible
diethylamine	$(CH_3CH_2)_2NH$	2°	73	55	miscible
butyl alcohol	$CH_3CH_2CH_2CH_2OH$	—	74	118	8
dipropylamine	$(CH_3CH_2CH_2)_2NH$	2°	101	111	4
triethylamine	$(CH_3CH_2)_3N$	3°	101	90	14
dipropyl ether	$(CH_3CH_2CH_2)_2O$	—	102	91	0.25


All three classes of amines can engage in hydrogen bonding with water (Figure **14.5.1***b*). Amines of low molar mass are quite soluble in water; the borderline of solubility in water is at five or six carbon atoms.

To Your Health: Amines in Death and Life

Amines have “interesting” odors. The simple ones smell very much like ammonia. Higher aliphatic amines smell like decaying fish. Or perhaps we should put it the other way around: Decaying fish give off odorous amines. The stench of rotting fish is due in part to two diamines: putrescine and cadaverine. They arise from the decarboxylation of ornithine and lysine, respectively, amino acids that are found in animal cells.

$$HOCH_2CH_2OH$$

Aromatic amines generally are quite toxic. They are readily absorbed through the skin, and workers must exercise caution when handling these compounds. Several aromatic amines, including β-naphthylamine, are potent carcinogens.



Key Takeaways

- Primary and secondary amines have higher boiling points than those of alkanes or ethers of similar molar mass because they can engage in intermolecular hydrogen bonding. Their boiling points are lower than those of alcohols because alcohol molecules have hydrogen atoms bonded to an oxygen atom, which is more electronegative.
- The boiling points of tertiary amines, which cannot engage in hydrogen bonding because they have no hydrogen atom on the nitrogen atom, are comparable to those of alkanes and ethers of similar molar mass.
- Because all three classes of amines can engage in hydrogen bonding with water, amines of low molar mass are quite soluble in water.

14.5: Physical Properties of Amines is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.11: Physical Properties of Amines** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.6: Physical Properties of Amides

Learning Objectives

- Compare the boiling points of amides with alcohols of similar molar mass.
- Compare the solubilities in water of amides of five or fewer carbon atoms with the solubilities of comparable alkanes and alcohols in water.

With the exception of formamide ($HCONH_2$), which is a liquid, all simple amides are solids (Table **14.6.1**). The lower members of the series are soluble in water, with borderline solubility occurring in those that have five or six carbon atoms. Like the esters, solutions of amides in water usually are neutral—neither acidic nor basic.

Table **14.6.1**: *Physical Constants of Some Unsubstituted Amides*

Condensed Structural Formula	Name	Melting Point (°C)	Boiling Point (°C)	Solubility in Water
$HCONH_2$	formamide	2	193	soluble
CH_3CONH_2	acetamide	82	222	soluble
$CH_3CH_2CONH_2$	propionamide	81	213	soluble
$CH_3CH_2CH_2CONH_2$	butyramide	115	216	soluble
$C_6H_5CONH_2$	benzamide	132	290	slightly soluble

The amides generally have high boiling points and melting points. These characteristics and their solubility in water result from the polar nature of the amide group and hydrogen bonding (Figure **14.6.1**). (Similar hydrogen bonding plays a critical role in determining the structure and properties of proteins, deoxyribonucleic acid [DNA], ribonucleic acid [RNA], and other giant molecules so important to life processes.




Figure **14.6.1**: Hydrogen Bonding in Amides. Amide molecules can engage in hydrogen bonding with water molecules (a). Those amides with a hydrogen atom on the nitrogen atom can also engage in hydrogen bonding (b). Both hydrogen bonding networks extend in all directions.

Key Takeaways

- Most amides are solids at room temperature; the boiling points of amides are much higher than those of alcohols of similar molar mass.
- Amides of five or fewer carbon atoms are soluble in water?.

14.6: Physical Properties of Amides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.14: Physical Properties of Amides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.7: Ionization of Carboxylic Acids

Learning Objectives

- Name the typical reactions that take place with carboxylic acids.
- Describe how carboxylic acids react with basic compounds.

Water-soluble carboxylic acids ionize slightly in water to form moderately acidic solutions.

$$\mathbf{RCOOH + H_2O \rightleftharpoons RCOO^- + H_3O^+}$$

Their aqueous solutions exhibit the typical properties of acids, such as changing litmus from blue to red.

The anion formed when a carboxylic acid dissociates is called the *carboxylate* anion ($RCOO^-$).

Whether soluble in water or not, carboxylic acids react with aqueous solutions of sodium hydroxide (NaOH), sodium carbonate (Na_2CO_3), and sodium bicarbonate ($NaHCO_3$) to form salts:

$$RCOOH + NaOH(aq) \rightarrow RCOO^-Na^+(aq) + H_2O$$

$$2RCOOH + Na_2CO_3(aq) \rightarrow 2RCOO^-Na^+(aq) + H_2O + CO_2(g)$$

$$RCOOH + NaHCO_3(aq) \rightarrow RCOO^-Na^+(aq) + H_2O + CO_2(g)$$

In these reactions, the carboxylic acids act like inorganic acids: they neutralize basic compounds. With solutions of carbonate (CO_3^{2-}) and bicarbonate (HCO_3^-) ions, they also form carbon dioxide gas.

Carboxylic acid salts are named in the same manner as inorganic salts: the name of the cation is followed by the name of the organic anion. The name of the anion is obtained by dropping the *-ic* ending of the acid name and replacing it with the suffix *-ate*. This rule applies whether we are using common names or International Union of Pure and Applied Chemistry (IUPAC) names:

$CH_3COO^-Li^+$	$CH_3CH_2CH_2COO^-K^+$	$C_6H_5COO^-Na^+$
Lithium acetate (lithium ethanoate)	Potassium butyrate (potassium butanoate)	Sodium benzoate

The salts of long-chain carboxylic acids are called soaps. We discuss the chemistry of soaps elsewhere.

$CH_3(CH_2)_{13}CH_2COO^-Na^+$

Sodium palmitate (a soap)

Example 14.7.1

Write an equation for each reaction.

1. the ionization of propionic acid in water (H_2O)
2. the neutralization of propionic acid with aqueous sodium hydroxide (NaOH)

Solution

Propionic acid has three carbon atoms, so its formula is CH_2CH_2COOH.

1. Propionic acid ionizes in water to form a propionate ion and a hydronium (H_3O^+) ion. $CH_3CH_2COOH(aq) + H_2O(\ell) \rightarrow CH_3CH_2COO^-(aq) + H_3O^+(aq)$
2. Propionic acid reacts with NaOH(aq) to form sodium propionate and water. $CH_3CH_2COOH(aq) + NaOH(aq) \rightarrow CH_3CH_2COO^-Na^+(aq) + H_2O(\ell)$

Exercise 14.7.1

Write an equation for each reaction.

a. the ionization of formic acid in water
b. the ionization of *p*-chlorobenzoic acid in water

Example 14.7.2

Write an equation for the reaction of decanoic acid with each compound.

a. aqueous sodium hydroxide (NaOH)
b. aqueous sodium bicarbonate ($NaHCO_3$)

Solution

a. Decanoic acid has 10 carbon atoms. It reacts with NaOH to form a salt and water (H_2O). $CH_3(CH_2)_8COOH + NaOH(aq) \rightarrow CH_3(CH_2)_8COO^-Na^+(aq) + H_2O(\ell)$
b. With $NaHCO_3$, the products are a salt, H_2O, and carbon dioxide (CO_2). $CH_3(CH_2)_8COOH + NaHCO_3(aq) \rightarrow CH_3(CH_2)_8COO^-Na^+(aq) + H_2O(\ell) + CO_2(g)$


Exercise 14.7.3

Write an equation for the reaction of benzoic acid with each compound.

a. aqueous sodium hydroxide (NaOH)
b. aqueous sodium bicarbonate ($NaHCO_3$)

To Your Health: Organic Salts as Preservatives

Some organic salts are used as preservatives in food products. They prevent spoilage by inhibiting the growth of bacteria and fungi. Calcium and sodium propionate, for example, are added to processed cheese and bakery goods; sodium benzoate is added to cider, jellies, pickles, and syrups; and sodium sorbate and potassium sorbate are added to fruit juices, sauerkraut, soft drinks, and wine. Look for them on ingredient labels the next time you shop for groceries.



Key Takeaways

- Soluble carboxylic acids are weak acids in aqueous solutions.
- Carboxylic acids neutralize bases to form salts.

14.7: Ionization of Carboxylic Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **15.4: Chemical Properties of Carboxylic Acids- Ionization and Neutralization** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.8: Amines as Bases

Learning Objectives

- Name the typical reactions that take place with amines.
- Describe heterocyclic amines.

Recall that ammonia (NH_3) acts as a base because the nitrogen atom has a lone pair of electrons that can accept a proton. Amines also have a lone electron pair on their nitrogen atoms and can accept a proton from water to form substituted ammonium (NH_4^+) ions and hydroxide (OH^-) ions:



As a specific example, methylamine reacts with water to form the methylammonium ion and the OH^- ion.

$$\underset{\text{Methylamine}}{CH_3NH_2(aq)} + H_2O \rightleftharpoons \underset{\text{Methylammonium ion}}{CH_3NH_3^+(aq)} + OH^-(aq)$$

Nearly all amines, including those that are not very soluble in water, will react with strong acids to form salts soluble in water.

$$\underset{\text{Octylamine (insoluble)}}{CH_3(CH_2)_6NH_2(l)} + HNO_3(aq) \longrightarrow \underset{\text{Octylammonium nitrate (soluble)}}{CH_3(CH_2)_6NH_3{}^+NO_3^-(aq)}$$

Amine salts are named like other salts: the name of the cation is followed by the name of the anion.

Example 14.8.1

What are the formulas of the acid and base that react to form $[CH_3NH_2CH_2CH_3]^+CH_3COO^-$?

Solution

The cation has two groups—methyl and ethyl—attached to the nitrogen atom. It comes from ethylmethylamine ($CH_3NHCH_2CH_3$). The anion is the acetate ion. It comes from acetic acid (CH_3COOH).

Exercise 14.8.1

What are the formulas of the acid and base that react to form $\mathbf{(CH_3CH_2CH_2)_3NH^+I^-}$?

To Your Health: Amine Salts as Drugs

Salts of aniline are properly named as *anilinium* compounds, but an older system, still in use for naming drugs, identifies the salt of aniline and hydrochloric acid as “aniline hydrochloride.” These compounds are ionic—they are salts—and the properties of the compounds (solubility, for example) are those characteristic of salts. Many drugs that are amines are converted to hydrochloride salts to increase their solubility in aqueous solution.

Heterocyclic Amines


Looking back at the various cyclic hydrocarbons discussed previously, we see that all the atoms in the rings of these compounds are carbon atoms. In other cyclic compounds, called **heterocyclic compounds** (Greek *heteros*, meaning “other”), nitrogen, oxygen, sulfur, or some other atom is incorporated in the ring. Many heterocyclic compounds are important in medicine and biochemistry.

Some compose part of the structure of the nucleic acids, which in turn compose the genetic material of cells and direct protein synthesis.

Many heterocyclic amines occur naturally in plants. Like other amines, these compounds are basic. Such a compound is an **alkaloid**, a name that means "like alkalis." Many alkaloids are physiologically active, including the familiar drugs caffeine, nicotine, and cocaine.


To Your Health: Three Well-Known Alkaloids

Caffeine is a stimulant found in coffee, tea, and some soft drinks. Its mechanism of action is not well understood, but it is thought to block the activity of adenosine, a heterocyclic base that acts as a neurotransmitter, a substance that carries messages across a tiny gap (synapse) from one nerve cell (neuron) to another cell. The effective dose of caffeine is about 200 mg, corresponding to about two cups of strong coffee or tea.




Caffeine

Nicotine acts as a stimulant by a different mechanism; it probably mimics the action of the neurotransmitter acetylcholine. People ingest this drug by smoking or chewing tobacco. Its stimulant effect seems transient, as this initial response is followed by depression. Nicotine is highly toxic to animals. It is especially deadly when injected; the lethal dose for a human is estimated to be about 50 mg. Nicotine has also been used in agriculture as a contact insecticide.



Nicotine

Cocaine acts as a stimulant by preventing nerve cells from taking up dopamine, another neurotransmitter, from the synapse. High levels of dopamine are therefore available to stimulate the pleasure centers of the brain. The enhancement of dopamine action is thought to be responsible for cocaine's "high" and its addictive properties. After the binge, dopamine is depleted in less than an hour. This leaves the user in a pleasureless state and (often) craving more cocaine.



Cocaine

Cocaine is used as the salt cocaine hydrochloride and in the form of broken lumps of the free (unneutralized) base, which is called *crack cocaine*.

$$\underbrace{C_{17}H_{21}O_4N}_{\text{cocaine (freebase)}} + HCl \longrightarrow \underbrace{C_{17}H_{21}O_4NH^+Cl^-}_{\text{cocaine hydrochloride}}$$

Because it is soluble in water, cocaine hydrochloride is readily absorbed through the watery mucous membranes of the nose when it is snorted. Crack cocaine is more volatile than cocaine hydrochloride. It vaporizes at the temperature of a burning cigarette. When smoked, cocaine reaches the brain in 15 s.

Summary

Amines are bases; they react with acids to form salts. Salts of aniline are properly named as *anilinium* compounds, but an older system is used to name drugs: the salts of amine drugs and hydrochloric acid are called “hydrochlorides.” Heterocyclic amines are cyclic compounds with one or more nitrogen atoms in the ring.

14.8: Amines as Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **15.12: Amines as Bases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

14.9: Chirality

http://chemwiki.ucdavis.edu/Organic_Chemistry/Chirality/Fundamentals_of_Chirality

http://chemwiki.ucdavis.edu/Organic_Chemistry/Organic_Chemistry_With_a_Biological_Emphasis/Chapter__3%3A_Conformations_and_Stereochemistry/Section_3.3%3A_Stereoisomerism_%E2%80%93_chirality,_stereocenters,_enantiomers

14.9: Chirality is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

14.10: End-of-Chapter Material

Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the summary and ask yourself how they relate to the topics in the chapter.

A **carboxylic acid** (RCOOH) contains the functional group COOH, called the **carboxyl group**, which has an OH group attached to a carbonyl carbon atom. An **ester** (RCOOR′) has an OR′ group attached to a carbonyl carbon atom. An **amine** is derived from ammonia (NH_3), with one, two, or all three of the hydrogen atoms of NH_3 replaced by an alkyl (or an aryl) group. The **amide** functional group has a carbonyl group joined to a nitrogen atom from NH_3 or an amine.

There are many familiar carboxylic acids. The R group may be a hydrogen atom (as in formic acid, HCOOH), an alkyl group (as in acetic acid, CH_2COOH), or an aryl group (as in benzoic acid, C_6H_5COOH). The location of substituents along the carbon chain is indicated by a Greek letter (for common names) or a number (for names from the International Union of Pure and Applied Chemistry).

A carboxylic acid is formed by the oxidation of an aldehyde with the same number of carbon atoms. Because aldehydes are formed from primary alcohols, these alcohols are also a starting material for carboxylic acids.

Carboxylic acids have strong, often disagreeable, odors. They are highly polar molecules and readily engage in hydrogen bonding, so they have relatively high boiling points.

Carboxylic acids are weak acids. They react with bases to form salts and with carbonates and bicarbonates to form carbon dioxide gas and the salt of the acid.

Esters are pleasant-smelling compounds that are responsible for the fragrances of flowers and fruits. They have lower boiling points than comparable carboxylic acids because, even though ester molecules are somewhat polar, they cannot engage in hydrogen bonding. However, with water, esters can engage in hydrogen bonding; consequently, the low molar mass esters are soluble in water. Esters can be synthesized by **esterification**, in which a carboxylic acid and an alcohol are combined under acidic conditions. Esters are neutral compounds that undergo **hydrolysis**, a reaction with water. Under acidic conditions, hydrolysis is essentially the reverse of esterification. When carried out under basic conditions, the process is called **saponification**.

Inorganic acids also react with alcohols to form esters. Some of the most important esters in biochemistry are those formed from phosphoric acid.


Amines are nitrogen-containing organic molecules derived from ammonia (NH_3). A **primary (1°) amine** (RNH_2) has one organic group bonded to the nitrogen atom, a **secondary (2°) amine** (R_2NH) has two organic groups bonded to the nitrogen atom, and a **tertiary (3°) amine** (R_3N) has three organic groups bonded to the nitrogen atom. Amines are basic compounds that react with strong acids to produce ammonium (NH_4^+) salts. A cyclic compound in which the ring contains one or more noncarbon atoms is called a **heterocyclic compound**. There are many heterocyclic amines, including many physiologically important ones. **Alkaloids** are heterocyclic amines found in many plants. Caffeine, nicotine, and cocaine are familiar alkaloids.

Organic compounds containing a carbonyl group bonded to a nitrogen atom are amides, and the carbon-to-nitrogen bond is an amide linkage (or a peptide linkage). Most amides are colorless and odorless, and the lighter ones are soluble in water. Because they are polar molecules, amides have comparatively high boiling points and melting points. Amides are synthesized from carboxylic acids and NH_3 or amines. Amides are neutral compounds. They resist hydrolysis in water, but acids, bases, and enzymes catalyze the reaction.


Additional Exercises

1. Of the families of organic compounds discussed in this chapter, which are known for their typically unpleasant odors? Which for their characteristically pleasant aromas?
2. What is esterification of a carboxylic acid? How does it differ from neutralization?
3. Like alcohols, phenols form esters with carboxylic acids. The hydrocarbon group from phenol is called phenyl. Draw the structure of phenyl acetate.

4. Describe the hydrogen bonding in carboxylic acids, both acid-acid and acid-water. How does this influence their physical properties?
5. Which compound is more soluble in water—benzoic acid or sodium benzoate? Explain.
6. Dicarboxylic acids have two carboxyl groups and are named with the ending *-dioic acid*. Give the equation for the reaction of 1,5-pentanedioic acid ($HOOCCH_2CH_2CH_2COOH$; common name, glutaric acid) with each of the following:
    1. 1 mol of NaOH
    2. 2 mol of NaOH
7. Without consulting tables, arrange the following compounds in order of increasing boiling point: butyl alcohol, methyl acetate, pentane, and propionic acid.
8. From which alcohol might each acid be prepared via oxidation with acidic dichromate?
    1. CH_3CH_2COOH
    2. $HCOOH$
    3. $HOOCH_2COOH$
    4. $(CH_3)_2CHCH_2COOH$
9. The distinctive aroma and flavor of oranges are due in part to octyl acetate, an ester formed from 1-octanol (octyl alcohol) and acetic acid. Write the condensed structural formula for octyl acetate.
10. A lactone is a cyclic ester. What product is formed in following reaction?



11. A lactam is a cyclic amide. What product is formed in the following reaction?




12. Draw the structures for the eight isomeric amines that have the molecular formula $C_4H_{11}N$. Give each a common name and classify it as primary, secondary, or tertiary.
13. Draw the structures for the five isomeric amines that have the molecular formula C_7H_9N and contain a benzene ring. Classify each compound as primary, secondary, or tertiary.
14. Cocaine is usually used in the form of the salt cocaine hydrochloride and sniffed up the nose. Some prefer to ingest their cocaine by smoking it (mixed with tobacco, for example). Before smoking, the cocaine hydrochloride must be converted back to the free base (that is, to the molecular form). Explain the choice of dosage form for each route of administration.
15. Draw the structures all the isomeric amides that have the molecular formula C_4H_9NO.
16. An ester with the molecular formula $C_6H_{12}O_2$ was hydrolyzed in aqueous acid to yield an acid Y and an alcohol Z. Oxidation of the alcohol with potassium dichromate ($K_2Cr_2O_7$) gave the identical acid Y. What is the condensed structural formula of the ester?

17. The neutralization of 125 mL of a 0.400 M NaOH solution requires 5.10 g of a monocarboxylic acid. Draw all the possible structures for the acid.
18. If 3.00 g of acetic acid reacts with excess methanol, how many grams of methyl acetate are formed?
19. How many milliliters of a 0.100 M barium hydroxide solution are required to neutralize 0.500 g of dichloroacetic acid?

Answers

1. unpleasant: carboxylic acids; pleasant: esters




3.

5. sodium benzoate because it is ionic and forms ion-dipole forces with water; benzoic acid can engage in hydrogen bonding only with water


7. pentane < methyl acetate < butyl alcohol < propionic acid

9. $CH_3COOCH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$


11. $H_3N^+CH_2CH_2COOH$




o-Methylaniline (1°)




m-Methylaniline (1°)



p-Methylaniline (1°)




Methylphenylamine (2°)




[Benzylamine] (1°)


13.




Butanamide




2-Methylpropanamide




N,N-dimethylethanamide




N-ethyl-N-methylmethanamide




N-methylpropanamide



N-propylmethanamide





N-ethylethanamide




N-isopropylmethanamide

15.







17.

19. 20.0 mL

14.10: End-of-Chapter Material is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

CHAPTER OVERVIEW

15: Carbohydrates

Glucose is one of the carbohydrates you will learn about in this chapter as we begin the study of biochemistry—the chemistry of molecules found in living organisms. Later we will study the other three major types of macromolecules found in living organisms: lipids, proteins, and nucleic acids.

15.1: Introduction to Carbohydrates
15.2: Classes of Monosaccharides
15.3: Important Hexoses
15.4: Cyclic Structures of Monosaccharides
15.5: Properties of Monosaccharides
15.6: Disaccharides
15.7: Polysaccharides
15.8: Diabetes
15.9: Chapter Summary


15: Carbohydrates is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

15.1: Introduction to Carbohydrates

Learning Objectives

- To recognize carbohydrates and classify them as mono-, di-, or polysaccharides.


All carbohydrates consist of carbon, hydrogen, and oxygen atoms and are polyhydroxy aldehydes or ketones or are compounds that can be broken down to form such compounds. Examples of carbohydrates include starch, fiber, the sweet-tasting compounds called sugars, and structural materials such as cellulose. The term *carbohydrate* had its origin in a misinterpretation of the molecular formulas of many of these substances. For example, because its formula is $C_6H_{12}O_6$, glucose was once thought to be a "carbon hydrate" with the structure $C_6{\cdot}6H_2O$.





Because glucose has an aldehyde functional group and several –OH (hydroxyl) groups, it is described as a polyhydroxy aldehyde.


Example 15.1.1

Which compounds would be classified as carbohydrates?

a. 

b. 

c. 

d. 

Solution

a. This is a carbohydrate because the molecule contains an aldehyde functional group with OH groups on the other two carbon atoms.





b. This is not a carbohydrate because the molecule does not contain an aldehyde or a ketone functional group.

c. This is a carbohydrate because the molecule contains a ketone functional group with OH groups on the other two carbon atoms.

d. This is not a carbohydrate; although it has a ketone functional group, one of the other carbons atoms does not have an OH group attached.

? Exercise 15.1.1

Which compounds would be classified as carbohydrates?

1. 
2. 
3. 
4. 

Green plants are capable of synthesizing glucose ($C_6H_{12}O_6$) from carbon dioxide (CO_2) and water (H_2O) by using solar energy in the process known as photosynthesis:

$$6\,CO_2 + 6\,H_2O + 686\ \text{kcal} \rightarrow C_6H_{12}O_6 + 6\,O_2 \tag{15.1.1}$$

(The 686 kcal come from solar energy.) Plants can use the glucose for energy or convert it to larger carbohydrates, such as starch or cellulose. Starch provides energy for later use, perhaps as nourishment for a plant's seeds, while cellulose is the structural material of plants. We can gather and eat the parts of a plant that store energy—seeds, roots, tubers, and fruits—and use some of that energy ourselves. Carbohydrates are also needed for the synthesis of nucleic acids and many proteins and lipids.

Animals, including humans, cannot synthesize carbohydrates from carbon dioxide and water and are therefore dependent on the plant kingdom to provide these vital compounds. We use carbohydrates not only for food (about 60%–65% by mass of the average diet) but also for clothing (cotton, linen, rayon), shelter (wood), fuel (wood), and paper (wood).

The simplest carbohydrates—those that cannot be hydrolyzed to produce even smaller carbohydrates—are called monosaccharides. Two or more monosaccharides can link together to form chains that contain from two to several hundred or thousand monosaccharide units. Prefixes are used to indicate the number of such units in the chains. Disaccharide molecules have two monosaccharide units, *trisaccharide* molecules have three units, and so on. Chains with many monosaccharide units joined together are called polysaccharides. All these so-called higher saccharides can be hydrolyzed back to their constituent monosaccharides.

> *Compounds that cannot be hydrolyzed will not react with water to form two or more smaller compounds.*

Summary

Carbohydrates are an important group of biological molecules that includes sugars and starches. Photosynthesis is the process by which plants use energy from sunlight to synthesize carbohydrates. A monosaccharide is the simplest carbohydrate and cannot be hydrolyzed to produce a smaller carbohydrate molecule. Disaccharides contain two monosaccharide units, and polysaccharides contain many monosaccharide units.

15.1: Introduction to Carbohydrates is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **16.1: Carbohydrates** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

15.2: Classes of Monosaccharides

Learning Objectives

- Classify monosaccharides as aldoses or ketoses and as trioses, tetroses, pentoses, or hexoses.
- Distinguish between a D sugar and an L sugar.

The naturally occurring monosaccharides contain three to seven carbon atoms per molecule. Monosaccharides of specific sizes may be indicated by names composed of a stem denoting the number of carbon atoms and the suffix *-ose*. For example, the terms *triose*, *tetrose*, *pentose*, and *hexose* signify monosaccharides with, respectively, three, four, five, and six carbon atoms. Monosaccharides are also classified as aldoses or ketoses. Those monosaccharides that contain an aldehyde functional group are called aldoses; those containing a ketone functional group on the second carbon atom are ketoses. Combining these classification systems gives general names that indicate both the type of carbonyl group *and* the number of carbon atoms in a molecule. Thus, monosaccharides are described as aldotetroses, aldopentoses, ketopentoses, ketoheptoses, and so forth. Glucose and fructose are specific examples of an aldohexose and a ketohexose, respectively.




Example 15.2.1

Draw an example of each type of compound.


a. a ketopentose
b. an aldotetrose

Solution

a. The structure must have five carbon atoms with the second carbon atom being a carbonyl group and the other four carbon atoms each having an OH group attached. Several structures are possible, but one example is shown.



- The structure must have four carbon atoms with the first carbon atom part of the aldehyde functional group. The other three carbon atoms each have an OH group attached. Several structures are possible, but one example is shown.



? Exercise 15.2.1

Draw an example of each type of compound.

a. an aldohexose
b. a ketotetrose

The simplest sugars are the trioses. The possible trioses are shown in part (a) of Figure **15.2.1**; glyceraldehyde is an aldotriose, while dihydroxyacetone is a ketotriose. Notice that two structures are shown for glyceraldehyde. These structures are stereoisomers, and hence are isomers having the same structural formula but differing in the arrangement of atoms or groups of atoms in three-dimensional space. If you make models of the two stereoisomers of glyceraldehyde, you will find that you cannot place one model on top of the other and have each functional group point in the same direction. However, if you place one of the models in front of a mirror, the image in the mirror will be identical to the second stereoisomer in part (b) of Figure **15.2.1**. Molecules that are nonsuperimposable (nonidentical) mirror images of each other are a type of stereoisomer called enantiomers (Greek *enantios*, meaning "opposite").

Note

These are another type of stereoisomers than the *cis-trans* (geometric) isomers previously discussed.




Figure **15.2.1**: Structures of the Trioses. (a) D- and L-glyceraldehyde are mirror images of each other and represent a pair of enantiomers. (b) A ball-and-stick model of D-glyceraldehyde is reflected in a mirror. Note that the reflection has the same structure as L-glyceraldehyde.

A key characteristic of enantiomers is that they have a carbon atom to which four different groups are attached. Note, for example, the four different groups attached to the central carbon atom of glyceraldehyde (part (a) of Figure **15.2.1**). A carbon atom that has four different groups attached is a chiral carbon. If a molecule contains one or more chiral carbons, it is likely to exist as two or more stereoisomers. Dihydroxyacetone does not contain a chiral carbon and thus does not exist as a pair of stereoisomers. Glyceraldehyde, however, has a chiral carbon and exists as a pair of enantiomers. Except for the direction in which each enantiomer rotates plane-polarized light, these two molecules have identical physical properties. One enantiomer has a specific rotation of +8.7°, while the other has a specific rotation of −8.7°.


H. Emil Fischer, a German chemist, developed the convention commonly used for writing two-dimensional representations of the monosaccharides, such as those in part (a) of Figure **15.2.1**. In these structural formulas, the aldehyde group is written at the top, and the hydrogen atoms and OH groups that are attached to each chiral carbon are written to the right or left. (If the monosaccharide is a ketose, the ketone functional group is the second carbon atom.) Vertical lines represent bonds pointing away from you, while horizontal lines represent bonds coming toward you. The formulas of chiral molecules represented in this manner are referred to as **Fischer projections**.

The two enantiomers of glyceraldehyde are especially important because monosaccharides with more than three carbon atoms can be considered as being derived from them. Thus, D- and L-glyceraldehyde provide reference points for designating and drawing all other monosaccharides. Sugars whose Fischer projections terminate in the same configuration as D-glyceraldehyde are designated as D sugars; those derived from L-glyceraldehyde are designated as L sugars.


> *By convention, the penultimate (next-to-last) carbon atom has been chosen as the carbon atom that determines if a sugar is D or L. It is the chiral carbon farthest from the aldehyde or ketone functional group.*

Looking Closer: Polarized Light

A beam of ordinary light can be pictured as a bundle of waves; some move up and down, some sideways, and others at all other conceivable angles. When a beam of light has been polarized, however, the waves in the bundle all vibrate in a single plane. Light altered in this way is called *plane-polarized light*. Much of what chemists know about stereoisomers comes from studying the effects they have on plane-polarized light. In this illustration, the light on the left is not polarized, while that on the right is polarized.



Sunlight, in general, is not polarized; light from an ordinary light bulb or an ordinary flashlight is not polarized. One way to polarize ordinary light is to pass it through Polaroid sheets, special plastic sheets containing carefully oriented organic compounds that permit only light vibrating in a single plane to pass through. To the eye, polarized light doesn't "look" any different from nonpolarized light. We can detect polarized light, however, by using a second sheet of polarizing material, as shown here.



In the photo on the left, two Polaroid sheets are aligned in the same direction; plane-polarized light from the first Polaroid sheet can pass through the second sheet. In the photo on the right, the top Polaroid sheet has been rotated 90° and now blocks the plane-polarized light that comes through the first Polaroid sheet.

Certain substances act on polarized light by rotating the plane of vibration. Such substances are said to be optically active. The extent of optical activity is measured by a polarimeter, an instrument that contains two polarizing lenses separated by a sample tube, as shown in the accompanying figure. With the sample tube empty, maximum light reaches the observer's eye when the two lenses are aligned so that both pass light vibrating in the same plane. When an optically active substance is placed in the sample tube, that substance rotates the plane of polarization of the light passing through it, so that the polarized light emerging from the

sample tube is vibrating in a different direction than when it entered the tube. To see the maximum amount of light when the sample is in place, the observer must rotate one lens to accommodate the change in the plane of polarization.




Figure **15.2.2**: Diagram of a Polarimeter

Some optically active substances rotate the plane of polarized light to the right (clockwise) from the observer's point of view. These compounds are said to be dextrorotatory; substances that rotate light to the left (counterclockwise) are levorotatory. To denote the direction of rotation, a positive sign (+) is given to dextrorotatory substances, and a negative sign (−) is given to levorotatory substances.

Summary

Monosaccharides can be classified by the number of carbon atoms in the structure and/or the type of carbonyl group they contain (aldose or ketose). Most monosaccharides contain at least one chiral carbon and can form stereoisomers. Enantiomers are a specific type of stereoisomers that are mirror images of each other.

15.2: Classes of Monosaccharides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **16.2: Classes of Monosaccharides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

15.3: Important Hexoses

Learning Objectives

- To identify the structures of D-glucose, D-galactose, and D-fructose and describe how they differ from each other.

Although a variety of monosaccharides are found in living organisms, three hexoses are particularly abundant: D-glucose, D-galactose, and D-fructose (Figure 15.3.1). Glucose and galactose are both aldohexoses, while fructose is a ketohexose.




Figure 15.3.1: Structures of Three Important Hexoses. Each hexose is pictured with a food source in which it is commonly found. Source: Photos © Thinkstock.

Glucose

D-Glucose, generally referred to as simply glucose, is the most abundant sugar found in nature; most of the carbohydrates we eat are eventually converted to it in a series of biochemical reactions that produce energy for our cells. It is also known by three other names: *dextrose*, from the fact that it rotates plane-polarized light in a clockwise (dextrorotatory) direction; *corn sugar* because in the United States cornstarch is used in the commercial process that produces glucose from the hydrolysis of starch; and *blood sugar* because it is the carbohydrate found in the circulatory system of animals. Normal blood sugar values range from 70 to 105 mg glucose/dL plasma, and normal urine may contain anywhere from a trace to 20 mg glucose/dL urine.




Figure 15.3.2: Fischer projection of D-glucose

The Fischer projection of D-glucose is given in Figure **15.3.2**. Glucose is a D sugar because the OH group on the fifth carbon atom (the chiral center farthest from the carbonyl group) is on the right. In fact, all the OH groups except the one on the third carbon atom are to the right.

Galactose

D-Galactose does not occur in nature in the uncombined state. It is released when lactose, a disaccharide found in milk, is hydrolyzed. The galactose needed by the human body for the synthesis of lactose is obtained by the metabolic conversion of D-glucose to D-galactose. Galactose is also an important constituent of the glycolipids that occur in the brain and the myelin sheath of nerve cells. For this reason it is also known as *brain sugar*. The structure of D-galactose is shown in Figure **15.3.1**. Notice that the configuration differs from that of glucose only at the fourth carbon atom.

Fructose

D-Fructose, also shown in Figure **15.3.1**, is the most abundant ketohexose. Note that from the third through the sixth carbon atoms, its structure is the same as that of glucose. It occurs, along with glucose and sucrose, in honey (which is 40% fructose) and sweet fruits. Fructose (from the Latin *fructus*, meaning "fruit") is also referred to as *levulose* because it has a specific rotation that is strongly levorotatory (−92.4°). It is the sweetest sugar, being 1.7 times sweeter than sucrose, although many nonsugars are several hundred or several thousand times as sweet (Table **15.3.1**).

Table **15.3.1**: The Relative Sweetness of Some Compounds (Sucrose = 100)

Compound	Relative Sweetness
lactose	16
maltose	32
glucose	74
sucrose	100
fructose	173
aspartame	18,000
acesulfame K	20,000
saccharin	30,000
sucralose	60,000


Looking Closer: Artificial Sweeteners

Although sweetness is commonly associated with mono- and disaccharides, it is not a property found only in sugars. Several other kinds of organic compounds have been synthesized that are far superior as sweetening agents. These so-called high-intensity or artificial sweeteners are useful for people with diabetes or other medical conditions that require them to control their carbohydrate intake. The synthetic compounds are noncaloric or used in such small quantities that they do not add significantly to the caloric value of food.


The first artificial sweetener—saccharin—was discovered by accident in 1879. It is 300 times sweeter than sucrose, but it passes through the body unchanged and thus adds no calories to the diet. After its discovery, saccharin was used until it was banned in the early 1900s. However, during the sugar-short years of World War I, the ban was lifted and was not reinstated at the war's end. One drawback to the use of saccharin is its bitter, metallic aftertaste. The initial solution to this problem was to combine saccharin with cyclamate, a second artificial sweetener discovered in 1937.

In the 1960s and 1970s, several clinical tests with laboratory animals implicated both cyclamate and saccharin as carcinogenic (cancer-causing) substances. The results from the cyclamate tests were completed first, and cyclamate was banned in the United States in 1969. Then a major study was released in Canada in 1977 indicating that saccharin increased the incidence of


bladder cancer in rats. The US Food and Drug Administration (FDA) proposed a ban on saccharin that raised immediate public opposition because saccharin was the only artificial sweetener still available. In response, Congress passed the Saccharin Study and Labeling Act in 1977, permitting the use of saccharin as long as any product containing it was labeled with a consumer warning regarding the possible elevation of the risk of bladder cancer. Today this warning is no longer required; moreover, the FDA is currently reviewing the ban on cyclamate, as 75 additional studies and years of usage in other countries, such as Canada, have failed to show that it has any carcinogenic effect.




Saccharin




Cyclamate



Aspartame



Acesulfame K



Sucralose

A third artificial sweetener, aspartame, was discovered in 1965. This white crystalline compound is about 180 times sweeter than sucrose and has no aftertaste. It was approved for use in 1981 and is used to sweeten a wide variety of foods because it blends well with other food flavors. Aspartame is not used in baked goods, however, because it is not heat stable.

In the body (or when heated), aspartame is initially hydrolyzed to three molecules: the amino acids aspartic acid and phenylalanine and an alcohol methanol. Repeated controversy regarding the safety of aspartame arises partly from the fact that the body metabolizes the released methanol to formaldehyde. It should be noted, though, that a glass of tomato juice has six times as much methanol as a similar amount of a diet soda containing aspartame. The only documented risk connected to aspartame use is for individuals with the genetic disease *phenylketonuria* (PKU); these individuals lack the enzyme needed to metabolize the phenylalanine released when aspartame is broken down by the body. Because of the danger to people with PKU, all products containing aspartame must carry a warning label.

Acesulfame K, discovered just two years after aspartame (1967), was approved for use in the United States in 1988. It is 200 times sweeter than sugar and, unlike aspartame, is heat stable. It has no lingering aftertaste.

One of the newest artificial sweeteners to gain FDA approval (April 1998) for use in the United States is sucralose, a white crystalline solid approximately 600 times sweeter than sucrose. Sucralose is synthesized from sucrose and has three chlorine atoms substituted for three OH groups. It is noncaloric because it passes through the body unchanged. It can be used in baking because it is heat stable.

All of the extensive clinical studies completed to date have indicated that these artificial sweeteners approved for use in the United States are safe for consumption by healthy individuals in moderate amounts.

Summary

Three abundant hexoses in living organisms are the aldohexoses D-glucose and D-galactose and the ketohexose D-fructose.


15.3: Important Hexoses is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

15.4: Cyclic Structures of Monosaccharides

Learning Objectives

- Define what is meant by anomers and describe how they are formed.
- Explain what is meant by mutarotation.

So far we have represented monosaccharides as linear molecules, but many of them also adopt cyclic structures. This conversion occurs because of the ability of aldehydes and ketones to react with alcohols:



You might wonder why the aldehyde reacts with the OH group on the fifth carbon atom rather than the OH group on the second carbon atom next to it. Recall that cyclic alkanes containing five or six carbon atoms in the ring are the most stable. The same is true for monosaccharides that form cyclic structures: rings consisting of five or six carbon atoms are the most stable.




Figure 15.4.1: Cyclization of D-Glucose. D-Glucose can be represented with a Fischer projection (a) or three dimensionally (b). By reacting the OH group on the fifth carbon atom with the aldehyde group, the cyclic monosaccharide (c) is produced.

When a straight-chain monosaccharide, such as any of the structures shown in Figure 15.4.1, forms a cyclic structure, the carbonyl oxygen atom may be pushed either up or down, giving rise to two stereoisomers, as shown in Figure 15.4.2. The structure shown on the left side of Figure 15.4.2, with the OH group on the first carbon atom projected downward, represent what is called the *alpha (α) form*. The structures on the right side, with the OH group on the first carbon atom pointed upward, is the *beta (β) form*. These two stereoisomers of a cyclic monosaccharide are known as anomers; they differ in structure around the anomeric carbon—that is, the carbon atom that was the carbonyl carbon atom in the straight-chain form.

It is possible to obtain a sample of crystalline glucose in which all the molecules have the α structure or all have the β structure. The α form melts at 146°C and has a specific rotation of +112°, while the β form melts at 150°C and has a specific rotation of +18.7°. When the sample is dissolved in water, however, a mixture is soon produced containing both anomers as well as the straight-chain form, in dynamic equilibrium (part (a) of Figure 15.4.2). You can start with a pure crystalline sample of glucose consisting entirely of either anomer, but as soon as the molecules dissolve in water, they open to form the carbonyl group and then reclose to form either the α or the β anomer. The opening and closing repeats continuously in an ongoing interconversion between anomeric forms and is referred to as mutarotation (Latin *mutare*, meaning “to change”). At equilibrium, the mixture consists of

about 36% α-D-glucose, 64% β-D-glucose, and less than 0.02% of the open-chain aldehyde form. The observed rotation of this solution is +52.7°.




Figure **15.4.2**: Monosaccharides. In an aqueous solution, monosaccharides exist as an equilibrium mixture of three forms. The interconversion between the forms is known as *mutarotation*, which is shown for D-glucose (a) and D-fructose (b).

Even though only a small percentage of the molecules are in the open-chain aldehyde form at any time, the solution will nevertheless exhibit the characteristic reactions of an aldehyde. As the small amount of free aldehyde is used up in a reaction, there is a shift in the equilibrium to yield more aldehyde. Thus, *all* the molecules may eventually react, even though very little free aldehyde is present at a time.

Commonly, (e.g., in Figures **15.4.1** and **15.4.2**) the cyclic forms of sugars are depicted using a convention first suggested by Walter N. Haworth, an English chemist. The molecules are drawn as planar hexagons with a darkened edge representing the side facing toward the viewer. The structure is simplified to show only the functional groups attached to the carbon atoms. Any group written to the right in a Fischer projection appears below the plane of the ring in a Haworth projection, and any group written to the left in a Fischer projection appears above the plane in a Haworth projection.

The difference between the α and the β forms of sugars may seem trivial, but such structural differences are often crucial in biochemical reactions. This explains why we can get energy from the starch in potatoes and other plants but not from cellulose, even though both starch and cellulose are polysaccharides composed of glucose molecules linked together.

Summary

Monosaccharides that contain five or more carbons atoms form cyclic structures in aqueous solution. Two cyclic stereoisomers can form from each straight-chain monosaccharide; these are known as anomers. In an aqueous solution, an equilibrium mixture forms between the two anomers and the straight-chain structure of a monosaccharide in a process known as mutarotation.

15.4: Cyclic Structures of Monosaccharides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **16.4: Cyclic Structures of Monosaccharides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


15.5: Properties of Monosaccharides

Learning Objectives

- To identify the physical and chemical properties of monosaccharides.

Monosaccharides such as glucose and fructose are crystalline solids at room temperature, but they are quite soluble in water, each molecule having several OH groups that readily engage in hydrogen bonding. The chemical behavior of these monosaccharides is likewise determined by their functional groups.

An important reaction of monosaccharides is the oxidation of the aldehyde group, one of the most easily oxidized organic functional groups. Aldehyde oxidation can be accomplished with any mild oxidizing agent, such as Tollens' reagent or Benedict's reagent. With the latter, complexed copper(II) ions are reduced to copper(I) ions that form a brick-red precipitate [copper(I) oxide; Figure **15.5.1**].



Any carbohydrate capable of reducing either Tollens' or Benedict's reagents without first undergoing hydrolysis is said to be a reducing sugar. Because both the Tollens' and Benedict's reagents are basic solutions, ketoses (such as fructose) also give positive tests due to an equilibrium that exists between ketoses and aldoses in a reaction known as *tautomerism*.




Figure **15.5.1**: Benedict's Test. Benedict's test was performed on three carbohydrates, depicted from left to right: fructose, glucose, and sucrose. The solution containing sucrose remains blue because sucrose is a nonreducing sugar.

These reactions have been used as simple and rapid diagnostic tests for the presence of glucose in blood or urine. For example, Clinitest tablets, which are used to test for sugar in the urine, contain copper(II) ions and are based on Benedict's test. A green color indicates very little sugar, whereas a brick-red color indicates sugar in excess of 2 g/100 mL of urine.

Summary

Monosaccharides are crystalline solids at room temperature and quite soluble in water. Monosaccharides are reducing sugars; they reduce mild oxidizing agents, such as Tollens' or Benedict's reagents.

15.5: Properties of Monosaccharides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **16.5: Properties of Monosaccharides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


https://chem.libretexts.org/@go/page/387630

15.6: Disaccharides

Learning Objectives

- Identify the structures of sucrose, lactose, and maltose.
- Identify the monosaccharides that are needed to form sucrose, lactose, and maltose

Previously, you learned that monosaccharides can form cyclic structures by the reaction of the carbonyl group with an OH group. These cyclic molecules can in turn react with another alcohol. Disaccharides ($C_{12}H_{22}O_{11}$) are sugars composed of two monosaccharide units that are joined by a carbon–oxygen-carbon linkage known as a glycosidic linkage. This linkage is formed from the reaction of the anomeric carbon of one cyclic monosaccharide with the OH group of a second monosaccharide.



The disaccharides differ from one another in their monosaccharide constituents and in the specific type of glycosidic linkage connecting them. There are three common disaccharides: maltose, lactose, and sucrose. All three are white crystalline solids at room temperature and are soluble in water. We'll consider each sugar in more detail.

Maltose

Maltose occurs to a limited extent in sprouting grain. It is formed most often by the partial hydrolysis of starch and glycogen. In the manufacture of beer, maltose is liberated by the action of malt (germinating barley) on starch; for this reason, it is often referred to as *malt sugar*. Maltose is about 30% as sweet as sucrose. The human body is unable to metabolize maltose or any other disaccharide directly from the diet because the molecules are too large to pass through the cell membranes of the intestinal wall. Therefore, an ingested disaccharide must first be broken down by hydrolysis into its two constituent monosaccharide units.

In the body, such hydrolysis reactions are catalyzed by enzymes such as *maltase*. The same reactions can be carried out in the laboratory with dilute acid as a catalyst, although in that case the rate is much slower, and high temperatures are required. Whether it occurs in the body or a glass beaker, the hydrolysis of maltose produces two molecules of D-glucose.

$$\text{maltose} \xrightarrow{H^+ \text{ or maltase}} 2\ \text{D-glucose}$$

Maltose is a reducing sugar. Thus, its two glucose molecules must be linked in such a way as to leave one anomeric carbon that can open to form an aldehyde group. The glucose units in maltose are joined in a *head-to-tail* fashion through an α-linkage from the first carbon atom of one glucose molecule to the fourth carbon atom of the second glucose molecule (that is, an α-1,4-glycosidic linkage; see Figure **15.6.1**). The bond from the anomeric carbon of the first monosaccharide unit is directed downward, which is why this is known as an α-glycosidic linkage. The OH group on the anomeric carbon of the second glucose can be in either the α or the β position, as shown in Figure **15.6.1**.





Figure **15.6.1**: An Equilibrium Mixture of Maltose Isomers

Lactose

Lactose is known as *milk sugar* because it occurs in the milk of humans, cows, and other mammals. In fact, the natural synthesis of lactose occurs only in mammary tissue, whereas most other carbohydrates are plant products. Human milk contains about 7.5% lactose, and cow's milk contains about 4.5%. This sugar is one of the lowest ranking in terms of sweetness, being about one-sixth as sweet as sucrose. Lactose is produced commercially from whey, a by-product in the manufacture of cheese. It is important as an infant food and in the production of penicillin.


Lactose is a reducing sugar composed of one molecule of D-galactose and one molecule of D-glucose joined by a β-1,4-glycosidic bond (the bond from the anomeric carbon of the first monosaccharide unit being directed upward). The two monosaccharides are obtained from lactose by acid hydrolysis or the catalytic action of the enzyme *lactase*:



Many adults and some children suffer from a deficiency of lactase. These individuals are said to be lactose intolerant because they cannot digest the lactose found in milk. A more serious problem is the genetic disease galactosemia, which results from the absence of an enzyme needed to convert galactose to glucose. Certain bacteria can metabolize lactose, forming lactic acid as one of the products. This reaction is responsible for the "souring" of milk.

Example 15.6.1

For this trisaccharide, indicate whether each glycosidic linkage is α or β.




Solution

The glycosidic linkage between sugars 1 and 2 is β because the bond is directed up from the anomeric carbon. The glycosidic linkage between sugars 2 and 3 is α because the bond is directed down from the anomeric carbon.

? Exercise 15.6.1

For this trisaccharide, indicate whether each glycosidic linkage is α or β.



To Your Health: Lactose Intolerance and Galactosemia

Lactose makes up about 40% of an infant's diet during the first year of life. Infants and small children have one form of the enzyme lactase in their small intestines and can digest the sugar easily; however, adults usually have a less active form of the enzyme, and about 70% of the world's adult population has some deficiency in its production. As a result, many adults experience a reduction in the ability to hydrolyze lactose to galactose and glucose in their small intestine. For some people the inability to synthesize sufficient enzyme increases with age. Up to 20% of the US population suffers some degree of lactose intolerance.

In people with lactose intolerance, some of the unhydrolyzed lactose passes into the colon, where it tends to draw water from the interstitial fluid into the intestinal lumen by osmosis. At the same time, intestinal bacteria may act on the lactose to produce organic acids and gases. The buildup of water and bacterial decay products leads to abdominal distention, cramps, and diarrhea, which are symptoms of the condition.


The symptoms disappear if milk or other sources of lactose are excluded from the diet or consumed only sparingly. Alternatively, many food stores now carry special brands of milk that have been pretreated with lactase to hydrolyze the lactose. Cooking or fermenting milk causes at least partial hydrolysis of the lactose, so some people with lactose intolerance are still able to enjoy cheese, yogurt, or cooked foods containing milk. The most common treatment for lactose intolerance, however, is the use of lactase preparations (e.g., Lactaid), which are available in liquid and tablet form at drugstores and grocery stores. These are taken orally with dairy foods—or may be added to them directly—to assist in their digestion.

Galactosemia is a condition in which one of the enzymes needed to convert galactose to glucose is missing. Consequently, the blood galactose level is markedly elevated, and galactose is found in the urine. An infant with galactosemia experiences a lack of appetite, weight loss, diarrhea, and jaundice. The disease may result in impaired liver function, cataracts, mental retardation, and even death. If galactosemia is recognized in early infancy, its effects can be prevented by the exclusion of milk and all other sources of galactose from the diet. As a child with galactosemia grows older, he or she usually develops an alternate pathway for metabolizing galactose, so the need to restrict milk is not permanent. The incidence of galactosemia in the United States is 1 in every 65,000 newborn babies.

Sucrose

Sucrose, probably the largest-selling pure organic compound in the world, is known as *beet sugar*, *cane sugar*, *table sugar*, or simply *sugar*. Most of the sucrose sold commercially is obtained from sugar cane and sugar beets (whose juices are 14%–20% sucrose) by evaporation of the water and recrystallization. The dark brown liquid that remains after the recrystallization of sugar is sold as molasses.

The sucrose molecule is unique among the common disaccharides in having an α-1,β-2-glycosidic (head-to-head) linkage. Because this glycosidic linkage is formed by the OH group on the anomeric carbon of α-D-glucose and the OH group on the anomeric carbon of β-D-fructose, it ties up the anomeric carbons of both glucose and fructose.



This linkage gives sucrose certain properties that are quite different from those of maltose and lactose. As long as the sucrose molecule remains intact, neither monosaccharide "uncyclizes" to form an open-chain structure. Thus, sucrose is incapable of mutarotation and exists in only one form both in the solid state and in solution. In addition, sucrose does not undergo reactions that are typical of aldehydes and ketones. Therefore, sucrose is a nonreducing sugar.

The hydrolysis of sucrose in dilute acid or through the action of the enzyme *sucrase* (also known as invertase) gives an equimolar mixture of glucose and fructose. This 1:1 mixture is referred to as *invert sugar* because it rotates plane-polarized light in the opposite direction than sucrose. The hydrolysis reaction has several practical applications. Sucrose readily recrystallizes from a solution, but invert sugar has a much greater tendency to remain in solution. In the manufacture of jelly and candy and in the canning of fruit, the recrystallization of sugar is undesirable. Therefore, conditions leading to the hydrolysis of sucrose are employed in these processes. Moreover, because fructose is sweeter than sucrose, the hydrolysis adds to the sweetening effect. Bees carry out this reaction when they make honey.

The average American consumes more than 100 lb of sucrose every year. About two-thirds of this amount is ingested in soft drinks, presweetened cereals, and other highly processed foods. The widespread use of sucrose is a contributing factor to obesity and tooth decay. Carbohydrates such as sucrose, are converted to fat when the caloric intake exceeds the body's requirements, and sucrose causes tooth decay by promoting the formation of plaque that sticks to teeth.

Summary

Maltose is composed of two molecules of glucose joined by an α-1,4-glycosidic linkage. It is a reducing sugar that is found in sprouting grain. Lactose is composed of a molecule of galactose joined to a molecule of glucose by a β-1,4-glycosidic linkage. It is a reducing sugar that is found in milk. Sucrose is composed of a molecule of glucose joined to a molecule of fructose by an α-1,β-2-glycosidic linkage. It is a nonreducing sugar that is found in sugar cane and sugar beets

15.6: Disaccharides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

15.7: Polysaccharides

Learning Objectives

- To compare and contrast the structures and uses of starch, glycogen, and cellulose.

The polysaccharides are the most abundant carbohydrates in nature and serve a variety of functions, such as energy storage or as components of plant cell walls. Polysaccharides are very large polymers composed of tens to thousands of monosaccharides joined together by glycosidic linkages. The three most abundant polysaccharides are starch, glycogen, and cellulose. These three are referred to as *homopolymers* because each yields only one type of monosaccharide (glucose) after complete hydrolysis. *Heteropolymers* may contain sugar acids, amino sugars, or noncarbohydrate substances in addition to monosaccharides. Heteropolymers are common in nature (gums, pectins, and other substances) but will not be discussed further in this textbook. The polysaccharides are nonreducing carbohydrates, are not sweet tasting, and do not undergo mutarotation.

Starch

Starch is the most important source of carbohydrates in the human diet and accounts for more than 50% of our carbohydrate intake. It occurs in plants in the form of granules, and these are particularly abundant in seeds (especially the cereal grains) and tubers, where they serve as a storage form of carbohydrates. The breakdown of starch to glucose nourishes the plant during periods of reduced photosynthetic activity. We often think of potatoes as a "starchy" food, yet other plants contain a much greater percentage of starch (potatoes 15%, wheat 55%, corn 65%, and rice 75%). Commercial starch is a white powder.

Starch is a mixture of two polymers: amylose and amylopectin. Natural starches consist of about 10%–30% amylose and 70%–90% amylopectin. Amylose is a linear polysaccharide composed entirely of D-glucose units joined by the α-1,4-glycosidic linkages we saw in maltose (part (a) of Figure **15.7.1**). Experimental evidence indicates that amylose is not a straight chain of glucose units but instead is coiled like a spring, with six glucose monomers per turn (part (b) of Figure **15.7.1**). When coiled in this fashion, amylose has just enough room in its core to accommodate an iodine molecule. The characteristic blue-violet color that appears when starch is treated with iodine is due to the formation of the amylose-iodine complex. This color test is sensitive enough to detect even minute amounts of starch in solution.




Figure **15.7.1**: Amylose. (a) Amylose is a linear chain of α-D-glucose units joined together by α-1,4-glycosidic bonds. (b) Because of hydrogen bonding, amylose acquires a spiral structure that contains six glucose units per turn.

Amylopectin is a branched-chain polysaccharide composed of glucose units linked primarily by α-1,4-glycosidic bonds but with occasional α-1,6-glycosidic bonds, which are responsible for the branching. A molecule of amylopectin may contain many thousands of glucose units with branch points occurring about every 25–30 units (Figure **15.7.2**). The helical structure of amylopectin is disrupted by the branching of the chain, so instead of the deep blue-violet color amylose gives with iodine, amylopectin produces a less intense reddish brown.




Figure **15.7.2**: Representation of the Branching in Amylopectin and Glycogen. Both amylopectin and glycogen contain branch points that are linked through α-1,6-linkages. These branch points occur more often in glycogen.

Dextrins are glucose polysaccharides of intermediate size. The shine and stiffness imparted to clothing by starch are due to the presence of dextrins formed when clothing is ironed. Because of their characteristic stickiness with wetting, dextrins are used as adhesives on stamps, envelopes, and labels; as binders to hold pills and tablets together; and as pastes. Dextrins are more easily digested than starch and are therefore used extensively in the commercial preparation of infant foods.

The complete hydrolysis of starch yields, in successive stages, glucose:

$$\text{starch} \rightarrow \text{dextrins} \rightarrow \text{maltose} \rightarrow \text{glucose}$$

In the human body, several enzymes known collectively as amylases degrade starch sequentially into usable glucose units.

Glycogen

Glycogen is the energy reserve carbohydrate of animals. Practically all mammalian cells contain some stored carbohydrates in the form of glycogen, but it is especially abundant in the liver (4%–8% by weight of tissue) and in skeletal muscle cells (0.5%–1.0%). Like starch in plants, glycogen is found as granules in liver and muscle cells. When fasting, animals draw on these glycogen reserves during the first day without food to obtain the glucose needed to maintain metabolic balance.

Glycogen is structurally quite similar to amylopectin, although glycogen is more highly branched (8–12 glucose units between branches) and the branches are shorter. When treated with iodine, glycogen gives a reddish brown color. Glycogen can be broken down into its D-glucose subunits by acid hydrolysis or by the same enzymes that catalyze the breakdown of starch. In animals, the enzyme phosphorylase catalyzes the breakdown of glycogen to phosphate esters of glucose.

> *About 70% of the total glycogen in the body is stored in muscle cells. Although the percentage of glycogen (by weight) is higher in the liver, the much greater mass of skeletal muscle stores a greater total amount of glycogen.*

Cellulose

Cellulose, a fibrous carbohydrate found in all plants, is the structural component of plant cell walls. Because the earth is covered with vegetation, cellulose is the most abundant of all carbohydrates, accounting for over 50% of all the carbon found in the vegetable kingdom. Cotton fibrils and filter paper are almost entirely cellulose (about 95%), wood is about 50% cellulose, and the dry weight of leaves is about 10%–20% cellulose. The largest use of cellulose is in the manufacture of paper and paper products. Although the use of noncellulose synthetic fibers is increasing, rayon (made from cellulose) and cotton still account for over 70% of textile production.

Like amylose, cellulose is a linear polymer of glucose. It differs, however, in that the glucose units are joined by β-1,4-glycosidic linkages, producing a more extended structure than amylose (part (a) of Figure **15.7.3**). This extreme linearity allows a great deal of hydrogen bonding between OH groups on adjacent chains, causing them to pack closely into fibers (part (b) of Figure **15.7.3**). As a result, cellulose exhibits little interaction with water or any other solvent. Cotton and wood, for example, are completely insoluble in water and have considerable mechanical strength. Because cellulose does not have a helical structure, it does not bind to iodine to form a colored product.




Figure **15.7.3**: Cellulose. (a) There is extensive hydrogen bonding in the structure of cellulose. (b) In this electron micrograph of the cell wall of an alga, the wall consists of successive layers of cellulose fibers in parallel arrangement.

Cellulose yields D-glucose after complete acid hydrolysis, yet humans are unable to metabolize cellulose as a source of glucose. Our digestive juices lack enzymes that can hydrolyze the β-glycosidic linkages found in cellulose, so although we can eat potatoes, we cannot eat grass. However, certain microorganisms can digest cellulose because they make the enzyme cellulase, which catalyzes the hydrolysis of cellulose. The presence of these microorganisms in the digestive tracts of herbivorous animals (such as cows, horses, and sheep) allows these animals to degrade the cellulose from plant material into glucose for energy. Termites also contain cellulase-secreting microorganisms and thus can subsist on a wood diet. This example once again demonstrates the extreme stereospecificity of biochemical processes.

Career Focus: Certified Diabetes Educator

Certified diabetes educators come from a variety of health professions, such as nursing and dietetics, and specialize in the education and treatment of patients with diabetes. A diabetes educator will work with patients to manage their diabetes. This involves teaching the patient to monitor blood sugar levels, make good food choices, develop and maintain an exercise program, and take medication, if required.



A certified diabetes educator at Naval Medical Center Portsmouth (left) and a registered dietician at the medical center (center), provide nutritional information to a diabetes patient and her mother at the Diabetes Boot Camp.

> Diabetes educators also work with hospital or nursing home staff to improve the care of diabetic patients. Educators must be willing to spend time attending meetings and reading the current literature to maintain their knowledge of diabetes medications, nutrition, and blood monitoring devices so that they can pass this information to their patients.

Summary

Starch is a storage form of energy in plants. It contains two polymers composed of glucose units: amylose (linear) and amylopectin (branched). Glycogen is a storage form of energy in animals. It is a branched polymer composed of glucose units. It is more highly branched than amylopectin. Cellulose is a structural polymer of glucose units found in plants. It is a linear polymer with the glucose units linked through β-1,4-glycosidic bonds.

15.7: Polysaccharides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **16.7: Polysaccharides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

15.8: Diabetes

Skills to Develop

- Summarize the long-term health implications and the dietary approaches to living with Type 1 and Type 2 diabetes.

Diabetes is one of the top three diseases in America. It affects millions of people and causes tens of thousands of deaths each year. Diabetes is a metabolic disease of insulin deficiency and glucose over-sufficiency. Like other diseases, genetics, nutrition, environment, and lifestyle are all involved in determining a person's risk for developing diabetes. One sure way to decrease your chances of getting diabetes is to maintain an optimal body weight by adhering to a diet that is balanced in carbohydrate, fat, and protein intake. There are three different types of diabetes: Type 1 diabetes, Type 2 diabetes, and gestational diabetes.

Type 1 Diabetes

Type 1 diabetes is a metabolic disease in which insulin-secreting cells in the pancreas are killed by an abnormal response of the immune system, causing a lack of insulin in the body. Its onset typically occurs before the age of thirty. The only way to prevent the deadly symptoms of this disease is to inject insulin under the skin. Before this treatment was discovered, people with Type 1 diabetes died rapidly after disease onset. Death was the result of extremely high blood-glucose levels affecting brain function and leading to coma and death. Up until 1921, patients with Type 1 diabetes, the majority of them children, spent their last days in a ward where they lapsed into a coma awaiting death. One of the most inspiring acts in medical history is that of the scientists who discovered, isolated, and purified insulin and then went on to find out that it relieved the symptoms of Type 1 diabetes, first in dogs and then in humans. Frederick Banting, Charles Best, and James Collip went into a hospital ward in Toronto, Canada and injected comatose children with insulin. Before they completed their rounds children were already awakening to the cheers of their families.

A person with Type 1 diabetes usually has a rapid onset of symptoms that include hunger, excessive thirst and urination, and rapid weight loss. Because the main function of glucose is to provide energy for the body, when insulin is no longer present there is no message sent to cells to take up glucose from the blood. Instead, cells use fat and proteins to make energy, resulting in weight loss. If Type 1 diabetes goes untreated individuals with the disease will develop a life-threatening condition called ketoacidosis. This condition occurs when the body uses fats and not glucose to make energy, resulting in a build-up of ketone bodies in the blood. It is a severe form of ketosis with symptoms of vomiting, dehydration, rapid breathing, and confusion and eventually coma and death. Upon insulin injection these severe symptoms are treated and death is avoided. Unfortunately, while insulin injection prevents death, it is not considered a cure. People who have this disease must adhere to a strict diet to prevent the development of serious complications. Type 1 diabetics are advised to consume a diet low in the types of carbohydrates that rapidly spike glucose levels (high-GI foods), to count the carbohydrates they eat, to consume healthy-carbohydrate foods, and to eat small meals frequently. These guidelines are aimed at preventing large fluctuations in blood glucose. Frequent exercise also helps manage blood-glucose levels. Type 1 diabetes accounts for between 5 and 10 percent of diabetes cases.

Type 2 Diabetes

The other 90 to 95 percent of diabetes cases are Type 2 diabetes, which is defined as a metabolic disease of insulin insufficiency, but it is also caused by muscle, liver, and fat cells no longer responding to the insulin in the body (Figure **15.8.1**). In brief, cells in the body have become resistant to insulin and no longer receive the full physiological message of insulin to take up glucose from the blood. Thus, similar to patients with Type 1 diabetes, those with Type 2 diabetes also have high blood-glucose levels.

For Type 2 diabetics, the onset of symptoms is more gradual and less noticeable than for Type 1 diabetics. The symptoms are increased thirst and urination, unexplained weight loss, and hunger. The first stage of Type 2 diabetes is characterized by high glucose and insulin levels. This is because the insulin-secreting cells in the pancreas attempt to compensate for insulin resistance by making more insulin. In the second stage of Type 2 diabetes, the insulin-secreting cells in the pancreas become exhausted and die. At this point, Type 2 diabetics also have to be treated with insulin injections. Healthcare providers is to prevent the second stage from happening. As with Type 1 diabetes, chronically high-glucose levels cause big detriments to health over time, so another goal for patients with Type 2 diabetes is to properly manage their blood-glucose levels. The front-line approach for treating Type 2 diabetes includes eating a healthy diet and increasing physical activity.




Figure* 15.8.1*: *Type 2 diabetes is a metabolic disease characterized by high blood-glucose levels. © Shutterstock*

The Centers for Disease Control Prevention (CDC) estimates that as of 2010, 25.8 million Americans have diabetes, which is 8.3 percent of the population. In 2007 the cost of diabetes to the United States was estimated at $174 billion. The incidence of Type 2 diabetes has more than doubled in America in the past thirty years and the rise is partly attributed to the increase in obesity in this country. Genetics, environment, nutrition, and lifestyle all play a role in determining a person's risk for Type 2 diabetes.

Video 15.8.1: Do You Have High Blood Sugar?

A more in-depth view of blood sugar and your health.(click to see video)

Summary

Diabetes is a disease of insulin deficiency and glucose oversufficiency. Like other diseases, genetics, nutrition, environment, and lifestyle are all involved in determining a person's risk for developing diabetes. Type 1 diabetes was once a death sentence, but now can be treated with insulin injections. However, insulin injections do not cure the disease, and diabetics can suffer many disease complications. Diabetes complications can be relieved by strictly managing blood-glucose levels, adhering to a healthy diet, and increasing physical activity. The incidence of Type 2 diabetes has more than doubled in America in the past thirty years and the rise is partly attributed to the increase in obesity. The front-line approach for treating Type 2 diabetes includes eating a healthy diet and increasing physical activity.

Discussion Starters

1. If you owned a grocery store what are some practices you could introduce to combat the epidemic of Type 2 diabetes in this country?
2. What are some options for you to intervene in your lifestyle and decrease your risk for Type 2 diabetes?

15.8: Diabetes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

15.9: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Carbohydrates, a large group of biological compounds containing carbon, hydrogen, and oxygen atoms, include sugars, starch, glycogen, and cellulose. All carbohydrates contain alcohol functional groups, and either an aldehyde or a ketone group (or a functional group that can be converted to an aldehyde or ketone). The simplest carbohydrates are **monosaccharides**. Those with two monosaccharide units are **disaccharides**, and those with many monosaccharide units are **polysaccharides**. Most sugars are either monosaccharides or disaccharides. Cellulose, glycogen, and starch are polysaccharides.

Many carbohydrates exist as **stereoisomers**, in which the three-dimensional spatial arrangement of the atoms in space is the only difference between the isomers. These particular stereoisomers contain at least one **chiral carbon**, a carbon atom that has four different groups bonded to it. A molecule containing a chiral carbon is nonsuperimposable on its mirror image, and two molecules that are nonsuperimposable mirror images of each other are a special type of stereoisomer called **enantiomers**. Enantiomers have the same physical properties, such as melting point, but differ in the direction they rotate polarized light.

A sugar is designated as being a D sugar or an L sugar according to how, in a Fischer projection of the molecule, the hydrogen atom and OH group are attached to the *penultimate* carbon atom, which is the carbon atom immediately before the terminal alcohol carbon atom. If the structure at this carbon atom is the same as that of D-glyceraldehyde (OH to the right), the sugar is a **D sugar**; if the configuration is the same as that of L-glyceraldehyde (OH to the left), the sugar is an **L sugar**.

Monosaccharides of five or more carbons atoms readily form cyclic structures when the carbonyl carbon atom reacts with an OH group on a carbon atom three or four carbon atoms distant. Consequently, glucose in solution exists as an equilibrium mixture of three forms, two of them cyclic (α- and β-) and one open chain. In Haworth projections, the *alpha* form is drawn with the OH group on the "former" carbonyl carbon atom (**anomeric carbon**) pointing downward; the *beta* form, with the OH group pointing upward; these two compounds are stereoisomers and are given the more specific term of **anomers**. Any solid sugar can be all alpha or all beta. Once the sample is dissolved in water, however, the ring opens up into the open-chain structure and then closes to form either the α- or the β-anomer. These interconversions occur back and forth until a dynamic equilibrium mixture is achieved in a process called **mutarotation**.

The carbonyl group present in monosaccharides is easily oxidized by Tollens' or Benedict's reagents (as well as others). Any mono- or disaccharide containing a free anomeric carbon is a **reducing sugar**. The disaccharide *maltose* contains two glucose units joined in an α-1,4-glycosidic linkage. The disaccharide *lactose* contains a galactose unit and a glucose unit joined by a β-1,4-glycosidic linkage. Both maltose and lactose contain a free anomeric carbon that can convert to an aldehyde functional group, so they are reducing sugars; they also undergo mutarotation. Many adults, and some children, have a deficiency of the enzyme lactase (which is needed to break down lactose) and are said to be **lactose intolerant**. A more serious problem is the genetic disease **galactosemia**, which results from the absence of an enzyme needed to convert galactose to glucose.

The disaccharide *sucrose* (table sugar) consists of a glucose unit and a fructose unit joined by a glycosidic linkage. The linkage is designated as an α-1,β-2-glycosidic linkage because it involves the OH group on the first carbon atom of glucose and the OH group on the second carbon atom of fructose. Sucrose is not a reducing sugar because it has no anomeric carbon that can reform a carbonyl group, and it cannot undergo mutarotation because of the restrictions imposed by this linkage.

Starch, the principal carbohydrate of plants, is composed of the polysaccharides **amylose** (10%–30%) and **amylopectin** (70%–90%). When ingested by humans and other animals, starch is hydrolyzed to glucose and becomes the body's energy source. *Glycogen* is the polysaccharide animals use to store excess carbohydrates from their diets. Similar in structure to amylopectin, glycogen is hydrolyzed to glucose whenever an animal needs energy for a metabolic process. The polysaccharide *cellulose* provides structure for plant cells. It is a linear polymer of glucose units joined by β-1,4-glycosidic linkages. It is indigestible in the human body but digestible by many microorganisms, including microorganisms found in the digestive tracts of many herbivores.

15.9: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **16.S: Carbohydrates (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

16: Lipids

The lipids are a large and diverse group of naturally occurring organic compounds that are related by their solubility in nonpolar organic solvents (e.g., ether, chloroform, acetone and benzene) and general insolubility in water. There is great structural variety among the lipids, as will be demonstrated in the following sections.

16.1: Introduction to Lipids
16.2: Fatty Acids
16.3: Fats and Oils
16.4: Membranes and Membrane Lipids
16.5: Steroids
16.6: Chapter Summary

16: Lipids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

16.1: Introduction to Lipids

On July 11, 2003, the Food and Drug Administration amended its food labeling regulations to require that manufacturers list the amount of *trans* fatty acids on Nutrition Facts labels of foods and dietary supplements, effective January 1, 2006. This amendment was a response to published studies demonstrating a link between the consumption of *trans* fatty acids and an increased risk of heart disease. *Trans* fatty acids are produced in the conversion of liquid oils to solid fats, as in the creation of many commercial margarines and shortenings. They have been shown to increase the levels of low-density lipoproteins (LDLs)—complexes that are often referred to as bad cholesterol—in the blood. In this chapter, you will learn about fatty acids and what is meant by a *trans* fatty acid, as well as the difference between fats and oils. You will also learn what cholesterol is and why it is an important molecule in the human body.

Fats and oils, found in many of the foods we eat, belong to a class of biomolecules known as lipids. Gram for gram, they pack more than twice the caloric content of carbohydrates: the oxidation of fats and oils supplies about 9 kcal of energy for every gram oxidized, whereas the oxidation of carbohydrates supplies only 4 kcal/g. Although the high caloric content of fats may be bad news for the dieter, it says something about the efficiency of nature's designs. Our bodies use carbohydrates, primarily in the form of glucose, for our *immediate* energy needs. Our capacity for storing carbohydrates for later use is limited to tucking away a bit of glycogen in the liver or in muscle tissue. We store our *reserve* energy in lipid form, which requires far less space than the same amount of energy stored in carbohydrate form. Lipids have other biological functions besides energy storage. They are a major component of the membranes of the 10 trillion cells in our bodies. They serve as protective padding and insulation for vital organs. Furthermore, without lipids in our diets, we would be deficient in the fat-soluble vitamins A, D, E, and K.

Lipids are not defined by the presence of specific functional groups, as carbohydrates are, but by a physical property—solubility. Compounds isolated from body tissues are classified as lipids if they are more soluble in organic solvents, such as dichloromethane, than in water. By this criterion, the lipid category includes not only fats and oils, which are esters of the trihydroxy alcohol glycerol and fatty acids, but also compounds that incorporate functional groups derived from phosphoric acid, carbohydrates, or amino alcohols, as well as steroid compounds such as cholesterol (Figure 16.1.1 presents one scheme for classifying the various kinds of lipids). We will discuss the various kinds of lipids by considering one subclass at a time and pointing out structural similarities and differences as we go.




Figure 16.1.1: Lipid Organization Based on Structural Relationships

Lipid categorized into fatty acids and steroids. Fatty acids are further separated into triglycerides, phospho-glycerides, waxes, and sphingolipids. Sphingolipids are separated into sphingo-myelins and glycolipids. Glycolipids are separated into cerebrosides and gangliosides.

16.1: Introduction to Lipids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.0: Prelude to Lipids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


https://chem.libretexts.org/@go/page/387636

16.2: Fatty Acids

Learning Objectives

- To recognize the structures of common fatty acids and classify them as saturated, monounsaturated, or polyunsaturated.

Fatty acids are carboxylic acids that are structural components of fats, oils, and all other categories of lipids, except steroids. More than 70 have been identified in nature. They usually contain an even number of carbon atoms (typically 12–20), are generally unbranched, and can be classified by the presence and number of carbon-to-carbon double bonds. Thus, saturated fatty acids contain no carbon-to-carbon double bonds, monounsaturated fatty acids contain one carbon-to-carbon double bond, and polyunsaturated fatty acids contain two or more carbon-to-carbon double bonds.



Cis fatty acids have two H atoms on the same side of the plane of the fatty acid while trans has the two H atoms on opposite sides.

Table **16.2.1** lists some common fatty acids and one important source for each. The atoms or groups around the double bonds in unsaturated fatty acids can be arranged in either the cis or trans isomeric form. Naturally occurring fatty acids are generally in the cis configuration.

Table **16.2.1**: Some Common Fatty Acids Found in Natural Fats


Name	Abbreviated Structural Formula	Condensed Structural Formula	Melting Point (°C)	Source
lauric acid	$C_{11}H_{23}COOH$	$CH_3(CH_2)_{10}COOH$	44	palm kernel oil
myristic acid	$C_{13}H_{27}COOH$	$CH_3(CH_2)_{12}COOH$	58	oil of nutmeg
palmitic acid	$C_{15}H_{31}COOH$	$CH_3(CH_2)_{14}COOH$	63	palm oil
palmitoleic acid	$C_{15}H_{29}COOH$	$CH_3(CH_2)_5CH=CH(CH_2)_7COOH$	0.5	macadamia oil
stearic acid	$C_{17}H_{35}COOH$	$CH_3(CH_2)_{16}COOH$	70	cocoa butter
oleic acid	$C_{17}H_{33}COOH$	$CH_3(CH_2)_7CH=CH(CH_2)_7COOH$	16	olive oil
linoleic acid	$C_{17}H_{31}COOH$	$CH_3(CH_2)_3(CH_2CH=CH)_2(CH_2)_7COOH$	−5	canola oil
α-linolenic acid	$C_{17}H_{29}COOH$	$CH_3(CH_2CH=CH)_3(CH_2)_7COOH$	−11	flaxseed
arachidonic acid	$C_{19}H_{31}COOH$	$CH_3(CH_2)_4(CH_2CH=CH)_4(CH_2)_2COOH$	−50	liver

Two polyunsaturated fatty acids—linoleic and α-linolenic acids—are termed essential fatty acids because humans must obtain them from their diets. Both substances are required for normal growth and development, but the human body does not synthesize them. The body uses linoleic acid to synthesize many of the other unsaturated fatty acids, such as arachidonic acid, a precursor for the synthesis of prostaglandins. In addition, the essential fatty acids are necessary for the efficient transport and metabolism of cholesterol. The average daily diet should contain about 4–6 g of the essential fatty acids.

To Your Health: Prostaglandins

Prostaglandins are chemical messengers synthesized in the cells in which their physiological activity is expressed. They are unsaturated fatty acids containing 20 carbon atoms and are synthesized from arachidonic acid—a polyunsaturated fatty acid—when needed by a particular cell. They are called *prostaglandins* because they were originally isolated from semen found in the prostate gland. It is now known that they are synthesized in nearly all mammalian tissues and affect almost all organs in the body. The five major classes of prostaglandins are designated as PGA, PGB, PGE, PGF, and PGI. Subscripts are attached at the end of these abbreviations to denote the number of double bonds outside the five-carbon ring in a given prostaglandin.

The prostaglandins are among the most potent biological substances known. Slight structural differences give them highly distinct biological effects; however, all prostaglandins exhibit some ability to induce smooth muscle contraction, lower blood pressure, and contribute to the inflammatory response. Aspirin and other nonsteroidal anti-inflammatory agents, such as ibuprofen, obstruct the synthesis of prostaglandins by inhibiting cyclooxygenase, the enzyme needed for the initial step in the conversion of arachidonic acid to prostaglandins.



Their wide range of physiological activity has led to the synthesis of hundreds of prostaglandins and their analogs. Derivatives of PGE_2 are now used in the United States to induce labor. Other prostaglandins have been employed clinically to lower or increase blood pressure, inhibit stomach secretions, relieve nasal congestion, relieve asthma, and prevent the formation of blood clots, which are associated with heart attacks and strokes.

Although we often draw the carbon atoms in a straight line, they actually have more of a zigzag configuration (Figure **16.2.2*a***). Viewed as a whole, however, the saturated fatty acid molecule is relatively straight (Figure **16.2.2*b***). Such molecules pack closely together into a crystal lattice, maximizing the strength of dispersion forces and causing fatty acids and the fats derived from them to have relatively high melting points. In contrast, each *cis* carbon-to-carbon double bond in an unsaturated fatty acid produces a pronounced bend in the molecule, so that these molecules do not stack neatly. As a result, the intermolecular attractions of unsaturated fatty acids (and unsaturated fats) are weaker, causing these substances to have lower melting points. Most are liquids at room temperature.





Figure **16.2.2**: The Structure of Saturated Fatty Acids. (a) There is a zigzag pattern formed by the carbon-to-carbon single bonds in the ball-and-stick model of a palmitic acid molecule. (b) A space-filling model of palmitic acid shows the overall straightness of a saturated fatty acid molecule.

Waxes are esters formed from long-chain fatty acids and long-chain alcohols. Most natural waxes are mixtures of such esters. Plant waxes on the surfaces of leaves, stems, flowers, and fruits protect the plant from dehydration and invasion by harmful microorganisms. Carnauba wax, used extensively in floor waxes, automobile waxes, and furniture polish, is largely myricyl cerotate, obtained from the leaves of certain Brazilian palm trees. Animals also produce waxes that serve as protective coatings, keeping the surfaces of feathers, skin, and hair pliable and water repellent. In fact, if the waxy coating on the feathers of a water bird is dissolved as a result of the bird swimming in an oil slick, the feathers become wet and heavy, and the bird, unable to maintain its buoyancy, drowns.



Myricyl cerotate
(found in carnauba wax)

Summary

Fatty acids are carboxylic acids that are the structural components of many lipids. They may be saturated or unsaturated. Most fatty acids are unbranched and contain an even number of carbon atoms. Unsaturated fatty acids have lower melting points than saturated fatty acids containing the same number of carbon atoms.

16.2: Fatty Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.1: Fatty Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

16.3: Fats and Oils


Learning Objectives

- Explain why fats and oils are referred to as triglycerides.
- Explain how the fatty acid composition of the triglycerides determines whether a substance is a fat or oil.
- Describe the importance of key reactions of triglycerides, such as hydrolysis, hydrogenation, and oxidation.


Fats and oils are the most abundant lipids in nature. They provide energy for living organisms, insulate body organs, and transport fat-soluble vitamins through the blood.

Structures of Fats and Oils


Fats and oils are called triglycerides (or *triacylcylgerols*) because they are esters composed of three fatty acid units joined to *glycerol*, a trihydroxy alcohol:



If all three OH groups on the glycerol molecule are esterified with the same fatty acid, the resulting ester is called a *simple triglyceride*. Although simple triglycerides have been synthesized in the laboratory, they rarely occur in nature. Instead, a typical triglyceride obtained from naturally occurring fats and oils contains two or three different fatty acid components and is thus termed a *mixed triglyceride*.



Tristearin
a simple triglyceride



a mixed triglyceride

A triglyceride is called a fat if it is a solid at 25°C; it is called an oil if it is a liquid at that temperature. These differences in melting points reflect differences in the degree of unsaturation and number of carbon atoms in the constituent fatty acids. Triglycerides obtained from animal sources are usually solids, while those of plant origin are generally oils. Therefore, we commonly speak of animal fats and vegetable oils.

No single formula can be written to represent the naturally occurring fats and oils because they are highly complex mixtures of triglycerides in which many different fatty acids are represented. Table 16.3.1 shows the fatty acid compositions of some common

fats and oils. The composition of any given fat or oil can vary depending on the plant or animal species it comes from as well as on dietetic and climatic factors. To cite just one example, lard from corn-fed hogs is more highly saturated than lard from peanut-fed hogs. Palmitic acid is the most abundant of the saturated fatty acids, while oleic acid is the most abundant unsaturated fatty acid.

Table **16.3.1**: Average Fatty Acid Composition of Some Common Fats and Oils (%)*

	Lauric	Myristic	Palmitic	Stearic	Oleic	Linoleic	Linolenic
Fats							
butter (cow)	3	11	27	12	29	2	1
tallow		3	24	19	43	3	1
lard		2	26	14	44	10	
Oils							
canola oil			4	2	62	22	10
coconut oil†	47	18	9	3	6	2	
corn oil			11	2	28	58	1
olive oil			13	3	71	10	1
peanut oil			11	2	48	32	
soybean oil			11	4	24	54	7


*Totals less than 100% indicate the presence of fatty acids with fewer than 12 carbon atoms or more than 18 carbon atoms.

†Coconut oil is highly saturated. It contains an unusually high percentage of the low-melting C_8, C_{10}, and C_{12} saturated fatty acids.


Terms such as *saturated fat* or *unsaturated oil* are often used to describe the fats or oils obtained from foods. Saturated fats contain a high proportion of saturated fatty acids, while unsaturated oils contain a high proportion of unsaturated fatty acids. The high consumption of saturated fats is a factor, along with the high consumption of cholesterol, in increased risks of heart disease.

Physical Properties of Fats and Oils

Contrary to what you might expect, *pure* fats and oils are colorless, odorless, and tasteless. The characteristic colors, odors, and flavors that we associate with some of them are imparted by foreign substances that are lipid soluble and have been absorbed by these lipids. For example, the yellow color of butter is due to the presence of the pigment carotene; the taste of butter comes from two compounds—diacetyl and 3-hydroxy-2-butanone—produced by bacteria in the ripening cream from which the butter is made.



Diacetyl




3-hydroxy-2-butanone

Fats and oils are lighter than water, having densities of about 0.8 g/cm^3. They are poor conductors of heat and electricity and therefore serve as excellent insulators for the body, slowing the loss of heat through the skin.

Chemical Reactions of Fats and Oils


Fats and oils can participate in a variety of chemical reactions—for example, because triglycerides are esters, they can be hydrolyzed in the presence of an acid, a base, or specific enzymes known as lipases. The hydrolysis of fats and oils in the presence of a base is used to make soap and is called saponification. Today most soaps are prepared through the hydrolysis of triglycerides (often from tallow, coconut oil, or both) using water under high pressure and temperature [700 lb/in^2 (~50 atm or 5,000 kPa) and 200°C]. Sodium carbonate or sodium hydroxide is then used to convert the fatty acids to their sodium salts (soap molecules):



Looking Closer: Soaps


Ordinary soap is a mixture of the sodium salts of various fatty acids, produced in one of the oldest organic syntheses practiced by humans (second only to the fermentation of sugars to produce ethyl alcohol). Both the Phoenicians (600 BCE) and the Romans made soap from animal fat and wood ash. Even so, the widespread production of soap did not begin until the 1700s. Soap was traditionally made by treating molten lard or tallow with a slight excess of alkali in large open vats. The mixture was heated, and steam was bubbled through it. After saponification was completed, the soap was precipitated from the mixture by the addition of sodium chloride (NaCl), removed by filtration, and washed several times with water. It was then dissolved in water and reprecipitated by the addition of more NaCl. The glycerol produced in the reaction was also recovered from the aqueous wash solutions.

Pumice or sand is added to produce scouring soap, while ingredients such as perfumes or dyes are added to produce fragrant, colored soaps. Blowing air through molten soap produces a floating soap. Soft soaps, made with potassium salts, are more expensive but produce a finer lather and are more soluble. They are used in liquid soaps, shampoos, and shaving creams.



Dirt and grime usually adhere to skin, clothing, and other surfaces by combining with body oils, cooking fats, lubricating greases, and similar substances that act like glues. Because these substances are not miscible in water, washing with water alone does little to remove them. Soap removes them, however, because soap molecules have a dual nature. One end, called the *head*, carries an ionic charge (a carboxylate anion) and therefore dissolves in water; the other end, the *tail*, has a hydrocarbon structure and dissolves in oils. The hydrocarbon tails dissolve in the soil; the ionic heads remain in the aqueous phase, and the soap breaks the oil into tiny soap-enclosed droplets called *micelles*, which disperse throughout the solution. The droplets repel each other because of their charged surfaces and do not coalesce. With the oil no longer “gluing” the dirt to the soiled surface (skin, cloth, dish), the soap-enclosed dirt can easily be rinsed away.

The double bonds in fats and oils can undergo hydrogenation and also oxidation. The hydrogenation of vegetable oils to produce semisolid fats is an important process in the food industry. Chemically, it is essentially identical to the catalytic hydrogenation reaction described for alkenes.



In commercial processes, the number of double bonds that are hydrogenated is carefully controlled to produce fats with the desired consistency (soft and pliable). Inexpensive and abundant vegetable oils (canola, corn, soybean) are thus transformed into margarine and cooking fats. In the preparation of margarine, for example, partially hydrogenated oils are mixed with water, salt, and nonfat dry milk, along with flavoring agents, coloring agents, and vitamins A and D, which are added to approximate the look, taste, and nutrition of butter. (Preservatives and antioxidants are also added.) In most commercial peanut butter, the peanut oil has been partially hydrogenated to prevent it from separating out. Consumers could decrease the amount of saturated fat in their diet by using the original unprocessed oils on their foods, but most people would rather spread margarine on their toast than pour oil on it.


Many people have switched from butter to margarine or vegetable shortening because of concerns that saturated animal fats can raise blood cholesterol levels and result in clogged arteries. However, during the hydrogenation of vegetable oils, an isomerization reaction occurs that produces the *trans* fatty acids mentioned in the opening essay. However, studies have shown that *trans* fatty acids also raise cholesterol levels and increase the incidence of heart disease. *Trans* fatty acids do not have the bend in their structures, which occurs in *cis* fatty acids and thus pack closely together in the same way that the saturated fatty acids do. Consumers are now being advised to use polyunsaturated oils and soft or liquid margarine and reduce their total fat consumption to less than 30% of their total calorie intake each day.

Fats and oils that are in contact with moist air at room temperature eventually undergo oxidation and hydrolysis reactions that cause them to turn rancid, acquiring a characteristic disagreeable odor. One cause of the odor is the release of volatile fatty acids by hydrolysis of the ester bonds. Butter, for example, releases foul-smelling butyric, caprylic, and capric acids. Microorganisms present in the air furnish lipases that catalyze this process. Hydrolytic rancidity can easily be prevented by covering the fat or oil and keeping it in a refrigerator.

Another cause of volatile, odorous compounds is the oxidation of the unsaturated fatty acid components, particularly the readily oxidized structural unit



in polyunsaturated fatty acids, such as linoleic and linolenic acids. One particularly offensive product, formed by the oxidative cleavage of both double bonds in this unit, is a compound called *malonaldehyde*.



Malonaldehyde

Rancidity is a major concern of the food industry, which is why food chemists are always seeking new and better antioxidants, substances added in very small amounts (0.001%–0.01%) to prevent oxidation and thus suppress rancidity. Antioxidants are compounds whose affinity for oxygen is greater than that of the lipids in the food; thus they function by preferentially depleting the supply of oxygen absorbed into the product. Because vitamin E has antioxidant properties, it helps reduce damage to lipids in the body, particularly to unsaturated fatty acids found in cell membrane lipids.

Summary

Fats and oils are composed of molecules known as triglycerides, which are esters composed of three fatty acid units linked to glycerol. An increase in the percentage of shorter-chain fatty acids and/or unsaturated fatty acids lowers the melting point of a fat or oil. The hydrolysis of fats and oils in the presence of a base makes soap and is known as saponification. Double bonds present in unsaturated triglycerides can be hydrogenated to convert oils (liquid) into margarine (solid). The oxidation of fatty acids can form compounds with disagreeable odors. This oxidation can be minimized by the addition of antioxidants.

16.3: Fats and Oils is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.2: Fats and Oils** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

16.4: Membranes and Membrane Lipids

Learning Objectives

- Identify the distinguishing characteristics of membrane lipids.
- Describe membrane components and how they are arranged.

All living cells are surrounded by a cell membrane. Plant cells (Figure **16.4.1*A***) and animal cells (Figure **16.4.1*B***) contain a cell nucleus that is also surrounded by a membrane and holds the genetic information for the cell. Everything between the cell membrane and the nuclear membrane—including intracellular fluids and various subcellular components such as the mitochondria and ribosomes—is called the cytoplasm. The membranes of all cells have a fundamentally similar structure, but membrane function varies tremendously from one organism to another and even from one cell to another within a single organism. This diversity arises mainly from the presence of different proteins and lipids in the membrane.




Figure **16.4.1**: (A) An Idealized Plant Cell. Not all the structures shown here occur in every type of plant cell. (B) An Idealized Animal Cell. The structures shown here will seldom all be found in a single animal cell.

The lipids in cell membranes are highly polar but have dual characteristics: part of the lipid is ionic and therefore dissolves in water, whereas the rest has a hydrocarbon structure and therefore dissolves in nonpolar substances. Often, the ionic part is referred to as hydrophilic, meaning “water loving,” and the nonpolar part as hydrophobic, meaning “water fearing” (repelled by water). When allowed to float freely in water, polar lipids spontaneously cluster together in any one of three arrangements: micelles, monolayers, and bilayers (Figure **16.4.2**).




Figure **16.4.2**: Spontaneously Formed Polar Lipid Structures in Water: Monolayer, Micelle, and Bilayer

Micelles are aggregations in which the lipids’ hydrocarbon tails—being hydrophobic—are directed toward the center of the assemblage and away from the surrounding water while the hydrophilic heads are directed outward, in contact with the water. Each micelle may contain thousands of lipid molecules. Polar lipids may also form a monolayer, a layer one molecule thick on the surface of the water. The polar heads face into water, and the nonpolar tails stick up into the air. Bilayers are double layers of lipids arranged so that the hydrophobic tails are sandwiched between an inner surface and an outer surface consisting of hydrophilic heads. The hydrophilic heads are in contact with water on either side of the bilayer, whereas the tails, sequestered inside the bilayer, are prevented from having contact with the water. Bilayers like this make up every cell membrane (Figure **16.4.3**).




Figure **16.4.3**: Schematic Diagram of a Cell Membrane. The membrane enclosing a typical animal cell is a phospholipid bilayer with embedded cholesterol and protein molecules. Short oligosaccharide chains are attached to the outer surface.

In the bilayer interior, the hydrophobic tails (that is, the fatty acid portions of lipid molecules) interact by means of dispersion forces. The interactions are weakened by the presence of unsaturated fatty acids. As a result, the membrane components are free to mill about to some extent, and the membrane is described as fluid.

The lipids found in cell membranes can be categorized in various ways. Phospholipids are lipids containing phosphorus. Glycolipids are sugar-containing lipids. The latter are found exclusively on the outer surface of the cell membrane, acting as distinguishing surface markers for the cell and thus serving in cellular recognition and cell-to-cell communication. Sphingolipids

are phospholipids or glycolipids that contain the unsaturated amino alcohol sphingosine rather than glycerol. Diagrammatic structures of representative membrane lipids are presented in Figure **16.4.4**.




Figure **16.4.4**: Component Structures of Some Important Membrane Lipids

Phosphoglyceride are composed of alcohol, phosphate, glycerol and 2 fatty acids. A sphingomyelin contains phosphate, sphingosine, and a fatty acid. A cerbroside contains sugar, sphingosine and a fatty acid.

Phosphoglycerides (also known as glycerophospholipids) are the most abundant phospholipids in cell membranes. They consist of a glycerol unit with fatty acids attached to the first two carbon atoms, while a phosphoric acid unit, esterified with an alcohol molecule (usually an amino alcohol, as in part (a) of Figure **16.4.5**) is attached to the third carbon atom of glycerol (part (b) of Figure **16.4.5**). Notice that the phosphoglyceride molecule is identical to a triglyceride up to the phosphoric acid unit (part (b) of Figure **16.4.5**).





Figure **16.4.5**: Phosphoglycerides. (a) Amino alcohols are commonly found in phosphoglycerides, which are evident in its structural formula (b).

Structural formula of ethanolamine and choline are shown. The structural formula of a phosphoglyceride is shown with the glycerol unit, phosphoric unit, and amino alcohol unit highlighted in different colors.

There are two common types of phosphoglycerides. Phosphoglycerides containing ethanolamine as the amino alcohol are called *phosphatidylethanolamines* or *cephalins*. Cephalins are found in brain tissue and nerves and also have a role in blood clotting. Phosphoglycerides containing choline as the amino alcohol unit are called *phosphatidylcholines* or *lecithins*. Lecithins occur in all

living organisms. Like cephalins, they are important constituents of nerve and brain tissue. Egg yolks are especially rich in lecithins. Commercial-grade lecithins isolated from soybeans are widely used in foods as emulsifying agents. An emulsifying agent is used to stabilize an emulsion—a dispersion of two liquids that do not normally mix, such as oil and water. Many foods are emulsions. Milk is an emulsion of butterfat in water. The emulsifying agent in milk is a protein called *casein*. Mayonnaise is an emulsion of salad oil in water, stabilized by lecithins present in egg yolk.



Sphingomyelins, the simplest sphingolipids, each contain a fatty acid, a phosphoric acid, sphingosine, and choline (Figure **16.4.6**). Because they contain phosphoric acid, they are also classified as phospholipids. Sphingomyelins are important constituents of the myelin sheath surrounding the axon of a nerve cell. Multiple sclerosis is one of several diseases resulting from damage to the myelin sheath.




Figure **16.4.6**: Sphingolipids. (a) Sphingosine, an amino alcohol, is found in all sphingolipids. (b) A sphingomyelin is also known as a phospholipid, as evidenced by the phosphoric acid unit in its structure.

Structural formula of sphingosine is shown. The general structure of a sphingolipid is shown with the sphingosine unit, fatty acid unit, phosphoric acid unit, and choline unit highlighted in different colors.

Most animal cells contain sphingolipids called cerebrosides (Figure **16.4.7**). Cerebrosides are composed of sphingosine, a fatty acid, and galactose or glucose. They therefore resemble sphingomyelins but have a sugar unit in place of the choline phosphate group. Cerebrosides are important constituents of the membranes of nerve and brain cells.




Figure 16.4.7: Cerebrosides. Cerebrosides are sphingolipids that contain a sugar unit.

General structure of a cerebroside with its sugar unit, sphingosine unit, and fatty acid unit highlighted in different colors.

The sphingolipids called gangliosides are more complex, usually containing a branched chain of three to eight monosaccharides and/or substituted sugars. Because of considerable variation in their sugar components, about 130 varieties of gangliosides have been identified. Most cell-to-cell recognition and communication processes (e.g., blood group antigens) depend on differences in the sequences of sugars in these compounds. Gangliosides are most prevalent in the outer membranes of nerve cells, although they also occur in smaller quantities in the outer membranes of most other cells. Because cerebrosides and gangliosides contain sugar groups, they are also classified as glycolipids.

Membrane Proteins

If membranes were composed only of lipids, very few ions or polar molecules could pass through their hydrophobic "sandwich filling" to enter or leave any cell. However, certain charged and polar species do cross the membrane, aided by proteins that move about in the lipid bilayer. The two major classes of proteins in the cell membrane are integral proteins, which span the hydrophobic interior of the bilayer, and peripheral proteins, which are more loosely associated with the surface of the lipid bilayer (Figure 16.4.3). Peripheral proteins may be attached to integral proteins, to the polar head groups of phospholipids, or to both by hydrogen bonding and electrostatic forces.

Small ions and molecules soluble in water enter and leave the cell by way of channels through the integral proteins. Some proteins, called *carrier proteins*, facilitate the passage of certain molecules, such as hormones and neurotransmitters, by specific interactions between the protein and the molecule being transported.

Summary

Lipids are important components of biological membranes. These lipids have dual characteristics: part of the molecule is hydrophilic, and part of the molecule is hydrophobic. Membrane lipids may be classified as phospholipids, glycolipids, and/or sphingolipids. Proteins are another important component of biological membranes. Integral proteins span the lipid bilayer, while peripheral proteins are more loosely associated with the surface of the membrane.

16.4: Membranes and Membrane Lipids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.3: Membranes and Membrane Lipids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

16.5: Steroids

Learning Objectives

- To identify the functions of steroids produced in mammals.

All the lipids discussed so far are *saponifiable*, reacting with aqueous alkali to yield simpler components, such as glycerol, fatty acids, amino alcohols, and sugars. Lipid samples extracted from cellular material, however, also contain a small but important fraction that does not react with alkali. The most important nonsaponifiable lipids are the steroids. These compounds include the *bile salts*, *cholesterol* and related compounds, and certain *hormones* (such as cortisone and the sex hormones).




Figure **16.5.1** Steroids. (a) The four-fused-ring steroid skeleton uses letter designations for each ring and the numbering of the carbon atoms. (b) The cholesterol molecule follows this pattern.

Steroids occur in plants, animals, yeasts, and molds but not in bacteria. They may exist in free form or combined with fatty acids or carbohydrates. All steroids have a characteristic structural component consisting of four fused rings. Chemists identify the rings by capital letters and number the carbon atoms as shown in Figure **16.5.1*a***. Slight variations in this structure or in the atoms or groups attached to it produce profound differences in biological activity.

Cholesterol

Cholesterol (Figure **16.5.1*b***) does not occur in plants, but it is the most abundant steroid in the human body (240 g is a typical amount). Excess cholesterol is believed to be a primary factor in the development of atherosclerosis and heart disease, which are major health problems in the United States today. About half of the body's cholesterol is interspersed in the lipid bilayer of cell membranes. Much of the rest is converted to cholic acid, which is used in the formation of bile salts. Cholesterol is also a precursor in the synthesis of sex hormones, adrenal hormones, and vitamin D.

Excess cholesterol not metabolized by the body is released from the liver and transported by the blood to the gallbladder. Normally, it stays in solution there until being secreted into the intestine (as a component of bile) to be eliminated. Sometimes, however, cholesterol in the gallbladder precipitates in the form of gallstones (Figure **16.5.2**). Indeed, the name *cholesterol* is derived from the Greek *chole*, meaning "bile," and *stereos*, meaning "solid."




Figure **16.5.2**: Numerous small gallstones made up largely of cholesterol, all removed in one patient. Grid scale 1 mm

To Your Health: Cholesterol and Heart Disease

Heart disease is the leading cause of death in the United States for both men and women. The Centers for Disease Control and Prevention reported that heart disease claimed 631,636 lives in the United States (26% of all reported deaths) in 2006.

Scientists agree that elevated cholesterol levels in the blood, as well as high blood pressure, obesity, diabetes, and cigarette smoking, are associated with an increased risk of heart disease. A long-term investigation by the National Institutes of Health showed that among men ages 30 to 49, the incidence of heart disease was five times greater for those whose cholesterol levels were above 260 mg/100 mL of serum than for those with cholesterol levels of 200 mg/100 mL or less. The cholesterol content of blood varies considerably with age, diet, and sex. Young adults average about 170 mg of cholesterol per 100 mL of blood, whereas males at age 55 may have cholesterol levels at 250 mg/100 mL or higher because the rate of cholesterol breakdown decreases with age. Females tend to have lower blood cholesterol levels than males.

To understand the link between heart disease and cholesterol levels, it is important to understand how cholesterol and other lipids are transported in the body. Lipids, such as cholesterol, are not soluble in water and therefore cannot be transported in the blood (an aqueous medium) unless they are complexed with proteins that are soluble in water, forming assemblages called *lipoproteins*. Lipoproteins are classified according to their density, which is dependent on the relative amounts of protein and lipid they contain. Lipids are less dense than proteins, so lipoproteins containing a greater proportion of lipid are less dense than those containing a greater proportion of protein.



Research on cholesterol and its role in heart disease has focused on serum levels of low-density lipoproteins (LDLs) and high-density lipoproteins (HDLs). One of the most fascinating discoveries is that high levels of HDLs reduce a person's risk of developing heart disease, whereas high levels of LDLs increase that risk. Thus the serum LDL:HDL ratio is a better predictor of heart disease risk than the overall level of serum cholesterol. Persons who, because of hereditary or dietary factors, have high LDL:HDL ratios in their blood have a higher incidence of heart disease.

How do HDLs reduce the risk of developing heart disease? No one knows for sure, but one role of HDLs appears to be the transport of excess cholesterol to the liver, where it can be metabolized. Therefore, HDLs aid in removing cholesterol from blood and from the smooth muscle cells of the arterial wall.






Dietary modifications and increased physical activity can help lower total cholesterol and improve the LDL:HDL ratio. The average American consumes about 600 mg of cholesterol from animal products each day and also synthesizes approximately 1 g of cholesterol each day, mostly in the liver. The amount of cholesterol synthesized is controlled by the cholesterol level in the blood; when the blood cholesterol level exceeds 150 mg/100 mL, the rate of cholesterol biosynthesis is halved. Hence, if cholesterol is present in the diet, a feedback mechanism suppresses its synthesis in the liver. However, the ratio of suppression is not a 1:1 ratio; the reduction in biosynthesis does not equal the amount of cholesterol ingested. Thus, dietary substitutions of unsaturated fat for saturated fat, as well as a reduction in consumption of *trans* fatty acids, is recommended to help lower serum cholesterol and the risk of heart disease.

Steroid Hormones

Hormones are chemical messengers that are released in one tissue and transported through the circulatory system to one or more other tissues. One group of hormones is known as steroid hormones because these hormones are synthesized from cholesterol, which is also a steroid. There are two main groups of steroid hormones: adrenocortical hormones and sex hormones.

The adrenocortical hormones, such as aldosterone and cortisol (Table **16.5.1**), are produced by the adrenal gland, which is located adjacent to each kidney. Aldosterone acts on most cells in the body, but it is particularly effective at enhancing the rate of reabsorption of sodium ions in the kidney tubules and increasing the secretion of potassium ions and/or hydrogen ions by the tubules. Because the concentration of sodium ions is the major factor influencing water retention in tissues, aldosterone promotes water retention and reduces urine output. Cortisol regulates several key metabolic reactions (for example, increasing glucose production and mobilizing fatty acids and amino acids). It also inhibits the inflammatory response of tissue to injury or stress. Cortisol and its analogs are therefore used pharmacologically as immunosuppressants after transplant operations and in the treatment of severe skin allergies and autoimmune diseases, such as rheumatoid arthritis.

Table **16.5.1**: Representative Steroid Hormones and Their Physiological Effects

Hormone	Effect
 Aldosterone	regulates salt metabolism; stimulates kidneys to retain sodium and excrete potassium
 Cortisol (Hydrocortisone)	stimulates the conversion of proteins to carbohydrates
 Progesterone	regulates the menstrual cycle; maintains pregnancy
 Estradiol (an estrogen)	stimulates female sex characteristics; regulates changes during the menstrual cycle
 Testosterone (an androgen)	stimulates and maintains male sex characteristics

The sex hormones are a class of steroid hormones secreted by the gonads (ovaries or testes), the placenta, and the adrenal glands. Testosterone and androstenedione are the primary male sex hormones, or *androgens*, controlling the primary sexual characteristics of males, or the development of the male genital organs and the continuous production of sperm. Androgens are also responsible for the development of secondary male characteristics, such as facial hair, deep voice, and muscle strength. Two kinds of sex

hormones are of particular importance in females: progesterone, which prepares the uterus for pregnancy and prevents the further release of eggs from the ovaries during pregnancy, and the estrogens, which are mainly responsible for the development of female secondary sexual characteristics, such as breast development and increased deposition of fat tissue in the breasts, the buttocks, and the thighs. Both males and females produce androgens and estrogens, differing in the amounts of secreted hormones rather than in the presence or absence of one or the other.

Sex hormones, both natural and synthetic, are sometimes used therapeutically. For example, a woman who has had her ovaries removed may be given female hormones to compensate. Some of the earliest chemical compounds employed in cancer chemotherapy were sex hormones. For example, estrogens are one treatment option for prostate cancer because they block the release and activity of testosterone. Testosterone enhances prostate cancer growth. Sex hormones are also administered in preparation for sex-change operations, to promote the development of the proper secondary sexual characteristics. Oral contraceptives are synthetic derivatives of the female sex hormones; they work by preventing ovulation.

Bile Salts

Bile is a yellowish green liquid (pH 7.8–8.6) produced in the liver. The most important constituents of bile are bile salts, which are sodium salts of amidelike combinations of bile acids, such as cholic acid (part (a) of Figure **16.5.3**) and an amine such as the amino acid glycine (part (b) of Figure **16.5.3**). They are synthesized from cholesterol in the liver, stored in the gallbladder, and then secreted in bile into the small intestine. In the gallbladder, the composition of bile gradually changes as water is absorbed and the other components become more concentrated.




Figure **16.5.3** Bile Acids. (a) Cholic acid is an example of a bile acid. (b) Sodium glycocholate is a bile salt synthesized from cholic acid and glycine.

Because they contain both hydrophobic and hydrophilic groups, bile salts are highly effective detergents and emulsifying agents; they break down large fat globules into smaller ones and keep those smaller globules suspended in the aqueous digestive environment. Enzymes can then hydrolyze fat molecules more efficiently. Thus, the major function of bile salts is to aid in the digestion of dietary lipids.

> Surgical removal is often advised for a gallbladder that becomes infected, inflamed, or perforated. This surgery does not seriously affect digestion because bile is still produced by the liver, but the liver's bile is more dilute and its secretion into the small intestine is not as closely tied to the arrival of food.

Summary

Steroids have a four-fused-ring structure and have a variety of functions. Cholesterol is a steroid found in mammals that is needed for the formation of cell membranes, bile acids, and several hormones. Bile salts are secreted into the small intestine to aid in the digestion of fats.

16.5: Steroids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.4: Steroids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

16.6: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Lipids, found in the body tissues of all organisms, are compounds that are more soluble in organic solvents than in water. Many of them contain **fatty acids**, which are carboxylic acids that generally contain an even number of 4–20 carbon atoms in an unbranched chain. **Saturated fatty acids** have no carbon-to-carbon double bonds. **Monounsaturated fatty acids** have a single carbon-to-carbon double bond, while **polyunsaturated fatty acids** have more than one carbon-to-carbon double bond. Linoleic and linolenic acid are known as **essential fatty acids** because the human body cannot synthesize these polyunsaturated fatty acids. The lipids known as **fats** and **oils** are triacylglycerols, more commonly called **triglycerides**—esters composed of three fatty acids joined to the trihydroxy alcohol glycerol. Fats are triglycerides that are solid at room temperature, and oils are triglycerides that are liquid at room temperature. Fats are found mainly in animals, and oils found mainly in plants. *Saturated triglycerides* are those containing a higher proportion of saturated fatty acid chains (fewer carbon-to-carbon double bonds); *unsaturated triglycerides* contain a higher proportion of unsaturated fatty acid chains.

Saponification is the hydrolysis of a triglyceride in a basic solution to form glycerol and three carboxylate anions or soap molecules. Other important reactions are the hydrogenation and oxidation of double bonds in unsaturated fats and oils.

Phospholipids are lipids containing phosphorus. In **phosphoglycerides**, the phosphorus is joined to an amino alcohol unit. Some phosphoglycerides, like lecithins, are used to stabilize an **emulsion**—a dispersion of two liquids that do not normally mix, such as oil and water. **Sphingolipids** are lipids for which the precursor is the amino alcohol sphingosine, rather than glycerol. A **glycolipid** has a sugar substituted at one of the OH groups of either glycerol or sphingosine. All are highly polar lipids found in cell membranes.

Polar lipids have dual characteristics: one part of the molecule is ionic and dissolves in water; the rest has a hydrocarbon structure and dissolves in nonpolar substances. Often, the ionic part is referred to as **hydrophilic** (literally, “water loving”) and the nonpolar part as **hydrophobic** (“water fearing”). When placed in water, polar lipids disperse into any one of three arrangements: *micelles*, *monolayers*, and *bilayers*. **Micelles** are aggregations of molecules in which the hydrocarbon tails of the lipids, being hydrophobic, are directed inward (away from the surrounding water), and the hydrophilic heads that are directed outward into the water. **Bilayers** are double layers arranged so that the hydrophobic tails are sandwiched between the two layers of hydrophilic heads, which remain in contact with the water.

Every living cell is enclosed by a *cell membrane* composed of a lipid bilayer. In animal cells, the bilayer consists mainly of phospholipids, glycolipids, and the steroid cholesterol. Embedded in the bilayer are **integral proteins**, and **peripheral proteins** are loosely associated with the surface of the bilayer. Everything between the cell membrane and the membrane of the cell nucleus is called the **cytoplasm**.

Most lipids can be saponified, but some, such as **steroids**, cannot be saponified. The steroid **cholesterol** is found in animal cells but never in plant cells. It is a main component of all cell membranes and a precursor for hormones, vitamin D, and bile salts. Bile salts are the most important constituents of **bile**, which is a yellowish-green liquid secreted by the gallbladder into the small intestine and is needed for the proper digestion of lipids.

16.6: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **17.S: Lipids (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

17: Proteins and Enzymes

Proteins may be defined as compounds of high molar mass consisting largely or entirely of chains of amino acids. Their masses range from several thousand to several million daltons (Da). In addition to carbon, hydrogen, and oxygen atoms, all proteins contain nitrogen and sulfur atoms, and many also contain phosphorus atoms and traces of other elements. Proteins serve a variety of roles in living organisms and are often classified by these biological roles. Muscle tissue is largely protein, as are skin and hair. Proteins are present in the blood, in the brain, and even in tooth enamel. Each type of cell in our bodies makes its own specialized proteins, as well as proteins common to all or most cells. We begin our study of proteins by looking at the properties and reactions of amino acids, which is followed by a discussion of how amino acids link covalently to form peptides and proteins. We end the chapter with a discussion of enzymes—the proteins that act as catalysts in the body.

17.1: Properties of Amino Acids
17.2: Reactions of Amino Acids
17.3: Peptides
17.4: Proteins
17.5: Enzymes
17.6: Enzymatic Catalysis
17.7: Enzymatic Activity
17.8: Enzyme Inhibition
17.9: Enzyme Cofactors and Vitamins
17.10: Chapter Summary

17: Proteins and Enzymes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


17.1: Properties of Amino Acids

Learning Objectives


- To recognize amino acids and classify them based on the characteristics of their side chains.

The proteins in all living species, from bacteria to humans, are constructed from the same set of 20 amino acids, so called because each contains an amino group attached to a carboxylic acid. The amino acids in proteins are α-amino acids, which means the amino group is attached to the α-carbon of the carboxylic acid. Humans can synthesize only about half of the needed amino acids; the remainder must be obtained from the diet and are known as essential amino acids. However, two additional amino acids have been found in limited quantities in proteins: Selenocysteine was discovered in 1986, while pyrrolysine was discovered in 2002.

The amino acids are colorless, nonvolatile, crystalline solids, melting and decomposing at temperatures above 200°C. These melting temperatures are more like those of inorganic salts than those of amines or organic acids and indicate that the structures of the amino acids in the solid state and in neutral solution are best represented as having both a negatively charged group and a positively charged group. Such a species is known as a zwitterion.



α-Amino acid drawn as a zwitterion























α-Amino acid drawn as an uncharged molecule; not an accurate respresentation of amino acid structure

Classification


In addition to the amino and carboxyl groups, amino acids have a side chain or R group attached to the α-carbon. Each amino acid has unique characteristics arising from the size, shape, solubility, and ionization properties of its R group. As a result, the side chains of amino acids exert a profound effect on the structure and biological activity of proteins. Although amino acids can be classified in various ways, one common approach is to classify them according to whether the functional group on the side chain at neutral pH is nonpolar, polar but uncharged, negatively charged, or positively charged. The structures and names of the 20 amino acids, their one- and three-letter abbreviations, and some of their distinctive features are given in Table **17.1.1**.

Table **17.1.1**: Common Amino Acids Found in Proteins

Common Name	Abbreviation	Structural Formula (at pH 6)	Molar Mass	Distinctive Feature
Amino acids with a nonpolar R group				
glycine	gly (G)		75	the only amino acid lacking a chiral carbon
alanine	ala (A)		89	—
valine	val (V)		117	a branched-chain amino acid
leucine	leu (L)		131	a branched-chain amino acid
isoleucine	ile (I)		131	an essential amino acid because most animals cannot synthesize branched-chain amino acids
phenylalanine	phe (F)		165	also classified as an aromatic amino acid
tryptophan	trp (W)		204	also classified as an aromatic amino acid
methionine	met (M)		149	side chain functions as a methyl group donor
proline	pro (P)		115	contains a secondary amine group; referred to as an *α-imino acid*
Amino acids with a polar but neutral R group				
serine	ser (S)		105	found at the active site of many enzymes
threonine	thr (T)		119	named for its similarity to the sugar threose
cysteine	cys (C)		121	oxidation of two cysteine molecules yields *cystine*


Common Name	Abbreviation	Structural Formula (at pH 6)	Molar Mass	Distinctive Feature
tyrosine	tyr (Y)		181	also classified as an aromatic amino acid
asparagine	asn (N)		132	the amide of aspartic acid
glutamine	gln (Q)		146	the amide of glutamic acid
Amino acids with a negatively charged R group				
aspartic acid	asp (D)		132	carboxyl groups are ionized at physiological pH; also known as aspartate
glutamic acid	glu (E)		146	carboxyl groups are ionized at physiological pH; also known as glutamate
Amino acids with a positively charged R group				
histidine	his (H)		155	the only amino acid whose R group has a pK_a (6.0) near physiological pH
lysine	lys (K)		147	—
arginine	arg (R)		175	almost as strong a base as sodium hydroxide

The first amino acid to be isolated was asparagine in 1806. It was obtained from protein found in asparagus juice (hence the name). Glycine, the major amino acid found in gelatin, was named for its sweet taste (Greek *glykys*, meaning “sweet”). In some cases an amino acid found in a protein is actually a derivative of one of the common 20 amino acids (one such derivative is hydroxyproline). The modification occurs *after* the amino acid has been assembled into a protein.



Configuration

Notice in Table 17.1.1 that glycine is the only amino acid whose α-carbon is *not* chiral. Therefore, with the exception of glycine, the amino acids could theoretically exist in either the D- or the L-enantiomeric form and rotate plane-polarized light. As with sugars, chemists used L-glyceraldehyde as the reference compound for the assignment of absolute configuration to amino acids. Its structure closely resembles an amino acid structure except that in the latter, an amino group takes the place of the OH group on the chiral carbon of the L-glyceraldehyde and a carboxylic acid replaces the aldehyde. Modern stereochemistry assignments using the Cahn-Ingold-Prelog priority rules used ubiquitously in chemistry show that all of the naturally occurring chiral amino acids are S except Cys which is R.



We learned that all naturally occurring sugars belong to the D series. It is interesting, therefore, that nearly all known plant and animal proteins are composed entirely of L-amino acids. However, certain bacteria contain D-amino acids in their cell walls, and several antibiotics (e.g., actinomycin D and the gramicidins) contain varying amounts of D-leucine, D-phenylalanine, and D-valine.

Summary

Amino acids can be classified based on the characteristics of their distinctive side chains as nonpolar, polar but uncharged, negatively charged, or positively charged. The amino acids found in proteins are L-amino acids.

17.1: Properties of Amino Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **18.1: Properties of Amino Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.2: Reactions of Amino Acids

Learning Objectives

- To explain how an amino acid can act as both an acid and a base.

The structure of an amino acid allows it to act as both an acid and a base. An amino acid has this ability because at a certain pH value (different for each amino acid) nearly all the amino acid molecules exist as zwitterions. If acid is added to a solution containing the zwitterion, the carboxylate group captures a hydrogen (H^+) ion, and the amino acid becomes positively charged. If base is added, ion removal of the H^+ ion from the amino group of the zwitterion produces a negatively charged amino acid. In both circumstances, the amino acid acts to maintain the pH of the system—that is, to remove the added acid (H^+) or base (OH^-) from solution.




Example 17.2.1

a. Draw the structure for the anion formed when glycine (at neutral pH) reacts with a base.
b. Draw the structure for the cation formed when glycine (at neutral pH) reacts with an acid.

Solution

a. The base removes H^+ from the protonated amine group.



- The acid adds H^+ to the carboxylate group.



Exercise 17.2.1

a. Draw the structure for the cation formed when valine (at neutral pH) reacts with an acid.
b. Draw the structure for the anion formed when valine (at neutral pH) reacts with a base.

The particular pH at which a given amino acid exists in solution as a zwitterion is called the isoelectric point (pI). At its pI, the positive and negative charges on the amino acid balance, and the molecule as a whole is electrically neutral. The amino acids whose side chains are always neutral have isoelectric points ranging from 5.0 to 6.5. The basic amino acids (which have positively charged side chains at neutral pH) have relatively high examples. Acidic amino acids (which have negatively charged side chains at neutral pH) have quite low examples (Table 17.2.1).

Table 17.2.1: ExampIes of Some Representative Amino Acids

Amino Acid	Classification	pI
alanine	nonpolar	6.0
valine	nonpolar	6.0
serine	polar, uncharged	5.7
threonine	polar, uncharged	6.5
arginine	positively charged (basic)	10.8
histidine	positively charged (basic)	7.6
lysine	positively charged (basic)	9.8
aspartic acid	negatively charged (acidic)	3.0
glutamic acid	negatively charged (acidic)	3.2

Amino acids undergo reactions characteristic of carboxylic acids and amines. The reactivity of these functional groups is particularly important in linking amino acids together to form peptides and proteins, as you will see later in this chapter. Simple chemical tests that are used to detect amino acids take advantage of the reactivity of these functional groups. An example is the ninhydrin test in which the amine functional group of α-amino acids reacts with ninhydrin to form purple-colored compounds. Ninhydrin is used to detect fingerprints because it reacts with amino acids from the proteins in skin cells transferred to the surface by the individual leaving the fingerprint.



Ninhydrin



Summary

Amino acids can act as both an acid and a base due to the presence of the amino and carboxyl functional groups. The pH at which a given amino acid exists in solution as a zwitterion is called the *isoelectric point* (pI).

17.2: Reactions of Amino Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **18.2: Reactions of Amino Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.3: Peptides


Learning Objectives

- Explain how a peptide is formed from individual amino acids.
- Explain why the sequence of amino acids in a protein is important.


Two or more amino acids can join together into chains called peptides. Previously, we discussed the reaction between ammonia and a carboxylic acid to form an amide. In a similar reaction, the amino group on one amino acid molecule reacts with the carboxyl group on another, releasing a molecule of water and forming an amide linkage:



An amide bond joining two amino acid units is called a peptide bond. Note that the product molecule still has a reactive amino group on the left and a reactive carboxyl group on the right. These can react with additional amino acids to lengthen the peptide. The process can continue until thousands of units have joined, resulting in large proteins.



A chain consisting of only two amino acid units is called a *dipeptide*; a chain consisting of three is a *tripeptide*. By convention, peptide and protein structures are depicted with the amino acid whose amino group is free (the N-terminal end) on the left and the amino acid with a free carboxyl group (the C-terminal end) to the right.



The general term *peptide* refers to an amino acid chain of unspecified length. However, chains of about 50 amino acids or more are usually called proteins or polypeptides. In its physiologically active form, a protein may be composed of one or more polypeptide chains.




Figure **17.3.1**: Space-filling model of bradykinin. (Public Domain; Fvasconcellos)

For peptides and proteins to be physiologically active, it is not enough that they incorporate certain amounts of specific amino acids. The order, or *sequence*, in which the amino acids are connected is also of critical importance. Bradykinin is a nine-amino acid peptide (Figure **17.3.1**) produced in the blood that has the following amino acid sequence:

arg-pro-pro-gly-phe-ser-pro-phe-arg

This peptide lowers blood pressure, stimulates smooth muscle tissue, increases capillary permeability, and causes pain. When the order of amino acids in bradykinin is reversed,

arg-phe-pro-ser-phe-gly-pro-pro-arg

the peptide resulting from this synthesis shows none of the activity of bradykinin.

Just as millions of different words are spelled with our 26-letter English alphabet, millions of different proteins are made with the 20 common amino acids. However, just as the English alphabet can be used to write gibberish, amino acids can be put together in the *wrong sequence* to produce nonfunctional proteins. Although the correct sequence is ordinarily of utmost importance, it is not always absolutely required. Just as you can sometimes make sense of incorrectly spelled English words, a protein with a small percentage of “incorrect” amino acids may continue to function. However, it rarely functions as well as a protein having the correct sequence. There are also instances in which seemingly minor errors of sequence have disastrous effects. For example, in some people, every molecule of hemoglobin (a protein in the blood that transports oxygen) has a single incorrect amino acid unit out of about 300 (a single valine replaces a glutamic acid). That “minor” error is responsible for sickle cell anemia, an inherited condition that usually is fatal.

Summary

The amino group of one amino acid can react with the carboxyl group on another amino acid to form a peptide bond that links the two amino acids together. Additional amino acids can be added on through the formation of addition peptide (amide) bonds. A sequence of amino acids in a peptide or protein is written with the N-terminal amino acid first and the C-terminal amino acid at the end (writing left to right).

17.3: Peptides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.3: Peptides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.4: Proteins

Learning Objectives

- Describe the four levels of protein structure.
- Identify the types of attractive interactions that hold proteins in their most stable three-dimensional structure.
- Explain what happens when proteins are denatured.
- Identify how a protein can be denatured.

Each of the thousands of naturally occurring proteins has its own characteristic amino acid composition and sequence that result in a unique three-dimensional shape. Since the 1950s, scientists have determined the amino acid sequences and three-dimensional conformation of numerous proteins and thus obtained important clues on how each protein performs its specific function in the body.

Proteins are compounds of high molar mass consisting largely or entirely of chains of amino acids. Because of their great complexity, protein molecules cannot be classified on the basis of specific structural similarities, as carbohydrates and lipids are categorized. The two major structural classifications of proteins are based on far more general qualities: whether the protein is (1) fiberlike and insoluble or (2) globular and soluble. Some proteins, such as those that compose hair, skin, muscles, and connective tissue, are fiberlike. These fibrous proteins are insoluble in water and usually serve structural, connective, and protective functions. Examples of fibrous proteins are keratins, collagens, myosins, and elastins. Hair and the outer layer of skin are composed of keratin. Connective tissues contain collagen. Myosins are muscle proteins and are capable of contraction and extension. Elastins are found in ligaments and the elastic tissue of artery walls.

Globular proteins, the other major class, are soluble in aqueous media. In these proteins, the chains are folded so that the molecule as a whole is roughly spherical. Familiar examples include egg albumin from egg whites and serum albumin in blood. Serum albumin plays a major role in transporting fatty acids and maintaining a proper balance of osmotic pressures in the body. Hemoglobin and myoglobin, which are important for binding oxygen, are also globular proteins.




Figure 17.4.1: Primary Structure of Human Insulin. Human insulin, whose amino acid sequence is shown here, is a hormone that is required for the proper metabolism of glucose.

Levels of Protein Structure

The structure of proteins is generally described as having four organizational levels. The first of these is the primary structure, which is the number and sequence of amino acids in a protein's polypeptide chain or chains, beginning with the free amino group

and maintained by the peptide bonds connecting each amino acid to the next. The primary structure of insulin, composed of 51 amino acids, is shown in Figure **17.4.1**.

A protein molecule is not a random tangle of polypeptide chains. Instead, the chains are arranged in unique but specific conformations. The term secondary structure refers to the fixed arrangement of the polypeptide backbone. On the basis of X ray studies, Linus Pauling and Robert Corey postulated that certain proteins or portions of proteins twist into a spiral or a helix. This helix is stabilized by *intrachain* hydrogen bonding between the carbonyl oxygen atom of one amino acid and the amide hydrogen atom four amino acids up the chain (located on the next turn of the helix) and is known as a right-handed α-helix. X ray data indicate that this helix makes one turn for every 3.6 amino acids, and the side chains of these amino acids project outward from the coiled backbone (Figure **17.4.2**). The α-keratins, found in hair and wool, are exclusively α-helical in conformation. Some proteins, such as gamma globulin, chymotrypsin, and cytochrome c, have little or no helical structure. Others, such as hemoglobin and myoglobin, are helical in certain regions but not in others.




Figure **17.4.2** A Ball-and-Stick Model of an α-Helix. This ball-and-stick model shows the intrachain hydrogen bonding between carbonyl oxygen atoms and amide hydrogen atoms. Each turn of the helix spans 3.6 amino acids. Note that the side chains (represented as green spheres) point out from the helix.

Another common type of secondary structure, called the *β-pleated sheet conformation*, is a sheetlike arrangement in which two or more extended polypeptide chains (or separate regions on the same chain) are aligned side by side. The aligned segments can run either parallel or antiparallel—that is, the N-terminals can face in the same direction on adjacent chains or in different directions—and are connected by *interchain* hydrogen bonding (Figure **17.4.3**). The β-pleated sheet is particularly important in structural proteins, such as silk fibroin. It is also seen in portions of many enzymes, such as carboxypeptidase A and lysozyme.




Figure **17.4.3**: A Ball-and-Stick Model of the β-Pleated Sheet Structure in Proteins. The side chains extend above or below the sheet and alternate along the chain. The protein chains are held together by interchain hydrogen bonding.

Tertiary structure refers to the unique three-dimensional shape of the protein as a whole, which results from the folding and bending of the protein backbone. The tertiary structure is intimately tied to the proper biochemical functioning of the protein. Figure **17.4.4** shows a depiction of the three-dimensional structure of insulin.





Figure 17.4.4: A Ribbon Model of the Three-Dimensional Structure of Insulin. The spiral regions represent sections of the polypeptide chain that have an α-helical structure, while the broad arrows represent β-pleated sheet structures.

Four major types of attractive interactions determine the shape and stability of the tertiary structure of proteins. You studied several of them previously.

1. Ionic bonding. Ionic bonds result from electrostatic attractions between positively and negatively charged side chains of amino acids. For example, the mutual attraction between an aspartic acid carboxylate ion and a lysine ammonium ion helps to maintain a particular folded area of a protein (part (a) of Figure 17.4.5).
2. Hydrogen bonding. Hydrogen bonding forms between a highly electronegative oxygen atom or a nitrogen atom and a hydrogen atom attached to another oxygen atom or a nitrogen atom, such as those found in polar amino acid side chains. Hydrogen bonding (as well as ionic attractions) is extremely important in both the intra- and intermolecular interactions of proteins (part (b) of Figure 17.4.5).
3. Disulfide linkages. Two cysteine amino acid units may be brought close together as the protein molecule folds. Subsequent oxidation and linkage of the sulfur atoms in the highly reactive sulfhydryl (SH) groups leads to the formation of cystine (part (c) of Figure 17.4.5). Intrachain disulfide linkages are found in many proteins, including insulin (yellow bars in Figure 17.4.1) and have a strong stabilizing effect on the tertiary structure.



4. Dispersion forces. Dispersion forces arise when a normally nonpolar atom becomes momentarily polar due to an uneven distribution of electrons, leading to an instantaneous dipole that induces a shift of electrons in a neighboring nonpolar atom. Dispersion forces are weak but can be important when other types of interactions are either missing or minimal (part (d) of Figure 17.4.5). This is the case with fibroin, the major protein in silk, in which a high proportion of amino acids in the protein have nonpolar side chains. The term *hydrophobic interaction* is often misused as a synonym for dispersion forces. Hydrophobic interactions arise because water molecules engage in hydrogen bonding with other water molecules (or groups in proteins capable of hydrogen bonding). Because nonpolar groups cannot engage in hydrogen bonding, the protein folds in such a way that these groups are buried in the interior part of the protein structure, minimizing their contact with water.




Figure **17.4.5**: Tertiary Protein Structure Interactions. Four interactions stabilize the tertiary structure of a protein: (a) ionic bonding, (b) hydrogen bonding, (c) disulfide linkages, and (d) dispersion forces.

When a protein contains more than one polypeptide chain, each chain is called a *subunit*. The arrangement of multiple subunits represents a fourth level of structure, the quaternary structure of a protein. Hemoglobin, with four polypeptide chains or subunits, is the most frequently cited example of a protein having quaternary structure (Figure **17.4.6**). The quaternary structure of a protein is produced and stabilized by the same kinds of interactions that produce and maintain the tertiary structure. A schematic representation of the four levels of protein structure is in Figure **17.4.7**.




Figure **17.4.6** The Quaternary Structure of Hemoglobin. Hemoglobin is a protein that transports oxygen throughout the body.

Source: Image from the RCSB PDB (www.pdb.org(opens in new window)) of PDB ID 1I3D (R.D. Kidd, H.M. Baker, A.J. Mathews, T. Brittain, E.N. Baker (2001) Oligomerization and ligand binding in a homotetrameric hemoglobin: two high-resolution crystal structures of hemoglobin Bart's (gamma(4)), a marker for alpha-thalassemia. Protein Sci. 1739–1749).




Figure **17.4.7**: Levels of Structure in Proteins

The *primary structure* consists of the specific amino acid sequence. The resulting peptide chain can twist into an α-helix, which is one type of *secondary structure*. This helical segment is incorporated into the *tertiary structure* of the folded polypeptide chain. The single polypeptide chain is a subunit that constitutes the *quaternary structure* of a protein, such as hemoglobin that has four polypeptide chains.

Denaturation of Proteins

The highly organized structures of proteins are truly masterworks of chemical architecture. But highly organized structures tend to have a certain delicacy, and this is true of proteins. Denaturation is the term used for any change in the three-dimensional structure of a protein that renders it incapable of performing its assigned function. A denatured protein cannot do its job. (Sometimes denaturation is equated with the precipitation or coagulation of a protein; our definition is a bit broader.) A wide variety of reagents and conditions, such as heat, organic compounds, pH changes, and heavy metal ions can cause protein denaturation (Figure **17.4.1**).

Figure **17.4.1**: Protein Denaturation Methods

Method	Effect on Protein Structure
Heat above 50°C or ultraviolet (UV) radiation	Heat or UV radiation supplies kinetic energy to protein molecules, causing their atoms to vibrate more rapidly and disrupting relatively weak hydrogen bonding and dispersion forces.
Use of organic compounds, such as ethyl alcohol	These compounds are capable of engaging in intermolecular hydrogen bonding with protein molecules, disrupting intramolecular hydrogen bonding within the protein.
Salts of heavy metal ions, such as mercury, silver, and lead	These ions form strong bonds with the carboxylate anions of the acidic amino acids or SH groups of cysteine, disrupting ionic bonds and disulfide linkages.
Alkaloid reagents, such as tannic acid (used in tanning leather)	These reagents combine with positively charged amino groups in proteins to disrupt ionic bonds.

Anyone who has fried an egg has observed denaturation. The clear egg white turns opaque as the albumin denatures and coagulates. No one has yet reversed that process. However, given the proper circumstances and enough time, a protein that has unfolded under sufficiently gentle conditions can refold and may again exhibit biological activity (Figure **17.4.8**). Such evidence suggests that, at least for these proteins, the primary structure determines the secondary and tertiary structure. A given sequence of amino acids seems to adopt its particular three-dimensional arrangement naturally if conditions are right.




Figure **17.4.8**: Denaturation and Renaturation of a Protein. The denaturation (unfolding) and renaturation (refolding) of a protein is depicted. The red boxes represent stabilizing interactions, such as disulfide linkages, hydrogen bonding, and/or ionic bonds.

The primary structures of proteins are quite sturdy. In general, fairly vigorous conditions are needed to hydrolyze peptide bonds. At the secondary through quaternary levels, however, proteins are quite vulnerable to attack, though they vary in their vulnerability to denaturation. The delicately folded globular proteins are much easier to denature than are the tough, fibrous proteins of hair and skin.

Summary

Proteins can be divided into two categories: fibrous, which tend to be insoluble in water, and globular, which are more soluble in water. A protein may have up to four levels of structure. The primary structure consists of the specific amino acid sequence. The resulting peptide chain can form an α-helix or β-pleated sheet (or local structures not as easily categorized), which is known as secondary structure. These segments of secondary structure are incorporated into the tertiary structure of the folded polypeptide chain. The quaternary structure describes the arrangements of subunits in a protein that contains more than one subunit. Four major types of attractive interactions determine the shape and stability of the folded protein: ionic bonding, hydrogen bonding, disulfide linkages, and dispersion forces. A wide variety of reagents and conditions can cause a protein to unfold or denature.

17.4: Proteins is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.4: Proteins** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.5: Enzymes

Learning Objectives

- Explain the functions of enzymes.
- Explain how enzymes are classified and named.

A catalyst is any substance that increases the *rate* or speed of a chemical reaction without being changed or consumed in the reaction. Enzymes are biological catalysts, and nearly all of them are proteins. The reaction rates attained by enzymes are truly amazing. In their presence, reactions occur at rates that are a million (10^6) or more times faster than would be attainable in their absence. What is even more amazing is that enzymes perform this function at body temperature (~37°C) and physiological pH (pH ~7), rather than at the conditions that are typically necessary to increase reaction rates (high temperature or pressure, the use of strong oxidizing or reducing agents or strong acids or bases, or a combination of any of these). In addition, enzymes are highly specific in their action; that is, each enzyme catalyzes only one type of reaction in only one compound or a group of structurally related compounds. The compound or compounds on which an enzyme acts are known as its substrates.

Hundreds of enzymes have been purified and studied in an effort to understand how they work so effectively and with such specificity. The resulting knowledge has been used to design drugs that inhibit or activate particular enzymes. An example is the intensive research to improve the treatment of or find a cure for acquired immunodeficiency syndrome (AIDS). AIDS is caused by the human immunodeficiency virus (HIV). Researchers are studying the enzymes produced by this virus and are developing drugs intended to block the action of those enzymes without interfering with enzymes produced by the human body. Several of these drugs have now been approved for use by AIDS patients.

Table **17.5.1**: Classes of Enzymes

Class	Type of Reaction Catalyzed	Examples
oxidoreductases	oxidation-reduction reactions	Dehydrogenases catalyze oxidation-reduction reactions involving hydrogen and reductases catalyze reactions in which a substrate is reduced.
transferases	transfer reactions of groups, such as methyl, amino, and acetyl	Transaminases catalyze the transfer of amino group, and kinases catalyze the transfer of a phosphate group.
hydrolases	hydrolysis reactions	Lipases catalyze the hydrolysis of lipids, and proteases catalyze the hydrolysis of proteins
lyases	reactions in which groups are removed without hydrolysis or addition of groups to a double bond	Decarboxylases catalyze the removal of carboxyl groups.
isomerases	reactions in which a compound is converted to its isomer	Isomerases may catalyze the conversion of an aldose to a ketose, and mutases catalyze reactions in which a functional group is transferred from one atom in a substrate to another.
ligases	reactions in which new bonds are formed between carbon and another atom; energy is required	Synthetases catalyze reactions in which two smaller molecules are linked to form a larger one.


The first enzymes to be discovered were named according to their source or method of discovery. The enzyme *pepsin*, which aids in the hydrolysis of proteins, is found in the digestive juices of the stomach (Greek *pepsis*, meaning “digestion”). *Papain*, another

enzyme that hydrolyzes protein (in fact, it is used in meat tenderizers), is isolated from papayas. As more enzymes were discovered, chemists recognized the need for a more systematic and chemically informative identification scheme. In the current numbering and naming scheme, under the oversight of the Nomenclature Commission of the International Union of Biochemistry, enzymes are arranged into six groups according to the general type of reaction they catalyze (Table **17.5.1**), with subgroups and secondary subgroups that specify the reaction more precisely.

Figure* 17.5.1*: *Structure of the alcohol dehydrogenase protein (E.C.1.1.1.1) (EE ISOZYME) complexed wtih nicotinamide adenini dinulceotide (NAD) and zinc (PDB: 1CDO).*

Each enzyme is assigned a four-digit number, preceded by the prefix EC—for enzyme classification—that indicates its group, subgroup, and so forth. This is demonstrated in Table **17.5.2** for alcohol dehydrogenase. Each enzyme is also given a name consisting of the root of the name of its substrate or substrates and the *-ase* suffix. Thus urease is the enzyme that catalyzes the hydrolysis of urea.

Table **17.5.2**: Assignment of an Enzyme Classification Number

Alcohol Dehydrogenase: EC 1.1.1.1	
The first digit indicates that this enzyme is an oxidoreductase; that is, an enzyme that catalyzes an oxidation-reduction reaction. The second digit indicates that this oxidoreductase catalyzes a reaction involving a primary or secondary alcohol. The third digit indicates that either the coenzyme NAD^+ or $NADP^+$ is required for this reaction.	
The fourth digit indicates that this was the first enzyme isolated, characterized, and named using this system of nomenclature.	
The systematic name for this enzyme is *alcohol:NAD^+ oxidoreductase*, while the recommended or common name is alcohol dehydrogenase.	
Reaction catalyzed:	

Summary

An enzyme is a biological catalyst, a substance that increases the rate of a chemical reaction without being changed or consumed in the reaction. A systematic process is used to name and classify enzymes.

17.5: Enzymes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.5: Enzymes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.6: Enzymatic Catalysis

Learning Objectives

- To describe the interaction between an enzyme and its substrate.

Enzyme-catalyzed reactions occur in at least two steps. In the first step, an enzyme molecule (E) and the substrate molecule or molecules (S) collide and react to form an intermediate compound called the *enzyme-substrate* (E–S) *complex*. (This step is reversible because the complex can break apart into the original substrate or substrates and the free enzyme.) Once the E–S complex forms, the enzyme is able to catalyze the formation of product (P), which is then released from the enzyme surface:

$$S + E \rightarrow E\text{–}S \tag{17.6.1}$$

$$E\text{–}S \rightarrow P + E \tag{17.6.2}$$

Hydrogen bonding and other electrostatic interactions hold the enzyme and substrate together in the complex. The structural features or functional groups on the enzyme that participate in these interactions are located in a cleft or pocket on the enzyme surface. This pocket, where the enzyme combines with the substrate and transforms the substrate to product is called the active site of the enzyme (Figure **17.6.1**).




Figure **17.6.1**: Substrate Binding to the Active Site of an Enzyme. The enzyme dihydrofolate reductase is shown with one of its substrates: $NADP^+$ (a) unbound and (b) bound. The $NADP^+$ (shown in red) binds to a pocket that is complementary to it in shape and ionic properties.

The active site of an enzyme possesses a unique conformation (including correctly positioned bonding groups) that is complementary to the structure of the substrate, so that the enzyme and substrate molecules fit together in much the same manner as a key fits into a tumbler lock. In fact, an early model describing the formation of the enzyme-substrate complex was called the lock-and-key model (Figure **17.6.2**). This model portrayed the enzyme as conformationally rigid and able to bond only to substrates that exactly fit the active site.




Figure **17.6.2**: The Lock-and-Key Model of Enzyme Action. (a) Because the substrate and the active site of the enzyme have complementary structures and bonding groups, they fit together as a key fits a lock. (b) The catalytic reaction occurs while the two are bonded together in the enzyme-substrate complex.

Working out the precise three-dimensional structures of numerous enzymes has enabled chemists to refine the original lock-and-key model of enzyme actions. They discovered that the binding of a substrate often leads to a large conformational change in the enzyme, as well as to changes in the structure of the substrate or substrates. The current theory, known as the induced-fit model, says that enzymes can undergo a change in conformation when they bind substrate molecules, and the active site has a shape complementary to that of the substrate only *after* the substrate is bound, as shown for hexokinase in Figure **17.6.3**. After catalysis, the enzyme resumes its original structure.




Figure **17.6.3**: The Induced-Fit Model of Enzyme Action. (a) The enzyme hexokinase without its substrate (glucose, shown in red) is bound to the active site. (b) The enzyme conformation changes dramatically when the substrate binds to it, resulting in additional interactions between hexokinase and glucose.

The structural changes that occur when an enzyme and a substrate join together bring specific parts of a substrate into alignment with specific parts of the enzyme's active site. Amino acid side chains in or near the binding site can then act as acid or base catalysts, provide binding sites for the transfer of functional groups from one substrate to another or aid in the rearrangement of a substrate. The participating amino acids, which are usually widely separated in the primary sequence of the protein, are brought close together in the active site as a result of the folding and bending of the polypeptide chain or chains when the protein acquires its tertiary and quaternary structure. Binding to enzymes brings reactants close to each other and aligns them properly, which has the same effect as increasing the concentration of the reacting compounds.

✔ Example 17.6.1

a. What type of interaction would occur between an OH group present on a substrate molecule and a functional group in the active site of an enzyme?

b. Suggest an amino acid whose side chain might be in the active site of an enzyme and form the type of interaction you just identified.

Solution

a. An OH group would most likely engage in hydrogen bonding with an appropriate functional group present in the active site of an enzyme.


b. Several amino acid side chains would be able to engage in hydrogen bonding with an OH group. One example would be asparagine, which has an amide functional group.

? Exercise 17.6.1

a. What type of interaction would occur between an COO^- group present on a substrate molecule and a functional group in the active site of an enzyme?

b. Suggest an amino acid whose side chain might be in the active site of an enzyme and form the type of interaction you just identified.

One characteristic that distinguishes an enzyme from all other types of catalysts is its *substrate specificity*. An inorganic acid such as sulfuric acid can be used to increase the reaction rates of many different reactions, such as the hydrolysis of disaccharides, polysaccharides, lipids, and proteins, with complete impartiality. In contrast, enzymes are much more specific. Some enzymes act on a single substrate, while other enzymes act on any of a group of related molecules containing a similar functional group or chemical bond. Some enzymes even distinguish between D- and L-stereoisomers, binding one stereoisomer but not the other. Urease, for example, is an enzyme that catalyzes the hydrolysis of a single substrate—urea—but not the closely related compounds methyl urea, thiourea, or biuret. The enzyme carboxypeptidase, on the other hand, is far less specific. It catalyzes the removal of nearly any amino acid from the carboxyl end of any peptide or protein.



Enzyme specificity results from the uniqueness of the active site in each different enzyme because of the identity, charge, and spatial orientation of the functional groups located there. It regulates cell chemistry so that the proper reactions occur in the proper place at the proper time. Clearly, it is crucial to the proper functioning of the living cell.

Summary

A substrate binds to a specific region on an enzyme known as the active site, where the substrate can be converted to product. The substrate binds to the enzyme primarily through hydrogen bonding and other electrostatic interactions. The induced-fit model says that an enzyme can undergo a conformational change when binding a substrate. Enzymes exhibit varying degrees of substrate specificity.

17.6: Enzymatic Catalysis is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.6: Enzyme Action** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.7: Enzymatic Activity

Learning Objectives

- To describe the interaction between an enzyme and its substrate.

Enzyme-catalyzed reactions occur in at least two steps. In the first step, an enzyme molecule (E) and the substrate molecule or molecules (S) collide and react to form an intermediate compound called the *enzyme-substrate* (E–S) *complex*. (This step is reversible because the complex can break apart into the original substrate or substrates and the free enzyme.) Once the E–S complex forms, the enzyme is able to catalyze the formation of product (P), which is then released from the enzyme surface:

$$S + E \rightarrow E\text{–}S \tag{17.7.1}$$

$$E\text{–}S \rightarrow P + E \tag{17.7.2}$$

Hydrogen bonding and other electrostatic interactions hold the enzyme and substrate together in the complex. The structural features or functional groups on the enzyme that participate in these interactions are located in a cleft or pocket on the enzyme surface. This pocket, where the enzyme combines with the substrate and transforms the substrate to product is called the active site of the enzyme (Figure **17.7.1**).




Figure **17.7.1**: Substrate Binding to the Active Site of an Enzyme. The enzyme dihydrofolate reductase is shown with one of its substrates: $NADP^+$ (a) unbound and (b) bound. The $NADP^+$ (shown in red) binds to a pocket that is complementary to it in shape and ionic properties.

The active site of an enzyme possesses a unique conformation (including correctly positioned bonding groups) that is complementary to the structure of the substrate, so that the enzyme and substrate molecules fit together in much the same manner as a key fits into a tumbler lock. In fact, an early model describing the formation of the enzyme-substrate complex was called the lock-and-key model (Figure **17.7.2**). This model portrayed the enzyme as conformationally rigid and able to bond only to substrates that exactly fit the active site.




Figure **17.7.2**: The Lock-and-Key Model of Enzyme Action. (a) Because the substrate and the active site of the enzyme have complementary structures and bonding groups, they fit together as a key fits a lock. (b) The catalytic reaction occurs while the two are bonded together in the enzyme-substrate complex.

Working out the precise three-dimensional structures of numerous enzymes has enabled chemists to refine the original lock-and-key model of enzyme actions. They discovered that the binding of a substrate often leads to a large conformational change in the enzyme, as well as to changes in the structure of the substrate or substrates. The current theory, known as the induced-fit model, says that enzymes can undergo a change in conformation when they bind substrate molecules, and the active site has a shape complementary to that of the substrate only *after* the substrate is bound, as shown for hexokinase in Figure **17.7.3**. After catalysis, the enzyme resumes its original structure.




Figure **17.7.3**: The Induced-Fit Model of Enzyme Action. (a) The enzyme hexokinase without its substrate (glucose, shown in red) is bound to the active site. (b) The enzyme conformation changes dramatically when the substrate binds to it, resulting in additional interactions between hexokinase and glucose.

The structural changes that occur when an enzyme and a substrate join together bring specific parts of a substrate into alignment with specific parts of the enzyme's active site. Amino acid side chains in or near the binding site can then act as acid or base catalysts, provide binding sites for the transfer of functional groups from one substrate to another or aid in the rearrangement of a substrate. The participating amino acids, which are usually widely separated in the primary sequence of the protein, are brought close together in the active site as a result of the folding and bending of the polypeptide chain or chains when the protein acquires its tertiary and quaternary structure. Binding to enzymes brings reactants close to each other and aligns them properly, which has the same effect as increasing the concentration of the reacting compounds.

✔ Example 17.7.1

a. What type of interaction would occur between an OH group present on a substrate molecule and a functional group in the active site of an enzyme?

b. Suggest an amino acid whose side chain might be in the active site of an enzyme and form the type of interaction you just identified.

Solution

a. An OH group would most likely engage in hydrogen bonding with an appropriate functional group present in the active site of an enzyme.


b. Several amino acid side chains would be able to engage in hydrogen bonding with an OH group. One example would be asparagine, which has an amide functional group.

? Exercise 17.7.1

a. What type of interaction would occur between an COO^- group present on a substrate molecule and a functional group in the active site of an enzyme?

b. Suggest an amino acid whose side chain might be in the active site of an enzyme and form the type of interaction you just identified.

One characteristic that distinguishes an enzyme from all other types of catalysts is its *substrate specificity*. An inorganic acid such as sulfuric acid can be used to increase the reaction rates of many different reactions, such as the hydrolysis of disaccharides, polysaccharides, lipids, and proteins, with complete impartiality. In contrast, enzymes are much more specific. Some enzymes act on a single substrate, while other enzymes act on any of a group of related molecules containing a similar functional group or chemical bond. Some enzymes even distinguish between D- and L-stereoisomers, binding one stereoisomer but not the other. Urease, for example, is an enzyme that catalyzes the hydrolysis of a single substrate—urea—but not the closely related compounds methyl urea, thiourea, or biuret. The enzyme carboxypeptidase, on the other hand, is far less specific. It catalyzes the removal of nearly any amino acid from the carboxyl end of any peptide or protein.



Enzyme specificity results from the uniqueness of the active site in each different enzyme because of the identity, charge, and spatial orientation of the functional groups located there. It regulates cell chemistry so that the proper reactions occur in the proper place at the proper time. Clearly, it is crucial to the proper functioning of the living cell.

Summary

A substrate binds to a specific region on an enzyme known as the active site, where the substrate can be converted to product. The substrate binds to the enzyme primarily through hydrogen bonding and other electrostatic interactions. The induced-fit model says that an enzyme can undergo a conformational change when binding a substrate. Enzymes exhibit varying degrees of substrate specificity.

17.7: Enzymatic Activity is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.6: Enzyme Action** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


17.8: Enzyme Inhibition

Learning Objectives

- Explain what an enzyme inhibitor is.
- Distinguish between reversible and irreversible inhibitors.
- Distinguish between competitive and noncompetitive inhibitors.


Previously, we noted that enzymes are inactivated at high temperatures and by changes in pH. These are nonspecific factors that would inactivate any enzyme. The activity of enzymes can also be regulated by more specific inhibitors. Many compounds are poisons because they bind covalently to particular enzymes or kinds of enzymes and inactivate them (Table **17.8.1**).

Table **17.8.1**: Poisons as Enzyme Inhibitors

Poison	Formula	Example of Enzyme Inhibited	Action
arsenate	AsO_4^{3-}	glyceraldehyde 3-phosphate dehydrogenase	substitutes for phosphate
iodoacetate	ICH_2COO^-	triose phosphate dehydrogenase	binds to cysteine **SH** group
diisopropylfluoro-phosphate (DIFP; a nerve poison)		acetylcholinesterase	binds to serine **OH** group

Irreversible Inhibition: Poisons

An irreversible inhibitor inactivates an enzyme by bonding covalently to a particular group at the active site. The inhibitor-enzyme bond is so strong that the inhibition cannot be reversed by the addition of excess substrate. The nerve gases, especially Diisopropyl fluorophosphate (DIFP), irreversibly inhibit biological systems by forming an enzyme-inhibitor complex with a specific OH group of serine situated at the active sites of certain enzymes. The peptidases trypsin and chymotrypsin contain serine groups at the active site and are inhibited by DIFP.



Reversible Inhibition

A reversible inhibitor inactivates an enzyme through noncovalent, more easily reversed, interactions. Unlike an irreversible inhibitor, a reversible inhibitor can dissociate from the enzyme. Reversible inhibitors include competitive inhibitors and noncompetitive inhibitors. (There are additional types of reversible inhibitors.) A competitive inhibitor is any compound that bears a structural resemblance to a particular substrate and thus competes with that substrate for binding at the active site of an enzyme. The inhibitor is not acted on by the enzyme but does prevent the substrate from approaching the active site.

The degree to which a competitive inhibitor interferes with an enzyme's activity depends on the relative concentrations of the substrate and the inhibitor. If the inhibitor is present in relatively large quantities, it will initially block most of the active sites. But because the binding is reversible, some substrate molecules will eventually bind to the active site and be converted to product. Increasing the substrate concentration promotes displacement of the inhibitor from the active site. Competitive inhibition can be completely reversed by adding substrate so that it reaches a much higher concentration than that of the inhibitor.

Studies of competitive inhibition have provided helpful information about certain enzyme-substrate complexes and the interactions of specific groups at the active sites. As a result, pharmaceutical companies have synthesized drugs that competitively inhibit metabolic processes in bacteria and certain cancer cells. Many drugs are competitive inhibitors of specific enzymes.

A classic example of competitive inhibition is the effect of malonate on the enzyme activity of succinate dehydrogenase (Figure 17.8.1). Malonate and succinate are the anions of dicarboxylic acids and contain three and four carbon atoms, respectively. The malonate molecule binds to the active site because the spacing of its carboxyl groups is not greatly different from that of succinate. However, no catalytic reaction occurs because malonate does not have a CH_2CH_2 group to convert to CH=CH. This reaction will also be discussed in connection with the Krebs cycle and energy production.




Figure 17.8.1: Competitive Inhibition. (a) Succinate binds to the enzyme succinate dehydrogenase. A dehydrogenation reaction occurs, and the product—fumarate—is released from the enzyme. (b) Malonate also binds to the active site of succinate dehydrogenase. In this case, however, no subsequent reaction occurs while malonate remains bound to the enzyme.

To Your Health: Penicillin

Chemotherapy is the strategic use of chemicals (that is, drugs) to destroy infectious microorganisms or cancer cells without causing excessive damage to the other, healthy cells of the host. From bacteria to humans, the metabolic pathways of all living organisms are quite similar, so the search for safe and effective chemotherapeutic agents is a formidable task. Many well-established chemotherapeutic drugs function by inhibiting a critical enzyme in the cells of the invading organism.

An *antibiotic* is a compound that kills bacteria; it may come from a natural source such as molds or be synthesized with a structure analogous to a naturally occurring antibacterial compound. Antibiotics constitute no well-defined class of chemically related substances, but many of them work by effectively inhibiting a variety of enzymes essential to bacterial growth.







Penicillin, one of the most widely used antibiotics in the world, was fortuitously discovered by Alexander Fleming in 1928, when he noticed antibacterial properties in a mold growing on a bacterial culture plate. In 1938, Ernst Chain and Howard Florey began an intensive effort to isolate penicillin from the mold and study its properties. The large quantities of penicillin needed for this research became available through development of a corn-based nutrient medium that the mold loved and through the discovery of a higher-yielding strain of mold at a United States Department of Agriculture research center near Peoria, Illinois. Even so, it was not until 1944 that large quantities of penicillin were being produced and made available for the treatment of bacterial infections.

Penicillin functions by interfering with the synthesis of cell walls of reproducing bacteria. It does so by inhibiting an enzyme—transpeptidase—that catalyzes the last step in bacterial cell-wall biosynthesis. The defective walls cause bacterial cells to burst. Human cells are not affected because they have cell membranes, not cell walls.

Several naturally occurring penicillins have been isolated. They are distinguished by different R groups connected to a common structure: a four-member cyclic amide (called a lactam ring) fused to a five-member ring. The addition of appropriate organic compounds to the culture medium leads to the production of the different kinds of penicillin.

The penicillins are effective against gram-positive bacteria (bacteria capable of being stained by Gram's stain) and a few gram-negative bacteria (including the intestinal bacterium *Escherichia coli*). They are effective in the treatment of diphtheria, gonorrhea, pneumonia, syphilis, many pus infections, and certain types of boils. Penicillin G was the earliest penicillin to be used on a wide scale. However, it cannot be administered orally because it is quite unstable; the acidic pH of the stomach

converts it to an inactive derivative. The major oral penicillins—penicillin V, ampicillin, and amoxicillin—on the other hand, are acid stable.

Penicillin Structure	R Group	Drug Name
		penicillin G
		penicillin V
		ampicillin
		amoxicillin
		methicillin

Some strains of bacteria become resistant to penicillin through a mutation that allows them to synthesize an enzyme—penicillinase—that breaks the antibiotic down (by cleavage of the amide linkage in the lactam ring). To combat these strains, scientists have synthesized penicillin analogs (such as methicillin) that are not inactivated by penicillinase.

Some people (perhaps 5% of the population) are allergic to penicillin and therefore must be treated with other antibiotics. Their allergic reaction can be so severe that a fatal coma may occur if penicillin is inadvertently administered to them. Fortunately, several other antibiotics have been discovered. Most, including aureomycin and streptomycin, are the products of microbial synthesis. Others, such as the semisynthetic penicillins and tetracyclines, are made by chemical modifications of antibiotics; and some, like chloramphenicol, are manufactured entirely by chemical synthesis. They are as effective as penicillin in destroying infectious microorganisms. Many of these antibiotics exert their effects by blocking protein synthesis in microorganisms.

Initially, antibiotics were considered miracle drugs, substantially reducing the number of deaths from blood poisoning, pneumonia, and other infectious diseases. Some seven decades ago, a person with a major infection almost always died. Today, such deaths are rare. Seven decades ago, pneumonia was a dreaded killer of people of all ages. Today, it kills only the very old or those ill from other causes. Antibiotics have indeed worked miracles in our time, but even miracle drugs have limitations. Not long after the drugs were first used, disease organisms began to develop strains resistant to them. In a race to stay ahead of resistant bacterial strains, scientists continue to seek new antibiotics. The penicillins have now been partially displaced by related compounds, such as the cephalosporins and vancomycin. Unfortunately, some strains of bacteria have already shown resistance to these antibiotics.

Some reversible inhibitors are noncompetitive. A noncompetitive inhibitor can combine with either the free enzyme or the enzyme-substrate complex because its binding site on the enzyme is distinct from the active site. Binding of this kind of inhibitor alters the three-dimensional conformation of the enzyme, changing the configuration of the active site with one of two results. Either the enzyme-substrate complex does not form at its normal rate, or, once formed, it does not yield products at the normal rate. Because the inhibitor does not structurally resemble the substrate, the addition of excess substrate does *not* reverse the inhibitory effect.




Figure **17.8.2**: Feedback Inhibition of Threonine Deaminase by Isoleucine. Threonine deaminase is the first enzyme in the conversion of threonine to isoleucine. Isoleucine inhibits threonine deaminase through feedback inhibition.

Feedback inhibition is a normal biochemical process that makes use of noncompetitive inhibitors to control some enzymatic activity. In this process, the final product inhibits the enzyme that catalyzes the first step in a series of reactions. Feedback inhibition is used to regulate the synthesis of many amino acids. For example, bacteria synthesize isoleucine from threonine in a series of five enzyme-catalyzed steps. As the concentration of isoleucine increases, some of it binds as a noncompetitive inhibitor to the first enzyme of the series (threonine deaminase), thus bringing about a decrease in the amount of isoleucine being formed (Figure **17.8.2**).

Summary

An irreversible inhibitor inactivates an enzyme by bonding covalently to a particular group at the active site. A reversible inhibitor inactivates an enzyme through noncovalent, reversible interactions. A competitive inhibitor competes with the substrate for binding at the active site of the enzyme. A noncompetitive inhibitor binds at a site distinct from the active site.

17.8: Enzyme Inhibition is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.8: Enzyme Inhibition** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

https://chem.libretexts.org/@go/page/387650

17.9: Enzyme Cofactors and Vitamins

Learning Objectives

- To explain why vitamins are necessary in the diet.

Many enzymes are simple proteins consisting entirely of one or more amino acid chains. Other enzymes contain a nonprotein component called a **cofactor** that is necessary for the enzyme's proper functioning. There are two types of cofactors: inorganic ions [e.g., zinc or Cu(I) ions] and organic molecules known as coenzymes. Most **coenzymes** are vitamins or are derived from vitamins.

Vitamins are organic compounds that are essential in very small (trace) amounts for the maintenance of normal metabolism. They generally cannot be synthesized at adequate levels by the body and must be obtained from the diet. The absence or shortage of a vitamin may result in a vitamin-deficiency disease. In the first half of the 20th century, a major focus of biochemistry was the identification, isolation, and characterization of vitamins. Despite accumulating evidence that people needed more than just carbohydrates, fats, and proteins in their diets for normal growth and health, it was not until the early 1900s that research established the need for trace nutrients in the diet.

Table **17.9.1**: Fat-Soluble Vitamins and Physiological Functions

Vitamin	Physiological Function	Effect of Deficiency
vitamin A (retinol)	formation of vision pigments; differentiation of epithelial cells	night blindness; continued deficiency leads to total blindness
vitamin D (cholecalciferol)	increases the body's ability to absorb calcium and phosphorus	osteomalacia (softening of the bones); known as rickets in children
vitamin E (tocopherol)	fat-soluble antioxidant	damage to cell membranes
vitamin K (phylloquinone)	formation of prothrombin, a key enzyme in the blood-clotting process	increases the time required for blood to clot

Because organisms differ in their synthetic abilities, a substance that is a vitamin for one species may not be so for another. Over the past 100 years, scientists have identified and isolated 13 vitamins required in the human diet and have divided them into two broad categories: the *fat-soluble vitamins*, which include vitamins A, D, E, and K, and the *water-soluble vitamins*, which are the B complex vitamins and vitamin C. All fat-soluble vitamins contain a high proportion of hydrocarbon structural components. There are one or two oxygen atoms present, but the compounds as a whole are nonpolar. In contrast, water-soluble vitamins contain large numbers of electronegative oxygen and nitrogen atoms, which can engage in hydrogen bonding with water. Most water-soluble vitamins act as coenzymes or are required for the synthesis of coenzymes. The fat-soluble vitamins are important for a variety of physiological functions. The key vitamins and their functions are found in Tables **17.9.1** and **17.9.2**.

Table **17.9.2**: Water-Soluble Vitamins and Physiological Functions

Vitamin	Coenzyme	Coenzyme Function	Deficiency Disease
vitamin B_1 (thiamine)	thiamine pyrophosphate	decarboxylation reactions	beri-beri
vitamin B_2 (riboflavin)	flavin mononucleotide or flavin adenine dinucleotide	oxidation-reduction reactions involving two hydrogen atoms	—
vitamin B_3 (niacin)	nicotinamide adenine dinucleotide or nicotinamide adenine dinucleotide phosphate	oxidation-reduction reactions involving the hydride ion (H^-)	pellagra
vitamin B_6 (pyridoxine)	pyridoxal phosphate	variety of reactions including the transfer of amino groups	—
vitamin B_{12} (cyanocobalamin)	methylcobalamin or deoxyadenoxylcobalamin	intramolecular rearrangement reactions	pernicious anemia
biotin	biotin	carboxylation reactions	—
folic acid	tetrahydrofolate	carrier of one-carbon units such as the formyl group	anemia
pantothenic Acid	coenzyme A	carrier of acyl groups	—
vitamin C (ascorbic acid)	none	antioxidant; formation of collagen, a protein found in tendons, ligaments, and bone	scurvy

Vitamins C and E, as well as the provitamin β-carotene can act as antioxidants in the body. Antioxidants prevent damage from free radicals, which are molecules that are highly reactive because they have unpaired electrons. Free radicals are formed not only through metabolic reactions involving oxygen but also by such environmental factors as radiation and pollution.

> *β-carotene is known as a provitamin because it can be converted to vitamin A in the body.*

Free radicals react most commonly react with lipoproteins and unsaturated fatty acids in cell membranes, removing an electron from those molecules and thus generating a new free radical. The process becomes a chain reaction that finally leads to the oxidative degradation of the affected compounds. Antioxidants react with free radicals to stop these chain reactions by forming a more stable molecule or, in the case of vitamin E, a free radical that is much less reactive (vitamin E is converted back to its original form through interaction with vitamin C).

Summary

Vitamins are organic compounds that are essential in very small amounts for the maintenance of normal metabolism. Vitamins are divided into two broad categories: fat-soluble vitamins and water-soluble vitamins. Most water-soluble vitamins are needed for the formation of coenzymes, which are organic molecules needed by some enzymes for catalytic activity.

17.9: Enzyme Cofactors and Vitamins is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.9: Enzyme Cofactors and Vitamins** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

17.10: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **protein** is a large biological polymer synthesized from **amino acids**, which are carboxylic acids containing an α-amino group. Proteins have a variety of important roles in living organisms, yet they are made from the same 20 L-amino acids. About half of these amino acids, the **essential amino acids**, cannot be synthesized by the human body and must be obtained from the diet. In the solid state and in neutral solutions, amino acids exist as **zwitterions**, species that are charged but electrically neutral. In this form, they behave much like inorganic salts. Each amino acid belongs to one of four classes depending on the characteristics of its R group or amino acid side chain: nonpolar, polar but neutral, positively charged, and negatively charged. Depending on the conditions, amino acids can act as either acids or bases, which means that proteins act as buffers. The pH at which an amino acid exists as the zwitterion is called the **isoelectric point (pI)**.

The amino acids in a protein are linked together by **peptide bonds**. Protein chains containing 10 or fewer amino acids are usually referred to as **peptides**, with a prefix such as di- or tri- indicating the number of amino acids. Chains containing more than 50 amino acid units are referred to as *proteins* or **polypeptides**. Proteins are classified globular or fibrous, depending on their structure and resulting solubility in water. **Globular proteins** are nearly spherical and are soluble in water; **fibrous proteins** have elongated or fibrous structures and are not soluble in water.

Protein molecules can have as many as four levels of structure. The **primary structure** is the sequence of amino acids in the chain. The **secondary structure** is the arrangement of adjacent atoms in the peptide chain; the most common arrangements are α-helices or β-pleated sheets. The **tertiary structure** is the overall three-dimensional shape of the molecule that results from the way the chain bends and folds in on itself. Proteins that consist of more than one chain have **quaternary structure**, which is the way the multiple chains are packed together.

Four types of intramolecular and intermolecular forces contribute to secondary, tertiary, and quaternary structure: (1) **hydrogen bonding** between an oxygen or a nitrogen atom and a hydrogen atom bound to an oxygen atom or a nitrogen atom, either on the same chain or on a neighboring chain; (2) **ionic bonding** between one positively charged side chain and one negatively charged side chain; (3) **disulfide linkages** between cysteine units; and (4) **dispersion forces** between nonpolar side chains.

Because of their complexity, protein molecules are delicate and easy to disrupt. A *denatured* protein is one whose conformation has been changed, in a process called **denaturation**, so that it can no longer do its physiological job. A variety of conditions, such as heat, ultraviolet radiation, the addition of organic compounds, or changes in pH can denature a protein.

An **enzyme** is an organic catalyst produced by a living cell. Enzymes are such powerful catalysts that the reactions they promote occur rapidly at body temperature. Without the help of enzymes, these reactions would require high temperatures and long reaction times.

The molecule or molecules on which an enzyme acts are called its **substrates**. An enzyme has an **active site** where its substrate or substrates bind to form an enzyme-substrate complex. The reaction occurs, and product is released:

$$E + S \rightarrow E\text{–}S \rightarrow E + P$$

The original **lock-and-key model** of enzyme and substrate binding pictured a rigid enzyme of unchanging configuration binding to the appropriate substrate. The newer **induced-fit model** describes the enzyme active site as changing its conformation after binding to the substrate.

Most enzymes have maximal activity in a narrow pH range centered on an **optimum pH**. In this pH range, the enzyme is correctly folded, and catalytic groups in the active site have the correct charge (positive, negative, or neutral). For most enzymes, the optimum pH is between 6 and 8.

Substances that interfere with enzyme function are called inhibitors. An **irreversible inhibitor** inactivates enzymes by forming covalent bonds to the enzyme, while a **reversible inhibitor** inactivates an enzyme by a weaker, noncovalent interaction that is easier to disrupt. A **competitive inhibitor** is a reversible inhibitor that is structurally similar to the substrate and binds to the active site. When the inhibitor is bound, the substrate is blocked from the active site and no reaction occurs. Because the binding of such

an inhibitor is reversible, a high substrate concentration will overcome the inhibition because it increases the likelihood of the substrate binding. A **noncompetitive inhibitor** binds reversibly at a site distinct from the active site. Thus, it can bind to either the enzyme or the enzyme-substrate complex. The inhibitor changes the conformation of the active site so that the enzyme cannot function properly. Noncompetitive inhibitors are important in **feedback inhibition**, in which the amount of product produced by a series of reactions is carefully controlled. The final product in a series of reactions acts as a noncompetitive inhibitor of the initial enzyme.

Simple enzymes consist entirely of one or more amino acid chains. Complex enzymes are composed of one or more amino acid chains joined to **cofactors**—inorganic ions or organic **coenzymes**. **Vitamins** are organic compounds that are essential in very small amounts for the maintenance of normal metabolism and generally cannot be synthesized at adequate levels by the body. Vitamins are divided into two broad categories: *fat-soluble* vitamins and *water-soluble* vitamins. Many of the water-soluble vitamins are used for the synthesis of coenzymes.

17.10: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **18.S: Amino Acids, Proteins, and Enzymes (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

18: Nucleic Acids

The blueprint for the reproduction and the maintenance of each organism is found in the nuclei of its cells, concentrated in elongated, threadlike structures called chromosomes. These complex structures, consisting of DNA and proteins, contain the basic units of heredity, called genes. The number of chromosomes (and genes) varies with each species. Human body cells have 23 pairs of chromosomes having 20,000–40,000 different genes.

Sperm and egg cells contain only a single copy of each chromosome; that is, they contain only one member of each chromosome pair. Thus, in sexual reproduction, the entire complement of chromosomes is achieved only when an egg and sperm combine. A new individual receives half its hereditary material from each parent. Calling the unit of heredity a “gene” merely gives it a name. But what really are genes and how is the information they contain expressed? One definition of a gene is that it is a segment of DNA that constitutes the code for a specific polypeptide. If genes are segments of DNA, we need to learn more about the structure and physiological function of DNA. We begin by looking at the small molecules needed to form DNA and RNA (ribonucleic acid)—the nucleotides.

18.1: Nucleotides
18.2: Nucleic Acid Structure
18.3: Replication and Expression of Genetic Information
18.4: Protein Synthesis and the Genetic Code
18.5: Mutations and Genetic Diseases
18.6: Viruses
18.7: Chapter Summary

18: Nucleic Acids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

18.1: Nucleotides


Learning Objectives

- To identify the different molecules that combine to form nucleotides.

The repeating, or monomer, units that are linked together to form nucleic acids are known as nucleotides. The deoxyribonucleic acid (DNA) of a typical mammalian cell contains about 3×10^9 nucleotides. Nucleotides can be further broken down to phosphoric acid (H_3PO_4), a pentose sugar (a sugar with five carbon atoms), and a nitrogenous base (a base containing nitrogen atoms).

$$\text{nucleic acids} \xrightarrow[\text{down into}]{\text{can be broken}} \text{nucleotides} \xrightarrow[\text{down into}]{\text{can be broken}} H_3PO_4 + \text{nitrogen base} + \text{pentose sugar}$$

If the pentose sugar is ribose, the nucleotide is more specifically referred to as a *ribonucleotide*, and the resulting nucleic acid is ribonucleic acid (RNA). If the sugar is 2-deoxyribose, the nucleotide is a *deoxyribonucleotide*, and the nucleic acid is DNA.



The nitrogenous bases found in nucleotides are classified as pyrimidines or purines. Pyrimidines are heterocyclic amines with two nitrogen atoms in a six-member ring and include uracil, thymine, and cytosine. Purines are heterocyclic amines consisting of a pyrimidine ring fused to a five-member ring with two nitrogen atoms. Adenine and guanine are the major purines found in nucleic acids (Figure **18.1.1**).




Figure **18.1.1**: The Nitrogenous Bases Found in DNA and RNA

The formation of a bond between C1′ of the pentose sugar and N1 of the pyrimidine base or N9 of the purine base joins the pentose sugar to the nitrogenous base. In the formation of this bond, a molecule of water is removed. Table **18.1.1** summarizes the similarities and differences in the composition of nucleotides in DNA and RNA.

> The numbering convention is that primed numbers designate the atoms of the pentose ring, and unprimed numbers designate the atoms of the purine or pyrimidine ring.

Table **18.1.1**: Composition of Nucleotides in DNA and RNA

Composition	DNA	RNA
purine bases	adenine and guanine	adenine and guanine
pyrimidine bases	cytosine and thymine	cytosine and uracil
pentose sugar	2-deoxyribose	ribose
inorganic acid	phosphoric acid (H_3PO_4)	H_3PO_4

The names and structures of the major ribonucleotides and one of the deoxyribonucleotides are given in Figure **18.1.2**.




Figure **18.1.2**: The Pyrimidine and Purine Nucleotides

Apart from being the monomer units of DNA and RNA, the nucleotides and some of their derivatives have other functions as well. Adenosine diphosphate (ADP) and adenosine triphosphate (ATP), shown in Figure **18.1.3**, have a role in cell metabolism. Moreover, a number of coenzymes, including flavin adenine dinucleotide (FAD), nicotinamide adenine dinucleotide (NAD^+), and coenzyme A, contain adenine nucleotides as structural components.




Figure **18.1.3**: Structures of Two Important Adenine-Containing Nucleotides

Summary

Nucleotides are composed of phosphoric acid, a pentose sugar (ribose or deoxyribose), and a nitrogen-containing base (adenine, cytosine, guanine, thymine, or uracil). Ribonucleotides contain ribose, while deoxyribonucleotides contain deoxyribose.

18.1: Nucleotides is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.1: Nucleotides** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

18.2: Nucleic Acid Structure

Skills to Develop

- Identify the two types of nucleic acids and the function of each type.
- Describe how nucleotides are linked together to form nucleic acids.
- Describe the secondary structure of DNA and the importance of complementary base pairing.

Nucleic acids are large polymers formed by linking nucleotides together and are found in every cell. Deoxyribonucleic acid (DNA) is the nucleic acid that stores genetic information. If all the DNA in a typical mammalian cell were stretched out end to end, it would extend more than 2 m. Ribonucleic acid (RNA) is the nucleic acid responsible for using the genetic information encoded in DNA to produce the thousands of proteins found in living organisms.

Primary Structure of Nucleic Acids

Nucleotides are joined together through the phosphate group of one nucleotide connecting in an ester linkage to the OH group on the third carbon atom of the sugar unit of a second nucleotide. This unit joins to a third nucleotide, and the process is repeated to produce a long nucleic acid chain (Figure 18.2.1). The backbone of the chain consists of alternating phosphate and sugar units (2-deoxyribose in DNA and ribose in RNA). The purine and pyrimidine bases branch off this backbone.

> *Each phosphate group has one acidic hydrogen atom that is ionized at physiological pH. This is why these compounds are known as nucleic acids.*




Figure 18.2.1 *Structure of a Segment of DNA. A similar segment of RNA would have OH groups on each C2', and uracil would replace thymine.*

Like proteins, nucleic acids have a primary structure that is defined as the sequence of their nucleotides. Unlike proteins, which have 20 different kinds of amino acids, there are only 4 different kinds of nucleotides in nucleic acids. For amino acid sequences in proteins, the convention is to write the amino acids in order starting with the N-terminal amino acid. In writing nucleotide sequences for nucleic acids, the convention is to write the nucleotides (usually using the one-letter abbreviations for the bases, shown in Figure 18.2.1) starting with the nucleotide having a free phosphate group, which is known as the 5′ end, and indicate the nucleotides in order. For DNA, a lowercase *d* is often written in front of the sequence to indicate that the monomers are

deoxyribonucleotides. The final nucleotide has a free OH group on the 3′ carbon atom and is called the *3′ end*. The sequence of nucleotides in the DNA segment shown in Figure 18.2.1 would be written 5′-dG-dT-dA-dC-3′, which is often further abbreviated to dGTAC or just GTAC.

Secondary Structure of DNA

The three-dimensional structure of DNA was the subject of an intensive research effort in the late 1940s to early 1950s. Initial work revealed that the polymer had a regular repeating structure. In 1950, Erwin Chargaff of Columbia University showed that the molar amount of adenine (A) in DNA was always equal to that of thymine (T). Similarly, he showed that the molar amount of guanine (G) was the same as that of cytosine (C). Chargaff drew no conclusions from his work, but others soon did.

At Cambridge University in 1953, James D. Watson and Francis Crick announced that they had a model for the secondary structure of DNA. Using the information from Chargaff's experiments (as well as other experiments) and data from the X ray studies of Rosalind Franklin (which involved sophisticated chemistry, physics, and mathematics), Watson and Crick worked with models that were not unlike a child's construction set and finally concluded that DNA is composed of two nucleic acid chains running antiparallel to one another—that is, side-by-side with the 5′ end of one chain next to the 3′ end of the other. Moreover, as their model showed, the two chains are twisted to form a double helix—a structure that can be compared to a spiral staircase, with the phosphate and sugar groups (the backbone of the nucleic acid polymer) representing the outside edges of the staircase. The purine and pyrimidine bases face the inside of the helix, with guanine always opposite cytosine and adenine always opposite thymine. These specific base pairs, referred to as complementary bases, are the steps, or treads, in our staircase analogy (Figure 18.2.2).




Figure 18.2.2 *DNA Double Helix. (a) This represents a computer-generated model of the DNA double helix. (b) This represents a schematic representation of the double helix, showing the complementary bases.*

The structure proposed by Watson and Crick provided clues to the mechanisms by which cells are able to divide into two identical, functioning daughter cells; how genetic data are passed to new generations; and even how proteins are built to required specifications. All these abilities depend on the pairing of complementary bases. Figure 18.2.3 shows the two sets of base pairs and illustrates two things. First, a pyrimidine is paired with a purine in each case, so that the long dimensions of both pairs are identical (1.08 nm).




***Figure* 18.2.3** *Complementary Base Pairing. Complementary bases engage in hydrogen bonding with one another: (a) thymine and adenine; (b) cytosine and guanine.*

If two pyrimidines were paired or two purines were paired, the two pyrimidines would take up less space than a purine and a pyrimidine, and the two purines would take up more space, as illustrated in Figure **18.2.4**. If these pairings were ever to occur, the structure of DNA would be like a staircase made with stairs of different widths. For the two strands of the double helix to fit neatly, a pyrimidine must always be paired with a purine. The second thing you should notice in Figure **18.2.3** is that the correct pairing enables formation of three instances of hydrogen bonding between guanine and cytosine and two between adenine and thymine. The additive contribution of this hydrogen bonding imparts great stability to the DNA double helix.




***Figure* 18.2.4** *Difference in Widths of Possible Base Pairs*

Summary

- DNA is the nucleic acid that stores genetic information. RNA is the nucleic acid responsible for using the genetic information in DNA to produce proteins.
- Nucleotides are joined together to form nucleic acids through the phosphate group of one nucleotide connecting in an ester linkage to the OH group on the third carbon atom of the sugar unit of a second nucleotide.
- Nucleic acid sequences are written starting with the nucleotide having a free phosphate group (the 5′ end).
- Two DNA strands link together in an antiparallel direction and are twisted to form a double helix. The nitrogenous bases face the inside of the helix. Guanine is always opposite cytosine, and adenine is always opposite thymine.

Concept Review Exercises


1.
   a. Name the two kinds of nucleic acids.
   b. Which type of nucleic acid stores genetic information in the cell?
2. What are complementary bases?
3. Why is it structurally important that a purine base always pair with a pyrimidine base in the DNA double helix?

Answers


1.
   a. deoxyribonucleic acid (DNA) and ribonucleic acid (RNA)
   b. DNA
2. the specific base pairings in the DNA double helix in which guanine is paired with cytosine and adenine is paired with thymine
3. The width of the DNA double helix is kept at a constant width, rather than narrowing (if two pyrimidines were across from each other) or widening (if two purines were across from each other).

Exercises

1. For this short RNA segment,
   a. identify the 5′ end and the 3′ end of the molecule.
   b. circle the atoms that comprise the backbone of the nucleic acid chain.
   c. write the nucleotide sequence of this RNA segment.



2. For this short DNA segment,
   a. identify the 5′ end and the 3′ end of the molecule.
   b. circle the atoms that comprise the backbone of the nucleic acid chain.
   c. write the nucleotide sequence of this DNA segment.



3. Which nitrogenous base in DNA pairs with each nitrogenous base?

   a. cytosine
   b. adenine
   c. guanine
   d. thymine

4. Which nitrogenous base in RNA pairs with each nitrogenous base?

   a. cytosine
   b. adenine
   c. guanine
   d. thymine

5. How many hydrogen bonds can form between the two strands in the short DNA segment shown below?


   5′ ATGCGACTA 3′ 3′ TACGCTGAT 5′

6. How many hydrogen bonds can form between the two strands in the short DNA segment shown below?

   5′ CGATGAGCC 3′ 3′ GCTACTCGG 5′

Answers

1.



c. ACU

3.

a. guanine
b. thymine
c. cytosine
d. adenine

5. 22 (2 between each AT base pair and 3 between each GC base pair)

18.2: Nucleic Acid Structure is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

18.3: Replication and Expression of Genetic Information

Learning Objectives

- Describe how a new copy of DNA is synthesized.
- Describe how RNA is synthesized from DNA.
- Identify the different types of RNA and the function of each type of RNA.

We previously stated that deoxyribonucleic acid (DNA) stores genetic information, while ribonucleic acid (RNA) is responsible for transmitting or expressing genetic information by directing the synthesis of thousands of proteins found in living organisms. But how do the nucleic acids perform these functions? Three processes are required: (1) *replication*, in which new copies of DNA are made; (2) *transcription*, in which a segment of DNA is used to produce RNA; and (3) *translation*, in which the information in RNA is translated into a protein sequence.

Replication

New cells are continuously forming in the body through the process of cell division. For this to happen, the DNA in a dividing cell must be copied in a process known as replication. The complementary base pairing of the double helix provides a ready model for how genetic replication occurs. If the two chains of the double helix are pulled apart, disrupting the hydrogen bonding between base pairs, each chain can act as a *template*, or pattern, for the synthesis of a new complementary DNA chain.

The nucleus contains all the necessary enzymes, proteins, and nucleotides required for this synthesis. A short segment of DNA is "unzipped," so that the two strands in the segment are separated to serve as templates for new DNA. DNA polymerase, an enzyme, recognizes each base in a template strand and matches it to the complementary base in a free nucleotide. The enzyme then catalyzes the formation of an ester bond between the 5′ phosphate group of the nucleotide and the 3′ OH end of the new, growing DNA chain. In this way, each strand of the original DNA molecule is used to produce a duplicate of its former partner (Figure 18.3.1). Whatever information was encoded in the original DNA double helix is now contained in each replicate helix. When the cell divides, each daughter cell gets one of these replicates and thus all of the information that was originally possessed by the parent cell.




Figure 18.3.1: A Schematic Diagram of DNA Replication. DNA replication occurs by the sequential unzipping of segments of the double helix. Each new nucleotide is brought into position by DNA polymerase and is added to the growing strand by the formation of a phosphate ester bond. Thus, two double helixes form from one, and each consists of one old strand and one new strand, an outcome called *semiconservative replications*. (This representation is simplified; many more proteins are involved in replication.)

Example 18.3.1

A segment of one strand from a DNA molecule has the sequence 5′-TCCATGAGTTGA-3′. What is the sequence of nucleotides in the opposite, or complementary, DNA chain?

Solution

Knowing that the two strands are antiparallel and that T base pairs with A, while C base pairs with G, the sequence of the complementary strand will be 3′-AGGTACTCAACT-5′ (can also be written as TCAACTCATGGA).

? Exercise 18.3.1

A segment of one strand from a DNA molecule has the sequence 5′-CCAGTGAATTGCCTAT-3′. What is the sequence of nucleotides in the opposite, or complementary, DNA chain?

What do we mean when we say information is encoded in the DNA molecule? An organism's DNA can be compared to a book containing directions for assembling a model airplane or for knitting a sweater. Letters of the alphabet are arranged into words, and these words direct the individual to perform certain operations with specific materials. If all the directions are followed correctly, a model airplane or sweater is produced.

In DNA, the particular sequences of nucleotides along the chains encode the directions for building an organism. Just as *saw* means one thing in English and *was* means another, the sequence of bases CGT means one thing, and TGC means something different. Although there are only four letters—the four nucleotides—in the genetic code of DNA, their sequencing along the DNA strands can vary so widely that information storage is essentially unlimited.

Transcription

For the hereditary information in DNA to be useful, it must be "expressed," that is, used to direct the growth and functioning of an organism. The first step in the processes that constitute DNA expression is the synthesis of RNA, by a template mechanism that is in many ways analogous to DNA replication. Because the RNA that is synthesized is a complementary copy of information contained in DNA, RNA synthesis is referred to as transcription. There are three key differences between replication and transcription:

1. RNA molecules are much shorter than DNA molecules; only a portion of one DNA strand is copied or transcribed to make an RNA molecule.
2. RNA is built from ribonucleotides rather than deoxyribonucleotides.
3. The newly synthesized RNA strand does not remain associated with the DNA sequence it was transcribed from.

The DNA sequence that is transcribed to make RNA is called the *template strand*, while the complementary sequence on the other DNA strand is called the *coding* or *informational strand*. To initiate RNA synthesis, the two DNA strands unwind at specific sites along the DNA molecule. Ribonucleotides are attracted to the uncoiling region of the DNA molecule, beginning at the 3′ end of the template strand, according to the rules of base pairing. Thymine in DNA calls for adenine in RNA, cytosine specifies guanine, guanine calls for cytosine, and adenine requires uracil. RNA polymerase—an enzyme—binds the complementary ribonucleotide and catalyzes the formation of the ester linkage between ribonucleotides, a reaction very similar to that catalyzed by DNA polymerase (Figure **18.3.2**). Synthesis of the RNA strand takes place in the 5′ to 3′ direction, antiparallel to the template strand. Only a short segment of the RNA molecule is hydrogen-bonded to the template strand at any time during transcription. When transcription is completed, the RNA is released, and the DNA helix reforms. The nucleotide sequence of the RNA strand formed during transcription is identical to that of the corresponding coding strand of the DNA, except that U replaces T.




Figure 18.3.2: A Schematic Diagram of RNA Transcription from a DNA Template. The representation of RNA polymerase is proportionately much smaller than the actual molecule, which encompasses about 50 nucleotides at a time.

Example 18.3.2

A portion of the template strand of a gene has the sequence 5′-TCCATGAGTTGA-3′. What is the sequence of nucleotides in the RNA that is formed from this template?

Solution

Four things must be remembered in answering this question: (1) the DNA strand and the RNA strand being synthesized are antiparallel; (2) RNA is synthesized in a 5′ to 3′ direction, so transcription begins at the 3′ end of the template strand; (3) ribonucleotides are used in place of deoxyribonucleotides; and (4) thymine (T) base pairs with adenine (A), A base pairs with uracil (U; in RNA), and cytosine (C) base pairs with guanine (G). The sequence is determined to be 3′-AGGUACUCAACU-5′ (can also be written as 5′-UCAACUCAUGGA-3′).

Exercise 18.3.2

A portion of the template strand of a gene has the sequence 5′-CCAGTGAATTGCCTAT-3′. What is the sequence of nucleotides in the RNA that is formed from this template?

Three types of RNA are formed during transcription: *messenger RNA* (mRNA), *ribosomal RNA* (rRNA), and *transfer RNA* (tRNA). These three types of RNA differ in function, size, and percentage of the total cell RNA (Table 18.3.1). mRNA makes up only a small percent of the total amount of RNA within the cell, primarily because each molecule of mRNA exists for a relatively short time; it is continuously being degraded and resynthesized. The molecular dimensions of the mRNA molecule vary according to the amount of genetic information a given molecule contains. After transcription, which takes place in the nucleus, the mRNA passes into the cytoplasm, carrying the genetic message from DNA to the ribosomes, the sites of protein synthesis. Elsewhere, we shall see how mRNA directly determines the sequence of amino acids during protein synthesis.

Table 18.3.1: Properties of Cellular RNA in Escherichia coli

Type	Function	Approximate Number of Nucleotides	Percentage of Total Cell RNA
mRNA	codes for proteins	100–6,000	~3
rRNA	component of ribosomes	120–2900	83
tRNA	adapter molecule that brings the amino acid to the ribosome	75–90	14

Ribosomes are cellular substructures where proteins are synthesized. They contain about 65% rRNA and 35% protein, held together by numerous noncovalent interactions, such as hydrogen bonding, in an overall structure consisting of two globular particles of unequal size.

Molecules of tRNA, which bring amino acids (one at a time) to the ribosomes for the construction of proteins, differ from one another in the kinds of amino acid each is specifically designed to carry. A set of three nucleotides, known as a codon, on the mRNA determines which kind of tRNA will add its amino acid to the growing chain. Each of the 20 amino acids found in proteins has at least one corresponding kind of tRNA, and most amino acids have more than one.




Figure 18.3.3: Transfer RNA. (a) In the two-dimensional structure of a yeast tRNA molecule for phenylalanine, the amino acid binds to the acceptor stem located at the 3′ end of the tRNA primary sequence. (The nucleotides that are not specifically identified here are slightly altered analogs of the four common ribonucleotides A, U, C, and G.) (b) In the three-dimensional structure of yeast phenylalanine tRNA, note that the anticodon loop is at the bottom and the acceptor stem is at the top right. (c) This shows a space-filling model of the tRNA.

The two-dimensional structure of a tRNA molecule has three distinctive loops, reminiscent of a cloverleaf (Figure 18.3.3). On one loop is a sequence of three nucleotides that varies for each kind of tRNA. This triplet, called the anticodon, is complementary to and pairs with the codon on the mRNA. At the opposite end of the molecule is the acceptor stem, where the amino acid is attached.

Summary

- In DNA replication, each strand of the original DNA serves as a template for the synthesis of a complementary strand.
- DNA polymerase is the primary enzyme needed for replication.
- In transcription, a segment of DNA serves as a template for the synthesis of an RNA sequence.
- RNA polymerase is the primary enzyme needed for transcription.
- Three types of RNA are formed during transcription: mRNA, rRNA, and tRNA.

18.3: Replication and Expression of Genetic Information is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.3: Replication and Expression of Genetic Information** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

18.4: Protein Synthesis and the Genetic Code

Learning Objectives

- describe the characteristics of the genetic code.
- describe how a protein is synthesized from mRNA.

One of the definitions of a gene is as follows: a segment of deoxyribonucleic acid (DNA) carrying the code for a specific polypeptide. Each molecule of messenger RNA (mRNA) is a transcribed copy of a gene that is used by a cell for synthesizing a polypeptide chain. If a protein contains two or more different polypeptide chains, each chain is coded by a different gene. We turn now to the question of how the sequence of nucleotides in a molecule of ribonucleic acid (RNA) is translated into an amino acid sequence.

How can a molecule containing just 4 different nucleotides specify the sequence of the 20 amino acids that occur in proteins? If each nucleotide coded for 1 amino acid, then obviously the nucleic acids could code for only 4 amino acids. What if amino acids were coded for by groups of 2 nucleotides? There are 4^2, or 16, different combinations of 2 nucleotides (AA, AU, AC, AG, UU, and so on). Such a code is more extensive but still not adequate to code for 20 amino acids. However, if the nucleotides are arranged in groups of 3, the number of different possible combinations is 4^3, or 64. Here we have a code that is extensive enough to direct the synthesis of the primary structure of a protein molecule.



***Video**: NDSU Virtual Cell Animations project animation "Translation". For more information, see VCell, NDSU(opens in new window) [vcell.ndsu.nodak.edu]*

The genetic code can therefore be described as the identification of each group of three nucleotides and its particular amino acid. The sequence of these triplet groups in the mRNA dictates the sequence of the amino acids in the protein. Each individual three-nucleotide coding unit, as we have seen, is called a *codon*.

Protein synthesis is accomplished by orderly interactions between mRNA and the other ribonucleic acids (transfer RNA [tRNA] and ribosomal RNA [rRNA]), the ribosome, and more than 100 enzymes. The mRNA formed in the nucleus during transcription is transported across the nuclear membrane into the cytoplasm to the ribosomes—carrying with it the genetic instructions. The process in which the information encoded in the mRNA is used to direct the sequencing of amino acids and thus ultimately to synthesize a protein is referred to as translation.




Figure 18.4.1: Binding of an Amino Acid to Its tRNA

Before an amino acid can be incorporated into a polypeptide chain, it must be attached to its unique tRNA. This crucial process requires an enzyme known as aminoacyl-tRNA synthetase (Figure 18.4.1). There is a specific aminoacyl-tRNA synthetase for each amino acid. This high degree of specificity is vital to the incorporation of the correct amino acid into a protein. After the amino acid molecule has been bound to its tRNA carrier, protein synthesis can take place. Figure 18.4.2 depicts a schematic stepwise representation of this all-important process.




Figure 18.4.2a: The Elongation Steps in Protein Synthesis - Protein synthesis is already in progress at the ribosome. The growing polypeptide chain is attached to the tRNA that brought in the previous amino acid (in this illustration, cys.)




Figure 18.4.2b: The Elongation Steps in Protein Synthesis - An activated tRNA, which has the anticodon AAA, binds to the ribosome next to the previous bound tRNA and interacts with the mRNA molecule though basepairing of the codon and anticodon. The amino acid Phe is being incorporated into the polypeptide chain by the formation of a peptide linkage between the carboxyl group of Cys and the amino acid group of the Phe. This reaction is catalyzed by the enzyme peptidyl transferase, a component of the ribosome.




Figure 18.4.2*c*: The Elongation Steps in Protein Synthesis - The Cys-Phe linkage is now complete, and the growing polypeptide chain remains attached to the tRNA for Phe.




Figure 18.4.2*d*: The Elongation Steps in Protein Synthesis - The ribosome moves to the right along the mRNA strand. This shift brings the next codon, GUC, into its correct position on the surface of the ribosome. Note that an activated tRNA molecule, containing the next amino acid to be attached to the chain is moving to the ribosome. Steps (b)-(d) will be repeated until the ribosome reaches a stop codon.

Early experimenters were faced with the task of determining which of the 64 possible codons stood for each of the 20 amino acids. The cracking of the genetic code was the joint accomplishment of several well-known geneticists—notably Har Khorana, Marshall Nirenberg, Philip Leder, and Severo Ochoa—from 1961 to 1964. The genetic dictionary they compiled, summarized in Figure 18.4.3, shows that 61 codons code for amino acids, and 3 codons serve as signals for the termination of polypeptide synthesis (much like the period at the end of a sentence). Notice that only methionine (AUG) and tryptophan (UGG) have single codons. All other amino acids have two or more codons.




Figure **18.4.3**: The Genetic Code

Example 18.4.1: Using the Genetic Code

A portion of an mRNA molecule has the sequence 5′-AUGCCACGAGUUGAC-3′. What amino acid sequence does this code for?

Solution

Use Figure **18.4.3** to determine what amino acid each set of three nucleotides (codon) codes for. Remember that the sequence is read starting from the 5′ end and that a protein is synthesized starting with the N-terminal amino acid. The sequence 5′-AUGCCACGAGUUGAC-3′ codes for met-pro-arg-val-asp.

Exercise 18.4.4

A portion of an RNA molecule has the sequence 5′-AUGCUGAAUUGCGUAGGA-3′. What amino acid sequence does this code for?

Further experimentation threw much light on the nature of the genetic code, as follows:

1. The code is virtually universal; animal, plant, and bacterial cells use the same codons to specify each amino acid (with a few exceptions).
2. The code is “degenerate”; in all but two cases (methionine and tryptophan), more than one triplet codes for a given amino acid.
3. The first two bases of each codon are most significant; the third base often varies. This suggests that a change in the third base by a mutation may still permit the correct incorporation of a given amino acid into a protein. The third base is sometimes called the “wobble” base.
4. The code is continuous and nonoverlapping; there are *no* nucleotides between codons, and adjacent codons do not overlap.
5. The three termination codons are read by special proteins called release factors, which signal the end of the translation process.
6. The codon AUG codes for methionine and is also the initiation codon. Thus methionine is the first amino acid in each newly synthesized polypeptide. This first amino acid is usually removed enzymatically before the polypeptide chain is completed; the vast majority of polypeptides do not begin with methionine.

Summary

In translation, the information in mRNA directs the order of amino acids in protein synthesis. A set of three nucleotides (codon) codes for a specific amino acid.

18.4: Protein Synthesis and the Genetic Code is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.4: Protein Synthesis and the Genetic Code** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

18.5: Mutations and Genetic Diseases

Learning Objectives

- To describe the causes of genetic mutations and how they lead to genetic diseases.

We have seen that the sequence of nucleotides in a cell's deoxyribonucleic acid (DNA) is what ultimately determines the sequence of amino acids in proteins made by the cell and thus is critical for the proper functioning of the cell. On rare occasions, however, the nucleotide sequence in DNA may be modified either spontaneously (by errors during replication, occurring approximately once for every 10 billion nucleotides) or from exposure to heat, radiation, or certain chemicals. Any chemical or physical change that alters the nucleotide sequence in DNA is called a mutation. When a mutation occurs in an egg or sperm cell that then produces a living organism, it will be inherited by all the offspring of that organism.

Common types of mutations include **substitution** (a different nucleotide is substituted), **insertion** (the addition of a new nucleotide), and **deletion** (the loss of a nucleotide). These changes within DNA are called point mutations because only one nucleotide is substituted, added, or deleted (Figure 18.5.1). Because an insertion or deletion results in a frame-shift that changes the reading of subsequent codons and, therefore, alters the entire amino acid sequence that follows the mutation, insertions and deletions are usually more harmful than a substitution in which only a single amino acid is altered.




Figure 18.5.1: Three Types of Point Mutations

The chemical or physical agents that cause mutations are called mutagens. Examples of physical mutagens are ultraviolet (UV) and gamma radiation. Radiation exerts its mutagenic effect either directly or by creating free radicals that in turn have mutagenic effects. Radiation and free radicals can lead to the formation of bonds between nitrogenous bases in DNA. For example, exposure to UV light can result in the formation of a covalent bond between two adjacent thymines on a DNA strand, producing a thymine dimer (Figure 18.5.2). If not repaired, the dimer prevents the formation of the double helix at the point where it occurs. The genetic disease *xeroderma pigmentosum* is caused by a lack of the enzyme that cuts out the thymine dimers in damaged DNA. Individuals affected by this condition are abnormally sensitive to light and are more prone to skin cancer than normal individuals.





Figure 18.5.2: An Example of Radiation Damage to DNA. (a) The thymine dimer is formed by the action of UV light. (b) When a defect in the double strand is produced by the thymine dimer, this defect temporarily stops DNA replication, but the dimer can be removed, and the region can be repaired by an enzyme repair system.

Sometimes gene mutations are beneficial, but most of them are detrimental. For example, if a point mutation occurs at a crucial position in a DNA sequence, the affected protein will lack biological activity, perhaps resulting in the death of a cell. In such cases the altered DNA sequence is lost and will not be copied into daughter cells. Nonlethal mutations in an egg or sperm cell may lead to metabolic abnormalities or hereditary diseases. Such diseases are called *inborn errors of metabolism* or genetic diseases. A partial listing of genetic diseases is presented in Figure 18.5.1, and two specific diseases are discussed in the following sections. In most cases, the defective gene results in a failure to synthesize a particular enzyme.


Figure 18.5.1: Some Representative Genetic Diseases in Humans and the Protein or Enzyme Responsible

Disease	Responsible Protein or Enzyme
alkaptonuria	homogentisic acid oxidase
galactosemia	galactose 1-phosphate uridyl transferase, galactokinase, or UDP galactose epimerase
Gaucher disease	glucocerebrosidase
gout and Lesch-Nyhan syndrome	hypoxanthine-guanine phosphoribosyl transferase
hemophilia	antihemophilic factor (factor VIII) or Christmas factor (factor IX)
homocystinuria	cystathionine synthetase
maple syrup urine disease	branched chain α-keto acid dehydrogenase complex
McArdle syndrome	muscle phosphorylase
Niemann-Pick disease	sphingomyelinase
phenylketonuria (PKU)	phenylalanine hydroxylase
sickle cell anemia	hemoglobin
Tay-Sachs disease	hexosaminidase A
tyrosinemia	fumarylacetoacetate hydrolase or tyrosine aminotransferase
von Gierke disease	glucose 6-phosphatase
Wilson disease	Wilson disease protein

PKU results from the absence of the enzyme phenylalanine hydroxylase. Without this enzyme, a person cannot convert phenylalanine to tyrosine, which is the precursor of the neurotransmitters dopamine and norepinephrine as well as the skin pigment melanin.



When this reaction cannot occur, phenylalanine accumulates and is then converted to higher than normal quantities of phenylpyruvate. The disease acquired its name from the high levels of phenylpyruvate (a phenyl ketone) in urine. Excessive amounts of phenylpyruvate impair normal brain development, which causes severe mental retardation.



PKU may be diagnosed by assaying a sample of blood or urine for phenylalanine or one of its metabolites. Medical authorities recommend testing every newborn's blood for phenylalanine within 24 h to 3 weeks after birth. If the condition is detected, mental retardation can be prevented by immediately placing the infant on a diet containing little or no phenylalanine. Because phenylalanine is plentiful in naturally produced proteins, the low-phenylalanine diet depends on a synthetic protein substitute plus very small measured amounts of naturally produced foods. Before dietary treatment was introduced in the early 1960s, severe mental retardation was a common outcome for children with PKU. Prior to the 1960s, 85% of patients with PKU had an intelligence quotient (IQ) less than 40, and 37% had IQ scores below 10. Since the introduction of dietary treatments, however, over 95% of children with PKU have developed normal or near-normal intelligence. The incidence of PKU in newborns is about 1 in 12,000 in North America.


> *Every state in the United States has mandated that screening for PKU be provided to all newborns.*

Several genetic diseases are collectively categorized as *lipid-storage diseases*. Lipids are constantly being synthesized and broken down in the body, so if the enzymes that catalyze lipid degradation are missing, the lipids tend to accumulate and cause a variety of medical problems. When a genetic mutation occurs in the gene for the enzyme hexosaminidase A, for example, gangliosides cannot be degraded but accumulate in brain tissue, causing the ganglion cells of the brain to become greatly enlarged and nonfunctional. This genetic disease, known as *Tay-Sachs disease*, leads to a regression in development, dementia, paralysis, and blindness, with death usually occurring before the age of three. There is currently no treatment, but Tay-Sachs disease can be diagnosed in a fetus by assaying the amniotic fluid (amniocentesis) for hexosaminidase A. A blood test can identify Tay-Sachs carriers—people who inherit a defective gene from only one rather than both parents—because they produce only half the normal amount of hexosaminidase A, although they do not exhibit symptoms of the disease.

Looking Closer: Recombinant DNA Technology

More than 3,000 human diseases have been shown to have a genetic component, caused or in some way modulated by the person's genetic composition. Moreover, in the last decade or so, researchers have succeeded in identifying many of the genes and even mutations that are responsible for specific genetic diseases. Now scientists have found ways of identifying and isolating genes that have specific biological functions and placing those genes in another organism, such as a bacterium, which can be easily grown in culture. With these techniques, known as *recombinant DNA technology*, the ability to cure many serious genetic diseases appears to be within our grasp.


Isolating the specific gene or genes that cause a particular genetic disease is a monumental task. One reason for the difficulty is the enormous amount of a cell's DNA, only a minute portion of which contains the gene sequence. Thus, the first task is to obtain smaller pieces of DNA that can be more easily handled. Fortunately, researchers are able to use *restriction enzymes* (also known as restriction endonucleases), discovered in 1970, which are enzymes that cut DNA at specific, known nucleotide sequences, yielding DNA fragments of shorter length. For example, the restriction enzyme *EcoRI* recognizes the nucleotide sequence shown here and cuts both DNA strands as indicated:



Once a DNA strand has been fragmented, it must be cloned; that is, multiple identical copies of each DNA fragment are produced to make sure there are sufficient amounts of each to detect and manipulate in the laboratory. Cloning is accomplished by inserting the individual DNA fragments into phages (bacterial viruses) that can enter bacterial cells and be replicated. When a bacterial cell infected by the modified phage is placed in an appropriate culture medium, it forms a colony of cells, all containing copies of the original DNA fragment. This technique is used to produce many bacterial colonies, each containing a different DNA fragment. The result is a *DNA library*, a collection of bacterial colonies that together contain the entire genome of a particular organism.

The next task is to screen the DNA library to determine which bacterial colony (or colonies) has incorporated the DNA fragment containing the desired gene. A short piece of DNA, known as a *hybridization probe*, which has a nucleotide sequence complementary to a known sequence in the gene, is synthesized, and a radioactive phosphate group is added to it as a "tag." You might be wondering how researchers are able to prepare such a probe if the gene has not yet been isolated. One way is to use a segment of the desired gene isolated from another organism. An alternative method depends on knowing all or part of the amino acid sequence of the protein produced by the gene of interest: the amino acid sequence is used to produce an approximate genetic code for the gene, and this nucleotide sequence is then produced synthetically. (The amino acid sequence used is carefully chosen to include, if possible, many amino acids such as methionine and tryptophan, which have only a single codon each.)

After a probe identifies a colony containing the desired gene, the DNA fragment is clipped out, again using restriction enzymes, and spliced into another replicating entity, usually a plasmid. *Plasmids* are tiny mini-chromosomes found in many bacteria, such as *Escherichia coli* (*E. coli*). A recombined plasmid would then be inserted into the host organism (usually the bacterium *E. coli*), where it would go to work to produce the desired protein.



Proponents of recombinant DNA research are excited about its great potential benefits. An example is the production of human growth hormone, which is used to treat children who fail to grow properly. Formerly, human growth hormone was available only in tiny amounts obtained from cadavers. Now it is readily available through recombinant DNA technology. Another gene that has been cloned is the gene for epidermal growth factor, which stimulates the growth of skin cells and can be used to speed the healing of burns and other skin wounds. Recombinant techniques are also a powerful research tool, providing enormous aid to scientists as they map and sequence genes and determine the functions of different segments of an organism's DNA.

In addition to advancements in the ongoing treatment of genetic diseases, recombinant DNA technology may actually lead to cures. When appropriate genes are successfully inserted into *E. coli*, the bacteria can become miniature pharmaceutical factories, producing great quantities of insulin for people with diabetes, clotting factor for people with hemophilia, missing enzymes, hormones, vitamins, antibodies, vaccines, and so on. Recent accomplishments include the production in *E. coli* of recombinant DNA molecules containing synthetic genes for tissue plasminogen activator, a clot-dissolving enzyme that can rescue heart attack victims, as well as the production of vaccines against hepatitis B (humans) and hoof-and-mouth disease (cattle).

Scientists have used other bacteria besides *E. coli* in gene-splicing experiments and also yeast and fungi. Plant molecular biologists use a bacterial plasmid to introduce genes for several foreign proteins (including animal proteins) into plants. The bacterium is *Agrobacterium tumefaciens*, which can cause tumors in many plants, but which can be treated so that its tumor-causing ability is eliminated. One practical application of its plasmids would be to enhance a plant's nutritional value by transferring into it the gene necessary for the synthesis of an amino acid in which the plant is normally deficient (for example, transferring the gene for methionine synthesis into pinto beans, which normally do not synthesize high levels of methionine).

> Restriction enzymes have been isolated from a number of bacteria and are named after the bacterium of origin. *EcoRI* is a restriction enzyme obtained from the R strain of *E. coli*. The roman numeral I indicates that it was the first restriction enzyme obtained from this strain of bacteria.

Summary

- The nucleotide sequence in DNA may be modified either spontaneously or from exposure to heat, radiation, or certain chemicals and can lead to mutations.
- Mutagens are the chemical or physical agents that cause mutations.
- Genetic diseases are hereditary diseases that occur because of a mutation in a critical gene.

18.5: Mutations and Genetic Diseases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.5: Mutations and Genetic Diseases** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

18.6: Viruses

Learning Objectives

- To explain how viruses reproduce in cells.

Viruses are visible only under an electron microscope. They come in a variety of shapes, ranging from spherical to rod shaped. The fact that they contain either deoxyribonucleic acid (DNA) or ribonucleic acid (RNA)—*but never both*—allows them to be divided into two major classes: DNA viruses and RNA viruses (Figure **18.6.1**).




Figure **18.6.1**: Viruses. Viruses come in a variety of shapes that are determined by their protein coats.

Most *RNA viruses* use their nucleic acids in much the same way as the DNA viruses, penetrating a host cell and inducing it to replicate the viral RNA and synthesize viral proteins. The new RNA strands and viral proteins are then assembled into new viruses. Some RNA viruses, however, called retroviruses (Figure **18.6.2**), synthesize DNA in the host cell, in a process that is the reverse of the DNA-to-RNA transcription that normally occurs in cells. The synthesis of DNA from an RNA template is catalyzed by the enzyme reverse transcriptase.






Figure **18.6.2**: Life Cycle of a Retrovirus

In 1987, **azidothymidine** (AZT, also known as zidovudine or the brand name Retrovir) became the first drug approved for the treatment of AIDS. It works by binding to reverse transcriptase in place of deoxythymidine triphosphate, after which, because AZT does not have a 3′OH group, further replication is blocked. In the past 10 years, several other drugs have been approved that also act by inhibiting the viral reverse transcriptase.



Raltegravir (Isentress) is a newer anti-AIDS drug that was approved by the FDA in October 2007. This drug inhibits the integrase enzyme that is needed to integrate the HIV DNA into cellular DNA, an essential step in the production of more HIV particles.



A major problem in treating HIV infections is that the virus can become resistant to any of these drugs. One way to combat the problem has been to administer a “cocktail” of drugs, typically a combination of two reverse transcriptase inhibitors along with a protease inhibitor. These treatments can significantly reduce the amount of HIV in an infected person.

Career Focus: Genetics Counselor

A genetics counselor works with individuals and families who have birth defects or genetic disorders or a family history of a disease, such as cancer, with a genetic link. A genetics counselor may work in a variety of health-care settings (such as a hospital) to obtain family medical and reproductive histories; explain how genetic conditions are inherited; explain the causes, diagnosis, and care of these conditions; interpret the results of genetic tests; and aid the individual or family in making decisions regarding genetic diseases or conditions. A certified genetics counselor must obtain a master’s degree from an accredited program. Applicants to these graduate programs usually have an undergraduate degree in biology, psychology, or genetics.



Photo courtesy of the United States National Institutes for Health, commons.wikimedia.org/wiki/File:Geneticcounseling.jpg.

Summary

Viruses are very small infectious agents that contain either DNA or RNA as their genetic material. The human immunodeficiency virus (HIV) causes acquired immunodeficiency syndrome (AIDS).

18.6: Viruses is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.6: Viruses** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

18.7: Chapter Summary

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A cell's hereditary information is encoded in **chromosomes** in the cell's nucleus. Each chromosome is composed of proteins and **deoxyribonucleic acid (DNA)**. The chromosomes contain smaller hereditary units called **genes,** which are relatively short segments of DNA. The hereditary information is expressed or used through the synthesis of **ribonucleic acid (RNA)**. Both **nucleic acids**—DNA and RNA—are polymers composed of monomers known as **nucleotides**, which in turn consist of phosphoric acid (H_3PO_4), a nitrogenous base, and a pentose sugar.

The two types of *nitrogenous bases* most important in nucleic acids are **purines**—adenine (A) and guanine (G)—and **pyrimidines**—cytosine (C), thymine (T), and uracil (U). DNA contains the nitrogenous bases adenine, cytosine, guanine, and thymine, while the bases in RNA are adenine, cytosine, guanine, and uracil. The sugar in the nucleotides of RNA is ribose; the one in DNA is 2-deoxyribose. The sequence of nucleotides in a nucleic acid defines the primary structure of the molecule.

RNA is a single-chain nucleic acid, whereas DNA possesses two nucleic-acid chains intertwined in a secondary structure called a **double helix**. The sugar-phosphate backbone forms the outside the double helix, with the purine and pyrimidine bases tucked inside. Hydrogen bonding between **complementary bases** holds the two strands of the double helix together; A always pairs with T and C always pairs with G.

Cell growth requires **replication**, or reproduction of the cell's DNA. The double helix unwinds, and hydrogen bonding between complementary bases breaks so that there are two single strands of DNA, and each strand is a *template* for the synthesis of a new strand. For protein synthesis, three types of RNA are needed: *messenger RNA* (mRNA), *ribosomal RNA* (rRNA), and *transfer RNA* (tRNA). All are made from a DNA template by a process called **transcription**. The double helix uncoils, and ribonucleotides base-pair to the deoxyribonucleotides on one DNA strand; however, RNA is produced using *uracil* rather than thymine. Once the RNA is formed, it dissociates from the template and leaves the nucleus, and the DNA double helix reforms.

Translation is the process in which proteins are synthesized from the information in mRNA. It occurs at structures called **ribosomes**, which are located outside the nucleus and are composed of rRNA and protein. The 64 possible three-nucleotide combinations of the 4 nucleotides of DNA constitute the **genetic code** that dictates the sequence in which amino acids are joined to make proteins. Each three-nucleotide sequence on mRNA is a **codon**. Each kind of tRNA molecule binds a specific amino acid and has a site containing a three-nucleotide sequence called an **anticodon**.

The general term for any change in the genetic code in an organism's DNA is **mutation**. A change in which a single base is substituted, inserted, or deleted is a **point mutation**. The chemical and/or physical agents that cause mutations are called **mutagens**. Diseases that occur due to mutations in critical DNA sequences are referred to as **genetic diseases**.

Viruses are infectious agents composed of a tightly packed central core of nucleic acids enclosed by a protective shell of proteins. Viruses contain either DNA or RNA as their genetic material but not both. Some RNA viruses, called **retroviruses**, synthesize DNA in the host cell from their RNA genome. The human immunodeficiency virus (HIV) causes acquired immunodeficiency syndrome (AIDS).

18.7: Chapter Summary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **19.S: Nucleic Acids (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

CHAPTER OVERVIEW

19: Energy Metabolism

Metabolism is the set of life-sustaining chemical transformations within the cells of living organisms. The three main purposes of metabolism are the conversion of food/fuel to energy to run cellular processes, the conversion of food/fuel to building blocks for proteins, lipids, nucleic acids, and some carbohydrates, and the elimination of nitrogenous wastes. These enzyme-catalyzed reactions allow organisms to grow and reproduce, maintain their structures, and respond to their environments. Metabolism is usually divided into two categories: **catabolism**, the breaking down of organic matter, for example, by cellular respiration, and **anabolism**, the building up of components of cells such as proteins and nucleic acids. Usually, breaking down releases energy and building up consumes energy.

19.1: Prelude to Energy Metabolism
19.2: ATP- the Universal Energy Currency
19.3: Stage I of Catabolism
19.4: Overview of Stage II of Catabolism
19.5: Stage III of Catabolism
19.6: Stage II of Carbohydrate Catabolism
19.7: Stage II of Lipid Catabolism
19.8: Stage II of Protein Catabolism
19.9: Oxidative Phosphorylation
19.10: Energy Metabolism (Exercises)
19.11: Energy Metabolism (Summary)

19: Energy Metabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

19.1: Prelude to Energy Metabolism

The discovery of the link between insulin and diabetes led to a period of intense research aimed at understanding exactly how insulin works in the body to regulate glucose levels. Hormones in general act by binding to some protein, known as the hormone's receptor, thus initiating a series of events that lead to a desired outcome. In the early 1970s, the insulin receptor was purified, and researchers began to study what happens after insulin binds to its receptor and how those events are linked to the uptake and metabolism of glucose in cells.

The insulin receptor is located in the cell membrane and consists of four polypeptide chains: two identical chains called α chains and two identical chains called β chains. The α chains, positioned on the outer surface of the membrane, consist of 735 amino acids each and contain the binding site for insulin. The β chains are integral membrane proteins, each composed of 620 amino acids. The binding of insulin to its receptor stimulates the β chains to catalyze the addition of phosphate groups to the specific side chains of tyrosine (referred to as phosphorylation) in the β chains and other cell proteins, leading to the activation of reactions that metabolize glucose. In this chapter we will look at the pathway that breaks down glucose—in response to activation by insulin—for the purpose of providing energy for the cell.

GLmol

Figure **19.1.1**: Model of the Structure of the Insulin Receptor (PDB code 4ZXB).

Life requires energy. Animals, for example, require heat energy to maintain body temperature, mechanical energy to move their limbs, and chemical energy to synthesize the compounds needed by their cells. Living cells remain organized and functioning properly only through a continual supply of energy. But only specific forms of energy can be used. Supplying a plant with energy by holding it in a flame will not prolong its life. On the other hand, a green plant is able to absorb radiant energy from the sun, the most abundant source of energy for life on the earth. Plants use this energy first to form glucose and then to make other carbohydrates, as well as lipids and proteins. Unlike plants, animals cannot directly use the sun's energy to synthesize new compounds. They must eat plants or other animals to get carbohydrates, fats, and proteins and the chemical energy stored in them. Once digested and transported to the cells, the nutrient molecules can be used in either of two ways: as building blocks for making new cell parts or repairing old ones or "burned" for energy.




Figure **19.1.2**: Some Energy Transformations in Living Systems. Plants and animals exist in a cycle; each requires products of the other.

The thousands of coordinated chemical reactions that keep cells alive are referred to collectively as **metabolism**. In general, metabolic reactions are divided into two classes: the breaking down of molecules to obtain energy is **catabolism**, and the building of new molecules needed by living systems is **anabolism**.

Definition: Metabolite

Any chemical compound that participates in a metabolic reaction is a *metabolite*.

Most of the energy required by animals is generated from lipids and carbohydrates. These fuels must be oxidized, or "burned," for the energy to be released. The oxidation process ultimately converts the lipid or carbohydrate to carbon dioxide (CO_2) and water (H_2O).

Carbohydrate:

$$C_6H_{12}O_6 + 6\,O_2 \longrightarrow 6\,CO_2 + 6\,H_2O + 670\,\text{kcal}$$

Lipid:

$$C_{16}H_{32}O_2 + 23\,O_2 \longrightarrow 16\,CO_2 + 16\,H_2O + 2{,}385\,\text{kcal}$$

These two equations summarize the biological combustion of a carbohydrate and a lipid by the cell through respiration. **Respiration** is the collective name for all metabolic processes in which gaseous oxygen is used to oxidize organic matter to carbon dioxide, water, and energy.

Like the combustion of the common fuels we burn in our homes and cars (wood, coal, gasoline), respiration uses oxygen from the air to break down complex organic substances to carbon dioxide and water. But the energy released in the burning of wood is manifested entirely in the form of heat, and excess heat energy is not only useless but also injurious to the living cell. Living organisms instead conserve much of the energy respiration releases by channeling it into a series of stepwise reactions that produce adenosine triphosphate (ATP) or other compounds that ultimately lead to the synthesis of ATP. The remainder of the energy is released as heat and manifested as body temperature.

19.1: Prelude to Energy Metabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


- **20.0: Prelude to Energy Metabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.2: ATP- the Universal Energy Currency


Learning Objectives

- To describe the importance of ATP as a source of energy in living organisms.

Adenosine triphosphate (ATP), a nucleotide composed of adenine, ribose, and three phosphate groups, is perhaps the most important of the so-called energy-rich compounds in a cell. Its concentration in the cell varies from 0.5 to 2.5 mg/mL of cell fluid.



Energy-rich compounds are substances having particular structural features that lead to a release of energy after hydrolysis. As a result, these compounds are able to supply energy for biochemical processes that require energy. The structural feature important in ATP is the phosphoric acid anhydride, or pyrophosphate, linkage:



The pyrophosphate bond, symbolized by a squiggle (~), is hydrolyzed when ATP is converted to adenosine diphosphate (ADP). In this hydrolysis reaction, the products contain less energy than the reactants; there is a release of energy (> 7 kcal/mol). One reason for the amount of energy released is that hydrolysis relieves the electron-electron repulsions experienced by the negatively charged phosphate groups when they are bonded to each other (Figure 20.1.1).




Figure **19.2.1**: Hydrolysis of ATP to Form ADP

Energy is released because the products (ADP and phosphate ion) have less energy than the reactants [ATP and water (H_2O)].

The general equation for ATP hydrolysis is as follows:





$$ATP + H_2O \rightarrow ADP + P_i + 7.4\ kcal/mol$$

If the hydrolysis of ATP releases energy, its synthesis (from ADP) requires energy. In the cell, ATP is produced by those processes that supply energy to the organism (absorption of radiant energy from the sun in green plants and breakdown of food in animals), and it is hydrolyzed by those processes that require energy (the syntheses of carbohydrates, lipids, proteins; the transmission of nerve impulses; muscle contractions). In fact, ATP is the principal medium of energy exchange in biological systems. Many scientists call it the energy currency of cells.

> *P_i is the symbol for the inorganic phosphate anions $H_2PO_4^-$ and HPO_4^{2-}.*

ATP is not the only high-energy compound needed for metabolism. Several others are listed in Table **19.2.1**. Notice, however, that the energy released when ATP is hydrolyzed is approximately midway between those of the high-energy and the low-energy phosphate compounds. This means that the hydrolysis of ATP can provide energy for the phosphorylation of the compounds below it in the table. For example, the hydrolysis of ATP provides sufficient energy for the phosphorylation of glucose to form glucose 1-phosphate. By the same token, the hydrolysis of compounds, such as creatine phosphate, that appear *above* ATP in the table can provide the energy needed to resynthesize ATP from ADP.

Table **19.2.1**: Energy Released by Hydrolysis of Some Phosphate Compounds

Type	Example	Energy Released (kcal/mol)
acyl phosphate 	1,3-bisphosphoglycerate (BPG)	−11.8
	acetyl phosphate	−11.3
guanidine phosphates 	creatine phosphate	−10.3
	arginine phosphate	−9.1
pyrophosphates 	$PP_{i*} \rightarrow 2P_i$	−7.8
	$ATP \rightarrow AMP + PP_i$	−7.7
	$ATP \rightarrow ADP + P_i$	−7.5
	$ADP \rightarrow AMP + P_i$	−7.5
sugar phosphates 	glucose 1-phosphate	−5.0
	fructose 6-phosphate	−3.8
	AMP → adenosine + P_i	−3.4
	glucose 6-phosphate	−3.3
	glycerol 3-phosphate	−2.2

*PP_i is the pyrophosphate ion.

Summary

The hydrolysis of ATP releases energy that can be used for cellular processes that require energy.

19.2: ATP- the Universal Energy Currency is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.1: ATP- the Universal Energy Currency** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.3: Stage I of Catabolism

Learning Objectives

- To describe how carbohydrates, fats, and proteins are broken down during digestion.

We have said that animals obtain chemical energy from the food—carbohydrates, fats, and proteins—they eat through reactions defined collectively as *catabolism*. We can think of catabolism as occurring in three stages (Figure **19.3.1**). In stage I, carbohydrates, fats, and proteins are broken down into their individual monomer units: carbohydrates into simple sugars, fats into fatty acids and glycerol, and proteins into amino acids. One part of stage I of catabolism is the breakdown of food molecules by hydrolysis reactions into the individual monomer units—which occurs in the mouth, stomach, and small intestine—and is referred to as digestion.

In stage II, these monomer units (or building blocks) are further broken down through different reaction pathways, one of which produces ATP, to form a common end product that can then be used in stage III to produce even more ATP. In this chapter, we will look at each stage of catabolism—as an overview and in detail.




Figure 19.3.1: Energy Conversions

The conversion of food into cellular energy (as ATP) occurs in three stages.

Digestion of Carbohydrates

Carbohydrate digestion begins in the mouth (Figure 19.3.2) where salivary α-amylase attacks the α-glycosidic linkages in starch, the main carbohydrate ingested by humans. Cleavage of the glycosidic linkages produces a mixture of dextrins, maltose, and glucose. The α-amylase mixed into the food remains active as the food passes through the esophagus, but it is rapidly inactivated in the acidic environment of the stomach.




Figure **19.3.2**: The Principal Events and Sites of Carbohydrate Digestion

The primary site of carbohydrate digestion is the small intestine. The secretion of α-amylase in the small intestine converts any remaining starch molecules, as well as the dextrins, to maltose. Maltose is then cleaved into two glucose molecules by maltase. Disaccharides such as sucrose and lactose are not digested until they reach the small intestine, where they are acted on by sucrase and lactase, respectively. The major products of the complete hydrolysis of disaccharides and polysaccharides are three monosaccharide units: glucose, fructose, and galactose. These are absorbed through the wall of the small intestine into the bloodstream.

Digestion of Proteins

Protein digestion begins in the stomach (Figure **19.3.3**), where the action of gastric juice hydrolyzes about 10% of the peptide bonds. Gastric juice is a mixture of water (more than 99%), inorganic ions, hydrochloric acid, and various enzymes and other proteins.

> The pain of a gastric ulcer is at least partially due to irritation of the ulcerated tissue by acidic gastric juice.




Figure **19.3.3**: The Principal Events and Sites of Protein Digestion

The hydrochloric acid (HCl) in gastric juice is secreted by glands in the stomach lining. The pH of freshly secreted gastric juice is about 1.0, but the contents of the stomach may raise the pH to between 1.5 and 2.5. HCl helps to denature food proteins; that is, it unfolds the protein molecules to expose their chains to more efficient enzyme action. The principal digestive component of gastric juice is pepsinogen, an inactive enzyme produced in cells located in the stomach wall. When food enters the stomach after a period of fasting, pepsinogen is converted to its active form—pepsin—in a series of steps initiated by the drop in pH. Pepsin catalyzes the hydrolysis of peptide linkages within protein molecules. It has a fairly broad specificity but acts preferentially on linkages involving the aromatic amino acids tryptophan, tyrosine, and phenylalanine, as well as methionine and leucine.

Protein digestion is completed in the small intestine. Pancreatic juice, carried from the pancreas via the pancreatic duct, contains inactive enzymes such as trypsinogen and chymotrypsinogen. They are activated in the small intestine as follows (Figure **19.3.4**): The intestinal mucosal cells secrete the proteolytic enzyme enteropeptidase, which converts trypsinogen to trypsin; trypsin then activates chymotrypsinogen to chymotrypsin (and also completes the activation of trypsinogen). Both of these active enzymes catalyze the hydrolysis of peptide bonds in protein chains. Chymotrypsin preferentially attacks peptide bonds involving the carboxyl groups of the aromatic amino acids (phenylalanine, tryptophan, and tyrosine). Trypsin attacks peptide bonds involving the carboxyl groups of the basic amino acids (lysine and arginine). Pancreatic juice also contains procarboxypeptidase, which is cleaved by trypsin to carboxypeptidase. The latter is an enzyme that catalyzes the hydrolysis of peptide linkages at the free carboxyl end of the peptide chain, resulting in the stepwise liberation of free amino acids from the carboxyl end of the polypeptide.




Figure 19.3.4: Activation of Some Pancreatic Enzymes in the Small Intestine

Aminopeptidases in the intestinal juice remove amino acids from the N-terminal end of peptides and proteins possessing a free amino group. Figure 19.3.5 illustrates the specificity of these protein-digesting enzymes. The amino acids that are released by protein digestion are absorbed across the intestinal wall into the circulatory system, where they can be used for protein synthesis.




Figure 19.3.5: Hydrolysis of a Peptide by Several Peptidases

This diagram illustrates where in a peptide the different peptidases we have discussed would catalyze hydrolysis the peptide bonds.

Digestion of Lipids

Lipid digestion begins in the upper portion of the small intestine (Figure 19.3.6). A hormone secreted in this region stimulates the gallbladder to discharge bile into the duodenum. The principal constituents of bile are the bile salts, which emulsify large, water-insoluble lipid droplets, disrupting some of the hydrophobic interactions holding the lipid molecules together and suspending the resulting smaller globules (micelles) in the aqueous digestive medium. These changes greatly increase the surface area of the lipid particles, allowing for more intimate contact with the lipases and thus rapid digestion of the fats. Another hormone promotes the secretion of pancreatic juice, which contains these enzymes.





Figure **19.3.6**: The Principal Events and Sites of Lipid (Primarily Triglyceride) Digestion

The lipases in pancreatic juice catalyze the digestion of triglycerides first to diglycerides and then to 2-monoglycerides and fatty acids:



The monoglycerides and fatty acids cross the intestinal lining into the bloodstream, where they are resynthesized into triglycerides and transported as lipoprotein complexes known as chylomicrons. Phospholipids and cholesteryl esters undergo similar hydrolysis in the small intestine, and their component molecules are also absorbed through the intestinal lining.

The further metabolism of monosaccharides, fatty acids, and amino acids released in stage I of catabolism occurs in stages II and III of catabolism.

Summary

During digestion, carbohydrates are broken down into monosaccharides, proteins are broken down into amino acids, and triglycerides are broken down into glycerol and fatty acids. Most of the digestion reactions occur in the small intestine.

19.3: Stage I of Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.2: Stage I of Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.4: Overview of Stage II of Catabolism

Learning Objectives

- To describe the role of acetyl-CoA in metabolism.

A metabolic pathway is a series of biochemical reactions by which an organism converts a given reactant to a specific end product. There are specific metabolic pathways—which are different for carbohydrates, triglycerides, and proteins—that break down the products of stage I of catabolism (monosaccharides, fatty acids, and amino acids) to produce a common end product, acetyl-coenzyme A (acetyl-CoA) in stage II of catabolism.




Figure 19.4.1: The Structure of Acetyl-Coenzyme A (Acetyl-CoA)

Acetyl-CoA is shown in Figure 19.4.1. The acetyl unit, derived (as we will see) from the breakdown of carbohydrates, lipids, and proteins, is attached to coenzyme A, making the acetyl unit more reactive. Acetyl-CoA is used in a myriad of biochemical pathways. For example, it may be used as the starting material for the biosynthesis of lipids (such as triglycerides, phospholipids, or cholesterol and other steroids). Most importantly for energy generation, it may enter the citric acid cycle and be oxidized to produce energy, if energy is needed and oxygen is available. The various fates or uses of acetyl-CoA are summarized in Figure 19.4.1.




Figure 19.4.2: Cell Chemistry. Acetyl-CoA plays a variety of roles in cell chemistry.

Glycolysis

Glycolysis is the catabolic process in which glucose is converted into pyruvate via ten enzymatic steps. There are three regulatory steps, each of which is highly regulated that are separated into two phases:

1. the "priming phase" because it requires an input of energy in the form of 2 ATPs per glucose molecule and
2. the "pay off phase" because energy is released in the form of 4 ATPs, 2 per glyceraldehyde molecule.

The end result of Glycolysis is two new pyruvate molecules which can then be fed into the Citric Acid cycle (also known as the Kreb's Cycle) if oxygen is present, or can be reduced to lactate or ethanol in the absence of of oxygen using a process known as fermentation.



Video 19.4.1: Glycolysis: An Overview. Glycolysis is a series of 10 reactions that converts sugars, like glucose, into 3-carbon molecules called pyruvate. This animation provides an overview of the energy consumed and produced by the pathway. NDSU VCell Production's animation; for more information please see Vcell, NDSU(opens in new window) [vcell.ndsu.edu].

Glycolysis occurs within almost all living cells and is the primary source of Acetyl-CoA, which is the molecule responsible for the majority of energy output under aerobic conditions. The structures of Glycolysis intermediates can be found in Figure 19.4.3.




Figure 19.4.3: Glycolysis pathway. (CC BY-SA 4.0; International; Thomas Shafee via Wikipedia)

Phase 1: The "Priming Step"

The first phase of Glycolysis requires an input of energy in the form of ATP (adenosine triphosphate).

1. alpha-D-Glucose is phosphorolated at the 6 carbon by ATP via the enzyme Hexokinase (Class: Transferase) to yield alpha-D-Glucose-6-phosphate (G-6-P). This is a regulatory step which is negatively regulated by the presence of glucose-6-phosphate.
2. alpha-D-Glucose-6-phosphate is then converted into D-Fructose-6-phosphate (F-6-P) by Phosphoglucoisomerase (Class: Isomerase)
3. D-Fructose-6-phosphate is once again phosphorolated this time at the 1 carbon position by ATP via the enzyme Phosphofructokinase (Class: Transferase) to yield D-Fructose-1,6-bisphosphate (FBP). This is the committed step of glycolysis because of its large ΔG value.
4. D-Fructose-1,6-bisphosphate is then cleaved into two, three carbon molecules; Dihydroxyacetone phosphate (DHAP) and D-Glyceraldehyde-3-phosphate (G-3-P) by the enzyme Fructose bisphosphate aldolase (Class: Lyase)

5. Because the next portion of Glycolysis requires the molecule D-Glyceraldehyde-3-phosphate to continue Dihydroxyacetone phosphate is converted into D-Glyceraldehyde-3-phosphate by the enzyme Triose phosphate isomerase (Class: Isomerase)

Phase 2: The "Pay Off Step"

The second phase of Glycolysis where 4 molecules of ATP are produced per molecule of glucose. Enzymes appear in red:

1. D-Glyceraldehyde-3-phosphate is phosphorolated at the 1 carbon by the enzyme Glyceraldehyde-3-phosphate dehodrogenase to yield the high energy molecule 1,3-Bisphosphoglycerate (BPG)
2. ADP is then phosphorolated at the expense of 1,3-Bisphosphoglycerate by the enzyme Phosphoglycerate kinase (Class: Transferase) to yield ATP and 3-Phosphoglycerate (3-PG)
3. 3-Phosphoglycerate is then converted into 2-Phosphoglycerate by Phosphoglycerate mutase in preparation to yield another high energy molecule
4. 2-Phosphoglycerate is then converted to phosphoenolpyruvate (PEP) by Enolase. H_2O, potassium, and magnesium are all released as a result.
5. ADP is once again phosphorolated, this time at the expense of PEP by the enzyme pyruvate kinase to yield another molecule of ATP and and pyruvate. This step is regulated by the energy in the cell. The higher the energy of the cell the more inhibited pyruvate kinase becomes. Indicators of high energy levels within the cell are high concentrations of ATP, Acetyl-CoA, Alanine, and cAMP.

Because Glucose is split to yield two molecules of D-Glyceraldehyde-3-phosphate, each step in the "Pay Off" phase occurs twice per molecule of glucose.

Beta-Oxidation

The best source of energy for eukaryotic organisms are fats. Glucose offers a ratio 6.3 moles of ATP per carbon while saturated fatty acids offer 8.1 ATP per carbon. Also the complete oxidation of fats yields enormous amounts of water for those organisms that do not have adequate access to drinkable water. Camels and killer whales are good example of this, they obtain their water requirements from the complete oxidation of fats.



Video 19.4.2: Fatty acid metabolism / beta oxidation / β-Oxidation

There are four distinct stages in the oxidation of fatty acids. Fatty acid degradation takes place within the mitochondria and requires the help of several different enzymes. In order for fatty acids to enter the mitochondria the assistance of two carrier proteins is required, Carnitine acyltransferase I and II. It is also interesting to note the similarities between the four steps of beta-oxidation and the later four steps of the TCA cycle.

Entry into Beta-oxidation

Most fats stored in eukaryotic organisms are stored as triglycerides as seen below. In order to enter into beta-oxidation bonds must be broken usually with the use of a Lipase. The end result of these broken bonds are a glycerol molecule and three fatty acids in the case of triglycerides. Other lipids are capable of being degraded as well.





Figure 19.4.4: Key molecules in beta-oxidation: (left) A triglyceride molecule, (middle) Glycerol, (right) Fatty Acids (unsaturated)

Activation Step

- Once the triglycerides are broken down into glycerol and fatty acids they must be activated before they can enter into the mitochondria and proceed on with beta-oxidation. This is done by Acyl-CoA synthetase to yield fatty acyl-CoA.
- After the fatty acid has been acylated it is now ready to enter into the mitochondria.
- There are two carrier proteins (Carnitine acyltransferase I and II), one located on the outer membrane and one on the inner membrane of the mitochondria. Both are required for entry of the Acyl-CoA into the mitochondria.
- Once inside the mitochondria the fatty acyl-CoA can enter into beta-oxidation.

Oxidation Step


A fatty acyl-CoA is oxidized by Acyl-CoA dehydrogenase to yield a trans alkene. This is done with the aid of an [FAD] prosthetic group.



Acyl-CoA trans-Δ^2-Enoyl-CoA

Hydration Step


The trans alkene is then hydrated with the help of Enoyl-CoA hydratase



trans-Δ^2-Enoyl-CoA Enoyl-CoA-Hydratase L-3-Hydroxyacyl-CoA

Oxidation Step


The alcohol of the hydroxyacly-CoA is then oxidized by NAD^+ to a carbonyl with the help of Hydroxyacyl-CoA dehydrogenase. NAD^+ is used to oxidize the alcohol rather then [FAD] because NAD^+ is capable of the alcohol while [FAD] is not.



L-3-Hydroxyacyl-CoA 3-Ketoacyl-CoA

Cleavage

Finally acetyl-CoA is cleaved off with the help of Thiolase to yield an Acyl-CoA that is two carbons shorter than before. The cleaved acetyl-CoA can then enter into the TCA and ETC because it is already within the mitochondria.



3-Ketoacyl-CoA | Acyl-CoA | Acetyl-CoA

Summary

Acetyl-CoA is formed from the breakdown of carbohydrates, lipids, and proteins. It is used in many biochemical pathways.

References

1. Garrett, H., Reginald and Charles Grisham. Biochemistry. Boston: Twayne Publishers, 2008.
2. Raven, Peter. Biology. Boston: Twayne Publishers, 2005.

19.4: Overview of Stage II of Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.3: Overview of Stage II of Catabolism** by Darik Benson is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.5: Stage III of Catabolism

Learning Objectives

- Describe the reactions of the citric acid cycle.
- Describe the function of the citric acid cycle and identify the products produced.
- Describe the role of the electron transport chain in energy metabolism.
- Describe the role of oxidative phosphorylation in energy metabolism.

The acetyl group enters a cyclic sequence of reactions known collectively as the citric acid cycle (or Krebs cycle or tricarboxylic acid [TCA] cycle). The cyclical design of this complex series of reactions, which bring about the oxidation of the acetyl group of acetyl-CoA to carbon dioxide and water, was first proposed by Hans Krebs in 1937. (He was awarded the 1953 Nobel Prize in Physiology or Medicine.) Acetyl-CoA's entrance into the citric acid cycle is the beginning of stage III of catabolism. The citric acid cycle produces adenosine triphosphate (ATP), reduced nicotinamide adenine dinucleotide (NADH), reduced flavin adenine dinucleotide ($FADH_2$), and metabolic intermediates for the synthesis of needed compounds.

Steps of the Citric Acid Cycle

At first glance, the citric acid cycle appears rather complex (Figure 19.5.1). All the reactions, however, are familiar types in organic chemistry: hydration, oxidation, decarboxylation, and hydrolysis. Each reaction of the citric acid cycle is numbered, and in Figure 19.5.1, the two acetyl carbon atoms are highlighted in red. Each intermediate in the cycle is a carboxylic acid, existing as an anion at physiological pH. All the reactions occur within the mitochondria, which are small organelles within the cells of plants and animals.





Figure 19.5.1: Reactions of the Citric Acid Cycle

1. In the first step, acetyl-CoA enters the citric acid cycle, and the acetyl group is transferred onto oxaloacetate, yielding citrate. Note that this step releases coenzyme A. The reaction is catalyzed by *citrate synthase*.

2. In the next step, *aconitase* catalyzes the isomerization of citrate to isocitrate. In this reaction, a tertiary alcohol, which cannot be oxidized, is converted to a secondary alcohol, which can be oxidized in the next step.
3. Isocitrate then undergoes a reaction known as oxidative decarboxylation because the alcohol is oxidized and the molecule is shortened by one carbon atom with the release of carbon dioxide (decarboxylation). The reaction is catalyzed by *isocitrate dehydrogenase*, and the product of the reaction is α-ketoglutarate. An important reaction linked to this is the reduction of the coenzyme nicotinamide adenine dinucleotide (NAD^+) to NADH. The NADH is ultimately reoxidized, and the energy released is used in the synthesis of ATP, as we shall see.
4. The fourth step is another oxidative decarboxylation. This time α-ketoglutarate is converted to succinyl-CoA, and another molecule of NAD^+ is reduced to NADH. The *α-ketoglutarate dehydrogenase complex* catalyzes this reaction. This is the only irreversible reaction in the citric acid cycle. As such, it prevents the cycle from operating in the reverse direction, in which acetyl-CoA would be synthesized from carbon dioxide.

> So far, in the first four steps, two carbon atoms have entered the cycle as an acetyl group, and two carbon atoms have been released as molecules of carbon dioxide. The remaining reactions of the citric acid cycle use the four carbon atoms of the succinyl group to resynthesize a molecule of oxaloacetate, which is the compound needed to combine with an incoming acetyl group and begin another round of the cycle.

In the fifth reaction, the energy released by the hydrolysis of the high-energy thioester bond of succinyl-CoA is used to form guanosine triphosphate (GTP) from guanosine diphosphate (GDP) and inorganic phosphate in a reaction catalyzed by *succinyl-CoA synthetase*. This step is the only reaction in the citric acid cycle that directly forms a high-energy phosphate compound. GTP can readily transfer its terminal phosphate group to adenosine diphosphate (ADP) to generate ATP in the presence of *nucleoside diphosphokinase*.



Succinate dehydrogenase then catalyzes the removal of two hydrogen atoms from succinate, forming fumarate. This oxidation-reduction reaction uses flavin adenine dinucleotide (FAD), rather than NAD^+, as the oxidizing agent. Succinate dehydrogenase is the only enzyme of the citric acid cycle located within the inner mitochondrial membrane. We will see soon the importance of this.

In the following step, a molecule of water is added to the double bond of fumarate to form L-malate in a reaction catalyzed by *fumarase*.

One revolution of the cycle is completed with the oxidation of L-malate to oxaloacetate, brought about by *malate dehydrogenase*. This is the third oxidation-reduction reaction that uses NAD^+ as the oxidizing agent. Oxaloacetate can accept an acetyl group from acetyl-CoA, allowing the cycle to begin again.



Video: "The Citric Acid Cycle: An Overview". In the matrix of the mitochondrion, the Citric Acid Cycle uses Acetyl CoA molecules to produce energy through eight chemical reactions. This animation provides an overview of the pathway and its products. NDSU VCell Production's animation; for more information please see Vcell, NDSU, Animations(opens in new window) [vcell.ndsu.edu].

Cellular Respiration

Respiration can be defined as the process by which cells oxidize organic molecules in the presence of gaseous oxygen to produce carbon dioxide, water, and energy in the form of ATP. We have seen that two carbon atoms enter the citric acid cycle from acetyl-CoA (step 1), and two different carbon atoms exit the cycle as carbon dioxide (steps 3 and 4). Yet nowhere in our discussion of the citric acid cycle have we indicated how oxygen is used. Recall, however, that in the four oxidation-reduction steps occurring in the citric acid cycle, the coenzyme NAD^+ or FAD is reduced to NADH or $FADH_2$, respectively. *Oxygen is needed to reoxidize these coenzymes*. Recall, too, that very little ATP is obtained directly from the citric acid cycle. Instead, oxygen participation and significant ATP production occur subsequent to the citric acid cycle, in two pathways that are closely linked: electron transport and oxidative phosphorylation.

All the enzymes and coenzymes for the citric acid cycle, the reoxidation of NADH and $FADH_2$, and the production of ATP are located in the mitochondria, which are small, oval organelles with double membranes, often referred to as the “power plants” of the cell (Figure **19.5.2**). A cell may contain 100–5,000 mitochondria, depending on its function, and the mitochondria can reproduce themselves if the energy requirements of the cell increase.




Figure **19.5.2**: Respiration

Cellular respiration occurs in the mitochondria

Figure **19.5.2** shows the mitochondrion’s two membranes: *outer* and *inner*. The inner membrane is extensively folded into a series of internal ridges called *cristae*. Thus there are two compartments in mitochondria: the *intermembrane space*, which lies between the membranes, and the *matrix*, which lies inside the inner membrane. The outer membrane is permeable, whereas the inner membrane is impermeable to most molecules and ions, although water, oxygen, and carbon dioxide can freely penetrate both membranes. The matrix contains all the enzymes of the citric acid cycle with the exception of succinate dehydrogenase, which is

embedded in the inner membrane. The enzymes that are needed for the reoxidation of NADH and $FADH_2$ and ATP production are also located in the inner membrane. They are arranged in specific positions so that they function in a manner analogous to a bucket brigade. This highly organized sequence of oxidation-reduction enzymes is known as the electron transport chain (or respiratory chain).

Electron Transport

Figure **19.5.3** illustrates the organization of the electron transport chain. The components of the chain are organized into four complexes designated I, II, III, and IV. Each complex contains several enzymes, other proteins, and metal ions. The metal ions can be reduced and then oxidized repeatedly as electrons are passed from one component to the next. Recall that a compound is reduced when it gains electrons or hydrogen atoms and is oxidized when it loses electrons or hydrogen atoms.






Figure **19.5.3**: The Mitochondrial Electron Transport Chain and ATP Synthase. The red line shows the path of electrons.

Electrons can enter the electron transport chain through either complex I or II. We will look first at electrons entering at complex I. These electrons come from NADH, which is formed in three reactions of the citric acid cycle. Let's use step 8 as an example, the reaction in which L-malate is oxidized to oxaloacetate and NAD^+ is reduced to NADH. This reaction can be divided into two half reactions:

- *Oxidation half-reaction*:



- *Reduction half-reaction*:




In the oxidation half-reaction, two hydrogen (H^+) ions and two electrons are removed from the substrate. In the reduction half-reaction, the NAD^+ molecule accepts both of those electrons and one of the H^+ ions. The other H^+ ion is transported from the matrix, across the inner mitochondrial membrane, and into the intermembrane space. The NADH diffuses through the matrix and is bound by complex I of the electron transport chain. In the complex, the coenzyme flavin mononucleotide (FMN) accepts both electrons from NADH. By passing the electrons along, NADH is oxidized back to NAD^+ and FMN is reduced to $FMNH_2$ (reduced form of flavin mononucleotide). Again, the reaction can be illustrated by dividing it into its respective half-reactions.

- *Oxidation half-reaction*:

$$NADH + H^+ \longrightarrow NAD^+ + 2H^+ + 2e^-$$

- *Reduction half-reaction*:



Complex I contains several proteins that have iron-sulfur (Fe·S) centers. The electrons that reduced FMN to $FMNH_2$ are now transferred to these proteins. The iron ions in the Fe·S centers are in the Fe(III) form at first, but by accepting an electron, each ion is reduced to the Fe(II) form. Because each Fe·S center can transfer only one electron, two centers are needed to accept the two electrons that will regenerate FMN.

- *Oxidation half-reaction*:

$$\mathbf{FMNH_2 \longrightarrow FMN + 2H^+ + 2e^-}$$

- *Reduction half-reaction*:

$$\mathbf{2\,Fe(III)\cdot S + 2e^- \longrightarrow 2\,Fe(II)\cdot S}$$

Electrons from $FADH_2$, formed in step 6 of the citric acid cycle, enter the electron transport chain through complex II. Succinate dehydrogenase, the enzyme in the citric acid cycle that catalyzes the formation of $FADH_2$ from FAD is part of complex II. The electrons from $FADH_2$ are then transferred to an Fe·S protein.

- *Oxidation half-reaction*:

$$\mathbf{FADH_2 \longrightarrow FAD + 2H^+ + 2e^-}$$

- *Reduction half-reaction*:

$$\mathbf{2\,Fe(III)\cdot S + 2e^- \longrightarrow 2\,Fe(II)\cdot S}$$


Electrons from complexes I and II are then transferred from the $\mathbf{Fe\cdot S}$ protein to coenzyme Q (CoQ), a mobile electron carrier that acts as the electron shuttle between complexes I or II and complex III.

> *Coenzyme Q is also called ubiquinone because it is ubiquitous in living systems.*

- *Oxidation half-reaction*:

$$\mathbf{2\,Fe(II)\cdot S \longrightarrow 2\,Fe(III)\cdot S + 2e^-}$$

- *Reduction half-reaction*:



Complexes III and IV include several iron-containing proteins known as cytochromes. The iron in these enzymes is located in substructures known as iron porphyrins (Figure **19.5.4**). Like the Fe·S centers, the characteristic feature of the cytochromes is the ability of their iron atoms to exist as either Fe(II) or Fe(III). Thus, each cytochrome in its oxidized form—Fe(III)—can accept one electron and be reduced to the Fe(II) form. This change in oxidation state is reversible, so the reduced form can donate its electron to the next cytochrome, and so on. Complex III contains cytochromes b and c, as well as Fe·S proteins, with cytochrome c acting as the electron shuttle between complex III and IV. Complex IV contains cytochromes a and a_3 in an enzyme known as *cytochrome oxidase*. This enzyme has the ability to transfer electrons to molecular oxygen, the last electron acceptor in the chain of electron transport reactions. In this final step, water (H_2O) is formed.

- *Oxidation half-reaction*:

$$4\,\text{Cyt}\ \ a_3\text{-Fe(II)} \longrightarrow 4\,\text{Cyt}\ \ a_3\text{-Fe(III)} + 4\,e^-$$

- *Reduction half-reaction*:

$$O_2 + 4\,H^+ + 4\,e^- \longrightarrow 2\,H_2O$$




Figure **19.5.4**: An Iron Porphyrin. Iron porphyrins are present in cytochromes as well as in myoglobin and hemoglobin.



Video: Cellular Respiration (Electron Transport Chain). Cellular respiration occurs in the mitochondria and provides both animals and plants with the energy needed to power other cellular processes. This section covers the electron transport chain.NDSU Virtual Cell Animations Project animation; ror more information please see Vcell, NDSU, Animations(opens in new window) [vcell.ndsu.edu]

Oxidative Phosphorylation

Each intermediate compound in the electron transport chain is reduced by the addition of one or two electrons in one reaction and then subsequently restored to its original form by delivering the electron(s) to the next compound along the chain. The successive electron transfers result in energy production. But how is this energy used for the synthesis of ATP? The process that links ATP synthesis to the operation of the electron transport chain is referred to as oxidative phosphorylation.

Electron transport is tightly coupled to oxidative phosphorylation. The coenzymes NADH and $FADH_2$ are oxidized by the respiratory chain *only* if ADP is simultaneously phosphorylated to ATP. The currently accepted model explaining how these two processes are linked is known as the *chemiosmotic hypothesis*, which was proposed by Peter Mitchell, resulting in Mitchell being awarded the 1978 Nobel Prize in Chemistry.

Looking again at Figure **19.5.3**, we see that as electrons are being transferred through the electron transport chain, hydrogen (H^+) ions are being transported across the inner mitochondrial membrane from the matrix to the intermembrane space. The concentration of H^+ is already higher in the intermembrane space than in the matrix, so energy is required to transport the additional H^+ there. This energy comes from the electron transfer reactions in the electron transport chain. But how does the extreme difference in H^+ concentration then lead to ATP synthesis? The buildup of H^+ ions in the intermembrane space results in an H^+ ion gradient that is a large energy source, like water behind a dam (because, given the opportunity, the protons will flow out of the intermembrane space and into the less concentrated matrix). Current research indicates that the flow of H^+ down this concentration gradient through a

fifth enzyme complex, known as ATP synthase, leads to a change in the structure of the synthase, causing the synthesis and release of ATP.

In cells that are using energy, the turnover of ATP is very high, so these cells contain high levels of ADP. They must therefore consume large quantities of oxygen continuously, so as to have the energy necessary to phosphorylate ADP to form ATP. Consider, for example, that resting skeletal muscles use about 30% of a resting adult's oxygen consumption, but when the same muscles are working strenuously, they account for almost 90% of the total oxygen consumption of the organism.

Experiment has shown that 2.5–3 ATP molecules are formed for every molecule of NADH oxidized in the electron transport chain, and 1.5–2 ATP molecules are formed for every molecule of $FADH_2$ oxidized. Table **19.5.1** summarizes the theoretical maximum yield of ATP produced by the complete oxidation of 1 mol of acetyl-CoA through the sequential action of the citric acid cycle, the electron transport chain, and oxidative phosphorylation.

Table **19.5.1**: Maximum Yield of ATP from the Complete Oxidation of 1 Mol of Acetyl-CoA

Reaction	Comments	Yield of ATP (moles)
Isocitrate → α-ketoglutarate + CO_2	produces 1 mol NADH	
α-ketoglutarate → succinyl-CoA + CO_2	produces 1 mol NADH	
Succinyl-CoA → succinate	produces 1 mol GTP	+1
Succinate → fumarate	produces 1 mol $FADH_2$	
Malate → oxaloacetate	produces 1 mol NADH	
1 $FADH_2$ from the citric acid cycle	yields 2 mol ATP	+2
3 NADH from the citric acid cycle	yields 3 mol ATP/NADH	+9
Net yield of ATP:		+12

Key Takeaways

- The acetyl group of acetyl-CoA enters the citric acid cycle. For each acetyl-CoA that enters the citric acid cycle, 2 molecules of carbon dioxide, 3 molecules of NADH, 1 molecule of ATP, and 1 molecule of $FADH_2$ are produced.
- The reduced coenzymes (NADH and $FADH_2$) produced by the citric acid cycle are reoxidized by the reactions of the electron transport chain. This series of reactions also produces a pH gradient across the inner mitochondrial membrane.
- The pH gradient produced by the electron transport chain drives the synthesis of ATP from ADP. For each NADH reoxidized, 2.5–3 molecules of ATP are produced; for each $FADH_2$ reoxidized, 1.5–2 molecules of ATP are produced.

19.5: Stage III of Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.4: Stage III of Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.6: Stage II of Carbohydrate Catabolism

Learning Objectives

- Describe the function of glycolysis and identify its major products.
- Describe how the presence or absence of oxygen determines what happens to the pyruvate and the NADH that are produced in glycolysis.
- Determine the amount of ATP produced by the oxidation of glucose in the presence and absence of oxygen.

In stage II of catabolism, the metabolic pathway known as glycolysis converts glucose into two molecules of pyruvate (a three-carbon compound with three carbon atoms) with the corresponding production of adenosine triphosphate (ATP). The individual reactions in glycolysis were determined during the first part of the 20th century. It was the first metabolic pathway to be elucidated, in part because the participating enzymes are found in soluble form in the cell and are readily isolated and purified. The pathway is structured so that the product of one enzyme-catalyzed reaction becomes the substrate of the next. The transfer of intermediates from one enzyme to the next occurs by diffusion.

Steps in Glycolysis

The 10 reactions of glycolysis, summarized in Figures **19.6.1** and **19.6.2**, can be divided into two phases. In the first 5 reactions—phase I—glucose is broken down into two molecules of glyceraldehyde 3-phosphate. In the last five reactions—phase II—each glyceraldehyde 3-phosphate is converted into pyruvate, and ATP is generated. Notice that all the intermediates in glycolysis are phosphorylated and contain either six or three carbon atoms.




Figure **19.6.1**: Phase 1 (preparatory phase) of Glycolysis

- When glucose enters a cell, it is immediately phosphorylated to form glucose 6-phosphate, in the first reaction of phase I. The phosphate donor in this reaction is ATP, and the enzyme—which requires magnesium ions for its activity—is *hexokinase*. In this reaction, ATP is being used rather than being synthesized. The presence of such a reaction in a catabolic pathway that is supposed to *generate* energy may surprise you. However, in addition to activating the glucose molecule, this initial reaction is essentially irreversible, an added benefit that keeps the overall process moving in the right direction. Furthermore, the addition of the negatively charged phosphate group prevents the intermediates formed in glycolysis from diffusing through the cell membrane, as neutral molecules such as glucose can do.
- In the next reaction, *phosphoglucose isomerase* catalyzes the isomerization of glucose 6-phosphate to fructose 6-phosphate. This reaction is important because it creates a primary alcohol, which can be readily phosphorylated.

- The subsequent phosphorylation of fructose 6-phosphate to form fructose 1,6-bisphosphate is catalyzed by *phosphofructokinase*, which requires magnesium ions for activity. ATP is again the phosphate donor.
- Fructose 1,6-bisphosphate is enzymatically cleaved by *aldolase* to form two triose phosphates: dihydroxyacetone phosphate and glyceraldehyde 3-phosphate.
- Isomerization of dihydroxyacetone phosphate into a second molecule of glyceraldehyde 3-phosphate is the final step in phase I. The enzyme catalyzing this reaction is *triose phosphate isomerase*.

When a molecule contains two phosphate groups on different carbon atoms, the convention is to use the prefix *bis*. When the two phosphate groups are bonded to each other on the same carbon atom (for example, adenosine diphosphate [ADP]), the prefix is *di*.

In steps 4 and 5, aldolase and triose phosphate isomerase effectively convert one molecule of fructose 1,6-bisphosphate into *two* molecules of glyceraldehyde 3-phosphate. Thus, phase I of glycolysis requires energy in the form of two molecules of ATP and releases none of the energy stored in glucose.




Figure **19.6.2**: Phase 2 (payoff phase) of Glycolysis

In the initial step of phase II (Figure **19.6.2**), glyceraldehyde 3-phosphate is both oxidized and phosphorylated in a reaction catalyzed by *glyceraldehyde-3-phosphate dehydrogenase*, an enzyme that requires nicotinamide adenine dinucleotide (NAD^+) as the oxidizing agent and inorganic phosphate as the phosphate donor. In the reaction, NAD^+ is reduced to reduced nicotinamide adenine dinucleotide (NADH), and 1,3-bisphosphoglycerate (BPG) is formed.

- BPG has a high-energy phosphate bond (Table **19.6.1**) joining a phosphate group to C1. This phosphate group is now transferred directly to a molecule of ADP, thus forming ATP and 3-phosphoglycerate. The enzyme that catalyzes the reaction is **phosphoglycerate kinase**, which, like all other kinases, requires magnesium ions to function. This is the first reaction to produce ATP in the pathway. Because the ATP is formed by a direct transfer of a phosphate group from a metabolite to ADP—that is, from one substrate to another—the process is referred to as substrate-level phosphorylation, to distinguish it from the *oxidative phosphorylation* discussed in Section 20.4.
- In the next reaction, the phosphate group on 3-phosphoglycerate is transferred from the OH group of C3 to the OH group of C2, forming 2-phosphoglycerate in a reaction catalyzed by *phosphoglyceromutase*.
- A dehydration reaction, catalyzed by *enolase*, forms phosphoenolpyruvate (PEP), another compound possessing a high-energy phosphate group.

- The final step is irreversible and is the second reaction in which substrate-level phosphorylation occurs. The phosphate group of PEP is transferred to ADP, with one molecule of ATP being produced per molecule of PEP. The reaction is catalyzed by *pyruvate kinase*, which requires both magnesium and potassium ions to be active.

Table **19.6.1**: Maximum Yield of ATP from the Complete Oxidation of 1 Mol of Glucose

Reaction	Comments	Yield of ATP (moles)
glucose → glucose 6-phosphate	consumes 1 mol ATP	−1
fructose 6-phosphate → fructose 1,6-bisphosphate	consumes 1 mol ATP	−1
glyceraldehyde 3-phosphate → BPG	produces 2 mol of cytoplasmic NADH	
BPG → 3-phosphoglycerate	produces 2 mol ATP	+2
phosphoenolpyruvate → pyruvate	produces 2 mol ATP	+2
pyruvate → acetyl-CoA + CO_2	produces 2 mol NADH	
isocitrate → α-ketoglutarate + CO_2	produces 2 mol NADH	
α-ketoglutarate → succinyl-CoA + CO_2	produces 2 mol NADH	
succinyl-CoA → succinate	produces 2 mol GTP	+2
succinate → fumarate	produces 2 mol $FADH_2$	
malate → oxaloacetate	produces 2 mol NADH	
2 cytoplasmic NADH from glycolysis	yields 2–3 mol ATP per NADH (depending on tissue)	+4 to +6
2 NADH from the oxidation of pyruvate	yields 3 mol ATP per NADH	+6
2 $FADH_2$ from the citric acid cycle	yields 2 ATP per $FADH_2$	+4
3 NADH from the citric acid cycle	yields 3 ATP per NADH	+18
Net yield of ATP:		+36 to +38

In phase II, two molecules of glyceraldehyde 3-phosphate are converted to two molecules of pyruvate, along with the production of four molecules of ATP and two molecules of NADH.

To Your Health: Diabetes

Although medical science has made significant progress against diabetes , it continues to be a major health threat. Some of the serious complications of diabetes are as follows:

- It is the leading cause of lower limb amputations in the United States.
- It is the leading cause of blindness in adults over age 20.
- It is the leading cause of kidney failure.
- It increases the risk of having a heart attack or stroke by two to four times.


Because a person with diabetes is unable to use glucose properly, excessive quantities accumulate in the blood and the urine. Other characteristic symptoms are constant hunger, weight loss, extreme thirst, and frequent urination because the kidneys excrete large amounts of water in an attempt to remove excess sugar from the blood.

There are two types of diabetes. In immune-mediated diabetes, insufficient amounts of insulin are produced. This type of diabetes develops early in life and is also known as *Type 1 diabetes*, as well as insulin-dependent or juvenile-onset diabetes.

Symptoms are rapidly reversed by the administration of insulin, and Type 1 diabetics can lead active lives provided they receive insulin as needed. Because insulin is a protein that is readily digested in the small intestine, it cannot be taken orally and must be injected at least once a day.

In Type 1 diabetes, insulin-producing cells of the pancreas are destroyed by the body's immune system. Researchers are still trying to find out why. Meanwhile, they have developed a simple blood test capable of predicting who will develop Type 1 diabetes several years before the disease becomes apparent. The blood test reveals the presence of antibodies that destroy the body's insulin-producing cells.

Type 2 diabetes, also known as noninsulin-dependent or adult-onset diabetes, is by far the more common, representing about 95% of diagnosed diabetic cases. (This translates to about 16 million Americans.) Type 2 diabetics usually produce sufficient amounts of insulin, but either the insulin-producing cells in the pancreas do not release enough of it, or it is not used properly because of defective insulin receptors or a lack of insulin receptors on the target cells. In many of these people, the disease can be controlled with a combination of diet and exercise alone. For some people who are overweight, losing weight is sufficient to bring their blood sugar level into the normal range, after which medication is not required if they exercise regularly and eat wisely.



Those who require medication may use oral antidiabetic drugs that stimulate the islet cells to secrete insulin. First-generation antidiabetic drugs stimulated the release of insulin. Newer second-generation drugs, such as glyburide, do as well, but they also increase the sensitivity of cell receptors to insulin. Some individuals with Type 2 diabetes do not produce enough insulin and thus do not respond to these oral medications; they must use insulin. In both Type 1 and Type 2 diabetes, the blood sugar level must be carefully monitored and adjustments made in diet or medication to keep the level as normal as possible (70–120 mg/dL).

Metabolism of Pyruvate

The presence or absence of oxygen determines the fates of the pyruvate and the NADH produced in glycolysis. When plenty of oxygen is available, pyruvate is completely oxidized to carbon dioxide, with the release of much greater amounts of ATP through the combined actions of the citric acid cycle, the electron transport chain, and oxidative phosphorylation. However, in the absence of oxygen (that is, under anaerobic conditions), the fate of pyruvate is different in different organisms. In vertebrates, pyruvate is converted to lactate, while other organisms, such as yeast, convert pyruvate to ethanol and carbon dioxide. These possible fates of pyruvate are summarized in Figure **19.6.2**. The conversion to lactate or ethanol under anaerobic conditions allows for the reoxidation of NADH to NAD^+ in the absence of oxygen.




Figure 19.6.2: Metabolic Fates of Pyruvate

ATP Yield from Glycolysis

The net energy yield from anaerobic glucose metabolism can readily be calculated in moles of ATP. In the initial phosphorylation of glucose (step 1), 1 mol of ATP is expended, along with another in the phosphorylation of fructose 6-phosphate (step 3). In step 7, 2 mol of BPG (recall that 2 mol of 1,3-BPG are formed for each mole of glucose) are converted to 2 mol of 3-phosphoglycerate, and 2 mol of ATP are produced. In step 10, 2 mol of pyruvate and 2 mol of ATP are formed per mole of glucose.

For every mole of glucose degraded, 2 mol of ATP are initially consumed and 4 mol of ATP are ultimately produced. The net production of ATP is thus 2 mol for each mole of glucose converted to lactate or ethanol. If 7.4 kcal of energy is conserved per mole of ATP produced, and the total amount of energy that can theoretically be obtained from the complete oxidation of 1 mol of glucose is 670 kcal (as stated in the chapter introduction), the energy conserved in the anaerobic catabolism of glucose to two molecules of lactate (or ethanol) is as follows:

$$\frac{2 \times 7.4\ \text{kcal}}{670\ \text{kcal}} \times 100 = 2.2\%$$

Thus anaerobic cells extract only a very small fraction of the total energy of the glucose molecule.

Contrast this result with the amount of energy obtained when glucose is completely oxidized to carbon dioxide and water through glycolysis, the citric acid cycle, the electron transport chain, and oxidative phosphorylation as summarized in Table 19.6.1. Note the indication in the table that a variable amount of ATP is synthesized, depending on the tissue, from the NADH formed in the cytoplasm during glycolysis. This is because NADH is not transported into the inner mitochondrial membrane where the enzymes for the electron transport chain are located. Instead, brain and muscle cells use a transport mechanism that passes electrons from the cytoplasmic NADH through the membrane to flavin adenine dinucleotide (FAD) molecules inside the mitochondria, forming reduced flavin adenine dinucleotide ($FADH_2$), which then feeds the electrons into the electron transport chain. This route lowers the yield of ATP to 1.5–2 molecules of ATP, rather than the usual 2.5–3 molecules. A more efficient transport system is found in liver, heart, and kidney cells where the formation of one cytoplasmic NADH molecule results in the formation of one mitochondrial NADH molecule, which leads to the formation of 2.5–3 molecules of ATP.The total amount of energy conserved in the aerobic catabolism of glucose in the liver is as follows:

$$\frac{38 \times 7.4\ \text{kcal}}{670\ \text{kcal}} \times 100 = 42\%$$

Conservation of 42% of the total energy released compares favorably with the efficiency of any machine. In comparison, automobiles are only about 20%–25% efficient in using the energy released by the combustion of gasoline.

As indicated earlier, the 58% of released energy that is not conserved enters the surroundings (that is, the cell) as heat that helps to maintain body temperature. If we are exercising strenuously and our metabolism speeds up to provide the energy needed for muscle contraction, more heat is produced. We begin to perspire to dissipate some of that heat. As the perspiration evaporates, the excess heat is carried away from the body by the departing water vapor.

Summary

- The monosaccharide glucose is broken down through a series of enzyme-catalyzed reactions known as glycolysis.
- For each molecule of glucose that is broken down, two molecules of pyruvate, two molecules of ATP, and two molecules of NADH are produced.
- In the absence of oxygen, pyruvate is converted to lactate, and NADH is reoxidized to NAD+. In the presence of oxygen, pyruvate is converted to acetyl-CoA and then enters the citric acid cycle.
- More ATP can be formed from the breakdown of glucose when oxygen is present.

19.6: Stage II of Carbohydrate Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.


19.7: Stage II of Lipid Catabolism

Learning Objectives

- To describe the reactions needed to completely oxidize a fatty acid to carbon dioxide and water.

Like glucose, the fatty acids released in the digestion of triglycerides and other lipids are broken down in a series of sequential reactions accompanied by the gradual release of usable energy. Some of these reactions are oxidative and require nicotinamide adenine dinucleotide (NAD^+) and flavin adenine dinucleotide (FAD). The enzymes that participate in fatty acid catabolism are located in the mitochondria, along with the enzymes of the citric acid cycle, the electron transport chain, and oxidative phosphorylation. This localization of enzymes in the mitochondria is of the utmost importance because it facilitates efficient utilization of energy stored in fatty acids and other molecules.

Fatty acid oxidation is initiated on the outer mitochondrial membrane. There the fatty acids, which like carbohydrates are relatively inert, must first be activated by conversion to an energy-rich fatty acid derivative of coenzyme A called *fatty acyl-coenzyme A* (CoA). The activation is catalyzed by *acyl-CoA synthetase*. For each molecule of fatty acid activated, one molecule of coenzyme A and one molecule of adenosine triphosphate (ATP) are used, equaling a net utilization of the two high-energy bonds in one ATP molecule (which is therefore converted to adenosine monophosphate [AMP] rather than adenosine diphosphate [ADP]):



The fatty acyl-CoA diffuses to the inner mitochondrial membrane, where it combines with a carrier molecule known as carnitine in a reaction catalyzed by *carnitine acyltransferase*. The acyl-carnitine derivative is transported into the mitochondrial matrix and converted back to the fatty acyl-CoA.

Steps in the β-Oxidation of Fatty Acids

Further oxidation of the fatty acyl-CoA occurs in the mitochondrial matrix via a sequence of four reactions known collectively as β-oxidation because the β-carbon undergoes successive oxidations in the progressive removal of two carbon atoms from the carboxyl end of the fatty acyl-CoA (Figure 19.7.1).




Figure **19.7.1**: Fatty Acid Oxidation. The fatty acyl-CoA formed in the final step becomes the substrate for the first step in the next round of β-oxidation. β-oxidation continues until two acetyl-CoA molecules are produced in the final step.

The first step in the catabolism of fatty acids is the formation of an alkene in an oxidation reaction catalyzed by *acyl-CoA dehydrogenase*. In this reaction, the coenzyme FAD accepts two hydrogen atoms from the acyl-CoA, one from the α-carbon and one from the β-carbon, forming reduced flavin adenine dinucleotide ($FADH_2$).


> The $FADH_2$ is reoxidized back to FAD via the electron transport chain that supplies energy to form 1.5–2 molecules of ATP.

Next, the *trans*-alkene is hydrated to form a secondary alcohol in a reaction catalyzed by *enoyl-CoA hydratase*. The enzyme forms only the L-isomer.

The secondary alcohol is then oxidized to a ketone by *β-hydroxyacyl-CoA dehydrogenase*, with NAD^+ acting as the oxidizing agent. The reoxidation of each molecule of NADH to NAD^+ by the electron transport chain furnishes 2.5–3 molecules of ATP.

The final reaction is cleavage of the β-ketoacyl-CoA by a molecule of coenzyme A. The products are acetyl-CoA and a fatty acyl-CoA that has been shortened by two carbon atoms. The reaction is catalyzed by *thiolase*.

The shortened fatty acyl-CoA is then degraded by repetitions of these four steps, each time releasing a molecule of acetyl-CoA. The overall equation for the β-oxidation of palmitoyl-CoA (16 carbon atoms) is as follows:




> *Because each shortened fatty acyl-CoA cycles back to the beginning of the pathway, β-oxidation is sometimes referred to as the fatty acid spiral.*

The fate of the acetyl-CoA obtained from fatty acid oxidation depends on the needs of an organism. It may enter the citric acid cycle and be oxidized to produce energy, it may be used for the formation of water-soluble derivatives known as ketone bodies, or it may serve as the starting material for the synthesis of fatty acids. For more information about the citric acid cycle, see Section 20.4.

Looking Closer: Ketone Bodies

In the liver, most of the acetyl-CoA obtained from fatty acid oxidation is oxidized by the citric acid cycle. However, some of the acetyl-CoA is used to synthesize a group of compounds known as *ketone bodies*: acetoacetate, β-hydroxybutyrate, and acetone. Two acetyl-CoA molecules combine, in a reversal of the final step of β-oxidation, to produce acetoacetyl-CoA. The acetoacetyl-CoA reacts with another molecule of acetyl-CoA and water to form β-hydroxy-β-methylglutaryl-CoA, which is then cleaved to acetoacetate and acetyl-CoA. Most of the acetoacetate is reduced to β-hydroxybutyrate, while a small amount is decarboxylated to carbon dioxide and acetone.



The acetoacetate and β-hydroxybutyrate synthesized by the liver are released into the blood for use as a metabolic fuel (to be converted back to acetyl-CoA) by other tissues, particularly the kidney and the heart. Thus, during prolonged starvation, ketone bodies provide about 70% of the energy requirements of the brain. Under normal conditions, the kidneys excrete about 20 mg of ketone bodies each day, and the blood levels are maintained at about 1 mg of ketone bodies per 100 mL of blood.

In starvation, diabetes mellitus, and certain other physiological conditions in which cells do not receive sufficient amounts of carbohydrate, the rate of fatty acid oxidation increases to provide energy. This leads to an increase in the concentration of acetyl-CoA. The increased acetyl-CoA cannot be oxidized by the citric acid cycle because of a decrease in the concentration of oxaloacetate, which is diverted to glucose synthesis. In response, the rate of ketone body formation in the liver increases further, to a level much higher than can be used by other tissues. The excess ketone bodies accumulate in the blood and the urine, a condition referred to as *ketosis*. When the acetone in the blood reaches the lungs, its volatility causes it to be expelled in the breath. The sweet smell of acetone, a characteristic of ketosis, is frequently noticed on the breath of severely diabetic patients.

Because two of the three kinds of ketone bodies are weak acids, their presence in the blood in excessive amounts overwhelms the blood buffers and causes a marked decrease in blood pH (to 6.9 from a normal value of 7.4). This decrease in pH leads to a serious condition known as *acidosis*. One of the effects of acidosis is a decrease in the ability of hemoglobin to transport oxygen in the blood. In moderate to severe acidosis, breathing becomes labored and very painful. The body also loses fluids and becomes dehydrated as the kidneys attempt to get rid of the acids by eliminating large quantities of water. The lowered oxygen supply and dehydration lead to depression; even mild acidosis leads to lethargy, loss of appetite, and a generally run-down feeling. Untreated patients may go into a coma. At that point, prompt treatment is necessary if the person's life is to be saved.

ATP Yield from Fatty Acid Oxidation

The amount of ATP obtained from fatty acid oxidation depends on the size of the fatty acid being oxidized. For our purposes here. we'll study palmitic acid, a saturated fatty acid with 16 carbon atoms, as a typical fatty acid in the human diet. Calculating its energy yield provides a model for determining the ATP yield of all other fatty acids.

The breakdown by an organism of 1 mol of palmitic acid requires 1 mol of ATP (for activation) and forms 8 mol of acetyl-CoA. Recall from Table 20.4.1 that each mole of acetyl-CoA metabolized by the citric acid cycle yields 10 mol of ATP. The complete degradation of 1 mol of palmitic acid requires the β-oxidation reactions to be repeated seven times. Thus, 7 mol of NADH and 7 mol of $FADH_2$ are produced. Reoxidation of these compounds through respiration yields 2.5–3 and 1.5–2 mol of ATP, respectively. The energy calculations can be summarized as follows:

ATP Yield from Fatty Acid Oxidation

1 mol of ATP is split to AMP and $2P_i$	−2 ATP
8 mol of acetyl-CoA formed (8 × 12)	96 ATP
7 mol of $FADH_2$ formed (7 × 2)	14 ATP
7 mol of NADH formed (7 × 3)	21 ATP
Total	129 ATP

> *The number of times β-oxidation is repeated for a fatty acid containing n carbon atoms is n/2 – 1 because the final turn yields two acetyl-CoA molecules.*

The combustion of 1 mol of palmitic acid releases a considerable amount of energy:

$$C_{16}H_{32}O_2 + 23O_2 \rightarrow 16CO_2 + 16H_2O + 2,340\ kcal$$

The percentage of this energy that is conserved by the cell in the form of ATP is as follows:

$$\frac{\text{energy conserved}}{\text{total energy available}} \times 100 = \frac{(129\ \text{ATP})(7.4\ \text{kcal/ATP})}{2,340\ \text{kcal}} \times 100 = 41\%$$

The efficiency of fatty acid metabolism is comparable to that of carbohydrate metabolism, which we calculated to be 42%. For more information about the efficiency of fatty acid metabolism, see II of Carbohydrate Catabolism" data-cke-saved-href="/Bookshelves/Introductory_Chemistry/Basics_of_General_Organic_and_Biological_Chemistry_(Ball_et_al.)/20:_Energy_Metabolism/20.05:_Stage_II_of_Carbohydrate_Catabolism" href="/Bookshelves/Introductory_Chemistry/Basics_of_General_Organic_and_Biological_Chemistry_(Ball_et_al.)/20:_Energy_Metabolism/20.05:_Stage_II_of_Carbohydrate_Catabolism" data-quail-id="91">Section 20.5.

> *The oxidation of fatty acids produces large quantities of water. This water, which sustains migratory birds and animals (such as the camel) for long periods of time.*

Summary

- Fatty acids, obtained from the breakdown of triglycerides and other lipids, are oxidized through a series of reactions known as β-oxidation.
- In each round of β-oxidation, 1 molecule of acetyl-CoA, 1 molecule of NADH, and 1 molecule of $FADH_2$ are produced.
- The acetyl-CoA, NADH, and $FADH_2$ are used in the citric acid cycle, the electron transport chain, and oxidative phosphorylation to produce ATP.

19.7: Stage II of Lipid Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.6: Stage II of Lipid Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.8: Stage II of Protein Catabolism

Learning Objectives

- To describe how excess amino acids are degraded.

The liver is the principal site of amino acid metabolism, but other tissues, such as the kidney, the small intestine, muscles, and adipose tissue, take part. Generally, the first step in the breakdown of amino acids is the separation of the amino group from the carbon skeleton, usually by a transamination reaction. The carbon skeletons resulting from the deaminated amino acids are used to form either glucose or fats, or they are converted to a metabolic intermediate that can be oxidized by the citric acid cycle. The latter alternative, amino acid catabolism, is more likely to occur when glucose levels are low—for example, when a person is fasting or starving.

Transamination

Transamination is an exchange of functional groups between any amino acid (except lysine, proline, and threonine) and an α-keto acid. The amino group is usually transferred to the keto carbon atom of pyruvate, oxaloacetate, or α-ketoglutarate, converting the α-keto acid to alanine, aspartate, or glutamate, respectively. Transamination reactions are catalyzed by specific transaminases (also called aminotransferases), which require pyridoxal phosphate as a coenzyme.




Figure 19.8.1).

> *In an α-keto acid, the carbonyl or keto group is located on the carbon atom adjacent to the carboxyl group of the acid.*






Figure 19.8.1: Two Transamination Reactions. In both reactions, the final acceptor of the amino group is α-ketoglutarate, and the final product is glutamate.

Oxidative Deamination

In the breakdown of amino acids for energy, the final acceptor of the α-amino group is α-ketoglutarate, forming glutamate. Glutamate can then undergooxidative deamination, in which it loses its amino group as an ammonium (NH_4^+) ion and is oxidized back to α-ketoglutarate (ready to accept another amino group):



This reaction occurs primarily in liver mitochondria. Most of the NH_4^+ ion formed by oxidative deamination of glutamate is converted to urea and excreted in the urine in a series of reactions known as the **urea cycle**.



The synthesis of glutamate occurs in animal cells by reversing the reaction catalyzed by glutamate dehydrogenase. For this reaction nicotinamide adenine dinucleotide phosphate (NADPH) acts as the reducing agent. The synthesis of glutamate is significant because it is one of the few reactions in animals that can incorporate inorganic nitrogen (NH_4^+) into an α-keto acid to form an amino acid. The amino group can then be passed on through transamination reactions, to produce other amino acids from the appropriate α-keto acids.

The Fate of the Carbon Skeleton

Any amino acid can be converted into an intermediate of the citric acid cycle. Once the amino group is removed, usually by transamination, the α-keto acid that remains is catabolized by a pathway unique to that acid and consisting of one or more reactions. For example, phenylalanine undergoes a series of six reactions before it splits into fumarate and acetoacetate. Fumarate is an intermediate in the citric acid cycle, while acetoacetate must be converted to acetoacetyl-coenzyme A (CoA) and then to acetyl-CoA before it enters the citric acid cycle.




Figure **19.8.2**: Fates of the Carbon Skeletons of Amino Acids

Those amino acids that can form any of the intermediates of carbohydrate metabolism can subsequently be converted to glucose via a metabolic pathway known as gluconeogenesis. These amino acids are called glucogenic amino acids. Amino acids that are converted to acetoacetyl-CoA or acetyl-CoA, which can be used for the synthesis of ketone bodies but not glucose, are called ketogenic amino acids. Some amino acids fall into both categories. Leucine and lysine are the only amino acids that are exclusively ketogenic. Figure **19.8.2** summarizes the ultimate fates of the carbon skeletons of the 20 amino acids.

Career Focus: Exercise Physiologist

An exercise physiologist works with individuals who have or wish to prevent developing a wide variety of chronic diseases, such as diabetes, in which exercise has been shown to be beneficial. Each individual must be referred by a licensed physician. An exercise physiologist works in a variety of settings, such as a hospital or in a wellness program at a commercial business, to design and monitor individual exercise plans. A registered clinical exercise physiologist must have an undergraduate degree in exercise physiology or a related degree. Some job opportunities require a master's degree in exercise physiology or a related degree.



Ergospirometry laboratory for the measurement of metabolic changes during a graded exercise test on a treadmill. from Wikipedia.

Summary

Generally the first step in the breakdown of amino acids is the removal of the amino group, usually through a reaction known as transamination. The carbon skeletons of the amino acids undergo further reactions to form compounds that can either be used for the synthesis of glucose or the synthesis of ketone bodies.

19.8: Stage II of Protein Catabolism is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.7: Stage II of Protein Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.9: Oxidative Phosphorylation

Skills to Develop

- Describe how electrons move through the electron transport chain and what happens to their energy levels
- Explain how a proton (H^+) gradient is established and maintained by the electron transport chain

You have just read about two pathways in glucose catabolism—glycolysis and the citric acid cycle—that generate ATP and coenzymes NADH and FADH2. Most of the ATP generated during the aerobic catabolism of glucose, however, is not generated directly from these pathways. Rather, it is derived from a process that begins with moving electrons (e-) from NADH and FADH2 through a series of electron transporters (Electron Trasport Chain, located in the inner mitochondrial membrane) that undergo redox reactions. The Electron Transport Chain catalyzes the reduction of O_2 into H_2O:

$4H^+ + 4e- + O_2$ --> $2H_2O$

or, equivalently,

$2H^+ + 2e- + 1/2\ O_2$ --> H_2O

Simultaneously with transporting electrons and reducing oxygen, some of the electrons transporters (complexes I, III, and IV, see figure Figure **19.9.1**) are able to pump protons (hydrogen ions, H+) from the matrix into the intermembrane space, and it is in this way that the hydrogen ion gradient is established and maintained between the two compartments separated by the inner mitochondrial membrane. This causes hydrogen ions to accumulate in the intermembrane space. Therefore, a concentration gradient forms. The energy accumulated by this gradient is used by the enzyme ATP synthase to synthesize ATP . While hydrogen ions diffuse back into the matrix by passing through ATP synthase, the flux of hydrogen ions powers the catalytic action of ATP synthase, which phosphorylates ADP, producing ATP. This mechanism is called **chemiosmosis**.

Electron Transport Chain

The electron transport chain (Figure **19.9.1**) is the last component of aerobic respiration and is the only part of glucose metabolism that uses atmospheric oxygen. Oxygen continuously diffuses into plants; in animals, it enters the body through the respiratory system. Electron transport is a series of redox reactions that resemble a relay race or bucket brigade in that electrons are passed rapidly from one component to the next, to the endpoint of the chain where the electrons reduce molecular oxygen, producing water. There are four complexes composed of proteins, labeled I through IV in Figure **19.9.1**, and the aggregation of these four complexes, together with associated mobile, accessory electron carriers, is called the **electron transport chain**. The electron transport chain is present in multiple copies in the inner mitochondrial membrane of eukaryotes and the plasma membrane of prokaryotes.




Figure **19.9.1**: The electron transport chain is a series of electron transporters embedded in the inner mitochondrial membrane that shuttles electrons from NADH and $FADH_2$ to molecular oxygen. In the process, protons are pumped from the mitochondrial matrix to the intermembrane space, and oxygen is reduced to form water.

Complex I

To start, two electrons are carried to the first complex aboard NADH. This complex, labeled I, is composed of flavin mononucleotide (FMN) and an iron-sulfur (Fe-S)-containing protein. FMN, which is derived from vitamin B_2, also called riboflavin, is one of several prosthetic groups or co-factors in the electron transport chain. A prosthetic group is a non-protein molecule required for the activity of a protein. Prosthetic groups are organic or inorganic, non-peptide molecules bound to a protein that facilitate its function; prosthetic groups include co-enzymes, which are the prosthetic groups of enzymes. The enzyme in complex I is NADH dehydrogenase and is a very large protein, containing 45 amino acid chains. Complex I can pump four hydrogen ions across the membrane from the matrix into the intermembrane space, and it is in this way that the hydrogen ion gradient is established and maintained between the two compartments separated by the inner mitochondrial membrane.

Q and Complex II

Complex II directly receives $FADH_2$, which does not pass through complex I. The compound connecting the first and second complexes to the third is ubiquinone (Q). The Q molecule is lipid soluble and freely moves through the hydrophobic core of the membrane. Once it is reduced, (QH_2), ubiquinone delivers its electrons to the next complex in the electron transport chain. Q receives the electrons derived from NADH from complex I and the electrons derived from $FADH_2$ from complex II, including succinate dehydrogenase. This enzyme and $FADH_2$ form a small complex that delivers electrons directly to the electron transport chain, bypassing the first complex. Since these electrons bypass and thus do not energize the proton pump in the first complex, fewer ATP molecules are made from the $FADH_2$ electrons. The number of ATP molecules ultimately obtained is directly proportional to the number of protons pumped across the inner mitochondrial membrane.

Complex III

The third complex is composed of cytochrome b, another Fe-S protein, Rieske center (2Fe-2S center), and cytochrome c proteins; this complex is also called cytochrome oxidoreductase. Cytochrome proteins have a prosthetic group of heme. The heme molecule is similar to the heme in hemoglobin, but it carries electrons, not oxygen. As a result, the iron ion at its core is reduced and oxidized as it passes the electrons, fluctuating between different oxidation states: Fe^{++} (reduced) and Fe^{+++} (oxidized). The heme molecules in the cytochromes have slightly different characteristics due to the effects of the different proteins binding them, giving slightly different characteristics to each complex. Complex III pumps protons through the membrane and passes its electrons to cytochrome c for transport to the fourth complex of proteins and enzymes (cytochrome c is the acceptor of electrons from Q; however, whereas Q carries pairs of electrons, cytochrome c can accept only one at a time).

Complex IV

The fourth complex is composed of cytochrome proteins c, a, and a_3. This complex contains two heme groups (one in each of the two cytochromes, a, and a_3) and three copper ions (a pair of Cu_A and one Cu_B in cytochrome a_3). The cytochromes hold an oxygen molecule very tightly between the iron and copper ions until the oxygen is completely reduced. The reduced oxygen then picks up two hydrogen ions from the surrounding medium to make water (H_2O). The removal of the hydrogen ions from the system contributes to the ion gradient used in the process of chemiosmosis.

Chemiosmosis

In chemiosmosis, the free energy from the series of redox reactions just described is used to pump hydrogen ions (protons) across the membrane. The uneven distribution of H^+ ions across the membrane establishes both concentration and electrical gradients (thus, an electrochemical gradient), owing to the hydrogen ions' positive charge and their aggregation on one side of the membrane.

If the membrane were open to diffusion by the hydrogen ions, the ions would tend to diffuse back across into the matrix, driven by their electrochemical gradient. Recall that many ions cannot diffuse through the nonpolar regions of phospholipid membranes without the aid of ion channels. Similarly, hydrogen ions in the matrix space can only pass through the inner mitochondrial membrane through an integral membrane protein called ATP synthase (Figure **19.9.2**). This complex protein acts as a tiny generator, turned by the force of the hydrogen ions diffusing through it, down their electrochemical gradient. The turning of parts of this molecular machine facilitates the addition of a phosphate to ADP, forming ATP, using the potential energy of the hydrogen ion gradient.




Figure **19.9.2**: ATP synthase is a complex, molecular machine that uses a proton (H+) gradient to form ATP from ADP and inorganic phosphate (Pi). (Credit: modification of work by Klaus Hoffmeier)

Chemiosmosis (Figure **19.9.3**) is used to generate 90 percent of the ATP made during aerobic glucose catabolism; it is also the method used in the light reactions of photosynthesis to harness the energy of sunlight in the process of photophosphorylation. Recall that the production of ATP using the process of chemiosmosis in mitochondria is called oxidative phosphorylation. The overall result of these reactions is the production of ATP from the energy of the electrons removed from hydrogen atoms. These

atoms were originally part of a glucose molecule. At the end of the pathway, the electrons are used to reduce an oxygen molecule to oxygen ions. The extra electrons on the oxygen attract hydrogen ions (protons) from the surrounding medium, and water is formed.




Figure 19.9.3: In oxidative phosphorylation, the pH gradient formed by the electron transport chain is used by ATP synthase to form ATP.

Summary

The electron transport chain is the portion of aerobic respiration that uses free oxygen as the final electron acceptor of the electrons removed from the intermediate compounds in glucose catabolism. The electron transport chain is composed of four large, multiprotein complexes embedded in the inner mitochondrial membrane and two small diffusible electron carriers shuttling electrons between them. The electrons are passed through a series of redox reactions, with a small amount of free energy used at three points to transport hydrogen ions across a membrane. This process contributes to the gradient used in chemiosmosis. The electrons passing through the electron transport chain gradually lose energy, High-energy electrons donated to the chain by either NADH or $FADH_2$ complete the chain, as low-energy electrons reduce oxygen molecules and form water. The level of free energy of the electrons drops from about 60 kcal/mol in NADH or 45 kcal/mol in $FADH_2$ to about 0 kcal/mol in water. The end products of the electron transport chain are water and ATP. A number of intermediate compounds of the citric acid cycle can be diverted into the anabolism of other biochemical molecules, such as nonessential amino acids, sugars, and lipids. These same molecules can serve as energy sources for the glucose pathways.

Contributors and Attributions

- Template:ContribOpenSTAXBio

This page titled 19.9: Oxidative Phosphorylation is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **7.4: Oxidative Phosphorylation** by OpenStax is licensed CC BY 4.0.

19.10: Energy Metabolism (Exercises)

20.1: ATP- the Universal Energy Currency

Concept Review Exercise

1. Why is ATP referred to as the energy currency of the cell?

Answer

1. ATP is the principal molecule involved in energy exchange reactions in biological systems.

Exercises

1. How do ATP and ADP differ in structure?
2. Why does the hydrolysis of ATP to ADP involve the release of energy?
3. Identify whether each compound would be classified as a high-energy phosphate compound.
   a. ATP
   b. glucose 6-phosphate
   c. creatine phosphate
4. Identify whether each compound would be classified as a high-energy phosphate compound.
   a. ADP
   b. AMP
   c. glucose 1-phosphate

Answers

1. ATP has a triphosphate group attached, while ADP has only a diphosphate group attached.

3. a. yes
   b. no
   c. yes

20.2: Stage I of Catabolism

Concept Review Exercises

1. Distinguish between each pair of compounds.
   a. pepsin and pepsinogen
   b. chymotrypsin and trypsin
   c. aminopeptidase and carboxypeptidase
2. What are the primary end products of each form of digestion?
   a. carbohydrate digestion
   b. lipid digestion
   c. protein digestion
3. In what section of the digestive tract does most of the carbohydrate, lipid, and protein digestion take place?

Answers

1. a. Pepsinogen is an inactive form of pepsin; pepsin is the active form of the enzyme.
   b. Both enzymes catalyze the hydrolysis of peptide bonds. Chymotrypsin catalyzes the hydrolysis of peptide bonds following aromatic amino acids, while trypsin catalyzes the hydrolysis of peptide bonds following lysine and arginine.
   c. Aminopeptidase catalyzes the hydrolysis of amino acids from the N-terminal end of a protein, while carboxypeptidase catalyzes the hydrolysis of amino acids from the C-terminal end of a protein.
2. a. glucose, fructose, and galactose

b. monoglycerides and fatty acids
c. amino acids

3. the small intestine

Exercises

1. What are the products of digestion (or stage I of catabolism)?
2. What is the general type of reaction used in digestion?
3. Give the site of action and the function of each enzyme.
   a. chymotrypsin
   b. lactase
   c. pepsin
   d. maltase
4. Give the site of action and the function of each enzyme.
   a. α-amylase
   b. trypsin
   c. sucrase
   d. aminopeptidase
5. a. What is the meaning of the following statement? “Bile salts act to emulsify lipids in the small intestine.”
   b. Why is emulsification important?
6. Using chemical equations, describe the chemical changes that triglycerides undergo during digestion.
7. What are the expected products from the enzymatic action of chymotrypsin on each amino acid segment?
   a. gly-ala-phe-thr-leu
   b. ala-ile-tyr-ser-arg
   c. val-trp-arg-leu-cys
8. What are the expected products from the enzymatic action of trypsin on each amino acid segment?
   a. leu-thr-glu-lys-ala
   b. phe-arg-ala-leu-val
   c. ala-arg-glu-trp-lys

Answers

1. proteins: amino acids; carbohydrates: monosaccharides; fats: fatty acids and glycerol
3. a. Chymotrypsin is found in the small intestine and catalyzes the hydrolysis of peptide bonds following aromatic amino acids.
   b. Lactase is found in the small intestine and catalyzes the hydrolysis of lactose.
   c. Pepsin is found in the stomach and catalyzes the hydrolysis of peptide bonds, primarily those that occur after aromatic amino acids.
   d. Maltase is found in the small intestine and catalyzes the hydrolysis of maltose.
5. a. Bile salts aid in digestion by dispersing lipids throughout the aqueous solution in the small intestine.
   b. Emulsification is important because lipids are not soluble in water; it breaks lipids up into smaller particles that can be more readily hydrolyzed by lipases.
7. a. gly-ala-phe and thr-leu
   b. ala-ile-tyr and ser-arg
   c. val-trp and arg-leu-cys

20.3: Overview of Stage II of Catabolism

Concept Review Exercises

1. What is a metabolic pathway?
2. What vitamin is required to make coenzyme A?
3. What is the net yield of Glycolysis as far as ATP?
4. Name the enzymes that are key regulatory sites in Glycolysis.
5. Why are the enzymes in the previous question targets for regulation?
6. Why is the priming phase necessary?
7. Draw the entire pathway for glycolysis including enzymes, reactants and products for each step.
8. Where does beta-oxidation occur?
9. What is the average net yield of ATP per carbon?
10. Where exactly is water formed during the process of fatty acid degradation? (Hint: H_2O is formed when when the one of the products of beta-oxidation is passed through another of the metabolic pathways)
11. During the process of beta-oxidation, why is it that [FAD] is used to oxidize an alkane to an alkene while NAD^+ is used to oxidize an alchol to a carbonyl
12. Draw out the entire process of the degradation of a triglyceride, include enzymes and products and reactants for each step.

Answers

1. A metabolic pathway is a series of biochemical reactions by which an organism converts a given reactant to a specific end product.
2. pantothenic acid

20.4: Stage III of Catabolism

Concept Review Exercises

1. What is the main function of the citric acid cycle?
2. Two carbon atoms are fed into the citric acid cycle as acetyl-CoA. In what form are two carbon atoms removed from the cycle?
3. What are mitochondria and what is their function in the cell?

Answers

1. the complete oxidation of carbon atoms to carbon dioxide and the formation of a high-energy phosphate compound, energy rich reduced coenzymes (NADH and $FADH_2$), and metabolic intermediates for the synthesis of other compounds
2. as carbon dioxide
3. Mitochondria are small organelles with a double membrane that contain the enzymes and other molecules needed for the production of most of the ATP needed by the body.

Exercises

1. Replace each question mark with the correct compound.
   a. $\text{?} \xrightarrow{\text{aconitase}} \text{isocitrate}$
   b. $\text{?} + \text{?} \xrightarrow{\text{citrate synthase}} \text{citrate} + \text{coenzyme A}$
   c. $\text{fumarate} \xrightarrow{\text{fumarase}} \text{?}$
   d. $\text{isocitrate} + \text{NAD}^+ \xrightarrow{\text{?}} \alpha\text{-ketoglurate} + \text{NADH} + \text{CO}_2$

2. Replace each question mark with the correct compound.

   a. $\text{malate} + \text{NAD}^+ \xrightarrow{?} \text{oxaloacetate} + \text{NADH}$

   b. $? + ? \xrightarrow{\text{nucleoside diphosphokinase}} \text{GDP} + \text{ATP}$

   c. $\text{succinyl-CoA} \xrightarrow{\text{succinyl-CoA synthetase}} ? + ?$

   d. $\text{succinate} + \text{FAD} \xrightarrow{\text{succinate dehydrogenase}} ? + \text{FADH}_2$

3. From the reactions in Exercises 1 and 2, select the equation(s) by number and letter in which each type of reaction occurs.

   a. isomerization

   b. hydration

   c. synthesis

4. From the reactions in Exercises 1 and 2, select the equation(s) by number and letter in which each type of reaction occurs.

   a. oxidation

   b. decarboxylation

   c. phosphorylation

5. What similar role do coenzyme Q and cytochrome c serve in the electron transport chain?

6. What is the electron acceptor at the end of the electron transport chain? To what product is this compound reduced?

7. What is the function of the cytochromes in the electron transport chain?

8. a. What is meant by this statement? “Electron transport is tightly coupled to oxidative phosphorylation.”

   b. How are electron transport and oxidative phosphorylation coupled or linked?

Answers

1. a. citrate

   b. oxaloacetate + acetyl-CoA

   c. malate

   d. α-ketoglutarate hydrogenase complex

3. a. reaction in 1a

   b. reaction in 1c

   c. reaction in 1b

5. Both molecules serve as electron shuttles between the complexes of the electron transport chain.

7. Cytochromes are proteins in the electron transport chain and serve as one-electron carriers.

20.5: Stage II of Carbohydrate Catabolism

- ### Concept Review Exercises

  1. In glycolysis, how many molecules of pyruvate are produced from one molecule of glucose?

  2. In vertebrates, what happens to pyruvate when

     a. plenty of oxygen is available?

     b. oxygen supplies are limited?

  3. In anaerobic glycolysis, how many molecules of ATP are produced from one molecule of glucose?

  #### Answers

  1. two

  2. a. Pyruvate is completely oxidized to carbon dioxide.

     b. Pyruvate is reduced to lactate, allowing for the reoxidation of NADH to NAD^+.

  3. There is a net production of two molecules of ATP.

Exercises

1. Replace each question mark with the correct compound.

   a. $\text{fructose 1, 6-bisphosphate} \xrightarrow{\text{aldolase}} ? + ?$

   b. $? + \text{ADP} \xrightarrow{\text{pyruvate kinase}} \text{pyruvate} + \text{ATP}$

   c. $\text{dihydroxyacetone phosphate} \xrightarrow{?} \text{glyceraldehyde 3-phosphate}$

   d. $\text{glucose} + \text{ATP} \xrightarrow{\text{hexokinase}} ? + \text{ADP}$

2. Replace each question mark with the correct compound.

   a. $\text{fructose 6-phosphate} + \text{ATP} \xrightarrow{?} \text{fructose 1, 6-bisphosphate} + \text{ADP}$

   b. $? \xrightarrow{\text{phosphoglucose isomerase}} \text{fructose 6-phosphate}$

   c. $\text{glyceraldehyde 3-phosphate} + \text{NAD}^+ + \text{P}_\text{i} \xrightarrow{?} \text{1, 3-bisphosphoglycerate} + \text{NADH}$

   d. $\text{3-phosphoglycerate} \xrightarrow{\text{phosphoglyceromutase}} ?$

3. From the reactions in Exercises 1 and 2, select the equation(s) by number and letter in which each type of reaction occurs.

   a. hydrolysis of a high-energy phosphate compound

   b. synthesis of ATP

4. From the reactions in Exercises 1 and 2, select the equation(s) by number and letter in which each type of reaction occurs.

   a. isomerization

   b. oxidation

5. What coenzyme is needed as an oxidizing agent in glycolysis?

6. Calculate

   a. the *total* number of molecules of ATP produced for each molecule of glucose converted to pyruvate in glycolysis.

   b. the number of molecules of ATP hydrolyzed in phase I of glycolysis.

   c. the *net* ATP production from glycolysis alone.

7. How is the NADH produced in glycolysis reoxidized when oxygen supplies are limited in

   a. muscle cells?

   b. yeast?

8. a. Calculate the number of moles of ATP produced by the aerobic oxidation of 1 mol of glucose in a liver cell.

   b. Of the total calculated in Exercise 9a, determine the number of moles of ATP produced in each process.

      a. glycolysis alone

      b. the citric acid cycle

      c. the electron transport chain and oxidative phosphorylation

Answers

1. a. glyceraldehyde 3-phosphate + dihydroxyacetone phosphate

   b. phosphoenolpyruvate

   c. triose phosphate isomerase

   d. glucose 6-phosphate

3. a. reactions 1b, 1d, and 2a

   b. reaction 1b

5. NAD^+

7. a. Pyruvate is reduced to lactate, and NADH is reoxidized to NAD^+.


   b. Pyruvate is converted to ethanol and carbon dioxide, and NADH is reoxidized to NAD^+.

20.6: Stage II of Lipid Catabolism

- ### Concept Review Exercises

1. How are fatty acids activated prior to being transported into the mitochondria and oxidized?
2. Draw the structure of hexanoic (caproic) acid and identify the α-carbon and the β-carbon.

Answers


1. They react with CoA to form fatty acyl-CoA molecules.
2. 


Key Takeaways


- Fatty acids, obtained from the breakdown of triglycerides and other lipids, are oxidized through a series of reactions known as β-oxidation.
- In each round of β-oxidation, 1 molecule of acetyl-CoA, 1 molecule of NADH, and 1 molecule of $FADH_2$ are produced.
- The acetyl-CoA, NADH, and $FADH_2$ are used in the citric acid cycle, the electron transport chain, and oxidative phosphorylation to produce ATP.

Exercises

1. For each reaction found in β-oxidation, identify the enzyme that catalyzes the reaction and classify the reaction as oxidation-reduction, hydration, or cleavage.

a. 

b. 

c. 

2. What are the products of β-oxidation?
3. How many rounds of β-oxidation are necessary to metabolize lauric acid (a saturated fatty acid with 12 carbon atoms)?
4. How many rounds of β-oxidation are necessary to metabolize arachidic acid (a saturated fatty acid with 20 carbon atoms)?
5. When myristic acid (a saturated fatty acid with 14 carbon atoms) is completely oxidized by β-oxidation, how many molecules of each are formed?
   a. acetyl-CoA
   b. $FADH_2$
   c. NADH
6. When stearic acid (a saturated fatty acid with 18 carbon atoms) is completely oxidized by β-oxidation, how many molecules of each are formed?
   a. acetyl-CoA

b. $FADH_2$

c. NADH

7. What is the net yield of ATP from the complete oxidation, in a liver cell, of one molecule of myristic acid?
8. What is the net yield of ATP from the complete oxidation, in a liver cell, of one molecule of stearic acid?

Answers

1. a. enoyl-CoA hydratase; hydration

   b. thiolase; cleavage

   c. acyl-CoA dehydrogenase; oxidation-reduction

3. five rounds
5. a. 7 molecules

   b. 6 molecules

   c. 6 molecules

7. 112 molecules

20.7: Stage II of Protein Catabolism

Concept Review Exercises


1. a. Write the equation for the transamination reaction between alanine and oxaloacetate.

   b. Name the two products that are formed.

2. What is the purpose of oxidative deamination?

Answers

1. a.



   b. pyruvate and aspartate

2. Oxidative deamination provides a reaction in which the amino group [as the ammonium (NH_4^+) ion] is removed from a molecule, not simply transferred from one molecule to another. Most of the NH_4^+ ion is converted to urea and excreted from the body.

Exercises

1. Write the equation for the transamination reaction between valine and pyruvate.
2. Write the equation for the transamination reaction between phenylalanine and oxaloacetate.
3. What products are formed in the oxidative deamination of glutamate?
4. Determine if each amino acid is glucogenic, ketogenic, or both.

   a. phenylalanine

   b. leucine

   c. serine

5. Determine if each amino acid is glucogenic, ketogenic, or both.

   a. asparagine

   b. tyrosine

   c. valine

Answers



1.

3. α-ketoglutarate, NADH, and NH_4^+

5. a. glucogenic
   b. both
   c. glucogenic

Additional Exercises

1. Hydrolysis of which compound—arginine phosphate or glucose 6-phosphate—would provide enough energy for the phosphorylation of ATP? Why?
2. If a cracker, which is rich in starch, is chewed for a long time, it begins to develop a sweet, sugary taste. Why?
3. Indicate where each enzymes would cleave the short peptide ala-ser-met-val-phe-gly-cys-lys-asp-leu.

   a. aminopeptidase
   b. chymotrypsin

4. Indicate where each enzymes would cleave the short peptide ala-ser-met-val-phe-gly-cys-lys-asp-leu.

   a. trypsin
   b. carboxypeptidase

5. If the methyl carbon atom of acetyl-CoA is labeled, where does the label appear after the acetyl-CoA goes through one round of the citric acid cycle?
6. If the carbonyl carbon atom of acetyl-CoA is labeled, where does the label appear after the acetyl-CoA goes through one round of the citric acid cycle?
7. The average adult consumes about 65 g of fructose daily (either as the free sugar or from the breakdown of sucrose). In the liver, fructose is first phosphorylated to fructose 1-phosphate, which is then split into dihydroxyacetone phosphate and glyceraldehyde. Glyceraldehyde is then phosphorylated to glyceraldehyde 3-phosphate, with ATP as the phosphate group donor. Write the equations (using structural formulas) for these three steps. Indicate the type of enzyme that catalyzes each step.
8. What critical role is played by both BPG and PEP in glycolysis?
9. How is the NADH produced in glycolysis reoxidized when oxygen supplies are abundant?
10. When a triglyceride is hydrolyzed to form three fatty acids and glycerol, the glycerol can be converted to glycerol 3-phosphate and then oxidized to form dihydroxyacetone phosphate, an intermediate of glycolysis. (In this reaction, NAD^+ is reduced to NADH.) If you assume that there is sufficient oxygen to completely oxidize the pyruvate formed from dihydroxyacetone phosphate, what is the maximum amount of ATP formed from the complete oxidation of 1 mol of glycerol?
11. How is the $FADH_2$ from β-oxidation converted back to FAD?
12. If 1 mol of alanine is converted to pyruvate in a muscle cell (through transamination) and the pyruvate is then metabolized via the citric acid cycle, the electron transport chain, and oxidative phosphorylation, how many moles of ATP are produced?
13. If the essential amino acid leucine (2-amino-4-methylpentanoic acid) is lacking in the diet, an α-keto acid can substitute for it. Give the structure of the α-keto acid and the probable reaction used to form leucine from this α-keto acid.

Answers


1. The hydrolysis of arginine phosphate releases more energy than is needed for the synthesis of ATP, while hydrolysis of glucose 6-phosphate does not.

3.

a. The enzyme will cleave off amino acids one at a time beginning with alanine (the N-terminal end).
b. following phenylalanine

5. Half of the label will be on the second carbon atom of oxaloacetate, while the other half will be on the third carbon atom.


7.



9. When oxygen is abundant, NADH is reoxidized through the reactions of the electron transport chain.

11. $FADH_2$ is reoxidized back to FAD via the electron transport chain.

13.



19.10: Energy Metabolism (Exercises) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.1: ATP- the Universal Energy Currency** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **20.3: Overview of Stage II of Catabolism** by Darik Benson is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **20.4: Stage III of Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **20.6: Stage II of Lipid Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **20.7: Stage II of Protein Catabolism** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

19.11: Energy Metabolism (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Metabolism is the general term for all chemical reactions in living organisms. The two types of metabolism are **catabolism**—those reactions in which complex molecules (carbohydrates, lipids, and proteins) are broken down to simpler ones with the concomitant release of energy—and **anabolism**—those reactions that consume energy to build complex molecules. Metabolism is studied by looking at individual **metabolic pathways**, which are a series of biochemical reactions in which a given reactant is converted to a desired end product.

The oxidation of fuel molecules (primarily carbohydrates and lipids), a process called **respiration**, is the source of energy used by cells. Catabolic reactions release energy from food molecules and use some of that energy for the synthesis of *adenosine triphosphate* (ATP); anabolic reactions use the energy in ATP to create new compounds. Catabolism can be divided into three stages. In stage I, carbohydrates, lipids, and proteins are broken down into their individual monomer units—simple sugars, fatty acids, and amino acids, respectively. In stage II, these monomer units are broken down by specific metabolic pathways to form a common end product *acetyl-coenzyme A* (CoA). In stage III, acetyl-CoA is completely oxidized to form carbon dioxide and water, and ATP is produced.

The **digestion** of carbohydrates begins in the mouth as α-amylase breaks glycosidic linkages in carbohydrate molecules. Essentially no carbohydrate digestion occurs in the stomach, and food particles pass through to the small intestine, where α-amylase and intestinal enzymes convert complex carbohydrate molecules (starches) to monosaccharides. The monosaccharides then pass through the lining of the small intestine and into the bloodstream for transport to all body cells.

Protein digestion begins in the stomach as pepsinogen in **gastric juice** is converted to pepsin, the enzyme that hydrolyzes peptide bonds. The partially digested protein then passes to the small intestine, where the remainder of protein digestion takes place through the action of several enzymes. The resulting amino acids cross the intestinal wall into the blood and are carried to the liver.

Lipid digestion begins in the small intestine. Bile salts emulsify the lipid molecules, and then lipases hydrolyze them to fatty acids and monoglycerides. The hydrolysis products pass through the intestine and are repackaged for transport in the bloodstream.

In cells that are operating aerobically, acetyl-CoA produced in stage II of catabolism is oxidized to carbon dioxide. The **citric acid cycle** describes this oxidation, which takes place with the formation of the coenzymes reduced nicotinamide adenine dinucleotide (NADH) and reduced flavin adenine dinucleotide ($FADH_2$). The sequence of reactions needed to oxidize these coenzymes and transfer the resulting electrons to oxygen is called the **electron transport chain**, or the **respiratory chain**. The compounds responsible for this series of oxidation-reduction reactions include proteins known as **cytochromes**, Fe·S proteins, and other molecules that ultimately result in the reduction of molecular oxygen to water. Every time a compound with two carbon atoms is oxidized in the citric acid cycle, a respiratory chain compound accepts the electrons lost in the oxidation (and so is reduced) and then passes them on to the next metabolite in the chain. The energy released by the electron transport chain is used to transport hydrogen (H^+) ions from the mitochondrial matrix to the intermembrane space. The flow of H^+ back through ATP synthase leads to the synthesis and release of ATP from adenosine diphosphate (ADP) and inorganic phosphate ions (P_i) in a process known as **oxidative phosphorylation**. Electron transport and oxidative phosphorylation are tightly coupled to each other. The enzymes and intermediates of the citric acid cycle, the electron transport chain, and oxidative phosphorylation are located in organelles called **mitochondria**.

The oxidation of carbohydrates is the source of over 50% of the energy used by cells. Glucose is oxidized to two molecules of pyruvate through a series of reactions known as **glycolysis**. Some of the energy released in these reactions is conserved by the formation of ATP from ADP. Glycolysis can be divided into two phases: phase I consists of the first five reactions and requires energy to “prime” the glucose molecule for phase II, the last five reactions in which ATP is produced through **substrate-level phosphorylation**.

The pyruvate produced by glycolysis has several possible fates, depending on the organism and whether or not oxygen is present. In animal cells, pyruvate can be further oxidized to acetyl-CoA and then to carbon dioxide (through the citric acid cycle) if oxygen supplies are sufficient. When oxygen supplies are insufficient, pyruvate is reduced to lactate. In yeast and other microorganisms, pyruvate is not converted to lactate in the absence of oxygen but instead is converted to ethanol and carbon dioxide.

The amount of ATP formed by the oxidation of glucose depends on whether or not oxygen is present. If oxygen is present, glucose is oxidized to carbon dioxide, and 36–38 ATP molecules are produced for each glucose molecule oxidized, using the combined pathways of glycolysis, the citric acid cycle, the electron transport chain, and oxidative phosphorylation. Thus, approximately 42% of the energy released by the complete oxidation of glucose is conserved by the synthesis of ATP. In the absence of oxygen, only 2 molecules of ATP are formed for each molecule of glucose converted to lactate (2 molecules), and the amount of energy conserved is much less (2%).

Fatty acids, released by the degradation of triglycerides and other lipids, are converted to fatty acyl-CoA, transported into the mitochondria, and oxidized by repeated cycling through a sequence of four reactions known as **β-oxidation**. In each round of β-oxidation, the fatty acyl-CoA is shortened by two carbon atoms as one molecule of acetyl-CoA is formed. The final round of β-oxidation, once the chain has been shortened to four carbon atoms, forms two molecules of acetyl-CoA. β-oxidation also forms the reduced coenzymes $FADH_2$ and NADH, whose reoxidation through the electron transport chain and oxidative phosphorylation leads to the synthesis of ATP. The efficiency of fatty acid oxidation in the human body is approximately 41%.

Amino acids from the breakdown of proteins can be catabolized to provide energy. Amino acids whose carbon skeletons are converted to intermediates that can be converted to glucose through gluconeogenesis are known as **glucogenic amino acids**. Amino acids whose carbon skeletons are broken down to compounds used to form ketone bodies are known as **ketogenic amino acids**.

The first step in amino acid catabolism is separation of the amino group from the carbon skeleton. In a **transamination**, the amino acid gives its NH_2 to pyruvate, α-ketoglutarate, or oxaloacetate. The products of this reaction are a new amino acid and an α-keto acid containing the carbon skeleton of the original amino acid. Pyruvate is transaminated to alanine, α-ketoglutarate to glutamate, and oxaloacetate to aspartate. The amino groups used to form alanine and aspartate are ultimately transferred to α-ketoglutarate, forming glutamate. The glutamate then undergoes **oxidative deamination** to yield α-ketoglutarate and ammonia.

Template:HideOrg

19.11: Energy Metabolism (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- **20.S: Energy Metabolism (Summary)** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

Index

A

addition reaction
13.1: Organic Reactions
ATP synthase
19.9: Oxidative Phosphorylation

C

catalyst
7.7: Factors Affecting Reaction Rates
condensation reaction
13.1: Organic Reactions

D

disaccharides
15.6: Disaccharides

E

elimination reaction
13.1: Organic Reactions
energy metabolism
19: Energy Metabolism

F

fatty acids
16.2: Fatty Acids

H

heterogeneous catalyst
7.8: Catalysis
homogeneous catalyst
7.8: Catalysis
hydrolysis
13.1: Organic Reactions

L

law of conservation of mass
6.2: The Law of Conservation of Matter
lipids
16.1: Introduction to Lipids

M

mass percentage
9.3: Other Units for Solution Concentrations
molarity
9.2: Solution Concentration- Molarity

O

oxidative phosphorylation
19.9: Oxidative Phosphorylation

P

parts per billion
9.3: Other Units for Solution Concentrations
parts per million
9.3: Other Units for Solution Concentrations
polymerization
13.1: Organic Reactions
polysaccharide
15.7: Polysaccharides
pressure
8.3: Gases and Pressure
prosthetic group
19.9: Oxidative Phosphorylation

R

radioactivity
3.1: Radioactivity
redox reaction
6.6: Oxidation-Reduction (Redox) Reactions
Redox Reactions (Organic Chemistry)
6.7: Redox Reactions in Organic Chemistry and Biochemistry

S

starch
15.7: Polysaccharides
stoichiometry
6.4: Quantitative Relationships Based on Chemical Equations
substitution reaction
13.1: Organic Reactions

U

ubiquinone
19.9: Oxidative Phosphorylation

V

volume percentage
9.3: Other Units for Solution Concentrations

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: CHEM 1121: General, Organic, and Biochemistry

Webpages: 185

Applicable Restrictions: Noncommercial

All licenses found:

- CC BY-NC-SA 4.0: 85.4% (158 pages)
- Undeclared: 6.5% (12 pages)
- CC BY-SA 4.0: 4.9% (9 pages)
- CC BY 4.0: 1.6% (3 pages)
- CK-12 License: 1.6% (3 pages)

By Page

- CHEM 1121: General, Organic, and Biochemistry - *Undeclared*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Measuring Matter and Energy - *CC BY-NC-SA 4.0*
    - 1.1: What is Chemistry - *CC BY-NC-SA 4.0*
    - 1.2: Classification of Matter - *CC BY-NC-SA 4.0*
    - 1.3: Introduction to Kinetic and Potential Energy - *CC BY-NC-SA 4.0*
    - 1.4: Measurements - *CC BY-NC-SA 4.0*
    - 1.5: Expressing Numbers- Scientific Notation - *CC BY-NC-SA 4.0*
    - 1.6: Expressing Numbers- Significant Figures - *CC BY-NC-SA 4.0*
    - 1.7: The International System of Units - *CC BY-NC-SA 4.0*
    - 1.8: Converting Units - *CC BY-NC-SA 4.0*
    - 1.9: Temperature Conversions - *CC BY-NC-SA 4.0*
    - 1.10: Specific Heat - *CC BY-NC-SA 4.0*
    - 1.11: Dosage Calculations - *CC BY-NC-SA 4.0*
    - 1.12: Chapter Summary - *CC BY-NC-SA 4.0*
  - 2: Atomic Structure - *CC BY-NC-SA 4.0*
    - 2.1: The Elements - *CC BY-NC-SA 4.0*
    - 2.2: Atomic Theory - *CC BY-NC-SA 4.0*
    - 2.3: The Structure of Atoms - *CC BY-NC-SA 4.0*
    - 2.4: Nuclei of Atoms - *CC BY-NC-SA 4.0*
    - 2.5: Arrangement of Electron (Shell Model) - *CC BY-NC-SA 4.0*
    - 2.6: The Periodic Table - *CC BY-NC-SA 4.0*
    - 2.7: Chapter Summary - *CC BY-NC-SA 4.0*
  - 3: Nuclear Chemistry - *CC BY-NC-SA 4.0*
    - 3.1: Radioactivity - *CC BY-NC-SA 4.0*
    - 3.2: Half-Life - *CC BY-NC-SA 4.0*
    - 3.3: Units of Radioactivity - *CC BY-NC-SA 4.0*
    - 3.4: Uses of Radioactive Isotopes - *CC BY-NC-SA 4.0*
    - 3.5: Nuclear Energy - *CC BY-NC-SA 4.0*
    - 3.6: 3.6 Chapter Summary - *CC BY-NC-SA 4.0*
    - 3.7: Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
  - 4: Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
    - 4.1: Two Types of Bonding - *CC BY-NC-SA 4.0*
    - 4.2: Ions - *CC BY-NC-SA 4.0*
    - 4.3: Formulas for Ionic Compounds - *CC BY-NC-SA 4.0*
    - 4.4: Ionic Nomenclature - *CC BY-NC-SA 4.0*
    - 4.5: Chapter Summary - *CC BY-NC-SA 4.0*
  - 5: Covalent Bonding and Simple Molecular Compounds - *CC BY-NC-SA 4.0*
    - 5.1: Covalent Bonds - *CC BY-NC-SA 4.0*
    - 5.2: Covalent Compounds- Formulas and Names - *CC BY-NC-SA 4.0*
    - 5.3: Multiple Covalent Bonds - *CC BY-NC-SA 4.0*
    - 5.4: Characteristics of Covalent Bonds - *CC BY-NC-SA 4.0*
    - 5.5: Characteristics of Molecules - *CC BY-NC-SA 4.0*
    - 5.6: Intermolecular Forces - *CC BY-NC-SA 4.0*
    - 5.7: Chapter Summary - *CC BY-NC-SA 4.0*
  - 6: Introduction to Chemical Reactions - *CC BY-SA 4.0*
    - 6.1: Prelude to Introduction to Chemical Reactions - *CC BY-SA 4.0*

- 6.2: The Law of Conservation of Matter - *CC BY-SA 4.0*
- 6.3: Chemical Equations - *CC BY-SA 4.0*
- 6.4: Quantitative Relationships Based on Chemical Equations - *CC BY-SA 4.0*
- 6.5: Types of Chemical Reactions - *Undeclared*
- 6.6: Oxidation-Reduction (Redox) Reactions - *CC BY-SA 4.0*
- 6.7: Redox Reactions in Organic Chemistry and Biochemistry - *CC BY-SA 4.0*
- 6.E: Introduction to Chemical Reactions (Exercises) - *CC BY-SA 4.0*
- 6.S: Introduction to Chemical Reactions (Summary) - *CC BY-SA 4.0*

- 7: Quantities in Chemical Reactions - *CC BY-NC-SA 4.0*
  - 7.1: Formula Mass - *CC BY-NC-SA 4.0*
  - 7.2: The Mole - *CC BY-NC-SA 4.0*
  - 7.3: Atomic and Molar Masses - *CC BY-NC-SA 4.0*
  - 7.4: Mole-Mass Conversions - *CC BY-NC-SA 4.0*
  - 7.5: Mole-Mole Relationships in Chemical Reactions - *CC BY-NC-SA 4.0*
  - 7.6: Energetics of Biochemical Reactions - *CC BY-NC-SA 4.0*
  - 7.7: Factors Affecting Reaction Rates - *CC BY 4.0*
  - 7.8: Catalysis - *CC BY 4.0*
  - 7.9: Equilbria and Le Chatelier's Principle - *CC BY-NC-SA 4.0*
  - 7.10: Chapter Summary - *CC BY-NC-SA 4.0*
- 8: States of Matter and the Gas Laws - *CC BY-NC-SA 4.0*
  - 8.1: Phase Changes - *CC BY-NC-SA 4.0*
  - 8.2: Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
  - 8.3: Gases and Pressure - *CC BY-NC-SA 4.0*
  - 8.4: Gas Laws - *CC BY-NC-SA 4.0*
  - 8.5: Chapter Summary - *CC BY-NC-SA 4.0*
- 9: Solutions - *CC BY-NC-SA 4.0*
  - 9.1: Introduction to Solutions - *CC BY-NC-SA 4.0*
  - 9.2: Solution Concentration- Molarity - *Undeclared*
  - 9.3: Other Units for Solution Concentrations - *CC BY 4.0*
  - 9.4: The Dissolution Process - *CC BY-NC-SA 4.0*
  - 9.5: Properties of Solutions - *CC BY-NC-SA 4.0*
  - 9.6: Dialysis and Tonicity - *CC BY-NC-SA 4.0*
  - 9.7: Chapter Summary - *CC BY-NC-SA 4.0*
- 10: Acids and Bases - *CC BY-NC-SA 4.0*
  - 10.1: Arrhenius Definition of Acids and Bases - *CC BY-NC-SA 4.0*
  - 10.2: Brønsted-Lowry Definition of Acids and Bases - *CC BY-NC-SA 4.0*
  - 10.3: Water is both an Acid and a Base - *CC BY-NC-SA 4.0*
  - 10.4: The Strengths of Acids and Bases - *CC BY-NC-SA 4.0*
  - 10.5: pH and Kw - *CC BY-NC-SA 4.0*
  - 10.6: Buffers - *CC BY-NC-SA 4.0*
  - 10.7: End-of-Chapter Material - *CC BY-NC-SA 4.0*
- 11: Organic Chemistry of Hydrocarbons - *CC BY-NC-SA 4.0*
  - 11.1: Introduction to Organic Chemistry - *CC BY-NC-SA 4.0*
  - 11.2: Organic Chemistry - *CC BY-NC-SA 4.0*
  - 11.3: Structures and Names of Alkanes - *CC BY-NC-SA 4.0*
  - 11.4: Branched-Chain Alkanes - *CC BY-NC-SA 4.0*
  - 11.5: Condensed Structural and Line-Angle Formulas - *CC BY-NC-SA 4.0*
  - 11.6: Cycloalkanes - *CC BY-NC-SA 4.0*
  - 11.7: IUPAC Nomenclature - *CC BY-NC-SA 4.0*
  - 11.8: Alkenes- Structure and Nomenclature - *CC BY-NC-SA 4.0*
  - 11.9: Alkenes- Cis-Tran Isomerization - *CC BY-NC-SA 4.0*
  - 11.10: Alkynes - *CC BY-NC-SA 4.0*
  - 11.11: Aromatic Compounds - *CC BY-NC-SA 4.0*
  - 11.12: Aromatic Compounds- Structure and Nomenclature - *CC BY-NC-SA 4.0*
  - 11.13: Physical Properties of Hydrocarbons - *CC BY-NC-SA 4.0*
  - 11.14: Chapter Summary - *CC BY-NC-SA 4.0*
- 12: Organic Functional Groups - Structure and Nomenclature - *CC BY-NC-SA 4.0*
  - 12.1: Organic Compounds with Functional Groups - *CC BY-NC-SA 4.0*
  - 12.2: 9.2 Alcohols and Phenols- Nomenclature and Classification - *CC BY-NC-SA 4.0*
  - 12.3: Glycols and Glycerol - *CC BY-NC-SA 4.0*
  - 12.4: Ethers - *CC BY-NC-SA 4.0*
  - 12.5: Aldehydes and Ketones- Structure and Names - *CC BY-NC-SA 4.0*
  - 12.6: Organic Sulfur Compounds - *CC BY-NC-SA 4.0*
  - 12.7: Carboxylic Acids- Structures and Names - *CC BY-NC-SA 4.0*
  - 12.8: Esters- Structures and Names - *CC BY-NC-SA 4.0*
  - 12.9: Phosphate Esters - *CC BY-NC-SA 4.0*
  - 12.10: Amines - *CC BY-NC-SA 4.0*
  - 12.11: Amides - *CC BY-NC-SA 4.0*
- 13: Organic Reactions - *CK-12 License*

- 13.1: Organic Reactions - *CK-12 License*
- 13.2: Organic Reactions (Exercises) - *CK-12 License*

- 14: Properties of Organic Functional Groups - *CC BY-NC-SA 4.0*
  - 14.1: Physical Properties of Alcohols - *CC BY-NC-SA 4.0*
  - 14.2: Physical Properties of Aldehydes and Ketones - *CC BY-NC-SA 4.0*
  - 14.3: Physical Properties of Carboxylic Acids - *CC BY-NC-SA 4.0*
  - 14.4: Physical Properties of Esters - *CC BY-NC-SA 4.0*
  - 14.5: Physical Properties of Amines - *CC BY-NC-SA 4.0*
  - 14.6: Physical Properties of Amides - *CC BY-NC-SA 4.0*
  - 14.7: Ionization of Carboxylic Acids - *CC BY-NC-SA 4.0*
  - 14.8: Amines as Bases - *CC BY-NC-SA 4.0*
  - 14.9: Chirality - *CC BY-NC-SA 4.0*
  - 14.10: End-of-Chapter Material - *CC BY-NC-SA 4.0*
- 15: Carbohydrates - *CC BY-NC-SA 4.0*
  - 15.1: Introduction to Carbohydrates - *CC BY-NC-SA 4.0*
  - 15.2: Classes of Monosaccharides - *CC BY-NC-SA 4.0*
  - 15.3: Important Hexoses - *CC BY-NC-SA 4.0*
  - 15.4: Cyclic Structures of Monosaccharides - *CC BY-NC-SA 4.0*
  - 15.5: Properties of Monosaccharides - *CC BY-NC-SA 4.0*
  - 15.6: Disaccharides - *CC BY-NC-SA 4.0*
  - 15.7: Polysaccharides - *CC BY-NC-SA 4.0*
  - 15.8: Diabetes - *CC BY-NC-SA 4.0*
  - 15.9: Chapter Summary - *CC BY-NC-SA 4.0*
- 16: Lipids - *CC BY-NC-SA 4.0*
  - 16.1: Introduction to Lipids - *CC BY-NC-SA 4.0*
  - 16.2: Fatty Acids - *CC BY-NC-SA 4.0*
  - 16.3: Fats and Oils - *CC BY-NC-SA 4.0*
  - 16.4: Membranes and Membrane Lipids - *CC BY-NC-SA 4.0*
  - 16.5: Steroids - *CC BY-NC-SA 4.0*
  - 16.6: Chapter Summary - *CC BY-NC-SA 4.0*
- 17: Proteins and Enzymes - *CC BY-NC-SA 4.0*
  - 17.1: Properties of Amino Acids - *CC BY-NC-SA 4.0*
  - 17.2: Reactions of Amino Acids - *CC BY-NC-SA 4.0*
  - 17.3: Peptides - *CC BY-NC-SA 4.0*
  - 17.4: Proteins - *CC BY-NC-SA 4.0*
  - 17.5: Enzymes - *CC BY-NC-SA 4.0*
  - 17.6: Enzymatic Catalysis - *CC BY-NC-SA 4.0*
  - 17.7: Enzymatic Activity - *CC BY-NC-SA 4.0*
  - 17.8: Enzyme Inhibition - *CC BY-NC-SA 4.0*
  - 17.9: Enzyme Cofactors and Vitamins - *CC BY-NC-SA 4.0*
  - 17.10: Chapter Summary - *CC BY-NC-SA 4.0*
- 18: Nucleic Acids - *CC BY-NC-SA 4.0*
  - 18.1: Nucleotides - *CC BY-NC-SA 4.0*
  - 18.2: Nucleic Acid Structure - *CC BY-NC-SA 4.0*
  - 18.3: Replication and Expression of Genetic Information - *CC BY-NC-SA 4.0*
  - 18.4: Protein Synthesis and the Genetic Code - *CC BY-NC-SA 4.0*
  - 18.5: Mutations and Genetic Diseases - *CC BY-NC-SA 4.0*
  - 18.6: Viruses - *CC BY-NC-SA 4.0*
  - 18.7: Chapter Summary - *CC BY-NC-SA 4.0*
- 19: Energy Metabolism - *CC BY-NC-SA 4.0*
  - 19.1: Prelude to Energy Metabolism - *CC BY-NC-SA 4.0*
  - 19.2: ATP- the Universal Energy Currency - *CC BY-NC-SA 4.0*
  - 19.3: Stage I of Catabolism - *CC BY-NC-SA 4.0*
  - 19.4: Overview of Stage II of Catabolism - *CC BY-NC-SA 4.0*
  - 19.5: Stage III of Catabolism - *CC BY-NC-SA 4.0*
  - 19.6: Stage II of Carbohydrate Catabolism - *CC BY-NC-SA 4.0*
  - 19.7: Stage II of Lipid Catabolism - *CC BY-NC-SA 4.0*
  - 19.8: Stage II of Protein Catabolism - *CC BY-NC-SA 4.0*
  - 19.9: Oxidative Phosphorylation - *Undeclared*
  - 19.10: Energy Metabolism (Exercises) - *CC BY-NC-SA 4.0*
  - 19.11: Energy Metabolism (Summary) - *CC BY-NC-SA 4.0*
- Back Matter - *CC BY-NC-SA 4.0*
  - Index - *Undeclared*
  - Glossary - *Undeclared*
  - Detailed Licensing - *Undeclared*