CHEMISTRY 305:INTRODUCTION TO CHEMISTRY



Caitlin Zumalt
American River College

LibreTexts™

ARC CHEM 305 (Zumalt) GOB

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 03/11/2026

TABLE OF CONTENTS

Licensing

1: Unit 1

- 1.1: The Scope of Chemistry
- 1.2: The Scientific Method - How Chemists Think
- 1.3: Hypothesis, Theories, and Laws
- 1.4: Taking Measurements
- 1.5: The Basic Units of Measurement
- 1.6: Units Raised to a Power
- 1.7: Scientific Notation - Writing Large and Small Numbers
- 1.8: Significant Figures - Writing Numbers to Reflect Precision
- 1.9: Significant Figures in Calculations
- 1.10: Problem Solving and Unit Conversions
- 1.11: Solving Multi-step Conversion Problems
- 1.12: Density
- 1.13: Temperature - Random Motion of Molecules and Atoms
- 1.E: Measurement and Problem Solving (Exercises)

2: Unit 2

- 2.1: In Your Room
- 2.2: What is Matter?
- 2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas
- 2.4: Classifying Matter According to Its Composition
- 2.5: Differences in Matter- Physical and Chemical Properties
- 2.6: Changes in Matter - Physical and Chemical Changes
- 2.7: Conservation of Mass - There is No New Matter
- 2.8: Cutting Aluminum until you get Atoms
- 2.9: Indivisible - The Atomic Theory
- 2.10: The Nuclear Atom
- 2.11: The Properties of Protons, Neutrons, and Electrons
- 2.12: Elements- Defined by Their Number of Protons
- 2.13: Looking for Patterns - The Periodic Table
- 2.14: Ions - Losing and Gaining Electrons
- 2.15: Isotopes - When the Number of Neutrons Varies
- 2.16: Atomic Mass - The Average Mass of an Element's Atoms
- 2.17: Sugar and Salt
- 2.18: Compounds Display Constant Composition
- 2.19: Chemical Formulas - How to Represent Compounds
- 2.20: A Molecular View of Elements and Compounds
- 2.21: Writing Formulas for Ionic Compounds
- 2.22: Nomenclature- Naming Compounds
- 2.23: Naming Ionic Compounds
- 2.24: Naming Molecular Compounds
- 2.25: Naming Acids
- 2.26: Nomenclature Summary

3: Unit 3

- 3.1: Evidence of a Chemical Reaction
- 3.2: The Chemical Equation
- 3.3: How to Write Balanced Chemical Equations
- 3.4: Classifying Chemical Reactions
- 3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water
- 3.6: Precipitation Reactions
- 3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
- 3.8: Prelude to Chemical Composition - How Much Sodium?
- 3.9: Counting Nails by the Pound
- 3.10: Counting Atoms by the Gram
- 3.11: Counting Molecules by the Gram
- 3.12: Chemical Formulas as Conversion Factors
- 3.13: Mass Percent Composition of Compounds
- 3.14: Mass Percent Composition from a Chemical Formula
- 3.15: Calculating Empirical Formulas for Compounds
- 3.16: Calculating Molecular Formulas for Compounds
- 3.17: Making Pancakes- Relationships Between Ingredients
- 3.18: Making Molecules- Mole-to-Mole Conversions
- 3.19: Making Molecules- Mass-to-Mass Conversions
- 3.20: Stoichiometry
- 3.21: Limiting Reactant and Theoretical Yield
- 3.22: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants

4: Unit 4

- 4.1: The Bohr Model - Atoms with Orbits
- 4.2: Quantum-Mechanical Orbitals and Electron Configurations
- 4.3: Electron Configurations and the Periodic Table
- 4.4: The Explanatory Power of the Quantum-Mechanical Model
- 4.5: Representing Valence Electrons with Dots
- 4.6: Lewis Structures of Ionic Compounds- Electrons Transferred
- 4.7: Covalent Lewis Structures- Electrons Shared
- 4.8: Writing Lewis Structures for Covalent Compounds
- 4.9: Resonance - Equivalent Lewis Structures for the Same Molecule
- 4.10: Predicting the Shapes of Molecules
- 4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix
- 4.12: Kinetic Molecular Theory- A Model for Gases
- 4.13: Pressure - The Result of Constant Molecular Collisions
- 4.14: Boyle's Law - Pressure and Volume
- 4.15: Charles's Law- Volume and Temperature
- 4.16: Gay-Lussac's Law- Temperature and Pressure
- 4.17: The Combined Gas Law- Pressure, Volume, and Temperature
- 4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles
- 4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen

5: Unit 5

- 5.1: Intermolecular Forces- Dispersion, Dipole–Dipole, Hydrogen Bonding, and Ion-Dipole
- 5.2: Solutions - Homogeneous Mixtures

- 5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy
- 5.4: Specifying Solution Concentration- Mass Percent
- 5.5: Specifying Solution Concentration- Molarity
- 5.6: Solution Stoichiometry
- 5.7: Sour Patch Kids and International Spy Movies
- 5.8: Acids- Properties and Examples
- 5.9: Bases- Properties and Examples
- 5.10: Molecular Definitions of Acids and Bases
- 5.11: Reactions of Acids and Bases
- 5.12: Acid–Base Titration
- 5.13: Strong and Weak Acids and Bases
- 5.14: Water - Acid and Base in One
- 5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity
- 5.16: Carbon - A Versitile Atom
- 5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
- 5.18: Alkanes- Saturated Hydrocarbons
- 5.19: Isomers- Same Formula, Different Structure
- 5.20: Naming Alkanes
- 5.21: Alkenes and Alkynes
- 5.22: Functional Groups
- 5.23: Carbohydrates
- 5.24: Lipids
- 5.25: Proteins

Index

Glossary

Detailed Licensing

Detailed Licensing

Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

CHAPTER OVERVIEW

1: Unit 1

1.1: The Scope of Chemistry
1.2: The Scientific Method - How Chemists Think
1.3: Hypothesis, Theories, and Laws
1.4: Taking Measurements
1.5: The Basic Units of Measurement
1.6: Units Raised to a Power
1.7: Scientific Notation - Writing Large and Small Numbers
1.8: Significant Figures - Writing Numbers to Reflect Precision
1.9: Significant Figures in Calculations
1.10: Problem Solving and Unit Conversions
1.11: Solving Multi-step Conversion Problems
1.12: Density
1.13: Temperature - Random Motion of Molecules and Atoms
1.E: Measurement and Problem Solving (Exercises)

1: Unit 1 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

1.1: The Scope of Chemistry

Learning Objectives

- To recognize the breadth, depth, and scope of chemistry.
- Define chemistry in relation to other sciences.
- Identify the main disciplines of chemistry.

Chemistry is the study of matter—what it consists of, what its properties are, and how it changes. Matter is anything that has mass and takes up space—that is, anything that is physically real. Some things are easily identified as matter—the screen on which you are reading this book, for example. Others are not so obvious. Because we move so easily through air, we sometimes forget that it, too, is matter. Because of this, chemistry is a science that has its fingers in just about everything. Being able to describe the ingredients in a cake and how they change when the cake is baked, for example, is chemistry!

Chemistry is one branch of science. Science is the process by which we learn about the natural universe by observing, testing, and then generating models that explain our observations. Because the physical universe is so vast, there are many different branches of science (Figure **1.1.1**). Thus, chemistry is the study of matter, biology is the study of living things, and geology is the study of rocks and the earth. Mathematics is the language of science, and we will use it to communicate some of the ideas of chemistry.




Figure **1.1.1**: The Relationships between Some of the Major Branches of Science. Chemistry lies more or less in the middle, which emphasizes its importance to many branches of science.

Although we divide science into different fields, there is much overlap among them. For example, some biologists and chemists work in both fields so much that their work is called biochemistry. Similarly, geology and chemistry overlap in the field called geochemistry. Figure **1.1.1** shows how many of the individual fields of science are related. At some level, all of these fields depend on matter because they all involve "stuff"; because of this, chemistry has been called the "central science", linking them all together.

> *There are many other fields of science, in addition to the ones (biology, medicine, etc.) listed here.*

Example 1.1.1: Science Fields

Which fields of study are branches of science? Explain.

a. sculpture
b. astronomy

Solution

a. Sculpture is not considered a science because it is not a study of some aspect of the natural universe.
b. Astronomy is the study of stars and planets, which are part of the natural universe. Astronomy is therefore a field of science.

Exercise 1.1.1

Which fields of study are branches of science?

a. physiology (the study of the function of an animal's or a plant's body)
b. geophysics
c. agriculture
d. politics

Answer a:

yes

Answer b:

yes

Answer c:

yes

Answer d:

no

Areas of Chemistry

The study of modern chemistry has many branches, but can generally be broken down into five main disciplines, or areas of study:

- **Physical chemistry:** Physical chemistry is the study of macroscopic properties, atomic properties, and phenomena in chemical systems. A physical chemist may study such things as the rates of chemical reactions, the energy transfers that occur in reactions, or the physical structure of materials at the molecular level.
- **Organic chemistry:** Organic chemistry is the study of chemicals containing carbon. Carbon is one of the most abundant elements on Earth and is capable of forming a tremendously vast number of chemicals (over twenty million so far). Most of the chemicals found in all living organisms are based on carbon.
- **Inorganic chemistry:** Inorganic chemistry is the study of chemicals that, in general, are not primarily based on carbon. Inorganic chemicals are commonly found in rocks and minerals. One current important area of inorganic chemistry deals with the design and properties of materials involved in energy and information technology.
- **Analytical chemistry:** Analytical chemistry is the study of the composition of matter. It focuses on separating, identifying, and quantifying chemicals in samples of matter. An analytical chemist may use complex instruments to analyze an unknown material in order to determine its various components.
- **Biochemistry:** Biochemistry is the study of chemical processes that occur in living things. Research may cover anything from basic cellular processes up to understanding disease states so that better treatments can be developed.



Figure **1.1.2**: (left) Measurement of trace metals using atomic spectroscopy. (right) Measurement of hormone concentrations.

In practice, chemical research is often not limited to just one of the five major disciplines. A particular chemist may use biochemistry to isolate a particular chemical found in the human body such as hemoglobin, the oxygen carrying component of red blood cells. He or she may then proceed to analyze the hemoglobin using methods that would pertain to the areas of physical or analytical chemistry. Many chemists specialize in areas that are combinations of the main disciplines, such as bioinorganic chemistry or physical organic chemistry.

History of Chemistry

The **history** of chemistry is an interesting and challenging one. Very early chemists were often motivated mainly by the achievement of a specific goal or product. Making perfume or soaps did not need a lot of **theory**, just a good recipe and careful attention to detail. There was no standard way of naming materials (and no periodic table that we could all agree on). It is often difficult to figure out exactly what a particular person was using. However, the science developed over the centuries by trial and error.

Major progress was made toward putting chemistry on a solid foundation when Robert Boyle (1637-1691) began his **research** in chemistry (Figure **1.1.3**). He developed the basic ideas about the behavior of gases. He could then describe gases mathematically. Boyle also helped form the idea that small particles could combine to form molecules. Many years later, John Dalton used these ideas to develop the atomic theory.



Figure **1.1.3**: Robert Boyle.

The field of chemistry began to develop rapidly in the 1700's. Joseph Priestley (1733-1804) isolated and characterized several gases: oxygen, carbon monoxide, and nitrous oxide. It was later discovered that nitrous oxide ("laughing gas") worked as an anesthetic. This gas was used for that purpose for the first time in 1844 during a tooth extraction. Other gases discovered during that time were chlorine, by C.W. Scheele (1742-1786) and nitrogen, by Antoine Lavoisier (1743-1794). Lavoisier has been considered by many scholars to be the "father of chemistry". Among other accomplishments, he discovered the role of oxygen in combustion and definitively formulated the law of conservation of matter.

Chemists continued to discover new compounds in the 1800's. The science also began to develop a more theoretical foundation. John Dalton (1766-1844) put forth his atomic theory in 1807. This idea allowed scientists to think about chemistry in a much more systematic way. Amadeo Avogadro (1776-1856) laid the groundwork for a more quantitative approach to chemistry by calculating the number of particles in a given amount of a gas. A lot of effort was put forth in studying chemical reactions. These efforts led to new materials being produced. Following the invention of the battery by Alessandro Volta (1745-1827), the field of electrochemistry (both theoretical and applications) developed through major contributions by Humphry Davy (1778-1829) and Michael Faraday (1791-1867). Other areas of the discipline also progressed rapidly.

It would take a large book to cover developments in chemistry during the twentieth century and up to today. One major area of expansion was in the area of the chemistry of living processes. Research in photosynthesis in plants, the discovery and

characterization of enzymes as biochemical catalysts, elucidation of the structures of biomolecules such as insulin and DNA—these efforts gave rise to an explosion of information in the field of biochemistry.

The practical aspects of chemistry were not ignored. The work of Volta, Davy, and Faraday eventually led to the development of batteries that provided a source of electricity to power a number of devices (Figure **1.1.4**).




Figure **1.1.4**: Battery developed by Volta. (CC BY-SA 3.0; (left) GuidoB and (right) Kkkdc).

Charles Goodyear (1800-1860) discovered the process of vulcanization, allowing a stable rubber product to be produced for the tires of all the vehicles we have today. Louis Pasteur (1822-1895) pioneered the use of heat sterilization to eliminate unwanted microorganisms in wine and milk. Alfred Nobel (1833-1896) invented dynamite (Figure **1.1.5**). After his death, the fortune he made from this product was used to fund the Nobel Prizes in science and the humanities. J.W. Hyatt (1837-1920) developed the first plastic. Leo Baekeland (1863-1944) developed the first synthetic resin, widely used for inexpensive and sturdy dinnerware.



Figure **1.1.5**: Dynamite explosion in Panama, Central America (1908).

Today, chemistry continues to be essential to the development of new materials and technologies, from semiconductors for electronics to powerful new medicines, and beyond.

Summary

- Chemistry is the study of matter and the changes it undergoes and considers both *macroscopic* and *microscopic* information.
- Matter is anything that has mass and occupies space.
- The five main disciplines of chemistry are physical chemistry, organic chemistry, inorganic chemistry, analytical chemistry and biochemistry.
- Many civilizations contributed to the growth of chemistry. A lot of early chemical research focused on practical uses. Basic chemistry theories were developed during the nineteenth century. New materials and batteries are a few of the products of modern chemistry.

1.1: The Scope of Chemistry is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **1.7: The Scope of Chemistry** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.2: The Scientific Method - How Chemists Think

Learning Objectives

- Identify the components of the scientific method.

Scientists search for answers to questions and solutions to problems by using a procedure called the scientific method. This procedure consists of making observations, formulating hypotheses, and designing experiments; which leads to additional observations, hypotheses, and experiments in repeated cycles (Figure 1.2.1).




Figure 1.2.1: The Steps in the Scientific Method.

Step 1: Make observations

Observations can be qualitative or quantitative. **Qualitative observations** describe properties or occurrences in ways that do not rely on numbers. Examples of qualitative observations include the following: "the outside air temperature is cooler during the winter season," "table salt is a crystalline solid," "sulfur crystals are yellow," and "dissolving a penny in dilute nitric acid forms a blue solution and a brown gas." **Quantitative observations** are measurements, which by definition consist of both a number and a unit. Examples of quantitative observations include the following: "the melting point of crystalline sulfur is 115.21° Celsius," and "35.9 grams of table salt—the chemical name of which is sodium chloride—dissolve in 100 grams of water at 20° Celsius." For the question of the dinosaurs' extinction, the initial observation was quantitative: iridium concentrations in sediments dating to 66 million years ago were 20–160 times higher than normal.

Step 2: Formulate a hypothesis

After deciding to learn more about an observation or a set of observations, scientists generally begin an investigation by forming a hypothesis, a tentative explanation for the observation(s). The hypothesis may not be correct, but it puts the scientist's understanding of the system being studied into a form that can be tested. For example, the observation that we experience alternating periods of light and darkness corresponding to observed movements of the sun, moon, clouds, and shadows is consistent with either one of two hypotheses:

a. Earth rotates on its axis every 24 hours, alternately exposing one side to the sun.
b. The sun revolves around Earth every 24 hours.

Suitable experiments can be designed to choose between these two alternatives. For the disappearance of the dinosaurs, the hypothesis was that the impact of a large extraterrestrial object caused their extinction. Unfortunately (or perhaps fortunately), this hypothesis does not lend itself to direct testing by any obvious experiment, but scientists can collect additional data that either support or refute it.

Step 3: Design and perform experiments

After a hypothesis has been formed, scientists conduct experiments to test its validity. Experiments are systematic observations or measurements, preferably made under controlled conditions—that is—under conditions in which a single variable changes.

Step 4: Accept or modify the hypothesis

A properly designed and executed experiment enables a scientist to determine whether or not the original hypothesis is valid. If the hypothesis is valid, the scientist can proceed to step 5. In other cases, experiments often demonstrate that the hypothesis is incorrect or that it must be modified and requires further experimentation.

Step 5: Development into a law and/or theory

More experimental data are then collected and analyzed, at which point a scientist may begin to think that the results are sufficiently reproducible (i.e., dependable) to merit being summarized in a law, a verbal or mathematical description of a phenomenon that allows for general predictions. A law simply states what happens; it does not address the question of why.

One example of a law, the law of definite proportions, which was discovered by the French scientist Joseph Proust (1754–1826), states that a chemical substance always contains the same proportions of elements by mass. Thus, sodium chloride (table salt) always contains the same proportion by mass of sodium to chlorine, in this case 39.34% sodium and 60.66% chlorine by mass, and sucrose (table sugar) is always 42.11% carbon, 6.48% hydrogen, and 51.41% oxygen by mass.

Whereas a law states only what happens, a theory attempts to explain why nature behaves as it does. Laws are unlikely to change greatly over time unless a major experimental error is discovered. In contrast, a theory, by definition, is incomplete and imperfect, evolving with time to explain new facts as they are discovered.

Because scientists can enter the cycle shown in Figure **1.2.1** at any point, the actual application of the scientific method to different topics can take many different forms. For example, a scientist may start with a hypothesis formed by reading about work done by others in the field, rather than by making direct observations.

✔ Example 1.2.1

Classify each statement as a law, a theory, an experiment, a hypothesis, an observation.

a. Ice always floats on liquid water.
b. Birds evolved from dinosaurs.
c. Hot air is less dense than cold air, probably because the components of hot air are moving more rapidly.
d. When 10 g of ice were added to 100 mL of water at 25°C, the temperature of the water decreased to 15.5°C after the ice melted.
e. The ingredients of Ivory soap were analyzed to see whether it really is 99.44% pure, as advertised.

Solution

a. This is a general statement of a relationship between the properties of liquid and solid water, so it is a law.
b. This is a possible explanation for the origin of birds, so it is a hypothesis.
c. This is a statement that tries to explain the relationship between the temperature and the density of air based on fundamental principles, so it is a theory.
d. The temperature is measured before and after a change is made in a system, so these are observations.
e. This is an analysis designed to test a hypothesis (in this case, the manufacturer's claim of purity), so it is an experiment.

? Exercise 1.2.1

Classify each statement as a law, a theory, an experiment, a hypothesis, a qualitative observation, or a quantitative observation.

a. Measured amounts of acid were added to a Rolaids tablet to see whether it really "consumes 47 times its weight in excess stomach acid."
b. Heat always flows from hot objects to cooler ones, not in the opposite direction.
c. The universe was formed by a massive explosion that propelled matter into a vacuum.

d. Michael Jordan is the greatest pure shooter to ever play professional basketball.
e. Limestone is relatively insoluble in water, but dissolves readily in dilute acid with the evolution of a gas.

Answer a
experiment
Answer b
law
Answer c
theory
Answer d
hypothesis
Answer e
observation

Summary

The scientific method is a method of investigation involving experimentation and observation to acquire new knowledge, solve problems, and answer questions. The key steps in the scientific method include the following:

- Step 1: Make observations.
- Step 2: Formulate a hypothesis.
- Step 3: Test the hypothesis through experimentation.
- Step 4: Accept or modify the hypothesis.
- Step 5: Develop into a law and/or a theory.

Contributions & Attributions

-
- Wikipedia

1.2: The Scientific Method - How Chemists Think is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **1.3: The Scientific Method - How Chemists Think** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.3: Hypothesis, Theories, and Laws

Learning Objectives

- Describe the difference between hypothesis and theory as scientific terms.
- Describe the difference between a theory and scientific law.

Although many have taken science classes throughout the course of their studies, people often have incorrect or misleading ideas about some of the most important and basic principles in science. Most students have heard of hypotheses, theories, and laws, but what do these terms really mean? Prior to reading this section, consider what you have learned about these terms before. What do these terms mean to you? What do you read that contradicts or supports what you thought?

What is a Fact?

A fact is a basic statement established by experiment or observation. All facts are true under the specific conditions of the observation.

What is a Hypothesis?

One of the most common terms used in science classes is a "hypothesis". The word can have many different definitions, depending on the context in which it is being used:

- An educated guess: a scientific hypothesis provides a suggested solution based on evidence.
- Prediction: if you have ever carried out a science experiment, you probably made this type of hypothesis when you predicted the outcome of your experiment.
- Tentative or proposed explanation: hypotheses can be suggestions about why something is observed. In order for it to be scientific, however, a scientist must be able to test the explanation to see if it works and if it is able to correctly predict what will happen in a situation. For example, "if my hypothesis is correct, we should see ___ result when we perform ___ test."

> *A hypothesis is very tentative; it can be easily changed.*

What is a Theory?

The *United States National Academy of Sciences* describes what a theory is as follows:

> "Some scientific explanations are so well established that no new evidence is likely to alter them. The explanation becomes a scientific theory. In everyday language a theory means a hunch or speculation. Not so in science. In science, the word **theory** refers to a comprehensive explanation of an important feature of nature supported by facts gathered over time. Theories also allow scientists to make predictions about as yet unobserved phenomena."
>
> "A scientific theory is a well-substantiated explanation of some aspect of the natural world, based on a body of facts that have been repeatedly confirmed through observation and experimentation. Such fact-supported theories are not "guesses" but reliable accounts of the real world. The theory of biological evolution is more than "just a theory." It is as factual an explanation of the universe as the atomic theory of matter (stating that everything is made of atoms) or the germ theory of disease (which states that many diseases are caused by germs). Our understanding of gravity is still a work in progress. But the phenomenon of gravity, like evolution, is an accepted fact.

Note some key features of theories that are important to understand from this description:

- Theories are explanations of natural phenomena. They aren't predictions (although we may use theories to make predictions). They are explanations as to why we observe something.
- Theories aren't likely to change. They have a large amount of support and are able to satisfactorily explain numerous observations. Theories can, indeed, be facts. Theories can change, but it is a long and difficult process. In order for a theory to change, there must be many observations or pieces of evidence that the theory cannot explain.
- Theories are not guesses. The phrase "just a theory" has no room in science. To be a scientific theory carries a lot of weight; it is not just one person's idea about something

> *Theories aren't likely to change.*

What is a Law?

Scientific laws are similar to scientific theories in that they are principles that can be used to predict the behavior of the natural world. Both scientific laws and scientific theories are typically well-supported by observations and/or experimental evidence. Usually scientific laws refer to rules for how nature will behave under certain conditions, frequently written as an equation. Scientific theories are more overarching explanations of how nature works and why it exhibits certain characteristics. As a comparison, theories explain why we observe what we do and laws describe what happens.

For example, around the year 1800, Jacques Charles and other scientists were working with gases to, among other reasons, improve the design of the hot air balloon. These scientists found, after many, many tests, that certain patterns existed in the observations on gas behavior. If the temperature of the gas is increased, the volume of the gas increased. This is known as a natural law. A law is a relationship that exists between variables in a group of data. Laws describe the patterns we see in large amounts of data, but do not describe why the patterns exist.

What is a Belief?

A **belief** is a statement that is not scientifically provable. Beliefs may or may not be incorrect; they just are outside the realm of science to explore.

Laws vs. Theories

A common misconception is that scientific theories are rudimentary ideas that will eventually graduate into scientific laws when enough data and evidence has accumulated. A theory does not change into a scientific law with the accumulation of new or better evidence. Remember, **theories are explanations** and **laws are patterns** we see in large amounts of data, frequently written as an equation. A theory will always remain a theory; a law will always remain a law.



Video **1.3.1**: What's the difference between a scientific law and theory?

Summary

- A hypothesis is a tentative explanation that can be tested by further investigation.
- A theory is a well-supported explanation of observations.
- A scientific law is a statement that summarizes the relationship between variables.
- An experiment is a controlled method of testing a hypothesis.

Contributions & Attributions

-
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

1.3: Hypothesis, Theories, and Laws is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **1.6: Hypothesis, Theories, and Laws** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.4: TAKING MEASUREMENTS

LEARNING OBJECTIVES

- Express quantities properly, using a number and a unit.

A coffee maker's instructions tell you to fill the coffee pot with 4 cups of water and to use 3 scoops of coffee. When you follow these instructions, you are measuring. When you visit a doctor's office, a nurse checks your temperature, height, weight, and perhaps blood pressure (Figure **1.4.1**); the nurse is also measuring.



Figure **1.4.1**: Measuring Blood Pressure. A nurse or a doctor measuring a patient's blood pressure is taking a measurement. (GFDL; Pia von Lützau).

Chemists measure the properties of matter and express these measurements as quantities. A quantity is an amount of something and consists of a number and a unit. The number tells us how many (or how much), and the unit tells us what the scale of measurement is. For example, when a distance is reported as "5 kilometers," we know that the quantity has been expressed in units of kilometers and that the number of kilometers is 5. If you ask a friend how far they walk from home to school, and the friend answers "12" without specifying a unit, you do not know whether your friend walks 12 kilometers, 12 miles, 12 furlongs, or 12 yards. *Both a number and a unit must be included to express a quantity properly.*

To understand chemistry, we need a clear understanding of the units chemists work with and the rules they follow for expressing numbers. The next two sections examine the rules for expressing numbers.

EXAMPLE 1.4.1

Identify the number and the unit in each quantity.

a. one dozen eggs
b. 2.54 centimeters
c. a box of pencils
d. 88 meters per second

SOLUTION

a. The number is one, and the unit is a dozen eggs.
b. The number is 2.54, and the unit is centimeter.
c. The number 1 is implied because the quantity is only *a* box. The unit is box of pencils.
d. The number is 88, and the unit is meters per second. Note that in this case the unit is actually a combination of two units: meters and seconds.

KEY TAKE AWAY

- Identify a quantity properly with a number and a unit.

1.4: Taking Measurements is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by LibreTexts.

- **2.1: Taking Measurements** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

1.5: The Basic Units of Measurement

Learning Objectives

- State the different measurement systems used in chemistry.
- Describe how prefixes are used in the metric system and identify how the prefixes milli-, centi-, and kilo- compare to the base unit.

How long is a yard? It depends on whom you ask and when you asked the question. Today we have a standard definition of the yard, which you can see marked on every football field. If you move the ball ten yards, you get a first down and it does not matter whether you are playing in Los Angeles, Dallas, or Green Bay. But at one time that yard was arbitrarily defined as the distance from the tip of the king's nose to the end of his outstretched hand. Of course, the problem there is simple: new king, new distance (and then you have to re-mark all of those football fields).




Figure **1.5.1**: Meter standard (left) and Kilogram standard (right).

SI Base Units

All measurements depend on the use of units that are well known and understood. The **English system** of measurement units (inches, feet, ounces, etc.) are not used in science because of the difficulty in converting from one unit to another. The **metric system** is used because all metric units are based on multiples of 10, making conversions very simple. The metric system was originally established in France in 1795. **The International System of Units** is a system of measurement based on the metric system. The acronym **SI** is commonly used to refer to this system and stands for the French term, *Le Système International d'Unités*. The SI was adopted by international agreement in 1960 and is composed of seven base units in Table **1.5.1**.

Table **1.5.1**: SI Base Units of Measurement

Quantity	SI Base Unit	Symbol
Length	meter	**m**
Mass	kilogram	**kg**
Temperature	kelvin	**K**
Time	second	**s**
Amount of a Substance	mole	**mol**
Electric Current	ampere	**A**
Luminous Intensity	candela	**cd**

The first units are frequently encountered in chemistry. All other measurement quantities, such as volume, force, and energy, can be derived from these seven base units.

Unfortunately, the Metric System is Not Ubiquitous

The map below shows the adoption of the SI units in countries around the world. The United States has legally adopted the metric system for measurements, but does not use it in everyday practice. Great Britain and much of Canada use a combination of metric and imperial units.




Figure 1.5.1: Areas of world using metric system (in green). Only a few countries are slow or resistant to adoption including the United States.

Prefix Multipliers

Conversions between metric system units are straightforward because the system is based on powers of ten. For example, meters, centimeters, and millimeters are all metric units of length. There are 10 millimeters in 1 centimeter and 100 centimeters in 1 meter. **Metric prefixes** are used to distinguish between units of different size. These prefixes all derive from either Latin or Greek terms. For example, *mega* comes from the Greek word μεγας, meaning "great". Table 1.5.2 lists the most common metric prefixes and their relationship to the central unit that has no prefix. Length is used as an example to demonstrate the relative size of each prefixed unit.

Table 1.5.2: SI Prefixes

Prefix	Unit Abbreviation	Meaning	Example
giga	G	1,000,000,000	1 gigameter $(\mathrm{Gm}) = 10^9\ \mathrm{m}$
mega	M	1,000,000	1 megameter $(\mathrm{Mm}) = 10^6\ \mathrm{m}$
kilo	k	1,000	1 kilometer $(\mathrm{km}) = 1,000\ \mathrm{m}$
hecto	h	100	1 hectometer $(\mathrm{hm}) = 100\ \mathrm{m}$
deka	da	10	1 dekameter $(\mathrm{dam}) = 10\ \mathrm{m}$
		1	1 meter (m)
deci	d	1/10	1 decimeter $(\mathrm{dm}) = 0.1\ \mathrm{m}$
centi	c	1/100	1 centimeter $(\mathrm{cm}) = 0.01\ \mathrm{m}$
milli	m	1/1,000	1 millimeter $(\mathrm{mm}) = 0.001\ \mathrm{m}$
micro	μ	1/1,000,000	1 micrometer $(\mu\mathrm{m}) = 10^{-6}\ \mathrm{m}$
nano	n	1/1,000,000,000	1 nanometer $(\mathrm{nm}) = 10^{-9}\ \mathrm{m}$
pico	p	1/1,000,000,000,000	1 picometer $(\mathrm{pm}) = 10^{-12}\ \mathrm{m}$

There are a couple of odd little practices with the use of metric abbreviations. Most abbreviations are lowercase. We use "**m**" for meter and not "**M**". However, when it comes to volume, the base unit "liter" is abbreviated as "**L**" and not "**l**". So we would write 3.5 milliliters as **3.5 mL**.

As a practical matter, whenever possible you should express the units in a small and manageable number. If you are measuring the weight of a material that weighs **6.5 kg**, this is easier than saying it weighs **6500 g** or **0.65 dag**. All three are correct, but the **kg** units in this case make for a small and easily managed number. However, if a specific problem needs grams instead of kilograms, go with the grams for consistency.

Example 1.5.1: Unit Abbreviations

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kiloliter
b. microsecond
c. decimeter
d. nanogram

Solutions

Solutions to Example 2.5.1

	Explanation	Answer
a	The prefix kilo means “1,000 ×,” so 1 kL equals 1,000 L.	kL
b	The prefix micro implies 1/1,000,000th of a unit, so 1 µs equals 0.000001 s.	µs
c	The prefix deci means 1/10th, so 1 dm equals 0.1 m.	dm
d	The prefix nano means 1/1000000000, so a nanogram is equal to 0.000000001 g.	ng

Exercise 1.5.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kilometer
b. milligram
c. nanosecond
d. centiliter

Answer a:

km

Answer b:

mg

Answer c:

ns

Answer d:

cL

Summary

- Metric prefixes derive from Latin or Greek terms. The prefixes are used to make the units manageable.
- The SI system is based on multiples of ten. There are seven basic units in the SI system. Five of these units are commonly used in chemistry.

1.5: The Basic Units of Measurement is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.5: The Basic Units of Measurement** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.6: Units Raised to a Power


Learning Objectives

- To convert a value reported in one unit raised to a power of 10, to a corresponding value in a different unit raised to the same power of 10, using conversion factors.

Conversion factors for area and volume can also be produced by the dimensional analysis method. Just remember that if a quantity is raised to a power of 10, both the number and the unit must be raised to the same power of 10. For example, to convert $\mathbf{1500\ cm^2}$ to $\mathbf{m^2}$, we need to start with the relationship between centimeter and meter. We know that 1 cm = 10^{-2} m or 100 cm =1 m, but since we are given the quantity in 1500 cm^2, then we have to use the relationship:

$$1\,cm^2 = (10^{-2}\,m)^2 = 10^{-4}\,m^2$$

CONCEPT MAP



To convert centimeters squared to meters squared, use the conversion factor 0.01 meters per 1 centimeter, squared overall

CALCULATION

$$1500\ \cancel{cm}^2 \times \left(\frac{10^{-2}\,m}{1\ \cancel{cm}}\right)^2 = 0.15\ m^2$$

or

$$1500\ \cancel{cm}^2 \times \left(\frac{1\,m}{100\ \cancel{cm}}\right)^2 = 0.15\ m^2$$

or


$$1500\ \cancel{cm}^2 \times \frac{1\,m^2}{10,000\ \cancel{cm^2}} = 0.15\ m^2$$

Example 1.6.1: Volume of a Sphere

What is the volume of a sphere (radius 4.30 inches) in cubic cm (cm^3)?

Solution

Solution for Example 2.8.1

Steps for Problem Solving	What is the volume of a sphere (radius 4.30 inches) in cubic cm (cm^3)?
Identify the "given" information and what the problem is asking you to "find."	Given: radius = 4.30 in Find: cm^3 (volume)
Determine other known quantities.	Volume of a sphere: V = $\frac{4}{3} \times \pi \times r^3$ = $\frac{4}{3} \times 3.1416 \times (4.30\underline{in})^3$ = $333.04 in^3$
Prepare a concept map.	 $\left(\frac{2.54\ cm}{1\ in}\right)^3$ To convert inches cubed to centimeters cubed, use conversion factor 2.54 centimeters per 1 inch, cubed overall
Calculate.	$333.04\ in^3 \left(\frac{2.54 cm}{1\ in}\right)^3 = 5.46 \times 10^3 cm^3$
Think about your result.	A centimeter is a smaller unit than an inch, so the answer in cubic centimeters is larger than the given value in cubic inches.

? Exercise 1.6.1

Lake Tahoe has a surface area of 191 square miles. What is the area in square km (km^2)?

Answer

495 km^2

1.6: Units Raised to a Power is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.8: Units Raised to a Power** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.7: Scientific Notation - Writing Large and Small Numbers

Learning Objectives

- Express a large number or a small number in scientific notation.
- Carry out arithmetical operations and express the final answer in scientific notation

Chemists often work with numbers that are exceedingly large or small. For example, entering the mass in grams of a hydrogen atom into a calculator would require a display with at least 24 decimal places. A system called **scientific notation** avoids much of the tedium and awkwardness of manipulating numbers with large or small magnitudes. In scientific notation, these numbers are expressed in the form

$$N \times 10^{n}$$

where N is greater than or equal to 1 and less than 10 ($1 \leq N < 10$), and n is a positive or negative integer ($10^0 = 1$). The number 10 is called the base because it is this number that is raised to the power n. Although a base number may have values other than 10, the base number in scientific notation is always 10.

A simple way to convert numbers to scientific notation is to move the decimal point as many places to the left or right as needed to give a number from 1 to 10 (N). The magnitude of n is then determined as follows:

- If the decimal point is moved to the left n places, n is positive.
- If the decimal point is moved to the right n places, n is negative.

Another way to remember this is to recognize that as the number N decreases in magnitude, the exponent increases and vice versa. The application of this rule is illustrated in Example **1.7.1**.

Example 1.7.1: Expressing Numbers in Scientific Notation

Convert each number to scientific notation.

a. 637.8
b. 0.0479
c. 7.86
d. 12,378
e. 0.00032
f. 61.06700
g. 2002.080
h. 0.01020

Solution

Solutions to Example 2.2.1

	Explanation	Answer
a	To convert 637.8 to a number from 1 to 10, we move the decimal point two places to the left: 637.8 Because the decimal point was moved two places to the left, n = 2.	6.378×10^{2}
b	To convert 0.0479 to a number from 1 to 10, we move the decimal point two places to the right: 0.0479 Because the decimal point was moved two places to the right, n = −2.	4.79×10^{-2}
c	This is usually expressed simply as 7.86. (Recall that $10^{0} = 1$.)	7.86×10^{0}
d	Because the decimal point was moved four places to the left, n = 4.	1.2378×10^{4}
e	Because the decimal point was moved four places to the right, n = −4.	3.2×10^{-4}
f	Because the decimal point was moved one place to the left, n = 1.	6.106700×10^{1}
g	Because the decimal point was moved three places to the left, n = 3.	2.002080×10^{3}
h	Because the decimal point was moved two places to the right, n = -2.	1.020×10^{-2}

Addition and Subtraction

Before numbers expressed in scientific notation can be added or subtracted, they must be converted to a form in which all the exponents have the same value. The appropriate operation is then carried out on the values of N. Example **1.7.2** illustrates how to do this.

Example 1.7.2: Expressing Sums and Differences in Scientific Notation

Carry out the appropriate operation and then express the answer in scientific notation.

a. $(1.36 \times 10^{2}) + (4.73 \times 10^{3})$
b. $(6.923 \times 10^{-3}) - (8.756 \times 10^{-4})$

Solution

Solutions to Example 2.2.2.

	Explanation	Answer
a	Both exponents must have the same value, so these numbers are converted to either $(1.36 \times 10^2) + (47.3 \times 10^2) = (1.36 + 47.3) \times 10^2 = 48.66 \times 10^2$ or $(0.136 \times 10^3) + (4.73 \times 10^3) = (0.136 + 4.73) \times 10^3) = 4.87 \times 10^3$. Choosing either alternative gives the same answer, reported to two decimal places. In converting 48.66 × 10^2 to scientific notation, n has become more positive by 1 because the value of N has decreased.	4.87×10^3
b	Converting the exponents to the same value gives either $(6.923 \times 10^{-3}) - (0.8756 \times 10^{-3}) = (6.923 - 0.8756) \times 10^{-3}$ or $(69.23 \times 10^{-4}) - (8.756 \times 10^{-4}) = (69.23 - 8.756) \times 10^{-4} = 60.474 \times 10^{-4}$. In converting 60.474 × 10^{-4} to scientific notation, n has become more positive by 1 because the value of N has decreased.	6.047×10^{-3}

Multiplication and Division

When multiplying numbers expressed in scientific notation, we multiply the values of N and add together the values of n. Conversely, when dividing, we divide N in the dividend (the number being divided) by N in the divisor (the number by which we are dividing) and then subtract n in the divisor from n in the dividend. In contrast to addition and subtraction, the exponents do not have to be the same in multiplication and division. Examples of problems involving multiplication and division are shown in Example **1.7.3**.

Example 1.7.3: Expressing Products and Quotients in Scientific Notation

Perform the appropriate operation and express your answer in scientific notation.

a. $(6.022 \times 10^{23})(6.42 \times 10^{-2})$

b. $\dfrac{1.67 \times 10^{-24}}{9.12 \times 10^{-28}}$

c. $\dfrac{(6.63 \times 10^{-34})(6.0 \times 10)}{8.52 \times 10^{-2}}$

Solution

Solution to Example 2.2.3

	Explanation	
a	In multiplication, we add the exponents: $(6.022 \times 10^{23})(6.42 \times 10^{-2}) = (6.022)(6.42) \times 10^{[23+(-2)]} = 38.7 \times 10^{21}$ In converting 38.7×10^{21} to scientific notation, n has become more positive by 1 because the value of N has decreased.	3.87×10^{22}
b	In division, we subtract the exponents: $\frac{1.67 \times 10^{-24}}{9.12 \times 10^{-28}} = \frac{1.67}{9.12} \times 10^{[-24-(-28)]} = 0.183 \times 10^{4}$ In converting 0.183×10^{4} to scientific notation, n has become more negative by 1 because the value of N has increased.	1.83×10^{3}
c	This problem has both multiplication and division: $\frac{(6.63 \times 10^{-34})(6.0 \times 10)}{(8.52 \times 10^{-2})} = \frac{39.78}{8.52} \times 10^{[-34+1-(-2)]}$	4.7×10^{-31}

1.7: Scientific Notation - Writing Large and Small Numbers is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.2: Scientific Notation - Writing Large and Small Numbers** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.8: Significant Figures - Writing Numbers to Reflect Precision

Learning Objectives

- Identify the number of significant figures in a reported value.

The **significant figures** in a measurement consist of all the certain digits in that measurement plus one uncertain or estimated digit. In the ruler illustration below, the bottom ruler gave a length with 2 significant figures, while the top ruler gave a length with 3 significant figures. In a correctly reported measurement, the final digit is significant but not certain. Insignificant digits are not reported. With either ruler, it would not be possible to report the length at $2.553\ \text{cm}$ as there is no possible way that the thousandths digit could be estimated. The 3 is not significant and would not be reported.




Figure 1.8.1: Measurement with two different rulers. Ruler A's measurement can be rounded to 2.55, with 2 certain digits, while Ruler B's measurement of 2.5 has 1 certain digit

Measurement Uncertainty

Some error or **uncertainty** always exists in any measurement. The amount of uncertainty depends both upon the skill of the measurer and upon the quality of the measuring tool. While some balances are capable of measuring masses only to the nearest $0.1\ \text{g}$, other highly sensitive balances are capable of measuring to the nearest $0.001\ \text{g}$ or even better. Many measuring tools such as rulers and graduated cylinders have small lines which need to be carefully read in order to make a measurement. Figure 1.8.1 shows two rulers making the same measurement of an object (indicated by the blue arrow).


With either ruler, it is clear that the length of the object is between 2 and $3\ \text{cm}$. The bottom ruler contains no millimeter markings. With that ruler, the tenths digit can be estimated and the length may be reported as $2.5\ \text{cm}$. However, another person may judge that the measurement is $2.4\ \text{cm}$ or perhaps $2.6\ \text{cm}$. While the 2 is known for certain, the value of the tenths digit is uncertain.

The top ruler contains marks for tenths of a centimeter (millimeters). Now the same object may be measured as $2.55\ \text{cm}$. The measurer is capable of estimating the hundredths digit because he can be certain that the tenths digit is a 5. Again, another measurer may report the length to be $2.54\ \text{cm}$ or $2.56\ \text{cm}$. In this case, there are two certain digits (the 2 and the 5), with the hundredths digit being uncertain. Clearly, the top ruler is a superior ruler for measuring lengths as precisely as possible.


Example 1.8.1: Reporting Measurements to the Proper Number of Significant Figures

Use each diagram to report a measurement to the proper number of significant figures.

a.



b.



Ruler measuring a rectangle in units of centimeters, with the rectangle's edge between 1.2 and 1.3 cm marks


Solutions

Solutions to Example 2.3.1

	Explanation	Answer
a.	The arrow is between 4.0 and 5.0, so the measurement is at least 4.0. The arrow is between the third and fourth small tick marks, so it's at least 0.3. We will have to estimate the last place. It looks like about one-third of the way across the space, so let us estimate the hundredths place as 3. The symbol psi stands for "pounds per square inch" and is a unit of pressure, like air in a tire. The measurement is reported to three significant figures.	4.33 psi
b.	The rectangle is at least 1.0 cm wide but certainly not 2.0 cm wide, so the first significant digit is 1. The rectangle's width is past the second tick mark but not the third; if each tick mark represents 0.1, then the rectangle is at least 0.2 in the next significant digit. We have to estimate the next place because there are no markings to guide us. It appears to be about halfway between 0.2 and 0.3, so we will estimate the next place to be a 5. Thus, the measured width of the rectangle is 1.25 cm. The measurement is reported to three significant figures.	1.25 cm

? Exercise 1.8.1

What would be the reported width of this rectangle?



Answer

1.25 cm

When you look at a reported measurement, it is necessary to be able to count the number of significant figures. The table below details the rules for determining the number of significant figures in a reported measurement. For the examples in the table, assume that the quantities are correctly reported values of a measured quantity.

Table **1.8.1**: Significant Figure Rules

Rule	Examples
1. All nonzero digits in a measurement are significant.	• 237 has three significant figures. • 1.897 has four significant figures.
2. Zeros that appear between other nonzero digits (middle zeros) are always significant.	• 39,004 has five significant figures. • 5.02 has three significant figures.
3. Zeros that appear in front of all of the nonzero digits are called leading zeros. Leading zeros are never significant.	• 0.008 has one significant figure. • 0.000416 has three significant figures.

4. Zeros that appear after all nonzero digits are called trailing zeros. A number with trailing zeros that lacks a decimal point may or may not be significant. **Use scientific notation to indicate the appropriate number of significant figures.**	• 1400 is ambiguous.
	  ◦ 1.4×10^3 has two significant figures.
	  ◦ 1.40×10^3 three significant figures.
	  ◦ 1.400×10^3 has four significant figures.

5. Trailing zeros in a number with a decimal point are significant. This is true whether the zeros occur before or after the decimal point.	• 620.0 has four significant figures. • 19.000 has five significant figures.

Exact Numbers

Integers obtained either by counting objects or from definitions are exact numbers, which are considered to have infinitely many significant figures. If we have counted four objects, for example, then the number 4 has an infinite number of significant figures (i.e., it represents 4.000…). Similarly, 1 foot (ft) is defined to contain 12 inches (in), so the number 12 in the following equation has infinitely many significant figures:

Example 1.8.2

Give the number of significant figures in each. Identify the rule for each.

a. 5.87
b. 0.031
c. 52.90
d. 00.2001
e. 500
f. 6 atoms

Solution

Solution to Example 2.3.2

	Explanation	Answer
a	All three numbers are significant (rule 1).	5.87, three significant figures
b	The leading zeros are not significant (rule 3). The 3 and the 1 are significant (rule 1).	0.031, two significant figures
c	The 5, the 2 and the 9 are significant (rule 1). The trailing zero is also significant (rule 5).	52.90, four significant figures
d	The leading zeros are not significant (rule 3). The 2 and the 1 are significant (rule 1) and the middle zeros are also significant (rule 2).	00.2001, four significant figures
e	The number is ambiguous. It could have one, two or three significant figures.	500, ambiguous
f	The 6 is a counting number. A counting number is an exact number.	6, infinite

? Exercise 1.8.2

Give the number of significant figures in each.

a. 36.7 m
b. 0.006606 s
c. 2,002 kg
d. 306,490,000 people
e. 3,800 g

Answer a

three significant figures

Answer b

four significant figures

Answer c

four significant figures

Answer d

infinite (exact number)

Answer e

Ambiguous, could be two, three or four significant figures.

Accuracy and Precision


Measurements may be accurate, meaning that the measured value is the same as the true value; they may be precise, meaning that multiple measurements give nearly identical values (i.e., reproducible results); they may be both accurate and precise; or they may be neither accurate nor precise. The goal of scientists is to obtain measured values that are both accurate and precise. The video below demonstrates the concepts of accuracy and precision.



Video 1.8.1: Difference between precision and accuracy.

Example 1.8.3

The following archery targets show marks that represent the results of four sets of measurements.



Which target shows

a. a precise, but inaccurate set of measurements?
b. a set of measurements that is both precise and accurate?
c. a set of measurements that is neither precise nor accurate?

Solution

a. Set **(a)** is precise, but inaccurate.
b. Set **(c)** is both precise and accurate.
c. Set **(d)** is neither precise nor accurate.

Summary

Uncertainty exists in all measurements. The degree of uncertainty is affected in part by the quality of the measuring tool. Significant figures give an indication of the certainty of a measurement. Rules allow decisions to be made about how many digits to use in any given situation.

1.8: Significant Figures - Writing Numbers to Reflect Precision is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew, Henry Agnew, Sridhar Budhi, & Sridhar Budhi.

- **2.3: Significant Figures - Writing Numbers to Reflect Precision** by Henry Agnew, Marisa Alviar-Agnew, Sridhar Budhi is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.9: Significant Figures in Calculations

Learning Objectives

- Use significant figures correctly in arithmetical operations.

Rounding

Before dealing with the specifics of the rules for determining the significant figures in a calculated result, we need to be able to round numbers correctly. To **round** a number, first decide how many significant figures the number should have. Once you know that, round to that many digits, starting from the left. If the number immediately to the right of the last significant digit is less than 5, it is dropped and the value of the last significant digit remains the same. If the number immediately to the right of the last significant digit is greater than or equal to 5, the last significant digit is increased by 1.

Consider the measurement **207.518 m**. Right now, the measurement contains six significant figures. How would we successively round it to fewer and fewer significant figures? Follow the process as outlined in Table **1.9.1**.

Table **1.9.1**: Rounding examples

Number of Significant Figures	Rounded Value	Reasoning
6	207.518	All digits are significant
5	207.52	8 rounds the 1 up to 2
4	207.5	2 is dropped
3	208	5 rounds the 7 up to 8
2	210	8 is replaced by a 0 and rounds the 0 up to 1
1	200	1 is replaced by a 0

Notice that the more rounding that is done, the less reliable the figure is. An approximate value may be sufficient for some purposes, but scientific work requires a much higher level of detail.

It is important to be aware of significant figures when you are mathematically manipulating numbers. For example, dividing 125 by 307 on a calculator gives 0.4071661238… to an infinite number of digits. But do the digits in this answer have any practical meaning, especially when you are starting with numbers that have only three significant figures each? When performing mathematical operations, there are two rules for limiting the number of significant figures in an answer—one rule is for addition and subtraction, and one rule is for multiplication and division.

> *In operations involving significant figures, the answer is reported in such a way that it reflects the reliability of the **least precise** operation. An answer is no more precise than the least precise number used to get the answer.*

Multiplication and Division

For multiplication or division, the rule is to count the number of significant figures in each number being multiplied or divided and then limit the significant figures in the answer to the lowest count. An example is as follows:

$$\underbrace{38.65}_{\text{4 sig figs}} \times \underbrace{105.93}_{\text{5 sig figs}} = \underbrace{4{,}094.1945}_{\text{reduce to 4 sig figs}}$$

The final answer, limited to four significant figures, is 4,094. The first digit dropped is 1, so we do not round up.

Scientific notation provides a way of communicating significant figures without ambiguity. You simply include all the significant figures in the leading number. For example, the number 450 has two significant figures and would be written in scientific notation as 4.5×10^2, whereas 450.0 has four significant figures and would be written as 4.500×10^2. In scientific notation, all significant figures are listed explicitly.

Example 1.9.1

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 23.096×90.300
b. 125×9.000

Solution

a

Explanation	Answer
The calculator answer is 2,085.5688, but we need to round it to five significant figures. Because the first digit to be dropped (in the tenths place) is greater than 5, we round up to 2,085.6.	$\mathbf{2.0856 \times 10^3}$

b

Explanation	Answer
The calculator gives 1,125 as the answer, but we limit it to three significant figures.	$\mathbf{1.13 \times 10^3}$

Addition and Subtraction

How are significant figures handled in calculations? It depends on what type of calculation is being performed. If the calculation is an addition or a subtraction, the rule is as follows: limit the reported answer to the rightmost column that all numbers have significant figures in common. For example, if you were to add 1.2 and 4.71, we note that the first number stops its significant figures in the tenths column, while the second number stops its significant figures in the hundredths column. We therefore limit our answer to the tenths column.

$$\begin{array}{l} 1.2 \\ \underline{4.41} \\ 5.61 \\ \quad \uparrow \text{ limit final answer to the tenths column: } 5.6 \end{array}$$

We drop the last digit—the 1—because it is not significant to the final answer.

The dropping of positions in sums and differences brings up the topic of rounding. Although there are several conventions, in this text we will adopt the following rule: the final answer should be rounded up if the first dropped digit is 5 or greater, and rounded down if the first dropped digit is less than 5.

$$\begin{array}{l} 77.2 \\ \underline{10.46} \\ 87.66 \\ \quad \uparrow \text{ limit final answer to the tenths column and round up: } 87.7 \end{array}$$

Example 1.9.2

a. 13.77 + 908.226
b. 1,027 + 611 + 363.06

Solution

a

Explanation	Answer
The calculator answer is 921.996, but because 13.77 has its farthest-right significant figure in the hundredths place, we need to round the final answer to the hundredths position. Because the first digit to be dropped (in the thousandths place) is greater than 5, we round up to 922.00	$922.00 = 9.2200 \times 10^2$

b

Explanation	Answer
The calculator gives 2,001.06 as the answer, but because 611 and 1027 has its farthest-right significant figure in the ones place, the final answer must be limited to the ones position.	$2,001.06 = 2.001 \times 10^3$

Exercise 1.9.2

Write the answer for each expression using scientific notation with the appropriate number of significant figures.

a. 217 ÷ 903
b. 13.77 + 908.226 + 515
c. 255.0 − 99
d. 0.00666 × 321

Answer a:

$0.240 = 2.40 \times 10^{-1}$

Answer b:

$1,437 = 1.437 \times 10^3$

Answer c:

$156 = 1.56 \times 10^2$

Answer d:

$2.14 = 2.14 \times 10^0$

Remember that calculators do not understand significant figures. *You* are the one who must apply the rules of significant figures to a result from your calculator.

Calculations Involving Multiplication/Division and Addition/Subtraction

In practice, chemists generally work with a calculator and carry all digits forward through subsequent calculations. When working on paper, however, we often want to minimize the number of digits we have to write out. Because successive rounding can compound inaccuracies, intermediate rounding needs to be handled correctly. When working on paper, always round an intermediate result so as to retain at least one more digit than can be justified and carry this number into the next step in the calculation. The final answer is then rounded to the correct number of significant figures at the very end.



Video 1.9.1: Significant figures in mixed operations (https://www.youtube.com/watch?v=yBntMndXQWA).



Video 1.9.2: https://www.youtube.com/watch?v=__csP0NtlGI

In the worked examples in this text, we will often show the results of intermediate steps in a calculation. In doing so, we will show the results to only the correct number of significant figures allowed for that step, in effect treating each step as a separate calculation. This procedure is intended to reinforce the rules for determining the number of significant figures, but in some cases it may give a final answer that differs in the last digit from that obtained using a calculator, where all digits are carried through to the last step.

Example 1.9.3

a. 2(1.008 g) + 15.99 g
b. 137.3 s + 2(35.45 s)
c. $\frac{118.7g}{2} - 35.5g$

Solution

a.

Explanation	Answer
2(1.008 g) + 15.99 g = Perform multiplication first. 2 (1.008 g *4 sig figs*) = **2.016 g** *4 sig figs* The number with the least number of significant figures is 1.008 g; *the number 2 is an exact number and therefore has an infinite number of significant figures*. **Then, perform the addition.** **2.016 g** *thousandths place* + 15.99 g *hundredths place (least precise)* = 18.006 g **Round the final answer.** Round the final answer to the hundredths place since 15.99 has its farthest right significant figure in the hundredths place (least precise).	18.01 g (rounding up)

b.

Explanation	Answer
137.3 s + 2(35.45 s) = **Perform multiplication first.** 2(35.45 s *4 sig figs*) = **70.90 s** *4 sig figs* The number with the least number of significant figures is 35.45; *the number 2 is an exact number and therefore has an infinite number of significant figures.* **Then, perform the addition.** 137.3 s *tenths place (least precise)* + **70.90 s** *hundredths place* = 208.20 s **Round the final answer.** Round the final answer to the tenths place based on 137.3 s.	208.2 s

c.

Explanation	Answer
$\frac{118.7g}{2} - 35.5g =$ **Perform division first.** $\frac{118.7g}{2}$ *4 sig figs* = **59.35 g** *4 sig figs* The number with the least number of significant figures is 118.7 g; *the number 2 is an exact number and therefore has an infinite number of significant figures.* **Perform subtraction next.** **59.35 g** *hundredths place* − 35.5 g *tenths place (least precise)* = 23.85 g **Round the final answer.** Round the final answer to the tenths place based on 35.5 g.	23.9 g (rounding up)

? Exercise 1.9.3

Complete the calculations and report your answers using the correct number of significant figures.

a. 5(1.008s) - 10.66 s
b. 99.0 cm+ 2(5.56 cm)

Answer a

-5.62 s

Answer b

110.2 cm

Summary

- Rounding
  - If the number to be dropped is greater than or equal to 5, increase the number to its left by 1 (e.g. 2.9699 rounded to three significant figures is 2.97).
  - If the number to be dropped is less than 5, there is no change (e.g. 4.00443 rounded to four significant figures is 4.004).
- The rule in multiplication and division is that the final answer should have the same number of significant figures as there are in the number with the fewest significant figures.
- The rule in addition and subtraction is that the answer is given the same number of decimal places as the term with the fewest decimal places.

1.9: Significant Figures in Calculations is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.4: Significant Figures in Calculations** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.10: PROBLEM SOLVING AND UNIT CONVERSIONS

LEARNING OBJECTIVES

- To convert a value reported in one unit to a corresponding value in a different unit using conversion factors.

During your studies of chemistry (and physics also), you will note that mathematical equations are used in many different applications. Many of these equations have a number of different variables with which you will need to work. You should also note that these equations will often require you to use measurements with their units. Algebra skills become very important here!

CONVERTING BETWEEN UNITS WITH CONVERSION FACTORS

A **conversion factor** is a factor used to convert one unit of measurement into another. A simple conversion factor can convert meters into centimeters, or a more complex one can convert miles per hour into meters per second. Since most calculations require measurements to be in certain units, you will find many uses for conversion factors. Always remember that a conversion factor has to represent a fact; this fact can either be simple or more complex. For instance, you already know that 12 eggs equal 1 dozen. A more complex fact is that the speed of light is 1.86×10^5 miles/sec. Either one of these can be used as a conversion factor depending on what type of calculation you are working with (Table 1.10.1).

Table 1.10.1: Conversion Factors from SI units to English Units

English Units	Metric Units	Quantity
1 ounce (oz)	28.35 grams (g)	*mass
1 fluid once (oz)	29.6 mL	volume
2.205 pounds (lb)	1 kilogram (kg)	*mass
1 inch (in)	2.54 centimeters (cm)	length
0.6214 miles (mi)	1 kilometer (km)	length
1 quarter (qt)	0.95 liters (L)	volume

*Pounds and ounces are technically units of force, not mass, but this fact is often ignored by the non-scientific community.

Of course, there are other ratios which are not listed in Table 1.10.1. They may include:

- Ratios embedded in the text of the problem (using words such as *per* or *in each*, or using symbols such as / or %).
- Conversions in the metric system, as covered earlier in this chapter.
- Common knowledge ratios (such as 60 seconds = 1 minute).

If you learned the SI units and prefixes described, then you know that 1 cm is 1/100th of a meter.

$$1\ \text{cm} = \frac{1}{100}\ \text{m} = 10^{-2}\text{m}$$

or

$$100\ \text{cm} = 1\ \text{m}$$

Suppose we divide both sides of the equation by 1m (both the number *and* the unit):

$$\frac{100\ \text{cm}}{1\ \text{m}} = \frac{1\ \text{m}}{1\ \text{m}}$$

As long as we perform the same operation on both sides of the equals sign, the expression remains an equality. Look at the right side of the equation; it now has the same quantity in the numerator (the top) as it has in the denominator (the bottom). Any fraction that has the same quantity in the numerator and the denominator has a value of 1:

$$\frac{100\ \text{cm}}{1\ \text{m}} = \frac{1000\ \text{mm}}{1\ \text{m}} = \frac{1 \times 10^6 \mu\text{m}}{1\ \text{m}} = 1$$

We know that 100 cm *is* 1 m, so we have the same quantity on the top and the bottom of our fraction, although it is expressed in different units.

PERFORMING DIMENSIONAL ANALYSIS

Dimensional analysis is amongst the most valuable tools that physical scientists use. Simply put, it is the conversion between an amount in one unit to the corresponding amount in a desired unit using various conversion factors. This is valuable because certain measurements are more accurate or easier to find than others. The use of units in a calculation to ensure that we obtain the final proper units is called *dimensional analysis*.

Here is a simple example. How many centimeters are there in 3.55 m? Perhaps you can determine the answer in your head. If there are 100 cm in every meter, then 3.55 m equals 355 cm. To solve the problem more formally with a conversion factor, we first write the quantity we are given, 3.55 m. Then we multiply this quantity by a conversion factor, which is the same as multiplying it by 1. We can write 1 as $\frac{100\text{ cm}}{1\text{ m}}$ and multiply:

$$3.55\text{ m} \times \frac{100\text{ cm}}{1\text{ m}}$$

The 3.55 m can be thought of as a fraction with a 1 in the denominator. Because m, the abbreviation for meters, occurs in both the numerator *and* the denominator of our expression, they cancel out:

$$\frac{3.55\ \cancel{\text{m}}}{1} \times \frac{100\text{ cm}}{1\ \cancel{\text{m}}}$$

The final step is to perform the calculation that remains once the units have been canceled:

$$\frac{3.55}{1} \times \frac{100\text{ cm}}{1} = 355\text{ cm}$$

In the final answer, we omit the 1 in the denominator. Thus, by a more formal procedure, we find that 3.55 m equals 355 cm. A generalized description of this process is as follows:

quantity (in old units) × conversion factor = quantity (in new units)

You may be wondering why we use a seemingly complicated procedure for a straightforward conversion. In later studies, the conversion problems you encounter *will not always be so simple*. If you master the technique of applying conversion factors, you will be able to solve a large variety of problems.

In the previous example, we used the fraction $\frac{100\text{ cm}}{1\text{ m}}$ as a conversion factor. Does the conversion factor $\frac{1\text{ m}}{100\text{ cm}}$ also equal 1? Yes, it does; it has the same quantity in the numerator as in the denominator (except that they are expressed in different units). Why did we not use *that* conversion factor? If we had used the second conversion factor, the original unit would not have canceled, and the result would have been meaningless. Here is what we would have gotten:

$$3.55\text{ m} \times \frac{1\text{ m}}{100\text{ cm}} = 0.0355\frac{\text{m}^2}{\text{cm}}$$

For the answer to be meaningful, we have to *construct the conversion factor in a form that causes the original unit to cancel out*. Figure **1.10.1** shows a concept map for constructing a proper conversion.




Figure **1.10.1**: A Concept Map for Conversions. This is how you construct a conversion factor to convert from one unit to another. Converting meters to centimeters: the unit you are converting is on the bottom of the fraction, and the unit to convert to is on top

GENERAL STEPS IN PERFORMING DIMENSIONAL ANALYSIS

1. Identify the "**given**" information in the problem. Look for a number with units to start this problem with.
2. What is the problem asking you to "**find**"? In other words, what unit will your answer have?

3. Use **ratios** and conversion factors to cancel out the units that aren't part of your answer, and leave you with units that are part of your answer.
4. When your units cancel out correctly, you are ready to do the **math**. You are multiplying fractions, so you multiply the top numbers and divide by the bottom numbers in the fractions.



SIGNIFICANT FIGURES IN CONVERSIONS

How do conversion factors affect the determination of significant figures?

- Numbers in conversion factors based on prefix changes, such as kilograms to grams, are *not* considered in the determination of significant figures in a calculation because the numbers in such conversion factors are exact.
- Exact numbers are defined or counted numbers, not measured numbers, and can be considered as having an infinite number of significant figures. (In other words, 1 kg is exactly 1,000 g, by the definition of kilo-.)
- Counted numbers are also exact. If there are 16 students in a classroom, the number 16 is exact.
- In contrast, conversion factors that come from measurements (such as density, as we will see shortly) or that are approximations have a limited number of significant figures and should be considered in determining the significant figures of the final answer.

EXAMPLE 1.10.1

Steps for Problem Solving for Example 2.6.1 and 2.6.2

	Example 1.10.1	**Example 1.10.2**
Steps for Problem Solving	The average volume of blood in an adult male is 4.7 L. What is this volume in milliliters?	A hummingbird can flap its wings once in 18 ms. How many seconds are in 18 ms?
Identify the "given" information and what the problem is asking you to "find."	Given: 4.7 L Find: mL	Given: 18 ms Find: s
List other known quantities.	$1\,mL = 10^{-3}L$	$1\,ms = 10^{-3}s$
Prepare a concept map and use the proper conversion factor.	 $\frac{1mL}{10^{-3}L}$	 $\frac{10^{-3}s}{1ms}$
Cancel units and calculate.	$4.7\,\cancel{L} \times \frac{1\text{ mL}}{10^{-3}\,\cancel{L}} = 4,700\text{ mL}$ or $4.7\,\cancel{L} \times \frac{1,000\text{ mL}}{1\,\cancel{L}} = 4,700\text{ mL}$ or 4.7 x 10^3 2SF, not ambiguous	$18\,\cancel{ms} \times \frac{10^{-3}\text{ s}}{1\,\cancel{ms}} = 0.018\text{ s}$ or $18\,\cancel{ms} \times \frac{1\text{ s}}{1,000\,\cancel{ms}} = 0.018\text{ s}$
Think about your result.	The amount in mL should be 1000 times larger than the given amount in L.	The amount in s should be 1/1000 the given amount in ms.

EXERCISE 1.10.1

Perform each conversion.

a. 101,000 ns to seconds
b. 32.08 kg to grams
c. 1.53 grams to cg

Answer a:

$1.01000x10^{-4}s$

Answer b:

$3.208x10^{4}g$

Answer c:

$1.53x10^{2}cg$

https://chem.libretexts.org/@go/page/492118

SUMMARY

- Conversion factors are used to convert one unit of measurement into another.
- Dimensional analysis (unit conversions) involves the use of conversion factors that will cancel unwanted units and produce the appropriate units.

1.10: Problem Solving and Unit Conversions is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.6: Problem Solving and Unit Conversions** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.


1.11: SOLVING MULTI-STEP CONVERSION PROBLEMS

MULTIPLE CONVERSIONS

Sometimes you will have to perform more than one conversion to obtain the desired unit. For example, suppose you want to convert 54.7 km into millimeters. We will set up a series of conversion factors so that each conversion factor produces the next unit in the sequence. We first convert the given amount in km to the base unit, which is meters. We know that 1,000 m =1 km.

Then we convert meters to mm, remembering that $1\ \text{mm} = 10^{-3}\ \text{m}$.

CONCEPT MAP



Convert kilometers to meters to millimeters: use conversion factors 1000 meters per 1 kilometer and 1 millimeter per 0.001 meter

CALCULATION


$$54.7\ \cancel{\text{km}} \times \frac{1{,}000\ \cancel{\text{m}}}{1\ \cancel{\text{km}}} \times \frac{1\ \text{mm}}{10^{-3}\ \cancel{\text{m}}} = 54{,}700{,}000\ \text{mm}$$
$$= 5.47 \times 10^{7}\ \text{mm}$$

> *In each step, the previous unit is canceled and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.*

✔ EXAMPLE 1.11.1: UNIT CONVERSION

Convert 58.2 ms to megaseconds in one multi-step calculation.


SOLUTION

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 58.2 ms Find: Ms
List other known quantities	$1ms = 10^{-3}s$ $1Ms = 10^{6}s$
Prepare a concept map.	 Convert milliseconds to seconds to microseconds: use conversion factors 0.001 second per millisecond and 1 microsecond per 1 million seconds
Calculate.	$58.2\ \cancel{\text{ms}} \times \frac{10^{-3}\ \cancel{s}}{1\ \cancel{\text{ms}}} \times \frac{1\ \text{Ms}}{1{,}000{,}000\ \cancel{s}} = 0.0000000582\ \text{Ms}$ $= 5.82 \times 10^{-8}\ \text{Ms}$ Neither conversion factor affects the number of significant figures in the final answer.

EXAMPLE 1.11.2: UNIT CONVERSION

How many seconds are in a day?

Solution

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 1 day Find: s
List other known quantities.	1 day = 24 hours 1 hour = 60 minutes 1 minute = 60 seconds
Prepare a concept map.	 $\frac{24\text{hr}}{1\text{ day}}$ $\frac{60\text{min}}{1\text{ hr}}$ $\frac{60\text{ s}}{1\ min}$ Convert day to hour to minute to second: use conversion factors 24 hours per day, 60 minutes per hour, and 60 seconds per minute
Calculate.	$1\text{ d} \times \frac{24\text{ hr}}{1\text{ d}} \times \frac{60\text{ min}}{1\text{ hr}} \times \frac{60\text{ s}}{1\text{ min}} = 86,400\text{ s}$

EXERCISE 1.11.1

Perform each conversion in one multi-step calculation.

a. 43.007 ng to kg
b. 1005 in to ft
c. 12 mi to km

Answer a

$4.3007 \times 10^{-11} kg$

Answer b

$83.75\ ft$

Answer c

$19\ km$

CAREER FOCUS: PHARMACIST

A pharmacist dispenses drugs that have been prescribed by a doctor. Although that may sound straightforward, pharmacists in the United States must hold a doctorate in pharmacy and be licensed by the state in which they work. Most pharmacy programs require four years of education in a specialty pharmacy school. Pharmacists must know a lot of chemistry and biology so they can understand the effects that drugs (which are chemicals, after all) have on the body. Pharmacists can advise physicians on the selection, dosage, interactions, and side effects of drugs. They can also advise patients on the proper use of their medications, including when and how to take specific drugs properly. Pharmacists can be found in drugstores, hospitals, and other medical facilities. Curiously, an outdated name for pharmacist is *chemist*, which was used when pharmacists formerly did a lot of drug preparation, or *compounding*. In modern times, pharmacists rarely compound their own drugs, but their knowledge of the sciences, including chemistry, helps them provide valuable services in support of everyone's health.



A woman consulting with a pharmacist. (Public Domain; Rhoda Baer via National Cancer Institute, an agency that is part of the National Institutes of Health.)

SUMMARY

In multi-step conversion problems, the previous unit is canceled for each step and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.

1.11: Solving Multi-step Conversion Problems is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.7: Solving Multi-step Conversion Problems** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

1.12: Density

Learning Objectives

- Define density.
- Use density as a conversion factor.

Density (ρ) is a physical property found by dividing the mass of an object by its volume. Regardless of the sample size, density is always constant. For example, the density of a pure sample of tungsten is always 19.25 grams per cubic centimeter. This means that whether you have one gram or one kilogram of the sample, the density will never vary. The equation, as we already know, is as follows:

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

or just

$$\rho = \frac{m}{V} \tag{1.12.1}$$

Based on this equation, it's clear that density can, and does, vary from element to element and substance to substance due to differences in the relationship of mass and volume. Pure water, for example, has a density of 0.998 g/cm^3 at 25° C. The average densities of some common substances are in Table **1.12.1**. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will "float."

Table **1.12.1**: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Density can be measured for all substances—solids, liquids and gases. For solids and liquids, density is often reported using the units of g/cm^3. Densities of gases, which are significantly lower than the densities of solids and liquids, are often given using units of g/L.

Example 1.12.1: Ethyl Alcohol

Calculate the density of a 30.2 mL sample of ethyl alcohol with a mass of 23.71002 g

Solution

This is a direct application of Equation 1.12.1:

$$\rho = \frac{23.71002\,g}{30.2\,mL} = 0.785\,g/mL$$

? Exercise 1.12.1

a. Find the density (in kg/L) of a sample that has a volume of 36.5 L and a mass of 10.0 kg.
b. If you have a 2.130 mL sample of acetic acid with mass 0.002234 kg, what is the density in kg/L?

Answer a

$0.274\ kg/L$

Answer b

$1.049\ kg/L$

Density as a Conversion Factor

Conversion factors can also be constructed for converting between different kinds of units. For example, density can be used to convert between the mass and the volume of a substance. Consider mercury, which is a liquid at room temperature and has a density of 13.6 g/mL. The density tells us that 13.6 g of mercury have a volume of 1 mL. We can write that relationship as follows:

13.6 g mercury = 1 mL mercury

This relationship can be used to construct two conversion factors:

$$\frac{13.6\ g}{1\ mL} = 1$$

and

$$\frac{1\ mL}{13.6\ g} = 1$$

Which one do we use? It depends, as usual, on the units we need to cancel and introduce. For example, suppose we want to know the mass of 2.0 mL of mercury. We would use the conversion factor that has milliliters on the bottom (so that the milliliter unit cancels) and grams on top, so that our final answer has a unit of mass:

$$2.0\ \cancel{mL} \times \frac{13.6\ g}{1\ \cancel{mL}} = 27.2\ g = 27\ g$$

In the last step, we limit our final answer to two significant figures because the volume quantity has only two significant figures; the 1 in the volume unit is considered an exact number, so it does not affect the number of significant figures. The other conversion factor would be useful if we were given a mass and asked to find volume, as the following example illustrates.


> *Density can be used as a conversion factor between mass and volume.*

✓ Example 1.12.2: Mercury Thermometer Steps for Problem Solving

A mercury thermometer for measuring a patient's temperature contains 0.750 g of mercury. What is the volume of this mass of mercury?

Solution

Solution to Example 2.9.2

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 0.750 g Find: mL
List other known quantities.	13.6 g/mL (density of mercury)
Prepare a concept map.	
Calculate.	$0.750\ \cancel{g} \times \frac{1\ \text{mL}}{13.6\ \cancel{g}} = 0.055147...\ \text{mL} \approx 0.0551\ \text{mL}$ We have limited the final answer to three significant figures.

? Exercise 1.12.2

What is the volume of 100.0 g of air if its density is 1.3 g/L?

Answer

$77\ L$

Summary

- Density is defined as the mass of an object divided by its volume.
- Density can be used as a conversion factor between mass and volume.

1.12: Density is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.9: Density** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.13: Temperature - Random Motion of Molecules and Atoms

Learning Objectives

- Identify the different between temperature and heat.
- Recognize the different scales used to measure temperature

The concept of temperature may seem familiar to you, but many people confuse temperature with heat. **Temperature** is a measure of how hot or cold an object is relative to another object (its thermal energy content), whereas **heat** is the flow of thermal energy between objects with different temperatures. Temperature is a measure of the average kinetic energy of the particles in matter. In everyday usage, temperature indicates a measure of how hot or cold an object is. Temperature is an important parameter in chemistry. When a substance changes from solid to liquid, it is because there was in increase in the temperature of the material. Chemical reactions usually proceed faster if the temperature is increased. Many unstable materials (such as enzymes) will be viable longer at lower temperatures.



Figure **1.13.1**: The glowing charcoal on the left represents high kinetic energy, while the snow and ice on the right are of much lower kinetic energy.

Three different scales are commonly used to measure temperature: Fahrenheit (expressed as °F), Celsius (°C), and Kelvin (K). Thermometers measure temperature by using materials that expand or contract when heated or cooled. Mercury or alcohol thermometers, for example, have a reservoir of liquid that expands when heated and contracts when cooled, so the liquid column lengthens or shortens as the temperature of the liquid changes.



Figure **1.13.2**: Daniel Gabriel Fahrenheit (left), Anders Celsius (center), and Lord Kelvin (right).

The Fahrenheit Scale

The first thermometers were glass and contained alcohol, which expanded and contracted as the temperature changed. The German scientist, Daniel Gabriel Fahrenheit used mercury in the tube, an idea put forth by Ismael Boulliau. The Fahrenheit scale was first developed in 1724 and tinkered with for some time after that. The main problem with this scale is the arbitrary definitions of temperature. The freezing point of water was defined as $32°F$ and the boiling point as $212°F$. The Fahrenheit scale is typically not used for scientific purposes.

The Celsius Scale

The Celsius scale of the metric system is named after Swedish astronomer Anders Celsius (1701-1744). The Celsius scale sets the freezing point and boiling point of water at $0°C$ and $100°C$ respectively. The distance between those two points is divided into 100 equal intervals, each of which is one degree. Another term sometimes used for the Celsius scale is "centigrade" because there are 100 degrees between the freezing and boiling points of water on this scale. However, the preferred term is "Celsius".

The Kelvin Scale

The Kelvin temperature scale is named after Scottish physicist and mathematician Lord Kelvin (1824-1907). It is based on molecular motion, with the temperature of $\mathbf{0\ K}$, also known as absolute zero, being the point where all molecular motion ceases. The freezing point of water on the Kelvin scale is $\mathbf{273.15\ K}$, while the boiling point is $\mathbf{373.15\ K}$. Notice that there is no "degree" used in the temperature designation. Unlike the Fahrenheit and Celsius scales where temperatures are referred to as "degrees **F**" or "degrees **C**", we simply designate temperatures in the Kelvin scale as kelvins.




Figure **1.13.1**: A Comparison of the Fahrenheit, Celsius, and Kelvin Temperature Scales. Because the difference between the freezing point of water and the boiling point of water is 100° on both the Celsius and Kelvin scales, the size of a degree Celsius (°C) and a kelvin (K) are precisely the same. In contrast, both a degree Celsius and a kelvin are 9/5 the size of a degree Fahrenheit (°F). (CC BY-SA-NC 3.0; anonymous)

Converting Between Scales

The Kelvin is the same size as the Celsius degree, so measurements are easily converted from one to the other. The freezing point of water is 0°C = 273.15 K; the boiling point of water is 100°C = 373.15 K. The Kelvin and Celsius scales are related as follows:

$$T\,(\text{in °C}) + 273.15 = T\,(\text{in K}) \tag{3.10.1}$$

$$T\ \ (\text{in K}) - 273.15 = T\ (\text{in °C}) \tag{3.10.2}$$

Degrees on the Fahrenheit scale, however, are based on an English tradition of using 12 divisions, just as 1 ft = 12 in. The relationship between degrees Fahrenheit and degrees Celsius is as follows: where the coefficient for degrees Fahrenheit is exact. (Some calculators have a function that allows you to convert directly between °F and °C.) There is only one temperature for which the numerical value is the same on both the Fahrenheit and Celsius scales: −40°C = −40°F. The relationship between the scales is as follows:

$$°C = \frac{(°F - 32)}{1.8} \tag{3.10.3}$$

$$°F = 1.8 \times (°C) + 32 \tag{3.10.4}$$

Example 1.13.1: Temperature Conversions

A student is ill with a temperature of 103.5°F. What is her temperature in °C and K?

Solution

Converting from Fahrenheit to Celsius requires the use of Equation 3.10.3:

$$°C = \frac{(103.5°F - 32)}{1.8} \tag{1.13.1}$$

$$= 39.7\,°C \tag{1.13.2}$$

Converting from Celsius to Kelvin requires the use of Equation 3.10.1:

$$K = 39.7\,°C + 273.15 \quad (1.13.3)$$
$$= 312.9\,K \quad (1.13.4)$$

? Exercise 1.13.1

Convert each temperature to °C and °F.

a. the temperature of the surface of the sun (5800 K)
b. the boiling point of gold (3080 K)
c. the boiling point of liquid nitrogen (77.36 K)

Answer (a)

5527 K, 9980 °F

Answer (b)

2807 K, 5084 °F

Answer (c)

-195.79 K, -320.42 °F

Summary

Three different scales are commonly used to measure temperature: Fahrenheit (expressed as °F), Celsius (°C), and Kelvin (K).

Contributions & Attributions

1.13: Temperature - Random Motion of Molecules and Atoms is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.10: Temperature - Random Motion of Molecules and Atoms** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

1.E: Measurement and Problem Solving (Exercises)

2.1: Measuring Global Temperatures

2.2: Scientific Notation: Writing Large and Small Numbers

2.3: Significant Figures: Writing Numbers to Reflect Precision

1. Define *significant figures*. Why are they important?
2. Define the different types of zeros found in a number and explain whether or not they are significant.
3. How many significant figures are in each number?
    a. 140
    b. 0.009830
    c. 15,050
    d. 221,560,000
    e. 5.67×10^3
    f. 2.9600×10^{-5}
4. How many significant figures are in each number?
    a. 1.05
    b. 9,500
    c. 0.0004505
    d. 0.00045050
    e. 7.210×10^6
    f. 5.00×10^{-6}
5. Round each number to three significant figures.
    a. 34,705
    b. 34,750
    c. 34,570

2.4: Significant Figures in Calculations

2.5: The Basic Units of Measurement

2.6: Problem Solving and Unit Conversions

2.7: Solving Multi-step Conversion Problems

2.8: Units Raised to a Power

2.9: Density

2.10: Numerical Problem-Solving Strategies and the Solution Map

1.E: Measurement and Problem Solving (Exercises) is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **2.E: Measurement and Problem Solving (Exercises)** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

2: Unit 2

2.1: In Your Room
2.2: What is Matter?
2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas
2.4: Classifying Matter According to Its Composition
2.5: Differences in Matter- Physical and Chemical Properties
2.6: Changes in Matter - Physical and Chemical Changes
2.7: Conservation of Mass - There is No New Matter
2.8: Cutting Aluminum until you get Atoms
2.9: Indivisible - The Atomic Theory
2.10: The Nuclear Atom
2.11: The Properties of Protons, Neutrons, and Electrons
2.12: Elements- Defined by Their Number of Protons
2.13: Looking for Patterns - The Periodic Table
2.14: Ions - Losing and Gaining Electrons
2.15: Isotopes - When the Number of Neutrons Varies
2.16: Atomic Mass - The Average Mass of an Element’s Atoms
2.17: Sugar and Salt
2.18: Compounds Display Constant Composition
2.19: Chemical Formulas - How to Represent Compounds
2.20: A Molecular View of Elements and Compounds
2.21: Writing Formulas for Ionic Compounds
2.22: Nomenclature- Naming Compounds
2.23: Naming Ionic Compounds
2.24: Naming Molecular Compounds
2.25: Naming Acids
2.26: Nomenclature Summary

2: Unit 2 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

2.1: In Your Room

Matter is any substance that has mass and takes up space. Matter includes atoms and anything made up of atoms, but not other energy phenomena or waves such as light or sound. While this simple definition is easily applied, the way people view matter is often broken down into two characteristic length scales: the macroscopic and the microscopic.



Figure **2.1.1**: A typical American university and college dormitory room in 2002 (CC BY-SA 3.0; Raul654).

The macroscopic scale is the length scale on which objects or phenomena are large enough to be visible almost practically with the naked eye, without magnifying optical instruments. Everything that one can see, touch, and handle in the dorm room of Figure **2.1.1** is within the macroscopic scale. To describe each of these objects, only a few macroscopic properties are required. However, each of these items can be decomposed into smaller microscopic scale properties.

The microscopic scale is the scale of objects and events smaller than those that can easily be seen by the naked eye, requiring a lens or microscope to see them clearly. All of the everyday objects that we can bump into, touch, or squeeze are ultimately composed of atoms. This ordinary atomic matter is in turn made up of interacting subatomic particles—usually a nucleus of protons and neutrons, and a cloud of orbiting electrons. Because of this, a large number of variables are needed to describe such a system which complicates the characterization.

Matter vs. Mass

Matter should not be confused with mass, as the two are not the same in modern physics. Matter is a physical substance of which systems may be composed, while mass is not a substance, but rather a quantitative property of matter and other substances or systems.

Contributions & Attributions

- Wikipedia

2.1: In Your Room is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.1: In Your Room** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.2: What is Matter?

Learning Objectives

- Define matter and explain how it is composed of building blocks known as "atoms".

We are all familiar with matter. The definition of **Matter** is anything that has mass and volume (takes up space). For most common objects that we deal with every day, it is fairly simple to demonstrate that they have mass and take up space. You might be able to imagine, however, the difficulty for people several hundred years ago to demonstrate that air had mass and volume. Air (and all other gases) are invisible to the eye, have very small masses compared to equal amounts of solids and liquids, and are quite easy to compress (change volume). Without sensitive equipment, it would have been difficult to convince people that gases are matter. Today, we can measure the mass of a small balloon when it is deflated and then blow it up, tie it off, and measure its mass again to detect the additional mass due to the air inside. The mass of air, under room conditions, that occupies a one quart jar is approximately 0.0002 pounds. This small amount of mass would have been difficult to measure in times before balances were designed to accurately measure very small masses. Later, scientists were able to compress gases into such a small volume that the gases turned into liquids, which made it clear that gases are matter.



Figure **2.2.1**: Everything from an ant, to a truck, to the earth, and even the entire galaxy is composed of matter. Images used with permission from Wikipedia (CC_SA-BY-3.0; credit High Contrast).

Even though the universe consists of "things" as wildly different as ants and galaxies, the matter that makes up all of these "things" is composed of a very limited number of building blocks. These building blocks are known as **atoms**, and so far, scientists have discovered or created a grand total of 118 different types of atoms. Scientists have given a name to each different type of atom. A substance that is composed of only one type of atom is called an element. At this point, what should amaze you is that all forms of matter in our universe are made with only 118 different building blocks. In some ways, it's sort of like cooking a gourmet, five-course meal using only three ingredients! How is it possible? To answer that question, you have to understand the ways in which different elements are put together to form matter.

The most important method that nature uses to organize atoms into matter is the formation of **molecules**. Molecules are groups of two or more atoms that have been bonded together. There are millions of different ways to bond atoms together, which means that

there are millions of different possible molecules. Each of these molecules has its own set of chemical properties, and it's these properties with which chemists are most concerned. You will learn a lot more about atoms and molecules, including how they were discovered, in a later part of the textbook.

Summary

All matter has mass and occupies space. All physical objects are made of matter. Matter itself is composed of tiny building blocks known as "atoms". There are only 118 different types of atoms known to man. Frequently, atoms are bonded together to form "molecules".

Contributions & Attributions

- Wikipedia

2.2: What is Matter? is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.2: What is Matter?** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas

Learning Objectives

- To describe the solid, liquid and gas phases.

Water can take many forms. At low temperatures (below **0°C**), it is a solid. When at "normal" temperatures (between **0°C** and **100°C**), it is a liquid. While at temperatures above **100°C**, water is a gas (steam). The state that water is in depends upon the temperature. Each state has its own unique set of physical properties. Matter typically exists in one of three states: **solid**, **liquid**, or **gas**.




Figure **2.3.1**: Matter is usually classified into three classical states, with plasma sometimes added as a fourth state. From left to right: quartz (solid), water (liquid), nitrogen dioxide (gas).

The state that a given substance exhibits is also a physical property. Some substances exist as gases at room temperature (oxygen and carbon dioxide), while others, like water and mercury metal, exist as liquids. Most metals exist as solids at room temperature. All substances can exist in any of these three states. Figure **2.3.2** shows the differences among solids, liquids, and gases at the molecular level. A solid has definite volume and shape, a liquid has a definite volume but no definite shape, and a gas has neither a definite volume nor shape.





Figure **2.3.2**: A Representation of the Solid, Liquid, and Gas States. (a) Solid O_2 has a fixed volume and shape, and the molecules are packed tightly together. (b) Liquid O_2 conforms to the shape of its container but has a fixed volume; it contains relatively densely packed molecules. (c) Gaseous O_2 fills its container completely—regardless of the container's size or shape—and consists of widely separated molecules.

Plasma: A Fourth State of Matter

Technically speaking, a fourth state of matter called plasma exists, but it does not naturally occur on earth, so we will omit it from our study here.



A plasma globe operating in a darkened room. (CC BY-SA 3.0; Chocolateoak).

Solids

In the solid state, the individual particles of a substance are in fixed positions with respect to each other because there is not enough thermal energy to overcome the intermolecular interactions between the particles. As a result, solids have a definite shape and volume. Most solids are hard, but some (like waxes) are relatively soft. Many solids composed of ions can also be quite brittle.

Solids are defined by the following characteristics:

- Definite shape (rigid)
- Definite volume
- Particles vibrate around fixed axes

If we were to cool liquid mercury to its freezing point of $-39°C$, and under the right pressure conditions, we would notice all of the liquid particles would go into the solid state. Mercury can be solidified when its temperature is brought to its freezing point. However, when returned to room temperature conditions, mercury does not exist in solid state for long, and returns back to its more common liquid form.

Solids usually have their constituent particles arranged in a regular, three-dimensional array of alternating positive and negative ions called a **crystal**. The effect of this regular arrangement of particles is sometimes visible macroscopically, as shown in Figure **2.3.3**. Some solids, especially those composed of large molecules, cannot easily organize their particles in such regular crystals and exist as amorphous (literally, “without form”) solids. Glass is one example of an amorphous solid.




Figure **2.3.3**: (left) The periodic crystalline lattice structure of quartz SiO_2 in two-dimensions. (right) The random network structure of glassy SiO_2 in two-dimensions. Note that, as in the crystal, each Silicon atom is bonded to 4 oxygen atoms, where the fourth oxygen atom is obscured from view in this plane. Images used with permission (public domain).

Liquids

If the particles of a substance have enough energy to partially overcome intermolecular interactions, then the particles can move about each other while remaining in contact. This describes the liquid state. In a liquid, the particles are still in close contact, so liquids have a definite volume. However, because the particles can move about each other rather freely, a liquid has no definite shape and takes a shape dictated by its container.

Liquids have the following characteristics:

- No definite shape (takes the shape of its container).
- Has definite volume.
- Particles are free to move over each other, but are still attracted to each other.

A familiar liquid is mercury metal. Mercury is an anomaly. It is the only metal we know of that is liquid at room temperature. Mercury also has an ability to stick to itself (surface tension)—a property that all liquids exhibit. Mercury has a relatively high surface tension, which makes it very unique. Here you see mercury in its common liquid form.



***Video* 2.3.1**: *Mercury boiling to become a gas.*

If we heat liquid mercury to its boiling point of $357^{\circ}C$ under the right pressure conditions, we would notice all particles in the liquid state go into the gas state.

Gases

If the particles of a substance have enough energy to completely overcome intermolecular interactions, then the particles can separate from each other and move about randomly in space. This describes the gas state, which we will consider in more detail elsewhere. Like liquids, gases have no definite shape, but unlike solids and liquids, gases have no definite volume either. The change from solid to liquid usually does not significantly change the volume of a substance. However, the change from a liquid to a gas significantly increases the volume of a substance, by a factor of 1,000 or more. Gases have the following characteristics:

- No definite shape (takes the shape of its container)
- No definite volume
- Particles move in random motion with little or no attraction to each other
- Highly compressible

Table **2.3.1**: Characteristics of the Three States of Matter

Characteristics	Solids	Liquids	Gases
shape	definite	indefinite	indefinite
volume	definite	definite	indefinite
relative intermolecular interaction strength	strong	moderate	weak
relative particle positions	in contact and fixed in place	in contact but not fixed	not in contact, random positions

Example 2.3.1

What state or states of matter does each statement, describe?

a. This state has a definite volume, but no definite shape.
b. This state has no definite volume.
c. This state allows the individual particles to move about while remaining in contact.

Solution

a. This statement describes the liquid state.
b. This statement describes the gas state.
c. This statement describes the liquid state.

? Exercise 2.3.1

What state or states of matter does each statement describe?

a. This state has individual particles in a fixed position with regard to each other.
b. This state has individual particles far apart from each other in space.
c. This state has a definite shape.

Answer a:

solid

Answer b:

gas

Answer c:

solid

Summary

- Three states of matter exist—solid, liquid, and gas.
- Solids have a definite shape and volume.
- Liquids have a definite volume, but take the shape of the container.
- Gases have no definite shape or volume.

2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.3: Classifying Matter According to Its State—Solid, Liquid, and Gas** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.4: Classifying Matter According to Its Composition

Learning Objectives

- Explain the difference between a pure substance and a mixture.
- Explain the difference between an element and a compound.
- Explain the difference between a homogeneous mixture and a heterogeneous mixture.

One useful way of organizing our understanding of matter is to think of a hierarchy that extends down from the most general and complex to the simplest and most fundamental (Figure **2.4.1**). Matter can be classified into two broad categories: pure substances and mixtures. A **pure substance** is a form of matter that has a constant composition (meaning that it is the same everywhere) and properties that are constant throughout the sample (meaning that there is only one set of properties such as melting point, color, boiling point, etc. throughout the matter). A material composed of two or more substances is a **mixture**. Elements and compounds are both examples of pure substances. A substance that cannot be broken down into chemically simpler components is an **element**. Aluminum, which is used in soda cans, is an element. A substance that can be broken down into chemically simpler components (because it has more than one element) is a **compound**. For example, water is a compound composed of the elements hydrogen and oxygen. Today, there are about 118 elements in the known universe. In contrast, scientists have identified tens of millions of different compounds to date.




Figure **2.4.1**: Relationships between the Types of Matter and the Methods Used to Separate Mixtures

Ordinary table salt is called sodium chloride. It is considered a **substance** because it has a uniform and definite composition. All samples of sodium chloride are chemically identical. Water is also a pure substance. Salt easily dissolves in water, but salt water cannot be classified as a substance because its composition can vary. You may dissolve a small amount of salt or a large amount into a given amount of water. A mixture is a physical blend of two or more components, each of which retains its own identity and

properties in the **mixture**. Only the form of the salt is changed when it is dissolved into water. It retains its composition and properties.

A **homogeneous** mixture is a mixture in which the composition is uniform throughout the mixture. The salt water described above is homogeneous because the dissolved salt is evenly distributed throughout the entire salt water sample. Often it is easy to confuse a homogeneous mixture with a pure substance because they are both uniform. The difference is that the composition of the substance is always the same. The amount of salt in the salt water can vary from one sample to another. All solutions are considered homogeneous because the dissolved material is present in the same amount throughout the solution.

A **heterogeneous mixture** is a mixture in which the composition is not uniform throughout the mixture. Vegetable soup is a heterogeneous mixture. Any given spoonful of soup will contain varying amounts of the different vegetables and other components of the soup.

Phase

A phase is any part of a sample that has a uniform composition and properties. By definition, a pure substance or a homogeneous mixture consists of a single phase. A heterogeneous mixture consists of two or more phases. When oil and water are combined, they do not mix evenly, but instead form two separate layers. Each of the layers is called a phase.

Example 2.4.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. filtered tea
b. freshly squeezed orange juice
c. a compact disc
d. aluminum oxide, a white powder that contains a 2:3 ratio of aluminum and oxygen atoms
e. selenium

Given: a chemical substance

Asked for: its classification

Strategy:

A. Decide whether a substance is chemically pure. If it is pure, the substance is either an element or a compound. If a substance can be separated into its elements, it is a compound.
B. If a substance is not chemically pure, it is either a heterogeneous mixture or a homogeneous mixture. If its composition is uniform throughout, it is a homogeneous mixture.

Solution

a. **A)** Tea is a solution of compounds in water, so it is not chemically pure. It is usually separated from tea leaves by filtration.
B) Because the composition of the solution is uniform throughout, it is ***a homogeneous mixture***.
b. **A)** Orange juice contains particles of solid (pulp) as well as liquid; it is not chemically pure.
B) Because its composition is not uniform throughout, orange juice is ***a heterogeneous mixture***.
c. **A)** A compact disc is a solid material that contains more than one element, with regions of different compositions visible along its edge. Hence, a compact disc is not chemically pure.
B) The regions of different composition indicate that a compact disc is ***a heterogeneous mixture.***
d. **A)** Aluminum oxide is a single, chemically ***pure compound***.
e. **A)** Selenium is one of the known ***elements***.

Exercise 2.4.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. white wine
b. mercury
c. ranch-style salad dressing
d. table sugar (sucrose)

Answer a:
homogeneous mixture (solution)

Answer b:
element

Answer c:
heterogeneous mixture

Answer d:
compound

Example 2.4.2

How would a chemist categorize each example of matter?

a. saltwater
b. soil
c. water
d. oxygen

Solution

a. Saltwater acts as if it were a single substance even though it contains two substances—salt and water. Saltwater is a homogeneous mixture, or a solution.
b. Soil is composed of small pieces of a variety of materials, so it is a heterogeneous mixture.
c. Water is a substance. More specifically, because water is composed of hydrogen and oxygen, it is a compound.
d. Oxygen, a substance, is an element.

Exercise 2.4.2

How would a chemist categorize each example of matter?

a. coffee
b. hydrogen
c. an egg

Answer a:
a homogeneous mixture (solution), assuming it is filtered coffee

Answer b:
element

Answer c:
heterogeneous mixture

Summary

Matter can be classified into two broad categories: pure substances and mixtures. A pure substance is a form of matter that has a constant composition and properties that are constant throughout the sample. Mixtures are physical combinations of two or more

elements and/or compounds. Mixtures can be classified as homogeneous or heterogeneous. Elements and compounds are both examples of pure substances. Compounds are substances that are made up of more than one type of atom. Elements are the simplest substances made up of only one type of atom.

Vocabulary

- Element: a substance that is made up of only one type of atom.
- Compound:a substance that is made up of more than one type of atom bonded together.
- Mixture: a combination of two or more elements or compounds which have not reacted to bond together; each part in the mixture retains its own properties.

Contributions & Attributions

-
- Stephen Lower, Professor Emeritus (Simon Fraser U.) Chem1 Virtual Textbook

---

2.4: Classifying Matter According to Its Composition is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew, Henry Agnew, Stephen Lower, & Stephen Lower.

- **3.4: Classifying Matter According to Its Composition** by Henry Agnew, Marisa Alviar-Agnew, Stephen Lower is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.5: Differences in Matter- Physical and Chemical Properties

Learning Objectives

To separate physical from chemical properties.

All matter has physical and chemical properties. Physical properties are characteristics that scientists can measure without changing the composition of the sample under study, such as mass, color, and volume (the amount of space occupied by a sample). Chemical properties describe the characteristic ability of a substance to react to form new substances; they include its flammability and susceptibility to corrosion. All samples of a pure substance have the same chemical and physical properties. For example, pure copper is always a reddish-brown solid (a physical property) and always dissolves in dilute nitric acid to produce a blue solution and a brown gas (a chemical property).

Physical Property

A *physical property* is a characteristic of a substance that can be observed or measured without changing the identity of the substance. Silver is a shiny metal that conducts electricity very well. It can be molded into thin sheets, a property called malleability. Salt is dull and brittle and conducts electricity when it has been dissolved into water, which it does quite easily. Physical properties of matter include color, hardness, malleability, solubility, electrical conductivity, density, melting point, and boiling point.

For the elements, color does not vary much from one element to the next. The vast majority of elements are colorless, silver, or gray. Some elements do have distinctive colors: sulfur and chlorine are yellow, copper is (of course) copper-colored, and elemental bromine is red. However, density can be a very useful parameter for identifying an element. Of the materials that exist as solids at room temperature, iodine has a very low density compared to zinc, chromium, and tin. Gold has a very high density, as does platinum. Pure water, for example, has a density of 0.998 g/cm^3 at 25°C. The average densities of some common substances are in Table **2.5.1**. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will "float."

Table **2.5.1**: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Hardness helps determine how an element (especially a metal) might be used. Many elements are fairly soft (silver and gold, for example) while others (such as titanium, tungsten, and chromium) are much harder. Carbon is an interesting example of hardness. In graphite, (the "lead" found in pencils) the carbon is very soft, while the carbon in a diamond is roughly seven times as hard.



Figure **2.5.1**: Pencil (left) and Diamond ring (right). Both are a form of carbon, but exhibit very different physical properties.

Melting and boiling points are somewhat unique identifiers, especially of compounds. In addition to giving some idea as to the identity of the compound, important information can be obtained about the purity of the material.

Chemical Properties

Chemical properties of matter describe its potential to undergo some chemical change or reaction by virtue of its composition. The elements, electrons, and bonds that are present give the matter potential for chemical change. It is quite difficult to define a chemical property without using the word "change". Eventually, after studying chemistry for some time, you should be able to look at the formula of a compound and state some chemical property. For example, hydrogen has the potential to ignite and explode given the right conditions—this is a chemical property. Metals in general have the chemical property of reacting with an acid. Zinc reacts with hydrochloric acid to produce hydrogen gas—this is a chemical property.



Figure **2.5.2**: Heavy rust on the links of a chain near the Golden Gate Bridge in San Francisco; it was continuously exposed to moisture and salt spray, causing surface breakdown, cracking, and flaking of the metal. (CC BY-SA 3.0; Marlith).

A chemical property of iron is its capability of combining with oxygen to form iron oxide, the chemical name of rust (Figure **2.5.2**). The more general term for rusting and other similar processes is corrosion. Other terms that are commonly used in descriptions of chemical changes are burn, rot, explode, decompose, and ferment. Chemical properties are very useful in identifying substances. However, unlike physical properties, chemical properties can only be observed as the substance is in the process of being changed into a different substance.

Table **2.5.2**: Contrasting Physical and Chemical Properties

Physical Properties	Chemical Properties
Gallium metal melts at 30 ^{o}C.	Iron metal rusts.
Mercury is a very dense liquid.	A green banana turns yellow when it ripens.
Gold is shiny.	A dry piece of paper burns.

Example 2.5.1

Which of the following is a chemical property of iron?

a. Iron corrodes in moist air.
b. Density = 7.874 g/cm^3
c. Iron is soft when pure.
d. Iron melts at 1808 K.

Solution

"Iron corrodes in moist air" is the only chemical property of iron from the list.

Exercise 2.5.1A

Which of the following is a physical property of matter?

a. corrosiveness
b. pH (acidity)
c. density
d. flammability

Answer

c

Exercise 2.5.1B

Which of the following is a chemical property?

a. flammability
b. melting point
c. boiling point
d. density

Answer

a

Summary

A physical property is a characteristic of a substance that can be observed or measured without changing the identity of the substance. Physical properties include color, density, hardness, and melting and boiling points. A chemical property describes the ability of a substance to undergo a specific chemical change. To identify a chemical property, we look for a chemical change. A chemical change always produces one or more types of matter that differ from the matter present before the change. The formation of rust is a chemical change because rust is a different kind of matter than the iron, oxygen, and water present before the rust formed.

2.5: Differences in Matter- Physical and Chemical Properties is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.5: Differences in Matter- Physical and Chemical Properties** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.6: Changes in Matter - Physical and Chemical Changes

Learning Objectives

- Label a change as chemical or physical.
- List evidence that can indicate a chemical change occurred.

Change is happening all around us all of the time. Just as chemists have classified elements and compounds, they have also classified types of changes. Changes are classified as either physical or chemical changes. Chemists learn a lot about the nature of matter by studying the changes that matter can undergo. Chemists make a distinction between two different types of changes that they study—physical changes and chemical changes.

Physical Change

Physical changes are changes in which no bonds are broken or formed. This means that the same types of compounds or elements that were there at the beginning of the change are there at the end of the change. Because the ending materials are the same as the beginning materials, the properties (such as color, boiling point, etc.) will also be the same. Physical changes involve moving molecules around, but not changing them. Some types of physical changes include:

- Changes of state (changes from a solid to a liquid or a gas and vice versa).
- Separation of a mixture.
- Physical deformation (cutting, denting, stretching).
- Making solutions (special kinds of mixtures).

As an ice cube melts, its shape changes as it acquires the ability to flow. However, its composition does not change. Melting is an example of a physical change. A physical change is a change to a sample of matter in which some properties of the material change, but the identity of the matter does not. When liquid water is heated, it changes to water vapor. However, even though the physical properties have changed, the molecules are exactly the same as before. We still have each water molecule containing two hydrogen atoms and one oxygen atom covalently bonded. When you have a jar containing a mixture of pennies and nickels and you sort the mixture so that you have one pile of pennies and another pile of nickels, you have not altered the identity of the pennies or the nickels—you've merely separated them into two groups. This would be an example of a physical change. Similarly, if you have a piece of paper, you don't change it into something other than a piece of paper by ripping it up. What was paper before you started tearing is still paper when you are done. Again, this is an example of a physical change.



Figure **2.6.1**: Ice melting is a physical change. When liquid water (H_2O) freezes into a solid state (ice), it appears changed; however, this change is only physical, as the composition of the constituent molecules is the same: 11.19% hydrogen and 88.81% oxygen by mass. (Public Domain; Moussa).

Physical changes can further be classified as reversible or irreversible. The melted ice cube may be refrozen, so melting is a reversible physical change. Physical changes that involve a change of state are all reversible. Other changes of state include **vaporization** (liquid to gas), **freezing** (liquid to solid), and **condensation** (gas to liquid). Dissolving is also a reversible physical change. When salt is dissolved into water, the salt is said to have entered the aqueous state. The salt may be regained by boiling off the water, leaving the salt behind.

Chemical Change

Chemical changes occur when bonds are broken and/or formed between molecules or atoms. This means that one substance with a certain set of properties (such as melting point, color, taste, etc) is turned into a different substance with different properties. Chemical changes are frequently harder to reverse than physical changes.

One good example of a chemical change is burning a candle. The act of burning paper actually results in the formation of new chemicals (carbon dioxide and water) from the burning of the wax. Another example of a chemical change is what occurs when natural gas is burned in your furnace. This time, on the left there is a molecule of methane, $\mathrm{CH_4}$, and two molecules of oxygen, $\mathrm{O_2}$; on the right are two molecules of water, $\mathrm{H_2O}$, and one molecule of carbon dioxide, $\mathrm{CO_2}$. In this case, not only has the appearance changed, but the structure of the molecules has also changed. The new substances do not have the same chemical properties as the original ones. Therefore, this is a chemical change.



Figure **2.6.2**: Burning of wax to generate water and carbon dioxide is a chemical reaction. (CC-SA-BY-3.0; Andrikkos)

We can't actually see molecules breaking and forming bonds, although that's what defines chemical changes. We have to make other observations to indicate that a chemical change has happened. Some of the evidence for chemical change will involve the energy changes that occur in chemical changes, but some evidence involves the fact that new substances with different properties are formed in a chemical change.

Observations that help to indicate chemical change include:

- Temperature changes (either the temperature increases or decreases).
- Light given off.
- Unexpected color changes (a substance with a different color is made, rather than just mixing the original colors together).
- Bubbles are formed (but the substance is not boiling—you made a substance that is a gas at the temperature of the beginning materials, instead of a liquid).
- Different smell or taste (do not taste your chemistry experiments, though!).
- A solid forms if two clear liquids are mixed (look for *floaties*—technically called a precipitate).

Example 2.6.1

Label each of the following changes as a physical or chemical change. Give evidence to support your answer.

a. Boiling water.
b. A nail rusting.
c. A green solution and colorless solution are mixed. The resulting mixture is a solution with a pale green color.
d. Two colorless solutions are mixed. The resulting mixture has a yellow precipitate.

Solution

a. Physical: boiling and melting are physical changes. When water boils, no bonds are broken or formed. The change could be written: $\mathrm{H_2O}\,(l) \rightarrow \mathrm{H_2O}\,(g)$

b. Chemical: The dark grey nail changes color to form an orange flaky substance (the rust); this must be a chemical change. Color changes indicate chemical change. The following reaction occurs: $Fe + O_2 \rightarrow Fe_2O_3$
c. Physical: because none of the properties changed, this is a physical change. The green mixture is still green and the colorless solution is still colorless. They have just been spread together. No color *change* occurred or other evidence of chemical change.
d. Chemical: the formation of a precipitate and the color change from colorless to yellow indicate a chemical change.

? Exercise 2.6.1

Label each of the following changes as a physical or chemical change.

a. A mirror is broken.
b. An iron nail corroded in moist air
c. Copper metal is melted.
d. A catalytic converter changes nitrogen dioxide to nitrogen gas and oxygen gas.

Answer a:

physical change

Answer b:

chemical change

Answer c:

physical change

Answer d:

chemical change

Separating Mixtures Through Physical Changes

Homogeneous mixtures (solutions) can be separated into their component substances by physical processes that rely on differences in some physical property, such as differences in their boiling points. Two of these separation methods are distillation and crystallization. Distillation makes use of differences in volatility, a measure of how easily a substance is converted to a gas at a given temperature. A simple distillation apparatus for separating a mixture of substances, at least one of which is a liquid. The most volatile component boils first and is condensed back to a liquid in the water-cooled condenser, from which it flows into the receiving flask. If a solution of salt and water is distilled, for example, the more volatile component, pure water, collects in the receiving flask, while the salt remains in the distillation flask.




Figure **2.6.3***: The Distillation of a Solution of Table Salt in Water. The solution of salt in water is heated in the distilling flask until it boils. The resulting vapor is enriched in the more volatile component (water), which condenses to a liquid in the cold condenser and is then collected in the receiving flask.*

Parts of a distillation setup: Bunsen burner, salt water in distilling flask, condenser with cool water in and warm water out, pure water in receiving flask

Mixtures of two or more liquids with different boiling points can be separated with a more complex distillation apparatus. One example is the refining of crude petroleum into a range of useful products: aviation fuel, gasoline, kerosene, diesel fuel, and lubricating oil (in the approximate order of decreasing volatility). Another example is the distillation of alcoholic spirits such as brandy or whiskey. This relatively simple procedure caused more than a few headaches for federal authorities in the 1920s during the era of Prohibition, when illegal stills proliferated in remote regions of the United States.

Another example for using physical properties to separate mixtures is filtration (Figure **2.6.4**). Filtration is any mechanical, physical or biological operation that separates solids from fluids (liquids or gases) by adding a medium through which only the fluid can pass. The fluid that passes through is called the filtrate. There are many different methods of filtration; all aim to attain the separation of substances. Separation is achieved by some form of interaction between the substance or objects to be removed and the filter. The substance that is to pass through the filter must be a fluid, i.e. a liquid or gas. Methods of filtration vary depending on the location of the targeted material, i.e. whether it is dissolved in the fluid phase or suspended as a solid.




Figure **2.6.4***: Filtration for the separation of solids from a hot solution. (CC BY-SA 4.0; Suman6395).*

Summary

- Chemists make a distinction between two different types of changes that they study—physical changes and chemical changes.
- Physical changes are changes that do not alter the identity of a substance.
- Chemical changes are changes that occur when one substance is turned into another substance.
- Chemical changes are frequently harder to reverse than physical changes. Observations that indicate a chemical change has occurred include color change, temperature change, light given off, formation of bubbles, formation of a precipitate, etc.

Contributions & Attributions

-
- Boundless (www.boundless.com)

2.6: Changes in Matter - Physical and Chemical Changes is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.6: Changes in Matter - Physical and Chemical Changes** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.7: Conservation of Mass - There is No New Matter

It may seem as though burning destroys matter, but the same amount, or mass, of matter still exists after a campfire as before. Look at Figure **2.7.1** below. It shows that when wood burns, it combines with oxygen and changes not only to ashes, but also to carbon dioxide and water vapor. The gases float off into the air, leaving behind just the ashes. Suppose you had measured the mass of the wood before it burned and the mass of the ashes after it burned. Also suppose you had been able to measure the oxygen used by the fire and the gases produced by the fire. What would you find? The total mass of matter after the fire would be the same as the total mass of matter before the fire.




Figure **2.7.1**: Burning is a chemical process. The flames are caused as a result of a fuel undergoing combustion (burning). (CC BY-SA 2.5; Einar Helland Berger for fire and Walter Siegmund for ash).

Law of Conservation of Mass

The law of conservation of mass was created in 1789 by a French chemist, Antoine Lavoisier. The **law of conservation of mass** states that matter cannot be created or destroyed in a chemical reaction. For example, when wood burns, the mass of the soot, ashes, and gases equals the original mass of the charcoal and the oxygen when it first reacted. So the mass of the product equals the mass of the reactant. A reactant is the chemical reaction of two or more elements to make a new substance, and a product is the substance that is formed as the result of a chemical reaction (Video **2.7.1**). Matter and its corresponding mass may not be able to be created or destroyed, but can change forms to other substances like liquids, gases, and solids.



Video **2.7.1**: This is a nice little demonstration showing the Conservation of Mass in action.

If you witness a 300 kg tree burn to the ground, there are only ashes left after the burn, and all of them together weigh 10 kg. It may make you wonder where the other 290 kg went. The missing 290 kg was released into the atmosphere as smoke, so the only thing left that you can see is the 10 kg of ash. If you know the law of conservation of mass, then you know that the other 290 kg has to go somewhere, because it has to equal the mass of the tree before it burnt down.

Example 2.7.1

If heating 10.0 grams of calcium carbonate ($CaCO_3$) produces 4.4 g of carbon dioxide (CO_2) and 5.6 g of calcium oxide (CaO), show that these observations are in agreement with the law of conservation of mass.

Solution

$$\text{Mass of the reactants} = \text{Mass of the products}$$
$$10.0\,\text{g of } CaCO_3 = 4.4\,\text{g of } CO_2 + 5.6\,\text{g of CaO}$$
$$10.0\,\text{g of reactant} = 10.0\,\text{g of products}$$

Because the mass of the reactant is equal to the mass of the products, the observations are in agreement with the law of conservation of mass.

Exercise 2.7.1

Potassium hydroxide (**KOH**) readily reacts with carbon dioxide (CO_2) to produce potassium carbonate (K_2CO_3) and water (H_2O). How many grams of potassium carbonate are produced if 224.4 g of **KOH** reacts with 88.0 g of CO_2? The reaction also produces 36.0 g of water.

Answer

276.4 g of potassium carbonate

The Law is also applicable to both chemical and physical changes. For example, if you have an ice cube that melts into a liquid and you heat that liquid up, it becomes a gas. It will appear to have disappeared, but is still there.

Summary

- Burning and other changes in matter do not destroy matter.
- The mass of matter is always the same before and after the changes occur.
- The law of conservation of mass states that matter cannot be created or destroyed.

Contributions & Attributions

- Binod Shrestha (University of Lorraine)

2.7: Conservation of Mass - There is No New Matter is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **3.7: Conservation of Mass - There is No New Matter** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.8: Cutting Aluminum until you get Atoms

Take some aluminum foil. Cut it in half. Now there are two smaller pieces of aluminum foil. Cut one of the pieces in half again. Cut one of those smaller pieces in half again. Continue cutting, making smaller and smaller pieces of aluminum foil. It should be obvious that the pieces are still aluminum foil; they are just becoming smaller and smaller. But how far can this exercise be taken, at least in theory? Can one continue cutting the aluminum foil into halves forever, making smaller and smaller pieces? Or is there some limit, some absolute smallest piece of aluminum foil? Thought experiments like this—and the conclusions based on them—were debated as far back as the fifth century BC by Democritus and other ancient Greek philosophers (Figure **2.8.1**).




Figure **2.8.1**: Democritus argued that matter, like an aluminum block, cannot be repeatedly cut in half perpetually. At some point, a limit is reached before the substance can no longer be called "aluminum"; this is the atomic limit. (CC BY-NC; Ümit Kaya via LibreTexts)

Most elements in their pure form exist as individual atoms. For example, a macroscopic chunk of iron metal is composed, microscopically, of individual atoms. Some elements, however, exist as groups of atoms called molecules. Several important elements exist as two-atom combinations and are called diatomic molecules. In representing a diatomic molecule, we use the symbol of the element and include the subscript 2 to indicate that two atoms of that element are joined together. The elements that exist as diatomic molecules are hydrogen (H_2), oxygen (O_2), nitrogen (N_2), fluorine (F_2), chlorine (Cl_2), bromine (Br_2), and iodine (I_2).

2.8: Cutting Aluminum until you get Atoms is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.1: Cutting Aluminum until you get Atoms** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.9: Indivisible - The Atomic Theory

Learning Objectives

- Give a short history of the concept of the atom.
- Describe the contributions of Democritus and Dalton to atomic theory.
- Summarize Dalton's atomic theory and explain its historical development.

You learned earlier that all matter in the universe is made out of tiny building blocks called atoms. All modern scientists accept the concept of the atom, but when the concept of the atom was first proposed about 2,500 years ago, ancient philosophers laughed at the idea. It has always been difficult to convince people of the existence of things that are too small to see. We will spend some time considering the evidence (observations) that convince scientists of the existence of atoms.

Democritus and the Greek Philosophers

About 2,500 years ago, early Greek philosophers believed the entire universe was a single, huge, entity. In other words, "everything was one." They believed that all objects, all matter, and all substances were connected as a single, big, unchangeable "thing." One of the first people to propose "atoms" was a man known as Democritus. As an alternative to the beliefs of the Greek philosophers, he suggested that **atomos**, or atomon—tiny, indivisible, solid objects—make up all matter in the universe.



Figure **2.9.1**: (left) Democritus by Hendrick ter Brugghen, 1628. Democritus was known as the "laughing philosopher." It was a good thing he liked to laugh, because most other philosophers were laughing at his theories. (right) British physicist and chemist John Dalton (1766-1844). Unlike the Greek philosophers, John Dalton believed in both logical thinking and experimentation.

Democritus then reasoned that changes occur when the many atomos in an object were reconnected or recombined in different ways. Democritus even extended this theory, suggesting that there were different varieties of atomos with different shapes, sizes, and masses. He thought, however, that shape, size, and mass were the only properties differentiating the different types of atomos. According to Democritus, other characteristics, like color and taste, did not reflect properties of the atomos themselves, but rather, resulted from the different ways in which the atomos were combined and connected to one another.

The early Greek philosophers tried to understand the nature of the world through reason and logic, but not through experiment and observation. As a result, they had some very interesting ideas, but they felt no need to justify their ideas based on life experiences. In a lot of ways, you can think of the Greek philosophers as being "all thought and no action." It's truly amazing how much they achieved using their minds, but because they never performed any experiments, they missed or rejected a lot of discoveries that they could have made otherwise. Greek philosophers dismissed Democritus' theory entirely. Sadly, it took over two millennia before the theory of atomos (or "atoms," as they are known today) was fully appreciated.

Greeks: "All Thought and No Action"

Greek philosophers were "all thought and no action" and did not feel the need to test their theories with reality. In contrast, Dalton's efforts were based on experimentation and testing ideas against reality.

While it must be assumed that many more scientists, philosophers, and others studied composition of matter after Democritus, a major leap forward in our understanding of the composition of matter took place in the 1800's with the work of the British

scientists John Dalton. He started teaching school at age twelve, and was primarily known as a teacher. In his twenties, he moved to the growing city of Manchester, where he was able to pursue some scientific studies. His work in several areas of science brought him a number of honors. When he died, over 40,000 people in Manchester marched at his funeral.


The modern atomic theory, proposed about 1803 by the English chemist John Dalton (Figure **2.9.1**), is a fundamental concept that states that all elements are composed of atoms. Previously, we defined an atom as the smallest part of an element that maintains the identity of that element. Individual atoms are extremely small; even the largest atom has an approximate diameter of only 5.4×10^{-10} m. With that size, it takes over 18 million of these atoms, lined up side by side, to equal the width of your little finger (about 1 cm).

Dalton studied the weights of various elements and compounds. He noticed that matter always combined in fixed ratios based on weight, or volume in the case of gases. Chemical compounds always contain the same proportion of elements by mass, regardless of amount, which provided further support for Proust's law of definite proportions. Dalton also observed that there could be more than one combination of two elements.

From his experiments and observations, as well as the work from peers of his time, Dalton proposed a new theory of the **atom**. This later became known as Dalton's **atomic theory**. The general tenets of this theory were as follows:

- All matter is composed of extremely small particles called atoms.
- Atoms of a given element are identical in size, mass, and other properties. Atoms of different elements differ in size, mass, and other properties.
- Atoms cannot be subdivided, created, or destroyed.
- Atoms of different elements can combine in simple whole number ratios to form chemical compounds.
- In chemical reactions, atoms are combined, separated, or rearranged.

Dalton's atomic theory has been largely accepted by the scientific community, with the exception of three changes. We know now that (1) an atom can be further subdivided, (2) all atoms of an element are not identical in mass, and (3) using nuclear fission and fusion techniques, we can create or destroy atoms by changing them into other atoms.


ELEMENTS. IV.

EXPLANATION OF THE PLATES. 219

PLATE IV. This plate contains the arbitrary marks or signs chosen to represent the several chemical elements or ultimate particles.

Figure **2.9.2**: Dalton's symbols from his text "A New System of Chemical Philosophy."

The evidence for atoms is so great that few doubt their existence. In fact, individual atoms are now routinely observed with state-of-the art technologies. Moreover, they can even be used for making pretty images; or as IBM research demonstrates in Video **2.9.1**, control of individual atoms can be use used create animations.



Video **2.9.1**: A Boy And His Atom: The World's Smallest Movie.

A Boy and His Atom is a 2012 stop-motion animated short film released by IBM Research. The movie tells the story of a boy and a wayward atom who meet and become friends. It depicts a boy playing with an atom that takes various forms. It was made by moving carbon monoxide molecules viewed with a scanning tunneling microscope, a device that magnifies them 100 million times. These molecules were moved to create images, which were then saved as individual frames to make the film.

Summary

- 2,500 years ago, Democritus suggested that all matter in the universe was made up of tiny, indivisible, solid objects he called "atomos." However, other Greek philosophers disliked Democritus' "atomos" theory because they felt it was illogical.
- Dalton's Atomic Theory is the first scientific theory to relate chemical changes to the structure, properties, and behavior of the atom. The general tenets of this theory are:
  - All matter is composed of extremely small particles called atoms.
  - Atoms of a given element are identical in size, mass, and other properties. Atoms of different elements differ in size, mass, and other properties.
  - Atoms cannot be subdivided, created, or destroyed.
  - Atoms of different elements can combine in simple whole number ratios to form chemical compounds.
  - In chemical reactions, atoms are combined, separated, or rearranged.

2.9: Indivisible - The Atomic Theory is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.2: Indivisible - The Atomic Theory** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.10: The Nuclear Atom

Learning Objectives

- Explain the observations that led to Thomson's discovery of the electron.
- Describe Thomson's "plum pudding" model of the atom and the evidence for it.
- Draw a diagram of Thomson's "plum pudding" model of the atom and explain why it has this name.
- Describe Rutherford's gold foil experiment and explain how this experiment altered the "plum pudding" model.
- Draw a diagram of the Rutherford model of the atom and label the nucleus and the electron cloud.

Dalton's Atomic Theory held up well to a lot of the different chemical experiments that scientists performed to test it. In fact, for almost 100 years, it seemed as if Dalton's Atomic Theory was the whole truth. However, in 1897, a scientist named J. J. Thomson conducted some research that suggested that Dalton's Atomic Theory was not the entire story. He suggested that the small, negatively charged particles making up the cathode ray were actually pieces of atoms. He called these pieces "corpuscles," although today we know them as **electrons**. Thanks to his clever experiments and careful reasoning, J. J. Thomson is credited with the discovery of the electron.



Figure **2.10.1**: J. J. Thomson (center) concluded experiments that suggested that Dalton's (left) atomic theory was not telling the entire story. Ernest Rutherford (right) later built on Thomson's work to argue for the model nuclear atom.

Electrons and Plums

The electron was discovered by J. J. Thomson in 1897. The existence of protons was also known, as was the fact that atoms were neutral in charge. Since the intact atom had no net charge and the electron and proton had opposite charges, the next step after the discovery of subatomic particles was to figure out how these particles were arranged in the atom. This was a difficult task because of the incredibly small size of the atom. Therefore, scientists set out to design a model of what they believed the atom could look like. The goal of each **atomic model** was to accurately represent all of the experimental evidence about atoms in the simplest way possible.

Following the discovery of the electron, J.J. Thomson developed what became known as the "**plum pudding**" model in 1904. Plum pudding is an English dessert similar to a blueberry muffin. In Thomson's plum pudding model of the atom, the electrons were embedded in a uniform sphere of positive charge like blueberries stuck into a muffin. The positive matter was thought to be jelly-like or similar to a thick soup. The electrons were somewhat mobile. As they got closer to the outer portion of the atom, the positive charge in the region was greater than the neighboring negative charges, and the electron would be pulled back more toward the center region of the atom.




Figure **2.10.2**: The "plum pudding" model.

However, this model of the atom soon gave way to a new model developed by New Zealander Ernest Rutherford (1871-1937) about five years later. Thomson did still receive many honors during his lifetime, including being awarded the Nobel Prize in Physics in 1906 and a knighthood in 1908.

Atoms and Gold

In 1911, Rutherford and coworkers Hans Geiger and Ernest Marsden initiated a series of groundbreaking experiments that would completely change the accepted model of the atom. They bombarded very thin sheets of gold foil with fast moving **alpha particles**. Alpha particles, a type of natural radioactive particle, are positively charged particles with a mass about four times that of a hydrogen atom.




Figure **2.10.3**: (A) The experimental setup for Rutherford's gold foil experiment: A radioactive element that emitted alpha particles was directed toward a thin sheet of gold foil that was surrounded by a screen which would allow detection of the deflected particles. (B) According to the plum pudding model (top), all of the alpha particles should have passed through the gold foil with little or no deflection. Rutherford found that a small percentage of alpha particles were deflected at large angles, which could be explained by an atom with a very small, dense, positively-charged nucleus at its center (bottom).

According to the accepted atomic model, in which an atom's mass and charge are uniformly distributed throughout the atom, the scientists expected that all of the alpha particles would pass through the gold foil with only a slight deflection or none at all. Surprisingly, while most of the alpha particles were indeed not deflected, a very small percentage (about 1 in 8000 particles) bounced off the gold foil at very large angles. Some were even redirected back toward the source. No prior knowledge had prepared them for this discovery. In a famous quote, Rutherford exclaimed that it was "as if you had fired a 15-inch [artillery] shell at a piece of tissue and it came back and hit you."

Rutherford needed to come up with an entirely new model of the atom in order to explain his results. Because the vast majority of the alpha particles had passed through the gold, he reasoned that most of the atom was empty space. In contrast, the particles that were highly deflected must have experienced a tremendously powerful force within the atom. He concluded that all of the positive charge and the majority of the mass of the atom must be concentrated in a very small space in the atom's interior, which he called the nucleus. The **nucleus** is the tiny, dense, central core of the atom and is composed of protons and neutrons.

Rutherford's atomic model became known as the **nuclear model**. In the nuclear atom, the protons and neutrons, which comprise nearly all of the mass of the atom, are located in the nucleus at the center of the atom. The electrons are distributed around the nucleus and occupy most of the volume of the atom. It is worth emphasizing just how small the nucleus is compared to the rest of the atom. If we could blow up an atom to be the size of a large professional football stadium, the nucleus would be about the size of a marble.

Rutherford's model proved to be an important step towards a full understanding of the atom. However, it did not completely address the nature of the electrons and the way in which they occupy the vast space around the nucleus. It was not until some years later that a full understanding of the electron was achieved. This proved to be the key to understanding the chemical properties of elements.

Atomic Nucleus

The **nucleus** (plural, nuclei) is a positively charged region at the center of the atom. It consists of two types of subatomic particles packed tightly together. The particles are protons, which have a positive electric charge, and neutrons, which are neutral in electric charge. Outside of the nucleus, an atom is mostly empty space, with orbiting negative particles called electrons whizzing through it. The figure below shows these parts of the atom.




Figure **2.10.4**: The nuclear atom.

The nucleus of the atom is extremely small. Its radius is only about 1/100,000 of the total radius of the atom. Electrons have virtually no mass, but protons and neutrons have a lot of mass for their size. As a result, the nucleus has virtually all the mass of an atom. Given its great mass and tiny size, the nucleus is very dense. If an object the size of a penny had the same density as the nucleus of an atom, its mass would be greater than 30 million tons!

Holding it all Together

Particles with opposite electric charges attract each other. This explains why negative electrons orbit the positive nucleus. Particles with the same electric charge repel each other. This means that the positive protons in the nucleus push apart from one another. So why doesn't the nucleus fly apart? An even stronger force—called the **strong nuclear force**—holds protons and neutrons together in the nucleus.

Summary

- Atoms are the ultimate building blocks of all matter.
- The modern atomic theory establishes the concepts of atoms and how they compose matter.
- Bombardment of gold foil with alpha particles showed that some particles were deflected.
- The nuclear model of the atom consists of a small and dense positively charged interior surrounded by a cloud of electrons.

This page titled 2.10: The Nuclear Atom is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew via source content that was edited to the style and standards of the LibreTexts platform.

- **4.3: The Nuclear Atom** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.11: The Properties of Protons, Neutrons, and Electrons

Learning Objectives

- Describe the locations, charges, and masses of the three main subatomic particles.
- Determine the number of protons and electrons in an atom.
- Define atomic mass unit (amu).

Dalton's Atomic Theory explained a lot about matter, chemicals, and chemical reactions. Nevertheless, it was not entirely accurate, because contrary to what Dalton believed, atoms can, in fact, be broken apart into smaller subunits or subatomic particles. We have been talking about the electron in great detail, but there are two other particles of interest to us: protons and neutrons. We already learned that J. J. Thomson discovered a negatively charged particle, called the **electron**. Rutherford proposed that these electrons orbit a positive nucleus. In subsequent experiments, he found that there is a smaller positively charged particle in the nucleus, called a **proton**. There is also a third subatomic particle, known as a neutron.

Electrons

Electrons are one of three main types of particles that make up atoms. Unlike protons and neutrons, which consist of smaller, simpler particles, electrons are fundamental particles that do not consist of smaller particles. They are a type of fundamental particle called leptons. All leptons have an electric charge of -1 or 0. Electrons are extremely small. The mass of an electron is only about 1/2000 the mass of a proton or neutron, so electrons contribute virtually nothing to the total mass of an atom. Electrons have an electric charge of -1, which is equal but opposite to the charge of a proton, which is $+1$. All atoms have the same number of electrons as protons, so the positive and negative charges "cancel out", making atoms electrically neutral.

Unlike protons and neutrons, which are located inside the nucleus at the center of the atom, electrons are found outside the nucleus. Because opposite electric charges attract one another, negative electrons are attracted to the positive nucleus. This force of attraction keeps electrons constantly moving through the otherwise empty space around the nucleus. The figure below is a common way to represent the structure of an atom. It shows the electron as a particle orbiting the nucleus, similar to the way that planets orbit the sun. However, this is an incorrect perspective, as quantum mechanics demonstrates that electrons are more complicated.



Figure **2.11.1**: Electrons are much smaller than protons or neutrons. If an electron was the mass of a penny, a proton or a neutron would have the mass of a large bowling ball!

Protons

A proton is one of three main particles that make up the atom. Protons are found in the nucleus of the atom. This is a tiny, dense region at the center of the atom. Protons have a positive electrical charge of one $(+1)$ and a mass of 1 atomic mass unit (amu), which is about 1.67×10^{-27} kilograms. Together with neutrons, they make up virtually all of the mass of an atom.

Neutrons

Atoms of all elements—except for most atoms of hydrogen—have neutrons in their nucleus. Unlike protons and electrons, which are electrically charged, neutrons have no charge—they are electrically neutral. That's why the neutrons in the diagram above are labeled n^0. The zero stands for "zero charge". The mass of a neutron is slightly greater than the mass of a proton, which is 1 atomic

mass unit (amu). (An atomic mass unit equals about 1.67×10^{-27} kilograms.) A neutron also has about the same diameter as a proton, or 1.7×10^{-15} meters.

As you might have already guessed from its name, the neutron is neutral. In other words, it has no charge whatsoever and is therefore neither attracted to nor repelled from other objects. Neutrons are in every atom (with one exception), and they are bound together with other neutrons and protons in the atomic nucleus.

Before we move on, we must discuss how the different types of subatomic particles interact with each other. When it comes to neutrons, the answer is obvious. Since neutrons are neither attracted to nor repelled from objects, they don't really interact with protons or electrons (beyond being bound into the nucleus with the protons).

Even though electrons, protons, and neutrons are all types of subatomic particles, they are not all the same size. When you compare the masses of electrons, protons, and neutrons, what you find is that electrons have an extremely small mass, compared to either protons or neutrons. On the other hand, the masses of protons and neutrons are fairly similar, although technically, the mass of a neutron is slightly larger than the mass of a proton. Because protons and neutrons are so much more massive than electrons, almost all of the mass of any atom comes from the nucleus, which contains all of the neutrons and protons.

Table 2.11.1: Properties of Subatomic Particles

Particle	Symbol	Mass (amu)	Relative Mass (proton = 1)	Relative Charge	Location
proton	p^+	1	1	+1	inside the nucleus
electron	e^-	5.45×10^{-4}	0.00055	−1	outside the nucleus
neutron	n^0	1	1	0	inside the nucleus

Table 2.11.1 gives the properties and locations of electrons, protons, and neutrons. The third column shows the masses of the three subatomic particles in "atomic mass units." An **atomic mass unit (amu)** is defined as one-twelfth of the mass of a carbon-12 atom. Atomic mass units (amu) are useful, because, as you can see, the mass of a proton and the mass of a neutron are almost exactly 1 in this unit system.

Negative and positive charges of equal magnitude cancel each other out. This means that the negative charge on an electron perfectly balances the positive charge on the proton. In other words, a neutral atom must have exactly one electron for every proton. If a neutral atom has 1 proton, it must have 1 electron. If a neutral atom has 2 protons, it must have 2 electrons. If a neutral atom has 10 protons, it must have 10 electrons. You get the idea. In order to be neutral, an atom must have the same number of electrons and protons.

Summary

- Electrons are a type of subatomic particle with a negative charge.
- Protons are a type of subatomic particle with a positive charge. Protons are bound together in an atom's nucleus as a result of the strong nuclear force.
- Neutrons are a type of subatomic particle with no charge (they are neutral). Like protons, neutrons are bound into the atom's nucleus as a result of the strong nuclear force.
- Protons and neutrons have approximately the same mass, but they are both much more massive than electrons (approximately 2,000 times as massive as an electron).
- The positive charge on a proton is equal in magnitude to the negative charge on an electron. As a result, a neutral atom must have an equal number of protons and electrons.
- The atomic mass unit (amu) is a unit of mass equal to one-twelfth the mass of a carbon-12 atom

2.11: The Properties of Protons, Neutrons, and Electrons is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.4: The Properties of Protons, Neutrons, and Electrons** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.12: Elements- Defined by Their Number of Protons

Learning Objectives

- Define atomic number.
- Define mass number.
- Determine the number of protons, neutrons, and electrons in an atom.

It's important to be able to distinguish atoms of one element from atoms of another element. Elements are pure substances that make up all other matter, so each one is given a unique name. The names of elements are also represented by unique one- or two-letter symbols, such as **H** for hydrogen, **C** for carbon, or **He** for helium. However, it would more powerful if these names could be used to identify the numbers of protons and neutrons in the atoms. That's where atomic number and mass number are useful.



Figure **2.12.1**: It is difficult to find qualities that differ between each element, and to distinguish one element from another. Each element, however, does have a unique number of protons. Sulfur has 16 protons, silicon has 14 protons, and gold has 79 protons. Images used with permission (public domain for sulfur and silicon, gold is licensed by CC-BY-SA-NC-ND; Alchemist-hp).

Atomic Number

Scientists distinguish between different elements by counting the number of protons in the nucleus (Table **2.12.1**). If an atom has only one proton, we know that it's a hydrogen atom. An atom with two protons is always a helium atom. If scientists count four protons in an atom, they know it's a beryllium atom. An atom with three protons is a lithium atom, an atom with five protons is a boron atom, an atom with six protons is a carbon atom . . . the list goes on.

Since an atom of one element can be distinguished from an atom of another element by the number of protons in its nucleus, scientists are always interested in this number, and how this number differs between different elements. The number of protons in an atom is called its **atomic number** (*Z*). This number is very important because it is unique for atoms of a given element. All atoms of an element have the same number of protons, and every element has a different number of protons in its atoms. For example, all helium atoms have two protons, and no other elements have atoms with two protons.

Table **2.12.1**: Atoms of the First Six Elements

Name	Protons	Neutrons	Electrons	Atomic Number (Z)	Mass Number (A)
Hydrogen	1	0	1	1	1
Helium	2	2	2	2	4
Lithium	3	4	3	3	7
Beryllium	4	5	4	4	9
Boron	5	6	5	5	11
Carbon	6	6	6	6	12

Of course, since neutral atoms have to have one electron for every proton, an element's atomic number also tells you how many electrons are in a neutral atom of that element. For example, hydrogen has an atomic number of 1. This means that an atom of hydrogen has one proton, and, if it's neutral, one electron as well. Gold, on the other hand, has an atomic number of 79, which means that an atom of gold has 79 protons, and, if it's neutral, 79 electrons as well.

Neutral Atoms

Atoms are neutral in electrical charge because they have the same number of negative electrons as positive protons (Table 2.12.1). Therefore, the atomic number of an atom also tells you how many electrons the atom has. This, in turn, determines many of the atom's chemical properties.

Mass Number

The **mass number** (A) of an atom is the total number of protons and neutrons in its nucleus. The mass of the atom is a unit called the atomic mass unit (amu). One atomic mass unit is the mass of a proton, or about 1.67×10^{-27} kilograms, which is an extremely small mass. A neutron has just a tiny bit more mass than a proton, but its mass is often assumed to be one atomic mass unit as well. Because electrons have virtually no mass, just about all the mass of an atom is in its protons and neutrons. Therefore, the total number of protons and neutrons in an atom determines its mass in atomic mass units (Table 2.12.1).

Consider helium again. Most helium atoms have two neutrons in addition to two protons. Therefore the mass of most helium atoms is 4 atomic mass units (2 amu for the protons + 2 amu for the neutrons). However, some helium atoms have more or less than two neutrons. Atoms with the same number of protons but different numbers of neutrons are called isotopes. Because the number of neutrons can vary for a given element, the mass numbers of different atoms of an element may also vary. For example, some helium atoms have three neutrons instead of two (these are called isotopes and are discussed in detail later on).

Why do you think that the "mass number" includes protons and neutrons, but not electrons? You know that most of the mass of an atom is concentrated in its nucleus. The mass of an atom depends on the number of protons and neutrons. You have already learned that the mass of an electron is very, very small compared to the mass of either a proton or a neutron (like the mass of a penny compared to the mass of a bowling ball). Counting the number of protons and neutrons tells scientists about the total mass of an atom.

$$\text{mass number } A = (\text{number of protons}) + (\text{number of neutrons})$$

An atom's mass number is very easy to calculate, provided that you know the number of protons and neutrons in an atom.

Example 4.5.1

What is the mass number of an atom of helium that contains 2 neutrons?

Solution

$(\text{number of protons}) = 2$ (Remember that an atom of helium always has 2 protons.)

$(\text{number of neutrons}) = 2$

$\text{mass number} = (\text{number of protons}) + (\text{number of neutrons})$

$$\text{mass number} = 2 + 2 = 4$$

A **chemical symbol** is a one- or two-letter designation of an element. Some examples of chemical symbols are **O** for oxygen, **Zn** for zinc, and **Fe** for iron. The first letter of a symbol is always capitalized. If the symbol contains two letters, the second letter is lower case. The majority of elements have symbols that are based on their English names. However, some of the elements that have been known since ancient times have maintained symbols that are based on their Latin names, as shown in Table 2.12.2.

Table 2.12.2: *Symbols and Latin Names for Elements*

Chemical Symbol	Name	Latin Name
Na	Sodium	Natrium
K	Potassium	Kalium
Fe	Iron	Ferrum
Cu	Copper	Cuprum
Ag	Silver	Argentum
Sn	Tin	Stannum
Sb	Antimony	Stibium
Au	Gold	Aurum
Pb	Lead	Plumbum

Summary

- Elements are pure substances that make up all matter, so each one is given a unique name.
- The names of elements are also represented by unique one- or two-letter symbols.
- Each element has a unique number of protons. An element's atomic number is equal to the number of protons in the nuclei of any of its atoms.
- The mass number of an atom is the sum of the protons and neutrons in the atom.
- Isotopes are atoms of the same element (same number of protons) that have different numbers of neutrons in their atomic nuclei.

2.12: Elements- Defined by Their Number of Protons is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.5: Elements- Defined by Their Number of Protons** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.13: Looking for Patterns - The Periodic Table

Learning Objectives

- Explain how elements are organized into the periodic table.
- Describe how some characteristics of elements relate to their positions on the periodic table.

In the 19th century, many previously unknown elements were discovered, and scientists noted that certain sets of elements had similar chemical properties. For example, chlorine, bromine, and iodine react with other elements (such as sodium) to make similar compounds. Likewise, lithium, sodium, and potassium react with other elements (such as oxygen) to make similar compounds. Why is this so?

In 1864, Julius Lothar Meyer, a German chemist, organized the elements by atomic mass and grouped them according to their chemical properties. Later that decade, Dmitri Mendeleev, a Russian chemist, organized all the known elements according to similar properties. He left gaps in his table for what he thought were undiscovered elements, and he made some bold predictions regarding the properties of those undiscovered elements. When elements were later discovered whose properties closely matched Mendeleev's predictions, his version of the table gained favor in the scientific community. Because certain properties of the elements repeat on a regular basis throughout the table (that is, they are periodic), it became known as the periodic table.

> *Mendeleev had to list some elements out of the order of their atomic masses to group them with other elements that had similar properties.*

The periodic table is one of the cornerstones of chemistry because it organizes all of the known elements on the basis of their chemical properties. A modern version is shown in Figure **2.13.1**. Most periodic tables provide additional data (such as atomic mass) in a box that contains each element's symbol. The elements are listed in order of atomic number.




Figure **2.13.1**: A Modern Periodic Table. A modern periodic table lists elements left to right by atomic number. An interactive Periodic table can be found Periodic Table of Elements. (Public Domain; PubChem via NIH)

Features of the Periodic Table

Elements that have similar chemical properties are grouped in columns called groups (or families). As well as being numbered, some of these groups have names—for example, *alkali metals* (the first column of elements), *alkaline earth metals* (the second column of elements), *halogens* (the next-to-last column of elements), and *noble gases* (the last column of elements).

> *The word halogen comes from the Greek for "salt maker" because these elements combine with other elements to form a group of compounds called salts.*

To Your Health: Radon

Radon is an invisible, odorless noble gas that is slowly released from the ground, particularly from rocks and soils whose uranium content is high. Because it is a noble gas, radon is not chemically reactive. Unfortunately, it is radioactive, and increased exposure to it has been correlated with an increased lung cancer risk.

Because radon comes from the ground, we cannot avoid it entirely. Moreover, because it is denser than air, radon tends to accumulate in basements, which if improperly ventilated can be hazardous to a building's inhabitants. Fortunately, specialized ventilation minimizes the amount of radon that might collect. Special fan-and-vent systems are available that draw air from below the basement floor, before it can enter the living space, and vent it above the roof of a house.

After smoking, radon is thought to be the second-biggest *preventable* cause of lung cancer in the United States. The American Cancer Society estimates that 10% of all lung cancers are related to radon exposure. There is uncertainty regarding what levels of exposure cause cancer, as well as what the exact causal agent might be (either radon or one of its breakdown products, many of which are also radioactive and, unlike radon, not gases). The US Environmental Protection Agency recommends testing every floor below the third floor for radon levels to guard against long-term health effects.

Each row of elements on the periodic table is called a period. Periods have different lengths; the first period has only 2 elements (hydrogen and helium), while the second and third periods have 8 elements each. The fourth and fifth periods have 18 elements each, and later periods are so long that a segment from each is removed and placed beneath the main body of the table.

Certain elemental properties become apparent in a survey of the periodic table as a whole. Every element can be classified as either a metal, a nonmetal, or a metalloid (or semi metal), as shown in Figure **2.13.2**. A **metal** is a substance that is shiny, typically (but not always) silvery in color, and an excellent conductor of electricity and heat. Metals are also malleable (they can be beaten into thin sheets) and ductile (they can be drawn into thin wires). A **nonmetal** is typically dull and a poor conductor of electricity and heat. Solid nonmetals are also very brittle. As shown in Figure **2.13.2**, metals occupy the left three-fourths of the periodic table, while nonmetals (except for hydrogen) are clustered in the upper right-hand corner of the periodic table. The elements with properties intermediate between those of metals and nonmetals are called **metalloids** (or **semi-metals**). Elements adjacent to the bold line in the right-hand portion of the periodic table have semimetal properties.




Figure **2.13.2**: Types of Elements. Elements are either metals, nonmetals, or metalloids (or semi metals). Each group is located in a different part of the periodic table. (CC BY-NC-SA; Anonymous by request)

Example 2.13.1

Based on its position in the periodic table, classify each element below as metal, a nonmetal, or a metalloid.

a. Se
b. Mg
c. Ge

Solution

a. In Figure **2.13.1**, selenium lies above and to the right of the diagonal line marking the boundary between metals and nonmetals, so it should be a nonmetal.
b. Magnesium lies to the left of the diagonal line marking the boundary between metals and nonmetals, so it should be a metal.
c. Germanium lies within the diagonal line marking the boundary between metals and nonmetals, so it should be a metalloid.

Exercise 2.13.1

Based on its location in the periodic table, do you expect indium to be a nonmetal, a metal, or a metalloid?

Answer

Indium is a metal.

Another way to categorize the elements of the periodic table is shown in Figure **2.13.3**. The first two columns on the left and the last six columns on the right are called the main group elements. The ten-column block between these columns contains the **transition metals**. The two rows beneath the main body of the periodic table contain the **inner transition metals**. The elements in these two rows are also referred to as, respectively, the **lanthanide metals** and the **actinide metals**.




Figure **2.13.3**: Special Names for Sections of the Periodic Table. (CC BY-NC-SA; Anonymous by request)

Sections of period table: Alkali metals, alkaline earth metals, transition metals, halogens, noble gases, lanthanide metals, actinide metals.

Descriptive Names

As previously noted, the periodic table is arranged so that elements with similar chemical behaviors are in the same group. Chemists often make general statements about the properties of the elements in a group using descriptive names with historical origins.

Group 1: The Alkali Metals

The alkali metals are lithium, sodium, potassium, rubidium, cesium, and francium. Hydrogen is unique in that it is generally placed in Group 1, but it is not a metal.

The compounds of the alkali metals are common in nature and daily life. One example is table salt (sodium chloride); lithium compounds are used in greases, in batteries, and as drugs to treat patients who exhibit manic-depressive, or bipolar, behavior. Although lithium, rubidium, and cesium are relatively rare in nature, and francium is so unstable and highly radioactive that it exists in only trace amounts, sodium and potassium are the seventh and eighth most abundant elements in Earth's crust, respectively.



Video: Alkali metals in water - Chemical elements: properties and reactions. (The Open University via https://youtu.be/6ZY6d6jrq-0)

Group 2: The Alkaline Earth Metals

The alkaline earth metals are beryllium, magnesium, calcium, strontium, barium, and radium. Beryllium, strontium, and barium are rare, and radium is unstable and highly radioactive. In contrast, calcium and magnesium are the fifth and sixth most abundant elements on Earth, respectively; they are found in huge deposits of limestone and other minerals.

Group 17: The Halogens

The halogens are fluorine, chlorine, bromine, iodine, and astatine. The name halogen is derived from the Greek words for "salt forming," which reflects that all of the halogens react readily with metals to form compounds, such as sodium chloride and calcium chloride (used in some areas as road salt).

Compounds that contain the fluoride ion are added to toothpaste and the water supply to prevent dental cavities. Fluorine is also found in Teflon coatings on kitchen utensils. Although chlorofluorocarbon propellants and refrigerants are believed to lead to the depletion of Earth's ozone layer and contain both fluorine and chlorine, the latter is responsible for the adverse effect on the ozone layer. Bromine and iodine are less abundant than chlorine, and astatine is so radioactive that it exists in only negligible amounts in nature.

Group 18: The Noble Gases

The noble gases are helium, neon, argon, krypton, xenon, and radon. Because the noble gases are composed of only single atoms, they are called monatomic. At room temperature and pressure, they are unreactive gases. Because of their lack of reactivity, for many years they were called inert gases or rare gases. However, the first chemical compounds containing the noble gases were prepared in 1962. Although the noble gases are relatively minor constituents of the atmosphere, natural gas contains substantial amounts of helium. Because of its low reactivity, argon is often used as an unreactive (inert) atmosphere for welding and in light bulbs. The red light emitted by neon in a gas discharge tube is used in neon lights.

Example 2.13.2: Groups

Provide the family or group name of each element.

a. Li
b. Ar
c. Cl

Solution

a. Lithium is an alkali metal (Group 1)
b. Argon is a noble gas (Group 18)
c. Chlorine is a halogen (Group 17)

? Exercise 2.13.2: Groups

Provide the family or group name of each element.

a. F
b. Ca
c. Kr

Answer a:

Fluorine is a halogen (Group 17).

Answer b:

Calcium is a alkaline earth metal (Group 2).

Answer c:

Krypton is a noble gas (Group 18).

✓ Example 2.13.3: Classification of Elements

Classify each element as metal, non metal, transition metal or inner transition metal.

a. Li
b. Ar
c. Am
d. Fe

Solution

a. Lithium is a metal.
b. Argon is a non metal.
c. Americium is an inner transition metal.
d. Iron is a transition metal.

? Exercise 2.13.3: Classification of Elements

Classify each element as metal, non metal, transition metal or inner transition metal.

a. F
b. U
c. Cu

Answer a:

Fluorine is a nonmetal.

Answer b:

Uranium is a metal (and a inner transition metal too).

Answer c:

Copper is a metal (and a transition metal too).

Summary

The periodic table is an arrangement of the elements in order of increasing atomic number. Elements that exhibit similar chemistry appear in vertical columns called groups (numbered 1–18 from left to right); the seven horizontal rows are called periods. Some of the groups have widely-used common names, including the alkali metals (Group 1) and the alkaline earth metals (Group 2) on the far left, and the halogens (Group 17) and the noble gases (Group 18) on the far right.

The elements can be broadly divided into metals, nonmetals, and semi metals. Semi metals exhibit properties intermediate between those of metals and nonmetals. Metals are located on the left of the periodic table, and nonmetals are located on the upper right. They are separated by a diagonal band of semi metals. Metals are lustrous, good conductors of electricity, and readily shaped (they are ductile and malleable). Solid nonmetals are generally brittle and poor electrical conductors. Other important groupings of elements in the periodic table are the main group elements, the transition metals, and the inner transition metals (the lanthanides, and the actinides).

References

1. Petrucci, Ralph H., William S. Harwood, F. G. Herring, and Jeffrey D. Madura. General Chemistry: Principles and Modern Applications. 9th ed. Upper Saddle River: Pearson Education, Inc., 2007.
2. Sisler, Harry H. Electronic structure, properties, and the periodic law. New york; Reinhold publishing corporation, 1963.
3. Petrucci, Ralph H., Carey Bissonnette, F. G. Herring, and Jeffrey D. Madura. General Chemistry: Principles and Modern Applications. Custom Edition for CHEM 2. Pearson Learning Solutions, 2010.

Contributions & Attributions

2.13: Looking for Patterns - The Periodic Table is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.6: Looking for Patterns - The Periodic Table** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.14: Ions - Losing and Gaining Electrons

Learning Objectives

- Define the two types of ions.

Atoms tend to lose, gain, or share some valance electrons, making bonds to acquire the electron configuration of the nearest noble gas, i.e., usually eight valence electrons for the main group elements, called an octet. Most atoms do not have eight electrons in their valence electron shell. Some atoms have only a few electrons in their outer shell, while others lack only one or two electrons to have an octet. In cases where an atom has three or fewer valence electrons, the atom may lose those valence electrons quite easily until what remains is a lower shell that contains an octet. Atoms that lose electrons acquire a positive charge because they are left with fewer negatively charged electrons to balance the positive charges of the protons in the nucleus. Positively charged ions are called **cations**. Most metals become cations when they make ionic compounds.

Cations

A neutral sodium atom is likely to achieve an octet in its outermost shell by losing its one valence electron.

$$Na \rightarrow Na^+ + e^-$$

The cation produced in this way, Na^+, is called the sodium ion to distinguish it from the element. The outermost shell of the sodium ion is the second electron shell, which has eight electrons in it. The octet rule has been satisfied. Figure **2.14.1** is a graphical depiction of this process.




Figure **2.14.1**: The Formation of a Sodium Ion. On the left, a sodium atom has 11 electrons. On the right, the sodium ion only has 10 electrons and a 1+ charge. Neutral sodium atom on left has 11 protons and 11 electrons. Sodium ion on right has 11 protons and 10 electrons, with a +1 overall charge.

Anions

Some atoms have nearly eight electrons in their valence shell and can gain additional valence electrons until they have an octet. When these atoms gain electrons, they acquire a negative charge because they now possess more electrons than protons. Negatively charged ions are called **anions**. Most nonmetals become anions when they make ionic compounds.

A neutral chlorine atom has seven electrons in its outermost shell. Only one more electron is needed to achieve an octet in chlorine's valence shell. (In table salt, this electron comes from the sodium atom.)

$$e^- + Cl \longrightarrow Cl^-$$

In this case, the ion has the same outermost shell as the original atom, but now that shell has eight electrons in it. Once again, the octet rule has been satisfied. The resulting anion, Cl^-, is called the chloride ion; note the slight change in the suffix (*-ide* instead of *-ine*) to create the name of this anion. Figure **2.14.2** is a graphical depiction of this process.




Figure **2.14.2**: The Formation of a Chlorine Ion. On the left, the chlorine atom has 17 electrons. On the right, the chloride ion has 18 electrons and has a 1− charge. Neutral chlorine atom on left has 17 protons and 17 electrons. Sodium ion on right has 17 protons and 18 electrons, with a -1 overall charge.

> *The names for positive and negative ions are pronounced CAT-eye-ons and ANN-eye-ons, respectively.*

In many cases, elements that belong to the same group (vertical column) on the periodic table form ions with the same charge because they have the same number of valence electrons. Thus, the periodic table becomes a tool for remembering the charges on many ions. For example, all ions made from alkali metals, the first column on the periodic table, have a 1+ charge. Ions made from alkaline earth metals, the second group on the periodic table, have a 2+ charge. On the other side of the periodic table, the next-to-last column, the halogens, form ions having a 1− charge. Figure **2.14.3** shows how the charge on many ions can be predicted by the location of an element on the periodic table. Note the convention of first writing the number and then the sign on a ion with multiple charges. The barium cation is written Ba^{2+}, not Ba^{+2}.




Figure **2.14.3**: Predicting Ionic Charges. The charge that an atom acquires when it becomes an ion is related to the structure of the periodic table. Within a group (family) of elements, atoms form ions of a certain charge.

2.14: Ions - Losing and Gaining Electrons is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.7: Ions - Losing and Gaining Electrons** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

2.15: Isotopes - When the Number of Neutrons Varies

Learning Objectives

- Explain what isotopes are and how an isotope affects an element's atomic mass.
- Determine the number of protons, electrons, and neutrons of an element with a given mass number.

All atoms of the same element have the same number of protons, but some may have different numbers of neutrons. For example, all carbon atoms have six protons, and most have six neutrons as well. But some carbon atoms have seven or eight neutrons instead of the usual six. Atoms of the same element that differ in their numbers of neutrons are called **isotopes**. Many isotopes occur naturally. Usually one or two isotopes of an element are the most stable and common. Different isotopes of an element generally have the same physical and chemical properties because they have the same numbers of protons and electrons.

An Example: Hydrogen Isotopes

Hydrogen is an example of an element that has isotopes. Three isotopes of hydrogen are modeled in Figure **2.15.1**. Most hydrogen atoms have just one proton, one electron, and lack a neutron. These atoms are just called hydrogen. Some hydrogen atoms have one neutron as well. These atoms are the isotope named deuterium. Other hydrogen atoms have two neutrons. These atoms are the isotope named tritium.




Figure **2.15.1**: The three most stable isotopes of hydrogen: protium (A = 1), deuterium (A = 2), and tritium (A = 3). (CC SA-BY 3.0; Balajijagadesh via Wikipedia).

For most elements other than hydrogen, isotopes are named for their mass number. For example, carbon atoms with the usual 6 neutrons have a mass number of 12 (6 protons + 6 neutrons = 12), so they are called carbon-12. Carbon atoms with 7 neutrons have an atomic mass of 13 (6 protons + 7 neutrons = 13). These atoms are the isotope called carbon-13.

Example 2.15.1: Lithium Isotopes

a. What is the atomic number and the mass number of an isotope of lithium containing 3 neutrons?
b. What is the atomic number and the mass number of an isotope of lithium containing 4 neutrons?

Solution

A lithium atom contains 3 protons in its nucleus irrespective of the number of neutrons or electrons.

a.

$$\begin{aligned}\text{atomic number} &= (\text{number of protons}) = 3\\ &\quad (\text{number of neutrons}) = 3\end{aligned}$$

$$\begin{aligned}\text{mass number} &= (\text{number of protons}) + (\text{number of neutrons})\\ \text{mass number} &= 3 + 3\\ &= 6\end{aligned}$$

b.

$$\begin{aligned}\text{atomic number} &= (\text{number of protons}) = 3\\ &\quad (\text{number of neutrons}) = 4\end{aligned}$$

$$\text{mass number} = (\text{number of protons}) + (\text{number of neutrons})$$
$$\text{mass number} = 3 + 4$$
$$= 7$$

Notice that because the lithium atom always has 3 protons, the atomic number for lithium is always 3. The mass number, however, is 6 in the isotope with 3 neutrons, and 7 in the isotope with 4 neutrons. In nature, only certain isotopes exist. For instance, lithium exists as an isotope with 3 neutrons, and as an isotope with 4 neutrons, but it doesn't exist as an isotope with 2 neutrons or as an isotope with 5 neutrons.

Stability of Isotopes

Atoms need a certain ratio of neutrons to protons to have a stable nucleus. Having too many or too few neutrons relative to protons results in an unstable, or radioactive, nucleus that will sooner or later break down to a more stable form. This process is called radioactive decay. Many isotopes have radioactive nuclei, and these isotopes are referred to as radioisotopes. When they decay, they release particles that may be harmful. This is why radioactive isotopes are dangerous and why working with them requires special suits for protection. The isotope of carbon known as carbon-14 is an example of a radioisotope. In contrast, the carbon isotopes called carbon-12 and carbon-13 are stable.

This whole discussion of isotopes brings us back to Dalton's Atomic Theory. According to Dalton, atoms of a given element are identical. But if atoms of a given element can have different numbers of neutrons, then they can have different masses as well! How did Dalton miss this? It turns out that elements found in nature exist as constant uniform mixtures of their naturally occurring isotopes. In other words, a piece of lithium always contains both types of naturally occurring lithium (the type with 3 neutrons and the type with 4 neutrons). Moreover, it always contains the two in the same relative amounts (or "relative abundance"). In a chunk of lithium, 93% will always be lithium with 4 neutrons, while the remaining 7% will always be lithium with 3 neutrons.

Dalton always experimented with large chunks of an element—chunks that contained all of the naturally occurring isotopes of that element. As a result, when he performed his measurements, he was actually observing the averaged properties of all the different isotopes in the sample. For most of our purposes in chemistry, we will do the same thing and deal with the average mass of the atoms. Luckily, aside from having different masses, most other properties of different isotopes are similar.

There are two main ways in which scientists frequently show the mass number of an atom they are interested in. It is important to note that the mass number is *not* given on the periodic table. These two ways include writing a nuclear symbol or by giving the name of the element with the mass number written.

To write a **nuclear symbol**, the mass number is placed at the upper left (superscript) of the chemical symbol and the atomic number is placed at the lower left (subscript) of the symbol. The complete nuclear symbol for helium-4 is drawn below:



Nuclear symbol for helium-4: The element symbol is He, the mass number to the top left is 4, and the atomic number to the bottom left is 2


The following nuclear symbols are for a nickel nucleus with 31 neutrons and a uranium nucleus with 146 neutrons.

$$^{59}_{28}\text{Ni}$$

$$^{238}_{92}\text{U}$$

In the nickel nucleus represented above, the atomic number 28 indicates that the nucleus contains 28 protons, and therefore, it must contain 31 neutrons in order to have a mass number of 59. The uranium nucleus has 92 protons, as all uranium nuclei do; and this particular uranium nucleus has 146 neutrons.

Another way of representing isotopes is by adding a hyphen and the mass number to the chemical name or symbol. Thus the two nuclei would be Nickel-59 or Ni-59 and Uranium-238 or U-238, where 59 and 238 are the mass numbers of the two atoms, respectively. Note that the mass numbers (not the number of neutrons) are given to the side of the name.



Example 2.15.2: Potassium-40

How many protons, electrons, and neutrons are in an atom of $^{40}_{19}\mathrm{K}$?

Solution

$$\text{atomic number} = (\text{number of protons}) = 19$$

For all atoms with no charge, the number of electrons is equal to the number of protons.

$$\text{number of electrons} = 19$$

The mass number, 40, is the sum of the protons and the neutrons.

To find the number of neutrons, subtract the number of protons from the mass number.

$$\text{number of neutrons} = 40 - 19 = 21.$$

Example 2.15.3: Zinc-65

How many protons, electrons, and neutrons are in an atom of zinc-65?

Solution

$$\text{number of protons} = 30$$

For all atoms with no charge, the number of electrons is equal to the number of protons.

$$\text{number of electrons} = 30$$

The mass number, 65, is the sum of the protons and the neutrons.

To find the number of neutrons, subtract the number of protons from the mass number.

$$\text{number of neutrons} = 65 - 30 = 35$$

Exercise 2.15.3

How many protons, electrons, and neutrons are in each atom?

a. $^{60}_{27}\mathrm{Co}$
b. Na-24
c. $^{45}_{20}\mathrm{Ca}$
d. Sr-90

Answer a:

27 protons, 27 electrons, 33 neutrons

Answer b:

11 protons, 11 electrons, 13 neutrons

Answer c:

20 protons, 20 electrons, 25 neutrons

Answer d:

38 protons, 38 electrons, 52 neutrons

Summary

- The number of protons is always the same in atoms of the same element.
- The number of neutrons can be different, even in atoms of the same element.
- Atoms of the same element that contain the same number of protons, but different numbers of neutrons, are known as **isotopes**.
- Isotopes of any given element all contain the same number of protons, so they have the same atomic number (for example, the atomic number of helium is always 2).
- Isotopes of a given element contain different numbers of neutrons, therefore, different isotopes have different mass numbers.

2.15: Isotopes - When the Number of Neutrons Varies is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.8: Isotopes - When the Number of Neutrons Varies** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.16: Atomic Mass - The Average Mass of an Element's Atoms

Learning Objectives

- Explain what is meant by the atomic mass of an element.
- Calculate the atomic mass of an element from the masses and relative percentages of the isotopes of the element.

In chemistry we very rarely deal with only one isotope of an element. We use a mixture of the isotopes of an element in chemical reactions and other aspects of chemistry, because all of the isotopes of an element react in the same manner. That means that we rarely need to worry about the mass of a specific isotope, but instead we need to know the average mass of the atoms of an element. Using the masses of the different isotopes and how abundant each isotope is, we can find the average mass of the atoms of an element. The **atomic mass** of an element is the weighted average mass of the atoms in a naturally occurring sample of the element. Atomic mass is typically reported in atomic mass units.

Calculating Atomic Mass

You can calculate the atomic mass (or average mass) of an element provided you know the **relative abundance** (the fraction of an element that is a given isotope), the element's naturally occurring isotopes, and the masses of those different isotopes. We can calculate this by the following equation:

$$\text{Atomic mass} = (\%_1)\,(\text{mass}_1) + (\%_2)\,(\text{mass}_2) + \cdots$$

Look carefully to see how this equation is used in the following examples.

Example 2.16.1: Boron Isotopes

Boron has two naturally occurring isotopes. In a sample of boron, 20% of the atoms are B-10, which is an isotope of boron with 5 neutrons and mass of $10\ \text{amu}$. The other 80% of the atoms are B-11, which is an isotope of boron with 6 neutrons and a mass of $11\ \text{amu}$. What is the atomic mass of boron?

Solution

Boron has two isotopes. We will use the equation:

$$\text{Atomic mass} = (\%_1)\,(\text{mass}_1) + (\%_2)\,(\text{mass}_2) + \cdots$$

- Isotope 1: $\%_1 = 0.20$ (Write all percentages as decimals), $\text{mass}_1 = 10$
- Isotope 2: $\%_2 = 0.80$, $\text{mass}_2 = 11$

Substitute these into the equation, and we get:

$$\text{Atomic mass} = (0.20)\,(10) + (0.80)\,(11)$$

$$\text{Atomic mass} = 10.8\ \text{amu}$$

The mass of an average boron atom, and thus boron's atomic mass, is $10.8\ \text{amu}$.

Example 2.16.2: Neon Isotopes

Neon has three naturally occurring isotopes. In a sample of neon, 90.92% of the atoms are Ne-20, which is an isotope of neon with 10 neutrons and a mass of $19.99\ \text{amu}$. Another 0.3% of the atoms are Ne-21, which is an isotope of neon with 11 neutrons and a mass of $20.99\ \text{amu}$. The final 8.85% of the atoms are Ne-22, which is an isotope of neon with 12 neutrons and a mass of $21.99\ \text{amu}$. What is the atomic mass of neon?

Solution

Neon has three isotopes. We will use the equation:

$$\text{Atomic mass} = (\%_1)\,(\text{mass}_1) + (\%_2)\,(\text{mass}_2) + \cdots$$

- Isotope 1: $\%_1 = 0.9092$ (write all percentages as decimals), $\text{mass}_1 = 19.99$

- Isotope 2: $\%_2 = 0.003$, $\text{mass}_2 = 20.99$
- Isotope 3: $\%_3 = 0.0885$, $\text{mass}_3 = 21.99$

Substitute these into the equation, and we get:

$$\text{Atomic mass} = (0.9092)(19.99) + (0.003)(20.99) + (0.0885)(21.99)$$

$$\text{Atomic mass} = 20.17\ \text{amu}$$

The mass of an average neon atom is $20.17\ \text{amu}$

The periodic table gives the atomic mass of each element. The atomic mass is a number that usually appears below the element's symbol in each square. Notice that the atomic mass of boron (symbol B) is 10.8, which is what we calculated in Example 2.16.1, and the atomic mass of neon (symbol Ne) is 20.8, which is what we calculated in Example 2.16.2. Take time to notice that not all periodic tables have the atomic number above the element's symbol and the mass number below it. If you are ever confused, remember that the atomic number should always be the smaller of the two and will be a whole number, while the atomic mass should always be the larger of the two and will be a decimal number.

? Exercise 2.16.1

Chlorine has two naturally occurring isotopes. In a sample of chlorine, 75.77% of the atoms are Cl-35, with a mass of 34.97 amu. Another 24.23% of the atoms are Cl-37, with a mass of 36.97 amu. What is the atomic mass of chlorine?

Answer

35.45 amu

Summary

- An element's atomic mass is the weighted average of the masses of the isotopes of an element
- An element's atomic mass can be calculated provided the relative abundance of the element's naturally occurring isotopes and the masses of those isotopes are known.
- The periodic table is a convenient way to summarize information about the different elements. In addition to the element's symbol, most periodic tables will also contain the element's atomic number and the element's atomic mass.

2.16: Atomic Mass - The Average Mass of an Element's Atoms is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **4.9: Atomic Mass - The Average Mass of an Element's Atoms** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.17: Sugar and Salt

Sodium chloride, also known as table salt, is an ionic compound with the chemical formula $\mathbf{NaCl}$, representing a 1:1 ratio of sodium and chloride ions. It is commonly used as a condiment and food preservative. Salt can be created by adding two very reactive elements together: sodium $(\mathbf{Na\,(s)}$ metal and chlorine $(\mathbf{Cl_2\,(g)}$ gas.

$$\mathbf{2\,Na\,(s) + Cl_2\,(g) \rightarrow 2\,NaCl\,(s)} \tag{2.17.1}$$

The element sodium (Figure **2.17.1***a*) is a very reactive metal; given the opportunity, it will react with the sweat on your hands and form sodium hydroxide, which is a very corrosive substance. The element chlorine (Figure **2.17.1***b*) is a pale yellow, corrosive gas that should not be inhaled due to its poisonous nature. Bring these two hazardous substances together, however, and they react to make the ionic compound sodium chloride (Figure **2.17.1***c*), known simply as salt.




Figure **2.17.1**: Sodium + Chlorine = Sodium Chloride (a) Sodium is a very reactive metal. (b) Chlorine is a pale yellow, noxious gas. (c) Together, sodium and chlorine make sodium chloride—salt—which is necessary for our survival. Source: Photo on the left courtesy of reenhorn1 and photo in the center courtesy of Benjah-bmm27. Photo on the right © Thinkstock.

Salt is necessary for life. $\mathbf{Na^+}$ ions are one of the main ions in the human body and are necessary to regulate the fluid balance in the body. $\mathbf{Cl^-}$ ions are necessary for proper nerve function and respiration. Both of these ions are supplied by salt. The taste of salt is one of the fundamental tastes; salt is probably the most ancient flavoring known, and one of the few rocks we eat. Clearly when the elemental sodium and chlorine combine (Equation 2.17.1), the resulting salt product has radically different properties (both physical and chemical). This reaction is spectacular to observe (Video **2.17.1**).



Video* 2.17.1: Making Table Salt using Sodium Metal and Chlorine gas*

Another compound is sugar, which is the generic name for sweet, soluble carbohydrates, many of which are used in food. Sugar has the chemical formulate $\mathbf{C_{12}H_{22}O_{11}}$ and is constructed from different elements than salt: carbon, hydrogen and oxygen. While sugar qualitatively resembles table salt (often confused in the kitchen), the two have distinctly different physical and chemical properties. There are various types of sugar derived from different sources. While sugar is made with carbon, hydrogen, and oxygen, it is considerably harder to synthesize from its constituent elements than table salt is (Equation 2.17.1). However, the thermal

decomposition is considerably easier and can be represented as a dehydration of sucrose to pure carbon and water vapor in Equation 2.17.2, and demonstrated in Video 2.17.2.

$$\mathrm{C_{12}H_{22}O_{11}\,(s) + heat \longrightarrow 12\,C\,(s) + 11\,H_2O\,(g)} \tag{2.17.2}$$



***Video* 2.17.2:** *A science experiment in the kitchen shows what happens to sugar molecules when they are heated. The experiment did not disappoint!*

As with salt, sugar has radically different properties (both physical and chemical) than its constituent elements. This difference in properties, of constituent elements and compounds, is a central feature of chemical reactions.

Contributions & Attributions

- Wikipedia

2.17: Sugar and Salt is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.1: Sugar and Salt** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.18: Compounds Display Constant Composition

When building a house, the starting point is a blueprint of what the house will look like. The plan states how many windows and what kind, how many doors and what style, how many rooms and what type (bedroom, kitchen, other). The blueprint shows how the different pieces will go together to make the house. As long as the blueprint is followed and exactly the same items are used, the house will be identical to its blueprint.

Compounds

A **compound** is a substance that contains two or more elements chemically combined in a fixed proportion. The elements carbon and hydrogen combine to form many different compounds. One of the simplest is called methane, in which there are always four times as many hydrogen particles as carbon particles. Methane is a pure substance because it always has the same composition. However, it is not an element because it can be broken down into simpler substances—carbon and hydrogen.

Recall that the components of a mixture can be separated from one another by physical means. This is not true for a compound. Table salt is a compound consisting of equal parts of the elements sodium and chlorine. Salt cannot be separated into its two elements by filtering, distillation, or any other physical process. Salt and other compounds can only be decomposed into their elements by a chemical process. A **chemical change** is a change that produces matter with a different composition. Many compounds can be decomposed into their elements by heating. When sugar is heated, it decomposes into carbon and water. Water is still a compound, but one which cannot be broken down into hydrogen and oxygen by heating. Instead, the passage of an electrical current through water will produce hydrogen and oxygen gases.

The properties of compounds are generally very different than the properties of the elements from which the compound is formed. Sodium is an extremely reactive soft metal that cannot be exposed to air or water. Chlorine is a deadly gas. The compound sodium chloride is a white solid which is essential for all living things (see below).




Figure **2.18.1**: (A) Sodium is so reactive that it must be stored under oil. (B) Chlorine is a poisonous yellow-green gas. (C) Salt crystals, a compound of sodium and chlorine.

Summary

- A compound is a substance that contains two or more elements chemically combined in a fixed proportion.
- A chemical change is a change that produces matter with a different composition.

2.18: Compounds Display Constant Composition is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.2: Compounds Display Constant Composition** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.19: Chemical Formulas - How to Represent Compounds

Learning Objectives

- Determine the number of different atoms in a formula.
- Define chemical formula, molecular formula, and empirical formula.

A **chemical formula** is an expression that shows the elements in a compound and the relative proportions of those elements. Water is composed of hydrogen and oxygen in a 2:1 ratio. The chemical formula for water is H_2O. Sulfuric acid is one of the most widely produced chemicals in the United States and is composed of the elements hydrogen, sulfur, and oxygen. The chemical formula for sulfuric acid is H_2SO_4.

Certain groups of atoms are bonded together to form what is called a polyatomic ion that acts as a single unit. Polyatomic ions are discussed in more detail in Section 5.5. Polyatomic ions are enclosed in parenthesis followed by a subscript if more than one of the same ion exist in a chemical formula. The formula $Ca_3(PO_4)_2$ represents a compound with the following:

$$3 \text{ Ca atoms} + 2 \ PO_4^{3-} \text{ ions}$$

To count the total number of atoms for formulas with polyatomic ions enclosed in parenthesis, use the subscript as a multiplier for each atom or number of atoms.

$$Ca_3(PO_4)_2$$

$$3 \text{ Ca} + 2 \text{ x1 P} + 2 \text{ x 4 O} = 3 \text{ Ca atoms} + 2 \text{ P atoms} + 8 \text{ O atoms}$$

Molecular Formula

A **molecular formula** is a chemical formula of a molecular compound that shows the kinds and numbers of atoms present in a molecule of the compound. Ammonia is a compound of nitrogen and hydrogen as shown below:




Figure 2.19.1: The molecular formula for ammonia. NH3. There is one atom of nitrogen and 3 atoms of hydrogen in a molecule of ammonia.

Note from the example that there are some standard rules to follow in writing molecular formulas. The arrangements of the elements depend on the particular structure, which is not of concern at this point. The number of atoms of each kind is indicated by a subscript following the atom. If there is only one atom, no number is written. If there is more than one atom of a specific kind, the number is written as a subscript following the atom. We would not write N_3H for ammonia, because that would mean that there are three nitrogen atoms and one hydrogen atom in the molecule, which is incorrect.

Empirical Formula

An **empirical formula** is a formula that shows the elements in a compound in their lowest whole-number ratio. Glucose is an important simple sugar that cells use as their primary source of energy. Its molecular formula is $C_6H_{12}O_6$. Since each of the subscripts is divisible by 6, the empirical formula for glucose is CH_2O. When chemists analyze an unknown compound, often the first step is to determine its empirical formula.

- molecular formula: $C_6H_{12}O_6$
- empirical formula: CH_2O

There are a great many compounds whose molecular and empirical formulas are the same. If the molecular formula cannot be simplified into a smaller whole-number ratio, as in the case of H_2O or P_2O_5, then the empirical formula is also the molecular formula.

Summary

- A chemical formula is an expression that shows the elements in a compound and the relative proportions of those elements.
- If only one atom of a specific type is present, no subscript is used.
- For atoms that have two or more of a specific type of atom present, a subscript is written after the symbol for that atom.
- Polyatomic ions in chemical formulas are enclosed in parentheses followed by a subscript if more than one of the same type of polyatomic ion exist.
- Molecular formulas do not indicate how the atoms are arranged in the molecule.
- The empirical formula tells the lowest whole-number ratio of elements in a compound. The empirical formula does not show the actual number of atoms.

2.19: Chemical Formulas - How to Represent Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.3: Chemical Formulas - How to Represent Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.20: A Molecular View of Elements and Compounds

Learning Objectives

- Classify substances as atomic elements, molecular elements, molecular compounds, or ionic compounds.

Atomic Elements

Most elements exist with **individual atoms** as their basic unit. It is assumed that there is only one atom in a formula if there is no numerical subscript on the right side of an element's symbol.

Molecular Elements

There are many substances that exist as two or more atoms connected together so strongly that they behave as a single particle. These multi-atom combinations are called **molecules**. A molecule is the smallest part of a substance that has the physical and chemical properties of that substance. In some respects, a molecule is similar to an atom. A molecule, however, is composed of more than one atom.

Table **2.20.1**: Elements That Exist as Diatomic Molecules

Hydrogen, H	Oxygen	Nitrogen	Fluorine	Chlorine	Bromine	Iodine

Some elements exist naturally as molecules. For example, hydrogen and oxygen exist as two-atom molecules. Other elements also exist naturally as diatomic molecules—a molecule with only two atoms (Table **2.20.1**). As with any molecule, these elements are labeled with a **molecular formula**, a formal listing of what and how many atoms are in a molecule. (Sometimes only the word *formula* is used, and its meaning is inferred from the context.) For example, the molecular formula for elemental hydrogen is H_2, with H being the symbol for hydrogen and the subscript 2 implying that there are two atoms of this element in the molecule. Other diatomic elements have similar formulas: O_2, N_2, and so forth. Other elements exist as molecules—for example, sulfur normally exists as an eight-atom molecule, S_8, while phosphorus exists as a four-atom molecule, P_4 (Figure **2.20.1**).




Figure **2.20.1**: Molecular Art of S_8 and P_4 Molecules. If each green ball represents a sulfur atom, then the diagram on the left represents an S_8 molecule. The molecule on the right shows that one form of elemental phosphorus exists, as a four-atom molecule.

Figure **2.20.1** shows two examples of how molecules will be represented in this text. An atom is represented by a small ball or sphere, which generally indicates where the nucleus is in the molecule. A cylindrical line connecting the balls represents the connection between the atoms that make this collection of atoms a molecule. This connection is called a chemical bond.

Ionic Compounds

The elements in the periodic table are divided into specific groupings; the metals, the non-metals, the semi-metals, and so on. These groupings are largely based on physical properties and on the tendency of the various elements to bond with other elements by forming either an ionic or a covalent bond. As a general rule of thumb, compounds that involve a metal binding with either a non-metal or a semi-metal will display ionic bonding. Thus, the compound formed from sodium and chlorine will be ionic (a metal and a non-metal). The basic unit of ionic compounds is the **formula unit**.

Molecular Compounds

Compounds that are composed of only non-metals or semi-metals with non-metals will display covalent bonding and will be classified as molecular compounds. Nitrogen monoxide (NO) will be a covalently bound molecule (two non-metals) and silicon dioxide (SiO_2) will also be a covalently bound molecule (a semi-metal and a non-metal). The basic unit of molecular compounds is the **molecule**.

Example 2.20.1

Provide the classification (i.e. atomic element, molecular element, molecular compound, or ionic compound) of each substance.

a. Fe
b. PCl_3
c. LiBr
d. P_4
e. oxygen gas

Solution

a. **Fe** (iron) is an element that is represented with no subscript, so it is an **atomic element**.
b. **PCl_3** is made up of two nonmetals, so it is a **molecular compound**.
c. **LiBr** is made up of lithium, a metal, and bromine, a nonmetal, so it is an **ionic compound**.
d. **P_4** is a substance that is made up of four atoms of the same element, so it is a **molecular element**.
e. The formula for **oxygen gas** is **O_2** so it is a molecular element.

Exercise 2.20.1

Provide the classification (i.e. atomic element, molecular element, molecular compound, or ionic compound) of each substance.

a. I_2
b. He
c. H_2O
d. Al
e. CuCl

Answer a:

molecular element

Answer b:

atomic element

Answer c:

molecular compound

Answer d:

atomic element

Answer e:

ionic compound

2.20: A Molecular View of Elements and Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.4: A Molecular View of Elements and Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.21: Writing Formulas for Ionic Compounds

Learning Objectives

- Write the correct formula for an ionic compound.
- Recognize polyatomic ions in chemical formulas.

Ionic compounds do not exist as molecules. In the solid state, ionic compounds are in crystal lattice containing many ions each of the cation and anion. An ionic formula, like $NaCl$, is an empirical formula. This formula merely indicates that sodium chloride is made of an equal number of sodium and chloride ions. Sodium sulfide, another ionic compound, has the formula Na_2S. This formula indicates that this compound is made up of twice as many sodium ions as sulfide ions. This section will teach you how to find the correct ratio of ions, so that you can write a correct formula.

If you know the name of a binary ionic compound, you can write its **chemical formula**. Start by writing the metal ion with its charge, followed by the nonmetal ion with its charge. Because the overall compound must be electrically neutral, decide how many of each ion is needed in order for the positive and negative charges to cancel each other out.

Example 2.21.1: Aluminum Nitride and Lithium Oxide

Write the formulas for aluminum nitride and lithium oxide.

Solution

Solution to Example 5.5.1

	Write the formula for aluminum nitride	Write the formula for lithium oxide
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Al^{3+} N^{3-}	Li^{+} O^{2-}
2. Use a multiplier to make the total charge of the cations and anions equal to each other.	total charge of cations = total charge of anions 1(3+) = 1(3-) +3 = -3	total charge of cations = total charge of anions 2(1+) = 1(2-) +2 = -2
3. Use the multipliers as subscript for each ion.	Al_1N_1	Li_2O_1
4. Write the final formula. Leave out all charges and all subscripts that are 1.	AlN	Li_2O

An alternative way to writing a correct formula for an ionic compound is to use the **crisscross method**. In this method, the numerical value of each of the ion charges is crossed over to become the subscript of the other ion. Signs of the charges are dropped.

Example 2.21.2: The Crisscross Method for Lead (IV) Oxide

Write the formula for lead (IV) oxide.

Solution

Solution to Example 5.5.2

Crisscross Method	Write the formula for lead (IV) oxide
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Pb^{4+} O^{2-}
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	
3. Reduce to the lowest ratio.	Pb_2O_4
4. Write the final formula. Leave out all subscripts that are 1.	PbO_2

Exercise 2.21.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the calcium ion and the oxygen ion
b. the 2+ copper ion and the sulfur ion
c. the 1+ copper ion and the sulfur ion

Answer a:

CaO

Answer b:

CuS

Answer c:

Cu_2S

> *Be aware that ionic compounds are empirical formulas and so must be written as the lowest ratio of the ions.*

Example 2.21.3: Sulfur Compound

Write the formula for sodium combined with sulfur.

Solution

Solution to Example 5.5.3

Crisscross Method	Write the formula for sodium combined with sulfur
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Na^{+} S^{2-}
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	
3. Reduce to the lowest ratio.	This step is not necessary.
4. Write the final formula. Leave out all subscripts that are 1.	Na_2S

Exercise 2.21.3

Write the formula for each ionic compound.

a. sodium bromide

b. lithium chloride
c. magnesium oxide

Answer a:
NaBr
Answer b:
LiCl
Answer c:
MgO

Polyatomic Ions

Some ions consist of groups of atoms bonded together and have an overall electric charge. Because these ions contain more than one atom, they are called polyatomic ions. Polyatomic ions have characteristic formulas, names, and charges that should be memorized. For example, NO_3^- is the nitrate ion; it has one nitrogen atom and three oxygen atoms and an overall 1− charge. Table **2.21.1** lists the most common polyatomic ions.

Table **2.21.1**: Some Polyatomic Ions

Name	Formula
ammonium ion	NH_4^+
acetate ion	$C_2H_3O_2^-$ (also written $CH_3CO_2^-$)
carbonate ion	CO_3^{2-}
chromate ion	CrO_4^{2-}
dichromate ion	$Cr_2O_7^{2-}$
hydrogen carbonate ion (bicarbonate ion)	HCO_3^-
cyanide ion	CN^-
hydroxide ion	OH^-
nitrate ion	NO_3^-
nitrite ion	NO_2^-
permanganate ion	MnO_4^-
phosphate ion	PO_4^{3-}
hydrogen phosphate ion	HPO_4^{2-}
dihydrogen phosphate ion	$H_2PO_4^-$
sulfate ion	SO_4^{2-}
hydrogen sulfate ion (bisulfate ion)	HSO_4^-
sulfite ion	SO_3^{2-}

The rule for constructing formulas for ionic compounds containing polyatomic ions is the same as for formulas containing monatomic (single-atom) ions: the positive and negative charges must balance. If more than one of a particular polyatomic ion is needed to balance the charge, the *entire formula* for the polyatomic ion must be enclosed in parentheses, and the numerical subscript is placed *outside* the parentheses. This is to show that the subscript applies to the entire polyatomic ion. An example is $Ba(NO_3)_2$.

Writing Formulas for Ionic Compounds Containing Polyatomic Ions

Writing a formula for ionic compounds containing polyatomic ions also involves the same steps as for a binary ionic compound. Write the symbol and charge of the cation followed by the symbol and charge of the anion.

Example 2.21.4: Calcium Nitrate

Write the formula for calcium nitrate.

Solution

Solution to Example 5.5.4

Crisscross Method	Write the formula for calcium nitrate
1. Write the symbol and charge of the cation (metal) first and the anion (nonmetal) second.	Ca^{2+} NO_3^-
2. Transpose only the number of the positive charge to become the subscript of the anion and the number only of the negative charge to become the subscript of the cation.	 The 2+ charge on Ca becomes the subscript of NO3 and the 1- charge on NO3 becomes the subscript of Ca.
3. Reduce to the lowest ratio.	$Ca_1(NO_3)_2$
4. Write the final formula. Leave out all subscripts that are 1. If there is only 1 of the polyatomic ion, leave off parentheses.	$Ca(NO_3)_2$

Example 2.21.5

Write the chemical formula for an ionic compound composed of the potassium ion and the sulfate ion.

Solution

Solution to Example 5.5.5

Explanation	Answer
Potassium ions have a charge of 1+, while sulfate ions have a charge of 2−. We will need two potassium ions to balance the charge on the sulfate ion, so the proper chemical formula is K_2SO_4.	K_2SO_4

Exercise 2.21.5

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the magnesium ion and the carbonate ion
b. the aluminum ion and the acetate ion

Answer a:

$MgCO_3$

Answer b:

$Al(CH_3COO)_3$

Recognizing Ionic Compounds

There are two ways to recognize ionic compounds.

Method 1

Compounds between metal and nonmetal elements are usually ionic. For example, $CaBr_2$ contains a metallic element (calcium, a group 2 [or 2A] metal) and a nonmetallic element (bromine, a group 17 [or 7A] nonmetal). Therefore, it is most likely an ionic compound (in fact, it *is* ionic). In contrast, the compound NO_2 contains two elements that are both nonmetals (nitrogen, from group 15 [or 5A], and oxygen, from group 16 [or 6A]. It is not an ionic compound; it belongs to the category of covalent compounds discussed elsewhere. Also note that this combination of nitrogen and oxygen has no electric charge specified, so it is *not* the nitrite ion.

Method 2

Second, if you recognize the formula of a polyatomic ion in a compound, the compound is ionic. For example, if you see the formula $Ba(NO_3)_2$, you may recognize the "NO_3" part as the nitrate ion, NO_3^-. (Remember that the convention for writing formulas for ionic compounds is not to include the ionic charge.) This is a clue that the other part of the formula, Ba, is actually the Ba^{2+} ion, with the 2+ charge balancing the overall 2− charge from the two nitrate ions. Thus, this compound is also ionic.

Example 2.21.6

Identify each compound as ionic or not ionic.

a. Na_2O
b. PCl_3
c. NH_4Cl
d. OF_2

Solution

Solution to Example 5.5.6

Explanation	Answer
a. Sodium is a metal, and oxygen is a nonmetal. Therefore, Na_2O is expected to be ionic via method 1.	Na_2O, ionic
b. Both phosphorus and chlorine are nonmetals. Therefore, PCl_3 is not ionic via method 1	PCl_3, not ionic
c. The NH_4 in the formula represents the ammonium ion, NH_4^+, which indicates that this compound is ionic via method 2	NH_4Cl, ionic
d. Both oxygen and fluorine are nonmetals. Therefore, OF_2 is not ionic via method 1	OF_2, not ionic

Exercise 2.21.6

Identify each compound as ionic or not ionic.

a. N_2O
b. $FeCl_3$
c. $(NH_4)_3PO_4$
d. $SOCl_2$

Answer a:

not ionic

Answer b:

ionic

Answer c:

ionic

Answer d:

not ionic

Summary

Formulas for ionic compounds contain the symbols and number of each atom present in a compound in the lowest whole number ratio.

2.21: Writing Formulas for Ionic Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.5: Writing Formulas for Ionic Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.22: Nomenclature- Naming Compounds

Nomenclature is the process of naming chemical compounds so that they can be easily identified as separate chemicals. The primary function of chemical nomenclature is to ensure that a spoken or written chemical name leaves no ambiguity concerning which chemical compound the name refers to—each chemical name should refer to a single substance. A less important aim is to ensure that each substance has a single name, although a limited number of alternative names is acceptable in some cases. Preferably, the name also conveys some information about the structure or chemistry of a compound. A common name will often suffice to identify a chemical compound in a particular set of circumstances. To be more generally applicable, the name should indicate at least the chemical formula. To be more specific still, the three-dimensional arrangement of the atoms may need to be specified.

Contributions & Attributions

- Wikipedia (CC-BY-SA-3.0)

2.22: Nomenclature- Naming Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.6: Nomenclature- Naming Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.23: Naming Ionic Compounds

Learning Objectives

- To use the rules for naming ionic compounds.

After learning a few more details about the names of individual ions, you will be one step away from knowing how to name ionic compounds. This section begins the formal study of nomenclature, the systematic naming of chemical compounds.

Naming Ions

The name of a monatomic cation is simply the name of the element followed by the word *ion*. Thus, Na^+ is the sodium ion, Al^{3+} is the aluminum ion, Ca^{2+} is the calcium ion, and so forth.

We have seen that some elements lose different numbers of electrons, producing ions of different charges (Figure 3.3). Iron, for example, can form two cations, each of which, when combined with the same anion, makes a different compound with unique physical and chemical properties. Thus, we need a different name for each iron ion to distinguish Fe^{2+} from Fe^{3+}. The same issue arises for other ions with more than one possible charge.

There are two ways to make this distinction. In the simpler, more modern approach, called the **Stock system**, an ion's positive charge is indicated by a roman numeral in parentheses after the element name, followed by the word *ion*. Thus, Fe^{2+} is called the iron(II) ion, while Fe^{3+} is called the iron(III) ion. This system is used only for elements that form more than one common positive ion. We do not call the Na^+ ion the sodium(I) ion because (I) is unnecessary. Sodium forms only a 1+ ion, so there is no ambiguity about the name *sodium ion*.

Table **2.23.1**: The Modern and Common System of Cation Names

Element	Stem	Charge	Modern Name	Common Name
iron	ferr-	2+	iron(II) ion	ferrous ion
		3+	iron(III) ion	ferric ion
copper	cupr-	1+	copper(I) ion	cuprous ion
		2+	copper(II) ion	cupric ion
tin	stann-	2+	tin(II) ion	stannous ion
		4+	tin(IV) ion	stannic ion
lead	plumb-	2+	lead(II) ion	plumbous ion
		4+	lead(IV) ion	plumbic ion
chromium	chrom-	2+	chromium(II) ion	chromous ion
		3+	chromium(III) ion	chromic ion
gold	aur-	1+	gold(I) ion	aurous ion
		3+	gold(III) ion	auric ion

The second system, called the **common system**, is not conventional but is still prevalent and used in the health sciences. This system recognizes that many metals have two common cations. The common system uses two suffixes (*-ic* and *-ous*) that are appended to the stem of the element name. The *-ic* suffix represents the greater of the two cation charges, and the *-ous* suffix represents the lower one. In many cases, the stem of the element name comes from the Latin name of the element. Table **2.23.1** lists the elements that use the common system, along with their respective cation names.

Table **2.23.2**: Some Monatomic Anions

Ion	Name
F^-	fluoride ion
Cl^-	chloride ion
Br^-	bromide ion
I^-	iodide ion
O^{2-}	oxide ion
S^{2-}	sulfide ion
P^{3-}	phosphide ion
N^{3-}	nitride ion

The name of a monatomic anion consists of the stem of the element name, the suffix *-ide*, and then the word *ion*. Thus, as we have already seen, Cl^- is "chlor-" + "-ide ion," or the chloride ion. Similarly, O^{2-} is the oxide ion, Se^{2-} is the selenide ion, and so forth. Table **2.23.2** lists the names of some common monatomic ions. The polyatomic ions have their own characteristic names, as discussed earlier.

Example 2.23.1

Name each ion.

a. Ca^{2+}
b. S^{2-}
c. SO_3^{2-}
d. NH_4^+
e. Cu^+

Solution

a. the calcium ion
b. the sulfide ion
c. the sulfite ion
d. the ammonium ion
e. the copper(I) ion or the cuprous ion

Exercise 2.23.1

Name each ion.

a. Fe^{2+}
b. Fe^{3+}
c. SO_4^{2-}
d. Ba^{2+}
e. HCO_3^-

Answer a:

iron(II) ion

Answer b:

iron(III) ion

Answer c:

sulfate ion

Answer d:

barium ion

Answer e:

hydrogen carbonate ion or bicarbonate ion

Example 2.23.2

Write the formula for each ion.

a. the bromide ion
b. the phosphate ion
c. the cupric ion
d. the magnesium ion

Solution

a. Br^-
b. PO_4^{3-}
c. Cu^{2+}
d. Mg^{2+}

Exercise 2.23.2

Write the formula for each ion.

a. the fluoride ion
b. the carbonate ion
c. the stannous ion
d. the potassium ion

Answer a:

F^-

Answer b:

CO_3^{2-}

Answer c:

Sn^{2+}

Answer d:

K^+

Naming Binary Ionic Compounds with a Metal that Forms Only One Type of Cation

A **binary** ionic compound is a compound composed of a **monatomic** metal **cation** and a monatomic nonmetal **anion**. The metal cation is named first, followed by the nonmetal anion as illustrated in Figure **2.23.1** for the compound $BaCl_2$. The word *ion* is dropped from both parts.




Figure **2.23.1**: Naming $BaCl_2$
Naming formula: Name of metal cation + base
name of nonmetal anion + suffix -ide. BaCl2 is named as barium chloride.

Subscripts in the formula do not affect the name.

✔ Example 2.23.3: Naming Ionic Compounds

Name each ionic compound.

a. $CaCl_2$
b. AlF_3
c. KCl

Solution

a. Using the names of the ions, this ionic compound is named calcium chloride.
b. The name of this ionic compound is aluminum fluoride.
c. The name of this ionic compound is potassium chloride

? Exercise 2.23.3

Name each ionic compound.

a. AgI
b. MgO
c. Ca_3P_2

Answer a:
silver iodide
Answer b:
magnesium oxide
Answer c:
calcium phosphide


Naming Binary Ionic Compounds with a Metal That Forms More Than One Type of Cation

If you are given a formula for an ionic compound whose cation can have more than one possible charge, you must first determine the charge on the cation before identifying its correct name. For example, consider $FeCl_2$ and $FeCl_3$. In the first compound, the iron

ion has a 2+ charge because there are two Cl^- ions in the formula (1− charge on each chloride ion). In the second compound, the iron ion has a 3+ charge, as indicated by the three Cl^- ions in the formula. These are two different compounds that need two different names. By the Stock system, the names are iron(II) chloride and iron(III) chloride (Figure **2.23.2**).

Table **2.23.3**: Naming the $FeCl_2$ and $FeCl_3$ Compounds in the Modern/Stock System.


Name of cation (metal) + (Roman Numeral in parenthesis) + Base name of anion (nonmetal) and *-ide*



Remove the word "ion"

iron (II) + chloride

iron (II) chloride



Remove the word "ion"

iron (III) + chloride


iron (III) chloride

If we were to use the stems and suffixes of the common system, the names would be ferrous chloride and ferric chloride, respectively (Figure **2.23.3**) .

Table **2.23.4**: Naming the $FeCl_2$ and $FeCl_3$ Compounds in the Old/Common System.

"Old" base name of cation (metal) and *-ic or -ous* + Base name of anion (nonmetal) and *-ide*

-ous (for ions with lower charge)




Remove the word "ion"

ferrous + chloride

ferrous chloride

-ic (for ions with higher charge)



Remove the word "ion"

ferric + chloride

ferric chloride

Example 2.23.4:

Name each ionic compound.

a. Co_2O_3
b. FeCl2

Solution

Solutions to Example 5.7.4

	Explanation	Answer
a	We know that cobalt can have more than one possible charge; we just need to determine what it is. • Oxide always has a 2− charge, so with three oxide ions, we have a total negative charge of 6−. • This means that the two cobalt ions have to contribute 6+, which for two cobalt ions means that each one is 3+. • Therefore, the proper name for this ionic compound is cobalt(III) oxide.	cobalt(III) oxide
b	Iron can also have more than one possible charge. • Chloride always has a 1− charge, so with two chloride ions, we have a total negative charge of 2−. • This means that the one iron ion must have a 2+ charge. • Therefore, the proper name for this ionic compound is iron(II) chloride.	iron(II) chloride

Exercise 2.23.4

Name each ionic compound.

a. $AuCl_3$
b. PbO_2
c. CuO

Answer a:

gold(III) chloride

Answer b:

lead(IV) oxide

Answer c:

copper(II) oxide

Naming Ionic Compounds with Polyatomic Ions

The process of naming ionic compounds with polyatomic ions is the same as naming binary ionic compounds. The cation is named first, followed by the anion. One example is the ammonium sulfate compound in Figure **2.23.6**.




Figure **2.23.2**: Naming Ionic Compounds with Polyatomic Ions

Example 2.23.5: Naming Ionic Compounds

Write the proper name for each ionic compound.

a. $(NH_4)_2S$
b. $AlPO_4$,
c. $Fe_3(PO_4)_2$

Solution

Solutions to Example 5.7.5

Explanation	Answer
a. The ammonium ion has a 1+ charge and the sulfide ion has a 2− charge. Two ammonium ions need to balance the charge on a single sulfide ion. The compound's name is ammonium sulfide.	ammonium sulfide
b. The ions have the same magnitude of charge, one of each (ion) is needed to balance the charges. The name of the compound is aluminum phosphate.	aluminum phosphate
c. Neither charge is an exact multiple of the other, so we have to go to the least common multiple of 6. To get 6+, three iron(II) ions are needed, and to get 6−, two phosphate ions are needed . The compound's name is iron(II) phosphate.	iron(II) phosphate

Exercise 2.23.5A

Write the proper name for each ionic compound.

a. $(NH_4)_3PO_4$
b. $Co(NO_2)_3$

Answer a:

ammonium phosphate

Answer b:

cobalt(III) nitrite

Figure **2.23.1** is a synopsis of how to name simple ionic compounds.




Figure **2.23.3**: A Guide to Naming Simple Ionic Compounds.

? Exercise 2.23.5*B*

Name each ionic compound.

a. $ZnBr_2$
b. Al_2O_3
c. $(NH_4)_3PO_4$
d. AuF_3
e. AgF

Answer a:

zinc bromide

Answer b:

aluminum oxide

Answer c:

ammonium phosphate

Answer d:

gold(III) fluoride or auric fluoride

Answer e:

silver fluoride

Summary

- Ionic compounds are named by stating the cation first, followed by the anion.
- Positive and negative charges must balance.
- Some anions have multiple forms and are named accordingly with the use of roman numerals in parentheses.
- Ternary compounds are composed of three or more elements.

2.23: Naming Ionic Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.7: Naming Ionic Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.24: Naming Molecular Compounds

Learning Objectives

- Determine the name of a simple molecular compound from its chemical formula.

Molecular Compounds

Molecular compounds are inorganic compounds that take the form of discrete **molecules**. Examples include such familiar substances as water (H_2O) and carbon dioxide (CO_2). These compounds are very different from ionic compounds like sodium chloride ($NaCl$). Ionic compounds are formed when metal atoms lose one or more of their electrons to nonmetal atoms. The resulting cations and anions are electrostatically attracted to each other.

So what holds the atoms of a molecule together? Rather than forming ions, the atoms of a molecule share their **electrons** in such a way that a **bond** forms between a pair of atoms. In a carbon dioxide molecule, there are two of these bonds, each occurring between the carbon atom and one of the two oxygen atoms.




Figure 2.24.1: Carbon dioxide molecules consist of a central carbon atom bonded to 2 oxygen atoms.

Larger molecules can have many, many bonds that serve to keep the molecule together. In a large sample of a given molecular compound, all of the individual molecules are identical.

Naming Binary Molecular Compounds

Recall that a molecular formula shows the number of atoms of each element that a molecule contains. A molecule of water contains two hydrogen atoms and one oxygen atom, so its formula is H_2O. A molecule of octane, which is a component of gasoline, contains 8 atoms of carbon and 18 atoms of hydrogen. The molecular formula of octane is C_8H_{18}.




Figure 2.24.2: Nitrogen dioxide (NO_2) is a reddish-brown toxic gas that is a prominent air pollutant produced by internal combustion engines.

Naming *binary* (two-element) molecular compounds is similar to naming simple ionic compounds. The first element in the formula is simply listed using the name of the element. The second element is named by taking the stem of the element name and adding the suffix *-ide*. A system of numerical prefixes is used to specify the number of atoms in a molecule. Table 2.24.1 lists these numerical prefixes.

Table 2.24.1: Numerical Prefixes for Naming Binary Covalent Compounds

Number of Atoms in Compound	Prefix on the Name of the Element
1	mono-*
2	di-
3	tri-
4	tetra-
5	penta-
6	hexa-
7	hepta-
8	octa-
9	nona-
10	deca-

*This prefix is not used for the first element's name.



Note

- Generally, the less electronegative element is written first in the formula, though there are a few exceptions. Carbon is always first in a formula and hydrogen is after nitrogen in a formula such as NH_3. The order of common nonmetals in binary compound formulas is C, P, N, H, S, I, Br, Cl, O, F.
- The *a* or *o* at the end of a prefix is usually dropped from the name when the name of the element begins with a vowel. As an example, four oxygen atoms, is tetr**o**xide instead of tetr**ao**xide.
- The prefix is "mono" is not added to the first element's name if there is only one atom of the first element in a molecule.

Some examples of molecular compounds are listed in Table **2.24.2**.

Table **2.24.2**

Formula	Name
NO	nitrogen monoxide
N_2O	dinitrogen monoxide
S_2Cl_2	disulfur dichloride
Cl_2O_7	dichlorine heptoxide

Notice that the *mono-* prefix is not used with the nitrogen in the first compound, but is used with the oxygen in both of the first two examples. The S_2Cl_2 emphasizes that the formulas for molecular compounds are not reduced to their lowest ratios. The *o* of the *mono-* and the *a* of *hepta-* are dropped from the name when paired with oxide.

? Exercise 2.24.1

Write the name for each compound.

a. CF_4
b. $SeCl_2$
c. SO_3

Answer a:

carbon tetrafluoride

Answer b:

selenium dichloride

Answer c:

sulfur trioxide

Simple molecular compounds with common names

For some simple covalent compounds, we use common names rather than systematic names. We have already encountered these compounds, but we list them here explicitly:

- H_2O: water
- NH_3: ammonia
- CH_4: methane
- H_2O_2: hydrogen peroxide

Methane is the simplest organic compound. Organic compounds are compounds with carbon atoms and are named by a separate nomenclature system.

Some Compounds Have Both Covalent and Ionic Bonds

If you recall the introduction of polyatomic ions, you will remember that the bonds that hold the polyatomic ions together are covalent bonds. Once the polyatomic ion is constructed with covalent bonds, it reacts with other substances as an ion. The bond between a polyatomic ion and another ion will be ionic. An example of this type of situation is in the compound sodium nitrate. Sodium nitrate is composed of a sodium ion and a nitrate ion. The nitrate ion is held together by covalent bonds and the nitrate ion is attached to the sodium ion by an ionic bond.

Summary

- A molecular compound is usually composed of two or more nonmetal elements.
- Molecular compounds are named with the first element first and then the second element by using the stem of the element name plus the suffix -ide. Numerical prefixes are used to specify the number of atoms in a molecule.

2.24: Naming Molecular Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.8: Naming Molecular Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.25: Naming Acids

A spot test for gold has been in use for decades. The sample is first treated with nitric acid. Other metals may react or dissolve in this acid, but gold will not. Then the sample is added to a mixture of nitric acid and hydrochloric acid. Gold will only dissolve in this mixture. The term "acid test" arose from the California gold rush in the late 1840's when this combination was used to test for the presence of real gold. It has since come to mean, "tested and approved" in a number of fields.

Acids

An **acid** can be defined in several ways. The most straightforward definition is that an acid is a molecular compound that contains one or more hydrogen atoms and produces hydrogen ions ($\mathbf{H^+}$) when dissolved in water.



Figure **2.25.1**: (A) Vinegar comes in a variety of types, but all contain acetic acid. (B) Citrus fruits like grapefruit contain citric and ascorbic acids.

This is a different type of compound than the others we have seen so far. Acids are molecular, which means that in their pure state they are individual molecules and do not adopt the extended three-dimensional structures of ionic compounds like $\mathbf{NaCl}$. However, when these molecules are dissolved in water, the chemical bond between the hydrogen atom and the rest of the molecule breaks, leaving a positively-charged hydrogen ion and an anion. This can be symbolized in a chemical equation:

$$\mathbf{HCl \rightarrow H^+ + Cl^-}$$


Since acids produce $\mathbf{H^+}$ cations upon dissolving in water, the $\mathbf{H}$ of an acid is written first in the formula of an inorganic acid. The remainder of the acid (other than the $\mathbf{H}$) is the anion after the acid dissolves. Organic acids are also an important class of compounds, but will not be discussed here.

Naming Acids

Since all acids contain hydrogen, the name of an acid is based on the anion that goes with it. These anions can either be monatomic or polyatomic.

Naming Binary acids (in aqueous form)

A **binary acid** is an acid that consists of hydrogen and one other element. The most common binary acids contain a halogen. The acid name begins with the prefix *hydro-*. followed by the base name of the anion, followed by the suffix *-ic*.




Formula for naming acids: Hydro- and Base name of nonmetal and -ic + acid. Example: HCl is hydrochloric acid.

Naming Oxyacids

An **oxyacid** is an acid that consists of hydrogen, oxygen, and a third element. The third element is usually a nonmetal.

a. Oxyanions with ***-ite*** ending.


The name of the acid is the root of the anion followed by the suffix ***-ous***. There is no prefix.



Formula for naming oxyanions with -ite ending: Base name of oxyanion and -ous + acid. Example: H2SO3 is sulfurous acid.

b. Oxyanions with ***-ate*** ending.

The name of the acid is the root of the anion followed by the suffix ***-ic***. There is no prefix.




Formula for naming oxyanions with -ate ending: Base name of oxyanion and -ic + acid. Example: H3PO4 is phosphoric acid.

> **Note**
>
> The base name for sulfur containing oxyacid is ***sulfur-*** instead of just ***sulf-***. The same is true for a phosphorus containing oxyacid. The base name is ***phosphor-*** instead of simply ***phosph-***.

Writing Formulas for Acids

Like other compounds that we have studied, acids are electrically neutral. Therefore, the charge of the anion part of the formula must be exactly balanced out by the H^+ ions. Another way to think about writing the correct formula is to utilize the crisscross method, shown below for sulfuric acid.



H2SO4: H has +1 charge and SO4 has -2 charge, so there must be 2 H+ and 1 SO4 for the charges to balance out.

Formula: H_2SO_4

Figure 2.25.2: *Crisscross approach to writing formula for sulfuric acid.*

Summary


- Acids are molecular compounds that release hydrogen ions.
- A binary acid consists of hydrogen and one other element.
- Oxyacids contain hydrogen, oxygen, and one other element.

- The name of the acid is based on the anion attached to the hydrogen.

2.25: Naming Acids is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.9: Naming Acids** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

2.26: Nomenclature Summary



Contributions & Attributions

2.26: Nomenclature Summary is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **5.10: Nomenclature Summary** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

3: Unit 3

3.1: Evidence of a Chemical Reaction
3.2: The Chemical Equation
3.3: How to Write Balanced Chemical Equations
3.4: Classifying Chemical Reactions
3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water
3.6: Precipitation Reactions
3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
3.8: Prelude to Chemical Composition - How Much Sodium?
3.9: Counting Nails by the Pound
3.10: Counting Atoms by the Gram
3.11: Counting Molecules by the Gram
3.12: Chemical Formulas as Conversion Factors
3.13: Mass Percent Composition of Compounds
3.14: Mass Percent Composition from a Chemical Formula
3.15: Calculating Empirical Formulas for Compounds
3.16: Calculating Molecular Formulas for Compounds
3.17: Making Pancakes- Relationships Between Ingredients
3.18: Making Molecules- Mole-to-Mole Conversions
3.19: Making Molecules- Mass-to-Mass Conversions
3.20: Stoichiometry
3.21: Limiting Reactant and Theoretical Yield
3.22: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants

3: Unit 3 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

3.1: Evidence of a Chemical Reaction

Learning Objectives

- Identify the evidence for chemical reactions.

In a chemical change, new substances are formed. In order for this to occur, the chemical bonds of the substances break, and the atoms that compose them separate and rearrange themselves into new substances with new chemical bonds. When this process occurs, we call it a chemical reaction. A **chemical reaction** is the process in which one or more substances are changed into one or more new substances.




Figure **3.1.1**: (a) Copper and nitric acid undergo a chemical change to form copper nitrate and brown, gaseous nitrogen dioxide. (b) During the combustion of a match, cellulose in the match and oxygen from the air undergo a chemical change to form carbon dioxide and water vapor. (c) Cooking red meat causes a number of chemical changes, including the oxidation of iron in myoglobin that results in the familiar red-to-brown color change. (d) A banana turning brown is a chemical change as new, darker (and less tasty) substances form. (Credit b: modification of work by Jeff Turner; credit c: modification of work by Gloria Cabada-Leman; credit d: modification of work by Roberto Verzo.)

To identify a chemical reaction, we look for a **chemical change**. A chemical change always produces one or more types of matter that differ from the matter present before the change. The formation of rust is a chemical change because rust is a different kind of matter than the iron, oxygen, and water present before the rust formed. The explosion of nitroglycerin is a chemical change because the gases produced are very different kinds of matter from the original substance. Other examples of chemical changes include: reactions that are performed in a lab (such as copper reacting with nitric acid), all forms of combustion (burning), and food being cooked, digested, or rotting (Figure **3.1.1**).



***Video*3.1.1*:** *Evidence of a Chemical Reaction*

Example 3.1.1: Evidence of a Chemical Reaction

Which of the following is a chemical reaction?

a. Freezing liquid mercury.
b. Adding yellow to blue to make green.
c. Cutting a piece of paper into two pieces.
d. Dropping a sliced orange into a vat of sodium dydroxide.
e. Filling a balloon with natural air.

Solution

A, B, C, & E involve only physical changes. A sliced orange has acid (citric acid) that can react with sodium hydroxide, so the answer is D.

Exercise 3.1.1

Which of the following is a chemical reaction?

a. Painting a wall blue.
b. A bicycle rusting.
c. Ice cream melting.
d. Scratching a key across a desk.
e. Making a sand castle.

Answer

B

Example 3.1.2: Evidence of a Chemical Reaction

Which of the following is not a chemical reaction?

a. Shattering glass with a baseball.
b. Corroding metal.
c. Fireworks exploding.
d. Lighting a match.
e. Baking a cake.

Solution

Shattering glass with a baseball results in glass broken into many pieces but no chemical change happens, so the answer is A.

? Exercise 3.1.2

Which of the following is NOT a chemical reaction?

a. Frying an egg.
b. Slicing carrots.
c. A Macbook falling out of a window.
d. Creating ATP in the human body.
e. Dropping a fizzy tablet into a glass of water.

Answer

B and C

Summary

Chemical reactions can be identified via a wide range of different observable factors including change in color, energy change (temperature change or light produced), gas production, formation of precipitate and change in properties.

This page titled 3.1: Evidence of a Chemical Reaction is shared under a CC BY 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew (OpenStax) via source content that was edited to the style and standards of the LibreTexts platform.

- **7.2: Evidence of a Chemical Reaction** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY 4.0. Original source: https://www.ck12.org/c/chemistry/.

3.2: The Chemical Equation

Learning Objectives

- Identify the reactants and products in any chemical reaction.
- Convert word equations into chemical equations.
- Use the common symbols, (s), (l), (g), (aq), and $\rightarrow$ appropriately when writing a chemical reaction.

In a chemical change, new substances are formed. In order for this to occur, the chemical bonds of the substances break, and the atoms that compose them separate and rearrange themselves into new substances with new chemical bonds. When this process occurs, we call it a chemical reaction. A **chemical reaction** is the process in which one or more substances are changed into one or more new substances.

Reactants and Products

To describe a chemical reaction, we need to indicate what substances are present at the beginning and what substances are present at the end. The substances that are present at the beginning are called **reactants** and the substances present at the end are called **products**.

Sometimes when reactants are put into a reaction vessel, a reaction will take place to produce products. Reactants are the starting materials, that is, whatever we have as our initial ingredients. The products are just that—what is produced—or the result of what happens to the reactants when we put them together in the reaction vessel. If we think about baking chocolate chip cookies, our reactants would be flour, butter, sugar, vanilla, baking soda, salt, egg, and chocolate chips. What would be the products? Cookies! The reaction vessel would be our mixing bowl.

$$\underbrace{\text{Flour} + \text{Butter} + \text{Sugar} + \text{Vanilla} + \text{Baking Soda} + \text{Eggs} + \text{Chocolate Chips}}_{\text{Ingredients = Reactants}} \rightarrow \underbrace{\text{Cookies}}_{\text{Product}}$$

Writing Chemical Equations

When sulfur dioxide is added to oxygen, sulfur trioxide is produced. Sulfur dioxide and oxygen, $SO_2 + O_2$, are reactants and sulfur trioxide, SO_3, is the product.

$$\underbrace{2\,SO_2\,(g) + O_2\,(g)}_{\text{Reactants}} \rightarrow \underbrace{2\,SO_3\,(g)}_{\text{Products}}$$

In chemical reactions, the reactants are found before the symbol "$\rightarrow$" and the products are found after the symbol "$\rightarrow$". The general equation for a reaction is:

$$\text{Reactants} \rightarrow \text{Products}$$


There are a few special symbols that we need to know in order to "talk" in chemical shorthand. In the table below is the summary of the major symbols used in chemical equations. Table **3.2.1** shows a listing of symbols used in chemical equations.

Table **3.2.1**: Symbols Used in Chemical Equations

Symbol	Description	Symbol	Description
$+$	used to separate multiple reactants or products	(s)	reactant or product in the solid state
$\rightarrow$	yield sign; separates reactants from products	(l)	reactant or product in the liquid state
$\rightleftharpoons$	replaces the yield sign for reversible reactions that reach equilibrium	(g)	reactant or product in the gas state
$\xrightarrow{\text{Pt}}$	formula written above the arrow is used as a catalyst in the reaction	(aq)	reactant or product in an aqueous solution (dissolved in water)
$\xrightarrow{\Delta}$	triangle indicates that the reaction is being heated		

Chemists have a choice of methods for describing a chemical reaction.

1. They could draw a picture of the chemical reaction.



2. They could write a word equation for the chemical reaction:
"Two molecules of hydrogen gas react with one molecule of oxygen gas to produce two molecules of water vapor."

3. They could write the equation in chemical shorthand.

$$2\text{H}_2\,(g) + \text{O}_2\,(g) \rightarrow 2\text{H}_2\text{O}\,(g)$$

In the symbolic equation, chemical formulas are used instead of chemical names for reactants and products, while symbols are used to indicate the phase of each substance. It should be apparent that the chemical shorthand method is the quickest and clearest method for writing chemical equations.

We could write that an aqueous solution of calcium nitrate is added to an aqueous solution of sodium hydroxide to produce solid calcium hydroxide and an aqueous solution of sodium nitrate. Or in shorthand we could write:

$$\text{Ca}(\text{NO}_3)_2\,(aq) + 2\text{NaOH}\,(aq) \rightarrow \text{Ca}(\text{OH})_2\,(s) + 2\text{NaNO}_3\,(aq)$$

How much easier is that to read? Let's try it in reverse. Look at the following reaction in shorthand and write the word equation for the reaction:

$$\text{Cu}\,(s) + \text{AgNO}_3\,(aq) \rightarrow \text{Cu}(\text{NO}_3)_2\,(aq) + \text{Ag}\,(s)$$

The word equation for this reaction might read something like "solid copper reacts with an aqueous solution of silver nitrate to produce a solution of copper (II) nitrate with solid silver."

To turn word equations into symbolic equations, we need to follow the given steps:

1. Identify the reactants and products. This will help you know which symbols go on each side of the arrow and where the $+$ signs go.
2. Write the correct formulas for all compounds. You will need to use the rules you learned in Chapter 5 (including making all ionic compounds charge balanced).

3. Write the correct formulas for all elements. Usually this is given straight off of the periodic table. However, there are seven elements that are considered diatomic, meaning that they are always found in pairs in nature. They include those elements listed in the table.

Table **3.2.1**: *Diatomic Elements*

Element Name	Hydrogen	Nitrogen	Oxygen	Fluorine	Chlorine	Bromine	Iodine
Formula	H_2	N_2	O_2	F_2	Cl_2	Br_2	I_2

Example 3.2.1

Transfer the following symbolic equations into word equations or word equations into symbolic equations.

a. $\mathrm{HCl}\,(aq) + \mathrm{NaOH}\,(aq) \rightarrow \mathrm{NaCl}\,(aq) + \mathrm{H_2O}\,(l)$

b. Gaseous propane, $\mathrm{C_3H_8}$, burns in oxygen gas to produce gaseous carbon dioxide and liquid water.

c. Hydrogen fluoride gas reacts with an aqueous solution of potassium carbonate to produce an aqueous solution of potassium fluoride, liquid water, and gaseous carbon dioxide.

Solution

a. An aqueous solution of hydrochloric acid reacts with an aqueous solution of sodium hydroxide to produce an aqueous solution of sodium chloride and liquid water.

b. Reactants: propane ($\mathrm{C_3H_8}$) and oxygen ($\mathrm{O_2}$)

Product: carbon dioxide ($\mathrm{CO_2}$) and water ($\mathrm{H_2O}$)

$$\mathrm{C_3H_8}\,(g) + \mathrm{O_2}\,(g) \rightarrow \mathrm{CO_2}\,(g) + \mathrm{H_2O}\,(l)$$

c. Reactants: hydrogen fluoride and potassium carbonate

Products: potassium fluoride, water, and carbon dioxide

$$\mathrm{HF}\,(g) + \mathrm{K_2CO_3}\,(aq) \rightarrow \mathrm{KF}\,(aq) + \mathrm{H_2O}\,(l) + \mathrm{CO_2}\,(g)$$

Exercise 3.2.1

Transfer the following symbolic equations into word equations or word equations into symbolic equations.

a. Hydrogen gas reacts with nitrogen gas to produce gaseous ammonia.

b. $\mathrm{HCl}\,(aq) + \mathrm{LiOH}\,(aq) \rightarrow \mathrm{LiCl}\,(aq) + \mathrm{H_2O}\,(l)$

c. Copper metal is heated with oxygen gas to produce solid copper(II) oxide.

Answer a

$H_2(g) + N_2(g) \rightarrow NH_3(g)$

Answer b

An aqueous solution of hydrochloric acid reacts with an aqueous solution of lithium hydroxide to produce an aqueous solution of lithium chloride and liquid water.

Answer c

$Cu(s) + O_2(g) \rightarrow CuO(s)$

Summary

- A chemical reaction is the process by which one or more substances are changed into one or more new substances.
- Chemical reactions are represented by chemical equations.
- Chemical equations have reactants on the left, an arrow that is read as "yields", and the products on the right.

3.2: The Chemical Equation is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.3: The Chemical Equation** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.3: How to Write Balanced Chemical Equations

Learning Objectives

- Explain the roles of subscripts and coefficients in chemical equations.
- Balance a chemical equation when given the unbalanced equation.
- Explain the role of the Law of Conservation of Mass in a chemical reaction.

Even though chemical compounds are broken up and new compounds are formed during a chemical reaction, atoms in the reactants do not disappear, nor do new atoms appear to form the products. In chemical reactions, atoms are never created or destroyed. The same atoms that were present in the reactants are present in the products—they are merely reorganized into different arrangements. In a complete chemical equation, the two sides of the equation must be present on the reactant and the product sides of the equation.

Coefficients and Subscripts

There are two types of numbers that appear in chemical equations. There are subscripts, which are part of the chemical formulas of the reactants and products; and there are coefficients that are placed in front of the formulas to indicate how many molecules of that substance is used or produced.





Figure **3.3.1**: Balancing Equations. You cannot change subscripts in a chemical formula to balance a chemical equation; you can change only the coefficients. Changing subscripts changes the ratios of atoms in the molecule and the resulting chemical properties. For example, water (H_2O) and hydrogen peroxide (H_2O_2) are chemically distinct substances. H_2O_2 decomposes to H_2O and O_2 gas when it comes in contact with the metal platinum, whereas no such reaction occurs between water and platinum.

The **subscripts** are part of the formulas and once the formulas for the reactants and products are determined, the subscripts may not be changed. The **coefficients** indicate the number of each substance involved in the reaction and may be changed in order to balance the equation. The equation above indicates that one mole of solid copper is reacting with two moles of aqueous silver nitrate to produce one mole of aqueous copper (II) nitrate and two atoms of solid silver.

Balancing a Chemical Equation

Because the identities of the reactants and products are fixed, the equation cannot be balanced by changing the subscripts of the reactants or the products. To do so would change the chemical identity of the species being described, as illustrated in Figure **3.3.1**.



Original molecule $\mathbf{H_2O}$: if the coefficient 2 is added in front, that makes 2 water molecules; but if the subscript 2 is added to make $\mathbf{H_2O_2}$, that's hydrogen peroxide.

The simplest and most generally useful method for balancing chemical equations is “inspection,” better known as trial and error. The following is an efficient approach to balancing a chemical equation using this method.

Steps in Balancing a Chemical Equation

1. Identify the most complex substance.
2. Beginning with that substance, choose an element(s) that appears in only one reactant and one product, if possible. Adjust the coefficients to obtain the same number of atoms of this element(s) on both sides.
3. Balance polyatomic ions (if present on both sides of the chemical equation) as a unit.
4. Balance the remaining atoms, usually ending with the least complex substance and using fractional coefficients if necessary. If a fractional coefficient has been used, multiply both sides of the equation by the denominator to obtain whole numbers for the coefficients.
5. Count the numbers of atoms of each kind on both sides of the equation to be sure that the chemical equation is balanced.

Example 3.3.1: Combustion of Heptane

Balance the chemical equation for the combustion of Heptane (C_7H_{16}).

$$C_7H_{16}(l) + O_2(g) \longrightarrow CO_2(g) + H_2O(g)$$

Solution

Solutions to Example **3.3.1**

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is the one with the largest number of different atoms, which is C_7H_{16}. We will assume initially that the final balanced chemical equation contains 1 molecule or formula unit of this substance.
2. **Adjust the coefficients.**	a. Because one molecule of n-heptane contains 7 carbon atoms, we need 7 CO_2 molecules, each of which contains 1 carbon atom, on the right side: $$C_7H_{16}(l) + O_2(g) \longrightarrow \underline{7}CO_2(g) + H_2O(g)$$ • 7 carbon atoms on both reactant and product sides b. Because one molecule of n-heptane contains 16 hydrogen atoms, we need 8 H_2O molecules, each of which contains 2 hydrogen atoms, on the right side: $$C_7H_{16}(l) + O_2(g) \longrightarrow 7\,CO_2(g) + \underline{8}H_2O(g)$$ • 16 hydrogen atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	There are no polyatomic ions to be considered in this reaction.
4. **Balance the remaining atoms.**	The carbon and hydrogen atoms are now balanced, but we have 22 oxygen atoms on the right side and only 2 oxygen atoms on the left. We can balance the oxygen atoms by adjusting the coefficient in front of the least complex substance, O_2, on the reactant side: $$C_7H_{16}(l) + \underline{11}O_2(g) \longrightarrow 7\,CO_2(g) + 8\,H_2O(g)$$ • 22 oxygen atoms on both reactant and product sides
5. **Check your work.**	The equation is now balanced, and there are no fractional coefficients: there are 7 carbon atoms, 16 hydrogen atoms, and 22 oxygen atoms on each side. Always check to be sure that a chemical equation is balanced.

Example 3.3.2: Combustion of Isooctane

Combustion of Isooctane (C_8H_{18})

$$C_8H_{18}\,(l) + O_2\,(g) \longrightarrow CO_2\,(g) + H_2O\,(g)$$

Solution

The assumption that the final balanced chemical equation contains only one molecule or formula unit of the most complex substance is not always valid, but it is a good place to start. The combustion of any hydrocarbon with oxygen produces carbon dioxide and water.

Solutions to Example 7.4.2

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is the one with the largest number of different atoms, which is C_8H_{18}. We will assume initially that the final balanced chemical equation contains 1 molecule or formula unit of this substance.
2. **Adjust the coefficients.**	a. The first element that appears only once in the reactants is carbon: 8 carbon atoms in isooctane means that there must be 8 CO_2 molecules in the products: $$C_8H_{18}(l) + O_2(g) \longrightarrow \underline{8}CO_2(g) + H_2O(g)$$ • 8 carbon atoms on both reactant and product sides b. 18 hydrogen atoms in isooctane means that there must be 9 H_2O molecules in the products: $$C_8H_{18}(l) + O_2(g) \longrightarrow 8\,CO_2(g) + \underline{9}H_2O(g)$$ • 18 hydrogen atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	There are no polyatomic ions to be considered in this reaction.
4. **Balance the remaining atoms.**	The carbon and hydrogen atoms are now balanced, but we have 25 oxygen atoms on the right side and only 2 oxygen atoms on the left. We can balance the least complex substance, O_2, but because there are 2 oxygen atoms per O_2 molecule, we must use a fractional coefficient ($\frac{25}{2}$) to balance the oxygen atoms: $$C_8H_{18}(l) + \underline{\frac{25}{2}}O_2\ (g) \longrightarrow 8\,CO_2(g) + 9\,H_2O(g)$$ • 25 oxygen atoms on both reactant and product sides The equation is now balanced, but we usually write equations with whole number coefficients. We can eliminate the fractional coefficient by multiplying all coefficients on both sides of the chemical equation by **2**: $$\underline{2}C_8H_{18}(l) + \underline{25}O_2(g) \longrightarrow \underline{16}CO_2(g) + \underline{18}H_2O(g)$$
5. **Check your work.**	The balanced chemical equation has 16 carbon atoms, 36 hydrogen atoms, and 50 oxygen atoms on each side. Balancing equations requires some practice on your part as well as some common sense. If you find yourself using very large coefficients or if you have spent several minutes without success, go back and make sure that you have written the formulas of the reactants and products correctly.

✔ Example 3.3.3: Precipitation of Lead (II) Chloride

Aqueous solutions of lead (II) nitrate and sodium chloride are mixed. The products of the reaction are an aqueous solution of sodium nitrate and a solid precipitate of lead (II) chloride. Write the balanced chemical equation for this reaction.

Solution

Solutions to Example 7.4.3

Steps	Example
1. **Identify the most complex substance.**	The most complex substance is lead (II) chloride. $\mathrm{Pb(NO_3)_2\,(aq) + NaCl\,(aq) \longrightarrow NaNO_3\,(aq) + PbCl_2\,(s)}$
2. **Adjust the coefficients.**	There are twice as many chloride ions in the reactants as there are in the products. Place a 2 in front of the NaCl in order to balance the chloride ions. $\mathrm{Pb(NO_3)_2\,(aq) + \underline{2}\,NaCl\,(aq) \longrightarrow NaNO_3\,(aq) + PbCl_2\,(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on reactant side, 1 Na atom on product side • 2 Cl atoms on both reactant and product sides
3. **Balance polyatomic ions as a unit.**	The nitrate ions are still unbalanced. Place a 2 in front of the $NaNO_3$. The result is: $\mathrm{Pb(NO_3)_2\,(aq) + 2\,NaCl\,(aq) \longrightarrow \underline{2}\,NaNO_3\,(aq) + PbCl_2\,(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on both reactant and product sides • 2 Cl atoms on both reactant and product sides • 2 NO_3^- atoms on both reactant and product sides
4. **Balance the remaining atoms.**	There is no need to balance the remaining atoms because they are already balanced.
5. **Check your work.**	$\mathrm{Pb(NO_3)_2\,(aq) + 2\,NaCl\,(aq) \longrightarrow 2\,NaNO_3\,(aq) + PbCl_2\,(s)}$ • 1 Pb atom on both reactant and product sides • 2 Na atoms on both reactant and product sides • 2 Cl atoms on both reactant and product sides • 2 NO_3^- atoms on both reactant and product sides

? Exercise 3.3.1

Is each chemical equation balanced?

a. $\mathrm{2\,Hg(\ell) + O_2\,(g) \rightarrow Hg_2O_2\,(s)}$
b. $\mathrm{C_2H_4\,(g) + 2\,O_2(g) \longrightarrow 2\,CO_2\,(g) + 2\,H_2O\,(g)}$
c. $\mathrm{Mg(NO_3)_2\,(s) + 2\,Li\,(s) \rightarrow Mg(s) + 2\,LiNO_3\,(s)}$

Answer a

yes

Answer b

no

Answer c

yes

Exercise 3.3.2

Balance the following chemical equations.

a. $\mathbf{N_2\,(g) + O_2\,(g) \longrightarrow NO_2\,(g)}$
b. $\mathbf{Pb(NO_3)_2\,(aq) + FeCl_3\,(aq) \longrightarrow Fe(NO_3)_3\,(aq) + PbCl_2\,(s)}$
c. $\mathbf{C_6H_{14}\,(l) + O_2(g) \longrightarrow CO_2\,(g) + H_2O\,(g)}$

Answer a

$N_2\,(g) + 2O_2\,(g) \rightarrow 2NO_2\,(g)$

Answer b

$3Pb(NO_3)_2(aq) + 2FeCl_3(aq) \rightarrow 2Fe(NO_3)_3(aq) + 3PbCl_2(s)$

Answer c

$2C_6H_{14}(l) + 19O_2(g) \rightarrow 12CO_2(g) + 14H_2O(g)$

Summary

- To be useful, chemical equations must always be balanced. Balanced chemical equations have the same number and type of each atom on both sides of the equation.
- The coefficients in a balanced equation must be the simplest whole number ratio. Mass is always conserved in chemical reactions.

Vocabulary

- **Chemical reaction** - The process in which one or more substances are changed into one or more new substances.
- **Reactants** - The starting materials in a reaction.
- **Products** - Materials present at the end of a reaction.
- **Balanced chemical equation** - A chemical equation in which the number of each type of atom is equal on the two sides of the equation.
- **Subscripts** - Part of the chemical formulas of the reactants and products that indicate the number of atoms of the preceding element.
- **Coefficient** - A small whole number that appears in front of a formula in a balanced chemical equation.

3.3: How to Write Balanced Chemical Equations is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.4: How to Write Balanced Chemical Equations** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

3.4: Classifying Chemical Reactions

Learning Objectives

- Classify a chemical reaction as a synthesis, decomposition, single replacement, double replacement, or a combustion reaction.
- Predict the products of simple reactions.

The chemical reactions we have described are only a tiny sampling of the infinite number of chemical reactions possible. How do chemists cope with this overwhelming diversity? How do they predict which compounds will react with one another and what products will be formed? The key to success is to find useful ways to categorize reactions. Familiarity with a few basic types of reactions will help you to predict the products that form when certain kinds of compounds or elements come in contact.

Most chemical reactions can be classified into one or more of five basic types: acid–base reactions, exchange reactions, condensation reactions (and the reverse, cleavage reactions), and oxidation–reduction reactions. The general forms of these five kinds of reactions are summarized in Table **3.4.1**, along with examples of each. It is important to note, however, that many reactions can be assigned to more than one classification, as you will see in our discussion.

Table **3.4.1:** *Basic Types of Chemical Reactions*

Name of Reaction	General Form	Examples
Oxidation–Reduction (redox)	oxidant + reductant → reduced oxidant + oxidized reductant	$C_7H_{16}(l) + 11O_2(g) \rightarrow 7CO_2(g) + 8H_2O(g)$
Acid–Base	acid + base → salt	$NaOH(aq) + HNO_3(aq) \rightarrow NaNO_3(aq) + H_2O(l)$
Exchange: Single Replacement	AB + C → AC + B	$ZnCl_2(aq) + Mg(s) \rightarrow MgCl_2(aq) + Zn(s)$
Exchange: Double Replacement	AB + CD → AD + CB	$BaCl_2(aq) + Na_2SO_4(aq) \rightarrow BaSO_4(s) + 2NaCl(aq)$
Combination (Synthesis)	A + B → AB	$CO_2(g) + H_2O(l) \rightarrow H_2CO_3(aq)$ $N_2(g) + 2O_2(g) \rightarrow 2NO_2(g)$
Decomposition	AB → A + B	$CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$

The classification scheme is only for convenience; the same reaction can be classified in different ways, depending on which of its characteristics is most important. Oxidation–reduction reactions, in which there is a net transfer of electrons from one atom to another, and condensation reactions are discussed in this section. Acid–base reactions are one kind of exchange reaction—the formation of an insoluble salt, such as barium sulfate, when solutions of two soluble salts are mixed together.

Combination Reactions

A **combination reaction** is a reaction in which two or more substances combine to form a single new substance. Combination reactions can also be called synthesis reactions. The general form of a combination reaction is:

$$\mathbf{A + B \rightarrow AB}$$

One combination reaction is two elements combining to form a compound. Solid sodium metal reacts with chlorine gas to produce solid sodium chloride.

$$\mathbf{2Na\,(\mathit{s}) + Cl_2\,(\mathit{g}) \rightarrow 2NaCl\,(\mathit{s})}$$

Notice that in order to write and balance the equation correctly, it is important to remember the seven elements that exist in nature as diatomic molecules ($\mathbf{H_2}$, $\mathbf{N_2}$, $\mathbf{O_2}$, $\mathbf{F_2}$, $\mathbf{Cl_2}$, $\mathbf{Br_2}$, and $\mathbf{I_2}$).

One type of combination reaction that occurs frequently is the reaction of an element with oxygen to form an oxide. Metals and nonmetals both react readily with oxygen under most conditions. Magnesium reacts rapidly and dramatically when ignited, combining with oxygen from the air to produce a fine powder of magnesium oxide:

$$2Mg\,(s) + O_2\,(g) \rightarrow 2MgO\,(s)$$

Sulfur reacts with oxygen to form sulfur dioxide:

$$S\,(s) + O_2\,(g) \rightarrow SO_2\,(g)$$

When nonmetals react with one another, the product is a molecular compound. Often, the nonmetal reactants can combine in different ratios and produce different products. Sulfur can also combine with oxygen to form sulfur trioxide:

$$2S\,(s) + 3O_2\,(g) \rightarrow 2SO_3\,(g)$$

Transition metals are capable of adopting multiple positive charges within their ionic compounds. Therefore, most transition metals are capable of forming different products in a combination reaction. Iron reacts with oxygen to form both iron (II) oxide and iron (III) oxide:

$$2Fe\,(s) + O_2\,(g) \rightarrow 2FeO\,(s)$$

$$4Fe\,(s) + 3O_2\,(g) \rightarrow 2Fe_2O_3\,(s)$$

Example 3.4.1: Combustion of Solid Potassium

Potassium is a very reactive alkali metal that must be stored under oil in order to prevent it from reacting with air. Write the balanced chemical equation for the combination reaction of potassium with oxygen.

Solution

Solutions to Example 7.10.1

Steps	Example Solution
Plan the problem.	Make sure formulas of all reactants and products are correct before balancing the equation. Oxygen gas is a diatomic molecule. Potassium oxide is an ionic compound and so its formula is constructed by the crisscross method. Potassium as an ion becomes K^+, while the oxide ion is O^{2-}.
Solve.	The skeleton (unbalanced) equation: $$K\,(s) + O_2\,(g) \rightarrow K_2O\,(s)$$ The equation is then easily balanced with coefficients. $$4K\,(s) + O_2\,(g) \rightarrow 2K_2O\,(s)$$
Think about your result.	Formulas are correct and the resulting combination reaction is balanced.

Combination reactions can also take place when an element reacts with a compound to form a new compound composed of a larger number of atoms. Carbon monoxide reacts with oxygen to form carbon dioxide according to the equation:

$$2CO\,(g) + O_2\,(g) \rightarrow 2CO_2\,(g)$$

Two compounds may also react to form a more complex compound. A very common example is the reactions of oxides with water. Calcium oxide reacts readily with water to produce an aqueous solution of calcium hydroxide:

$$CaO\,(s) + H_2O\,(l) \rightarrow Ca(OH)_2\,(aq)$$

Sulfur trioxide gas reacts with water to form sulfuric acid. This is an unfortunately common reaction that occurs in the atmosphere in some places where oxides of sulfur are present as pollutants. The acid formed in the reaction falls to the ground as acid rain.

$$SO_3\,(g) + H_2O\,(l) \rightarrow H_2SO_4\,(aq)$$



Figure **3.4.1**: Acid rain has severe consequences on both natural and manmade objects. Acid rain degrades marble statues like the one on the left (A). The trees in the forest on the right (B) have been killed by acid rain.

? Exercise 3.4.1

a. Write the chemical equation for the synthesis of silver bromide, **AgBr**.
b. Predict the products for the following reaction: $CO_2\,(g) + H_2O\,(l)$

Answer a:

$2Ag + Br_2 \rightarrow 2AgBr$

Answer b:

$CO_2\,(g) + H_2O\,(l) \rightarrow H_2CO_3$

Decomposition Reactions

A **decomposition reaction** is a reaction in which a compound breaks down into two or more simpler substances. The general form of a decomposition reaction is:

$$AB \rightarrow A + B$$

Most decomposition reactions require an input of energy in the form of heat, light, or electricity.

Binary compounds are compounds composed of just two elements. The simplest kind of decomposition reaction is when a binary compound decomposes into its elements. Mercury (II) oxide, a red solid, decomposes when heated to produce mercury and oxygen gas:

$$2HgO\,(s) \rightarrow 2Hg\,(l) + O_2\,(g)$$



***Video* 3.4.2:** *Mercury (II) oxide is a red solid. When it is heated, it decomposes into mercury metal and oxygen gas.*

A reaction is also considered to be a decomposition reaction even when one or more of the products are still compounds. A metal carbonate decomposes into a metal oxide and carbon dioxide gas. For example, calcium carbonate decomposes into calcium oxide and carbon dioxide:

$$CaCO_3\,(s) \rightarrow CaO\,(s) + CO_2\,(g)$$

Metal hydroxides decompose on heating to yield metal oxides and water. Sodium hydroxide decomposes to produce sodium oxide and water:

$$2NaOH\,(s) \rightarrow Na_2O\,(s) + H_2O\,(g)$$

Some unstable acids decompose to produce nonmetal oxides and water. Carbonic acid decomposes easily at room temperature into carbon dioxide and water:

$$H_2CO_3\,(aq) \rightarrow CO_2\,(g) + H_2O\,(l)$$

Example 3.4.2: Electrolysis of Water

When an electric current is passed through pure water, it decomposes into its elements. Write a balanced equation for the decomposition of water.

Solution

Solutions to Example 7.10.2

Steps	Example Solution
Plan the problem.	Water is a binary compound composed of hydrogen and oxygen. The hydrogen and oxygen gases produced in the reaction are both diatomic molecules.
Solve.	The skeleton (unbalanced) equation: $$H_2O\,(l) \xrightarrow{elec} H_2\,(g) + O_2\,(g)$$ Note the abbreviation "**elec**" above the arrow to indicate the passage of an electric current to initiate the reaction. Balance the equation. $$2H_2O\,(l) \xrightarrow{elec} 2H_2\,(g) + O_2\,(g)$$
Think about your result.	The products are elements and the equation is balanced.

Exercise 3.4.2

Write the chemical equation for the decomposition of:

a. Al_2O_3
b. Ag_2S

Answer a

$$2Al_2O_3 \rightarrow 4Al + 3O_2$$

Answer b

$$Ag_2S \rightarrow 2Ag + S$$

Single Replacement Reactions

A third type of reaction is the single replacement reaction, in which one element replaces a similar element in a compound. The general form of a single-replacement (also called single-displacement) reaction is:

$$A + BC \rightarrow AC + B$$

In this general reaction, element **A** is a metal and replaces element **B**, also a metal, in the compound. When the element that is doing the replacing is a nonmetal, it must replace another nonmetal in a compound, and the general equation becomes:

$$\mathbf{Y + XZ \rightarrow XY + Z}$$

where **Y** is a nonmetal and replaces the nonmetal **Z** in the compound with **X**.

Metal Replacement

Magnesium is a more reactive metal than copper. When a strip of magnesium metal is placed in an aqueous solution of copper (II) nitrate, it replaces the copper. The products of the reaction are aqueous magnesium nitrate and solid copper metal.

$$\mathrm{Mg\,(\mathit{s})} + \mathrm{Cu(NO_3)_2\,(\mathit{aq})} \rightarrow \mathrm{Mg(NO_3)_2\,(\mathit{aq})} + \mathrm{Cu\,(\mathit{s})}$$

This subcategory of single-replacement reactions is called a metal replacement reaction because it is a metal that is being replaced (copper).

Hydrogen Replacement

Many metals react easily with acids and when they do so, one of the products of the reaction is hydrogen gas. Zinc reacts with hydrochloric acid to produce aqueous zinc chloride and hydrogen (figure below).

$$\mathrm{Zn\,(\mathit{s})} + \mathrm{2HCl\,(\mathit{aq})} \rightarrow \mathrm{ZnCl_2\,(\mathit{aq})} + \mathrm{H_2\,(\mathit{g})}$$

In a hydrogen replacement reaction, the hydrogen in the acid is replaced by an active metal. Some metals are so reactive that they are capable of replacing the hydrogen in water. The products of such a reaction are the metal hydroxide and hydrogen gas. All Group 1 metals undergo this type of reaction. Sodium reacts vigorously with water to produce aqueous sodium hydroxide and hydrogen (see figure below).

$$\mathrm{2Na\,(\mathit{s})} + \mathrm{2H_2O\,(\mathit{l})} \rightarrow \mathrm{2NaOH\,(\mathit{aq})} + \mathrm{H_2\,(\mathit{g})}$$



Figure **3.4.2**: (First image) Zinc metal reacts with hydrochloric acid to give off hydrogen gas in a single-displacement reaction. (Second image) Sodium metal reacts vigorously with water, giving off hydrogen gas. A large piece of sodium will often generate so much heat that the hydrogen will ignite.

Halogen Replacement

The element chlorine reacts with an aqueous solution of sodium bromide to produce aqueous sodium chloride and elemental bromine:

$$\mathrm{Cl_2\,(\mathit{g})} + \mathrm{2NaBr\,(\mathit{aq})} \rightarrow \mathrm{2NaCl\,(\mathit{aq})} + \mathrm{Br_2\,(\mathit{l})}$$

The reactivity of the halogen group (group 17) decreases from top to bottom within the group. Fluorine is the most reactive halogen, while iodine is the least. Since chlorine is above bromine, it is more reactive than bromine and can replace it in a halogen replacement reaction.

Example 3.4.3

What are the products of the reaction between solid aluminum (**Al**) and iron (III) oxide ($\mathbf{Fe_2O_3}$)?

Solution

Solutions to Example 7.10.3

Steps	Example Solution
Plan the problem.	To predict the products, we need to know that aluminum will replace iron and form aluminum oxide (the metal will replace the metal ion in the compound). Aluminum has a charge of $+3$ and oxygen has a charge of -2. The compound formed between aluminum and oxygen, therefore, will be Al_2O_3. Since iron is replaced in the compound by aluminum, the iron will now be the single element in the products.
Solve.	The unbalanced equation will be: $$Al + Fe_2O_3 \rightarrow Al_2O_3 + Fe$$ and the balanced equation will be: $$2Al + Fe_2O_3 \rightarrow Al_2O_3 + 2Fe$$
Think about your result.	This is a single replacement reaction, and when balanced the coefficients accurately reflect that the iron and aluminum have the same charge in this reaction.

? Exercise 3.4.3

a. Write the chemical equation for the single replacement reaction between zinc solid and lead (II) nitrate solution to produce zinc nitrate solution and solid lead. (Note that zinc forms ions with a $+2$ charge.)

b. Predict the products for the following reaction: $Fe + CuSO_4$. (In this reaction, assume iron forms ions with a $+2$ charge.)

Answer a

$$Zn + Pb(NO_3)_2 \rightarrow Pb + Zn(NO_3)_2$$

Answer b

$$Fe + CuSO_4 \rightarrow Cu + FeSO_4$$

Double Replacement Reactions

A **double-replacement reaction** is a reaction in which the positive and negative ions of two ionic compounds exchange places to form two new compounds. The general form of a double-replacement (also called double-displacement) reaction is:

$$AB + CD \rightarrow AD + BC$$

In this reaction, A and C are positively-charged cations, while B and D are negatively-charged anions. Double-replacement reactions generally occur between substances in aqueous solution. In order for a reaction to occur, one of the products is usually a solid precipitate, a gas, or a molecular compound such as water.

Formation of a Precipitate

A precipitate forms in a double-replacement reaction when the cations from one of the reactants combine with the anions from the other reactant to form an insoluble ionic compound. When aqueous solutions of potassium iodide and lead (II) nitrate are mixed, the following reaction occurs:

$$2KI\,(aq) + Pb(NO_3)_2\,(aq) \rightarrow 2KNO_3\,(aq) + PbI_2\,(s) \qquad (3.4.1)$$

There are very strong attractive forces that occur between Pb^{2+} and I^- ions and the result is a brilliant yellow precipitate (Figure **3.4.3**). The other product of the reaction, potassium nitrate, remains soluble.



Figure **3.4.3**: Lead (II) iodide precipitates when potassium iodide is mixed with lead (II) nitrate (Equation **3.4.1**). (CC BY-SA 3.0; PRHaney).

Formation of a Gas

Some double-replacement reactions produce a gaseous product which then bubbles out of the solution and escapes into the air. When solutions of sodium sulfide and hydrochloric acid are mixed, the products of the reaction are aqueous sodium chloride and hydrogen sulfide gas:

$$\mathrm{Na_2S}\,(aq) + 2\mathrm{HCl}\,(aq) \rightarrow 2\mathrm{NaCl}\,(aq) + \mathrm{H_2S}\,(g)$$

Formation of a Molecular Compound

Another kind of double-replacement reaction is one that produces a molecular compound as one of its products. Many examples in this category are reactions that produce water. When aqueous hydrochloric acid is reacted with aqueous sodium hydroxide, the products are aqueous sodium chloride and water:

$$\mathrm{HCl}\,(aq) + \mathrm{NaOH}\,(aq) \rightarrow \mathrm{NaCl}\,(aq) + \mathrm{H_2O}\,(l)$$

Example 3.4.4

Write a complete and balanced chemical equation for the double-replacement reaction $\mathrm{NaCN}\,(aq) + \mathrm{HBr}\,(aq) \rightarrow$ (hydrogen cyanide gas is formed).

Solution

Solutions to Example 7.10.4

Steps	Example Solution
Plan the problem.	The production of a gas drives the reaction.
Solve.	The cations of both reactants are $+1$ charged ions, while the anions are -1 charged ions. After exchanging partners, the balanced equation is: $\mathrm{NaCN}\,(aq) + \mathrm{HBr}\,(aq) \rightarrow \mathrm{NaBr}\,(aq) + \mathrm{HCN}\,(g)$
Think about your result.	This is a double replacement reaction. All formulas are correct and the equation is balanced.

Exercise 3.4.4

Write a complete and balanced chemical equation for the double-replacement reaction $(\mathrm{NH_4})_2\mathrm{SO_4}\,(aq) + \mathrm{Ba(NO_3)_2}\,(aq) \rightarrow$ (a precipitate of barium sulfate forms).

Answer a:

$$(\mathrm{NH_4})_2\mathrm{SO_4}\,(aq) + \mathrm{Ba(NO_3)_2}\,(aq) \rightarrow 2\mathrm{NH_4NO_3}\,(aq) + \mathrm{BaSO_4}\,(s)$$

Occasionally, a reaction will produce both a gas and a molecular compound. The reaction of a sodium carbonate solution with hydrochloric acid produces aqueous sodium chloride, carbon dioxide gas, and water:

$$\mathrm{Na_2CO_3}\,(aq) + \mathrm{2HCl}\,(aq) \rightarrow 2\ \text{DEL}\ \mathrm{NaCl}\,(aq) + \mathrm{CO_2}\,(g) + \mathrm{H_2O}\,(l)$$

3.4: Classifying Chemical Reactions is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.10: Classifying Chemical Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water

Learning Objectives

- Define and give examples of electrolytes.

When some substances are dissolved in water, they undergo either a physical or a chemical change that yields ions in solution. These substances constitute an important class of compounds called electrolytes. Substances that do not yield ions when dissolved are called nonelectrolytes. If the physical or chemical process that generates the ions is essentially 100% efficient (all of the dissolved compound yields ions), then the substance is known as a strong electrolyte. If only a relatively small fraction of the dissolved substance undergoes the ion-producing process, it is called a weak electrolyte.

Substances may be identified as strong, weak, or nonelectrolytes by measuring the electrical conductance of an aqueous solution containing the substance. To conduct electricity, a substance must contain freely mobile, charged species. Most familiar is the conduction of electricity through metallic wires, in which case the mobile, charged entities are electrons. Solutions may also conduct electricity if they contain dissolved ions, with conductivity increasing as ion concentration increases. Applying a voltage to electrodes immersed in a solution permits assessment of the relative concentration of dissolved ions, either quantitatively, by measuring the electrical current flow, or qualitatively, by observing the brightness of a light bulb included in the circuit (Figure 3.5.1).




Figure 3.5.1: Solutions of nonelectrolytes, such as ethanol, do not contain dissolved ions and cannot conduct electricity. Solutions of electrolytes contain ions that permit the passage of electricity. The conductivity of an electrolyte solution is related to the strength of the electrolyte. This diagram shows three separate beakers. Each has a wire plugged into a wall outlet. In each case, the wire leads from the wall to the beaker and is split resulting in two ends. One end leads to a light bulb and continues on to a rectangle labeled with a plus sign. The other end leads to a rectangle labeled with a minus sign. The rectangles are in a solution. In the first beaker, labeled “Ethanol No Conductivity,” four pairs of linked small green spheres suspended in the solution between the rectangles. In the second beaker, labeled “K C l Strong Conductivity,” six individual green spheres, three labeled plus and three labeled minus are suspended in the solution. Each of the six spheres has an arrow extending from it pointing to the rectangle labeled with the opposite sign. In the third beaker, labeled “Acetic acid solution Weak conductivity,” two pairs of joined green spheres and two individual spheres, one labeled plus and one labeled minus are shown suspended between the two rectangles. The plus labeled sphere has an arrow pointing to the rectangle labeled minus and the minus labeled sphere has an arrow pointing to the rectangle labeled plus.

Ionic Electrolytes

Water and other polar molecules are attracted to ions, as shown in Figure 3.5.2. The electrostatic attraction between an ion and a molecule with a dipole is called an ion-dipole attraction. These attractions play an important role in the dissolution of ionic compounds in water.




Figure 3.5.2: As potassium chloride (KCl) dissolves in water, the ions are hydrated. The polar water molecules are attracted by the charges on the K^+ and Cl^- ions. Water molecules in front of and behind the ions are not shown. The diagram shows eight purple spheres labeled K superscript plus and eight green spheres labeled C l superscript minus mixed and touching near the center of the diagram. Outside of this cluster of spheres are seventeen clusters of three spheres, which include one red and two white spheres. A red sphere in one of these clusters is labeled O. A white sphere is labeled H. Two of the green C l superscript minus spheres are surrounded by three of the red and white clusters, with the red spheres closer to the green spheres than the white spheres. One of the K superscript plus purple spheres is surrounded by four of the red and white clusters. The white spheres of these clusters are closest to the purple spheres.

When ionic compounds dissolve in water, the ions in the solid separate and disperse uniformly throughout the solution because water molecules surround and solvate the ions, reducing the strong electrostatic forces between them. This process represents a physical change known as dissociation. Under most conditions, ionic compounds will dissociate nearly completely when dissolved, and so they are classified as strong electrolytes.

Let us consider what happens at the microscopic level when we add solid KCl to water. Ion-dipole forces attract the positive (hydrogen) end of the polar water molecules to the negative chloride ions at the surface of the solid, and they attract the negative (oxygen) ends to the positive potassium ions. The water molecules penetrate between individual K^+ and Cl^- ions and surround them, reducing the strong interionic forces that bind the ions together and letting them move off into solution as solvated ions, as Figure 3.5.2 shows. The reduction of the electrostatic attraction permits the independent motion of each hydrated ion in a dilute solution, resulting in an increase in the disorder of the system, as the ions change from their fixed and ordered positions in the crystal to mobile and much more disordered states in solution. This increased disorder is responsible for the dissolution of many ionic compounds, including KCl, which dissolve with absorption of heat.

In other cases, the electrostatic attractions between the ions in a crystal are so large, or the ion-dipole attractive forces between the ions and water molecules are so weak, that the increase in disorder cannot compensate for the energy required to separate the ions, and the crystal is insoluble. Such is the case for compounds such as calcium carbonate (limestone), calcium phosphate (the inorganic component of bone), and iron oxide (rust).

Solubility Rules

Some combinations of aqueous reactants result in the formation of a solid precipitate as a product. However, some combinations will not produce such a product. If solutions of sodium nitrate and ammonium chloride are mixed, no reaction occurs. One could write a molecular equation showing a double-replacement reaction, but both products, sodium chloride and ammonium nitrate, are soluble and would remain in the solution as ions. Every ion is a spectator ion and there is no net ionic equation at all. It is useful to be able to predict when a precipitate will occur in a reaction. To do so, you can use a set of guidelines called the **solubility rules** (Tables 3.5.1 and 3.5.2).

Table 3.5.1: Solubility Rules for Soluble Substances

Soluble in Water	Important Exceptions (Insoluble)
All Group IA and NH_4^+ salts	none
All nitrates, chlorates, perchlorates and acetates	none
All sulfates	$CaSO_4$, $BaSO_4$, $SrSO_4$, $PbSO_4$
All chlorides, bromides, and iodides	AgX, Hg_2X_2, PbX_2 (X= Cl, Br, or I)

Table **3.5.2**: Solubility Rules for Sparingly Soluble Substances

Sparingly Soluble in Water	Important Exceptions (Soluble)
All carbonates and phosphates	Group IA and NH_4^+ salts
All hydroxides	Group IA and NH_4^+ salts; Ba^{2+}, Sr^{2+}, Ca^{2+} sparingly soluble
All sulfides	Group IA, IIA and NH_4^+ salts; MgS, CaS, BaS sparingly soluble
All oxalates	Group IA and NH_4^+ salts
Special note: The following electrolytes are of only moderate solubility in water: CH_3COOAg, Ag_2SO_4, $KClO_4$. They will precipitate only if rather concentrated solutions are used.	

As an example on how to use the solubility rules, predict if a precipitate will form when solutions of cesium bromide and lead (II) nitrate are mixed.

$$Cs^+(aq) + Br^-(aq) + Pb^{2+}(aq) + 2NO_3^-(aq) \rightarrow ?$$

The potential precipitates from a double-replacement reaction are cesium nitrate and lead (II) bromide. According to the solubility rules table, cesium nitrate is soluble because all compounds containing the nitrate ion, as well as all compounds containing the alkali metal ions, are soluble. Most compounds containing the bromide ion are soluble, but lead (II) is an exception. Therefore, the cesium and nitrate ions are spectator ions and the lead (II) bromide is a precipitate. The balanced net ionic reaction is:

$$Pb^{2+}(aq) + 2Br^-(aq) \rightarrow PbBr_2(s)$$

✔ Example 3.5.1: Solubility

Classify each compound as soluble or insoluble

a. $Zn(NO_3)_2$
b. $PbBr_2$
c. $Sr_3(PO_4)_2$

Solution

a. All nitrates are soluble in water, so $Zn(NO_3)_2$ is soluble.
b. All bromides are soluble in water, except those combined with Pb^{2+}, so $PbBr_2$ is insoluble.
c. All phosphates are insoluble, so $Sr_3(PO_4)_2$ is insoluble.

? Exercise 3.5.1: Solubility

Classify each compound as soluble or insoluble.

a. $Mg(OH)_2$
b. KBr
c. $Pb(NO_3)_2$

Answer a

insoluble

Answer b

soluble

Answer c

soluble

Summary

Substances that dissolve in water to yield ions are called electrolytes. Nonelectrolytes are substances that do not produce ions when dissolved in water. Solubility rules allow prediction of what products will be insoluble in water.

Contributions & Attributions

- Paul Flowers (University of North Carolina - Pembroke), Klaus Theopold (University of Delaware) and Richard Langley (Stephen F. Austin State University) with contributing authors. Textbook content produced by OpenStax College is licensed under a Creative Commons Attribution License 4.0 license. Download for free at http://cnx.org/contents/85abf193-2bd...a7ac8df6@9.110).
-

3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.6: Precipitation Reactions

Learning Objectives

- To identify a precipitation reaction and predict solubility.

A precipitation reaction is a reaction that yields an insoluble product—a precipitate—when two solutions are mixed. When a colorless solution of silver nitrate is mixed with a yellow-orange solution of potassium dichromate, a reddish precipitate of silver dichromate is produced.

$$\mathrm{AgNO_3\,(aq) + K_2Cr_2O_7\,(aq) \rightarrow Ag_2Cr_2O_7\,(s) + KNO_3\,(aq)} \tag{3.6.1}$$

This unbalanced equation has the general form of an exchange reaction:

$$\mathrm{AC + BD \rightarrow \underset{insoluble}{AD} + BC} \tag{3.6.2}$$

Thus precipitation reactions are a subclass of exchange reactions that occur between ionic compounds when one of the products is insoluble. Because both components of each compound change partners, such reactions are sometimes called **double-displacement reactions**. Precipitation reactions are used to isolate metals that have been extracted from their ores, and to recover precious metals for recycling.



Video: Mixing potassium dichromate and silver nitrate together to initiate a precipitation reaction (Equation 3.6.1).

Just as important as predicting the product of a reaction is knowing when a chemical reaction will *not* occur. Simply mixing solutions of two different chemical substances does *not* guarantee that a reaction will take place. For example, if 500 mL of aqueous NaCl solution is mixed with 500 mL of aqueous KBr solution, the final solution has a volume of 1.00 L and contains $\mathrm{Na^+\,(aq)}$, $\mathrm{Cl^-\,(aq)}$, $\mathrm{K^+\,(aq)}$, and $\mathrm{Br^-\,(aq)}$. As you will see in (Figure 3.6.1), none of these species reacts with any of the others. When these solutions are mixed, the only effect is to dilute each solution with the other.





Figure **3.6.1**: The Effect of Mixing Aqueous **KBr** and **NaCl** Solutions. Because no net reaction occurs, the only effect is to dilute each solution with the other. (Water molecules are omitted from molecular views of the solutions for clarity.)

Predicting Precipitation Reactions

A precipitation reaction occurs when a solid precipitate forms after mixing two strong electrolyte solutions. As stated previously, if none of the species in the solution reacts then no net reaction occurred.

Predict what will happen when aqueous solutions of barium chloride and lithium sulfate are mixed.

Change the partners of the anions and cations on the reactant side to form new compounds (products):



Chemical equation of the reactants barium chloride and lithium sulfate forming the products barium sulfate and lithium chloride.

Because barium chloride and lithium sulfate are strong electrolytes, each dissociates completely in water to give a solution that contains the constituent anions and cations. Mixing the two solutions *initially* gives an aqueous solution that contains Ba^{2+}, Cl^-, Li^+, and SO_4^{2-} ions. The only possible exchange reaction is to form **LiCl** and **$BaSO_4$**.

Correct the formulas of the products based on the charges of the ions.

No need to correct the formula as both compounds already have their charges balanced.

$$BaCl_2\,(aq) + Li_2SO_4\,(aq) \rightarrow BaSO_4 + LiCl$$

Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.

$$BaCl_2\,(aq) + Li_2SO_4\,(aq) \rightarrow BaSO_4\,(s) + LiCl\,(aq)$$

Table 7.5.1 from the previous section shows that **LiCl** is soluble in water, but **$BaSO_4$** is not soluble in water.

Balance the equation:

$$BaCl_2\,(aq) + Li_2SO_4\,(aq) \rightarrow BaSO_4\,(s) + 2\,LiCl\,(aq)$$

Although soluble barium salts are toxic, **$BaSO_4$** is so insoluble that it can be used to diagnose stomach and intestinal problems without being absorbed into tissues. An outline of the digestive organs appears on x-rays of patients who have been given a "barium milkshake" or a "barium enema"—a suspension of very fine **$BaSO_4$** particles in water.




Figure **3.6.2**: An x-ray of the digestive organs of a patient who has swallowed a “barium milkshake.” A barium milkshake is a suspension of very fine $BaSO_4$ particles in water; the high atomic mass of barium makes it opaque to x-rays. (Public Domain; Glitzy queen00 via Wikipedia).

Example 3.6.1

Predict what will happen if aqueous solutions of rubidium hydroxide and cobalt(II) chloride are mixed.

Solution

Solutions to Example 7.6.1


Steps	Example
Change the partners of the anions and cations on the reactant side to form new compounds (products).	 Chemical equation of the reactants rubidium hydroxide and cobalt(II) chloride forming the products rubidium chloride and cobalt hydroxide.
Correct the formulas of the products based on the charges of the ions.	$\mathbf{RbOH\,(aq) + CoCl_2\,(aq) \rightarrow RbCl + Co(OH)_2}$
Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.	$\mathbf{RbOH\,(aq) + CoCl_2\,(aq) \rightarrow RbCl\,(aq) + Co(OH)_2\,(s)}$
Balance the equation.	Coefficients already balanced. $\mathbf{RbOH\,(aq) + CoCl_2\,(aq) \rightarrow RbCl\,(aq) + Co(OH)_2\,(s)}$

Example 3.6.2

Predict what will happen if aqueous solutions of strontium bromide and aluminum nitrate are mixed.

Solution

Solutions for Example 7.6.2

Steps	Example
Change the partners of the anions and cations on the reactant side to form new compounds (products).	 Chemical equation of the reactants strontium bromide and aluminum nitrate forming the products strontium nitrate and aluminum bromide.
Correct the formulas of the products based on the charges of the ions.	$\mathbf{SrBr_2\,(aq) + Al(NO_3)_3\,(aq) \rightarrow Sr(NO_3)_2 + AlBr_3}$
Refer to the solubility rules table to determine insoluble products which will therefore form a precipitate.	$\mathbf{SrBr_2\,(aq) + Al(NO_3)_3\,(aq) \rightarrow Sr(NO_3)_2\,(aq) + AlBr_3\,(aq)}$ According to Table 7.5.1 from the previous section, both $\mathbf{AlBr_3}$ (rule 4) and $\mathbf{Sr(NO_3)_2}$ (rule 2) are soluble.
If all possible products are soluble, then no net reaction will occur.	$\mathbf{SrBr_2\,(aq) + Al(NO_3)_3\,(aq) \rightarrow}$ **NO REACTION**

? Exercise 3.6.2

Using the information in Table 7.5.1 from the previous section, predict what will happen in each case involving strong electrolytes.

a. An aqueous solution of strontium hydroxide is added to an aqueous solution of iron(II) chloride.
b. Solid potassium phosphate is added to an aqueous solution of mercury(II) perchlorate.
c. Solid sodium fluoride is added to an aqueous solution of ammonium formate.
d. Aqueous solutions of calcium bromide and cesium carbonate are mixed.

Answer a

$Fe(OH)_2$ precipitate is formed.

Answer b

$Hg_3(PO_4)_2$ precipitate is formed.

Answer c

No Reaction.

Answer d

$CaCO_3$ is precipitate formed.

Summary

In a **precipitation reaction**, a subclass of exchange reactions, an insoluble material (a **precipitate**) forms when two electrolyte solutions are mixed. To predict the product of a precipitation reaction, all species initially present in the solutions are identified, as are any combinations likely to produce an insoluble salt.

Contributions & Attributions

- Modified by Joshua Halpern (Howard University)

3.6: Precipitation Reactions is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.6: Precipitation Reactions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

A typical precipitation reaction occurs when an aqueous solution of barium chloride is mixed with one containing sodium sulfate. The **complete chemical equation** can be written to describe what happens, and such an equation is useful in making chemical calculations.

$$\underbrace{\mathrm{BaCl_2\,(aq) + Na_2SO_4\,(aq) \longrightarrow BaSO_4\,(s) + 2\,NaCl\,(aq)}}_{\text{Complete Chemical Equation}} \tag{3.7.1}$$

However, Equation 3.7.1 does not really represent the microscopic particles (that is, the ions) present in the solution. Below is the complete ionic equation:

$$\underbrace{\mathrm{Ba^{2+}\,(aq) + \overbrace{2Cl^-(aq)}^{\text{spectator}} + \overbrace{2Na^+(aq)}^{\text{spectator}} + SO_4{}^{2-}\,(aq) \longrightarrow BaSO_4\,(s) + \overbrace{2Na^+(aq)}^{\text{spectator}} + \overbrace{Cl^-(aq)}^{\text{spectator}}}}_{\text{Complete Ionic Equation}} \tag{3.7.2}$$

Equation 3.7.2 is rather cumbersome and includes so many different ions that it may be confusing. In any case, we are often interested in the independent behavior of ions, not the specific compound from which they came. A precipitate of $\mathrm{BaSO_4\,(s)}$ will form when *any* solution containing $\mathrm{Ba^{2+}\,(aq)}$ is mixed with *any* solution containing $\mathrm{SO_4{}^{2-}\,(aq)}$ (provided concentrations are not extremely small). This happens independently of the $\mathrm{Cl^-\,(aq)}$ and $\mathrm{Na^+\,(aq)}$ ions in Equation 3.7.2. These ions are called **spectator ions** because they do not participate in the reaction. When we want to emphasize the independent behavior of ions, a **net ionic equation** is written, omitting the spectator ions. For precipitation of $\mathrm{BaSO_4}$ the net ionic equation is

$$\underbrace{\mathrm{Ba^{2+}\,(aq) + SO_4{}^{2-}\,(aq) \longrightarrow BaSO_4\,(s)}}_{\text{Net Ionic Equation}} \tag{3.7.3}$$

Example 3.7.1

a. When a solution of $\mathrm{AgNO_3}$ is added to a solution of $\mathrm{CaCl_2}$, insoluble AgCl precipitates. Write three equations (complete chemical equation, complete ionic equation, and net ionic equation) that describe this process.

b. Write the balanced net ionic equation to describe any reaction that occurs when the solutions of $\mathrm{Na_2SO_4}$ and $\mathrm{NH_4I}$ are mixed.

Solution

Solutions to Example 7.7.1

Equation Type	Example 3.7.1*a*	Example 3.7.1*b*
Complete Chemical Equation	$\mathrm{2\,AgNO_3\,(aq) + CaCl_2\,(aq) \longrightarrow 2\,AgCl\,(s) + Ca(NO_3)_2\,(aq)}$ The proper states and formulas of all products are written and the chemical equation is balanced.	$\mathrm{Na_2SO_4\,(aq) + 2\,NH_4I\,(aq) \longrightarrow 2\,NaI\,(aq) + (NH_4)_2SO_4\,(aq)}$ Both products are aqueous so there is no net ionic equation that can be written.
Complete Ionic Equation	$\mathrm{2\,Ag^+\,(aq) + 2\,NO_3{}^-\,(aq) + Ca^{2+}\,(aq) + 2\,Cl^-\,(aq) \longrightarrow 2\,AgCl\,(s) + Ca^{2+}\,(aq) + 2\,NO_3{}^-\,(aq)}$ AgCl is a solid so it does not break up into ions in solution.	
Net Ionic Equation	$\mathrm{Ag^+\,(aq) + Cl^-\,(aq) \longrightarrow AgCl\,(s)}$ All spectator ions are removed. The chemical equation is written using the lowest common coefficients.	NaI and $\mathrm{(NH_4)_2SO_4}$ are both soluble. No net ionic equation. There is no reaction.

The occurrence or nonoccurrence of precipitates can be used to detect the presence or absence of various species in solution. A $BaCl_2$ solution, for instance, is often used as a test for the presence of $SO_4^{2-}(aq)$ ions. There are several insoluble salts of Ba, but they all dissolve in dilute acid except for $BaSO_4$. Thus, if $BaCl_2$ solution is added to an unknown solution which has previously been acidified, the occurrence of a white precipitate is proof of the presence of the SO_4^{2-} ion.



Figure 3.7.1: The three common silver halide precipitates: AgI, AgBr and AgCl (left to right). The silver halides precipitate out of solution, but often form suspensions before settling. (CC BY-SA 3.0; Cychr).

$AgNO_3$ solutions are often used in a similar way to test for halide ions. If $AgNO_3$ solution is added to an acidified unknown solution, a white precipitate indicates the presence of Cl^- ions, a cream-colored precipitate indicates the presence of Br^- ions, and a yellow precipitate indicates the presence of I^- ions (Figure 3.7.1). Further tests can then be made to see whether perhaps a mixture of these ions is present. When $AgNO_3$ is added to tap water, a white precipitate is almost always formed. The Cl^- ions in tap water usually come from the Cl_2 which is added to municipal water supplies to kill microorganisms.

? Exercise 3.7.1

Write balanced net ionic equations to describe any reaction that occurs when the following solutions are mixed.

a. $K_2CO_3 + SrCl_2$
b. $FeSO_4 + Ba(NO_3)_2$

Answer a

$$Sr^{2+}(aq) + CO_3^{2-}(aq) \longrightarrow SrCO_3(s)$$

Answer b

$$Ba^{2+}(aq) + SO_4^{2-}(aq) \longrightarrow Ba(SO_4)(s)$$

Precipitates are also used for quantitative analysis of solutions, that is, to determine the amount of solute or the mass of solute in a given solution. For this purpose it is often convenient to use the first of the three types of equations described above. Then the rules of stoichiometry may be applied.

Contributions & Attributions

- Ed Vitz (Kutztown University), John W. Moore (UW-Madison), Justin Shorb (Hope College), Xavier Prat-Resina (University of Minnesota Rochester), Tim Wendorff, and Adam Hahn.

3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **7.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.8: PRELUDE TO CHEMICAL COMPOSITION - HOW MUCH SODIUM?

Why is knowledge of composition important? Everything in nature is either chemically or physically combined with other substances. To find the amount of a material in a sample, you need to know what fraction of the sample it is. Some simple applications of composition are: the amount of sodium in sodium chloride for a diet, the amount of iron in iron ore for steel production, the amount of hydrogen in water for hydrogen fuel, and the amount of chlorine in freon to estimate ozone depletion.



How much salt is in this salt packet? (CC-BY-SA; 3.0; Swilliams).

3.8: Prelude to Chemical Composition - How Much Sodium? is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.1: Prelude to Chemical Composition - How Much Sodium?** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-SA 4.0.

3.9: COUNTING NAILS BY THE POUND

COUNTING BY WEIGHING AND AVOGADRO'S NUMBER

The size of molecule is so small that it is physically difficult, if not impossible, to directly count out molecules (Figure **3.9.1**). However, we can count them indirectly by using a common trick of "counting by weighing".




Figure **3.9.1**: A comparison of the scales of various biological and technological objects. (CC BY-SA 3.0; Wikipedia)

Consider the example of counting nails in a big box at a hardware store. You need to estimate the number of nails in a box. The weight of an empty box is **213 *g*** and the weight of the box plus a bunch of big nails is **1340 *g***. Assume that we know that the weight of one big nail is 0.450 g. Hopefully it's not necessary to tear open the package and count the nails. We agree that

$$\textbf{mass of big nails} = 1340\,g - 213\,g = 1227\,g$$

Therefore

$$\textbf{Number of big nails in box} = \frac{1227\,g}{0.450\,g/\textbf{big nail}} = 2,726.6\,\textbf{big nails} = 2,730\,\textbf{big nails.} \tag{3.9.1}$$

You have just counted the number of big nails in the box by weighing them (rather than by counting them individually).



Figure **3.9.2**: Galvanized nails. Individually counting nails in a box would would require significant effort. Alternatively, we can count them by weighing. (Public Domain; Wikipedia).

Now consider if the box of nails weighed the same, but the box were filled with small nails with an individual mass of **0.23 *g*/small nail** instead? You would do the same math, but use a different denominator in Equation 3.9.1:

$$\textbf{Number of small nails in box} = \frac{1227\,g}{0.230\,g/\textbf{small nail}} = 5,$$
$$334.7\,\textbf{small nails} = 5,335\,\textbf{small nails.} \tag{3.9.2}$$

The individual mass is the conversion factor used in the calculation and changes, based on the nature of the nail (big or small). Let's ask a different question: how many *dozens* of nails are there in the same box of small nails described above?

If we know the information from Equation 3.9.2, we can just use the conversion of how many nails are in a dozen:

$$\frac{5,335\,\text{small nails}}{12\,\text{small nails/dozen}} = 444.6\,\text{dozen small nails} \tag{3.9.3}$$

If we want to get this value from weighing, we use the "dozen mass" instead of individual mass:

$$12 \times 0.23g = 2.76\,g/\text{dozen small nails.} \tag{3.9.4}$$

So following Equation 3.9.2, we get:

$$\begin{aligned}\text{Number of dozens of small nails} &= \frac{1227\,g}{2.76\,g/\text{dozen small nails}} \\ &= 444.6\,\text{dozen small nails}\end{aligned} \tag{3.9.5}$$

and this is the same result as Equation 3.9.3. These calculations demonstrate the difference between individual mass (i.e., per individual) and collective mass (e.g., per dozen or per gross). The collective mass of most importance to chemistry is *molar mass* (i.e., mass per mole or mass per 6.022×10^{23}).

AVOGADRO'S NUMBER

Avogadro's number is an accident of nature. It is the number of particles that delivers a mole of a substance. Avogadro's number = 6.022×10^{23}. The reason why the value is an accident of nature is that the mole is tied to the gram mass unit. The gram is a convenient mass unit because it matches human sizes. If we were a thousand times greater in size (like Paul Bunyan) we would find it handy to use kilogram amounts. This means the kilogram mole would be convenient. The number of particles handled in a kilogram mole is 1000 times greater. The kilo Avogadro number for the count of particles in a kilomole is 6.022×10^{26}.

If humans were tiny creatures (like Lilliputians) only 1/1000 our present size, milligrams would be more convenient. This means the milligram mole would be more useful. The number of particles handled in a milligram mole (millimole) would be 1/1000 times smaller. The milli Avogadro number for the count of particles in a millimole is 6.022×10^{20}.

What do you think would happen to Avogadro's number if the American system was used and amounts were measured in pound moles? Remember 1 pound = 454 grams. Avogadro's number would be larger by a factor of 454. A pound mole of hydrogen would weigh 1 pound, which is 454 grams. A gram mole of hydrogen weighs 1 gram and contains 6.022×10^{23} H atoms.

MOLAR MASS FOR ELEMENTS

You are able to read the periodic table and determine the average atomic mass for an element like carbon. The average mass is 12.01 amu. This mass is a ridiculously tiny number of grams. It is too small to handle normally. The molar mass of carbon is defined as the mass in grams that is numerically equal to the average atomic weight. This means

$$1g/molecarbon = 12.01\,g\,carbon$$

this is commonly written

$$1\,mol\,carbon = 12.01\,grams\,carbon.$$

This is the mass of carbon that contains 6.022×10^{23} carbon atoms.

- Avogadro's number is 6.022×10^{23} particles.

This same process gives us the molar mass of any element. For example:

- $1\,mol\,neon = 20.18\,g\,neon\,Ne$
- $1\,mol\,sodium = 22.99\,g\,sodium\,Na$

MOLAR MASS FOR COMPOUNDS

EXAMPLE 3.9.1: MOLAR MASS OF WATER

The formulas for compounds are familiar to you. You know the formula for water is H_2O. It should be reasonable that the weight of a formula unit can be calculated by adding up the weights for the atoms in the formula.

Solution

The formula weight for water

weight from hydrogen + weight from oxygen

The formula weight for water

2 H atoms x 1.008 amu + 1 O atom x 16.00 amu = 18.016 amu

The molar mass for water

18.016 grams water or 18 grams to the nearest gram

EXAMPLE 3.9.2: MOLAR MASS OF METHANE

The formula for methane, the major component in natural gas, is CH_4.

Solution

The formula weight for methane

weight from hydrogen + weight from carbon

The formula weight for methane

4 H atoms x 1.008 amu + 1 C atom x 12.01 amu = 16.04 amu

The molar mass for methane

16.04 grams per mole of methane

EXAMPLE 3.9.3: MOLAR MASS OF ETHYL CHLORIDE

What is its molar mass for ethyl chloride CH_3CH_2Cl?

Solution

The *formula weight*

weight from hydrogen + weight from carbon + weight from chlorine

The formula weight

5 H atoms x 1.008 amu + 2 C atom x 12.01 amu + 35.5 amu = 64.5 amu

The molar mass for ethyl chloride

64.5 grams per mole of ethyl chloride

EXAMPLE 3.9.4: MOLAR MASS OF SULFUR DIOXIDE

What is the molar mass for sulfur dioxide, $SO_2(g)$, a gas used in bleaching and disinfection processes?

Solution

Look up the atomic weight for each of the elements in the formula.

- 1 sulfur atom = 32.07 amu
- 1 oxygen atom = 16.00 amu

Count the atoms of each element in the formula unit.

- one sulfur atom
- two oxygen atoms

The formula weight

weight from sulfur + weight from oxygen

The formula weight

1 sulfur atom x (32. 07 amu) + 2 oxygen atoms x (16.00 amu)

The formula weight

SO_2 = 32. 07 amu + 32.00 amu = 64.07 amu = 64 amu SO_2

The molar mass for SO_2 is

64.07 grams of SO_2; 1 mol SO_2 = 64 grams per mole of SO_2

EXERCISE 3.9.1

What is the formula weight and molar mass for alum, $KAl(SO_4)_2 \cdot 12H_2O$?

Answer

1. Check the periodic table for the atomic masses for each atom in the formula.
2. Count the number of each type of atom in the formula.
3. Multiply the number of atoms by the atomic mass for each element.
4. Add up the masses for all of the elements.

Table **3.9.1**: Masses of each element in alum, $KAl(SO_4)_2 \cdot 12H_2O$

element	average atomic mass	number of atoms in formula	rounded to nearest one unit for simplicity
potassium k	39.1 amu	1	39. amu
aluminum	26.98 amu	1	27. amu
sulfur	32.07 amu	2	64. amu
oxygen	16.00 amu	8 + 12 = 20	320. amu
hydrogen	1.008 amu	2 x 12 = 24	24. amu

Molar mass is 474 grams (add up the amu of each element to find the total of 474 amu). This is a mass in grams that is numerically (474) the same as the formula weight.

1 mole alum $KAl(SO_4)_2 \cdot 12H_2O$ = 474 grams alum $KAl(SO_4)_2 \cdot 12H_2O$

This page titled 3.9: Counting Nails by the Pound is shared under a CC BY-SA 3.0 license and was authored, remixed, and/or curated by Delmar Larsen, Marisa Alviar-Agnew, Henry Agnew, & Henry Agnew.

- **6.2: Counting Nails by the Pound** by Delmar Larsen, Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-SA 3.0.

3.10: Counting Atoms by the Gram

Learning Objectives

- Use Avogadro's number to convert to moles and vice versa given the number of particles of an element.
- Use the molar mass to convert to grams and vice versa given the number of moles of an element.

When objects are very small, it is often inconvenient, inefficient, or even impossible to deal with the objects one at a time. For these reasons, we often deal with very small objects in groups, and have even invented names for various numbers of objects. The most common of these is "dozen" which refers to 12 objects. We frequently buy objects in groups of 12, like doughnuts or pencils. Even smaller objects such as straight pins or staples are usually sold in boxes of 144, or a dozen dozen. A group of 144 is called a "gross".

This problem of dealing with things that are too small to operate with as single items also occurs in chemistry. Atoms and molecules are too small to see, let alone to count or measure. Chemists needed to select a group of atoms or molecules that would be convenient to operate with.

Avogadro's Number and Mole

In chemistry, it is impossible to deal with a single atom or molecule because we can't see them, count them, or weigh them. Chemists have selected a number of particles with which to work that is convenient. Since molecules are extremely small, you may suspect this number is going to be very large, and you are right. The number of particles in this group is $\mathbf{6.02 \times 10^{23}}$ particles and the name of this group is the **mole** (the abbreviation for ***mole*** is $\mathbf{mol}$). One mole of any object is $\mathbf{6.02 \times 10^{23}}$ of those objects. There is a particular reason that this number was chosen and this reason will become clear as we proceed.

When chemists are carrying out chemical reactions, it is important that the relationship between the numbers of particles of each reactant is known. Any readily measurable mass of an element or compound contains an extraordinarily large number of atoms, molecules, or ions, so an extremely large numerical unit is needed to count them. The mole is used for this purpose.

The **mole** (symbol: **mol**) is the base unit of amount of substance ("number of substance") in the International System of Units or System International (SI), defined as exactly $6.02214076 \times 10^{23}$ particles, e.g., atoms, molecules, ions or electrons. The current definition was adopted in November 2018, revising its old definition based on the number of atoms in 12 grams of carbon-12 (^{12}C) (the isotope of carbon with relative atomic mass 12 Daltons, by definition). For most purposes, 6.022×10^{23} provides an adequate number of significant figures. Just as 1 mole of atoms contains 6.022×10^{23} atoms, 1 mole of eggs contains 6.022×10^{23} eggs. This number is called Avogadro's number, after the 19th-century Italian scientist who first proposed a relationship between the volumes of gases and the numbers of particles they contain.

It is not obvious why eggs come in dozens rather than 10s or 14s, or why a ream of paper contains 500 sheets rather than 400 or 600. The definition of a mole—that is, the decision to base it on 12 g of carbon-12—is also arbitrary. The important point is that 1 mole of carbon—or of anything else, whether atoms, compact discs, or houses—always has the same number of objects: 6.022×10^{23}.

Converting Between Number of Atoms to Moles and Vice Versa

We can use Avogadro's number as a conversion factor, or ratio, in dimensional analysis problems. If we are given the number of atoms of an element X, we can convert it into moles by using the relationship

$$\mathbf{1\ mol\ X = 6.022 \times 10^{23}\ X\ atoms.}$$

Example 3.10.1: Moles of Carbon

The element carbon exists in two primary forms: graphite and diamond. How many moles of carbon atoms is $\mathbf{4.72 \times 10^{24}}$ atoms of carbon?

Solution

Solutions to Example 6.3.1

Steps for Problem Solving	The element carbon exists in two primary forms: graphite and diamond. How many moles of carbon atoms is 4.72×10^{24} atoms of carbon?
Identify the "given" information and what the problem is asking you to "find."	Given: 4.72×10^{24} C atoms Find: mol C
List other known quantities.	$1\,mol = 6.022 \times 10^{23}$ C atoms
Prepare a concept map and use the proper conversion factor.	# C atoms → mol C $\frac{1\,mol\,C}{6.022 \times 10^{23} C\,atoms}$
Cancel units and calculate.	$4.72 \times 10^{24}\ \cancel{\text{C atoms}} \times \frac{1\ \text{mol C}}{6.02 \times 10^{23}\ \cancel{\text{C atoms}}} = 7.84\ \text{mol C}$
Think about your result.	The given number of carbon atoms was greater than Avogadro's number, so the number of moles of C atoms is greater than 1 mole. Since Avogadro's number is a measured quantity with three significant figures, the result of the calculation is rounded to three significant figures.

Molar Mass

Molar mass is defined as the mass of one mole of representative particles of a substance. By looking at a periodic table, we can conclude that the molar mass of the element lithium is 6.94g, the molar mass of zinc is 65.38g, and the molar mass of gold is 196.97g. Each of these quantities contains 6.022×10^{23} atoms of that particular element. The units for molar mass are grams per mole or g/mol. $1.00\ \text{mol}$ of carbon-12 atoms has a mass of $12.0\ \text{g}$ and contains 6.022×10^{23} atoms. 1.00 mole of any element has a mass numerically equal to its atomic mass in grams and contains 6.022×10^{23} particles. The mass, in grams, of 1 mole of particles of a substance is now called the **molar mass** (mass of 1.00 mole).

Converting Grams to Moles of an Element and Vice Versa

We can also convert back and forth between grams of an element and moles. The conversion factor for this is the molar mass of the substance. The **molar mass** is the ratio giving the number of grams for each one mole of the substance. This ratio is easily found by referring to the atomic mass of the element using the periodic table. This ratio has units of grams per mole or g/mol.

Conversions like this are possible for any substance, as long as the proper atomic mass, formula mass, or molar mass is known (or can be determined) and expressed in grams per mole. Figure 6.4.1 illustrates what conversion factor is needed and two examples are given below.




Figure 3.10.1: A Simple Flowchart for Converting Between Mass and Moles of an Element.
Flowchart: to convert moles of Element A to mass of Element A, use g/mol, and to convert vice versa, use mol/g.

Example 3.10.2: Chromium

Chromium metal is used for decorative electroplating of car bumpers and other surfaces. Find the mass of 0.560 moles of chromium.

Solution

Solutions to Example 6.3.2

Steps for Problem Solving	**Chromium metal is used for decorative electroplating of car bumpers and other surfaces. Find the mass of 0.560 moles of chromium.**
Identify the "given" information and what the problem is asking you to "find."	Given: 0.560 mol Cr Find: g Cr
List other known quantities.	1 mol Cr = 52.00g Cr
Prepare a concept map and use the proper conversion factor.	mol Cr → g Cr $\frac{52.00g\ Cr}{1\ mol\ Cr}$
Cancel units and calculate.	$0.560\ \cancel{mol\ Cr} \times \frac{52.00\ g\ Cr}{1\ \cancel{mol\ Cr}} = 29.1\ g\ Cr$
Think about your result.	Since the desired amount was slightly more than one half of a mole, the mass should be slightly more than one half of the molar mass. The answer has three significant figures because of the **0.560 mol**

Example 3.10.3: Silicon

How many moles are in 107.6g of Si?

Solution

Solutions to Example 6.3.3

Steps for Problem Solving	**How many moles are in 107.6g of Si.**
Identify the "given" information and what the problem is asking you to "find."	Given: 107.6g Si Find: mol Si
List other known quantities.	1 mol Si = 28.09g Si
Prepare a concept map and use the proper conversion factor.	g Si → mol Si $\frac{1\ mol\ Si}{28.09g\ Si}$
Cancel units and calculate.	$107.6\ \cancel{g\ Si} \times \frac{1\ mol\ Si}{28.09\ \cancel{g\ Si}} = 3.83\ mol\ Si$
Think about your result.	Since 1 mol of Si is 28.09g, 107.6 should be about 4 moles.

? Exercise 3.10.1

a. How many moles are present in 100.0 g of Al?
b. What is the mass of 0.552 mol of Ag metal?

Answer a:

3.706 mol Al

Answer b:

59.5 g Ag

Summary

- A mole is defined as exactly $6.02214076 \times 10^{23}$ particles, e.g., atoms, molecules, ions or electrons.
- There are $\mathbf{6.02214076 \times 10^{23}}$ particles in 1.00 mole. This number is called Avogadro's number.
- The molar mass of an element can be found by referring to the atomic mass on a periodic table with units of g/mol.
- Using dimensional analysis, it is possible to convert between grams, moles, and the number of atoms or molecules.

Further Reading/Supplemental Links

- learner.org/resources/series61.html - The ***learner.org*** website allows users to view streaming videos of the Annenberg series of chemistry videos. You are required to register before you can watch the videos, but there is no charge. The website has one video that relates to this lesson called ***The Mole***.
- Using Avogadro's law, the mass of a substance can be related to the number of particles contained in that mass. The Mole: (www.learner.org/vod/vod_window.html?pid=803)
- Vision Learning tutorial: The Mole http://visionlearning.com/library/mo...p?mid-53&1=&c3=

Contributions & Attributions

-
- Wikipedia

3.10: Counting Atoms by the Gram is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.3: Counting Atoms by the Gram** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.11: COUNTING MOLECULES BY THE GRAM

LEARNING OBJECTIVES

- Define molecular mass and formula mass.
- Perform conversions between mass and moles of a compound.
- Perform conversions between mass and number of particles.

MOLECULAR AND FORMULA MASSES

The molecular mass of a substance is the sum of the average masses of the atoms in one molecule of a substance. It is calculated by adding together the atomic masses of the elements in the substance, each multiplied by its subscript (written or implied) in the molecular formula. Because the units of atomic mass are atomic mass units, the units of molecular mass are also atomic mass units. The procedure for calculating molecular masses is illustrated in Example **3.11.1**.

EXAMPLE 3.11.1: ETHANOL

Calculate the molecular mass of ethanol, whose condensed structural formula is $\mathbf{CH_3CH_2OH}$. Among its many uses, ethanol is a fuel for internal combustion engines

Solution

Solutions to Example 6.4.1

Steps for Problem Solving	Calculate the molecular mass of ethanol, whose condensed structural formula is $\mathbf{CH_3CH_2OH}$
Identify the "given"information and what the problem is asking you to "find."	Given: Ethanol molecule (CH_3CH_2OH) Find: molecular mass
Determine the number of atoms of each element in the molecule.	The molecular formula of ethanol may be written in three different ways: • CH_3CH_2OH (which illustrates the presence of an ethyl group • CH_3CH_2–, and an −OH group) • C_2H_5OH, and C_2H_6O; All show that ethanol has two carbon atoms, six hydrogen atoms, and one oxygen atom.
Obtain the atomic masses of each element from the periodic table and multiply the atomic mass of each element by the number of atoms of that element.	1 C atom = 12.011 amu 1 H atom = 1.0079 amu 1 O atom = 15.9994 amu
Add the masses together to obtain the molecular mass.	2C: (2 atoms)(12.011amu/atom) = 24.022 amu 6H: (6 atoms)(1.0079amu/atom) = 6.0474amu +1O: (1 atoms)(15.9994amu/atom) =15.9994amu C_2H_6O : molecular mass of ethanol = 46.069amu

EXERCISE 3.11.1: FREON

Calculate the molecular mass of trichlorofluoromethane, also known as Freon-11, which has a condensed structural formula of $\mathbf{CCl_3F}$. Until recently, it was used as a refrigerant. The structure of a molecule of Freon-11 is as follows:




Figure **3.11.1**: Molecular structure of freon-11, $\mathbf{CCl_3F}$.

Answer

Answer

137.37 amu

Unlike molecules, which form covalent bonds, ionic compounds do not have a readily identifiable molecular unit. Therefore, for ionic compounds, the **formula mass** (also called the empirical formula mass) of the compound is used instead of the molecular mass. The formula mass is the sum of the atomic masses of all the elements in the empirical formula, each multiplied by its subscript (written or implied). It is directly analogous to the molecular mass of a covalent compound. The units are atomic mass units.

> *Atomic mass, molecular mass, and formula mass all have the same units: atomic mass units.*

EXAMPLE 3.11.2: CALCIUM PHOSPHATE

Calculate the formula mass of $\mathbf{Ca_3(PO_4)_2}$, commonly called calcium phosphate. This compound is the principal source of calcium found in bovine milk.

Solution

Solutions to Example 6.4.2

Steps for Problem Solving	Calculate the formula mass of $\mathbf{Ca_3(PO_4)_2}$, commonly called calcium phosphate.
Identify the "given" information and what the problem is asking you to "find."	Given: Calcium phosphate [$Ca_3(PO_4)_2$] formula unit Find: formula mass
Determine the number of atoms of each element in the molecule.	• The empirical formula—$Ca_3(PO_4)_2$—indicates that the simplest electrically neutral unit of calcium phosphate contains three Ca^{2+} ions and two PO_4^{3-} ions. • The formula mass of this molecular unit is calculated by adding together the atomic masses of three calcium atoms, two phosphorus atoms, and eight oxygen atoms.
Obtain the atomic masses of each element from the periodic table and multiply the atomic mass of each element by the number of atoms of that element.	1 Ca atom = 40.078 amu 1 P atom = 30.973761 amu 1 O atom = 15.9994 amu
Add together the masses to give the formula mass.	3Ca: (3 atoms) (40.078 amu/atom)=120.234amu 2P: (2 atoms) (30.973761amu/atom)=61.947522amu + 8O: (8 atoms)(15.9994amu/atom)=127.9952amu Formula mass of $Ca_3(PO_4)_2$=310.177amu

EXERCISE 3.11.2: SILICON NITRIDE

Calculate the formula mass of $\mathbf{Si_3N_4}$, commonly called silicon nitride. It is an extremely hard and inert material that is used to make cutting tools for machining hard metal alloys.



Figure 3.11.2: $\mathbf{Si_3N_4}$ bearing parts. (Public Domain; David W. Richerson and Douglas W. Freitag; Oak Ridge National Laboratory).

Answer

140.29 amu

MOLAR MASS

The molar mass of a substance is defined as the mass in grams of 1 mole of that substance. One mole of isotopically pure carbon-12 has a mass of 12 g. For an element, the molar mass is the mass of 1 mol of atoms of that element; for a covalent molecular compound, it is the mass of 1 mol of molecules of that compound; for an ionic compound, it is the mass of 1 mol of formula units. That is, the molar mass of a

substance is the mass (in grams per mole) of 6.022×10^{23} atoms, molecules, or formula units of that substance. In each case, the number of grams in 1 mol is the same as the number of atomic mass units that describe the atomic mass, the molecular mass, or the formula mass, respectively.

> *The molar mass of any substance is its atomic mass, molecular mass, or formula mass in grams per mole.*

The periodic table lists the atomic mass of carbon as 12.011 amu; the average molar mass of carbon—the mass of 6.022×10^{23} carbon atoms—is therefore 12.011 g/mol:

Table **3.11.1**: Molar Mass of Select Substances

Substance (formula)	Basic Unit	Atomic, Molecular, or Formula Mass (amu)	Molar Mass (g/mol)
carbon (C)	atom	12.011 (atomic mass)	12.011
ethanol (C_2H_5OH)	molecule	46.069 (molecular mass)	46.069
calcium phosphate [$Ca_3(PO_4)_2$]	formula unit	310.177 (formula mass)	310.177

CONVERTING BETWEEN GRAMS AND MOLES OF A COMPOUND

The molar mass of any substance is the mass in grams of one mole of representative particles of that substance. The representative particles can be atoms, molecules, or formula units of ionic compounds. This relationship is frequently used in the laboratory. Suppose that for a certain experiment you need 3.00 moles of calcium chloride ($\mathbf{CaCl_2}$). Since calcium chloride is a solid, it would be convenient to use a balance to measure the mass that is needed. Dimensional analysis will allow you to calculate the mass of $\mathbf{CaCl_2}$ that you should measure as shown in Example **3.11.3**.

EXAMPLE 3.11.3: CALCIUM CHLORIDE

Calculate the mass of 3.00 moles of calcium chloride ($CaCl_2$).



Figure **3.11.3**: Calcium chloride is used as a drying agent and as a road deicer.

Solution

Solutions to Example 6.4.3

Steps for Problem Solving	Calculate the mass of 3.00 moles of calcium chloride
Identify the "given" information and what the problem is asking you to "find."	Given: 3.00 moles of $\mathbf{CaCl_2}$ Find: g $\mathbf{CaCl_2}$
List other known quantities.	1 mol $\mathbf{CaCl_2}$ = 110.98 g $\mathbf{CaCl_2}$
Prepare a concept map and use the proper conversion factor.	mol $CaCl_2$ → g $CaCl_2$ $\frac{110.98\ g\ CaCl_2}{1\ mol\ CaCl_2}$
Cancel units and calculate.	$3.00\ \cancel{mol\ CaCl_2} \times \frac{110.98\ g\ CaCl_2}{1\ \cancel{mol\ CaCl_2}} = 333\ g\ CaCl_2$
Think about your result.	

EXERCISE 3.11.3: CALCIUM OXIDE

What is the mass of **7.50 mol** of (calcium oxide) **CaO**?

Answer

420.60 g

EXAMPLE 3.11.4: WATER

How many moles are present in 108 grams of water?

Solution

Solutions to Example 6.4.4

Steps for Problem Solving	How many moles are present in 108 grams of water?
Identify the "given" information and what the problem is asking you to "find."	Given: 108 g $\mathbf{H_2O}$ Find: mol $\mathbf{H_2O}$
List other known quantities.	$\mathbf{1\ mol\ H_2O = 18.02\ g}$ H2O
Prepare a concept map and use the proper conversion factor.	g H_2O → mol H_2O $\frac{1\ mol\ H_2O}{18.02\ g\ H_2O}$
Cancel units and calculate.	$\mathbf{108\ \cancel{g\ H_2O} \times \frac{1\ mol\ H_2O}{18.02\ \cancel{g\ H_2O}} = 5.99\ mol\ H_2O}$
Think about your result.	

EXERCISE 3.11.4: NITROGEN GAS

What is the mass of **7.50 mol** of Nitrogen gas $\mathbf{N_2}$?

Answer

210 g

CONVERSIONS BETWEEN MASS AND NUMBER OF PARTICLES

In "Conversions Between Moles and Mass", you learned how to convert back and forth between moles and the number of representative particles. Now you have seen how to convert back and forth between moles and mass of a substance in grams. We can combine the two types of problems into one. Mass and number of particles are both related to moles. To convert from mass to number of particles or vice-versa, it will first require a conversion to moles as shown in Figure **3.11.1** and Example **3.11.5**.





Figure **3.11.4**: Conversion from number of particles to mass, or from mass to number of particles, requires two steps. To convert from number of particles to moles, use mol/Avogrado's #, an d to convert from moles to mass, use g/mol.

EXAMPLE 3.11.5: CHLORINE

How many molecules is **20.0 g** of chlorine gas, Cl_2?

Solution

Solutions to Example 6.4.5

Steps for Problem Solving	How many molecules is **20.0 g** of chlorine gas, Cl_2?
Identify the "given" information and what the problem is asking you to "find."	Given: 20.0 g Cl_2 Find: # Cl_2 molecules
List other known quantities.	• 1 mol Cl_2 = 70.90 g Cl_2, • 1mol Cl_2 = 6.022×10^{23} Cl_2 molecules
Prepare a concept map and use the proper conversion factor.	 $\frac{1\ mol\ Cl_2}{70.90\ g\ Cl_2}$ $\frac{6.022 \times 10^{23}\ Cl_2\ molecules}{1\ mol\ Cl_2}$ The conversion factors are 1 mole Cl_2 over 70.90 grams Cl_2, and 6.022×10^{23} Cl_2 molecules over 1 mole Cl_2.
Cancel units and calculate.	$20.0\ \cancel{g\ Cl_2} \times \frac{1\ \cancel{mol\ Cl_2}}{70.90\ \cancel{g\ Cl_2}} \times \frac{6.02 \times 10^{23}\ molecules\ Cl_2}{1\ \cancel{mol\ Cl_2}}$ $= 1.70 \times 10^{23}\ molecules\ Cl_2$
Think about your result.	Since the given mass is less than half of the molar mass of chlorine, the resulting number of molecules is less than half of Avogadro's number.

EXERCISE 3.11.5: CALCIUM CHLORIDE

How many formula units are in 25.0 g of $CaCl_2$?

Answer

1.36 x 10^{23} $CaCl_2$ formula units

SUMMARY

- Calculations for formula mass and molecular mass are described.
- Calculations involving conversions between moles of a material and the mass of that material are described.
- Calculations are illustrated for conversions between mass and number of particles.

This page titled 3.11: Counting Molecules by the Gram is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.4: Counting Molecules by the Gram** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

3.12: Chemical Formulas as Conversion Factors

Learning Objectives

- Use chemical formulas as conversion factors.

Figure **3.12.1** shows that we need 2 hydrogen atoms and 1 oxygen atom to make one water molecule. If we want to make two water molecules, we will need 4 hydrogen atoms and 2 oxygen atoms. If we want to make five molecules of water, we need 10 hydrogen atoms and 5 oxygen atoms. The ratio of atoms we will need to make any number of water molecules is the same: 2 hydrogen atoms to 1 oxygen atom.




Figure **3.12.1** Water Molecules. The ratio of hydrogen atoms to oxygen atoms used to make water molecules is always 2:1, no matter how many water molecules are being made.


Using formulas to indicate how many atoms of each element we have in a substance, we can relate the number of moles of molecules to the number of moles of atoms. For example, in 1 mol of water (H_2O) we can construct the relationships given in (Table **3.12.1**).

Table **3.12.1**: Molecular Relationships for Water

1 Molecule of H_2O Has	1 Mol of H_2O Has	Molecular Relationships
2 H atoms	2 mol of H atoms	$\frac{2\text{ mol H atoms}}{1\text{ mol }H_2O\text{ molecules}}$ or $\frac{1\text{ mol }H_2O\text{ molecules}}{2\text{ mol H atoms}}$
1 O atom	1 mol of O atoms	$\frac{1\text{ mol O atoms}}{1\text{ mol }H_2O\text{ molecules}}$ or $\frac{1\text{ mol }H_2O\text{ molecules}}{1\text{ mol O atoms}}$

The Mole is big

A mole represents a very large number! The number 602,214,129,000,000,000,000,000 looks about twice as long as a trillion, which means it's about a trillion trillion.



(CC BY-SA NC; what if? [what-if.xkcd.com]).

A trillion trillion kilograms is how much a planet weighs. If 1 mol of quarters were stacked in a column, it could stretch back and forth between Earth and the sun *6.8 billion* times.

Table **3.12.2**: Molecular and Mass Relationships for Ethanol

1 Molecule of C_2H_6O Has	1 Mol of C_2H_6O Has	Molecular and Mass Relationships
2 C atoms	2 mol of C atoms	$\frac{2\text{ mol C atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{2\text{ mol C atoms}}$
6 H atoms	6 mol of H atoms	$\frac{6\text{ mol H atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{6\text{ mol H atoms}}$
1 O atom	1 mol of O atoms	$\frac{1\text{ mol O atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{1\text{ mol O atoms}}$
2 (12.01 amu) C 24.02 amu C	2 (12.01 g) C 24.02 g C	$\frac{24.02\text{ g C}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{24.02\text{ g C}}$
6 (1.008 amu) H 6.048 amu H	6 (1.008 g) H 6.048 g H	$\frac{6.048\text{ g H}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{6.048\text{ g H}}$
1 (16.00 amu) O 16.00 amu O	1 (16.00 g) O 16.00 g O	$\frac{16.00\text{ g O}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{16.00\text{ g O}}$

The following example illustrates how we can use the relationships in Table **3.12.2** as conversion factors.

Example 3.12.1: Ethanol

If a sample consists of 2.5 mol of ethanol (C_2H_6O), how many moles of carbon atoms does it have?

Solution

Solutions to Example 6.5.1

Steps for Problem Solving	If a sample consists of 2.5 mol of ethanol (C_2H_6O), how many moles of carbon atoms does it have?
Identify the "given" information and what the problem is asking you to "find."	Given: 2.5 mol C_2H_6O Find: mol C atoms
List other known quantities.	1 mol C_2H_6O = 2 mol C
Prepare a concept map and use the proper conversion factor.	mol C_2H_6O → mol C atom $\frac{2\text{ mol }C}{1\text{ mol }C_2H_6O}$
Cancel units and calculate.	Note how the unit *mol C_2H_6O molecules* cancels algebraically. $2.5\ \cancel{\text{mol }C_2H_6O\text{ molecules}} \times \frac{2\text{ mol C atoms}}{1\ \cancel{\text{mol }C_2H_6O\text{ molecules}}} = 5.0\text{ mol C atoms}$
Think about your result.	There are twice as many C atoms in one C_2H_6O molecule, so the final amount should be double.

Exercise 3.12.1

If a sample contains 6.75 mol of Na_2SO_4, how many moles of sodium atoms, sulfur atoms, and oxygen atoms does it have?

Answer

13.5 mol Na atoms, 6.75 mol S atoms, and 27.0 mol O atoms


The fact that 1 mol equals 6.022×10^{23} items can also be used as a conversion factor.

Example 3.12.2: Oxygen Mass

Determine the mass of Oxygen in 75.0g of C_2H_6O.

Solution

Solutions to Example 6.5.2

Steps for Problem Solving	Determine the mass of Oxygen in 75.0g of C_2H_6O
Identify the "given" information and what the problem is asking you to "find."	Given: 75.0g C_2H_6O Find: g O
List other known quantities.	1 mol O = 16.0g O 1 mol C_2H_6O = 1 mol O 1 mol C_2H_6O = 46.07g C_2H_6O
Prepare a concept map and use the proper conversion factor.	 The conversion factors are 1 mol C2H6O over 46.07 g C2H6O, 1 mol O over 1 mol C2H6O, and 16.00 g O over 1 mole O.
Cancel units and calculate.	$75.0\ g\ C_2H_6O \times \frac{1\ mol\ C_2H_6O}{46.07\ g\ C_2H_6O} \times \frac{1\ mol\ O}{1\ mol\ C_2H_6O} \times \frac{16.00\ g\ O}{1\ mol\ O} = 26.0\ g$
Think about your result.	

Exercise 3.12.2

a. How many molecules are present in 16.02 mol of C_4H_{10}? How many C atoms are in 16.02 mol?
b. How many moles of each type of atom are in 2.58 mol of Na_2SO_4?

Answer a:

9.647×10^{24} C_4H_{10} molecules and 3.859×10^{25} C atoms

Answer b:

5.16 mol Na atoms, 2.58 mol S atoms, and 10.3 mol O atoms

Summary

In any given formula, the ratio of the number of moles of molecules (or formula units) to the number of moles of atoms can be used as a conversion factor.

3.12: Chemical Formulas as Conversion Factors is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.5: Chemical Formulas as Conversion Factors** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.13: Mass Percent Composition of Compounds

Learning Objectives

- Determine percent composition of each element in a compound based on mass.

Packaged foods that you eat typically have nutritional information provided on the label. The label on a jar of peanut butter reveals that one serving size is considered to be $32\ \text{g}$. The label also gives the masses of various types of compounds that are present in each serving. One serving contains $7\ \text{g}$ of protein, $15\ \text{g}$ of fat, and $3\ \text{g}$ of sugar. By calculating the fraction of protein, fat, or sugar in one serving size of peanut butter and converting to percent values, we can determine the composition of peanut butter on a percent by mass basis.

Percent Composition

Chemists often need to know what elements are present in a compound and in what percentage. The **percent composition** is the percent by mass of each element in a compound. It is calculated in a similar way to that of the composition of the peanut butter.

$$\%\ \text{by mass} = \frac{\text{mass of element}}{\text{mass of compound}} \times 100\%$$

The sample problem below shows the calculation of the percent composition of a compound based on mass data.

Example 3.13.1: Percent Composition from Mass Data

A certain newly synthesized compound is known to contain the elements zinc and oxygen. When a $20.00\ \text{g}$ sample of the compound is decomposed, $16.07\ \text{g}$ of zinc remains. Determine the percent composition of the compound.

Solution

Solutions to Example 6.6.1

Steps for Problem Solving	When a 20.00 g sample of the zinc-and-oxygen compound is decomposed, 16.07 g of zinc remains. Determine the percent composition of the compound.
Identify the "given" information and what the problem is asking you to "find."	Given : Mass of compound = 20.00 g Mass of Zn = 16.07 g Find: % Composition (% Zn and %O)
List other known quantities.	Subtract to find the mass of oxygen in the compound. Divide each element's mass by the mass of the compound to find the percent by mass. Mass of oxygen = 20.00 g - 16.07 g = 3.93 g O
Cancel units and calculate.	$\%\ \text{Zn} = \frac{16.07\ \text{g Zn}}{20.00\ \text{g}} \times 100\% = 80.35\%\ \text{Zn}$ $\%\ \text{O} = \frac{3.93\ \text{g O}}{20.00\ \text{g}} \times 100\% = 19.65\%\ \text{O}$ Calculate the percent by mass of each element by dividing the mass of that element by the mass of the compound and multiplying by 100%.
Think about your result.	The calculations make sense because the sum of the two percentages adds up to 100%. By mass, the compound is mostly zinc.

? Exercise 3.13.1

Sulfuric acid, H_2SO_4 is a very useful chemical in industrial processes. If 196.0 g of sulfuric acid contained 64.0g oxygen and 4.0 g of hydrogen, what is the percent composition of the compound?

Answer

2.04% H, 32.65% S, and 65.3% O

Summary

- Processes are described for calculating the percent composition of a compound based on mass.

3.13: Mass Percent Composition of Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.6: Mass Percent Composition of Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.14: Mass Percent Composition from a Chemical Formula

Learning Objectives

- Determine the percent composition of each element in a compound from the chemical formula.

The percent composition of a compound can also be determined from the formula of the compound. The subscripts in the formula are first used to calculate the mass of each element in one mole of the compound. This is divided by the molar mass of the compound and multiplied by 100%.

$$\%\text{ by mass} = \frac{\text{mass of element in 1 mol}}{\text{molar mass of compound}} \times 100\%$$

The percent composition of a given compound is always the same, given that the compound is pure.

Example 3.14.1

Dichlorine heptoxide (Cl_2O_7) is a highly reactive compound used in some organic synthesis reactions. Calculate the percent composition of dichlorine heptoxide.

Solution

Solutions to Example 6.7.1

Steps for Problem Solving	**Calculate the percent composition of dichlorine heptoxide (Cl_2O_7).**
Identify the "given" information and what the problem is asking you to "find."	Given : Cl_2O_7 Find: % Composition (% Cl and %O)
List other known quantities.	Mass of Cl in 1 mol Cl_2O_7 , 2 Cl : 2 x 35.45 g = 70.90 g Mass of O in 1 mol Cl_2O_7 , 7 O: 7 x 16.00 g = 112.00 g Molar mass of Cl_2O_7 = 182.90 g/mol
Cancel units and calculate.	$$\%Cl = \frac{70.90\text{ g Cl}}{182.90\text{ g}} \times 100\% = 38.76\%\text{ Cl}$$ $$\%\text{ O} = \frac{112.00\text{ g O}}{182.90\text{ g}} \times 100\% = 61.24\%\text{ O}$$ Calculate the percent by mass of each element by dividing the mass of that element in 1 mole of the compound by the molar mass of the compound and multiplying by 100%.
Think about your result.	The percentages add up to 100%.

Percent composition can also be used to determine the mass of a certain element that is contained in any mass of a compound. In the previous sample problem, it was found that the percent composition of dichlorine heptoxide is 38.76% Cl and 61.24% O. Suppose that you needed to know the masses of chlorine and oxygen present in a 12.50 g sample of dichlorine heptoxide. You can set up a conversion factor based on the percent by mass of each element.

$$12.50\text{ g } Cl_2O_7 \times \frac{38.76\text{ g Cl}}{100\text{ g } Cl_2O_7} = 4.845\text{ g Cl}$$

$$12.50\text{ g } Cl_2O_7 \times \frac{61.24\text{ g O}}{100\text{ g } Cl_2O_7} = 7.655\text{ g O}$$

The sum of the two masses is 12.50 g, the mass of the sample size.

? Exercise 3.14.1

Barium fluoride is a transparent crystal that can be found in nature as the mineral frankdicksonite. Determine the percent composition of barium fluoride.

Answer a:

78.32% Ba and 21.67% F

Summary

- Processes are described for calculating the percent composition of a compound based on the chemical formula.

3.14: Mass Percent Composition from a Chemical Formula is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.7: Mass Percent Composition from a Chemical Formula** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.15: Calculating Empirical Formulas for Compounds

Learning Objectives

- Define empirical formula.
- Determine empirical formula from percent composition of a compound.

In the early days of chemistry, there were few tools for the detailed study of compounds. Much of the information regarding the composition of compounds came from the elemental analysis of inorganic materials. The "new" field of organic chemistry (the study of carbon compounds) faced the challenge of not being able to characterize a compound completely. The relative amounts of elements could be determined, but so many of these materials had carbon, hydrogen, oxygen, and possibly nitrogen in simple ratios. We did not know exactly how many of these atoms were actually in a specific molecule.

Determining Empirical Formulas

An empirical formula tells us the relative ratios of different atoms in a compound. The ratios hold true on the *molar* level as well. Thus, H_2O is composed of two atoms of hydrogen and 1 atom of oxygen. Likewise, **1.0 mole of H_2O** is composed **of 2.0 moles of hydrogen** and **1.0 mole of oxygen**. We can also work backwards from molar ratios because *if we know the molar amounts of each element in a compound, we can determine the empirical formula.*

> *In a procedure called elemental analysis, an unknown compound can be analyzed in the laboratory in order to determine the percentages of each element contained within it. These percentages can be transformed into the mole ratio of the elements, which leads to the empirical formula.*

Empirical Formula: In Steps

Steps to determine empirical formula:

1. Assume a **100 g** sample of the compound so that the given percentages can be directly converted into grams.
2. Use each element's molar mass to convert the grams of each element to moles.
3. In order to find a whole-number ratio, divide the moles of each element by whichever of the moles from step 2 is the smallest.
4. If all the moles at this point are whole numbers (or very close), the empirical formula can be written with the moles as the subscript of each element.
5. In some cases, one or more of the moles calculated in step 3 will not be whole numbers. Multiply each of the moles by the smallest whole number that will convert each into a whole number. Write the empirical formula.

Example 3.15.1

A compound of iron and oxygen is analyzed and found to contain **69.94%** iron and **30.06%** oxygen. Find the empirical formula of the compound.

Solution

Solutions to Example 6.8.1

Steps for Problem Solving	**Find the empirical formula of a compound of 69.94% iron and 30.06% oxygen.**
Identify the "given" information and what the problem is asking you to "find."	Given: % of $\mathbf{Fe} = 69.94\%$ % of $\mathbf{O} = 30.06\%$ Find: Empirical formula $= \mathbf{Fe_?O_?}$
Calculate	
a. Assume a $\mathbf{100\ g}$ sample, convert the same % values to grams.	$69.94\ \mathbf{g\ Fe}$ $30.06\ \mathbf{g\ O}$
b. Convert to moles.	$69.94\ \mathbf{g\ Fe} \times \frac{1\ \mathbf{mol\ Fe}}{55.85\ \mathbf{g\ Fe}} = 1.252\ \mathbf{mol\ Fe}$ $30.06\ \mathbf{g\ O} \times \frac{1\ \mathbf{mol\ O}}{16.00\ \mathbf{g\ O}} = 1.879\ \mathbf{mol\ O}$
c. Divide both moles by the smallest of the results.	$\mathbf{Fe}: \frac{1.252\ \mathbf{mol}}{1.252}$ $\mathbf{O}: \frac{1.879\ \mathbf{mol}}{1.252}$ The "non-whole number" empirical formula of the compound is $\mathbf{Fe_1O_{1.5}}$
Multiply each of the moles by the smallest whole number that will convert each into a whole number.	Fe:O = 2 (1:1.5) = 2:3 Since the moles of $\mathbf{O}$ is still not a whole number, both moles can be multiplied by 2, while rounding to a whole number.
Write the empirical formula.	The empirical formula of the compound is $\mathbf{Fe_2O_3}$.
Think about your result.	The subscripts are whole numbers and represent the mole ratio of the elements in the compound. The compound is the ionic compound iron (III) oxide.

? Exercise 3.15.1

Mercury forms a compound with chlorine that is 73.9% mercury and 26.1% chlorine by mass. What is the empirical formula?

Answer

$HgCl_2$

Summary

- A process is described for the calculation of the empirical formula of a compound, based on the percent composition of that compound.

3.15: Calculating Empirical Formulas for Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.8: Calculating Empirical Formulas for Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.16: Calculating Molecular Formulas for Compounds

Learning Objectives

- Understand the difference between empirical formulas and molecular formulas.
- Determine molecular formula from percent composition and molar mass of a compound.

Below, we see two carbohydrates: glucose and sucrose. Sucrose is almost exactly twice the size of glucose, although their empirical formulas are very similar. Some people can distinguish them on the basis of taste, but it's not a good idea to go around tasting chemicals. The best way to tell glucose and sucrose apart is to determine the molar masses—this approach allows you to easily tell which compound is which.




Figure **3.16.1**: (A) the molecular structure of glucose and (B) the molecular structure of sucrose.

Molecular Formulas

Molecular formulas give the kind and number of atoms of each element present in the molecular compound. In many cases, the molecular formula is the same as the empirical formula. The chemical formula will always be some *integer multiple* (n) of the empirical formula (i.e. integer multiples of the subscripts of the empirical formula).

$$\text{Molecular Formula} = n(\text{Empirical formula})$$

therefore

$$n = \frac{\text{Molecular Formula}}{\text{Empirical Formula}}$$

The integer multiple, n, can also be obtained by dividing the molar mass, MM, of the compound by the empirical formula mass, EFM (the molar mass represented by the empirical formula).

$$n = \frac{MM(molar mass)}{EFM(empirical formula molar mass)}$$

Table **3.16.1** shows the comparison between the empirical and molecular formula of methane, acetic acid, and glucose, and the different values of n. The molecular formula of methane is CH_4 and because it contains only one carbon atom, that is also its empirical formula. Sometimes, however, the molecular formula is a simple whole number multiple of the empirical formula. Acetic acid is an organic acid that is the main component of vinegar. Its molecular formula is $C_2H_4O_2$. Glucose is a simple sugar that cells use as a primary source of energy. Its molecular formula is $C_6H_{12}O_6$. The structures of both molecules are shown in Figure **3.16.2**. They are very different compounds, yet both have the same empirical formula of CH_2O.

Table **3.16.1**: Molecular Formula and Empirical Formula of Various Compounds.

Name of Compound	Molecular Formula	Empirical Formula	n
Methane	CH_4	CH_4	1
Acetic acid	$C_2H_4O_2$	CH_2O	2
Glucose	$C_6H_{12}O_6$	CH_2O	6




Figure 3.16.2: Acetic acid (left) has a molecular formula of $C_2H_4O_2$, while glucose (right) has a molecular formula of $C_6H_{12}O_6$. Both have the empirical formula CH_2O.

Empirical formulas can be determined from the percent composition of a compound as discussed in section 6.8. In order to determine its molecular formula, it is necessary to know the molar mass of the compound. Chemists use an instrument called a mass spectrometer to determine the molar mass of compounds. In order to go from the empirical formula to the molecular formula, follow these steps:

1. Calculate the **empirical formula molar mass (EFM)**.
2. Divide the molar mass of the compound by the empirical formula molar mass. The result should be a whole number or very close to a whole number.
3. Multiply all the subscripts in the empirical formula by the whole number found in step 2. The result is the molecular formula.

Example 3.16.1

The empirical formula of a compound of boron and hydrogen is BH_3. Its molar mass is $27.7\ g/mol$. Determine the molecular formula of the compound.

Solution

Solutions to Example 6.9.1

Steps for Problem Solving	Determine the molecular formula of BH_3.
Identify the "given" information and what the problem is asking you to "find."	Given: Empirical formula $= BH_3$ Molar mass $= 27.7\ g/mol$ Find: Molecular formula $=?$
Calculate the empirical formula mass (EFM).	$\text{Empirical formula molar mass (EFM)} = 13.84\ g/mol$
Divide the molar mass of the compound by the empirical formula mass. The result should be a whole number or very close to a whole number.	$\frac{\text{molar mass}}{\text{EFM}} = \frac{27.7 g/mol}{13.84 g/mol} = 2$
Multiply all the subscripts in the empirical formula by the whole number found in step 2. The result is the molecular formula.	$BH_3 \times 2 = B_2H_6$
Write the molecular formula.	The molecular formula of the compound is B_2H_6.
Think about your result.	The molar mass of the molecular formula matches the molar mass of the compound.

Exercise 3.16.1

Vitamin C (ascorbic acid) contains 40.92 % C, 4.58 % H, and 54.50 % O, by mass. The experimentally determined molecular mass is 176 amu. What are the empirical and chemical formulas for ascorbic acid?

Answer Empirical Formula

$C_3H_4O_3$

Answer Molecular Formula

$C_6H_8O_6$

Summary

- A procedure is described that allows the calculation of the exact molecular formula for a compound.

3.16: Calculating Molecular Formulas for Compounds is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **6.9: Calculating Molecular Formulas for Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.17: Making Pancakes- Relationships Between Ingredients

3.17: Making Pancakes- Relationships Between Ingredients is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

3.18: MAKING MOLECULES- MOLE-TO-MOLE CONVERSIONS

/;LEARNING OBJECTIVES

- Use a balanced chemical equation to determine molar relationships between substances.

Previously, you learned to balance chemical equations by comparing the numbers of each type of atom in the reactants and products. The coefficients in front of the chemical formulas represent the numbers of molecules or formula units (depending on the type of substance). As follows, we will extend the meaning of the coefficients in a chemical equation.

Consider the simple chemical equation:

$$2H_2 + O_2 \rightarrow 2H_2O$$

The convention for writing balanced chemical equations is to use the lowest whole-number ratio for the coefficients. However, the equation is balanced as long as the coefficients are in a 2:1:2 ratio. For example, this equation is also balanced if we write it as

$$4H_2 + 2O_2 \rightarrow 4H_2O$$

The ratio of the coefficients is 4:2:4, which reduces to 2:1:2. The equation is also balanced if we were to write it as

$$22H_2 + 11O_2 \rightarrow 22H_2O$$

because 22:11:22 also reduces to 2:1:2.

Suppose we want to use larger numbers. Consider the following coefficients:

$$12.044 \times 10^{23}\ \mathrm{H_2} + 6.022 \times 10^{23}\ \mathrm{O_2} \rightarrow 12.044 \times 10^{23}\ \mathrm{H_2O}$$

These coefficients also have the ratio 2:1:2 (check it and see), so this equation is balanced. But 6.022×10^{23} is 1 mol, while 12.044×10^{23} is 2 mol (and the number is written that way to make this more obvious), so we can simplify this version of the equation by writing it as

$$2\ mol\ \mathrm{H_2} + 1\ mol\ \mathrm{O_2} \rightarrow 2\ mol\ \mathrm{H_2O}$$

We can leave out the word *mol* and not write the 1 coefficient (as is our habit), so the final form of the equation, still balanced, is

$$2\,\mathrm{H_2} + \mathrm{O_2} \longrightarrow 2\,\mathrm{H_2O}$$

Now we interpret the coefficients as referring to molar amounts, not individual molecules. The lesson? *Balanced chemical equations are balanced not only at the molecular level, but also in terms of molar amounts of reactants and products.* Thus, we can read this reaction as "two moles of hydrogen react with one mole of oxygen to produce two moles of water."

By the same token, the ratios we constructed to describe a molecular reaction can also be constructed in terms of moles rather than molecules. For the reaction in which hydrogen and oxygen combine to make water, for example, we can construct the following ratios:

$$\frac{2\ \mathrm{mol\ H_2}}{1\ \mathrm{mol\ O_2}} \text{ or } \frac{1\ \mathrm{mol\ O_2}}{2\ \mathrm{mol\ H_2}}$$

$$\frac{2\ \mathrm{mol\ H_2O}}{1\ \mathrm{mol\ O_2}} \text{ or } \frac{1\ \mathrm{mol\ O_2}}{2\ \mathrm{mol\ H_2O}}$$

$$\frac{2\ \mathrm{mol\ H_2}}{2\ \mathrm{mol\ H_2O}} \text{ or } \frac{2\ \mathrm{mol\ H_2O}}{2\ \mathrm{mol\ H_2}}$$


We can use these ratios to determine what amount of a substance, in moles, will react with or produce a given number of moles of a different substance. The study of the numerical relationships between the reactants and the products in balanced chemical reactions is called *stoichiometry*.

EXAMPLE 3.18.1

How many moles of oxygen react with hydrogen to produce 27.6 mol of $\mathrm{H_2O}$?

Solution

Solutions to Example 8.3.1

Steps for Problem Solving	How many moles of oxygen react with hydrogen to produce 27.6 mol of H_2O?
Find a balanced equation that describes the reaction.	Unbalanced: $H_2 + O_2 \rightarrow H_2O$ Balanced: $\underline{2}H_2 + O_2 \rightarrow \underline{2}H_2O$
Identify the "given" information and what the problem is asking you to "find."	Given: moles H_2O Find: moles oxygen
List other known quantities.	1 mol O_2 = 2 mol H_2O
Prepare a concept map and use the proper conversion factor.	 $\frac{1\ mol\ O_2}{2\ mol\ H_2O}$
Cancel units and calculate.	$\cancel{27.6\ mol\ H_2O} \times \frac{1\ mol\ O_2}{\cancel{2\ mol\ H_2O}} = 13.8\ mol\ O_2$ To produce 27.6 mol of H_2O, 13.8 mol of O_2 react.
Think about your result.	Since each mole of oxygen produces twice as many moles of water, it makes sense that the produced amount is greater than the reactant amount

EXAMPLE 3.18.2

How many moles of ammonia are produced if 4.20 moles of hydrogen are reacted with an excess of nitrogen?

Solution

Solutions to Example 8.3.2

Steps for Problem Solving	How many moles of ammonia are produced if 4.20 moles of hydrogen are reacted with an excess of nitrogen?
Find a balanced equation that describes the reaction.	Unbalanced: $N_2 + H_2 \rightarrow NH_3$ Balanced: $N_2 + \underline{3}H_2 \rightarrow \underline{2}NH_3$
Identify the "given" information and what the problem is asking you to "find."	Given: $\mathbf{H_2 = 4.20\ mol}$ Find: mol of $\mathbf{NH_3}$
List other known quantities.	3 mol H_2 = 2 mol NH_3
Prepare a concept map and use the proper conversion factor.	 $\frac{2\ mol\ NH_3}{3\ mol\ H_2}$
Cancel units and calculate.	$\cancel{4.20\ mol\ H_2} \times \frac{2\ mol\ NH_3}{\cancel{3\ mol\ H_2}} = 2.80\ mol\ NH_3$ The reaction of **4.20 mol** of hydrogen with excess nitrogen produces **2.80 mol** of ammonia.
Think about your result.	The result corresponds to the 3:2 ratio of hydrogen to ammonia from the balanced equation.

EXERCISE 3.18.3

a. Given the following balanced chemical equation:

$$\mathbf{C_5H_{12} + 8\,O_2 \longrightarrow 5\,CO_2 + 6\,H_2O}$$

, How many moles of $\mathbf{H_2O}$ can be formed if 0.0652 mol of $\mathbf{C_5H_{12}}$ were to react?

b. Balance the following unbalanced equation and determine how many moles of $\mathbf{H_2O}$ are produced when 1.65 mol of NH_3 react:

$$\mathbf{NH_3 + O_2 \longrightarrow N_2 + H_2O}$$

Answer a

0.391 mol H_2O

Answer b

$4NH_3 + 3O_2 \rightarrow 2N_2 + 6H_2O$; 2.48 mol H_2O

SUMMARY

- The balanced chemical reaction can be used to determine molar relationships between substances.

3.18: Making Molecules- Mole-to-Mole Conversions is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **8.3: Making Molecules- Mole-to-Mole Conversions** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.


3.19: MAKING MOLECULES- MASS-TO-MASS CONVERSIONS

LEARNING OBJECTIVES

- Convert from mass or moles of one substance to mass or moles of another substance in a chemical reaction.

MOLE TO MASS CONVERSIONS

We have established that a balanced chemical equation is balanced in terms of moles, as well as atoms or molecules. We have used balanced equations to set up ratios, in terms of moles of materials, that we can use as conversion factors to answer stoichiometric questions—such as how many moles of substance A react with so many moles of reactant B. We can extend this technique even further. Recall that we can relate a molar amount to a mass amount using molar mass. We can use that relation to answer stoichiometry questions in terms of the masses of a particular substance, in addition to moles. We do this using the following sequence:



Collectively, these conversions are called mole-mass calculations. Flowchart of mole mass calculations: To convert from moles of substance A to moles of substance B, use the mole ratio conversion factor, and to convert from moles to grams of substance B, use molar mass conversion factor

As an example, consider the balanced chemical equation

$$\mathrm{Fe_2O_3 + 3\,SO_3 \rightarrow Fe_2(SO_4)_3} \tag{3.19.1}$$

If we have 3.59 mol of $\mathrm{Fe_2O_3}$, how many grams of $\mathrm{SO_3}$ can react with it? Using the mole-mass calculation sequence, we can determine the required mass of $\mathrm{SO_3}$ in two steps. First, we construct the appropriate molar ratio, determined from the balanced chemical equation, to calculate the number of moles of $\mathrm{SO_3}$ needed. Then, using the molar mass of $\mathrm{SO_3}$ as a conversion factor, we determine the mass that this number of moles of $\mathrm{SO_3}$ has.

As usual, we start with the quantity we were given:

$$3.59\ \cancel{\mathrm{mol\ Fe_2O_3}} \times \left(\frac{3\ \mathrm{mol\ SO_3}}{1\ \cancel{\mathrm{mol\ Fe_2O_3}}}\right) = 10.77\ \mathrm{mol\ SO_3} \tag{3.19.2}$$

The mol $\mathrm{Fe_2O_3}$ units cancel, leaving mol $\mathrm{SO_3}$ unit. Now, we take this answer and convert it to grams of $\mathrm{SO_3}$, using the molar mass of $\mathrm{SO_3}$ as the conversion factor:

$$10.77\ \cancel{\mathrm{mol\ SO_3}} \times \left(\frac{80.06\ \mathrm{g\ SO_3}}{1\ \cancel{\mathrm{mol\ SO_3}}}\right) = 862\ \mathrm{g\ SO_3} \tag{3.19.3}$$

Our final answer is expressed to three significant figures. Thus, in a two-step process, we find that 862 g of $\mathrm{SO_3}$ will react with 3.59 mol of $\mathrm{Fe_2O_3}$. Many problems of this type can be answered in this manner.

The same two-step problem can also be worked out in a single line, rather than as two separate steps, as follows:

$$3.59\ \cancel{mol\ Fe_2O_3} \times \underbrace{\left(\frac{3\ \cancel{mol\ SO_3}}{1\ \cancel{mol\ Fe_2O_3}}\right)}_{\text{converts to moles of } SO_3} \times \underbrace{\left(\frac{80.06\ g\ SO_3}{1\ \cancel{mol\ SO_3}}\right)}_{\text{converts to grams of } SO_3} = 862\ g\ SO_3$$

We get exactly the same answer when combining all math steps together.


EXAMPLE 3.19.1: GENERATION OF ALUMINUM OXIDE

How many moles of $\mathbf{HCl}$ will be produced when 249 g of $\mathbf{AlCl_3}$ are reacted according to this chemical equation?

$$\mathbf{2\,AlCl_3 + 3\,H_2O(\ell) \longrightarrow Al_2O_3 + 6\,HCl\,(g)}$$

Solution

Solutions to Example 8.5.1

Steps for Problem Solving	Example 3.19.1
Identify the "given" information and what the problem is asking you to "find."	Given: 249 g AlCl3 Find: moles HCl
List other known quantities.	1 mol AlCl3 = 133.33 g AlCl3 6 mol of HCl to 2 mol $AlCl_3$
Prepare a concept map and use the proper conversion factor.	 Flowchart of needed conversion factors: 1 mole AlCl3 to 133.33 grams AlCl3, and 6 moles HCl to 2 moles AlCl3
Cancel units and calculate.	$249\ \cancel{g\ AlCl_3} \times \frac{1\ \cancel{mol\ AlCl_3}}{133.33\ \cancel{g\ AlCl_3}} \times \frac{6\,mol\,HCl}{2\ \cancel{mol\ AlCl_3}} = 5.60\,mol\,HCl$
Think about your result.	Since 249 g of $AlCl_3$ is less than 266.66 g, the mass for 2 moles of $AlCl_3$ and the relationship is 6 mol of HCl to 2 mol $AlCl_3$, the answer should be less than 6 moles of HCl.

EXERCISE 3.19.1: GENERATION OF ALUMINUM OXIDE

How many moles of $\mathbf{Al_2O_3}$ will be produced when 23.9 g of $\mathbf{H_2O}$ are reacted according to this chemical equation?


$$\mathbf{2\,AlCl_3 + 3\,H_2O(\ell) \longrightarrow Al_2O_3 + 6\,HCl\,(g)}$$

Answer

0.442 mol Al2O3

MASS TO MASS CONVERSIONS

It is a small step from mole-mass calculations to mass-mass calculations. If we start with a known mass of one substance in a chemical reaction (instead of a known number of moles), we can calculate the corresponding masses of other substances in the reaction. The first step in this case is to convert the known mass into moles, using the substance's molar mass as the conversion factor. Then—and only then—we use the balanced chemical equation to construct a conversion factor to convert that quantity to moles of another substance, which in turn can be converted to a corresponding mass. Sequentially, the process is as follows:



This three-part process can be carried out in three discrete steps or combined into a single calculation that contains three conversion factors. The following example illustrates both techniques. Flowchart of mole mass calculations: To convert from grams to moles of substance A, use molar mass conversion factor; To convert from moles of substance A to moles of substance B, use the mole ratio conversion factor, and to convert from moles to grams of substance B, use molar mass conversion factor



✔ EXAMPLE 3.19.2: DECOMPOSITION OF AMMONIUM NITRATE

Ammonium nitrate decomposes to dinitrogen monoxide and water according to the following equation.

$$\mathrm{NH_4NO_3}\,(s) \rightarrow \mathrm{N_2O}\,(g) + 2\mathrm{H_2O}\,(l)$$

In a certain experiment, **45.7 g** of ammonium nitrate is decomposed. Find the mass of each of the products formed.

Solutions to Example 8.5.2

Steps for Problem Solving	Example 3.19.2
Identify the "given" information and what the problem is asking you to "find."	Given: **45.7 g NH_4NO_3** Find: Mass **N_2O =? g** Mass **H_2O =? g**
List other known quantities.	1 mol **NH_4NO_3 = 80.06 g** 1 mol **N_2O = 44.02 g** 1 mol **H_2O = 18.02 g** 1 mol NH_4NO_3 to 1 mol N_2O to 2 mol H_2O
Prepare two concept maps and use the proper conversion factor.	 $\frac{1\ mol\ NH_4NO_3}{80.06g\ NH_4NO_3}$ $\frac{1\ mol\ N_2O}{1\ mol\ NH_4NO_3}$ $\frac{44.02g\ N_2O}{1\ mol\ N_2O}$ Flowchart of conversion factors: 1 mole NH4NO3 to 80.06 grams NH4NO3, 1 mole N2O to 1 mole NH4NO3, 44.02 grams N2O to 1 mole N2O  $\frac{1\ mol\ NH_4NO_3}{80.06g\ NH_4NO_3}$ $\frac{2\ mol\ H_2O}{1\ mol\ NH_4NO_3}$ $\frac{18.02g\ H_2O}{1\ mol\ H_2O}$ Flowchart of conversion factors: 1 mole NH4NO3 to 80.06 grams NH4NO3, 2 moles H2O to 1 mole NH4NO3, 18.02 grams H2O to 1 mole H2O
Cancel units and calculate.	$45.7\ g\ NH_4NO_3 \times \frac{1\ mol\ NH_4NO_3}{80.06\ g\ NH_4NO_3} \times \frac{1\ mol\ N_2O}{1\ mol\ NH_4NO_3} \times \frac{44.02\ g\ N_2O}{1\ mol\ N_2O} = 25.1\ g\ N_2O$ $45.7\ g\ NH_4NO_3 \times \frac{1\ mol\ NH_4NO_3}{80.06\ g\ NH_4NO_3} \times \frac{2\ mol\ H_2O}{1\ mol\ NH_4NO_3} \times \frac{18.02\ g\ H_2O}{1\ mol\ H_2O} = 20.6\ g\ H_2O$
Think about your result.	The total mass of the two products is equal to the mass of ammonium nitrate which decomposed, demonstrating the law of conservation of mass. Each answer has three significant figures.

? EXERCISE 3.19.2: CARBON TETRACHLORIDE

Methane can react with elemental chlorine to make carbon tetrachloride (CCl_4). The balanced chemical equation is as follows:

$$CH_4(g) + 4\,Cl_2(g) \longrightarrow CCl_2(l) + 4\,HCl(l)$$

How many grams of HCl are produced by the reaction of 100.0 g of CH_4?

Answer

908.7g HCl

SUMMARY

- Calculations involving conversions between moles of a substance and the mass of that substance can be done using conversion factors.
- A balanced chemical reaction can be used to determine molar and mass relationships between substances.

3.19: Making Molecules- Mass-to-Mass Conversions is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **8.4: Making Molecules- Mass-to-Mass Conversions** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

3.20: Stoichiometry

Learning Objectives

- Explain the meaning of the term "stoichiometry".
- Determine the relative amounts of each substance in chemical equations.

You have learned that chemical equations provide us with information about the types of particles that react to form products. Chemical equations also provide us with the relative number of particles and moles that react to form products. In this section you will explore the quantitative relationships that exist between the quantities of reactants and products in a balanced equation. This is known as **stoichiometry**.

Stoichiometry, by definition, is the calculation of the quantities of reactants or products in a chemical reaction using the relationships found in the balanced chemical equation. The word stoichiometry is actually Greek from two words: *στοικηιον*, which means "element", and \(\mu \epsilon \tau \rho \omicron \nu), which means "measure".

Interpreting Chemical Equations

The mole, as you remember, is a quantitative measure that is equivalent to Avogadro's number of particles. So how does this relate to the chemical equation? Look at the chemical equation below.

$$2CuSO_4 + 4KI \rightarrow 2CuI + 2K_2SO_4 + I_2$$

The coefficients used, as we have learned, tell us the relative amounts of each substance in the equation. So for every 2 units of copper (II) sulfate ($CuSO_4$) we have, we need to have 4 units of potassium iodide (KI). For every two dozen copper (II) sulfates, we need 4 dozen potassium iodides. Because the unit "mole" is also a counting unit, we can interpret this equation in terms of moles, as well: For every two moles of copper (II) sulfate, we need 4 moles potassium iodide.

The production of ammonia (NH_3) from nitrogen and hydrogen gases is an important industrial reaction called the Haber process, after German chemist Fritz Haber.

$$N_2\,(g) + 3H_2\,(g) \rightarrow 2NH_3\,(g)$$

The balanced equation can be analyzed in several ways, as shown in the figure below.




Figure 3.20.1: This representation of the production of ammonia from nitrogen and hydrogen show several ways to interpret the quantitative information of a chemical reaction.

We see that 1 molecule of nitrogen reacts with 3 molecules of hydrogen to form 2 molecules of ammonia. This is the smallest possible relative amount of the reactants and products. To consider larger relative amounts, each coefficient can be multiplied by the same number. For example, 10 molecules of nitrogen would react with 30 molecules of hydrogen to produce 20 molecules of ammonia.

The most useful quantity for counting particles is the mole. So if each coefficient is multiplied by a mole, the balanced chemical equation tells us that 1 mole of nitrogen reacts with 3 moles of hydrogen to produce 2 moles of ammonia. This is the conventional way to interpret any balanced chemical equation.

Finally, if each mole quantity is converted to grams by using the molar mass, we can see that the law of conservation of mass is followed. $1\ \text{mol}$ of nitrogen has a mass of $28.02\ \text{g}$, while $3\ \text{mol}$ of hydrogen has a mass of $6.06\ \text{g}$, and $2\ \text{mol}$ of ammonia has a mass of $34.08\ \text{g}$.

$$28.02\ \text{g}\ \text{N}_2 + 6.06\ \text{g}\ \text{H}_2 \rightarrow 34.08\ \text{g}\ \text{NH}_3$$

Mass and the number of atoms must be conserved in any chemical reaction. The number of molecules is not necessarily conserved.

Example 3.20.1

The equation for the combustion of ethane (C_2H_6) is

$$2_2\text{CH}_6 + 7\text{O}_2 \rightarrow 4\text{CO}_2 + 6\text{H}_2\text{O}$$

a. Indicate the number of formula units or molecules in the balanced equation.
b. Indicate the number of moles present in the balanced equation.

Solution

a. Two molecules of C_2H_6 plus seven molecules of O_2 yields four molecules of CO_2 plus six molecules of H_2O.
b. Two moles of C_2H_6 plus seven moles of O_2 yields four moles of CO_2 plus six moles of H_2O.

Exercise 3.20.1

For the following equation below, indicate the number of formula units or molecules, and the number of moles present in the balanced equation.

$$\text{KBrO}_3 + 6\text{KI} + 6\text{HBr} \rightarrow 7\text{KBr} + 3\text{H}_2\text{O}$$

Answer

One molecules of KBrO_3 plus six molecules of KI plus six molecules of HBr yields seven molecules of KBr plus three molecules of I_2 and three molecules of H_2O. One mole of KBrO_3 plus six moles of KI plus six moles of HBr yields seven moles of KBr plus three moles of I_2 plus three moles of H_2O.

Summary

- Stoichiometry is the calculation of the quantities of reactants or products in a chemical reaction using the relationships found in a balanced chemical equation.
- The coefficients in a balanced chemical equation represent the reacting ratios of the substances in the reaction.
- The coefficients of a balanced equation can be used to determine the ratio of moles of all substances in the reaction.

Vocabulary

- **Stoichiometry** - The calculation of quantitative relationships of the reactants and products in a balanced chemical equation.
- **Formula unit** - The empirical formula of an ionic compound.
- **Mole ratio** - The ratio of the moles of one reactant or product to the moles of another reactant or product according to the coefficients in the balanced chemical equation.

3.20: Stoichiometry is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **8.5: Stoichiometry** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

3.21: LIMITING REACTANT AND THEORETICAL YIELD

LEARNING OBJECTIVES

- Identify the limiting reactant (limiting reagent) in a given chemical reaction.
- Calculate how much product will be produced from the limiting reactant.
- Calculate how much reactant(s) remains when the reaction is complete.

In all examples discussed thus far, the reactants were assumed to be present in stoichiometric quantities. Consequently, none of the reactants were left over at the end of the reaction. This is often desirable—as in the case of a space shuttle—where excess oxygen or hydrogen is not only extra freight to be hauled into orbit, but also an explosion hazard. More often, however, reactants are present in mole ratios that are not the same as the ratio of the coefficients in the balanced chemical equation. As a result, one or more of them will not be used up completely, but will be left over when the reaction is completed. In this situation, the amount of product that can be obtained is limited by the amount of only one of the reactants. The reactant that restricts the amount of product obtained is called the limiting reactant. The reactant that remains after a reaction has gone to completion is in excess.

Consider a nonchemical example. Assume you have invited some friends for dinner and want to bake brownies for dessert. You find two boxes of brownie mix in your pantry and see that each package requires two eggs. The balanced equation for brownie preparation is:

$$\textbf{1 box mix} + \textbf{2 eggs} \rightarrow \textbf{1 batch brownies} \tag{3.21.1}$$

If you have a dozen eggs, which ingredient will determine the number of batches of brownies that you can prepare? Because each box of brownie mix requires two eggs and you have two boxes, you need four eggs. Twelve eggs is eight more eggs than you need. Although the ratio of eggs to boxes in is 2:1, the ratio in your possession is 6:1. Hence the eggs are the ingredient (reactant) present in excess, and the brownie mix is the limiting reactant. Even if you had a refrigerator full of eggs, you could make only two batches of brownies.





Figure **3.21.1**: The Concept of a Limiting Reactant in the Preparation of Brownies. For a chemist, the **balanced** chemical equation is the recipe that must be followed. 2 boxes of brownie mix and 12 eggs results in 2 batches of brownies and 8 eggs; in this case the 8 eggs are reactant present in excess

PHET SIMULATION: REACTANTS, PRODUCTS AND LEFTOVERS

View this interactive simulation illustrating the concepts of limiting and excess reactants.



Consider this concept now with regard to a chemical process, the reaction of hydrogen with chlorine to yield hydrogen chloride:

$$H_2 + Cl_2(g) \rightarrow 2\,HCl(g)$$

The balanced equation shows that hydrogen and chlorine react in a 1:1 stoichiometric ratio. If these reactants are provided in any other amounts, one of the reactants will nearly always be entirely consumed, thus limiting the amount of product that may be generated. This substance is the limiting reactant, and the other substance is the excess reactant. Identifying the limiting and excess reactants for a given situation

requires computing the molar amounts of each reactant provided and comparing them to the stoichiometric amounts represented in the balanced chemical equation. For example, imagine combining 3 moles of H_2 and 2 moles of Cl_2. This represents a 3:2 (or 1.5:1) ratio of hydrogen to chlorine present for reaction, which is greater than the stoichiometric ratio of 1:1. Hydrogen, therefore, is present in excess, and chlorine is the limiting reactant. Reaction of all the provided chlorine (2 mol) will consume 2 mol of the 3 mol of hydrogen provided, leaving 1 mol of hydrogen non-reacted.

An alternative approach to identifying the limiting reactant involves comparing the amount of product expected for the complete reaction of each reactant. Each reactant amount is used to separately calculate the amount of product that would be formed per the reaction's stoichiometry. The reactant yielding the lesser amount of product is the limiting reactant. For the example, in the previous paragraph, complete reaction of the hydrogen would yield:

$$\text{mol HCl produced} = 3\ \text{mol } H_2 \times \frac{2\ \text{mol HCl}}{1\ \text{mol } H_2} = 6\ \text{mol HCl}$$

Complete reaction of the provided chlorine would produce:

$$\text{mol HCl produced} = 2\ \text{mol } Cl_2 \times \frac{2\ \text{mol HCl}}{1\ \text{mol } Cl_2} = 4\ \text{mol HCl}$$

The chlorine will be completely consumed once 4 moles of HCl have been produced. Since enough hydrogen was provided to yield 6 moles of HCl, there will be non-reacted hydrogen remaining once this reaction is complete. Chlorine, therefore, is the limiting reactant and hydrogen is the excess reactant (Figure **3.21.2**).




Figure **3.21.2**: When H_2 and Cl_2 are combined in nonstoichiometric amounts, one of these reactants will limit the amount of HCl that can be produced. This illustration shows a reaction in which hydrogen is present in excess and chlorine is the limiting reactant. The figure shows a space-filling molecular models reacting. There is a reaction arrow pointing to the right in the middle. To the left of the reaction arrow there are three molecules each consisting of two green spheres bonded together. There are also five molecules each consisting of two smaller, white spheres bonded together. Above these molecules is the label, "Before reaction," and below these molecules is the label, "6 H subscript 2 and 4 C l subscript 2." To the right of the reaction arrow, there are eight molecules each consisting of one green sphere bonded to a smaller white sphere. There are also two molecules each consisting of two white spheres bonded together. Above these molecules is the label, "After reaction," and below these molecules is the label, "8 H C l and 2 H subscript 2."

A similar situation exists for many chemical reactions: you usually run out of one reactant before all of the other reactant has reacted. The reactant you run out of is called the limiting reactant; the other reactant or reactants are considered to be *in excess*. A crucial skill in evaluating the conditions of a chemical process is to determine which reactant is the limiting reactant and which is in excess.

HOW TO IDENTIFY THE LIMITING REACTANT (LIMITING REAGENT)

There are two ways to determine the limiting reactant. One method is to find and compare the mole ratio of the reactants used in the reaction (Approach 1). Another way is to calculate the grams of products produced from the given quantities of reactants; the reactant that produces the **smallest** amount of product is the limiting reactant (Approach 2). This section will focus more on the second method.

Approach 1 (The "Reactant Mole Ratio Method"): Find the limiting reactant by looking at the number of moles of each reactant.

1. Determine the balanced chemical equation for the chemical reaction.
2. Convert all given information into moles (most likely, through the use of molar mass as a conversion factor).
3. Calculate the mole ratio from the given information. Compare the calculated ratio to the actual ratio.
4. Use the amount of limiting reactant to calculate the amount of product produced.
5. If necessary, calculate how much is left in excess of the non-limiting (excess) reactant.

Approach 2 (The "The Product Method"): Find the limiting reactant by calculating and comparing the amount of product that each reactant will produce.

1. Balance the chemical equation for the chemical reaction.
2. Convert the given information into moles.
3. Use stoichiometry for each individual reactant to find the mass of product produced.
4. The reactant that produces a lesser amount of product is the limiting reactant.
5. The reactant that produces a larger amount of product is the excess reactant.
6. To find the amount of remaining excess reactant, subtract the mass of excess reactant consumed from the total mass of excess reactant given.

The key to recognizing which reactant is the limiting reactant is based on a mole-mass or mass-mass calculation: whichever reactant gives the *lesser* amount of product is the limiting reactant. What we need to do is determine an amount of one product (either moles or mass) assuming all of each reactant reacts. Whichever reactant gives the least amount of that particular product is the limiting reactant. It does not matter which product we use, as long as we use the same one each time. It does not matter whether we determine the number of moles or grams of that product; however, we will see shortly that knowing the final mass of product can be useful.

EXAMPLE 3.21.1: IDENTIFYING THE LIMITING REACTANT

As an example, consider the balanced equation

$$4\,C_2H_3Br_3 + 11\,O_2 \rightarrow 8\,CO_2 + 6\,H_2O + 6\,Br_2$$

What is the limiting reactant if 76.4 grams of $C_2H_3Br_3$ reacted with 49.1 grams of O_2?

Solution

Using Approach 1:

Step 1: *Balance the chemical equation.*

The equation is already balanced with the relationship

4 mol $C_2H_3Br_3$ to 11 mol O_2 to 6 mol H_2O to 6 mol Br

Step 2: Convert all given information into moles.

$$76.4\ \cancel{\text{g } C_2H_3Br_3} \times \frac{1\ \text{mol } C_2H_3Br_3}{266.72\ \cancel{\text{g } C_2H_3Br_3}} = 0.286\ \text{mol } C_2H_3Br_3$$

$$49.1\ \cancel{\text{g } O_2} \times \frac{1\ \text{mol } O_2}{32.00\ \cancel{\text{g } O_2}} = 1.53\ \text{mol } O_2$$

Step 3: Calculate the mole ratio from the given information. Compare the calculated ratio to the actual ratio.

Assuming that all of the oxygen is used up,

$$1.53\ \cancel{\text{mol } O_2} \times \frac{4\ \text{mol } C_2H_3Br_3}{11\ \cancel{\text{mol } O_2}} = 0.556\ \text{mol } C_2H_3Br_3 \text{ are required.}$$

Because 0.556 moles of $C_2H_3Br_3$ required > 0.286 moles of $C_2H_3Br_3$ available, $C_2H_3Br_3$ is the limiting reactant.

Using Approach 2:

Step 1: *Balance the chemical equation.*

The equation is already balanced with the relationship

4 mol $C_2H_3Br_3$ to 11 mol O_2 to 6 mol H_2O to 6 mol Br_2

Step 2 and Step 3: Convert mass to moles and stoichiometry.

$$76.4\ \text{g}\ C_2H_3Br_3 \times \frac{1\ \text{mol}\ C_2H_3Br_3}{266.72\ \text{g}\ C_2H_3Br_3} \times \frac{8\ \text{mol}\ CO_2}{4\ \text{mol}\ C_2H_3Br_3} \times \frac{44.01\ \text{g}\ CO_2}{1\ \text{mol}\ CO_2} = 25.2\ \text{g}\ CO_2$$

$$49.1\ \text{g}\ O_2 \times \frac{1\ \text{mol}\ O_2}{32.00\ \text{g}\ O_2} \times \frac{8\ \text{mol}\ CO_2}{11\ \text{mol}\ O_2} \times \frac{44.01\ \text{g}\ CO_2}{1\ \text{mol}\ CO_2} = 49.1\ \text{g}\ CO_2$$

Step 4: The reactant that produces a smaller amount of product is the limiting reactant.

Therefore, by either method, $C_2H_3Br_3$ is the limiting reactant.

EXAMPLE 3.21.2: IDENTIFYING THE LIMITING REACTANT AND THE MASS OF EXCESS REACTANT

For example, in the reaction of magnesium metal and oxygen, calculate the mass of magnesium oxide that can be produced if 2.40 g ***Mg*** reacts with 10.0 g ***O_2***. Also determine the amount of excess reactant. **MgO** is the only product in the reaction.

Solution

Following Approach 1:

Step 1: Balance the chemical equation.

$$2\ \text{Mg (s)} + O_2\ \text{(g)} \rightarrow 2\ \text{MgO (s)}$$

The balanced equation provides the relationship of 2 mol Mg to 1 mol O_2 to 2 mol MgO

Step 2 and Step 3: Convert mass to moles and stoichiometry.

$$2.40\ \text{g}\ Mg \times \frac{1\ \text{mol}\ Mg}{24.31\ \text{g}\ Mg} \times \frac{2\ \text{mol}\ MgO}{2\ \text{mol}\ Mg} \times \frac{40.31\ \text{g}\ MgO}{1\ \text{mol}\ MgO} = 3.98\ \text{g}\ MgO$$

$$10.0\ \text{g}\ O_2 \times \frac{1\ \text{mol}\ O_2}{32.00\ \text{g}\ O_2} \times \frac{2\ \text{mol}\ MgO}{1\ \text{mol}\ O_2} \times \frac{40.31\ \text{g}\ MgO}{1\ \text{mol}\ MgO} = 25.2\ \text{g}\ MgO$$

Step 4: The reactant that produces a smaller amount of product is the limiting reactant.

Mg produces less MgO than does O_2 (3.98 g MgO vs. 25.2 g MgO), therefore Mg is the limiting reactant in this reaction.

Step 5: The reactant that produces a larger amount of product is the excess reactant.

O_2 produces more amount of MgO than Mg (25.2g MgO vs. 3.98 MgO), therefore O_2 is the excess reactant in this reaction.

Step 6: Find the amount of remaining excess reactant by subtracting the mass of the excess reactant consumed from the total mass of excess reactant given.

Mass of excess reactant calculated using the limiting reactant:

$$2.40\ \text{g}\ Mg \times \frac{1\ \text{mol}\ Mg}{24.31\ \text{g}\ Mg} \times \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ Mg} \times \frac{32.00\ \text{g}\ O_2}{1\ \text{mol}\ O_2} = 1.58\ \text{g}\ O_2$$

OR

Mass of excess reactant calculated using the mass of the product:

$$3.98\ \text{g}\ MgO \times \frac{1\ \text{mol}\ MgO}{40.31\ \text{g}\ MgO} \times \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ MgO} \times \frac{32.0\ \text{g}\ O_2}{1\ \text{mol}\ O_2} = 1.58\ \text{g}\ O_2$$

Mass of total excess reactant given – mass of excess reactant consumed in the reaction:

10.0g O_2 - (available) 1.58g O_2 (used) = 8.42g O_2 (excess)

Therefore, O_2 is in excess.


EXAMPLE 3.21.3: LIMITING REACTANT

What is the limiting reactant if 78.0 grams of Na_2O_2 were reacted with 29.4 grams of H_2O? The unbalanced chemical equation is

$$Na_2O_2\,(s) + H_2O\,(l) \longrightarrow NaOH\,(aq) + H_2O_2\,(l)$$

Solution

Solutions to Example 8.4.3

Steps for Problem Solving- The Product Method	Example 3.21.1
Identify the "given" information and what the problem is asking you to "find."	Given: 78.0 grams of Na_2O_2 29.4 g H_2O Find: limiting reactant
List other known quantities.	1 mol Na_2O_2= 77.96 g/mol 1 mol H_2O = 18.02 g/mol Since the amount of product in grams is not required, only the molar mass of the reactants is needed.
Balance the equation.	Na_2O_2 (s) + $\underline{2}H_2O$ (l) → $\underline{2}$NaOH (aq) + H_2O_2 (l) The balanced equation provides the relationship of 1 mol Na_2O_2 to 2 mol H_2O 2mol NaOH to 1 mol H_2O_2
Prepare a concept map and use the proper conversion factor.	  Because the question only asks for the limiting reactant, we can perform two **mass-mole** calculations and determine which amount is less.
Cancel units and calculate.	$78.0\ g\ Na_2O_2 \times \frac{1\ mol\ Na_2O_2}{77.96\ g\ Na_2O_2} \times \frac{2\ mol\ NaOH}{1\ mol\ Na_2O_2} \times \frac{40\ g\ NaOH}{1\ mol\ NaOH} = 2.00\ mol\ NaOH$ $29.4\ g\ H_2O \times \frac{1\ mol\ H_2O}{18.02\ g\ H_2O} \times \frac{2\ mol\ NaOH}{2\ mol\ Na_2O_2} \times \frac{40\ g\ NaOH}{1\ mol\ NaOH} = 1.63\ mol\ NaOH$ Therefore, H_2O is the *limiting reactant*.

Think about your result.

✔ EXAMPLE 3.21.4: LIMITING REACTANT AND MASS OF EXCESS REACTANT




A 5.00 g quantity of **Rb** is combined with 3.44 g of $\mathbf{MgCl_2}$ according to this chemical reaction:

$$2Rb(s) + MgCl_2(s) \rightarrow Mg(s) + 2RbCl(s)$$

What mass of **Mg** is formed, and what mass of remaining reactant is left over?

Solution

Solutions to Example 8.4.10

Steps for Problem Solving- The Product Method	Example 3.21.2
Steps for Problem Solving	A 5.00 g quantity of Rb is combined with 3.44 g of $MgCl_2$ according to this chemical reaction: $$2Rb(s) + MgCl_2(s) \rightarrow Mg(s) + 2RbCl(s)$$ What mass of Mg is formed, and what mass of remaining reactant is left over?
Identify the "given" information and what the problem is asking you to "find."	Given: 5.00g Rb, 2.44g MgCl2 Find: mass of Mg formed, mass of remaining reactant
List other known quantities.	• molar mass: Rb = 85.47 g/mol • molar mass: $MgCl_2$ = 95.21 g/mol • molar mass: Mg = 24.31 g/mol
Prepare concept maps and use the proper conversion factor.	Find mass Mg formed based on mass of Rb  Find mass of Mg formed based on mass of $MgCl_2$  Conversion factors: 1 mole MgCl2 to 95.21 grams MgCl2, 1 mole Mg to 1 mole MgCl2, 24.31 grams Mg to 1 mole Mg Use limiting reactant to determine amount of excess reactant consumed  Conversion factors: 1 mole Rb to 85.47 grams Rb, 1 mole MgCl2 to 2 moles Rb, 95.21 grams MgCl2 to 1 mole MgCl2
Cancel units and calculate.	Because the question asks what mass of magnesium is formed, we can perform two mass-mass calculations and determine which amount is less. $$5.00\ g\,Rb \times \frac{1\ mol\ Rb}{85.47\ g\,Rb} \times \frac{1\ mol\ Mg}{2\ mol\ Rb} \times \frac{24.31\ g\ Mg}{1\ mol\ Mg} = 0.711\ g\ Mg$$ $$3.44\ g\,MgCl_2 \times \frac{1\ mol\ MgCl_2}{95.21\ g\,MgCl_2} \times \frac{1\ mol\ Mg}{1\ mol\ MgCl_2} \times \frac{24.31\ g\ Mg}{1\ mol\ Mg} = 0.878\ g\ Mg$$ The 0.711 g of Mg is the lesser quantity, so the associated reactant—5.00 g of Rb—is the limiting reactant. To determine how much of the other reactant is left, we have to do one more mass-mass calculation to determine what mass of $MgCl_2$ reacted with the 5.00 g of Rb, and then subtract the amount reacted from the original amount. $$5.00\ g\,Rb \times \frac{1\ mol\ Rb}{85.47\ g\,Rb} \times \frac{1\ mol\ MgCl_2}{2\ mol\ Rb} \times \frac{95.21\ g\ MgCl_2}{1\ mol\ MgCl_2} = 2.78\ g\ MgCl_2\ reacted$$ Because we started with 3.44 g of $MgCl_2$, we have 3.44 g $MgCl_2$ − 2.78 g $MgCl_2$ reacted = 0.66 g $MgCl_2$ left
Think about your result.	It usually is not possible to determine the limiting reactant using just the initial masses, as the reagents have different molar masses and coefficients.

? EXERCISE 3.21.1

Given the initial amounts listed, what is the limiting reactant, and what is the mass of the leftover reactant?

$$\underset{MgO(s)}{22.7\ g} + \underset{H_2S}{17.9\ g} \rightarrow MgS(s) + H_2O(l)$$

Answer

H_2S is the limiting reagent; 1.5 g of MgO are left over.

3.21: Limiting Reactant and Theoretical Yield is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Sarick Shah, Marisa Alviar-Agnew, Henry Agnew, & Henry Agnew.

- **8.6: Limiting Reactant and Theoretical Yield** by Henry Agnew, Marisa Alviar-Agnew, Sarick Shah is licensed CC BY-NC-SA 3.0.

https://chem.libretexts.org/@go/page/492172



https://chem.libretexts.org/@go/page/492172

3.22: LIMITING REACTANT, THEORETICAL YIELD, AND PERCENT YIELD FROM INITIAL MASSES OF REACTANTS

LEARNING OBJECTIVES

- Calculate percentage or actual yields from known amounts of reactants.

The world of pharmaceutical production is an expensive one. Many drugs have several steps in their synthesis and use costly chemicals. A great deal of research takes place to develop better ways to make drugs faster and more efficiently. Studying how much of a compound is produced in any given reaction is an important part of cost control.

PERCENT YIELD

Chemical reactions in the real world don't always go exactly as planned on paper. In the course of an experiment, many things will contribute to the formation of less product than predicted. Besides spills and other experimental errors, there are usually losses due to an incomplete reaction, undesirable side reactions, etc. Chemists need a measurement that indicates how successful a reaction has been. This measurement is called the percent yield.

To compute the percent yield, it is first necessary to determine how much of the product should be formed based on stoichiometry. This is called the **theoretical yield**, the maximum amount of product that can be formed from the given amounts of reactants. The **actual yield** is the amount of product that is actually formed when the reaction is carried out in the laboratory. The **percent yield** is the ratio of the actual yield to the theoretical yield, expressed as a percentage.

$$\text{Percent Yield} = \frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100\%$$

Percent yield is very important in the manufacture of products. Much time and money is spent improving the percent yield for chemical production. When complex chemicals are synthesized by many different reactions, one step with a low percent yield can quickly cause a large waste of reactants and unnecessary expense.

Typically, percent yields are understandably less than 100% because of the reasons indicated earlier. However, percent yields greater than 100% are possible if the measured product of the reaction contains impurities that cause its mass to be greater than it actually would be if the product was pure. When a chemist synthesizes a desired chemical, he or she is always careful to purify the products of the reaction. Example **3.22.1** illustrates the steps for determining percent yield.

EXAMPLE 3.22.1: DECOMPOSITION OF POTASSIUM CHLORATE

Potassium chlorate decomposes upon slight heating in the presence of a catalyst, according to the reaction below:

$$2KClO_3\,(s) \rightarrow 2KCl\,(s) + 3O_2\,(g)$$

In a certain experiment, $40.0\ g\ KClO_3$ is heated until it completely decomposes. The experiment is performed and the oxygen gas is collected and its mass is found to be $14.9\ g$.

a. What is the theoretical yield of oxygen gas?
b. What is the percent yield for the reaction?

Solution

a. Calculation of theoretical yield

First, we will calculate the theoretical yield based on the stoichiometry.

Step 1: Identify the "given" information and what the problem is asking you to "find".

Given: Mass of $KClO_3 = 40.0\ g$

Mass of O_2 collected = 14.9g


Find: Theoretical yield, g O_2

Step 2: List other known quantities and plan the problem.

1 mol $KClO_3$ = 122.55 g/mol

1 mol O_2 = 32.00 g/mol

Step 3: Apply stoichiometry to convert from the mass of a reactant to the mass of a product:



Step 4: Solve.

$$40.0\ \cancel{\text{g KClO}_3} \times \frac{1\ \cancel{\text{mol KClO}_3}}{122.55\ \cancel{\text{g KClO}_3}} \times \frac{3\ \cancel{\text{mol O}_2}}{2\ \cancel{\text{mol KClO}_3}} \times \frac{32.00\ \text{g O}_2}{1\ \cancel{\text{mol O}_2}} = 15.7\ \text{g O}_2$$

The theoretical yield of O_2 is **15.7 g**, 15.67 g unrounded.

Step 5: Think about your result.

The mass of oxygen gas must be less than the **40.0 g** of potassium chlorate that was decomposed.

b. Calculation of percent yield

Now we will use the actual yield and the theoretical yield to calculate the percent yield.

Step 1: Identify the "given" information and what the problem is asking you to "find".

Given: Theoretical yield =15.67 g, use the un-rounded number for the calculation.

Actual yield = 14.9g

Find: Percent yield, % Yield

Step 2: List other known quantities and plan the problem.

No other quantities needed.

Step 3: Use the percent yield equation below.

$$\text{Percent Yield} = \frac{\text{Actual Yield}}{\text{Theoretical Yield}} \times 100\%$$

Step 4: Solve.

$$\text{Percent Yield} = \frac{14.9\ \text{g}}{15.67\ \text{g}} \times 100\% = 94.9\%$$

Step 5: Think about your result.

Since the actual yield is slightly less than the theoretical yield, the percent yield is just under **100%**.

EXAMPLE 3.22.2: OXIDATION OF ZINC


Upon reaction of 1.274 g of copper sulfate with excess zinc metal, 0.392 g copper metal was obtained according to the equation:

$$CuSO_4(aq) + Zn(s) \rightarrow Cu(s) + ZnSO_4(aq)$$

What is the percent yield?

Solution

Solutions to Example 8.6.2

Steps for Problem Solving-The Product Method	Example 3.22.1
Identify the "given" information and what the problem is asking you to "find."	Given: 1.274 g CuSO4 Actual yield = 0.392 g Cu Find: Percent yield
List other known quantities.	1 mol CuSO4= 159.62 g/mol 1 mol Cu = 63.55 g/mol Since the amount of product in grams is not required, only the molar mass of the reactants is needed.
Balance the equation.	The chemical equation is already balanced. The balanced equation provides the relationship of 1 mol CuSO4 to 1 mol Zn to 1 mol Cu to 1 mol ZnSO4.
Prepare a concept map and use the proper conversion factor.	 The provided information identifies copper sulfate as the limiting reactant, and so the theoretical yield (g Cu) is found by performing **mass-mass** calculation based on the initial amount of $CuSO_4$.
Cancel units and calculate.	Double subscripts: use braces to clarify Using this theoretical yield and the provided value for actual yield, the percent yield is calculated to be: $\text{percent yield} = \left(\frac{\text{actual yield}}{\text{theoretical yield}}\right) \times 100$ $= \left(\frac{0.392\ g\ Cu}{0.5072\ g\ Cu}\right) \times 100$ $= 77.3\%$
Think about your result.	Since the actual yield is slightly less than the theoretical yield, the percent yield is just under 100%.

EXERCISE 3.22.1

What is the percent yield of a reaction that produces 12.5 g of the Freon CF_2Cl_2 from 32.9 g of CCl_4 and excess HF?

$$CCl_4 + 2\,HF \rightarrow CF_2Cl_2 + 2\,HCl$$

Answer

48.3%

SUMMARY

Theoretical yield is calculated based on the stoichiometry of the chemical equation. The actual yield is experimentally determined. The percent yield is determined by calculating the ratio of actual yield to theoretical yield.

3.22: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **8.7: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

CHAPTER OVERVIEW

4: Unit 4

4.1: The Bohr Model - Atoms with Orbits
4.2: Quantum-Mechanical Orbitals and Electron Configurations
4.3: Electron Configurations and the Periodic Table
4.4: The Explanatory Power of the Quantum-Mechanical Model
4.5: Representing Valence Electrons with Dots
4.6: Lewis Structures of Ionic Compounds- Electrons Transferred
4.7: Covalent Lewis Structures- Electrons Shared
4.8: Writing Lewis Structures for Covalent Compounds
4.9: Resonance - Equivalent Lewis Structures for the Same Molecule
4.10: Predicting the Shapes of Molecules
4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix
4.12: Kinetic Molecular Theory- A Model for Gases
4.13: Pressure - The Result of Constant Molecular Collisions
4.14: Boyle’s Law - Pressure and Volume
4.15: Charles’s Law- Volume and Temperature
4.16: Gay-Lussac's Law- Temperature and Pressure
4.17: The Combined Gas Law- Pressure, Volume, and Temperature
4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles
4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen

4: Unit 4 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

4.1: The Bohr Model - Atoms with Orbits

Learning Objectives

- Define an energy level in terms of the Bohr model.
- Discuss how the Bohr model can be used to explain atomic spectra.

Electric light bulbs contain a very thin wire in them that emits light when heated. The wire is called a filament. The particular wire used in light bulbs is made of tungsten. A wire made of any metal would emit light under these circumstances, but tungsten was chosen because the light it emits contains virtually every frequency and therefore, the light emitted by tungsten appears white. A wire made of some other element would emit light of some color that was not convenient for our uses. Every element emits light when energized by heating or passing electric current through it. Elements in solid form begin to glow when they are heated sufficiently, and elements in gaseous form emit light when electricity passes through them. This is the source of light emitted by neon signs and is also the source of light in a fire.



Figure **4.1.1**: Human/Need/Desire. Neon sculpture by Bruce Nauman (1983), who has been characterized as a conceptual artist.

Each Element Has a Unique Spectrum

The light frequencies emitted by atoms are mixed together by our eyes so that we see a blended color. Several physicists, including Angstrom in 1868 and Balmer in 1875, passed the light from energized atoms through glass prisms in such a way that the light was spread out so they could see the individual frequencies that made up the light. The **emission spectrum** (or **atomic spectrum**) of a chemical element is the unique pattern of light obtained when the element is subjected to heat or electricity.



Figure **4.1.2**: Atomic Emission Spectrum of Hydrogen.

When hydrogen gas is placed into a tube and electric current passed through it, the color of emitted light is pink. But when the color is spread out, we see that the hydrogen spectrum is composed of four individual frequencies. The pink color of the tube is the result of our eyes blending the four colors. Every atom has its own characteristic spectrum; no two atomic spectra are alike. The image below shows the emission spectrum of iron. Because each element has a unique emission spectrum, elements can be defined using them.




Figure **4.1.3**: Atomic Emission Spectrum of Iron.

You may have heard or read about scientists discussing what elements are present in the sun or some more distant star, and after hearing that, wondered how scientists could know what elements were present in a place no one has ever been. Scientists determine what elements are present in distant stars by analyzing the light that comes from stars and finding the atomic spectrum of elements in that light. If the exact four lines that compose hydrogen's atomic spectrum are present in the light emitted from the star, that element contains hydrogen.

Bohr's Model of the Atom

By 1913, the concept of the atom had evolved from Dalton's indivisible spheres idea, to J. J. Thomson's plum pudding model, and then to Rutherford's nuclear atom theory. Rutherford, in addition to carrying out the brilliant experiment that demonstrated the presence of the atomic nucleus, also proposed that the electrons circled the nucleus in a planetary type motion. The solar system or planetary model of the atom was attractive to scientists because it was similar to something with which they were already familiar, namely the solar system.



Figure **4.1.3**: Niels Bohr with Albert Einstein at Paul Ehrenfest's home in Leiden (December 1925).

Unfortunately, there was a serious flaw in the planetary model. It was already known that when a charged particle (such as an electron) moves in a curved path, it gives off some form of light and loses energy in doing so. This is, after all, how we produce TV signals. If the electron circling the nucleus in an atom loses energy, it would necessarily have to move closer to the nucleus as it loses energy, and would eventually crash into the nucleus. Furthermore, Rutherford's model was unable to describe how electrons give off light forming each element's unique atomic spectrum. These difficulties cast a shadow on the planetary model and indicated that, eventually, it would have to be replaced.

In 1913, the Danish physicist Niels Bohr proposed a model of the electron cloud of an atom in which electrons orbit the nucleus and were able to produce atomic spectra. Understanding Bohr's model requires some knowledge of electromagnetic radiation (or light).

Energy Levels

Bohr's key idea in his model of the atom is that electrons occupy definite orbitals that require the electron to have a specific amount of energy. In order for an electron to be in the electron cloud of an atom, it must be in one of the allowable orbitals and it must have the precise energy required for that orbit. Orbits closer to the nucleus would require smaller amounts of energy for an electron, and orbits farther from the nucleus would require the electron to have a greater amount of energy. The possible orbits are known as **energy levels**. One of the weaknesses of Bohr's model was that he could not offer a reason why only certain energy levels or orbits were allowed.




Figure **4.1.4**: The energy levels of the electrons can be viewed as rungs on a ladder. Note that the spacing between rungs gets smaller at higher energies (CC BY-NC; Ümit Kaya)

Bohr hypothesized that the only way electrons could gain or lose energy would be to move from one energy level to another, thus gaining or losing precise amounts of energy. The energy levels are **quantized**, meaning that only specific amounts are possible. It would be like a ladder that had rungs only at certain heights. The only way you can be on that ladder is to be on one of the rungs, and the only way you could move up or down would be to move to one of the other rungs. Suppose we had such a ladder with 10 rungs. Other rules for the ladder are that only one person can be on a rung in the normal state, and the ladder occupants must be on the lowest rung available. If the ladder had five people on it, they would be on the lowest five rungs. In this situation, no person could move down because all of the lower rungs are full. Bohr worked out rules for the maximum number of electrons that could be in each energy level in his model, and required that an atom in its normal state (ground state) had all electrons in the lowest energy levels available. Under these circumstances, no electron could lose energy because no electron could move down to a lower energy level. In this way, Bohr's model explained why electrons circling the nucleus did not emit energy and spiral into the nucleus.




Figure **4.1.5**: In Bohr's Model of the atom, electrons absorb energy to move to a higher level and release energy to move to lower levels. (CC BY-SA 3.0; Kurzon).

Bohr's Model and Atomic Spectra

The evidence used to support Bohr's model came from the atomic spectra. He suggested that an atomic spectrum is made by the electrons in an atom moving energy levels. The electrons typically have the lowest energy possible, called the **ground state**. If the electrons are given energy (through heat, electricity, light, etc.) the electrons in an atom could absorb energy by jumping to a higher energy level, or **excited state**. The electrons then give off the energy in the form of a piece of light—called a **photon**—that they had absorbed, to fall back to a lower energy level. The energy emitted by electrons dropping back to lower energy levels will always be precise amounts of energy, because the differences in energy levels are precise. This explains why you see specific lines of light when looking at an atomic spectrum—each line of light matches a specific "step down" that an electron can take in that atom. This also explains why each element produces a different atomic spectrum. Because each element has different acceptable energy levels for its electrons, the possible steps each element's electrons can take differ from all other elements.

Summary

- Bohr's model suggests each atom has a set of unchangeable energy levels, and electrons in the electron cloud of that atom must be in one of those energy levels.
- Bohr's model suggests that the atomic spectra of atoms is produced by electrons gaining energy from some source, jumping up to a higher energy level, then immediately dropping back to a lower energy level and emitting the energy difference between the two energy levels.
- The existence of the atomic spectra is support for Bohr's model of the atom.
- Bohr's model was only successful in calculating energy levels for the hydrogen atom.

Vocabulary

- **Emission spectrum** (or **atomic spectrum**) - The unique pattern of light given off by an element when it is given energy.
- **Energy levels** - Possible orbits that an electron can have in the electron cloud of an atom.
- **Ground state** - To be in the lowest energy level possible.
- **Excited state** - To be in a higher energy level.
- **Photon** - A piece of electromagnetic radiation, or light, with a specific amount of energy.
- **Quantized** - To have a specific amount.

4.1: The Bohr Model - Atoms with Orbits is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **9.4: The Bohr Model - Atoms with Orbits** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.2: Quantum-Mechanical Orbitals and Electron Configurations

Learning Objectives

- Represent the organization of electrons by an electron configuration and orbital diagram.

The flight path of a commercial airliner is carefully regulated by the Federal Aviation Administration. Each airplane must maintain a distance of five miles from another plane flying at the same altitude and 2,000 feet above and below another aircraft (1,000 feet if the altitude is less than 29,000 feet). So, each aircraft only has certain positions it is allowed to maintain while it flies. As we explore quantum mechanics, we see that electrons have similar restrictions on their locations.

Orbitals

We can apply our knowledge of quantum numbers to describe the arrangement of electrons for a given atom. We do this with something called **electron configurations**. They are effectively a map of the electrons for a given atom. We look at the four quantum numbers for a given electron and then assign that electron to a specific orbital below.

s Orbitals

For any value of n, a value of $l = 0$ places that electron in an s **orbital**. This orbital is spherical in shape:




Figure 4.2.1: s orbitals have no orientational preference and resemble spheres.

p Orbitals

For the table below, we see that we can have three possible orbitals when $l = 1$. These are designated as p **orbitals** and have dumbbell shapes. Each of the p orbitals has a different orientation in three-dimensional space.




Figure 4.2.2: p orbitals have an orientational preference and resemble dumbbells.

d Orbitals

When $l = 2$, m_l values can be $-2, -1, 0, +1, +2$ for a total of five d **orbitals**. Note that all five of the orbitals have specific three-dimensional orientations.




Figure 4.2.3: d orbitals have an orientational preference and exhibit complex structures.

f Orbitals

The most complex set of orbitals are the f **orbitals**. When $l = 3$, m_l values can be $-3, -2, -1, 0, +1, +2, +3$ for a total of seven different orbital shapes. Again, note the specific orientations of the different f orbitals.




Figure 4.2.4: f orbitals have an orientational preference and exhibit quite complex structures.

Orbitals that have the same value of the principal quantum number form a **shell**. Orbitals within a shell are divided into **subshells** that have the same value of the angular quantum number. Some of the allowed combinations of quantum numbers are compared in Table 4.2.1.

Table 4.2.1: Electron Arrangement Within Energy Levels

Principal Quantum Number (n)	Allowable Sublevels	Number of Orbitals per Sublevel	Number of Orbitals per Principal Energy Level	Number of Electrons per Sublevel	Number of Electrons per Principal Energy Level
1	s	1	1	2	2
2	s	1	4	2	8
	p	3		6	
3	s	1	9	2	18
	p	3		6	
	d	5		10	
4	s	1	16	2	32
	p	3		6	
	d	5		10	
	f	7		14	

Electron Configurations

Can you name one thing that easily distinguishes you from the rest of the world? And we're not talking about DNA—that's a little expensive to sequence. For many people, it is their email address. Your email address allows people all over the world to contact you. It does not belong to anyone else, but serves to identify you. Electrons also have a unique set of identifiers in the quantum numbers that describe their location and spin. Chemists use an **electronic configuration** to represent the organization of electrons in shells and subshells in an atom. An electron configuration simply lists the shell and subshell labels, with a right superscript giving the number of electrons in that subshell. The shells and subshells are listed in the order of filling. Electrons are typically organized around an atom by starting at the lowest possible quantum numbers first, which are the shells-subshells with lower energies.

For example, an H atom has a single electron in the 1s subshell. Its electron configuration is

$$\mathbf{H:}\ 1s^1$$

He has two electrons in the 1s subshell. Its electron configuration is

$$\mathbf{He:}\ 1s^2$$

The three electrons for Li are arranged in the 1s subshell (two electrons) and the 2s subshell (one electron). The electron configuration of Li is

$$\mathbf{Li:}\ 1s^2 2s^1$$

Be has four electrons, two in the 1s subshell and two in the 2s subshell. Its electron configuration is

$$\text{Be}: 1s^2 2s^2$$

Now that the 2s subshell is filled, electrons in larger atoms must go into the 2p subshell, which can hold a maximum of six electrons. The next six elements progressively fill up the 2p subshell:

- B: $1s^2 2s^2 2p^1$
- C: $1s^2 2s^2 2p^2$
- N: $1s^2 2s^2 2p^3$
- O: $1s^2 2s^2 2p^4$
- F: $1s^2 2s^2 2p^5$
- Ne: $1s^2 2s^2 2p^6$

Now that the 2p subshell is filled (all possible subshells in the n = 2 shell), the next electron for the next-larger atom must go into the n = 3 shell, s subshell.

Second Period Elements

Periods refer to the horizontal rows of the periodic table. Looking at a periodic table you will see that the first period contains only the elements hydrogen and helium. This is because the first principal energy level consists of only the s sublevel and so only two electrons are required in order to fill the entire principal energy level. Each time a new principal energy level begins, as with the third element lithium, a new period is started on the periodic table. As one moves across the second period, electrons are successively added. With beryllium ($Z = 4$), the $2s$ sublevel is complete and the $2p$ sublevel begins with boron ($Z = 5$). Since there are three $2p$ orbitals and each orbital holds two electrons, the $2p$ sublevel is filled after six elements. Table 4.2.1 shows the electron configurations of the elements in the second period.

Table 4.2.2: Electron Configurations of Second-Period Elements

Element Name	Symbol	Atomic Number	Electron Configuration
Lithium	Li	3	$1s^2 2s^1$
Beryllium	Be	4	$1s^2 2s^2$
Boron	B	5	$1s^2 2s^2 2p^1$
Carbon	C	6	$1s^2 2s^2 2p^2$
Nitrogen	N	7	$1s^2 2s^2 2p^3$
Oxygen	O	8	$1s^2 2s^2 2p^4$
Fluorine	F	9	$1s^2 2s^2 2p^5$
Neon	Ne	10	$1s^2 2s^2 2p^6$

Aufbau Principle

Construction of a building begins at the bottom. The foundation is laid and the building goes up step by step. You obviously cannot start with the roof since there is no place to hang it. The building goes from the lowest level to the highest level in a systematic way. In order to create ground state electron configurations for any element, it is necessary to know the way in which the atomic sublevels are organized in order of increasing energy. Figure 4.2.5 shows the order of increasing energy of the sublevels.

The lowest energy sublevel is always the $1s$ sublevel, which consists of one orbital. The single electron of the hydrogen atom will occupy the $1s$ orbital when the atom is in its ground state. As we proceed with atoms with multiple electrons, those electrons are added to the next lowest sublevel: $2s$, $2p$, $3s$, and so on. The **Aufbau principle** states that an electron occupies orbitals in order from lowest energy to highest. The Aufbau (German: "building up, construction") principle is sometimes referred to as the "building up" principle. It is worth noting that in reality atoms are not built by adding protons and electrons one at a time, and that this method is merely an aid for us to understand the end result.




Figure 4.2.5: Electrons are added to atomic orbitals in order from low energy (bottom of the graph) to high (top of the graph) according to the Aufbau principle. Principle energy levels are color coded, while sublevels are grouped together and each circle represents an orbital capable of holding two electrons.

As seen in the figure above, the energies of the sublevels in different principal energy levels eventually begin to overlap. After the **3*p*** sublevel, it would seem logical that the **3*d*** sublevel should be the next lowest in energy. However, the **4*s*** sublevel is slightly lower in energy than the **3*d*** sublevel and thus fills first. Following the filling of the **3*d*** sublevel is the **4*p***, then the **5*s*** and the **4*d***. Note that the **4*f*** sublevel does not fill until just after the **6*s*** sublevel. Figure **4.2.6** is a useful and simple aid for keeping track of the order of fill of the atomic sublevels.




Figure 4.2.6: The arrow leads through each subshell in the appropriate filling order for electron configurations. This chart is straightforward to construct. Simply make a column for all the s orbitals with each n shell on a separate row. Repeat for p, d, and f. Be sure to only include orbitals allowed by the quantum numbers (no 1p or 2d, and so forth). Finally, draw diagonal lines from top to bottom as shown.



Video **4.2.1**: Energy levels, sublevels and orbitals.

Example 4.2.1: Nitrogen Atoms

Nitrogen has 7 electrons. Write the electron configuration for nitrogen.

Solution:

Take a close look at Figure **4.2.5**, and use it to figure out how many electrons go into each sublevel, and also the order in which the different sublevels get filled.

1. Begin by filling up the 1*s* sublevel. **This gives $1s^2$**. Now all of the orbitals in the red $n = 1$ block are filled.

   Since we used 2 electrons, there are $7 - 2 =$ **5 electrons left**

2. Next, fill the 2*s* sublevel. This gives **$1s^2 2s^2$**. Now all of the orbitals in the *s* sublevel of the orange $n = 2$ block are filled.

   Since we used another 2 electrons, there are $5 - 2 =$ **3 electrons left**

3. Notice that we haven't filled the entire $n = 2$ block yet… there are still the *p* orbitals!

   The final 3 electrons go into the 2*p* sublevel. This gives **$1s^2 2s^2 2p^3$**

The overall electron configuration is: **$1s^2 2s^2 2p^3$**.

Example 4.2.2: Potassium Atoms

Potassium has 19 electrons. Write the electron configuration code for potassium.

Solution

This time, take a close look at Figure **4.2.5**.

1. Begin by filling up the 1*s* sublevel. **This gives $1s^2$**. Now the $n = 1$ level is filled.

   Since we used 2 electrons, there are $19 - 2 =$ **17 electrons left**

2. Next, fill the 2*s* sublevel. **This gives $1s^2 2s^2$**

   Since we used another 2 electrons, there are $17 - 2 =$ **15 electrons left**

3. Next, fill the 2*p* sublevel. **This gives $1s^2 2s^2 2p^6$.** Now the $n = 2$ level is filled.

   Since we used another 6 electrons, there are $15 - 6 =$ **9 electrons left**

4. Next, fill the 3*s* sublevel. **This gives $1s^2 2s^2 2p^6 3s^2$**

Since we used another 2 electrons, there are 9 − 2 = **7 electrons left**

5. Next, fill the 3*p* sublevel. **This gives $1s^2 2s^2 2p^6 3s^2 3p^6$**

Since we used another 6 electrons, there are 7 − 6 = **1 electron left**

Here's where we have to be careful – right after $3p^6$!

Remember, 4*s* comes before 3*d*

6. The final electron goes into the 4*s* sublevel. **This gives $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$**

The overall electron configuration is: **$1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$**

? Exercise 4.2.1: Magnesium and Sodium Atoms

What is the electron configuration for Mg and Na?

Answer Mg

Mg: $1s^2 2s^2 2p^6 3s^2$

Answer Na

Na: $1s^2 2s^2 2p^6 3s^1$

Pauli Exclusion Principle

When we look at the orbital possibilities for a given atom, we see that there are different arrangements of electrons for each different type of atom. Since each electron must maintain its unique identity, we intuitively sense that the four quantum numbers for any given electron must not match up exactly with the four quantum numbers for any other electron in that atom.

For the hydrogen atom, there is no problem since there is only one electron in the **H** atom. However, when we get to helium we see that the first three quantum numbers for the two electrons are the same: same energy level, same spherical shape. What differentiates the two helium electrons is their spin. One of the electrons has a $+\frac{1}{2}$ spin while the other electron has a $-\frac{1}{2}$ spin. So the two electrons in the **1*s*** orbital are each unique and distinct from one another because their spins are different. This observation leads to the **Pauli exclusion principle**, which states that no two electrons in an atom can have the same set of four quantum numbers. The energy of the electron is specified by the principal, angular momentum, and magnetic quantum numbers. If those three numbers are identical for two electrons, the spin numbers must be different in order for the two electrons to be differentiated from one another. The two values of the spin quantum number allow each orbital to hold two electrons. Figure **4.2.7** shows how the electrons are indicated in a diagram.




Figure **4.2.7**: In an orbital filling diagram, a square represents an orbital, while arrows represent electrons. An arrow pointing upward represents one spin direction, while an arrow pointing downward represents the other spin direction.

Hund's Rule

The last of the three rules for constructing electron arrangements requires electrons to be placed one at a time in a set of orbitals within the same sublevel. This minimizes the natural repulsive forces that one electron has for another. **Hund's rule** states that orbitals of equal energy are each occupied by one electron before any orbital is occupied by a second electron and that each of the single electrons must have the same spin. The figure below shows how a set of three ***p*** orbitals is filled with one, two, three, and four electrons.




Figure 4.2.8: The $2p$ sublevel, for the elements boron ($Z = 5$), carbon ($Z = 6$), nitrogen ($Z = 7$), and oxygen ($Z = 8$). According to Hund's rule, as electrons are added to a set of orbitals of equal energy, one electron enters each orbital before any orbital receives a second electron.

Orbital Filling Diagrams

An **orbital filling diagram** is the more visual way to represent the arrangement of all the electrons in a particular atom. In an orbital filling diagram, the individual orbitals are shown as circles (or squares) and orbitals within a sublevel are drawn next to each other horizontally. Each sublevel is labeled by its principal energy level and sublevel. Electrons are indicated by arrows inside of the circles. An arrow pointing upwards indicates one spin direction, while a downward pointing arrow indicates the other direction. The orbital filling diagrams for hydrogen, helium, and lithium are shown in the figure below.




Figure 4.2.9: Orbital filling diagrams for hydrogen, helium, and lithium.

According to the Aufbau process, sublevels and orbitals are filled with electrons in order of increasing energy. Since the s sublevel consists of just one orbital, the second electron simply pairs up with the first electron as in helium. The next element is lithium and necessitates the use of the next available sublevel, the $2s$.

The filling diagram for carbon is shown in Figure 4.2.10. There are two $2p$ electrons for carbon and each occupies its own $2p$ orbital.




Figure 4.2.10: Orbital filling diagram for carbon.

Oxygen has four $2p$ electrons. After each $2p$ orbital has one electron in it, the fourth electron can be placed in the first $2p$ orbital with a spin opposite that of the other electron in that orbital.




Figure 4.2.11: Orbital filling diagram for oxygen.

If you keep your papers in manila folders, you can pick up a folder and see how much it weighs. If you want to know how many different papers (articles, bank records, or whatever else you keep in a folder), you have to take everything out and count. A computer directory, on the other hand, tells you exactly how much you have in each file. We can get the same information on atoms. If we use an orbital filling diagram, we have to count arrows. When we look at electron configuration data, we simply add up the numbers.

Example 4.2.3: Carbon Atoms

Draw the orbital filling diagram for carbon and write its electron configuration.

Solution


Step 1: List the known quantities and plan the problem.

Known

- Atomic number of carbon, Z=6

Use the order of fill diagram to draw an orbital filling diagram with a total of six electrons. Follow Hund's rule. Write the electron configuration.

Step 2: Construct the diagram.



Orbital filling diagram for carbon.

Electron configuration $1s^2 2s^2 2p^2$

Step 3: Think about your result.

Following the 2s sublevel is the 2p, and p sublevels always consist of three orbitals. All three orbitals need to be drawn even if one or more is unoccupied. According to Hund's rule, the sixth electron enters the second of those p orbitals and has the same spin as the fifth electron.


Exercise 4.2.2: Electronic Configurations

Write the electron configurations and orbital diagrams for

a. Potassium atom: **K**
b. Arsenic atom: **As**
c. Phosphorus atom: **P**

Answer a:

Potassium: $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$




Answer b:

Arsenic: $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^3$



Answer c:

Phosphorus $1s^2 2s^2 2p^6 3s^2 3p^3$



The Atom Neighborhood



Figure **4.2.12**: The atom neighborhood. Source: Dr. Binh Dao, Sacramento City College.

Summary

There are four different classes of electron orbitals. These orbitals are determined by the value of the angular momentum quantum number ℓ. An orbital is a wave function for an electron defined by the three quantum numbers, n, ℓ and m_ℓ. Orbitals define regions in space where you are likely to find electrons. *s* orbitals ($\ell = 0$) are spherical shaped. *p* orbitals ($\ell = 1$) are dumb-bell shaped. The three possible *p* orbitals are always perpendicular to each other.

Electron configuration notation simplifies the indication of where electrons are located in a specific atom. Superscripts are used to indicate the number of electrons in a given sublevel. The Aufbau principle gives the order of electron filling in an atom. It can be used to describe the locations and energy levels of every electron in a given atom. Hund's rule specifies the order of electron filling within a set of orbitals. Orbital filling diagrams are a way of indicating electron locations in orbitals. The Pauli exclusion principle specifies limits on how identical quantum numbers can be for two electrons in the same atom.

Vocabulary

principal quantum number (*n*)
: Defines the energy level of the wave function for an electron, the size of the electron's standing wave, and the number of nodes in that wave.

quantum numbers
: Integer numbers assigned to certain quantities in the electron wave function. Because electron standing waves must be continuous and must not "double over" on themselves, quantum numbers are restricted to integer values.

Contributions & Attributions

-
- Paul Flowers (University of North Carolina - Pembroke), Klaus Theopold (University of Delaware) and Richard Langley (Stephen F. Austin State University) with contributing authors. Textbook content produced by OpenStax College is licensed

under a Creative Commons Attribution License 4.0 license. Download for free at http://cnx.org/contents/85abf193-2bd...a7ac8df6@9.110).

4.2: Quantum-Mechanical Orbitals and Electron Configurations is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **9.6: Quantum-Mechanical Orbitals and Electron Configurations** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.3: Electron Configurations and the Periodic Table

Learning Objectives

- Relate the electron configurations of the elements to the shape of the periodic table.
- Determine the expected electron configuration of an element by its place on the periodic table.

Previously, we introduced the periodic table as a tool for organizing the known chemical elements. A periodic table is shown in Figure **4.3.1**. The elements are listed by atomic number (the number of protons in the nucleus), and elements with similar chemical properties are grouped together in columns.




Figure **4.3.1**: The Periodic Table

Why does the periodic table have the structure it does? The answer is rather simple, if you understand electron configurations: the shape of the periodic table mimics the filling of the subshells with electrons.

> *The shape of the periodic table mimics the filling of the subshells with electrons.*

Let us start with H and He. Their electron configurations are $1s^1$ and $1s^2$, respectively; with He, the $n = 1$ shell is filled. These two elements make up the first row of the periodic table (Figure **4.3.2**)




Figure **4.3.2**: The 1*s* Subshell. H and He represent the filling of the 1*s* subshell.

The next two electrons, for Li and Be, would go into the 2*s* subshell. Figure **4.3.3** shows that these two elements are adjacent on the periodic table.




Figure 4.3.3: The 2*s* Subshell. In Li and Be, the 2*s* subshell is being filled.

For the next six elements, the 2*p* subshell is being occupied with electrons. On the right side of the periodic table, these six elements (B through Ne) are grouped together (Figure 4.3.4).




Figure 4.3.4: The 2*p* Subshell. For B through Ne, the 2*p* subshell is being occupied.

The next subshell to be filled is the 3*s* subshell. The elements when this subshell is being filled, Na and Mg, are back on the left side of the periodic table (Figure 4.3.5).




Figure 4.3.5: The 3*s* Subshell. Now the 3*s* subshell is being occupied.

Next, the 3*p* subshell is filled with the next six elements (Figure 4.3.6).




Figure **4.3.6**: The 3*p* Subshell. Next, the 3*p* subshell is filled with electrons.

Instead of filling the 3*d* subshell next, electrons go into the 4*s* subshell (Figure **4.3.7**).




Figure **4.3.7**: The 4*s* Subshell. The 4*s* subshell is filled before the 3*d* subshell. This is reflected in the structure of the periodic table.

After the 4*s* subshell is filled, the 3*d* subshell is filled with up to 10 electrons. This explains the section of 10 elements in the middle of the periodic table (Figure **4.3.8**).




Figure **4.3.8**: The 3*d* Subshell. The 3*d* subshell is filled in the middle section of the periodic table.

...And so forth. As we go across the rows of the periodic table, the overall shape of the table outlines how the electrons are occupying the shells and subshells.

The first two columns on the left side of the periodic table are where the *s* subshells are being occupied. Because of this, the first two rows of the periodic table are labeled the **s block**. Similarly, the **p block** are the right-most six columns of the periodic table, the **d block** is the middle 10 columns of the periodic table, while the **f block** is the 14-column section that is normally depicted as detached from the main body of the periodic table. It could be part of the main body, but then the periodic table would be rather long and cumbersome. Figure **4.3.9** shows the blocks of the periodic table.




Figure **4.3.9**: Blocks on the Periodic Table. The periodic table is separated into blocks depending on which subshell is being filled for the atoms that belong in that section.

The electrons in the highest-numbered shell, plus any electrons in the last unfilled subshell, are called **valence electrons**; the highest-numbered shell is called the **valence shell**. (The inner electrons are called *core electrons*.) The valence electrons largely control the chemistry of an atom. If we look at just the valence shell's electron configuration, we find that in each column, the valence shell's electron configuration is the same. For example, take the elements in the first column of the periodic table: H, Li, Na, K, Rb, and Cs. Their electron configurations (abbreviated for the larger atoms) are as follows, with the valence shell electron configuration highlighted:

Electrons, electron configurations, and the valence shell electron configuration highlighted.

H:	$1s^{\mathbf{1}}$
Li:	$1s^2\mathbf{2s^1}$
Na:	$[Ne]\mathbf{3s^1}$
K:	$[Ar]\mathbf{4s^1}$
Rb:	$[Kr]\mathbf{5s^1}$
Cs:	$[Xe]\mathbf{6s^1}$

They all have a similar electron configuration in their valence shells: a single *s* electron. Because much of the chemistry of an element is influenced by valence electrons, we would expect that these elements would have similar chemistry—*and they do*. The organization of electrons in atoms explains not only the shape of the periodic table, but also the fact that elements in the same column of the periodic table have similar chemistry.

The same concept applies to the other columns of the periodic table. Elements in each column have the same valence shell electron configurations, and the elements have some similar chemical properties. This is strictly true for all elements in the *s* and *p* blocks. In the *d* and *f* blocks, because there are exceptions to the order of filling of subshells with electrons, similar valence shells are not absolute in these blocks. However, many similarities do exist in these blocks, so a similarity in chemical properties is expected.

Similarity of valence shell electron configuration implies that we can determine the electron configuration of an atom solely by its position on the periodic table. Consider Se, as shown in Figure **4.3.10**. It is in the fourth column of the *p* block. This means that its electron configuration should end in a p^4 electron configuration. Indeed, the electron configuration of Se is $[Ar]4s^23d^{10}4p^4$, as expected.




Figure 4.3.10: Selenium on the Periodic Table

Example 4.3.1: Predicting Electron Configurations

From the element's position on the periodic table, predict the valve shell electron configuration for each atom (Figure 4.3.11).




Figure 4.3.11: Various Elements on the Periodic Table

a. Ca
b. Sn

Solution

a. Ca is located in the second column of the *s* block. We expect that its electron configuration should end with s^2. Calcium's electron configuration is $[Ar]4s^2$.
b. Sn is located in the second column of the *p* block, so we expect that its electron configuration would end in p^2. Tin's electron configuration is $[Kr]5s^24d^{10}5p^2$.

Exercise 4.3.1

From the element's position on the periodic table, predict the valence shell electron configuration for each atom. Figure 4.3.11.

a. Ti
b. Cl

Answer a

$[Ar]4s^23d^2$

Answer b

$[Ne]3s^23p^5$

https://chem.libretexts.org/@go/page/492177

Summary

The arrangement of electrons in atoms is responsible for the shape of the periodic table. Electron configurations can be predicted by the position of an atom on the periodic table.

4.3: Electron Configurations and the Periodic Table is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **9.7: Electron Configurations and the Periodic Table** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.4: The Explanatory Power of the Quantum-Mechanical Model

Learning Objectives

- Give the name and location of specific groups on the periodic table, including alkali metals, alkaline earth metals, noble gases, halogens, and transition metals.
- Explain the relationship between the chemical behavior of families in the periodic table and their electron configurations.
- Identify elements that will have the most similar properties to a given element.

The chemical behavior of atoms is controlled by their electron configuration. Since the families of elements were organized by their chemical behavior, it is predictable that the individual members of each chemical family will have similar electron configurations.


Families of the Periodic Table

Remember that Mendeleev arranged the periodic table so that elements with the most similar properties were placed in the same group. A **group** is a vertical column of the periodic table. All of the 1A elements have one valence electron. This is what causes these elements to react in the same ways as the other members of the family. The elements in 1A are all very reactive and form compounds in the same ratios with similar properties with other elements. Because of their similarities in their chemical properties, Mendeleev put these elements into the same group. Group 1A is also known as the **alkali metals**. Although most metals tend to be very hard, these metals are actually soft and can be easily cut.

Group 2A is also called the **alkaline earth metals**. Once again, because of their similarities in electron configurations, these elements have similar properties to each other. The same pattern is true of other groups on the periodic table. Remember, Mendeleev arranged the table so that elements with the most similar properties were in the same group on the periodic table.

It is important to recognize a couple of other important groups on the periodic table by their group name. Group 7A (or 17) elements are also called **halogens**. This group contains very reactive nonmetal elements.

The **noble gases** are in group 8A. These elements also have similar properties to each other, the most significant property being that they are extremely unreactive, rarely forming compounds. The reason for this will be communicated later, when we discuss how compounds form. The elements in this group are also gases at room temperature.




Families of the periodic table.
CC – Tracy Poulsen

An alternate numbering system numbers all of the ***s***, ***p***, and ***d*** block elements from 1-18. In this numbering system, group 1A is group 1; group 2A is group 2; the halogens (7A) are group 17; and the noble gases (8A) are group 18. You will come across periodic tables with both numbering systems. It is important to recognize which numbering system is being used, and to be able to find the number of valence electrons in the main block elements, regardless of which numbering system is being used.

Periods of the Periodic Table

If you can locate an element on the Periodic Table, you can use the element's position to figure out the energy level of the element's valence electrons. A **period** is a horizontal row of elements on the periodic table. For example, the elements sodium (**Na**) and magnesium (**Mg**) are both in period 3. The elements astatine (**At**) and radon (**Rn**) are both in period 6.



CC – Tracy Poulsen

Summary

- The vertical columns on the periodic table are called groups or families because of their similar chemical behavior.
- All the members of a family of elements have the same number of valence electrons and similar chemical properties.
- The horizontal rows on the periodic table are called periods.

Vocabulary

- **Group (family)** - A vertical column of the periodic table.
- **Alkali metals** - Group 1A of the periodic table.
- **Alkaline earth metals** - Group 2A of the periodic table.
- **Halogens** - Group 7A of the periodic table.
- **Noble gases** - Group 8A of the periodic table.
- **Transition elements** - Groups 3 to 12 of the periodic table.

4.4: The Explanatory Power of the Quantum-Mechanical Model is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **9.8: The Explanatory Power of the Quantum-Mechanical Model** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.5: REPRESENTING VALENCE ELECTRONS WITH DOTS

LEARNING OBJECTIVE

- Draw a Lewis electron dot diagram for an atom or a monatomic ion.

In almost all cases, chemical bonds are formed by interactions of valence electrons in atoms. To facilitate our understanding of how valence electrons interact, a simple way of representing those valence electrons would be useful.

A **Lewis electron dot diagram** (or electron dot diagram, or a Lewis diagram, or a Lewis structure) is a representation of the valence electrons of an atom that uses dots around the symbol of the element. The number of dots equals the number of valence electrons in the atom. These dots are arranged to the right and left and above and below the symbol, with no more than two dots on a side. (The order in which the positions are used does not matter.) For example, the Lewis electron dot diagram for hydrogen is simply

$$\mathbf{H}\cdot$$

Because the side is not important, the Lewis electron dot diagram could also be drawn as follows:

$$\dot{\mathbf{H}} \quad \textit{or} \quad \cdot\mathbf{H} \quad \textit{or} \quad \underset{\cdot}{\mathbf{H}}$$

The electron dot diagram for helium, with two valence electrons, is as follows:

$$\mathbf{He:}$$

By putting the two electrons together on the same side, we emphasize the fact that these two electrons are both in the 1*s* subshell; this is the common convention we will adopt, although there will be exceptions later. The next atom, lithium, has an electron configuration of $1s^2 2s^1$, so it has only one electron in its valence shell. Its electron dot diagram resembles that of hydrogen, except the symbol for lithium is used:

$$\mathbf{Li}\cdot$$

Beryllium has two valence electrons in its 2*s* shell, so its electron dot diagram is like that of helium:

$$\mathbf{Be:}$$

The next atom is boron. Its valence electron shell is $2s^2 2p^1$, so it has three valence electrons. The third electron will go on another side of the symbol:

$$\dot{\mathbf{B}}\mathbf{:}$$

Again, it does not matter on which sides of the symbol the electron dots are positioned.

For carbon, there are four valence electrons, two in the 2*s* subshell and two in the 2*p* subshell. As usual, we will draw two dots together on one side, to represent the 2*s* electrons. However, conventionally, we draw the dots for the two *p* electrons on different sides. As such, the electron dot diagram for carbon is as follows:

$$\cdot\dot{\mathbf{C}}\mathbf{:}$$

With N, which has three *p* electrons, we put a single dot on each of the three remaining sides:

$$\cdot\underset{\cdot}{\dot{\mathbf{N}}}\mathbf{:}$$

For oxygen, which has four *p* electrons, we now have to start doubling up on the dots on one other side of the symbol. When doubling up electrons, make sure that each side has no more than two electrons.

$$\cdot\underset{\cdot}{\ddot{\mathbf{O}}}\mathbf{:}$$

Fluorine and neon have seven and eight dots, respectively:

$$:\underset{\cdot}{\ddot{\mathbf{F}}}\mathbf{:}$$

$$:\underset{\cdot\,\cdot}{\ddot{\mathbf{Ne}}}\mathbf{:}$$

With the next element, sodium, the process starts over with a single electron because sodium has a single electron in its highest-numbered shell, the $n = 3$ shell. By going through the periodic table, we see that the Lewis electron dot diagrams of atoms will never have more than eight dots around the atomic symbol.

EXAMPLE 4.5.1: LEWIS DOT DIAGRAMS

What is the Lewis electron dot diagram for each element?

a. aluminum
b. selenium

Solution

a. The valence electron configuration for aluminum is $3s^2 3p^1$. So it would have three dots around the symbol for aluminum, two of them paired to represent the 3*s* electrons:



2. The valence electron configuration for selenium is $4s^2 4p^4$. In the highest-numbered shell, the $n = 4$ shell, there are six electrons. Its electron dot diagram is as follows:



EXERCISE 4.5.1

What is the Lewis electron dot diagram for each element?

a. phosphorus
b. argon

Answer a



Answer b



SUMMARY

- Lewis electron dot diagrams use dots to represent valence electrons around an atomic symbol.
- Lewis electron dot diagrams for ions have less (for cations) or more (for anions) dots than the corresponding atom.

4.5: Representing Valence Electrons with Dots is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.2: Representing Valence Electrons with Dots** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

4.6: LEWIS STRUCTURES OF IONIC COMPOUNDS- ELECTRONS TRANSFERRED

LEARNING OBJECTIVES

- State the octet rule.
- Define *ionic bond*.
- Draw Lewis structures for ionic compounds.

In Section 4.7, we demonstrated that ions are formed by losing electrons to make cations, or by gaining electrons to form anions. The astute reader may have noticed something: many of the ions that form have eight electrons in their valence shell. Either atoms gain enough electrons to have eight electrons in the valence shell and become the appropriately charged anion, or they lose the electrons in their original valence shell; the *lower* shell, now the valence shell, has eight electrons in it, so the atom becomes positively charged. For whatever reason, having eight electrons in a valence shell is a particularly energetically stable arrangement of electrons. The **octet rule** explains the favorable trend of atoms having eight electrons in their valence shell. When atoms form compounds, the octet rule is not always satisfied for all atoms at all times, but it is a very good rule of thumb for understanding the kinds of bonding arrangements that atoms can make.

It is not impossible to violate the octet rule. Consider sodium: in its elemental form, it has one valence electron and is stable. It is rather reactive, however, and does not require a lot of energy to remove that electron to make the Na^+ ion. We *could* remove another electron by adding even more energy to the ion, to make the Na^{2+} ion. However, that requires much more energy than is normally available in chemical reactions, so sodium stops at a 1+ charge after losing a single electron. It turns out that the Na^+ ion has a complete octet in its new valence shell, the $n = 2$ shell, which satisfies the octet rule. The octet rule is a result of trends in energies and is useful in explaining why atoms form the ions that they do.

Now consider an Na atom in the presence of a Cl atom. The two atoms have these Lewis electron dot diagrams and electron configurations:

$$\mathbf{Na}\cdot \qquad \cdot\ddot{\mathbf{Cl}}\mathbf{:}$$

$$[Ne]\,3s^1 \qquad [Ne]\,3s^2 3p^5$$

For the Na atom to obtain an octet, it must lose an electron; for the Cl atom to gain an octet, it must gain an electron. An electron transfers from the Na atom to the Cl atom:

$$\mathbf{Na}\cdot \rightsquigarrow \cdot\ddot{\mathbf{Cl}}\mathbf{:}$$

resulting in two ions—the Na^+ ion and the Cl^- ion:

$$\mathbf{Na}^+ \qquad \mathbf{:}\ddot{\mathbf{Cl}}\mathbf{:}^-$$

$$[Ne] \qquad [Ne]\,3s^2 3p^6$$

Both species now have complete octets, and the electron shells are energetically stable. From basic physics, we know that opposite charges attract. This is what happens to the Na^+ and Cl^- ions:

$$\mathbf{Na}^+ + \mathbf{:}\ddot{\mathbf{Cl}}\mathbf{:}^- \rightarrow Na^+Cl^- \;\; or \;\; NaCl$$

where we have written the final formula (the formula for sodium chloride) as per the convention for ionic compounds, without listing the charges explicitly. The attraction between oppositely charged ions is called an **ionic bond,** and it is one of the main types of chemical bonds in chemistry. Ionic bonds are caused by electrons *transferring* from one atom to another.

In electron transfer, the number of electrons lost must equal the number of electrons gained. We saw this in the formation of NaCl. A similar process occurs between Mg atoms and O atoms, except in this case two electrons are transferred:



The two ions each have octets as their valence shell, and the two oppositely charged particles attract, making an ionic bond:


$$Mg^{2+} + \left[:\ddot{O}:\right]^{2-} \quad Mg^{2+}O^{2-} \text{ or } MgO$$

Remember, in the final formula for the ionic compound, we do not write the charges on the ions.

What about when an Na atom interacts with an O atom? The O atom needs two electrons to complete its valence octet, but the Na atom supplies only one electron:

$$Na\cdot \rightsquigarrow \cdot\ddot{O}:$$

The O atom still does not have an octet of electrons. What we need is a second Na atom to donate a second electron to the O atom:




These three ions attract each other to give an overall neutral-charged ionic compound, which we write as Na_2O. The need for the number of electrons lost being equal to the number of electrons gained explains why ionic compounds have the ratio of cations to anions that they do. This is required by the law of conservation of matter as well.

✔ EXAMPLE 4.6.1: SYNTHESIS OF CALCIUM CHLORIDE FROM ELEMENTS

With arrows, illustrate the transfer of electrons to form calcium chloride from *Ca* atoms and *Cl* atoms.

Solution

A *Ca* atom has two valence electrons, while a *Cl* atom has seven electrons. A *Cl* atom needs only one more to complete its octet, while *Ca* atoms have two electrons to lose. Thus we need two *Cl* atoms to accept the two electrons from one *Ca* atom. The transfer process looks as follows:




The oppositely charged ions attract each other to make $CaCl_2$.

? EXERCISE 4.6.1

With arrows, illustrate the transfer of electrons to form potassium sulfide from *K* atoms and *S* atoms.

Answer



SUMMARY

- The tendency to form species that have eight electrons in the valence shell is called the octet rule.
- The attraction of oppositely charged ions caused by electron transfer is called an ionic bond.
- The strength of ionic bonding depends on the magnitude of the charges and the sizes of the ions.

4.6: Lewis Structures of Ionic Compounds- Electrons Transferred is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.3: Lewis Structures of Ionic Compounds- Electrons Transferred** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

https://chem.libretexts.org/@go/page/492180

4.7: Covalent Lewis Structures- Electrons Shared

Learning Objectives

- Define *covalent bond*.
- Illustrate covalent bond formation with Lewis electron dot diagrams.

Ionic bonding typically occurs when it is easy for one atom to lose one or more electrons and another atom to gain one or more electrons. However, some atoms won't give up or gain electrons easily. Yet they still participate in compound formation. How? There is another mechanism for obtaining a complete valence shell: *sharing* electrons. When electrons are shared between two atoms, they make a bond called a **covalent bond**.


Let us illustrate a covalent bond by using H atoms, with the understanding that H atoms need only two electrons to fill the 1*s* subshell. Each H atom starts with a single electron in its valence shell:

$$\mathbf{H}\cdot \quad \cdot\mathbf{H}$$

The two H atoms can share their electrons:

$$\mathbf{H:H}$$


We can use circles to show that each H atom has two electrons around the nucleus, completely filling each atom's valence shell:



Because each H atom has a filled valence shell, this bond is stable, and we have made a diatomic hydrogen molecule. (This explains why hydrogen is one of the diatomic elements.) For simplicity's sake, it is not unusual to represent the covalent bond with a dash, instead of with two dots:

H–H


Because two atoms are sharing one pair of electrons, this covalent bond is called a **single bond**. As another example, consider fluorine. F atoms have seven electrons in their valence shell:



These two atoms can do the same thing that the H atoms did; they share their unpaired electrons to make a covalent bond.



Note that each F atom has a complete octet around it now:




We can also write this using a dash to represent the shared electron pair:




There are two different types of electrons in the fluorine diatomic molecule. The **bonding electron pair** makes the covalent bond. Each F atom has three other pairs of electrons that do not participate in the bonding; they are called **lone pair electrons**. Each F

atom has one bonding pair and three lone pairs of electrons.

Covalent bonds can be made between different elements as well. One example is HF. Each atom starts out with an odd number of electrons in its valence shell:



The two atoms can share their unpaired electrons to make a covalent bond:



We note that the H atom has a full valence shell with two electrons, while the F atom has a complete octet of electrons.

Example 4.7.1:


Use Lewis electron dot diagrams to illustrate the covalent bond formation in HBr.

Solution

HBr is very similar to HF, except that it has Br instead of F. The atoms are as follows:




The two atoms can share their unpaired electron:



Exercise 4.7.1

Use Lewis electron dot diagrams to illustrate the covalent bond formation in Cl_2.

Answer



When working with covalent structures, it sometimes looks like you have leftover electrons. You apply the rules you learned so far, and there are still some electrons that remain unattached. You can't just leave them there. So where do you put them?

Multiple Covalent Bonds

Some molecules are not able to satisfy the octet rule by making only single covalent bonds between the atoms. Consider the compound ethene, which has a molecular formula of $\mathbf{C_2H_4}$. The carbon atoms are bonded together, with each carbon also bonded to two hydrogen atoms.

$$\text{two } \mathbf{C} \text{ atoms } = 2 \times 4 = 8 \text{ valence electrons}$$

$$\text{four } \mathbf{H} \text{ atoms } = 4 \times 1 = 4 \text{ valence electrons}$$

$$\text{total of } 12 \text{ valence electrons in the molecule}$$

If the Lewis electron dot structure was drawn with a single bond between the carbon atoms and with the octet rule followed, it would look like this:




Figure 4.7.1: Incorrect dot structure of ethene. (CK12 License)

This Lewis structure is incorrect because it contains a total of 14 electrons. However, the Lewis structure can be changed by eliminating the lone pairs on the carbon atoms and having to share two pairs instead of only one pair.




Figure 4.7.2: Correct dot structure for ethene. (CK12 License)

A **double covalent bond** is a covalent bond formed by atoms that share two pairs of electrons. The double covalent bond that occurs between the two carbon atoms in ethane can also be represented by a structural formula and with a molecular model as shown in the figure below.




Figure 4.7.3: (A) The structural model for C_2H_4 consists of a double covalent bond between the two carbon atoms and single bonds to the hydrogen atoms. (B) Molecular model of C_2H_4.

A **triple covalent bond** is a covalent bond formed by atoms that share three pairs of electrons. The element nitrogen is a gas that composes the majority of Earth's atmosphere. A nitrogen atom has five valence electrons, which can be shown as one pair and three single electrons. When combining with another nitrogen atom to form a diatomic molecule, the three single electrons on each atom combine to form three shared pairs of electrons.




Figure 4.7.4: Triple bond in N_2.

Each nitrogen atom follows the octet rule with one lone pair of electrons, and six electrons that are shared between the atoms.

Summary

- Covalent bonds are formed when atoms share electrons.
- Lewis electron dot diagrams can be drawn to illustrate covalent bond formation.
- Double bonds or triple bonds between atoms may be necessary to properly illustrate the bonding in some molecules.

Contributions & Attributions

-
- Anonymous by request

4.7: Covalent Lewis Structures- Electrons Shared is shared under a mixed license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.4: Covalent Lewis Structures- Electrons Shared** by Henry Agnew, Marisa Alviar-Agnew is licensed notset.

4.8: Writing Lewis Structures for Covalent Compounds

Learning Objectives

- Draw Lewis structures for covalent compounds.

The following procedure can be used to construct Lewis electron structures for more complex molecules and ions.

How-to: Constructing Lewis electron structures

1. Determine the total number of valence electrons in the molecule or ion.

- Add together the valence electrons from each atom. (Recall that the number of valence electrons is indicated by the position of the element in the periodic table.)
- If the species is a polyatomic ion, remember to add or subtract the number of electrons necessary to give the total charge on the ion.

  For CO_3^{2-}, for example, we add two electrons to the total because of the -2 charge.

2. Arrange the atoms to show specific connections.

- When there is a central atom, it is usually the least electronegative element in the compound. Chemists usually list this central atom first in the chemical formula (as in CCl_4 and CO_3^{2-}, which both have C as the central atom), which is another clue to the compound's structure.
- Hydrogen and the halogens are almost always connected to only one other atom, so they are usually *terminal* rather than central.

3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond.

- In H_2O, for example, there is a bonding pair of electrons between oxygen and each hydrogen.

4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).

- These electrons will usually be lone pairs.

5. If any electrons are left over, place them on the central atom.

- We will explain later that some atoms are able to accommodate more than eight electrons.

6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.

- This will not change the number of electrons on the terminal atoms.

7. Final check

- Always make sure all valence electrons are accounted for and that each atom has an octet of electrons, except for hydrogen (with two electrons).
- The central atom is usually the least electronegative element in the molecule or ion; hydrogen and the halogens are usually terminal.

Now let's apply this procedure to some particular compounds, beginning with one we have already discussed.

Example 4.8.1: Water

Write the Lewis Structure for H_2O.

Solution

Solutions to Example 10.4.1



Steps for Writing Lewis Structures	Example 4.8.1
1. Determine the total number of valence electrons in the molecule or ion.	Each H atom (group 1) has 1 valence electron, and the O atom (group 16) has 6 valence electrons, for a total of 8 valence electrons.
2. Arrange the atoms to show specific connections.	H O H Because H atoms are almost always terminal, the arrangement within the molecule must be HOH.
3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond. 4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).	Placing one bonding pair of electrons between the O atom and each H atom gives H -O- H with 4 electrons left over. Each H atom has a full valence shell of 2 electrons.
5. If any electrons are left over, place them on the central atom.	Adding the remaining 4 electrons to the oxygen (as two lone pairs) gives the following structure: 
6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.	Not necessary.
7. Final check.	The Lewis structure gives oxygen an octet and each hydrogen 2 electrons.

Example 4.8.2

Write the Lewis structure for the CH_2O molecule

Solution


Solutions to Example 10.4.2

Steps for Writing Lewis Structures	Example 4.8.2
1. Determine the total number of valence electrons in the molecule or ion.	Each hydrogen atom (group 1) has 1 valence electron, carbon (group 14) has 4 valence electrons, and oxygen (group 16) has 6 valence electrons, for a total of [(2)(1) + 4 + 6] = 12 valence electrons.
2. Arrange the atoms to show specific connections.	 Because carbon is less electronegative than oxygen and hydrogen is normally terminal, C must be the central atom.
3. Place a bonding pair of electrons between each pair of adjacent atoms to give a single bond.	Placing a bonding pair of electrons between each pair of bonded atoms gives the following:  6 electrons are used, and 6 are left over.
4. Beginning with the terminal atoms, add enough electrons to each atom to give each atom an octet (two for hydrogen).	Adding all 6 remaining electrons to oxygen (as three lone pairs) gives the following:  Although oxygen now has an octet and each hydrogen has 2 electrons, carbon has only 6 electrons.
5. If any electrons are left over, place them on the central atom.	Not necessary. There are no electrons left to place on the central atom.
6. If the central atom has fewer electrons than an octet, use lone pairs from terminal atoms to form multiple (double or triple) bonds to the central atom to achieve an octet.	To give carbon an octet of electrons, we use one of the lone pairs of electrons on oxygen to form a carbon–oxygen double bond: 
7. Final check	Both the oxygen and the carbon now have an octet of electrons, so this is an acceptable Lewis electron structure. The O has two bonding pairs and two lone pairs, and C has four bonding pairs. This is the structure of formaldehyde, which is used in embalming fluid.

? Exercise 4.8.1

Write Lewis electron structures for CO_2 and SCl_2, a vile-smelling, unstable red liquid that is used in the manufacture of rubber.


Answer CO_2





Carbon dioxide

Answer SCl_2



The United States Supreme Court has the unenviable task of deciding what the law is. This responsibility can be a major challenge when there is no clear principle involved or where there is a new situation not encountered before. Chemistry faces the same challenge in extending basic concepts to fit a new situation. Drawing of Lewis structures for polyatomic ions uses the same approach, but tweaks the process a little to fit a somewhat different set of circumstances.

Writing Lewis Structures for Polyatomic Ions (CK-12)

Recall that a polyatomic ion is a group of atoms that are covalently bonded together and which carry an overall electrical charge. The ammonium ion, $\mathbf{NH_4^+}$, is formed when a hydrogen ion ($\mathbf{H^+}$) attaches to the lone pair of an ammonia ($\mathbf{NH_3}$) molecule in a coordinate covalent bond.




Figure **4.8.3**: The ammonium ion. (CK12 License)

When drawing the Lewis structure of a polyatomic ion, the charge of the ion is reflected in the number of total valence electrons in the structure. In the case of the ammonium ion:

$$\mathbf{1\ N}\text{ atom } = \mathbf{5}\text{ valence electrons}$$

$$\mathbf{4\ H}\text{ atoms } = \mathbf{4 \times 1 = 4}\text{ valence electrons}$$

subtract 1 electron for the $\mathbf{1+}$ charge of the ion


total of 8 valence electrons in the ion

It is customary to put the Lewis structure of a polyatomic ion into a large set of brackets, with the charge of the ion as a superscript outside of the brackets.

Exercise 4.8.2

Draw the Lewis electron dot structure for the sulfate ion.

Answer (CK12 License)



Exceptions to the Octet Rule (BC Campus)


As important and useful as the octet rule is in chemical bonding, there are some well-known violations. This does not mean that the octet rule is useless—quite the contrary. As with many rules, there are exceptions, or violations.

There are three violations to the octet rule. Odd-electron molecules represent the first violation to the octet rule. Although they are few, some stable compounds have an odd number of electrons in their valence shells. With an odd number of electrons, at least one atom in the molecule will have to violate the octet rule. Examples of stable odd-electron molecules are NO, NO_2, and ClO_2. The Lewis electron dot diagram for NO is as follows:




Although the O atom has an octet of electrons, the N atom has only seven electrons in its valence shell. Although NO is a stable compound, it is very chemically reactive, as are most other odd-electron compounds.


Electron-deficient molecules represent the second violation to the octet rule. These stable compounds have less than eight electrons around an atom in the molecule. The most common examples are the covalent compounds of beryllium and boron. For example, beryllium can form two covalent bonds, resulting in only four electrons in its valence shell:



Boron commonly makes only three covalent bonds, resulting in only six valence electrons around the B atom. A well-known example is BF_3:



The third violation to the octet rule is found in those compounds with more than eight electrons assigned to their valence shell. These are called expanded valence shell molecules. Such compounds are formed only by central atoms in the third row of the periodic table or beyond that have empty *d* orbitals in their valence shells that can participate in covalent bonding. One such compound is PF_5. The only reasonable Lewis electron dot diagram for this compound has the P atom making five covalent bonds:



Formally, the P atom has 10 electrons in its valence shell.

Example 4.8.3: Octet Violations

Identify each violation to the octet rule by drawing a Lewis electron dot diagram.


a. ClO
b. SF_6

Solution

a. With one Cl atom and one O atom, this molecule has 6 + 7 = 13 valence electrons, so it is an odd-electron molecule. A Lewis electron dot diagram for this molecule is as follows:



b. In SF_6, the central S atom makes six covalent bonds to the six surrounding F atoms, so it is an expanded valence shell molecule. Its Lewis electron dot diagram is as follows:



? Exercise 4.8.3: Xenon Difluoride

Identify the violation to the octet rule in XeF_2 by drawing a Lewis electron dot diagram.

Answer



The Xe atom has an expanded valence shell with more than eight electrons around it.

Summary

Lewis dot symbols provide a simple rationalization of why elements form compounds with the observed stoichiometries. A plot of the overall energy of a covalent bond as a function of internuclear distance is identical to a plot of an ionic pair because both result from attractive and repulsive forces between charged entities. In Lewis electron structures, we encounter **bonding pairs**, which are shared by two atoms, and **lone pairs**, which are not shared between atoms. Lewis structures for polyatomic ions follow the same rules as those for other covalent compounds. There are three violations to the octet rule: odd-electron molecules, electron-deficient molecules, and expanded valence shell molecules.

4.8: Writing Lewis Structures for Covalent Compounds is shared under a mixed license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.5: Writing Lewis Structures for Covalent Compounds** by Henry Agnew, Marisa Alviar-Agnew is licensed notset. Original source: https://www.ck12.org/c/chemistry/.

4.9: Resonance - Equivalent Lewis Structures for the Same Molecule

Learning Objectives

- Explain the concept of resonance and how it works with within molecules.

Resonance

There are some cases in which more than one viable Lewis structure can be drawn for a molecule. An example is the ozone (O_3) molecule in Figure **4.9.1**. There are a total of 18 electrons in the structure and so the following two structures are possible.




Figure **4.9.1**: Resonance forms of ozone. Note the use of the double-headed arrow.

The structure on the left (**4.9.1**) can be converted to the structure on the right by a shifting of electrons without altering the positions of the atoms.

It was once thought that the structure of a molecule such as O_3 consisted of one single bond and one double bond which then shifted back and forth as shown above. However, further studies showed that the two bonds are identical. Any double covalent bond between two given atoms is typically shorter than a single covalent bond. Studies of the O_3 and other similar molecules showed that the bonds were identical in length. Interestingly, the length of the bond is in between the lengths expected for an O–O single bond and a double bond.

Resonance is the use of two or more Lewis structures to represent the covalent bonding in a molecule. One of the valid structures is referred to as a resonance structure. It is now understood that the true structure of a molecule which displays resonance is that of an average or a hybrid of all the resonance structures. In the case of the O_3 molecule, each of the covalent bonds between O atoms are best thought of as being "one and a half" bonds, as opposed to either a pure single bond or a pure double bond. This "half-bond" can be shown as a dotted line in both the Lewis structure and the molecular model (Figure **4.9.2**).




Figure **4.9.2**: "Half-bond" model of ozone molecule. This is a better description of the electronic structure of ozone than either of the resonance structures in Figure **4.9.1**.

Many polyatomic ions also display resonance. In some cases, the true structure may be an average of three valid resonance structures, as in the case of the nitrate ion, NO_3^- in Figure **4.9.3**.




Figure **4.9.3**: Resonance structure of nitrate anion.

The bond lengths between the central N atom and each O atom are identical and the bonds can be approximated as being equal to one and one-third bonds.

Summary

- Resonance structures are averages of different Lewis structure possibilities.
- Bond lengths are intermediate between covalent bonds and covalent double bonds.

4.9: Resonance - Equivalent Lewis Structures for the Same Molecule is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.6: Resonance - Equivalent Lewis Structures for the Same Molecule** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.10: PREDICTING THE SHAPES OF MOLECULES

LEARNING OBJECTIVE

- Determine the shape of simple molecules.

Molecules have shapes. There is an abundance of experimental evidence to that effect—from their physical properties to their chemical reactivity. Small molecules—molecules with a single central atom—have shapes that can be easily predicted. The basic idea in molecular shapes is called **valence shell electron pair repulsion (VSEPR).** VSEPR says that electron pairs, being composed of negatively charged particles, repel each other to get as far away from one another as possible. VSEPR makes a distinction between *electron group geometry*, which expresses how electron groups (bonds and nonbonding electron pairs) are arranged, and *molecular geometry*, which expresses how the atoms in a molecule are arranged. However, the two geometries are related.

There are two types of **electron groups**: any type of bond—single, double, or triple—and lone electron pairs. When applying VSEPR to simple molecules, the first thing to do is to count the number of electron groups around the central atom. Remember that a multiple bond counts as only *one* electron group.

Any molecule with only two atoms is linear. A molecule whose central atom contains only two electron groups orients those two groups as far apart from each other as possible—180° apart. When the two electron groups are 180° apart, the atoms attached to those electron groups are also 180° apart, so the overall molecular shape is linear. Examples include BeH_2 and CO_2:



Figure **4.10.1**: Beryllium hydride and carbon dioxide bonding.

The two molecules, shown in the figure below in a "ball and stick" model.




Figure **4.10.2**: Beryllium hydride and carbon dioxide models. (CK12 Licence)

A molecule with three electron groups orients the three groups as far apart as possible. They adopt the positions of an equilateral triangle—120° apart and in a plane. The shape of such molecules is *trigonal planar*. An example is BF_3:




Figure **4.10.3**: Boron trifluoride bonding. (CK12 Licence)

Some substances have a trigonal planar electron group distribution but have atoms bonded to only two of the three electron groups. An example is GeF_2:




Figure **4.10.4**: Germanium difluoride bonding.

From an electron group geometry perspective, GeF_2 has a trigonal planar shape, but its real shape is dictated by the positions of the atoms. This shape is called *bent* or *angular*.

A molecule with four electron groups about the central atom orients the four groups in the direction of a tetrahedron, as shown in Figure **4.10.1** Tetrahedral Geometry. If there are four atoms attached to these electron groups, then the molecular shape is also *tetrahedral*. Methane (CH_4) is an example.




Figure **4.10.5**: Tetrahedral structure of methane. (CK12 Licence)

This diagram of CH_4 illustrates the standard convention of displaying a three-dimensional molecule on a two-dimensional surface. The straight lines are in the plane of the page, the solid wedged line is coming out of the plane toward the reader, and the dashed wedged line is going out of the plane away from the reader.




Figure **4.10.6**: Methane bonding. (CK12 Licence)

NH_3 is an example of a molecule whose central atom has four electron groups, but only three of them are bonded to surrounding atoms.




Figure **4.10.7**: Ammonia bonding. (CK12 Licence)

Although the electron groups are oriented in the shape of a tetrahedron, from a molecular geometry perspective, the shape of NH_3 is *trigonal pyramidal*.

H_2O is an example of a molecule whose central atom has four electron groups, but only two of them are bonded to surrounding atoms.



Figure **4.10.8**: Water bonding.

Although the electron groups are oriented in the shape of a tetrahedron, the shape of the molecule is *bent* or *angular*. A molecule with four electron groups about the central atom, but only one electron group bonded to another atom, is linear because there are only two atoms in the molecule.

Double or triple bonds count as a single electron group. The Lewis electron dot diagram of formaldehyde (CH_2O) is shown in Figure **4.10.9**.



Figure **4.10.9**: Lewis Electron Dot Diagram of Formaldehyde.

The central C atom has three electron groups around it because the double bond counts as one electron group. The three electron groups repel each other to adopt a trigonal planar shape.




Figure **4.10.10**: Formaldehyde bonding.

(The lone electron pairs on the O atom are omitted for clarity.) The molecule will not be a perfect equilateral triangle because the C–O double bond is different from the two C–H bonds, but both planar and triangular describe the appropriate approximate shape of this molecule.

Table **4.10.1** summarizes the shapes of molecules based on the number of electron groups and surrounding atoms.

Table 4.10.1: Summary of Molecular Shapes

Number of Electron Groups on Central Atom	Number of Bonding Groups	Number of Lone Pairs	Electron Geometry	Molecular Shape
2	2	0	linear	linear
3	3	0	trigonal planar	trigonal planar
3	2	1	trigonal planar	bent
4	4	0	tetrahedral	tetrahedral
4	3	1	tetrahedral	trigonal pyramidal
4	2	2	tetrahedral	bent

EXAMPLE 4.10.1


What is the approximate shape of each molecule?

a. PCl_3
b. NOF

Solution


The first step is to draw the Lewis structure of the molecule.

For PCl_3, the electron dot diagram is as follows:



The lone electron pairs on the Cl atoms are omitted for clarity. The P atom has four electron groups with three of them bonded to surrounding atoms, so the molecular shape is trigonal pyramidal.

The electron dot diagram for NOF is as follows:



The N atom has three electron groups on it, two of which are bonded to other atoms. The molecular shape is bent.

EXERCISE 4.10.1

What is the approximate molecular shape of CH_2Cl_2?

Answer

Tetrahedral

EXERCISE 4.10.2

Ethylene (C_2H_4) has two central atoms. Determine the geometry around each central atom and the shape of the overall molecule. (Hint: hydrogen is a terminal atom.)

Answer

Trigonal planar about both central C atoms.

SUMMARY

The approximate shape of a molecule can be predicted from the number of electron groups and the number of surrounding atoms.

4.10: Predicting the Shapes of Molecules is shared under a mixed 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.7: Predicting the Shapes of Molecules** by Henry Agnew, Marisa Alviar-Agnew is licensed notset.

4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix

Learning Objectives

- Explain how polar compounds differ from nonpolar compounds.
- Determine if a molecule is polar or nonpolar.
- Given a pair of compounds, predict which would have a higher melting or boiling point.

Bond Polarity

The ability of an atom in a molecule to attract shared electrons is called **electronegativity**. When two atoms combine, the difference between their electronegativities is an indication of the type of bond that will form. If the difference between the electronegativities of the two atoms is small, neither atom can take the shared electrons completely away from the other atom, and the bond will be covalent. If the difference between the electronegativities is large, the more electronegative atom will take the bonding electrons completely away from the other atom (electron transfer will occur), and the bond will be ionic. This is why metals (low electronegativities) bonded with nonmetals (high electronegativities) typically produce ionic compounds.

A bond may be so polar that an electron actually transfers from one atom to another, forming a true ionic bond. How do we judge the degree of polarity? Scientists have devised a scale called **electronegativity**, a scale for judging how much atoms of any element attract electrons. Electronegativity is a unitless number; the higher the number, the more an atom attracts electrons. A common scale for electronegativity is shown in Figure **4.11.1**.




Figure **4.11.1**: Electronegativities of the Elements. Electronegativities are used to determine the polarity of covalent bonds.

The polarity of a covalent bond can be judged by determining the *difference* of the electronegativities of the two atoms involved in the covalent bond, as summarized in the following table:

difference of the electronegativities of the two atoms involved in the covalent bond

Electronegativity Difference	Bond Type
0–0.4	pure covalent
0.5–2.0	polar covalent
>2.0	likely ionic

Nonpolar Covalent Bonds

A bond in which the electronegativity difference is less than 1.9 is considered to be mostly covalent in character. However, at this point we need to distinguish between two general types of covalent bonds. A **nonpolar covalent bond** is a covalent bond in which the bonding electrons are shared equally between the two atoms. In a nonpolar covalent bond, the distribution of electrical charge is balanced between the two atoms.




Figure **4.11.2**: A nonpolar covalent bond is one in which the distribution of electron density between the two atoms is equal.

The two chlorine atoms share the pair of electrons in the single covalent bond equally, and the electron density surrounding the $\mathbf{Cl_2}$ molecule is symmetrical. Also note that molecules in which the electronegativity difference is very small (<0.5) are also considered nonpolar covalent. An example would be a bond between chlorine and bromine ($\mathbf{\Delta EN = 3.0 - 2.8 = 0.2}$).

Polar Covalent Bonds

A bond in which the electronegativity difference between the atoms is between 0.5 and 2.0 is called a polar covalent bond. A **polar covalent bond** is a covalent bond in which the atoms have an unequal attraction for electrons and so the sharing is unequal. In a polar covalent bond, sometimes simply called a polar bond, the distribution of electrons around the molecule is no longer symmetrical.




Figure **4.11.3**: In the polar covalent bond of **HF**, the electron density is unevenly distributed. There is a higher density (red) near the fluorine atom, and a lower density (blue) near the hydrogen atom.

An easy way to illustrate the uneven electron distribution in a polar covalent bond is to use the Greek letter delta (δ).




Figure **4.11.4**: Use of δ to indicate partial charge.

The atom with the greater electronegativity acquires a partial negative charge, while the atom with the lesser electronegativity acquires a partial positive charge. The delta symbol is used to indicate that the quantity of charge is less than one. A crossed arrow can also be used to indicate the direction of greater electron density.



Figure **4.11.5**: Use of crossed arrow to indicate polarity.



Electronegativity differences in bonding using Pauling scale. Differences in electronegativity classify bonds as covalent, polar covalent, or ionic.

Example 4.11.1: Bond Polarity

What is the polarity of each bond?

a. C–H
b. O–H

Solution

Using Figure **4.11.1**, we can calculate the difference of the electronegativities of the atoms involved in the bond.

a. For the C–H bond, the difference in the electronegativities is 2.5 − 2.1 = 0.4. Thus we predict that this bond will be nonpolar covalent.
b. For the O–H bond, the difference in electronegativities is 3.5 − 2.1 = 1.4, so we predict that this bond will be polar covalent.

Exercise 4.11.1

What is the polarity of each bond?

a. Rb–F
b. P–Cl

Answer a

likely ionic

Answer b

polar covalent

Molecular Polarity

To determine if a molecule is polar or nonpolar, it is generally useful to look at Lewis structures. **Nonpolar compounds** will be symmetric, meaning all of the sides around the central atom are identical—bonded to the same element with no unshared pairs of electrons. **Polar molecules** are asymmetric, either containing lone pairs of electrons on a central atom or having atoms with different electronegativities bonded. This works pretty well, as long as you can visualize the molecular geometry. That's the hard part. To know how the bonds are oriented in space, you have to have a strong grasp of Lewis structures and VSEPR theory. Assuming that you do, you can look at the structure of each one and decide if it is polar or not, *whether or not you know the individual atom's electronegativity*. This is because you know that all bonds between dissimilar elements are polar, and in these particular examples, it doesn't matter which direction the dipole moment vectors are pointing (out or in).

A **polar molecule** is a molecule in which one end of the molecule is slightly positive, while the other end is slightly negative. A diatomic molecule that consists of a polar covalent bond, such as **HF**, is a polar molecule. The two electrically charged regions on either end of the molecule are called poles, similar to a magnet having a north and a south pole. A molecule with two poles is called a **dipole** (see figure below). Hydrogen fluoride is a dipole.



Figure **4.11.6**: A dipole is any molecule with a positive end and a negative end, resulting from unequal distribution of electron density throughout the molecule.

For molecules with more than two atoms, the molecular geometry must also be taken into account when determining if the molecule is polar or nonpolar. The figure below shows a comparison between carbon dioxide and water. Carbon dioxide (CO_2) is a linear molecule. The oxygen atoms are more electronegative than the carbon atom, so there are two individual dipoles pointing outward from the **C** atom to each **O** atom. However, since the dipoles are of equal strength and are oriented this way, they cancel out and the overall molecular polarity of CO_2 is zero.

Water is a bent molecule because of the two lone pairs on the central oxygen atom. The individual dipoles point from the **H** atoms toward the **O** atom. Because of the shape, the dipoles do not cancel each other out and the water molecule is polar. In the figure below, the net dipole is shown in blue and points upward.




Figure **4.11.7**: The molecular geometry of a molecule affects its polarity. Each CO bond has a dipole moment, but they point in opposite directions so that the net CO2 molecule is nonpolar. In contrast, water is polar because the OH bond moments do not cancel out.

Some other molecules are shown in the figure below. Notice that a tetrahedral molecule such as CH_4 is nonpolar. However, if one of the peripheral **H** atoms is replaced with another atom that has a different electronegativity, the molecule becomes polar. A trigonal planar molecule (BF_3) may be nonpolar if all three peripheral atoms are the same, but a trigonal pyramidal molecule (NH_3) is polar.




Figure **4.11.8**: Some examples of polar and nonpolar molecules based on molecular geometry.

To summarize, to be polar, a molecule must:

1. Contain at least one polar covalent bond.
2. Have a molecular structure such that the sum of the vectors of each bond dipole moment do not cancel.

Steps to Identify Polar Molecules

1. Draw the Lewis structure.
2. Figure out the geometry (using VSEPR theory).
3. Visualize or draw the geometry.
4. Find the net dipole moment (you don't have to actually do calculations if you can visualize it).
5. If the net dipole moment is zero, it is non-polar. Otherwise, it is polar.

Properties of Polar Molecules

Polar molecules tend to align when placed in an electric field with the positive end of the molecule oriented toward the negative plate and the negative end toward the positive plate (Figure **4.11.14**). We can use an electrically charged object to attract polar molecules, but nonpolar molecules are not attracted. Also, polar solvents are better at dissolving polar substances, and nonpolar solvents are better at dissolving nonpolar substances.




Figure **4.11.9**: (a) Molecules are always randomly distributed in the liquid state in the absence of an electric field. (b) When an electric field is applied, polar molecules like HF will align to the dipoles with the field direction. (OpenStax CC-BY-SA);

While molecules can be described as "polar covalent" or "ionic", it must be noted that this is often a relative term, with one molecule simply being *more polar* or *less polar* than another. However, the following properties are typical of such molecules. Polar molecules tend to:

- have higher melting points than nonpolar molecules
- have higher boiling points than nonpolar molecules
- be more soluble in water (dissolve better) than nonpolar molecules
- have lower vapor pressures than nonpolar molecules

Example 4.11.2:

Label each of the following as polar or nonpolar.

a. Water, H_2O:

b. Methanol, CH_3OH:

c. Hydrogen Cyanide, HCN: H–C≡N

d. Oxygen, O_2:

e. Propane, C_3H_8: 

Solution

a. Water is polar. Any molecule with lone pairs of electrons around the central atom is polar.
b. Methanol is polar. This is not a symmetric molecule. The $-OH$ side is different from the other 3 $-H$ sides.
c. Hydrogen cyanide is polar. The molecule is not symmetric. The nitrogen and hydrogen have different electronegativities, creating an uneven pull on the electrons.
d. Oxygen is nonpolar. The molecule is symmetric. The two oxygen atoms pull on the electrons by exactly the same amount.
e. Propane is nonpolar, because it is symmetric, with H atoms bonded to every side around the central atoms and no unshared pairs of electrons.

? Exercise 4.11.2

Label each of the following as polar or nonpolar.

a. SO_3
b. NH_3

Answer a

nonpolar

Answer b

polar

Contributions & Attributions

- StackExchange (thomij).

4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **10.8: Electronegativity and Polarity - Why Oil and Water Do not Mix** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.12: Kinetic Molecular Theory- A Model for Gases

Learning Objectives

- State the major concepts behind the kinetic theory of gases.
- Relate the general properties of gases to the kinetic theory.

Gases were among the first substances studied in terms of the modern scientific method, which was developed in the 1600s. It did not take long to recognize that gases all shared certain physical behaviors, suggesting that all gases could be described by one all-encompassing theory. Today, that theory is the **kinetic theory of gases**. It is based on the following statements:

1. Gases consist of tiny particles of matter that are in constant motion.
2. Gas particles are constantly colliding with each other and the walls of a container. These collisions are elastic; that is, there is no net loss of energy from the collisions.
3. Gas particles are separated by large distances, with the size of a gas particle tiny compared to the distances that separate them.
4. There are no interactive forces (i.e., attraction or repulsion) between the particles of a gas.
5. The average speed of gas particles is dependent on the temperature of the gas.

Figure **4.12.1** shows a representation of how we mentally picture the gas phase.




Figure **4.12.1**: The Kinetic Theory of Gases. The kinetic theory of gases describes this state of matter as composed of tiny particles in constant motion with a lot of distance between the particles.

This model of gases explains some of the physical properties of gases. Because most of a gas is empty space, a gas has a low density and can expand or contract under the appropriate influence. The fact that gas particles are in constant motion means that two or more gases will always mix, as the particles from the individual gases move and collide with each other.

An **ideal gas** is a gas that exactly follows the statements of the kinetic theory. Unfortunately, *real gases* are not ideal. Many gases deviate slightly from agreeing perfectly with the kinetic theory of gases. However, most gases adhere to the statements so well that the kinetic theory of gases is well accepted by the scientific community.

- The physical behavior of gases is explained by the kinetic theory of gases.
- An ideal gas adheres exactly to the kinetic theory of gases.

4.12: Kinetic Molecular Theory- A Model for Gases is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.2: Kinetic Molecular Theory- A Model for Gases** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0. Original source: https://www.ck12.org/c/chemistry/.

4.13: Pressure - The Result of Constant Molecular Collisions

Learning Objectives

- Define *pressure*.
- Learn the units of pressure and how to convert between them.

The kinetic theory of gases indicates that gas particles are always in motion and are colliding with other particles and the walls of the container holding them. Although collisions with container walls are elastic (i.e., there is no net energy gain or loss because of the collision), a gas particle does exert a force on the wall during the collision. The accumulation of all these forces distributed over the area of the walls of the container causes something we call pressure. **Pressure** (***P***) is defined as the force of all the gas particle/wall collisions divided by the area of the wall:

$$\text{pressure} = \frac{\text{force}}{\text{area}}$$

All gases exert pressure; it is one of the fundamental measurable quantities of this phase of matter. Even our atmosphere exerts pressure—in this case, the gas is being "held in" by the earth's gravity, rather than the gas being in a container. The pressure of the atmosphere is about 14.7 pounds of force for every square inch of surface area: 14.7 lb/in^2.

Pressure has a variety of units. The formal, SI-approved unit of pressure is the *pascal* (Pa), which is defined as 1 N/m^2 (one newton of force over an area of one square meter). However, this is usually too small in magnitude to be useful. A common unit of pressure is the **atmosphere** (atm), which was originally defined as the average atmospheric pressure at sea level.

However, "average atmospheric pressure at sea level" is difficult to pinpoint because of atmospheric pressure variations. A more reliable and common unit is **millimeters of mercury** (mmHg), which is the amount of pressure exerted by a column of mercury exactly 1 mm high. An equivalent unit is the **torr**, which equals 1 mmHg. (The torr is named after Evangelista Torricelli, a seventeenth-century Italian scientist who invented the mercury barometer.) With these definitions of pressure, the atmosphere unit is redefined: 1 atm is defined as exactly 760 mmHg, or 760 torr. We thus have the following equivalents:

$$1 \text{ atm} = 760 \text{ mmHg} = 760 \text{ torr}$$


We can use these equivalents as with any equivalence—to perform conversions from one unit to another. Relating these to the formal SI unit of pressure, 1 atm = 101,325 Pa.

Example 4.13.1: Pressure Conversion

How many atmospheres are there in 595 torr?

Solution

Solutions to Example 11.3.1

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 595 torr Find: ? atm
List other known quantities.	1 atm = 760 torr
Prepare a concept map.	 $\frac{1\ atm}{760\ torr}$
Cancel units and calculate.	$595\ \cancel{torr} \times \frac{1\ atm}{760\ \cancel{torr}} = 0.783\ atm$
Think about your result.	595 torr is less than 760 torr so the final answer should be less than 1 atm.

Exercise 4.13.1

How many atmospheres are there in 1,022 torr?

Answer


1.345 atm

Example 4.13.2: Mars

The atmosphere on Mars is largely CO_2 at a pressure of 6.01 mmHg. What is this pressure in atmospheres?

Solution

Solutions to Example 11.3.2

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 6.01mmHg Find: ? atm
List other known quantities.	1 atm = 760 mmHg
Prepare a concept map.	 $\frac{1\ atm}{760\ mmHg}$
Cancel units and calculate.	$6.01\ \cancel{mmHg} \times \frac{1\ atm}{760\ \cancel{mmHg}} = 0.00791\ atm = 7.91 \times 10^{-3} atm$
Think about your result.	6.01 is a very small number relative to 760 mmHg, just like the value in atmospheres.

Exercise 4.13.2

Atmospheric pressure is low in the eye of a hurricane. In a 1979 hurricane in the Pacific Ocean, a pressure of 0.859 atm was reported inside the eye. What is this pressure in torr?

Answer

652 torr

Summary

- Pressure is a force exerted over an area.
- Pressure has several common units that can be converted.

4.13: Pressure - The Result of Constant Molecular Collisions is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.3: Pressure - The Result of Constant Molecular Collisions** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.14: Boyle's Law - Pressure and Volume

Learning Objectives

- Learn what is meant by the term *gas laws*.
- Learn and apply Boyle's Law.

When seventeenth-century scientists began studying the physical properties of gases, they noticed some simple relationships between some of the measurable properties of the gas. Take pressure (P) and volume (V), for example. Scientists noted that for a given amount of a gas (usually expressed in units of moles [n]), if the temperature (T) of the gas was kept constant, pressure and volume were related: as one increases, the other decreases. As one decreases, the other increases. This means that pressure and volume are *inversely related*.

There is more to it, however: pressure and volume of a given amount of gas at constant temperature are *numerically* related. If you take the pressure value and multiply it by the volume value, the product is a constant for a given amount of gas at a constant temperature:

$$P \times V = \text{ constant at constant n and T}$$

If either volume or pressure changes while amount and temperature stay the same, then the other property must change so that the product of the two properties still equals that same constant. That is, if the original conditions are labeled P_1 and V_1 and the new conditions are labeled P_2 and V_2, we have

$$P_1V_1 = \text{constant} = P_2V_2$$

where the properties are assumed to be multiplied together. Leaving out the middle part, we have simply

$$P_1V_1 = P_2V_2 \text{ at constant n and T}$$

This equation is an example of a gas law. A **gas law** is a simple mathematical formula that allows you to model, or predict, the behavior of a gas. This particular gas law is called **Boyle's Law**, after the English scientist Robert Boyle, who first announced it in 1662. Figure **4.14.1** shows two representations of how Boyle's Law works.




Figure **4.14.1**: Boyle's Law. A piston having a certain pressure and volume (left piston) will have half the volume when its pressure is twice as much (right piston). One can also plot P versus V for a given amount of gas at a certain temperature; such a plot will look like the graph on the right.

Boyle's Law is an example of a second type of mathematical problem we see in chemistry—one based on a mathematical formula. Tactics for working with mathematical formulas are different from tactics for working with conversion factors. First, most of the questions you will have to answer using formulas are word-type questions, so the first step is to identify what quantities are known and assign them to variables. Second, in most formulas, some mathematical rearrangements (i.e., algebra) must be performed to solve for an unknown variable. The rule is that to find the value of the unknown variable, you must mathematically isolate the unknown variable *by itself and in the numerator* of one side of the equation. Finally, units must be consistent. For example, in Boyle's Law there are two pressure variables; they must have the same unit. There are also two volume variables; they also must have the same unit. In most cases, it won't matter *what* the unit is, but the unit must be the *same* on both sides of the equation.

Example 4.14.1

A sample of gas has an initial pressure of 2.44 atm and an initial volume of 4.01 L. Its pressure changes to 1.93 atm. What is the new volume if temperature and amount are kept constant?

Solution

Solutions to Example 11.8.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: P_1 = 2.44 atm and V_1 = 4.01 L P_2 = 1.93 atm Find: V_2 = ? L
List other known quantities.	none
Plan the problem.	First, rearrange the equation algebraically to solve for V_2. $$V_2 = \frac{P_1 \times V_1}{P_2}$$
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $$V_2 = \frac{2.44 \cancel{\text{ atm}} \times 4.01 \text{ L}}{1.93 \cancel{\text{ atm}}} = 5.07 \text{ L}$$
Think about your result.	We know that pressure and volume are inversely related; as one decreases, the other increases. Pressure is decreasing (from 2.44 atm to 1.93 atm), so volume should be increasing to compensate, and it is (from 4.01 L to 5.07 L). So the answer makes sense based on Boyle's Law.

Exercise 4.14.1

If P_1 = 334 torr, V_1 = 37.8 mL, and P_2 = 102 torr, what is V_2?

Answer

124 mL

As mentioned, you can use any units for pressure and volume, but both pressures must be expressed in the same units, and both volumes must be expressed in the same units.

Example 4.14.2:

A sample of gas has an initial pressure of 722 torr and an initial volume of 88.8 mL. Its volume changes to 0.663 L. What is the new pressure?

Solution

Solutions to Example 11.8.2

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: P_1 = 722 torr and V_1 = 88.8 mL V_2 = 0.633 L Find: P_2 = ? torr
List other known quantities.	1 L = 1000 mL to have the same units for volume.
Plan the problem.	1. Perform the conversion of the second volume unit from L to mL. 2. Rearrange the equation algebraically to solve for P_2. $$P_2 = \frac{P_1 \times V_1}{V_2}$$
Cancel units and calculate.	1. $$0.663 \cancel{L} \times \frac{1000\,ml}{1\,\cancel{L}} = 663\,ml$$ 2. Substitute the known quantities into the equation and solve. $$P_2 = \frac{722\text{ torr} \times 88.8\ \cancel{mL}}{663\ \cancel{mL}} = 96.7\text{ torr}$$
Think about your result.	When the volume increased, the pressure decreased, which is as expected for Boyle's Law.

? Exercise 4.14.2

If V_1 = 456 mL, P_1 = 308 torr, and P_2 = 1.55 atm, what is V_2?

Answer

119 mL

Summary

- The behavior of gases can be modeled with gas laws.
- Boyle's Law relates the pressure and volume of a gas at constant temperature and amount.

4.14: Boyle's Law - Pressure and Volume is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.4: Boyle's Law - Pressure and Volume** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.15: Charles's Law- Volume and Temperature

Learning Objectives

- Learn and apply Charles's Law.

Everybody enjoys the smell and taste of freshly-baked bread. It is light and fluffy as a result of the action of yeast on sugar. The yeast converts the sugar to carbon dioxide, which at high temperatures causes the dough to expand. The end result is an enjoyable treat, especially when covered with melted butter.

Charles's Law

French physicist Jacques Charles (1746-1823) studied the effect of temperature on the volume of a gas at constant pressure. **Charles's Law** states that the volume of a given mass of gas varies directly with the absolute temperature of the gas when pressure is kept constant. The absolute temperature is temperature measured with the Kelvin scale. The Kelvin scale must be used because zero on the Kelvin scale corresponds to a complete stop of molecular motion.




Figure 4.15.1: As a container of confined gas is heated, its molecules increase in kinetic energy and push the movable piston outward, resulting in an increase in volume.

Mathematically, the direct relationship of Charles's Law can be represented by the following equation:

$$\frac{V}{T} = k$$

As with Boyle's Law, k is constant only for a given gas sample. The table below shows temperature and volume data for a set amount of gas at a constant pressure. The third column is the constant for this particular data set and is always equal to the volume divided by the Kelvin temperature.

Table 4.15.1: Temperature-Volume Data

Temperature (K)	Volume (mL)	$\frac{V}{T} = k \left(\frac{\text{mL}}{\text{K}}\right)$
50	20	0.40
100	40	0.40
150	60	0.40
200	80	0.40
300	120	0.40
500	200	0.40
1000	400	0.40

When this data is graphed, the result is a straight line, indicative of a direct relationship, shown in the figure below.




Figure 4.15.2:The volume of a gas increases as the Kelvin temperature increases.

Notice that the line goes exactly toward the origin, meaning that as the absolute temperature of the gas approaches zero, its volume approaches zero. However, when a gas is brought to extremely cold temperatures, its molecules would eventually condense into the liquid state before reaching absolute zero. The temperature at which this change into the liquid state occurs varies for different gases.

Charles's Law can also be used to compare changing conditions for a gas. Now we use V_1 and T_1 to stand for the initial volume and temperature of a gas, while V_2 and T_2 stand for the final volume and temperature. The mathematical relationship of Charles's Law becomes:

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

This equation can be used to calculate any one of the four quantities if the other three are known. The direct relationship will only hold if the temperatures are expressed in Kelvin. Temperatures in Celsius will not work. Recall the relationship that $\mathrm{K} = {}^\circ\mathrm{C} + 273$.

Example 4.15.1:

A balloon is filled to a volume of $2.20\ \mathrm{L}$ at a temperature of $22^\circ\mathrm{C}$. The balloon is then heated to a temperature of $71^\circ\mathrm{C}$. Find the new volume of the balloon.

Solution

Solutions to Example 11.5.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: $V_1 = 2.20\ \text{L}$ and $T_1 = 22°\text{C} = 295\ \text{K}$ $T_2 = 71°\text{C} = 344\ \text{K}$ Find: V_2 = ? L
List other known quantities.	The temperatures have first been converted to Kelvin.
Plan the problem.	First, rearrange the equation algebraically to solve for V_2. $$V_2 = \frac{V_1 \times T_2}{T_1}$$
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $$V_2 = \frac{2.20\ \text{L} \times 344\ \cancel{\text{K}}}{295\ \cancel{\text{K}}} = 2.57\ \text{L}$$
Think about your result.	The volume increases as the temperature increases. The result has three significant figures.

? Exercise 4.15.1

If V_1 = 3.77 L and T_1 = 255 K, what is V_2 if T_2 = 123 K?

Answer

1.82 L

✔ Example 4.15.2:

A sample of a gas has an initial volume of 34.8 L and an initial temperature of −67°C. What must be the temperature of the gas for its volume to be 25.0 L?

Solution

Solutions to Example 11.5.2

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: Given:T_1 = -27°C and V_1 = 34.8 L V_2 = 25.0 L Find: T_2 = ? K
List other known quantities.	K = -27°C + 273
Plan the problem.	1. Convert the initial temperature to Kelvin 2. Rearrange the equation algebraically to solve for T_2. $$T_2 = \frac{V_2 \times T_1}{V_1}$$
Cancel units and calculate.	1. −67°C + 273 = 206 K 2. Substitute the known quantities into the equation and solve. $$T_2 = \frac{25.0 \cancel{\text{L}} \times 206 \text{ K}}{34.8 \cancel{\text{L}}} = 148 \text{ K}$$
Think about your result.	This is also equal to −125°C. As temperature decreases, volume decreases—which it does in this example.

? Exercise 4.15.2

If V_1 = 623 mL, T_1 = 255°C, and V_2 = 277 mL, what is T_2?

Answer

235 K, or −38°C

Summary

- Charles's Law relates the volume and temperature of a gas at constant pressure and amount.

4.15: Charles's Law- Volume and Temperature is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.5: Charles's Law- Volume and Temperature** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.16: Gay-Lussac's Law- Temperature and Pressure

Learning Objectives

- Explain Gay-Lussac's Law.

Propane tanks are widely used with barbeque grills. But it's not fun to find out halfway through grilling that you have run out of gas. You can buy gauges that measure the pressure inside the tank to see how much is left. The gauge measures pressure and will register a higher pressure on a hot day than it will on a cold day. So you need to take the air temperature into account when you decide whether or not to refill the tank before your next cook-out.

Gay-Lussac's Law

When the temperature of a sample of gas in a rigid container is increased, the pressure of the gas increases as well. The increase in kinetic energy results in the molecules of gas striking the walls of the container with more force, resulting in a greater pressure. The French chemist Joseph Gay-Lussac (1778-1850) discovered the relationship between the pressure of a gas and its absolute temperature. **Gay-Lussac's Law** states that the pressure of a given mass of gas varies directly with the absolute temperature of the gas, when the volume is kept constant. Gay-Lussac's Law is very similar to Charles's Law, with the only difference being the type of container. Whereas the container in a Charles's Law experiment is flexible, it is rigid in a Gay-Lussac's Law experiment.



Figure **4.16.1**: Joseph Gay-Lussac.

The mathematical expressions for Gay-Lussac's Law are likewise similar to those of Charles's Law:

$$\frac{P}{T} \text{ and } \frac{P_1}{T_1} = \frac{P_2}{T_2}$$

A graph of pressure vs. temperature also illustrates a direct relationship. As a gas is cooled at constant volume, its pressure continually decreases until the gas condenses to a liquid.

Example 4.16.1

The gas in an aerosol can is under a pressure of $3.00\ \text{atm}$ at a temperature of $25°\text{C}$. It is dangerous to dispose of an aerosol can by incineration. What would the pressure in the aerosol can be at a temperature of $845°\text{C}$?

Solution

Solutions to Example 11.10.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: $P_1 = 3.00\ \text{atm}$ $T_1 = 25°\text{C} = 298\ \text{K}$ $T_2 = 845°\text{C} = 1118\ \text{K}$ Find: $P_2 = ?\ \text{atm}$
List other known quantities.	The temperatures have first been converted to Kelvin.
Plan the problem.	First, rearrange the equation algebraically to solve for P_2. $$P_2 = \frac{P_1 \times T_2}{T_1}$$
Calculate.	Now substitute the known quantities into the equation and solve. $$P_2 = \frac{3.00\ \text{atm} \times 1118\ \cancel{\text{K}}}{298\ \cancel{\text{K}}} = 11.3\ \text{atm}$$
Think about your result.	The pressure increases dramatically due to a large increase in temperature.

Summary

- Pressure and temperature at constant volume are directly proportional.

4.16: Gay-Lussac's Law- Temperature and Pressure is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.11: Gay-Lussac's Law- Temperature and Pressure** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.17: The Combined Gas Law- Pressure, Volume, and Temperature

Learning Objectives

- Learn and apply the Combined Gas Law.

One thing we notice about all the gas laws is that, collectively, volume and pressure are always in the numerator, and temperature is always in the denominator. This suggests that we can propose a gas law that combines pressure, volume, and temperature. This gas law is known as the **Combined Gas Law**, and its mathematical form is

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2} \; at \; constant \; n$$

This allows us to follow changes in all three major properties of a gas. Again, the usual warnings apply about how to solve for an unknown algebraically (isolate it on one side of the equation in the numerator), units (they must be the same for the two similar variables of each type), and units of temperature must be in Kelvin.

Example 4.17.1:

A sample of gas at an initial volume of 8.33 L, an initial pressure of 1.82 atm, and an initial temperature of 286 K simultaneously changes its temperature to 355 K and its volume to 5.72 L. What is the final pressure of the gas?

Solution

Solutions to Example 11.4.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: V_1 = 8.33 L, P_1 = 1.82 atm, and T_1 = 286 K V2 = 5.72 L and T_2 = 355 K Find: $P2$ = ? atm
List other known quantities.	none
Plan the problem.	First, rearrange the equation algebraically to solve for V_2. $P_2 = \frac{P_1V_1T_2}{T_1V_2}$
Calculate.	Now substitute the known quantities into the equation and solve. $P_2 = \frac{(1.82\,atm)(8.33\,\cancel{L})(355\,\cancel{K})}{(286\,\cancel{K})(5.72\,\cancel{L})} = 3.22atm$
Think about your result.	Ultimately, the pressure increased, which would have been difficult to predict because two properties of the gas were changing.

Exercise 4.17.1

If P_1 = 662 torr, V_1 = 46.7 mL, T_1 = 266 K, P_2 = 409 torr, and T_2 = 371 K, what is V_2?

Answer

105 mL

As with other gas laws, if you need to determine the value of a variable in the denominator of the combined gas law, you can either cross-multiply all the terms or just take the reciprocal of the combined gas law. Remember, the variable you are solving for must be in the numerator and all by itself on one side of the equation.

Summary

- The Combined Gas Law relates pressure, volume, and temperature of a gas.

4.17: The Combined Gas Law- Pressure, Volume, and Temperature is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.6: The Combined Gas Law- Pressure, Volume, and Temperature** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles

Learning Objectives

- Explain the Ideal Gas Law.

There are a number of chemical reactions that require ammonia. In order to carry out the reaction efficiently, we need to know how much ammonia we have for stoichiometric purposes. Using gas laws, we can determine the number of moles present in the tank if we know the volume, temperature, and pressure of the system.

Ideal Gas Law

The Combined Gas Law shows that the pressure of a gas is inversely proportional to volume and directly proportional to temperature. Avogadro's Law shows that volume or pressure is directly proportional to the number of moles of gas. Putting these together leaves us with the following equation:

$$\frac{P_1 \times V_1}{T_1 \times n_1} = \frac{P_2 \times V_2}{T_2 \times n_2}$$

As with the other gas laws, we can also say that $\frac{(P \times V)}{(T \times n)}$ is equal to a constant. The constant can be evaluated provided that the gas being described is considered to be ideal.

The **Ideal Gas Law** is a single equation which relates the pressure, volume, temperature, and number of moles of an ideal gas. If we substitute in the variable R for the constant, the equation becomes:

$$\frac{P \times V}{T \times n} = R$$

The Ideal Gas Law is conveniently rearranged to look this way, with the multiplication signs omitted:

$$PV = nRT$$

The variable R in the equation is called the **ideal gas constant**.

Evaluating the Ideal Gas Constant

The value of R, the ideal gas constant, depends on the units chosen for pressure, temperature, and volume in the ideal gas equation. It is necessary to use Kelvin for the temperature and it is conventional to use the SI unit of liters for the volume. However, pressure is commonly measured in one of three units: $\mathbf{kPa}$, $\mathbf{atm}$, or $\mathbf{mm\ Hg}$. Therefore, R can have three different values.

We will demonstrate how R is calculated when the pressure is measured in $\mathbf{kPa}$. The volume of $\mathbf{1.00\ mol}$ of any gas at STP (Standard temperature, 273.15 K and pressure, 1 atm) is measured to be $\mathbf{22.414\ L}$. We can substitute $\mathbf{101.325\ kPa}$ for pressure, $\mathbf{22.414\ L}$ for volume, and $\mathbf{273.15\ K}$ for temperature into the ideal gas equation and solve for R.

$$\begin{aligned} R &= \frac{PV}{nT} \\ &= \frac{101.325\ \text{kPa} \times 22.414\ \text{L}}{1.000\ \text{mol} \times 273.15\ \text{K}} \\ &= 8.314\ \text{kPa} \cdot \text{L/K} \cdot \text{mol} \end{aligned}$$

This is the value of R that is to be used in the ideal gas equation when the pressure is given in $\mathbf{kPa}$. The table below shows a summary of this and the other possible values of R. It is important to choose the correct value of R to use for a given problem.

Table 4.18.1: Values of the Ideal Gas Constant

Unit of P	Unit of V	Unit of n	Unit of T	Value and Unit of R
kPa	L	mol	K	$8.314\ \text{J/K} \cdot \text{mol}$
atm	L	mol	K	$0.08206\ \text{L} \cdot \text{atm/K} \cdot \text{mol}$
mm Hg	L	mol	K	$62.36\ \text{L} \cdot \text{mm Hg/K} \cdot \text{mol}$

Notice that the unit for R when the pressure is in kPa has been changed to $\text{J/K} \cdot \text{mol}$. A kilopascal multiplied by a liter is equal to the SI unit for energy, a joule (J).

Example 4.18.1 Oxygen Gas

What volume is occupied by 3.76 g of oxygen gas at a pressure of 88.4 kPa and a temperature of 19°C? Assume the oxygen is ideal.

Solution

Solutions to Example 11.5.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: • $P = 88.4\ \text{kPa}$ • $T = 19°\text{C} = 292\ \text{K}$ Mass $\text{O}_2 = 3.76\ \text{g}$ Find: V = ? L
List other known quantities.	$\text{O}_2 = 32.00\ \text{g/mol}$ $R = 8.314\ \text{J/K} \cdot \text{mol}$
Plan the problem.	1. First, determine the number of moles of O_2 from the given mass and the molar mass. 2. Then, rearrange the equation algebraically to solve for V $$V = \frac{nRT}{P}$$
Calculate.	1. $$3.76\ \cancel{g} \times \frac{1\ \text{mol O}_2}{32.00\ \cancel{g}\ \text{O}_2} = 0.1175\ \text{mol O}_2$$ 2. Now substitute the known quantities into the equation and solve. $$V = \frac{nRT}{P}$$ $$= \frac{0.1175\ \cancel{\text{mol}} \times 8.314\ \cancel{\text{J/K}} \cdot \cancel{\text{mol}} \times 292\ \cancel{\text{K}}}{88.4\ \cancel{\text{kPa}}}$$ $$= 3.23\ \text{L O}_2$$
Think about your result.	The number of moles of oxygen is far less than one mole, so the volume should be fairly small compared to molar volume (22.4 L/mol) since the pressure and temperature are reasonably close to standard. The result has three significant figures because of the values for T and P. Since a joule $(\text{J}) = \text{kPa} \cdot \text{L}$, the units cancel out correctly, leaving a volume in liters.

Example 4.18.2: Argon Gas

A 4.22 mol sample of Ar has a pressure of 1.21 atm and a temperature of 34°C. What is its volume?

Solution

Solutions to Example 11.5.2

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: n = 4.22 mol P = 1.21 atm T = 34°C Find: V = ? L
List other known quantities.	none
Plan the problem.	1. The first step is to convert temperature to Kelvin. 2. Then, rearrange the equation algebraically to solve for V $$V = \frac{nRT}{P}$$
Calculate.	1. 34 + 273 = 307 K 2. Now substitute the known quantities into the equation and solve. $$V = \frac{(4.22\ \cancel{mol})(0.08205\frac{L.\cancel{atm}}{\cancel{mol.K}})(307\ \cancel{K})}{1.21\ \cancel{atm}}$$ $$= 87.9\ L$$
Think about your result.	The number of moles of Ar is large so the expected volume should also be large.

Exercise 4.18.1

A 0.0997 mol sample of O_2 has a pressure of 0.692 atm and a temperature of 333 K. What is its volume?

Answer

3.94 L

Exercise 4.18.2

For a 0.00554 mol sample of H_2, P = 23.44 torr and T = 557 K. What is its volume?

Answer

8.21 L

Summary

- The Ideal Gas Law is a single equation which relates the pressure, volume, temperature, and number of moles of an ideal gas.

4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.8: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

https://chem.libretexts.org/@go/page/492192

4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen

Learning Objectives

- Explain Dalton's Law of Partial Pressures.

The atmosphere of Venus is markedly different from that of Earth. The gases in the Venusian atmosphere are 96.5% carbon dioxide and 3% nitrogen. The atmospheric pressure on Venus is roughly 92 times that of Earth, so the amount of nitrogen on Venus would contribute a pressure well over $2700\ \text{mm Hg}$. And there is no oxygen present, so we couldn't breathe there. Not that we would want to go to Venus, as the surface temperature is usually over 460°C.

Dalton's Law of Partial Pressures

Gas pressure results from collisions between gas particles and the inside walls of their container. If more gas is added to a rigid container, the gas pressure increases. The identities of the two gases do not matter. John Dalton, the English chemist who proposed the atomic theory, also studied mixtures of gases. He found that each gas in a mixture exerts a pressure independently of every other gas in the mixture. For example, our atmosphere is composed of about 78% nitrogen and 21% oxygen, with smaller amounts of several other gases making up the rest. Since nitrogen makes up 78% of the gas particles in a given sample of air, it exerts 78% of the pressure. If the overall atmospheric pressure is $1.00\ \text{atm}$, then the pressure of just the nitrogen in the air is $0.78\ \text{atm}$. The pressure of the oxygen in the air is $0.21\ \text{atm}$.

The **partial pressure** of a gas is the contribution that gas makes to the total pressure when the gas is part of a mixture. The partial pressure of nitrogen is represented by P_{N_2}. **Dalton's Law of Partial Pressures** states that the total pressure of a mixture of gases is equal to the sum of all of the partial pressures of the component gases. Dalton's Law can be expressed with the following equation:

$$P_{\text{total}} = P_1 + P_2 + P_3 + \cdots$$

The figure below shows two gases that are in separate, equal-sized containers at the same temperature and pressure. Each exerts a different pressure, P_1 and P_2, reflective of the number of particles in the container. On the right, the two gases are combined into the same container, with no volume change. The total pressure of the gas mixture is equal to the sum of the individual pressures. If $P_1 = 300\ \text{mm Hg}$ and $P_2 = 500\ \text{mm Hg}$, then $P_{\text{total}} = 800\ \text{mm Hg}$.




Figure 4.19.1: Dalton's Law states that the pressure of a gas mixture is equal to the partial pressures of the combining gases.

Collecting Gases Over Water

You need to do a lab experiment where hydrogen gas is generated. In order to calculate the yield of gas, you have to know the pressure inside the tube where the gas is collected. But how can you get a barometer in there? Very simple: you don't. All you need

is the atmospheric pressure in the room. As the gas pushes out the water, it is pushing against the atmosphere, so the pressure inside is equal to the pressure outside.

Gas Collection by Water Displacement

Gases that are produced in laboratory experiments are often collected by a technique called **water displacement** (Figure 4.19.2). A bottle is filled with water and placed upside-down in a pan of water. The reaction flask is fitted with rubber tubing, which is then fed under the bottle of water. As the gas is produced in the reaction flask, it exits through the rubber tubing and displaces the water in the bottle. When the bottle is full of the gas, it can be sealed with a lid.



Figure 4.19.2: A gas produced in a chemical reaction can be collected by water displacement.

Because the gas is collected over water, it is not pure, but is mixed with vapor from the evaporation of the water. Dalton's Law can be used to calculate the amount of the desired gas by subtracting the contribution of the water vapor.

$$P_{\text{Total}} = P_g + P_{H_2O}$$

where P_g is the pressure of the desired gas, which can be solved for:

$$P_g = P_{Total} - P_{H_2O}$$

In order to solve a problem, it is necessary to know the vapor pressure of water at the temperature of the reaction (see table below). The sample problem illustrates the use of Dalton's Law when a gas is collected over water.

Table 4.19.1: Vapor Pressure of Water (mm Hg) at Selected Temperatures (°C)

0	5	10	15	20	25	30	35	40	45	50	55	60
4.58	6.54	9.21	12.79	17.54	23.76	31.82	42.18	55.32	71.88	92.51	118.04	149.38

Example 14.14.1

A certain experiment generates 2.58 L of hydrogen gas, which is collected over water. The temperature is 20°C and the atmospheric pressure is 98.60 kPa. Find the volume that the dry hydrogen would occupy at STP.

Solution

Step 1: List the known quantities and plan the problem.

Known

- $V_{\text{Total}} = 2.58\ \text{L}$
- $T = 20°\text{C} = 293\ \text{K}$
- $P_{\text{Total}} = 98.60\ \text{kPa} = 739.7\ \text{mm Hg}$

Unknown

- V_{H_2} at STP $= ?\ \text{L}$

The atmospheric pressure is converted from kPa to mm Hg in order to match units with the table. The sum of the pressures of the hydrogen and the water vapor is equal to the atmospheric pressure. The pressure of the hydrogen is found by subtraction. Then, the volume of the gas at STP can be calculated by using the combined gas law.

Step 2: Solve.

$$\begin{aligned} P_{H_2} &= P_{\text{Total}} - P_{H_2O} \\ &= 739,7 \text{ mm Hg} - 17.54 \text{ mm Hg} \\ &= 722.2 \text{ mm Hg} \end{aligned}$$

Now the combined gas law is used, solving for V_2, the volume of hydrogen at STP.

$$\begin{aligned} V_2 &= \frac{P_1 \times V_1 \times T_2}{P_2 \times T_1} \\ &= \frac{722.2 \text{ mm Hg} \times 2.58 \text{ L} \times 273 \text{ K}}{760 \text{ mm Hg} \times 293 \text{ K}} \\ &= 2.28 \text{ L } H_2 \end{aligned}$$

Step 3: Think about your result.

If the hydrogen gas were to be collected at STP and without the presence of the water vapor, its volume would be **2.28 L**. This is less than the actual collected volume because some of that is water vapor. The conversion using STP is useful for stoichiometry purposes.

Summary

- Dalton's Law of Partial Pressures states that the total pressure in a system is equal to the sum of the partial pressures of the gases present.
- The vapor pressure due to water in a sample can be corrected for, in order to get the true value for the pressure of the gas.

4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **11.9: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

CHAPTER OVERVIEW

5: Unit 5

5.1: Intermolecular Forces- Dispersion, Dipole–Dipole, Hydrogen Bonding, and Ion-Dipole
5.2: Solutions - Homogeneous Mixtures
5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy
5.4: Specifying Solution Concentration- Mass Percent
5.5: Specifying Solution Concentration- Molarity
5.6: Solution Stoichiometry
5.7: Sour Patch Kids and International Spy Movies
5.8: Acids- Properties and Examples
5.9: Bases- Properties and Examples
5.10: Molecular Definitions of Acids and Bases
5.11: Reactions of Acids and Bases
5.12: Acid–Base Titration
5.13: Strong and Weak Acids and Bases
5.14: Water - Acid and Base in One
5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity
5.16: Carbon - A Versitile Atom
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
5.18: Alkanes- Saturated Hydrocarbons
5.19: Isomers- Same Formula, Different Structure
5.20: Naming Alkanes
5.21: Alkenes and Alkynes
5.22: Functional Groups
5.23: Carbohydrates
5.24: Lipids
5.25: Proteins

5: Unit 5 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

5.1: INTERMOLECULAR FORCES- DISPERSION, DIPOLE–DIPOLE, HYDROGEN BONDING, AND ION-DIPOLE

LEARNING OBJECTIVES

- To describe the intermolecular forces in liquids.

The properties of liquids are intermediate between those of gases and solids, but are more similar to solids. In contrast to *intra*molecular forces, such as the covalent bonds that hold atoms together in molecules and polyatomic ions, *inter*molecular forces hold molecules together in a liquid or solid. Intermolecular forces are generally much weaker than covalent bonds. For example, it requires 927 kJ to overcome the intramolecular forces and break both O–H bonds in 1 mol of water, but it takes only about 41 kJ to overcome the intermolecular attractions and convert 1 mol of liquid water to water vapor at 100°C. (Despite this seemingly low value, the intermolecular forces in liquid water are among the strongest such forces known!) Given the large difference in the strengths of intra- and intermolecular forces, changes between the solid, liquid, and gaseous states almost invariably occur for molecular substances *without breaking covalent bonds*.

> *The properties of liquids are intermediate between those of gases and solids, but are more similar to solids.*

Intermolecular forces determine bulk properties, such as the melting points of solids and the boiling points of liquids. Liquids boil when the molecules have enough thermal energy to overcome the intermolecular attractive forces that hold them together, thereby forming bubbles of vapor within the liquid. Similarly, solids melt when the molecules acquire enough thermal energy to overcome the intermolecular forces that lock them into place in the solid.

Intermolecular forces are electrostatic in nature; that is, they arise from the interaction between positively and negatively charged species. Like covalent and ionic bonds, intermolecular interactions are the sum of both attractive and repulsive components. Because electrostatic interactions fall off rapidly with increasing distance between molecules, intermolecular interactions are most important for solids and liquids, where the molecules are close together. These interactions become important for gases only at very high pressures, where they are responsible for the observed deviations from the ideal gas law at high pressures.

In this section, we explicitly consider three kinds of intermolecular interactions. There are two additional types of electrostatic interaction that you are already familiar with: the ion–ion interactions that are responsible for ionic bonding, and the ion–dipole interactions that occur when ionic substances dissolve in a polar substance such as water. The first two are often described collectively as van der Waals forces.

DIPOLE–DIPOLE INTERACTIONS

Polar covalent bonds behave as if the bonded atoms have localized fractional charges that are equal but opposite (i.e., the two bonded atoms generate a *dipole*). If the structure of a molecule is such that the individual bond dipoles do not cancel one another, then the molecule has a net dipole moment. Molecules with net dipole moments tend to align themselves so that the positive end of one dipole is near the negative end of another and vice versa, as shown in Figure **5.1.1***a*.




Figure **5.1.1**: Attractive and Repulsive Dipole–Dipole Interactions. (a and b) Molecular orientations in which the positive end of one dipole (δ^+) is near the negative end of another (δ^-) (and vice versa) produce attractive interactions. (c and d) Molecular orientations that juxtapose the positive or negative ends of the dipoles on adjacent molecules produce repulsive interactions. (CC BY-SA-NC; anonymous)

These arrangements are more stable than arrangements in which two positive or two negative ends are adjacent (Figure **5.1.1***c*). Hence dipole–dipole interactions, such as those in Figure **5.1.1***b*, are *attractive intermolecular interactions*, whereas those in Figure **5.1.1***d* are *repulsive intermolecular interactions*. Because molecules in a liquid move freely and continuously, molecules always experience both

attractive and repulsive dipole–dipole interactions simultaneously, as shown in Figure **5.1.2**. On average, however, the attractive interactions dominate.




Figure **5.1.2**: Both attractive and repulsive dipole–dipole interactions occur in a liquid sample with many molecules. (CC BY-SA-NC; anonymous)

The green arrows pointing towards each other represent attraction. The gray arrows pointing away from each other represent repulsion

Because each end of a dipole possesses only a fraction of the charge of an electron, dipole–dipole interactions are substantially weaker than the interactions between two ions, each of which has a charge of at least ±1, or between a dipole and an ion, in which one of the species has at least a full positive or negative charge. In addition, the attractive interaction between dipoles falls off much more rapidly with increasing distance than do the ion–ion interactions. Recall that the attractive energy between two ions is proportional to $1/r$, where r is the distance between the ions. Doubling the distance ($r \rightarrow 2r$) decreases the attractive energy by one-half. In contrast, the energy of the interaction of two dipoles is proportional to $1/r^3$, so doubling the distance between the dipoles decreases the strength of the interaction by 2^3, or 8-fold. Thus a substance such as **HCl**, which is partially held together by dipole–dipole interactions, is a gas at room temperature and 1 atm pressure. Conversely, **NaCl**, which is held together by interionic interactions, is a high-melting-point solid. Within a series of compounds of similar molar mass, the strength of the intermolecular interactions increases as the dipole moment of the molecules increases, as shown in Table **5.1.1**.

Table **5.1.1**: Relationships Between the Dipole Moment and the Boiling Point for Organic Compounds of Similar Molar Mass

Compound	Molar Mass (g/mol)	Dipole Moment (D)	Boiling Point (K)
C_3H_6 (cyclopropane)	42	0	240
CH_3OCH_3 (dimethyl ether)	46	1.30	248
CH_3CN (acetonitrile)	41	3.9	355


> *The attractive energy between two ions is proportional to 1/r, whereas the attractive energy between two dipoles is proportional to 1/r6.*



Video Discussing Dipole Intermolecular Forces. Source: Dipole Intermolecular Force, YouTube(opens in new window) [youtu.be]

EXAMPLE 5.1.1

Arrange ethyl methyl ether ($CH_3OCH_2CH_3$), 2-methylpropane [isobutane, $(CH_3)_2CHCH_3$], and acetone (CH_3COCH_3) in order of increasing boiling points. Their structures are as follows:



2-Methylpropane Ethyl methyl ether Acetone

Given: compounds.

Asked for: order of increasing boiling points.

Strategy:

Compare the molar masses and the polarities of the compounds. Compounds with higher molar masses and that are polar will have the highest boiling points.

Solution:

The three compounds have essentially the same molar mass (58–60 g/mol), so we must look at differences in polarity to predict the strength of the intermolecular dipole–dipole interactions and thus the boiling points of the compounds.

The first compound, 2-methylpropane, contains only C–H bonds, which are not very polar because C and H have similar electronegativities. It should therefore have a very small (but nonzero) dipole moment and a very low boiling point.

Ethyl methyl ether has a structure similar to H_2O; it contains two polar C–O single bonds oriented at about a 109° angle to each other, in addition to relatively nonpolar C–H bonds. As a result, the C–O bond dipoles partially reinforce one another and generate a significant dipole moment that should give a moderately high boiling point.

Acetone contains a polar C=O double bond oriented at about 120° to two methyl groups with nonpolar C–H bonds. The C–O bond dipole therefore corresponds to the molecular dipole, which should result in both a rather large dipole moment and a high boiling point.

Thus we predict the following order of boiling points:

2-methylpropane < ethyl methyl ether < acetone

This result is in good agreement with the actual data: 2-methylpropane, boiling point = −11.7°C, and the dipole moment (μ) = 0.13 D; methyl ethyl ether, boiling point = 7.4°C and μ = 1.17 D; acetone, boiling point = 56.1°C and μ = 2.88 D.

? EXERCISE 5.1.1

Arrange carbon tetrafluoride (CF_4), ethyl methyl sulfide ($CH_3SC_2H_5$), dimethyl sulfoxide [$(CH_3)_2S{=}O$], and 2-methylbutane [isopentane, $(CH_3)_2CHCH_2CH_3$] in order of decreasing boiling points.

Answer

dimethyl sulfoxide (boiling point = 189.9°C) > ethyl methyl sulfide (boiling point = 67°C) > 2-methylbutane (boiling point = 27.8°C) > carbon tetrafluoride (boiling point = −128°C)

LONDON DISPERSION FORCES

Thus far, we have considered only interactions between polar molecules. Other factors must be considered to explain why many nonpolar molecules, such as bromine, benzene, and hexane, are liquids at room temperature; why others, such as iodine and naphthalene, are solids. Even the noble gases can be liquefied or solidified at low temperatures, high pressures, or both (Table **5.1.2**).

What kind of attractive forces can exist between nonpolar molecules or atoms? This question was answered by Fritz London (1900–1954), a German physicist who later worked in the United States. In 1930, London proposed that temporary fluctuations in the electron distributions within atoms and nonpolar molecules could result in the formation of short-lived instantaneous dipole moments, which produce attractive forces called London dispersion forces between otherwise nonpolar substances.

Table **5.1.2**: Normal Melting and Boiling Points of Some Elements and Nonpolar Compounds

Substance	Molar Mass (g/mol)	Melting Point (°C)	Boiling Point (°C)
Ar	40	−189.4	−185.9
Xe	131	−111.8	−108.1
N_2	28	−210	−195.8
O_2	32	−218.8	−183.0
F_2	38	−219.7	−188.1
I_2	254	113.7	184.4
CH_4	16	−182.5	−161.5

Consider a pair of adjacent He atoms, for example. On average, the two electrons in each He atom are uniformly distributed around the nucleus. Because the electrons are in constant motion, however, their distribution in one atom is likely to be asymmetrical at any given instant, resulting in an instantaneous dipole moment. As shown in part (a) in Figure **5.1.3**, the instantaneous dipole moment on one atom can interact with the electrons in an adjacent atom, pulling them toward the positive end of the instantaneous dipole or repelling them from the negative end. The net effect is that the first atom causes the temporary formation of a dipole, called an induced dipole, in the second. Interactions between these temporary dipoles cause atoms to be attracted to one another. These attractive interactions are weak and fall off rapidly with increasing distance. London was able to show with quantum mechanics that the attractive energy between molecules due to temporary dipole–induced dipole interactions falls off as $1/r^6$. Doubling the distance therefore decreases the attractive energy by 2^6, or 64-fold.




Figure **5.1.3**: Instantaneous Dipole Moments. The formation of an instantaneous dipole moment on one He atom (a) or an H_2 molecule (b) results in the formation of an induced dipole on an adjacent atom or molecule.

Instantaneous dipole–induced dipole interactions between nonpolar molecules can produce intermolecular attractions just as they produce interatomic attractions in monatomic substances like Xe. This effect, illustrated for two H_2 molecules in part (b) in Figure **5.1.3**, tends to become more pronounced as atomic and molecular masses increase (Table **5.1.2**). For example, Xe boils at −108.1°C, whereas He boils at −269°C. The reason for this trend is that the strength of London dispersion forces is related to the ease with which the electron distribution in a given atom can be perturbed. In small atoms such as He, the two 1*s* electrons are held close to the nucleus in a very small volume, and electron–electron repulsions are strong enough to prevent significant asymmetry in their distribution. In larger atoms such as Xe, however, the outer electrons are much less strongly attracted to the nucleus because of filled intervening shells. As a result, it is relatively easy to temporarily deform the electron distribution to generate an instantaneous or induced dipole. The ease of deformation of the electron distribution in an atom or molecule is called its polarizability. Because the electron distribution is more easily perturbed in large, heavy species than in small, light species, we say that heavier substances tend to be much more *polarizable* than lighter ones.

> *For similar substances, London dispersion forces get stronger with increasing molecular size.*

The polarizability of a substance also determines how it interacts with ions and species that possess permanent dipoles. Thus, London dispersion forces are responsible for the general trend toward higher boiling points with increased molecular mass and greater surface area in a homologous series of compounds, such as the alkanes (part (a) in Figure **5.1.4**). The strengths of London dispersion forces also depend significantly on molecular shape because shape determines how much of one molecule can interact with its neighboring molecules at any given time. For example, part (b) in Figure **5.1.4** shows 2,2-dimethylpropane (neopentane) and *n*-pentane, both of which have the empirical formula C_5H_{12}. Neopentane is almost spherical, with a small surface area for intermolecular interactions, whereas *n*-pentane has an extended conformation that enables it to come into close contact with other *n*-pentane molecules. As a result, the boiling point of neopentane (9.5°C) is more than 25°C lower than the boiling point of *n*-pentane (36.1°C).




Figure **5.1.4**: Mass and Surface Area Affect the Strength of London Dispersion Forces. (a) In this series of four simple alkanes, larger molecules have stronger London forces between them than smaller molecules do, and consequently have higher boiling points. (b) Linear *n*-pentane molecules have a larger surface area and stronger intermolecular forces than spherical neopentane molecules. As a result, neopentane is a gas at room temperature, whereas *n*-pentane is a volatile liquid.

All molecules, whether polar or nonpolar, are attracted to one another by London dispersion forces in addition to any other attractive forces that may be present. In general, however, dipole–dipole interactions in small polar molecules are significantly stronger than London dispersion forces, so the former predominate.



Video Discussing London/Dispersion Intermolecular Forces. Source: Dispersion Intermolecular Force, YouTube(opens in new window) [youtu.be]

EXAMPLE 5.1.2

Arrange *n*-butane, propane, 2-methylpropane [isobutene, $(CH_3)_2CHCH_3$], and *n*-pentane in order of increasing boiling points.

Given: compounds

Asked for: order of increasing boiling points

Strategy:

Determine the intermolecular forces in the compounds, and then arrange the compounds according to the strength of those forces. The substance with the weakest forces will have the lowest boiling point.

Solution:

The four compounds are alkanes and nonpolar, so London dispersion forces are the only important intermolecular forces. These forces are generally stronger with increasing molecular mass, so propane should have the lowest boiling point and *n*-pentane should have the highest, with the two butane isomers falling in between. Of the two butane isomers, 2-methylpropane is more compact, and *n*-butane has the more extended shape. Consequently, we expect intermolecular interactions for *n*-butane to be stronger due to its larger surface area, resulting in a higher boiling point. The overall order is thus as follows, with actual boiling points in parentheses: propane (−42.1°C) < 2-methylpropane (−11.7°C) < *n*-butane (−0.5°C) < *n*-pentane (36.1°C).

? EXERCISE 5.1.2

Arrange GeH_4, $SiCl_4$, SiH_4, CH_4, and $GeCl_4$ in order of decreasing boiling points.

Answer

$GeCl_4$ (87°C) > $SiCl_4$ (57.6°C) > GeH_4 (−88.5°C) > SiH_4 (−111.8°C) > CH_4 (−161°C)

HYDROGEN BONDS

Molecules with hydrogen atoms bonded to electronegative atoms such as O, N, and F (and to a much lesser extent, Cl and S) tend to exhibit unusually strong intermolecular interactions. These result in much higher boiling points than are observed for substances in which London dispersion forces dominate, as illustrated for the covalent hydrides of elements of groups 14–17 in Figure **5.1.5**. Methane and its heavier congeners in group 14 form a series whose boiling points increase smoothly with increasing molar mass. This is the expected trend in nonpolar molecules, for which London dispersion forces are the exclusive intermolecular forces. In contrast, the hydrides of the lightest members of groups 15–17 have boiling points that are more than 100°C greater than predicted on the basis of their molar masses. The effect is most dramatic for water: if we extend the straight line connecting the points for H_2Te and H_2Se to the line for period 2, we obtain an estimated boiling point of −130°C for water! Imagine the implications for life on Earth if water boiled at −130°C rather than 100°C.




Figure **5.1.5**: The Effects of Hydrogen Bonding on Boiling Points. These plots of the boiling points of the covalent hydrides of the elements of groups 14–17 show that the boiling points of the lightest members of each series for which hydrogen bonding is possible (HF, NH_3, and H_2O) are anomalously high for compounds with such low molecular masses.

Group 14 is in purple, group 15 is in green, group 16 is red, and group 17 is blue. Graph of boiling point against period.

Why do strong intermolecular forces produce such anomalously high boiling points and other unusual properties, such as high enthalpies of vaporization and high melting points? The answer lies in the highly polar nature of the bonds between hydrogen and very electronegative elements such as O, N, and F. The large difference in electronegativity results in a large partial positive charge on hydrogen and a correspondingly large partial negative charge on the O, N, or F atom. Consequently, H–O, H–N, and H–F bonds have very large bond dipoles that can interact strongly with one another. Because a hydrogen atom is so small, these dipoles can also approach one another more closely than most other dipoles. The combination of large bond dipoles and short dipole–dipole distances results in very strong dipole–dipole interactions called hydrogen bonds, as shown for ice in Figure **5.1.6**. A hydrogen bond is usually indicated by a dotted line between the hydrogen atom attached to O, N, or F (the *hydrogen bond donor*) and the atom that has the lone pair of electrons (the *hydrogen bond acceptor*). Because each water molecule contains two hydrogen atoms and two lone pairs, a tetrahedral arrangement maximizes the number of hydrogen bonds that can be formed. In the structure of ice, each oxygen atom is surrounded by a distorted tetrahedron of hydrogen atoms that form bridges to the oxygen atoms of adjacent water molecules. The bridging hydrogen atoms are *not* equidistant from the two oxygen atoms they connect, however. Instead, each hydrogen atom is 101 pm from one oxygen and 174 pm from the other. In contrast, each oxygen atom is bonded to two H atoms at the shorter distance and two at the longer distance, corresponding to two O–H covalent bonds and two O···H hydrogen bonds from adjacent water molecules, respectively. The resulting open, cagelike structure of ice means that the solid is actually slightly less dense than the liquid, which explains why ice floats on water, rather than sinks.




Figure **5.1.6**: The Hydrogen-Bonded Structure of Ice

Each water molecule accepts two hydrogen bonds from two other water molecules and donates two hydrogen atoms to form hydrogen bonds with two more water molecules, producing an open, cage like structure. The structure of liquid water is very similar, but in the liquid, the hydrogen bonds are continually broken and formed because of rapid molecular motion.

> *Hydrogen bond formation requires both a hydrogen bond donor and a hydrogen bond acceptor.*

Because ice is less dense than liquid water, rivers, lakes, and oceans freeze from the top down. In fact, the ice forms a protective surface layer that insulates the rest of the water, allowing fish and other organisms to survive in the lower levels of a frozen lake or sea. If ice were denser than the liquid, the ice formed at the surface in cold weather would sink as fast as it formed. Bodies of water would freeze from the bottom up, which would be lethal for most aquatic creatures. The expansion of water when freezing also explains why automobile or boat engines must be protected by “antifreeze” and why unprotected pipes in houses break if they are allowed to freeze.



Video Discussing Hydrogen Bonding Intermolecular Forces. Source: Hydrogen Bonding Intermolecular Force, YouTube(opens in new window) [youtu.be]

EXAMPLE 5.1.3

Considering CH_3OH, C_2H_6, Xe, and $(CH_3)_3N$, which can form hydrogen bonds with themselves? Draw the hydrogen-bonded structures.

Given: compounds

Asked for: formation of hydrogen bonds and structure

Strategy:


A. Identify the compounds with a hydrogen atom attached to O, N, or F. These are likely to be able to act as hydrogen bond donors.

B. Of the compounds that can act as hydrogen bond donors, identify those that also contain lone pairs of electrons, which allow them to be hydrogen bond acceptors. If a substance is both a hydrogen donor and a hydrogen bond acceptor, draw a structure showing the hydrogen bonding.

Solution:

A. Of the species listed, xenon (Xe), ethane (C_2H_6), and trimethylamine [$(CH_3)_3N$] do not contain a hydrogen atom attached to O, N, or F; hence they cannot act as hydrogen bond donors.

B. The one compound that can act as a hydrogen bond donor, methanol (CH_3OH), contains both a hydrogen atom attached to O (making it a hydrogen bond donor) and two lone pairs of electrons on O (making it a hydrogen bond acceptor); methanol can thus form hydrogen bonds by acting as either a hydrogen bond donor or a hydrogen bond acceptor. The hydrogen-bonded structure of methanol is as follows:




Hydrogen bonding in methanol

EXERCISE 5.1.3


Considering CH_3CO_2H, $(CH_3)_3N$, NH_3, and CH_3F, which can form hydrogen bonds with themselves? Draw the hydrogen-bonded structures.

Answer

CH_3CO_2H and NH_3;



Hydrogen bonding in ammonia



Hydrogen bonding in acetic acid

Hydrogen bonding in ammonia between nitrogen and hydrogen. hydrogen bonding in acetic acid is between oxygen and hydrogen.

Although hydrogen bonds are significantly weaker than covalent bonds, with typical dissociation energies of only 15–25 kJ/mol, they have a significant influence on the physical properties of a compound. Compounds such as HF can form only two hydrogen bonds at a time as can, on average, pure liquid NH_3. Consequently, even though their molecular masses are similar to that of water, their boiling points are significantly lower than the boiling point of water, which forms *four* hydrogen bonds at a time.

EXAMPLE 5.1.4: BUCKYBALLS

Arrange C_{60} (buckminsterfullerene, which has a cage structure), NaCl, He, Ar, and N_2O in order of increasing boiling points.

Given: compounds.

Asked for: order of increasing boiling points.

Strategy:

Identify the intermolecular forces in each compound and then arrange the compounds according to the strength of those forces. The substance with the weakest forces will have the lowest boiling point.

Solution

Electrostatic interactions are strongest for an ionic compound, so we expect NaCl to have the highest boiling point. To predict the relative boiling points of the other compounds, we must consider their polarity (for dipole–dipole interactions), their ability to form hydrogen bonds, and their molar mass (for London dispersion forces). Helium is nonpolar and by far the lightest, so it should have the lowest boiling point. Argon and N_2O have very similar molar masses (40 and 44 g/mol, respectively), but N_2O is polar while Ar is not. Consequently, N_2O should have a higher boiling point. A C_{60} molecule is nonpolar, but its molar mass is 720 g/mol, much greater than that of Ar or N_2O. Because the boiling points of nonpolar substances increase rapidly with molecular mass, C_{60} should boil at a higher temperature than the other nonionic substances. The predicted order is thus as follows, with actual boiling points in parentheses:

$$\text{He } (-269^\circ\text{C}) < \text{Ar } (-185.7^\circ\text{C}) < N_2O\ (-88.5^\circ\text{C}) < C_{60}\ (>280^\circ\text{C}) < \text{NaCl } (1465^\circ\text{C}).$$

? EXERCISE 5.1.4

Arrange 2,4-dimethylheptane, Ne, CS_2, Cl_2, and KBr in order of decreasing boiling points.

Answer

KBr (1435°C) > 2,4-dimethylheptane (132.9°C) > CS_2 (46.6°C) > Cl_2 (−34.6°C) > Ne (−246°C)

✓ EXAMPLE 5.1.5

Identify the most significant intermolecular force in each substance.

a. C_3H_8
b. CH_3OH
c. H_2S

Solution

a. Although C–H bonds are polar, they are only minimally polar. The most significant intermolecular force for this substance would be dispersion forces.
b. This molecule has an H atom bonded to an O atom, so it will experience hydrogen bonding.
c. Although this molecule does not experience hydrogen bonding, the Lewis electron dot diagram and VSEPR indicate that it is bent, so it has a permanent dipole. The most significant force in this substance is dipole-dipole interaction.

? EXERCISE 5.1.6

Identify the most significant intermolecular force in each substance.

a. HF
b. HCl

Answer a

hydrogen bonding

Answer b

dipole-dipole interactions

SUMMARY

Intermolecular forces are electrostatic in nature and include van der Waals forces and hydrogen bonds. Molecules in liquids are held to other molecules by intermolecular interactions, which are weaker than the intramolecular interactions that hold the atoms together within molecules and polyatomic ions. Transitions between the solid and liquid, or the liquid and gas phases, are due to changes in intermolecular interactions, but do not affect intramolecular interactions. The three major types of intermolecular interactions are dipole–dipole interactions, London dispersion forces (these two are often referred to collectively as **van der Waals forces**), and hydrogen bonds. **Dipole–dipole interactions** arise from the electrostatic interactions of the positive and negative ends of molecules with permanent dipole moments; their strength is proportional to the magnitude of the dipole moment and to $1/r^3$, where r is the distance between dipoles. **London dispersion forces** are due to the formation of **instantaneous dipole moments** in polar or nonpolar molecules as a result of short-lived fluctuations of electron charge distribution, which in turn cause the temporary formation of an **induced dipole** in adjacent molecules; their

energy falls off as $1/r^6$. Larger atoms tend to be more **polarizable** than smaller ones, because their outer electrons are less tightly bound and are therefore more easily perturbed. **Hydrogen bonds** are especially strong dipole–dipole interactions between molecules that have hydrogen bonded to a highly electronegative atom, such as O, N, or F. The resulting partially positively charged H atom on one molecule (the *hydrogen bond donor*) can interact strongly with a lone pair of electrons of a partially negatively charged O, N, or F atom on adjacent molecules (the *hydrogen bond acceptor*). Because of strong O···H hydrogen bonding between water molecules, water has an unusually high boiling point, and ice has an open, cage like structure that is less dense than liquid water.

5.1: Intermolecular Forces- Dispersion, Dipole–Dipole, Hydrogen Bonding, and Ion-Dipole is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by LibreTexts.

- **11.2: Intermolecular Forces** is licensed CC BY-NC-SA 3.0.

5.2: SOLUTIONS - HOMOGENEOUS MIXTURES

LEARNING OBJECTIVES

- Learn terminology involving solutions.
- Explain the significance of the statement "like dissolves like."
- Explain why certain substances dissolve in other substances.

The major component of a solution is called the **solvent.** The minor component of a solution is called the **solute**. By major and minor we mean whichever component has the greater or lesser presence by mass or by moles. Sometimes this becomes confusing, especially with substances with very different molar masses. However, here we will confine the discussion to solutions for which the major component and the minor component are obvious.



Figure 5.2.1: Making a saline water solution by dissolving table salt (NaCl) in water. The salt is the solute and the water the solvent. (CC-BY-SA 3.0; Chris 73).

Solutions exist for every possible phase of the solute and the solvent. Salt water, for example, is a solution of solid $\mathbf{NaCl}$ in liquid water, while air is a solution of a gaseous solute ($\mathbf{O_2}$) in a gaseous solvent ($\mathbf{N_2}$). In all cases, however, the overall phase of the solution is the same phase as the solvent. Table 5.2.1 lists some common types of solutions, with examples of each.

Table 5.2.1: Types of Solutions

Solvent Phase	Solute Phase	Example
gas	gas	air
liquid	gas	carbonated beverages
liquid	liquid	ethanol (C_2H_5OH) in H_2O (alcoholic beverages)
liquid	solid	salt water
solid	gas	H_2 gas absorbed by Pd metal
solid	liquid	Hg(ℓ) in dental fillings
solid	solid	steel alloys

EXAMPLE 5.2.1: SUGAR AND WATER

A solution is made by dissolving 1.00 g of sucrose ($\mathbf{C_{12}H_{22}O_{11}}$) in 100.0 g of liquid water. Identify the solvent and solute in the resulting solution.

Solution

Either by mass or by moles, the obvious minor component is **sucrose**, so it is **the solute**. **Water**—the majority component—is **the solvent**. The fact that the resulting solution is the same phase as water also suggests that water is the solvent.

EXERCISE 5.2.1

A solution is made by dissolving 3.33 g of $\mathbf{HCl(g)}$ in 40.0 g of liquid methyl alcohol ($\mathbf{CH_3OH}$). Identify the solvent and solute in the resulting solution.

Answer

solute: $\mathbf{HCl(g)}$

solvent: CH_3OH

LIKE DISSOLVES LIKE

A simple way to predict which compounds will dissolve in other compounds is the phrase "like dissolves like". What this means is that polar compounds dissolve polar compounds, nonpolar compounds dissolve nonpolar compounds, but polar and nonpolar do not dissolve in each other.

Even some nonpolar substances dissolve in water but only to a limited degree. Have you ever wondered why fish are able to breathe? Oxygen gas, a nonpolar molecule, does dissolve in water—it is this oxygen that the fish take in through their gills. The reason we can enjoy carbonated sodas is also due to a nonpolar compound that dissolves in water. Pepsi-cola and all the other sodas have carbon dioxide gas, CO_2, a nonpolar compound, dissolved in a sugar-water solution. In this case, to keep as much gas in solution as possible, the sodas are kept under pressure.

This general trend of "like dissolves like" is summarized in the following table:

Table 5.2.2: Summary of Solubilities

Solute (Polarity of Compound)	Solvent (Polarity of Compound)	Dominant Intermolecular Force	Is Solution Formed?
Polar	Polar	Dipole-Dipole Force and/or Hydrogen Bond	yes
Non-polar	Non-polar	Dispersion Force	yes
Polar	Non-polar		no
Non-polar	Polar		no
Ionic	Polar	Ion-Dipole	yes
Ionic	Non-polar		no

Note that every time charged particles (ionic compounds or polar substances) are mixed, a solution is formed. When particles with no charges (nonpolar compounds) are mixed, they will form a solution. However, if substances with charges are mixed with other substances without charges, a solution does not form. When an ionic compound is considered "insoluble", it doesn't necessarily mean the compound is completely untouched by water. All ionic compounds dissolve to some extent. An insoluble compound just doesn't dissolve in any noticeable or appreciable amount.

What is it that makes a solute soluble in some solvents but not others?

The answer is intermolecular interactions. The intermolecular interactions include London dispersion forces, dipole-dipole interactions, and hydrogen bonding (as described in Chapter 10). From experimental studies, it has been determined that if molecules of a solute experience the same intermolecular forces that the solvent does, the solute will likely dissolve in that solvent. So, NaCl—a very polar substance because it is composed of ions—dissolves in water, which is very polar, but not in oil, which is generally nonpolar. Nonpolar wax dissolves in nonpolar hexane, but not in polar water.



Figure 5.2.2: Water (clear liquid) and oil (yellow) do not form liquid solutions. (CC BY-SA 1.0 Generic; Victor Blacus)

EXAMPLE 5.2.2: POLAR AND NONPOLAR SOLVENTS

Would I_2 be more soluble in CCl_4 or H_2O? Explain your answer.

Solution

I_2 is nonpolar. Of the two solvents, CCl_4 is nonpolar and H_2O is polar, so I_2 would be expected to be more soluble in CCl_4.

? EXERCISE 5.2.2

Would C_3H_7OH be more soluble in CCl_4 or H_2O? Explain your answer.

Answer

H_2O, because both experience hydrogen bonding.

✓ EXAMPLE 5.2.3

Water is considered a polar solvent. Which substances should dissolve in water?

a. methanol (CH_3OH)
b. sodium sulfate (\ce{Na2SO4}\))
c. octane (C_8H_{18})

Solution

Because water is polar, substances that are polar or ionic will dissolve in it.

a. Because of the OH group in methanol, we expect its molecules to be polar. Thus, we expect it to be soluble in water. As both water and methanol are liquids, the word *miscible* can be used in place of *soluble*.
b. Sodium sulfate is an ionic compound, so we expect it to be soluble in water.
c. Like other hydrocarbons, octane is nonpolar, so we expect that it would not be soluble in water.

? EXERCISE 5.2.3: TOLUENE

Toluene ($C_6H_5CH_3$) is widely used in industry as a nonpolar solvent. Which substances should dissolve in toluene?

a. water (H_2O)
b. sodium sulfate (Na_2SO_4)
c. octane (C_8H_{18})

Answer

Octane (C_8H_{18}) will dissolve. It is also non-polar.

SUMMARY

- Solutions are composed of a solvent (major component) and a solute (minor component).
- “Like dissolves like” is a useful rule for deciding if a solute will be soluble in a solvent.

5.2: Solutions - Homogeneous Mixtures is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **13.2: Solutions - Homogeneous Mixtures** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-NC-SA 3.0.

5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy

Learning Objectives

- Define electrolytes and non electrolytes
- Explain why solutions form.
- Discuss the idea of water as the "universal solvent".
- Explain how water molecules attract ionic solids when they dissolve in water.

We have learned that solutions can be formed in a variety of combinations using solids, liquids, and gases. We also know that solutions have constant composition, and that this composition can be varied up to a point to maintain the homogeneous nature of the solution. But how exactly do solutions form? Why is it that oil and water will not form a solution, and yet vinegar and water will? Why could we dissolve table salt in water, but not in vegetable oil? The reasons why solutions will form will be explored in this section, along with a discussion of why water is used most frequently to dissolve substances of various types.

Solubility and Saturation

Table salt (**NaCl**) readily dissolves in water. In most cases, only a certain maximum amount of solute can be dissolved in a given amount of solvent. This maximum amount is specified as the **solubility** of the solute. It is usually expressed in terms of the amount of solute that can dissolve in 100 g of the solvent at a given temperature. Table **5.3.1** lists the solubilities of some simple ionic compounds. These solubilities vary widely. NaCl can dissolve up to 31.6 g per 100 g of H_2O, while AgCl can dissolve only 0.00019 g per 100 g of H_2O.

Table **5.3.1**: Solubilities of Some Ionic Compounds

Solute	Solubility (g per 100 g of H_2O at 25°C)
AgCl	0.00019
$CaCO_3$	0.0006
KBr	70.7
NaCl	36.1
$NaNO_3$	94.6

When the maximum amount of solute has been dissolved in a given amount of solvent, we say that the solution is **saturated** with solute. When less than the maximum amount of solute is dissolved in a given amount of solute, the solution is **unsaturated**. These terms are also qualitative terms because each solute has its own solubility. A solution of 0.00019 g of AgCl per 100 g of H_2O may be saturated, but with so little solute dissolved, it is also rather dilute. A solution of 36.1 g of NaCl in 100 g of H_2O is also saturated, but rather concentrated. In some circumstances, it is possible to dissolve more than the maximum amount of a solute in a solution. Usually, this happens by heating the solvent, dissolving more solute than would normally dissolve at regular temperatures, and letting the solution cool down slowly and carefully. Such solutions are called **supersaturated** solutions and are not stable; given an opportunity (such as dropping a crystal of solute in the solution), the excess solute will precipitate from the solution.The figure below illustrates the above process and shows the distinction between unsaturated and saturated.




Figure 5.3.1: When 30.0 g of NaCl is added to 100 mL, it all dissolves, forming an unsaturated solution. When 40.0 g is added, 36.0 g dissolves and 4.0 g remains undissolved, forming a saturated solution.

How can you tell if a solution is saturated or unsaturated? If more solute is added and it does not dissolve, then the original solution was saturated. If the added solute dissolves, then the original solution was unsaturated. A solution that has been allowed to reach equilibrium, but which has extra undissolved solute at the bottom of the container, must be saturated.



Electrolyte Solutions: Dissolved Ionic Solids

When some substances are dissolved in water, they undergo either a physical or a chemical change that yields ions in solution. These substances constitute an important class of compounds called **electrolytes**. Substances that do not yield ions when dissolved are called **nonelectrolytes**. If the physical or chemical process that generates the ions is essentially 100% efficient (all of the dissolved compound yields ions), then the substance is known as a strong electrolyte (good conductor). If only a relatively small fraction of the dissolved substance undergoes the ion-producing process, the substance is a weak electrolyte (does not conduct electricity as well).

Substances may be identified as strong, weak, or nonelectrolytes by measuring the electrical conductance of an aqueous solution containing the substance. To conduct electricity, a substance must contain freely mobile, charged species. Most familiar is the conduction of electricity through metallic wires, in which case the mobile, charged entities are electrons. Solutions may also conduct electricity if they contain dissolved ions, with conductivity increasing as ion concentration increases. Applying a voltage to electrodes immersed in a solution permits assessment of the relative concentration of dissolved ions, either quantitatively, by measuring the electrical current flow, or qualitatively, by observing the brightness of a light bulb included in the circuit (Figure 5.3.1).




Figure **5.3.1**: Solutions of nonelectrolytes, such as ethanol, do not contain dissolved ions and cannot conduct electricity. Solutions of electrolytes contain ions that permit the passage of electricity. The conductivity of an electrolyte solution is related to the strength of the electrolyte. This diagram shows three separate beakers. Each has a wire plugged into a wall outlet. In each case, the wire leads from the wall to the beaker and is split resulting in two ends. One end leads to a light bulb and continues on to a rectangle labeled with a plus sign. The other end leads to a rectangle labeled with a minus sign. The rectangles are in a solution. In the first beaker, labeled “Ethanol No Conductivity,” four pairs of linked small green spheres suspended in the solution between the rectangles. In the second beaker, labeled “K C l Strong Conductivity,” six individual green spheres, three labeled plus and three labeled minus are suspended in the solution. Each of the six spheres has an arrow extending from it pointing to the rectangle labeled with the opposite sign. In the third beaker, labeled “Acetic acid solution Weak conductivity,” two pairs of joined green spheres and two individual spheres, one labeled plus and one labeled minus are shown suspended between the two rectangles. The plus labeled sphere has an arrow pointing to the rectangle labeled minus and the minus labeled sphere has an arrow pointing to the rectangle labeled plus.

Water and other polar molecules are attracted to ions, as shown in Figure **5.3.2**. The electrostatic attraction between an ion and a molecule with a dipole is called an ion-dipole attraction. These attractions play an important role in the dissolution of ionic compounds in water.




Figure **5.3.2**: As potassium chloride (KCl) dissolves in water, the ions are hydrated. The polar water molecules are attracted by the charges on the K^+ and Cl^- ions. Water molecules in front of and behind the ions are not shown. The diagram shows eight purple spheres labeled K superscript plus and eight green spheres labeled C l superscript minus mixed and touching near the center of the diagram. Outside of this cluster of spheres are seventeen clusters of three spheres, which include one red and two white spheres. A red sphere in one of these clusters is labeled O. A white sphere is labeled H. Two of the green C l superscript minus spheres are surrounded by three of the red and white clusters, with the red spheres closer to the green spheres than the white spheres. One of the K superscript plus purple spheres is surrounded by four of the red and white clusters. The white spheres of these clusters are closest to the purple spheres.

When ionic compounds dissolve in water, the ions in the solid separate and disperse uniformly throughout the solution because water molecules surround and solvate the ions, reducing the strong electrostatic forces between them. This process represents a physical change known as dissociation. Under most conditions, ionic compounds will dissociate nearly completely when dissolved, and so they are classified as strong electrolytes.

Example 5.3.1: Identifying Ionic Compounds

Which compound(s) will dissolve in solution to separate into ions?

a. **LiF**
b. **P_2F_5**
c. **C_2H_5OH**

Solution

LiF will separate into ions when dissolved in solution, because it is an ionic compound. **P_2F_5** and **C_2H_5OH** are both covalent and will stay as molecules in a solution.

Exercise 5.3.1

Which compounds will dissolve in solution to separate into ions?

a. $C_6H_{12}O_{11}$, glucose
b. CCl_4
c. $CaCl_2$
d. $AgNO_3$

Answer

c & d

How Temperature Influences Solubility

The **solubility** of a substance is the amount of that substance that is required to form a saturated solution in a given amount of solvent at a specified temperature. Solubility is often measured as the grams of solute per **100 g** of solvent. The solubility of sodium chloride in water is **36.0 g** per **100 g** water at **20°C**. The temperature must be specified because solubility varies with temperature. For gases, the pressure must also be specified. Solubility is specific for a particular solvent. We will consider solubility of material in water as solvent.

The solubility of the majority of solid substances increases as the temperature increases. However, the effect is difficult to predict and varies widely from one solute to another. The temperature dependence of solubility can be visualized with the help of a **solubility curve**, a graph of the solubility vs. temperature (Figure 5.3.4).




Figure 5.3.4: Solubility curves for several compounds.

Notice how the temperature dependence of $NaCl$ is fairly flat, meaning that an increase in temperature has relatively little effect on the solubility of $NaCl$. The curve for KNO_3, on the other hand, is very steep and so an increase in temperature dramatically increases the solubility of KNO_3.

Several substances—HCl, NH_3, and SO_2—have solubility that decreases as temperature increases. They are all gases at standard pressure. When a solvent with a gas dissolved in it is heated, the kinetic energy of both the solvent and solute increase. As the kinetic energy of the gaseous solute increases, its molecules have a greater tendency to escape the attraction of the solvent molecules and return to the gas phase. Therefore, the solubility of a gas decreases as the temperature increases.

Solubility curves can be used to determine if a given solution is saturated or unsaturated. Suppose that $80\ g$ of KNO_3 is added to $100\ g$ of water at $30°C$. According to the solubility curve, approximately $48\ g$ of KNO_3 will dissolve at $30°C$. This means that the solution will be saturated since $48\ g$ is less than $80\ g$. We can also determine that there will be $80 - 48 = 32\ g$ of undissolved KNO_3 remaining at the bottom of the container. Now suppose that this saturated solution is heated to $60°C$. According to the curve, the solubility of KNO_3 at $60°C$ is about $107\ g$. Now the solution is unsaturated since it contains only the original $80\ g$ of dissolved solute. Now suppose the solution is cooled all the way down to $0°C$. The solubility at $0°C$ is about $14\ g$, meaning that $80 - 14 = 66\ g$ of the KNO_3 will re-crystallize.

Summary

- Solubility is the specific amount of solute that can dissolve in a given amount of solvent.
- Saturated and unsaturated solutions are defined.
- Ionic compounds dissolve in polar solvents, especially water. This occurs when the positive cation from the ionic solid is attracted to the negative end of the water molecule (oxygen) and the negative anion of the ionic solid is attracted to the positive end of the water molecule (hydrogen).
- Water is considered the universal solvent since it can dissolve both ionic and polar solutes, as well as some nonpolar solutes (in very limited amounts).
- The solubility of a solid in water increases with an increase in temperature.

Vocabulary

- **Miscible** - Liquids that have the ability to dissolve in each other.
- **Immiscible** - Liquids that do not have the ability to dissolve in each other.
- **Electrostatic attraction** - The attraction of oppositely charged particles.

5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **13.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.4: Specifying Solution Concentration- Mass Percent

Learning Objectives

- Express the amount of solute in a solution in various concentration units.

To define a solution precisely, we need to state its concentration: how much solute is dissolved in a certain amount of solvent. Words such as *dilute* or *concentrated* are used to describe solutions that have a little or a lot of dissolved solute, respectively, but these are relative terms with meanings that depend on various factors.

Introduction

Concentration is the measure of how much of a given substance is mixed with another substance. Solutions are said to be either dilute or concentrated. When we say that vinegar is 5% acetic acid in water, we are giving the concentration. If we said the mixture was 10% acetic acid, this would be more concentrated than the vinegar solution.




Figure 5.4.1: The solution on the left is more concentrated than the solution on the right because there is a greater ratio of solute (red balls) to solvent (blue balls) particles. The solution particles are closer together. The solution on the right is more dilute (less concentrated). (CC-SA-BY-3.0 Tracy Poulsen).

A **concentrated** solution is one in which there is a large amount of solute in a given amount of solvent. A **dilute** solution is one in which there is a small amount of solute in a given amount of solvent. A dilute solution is a concentrated solution that has been, in essence, watered down. Think of the frozen juice containers you buy in the grocery store. To make juice, you have to mix the frozen juice concentrate from inside these containers with three or four times the container size full of water. Therefore, you are diluting the concentrated juice. In terms of solute and solvent, the concentrated solution has a lot of solute versus the dilute solution that would have a smaller amount of solute.

The terms "concentrated" and "dilute" provide qualitative methods of describing concentration. Although qualitative observations are necessary and have their place in every part of science, including chemistry, we have seen throughout our study of science that there is a definite need for quantitative measurements in science. This is particularly true in solution chemistry. In this section, we will explore some quantitative methods of expressing solution concentration.

Mass Percent

There are several ways of expressing the concentration of a solution by using a percentage. The mass/mass percent (% m/m) is defined as the mass of a solute divided by the mass of a solution times 100:

$$\%\,\text{m/m} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 100\%$$

mass of solution = mass of solute + mass solvent

If you can measure the masses of the solute and the solution, determining the mass/mass percent is easy. Each mass must be expressed in the same units to determine the proper concentration.

Suppose that a solution was prepared by dissolving $25.0\ \text{g}$ of sugar into $100.0\ \text{g}$ of water.

The mass of the solution is

mass of solution = 25.0g sugar + 100.0g water = 125.0 g

The percent by mass would be calculated by:

$$\text{Percent by mass} = \frac{25.0\ \text{g sugar}}{125.0\ \text{g solution}} \times 100\% = 20.0\%\ \text{sugar}$$

Example 5.4.1

A saline solution with a mass of 355 g has 36.5 g of NaCl dissolved in it. What is the mass/mass percent concentration of the solution?

Solution

We can substitute the quantities given in the equation for mass/mass percent:

$$\%\,\mathrm{m/m} = \frac{36.5\ \mathrm{g}}{355\ \mathrm{g}} \times 100\% = 10.3\%$$

Exercise 5.4.1

A dextrose (also called D-glucose, $C_6H_{12}O_6$) solution with a mass of 2.00×10^2 g has 15.8 g of dextrose dissolved in it. What is the mass/mass percent concentration of the solution?

Answer

7.90 %

Using Mass Percent in Calculations

Sometimes you may want to make up a particular mass of solution of a given percent by mass and need to calculate what mass of the solute to use. Using mass percent as a conversion can be useful in this type of problem. The mass percent can be expressed as a conversion factor in the form $\frac{g\ \text{solute}}{100\ \text{gsolution}}$ or $\frac{100\ \text{gsolution}}{g\ \text{solute}}$

For example, if you need to make $3000.0\ \mathrm{g}$ of a 5.00% solution of sodium chloride, the mass of solute needs to be determined.

Solution


Given: 3000.0 g NaCl solution

5.00% NaCl solution

Find: mass of solute = ? g NaCl

Other known quantities: 5.00 g NaCl is to 100 g solution

The appropriate conversion factor (based on the given mass percent) can be used follows:



$$\frac{5.00\ g\ NaCl}{100\ g\ NaCl\ solution}$$

To solve for the mass of NaCl, the given mass of solution is multiplied by the conversion factor.

$$gNaCl = 3,000.0\ \cancel{g\ NaCl\ solution} \times \frac{5.00\ g\ NaCl}{100\ \cancel{g\ NaCl\ solution}} = 150.0 g\ NaCl$$

You would need to weigh out $150\ \mathrm{g}$ of NaCl and add it to $2850\ \mathrm{g}$ of water. Notice that it was necessary to subtract the mass of the NaCl $(150\ \mathrm{g})$ from the mass of solution $(3000\ \mathrm{g})$ to calculate the mass of the water that would need to be added.

Exercise 5.4.1

What is the amount (in g) of hydrogen peroxide (H_2O_2) needed to make a 6.00 kg, 3.00 % (by mass) H_2O_2 solution?

Answer

180 g H_2O_2

Contributors and Attributions

5.4: Specifying Solution Concentration- Mass Percent is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **13.5: Specifying Solution Concentration- Mass Percent** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.5: Specifying Solution Concentration- Molarity

Learning Objectives

- Use molarity to determine quantities in chemical reactions.
- Use molarity as a conversion factor in calculations.

Another way of expressing concentration is to give the number of moles of solute per unit volume of solution. Of all the quantitative measures of concentration, molarity is the one used most frequently by chemists. **Molarity** is defined as the number of moles of solute per liter of solution.

$$\text{molarity} = \frac{\text{number of moles of solute}}{\text{number of liters of solution}} \tag{5.5.1}$$

The symbol for molarity is **M** or moles/liter. Chemists also use square brackets to indicate a reference to the molarity of a substance. For example, the expression $[Ag^+]$ refers to the molarity of the silver ion in solution. Solution concentrations expressed in molarity are the easiest to perform calculations with, but the most difficult to make in the lab. Such concentration units are useful for discussing chemical reactions in which a solute is a product or a reactant. Molar mass can then be used as a conversion factor to convert amounts in moles to amounts in grams.

It is important to remember that “mol” in this expression refers to moles of solute and that “L” refers to liters of solution. For example, if you have 1.5 mol of NaCl dissolved in 0.500 L of solution, its molarity is

$$\frac{1.5 \text{ mol NaCl}}{0.500 \text{ L solution}} = 3.0 \text{ M NaCl}$$

Sometimes (aq) is added when the solvent is water, as in “3.0 M NaCl (aq).” This is read as "a 3.00 ***molar*** sodium chloride solution," meaning that there are 3.00 moles of NaOH dissolved per one liter of solution.

> *Be sure to note that molarity is calculated as the total volume of the **entire** solution, not just volume of solvent! The solute contributes to total volume.*

If the quantity of the solute is given in mass units, you must convert mass units to mole units before using the definition of molarity to calculate concentration. For example, what is the molar concentration of a solution of 22.4 g of HCl dissolved in 1.56 L?

Step 1: First, convert the mass of solute to moles using the molar mass of HCl (36.5 g/mol):

$$22.4\,\cancel{g HCl} \times \frac{1\,mol\,HCl}{36.5\,\cancel{g HCl}} = 0.614\,mol\,HCl$$

Step 2: Now we can use the definition of molarity to determine a concentration:

$$M = \frac{0.614\,mol\,HCl}{1.56 L\,solution} = 0.394\,M HCl$$


Before a molarity concentration can be calculated, the amount of the solute must be expressed in moles, and the volume of the solution must be expressed in liters, as demonstrated in the following example.

Example 5.5.1

A solution is prepared by dissolving **42.23 g** of NH_4Cl into enough water to make **500.0 mL** of solution. Calculate its molarity.

Solution

Solutions to Example 13.6.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: Mass $= 42.23\ \text{g}\ NH_4Cl$ Volume solution $= 500.0\ \text{mL} = 0.5000\ \text{L}$ Find: Molarity = ? M
List other known quantities.	Molar mass $NH_4Cl = 53.50\ \text{g/mol}$
Plan the problem.	1. The mass of the ammonium chloride is first converted to moles.  $\frac{1\ mol\ NH_4Cl}{53.50g\ NH_4Cl}$ 2. Then the molarity is calculated by dividing by liters. Note the given volume has been converted to liters. $\text{M} = \frac{\text{mol } NH_4Cl}{\text{L solution}}$
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $42.23\ \cancel{\text{g } NH_4Cl} \times \frac{1\ \text{mol } NH_4Cl}{53.50\ \cancel{\text{g } NH_4Cl}} = 0.7893\ \text{mol } NH_4Cl$ $\frac{0.7893\ \text{mol } NH_4Cl}{0.5000\ \text{L solution}} = 1.579\ \text{M}$
Think about your result.	The molarity is $1.579\ \text{M}$, meaning that a liter of the solution would contain $1.579\ \text{mol } NH_4Cl$. Four significant figures are appropriate.

? Exercise 5.5.1*A*

What is the molarity of a solution made when 66.2 g of $C_6H_{12}O_6$ are dissolved to make 235 mL of solution?

Answer

1.57 M $C_6H_{12}O_6$

? Exercise 5.5.1*B*

What is the concentration, in mol/L, where $137\ \text{g}$ of $NaCl$ has been dissolved in enough water to make $500\ \text{mL}$ of solution?

Answer

4.69 M NaCl

Using Molarity in Calculations


Concentration can be a conversion factor between the amount of solute and the amount of solution or solvent (depending on the definition of the concentration unit). As such, concentrations can be useful in a variety of stoichiometry problems. In many cases, it is best to use the original definition of the concentration unit; it is that definition that provides the conversion factor.

A simple example of using a concentration unit as a conversion factor is one in which we use the definition of the concentration unit and rearrange; we can do the calculation again as a unit conversion, rather than as a definition.

Example 5.5.2: Determining Moles of Solute, Given the Concentration and Volume of a Solution

For example, suppose we ask how many moles of solute are present in 0.108 L of a 0.887 M NaCl solution. Because 0.887 M means 0.887 mol/L, we can use this second expression for the concentration as a conversion factor:

Solution



$$0.108 \cancel{L\ solution} \times \frac{0.887\ mol\ NaCl}{1 \cancel{L\ solution}} = 0.0958\ mol\ NaCl$$

If we used the definition approach, we get the same answer, but now we are using conversion factor skills.


Like any other conversion factor that relates two different types of units, the reciprocal of the concentration can be also used as a conversion factor.

Example 5.5.3: Determining Volume of a Solution, Given the Concentration and Moles of Solute

Using concentration as a conversion factor, how many liters of 2.35 M $CuSO_4$ are needed to obtain 4.88 mol of $CuSO_4$?

Solution

This is a one-step conversion, but the concentration must be written as the reciprocal for the units to work out:



$$4.88 \cancel{mol\ CuSO_4} \times \frac{1\ L\ solution}{2.35 \cancel{mol\ CuSO_4}} = 2.08\ L\ of\ solution$$


In a laboratory situation, a chemist must frequently prepare a given volume of solutions of a known molarity. The task is to calculate the mass of the solute that is necessary. The molarity equation can be rearranged to solve for moles, which can then be converted to grams. The following example illustrates this.

Example 5.5.4

A chemist needs to prepare **3.00 L** of a **0.250 M** solution of potassium permanganate (**$KMnO_4$**). What mass of **$KMnO_4$** does she need to make the solution?

Solution

Solutions to Example 13.6.4

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: Molarity $= 0.250\ \mathrm{M}$ Volume $= 3.00\ \mathrm{L}$ Find: Mass $\mathrm{KMnO_4} = ?\ \mathrm{g}$
List other known quantities.	Molar mass $\mathrm{KMnO_4} = 158.04\ \mathrm{g/mol}$ 0.250 mol $KMnO_4$ to 1 L of $KMnO_4$ solution
Plan the problem.	
Cancel units and calculate.	Now substitute the known quantities into the equation and solve. $\mathrm{mol\ KMnO_4} = 0.250\ \mathrm{M\ KMnO_4} \times 3.00\ \mathrm{L} = 0.750\ \mathrm{mol\ KMnO_4}$ $3.00\ \cancel{L\ solution} \times \frac{0.250\ \cancel{\mathrm{mol\ KMnO_4}}}{1\ \cancel{L\ solution}} \times \frac{158.04\ \mathrm{g\ KMnO_4}}{1\ \cancel{\mathrm{mol\ KMnO_4}}} = 119\ \mathrm{g\ KMnO_4}$
Think about your result.	When $119\ \mathrm{g}$ of potassium permanganate is dissolved into water to make $3.00\ \mathrm{L}$ of solution, the molarity is $0.250\ \mathrm{M}$.

Exercise 5.5.4A

Using concentration as a conversion factor, how many liters of 0.0444 M CH_2O are needed to obtain 0.0773 mol of CH_2O?

Answer

1.74 L

Exercise 5.5.4B

Answer the problems below using concentration as a conversion factor.

a. What mass of solute is present in 1.08 L of 0.0578 M H_2SO_4?
b. What volume of 1.50 M HCl solution contains 10.0 g of hydrogen chloride?

Answer a

6.12 g

Answer b

183 mL or 0.183L

How to Indicate Concentration

- Square brackets are often used to represent concentration, e.g., [NaOH] = 0.50 M.
- Use the capital letter M for molarity, not a lower case m (this is a different concentration unit called *molality*).



Watch as the Flinn Scientific Tech Staff demonstrates "How To Prepare Solutions."

It is important to note that there are many different ways you can set up and solve your chemistry equations. Some students prefer to answer multi-step calculations in one long step, while others prefer to work out each step individually. Neither method is necessarily better or worse than the other method—whichever makes the most sense to ***you*** is the one that you should use. In this text, we will typically use unit analysis (also called dimension analysis or factor analysis).

Contributors and Attributions

- Peggy Lawson (Oxbow Prairie Heights School). Funded by Saskatchewan Educational Technology Consortium.

5.5: Specifying Solution Concentration- Molarity is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.


- **13.6: Specifying Solution Concentration- Molarity** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.6: SOLUTION STOICHIOMETRY

LEARNING OBJECTIVES

- Determine amounts of reactants or products in aqueous solutions.

As we learned previously, double replacement reactions involve the reaction between ionic compounds in solution and, in the course of the reaction, the ions in the two reacting compounds are "switched" (they *replace* each other). Because these reactions occur in aqueous solution, we can use the concept of molarity to directly calculate the number of moles of reactants or products that will be formed, and therefore their amounts (i.e. volume of solutions or mass of precipitates).



As an example, lead (II) nitrate and sodium chloride react to form sodium nitrate and the *insoluble* compound, lead (II) chloride.

$$\mathrm{Pb(NO_3)_2\,(aq) + 2\,NaCl\,(aq) \longrightarrow PbCl_2\,(s) + 2\,NaNO_3\,(aq)} \tag{5.6.1}$$

In the reaction shown above, if we mixed 0.123 L of a 1.00 M solution of NaCl with 1.50 M solution of $\mathrm{Pb(NO_3)_2}$, we could calculate the volume of $\mathrm{Pb(NO_3)_2}$ solution needed to completely precipitate the $\mathrm{Pb^{2+}}$ ions.

The molar concentration can also be expressed as the following:

$$1.00\,M\,\mathrm{NaCl} = \frac{1.00\,mol\,\mathrm{NaCl}}{1\,L\,\mathrm{NaCl\ solution}}$$

and

$$1.50\,M\,\mathrm{Pb(NO_3)_2} = \frac{1.50\,mol\,\mathrm{Pb(NO_3)_2}}{1\,L\,\mathrm{Pb(NO_3)_2 solution}}$$

First, we must examine the reaction stoichiometry in the balanced reaction (Equation 5.6.1). In this reaction, one mole of $\mathrm{Pb(NO_3)_2}$ reacts with two moles of NaCl to give one mole of $\mathrm{PbCl_2}$ precipitate. Thus, the concept map utilizing the stoichiometric ratios is:



so the volume of lead (II) nitrate that reacted is calculated as:

$$0.123\,L\,\mathrm{NaCl\,solution} \times \frac{1.00\,mol\,\mathrm{NaCl}}{1\,L\,\mathrm{NaCl\ solution}} \times \frac{1\,mol\,\mathrm{Pb(NO_3)_2}}{2\,mol\,\mathrm{NaCl}} \times \frac{1\,L\,\mathrm{Pb(NO_3)_2\ solution}}{1.5\,mol\,\mathrm{Pb(NO_3)_2}}$$
$$= 0.041\,\mathrm{Pb(NO_3)_2}\,L\,\mathrm{solution}$$


This volume makes intuitive sense for two reasons: (1) the number of moles of $\mathrm{Pb(NO_3)_2}$ required is half of the number of moles of NaCl, based off of the stoichiometry in the balanced reaction (Equation 5.6.1); (2) the concentration of $\mathrm{Pb(NO_3)_2}$ solution is 50% greater than the NaCl solution, so less volume is needed.

EXAMPLE 5.6.1

What volume (in L) of 0.500 M sodium sulfate will react with 275 mL of 0.250 M barium chloride to completely precipitate all $\mathrm{Ba^{2+}}$ in the solution?

Solution

Solutions to Example 13.8.1

Steps for Problem Solving	Example 5.6.1
Identify the "given" information and what the problem is asking you to "find."	Given: 275 mL $BaCl_2$ 0.250 M $\mathbf{BaCl_2}$ or $\frac{0.250\ mol BaCl_2}{1\ L\ BaCl_2\ solution}$ 0.500 M $\mathbf{Na_2SO_4}$ or $\frac{0.500\ mol Na_2SO_4}{1\ L\ Na_2SO_4\ solution}$ Find: Volume $\mathbf{Na_2SO_4}$ solution.
Set up and balance the chemical equation.	$\mathbf{Na_2SO_4\,(aq) + BaCl_2\,(aq) \longrightarrow BaSO_4\,(s) + 2NaCl\,(aq)}$ An insoluble product is formed after the reaction.
List other known quantities.	1 mol of Na_2SO_4 to 1 mol $BaCl_2$ 1000 mL = 1 L
Prepare a concept map and use the proper conversion factor.	
Cancel units and calculate.	$275\ mL\ BaCl_2\ solution \times \frac{1\ L}{1000\ mL} \times \frac{0.250\ mol\ BaCl_2}{1\ L\ BaCl_2\ solution} \times \frac{1\ mol\ Na_2SO_4}{1\ mol\ BaCl_2} \times \frac{1\ L\ Na_2SO_4\ solution}{0.500\ mol\ Na_2SO_4} = 0.1375$ L sodium sulfate
Think about your result.	The lesser amount (almost half) of sodium sulfate is to be expected as it is more concentrated than barium chloride. Also, the units are correct.

? EXERCISE 5.6.1

What volume of 0.250 M lithium hydroxide will completely react with 0.500 L of 0.250 M of sulfuric acid solution?

Answer

0.250 L $\mathbf{LiOH}$ solution

This page titled 5.6: Solution Stoichiometry is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew, Henry Agnew, Paul Young, & Paul Young (ChemistryOnline.com) .

- **13.8: Solution Stoichiometry** by Henry Agnew, Marisa Alviar-Agnew, Paul Young is licensed CC BY-SA 4.0.

5.7: Sour Patch Kids and International Spy Movies

Sour Patch Kids are a soft candy with a coating of invert sugar and sour sugar. The candy's slogan, "Sour. Sweet. Gone.", refers to its sour-to-sweet taste.



Figure **5.7.1**: Sour Patch Kids candies. Image courtesy of Evan-Amos (public domain).

Sour sugar is a food ingredient that is used to impart a sour flavor, made from citric or tartaric acid and sugar. It is used to coat sour candies like Sour Patch Kids. Eating large amounts of sour sugar can cause irritation of the tongue because of the acid. It can also cause irreversible dental erosion.

This page titled 5.7: Sour Patch Kids and International Spy Movies is shared under a CC BY-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew via source content that was edited to the style and standards of the LibreTexts platform.

- **14.1: Sour Patch Kids and International Spy Movies** by Henry Agnew, Marisa Alviar-Agnew is licensed CC BY-SA 4.0. Original source: https://www.ck12.org/c/chemistry/.

5.8: Acids- Properties and Examples

Learning Objectives

- Examine properties of acids.

Many people enjoy drinking coffee. A cup first thing in the morning helps start the day. But keeping the coffee maker clean can be a problem. Lime deposits build up after a while and slow down the brewing process. The best cure for this is to put vinegar (dilute acetic acid) in the pot and run it through the brewing cycle. The vinegar dissolves the deposits and cleans the maker, which will speed up the brewing process back to its original rate. Just be sure to run water through the brewing process after the vinegar, or you will get some really horrible coffee.

Acids

Acids are very common in some of the foods that we eat. Citrus fruits such as oranges and lemons contain citric acid and ascorbic acid, which is better known as vitamin C. Carbonated sodas contain phosphoric acid. Vinegar contains acetic acid. Your own stomach utilizes hydrochloric acid to digest food. Acids are a distinct class of compounds because of the properties of their aqueous solutions as outlined below:

1. Aqueous solutions of acids are electrolytes, meaning that they conduct electrical current. Some acids are strong electrolytes because they ionize completely in water, yielding a great many ions. Other acids are weak electrolytes that exist primarily in a non-ionized form when dissolved in water.
2. Acids have a sour taste. Lemons, vinegar, and sour candies all contain acids.
3. Acids change the color of certain acid-base indicates. Two common indicators are litmus and phenolphthalein. Blue litmus turns red in the presence of an acid, while phenolphthalein turns colorless.
4. Acids react with active metals to yield hydrogen gas. Recall that an activity series is a list of metals in descending order of reactivity. Metals that are above hydrogen in the activity series will replace the hydrogen from an acid in a single-replacement reaction, as shown below:

$$\mathrm{Zn}\,(s) + \mathrm{H_2SO_4}\,(aq) \rightarrow \mathrm{ZnSO_4}\,(aq) + \mathrm{H_2}\,(g) \tag{5.8.1}$$

5. Acids react with bases to produce a salt compound and water. When equal moles of an acid and a base are combined, the acid is neutralized by the base. The products of this reaction are an ionic compound, which is labeled as a salt, and water.

It should not be hard for you to name several common acids (but you might find that listing bases is a little more difficult). Below is a partial list of some common acids, along with some chemical formulas:

Table **5.8.1**: Common Acids and Their Uses

Chemist Name	Common Name	Uses
hydrochloric acid, HCl	muriatic acid (used in pools) and stomach acid is HCl	Used in cleaning (refining) metals, in maintenance of swimming pools, and for household cleaning.
sulfuric acid, H_2SO_4		Used in car batteries, and in the manufacture of fertilizers.
nitric acid, HNO_3		Used in the manufacture of fertilizers, explosives and in extraction of gold.
acetic acid, $HC_2H_3O_2$	vinegar	Main ingredient in vinegar.
carbonic acid, H_2CO_3	responsible for the "fizz" in carbonated drinks	As an ingredient in carbonated drinks.
citric acid, $C_6H_8O_7$		Used in food and dietary supplements. Also added as an acidulant in creams, gels, liquids, and lotions.
acetylsalicylic acid, $C_6H_4(OCOCH_3)CO_2H$	aspirin	The active ingredient in aspirin.

What exactly makes an acid an acid, and what makes a base act as a base? Take a look at the formulas given in the above table and take a guess.

Hydrochloric Acid

Hydrochloric acid is a corrosive, strong mineral acid with many industrial uses. A colorless, highly pungent solution of hydrogen chloride (HCl) in water. Hydrochloric acid is usually prepared by treating $\mathbf{HCl}$ with water.

$$\mathbf{HCl\,(g) + H_2O\,(l) \longrightarrow H_3O^+\,(aq) + Cl^-\,(aq)}$$

Hydrochloric acid can therefore be used to prepare chloride salts. Hydrochloric acid is a strong acid, since it is completely dissociated in water. Hydrochloric acid is the preferred acid in titration for determining the amount of bases.

Sulfuric Acid

Sulfuric acid is a highly corrosive strong mineral acid with the molecular formula $\mathbf{H_2SO_4}$. Sulfuric acid is a diprotic acid and has a wide range of applications including use in domestic acidic drain cleaners,[] as an electrolyte in lead-acid batteries, and in various cleaning agents. It is also a central substance in the chemical industry.



Figure 5.8.1: Drops of concentrated sulfuric acid rapidly decompose a piece of cotton towel by dehydration. (CC BY-SA 3.0; Toxic Walker).

Because the hydration of sulfuric acid is thermodynamically favorable (and is highly exothermic) and the affinity of it for water is sufficiently strong, sulfuric acid is an excellent dehydrating agent. Concentrated sulfuric acid has a very powerful dehydrating

property, removing water (H_2O) from other compounds including sugar and other carbohydrates and producing carbon, heat, steam. Sulfuric acid behaves as a typical acid in its reaction with most metals by generating hydrogen gas (Equation 5.8.2).

$$M + H_2SO_4 \longrightarrow M(SO_4) + H_2 \tag{5.8.2}$$

Nitric Acid

Nitric acid (HNO_3) is a highly corrosive mineral acid and is also commonly used as a strong oxidizing agent. Nitric acid is normally considered to be a strong acid at ambient temperatures. Nitric acid can be made by reacting nitrogen dioxide ($NO_2\,(g)$) with water.

$$3\,NO_2\,(g) + H_2O\,(l) \longrightarrow 2\,HNO_3\,(ag) + NO\,(g)$$

Nitric acid reacts with most metals, but the details depend on the concentration of the acid and the nature of the metal. Dilute nitric acid behaves as a typical acid in its reaction with most metals (e.g., nitric acid with magnesium, manganese or zinc will liberate H_2 gas):

$$Mg + 2\,HNO_3 \longrightarrow Mg(NO_3)_2 + H_2$$

$$Mn + 2\,HNO_3 \longrightarrow Mn(NO_3)_2 + H_2$$

$$Zn + 2\,HNO_3 \longrightarrow Zn(NO_3)_2 + H_2$$

Nitric acid is a corrosive acid and a powerful oxidizing agent. The major hazard it poses is chemical burn, as it carries out acid hydrolysis with proteins (amide) and fats (ester) which consequently decomposes living tissue (Figure 5.8.2). Concentrated nitric acid stains human skin yellow due to its reaction with the keratin



Figure 5.8.2: Second degree burn caused by nitric acid. (CC BY-SA 3.0; Alcamán).

Carbonic Acid

Carbonic acid is a chemical compound with the chemical formula H_2CO_3 and is also a name sometimes given to solutions of carbon dioxide in water (carbonated water), because such solutions contain small amounts of $H_2CO_3\,(aq)$. Carbonic acid, which is a weak acid, forms two kinds of salts: the carbonates and the bicarbonates. In geology, carbonic acid causes limestone to dissolve, producing calcium bicarbonate—which leads to many limestone features such as stalactites and stalagmites. Carbonic acid is a polyprotic acid, specifically it is diprotic, meaning that it has two protons which may dissociate from the parent molecule.

When carbon dioxide dissolves in water, it exists in chemical equilibrium (discussed in Chapter 15), producing carbonic acid:

$$CO_2 + H_2O \rightleftharpoons H_2CO_3$$

The reaction can be pushed to favor the reactants to generate $CO_2\,(g)$ from solution, which is key to the bubbles observed in carbonated beverages (Figure 5.8.3).



Figure 5.8.3: A glass of sparkling water. (CC BY-SA 3.0; Nevit Dilmen).

Formic Acid

Formic acid (HCO_2H) is the simplest carboxylic acid and is an important intermediate in chemical synthesis and occurs naturally, most notably in some ants. The word "formic" comes from the Latin word for ant, formica, referring to its early isolation by the distillation of ant bodies. Formic acid occurs widely in nature as its conjugate base formate.

Citric Acid

Citric acid ($C_6H_8O_7$) is a weak organic tricarboxylic acid that occurs naturally in citrus fruits. The citrate ion is an intermediate in the TCA cycle (Krebs cycle), a central metabolic pathway for animals, plants and bacteria. Because it is one of the stronger edible acids, the dominant use of citric acid is used as a flavoring and preservative in food and beverages, especially soft drinks.




Figure 5.8.4: Lemons, oranges, limes, and other citrus fruits possess high concentrations of citric acid (CC BY-SA 2.5; André Karwath).

Acetylsalicylic Acid

Acetylsalicylic acid (also known as aspirin) is a medication used to treat pain, fever, and inflammation. Aspirin, in the form of leaves from the willow tree, has been used for its health effects for at least 2,400 years.




Figure 5.8.5: Ball-and-stick model of the aspirin molecule. (Public Domain; Ben Mills).

Aspirin is a white, crystalline, weakly acidic substance.

Summary

A brief summary of key aspects of several acids commonly encountered by students was given. Acids are a distinct class of compounds because of the properties of their aqueous solutions.

Contributions & Attributions

-
- Peggy Lawson (Oxbow Prairie Heights School). Funded by Saskatchewan Educational Technology Consortium.

5.8: Acids- Properties and Examples is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.2: Acids- Properties and Examples** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.9: Bases- Properties and Examples

Learning Objectives

- Examine properties of bases.

Perhaps you have eaten too much pizza and felt very uncomfortable hours later. This feeling is due to excess stomach acid being produced. The discomfort can be dealt with by taking an antacid. The base in the antacid will react with the **HCl** in the stomach and neutralize it, taking care of that unpleasant feeling.

Bases

Bases have properties that mostly contrast with those of acids.

1. Aqueous solutions of bases are also electrolytes. Bases can be either strong or weak, just as acids can.
2. Bases often have a bitter taste and are found in foods less frequently than acids. Many bases, like soaps, are slippery to the touch.
3. Bases also change the color of indicators. Litmus turns blue in the presence of a base, while phenolphthalein turns pink.
4. Bases do not react with metals in the way that acids do.
5. Bases react with acids to produce a salt and water.



Figure **5.9.1**: Phenolphthalein indicator in presence of base.

Warning!

Tasting chemicals and touching them are NOT good lab practices and should be avoided—in other words—don't do this at home.

Bases are less common as foods, but they are nonetheless present in many household products. Many cleaners contain ammonia, a base. Sodium hydroxide is found in drain cleaner. **Antacids**, which combat excess stomach acid, are comprised of bases such as magnesium hydroxide or sodium hydrogen carbonate. Various common bases and corresponding uses are given in Table **5.9.2**.

Table **5.9.1**: Common Bases and Corresponding Uses

Some Common Bases	Uses
sodium hydroxide, NaOH (lye or caustic soda)	Used in the manufacture of soaps and detergents, and as the main ingredient in oven and drain cleaners.
potassium hydroxide, KOH (lye or caustic potash)	Used in the production of liquid soaps and soft soaps. Used in alkaline batteries.
magnesium hydroxide, $Mg(OH)_2$ (milk of magnesia)	Used as an ingredient in laxatives, antacids, and deodorants. Also used in the neutralization of acidic wastewater.
calcium hydroxide, $Ca(OH)_2$ (slaked lime)	Used in the manufacture of cement and lime water. Also, added to neutralize acidic soil.
aluminum hydroxide	Used in water purification and as an ingredient in antacids.
ammonia, NH_3	Used as a building block for the synthesis of many pharmaceutical products and in many commercial cleaning products. Used in the manufacture of fertilizers.

Sodium Hydroxide

Sodium hydroxide, also known as lye and caustic soda, is an inorganic compound with formula $\mathbf{NaOH}$. It is a white solid ionic compound consisting of sodium cations $\mathbf{Na^+}$ and hydroxide anions $\mathbf{OH^-}$.

Dissolution of solid sodium hydroxide in water is a highly exothermic reaction:

$$\mathbf{NaOH\,(s) \rightarrow Na^+\,(aq) + OH^-\,(aq)}$$

The resulting solution is usually colorless and odorless and feels slippery when it comes in contact with skin.



Figure 5.9.1: Sample of sodium hydroxide as pellets in a watch glass. (Public Domain; Walkerma.)

Potassium Hydroxide

Potassium hydroxide is an inorganic compound with the formula $\mathbf{KOH}$, and is commonly called caustic potash. Along with sodium hydroxide (NaOH), this colorless solid is a prototypical strong base. It has many industrial and niche applications, most of which exploit its corrosive nature and its reactivity toward acids. Its dissolution in water is strongly exothermic.

$$\mathbf{KOH\,(s) \rightarrow K^+\,(aq) + OH^-\,(aq)}$$

Concentrated aqueous solutions are sometimes called *potassium lyes*.

Magnesium Hydroxide

Magnesium hydroxide is the inorganic compound with the chemical formula $\mathbf{Mg(OH)_2}$. Magnesium hydroxide is a common component of antacids, such as milk of magnesia, as well as laxatives.



Figure 5.9.1: Bottle of Antacid tablets. (CC BY 2.,5; Midnightcomm).

It is a white solid with low solubility in water. Combining a solution of many magnesium salts with basic water induces precipitation of solid $\mathbf{Mg(OH)_2}$. However, a weak concentration of dissociated ions can be found in solution:

$$\mathrm{Mg(OH)_2\,(s) \rightleftharpoons Mg^{2+}\,(aq) + 2\,OH^-\,(aq)}$$

Calcium Hydroxide

Calcium hydroxide (traditionally called slaked lime) is an inorganic compound with the chemical formula $\mathbf{Ca(OH)_2}$. It is a colorless crystal or white powder. It has many names including hydrated lime, caustic lime, builders' lime, slaked lime, cal, or pickling lime. Calcium hydroxide is used in many applications, including food preparation. Limewater is the common name for a saturated solution of calcium hydroxide.

Calcium hydroxide is relatively insoluble in water, but is large enough that its solutions are basic according to the following reaction:

$$\mathrm{Ca(OH)_2\,(s) \rightleftharpoons Ca^{2+}\,(aq) + 2\,OH^-\,(aq)}$$

Ammonia

Ammonia is a compound of nitrogen and hydrogen with the formula $\mathbf{NH_3}$ and is a colorless gas with a characteristic pungent smell. It is the active product of “smelling salts,” and can quickly revive the faint of heart and light of head. Although common in nature and in wide use, ammonia is both caustic and hazardous in its concentrated form.




Figure 5.9.1: Ball-and-stick model of the ammonia molecule. (Public Domain; Ben Mills).

In aqueous solution, ammonia acts as a base, acquiring hydrogen ions from $\mathbf{H_2O}$ to yield ammonium and hydroxide ions:

$$\mathrm{NH_3\,(g) + H_2O\,(l) \rightleftharpoons NH_4^+\,(aq) + OH^-\,(aq)}$$

Ammonia is also a building block for the synthesis of many pharmaceutical products and is used in many commercial cleaning products.

Summary

- A brief summary of properties of bases was given.
- The properties of bases mostly contrast those of acids.
- Bases have many, varied uses.

5.9: Bases- Properties and Examples is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.3: Bases- Properties and Examples** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.10: Molecular Definitions of Acids and Bases

Learning Objectives

- Identify an Arrhenius acid and an Arrhenius base.
- Identify a Brønsted-Lowry acid and a Brønsted-Lowry base.
- Identify conjugate acid-base pairs in an acid-base reaction.

There are three major classifications of substances known as acids or bases. The theory developed by Svante Arrhenius in 1883, the Arrhenius definition, states that an acid produces H^+ in solution and a base produces OH^-. Later, two more sophisticated and general theories were proposed. These theories are the Brønsted-Lowry and Lewis definitions of acids and bases. This section will cover the Arrhenius and Brønsted-Lowry theories; the Lewis theory is discussed elsewhere.

The Arrhenius Theory of Acids and Bases

In 1884, the Swedish chemist Svante Arrhenius proposed two specific classifications of compounds, termed acids and bases. When dissolved in an aqueous solution, certain ions were released into the solution. An **Arrhenius acid** is a compound that **increases the concentration of H^+ ions** that are present when added to water. These H^+ ions form the hydronium ion (H_3O^+) when they combine with water molecules. This process is represented in a chemical equation by adding H_2O to the reactants side.

$$\mathrm{HCl\,(aq)} \rightarrow \mathrm{H^+\,(aq) + Cl^-\,(aq)}$$

In this reaction, hydrochloric acid (*HCl*) dissociates completely into hydrogen (H^+) and chlorine (Cl^-) ions when dissolved in water, thereby releasing H^+ ions into solution. Formation of the hydronium ion equation:

$$\mathrm{HCl\,(aq) + H_2O\,(l)} \rightarrow \mathrm{H_3O^+\,(aq) + Cl^-\,(aq)}$$

An **Arrhenius base** is a compound that **increases the concentration of OH^- ions** that are present when added to water. The dissociation is represented by the following equation:

$$\mathrm{NaOH\ \ (aq)} \rightarrow \mathrm{Na^+\ \ (aq) + OH^-\ \ (aq)}$$

In this reaction, sodium hydroxide (NaOH) disassociates into sodium (Na^+) and hydroxide (OH^-) ions when dissolved in water, thereby releasing OH^- ions into solution.

> *Arrhenius acids are substances which produce hydrogen ions in solution and Arrhenius bases are substances which produce hydroxide ions in solution.*

Limitations to the Arrhenius Theory

The Arrhenius theory has many more limitations than the other two theories. The theory does not explain the weak base ammonia (NH_3), which in the presence of water, releases hydroxide ions into solution, but does not contain OH- itself. The Arrhenius definition of acid and base is also limited to aqueous (i.e., water) solutions.

The Brønsted-Lowry Theory of Acids and Bases


In 1923, Danish chemist Johannes Brønsted and English chemist Thomas Lowry independently proposed new definitions for acids and bases, ones that focus on proton transfer. A **Brønsted-Lowry acid** is any species that can donate a proton (H^+) to another molecule. A **Brønsted-Lowry base** is any species that can accept a proton from another molecule. In short, **a Brønsted-Lowry acid** is a **proton donor (PD)**, while **a Brønsted-Lowry base** is a **proton acceptor (PA)**.

> *A Brønsted-Lowry acid is a proton donor, while a Brønsted-Lowry base is a proton acceptor.*

Let us use the reaction of ammonia in water to demonstrate the Brønsted-Lowry definitions of an acid and a base. Ammonia and water molecules are reactants, while the ammonium ion and the hydroxide ion are products:

$$NH_3\,(aq) + H_2O\,(\ell) \rightleftharpoons NH_4^+\,(aq) + OH^-\,(aq) \tag{5.10.1}$$

What has happened in this reaction is that the original water molecule has donated a hydrogen ion to the original ammonia molecule, which in turn has accepted the hydrogen ion. We can illustrate this as follows:




Because the water molecule donates a hydrogen ion to the ammonia, it is the Brønsted-Lowry acid, while the ammonia molecule—which accepts the hydrogen ion—is the Brønsted-Lowry base. Thus, ammonia acts as a base in both the Arrhenius sense and the Brønsted-Lowry sense.

Is an Arrhenius acid like hydrochloric acid still an acid in the Brønsted-Lowry sense? Yes, but it requires us to understand what really happens when HCl is dissolved in water. Recall that the hydrogen *atom* is a single proton surrounded by a single electron. To make the hydrogen *ion*, we remove the electron, leaving a bare proton. Do we *really* have bare protons floating around in aqueous solution? No, we do not. What really happens is that the H^+ ion attaches itself to H_2O to make H_3O^+, which is called the *hydronium ion*. For most purposes, H^+ and H_3O^+ represent the same species, but writing H_3O^+ instead of H^+ shows that we understand that there are no bare protons floating around in solution. Rather, these protons are actually attached to solvent molecules.

The Hydronium Ion


A proton in aqueous solution may be surrounded by more than one water molecule, leading to formulas like $H_5O_2^+$ or $H_9O_4^+$ rather than H_3O^+. It is simpler, however, to use H_3O^+ to represent the hydronium ion.



With this in mind, how do we define HCl as an acid in the Brønsted-Lowry sense? Consider what happens when HCl is dissolved in H_2O:

$$HCl\,(g) + H_2O\,(\ell) \rightarrow H_3O^+\,(aq) + Cl^-\,(aq) \tag{5.10.2}$$

We can depict this process using Lewis electron dot diagrams:



Now we see that a hydrogen ion is transferred from the HCl molecule to the H_2O molecule to make chloride ions and hydronium ions. As the hydrogen ion donor, HCl acts as a Brønsted-Lowry acid; as a hydrogen ion acceptor, H_2O is a Brønsted-Lowry base. So HCl is an acid not just in the Arrhenius sense, but also in the Brønsted-Lowry sense. Moreover, by the Brønsted-Lowry definitions, H_2O is a base in the formation of aqueous HCl. So the Brønsted-Lowry definitions of an acid and a base classify the dissolving of HCl in water as a reaction between an acid and a base—although the Arrhenius definition would not have labeled H_2O a base in this circumstance.

- A Brønsted-Lowry acid is a proton (hydrogen ion) donor.
- A Brønsted-Lowry base is a proton (hydrogen ion) acceptor.
- All Arrhenius acids and bases are Brønsted-Lowry acids and bases as well. However, not all Brønsted-Lowry acids and bases are Arrhenius acids and bases.

Example 5.10.1

Aniline ($C_6H_5NH_2$) is slightly soluble in water. It has a nitrogen atom that can accept a hydrogen ion from a water molecule, just like the nitrogen atom in ammonia does. Write the chemical equation for this reaction and identify the Brønsted-Lowry acid and base.

Solution

$C_6H_5NH_2$ and H_2O are the reactants. When $C_6H_5NH_2$ accepts a proton from H_2O, it gains an extra H and a positive charge and leaves an OH^- ion behind. The reaction is as follows:

$$\mathbf{C_6H_5NH_2\,(aq) + H_2O(\ell) \rightleftharpoons C_6H_5NH_3^+\,(aq) + OH^-\,(aq)}$$

Because $C_6H_5NH_2$ accepts a proton, it is the Brønsted-Lowry base. The H_2O molecule, because it donates a proton, is the Brønsted-Lowry acid.

Exercise 5.10.1

Identify the Brønsted-Lowry acid and the Brønsted-Lowry base in this chemical equation.

$$\mathbf{H_2PO_4^- + H_2O \rightleftharpoons HPO_4^{2-} + H_3O^+}$$

Answer

Brønsted-Lowry acid: $H_2PO_4^-$; Brønsted-Lowry base: H_2O

Exercise 5.10.2

Which of the following compounds is a Bronsted-Lowry base?

a. HCl
b. HPO_4^{2-}
c. H_3PO_4
d. NH_4^+
e. $CH_3NH_3^+$

Answer

A Brønsted-Lowry Base is a proton acceptor, which means it will take in an H^+. This eliminates $\mathbf{HCl}$, $\mathbf{H_3PO_4}$, $\mathbf{NH_4^+}$ and $\mathbf{CH_3NH_3^+}$ because they are Bronsted-Lowry acids. They all give away protons. In the case of $\mathbf{HPO_4^{2-}}$, consider the following equation:

$$\mathbf{HPO_4^{2-}\,(aq) + H_2O\,(l) \rightarrow PO_4^{3-}\,(aq) + H_3O^+\,(aq)}$$

Here, it is clear that HPO_4^{2-} is the acid since it donates a proton to water to make H_3O^+ and PO_4^{3-}. Now consider the following equation:


$$\mathbf{HPO_4^{2-}\,(aq) + H_2O\,(l) \rightarrow H_2PO_4^- + OH^-\,(aq)}$$

In this case, HPO_4^{2-} is the base since it accepts a proton from water to form $H_2PO_4^-$ and OH^-. Thus, HPO_4^{2-} is an acid and base together, making it amphoteric.

Since HPO_4^{2-} is the only compound from the options that can act as a base, the answer is **(b) $\mathbf{HPO_4^{2-}}$.**

Conjugate Acid-Base Pair

In reality, all acid-base reactions involve the transfer of protons between acids and bases. For example, consider the acid-base reaction that takes place when ammonia is dissolved in water. A water molecule (functioning as an acid) transfers a proton to an ammonia molecule (functioning as a base), yielding the conjugate base of water, $\mathbf{OH^-}$, and the conjugate acid of ammonia, $\mathbf{NH_4^+}$:



This figure has three parts in two rows. In the first row, two diagrams of acid-base pairs are shown. On the left, a space filling model of H subscript 2 O is shown with a red O atom at the center and two smaller white H atoms attached in a bent shape. Above this model is the label “H subscript 2 O (acid)” in purple. An arrow points right, which is labeled “Remove H superscript plus.” To the right is another space filling model with a single red O atom to which a single smaller white H atom is attached. The label in purple above this model reads, “O H superscript negative (conjugate base).” Above both of these red and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” To the right is a space filling model with a central blue N atom to which three smaller white H atoms are attached in a triangular pyramid arrangement. A label in green above reads “N H subscript 3 (base).” An arrow labeled “Add H superscript plus” points right. To the right of the arrow is another space filling model with a blue central N atom and four smaller white H atoms in a tetrahedral arrangement. The green label above reads “N H subscript 3 superscript plus (conjugate acid).” Above both of these blue and white models is an upward pointing bracket that is labeled “Conjugate acid-base pair.” The second row of the figure shows the chemical reaction, H subscript 2 O (l) is shown in purple, and is labeled below in purple as “acid,” plus N H subscript 3 (a q) in green, labeled below in green as “base,” followed by a double sided arrow arrow and O H superscript negative (a q) in purple, labeled in purple as “conjugate base,” plus N H subscript 4 superscript plus (a q)” in green, which is labeled in green as “conjugate acid.” The acid on the left side of the equation is connected to the conjugate base on the right with a purple line. Similarly, the base on the left is connected to the conjugate acid on the right side.

In the reaction of ammonia with water to give ammonium ions and hydroxide ions, ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The conjugate acid–base pairs for this reaction are $\boldsymbol{NH_4^+/NH_3}$ and $\boldsymbol{H_2O/OH^-}$.




Figure 5.10.1. The pairing of parent acids and bases with conjugate acids and bases.

https://chem.libretexts.org/@go/page/492204




Figure 5.10.1: The Relative Strengths of Some Common Conjugate Acid-Base Pairs. The strongest acids are at the bottom left, and the strongest bases are at the top right. The conjugate base of a strong acid is a very weak base, and, conversely, the conjugate acid of a strong base is a very weak acid.

Example 5.10.2

Identify the conjugate acid-base pairs in this equilibrium.


$$\mathrm{CH_3CO_2H + H_2O \rightleftharpoons H_3O^+ + CH_3CO_2^-}$$

Solution

Similarly, in the reaction of acetic acid with water, acetic acid **donates** a proton to water, which acts as the base. In the reverse reaction, $\mathrm{H_3O^+}$ is the acid that donates a proton to the acetate ion, which acts as the base.

Once again, we have two conjugate acid-base pairs:

- the parent acid and its conjugate base ($\mathrm{CH_3CO_2H/CH_3CO_2^-}$) and
- the parent base and its conjugate acid ($\mathrm{H_3O^+/H_2O}$).



Example 5.10.3

Identify the conjugate acid-base pairs in this equilibrium.

$$(CH_3)_3N + H_2O \rightleftharpoons (CH_3)_3NH^+ + OH^-$$

Solution

One pair is H_2O and OH^-, where H_2O has one more H^+ and is the conjugate acid, while OH^- has one less H^+ and is the conjugate base.

The other pair consists of $(CH_3)_3N$ and $(CH_3)_3NH^+$, where $(CH_3)_3NH^+$ is the conjugate acid (it has an additional proton) and $(CH_3)_3N$ is the conjugate base.

Exercise 5.10.3

Identify the conjugate acid-base pairs in this equilibrium.

$$NH_2^- + H_2O \rightleftharpoons NH_3 + OH^-$$

Answer

H_2O (acid) and OH^- (base); NH_2^- (base) and NH_3 (acid)

5.10: Molecular Definitions of Acids and Bases is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.4: Molecular Definitions of Acids and Bases** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.11: Reactions of Acids and Bases

Learning Objectives

- Write acid-base neutralization reactions.
- Write reactions of acids with metals.
- Write reactions of bases with metals.

Neutralization Reactions

The reaction that happens when an acid, such as **HCl**, is mixed with a base, such as **NaOH**:

$$\mathrm{HCl\,(aq) + NaOH\,(aq) \longrightarrow NaCl\,(aq) + H_2O\,(l)}$$

When an acid and a base are combined, water and a salt are the products. Salts are ionic compounds containing a positive ion other than $\mathrm{H^+}$ and a negative ion other than the hydroxide ion, $\mathrm{OH^-}$. Double displacement reactions of this type are called **neutralization reactions**. We can write an expanded version of this equation, with aqueous substances written in their longer form:

$$\mathrm{H^+\,(aq) + Cl^-\,(aq) + Na^+\,(aq) + OH^-\,(aq) \longrightarrow Na^+\,(aq) + Cl^-\,(aq) + H_2O\,(l)}$$

After removing the spectator ions, we get the net ionic equation:

$$\mathrm{H^+\,(aq) + OH^-\,(aq) \longrightarrow H_2O\,(l)}$$

When a strong acid and a strong base are combined in the proper amounts—when $[\mathrm{H^+}]$ equals $[\mathrm{OH^-}]$\)—a neutral solution results in which pH = 7. The acid and base have neutralized each other, and the acidic and basic properties are no longer present.

Salt solutions do not always have a pH of 7, however. Through a process known as **hydrolysis**, the ions produced when an acid and base combine may react with the water molecules to produce a solution that is slightly acidic or basic. As a general concept, if a strong acid is mixed with a weak base, the resulting solution will be slightly acidic. If a strong base is mixed with a weak acid, the solution will be slightly basic.



Video: Equimolar (~0.01 M) and equivolume solutions of **HCl** and **NaOH** are combined to make salt water. https://youtu.be/TS-I9KrUjB0

Example 5.11.1: Propionic Acid + Calcium Hydroxide

Calcium propionate is used to inhibit the growth of molds in foods, tobacco, and some medicines. Write a balanced chemical equation for the reaction of aqueous propionic acid ($\mathrm{CH_3CH_2CO_2H}$) with aqueous calcium hydroxide [$\mathrm{Ca(OH)_2}$].

Solution

Solutions to Example 14.5.1

Steps	Reaction
Write the unbalanced equation. This is a double displacement reaction, so the cations and anions swap to create the water and the salt.	$CH_3CH_2CO_2H(aq) + Ca(OH)_2(aq) \rightarrow (CH_3CH_2CO_2)_2Ca(aq) + H_2O(l)$
Balance the equation. Because there are two OH^- ions in the formula for $Ca(OH)_2$, we need two moles of propionic acid, $CH_3CH_2CO_2H$, to provide H^+ ions.	$\underline{2}CH_3CH_2CO_2H(aq) + Ca(OH)_2(aq) \rightarrow (CH_3CH_2CO_2)_2Ca(aq) + \underline{2}H_2O(l)$

? Exercise 5.11.1

Write a balanced chemical equation for the reaction of solid barium hydroxide with dilute acetic acid.

Answer

$$Ba(OH)_2\,(s) + 2\,CH_3CO_2H\,(aq) \longrightarrow Ba(CH_3CO_2)_2\,(aq) + 2\,H_2O\,(l)$$

Acids and Bases React with Metals

Acids react with most metals to form a salt and hydrogen gas. As discussed previously, metals that are more active than acids can undergo a **single displacement reaction**. For example, zinc metal reacts with hydrochloric acid, producing zinc chloride and hydrogen gas.

$$Zn\,(s) + 2\,HCl\,(aq) \longrightarrow ZnCl_2\,(aq) + H_2\,(g)$$

Bases also react with certain metals, like zinc or aluminum, to produce hydrogen gas. For example, sodium hydroxide reacts with zinc and water to form sodium zincate and hydrogen gas.

$$Zn\,(s) + 2\,NaOH\,(aq) + 2\,H_2O\,(l) \longrightarrow Na_2Zn(OH)_4\,(aq) + H_2\,(g).$$

5.11: Reactions of Acids and Bases is shared under a Public Domain license and was authored, remixed, and/or curated by Marisa Alviar-Agnew, Henry Agnew, Peggy Lawson, & Peggy Lawson.

- **14.5: Reactions of Acids and Bases** by Henry Agnew, Marisa Alviar-Agnew, Peggy Lawson is licensed Public Domain. Original source: https://sites.prairiesouth.ca/legacy/chemistry/chem30.

5.12: Acid–Base Titration

Learning Objectives

- Understand the basics of acid-base titrations.
- Understand the use of indicators.
- Perform a titration calculation correctly.

The reaction of an acid with a base to make a salt and water is a common reaction in the laboratory, partly because so many compounds can act as acids or bases. Another reason that acid-base reactions are so prevalent is because they are often used to determine quantitative amounts of one or the other. Performing chemical reactions quantitatively to determine the exact amount of a reagent is called a **titration**. A titration can be performed with almost any chemical reaction for which the balanced chemical equation is known. Here, we will consider titrations that involve acid-base reactions.

During an acid-base titration, an acid with a known concentration (a **standard solution**) is slowly added to a base with an unknown concentration (or vice versa). A few drops of indicator solution are added to the base. The indicator will signal, by color change, when the base has been neutralized (when $[H^+] = [OH^-]$). At that point—called the **equivalence point,** or **end point**—the titration is stopped. By knowing the volumes of acid and base used, and the concentration of the standard solution, calculations allow us to determine the concentration of the other solution.

It is important to accurately measure volumes when doing titrations. The instrument you would use is called a burette (or buret).



Figure 5.12.1: Equipment for Titrations. A burette is a type of liquid dispensing system that can accurately indicate the volume of liquid dispensed.

For example, suppose 25.66 mL (or 0.02566 L) of 0.1078 M HCl was used to titrate an unknown sample of NaOH. What mass of NaOH was in the sample? We can calculate the number of moles of HCl reacted:

$$\text{\# mol HCl} = (0.02566\ \text{L})(0.1078\ \text{M}) = 0.002766\ \text{mol HCl}$$

We also have the balanced chemical reaction between HCl and NaOH:

$$\mathrm{HCl + NaOH \longrightarrow NaCl + H_2O}$$

So we can construct a conversion factor to convert to number of moles of NaOH reacted:

$$0.002766\ \cancel{mol\ HCl} \times \frac{1\ mol\ NaOH}{1\ \cancel{mol\ HCl}} = 0.002766\ mol\ NaOH$$

Then we convert this amount to mass, using the molar mass of NaOH (40.00 g/mol):

$$0.002766\ \cancel{mol\ HCl} \times \frac{40.00\ g\ NaOH}{1\ \cancel{mol\ HCl}} = 0.1106\ g\ NaOH$$

This type of calculation is performed as part of a titration.

Example 5.12.1: Equivalence Point

What mass of $Ca(OH)_2$ is present in a sample if it is titrated to its equivalence point with 44.02 mL of 0.0885 M HNO_3? The balanced chemical equation is as follows:

$$\mathrm{2\,HNO_3 + Ca(OH)_2 \longrightarrow Ca(NO_3)_2 + 2\,H_2O}$$

Solution

In liters, the volume is 0.04402 L. We calculate the number of moles of titrant:

moles HNO_3 = (0.04402 L)(0.0885 M) = 0.00390 mol HNO_3

Using the balanced chemical equation, we can determine the number of moles of $Ca(OH)_2$ present in the analyte:

$$0.00390\ \cancel{mol\ HNO_3} \times \frac{1\ mol\ Ca(OH)_2}{2\ \cancel{mol\ HNO_3}} = 0.00195\ mol\ Ca(OH)_2$$

Then we convert this to a mass using the molar mass of $Ca(OH)_2$:

$$0.00195\ \cancel{mol\ Ca(OH)_2} \times \frac{74.1\ g\ Ca(OH)_2}{\cancel{mol\ Ca(OH)_2}} = 0.144\ g\ Ca(OH)_2$$

? Exercise 5.12.1

What mass of $H_2C_2O_4$ is present in a sample if it is titrated to its equivalence point with 18.09 mL of 0.2235 M NaOH? The balanced chemical reaction is as follows:

$$H_2C_2O_4 + 2\,NaOH \longrightarrow Na_2C_2O_4 + 2\,H_2O$$

Answer

0.182 g

? Exercise 5.12.2

If 25.00 mL of HCl solution with a concentration of 0.1234 M is neutralized by 23.45 mL of NaOH, what is the concentration of the base?

Answer

0.1316 M NaOH

? Exercise 5.12.3

A 20.0 mL solution of strontium hydroxide, $Sr(OH)_2$, is placed in a flask and a drop of indicator is added. The solution turns color after 25.0 mL of a standard 0.0500 M HCl solution is added. What was the original concentration of the $Sr(OH)_2$ solution?

Answer

$3.12 \times 10^{-2}\ M$ $Sr(OH)_2$

Indicator Selection for Titrations

The indicator used depends on the type of titration performed. The indicator of choice should change color when enough of one substance (acid or base) has been added to exactly use up the other substance. Only when a strong acid and a strong base are produced will the resulting solution be neutral. The three main types of acid-base titrations, and suggested indicators, are:

The three main types of acid-base titrations, suggested indicators, and explanations

Titration between . . .	Indicator	Explanation
strong acid and strong base	any	
strong acid and weak base	methyl orange	changes color in the acidic range (3.2 - 4.4)
weak acid and strong base	phenolphthalein	changes color in the basic range (8.2 - 10.6)

Summary

A titration is the quantitative reaction of an acid and a base. Indicators are used to show that all the analyte has reacted with the titrant.

Contributions & Attributions

- Peggy Lawson (Oxbow Prairie Heights School). Funded by Saskatchewan Educational Technology Consortium.

5.12: Acid–Base Titration is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.6: Acid–Base Titration** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.13: Strong and Weak Acids and Bases

Learning Objectives

- Define a strong and a weak acid and base.
- Recognize an acid or a base as strong or weak.
- Determine if a salt produces an acidic or a basic solution.

Strong and Weak Acids

Except for their names and formulas, so far we have treated all acids as equals, especially in a chemical reaction. However, acids can be very different in a very important way. Consider HCl(aq). When HCl is dissolved in H_2O, it completely dissociates into H^+(aq) and Cl^-(aq) ions; all the HCl molecules become ions:

$$HCl \xrightarrow{100\%} H^+(aq) + Cl^-(aq)$$

Any acid that dissociates 100% into ions is called a **strong acid**. If it does not dissociate 100%, it is a **weak acid**. $HC_2H_3O_2$ is an example of a weak acid:

$$HC_2H_3O_2 \xrightarrow{\sim 6\%} H^+(aq) + C_2H_3O_2^-(aq)$$

Because this reaction does not go 100% to completion, it is more appropriate to write it as **a reversible reaction**:

$$HC_2H_3O_2 \rightleftharpoons H^+(aq) + C_2H_3O_2^-(aq)$$

As it turns out, there are very few strong acids, which are given in Table **5.13.1**. If an acid is not listed here, it is a weak acid. It may be 1% ionized or 99% ionized, but it is still classified as a weak acid.

> *Any acid that dissociates 100% into ions is called a strong acid. If it does not dissociate 100%, it is a weak acid.*

Table **5.13.1**: Strong Acids and Bases

Acids	Bases
HCl	LiOH
HBr	NaOH
HI	KOH
HNO_3	RbOH
H_2SO_4	CsOH
$HClO_3$	$Mg(OH)_2$
$HClO_4$	$Ca(OH)_2$
	$Sr(OH)_2$
	$Ba(OH)_2$

Strong and Weak Bases

The issue is similar with bases: a **strong base** is a base that is 100% ionized in solution. If it is less than 100% ionized in solution, it is a **weak base**. There are very few strong bases (Table **5.13.1**); any base not listed is a weak base. All strong bases are OH^- compounds. So a base based on some other mechanism, such as NH_3 (which does not contain OH^- ions as part of its formula), will be a weak base.

Example 5.13.1: Identifying Strong and Weak Acids and Bases

Identify each acid or base as strong or weak.

a. HCl
b. $Mg(OH)_2$
c. C_5H_5N

Solution

a. Because HCl is listed in Table **5.13.1**, it is a strong acid.
b. Because $Mg(OH)_2$ is listed in Table **5.13.1**, it is a strong base.
c. The nitrogen in C_5H_5N would act as a proton acceptor and therefore can be considered a base, but because it does not contain an OH compound, it cannot be considered a strong base; it is a weak base.

Exercise 5.13.1

Identify each acid or base as strong or weak.

a. **RbOH**
b. **HNO_2**

Answer a

strong base

Answer b

weak acid

Example 5.13.2: Characterizing Base Ionization

Write the balanced chemical equation for the dissociation of $Ca(OH)_2$ and indicate whether it proceeds 100% to products or not.

Solution

This is an ionic compound of Ca^{2+} ions and OH^- ions. When an ionic compound dissolves, it separates into its constituent ions:

$$Ca(OH)_2 \longrightarrow Ca^{2+}(aq) + 2\,OH^-(aq)$$

Because $Ca(OH)_2$ is listed in Table **5.13.1**, this reaction proceeds 100% to products.

? Exercise 5.13.2

Write the balanced chemical equation for the dissociation of hydrazoic acid (HN_3) and indicate whether it proceeds 100% to products or not.

Answer a

The reaction is as follows:

$$HN_3 \longrightarrow H^+(aq) + N_3^-(aq)$$

It does not proceed 100% to products because hydrazoic acid is not a strong acid.

Key Takeaways

- Strong acids and bases are 100% ionized in aqueous solution.
- Weak acids and bases are less than 100% ionized in aqueous solution.
- Salts of weak acids or bases can affect the acidity or basicity of their aqueous solutions.

Contributions & Attributions

5.13: Strong and Weak Acids and Bases is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.7: Strong and Weak Acids and Bases** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.14: Water - Acid and Base in One

Learning Objectives

- Describe the autoionization of water.
- Calculate the concentrations of H_3O^+ and OH^- in aqueous solutions, knowing the other concentration.

We have already seen that H_2O can act as an acid or a base:

$$\underbrace{NH_3}_{\text{base}} + \underbrace{H_2O}_{\text{acid}} \rightleftharpoons NH_4^+ + OH^-$$

where H_2O acts as an acid (in red).

$$\underbrace{HCl}_{\text{acid}} + \underbrace{H_2O}_{\text{base}} \longrightarrow H_3O^+ + Cl^-$$

where H_2O acts as an base (in blue).

It may not surprise you to learn, then, that within any given sample of water, some H_2O molecules are acting as acids, and other H_2O molecules are acting as bases. The chemical equation is as follows:

$$\underbrace{H_2O}_{\text{acid}} + \underbrace{H_2O}_{\text{base}} \rightleftharpoons H_3O^+ + OH^- \tag{5.14.1}$$

This occurs only to a very small degree: only about 6 in 10^8 H_2O molecules are participating in this process, which is called the **autoionization of water**.




Figure 5.14.1: Autoionization of water, resulting in hydroxide and hydronium ions.

At this level, the concentration of both $H_3O^+(aq)$ and $OH^-(aq)$ in a sample of pure H_2O is about $1.0 \times 10^{-7}\ M$ (at room temperature). If we use square brackets—[ ]—around a dissolved species to imply the molar concentration of that species, we have

$$[H_3O^+]=[OH^-] = 1.0 \times 10^{-7} \tag{5.14.2}$$

for *any* sample of pure water because H_2O can act as both an acid and a base. The product of these two concentrations is 1.0×10^{-14}:

$$[H_3O^+] \times [OH^-] = (1.0 \times 10^{-7})(1.0 \times 10^{-7}) = 1.0 \times 10^{-14}$$

- For acids, the concentration of $H_3O^+(aq)$ (i.e., $[H_3O^+]$) is greater than $1.0 \times 10^{-7}\ M$.
- For bases the concentration of $OH^-(aq)$ (i.e., $[OH^-]$) is greater than $1.0 \times 10^{-7}\ M$.

However, the *product* of the two concentrations—$[H_3O^+][OH^-]$—is *always* equal to 1.0×10^{-14}, no matter whether the aqueous solution is an acid, a base, or neutral:

$$[H_3O^+][OH^-] = 1.0 \times 10^{-14}$$

This value of the product of concentrations is so important for aqueous solutions that it is called the **autoionization constant of water** and is denoted K_w:

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \tag{5.14.3}$$

This means that if you know $[H_3O^+]$ for a solution, you can calculate what $[OH^-]$ has to be for the product to equal 1.0×10^{-14}; or if you know $[OH^-]$, you can calculate $[H_3O^+]$. This also implies that as one concentration goes up, the other must go down to compensate so that their product always equals the value of K_w.

Warning: Temperature Matters

The degree of autoionization of water (Equation 5.14.1)—and hence the value of K_w—changes with temperature, so Equations 5.14.2 - 5.14.3 are accurate only at room temperature.

Example 5.14.1: Hydroxide Concentration

What is $[OH^-]$) of an aqueous solution if $[H_3O^+]$ is $1.0 \times 10^{-4} M$?

Solution

Solutions to Example 14.7.1

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: $[H_3O^+] = 1.0 \times 10^{-4}\ M$ Find: $[OH^-]$ = ? M
List other known quantities.	none
Plan the problem.	Using the expression for K_w, (Equation 5.14.3), rearrange the equation algebraically to solve for $[OH^-]$. $$[OH^-] = \frac{1.0 \times 10^{-14}}{[H_3O^+]}$$
Calculate.	Now substitute the known quantities into the equation and solve. $$[OH^-] = \frac{1.0 \times 10^{-14}}{1.0 \times 10^{-4}} = 1.0 \times 10^{-10} M$$ It is assumed that the concentration unit is molarity, so $[OH^-]$ is 1.0×10^{-10} M.
Think about your result.	The concentration of the acid is high (> 1 x 10^{-7} M), so $[OH^-]$ should be low.

Exercise 5.14.1

What is $[OH^-]$ in a 0.00032 M solution of H_2SO_4?

Hint

Assume **both** protons ionize from the molecule...although this is not the case.

Answer

$3.1 \times 10^{-11}\ M$

When you have a solution of a particular acid or base, you need to look at the formula of the acid or base to determine the number of H_3O^+ or OH^- ions in the formula unit because $[H_3O^+]$ or $[OH^-]$) may not be the same as the concentration of the acid or base itself.

Example 5.14.2: Hydronium Concentration

What is $[H_3O^+]$ in a 0.0044 M solution of $Ca(OH)_2$?

Solution

Solutions to Example 14.7.2

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: $[Ca(OH)_2] = 0.0044\,M$ Find: $[H_3O^+]$ = ? M
List other known quantities.	We begin by determining $[OH^-]$. The concentration of the solute is 0.0044 M, but because $Ca(OH)_2$ is a strong base, there are two OH^- ions in solution for every formula unit dissolved, so the actual $[OH^-]$ is two times this: $$[OH^-] = 2 \times 0.0044\ \text{M} = 0.0088\,\text{M}\cdot$$
Plan the problem.	Use the expression for K_w (Equation 5.14.3) and rearrange the equation algebraically to solve for $[H_3O^+]$. $$[H_3O^+] = \frac{1.0 \times 10^{-14}}{[OH^-]}$$
Calculate.	Now substitute the known quantities into the equation and solve. $$[H_3O^+] = \frac{1.0 \times 10^{-14}}{(0.0088)} = 1.1 \times 10^{-12} M$$ $[H_3O^+]$ has decreased significantly in this basic solution.
Think about your result.	The concentration of the base is high (> 1 x 10^{-7} M) so Extra close brace or missing open brace should be low.

? Exercise 5.14.2

What is $[H_3O^+]$ of an aqueous solution if $[OH^-]$ is $1.0 \times 10^{-9}\,M$?

Answer

1.0×10^{-5} M

In any aqueous solution, the product of $[H_3O^+]$ and $[OH^-]$ equals 1.0×10^{-14} (at room temperature).

Contributions & Attributions

5.14: Water - Acid and Base in One is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.8: Water - Acid and Base in One** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity

Learning Objectives

- Define pH and pOH.
- Determine the pH of acidic and basic solutions.
- Determine the hydronium ion concentration and pOH from pH.

As we have seen, $[H_3O^+]$ and $[OH^-]$ values can be markedly different from one aqueous solution to another. So chemists defined a new scale that succinctly indicates the concentrations of either of these two ions.

pH is a logarithmic function of $[H_3O^+]$:

$$pH = -\log[H_3O^+] \tag{5.15.1}$$

pH is usually (but not always) between 0 and 14. Knowing the dependence of pH on $[H_3O^+]$, we can summarize as follows:

- If pH < 7, then the solution is acidic.
- If pH = 7, then the solution is neutral.
- If pH > 7, then the solution is basic.

This is known as the pH scale. The pH scale is the range of values from 0 to 14 that describes the acidity or basicity of a solution. You can use pH to make a quick determination whether a given aqueous solution is acidic, basic, or neutral. Figure 5.15.1 illustrates this relationship, along with some examples of various solutions. Because hydrogen ion concentrations are generally less than one (for example $1.3 \times 10^{-3}\ M$), the log of the number will be a negative number. To make pH even easier to work with, pH is defined as the ***negative* log of** $[H_3O^+]$, which will give a positive value for pH.




Figure 5.15.1: The pH values for several common materials.

Example 5.15.1

Label each solution as acidic, basic, or neutral based only on the stated pH.

a. milk of magnesia, pH = 10.5
b. pure water, pH = 7
c. wine, pH = 3.0

Answer

a. With a pH greater than 7, milk of magnesia is basic. (Milk of magnesia is largely $Mg(OH)_2$.)

b. Pure water, with a pH of 7, is neutral.
c. With a pH of less than 7, wine is acidic.

? Exercise 5.15.1

Identify each substance as acidic, basic, or neutral based only on the stated pH.

a. human blood with pH = 7.4
b. household ammonia with pH = 11.0
c. cherries with pH = 3.6

Answer a

basic

Answer b

basic

Answer c

acidic

Calculating pH from Hydronium Concentration

The pH of solutions can be determined by using logarithms as illustrated in the next example for stomach acid. Stomach acid is a solution of HCl with a hydronium ion concentration of $1.2 \times 10^{-3}\ M$, what is the pH of the solution?

$$\begin{aligned} \text{pH} &= -\log[\text{H}_3\text{O}^+] \\ &= -\log(1.2 \times 10^{-3}) \\ &= -(-2.92) = 2.92 \end{aligned}$$

Logarithms

To get the **log value** on your calculator, enter the number (in this case, the hydronium ion concentration) first, then press the LOG key.

If the number is 1.0 x 10^{-5} (for $[H_3O^+]$ = 1.0 x 10^{-5} M) you should get an answer of "-5".

If you get a different answer, or an error, try pressing the LOG key before you enter the number.


Example 5.15.2: Converting Ph to Hydronium Concentration

Find the pH, given the $[H_3O^+]$ of the following:

a. 1 ×10^{-3} M
b. 2.5 ×10^{-11} M
c. 4.7 ×10^{-9} M

Solution

Solutions to Example 14.9.2

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: a. $[H_3O^+] = 1 \times 10^{-3}$ M b. $[H_3O^+] = 2.5 \times 10^{-11}$ M c. $[H_3O^+] = 4.7 \times 10^{-9}$ M Find: ? pH
Plan the problem.	Need to use the expression for pH (Equation 5.15.1). $pH = - \log [H_3O^+]$
Calculate.	Now substitute the known quantity into the equation and solve. a. $pH = - \log [1 \times 10^{-3}] = 3.0$ (1 decimal places since 1 has 1 significant figure) b. $pH = - \log [2.5 \times 10^{-11}] = 10.60$ (2 decimal places since 2.5 has 2 significant figures) c. $pH = - \log [4.7 \times 10^{-9}] = 8.30$ (2 decimal places since 4.7 has 2 significant figures) The other issue that concerns us here is significant figures. Because the number(s) before the decimal point in a logarithm relate to the power on 10, the number of digits *after* the decimal point is what determines the number of significant figures in the final answer: 

? Exercise 5.15.2

Find the pH, given $[H_3O^+]$ of the following:

a. 5.8×10^{-4} M
b. 1.0×10^{-7}

Answer a

3.22

Answer b

7.00

Calculating Hydronium Concentration from pH

Sometimes you need to work "backwards"—you know the pH of a solution and need to find $[H_3O^+]$, or even the concentration of the acid solution. How do you do that? To convert pH into $[H_3O^+]$ we solve Equation 5.15.1 for $[H_3O^+]$. This involves taking the antilog (or inverse log) of the negative value of pH .

$$[H_3O^+] = \text{antilog}(-pH)$$

or

$$[H_3O^+] = 10^{-pH} \tag{5.15.2}$$

As mentioned above, different calculators work slightly differently—make sure you can do the following calculations using ***your*** calculator.

Calculator Skills

We have a solution with a pH = 8.3. What is $[H_3O^+]$?

With some calculators you will do things in the following order:

1. Enter 8.3 as a negative number (use the key with both the +/- signs, not the subtraction key).
2. Use your calculator's 2nd or Shift or INV (inverse) key to type in the symbol found above the LOG key. The shifted function should be 10^x.
3. You should get the answer $\mathbf{5.0 \times 10^{-9}}$.

Other calculators require you to enter keys in the order they appear in the equation.

1. Use the Shift or second function to key in the 10^x function.
2. Use the +/- key to type in a negative number, then type in 8.3.
3. You should get the answer $\mathbf{5.0 \times 10^{-9}}$.

If neither of these methods work, try rearranging the order in which you type in the keys. Don't give up—you must master your calculator!

Example 5.15.3: Calculating Hydronium Concentration from pH

Find the hydronium ion concentration in a solution with a pH of 12.6. Is this solution an acid or a base? How do you know?

Solution

Solutions to Example 14.9.3

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: pH = 12.6 Find: $[H_3O^+]$ = ? M
Plan the problem.	Need to use the expression for $[H_3O^+]$ (Equation 5.15.2). $[H_3O^+]$ = antilog (-pH) or $[H_3O^+] = 10^{-pH}$
Calculate.	Now substitute the known quantity into the equation and solve. $[H_3O^+]$ = antilog (12.60) = 2.5 x 10^{-13} M (2 significant figures since 4.7 has 12.60 2 decimal places) or $[H_3O^+] = 10^{-12.60} = 2.5 \times 10^{-13}$ M (2 significant figures since 4.7 has 12.60 2 decimal places) The other issue that concerns us here is significant figures. Because the number(s) before the decimal point in a logarithm relate to the power on 10, the number of digits *after* the decimal point is what determines the number of significant figures in the final answer: X.YYY Y.YY x 10^x

Exercise 5.15.3

If moist soil has a pH of 7.84, what is $[H_3O^+]$ of the soil solution?

Answer

1.5 x 10^{-8} M

The pOH scale

As with the hydrogen-ion concentration, the concentration of the hydroxide ion can be expressed logarithmically by the pOH. The **pOH** of a solution is the negative logarithm of the hydroxide-ion concentration.

$$\mathrm{pOH} = -\log\left[\mathrm{OH}^-\right]$$

The relation between the hydronium and hydroxide ion concentrations expressed as p-functions is easily derived from the K_w expression:

$$K_\mathrm{w} = [\mathrm{H_3O^+}][\mathrm{OH}^-] \tag{5.15.3}$$

$$-\log K_\mathrm{w} = -\log([\mathrm{H_3O^+}][\mathrm{OH}^-]) = -\log[\mathrm{H_3O^+}] + -\log[\mathrm{OH}^-] \tag{5.15.4}$$

$$\mathrm{p}K_\mathrm{w} = \mathrm{pH} + \mathrm{pOH} \tag{5.15.5}$$

At 25 °C, the value of K_w is 1.0×10^{-14}, and so:

$$14.00 = \mathrm{pH} + \mathrm{pOH} \tag{5.15.6}$$

The hydronium ion molarity in pure water (or any neutral solution) is $1.0 \times 10^{-7}\ M$ at 25 °C. The pH and pOH of a neutral solution at this temperature are therefore:

$$\mathrm{pH} = -\log[\mathrm{H_3O^+}] = -\log(1.0 \times 10^{-7}) = 7.00 \tag{5.15.7}$$

$$\mathrm{pOH} = -\log[\mathrm{OH}^-] = -\log(1.0 \times 10^{-7}) = 7.00 \tag{5.15.8}$$

And so, at this temperature, acidic solutions are those with hydronium ion molarities greater than $1.0 \times 10^{-7}\ M$ and hydroxide ion molarities less than $1.0 \times 10^{-7}\ M$ (corresponding to pH values less than 7.00 and pOH values greater than 7.00). Basic solutions are those with hydronium ion molarities less than $1.0 \times 10^{-7}\ M$ and hydroxide ion molarities greater than $1.0 \times 10^{-7}\ M$ (corresponding to pH values greater than 7.00 and pOH values less than 7.00).

Example 5.15.4:

Find the pOH of a solution with a pH of 4.42.

Solution

Solutions to Example 14.9.4

Steps for Problem Solving	
Identify the "given" information and what the problem is asking you to "find."	Given: pH =4.42 Find: ? pOH
Plan the problem.	Need to use the expression pOH = 14 - pH
Calculate.	Now substitute the known quantity into the equation and solve. pOH=14−4.42=9.58
Think about your result.	The pH is that of an acidic solution, and the resulting pOH is the difference after subtracting from 14. The answer has two significant figures because the given pH has two decimal places.

Exercise 5.15.4

The pH of a solution is 8.22. What is the pOH?

Answer

5.78

The diagram below shows all of the interrelationships between [H3O+][H3O+], [OH−][OH−], pH, and pOH.




Figure **5.15.1**: Relationships between hydrogen ion concentration, hydroxide ion concentration, pH and pOH.

Contributions & Attributions

- Peggy Lawson (Oxbow Prairie Heights School). Funded by Saskatchewan Educational Technology Consortium.
-
- Template:OpenStax

5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity is shared under a CK-12 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew & Henry Agnew.

- **14.9: The pH and pOH Scales - Ways to Express Acidity and Basicity** by Henry Agnew, Marisa Alviar-Agnew is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

5.16: Carbon - A Versitile Atom

5.16: Carbon - A Versitile Atom is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

Learning Objectives

- Identify alkanes, alkenes, alkynes, and aromatic compounds.
- List some properties of hydrocarbons.

The simplest organic compounds are those composed of only two elements: carbon and hydrogen. These compounds are called **hydrocarbons**. Hydrocarbons themselves are separated into two types: aliphatic hydrocarbons and aromatic hydrocarbons.

Aliphatic hydrocarbons are hydrocarbons based on chains of C atoms. There are three types of aliphatic hydrocarbons. **Alkanes** are aliphatic hydrocarbons with only single covalent bonds. **Alkenes** are hydrocarbons that contain at least one C–C double bond, and **Alkynes** are hydrocarbons that contain a C–C triple bond. Occasionally, we find an aliphatic hydrocarbon with a ring of C atoms; these hydrocarbons are called *cycloalkanes* (or *cycloalkenes* or *cycloalkynes*).

Aromatic hydrocarbons have a special six-carbon ring called a *benzene* ring. Electrons in the benzene ring have special energetic properties that give benzene physical and chemical properties that are markedly different from alkanes. Originally, the term *aromatic* was used to describe this class of compounds because they were particularly fragrant. However, in modern chemistry the term *aromatic* denotes the presence of a six-membered ring that imparts different and unique properties to a molecule.

The simplest alkanes have their C atoms bonded in a straight chain; these are called *normal* alkanes. They are named according to the number of C atoms in the chain. The smallest alkane is methane:




Figure **5.17.1** - Three-Dimensional Representation of Methane.






Figure **5.17.1** Three-Dimensional Representation of Methane © Thinkstock. The methane molecule is three dimensional, with the H atoms in the positions of the four corners of a tetrahedron.

The next-largest alkane has two C atoms that are covalently bonded to each other. For each C atom to make four covalent bonds, each C atom must be bonded to three H atoms. The resulting molecule, whose formula is C_2H_6, is ethane:



Propane has a backbone of three C atoms surrounded by H atoms. You should be able to verify that the molecular formula for propane is C_3H_8:




The diagrams representing alkanes are called **structural formulas** because they show the structure of the molecule. As molecules get larger, structural formulas become more and more complex. One way around this is to use a **condensed structural formula**, which lists the formula of each C atom in the backbone of the molecule. For example, the condensed structural formula for ethane is CH_3CH_3, while for propane it is $CH_3CH_2CH_3$. Table **5.17.1** - The First 10 Alkanes, gives the molecular formulas, the condensed structural formulas, and the names of the first 10 alkanes.

Table **5.17.1** The First 10 Alkanes


Molecular Formula	Condensed Structural Formula	Name
CH_4	CH_4	methane
C_2H_6	CH_3CH_3	ethane
C_3H_8	$CH_3CH_2CH_3$	propane
C_4H_{10}	$CH_3CH_2CH_2CH_3$	butane
C_5H_{12}	$CH_3CH_2CH_2CH_2CH_3$	pentane
C_6H_{14}	$CH_3(CH_2)_4CH_3$	hexane
C_7H_{16}	$CH_3(CH_2)_5CH_3$	heptane
C_8H_{18}	$CH_3(CH_2)_6CH_3$	octane
C_9H_{20}	$CH_3(CH_2)_7CH_3$	nonane
$C_{10}H_{22}$	$CH_3(CH_2)_8CH_3$	decane

Because alkanes have the maximum number of H atoms possible according to the rules of covalent bonds, alkanes are also referred to as **saturated hydrocarbons**.

Alkenes have a C–C double bond. Because they have less than the maximum number of H atoms possible, they are **unsaturated hydrocarbons**. The smallest alkene—ethene—has two C atoms and is also known by its common name ethylene:




The next largest alkene—propene—has three C atoms with a C–C double bond between two of the C atoms. It is also known as propylene:



What do you notice about the names of alkanes and alkenes? The names of alkenes are the same as their corresponding alkanes except that the ending is *-ene*, rather than *-ane*. Using a stem to indicate the number of C atoms in a molecule and an ending to represent the type of organic compound is common in organic chemistry, as we shall see.

With the introduction of the next alkene, butene, we begin to see a major issue with organic molecules: choices. With four C atoms, the C–C double bond can go between the first and second C atoms or between the second and third C atoms:




2 structural formulas for butene, with the first butene having the double bond on the first and second carbon from the left and the latter having its double bond on the second and third carbon from the left.

(A double bond between the third and fourth C atoms is the same as having it between the first and second C atoms, only flipped over.) The rules of naming in organic chemistry require that these two substances have different names. The first molecule is named *1-butene*, while the second molecule is named *2-butene*. The number at the beginning of the name indicates where the double bond originates. The lowest possible number is used to number a feature in a molecule; hence, calling the second molecule 3-butene would be incorrect. Numbers are common parts of organic chemical names because they indicate which C atom in a chain contains a distinguishing feature.

The compounds 1-butene and 2-butene have different physical and chemical properties, even though they have the same molecular formula—C_4H_8. Different molecules with the same molecular formula are called **isomers**. Isomers are common in organic chemistry and contribute to its complexity.

Example 5.17.1

Based on the names for the butene molecules, propose a name for this molecule.




A structural formula of a five carbon molecule with a double bond on the third and fourth carbon from the left. There are ten hydrogen atoms in total.

Solution

With five C atoms, we will use the *pent-* stem, and with a C–C double bond, this is an alkene, so this molecule is a pentene. In numbering the C atoms, we use the number *2* because it is the lower possible label. So this molecule is named 2-pentene.

Exercise 5.17.1

Based on the names for the butene molecules, propose a name for this molecule.



A structural formula of a six carbon molecule with a double bond on the third and fourth carbon from the left. There are twelve hydrogen atoms in total.


Answer

3-hexene

Alkynes, with a C–C triple bond, are named similarly to alkenes except their names end in *-yne*. The smallest alkyne is ethyne, which is also known as acetylene:




Propyne has the structure




Structural formula showing three carbon molecules with a triple bond present between the first and second carbon atom. The appropriate number of hydrogen atoms is attached to each carbon atom.

With butyne, we need to start numbering the position of the triple bond, just as we did with alkenes:



Two structural formula of butyne. One butyne has a triple bond between the first and second carbon atom, while two butyne has the triple bond between the second and third carbon atom.

Aromatic compounds contain the benzene unit. Benzene itself is composed of six C atoms in a ring, with alternating single and double C–C bonds:



The six carbons are arranged in a hexagon pattern with one hydrogen atom emerging outwards from each carbon atom. The presence of a double bond is alternated between every other carbon atom.

The alternating single and double C–C bonds give the benzene ring a special stability, and it does not react like an alkene as might be suspected. Benzene has the molecular formula C_6H_6; in larger aromatic compounds, a different atom replaces one or more of the H atoms.


As fundamental as hydrocarbons are to organic chemistry, their properties and chemical reactions are rather mundane. Most hydrocarbons are nonpolar because of the close electronegativities of the C and H atoms. As such, they dissolve only sparingly in H_2O and other polar solvents. Small hydrocarbons, such as methane and ethane, are gases at room temperature, while larger hydrocarbons, such as hexane and octane, are liquids. Even larger hydrocarbons are solids at room temperature and have a soft, waxy consistency.

Hydrocarbons are rather unreactive, but they do participate in some classic chemical reactions. One common reaction is substitution with a halogen atom by combining a hydrocarbon with an elemental halogen. Light is sometimes used to promote the reaction, such as this one between methane and chlorine:

$$CH_4 + Cl_2 \overset{light}{\rightarrow} CH_3Cl + HCl$$

Halogens can also react with alkenes and alkynes, but the reaction is different. In these cases, the halogen reacts with the C–C double or triple bond and inserts itself onto each C atom involved in the multiple bonds. This reaction is called an **addition**

reaction. One example is



Structural formula showing the reaction of ethylene with a chlorine molecule to form ethylene dichloride.

The reaction conditions are usually mild; in many cases, the halogen reacts spontaneously with an alkene or an alkyne.

Hydrogen can also be added across a multiple bond; this reaction is called a **hydrogenation reaction**. In this case, however, the reaction conditions may not be mild; high pressures of H_2 gas may be necessary. A platinum or palladium catalyst is usually employed to get the reaction to proceed at a reasonable pace:

$$CH_2 = CH_2 + H_2 \overset{metal\ catalyst}{\rightarrow} CH_3CH_3$$

By far the most common reaction of hydrocarbons is combustion, which is the combination of a hydrocarbon with O_2 to make CO_2 and H_2O. The combustion of hydrocarbons is accompanied by a release of energy and is a primary source of energy production in our society (Figure **5.17.2** - Combustion). The combustion reaction for gasoline, for example, which can be represented by C_8H_{18}, is as follows:

$$2C^8H_{18} + 25O_2 \rightarrow 16CO_2 + 18H_2O + \sim 5060kJ$$




Figure **5.17.2** Combustion © Thinkstock. The combustion of hydrocarbons is a primary source of energy in our society.


Key Takeaways


- The simplest organic compounds are hydrocarbons and are composed of carbon and hydrogen.
- Hydrocarbons can be aliphatic or aromatic; aliphatic hydrocarbons are divided into alkanes, alkenes, and alkynes.
- The combustion of hydrocarbons is a primary source of energy for our society.

? Exercise 5.17.2


1. Define *hydrocarbon*. What are the two general types of hydrocarbons?
2. What are the three different types of aliphatic hydrocarbons? How are they defined?
3. Indicate whether each molecule is an aliphatic or an aromatic hydrocarbon; if aliphatic, identify the molecule as an alkane, an alkene, or an alkyne.


a. 


b. 

c. 


4. Indicate whether each molecule is an aliphatic or an aromatic hydrocarbon; if aliphatic, identify the molecule as an alkane, an alkene, or an alkyne.

a. 


b. 


c. 

5. Indicate whether each molecule is an aliphatic or an aromatic hydrocarbon; if aliphatic, identify the molecule as an alkane, an alkene, or an alkyne.


a. 


https://chem.libretexts.org/@go/page/492211


b. 

c. 

6. Indicate whether each molecule is an aliphatic or an aromatic hydrocarbon; if aliphatic, identify the molecule as an alkane, an alkene, or an alkyne.


a. 


b. 


c. 


7. Name and draw the structural formulas for the four smallest alkanes.
8. Name and draw the structural formulas for the four smallest alkenes.
9. What does the term *aromatic* imply about an organic molecule?
10. What does the term *normal* imply when used for alkanes?
11. Explain why the name *1-propene* is incorrect. What is the proper name for this molecule?
12. Explain why the name *3-butene* is incorrect. What is the proper name for this molecule?
13. Name and draw the structural formula of each isomer of pentene.
14. Name and draw the structural formula of each isomer of hexyne.
15. Write a chemical equation for the reaction between methane and bromine.
16. Write a chemical equation for the reaction between ethane and chlorine.
17. Draw the structure of the product of the reaction of bromine with propene.
18. Draw the structure of the product of the reaction of chlorine with 2-butene.
19. Draw the structure of the product of the reaction of hydrogen with 1-butene.
20. Draw the structure of the product of the reaction of hydrogen with 1-butene.
21. Write the balanced chemical equation for the combustion of heptane.
22. Write the balanced chemical equation for the combustion of nonane.


Answers


1. an organic compound composed of only carbon and hydrogen; aliphatic hydrocarbons and aromatic hydrocarbons
2.
3. a. aliphatic; alkane
   b. aromatic
   c. aliphatic; alkene
4.
5. a. aliphatic; alkane
   b. aliphatic; alkene
   c. aromatic
6. 








7.
8.
9. Aromatic means that the molecule has a benzene ring.
10.
11. The 1 is not necessary. The name of the compound is simply *propene*.
12. 




13.
14.
15. $CH_4 + Br_2 \rightarrow CH_3Br + HBr$

16.

17. 

18.

19. 

20.

21. $C_7H_{16} + 11O_2 \rightarrow 7CO_2 + 8H_2O$

This page titled 5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon is shared under a CC BY-NC-SA 3.0 license and was authored, remixed, and/or curated by Anonymous via source content that was edited to the style and standards of the LibreTexts platform.

- **16.2: Hydrocarbons** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/beginning-chemistry.

5.18: Alkanes- Saturated Hydrocarbons

Learning Objectives

- To identify and name simple (straight-chain) alkanes given formulas and write formulas for straight-chain alkanes given their names.

We begin our study of organic chemistry with the hydrocarbons, the simplest organic compounds, which are composed of carbon and hydrogen atoms only. As we noted, there are several different kinds of hydrocarbons. They are distinguished by the types of bonding between carbon atoms and the properties that result from that bonding. Hydrocarbons with only carbon-to-carbon single bonds (C–C) and existing as a continuous chain of carbon atoms also bonded to hydrogen atoms are called alkanes (or saturated hydrocarbons). *Saturated*, in this case, means that each carbon atom is bonded to four other atoms (hydrogen or carbon)—the most possible; there are no double or triple bonds in the molecules.

> The word *saturated* has the same meaning for hydrocarbons as it does for the dietary fats and oils: the molecule has no carbon-to-carbon double bonds (C=C).

We previously introduced the three simplest alkanes—methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) and they are shown again in Figure **5.18.1**.




Figure **5.18.1**: The Three Simplest Alkanes

The flat representations shown do not accurately portray bond angles or molecular geometry. Methane has a tetrahedral shape that chemists often portray with wedges indicating bonds coming out toward you and dashed lines indicating bonds that go back away from you. An ordinary solid line indicates a bond in the plane of the page. Recall that the VSEPR theory correctly predicts a tetrahedral shape for the methane molecule (Figure **5.18.2**).




Figure **5.18.2**: The Tetrahedral Methane Molecule

Methane (CH_4), ethane (C_2H_6), and propane (C_3H_8) are the beginning of a series of compounds in which any two members in a sequence differ by one carbon atom and two hydrogen atoms—namely, a CH_2 unit. The first 10 members of this series are given in Table **5.18.1**.

Table **5.18.1**: The First 10 Straight-Chain Alkanes

Name	Molecular Formula (C_nH_{2n+2})	Condensed Structural Formula	Number of Possible Isomers
methane	CH_4	CH_4	—
ethane	C_2H_6	CH_3CH_3	—
propane	C_3H_8	$CH_3CH_2CH_3$	—
butane	C_4H_{10}	$CH_3CH_2CH_2CH_3$	2
pentane	C_5H_{12}	$CH_3CH_2CH_2CH_2CH_3$	3
hexane	C_6H_{14}	$CH_3CH_2CH_2CH_2CH_2CH_3$	5
heptane	C_7H_{16}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_3$	9
octane	C_8H_{18}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	18
nonane	C_9H_{20}	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	35
decane	$C_{10}H_{22}$	$CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_2CH_3$	75

Consider the series in Figure **5.18.3**. The sequence starts with C_3H_8, and a CH_2 unit is added in each step moving up the series. Any family of compounds in which adjacent members differ from each other by a definite factor (here a CH_2 group) is called a homologous series. The members of such a series, called *homologs*, have properties that vary in a regular and predictable manner. The principle of *homology* gives organization to organic chemistry in much the same way that the periodic table gives organization to inorganic chemistry. Instead of a bewildering array of individual carbon compounds, we can study a few members of a homologous series and from them deduce some of the properties of other compounds in the series.




Figure **5.18.3**: Members of a Homologous Series. Each succeeding formula incorporates one carbon atom and two hydrogen atoms more than the previous formula.

The principle of homology allows us to write a general formula for alkanes: C_nH_{2n+2}. Using this formula, we can write a molecular formula for any alkane with a given number of carbon atoms. For example, an alkane with eight carbon atoms has the molecular formula $C_8H_{(2\times 8)+2} = C_8H_{18}$.

Key Takeaway

- Simple alkanes exist as a homologous series, in which adjacent members differ by a CH_2 unit.

5.18: Alkanes- Saturated Hydrocarbons is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **12.2: Structures and Names of Alkanes** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.19: Isomers- Same Formula, Different Structure

5.19: Isomers- Same Formula, Different Structure is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

5.20: Naming Alkanes


5.20: Naming Alkanes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

5.21: Alkenes and Alkynes

Learning Objectives

- To name alkenes given formulas and write formulas for alkenes given names.

As noted before, alkenes are hydrocarbons with carbon-to-carbon double bonds ($R_2C{=}CR_2$) and alkynes are hydrocarbons with carbon-to-carbon triple bonds (R–C≡C–R). Collectively, they are called unsaturated hydrocarbons because they have fewer hydrogen atoms than does an alkane with the same number of carbon atoms, as is indicated in the following general formulas:




Some representative alkenes—their names, structures, and physical properties—are given in Table **5.21.1**.

Table **5.21.1**: Physical Properties of Some Selected Alkenes

IUPAC Name	Molecular Formula	Condensed Structural Formula	Melting Point (°C)	Boiling Point (°C)
ethene	C_2H_4	$CH_2{=}CH_2$	–169	–104
propene	C_3H_6	$CH_2{=}CHCH_3$	–185	–47
1-butene	C_4H_8	$CH_2{=}CHCH_2CH_3$	–185	–6
1-pentene	C_5H_{10}	$CH_2{=}CH(CH_2)_2CH_3$	–138	30
1-hexene	C_6H_{12}	$CH_2{=}CH(CH_2)_3CH_3$	–140	63
1-heptene	C_7H_{14}	$CH_2{=}CH(CH_2)_4CH_3$	–119	94
1-octene	C_8H_{16}	$CH_2{=}CH(CH_2)_5CH_3$	–102	121

We used only condensed structural formulas in Table **5.21.1**. Thus, $CH_2{=}CH_2$ stands for



The double bond is shared by the two carbons and does not involve the hydrogen atoms, although the condensed formula does not make this point obvious. Note that the molecular formula for ethene is C_2H_4, whereas that for ethane is C_2H_6.

The first two alkenes in Table **5.21.1**, ethene and propene, are most often called by their common names—ethylene and propylene, respectively (Figure **5.21.1**). Ethylene is a major commercial chemical. The US chemical industry produces about 25 billion kilograms of ethylene annually, more than any other synthetic organic chemical. More than half of this ethylene goes into the manufacture of polyethylene, one of the most familiar plastics. Propylene is also an important industrial chemical. It is converted to plastics, isopropyl alcohol, and a variety of other products.





Figure **5.21.1**: Ethene and Propene. The ball-and-spring models of ethene/ethylene (a) and propene/propylene (b) show their respective shapes, especially bond angles.

Although there is only one alkene with the formula C_2H_4 (ethene) and only one with the formula C_3H_6 (propene), there are several alkenes with the formula C_4H_8.


Here are some basic rules for naming alkenes from the International Union of Pure and Applied Chemistry (IUPAC):


1. The longest chain of carbon atoms *containing the double bond* is considered the parent chain. It is named using the same stem as the alkane having the same number of carbon atoms but ends in *-ene* to identify it as an alkene. Thus the compound CH_2=$CHCH_3$ is *propene*.
2. If there are four or more carbon atoms in a chain, we must indicate the position of the double bond. The carbons atoms are numbered so that the first of the two that are doubly bonded is given the lower of the two possible numbers.The compound CH_3CH=$CHCH_2CH_3$, for example, has the double bond between the second and third carbon atoms. Its name is 2-pentene (not 3-pentene).
3. Substituent groups are named as with alkanes, and their position is indicated by a number. Thus, the structure below is 5-methyl-2-hexene. Note that the numbering of the parent chain is always done in such a way as to give the double bond the lowest number, even if that causes a substituent to have a higher number. *The double bond always has priority in numbering.*



Example 5.21.1

Name each compound.

a. 

b. 

Solution


a. The longest chain containing the double bond has five carbon atoms, so the compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the fourth carbon atom (rule 3), so the compound's name is 4-methyl-2-pentene.

b. The longest chain containing the double bond has five carbon atoms, so the parent compound is a *pentene* (rule 1). To give the first carbon atom of the double bond the lowest number (rule 2), we number from the left, so the compound is a 2-pentene. There is a methyl group on the third carbon atom (rule 3), so the compound's name is 3-methyl-2-pentene.

Exercise 5.21.1

Name each compound.

a. $CH_3CH_2CH_2CH_2CH_2CH{=}CHCH_3$

b. 

Answer

Just as there are cycloalkanes, there are *cycloalkenes*. These compounds are named like alkenes, but with the prefix *cyclo-* attached to the beginning of the parent alkene name.

Example 5.21.2


Draw the structure for each compound.

a. 3-methyl-2-pentene
b. cyclohexene


Solution

a.

First write the parent chain of five carbon atoms: C–C–C–C–C. Then add the double bond between the second and third carbon atoms:




Now place the methyl group on the third carbon atom and add enough hydrogen atoms to give each carbon atom a total of four bonds.



b

First, consider what each of the three parts of the name means. *Cyclo* means a ring compound, *hex* means 6 carbon atoms, and *-ene* means a double bond.



Exercise 5.21.2

Draw the structure for each compound.

a. 2-ethyl-1-hexene
b. cyclopentene

Key Takeaway

- Alkenes are hydrocarbons with a carbon-to-carbon double bond.

5.21: Alkenes and Alkynes is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.




- **13.1: Alkenes- Structures and Names** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.22: Functional Groups

Learning Objectives

- To know the major classes of organic compounds and identify important functional groups.


You were previously introduced to several structural units that chemists use to classify organic compounds and predict their reactivities. These functional groups, which determine the chemical reactivity of a molecule under a given set of conditions, can consist of a single atom (such as Cl) or a group of atoms (such as CO_2H). The major families of organic compounds are characterized by their functional groups. Figure **5.22.1** summarizes five families introduced in earlier chapters, gives examples of compounds that contain each functional group, and lists the suffix or prefix used in the systematic nomenclature of compounds that contain each functional group.

Class	General Formula	Example	Common Name (Systematic Name)	Common Suffix/Prefix (Systematic)
Hydrocarbons				
Alkanes	RH	CH_3CH_3	ethane	-ane
Alkenes	$RR'C{=}CR''R'''$	$H_2C{=}CH_2$	ethylene (ethene)	-ene
Alkynes	$RC{\equiv}CR'$	$HC{\equiv}CH$	acetylene (ethyne)	(-yne)
Arenes	ArH[a]		benzene	-ene
Halogen-Containing Compounds				
Alkyl halides	RX	CH_3CH_2Cl	ethyl chloride (chloroethane)	halide (halo-)
Aryl halides	ArX[a]		chlorobenzene	halo-
Oxygen-Containing Compounds				
Alcohols	ROH[a]	CH_3CH_2OH	ethyl alcohol (ethanol)	-ol
Phenols	ArOH[b]		phenol	-ol
Ethers	ROR′	$H_3CH_2COCH_2CH_3$	diethyl ether	ether
Aldehydes	RCHO	$CH_3\overset{\overset{O}{\|}}{C}H$	acetaldehyde (ethanal)	-aldehyde (-al)
Ketones	RR′C=O	$CH_3\overset{\overset{O}{\|}}{C}CH_3$	acetone (2-propanone)	-one
Carboxylic acids	RCO_2H	$CH_3\overset{\overset{O}{\|}}{C}OH$	acetic acid (ethanoic acid)	-ic acid (-oic acid)
Carboxylic Acid Derivatives				
Esters	RCO_2R'	$CH_3\overset{\overset{O}{\|}}{C}OCH_3$	methyl acetate (methyl ethanoate)	-ate (-oate)
Amides	RCONHR′	$CH_3\overset{\overset{O}{\|}}{C}NHCH_3$	*N*-methylacetamide	-amide
Nitrogen-Containing Compounds				
Amines	RNH_2, RNHR′, RNR′R″	$CH_3CH_2NH_2$	ethylamine	-amine
Nitriles	$RC{\equiv}N$	$H_3CC{\equiv}N$	acetonitrile	-nitrile
Nitro compounds	$ArNO_2$[a]		nitrobenzene	nitro-

[a]R indicates an alkyl group [b]Ar indicates an aryl group Figure 5.22.1: Major Classes of Organic Compounds

The first family listed in Figure 5.22.1 is the hydrocarbons. These include alkanes, with the general molecular formula C_nH_{2n+2} where n is an integer; alkenes, represented by C_nH_{2n}; alkynes, represented by $C_nH_{2n−2}$; and arenes. Halogen-substituted alkanes, alkenes, and arenes form a second major family of organic compounds, which include the alkyl halides and the aryl halides. Oxygen-containing organic compounds, a third family, may be divided into two main types: those that contain at least one C–O bond, which include alcohols, phenols (derivatives of benzene), and ethers, and those that contain a carbonyl group (C=O), which include aldehydes, ketones, and carboxylic acids. Carboxylic acid derivatives, the fourth family listed, are compounds in which the OH of the $–CO_2H$ functional group is replaced by either an alkoxy (–OR) group, producing an ester, or by an amido (–NRR′, where R and R′ can be H and/or alkyl groups), forming an amide. Nitrogen-containing organic compounds, the fifth family, include amines; nitriles, which have a C≡N bond; and nitro compounds, which contain the $–NO_2$ group.

The systematic nomenclature of organic compounds indicates the positions of substituents using the lowest numbers possible to identify their locations in the carbon chain of the parent compound. If two compounds have the same systematic name, then they are the same compound. Although systematic names are preferred because they are unambiguous, many organic compounds are known by their common names rather than their systematic names. Common nomenclature uses the prefix form—for a compound that contains no carbons other than those in the functional group, and acet—for those that have one carbon atom in addition [two in the case of acetone, $(CH_3)_2C=O$]. Thus methanal and ethanal, respectively, are the systematic names for formaldehyde and acetaldehyde.



Recall that in the systematic nomenclature of aromatic compounds, the positions of groups attached to the aromatic ring are indicated by numbers, starting with 1 and proceeding around the ring in the direction that produces the lowest possible numbers. For example, the position of the first CH_3 group in dimethyl benzene is indicated with a 1, but the second CH_3 group, which can be placed in any one of three positions, produces 1,2-dimethylbenzene, 1,3-dimethylbenzene, or 1,4-dimethylbenzene (Figure 5.22.2). In common nomenclature, in contrast, the prefixes ortho-, meta-, and para- are used to describe the relative positions of groups attached to an aromatic ring. If the CH_3 groups in dimethylbenzene, whose common name is xylene, are adjacent to each other, the compound is commonly called ortho-xylene, abbreviated o-xylene. If they are across from each other on the ring, the compound is commonly called para-xylene or p-xylene. When the arrangement is intermediate between those of ortho- and para- compounds, the name is meta-xylene or m-xylene.




Prefix	Positions of Methyl Groups	Example	Common Name (Systematic Name)
ortho-	1,2		*o*-xylene (1,2-dimethylbenzene)
meta-	1,3		*m*-xylene (1,3-dimethylbenzene)
para-	1,4		*p*-xylene (1,4-dimethylbenzene)

Figure 5.22.2: Common Nomenclature for Aromatic Ring Substitutions

We begin our discussion of the structure and reactivity of organic compounds by exploring structural variations in the simple saturated hydrocarbons known as alkanes. These compounds serve as the scaffolding to which the various functional groups are most often attached.

Summary

Functional groups determine the chemical reactivity of an organic molecule. Functional groups are structural units that determine the chemical reactivity of a molecule under a given set of conditions. Organic compounds are classified into several major categories based on the functional groups they contain. In the systematic names of organic compounds, numbers indicate the positions of functional groups in the basic hydrocarbon framework. Many organic compounds also have common names, which use the prefix form—for a compound that contains no carbons other than those in the functional group and acet—for those that have one additional carbon atom.

Conceptual Problems

1. Can two substances have the same systematic name and be different compounds?
2. Is a carbon–carbon multiple bond considered a functional group?

5.22: Functional Groups is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.


- **23.2: Functional Groups and Classes of Organic Compounds** by Anonymous is licensed CC BY-NC-SA 3.0.

5.23: Carbohydrates

Learning Objectives

- To recognize carbohydrates and classify them as mono-, di-, or polysaccharides.


All carbohydrates consist of carbon, hydrogen, and oxygen atoms and are polyhydroxy aldehydes or ketones or are compounds that can be broken down to form such compounds. Examples of carbohydrates include starch, fiber, the sweet-tasting compounds called sugars, and structural materials such as cellulose. The term *carbohydrate* had its origin in a misinterpretation of the molecular formulas of many of these substances. For example, because its formula is $C_6H_{12}O_6$, glucose was once thought to be a "carbon hydrate" with the structure $C_6 \cdot 6H_2O$.





Because glucose has an aldehyde functional group and several –OH (hydroxyl) groups, it is described as a polyhydroxy aldehyde.


Example 5.23.1

Which compounds would be classified as carbohydrates?

a. 

b. 

c. 

d. 

Solution

a. This is a carbohydrate because the molecule contains an aldehyde functional group with OH groups on the other two carbon atoms.





b. This is not a carbohydrate because the molecule does not contain an aldehyde or a ketone functional group.

c. This is a carbohydrate because the molecule contains a ketone functional group with OH groups on the other two carbon atoms.

d. This is not a carbohydrate; although it has a ketone functional group, one of the other carbons atoms does not have an OH group attached.

? Exercise 5.23.1

Which compounds would be classified as carbohydrates?

1. 
2. 
3. 
4. 

Green plants are capable of synthesizing glucose ($C_6H_{12}O_6$) from carbon dioxide (CO_2) and water (H_2O) by using solar energy in the process known as photosynthesis:

$$6\,CO_2 + 6\,H_2O + 686\ \text{kcal} \rightarrow C_6H_{12}O_6 + 6\,O_2 \tag{5.23.1}$$

(The 686 kcal come from solar energy.) Plants can use the glucose for energy or convert it to larger carbohydrates, such as starch or cellulose. Starch provides energy for later use, perhaps as nourishment for a plant's seeds, while cellulose is the structural material of plants. We can gather and eat the parts of a plant that store energy—seeds, roots, tubers, and fruits—and use some of that energy ourselves. Carbohydrates are also needed for the synthesis of nucleic acids and many proteins and lipids.

Animals, including humans, cannot synthesize carbohydrates from carbon dioxide and water and are therefore dependent on the plant kingdom to provide these vital compounds. We use carbohydrates not only for food (about 60%–65% by mass of the average diet) but also for clothing (cotton, linen, rayon), shelter (wood), fuel (wood), and paper (wood).

The simplest carbohydrates—those that cannot be hydrolyzed to produce even smaller carbohydrates—are called monosaccharides. Two or more monosaccharides can link together to form chains that contain from two to several hundred or thousand monosaccharide units. Prefixes are used to indicate the number of such units in the chains. Disaccharide molecules have two monosaccharide units, *trisaccharide* molecules have three units, and so on. Chains with many monosaccharide units joined together are called polysaccharides. All these so-called higher saccharides can be hydrolyzed back to their constituent monosaccharides.

> *Compounds that cannot be hydrolyzed will not react with water to form two or more smaller compounds.*

Summary

Carbohydrates are an important group of biological molecules that includes sugars and starches. Photosynthesis is the process by which plants use energy from sunlight to synthesize carbohydrates. A monosaccharide is the simplest carbohydrate and cannot be hydrolyzed to produce a smaller carbohydrate molecule. Disaccharides contain two monosaccharide units, and polysaccharides contain many monosaccharide units.

5.23: Carbohydrates is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **16.1: Carbohydrates** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.24: Lipids

On July 11, 2003, the Food and Drug Administration amended its food labeling regulations to require that manufacturers list the amount of *trans* fatty acids on Nutrition Facts labels of foods and dietary supplements, effective January 1, 2006. This amendment was a response to published studies demonstrating a link between the consumption of *trans* fatty acids and an increased risk of heart disease. *Trans* fatty acids are produced in the conversion of liquid oils to solid fats, as in the creation of many commercial margarines and shortenings. They have been shown to increase the levels of low-density lipoproteins (LDLs)—complexes that are often referred to as bad cholesterol—in the blood. In this chapter, you will learn about fatty acids and what is meant by a *trans* fatty acid, as well as the difference between fats and oils. You will also learn what cholesterol is and why it is an important molecule in the human body.

Fats and oils, found in many of the foods we eat, belong to a class of biomolecules known as lipids. Gram for gram, they pack more than twice the caloric content of carbohydrates: the oxidation of fats and oils supplies about 9 kcal of energy for every gram oxidized, whereas the oxidation of carbohydrates supplies only 4 kcal/g. Although the high caloric content of fats may be bad news for the dieter, it says something about the efficiency of nature's designs. Our bodies use carbohydrates, primarily in the form of glucose, for our *immediate* energy needs. Our capacity for storing carbohydrates for later use is limited to tucking away a bit of glycogen in the liver or in muscle tissue. We store our *reserve* energy in lipid form, which requires far less space than the same amount of energy stored in carbohydrate form. Lipids have other biological functions besides energy storage. They are a major component of the membranes of the 10 trillion cells in our bodies. They serve as protective padding and insulation for vital organs. Furthermore, without lipids in our diets, we would be deficient in the fat-soluble vitamins A, D, E, and K.

Lipids are not defined by the presence of specific functional groups, as carbohydrates are, but by a physical property—solubility. Compounds isolated from body tissues are classified as lipids if they are more soluble in organic solvents, such as dichloromethane, than in water. By this criterion, the lipid category includes not only fats and oils, which are esters of the trihydroxy alcohol glycerol and fatty acids, but also compounds that incorporate functional groups derived from phosphoric acid, carbohydrates, or amino alcohols, as well as steroid compounds such as cholesterol (Figure 5.24.1 presents one scheme for classifying the various kinds of lipids). We will discuss the various kinds of lipids by considering one subclass at a time and pointing out structural similarities and differences as we go.





Figure 5.24.1: Lipid Organization Based on Structural Relationships

Lipid categorized into fatty acids and steroids. Fatty acids are further separated into triglycerides, phospho-glycerides, waxes, and sphingolipids. Sphingolipids are separated into sphingo-myelins and glycolipids. Glycolipids are separated into cerebrosides and gangliosides.

Learning Objectives

- To recognize the structures of common fatty acids and classify them as saturated, monounsaturated, or polyunsaturated.

Fatty acids are carboxylic acids that are structural components of fats, oils, and all other categories of lipids, except steroids. More than 70 have been identified in nature. They usually contain an even number of carbon atoms (typically 12–20), are generally unbranched, and can be classified by the presence and number of carbon-to-carbon double bonds. Thus, saturated fatty acids contain no carbon-to-carbon double bonds, monounsaturated fatty acids contain one carbon-to-carbon double bond, and polyunsaturated fatty acids contain two or more carbon-to-carbon double bonds.



Cis fatty acids have two H atoms on the same side of the plane of the fatty acid while trans has the two H atoms on opposite sides.

Table **5.24.1** lists some common fatty acids and one important source for each. The atoms or groups around the double bonds in unsaturated fatty acids can be arranged in either the cis or trans isomeric form. Naturally occurring fatty acids are generally in the cis configuration.

Table **5.24.1**: Some Common Fatty Acids Found in Natural Fats


Name	Abbreviated Structural Formula	Condensed Structural Formula	Melting Point (°C)	Source
lauric acid	$C_{11}H_{23}COOH$	$CH_3(CH_2)_{10}COOH$	44	palm kernel oil
myristic acid	$C_{13}H_{27}COOH$	$CH_3(CH_2)_{12}COOH$	58	oil of nutmeg
palmitic acid	$C_{15}H_{31}COOH$	$CH_3(CH_2)_{14}COOH$	63	palm oil
palmitoleic acid	$C_{15}H_{29}COOH$	$CH_3(CH_2)_5CH=CH(CH_2)_7COOH$	0.5	macadamia oil
stearic acid	$C_{17}H_{35}COOH$	$CH_3(CH_2)_{16}COOH$	70	cocoa butter
oleic acid	$C_{17}H_{33}COOH$	$CH_3(CH_2)_7CH=CH(CH_2)_7COOH$	16	olive oil
linoleic acid	$C_{17}H_{31}COOH$	$CH_3(CH_2)_3(CH_2CH=CH)_2(CH_2)_7COOH$	−5	canola oil
α-linolenic acid	$C_{17}H_{29}COOH$	$CH_3(CH_2CH=CH)_3(CH_2)_7COOH$	−11	flaxseed
arachidonic acid	$C_{19}H_{31}COOH$	$CH_3(CH_2)_4(CH_2CH=CH)_4(CH_2)_2COOH$	−50	liver

Two polyunsaturated fatty acids—linoleic and α-linolenic acids—are termed essential fatty acids because humans must obtain them from their diets. Both substances are required for normal growth and development, but the human body does not synthesize them. The body uses linoleic acid to synthesize many of the other unsaturated fatty acids, such as arachidonic acid, a precursor for the synthesis of prostaglandins. In addition, the essential fatty acids are necessary for the efficient transport and metabolism of cholesterol. The average daily diet should contain about 4–6 g of the essential fatty acids.

To Your Health: Prostaglandins

Prostaglandins are chemical messengers synthesized in the cells in which their physiological activity is expressed. They are unsaturated fatty acids containing 20 carbon atoms and are synthesized from arachidonic acid—a polyunsaturated fatty acid—when needed by a particular cell. They are called *prostaglandins* because they were originally isolated from semen found in the prostate gland. It is now known that they are synthesized in nearly all mammalian tissues and affect almost all organs in the body. The five major classes of prostaglandins are designated as PGA, PGB, PGE, PGF, and PGI. Subscripts are attached at the end of these abbreviations to denote the number of double bonds outside the five-carbon ring in a given prostaglandin.

The prostaglandins are among the most potent biological substances known. Slight structural differences give them highly distinct biological effects; however, all prostaglandins exhibit some ability to induce smooth muscle contraction, lower blood pressure, and contribute to the inflammatory response. Aspirin and other nonsteroidal anti-inflammatory agents, such as ibuprofen, obstruct the synthesis of prostaglandins by inhibiting cyclooxygenase, the enzyme needed for the initial step in the conversion of arachidonic acid to prostaglandins.



Their wide range of physiological activity has led to the synthesis of hundreds of prostaglandins and their analogs. Derivatives of PGE_2 are now used in the United States to induce labor. Other prostaglandins have been employed clinically to lower or increase blood pressure, inhibit stomach secretions, relieve nasal congestion, relieve asthma, and prevent the formation of blood clots, which are associated with heart attacks and strokes.

Although we often draw the carbon atoms in a straight line, they actually have more of a zigzag configuration (Figure **5.24.2*a***). Viewed as a whole, however, the saturated fatty acid molecule is relatively straight (Figure **5.24.2*b***). Such molecules pack closely together into a crystal lattice, maximizing the strength of dispersion forces and causing fatty acids and the fats derived from them to have relatively high melting points. In contrast, each *cis* carbon-to-carbon double bond in an unsaturated fatty acid produces a pronounced bend in the molecule, so that these molecules do not stack neatly. As a result, the intermolecular attractions of unsaturated fatty acids (and unsaturated fats) are weaker, causing these substances to have lower melting points. Most are liquids at room temperature.





Figure 5.24.2: The Structure of Saturated Fatty Acids. (a) There is a zigzag pattern formed by the carbon-to-carbon single bonds in the ball-and-stick model of a palmitic acid molecule. (b) A space-filling model of palmitic acid shows the overall straightness of a saturated fatty acid molecule.

Waxes are esters formed from long-chain fatty acids and long-chain alcohols. Most natural waxes are mixtures of such esters. Plant waxes on the surfaces of leaves, stems, flowers, and fruits protect the plant from dehydration and invasion by harmful microorganisms. Carnauba wax, used extensively in floor waxes, automobile waxes, and furniture polish, is largely myricyl cerotate, obtained from the leaves of certain Brazilian palm trees. Animals also produce waxes that serve as protective coatings, keeping the surfaces of feathers, skin, and hair pliable and water repellent. In fact, if the waxy coating on the feathers of a water bird is dissolved as a result of the bird swimming in an oil slick, the feathers become wet and heavy, and the bird, unable to maintain its buoyancy, drowns.



Myricyl cerotate
(found in carnauba wax)

Summary

Fatty acids are carboxylic acids that are the structural components of many lipids. They may be saturated or unsaturated. Most fatty acids are unbranched and contain an even number of carbon atoms. Unsaturated fatty acids have lower melting points than saturated fatty acids containing the same number of carbon atoms.

Learning Objectives

- Identify the distinguishing characteristics of membrane lipids.
- Describe membrane components and how they are arranged.

All living cells are surrounded by a cell membrane. Plant cells (Figure 5.24.1*A*) and animal cells (Figure 5.24.1*B*) contain a cell nucleus that is also surrounded by a membrane and holds the genetic information for the cell. Everything between the cell membrane and the nuclear membrane—including intracellular fluids and various subcellular components such as the mitochondria and ribosomes—is called the cytoplasm. The membranes of all cells have a fundamentally similar structure, but membrane function varies tremendously from one organism to another and even from one cell to another within a single organism. This diversity arises mainly from the presence of different proteins and lipids in the membrane.




Figure **5.24.1**: (A) An Idealized Plant Cell. Not all the structures shown here occur in every type of plant cell. (B) An Idealized Animal Cell. The structures shown here will seldom all be found in a single animal cell.

The lipids in cell membranes are highly polar but have dual characteristics: part of the lipid is ionic and therefore dissolves in water, whereas the rest has a hydrocarbon structure and therefore dissolves in nonpolar substances. Often, the ionic part is referred to as hydrophilic, meaning “water loving,” and the nonpolar part as hydrophobic, meaning “water fearing” (repelled by water). When allowed to float freely in water, polar lipids spontaneously cluster together in any one of three arrangements: micelles, monolayers, and bilayers (Figure **5.24.2**).




Figure 5.24.2: Spontaneously Formed Polar Lipid Structures in Water: Monolayer, Micelle, and Bilayer

Micelles are aggregations in which the lipids' hydrocarbon tails—being hydrophobic—are directed toward the center of the assemblage and away from the surrounding water while the hydrophilic heads are directed outward, in contact with the water. Each micelle may contain thousands of lipid molecules. Polar lipids may also form a monolayer, a layer one molecule thick on the surface of the water. The polar heads face into water, and the nonpolar tails stick up into the air. Bilayers are double layers of lipids arranged so that the hydrophobic tails are sandwiched between an inner surface and an outer surface consisting of hydrophilic heads. The hydrophilic heads are in contact with water on either side of the bilayer, whereas the tails, sequestered inside the bilayer, are prevented from having contact with the water. Bilayers like this make up every cell membrane (Figure 5.24.3).




Figure 5.24.3: Schematic Diagram of a Cell Membrane. The membrane enclosing a typical animal cell is a phospholipid bilayer with embedded cholesterol and protein molecules. Short oligosaccharide chains are attached to the outer surface.

In the bilayer interior, the hydrophobic tails (that is, the fatty acid portions of lipid molecules) interact by means of dispersion forces. The interactions are weakened by the presence of unsaturated fatty acids. As a result, the membrane components are free to mill about to some extent, and the membrane is described as fluid.

The lipids found in cell membranes can be categorized in various ways. Phospholipids are lipids containing phosphorus. Glycolipids are sugar-containing lipids. The latter are found exclusively on the outer surface of the cell membrane, acting as distinguishing surface markers for the cell and thus serving in cellular recognition and cell-to-cell communication. Sphingolipids are phospholipids or glycolipids that contain the unsaturated amino alcohol sphingosine rather than glycerol. Diagrammatic structures of representative membrane lipids are presented in Figure 5.24.4.




Figure **5.24.4**: Component Structures of Some Important Membrane Lipids

Phosphoglyceride are composed of alcohol, phosphate, glycerol and 2 fatty acids. A sphingomyelin contains phosphate, sphingosine, and a fatty acid. A cerbroside contains sugar, sphingosine and a fatty acid.

Phosphoglycerides (also known as glycerophospholipids) are the most abundant phospholipids in cell membranes. They consist of a glycerol unit with fatty acids attached to the first two carbon atoms, while a phosphoric acid unit, esterified with an alcohol molecule (usually an amino alcohol, as in part (a) of Figure **5.24.5**) is attached to the third carbon atom of glycerol (part (b) of Figure **5.24.5**). Notice that the phosphoglyceride molecule is identical to a triglyceride up to the phosphoric acid unit (part (b) of Figure **5.24.5**).





Figure **5.24.5**: Phosphoglycerides. (a) Amino alcohols are commonly found in phosphoglycerides, which are evident in its structural formula (b).

Structural formula of ethanolamine and choline are shown. The structural formula of a phosphoglyceride is shown with the glycerol unit, phosphoric unit, and amino alcohol unit highlighted in different colors.

There are two common types of phosphoglycerides. Phosphoglycerides containing ethanolamine as the amino alcohol are called *phosphatidylethanolamines* or *cephalins*. Cephalins are found in brain tissue and nerves and also have a role in blood clotting. Phosphoglycerides containing choline as the amino alcohol unit are called *phosphatidylcholines* or *lecithins*. Lecithins occur in all living organisms. Like cephalins, they are important constituents of nerve and brain tissue. Egg yolks are especially rich in lecithins. Commercial-grade lecithins isolated from soybeans are widely used in foods as emulsifying agents. An emulsifying agent

is used to stabilize an emulsion—a dispersion of two liquids that do not normally mix, such as oil and water. Many foods are emulsions. Milk is an emulsion of butterfat in water. The emulsifying agent in milk is a protein called *casein*. Mayonnaise is an emulsion of salad oil in water, stabilized by lecithins present in egg yolk.



Sphingomyelins, the simplest sphingolipids, each contain a fatty acid, a phosphoric acid, sphingosine, and choline (Figure **5.24.6**). Because they contain phosphoric acid, they are also classified as phospholipids. Sphingomyelins are important constituents of the myelin sheath surrounding the axon of a nerve cell. Multiple sclerosis is one of several diseases resulting from damage to the myelin sheath.




Figure **5.24.6**: Sphingolipids. (a) Sphingosine, an amino alcohol, is found in all sphingolipids. (b) A sphingomyelin is also known as a phospholipid, as evidenced by the phosphoric acid unit in its structure.

Structural formula of sphingosine is shown. The general structure of a sphingolipid is shown with the sphingosine unit, fatty acid unit, phosphoric acid unit, and choline unit highlighted in different colors.

Most animal cells contain sphingolipids called cerebrosides (Figure **5.24.7**). Cerebrosides are composed of sphingosine, a fatty acid, and galactose or glucose. They therefore resemble sphingomyelins but have a sugar unit in place of the choline phosphate group. Cerebrosides are important constituents of the membranes of nerve and brain cells.




Figure 5.24.7: Cerebrosides. Cerebrosides are sphingolipids that contain a sugar unit.
General structure of a cerebroside with its sugar unit, sphingosine unit, and fatty acid unit highlighted in different colors.

The sphingolipids called gangliosides are more complex, usually containing a branched chain of three to eight monosaccharides and/or substituted sugars. Because of considerable variation in their sugar components, about 130 varieties of gangliosides have been identified. Most cell-to-cell recognition and communication processes (e.g., blood group antigens) depend on differences in the sequences of sugars in these compounds. Gangliosides are most prevalent in the outer membranes of nerve cells, although they also occur in smaller quantities in the outer membranes of most other cells. Because cerebrosides and gangliosides contain sugar groups, they are also classified as glycolipids.

Membrane Proteins

If membranes were composed only of lipids, very few ions or polar molecules could pass through their hydrophobic “sandwich filling” to enter or leave any cell. However, certain charged and polar species do cross the membrane, aided by proteins that move about in the lipid bilayer. The two major classes of proteins in the cell membrane are integral proteins, which span the hydrophobic interior of the bilayer, and peripheral proteins, which are more loosely associated with the surface of the lipid bilayer (Figure 5.24.3). Peripheral proteins may be attached to integral proteins, to the polar head groups of phospholipids, or to both by hydrogen bonding and electrostatic forces.

Small ions and molecules soluble in water enter and leave the cell by way of channels through the integral proteins. Some proteins, called *carrier proteins*, facilitate the passage of certain molecules, such as hormones and neurotransmitters, by specific interactions between the protein and the molecule being transported.

Summary

Lipids are important components of biological membranes. These lipids have dual characteristics: part of the molecule is hydrophilic, and part of the molecule is hydrophobic. Membrane lipids may be classified as phospholipids, glycolipids, and/or sphingolipids. Proteins are another important component of biological membranes. Integral proteins span the lipid bilayer, while peripheral proteins are more loosely associated with the surface of the membrane.

5.24: Lipids is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **17.0: Prelude to Lipids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **17.1: Fatty Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **17.3: Membranes and Membrane Lipids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


5.25: Proteins

Learning Objectives


- To recognize amino acids and classify them based on the characteristics of their side chains.

The proteins in all living species, from bacteria to humans, are constructed from the same set of 20 amino acids, so called because each contains an amino group attached to a carboxylic acid. The amino acids in proteins are α-amino acids, which means the amino group is attached to the α-carbon of the carboxylic acid. Humans can synthesize only about half of the needed amino acids; the remainder must be obtained from the diet and are known as essential amino acids. However, two additional amino acids have been found in limited quantities in proteins: Selenocysteine was discovered in 1986, while pyrrolysine was discovered in 2002.

The amino acids are colorless, nonvolatile, crystalline solids, melting and decomposing at temperatures above 200°C. These melting temperatures are more like those of inorganic salts than those of amines or organic acids and indicate that the structures of the amino acids in the solid state and in neutral solution are best represented as having both a negatively charged group and a positively charged group. Such a species is known as a zwitterion.



α-Amino acid drawn as a zwitterion























α-Amino acid drawn as an uncharged molecule; not an accurate respresentation of amino acid structure

Classification


In addition to the amino and carboxyl groups, amino acids have a side chain or R group attached to the α-carbon. Each amino acid has unique characteristics arising from the size, shape, solubility, and ionization properties of its R group. As a result, the side chains of amino acids exert a profound effect on the structure and biological activity of proteins. Although amino acids can be classified in various ways, one common approach is to classify them according to whether the functional group on the side chain at neutral pH is nonpolar, polar but uncharged, negatively charged, or positively charged. The structures and names of the 20 amino acids, their one- and three-letter abbreviations, and some of their distinctive features are given in Table **5.25.1**.

Table **5.25.1**: Common Amino Acids Found in Proteins

Common Name	Abbreviation	Structural Formula (at pH 6)	Molar Mass	Distinctive Feature
Amino acids with a nonpolar R group				
glycine	gly (G)		75	the only amino acid lacking a chiral carbon
alanine	ala (A)		89	—
valine	val (V)		117	a branched-chain amino acid
leucine	leu (L)		131	a branched-chain amino acid
isoleucine	ile (I)		131	an essential amino acid because most animals cannot synthesize branched-chain amino acids
phenylalanine	phe (F)		165	also classified as an aromatic amino acid
tryptophan	trp (W)		204	also classified as an aromatic amino acid
methionine	met (M)		149	side chain functions as a methyl group donor
proline	pro (P)		115	contains a secondary amine group; referred to as an *α-imino acid*
Amino acids with a polar but neutral R group				
serine	ser (S)		105	found at the active site of many enzymes
threonine	thr (T)		119	named for its similarity to the sugar threose
cysteine	cys (C)		121	oxidation of two cysteine molecules yields *cystine*


Common Name	Abbreviation	Structural Formula (at pH 6)	Molar Mass	Distinctive Feature
tyrosine	tyr (Y)		181	also classified as an aromatic amino acid
asparagine	asn (N)		132	the amide of aspartic acid
glutamine	gln (Q)		146	the amide of glutamic acid
Amino acids with a negatively charged R group				
aspartic acid	asp (D)		132	carboxyl groups are ionized at physiological pH; also known as aspartate
glutamic acid	glu (E)		146	carboxyl groups are ionized at physiological pH; also known as glutamate
Amino acids with a positively charged R group				
histidine	his (H)		155	the only amino acid whose R group has a pK_a (6.0) near physiological pH
lysine	lys (K)		147	—
arginine	arg (R)		175	almost as strong a base as sodium hydroxide

The first amino acid to be isolated was asparagine in 1806. It was obtained from protein found in asparagus juice (hence the name). Glycine, the major amino acid found in gelatin, was named for its sweet taste (Greek *glykys*, meaning “sweet”). In some cases an amino acid found in a protein is actually a derivative of one of the common 20 amino acids (one such derivative is hydroxyproline). The modification occurs *after* the amino acid has been assembled into a protein.



Configuration

Notice in Table 5.25.1 that glycine is the only amino acid whose α-carbon is *not* chiral. Therefore, with the exception of glycine, the amino acids could theoretically exist in either the D- or the L-enantiomeric form and rotate plane-polarized light. As with sugars, chemists used L-glyceraldehyde as the reference compound for the assignment of absolute configuration to amino acids. Its structure closely resembles an amino acid structure except that in the latter, an amino group takes the place of the OH group on the chiral carbon of the L-glyceraldehyde and a carboxylic acid replaces the aldehyde. Modern stereochemistry assignments using the Cahn-Ingold-Prelog priority rules used ubiquitously in chemistry show that all of the naturally occurring chiral amino acids are S except Cys which is R.



We learned that all naturally occurring sugars belong to the D series. It is interesting, therefore, that nearly all known plant and animal proteins are composed entirely of L-amino acids. However, certain bacteria contain D-amino acids in their cell walls, and several antibiotics (e.g., actinomycin D and the gramicidins) contain varying amounts of D-leucine, D-phenylalanine, and D-valine.

Summary

Amino acids can be classified based on the characteristics of their distinctive side chains as nonpolar, polar but uncharged, negatively charged, or positively charged. The amino acids found in proteins are L-amino acids.

Learning Objectives

- Describe the four levels of protein structure.
- Identify the types of attractive interactions that hold proteins in their most stable three-dimensional structure.
- Explain what happens when proteins are denatured.
- Identify how a protein can be denatured.

Each of the thousands of naturally occurring proteins has its own characteristic amino acid composition and sequence that result in a unique three-dimensional shape. Since the 1950s, scientists have determined the amino acid sequences and three-dimensional conformation of numerous proteins and thus obtained important clues on how each protein performs its specific function in the body.

Proteins are compounds of high molar mass consisting largely or entirely of chains of amino acids. Because of their great complexity, protein molecules cannot be classified on the basis of specific structural similarities, as carbohydrates and lipids are categorized. The two major structural classifications of proteins are based on far more general qualities: whether the protein is (1) fiberlike and insoluble or (2) globular and soluble. Some proteins, such as those that compose hair, skin, muscles, and connective tissue, are fiberlike. These fibrous proteins are insoluble in water and usually serve structural, connective, and protective functions. Examples of fibrous proteins are keratins, collagens, myosins, and elastins. Hair and the outer layer of skin are composed of keratin. Connective tissues contain collagen. Myosins are muscle proteins and are capable of contraction and extension. Elastins are found in ligaments and the elastic tissue of artery walls.

Globular proteins, the other major class, are soluble in aqueous media. In these proteins, the chains are folded so that the molecule as a whole is roughly spherical. Familiar examples include egg albumin from egg whites and serum albumin in blood. Serum albumin plays a major role in transporting fatty acids and maintaining a proper balance of osmotic pressures in the body. Hemoglobin and myoglobin, which are important for binding oxygen, are also globular proteins.




Figure 5.25.1: Primary Structure of Human Insulin. Human insulin, whose amino acid sequence is shown here, is a hormone that is required for the proper metabolism of glucose.

Levels of Protein Structure

The structure of proteins is generally described as having four organizational levels. The first of these is the primary structure, which is the number and sequence of amino acids in a protein's polypeptide chain or chains, beginning with the free amino group and maintained by the peptide bonds connecting each amino acid to the next. The primary structure of insulin, composed of 51 amino acids, is shown in Figure 5.25.1.

A protein molecule is not a random tangle of polypeptide chains. Instead, the chains are arranged in unique but specific conformations. The term secondary structure refers to the fixed arrangement of the polypeptide backbone. On the basis of X ray studies, Linus Pauling and Robert Corey postulated that certain proteins or portions of proteins twist into a spiral or a helix. This helix is stabilized by *intrachain* hydrogen bonding between the carbonyl oxygen atom of one amino acid and the amide hydrogen atom four amino acids up the chain (located on the next turn of the helix) and is known as a right-handed α-helix. X ray data indicate that this helix makes one turn for every 3.6 amino acids, and the side chains of these amino acids project outward from the coiled backbone (Figure 5.25.2). The α-keratins, found in hair and wool, are exclusively α-helical in conformation. Some proteins, such as gamma globulin, chymotrypsin, and cytochrome c, have little or no helical structure. Others, such as hemoglobin and myoglobin, are helical in certain regions but not in others.




Figure **5.25.2** A Ball-and-Stick Model of an α-Helix. This ball-and-stick model shows the intrachain hydrogen bonding between carbonyl oxygen atoms and amide hydrogen atoms. Each turn of the helix spans 3.6 amino acids. Note that the side chains (represented as green spheres) point out from the helix.

Another common type of secondary structure, called the *β-pleated sheet conformation*, is a sheetlike arrangement in which two or more extended polypeptide chains (or separate regions on the same chain) are aligned side by side. The aligned segments can run either parallel or antiparallel—that is, the N-terminals can face in the same direction on adjacent chains or in different directions—and are connected by *interchain* hydrogen bonding (Figure **5.25.3**). The β-pleated sheet is particularly important in structural proteins, such as silk fibroin. It is also seen in portions of many enzymes, such as carboxypeptidase A and lysozyme.




Figure **5.25.3**: A Ball-and-Stick Model of the β-Pleated Sheet Structure in Proteins. The side chains extend above or below the sheet and alternate along the chain. The protein chains are held together by interchain hydrogen bonding.

Tertiary structure refers to the unique three-dimensional shape of the protein as a whole, which results from the folding and bending of the protein backbone. The tertiary structure is intimately tied to the proper biochemical functioning of the protein. Figure **5.25.4** shows a depiction of the three-dimensional structure of insulin.





Figure **5.25.4**: A Ribbon Model of the Three-Dimensional Structure of Insulin. The spiral regions represent sections of the polypeptide chain that have an α-helical structure, while the broad arrows represent β-pleated sheet structures.

Four major types of attractive interactions determine the shape and stability of the tertiary structure of proteins. You studied several of them previously.

1. Ionic bonding. Ionic bonds result from electrostatic attractions between positively and negatively charged side chains of amino acids. For example, the mutual attraction between an aspartic acid carboxylate ion and a lysine ammonium ion helps to maintain a particular folded area of a protein (part (a) of Figure **5.25.5**).
2. Hydrogen bonding. Hydrogen bonding forms between a highly electronegative oxygen atom or a nitrogen atom and a hydrogen atom attached to another oxygen atom or a nitrogen atom, such as those found in polar amino acid side chains. Hydrogen bonding (as well as ionic attractions) is extremely important in both the intra- and intermolecular interactions of proteins (part (b) of Figure **5.25.5**).
3. Disulfide linkages. Two cysteine amino acid units may be brought close together as the protein molecule folds. Subsequent oxidation and linkage of the sulfur atoms in the highly reactive sulfhydryl (SH) groups leads to the formation of cystine (part (c) of Figure **5.25.5**). Intrachain disulfide linkages are found in many proteins, including insulin (yellow bars in Figure **5.25.1**) and have a strong stabilizing effect on the tertiary structure.



4. Dispersion forces. Dispersion forces arise when a normally nonpolar atom becomes momentarily polar due to an uneven distribution of electrons, leading to an instantaneous dipole that induces a shift of electrons in a neighboring nonpolar atom. Dispersion forces are weak but can be important when other types of interactions are either missing or minimal (part (d) of Figure **5.25.5**). This is the case with fibroin, the major protein in silk, in which a high proportion of amino acids in the protein have nonpolar side chains. The term *hydrophobic interaction* is often misused as a synonym for dispersion forces. Hydrophobic interactions arise because water molecules engage in hydrogen bonding with other water molecules (or groups in proteins capable of hydrogen bonding). Because nonpolar groups cannot engage in hydrogen bonding, the protein folds in such a way that these groups are buried in the interior part of the protein structure, minimizing their contact with water.




Figure 5.25.5: Tertiary Protein Structure Interactions. Four interactions stabilize the tertiary structure of a protein: (a) ionic bonding, (b) hydrogen bonding, (c) disulfide linkages, and (d) dispersion forces.

When a protein contains more than one polypeptide chain, each chain is called a *subunit*. The arrangement of multiple subunits represents a fourth level of structure, the quaternary structure of a protein. Hemoglobin, with four polypeptide chains or subunits, is the most frequently cited example of a protein having quaternary structure (Figure 5.25.6). The quaternary structure of a protein is produced and stabilized by the same kinds of interactions that produce and maintain the tertiary structure. A schematic representation of the four levels of protein structure is in Figure 5.25.7.




Figure 5.25.6 The Quaternary Structure of Hemoglobin. Hemoglobin is a protein that transports oxygen throughout the body.

Source: Image from the RCSB PDB (www.pdb.org(opens in new window)) of PDB ID 1I3D (R.D. Kidd, H.M. Baker, A.J. Mathews, T. Brittain, E.N. Baker (2001) Oligomerization and ligand binding in a homotetrameric hemoglobin: two high-resolution crystal structures of hemoglobin Bart's (gamma(4)), a marker for alpha-thalassemia. Protein Sci. 1739–1749).




Figure **5.25.7**: Levels of Structure in Proteins

The *primary structure* consists of the specific amino acid sequence. The resulting peptide chain can twist into an α-helix, which is one type of *secondary structure*. This helical segment is incorporated into the *tertiary structure* of the folded polypeptide chain. The single polypeptide chain is a subunit that constitutes the *quaternary structure* of a protein, such as hemoglobin that has four polypeptide chains.

Denaturation of Proteins

The highly organized structures of proteins are truly masterworks of chemical architecture. But highly organized structures tend to have a certain delicacy, and this is true of proteins. Denaturation is the term used for any change in the three-dimensional structure of a protein that renders it incapable of performing its assigned function. A denatured protein cannot do its job. (Sometimes denaturation is equated with the precipitation or coagulation of a protein; our definition is a bit broader.) A wide variety of reagents and conditions, such as heat, organic compounds, pH changes, and heavy metal ions can cause protein denaturation (Figure **5.25.1**).

Figure **5.25.1**: Protein Denaturation Methods

Method	Effect on Protein Structure
Heat above 50°C or ultraviolet (UV) radiation	Heat or UV radiation supplies kinetic energy to protein molecules, causing their atoms to vibrate more rapidly and disrupting relatively weak hydrogen bonding and dispersion forces.
Use of organic compounds, such as ethyl alcohol	These compounds are capable of engaging in intermolecular hydrogen bonding with protein molecules, disrupting intramolecular hydrogen bonding within the protein.
Salts of heavy metal ions, such as mercury, silver, and lead	These ions form strong bonds with the carboxylate anions of the acidic amino acids or SH groups of cysteine, disrupting ionic bonds and disulfide linkages.
Alkaloid reagents, such as tannic acid (used in tanning leather)	These reagents combine with positively charged amino groups in proteins to disrupt ionic bonds.

Anyone who has fried an egg has observed denaturation. The clear egg white turns opaque as the albumin denatures and coagulates. No one has yet reversed that process. However, given the proper circumstances and enough time, a protein that has unfolded under sufficiently gentle conditions can refold and may again exhibit biological activity (Figure **5.25.8**). Such evidence suggests that, at least for these proteins, the primary structure determines the secondary and tertiary structure. A given sequence of amino acids seems to adopt its particular three-dimensional arrangement naturally if conditions are right.




Figure **5.25.8**: Denaturation and Renaturation of a Protein. The denaturation (unfolding) and renaturation (refolding) of a protein is depicted. The red boxes represent stabilizing interactions, such as disulfide linkages, hydrogen bonding, and/or ionic bonds.

The primary structures of proteins are quite sturdy. In general, fairly vigorous conditions are needed to hydrolyze peptide bonds. At the secondary through quaternary levels, however, proteins are quite vulnerable to attack, though they vary in their vulnerability to denaturation. The delicately folded globular proteins are much easier to denature than are the tough, fibrous proteins of hair and skin.

Summary

Proteins can be divided into two categories: fibrous, which tend to be insoluble in water, and globular, which are more soluble in water. A protein may have up to four levels of structure. The primary structure consists of the specific amino acid sequence. The resulting peptide chain can form an α-helix or β-pleated sheet (or local structures not as easily categorized), which is known as secondary structure. These segments of secondary structure are incorporated into the tertiary structure of the folded polypeptide chain. The quaternary structure describes the arrangements of subunits in a protein that contains more than one subunit. Four major types of attractive interactions determine the shape and stability of the folded protein: ionic bonding, hydrogen bonding, disulfide linkages, and dispersion forces. A wide variety of reagents and conditions can cause a protein to unfold or denature.

5.25: Proteins is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **18.1: Properties of Amino Acids** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.
- **18.4: Proteins** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

Index

A

acid–base titration
5.12: Acid–Base Titration
addition reaction
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
Aliphatic hydrocarbons
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
alkanes
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
alkenes
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
alkynes
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
anion
2.14: Ions - Losing and Gaining Electrons
aromatic
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
aromatic hydrocarbons
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
atomic mass unit
2.11: The Properties of Protons, Neutrons, and Electrons
atomic theory
2.9: Indivisible - The Atomic Theory
autoionization of water
5.14: Water - Acid and Base in One

B

balanced chemical equation
3.18: Making Molecules- Mole-to-Mole Conversions
Balancing a Chemical Equation
3.3: How to Write Balanced Chemical Equations
Bohr model
4.1: The Bohr Model - Atoms with Orbits
Boyle’s law
4.14: Boyle’s Law - Pressure and Volume

C

cation
2.14: Ions - Losing and Gaining Electrons
Charles's Law
4.15: Charles’s Law- Volume and Temperature
chemical change
2.6: Changes in Matter - Physical and Chemical Changes
chemical property
2.5: Differences in Matter- Physical and Chemical Properties
coefficient
3.18: Making Molecules- Mole-to-Mole Conversions
Coefficients and Subscripts
3.3: How to Write Balanced Chemical Equations
combined gas law
4.17: The Combined Gas Law- Pressure, Volume, and Temperature
complete ionic equation
3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
compound
2.4: Classifying Matter According to Its Composition
Condensed Structural Formula
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
conservation of mass
2.7: Conservation of Mass - There is No New Matter
crisscross method
2.21: Writing Formulas for Ionic Compounds

D

Dalton’s law of partial pressures
4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen
democritus
2.8: Cutting Aluminum until you get Atoms
2.9: Indivisible - The Atomic Theory
density
1.12: Density
dimensional analysis
1.10: Problem Solving and Unit Conversions

E

electron
2.11: The Properties of Protons, Neutrons, and Electrons
electronegativity
4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix
element
2.4: Classifying Matter According to Its Composition
empirical formula
3.16: Calculating Molecular Formulas for Compounds
empirical formula mass
3.16: Calculating Molecular Formulas for Compounds
Exceptions to the Octet Rule
4.8: Writing Lewis Structures for Covalent Compounds

F

formula unit
3.20: Stoichiometry

H

hydrocarbons
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
hydrogenation reaction
5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon
hydrolysis
5.11: Reactions of Acids and Bases
hypothesis
1.3: Hypothesis, Theories, and Laws

I

ions
2.14: Ions - Losing and Gaining Electrons
isotopes
2.15: Isotopes - When the Number of Neutrons Varies

L

law
1.3: Hypothesis, Theories, and Laws

M

matter
2.2: What is Matter?
mixture
2.4: Classifying Matter According to Its Composition
molarity
5.5: Specifying Solution Concentration- Molarity
mole ratio
3.20: Stoichiometry
molecular formula
3.16: Calculating Molecular Formulas for Compounds

N

net ionic equation
3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations
neutralization reaction
5.11: Reactions of Acids and Bases
neutron
2.11: The Properties of Protons, Neutrons, and Electrons

P

pascal (unit)
4.13: Pressure - The Result of Constant Molecular Collisions
pH scale
5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity
physical change
2.6: Changes in Matter - Physical and Chemical Changes
physical property
2.5: Differences in Matter- Physical and Chemical Properties
pOH
5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity
polarity
4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix
precipitate
3.6: Precipitation Reactions
precipitation reaction
3.6: Precipitation Reactions
proton
2.11: The Properties of Protons, Neutrons, and Electrons

R

relative abundances

2.16: Atomic Mass - The Average Mass of an Element’s Atoms

resonance

4.9: Resonance - Equivalent Lewis Structures for the Same Molecule

rounding

1.9: Significant Figures in Calculations

S

saturated

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

saturated hydrocarbons

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

scienfitic notation

1.7: Scientific Notation - Writing Large and Small Numbers

scientific method

1.2: The Scientific Method - How Chemists Think

Separation of Mixtures

2.6: Changes in Matter - Physical and Chemical Changes

significant figures

1.8: Significant Figures - Writing Numbers to Reflect Precision

1.9: Significant Figures in Calculations

solute

5.2: Solutions - Homogeneous Mixtures

solution stoichiometry

5.6: Solution Stoichiometry

solvent

5.2: Solutions - Homogeneous Mixtures

spectator ions

3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations

Stability of Isotopes

2.15: Isotopes - When the Number of Neutrons Varies

stock system

2.23: Naming Ionic Compounds

stoichiometry

3.20: Stoichiometry

structural formulas

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

T

temperature

1.13: Temperature - Random Motion of Molecules and Atoms

theory

1.3: Hypothesis, Theories, and Laws

U

unit conversions

1.10: Problem Solving and Unit Conversions

unsaturated

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

unsaturated hydrocarbons

5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon

V

valence shell electron pair repulsion theory

4.10: Predicting the Shapes of Molecules

VSEPR

4.10: Predicting the Shapes of Molecules

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: CHEM 305: Introduction to Chemistry (Zumalt)

Webpages: 122

Applicable Restrictions: Noncommercial

All licenses found:

- CK-12 License: 59% (72 pages)
- Undeclared: 20.5% (25 pages)
- CC BY-NC-SA 3.0: 12.3% (15 pages)
- CC BY-SA 4.0: 3.3% (4 pages)
- Unknown License: 2.5% (3 pages)
- CC BY 4.0: 0.8% (1 page)
- CC BY-SA 3.0: 0.8% (1 page)
- Public Domain: 0.8% (1 page)
- Unknown License: 0% (0 page)
- Unknown License: 0% (0 page)

By Page

- CHEM 305: Introduction to Chemistry (Zumalt) - *CC BY-SA 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Unit 1 - *Undeclared*
    - 1.1: The Scope of Chemistry - *CK-12 License*
    - 1.2: The Scientific Method - How Chemists Think - *CK-12 License*
    - 1.3: Hypothesis, Theories, and Laws - *CK-12 License*
    - 1.4: Taking Measurements - *CC BY-NC-SA 3.0*
    - 1.5: The Basic Units of Measurement - *CK-12 License*
    - 1.6: Units Raised to a Power - *CK-12 License*
    - 1.7: Scientific Notation - Writing Large and Small Numbers - *CK-12 License*
    - 1.8: Significant Figures - Writing Numbers to Reflect Precision - *CK-12 License*
    - 1.9: Significant Figures in Calculations - *CK-12 License*
    - 1.10: Problem Solving and Unit Conversions - *CC BY-NC-SA 3.0*
    - 1.11: Solving Multi-step Conversion Problems - *CC BY-NC-SA 3.0*
    - 1.12: Density - *CK-12 License*
    - 1.13: Temperature - Random Motion of Molecules and Atoms - *CK-12 License*
    - 1.E: Measurement and Problem Solving (Exercises) - *CK-12 License*
  - 2: Unit 2 - *Undeclared*
    - 2.1: In Your Room - *CK-12 License*
    - 2.2: What is Matter? - *CK-12 License*
    - 2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas - *CK-12 License*
    - 2.4: Classifying Matter According to Its Composition - *CK-12 License*
    - 2.5: Differences in Matter- Physical and Chemical Properties - *CK-12 License*
    - 2.6: Changes in Matter - Physical and Chemical Changes - *CK-12 License*
    - 2.7: Conservation of Mass - There is No New Matter - *CK-12 License*
    - 2.8: Cutting Aluminum until you get Atoms - *CK-12 License*
    - 2.9: Indivisible - The Atomic Theory - *CK-12 License*
    - 2.10: The Nuclear Atom - *CK-12 License*
    - 2.11: The Properties of Protons, Neutrons, and Electrons - *CK-12 License*
    - 2.12: Elements- Defined by Their Number of Protons - *CK-12 License*
    - 2.13: Looking for Patterns - The Periodic Table - *CK-12 License*

- 2.14: Ions - Losing and Gaining Electrons - *CK-12 License*
- 2.15: Isotopes - When the Number of Neutrons Varies - *CK-12 License*
- 2.16: Atomic Mass - The Average Mass of an Element's Atoms - *CK-12 License*
- 2.17: Sugar and Salt - *CK-12 License*
- 2.18: Compounds Display Constant Composition - *CK-12 License*
- 2.19: Chemical Formulas - How to Represent Compounds - *CK-12 License*
- 2.20: A Molecular View of Elements and Compounds - *CK-12 License*
- 2.21: Writing Formulas for Ionic Compounds - *CK-12 License*
- 2.22: Nomenclature- Naming Compounds - *CK-12 License*
- 2.23: Naming Ionic Compounds - *CK-12 License*
- 2.24: Naming Molecular Compounds - *CK-12 License*
- 2.25: Naming Acids - *CK-12 License*
- 2.26: Nomenclature Summary - *CK-12 License*

3: Unit 3 - *Undeclared*

- 3.1: Evidence of a Chemical Reaction - *CC BY 4.0*
- 3.2: The Chemical Equation - *CK-12 License*
- 3.3: How to Write Balanced Chemical Equations - *CC BY-NC-SA 3.0*
- 3.4: Classifying Chemical Reactions - *CK-12 License*
- 3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water - *CK-12 License*
- 3.6: Precipitation Reactions - *CK-12 License*
- 3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations - *CK-12 License*
- 3.8: Prelude to Chemical Composition - How Much Sodium? - *CC BY-SA 4.0*
- 3.9: Counting Nails by the Pound - *CC BY-SA 3.0*
- 3.10: Counting Atoms by the Gram - *CK-12 License*
- 3.11: Counting Molecules by the Gram - *CC BY-NC-SA 3.0*
- 3.12: Chemical Formulas as Conversion Factors - *CK-12 License*
- 3.13: Mass Percent Composition of Compounds - *CK-12 License*
- 3.14: Mass Percent Composition from a Chemical Formula - *CK-12 License*
- 3.15: Calculating Empirical Formulas for Compounds - *CK-12 License*
- 3.16: Calculating Molecular Formulas for Compounds - *CK-12 License*
- 3.17: Making Pancakes- Relationships Between Ingredients - *Undeclared*
- 3.18: Making Molecules- Mole-to-Mole Conversions - *CC BY-NC-SA 3.0*
- 3.19: Making Molecules- Mass-to-Mass Conversions - *CC BY-NC-SA 3.0*
- 3.20: Stoichiometry - *CK-12 License*
- 3.21: Limiting Reactant and Theoretical Yield - *CC BY-NC-SA 3.0*
- 3.22: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants - *CC BY-NC-SA 3.0*

4: Unit 4 - *Undeclared*

- 4.1: The Bohr Model - Atoms with Orbits - *CK-12 License*
- 4.2: Quantum-Mechanical Orbitals and Electron Configurations - *CK-12 License*
- 4.3: Electron Configurations and the Periodic Table - *CK-12 License*
- 4.4: The Explanatory Power of the Quantum-Mechanical Model - *CK-12 License*
- 4.5: Representing Valence Electrons with Dots - *CC BY-NC-SA 3.0*
- 4.6: Lewis Structures of Ionic Compounds- Electrons Transferred - *CC BY-NC-SA 3.0*
- 4.7: Covalent Lewis Structures- Electrons Shared - *Unknown License*
- 4.8: Writing Lewis Structures for Covalent Compounds - *Unknown License*
- 4.9: Resonance - Equivalent Lewis Structures for the Same Molecule - *CK-12 License*
- 4.10: Predicting the Shapes of Molecules - *Unknown License*
- 4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix - *CK-12 License*
- 4.12: Kinetic Molecular Theory- A Model for Gases - *CC BY-NC-SA 3.0*
- 4.13: Pressure - The Result of Constant Molecular Collisions - *CK-12 License*
- 4.14: Boyle's Law - Pressure and Volume - *CK-12 License*
- 4.15: Charles's Law- Volume and Temperature - *CK-12 License*
- 4.16: Gay-Lussac's Law- Temperature and Pressure - *CK-12 License*
- 4.17: The Combined Gas Law- Pressure, Volume, and Temperature - *CK-12 License*

- 4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles - *CK-12 License*
- 4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen - *CK-12 License*

5: Unit 5 - *Undeclared*

- 5.1: Intermolecular Forces- Dispersion, Dipole–Dipole, Hydrogen Bonding, and Ion-Dipole - *CC BY-NC-SA 3.0*
- 5.2: Solutions - Homogeneous Mixtures - *CC BY-NC-SA 3.0*
- 5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy - *CK-12 License*
- 5.4: Specifying Solution Concentration- Mass Percent - *CK-12 License*
- 5.5: Specifying Solution Concentration- Molarity - *CK-12 License*
- 5.6: Solution Stoichiometry - *CC BY-SA 4.0*
- 5.7: Sour Patch Kids and International Spy Movies - *CC BY-SA 4.0*
- 5.8: Acids- Properties and Examples - *CK-12 License*
- 5.9: Bases- Properties and Examples - *CK-12 License*
- 5.10: Molecular Definitions of Acids and Bases - *CK-12 License*
- 5.11: Reactions of Acids and Bases - *Public Domain*
- 5.12: Acid–Base Titration - *CK-12 License*
- 5.13: Strong and Weak Acids and Bases - *CK-12 License*
- 5.14: Water - Acid and Base in One - *CK-12 License*
- 5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity - *CK-12 License*
- 5.16: Carbon - A Versitile Atom - *Undeclared*
- 5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon - *CC BY-NC-SA 3.0*
- 5.18: Alkanes- Saturated Hydrocarbons - *Undeclared*
- 5.19: Isomers- Same Formula, Different Structure - *Undeclared*
- 5.20: Naming Alkanes - *Undeclared*
- 5.21: Alkenes and Alkynes - *Undeclared*
- 5.22: Functional Groups - *Undeclared*
- 5.23: Carbohydrates - *Undeclared*
- 5.24: Lipids - *Undeclared*
- 5.25: Proteins - *Undeclared*

Back Matter - *Undeclared*

- Index - *Undeclared*
- Glossary - *Undeclared*
- Detailed Licensing - *Undeclared*
- Detailed Licensing - *Undeclared*

Detailed Licensing

Overview

Title: CHEM 305: Introduction to Chemistry (Zumalt)

Webpages: 122

Applicable Restrictions: Noncommercial

All licenses found:

- CK-12 License: 59% (72 pages)
- Undeclared: 20.5% (25 pages)
- CC BY-NC-SA 3.0: 12.3% (15 pages)
- CC BY-SA 4.0: 3.3% (4 pages)
- Unknown License: 2.5% (3 pages)
- CC BY 4.0: 0.8% (1 page)
- CC BY-SA 3.0: 0.8% (1 page)
- Public Domain: 0.8% (1 page)
- Unknown License: 0% (0 page)
- Unknown License: 0% (0 page)

By Page

- CHEM 305: Introduction to Chemistry (Zumalt) - *CC BY-SA 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Unit 1 - *Undeclared*
    - 1.1: The Scope of Chemistry - *CK-12 License*
    - 1.2: The Scientific Method - How Chemists Think - *CK-12 License*
    - 1.3: Hypothesis, Theories, and Laws - *CK-12 License*
    - 1.4: Taking Measurements - *CC BY-NC-SA 3.0*
    - 1.5: The Basic Units of Measurement - *CK-12 License*
    - 1.6: Units Raised to a Power - *CK-12 License*
    - 1.7: Scientific Notation - Writing Large and Small Numbers - *CK-12 License*
    - 1.8: Significant Figures - Writing Numbers to Reflect Precision - *CK-12 License*
    - 1.9: Significant Figures in Calculations - *CK-12 License*
    - 1.10: Problem Solving and Unit Conversions - *CC BY-NC-SA 3.0*
    - 1.11: Solving Multi-step Conversion Problems - *CC BY-NC-SA 3.0*
    - 1.12: Density - *CK-12 License*
    - 1.13: Temperature - Random Motion of Molecules and Atoms - *CK-12 License*
    - 1.E: Measurement and Problem Solving (Exercises) - *CK-12 License*
  - 2: Unit 2 - *Undeclared*
    - 2.1: In Your Room - *CK-12 License*
    - 2.2: What is Matter? - *CK-12 License*
    - 2.3: Classifying Matter According to Its State—Solid, Liquid, and Gas - *CK-12 License*
    - 2.4: Classifying Matter According to Its Composition - *CK-12 License*
    - 2.5: Differences in Matter- Physical and Chemical Properties - *CK-12 License*
    - 2.6: Changes in Matter - Physical and Chemical Changes - *CK-12 License*
    - 2.7: Conservation of Mass - There is No New Matter - *CK-12 License*
    - 2.8: Cutting Aluminum until you get Atoms - *CK-12 License*
    - 2.9: Indivisible - The Atomic Theory - *CK-12 License*
    - 2.10: The Nuclear Atom - *CK-12 License*
    - 2.11: The Properties of Protons, Neutrons, and Electrons - *CK-12 License*
    - 2.12: Elements- Defined by Their Number of Protons - *CK-12 License*
    - 2.13: Looking for Patterns - The Periodic Table - *CK-12 License*

- 2.14: Ions - Losing and Gaining Electrons - *CK-12 License*
- 2.15: Isotopes - When the Number of Neutrons Varies - *CK-12 License*
- 2.16: Atomic Mass - The Average Mass of an Element's Atoms - *CK-12 License*
- 2.17: Sugar and Salt - *CK-12 License*
- 2.18: Compounds Display Constant Composition - *CK-12 License*
- 2.19: Chemical Formulas - How to Represent Compounds - *CK-12 License*
- 2.20: A Molecular View of Elements and Compounds - *CK-12 License*
- 2.21: Writing Formulas for Ionic Compounds - *CK-12 License*
- 2.22: Nomenclature- Naming Compounds - *CK-12 License*
- 2.23: Naming Ionic Compounds - *CK-12 License*
- 2.24: Naming Molecular Compounds - *CK-12 License*
- 2.25: Naming Acids - *CK-12 License*
- 2.26: Nomenclature Summary - *CK-12 License*

3: Unit 3 - *Undeclared*

- 3.1: Evidence of a Chemical Reaction - *CC BY 4.0*
- 3.2: The Chemical Equation - *CK-12 License*
- 3.3: How to Write Balanced Chemical Equations - *CC BY-NC-SA 3.0*
- 3.4: Classifying Chemical Reactions - *CK-12 License*
- 3.5: Aqueous Solutions and Solubility - Compounds Dissolved in Water - *CK-12 License*
- 3.6: Precipitation Reactions - *CK-12 License*
- 3.7: Writing Chemical Equations for Reactions in Solution- Molecular, Complete Ionic, and Net Ionic Equations - *CK-12 License*
- 3.8: Prelude to Chemical Composition - How Much Sodium? - *CC BY-SA 4.0*
- 3.9: Counting Nails by the Pound - *CC BY-SA 3.0*
- 3.10: Counting Atoms by the Gram - *CK-12 License*
- 3.11: Counting Molecules by the Gram - *CC BY-NC-SA 3.0*
- 3.12: Chemical Formulas as Conversion Factors - *CK-12 License*
- 3.13: Mass Percent Composition of Compounds - *CK-12 License*
- 3.14: Mass Percent Composition from a Chemical Formula - *CK-12 License*
- 3.15: Calculating Empirical Formulas for Compounds - *CK-12 License*
- 3.16: Calculating Molecular Formulas for Compounds - *CK-12 License*
- 3.17: Making Pancakes- Relationships Between Ingredients - *Undeclared*
- 3.18: Making Molecules- Mole-to-Mole Conversions - *CC BY-NC-SA 3.0*
- 3.19: Making Molecules- Mass-to-Mass Conversions - *CC BY-NC-SA 3.0*
- 3.20: Stoichiometry - *CK-12 License*
- 3.21: Limiting Reactant and Theoretical Yield - *CC BY-NC-SA 3.0*
- 3.22: Limiting Reactant, Theoretical Yield, and Percent Yield from Initial Masses of Reactants - *CC BY-NC-SA 3.0*

4: Unit 4 - *Undeclared*

- 4.1: The Bohr Model - Atoms with Orbits - *CK-12 License*
- 4.2: Quantum-Mechanical Orbitals and Electron Configurations - *CK-12 License*
- 4.3: Electron Configurations and the Periodic Table - *CK-12 License*
- 4.4: The Explanatory Power of the Quantum-Mechanical Model - *CK-12 License*
- 4.5: Representing Valence Electrons with Dots - *CC BY-NC-SA 3.0*
- 4.6: Lewis Structures of Ionic Compounds- Electrons Transferred - *CC BY-NC-SA 3.0*
- 4.7: Covalent Lewis Structures- Electrons Shared - *Unknown License*
- 4.8: Writing Lewis Structures for Covalent Compounds - *Unknown License*
- 4.9: Resonance - Equivalent Lewis Structures for the Same Molecule - *CK-12 License*
- 4.10: Predicting the Shapes of Molecules - *Unknown License*
- 4.11: Electronegativity and Polarity - Why Oil and Water Do not Mix - *CK-12 License*
- 4.12: Kinetic Molecular Theory- A Model for Gases - *CC BY-NC-SA 3.0*
- 4.13: Pressure - The Result of Constant Molecular Collisions - *CK-12 License*
- 4.14: Boyle's Law - Pressure and Volume - *CK-12 License*
- 4.15: Charles's Law- Volume and Temperature - *CK-12 License*
- 4.16: Gay-Lussac's Law- Temperature and Pressure - *CK-12 License*
- 4.17: The Combined Gas Law- Pressure, Volume, and Temperature - *CK-12 License*

- 4.18: The Ideal Gas Law- Pressure, Volume, Temperature, and Moles - *CK-12 License*
- 4.19: Mixtures of Gases - Why Deep-Sea Divers Breathe a Mixture of Helium and Oxygen - *CK-12 License*

5: Unit 5 - *Undeclared*

- 5.1: Intermolecular Forces- Dispersion, Dipole–Dipole, Hydrogen Bonding, and Ion-Dipole - *CC BY-NC-SA 3.0*
- 5.2: Solutions - Homogeneous Mixtures - *CC BY-NC-SA 3.0*
- 5.3: Solutions of Solids Dissolved in Water- How to Make Rock Candy - *CK-12 License*
- 5.4: Specifying Solution Concentration- Mass Percent - *CK-12 License*
- 5.5: Specifying Solution Concentration- Molarity - *CK-12 License*
- 5.6: Solution Stoichiometry - *CC BY-SA 4.0*
- 5.7: Sour Patch Kids and International Spy Movies - *CC BY-SA 4.0*
- 5.8: Acids- Properties and Examples - *CK-12 License*
- 5.9: Bases- Properties and Examples - *CK-12 License*
- 5.10: Molecular Definitions of Acids and Bases - *CK-12 License*
- 5.11: Reactions of Acids and Bases - *Public Domain*
- 5.12: Acid–Base Titration - *CK-12 License*
- 5.13: Strong and Weak Acids and Bases - *CK-12 License*
- 5.14: Water - Acid and Base in One - *CK-12 License*
- 5.15: The pH and pOH Scales - Ways to Express Acidity and Basicity - *CK-12 License*
- 5.16: Carbon - A Versitile Atom - *Undeclared*
- 5.17: Hydrocarbons- Compounds Containing Only Carbon and Hydrocarbon - *CC BY-NC-SA 3.0*
- 5.18: Alkanes- Saturated Hydrocarbons - *Undeclared*
- 5.19: Isomers- Same Formula, Different Structure - *Undeclared*
- 5.20: Naming Alkanes - *Undeclared*
- 5.21: Alkenes and Alkynes - *Undeclared*
- 5.22: Functional Groups - *Undeclared*
- 5.23: Carbohydrates - *Undeclared*
- 5.24: Lipids - *Undeclared*
- 5.25: Proteins - *Undeclared*

Back Matter - *Undeclared*

- Index - *Undeclared*
- Glossary - *Undeclared*
- Detailed Licensing - *Undeclared*
- Detailed Licensing - *Undeclared*