CH 104Z - INTRODUCTION TO CHEMISTRY



Stephanie Bryan & Kenneth Friedrich
Portland Community College

LibreTexts™

CH 104Z - Intro to Chemistry

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 08/24/2026

TABLE OF CONTENTS

Licensing

1: Introduction to Chemistry, the Scientific Method, and Measurement

1.1: What is Chemistry?

1.2: The Scientific Method

1.3: Hypothesis, Theories, and Laws

1.4: Types of Observations, Precision and Measurement

1.5: Experimental Design

1.6: Scientific Notation

1.7: The International System of Units

2: Unit Conversion

2.1: Converting Units

2.2: More Multi-Step Conversion Problems

2.3: Units in the Numerator and the Denominator

2.4: Density

2.5: Temperature

2.6: Dosage Calculations

2.S: Chemistry, Matter, and Measurement (Summary)

3: Classification of Matter- Properties and Changes

3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases

3.2: Classifying Matter- the Macroscopic Level

3.3: Classifying Matter- the Nanoscopic Level

3.4: The Elements

3.5: Physical and Chemical Changes- the Macroscopic Level

3.6: Physcial and Chemical Changes- the Nanoscopic Level

3.7: Separating Mixtures through Physcial Changes

3.8: Differences in Matter- Physical and Chemical Properties

4: Atoms and the Periodic Table

4.1: Prelude to Atoms, and the Periodic Table

4.2: Atomic Theory

4.3: The Structure of Atoms

4.4: Nuclei of Atoms

4.5: Atomic Masses

4.6: Bohr Diagrams of Atoms and Ions

4.7: Electron Arrangements

4.8: The Periodic Table

4.S: Elements, Atoms, and the Periodic Table (Summary)

5: Ionic Bonding and Simple Ionic Compounds

5.1: Prelude to Ionic Bonding and Simple Ionic Compounds

5.2: Two Types of Bonding

5.3: Ions

5.4: Formulas for Ionic Compounds

5.5: Ionic Nomenclature

5.6: Formula Mass

5.7: Characteristics of Ionic Compounds

5.S: Ionic Bonding and Simple Ionic Compounds (Summary)

6: Covalent Bonding and Simple Molecular Compounds

6.1: Prelude to Covalent Bonding and Simple Molecular Compounds

6.2: Covalent Bonds

6.3: Covalent Compounds - Formulas and Names

6.4: Drawing Lewis Structures

6.5: Characteristics of Covalent Bonds

6.6: Characteristics of Molecules and VSEPR

6.7: Organic Chemistry

6.S: Covalent Bonding and Simple Molecular Compounds (Summary)

7: Introduction to Chemical Reactions

7.1: Prelude to Introduction to Chemical Reactions

7.2: The Law of Conservation of Matter

7.3: Chemical Equations

7.4: Quantitative Relationships Based on Chemical Equations

7.5: Some Types of Chemical Reactions

7.6: Oxidation-Reduction (Redox) Reactions

7.7: Redox Reactions in Organic Chemistry and Biochemistry

7.S: Introduction to Chemical Reactions (Summary)

8: Quantities in Chemical Reactions

8.1: Prelude to Quantities in Chemical Reactions

8.2: The Mole

8.3: Atomic and Molar Masses

8.4: Mole-Mass Conversions

8.5: Mole-Mole Relationships in Chemical Reactions

8.6: Mole-Mass and Mass-Mass Problems

8.S: Quantities in Chemical Reactions (Summary)

9: Solids, Liquids, and Gases

9.1: Prelude to Solids, Liquids, and Gases

9.2: Intermolecular Interactions

9.3: Solids and Liquids

9.4: Gases and Pressure
9.5: Gas Laws
9.S: Solids, Liquids, and Gases (Summary)

10: Solutions

10.1: Prelude to Solutions
10.2: Solutions
10.3: Concentration
10.4: The Dissolution Process
10.5: Properties of Solutions
10.6: Chemical Equilibrium
10.7: Le Chatelier's Principle
10.8: Osmosis and Diffusion
10.S: Solutions (Summary)

11: Acids and Bases

11.1: Prelude to Acids and Bases
11.2: Arrhenius Definition of Acids and Bases
11.3: Brønsted-Lowry Definition of Acids and Bases
11.4: Water - Both an Acid and a Base
11.5: The Strengths of Acids and Bases
11.6: Buffers
11.S: Acids and Bases (Summary)

Index

Glossary

Detailed Licensing

Detailed Licensing

This page titled CH 104Z - Introduction to Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Stephanie Bryan & Kenneth Friedrich.

Licensing

A detailed breakdown of this resource's licensing can be found in ***Back Matter/Detailed Licensing***.

This page titled Licensing is shared under a license and was authored, remixed, and/or curated by LibreTexts.

1: INTRODUCTION TO CHEMISTRY, THE SCIENTIFIC METHOD, AND MEASUREMENT

1.1: What is Chemistry?

1.2: The Scientific Method

1.3: Hypothesis, Theories, and Laws

1.4: Types of Observations, Precision and Measurement

1.5: Experimental Design

1.6: Scientific Notation

1.7: The International System of Units

This page titled 1: Introduction to Chemistry, the Scientific Method, and Measurement is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

1.1: What is Chemistry?

Learning Objectives

- Understand the scope of chemistry as a scientific discipline.

Matter is any substance that has mass and takes up space. Matter includes atoms and anything made up of atoms. It does not include various forms of energy such as light, sound, or heat. Chemists often view matter on two characteristic length scales: the macroscopic and the nanoscopic. The nanoscopic level can also be called the particle level.



Figure 1.1.1: Trees in a forest are macroscopic objects.

The macroscopic scale is the length scale on which objects or phenomena are large enough to sense directly. Everything that one can see, touch, and handle in the forest of Figure 1.1.1 is within the macroscopic scale. However, each of these items can be decomposed into smaller nanoscopic scale particles.

The nanoscopic scale is the scale of atoms and molecules. All of the everyday objects that we can see or touch are ultimately composed of atoms. This ordinary atomic matter is in turn made up of interacting subatomic particles—usually a nucleus of protons and neutrons, and a cloud of orbiting electrons. Chemistry is an attempt to understand how the nanoscopic level of matter controls the macroscopic level of matter.

Matter vs. Mass

Matter should not be confused with mass, as the two are not the same in modern physics. Matter is a physical substance of which systems may be composed, while mass is not a substance, but rather a quantitative property of matter and other substances or systems.

Chemistry is the study of matter—what it consists of, what its properties are, and how it changes. Being able to describe the ingredients in a cake and how they change when the cake is baked is called chemistry. Matter is anything that has mass and takes up space—that is, anything that is physically real. Some things are easily identified as matter—this book, for example. Others are not so obvious. Because we move so easily through air, we sometimes forget that it, too, is matter.

Chemistry is one branch of science. Science is the process by which we learn about the natural universe by observing, testing, and then generating models that explain our observations. Because the physical universe is so vast, there are many different branches

of science (Figure 1.1.2). Thus, chemistry is the study of matter, biology is the study of living things, and geology is the study of rocks and the earth. Mathematics is the language of science, and we will use it to communicate some of the ideas of chemistry.




Figure 1.1.2: The Relationships between Some of the Major Branches of Science. Chemistry lies more or less in the middle, which emphasizes its importance to many branches of science.

Although we divide science into different fields, there is much overlap among them. For example, some biologists and chemists work in both fields so much that their work is called biochemistry. Similarly, geology and chemistry overlap in the field called geochemistry. Figure 1.1.2 shows how many of the individual fields of science are related.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Wikipedia

This page titled 1.1: What is Chemistry? is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 1.1: What is Chemistry? by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.2: The Scientific Method

Learning Objectives

- Define the steps of the scientific method.

How do scientists work? Generally, they follow a process called the scientific method. The scientific method is an organized procedure for learning answers to questions and making explanations for observations. To find the answer to a question (for example, "Why do birds fly toward Earth's equator during the cold months?"), a scientist goes through the following steps, which are also illustrated in Figure 1.2.1:




Figure 1.2.1: The General Steps of the Scientific Method. After an observation is made or a question is identified, a hypothesis is made and experiments are designed to test the hypothesis.

The steps may not be as clear-cut in real life as described here, but most scientific work follows this general outline.

1. **Propose a hypothesis.** A scientist generates a testable idea, or hypothesis, to try to answer a question or explain an observation about how the natural universe works. Some people use the word *theory* in place of hypothesis, but the word hypothesis is the proper word in science. For scientific applications, the word theory is a general statement that describes a large set of observations and data. A theory represents the highest level of scientific understanding.
2. **Test the hypothesis.** A scientist evaluates the hypothesis by devising and carrying out experiments to test it. If the hypothesis passes the test, it may be a proper answer to the question. If the hypothesis does not pass the test, it may not be a good answer.
3. **Refine the hypothesis if necessary.** Depending on the results of experiments, a scientist may want to modify the hypothesis and then test it again. Sometimes the results show the original hypothesis to be completely wrong, in which case a scientist will have to devise a new hypothesis.

Not all scientific investigations are simple enough to be separated into these three discrete steps. But these steps represent the general method by which scientists learn about our natural universe.

This page titled 1.2: The Scientific Method is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 1.1: What is Chemistry? by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.3: Hypothesis, Theories, and Laws

Learning Objectives

- Define the scientific means of hypothesis, theory, and law.

Although many have taken science classes throughout the course of their studies, people often have incorrect or misleading ideas about some of the most important and basic principles in science. Most students have heard of hypotheses, theories, and laws, but what do these terms really mean? Prior to reading this section, consider what you have learned about these terms before. What do these terms mean to you? What do you read that contradicts or supports what you thought?

What is a Fact?

A fact is a basic statement established by experiment or observation. All facts are true under the specific conditions of the observation.

What is a Hypothesis?

One of the most common terms used in science classes is a "hypothesis". The word can have many different definitions, depending on the context in which it is being used:

- An educated guess: a scientific hypothesis provides a suggested solution based on evidence.
- Prediction: if you have ever carried out a science experiment, you probably made this type of hypothesis when you predicted the outcome of your experiment.
- Tentative or proposed explanation: hypotheses can be suggestions about why something is observed. In order for it to be scientific, however, a scientist must be able to test the explanation to see if it works and if it is able to correctly predict what will happen in a situation. For example, "if my hypothesis is correct, we should see ___ result when we perform ___ test."

  A hypothesis is very tentative; it can be easily changed.

What is a Theory?

The *United States National Academy of Sciences* describes what a theory is as follows:

> "Some scientific explanations are so well established that no new evidence is likely to alter them. The explanation becomes a scientific theory. In everyday language a theory means a hunch or speculation. Not so in science. In science, the word **theory** refers to a comprehensive explanation of an important feature of nature supported by facts gathered over time. Theories also allow scientists to make predictions about as yet unobserved phenomena."
>
> "A scientific theory is a well-substantiated explanation of some aspect of the natural world, based on a body of facts that have been repeatedly confirmed through observation and experimentation. Such fact-supported theories are not "guesses" but reliable accounts of the real world. The theory of biological evolution is more than "just a theory." It is as factual an explanation of the universe as the atomic theory of matter (stating that everything is made of atoms) or the germ theory of disease (which states that many diseases are caused by germs). Our understanding of gravity is still a work in progress. But the phenomenon of gravity, like evolution, is an accepted fact.

Note some key features of theories that are important to understand from this description:

- Theories are explanations of natural phenomena. They aren't predictions (although we may use theories to make predictions). They are explanations as to why we observe something.
- Theories aren't likely to change. They have a large amount of support and are able to satisfactorily explain numerous observations. Theories can, indeed, be facts. Theories can change, but it is a long and difficult process. In order for a theory

to change, there must be many observations or pieces of evidence that the theory cannot explain.
- Theories are not guesses. The phrase "just a theory" has no room in science. To be a scientific theory carries a lot of weight; it is not just one person's idea about something

  Theories aren't likely to change.

What is a Law?

Scientific laws are similar to scientific theories in that they are principles that can be used to predict the behavior of the natural world. Both scientific laws and scientific theories are typically well-supported by observations and/or experimental evidence. Usually scientific laws refer to rules for how nature will behave under certain conditions, frequently written as an equation. Scientific theories are more overarching explanations of how nature works and why it exhibits certain characteristics. As a comparison, theories explain why we observe what we do and laws describe what happens.

For example, around the year 1800, Jacques Charles and other scientists were working with gases to, among other reasons, improve the design of the hot air balloon. These scientists found, after many, many tests, that certain patterns existed in the observations on gas behavior. If the temperature of the gas is increased, the volume of the gas increased. This is known as a natural law. A law is a relationship that exists between variables in a group of data. Laws describe the patterns we see in large amounts of data, but do not describe why the patterns exist.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

This page titled 1.3: Hypothesis, Theories, and Laws is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew and Henry Agnew.

1.4: Types of Observations, Precision and Measurement

Learning Objectives

- Distinguish between quantitative and qualitative observations.
- Given an instrument and an object, make a measurement including units with the appropriate level of uncertainty.
- Given a physical quantity, identify the correct number of significant figures.

Types of Observations

Observations can be qualitative or quantitative. Qualitative observations describe properties or occurrences in ways that do not rely on numbers. Examples of qualitative observations include the following: the outside air temperature is cooler during the winter season, table salt is a crystalline solid, sulfur crystals are yellow, and dissolving a penny in concentrated nitric acid forms a blue solution and a brown gas. Quantitative observations are measurements, which by definition consist of both a number and a unit. Examples of quantitative observations include the following: the melting point of crystalline sulfur is 115.21 °C, and 35.9 grams of table salt—whose chemical name is sodium chloride—dissolve in 100 grams of water at 20 °C.

Exercise 1.4.1

Classify each statement as a law, a theory, an experiment, a hypothesis, a qualitative observation, or a quantitative observation.

a. When 10 g of ice were added to 100 mL of water at 25 °C, the temperature of the water decreased to 15.5 °C after the ice melted.
b. Litmus paper dipped in lemon juice turns red.
c. A prism separates white light into a spectrum of colors.
d. Limestone is relatively insoluble in water but dissolves readily in dilute acid with the evolution of a gas.
e. Gas mixtures that contain more than 4% hydrogen in air are potentially explosive.

Answer a available at the end of this section.

Quantitative Observations or Measurements

Quantitative observations have both a number and a unit. The number communicates how precise the measurement is. Scientists have established certain conventions for communicating this degree of **precision** of a measurement, which is dependent on the measuring device used. Imagine, for example, that you are using a meterstick to measure the width of a table. The centimeters (cm) marked on the meterstick, tell you how many *centimeter*s wide the table is. Many metersticks also have markings for millimeters (mm), so we can measure the table to the nearest *millimeter*. Most metersticks do not have any smaller (or more precise) markings indicated, so you cannot report the measured width of the table any more precise than to the nearest millimeter. However, you can *estimate* one past the smallest marking, in this case the millimeter, to the next decimal place in the measurement (Figure 1.4.1).




Figure 1.4.1: Measuring an Object to the Correct Number of Digits. How many digits should be reported for the length of this object?

Reading a Measurement Device

When reading a measuring device, first determine the value represented by the smallest marked division. Record all digits that can be read with certainty from the device, and then estimate one additional digit between the smallest markings. This final estimated digit communicates the precision of the measuring device and is included as a significant figure. A length halfway between 4.2 cm and 4.3 cm might reasonably be recorded as 4.25 cm. A measurement reported with more decimal places generally came from a more precise measuring device.

For a digital instrument, such as an electronic balance, record all digits displayed because the instrument has already estimated the final digit. Always report the measurement with the appropriate unit. When reading liquid volume in a graduated cylinder, place your eyes at the same height as the liquid surface and read the bottom of the curve in the liquid surface, which is called the meniscus.

Significant Digits

The concept of **significant figures** takes this limitation into account. The significant figures of a measured quantity are defined as all the digits known with *certainty* (those indicated by the markings on the measuring device) ***and*** the first uncertain, or estimated, digit (one digit past the smallest marking on the measuring device). It makes no sense to report any digits after the first uncertain one, so it is the last digit reported in a measurement. Zeros are used when needed to place the significant figures in their correct positions. Thus, zeros are sometimes counted as significant figures but are sometimes only used as placeholders.

> "Sig figs" is a common abbreviation for significant figures.

Consider the earlier example of measuring the width of a table with a meterstick. If the table is measured and reported as being 1,357 mm wide, the number 1,357 has four significant figures. The 1 (thousands place), the 3 (hundreds place), and the 5 (tens place) are certain; the 7 (ones place) is assumed to have been estimated. It would make no sense to report such a measurement as 1,357.0 (five Sig Figs) or 1,357.00 (six Sig Figs) because that would suggest the measuring device was able to determine the width to the nearest tenth or hundredth of a millimeter, when in fact it shows only tens of millimeters and therefore the ones place was estimated.

On the other hand, if a measurement is reported as 150 mm, the 1 (hundreds) and the 5 (tens) are known to be significant, but how do we know whether the zero is or is not significant? The measuring device could have had marks indicating every 100 mm or marks indicating every 10 mm. How can you determine if the zero is significant (the estimated digit), or if the 5 is significant and the zero a value placeholder?

The **rules** for deciding which digits in a measurement are significant are as follows:

1. All nonzero digits are significant. In 1,357 mm, all the digits are significant.
2. *Sandwiched (or embedded) zeros*, those between significant digits, are significant. Thus, 405 g has *three* significant figures.
3. *Leading zeros*, which are zeros at the beginning of a decimal number less than 1, are not significant. In 0.000458 mL, the first four digits are leading zeros and are not significant. The zeros serve only to put the digits 4, 5, and 8 in the correct decimal positions. This number has three significant figures.
4. *Trailing zeros*, which are zeros at the end of a number, are significant only if the number has a decimal point. Thus, in 1,500 m, the two trailing zeros are not significant because the number is written without a decimal point; the number has two significant figures. However, in 1,500.00 m, all six digits are significant because the number has a decimal point.

Example 1.4.1

How many significant figures does each number have?

a. 6,798,000
b. 6,000,798
c. 6,000,798.00
d. 0.0006798

Answer a available at the end of this section.

Exercise 1.4.2

How many significant figures does each number have?

a. 2.1828
b. 0.005505
c. 55,050
d. 5
e. 500

Answer a available at the end of this section.

Rounding off numbers

Before dealing with the specifics of the rules for determining the significant figures in a calculated result, we need to be able to round numbers correctly. To **round** a number, first decide how many significant figures the number should have. Once you know that, round to that many digits, starting from the left. **If the number immediately to the right of the last significant digit is less than 5, it is dropped and the value of the last significant digit remains the same. If the number immediately to the right of the last significant digit is greater than or equal to 5, the last significant digit is increased by 1.**

Consider the measurement 207.518m. Right now, the measurement contains six significant figures. How would we successively round it to fewer and fewer significant figures? Follow the process as outlined in Table 1.5.1.

Table 1.5.1: Rounding examples

Number of Significant Figures	Rounded Value	Reasoning
6	207.518	All digits are significant
5	207.52	8 rounds the 1 up to 2
4	207.5	2 is dropped
3	208	5 rounds the 7 up to 8
2	210	8 is replaced by a 0 and rounds the 0 up to 1
1	200	1 is replaced by a 0

Notice that the more rounding that is done, the less reliable the figure is. An approximate value may be sufficient for some purposes, but scientific work requires a much higher level of detail.

It is important to be aware of significant figures when you are mathematically manipulating numbers. For example, dividing 125 by 307 on a calculator gives 0.4071661238... to an infinite number of digits. But do the digits in this answer have any practical meaning, especially when you are starting with numbers that have only three significant figures each? When performing mathematical operations, there are two rules for limiting the number of significant figures in an answer—one rule is for addition and subtraction, and one rule is for multiplication and division. We will not be learning these rules in this class but you will need to round your answer to the given number of significant digits because these rules exist.

Answers

Exercise 1.4.1

quantitative observation
qualitative observation
qualitative observation
qualitative observation
quantitative observation

Example 1.4.1

four (by rules 1 and 4)
seven (by rules 1 and 2)
nine (by rules 1, 2, and 4)
four (by rules 1 and 3)

Exercise 1.4.2

five
four
four
one
one

This page titled 1.4: Types of Observations, Precision and Measurement is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Anonymous.

- 1.5: Expressing Numbers - Significant Figures by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.5: Experimental Design

Learning Objectives

- Classify dependent, independent, and controlled variables.

An **experiment** is a controlled method of testing a hypothesis. Experiments are designed to provide an opportunity to make observation that will help test a hypothesis. Experiments are best understood in term of three types of variables: independent variables, dependent variables, and controlled variables. It may be helpful to think of a variable simply as something that can be measured.

Most experiments are repeated multiple times with slight variations. These repetitions are often called trials. A variable that is purposely altered between trials is called an **independent variable**. Usually, it is a best practice to have a single independent variable in an experiment. It may be helpful to think of the independent variable as the "input" of the experiment.

If the independent variable is the "input" of an experiment, than the **dependent variable** is the "output." Dependent variables change in response to the independent variable (their name comes from the fact that they depend on the independent variable).

Finally, some variables, called **controlled variables**, are kept constant through all of the trials. Controlled variables are kept constant so that their fluctuations do not alter the dependent variable and cloud its relationship with the independent variable.

Sometimes a scatter plot (a mathematical diagram in which data is graphed against an x and a y axis) is used to display results from an experiment. If a scatter plot is used, the independent variable is generally plotted on the horizontal axis and the dependent variable is plotted on the vertical axis.

This page titled 1.5: Experimental Design is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Anonymous.

1.6: Scientific Notation

Learning Objectives

- Convert numbers between scientific notation and decimal form.

The instructions for making a pot of coffee specified 3 scoops (rather than 12,000 grounds) because any measurement is expressed more efficiently with units that are appropriate in size. In science, however, we often must deal with quantities that are extremely small or incredibly large. For example, you may have 5,000,000,000,000 red blood cells in a liter of blood, and the diameter of an iron atom is 0.000000014 inches. Numbers with many zeros can be cumbersome to work with, so scientists use scientific notation.

Scientific notation is a system for expressing very large or very small numbers in a compact manner. It uses the idea that such numbers can be rewritten as a simple number multiplied by 10 raised to a certain exponent, or power.

Let us look first at very large numbers. Suppose a spacecraft is 1,500,000 miles from Mars. The number 1,500,000 can be thought of as follows:

$$1.5 \times \underbrace{1{,}000{,}000}_{\underbrace{10 \times 10 \times 10 \times 10 \times 10 \times 10}_{10^6}} = 1.5 \times 10^6$$

That is, 1,500,000 is the same as 1.5 times 1 million, and 1 million is 10 × 10 × 10 × 10 × 10 × 10, or 10^6 (which is read as "ten to the sixth power"). Therefore, 1,500,000 can be rewritten as 1.5 times 10^6, or 1.5 × 10^6. The distance of the spacecraft from Mars can therefore be expressed as 1.5 × 10^6 miles.

Recall that:

- $10^0 = 1$
- $10^1 = 10$
- $10^2 = 100$
- $10^3 = 1{,}000$
- $10^4 = 10{,}000$
- and so forth

The standard convention for expressing numbers in scientific notation is to write a single *nonzero* first digit, a decimal point, and the rest of the digits, excluding any trailing zeros (see rules for significant figures in the next section for more details on what to exclude). This number is followed by a multiplication sign and then by 10 raised to the power necessary to reproduce the original number. For example, although 1,500,000 can also be written as 15. × 10^5 (which would be 15. × 100,000), the convention is to have only one digit before the decimal point. How do we know to what power 10 is raised? The power is the number of places you have to move the decimal point to the *left* to place it after the first digit, so that the number being multiplied is *between 1 and 10*:

$$1{,}500{,}000 = 1.5 \times 10^6$$

Example 1.6.1: Scientific Notation

Express each number in scientific notation.

a. 67,000,000,000
b. 1,689
c. 12.6

Answer a available at the end of this section.

Exercise 1.6.1

Express each number in scientific notation.

a. 1,492
b. 102,000,000
c. 101,325

Answer a available at the end of this section.

To change a number in **scientific notation** to **standard form**, we reverse the process, moving the decimal point to the right. Add zeros to the end of the number being converted, if necessary, to produce a number of the proper magnitude. Lastly, we drop the number 10 and its power.

$$1.5 \times 10^6 = 1.500000. = 1{,}500{,}000$$

Example 1.6.2

Express each number in standard, or conventional notation.

a. 5.27×10^4
b. 1.0008×10^6

Answer a available at the end of this section.

Exercise 1.6.2

Express each number in standard, or conventional notation.

a. 6.98×10^8
b. 1.005×10^2

Answer a available at the end of this section.

We can also use scientific notation to express numbers whose magnitudes are less than 1. For example, the quantity 0.006 centimeters can be expressed as follows:

$$6 \times \underbrace{\frac{1}{1{,}000}}_{\underbrace{\frac{1}{10} \times \frac{1}{10} \times \frac{1}{10}}_{10^{-3}}} = 6 \times 10^{-3}$$

That is, 0.006 centimeters is the same as 6 *divided by* one thousand, which is the same as 6 *divided* by 10 x 10 x 10 or 6 *times* 10^{-3} (which is read as "ten to the negative third power"). Therefore, 0.006 centimeters can be rewritten as 6 times 10^{-3}, or 6 × 10^{-3} centimeters.

Recall that:

- $10^{-1} = 1/10$
- $10^{-2} = 1/100$
- $10^{-3} = 1/1{,}000$
- $10^{-4} = 1/10{,}000$
- $10^{-5} = 1/100{,}000$
- and so forth

We use a negative number as the power to indicate the number of places we have to move the decimal point to the right to make it follow the first nonzero digit so that the number is between 1 and 10. This is illustrated as follows:

$$0.\underbrace{006} = 6 \times 10^{-3}$$

Note:

In writing scientific notations, the convention is to have only one digit before the decimal point.

- Numbers that are greater than one have a positive power in scientific notation. If the decimal point is moved to the left n places, the power (n) of 10 is positive.

$$1,\underbrace{500,}_{\text{6 places}}\underbrace{000}_{\text{6 places}} = 1.5 \times 10^{6}$$

- Numbers that are less than one have a negative power in scientific notation. If the decimal point is moved to the right n places, the power (n) of 10 is negative. $0.\underbrace{006} = 6 \times 10^{-3}$

Example 1.6.3

Express each number in scientific notation.

a. 0.000006567
b. −0.0004004
c. 0.000000000000123

Answer a available at the end of this section.

Exercise 1.6.3

Express each number in scientific notation.

a. 0.000355
b. 0.314159
c. −0.051204

Answer a available at the end of this section.

As with numbers with positive powers of 10, when changing from **scientific** notation to **standard or conventional** format, we reverse the process.

$$6 \times 10^{-3} = .006 = 0.006$$

Note

Changing a number in scientific notation to standard form:

- If the scientific notation has a positive power, the standard number is greater than one. Example: $8 \times 10^{4} = 80{,}000$
- If the scientific notation has a negative power, then the standard number is less than one. Example: $8 \times 10^{-2} = 0.08$

Example 1.6.4

Change the number in scientific notation to standard form.

a. 6.22×10^{-2}
b. 9.9×10^{-9}

Answer a available at the end of this section.

Exercise 1.6.4

Change the number in scientific notation to standard form.

a. 9.98×10^{-5}
b. 5.109×10^{-8}

Answer a available at the end of this section.

Although calculators can show 8 to 10 digits in their display windows, that is not always enough when working with very large or very small numbers. For this reason, many calculators are designed to handle scientific notation. The method for entering scientific notation differs for each calculator model, so take the time to learn how to do it properly on your calculator, *asking your instructor for assistance if necessary*. If you do not learn to enter scientific notation into your calculator properly, you will not get the correct final answer when performing a calculation.

Answers

Example 1.6.1: Scientific Notation

Moving the decimal point 10 places to the left gives 6.7×10^{10}.

The decimal point is assumed to be at the end of the number, so moving it three places to the left gives 1.689×10^{3}.

In this case, we need to move the decimal point only one place to the left, which yields 1.26×10^{1}.

Exercise 1.6.1

Moving the decimal point 3 places to the left gives 1.492×10^{3}.

The decimal point is assumed to be at the end of the number, so moving it 8 places to the left gives 1.02×10^{8}.

Moving the decimal point 5 places to the left yields 1.01325×10^{5}.

Example 1.6.2

Moving the decimal four places to the right and adding zeros give 52,700.

Moving the decimal six places to the right and adding zeros give 1,000,800.

Exercise 1.6.2

Moving the decimal point eight places to the right and adding zeros give 698,000,000.

Moving the decimal point two places to the right gives 100.5

Example 1.6.3

Move the decimal point six places to the right to get 6.567×10^{-6}.

Move the decimal point four places to the right to get -4.004×10^{-4}. The negative sign on the number itself does not affect how we apply the rules of scientific notation.

Move the decimal point 13 places to the right to get 1.23×10^{-13}.

Exercise 1.6.3

Moving the decimal point four places to the right gives 3.55×10^{-4}.

Moving the decimal point one place to the right gives 3.14159×10^{-1}.

Moving the decimal point one place to the right gives -5.1204×10^{-2}.

Example 1.6.4

0.0622

0.0000000099

Exercise 1.6.4

0.0000998

0.00000005109

This page titled 1.6: Scientific Notation is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Anonymous.

- 1.4: Expressing Numbers - Scientific Notation by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

1.7: The International System of Units

Learning Objectives

- List SI units of length, mass, time, and temperature.
- List the multipliers, symbols, and numerical meanings for all SI units.

People who live in the United States measure weight in pounds, height in feet and inches, and a car's speed in miles per hour. In contrast, chemistry and other branches of science use the International System of Units (also known as **SI** after *Système Internationale d'Unité*s), which was established so that scientists around the world could communicate efficiently with each other. Many countries have also adopted SI units for everyday use as well. The United States is one of the few countries that has not.

Base SI Units

Base (or basic) units, are the fundamental units of SI. There are seven base units, which are listed in Table 1.7.1, Chemistry uses five of the base units: the mole for amount, the kilogram for mass, the meter for length, the second for time, and the Kelvin for temperature. The degree Celsius (°C) is also commonly used for temperature. The numerical relationship between Kelvins and degrees Celsius is as follows:

$$K = {}^{\circ}C + 273 \tag{1.7.1}$$

Table 1.7.1: The Seven Base SI Units

Property	Unit	Abbreviation
length	meter	m
mass	kilogram	kg
time	second	s
amount	mole	mol
temperature	Kelvin	K
electrical current	ampere	amp
luminous intensity	candela	cd

The United States uses the English (sometimes called Imperial) system of units for many quantities. Inches, feet, miles, gallons, pounds, and so forth, are all units connected with the English system of units. There have been many mistakes due to the improper conversion of units between the SI and English systems.

The size of each base unit is defined by international convention. For example, the *kilogram* is defined as the quantity of mass of a special metal cylinder kept in a vault in France (Figure 1.7.1). The other base units have similar definitions and standards. The sizes of the base units are not always convenient for all measurements. For example, a meter is a rather large unit for describing the width of something as narrow as human hair. Instead of reporting the diameter of hair as 0.00012 m or as 1.2×10^{-4} m using scientific notation as discussed in section 1.4, SI also provides a series of **prefixes** that can be attached to the units, creating units that are larger or smaller by powers of 10.




Figure 1.7.1: The Kilogram. The standard for the kilogram is a platinum-iridium cylinder kept in a special vault in France. Source: Photo reproduced by permission of the Bureau International des Poids et Mesures, who retain full internationally protected copyright.

Common prefixes and their multiplicative factors are listed in Table 1.7.2. (Perhaps you have already noticed that the base unit *kilogram* is a combination of a prefix, kilo- meaning 1,000 ×, and a unit of mass, the gram.) Some prefixes create a multiple of the original unit: 1 kilogram equals 1,000 grams, and 1 megameter equals 1,000,000 meters. Other prefixes create a fraction of the original unit. Thus, 1 centimeter equals 1/100 of a meter, 1 millimeter equals 1/1,000 of a meter, 1 microgram equals 1/1,000,000 of a gram, and so forth.

Table 1.7.2: Prefixes Used with SI Units

Prefix	Abbreviation	Multiplicative Factor	Multiplicative Factor in Scientific Notation
giga-	G	1,000,000,000 ×	10^{9} ×
mega-	M	1,000,000 ×	10^{6} ×
kilo-	k	1,000 ×	10^{3} ×
deca-	D	10 ×	10^{1} ×
deci-	d	1/10 ×	10^{-1} ×
centi-	c	1/100 ×	10^{-2} ×
milli-	m	1/1,000 ×	10^{-3} ×
micro-	μ*	1/1,000,000 ×	10^{-6} ×
nano-	n	1/1,000,000,000 ×	10^{-9} ×
***The letter μ is the Greek lowercase letter for *m* and is called "mu," which is pronounced "myoo."**			

Both SI units and prefixes have abbreviations, and the combination of a prefix abbreviation with a base unit abbreviation gives the abbreviation for the modified unit. For example, kg is the abbreviation for kilogram. We will be using these abbreviations throughout this book.

The Difference Between Mass and Weight

The mass of a body is a measure of its inertial property or how much matter it contains. The weight of a body is a measure of the force exerted on it by gravity or the force needed to support it. Gravity on earth gives a body a downward acceleration of about 9.8 m/s^2. In common parlance, weight is often used as a synonym for mass in weights and measures. For instance, the verb “to weigh” means “to determine the mass of” or “to have a mass of.” The incorrect use of weight in place of mass should be phased out, and the term mass used when mass is meant. The SI unit of mass is the kilogram (kg). In science and technology, the weight of a body in a particular reference frame is defined as the force that gives the body an acceleration equal to the local acceleration of free fall in that reference frame. Thus, the SI unit of the quantity weight defined in this way (force) is the newton (N).

Derived SI Units

Derived units are combinations of SI base units. Units can be multiplied and divided, just as numbers can be multiplied and divided. For example, the area of a square having a side of 2 cm is 2 cm × 2 cm, or 4 cm^2 (read as “four centimeters squared” or “four square centimeters”). Notice that we have squared a length unit, the centimeter, to get a derived unit for area, the square centimeter.

Volume is an important quantity that uses a derived unit. **Volume** is the amount of space that a given substance occupies and is defined geometrically as length × width × height. Each distance can be expressed using the meter unit, so volume has the derived unit m × m × m, or m^3 (read as “meters cubed” or “cubic meters”). A cubic meter is a rather large volume, so scientists typically express volumes in terms of 1/1,000 of a cubic meter. This unit has its own name—the liter (L). A liter is a little larger than 1 US quart in volume. Below are approximate equivalents for some of the units used in chemistry.

Approximate Equivalents to Some SI Units

- 1 m ≈ 39.36 in. ≈ 3.28 ft ≈ 1.09 yd
- 1 in. = 2.54 cm (this is exact rather than an approximation)
- 1 km ≈ 0.6214 mi
- 1 kg ≈ 2.205 lb
- 1 lb ≈ 454 g
- 1 L ≈ 1.06 qt
- 1 qt ≈ 0.946 L




Figure 1.7.2: The relative lengths of 1 m, 1 yd, 1 cm, and 1 in. are shown (not actual size), as well as comparisons of 2.54 cm and 1 in., and of 1 m and 1.094 yd. (CC BY 4.0; OpenStax)

As shown in Figure 1.7.3, a liter is also 1,000 cm^3. By definition, there are 1,000 mL in 1 L, so 1 milliliter and 1 cubic centimeter represent the same volume.

$$1\ mL = 1\ cm^3 \tag{1.7.2}$$




Figure 1.7.3: Units of Volume. (a) The relative volumes are shown for cubes of 1 m^3, 1 dm^3 (1 L), and 1 cm^3 (1 mL) (not to scale). A liter (L) is defined as a cube 1 dm (1/10th of a meter) on a side. A milliliter (mL), 1/1,000th of a liter, is equal to 1 cubic centimeter. (b) The diameter of a dime is compared relative to the edge length of a 1-cm^3 (1-mL) cube. (CC BY 4.0; OpenStax)

Example 1.7.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kiloliter
b. microsecond
c. decimeter
d. nanogram

Answer a available at the end of this section.

Exercise 1.7.1

Give the abbreviation for each unit and define the abbreviation in terms of the base unit.

a. kilometer
b. milligram
c. nanosecond
d. centiliter

Answer a available at the end of this section.

Energy, another important quantity in chemistry, is the ability to perform work, such as moving a box of books from one side of a room to the other side. It has a derived unit of kg•m^2/s^2. (The dot between the kg and m units implies the units are multiplied together.) Because this combination is cumbersome, this collection of units is redefined as a **joule** (J). An older unit of energy, but likely more familiar to you, the calorie (cal), is also widely used. There are 4.184 J in 1 cal. Energy changes occur during all chemical processes and will be discussed in a later chapter.

Answers

Example 1.7.1

The abbreviation for a kiloliter is kL. Because kilo means “1,000 ×,” 1 kL equals 1,000 L.

The abbreviation for microsecond is µs. Micro implies 1/1,000,000th of a unit, so 1 µs equals 0.000001 s.

The abbreviation for decimeter is dm. Deci means 1/10th, so 1 dm equals 0.1 m.

The abbreviation for nanogram is ng and equals 0.000000001 g.

Exercise 1.7.1

km (1,000 m)

mg (0.001 g)

ns (0.000000001 s)

cL (0.01L)

This page titled 1.7: The International System of Units is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 1.6: The International System of Units by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2: UNIT CONVERSION

The study of chemistry will open your eyes to a fascinating world. Chemical processes are continuously at work all around us. They happen as you cook and eat food, strike a match, shampoo your hair, and even read this page. Chemistry is called the central science because a knowledge of chemical principles is essential for other sciences. You might be surprised at the extent to which chemistry pervades your life.

2.1: Converting Units

2.2: More Multi-Step Conversion Problems

2.3: Units in the Numerator and the Denominator

2.4: Density

2.5: Temperature

2.6: Dosage Calculations

2.S: Chemistry, Matter, and Measurement (Summary)

Template:HideTOC

This page titled 2: Unit Conversion is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 1: Chemistry, Matter, and Measurement by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.1: Converting Units

Learning Objectives

- Convert a value reported in one unit to a corresponding value in a different unit.

The ability to convert from one unit to another is an important skill. For example, a nurse with 50 mg aspirin tablets who must administer 0.2 g of aspirin to a patient needs to know that 0.2 g equals 200 mg, so 4 tablets are needed. Fortunately, there is a simple way to convert from one unit to another.

Conversion Factors

If you learned the SI units and prefixes described, then you know that 1 cm is 1/100th of a meter.

$$1 \text{ cm} = \frac{1}{100} \text{ m}$$

or

$$100 \text{ cm} = 1 \text{ m}$$

Suppose we divide both sides of the equation by 1 m (both the number *and* the unit):

$$\frac{100 \text{ cm}}{1 \text{ m}} = \frac{1 \text{ m}}{1 \text{ m}}$$

As long as we perform the same operation on both sides of the equals sign, the expression remains an equality. Look at the right side of the equation; it now has the same quantity in the numerator (the top) as it has in the denominator (the bottom). Any fraction that has the same quantity in the numerator and the denominator has a value of 1:

$$\text{same quantity} \left\langle \frac{100 \text{ cm}}{1 \text{ m}} = 1 \right.$$

We know that 100 cm *is* 1 m, so we have the same quantity on the top and the bottom of our fraction, although it is expressed in different units. A fraction that has equivalent quantities in the numerator and the denominator but expressed in *different units* is called a **conversion factor**.

Here is a simple example. How many centimeters are there in 3.55 m? Perhaps you can determine the answer in your head. If there are 100 cm in every meter, then 3.55 m equals 355 cm. To solve the problem more formally with a conversion factor, we first write the quantity we are given, 3.55 m. Then we multiply this quantity by a conversion factor, which is the same as multiplying it by 1. We can write 1 as $\frac{100 \text{ cm}}{1 \text{ m}}$ and multiply:

$$3.55 \text{ m} \times \frac{100 \text{ cm}}{1 \text{ m}}$$

The 3.55 m can be thought of as a fraction with a 1 in the denominator. Because m, the abbreviation for meters, occurs in both the numerator *and* the denominator of our expression, they cancel out:

$$\frac{3.55 \cancel{\text{ m}}}{1} \times \frac{100 \text{ cm}}{1 \cancel{\text{ m}}}$$

The final step is to perform the calculation that remains once the units have been canceled:

$$\frac{3.55}{1} \times \frac{100 \text{ cm}}{1} = 355 \text{ cm} \tag{2.1.1}$$

In the final answer, we omit the 1 in the denominator. Thus, by a more formal procedure, we find that 3.55 m equals 355 cm. A generalized description of this process is as follows:

$$\text{quantity (in old units)} \times \text{conversion factor} = \text{quantity (in new units)}$$

You may be wondering why we use a seemingly complicated procedure for a straightforward conversion. In later studies, the conversion problems you will encounter *will not always be so simple*. If you can master the technique of applying conversion factors, you will be able to solve a large variety of problems.

In the previous example (Equation 2.1.1), we used the fraction $\frac{100\text{ cm}}{1\text{ m}}$ as a conversion factor. Does the conversion factor $\frac{1\text{ m}}{100\text{ cm}}$ also equal 1? Yes, it does; it has the same quantity in the numerator as in the denominator (except that they are expressed in different units). Why did we not use *that* conversion factor? If we had used the second conversion factor, the original unit would not have canceled, and the result would have been meaningless. Here is what we would have gotten:

$$3.55\text{ m} \times \frac{1\text{ m}}{100\text{ cm}} = 0.0355\frac{\text{m}^2}{\text{cm}}$$

For the answer to be meaningful, we have to *construct the conversion factor in a form that causes the original unit to cancel out*. Figure 2.1.1 shows a **concept map** for constructing a proper conversion.




Figure 2.1.1: A Concept Map for Conversions. This is how you construct a conversion factor to convert from one unit to another.

Meters converted to centimeters. A fraction showing 100 cm over 1 m. The unit you convert to is on top, and the unit you convert from is on bottom.

Significant Figures in Conversions

How do conversion factors affect the determination of significant figures? Numbers in conversion factors based on prefix changes, such as kilograms to grams, are *not* considered in the determination of significant figures in a calculation because the numbers in such conversion factors are exact. **Exact numbers** are *defined* or *counted* numbers, not measured numbers, and can be considered as having an infinite number of significant figures. (In other words, 1 kg is exactly 1,000 g, by the definition of kilo-.) Counted numbers are also exact. If there are 16 students in a classroom, the number 16 is exact. In contrast, conversion factors that come from measurements (such as density, as we will see shortly) or are approximations have a limited number of significant figures and should be considered in determining the significant figures of the final answer.

Example 2.1.1


a. The average volume of blood in an adult male is 4.7 L. What is this volume in milliliters?
b. A hummingbird can flap its wings once in 18 ms. How many seconds are in 18 ms?

Solution

a. We start with what we are given, 4.7 L. We want to change the unit from liters to milliliters. There are 1,000 mL in 1 L. From this relationship, we can construct two conversion factors:

$$\frac{1\text{ L}}{1,000\text{ mL}} \text{ or } \frac{1,000\text{ mL}}{1\text{ L}}$$

We use the conversion factor that will cancel out the original unit, liters, and introduce the unit we are converting to, which is milliliters. The conversion factor that does this is the one on the right.




$$4.7\cancel{\text{L}} \times \frac{1,000\text{ mL}}{1\ \cancel{\text{L}}} = 4,700\text{ mL}$$

Because the numbers in the conversion factor are exact, we do not consider them when determining the number of significant figures in the final answer. Thus, we report two significant figures in the final answer.

b. We can construct two conversion factors from the relationships between milliseconds and seconds:

$$\frac{1,000\text{ ms}}{1\text{ s}} \text{ or } \frac{1\text{ s}}{1,000\text{ ms}}$$

To convert 18 ms to seconds, we choose the conversion factor that will cancel out milliseconds and introduce seconds. The conversion factor on the right is the appropriate one. We set up the conversion as follows:



$$18\ \cancel{\text{ms}} \times \frac{1\text{ s}}{1,000\ \cancel{\text{ms}}} = 0.018\text{ s}$$

The conversion factor's numerical values do not affect our determination of the number of significant figures in the final answer.

Exercise 2.1.1

Perform each conversion.

a. 101,000 ns to seconds
b. 32.08 kg to grams

Answer a available at the end of this section.

Conversion Factors From Different Units

Conversion factors can also be constructed for converting between different kinds of units. For example, density can be used to convert between the mass and the volume of a substance. Consider mercury, which is a liquid at room temperature and has a density of 13.6 g/mL. The density tells us that 13.6 g of mercury have a volume of 1 mL. We can write that relationship as follows:

13.6 g mercury = 1 mL mercury

This relationship can be used to construct two conversion factors:

$$\frac{13.6\ \text{g}}{1\ \text{mL}} \text{ and } \frac{1\ \text{mL}}{13.6\ \text{g}}$$

Which one do we use? It depends, as usual, on the units we need to cancel and introduce. For example, suppose we want to know the mass of 16 mL of mercury. We would use the conversion factor that has milliliters on the bottom (so that the milliliter unit cancels) and grams on top so that our final answer has a unit of mass:

$$16\ \cancel{\text{mL}} \times \frac{13.6\ \text{g}}{1\ \cancel{\text{mL}}} = 217.6\ \text{g}$$

$$\approx 220\ \text{g}$$

In the last step, we limit our final answer to two significant figures because the volume quantity has only two significant figures; the 1 in the volume unit is considered an exact number, so it does not affect the number of significant figures. The other conversion factor would be useful if we were given a mass and asked to find volume, as the following example illustrates.

Density can be used as a conversion factor between mass and volume.

Example 2.1.2: Mercury Thermometer

A mercury thermometer for measuring a patient's temperature contains 0.750 g of mercury. What is the volume of this mass of mercury?

Solution

Because we are starting with grams, we want to use the conversion factor that has grams in the denominator. The gram unit will cancel algebraically, and milliliters will be introduced in the numerator.

$$0.750\ \cancel{\text{g}} \times \frac{1\ \text{mL}}{13.6\ \cancel{\text{g}}} = 0.055147\ldots\ \text{mL}$$

$$\approx 0.0551\ \text{mL}$$

We have limited the final answer to three significant figures.

Exercise 2.1.2

What is the volume of 100.0 g of air if its density is 1.3 g/L?

Answer available at the end of this section.

Looking Closer: Density and the Body

The densities of many components and products of the body have a bearing on our health.

Bones. Bone density is important because bone tissue of lower-than-normal density is mechanically weaker and susceptible to breaking. The density of bone is, in part, related to the amount of calcium in one's diet; people who have a diet deficient in calcium, which is an important component of bones, tend to have weaker bones. Dietary supplements or adding dairy products to the diet seems to help strengthen bones. As a group, women experience a decrease in bone density as they age. It has been estimated that fully half of women over age 50 suffer from excessive bone loss, a condition known as osteoporosis. Exact bone densities vary within the body, but for a healthy 30-year-old female, it is about 0.95–1.05 g/cm^3. Osteoporosis is diagnosed if the bone density is below 0.6–0.7 g/cm^3.

Urine. The density of urine can be affected by a variety of medical conditions. Sufferers of diabetes produce an abnormally large volume of urine with a relatively low density. In another form of diabetes, called diabetes mellitus, there is excess glucose dissolved in the urine, so that the density of urine is abnormally high. The density of urine may also be abnormally high because of excess protein in the urine, which can be caused by congestive heart failure or certain renal (kidney) problems. Thus, a urine density test can provide clues to various kinds of health problems. The density of urine is commonly expressed as a specific gravity, which is a unitless quantity defined as

$$\frac{\text{density of some material}}{\text{density of water}}$$

Normal values for the specific gravity of urine range from 1.002 to 1.028.

Body Fat. The overall density of the body is one indicator of a person's total body fat. Fat is less dense than muscle and other tissues, so as it accumulates, the overall density of the body decreases. Measurements of a person's weight and volume provide the overall body density, which can then be correlated to the percentage of body fat. (The body's volume can be measured by immersion in a large tank of water. The amount of water displaced is equal to the volume of the body.)

Problem Solving With Multiple Conversions

Sometimes you will have to perform more than one conversion to obtain the desired unit. For example, suppose you want to convert 54.7 km into millimeters. You can either memorize the relationship between kilometers and millimeters, or you can do the conversion in two steps. Most people prefer to convert in steps.

To do a stepwise conversion, we first convert the given amount to the base unit. In this example, the base unit is meters. We know that there are 1,000 m in 1 km:

$$54.7 \cancel{\text{ km}} \times \frac{1,000 \text{ m}}{1 \cancel{\text{ km}}} = 54,700 \text{ m}$$


Then we take the result (54,700 m) and convert it to millimeters, remembering that there are $1,000 \text{ mm}$ for every 1 m:

$$54,700 \cancel{\text{ m}} \times \frac{1,000 \text{ mm}}{1 \cancel{\text{ m}}} = 54,700,000 \text{ mm}$$
$$= 5.47 \times 10^7 \text{ mm}$$

We have expressed the final answer in scientific notation.

As a shortcut, both steps in the conversion can be combined into a single, multistep expression:

Concept Map



Calculation

$$54.7 \cancel{\text{km}} \times \frac{1,000 \cancel{\text{m}}}{1 \cancel{\text{km}}} \times \frac{1,000 \text{ mm}}{1 \cancel{\text{m}}} = 54,700,000 \text{ mm}$$

$$= 5.47 \times 10^7 \text{ mm}$$

In each step, the previous unit is canceled and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.

Either method—one step at a time or all the steps together—is acceptable. If you do all the steps together, the restriction for the proper number of significant figures should be done after the last step. As long as the math is performed correctly, you should get the same answer no matter which method you use.


Example 2.1.3

Convert 58.2 ms to megaseconds in one multistep calculation.

Solution

First, convert the given unit (ms) to the base unit—in this case, seconds—and then convert seconds to the final unit, megaseconds:

Concept Map



Calculation

$$58.2 \cancel{\text{ms}} \times \frac{1 \cancel{\text{s}}}{1,000 \cancel{\text{ms}}} \times \frac{1 \text{ Ms}}{1,000,000 \cancel{\text{s}}} = 0.0000000582 \text{ Ms}$$

$$= 5.82 \times 10^{-8} \text{ Ms}$$

Neither conversion factor affects the number of significant figures in the final answer.

Exercise 2.1.3

Convert 43.007 mg to kilograms in one multistep calculation.

Answer available at the end of this section.

Career Focus: Pharmacist

A pharmacist dispenses drugs that have been prescribed by a doctor. Although that may sound straightforward, pharmacists in the United States must hold a doctorate in pharmacy and be licensed by the state in which they work. Most pharmacy programs require four years of education in a specialty pharmacy school.

Pharmacists must know a lot of chemistry and biology so they can understand the effects that drugs (which are chemicals, after all) have on the body. Pharmacists can advise physicians on the selection, dosage, interactions, and side effects of drugs. They can also advise patients on the proper use of their medications, including when and how to take specific drugs properly. Pharmacists can be found in drugstores, hospitals, and other medical facilities.

Curiously, an outdated name for pharmacist is *chemist*, which was used when pharmacists formerly did a lot of drug preparation, or *compounding*. In modern times, pharmacists rarely compound their own drugs, but their knowledge of the sciences, including chemistry, helps them provide valuable services in support of everyone's health.

Key Takeaway

- A unit can be converted to another unit of the same type with a conversion factor.

Concept Review Exercises

1. How do you determine which quantity in a conversion factor goes in the denominator of the fraction?
2. State the guidelines for determining significant figures when using a conversion factor.
3. Write a concept map (a plan) for how you would convert 1.0×10^{12} *nano*liters (nL) to *kilo*liters (kL).

Answers

1. The unit you want to cancel from the numerator goes in the denominator of the conversion factor.
2. Exact numbers that appear in many conversion factors do not affect the number of significant figures; otherwise, the normal rules of multiplication and division for significant figures apply.
3. Concept Map: Convert the given (nanoliters, nL) to liters; then convert liters to kiloliters.



Answers

Exercise 2.1.1

$$101,000 \cancel{\mu s} \times \frac{1 \text{ s}}{1,000,000,000 \cancel{\mu s}} = 0.000101 \text{ s}$$

$$32.08 \cancel{\text{kg}} \times \frac{1,000 \text{ g}}{1 \cancel{\text{kg}}} = 32,080 \text{ g}$$

Exercise 2.1.2

$$100.0 \cancel{g} \times \frac{1 \text{ L}}{1.3 \cancel{g}} = 76.92307692 \text{ L} \approx 77\text{L}$$

Because the density (1.3 g/L) has only 2 significant figures, we are rounding off the final answer to 2 significant figures.

Exercise 2.1.3

$$43.007 \cancel{\text{mg}} \times \frac{\cancel{1\text{g}}}{1,000 \cancel{\text{mg}}} \times \frac{1 \text{ kg}}{1,000 \cancel{g}} = 0.000043007 \text{ kg}$$
$$= 4.3007 \times 10^{-5} \text{ kg}$$

.

Neither conversion factor affects the number of significant figures in the final answer.

This page titled 2.1: Converting Units is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 1.7: Converting Units by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.2: More Multi-Step Conversion Problems

Learning Objectives


- Given a quantity, convert from one set of units to another using dimensional analysis showing canceling on units. This includes multiple factor conversions.

Multiple Conversions

Sometimes you will have to perform more than one conversion to obtain the desired unit. For example, suppose you want to convert 54.7 km into millimeters. We will set up a series of conversion factors so that each conversion factor produces the next unit in the sequence. We first convert the given amount in km to the base unit, which is meters. We know that 1,000 m =1 km.

Then we convert meters to mm, remembering that $1\ \text{mm} = 10^{-3}\ \text{m}$.

Concept Map



Calculation


$$54.7\ \cancel{\text{km}} \times \frac{1,000\ \cancel{\text{m}}}{1\ \cancel{\text{km}}} \times \frac{1\ \text{mm}}{10^{-3}\ \cancel{\text{m}}} = 54,700,000\ \text{mm}$$
$$= 5.47 \times 10^{7}\ \text{mm}$$

In each step, the previous unit is canceled and the next unit in the sequence is produced, each successive unit canceling out until only the unit needed in the answer is left.

Example 2.2.1: Unit Conversion

Convert 58.2 ms to megaseconds in one multi-step calculation.

Solution

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 58.2 ms Find: Ms
List other known quantities	$1ms = 10^{-3}s$ $1Ms = 10^{6}s$
Prepare a concept map.	 $\frac{10^{-3}s}{1\ ms}$ $\frac{1\ Ms}{10^{6}\ s}$
Calculate.	$58.2\ \cancel{ms} \times \frac{10^{-3}\ \cancel{s}}{1\ \cancel{ms}} \times \frac{1\ \text{Ms}}{1,000,000\ \cancel{s}} = 0.0000000582\ \text{Ms}$ $= 5.82 \times 10^{-8}\ \text{Ms}$ Neither conversion factor affects the number of significant figures in the final answer.

Example 2.2.2: Unit Conversion

How many seconds are in 2.50 days?

Solution

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 2.50 days Find: s
List other known quantities.	1 day = 24 hours 1 hour = 60 minutes 1 minute = 60 seconds
Prepare a concept map.	 $\frac{24\text{hr}}{1\text{ day}}$ $\frac{60\text{min}}{1\text{ hr}}$ $\frac{60\text{ s}}{1\ min}$
Calculate.	$2.50\text{ d} \times \frac{24\text{ hr}}{1\text{ d}} \times \frac{60\text{ min}}{1\text{ hr}} \times \frac{60\text{ s}}{1\text{ min}} = 216,000\text{ s}$ 2.16×10^5 s 3SF, not ambiguous

Exercise 2.2.1

Perform each conversion in one multi-step calculation.

a. 43.007 ng to kg
b. 1005 in to yd

Answer a available at the end of this section.

Answer b available at the end of this section.

Contributions

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 2.2.1

$4.3007 \times 10^{-11} kg$

Exercise 2.2.1

$27.92\ yd$

This page titled 2.2: More Multi-Step Conversion Problems is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew and Henry Agnew.

2.3: Units in the Numerator and the Denominator

Learning Objectives


- Given a quantity, convert from one set of units to another using dimensional analysis showing canceling on units. This includes compound units (eg: mph to kps).

Compound Units

There are many units that combine a unit in the numerator with a unit in the denominator. Common examples are miles per hour, meters per second, and miles per gallon. The "per" that appears when we talk about the compound unit or when we write it in text form translates to the divider between numerator and denominator when we write it symbolically. Therefore miles per hour would be written mi/h, meters per second would be written m/s, and miles per gallon would be written mi/gal.

For example, suppose we want to convert 0.0128 kilometers per minute to meters per hour.

Concept Map



Calculation


$$\frac{0.0128\ \cancel{\text{km}}}{1\ \cancel{\text{min}}} \times \frac{60\ \cancel{\text{min}}}{1\ \text{h}} \times \frac{1,000\ \text{m}}{1\ \cancel{\text{km}}} = \frac{768\ \text{m}}{\text{h}}$$
$$= 768\ \text{m/h}$$

Which we would write or say as 768 meters per hour.

Example 2.3.1: Unit Conversion

Convert 45 miles per gallon to kilometers per liter in one multi-step calculation.

Solution

Steps for Problem Solving	Unit Conversion
Identify the "given" information and what the problem is asking you to "find."	Given: 45 mi/gal Find: km/L
List other known quantities	$1km = 0.6214mi$ $1gal = 3.785L$
Prepare a concept map.	
Calculate.	$\frac{45\ \cancel{\text{mi}}}{1\ \cancel{\text{gal}}} \times \frac{1\ \cancel{\text{gal}}}{3.785\ \text{L}} \times \frac{1\ \text{km}}{0.6214\ \cancel{\text{mi}}} = \frac{19\ \text{km}}{\text{L}}$ $= 19\ \text{km/L}$

Exercise 2.3.1

Convert 1.25 cm/s to in/min.

Answer available at the end of this section.

Answers

Exercise 2.3.1

$29.5\ in/min$

This page titled 2.3: Units in the Numerator and the Denominator is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

2.4: Density

Learning Objectives

- Given a mass and volume calculate the density of a substance.

Density (ρ) is a physical property found by dividing the mass of an object by its volume. Regardless of the sample size, density is always constant. For example, the density of a pure sample of tungsten is always 19.25 grams per cubic centimeter. This means that whether you have one gram or one kilogram of the sample, the density will never vary. The equation is as follows:

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}} \tag{2.4.1}$$

or just

$$\rho = \frac{m}{V} \tag{2.4.2}$$

Based on this equation, it's clear that density can, and does, vary from element to element and substance to substance due to differences in the relationship of mass and volume. Pure water, for example, has a density of 0.998 g/cm^3 at 25° C. The average densities of some common substances are in Table 2.4.1. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will “float.”

Table 2.4.1: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Density can be measured for all substances—solids, liquids and gases. For solids and liquids, density is often reported using the units of g/cm^3 or g/mL. Densities of gases, which are significantly lower than the densities of solids and liquids, are often given using units of g/L.

Example 2.4.1: Ethyl Alcohol

Calculate the density of a 30.2 mL sample of ethyl alcohol with a mass of 23.71002 g

Solution

This is a direct application of Equation 2.4.2:

$$\rho = \frac{23.71002\,g}{30.2\,mL} = 0.785\,g/mL$$

Exercise 2.4.1

a. Find the density (in kg/L) of a sample that has a volume of 36.5 L and a mass of 10.0 kg.
b. If you have a 2.130 mL sample of acetic acid with mass 0.002234 kg, what is the density in kg/L?

Answer a available at the end of this section.

Density as a Conversion Factor

Conversion factors can also be constructed for converting between different kinds of units. For example, density can be used to convert between the mass and the volume of a substance. Consider mercury, which is a liquid at room temperature and has a density of 13.6 g/mL. The density tells us that 13.6 g of mercury have a volume of 1 mL. We can write that relationship as follows:

13.6 g mercury = 1 mL mercury

This relationship can be used to construct two conversion factors:

$$\frac{13.6\,g}{1\,mL} = 1$$

and

$$\frac{1\,mL}{13.6\,g} = 1$$

Which one do we use? It depends, as usual, on the units we need to cancel and introduce. For example, suppose we want to know the mass of 2.0 mL of mercury. We would use the conversion factor that has milliliters on the bottom (so that the milliliter unit cancels) and grams on top, so that our final answer has a unit of mass:

$$2.0\ \cancel{\text{mL}} \times \frac{13.6\ \text{g}}{1\ \cancel{\text{mL}}} = 27.2\ \text{g} = 27\ \text{g}$$

In the last step, we limit our final answer to two significant figures because the volume quantity has only two significant figures; the 1 in the volume unit is considered an exact number, so it does not affect the number of significant figures. The other conversion factor would be useful if we were given a mass and asked to find volume, as the following example illustrates.


Density can be used as a conversion factor between mass and volume.

Example 2.4.2: Mercury Thermometer Steps for Problem Solving

A mercury thermometer for measuring a patient's temperature contains 0.750 g of mercury. What is the volume of this mass of mercury?

Solution

Steps for Problem Solving	Unit Conversion

Identify the "given" information and what the problem is asking you to "find."	Given: 0.750 g Find: mL
List other known quantities.	13.6 g/mL (density of mercury)
Prepare a concept map.	
Calculate.	$0.750\ \cancel{g} \times \frac{1\ \text{mL}}{13.6\ \cancel{g}} = 0.0551\ \text{mL}$

Density is a measurement, and like any measurement it contains some error. The significant figures in the density must therefore be considered when determining the significant figures in any answer calculated using it. Consider, for example, if we had tried calculating the volume of 0.7500 g of mercury in the previous example. Although the mass we are starting with now has four significant figures, the density we are using as a conversion factor still only has three significant figures. We would therefore still need to report the answer to three significant figures, 0.0551 mL.

Exercise 2.4.2

What is the volume of 100.0 g of air if its density is 1.3 g/L?

Answer available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 2.4.1

$0.274\ kg/L$

$1.049\ kg/L$

Exercise 2.4.2

$77\ L$

This page titled 2.4: Density is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

2.5: Temperature

Learning Objectives

- Convert temperatures between Celsius, Kelvin, and Fahrenheit scales.

The concept of temperature may seem familiar to you, but many people confuse temperature with heat. **Temperature** is a measure of the average kinetic energy of the particles in a sample of matter. As temperature increases, the kinetic energy of the particles also increases. **Heat**, on the other hand, is the flow of thermal energy from an object with a higher temperature to an object with a lower temperature. Temperature is an important parameter in chemistry. When a substance changes from solid to liquid, it is almost always because there was an increase in the temperature of the material. Chemical reactions usually proceed faster if the temperature is increased. Many unstable materials (such as enzymes) will be viable longer at lower temperatures.



Figure 2.5.1: The glowing charcoal on the left represents high kinetic energy, while the snow and ice on the right are of much lower kinetic energy.

Three different scales are commonly used to measure temperature: Fahrenheit (expressed as °F), Celsius (°C), and Kelvin (K). Thermometers measure temperature by using materials that expand or contract when heated or cooled. Mercury or alcohol thermometers, for example, have a reservoir of liquid that expands when heated and contracts when cooled, so the liquid column lengthens or shortens as the temperature of the liquid changes.

The Fahrenheit Scale

The main problem with the Fahrenheit scale is the arbitrary definitions of temperature. The freezing point of water was defined as $32^{\circ}\mathrm{F}$ and the boiling point as $212^{\circ}\mathrm{F}$. The Fahrenheit scale is typically not used for scientific purposes.

The Celsius Scale

The Celsius scale sets the freezing point and boiling point of water at $0^{\circ}\mathrm{C}$ and $100^{\circ}\mathrm{C}$ respectively. The distance between those two points is divided into 100 equal intervals, each of which is one degree. Another term sometimes used for the Celsius scale is "centigrade" because there are 100 degrees between the freezing and boiling points of water on this scale. However, the preferred term is "Celsius".

The Kelvin Scale

The Kelvin temperature scale is based on molecular motion, with the temperature of $0\ \mathrm{K}$, also known as absolute zero, being the point where all molecular motion ceases. The freezing point of water on the Kelvin scale is $273.15\ \mathrm{K}$, while the boiling point is $373.15\ \mathrm{K}$. Notice that there is no "degree" used in the temperature designation. Unlike the Fahrenheit and Celsius scales where temperatures are referred to as "degrees F" or "degrees C", we simply designate temperatures in the Kelvin scale as Kelvins.




Figure 2.5.2: A Comparison of the Fahrenheit, Celsius, and Kelvin Temperature Scales. Because the difference between the freezing point of water and the boiling point of water is 100° on both the Celsius and Kelvin scales, the size of a degree Celsius (°C) and a kelvin (K) are precisely the same. In contrast, both a degree Celsius and a kelvin are 9/5 the size of a degree Fahrenheit (°F). (CC BY-SA-NC 3.0; anonymous)

Converting Between Scales

The Kelvin is the same size as the Celsius degree, so measurements are easily converted from one to the other. The freezing point of water is 0°C = 273.15 K; the boiling point of water is 100°C = 373.15 K. The Kelvin and Celsius scales are related as follows:

$$T\,(\text{in °C}) + 273.15 = T\,(\text{in K}) \tag{2.5.1}$$

$$T\ \ (\text{in K}) - 273.15 = T\ (\text{in °C}) \tag{2.5.2}$$

Degrees on the Fahrenheit scale, however, are based on an English tradition of using 12 divisions, just as 1 ft = 12 in. The relationship between degrees Fahrenheit and degrees Celsius is as follows:

$$°C = \frac{(°F - 32)}{1.8} \tag{2.5.3}$$

$$°F = 1.8 \times (°C) + 32 \tag{2.5.4}$$

Example 2.5.1: Temperature Conversions

A student is ill with a temperature of 103.5°F. What is her temperature in °C and K?

Solution

Converting from Fahrenheit to Celsius:

$$°C = \frac{(103.5°F - 32)}{1.8} = 39.7°C \tag{2.5.5}$$

Converting from Celsius to Kelvin:

$$K = 39.7°C + 273.15 = 312.9K \tag{2.5.6}$$

Exercise 2.5.1

Convert each temperature to °C and °F.

a. the temperature of the surface of the sun (5800 K)
b. the boiling point of gold (3080 K)
c. the boiling point of liquid nitrogen (77.36 K)

Answer (a) available at the end of this section.

Answer (b) available at the end of this section.

Answer (c) available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 2.5.1
5527 °C, 9981 °F

Exercise 2.5.1
2807 °C, 5085 °F

Exercise 2.5.1
-195.79 °C, -320.42 °F

This page titled 2.5: Temperature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew and Henry Agnew.

2.6: Dosage Calculations

Learning Objectives

- Calculate drug dosages using conversion factors.

To Your Health: Dosages

A medicine can be more harmful than helpful if it is not taken in the proper dosage. A dosage (or dose) is the specific amount of a medicine that is known to be therapeutic for an ailment in a patient of a certain size. Dosages of the active ingredient in medications are usually described by units of mass, typically grams or milligrams, and generally are equated with a number of capsules or teaspoonfuls to be swallowed or injected. The amount of the active ingredient in a medicine is carefully controlled so that the proper number of pills or spoonfuls contains the proper dose.

Most drugs must be taken in just the right amount. If too little is taken, the desired effects will not occur (or will not occur fast enough for comfort); if too much is taken, there may be potential side effects that are worse than the original ailment. Some drugs are available in multiple dosages. For example, tablets of the medication levothyroxine sodium, a synthetic thyroid hormone for those suffering from decreased thyroid gland function, are available in 11 different doses, ranging from 25 micrograms (μg) to 300 μg. It is a doctor's responsibility to prescribe the correct dosage for a patient, and it is a pharmacist's responsibility to provide the patient with the correct medicine at the dosage prescribed. Thus, proper quantities—which are expressed using numbers and their associated units—are crucial for keeping us healthy.

Effects are dose-dependent

Chemicals are the most common things for which doses are measured, but there are others, such as radiation exposure. For humans, most doses of micronutrients and medications are measured in milligrams (mg), but some are measured in micrograms because of their potency. Nonmedicinal poisons span the measurement scale; some poisons are so dangerous that a single microgram of it could be deadly, whereas other substances take much more. For example, even water is toxic when consumed in large enough quantities.

Dosage (the size of each dose) determines the strength and duration of the health benefits of nutrients, and also of the therapeutic effects of medical treatments. Dosage also determines the severity of adverse effects of treatments and toxins.

Duration of exposure, that is, the period of time over which the dose was received (all at once or gradually) also determines its effects (the body may build tolerance to gradual exposure to a drug, while a large immediate dose could be deadly).

The route by which a dose is exposed to, may affect the outcome, because some medications have different effects depending on whether they are inhaled, ingested, taken transdermally, injected, or inserted.

The dosage, route, concentration, and division over time may all be critical considerations in the administering of drugs, or in responding to exposure to a toxin. In nutrition, the route is usually a given, as nutrients are generally eaten; while dosage and the frequency of ingestion of nutrients are very important variables in preventing disease and promoting overall health.

Calculation of dose

Calculating drug dosages for humans based on the doses used in animal studies can be based on weight (e.g., mg/kg) or surface area (e.g., mg/m^2) based on weight. Drug dosage calculation is required if the physician's order is different from what is available.

Example 2.6.1

a. A physician ordered 100 mg of Demerol. Demerol is available as 50 mg per tablet. How many tablets should the nurse administer?
b. The doctor's order is 1.2 g of Folic Acid. Folic Acid is available as 800 mg per tablet. How many tablets should be taken?

Solution

a. We start with the given, 100 mg. We want to change the unit from *mg* to *tablets*. There are 50 mg in 1 tablet (Remember that *per* tablet means *one* tablet. From this relationship, we can construct the conversion factor. We use the conversion factor that will cancel out the original unit, *mg*, and introduce the unit we are converting to, which is *tablet*.

$$100 \cancel{\text{mg}} \times \frac{1 \text{ tablet}}{50 \cancel{\text{mg}}} = 2 \text{ tablets}$$

Hence, the nurse should administer 2 tablets.

b. We start with the given, 1.2 g and we want to change *grams* to number of *tablets*. First, we convert *1.2 g* to *mg* and then convert *mg to tablets*. We need a conversion factor for each step.

$$1.2 \cancel{\text{g}} \times \frac{1{,}000 \cancel{\text{mg}}}{1 \cancel{\text{g}}} \times \frac{1 \text{ tablet}}{800 \cancel{\text{mg}}} = 1.5 \text{ tablets}$$

Hence, 1.5 tablets should be taken.

Exercise 2.6.1

Calculate each of the following.

a. The physician ordered 20 mg of Valium. Valium is available as 10 mg per tablet. How many tablets should the nurse administer?
b. The doctor's order is 1 g of Calcium. What is on hand is Calcium as 500 mg per tablet. How many tablets should be taken?

Answer a available at the end of this section.

Drug Dosage Calculation based on Body Weight. Many drugs (especially in children) are dosed according to body weight (mg/kg). These calculations are carried in 3-step conversions. The first step is to convert the body weight from *pounds (lbs)* to *kg*. The second step is to convert *kg* to *mg* (the total *mg dose* calculated based on body weight). And, finally, the *mg dose* is converted to the number of *tablets*. (as in Example 1.8.1).

Example 2.6.2

a. Demerol is ordered 1.5mg/kg for a patient that is 220 lbs. Demerol is available as 50 mg per tablet. How many tablets should the nurse administer?
b. A doctor prescribes amoxicillin 30mg/kg to a child weighing 73.5 lbs. Amoxicillin is available as 500 mg tablets. How many tablets should the nurse administer?

Solution

a. We start with the given, 220 *lbs*. We want to change the unit from *lbs* to *kg,* and then, from *kg* to *total dose (mg)* and then *the mg dose to tablets*. The first conversion factor will cancel out the original unit, *lbs*, and introduce the unit we are converting to, which is *kg*. The second conversion factor will cancel out *kg*, and introduce the unit of the *dose* (usually *mg*) and then *mg* to *tablet*.

$$220\ \cancel{\text{lbs}} \times \frac{1\ \cancel{\text{kg}}}{2.2\ \cancel{\text{lbs}}} \times \frac{1.5\ \cancel{\text{mg}}}{1\ \cancel{\text{kg}}} \times \frac{1\ \text{tablet}}{50\ \cancel{\text{mg}}} = 3\ \text{tablets}$$

Hence, the nurse administers 3 tablets.

b. Start with the given, 73.5 *lbs*. We want to change the unit from *lbs* to *kg*, and then, from *kg* to total dose (*mg*) and then *mg* to *tablets*. The first conversion factor will cancel out the original unit, *lbs,* and convert to *kg*. The second conversion factor will cancel out *kg*, and convert to the total *mg dose* and the final conversion will cancel *mg* to introduce the final unit, *tablet*.

$$73.5\ \cancel{\text{lbs}} \times \frac{1\ \cancel{\text{kg}}}{2.2\ \cancel{\text{lbs}}} \times \frac{30\ \cancel{\text{mg}}}{1\ \cancel{\text{kg}}} \times \frac{1\ \text{tablet}}{500\ \cancel{\text{mg}}} = 2\ \text{tablets}$$

Hence, the nurse administers 2 tablets.

Exercise 2.6.2

Calculate each of the following.

a. Vancomycin is ordered 15mg/kg for a patient that is 110 lbs. Vancomycin is available as 250 mg per capsule. How many capsules should the nurse administer? (ans. 3 capsules)
b. A doctor prescribes ampicillin 40mg/kg to a patient weighing 55 lbs. Ampicillin is available as 500 mg tablets. How many tablets should the nurse administer? (ans. 2 tablets)

Answer a available at the end of this section.

References

1. The Use of Body Surface Area as a Criterion of Drug Dosage in Cancer Chemotherapy(opens in new window) [cancerres.aacrjournals.org] "The Use of Body Surface Area as a Criterion of Drug Dosage in Cancer Chemotherapy" D Pinkel. Cancer Research 1958

Contributors and Attributions

- Wikipedia
- Leticia Colmenares (UH-WCC)

Answers

Exercise 2.6.1

Start with 20 mg of Valium. For the conversion factor, we know that 10 mg Valium = 1 tablet

$$20\ \cancel{\text{mg}} \times \frac{1\ \text{tablet}}{10\ \cancel{\text{mg}}} = 2\ \text{tablets}$$

Start with 1 g (1000 mg) of Calcium. For the conversion factor, we know that 500 mg Calcium = 1 tablet

$$1000\ \cancel{\text{mg}} \times \frac{1\ \text{tablet}}{500\ \cancel{\text{mg}}} = 2\ \text{tablets}$$

Exercise 2.6.2

$$110\ \cancel{\text{lbs}} \times \frac{1\ \cancel{\text{kg}}}{2.2\ \cancel{\text{lbs}}} \times \frac{15\ \cancel{\text{mg}}}{1\ \cancel{\text{kg}}} \times \frac{1\ \text{tablet}}{250\ \cancel{\text{mg}}} = 3\ \text{tablets}$$

$$55\ \cancel{\text{lbs}} \times \frac{1\ \cancel{\text{kg}}}{2.2\ \cancel{\text{lbs}}} \times \frac{40\ \cancel{\text{mg}}}{1\ \cancel{\text{kg}}} \times \frac{1\ \text{tablet}}{500\ \cancel{\text{mg}}} = 2\ \text{tablets}$$

This page titled 2.6: Dosage Calculations is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 1.8: Dosage Calculations by Anonymous is licensed CC BY-SA 4.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

2.S: Chemistry, Matter, and Measurement (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Chemistry is the study of **matter**, which is anything that has mass and takes up space. Chemistry is one branch of **science**, which is the study of the natural universe. Like all branches of science, chemistry relies on the **scientific method**, which is a process of learning about the world around us. In the scientific method, a guess or **hypothesis** is tested through experiment and measurement.

Matter can be described in a number of ways. **Physical properties** describe characteristics of a sample that do not change the chemical identity of the material (size, shape, color, and so on), while **chemical properties** describe how a sample of matter changes its chemical composition. A **substance** is any material that has the same physical and chemical properties throughout. An **element** is a substance that cannot be broken down into chemically simpler components. The smallest chemically identifiable piece of an element is an **atom**. A substance that can be broken down into simpler chemical components is a **compound**. The smallest chemically identifiable piece of a compound is a **molecule**. Two or more substances combine physically to make a **mixture**. If the mixture is composed of discrete regions that maintain their own identity, the mixture is a **heterogeneous mixture**. If the mixture is so thoroughly mixed that the different components are evenly distributed throughout, it is a **homogeneous mixture**. Another name for a homogeneous mixture is a **solution**. Substances can also be described by their **phase**: solid, liquid, or gas.

Scientists learn about the universe by making measurements of **quantities**, which consist of **numbers** (how many) and **units** (of what). The numerical portion of a quantity can be expressed using **scientific notation**, which is based on **powers**, or exponents, of 10. Large numbers have positive powers of 10, while numbers less than 1 have negative powers of 10. The proper reporting of a measurement requires proper use of **significant figures**, which are all the known digits of a measurement plus the first estimated digit. The number of figures to report in the result of a calculation based on measured quantities depends on the numbers of significant figures in those quantities. For addition and subtraction, the number of significant figures is determined by position; for multiplication and division, it is decided by the number of significant figures in the original measured values. Nonsignificant digits are dropped from a final answer in accordance with the rules of **rounding**.

Chemistry uses SI, a system of units based on seven **basic units**. The most important ones for chemistry are the units for length, mass, amount, time, and temperature. Basic units can be combined with numerical prefixes to change the size of the units. They can also be combined with other units to make **derived units**, which are used to express other quantities such as **volume**, **density**, or **energy**. A formal conversion from one unit to another uses a **conversion factor**, which is constructed from the relationship between the two units. Numbers in conversion factors may affect the number of significant figures in a calculated quantity, depending on whether the conversion factor is **exact**. Conversion factors can be applied in separate computations, or several can be used at once in a single, longer computation. Conversion factors are very useful in calculating **dosages**.

This page titled 2.S: Chemistry, Matter, and Measurement (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 1.S: Chemistry, Matter, and Measurement (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3: CLASSIFICATION OF MATTER- PROPERTIES AND CHANGES

3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases

3.2: Classifying Matter- the Macroscopic Level

3.3: Classifying Matter- the Nanoscopic Level

3.4: The Elements

3.5: Physical and Chemical Changes- the Macroscopic Level

3.6: Physcial and Chemical Changes- the Nanoscopic Level

3.7: Separating Mixtures through Physcial Changes

3.8: Differences in Matter- Physical and Chemical Properties

This page titled 3: Classification of Matter- Properties and Changes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases

Learning Objectives

- To describe the solid, liquid and gas phases.

Water can take many forms. At low temperatures (below $0^{\circ}\mathrm{C}$), it is a solid. When at "normal" temperatures (between $0^{\circ}\mathrm{C}$ and $100^{\circ}\mathrm{C}$), it is a liquid. While at temperatures above $100^{\circ}\mathrm{C}$, water is a gas (steam). The state that water is in depends upon the temperature. Each state has its own unique set of physical properties. Matter typically exists in one of three states: **solid**, **liquid**, or **gas**.




Figure 3.1.1: Matter is usually classified into three classical states, with plasma sometimes added as a fourth state. From left to right: quartz (solid), water (liquid), nitrogen dioxide (gas).

Some substances exist as gases at room temperature (oxygen and carbon dioxide), while others, like water and mercury metal, exist as liquids. Most metals exist as solids at room temperature. Most substances can exist in any of these three states. Figure 3.1.2 shows the differences among solids, liquids, and gases. On the macroscopic level, a solid has definite volume and shape, a liquid has a definite volume but no definite shape, and a gas has neither a definite volume nor shape.





Figure 3.1.2: A Representation of the Solid, Liquid, and Gas States. (a) Solid O_2 has a fixed volume and shape, and the molecules are packed tightly together. (b) Liquid O_2 conforms to the shape of its container but has a fixed volume; it contains relatively densely packed molecules. (c) Gaseous O_2 fills its container completely—regardless of the container's size or shape—and consists of widely separated molecules.

Plasma: A Fourth State of Matter

Technically speaking, a fourth state of matter called plasma exists, but it does not naturally occur on earth, so we will omit it from our study here.



A plasma globe operating in a darkened room. (CC BY-SA 3.0; Chocolateoak).

Solids

In the solid state, the individual particles of a substance are in fixed positions with respect to each other because there is not enough thermal energy to overcome the attractive forces between the particles (known as intermolecular forces). As a result, solids have a definite shape and volume. Most solids are hard, but some (like waxes) are relatively soft. Many solids composed of ions can also be quite brittle.

Solids are defined by the following characteristics:

- Definite shape (rigid)
- Definite volume
- Particles vibrate around fixed axes

If we were to cool liquid mercury to its freezing point of $-39^{\circ}\mathrm{C}$, and under the right pressure conditions, we would notice all of the liquid particles would go into the solid state. Mercury can be solidified when its temperature is brought to its freezing point. However, when returned to room temperature conditions, mercury does not exist in solid state for long, and returns back to its more common liquid form.

Solids often have their constituent particles arranged in a regular, three-dimensional array called a **crystal**, as shown in Figure 3.1.3. Some solids, especially those composed of large molecules, cannot easily organize their particles in such regular crystals and exist as amorphous (literally, “without form”) solids. Glass is one example of an amorphous solid.




Figure 3.1.3: (left) The periodic crystalline lattice structure of quartz SiO_2 in two-dimensions. (right) The random network structure of glassy SiO_2 in two-dimensions. Note that, as in the crystal, each Silicon atom is bonded to 4 oxygen atoms, where the fourth oxygen atom is obscured from view in this plane. Images used with permission (public domain).

Liquids

If the particles of a substance have enough energy to partially overcome intermolecular interactions, then the particles can move about each other while remaining in contact. This describes the liquid state. In a liquid, the particles are still in close contact, so liquids have a definite volume. However, because the particles can move about each other rather freely, a liquid has no definite shape and takes a shape dictated by its container.

Liquids have the following characteristics:

- No definite shape (takes the shape of its container).
- Has definite volume.
- Particles are free to move over each other, but are still attracted to each other.

A familiar liquid is mercury metal. Mercury is an anomaly. It is the only metal we know of that is liquid at room temperature. Mercury also has an ability to stick to itself (surface tension)—a property that all liquids exhibit. Mercury has a relatively high surface tension, which makes it very unique. Here you see mercury in its common liquid form.

YouTube Video

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Video 3.1.1*: Mercury boiling to become a gas.*

If we heat liquid mercury to its boiling point of $357^{\circ}\mathrm{C}$ under the right pressure conditions, we would notice all particles in the liquid state go into the gas state.

Gases

If the particles of a substance have enough energy to completely overcome intermolecular interactions, then the particles can separate from each other and move about randomly in space. This describes the gas state. Like liquids, gases have no definite shape, but unlike solids and liquids, gases have no definite volume either. The change from solid to liquid usually does not significantly change the volume of a substance. However, the change from a liquid to a gas significantly increases the volume of a substance, by a factor of 1,000 or more. Gases have the following characteristics:

- No definite shape (takes the shape of its container)
- No definite volume (another way of saying this is that a gas is highly compressible)
- Particles move in random motion with little or no attraction to each other

Table 3.1.1*: Characteristics of the Three States of Matter*

Characteristics	Solids	Liquids	Gases
shape	definite	indefinite	indefinite
volume	definite	definite	indefinite
strength of intermolecular forces relative to kinetic energy	strong	moderate	weak
relative particle positions	in contact and fixed in place	in contact but not fixed	not in contact, random positions

Example 3.1.1

What state or states of matter does each statement, describe?

a. This state has a definite volume, but no definite shape.
b. This state has no definite volume.
c. This state allows the individual particles to move about while remaining in contact.

Solution

a. This statement describes the liquid state.
b. This statement describes the gas state.

c. This statement describes the liquid state.

Exercise 3.1.1

What state or states of matter does each statement describe?

a. This state has individual particles in a fixed position with regard to each other.
b. This state has individual particles far apart from each other in space.
c. This state has a definite shape.

Answer a: available at the end of this section.

Answer b: available at the end of this section.

Answer c: available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 3.1.1
solid

Exercise 3.1.1
gas

Exercise 3.1.1
solid

This page titled 3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew and Henry Agnew.

3.2: Classifying Matter- the Macroscopic Level

Learning Objectives

- Given a macroscopic definition classify the type of matter.

Matter can be classified using a hierarchy (Figure 3.2.1). In this section, we will think about this hierarchy in terms of macroscopic properties. In the next section, we will revisit this hierarchy based on nanoscopic structure.




Figure 3.2.1: Relationships between the Types of Matter and the Methods Used to Separate Mixtures

A **pure substance** always has the same composition and same physical properties regardless of the source of the sample. For example, pure water is always composed of two atoms of hydrogen for every one atom of oxygen. This means its composition is always the same. Also, the physical properties of pure water will always be the same: at "normal" pressures it will boil at $100^{\circ}\mathrm{C}$, freeze at $0^{\circ}\mathrm{C}$, and have a density of 1.00 g/ml.

Contrast this with salt water. The amount of salt in the water can vary, meaning the composition of salt water is not always the same. Additionally, the density, freezing print, and boiling point will change depending how much salt is in the water. Variable composition and variable properties means that salt water is not a pure substance. In fact, it is composed of two pure substances: water and sodium chloride. Any matter composed of multiple pure substances is a **mixture**. Mixtures can be separated into pure substances by means of physical separations.

There are two types of pure substances: compounds and elements. **Compounds** can be broken into simpler pure substances by chemical means. When a current with a sufficient voltage is passed through water, for example, it can be broken into oxygen and hydrogen. This process is called hydrolysis and is one example of a chemical separation. Since water is a pure substance that can be broken into simpler pure substances by chemical means, water is a compound.

A pure substance that can not be broken down into simpler substances by chemical means is an element. You might already know from a previous science course that the periodic table contains all the known elements. The oxygen and hydrogen that form when water undergoes hydrolysis are examples of elements. They cannot be broken down into simpler substances by any means. Today, there are about 118 elements in the known universe. In contrast, scientists have identified tens of millions of different compounds to date.

Mixtures can also be further divided: some are homogeneous and some are heterogeneous.

A **homogeneous mixture** is a mixture in which the composition is uniform throughout the mixture. The salt water described

above is homogeneous because the dissolved salt is evenly distributed throughout the entire salt water sample. Often it is easy to confuse a homogeneous mixture with a pure substance because they are both uniform. The difference is that the composition of the substance is always the same. The amount of salt in the salt water can vary from one sample to another. All solutions are considered homogeneous because the dissolved material is present in the same amount throughout the solution.

A **heterogeneous mixture** is a mixture in which the composition is not uniform throughout the mixture. Vegetable soup is a heterogeneous mixture. Any given spoonful of soup will contain varying amounts of the different vegetables and other components of the soup.

Phase

A phase is any part of a sample that has a uniform composition and properties. By definition, a pure substance or a homogeneous mixture consists of a single phase. A heterogeneous mixture consists of two or more phases. When oil and water are combined, they do not mix evenly, but instead form two separate layers. Each of the layers is called a phase.

Example 3.2.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. filtered tea
b. freshly squeezed orange juice
c. a compact disc
d. aluminum oxide, a white powder that contains a 2:3 ratio of aluminum and oxygen atoms
e. selenium

Given: a chemical substance

Asked for: its classification

Strategy:

A. Decide whether a substance is chemically pure. If it is pure, the substance is either an element or a compound. If a substance can be separated into its elements, it is a compound.
B. If a substance is not chemically pure, it is either a heterogeneous mixture or a homogeneous mixture. If its composition is uniform throughout, it is a homogeneous mixture.

Solution

a. **A)** Tea is a solution of compounds in water, so it is not chemically pure. It is usually separated from tea leaves by filtration.
B) Because the composition of the solution is uniform throughout, it is ***a homogeneous mixture***.
b. **A)** Orange juice contains particles of solid (pulp) as well as liquid; it is not chemically pure.
B) Because its composition is not uniform throughout, orange juice is ***a heterogeneous mixture***.
c. **A)** A compact disc is a solid material that contains more than one element, with regions of different compositions visible along its edge. Hence, a compact disc is not chemically pure.
B) The regions of different composition indicate that a compact disc is ***a heterogeneous mixture.***
d. **A)** Aluminum oxide is a single, chemically ***pure compound***.
e. **A)** Selenium is one of the known ***elements***.

Exercise 3.2.1

Identify each substance as a compound, an element, a heterogeneous mixture, or a homogeneous mixture (solution).

a. white wine
b. mercury

c. ranch-style salad dressing
d. table sugar (sucrose)

Answer a: available at the end of this section.

Example 3.2.2

How would a chemist categorize each example of matter?

a. saltwater
b. soil
c. water
d. oxygen

Solution

a. Saltwater acts as if it were a single substance even though it contains two substances—salt and water. Saltwater is a homogeneous mixture, or a solution.
b. Soil is composed of small pieces of a variety of materials, so it is a heterogeneous mixture.
c. Water is a substance. More specifically, because water is composed of hydrogen and oxygen, it is a compound.
d. Oxygen, a substance, is an element.

Exercise 3.2.2

How would a chemist categorize each example of matter?

a. coffee
b. hydrogen
c. an egg

Answer a: available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Stephen Lower, Professor Emeritus (Simon Fraser U.) Chem1 Virtual Textbook
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 3.2.1

homogeneous mixture (solution)
element
heterogeneous mixture
compound

Exercise 3.2.2

a homogeneous mixture (solution), assuming it is filtered coffee
element
heterogeneous mixture

This page titled 3.2: Classifying Matter- the Macroscopic Level is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.3: Classifying Matter- the Nanoscopic Level

Learning Objectives

- Given a particle level picture or chemical formula classify the type of matter.

Particles

In the previous section, matter was classified based on macroscopic observations such as the ability of a sample to be separated into simpler substances or whether the appearance of the sample is consistent throughout all its parts. These macroscopic observations are manifestations of the nanoscopic structure. In this section, different shapes will be used to represent atoms of different elements. We will use the term **particle** to refer to an atom or group of atoms that are chemically bonded together.

Pure Substances

The drawings below all represent pure substances on the nanoscopic level.




Figure 3.3.1: Nanoscopic Level Representations of Pure Substances

In a pure substance, all the particles are the same. Notice that the particle may be composed of a single atom or multiple atoms. If there are multiple atoms, the atoms may be all the same element or they may be different elements. The important feature is that the same particle is repeated over and over again throughout the sample.

Mixtures

Notice how the drawings below, which all represent mixtures, differ from the drawings of pure substances above.




Figure 3.3.2: Nanoscopic Level Representations of Mixtures

In a mixture, all the particles are not the same. That is to say, a mixture is composed of two or more pure substances. In homogeneous mixtures, the composition of the particles is the same throughout. In heterogeneous mixtures, particular regions are richer in one type of particle relative to other regions in the sample.

Elements

Recall that pure substances can be further subdivided into elements and compounds. The following drawings all represent sample of elements.




Figure 3.3.3: Nanoscopic Level Representations of Elements

In a sample of an element, not only are all the particles the same, but all of the atoms that compose the particle are the same. Notice that the particle may contain a single atom or multiple atoms.

Compounds

Compound might be the most important chemistry term that is routinely misunderstood by beginning chemistry students.

Mistakenly, mixtures are often misidentified as compounds. Play close attention to the drawings of compounds below.




Figure 3.3.4: Nanoscopic Level Representations of Compounds

Two requirements must be met for a sample of matter to be called a compound. First, all of the particles must be the same. This is another way of saying that a compound is a pure substance. Second, the particle in a compound must include at least two different types of atoms.

Chemical Formulas



A chemical formula is a symbolic way of representing the atoms that compose a particle. A formula indicates both the elements present in a particle and the number of atoms of each element. We cannot write a chemical formula for a mixture since there are a multiple particles with different compositions present.



Chemical formulas consist of the symbols for the elements present, and each symbol is followed by a subscript. Symbols for elements are either a single upper case letter or an upper case letter followed by a lower case letter. Subscripts (small numbers written to the right of the symbol and sitting slightly lower than the symbol) indicate the number of that type of atom in the particle. Subscript of one are not written, so that a missing subscript is understood to mean there is only one of a particular type of atom in the particle.


Consider the following three fictitious elements and their symbols:

element			
symbol	Z	T	X

These symbols can be used to write the chemical formulas for some of the pure substances represented by particle-level drawings in this section:

particle-level drawing	chemical formula	type of pure substance
	X	element
	X_2	element

particle-level drawing	chemical formula	type of pure substance
	T_3	element
	ZTX	compound

particle-level drawing	chemical formula	type of pure substance
	ZT_2	compound

This page titled 3.3: Classifying Matter- the Nanoscopic Level is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.4: The Elements

Learning Objectives

- Define a chemical element.
- Represent a chemical element with a chemical symbol.

An element is a substance that cannot be broken down into simpler chemical substances. There are about 90 naturally occurring elements known on Earth. Using technology, scientists have been able to create nearly 30 additional elements that do not occur in nature. Today, chemistry recognizes 118 elements—some of which were created an atom at a time. Figure 3.4.1 shows some of the chemical elements.



Figure 3.4.1: Samples of Elements. Gold is a yellowish solid, iron is a silvery solid, while mercury is a silvery liquid at room temperature. © Thinkstock

Abundance

The elements vary widely in abundance. In the universe as a whole, the most common element is hydrogen (about 90% of atoms), followed by helium (most of the remaining 10%). All other elements are present in relatively minuscule amounts, as far as we can detect.

Table 3.4.1: Elemental Composition of Earth

Earth's Crust		Earth (overall)	
Element	**Percentage**	**Element**	**Percentage**
oxygen	46.1	iron	34.6
silicon	28.2	oxygen	29.5
aluminum	8.23	silicon	15.2
iron	5.53	magnesium	12.7
calcium	4.15	nickel	2.4
sodium	2.36	sulfur	1.9
magnesium	2.33	all others	3.7

Earth's Crust		Earth (overall)	
Element	Percentage	Element	Percentage
potassium	2.09		
titanium	0.565		
hydrogen	0.14		
phosphorus	0.105		
all others	0.174		
Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 14–17.			

On the planet Earth, however, the situation is rather different. Oxygen makes up 46.1% of the mass of Earth's crust (the relatively thin layer of rock forming Earth's surface), mostly in combination with other elements, while silicon makes up 28.2%. Hydrogen, the most abundant element in the universe, makes up only 0.14% of Earth's crust. Table 3.4.1 lists the relative abundances of elements on Earth as a whole and in Earth's crust. Table 3.4.2 lists the relative abundances of elements in the human body. If you compare Table 3.4.1 and Table 3.4.2, you will find disparities between the percentage of each element in the human body and on Earth. Oxygen has the highest percentage in both cases, but carbon, the element with the second highest percentage in the body, is relatively rare on Earth and does not even appear as a separate entry in Table 3.4.1; carbon is part of the 0.174% representing "other" elements. How does the human body concentrate so many apparently rare elements?

Table 3.4.2: Elemental Composition of a Human Body

Element	Percentage by Mass
oxygen	61
carbon	23
hydrogen	10
nitrogen	2.6
calcium	1.4
phosphorus	1.1
sulfur	0.20
potassium	0.20
sodium	0.14
chlorine	0.12
magnesium	0.027
silicon	0.026
iron	0.006
fluorine	0.0037
zinc	0.0033
all others	0.174
Source: D. R. Lide, ed. *CRC Handbook of Chemistry and Physics*, 89th ed. (Boca Raton, FL: CRC Press, 2008–9), 7–24.	

The relative amounts of elements in the body have less to do with their abundances on Earth than with their availability in a form we can assimilate. We obtain oxygen from the air we breathe and the water we drink. We also obtain hydrogen from water. On the other hand, although carbon is present in the atmosphere as carbon dioxide, and about 80% of the atmosphere is nitrogen, we obtain those two elements from the food we eat, not the air we breathe.

Names and Symbols

Each element has a name. Some of these names date from antiquity, while others are quite new. Today, the names for new elements are proposed by their discoverers but must be approved by the International Union of Pure and Applied Chemistry, an international organization that makes recommendations concerning all kinds of chemical terminology.

The names of the elements can be cumbersome to write in full, especially when combined to form the names of compounds. Therefore, each element name is abbreviated as a one- or two-letter chemical symbol. By convention, the first letter of a chemical symbol is a capital letter, while the second letter (if there is one) is a lowercase letter. The first letter of the symbol is usually the first letter of the element's name, while the second letter is some other letter from the name. Some elements have symbols that derive from earlier, mostly Latin names, so the symbols may not contain any letters from the English name. Table 3.4.3 lists the names and symbols of some of the most familiar elements.

Table 3.4.3: Element Names and Symbols

aluminum	Al	magnesium	Mg
argon	Ar	manganese	Mn
arsenic	As	mercury	Hg*
barium	Ba	neon	Ne
bismuth	Bi	nickel	Ni
boron	B	nitrogen	N
bromine	Br	oxygen	O
calcium	Ca	phosphorus	P
carbon	C	platinum	Pt
chlorine	Cl	potassium	K*
chromium	Cr	silicon	Si
copper	Cu*	silver	Ag*
fluorine	F	sodium	Na*
gold	Au*	strontium	Sr
helium	He	sulfur	S
hydrogen	H	tin	Sn*
iron	Fe	tungsten	W†
iodine	I	uranium	U
lead	Pb*	zinc	Zn
lithium	Li	zirconium	Zr
***The symbol comes from the Latin name of element.**			
†The symbol for tungsten comes from its German name—*wolfram*.			

Symbols for elements that seem odd for English speakers in many cases make sense to speakers of languages such as Spanish and French that are descended from Latin. For example, gold is *oro* in Spanish and *or* in French (close to the Latin *aurum*), tin is *estaño* in Spanish (compare to *stannum*), lead is *plomo* in Spanish and *plomb* in French (compare to *plumbum*), silver is *argent* in French (compare to *argentum*), and iron is *fer* in French and *hierro* in Spanish (compare to *ferrum*). The closeness is even more apparent in pronunciation than in spelling.

Example 3.4.1

Write the chemical symbol for each element without consulting Table 3.4.3 "Element Names and Symbols".

a. bromine
b. boron
c. carbon
d. calcium
e. gold

Answer a available at the end of this section.

Exercise 3.4.1

Write the chemical symbol for each element without consulting Table 3.4.3.

a. manganese
b. magnesium
c. neon
d. nitrogen
e. silver

Answer a available at the end of this section.

Example 3.4.2

What element is represented by each chemical symbol?

a. Na
b. Hg
c. P
d. K
e. I

Answer a available at the end of this section.

Exercise 3.4.2

What element is represented by each chemical symbol?

a. Pb
b. Sn
c. U
d. O
e. F

Answer a available at the end of this section.

Contributors and Attributions

- Anonymous

Answers

Example 3.4.1

Br
B
C
Ca
Au

Exercise 3.4.1

Mn
Mg
Ne
N
Ag

Example 3.4.2

sodium
mercury
phosphorus
potassium
iodine

Exercise 3.4.2

lead
tin
uranium
oxygen
fluorine

This page titled 3.4: The Elements is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Anonymous.

- 2.1: The Elements by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

3.5: Physical and Chemical Changes- the Macroscopic Level

Learning Objectives

- Recognize physical evidence that a chemical reaction has occurred.

Change is happening all around us all of the time. Just as chemists have classified elements and compounds, they have also classified types of changes. Changes are classified as either physical or chemical changes. Chemists learn a lot about the nature of matter by studying the changes that matter can undergo. Chemists make a distinction between two different types of changes that they study—physical changes and chemical changes.

Physical Change

Physical changes are changes in which no chemical bonds are broken or formed. This means that the same types of compounds or elements that were there at the beginning of the change are there at the end of the change. Physical changes involve moving molecules around, but not changing them. Some types of physical changes include:

- Changes of state (changes from a solid to a liquid or a gas and vice versa).
- Separation of a mixture.
- Physical deformation (cutting, denting, stretching).
- Making solutions (dissolving one substance in another).

As an ice cube melts, its shape changes as it acquires the ability to flow. However, its composition does not change. Melting is an example of a physical change. A physical change is a change to a sample of matter in which some properties of the material change, but the identity of the matter does not. When liquid water is heated, it changes to water vapor. However, even though the physical properties have changed, the molecules are exactly the same as before. We still have each water molecule containing two hydrogen atoms and one oxygen atom covalently bonded. When you have a jar containing a mixture of pennies and nickels and you sort the mixture so that you have one pile of pennies and another pile of nickels, you have not altered the identity of the pennies or the nickels—you've merely separated them into two groups. This would be an example of a physical change. Similarly, if you have a piece of paper, you don't change it into something other than a piece of paper by ripping it up. What was paper before you started tearing is still paper when you are done. Again, this is an example of a physical change.



Figure 3.5.1: Ice melting is a physical change. When liquid water (H_2O) freezes into a solid state (ice), it appears changed; however, this change is only physical, as the composition of the constituent molecules is the same: 11.19% hydrogen and 88.81% oxygen by mass. (Public Domain; Moussa).

Physical changes can further be classified as reversible or irreversible. The melted ice cube may be refrozen, so melting is a reversible physical change. Physical changes that involve a change of state are all reversible. Other changes of state include **vaporization** (liquid to gas), **freezing** (liquid to solid), and **condensation** (gas to liquid). Dissolving is also a reversible physical change. When salt is dissolved into water, the salt is said to have entered the aqueous state. The salt may be regained by boiling off the water, leaving the salt behind.

Chemical Change

Chemical changes occur when bonds are broken and/or formed between molecules or atoms. This means that one substance with a certain set of properties (such as melting point, color, taste, etc) is turned into a different substance with different properties. Chemical changes are frequently harder to reverse than physical changes.

One good example of a chemical change is burning a candle. The act of burning a candle results in the formation of new compounds (carbon dioxide and water) from the burning of the wax and oxygen in the air. Not only has the appearance changes, but the structure of the molecules have also changed. The new substances do not have the same chemical properties as the original ones.



Figure 3.5.2: Burning of wax to generate water and carbon dioxide is a chemical reaction. (CC-SA-BY-3.0; Andrikkos)

We can't actually see molecules breaking and forming bonds, although that's what defines chemical changes. We have to make other observations to indicate that a chemical change has happened. Some of the evidence for chemical change will involve the energy changes that occur in chemical changes, but some evidence involves the fact that new substances with different properties are formed in a chemical change.

Observations that help to indicate chemical change include:

- Temperature changes (either the temperature increases or decreases).
- Light given off.
- Unexpected color changes (a substance with a different color is made, rather than just mixing the original colors together).
- Bubbles are formed (but the substance is not boiling—you made a substance that is a gas at the temperature of the beginning materials, instead of a liquid).
- Different smell or taste (do not taste your chemistry experiments, though!).
- A solid forms if two clear liquids are mixed (look for *floaties*—technically called a precipitate).

Example 3.5.1

Label each of the following changes as a physical or chemical change. Give evidence to support your answer.

a. Boiling water.
b. A nail rusting.
c. A green solution and colorless solution are mixed. The resulting mixture is a solution with a pale green color.
d. Two colorless solutions are mixed. The resulting mixture has a yellow precipitate.

Solution

a. Physical: boiling and melting are physical changes. When water boils, no bonds are broken or formed. The change could be written: $\mathrm{H_2O}\,(l) \rightarrow \mathrm{H_2O}\,(g)$
b. Chemical: The dark grey nail changes color to form an orange flaky substance (the rust); this must be a chemical change. Color changes indicate chemical change. The following reaction occurs: $\mathrm{Fe} + \mathrm{O_2} \rightarrow \mathrm{Fe_2O_3}$
c. Physical: because none of the properties changed, this is a physical change. The green mixture is still green and the colorless solution is still colorless. They have just been spread together. No color *change* occurred or other evidence of chemical change.
d. Chemical: the formation of a precipitate and the color change from colorless to yellow indicate a chemical change.

Exercise 3.5.1

Label each of the following changes as a physical or chemical change.

a. A mirror is broken.
b. An iron nail corroded in moist air
c. Copper metal is melted.
d. A catalytic converter changes nitrogen dioxide to nitrogen gas and oxygen gas.

Answer a: available at the end of this section.

Answer b: available at the end of this section.

Answer c: available at the end of this section.

Answer d: available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

-

- Boundless (www.boundless.com)
- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise 3.5.1
physical change

Exercise 3.5.1
chemical change

Exercise 3.5.1
physical change

Exercise 3.5.1
chemical change

This page titled 3.5: Physical and Chemical Changes- the Macroscopic Level is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.6: Physcial and Chemical Changes- the Nanoscopic Level

Learning Objectives

- Classify changes in matter as physical or chemical on the particle level.

Physical Changes

Chemical and physical changes can also be understood using nanoscopic level illustrations. Representations of three physical changes are shown below. In all of these illustrations, the arrow represents the process of transformation. What is to the left of the arrow can be thought of as a "before" snapshot and what is to the right of the arrow can be thought of as an "after" snapshot.




Figure 3.6.1: Nanoscopic Level Representations of Physical Changes

The first illustration represents a phase change. The left square represents a solid and the right square represents a gas. Since the same particle--consisting of two hexagons--makes up both the solid and the gas, this is a physical change. The second illustration represents a solid being broken into smaller pieces. The particle in both the left square and the right square consists of a single circle. Finally, the third illustration represents a solid dissolving in a liquid. Neither particle represented--the one formed from a single circle or the one formed from two squares--changes during the process of dissolving.

What all of these changes have in common is that the particles themselves remain unaltered though the transformation. Whatever particles you start with in the left square, you end with in the right square. Another way of saying this is that no new pure substances are formed.

Chemical Changes

Nanoscopic level representations of three chemical changes are shown below.




Figure 3.6.2: Nanoscopic Level Representations of Chemical Changes

What all these illustrations have in common is that new particles are formed while moving from the left square to the right square. In the first transformation a particle with a circle and a triangle is formed that did not originally exist. In the second illustration, a particle is formed that contains two triangles and another particle is formed that contains a single square. Originally, the only particle present contained a square and a triangle. Finally, a particle containing two hexagons is formed from a sample that originally contained particles with three hexagons.

The first illustration shows that a chemical change can sometimes include a change in state: here a solid disappears as it reacts and becomes part of the gas phase. The final illustration shows that it is possible for one sample of an element to change to another sample of an element. Even though the atoms themselves do not change, the way they bond together to form a particle is altered.

This page titled 3.6: Physcial and Chemical Changes- the Nanoscopic Level is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.7: Separating Mixtures through Physcial Changes

There are many techniques that use physical changes to separate mixtures. Here are a number of common separation techniques:

Distillation

Distillation is an effective method to separate mixtures comprised of two or more pure liquids. Distillation is a purification process where one component of a liquid mixture is vaporized and then condensed and isolated. In simple distillation Figure 3.7.1 , a mixture is heated and the most volatile component vaporizes at the lowest temperature. The vapor passes through a cooled tube (a condenser), where it condenses back into its liquid state. The condensate that is collected is called distillate.

Outside the chemistry lab, distillation is commonly used to increase the alcohol content of liquors such as vodka, whiskey, and brandy.




Figure 3.7.1: Distillation apparatus.

Evaporation

Evaporation is a technique used to separate out a soluble solid from a liquid. The method drives off the liquid components from the solid components. The process typically involves heating the mixture until no more liquid remains. Evaporation can be used, for example, to obtain table salt from sea water, Figure 3.7.2. The heat for the process comes from the sun.



Figure 3.7.2: Once the sea water in these evaporation ponds has evaporated, the salt can be harvested.

Filtration

Filtration is a separation method used to separate out pure substances in mixtures comprised of particles, some of which are large enough in size to be captured with a porous material. Particle size can vary considerably, given the type of mixture. For instance, stream water is a mixture that contains naturally occurring biological organisms like bacteria, viruses, and protozoans. Some water filters can filter out bacteria, the length of which is on the order of 1 micron. Other mixtures, like soil, have relatively large particle sizes, which can be filtered through something like a coffee filter.




Figure 3.7.3: Filtration.

Chromatography

Chromatography is the separation of a liquid or gas mixture by passing it through a medium in which the components move at different rates. Thin-layer chromatography is perhaps the simplest form of chromatography. It uses a plate--a plastic or glass slide covered on one side with silica (essentially very fine, purified sand), alumina, or some other solid. The plate is placed so that the bottom of it is submerged in a solvent, and the solvent then moves upward on the plate. If a drop of a mixture is placed a little above the level of the solvent, different components of the mixture will be carried upward at different rates depending on

how strongly they interact with the silica. Figure 3.7.4 shows separation of the chromophores (colored compounds) in a spinach leaf by thin layer chromatography.The pencil line at the bottom of the plate shows the original position of the mixture before the solvent started traveling up the plate.




Figure 3.7.4: Separation of compounds in a spinach leaf. (photo credit: Heather Coleman)

This page titled 3.7: Separating Mixtures through Physcial Changes is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

3.8: Differences in Matter- Physical and Chemical Properties

Learning Objectives

Given a property of matter, classify as physical or chemical.

All matter has physical and chemical properties. Physical properties are characteristics that scientists can measure without changing the composition of the sample under study, such as mass, color, and volume (the amount of space occupied by a sample). Chemical properties describe the characteristic ability of a substance to react to form new substances; they include its flammability and susceptibility to corrosion. All samples of a pure substance have the same chemical and physical properties. For example, pure copper is always a reddish-brown solid (a physical property) and always dissolves in dilute nitric acid to produce a blue solution and a brown gas (a chemical property).

Physical Property

A *physical property* is a characteristic of a substance that can be observed or measured without changing the identity of the substance. Silver is a shiny metal that conducts electricity very well. It can be molded into thin sheets, a property called malleability. Salt is dull and brittle and conducts electricity when it has been dissolved into water, which it does quite easily. Physical properties of matter include color, hardness, malleability, solubility, electrical conductivity, density, melting point, and boiling point.

For the elements, color does not vary much from one element to the next. The vast majority of elements are colorless, silver, or gray. Some elements do have distinctive colors: sulfur and chlorine are yellow, copper is (of course) copper-colored, and elemental bromine is red. However, density can be a very useful parameter for identifying an element. Of the materials that exist as solids at room temperature, iodine has a very low density compared to zinc, chromium, and tin. Gold has a very high density, as does platinum. Pure water, for example, has a density of 0.998 g/cm^3 at 25°C. The average densities of some common substances are in Table 3.8.1. Notice that corn oil has a lower mass to volume ratio than water. This means that when added to water, corn oil will "float."

Table 3.8.1: Densities of Common Substances

Substance	Density at 25°C (g/cm3)
blood	1.035
body fat	0.918
whole milk	1.030
corn oil	0.922
mayonnaise	0.910
honey	1.420

Hardness helps determine how an element (especially a metal) might be used. Many elements are fairly soft (silver and gold, for example) while others (such as titanium, tungsten, and chromium) are much harder. Carbon is an interesting example of hardness. In graphite, (the "lead" found in pencils) the carbon is very soft, while the carbon in a diamond is roughly seven times as hard.



Figure 3.8.1: The graphite in a pencil (left) and the diamond in a diamond ring (right). Both are a form of carbon, but exhibit very different physical properties.

Melting and boiling points are somewhat unique identifiers, especially of compounds. In addition to giving some idea as to the identity of the compound, important information can be obtained about the purity of the material.

Chemical Properties

Chemical properties of matter describe its potential to undergo some chemical change or reaction by virtue of its composition. The elements, electrons, and bonds that are present give the matter potential for chemical change. It is quite difficult to define a chemical property without using the word "change". Eventually, after studying chemistry for some time, you should be able to look at the formula of a compound and state some chemical property. For example, hydrogen has the potential to ignite and explode given the right conditions—this is a chemical property. Metals in general have the chemical property of reacting with an acid. Zinc reacts with hydrochloric acid to produce hydrogen gas—this is a chemical property.



Figure 3.8.2: Heavy rust on the links of a chain near the Golden Gate Bridge in San Francisco; it was continuously exposed to moisture and salt spray, causing surface breakdown, cracking, and flaking of the metal. (CC BY-SA 3.0; Marlith).

A chemical property of iron is its capability of combining with oxygen to form iron oxide, the chemical name of rust (Figure 3.8.2).

The more general term for rusting and other similar processes is corrosion. Other terms that are commonly used in descriptions of chemical changes are burn, rot, explode, decompose, and ferment. Chemical properties are very useful in identifying substances. However, unlike physical properties, chemical properties can only be observed as the substance is in the process of being changed into a different substance.

Table 3.8.2: Contrasting Physical and Chemical Properties

Physical Properties	Chemical Properties
Gallium metal melts at 30 °C.	Iron metal rusts.
Mercury is a very dense liquid.	A green banana turns yellow when it ripens.
Gold is shiny.	A dry piece of paper burns.

Example 3.8.1

Which of the following is a chemical property of iron?

a. Iron corrodes in moist air.
b. Density = 7.874 g/cm^3
c. Iron is soft when pure.
d. Iron melts at 1808 K.

Solution

"Iron corrodes in moist air" is the only chemical property of iron from the list.

Exercise 3.8.1A

Which of the following is a physical property of matter?

a. corrosiveness
b. pH (acidity)
c. density
d. flammability

Answer available at the end of this section.

Exercise 3.8.1B

Which of the following is a chemical property?

a. flammability
b. melting point
c. boiling point
d. density

Answer available at the end of this section.

Contributions & Attributions

This page was constructed from content via the following contributor(s) and edited (topically or extensively) by the LibreTexts development team to meet platform style, presentation, and quality:

- Marisa Alviar-Agnew (Sacramento City College)
- Henry Agnew (UC Davis)

Answers

Exercise $3.8.1A$

c

Exercise $3.8.1B$

a

This page titled 3.8: Differences in Matter- Physical and Chemical Properties is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Marisa Alviar-Agnew and Henry Agnew.

4: ATOMS AND THE PERIODIC TABLE

Just as a language has an alphabet from which words are built, chemistry has an alphabet from which matter is described. However, the chemical alphabet is larger than the one we use for spelling. You may have already figured out that the chemical alphabet consists of the chemical elements. Their role is central to chemistry, for they combine to form the millions and millions of known compounds.

4.1: Prelude to Atoms, and the Periodic Table

4.2: Atomic Theory

4.3: The Structure of Atoms

4.4: Nuclei of Atoms

4.5: Atomic Masses

4.6: Bohr Diagrams of Atoms and Ions

4.7: Electron Arrangements

4.8: The Periodic Table

4.S: Elements, Atoms, and the Periodic Table (Summary)

Template:HideTOC

This page titled 4: Atoms and the Periodic Table is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2: Elements, Atoms, and the Periodic Table by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.1: Prelude to Atoms, and the Periodic Table

The hardest material in the human body is tooth enamel. It has to be hard so that our teeth can serve us for a lifetime of biting and chewing; however, tough as it is, tooth enamel is susceptible to chemical attack. Acids found in some foods or made by bacteria that feed on food residues on our teeth are capable of dissolving enamel. Unprotected by enamel, a tooth will start to decay, thus developing cavities and other dental problems.

In the early 1900s, a dentist in Colorado Springs, Colorado, noted that many people who lived in the area had brown-stained teeth that, while unsightly, were surprisingly resistant to decay. After years of study, excess fluorine compounds in the drinking water were discovered to be the cause of both these effects. Research continued, and in the 1930s, the US Public Health Service found that low levels of fluorine in water would provide the benefit of resisting decay without discoloring teeth.

The protective effects of fluorine have a simple chemical explanation. Tooth enamel consists mostly of a mineral called hydroxyapatite, which is composed of calcium, phosphorus, oxygen, and hydrogen. We know now that fluorine combines with hydroxyapatite to make fluorapatite, which is more resistant to acidic decay than hydroxyapatite is. Currently about 50% of the US population drinks water that has some fluorine added (in the form of sodium fluoride, NaF) to reduce tooth decay. This intentional fluoridation, coupled with the use of fluoride-containing toothpastes and improved oral hygiene, has reduced tooth decay by as much as 60% in children. The nationwide reduction of tooth decay has been cited as an important public health advance in history. (Another important advance was the eradication of polio.)

This page titled 4.1: Prelude to Atoms, and the Periodic Table is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.0: Prelude to Elements, Atoms, and the Periodic Table by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.2: Atomic Theory

Learning Objectives

- Explain how all matter is composed of atoms.
- Describe the modern atomic theory.
- Recognize which elements exist as diatomic molecules.

Take some aluminum foil. Cut it in half. Now you have two smaller pieces of aluminum foil. Cut one of the pieces in half again. Cut one of those smaller pieces in half again. Continue cutting, making smaller and smaller pieces of aluminum foil.




Figure 4.2.1: Aluminum foil is made up of aluminium atoms. (CC BY-NC-SA 2.0; James Milstid via Flickr)

It should be obvious that the pieces are still aluminum foil; they are just becoming smaller and smaller. But how far can you take this exercise, at least in theory? Can you continue cutting the aluminum foil into halves forever, making smaller and smaller pieces? Or is there some limit, some absolute smallest piece of aluminum foil? (Thought experiments like this—and the conclusions based on them—were debated as far back as the fifth century BC.)

The modern atomic theory, proposed about 1803 by the English chemist John Dalton (Figure 4.2.2), is a fundamental concept that states that all elements are composed of atoms. Previously, we defined an atom as the smallest part of an element that maintains the identity of that element. Individual atoms are extremely small; even the largest atom has an approximate diameter of only 5.4×10^{-10} m. With that size, it takes over 18 million of these atoms, lined up side by side, to equal the width of your little finger (about 1 cm).




Figure 4.2.2 John Dalton was an English scientist who enunciated the modern atomic theory.

Most elements in their pure form exist as **individual atoms**. For example, a macroscopic chunk of iron (Fe) metal is composed, microscopically, of individual atoms. Some elements, however, exist as groups of atoms called molecules. Several important elements exist as two-atom combinations and are called **diatomic molecules**. In representing a diatomic molecule, we use the symbol of the element and include the ***subscript*** "2" to indicate that two atoms of that element are joined together. The elements that exist as diatomic molecules are hydrogen (H_2), nitrogen (N_2), fluorine (F_2), oxygen (O_2), iodine (I_2), chlorine (Cl_2) and bromine (Br_2). A few other elements can exist as 3-atom molecules like ozone (O_3) and 4-atom molecules like phosphorus (P_4). The most common form of the element sulfur is composed of molecules that consist of eight atoms of sulfur; its molecular formula is S_8 (Figure 4.2.3).




Figure 4.2.3: A molecule of sulfur is composed of eight sulfur atoms and is therefore written as S_8. It can be represented as (a) a structural formula, (b) a ball-and-stick model, and (c) a space-filling model. Sulfur atoms are represented by yellow spheres.

It is important to note that a subscript following a symbol and a number in front of a symbol do not represent the same thing; for example, H_2 and 2H represent distinctly different species. H_2 is a molecular formula; it represents a diatomic molecule of hydrogen, consisting of two atoms of the element that are chemically bonded together. The expression 2H, on the other hand, indicates two separate hydrogen atoms that are **not** combined as a unit. The expression $2H_2$ represents two molecules of diatomic hydrogen (Figure 4.2.4).




Figure 4.2.4: The symbols H, 2H, H_2, and $2H_2$ represent very different entities.

Example 4.2.1

Write the chemical formula of the following elements:

a. oxygen
b. carbon
c. potassium

Solution

Memorizing the diatomic molecules is worthwhile for our future endeavors. A mnemonic device to help in the memorization of the diatomic elements is as follows: **H**ave **N**o **F**ear **O**f **I**ce **C**old **B**eer. (hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine).

a. Oxygen is one of the seven diatomic molecular elements. Its chemical formula is O_2.
b. Carbon is monatomic, hence its formula is C.
c. Potassium is monatomic hence its formula is K.

Exercise 4.2.1

Write the chemical formula of the following elements:

a. hydrogen
b. nitrogen
c. neon

Answer a available at the end of this section.

Looking Closer: Atomic Theory

Dalton's ideas are called the *modern* atomic theory because the concept of atoms is very old. The Greek philosophers Leucippus and Democritus originally introduced atomic concepts in the fifth century BC. (The word *atom* comes from the Greek word *atomos*, which means "indivisible" or "uncuttable.") Dalton had something that the ancient Greek philosophers didn't have, however; he had experimental evidence, such as the formulas of simple chemicals and the behavior of gases. In the 150 years or so before Dalton, natural philosophy had been maturing into modern science, and the scientific method was being used to study nature. So when Dalton announced a modern atomic theory, he was proposing a fundamental theory to describe many previous observations of the natural world; he was not just participating in a philosophical discussion.

Concept Review Exercises

1. What is the modern atomic theory?

2. What are atoms?

Answers

1. The modern atomic theory states that all matter is composed of atoms.
2. Atoms are the smallest parts of an element that maintain the identity of that element.

Key Takeaways

- **Atoms** are the ultimate building blocks of all matter.
- The modern atomic theory establishes the concepts of **atoms** and how they compose matter.
- The **diatomic** elements are hydrogen, nitrogen, fluorine, oxygen, iodine, chlorine and bromine.

Answers

Exercise 4.2.1

Hydrogen is one of the seven diatomic molecular elements. Its chemical formula is H_2.

Nitrogen is one of the seven diatomic molecular elements. Its chemical formula is N_2.

Neon is a monatomic element, hence its formula is Ne.

This page titled 4.2: Atomic Theory is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.2: Atomic Theory by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.3: The Structure of Atoms

Learning Objectives

- Describe the three main subatomic particles.
- State how the subatomic particles are arranged in atoms.

There have been several minor but important modifications to Dalton's atomic theory. For one thing, Dalton considered atoms to be indivisible. We know now that atoms not only can be divided but also are composed of three different kinds of particle, *subatomic particles*, with their own properties that are different from the chemical properties of atoms.

Subatomic Particles

The first subatomic particle was identified in 1897 and called the electron. It is an extremely tiny particle, with a mass of about 9.109×10^{-31} kg. Experiments with magnetic fields showed that the electron has a negative electrical charge. By 1920, experimental evidence indicated the existence of a second particle. A proton has the same amount of charge as an electron, but its charge is positive, not negative. Another major difference between a proton and an electron is mass. Although still incredibly small, the mass of a proton is 1.673×10^{-27} kg, which is almost 2,000 times greater than the mass of an electron. Because opposite charges attract each other (while like charges repel each other), protons attract electrons (and vice versa).

Finally, additional experiments pointed to the existence of a third particle. Evidence produced in 1932 established the existence of the neutron, a particle with about the same mass as a proton but with no electrical charge, it is *neutral*. We understand now that all atoms can be broken down into subatomic particles: protons, neutrons, and electrons. Table $4.3.1$ lists some of their important characteristics and the symbols used to represent each particle.

Table $4.3.1$: Properties of the Subatomic Particles

Particle	Symbol	Mass (kg)	Relative Mass (proton = 1)	Relative Charge	Location
proton	p^+	1.673×10^{-27}	1	+1	in the nucleus
neutron	n^0	1.675×10^{-27}	1	0	in the nucleus
electron	e^-	9.109×10^{-31}	0.00055	–1	outside the nucleus

The Nucleus

How are these subatomic particles arranged? Between 1909 and 1911, Ernest Rutherford, a Cambridge physicist, and his associates Hans Geiger and Ernest Marsden performed experiments that provided strong evidence concerning the internal structure of an atom. They took a very thin metal foil, such as gold or platinum, and aimed a beam of positively charged particles (called alpha particles, which are combinations of two protons and two neutrons) from a radioactive source toward the foil. Surrounding the foil was a detector—either a scintillator (a material that glows when hit by such particles) or some unexposed film (which is exposed where the particles hit it). The detector allowed the scientists to determine the distribution of the alpha particles after they interacted with the foil. Figure $4.3.1$ shows a diagram of the experimental setup.




Figure 4.3.1: The Geiger-Marsden Experimental Setup. Experiments using this setup were used to investigate the structure of atoms. In this experiment, most of the particles traveled straight through the foil, but some alpha particles were deflected off to one side. Some were even deflected back toward the source. This was unexpected. Rutherford once said, "It was almost as incredible as if you fired a 15-inch shell at a piece of tissue paper and it came back and hit you." (CC SA-BY 3.0; Kurzon).

Rutherford proposed the following model to explain these experimental results. Protons and neutrons are concentrated in a central region he called the nucleus (plural, *nuclei*) of the atom. Electrons are outside the nucleus and orbit about it because they are attracted to the positive charge in the nucleus. Most of the mass of an atom is in the nucleus, while the orbiting electrons account for an atom's size. As a result, an atom consists largely of empty space. Rutherford called his description the "planetary model" of the atom. Figure 4.3.2 shows how this model explains the experimental results.




Figure 4.3.2: Rutherford's Metal Foil Experiments. Rutherford explained the results of the metal foil experiments by proposing that most of the mass and the positive charge of an atom are located in its nucleus, while the relatively low-mass electrons orbit about the nucleus. Most alpha particles go straight through the empty space, a few particles are deflected, and fewer still ricochet back toward the source. The nucleus is much smaller proportionately than depicted here.

> The planetary model of the atom replaced the plum pudding model, which had electrons floating around aimlessly like plums in a "pudding" of positive charge.

Rutherford's model is essentially the same model that we use today to describe atoms but with one important modification. The planetary model suggests that electrons occupy certain specific, circular orbits about the nucleus. We know now that this model is

overly simplistic. A better description is that electrons form fuzzy clouds around nuclei. Figure 4.3.3 shows a more modern version of our understanding of atomic structure.




Figure 4.3.3: A Modern Depiction of Atomic Structure. A more modern understanding of atoms, reflected in these representations of the electron in a hydrogen atom, is that electrons occupy regions of space about the nucleus; they are not in discrete orbits like planets around the sun. (a) The darker the color, the higher the probability that an electron will be at that point. (b) In a two-dimensional cross section of the electron in a hydrogen atom, the more crowded the dots, the higher the probability that an electron will be at that point. In both (a) and (b), the nucleus is in the center of the diagram.

Concept Review Exercises

1. What are the charges and the relative masses of the three subatomic particles?
2. Describe the structure of an atom in terms of its protons, neutrons, and electrons.

Answers

1. proton: +1, large; neutron: 0, large; electron: –1, small
2. Protons and neutrons are located in a central nucleus, while electrons orbit about the nucleus.

Key Takeaways

- Atoms are composed of three main subatomic particles: protons, neutrons, and electrons.
- Protons and neutrons are grouped together in the nucleus of an atom, while electrons orbit about the nucleus.

This page titled 4.3: The Structure of Atoms is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.3: The Structure of Atoms by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.4: Nuclei of Atoms

Learning Objectives

- Define and differentiate between the atomic number and the mass number of an element.
- Explain how isotopes differ from one another.

Now that we know how atoms are generally constructed, what do atoms of any particular element look like? How many protons, neutrons, and electrons are in a specific kind of atom? First, if an atom is electrically neutral overall, then the number of protons equals the number of electrons. Because these particles have the same but opposite charges, equal numbers cancel out, producing a neutral atom.

Atomic Number

In the 1910s, experiments with x-rays led to this useful conclusion: the magnitude of the positive charge in the nucleus of every atom of a particular element is the same. In other words, all atoms of the same element have the same number of protons. Furthermore, different elements have a different number of protons in their nuclei, so the number of protons in the nucleus of an atom is characteristic of a particular element. This discovery was so important to our understanding of atoms that the number of protons in the nucleus of an atom is called the atomic number (Z).

For example, hydrogen has the atomic number 1; all hydrogen atoms have 1 proton in their nuclei. Helium has the atomic number 2; all helium atoms have 2 protons in their nuclei. There is no such thing as a hydrogen atom with 2 protons in its nucleus; a nucleus with 2 protons would be a helium atom. The atomic number *defines* an element. Table $4.4.1$ lists some common elements and their atomic numbers. Based on its **atomic number**, you can determine the **number of protons in the nucleus** of an atom. The largest atoms have over 100 protons in their nuclei.

Example $4.4.1$

What is the number of protons in the nucleus of each element?

a. aluminum
b. iron
c. carbon

Answer a available at the end of this section.

Exercise $4.4.1$

What is the number of protons in the nucleus of each element? Use Table 2.4.1.

a. sodium
b. oxygen
c. chlorine

Answer a available at the end of this section.

How many electrons are in an atom? Previously we said that for an electrically neutral atom, the **number of electrons equals the number of protons**, so the total opposite charges cancel. Thus, the **atomic number** of an element also gives the **number of electrons** in an atom of that element. (Later we will find that some elements may gain or lose electrons from their atoms, so those

atoms will no longer be electrically neutral. Thus we will need a way to differentiate the number of electrons for those elements.)

Example 4.4.2

How many electrons are present in the atoms of each element?

a. sulfur
b. tungsten
c. argon

Answer a available at the end of this section.

Exercise 4.4.2

How many electrons are present in the atoms of each element?

a. magnesium
b. potassium
c. iodine

Answer a available at the end of this section.

Table 4.4.1: Some Common Elements and Their Atomic Numbers

Element	Atomic Number	Element	Atomic Nmbers
aluminum (Al)	13	magnesium (Mg)	12
americium (Am)	95	manganese (Mn)	25
argon (Ar)	18	mercury (Hg)	80
barium (Ba)	56	neon (Ne)	10
beryllium (Be)	4	nickel (Ni)	28
bromine (Br)	35	nitrogen (N)	7
calcium (Ca)	20	oxygen (O)	8
carbon (C)	6	phosphorus (P)	15
chlorine (Cl)	17	platinum (Pt)	78
chromium (Cr)	24	potassium (K)	19
cesium (Cs)	55	radon (Rn)	86
fluorine (F)	9	silver (Ag)	47
gallium (Ga)	31	sodium (Na)	11
gold (Au)	79	strontium (Sr)	38
helium (He)	2	sulfur (S)	16
hydrogen (H)	1	titanium (Ti)	22
iron (Fe)	26	tungsten (W)	74
iodine (I)	53	uranium (U)	92
lead (Pb)	82	zinc (Zn)	30
lithium (Li)	3	zirconium (Zr)	40

Isotopes

How many neutrons are in atoms of a particular element? At first it was thought that the number of neutrons in a nucleus was also characteristic of an element. However, it was found that atoms of the same element can have *different* numbers of neutrons. Atoms of the same element (i.e., same atomic number, Z) that have different numbers of neutrons are called isotopes. For example, 99% of the carbon atoms on Earth have 6 neutrons and 6 protons in their nuclei; about 1% of the carbon atoms have 7 neutrons in their nuclei. Naturally occurring carbon on Earth, therefore, is actually a mixture of isotopes, albeit a mixture that is 99% carbon with 6 neutrons in each nucleus.




Figure 4.4.1: Isotopes of Hydrogen. Most hydrogen atoms have only a proton in the nucleus (a). A small amount of hydrogen exists as the isotope deuterium, which has one proton and one neutron in its nucleus (b). A tiny amount of the hydrogen isotope tritium, with one proton and two neutrons in its nucleus, also exists on Earth (c). The nuclei and electrons are proportionately much smaller than depicted here.

An important series of isotopes is found with hydrogen atoms. Most hydrogen atoms have a nucleus with only a single proton. About 1 in 10,000 hydrogen nuclei, however, also has a neutron; this particular isotope is called *deuterium*. An extremely rare hydrogen isotope, *tritium*, has 1 proton and 2 neutrons in its nucleus. Figure 4.4.1 compares the three isotopes of hydrogen.

> The discovery of isotopes required a minor change in Dalton's atomic theory. Dalton thought that all atoms of the same element were exactly the same.

Most elements exist as mixtures of isotopes. In fact, there are currently over 3,500 isotopes known for all the elements. When scientists discuss individual isotopes, they need an efficient way to specify the number of neutrons in any particular nucleus. The *mass number* (A) of an atom is the sum of the numbers of protons and neutrons in the nucleus. Given the mass number for a nucleus (and knowing the atomic number of that particular atom), you can determine the number of neutrons by subtracting the atomic number from the mass number.

A simple way of indicating the mass number of a particular isotope is to list it as a superscript on the left side of an element's symbol. Atomic numbers are often listed as a subscript on the left side of an element's symbol. Thus, we might see

$$\begin{matrix}\text{mass number}\longrightarrow 56\\ \text{atomic number}\longrightarrow 26\end{matrix}\mathrm{Fe} \tag{4.4.1}$$

which indicates a particular isotope of iron. The 26 is the atomic number (which is the same for all iron atoms), while the 56 is the mass number of the isotope. To determine the number of neutrons in this isotope, we subtract 26 from 56: 56 – 26 = 30, so there are 30 neutrons in this atom.

Example 4.4.3

How many protons and neutrons are in each atom?

a. $^{35}_{17}\mathrm{Cl}$

b. $^{127}_{53}\mathrm{I}$

Answer a available at the end of this section.

Exercise 4.4.3

How many protons and neutrons are in each atom?

a. $^{197}_{79}\mathrm{Au}$
b. $^{23}_{11}\mathrm{Na}$

Answer a available at the end of this section.

It is not absolutely necessary to indicate the atomic number as a subscript because each element has its own unique atomic number. Many isotopes are indicated with a superscript only, such as ^{13}C or ^{235}U. You may also see isotopes represented in print as, for example, carbon-13 or uranium-235.

Summary

The atom consists of discrete particles that govern its chemical and physical behavior. Each atom of an element contains the same number of protons, which is the **atomic number (Z)**. Neutral atoms have the same number of electrons and protons. Atoms of an element that contain different numbers of neutrons are called **isotopes**. Each isotope of a given element has the same atomic number but a different **mass number (A)**, which is the **sum of the numbers of protons and neutrons**.

Almost all of the mass of an atom is from the total protons and neutrons contained within a tiny (and therefore very dense) nucleus. The majority of the volume of an atom is the surrounding space in which the electrons reside. A representation of a carbon-12 atom is shown below in Figure 4.4.2.




Figure 4.4.2: Formalism used for identifying specific nuclide (any particular kind of nucleus)

The atomic symbol, X, is the abbreviation used to represent an atom in chemical formulas. The mass number, A, is the number of protons and neutrons in the atom, which is to the left of the X. The atomic number, Z, is the number of protons in the atom, which is to the left of the X and below the A.

Concept Review Exercises

1. Why is the atomic number so important to the identity of an atom?
2. What is the relationship between the number of protons and the number of electrons in an atom?
3. How do isotopes of an element differ from each other?
4. What is the mass number of an element?

Answers

1. The atomic number defines the identity of an element. It describes the number of protons in the nucleus.
2. In an electrically neutral atom, the number of protons equals the number of electrons.
3. Isotopes of an element have the same number of protons but have different numbers of neutrons in their nuclei.
4. The mass number is the sum of the numbers of protons and neutrons in the nucleus of an atom.

Key Takeaways

- Each element is identified by its atomic number. The atomic number provides the element's location on the periodic table
- The isotopes of an element have different masses and are identified by their mass numbers.

Answers

Example 4.4.1

According to Table 2.4.1, aluminum has an atomic number of 13. Therefore, every aluminum atom has 13 protons in its nucleus.

Iron has an atomic number of 26. Therefore, every iron atom has 26 protons in its nucleus.

Carbon has an atomic number of 6. Therefore, every carbon atom has 6 protons in its nucleus.

Exercise 4.4.1

Sodium has 11 protons in its nucleus.

Oxygen has 8 protons in its nucleus.

Chlorine has 17 protons in its nucleus

Example 4.4.2

The atomic number of sulfur is 16. Therefore, in a neutral atom of sulfur, there are 16 electrons.

The atomic number of tungsten is 74. Therefore, in a neutral atom of tungsten, there are 74 electrons.

The atomic number of argon is 18. Therefore, in a neutral atom of argon, there are 18 electrons.

Exercise 4.4.2

Mg has 12 electrons.

K has 19 electrons.

I has 53 electrons.

Example 4.4.3

In $^{35}_{17}\mathrm{Cl}$ there are 17 protons, and 35 – 17 = 18 neutrons in each nucleus.

In $^{127}_{53}\mathrm{I}$ there are 53 protons, and 127 – 53 = 74 neutrons in each nucleus.

Exercise 4.4.3

In $^{197}_{79}\mathrm{Au}$ there are 79 protons, and 197 – 79 = 118 neutrons in each nucleus.

In $^{23}_{11}\mathrm{Na}$ there are 11 protons, and 23 – 11 = 12 neutrons in each nucleus.

This page titled 4.4: Nuclei of Atoms is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.4: Nuclei of Atoms by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.5: Atomic Masses

Learning Objectives

- Define atomic mass and atomic mass unit.
- Calculate atomic mass.

Even though atoms are very tiny pieces of matter, they have mass. Their masses are so small, however, that chemists often use a unit other than grams to express them—the atomic mass unit.

Atomic Mass Unit

The atomic mass unit (abbreviated u, although amu is also used) is defined as 1/12 of the mass of a ^{12}C atom:

$$1\ \text{u} = \frac{1}{12}\ \text{the mass of}\ {}^{12}\text{C}\ \text{atom} \tag{4.5.1}$$

It is equal to 1.661×10^{-24} g.

Masses of other atoms are expressed with respect to the atomic mass unit. For example, the mass of an atom of ^{1}H is 1.008 u, the mass of an atom of ^{16}O is 15.995 u, and the mass of an atom of ^{32}S is 31.97 u. Note, however, that these masses are for particular isotopes of each element. Because most elements exist in nature as a mixture of isotopes, any sample of an element will actually be a mixture of atoms having slightly different masses (because neutrons have a significant effect on an atom's mass). How, then, do we describe the mass of a given element? By calculating an average of an element's atomic masses, weighted by the natural abundance of each isotope, we obtain a weighted average mass called the atomic mass (also commonly referred to as the *atomic weight*) of an element.

Atomic Mass is the Weighted Average Mass of Isotopes

As stated above, most elements occur naturally as a mixture of two or more isotopes. Listed below (Table 4.5.1) are the naturally occurring isotopes of selected elements along with the percent natural abundance of each.

Table 4.5.1: Atomic Masses and Percent Abundances of Some Natural Isotopes

Element	Isotope (Symbol)	Percent Natural Abundance	Atomic Mass (amu)	Average Atomic Mass (amu)
Hydrogen	$^{1}_{1}H$	99.985	1.0078	1.0079
	$^{2}_{1}H$	0.015	2.0141	
	$^{3}_{1}H$	negligible	3.0160	
Carbon	$^{12}_{6}C$	98.89	12.000	12.011
	$^{13}_{6}C$	1.11	13.003	
	$^{14}_{6}C$	trace	14.003	
Oxygen	$^{16}_{8}O$	99.759	15.995	15.999
	$^{17}_{8}O$	0.037	16.995	
	$^{18}_{8}O$	0.204	17.999	
Chlorine	$^{35}_{17}Cl$	75.77	34.969	35.453
	$^{38}_{17}Cl$	24.23	36.966	

Element	Isotope (Symbol)	Percent Natural Abundance	Atomic Mass (amu)	Average Atomic Mass (amu)
Copper	$^{63}_{29}\mathrm{Cu}$	69.17	62.930	63.546
	$^{65}_{29}\mathrm{Cu}$	30.83	64.928	

For some elements, one particular isotope is much more abundant than any other isotopes. For example, naturally occurring hydrogen is nearly all hydrogen-1, and naturally occurring oxygen is nearly all oxygen-16. For many other elements, however, more than one isotope may exist in substantial quantities. Chlorine (atomic number 17) is yellowish-green toxic gas. About three quarters of all chlorine atoms have 18 neutrons, giving those atoms a mass number of 35. About one quarter of all chlorine atoms have 20 neutrons, giving those atoms a mass number of 37. Were you to simply calculate the arithmetic average of the precise atomic masses, you would get approximately 36.

$$\frac{34.969\,u + 36.966\,u}{2} = 35.968\,u$$

As you can see, the average atomic mass given in the last column of the table above (35.453) is significantly lower. Why? The reason is that we need to take into account the natural abundance percentages of each isotope in order to calculate what is called the **weighted average**. The atomic mass *of an element is the weighted average of the atomic masses of the naturally occurring isotopes of that element.*

$$0.7577\,(34.969\,u) + 0.2423\,(36.966\,u) = 35.453\,u$$

The weighted average is determined by multiplying the percent of natural abundance by the actual mass of the isotope. This is repeated until there is a term for each isotope. For chlorine, there are only two naturally occurring isotopes so there are only two terms.

$$\text{Atomic mass} = (\%1)(\text{mass 1}) + (\%2)(\text{mass 2}) + \cdots$$

Another example: oxygen exists as a mixture that is 99.759% ^{16}O, 0.037% ^{17}O and 0.204% ^{18}O. The atomic mass of oxygen (use percent natural abundance data from Table 2.5.1) would be calculated as follows:

$$\text{Atomic mass} = (\%1)(\text{mass 1}) + (\%2)(\text{mass 2}) + (\%3)(\text{mass 3})$$

$$0.99759\,(15.995u) + 0.00037\,(16.995u) + 0.00204\,(17.999u) = 15.999u$$

To confirm your answer, compare the calculated value to the weighted mass displayed on the periodic table.

Example 4.5.1

Calculate the atomic mass of oxygen. Oxygen exists as a mixture of 3 isotopes. Their respective masses and natural abundance are shown below.

- ^{16}O: 15.995 u (99.759%)
- ^{17}O: 16.995 u (0.037%)
- ^{18}O: 17.999 u (0.204%)

Solution

Multiply the isotope abundance by the actual mass of the isotope, and then sum up the products.

$$0.99759\,(15.995\,u) + 0.00037\,(16.995\,u) + 0.00204\,(17.999\,u) = 15.999\,u$$

Exercise 4.5.1

Calculate the atomic mass of copper. Copper exists as a mixture of 2 isotopes. Their respective masses and natural abundance are shown below.

- ^{63}Cu: 62.930 u (69.17%)
- ^{65}Cu: 64.928 u (30.83%)

Answer available at the end of this section.

The **atomic mass** of each element is found under the element symbol in the **periodic table.** Examples are shown below. The atomic mass of tin (Sn) is 118.71 u while the atomic mass of carbon (C) is 12.011 u. On the other hand, the **atomic number (Z)** of each element is found **above** the atomic symbol.




Figure 4.5.1: Atomic mass indicated on entries of the Periodic Table. (public Domain; Pubchem)

The **periodic table** is found in this link:

https://pubchem.ncbi.nlm.nih.gov/periodic-table/png/Periodic_Table_of_Elements_w_Atomic_Mass_PubChem.png

Example 4.5.2: Mass of Carbon

What is the average mass of a carbon atom in grams? The atomic mass is found in the Periodic Table. Please use two decimal places.

Solution

This is a simple one-step conversion, similar to conversions we did in Chapter 1. We use the fact that 1 u = 1.661 × 10^{-24} g:

$$12.01 \cancel{\text{u}} \times \frac{1.661 \times 10^{-24}\ \text{g}}{1\ \cancel{\text{u}}} = 1.995 \times 10^{-23}\ \text{g}$$

This is an extremely small mass, which illustrates just how small individual atoms are.

Exercise 4.5.2: Mass of Tin

What is the average mass of a tin atom in grams? The average atomic mass is found in the Periodic Table. Please use two decimal places.

Answer available at the end of this section.

Concept Review Exercises

1. Define atomic mass. Why is it considered a weighted average?

2. What is an atomic mass unit?

Answers

1. The atomic mass is an average of an element's atomic masses, weighted by the natural abundance of each isotope of that element. It is a weighted average because different isotopes have different masses.
2. An atomic mass unit is 1/12th of the mass of a ^{12}C atom.

Key Takeaway

- Atoms have a mass that is based largely on the number of protons and neutrons in their nucleus.
- The atomic mass of each element in the Periodic Table is the weighted average of the mass of all its isotopes.

Answers

Exercise $4.5.1$

63.546 u

Exercise $4.5.2$**: Mass of Tin**

$$118.71 \cancel{u} \times \frac{1.661 \times 10^{-24}\ \text{g}}{1\ \cancel{u}} = 1.972 \times 10^{-22}\ \text{g}$$

This page titled 4.5: Atomic Masses is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.5: Atomic Masses by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.6: Bohr Diagrams of Atoms and Ions

Learning Objectives

- Draw the Bohr Diagram of an atom with 18 electrons or fewer.

Electron Shells

Niels Bohr proposed an early model of the atom as a central nucleus containing protons and neutrons being orbited by electrons in shells. As previously discussed, there is a connection between the number of protons in an element, the atomic number that distinguishes one element from another, and the number of electrons it has. In all electrically-neutral atoms, the number of electrons is the same as the number of protons. Each element, when electrically neutral, has a number of electrons equal to its atomic number.

An early model of the atom was developed in 1913 by Danish scientist Niels Bohr (1885–1962). The Bohr model shows the atom as a central nucleus containing protons and neutrons with the electrons in circular orbitals at specific distances from the nucleus (Figure 4.6.1). These orbits form electron shells or energy levels, which are a way of visualizing the number of electrons in the various shells. These energy levels are designated by a number and the symbol "n." For example, the 1n shell represents the first energy level located closest to the nucleus.




Figure 4.6.1: The Bohr model postulated that electron orbited the nucleus in shells of fixed distance.

An electron normally exists in the lowest energy shell available, which is the one closest to the nucleus. Energy from a photon of light can bump it up to a higher energy shell, but this situation is unstable and the electron quickly decays back to the ground state.

Bohr Diagrams

Bohr diagrams show electrons orbiting the nucleus of an atom somewhat like planets orbit around the sun. In the Bohr model, electrons are pictured as traveling in circles at different shells, depending on which element you have. Figure 4.6.2 contrast the Bohr diagrams for lithium, fluorine and aluminum atoms. The shell closest to the nucleus is called the K shell, next is the L shell, next is the M shell.




Figure 4.6.2: Bohr diagrams for neutral lithium, fluorine and aluminum atoms.

Each shell can only hold certain number of electrons. The first shell can have 2, the second shell can have 8 , and the third shell can have 18 electrons and so on.

- Lithium has three electrons:
  - two go shell 1 and
  - the remaining one goes to shell 2.
- Fluorine has nine electrons:
  - two go to shell 1 and
  - the remaining seven go to shell 2.
- Aluminum has thirteen electrons:
  - two go to shell 1,
  - eight go to shell 2, and
  - remaining three go to shell 3.

Orbitals in the Bohr model

Electrons fill orbit shells in a consistent order. Under standard conditions, atoms fill the inner shells (closer to the nucleus) first, often resulting in a variable number of electrons in the outermost shell. The innermost shell has a maximum of two electrons, but the next two electron shells can each have a maximum of eight electrons. This is known as the octet rule which states that, with the exception of the innermost shell, atoms are more stable energetically when they have eight electrons in their valence shell, the outermost electron shell. Examples of some neutral atoms and their electron configurations are shown in Figure 4.6.3 . As shown, helium has a complete outer electron shell, with two electrons filling its first and only shell. Similarly, neon has a complete outer 2n shell containing eight electrons. In contrast, chlorine and sodium have seven and one electrons in their outer shells, respectively. Theoretically, they would be more energetically stable if they followed the octet rule and had eight.




Figure 4.6.3:

Bohr diagrams

Bohr diagrams indicate how many electrons fill each principal shell. Group 18 elements (helium, neon, and argon are shown) have a full outer, or valence, shell. A full valence shell is the most stable electron configuration. Elements in other groups have partially-filled valence shells and gain or lose electrons to achieve a stable electron configuration.

An atom may gain or lose electrons to achieve a full valence shell, the most stable electron configuration. The periodic table is arranged in columns and rows based on the number of electrons and where these electrons are located, providing a tool to understand how electrons are distributed in the outer shell of an atom. As shown in , the group 18 atoms helium (He), neon (Ne), and argon (Ar) all have filled outer electron shells, making it unnecessary for them to gain or lose electrons to attain stability; they are highly stable as single atoms. Their non-reactivity has resulted in their being named the inert gases (or noble gases). In comparison, the group 1 elements, including hydrogen (H), lithium (Li), and sodium (Na), all have one electron in their outermost shells. This means that they can achieve a stable configuration and a filled outer shell by donating or losing an electron. As a result of losing a negatively-charged electron, they become positively-charged ions. When an atom loses an electron to become a positively-charged ion, this is indicated by a plus sign after the element symbol; for example, Na^{+}. Group 17 elements, including fluorine and chlorine, have seven electrons in their outermost shells; they tend to fill this shell by gaining an electron from other atoms, making them negatively-charged ions. When an atom gains an electron to become a negatively-charged ion this is indicated by a minus sign after the element symbol; for example, F^{-}. Thus, the columns of the periodic table represent the potential shared state of these elements' outer electron shells that is responsible for their similar chemical characteristics.

Contributors and Attributions

Boundless (www.boundless.com)

- Manish Bhardwaj

This page titled 4.6: Bohr Diagrams of Atoms and Ions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

- 2.5: Atomic Masses by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.7: Electron Arrangements

Learning Outcomes

- Determine the energy levels of electrons for the first 20 elements.
- Explain the relevance of valence electrons in chemical processes.
- Identify the number of valence electrons in an element.
- Describe the stability of an atom as a result of following the octet rule.

The structure of the atom was discussed in the previous unit and now we will focus on the role that electrons play in the formation of compounds. Regardless of the type of compound or the number of atoms or electrons involved, it is the electrons of those atoms that interact to form a compound.

Electron Arrangement

Electrons are not randomly arranged in an atom and their position within the atom can be described using **electron arrangements**, which are a simplified version of electron configurations. For each element of interest, we look at the number of electrons in a single atom and then determine how those electrons are arranged based on the atomic model. The main idea behind electron arrangements is that electrons can only exist at certain energy levels. By understanding the energy levels of electrons in an atom, we can predict properties and understand behavior of the atom.

As shown in the figure below, there are multiple energy levels where electrons can be found. As the energy level increases, the energy difference between them decreases. A maximum of two electrons can be found in the $n = 1$ level; eight electrons can be in the $n = 2$ level. Although the $n = 3$ and $n = 4$ levels show only eight electrons in this diagram, those energy levels can hold more but not until we start looking at the transition metals. We will only be concerned with the electron arrangements of elements through calcium $(Z = 20)$ so we will put a maximum of eight electrons in the $n = 3$ level and two in the $n = 4$ level.




Figure 4.7.1: Energy levels of electrons.

Example 4.7.1

What is the electron arrangement of oxygen?

Solution

Oxygen has eight electrons. The first two electrons will go in the $n = 1$ level. Two is the maximum number of electrons for the level so the other electrons will have to go in a higher energy level. The $n = 2$ level can hold up to eight electrons so the remaining six electrons will go in the $n = 2$ level. The electron arrangement of oxygen is (2, 6).

Example 4.7.2

What is the electron arrangement of chlorine?

Solution

Chlorine has 17 electrons. The first two electrons will go in the $n = 1$ level. Two is the maximum number of electrons for the level so the other electrons will have to go in higher energy levels. The $n = 2$ level can hold up to eight electrons so the next 8 electrons will go in the $n = 2$ level. The remaining 7 electrons can go in the $n = 3$ level since it holds a maximum of 8 electrons. The electron arrangement of chlorine is (2, 8, 7).

The electron arrangement also provides information about the number of **valence electrons**. The valence electrons are the electrons in the highest energy level and the ones involved in ion and bond formation. Knowing the number of valence electrons will allow us to predict how a particular element will interact with other elements. Electrons in lower energy levels are called the **core electrons**.

Let's look at the figure below which shows the electron diagram for magnesium and its 12 electrons. The first two electrons are found in the $n = 1$ energy level, the next eight electrons are found in the $n = 2$ level, and the remaining two electrons are found in the $n = 3$ level. The electrons always fill the lowest energy levels available until that level is filled, then electrons fill the next energy level until it is filled. This continues for all of the electrons in an atom. We can show the electron arrangement as (2, 8, 2) representing the electrons in the $n = 1$, $n = 2$, and $n = 3$ levels, respectively.




Figure 4.7.2: Electron diagram for magnesium.

The electron arrangement also shows the number of valence electrons which is two for magnesium because there are two electrons in the $n = 3$ energy level which is the highest occupied energy level for magnesium. This corresponds to the $2+$ charge formed when magnesium forms an ion. It is willing to lose 2 electrons so that it has the same electron arrangements as the nearest noble gas, which is neon (2, 8). Atoms will gain or lose electrons to look like the nearest noble gas because the noble gases are unreactive due to the stability of having eight electrons in the highest energy level. This desire of atoms to have eight electrons in their outermost shell is known as the **octet rule**.

Example 4.7.3

What is the electron arrangement of aluminum? How many valence electrons does it have?

Solution

Aluminum has 13 electrons so it will have the electron arrangement (2, 8, 3) which represents two electrons in the $n = 1$ energy level, eight electrons in the $n = 2$ level, and three electrons in the $n = 3$ level. Aluminum has three valence electrons (indicated by the three electrons in the $n = 3$ level).

Example 4.7.4

How many valence electrons does chlorine have? How many electrons will chlorine gain or lose to form an ion?

Solution

Chlorine has 7 electrons in its valence shell. To meet the octet rule, it must either gain one electron or lose seven electrons. Gaining one is easier than losing seven so it will gain one electron to have a total of eight electrons when it forms an ion (i.e. charged particle).

Contributors and Attributions

-

- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)

This page titled 4.7: Electron Arrangements is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Allison Soult.

4.8: The Periodic Table

Learning Objectives

- Explain how elements are organized into the periodic table.
- Describe how some characteristics of elements relate to their positions on the periodic table.

In the 19th century, many previously unknown elements were discovered, and scientists noted that certain sets of elements had similar chemical properties. For example, chlorine, bromine, and iodine react with other elements (such as sodium) to make similar compounds. Likewise, lithium, sodium, and potassium react with other elements (such as oxygen) to make similar compounds. Why is this so?

In 1864, Julius Lothar Meyer, a German chemist, organized the elements by atomic mass and grouped them according to their chemical properties. Later that decade, Dmitri Mendeleev, a Russian chemist, organized all the known elements according to similar properties. He left gaps in his table for what he thought were undiscovered elements, and he made some bold predictions regarding the properties of those undiscovered elements. When elements were later discovered whose properties closely matched Mendeleev's predictions, his version of the table gained favor in the scientific community. Because certain properties of the elements repeat on a regular basis throughout the table (that is, they are periodic), it became known as the periodic table.

> Mendeleev had to list some elements out of the order of their atomic masses to group them with other elements that had similar properties.

The periodic table is one of the cornerstones of chemistry because it organizes all the known elements on the basis of their chemical properties. A modern version is shown in Figure 4.8.1. Most periodic tables provide additional data (such as atomic mass) in a box that contains each element's symbol. The elements are listed in order of atomic number.




Figure 4.8.1: Modern Periodic Table. (Public Domain; PubChem modified Leticia Colmenares).

Features of the Periodic Table

Elements that have similar chemical properties are grouped in columns called **groups** (or families). The groups are numbered from 1 to 18. As well as being numbered, some of these groups have names—for example, *alkali metals* (the first column of elements), *alkaline earth metals* (the second column of elements), *halogens* (the next-to-last column of elements), and *noble gases* (the last column of elements).

Each row of elements on the periodic table is called a **period**. The periods are numbered from 1 to 7 from top to bottom. Periods have different lengths; the first period has only 2 elements (hydrogen and helium), while the second and third periods have 8 elements each. The fourth and fifth periods have 18 elements each, and later periods are so long that a segment from each is removed and placed beneath the main body of the table.

Metals, Nonmetals and Metalloids

Certain elemental properties become apparent in a survey of the periodic table as a whole. Every element can be classified as either a metal, a nonmetal, or a semimetal, as shown in Figure 4.8.2. A metal is a substance that is shiny, typically (but not always) silvery in color, and an excellent conductor of electricity and heat. Metals are also malleable (they can be beaten into thin sheets) and ductile (they can be drawn into thin wires). A nonmetal is typically dull and a poor conductor of electricity and heat. Solid nonmetals are also very brittle. As shown in Figure 4.8.2, **metals** occupy the left three-fourths of the periodic table, while **nonmetals** (except for hydrogen) are clustered in the upper right-hand corner of the periodic table. The elements with properties intermediate between those of metals and nonmetals are called **semimetals** (or **metalloids**). Elements adjacent to the **bold zigzag** line in the right-hand portion of the periodic table have semimetal properties.




Figure 4.8.2: Types of Elements. Elements are either metals, nonmetals, or semimetals. Each group is located in a different part of the periodic table.

Example 4.8.1

Based on its position in the periodic table, do you expect selenium (Se) to be a metal, a nonmetal, or a semimetal?

Solution

The atomic number of selenium is 34, which places it in period 4 and group 16. In Figure 2.7.1, selenium lies above and to the right of the diagonal line marking the boundary between metals and nonmetals, so it should be a nonmetal. Note, however, that because selenium is close to the metal-nonmetal dividing line, it would not be surprising if selenium were similar to a semimetal in some of its properties.

4.8.1 Exerc

Based on its location in the periodic table, do you expect indium (In) to be a nonmetal, a metal, or a semimetal?

Answer available at the end of this section.

Main Group, Transition and Inner-transition Elements

Another way to categorize the elements of the periodic table is shown in Figure 4.8.3. The first two columns on the left and the last six columns on the right are called the **main group** or **representative** elements. The ten-column block between these columns contains the **transition metals**. The two rows beneath the main body of the periodic table contain the inner transition metals. The elements in these two rows are also referred to as, respectively, the **lanthanide metals** and the **actinide metals**.




Figure 4.8.3: Special Names for Sections of the Periodic Table. Some sections of the periodic table have special names. The elements lithium, sodium, potassium, rubidium, cesium, and francium are collectively known as alkali metals.

Alkali metals are the first column. Alkaline earth metals are the second. Halogens are the second to last column. Noble gases are the last column.

To Your Health: Transition Metals in the Body

Most of the elemental composition of the human body consists of main group elements. The first element appearing on the list that is not a main group element is iron, at 0.006 percentage by mass. Because iron has relatively massive atoms, it would appear even lower on a list organized in terms of percent by *atoms* rather than percent by mass.

Iron is a transition metal. Transition metals have interesting chemical properties, partially because some of their electrons are in *d* subshells. The chemistry of iron makes it a key component in the proper functioning of red blood cells.

Red blood cells are cells that transport oxygen from the lungs to cells of the body and then transport carbon dioxide from the cells to the lungs. Without red blood cells, animal respiration as we know it would not exist. The critical part of the red blood cell is a protein called *hemoglobin*. Hemoglobin combines with oxygen and carbon dioxide, transporting these gases from one location to another in the body. Hemoglobin is a relatively large molecule, with a mass of about 65,000 u.

The crucial atom in the hemoglobin protein is iron. Each hemoglobin molecule has four iron atoms, which act as binding sites for oxygen. It is the presence of this particular transition metal in your red blood cells that allows you to use the oxygen you inhale.

Other transition metals have important functions in the body, despite being present in low amounts. Zinc is needed for the body's immune system to function properly, as well as for protein synthesis and tissue and cell growth. Copper is also needed for several proteins to function properly in the body. Manganese is needed for the body to metabolize oxygen properly. Cobalt is a necessary component of vitamin B-12, a vital nutrient. These last three metals are not listed explicitly in Table 2.1.2, so they are present in the body in very small quantities. However, even these small quantities are required for the body to function properly.

As previously noted, the periodic table is arranged so that elements with similar chemical behaviors are in the same group.

Chemists often make general statements about the properties of the elements in a group using descriptive names with historical origins. For example, the elements of Group 1 are known as the alkali metals, Group 2 are the alkaline earth metals, Group 17 are the halogens, and Group 18 are the noble gases.

Group 1: The Alkali Metals

The alkali metals are lithium, sodium, potassium, rubidium, cesium, and francium. Hydrogen is unique in that it is generally placed in Group 1, but it is not a metal. The compounds of the alkali metals are common in nature and daily life. One example is table salt (sodium chloride); lithium compounds are used in greases, in batteries, and as drugs to treat patients who exhibit manic-depressive, or bipolar, behavior. Although lithium, rubidium, and cesium are relatively rare in nature, and francium is so unstable and highly radioactive that it exists in only trace amounts, sodium and potassium are the seventh and eighth most abundant elements in Earth's crust, respectively.

Group 2: The Alkaline Earth Metals

The alkaline earth metals are beryllium, magnesium, calcium, strontium, barium, and radium. Beryllium, strontium, and barium are rare, and radium is unstable and highly radioactive. In contrast, calcium and magnesium are the fifth and sixth most abundant elements on Earth, respectively; they are found in huge deposits of limestone and other minerals.

Group 17: The Halogens

The halogens are fluorine, chlorine, bromine, iodine, and astatine. The name halogen is derived from the Greek words for "salt forming," which reflects that all the halogens react readily with metals to form compounds, such as sodium chloride and calcium chloride (used in some areas as road salt).

Compounds that contain the fluoride ion are added to toothpaste and the water supply to prevent dental cavities. Fluorine is also found in Teflon coatings on kitchen utensils. Although chlorofluorocarbon propellants and refrigerants are believed to lead to the depletion of Earth's ozone layer and contain both fluorine and chlorine, the latter is responsible for the adverse effect on the ozone layer. Bromine and iodine are less abundant than chlorine, and astatine is so radioactive that it exists in only negligible amounts in nature.

Group 18: The Noble Gases

The noble gases are helium, neon, argon, krypton, xenon, and radon. Because the noble gases are composed of only single atoms, they are called monatomic. At room temperature and pressure, they are **unreactive gases**. Because of their lack of reactivity, for many years they were called **inert gases** or rare gases. However, the first chemical compounds containing the noble gases were prepared in 1962. Although the noble gases are relatively minor constituents of the atmosphere, natural gas contains substantial amounts of helium. Because of its low reactivity, argon is often used as an unreactive (inert) atmosphere for welding and in light bulbs. The red light emitted by neon in a gas discharge tube is used in neon lights.

To Your Health: Radon

Radon is an invisible, odorless noble gas that is slowly released from the ground, particularly from rocks and soils whose uranium content is high. Because it is a noble gas, radon is not chemically reactive. Unfortunately, it is radioactive, and increased exposure to it has been correlated with an increased lung cancer risk.

Because radon comes from the ground, we cannot avoid it entirely. Moreover, because it is denser than air, radon tends to accumulate in basements, which if improperly ventilated can be hazardous to a building's inhabitants. Fortunately, specialized ventilation minimizes the amount of radon that might collect. Special fan-and-vent systems are available that draw air from below the basement floor, before it can enter the living space, and vent it above the roof of a house.

After smoking, radon is thought to be the second-biggest *preventable* cause of lung cancer in the United States. The American Cancer Society estimates that 10% of all lung cancers are related to radon exposure. There is uncertainty regarding what levels of exposure cause cancer, as well as what the exact causal agent might be (either radon or one of its breakdown products, many of which are also radioactive and, unlike radon, not gases). The US Environmental Protection Agency recommends testing every floor below the third floor for radon levels to guard against long-term health effects.

Valence Electrons and Group Number

The number of valence electrons of an element can be determined by the periodic table group (vertical column) in which the element is categorized. With the exception of groups 3–12 (the transition metals), the units digit of the group number identifies how many valence electrons are associated with a neutral atom of an element listed under that particular column.

Table 4.8.1. *The Group number and the number of valence electrons.*

Periodic table group	Valence electrons
Group 1 (I) (alkali metals)	1
Group 2 (II) (alkaline earth metals)	2
Groups 3-12 (transition metals)	2*
Group 13 (III) (boron group)	3
Group 14 (IV) (carbon group)	4
Group 15 (V) (pnictogens)	5
Group 16 (VI) (chalcogens)	6
Group 17 (VII) (halogens)	7
Group 18 (VIII or 0) (noble gases)	8**

* *The general method for counting valence electrons is generally not useful for transition metals.*

** *Except for helium, which has only two valence electrons.*

Atomic Radius

The periodic table is useful for understanding atomic properties that show periodic trends. One such property is the atomic radius (Figure 4.8.5). The atomic radius is defined as one-half the distance between the nuclei of identical atoms that are bonded together. The units for atomic radii are picometers, equal to 10^{-12} meters. As an example, the internuclear distance between the two hydrogen atoms in an H_2 molecule is measured to be $74\ \text{pm}$. Therefore, the atomic radius of a hydrogen atom is $\frac{74}{2} = 37\ \text{pm}$.

As mentioned earlier, the higher the shell number, the farther from the nucleus the electrons in that shell are likely to be. In other words, the size of an atom is generally determined by the number of the valence electron shell. Therefore, as we go down a column on the periodic table, the atomic radius increases. As we go *across* a period on the periodic table, however, electrons are being added to the *same* valence shell; meanwhile, more protons are being added to the nucleus, so the positive charge of the nucleus is increasing. The increasing positive charge attracts the electrons more strongly, pulling them closer to the nucleus. Consequently, as we go across a period, from left to right, the atomic radius decreases. These trends are seen clearly in Figure 4.8.5




Figure 4.8.5 Trends on the Periodic Table. Atomic radii of the representative elements measured in picometers. The relative sizes of the atoms show several trends with regard to the structure of the periodic table. Atoms become larger going down a group and going from right to left across a period.

Example 4.8.2

Using the periodic table (rather than Figure 4.8.5), which atom is larger?

a. N or Bi
b. Mg or Cl

Answer a available at the end of this section.

Exercise 4.8.2

Using the periodic table (rather than Figure 4.8.5), which atom is larger?

a. Li or F
b. Na or K

Answer a available at the end of this section.

Career Focus: Clinical Chemist

Clinical chemistry is the area of chemistry concerned with the analysis of body fluids to determine the health status of the human body. Clinical chemists measure a variety of substances, ranging from simple elements such as sodium and potassium to complex molecules such as proteins and enzymes, in blood, urine, and other body fluids. The absence or presence, or abnormally low or high amounts, of a substance can be a sign of some disease or an indication of health. Many clinical chemists use sophisticated equipment and complex chemical reactions in their work, so they not only need to understand basic chemistry, but also be familiar with special instrumentation and how to interpret test results.

Concept Review Exercises

1. How are the elements organized into the periodic table?
2. Looking at the periodic table, where do the following elements appear?

a. the metals
b. the nonmetals
c. the halogens
d. the transition metals
e. the noble gases

3. Describe the trends in atomic radii as related to an element's position on the periodic table.

Answers

1. Elements are organized in order of increasing atomic number.
2. a. the left three-quarters of the periodic table (to the left of the zigzag band)
   b. the right quarter of the periodic table (to the right of the zigzag band)
   c. the next-to-last column of the periodic table
   d. the middle section of the periodic table
   e. the last column of the periodic table
3. As you go across the periodic table, atomic radii decrease; as you go down the periodic table, atomic radii increase.

Key Takeaways

- The chemical elements are arranged in a chart called the periodic table.
- Some characteristics of the elements are related to their position on the periodic table.
- The number of valence electrons of an element can be determined by the group (vertical column) number in the Periodic Table. Elements with the same number of valence electrons have similar chemical properties.

2.7: The Periodic Table

Answers

metal

Example 4.8.2

Bi is below N in Group 5A in the periodic table and has electrons in higher-numbered shells, hence we expect that Bi atoms are larger than N atoms.

Both Mg and Cl are in period 3 of the periodic table, but Cl lies farther to the right. Therefore we expect Mg atoms to be larger than Cl atoms.

Exercise 4.8.2

Li and F are on the same period, but F lies farther to the right. Therefore, we expect Li to be larger than F atoms.

K lies below Na in Group 1A, hence has more electron shells, making it larger than Na.

This page titled 4.8: The Periodic Table is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.7: The Periodic Table by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

4.S: Elements, Atoms, and the Periodic Table (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms and ask yourself how they relate to the topics in the chapter.

An element is a substance that cannot be broken down into simpler chemical substances. Only about 90 naturally occurring elements are known. They have varying abundances on Earth and in the body. Each element has a one- or two-letter **chemical symbol**.

The **modern atomic theory** states that the smallest piece of an element is an **atom**. Individual atoms are extremely small, on the order of 10^{-10} m across. Most elements exist in pure form as individual atoms, but some exist as **diatomic molecules**. Atoms themselves are composed of subatomic particles. The **electron** is a tiny subatomic particle with a negative charge. The **proton** has a positive charge and, while small, is much larger than the electron. The **neutron** is also much larger than an electron but has no electrical charge.

Protons, neutrons, and electrons have a specific arrangement in an atom. The protons and neutrons are found in the center of the atom, grouped together into a **nucleus**. The electrons are found in fuzzy clouds around the nucleus.

Each element has a characteristic number of protons in its nucleus. This number of protons is the **atomic number** of the element. An element may have different numbers of neutrons in the nuclei of its atoms; such atoms are referred to as **isotopes**. Two isotopes of hydrogen are deuterium, with a proton and a neutron in its nucleus, and tritium, with a proton and two neutrons in its nucleus. The sum of the numbers of protons and neutrons in a nucleus is called the **mass number** and is used to distinguish isotopes from each other.

Masses of individual atoms are measured in **atomic mass units**. An atomic mass unit is equal to 1/12th of the mass of a single carbon-12 atom. Because different isotopes of an element have different masses, the **atomic mass** of an element is a weighted average of the masses of all the element's naturally occurring isotopes.

The modern theory of electron behavior is called **quantum mechanics**. According to this theory, electrons in atoms can only have specific, or **quantized**, energies. Electrons are grouped into general regions called **shells**, and within these into more specific regions called **subshells**. There are four types of subshells, and each type can hold up to a maximum number of electrons. The distribution of electrons into shells and subshells is the **electron configuration** of an atom. Chemistry typically occurs because of interactions between the electrons of the outermost shell of different atoms, called the valence shell electrons. Electrons in inner shells are called core electrons.

Elements are grouped together by similar chemical properties into a chart called the **periodic table**. Vertical columns of elements are called **groups** or **families**. Some of the groups of elements have names, like the alkali metals, the alkaline earth metals, the halogens, and the noble gases. A horizontal row of elements is called a **period**. Periods and groups have differing numbers of elements in them. The periodic table separates elements into **metals**, **nonmetals**, and **semimetals**. The periodic table is also separated into **main group elements**, **transition metals**, **lanthanide elements**, and **actinide elements**. The lanthanide and actinide elements are also referred to as **inner transition metal elements**. The shape of the periodic table reflects the sequential filling of shells and subshells in atoms.

The periodic table helps us understand trends in some of the properties of atoms. One such property is the **atomic radius** of atoms. From top to bottom of the periodic table, atoms get bigger because electrons are occupying larger and bigger shells. From left to right across the periodic table, electrons are filling the same shell but are being attracted by an increasing positive charge from the nucleus, and thus the atoms get smaller.

This page titled 4.S: Elements, Atoms, and the Periodic Table (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 2.S: Elements, Atoms, and the Periodic Table (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5: IONIC BONDING AND SIMPLE IONIC COMPOUNDS

There are only 118 known chemical elements but tens of millions of known chemical compounds. Compounds can be very complex combinations of atoms, but many important compounds are fairly simple. Table salt, as we have seen, consists of only two elements: sodium and chlorine. Nevertheless, the compound has properties completely different from either elemental sodium (a chemically reactive metal) or elemental chlorine (a poisonous, green gas). We will see additional examples of such differences in this chapter as we consider how atoms combine to form compounds.

5.1: Prelude to Ionic Bonding and Simple Ionic Compounds

5.2: Two Types of Bonding

5.3: Ions

5.4: Formulas for Ionic Compounds

5.5: Ionic Nomenclature

5.6: Formula Mass

5.7: Characteristics of Ionic Compounds

5.S: Ionic Bonding and Simple Ionic Compounds (Summary)

Template:HideTOC

This page titled 5: Ionic Bonding and Simple Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3: Ionic Bonding and Simple Ionic Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.1: Prelude to Ionic Bonding and Simple Ionic Compounds

We will see that the word salt has a specific meaning in chemistry, but to most people, this word refers to table salt. This kind of salt is used as a condiment throughout the world, but it was not always so abundant. Two thousand years ago, Roman soldiers received part of their pay as salt, which explains why the words salt and salary come from the same Latin root (salarium). Today, table salt is either mined or obtained from the evaporation of saltwater.

Table salt is sodium chloride (NaCl), which is a simple compound of two elements that are necessary for the human body to function properly. Sodium, for example, is important for nerve conduction and fluid balance. In fact, human blood is about a 0.9% sodium chloride solution, and a solution called normal saline is commonly administered intravenously in hospitals.

Although some salt in our diets is necessary to replenish the sodium and chloride ions that we excrete in urine and sweat, too much is unhealthy, and many people may be ingesting more salt than their bodies need. The RDI of sodium is 2,400 mg—the amount in about 1 teaspoon of salt—but the average intake of sodium in the United States is between 4,000 mg and 5,000 mg, partly because salt is a common additive in many prepared foods. Previously, the high ingestion of salt was thought to be associated with high blood pressure, but current research does not support this link. Even so, some doctors still recommend a low-salt diet (never a “no-salt” diet) for patients with high blood pressure, which may include using a salt substitute. Most salt substitutes use potassium instead of sodium, but some people complain that the potassium imparts a slightly bitter taste.

This page titled 5.1: Prelude to Ionic Bonding and Simple Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.0: Prelude to Ionic Bonding and Simple Ionic Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.2: Two Types of Bonding

Learning Objectives

- Define the octet rule.
- Describe how ionic bonds are formed.

Atoms can join together by forming a chemical bond, which is a very strong attraction between two atoms. Chemical bonds are formed when electrons in different atoms interact with each other to make an arrangement that is more stable than when the atoms are apart.

What causes atoms to make a chemical bond with other atoms, rather than remaining as individual atoms? A clue comes by considering the noble gas elements, the rightmost column of the periodic table. These elements—helium, neon, argon, krypton, xenon, and radon—do not form compounds very easily, which suggests that they are especially stable as lone atoms. What else do the noble gas elements have in common? Except for helium, they all have eight valence electrons. Chemists have concluded that atoms are especially stable if they have eight electrons in their outermost shell. This useful rule of thumb is called the **octet rule**, and it is a key to understanding why compounds form.

> Of the noble gases, only krypton, xenon, and radon have been found to make compounds.

There are two ways for an atom that does not have an octet of valence electrons to obtain an octet in its outer shell. One way is the transfer of electrons between two atoms until all atoms have octets. Because some atoms will lose electrons and some atoms will gain electrons, there is no overall change in the number of electrons, but individual atoms acquire a nonzero electric charge. Those that lose electrons become positively charged, and those that gain electrons become negatively charged. Charged atoms are called ions. Because opposite charges attract (while like charges repel), these oppositely charged ions attract each other, forming **ionic bonds**. The resulting compounds are called **ionic compounds** and are the primary subject of this chapter.

The second way for an atom to obtain an octet of electrons is by sharing electrons with another atom. These shared electrons simultaneously occupy the outermost shell of more than one atom. The bond made by electron sharing is called a **covalent bond**.

> Despite our focus on the octet rule, we must remember that for small atoms, such as hydrogen, helium, and lithium, the first shell is, or becomes, the outermost shell and hold only two electrons. Therefore, these atoms satisfy a "**duet rule**" rather than the octet rule.

Example 5.2.1

A sodium atom has one valence electron. Do you think it is more likely for a sodium atom to lose one electron or gain seven electrons to obtain an octet?

Solution

Although either event is possible, a sodium atom is more likely to lose its single valence electron. When that happens, it becomes an ion with a net positive charge. This can be illustrated as follows:

Table explaining the solution to the example.

Sodium atom		Sodium ion	
11 protons	11+	11 protons	11+
11 electrons	11–	10 electrons	10–
	0 overall charge		+1 overall charge

Exercise 5.2.1

A fluorine atom has seven valence electrons. Do you think it is more likely for a fluorine atom to lose seven electrons or gain one electron to obtain an octet? Write the formula of the resulting ion.

Answer available at the end of this section.

Key Takeaways

- Atoms have a tendency to have eight electrons in their valence shell.
- The attraction of oppositely charged ions is what makes ionic bonds.

Exercises

1. What is the octet rule?

2. How are ionic bonds formed?

3. Why is an ionic compound unlikely to consist of two positively charged ions?

4. Why is an ionic compound unlikely to consist of two negatively charged ions?

5. A calcium atom has two valence electrons. Do you think it will lose two electrons or gain six electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

6. An aluminum atom has three valence electrons. Do you think it will lose three electrons or gain five electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

7. A selenium atom has six valence electrons. Do you think it will lose six electrons or gain two electrons to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

8. An iodine atom has seven valence electrons. Do you think it will lose seven electrons or gain one electron to obtain an octet in its outermost electron shell? Write the formula of the resulting ion.

Answers

1. The octet rule is the concept that atoms tend to have eight electrons in their valence electron shell.

2. Ionic bonds are formed by the attraction between oppositely charged ions.

3. Positive charges repel each other, so an ionic compound is not likely between two positively charged ions.

4. Negative charges repel each other also. 5. Ca atom is more likely to lose two electrons. It will become Ca^{2+} ion. 6. An Al atom is more likely to lose three electrons. It will become Al^{3+} ion.

7. Selenium is more likely to gain two electrons. It will become Se^{2-} ion.

8. Iodine is more likely to gain one electron. It will become I^{-} ion.

Answers

Exercise 5.2.1

The process that involves less number of electrons is more favorable. Fluorine would gain one electron. The formula of the resulting ion is F^{-}.

This page titled 5.2: Two Types of Bonding is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.1: Two Types of Bonding by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.3: Ions

Learning Objectives

- Define the two types of ions.
- Use Lewis diagrams to illustrate ion formation.

Most atoms do not have eight electrons in their valence electron shell. Some atoms have only a few electrons in their outer shell, while some atoms lack only one or two electrons to have an octet. In cases where an atom has three or fewer valence electrons, the atom may lose those valence electrons quite easily until what remains is a lower shell that contains an octet. Atoms that lose electrons acquire a positive charge as a result because they are left with fewer negatively charged electrons to balance the positive charges of the protons in the nucleus. Positively charged ions are called cations. Most metals become cations when they make ionic compounds.

Some atoms have nearly eight electrons in their valence shell and can gain additional valence electrons until they have an octet. When these atoms gain electrons, they acquire a negative charge because they now possess more electrons than protons. Negatively charged ions are called anions. Most nonmetals become anions when they make ionic compounds.

> The names for positive and negative ions are pronounced CAT-eye-ons (cations) and ANN-eye-ons (anions), respectively.

Electron Transfer

We can look at the electron arrangement to illustrate the electron transfer process between sodium atoms and chlorine atoms.

As demonstrated here, a sodium atom (Na) has one valence electron in the third shell. It is likely to achieve an octet in its outermost shell by losing its one valence electron. The cation produced in this way, Na^+, is called the sodium ion to distinguish it from the element. The sodium ion, Na^+, has the electron arrangement with an octet of electrons from the second principal energy level. It is now the same as that of the noble gas neon. The term **isoelectronic** refers to an atom and an ion of a different atom (or two different ions) that have the same number of electrons in each shell. The sodium ion is isoelectronic with the neon atom. The equation below illustrates this process.

$$\mathrm{Na} \rightarrow \mathrm{Na^+} + \mathrm{e^-} \tag{5.3.1}$$

Figure $5.3.1$ is a graphical depiction of this process.




Figure 5.3.1: The Formation of a Sodium Ion. On the left, a sodium atom has 11 electrons. On the right, the sodium ion only has 10 electrons and a 1+ charge.

Now, let's consider chlorine atom. Chlorine has 7 electrons in its outer shell.

Only one more electron is needed to achieve an octet in chlorine's valence shell. When a chlorine atom gains an electron, its outermost principal energy level achieves an octet. In this case, the ion has the same outermost shell as the original atom, but now that shell has eight electrons in it. Once again, the octet rule has been satisfied. The resulting anion, Cl^-, is called the chloride ion; note the slight change in the suffix (*-ide* instead of *-ine*) to create the name of this anion. This process is illustrated below. (In table salt, this electron comes from the sodium atom.)

$$Cl + e^- \rightarrow Cl^- \tag{5.3.2}$$

Figure 5.3.2 is a graphical depiction of this process.




Figure 5.3.2: The Formation of a Chlorine Ion. On the left, the chlorine atom has 17 electrons. On the right, the chloride ion has 18 electrons and has a 1– charge.

With two oppositely charged ions, there is an electrostatic attraction between them because opposite charges attract. The resulting combination is the compound sodium chloride. Notice that there are no leftover electrons. The number of electrons lost by the sodium atom (one) equals the number of electrons gained by the chlorine atom (one), so the compound is electrically neutral. In macroscopic samples of sodium chloride, there are billions and billions of sodium and chloride ions, although there is

always the same number of cations and anions.

Example 5.3.1

How many electrons must Al lose/gain to achieve octet? Write the formula of the resulting ion.

Solution

A neutral Al atom has 13 electrons. The second shell has 8 electrons while the valence shell has 3 electrons. Al can achieve octet by losing the 3 valence electrons. The resulting cation is Al^{3+}

Exercise 5.3.1

How many electrons must O lose/gain to achieve octet? Write the formula of the resulting ion.

Answer available at the end of this section.

In many cases, elements that belong to the same group (vertical column) on the periodic table form ions with the same charge because they have the same number of valence electrons. Thus, the periodic table becomes a tool for remembering the charges on many ions. For example, all ions made from alkali metals, the first column on the periodic table, have a 1+ charge. Ions made from alkaline earth metals, the second group on the periodic table, have a 2+ charge. On the other side of the periodic table, the next-to-last column, the halogens, form ions having a 1– charge. Figure 5.3.3 shows how the charge on many ions can be predicted by the location of an element on the periodic table. Note the convention of first writing the number and then the sign on a multiply charged ion. The barium cation is written Ba^{2+}, not Ba^{+2}.




Figure 5.3.3: Predicting Ionic Charges. The charge that an atom acquires when it becomes an ion is related to the structure of the periodic table. Within a group (family) of elements, atoms form ions of a certain charge.

Example 5.3.2

Which of these ions is *not likely* to form?

a. Mg^{+}
b. K^{+}

Solution

(a) Mg is in Group 2A and has two valence electrons. It achieves octet by losing two electrons to form Mg^{2+} cation. Losing only one electron to form Mg^{+} does not make an octet, hence, Mg^{+} is not likely to form.

Exercise 5.3.2

Which of these ions is *not likely* to form?

a. S^{3-}
b. N^{3-}

Answer available at the end of this section.

Lewis Diagrams

Chemists use simple diagrams to show an atom's valence electrons and how they transfer. These diagrams have two advantages over the electron shell diagrams. First, they show only valence electrons. Second, instead of having a circle around the chemical symbol to represent the electron shell, they have up to eight dots around the symbol; each dot represents a valence electron. These dots are arranged to the right and left and above and below the symbol, **with no more than two dots on a side**. For example, the representation for sodium is as follows:



and the representation for chlorine is as follows:



For the above diagrams, it does not matter what sides the dots are placed on in Lewis diagrams as long as each side has a maximum of two dots.

These diagrams are called **Lewis electron dot diagrams**, or simply Lewis diagrams, after Gilbert N. Lewis, the American chemist who introduced them. To write an element's Lewis dot symbol, place the dots representing its valence electrons, one at a time, around the element's chemical symbol. Up to four dots are placed above, below, to the left, and to the right of the symbol (in any order, as long as elements with four or fewer valence electrons have no more than one dot in each position). The next dots, for elements with more than four valence electrons, are again distributed one at a time, each paired with one of the first four. In other words, **place the dots singly on each side before pairing them**. The Lewis electron dot diagram of fluorine, for example, with seven valence electrons, is constructed as follows:





Figure 5.3.4 shows the electron configurations and Lewis diagrams of the elements lithium through neon, which is the entire second period of the periodic table. For the **main group** elements, the **number of valence electrons** is the **same as the group number** listed at the top of the periodic table.



Figure 5.3.4: Going across the periodic table from Lithium through Neon, the Lewis Dot structures gain electrons.

The transfer of electrons can be illustrated easily with Lewis diagrams:




In representing the final formula, the dots are omitted.

Example 5.3.3

Starting with lithium and bromine atoms, use Lewis diagrams to show the formation of the ionic compound LiBr.

Solution

From the periodic table, we see that lithium is in the same column as sodium, so it will have the same valence shell electron configuration. That means that the neutral lithium atom will have the same Lewis diagram that the sodium atom has. Similarly, bromine is in the same column as chlorine, so it will have the same Lewis diagram that chlorine has. Therefore,




Exercise 5.3.3

Starting with magnesium and oxygen atoms, use Lewis diagrams to show the formation of the ionic compound MgO.

Answer available at the end of this section.

Some ionic compounds have different numbers of cations and anions. In those cases, electron transfer occurs between more than one atom. For example, here is the formation of $MgBr_2$:



Notice that in this example there are two bromide ions (1– charge) needed for every one magnesium ion (2+ charge) in order for the overall charge of the compound to equal zero. This is called **charge balance**. The number of each type of ion is indicated in the formula by the subscript.

Most of the elements that make ionic compounds form an ion that has a characteristic charge. For example, sodium makes ionic compounds in which the sodium ion always has a 1+ charge. Chlorine makes ionic compounds in which the chloride ion always has a 1– charge. Some elements, especially transition metals, can form ions of multiple charges. Figure 5.3.5 shows the characteristic charges for some of these ions. As we saw in Figure 5.3.1, there is a pattern to the charges on many of the main group ions, but there is no simple pattern for transition metal ions (or for the larger main group elements).




Figure 5.3.5: Charges of the Monatomic Ions. Note that some atoms commonly form ions of variable charges.

Key Takeaways

- Ions can be positively charged or negatively charged.
- A Lewis diagram is used to show how electrons are transferred to make ions and ionic compounds.

Exercises

1. What are the two types of ions?
2. Use Lewis diagrams to illustrate the formation of an ionic compound from a potassium atom and an iodine atom.
3. When the following atoms become ions, what charges do they acquire?

    a. Li

    b. S

    c. Ca

    d. F

4. Identify each as a cation, an anion, or neither.

    a. H^+
    b. Cl^-
    c. O_2
    d. Ba^{2+}
    e. CH_4
    f. CS_2

5. Identify each as a cation, an anion, or neither.

    a. NH_3
    b. Br^-
    c. H^-
    d. Hg^{2+}
    e. CCl_4
    f. SO_3

6. Draw Lewis diagrams for the ions listed in Exercise 6. Also include Lewis diagrams for the respective neutral atoms as a comparison.

7. What characteristic charge do atoms in the first column of the periodic table have when they become ions?


8. What characteristic charge do atoms in the second column of the periodic table have when they become ions?

9. What characteristic charge do atoms in the third-to-last column of the periodic table have when they become ions?

10. What characteristic charge do atoms in the next-to-last column of the periodic table have when they become ions?

Answers


1. Cations (positive charged) and anions (negative charged).


2. 


3.
   a. 1+
   b. 2–
   c. 2+
   d. 1–


4.
   a. cation
   b. anion
   c. neither
   d. cation
   e. neither
   f. neither

5.
   a. neither
   b. anion
   c. anion
   d. cation
   e. neither
   f. neither

6. 

   b. 

   c. 

   d. 

7. 1+

8. 2+

9. 2–

10. 1–

Answers


Exercise 5.3.1

Oxygen has 8 electrons. Two electrons in shell 1 and 6 electrons in its valence shell. Oxygen will gain 2 electrons to get an octet. The resulting anion is O^{2-} .

Exercise 5.3.2

(a) S is in Group 6A and has six valence electrons. It achieves octet by gaining two electrons to form S^{2-} anion. Gaining three electrons to form S^{3-}does not make it octet, hence, S^{3-} is not likely to form.

Exercise 5.3.3



This page titled 5.3: Ions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.2: Ions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.4: Formulas for Ionic Compounds

Learning Objectives

- Write the chemical formula for a simple ionic compound.
- Recognize polyatomic ions in chemical formulas.

We have already encountered some chemical formulas for simple ionic compounds. A chemical formula is a concise list of the elements in a compound and the ratios of these elements. To better understand what a chemical formula means, we must consider how an ionic compound is constructed from its ions.

Ionic compounds exist as alternating positive and negative ions in regular, three-dimensional arrays called crystals (Figure 5.4.1). As you can see, there are no individual NaCl "particles" in the array; instead, there is a continuous lattice of alternating sodium and chloride ions. However, we can use the ratio of sodium ions to chloride ions, expressed in the lowest possible whole numbers, as a way of describing the compound. In the case of sodium chloride, the ratio of sodium ions to chloride ions, expressed in lowest whole numbers, is 1:1, so we use NaCl (one Na symbol and one Cl symbol) to represent the compound. Thus, NaCl is the chemical formula for sodium chloride, which is a concise way of describing the relative number of different ions in the compound. A macroscopic sample is composed of myriads of NaCl pairs; each individual pair called a **formula unit**. Although it is convenient to think that NaCl crystals are composed of individual NaCl units, Figure 5.4.1 shows that no single ion is exclusively associated with any other single ion. Each ion is surrounded by ions of opposite charge.




Figure 5.4.1: A Sodium Chloride Crystal. A crystal contains a three-dimensional array of alternating positive and negative ions. The precise pattern depends on the compound. A crystal of sodium chloride, shown here, is a collection of alternating sodium and chlorine ions.

The formula for an ionic compound follows several conventions. First, the **cation** is written **before** the **anion**. Because most metals form cations and most nonmetals form anions, formulas typically list the metal first and then the nonmetal. Second, **charges** are **not written** in a formula. Remember that in an ionic compound, the component species are ions, not neutral atoms, even though the formula does not contain charges. Finally, the proper formula for an ionic compound always has ***a net zero charge,*** meaning the total positive charge must equal the total negative charge. To determine the proper formula of any combination of ions, determine how many of each ion is needed to balance the total positive and negative charges in the compound.

This rule is ultimately based on the fact that matter is, overall, electrically neutral.

> By convention, assume that there is only one atom if a subscript is not present. We do not use 1 as a subscript.

If we look at the ionic compound consisting of lithium ions and bromide ions, we see that the lithium ion has a 1+ charge and the

bromide ion has a 1– charge. Only one ion of each is needed to balance these charges. The formula for lithium bromide is LiBr.

When an ionic compound is formed from magnesium and oxygen, the magnesium ion has a 2+ charge, and the oxygen atom has a 2– charge. Although both of these ions have higher charges than the ions in lithium bromide, they still balance each other in a one-to-one ratio. Therefore, the proper formula for this ionic compound is MgO.

Now consider the ionic compound formed by magnesium and chlorine. A magnesium ion has a 2+ charge, while a chlorine ion has a 1– charge:

$$\mathrm{Mg^{2+}Cl^{-}}$$

Combining one ion of each does not completely balance the positive and negative charges. The easiest way to balance these charges is to assume the presence of *two* chloride ions for each magnesium ion:

$$\mathrm{Mg^{2+}\ Cl^{-}\ Cl^{-}}$$

Now the positive and negative charges are balanced. We could write the chemical formula for this ionic compound as MgClCl, but the convention is to use a numerical subscript when there is more than one ion of a given type—$\mathrm{MgCl_2}$. This chemical formula says that there are one magnesium ion and *two* chloride ions in this formula. (Do not read the "$\mathrm{Cl_2}$" part of the formula as a molecule of the diatomic elemental chlorine. Chlorine does not exist as a diatomic element in this compound. Rather, it exists as two individual chloride ions.) By convention, the **lowest whole number ratio** is used in the formulas of ionic compounds. The formula $\mathrm{Mg_2Cl_4}$ has balanced charges with the ions in a 1:2 ratio, but it is not the lowest whole number ratio.

> By convention, the lowest whole-number ratio of the ions is used in ionic formulas. There are exceptions for certain ions, such as $\mathrm{Hg_2^{2+}}$.

For compounds in which the ratio of ions is not as obvious, the subscripts in the formula can be obtained by **crossing charges**: use the absolute value of the charge on one ion as the subscript for the other ion. This method is shown schematically in Figure $5.4.2$ and Figure $5.4.3$.




Figure $5.4.2$: Crossing charges. One method for obtaining subscripts in the empirical formula is by crossing charges.




Figure 5.4.3: Using the method above to determine the formula for aluminum oxide.

When crossing charges, it is sometimes necessary to reduce the subscripts to their simplest ratio to write the empirical formula. Consider, for example, the compound formed by Pb^{4+} and O^{2-}. Using the absolute values of the charges on the ions as subscripts gives the formula Pb_2O_4. This simplifies to its correct empirical formula **PbO_2**. The empirical formula has one Pb^{4+} ion and two O^{2-} ions.

Example 5.4.1

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the sodium ion and the sulfur ion
b. the aluminum ion and the fluoride ion
c. the 3+ iron ion and the oxygen ion

Solution

a. To obtain a valence shell octet, sodium forms an ion with a 1+ charge, while the sulfur ion has a 2– charge. Two sodium 1+ ions are needed to balance the 2– charge on the sulfur ion. Rather than writing the formula as NaNaS, we shorten it by convention to $\mathrm{Na_2S}$.
b. The aluminum ion has a 3+ charge, while the fluoride ion formed by fluorine has a 1– charge. Three fluorine 1– ions are needed to balance the 3+ charge on the aluminum ion. This combination is written as $\mathrm{AlF_3}$.
c. Iron can form two possible ions, but the ion with a 3+ charge is specified here. The oxygen atom has a 2– charge as an ion. To balance the positive and negative charges, we look to the least common multiple—6: two iron 3+ ions will give 6+, while three 2– oxygen ions will give 6–, thereby balancing the overall positive and negative charges. Thus, the formula for this ionic compound is $\mathrm{Fe_2O_3}$. Alternatively, use the crossing charges method shown in Figure 3.3.2.

Exercise 5.4.1

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the calcium ion and the oxygen ion
b. the 2+ copper ion and the sulfur ion
c. the 1+ copper ion and the sulfur ion

Answer a: available at the end of this section.

Polyatomic Ions

Some ions consist of groups of atoms ***covalently*** bonded together and have an **overall electric charge**. Because these ions

contain more than one atom, they are called polyatomic ions. The Lewis structures, names and formulas of some polyatomic ions are found in Figure 5.4.4.




Figure 5.4.4: A visual guide to the polyatomic ions.

Some Polyatomic Ions

Polyatomic ions have defined formulas, names, and charges that cannot be modified in any way. Table 5.4.1 lists the ion names and ion formulas of the most common polyatomic ions. For example, NO_3^- is the nitrate ion; it has one nitrogen atom and three oxygen atoms and an overall 1– charge. Figure 5.4.4 lists the most common polyatomic ions.

Table 5.4.1: Ion Names and Ion Formulas of Common Polyatomic Ions

Ion Name	Ion Formula
ammonium ion	NH_4^{+1}
hydroxide ion	OH^{-1}
cyanide ion	CN^{-1}
carbonate ion	CO_3^{-2}
bicarbonate or hydrogen carbonate	HCO_3^-
acetate ion	$C_2H_3O_2^{-1}$ **or** $CH_3CO_2^{-1}$
nitrate ion	NO_3^{-1}
nitrite ion	NO_2^{-1}
sulfate ion	SO_4^{-2}
sulfite ion	SO_3^{-2}
phosphate ion	PO_4^{-3}
phosphite ion	PO_3^{-3}

Note that only one polyatomic ion in this Table, the ammonium ion ($NH_4^{+1)}$, is a cation. This polyatomic ion contains one nitrogen

and four hydrogens that collectively bear a +1 charge. The remaining polyatomic ions are all negatively-charged and, therefore, are classified as anions. However, only two of these, the hydroxide ion and the cyanide ion, are named using the "-ide" suffix that is typically indicative of negatively-charged particles. The remaining polyatomic anions, which all contain oxygen, in combination with another non-metal, exist as part of a series in which the number of oxygen atoms within the polyatomic unit can vary. As has been repeatedly emphasized in several sections of this text, no two chemical formulas should share a common chemical name. A single suffix, "-ide," is insufficient for distinguishing the names of the anions in a related polyatomic series. Therefore, "-ate" and "-ite" suffixes are employed, in order to denote that the corresponding polyatomic ions are part of a series. Additionally, these suffixes also indicate the relative number of oxygen atoms that are contained within the polyatomic ions. Note that all of the polyatomic ions whose names end in "-ate" contain one more oxygen than those polyatomic anions whose names end in "-ite." Unfortunately, much like the common system for naming transition metals, these suffixes only indicate the *relative* number of oxygens that are contained within the polyatomic ions. For example, the nitrate ion, which is symbolized as NO_3^{-1}, has one more oxygen than the nitrite ion, which is symbolized as NO_2^{-1}. However, the sulfate ion is symbolized as SO_4^{-2}. While both the nitrate ion and the sulfate ion share an "-ate" suffix, the former contains three oxygens, but the latter contains four. Additionally, both the nitrate ion and the sulfite ion contain three oxygens, but these polyatomic ions do not share a common suffix. Unfortunately, the relative nature of these suffixes mandates that the ion formula/ion name combinations of the polyatomic ions must simply be memorized.

The rule for constructing formulas for ionic compounds containing polyatomic ions is the same as for formulas containing monatomic (single-atom) ions: the positive and negative charges must balance. If more than one of a particular polyatomic ion is needed to balance the charge, the *entire formula* for the polyatomic ion must be enclosed in parentheses, and the numerical subscript is placed *outside* the parentheses. This is to show that the subscript applies to the entire polyatomic ion. Two examples are shown below:




Figure 5.4.4:Formulas with polyatomic ions can be found with the charge crossing method used previously.

Example 5.4.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the potassium ion and the sulfate ion
b. the calcium ion and the nitrate ion

Solution

a. Potassium ions have a charge of 1+, while sulfate ions have a charge of 2–. We will need two potassium ions to balance the charge on the sulfate ion, so the proper chemical formula is K_2SO_4.

b. Calcium ions have a charge of 2+, while nitrate ions have a charge of 1–. We will need two nitrate ions to balance the charge on each calcium ion. The formula for nitrate must be enclosed in parentheses. Thus, we write $\mathrm{Ca(NO_3)_2}$ as the formula for this ionic compound.

Exercise 5.4.2

Write the chemical formula for an ionic compound composed of each pair of ions.

a. the magnesium ion and the carbonate ion
b. the aluminum ion and the acetate ion

Answer a: available at the end of this section.

Recognizing Ionic Compounds

There are two ways to recognize ionic compounds. First, compounds between **metal and nonmetal** elements are usually **ionic**. For example, $CaBr_2$ contains a metallic element (calcium, a group 2A metal) and a nonmetallic element (bromine, a group 7A nonmetal). Therefore, it is most likely an ionic compound. (In fact, it *is* ionic.) In contrast, the compound NO_2 contains two elements that are both nonmetals (nitrogen, from group 5A, and oxygen, from group 6A). It is not an ionic compound; it belongs to the category of covalent compounds discuss elsewhere. Also note that this combination of nitrogen and oxygen has no electric charge specified, so it is *not* the nitrite ion.

Second, if you recognize the formula of a **polyatomic ion** in a compound, the compound is **ionic**. For example, if you see the formula $\mathrm{Ba(NO_3)_2}$, you may recognize the "NO_3" part as the nitrate ion, $\mathrm{NO_3^-}$. (Remember that the convention for writing formulas for ionic compounds is not to include the ionic charge.) This is a clue that the other part of the formula, Ba, is actually the $\mathrm{Ba^{2+}}$ ion, with the 2+ charge balancing the overall 2– charge from the two nitrate ions. Thus, this compound is also ionic.

Example 5.4.3

Identify each compound as ionic or not ionic.

a. $\mathrm{Na_2O}$
b. $\mathrm{PCl_3}$
c. $\mathrm{NH_4Cl}$
d. $\mathrm{OF_2}$

Solution

a. Sodium is a metal, and oxygen is a nonmetal; therefore, $\mathrm{Na_2O}$ is expected to be ionic.
b. Both phosphorus and chlorine are nonmetals. Therefore, $\mathrm{PCl_3}$ is not ionic.
c. The $\mathrm{NH_4}$ in the formula represents the ammonium ion, $\mathrm{NH_4^+}$, which indicates that this compound is ionic.
d. Both oxygen and fluorine are nonmetals. Therefore, $\mathrm{OF_2}$ is not ionic.

Exercise 5.4.3

Identify each compound as ionic or not ionic.

a. $\mathrm{N_2O}$
b. $\mathrm{FeCl_3}$
c. $\mathrm{(NH_4)_3PO_4}$
d. $\mathrm{SOCl_2}$

Answer a: available at the end of this section.

Looking Closer: Blood and Seawater

Science has long recognized that blood and seawater have similar compositions. After all, both liquids have ionic compounds dissolved in them. The similarity may be more than mere coincidence; many scientists think that the first forms of life on Earth arose in the oceans. A closer look, however, shows that blood and seawater are quite different. A 0.9% solution of sodium chloride approximates the salt concentration found in blood. In contrast, seawater is principally a 3% sodium chloride solution, over three times the concentration in blood. Here is a comparison of the amounts of ions in blood and seawater:

Table showing a comparison of the amounts of ions in blood and seawater.

Ion	Percent in Seawater	Percent in Blood
Na^+	2.36	0.322
Cl^-	1.94	0.366
Mg^{2+}	0.13	0.002
SO_4^{2-}	0.09	—
K^+	0.04	0.016
Ca^{2+}	0.04	0.0096
HCO_3^-	0.002	0.165
HPO_4^{2-}, $H_2PO_4^-$	—	0.01

Most ions are more abundant in seawater than they are in blood, with some important exceptions. There are far more hydrogen carbonate ions (HCO_3^-) in blood than in seawater. This difference is significant because the hydrogen carbonate ion and some related ions have a crucial role in controlling the acid-base properties of blood. The amount of hydrogen phosphate ions—HPO_4^{2-} and $H_2PO_4^-$—in seawater is very low, but they are present in higher amounts in blood, where they also affect acid-base properties. Another notable difference is that blood does not have significant amounts of the sulfate ion (SO_4^{2-}), but this ion is present in seawater.

Key Takeaways

- Proper chemical formulas for ionic compounds balance the total positive charge with the total negative charge.
- Groups of atoms with an overall charge, called polyatomic ions, also exist.

EXERCISES

1. What information is contained in the formula of an ionic compound?

2. Why do the chemical formulas for some ionic compounds contain subscripts, while others do not?

3. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Mg^{2+} and I^-
   b. Na^+ and O^{2-}

4. Write the chemical formula for the ionic compound formed by each pair of ions.

   a. Na^+ and Br^-

b. Mg^{2+} and Br^{-}
c. Mg^{2+} and S^{2-}

5. Write the chemical formula for the ionic compound formed by each pair of ions.

a. K^{+} and Cl^{-}
b. Mg^{2+} and Cl^{-}
c. Mg^{2+} and Se^{2-}

6. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Na^{+} and N^{3-}
b. Mg^{2+} and N^{3-}
c. Al^{3+} and S^{2-}

7. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Li^{+} and N^{3-}
b. Mg^{2+} and P^{3-}
c. Li^{+} and P^{3-}

8. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Fe^{3+} and Br^{-}
b. Fe^{2+} and Br^{-}
c. Au^{3+} and S^{2-}
d. Au^{+} and S^{2-}

9. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Cr^{3+} and O^{2-}
b. Cr^{2+} and O^{2-}
c. Pb^{2+} and Cl^{-}
d. Pb^{4+} and Cl^{-}

10. Write the chemical formula for the ionic compound formed by each pair of ions.

a. Cr^{3+} and NO_3^{-}
b. Fe^{2+} and PO_4^{3-}
c. Ca^{2+} and CrO_4^{2-}
d. Al^{3+} and OH^{-}

11. Write the chemical formula for the ionic compound formed by each pair of ions.

a. NH_4^{+} and NO_3^{-}
b. H^{+} and $Cr_2O_7^{2-}$
c. Cu^{+} and CO_3^{2-}
d. Na^{+} and HCO_3^{-}

12. For each pair of elements, determine the charge for their ions and write the proper formula for the resulting ionic compound between them.

a. Ba and S
b. Cs and I

13. For each pair of elements, determine the charge for their ions and write the proper formula for the resulting ionic compound between them.

a. K and S
b. Sc and Br

14. Which compounds would you predict to be ionic?

a. Li_2O
b. $(NH_4)_2O$
c. CO_2
d. $FeSO_3$
e. C_6H_6
f. C_2H_6O

15. Which compounds would you predict to be ionic?

a. $Ba(OH)_2$
b. CH_2O
c. NH_2CONH_2
d. $(NH_4)_2CrO_4$
e. C_8H_{18}
f. NH_3

Answers

1. the ratio of each kind of ion in the compound

2. Sometimes more than one ion is needed to balance the charge on the other ion in an ionic compound.

3.

a. MgI_2
b. Na_2O

4.

a. NaBr
b. $MgBr_2$
c. MgS

5.

a. KCl
b. $MgCl_2$
c. MgSe

6.

a. Na_3N
b. Mg_3N_2
c. Al_2S_3

7.

a. Li_3N
b. Mg_3P_2
c. Li_3P

8.

a. $FeBr_3$
b. $FeBr_2$
c. Au_2S_3
d. Au_2S

9.

a. Cr_2O_3
b. CrO
c. $PbCl_2$
d. $PbCl_4$

10.

a. $Cr(NO_3)_3$
b. $Fe_3(PO_4)_2$
c. $CaCrO_4$
d. $Al(OH)_3$

11.

a. NH_4NO_3
b. $H_2Cr_2O_7$
c. Cu_2CO_3
d. $NaHCO_3$

12.

a. Ba^{2+}, S^{2-}, BaS
b. Cs^{+}, I^{-}, CsI

13.

a. K^{+}, S^{2-}, K_2S
b. Sc^{3+}, Br^{-}, $ScBr_3$

14.

a. ionic
b. ionic
c. not ionic
d. ionic
e. not ionic
f. not ionic

15.

a. ionic
b. not ionic
c. not ionic
d. ionic

e. not ionic
f. not ionic

Answers

Exercise 5.4.1

CaO
CuS
Cu_2S

Exercise 5.4.2

Mg^{2+} and CO_3^{2-} = $MgCO_3$

Al^{3+} and $C_2H_3O_2^-$ = $Al(C_2H_3O_2)_3$

Exercise 5.4.3

not ionic
ionic
ionic
not ionic

This page titled 5.4: Formulas for Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.3: Formulas for Ionic Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.5: Ionic Nomenclature

Learning Objectives

- To use the rules for naming ionic compounds

After learning a few more details about the names of individual ions, you will be a step away from knowing how to name ionic compounds. This section begins the formal study of nomenclature, the systematic naming of chemical compounds.

Naming Ions

The name of a monatomic cation is simply the name of the element followed by the word *ion*. Thus, Na^+ is the sodium ion, Al^{3+} is the aluminum ion, Ca^{2+} is the calcium ion, and so forth.

We have seen that some elements lose different numbers of electrons, producing ions of different charges. Iron, for example, can form two cations, each of which, when combined with the same anion, makes a different compound with unique physical and chemical properties. Thus, we need a different name for each iron ion to distinguish Fe^{2+} from Fe^{3+}. The same issue arises for other ions with more than one possible charge.

There are two ways to make this distinction. In the simpler, more modern approach, called the ***Stock system***, an ion's positive charge is indicated by a roman numeral in parentheses after the element name, followed by the word *ion*. Thus, Fe^{2+} is called the iron(II) ion, while Fe^{3+} is called the iron(III) ion. This system is used only for elements that form more than one common positive ion. We do not call the Na^+ ion the sodium(I) ion because (I) is unnecessary. Sodium forms only a 1+ ion, so there is no ambiguity about the name *sodium ion*.

Table 5.5.1: The Common System of Cation Names

Element	Stem	Charge	Name
iron	ferr-	2+	ferrous ion
		3+	ferric ion
copper	cupr-	1+	cuprous ion
		2+	cupric ion
tin	stann-	2+	stannous ion
		4+	stannic ion
lead	plumb-	2+	plumbous ion
		4+	plumbic ion
chromium	chrom-	2+	chromous ion
		3+	chromic ion
gold	aur-	1+	aurous ion
		3+	auric ion

The second system, called the **common system**, is not conventional but is still prevalent and used in the health sciences. This system recognizes that many metals have two common cations. The common system uses two suffixes (*-ic* and *-ous*) that are appended to the stem of the element name. The *-ic* suffix represents the greater of the two cation charges, and the *-ous* suffix represents the lower one. In many cases, the stem of the element name comes from the Latin name of the element. Table 5.5.1 lists the elements that use the common system, along with their respective cation names.

The name of a monatomic anion consists of the stem of the element name, the suffix *-ide*, and then the word *ion*. Thus, as we have already seen, Cl^- is "chlor-" + "-ide ion," or the chloride ion. Similarly, O^{2-} is the oxide ion, Se^{2-} is the selenide ion, and so forth. Table 5.5.2 lists the names of some common monatomic ions.

Table 5.5.2: Some Monatomic Anions

Ion	Name
F^-	fluoride ion
Cl^-	chloride ion
Br^-	bromide ion
I^-	iodide ion
O^{2-}	oxide ion
S^{2-}	sulfide ion
P^{3-}	phosphide ion
N^{3-}	nitride ion

The polyatomic ions have their own characteristic names, as discussed earlier.

Example 5.5.1

Name each ion.

a. Ca^{2+}
b. S^{2-}
c. SO_3^{2-}
d. NH_4^+
e. Cu^+

Answer a available at the end of this section.

Exercise 5.5.1

Name each ion.

a. Fe^{2+}
b. Fe^{3+}
c. SO_4^{2-}
d. Ba^{2+}
e. HCO_3^-

Answer a available at the end of this section.

Example 5.5.2

Write the formula for each ion.

a. the bromide ion
b. the phosphate ion
c. the cupric ion
d. the magnesium ion

Answer a available at the end of this section.

Exercise 5.5.2


Write the formula for each ion.

a. the fluoride ion
b. the carbonate ion
c. the ferrous ion
d. the potassium ion

Answer a available at the end of this section.

Naming Compounds

Now that we know how to name ions, we are ready to name ionic compounds. We do so by placing the name of the cation first, followed by the name of the anion, and dropping the word *ion* from both parts. For example, what is the name of the compound whose formula is $\mathrm{Ba(NO_3)_2}$?



The compound's name does not indicate that there are two nitrate ions for every barium ion. You must determine the relative numbers of ions by balancing the positive and negative charges.

If you are given a formula for an ionic compound whose cation can have more than one possible charge, you must first determine the charge on the cation before identifying its correct name. For example, consider $\mathrm{FeCl_2}$ and $\mathrm{FeCl_3}$. In the first compound, the iron ion has a 2+ charge because there are two $\mathrm{Cl^-}$ ions in the formula (1– charge on each chloride ion). In the second compound, the iron ion has a 3+ charge, as indicated by the three $\mathrm{Cl^-}$ ions in the formula. These are two different compounds that need two different names. By the Stock system, the names are iron(II) chloride and iron(III) chloride. If we were to use the stems and suffixes of the common system, the names would be ferrous chloride and ferric chloride, respectively.

Example 5.5.3

Name each ionic compound, using both Stock and common systems if necessary.

a. $Ca_3(PO_4)_2$
b. $(NH_4)_2Cr_2O_7$
c. KCl
d. CuCl
e. SnF_2

Answer a available at the end of this section.

Exercise 5.5.3

Name each ionic compound, using both Stock and common systems if necessary.

a. $ZnBr_2$
b. $Fe(NO_3)_3$
c. Al_2O_3
d. CuF_2
e. AgF

Answer a available at the end of this section.

Figure 5.5.1 is a synopsis of how to name simple ionic compounds.




Figure 5.5.1: A Guide to Naming Simple Ionic Compounds. Follow these steps to name a simple ionic compound.

KEY TAKEAWAY

- Each ionic compound has its own unique name that comes from the names of the ions.

EXERCISES

1. Briefly describe the process for naming an ionic compound.

- In what order do the names of ions appear in the names of ionic compounds?

3. Which ionic compounds can be named using two different systems? Give an example.

4. Name each ion.

a. Ra^{2+}
b. P^{3-}
c. $H_2PO_4^-$
d. Sn^{4+}

5. Name each ion.

a. Cs^+
b. As^{3-}
c. HSO_4^-
d. Sn^{2+}

6. Name the ionic compound formed by each pair of ions.

a. Na^+ and Br^-
b. Mg^{2+} and Br^-
c. Mg^{2+} and S^{2-}

7. Name the ionic compound formed by each pair of ions.

a. K^+ and Cl^-
b. Mg^{2+} and Cl^-
c. Mg^{2+} and Se^{2-}

8. Name the ionic compound formed by each pair of ions.

a. Na^+ and N^{3-}
b. Mg^{2+} and N^{3-}
c. Al^{3+} and S^{2-}

9. Name the ionic compound formed by each pair of ions.

a. Li^+ and N^{3-}
b. Mg^{2+} and P^{3-}
c. Li^+ and P^{3-}

10. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

a. Fe^{3+} and Br^-
b. Fe^{2+} and Br^-
c. Au^{3+} and S^{2-}
d. Au^+ and S^{2-}

11. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

a. Cr^{3+} and O^{2-}
b. Cr^{2+} and O^{2-}
c. Pb^{2+} and Cl^-
d. Pb^{4+} and Cl^-

12. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

a. Cr^{3+} and NO_3^-
b. Fe^{2+} and PO_4^{3-}
c. Ca^{2+} and CrO_4^{2-}
d. Al^{3+} and OH^-

13. Name the ionic compound formed by each pair of ions. Use both the Stock and common systems, where appropriate.

a. NH_4^+ and NO_3^-
b. K^+ and $Cr_2O_7^{2-}$
c. Cu^+ and CO_3^{2-}
d. Na^+ and HCO_3^-

14. Give two names for each compound.

a. $Al(HSO_4)_3$
b. $Mg(HSO_4)_2$

15. Give two names for each compound.

a. $Co(HCO_3)_2$
b. $LiHCO_3$

AnswerS

1. Name the cation and then the anion but don't use numerical prefixes.
2. the cation name followed by the anion name
3. Ionic compounds in which the cation can have more than one possible charge have two naming systems. $FeCl_3$ is either iron(III) chloride or ferric chloride (answers will vary).

4.

a. the radium ion
b. the phosphide ion
c. the dihydrogen phosphate ion
d. the tin(IV) ion or the stannic ion

5.

a. the cesium ion
b. the arsenide ion
c. the hydrogen sulfate ion
d. the tin(II) ion or the stannous ion

6.

a. sodium bromide
b. magnesium bromide
c. magnesium sulfide

7.

a. potassium chloride
b. magnesium chloride
c. magnesium selenide

8.

a. sodium nitride
b. magnesium nitride
c. aluminum sulfide

9.

a. lithium nitride
b. magnesium phosphide
c. lithium phosphide

10.

a. iron(III) bromide or ferric bromide
b. iron(II) bromide or ferrous bromide
c. gold(III) sulfide or auric sulfide
d. gold(I) sulfide or aurous sulfide

11.

a. chromium(III) oxide or chromic oxide
b. chromium(II) oxide or chromous oxide
c. lead(II) chloride or plumbous chloride
d. lead(IV) chloride or plumbic chloride

12.

a. chromium(III) nitrate or chromic nitrate
b. iron(II) phosphate or ferrous phosphate
c. calcium chromate
d. aluminum hydroxide

13.

a. ammonium nitrate
b. potassium dichromate
c. copper(I) carbonate or cuprous carbonate
d. sodium hydrogen carbonate or sodium bicarbonate

14.

a. aluminum hydrogen sulfate or aluminum bisulfate
b. magnesium hydrogen sulfate or magnesium bisulfate

15.

a. cobalt hydrogen carbonate or cobalt bicarbonate
b. lithium hydrogen carbonate or lithium bicarbonate

Answers

Example 5.5.1

the calcium ion
the sulfide ion (from Table 5.5.2)
the sulfite ion
the ammonium ion
the copper(I) ion or the cuprous ion (copper can form cations with either a 1+ or 2+ charge, so we have to specify which charge this ion has

Exercise 5.5.1

the iron (II) or ferrous ion
the iron (III) or ferric ion
the sulfate ion

the barium ion
the bicarbonate ion or hydrogen carbonate ion

Example 5.5.2

Br^-

PO_4^{3-}

Cu^{2+}

Mg^{2+}

Exercise 5.5.2

F^-

CO_3^{2-}

Fe^{2+}

K^+

Example 5.5.3

calcium phosphate
ammonium dichromate (the prefix *di-* is part of the name of the anion)
potassium chloride
copper(I) chloride or cuprous chloride
tin(II) fluoride or stannous fluoride

Exercise 5.5.3

zinc bromide
iron (III) nitrate or ferric nitrate
aluminum oxide
copper (II) fluoride or cupric fluoride
silver fluoride

This page titled 5.5: Ionic Nomenclature is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.4: Ionic Nomenclature by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.6: Formula Mass

Learning Objectives

- To determine the formula mass of an ionic compound.

One skill needed in future chapters is the ability to determine the mass of the formula of an ionic compound. This quantity is called the formula mass. The formula mass is obtained by adding the masses of each individual atom in the formula of the compound. Because a proper formula is electrically neutral (with no net electrons gained or lost), the ions can be considered atoms for the purpose of calculating the formula mass.

Let us start by calculating the formula mass of sodium chloride (NaCl). This formula mass is the sum of the atomic masses of one sodium atom and one chlorine atom, which we find from the periodic table; here, we use the masses to two decimal places:

Table shows how to calculate the formula mass of sodium chloride by using their atomic masses.

Na:	22.99 amu
Cl:	+ 35.45 amu
Total:	58.44 amu

To two decimal places, the formula mass of NaCl is 58.44 amu.

When an ionic compound has more than one anion or cation, you must remember to use the proper multiple of the atomic mass for the element in question. For the formula mass of calcium fluoride (CaF_2), we must multiply the mass of the fluorine atom by 2 to account for the two fluorine atoms in the chemical formula:

Table shows how to find the formula mass of calcium fluoride by using their atomic masses.

Ca:	1 × 40.08	40.08 amu
F:	2 × 19.00 =	+ 38.00 amu
Total:		78.08 amu

The formula mass of CaF_2 is 78.08 amu.

For ionic compounds with polyatomic ions, the sum must include the number and mass of each atom in the formula for the polyatomic ion. For example, potassium nitrate (KNO_3) has one potassium atom, one nitrogen atom, and three oxygen atoms:

Table shows how to find the formula mass of potassium nitrate by using their atomic masses.

K:	1 × 39.10	39.10 amu
N:	1 × 14.01	+ 14.01 amu
O:	3 × 16.00 =	+ 48.00 amu
Total:		101.11 amu

The formula mass of KNO_3 is 101.11 amu.

> Potassium nitrate is a key ingredient in gunpowder and has been used clinically as a diuretic.

When a formula contains more than one polyatomic unit in the chemical formula, as in $Ca(NO_3)_2$, do not forget to multiply the

atomic mass of every atom inside the parentheses by the subscript outside the parentheses. This is necessary because the subscript refers to the *entire polyatomic ion*. Thus, for $Ca(NO_3)_2$, the subscript 2 implies two complete nitrate ions, so we must sum the masses of two (1 × 2) nitrogen atoms and six (3 × 2) oxygen atoms, along with the mass of a single calcium atom:

Table shows how to find the formula mass of $Ca(NO_3)_2$ by using their atomic masses.

Ca:	1 × 40.08	40.08 amu
N:	2 × 14.01 =	+ 28.02 amu
O:	6 × 16.00 =	+ 96.00 amu
Total:		164.10 amu

The key to calculating the formula mass of an ionic compound is to correctly count each atom in the formula and multiply the atomic masses of its atoms accordingly.

The Periodic Table is found in this link: Periodic Table of Elements with Atomic Mass [pubchem.ncbi.nlm.nih.gov]

Example 5.6.1

Use the atomic masses (**rounded to two decimal places**) from the **Periodic Table** to determine the formula mass for each ionic compound.

a. $FeCl_3$
b. $(NH_4)_3PO_4$

Solution

a.

Table shows how to find the formula mass of $FeCl_3$ by using their atomic masses.

Fe:		55.84 amu
Cl:	3 × 35.45 =	+ 106.35 amu
Total:		162.19 amu

The formula mass of $FeCl_3$ is 162.19 amu.

a. When we distribute the subscript 3 through the parentheses containing the formula for the ammonium ion, we see that we have 3 nitrogen atoms and 12 hydrogen atoms. Thus, we set up the sum as follows:

Table shows how to find the formula mass of $(NH_4)_3PO_4$ by using their atomic masses.

N:	3 × 14.01 =	42.03 amu
H:	12 × 1.01 =	+ 12.12 amu
P:		+ 30.97 amu
O:	4 × 16.00 =	+ 64.00 amu
Total:		149.12 amu

The formula mass for $(NH_4)_3PO_4$ is 149.12 amu.

Exercise 5.6.1

Use the atomic masses (**rounded to two decimal places**) from the **Periodic Table** to determine the formula mass for each ionic compound.

a. TiO_2
b. AgBr
c. $Au(NO_3)_3$
d. $Fe_3(PO_4)_2$

Answer a: available at the end of this section.

To Your Health: Hydrates

Some ionic compounds have water (H_2O) incorporated within their formula unit. These compounds, called *hydrates*, have a characteristic number of water units associated with each formula unit of the compound. Hydrates are solids, not liquids or solutions, despite the water they contain.

To write the chemical formula of a hydrate, write the number of water units per formula unit of compound after its chemical formula. The two chemical formulas are separated by a vertically centered dot. The hydrate of copper(II) sulfate has five water units associated with each formula unit, so it is written as $CuSO_4 \cdot 5\,H_2O$. The name of this compound is copper(II) sulfate pentahydrate, with the *penta-* prefix indicating the presence of five water units per formula unit of copper(II) sulfate.



Magnesium sulfate heptahydrate. (Public Domain; Chemicalinterest).

Hydrates have various uses in the health industry. Calcium sulfate hemihydrate ($CaSO_4 \cdot ½H_2O$), known as plaster of Paris, is used to make casts for broken bones. Epsom salt ($MgSO_4 \cdot 7\,H_2O$) is used as a bathing salt and a laxative. Aluminum chloride hexahydrate is an active ingredient in antiperspirants. The accompanying table lists some useful hydrates.

Table 5.6.1: Names and Formulas of Some Widely Used Hydrates

Formula	Name	Uses
$AlCl_3•6H_2O$	aluminum chloride hexahydrate	antiperspirant
$CaSO_4•½H_2O$	calcium sulfate hemihydrate (plaster of Paris)	casts (for broken bones and castings)
$CaSO_4•2H_2O$	calcium sulfate dihydrate (gypsum)	drywall component
$CoCl_2•6H_2O$	cobalt(II) chloride hexahydrate	drying agent, humidity indicator

Formula	Name	Uses
$CuSO_4•5H_2O$	copper(II) sulfate pentahydrate	fungicide, algicide, herbicide
$MgSO_4•7H_2O$	magnesium sulfate heptahydrate (Epsom salts)	laxative, bathing salt
$Na_2CO_3•10H_2O$	sodium carbonate decahydrate (washing soda)	laundry additive/cleaneKEY TAKEAWAY

KEY TAKEAWAY

- Formula masses of ionic compounds can be determined from the masses of the atoms in their formulas.

EXERCISES

1. What is the relationship between atomic mass and formula mass?

2. How are subscripts used to determine a formula mass when more than one polyatomic ion is present in a chemical formula?

3. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Na^+ and Br^-
   b. Mg^{2+} and Br^-
   c. Mg^{2+} and S^{2-}

4. What is the formula mass for the ionic compound formed by each pair of ions?

   a. K^+ and Cl^-
   b. Mg^{2+} and Cl^-
   c. Mg^{2+} and Se^{2-}

5. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Na^+ and N^{3-}
   b. Mg^{2+} and N^{3-}
   c. Al^{3+} and S^{2-}

6. What is the formula mass for the ionic compound formed by each pair of ions?

   a. Li^+ and N^{3-}
   b. Mg^{2+} and P^{3-}
   c. Li^+ and P^{3-}

7. What is the formula mass for each compound?

   a. $FeBr_3$
   b. $FeBr_2$
   c. Au_2S_3
   d. Au_2S

8. What is the formula mass for each compound?

   a. Cr_2O_3
   b. CrO
   c. $PbCl_2$

d. $PbCl_4$

9. What is the formula mass for each compound?

a. $Cr(NO_3)_3$
b. $Fe_3(PO_4)_2$
c. $CaCrO_4$
d. $Al(OH)_3$

10. What is the formula mass for each compound?

a. NH_4NO_3
b. $K_2Cr_2O_7$
c. Cu_2CO_3
d. $NaHCO_3$

11. What is the formula mass for each compound?

a. $Al(HSO_4)_3$
b. $Mg(HSO_4)_2$

12. What is the formula mass for each compound?

a. $Co(HCO_3)_2$
b. $LiHCO_3$

Answers

1. The formula mass is the sum of the atomic masses of the atoms in the formula.
2. The subscript is distributed throughout the parentheses to determine the total number of atoms in the formula.

3.

a. 102.89 amu
b. 184.11 amu
c. 56.38 amu

4.

a. 74.55 amu
b. 95.21 amu
c. 103.28 amu

5.

a. 82.98 amu
b. 100.95 amu
c. 150.17 amu

6.

a. 35.01 amu
b. 134.87 amu
c. 51.79 amu

7.

a. 295.55 amu

b. 215.65 amu
c. 490.15 amu
d. 426.01 amu

8.

a. 152.00 amu
b. 68.00 amu
c. 278.10 amu
d. 349.00 amu

9.

a. 238.03 amu
b. 357.49 amu
c. 156.08 amu
d. 78.01 amu

10.

a. 80.06 amu
b. 294.20 amu
c. 187.11 amu
d. 84.01 amu

11.

a. 318.22 amu
b. 218.47 amu

12.

a. 180.97 amu
b. 67.96 amu

Answers

Exercise 5.6.1

47.87 + 2 (16.00) = 79.87 amu
107.87 + 79.90 = 187.77 amu
196.97 + 3(14.01) + 9(16.00) = 383.00 amu
3(55.84) + 2(30.97) + 8(16.00) = 357.46 amu

This page titled 5.6: Formula Mass is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.5: Formula Mass by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.7: Characteristics of Ionic Compounds

Learning Objectives

- To describe the physical properties of ionic compounds.

The figure below shows just a few examples of the color and brilliance of naturally occurring ionic crystals. The regular and orderly arrangement of ions in the crystal lattice is responsible for the various shapes of these crystals, while transition metal ions give rise to the colors.




Figure 3.6.1: *In nature, the ordered arrangement of ionic solids gives rise to beautiful crystals. (A) Amethyst - a form of quartz, SiO_2, whose purple color comes from iron ions. (B) Cinnabar - the primary ore of mercury is mercury (II) sulfide, HgS (C) Azurite - a copper mineral, $Cu_3(CO_3)_2(OH)_2$. (D) Vanadinite - the primary ore of vanadium, $Pb_3(VO_4)_3Cl$.*

Physical Properties of Ionic Compounds

Melting Points

Because of the many simultaneous attractions between cations and anions that occur, ionic crystal lattices are very strong. The process of melting an ionic compound requires the addition of large amounts of energy in order to break all of the ionic bonds in the crystal. For example, sodium chloride has a melting temperature of about 800°C. As a comparison, the molecular compound water melts at 0 °C.

Shattering

Ionic compounds are generally hard, but **brittle.** Why? It takes a large amount of mechanical force, such as striking a crystal with a hammer, to force one layer of ions to shift relative to its neighbor. However, when that happens, it brings ions of the same charge next to each other (see below). The repulsive forces between like-charged ions cause the crystal to shatter. When an ionic crystal breaks, it tends to do so along smooth planes because of the regular arrangement of the ions.




Figure 3.6.2: *(A) The sodium chloride crystal is shown in two dimensions. (B) When struck by a hammer, the negatively-charged chloride ions are forced near each other and the repulsive force causes the crystal to shatter.*

Conductivity

Another characteristic property of ionic compounds is their **electrical conductivity**. The figure below shows three experiments in which two electrodes that are connected to a light bulb are placed in beakers containing three different substances.




Figure 3.6.3: *(A) Distilled water does not conduct electricity. (B) A solid ionic compound also does not conduct. (C) A water solution of an ionic compound conducts electricity well.*

In the first beaker, distilled water does not conduct a current because water is a molecular compound. In the second beaker, solid sodium chloride also does not conduct a current. Despite being ionic and thus composed of charged particles, the solid crystal lattice does not allow the ions to move between the electrodes. Mobile charged particles are required for the circuit to be complete and the light bulb to light up. In the third beaker, the NaCl has been dissolved into the distilled water. Now the crystal lattice has been broken apart and the individual positive and negative ions can move. Cations move to one electrode, while anions move to the other, allowing electricity to flow (see figure below). Melting an ionic compound also frees the ions to conduct a current. Ionic compounds conduct an electric current when melted or dissolved in water. The dissolution of ionic compounds in water will be discussed in Section 9.3.




Figure 3.6.4: *In an ionic solution, the A^+ ions migrate toward the negative electrode, while the B^- ions migrate toward the positive electrode.*

Example 5.7.1

Write the dissociation equation of solid NaCl in water.

Solution

$NaCl(s) \rightarrow Na^+(aq) + Cl^-(aq)$

5.7.1 Exerc

Write the dissociation equation of solid NH_4NO_3 in water.

Answer available at the end of this section.

Key Takeaways

- Ionic compounds have high melting points.
- Ionic compounds are hard and brittle.
- Ionic compounds dissociate into ions when dissolved in water.
- Solutions of ionic compounds and melted ionic compounds conduct electricity, but solid materials do not.
- An ionic compound can be identified by its chemical formula: metal + nonmetal or polyatomic ions.

Contributors and Attributions

- CK-12 Foundation by Sharon Bewick, Richard Parsons, Therese Forsythe, Shonna Robinson, and Jean Dupon.

Exercises

1. Explain how the bonding in an ionic solid explains some of the properties of these solids.

2. Which type(s) of solid conduct(s) electricity in their liquid state but not in their solid state?

3. Based on chemical formula, identify which of the following is an ionic solid?

   a. Hg

   b. PH_3

   c. $Ba(NO_3)_2$

Answers

$NH_4NO_3(s) \rightarrow NH_4^+(aq) + NO_3^-(aq)$

This page titled 5.7: Characteristics of Ionic Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.6: Characteristics of Ionic Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

5.S: Ionic Bonding and Simple Ionic Compounds (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms and ask yourself how they relate to the topics in the chapter.

Atoms combine into compounds by forming **chemical bonds**. A survey of stable atoms and molecules leads to the **octet rule**, which says that stable atoms tend to have eight electrons in their outermost, or valence, shell. One way atoms obtain eight electrons in the valence shell is for some atoms to lose electrons while other atoms gain them. When this happens, the atoms take on an electrical charge. Charged atoms are called **ions**. Ions having opposite charges attract each other. This attraction is called **ionic bonding**, and the compounds formed are called **ionic compounds**.

Positively charged ions are called **cations**, while negatively charged ions are called **anions**. The formation of both cations and anions can be illustrated using electron configurations. Because elements in a column of the periodic table have the same valence shell electron configuration, atoms in the same column of the periodic table tend to form ions having the same charge. **Electron dot diagrams**, or **Lewis diagrams**, can also be used to illustrate the formation of cations and anions.

Ionic compounds are represented in writing by a **chemical formula**, which gives the lowest ratio of cations and anions present in the compound. In a formula, the symbol of the cation is written first, followed by the symbol of the anion. **Formula unit** is considered the basic unit of an ionic compound because ionic compounds do not exist as discrete units. Instead, they exist as **crystals**, three-dimensional arrays of ions, with cations surrounded by anions and anions surrounded by cations. Chemical formulas for ionic compounds are determined by balancing the positive charge from the cation(s) with the negative charge from the anion(s). A subscript to the right of the ion indicates that more than one of that ion is present in the chemical formula.

Some ions are groups of atoms bonded together and having an overall electrical charge. These are called **polyatomic ions**. Writing formulas with polyatomic ions follows the same rules as with monatomic ions, except that when more than one polyatomic ion is present in a chemical formula, the polyatomic ion is enclosed in parentheses and the subscript is outside the right parenthesis. Ionic compounds typically form between metals and nonmetals or between polyatomic ions.

Names of ionic compounds are derived from the names of the ions, with the name of the cation coming first, followed by the name of the anion. If an element can form cations of different charges, there are two alternate systems for indicating the compound's name. In the **Stock system**, a roman numeral in parentheses indicates the charge on the cation. An example is the name for $FeCl_2$, which is iron(II) chloride. In the common system, the suffixes *-ous* and *-ic* are used to stand for the lower and higher possible charge of the cation, respectively. These suffixes are attached to a stem representing the element (which frequently comes from the Latin form of the element name). An example is the common name for $FeCl_2$, which is ferrous chloride.

The **formula mass** of an ionic compound is the sum of the masses of each individual atom in the formula. Care must be taken when calculating formula masses for formulas containing multiple polyatomic ions because the subscript outside the parentheses refers to all the atoms in the polyatomic ion.

Ionic compounds are hard, brittle and have very high melting points. When in solution, the ions separate and form **electrolyte** solutions.

This page titled 5.S: Ionic Bonding and Simple Ionic Compounds (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 3.S: Ionic Bonding and Simple Ionic Compounds (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6: COVALENT BONDING AND SIMPLE MOLECULAR COMPOUNDS

Ionic bonding results from the transfer of electrons among atoms or groups of atoms. In this chapter, we will consider another type of bonding—covalent bonding. We will examine how atoms share electrons to form these bonds, and we will begin to explore how the resulting compounds, such as cholesterol, are different from ionic compounds.

6.1: Prelude to Covalent Bonding and Simple Molecular Compounds

6.2: Covalent Bonds

6.3: Covalent Compounds - Formulas and Names

6.4: Drawing Lewis Structures

6.5: Characteristics of Covalent Bonds

6.6: Characteristics of Molecules and VSEPR

6.7: Organic Chemistry

6.S: Covalent Bonding and Simple Molecular Compounds (Summary)

Template:HideTOC

This page titled 6: Covalent Bonding and Simple Molecular Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4: Covalent Bonding and Simple Molecular Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.1: Prelude to Covalent Bonding and Simple Molecular Compounds

Cholesterol, a compound that is sometimes in the news, is a white, waxy solid produced in the liver of every animal, including humans. It is important for building cell membranes and in producing certain hormones (chemicals that regulate cellular activity in the body). As such, it is necessary for life, but why is cholesterol the object of attention?

Besides producing cholesterol, we also ingest some whenever we eat meat or other animal-based food products. People who eat such products in large quantities, or whose metabolisms are unable to handle excess amounts, may experience an unhealthy buildup of cholesterol in their blood. Deposits of cholesterol, called plaque, may form on blood vessel walls, eventually blocking the arteries and preventing the delivery of oxygen to body tissues. Heart attacks, strokes, and other circulatory problems can result.

Most medical professionals recommend diets that minimize the amount of ingested cholesterol as a way of preventing heart attacks and strokes. Tests are available to measure cholesterol in the blood, and there are several drugs capable of lowering cholesterol levels.




Figure 6.1.1: A Molecular Model of Cholesterol

This page titled 6.1: Prelude to Covalent Bonding and Simple Molecular Compounds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.0: Prelude to Covalent Bonding and Simple Molecular Compounds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.2: Covalent Bonds

Learning Objectives

- To describe how a covalent bond forms.
- To apply the octet rule to covalent compounds

You have already seen examples of substances that contain covalent bonds. One substance mentioned previously was water H_2O. You can tell from its formula that it is not an ionic compound; it is not composed of a metal and a nonmetal. Consequently, its properties are different from those of ionic compounds.


Electron Sharing

Previously, we discussed ionic bonding where electrons can be transferred from one atom to another so that both atoms have an energy-stable outer electron shell. Because most filled electron shells have eight electrons in them, chemists called this tendency the octet rule. However, there is another way an atom can achieve a full valence shell: atoms can *share* electrons.

This concept can be illustrated by using two hydrogen atoms, each of which has a single electron in its valence shell. (For small atoms such as hydrogen atoms, the valence shell will be the first shell, which holds only two electrons.) We can represent the two individual hydrogen atoms as follows:



In contrast, when two hydrogen atoms get close enough together to share their electrons, they can be represented as follows:



By sharing their valence electrons, both hydrogen atoms now have two electrons in their respective valence shells. Because each valence shell is now filled, this arrangement is more stable than when the two atoms are separate. The sharing of electrons between atoms is called a covalent bond, and the two electrons that join atoms in a covalent bond are called a bonding pair of

electrons. A discrete group of atoms connected by covalent bonds is called a molecule—the smallest part of a compound that retains the chemical identity of that compound.

Chemists frequently use Lewis diagrams to represent covalent bonding in molecular substances. For example, the Lewis diagrams of two separate hydrogen atoms are as follows:



The Lewis diagram of two hydrogen atoms sharing electrons looks like this:



This depiction of molecules is simplified further by using a dash to represent a covalent bond. The hydrogen molecule is then represented as follows:



Remember that the dash, also referred to as a single bond, represents a *pair* of electrons.

The bond in a hydrogen molecule, measured as the distance between the two nuclei, is about 7.4×10^{-11} m, or 74 picometers (pm; 1 pm = 1×10^{-12} m). This particular bond length represents a balance between several forces: the attractions between oppositely charged electrons and nuclei, the repulsion between two negatively charged electrons, and the repulsion between two positively charged nuclei. If the nuclei were closer together, they would repel each other more strongly; if the nuclei were farther apart, there would be less attraction between the positive and negative particles.

Fluorine is another element whose atoms bond together in pairs to form *diatomic* (two-atom) molecules. Two separate fluorine atoms have the following electron dot diagrams:




Each fluorine atom contributes one valence electron, making a single bond and giving each atom a complete valence shell, which fulfills the **octet** rule:



Two F's with two dots in between them, and two dots on the top, left/right side, and bottoms of each F.

The circles show that each fluorine atom has eight electrons around it. As with hydrogen, we can represent the fluorine molecule with a dash in place of the bonding electrons:



Two F's with a long dash in between them, and two dots on the top, left/right side, and bottoms of each F.


Each fluorine atom has six electrons, or three pairs of electrons, that are not participating in the covalent bond. Rather than being shared, they are considered to belong to a single atom. These are called nonbonding pairs (or lone pairs) of electrons.

Covalent Bonds between Different Atoms

Now that we have looked at electron sharing between atoms of the same element, let us look at covalent bond formation between atoms of different elements. Consider a molecule composed of one hydrogen atom and one fluorine atom:



Each atom needs one additional electron to complete its valence shell. By each contributing one electron, they make the following molecule:



In this molecule, the hydrogen atom does not have nonbonding electrons, while the fluorine atom has six nonbonding electrons (three lone electron pairs). The circles show how the valence electron shells are filled for both atoms.

Example 6.2.1


Use Lewis diagrams to indicate the formation of the following:

a. Cl_2
b. HBr


Solution

a. When two chlorine atoms form a chlorine molecule, they share one pair of electrons. In Cl_2 molecule, each chlorine atom is surrounded by an **octet** number of electrons.

The Lewis diagram for a Cl_2 molecule is similar to the one for F_2 (shown above).



b. When a hydrogen atom and a bromine atom form HBr, they share one pair of electrons. In the HBr molecule, H achieves a full valence of two electrons (**duet**) while Br achieves an **octet**. The Lewis diagram for HBr is similar to that for HF shown above.



Exercise 6.2.1


Draw the Lewis diagram for each compound.

a. a molecule composed of one chlorine atom and one fluorine atom
b. a molecule composed of one hydrogen atom and one iodine atom

Answer a: available at the end of this section.


Covalent Bonds in Larger Molecules

The formation of a water molecule from two hydrogen atoms and an oxygen atom can be illustrated using Lewis dot symbols (shown below).




H with one dot + O with 6 dots + H with one dot turns into H connected to O with two dots connected to H with two dots. O also has two dots on top and top dots on bottom.

The structure shown is the *Lewis electron structure*, or *Lewis structure*, for $\mathrm{H_2O}$. With two bonding pairs and two lone pairs, the oxygen atom has now completed its **octet**. Moreover, by sharing a bonding pair with oxygen, each hydrogen atom now has a full valence shell of two electrons. Chemists usually indicate a bonding pair by a single line, as shown (below).




Water

Other large molecules are constructed in a similar fashion, with some atoms participating in more than one covalent bond. For example, methane ($\mathrm{CH_4}$), the central carbon atom bonded to four hydrogen atoms, can be represented using either of the Lewis structures below. Again, sharing electrons between C and H atoms results in C achieving and octet while H achieving a duet number of electrons.



How Many Covalent Bonds Are Formed?

The number of bonds that an atom can form can often be predicted from the number of electrons needed to reach an octet (eight valence electrons). In the Lewis structure, the number of bonds formed by an element in a neutral compound is the same as the number of unpaired electrons it must share with other atoms to complete its octet of electrons. For example, each atom of a **group 4A** (14) element has four electrons in its outermost shell and therefore requires four more electrons to reach an octet. These four electrons can be gained by forming **four covalent bonds**, as illustrated here for carbon in $\mathrm{CH_4}$ (methane). **Group 5A** (15) elements such as nitrogen have five valence electrons in the atomic Lewis symbol: one lone pair and three unpaired electrons. To obtain an octet, these atoms form **three covalent bonds**, as in $\mathrm{NH_3}$ (ammonia). Oxygen and other atoms in **group 6A** (16) obtain an octet by forming **two covalent bonds**. Fluorine and the other halogens in **group 7A** (17) have seven valence electrons and can obtain an octet by forming **one covalent bond**.



Typically, the atoms of group 4A form 4 covalent bonds; group 5A form 3 bonds; group 6A form 2 bonds; and group 7A form one bond. The number of electrons required to obtain an octet determines the number of covalent bonds an atom can form. This is summarized in the table below. In each case, the sum of the number of bonds and the number of lone pairs is 4, which is equivalent to eight (octet) electrons.

This table shows atoms and their group numbers, and how many bonds and lone pairs each has.

Atom (Group number)	Number of Bonds	Number of Lone Pairs
Carbon (Group 14 or 4A)	4	0
Nitrogen (Group 15 or 5A)	3	1
Oxygen (Group 16 or 6A)	2	2
Fluorine (Group 17 or 7A)	1	3

Because hydrogen only needs two electrons to fill its valence shell, it follows the duet rule. It is an **exception to the octet rule.** Hydrogen only needs to form one bond. This is the reason why H is always a terminal atom and never a central atom. Figure 6.2.1 shows the number of covalent bonds various atoms typically form.

The transition elements and inner transition elements also do not follow the octet rule since they have d and f electrons involved in their valence shells.





Figure 6.2.1: How Many Covalent Bonds Are Formed? In molecules, there is a pattern to the number of covalent bonds that different atoms can form. Each block with a number indicates the number of covalent bonds formed by that atom in neutral compounds.

Example 6.2.2

Examine the Lewis structure of OF_2 below. Count the number of bonds formed by each element. Based on the element's location in the periodic table, does it correspond to the expected number of bonds shown in Table 4.1? Does the Lewis structure below follow the octet rule?




Solution

Yes. F (group 7A) forms one bond and O (group 6A) forms 2 bonds. Each atom is surrounded by 8 electrons. This structure satisfies the octet rule.

6.2.2Exerc

Examine the Lewis structure of NCl_3 below. Count the number of bonds formed by each element. Based on the element's location in the periodic table, does it correspond to the expected number of bonds shown in Table 4.1? Does the Lewis structure below follow the octet rule?



Answer available at the end of this section.


Key Takeaways

- A covalent bond is formed between two atoms by sharing electrons.
- The number of bonds an element forms in a covalent compound is determined by the number of electrons it needs to reach octet.
- Hydrogen is an exception to the octet rule. H forms only one bond because it needs only two electrons.

Answers

Exercise 6.2.1





Both Cl and N form the expected number of bonds. Cl (group 7A) has one bond and 3 lone pairs. The central atom N (group 5A) has 3 bonds and one lone pair. Yes, the Lewis structure of NCl_3 follows the octet rule.

This page titled 6.2: Covalent Bonds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.1: Covalent Bonds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.3: Covalent Compounds - Formulas and Names

Learning Objectives


- Identify covalent and ionic compounds.
- Determine the chemical formula of a simple covalent compound from its name.
- Determine the name of a simple covalent compound from its chemical formula.

COVALENT AND IONIC COMPOUNDS


What elements make covalent bonds? Covalent bonds form when two or more nonmetals combine. For example, both hydrogen and oxygen are nonmetals, and when they combine to make water, they do so by forming covalent bonds. Compounds that are composed of only non-metals or semi-metals with non-metals will display covalent bonding and will be classified as molecular compounds.

As a general rule of thumb, compounds that involve a metal binding with either a non-metal or a semi-metal will display ionic bonding. Thus, the compound formed from sodium and chlorine will be ionic (a metal and a non-metal). Nitrogen monoxide (NO) will be a covalently bound molecule (two non-metals), silicon dioxide (SiO_2) will be a covalently bound molecule (a semi-metal and a non-metal) and $MgCl_2$ will be ionic (a metal and a non-metal).

A polyatomic ion is an ion composed of two or more atoms that have a charge as a group (poly = many). The ammonium ion (see figure below) consists of one nitrogen atom and four hydrogen atoms. Together, they comprise a single ion with a 1+ charge and a formula of NH_4^+. The carbonate ion (see figure below) consists of one carbon atom and three oxygen atoms and carries an overall charge of 2–. The formula of the carbonate ion is CO_3^{2-}.



ammonium ion



carbonate ion

The atoms of a polyatomic ion are tightly bonded together and so the entire ion behaves as a single unit. Several examples are found in Table 3.3.1. Nonmetal atoms in polyatomic ions are joined by covalent bonds, but the ion as a whole participates in ionic bonding. For example, ammonium chloride (NH_4Cl) has ionic bonding between a polyatomic ion, NH_4^+, and Cl^- ions, but within the ammonium ion (NH_4^+), the nitrogen and hydrogen atoms are connected by covalent bonds (shown above).

Both ionic and covalent bonding are also found in calcium carbonate. Calcium carbonate ($CaCO_3$) has ionic bonding between calcium ion Ca^{2+} and a polyatomic ion, CO_3^{2-}, but within the carbonate ion (CO_3^{2-}), the carbon and oxygen atoms are connected by covalent bonds (shown above).

Characteristics of Covalent (Molecular) Compounds

Compounds that contain covalent bonds (also called molecular compounds) exhibit different physical properties than ionic

compounds. Because the attraction between molecules, which are electrically neutral, is weaker than that between electrically charged ions, covalent compounds generally have much lower melting and boiling points than ionic compounds (discussed in Section 3.6). For example, water (molecular compound) boils at 100 °C while sodium chloride (ionic compound) boils at 1413 °C. In fact, many covalent compounds are liquids or gases at room temperature, and, in their solid states, they are typically much softer than ionic solids. Furthermore, whereas ionic compounds are good conductors of electricity when dissolved in water, most covalent compounds, being electrically neutral, are poor conductors of electricity in any state. The attraction between molecules (called intermolecular forces) will be discussed in more detail in Section 8.1

Example 6.3.1

Is each compound formed from ionic bonds, covalent bonds, or both?

a. Na_2O
b. Na_3PO_4
c. N_2O_4

Answer a available at the end of this section.

Exercise 6.3.1


Is each compound are formed from ionic bonds, covalent bonds, or both?

a. $Ba(OH)_2$
b. F_2
c. PCl_3

Answer a: available at the end of this section.

MOLECULAR FORMULAS

The chemical formulas for covalent compounds are referred to as **molecular formulas** because these compounds exist as separate, discrete molecules. Typically, a molecular formula begins with the nonmetal that is closest to the lower left corner of the periodic table. Then the other nonmetal symbols are listed. Numerical subscripts are used if there is more than one of a particular atom. For example, we have already seen CH_4, the molecular formula for methane. Below is the molecular formula of ammonia, NH_3.



NAMING COVALENT COMPOUNDS

Naming *binary* (two-element) covalent compounds is similar to naming simple ionic compounds. The first element in the formula is simply listed using the name of the element. The second element is named by taking the stem of the element name and adding the suffix *-ide*. A system of numerical prefixes is used to specify the number of atoms in a molecule. Table 6.3.1 lists these numerical prefixes. Normally, no prefix is added to the first element's name if there is only one atom of the first element in a molecule. If the second element is oxygen, the trailing vowel is usually omitted from the end of a polysyllabic prefix but not a monosyllabic one (that is, we would say "monoxide" rather than "monooxide" and "trioxide" rather than "troxide").

Table 6.3.1: Numerical Prefixes for Naming Binary Covalent Compounds

Number of Atoms in Compound	Prefix on the Name of the Element
1	mono-*
2	di-
3	tri-
4	tetra-
5	penta-
6	hexa-
7	hepta-
8	octa-
9	nona-
10	deca-
***This prefix is not used for the first element's name.**	

Let us practice by naming the compound whose molecular formula is CCl_4. The name begins with the name of the first element—carbon. The second element, chlor*ine*, becomes chlor*ide*, and we attach the correct numerical prefix ("tetra-") to indicate that the molecule contains four chlorine atoms. Putting these pieces together gives the name *carbon tetrachloride* for this compound.

Example 6.3.2

Write the molecular formula for each compound.

a. chlorine trifluoride
b. phosphorus pentachloride
c. sulfur dioxide
d. dinitrogen pentoxide

Solution

If there is no numerical prefix on the first element's name, we can assume that there is only one atom of that element in a molecule.

a. ClF_3
b. PCl_5
c. SO_2
d. N_2O_5 (The *di-* prefix on nitrogen indicates that two nitrogen atoms are present.)

Exercise 6.3.2

Write the molecular formula for each compound.

a. nitrogen dioxide
b. dioxygen difluoride
c. sulfur hexafluoride
d. selenium monoxide

Answer a: available at the end of this section.

Because it is so unreactive, sulfur hexafluoride is used as a spark suppressant in electrical devices such as transformers.

Example 6.3.3

Write the name for each compound.

a. BrF_5
b. S_2F_2
c. CO

Solution

a. bromine pentafluoride
b. disulfur difluoride
c. carbon monoxide

Exercise 6.3.3

Write the name for each compound.

a. CF_4
b. $SeCl_2$
c. SO_3

Answer a: available at the end of this section.

For some simple covalent compounds, we use common names rather than systematic names. We have already encountered these compounds, but we list them here explicitly:

- H_2O: water
- NH_3: ammonia
- CH_4: methane

Methane is the simplest organic compound. Organic compounds are compounds with carbon atoms and are named by a separate nomenclature system that we will introduce in in a separate section.

Key Takeaways

- The chemical formula of a simple covalent compound can be determined from its name.
- The name of a simple covalent compound can be determined from its chemical formula.

Answers

Example 6.3.1

The elements in Na_2O are a metal and a nonmetal, which form ionic bonds.

Because sodium is a metal and we recognize the formula for the phosphate ion, we know that this compound is ionic. However, within the polyatomic phosphate ion, the atoms are held together by covalent bonds, so this compound contains both ionic and covalent bonds.

The elements in $N_2O_4|$ are both nonmetals, rather than a metal and a nonmetal. Therefore, the atoms form covalent bonds.

Exercise 6.3.1

both

covalent

covalent

Exercise 6.3.2

a. NO_2

O_2F_2

SF_6

SeO

Exercise 6.3.3

carbon tetrafluoride

selenium dichloride

sulfur trioxide

This page titled 6.3: Covalent Compounds - Formulas and Names is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.2: Covalent Compounds - Formulas and Names by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.


6.4: Drawing Lewis Structures

Learning Objectives

- To draw Lewis structures.
- To recognize molecules that are likely to have multiple covalent bonds.

DRAWING LEWIS STRUCTURES

For very simple molecules and molecular ions, we can write the Lewis structures by merely pairing up the unpaired electrons on the constituent atoms. See these examples:



For more complicated molecules and molecular ions, it is helpful to follow the step-by-step procedure outlined here:

1. Determine the total number of valence (outer shell) electrons among all the atoms. For cations, subtract one electron for each positive charge. For anions, add one electron for each negative charge.
2. Draw a skeleton structure of the molecule or ion, arranging the atoms around a central atom. (Generally, the least electronegative element should be placed in the center.) Connect each atom to the central atom with a single bond (one electron pair).
3. Distribute the remaining electrons as lone pairs on the terminal atoms (except hydrogen), completing an octet around each atom.
4. Place all remaining electrons on the central atom.
5. Rearrange the electrons of the outer atoms to make multiple bonds with the central atom in order to obtain octets wherever possible.

Let us determine the Lewis structures of OF_2 and HCN as examples in following this procedure:

1. Determine the total number of valence (outer shell) electrons in the molecule or ion. For a molecule, we add the number of valence electrons (use the main group number) on each atom in the molecule. This is the total number of electrons that must be used in the Lewis structure.

$$O + 2\,(F) = OF_2$$

$$6e^- + (2 \times 7e^-) = 20e^-$$

$$H + C + N = HCN$$


$1e^- + 4e^- + 5e^- = 10e^-$

2. Draw a skeleton structure of the molecule or ion, arranging the atoms around a central atom and connecting each atom to the central atom with a single (one electron pair) bond. Note that H and F can only form one bond, and are always on the periphery rather than the central atom.




3. Distribute the remaining electrons as lone pairs on the terminal atoms (except hydrogen) to complete their valence shells with an octet of electrons.

- In OF_2, six electrons are placed on each F.
- In HCN, six electrons placed on N




4. Place all remaining electrons on the central atom.


- In OF_2, 4 electrons are placed on O.
- In HCN: no electrons remain (the total valence of $10e^-$ is reached) so nothing changes.



5. Rearrange the electrons of the outer atoms to make multiple bonds with the central atom in order to obtain octets wherever possible.

- In OF_2, each atom has an octet as drawn, so nothing changes.
- In HCN, form two more C–N bonds






Finally, check to see if the total number of valence electrons are present in the Lewis structure. And then, inspect if the H atom has 2 electrons surrounding it and if each of the main group atoms is surrounded by 8 electrons.

MULTIPLE BONDS


In many molecules, the octet rule would not be satisfied if each pair of bonded atoms shares only two electrons. Review HCN in Step 5 above. Another example is carbon dioxide (CO_2). CO_2 has a total valence of $4e^- + (2 \times 6e^-) = 16e^-$. Following steps 1 to 4, we draw the following:




This does not give the carbon atom a complete octet; only four electrons are in its valence shell. This arrangement of shared electrons is far from satisfactory.




In this case, more than one pair of electrons must be shared between two atoms for both atoms to have an octet. A second electron pair from each oxygen atom must be shared with the central carbon atom shown by the arrows above. A lone pair from each O must be converted into a bonding pair of electrons.



In this arrangement, the carbon atom shares four electrons (two pairs) with the oxygen atom on the left and four electrons with the oxygen atom on the right. There are now eight electrons around each atom. Two pairs of electrons shared between two atoms make a double bond between the atoms, which is represented by a double dash:



Some molecules contain triple bonds (like HCN, shown above). Triple bonds are covalent bonds in which *three* pairs of electrons are shared by two atoms. Another compound that has a triple bond is acetylene (C_2H_2), whose Lewis diagram is as follows:



Example 6.4.1

Draw the Lewis diagram for each molecule.


1. N_2
2. CH_2O (The carbon atom is the central atom.) One application of CH_2O, also called formaldehyde, is the preservation of biological specimens. Aqueous solutions of CH_2O are called formalin and have a sharp, characteristic (pungent) odor.

Solution

1. The total number of electrons is 2 x 5 = 10 electrons. The bond between the two nitrogen atoms is a triple bond. The Lewis diagram for $N_2$is as follows:



2. The total number of electrons is 4 x 2(1) + 6 = 12 electrons. In CH_2O, the central atom is surrounded by two different types of atoms. The Lewis diagram that fills each atom's valence electron shell is as follows:



Exercise 6.4.1


Draw the Lewis diagram for each molecule.


a. O_2
b. C_2H_4


Answer a: available at the end of this section.

Example 6.4.2

Which is the correct Lewis structure for N_2H_2?

A. 

B. 

C. 

Solution


Lewis structure A is the correct answer. It has a total of (2 x $5e^-$) + (2 x $1e^-$) = $12e^-$. Each of the N atoms satisfy the octet requirement and the H atoms follow the duet rule.


Structure B is electron deficient. It has only $10e^-$ instead of 12.


Structure C has 14 (2 extra) electrons. The N atoms do not satisfy the octet.

6.4.2Exerc

Which is the correct Lewis structure for NOCl?

A. 

B. 

C. 


Answer available at the end of this section.

Key Takeaways

- A Lewis structure shows the bonding and nonbonding electrons around individual atoms in a molecule.
- Some molecules must have multiple covalent bonds between atoms to satisfy the octet rule.
- A double bond contains four electrons and a triple bond contains six electrons.

Answers

Exercise 6.4.1



Structure A violates the octet rule; N is surrounded by only $6e^-$.

Structure B violates the octet rule; Cl has $10e^-$ around it. Furthermore, there are a total of $20e^-$ instead of $18e^-$.

Structure C is the correct structure. It has a total of $6e^-$ + $5e^-$ + $7e^-$ = $18e^-$. Each atom is surrounded by 8 electrons (octet rule).

This page titled 6.4: Drawing Lewis Structures is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.3: Drawing Lewis Structures by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.5: Characteristics of Covalent Bonds

Learning Objectives

- Recognize bond characteristics of covalent compounds: bond length and bond polarity.
- Use electronegativity values to predict bond polarity.

Bond Length

We previously stated that the covalent bond in the hydrogen molecule (H_2) has a certain length (about 7.4×10^{-11} m). Other covalent bonds also have known bond lengths, which are dependent on both the identities of the atoms in the bond and whether the bonds are single, double, or triple bonds. Table 6.5.1 lists the approximate bond lengths for some single covalent bonds. The exact bond length may vary depending on the identity of the molecule but will be close to the value given in the table.

Table 6.5.1: Approximate Bond Lengths of Some Single Bonds

Bond	Length ($\times 10^{-12}$ m)
H–H	74
H–C	110
H–N	100
H–O	97
H–I	161
C–C	154
C–N	147
C–O	143
N–N	145
O–O	145

Table 6.5.2 compares the lengths of single covalent bonds with those of double and triple bonds between the same atoms. Without exception, as the number of covalent bonds between two atoms increases, the bond length decreases. With more electrons between the two nuclei, the nuclei can get closer together before the internuclear repulsion is strong enough to balance the attraction.

Table 6.5.2: Comparison of Bond Lengths for Single and Multiple Bonds

Bond	Length ($\times 10^{-12}$ m)
C–C	154
C=C	134
C≡C	120
C–N	147
C=N	128
C≡N	116
C–O	143
C=O	120

Bond	Length ($\times 10^{-12}\ m$)
C≡O	113
N–N	145
N=N	123
N≡N	110
O–O	145
O=O	121

Electronegativity and Bond Polarity

Although we defined covalent bonding as electron sharing, the electrons in a covalent bond are not always shared equally by the two bonded atoms. Unless the bond connects two atoms of the same element, as in H_2, there will always be one atom that attracts the electrons in the bond more strongly than the other atom does, as in HCl, shown in Figure $6.5.1$. A covalent bond that has an equal sharing of electrons (Figure $6.5.1a$) is called a **nonpolar covalent bond**. A covalent bond that has an unequal sharing of electrons, as in Figure $6.5.1b$, is called a **polar covalent bond**.




Figure $6.5.1$ Polar versus Nonpolar Covalent Bonds. (a) The electrons in the covalent bond are equally shared by both hydrogen atoms. This is a nonpolar covalent bond. (b) The chlorine atom attracts the electrons in the bond more than the hydrogen atom does, leading to an imbalance in the electron distribution. This is a polar covalent bond.

The distribution of electron density in a polar bond is uneven. It is greater around the atom that attracts the electrons more than the other. For example, the electrons in the H–Cl bond of a hydrogen chloride molecule spend more time near the chlorine atom than near the hydrogen atom. Note that the shaded area around Cl in Figure $6.5.1b$ is much larger than it is around H.

This imbalance in electron density results in a buildup of partial negative charge (designated as δ–) on one side of the bond (Cl) and a partial positive charge (designated δ+) on the other side of the bond (H). This is seen in Figure $6.5.2a$. The separation of charge in a polar covalent bond results in an electric dipole (two poles), represented by the arrow in Figure $6.5.2b$. The direction of the arrow is pointed toward the δ– end while the + tail of the arrow indicates the δ+ end of the bond.




Figure 6.5.2: (a) Unequal sharing of the bonding pair of electrons between H and Cl leads to partial positive charge on the H atom and partial negative charge on the Cl. Symbols δ+ and δ– indicate the polarity of the H–Cl bond. (b) The dipole is represented by an arrow with a cross at the tail. The cross is near the δ+ end and the arrowhead coincides with the δ–.

Any covalent bond between atoms of different elements is a polar bond, but the degree of polarity varies widely. Some bonds between different elements are only minimally polar, while others are strongly polar. Ionic bonds can be considered the ultimate in polarity, with electrons being transferred rather than shared. To judge the relative polarity of a covalent bond, chemists use **electronegativity**, which is a relative measure of how strongly an atom attracts electrons when it forms a covalent bond. There are various numerical scales for rating electronegativity. Figure 6.5.3 shows one of the most popular—the Pauling scale.





Figure 6.5.3 The electronegativity values derived by Pauling follow predictable periodic trends with the higher electronegativities toward the upper right of the periodic table. Fluorine has the highest value (4.0).

Looking Closer: Linus Pauling

Arguably the most influential chemist of the 20th century, Linus Pauling (1901–94) is the only person to have won two individual (that is, unshared) Nobel Prizes. In the 1930s, Pauling used new mathematical theories to enunciate some fundamental principles of the chemical bond. His 1939 book *The Nature of the Chemical Bond* is one of the most significant books ever published in chemistry.


By 1935, Pauling's interest turned to biological molecules, and he was awarded the 1954 Nobel Prize in Chemistry for his work on protein structure. (He was very close to discovering the double helix structure of DNA when James Watson and James Crick announced their own discovery of its structure in 1953.) He was later awarded the 1962 Nobel Peace Prize for his efforts to ban the testing of nuclear weapons.



Linus Pauling was one of the most influential chemists of the 20th century.

In his later years, Pauling became convinced that large doses of vitamin C would prevent disease, including the common cold. Most clinical research failed to show a connection, but Pauling continued to take large doses daily. He died in 1994, having spent a lifetime establishing a scientific legacy that few will ever equal.

The polarity of a covalent bond can be judged by determining the difference in the electronegativities of the two atoms making the bond. The greater the difference in electronegativities, the greater the imbalance of electron sharing in the bond. Although there are no hard and fast rules, the general rule is if the difference in electronegativities is **less than about 0.4,** the bond is considered **nonpolar;** if the difference is **greater than 0.4,** the bond is considered **polar.** If the difference in electronegativities is large enough **(**generally **greater than about 1.8),** the resulting compound is considered **ionic** rather than covalent**.** An electronegativity difference of **zero,** of course, indicates a **nonpolar covalent bond.**



Bond Type	Electronegativity Difference
pure covalent	< 0.4
polar covalent	between 0.4 and 1.8
ionic	> 1.8

Figure 6.5.4: As the electronegativity difference increases between two atoms, the bond becomes more ionic.

Example 6.5.1

Describe the electronegativity difference between each pair of atoms and the resulting polarity (or bond type).

a. C and H
b. H and H

c. Na and Cl
d. O and H

Solution

a. Carbon has an electronegativity of 2.5, while the value for hydrogen is 2.1. The difference is 0.4, which is rather small. The C–H bond is therefore considered nonpolar.
b. Both hydrogen atoms have the same electronegativity value—2.1. The difference is zero, so the bond is nonpolar.
c. Sodium's electronegativity is 0.9, while chlorine's is 3.0. The difference is 2.1, which is rather high, and so sodium and chlorine form an ionic compound.
d. With 2.1 for hydrogen and 3.5 for oxygen, the electronegativity difference is 1.4. We would expect a very polar bond. The sharing of electrons between O and H is unequal with the electrons more strongly drawn towards O.

Exercise 6.5.1

Describe the electronegativity (EN) difference between each pair of atoms and the resulting polarity (or bond type).

a. C and O
b. K and Br
c. N and N
d. Cs and F

Answer a: available at the end of this section.

Bond Polarity and Molecular Polarity

If there is only one bond in the molecule, the bond polarity determines the molecular polarity. Any diatomic molecule in which the two atoms are the same element must be a **nonpolar** molecule. A diatomic molecule that consists of a polar covalent bond, such as HF, is a **polar molecule**. A **polar molecule** *is a molecule in which one end of the molecule is slightly positive, while the other end is slightly negative*. The two electrically charged regions on either end of the molecule are called poles, similar to a magnet having a north and a south pole. Hence, a *molecule with two poles is called a* **dipole**. A simplified way to depict polar molecules like HF is pictured below (see figure below).




Figure 6.5.5: A molecular dipole results from the unequal distribution of electron density throughout a molecule.

When placed between oppositely charged plates, polar molecules orient themselves so that their positive ends are closer to the negative plate and their negative ends are closer to the positive plate (see Figure 4.4.6 below).




Figure 6.5.6: Polar molecules are randomly oriented in the absence of an applied electric field (top). In an electric field, the molecules orient themselves to maximize the attraction between opposite charges (bottom).

Experimental techniques involving electric fields can be used to determine if a certain substance is composed of polar molecules and to measure the degree of polarity.

For molecules with more than two atoms, the **molecular geometry** must also be taken into account when determining if the molecule is polar or nonpolar. (Figure 6.5.7) is a comparison between carbon dioxide and water. Carbon dioxide (CO_2) is a **linear** molecule. The oxygen atoms are more electronegative than the carbon atom, so there are two individual dipoles pointing outward from the C atom to each O atom. However, since the dipoles are of equal strength and are oriented directly opposite each other, they cancel each other out, and the overall molecular polarity of CO_2 is zero. CO_2 is a **nonpolar** molecule. In H_2O, the orientation of the two O–H bonds is **bent**: one end of the molecule has a partial positive charge, and the other end has a partial negative charge. In short, the molecule itself is **polar**. The polarity of water has an enormous impact on its physical and chemical properties. For example, the boiling point of water (100°C) is high for such a small molecule due to the fact that polar molecules attract each other strongly. On the other hand, the nonpolar carbon dioxide becomes a gas at –77°C, almost 200° lower than the temperature at which water boils. More details will be presented in the next section.



Figure 6.5.7 Physical Properties and Polarity. The physical properties of water and carbon dioxide are affected by their polarities.

Key Takeaways

- Covalent bonds between different atoms have different bond lengths.
- Covalent bonds can be polar or nonpolar, depending on the electronegativity difference between the atoms involved.

Answers

Exercise 6.5.1

The EN difference is 1.0 , hence polar. The sharing of electrons between C and O is unequal with the electrons more strongly drawn towards O.

The EN difference is greater than 1.8, hence ionic.

Identical atoms have zero EN difference, hence nonpolar.
The EN difference is greater than 1.8, hence ionic.

This page titled 6.5: Characteristics of Covalent Bonds is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.4: Characteristics of Covalent Bonds by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.6: Characteristics of Molecules and VSEPR

Learning Objectives

- Determine the molecular mass of a molecule.
- Predict the general shape of a simple covalent molecule.
- Predict the polarity of the molecule.
- Compare the properties of ionic and molecular compounds.

Unlike the ions in ionic compounds, which are arranged in lattices called crystals, molecules of covalent compounds exist as discrete units with a characteristic mass and a certain three-dimensional shape.

Molecular Mass

The mass of a molecule—the molecular mass (sometimes called the molecular weight)—is simply the sum of the masses of its atoms. As with formula masses, it is important that you keep track of the number of atoms of each element in the molecular formula to obtain the correct molecular mass.

Example 6.6.1

What is the molecular mass of each covalent compound?

a. H_2O
b. C_6H_6
c. NO_2
d. N_2O_5

Solution

Use the atomic masses from the Periodic Table in Section 2.7.

The molecular formula H_2O indicates that there are two hydrogen atoms and one oxygen atom in each molecule. Summing the masses of these atoms,

This table finds the mass of atoms.

2 H:	2 × 1.01 =	2.02 amu
1 O:		+ 16.00 amu
Total:		18.02 amu

The molecular mass of H_2O is **18.02** amu.

This table finds the mass of atoms.

6 C:	6 × 12.01 =	72.06 amu
6 H:	6 × 1.01 =	+ 6.06 amu
Total:		78.12 amu

The molecular mass of C_6H_6 is **78.12** amu.

This table finds the mass of atoms.

1 N:		14.01 amu
2 O:	2 × 16.00 =	+ 32.00 amu
Total:		46.01 amu

The molecular mass of NO_2 is **46.01** amu.

This table finds the mass of atoms.

2 N:	2 × 14.01 =	28.02 amu
5 O:	5 × 16.00 =	+ 80.00 amu
Total:		108.02 amu

The molecular mass of N_2O_5 is **108.02** amu.

Note that the two different nitrogen and oxygen compounds in these examples have different molecular masses.

Exercise 6.6.1


What is the molecular mass of each covalent compound?

a. C_2H_2
b. CO
c. CO_2
d. BF_3


Answer a: available at the end of this section.

Molecular Shape: VSEPR Theory

Unlike ionic compounds, with their extended crystal lattices, covalent molecules are discrete units with specific three-dimensional shapes. The shape of a molecule is determined by the fact that covalent bonds, which are composed of negatively charged electrons, tend to repel one another. This concept is called the **valence shell electron pair repulsion (VSEPR) theory**. For example, the two covalent bonds in $BeCl_2$ **stay as far from each other** as possible, ending up 180° apart from each other. The result is a ***linear*** molecule:



The three covalent bonds in BF_3 repel each other to form 120° angles in a plane, in a shape called ***trigonal planar***:



The molecules $BeCl_2$ and BF_3 actually violate the octet rule; however, such exceptions are rare and will not be discussed in this text.


> Try sticking three toothpicks into a marshmallow or a gumdrop and see if you can find different positions where your "bonds" are farther apart than the planar 120° orientation.

The four covalent bonds in CCl_4 arrange themselves three dimensionally, pointing toward the corner of a tetrahedron and making bond angles of 109.5°. CCl_4 is said to have a ***tetrahedral*** shape:

Table 6.6.1: Describes how to use the number of atoms around the central atom to determine the geometry of a molecule, as well as examples.


Atoms Around Central Atom	Geometry	Example
2 AB_2	Linear	$BeCl_2$
3 AB_3	Trigonal Planar	BF_3
4 AB_4	Tetrahedral	CCl_4

In determining the shapes of molecules, it is useful to first determine the Lewis diagram for a molecule. The shapes of molecules with multiple bonds are determined by treating the multiple bonds as one bond. Thus, formaldehyde (CH_2O) has a shape similar to that of BF_3. It is ***trigonal planar***.




Molecules With Lone Pairs Around Central Atom

Molecules with lone electron pairs around the central atom have a shape based on the position of the atoms, not the electron pairs. For example, NH_3 has **one lone electron pair** and **three bonded electron pairs**. These four electron pairs repel each other and adopt a tetrahedral arrangement. However, the shape of the molecule is described in terms of the positions of the atoms, not the lone electron pairs. Thus, NH_3 is said to have a ***trigonal pyramidal*** shape, not a tetrahedral one.



Similarly, H_2O has **two lone pairs** of electrons around the central oxygen atom and **two bonded electron pairs**. Although the four electron pairs adopt a tetrahedral arrangement, the shape of the molecule is described by the positions of the atoms only. The shape of H_2O is ***bent*** with an approximate 109.5° angle.



In summary, to determine the molecular geometry:

Step 1: Draw the Lewis structure.

Step 2: Count the number of bonds (a double/triple bond counts as one) and lone pairs around the **central atom**.


Step 3: Use Figure 6.6.1 to determine the molecular geometry.

Number of bonds and lone pairs	Number of bonds 0 lone pair	1 lone pair	2 lone pairs
2	180° X–A–X Linear		
3	120° Trigonal planar	<120° Bent or angular	Lone Pair
4	109° Tetrahedral	<109° Trigonal pyramid	<<109° Bent or angular

Figure 6.6.1: The molecular geometry depends on the number of bonds and lone pairs around the **central atom, A.**


Example 6.6.2

What is the geometry of the ammonium ion, NH_4^+? Its Lewis structure is shown below. How is this different from ammonia, NH_3?




Solution

In ammonium ion, the central atom N has 4 bonds and no lone pair. It is equivalent to the below in Table 4.5.1. Hence, this is *tetrahedral*.



In ammonia (NH_3), shown below, N has 3 bonds and one lone pair.




It is equivalent to the below in Table 4.5.1. Hence, the shape of this molecule is *trigonal pyramid*.



Exercise 6.6.2

What is the molecular shape of nitrosyl chloride, a highly corrosive, reddish-orange gas? Its Lewis structure is shown below.



Answer available at the end of this section.

Molecular Polarity

In general, a molecule is nonpolar if all its bonds are nonpolar. Examples are I_2, O_2, H_2, CH_4, C_2H_6 and C_3H_8.

In general, a molecule is polar if it contains polar bonds EXCEPT when the bond polarities cancel each other. As mentioned in Section 4.4, the shape of the CO_2 molecule (linear) orients the two C=O polar bonds **directly opposite** each other, thus **cancelling** each other's effect. **Carbon dioxide** (CO_2) is a ***nonpolar*** molecule.

On the other hand, water (as discussed above) is a bent molecule because of the two lone pairs on the central oxygen atom. Because of the bent shape, the dipoles do not cancel each other out and the water molecule is ***polar***. In the figure below, the individual H-O polar bonds represented by the two red arrows are not directly opposite each other. These two dipoles don't cancel each other out. In fact, the net dipole (blue arrow) points upward. There is a resultant partial positive charge at one end (between the two H atoms) and a partial negative charge on the other end (where O is located). The uneven distribution of charge or the overall dipole is shown by the blue arrow below (Figure 6.6.2). Hence, **water is *polar*** (has + and - poles) while **carbon dioxide is *nonpolar***.




Figure 6.6.2: The molecular geometry of a molecule (linear vs. bent) affects its polarity.

Similarly, in BF_3 (trigonal planar), the effect of a B-F bond is cancelled by the sum of the other two B-F bonds (see video). Hence, a trigonal planar molecule (BF_3) is nonpolar because the bond polarities cancel each other, but a trigonal pyramidal molecule (NH_3) is polar.




Figure 6.6.3: The molecular geometry of a molecule (trigonal planar vs. trigonal pyramid) affects its polarity.

Some other molecules are shown in the figure below. Notice that a tetrahedral molecule such as CCl_4 is ***nonpolar***. However, if the peripheral atoms are not of the same electronegativity, the bond polarities don't cancel and the molecule becomes ***polar***, as in CH_3Cl.




Figure 6.6.4: *The same molecular geometry but peripheral bonds are of different electronegativity. CCl_4 is nonpolar but CH_3Cl is polar.*




Figure 6.6.5: Molecules with Polar Bonds. Individual bond dipole moments are indicated in red. Due to their different three-dimensional geometry, some molecules with polar bonds have a net dipole moment (HCl, CH_2O, NH_3, and $CHCl_3$), indicated in blue, whereas others do not because the bond dipoles cancel due to symmetry (BCl_3 and CCl_4).

Physical Properties of Molecular Compounds

The physical state and properties of a particular compound depend in large part on the type of chemical bonding it displays. **Molecular compounds**, sometimes called covalent compounds, display a wide range of physical properties due to the different types of intermolecular attractions such as different kinds of polar interactions. The melting and boiling points of molecular compounds are generally quite low compared to those of **ionic compounds**. This is because the energy required to disrupt the **intermolecular forces** (discussed in Section 8.1) between molecules is far less than the energy required to break the ionic bonds in a crystalline ionic compound. Since molecular compounds are composed of neutral molecules, their electrical conductivity is generally quite poor, whether in the solid or liquid state. Ionic compounds do not conduct electricity in the solid state because of their rigid structure, but conduct well when either molten or dissolved into a solution. The water solubility of molecular compounds is variable and depends primarily on the type of intermolecular forces involved. Substances that exhibit hydrogen bonding or dipole-dipole forces are generally water soluble, whereas those that exhibit only London dispersion forces are generally insoluble. Most, but not all, ionic compounds are quite soluble in water. The table below summarizes some of the differences between ionic and molecular compounds.

Table 6.6.2: Comparison of Ionic and Molecular Compounds

Property	Ionic Compounds	Molecular Compounds
Type of elements	Metal and nonmetal	Nonmetals only
Bonding	Ionic - transfer of electron(s) between atoms	Covalent - sharing of pair(s) of electrons between atoms
Representative unit	Formula unit	Molecule
Physical state at room temperature	Solid	Gas, liquid, or solid
Water solubility	Usually high	Variable

Property	Ionic Compounds	Molecular Compounds
Melting and boiling temperatures	Generally high	Generally low
Electrical conductivity	Good when molten or in solution	Poor

In summary, covalent compounds are softer, have lower boiling and melting points, are more flammable, are less soluble in water and do not conduct electricity compared to ionic compounds. The individual melting and boiling points, solubility and other physical properties of molecular compounds depend on molecular polarity.


Example 6.6.3

Describe the shape of each molecule. Is it polar or nonpolar?

a. PCl_3
b. CO_2


Solution

a. The Lewis diagram for PCl_3 is as follows:



Focus on the central atom, P that has 3 bonds and one lone pair. The four electron pairs arrange themselves tetrahedrally, but the lone electron pair is not considered in describing the molecular shape. Like NH_3, this molecule is *pyramidal*. The 3 P-Cl bonds don't cancel each other. This is *polar*.

b. The Lewis diagram for CO_2 is as follows:



Focus on the central atom, C. The multiple bonds are treated as one group, hence C has 2 bonds and zero lone pair. CO_2 has only two groups of electrons that repel each other. They will direct themselves 180° apart from each other, so CO_2 molecules are *linear*. This is highly symmetrical, with the two opposite dipoles cancelling each other. The CO_2 molecule is *nonpolar*.

Exercise 6.6.3

Describe the shape of each molecule. Is it polar or nonpolar?

a. CBr_4
b. BCl_3

Answer a: available at the end of this section.

Key Takeaways

- A molecule has a certain mass, called the molecular mass.
- Simple molecules have geometries that can be determined from VSEPR theory.
- Polar molecules result from differences in electronegativity of the atoms in the molecule.
- Dipoles that are directly opposite one another cancel each other out.

Answers

Exercise 6.6.1

26.04 amu


28.01 amu

44.01 amu


67.81 amu

Exercise 6.6.2


Focus on the central atom, N. It has a double bond to O, count this as one bond. It also has a single bond to Cl. Thus, N has 2 bonds and one lone pair. These 3 electron pairs will spread out 120 degrees from each other. But, since the shape is defined by the arrangement of the atoms only, the shape is bent or angular. If you consult Table 4.5.1, this molecule is equivalent to the below. Hence, two bonds and one lone pair has a *bent or angular* shape.




Exercise 6.6.3



or



The Lewis structure shows 4 groups attached to the central atom, hence *tetrahedral*. All the 4 groups are identical and the shape is symmetrical. Hence, it is *nonpolar*.



The Lewis diagram shows 3 groups attached to the central atom, hence *trigonal planar*. All the 3 groups are identical and shape is symmetrical, hence, it is *nonpolar*.

This page titled 6.6: Characteristics of Molecules and VSEPR is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.5: Characteristics of Molecules by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.7: Organic Chemistry

Learning Objectives

- Define *organic chemistry*.
- Identify organic molecules as alkanes, alkenes, alkynes, alcohols, or carboxylic acids.

When methane was mentioned previously, we described it as the simplest organic compound. In this section, we introduce organic chemistry more formally. Organic chemistry is the study of the chemistry of carbon compounds. Carbon is singled out because it has a chemical diversity unrivaled by any other chemical element. Its diversity is based on the following:

- Carbon atoms bond reasonably strongly with other carbon atoms.
- Carbon atoms bond reasonably strongly with atoms of other elements.
- Carbon atoms make a large number of covalent bonds (four).

Curiously, elemental carbon is not particularly abundant. It does not even appear in the list of the most common elements in Earth's crust. Nevertheless, all living things consist of organic compounds. Most organic chemicals are covalent compounds, which is why we introduce organic chemistry here. By convention, compounds containing carbonate ions and bicarbonate ions, as well as carbon dioxide and carbon monoxide, are not considered part of organic chemistry, even though they contain carbon.

Alkanes

The simplest organic compounds are the **hydrocarbons**, compounds composed of carbon and hydrogen atoms only. Some hydrocarbons have only single bonds and appear as a chain (which can be a straight chain or can have branches) of carbon atoms also bonded to hydrogen atoms. These hydrocarbons are called **alkanes (saturated hydrocarbons**). Each alkane has a characteristic, systematic name depending on the number of carbon atoms in the molecule. These names consist of a stem that indicates the number of carbon atoms in the chain plus the ending -ane. The stem meth- means one carbon atom, so methane is an alkane with one carbon atom. Similarly, the stem eth- means two carbon atoms; ethane is an alkane with two carbon atoms. Continuing, the stem prop- means three carbon atoms, so propane is an alkane with three carbon atoms. The stem but- means four carbon atoms; butane is an alkane with four carbon atoms. Figure 6.7.1 gives the Lewis structures, condensed structural formulas and molecular formulas of the four simplest alkanes. In the **condensed structural formula**, the covalent bonds are understood to exist between each carbon and the hydrogens associated with it, as well as between carbon atoms.




Figure 6.7.1: Formulas and Molecular Models of the Four Simplest Alkanes. The four smallest alkanes are methane, ethane, propane and butane.

Alkenes

Some hydrocarbons have one or more carbon–carbon double bonds (denoted C=C). These hydrocarbons are called alkenes. Figure 6.7.2 shows the formulas and the molecular models of the two simplest alkenes. Note that the names of alkenes have the

same stem as the alkane with the same number of carbon atoms in its chain but have the ending *-ene*. Thus, ethene is an alkene with two carbon atoms per molecule, and propene is a compound with three carbon atoms and one double bond.





Figure 6.7.2: Formulas and Molecular Models of the Two Simplest Alkenes. Ethene is commonly called ethylene, while propene is commonly called propylene.

Alkynes

Alkynes are hydrocarbons with a carbon–carbon triple bond (denoted C≡C) as part of their carbon skeleton. Below is the formula and the molecular model of the simplest alkyne and its systematic name. Its common name is acetylene. Its chemical formula is **C_2H_2**.



Ethyne

The names for alkynes have the same stems as for alkanes but with the ending *-yne*.

To Your Health: Saturated and Unsaturated Fats

Hydrocarbons are not the only compounds that can have carbon–carbon double bonds. A group of compounds called fats can have them as well, and their presence or absence in the human diet is becoming increasingly correlated with health issues.

Fats are combinations of long-chain organic compounds (fatty acids) and glycerol ($C_3H_8O_3$). (For more information on fats, see Chapter 17) The long carbon chains can have either all single bonds, in which case the fat is classified as *saturated*, or one or more double bonds, in which case it is a *monounsaturated* or a *polyunsaturated* fat, respectively. Saturated fats are typically solids at room temperature; beef fat (tallow) is one example. Mono- or polyunsaturated fats are likely to be liquids at room temperature and are often called oils. Olive oil, flaxseed oil, and many fish oils are mono- or polyunsaturated fats.

Studies have linked higher amounts of saturated fats in people's diets with a greater likelihood of developing heart disease, high cholesterol, and other diet-related diseases. In contrast, increases in unsaturated fats (either mono- or polyunsaturated) have been linked to a lower incidence of certain diseases. Thus, there have been an increasing number of recommendations by government bodies and health associations to decrease the proportion of saturated fat and increase the proportion of unsaturated fat in the diet. Most of these organizations also recommend decreasing the total amount of fat in the diet.

Recently, certain fats called trans fats have been implicated in the presence of heart disease. These are fats from animal sources and are also produced when liquid oils are exposed to partial hydrogenation, an industrial process that increases their saturation. Trans fats are used in many prepared and fried foods. Because they bring with them the health risks that naturally occurring saturated fats do, there has been some effort to better quantify the presence of trans fats in food products. US law now requires that food labels list the amount of trans fat in each serving.

Since the early 1900's, the US Department of Agriculture has been providing science-based dietary guidelines for the public. The most current version is the MyPlate illustration that gives a simple, visual picture of how much of what kind of foods make up a good, balanced diet. It recommends minimizing daily intake of sugars, the "bad fats", trans and saturated fat, and sodium. "Good fats", unsaturated fats or oils, are not considered a food group but do contain essential nutrients and therefore are included as part of a healthy eating pattern. The difference as simple as the difference between a single and double carbon–carbon bond, good and bad fats, can have a significant impact on health.




Figure 6.7.3: MyPlate is a reminder that everything you eat and drink matters. Source: Image courtesy of the USDA, www.mypyramid.gov/downloads/MiniPoster.pdf.

Functional Groups

The carbon–carbon double and triple bonds are examples of functional groups in organic chemistry. A functional group is a specific structural arrangement of atoms or bonds that imparts a characteristic chemical reactivity to a molecule. Alkanes have no functional group. A carbon–carbon double bond is considered a functional group because carbon–carbon double bonds chemically react in specific ways that differ from reactions of alkanes (for example, under certain circumstances, alkenes react with water); a carbon–carbon triple bond also undergoes certain specific chemical reactions. In the remainder of this section, we introduce two other common functional groups.

If an OH group (also called a hydroxyl group) is substituted for a hydrogen atom in a hydrocarbon molecule, the compound is an alcohol. Alcohols are named using the parent hydrocarbon name but with the final *-e* dropped and the suffix *-ol* attached. The two simplest alcohols are methanol and ethanol. Figure 6.7.4 shows their formulas along with a molecular model of each.




Figure 6.7.4: The Two Simplest Organic Alcohol Compounds. Alcohols have an OH functional group in the molecule.

Cholesterol, described in the chapter-opening essay, has an alcohol functional group, as its name implies.

Alcohol

Ethanol (also called ethyl alcohol) is the alcohol in alcoholic beverages. Other alcohols include methanol (or methyl alcohol), which is used as a solvent and a cleaner, and isopropyl alcohol (or rubbing alcohol), which is used as a medicinal disinfectant. Neither methanol nor isopropyl alcohol should be ingested, as they are toxic even in small quantities.

Another important family of organic compounds has a carboxyl group, in which a carbon atom is double-bonded to an oxygen atom and to an OH group. Compounds with a carboxyl functional group are called carboxylic acids, and their names end in *-oic acid*. Figure 6.7.5 shows the formulas and the molecular models of the two simplest carboxylic acids, perhaps best known by the common names formic acid and acetic acid. The carboxyl group is sometimes written in molecules as COOH.




Figure 6.7.5: The Two Smallest Organic Acids. The two smallest carboxylic acids are formic acid (found in the stingers of ants) and acetic acid (found in vinegar).


The condensed structures of methanoic acid and ethanoic acid are **HCOOH** and **CH_3COOH**, respectively.


Many organic compounds are considerably more complex than the examples described here. Many compounds, such as


cholesterol discussed in the chapter-opening essay, contain more than one functional group. The formal names can also be quite complex.

Example 6.7.1

Identify the functional group(s) in each molecule as a double bond, a triple bond, an alcohol, or a carboxyl.

a. 

b. 

c. 


d. $CH_3CH_2CH_2CH_2OH$


Answer a available at the end of this section.


Exercise 6.7.1

Identify the functional group(s) in each molecule as a double bond, a triple bond, an alcohol, or a carboxyl.

a. 

b. 

c. 

d. 

Answer a: available at the end of this section.

Career Focus: Forensic Chemist

The main job of a forensic chemist is to identify unknown materials and their origins. Although forensic chemists are most closely associated in the public mind with crime labs, they are employed in pursuits as diverse as tracing evolutionary patterns in living organisms, identifying environmental contaminants, and determining the origin of manufactured chemicals.

In a crime lab, the forensic chemist has the job of identifying the evidence so that a crime can be solved. The unknown samples may consist of almost anything—for example, paint chips, blood, glass, cloth fibers, drugs, or human remains. The forensic chemist subjects them to a variety of chemical and instrumental tests to discover what the samples are. Sometimes these samples are extremely small, but sophisticated forensic labs have state-of-the-art equipment capable of identifying the smallest amount of unknown sample.

Another aspect of a forensic chemist's job is testifying in court. Judges and juries need to be informed about the results of forensic analyses, and it is the forensic chemist's job to explain those results. Good public-speaking skills, along with a broad background in chemistry, are necessary to be a successful forensic chemist.

Key Takeaways

- Organic chemistry is the study of the chemistry of carbon compounds.
- Organic molecules can be classified according to the types of elements and bonds in the molecules.

Answers

Example 6.7.1

This molecule has a double bond and a carboxyl functional group.

This molecule has an alcohol functional group.

This molecule has a carbon-carbon double bond and a carboxyl functional group.

This molecule has an alcohol functional group.

Exercise 6.7.1

triple bond (alkyne)

carboxyl group

alcohol group

double bond (alkene) and carboxyl group

This page titled 6.7: Organic Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.6: Organic Chemistry by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

6.S: Covalent Bonding and Simple Molecular Compounds (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Atoms can share pairs of valence electrons to obtain a valence shell octet. This sharing of electrons is a **covalent bond**. A species formed from covalently bonded atoms is a **molecule** and is represented by a **molecular formula**, which gives the number of atoms of each type in the molecule. The two electrons shared in a covalent bond are called a **bonding pair of electrons**. The electrons that do not participate in covalent bonds are called **nonbonding pairs** (or **lone pairs**) **of electrons**. A covalent bond consisting of one pair of shared electrons is called a **single bond**.

Covalent bonds occur between nonmetal atoms. Naming simple covalent compounds follows simple rules similar to those for ionic compounds. However, for covalent compounds, numerical prefixes are used as necessary to specify the number of atoms of each element in the compound.

In some cases, more than one pair of electrons is shared to satisfy the octet rule. Two pairs of electrons are shared by two atoms to make a **double bond**. Three pairs of atoms are shared to make a **triple bond**. Single, double, and triple covalent bonds may be represented by one, two, or three dashes, respectively, between the symbols of the atoms.

The distance between two covalently bonded atoms is the **bond length**. Bond lengths depend on the types of atoms participating in the bond as well as the number of electron pairs being shared. A covalent bond can be a **polar covalent bond** if the electron sharing between the two atoms is unequal. If the sharing is equal, the bond is a **nonpolar covalent bond**. Because the strength of an atom's attraction for electrons in a bond is rated by the atom's **electronegativity**, the difference in the two atoms' electronegativities indicates how polar a covalent bond between those atoms will be.

The mass of a molecule is called its **molecular mass** and is the sum of the masses of the atoms in the molecule. The shape of a molecule can be predicted using **valence shell electron pair repulsion (VSEPR)**, which uses the fact that the negative electrons in covalent bonds repel each other as much as possible. Molecules with polar bonds are **polar** except when the bond polarities cancel due to symmetry.

Organic chemistry is the chemistry of carbon compounds. Carbon forms covalent bonds with other carbon atoms and with the atoms of many other elements. The simplest organic compounds are **hydrocarbons**, which consist solely of carbon and hydrogen. Hydrocarbons containing only single bonds are called **alkanes (saturated hydrocarbons)**. Hydrocarbons containing carbon–carbon double bonds are **alkenes**, while hydrocarbons with carbon–carbon triple bonds are **alkynes**. Carbon-carbon double and triple bonds are examples of **functional groups**, atoms or bonds that impart a characteristic chemical function to the molecule. Other functional groups include the alcohol functional group (OH) and the **carboxyl functional group** (COOH). They are the characteristic functional group in organic compounds called **alcohols** and **carboxylic acids**.

This page titled 6.S: Covalent Bonding and Simple Molecular Compounds (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 4.S: Covalent Bonding and Simple Molecular Compounds (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7: INTRODUCTION TO CHEMICAL REACTIONS

Chemical change is a central concept in chemistry. The goal of chemists is to know how and why a substance changes in the presence of another substance or even by itself. Because there are tens of millions of known substances, there are a huge number of possible chemical reactions. In this chapter, we will find that many of these reactions can be classified into a small number of categories according to certain shared characteristics.

7.1: Prelude to Introduction to Chemical Reactions

7.2: The Law of Conservation of Matter

7.3: Chemical Equations

7.4: Quantitative Relationships Based on Chemical Equations

7.5: Some Types of Chemical Reactions

7.6: Oxidation-Reduction (Redox) Reactions

7.7: Redox Reactions in Organic Chemistry and Biochemistry

7.S: Introduction to Chemical Reactions (Summary)

Template:HideTOC

http://en.Wikipedia.org/wiki/Exother...teReaction.jpg

This page titled 7: Introduction to Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5: Introduction to Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.1: Prelude to Introduction to Chemical Reactions

Although yeast has been used for thousands of years, its true nature has been known only for the last two centuries. Yeasts are single-celled fungi. About 1,000 species are recognized, but the most common species is *Saccharomyces cerevisiae*, which is used in bread making. Other species are used for the fermentation of alcoholic beverages. Some species can cause infections in humans.

Yeasts live primarily on sugars, such as glucose ($\mathrm{C_6H_{12}O_6}$). They convert glucose into carbon dioxide ($\mathrm{CO_2}$) and ethanol ($\mathrm{C_2H_5OH}$) in a chemical transformation that is represented as follows:

$$\mathrm{C_6H_{12}O_6\,(s) \longrightarrow 2\,CO_2\,(g) + 2\,C_2H_5OH(\ell)}$$

Bread making depends on the production of carbon dioxide. The gas, which is produced in tiny pockets in bread dough, acts as a leavening agent: it expands during baking and makes the bread rise. Leavened bread is softer, lighter, and easier to eat and chew than unleavened bread. The other major use of yeast, fermentation, depends on the production of ethanol, which results from the same chemical transformation. Some alcoholic beverages, such as champagne, can also be carbonated using the carbon dioxide produced by the yeast.

Yeast is among the simplest life forms on Earth, yet it is absolutely necessary for at least two major food industries. Without yeast to turn dough into bread and juice into wine, these foods and food industries would not exist today.

This page titled 7.1: Prelude to Introduction to Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.0: Prelude to Introduction to Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.2: The Law of Conservation of Matter

Learning Objectives

- Correctly define a law as it pertains to science.
- State the law of conservation of matter.

In science, a law is a general statement that explains a large number of observations. Before being accepted, a law must be verified many times under many conditions. Laws are therefore considered the highest form of scientific knowledge and are generally thought to be inviolable. Scientific laws form the core of scientific knowledge. One scientific law that provides the foundation for understanding in chemistry is the law of conservation of matter. It states that in any given system that is closed to the transfer of matter (in and out), the amount of matter in the system stays constant. A concise way of expressing this law is to say that the amount of matter in a system is *conserved*.

With the development of more precise ideas on elements, compounds and mixtures, scientists began to investigate how and why substances react. French chemist A. Lavoisier laid the foundation to the scientific investigation of matter by describing that substances react by following certain laws. These laws are called the laws of chemical combination. These eventually formed the basis of Dalton's Atomic Theory of Matter.

Law of Conservation of Mass

According to this law, during any physical or chemical change, the total mass of the products remains equal to the total mass of the reactants.

$$\underbrace{\overbrace{\mathrm{HgO\,(s)}}^{\text{Mercuric oxide}}}_{100\text{ g}} \rightarrow \underbrace{\overbrace{\mathrm{Hg\,(l)}}^{\text{Mercury}}}_{92.6\text{ g}} + \underbrace{\overbrace{\mathrm{O_2\,(g)}}^{\text{Oxygen}}}_{7.4\text{ g}}$$

Another way of stating this is, "In a chemical reaction, matter is neither created nor destroyed." The law of conservation of mass is also known as the "law of indestructibility of matter."

Example 7.2.1

If heating 10 grams of $\mathrm{CaCO_3}$ produces 4.4 g of $\mathrm{CO_2}$ and 5.6 g of CaO, show that these observations are in agreement with the law of conservation of mass.



A sample of calcium carbonate (CaCO3). (Public Domain; Walkerma).

Solution

- Mass of the reactants, $\mathrm{CaCO_3}$: $10\ g$
- Mass of the products, $\mathrm{CO_2}$ and CaO: $4.4\ g + 5.6\ g = 10\ g$.

Because the mass of the reactants = the mass of the products, the observations are in agreement with the law of conservation of mass.

What does this mean for chemistry? In any chemical change, one or more initial substances change into a different substance or substances. Both the initial and final substances are composed of atoms because all matter is composed of atoms. According to the law of conservation of matter, matter is neither created nor destroyed, so we must have the **same number and kind of atoms** after the chemical change as were present before the chemical change.

It may seem as though burning destroys matter, but the same amount, or mass, of matter still exists after a campfire as before. Figure 5.1.1 shows that when wood burns, it combines with oxygen and changes not only to ashes, but also to carbon dioxide and water vapor. The gases float off into the air, leaving behind just the ashes. Suppose we had measured the mass of the wood before it burned and the mass of the ashes after it burned. Also suppose we had been able to measure the oxygen used by the fire and the gases produced by the fire. What would we find? The total mass of matter after the fire would be the same as the total mass of matter before the fire.




Figure 7.2.1: Burning is a chemical process. The flames are caused as a result of a fuel undergoing combustion (burning). Images used with permission (CC BY-SA 2.5; Einar Helland Berger for fire and for ash).

Exercise 7.2.1

a. What is the law of conservation of matter?
b. How does the law of conservation of matter apply to chemistry?

Answer a: available at the end of this section.

Key Takeaway

The amount of matter in a closed system is conserved.

Answers

Exercise 7.2.1

The law of conservation of matter states that in any given system that is closed to the transfer of matter, the amount of matter in the system stays constant
The law of conservation of matter says that in chemical reactions, the total mass of the products must equal the total mass of the reactants.

This page titled 7.2: The Law of Conservation of Matter is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Binod Shrestha via source content that was edited to the style and standards of the LibreTexts platform.

- 5.1: The Law of Conservation of Matter by Binod Shrestha is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.3: Chemical Equations

Learning Objectives

- Define chemical reaction.
- Use a balanced chemical equation to represent a chemical reaction.

Water (H_2O) is composed of hydrogen and oxygen. Suppose we imagine a process in which we take some elemental hydrogen (H_2) and elemental oxygen (O_2) and let them react to make water. The statement

"*hydrogen and oxygen react to make water*"

is one way to represent that process, which is called a chemical reaction. Figure 7.3.1 shows a rather dramatic example of this very reaction.




Figure 7.3.1: The Formation of Water. Hydrogen and oxygen combine to form water. Here, the hydrogen gas in the zeppelin SS *Hindenburg* reacts with oxygen in the air to make water. Source: Photo courtesy of the US Navy. For a video of this see www.youtube.com/watch?v=CgWHbpMVQ1U.

To simplify the writing of reactions, we use formulas instead of names when we describe a reaction. We can also use symbols to represent other words in the reaction. A **plus sign** connects the initial substances (and final substances, if there is more than one), and an **arrow** ($\rightarrow$) represents the chemical change:

$$\mathrm{H_2 + O_2 \ \rightarrow H_2O} \tag{7.3.1}$$

This statement is one example of a **chemical equation**, an abbreviated way of using symbols to represent a **chemical change**. The substances on the left side of the arrow are called **reactants**, and the substances on the right side of the arrow are called **products**. It is not uncommon to include a phase label with each formula—(s) for solid, (ℓ) for liquid, (g) for gas, and (aq) for a substance dissolved in water, also known as an *aqueous solution*. If we included phase labels for the reactants and products, under normal environmental conditions, the reaction would be as follows:

$$\mathrm{H_2\,(g) + O_2\,(g) \ \rightarrow H_2O\ (\ell)} \tag{7.3.2}$$

> Chemical equations can also be used to describe physical changes. We will see examples of this soon.

This equation is still not complete because *it does not satisfy the* ***law of conservation of matter***. Count the **number of atoms of each element** on each side of the arrow. On the reactant side, there are two H atoms and two O atoms; on the product side, there are two H atoms and only one oxygen atom. The equation is **not balanced** because the number of oxygen atoms on each

side is not the same (Figure 7.3.2).




Figure 7.3.2: Balanced—Yes or No?. By counting the atoms of each element, we can see that the reaction is not balanced as written.

To make this chemical equation conform to the law of conservation of matter, we must revise the amounts of the reactants and the products as necessary to get the same number of atoms of a given element on each side. Because every substance has a characteristic chemical formula, we cannot change the chemical formulas of the individual substances. For example, we cannot change the formula for elemental oxygen to O. However, we can assume that different numbers of reactant molecules or product molecules may be involved. For instance, perhaps two water molecules are produced, not just one:

$$\mathrm{H_2\,(g) + O_2\,(g)\ \rightarrow 2\,H_2O\ (\ell)} \tag{7.3.3}$$

The **2** preceding the formula for water is called a **coefficient**. It implies that two water molecules are formed. There are now two oxygen atoms on each side of the equation.

> This point is so important that we should repeat it. You **cannot** change the formula of a chemical substance to balance a chemical reaction! You **must** use the proper chemical formula of the substance.

Unfortunately, by inserting the coefficient 2 in front of the formula for water, we have also changed the number of hydrogen atoms on the product side as well. As a result, we no longer have the same number of hydrogen atoms on each side. This can be easily fixed, however, by putting a coefficient of 2 in front of the diatomic hydrogen reactant:

$$\mathrm{2\,H_2\,(g) + O_2\,(g)\ \rightarrow 2\,H_2O\ (\ell)} \tag{7.3.4}$$

Now we have four hydrogen atoms and two oxygen atoms on each side of the equation. The **law of conservation of matter is satisfied** because we now have the same number of atoms of each element in the reactants and in the products. We say that the reaction is **balanced** (Figure 7.3.3). The diatomic oxygen has a coefficient of 1, which typically is not written but assumed in balanced chemical equations.




Figure 7.3.3: Balanced—Yes or No?. By counting the atoms of each element, we can see that the reaction is now balanced.

Proper chemical equations should be balanced. Writing balanced reactions is a chemist's way of acknowledging the law of conservation of matter.

Example 7.3.1

Is each chemical equation balanced?

a. $\mathrm{2\,Na\,(s) + O_2\,(g) \longrightarrow 2\,Na_2O\,(s)}$

b. $CH_4(g) + 2\,O_2(g) \longrightarrow CO_2(g) + 2\,H_2O(\ell)$
c. $AgNO_3(aq) + 2\,KCl(aq) \longrightarrow AgCl(s) + KNO_3(aq)$

Solution

a. By counting, we find two sodium atoms and two oxygen atoms in the reactants and four sodium atoms and two oxygen atoms in the products. This equation is not balanced.
b. The reactants have one carbon atom, four hydrogen atoms, and four oxygen atoms. The products have one carbon atom, four hydrogen atoms, and four oxygen atoms. This equation is balanced.
c. The reactants have one silver atom, one nitrogen atom, three oxygen atoms, two potassium atoms, and two chlorine atoms. The products have one silver atom, one chlorine atom, one potassium atom, one nitrogen atom, and three oxygen atoms. Because there are different numbers of chlorine and potassium atoms, this equation is not balanced.

Exercise 7.3.1

Is each chemical equation balanced?

a. $2\,Hg(l) + O_2(g) \longrightarrow Hg_2O_2(s)$
b. $C_2H_4(g) + 2\,O_2(g) \longrightarrow 2\,CO_2(g) + 2\,H_2O(l)$
c. $Mg(NO_3)_2(s) + 2\,Li(s) \longrightarrow Mg(s) + 2\,LiNO_3(s)$.

Answer a: available at the end of this section.

How does one balance a chemical equation, starting with the correct formulas of the reactants and products? Basically, a **back-and-forth approach** is adopted, counting the number of atoms of one element on one side, checking the number of atoms of that element on the other side, and changing a coefficient if necessary. Then check another element, going back and forth from one side of the equation to another, until each element has the same number of atoms on both sides of the arrow. In many cases, it does not matter which element is balanced first and which is balanced last, as long as all elements have the same number of atoms on each side of the equation.

Below are guidelines for writing and balancing chemical equations.

Writing and Balancing Chemical Equations

1. Determine the correct chemical formulas for each reactant and product. Write the skeleton equation.
2. Count the number of atoms of each element that appears as a reactant and as a product. If a polyatomic ion is unchanged on both sides of the equation, count it as a unit.
3. Balance each element one at a time by placing coefficients in front of the formulas. No coefficient is written for a 1. It is best to begin by balancing elements that only appear in one chemical formula on each side of the equation. NEVER change the subscripts in a chemical formula - you can only balance equations by using coefficients.
4. Check each atom or polyatomic ion to be sure that they are equal on both sides of the equation.
5. Make sure that all coefficients are in the lowest possible ratio. If necessary, reduce to the lowest ratio.

For example, to balance the equation

$$CH_4 + Cl_2 \rightarrow CCl_4 + HCl$$

Step 1: Write the skeleton equation with the correct formulas.

$$CH_4 + Cl_2 \rightarrow CCl_4 + HCl \tag{7.3.5}$$

Step 2: Count the number of each atom or polyatomic ion on both sides of the equation.

Reactants	Products
1 C atom	1 C atom
4 H ions	1 H ions
2 Cl atom	5 Cl atoms

Step 3: We find that both sides are already balanced with one carbon atom. So we proceed to balance the hydrogen atoms. We find that the reactant side has four hydrogen atoms, so the product side must also have four hydrogen atoms. This is balanced by putting a 4 in front of the HCl:

$$\mathrm{CH_4 + Cl_2 \rightarrow CCl_4 + 4\,HCl} \tag{7.3.6}$$

Reactants	Products
1 C atom	1 C atom
4 H ions	4 H ions
2 Cl atom	8 Cl atoms

Now each side has four hydrogen atoms. The product side has a total of eight chlorine atoms (four from the $\mathrm{CCl_4}$ and four from the four molecules of HCl), so we need eight chlorine atoms as reactants. Because elemental chlorine is a diatomic molecule, we need four chlorine molecules to get a total of eight chlorine atoms. We add another 4 in front of the $\mathrm{Cl_2}$ reactant:

$$\mathrm{CH_4 + 4\,Cl_2 \rightarrow CCl_4 + 4\,HCl} \tag{7.3.7}$$

Reactants	Products
1 C atom	1 C atom
4 H ions	4 H ions
8 Cl atom	8 Cl atoms

Step 3: Now we check: each side has one carbon atom, four hydrogen atoms, and eight chlorine atoms. The chemical equation is balanced. And, the coefficients are in the lowest possible ratio.

Example 7.3.2

Aqueous solutions of lead (II) nitrate and sodium chloride are mixed. The products of the reaction are an aqueous solution of sodium nitrate and a solid precipitate of lead (II) chloride. Write the balanced chemical equation for this reaction.

Solution

Step 1: Write the skeleton equation with the correct formulas.

$$\mathrm{Pb(NO_3)_2\,(aq) + NaCl\,(aq) \rightarrow NaNO_3\,(aq) + PbCl_2\,(s)}$$

Step 2: Count the number of each atom or polyatomic ion on both sides of the equation.

Reactants	Products
1 Pb atom	1 Pb atom
2 $\mathrm{NO_3^-}$ ions	1 $\mathrm{NO_3^-}$ ions
1 Na atom	1 Na atom
1 Cl atom	2 Cl atoms

Step 3: Solve. The nitrate ions and the chlorine atoms are unbalanced. Start by placing a 2 in front of the NaCl. This increases the reactant counts to 2 Na atoms and 2 Cl atoms. Then place a 2 in front of the $\mathrm{NaNO_3}$. The result is:

$$\mathrm{Pb(NO_3)_2\,(aq) + 2NaCl\,(aq) \rightarrow 2NaNO_3\,(aq) + PbCl_2\,(s)}$$

Step 4: The new count for each atom and polyatomic ion becomes:

Reactants	Products
1 Pb atom	1 Pb atom
2 NO_3^- ions	2 NO_3^- ions
2 Na atom	2 Na atom
2 Cl atom	2 Cl atoms

Step 5: Think about the result.

The equation is now balanced since there are equal numbers of atoms of each element on both sides of the equation. And, the coefficients are in the lowest possible ratio.

Exercise 7.3.2

Fermentation is a biochemical process that enables yeast cells to live in the absence of oxygen. Humans have exploited it for centuries to produce wine and beer and make bread rise. In fermentation, sugars such as glucose ($C_6H_{12}O_6$) are converted to ethanol (C_2H_5OH) and carbon dioxide CO_2. Write a balanced chemical reaction for the fermentation of glucose.



Commercial use of fermentation. (a) Microbrewery vats are used to prepare beer. (b) The fermentation of glucose by yeast cells is the reaction that makes beer production possible.

Answer available at the end of this section.

Key Takeaway

Chemical reactions are represented by chemical equations that list reactants and products. Proper chemical equations are balanced; the same number of each element's atoms appears on each side of the equation.

Answers

Exercise 7.3.1

balanced

O is not balanced; the 4 atoms of oxygen on the left does not balance with the 6 oxygen atoms on the right

balanced

Exercise 7.3.2

$$C_6H_{12}O_6\,(s) \longrightarrow 2\,C_2H_5OH\,(l) + 2\,CO_2\,(g)$$

This page titled 7.3: Chemical Equations is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.2: Chemical Equations by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.4: Quantitative Relationships Based on Chemical Equations

Learning Objectives

- To calculate the amount of one substance that will react with or be produced from a given amount of another substance.

A balanced chemical equation not only describes some of the chemical properties of substances—by showing us what substances react with what other substances to make what products—but also shows numerical relationships between the reactants and the products. The study of these numerical relationships is called stoichiometry. The stoichiometry of chemical equations revolves around the coefficients in the balanced chemical equation because these coefficients determine the molecular ratio in which reactants react and products are made.

> The word *stoichiometry* is pronounced "stow-eh-key-OM-et-tree." It is of mixed Greek and English origins, meaning roughly "measure of an element."

Looking Closer: Stoichiometry in Cooking

Let us consider a stoichiometry analogy from the kitchen. A recipe that makes 1 dozen biscuits needs 2 cups of flour, 1 egg, 4 tablespoons of shortening, 1 teaspoon of salt, 1 teaspoon of baking soda, and 1 cup of milk. If we were to write this as a chemical equation, we would write

$$\text{2 c flour + 1 egg + 4 tbsp shortening + 1 tsp salt + 1 tsp baking soda + 1 c milk} \rightarrow \text{12 biscuits}$$

(Unlike true chemical reactions, this one has all 1 coefficients written explicitly—partly because of the many different units here.) This equation gives us ratios of how much of what reactants are needed to make how much of what product. Two cups of flour, when combined with the proper amounts of the other ingredients, will yield 12 biscuits. One teaspoon of baking soda (when also combined with the right amounts of the other ingredients) will make 12 biscuits. One egg must be combined with 1 cup of milk to yield the product food. Other relationships can also be expressed.

We can use the ratios we derive from the equation for predictive purposes. For instance, if we have 4 cups of flour, how many biscuits can we make if we have enough of the other ingredients? It should be apparent that we can make a double recipe of 24 biscuits.

But how would we find this answer formally, that is, mathematically? We would set up a conversion factor, much like we did in Chapter 1. Because 2 cups of flour make 12 biscuits, we can set up an equivalency ratio:

$$\frac{\text{12 biscuits}}{\text{2 c flour}}$$

We then can use this ratio in a formal conversion of flour to biscuits:

$$\text{4 c flour} \times \frac{\text{12 biscuits}}{\text{2 c flour}} = \text{24 biscuits}$$

Similarly, by constructing similar ratios, we can determine how many biscuits we can make from any amount of ingredient. When you are doubling or halving a recipe, you are doing a type of stoichiometry. Applying these ideas to chemical reactions should not be difficult if you use recipes when you cook.



A recipe shows how much of each ingredient is needed for the proper reaction to take place. (CC BY-SA 3.0 Unported; Peng via Wikipedia)

Consider the following balanced chemical equation:

$$2\,C_2H_2 + 5\,O_2 \rightarrow 4\,CO_2 + 2\,H_2O \tag{7.4.1}$$

The coefficients on the chemical formulas give the ratios in which the reactants combine and the products form. Thus, we can make the following statements and construct the following ratios:

Table 7.4.1: Uses a statement from the balanced chemical reaction to show ratios and inverse ratios.

Statement from the Balanced Chemical Reaction	Ratio	Inverse Ratio
two C_2H_2 molecules react with five O_2 molecules	$\frac{2C_2H_2}{5O_2}$	$\frac{5O_2}{2C_2H_2}$
two C_2H_2 molecules react to make four CO_2 molecules	$\frac{2C_2H_2}{4CO_2}$	$\frac{4CO_2}{2C_2H_2}$
five O_2 molecules react to make two H_2O molecules	$\frac{5O_2}{2H_2O}$	$\frac{2H_2O}{5O_2}$
four CO_2 molecules are made at the same time as two H_2O molecules	$\frac{2H_2O}{4CO_2}$	$\frac{4CO_2}{2H_2O}$

Other relationships are possible; in fact, 12 different conversion factors can be constructed from this balanced chemical equation. In each ratio, the unit is assumed to be molecules because that is how we are interpreting the chemical equation.

Any of these fractions can be used as a conversion factor to relate an amount of one substance to an amount of another substance. For example, suppose we want to know how many CO_2 molecules are formed when 26 molecules of C_2H_2 are reacted. As usual with a conversion problem, we start with the amount we are given—$26C_2H_2$—and multiply it by a conversion factor that cancels out our original unit and introduces the unit we are converting to—in this case, CO_2. That conversion factor is $\frac{4CO_2}{2C_2H_2}$, which is composed of terms that come directly from the balanced chemical equation. Thus, we have

$$26C_2H_2 \times \frac{4CO_2}{2C_2H_2}$$

The molecules of C_2H_2 cancel, and we are left with molecules of CO_2. Multiplying through, we get

$$26C_2H_2 \times \frac{4CO_2}{2C_2H_2} = 52CO_2$$

Thus, 52 molecules of CO_2 are formed.

This application of stoichiometry is extremely powerful in its predictive ability, as long as we begin with a balanced chemical equation. Without a balanced chemical equation, the predictions made by simple stoichiometric calculations will be incorrect.

Example 7.4.1

Start with this balanced chemical equation.

$$\mathrm{KMnO_4 + 8\,HCl + 5\,FeCl_2 \longrightarrow 5\,FeCl_3 + MnCl_2 + 4\,H_2O + KCl}$$

a. Verify that the equation is indeed balanced.
b. Give 2 ratios that give the relationship between HCl and $\mathrm{FeCl_3}$.

Solution a

Each side has 1 K atom and 1 Mn atom. The 8 molecules of HCl yield 8 H atoms, and the 4 molecules of H_2O also yield 8 H atoms, so the H atoms are balanced. The Fe atoms are balanced, as we count 5 Fe atoms from 5 $FeCl_2$ reactants and 5 $FeCl_3$ products. As for Cl, on the reactant side, there are 8 Cl atoms from HCl and 10 Cl atoms from the 5 $FeCl_2$ formula units, for a total of 18 Cl atoms. On the product side, there are 15 Cl atoms from the 5 $FeCl_3$ formula units, 2 from the $MnCl_2$ formula unit, and 1 from the KCl formula unit. This is a total of 18 Cl atoms in the products, so the Cl atoms are balanced. All the elements are balanced, so the entire chemical equation is balanced.

Solution b

Because the balanced chemical equation tells us that 8 HCl molecules react to make 5 $FeCl_3$ formula units, we have the following 2 ratios: $\frac{8\mathrm{HCl}}{5\mathrm{FeCl_3}}$ and $\frac{5\mathrm{FeCl_3}}{8\mathrm{HCl}}$. There are a total of 42 possible ratios. Can you find the other 40 relationships?

Exercise 7.4.1

Start with this balanced chemical equation.

$$\mathrm{2KMnO_4 + 3CH_2{=}CH_2 + 4H_2O \rightarrow 2MnO_2 + 3HOCH_2CH_2OH + 2KOH}$$

a. Verify that the equation is balanced.
b. Give 2 ratios that give the relationship between $KMnO_4$ and $CH_2{=}CH_2$. (A total of 30 relationships can be constructed from this chemical equation. Can you find the other 28?)

Answer a: available at the end of this section.

Key Takeaway

The coefficients in a balanced chemical equation give the ratios in which molecules of substances react and are produced in a chemical reaction.

Answers

Exercise 7.4.1

Each side has 2 K atoms and 2 Mn atoms. On the reactant side, 3 $CH_2{=}CH_2$ yield 6 C atoms and on the product side, 3 $HOCH_2CH_2OH$ also yield 6 C atoms, so the C atoms are balanced. There are 20 H atoms on the reactants side: 12 H atoms from 3 $CH_2{=}CH_2$ and 8 H atoms from 4 H_2O. On the product side, there are also 20 H atoms: 18 H atoms from 3 $HOCH_2CH_2OH$ and 2 H atoms from 2 KOH. So, the H atoms are balanced. As for O, on the reactant side, there are 8 O atoms from 2 $KMnO_4$ and 4 O atoms from 4 H_2O, for a total of 12 O atoms. On the product side, there are 4 O atoms from the 2 MnO_2 formula units, 6 O atoms from 3 $HOCH_2CH_2OH$, and 2 O atoms from the 2 KOH formula units. This is a total of 12 O atoms in the products, so the O atoms are balanced. All the elements are balanced, so the entire chemical equation is balanced.

Because the balanced chemical equation tells us that 2 $KMnO_4$ formula units react with 3 $CH_2{=}CH_2$ molecules, we have the following 2 ratios:

$\frac{2\mathrm{KMnO_4}}{3\mathrm{CH_2{=}CH_2}}$ and $\frac{3\mathrm{CH_2{=}CH_2}}{2\mathrm{KMnO_4}}$. There are a total of 30 possible ratios. Can you find the other 28 relationships?

This page titled 7.4: Quantitative Relationships Based on Chemical Equations is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.3: Quantitative Relationships Based on Chemical Equations by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.5: Some Types of Chemical Reactions

Learning Objectives

- To classify a given chemical reaction into a variety of types.

Although there are untold millions of possible chemical reactions, most can be classified into a small number of general reaction types. Classifying reactions has two purposes: it helps us to recognize similarities among them, and it enables us to predict the products of certain reactions. A particular reaction may fall into more than one of the categories that we will define in this book.

Combination (composition) Reactions

A combination (composition) reaction is a chemical reaction that makes a single substance from two or more reactants. There may be more than one molecule of product in the balanced chemical equation, but there is only one substance produced. For example, the equation

$$4\,\mathrm{Fe} + 3\,\mathrm{O_2} \rightarrow 2\,\mathrm{Fe_2O_3}$$

is a combination reaction that produces $\mathrm{Fe_2O_3}$ from its constituent elements — Fe and $\mathrm{O_2}$. Combination reactions do not have to combine elements, however. The chemical equation

$$\mathrm{Fe_2O_3} + 3\,\mathrm{SO_3} \rightarrow \mathrm{Fe_2(SO_4)_3}$$

shows a combination reaction in which $\mathrm{Fe_2O_3}$ combines with three molecules of $\mathrm{SO_3}$ to make $\mathrm{Fe_2(SO_4)_3}$.

Example 7.5.1

Which equations are combination reactions?

a. $\mathrm{Co\,(s) + Cl_2\,(g) \longrightarrow CoCl_2\,(s)}$
b. $\mathrm{CO\,(g) + Cl_2\,(g) \longrightarrow COCl_2\,(g)}$
c. $\mathrm{N_2H_4(\ell) + O_2\,(g) \longrightarrow N_2\,(g) + 2\,H_2O(\ell)}$

Solution

a. This is a combination reaction.
b. This is a combination reaction. (The compound $\mathrm{COCl_2}$ is called phosgene and, in the past, was used as a gassing agent in chemical warfare.)
c. This is not a combination reaction.

Exercise 7.5.1

Which equations are combination reactions?

a. $\mathrm{P_4\,(s) + 6\,Cl_2\,(g) \longrightarrow 4\,PCl_3\,(g)}$
b. $\mathrm{SO_3(\ell) + H_2O(\ell) \longrightarrow H_2SO_4(\ell)}$
c. $\mathrm{NaOH\,(s) + HCl\,(g) \longrightarrow NaCl\,(s) + H_2O(\ell)}$

Answer available at the end of this section.

Decomposition Reactions

A decomposition reaction is the reverse of a combination reaction. In a decomposition reaction, a single substance is converted into two or more products. There may be more than one molecule of the reactant, but there is only one substance initially. For example, the equation

$$2\,\mathrm{NaHCO_3\,(s)} \rightarrow \mathrm{Na_2CO_3\,(s)} + \mathrm{CO_2\,(g)} + \mathrm{H_2O}\ (\ell)$$

is a decomposition reaction that occurs when $\mathrm{NaHCO_3}$ is exposed to heat. Another example is the decomposition of $\mathrm{KClO_3}$:

$$2\,\mathrm{KClO_3\,(s)} \rightarrow 2\,\mathrm{KCl\,(s)} + 3\,\mathrm{O_2\,(g)}$$

This reaction was once commonly used to generate small amounts of oxygen in the chemistry lab.

> The decomposition reaction of $\mathrm{NaHCO_3}$ is the reaction that occurs when baking soda is poured on a small kitchen fire. The intent is that the $\mathrm{H_2O}$ and $\mathrm{CO_2}$ produced by the decomposition will smother the flames.

Combustion Reactions

A combustion reaction occurs when a substance combines with molecular oxygen to make oxygen-containing compounds of other elements in the reaction. Many combustion reactions occur with a hydrocarbon, a compound made up solely of carbon and hydrogen. The products of the combustion of hydrocarbons are carbon dioxide and water. Many hydrocarbons are used as fuel because their combustion releases very large amounts of heat energy. An example is propane ($\mathrm{C_3H_8}$), a gaseous hydrocarbon that is commonly used as the fuel source in gas grills.

$$\mathrm{C_3H_8\,(g)} + 5\,\mathrm{O_2\,(g)} \longrightarrow 3\,\mathrm{CO_2\,(g)} + 4\,\mathrm{H_2O\,(g)}$$

Another example is the burning of acetylene ($\mathrm{C_2H_2}$) in welding torches:


$$2\,\mathrm{C_2H_2} + 5\,\mathrm{O_2} \rightarrow 4\,\mathrm{CO_2} + 2\,\mathrm{H_2O}$$

Oxygen (in its elemental form) is a crucial reactant in combustion reactions.

Energy in the form of heat is usually given off as a product in a combustion reaction as well.

Example 7.5.2

Carbon dioxide ($\mathrm{CO_2}$) is an important heat-trapping (greenhouse) gas, which is released through human activities such as deforestation and burning fossil fuels, as well as natural processes such as respiration and volcanic eruptions. The graph above shows $\mathrm{CO_2}$ levels during the last three glacial cycles, as reconstructed from ice cores.



Carbon dioxide is the primary greenhouse gas emitted through human activities. In 2015, CO_2 accounted for about 82.2% of all U.S. greenhouse gas emissions from human activities. Carbon dioxide is naturally present in the atmosphere as part of the Earth's carbon cycle (the natural circulation of carbon among the atmosphere, oceans, soil, plants, and animals). Human activities are altering the carbon cycle–both by adding more CO_2 to the atmosphere and by influencing the ability of natural sinks, like forests, to remove CO_2 from the atmosphere. While CO_2 emissions come from a variety of natural sources, human-related emissions are responsible for the increase that has occurred in the atmosphere since the industrial revolution.

The main human activity that emits CO_2 is the combustion of fossil fuels (coal, natural gas, and oil) for energy and transportation, although certain industrial processes and land-use changes also emit CO_2. An example of how CO_2 can be generated is the **combustion of octane** (C_8H_{10}), a component of gasoline.

Write the balanced equation to represent the combustion of octane.

Solution

$$2\,C_8H_{18}\,(l) + 25\,O_2\,(g) \longrightarrow 16\,CO_2\,(g) + 18\,H_2O\,(g)$$

The balanced reaction argues that for every two molecules of octane that are burned, 16 molecules of CO_2 are generated.

Example 7.5.2

Identify each type of reaction.

a. $2\,K\,(s) + S\,(s) + 2\,O_2\,(g) \longrightarrow K_2SO_4\,(s)$
b. $(NH_4)_2Cr_2O_7\,(s) \longrightarrow N_2\,(g) + Cr_2O_3\,(s) + 4\,H_2O(\ell)$
c. $CH_4\,(g) + 2\,O_2\,(g) \longrightarrow CO_2\,(g) + 2\,H_2O(\ell)$

Solution

a. Multiple reactants are combining to make a single product, so this reaction is a combination reaction.
b. A single substance reacts to make several products, so we have a decomposition reaction.

c. Oxygen reacts with a compound to make carbon dioxide (an oxide of carbon) and water (an oxide of hydrogen). This is a combustion reaction.

Exercise 7.5.2

Identify each type of reaction.

a. $C_2H_5OH + 3\,O_2 \longrightarrow 2\,CO_2 + 3\,H_2O$
b. $2\,Mg\,(s) + O_2\,(g) \longrightarrow 2\,MgO\,(s)$
c. $CaCO_3\,(s) \longrightarrow CaO\,(s) + CO_2\,(g)$

Answer a available at the end of this section.

Key Takeaway

There are several recognizable types of chemical reactions: combination, decomposition, and combustion reactions are examples.

Answers

Exercise 7.5.1
a and b are combination reactions

Exercise 7.5.2
combustion
combination (also combustion)
decomposition

This page titled 7.5: Some Types of Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.4: Some Types of Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.6: Oxidation-Reduction (Redox) Reactions

Learning Objectives

- To identify a chemical reaction as an oxidation-reduction reaction.

When zinc metal is submerged into a quantity of aqueous HCl, the following reaction occurs (Figure $7.6.1$):

$$\mathrm{Zn\,(s) + 2\,HCl\,(aq) \longrightarrow H_2\,(g) + ZnCl_2\,(aq)} \tag{7.6.1}$$

This is one example of what is sometimes called a *single replacement reaction* because Zn replaces H in combination with Cl.




Figure $7.6.1$: Zinc Metal plus Hydrochloric Acid. It is fairly obvious that zinc metal reacts with aqueous hydrochloric acid! The bubbles are hydrogen gas (right side of Equation $7.6.1$).

Because some of the substances in this reaction are aqueous, we can separate them into ions:

$$\mathrm{Zn\,(s) + 2\,H^+\,(aq) + 2\,Cl^-\,(aq) \longrightarrow H_2\,(g) + Zn^{2+}\,(aq) + 2\,Cl^-\,(aq)}$$

Viewed this way, the net reaction seems to be a charge transfer between zinc and hydrogen atoms. (There is no net change experienced by the chloride ion.) In fact, electrons are being transferred from the zinc atoms to the hydrogen atoms (which ultimately make a molecule of diatomic hydrogen), changing the charges on both elements.

To understand electron-transfer reactions like the one between zinc metal and hydrogen ions, chemists separate them into two parts: one part focuses on the loss of electrons, and one part focuses on the gain of electrons. The loss of electrons is called oxidation. The gain of electrons is called reduction. Because any loss of electrons by one substance must be accompanied by a gain in electrons by something else, oxidation and reduction always occur together. As such, electron-transfer reactions are also called oxidation-reduction reactions, or simply **redox reactions**. The atom that loses electrons is **oxidized**, and the atom that gains electrons is **reduced**. Also, because we can think of the species being oxidized as causing the reduction, the species being oxidized is called the **reducing agent**, and the species being reduced is called the **oxidizing agent**.

Because batteries are used as sources of electricity (that is, of electrons), all batteries are based on redox reactions.

Although the two reactions occur together, it can be helpful to write the oxidation and reduction reactions separately as half reactions. In half reactions, we include only the reactant being oxidized or reduced, the corresponding product species, any other species needed to balance the half reaction, and the electrons being transferred. Electrons that are lost are written as products; electrons that are gained are written as reactants. For example, in our earlier equation, now written without the chloride ions,

$$\mathrm{Zn\,(s) + 2\,H^+\,(aq) \longrightarrow Zn^{2+}\,(aq) + H_2\,(g)}$$

zinc atoms are oxidized to Zn^{2+}. The **half reaction for the oxidation** reaction, omitting phase labels, is as follows:

$$\mathrm{Zn \longrightarrow Zn^{2+} + 2\,e^-}$$

This half reaction is balanced in terms of the number of zinc atoms, and it also shows the two electrons that are needed as products to account for the zinc atom losing two negative charges to become a 2+ ion. With half reactions, there is one more item to balance: the overall charge on each side of the reaction. If you check each side of this reaction, you will note that both sides have a zero net charge.

Hydrogen is reduced in the reaction. The balanced **reduction half reaction** is as follows:

$$\mathrm{2\,H^+ + 2\,e^- \longrightarrow H_2}$$

There are two hydrogen atoms on each side, and the two electrons written as reactants serve to neutralize the 2+ charge on the reactant hydrogen ions. Again, the overall charge on both sides is zero.

The overall reaction is simply the combination of the two half reactions and is shown by adding them together.

$$\begin{aligned} \mathrm{Zn} &\rightarrow \mathrm{Zn^{2+}} + \cancel{2e^-} \\ \mathrm{2H^+} + \cancel{2e^-} &\rightarrow \mathrm{H_2} \\ \hline \mathrm{Zn + 2H^+} &\rightarrow \mathrm{Zn^{2+} + H_2} \end{aligned}$$

Because we have two electrons on each side of the equation, they can be canceled. This is the key criterion for a balanced redox reaction: the electrons have to cancel exactly. If we check the charge on both sides of the equation, we see they are the same—2+. (In reality, this positive charge is balanced by the negative charges of the chloride ions, which are not included in this reaction because chlorine does not participate in the charge transfer.)

Redox reactions are often balanced by balancing each individual half reaction and then combining the two balanced half reactions. **Sometimes a half reaction must have all of its coefficients multiplied by some integer for all the electrons to cancel**. The following example demonstrates this process.

Example 7.6.1: Reducing Silver Ions

Write and balance the redox reaction that has silver ions and aluminum metal as reactants and silver metal and aluminum ions as products. Identify the substance oxidized, substance reduced, reducing agent and reducing agent.

Solution

We start by using symbols of the elements and ions to represent the reaction:

$$\mathrm{Ag^+ + Al \longrightarrow Ag + Al^{3+}}$$

The equation looks balanced as it is written. However, when we compare the overall charges on each side of the equation, we find a charge of +1 on the left but a charge of +3 on the right. This equation is not properly balanced. To balance it, let us write the two half reactions. Silver ions are reduced, and it takes one electron to change $\mathrm{Ag^+}$ to Ag:

Reduction half-reaction:

$$\mathrm{Ag^+ + e^- \longrightarrow Ag}$$

Aluminum is oxidized, losing three electrons to change from Al to $\mathrm{Al^{3+}}$:

Oxidation half-reaction:

$$\mathrm{Al} \longrightarrow \mathrm{Al}^{3+} + 3\,\mathrm{e}^-$$

To combine these two half reactions and cancel out all the electrons, we need to multiply the silver reduction reaction by 3:

$$3\left(\mathrm{Ag}^+ + \mathrm{e}^- \rightarrow \mathrm{Ag}\right)$$
$$\mathrm{Al} \rightarrow \mathrm{Al}^{3+} + 3\mathrm{e}^-$$
$$3\mathrm{Ag}^+ + \mathrm{Al} \rightarrow 3\mathrm{Ag} + \mathrm{Al}^{3+}$$

Now the equation is balanced, not only in terms of elements but also in terms of charge.

- The **substance oxidized** is the reactant that had undergone oxidation: Al
- The **substance reduced** is the reactant that had undergone reduction: Ag^+
- The **reducing agent** is the same as the substance oxidized: Al
- The **oxidizing agent** is the same as the substance reduced: Ag^+

Exercise 7.6.1

Write and balance the redox reaction that has calcium ions and potassium metal as reactants and calcium metal and potassium ions as products. Identify the substance oxidized, substance reduced, reducing agent and reducing agent.

Answer available at the end of this section.

To Your Health: Redox Reactions and Pacemaker Batteries

All batteries use redox reactions to supply electricity because electricity is basically a stream of electrons being transferred from one substance to another. Pacemakers—surgically implanted devices for regulating a person's heartbeat—are powered by tiny batteries, so the proper operation of a pacemaker depends on a redox reaction.

Pacemakers used to be powered by NiCad batteries, in which nickel and cadmium (hence the name of the battery) react with water according to this redox reaction:

$$\mathrm{Cd\,(s)} + 2\,\mathrm{NiOOH\,(s)} + 2\,\mathrm{H_2O}(\ell) \longrightarrow \mathrm{Cd(OH)_2\,(s)} + 2\,\mathrm{Ni(OH)_2\,(s)}$$

The cadmium is oxidized, while the nickel atoms in NiOOH are reduced. Except for the water, all the substances in this reaction are solids, allowing NiCad batteries to be recharged hundreds of times before they stop operating. Unfortunately, NiCad batteries are fairly heavy batteries to be carrying around in a pacemaker. Today, the lighter lithium/iodine battery is used instead. The iodine is dissolved in a solid polymer support, and the overall redox reaction is as follows:

$$2\,\mathrm{Li\,(s)} + \mathrm{I_2\,(s)} \longrightarrow 2\,\mathrm{LiI\,(s)}$$

Lithium is oxidized, and iodine is reduced. Although the lithium/iodine battery cannot be recharged, one of its advantages is that it lasts up to 10 years. Thus, a person with a pacemaker does not have to worry about periodic recharging; about once per decade a person requires minor surgery to replace the pacemaker/battery unit. Lithium/iodine batteries are also used to power calculators and watches.



Figure 7.6.2: A small button battery like this is used to power a watch, pacemaker, or calculator. (CC BY-SA; Gerhard H Wrodnigg via Wikipedia)

Oxidation and reduction can also be defined in terms of changes in composition. The original meaning of **oxidation** was "**adding oxygen**," so when oxygen is added to a molecule, the molecule is being oxidized. The reverse is true for **reduction**: if a molecule **loses oxygen** atoms, the molecule is being reduced. For example, the acetaldehyde (CH_3CHO) molecule takes on an oxygen atom to become acetic acid (CH_3COOH).

$$2\,CH_3CHO + O_2 \longrightarrow 2\,CH_3COOH$$

Thus, acetaldehyde is being oxidized.

Similarly, **reduction and oxidation can be defined in terms of the gain or loss of hydrogen atoms**. If a molecule adds hydrogen atoms, it is being reduced. If a molecule loses hydrogen atoms, the molecule is being oxidized. For example, in the conversion of acetaldehyde into ethanol (CH_3CH_2OH), hydrogen atoms are added to acetaldehyde, so the acetaldehyde is being reduced:

$$CH_3CHO + H_2 \longrightarrow CH_3CH_2OH$$

Table 7.6.1: shows how each process affects the change in oxygen and change in hydrogen.

Process	Change in oxygen (some reactions)	Change in hydrogen (some reactions)
Oxidation	gain	lose
Reduction	lose	gain

Example 7.6.2

In each conversion, indicate whether oxidation or reduction is occurring.

a. $N_2 \longrightarrow NH_3$
b. $CH_3CH_2OHCH_3 \longrightarrow CH_3COCH_3$
c. $HCHO \longrightarrow HCOOH$

Solution

a. Hydrogen is being added to the original reactant molecule, so reduction is occurring.
b. Hydrogen is being removed from the original reactant molecule, so oxidation is occurring.

c. Oxygen is being added to the original reactant molecule, so oxidation is occurring.

Exercise 7.6.2

In each conversion, indicate whether oxidation or reduction is occurring.

a. $\mathrm{CH_4} \longrightarrow \mathrm{CO_2} + \mathrm{H_2O}$
b. $\mathrm{NO_2} \longrightarrow \mathrm{N_2}$
c. $\mathrm{CH_2{=}CH_2} \longrightarrow \mathrm{CH_3CH_3}$

Answer a: available at the end of this section.

Answer b: available at the end of this section.

Answer a: available at the end of this section.

Key Takeaway

Chemical reactions in which electrons are transferred are called oxidation-reduction, or redox, reactions. Oxidation is the loss of electrons. Reduction is the gain of electrons. Oxidation and reduction always occur together, even though they can be written as separate chemical equations.

Answers

Exercise 7.6.1

Reduction: $\mathrm{Ca^{2+} + 2\,e^- \longrightarrow Ca}$

Oxidation: \(2 \times (\ce{K -> K^{+} + e^{-}}\)

Combined: $\mathrm{Ca^{2+} + 2\,K \longrightarrow Ca + 2\,K^+}$

- The **substance oxidized** is the reactant that had undergone oxidation: K
- The **substance reduced** is the reactant that had undergone reduction: $\mathrm{Ca^{2+}}$
- The **reducing agent** is the same as the substance oxidized: K
- The **oxidizing agent** is the same as the substance reduced: $\mathrm{Ca^{2+}}$

Potassium has been used as a reducing agent to obtain various metals in their elemental form.

Exercise 7.6.2

Oxygen is being added. Oxidation is occurring.

Exercise 7.6.2

Oxygen is being removed. Reduction is occurring.

Exercise 7.6.2

Hydrogen is being added. Reduction is occurring.

This page titled 7.6: Oxidation-Reduction (Redox) Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.5: Oxidation-Reduction (Redox) Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.7: Redox Reactions in Organic Chemistry and Biochemistry

Learning Objectives

- To identify oxidation-reduction reactions with organic compounds.

Oxidation-reduction reactions are of central importance in organic chemistry and biochemistry. The burning of fuels that provides the energy to maintain our civilization and the metabolism of foods that furnish the energy that keeps us alive both involve redox reactions.



Figure 7.7.1: The Burning of Natural Gas. The burning of natural gas is not only a combustion reaction but also a redox reaction. Similar reactions include the burning of gasoline and coal. These are also redox reactions. (CC BY-SA 2.0 Generic; Simon Law via Flickr).

All combustion reactions are also redox reactions. A typical combustion reaction is the burning of methane, the principal component of natural gas (Figure 7.7.1).

$$\mathrm{CH_4 + 2\,O_2 \rightarrow CO_2 + 2\,H_2O} \tag{7.7.1}$$

In respiration, the biochemical process by which the oxygen we inhale in air oxidizes foodstuffs to carbon dioxide and water, redox reactions provide energy to living cells. A typical respiratory reaction is the oxidation of glucose ($\mathrm{C_6H_{12}O_6}$), the simple sugar we encountered in the chapter-opening essay that makes up the diet of yeast:

$$\mathrm{C_6H_{12}O_6 + 6\,O_2 \rightarrow 6\,CO_2 + 6\,H_2O} \tag{7.7.2}$$

Organic chemists use a variety of redox reactions. For example, potassium dichromate ($\mathrm{K_2Cr_2O_7}$) is a common oxidizing agent that can be used to oxidize alcohols (symbolized by the general formula ROH). The product of the reaction depends on the location of the OH functional group in the alcohol molecule, the relative proportions of alcohol and the dichromate ion, and reaction conditions such as temperature. If the OH group is attached to a terminal carbon atom and the product is distilled off as it forms, the product is an aldehyde, which has a terminal *carbonyl group* (C=O) and is often written as RCHO. One example is the reaction used by the Breathalyzer to detect ethyl alcohol ($\mathrm{CH_3CH_2OH}$) in a person's breath:

$$\mathrm{3\,CH_3CH_2OH + Cr_2O_7^{2-} + 8\,H^+ \rightarrow 3\,CH_3CHO + 2\,Cr^{3+} + 7\,H_2O} \tag{7.7.3}$$


If the product acetaldehyde ($\mathrm{CH_3CHO}$) is not removed as it forms, it is further oxidized to acetic acid ($\mathrm{CH_3COOH}$). In this case, the overall reaction is as follows:

$$\mathrm{3\,CH_3CH_2OH + 2\,Cr_2O_7^{2-} + 16\,H^+ \rightarrow 3\,CH_3COOH + 4\,Cr^{3+} + 11\,H_2O} \tag{7.7.4}$$

In this reaction, the chromium atom is reduced because it lost oxygen atoms from $Cr_2O_7^{2-}$ to Cr^{3+}. On the other hand, the carbon atom in ethanol is oxidized. In the **oxidation** of ethyl alcohol (CH_3CH_2OH, a.k.a. ethanol) to form acetaldehyde (CH_3CHO, a.k.a. ethanal), the **number of bonds to oxygen has increased and the number of hydrogen atoms has decreased** from six to four. Either or both of these indicate that an oxidation has occurred.

the reaction of ethanol to ethanal

In the oxidation of acetaldehyde to acetic acid (a.k.a. ethanoic acid), the carbon atom that gained an additional oxygen is the element oxidized.



When the OH group of the alcohol is bonded to an interior carbon atom, the oxidation of an alcohol will produce a ketone (the formulas of ketones are often written as RCOR, and the carbon–oxygen bond is a double bond). The simplest ketone is derived from 2-propanol ($CH_3CHOHCH_3$). It is the common solvent acetone [$(CH_3)_2CO$], which is used in varnishes, lacquers, rubber cement, and nail polish remover. Acetone can be formed by the following redox reaction:

$$3\,CH_3CHOHCH_3 + Cr_2O_7{}^{2-} + 8\,H^+ \rightarrow 3\,(CH_3)_2CO + 2\,Cr^{3+} + 7\,H_2O \tag{7.7.5}$$

As we have just seen, aldehydes and ketones can be formed by the oxidation of alcohols. Conversely, aldehydes and ketones can be reduced to alcohols. Reduction of the carbonyl group is important in living organisms. For example, in anaerobic metabolism, in which biochemical processes take place in the absence of oxygen, pyruvic acid ($CH_3COCOOH$) is reduced to lactic acid ($CH_3CHOHCOOH$) in the muscles.

$$CH_3COCOOH \rightarrow CH_3CHOHCOOH \tag{7.7.6}$$

(Pyruvic acid is both a carboxylic acid and a ketone; only the ketone group is reduced.) The buildup of lactic acid during vigorous exercise is responsible in large part for the fatigue that we experience.

In food chemistry, the substances known as antioxidants are reducing agents. Ascorbic acid (vitamin C; $C_6H_8O_6$) is thought to retard potentially damaging oxidation of living cells. In the process, it is oxidized to dehydroascorbic acid ($C_6H_6O_6$). In the stomach, ascorbic acid reduces the nitrite ion ($NO_2{}^-$) to nitric oxide (NO):

$$C_6H_8O_6 + 2\,H^+ + 2\,NO_2{}^- \rightarrow C_6H_6O_6 + 2\,H_2O + 2\,NO \tag{7.7.7}$$

> If reaction in Equation 7.7.7 did not occur, nitrite ions from foods would oxidize the iron in hemoglobin, destroying its ability to carry oxygen.

Tocopherol (vitamin E) is also an antioxidant. In the body, **vitamin E** is thought to act by scavenging harmful by-products of metabolism, such as the highly reactive molecular fragments called free radicals. In foods, vitamin E acts to prevent fats from being oxidized and thus becoming rancid. **Vitamin C** is also a good antioxidant (Figure 7.7.2).



Figure 7.7.2: Citrus Fruits. Citrus fruits, such as oranges, lemons, and limes, are good sources of vitamin C, which is an antioxidant. (Pexels License; Scott Webb via Pexels)

Finally, and of greatest importance, green plants carry out the redox reaction that makes possible almost all life on Earth. They do this through a process called photosynthesis, in which carbon dioxide and water are converted to glucose ($C_6H_{12}O_6$). The synthesis of glucose requires a variety of proteins called enzymes and a green pigment called chlorophyll that converts sunlight into chemical energy (Figure 7.7.3). The overall change that occurs is as follows:

$$6\,CO_2 + 6\,H_2O \rightarrow C_6H_{12}O_6 + 6\,O_2 \tag{7.7.8}$$

In this reaction, carbon dioxide is reduced to glucose, and water is oxidized to oxygen gas. Other reactions convert the glucose to more complex carbohydrates, plant proteins, and oils.



Figure 7.7.3: Life on Earth. Photosynthesis is the fundamental process by which plants use sunlight to convert carbon dioxide and water into glucose and oxygen. Then plants make more complex carbohydrates. It is the ultimate source of all food on Earth, and it is a redox reaction. (Public Domain; Wikipedia).

Summary

Redox reactions are common in organic and biological chemistry, including the combustion of organic chemicals, respiration,

and photosynthesis.

This page titled 7.7: Redox Reactions in Organic Chemistry and Biochemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 5.6: Redox Reactions in Organic Chemistry and Biochemistry by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

7.S: Introduction to Chemical Reactions (Summary)

This page titled 7.S: Introduction to Chemical Reactions (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

8: QUANTITIES IN CHEMICAL REACTIONS

So far, we have talked about chemical reactions in terms of individual atoms and molecules. Although this works, most of the reactions occurring around us involve much larger amounts of chemicals. Even a tiny sample of a substance will contain millions, billions, or a hundred billion billions of atoms and molecules. How do we compare amounts of substances to each other in chemical terms when it is so difficult to count to a hundred billion billion? Actually, there are ways to do this, which we will explore in this chapter. In doing so, we will increase our understanding of stoichiometry, which is the study of the numerical relationships between the reactants and the products in a balanced chemical reaction.

8.1: Prelude to Quantities in Chemical Reactions

8.2: The Mole

8.3: Atomic and Molar Masses

8.4: Mole-Mass Conversions

8.5: Mole-Mole Relationships in Chemical Reactions

8.6: Mole-Mass and Mass-Mass Problems

8.S: Quantities in Chemical Reactions (Summary)

Template:HideTOC

This page titled 8: Quantities in Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6: Quantities in Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.1: Prelude to Quantities in Chemical Reactions

> When the disengaged gasses are carefully examined, they are found to weigh **113.7 grs.**; these are of two kinds, viz. **144 cubical inches** of carbonic acid gas, weighing **100 grs.** and **380 cubical inches** of a very light gas, weighing only **13.7 grs.**...and, when the water which has passed over into the bottle [labeled] H is carefully examined, it is found to have lost **85.7 grs.** of its weight. Thus, in this experiment, **85.7 grs.** of water, joined to **28 grs.** of charcoal, have combined in such a way as to form **100 grs.** of carbonic acid, and **13.7 grs.** of a particular gas capable of being burnt. (Bold emphasis added.)

In this paragraph from the *Elements of Chemistry*, Antoine Lavoisier (1743–94) is explaining an experiment in which he was trying to demonstrate that water is not an element but instead is composed of hydrogen (the gas "capable of being burnt") and oxygen. This is a historical account of a groundbreaking experiment and illustrates the importance of amounts in chemistry. Lavoisier was pointing out that the initial total mass of water and charcoal, 85.7 g plus 28 g, equals the final total mass of carbonic acid and the particular gas, 100 g plus 13.7 g. In this way, he was illustrating the law of conservation of matter. It is another way of saying that *amounts matter*.

Amounts do matter and in a variety of circumstances. The chapter-opening essay in Chapter 1 tells the story of a nurse who mistakenly read "2–3 mg" as "23 mg" and administered the higher and potentially fatal dose of morphine to a child. Food scientists who work in test kitchens must keep track of specific amounts of ingredients as they develop new products for us to eat. Quality control technicians measure amounts of substances in manufactured products to ensure that the products meet company or government standards. Supermarkets routinely weigh meat and produce and charge consumers by the ounce or the pound.

This page titled 8.1: Prelude to Quantities in Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.0: Prelude to Quantities in Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.2: The Mole

Learning Objectives

- To define the mole unit.

Figure 8.2.1 shows that we need 2 hydrogen atoms and 1 oxygen atom to make 1 water molecule. If we want to make 2 water molecules, we will need 4 hydrogen atoms and 2 oxygen atoms. If we want to make 5 molecules of water, we need 10 hydrogen atoms and 5 oxygen atoms. The ratio of atoms we will need to make any number of water molecules is the same: 2 hydrogen atoms to 1 oxygen atom.




Figure 8.2.1 Water Molecules. The ratio of hydrogen atoms to oxygen atoms used to make water molecules is always 2:1, no matter how many water molecules are being made.

One problem we have, however, is that it is extremely difficult, if not impossible, to organize atoms one at a time. As stated in the introduction, we deal with billions of atoms at a time. How can we keep track of so many atoms (and molecules) at a time? We do it by using mass rather than by counting individual atoms.

A hydrogen atom has a mass of approximately 1 u. An oxygen atom has a mass of approximately 16 u. The ratio of the mass of an oxygen atom to the mass of a hydrogen atom is therefore approximately 16:1.

If we have 2 atoms of each element, the ratio of their masses is approximately 32:2, which reduces to 16:1—the same ratio. If we have 12 atoms of each element, the ratio of their total masses is approximately (12 × 16):(12 × 1), or 192:12, which also reduces to 16:1. If we have 100 atoms of each element, the ratio of the masses is approximately 1,600:100, which again reduces to 16:1. As long as we have equal numbers of hydrogen and oxygen atoms, the ratio of the masses will always be 16:1.

The same consistency is seen when ratios of the masses of other elements are compared. For example, the ratio of the masses of silicon atoms to equal numbers of hydrogen atoms is always approximately 28:1, while the ratio of the masses of calcium atoms to equal numbers of lithium atoms is approximately 40:7.

So we have established that the masses of atoms are constant with respect to each other, as long as we have the same number of each type of atom. Consider a more macroscopic example. If a sample contains 40 g of Ca, this sample has the same number of atoms as there are in a sample of 7 g of Li. What we need, then, is a number that represents a convenient quantity of atoms so we can relate macroscopic quantities of substances. Clearly even 12 atoms are too few because atoms themselves are so small. We need a number that represents billions and billions of atoms.

Chemists use the term **mole** to represent a large number of atoms or molecules. Just as a dozen implies 12 things, a **mole** (abbreviated as ***mol***) represents **6.022×10^{23} things**. The number 6.022×10^{23}, called Avogadro's number after the 19th-century chemist Amedeo Avogadro, is the number we use in chemistry to represent macroscopic amounts of atoms and molecules. Thus, if we have 6.022×10^{23} Na atoms, we say we have 1 mol of Na atoms. If we have 2 mol of Na atoms, we have $2 \times (6.022 \times 10^{23})$ Na atoms, or 1.2044×10^{24} Na atoms. Similarly, if we have 0.5 mol of benzene (C_6H_6) molecules, we have $0.5 \times (6.022 \times 10^{23})$ C_6H_6 molecules, or 3.011×10^{23} C_6H_6 molecules.

> A mole represents a very large number! If 1 mol of quarters were stacked in a column, it could stretch back and forth between Earth and the sun *6.8 billion* times.

Notice that we are applying the mole unit to different types of **chemical entities**. The word *mole* represents a number of things—6.022×10^{23} of them—but does not by itself specify what "they" are. The chemical entities can be ***atoms***, ***molecules***, ***formula units*** and ***ions***. This specific information needs to be specified accurately. Most students find this confusing hence, we need to review the composition of elements, covalent and ionic compounds.

Most elements are made up of individual ***atoms***, such as helium. However, some elements consist of ***molecules***, such as the diatomic elements, nitrogen, hydrogen, oxygen, etc. One mole of He consists of 6.022×10^{23} He ***atoms*** but one mole of nitrogen contains 6.022×10^{23} N_2 ***molecules***. The basic units of covalent (molecular) compounds are ***molecules*** as well. The molecules of "compounds" consist of different kinds of atoms while the molecules of "elements" consist of only one type of atom. For example, the molecules of ammonia (NH_3) consist of nitrogen and hydrogen atoms while N_2 molecules have N atoms only. Compounds that are ionic, like NaCl, are represented by ionic formulas. One mole of NaCl, for example, refers to 6.022×10^{23} ***formula units*** of NaCl. And, one formula unit of NaCl consists of one sodium ***ion*** and one chloride ***ion***. Figure 6.1.2 summarizes the basic units of elements, covalent and ionic compounds




Figure 8.2.2: The basic units of elements (atoms or molecules), covalent compounds (molecules) and ionic compounds (formula units of ions).

Conversion Between Moles and Atoms, Molecules and Ions

Using our unit conversion techniques learned in Chapter 1, we can use the mole relationship and the chemical formula to convert back and forth between the moles and the number of chemical entities (atoms, molecules or ions).

Because 1 N_2 molecule contains 2 N atoms, 1 mol of N_2 molecules (6.022×10^{23} molecules) has 2 mol of N atoms. Using formulas to indicate how many atoms of each element we have in a substance, we can relate the number of moles of molecules to the number of moles of atoms. For example, in 1 mol of ethanol (C_2H_6O), we can construct the following relationships (Table 8.2.1):

Table 8.2.1: Molecular Relationships

1 Molecule of C_2H_6O Has	1 Mol of C_2H_6O Has	Molecular Relationships
2 C atoms	2 mol of C atoms	$\frac{2\text{ mol C atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{2\text{ mol C atoms}}$
6 H atoms	6 mol of H atoms	$\frac{6\text{ mol H atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{6\text{ mol H atoms}}$
1 O atom	1 mol of O atoms	$\frac{1\text{ mol O atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}$ or $\frac{1\text{ mol }C_2H_6O\text{ molecules}}{1\text{ mol O atoms}}$

The following example illustrates how we can use these relationships as conversion factors.

Example 8.2.1

If a sample consists of 2.5 mol of ethanol (C_2H_6O), how many moles of carbon atoms, hydrogen atoms, and oxygen atoms does it have?

Solution

Using the relationships in Table 8.2.1, we apply the appropriate conversion factor for each element:

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \underbrace{\frac{2\text{ mol C atoms}}{1\text{ mol }C_2H_6O\text{ molecules}}}_{\text{conversion factor}} = 5.0\text{ mol C atoms}$$

Note how the unit *mol C_2H_6O molecules* cancels algebraically. Similar equations can be constructed for determining the number of H and O atoms:

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \frac{6\text{ mol H atoms}}{1\text{ mol }C_2H_6O\text{ molecules}} = 15\text{ mol H atoms}$$

$$2.5\text{ mol }C_2H_6O\text{ molecules} \times \frac{1\text{ mol O atoms}}{1\text{ mol }C_2H_6O\text{ molecules}} = 2.5\text{ mol O atoms}$$

Exercise 8.2.1

If a sample contains 6.75 mol of Na_2SO_4, how many moles of sodium atoms, sulfur atoms, and oxygen atoms does it have?

Answer available at the end of this section.

We can use Avogadro's number as a conversion factor, or ratio, in dimensional analysis problems. For example, if we are dealing with element X, the mole relationship is expressed as follows:

$$1\text{ mol X} = 6.022 \times 10^{23}\text{ X atoms}$$

We can convert this relationship into two possible conversion factors shown below:

$$\frac{1\text{ mol X}}{6.022 \times 10^{23}\text{ X atoms}} \text{ or } \frac{6.022 \times 10^{23}\text{ X atoms}}{1\text{ mol X}}$$

If the number of "atoms of element X" is given, we can convert it into "moles of X" by multiplying the given value with the conversion factor at the left. However, if the number of "mol of X" is given, the appropriate conversion factor to use is the one at the right.

If we are dealing with a molecular compound (such as C_4H_{10}), the mole relationship is expressed as follows:

$$1 \text{ mol } C4H10 = 6.022 \times 10^{23} \text{ } C4H10 \text{ molecules}$$

If working with ionic compounds (such as NaCl), the mole relationship is expressed as follows:

$$1 \text{ mol NaCl} = 6.022 \times 10^{23} \text{ NaCl formula units}$$

Example 8.2.2

How many formula units are present in 2.34 mol of NaCl? How many ions are in 2.34 mol?

Solution

Typically in a problem like this, we start with what we are given and apply the appropriate conversion factor. Here, we are given a quantity of 2.34 mol of NaCl, to which we can apply the definition of a **mole as a conversion factor**:




Figure 8.2.3 The process for converting moles of NaCl to formula units of NaCl

$$2.34 \text{ mol NaCl} \times \frac{6.022 \times 10^{23} \text{ NaCl units}}{1 \text{ mol NaCl}} = 1.41 \times 10^{24} \text{ NaCl units}$$

Because there are two ions per formula unit, there are

$$1.41 \times 10^{24} \text{ NaCl units} \times \frac{2 \text{ ions}}{\text{NaCl units}} = 2.82 \times 10^{24} \text{ ions}$$

in the sample.

Exercise 8.2.2

How many molecules are present in 16.02 mol of C_4H_{10}? How many atoms are in 16.02 mol?

Answer available at the end of this section.

Key Takeaway

- A mole is 6.022×10^{23} things.

Answers

Exercise 8.2.1

13.5 mol Na, 6.75 mol S and 27 mol O.

Exercise 8.2.2

9.647 x 10^{24}molecules, 1.351 x 10^{26} atoms.

This page titled 8.2: The Mole is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.1: The Mole by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.3: Atomic and Molar Masses

Learning Objectives

- To learn how the masses of moles of atoms and molecules are expressed.

Now that we have introduced the mole and practiced using it as a conversion factor, we ask the obvious question: why is the mole *that particular* number of things? Why is it 6.022×10^{23} and not 1×10^{23} or even 1×10^{20}?

The number in a mole, Avogadro's number, is related to the relative sizes of the **atomic mass unit** and **gram mass units**. Whereas one hydrogen atom has a mass of approximately 1 u, **1 mol of H atoms** has a mass of approximately **1 *gram***. And whereas one sodium atom has an approximate mass of 23 u, 1 mol of Na atoms has an approximate mass of 23 *grams*.

One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units. The numbers in the periodic table that we identified as the atomic masses of the atoms not only tell us the mass of one atom in u but also tell us the mass of 1 mol of atoms in grams.

One mole of a substance has the same mass in grams that one atom or molecule has in atomic mass units.

Example 8.3.1: Molar Mass of Elements

What is the mass of each quantity?

a. 1 mol of Al atoms
b. 2 mol of U atoms

Solution

a. One mole of Al atoms has a mass in grams that is numerically equivalent to the atomic mass of aluminum. The periodic table shows that the atomic mass (rounded to two decimal points) of Al is 26.98, so 1 mol of Al atoms has a mass of 26.98 g.
b. According to the periodic table, 1 mol of U has a mass of 238.0 g, so the mass of 2 mol is twice that, or 476.0 g.

Exercise 8.3.1: Molar Mass of Elements

What is the mass of each quantity?

a. 1 mol of Au atoms
b. 5 mol of Br atoms

Answer a available at the end of this section.

Answer b available at the end of this section.

The mole concept can be extended to masses of formula units and molecules as well. The mass of 1 mol of molecules (or formula units) in grams is numerically equivalent to the mass of one molecule (or formula unit) in atomic mass units. For example, a single molecule of O_2 has a mass of 32.00 u, and 1 mol of O_2 molecules has a mass of 32.00 g. As with atomic mass unit–based masses, to obtain the mass of 1 mol of a substance, we simply sum the masses of the individual atoms in the formula of that substance. The mass of 1 mol of a substance is referred to as its **molar mass**, whether the substance is an element, an ionic compound, or a covalent compound.

Example 8.3.2: Molar Mass of Compounds

What is the mass of 1 mol of each substance?

a. NaCl
b. bilirubin ($C_{33}H_{36}N_4O_6$), the principal pigment present in bile (a liver secretion)

Solution a

Summing the molar masses of the atoms in the NaCl formula unit gives

Solutions to Example 6.2.2, Summing the molar masses of the atoms in the NaCl formula unit gives

1 Na molar mass:	22.99 g
1 Cl molar mass:	35.45 g
Total:	58.44 g

The mass of 1 mol of NaCl is 58.44 g.

Solution b

Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

Solutions to Example 6.2.2, Multiplying the molar mass of each atom by the number of atoms of that type in bilirubin's formula and adding the results, we get

33 C molar mass:	33 × 12.01 g	396.33 g
36 H molar mass:	36 × 1.01 =	36.36 g
4 N molar mass:	4 × 14.01 =	56.04 g
6 O molar mass:	6 × 16.00 =	96.00 g
Total:		584.73 g

The mass of 1 mol of bilirubin is 584.73 g.

Exercise 8.3.2: Molar Mass of Compounds

What is the mass of 1 mol of each substance?

a. barium sulfate ($BaSO_4$), used to take X rays of the gastrointestional tract
b. adenosine ($C_{10}H_{13}N_5O_4$), a component of cell nuclei crucial for cell division

Answer a available at the end of this section.

Be careful when counting atoms. In formulas with polyatomic ions in parentheses, the subscript outside the parentheses is applied to every atom inside the parentheses. For example, the molar mass of $Ba(OH)_2$ requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

The molar mass of Ba(OH)2 requires the sum of 1 mass of Ba, 2 masses of O, and 2 masses of H:

1 Ba molar mass:	1 × 137.3 g =	137.3 g
2 O molar mass:	2 × 16.00 g =	32.00 g

2 H molar mass:	2 × 1.01 g =	2.02 g
Total:		171.32 g

Because molar mass is defined as the mass for 1 mol of a substance, we can refer to molar mass as grams per mole (g/mol). The division sign (/) implies “per,” and “1” is implied in the denominator. Thus, the molar mass of bilirubin can be expressed as 584.73 g/mol, which is read as “five hundred eighty four point seventy three grams per mole.”

Key Takeaway

- The mass of moles of atoms and molecules is expressed in units of grams.

Answers

Exercise 8.3.1: Molar Mass of Elements

197.0 g

Exercise 8.3.1: Molar Mass of Elements

5 mol Br atoms x 79.90 g/mol = 399.5 g

Exercise 8.3.2: Molar Mass of Compounds

233.36 g

267.28 g

This page titled 8.3: Atomic and Molar Masses is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.2: Atomic and Molar Masses by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.4: Mole-Mass Conversions

Learning Objectives

- To convert quantities between mass units and mole units.

A previous Example stated that the mass of 2 mol of U is twice the molar mass of uranium. Such a straightforward exercise does not require any formal mathematical treatment. Many questions concerning mass are not so straightforward, however, and require some mathematical manipulations.

The simplest type of manipulation using molar mass as a conversion factor is a mole-mass conversion (or its reverse, a mass-mole conversion). In such a conversion, we use the molar mass of a substance as a conversion factor to convert mole units into mass units (or, conversely, mass units into mole units).

We also established that 1 mol of Al has a mass of 26.98 g (Example 8.4.1). Stated mathematically,

$$1 \text{ mol Al} = 26.98 \text{ g Al}$$

We can divide both sides of this expression by either side to get one of two possible conversion factors:

$$\frac{1 \text{ mol Al}}{26.98 \text{ g Al}} \quad \text{and} \quad \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$

The first conversion factor can be used to convert from mass to moles, and the second converts from moles to mass. Both can be used to solve problems that would be hard to do “by eye.”

Example 8.4.1

What is the mass of 3.987 mol of Al?

Solution

The first step in a conversion problem is to decide what conversion factor to use. Because we are starting with mole units, we want a conversion factor that will cancel the mole unit and introduce the unit for mass in the numerator. Therefore, we should use the $\frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$ conversion factor. We start with the given quantity and multiply by the conversion factor:




Figure 8.4.1 An example for Converting between Mass and Moles of a Substance. It takes one mathematical step to convert from moles to mass or from mass to moles.

$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}}$$

Note that the mol units cancel algebraically. (The quantity 3.987 mol is understood to be in the numerator of a fraction that has 1 in the unwritten denominator.) Canceling and solving gives

$$3.987 \text{ mol Al} \times \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = 107.6 \text{ g Al}$$

Our final answer is expressed to four significant figures.

Exercise 8.4.1

How many moles are present in 100.0 g of Al? (Hint: you will have to use the other conversion factor we obtained for aluminum.)

Answer available at the end of this section.

Conversions like this are possible for any substance, as long as the proper atomic mass, formula mass, or molar mass is known (or can be determined) and expressed in grams per mole. Figure 8.4.1 is a chart for determining what conversion factor is needed, and Figure 8.4.2 is a flow diagram for the steps needed to perform a conversion.




Figure 8.4.2 A Simple Flowchart for Converting between Mass and Moles of a Substance. It takes one mathematical step to convert from moles to mass or from mass to moles.




Figure 8.4.3 A Flowchart Illustrating the Steps in Performing a Unit Conversion. When performing many unit conversions, the same logical steps can be taken.

Example 8.4.2

A biochemist needs 0.00655 mol of bilirubin ($C_{33}H_{36}N_4O_6$) for an experiment. How many grams of bilirubin will that be?

Solution

To convert from moles to mass, we need the molar mass of bilirubin, which we can determine from its chemical formula:

Solutions to Example 6.3.2

33 C molar mass:	33 × 12.01 g =	396.33 g
36 H molar mass:	36 × 1.01 g =	36.36 g
4 N molar mass:	4 × 14.01 g =	56.04 g
6 O molar mass:	6 × 16.00 g =	96.00 g
Total:		584.73 g

The molar mass of bilirubin is 584.73 g. Using the relationship

$$1 \text{ mol bilirubin} = 584.73 \text{ g bilirubin}$$

we can construct the appropriate conversion factor for determining how many grams there are in 0.00655 mol. Following the steps from Figure 8.4.2:

$$0.00655 \text{ mol bilirubin} \times \frac{584.73 \text{ g bilirubin}}{\text{mol bilirubin}} = 3.83 \text{ g bilirubin}$$

The mol bilirubin unit cancels. The biochemist needs 3.83 g of bilirubin.

Exercise 8.4.2

A chemist needs 457.8 g of $KMnO_4$ to make a solution. How many moles of $KMnO_4$ is that?

Answer available at the end of this section.

To Your Health: Minerals

For our bodies to function properly, we need to ingest certain substances from our diets. Among our dietary needs are minerals, the noncarbon elements our body uses for a variety of functions, such developing bone or ensuring proper nerve transmission. The US Department of Agriculture has established some recommendations for the RDIs of various minerals. The accompanying table lists the RDIs for minerals, both in mass and moles, assuming a 2,000-calorie daily diet.

Table 8.4.1: Essential Minerals and their Composition in Humans

Mineral	Male (age 19–30 y)		Female (age 19–30 y)	
Ca	1,000 mg	0.025 mol	1,000 mg	0.025 mol
Cr	35 µg	6.7×10^{-7} mol	25 µg	4.8×10^{-7} mol
Cu	900 µg	1.4×10^{-5} mol	900 µg	1.4×10^{-5} mol
F	4 mg	2.1×10^{-4} mol	3 mg	1.5×10^{-4} mol
I	150 µg	1.2×10^{-6} mol	150 µg	1.2×10^{-6} mol
Fe	8 mg	1.4×10^{-4} mol	18 mg	3.2×10^{-4} mol
K	3,500 mg	9.0×10^{-2} mol	3,500 mg	9.0×10^{-2} mol
Mg	400 mg	1.6×10^{-2} mol	310 mg	1.3×10^{-2} mol
Mn	2.3 mg	4.2×10^{-5} mol	1.8 mg	3.3×10^{-5} mol
Mo	45 mg	4.7×10^{-7} mol	45 mg	4.7×10^{-7} mol

Mineral	Male (age 19–30 y)		Female (age 19–30 y)	
Na	2,400 mg	1.0×10^{-1} mol	2,400 mg	1.0×10^{-1} mol
P	700 mg	2.3×10^{-2} mol	700 mg	2.3×10^{-2} mol
Se	55 µg	7.0×10^{-7} mol	55 µg	7.0×10^{-7} mol
Zn	11 mg	1.7×10^{-4} mol	8 mg	1.2×10^{-4} mol

Table 8.4.1 illustrates several things. First, the needs of men and women for some minerals are different. The extreme case is for iron; women need over twice as much as men do. In all other cases where there is a different RDI, men need more than women.

Second, the amounts of the various minerals needed on a daily basis vary widely—both on a mass scale and a molar scale. The average person needs 0.1 mol of Na a day, which is about 2.5 g. On the other hand, a person needs only about 25–35 µg of Cr per day, which is under one millionth of a mole. As small as this amount is, a deficiency of chromium in the diet can lead to diabetes-like symptoms or neurological problems, especially in the extremities (hands and feet). For some minerals, the body does not require much to keep itself operating properly.

Although a properly balanced diet will provide all the necessary minerals, some people take dietary supplements. However, too much of a good thing, even minerals, is not good. Exposure to too much chromium, for example, causes a skin irritation, and certain forms of chromium are known to cause cancer (as presented in the movie *Erin Brockovich*).

Key Takeaway

- It is possible to convert between moles of material and mass of material.

Answers

Exercise 8.4.1

$$100.0 \text{ g Al} \times \frac{1 \text{ mol Al}}{26.98 \text{ g Al}} = 3.706 \text{ mol Al}$$

Exercise 8.4.2

$$457.8 \text{ g } KMnO_4 \times \frac{1 \text{ mol } KMnO_4}{158.04 \text{ g } KMnO_4} = 2.897 \text{ mol } KMnO_4$$

This page titled 8.4: Mole-Mass Conversions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.3: Mole-Mass Conversions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.5: Mole-Mole Relationships in Chemical Reactions

Learning Objectives

- To use a balanced chemical reaction to determine molar relationships between the substances.

Coefficients as Molar Relationships

Previously, you learned to balance chemical equations by comparing the numbers of each type of atom in the reactants and products. The coefficients in front of the chemical formulas represent the numbers of molecules or formula units (depending on the type of substance). Here, we will extend the meaning of the coefficients in a chemical equation.

Consider the simple chemical equation

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

The convention for writing balanced chemical equations is to use the lowest whole-number ratio for the coefficients. However, the equation is balanced as long as the coefficients are in a 2:1:2 ratio. For example, this equation is also balanced if we write it as

$$4\,H_2 + 2\,O_2 \longrightarrow 4\,H_2O$$

The ratio of the coefficients is 4:2:4, which reduces to 2:1:2. The equation is also balanced if we were to write it as

$$22\,H_2 + 11\,O_2 \longrightarrow 22\,H_2O$$

because 22:11:22 also reduces to 2:1:2.

Suppose we want to use larger numbers. Consider the following coefficients:

$$12.044 \times 10^{23}\ H_2 + 6.022 \times 10^{23}\ O_2 \rightarrow 12.044 \times 10^{23}\ H_2O$$

These coefficients also have the ratio 2:1:2 (check it and see), so this equation is balanced. But 6.022×10^{23} is 1 mol, while 12.044×10^{23} is 2 mol (and the number is written that way to make this more obvious), so we can simplify this version of the equation by writing it as

$$2\ \text{mol}\ \ H_2 + 1\ \ \text{mol}\ \ O_2 \longrightarrow 2\ \text{mol}\ \ H_2O$$

We can leave out the word *mol* and not write the 1 coefficient (as is our habit), so the final form of the equation, still balanced, is

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

Now we interpret the coefficients as referring to molar amounts, not individual molecules. The lesson? *Balanced chemical equations are balanced not only at the molecular level but also in terms of molar amounts of reactants and products.* Thus, we can read this reaction as “two moles of hydrogen react with one mole of oxygen to produce two moles of water.”




Figure 8.5.1 A molecular model diagram illustrating the chemical reaction between hydrogen and oxygen to form water molecules.

2 molecules H_2 1 molecule O_2 2 molecules H_2O

2 moles H_2 1 mole O_2 2 moles H_2O

2 x 2.02 g=4.04 g H_2 32.0 g O_2 2 x 18.02 g=36.04 g H_2O

Figure 8.5.1: This representation of the production of water from oxygen and hydrogen show several ways to interpret the quantitative information of a chemical reaction.

By the same token, the ratios we constructed to describe molecules reaction can also be constructed in terms of moles rather than molecules. For the reaction in which hydrogen and oxygen combine to make water, for example, we can construct the following ratios:

$$\frac{2\ \text{mol}\ H_2}{1\ \text{mol}\ O_2}\ \text{or}\ \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2}$$

$$\frac{2\ \text{mol}\ H_2O}{1\ \text{mol}\ O_2}\ \text{or}\ \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2O}$$

$$\frac{2\ \text{mol}\ H_2}{2\ \text{mol}\ H_2O}\ \text{or}\ \frac{2\ \text{mol}\ H_2O}{2\ \text{mol}\ H_2}$$

We can use these ratios to determine what amount of a substance, in moles, will react with or produce a given number of moles of a different substance. The study of the numerical relationships between the reactants and the products in balanced chemical reactions is called ***stoichiometry*.** The ratio of coefficients in a balanced chemical equation, used in computations relating amounts of reactants and products is called the **stoichiometric factor.**

Example 8.5.1

How many moles of oxygen react with hydrogen to produce 27.6 mol of H_2O? The balanced equation is as follows:

$$2\,H_2 + O_2 \longrightarrow 2\,H_2O$$

Solution

Because we are dealing with quantities of H_2O and O_2, we will use the **stoichiometric ratio** that relates those two substances. Because we are given an amount of H_2O and want to determine an amount of O_2, we will use the ratio that has H_2O in the denominator (so it cancels) and O_2 in the numerator (so it is introduced in the answer). Thus,




Figure 8.5.2 A flow diagram illustrating the stoichiometric conversion between moles of water and moles of oxygen using the molar ratio, stoichometric ratio, derived from a balanced chemical equation.

$$27.6\ \text{mol}\ H_2O \times \frac{1\ \text{mol}\ O_2}{2\ \text{mol}\ H_2O} = 13.8\ \text{mol}\ O_2$$

To produce 27.6 mol of H_2O, 13.8 mol of O_2 react.

Exercise 8.5.1

Using $2H_2 + O_2 \rightarrow 2H_2O$, how many moles of hydrogen react with 3.07 mol of oxygen to produce H_2O?

Answer available at the end of this section.

Key Takeaway

- The balanced chemical reaction can be used to determine molar relationships between substances.

Answers

Exercise 8.5.1

$$3.07 \text{ mol } O_2 \times \frac{2 \text{ mol } H_2}{1 \text{ mol } O_2} = 6.14 \text{ mol } H_2$$

This page titled 8.5: Mole-Mole Relationships in Chemical Reactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.4: Mole-Mole Relationships in Chemical Reactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.6: Mole-Mass and Mass-Mass Problems

Learning Objectives

- To convert from mass or moles of one substance to mass or moles of another substance in a chemical reaction.

Mole-Mass Calculations

We have established that a balanced chemical equation is balanced in terms of moles as well as atoms or molecules. We have used balanced equations to set up ratios, now in terms of moles of materials, that we can use as conversion factors to answer stoichiometric questions, such as how many moles of substance A react with so many moles of reactant B. We can extend this technique even further. Recall that we can relate a molar amount to a mass amount using molar mass. We can use that ability to answer stoichiometry questions in terms of the masses of a particular substance, in addition to moles. We do this using the following sequence:

laboratory.




Figure 8.6.1 A stoichiometry flow chart demonstrating the multi-step process of converting moles of one substance into the mass (grams) of a different substance.

Collectively, these conversions are called mole-mass calculations.

As an example, consider the balanced chemical equation

$$\mathrm{Fe_2O_3} + 3\,\mathrm{SO_3} \rightarrow \mathrm{Fe_2(SO_4)_3}$$

If we have 3.59 mol of Fe_2O_3, how many grams of SO_3 can react with it? Using the mole-mass calculation sequence, we can determine the required mass of SO_3 in two steps. First, we construct the appropriate molar ratio, determined from the balanced chemical equation, to calculate the number of moles of SO_3 needed. Then using the molar mass of SO_3 as a conversion factor, we determine the mass that this number of moles of SO_3 has.

The first step resembles the exercises we did in Section 6.4. As usual, we start with the quantity we were given:

$$3.59\ \mathrm{mol\ Fe_2O_3} \times \frac{3\ \mathrm{mol\ SO_3}}{1\ \mathrm{mol\ Fe_2O_3}} = 10.77\ \mathrm{mol\ SO_3}$$

The mol Fe_2O_3 units cancel, leaving mol SO_3 unit. Now, we take this answer and convert it to grams of SO_3, using the molar mass of SO_3 as the conversion factor:

$$10.77\ \text{mol SO}_3 \times \frac{80.07\ \text{g SO}_3}{1\ \text{mol SO}_3} = 862.4\ \text{g SO}_3$$

Our final answer is expressed to three significant figures. Thus, in a two-step process, we find that 862 g of SO_3 will react with 3.59 mol of Fe_2O_3. Many problems of this type can be answered in this manner.

The same two-step problem can also be worked out in a single line, rather than as two separate steps, as follows:




Figure 8.6.2 A stoichiometry flow diagram illustrating the conversion of moles of ferric oxide to the mass of sulfur trioxide using a stoichiometric ratio and molar mass.

$$3.59\ \text{mol } \cancel{Fe_2O_3} \times \underbrace{\frac{3\ \text{mol } \cancel{SO_3}}{1\ \text{mol } \cancel{Fe_2O_3}}}_{\text{converts to moles of } SO_3} \times \underbrace{\frac{80.07\ \text{g } SO_3}{1\ \text{mol } \cancel{SO_3}}}_{\text{converts to grams of } SO_3} = 862.4\ \text{g SO}_3$$

Figure 8.6.3 AA dimensional aalysis calculation showing the multi-step conversion from moles of iron (III) oxide to grams of sulfur trioxide, demonstrating the cancellation of units.

We get exactly the same answer when combining all the math steps together as we do when we calculate one step at a time.

Example 8.6.1

How many grams of CO_2 are produced if 2.09 mol of HCl are reacted according to this balanced chemical equation?

$$CaCO_3 + 2\,HCl \rightarrow CaCl_2 + CO_2 + H_2O$$

Solution

Our strategy will be to convert from moles of HCl to moles of CO_2 and then from moles of CO_2 to grams of CO_2. We will need the molar mass of CO_2, which is 44.01 g/mol. Performing these two conversions in a single-line gives 46.0 g of CO_2:




Figure 8.6.4 A stoichiometry calculation using dimensional analysis to convert moles of hydrochloric acid to the mass of carbon dioxide produced.

The molar ratio between CO_2 and HCl comes from the balanced chemical equation.

Exercise

How many grams of glucose ($C_6H_{12}O_6$) are produced if 17.3 mol of H_2O are reacted according to this balanced chemical equation?

$$6\,CO_2 + 6\,H_2O \rightarrow C_6H_{12}O_6 + 6\,O_2$$

Answer available at the end of this section.

Mass-Mass Calculations

It is a small step from mole-mass calculations to mass-mass calculations. If we start with a known mass of one substance in a chemical reaction (instead of a known number of moles), we can calculate the corresponding masses of other substances in the reaction. The first step in this case is to convert the known mass into moles, using the substance's molar mass as the conversion factor. Then—and only then—we use the balanced chemical equation to construct a conversion factor to convert that quantity to moles of another substance, which in turn can be converted to a corresponding mass. Sequentially, the process is as follows:




Figure 8.6.5 A comprehensive stoichiometry flow chart illustrating the three-step process to convert the mass (grams) of one substance into the mass (grams) of a different substance.

This three-part process can be carried out in three discrete steps or combined into a single calculation that contains three conversion factors. The following example illustrates both techniques.

Example 8.6.2: Chlorination of Carbon

Methane can react with elemental chlorine to make carbon tetrachloride (CCl_4). The balanced chemical equation is as follows:

$$CH_4 + 4\,Cl_2 \rightarrow CCl_4 + 4\,HCl$$

How many grams of HCl are produced by the reaction of 100.0 g of CH_4?

Solution

First, let us work the problem in stepwise fashion. We begin by converting the mass of CH_4 to moles of CH_4, using the molar mass of CH_4 (16.05 g/mol) as the conversion factor:

$$100.0\text{ g } CH_4 \times \frac{1\text{ mol } CH_4}{16.05\text{ g } CH_4} = 6.231\text{ mol } CH_4$$

Note that we inverted the molar mass so that the gram units cancel, giving us an answer in moles. Next, we use the balanced chemical equation to determine the ratio of moles CCl_4 and moles HCl and convert our first result into moles of HCl:

$$6.231\text{ mol } CH_4 \times \frac{4\text{ mol HCl}}{1\text{ mol } CH_4} = 24.92\text{ mol HCl}$$

Finally, we use the molar mass of HCl (36.46 g/mol) as a conversion factor to calculate the mass of 24.92 mol of HCl:

$$24.92\text{ mol HCl} \times \frac{36.46\text{ g HCl}}{1\text{ mol HCl}} = 908.5\text{ g HCl}$$

In each step, we have limited the answer to the proper number of significant figures. If desired, we can do all three conversions on a single line:




Figure 8.6.6 A stoichiometry flow diagram illustrating the three-step conversion from the mass of methane to the mass of hydrochloric acid using molar masses and a stoichiometric ratio.

$$100.0\text{ g } CH_4 \times \frac{1\text{ mol } CH_4}{16.05\text{ g } CH_4} \times \frac{4\text{ mol HCl}}{1\text{ mol } CH_4} \times \frac{36.46\text{ g HCl}}{1\text{ mol HCl}} = 908.7\text{ g HCl}$$

This final answer is slightly different from our first answer because only the final answer is restricted to the proper number of significant figures. In the first answer, we limited each intermediate quantity to the proper number of significant figures. As you can see, both answers are essentially the same.

Exercise 8.6.2: Oxidation of Propanal

The oxidation of propanal (CH_3CH_2CHO) to propionic acid (CH_3CH_2COOH) has the following chemical equation:

$$CH_3CH_2CHO + 2K_2Cr_2O_7 \rightarrow CH_3CH_2COOH + \text{other products}$$

How many grams of propionic acid are produced by the reaction of 135.8 g of $K_2Cr_2O_7$?

Answer available at the end of this section.

To Your Health: The Synthesis of Taxol

Taxol is a powerful anticancer drug that was originally extracted from the Pacific yew tree (*Taxus brevifolia*). As you can see from the accompanying figure, taxol is a very complicated molecule, with a molecular formula of $C_{47}H_{51}NO_{14}$. Isolating taxol from its natural source presents certain challenges, mainly that the Pacific yew is a slow-growing tree, and the equivalent of six trees must be harvested to provide enough taxol to treat a single patient. Although related species of yew trees also produce taxol in small amounts, there is significant interest in synthesizing this complex molecule in the laboratory.




Figure 8.6.7 The Structure of the Cancer Drug Taxol. Because of the complexity of the molecule, hydrogen atoms are not shown, but they are present on every atom to give the atom the correct number of covalent bonds (four bonds for each carbon atom).

After a 20-year effort, two research groups announced the complete laboratory synthesis of taxol in 1994. However, each synthesis required over 30 separate chemical reactions, with an overall efficiency of less than 0.05%. To put this in perspective, to obtain a single 300 mg dose of taxol, you would have to begin with 600 g of starting material. To treat the 26,000 women who are diagnosed with ovarian cancer each year with one dose, almost 16,000 kg (over 17 tons) of starting material must be converted to taxol. Taxol is also used to treat breast cancer, with which 200,000 women in the United States are diagnosed every year. This only increases the amount of starting material needed.

Clearly, there is intense interest in increasing the overall efficiency of the taxol synthesis. An improved synthesis not only will be easier but also will produce less waste materials, which will allow more people to take advantage of this potentially life-saving drug.

Key Takeaway

- A balanced chemical equation can be used to relate masses or moles of different substances in a reaction.

Answers

Exercise

$$17.3 \text{ mol } H_2O \times \frac{1 \text{ mol } C_6H_{12}O_6}{6 \text{ mol } H_2O} \times \frac{180.18 \text{ g } C_6H_{12}O_6}{1 \text{ mol } C_6H_{12}O_6} = 520 \text{ g } C_6H_{12}O_6$$

Exercise 8.6.2: Oxidation of Propanal

$$135.8 \text{ g } K_2Cr_2O_7 \times \frac{1 \text{ mol } K_2Cr_2O_7}{294.20 \text{ g } K_2Cr_2O_7} \times \frac{1 \text{ mol } CH_3CH_2COOH}{2 \text{ mol } K_2Cr_2O_7} \times \frac{74.09 \text{ g } CH_3CH_2COOH}{1 \text{ mol } CH_3CH_2COOH} = 17.10 \text{ g } CH_3CH_2COOH$$

This page titled 8.6: Mole-Mass and Mass-Mass Problems is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.5: Mole-Mass and Mass-Mass Problems by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

8.S: Quantities in Chemical Reactions (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

Chemical reactions relate quantities of reactants and products. Chemists use the **mole** unit to represent 6.022×10^{23} things, whether the things are atoms of elements or molecules of compounds. This number, called **Avogadro's number**, is important because this number of atoms or molecules has the same mass in grams as one atom or molecule has in atomic mass units. **Molar masses** of substances can be determined by summing the appropriate masses from the periodic table; the final molar mass will have units of grams.

Because one mole of a substance will have a certain mass, we can use that relationship to construct conversion factors that will convert a mole amount into a mass amount, or vice versa. Such **mole-mass conversions** typically take one algebraic step.

Chemical reactions list reactants and products in molar amounts, not just molecular amounts. We can use the coefficients of a balanced chemical equation to relate moles of one substance in the reaction to moles of other substances (stoichiometry). Chemical reactions obey the Law of Conservation of Mass. To balance a chemical reaction, the coefficients in front of each compound can be adjusted until the total number of atoms of each elements is equal on both sides of the reaction arrow.

Collision Theory can be used to described the energetic aspects of chemical reactions. The reactants and products of a chemical reaction store potential energy in the form of chemical bonds and intermolecular forces. The energy difference in the bond energies of the reactants and products is called "Enthalpy". When the products are lower in potential energy than the reactants, then this excess energy is released as heat and the reaction is described as exothermic. When the products are higher in potential energy than the reactants, then energy must be added to the reaction as heat for it to occur and the reaction is described as endothermic.

The rate of a chemical reaction is influenced by its activation energy. The larger the activation energy, the slower the reaction rate. Catalysts can be added to reactions to lower the activation energy and increase the reaction rate.

Not all reactions go to completion. For some reactions, both the forward and reverse reaction can occur simultaneously. When the forward and reverse reaction rates are equal, the reaction is described as being "at equilibrium". Another way to recognize when a system is at equilibrium is when the concentration of reactants and products remain constant. Le Chatlier's principle states that the direction of an equilibrium reaction will shift to reduce the stress.

This page titled 8.S: Quantities in Chemical Reactions (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 6.S: Quantities in Chemical Reactions (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9: SOLIDS, LIQUIDS, AND GASES

Most of us are familiar with the three phases of matter: solid, liquid, and gas. Indeed, we addressed the energy changes involved in phase changes. The substance we are probably most familiar with has those three phases: water. In everyday life, we commonly come in contact with water as a solid (ice), as a liquid, and as a gas (steam). All we have to do is change the conditions of the substance—typically temperature—and we can change the phase from solid to liquid to gas and back again. Under the proper conditions of temperature and pressure, many substances—not only water—can experience the three different phases. An understanding of the phases of matter is important for our understanding of all matter. In this chapter, we will explore the three phases of matter.

9.1: Prelude to Solids, Liquids, and Gases

9.2: Intermolecular Interactions

9.3: Solids and Liquids

9.4: Gases and Pressure

9.5: Gas Laws

9.S: Solids, Liquids, and Gases (Summary)

Template:HideTOC

This page titled 9: Solids, Liquids, and Gases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8: Solids, Liquids, and Gases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.1: Prelude to Solids, Liquids, and Gases

We normally experience carbon dioxide (CO_2) as a gas, but if it were cooled down to about –78°C, it would become a solid. The everyday term for solid carbon dioxide is dry ice.

Why “dry” ice? Solid carbon dioxide is called dry ice because it converts from a solid to a gas directly, without going through the liquid phase, in a process called sublimation. Thus, there is no messy liquid phase to worry about. Although it is a novelty, dry ice has some potential dangers. Because it is so cold, it can freeze living tissues very quickly, so people handling dry ice should wear special protective gloves. The cold carbon dioxide gas is also heavier than air (because it is cold and more dense), so people in the presence of dry ice should be in a well-ventilated area.



Figure 9.1.1: Dry ice. (CC BY SA 3.0 unported; ProjectManhattan).

Dry ice has several common uses. Because it is so cold, it is used as a refrigerant to keep other things cold or frozen (e.g., meats or ice cream). In the medical field, dry ice is used to preserve medical specimens, blood products, and drugs. It also has dermatological applications (e.g., freezing off warts). Organs for transplant are kept cool with dry ice until the recipient of the new organ is ready for surgery. In this respect, carbon dioxide is much like water—more than one phase of the same substance has significant uses in the real world.

This page titled 9.1: Prelude to Solids, Liquids, and Gases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.0: Prelude to Solids, Liquids, and Gases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.2: Intermolecular Interactions

Learning Objectives

- Define *phase*.
- Identify the types of interactions between molecules.

A phase is a certain form of matter that includes a specific set of physical properties. That is, the atoms, the molecules, or the ions that make up the phase do so in a consistent manner throughout the phase. Science recognizes three stable phases: the ***solid*** *phase*, in which individual particles can be thought of as in contact and held in place; the ***liquid*** *phase*, in which individual particles are in contact but moving with respect to each other; and the ***gas*** *phase*, in which individual particles are separated from each other by relatively large distances (see Figure 8.1.1). Not all substances will readily exhibit all phases. For example, carbon dioxide does not exhibit a liquid phase unless the pressure is greater than about six times normal atmospheric pressure. Other substances, especially complex organic molecules, may decompose at higher temperatures, rather than becoming a liquid or a gas.

> For many substances, there are different arrangements the particles can take in the solid phase, depending on temperature and pressure.

Which phase a substance adopts depends on the pressure and the temperature it experiences. Of these two conditions, temperature variations are more obviously related to the phase of a substance. When it is very cold, H_2O exists in the solid form as ice. When it is warmer, the liquid phase of H_2O is present. At even higher temperatures, H_2O boils and becomes steam.

Pressure changes can also affect the presence of a particular phase (as we indicated for carbon dioxide), but its effects are less obvious most of the time. We will mostly focus on the temperature effects on phases, mentioning pressure effects only when they are important. Most chemical substances follow the same pattern of phases when going from a low temperature to a high temperature: the solid phase, then the liquid phase, and then the gas phase. However, the temperatures at which these phases are present differ for all substances and can be rather extreme. Table $9.2.1$ shows the temperature ranges for solid, liquid, and gas phases for three substances. As you can see, there is extreme variability in the temperature ranges.

Table $9.2.1$: Temperature Ranges for the Three Phases of Various Substances

Substance	Solid Phase Below	Liquid Phase Above	Gas Phase Above
hydrogen (H_2)	–259°C	–259°C	–253°C
water (H_2O)	0°C	0°C	100°C
sodium chloride (NaCl)	801°C	801°C	1413°C

> The *melting point* of a substance is the temperature that separates a solid and a liquid. The *boiling point* of a substance is the temperature that separates a liquid and a gas.

What accounts for this variability? Why do some substances become liquids at very low temperatures, while others require very high temperatures before they become liquids? It all depends on the strength of the **intermolecular forces (IMF)** between the **particles** of substances and the **kinetic energies** (KE) of its molecules. (Although ionic compounds are not composed of discrete molecules, we will still use the term *intermolecular* to include interactions between the ions in such compounds.) Substances that experience **strong intermolecular interactions** require **higher temperatures to become liquids and, finally, gases.** Substances that experience **weak intermolecular interactions** do not need much energy (as measured by temperature) to become liquids and gases and will exhibit these phases at **lower temperatures.**




Figure 9.2.1: Transitions between solid, liquid, and gaseous states of a substance occur when conditions of temperature or pressure favor the associated changes in intermolecular forces. (Note: The space between particles in the gas phase is much greater than shown.) (CC BY 4.0; OpenStax)

Covalent Network Materials

Substances with the highest melting and boiling points have **covalent network** bonding. This type of intermolecular interaction is actually a covalent bond. In these substances, all the atoms in a sample are covalently bonded to one another; in effect, the entire sample is essentially **one giant molecule**. Many of these substances are solid over a large temperature range because it takes a lot of energy to disrupt all the covalent bonds at once. One example of a substance that shows covalent network bonding is diamond (Figure 9.2.2). Diamond is composed entirely of carbon atoms, each bonded to four other carbon atoms in a tetrahedral geometry. Melting a covalent network solid is not accomplished by overcoming the relatively weak intermolecular forces. Rather, all of the covalent bonds must be broken, a process that requires extremely high temperatures. Diamond, in fact, does not melt at all. Instead, it vaporizes to a gas at temperatures above 3,500°C. Diamond is extremely hard and is one of the few materials that can cut glass.




Figure 9.2.2: Diamond. Diamond, a form of pure carbon, has covalent network bonding. It takes a very high temperature—over 3,500°C—for diamond to leave the solid state. Source: Photo © Thinkstock

Ionic Compounds

The strongest force between any two particles is the ionic bond, in which two ions of opposing charge are attracted to each other. Thus, ionic interactions between particles are another type of intermolecular interaction. Substances that contain ionic

interactions are relatively strongly held together, so these substances typically have high melting and boiling points. Sodium chloride (Figure 9.2.3) is an example of a substance whose particles experience ionic interactions (Table 9.2.1). These attractive forces are sometimes referred to as ion-ion interactions.




Figure 9.2.3: Sodium Chloride. Solid NaCl is held together by ionic interactions. Source: Photo © Thinkstock

Covalent Molecular Compounds

There are two different covalent structures: molecular and network. Covalent network compounds like SiO_2 (quartz) have structures of atoms in a network like diamond described earlier. In this section, we are dealing with the molecular type that contains individual molecules. The bonding between atoms in the individual molecule is covalent but the attractive forces between the molecules are called ***intermolecular forces (IMF)***.

In contrast to *intramolecular* forces (see Figure 8.1.4), such as the covalent bonds that hold atoms together in molecules and polyatomic ions, intermolecular forces hold molecules together in a **liquid or solid**. Intermolecular forces are generally much weaker than covalent bonds. For example, it requires 927 kJ to overcome the intramolecular forces and break both O–H bonds in 1 mol of water, but it takes only about 41 kJ to overcome the intermolecular attractions and convert 1 mol of liquid water to water vapor at 100°C. (Despite this seemingly low value, the intermolecular forces in liquid water are among the strongest such forces known!) Given the large difference in the strengths of intra- and intermolecular forces, changes between the solid, liquid, and gaseous states almost invariably occur for molecular substances ***without breaking covalent bonds*.**




Figure 9.2.4: **Intramolecular** forces keep a molecule intact. **Intermolecular** forces hold multiple molecules together and determine many of a substance's properties. (CC BY 4.0; OpenStax)

In this section, we will discuss the three types of IMF in molecular compounds: dipole-dipole, hydrogen bonding and London dispersion forces.

Dipole-dipole Intermolecular Forces

As discussed in the section on covalent molecules, covalent bond that has an unequal sharing of electrons is called a **polar covalent** bond. (A covalent bond that has an equal sharing of electrons, as in a covalent bond with the same atom on each side, is called a **nonpolar covalent** bond.) A molecule with a net unequal distribution of electrons in its covalent bonds is a polar molecule. HF is an example of a polar molecule (see Figure 8.1.5).




Figure 9.2.5: Polar Covalent Bonds. The electrons in the HF molecule are not equally shared by the two atoms in the bond. Because the fluorine atom has nine protons in its nucleus, it attracts the negatively charged electrons in the bond more than the hydrogen atom does with its one proton in its nucleus. Thus, electrons are more strongly attracted to the fluorine atom, leading to an imbalance in the electron distribution between the atoms. The fluorine side of the bond picks up a partial overall negative charge (represented by the δ– in the diagram), while the hydrogen side of the bond has an overall partial positive charge (represented by the δ+ in the diagram). Such a bond is called a *polar covalent bond*.

The charge separation in a **polar covalent bond** is not as extreme as is found in ionic compounds, but there is a related result: oppositely charged ends of different molecules will attract each other. This type of intermolecular interaction is called a **dipole-dipole interaction**. Many molecules with polar covalent bonds experience dipole-dipole interactions.

The covalent bonds in some molecules are oriented in space in such a way that the bonds in the molecules cancel each other out. The individual bonds are polar, but due to molecular symmetry, the overall molecule is not polar; rather, the molecule is ***nonpolar*.** Such molecules experience little or no dipole-dipole interactions. Carbon dioxide (CO_2) and carbon tetrachloride (CCl_4) are examples of such molecules (Figure 9.2.6).




Figure 9.2.6: Nonpolar Molecules. Although the individual bonds in both CO_2 and CCl_4 are polar, their effects cancel out because of the spatial orientation of the bonds in each molecule. As a result, such molecules experience little or no dipole-dipole interaction.

Recall from the previous sections on chemical bonding and molecular geometry that *polar* molecules have a partial positive charge on one side and a partial negative charge on the other side of the molecule—a separation of charge called a ***dipole*.** Consider a polar molecule such as hydrogen chloride, HCl. In the HCl molecule, the more electronegative Cl atom bears the partial negative charge, whereas the less electronegative H atom bears the partial positive charge. An attractive force between HCl molecules results from the attraction between the positive end of one HCl molecule and the negative end of another. This attractive force is called a dipole-dipole attraction—the electrostatic force between the partially positive end of one polar molecule and the partially negative end of another, as illustrated in Figure 9.2.7.




Figure 9.2.7: This image shows two arrangements of polar molecules, such as HCl, that allow an attraction between the partial negative end of one molecule and the partial positive end of another.

The effect of a dipole-dipole attraction is apparent when we compare the properties of HCl molecules to nonpolar F_2 molecules. Both HCl and F_2 consist of the same number of atoms and have approximately the same molecular mass. At a temperature of 150 K, molecules of both substances would have the same average kinetic energy. However, the dipole-dipole attractions between HCl molecules are sufficient to cause them to "stick together" to form a liquid, whereas the relatively weaker dispersion forces between nonpolar F_2 molecules are not, and so this substance is gaseous at this temperature. The higher normal boiling point of HCl (188 K) compared to F_2 (85 K) is a reflection of the greater strength of dipole-dipole attractions between HCl molecules, compared to the attractions between nonpolar F_2 molecules. We will often use values such as boiling or freezing points as indicators of the relative strengths of IMFs of attraction present within different substances.

Example 9.2.1

Predict which will have the higher boiling point: N_2 or CO. Explain your reasoning.

Solution

CO and N_2 are both diatomic molecules with masses of about 28 amu, so they experience similar London dispersion forces. Because CO is a polar molecule, it experiences dipole-dipole attractions. Because N_2 is nonpolar, its molecules cannot exhibit dipole-dipole attractions. The dipole-dipole attractions between CO molecules are comparably stronger than the dispersion forces between nonpolar N_2 molecules, so CO is expected to have the higher boiling point.

Exercise 9.2.1

Predict which will have the higher boiling point: ICl or $\mathrm{Br_2}$. Explain your reasoning.

Answer available at the end of this section.

Hydrogen Bonding Intermolecular Forces

Molecules with **hydrogen atoms** bonded to electronegative atoms such as **O, N, and F** tend to exhibit unusually strong intermolecular interactions due to a particularly strong type of dipole-dipole attraction called **hydrogen bonding**. The very large difference in electronegativity between the H atom (2.1) and the atom to which it is bonded (4.0 for an F atom, 3.5 for an O atom, or 3.0 for a N atom), combined with the very small size of a H atom and the relatively small sizes of F, O, or N atoms, leads to ***highly concentrated partial charges*** with these atoms. Because the hydrogen atom does not have any electrons other than the ones in the covalent bond, its positively charged nucleus is almost completely exposed, allowing strong attractions to other nearby lone pairs of electrons.

Examples of hydrogen bonds include HF···HF, H_2O···HOH, and H_3N···HNH_2, in which the **hydrogen bonds** are denoted by **dots**. Figure 9.2.8 illustrates hydrogen bonding between water molecules. The physical properties of water, which has two O–H bonds, are strongly affected by the presence of hydrogen bonding between water molecules. Most molecular compounds that have a mass similar to water are gases at room temperature. However, because of the strong hydrogen bonds, water molecules are able to stay condensed in the liquid state.




Figure 9.2.8: Water molecules participate in multiple hydrogen-bonding interactions with nearby water molecules.
Five water molecules are shown near one another, but not touching. A dotted line lies between many of the hydrogen atoms on one molecule and the oxygen atom on another molecule.

A **hydrogen bond** is an intermolecular attractive force in which *a* **hydrogen atom**, that is covalently bonded to a small, highly electronegative atom, is attracted to a **lone pair of electrons** on an atom in a neighboring molecule. Figure 9.2.9 shows how methanol (CH_3OH) molecules experience hydrogen bonding. Methanol contains both a hydrogen atom attached to O; methanol can thus form hydrogen bonds by acting as either a hydrogen bond donor or a hydrogen bond acceptor (lone pair). The hydrogen-bonded structure of methanol is as follows:




Figure 9.2.9: Methanol molecules participate in multiple hydrogen-bonding interactions with nearby water molecules.

Despite use of the word “bond,” keep in mind that hydrogen bonds are *intermolecular* attractive forces, not *intramolecular* attractive forces (covalent bonds). Hydrogen bonds are much weaker than covalent bonds, but are generally much stronger than other dipole-dipole attractions and dispersion forces.

Example \({9{.}2{.}\eatSpaces{2\)

Consider the compounds dimethylether (CH_3OCH_3), ethanol (CH_3CH_2OH), and propane ($CH_3CH_2CH_3$). Their boiling points, not necessarily in order, are –42.1 °C, –24.8 °C, and 78.4 °C. Match each compound with its boiling point. Explain your reasoning.

Solution

The shapes of CH_3OCH_3, CH_3CH_2OH, and $CH_3CH_2CH_3$ are similar, as are their molar masses (46 g/mol, 46 g/mol, and 44 g/mol, respectively), so they will exhibit similar dispersion forces. Since $CH_3CH_2CH_3$ is nonpolar, it may exhibit *only* dispersion forces. Because CH_3OCH_3 is polar, it will also experience dipole-dipole attractions. Finally, CH_3CH_2OH has an –OH group, and so it will experience the uniquely strong dipole-dipole attraction known as hydrogen bonding. So the ordering in terms of strength of IMFs, and thus boiling points, is $CH_3CH_2CH_3 < CH_3OCH_3 < CH_3CH_2OH$. The boiling point of propane is –42.1 °C, the boiling point of dimethylether is –24.8 °C, and the boiling point of ethanol is 78.5 °C.

Exercise Extra close brace or missing open brace

Ethane (CH_3CH_3) has a melting point of –183 °C and a boiling point of –89 °C. Predict the melting and boiling points for methylamine (CH_3NH_2). Explain your reasoning.

Answer available at the end of this section.

Hydrogen Bonding and DNA

Deoxyribonucleic acid (DNA) is found in every living organism and contains the genetic information that determines the organism's characteristics, provides the blueprint for making the proteins necessary for life, and serves as a template to pass this information on to the organism's offspring. A DNA molecule consists of two (anti-)parallel chains of repeating nucleotides, which form its well-known double helical structure, as shown in Figure 9.2.10.




Figure 9.2.10: Two separate DNA molecules form a double-stranded helix in which the molecules are held together via hydrogen bonding. (credit: modification of work by Jerome Walker, Dennis Myts)

Each nucleotide contains a (deoxyribose) sugar bound to a phosphate group on one side, and one of four nitrogenous bases on the other. Two of the bases, cytosine (C) and thymine (T), are single-ringed structures known as pyrimidines. The other two, adenine (A) and guanine (G), are double-ringed structures called purines. These bases form complementary base pairs consisting of one purine and one pyrimidine, with adenine pairing with thymine, and cytosine with guanine. Each base pair is held together by

hydrogen bonding. A and T share two hydrogen bonds, C and G share three, and both pairings have a similar shape and structure Figure 9.2.11




Figure 9.2.11: The geometries of the base molecules result in maximum hydrogen bonding between adenine and thymine (AT) and between guanine and cytosine (GC), so-called “complementary base pairs.”

The cumulative effect of millions of hydrogen bonds effectively holds the two strands of DNA together. Importantly, the two strands of DNA can relatively easily “unzip” down the middle since hydrogen bonds are relatively weak compared to the covalent bonds that hold the atoms of the individual DNA molecules together. This allows both strands to function as a template for replication.

London Dispersion Forces

Finally, there are forces between all molecules that are caused by electrons being in different places in a molecule at any one time, which sets up a temporary separation of charge that disappears almost as soon as it appears. These are very weak intermolecular interactions and are called **dispersion forces (or London forces)**. (An alternate name is London dispersion forces.) Molecules that experience no other type of intermolecular interaction will at least experience dispersion forces. Substances that experience only dispersion forces are typically soft in the solid phase and have relatively low melting points. Examples include *waxes*, which are long hydrocarbon chains that are solids at room temperature because the molecules have so many electrons. The resulting dispersion forces between these molecules make them assume the solid phase at normal temperatures.

Dispersion forces that develop between atoms in different molecules can attract the two molecules to each other. The forces are relatively weak, however, and become significant only when the molecules are very close. **Larger and heavier atoms and molecules** exhibit **stronger dispersion forces** than do smaller and lighter atoms and molecules. F_2 and Cl_2 are gases at room temperature (reflecting weaker attractive forces); Br_2 is a liquid, and I_2 is a solid (reflecting stronger attractive forces). Trends in observed melting and boiling points for the halogens clearly demonstrate this effect, as seen in Table 8.1.2.

Table 9.2.2: ***Melting and Boiling Points of the Halogens***

Halogen	Molar Mass	Atomic Radius	Melting Point	Boiling Point
fluorine, F_2	38 g/mol	72 pm	53 K	85 K
chlorine, Cl_2	71 g/mol	99 pm	172 K	238 K
bromine, Br_2	160 g/mol	114 pm	266 K	332 K
iodine, I_2	254 g/mol	133 pm	387 K	457 K
astatine, At_2	420 g/mol	150 pm	575 K	610 K

The increase in melting and boiling points with increasing atomic/molecular size may be rationalized by considering how the

strength of dispersion forces is affected by the electronic structure of the atoms or molecules in the substance. In a larger atom, the valence electrons are, on average, farther from the nuclei than in a smaller atom. Thus, they are less tightly held and can more easily form the temporary dipoles that produce the attraction. The measure of how easy or difficult it is for another electrostatic charge (for example, a nearby ion or polar molecule) to distort a molecule's charge distribution (its electron cloud) is known as **polarizability**. A molecule that has a charge cloud that is easily distorted is said to be very polarizable and will have large dispersion forces; one with a charge cloud that is difficult to distort is not very polarizable and will have small dispersion forces.

Example 9.2.3

Order the following compounds of a group 14 element and hydrogen from lowest to highest boiling point: CH_4, SiH_4, GeH_4, and SnH_4. Explain your reasoning.

Solution

Applying the skills acquired in the chapter on chemical bonding and molecular geometry, all of these compounds are predicted to be nonpolar, so they may experience only dispersion forces: the smaller the molecule, the less polarizable and the weaker the dispersion forces; the larger the molecule, the larger the dispersion forces. The molar masses of CH_4, SiH_4, GeH_4, and SnH_4 are approximately 16 g/mol, 32 g/mol, 77 g/mol, and 123 g/mol, respectively. Therefore, CH_4 is expected to have the lowest boiling point and SnH_4 the highest boiling point. The ordering from lowest to highest boiling point is expected to be

$CH_4 < SiH_4 < GeH_4 < SnH_4$

A graph of the actual boiling points of these compounds versus the period of the group 14 elementsshows this prediction to be correct:




Figure 9.2.11: Graph showing the relationship between temperature (°C) and time (minutes), with a positive trend indicating increasing temperature.

Exercise 9.2.3

Order the following hydrocarbons from lowest to highest boiling point: C_2H_6, C_3H_8, and C_4H_{10}.

Answer available at the end of this section.

Answer available at the end of this section.

Applications: Geckos and Intermolecular Forces

Geckos have an amazing ability to adhere to most surfaces. They can quickly run up smooth walls and across ceilings that have no toe-holds, and they do this without having suction cups or a sticky substance on their toes. And while a gecko can lift its feet easily as it walks along a surface, if you attempt to pick it up, it sticks to the surface. How are geckos (as well as spiders and some other insects) able to do this? Although this phenomenon has been investigated for hundreds of years, scientists only recently uncovered the details of the process that allows geckos' feet to behave this way.




Figure 9.2.12: Hierarchical magnification of a gecko's toe, showing the branching of microscopic **setae** into thousands of tiny **spatulae** that facilitate surface adhesion.

Geckos' toes are covered with hundreds of thousands of tiny hairs known as *setae*, with each seta, in turn, branching into hundreds of tiny, flat, triangular tips called *spatulae*. The huge numbers of spatulae on its setae provide a gecko, shown in Figure 8.1.12, with a large total surface area for sticking to a surface. In 2000, Kellar Autumn, who leads a multi-institutional gecko research team, found that geckos adhered equally well to both polar silicon dioxide and nonpolar gallium arsenide. This proved that geckos stick to surfaces because of dispersion forces—weak intermolecular attractions arising from temporary, synchronized charge distributions between adjacent molecules. Although dispersion forces are very weak, the total attraction over millions of spatulae is large enough to support many times the gecko's weight.

In 2014, two scientists developed a model to explain how geckos can rapidly transition from "sticky" to "non-sticky." Alex Greaney and Congcong Hu at Oregon State University described how geckos can achieve this by changing the angle between their spatulae and the surface. Geckos' feet, which are normally nonsticky, become sticky when a small shear force is applied. By curling and uncurling their toes, geckos can alternate between sticking and unsticking from a surface, and thus easily move across it. Further investigations may eventually lead to the development of better adhesives and other applications.

Boiling Points and Bonding Types

In order for a substance to enter the gas phase, its particles must completely overcome the intermolecular forces holding them together. Therefore, a comparison of boiling points is essentially equivalent to comparing the strengths of the attractive intermolecular forces exhibited by the individual molecules. For small molecular compounds, London dispersion forces are the weakest intermolecular forces. Dipole-dipole forces are somewhat stronger, and hydrogen bonding is a particularly strong form of dipole-dipole interaction. However, when the mass of a nonpolar molecule is sufficiently large, its dispersion forces can be stronger than the dipole-dipole forces in a lighter polar molecule. Thus, nonpolar Cl_2 has a higher boiling point than polar HCl.

Table 9.2.3: Intermolecular Forces and Boiling Points

Substance	Strongest Intermolecular Force	Boiling Point (°C)
H_2	dispersion	-253
Ne	dispersion	-246

Substance	Strongest Intermolecular Force	Boiling Point (°C)
O_2	dispersion	-183
Cl_2	dispersion	-34
HCl	dipole-dipole	-85
HBr	dipole-dipole	-66
H_2S	dipole-dipole	-61
NH_3	hydrogen bonding	-33
HF	hydrogen bonding	20
H_2O	hydrogen bonding	100

Example 9.2.4: Intermolecular Forces

What intermolecular forces besides dispersion forces, if any, exist in each substance? Are any of these substances solids at room temperature?

a. potassium chloride (KCl)
b. ethanol (C_2H_5OH)
c. bromine (Br_2)

Solution

a. Potassium chloride is composed of ions, so the intermolecular interaction in potassium chloride is ionic forces. Because ionic interactions are strong, it might be expected that potassium chloride is a solid at room temperature.
b. Ethanol has a hydrogen atom attached to an oxygen atom, so it would experience hydrogen bonding. If the hydrogen bonding is strong enough, ethanol might be a solid at room temperature, but it is difficult to know for certain. (Ethanol is actually a liquid at room temperature.)
c. Elemental bromine has two bromine atoms covalently bonded to each other. Because the atoms on either side of the covalent bond are the same, the electrons in the covalent bond are shared equally, and the bond is a nonpolar covalent bond. Thus, diatomic bromine does not have any intermolecular forces other than dispersion forces. It is unlikely to be a solid at room temperature unless the dispersion forces are strong enough. Bromine is a liquid at room temperature.

Exercise 9.2.4

What intermolecular forces besides dispersion forces, if any, exist in each substance? Are any of these substances solids at room temperature?

a. methylamine (CH_3NH_2)
b. calcium sulfate ($CaSO_4$)
c. carbon monoxide (CO)

Answer available at the end of this section.

Key Takeaways

- A phase is a form of matter that has the same physical properties throughout.
- Molecules interact with each other through various forces: dipole-dipole interactions, hydrogen bonding, and dispersion forces.
- Dipole-dipole attractions result from the electrostatic attraction of the partial negative end of one dipolar molecule for the partial positive end of another.
- Hydrogen bonds are a special type of dipole-dipole attraction that results when hydrogen is bonded to one of the three most

electronegative elements: F, O, or N.
- The temporary dipole that results from the motion of the electrons in an atom can induce a dipole in an adjacent atom and give rise to the London dispersion force. London forces increase with increasing molecular size.

Answers

Exercise 9.2.1

ICl. ICl and Br_2 have similar masses (~160 amu) and therefore experience similar London dispersion forces. ICl is polar and thus also exhibits dipole-dipole attractions; Br_2 is nonpolar and does not. The relatively stronger dipole-dipole attractions require more energy to overcome, so ICl will have the higher boiling point.

Exercise 9.2.2

The melting point and boiling point for methylamine are predicted to be significantly greater than those of ethane. CH_3CH_3 and CH_3NH_2 are similar in size and mass, but methylamine possesses an –NH group and therefore may exhibit hydrogen bonding. This greatly increases its IMFs, and therefore its melting and boiling points. It is difficult to predict values, but the known values are a melting point of –93 °C and a boiling point of –6 °C.

Exercise 9.2.3

Exercise 9.2.3

All of these compounds are nonpolar and only have London dispersion forces: the larger the molecule, the larger the dispersion forces and the higher the boiling point. The ordering from lowest to highest boiling point is therefore

$$C_2H_6 < C_3H_8 < C_4H_{10}.$$

Exercise 9.2.4

a. dipole-dipole, hydrogen bonding

b. ionic forces (solid at room temperature)

c. dipole-dipole

This page titled 9.2: Intermolecular Interactions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.1: Intermolecular Interactions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.3: Solids and Liquids

Learning Objectives

- To describe the solid and liquid phases.

Solids and liquids are collectively called *condensed phases* because their particles are in virtual contact. The two states share little else, however.

Solids

In the solid state, the individual particles of a substance are in fixed positions with respect to each other because there is not enough thermal energy to overcome the intermolecular interactions between the particles. As a result, solids have a definite shape and volume. Most solids are hard, but some (like waxes) are relatively soft. Many solids composed of ions can also be quite brittle.



Figure 9.3.1: Crystalline Arrangement of Quartz crystal cluster. Some large crystals look the way they do because of the regular arrangement of atoms (ions) in their crystal structure. from Wikipedia.

Solids usually have their constituent particles arranged in a regular, three-dimensional array of alternating positive and negative ions called a crystal. The effect of this regular arrangement of particles is sometimes visible macroscopically, as shown in Figure 9.3.1. Some solids, especially those composed of large molecules, cannot easily organize their particles in such regular crystals and exist as amorphous (literally, “without form”) solids. Glass is one example of an amorphous solid.

Liquids

If the particles of a substance have enough energy to partially overcome intermolecular interactions, then the particles can move about each other while remaining in contact. This describes the liquid state. In a liquid, the particles are still in close contact, so liquids have a definite volume. However, because the particles can move about each other rather freely, a liquid has no definite shape and takes a shape dictated by its container.



Figure 9.3.2: The formation of a spherical droplet of liquid water minimizes the surface area, which is the natural result of surface tension in liquids. from Wikipedia.

Gases

If the particles of a substance have enough energy to completely overcome intermolecular interactions, then the particles can separate from each other and move about randomly in space. Like liquids, gases have no definite shape, but unlike solids and liquids, gases have no definite volume either.




Figure 9.3.3: A Representation of the Solid, Liquid, and Gas States. A solid has definite volume and shape, a liquid has a definite volume but no definite shape, and a gas has neither a definite volume nor shape.

The change from solid to liquid usually does not significantly change the volume of a substance. However, the change from a liquid to a gas significantly increases the volume of a substance, by a factor of 1,000 or more. Figure 9.3.3 shows the differences among solids, liquids, and gases at the molecular level, while Table 9.3.1 lists the different characteristics of these states.

Table 9.3.1: Characteristics of the Three States of Matter

Characteristic	Solid	Liquid	Gas
shape	definite	indefinite	indefinite
volume	definite	definite	indefinite
relative intermolecular interaction strength	strong	moderate	weak

Characteristic	Solid	Liquid	Gas
relative particle positions	in contact and fixed in place	in contact but not fixed	not in contact, random positions

Example 9.3.1

What state or states of matter does each statement, describe?

a. This state has a definite volume.
b. This state has no definite shape.
c. This state allows the individual particles to move about while remaining in contact.

Solution

a. This statement describes either the liquid state or the solid state.
b. This statement describes either the liquid state or the gas state.
c. This statement describes the liquid state.

Exercise 9.3.1

What state or states of matter does each statement describe?

a. This state has individual particles in a fixed position with regard to each other.
b. This state has individual particles far apart from each other in space.
c. This state has a definite shape.

Answer available at the end of this section.

Looking Closer: Water, the Most Important Liquid

Earth is the only known body in our solar system that has liquid water existing freely on its surface. That is a good thing because life on Earth would not be possible without the presence of liquid water.

Water has several properties that make it a unique substance among substances. It is an excellent solvent; it dissolves many other substances and allows those substances to react when in solution. In fact, water is sometimes called the *universal solvent* because of this ability. Water has unusually high melting and boiling points (0°C and 100°C, respectively) for such a small molecule. The boiling points for similar-sized molecules, such as methane (BP = –162°C) and ammonia (BP = –33°C), are more than 100° lower. Though a liquid at normal temperatures, water molecules experience a relatively strong intermolecular interaction that allows them to maintain the liquid phase at higher temperatures than expected.

Unlike most substances, the solid form of water is less dense than its liquid form, which allows ice to float on water. The most energetically favorable configuration of H_2O molecules is one in which each molecule is hydrogen-bonded to four neighboring molecules. Owing to the thermal motions, this ideal is never achieved in the liquid, but when water freezes to ice, the molecules settle into exactly this kind of an arrangement in the ice crystal. This arrangement requires that the molecules be somewhat farther apart than would otherwise be the case; as a consequence, ice, in which hydrogen bonding is at its maximum, has a more open structure, and thus a lower density than water.




Figure 9.3.4: Three-dimensional views of a typical local structure of liquid water (left) and ice (right).

Here are three-dimensional views of a typical local structure of water (left) and ice (right.) Notice the greater openness of the ice structure which is necessary to ensure the strongest degree of hydrogen bonding in a uniform, extended crystal lattice. The structure of liquid water is very similar, but in the liquid, the hydrogen bonds are continually broken and formed because of rapid molecular motion. Because ice is less dense than liquid water, rivers, lakes, and oceans freeze from the top down. In fact, the ice forms a protective surface layer that insulates the rest of the water, allowing fish and other organisms to survive in the lower levels of a frozen lake or sea. If ice were denser than the liquid, the ice formed at the surface in cold weather would sink as fast as it formed. Bodies of water would freeze from the bottom up, which would be lethal for most aquatic creatures. The expansion of water when freezing also explains why automobile or boat engines must be protected by "antifreeze" and why unprotected pipes in houses break if they are allowed to freeze.

Water also requires an unusually large amount of energy to change temperature. While 100 J of energy will change the temperature of 1 g of Fe by 230°C, this same amount of energy will change the temperature of 1 g of H_2O by only 100°C. Thus, water changes its temperature slowly as heat is added or removed. This has a major impact on weather, as storm systems like hurricanes can be impacted by the amount of heat that ocean water can store.

Water's influence on the world around us is affected by these properties. Isn't it fascinating that such a small molecule can have such a big impact?

Key Takeaway

- Solids and liquids are phases that have their own unique properties.

Answers

Exercise 9.3.1

a. solid
b. gas
c. solid

This page titled 9.3: Solids and Liquids is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.2: Solids and Liquids by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.4: Gases and Pressure

Learning Objectives

- To describe the gas phase.

The gas phase is unique among the three states of matter in that there are some simple models we can use to predict the physical behavior of all gases—independent of their identities. We cannot do this for the solid and liquid states. In fact, the development of this understanding of the behavior of gases represents the historical dividing point between alchemy and modern chemistry. Initial advances in the understanding of gas behavior were made in the mid 1600s by Robert Boyle, an English scientist who founded the Royal Society (one of the world's oldest scientific organizations).

Kinetic Theory of Gases

How is it that we can model all gases independent of their chemical identity? The answer is in a group of statements called the kinetic theory of gases:

- Gases are composed of tiny particles that are separated by large distances.
- Gas particles are constantly moving, experiencing collisions with other gas particles and the walls of their container.
- The velocity of gas particles is related to the temperature of a gas.
- Gas particles do not experience any force of attraction or repulsion with each other.

Did you notice that none of these statements relates to the identity of the gas? This means that all gases should behave similarly. A gas that follows these statements perfectly is called an *ideal gas*. Most gases show slight deviations from these statements and are called *real gases*. However, the existence of real gases does not diminish the importance of the kinetic theory of gases.

Gas Pressure

One of the statements of the kinetic theory mentions collisions. As gas particles are constantly moving, they are also constantly colliding with each other and with the walls of their container. There are forces involved as gas particles bounce off the container walls (Figure 9.4.1). The force generated by gas particles divided by the area of the container walls yields pressure. Pressure is a property we can measure for a gas, but we typically do not consider pressure for solids or liquids.




Figure 9.4.1: Gas Pressure. Pressure is what results when gas particles rebound off the walls of their container.

The basic unit of pressure is the newton per square meter (N/m^2). This combined unit is redefined as a pascal (Pa). One pascal is not a very large amount of pressure. A more useful unit of pressure is the bar, which is 100,000 Pa (1 bar = 100,000 Pa). Other common units of pressure are the atmosphere (atm), which was originally defined as the average pressure of Earth's atmosphere at sea level; and mmHg (millimeters of mercury), which is the pressure generated by a column of mercury 1 mm high. The unit millimeters of mercury is also called a torr, named after the Italian scientist Evangelista Torricelli, who invented the barometer in the mid-1600s. A more precise definition of atmosphere, in terms of torr, is that there are exactly 760 torr in 1 atm. A bar equals 1.01325 atm. Given all the relationships between these pressure units, the ability to convert from one pressure unit to another is a useful skill.

Example 9.4.1: Converting Pressures

Write a conversion factor to determine how many atmospheres are in 1,547 mmHg.

Solution

Because 1 mmHg equals 1 torr, the given pressure is also equal to 1,547 torr. Because there are 760 torr in 1 atm, we can use this conversion factor to do the mathematical conversion:

$$1,547 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 2.04 \text{ atm}$$

Note how the torr units cancel algebraically.

Exercise 9.4.1: Converting Pressures

Write a conversion factor to determine how many millimeters of mercury are in 9.65 atm.

Answer available at the end of this section.

The kinetic theory also states that there is no interaction between individual gas particles. Although we know that there are, in fact, intermolecular interactions in real gases, the kinetic theory assumes that gas particles are so far apart that the individual particles don't "feel" each other. Thus, we can treat gas particles as tiny bits of matter whose identity isn't important to certain physical properties.

Key Takeaway

- The gas phase has certain general properties characteristic of that phase.

Answers

Exercise 9.4.1: Converting Pressures

$$9.65 \text{ atm} \times \frac{760 \text{ mmHg}}{1 \text{ atm}} = 7,334 \text{ mmHg}.$$

This page titled 9.4: Gases and Pressure is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.3: Gases and Pressure by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.5: Gas Laws

Learning Objectives

- To predict the properties of gases using the gas laws.

Gas Properties

Experience has shown that several properties of a gas can be related to each other under certain conditions. The properties are pressure (*P*), volume (*V*), temperature (*T*, in kelvins), and amount of material expressed in moles (*n*). What we find is that a sample of gas cannot have any random values for these properties. Instead, only certain values, dictated by some simple mathematical relationships, will occur.

Boyle's Law

The first simple relationship, referred to as a gas law, is between the pressure of a gas and its volume. If the amount of gas in a sample and its temperature are kept constant, then as the pressure of a gas is increased, the volume of the gas decreases proportionately. Mathematically, this is written as

$$\mathrm{P} \propto \frac{1}{\mathrm{V}}$$

where the "$\propto$" symbol means "is proportional to." This is one form of Boyle's law, which relates the pressure of a gas to its volume.

A more useful form of Boyle's law involves a change in conditions of a gas. For a given amount of gas at a constant temperature, if we know the initial pressure and volume of a gas sample and the pressure or volume changes, we can calculate what the new volume or pressure will be. That form of Boyle's law is written

$$P_i V_i = P_f V_f \tag{9.5.1}$$

where the subscript i refers to initial conditions and the subscript f refers to final conditions.

To use 9.5.1, you need to know any three of the variables so that you can algebraically calculate the fourth variable. Also, the pressure quantities must have the same units, as must the two volume quantities. If the two similar variables don't have the same variables, one value must be converted to the other value's unit.

Example 9.5.1: Increasing Pressure in a Gas

What happens to the volume of a gas if its pressure is increased? Assume all other conditions remain the same.

Solution

If the pressure of a gas is increased, the volume decreases in response.

Exercise 9.5.1: Increasing Volume in a Gas

What happens to the pressure of a gas if its volume is increased? Assume all other conditions remain the same.

Answer available at the end of this section.

Example 9.5.2: Gas Compression

If a sample of gas has an initial pressure of 1.56 atm and an initial volume of 7.02 L, what is the final volume if the pressure is reduced to 0.987 atm? Assume that the amount and the temperature of the gas remain constant.

Solution

The key in problems like this is to be able to identify which quantities represent which variables from the relevant equation. The way the question is worded, you should be able to tell that 1.56 atm is P_i, 7.02 L is V_i, and 0.987 atm is P_f. What we are looking for is the final volume—V_f. Therefore, substituting these values into $P_iV_i = P_fV_f$:

$$(1.56 \text{ atm})(7.02 \text{ L}) = (0.987 \text{ atm}) \times V_f$$

The expression has atmospheres on both sides of the equation, so they cancel algebraically:

$$(1.56)(7.02 \text{ L}) = (0.987) \times V_f$$

Now we divide both sides of the expression by 0.987 to isolate V_f, the quantity we are seeking:

$$\frac{(1.56)(7.02 \text{ L})}{0.987} = \text{V}_\text{f}$$

Performing the multiplication and division, we get the value of V_f, which is 11.1 L. The volume increases. This should make sense because the pressure decreases, so pressure and volume are inversely related.

Exercise 9.5.2

If a sample of gas has an initial pressure of 3.66 atm and an initial volume of 11.8 L, what is the final pressure if the volume is reduced to 5.09 L? Assume that the amount and the temperature of the gas remain constant.

Answer available at the end of this section.

If the units of similar quantities are not the same, one of them must be converted to the other quantity's units for the calculation to work out properly. It does not matter which quantity is converted to a different unit; the only thing that matters is that the conversion and subsequent algebra are performed properly. The following example illustrates this process.

Example 9.5.3

If a sample of gas has an initial pressure of 1.56 atm and an initial volume of 7.02 L, what is the final volume if the pressure is changed to 1,775 torr? Does the answer make sense? Assume that the amount and the temperature of the gas remain constant.

Solution

This example is similar to Example 9.5.2, except now the final pressure is expressed in torr. For the math to work out properly, one of the pressure values must be converted to the other unit. Let us change the initial pressure to torr:

$$1.56 \text{ atm} \times \frac{760 \text{ torr}}{1 \text{ atm}} = 1,190 \text{ torr}$$

Now we can use Boyle's law:

$$(1,190 \text{ torr})(7.02 \text{ L}) = (1,775 \text{ torr}) \times V_f$$

Torr cancels algebraically from both sides of the equation, leaving

$$(1{,}190)(7.02\ \text{L}) = (1{,}775) \times V_f$$

Now we divide both sides of the equation by 1,775 to isolate V_f on one side. Solving for the final volume,

$$\text{V}_\text{f} = \frac{(1,190)(7.02\ \text{L})}{1,775} = 4.71\ \text{L}$$

Because the pressure increases, it makes sense that the volume decreases.

The answer for the final volume is essentially the same if we converted the 1,775 torr to atmospheres: $1,775\ \text{torr} \times \frac{1\ \text{atm}}{760\ \text{torr}} = 2.336\ \text{atm}$. Using Boyle's law: (1.56 atm)(7.02 L) = (2.335 atm) × V_f; $\text{V}_\text{f} = \frac{(1.56\ \text{atm})(7.02\ \text{L})}{2.336\ \text{atm}} = 4.69\ \text{L}$.

Exercise 9.5.3

If a sample of gas has an initial pressure of 375 torr and an initial volume of 7.02 L, what is the final pressure if the volume is changed to 4,577 mL? Does the answer make sense? Assume that amount and the temperature of the gas remain constant.

Answer available at the end of this section.

To Your Health: Breathing

Breathing certainly is a major contribution to your health! Without breathing, we could not survive. Curiously, the act of breathing itself is little more than an application of Boyle's law.

The lungs are a series of ever-narrowing tubes that end in a myriad of tiny sacs called alveoli. It is in the alveoli that oxygen from the air transfers to the bloodstream and carbon dioxide from the bloodstream transfers to the lungs for exhalation. For air to move in and out of the lungs, the pressure inside the lungs must change, forcing the lungs to change volume—just as predicted by Boyle's law.

The pressure change is caused by the diaphragm, a muscle that covers the bottom of the lungs. When the diaphragm moves down, it expands the size of our lungs. When this happens, the air pressure inside our lungs decreases slightly. This causes new air to rush in, and we inhale. The pressure decrease is slight—only 3 torr, or about 0.4% of an atmosphere. We inhale only 0.5–1.0 L of air per normal breath.

Exhaling air requires that we relax the diaphragm, which pushes against the lungs and slightly decreases the volume of the lungs. This slightly increases the pressure of the air in the lungs, and air is forced out; we exhale. Only 1–2 torr of extra pressure is needed to exhale. So with every breath, our own bodies are performing an experimental test of Boyle's law.

Charles's Law

Another simple gas law relates the volume of a gas to its temperature. Experiments indicate that as the temperature of a gas sample is increased, its volume increases as long as the pressure and the amount of gas remain constant. The way to write this mathematically is

$$\text{V} \propto T$$

At this point, the concept of temperature must be clarified. Although the Kelvin scale is the preferred temperature scale, the Celsius scale is also a common temperature scale used in science. The Celsius scale is based on the melting and boiling points of water and is actually the common temperature scale used by most countries around the world (except for the United States, which still uses the Fahrenheit scale). The value of a Celsius temperature is directly related to its Kelvin value by a simple expression:

$$\text{Kelvin temperature} = \text{Celsius temperature} + 273.15$$

Thus, it is easy to convert from one temperature scale to another.

The Kelvin scale is sometimes referred to as the absolute scale because the zero point on the Kelvin scale is at absolute zero, the coldest possible temperature. On the other temperature scales, absolute zero is –273.15°C or –459.67°F.

The expression relating a gas volume to its temperature begs the following question: to which temperature scale is the volume of a gas related? The answer is that gas volumes are directly related to the *Kelvin temperature*. Therefore, the temperature of a gas sample should always be expressed in (or converted to) a Kelvin temperature.

Example 9.5.4: Increasing Temperature

What happens to the volume of a gas if its temperature is decreased? Assume that all other conditions remain constant.

Solution

If the temperature of a gas sample is decreased, the volume decreases as well.

Exercise 9.5.4

What happens to the temperature of a gas if its volume is increased? Assume that all other conditions remain constant.

Answer available at the end of this section.

As with Boyle's law, the relationship between volume and temperature can be expressed in terms of initial and final values of volume and temperature, as follows:

$$\frac{\mathrm{V_i}}{\mathrm{T_i}} = \frac{\mathrm{V_f}}{\mathrm{T_f}}$$

where V_i and T_i are the initial volume and temperature, and V_f and T_f are the final volume and temperature. This is Charles's law. The restriction on its use is that the pressure of the gas and the amount of gas must remain constant. (Charles's law is sometimes referred to as Gay-Lussac's law, after the scientist who promoted Charles's work.)

Example 9.5.5

A gas sample at 20°C has an initial volume of 20.0 L. What is its volume if the temperature is changed to 60°C? Does the answer make sense? Assume that the pressure and the amount of the gas remain constant.

Solution

Although the temperatures are given in degrees Celsius, we must convert them to the kelvins before we can use Charles's law. Thus,

20°C + 273 = 293 K = T_i 60°C + 273 = 333 K = T_f

Now we can substitute these values into Charles's law, along with the initial volume of 20.0 L:

$$\frac{20.0\ \mathrm{L}}{293\ \mathrm{K}} = \frac{\mathrm{V_f}}{333\ \mathrm{K}}$$

Multiplying the 333 K to the other side of the equation, we see that our temperature units will cancel:

$$\frac{(333\ \mathrm{K})(20.0\ \mathrm{L})}{293\ \mathrm{K}} = \mathrm{V_f}$$

Solving for the final volume, V_f = 22.7 L. So, as the temperature is increased, the volume increases. This makes sense because volume is directly proportional to the absolute temperature (as long as the pressure and the amount of the remain constant).

Exercise 9.5.5

A gas sample at 35°C has an initial volume of 5.06 L. What is its volume if the temperature is changed to –35°C? Does the answer make sense? Assume that the pressure and the amount of the gas remain constant.

Answer available at the end of this section.

Combined Gas Law

Other gas laws can be constructed, but we will focus on only two more. The combined gas law brings Boyle's and Charles's laws together to relate pressure, volume, and temperature changes of a gas sample:

$$\frac{(\mathrm{V_i})(\mathrm{T_i})}{\mathrm{P_i}} = \frac{(\mathrm{V_f})(\mathrm{T_f})}{\mathrm{P_f}}$$

To apply this gas law, the amount of gas should remain constant. As with the other gas laws, the temperature must be expressed in kelvins, and the units on the similar quantities should be the same. Because of the dependence on three quantities at the same time, it is difficult to tell in advance what will happen to one property of a gas sample as two other properties change. The best way to know is to work it out mathematically.

Example 9.5.6

A sample of gas has P_i = 1.50 atm, V_i = 10.5 L, and T_i = 300 K. What is the final volume if P_f = 0.750 atm and T_f = 350 K?

Solution

Using the combined gas law, substitute for five of the quantities:

$$\frac{(1.50\ \mathrm{atm})(10.5\ \mathrm{L})}{300\ \mathrm{K}} = \frac{(0.750\ \mathrm{atm})(\mathrm{V_f})}{350\ \mathrm{K}}$$

We algebraically rearrange this expression to isolate V_f on one side of the equation:

$$\mathrm{V_f} = \frac{(1.50\ \mathrm{atm})(10.5\ \mathrm{L})(350\ \mathrm{K})}{(300\ \mathrm{K})(0.750\ \mathrm{atm})} = 24.5\ \mathrm{L}$$

Note how all the units cancel except the unit for volume.

Exercise 9.5.6

A sample of gas has P_i = 0.768 atm, V_i = 10.5 L, and T_i = 300 K. What is the final pressure if V_f = 7.85 L and T_f = 250 K?

Answer available at the end of this section.

Example 9.5.7

A balloon containing a sample of gas has a temperature of 22°C and a pressure of 1.09 atm in an airport in Cleveland. The balloon has a volume of 1,070 mL. The balloon is transported by plane to Denver, where the temperature is 11°C and the pressure is 655 torr. What is the new volume of the balloon?

Solution

The first task is to convert all quantities to the proper and consistent units. The temperatures must be expressed in kelvins, and the pressure units are different so one of the quantities must be converted. Let us convert the atmospheres to torr:

$$22°C + 273 = 295\ K = T_i$$

$$11°C + 273 = 284\ K = T_f$$

$$1.09\ \text{atm} \times \frac{760\ \text{torr}}{1\ \text{atm}} = 828\ \text{torr} = \text{P}_\text{i}$$

Now we can substitute the quantities into the combined has law:

$$\frac{(828\ \text{torr})(1,070\ \text{mL})}{295\ \text{K}} = \frac{(655\ \text{torr}) \times \text{V}_\text{f}}{284\ \text{K}}$$

To solve for V_f, we multiply the 284 K in the denominator of the right side into the numerator on the left, and we divide 655 torr in the numerator of the right side into the denominator on the left:

$$\frac{(828\ \text{torr})(1,070\ \text{mL})(284\ \text{K})}{(295\ \text{K})(655\ \text{torr})} = \text{V}_\text{f}$$

Notice that torr and kelvins cancel, as they are found in both the numerator and denominator. The only unit that remains is milliliters, which is a unit of volume. So V_f = 1,300 mL. The overall change is that the volume of the balloon has increased by 230 mL.

Exercise 9.5.7

A balloon used to lift weather instruments into the atmosphere contains gas having a volume of 1,150 L on the ground, where the pressure is 0.977 atm and the temperature is 18°C. Aloft, this gas has a pressure of 6.88 torr and a temperature of –15°C. What is the new volume of the gas?

Answer available at the end of this section.

The Ideal Gas Law

So far, the gas laws we have used have focused on changing one or more properties of the gas, such as its volume, pressure, or temperature. There is one gas law that relates all the independent properties of a gas under any particular condition, rather than a change in conditions. This gas law is called the ideal gas law. The formula of this law is as follows:

$$PV = nRT$$

In this equation, *P* is pressure, *V* is volume, *n* is amount of moles, and *T* is temperature. *R* is called the ideal gas law constant and is a proportionality constant that relates the values of pressure, volume, amount, and temperature of a gas sample. The variables in this equation do not have the subscripts *i* and *f* to indicate an initial condition and a final condition. The ideal gas law relates the four independent properties of a gas under *any* conditions.

The value of *R* depends on what units are used to express the other quantities. If volume is expressed in liters and pressure in atmospheres, then the proper value of *R* is as follows:

$$\text{R} = 0.08205\ \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}$$

This may seem like a strange unit, but that is what is required for the units to work out algebraically.

Example 9.5.8

What is the volume in liters of 1.45 mol of N_2 gas at 298 K and 3.995 atm?

Solution

Using the ideal gas law where P = 3.995 atm, n = 1.45, and T = 298,

$$(3.995\ \text{atm}) \times \text{V} = (1.45\ \text{mol}) \left(0.08205\ \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (298\ \text{K})$$

On the right side, the moles and kelvins cancel. Also, because atmospheres appear in the numerator on both sides of the equation, they also cancel. The only remaining unit is liters, a unit of volume. So

$$3.995 \times V = (1.45)(0.08205)(298)\ \text{L}$$

Dividing both sides of the equation by 3.995 and evaluating, we get V = 8.87 L. Note that the conditions of the gas are not changing. Rather, the ideal gas law allows us to determine what the fourth property of a gas (here, volume) *must* be if three other properties (here, amount, pressure, and temperature) are known.

Exercise 9.5.8

What is the pressure of a sample of CO_2 gas if 0.557 mol is held in a 20.0 L container at 451 K?

Answer available at the end of this section.

For convenience, scientists have selected 273 K (0°C) and 1.00 atm pressure as a set of standard conditions for gases. This combination of conditions is called standard temperature and pressure (STP). Under these conditions, 1 mol of any gas has about the same volume. We can use the ideal gas law to determine the volume of 1 mol of gas at STP:

$$(1.00\ \text{atm}) \times \text{V} = (1.00\ \text{mol}) \left(0.08205\ \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (273\ \text{K})$$

This volume is 22.4 L. Because this volume is independent of the identity of a gas, the idea that 1 mol of gas has a volume of 22.4 L at STP makes a convenient conversion factor:

1 mol gas = 22.4 L (at STP)

Example 9.5.9

Cyclopropane (C_3H_6) is a gas that formerly was used as an anesthetic. How many moles of gas are there in a 100.0 L sample if the gas is at STP?

Solution

We can set up a simple, one-step conversion that relates moles and liters:

$$100.0\ \text{L}\ \text{C}_3\text{H}_6 \times \frac{1\ \text{mol}}{22.4\ \text{L}} = 4.46\ \text{mol}\ \text{C}_3\text{H}_6$$

There are almost 4.5 mol of gas in 100.0 L.

Note: Because of its flammability, cyclopropane is no longer used as an anesthetic gas.

Exercise 9.5.9

Freon is a trade name for a series of fluorine- and chlorine-containing gases that formerly were used in refrigeration systems. What volume does 8.75 mol of Freon have at STP?

Note: Many gases known as Freon are no longer used because their presence in the atmosphere destroys the ozone layer, which protects us from ultraviolet light from the sun.

Answer available at the end of this section.

Airbags

Airbags (Figure 9.5.1) are a safety feature provided in most automobiles since the 1990s. The effective operation of an airbag requires that it be rapidly inflated with an appropriate amount (volume) of gas when the vehicle is involved in a collision. This requirement is satisfied in many automotive airbag systems through use of explosive chemical reactions, one common choice being the decomposition of sodium azide, NaN_3. When sensors in the vehicle detect a collision, an electrical current is passed through a carefully measured amount of NaN_3 to initiate its decomposition:

$$2\,\mathrm{NaN_3}(s) \rightarrow 3\,\mathrm{N_2}(g) + 2\,\mathrm{Na}(s)$$

This reaction is very rapid, generating gaseous nitrogen that can deploy and fully inflate a typical airbag in a fraction of a second (~0.03–0.1 s). Among many engineering considerations, the amount of sodium azide used must be appropriate for generating enough nitrogen gas to fully inflate the air bag and ensure its proper function. For example, a small mass (~100 g) of NaN_3 will generate approximately 50 L of N_2.



Figure 9.5.1: Airbags deploy upon impact to minimize serious injuries to passengers. (credit: Jon Seidman)

Dalton's Law of Partial Pressures

The ideal gas equation of state applies to mixtures just as to pure gases. It was in fact with a gas mixture, ordinary air, that Boyle, Gay-Lussac and Charles did their early experiments. The only new concept we need in order to deal with gas mixtures is the *partial pressure*, a concept invented by the famous English chemist John Dalton (1766-1844). Dalton reasoned that the low density and high compressibility of gases indicates that they consist mostly of empty space; from this it follows that when two or more different gases occupy the same volume, they behave entirely independently. The contribution that each component of a gaseous mixture makes to the total pressure of the gas is known as the *partial pressure* of that gas.

The definition of Dalton's Law of Partial Pressures that address this is:

The total pressure of a gas is the sum of the partial pressures of its components

which is expressed algebraically as

$$P_{total} = P_1 + P_2 + P_3... = \sum_i P_i$$

or, equivalently

$$P_{total} = \frac{RT}{V} \sum_i n_i$$

There is also a similar relationship based on *volume fractions*, known as *Amagat's law of partial volumes*. It is exactly analogous to Dalton's law, in that it states that the total volume of a mixture is just the sum of the partial volumes of its components. But there are two important differences: Amagat's law holds only for ideal gases which must all be at the same temperature and pressure. Dalton's law has neither of these restrictions. Although Amagat's law seems intuitively obvious, it sometimes proves useful in chemical engineering applications. We will make no use of it in this course.

Example 9.5.10

Three flasks having different volumes and containing different gases at various pressures are connected by stopcocks as shown. When the stopcocks are opened,

a. What will be the pressure in the system?
b. Which gas will be most abundant in the mixture?

Assume that the temperature is uniform and that the volume of the connecting tubes is negligible.




Figure 9.5.2: Airbags deploy upon impact to minimize serious injuries to passengers. (credit: Jon Seidman)

Solution

The trick here is to note that the total number of moles n_T and the temperature remain unchanged, so we can make use of Boyle's law *PV* = constant. We will work out the details for CO_2 only, denoted by subscripts *a*.

For CO_2,

$$P_a V_a = (2.13\ atm)(1.50\ L) = 3.20\ L \cdot atm$$

Adding the *PV* products for each separate container, we obtain

\[\sum_i P_i V_i = 6.36\; L \cdot atm = n_T RT \nonumber \]

We will call this sum P_1V_1. After the stopcocks have been opened and the gases mix, the new conditions are denoted by P_2V_2.

From Boyle's law (9.5.1,

$$P_1V_1 = P_2V_2 = 6.36\ L \cdot atm$$

$$V_2 = \sum_i V_i = 4.50\ L$$

Solving for the final pressure P_2 we obtain (6.36 L-atm)/(4.50 L) = **1.41 atm**.

For part *(b)*, note that the number of moles of each gas is $n = PV/RT$. The mole fraction of any one gas is $X_i = n_i / n_T$. For CO_2, this works out to (3.19/*RT*) / (6.36/*RT*) = 0.501. Because this exceeds 0.5, we know that this is the most abundant gas in the final mixture.

Dalton's law states that in a gas mixture (P_{total}) each gas will exert a pressure independent of the other gases (P_n) and each gas will behave as if it alone occupies the total volume. By extension, the partial pressure of each gas can be calculated by multiplying the total pressure (P_{total}) by the gas percentage (%).

$$P_{Total} = P_1 + P_2 + P_3 + P_4 + ... + P_n$$

or

$$P_n = \frac{^n}{P_{Total}}$$

Table 9.5.1: Partial Pressures for the gases in air on a typical day

Gas	Partial Pressure (mm Hg)	Percentage (%)
Nitrogen, (N_2\)	P_{N_2} = 594	78
Oxygen, O_2	P_{O_2}= 160	21
Carbon Dioxide, CO_2	P_{CO_2} = 0.25	0.033
Water Vapor, H_2O	P_{H_2O} = 5.7	0.75
Other trace gases	P_{Other} = 0.05	0.22
Total air	P_{Total} = 760	1

Application of Dalton's Law: Collecting Gases over Water

A common laboratory method of collecting the gaseous product of a chemical reaction is to conduct it into an inverted tube or bottle filled with water, the opening of which is immersed in a larger container of water. This arrangement is called a *pneumatic trough*, and was widely used in the early days of chemistry. As the gas enters the bottle it displaces the water and becomes trapped in the upper part.




Figure 9.5.3:An Apparatus for Collecting Gases by the Displacement of Water

The volume of the gas can be observed by means of a calibrated scale on the bottle, but what about its pressure? The total pressure confining the gas is just that of the atmosphere transmitting its force through the water. (An exact calculation would also have to take into account the height of the water column in the inverted tube.) But liquid water itself is always in equilibrium with its vapor, so the space in the top of the tube is a mixture of two gases: the gas being collected, and gaseous H_2O. The partial pressure of H_2O is known as the vapor pressure of water and it depends on the temperature. In order to determine the quantity of gas we have collected, we must use Dalton's Law to find the partial pressure of that gas.

Example 9.5.11

Oxygen gas was collected over water as shown above. The atmospheric pressure was 754 torr, the temperature was 22°C, and the volume of the gas was 155 mL. The vapor pressure of water at 22°C is 19.8 torr. Use this information to estimate the number of moles of O_2 produced.

Solution

From Dalton's law,

$$P_{O_2} = P_{total} – P_{H_2O} = 754 – 19.8 = 734\ torr = 0.966\ atm$$

Now use the Ideal Gas Law to convert to moles

$$n = \frac{PV}{RT} = \frac{(0.966\ atm)(0.155\ L)}{(0.082\ Latmmol^{-1}K^{-1})(295\ K)} = 0.00619\ mol$$

Henry's Law

Henry's law is one of the gas laws formulated by William Henry in 1803. It states: "At a constant temperature, the amount of a given gas that dissolves in a given type and volume of liquid is directly proportional to the partial pressure of that gas in equilibrium with that liquid." An equivalent way of stating the law is that the solubility of a gas in a liquid is directly proportional to the partial pressure of the gas above the liquid.

To explain this law, Henry derived the equation:

$$C = kP_{gas}$$

where

Henry's Law tells us that the greater the pressure of gas above the surface of a liquid, the higher the concentration of the gas in the liquid. Also, Henry's law tells us that gases diffuse from areas of high gas concentration to areas of low gas concentration.

Applicability of Henry's Law

- Henry's law only works if the molecules are at equilibrium.
- Henry's law does not work for gases at high pressures (e.g., $N_{2\ (g)}$ at high pressure becomes very soluble and harmful when in the blood supply).
- Henry's law does not work if there is a chemical reaction between the solute and solvent (e.g., $HCl_{(g)}$ reacts with water by a dissociation reaction to generate H_3O^+ and Cl^- ions).

Application of Henry's Law: Scuba diving

Our respiratory systems are designed to maintain the proper oxygen concentration in the blood when the partial pressure of O_2 is 0.21 atm, its normal sea-level value. Below the water surface, the pressure increases by 1 atm for each 10.3 m increase in depth; thus a scuba diver at 10.3 m experiences a total of 2 atm pressure pressing on the body. In order to prevent the lungs from collapsing, the air the diver breathes should also be at about the same pressure.




Figure 9.5.4: Scuba Diving actively takes into account both Henry's and Dalton's Laws

But at a total pressure of 2 atm, the partial pressure of O_2 in ordinary air would be 0.42 atm; at a depth of 100 ft (about 30 m), the O_2 pressure of 0.8 atm would be far too high for health. For this reason, the air mixture in the pressurized tanks that scuba divers wear must contain a smaller fraction of O_2. This can be achieved most simply by raising the nitrogen content, but high partial pressures of N_2 can also be dangerous, resulting in a condition known as nitrogen narcosis. The preferred diluting agent for sustained deep diving is helium, which has very little tendency to dissolve in the blood even at high pressures.

Career Focus: Respiratory Therapist

Certain diseases—such as emphysema, lung cancer, and severe asthma—primarily affect the lungs. Respiratory therapists help patients with breathing-related problems. They can evaluate, help diagnose, and treat breathing disorders and even help provide emergency assistance in acute illness where breathing is compromised.

Most respiratory therapists must complete at least two years of college and earn an associate's degree, although therapists can assume more responsibility if they have a college degree. Therapists must also pass state or national certification exams. Once certified, respiratory therapists can work in hospitals, doctor's offices, nursing homes, or patient's homes. Therapists work with equipment such as oxygen tanks and respirators, may sometimes dispense medication to aid in breathing, perform tests, and educate patients in breathing exercises and other therapy.

Because respiratory therapists work directly with patients, the ability to work well with others is a must for this career. It is an important job because it deals with one of the most crucial functions of the body.

Key Takeaway

- The physical properties of gases are predictable using mathematical formulas known as gas laws.
- C is the solubility of a gas at a fixed temperature in a particular solvent (in units of M or mL gas/L)
- k is Henry's law constant (often in units of M/atm)
- P_{gas} is the partial pressure of the gas (often in units of Atm)

Answers

Exercise $9.5.1$**: Increasing Volume in a Gas**

If the volume of a gas is increased, the pressure decreases.

Exercise $9.5.2$

8.48 atm

Exercise $9.5.3$

575 torr

Exercise $9.5.4$

The temperature increases.

Exercise $9.5.5$

3.91 L

Exercise $9.5.6$

0.856 atm

Exercise $9.5.7$

110,038 L

Exercise $9.5.8$

1.03 atm

Exercise $9.5.9$

196 L

This page titled 9.5: Gas Laws is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.4: Gas Laws by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

9.S: Solids, Liquids, and Gases (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the following bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **phase** is a certain form of matter that has the same physical properties throughout. Three phases are common: the solid, the liquid, and the gas phase. What determines the phase of a substance? Generally, the strength of the **intermolecular interactions** determines whether a substance is a solid, liquid, or gas under any particular conditions. **Covalent network bonding** is a very strong form of intermolecular interaction. Diamond is one example of a substance that has this intermolecular interaction. **Ionic interactions**, the forces of attraction due to oppositely charged ions, are also relatively strong. Covalent bonds are another type of interaction within molecules, but if the bonds are **polar covalent bonds**, then the unequal sharing of electrons can cause charge imbalances within molecules that cause interactions between molecules. These molecules are described as **polar**, and these interactions are called **dipole-dipole interactions**. A certain rather strong type of dipole-dipole interaction, involving a hydrogen atom, is called **hydrogen bonding**. On the other hand, equal sharing of electrons forms **nonpolar covalent bonds**, and the interactions between different molecules is less because the molecules are nonpolar. All substances have very weak **dispersion forces** (also called **London forces**) caused by the movement of electrons within the bonds themselves.

In the solid phase, intermolecular interactions are so strong that they hold the individual atoms or molecules in place. In many solids, the regular three-dimensional arrangement of particles makes a **crystal**. In other solids, the irregular arrangement of particles makes an **amorphous** solid. In liquids, the intermolecular interactions are strong enough to keep the particles of substance together but not in place. Thus, the particles are free to move over each other but still remain in contact.

In gases, the intermolecular interactions are weak enough that the individual particles are separated from each other in space. The **kinetic theory of gases** is a collection of statements that describe the fundamental behavior of all gases. Among other properties, gases exert a **pressure** on their container. Pressure is measured using units like **pascal**, **bar**, **atmosphere**, or **mmHg** (also called a **torr**).

There are several simple relationships between the variables used to describe a quantity of gas. These relationships are called **gas laws**. **Boyle's law** relates the pressure and volume of a gas, while **Charles's law** relates the volume and absolute temperature of a gas. The **combined gas law** relates the volume, pressure, and absolute temperature of a gas sample. All of these gas laws allow us to understand the changing conditions of a gas. The **ideal gas law** relates the pressure, volume, amount, and absolute temperature of a gas under any conditions. These four variables are related to the **ideal gas law constant**, which is the proportionality constant used to calculate the conditions of a gas. Because the conditions of a gas can change, a set of benchmark conditions called **standard temperature and pressure (STP)** is defined. Standard temperature is 0°C, and standard pressure is 1.00 atm.

This page titled 9.S: Solids, Liquids, and Gases (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 8.S: Solids, Liquids, and Gases (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10: SOLUTIONS

Solutions are all around us. Air, for example, is a solution. If you live near a lake, a river, or an ocean, that body of water is not pure H_2O but most probably a solution. Much of what we drink—for example, soda, coffee, tea, and milk—is at least in part a solution. Solutions are a large part of everyday life. A lot of the chemistry occurring around us happens in solution. In fact, much of the chemistry that occurs in our own bodies takes place in solution, and many solutions—such as the Ringer's lactate IV solution—are important for our health. In our understanding of chemistry, we need to understand a little bit about solutions. In this chapter, you will learn about the special characteristics of solutions, how solutions are characterized, and some of their properties.

10.1: Prelude to Solutions

10.2: Solutions

10.3: Concentration

10.4: The Dissolution Process

10.5: Properties of Solutions

10.6: Chemical Equilibrium

10.7: Le Chatelier's Principle

10.8: Osmosis and Diffusion

10.S: Solutions (Summary)

Template:HideTOC

This page titled 10: Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9: Solutions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.1: Prelude to Solutions

If you watch any of the medical dramas on television, you may have heard a doctor (actually an actor) call for an intravenous solution of “Ringer’s lactate” (or “lactated Ringer’s” or "Ri-Lac"). So what is Ringer’s lactate?

Intravenous (IV) solutions are administered for two main reasons:

1. to introduce necessary substances into the bloodstream, such as ions for proper body function, sugar and other food substances for energy, or drugs to treat a medical condition, and
2. to increase the volume of the bloodstream.

Many people with acute or long-term medical conditions have received some type of an IV solution.

One basic IV solution, called a *normal saline solution*, is simply a dilute solution of NaCl dissolved in water. Normal saline is 9.0 g of NaCl dissolved in each liter of solution. *Ringer’s lactate* is a normal saline solution that also has small amounts of potassium and calcium ions mixed in. In addition, it contains about 2.5 g of lactate ions ($C_3H_5O_3^-$) per liter of solution. The liver metabolizes lactate ions into bicarbonate (HCO_3^-) ions, which help maintain the acid-base balance of blood. Many medical problems, such as heart attacks and shock, affect the acid-base balance of blood, and the presence of lactate in the IV solution eases problems caused by this imbalance.

Physicians can select from a range of premade IV solutions, in accordance with a patient’s particular needs. Ringer’s lactate is commonly used when a patient’s blood volume must be increased quickly. Another frequently used IV solution, called D5W, is a 5% solution of dextrose (a form of sugar) in water.

This page titled 10.1: Prelude to Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.0: Prelude to Solutions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.2: Solutions

Learning Objectives

- To understand what causes solutions to form.

A solution is another name for a homogeneous mixture. A *mixture* as a material composed of two or more substances. In a solution, the combination is so intimate that the different substances cannot be differentiated by sight, even with a microscope. Compare, for example, a mixture of salt and pepper and another mixture consisting of salt and water. In the first mixture, we can readily see individual grains of salt and the flecks of pepper. A mixture of salt and pepper is not a solution. However, in the second mixture, no matter how carefully we look, we cannot see two different substances. Salt dissolved in water is a solution.

The major component of a solution, called the **solvent**, is typically the same phase as the solution itself. The solvent is the component present in the largest amount. Each minor component of a solution (and there may be more than one) is called the **solute**. In most of the solutions we will describe in this textbook, there will be no ambiguity about whether a component is the solvent or the solute. For example, in a solution of salt in water, the solute is salt, and solvent is water.

Solutions come in all phases, and the solvent and the solute do not have to be in the same phase to form a solution (such as salt and water). For example, air is a gaseous solution of about 80% nitrogen and about 20% oxygen, with some other gases present in much smaller amounts. An alloy is a solid solution consisting of a metal (like iron) with some other metals or nonmetals dissolved in it. Steel, an alloy of iron and carbon and small amounts of other metals, is an example of a solid solution. Table 10.2.1 lists some common types of solutions, with examples of each.

Table 10.2.1: Types of Solutions

Solvent Phase	Solute Phase	Example
gas	gas	air
liquid	gas	carbonated beverages
liquid	liquid	ethanol (C_2H_5OH) in H_2O (alcoholic beverages)
liquid	solid	saltwater
solid	gas	H_2 gas absorbed by Pd metal
solid	liquid	Hg(ℓ) in dental fillings
solid	solid	steel alloys

What Causes a Solution to Form?

The simple answer is that the solvent and the solute must have similar intermolecular interactions. When this is the case, the individual particles of solvent and solute can easily mix so intimately that each particle of solute is surrounded by particles of solvent, forming a solution. However, if two substances have very different intermolecular interactions, large amounts of energy are required to force their individual particles to mix intimately, so a solution does not form. Thus two alkanes like *n*-heptane, C_7H_{16}, and *n*-hexane, C_6H_{14}, are completely miscible in all proportions. The C_7H_{16} and C_6H_{14} molecules are so similar (recall Section 4.6) that there are only negligible differences in intermolecular forces.

For a similar reason, methanol, CH_3OH, is completely miscible with water. In this case both molecules are polar and can form hydrogen bonds among themselves, and so there are strong intermolecular attractions within each liquid. However, CH_3OH dipoles can align with H_2O dipoles, and CH_3OH molecules can hydrogen bond to H_2O molecules, and so the attractions among unlike molecules in the solution are similar to those among like molecules in each pure liquid.

Like Dissolves Like

This process leads to a simple rule of thumb: *like dissolves like*. Solvents that are very polar will dissolve solutes that are very polar or even ionic. Solvents that are nonpolar will dissolve nonpolar solutes. Thus water, being polar, is a good solvent for ionic compounds and polar solutes like ethanol (C_2H_5OH). However, water does not dissolve nonpolar solutes, such as many oils and greases (Figure 10.2.1).



Figure 10.2.1: Oil (upper layer) which is nonpolar will not dissolve in polar water (lower layer). Figure used with permission from Wikipedia.

We use the word soluble to describe a solute that dissolves in a particular solvent, and the word insoluble for a solute that does not dissolve in a solvent. Thus, we say that sodium chloride is soluble in water but insoluble in hexane (C_6H_{14}). If the solute and the solvent are both liquids and soluble in any proportion, we use the word miscible, and the word immiscible if they are not.

Example 10.2.1

Water is considered a polar solvent. Which substances should dissolve in water?

1. methanol (CH_3OH)
2. sodium sulfate (Na_2SO_4)
3. octane (C_8H_{18})

Solution

Because water is polar, substances that are polar or ionic will dissolve in it.

1. Because of the OH group in methanol, we expect its molecules to be polar. Thus, we expect it to be soluble in water. As both water and methanol are liquids, the word *miscible* can be used in place of *soluble*.
2. Sodium sulfate is an ionic compound, so we expect it to be soluble in water.
3. Like other hydrocarbons, octane is nonpolar, so we expect that it would not be soluble in water.

Exercise 10.2.1

Toluene ($C_6H_5CH_3$) is widely used in industry as a nonpolar solvent. Which substances should dissolve in toluene?

a. water (H_2O)
b. sodium sulfate (Na_2SO_4)

c. octane (C_8H_{18})

Answer available at the end of this section.

Example 10.2.2

Predict which of the following compounds will be most soluble in water:

a. $\underset{\text{Ethanol}}{CH_3CH_2OH}$
b. $\underset{\text{Hexanol}}{CH_3CH_2CH_2CH_2CH_2CH_2OH}$

Solution

Since ethanol contains an **OH** group, it can hydrogen bond to water. Although the same is true of hexanol, the OH group is found only at one end of a fairly large molecule. The rest of the molecule can be expected to behave much as though it were a nonpolar alkane. This substance should thus be much less soluble than the first. Experimentally we find that ethanol is completely miscible with water, while only 0.6 g hexanol dissolves in 100 g water.

Exercise 10.2.2

Would I_2 be more soluble in CCl_4 or H_2O?

Answer available at the end of this section.

Key Takeaway

- Solutions form because a solute and a solvent experience similar intermolecular interactions.

Answers

Exercise 10.2.1

Octane only.

Exercise 10.2.2

I_2 is nonpolar. Of the two solvents, CCl_4 is nonpolar and H_2O is polar, so I_2 would be expected to be more soluble in CCl_4.

This page titled 10.2: Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.1: Solutions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.3: Concentration

Learning Objectives

- Calculate percentage concentration (m/m, v/v, m/v), ppm and ppb.
- Calculate the molarity of a solution.
- Use concentration units to calculate the amount of solute in a solution.
- Use molarity to determine quantities in chemical reactions.
- Determine the resulting concentration of a diluted solution.

To define a solution precisely, we need to state its concentration: how much solute is dissolved in a certain amount of solvent. Words such as *dilute* or *concentrated* are used to describe solutions that have a little or a lot of dissolved solute, respectively, but these are relative terms whose meanings depend on various factors.

Solubility

There is usually a limit to how much solute will dissolve in a given amount of solvent. This limit is called the **solubility** of the solute. Some solutes have a very small solubility, while other solutes are soluble in all proportions. Table 10.3.1 lists the solubilities of various solutes in water. Solubilities vary with temperature, so Table 10.3.1 includes the temperature at which the solubility was determined.

Table 10.3.1: Solubilities of Various Solutes in Water at 25°C (Except as Noted)

Substance	Solubility (g in 100 mL of H2O)
AgCl(s)	0.019
$C_6H_6(\ell)$ (benzene)	0.178
$CH_4(g)$	0.0023
$CO_2(g)$	0.150
$CaCO_3(s)$	0.058
$CaF_2(s)$	0.0016
$Ca(NO_3)_2(s)$	143.9
$C_6H_{12}O_6$ (glucose)	120.3 (at 30°C)
KBr(s)	67.8
$MgCO_3(s)$	2.20
NaCl(s)	36.0
$NaHCO_3(s)$	8.41
$C_{12}H_{22}O_{11}$ (sucrose)	204.0 (at 20°C)

If a solution contains so much solute that its solubility limit is reached, the solution is said to be **saturated**, and its concentration is known from information contained in Table 10.3.1. If a solution contains less solute than the solubility limit, it is **unsaturated**. Under special circumstances, more solute can be dissolved even after the normal solubility limit is reached; such solutions are called ***supersaturated*** and are not stable. If the solute is solid, excess solute can easily recrystallize. If the solute is a gas, it can

bubble out of solution uncontrollably, like what happens when you shake a soda can and then immediately open it.

Precipitation from Supersaturated Solutions

Recrystallization of excess solute from a supersaturated solution usually gives off energy as heat. Commercial heat packs containing supersaturated sodium acetate ($NaC_2H_3O_2$) take advantage of this phenomenon. You can probably find them at your local drugstore.

This interactive element is not available in this format. Visit the link or scan the QR code to access it online.

View this YouTube Video online

Video $10.3.1$***:*** Watered-down sodium acetate trihydrate. Needle crystal is truly wonderful structures

Most solutions we encounter are unsaturated, so knowing the solubility of the solute does not accurately express the amount of solute in these solutions. There are several common ways of specifying the concentration of a solution.

Percent Composition

There are several ways of expressing the concentration of a solution by using a percentage. The **mass/mass percent** (% m/m) is defined as the mass of a solute divided by the mass of a solution times 100:

$$\%\ \mathrm{m/m} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 100\%$$

If you can measure the masses of the solute and the solution, determining the mass/mass percent is easy. Each mass must be expressed in the same units to determine the proper concentration.

Example $10.3.1$

A saline solution with a mass of 355 g has 36.5 g of NaCl dissolved in it. What is the mass/mass percent concentration of the solution?

Solution

We can substitute the quantities given in the equation for mass/mass percent:

$$\%\ \mathrm{m/m} = \frac{36.5\ \mathrm{g}}{355\ \mathrm{g}} \times 100\% = 10.3\%$$

Exercise $10.3.1$

A dextrose (also called D-glucose, $C_6H_{12}O_6$) solution with a mass of 2.00×10^2 g has 15.8 g of dextrose dissolved in it. What is the mass/mass percent concentration of the solution?

Answer available at the end of this section.

For gases and liquids, volumes are relatively easy to measure, so the concentration of a liquid or a gas solution can be expressed as a **volume/volume percent** (% v/v): the volume of a solute divided by the volume of a solution times 100:

$$\% \text{ v/v} = \frac{\text{volume of solute}}{\text{volume of solution}} \times 100\%$$

Again, the units of the solute and the solution must be the same. A hybrid concentration unit, **mass/volume percent** (% m/v), is commonly used for intravenous (IV) fluids (Figure 10.3.1). It is defined as the mass in grams of a solute, divided by volume in milliliters of solution times 100:

$$\% \text{ m/v} = \frac{\text{mass of solute (g)}}{\text{volume of solution (mL)}} \times 100\%$$



Figure 10.3.1: Mass/Volume Percent.The 0.9% NaCl concentration on this IV bag is mass/volume percent (left). Such solution is used for other purposes and available in bottles (right). Figures used with permission from Wikipedia

Using Percent Composition in Calculations

The **percent concentration** can be used to produce a **conversion factor** between the **amount of solute** and the **amount of solution**. As such, concentrations can be useful in a variety of stoichiometry problems as discussed in Chapter 6. In many cases, it is best to use the original definition of the concentration unit; it is that definition that provides the conversion factor.

As an example, if the given concentration is **5% v/v solution of alcohol**, this means that there are 5 mL of alcohol dissolved in every 100 mL solution.

5 mL alcohol = 100 mL solution

The two possible conversion factors are written as follows:

$$\frac{5 \text{ mL alcohol}}{100 \text{ mL solution}} \text{ or } \frac{100 \text{ mL solution}}{5 \text{ mL alcohol}}$$

Use the first conversion factor to convert from a given amount of solution to amount of solute. The second conversion factor is used to convert from a given amount of solute to amount of solution. Given any two quantities in any percent composition, the third quantity can be calculated, as the following example illustrates.

Example 10.3.2

A sample of 45.0% v/v solution of ethanol (C_2H_5OH) in water has a volume of 115 mL. What volume of ethanol solute does the sample contain?

Solution

A percentage concentration is simply the number of parts of solute per 100 parts of solution. Thus, the percent concentration of 45.0% v/v implies the following:

$$45.0\%\ \text{v/v} \rightarrow \frac{45\ \text{mL}\ C_2H_5OH}{100\ \text{mL solution}}$$

That is, there are 45 mL of C_2H_5OH for every 100 mL of solution. We can use this fraction as a **conversion factor** to determine the amount of C_2H_5OH in 115 mL of solution:

$$115\ \text{mL solution} \times \frac{45\ \text{mL}\ C_2H_5OH}{100\ \text{mL solution}} = 51.8\ \text{mL}\ C_2H_5OH$$

Exercise 10.3.2

What volume of a 12.75% m/v solution of glucose ($C_6H_{12}O_6$) in water is needed to obtain 50.0 g of $C_6H_{12}O_6$?

Answer available at the end of this section.

Example 10.3.3

A normal saline IV solution contains 9.0 g of NaCl in every liter of solution. What is the mass/volume percent of normal saline?

Solution

We can use the definition of mass/volume percent, but first we have to express the volume in milliliter units:

1 L = 1,000 mL

Because this is an exact relationship, it does not affect the significant figures of our result.

$$\%\ \text{m/v} = \frac{9.0\ \text{g NaCl}}{1,000\ \text{mL solution}} \times 100\% = 0.90\%\ \text{m/v}$$

Exercise 10.3.3

The chlorine bleach that you might find in your laundry room is typically composed of 27.0 g of sodium hypochlorite (NaOCl), dissolved to make 500.0 mL of solution. What is the mass/volume percent of the bleach?

Answer available at the end of this section.

In addition to percentage units, the units for expressing the concentration of extremely dilute solutions are parts per million (ppm) and parts per billion (ppb). Both of these units are mass based and are defined as follows:

$$\text{ppm} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 1,000,000$$

$$\text{ppb} = \frac{\text{mass of solute}}{\text{mass of solution}} \times 1,000,000,000$$

Similar to parts per million and parts per billion, related units include parts per thousand (ppth) and parts per trillion (ppt).

Concentrations of *trace elements* in the body—elements that are present in extremely low concentrations but are nonetheless necessary for life—are commonly expressed in parts per million or parts per billion. Concentrations of poisons and pollutants are also described in these units. For example, cobalt is present in the body at a concentration of 21 ppb, while the State of Oregon's

Department of Agriculture limits the concentration of arsenic in fertilizers to 9 ppm.

In aqueous solutions, 1 ppm is essentially equal to 1 mg/L, and 1 ppb is equivalent to 1 µg/L.

Example 10.3.4

If the concentration of cobalt in a human body is 21 ppb, what mass in grams of Co is present in a body having a mass of 70.0 kg?

Solution

A concentration of 21 ppb means "21 g of solute per 1,000,000,000 g of solution." Written as a **conversion factor,** this concentration of Co is as follows:

$$21 \text{ ppb Co} \rightarrow \frac{21 \text{ g Co}}{1,000,000,000 \text{ g solution}}$$

We can use this as a conversion factor, but first we must convert 70.0 kg to gram units:

$$70.0 \text{ kg} \times \frac{1,000 \text{ g}}{1 \text{ kg}} = 7.00 \times 10^4 \text{ g}$$

Now we determine the amount of Co:

$$7.00 \times 10^4 \text{ g solution} \times \frac{21 \text{ g Co}}{1,000,000,000 \text{ g solution}} = 0.0015 \text{ g Co}$$

This is only 1.5 mg.

Exercise 10.3.4

An 85 kg body contains 0.012 g of Ni. What is the concentration of Ni in parts per million?

Answer available at the end of this section.

Molarity

Another way of expressing concentration is to give the number of moles of solute per unit volume of solution. Such concentration units are useful for discussing chemical reactions in which a solute is a product or a reactant. Molar mass can then be used as a conversion factor to convert amounts in moles to amounts in grams.

Molarity is defined as the number of moles of a solute dissolved per liter of solution:

$$\text{molarity} = \frac{\text{number of moles of solute}}{\text{number of liters of solution}}$$

Molarity is abbreviated M (often referred to as "molar"), and the units are often abbreviated as mol/L. It is important to remember that "mol" in this expression refers to moles of solute and that "L" refers to liters of solution. For example, if you have 1.5 mol of NaCl dissolved in 0.500 L of solution, its molarity is therefore

$$\frac{1.5 \text{ mol NaCl}}{0.500 \text{ L solution}} = 3.0 \text{ M NaCl}$$

which is read as "three point oh **molar** sodium chloride." Sometimes (aq) is added when the solvent is water, as in "3.0 M NaCl(aq)."

Before a molarity concentration can be calculated, the amount of the **solute must be expressed in moles**, and the **volume of the solution** must be expressed in **liters**.

If the quantity of the solute is given in mass units, you must convert mass units to mole units before using the definition of molarity to calculate concentration. For example, what is the molar concentration of a solution of 22.4 g of HCl dissolved in 1.56 L?




Figure 10.3.2: Determination of solution molarity from mass of hydrochloric acid.

Step 1: convert the **mass of solute** to **moles** using the **molar mass of HCl (36.46 g/mol):**

$$22.4\ \cancel{g HCl} \times \frac{1\ mol\ HCl}{36.46\ \cancel{g HCl}} = 0.614\ mol\ HCl$$

Step 2: use the **definition of molarity** to determine the concentration:

$$M\ =\ \frac{0.614\ mol\ HCl}{1.56 L\ solution} = 0.394\ M HCl$$

Example 10.3.5

What is the molarity of an aqueous solution of 25.0 g of NaOH in 750 mL?

Solution

Before we substitute these quantities into the definition of molarity, we must convert them to the proper units. The mass of NaOH must be converted to moles of NaOH. The molar mass of NaOH is 40.00 g/mol:

$$25.0\ \text{g NaOH} \times \frac{1\ \text{mol NaOH}}{40.00\ \text{g NaOH}} = 0.625\ \text{mol NaOH}$$

Next, we convert the volume units from milliliters to liters:

$$750\ \text{mL} \times \frac{1\ \text{L}}{1,000\ \text{mL}} = 0.750\ \text{L}$$

Now that the quantities are expressed in the proper units, we can substitute them into the definition of molarity:

$$\text{M} = \frac{0.625\ \text{mol NaOH}}{0.750\ \text{L}} = 0.833\ \text{M NaOH}$$

Exercise 10.3.5

If a 350 mL cup of coffee contains 0.150 g of caffeine ($C_8H_{10}N_4O_2$), what is the molarity of this caffeine solution?

Answer available at the end of this section.

Using Molarity in Calculations

The definition of molarity can also be used to calculate a needed volume of solution, given its concentration and the number of moles desired, or the number of moles of solute (and subsequently, the mass of the solute), given its concentration and volume. As in the percent concentration, molarity can also be expressed as a **conversion factor**.

Molarity is defined as moles solute per liter solution. There is an understood 1 in the denominator of the conversion factor. For example, a **3.0 M solution of sucrose** means that there are three moles of sucrose dissolved in every **liter of solution**. Mathematically, this is stated as follows:

3.0 moles sucrose = 1 L solution

Dividing both sides of this expression by either side, we generate two possible conversion factors:

$$\frac{3.0\text{ mol sucrose}}{1\text{ L solution}} \text{ or } \frac{1\text{ L solution}}{3.0\text{ mol sucrose}}$$

The first conversion factor can be used to convert from *volume (L) of solution to moles solute*, and the *second converts from moles of solute to volume (L) of solution.*




Figure 10.3.3: Diagram showing that molarity can be sue to convert between liters of solution and moles of solute.

For example, suppose we are asked how many moles of sucrose are present in 0.108 L of a 3.0 M sucrose solution. The given volume (0.108 L) is multiplied by the first conversion factor to cancel the L units, and find that 0.32 moles of sucrose are present.

$$0.108\ \cancel{L\ solution} \times \frac{3.0\ mol\ sucrose}{1\cancel{L\ solution}} = 0.32\ mol\ sucrose$$

How many liters of 3.0 M sucrose solution are needed to obtain 4.88 mol of sucrose? In such a conversion, we multiply the given (4.88 moles sucrose) with the second conversion factor. This cancels the moles units and converts it to liters of solution.

$$4.88\ \cancel{mol\ sucrose} \times \frac{1\ L\ solution}{3.0\cancel{mol\ sucrose}} = 1.63\ L\ solution$$

Example 10.3.6

1. What volume of a 0.0753 M solution of dimethylamine [$(CH_3)_2NH$] is needed to obtain 0.450 mol of the compound?
2. Ethylene glycol ($C_2H_6O_2$) is mixed with water to make auto engine coolants. How many grams of $C_2H_6O_2$ are in 5.00 L of a 6.00 M aqueous solution?

Solution

1. To solve for the volume, multiply the "given" (0.450 mol of dimethylamine) with the molarity conversion factor (0.0753 M). Use the proper conversion factor to cancel the unit "mol" and get the unit volume (L) of solution:

$$0.450 \text{ mol dimethylamine} \times \frac{1 \text{ L solution}}{0.0753 \text{ mol dimethylamine}} = 5.98 \text{ L solution}$$

2. The strategy in solving this problem is to convert the given volume (5.00 L) using the 6.00 M (conversion factor) to solve for moles of ethylene glycol, which can then be converted to grams.

Step 1: Convert the given volume (5.00 L) to moles ethylene glycol.

$$5.00 \text{ L solution} \times \frac{6.00 \text{ mol } C_2H_6O_2}{1 \text{ L solution}} = 30.0 \text{ mol } C_2H_6O_2$$

Step 2: Convert 30.0 mols $C_2H_6O_2$ to grams $C_2H_6O_2$. Molar mass of $C_2H_6O_2$= 62.08 g/mol

$$30.0 \text{ mol } C_2H_6O_2 \times \frac{62.08 \text{ g } C_2H_6O_2}{1 \text{ mol } C_2H_6O_2} = 1,860 \text{ g } C_2H_6O_2$$

The same two-step problem can also be worked out in a single line, rather than as two separate steps, as follows:




Figure 10.3.4: Determination of mass of solute in a solution of known molarity.

$$5.00 \text{ L solution} \times \frac{6.00 \text{ mol } C_2H_6O_2}{1 \text{ L solution}} \times \frac{62.08 \text{ g } C_2H_6O_2}{1 \text{ mol } C_2H_6O_2} = 1,860 \text{ g } C_2H_6O_2$$

The final answer is rounded off to 3 significant figures. Thus, there are 1,860 g of $C_2H_6O_2$ in the specified amount of engine coolant.

Note: Dimethylamine has a “fishy” odor. In fact, organic compounds called amines cause the odor of decaying fish.

Example 10.3.6A

1. What volume of a 0.0753 M solution of dimethylamine [$(CH_3)_2NH$] is needed to obtain 0.450 mol of the compound?
2. Ethylene glycol ($C_2H_6O_2$) is mixed with water to make auto engine coolants. How many grams of $C_2H_6O_2$ are in 5.00 L of a 6.00 M aqueous solution?

Solution

This is an alternative method in case you don't want to use the conversion factor for molarity. In both parts, we will use the definition of molarity to solve for the desired quantity.

1. $0.0753 \text{ M} = \frac{0.450 \text{ mol } (CH_3)_2NH}{\text{volume of solution}}$

To solve for the volume of solution, we multiply both sides by volume of solution and divide both sides by the molarity value to isolate the volume of solution on one side of the equation:

$$\text{volume of solution} = \frac{0.450 \text{ mol } (CH_3)_2NH}{0.0753 \text{ M}} = 5.98 \text{ L}$$

Note that because the definition of molarity is mol/L, the division of mol by M yields L, a unit of volume.

2. The molar mass of $C_2H_6O_2$ is 62.08 g/mol., so

$$6.00 \text{ M} = \frac{\text{moles of solute}}{5.00 \text{ L}}$$

To solve for the number of moles of solute, we multiply both sides by the volume:

$$\text{moles of solute} = (6.00 \text{ M})(5.00 \text{ L}) = 30.0 \text{ mol}$$

Note that because the definition of molarity is mol/L, the product M × L gives mol, a unit of amount. Now, using the molar mass of $C_3H_8O_3$, we convert mol to g:

$$30.0 \text{ mol} \times \frac{62.08 \text{ g}}{\text{mol}} = 1,860 \text{ g}$$

Thus, there are 1,860 g of $C_2H_6O_2$ in the specified amount of engine coolant.

Exercise 10.3.6

a. What volume of a 0.0902 M solution of formic acid (HCOOH) is needed to obtain 0.888 mol of HCOOH?
b. Acetic acid ($HC_2H_3O_2$) is the acid in vinegar. How many grams of $HC_2H_3O_2$ are in 0.565 L of a 0.955 M solution?

Answer available at the end of this section.

Using Molarity in Stoichiometry Problems

Of all the ways of expressing concentration, molarity is the one most commonly used in stoichiometry problems because it is directly related to the mole unit. Consider the following chemical equation:

$$HCl(aq) + NaOH(s) \rightarrow H_2O(\ell) + NaCl(aq)$$

Suppose we want to know how many liters of aqueous HCl solution will react with a given mass of NaOH. A typical approach to answering this question is as follows:




Figure 10.3.5*: Typical approach to solving Molarity problems*

In itself, each step is a straightforward conversion. It is the combination of the steps that is a powerful quantitative tool for problem solving.

Example 10.3.7

How many milliliters of a 2.75 M HCl solution are needed to react with 185 g of NaOH? The balanced chemical equation for this reaction is as follows:

$$HCl(aq) + NaOH(s) \rightarrow H_2O(\ell) + NaCl(aq)$$

Solution

We will follow the flowchart to answer this question. First, we convert the mass of NaOH to moles of NaOH using its molar mass, 40.00 g/mol:

$$185 \text{ g NaOH} \times \frac{1 \text{ mol NaOH}}{40.00 \text{ g NaOH}} = 4.63 \text{ mol NaOH}$$

Using the balanced chemical equation, we see that there is a one-to-one ratio of moles of HCl to moles of NaOH. We use this to determine the number of moles of HCl needed to react with the given amount of NaOH:

$$4.63 \text{ mol NaOH} \times \frac{1 \text{ mol HCl}}{1 \text{ mol NaOH}} = 4.63 \text{ mol HCl}$$

Finally, we use the definition of molarity to determine the volume of 2.75 M HCl needed:

$$2.75 \text{ M HCl} = \frac{4.63 \text{ mol HCl}}{\text{volume of HCl solution}}$$

$$\text{volume of HCl} = \frac{4.63 \text{ mol HCl}}{2.75 \text{ M HCl}} = 1.68 \text{ L} \times \frac{1,000 \text{ mL}}{1 \text{ L}} = 1,680 \text{ mL}$$

We need 1,680 mL of 2.75 M HCl to react with the NaOH.

The same multi-step problem can also be worked out in a single line, rather than as separate steps, as follows:




Figure 10.3.6: Determining the volume (mL) of 2.75 M HCl needed to reactions with a mass of NaOH

$$185\text{ g NaOH} \times \frac{1\text{ mol NaOH}}{40.00\text{ g NaOH}} \times \frac{1\text{ mol HCl}}{1\text{ mol NaOH}} \times \frac{1\text{ L HCl solution}}{2.75\text{ mol HCl}} \times \frac{1000\text{ mL HCl solution}}{1\text{ L HCl solution}} = 1,680\text{ mL HCl solution}$$

Our final answer (rounded off to three significant figures) is 1,680 mL HCl solution.

Exercise 10.3.7

How many milliliters of a 1.04 M H_2SO_4 solution are needed to react with 98.5 g of $Ca(OH)_2$? The balanced chemical equation for the reaction is as follows:

$$H_2SO_{4(aq)} + Ca(OH)_{2(s)} \rightarrow 2H_2O_{(\ell)} + CaSO_{4(aq)}$$

Answer available at the end of this section.

The general steps for performing stoichiometry problems such as this are shown in Figure 10.3.3. You may want to consult this figure when working with solutions in chemical reactions. The double arrows in Figure 10.3.3 indicate that you can start at either end of the chart and, after a series of simple conversions, determine the quantity at the other end.




Figure 10.3.7: Diagram of Steps for Using Molarity in Stoichiometry Calculations. When using molarity in stoichiometry calculations, a specific sequence of steps usually leads you to the correct answer.

Many of the fluids found in our bodies are solutions. The solutes range from simple ionic compounds to complex proteins. Table 10.3.2 lists the typical concentrations of some of these solutes.

*Table 10.3.2: Approximate Concentrations of Various Solutes in Some Solutions in the Body**

Solution	Solute	Concentration (M)
blood plasma	Na^+	0.138
	K^+	0.005
	Ca^{2+}	0.004
	Mg^{2+}	0.003
	Cl^-	0.110
	HCO_3^-	0.030
stomach acid	HCl	0.10
urine	NaCl	0.15
	PO_4^{3-}	0.05
	NH_2CONH_2 (urea)	0.30
***Note: Concentrations are approximate and can vary widely.**		

Looking Closer: The Dose Makes the Poison

Why is it that we can drink 1 qt of water when we are thirsty and not be harmed, but if we ingest 0.5 g of arsenic, we might die? There is an old saying: *the dose makes the poison*. This means that what may be dangerous in some amounts may not be dangerous in other amounts.

Take arsenic, for example. Some studies show that arsenic deprivation limits the growth of animals such as chickens, goats, and pigs, suggesting that arsenic is actually an essential trace element in the diet. Humans are constantly exposed to tiny amounts of arsenic from the environment, so studies of completely arsenic-free humans are not available; if arsenic is an essential trace mineral in human diets, it is probably required on the order of 50 ppb or less. A toxic dose of arsenic corresponds to about 7,000 ppb and higher, which is over 140 times the trace amount that may be required by the body. Thus, arsenic is not poisonous in and of itself. Rather, it is the amount that is dangerous: the dose makes the poison.

Similarly, as much as water is needed to keep us alive, too much of it is also risky to our health. Drinking too much water too fast can lead to a condition called water intoxication, which may be fatal. The danger in water intoxication is not that water itself becomes toxic. It is that the ingestion of too much water too fast dilutes sodium ions, potassium ions, and other salts in the bloodstream to concentrations that are not high enough to support brain, muscle, and heart functions. Military personnel, endurance athletes, and even desert hikers are susceptible to water intoxication if they drink water but do not replenish the salts lost in sweat. As this example shows, even the right substances in the wrong amounts can be dangerous!

Equivalents

Concentrations of ionic solutes are occasionally expressed in units called equivalents (Eq). One equivalent equals 1 mol of positive or negative charge. Thus, 1 mol/L of $Na^+(aq)$ is also 1 Eq/L because sodium has a 1+ charge. A 1 mol/L solution of $Ca^{2+}(aq)$ ions has a concentration of 2 Eq/L because calcium has a 2+ charge. Dilute solutions may be expressed in milliequivalents (mEq)—for example, human blood plasma has a total concentration of about 150 mEq/L. (For more information about the ions present in blood plasma, see Chapter 3, Section 3.3.)

Dilutions

When additional water is added to an aqueous solution, the concentration of that solution decreases. This is because the number of **moles of the solute does not change**, while the **volume of the solution increases**. We can set up an equality between the moles of the solute before the dilution (1) and the moles of the solute after the dilution (2).

$$\text{mol}_1 = \text{mol}_2$$

Since the moles of solute in a solution is equal to the molarity multiplied by the liters, we can set those equal.

$$M_1 \times L_1 = M_2 \times L_2$$

Finally, because the two sides of the equation are set equal to one another, the volume can be in any units we choose, as long as that unit is the same on both sides. Our equation for calculating the molarity of a diluted solution becomes:

$$M_1 \times V_1 = M_2 \times V_2$$

Suppose that you have $100.\ \text{mL}$ of a $2.0\ \text{M}$ solution of HCl. You dilute the solution by adding enough water to make the solution volume $500.\ \text{mL}$. The new molarity can easily be calculated by using the above equation and solving for M_2.

$$M_2 = \frac{M_1 \times V_1}{V_2} = \frac{2.0\ \text{M} \times 100.\ \text{mL}}{500.\ \text{mL}} = 0.40\ \text{M HCl}$$

The solution has been diluted by one-fifth since the new volume is five times as great as the original volume. Consequently, the molarity is one-fifth of its original value.




Figure 10.3.8: The process of dilution shown at both the macroscopic and particle levels, where adding solvent increases the total volume (V) and decreases the molarity (M) while the total amount of solute remains constant. CC BY-SA 4.0.

Dilution Equations

Any units of concentration and volume can be used, as long as both concentrations and both volumes have the same unit. For example, if we are using M (molarity), then we can express the equation as follows:

$$\text{Molarity}_{\text{initial}} \times \text{volume}_{\text{initial}} = \text{Molarity}_{\text{final}} \times \text{volume}_{\text{final}} \text{ or}$$

$$\boldsymbol{M_1V_1 = M_2V_2}$$

If we are using percent, the dilution equation is as follows:

$$\%_{initial} \times volume_{initial} = \%_{final} \times volume_{final} \text{ or}$$

$$\%_1V_1 = \%_2V_2$$

Example 10.3.8

A 125 mL sample of 0.900 M NaCl is diluted to 1,125 mL. What is the final concentration of the diluted solution?

Solution

Because the volume units are the same, and we are looking for the molarity of the final solution, we can use (concentration × volume)$_{initial}$ = (concentration × volume)$_{final}$:

$$(0.900 \text{ M} \times 125 \text{ mL}) = (\text{concentration} \times 1{,}125 \text{ mL})$$

We solve by isolating the unknown concentration by itself on one side of the equation. Dividing by 1,125 mL gives

$$\text{concentration} = \frac{0.900 \text{ M} \times 125 \text{ mL}}{1,125 \text{ mL}} = 0.100 \text{ M}$$

as the final concentration.

Exercise 10.3.8

a. A nurse uses a syringe to inject 5.00 mL of 0.550 M heparin solution (heparin is an anticoagulant drug) into a 250 mL IV bag, for a final volume of 255 mL. What is the concentration of the resulting heparin solution?

b. A 0.885 M solution of KBr whose initial volume is 76.5 mL has more water added until its concentration is 0.500 M. What is the new volume of the solution?

Answer available at the end of this section.

Preparing IV Solutions

In a hospital emergency room, a physician orders an intravenous (IV) delivery of 100 mL of 0.5% KCl for a patient suffering from hypokalemia (low potassium levels). Does an aide run to a supply cabinet and take out an IV bag containing this concentration of KCl?

Not likely. It is more probable that the aide must make the proper solution from an IV bag of sterile solution and a more concentrated, sterile solution, called a *stock solution*, of KCl. The aide is expected to use a syringe to draw up some stock solution and inject it into the waiting IV bag and dilute it to the proper concentration. Thus the aide must perform a dilution calculation.



Figure 10.3.9:*Preparing IV Solution © Thinkstock. Medical personnel commonly must perform dilutions for IV solutions.*

If the stock solution is 10.0% KCl and the final volume and concentration need to be 100 mL and 0.50%, respectively, then it is an easy calculation to determine how much stock solution to use:

$$(10\%)V_1 = (0.50\%)(100\text{ mL})$$

$$V_1 = 5\text{ mL}$$

Of course, the addition of the stock solution affects the total volume of the diluted solution, but the final concentration is likely close enough even for medical purposes.

Medical and pharmaceutical personnel are constantly dealing with dosages that require concentration measurements and dilutions. It is an important responsibility: calculating the *wrong* dose can be useless, harmful, or even fatal!

Key Takeaways

- Various concentration units are used to express the amounts of solute in a solution.
- Concentration units can be used as conversion factors in stoichiometry problems.
- New concentrations can be easily calculated if a solution is diluted.

Answers

Exercise 10.3.1

7.90%

Exercise 10.3.2

$$50.0\text{ g } C_6H_12O_6 \times \frac{100\text{ mL solution}}{12.75\text{ g } C_6H_12O_6} = 392\text{ mL solution}$$

Exercise 10.3.3

$$\%\text{ m/v} = \frac{27.0\text{ g NaOCl}}{500.0\text{ mL solution}} \times 100\% = 5.40\%\text{ m/v}$$

Exercise 10.3.4

0.14 ppm

Exercise 10.3.5

0.00221 M

Exercise 10.3.6

a. 9.84 L

b. 32.4 g

Exercise 10.3.7

1,280 mL

Exercise 10.3.8

a. 0.0108 M

b. 135.4 mL

This page titled 10.3: Concentration is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.2: Concentration by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.4: The Dissolution Process

Learning Objectives

- To describe the dissolution process at the molecular level

The Dissolution Process

What occurs at the molecular level to cause a solute to dissolve in a solvent? The answer depends in part on the solute, but there are some similarities common to all solutes.

Recall the rule that *like dissolves like*. This means that substances must have similar intermolecular forces to form solutions. When a soluble solute is introduced into a solvent, the particles of solute can interact with the particles of solvent. In the case of a solid or liquid solute, the interactions between the solute particles and the solvent particles are so strong that the individual solute particles separate from each other and, surrounded by solvent molecules, enter the solution. (Gaseous solutes already have their constituent particles separated, but the concept of being surrounded by solvent particles still applies.) This process is called **solvation** and is illustrated in Figure 10.4.1. When the solvent is water, the word **hydration**, rather than solvation, is used.




Figure 10.4.1: Solvation. When a solute dissolves, the individual particles of solute become surrounded by solvent particles. Eventually the particle detaches from the remaining solute, surrounded by solvent molecules in solution. Source: Photo © Thinkstock

Ionic Compounds and Covalent Compounds as Solutes

In the case of molecular solutes like glucose, the solute particles are individual molecules. However, if the solute is ionic, the individual ions separate from each other and become surrounded by solvent particles. That is, the cations and anions of an ionic solute separate when the solute dissolves. This process is referred to as **dissociation** (Figure 10.4.1).

The dissociation of soluble ionic compounds gives solutions of these compounds an interesting property: they conduct electricity. Because of this property, soluble ionic compounds are referred to as electrolytes. Many ionic compounds dissociate completely and are therefore called **strong electrolytes**. Sodium chloride is an example of a strong electrolyte. Some compounds dissolve but dissociate only partially, and solutions of such solutes may conduct electricity only weakly. These solutes are called **weak electrolytes**. Acetic acid (CH_3COOH), the compound in vinegar, is a weak electrolyte. Solutes that dissolve into individual neutral molecules without dissociation do not impart additional electrical conductivity to their solutions and are called nonelectrolytes. Table sugar ($C_{12}H_{22}O_{11}$) is an example of a nonelectrolyte.

The term *electrolyte* is used in medicine to mean any of the important ions that are dissolved in aqueous solution in the body. Important physiological electrolytes include Na^+, K^+, Ca^{2+}, Mg^{2+}, and Cl^-.

Example 10.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

1. potassium chloride (KCl)
2. fructose ($C_6H_{12}O_6$)
3. isopropyl alcohol [$CH_3CH(OH)CH_3$]
4. magnesium hydroxide [$Mg(OH)_2$]

Solution

Each substance can be classified as an ionic solute or a nonionic solute. Ionic solutes are electrolytes, and nonionic solutes are nonelectrolytes.

1. Potassium chloride is an ionic compound; therefore, when it dissolves, its ions separate, making it an electrolyte.
2. Fructose is a sugar similar to glucose. (In fact, it has the same molecular formula as glucose.) Because it is a molecular compound, we expect it to be a nonelectrolyte.
3. Isopropyl alcohol is an organic molecule containing the alcohol functional group. The bonding in the compound is all covalent, so when isopropyl alcohol dissolves, it separates into individual molecules but not ions. Thus, it is a nonelectrolyte
4. Magnesium hydroxide is an ionic compound, so when it dissolves it dissociates. Thus, magnesium hydroxide is an electrolyte.

Exercise 10.4.1

The following substances all dissolve to some extent in water. Classify each as an electrolyte or a nonelectrolyte.

a. acetone (CH_3COCH_3)
b. iron(III) nitrate [$Fe(NO_3)_3$]
c. elemental bromine (Br_2)
d. sodium hydroxide (NaOH)

Answer available at the end of this section.

Electrolytes in Body Fluids

Our body fluids are solutions of electrolytes and many other things. The combination of blood and the circulatory system is the *river of life*, because it coordinates all the life functions. When the heart stops pumping in a heart attack, the life ends quickly. Getting the heart restarted as soon as one can is crucial in order to maintain life.

The primary electrolytes required in the body fluid are cations (of calcium, potassium, sodium, and magnesium) and anions (of chloride, carbonates, aminoacetates, phosphates, and iodide). These are nutritionally called **macrominerals**.

Electrolyte balance is crucial to many body functions. Here's some extreme examples of what can happen with an imbalance of electrolytes: elevated potassium levels may result in cardiac arrhythmias; decreased extracellular potassium produces paralysis; excessive extracellular sodium causes fluid retention; and decreased plasma calcium and magnesium can produce muscle spasms of the extremities.

When a patient is dehydrated, a carefully prepared (commercially available) electrolyte solution is required to maintain health and well being. In terms of child health, oral electrolyte is given when a child is dehydrated due to diarrhea. The use of oral electrolyte maintenance solutions, which is responsible for saving millions of lives worldwide over the last 25 years, is one of the most important medical advances in protecting the health of children in the century, explains Juilus G.K. Goepp, MD, assistant

director of the Pediatric Emergency Department of the Children's Center at Johns Hopkins Hospital. If a parent provides an oral electrolyte maintenance solution at the very start of the illness, dehydration can be prevented. The functionality of electrolyte solutions is related to their properties, and interest in electrolyte solutions goes far beyond chemistry.

Sports drinks are designed to rehydrate the body after excessive fluid depletion. Electrolytes in particular promote normal rehydration to prevent fatigue during physical exertion. Are they a good choice for achieving the recommended fluid intake? Are they performance and endurance enhancers like they claim? Who should drink them?

Typically, eight ounces of a sports drink provides between fifty and eighty calories and 14 to 17 grams of carbohydrate, mostly in the form of simple sugars. Sodium and potassium are the most commonly included electrolytes in sports drinks, with the levels of these in sports drinks being highly variable. The American College of Sports Medicine says a sports drink should contain 125 milligrams of sodium per 8 ounces as it is helpful in replenishing some of the sodium lost in sweat and promotes fluid uptake in the small intestine, improving hydration.

Gatorade

In the summer of 1965, the assistant football coach of the University of Florida Gators requested scientists affiliated with the university study why the withering heat of Florida caused so many heat-related illnesses in football players and provide a solution to increase athletic performance and recovery post-training or game. The discovery was that inadequate replenishment of fluids, carbohydrates, and electrolytes was the reason for the “wilting” of their football players. Based on their research, the scientists concocted a drink for the football players containing water, carbohydrates, and electrolytes and called it “Gatorade.” In the next football season the Gators were nine and two and won the Orange Bowl. The Gators’ success launched the sports-drink industry, which is now a multibillion-dollar industry that is still dominated by Gatorade.



Figure 10.4.2: University of Florida football player Chip Hinton testing Gatorade in 1965, pictured next to the leader of its team of inventors, Robert Cade.

Key Takeaway

- When a solute dissolves, its individual particles are surrounded by solvent molecules and are separated from each other.

Answers

Exercise 10.4.1

a. nonelectrolyte
b. electrolyte
c. nonelectrolyte
d. electrolyte

This page titled 10.4: The Dissolution Process is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.3: The Dissolution Process by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.5: Properties of Solutions

Learning Objectives

- To describe how the properties of solutions differ from those of pure solvents.

Solutions are likely to have properties similar to those of their major component—usually the solvent. However, some solution properties differ significantly from those of the solvent. Here, we will focus on liquid solutions that have a solid solute, but many of the effects we will discuss in this section are applicable to all solutions.

Colligative Properties

Solutes affect some properties of solutions that depend only on the concentration of the dissolved particles. These properties are called **colligative properties**. Four important colligative properties that we will examine here are vapor pressure depression, boiling point elevation, freezing point depression, and osmotic pressure.

Molecular compounds separate into individual molecules when they are dissolved, so for every 1 mol of molecules dissolved, we get 1 mol of particles. In contrast, **ionic** compounds separate into their constituent ions when they dissolve, so 1 mol of an ionic compound will produce more than 1 mol of dissolved particles. For example, **every mole of NaCl** that dissolves yields **1 mol of Na^+ ions** and **1 mol of Cl^- ions**, for a **total of 2 mol** of particles in solution. Thus, the effect on a solution's properties by dissolving NaCl may be twice as large as the effect of dissolving the same amount of moles of glucose ($C_6H_{12}O_6$).

Vapor Pressure Depression

All liquids evaporate. In fact, given enough volume, a liquid will turn completely into a vapor. If enough volume is not present, a liquid will evaporate only to the point where the rate of evaporation equals the rate of vapor condensing back into a liquid. The pressure of the vapor at this point is called the **vapor pressure** of the liquid.

The presence of a dissolved solid lowers the characteristic vapor pressure of a liquid so that it evaporates more slowly. (The exceptions to this statement are if the solute itself is a liquid or a gas, in which case the solute will also contribute something to the evaporation process. We will not discuss such solutions here.) This property is called vapor pressure depression and is depicted in Figure 10.5.1.




Figure 10.5.1: Vapor Pressure Lowering in a Solution. A comparison between a pure solvent (left) and a solution containing a non-volatile solute (right). The solute particles (orange) occupy a portion of the liquid's surface area, reducing the number of solvent molecules available to evaporate, which results in a lower equilibrium vapor pressure.

Boiling Point and Freezing Point Effects

A related property of solutions is that their boiling points are higher than the boiling point of the pure solvent. Because the presence of solute particles decreases the vapor pressure of the liquid solvent, a higher temperature is needed to reach the boiling point. This phenomenon is called boiling point elevation. **For every mole of particles dissolved in a liter of water, the boiling point of water increases by about 0.5°C.** The addition of one mole of sucrose (molecular compound) in one liter of water will raise the boiling point from 100^{0}C to 100.5^{0}C but the addition of one mole of NaCl in one liter of water will raise the boiling point by 2 x 0.5^{0}C = 1^{0}C. Furthermore, the addition of one mole of $\mathrm{CaCl_2}$ in one liter of water will raise the boiling point by 3 x 0.5^{0}C = 1.5^{0}C.

> Some people argue that putting a pinch or two of salt in water used to cook spaghetti or other pasta makes a solution that has a higher boiling point, so the pasta cooks faster. In actuality, the amount of solute is so small that the boiling point of the water is practically unchanged.

The presence of solute particles has the opposite effect on the freezing point of a solution. When a solution freezes, only the solvent particles come together to form a solid phase, and the presence of solute particles interferes with that process. Therefore, for the liquid solvent to freeze, more energy must be removed from the solution, which lowers the temperature. Thus, solutions have lower freezing points than pure solvents do. This phenomenon is called freezing point depression. **For every mole of particles in a liter of water, the freezing point decreases by about 1.9°C.**




Figure 10.5.2: Comparison of boiling and freezing points of a pure liquid (right side) with a solution (left side).

Both boiling point elevation and freezing point depression have practical uses. For example, solutions of water and ethylene glycol ($C_2H_6O_2$) are used as coolants in automobile engines because the boiling point of such a solution is greater than 100°C, the normal boiling point of water. In winter, salts like NaCl and $CaCl_2$ are sprinkled on the ground to melt ice or keep ice from forming on roads and sidewalks (Figure 10.5.2). This is because the solution made by dissolving sodium chloride or calcium chloride in water has a lower freezing point than pure water, so the formation of ice is inhibited.



Figure 10.5.2: Effect of Freezing Point Depression. The salt sprinkled on this sidewalk makes the water on the sidewalk have a lower freezing point than pure water, so it does not freeze as easily. This makes walking on the sidewalk less hazardous in winter. © Thinkstock

Example 10.5.1

Which solution's freezing point deviates more from that of pure water—a 1 M solution of NaCl or a 1 M solution of $CaCl_2$?

Solution

Colligative properties depend on the number of dissolved particles, so the solution with the greater number of particles in solution will show the greatest deviation. When NaCl dissolves, it separates into two ions, Na^+ and Cl^-. But when $CaCl_2$ dissolves, it separates into three ions—one Ca^{2+} ion and two Cl^- ions. Thus, mole for mole, $CaCl_2$ will have 50% more impact on freezing point depression than NaCl.

Exercise 10.5.1

Which solution's boiling point deviates more from that of pure water—a 1 M solution of $CaCl_2$ or a 1 M solution of $MgSO_4$?

Answer available at the end of this section.

Example 10.5.2

Estimate the boiling point of 0.2 M $CaCl_2$ solution.

Solution

The boiling point increases 0.5^0C for every mole of solute per liter of water. For this estimation, let's assume that 1 liter of solution is roughly the same volume as 1 liter of water. A 0.2 M $CaCl_2$ solution contains 0.2 moles of $CaCl_2$ solution formula units per liter of solution. Each $CaCl_2$ unit separates into three ions.

$$0.2 \text{ mol } CaCl_2 \times \frac{3 \text{ mol ions}}{1 \text{ mol } CaCl_2} \times \frac{0.5 \text{ deg C}}{1 \text{ mol ion}} = 0.3 \text{ deg C}$$

The normal boiling point of water is 100^0C, so the boiling point of the solution is raised to 100.3^0C.

Exercise 10.5.2

Estimate the freezing point of 0.3 M $CaCl_2$ solution.

Answer available at the end of this section.

Osmotic Pressure

The last colligative property of solutions we will consider is a very important one for biological systems. It involves **osmosis**, the process by which solvent molecules can pass through certain membranes but solute particles cannot. When two solutions of different concentration are present on either side of these membranes (called *semipermeable membranes*), there is a tendency for solvent molecules to move from the more dilute solution to the more concentrated solution until the concentrations of the two solutions are equal. This tendency is called **osmotic pressure**. External pressure can be exerted on a solution to counter the flow of solvent; the pressure required to halt the osmosis of a solvent is equal to the osmotic pressure of the solution.

Osmolarity (osmol) is a way of reporting the total number of particles in a solution to determine osmotic pressure. It is defined as the molarity of a solute times the number of particles a formula unit of the solute makes when it dissolves (represented by i):

$$osmol = M \times i \tag{10.5.1}$$

If more than one solute is present in a solution, the individual osmolarities are additive to get the total osmolarity of the solution. Solutions that have the same osmolarity have the same osmotic pressure. If solutions of differing osmolarities are present on opposite sides of a semipermeable membrane, solvent will transfer from the lower-osmolarity solution to the higher-osmolarity solution. Counterpressure exerted on the high-osmolarity solution will reduce or halt the solvent transfer. An even higher pressure can be exerted to force solvent from the high-osmolarity solution to the low-osmolarity solution, a process called *reverse osmosis*.

Reverse osmosis is used to make potable water from saltwater where sources of fresh water are scarce.

Example 10.5.3

A 0.50 M NaCl aqueous solution and a 0.30 M $Ca(NO_3)_2$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Solution

The solvent will flow into the solution of higher osmolarity. The NaCl solute separates into two ions—Na^+ and Cl^-—when it dissolves, so its osmolarity is as follows:

$$\text{osmol (NaCl)} = 0.50\ \text{M} \times 2 = 1.0\ \text{osmol}$$

The $Ca(NO_3)_2$ solute separates into three ions—one Ca^{2+} and two NO_3^-—when it dissolves, so its osmolarity is as follows:

$$\text{osmol } [Ca(NO_3)_2] = 0.30\ \text{M} \times 3 = 0.90\ \text{osmol}$$

The osmolarity of the $Ca(NO_3)_2$ solution is lower than that of the NaCl solution, so water will transfer through the membrane from the $Ca(NO_3)_2$ solution to the NaCl solution.

Exercise 10.5.3

A 1.5 M $C_6H_{12}O_6$ aqueous solution and a 0.40 M $Al(NO_3)_3$ aqueous solution are placed on opposite sides of a semipermeable membrane. Determine the osmolarity of each solution and predict the direction of solvent flow.

Answer available at the end of this section.

To Your Health: Dialysis

The main function of the kidneys is to filter the blood to remove wastes and extra water, which are then expelled from the body as urine. Some diseases rob the kidneys of their ability to perform this function, causing a buildup of waste materials in the bloodstream. If a kidney transplant is not available or desirable, a procedure called dialysis can be used to remove waste materials and excess water from the blood.

In one form of dialysis, called *hemodialysis*, a patient's blood is passed though a length of tubing that travels through an *artificial kidney machine* (also called a *dialysis machine*). A section of tubing composed of a semipermeable membrane is immersed in a solution of sterile water, glucose, amino acids, and certain electrolytes. The osmotic pressure of the blood forces waste molecules and excess water through the membrane into the sterile solution. Red and white blood cells are too large to pass through the membrane, so they remain in the blood. After being cleansed in this way, the blood is returned to the body.



Figure 10.5.3: A patient undergoing hemodialysis depends on osmosis to cleanse the blood of waste products that the kidneys are incapable of removing due to disease. from Wikipedia.

Dialysis is a continuous process, as the osmosis of waste materials and excess water takes time. Typically, 5–10 lb of waste-containing fluid is removed in each dialysis session, which can last 2–8 hours and must be performed several times a week. Although some patients have been on dialysis for 30 or more years, dialysis is always a temporary solution because waste materials are constantly building up in the bloodstream. A more permanent solution is a kidney transplant.

Cell walls are semipermeable membranes, so the osmotic pressures of the body's fluids have important biological consequences. If solutions of different osmolarity exist on either side of the cells, solvent (water) may pass into or out of the cells, sometimes with disastrous results. Consider what happens if red blood cells are placed in a *hypotonic* solution, meaning a solution of lower osmolarity than the liquid inside the cells. The cells swell up as water enters them, disrupting cellular activity and eventually causing the cells to burst. This process is called *hemolysis*. If red blood cells are placed in a *hypertonic* solution, meaning one having a higher osmolarity than exists inside the cells, water leaves the cells to dilute the external solution, and the red blood cells shrivel and die. This process is called *crenation*. Only if red blood cells are placed in *isotonic* solutions that have the same osmolarity as exists inside the cells are they unaffected by negative effects of osmotic pressure. **Glucose solutions of about 0.31 M, or sodium chloride solutions of about 0.16 M, are isotonic with blood plasma.**

The concentration of an isotonic sodium chloride (NaCl) solution is only half that of an isotonic glucose ($C_6H_{12}O_6$) solution because NaCl produces two ions when a formula unit dissolves, while molecular $C_6H_{12}O_6$ produces only one particle when a formula unit dissolves. The osmolarities are therefore the same even though the concentrations of the two solutions are different.

Osmotic pressure explains why you should not drink seawater if you are abandoned in a life raft in the middle of the ocean. Its osmolarity is about three times higher than most bodily fluids. You would actually become thirstier as water from your cells was drawn out to dilute the salty ocean water you ingested. Our bodies do a better job coping with hypotonic solutions than with hypertonic ones. The excess water is collected by our kidneys and excreted.

Osmotic pressure effects are used in the food industry to make pickles from cucumbers and other vegetables and in brining meat to make corned beef. It is also a factor in the mechanism of getting water from the roots to the tops of trees!

Career Focus: Perfusionist

A perfusionist is a medical technician trained to assist during any medical procedure in which a patient's circulatory or breathing functions require support. The use of perfusionists has grown rapidly since the advent of open-heart surgery in 1953.

Most perfusionists work in operating rooms, where their main responsibility is to operate heart-lung machines. During many heart surgeries, the heart itself must be stopped. In these situations, a heart-lung machine keeps the patient alive by aerating the blood with oxygen and removing carbon dioxide. The perfusionist monitors both the machine and the status of the blood, notifying the surgeon and the anesthetist of any concerns and taking corrective action if the status of the blood becomes abnormal.

Despite the narrow parameters of their specialty, perfusionists must be highly trained. Certified perfusion education programs require a student to learn anatomy, physiology, pathology, chemistry, pharmacology, math, and physics. A college degree is usually required. Some perfusionists work with other external artificial organs, such as hemodialysis machines and artificial livers.

Answers

Exercise 10.5.1

$CaCl_2$

Exercise 10.5.2

minus 1.7°C

Exercise 10.5.3

osmol $C_6H_{12}O_6$ = 1.5; osmol $Al(NO_3)_3$ = 1.6

The solvent flows from $C_6H_{12}O_6$ solution (lower osmolarity) to $Al(NO_3)_3$ solution (higher osmolarity).

This page titled 10.5: Properties of Solutions is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.4: Properties of Solutions by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.6: Chemical Equilibrium

Learning Outcomes

- Explain chemical equilibrium.
- Write expression for calculating K.
- Calculate and compare Q and K values.
- Predict relative amounts of reactants and products based on equilibrium constant K.

Hydrogen and iodine gases react to form hydrogen iodide according to the following reaction:

$$\mathrm{H_2\,(g) + I_2\,(g) \rightleftharpoons 2HI\,(g)}$$

$$\text{Forward reaction: } \mathrm{H_2\,(g) + I_2\,(g) \rightarrow 2HI\,(g)} \tag{10.6.1}$$
$$\text{Reverse reaction: } \mathrm{2HI\,(g) \rightarrow H_2\,(g) + I_2\,(g)} \tag{10.6.2}$$

Initially, only the forward reaction occurs because no HI is present. As soon as some HI has formed, it begins to decompose back into $\mathrm{H_2}$ and $\mathrm{I_2}$. Gradually, the rate of the forward reaction decreases while the rate of the reverse reaction increases. Eventually the rate of combination of $\mathrm{H_2}$ and $\mathrm{I_2}$ to produce HI becomes equal to the rate of decomposition of HI into $\mathrm{H_2}$ and $\mathrm{I_2}$. When the rates of the forward and reverse reactions have become equal to one another, the reaction has achieved a state of balance. **Chemical equilibrium** is the state of a system in which the rate of the forward reaction is equal to the rate of the reverse reaction.




Figure 10.6.1: Equilibrium in reaction: $\mathrm{H_2\,(g) + I_2\,(g) \rightleftharpoons 2HI\,(g)}$.

Chemical equilibrium can be attained whether the reaction begins with all reactants and no products, all products and no reactants, or some of both. The figure below shows changes in concentration of $\mathrm{H_2}$, $\mathrm{I_2}$, and HI for two different reactions. In the reaction depicted by the graph on the left (A), the reaction begins with only $\mathrm{H_2}$ and $\mathrm{I_2}$ present. There is no HI initially. As the reaction proceeds towards equilibrium, the concentrations of the $\mathrm{H_2}$ and $\mathrm{I_2}$ gradually decrease, while the concentration of the HI gradually increases. When the curve levels out and the concentrations all become constant, equilibrium has been reached. At equilibrium, concentrations of all substances are constant.

In reaction B, the process begins with only HI and no $\mathrm{H_2}$ or $\mathrm{I_2}$. In this case, the concentration of HI gradually decreases while the concentrations of $\mathrm{H_2}$ and $\mathrm{I_2}$ gradually increase until equilibrium is again reached. Notice that in both cases, the relative position of equilibrium is the same, as shown by the relative concentrations of reactants and products. The concentration of HI at equilibrium is significantly higher than the concentrations of $\mathrm{H_2}$ and $\mathrm{I_2}$. This is true whether the reaction began with all reactants or all products. The position of equilibrium is a property of the particular reversible reaction and does not depend upon how

equilibrium was achieved.




Figure 10.6.2: Equilibrium between reactants and products is achieved regardless of whether the reaction starts with the reactants or products.

Conditions for Equilibrium and Types of Equilibrium

It may be tempting to think that once equilibrium has been reached, the reaction stops. Chemical equilibrium is a dynamic process. The forward and reverse reactions continue to occur even after equilibrium has been reached. However, because the rates of the reactions are the same, there is no change in the relative concentrations of reactants and products for a reaction that is at equilibrium. The conditions and properties of a system at equilibrium are summarized below.

1. The system must be closed, meaning no substances can enter or leave the system.
2. Equilibrium is a dynamic process. Even though we don't necessarily see the reactions, both forward and reverse are taking place.
3. The rates of the forward and reverse reactions must be equal.
4. The amount of reactants and products do not have to be equal. However, after equilibrium is attained, the amounts of reactants and products will be constant.

The description of equilibrium in this concept refers primarily to equilibrium between reactants and products in a chemical reaction. Other types of equilibrium include phase equilibrium and solution equilibrium. A phase equilibrium occurs when a substance is in equilibrium between two states. For example, a stoppered flask of water attains equilibrium when the rate of evaporation is equal to the rate of condensation. A solution equilibrium occurs when a solid substance is in a saturated solution. At this point, the rate of dissolution is equal to the rate of recrystallization. Although these are all different types of transformations, most of the rules regarding equilibrium apply to any situation in which a process occurs reversibly.




Figure 10.6.3: These jugglers provide an illustration of dynamic equilibrium. Each throws clubs to the other at the same rate at which he receives clubs from that person. Because clubs are thrown continuously in both directions, the number of clubs moving in each direction is constant, and the number of clubs each juggler has at a given time remains (roughly) constant.

Red blood cells transport oxygen to the tissues so they can function. In the absence of oxygen, cells cannot carry out their biochemical responsibilities. Oxygen moves to the cells attached to hemoglobin, a protein found in the red cells. In cases of carbon monoxide poisoning, CO binds much more strongly to the hemoglobin, blocking oxygen attachment and lowering the amount of oxygen reaching the cells. Treatment involves the patient breathing pure oxygen to displace the carbon monoxide. The equilibrium reaction shown below illustrates the shift toward the right when excess oxygen is added to the system:

$$\mathrm{Hb(CO)_4}\,(aq) + 4\mathrm{O_2}\,(g) \rightleftharpoons \mathrm{Hb(O_2)_4}\,(aq) + 4\mathrm{CO}\,(g)$$

Equilibrium Constant

Consider the hypothetical reversible reaction in which reactants A and B react to form products C and D. This equilibrium can be shown below, where the lowercase letters represent the coefficients of each substance.

$$a\mathrm{A} + b\mathrm{B} \rightleftharpoons c\mathrm{C} + d\mathrm{D}$$

As we have established, the rates of the forward and reverse reactions are the same at equilibrium, and so the concentrations of all of the substances are constant. Since that is the case, it stands to reason that a ratio of the concentration for any given reaction at equilibrium maintains a constant value. The **equilibrium constant** (K_{eq}) is the ratio of the mathematical product of the products of a reaction to the mathematical product of the concentrations of the reactants of the reaction. Each concentration is raised to the power of its coefficient in the balanced chemical equation. For the general reaction above, the equilibrium constant expression is written as follows:

$$K_{\mathrm{eq}} = \frac{[\mathrm{C}]^c\,[\mathrm{D}]^d}{[\mathrm{A}]^a\,[\mathrm{B}]^b}$$

The concentrations of each substance, indicated by the square brackets around the formula, are measured in molarity units $(\mathrm{mol/L})$.

The value of the equilibrium constant for any reaction is only determined by experiment. As detailed in the above section, the position of equilibrium for a given reaction does not depend on the starting concentrations and so the value of the equilibrium constant is truly constant. It does, however, depend on the temperature of the reaction. This is because equilibrium is defined as a condition resulting from the rates of forward and reverse reactions being equal. If the temperature changes, the corresponding change in those reaction rates will alter the equilibrium constant. For any reaction in which a K_{eq} is given, the temperature should be specified.

The equilibrium constant can vary over a wide range of values. Because products are in the numerator of the equilibrium constant expression and reactants are in the denominator, values of K greater than 10^3 indicate a strong tendency for reactants to form products. In this case, chemists say that equilibrium lies to the right as written, favoring the formation of products. An example is the reaction between H_2 and Cl_2 to produce HCl, which has an equilibrium constant of 1.6×10^{33} at 300 K. Because H_2 is a good reducing agent and Cl_2 is a good oxidizing agent, the reaction proceeds essentially to completion.

$$H_{2(g)} + Cl_{2(g)} \rightleftharpoons 2HCl_{(g)} \quad K=1.6 \times 10^{33} \text{ at } 300K$$

In contrast, values of K less than 10^{-3} indicate that the ratio of products to reactants at equilibrium is very small. That is, reactants do not tend to form products readily, and the equilibrium lies to the left as written, favoring the formation of reactants. An example is the decomposition of water. No wonder you won't find water a very good source of oxygen gas and hydrogen gas at ordinary temperatures!

$$H_2O_{(g)} \rightleftharpoons H_{2(g)} + \tfrac{1}{2}O_{2(g)} \quad K=8 \times 10^{-41} \text{ at } 25°C$$

Figure 10.6.4 summarizes the relationship between the magnitude of K and the relative concentrations of reactants and products at equilibrium for a general reaction, written as reactants ⇌ products.




Figure 10.6.4: The Relationship between the Composition of the Mixture at Equilibrium and the Magnitude of the Equilibrium Constant. The larger the K, the farther the reaction proceeds to the right before equilibrium is reached, and the greater the ratio of products to reactants at equilibrium.

A large value of the equilibrium constant K means that products predominate at equilibrium; a small value means that reactants predominate at equilibrium.

- When K is a very **large** number, the reaction is essentially irreversible, products predominate at equilibrium.
- When K is between 1 and 1000, more products than reactants are present at equilibrium
- When K is between 1 and 0.001, more reactants than products are present at equilibrium.
- When K is a very **small** number, the reaction produces almost no products; only reactants are present at equilibrium.

Example 10.6.1

Write the expression for the equilibrium constant K for the following reaction:

N2 *(g)* + 3 H2 *(g)* ⇄ 2 NH3 *(g)*

Solution

The equilibrium constant equation is the ratio of the concentration of the products (NH_3) to the concentration of the reactants (N_2 and H_2), each raised to the power of its coefficient in the balanced chemical equation.

$$K = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

Exercise 10.6.1

Write the expression of the equilibrium constant K for the following reaction:.

2 NOCl *(g)* ⇄ 2 NO *(g)* + Cl_2 *(g)*

Answer available at the end of this section.

Example 10.6.2

Predict which systems at equilibrium will (a) contain essentially only products, (b) contain essentially only reactants, and (c) contain appreciable amounts of both products and reactants.

1. $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad K_{(700K)} = 54$
2. $2CO_{2(g)} \rightleftharpoons 2CO_{(g)} + O_{2(g)} \qquad K_{(1200K)} = 3.1 \times 10^{-18}$
3. $PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)} \qquad K_{(613K)} = 97$
4. $2O_{3(g)} \rightleftharpoons 3O_{2(g)} \qquad K_{(298K)} = 5.9 \times 10^{55}$

Solution

Use the value of the equilibrium constant to determine whether the equilibrium mixture will contain essentially only products, essentially only reactants, or significant amounts of both.

a. Only system 4 has $K \gg 10^3$, so at equilibrium it will consist of essentially only products.
b. System 2 has $K \ll 10^{-3}$, so the reactants have little tendency to form products under the conditions specified; thus, at equilibrium the system will contain essentially only reactants.
c. Both systems 1 and 3 have equilibrium constants in the range $10^3 \geq K \geq 10^{-3}$, indicating that the equilibrium mixtures will contain appreciable amounts of both products and reactants.

Exercise 10.6.2

Hydrogen and nitrogen react to form ammonia according to the following balanced chemical equation:

$$3H_{2(g)} + N_{2(g)} \rightleftharpoons 2NH_{3(g)}$$

Values of the equilibrium constant at various temperatures were reported as

- $K_{25°C} = 3.3 \times 10^8$,
- $K_{177°C} = 2.6 \times 10^3$, and
- $K_{327°C} = 4.1$.

a. At which temperature would you expect to find the highest proportion of H_2 and N_2 in the equilibrium mixture?
b. At which temperature would you expect to find the highest proportion of ammonia in the equilibrium mixture?

Answer available at the end of this section.

Reaction Quotient

The reaction quotient, Q, is used when questioning if we are at equilibrium. The calculation for Q is **exactly** the same as for K but we can only use K when we know we are at equilibrium. Comparing Q and K allows the direction of the reaction to be predicted.

- Q = K equilibrium
- Q < K reaction proceeds to the right to form more products and decrease amount of reactants so value of Q will increase
- Q > K reaction proceeds to the left to form more reactants and decrease amount of products so value of Q will decrease

Key Takeaway

- As a chemical change proceeds, the quantities of the reactants/products will decrease, and those of the products/reactants will increase. Eventually the reaction slows down and the composition of the system stops changing. At this point the reaction is in its *equilibrium state*, and no further change in composition will occur as long as the system is left undisturbed.

Contributors and Attributions

-

- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)

Answers

Exercise $10.6.1$

$$K = \frac{[NO]^2[Cl_2]}{[NOCl]^2}$$

Exercise $10.6.2$

a. 327°C, where K is smallest

b. 25°C

This page titled 10.6: Chemical Equilibrium is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.5: Chemical Equilibrium by CK12 is licensed CK-12. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

10.7: Le Chatelier's Principle

Learning Outcomes

- Define Le Chatelier's principle.
- Predict how the change in amounts of substances, temperature, or pressure will affect amounts of reactants and products present at equilibrium.

Le Chatelier's Principle

Chemical equilibrium was studied by the French chemist Henri Le Chatelier (1850 - 1936) and his description of how a system responds to a stress to equilibrium has become known as **Le Chatelier's principle**: When a chemical system that is at equilibrium is disturbed by a stress, the system will respond in order to relieve the stress. Stresses to a chemical system involve changes in the concentrations of reactants or products, changes in the temperature of the system, or changes in the pressure of the system. We will discuss each of these stresses separately. The change to the equilibrium position in every case is either a favoring of the forward reaction or a favoring of the reverse reaction. When the forward reaction is favored, the concentrations of products increase, while the concentrations of reactants decrease. When the reverse reaction is favored, the concentrations of the products decrease, while the concentrations of reactants increase.

Original Equilibrium	Favored Reaction	Result
$\mathrm{A} \rightleftharpoons \mathrm{B}$	Forward: $\mathrm{A} \rightarrow \mathrm{B}$	[A] decreases; [B] increases
$\mathrm{A} \rightleftharpoons \mathrm{B}$	Reverse: $\mathrm{A} \leftarrow \mathrm{B}$	[A] increases; [B] decreases



Figure 10.7.1: *Henri Le Chatelier.*

Effect of Concentration

A change in concentration of one of the substances in an equilibrium system typically involves either the addition or the removal of one of the reactants or products. Consider the Haber-Bosch process for the industrial production of ammonia from nitrogen and hydrogen gases.

$$\mathrm{N_2}\,(g) + 3\mathrm{H_2}\,(g) \rightleftharpoons 2\mathrm{NH_3}\,(g)$$

If the concentration of one substance in a system is increased, the system will respond by favoring the reaction that removes that substance. When more $\mathrm{N_2}$ is added, the forward reaction will be favored because the forward reaction uses up $\mathrm{N_2}$ and converts it to $\mathrm{NH_3}$. The forward reaction speeds up temporarily as a result of the addition of a reactant. The position of equilibrium shifts as more $\mathrm{NH_3}$ is produced. The concentration of $\mathrm{NH_3}$ increases, while the concentrations of $\mathrm{N_2}$ and $\mathrm{H_2}$ decrease. After some time passes, equilibrium is reestablished with new concentrations of all three substance. As can be seen in the figure below, if more $\mathrm{N_2}$ is added, a new equilibrium is achieved by the system. The new concentration of $\mathrm{NH_3}$ is higher because of the favoring of the forward reaction. The new concentration of the $\mathrm{H_2}$ is lower .The concentration of $\mathrm{N_2}$ is higher than in the original equilibrium, but

went down slightly following the addition of the N_2 that disturbed the original equilibrium. By responding in this way, the value of the equilibrium constant for the reaction, K_{eq}, does not change as a result of the stress to the system.

In other words, the amount of each substance is different but the ratio of the amount of each remains the same.

If more NH_3 were added, the reverse reaction would be favored. This "favoring" of a reaction means temporarily speeding up the reaction in that direction until equilibrium is reestablished. Recall that once equilibrium is reestablished, the rates of the forward and reverse reactions are again equal. The addition of NH_3 would result in increased formation of the reactants, N_2 and H_2.




Figure 10.7.2: The Haber-Bosch process is an equilibrium between reactants N_2 and H_2 and product NH_3.

An equilibrium can also be disrupted by the removal of one of the substances. If the concentration of a substance is decreased, the system will respond by favoring the reaction that replaces that substance. In the industrial Haber-Bosch process, NH_3 is removed from the equilibrium system as the reaction proceeds. As a result, the forward reaction is favored so that more NH_3 is produced. The concentrations of N_2 and H_2 decrease. Continued removal of NH_3 will eventually force the reaction to go to completion until all of the reactants are used up. If either N_2 or H_2 were removed from the equilibrium system, the reverse reaction would be favored and the concentration of NH_3 would decrease.

The effect of changes in concentration on an equilibrium system according to Le Chatelier's principle is summarized in the table below.

Table 10.7.1*:*

Stress	Response
addition of reactant	forward reaction favored
addition of product	reverse reaction favored
removal of reactant	reverse reaction favored
removal of product	forward reaction favored

Example 10.7.1

Given this reaction at equilibrium:

$$N_2 + 3H_2 \rightleftharpoons 2NH_3$$

How will it affect the reaction if the equilibrium is stressed by each change?

1. H2 is added.
2. NH3 is added.
3. NH3 is removed.

Solution

1. If H2 is added, there is now more reactant, so the reaction will shift to the right (toward products) to reduce the added H2.
2. If NH3 is added, there is now more product, so the reaction will shift to the left (toward reactants) to reduce the added NH3.
3. If NH3 is removed, there is now less product, so the reaction will shift to the right (toward products) to replace the product removed.

Exercise 10.7.1

Given this reaction at equilibrium:

$$CO(g) + Br_2(g) \rightleftharpoons COBr_2(g)$$

How will it affect the reaction if the equilibrium is stressed by each change?

1. Br2 is removed.
2. COBr2 is added.

Answer available at the end of this section.

Effect of Temperature

Increasing or decreasing the temperature of a system at equilibrium is also a stress to the system. The equation for the Haber-Bosch process is written again below, as a thermochemical equation (i.e. it contains information about the energy gained or lost when the reaction occurs).

$$\mathrm{N_2}\,(g) + 3\mathrm{H_2}\,(g) \rightleftharpoons 2\mathrm{NH_3}\,(g) + 91\ \mathrm{kJ}$$

The forward reaction is the exothermic direction: the formation of $\mathrm{NH_3}$ releases heat which is why that is shown as a product. The reverse reaction is the endothermic direction: as $\mathrm{NH_3}$ decomposes to $\mathrm{N_2}$ and $\mathrm{H_2}$, heat is absorbed. An increase in the temperature for this is like adding a product because heat is being released by the reaction. If we add a product then the reaction proceeds towards the formation of more reactants. Reducing the temperature for this system would be similar to removing a product which would favor the formation of more products. The amount of $\mathrm{NH_3}$ will increase and the amount of $\mathrm{N_2}$ and $\mathrm{H_2}$ will decrease.

For changes in concentration, the system responds in such a way that the value of the equilibrium constant, K_{eq} is unchanged. However, a change in temperature shifts the equilibrium and the K_{eq} value either increases or decreases. As discussed in the previous section, values of K_{eq} are dependent on the temperature. When the temperature of the system for the Haber-Bosch process is increased, the resultant shift in equilibrium towards the reactants means that the K_{eq} value decreases. When the temperature is decreased, the shift in equilibrium towards the products means that the K_{eq} value increases.

Le Chatelier's principle as related to temperature changes can be illustrated easily be the reaction in which dinitrogen tetroxide is in equilibrium with nitrogen dioxide.

$$\mathrm{N_2O_4}\,(g) + \mathrm{heat} \rightleftharpoons 2\mathrm{NO_2}\,(g)$$

Dinitrogen tetroxide $(\mathrm{N_2O_4})$ is colorless, while nitrogen dioxide $(\mathrm{NO_2})$ is dark brown in color. When $\mathrm{N_2O_4}$ breaks down into $\mathrm{NO_2}$, heat is absorbed (endothermic) according to the forward reaction above. Therefore, an increase in temperature (adding heat) of the system will favor the forward reaction. Conversely, a decrease in temperature (removing heat) will favor the reverse reaction.

Example 10.7.2

Predict the effect of increasing the temperature on this equilibrium.

$$PCl_3 + Cl_2 \rightleftharpoons PCl_5 + 60kJ$$

Solution

Because energy is listed as a product, it is being produced, so the reaction is exothermic. If the temperature is increasing, a product is being added to the equilibrium, so the equilibrium shifts to minimize the addition of extra product: it shifts to the left (back toward reactants).

Exercise 10.7.2

Predict the effect of decreasing the temperature on this equilibrium.

$$N_2O_4 + 57kJ \rightleftharpoons 2NO_2$$

Answer available at the end of this section.

Effect of Pressure

Changing the pressure of an equilibrium system in which gases are involved is also a stress to the system. A change in the pressure on a liquid or a solid has a negligible effect. We will return again the equilibrium for the Haber-Bosch process. Imagine the gases are contained in a closed system in which the volume of the system is controlled by an adjustable piston as shown in the figure below.




Figure 10.7.3: Effect of pressure on equilibrium. (A) is at equilibrium, (B) increase of pressure through decreased volume, and (C) equilibrium is reestablished.

On the far left, the reaction system contains primarily $\mathrm{N_2}$ and $\mathrm{H_2}$, with only one molecule of $\mathrm{NH_3}$ present. As the piston is pushed inwards, the pressure of the system increases according to Boyle's law. This is a stress to the equilibrium. In the middle image, the same number of molecules is now confined in a smaller space and so the pressure has increased. According to Le Chatelier's principle, the system responds in order to relieve the stress. In the image on the right, the forward reaction has been favored and more $\mathrm{NH_3}$ is produced. The overall result is a decrease in the number of gas molecules in the entire system. This in turn decreases the pressure and provides a relief to the original stress of a pressure increase. An increase in pressure on an equilibrium system favors the reaction which products fewer total moles of gas. In this case, it is the forward reaction that is favored.

A decrease in pressure on the above system could be achieved by pulling the piston outward, increasing the container volume. The equilibrium would respond by favoring the reverse reaction in which $\mathrm{NH_3}$ decomposes to $\mathrm{N_2}$ and $\mathrm{H_2}$. This is because the overall number of gas molecules would increase and so would the pressure. A decrease in pressure on an equilibrium system favors the reaction which produces more total moles of gas. This is summarized in the table below.

Table 9.6.2

Stress	Response
pressure increase	reaction produces fewer gas molecules
pressure decrease	reaction produces more gas molecules

Like changes in concentration, the K_{eq} value for a given reaction is unchanged by a change in pressure. The amounts of each substance will change but the ratio will not. It is important to remember when analyzing the effect of a pressure change on equilibrium that only gases are affected. If a certain reaction involves liquids or solids, they should be ignored. For example, calcium carbonate decomposes according to the equilibrium reaction:

$$\mathrm{CaCO_3\,(s) \rightleftharpoons CaO\,(s) + O_2\,(g)}$$

Oxygen is the only gas in the system. An increase in the pressure of the system slows the rate of decomposition of $\mathrm{CaCO_3}$ because the reverse reaction is favored. When a system contains equal moles of gas on both sides of the equation, pressure has no effect on the equilibrium position, as in the formation of HCl from $\mathrm{H_2}$ and $\mathrm{Cl_2}$.

$$\mathrm{H_2\,(g) + Cl_2\,(g) \rightleftharpoons 2HCl\,(g)}$$

Example 10.7.3

What is the effect on this equilibrium if pressure is increased?

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

Solution

According to Le Chatelier's principle, if pressure is increased, then the equilibrium shifts to the side with the fewer number of moles of gas. This particular reaction shows a total of 4 mol of gas as reactants and 2 mol of gas as products, so the reaction shifts to the right (toward the products side).

Exercise 10.7.3

What is the effect on this equilibrium if pressure is decreased?

$$3O_2(g) \rightleftharpoons 2O_3(g)$$

Answer available at the end of this section.

Application of Le Chatelier's Principle

Oxygen transport by the blood

In aerobic respiration, oxygen is transported to the cells where it is combined with glucose and metabolized to carbon dioxide, which then moves back to the lungs from which it is expelled.

hemoglobin + O_2 $\rightleftharpoons$ oxyhemoglobin

The partial pressure of O_2 in the air is 0.2 atm, sufficient to allow these molecules to be taken up by hemoglobin (the red pigment of blood) in which it becomes loosely bound in a complex known as oxyhemoglobin. At the ends of the capillaries which deliver

the blood to the tissues, the O_2 concentration is reduced by about 50% owing to its consumption by the cells. This shifts the equilibrium to the left, releasing the oxygen so it can diffuse into the cells.

Key Takeaways

- In a reaction at equilibrium, the introduction of more products will shift the mass balance towards more reactants, and the introduction of more reactants will lead to the formation of more products, but the *ratio* of *Products/Reactants* (*equilibrium constant*), *K* is unchanged.
- If temperature is changed, the numeric value *K* will change. If a reaction is *exothermic* (releases heat), an increase in the **temperature** will force the equilibrium to the left, causing the system to absorb heat and thus partially offsetting the rise in temperature. The opposite effect occurs for *endothermic* reactions, which are shifted to the right by rising temperature.
- The effect of **pressure** on an equilibrium is significant only for reactions which involve different numbers of moles of gases on the two sides of the equation. An increase in the total pressure will shift to the side with fewer moles of gas. A decrease in pressure will shift to the side with more moles of gas.

Contributors and Attributions

-

- Allison Soult, Ph.D. (Department of Chemistry, University of Kentucky)

Answers

Exercise 10.7.1

1. shift to the left (toward reactants)
2. shift to the left (toward reactants)

Exercise 10.7.2

Equilibrium shifts to the left (toward reactants).

Exercise 10.7.3

Reaction shifts to the left (toward reactants).

This page titled 10.7: Le Chatelier's Principle is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis.

- 9.7 Le Chatelier's Principle is licensed CC BY-SA 4.0.

10.8: Osmosis and Diffusion

Learning Outcomes

- Define osmosis and diffusion.
- Distinguish among hypotonic, hypertonic, and isotonic solutions.
- Describe a semipermeable membrane.
- Predict behavior of blood cells in different solution types.
- Describe flow of solvent molecules across a membrane.
- Identify the polar and nonpolar regions of a cell membrane.
- Explain the components present in a phospholipid.

Fish cells, like all cells, have semipermeable membranes. Eventually, the concentration of "stuff" on either side of them will even out. A fish that lives in salt water will have somewhat salty water inside itself. Put it in freshwater, and the freshwater will, through osmosis, enter the fish, causing its cells to swell, and the fish will die. What will happen to a freshwater fish in the ocean?

Osmosis

Imagine you have a cup that has $100\ \mathrm{mL}$ water, and you add $15\ \mathrm{g}$ of table sugar to the water. The sugar dissolves and the mixture that is now in the cup is made up of a **solute** (the sugar) that is dissolved in the **solvent** (the water). The mixture of a solute in a solvent is called a **solution**.

Imagine now that you have a second cup with $100\ \mathrm{mL}$ of water, and you add $45\ \mathrm{g}$ of table sugar to the water. Just like the first cup, the sugar is the solute, and the water is the solvent. But now you have two mixtures of different solute concentrations. In comparing two solutions of unequal solute concentration, the solution with the higher solute concentration is **hypertonic**, and the solution with the lower solute concentration is **hypotonic**. Solutions of equal solute concentration are **isotonic**. The first sugar solution is hypotonic to the second solution. The second sugar solution is hypertonic to the first.

You now add the two solutions to a beaker that has been divided by a semipermeable membrane, with pores that are too small for the sugar molecules to pass through, but are big enough for the water molecules to pass through. The hypertonic solution is one side of the membrane and the hypotonic solution on the other. The hypertonic solution has a lower water concentration than the hypotonic solution, so a concentration gradient of water now exists across the membrane. Water molecules will move from the side of higher water concentration to the side of lower concentration until both solutions are isotonic. At this point, **equilibrium** is reached.

Red blood cells behave the same way (see figure below). When red blood cells are in a hypertonic (higher concentration) solution, water flows out of the cell faster than it comes in. This results in *crenation* (shriveling) of the blood cell. On the other extreme, a red blood cell that is hypotonic (lower concentration outside the cell) will result in more water flowing into the cell than out. This results in swelling of the cell and potential *hemolysis* (bursting) of the cell. In an isotonic solution, the flow of water in and out of the cell is happening at the same rate.




Figure 10.8.1: Red blood cells in hypertonic, isotonic, and hypotonic solutions.

Osmosis is the diffusion of water molecules across a semipermeable membrane from an area of lower concentration solution (i.e., higher concentration of water) to an area of higher concentration solution (i.e., lower concentration of water). Water moves into and out of cells by osmosis.

- If a cell is in a hypertonic solution, the solution has a lower water concentration than the cell cytosol, and water moves out of the cell until both solutions are isotonic.
- Cells placed in a hypotonic solution will take in water across their membranes until both the external solution and the cytosol are isotonic.

A red blood cell will swell and undergo hemolysis (burst) when placed in a hypotonic solution. When placed in a hypertonic solution, a red blood cell will lose water and undergo *crenation* (shrivel). Animal cells tend to do best in an isotonic environment, where the flow of water in and out of the cell is occurring at equal rates.

Diffusion

Passive transport is a way that small molecules or ions move across the cell membrane without input of energy by the cell. The three main kinds of passive transport are diffusion (or simple diffusion), osmosis, and facilitated diffusion. Simple diffusion and osmosis do not involve transport proteins. Facilitated diffusion requires the assistance of proteins.

Diffusion is the movement of molecules from an area of high concentration of the molecules to an area with a lower concentration. For cell transport, diffusion is the movement of small molecules across the cell membrane. The difference in the concentrations of the molecules in the two areas is called the **concentration gradient**. The kinetic energy of the molecules results in random motion, causing diffusion. In simple diffusion, this process proceeds without the aid of a transport protein. It is the random motion of the molecules that causes them to move from an area of high concentration to an area with a lower concentration.

Diffusion will continue until the concentration gradient has been eliminated. Since diffusion moves materials from an area of higher concentration to the lower, it is described as moving solutes "down the concentration gradient". The end result is an equal concentration, or **equilibrium**, of molecules on both sides of the membrane. At equilibrium, movement of molecules does not stop. At equilibrium, there is equal movement of materials in both directions.

Not everything can make it into your cells. Your cells have a plasma membrane that helps to guard your cells from unwanted intruders.

The Plasma Membrane and Cytosol

If the outside environment of a cell is water-based, and the inside of the cell is also mostly water, something has to make sure

the cell stays intact in this environment. What would happen if a cell dissolved in water, like sugar does? Obviously, the cell could not survive in such an environment. So something must protect the cell and allow it to survive in its water-based environment. All cells have a barrier around them that separates them from the environment and from other cells. This barrier is called the **plasma membrane**, or cell membrane.

The Plasma Membrane

The plasma membrane (see figure below) is made of a double layer of special lipids, known as **phospholipids**. The phospholipid is a lipid molecule with a hydrophilic ("water-loving") head and two hydrophobic ("water-hating") tails. Because of the hydrophilic and hydrophobic nature of the phospholipid, the molecule must be arranged in a specific pattern as only certain parts of the molecule can physically be in contact with water. Remember that there is water outside the cell, and the **cytoplasm** inside the cell is mostly water as well. So the phospholipids are arranged in a double layer (a bilayer) to keep the cell separate from its environment. Lipids do not mix with water (recall that oil is a lipid), so the phospholipid bilayer of the cell membrane acts as a barrier, keeping water out of the cell, and keeping the cytoplasm inside the cell. The cell membrane allows the cell to stay structurally intact in its water-based environment.

The function of the plasma membrane is to control what goes in and out of the cell. Some molecules can go through the cell membrane to enter and leave the cell, but some cannot. The cell is therefore not completely permeable. "Permeable" means that anything can cross a barrier. An open door is completely permeable to anything that wants to enter or exit through the door. The plasma membrane is **semipermeable**, meaning that some things can enter the cell, and some things cannot.

Molecules that cannot easily pass through the bilayer include ions and small hydrophilic molecules, such as glucose, and macromolecules, including proteins and RNA. Examples of molecules that can easily diffuse across the plasma membrane include carbon dioxide and oxygen gas. These molecules diffuse freely in and out of the cell, along their concentration gradient. Though water is a polar molecule, it can also diffuse through the plasma membrane.




Figure 10.8.2: Plasma membranes are primarily made up of phospholipids (orange). The hydrophilic ("water-loving") head and two hydrophobic ("water-hating") tails are shown. The phospholipids form a bilayer (two layers). The middle of the bilayer is an area without water. There can be water on either side of the bilayer. There are many proteins throughout the membrane.

Cytosol

The inside of all cells also contain a jelly-like substance called **cytosol**. Cytosol is composed of water and other molecules, including **enzymes**, which are proteins that speed up the cell's chemical reactions. Everything in the cell sits in the cytosol, like fruit in a Jell-o mold. The term cytoplasm refers to the cytosol and all of the organelles, the specialized compartments of the cell. The cytoplasm does not include the nucleus. As a prokaryotic cell does not have a nucleus, the DNA is in the cytoplasm.

Supplemental Resources

- The Plasma Membrane: www.youtube.com/watch?v=moPJkCbKjBs

Key Takeaways

- Water moves into and out of cells by osmosis.
- Water (solvent) moves from an area of lower concentration solution (i.e., higher concentration of water) to an area of higher concentration solution (i.e., lower concentration of water).

This page titled 10.8: Osmosis and Diffusion is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis.

- 8.4: Osmosis and Diffusion by CK12 is licensed CK-12. Original source: https://www.ck12.org/c/chemistry/.

10.S: Solutions (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

A **solution** is a homogeneous mixture. The major component is the **solvent**, while the minor component is the **solute**. Solutions can have any phase; for example, an **alloy** is a solid solution. Solutes are **soluble** or **insoluble**, meaning they dissolve or do not dissolve in a particular solvent. The terms **miscible** and **immiscible**, instead of soluble and insoluble, are used for liquid solutes and solvents. The statement *like dissolves like* is a useful guide to predicting whether a solute will dissolve in a given solvent.

The amount of solute in a solution is represented by the **concentration** of the solution. The maximum amount of solute that will dissolve in a given amount of solvent is called the **solubility** of the solute. Such solutions are **saturated**. Solutions that have less than the maximum amount are **unsaturated**. Most solutions are unsaturated, and there are various ways of stating their concentrations. **Mass/mass percent**, **volume/volume percent**, and **mass/volume percent** indicate the percentage of the overall solution that is solute. **Parts per million (ppm)** and **parts per billion (ppb)** are used to describe very small concentrations of a solute. **Molarity**, defined as the number of moles of solute per liter of solution, is a common concentration unit in the chemistry laboratory. **Equivalents** express concentrations in terms of moles of charge on ions. When a solution is diluted, we use the fact that the amount of solute remains constant to be able to determine the volume or concentration of the final diluted solution.

Dissolving occurs by **solvation**, the process in which particles of a solvent surround the individual particles of a solute, separating them to make a solution. For water solutions, the word **hydration** is used. If the solute is molecular, it dissolves into individual molecules. If the solute is ionic, the individual ions separate from each other, forming a solution that conducts electricity. Such solutions are called **electrolytes**. If the dissociation of ions is complete, the solution is a **strong electrolyte**. If the dissociation is only partial, the solution is a **weak electrolyte**. Solutions of molecules do not conduct electricity and are called **nonelectrolytes**.

Solutions have properties that differ from those of the pure solvent. Some of these are **colligative** properties, which are due to the number of solute particles dissolved, not the chemical identity of the solute. Colligative properties include **vapor pressure depression**, **boiling point elevation**, **freezing point depression**, and **osmotic pressure**. Osmotic pressure is particularly important in biological systems. It is caused by **osmosis**, the passage of solvents through certain membranes like cell walls. The **osmolarity** of a solution is the product of a solution's molarity and the number of particles a solute separates into when it dissolves. Osmosis can be reversed by the application of pressure; this reverse osmosis is used to make fresh water from saltwater in some parts of the world. Because of osmosis, red blood cells placed in hypotonic or hypertonic solutions lose function through either hemolysis or crenation. If they are placed in isotonic solutions, however, the cells are unaffected because osmotic pressure is equal on either side of the cell membrane.

This page titled 10.S: Solutions (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 9.S: Solutions (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11: ACIDS AND BASES

Many of us are familiar with the group of chemicals called acids. But do you know what it takes for a compound to be an acid? Actually, there are several different definitions of acid that chemistry uses, and each definition is appropriate under different circumstances. Less familiar—but just as important to chemistry and ultimately to us—is the group of chemicals known as bases. Both acids and bases are important enough that we devote an entire chapter to them—their properties and their reactions.

11.1: Prelude to Acids and Bases

11.2: Arrhenius Definition of Acids and Bases

11.3: Brønsted-Lowry Definition of Acids and Bases

11.4: Water - Both an Acid and a Base

11.5: The Strengths of Acids and Bases

11.6: Buffers

11.S: Acids and Bases (Summary)

Template:HideTOC

This page titled 11: Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10: Acids and Bases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.1: Prelude to Acids and Bases

One of the most concentrated acids in the body is stomach acid, which can be approximated as a 0.05 M hydrochloric acid solution. Special cells in the stomach wall secrete this acid, along with special enzymes, as part of the digestion process. In a laboratory, a 0.05 M solution of hydrochloric acid would dissolve some metals. How does the stomach survive the presence of such a reactive acid?

Actually, the stomach has several mechanisms for withstanding this chemical onslaught. First, the lining of the stomach is coated with a thin layer of mucus that contains some bicarbonate ions (HCO_3^-). These react with the hydrochloric acid to produce water, carbon dioxide, and harmless chloride ions. If any acid penetrates through the mucus, it can attack the surface layer of stomach cells, called the *gastric epithelium*. Cells in the gastric epithelium are being constantly shed, so damaged cells are quickly removed and replaced with healthy cells.




Figure 11.1.1: An anatomical diagram of the stomach and esophagus, highlighting the lower esophageal sphincter in its closed position to prevent the backflow of stomach contents. (CC BY-SA 4.0; BruceBlaus).

However, if the gastric epithelium is destroyed faster than it can be replaced, the acid may reach the wall of the stomach, resulting in ulcers. If an ulcer grows large enough, it can expose blood vessels in the stomach wall, causing bleeding. In extreme situations, the loss of blood through a severe ulcer can threaten a person's health.

Ulcers can also result from the presence of a certain bacterium—*Helicobacter pylori*—in the stomach. The mechanism for this ulcer formation is not the same as that for ulcers caused by stomach acid and is not completely understood. However, there are two main treatments for ulcers: (1) *antacids* to react chemically with excess hydrochloric acid in the stomach and (2) *antibiotics* to destroy the *H. pylori* bacteria in the stomach.

This page titled 11.1: Prelude to Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.0: Prelude to Acids and Bases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.2: Arrhenius Definition of Acids and Bases

Learning Objectives

- To recognize a compound as an Arrhenius acid or an Arrhenius base.
- To describe characteristics of acids and bases.
- To write equations of neutralization reactions.

One way to define a class of compounds is by describing the various characteristics its members have in common. In the case of the compounds known as acids, the common characteristics include a sour taste, the ability to change the color of the vegetable dye *litmus* to red, and the ability to dissolve certain metals and simultaneously produce hydrogen gas. For the compounds called bases, the common characteristics are a slippery texture, a bitter taste, and the ability to change the color of litmus to blue. Acids and bases also react with each other to form compounds generally known as salts.




Figure 11.2.1: A Common characteristics of acids and bases, illustrated by a lemon (acidic) and soap (basic), highlighting their effects on litmus paper, taste, and chemical reactivity.

Although we include their tastes among the common characteristics of acids and bases, we never advocate tasting an unknown chemical!

Chemists prefer, however, to have definitions for acids and bases in chemical terms. The Swedish chemist Svante Arrhenius developed the first chemical definitions of acids and bases in the late 1800s. Arrhenius defined an acid as a compound that increases the concentration of hydrogen ion (H^+) in aqueous solution. Many acids are simple compounds that release a hydrogen cation into solution when they dissolve. Similarly, Arrhenius defined a base as a compound that increases the concentration of hydroxide ion (OH^-) in aqueous solution. Many bases are ionic compounds that have the hydroxide ion as their anion, which is released when the base dissolves in water.

Table 11.2.1: Formulas and Names for Some Acids and Bases

Acids		Bases	
Formula	Name	Formula	Name
HCl(aq)	hydrochloric acid	NaOH(aq)	sodium hydroxide
HBr(aq)	hydrobromic acid	KOH(aq)	potassium hydroxide
HI(aq)	hydriodic acid	$Mg(OH)_2$(aq)	magnesium hydroxide
H_2S(aq)	hydrosulfuric acid	$Ca(OH)_2$(aq)	calcium hydroxide
$HC_2H_3O_2$(aq)	acetic acid	NH_3(aq)	ammonia
HNO_3(aq)	nitric acid	$NaHCO_3$ (aq)	sodium bicarbonate

Acids		Bases	
$HNO_2(aq)$	nitrous acid	$CaCO_3$ (aq)	calcium carbonate
$H_2SO_4(aq)$	sulfuric acid		
$H_2SO_3(aq)$	sulfurous acid		
$HClO_3(aq)$	chloric acid		
$HClO_4(aq)$	perchloric acid		
$HClO_2(aq)$	chlorous acid		
$H_3PO_4(aq)$	phosphoric acid		
$H_3PO_3(aq)$	phosphorous acid		
$H_2CO_3(aq)$	carbonic acid		

Many bases and their aqueous solutions are named using the normal rules of ionic compounds that were presented previously; that is, they are named as hydroxide compounds. For example, the base sodium hydroxide (NaOH) is both an ionic compound and an aqueous solution. However, aqueous solutions of acids have their own naming rules. The names of *binary acids* (compounds with hydrogen and one other element in their formula) are based on the root of the name of the other element preceded by the prefix *hydro-* and followed by the suffix *-ic acid*. Thus, an aqueous solution of HCl [designated "HCl(aq)"] is called hydrochloric acid, $H_2S(aq)$ is called hydrosulfuric acid, and so forth. Acids composed of more than two elements (typically hydrogen and oxygen and some other element) have names based on the name of the other element, followed by the suffix *-ic acid* or *-ous acid*, depending on the number of oxygen atoms in the acid's formula. Other prefixes, like *per-* and *hypo-*, also appear in the names for some acids. Unfortunately, there is no strict rule for the number of oxygen atoms that are associated with the *-ic acid* suffix; the names of these acids are best memorized. Table 11.2.1 lists some acids and bases and their names. Note that acids have hydrogen written first, as if it were the cation, while most bases have the negative hydroxide ion, if it appears in the formula, written last.

> The name oxygen comes from the Latin meaning "acid producer" because its discoverer, Antoine Lavoisier, thought it was the essential element in acids. Lavoisier was wrong, but it is too late to change the name now.

Example 11.2.1

Name each substance.

a. $HF(aq)$
b. $Sr(OH)_2(aq)$

Solution

a. This acid has only two elements in its formula, so its name includes the *hydro-* prefix. The stem of the other element's name, fluorine, is *fluor*, and we must also include the *-ic acid* ending. Its name is hydrofluoric acid.
b. This base is named as an ionic compound between the strontium ion and the hydroxide ion: strontium hydroxide.

Exercise 11.2.1

Name each substance.

a. $H_2Se(aq)$
b. $Ba(OH)_2(aq)$

Answer available at the end of this section.

Notice that one base listed in Table 11.2.1—ammonia—does not have hydroxide as part of its formula. How does this compound

increase the amount of hydroxide ion in aqueous solution? Instead of dissociating into hydroxide ions, ammonia molecules react with water molecules by taking a hydrogen ion from the water molecule to produce an ammonium ion and a hydroxide ion:

$$NH_{3(aq)} + H_2O_{(\ell)} \rightarrow NH^+_{4(aq)} + OH^-_{(aq)} \tag{11.2.1}$$

Because this reaction of ammonia with water causes an increase in the concentration of hydroxide ions in solution, ammonia satisfies the Arrhenius definition of a base. Many other nitrogen-containing compounds are bases because they too react with water to produce hydroxide ions in aqueous solution.

Neutralization

As we noted previously, acids and bases react chemically with each other to form *salts*. A salt is a general chemical term for any ionic compound formed from an acid and a base. In reactions where the acid is a hydrogen ion containing compound and the base is a hydroxide ion containing compound, water is also a product. The general reaction is as follows:

$$\text{acid + base} \rightarrow \text{water + salt}$$

The reaction of acid and base to make water and a salt is called **neutralization**. Like any chemical equation, a neutralization chemical equation must be properly balanced. For example, the neutralization reaction between sodium hydroxide and hydrochloric acid is as follows:

$$NaOH(aq) + HCl_{(aq)} \rightarrow NaCl_{(aq)} + H_2O_{(\ell)} \tag{11.2.2}$$

with coefficients all understood to be one. The neutralization reaction between sodium hydroxide and sulfuric acid is as follows:

$$2NaOH_{(aq)} + H_2SO_{4(aq)} \rightarrow Na_2SO_{4(aq)} + 2H_2O_{(\ell)} \tag{11.2.3}$$

Once a neutralization reaction is properly balanced, we can use it to perform stoichiometry calculations, such as the ones we practiced earlier.

There are a number of examples of acid-base chemistry in everyday life. One example is the use of baking soda, or sodium bicarbonate in baking. $NaHCO_3$ is a base. When it reacts with an acid such as lemon juice, buttermilk, or sour cream in a batter, bubbles of carbon dioxide gas are formed from decomposition of the resulting carbonic acid, and the batter "rises." Baking powder is a combination of sodium bicarbonate, and one or more acid salts that react when the two chemicals come in contact with water in the batter.

$$HCO^-_3(aq) + H^+(aq) \rightarrow H_2CO_3(aq) \tag{11.2.4}$$

$$H_2CO_3(aq) \rightarrow CO_2(g) + H_2O(l)$$

Example 11.2.2

Nitric acid [$HNO_3(aq)$] can be neutralized by calcium hydroxide [$Ca(OH)_2(aq)$].

a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 16.8 g of HNO_3 is present initially. How many grams of $Ca(OH)_2$ are needed to neutralize that much HNO_3?
c. In a second reaction, 805 mL of 0.672 M $Ca(OH)_2$ is present initially. What volume of 0.432 M HNO_3 solution is necessary to neutralize the $Ca(OH)_2$ solution?

Solution

a. Because there are two OH^- ions in the formula for $Ca(OH)_2$, we need two moles of HNO_3 to provide H^+ ions. The balanced chemical equation is as follows: $Ca(OH)_2(aq) + 2HNO_3(aq) \rightarrow Ca(NO_3)_2(aq) + 2H_2O(\ell)$

   The salt formed is calcium nitrate.

b. This calculation is much like the calculations we did in Chapter 6. First we convert the mass of HNO_3 to moles using its molar mass of 1.01 + 14.01 + 3(16.00) = 63.02 g/mol; then we use the balanced chemical equation to determine the related number of moles of $Ca(OH)_2$ needed to neutralize it; and then we convert that number of moles of $Ca(OH)_2$ to the mass of $Ca(OH)_2$ using its molar mass of 40.08 + 2(1.01) + 2(16.00) = 74.10 g/mol.




Figure 11.2.2: A stoichiometry map showing the calculation path to determine the required mass of calcium hydroxide from a given mass of nitric acid, utilizing molar masses and the balanced equation's mole ratio.

$$16.8\ \text{g}\ HNO_3 \times \frac{1\ \text{mol}\ HNO_3}{63.02\ \text{g}\ HNO_3} \times \frac{1\ \text{mol}\ Ca(OH)_2}{2\ \text{mol}\ HNO_3} \times \frac{74.10\ \text{g}\ Ca(OH)_2}{1\ \text{mol}\ Ca(OH)_2} = 9.88\ \text{g}\ Ca(OH)_2\ \text{needed}$$

c. Having concentration information allows us to employ the skills we developed in Chapter 9. We have two alternative solutions: the multi-step process and the combined one-line process (found at the bottom).

First, we use the concentration and volume data to determine the number of moles of $Ca(OH)_2$ present. Recognizing that 805 mL = 0.805 L,

$$0.672\ \text{M}\ Ca(OH)_2 = \frac{\text{mol}\ Ca(OH)_2}{0.805\ \text{L soln}}$$

0.672 M $CaOH)_2$ × (0.805 L soln) = mol $Ca(OH)_2$ = 0.541 mol $Ca(OH)_2$

We combine this information with the proper ratio from the balanced chemical equation to determine the number of moles of HNO_3 needed:

$$0.541\ \text{mol}\ Ca(OH)_2 \times \frac{2\ \text{mol}\ HNO_3}{1\ \text{mol}\ Ca(OH)_2} = 1.08\ \text{mol}\ HNO_3$$

Now, using the definition of molarity one more time, we determine the volume of acid solution needed:

$$0.432\ \text{M}\ HNO_3 = \frac{1.08\ \text{mol}\ HNO_3}{\text{volume of}\ HNO_3}$$

$$\text{volume of}\ HNO_3 = \frac{1.08\ \text{mol}\ HNO_3}{0.432\ \text{M}\ HNO_3} = 2.50\ \text{L} = 2.50 \times 10^3\ \text{mL}\ HNO_3$$




Figure 11.2.3: A stoichiometry map showing the calculation path to determine the required liters of nitric acid solucion from a given liters of a calcium hydroxide solution, utilizing molarities and the balanced equation's mole ratio.

$$0.805\ \text{L Ca(OH)}_2\ \text{soln} \times \frac{0.672\ \text{mol Ca(OH)}_2}{1\ \text{L Ca(OH)}_2\ \text{soln}} \times \frac{2\ \text{mol HNO}_3}{1\ \text{mol Ca(OH)}_2} \times \frac{1\ \text{L HNO}_3\ \text{soln}}{0.432\ \text{mol HNO}_3} = 2.50\ \text{L HNO}_3\ \text{soln needed}$$

Exercise 11.2.2

Hydrocyanic acid [HCN(aq)] can be neutralized by potassium hydroxide [KOH(aq)].

a. Write a balanced chemical equation for the reaction between these two compounds and identify the salt it produces.
b. For one reaction, 37.5 g of HCN is present initially. How many grams of KOH are needed to neutralize that much HCN?
c. In a second reaction, 43.0 mL of 0.0663 M KOH is present initially. What volume of 0.107 M HCN solution is necessary to neutralize the KOH solution?

Answer available at the end of this section.

Hydrocyanic acid (HCN) is one exception to the acid-naming rules that specify using the prefix *hydro-* for binary acids (acids composed of hydrogen and only one other element).

Stomach Antacids

Our stomachs contain a solution of roughly 0.03 *M* HCl, which helps us digest the food we eat. The burning sensation associated with heartburn is a result of the acid of the stomach leaking through the muscular valve at the top of the stomach into the lower reaches of the esophagus. The lining of the esophagus is not protected from the corrosive effects of stomach acid the way the lining of the stomach is, and the results can be very painful. When we have heartburn, it feels better if we reduce the excess acid in the esophagus by taking an antacid. As you may have guessed, antacids are bases. One of the most common antacids is calcium carbonate, $CaCO_3$. The reaction,

$$CaCO_3(s) + 2HCl(aq) \rightleftharpoons CaCl_2(aq) + H_2O(l) + CO_2(g)$$

not only neutralizes stomach acid, it also produces $CO_2(g)$, which may result in a satisfying belch.

Milk of Magnesia is a suspension of the sparingly soluble base magnesium hydroxide, $Mg(OH)_2$. It works according to the reaction:

$$Mg(OH)_2(s) \rightleftharpoons Mg^{2+}(aq) + 2OH^-(aq)$$

The hydroxide ions generated in this equilibrium then go on to react with the hydronium ions from the stomach acid, so that :

$$H_3O^+ + OH^- \rightleftharpoons 2H_2O(l)$$

This reaction does not produce carbon dioxide, but magnesium-containing antacids can have a laxative effect. Several antacids have aluminum hydroxide, $Al(OH)_3$, as an active ingredient. The aluminum hydroxide tends to cause constipation, and some antacids use aluminum hydroxide in concert with magnesium hydroxide to balance the side effects of the two substances.

Example 11.2.3

Assume that the stomach of someone suffering from acid indigestion contains 75 mL of 0.20 M HCl. How many Tums tablets are required to neutralize 90% of the stomach acid, if each tablet contains 500 mg of $CaCO_3$? (Neutralizing all of the stomach acid is not desirable because that would completely shut down digestion.)

Solution

A. Write the balanced chemical equation for the reaction and then decide whether the reaction will go to completion.
B. Calculate the number of moles of acid present. Multiply the number of moles by the percentage to obtain the quantity of acid that must be neutralized. Using mole ratios, calculate the number of moles of base required to neutralize the acid.

C. Calculate the number of moles of base contained in one tablet by dividing the mass of base by the corresponding molar mass. Calculate the number of tablets required by dividing the moles of base by the moles contained in one tablet.

A. We first write the balanced chemical equation for the reaction:

$$2HCl(aq) + CaCO_3(s) \rightarrow CaCl_2(aq) + H_2CO_3(aq)$$

Each carbonate ion can react with 2 mol of H^+ to produce H_2CO_3, which rapidly decomposes to H_2O and CO_2. Because HCl is a strong acid and CO_3^{2-} is a weak base, the reaction will go to completion.

B. Next we need to determine the number of moles of HCl present:

$$75 \ \cancel{mL}\left(\frac{1 \ \cancel{L}}{1000 \ \cancel{mL}}\right)\left(\frac{0.20 \, mol \, HCl}{\cancel{L}}\right) = 0.015 \ mol \ HCl$$

Because we want to neutralize only 90% of the acid present, we multiply the number of moles of HCl by 0.90:

$$(0.015 \ mol \ HCl)(0.90) = 0.014 \ mol \ HCl$$

We know from the stoichiometry of the reaction that each mole of $CaCO_3$ reacts with 2 mol of HCl, so we need

$$moles \ CaCO_3 = 0.014 \ \cancel{mol \, HCl}\left(\frac{1 \, mol \, CaCO_3}{2 \ \cancel{mol \, HCl}}\right) = 0.0070 \ mol \ CaCO_3$$

C. Each Tums tablet contains

$$\left(\frac{500 \ \cancel{mg \, CaCO_3}}{1 \, Tums \, tablet}\right)\left(\frac{1 \ \cancel{g}}{1000 \ \cancel{mg \, CaCO_3}}\right)\left(\frac{1 \, mol \, CaCO_3}{100.1 \ \cancel{g}}\right) = 0.00500 \ mol \ CaCO_3$$

Thus we need $\dfrac{0.0070 \ \cancel{mol \, CaCO_3}}{0.00500 \ \cancel{mol \, CaCO_3}} = 1.4$ Tums tablets.

Exercise 11.2.3

Assume that as a result of overeating, a person's stomach contains 300 mL of 0.25 M HCl. How many Rolaids tablets must be consumed to neutralize 95% of the acid, if each tablet contains 400 mg of $NaAl(OH)_2CO_3$? The neutralization reaction can be written as follows:

$$NaAl(OH)_2CO_3(s) + 4HCl(aq) \rightarrow AlCl_3(aq) + NaCl(aq) + CO_2(g) + 3H_2O(l)$$

Answer available at the end of this section.

Key Takeaway

- An Arrhenius acid increases the H^+ ion concentration in water, while an Arrhenius base increases the OH^- ion concentration in water.

Answers

Exercise 11.2.1

a. hydroselenic acid

b. barium hydroxide

Exercise 11.2.2

a. $KOH(aq) + HCN(aq) \rightarrow KCN(aq) + H_2O(\ell)$ KCN is the salt.

b. 77.8 g

c. 0.0266 L or 26.6 mL

Exercise 11.2.3

6.4 tablets

This page titled 11.2: Arrhenius Definition of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.1: Arrhenius Definition of Acids and Bases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.3: Brønsted-Lowry Definition of Acids and Bases

Learning Objectives

1. Recognize a compound as a Brønsted-Lowry acid or a Brønsted-Lowry base.
2. Illustrate the proton transfer process that defines a Brønsted-Lowry acid-base reaction.

Ammonia (NH_3) increases the hydroxide ion concentration in aqueous solution by reacting with water rather than releasing hydroxide ions directly. In fact, the Arrhenius definitions of an acid and a base focus on hydrogen ions and hydroxide ions. Are there more fundamental definitions for acids and bases?

In 1923, the Danish scientist Johannes Brønsted and the English scientist Thomas Lowry independently proposed new definitions for acids and bases. Rather than considering both hydrogen and hydroxide ions, they focused on the hydrogen ion only. A Brønsted-Lowry acid is a compound that supplies a hydrogen ion in a reaction. A Brønsted-Lowry base, conversely, is a compound that accepts a hydrogen ion in a reaction. Thus, the Brønsted-Lowry definitions of an acid and a base focus on the movement of hydrogen ions in a reaction, rather than on the production of hydrogen ions and hydroxide ions in an aqueous solution.

Let us use the reaction of ammonia in water to demonstrate the Brønsted-Lowry definitions of an acid and a base. Ammonia and water molecules are reactants, while the ammonium ion and the hydroxide ion are products:

$$NH_3(aq) + H_2O(\ell) \rightarrow NH_4^+(aq) + OH^-(aq) \tag{11.3.1}$$

What has happened in this reaction is that the original water molecule has donated a hydrogen ion to the original ammonia molecule, which in turn has accepted the hydrogen ion. We can illustrate this as follows:




Figure 11.3.1: A Brønsted-Lowry acid-base reaction showing ammonia acting as a base by accepting a proton from water H_2O, the acid, to form ammonium and hydroxide.

Because the water molecule donates a hydrogen ion to the ammonia, it is the Brønsted-Lowry acid, while the ammonia molecule—which accepts the hydrogen ion—is the Brønsted-Lowry base. Thus, ammonia acts as a base in both the Arrhenius sense and the Brønsted-Lowry sense.

Is an Arrhenius acid like hydrochloric acid still an acid in the Brønsted-Lowry sense? Yes, but it requires us to understand what really happens when HCl is dissolved in water. Recall that the hydrogen *atom* is a single proton surrounded by a single electron. To make the hydrogen *ion*, we remove the electron, leaving a bare proton. Do we *really* have bare protons floating around in aqueous solution? No, we do not. What really happens is that the H^+ ion attaches itself to H_2O to make H_3O^+, which is called the *hydronium ion*. For most purposes, H^+ and H_3O^+ represent the same species, but writing H_3O^+ instead of H^+ shows that we understand that there are no bare protons floating around in solution. Rather, these protons are actually attached to solvent molecules.

With this in mind, how do we define HCl as an acid in the Brønsted-Lowry sense? Consider what happens when HCl is dissolved in H_2O:

$$HCl(g) + H_2O(\ell) \rightarrow H_3O^+(aq) + Cl^-(aq) \tag{11.3.2}$$

We can depict this process using Lewis electron dot diagrams:




Figure 11.3.2: The reaction of hydrochloric acid with water.

Now we see that a hydrogen ion is transferred from the HCl molecule to the H_2O molecule to make chloride ions and hydronium ions. As the hydrogen ion donor, **HCl acts as a Brønsted-Lowry acid**; as a hydrogen ion acceptor, **H_2O is a Brønsted-Lowry base**. So HCl is an acid not just in the Arrhenius sense but also in the Brønsted-Lowry sense. Moreover, by the Brønsted-Lowry definitions, H_2O is a base in the formation of aqueous HCl. So the Brønsted-Lowry definitions of an acid and a base classify the dissolving of HCl in water as a reaction between an acid and a base—although the Arrhenius definition would not have labeled H_2O a base in this circumstance.

> All Arrhenius acids and bases are Brønsted-Lowry acids and bases as well. However, not all Brønsted-Lowry acids and bases are Arrhenius acids and bases.

Example 11.3.1

Aniline ($C_6H_5NH_2$) is slightly soluble in water. It has a nitrogen atom that can accept a hydrogen ion from a water molecule just like the nitrogen atom in ammonia does. Write the chemical equation for this reaction and identify the Brønsted-Lowry acid and base.

Solution

$C_6H_5NH_2$ and H_2O are the reactants. When $C_6H_5NH_2$ accepts a proton from H_2O, it gains an extra H and a positive charge and leaves an OH^- ion behind. The reaction is as follows:

$$C_6H_5NH_2(aq) + H_2O(\ell) \rightarrow C_6H_5NH_3^+(aq) + OH^-(aq)$$

Because $C_6H_5NH_2$ accepts a proton, it is the Brønsted-Lowry base. The H_2O molecule, because it donates a proton, is the Brønsted-Lowry acid.

Exercise 11.3.1

Caffeine ($C_8H_{10}N_4O_2$) is a stimulant found in coffees and teas. When dissolved in water, it can accept a proton from a water molecule. Write the chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer available at the end of this section.

The Brønsted-Lowry definitions of an acid and a base can be applied to chemical reactions that occur in solvents other than water. The following example illustrates.

Example 11.3.2

Sodium amide ($NaNH_2$) dissolves in methanol (CH_3OH) and separates into sodium ions and amide ions (NH_2^-). The amide ions react with methanol to make ammonia and the methoxide ion (CH_3O^-). Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Solution

The equation for the reaction is between NH_2^- and CH_3OH to make NH_3 and CH_3O^- is as follows:

$$NH_2^-(solv) + CH_3OH(\ell) \rightarrow NH_3(solv) + CH_3O^-(solv)$$

The label *(solv)* indicates that the species are dissolved in some solvent, in contrast to *(aq)*, which specifies an aqueous (H_2O) solution. In this reaction, we see that the NH_2^- ion accepts a proton from a CH_3OH molecule to make an NH_3 molecule. Thus, as the proton acceptor, NH_2^- is the Brønsted-Lowry base. As the proton donor, CH_3OH is the Brønsted-Lowry acid.

Exercise 11.3.2

Pyridinium chloride (C_5H_5NHCl) dissolves in ethanol (C_2H_5OH) and separates into pyridinium ions ($C_5H_5NH^+$) and chloride ions. The pyridinium ion can transfer a hydrogen ion to a solvent molecule. Write a balanced chemical equation for this process and identify the Brønsted-Lowry acid and base.

Answer available at the end of this section.

Application in Everyday Life

Many people like to put lemon juice or vinegar, both of which are acids, on cooked fish (Figure 11.3.3). It turns out that fish have volatile amines (bases) in their systems, which are neutralized by the acids to yield involatile ammonium salts. This reduces the odor of the fish, and also adds a “sour” taste that we seem to enjoy.



Figure 11.3.3: A neutralization reaction takes place between citric acid or acetic acid (proton donors) in lemons or vinegar and putrescine (proton acceptor) in the flesh of fish. (CC BY 4.0; OpenStax)

Pickling is a method used to preserve vegetables using a naturally produced acidic environment. The vegetable, such as a

cucumber, is placed in a sealed jar submerged in a brine solution. The brine solution favors the growth of beneficial bacteria and suppresses the growth of harmful bacteria. The beneficial bacteria feed on starches in the cucumber and produce lactic acid as a waste product in a process called fermentation. The lactic acid eventually increases the acidity of the brine to a level that kills any harmful bacteria, which require a basic environment. Without the harmful bacteria consuming the cucumbers they are able to last much longer than if they were unprotected. A byproduct of the pickling process changes the flavor of the vegetables with the acid making them taste sour.

To Your Health: Brønsted-Lowry Acid-Base Reactions in Pharmaceuticals

There are many interesting applications of Brønsted-Lowry acid-base reactions in the pharmaceutical industry. For example, drugs often need to be water soluble for maximum effectiveness. However, many complex organic compounds are not soluble or are only slightly soluble in water. Fortunately, those drugs that contain proton-accepting nitrogen atoms (and there are a lot of them) can be reacted with dilute hydrochloric acid [HCl(aq)]. The nitrogen atoms—acting as Brønsted-Lowry bases—accept the hydrogen ions from the acid to make an ion, which is usually much more soluble in water. The modified drug molecules can then be isolated as chloride salts:

$$\mathrm{RN(aq) + H^+}(aq) \rightarrow \mathrm{RNH^+}(aq) \xrightarrow{\mathrm{Cl^-}(aq)} \mathrm{RNH_2Cl(s)} \tag{11.3.3}$$

where RN represents some organic compound containing nitrogen. The label *(sl aq)* means “slightly aqueous,” indicating that the compound RN is only slightly soluble. Drugs that are modified in this way are called *hydrochloride salts*. Examples include the powerful painkiller codeine, which is commonly administered as codeine hydrochloride. Acids other than hydrochloric acid are also used. Hydrobromic acid, for example, gives *hydrobromide salts*. Dextromethorphan, an ingredient in many cough medicines, is dispensed as dextromethorphan hydrobromide. The accompanying figure shows another medication (lidocaine) as a hydrochloride salt.




Figure 11.3.4: Lidocaine. (Figure used with permission from Wikipedia.)

Conjugate Acid-Base Pairs

According to the Bronsted-Lowry theory of acids and bases, an acid is a proton donor and a base is a proton acceptor. Once an acid has given up a proton, the part that remains is called the acid's **conjugate base**. This species is a base because it can accept a proton (to re-form the acid). The conjugate base of HF (first example below) is fluoride ion, F^-.

$$\mathrm{Acid} = \mathrm{H^+} + \mathrm{Conjugate\ base\ of\ Acid^-}$$

$$\mathrm{HF} \rightleftharpoons \mathrm{H^+} + \mathrm{F^-}$$

$$\mathrm{H_2O} \rightleftharpoons \mathrm{H^+} + \mathrm{OH^-}$$

$$NH_4^+ \rightleftharpoons H^+ + NH_3$$

Similarly, the part of the base that remains after a base accepts a proton is called the base's **conjugate acid**. This species is an acid because it can give up a proton (and thus re-form the base). The conjugate acid of fluoride ion, F^- (first example below) is HF.

$$H^+ + \text{Base} = \text{Conjugate acid of Base}^+$$

$$H^+ + F^- \rightleftharpoons HF$$

$$H^+ + OH^- \rightleftharpoons H_2O$$

$$H^+ + H_2O \rightleftharpoons H_3O^+$$

$$H^+ + NH_3 \rightleftharpoons NH_4^+$$

To summarize, the conjugate base of HF is fluoride ion, F^-, and the conjugate acid of fluoride ion, F^-, is HF. The HF/F- pair is referred to as a conjugate acid-base pair. The difference in the formulas of a conjugate acid-base pair (example: HF and F^-) is H^+. The table below lists conjugate acid-base pairs for your reference so that you can figure out the strategy of identifying them. For any given acid or base, you should be able to give its conjugate base or conjugate acid. The formula of an acid's conjugate base is generated by **removing a proton** (H^+) from the acid formula. The formula of the base's conjugate acid is formed by **adding a proton** (H^+) to the formula of the base.

Table 11.3.1. *Conjugate acid-base pairs.*

Conjugate Acid	Conjugate Base
H_3O^+	H_2O
H_2O	OH^-
H_2SO_4	HSO_4^-
HSO_4^-	SO_4^{2-}
NH_4^+	NH_3
NH_3	NH_2^-
CH_3COOH	CH_3COO^-
$CH_3NH_3^+$	CH_3NH_2

Example 11.3.3 : Conjugate Pairs

Write the formula of the conjugate base of (a) HCl and (b) HCO_3^-.

Write the formula of the conjugate acid of (c) CH_3NH_2 and (d) OH^-.

Solution:

A conjugate base is formed by removing a proton (H^+). A conjugate acid is formed by accepting a proton (H^+).

a. After HCl donates a proton, a Cl^- ion is produced, and so Cl^- is the conjugate base.
b. After hydrogen carbonate ion, HCO_3^-, donates a proton, its conjugate base, CO_3^{2-} is produced.
c. After accepting a proton (H^+), CH_3NH_2 is converted to $CH_3NH_3^+$, its conjugate acid.
d. After accepting a proton (H^+), OH^- is converted to H_2O, its conjugate acid.

Exercise 11.3.3: Conjugate Pairs

Write the formula of the conjugate base of (a) HNO_2 and (b) H_2CO_3.

Write the formula of the conjugate acid of (c) C_6H5NH_2 and (d) HCO_3^-.

Answer available at the end of this section.

Identifying Conjugate Acid-Base Pairs in Reactions

In the reaction of ammonia with water to give ammonium ions and hydroxide ions, ammonia acts as a base by accepting a proton from a water molecule, which in this case means that water is acting as an acid. In the reverse reaction, an ammonium ion acts as an acid by donating a proton to a hydroxide ion, and the hydroxide ion acts as a base. The **conjugate acid–base pairs** for this reaction are NH_4^+/NH_3 and H_2O/OH^-. This means that the conjugate acid of the base NH_3 is NH_4^+ while the conjugate base of the acid NH_4^+ is NH_3. Similarly, the conjugate base of the acid H_2O is OH^-, and the conjugate acid of the base OH^- is H_2O.




Figure 11.3.5: Diagram illustrating conjugate acid-base pairs, showing the removal and addition of H⃟ ions between pairs.

In the **forward** reaction, the parent acid is H_2O and and the parent base is NH_3 (shown in the illustration below). The acid H_2O loses a proton (H^+) to form its conjugate base OH^-. The base NH_3 gains a proton, to produce its conjugate acid NH_4^+. In the **reverse** reaction, the acid NH_4^+ loses a proton (H^+) to form its conjugate base NH_3. The base OH^- gains a proton, to produce its conjugate acid H_2O.




Figure 11.3.6: A reversible acid-base reaction between water and ammonia illustrating how the parent acid becomes a conjugate base and the parent base becomes a conjugate acid through the transfer of a proton.

When hydrogen fluoride (HF) dissolves in water and ionizes, protons are transferred from hydrogen fluoride (parent acid) molecules to water (parent base) molecules, yielding hydronium ions (conjugate acid of water) and fluoride ions (conjugate base of HF):

$$\mathrm{HF} + H_2O \rightleftharpoons H_3O^+ + \mathrm{F}^-$$




Figure 11.3.7: The reversible reaction of hydrofluoric acid with water, identifying the proton transfer that creates the conjugate acid-base pairs HF/F^- and H_3O^+/H_2O

Example 11.3.4

Identify the conjugate acid-base pairs in this equilibrium.

$$\mathrm{CH_3CO_2H} + \mathrm{H_2O} \rightleftharpoons \mathrm{H_3O^+} + \mathrm{CH_3CO_2^-}$$

Solution

Similarly, in the reaction of acetic acid with water, acetic acid **donates** a proton to water, which acts as the base. In the reverse reaction, H_3O^+ is the acid that donates a proton to the acetate ion, which acts as the base.

Once again, we have two conjugate acid–base pairs:

- the parent acid and its conjugate base ($CH_3CO_2H/CH_3CO_2^-$) and
- the parent base and its conjugate acid (H_3O^+/H_2O).




Figure 11.3.8: The reversible ionization of acetic acid in water, demonstrating the relationship between the parent acid and its conjugate base, and the parent base and its conjugate acid via proton transfer.

Example 11.3.5

Identify the conjugate acid-base pairs in this equilibrium.

$$(CH_3)_3N + H_2O \rightleftharpoons (CH_3)_3NH^+ + OH^-$$

Solution

One pair is H_2O and OH^-, where H_2O has one more H^+ and is the conjugate acid, while OH^- has one less H^+ and is the conjugate base.

The other pair consists of $(CH_3)_3N$ and $(CH_3)_3NH^+$, where $(CH_3)_3NH^+$ is the conjugate acid (it has an additional proton) and $(CH_3)_3N$ is the conjugate base.

Exercise 11.3.4

Identify the conjugate acid-base pairs in this equilibrium.

$$NH_2^- + H_2O \rightleftharpoons NH_3 + OH^-$$

Answer: available at the end of this section.

The use of conjugate acid-base pairs allows us to make a very simple statement about relative strengths of acids and bases. *The stronger an acid, the weaker its conjugate base*, and, conversely, *the stronger a base, the weaker its conjugate acid.*

Key Takeaways

- A Brønsted-Lowry acid is a proton donor, and a Brønsted-Lowry base is a proton acceptor.
- Brønsted-Lowry acid-base reactions are essentially proton transfer reactions.

Answers

Exercise 11.3.1

$C_8H_{10}N_4O_2(aq) + H_2O(\ell) \rightarrow C_8H_{11}N_4O_2^+(aq) + OH^-(aq)$

B-L base B-L acid

Exercise 11.3.2

$C_5H_5NH^+(solv) + C_2H_5OH(\ell) \rightarrow C_5H_5N(solv) + C_2H_5OH_2^+(solv)$

B-L acid B-L base

Exercise 11.3.3**: Conjugate Pairs**

a. NO_2^- is the conjugate base of HNO_2.

b. HCO_3^- is the conjugate base of H_2CO_3

c. $C_6H5NH_3^+$ is the conjugate acid of C_6H5NH_2.

d. H_2CO_3 is the conjugate acid of HCO_3^-

Exercise 11.3.4

$H2O$ (acid) and OH^- (base); $NH2^-$ (base) and $NH3$ (acid)

This page titled 11.3: Brønsted-Lowry Definition of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.2: Brønsted-Lowry Definition of Acids and Bases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.4: Water - Both an Acid and a Base

Learning Objectives

- To write chemical equations for water acting as an acid and as a base.

Water as a Brønsted-Lowry Base

Water (H_2O) is an interesting compound in many respects. Here, we will consider its ability to behave as an acid or a base.

In some circumstances, a water molecule will accept a proton and thus act as a **Brønsted-Lowry base**. We saw an example in the dissolving of HCl in H_2O:

$$\mathrm{HCl} + \mathrm{H_2O}(\ell) \rightarrow \mathrm{H_3O^+(aq)} + \mathrm{Cl^-(aq)} \tag{11.4.1}$$

Water as a Brønsted-Lowry Acid

In other circumstances, a water molecule can donate a proton and thus act as a **Brønsted-Lowry acid**. For example, in the presence of the amide ion (see Example 4 in Section 11.2), a water molecule donates a proton, making ammonia as a product:

$$H_2O(\ell) + NH_2^-(aq) \rightarrow OH^-(aq) + NH_3(aq) \tag{11.4.2}$$

In this case, NH_2^- is a Brønsted-Lowry base (the proton acceptor).

Water Can Act as Either an Acid or a Base

So, depending on the circumstances, H_2O can act as either a Brønsted-Lowry acid or a Brønsted-Lowry base. Water is not the only substance that can react as an acid in some cases or a base in others, but it is certainly the most common example—and the most important one. A substance that can either donate or accept a proton, depending on the circumstances, is called an **amphiprotic** compound. A substance that can act as either an acid or a base is called **amphoteric**.

The Autoionization of Water

A water molecule can act as an acid or a base even in a sample of pure water. About 6 in every 100 million (6 in 10^8) water molecules undergo the following reaction:

$$H_2O(\ell) + H_2O(\ell) \rightarrow H_3O^+(aq) + OH^-(aq) \tag{11.4.3}$$

This process is called the **autoionization of water** (Figure 11.4.1) and occurs in every sample of water, whether it is pure or part of a solution. Autoionization occurs to some extent in any amphiprotic liquid. (For comparison, liquid ammonia undergoes autoionization as well, but only about 1 molecule in a million billion (1 in 10^{15}) reacts with another ammonia molecule.)

Note

It is rare to truly have pure water. Water exposed to air will usually be slightly acidic because dissolved carbon dioxide gas, or carbonic acid, decreases the pH slightly below 7. Alternatively, dissolved minerals, like calcium carbonate (limestone), can make water slightly basic.




Figure 11.4.1 Autoionization. A small fraction of water molecules—approximately 6 in 100 million—ionize spontaneously into hydronium ions and hydroxide ions. This picture necessarily overrepresents the amount of autoionization that really occurs in pure water.

Example 11.4.1

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $H_2O(\ell) + NO_2^-(aq) \rightarrow HNO_2(aq) + OH^-(aq)$
2. $HC_2H_3O_2(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + C_2H_3O_2^-(aq)$

Solution

1. In this reaction, the water molecule donates a proton to the NO_2^- ion, making $OH^-(aq)$. As the proton donor, H_2O acts as a Brønsted-Lowry acid.
2. In this reaction, the water molecule accepts a proton from $HC_2H_3O_2$, becoming $H_3O^+(aq)$. As the proton acceptor, H_2O is a Brønsted-Lowry base.

Exercise 11.4.1

Identify water as either a Brønsted-Lowry acid or a Brønsted-Lowry base.

1. $HCOOH(aq) + H_2O(\ell) \rightarrow H_3O^+(aq) + HCOO^-(aq)$
2. $H_2O(\ell) + PO_4^{3-}(aq) \rightarrow OH^-(aq) + HPO_4^{2-}(aq)$

Answer available at the end of this section.

Key Takeaway

- Water molecules can act as both an acid and a base, depending on the conditions.

Answers

Exercise 11.4.1

1. H_2O acts as the proton acceptor (Brønsted-Lowry base)

2. H_2O acts as the proton donor (Brønsted-Lowry acid)

This page titled 11.4: Water - Both an Acid and a Base is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.3: Water - Both an Acid and a Base by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.5: The Strengths of Acids and Bases

Learning Objectives

- Describe the difference between strong and weak acids and bases.
- Describe how a chemical reaction reaches chemical equilibrium.
- Define the pH scale and use it to describe acids and bases.

Acids and bases do not all demonstrate the same degree of chemical activity in solution. Different acids and bases have different strengths.

Strong and Weak Acids

Let us consider the strengths of acids first. A small number of acids ionize completely in aqueous solution. For example, when HCl dissolves in water, every molecule of HCl separates into a hydronium ion and a chloride ion:

$$\mathrm{HCl\,(g) + H_2O\,(l) \xrightarrow{\sim 100\,\%} H_3O^+\,(aq) + Cl^-\,(aq)} \tag{11.5.1}$$

HCl(aq) is one example of a **strong acid**, which is a compound that is essentially **100% ionized** in aqueous solution. There are very few strong acids. The important ones are listed in Table 11.5.1.

Table 11.5.1: Strong Acids and Bases (All in Aqueous Solution)

Acids	Bases
HCl	LiOH
HBr	NaOH
HI	KOH
HNO_3	$Mg(OH)_2$
H_2SO_4	$Ca(OH)_2$
$HClO_4$	

By analogy, a strong base is a compound that is essentially 100% ionized in aqueous solution. As with acids, there are only a few strong bases, which are also listed in Table 11.5.1.

If an acid is **not listed** in Table 11.5.1, it is likely a **weak acid**, which is a compound that is not 100% ionized in aqueous solution. Similarly, a weak base is a compound that is not 100% ionized in aqueous solution. For example, acetic acid ($HC_2H_3O_2$) is a weak acid. The ionization reaction for acetic acid is as follows:

$$\mathrm{HC_2H_3O_2\,(aq) + H_2O(\ell) \rightarrow H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)} \tag{11.5.2}$$

Depending on the concentration of [$\mathrm{HC_2H_3O_2\,(aq)}$], the ionization reaction may occur only for 1%–5% of the acetic acid molecules.

Looking Closer: Household Acids and Bases

Many household products are acids or bases. For example, the owner of a swimming pool may use muriatic acid to clean the pool. Muriatic acid is another name for hydrochloric acid [$\mathrm{HCl\,(aq)}$]. Vinegar has already been mentioned as a dilute solution of acetic acid [$\mathrm{HC_2H_3O_2\,(aq)}$]. In a medicine chest, one may find a bottle of vitamin C tablets; the chemical name of vitamin C is ascorbic acid ($HC_6H_7O_6$).

One of the more familiar household bases is ammonia (NH_3), which is found in numerous cleaning products. As we mentioned previously, ammonia is a base because it increases the hydroxide ion concentration by reacting with water:

$$\mathrm{NH_3\,(aq) + H_2O(\ell)\ \rightarrow NH_4^+\,(aq) + OH^-\,(aq)} \tag{11.5.3}$$

Many soaps are also slightly basic because they contain compounds that act as Brønsted-Lowry bases, accepting protons from water and forming excess hydroxide ions. This is one reason that soap solutions are slippery.



Figure 11.5.1: (left) Bottles of alkaline drain cleaners containing sodium hydroxide can dissolve greases and hair. (right) Chemical burns caused by sodium hydroxide solution photographed 44 hours after exposure. Figures used with permission from Wikipedia.

Perhaps the most dangerous household chemical is the lye-based drain cleaner. Lye is a common name for sodium hydroxide, although it is also used as a synonym for potassium hydroxide. Lye is an extremely caustic chemical that can react with grease, hair, food particles, and other substances that may build up and form a clog in a pipe. Unfortunately, lye can also attack tissues and other substances in our bodies. Thus, when we use lye-based drain cleaners, we must be very careful not to touch any of the solid drain cleaner or spill the water it was poured into. Safer, nonlye drain cleaners use peroxide compounds to react on the materials in the clog and clear the drain.

Chemical Equilibrium in Weak Acids and Bases

The behavior of **weak acids and bases** illustrates a key concept in chemistry. Does the chemical reaction describing the ionization of a weak acid or base just stop when the acid or base is done ionizing? Actually, no. Rather, the reverse process—the reformation of the molecular form of the acid or base—occurs, ultimately at the same rate as the ionization process. For example, the ionization of the weak acid $\mathrm{HC_2H_3O_2\,(aq)}$ is as follows:

$$\mathrm{HC_2H_3O_2\,(aq) + H_2O(\ell)\ \rightarrow H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)} \tag{11.5.4}$$

The reverse process also begins to occur:

$$\mathrm{H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)\ \rightarrow HC_2H_3O_2\,(aq) + H_2O(\ell)} \tag{11.5.5}$$

Eventually, there is a balance between the two opposing processes, and no additional change occurs. The chemical reaction is better represented at this point with a double arrow:

$$\mathrm{HC_2H_3O_2\,(aq) + H_2O(\ell) \rightleftharpoons H_3O^+\,(aq) + C_2H_3O_2^-\,(aq)} \tag{11.5.6}$$

The $\rightleftharpoons$ implies that **both the forward and reverse reactions** are occurring, and their effects cancel each other out. A process at this point is considered to be at **chemical equilibrium** (or equilibrium). It is important to note that the processes do not stop. They balance out each other so that there is no further net change; that is, chemical equilibrium is a *dynamic equilibrium*.

Example 11.5.1: Partial Ionization

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HNO_2(aq)$
b. $C_5H_5N(aq)$

Solution

a. $HNO_2(aq) + H_2O(\ell) \rightleftharpoons NO_2^-(aq) + H_3O^+(aq)$
b. $C_5H_5N(aq) + H_2O(\ell) \rightleftharpoons C_5H_5NH^+(aq) + OH^-(aq)$

Exercise 11.5.1

Write the equilibrium chemical equation for the partial ionization of each weak acid or base.

a. $HF_{(aq)}$
b. $AgOH_{(aq)}$
c. $CH_3NH_2(aq)$

Answer available at the end of this section.

Acid Ionization Constant, K_a

The ionization for a general weak acid, HA, can be written as follows:

$$\mathrm{HA}\,(aq) \rightleftharpoons \mathrm{H^+}\,(aq) + \mathrm{A^-}\,(aq)$$

Because the acid is weak, an equilibrium expression can be written. An **acid ionization constant** (K_a) is the equilibrium constant for the ionization of an acid.

$$K_a = \frac{[\mathrm{H^+}]\,[\mathrm{A^-}]}{[\mathrm{HA}]}$$

The acid ionization represents the fraction of the original acid that has been ionized in solution. Therefore, the numerical value of K_a is a reflection of the strength of the acid. Weak acids with relatively higher K_a values are stronger than acids with relatively lower K_a values. Because strong acids are essentially 100% ionized, the concentration of the acid in the denominator is nearly zero and the K_a value approaches infinity. For this reason, K_a values are generally reported for weak acids only.

The table below is a listing of acid ionization constants for several acids. Note that polyprotic acids have a distinct ionization constant for each ionization step, with each successive ionization constant being smaller than the previous one.

Table 11.5.2: Acid Ionization Constants at $25^\circ\mathrm{C}$

Name of Acid	Ionization Equation	K_a
Sulfuric acid	$\mathrm{H_2SO_4} \rightleftharpoons \mathrm{H^+} + \mathrm{HSO_4^-}$ $\mathrm{HSO_4} \rightleftharpoons \mathrm{H^+} + \mathrm{SO_4^{2-}}$	very large 1.3×10^{-2}
Hydrofluoric acid	$\mathrm{HF} \rightleftharpoons \mathrm{H^+} + \mathrm{F^-}$	7.1×10^{-4}
Nitrous acid	$\mathrm{HNO_2} \rightleftharpoons \mathrm{H^+} + \mathrm{NO_2^-}$	4.5×10^{-4}
Benzoic acid	$\mathrm{C_6H_5COOH} \rightleftharpoons \mathrm{H^+} + \mathrm{C_6H_5COO^-}$	6.5×10^{-5}

Name of Acid	Ionization Equation	K_a
Acetic acid	$CH_3COOH \rightleftharpoons H^+ + CH_3COO^-$	1.8×10^{-5}
Carbonic acid	$H_2CO_3 \rightleftharpoons H^+ + HCO_3^-$ $HCO_3^- \rightleftharpoons H^+ + CO_3^{2-}$	4.2×10^{-7} 4.8×10^{-11}

Hydrofluoric acid $HF_{(aq)}$ reacts directly with glass (very few chemicals react with glass). Hydrofluoric acid is used in glass etching.

Strong and Weak Bases and Base Ionization Constant, K_b

As with acids, bases can either be strong or weak, depending on their extent of ionization. A **strong base** is a base, which ionizes completely in an aqueous solution. The most common strong bases are soluble metal hydroxide compounds such as potassium hydroxide. Some metal hydroxides are not as strong simply because they are not as soluble. Calcium hydroxide is only slightly soluble in water, but the portion that does dissolve also dissociates into ions.

A **weak base** is a base that ionizes only slightly in an aqueous solution. Recall that a base can be defined as a substance, which accepts a hydrogen ion from another substance. When a weak base such as ammonia is dissolved in water, it accepts an H^+ ion from water, forming the hydroxide ion and the conjugate acid of the base, the ammonium ion.

$$NH_3\,(aq) + H_2O\,(l) \rightleftharpoons NH_4{}^+\,(aq) + OH^-\,(aq)$$

The equilibrium greatly favors the reactants and the extent of ionization of the ammonia molecule is very small.

An equilibrium expression can be written for the reactions of weak bases with water. Because the concentration of water is extremely large and virtually constant, the water is not included in the expression. A **base ionization constant** (K_b) is the equilibrium constant for the ionization of a base. For ammonia the expression is:

$$K_b = \frac{[NH_4{}^+]\,[OH^-]}{[NH_3]}$$

The numerical value of K_b is a reflection of the strength of the base. Weak bases with relatively higher K_b values are stronger than bases with relatively lower K_b values. Table $11.5.3$ is a listing of base ionization constants for several weak bases.

Table $11.5.3$*: Base Ionization Constants at* 25^oC

Name of Base	Ionization Equation	K_b
Methylamine	$CH_3NH_2 + H_2O \rightleftharpoons CH_3NH_3{}^+ + OH^-$	5.6×10^{-4}
Ammonia	$NH_3 + H_2O \rightleftharpoons NH_4{}^+ + OH^-$	1.8×10^{-5}
Pyridine	$C_5H_5N + H_2O \rightleftharpoons C_5H_5NH^+ + OH^-$	1.7×10^{-9}
Acetate ion	$CH_3COO^- + H_2O \rightleftharpoons CH_3COOH + OH^-$	5.6×10^{-10}
Fluoride ion	$F^- + H_2O \rightleftharpoons HF + OH^-$	1.4×10^{-11}
Urea	$H_2NCONH_2 + H_2O \rightleftharpoons H_2NCONH_3{}^+ + OH^-$	1.5×10^{-14}

The Ion-Product of Water

As we have already seen, H_2O can act as an acid or a base. Within any given sample of water, some H_2O molecules are acting as acids, and other H_2O molecules are acting as bases. The chemical equation is as follows:

$$\underbrace{H_2O}_{\text{acid}} + \underbrace{H_2O}_{\text{base}} \rightleftharpoons H_3O^+ + OH^- \tag{11.5.7}$$




Figure 11.5.2: Autoionization of water, giving hydroxide and hydronium ions.

Similar to a weak acid, the autoionization of water is an equilibrium process, and is more properly written as follows:

$$H_2O(\ell) + H_2O(\ell) \rightleftharpoons H_3O^+(aq) + OH^-(aq) \tag{11.5.8}$$

We often use the simplified form of the reaction:

$$H_2O(l) \rightleftharpoons H^+(aq) + OH^-(aq)$$

The equilibrium constant for the autoionization of water is referred to as the ion-product for water and is given the symbol Kw.

$$K_w = [H^+][OH^-]$$

The **ion-product of water (*Kw*)** is the mathematical product of the concentration of hydrogen ions and hydroxide ions. Note that H_2O is not included in the ion-product expression because it is a pure liquid. The value of Kw is very small, in accordance with a reaction that favors the reactants. At 25°C, the experimentally determined value of K_w in pure water is 1.0×10^{-14}.

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14}$$

In a sample of pure water, the concentrations of hydrogen and hydroxide ions are equal to one another. Pure water or any other aqueous solution in which this ratio holds is said to be **neutral.** To find the molarity of each ion, the square root of Kw is taken.

$$[H^+] = [OH^-] = 1.0\times10^{-7}$$

The product of these two concentrations is 1.0×10^{-14}

$$[H^+]\times[OH^-] = (1.0 \times 10^{-7})(1.0 \times 10^{-7}) = 1.0 \times 10^{-14}$$

- For acids, the concentration of H^+ or $[H^+]$) is greater than 1.0×10^{-7} M
- For bases, the concentration of OH^- or $[OH^-]$ is greater than 1.0×10^{-7} M.

Aqueous HCl is an example of acidic solution. Hydrogen chloride (HCl) ionizes to produce H^+ and Cl^- ions upon dissolving in water. This **increases the concentration of H^+ ions** in the solution. According to **Le Chatelier's principle**, the equilibrium represented by

$$H_2O(l) \rightleftharpoons H^+(aq) + OH^-(aq)$$

$$HCl(g) \longrightarrow H^+(aq) + Cl^-(aq)$$

is **forced to the left**, towards the reactant. As a result, the **concentration of the hydroxide ion decreases**.

Now, consider KOH (aq), a **basic solution**. Solid potassium hydroxide (KOH) dissociates in water to yield potassium ions and hydroxide ions.

$$KOH(s) \rightarrow K^+(aq) + OH^-(aq)$$

The **increase in concentration of the OH^- ions** will cause a decrease in the concentration of the H^+ ions.

No matter whether the aqueous solution is an acid, a base, or neutral:and the ion-product of $[H^+][OH^-]$ remains constant.

- For acidic solutions, $[H^+]$ is greater than $[OH^-]$.
- For basic solutions, [OH–] is greater than $[H^+]$.
- For neutral solutions, $[H_3O^+]$ = [OH–] = 1.0×10^{-7}M

This means that if you know $[H^+]$ for a solution, you can calculate what $[OH^-]$) has to be for the product to equal 1.0×10^{-14}, or if you know $[OH^-]$), you can calculate $[H^+]$. This also implies that as one concentration goes up, the other must go down to compensate so that their product always equals the value of K_w.

$$K_w = [H_3O^+][OH^-] = 1.0 \times 10^{-14} \tag{11.5.9}$$

Example 11.5.2

Hydrochloric acid (HCl) is a strong acid, meaning it is 100% ionized in solution. What is the $[H^+]$ and the $[OH^-]$ in a solution of 2.0×10^{-3} M HCl?

Solution

Step 1: List the known values and plan the problem.

Known

- [HCl] = 2.0×10^{-3} M
- Kw = 1.0×10^{-14}

Unknown

- $[H^+]$=?M
- $[OH^-]$=?M

Because HCl is 100% ionized, the concentration of H+ ions in solution will be equal to the original concentration of HCl. Each HCl molecule that was originally present ionizes into one H+ ion and one Cl– ion. The concentration of OH– can then be determined from the [H+] and Kw.

Step 2: Solve.

$$[H^+]=2.0\times10^{-3}\ M$$

$$Kw = [H^+][OH^-] = 1.0\times10^{-14}$$

$$[OH^-] = Kw/[H^+] = 1.0\times10^{-14}/2.0\times10^{-3} = 5.0\times10^{-12}\ M$$

Step 3: Think about your result.

The $[H^+]$ is much higher than the $[OH^-]$ because the solution is acidic. As with other equilibrium constants, the unit for Kw is customarily omitted.

Exercise 11.5.2

Sodium hydroxide (NaOH) is a strong base. What is the $[H^+]$ and the $[OH^-]$ in a 0.001 M NaOH solution at 25 °C?

Answer available at the end of this section.

The pH Scale

One qualitative measure of the strength of an acid or a base solution is the pH scale, which is based on the concentration of the hydronium (or hydrogen) ion in aqueous solution.

$$pH = -\log[H^+]$$

or

$$pH = -\log[H_3O^+]$$

Figure 11.5.3 illustrates this relationship, along with some examples of various solutions. Because hydrogen ion concentrations are generally less than one (for example $1.3 \times 10^{-3}\ M$), the log of the number will be a negative number. To make pH even easier to work with, pH is defined as the ***negative* log of** $[H_3O^+]$, which will give a positive value for pH.




Figure 11.5.3: The relationship between [H$^+$] and pH *values for several common materials.*

A neutral (neither acidic nor basic) solution has a pH of 7. A pH below 7 means that a solution is acidic, with lower values of pH corresponding to increasingly acidic solutions. A pH greater than 7 indicates a basic solution, with higher values of pH corresponding to increasingly basic solutions. Thus, given the pH of several solutions, you can state which ones are acidic, which ones are basic, and which are more acidic or basic than others. These are summarized in Table 11.5.4.

Table 11.5.4: Acidic, Basic and Neutral pH Values

Acidic, Basic and Neutral pH Values

Classification	Relative Ion Concentrations	pH at 25 °C
acidic	[H$^+$] > [OH$^-$]	pH < 7
neutral	[H$^+$] = [OH$^-$]	pH = 7
basic	[H$^+$] < [OH$^-$]	pH > 7

Example 11.5.3

Find the pH, given the $[H^+]$ of the following:

a. 1×10^{-3} M

b. 2.5×10^{-11} M
c. 4.7×10^{-9} M

Solution

pH = - log $[H_3O^+]$

Substitute the known quantity into the equation and solve. Use a scientific calculator for b and c.

a. pH = - log $[1 \times 10^{-3}]$ = 3.0 (1 decimal place since 1 has 1 significant figure)
b. pH = - log $[2.5 \times 10^{-11}]$ = 10.60 (2 decimal places since 2.5 has 2 significant figures)
c. pH = - log $[4.7 \times 10^{-9}]$ = 8.33 (2 decimal places since 4.7 has 2 significant figures)

Note on significant figures:

Because the number(s) before the decimal point in the pH value relate to the power on 10, the number of digits *after* the decimal point (underlined) is what determines the number of significant figures in the final answer.

Exercise 11.5.3

Find the pH, given $[H^+]$ of the following:

a. 5.8×10^{-4} M
b. 1.0×10^{-7} M

Answer available at the end of this section.

Table 11.5.5 lists the pH of several common solutions. The most acidic among the listed solutions is battery acid with the lowest pH value (0.3). The most basic is 1M NaOH solution with the highest pH value of 14.0. Notice that some biological fluids (stomach acid and urine) are nowhere near neutral. You may also notice that many food products are slightly acidic. They are acidic because they contain solutions of weak acids. If the acid components of these foods were strong acids, the food would likely be inedible.

Table 11.5.5: The pH Values of Some Common Solutions

Solution	pH
battery acid	0.3
stomach acid	1–2
lemon or lime juice	2.1
vinegar	2.8–3.0
Coca-Cola	3
wine	2.8–3.8
beer	4–5
coffee	5
milk	6
urine	6
pure H_2O	7
(human) blood	7.3–7.5
sea water	8

Solution	pH
antacid (milk of magnesia)	10.5
NH_3 (1 M)	11.6
bleach	12.6
NaOH (1 M)	14.0

Example 11.5.4

Label each solution as acidic, basic, or neutral based only on the stated pH.

a. milk of magnesia, pH = 10.5
b. pure water, pH = 7
c. wine, pH = 3.0

Solution

a. With a pH greater than 7, milk of magnesia is basic. (Milk of magnesia is largely $Mg(OH)_2$.)
b. Pure water, with a pH of 7, is neutral.
c. With a pH of less than 7, wine is acidic.

Exercise 11.5.4

Identify each substance as acidic, basic, or neutral based only on the stated pH.

a. human blood with pH = 7.4
b. household ammonia with pH = 11.0
c. cherries with pH = 3.6

Answer available at the end of this section.

Acid Rain

Normal rainwater has a pH between 5 and 6 due to the presence of dissolved CO_2 which forms carbonic acid:

$$\mathrm{H_2O\,(l) + CO_2\,(g) \longrightarrow H_2CO_3\,(aq)} \tag{11.5.10}$$

$$\mathrm{H_2CO_3\,(aq) \rightleftharpoons H^+\,(aq) + HCO_3{}^-\,(aq)} \tag{11.5.11}$$

Acid rain is rainwater that has a pH of less than 5, due to a variety of nonmetal oxides, including CO_2, SO_2, SO_3, NO, and NO_2 being dissolved in the water and reacting with it to form not only carbonic acid, but sulfuric acid and nitric acid. The formation and subsequent ionization of sulfuric acid are shown here:

$$\mathrm{H_2O\,(l) + SO_3\,(g) \longrightarrow H_2SO_4\,(aq)} \tag{11.5.12}$$

$$\mathrm{H_2SO_4\,(aq) \longrightarrow H^+\,(aq) + HSO_4{}^-\,(aq)} \tag{11.5.13}$$

Carbon dioxide is naturally present in the atmosphere because we and most other organisms produce it as a waste product of metabolism. Carbon dioxide is also formed when fires release carbon stored in vegetation or when we burn wood or fossil fuels. Sulfur trioxide in the atmosphere is naturally produced by volcanic activity, but it also stems from burning fossil fuels, which have traces of sulfur, and from the process of “roasting” ores of metal sulfides in metal-refining processes. Oxides of nitrogen are formed in internal combustion engines where the high temperatures make it possible for the nitrogen and oxygen in air to chemically combine.

Acid rain is a particular problem in industrial areas where the products of combustion and smelting are released into the air without being stripped of sulfur and nitrogen oxides. In North America and Europe until the 1980s, it was responsible for the destruction of forests and freshwater lakes, when the acidity of the rain actually killed trees, damaged soil, and made lakes uninhabitable for all but the most acid-tolerant species. Acid rain also corrodes statuary and building facades that are made of marble and limestone (Figure 11.5.4). Regulations limiting the amount of sulfur and nitrogen oxides that can be released into the atmosphere by industry and automobiles have reduced the severity of acid damage to both natural and manmade environments in North America and Europe. It is now a growing problem in industrial areas of China and India.



Figure 11.5.4: (a) Acid rain makes trees more susceptible to drought and insect infestation, and depletes nutrients in the soil. (b) It also is corrodes statues that are carved from marble or limestone. (credit a: modification of work by Chris M Morris; credit b: modification of work by "Eden, Janine and Jim"/Flickr)

Key Takeaways

- Acids and bases can be strong or weak depending on the extent of ionization in solution.
- Most chemical reactions reach equilibrium at which point there is no net change.
- The ion-product of $[H^+][OH^-]$ in an aqueous solution remains constant.
- A ***p*H** value is simply the *negative* of the logarithm of the H^+ ion concentration ($-\log[H^+]$).
- The pH scale is used to succinctly communicate the acidity or basicity of a solution.
- A solution is **acidic** if *p*H < 7.
- A solution is **basic** if *p*H > 7.
- A solution is **neutral** if *p*H = 7.

Answers

Exercise 11.5.1

a. $HF(aq) + H_2O(\ell) \rightleftharpoons F^-(aq) + H_3O^+(aq)$

b. $AgOH(aq) \rightleftharpoons Ag^+(aq) + OH^-(aq)$

c. $CH_3NH_2(aq) + H_2O(\ell) \rightleftharpoons CH_3NH_3^+(aq) + OH^-(aq)$

Exercise 11.5.2

$[OH^-] = 0.001M$ or $1 \times 10^{-3}M$; $[H^+]=1\times10^{-11}M$.

Exercise 11.5.3

a. 3.24

b. 7.00

Exercise 11.5.4

a. slightly basic

b. basic

c. acidic

This page titled 11.5: The Strengths of Acids and Bases is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.4: The Strengths of Acids and Bases by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.6: Buffers

Learning Objectives

- To define *buffer* and describe how it reacts with an acid or a base.

Weak acids are relatively common, even in the foods we eat. But we occasionally come across a strong acid or base, such as stomach acid, that has a strongly acidic pH of 1–2. By definition, strong acids and bases can produce a relatively large amount of hydrogen or hydroxide ions and, as a consequence, have a marked chemical activity. In addition, very small amounts of strong acids and bases can change the pH of a solution very quickly. If 1 mL of stomach acid [which we will approximate as 0.05 M HCl(aq)] is added to the bloodstream, and if no correcting mechanism is present, the pH of the blood would go from about 7.4 to about 4.9—a pH that is not conducive to continued living. Fortunately, the body has a mechanism for minimizing such dramatic pH changes.

The mechanism involves a **buffer**, a solution that resists dramatic changes in pH. A *buffer* (or *buffered*) solution is one that resists a change in its pH when H^+ or OH^- ions are added or removed owing to some other reaction taking place in the same solution. Buffers do so by being composed of certain pairs of solutes: either a **weak acid plus its conjugate base** or a **weak base plus its conjugate acid**.

For example, a buffer can be composed of dissolved **acetic acid ($HC_2H_3O_2$, a weak acid) and sodium acetate** ($NaC_2H_3O_2$). Sodium acetate is a salt that dissociates into sodium ions and acetate ions in solution. For as long as acetic acid and acetate ions are present in significant amounts a solution, this can resist dramatic pH changes. Another example of a buffer is a solution containing **ammonia (NH_3, a weak base) and ammonium chloride** (NH_4Cl). Ammonium acetate is also a salt that dissociates into ammonium ions and chloride ions in solution. The presence of ammonium ions with ammonia molecules satisfies the requisite condition for a buffer solution.

How Buffers Work

The essential component of a buffer system is a **conjugate acid-base pair** whose concentration is fairly high in relation to the concentrations of added H^+ or OH^- it is expected to buffer against. Let us use an acetic acid–sodium acetate buffer to demonstrate how buffers work. If a strong base—a source of OH^-(aq) ions—is added to the buffer solution, those hydroxide ions will react with the acetic acid in an acid-base reaction:

$$HC_2H_3O_2\,(aq) + OH^-\,(aq) \longrightarrow H_2O(\ell) + C_2H_3O_2^-\,(aq) \tag{11.6.1}$$

Rather than changing the pH dramatically by making the solution basic, the added hydroxide ions react to make water, and the pH does not change much.

> Many people are aware of the concept of buffers from *buffered aspirin*, which is aspirin that also has magnesium carbonate, calcium carbonate, magnesium oxide, or some other salt. The salt acts like a base, while aspirin is itself a weak acid.

If a strong acid—a source of H^+ ions—is added to the buffer solution, the H^+ ions will react with the anion from the salt. Because $HC_2H_3O_2$ is a weak acid, it is not ionized much. This means that if lots of hydrogen ions and acetate ions (from sodium acetate) are present in the same solution, they will come together to make acetic acid:

$$H^+\,(aq) + C_2H_3O_2^-\,(aq) \longrightarrow HC_2H_3O_2\,(aq) \tag{11.6.2}$$

Rather than changing the pH dramatically and making the solution acidic, the added hydrogen ions react to make molecules of a weak acid. Figure $11.6.1$ illustrates both actions of a buffer.




Figure 11.6.1: The Action of Buffers. Buffers can react with both strong acids (top) and strong bases (bottom) to minimize large changes in pH.

A simple buffer system might be a 0.2 *M* solution of sodium acetate; the conjugate pair here is acetic acid HAc and its conjugate base, the acetate ion Ac^-. The idea is that this conjugate pair "pool" will be available to gobble up any small ($\leq 10^{-3}$ *M*) addition of H+ or OH^- that may result from other processes going on in the solution.




Figure 11.6.2: How HAc/Ac buffer works

Buffers work well only for limited amounts of added strong acid or base. Once either solute is all reacted, the solution is no longer a buffer, and rapid changes in pH may occur. We say that a buffer has a certain capacity. Buffers that have more solute dissolved in them to start with have larger capacities, as might be expected.

Buffers made from weak bases and salts of weak bases act similarly. For example, in a buffer containing NH_3 and NH_4Cl, ammonia molecules can react with any excess hydrogen ions introduced by strong acids:

$$NH_{3(aq)} + H^+_{(aq)} \rightarrow NH^+_{4(aq)} \tag{11.6.3}$$

while the ammonium ion [NH_4^+(aq)] can react with any hydroxide ions introduced by strong bases:

$$NH^+_{4(aq)} + OH^-_{(aq)} \rightarrow NH_{3(aq)} + H_2O_{(\ell)} \tag{11.6.4}$$




Figure 11.6.3: How NH_4^+/NH_3 buffer works

Example 11.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $HCHO_2$ and $NaCHO_2$
b. HCl and NaCl
c. CH_3NH_2 and CH_3NH_3Cl
d. NH_3 and NaOH

Solution

a. Formic acid ($HCHO_2$) is a weak acid, while $NaCHO_2$ is a salt supplying —formate ion (CHO_2^-), the conjugate base of $HCHO_2$. The combination of these two solutes would make a buffer solution.
b. Hydrochloric acid (HCl) is a strong acid, not a weak acid, so the combination of these two solutes would not make a buffer solution.
c. Methylamine (CH_3NH_2) is like ammonia, a weak base. The compound CH_3NH_3Cl is a salt supplying $CH_3NH_3^+$, the conjugate acid of CH_3NH_2. The combination of these two solutes would make a buffer solution.
d. Ammonia (NH_3) is a weak base, but NaOH is a strong base. The combination of these two solutes would not make a buffer solution.

Exercise 11.6.1

Which solute combinations can make a buffer solution? Assume all are aqueous solutions.

a. $NaHCO_3$ and NaCl
b. H_3PO_4 and NaH_2PO_4
c. NH_3 and $(NH_4)_3PO_4$
d. NaOH and NaCl

Answer available at the end of this section.

Food and Drink App: The Acid That Eases Pain

Although medicines are not exactly "food and drink," we do ingest them, so let's take a look at an acid that is probably the most common medicine: acetylsalicylic acid, also known as aspirin. Aspirin is well known as a pain reliever and antipyretic (fever reducer).

The structure of aspirin is shown in the accompanying figure. The acid part is circled; it is the H atom in that part that can be donated as aspirin acts as a Brønsted-Lowry acid. Because it is not given in Table 10.5.1, acetylsalicylic acid is a weak acid. However, it is still an acid, and given that some people consume relatively large amounts of aspirin daily, its acidic nature can cause problems in the stomach lining, despite the stomach's defenses against its own stomach acid.




Figure 11.6.4 The Molecular Structure of Aspirin. The circled atoms are the acid part of the molecule.

Because the acid properties of aspirin may be problematic, many aspirin brands offer a "buffered aspirin" form of the medicine. In these cases, the aspirin also contains a buffering agent-usually MgO-that regulates the acidity of the aspirin to minimize its acidic side effects.

As useful and common as aspirin is, it was formally marketed as a drug starting in 1899. The US Food and Drug Administration (FDA), the governmental agency charged with overseeing and approving drugs in the United States, wasn't formed until 1906. Some have argued that if the FDA had been formed before aspirin was introduced, aspirin may never have gotten approval due to its potential for side effects-gastrointestinal bleeding, ringing in the ears, Reye's syndrome (a liver problem), and some allergic reactions. However, recently aspirin has been touted for its effects in lessening heart attacks and strokes, so it is likely that aspirin is here to stay.

Buffer solutions are essential components of all living organisms.

- Our blood is buffered to maintain a pH of 7.4 that must remain unchanged as metabolically-generated CO_2 (carbonic acid) is added and then removed by our lungs.
- Buffers in the oceans, in natural waters such as lakes and streams, and within soils help maintain their environmental stability against acid rain and increases in atmospheric CO_2.
- Many industrial processes, such as brewing, require buffer control, as do research studies in biochemistry and physiology that involve enzymes, are active only within certain pH ranges.

The pH in living systems (Table \({11{.}6{.}\eatSpaces{1} \endSpaces}) is maintained by buffer systems.

(Table \({11{.}6{.}\eatSpaces{1} \endSpaces}): pH in Living Systems

Compartment	pH
Gastric Acid	1
Lysosomes	4.5
Granules of Chromaffin Cells	5.5
Human Skin	5.5
Urine	6
Neutral H_2O at 37 °C	6.81
Cytosol	7.2

Compartment	pH
Cerebrospinal Fluid	7.3
Blood	7.43-7.45
Mitochondrial Matrix	7.5
Pancreas Secretions	8.1

Human blood has a buffering system to minimize extreme changes in pH. One buffer in blood is based on the presence of HCO_3^- and H_2CO_3 [H_2CO_3 is another way to write $CO_2(aq)$]. With this buffer present, even if some stomach acid were to find its way directly into the bloodstream, the change in the pH of blood would be minimal. Inside many of the body's cells, there is a buffering system based on phosphate ions.

Medicine: The Buffer System in Blood

The normal pH of human blood is about 7.4. The carbonate buffer system in the blood uses the following equilibrium reaction:

$$\mathrm{CO_2}(g) + 2\,\mathrm{H_2O}(l) \rightleftharpoons \mathrm{H_2CO_3}(aq) \rightleftharpoons \mathrm{HCO_3}^-(aq) + \mathrm{H_3O}^+(aq)$$

The concentration of carbonic acid, H_2CO_3 is approximately 0.0012 *M*, and the concentration of the hydrogen carbonate ion, $\mathrm{HCO_3}^-$, is around 0.024 *M*. Using the Henderson-Hasselbalch equation and the pK_a of carbonic acid at body temperature, we can calculate the pH of blood:

$$\mathrm{pH} = \mathrm{p}K_\mathrm{a} + \log\frac{[\mathrm{base}]}{[\mathrm{acid}]} = 6.1 + \log\frac{0.024}{0.0012} = 7.4$$

The fact that the H_2CO_3 concentration is significantly lower than that of the $\mathrm{HCO_3}^-$ ion may seem unusual, but this imbalance is due to the fact that most of the by-products of our metabolism that enter our bloodstream are acidic. Therefore, there must be a larger proportion of base than acid, so that the capacity of the buffer will not be exceeded.

Lactic acid is produced in our muscles when we exercise. As the lactic acid enters the bloodstream, it is neutralized by the $\mathrm{HCO_3}^-$ ion, producing H_2CO_3. An enzyme then accelerates the breakdown of the excess carbonic acid to carbon dioxide and water, which can be eliminated by breathing. In fact, in addition to the regulating effects of the carbonate buffering system on the pH of blood, the body uses breathing to regulate blood pH. If the pH of the blood decreases too far, an increase in breathing removes CO_2 from the blood through the lungs driving the equilibrium reaction such that [H_3O^+] is lowered. If the blood is too alkaline, a lower breath rate increases CO_2 concentration in the blood, driving the equilibrium reaction the other way, increasing [H^+] and restoring an appropriate pH.

Career Focus: Blood Bank Technology Specialist

At this point in this text, you should have the idea that the chemistry of blood is fairly complex. Because of this, people who work with blood must be specially trained to work with it properly.

A blood bank technology specialist is trained to perform routine and special tests on blood samples from blood banks or transfusion centers. This specialist measures the pH of blood, types it (according to the blood's ABO+/– type, Rh factors, and other typing schemes), tests it for the presence or absence of various diseases, and uses the blood to determine if a patient has any of several medical problems, such as anemia. A blood bank technology specialist may also interview and prepare donors to give blood and may actually collect the blood donation.

Blood bank technology specialists are well trained. Typically, they require a college degree with at least a year of special training in blood biology and chemistry. In the United States, training must conform to standards established by the American Association of Blood Banks.

Key Takeaway

- A buffer is a solution that resists sudden changes in pH.

Concept Review Exercise

1. Explain how a buffer prevents large changes in pH.

Answer

1. A buffer has components that react with both strong acids and strong bases to resist sudden changes in pH.

Exercises

1. Describe a buffer. What two related chemical components are required to make a buffer?
2. Can a buffer be made by combining a strong acid with a strong base? Why or why not?
3. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. HCl and NaCl
    2. HNO_2 and $NaNO_2$
    3. NH_4NO_3 and HNO_3
    4. NH_4NO_3 and NH_3
4. Which solute combinations can make a buffer? Assume all are aqueous solutions.
    1. H_3PO_4 and Na_3PO_4
    2. $NaHCO_3$ and Na_2CO_3
    3. $NaNO_3$ and $Ca(NO_3)_2$
    4. HN_3 and NH_3
5. For each combination in Exercise 3 that is a buffer, write the chemical equations for the reactions of the buffer components when a strong acid and a strong base is added.
6. For each combination in Exercise 4 that is a buffer, write the chemical equations for the reaction of the buffer components when a strong acid and a strong base is added.
7. The complete phosphate buffer system is based on four substances: H_3PO_4, $H_2PO_4^-$, HPO_4^{2-}, and PO_4^{3-}. What different buffer solutions can be made from these substances?
8. Explain why NaBr cannot be a component in either an acidic or a basic buffer.
9. Explain why $Mg(NO_3)_2$ cannot be a component in either an acidic or a basic buffer.

Answers

1. A buffer resists sudden changes in pH. It has a weak acid or base and a salt of that weak acid or base.
2. No. Combining a strong acid and a strong base will produce salt and water. Excess strong acid or strong base will not act as a buffer.

3.
    1. not a buffer
    2. buffer
    3. not a buffer
    4. buffer

4. 1. not a buffer

2. buffer

3. not a buffer

4. not buffer

5. 3b: strong acid: $H^+ + NO_2^- \rightarrow HNO_2$; strong base: $OH^- + HNO_2 \rightarrow H_2O + NO_2^-$; 3d: strong acid: $H^+ + NH_3 \rightarrow NH_4^+$; strong base: $OH^- + NH_4^+ \rightarrow H_2O + NH_3$
6. 4b: strong acid: $H^+ + CO_3^{2-} \rightarrow HCO3^-$; strong base: $OH^- + HCO_3^- \rightarrow H_2O + CO_3^{2-}$;
7. Buffers can be made by combining H_3PO_4 and $H_2PO_4^-$, $H_2PO_4^-$ and HPO_4^{2-}, and HPO_4^{2-} and PO_4^{3-}.
8. NaBr splits up into two ions in solution, Na^+ and Br^-. Na^+ will not react with any added base knowing that NaOH is a strong base. Br- will not react with any added acid knowing that HBr is a strong acid. Because NaBr will not react with any added base or acid, it does not resist change in pH and is not a buffer.
9. $Mg(NO_3)_2$ includes two types of ions, Mg^{2+} and NO_3^-. $Mg(OH)_2$ is strong base and completely dissociates (100% falls apart), so Mg^{2+} will not react with any added base (0% combines with OH^-). HNO_3 is strong acid and completely dissociates (100% falls apart), so NO_3^- will not react with any added acid (0% combines with H^+). Because $Mg(NO_3)_2$ will not react with any added base or acid, it does not resist change in pH and is not a buffer.

Answers

Exercise 11.6.1

b. H_3PO_4 (weak acid) and $H_2PO_4^-$ (conjugate base of H_3PO_4) make a buffer.

c. NH_3 (weak base) and NH_4^+ (conjugate acid of NH_3) make a buffer

This page titled 11.6: Buffers is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.5: Buffers by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

11.S: Acids and Bases (Summary)

To ensure that you understand the material in this chapter, you should review the meanings of the bold terms in the following summary and ask yourself how they relate to the topics in the chapter.

The earliest chemical definition of an acid, the **Arrhenius definition**, says that an **acid** is a compound that increases the amount of hydrogen ion (H^+) in aqueous solution. An **Arrhenius base** is a compound that increases the amount of hydroxide ion (OH^-) in aqueous solution. While most bases are named as ionic hydroxide compounds, aqueous acids have a naming system unique to acids. Acids and bases react together in a characteristic chemical reaction called **neutralization**, in which the products are water and a salt. The principles of stoichiometry, along with the balanced chemical equation for a reaction between an acid and a base, can be used to determine how much of one compound will react with a given amount of the other.

A **Brønsted-Lowry acid** is any substance that donates a proton to another substance. A **Brønsted-Lowry base** is any substance that accepts a proton from another substance. The reaction of ammonia with water to make ammonium ions and hydroxide ions can be used to illustrate Brønsted-Lowry acid and base behavior.

Some compounds can either donate or accept protons, depending on the circumstances. Such compounds are called **amphiprotic**. Water is one example of an amphiprotic compound. One result of water being amphiprotic is that a water molecule can donate a proton to another water molecule to make a hydronium ion and a hydroxide ion. This process is called the **autoionization of water** and occurs in any sample of water.

Not all acids and bases are equal in chemical strength. A **strong acid** is an acid whose molecules are all dissociated into ions in aqueous solution. Hydrochloric acid is an example of a strong acid. Similarly, a **strong base** is a base whose molecules are dissociated into ions in aqueous solution. Sodium hydroxide is an example of a strong base. Any acid or base whose molecules are not all dissociated into ions in aqueous solution is a **weak acid** or a **weak base**. Solutions of weak acids and weak bases reach a **chemical equilibrium** between the un-ionized form of the compound and the dissociated ions. It is a dynamic equilibrium because acid and base molecules are constantly dissociating into ions and reassociating into neutral molecules.

The **pH** scale is a scale used to express the concentration of hydrogen ions in solution. A neutral solution, neither acidic nor basic, has a pH of 7. Acidic solutions have a pH lower than 7, while basic solutions have a pH higher than 7.

Buffers are solutions that resist dramatic changes in pH when an acid or a base is added to them. They contain a weak acid and a salt of that weak acid, or a weak base and a salt of that weak base. When a buffer is present, any strong acid reacts with the anion of the salt, forming a weak acid and minimizing the presence of hydrogen ions in solution. Any strong base reacts with the weak acid, minimizing the amount of additional hydroxide ions in solution. However, buffers only have limited **capacity**; there is a limit to the amount of strong acid or strong base any given amount of buffer will react with.

This page titled 11.S: Acids and Bases (Summary) is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Eden Francis via source content that was edited to the style and standards of the LibreTexts platform.

- 10.S: Acids and Bases (Summary) by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/introduction-to-chemistry-general-organic-and-biological.

INDEX

A

acid strength
11.5: The Strengths of Acids and Bases
Arrhenius acid
11.2: Arrhenius Definition of Acids and Bases
Arrhenius base
11.2: Arrhenius Definition of Acids and Bases
atomic mass
4.4: Nuclei of Atoms
atomic model
4.7: Electron Arrangements

B

base strength
11.5: The Strengths of Acids and Bases
Bohr Atom
4.6: Bohr Diagrams of Atoms and Ions
Bohr Diagram
4.6: Bohr Diagrams of Atoms and Ions
Boyle's law
9.5: Gas Laws

C

charge balance
5.3: Ions
Charles's law
9.5: Gas Laws
chemical change
3.5: Physical and Chemical Changes- the Macroscopic Level
chemical processes
4.7: Electron Arrangements
chemical property
3.8: Differences in Matter- Physical and Chemical Properties
chemistry equations
7.3: Chemical Equations
colligative properties
10.5: Properties of Solutions
combination reaction
7.5: Some Types of Chemical Reactions
combustion reaction
7.5: Some Types of Chemical Reactions
common system
5.5: Ionic Nomenclature
compound
3.2: Classifying Matter- the Macroscopic Level
conversion factor
2.1: Converting Units
8.4: Mole-Mass Conversions
covalent bond
6: Covalent Bonding and Simple Molecular Compounds
6.2: Covalent Bonds

D

Dalton's Law
9.5: Gas Laws
decomposition reaction
7.5: Some Types of Chemical Reactions
dissociation
10.4: The Dissolution Process
duet rule
5.2: Two Types of Bonding
dynamic equilibrium
11.5: The Strengths of Acids and Bases

E

electron arrangement
4.7: Electron Arrangements
electron configurations
4.7: Electron Arrangements
element
3.2: Classifying Matter- the Macroscopic Level
energy levels
4.7: Electron Arrangements

H

hydration
10.4: The Dissolution Process
hypothesis
1.3: Hypothesis, Theories, and Laws

I

ideal gas law
9.5: Gas Laws
ion formation
4.7: Electron Arrangements
ionic bonding
5: Ionic Bonding and Simple Ionic Compounds
ionic nomenclature
5.5: Ionic Nomenclature
isotopes
4.4: Nuclei of Atoms

L

law
1.3: Hypothesis, Theories, and Laws
law of conservation of mass
7.2: The Law of Conservation of Matter
Le Chatelier's principle
10.7: Le Chatelier's Principle
learning outcomes
4.7: Electron Arrangements

M

mixture
3.2: Classifying Matter- the Macroscopic Level
molecules
6: Covalent Bonding and Simple Molecular Compounds
monatomic anions
5.5: Ionic Nomenclature

N

neutralization
11.2: Arrhenius Definition of Acids and Bases

O

octet rule
4.7: Electron Arrangements
osmosis
10.8: Osmosis and Diffusion
osmotic pressure
10.5: Properties of Solutions

P

physical change
3.5: Physical and Chemical Changes- the Macroscopic Level
physical property
3.8: Differences in Matter- Physical and Chemical Properties

R

redox reaction
7.6: Oxidation-Reduction (Redox) Reactions
Redox Reactions (Organic Chemistry)
7.7: Redox Reactions in Organic Chemistry and Biochemistry
Ringer's lactate
10.1: Prelude to Solutions

S

scientific notation
1.6: Scientific Notation
Separation of Mixtures
3.5: Physical and Chemical Changes- the Macroscopic Level
significant figures
2.1: Converting Units
solubility
10.3: Concentration
solvation
10.4: The Dissolution Process
stability of atom
4.7: Electron Arrangements
stock system
5.5: Ionic Nomenclature
stoichiometric factor
8.5: Mole-Mole Relationships in Chemical Reactions
stoichiometry
7.4: Quantitative Relationships Based on Chemical Equations
8.5: Mole-Mole Relationships in Chemical Reactions
strong electrolytes
10.4: The Dissolution Process

T

taxol
8.6: Mole-Mass and Mass-Mass Problems

temperature
2.5: Temperature
theory
1.3: Hypothesis, Theories, and Laws

V

valence electrons
4.7: Electron Arrangements
vapor pressure depression
10.5: Properties of Solutions
VSEPR
6.6: Characteristics of Molecules and VSEPR

W

weak electrolytes
10.4: The Dissolution Process

This page titled Index is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

GLOSSARY

S

Sample Word 1 | Sample Definition 1

This page titled Glossary is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by LibreTexts.

DETAILED LICENSING

Overview

Title: CH 104Z - Introduction to Chemistry

Webpages: 106

Applicable Restrictions: Noncommercial

All licenses found:

- CC BY-NC-SA 4.0: 94.3% (100 pages)
- Undeclared: 5.7% (6 pages)

By Page

- CH 104Z - Introduction to Chemistry - *CC BY-NC-SA 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Introduction to Chemistry, the Scientific Method, and Measurement - *CC BY-NC-SA 4.0*
    - 1.1: What is Chemistry? - *CC BY-NC-SA 4.0*
    - 1.2: The Scientific Method - *CC BY-NC-SA 4.0*
    - 1.3: Hypothesis, Theories, and Laws - *CC BY-NC-SA 4.0*
    - 1.4: Types of Observations, Precision and Measurement - *CC BY-NC-SA 4.0*
    - 1.5: Experimental Design - *CC BY-NC-SA 4.0*
    - 1.6: Scientific Notation - *CC BY-NC-SA 4.0*
    - 1.7: The International System of Units - *CC BY-NC-SA 4.0*
  - 2: Unit Conversion - *CC BY-NC-SA 4.0*
    - 2.1: Converting Units - *CC BY-NC-SA 4.0*
    - 2.2: More Multi-Step Conversion Problems - *CC BY-NC-SA 4.0*
    - 2.3: Units in the Numerator and the Denominator - *CC BY-NC-SA 4.0*
    - 2.4: Density - *CC BY-NC-SA 4.0*
    - 2.5: Temperature - *CC BY-NC-SA 4.0*
    - 2.6: Dosage Calculations - *CC BY-NC-SA 4.0*
    - 2.S: Chemistry, Matter, and Measurement (Summary) - *CC BY-NC-SA 4.0*
  - 3: Classification of Matter- Properties and Changes - *CC BY-NC-SA 4.0*
    - 3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
    - 3.2: Classifying Matter- the Macroscopic Level - *CC BY-NC-SA 4.0*
    - 3.3: Classifying Matter- the Nanoscopic Level - *CC BY-NC-SA 4.0*
    - 3.4: The Elements - *CC BY-NC-SA 4.0*
    - 3.5: Physical and Chemical Changes- the Macroscopic Level - *CC BY-NC-SA 4.0*
    - 3.6: Physcial and Chemical Changes- the Nanoscopic Level - *CC BY-NC-SA 4.0*
    - 3.7: Separating Mixtures through Physcial Changes - *CC BY-NC-SA 4.0*
    - 3.8: Differences in Matter- Physical and Chemical Properties - *CC BY-NC-SA 4.0*
  - 4: Atoms and the Periodic Table - *CC BY-NC-SA 4.0*
    - 4.1: Prelude to Atoms, and the Periodic Table - *CC BY-NC-SA 4.0*
    - 4.2: Atomic Theory - *CC BY-NC-SA 4.0*
    - 4.3: The Structure of Atoms - *CC BY-NC-SA 4.0*
    - 4.4: Nuclei of Atoms - *CC BY-NC-SA 4.0*
    - 4.5: Atomic Masses - *CC BY-NC-SA 4.0*
    - 4.6: Bohr Diagrams of Atoms and Ions - *CC BY-NC-SA 4.0*
    - 4.7: Electron Arrangements - *CC BY-NC-SA 4.0*
    - 4.8: The Periodic Table - *CC BY-NC-SA 4.0*
    - 4.S: Elements, Atoms, and the Periodic Table (Summary) - *CC BY-NC-SA 4.0*
  - 5: Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.1: Prelude to Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.2: Two Types of Bonding - *CC BY-NC-SA 4.0*
    - 5.3: Ions - *CC BY-NC-SA 4.0*
    - 5.4: Formulas for Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.5: Ionic Nomenclature - *CC BY-NC-SA 4.0*
    - 5.6: Formula Mass - *CC BY-NC-SA 4.0*
    - 5.7: Characteristics of Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.S: Ionic Bonding and Simple Ionic Compounds (Summary) - *CC BY-NC-SA 4.0*
  - 6: Covalent Bonding and Simple Molecular Compounds - *CC BY-NC-SA 4.0*
    - 6.1: Prelude to Covalent Bonding and Simple

Molecular Compounds - *CC BY-NC-SA 4.0*
- 6.2: Covalent Bonds - *CC BY-NC-SA 4.0*
- 6.3: Covalent Compounds - Formulas and Names - *CC BY-NC-SA 4.0*
- 6.4: Drawing Lewis Structures - *CC BY-NC-SA 4.0*
- 6.5: Characteristics of Covalent Bonds - *CC BY-NC-SA 4.0*
- 6.6: Characteristics of Molecules and VSEPR - *CC BY-NC-SA 4.0*
- 6.7: Organic Chemistry - *CC BY-NC-SA 4.0*
- 6.S: Covalent Bonding and Simple Molecular Compounds (Summary) - *CC BY-NC-SA 4.0*

◦ 7: Introduction to Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.1: Prelude to Introduction to Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.2: The Law of Conservation of Matter - *CC BY-NC-SA 4.0*
- 7.3: Chemical Equations - *CC BY-NC-SA 4.0*
- 7.4: Quantitative Relationships Based on Chemical Equations - *CC BY-NC-SA 4.0*
- 7.5: Some Types of Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.6: Oxidation-Reduction (Redox) Reactions - *CC BY-NC-SA 4.0*
- 7.7: Redox Reactions in Organic Chemistry and Biochemistry - *CC BY-NC-SA 4.0*
- 7.S: Introduction to Chemical Reactions (Summary) - *CC BY-NC-SA 4.0*

◦ 8: Quantities in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.1: Prelude to Quantities in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.2: The Mole - *CC BY-NC-SA 4.0*
- 8.3: Atomic and Molar Masses - *CC BY-NC-SA 4.0*
- 8.4: Mole-Mass Conversions - *CC BY-NC-SA 4.0*
- 8.5: Mole-Mole Relationships in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.6: Mole-Mass and Mass-Mass Problems - *CC BY-NC-SA 4.0*
- 8.S: Quantities in Chemical Reactions (Summary) - *CC BY-NC-SA 4.0*

◦ 9: Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
- 9.1: Prelude to Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
- 9.2: Intermolecular Interactions - *CC BY-NC-SA 4.0*
- 9.3: Solids and Liquids - *CC BY-NC-SA 4.0*
- 9.4: Gases and Pressure - *CC BY-NC-SA 4.0*
- 9.5: Gas Laws - *CC BY-NC-SA 4.0*
- 9.S: Solids, Liquids, and Gases (Summary) - *CC BY-NC-SA 4.0*

◦ 10: Solutions - *CC BY-NC-SA 4.0*
- 10.1: Prelude to Solutions - *CC BY-NC-SA 4.0*
- 10.2: Solutions - *CC BY-NC-SA 4.0*
- 10.3: Concentration - *CC BY-NC-SA 4.0*
- 10.4: The Dissolution Process - *CC BY-NC-SA 4.0*
- 10.5: Properties of Solutions - *CC BY-NC-SA 4.0*
- 10.6: Chemical Equilibrium - *CC BY-NC-SA 4.0*
- 10.7: Le Chatelier's Principle - *CC BY-NC-SA 4.0*
- 10.8: Osmosis and Diffusion - *CC BY-NC-SA 4.0*
- 10.S: Solutions (Summary) - *CC BY-NC-SA 4.0*

◦ 11: Acids and Bases - *CC BY-NC-SA 4.0*
- 11.1: Prelude to Acids and Bases - *CC BY-NC-SA 4.0*
- 11.2: Arrhenius Definition of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.3: Brønsted-Lowry Definition of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.4: Water - Both an Acid and a Base - *CC BY-NC-SA 4.0*
- 11.5: The Strengths of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.6: Buffers - *CC BY-NC-SA 4.0*
- 11.S: Acids and Bases (Summary) - *CC BY-NC-SA 4.0*

◦ Back Matter - *CC BY-NC-SA 4.0*
- Index - *CC BY-NC-SA 4.0*
- Glossary - *CC BY-NC-SA 4.0*
- Detailed Licensing - *CC BY-NC-SA 4.0*
- Detailed Licensing - *Undeclared*

This page titled CH 104Z - Introduction to Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Stephanie Bryan & Kenneth Friedrich.

DETAILED LICENSING

Overview

Title: CH 104Z - Introduction to Chemistry

Webpages: 106

Applicable Restrictions: Noncommercial

All licenses found:

- CC BY-NC-SA 4.0: 94.3% (100 pages)
- Undeclared: 5.7% (6 pages)

By Page

- CH 104Z - Introduction to Chemistry - *CC BY-NC-SA 4.0*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Introduction to Chemistry, the Scientific Method, and Measurement - *CC BY-NC-SA 4.0*
    - 1.1: What is Chemistry? - *CC BY-NC-SA 4.0*
    - 1.2: The Scientific Method - *CC BY-NC-SA 4.0*
    - 1.3: Hypothesis, Theories, and Laws - *CC BY-NC-SA 4.0*
    - 1.4: Types of Observations, Precision and Measurement - *CC BY-NC-SA 4.0*
    - 1.5: Experimental Design - *CC BY-NC-SA 4.0*
    - 1.6: Scientific Notation - *CC BY-NC-SA 4.0*
    - 1.7: The International System of Units - *CC BY-NC-SA 4.0*
  - 2: Unit Conversion - *CC BY-NC-SA 4.0*
    - 2.1: Converting Units - *CC BY-NC-SA 4.0*
    - 2.2: More Multi-Step Conversion Problems - *CC BY-NC-SA 4.0*
    - 2.3: Units in the Numerator and the Denominator - *CC BY-NC-SA 4.0*
    - 2.4: Density - *CC BY-NC-SA 4.0*
    - 2.5: Temperature - *CC BY-NC-SA 4.0*
    - 2.6: Dosage Calculations - *CC BY-NC-SA 4.0*
    - 2.S: Chemistry, Matter, and Measurement (Summary) - *CC BY-NC-SA 4.0*
  - 3: Classification of Matter- Properties and Changes - *CC BY-NC-SA 4.0*
    - 3.1: Introduction to the Nanoscopic Level- Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
    - 3.2: Classifying Matter- the Macroscopic Level - *CC BY-NC-SA 4.0*
    - 3.3: Classifying Matter- the Nanoscopic Level - *CC BY-NC-SA 4.0*
    - 3.4: The Elements - *CC BY-NC-SA 4.0*
    - 3.5: Physical and Chemical Changes- the Macroscopic Level - *CC BY-NC-SA 4.0*
    - 3.6: Physcial and Chemical Changes- the Nanoscopic Level - *CC BY-NC-SA 4.0*
    - 3.7: Separating Mixtures through Physcial Changes - *CC BY-NC-SA 4.0*
    - 3.8: Differences in Matter- Physical and Chemical Properties - *CC BY-NC-SA 4.0*
  - 4: Atoms and the Periodic Table - *CC BY-NC-SA 4.0*
    - 4.1: Prelude to Atoms, and the Periodic Table - *CC BY-NC-SA 4.0*
    - 4.2: Atomic Theory - *CC BY-NC-SA 4.0*
    - 4.3: The Structure of Atoms - *CC BY-NC-SA 4.0*
    - 4.4: Nuclei of Atoms - *CC BY-NC-SA 4.0*
    - 4.5: Atomic Masses - *CC BY-NC-SA 4.0*
    - 4.6: Bohr Diagrams of Atoms and Ions - *CC BY-NC-SA 4.0*
    - 4.7: Electron Arrangements - *CC BY-NC-SA 4.0*
    - 4.8: The Periodic Table - *CC BY-NC-SA 4.0*
    - 4.S: Elements, Atoms, and the Periodic Table (Summary) - *CC BY-NC-SA 4.0*
  - 5: Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.1: Prelude to Ionic Bonding and Simple Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.2: Two Types of Bonding - *CC BY-NC-SA 4.0*
    - 5.3: Ions - *CC BY-NC-SA 4.0*
    - 5.4: Formulas for Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.5: Ionic Nomenclature - *CC BY-NC-SA 4.0*
    - 5.6: Formula Mass - *CC BY-NC-SA 4.0*
    - 5.7: Characteristics of Ionic Compounds - *CC BY-NC-SA 4.0*
    - 5.S: Ionic Bonding and Simple Ionic Compounds (Summary) - *CC BY-NC-SA 4.0*
  - 6: Covalent Bonding and Simple Molecular Compounds - *CC BY-NC-SA 4.0*
    - 6.1: Prelude to Covalent Bonding and Simple

Molecular Compounds - *CC BY-NC-SA 4.0*
- 6.2: Covalent Bonds - *CC BY-NC-SA 4.0*
- 6.3: Covalent Compounds - Formulas and Names - *CC BY-NC-SA 4.0*
- 6.4: Drawing Lewis Structures - *CC BY-NC-SA 4.0*
- 6.5: Characteristics of Covalent Bonds - *CC BY-NC-SA 4.0*
- 6.6: Characteristics of Molecules and VSEPR - *CC BY-NC-SA 4.0*
- 6.7: Organic Chemistry - *CC BY-NC-SA 4.0*
- 6.S: Covalent Bonding and Simple Molecular Compounds (Summary) - *CC BY-NC-SA 4.0*

◦ 7: Introduction to Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.1: Prelude to Introduction to Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.2: The Law of Conservation of Matter - *CC BY-NC-SA 4.0*
- 7.3: Chemical Equations - *CC BY-NC-SA 4.0*
- 7.4: Quantitative Relationships Based on Chemical Equations - *CC BY-NC-SA 4.0*
- 7.5: Some Types of Chemical Reactions - *CC BY-NC-SA 4.0*
- 7.6: Oxidation-Reduction (Redox) Reactions - *CC BY-NC-SA 4.0*
- 7.7: Redox Reactions in Organic Chemistry and Biochemistry - *CC BY-NC-SA 4.0*
- 7.S: Introduction to Chemical Reactions (Summary) - *CC BY-NC-SA 4.0*

◦ 8: Quantities in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.1: Prelude to Quantities in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.2: The Mole - *CC BY-NC-SA 4.0*
- 8.3: Atomic and Molar Masses - *CC BY-NC-SA 4.0*
- 8.4: Mole-Mass Conversions - *CC BY-NC-SA 4.0*
- 8.5: Mole-Mole Relationships in Chemical Reactions - *CC BY-NC-SA 4.0*
- 8.6: Mole-Mass and Mass-Mass Problems - *CC BY-NC-SA 4.0*
- 8.S: Quantities in Chemical Reactions (Summary) - *CC BY-NC-SA 4.0*

◦ 9: Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
- 9.1: Prelude to Solids, Liquids, and Gases - *CC BY-NC-SA 4.0*
- 9.2: Intermolecular Interactions - *CC BY-NC-SA 4.0*
- 9.3: Solids and Liquids - *CC BY-NC-SA 4.0*
- 9.4: Gases and Pressure - *CC BY-NC-SA 4.0*
- 9.5: Gas Laws - *CC BY-NC-SA 4.0*
- 9.S: Solids, Liquids, and Gases (Summary) - *CC BY-NC-SA 4.0*

◦ 10: Solutions - *CC BY-NC-SA 4.0*
- 10.1: Prelude to Solutions - *CC BY-NC-SA 4.0*
- 10.2: Solutions - *CC BY-NC-SA 4.0*
- 10.3: Concentration - *CC BY-NC-SA 4.0*
- 10.4: The Dissolution Process - *CC BY-NC-SA 4.0*
- 10.5: Properties of Solutions - *CC BY-NC-SA 4.0*
- 10.6: Chemical Equilibrium - *CC BY-NC-SA 4.0*
- 10.7: Le Chatelier's Principle - *CC BY-NC-SA 4.0*
- 10.8: Osmosis and Diffusion - *CC BY-NC-SA 4.0*
- 10.S: Solutions (Summary) - *CC BY-NC-SA 4.0*

◦ 11: Acids and Bases - *CC BY-NC-SA 4.0*
- 11.1: Prelude to Acids and Bases - *CC BY-NC-SA 4.0*
- 11.2: Arrhenius Definition of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.3: Brønsted-Lowry Definition of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.4: Water - Both an Acid and a Base - *CC BY-NC-SA 4.0*
- 11.5: The Strengths of Acids and Bases - *CC BY-NC-SA 4.0*
- 11.6: Buffers - *CC BY-NC-SA 4.0*
- 11.S: Acids and Bases (Summary) - *CC BY-NC-SA 4.0*

◦ Back Matter - *CC BY-NC-SA 4.0*
- Index - *CC BY-NC-SA 4.0*
- Glossary - *CC BY-NC-SA 4.0*
- Detailed Licensing - *CC BY-NC-SA 4.0*
- Detailed Licensing - *Undeclared*

This page titled CH 104Z - Introduction to Chemistry is shared under a CC BY-NC-SA 4.0 license and was authored, remixed, and/or curated by Stephanie Bryan & Kenneth Friedrich.