EARTH SCIENCE



Lumen Learning
Lumen Learning



Book: Earth Science (Lumen)

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 05/21/2026

TABLE OF CONTENTS

Licensing

1: Introduction to Earth Science

- 1.1: Principles of Science
- 1.2: Earth Science and Its Branches

2: Minerals

- 2.1: Materials of Earth's Crust
- 2.2: Atoms to Molecules
- 2.3: States of Matter
- 2.4: Mineral Identification
- 2.5: Minerals and Mineral Groups
- 2.6: Mineral Formation
- 2.7: Mining and Mineral Use

3: Rocks

- 3.1: Igneous Rocks
- 3.2: Sedimentary Rocks
- 3.3: Metamorphic Rocks

4: Earth's Interior Processes

- 4.1: Earth's Interior
- 4.2: Plate Tectonics
- 4.3: Volcanoes
- 4.4: Earthquakes

5: Volcanoes

- 5.1: Volcanic Activity
- 5.2: Types of Volcanoes
- 5.3: Volcanic Landforms and Geothermal Activity
- 5.4: Volcanic Eruptions

6: Surface Water

- 6.1: Earth's Fresh Water
- 6.2: Streams and Rivers
- 6.3: Groundwater
- 6.4: Water Resources
- 6.5: Floods
- 6.6: Wetlands

7: The Ocean

- 7.1: Introduction to the Oceans
- 7.2: Composition of Ocean Water

- 7.3: The Seafloor
- 7.4: Ocean Currents
- 7.5: Oceans and Coastal Environments
- 7.6: Tides
- 7.7: Currents

8: Coastal Processes
- 8.1: Waves and Coastal Features

9: Mountain Building
- 9.1: Tectonic Stress and Geologic Structures
- 9.2: Mountain Formation

10: Geologic History
- 10.1: Relative Ages of Rocks
- 10.2: Absolute Ages of Rocks
- 10.3: Fossils
- 10.4: History of Earth's Life Forms
- 10.5: Early Earth
- 10.6: Geologic Time Scale

11: Ocean Water and Life
- 11.1: Thermohaline circulation
- 11.2: Thermocline
- 11.3: Tsunamis
- 11.4: Ocean Organisms

12: The Atmosphere
- 12.1: The Atmosphere

13: Weather
- 13.1: Weather Processes and Systems

14: Astronomy
- 14.1: The Universe

15: Stellar Evolution and Our Sun
- 15.1: Stars
- 15.2: Stellar Life Cycle
- 15.3: The Sun
- 15.4: The Sun and the Earth-Moon System
- 15.5: Sun – Earth Relationship- The Seasons

16: Light and the Sun
- 16.1: Electromagnetic Spectrum
- 16.2: Doppler Effect

- 16.3: Telescopes
- 16.4: Nuclear Fusion

17: The Solar System

- 17.1: Introduction to the Solar System
- 17.2: Inner Planets
- 17.3: Outer Planets
- 17.4: Other Objects in the Solar System

18: Climate and Climate Change

- 18.1: Climate Systems and Change
- 18.2: Climate and Weather
- 18.3: Global Climate Change

19: Deserts

- 19.1: Deserts
- 19.2: Desertification
- 19.3: Dunes
- 19.4: Video- Desertification

20: Glaciers

- 20.1: Glacial Erosion and Deposition
- 20.2: Glaciers

Index

Glossary

Detailed Licensing

Licensing

*A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.*

CHAPTER OVERVIEW

1: Introduction to Earth Science

1.1: Principles of Science

1.2: Earth Science and Its Branches

This page titled 1: Introduction to Earth Science is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

1.1: Principles of Science

Science is Different from Other Ways of Understanding the World

If someone asks you “what is science?” you might say that it’s a bunch of facts and explanations. But that’s only part of the story. Science is a knowledge base. But science is also a way of learning about the world.

Science is different from other types of information because scientists follow rigorous methods to learn about the world. A scientific idea must be testable and falsifiable. Ideas that are not supported by observations and data are revised or thrown out. The distinction between science and other ways of understanding the world is important because scientific information has been obtained with much more rigor than ideas that are the result of opinion, gut feelings, or faith. Scientists use the scientific method to answer questions about the natural world. The scientific method is not linear but takes on this basic structure: Ask a question, do background research, propose a hypothesis, test the hypothesis using data from observations and experiments, continue testing the hypothesis if it holds up or find a new hypothesis if it does not, eventually create a theory. A theory is an explanation of a complicated set of phenomena that fits virtually all of the available data. The theories of evolution, plate tectonics, and climate change are crucial to understanding earth science.

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-Concepts-For-High-School/section/1.0/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 1.1: Principles of Science is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

1.2: Earth Science and Its Branches

Lesson Objectives

- Define and describe Earth Science as a general field with many branches.
- Identify the field of geology as a branch of Earth Science that deals with the solid part of the Earth.
- Describe the field of oceanography as a branch of Earth Science that has several subdivisions that deal with the various aspects of the ocean.
- Define the field of meteorology as a branch of Earth Science that deals with the atmosphere.
- Understand that astronomy is an extension of Earth Science that examines other parts of the solar system and universe.
- List some of the other branches of Earth Science, and how they relate to the study of the Earth.

Overview of Earth Science



Figure 1.9: Earth as seen from Apollo 17.

Earth is the mighty planet upon which we all live. Only recently have humans begun to understand the complexity of this planet. In fact, it was only a few hundred years ago that we discovered that Earth was just a tiny part of an enormous galaxy, which in turn is a small part of an even greater universe. Earth Science deals with any and all aspects of the Earth. Our Earth has molten lava, icy mountain peaks, steep canyons and towering waterfalls. Earth scientists study the atmosphere high above us as well as the planet's core far beneath us. Earth scientists study parts of the Earth as big as continents and as small as the tiniest atom. In all its wonder, Earth scientists seek to understand the beautiful sphere on which we thrive (Figure 1.9).

Because the Earth is so large and science is so complex, Earth scientists specialize in studying just a small aspect of our Earth. Since all of the branches are connected together, specialists work together to answer complicated questions. Let's look at some important branches of Earth Science.

Geology

Geology is the study of the solid matter that makes up Earth. Anything that is solid, like rocks, minerals, mountains, and canyons is part of geology. Geologists study the way that these objects formed, their composition, how they interact with one another, how they erode, and how humans can use them. Geology has so many branches that most geologists become specialists in one area. For example, a mineralogist studies the composition and structure of minerals such as halite (rock salt), quartz, calcite, and magnetite (Figure 1.10).



Figure 1.10: Mineralogists focus on all kinds of minerals.



Figure 1.11: Seismographs are used to measure earthquakes and pinpoint their origins.

A volcanologist braves the high temperatures and molten lava of volcanoes. Seismologists study earthquakes and the forces of the Earth that create them. Seismologists monitor earthquakes worldwide to help protect people and property from harm (Figure 1.11). Scientists interested in fossils are paleontologists, while scientists who compare other planets' geologies to that of the Earth are called planetary geologists. There are geologists who only study the Moon. Some geologists look for petroleum, others are specialists on soil. Geochronologists study how old rocks are and determine how different rock layers formed. There are so many specialties in geology that there is probably an expert in almost anything you can think of related to the Earth (Figure 1.12).



Figure 1.12: Geology is the study of the solid Earth and its processes.

Oceanography

Oceanography is the study of everything in the ocean environment. More than 70% of the Earth's surface is covered with water. Most of that water is found in the oceans. Recent technology has allowed us to go to the deepest parts of the ocean, yet much of the ocean remains truly unexplored. Some people call the ocean the last frontier. But it is a frontier already deeply influence by human

activity. As the human population gets ever bigger, we are affecting the ocean in many ways. Populations of fish and other marine species have plummeted because of overfishing; contaminants are polluting the waters, and global warming caused by greenhouse gases is melting the thick ice caps. As ocean waters warm, the water expands and, along with the melting ice caps, causes sea levels to rise.

Climatologists help us understand the climate and how it will change in the future in response to global warming. Oceanographers study the vast seas and help us to understand all that happens in the water world. As with geology, there are many branches of oceanography. Physical oceanography is the study of the processes in the ocean itself, like waves and ocean currents (Figure 1.13). Marine geology uses geology to study ocean earthquakes, mountains, and trenches. Chemical oceanography studies the natural elements in ocean water and pollutants.



Figure 1.13: Physical oceanography studies things like currents and waves.

Climatology and Meteorology

Meteorologists don't study meteors — they study the atmosphere! Perhaps this branch of Earth Science is strangely named but it is very important to living creatures like humans. **Meteorology** includes the study of weather patterns, clouds, hurricanes, and tornadoes. Using modern technology like radars and satellites, meteorologists work to predict or forecast the weather. Because of more accurate forecasting techniques, meteorologists can help us to prepare for major storms, as well as help us know when we should go on picnics.

Climatologists and other atmospheric scientists study the whole atmosphere, which is a thin layer of gas that surrounds the Earth. Most of it is within about 10 – 11 kilometers of the Earth's surface. Earth's atmosphere is denser than Mars's thin atmosphere, where the average temperature is -63° C, and not as thick as the dense atmosphere on Venus, where carbon dioxide in the atmosphere makes it hot and sulfuric acid rains in the upper atmosphere. The atmosphere on Earth is just dense enough to even out differences in temperature from the equator to the poles, and contains enough oxygen for animals to breathe.

Over the last several decades, climatologists studying the gases in our atmosphere have found that humans are putting higher levels of carbon dioxide into the air by burning fossil fuels (Figure 1.14). Normally, the atmosphere contains small amounts of carbon dioxide, however, with increases in the burning of fossil fuels more than normal amounts are present. These higher concentrations of carbon dioxide can lead to higher surface temperatures. Much of climate change science is based on the increases of greenhouse gases, like carbon dioxide, in the atmosphere and the effect those higher concentrations have on global temperatures. Climatologists can help us better understand the climate and how it may change in the future in response to different amounts of greenhouse gases and other factors (Figure 1.15).



Figure 1.14: Man-made carbon dioxide released into the atmosphere has been linked to rises in atmospheric temperatures.



Figure 1.15: When hurricanes are accurately forecast by meteorologists, many lives can be saved.

Astronomy



Figure 1.16: The Hubble Space Telescope

Astronomers have proven that our Earth and solar system are not the only set of planets in the universe. As of June 2015, over a thousand planets outside our solar system had been discovered. Although no one can be sure how many there are, astronomers estimate that there are billions of other planets. In addition, the universe contains black holes, other galaxies, asteroids, comets, and nebula. As big as Earth seems to us, the entire universe is vastly greater. Our Earth is an infinitesimally small part of our universe.

Astronomers use resources on the Earth to study physical things beyond the Earth. They use a variety of instruments like optical telescopes and radio telescopes to see things far beyond what the human eye can see. Spacecraft travel great distances in space to send us information on faraway places, while telescopes in orbit observe astronomical bodies from the darkness of space (Figure 1.16).

Astronomers ask a wide variety of questions. Astronomers could study how an object or energy outside of Earth could affect us. An impact from an asteroid could have terrible effects for life on Earth. Strong bursts of energy from the sun, called solar flares, can knock out a power grid or disturb radio, television or cell phone communications. But astronomers ask bigger questions too. How was the universe created? Are there other planets on which we might live? Are there resources that we could use? Is there other life out there? Astronomy also relies on Earth Science, when scientists compare what we know about life on Earth to the chances of finding life beyond this planet.

Other Branches of Earth Science

Geology, oceanography, and meteorology represent a large part of Earth science, while astronomy represents science beyond Earth. However, there are still many smaller branches of science that deal with the Earth or interact greatly with Earth sciences. Most branches of science are connected with other branches of science in some way or another. A biologist who studies monkeys in rainforests must be concerned with the water cycle that brings the rain to the rainforests. She must understand the organic chemistry of the food the monkeys eat, as well as the behavior between the monkeys. She might examine the soil in which the trees of the rainforest grow. She must even understand the economy of the rainforest to understand reasons for its destruction. This is just one example of how all branches of science are connected.

Below are examples of a few branches of science that are directly related to Earth science. Environmental scientists study the ways that humans interact with the Earth and the effects of that interaction. We hope to find better ways of sustaining the environment. Biogeography is a branch of science that investigates changes in populations of organisms in relation to place over time. These scientists attempt to explain the causes of species' movement in history. Ecologists focus on ecosystems, the complex relationship of all life forms and the environment in a given place (Figure 1.17). They try to predict the chain reactions that could occur when one part of the ecosystem is disrupted.




Figure 1.17: In a marine ecosystem, coral, fish, and other sea life depend on each other for survival.

As opposed to an oceanographer, a limnologist studies inland waters like rivers and lakes. A hydrogeologist focuses on underground water found between soil and rock particles, while glaciologists study glaciers and ice.

None of these scientific endeavors would be possible without geographers who explore the features of the surface and work with cartographers, who make maps. Stratigraphy is another area of Earth science which examines layers of rock beneath the surface (Figure 1.18). This helps us to understand the geological history of the Earth. There is a branch of science for every interest and each is related to the others.




Figure 1.18: Folded strata are layers in the rock that have bent over time. Stratigraphy attempts to explain these layers and the geologic history of the area.

Review Questions

1. What are three major branches of Earth science?
2. What branch of science deals with stars & galaxies beyond the Earth?
3. List important functions of Earth scientists.
4. What do you think is the focus of a meteorologist?
5. An ecologist notices that an important coral reef is dying off. She believes that it has to do with some pollution from a local electric plant. What type of scientist might help her analyze the water for contamination?
6. Design an experiment that you could conduct in any branch of Earth science. Identify the independent variable and dependent variable. What safety precautions would you have to take?

Vocabulary

astronomers
Scientists who study the universe, galaxies, and stars.

geology
The study of the rocks, processes, and history of Earth.

meteorology
Study of the atmosphere, weather, and storms.

oceanography
Study of the ocean realm in all its aspects.

atmospheric science
the study of the atmosphere and its variations. Includes meteorology and aeronomy.

ecology
the study of the life forms that inhabit Earth, their history, and their interaction with the above.

- Therefore, we must utilize the processes of System Science, in order to fully understand the Earth Systems and its variations as a whole.

Points to Consider

- Why is Earth science so important?
- Which branch of Earth science would you most like to explore?
- What is the biggest problem that we face today? Which Earth scientists may help us to solve the problem?
- What other branches of science or society are related to and necessary for Earth science?

CC licensed content, Shared previously

- **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Earth_Science_and_Its_Branches. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 1.2: Earth Science and Its Branches is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

2: Minerals

2.1: Materials of Earth’s Crust
2.2: Atoms to Molecules
2.3: States of Matter
2.4: Mineral Identification
2.5: Minerals and Mineral Groups
2.6: Mineral Formation
2.7: Mining and Mineral Use

This page titled 2: Minerals is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.1: Materials of Earth's Crust

What is Earth's crust made of?

The best way to learn about Earth's crust would be to travel around the world, viewing minerals, rocks and structures in a variety of places to see what they are and how they can be coaxed into telling Earth's story. A simpler thing to do to learn a lot about Earth materials is to visit a museum. In a museum you can see lots of samples with good explanations of what they are and, more importantly, what they tell scientists about our planet. So the next time you're in a major city, find a way to spend a few hours in a natural history museum! In this chapter we'll learn about a variety of Earth materials.

All matter is made of tiny particles. Protons, neutrons, and electrons form atoms that bond together to create molecules. Atoms are the smallest units that have the properties of the element they are and molecules are the smallest units of a compound. For example, water is made of hydrogen and oxygen, but a molecule of water is very different from an atom of hydrogen or an atom of oxygen. The atoms combine to form molecules by different types of chemical bonding. Molecules bond into structures as well. The structures created by molecules form the different types of minerals, most importantly silicates, which are the substances that make up most of Earth's crust. Other important minerals are carbonates and native elements, which are some of the most important materials used by society. Minerals come together to create the three major rock types, igneous, sedimentary, and metamorphic. These rocks are the material part of the rock cycle. Different processes can convert any type of rock into any other type of rock. These processes include weathering and erosion, melting and cooling, and burial and pressure, among others. Each rock contains a story of how it formed and what it formed from. Geologists piece together these stories to understand the geologic past of any region of our planet and of the planet as a whole.

This page titled 2.1: Materials of Earth's Crust is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.2: Atoms to Molecules

What is your brain made of?

Everything you can see, touch, smell, feel, and taste is made of atoms. **Atoms are the basic building-block of all matter** (including you and me, and everyone else you'll ever meet), so if we want to know about what Earth is made of, then we have to know a few things about these incredibly small objects.


Atoms

Everyday experience should convince you that matter is found in myriad forms, yet all the matter you have ever seen is made of atoms, or atoms stuck together in configurations of dizzying complexity. A chemical **element** is a substance that cannot be made into a simpler form by ordinary chemical means. The smallest unit of a chemical element is an **atom** , and all atoms of a particular element are identical.

Parts of an Atom

There are two parts to an atom (**Figure** below):

- At the center of an atom is a **nucleus** made up of two types of particles called protons and neutrons.
  - **Protons** have a positive electrical charge. The number of protons in the nucleus determines what element the atom is.
  - **Neutrons** are about the size of protons but have no charge.
- **Electrons** , much smaller than protons or neutrons, have a negative electrical charge, move at nearly the speed of light, and orbit the nucleus at exact distances, depending on their energy.



Major parts of an atom. What chemical element is this? (Hint: 3 protons, 3 electrons)

Atomic Mass

Because electrons are minuscule compared with protons and neutrons, the number of protons plus neutrons gives the atom its **atomic mass** . All atoms of a given element always have the same number of protons, but may differ in the number of neutrons found in the nucleus.

Ions

Atoms are stable when they have a full outermost electron energy level. To fill its outermost shell, an atom will give, take, or share electrons. When an atom either gains or loses electrons, this creates an **ion** . Ions have either a positive or a negative electrical charge. What is the charge of an ion if the atom loses an electron? An atom with the same number of protons and electrons has no

overall charge, so if an atom loses the negatively charged electron, it will have more protons, therefore, a positive charge. Ions with a positive charge are referred to as a **cation**.What is the charge of an ion if the atom gains an electron? If the atom gains an electron, there will be more electrons and will have a negative charge. **Anions** are ions that have a negative charge.

Molecules

In the previous section we said that many atoms are more stable when they have a net charge: they are more stable as ions. When a cation gets close to an anion, they link up because of their different net charges — positive charges attract negative charges and vice versa. When two or more atoms link up, they create a **molecule** . A molecule of water is made of two atoms of hydrogen (H) and one atom of oxygen (O). The **molecular mass** is the sum of the masses of all the atoms in the molecule. A collection of molecules is called a compound.

Summary

- An atom has negatively-charged electrons in orbit around its nucleus, which is composed of positively-charged protons and neutrons, which have no charge.
- An atom that gains or loses electrons is an ion. Positively charged ions are cations, negatively charged ions are anions.

Lumen Learning authored content

- **License**: *Public Domain: No Known Copyright*

This page titled 2.2: Atoms to Molecules is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.3: States of Matter

A **state of matter** is one of the distinct forms that different phases of matter take on. **Four states of matter are observable in everyday life: solid, liquid, gas, and plasma**. Many other states are known such as Bose–Einstein condensates and neutron-degenerate matter but these only occur in extreme situations such as ultra cold or ultra dense matter. Other states, such as quark–gluon plasmas, are believed to be possible but remain theoretical for now.

Historically, the distinction is made based on qualitative differences in properties. Matter in the solid state maintains a fixed volume and shape, with component particles (atoms, molecules or ions) close together and fixed into place. Matter in the liquid state maintains a fixed volume, but has a variable shape that adapts to fit its container. Its particles are still close together but move freely. Matter in the gaseous state has both variable volume and shape, adapting both to fit its container. Its particles are neither close together nor fixed in place. Matter in the plasma state has variable volume and shape, but as well as neutral atoms, it contains a significant number of ions and electrons, both of which can move around freely. Plasma is the most common form of visible matter in the universe.[1]



The four fundamental states of matter. Clockwise from top left, they are solid, liquid, plasma and gas, represented by an ice sculpture, a drop of water, electrical arcing from a tesla coil, and the air around clouds respectively.

The four fundamental states

Solid


In a solid, the particles (ions, atoms or molecules) are closely packed together. The forces between particles are strong so that the **particles cannot move freely** but can only vibrate. As a result, a **solid has a stable, definite shape, and a definite volume**. Solids can only change their shape by force, as when broken or cut.

In crystalline solids, the particles (atoms, molecules, or ions) are packed in a regularly ordered, repeating pattern. There are various different crystal structures, and the same substance can have more than one structure (or solid phase). For example, iron has a body-centred cubic structure at temperatures below 912 °C, and a face-centred cubic structure between 912 and 1394 °C. Ice has fifteen known crystal structures, or fifteen solid phases, which exist at various temperatures and pressures.[2]

Glasses and other non-crystalline, amorphous solids without long-range order are not thermal equilibrium ground states; therefore they are described below as nonclassical states of matter.

Solids can be transformed into liquids by melting and can also change directly into gases through the process of sublimation.


Liquid



Structure of a classical single atom liquid. Atoms have many nearest neighbors in contact, yet no long-range order is present.

A liquid is a nearly incompressible fluid that conforms to the shape of its container but retains a (nearly) constant volume independent of pressure. The volume is definite if the temperature and pressure are constant. When a solid is heated above its melting point, it becomes liquid, given that the pressure is higher than the triple point of the substance. Intermolecular (or interatomic or interionic) forces are still important, but the molecules have enough energy to move relative to each other and the structure is mobile. This means that the shape of a liquid is not definite but is determined by its container. The volume is usually greater than that of the corresponding solid, the best known exception being water, H_2O. The highest temperature at which a given liquid can exist is its critical temperature.[3]

Gas



The spaces between gas molecules are very big. Gas molecules have very weak or no bonds at all. The molecules in “gas” can move freely and fast.

Main article: Gas


A gas is a compressible fluid. Not only will a gas conform to the shape of its container but it will also expand to fill the container.

In a gas, the molecules have enough kinetic energy so that the effect of intermolecular forces is small (or zero for an ideal gas), and the typical distance between neighboring molecules is much greater than the molecular size. A gas has no definite shape or volume, but occupies the entire container in which it is confined. A liquid may be converted to a gas by heating at constant pressure to the boiling point, or else by reducing the pressure at constant temperature.

At temperatures below its critical temperature, a gas is also called a vapor, and can be liquefied by compression alone without cooling. A vapour can exist in equilibrium with a liquid (or solid), in which case the gas pressure equals the vapor pressure of the liquid (or solid).

A supercritical fluid (SCF) is a gas whose temperature and pressure are above the critical temperature and critical pressure respectively. In this state, the distinction between liquid and gas disappears. A supercritical fluid has the physical properties of a gas, but its high density confers solvent properties in some cases, which leads to useful applications. For example, supercritical carbon dioxide is used to extract caffeine in the manufacture of decaffeinated coffee.[4]

Plasma



In a plasma, electrons are ripped away from their nuclei, forming an electron “sea”. This gives it the ability to conduct electricity.

Main article: Plasma (physics)


Like a gas, plasma does not have definite shape or volume. Unlike gases, plasmas are electrically conductive, produce magnetic fields and electric currents, and respond strongly to electromagnetic forces. Positively charged nuclei swim in a “sea” of freely-moving disassociated electrons, similar to the way such charges exist in conductive metal. In fact it is this electron “sea” that allows matter in the plasma state to conduct electricity.

The plasma state is often misunderstood, but it is actually quite common on Earth, and the majority of people observe it on a regular basis without even realizing it. Lightning, electric sparks, fluorescent lights, neon lights, plasma televisions, some types of flame and the stars are all examples of illuminated matter in the plasma state.

A gas is usually converted to a plasma in one of two ways, either from a huge voltage difference between two points, or by exposing it to extremely high temperatures.

Heating matter to high temperatures causes electrons to leave the atoms, resulting in the presence of free electrons. At very high temperatures, such as those present in stars, it is assumed that essentially all electrons are “free”, and that a very high-energy plasma is essentially bare nuclei swimming in a sea of electrons.

Phase transitions

Main article: Phase transitions 

This diagram illustrates transitions between the four fundamental states of matter.

A state of matter is also characterized by phase transitions. A phase transition indicates a change in structure and can be recognized by an abrupt change in properties. A distinct state of matter can be defined as any set of states distinguished from any other set of states by a phase transition. Water can be said to have several distinct solid states.[5] The appearance of superconductivity is associated with a phase transition, so there are superconductive states. Likewise, ferromagnetic states are demarcated by phase transitions and have distinctive properties. When the change of state occurs in stages the intermediate steps are called mesophases. Such phases have been exploited by the introduction of liquid crystal technology. [6][7]


The state or *phase* of a given set of matter can change depending on pressure and temperature conditions, transitioning to other phases as these conditions change to favor their existence; for example, solid transitions to liquid with an increase in temperature. Near absolute zero, a substance exists as a solid. As heat is added to this substance it melts into a liquid at its melting point, boils into a gas at its boiling point, and if heated high enough would enter a plasma state in which the electrons are so energized that they leave their parent atoms.

Forms of matter that are not composed of molecules and are organized by different forces can also be considered different states of matter. Superfluids (like Fermionic condensate) and the quark–gluon plasma are examples.

In a chemical equation, the state of matter of the chemicals may be shown as (s) for solid, (l) for liquid, and (g) for gas. An aqueous solution is denoted (aq). Matter in the plasma state is seldom used (if at all) in chemical equations, so there is no standard symbol to denote it. In the rare equations that plasma is used in plasma is symbolized as (p).

Non-classical states

Glass

Main article: Glass 

Schematic representation of a random-network glassy form (left) and ordered crystalline lattice (right) of identical chemical composition.

Glass is a non-crystalline or amorphous solid material that exhibits a glass transition when heated towards the liquid state. Glasses can be made of quite different classes of materials: inorganic networks (such as window glass, made of silicate plus additives),

metallic alloys, ionic melts, aqueous solutions, molecular liquids, and polymers. Thermodynamically, a glass is in a metastable state with respect to its crystalline counterpart. The conversion rate, however, is practically zero.

Crystals with some degree of disorder

A plastic crystal is a molecular solid with long-range positional order but with constituent molecules retaining rotational freedom; in an orientational glass this degree of freedom is frozen in a quenched disordered state.

Similarly, in a spin glass magnetic disorder is frozen.

Liquid crystal states

Main article: Liquid crystal

Liquid crystal states have properties intermediate between mobile liquids and ordered solids. Generally, they are able to flow like a liquid, but exhibiting long-range order. For example, the nematic phase consists of long rod-like molecules such as para-azoxyanisole, which is nematic in the temperature range 118–136 °C.[8] In this state the molecules flow as in a liquid, but they all point in the same direction (within each domain) and cannot rotate freely.

Other types of liquid crystals are described in the main article on these states. Several types have technological importance, for example, in liquid crystal displays.

Magnetically ordered

Transition metal atoms often have magnetic moments due to the net spin of electrons that remain unpaired and do not form chemical bonds. In some solids the magnetic moments on different atoms are ordered and can form a ferromagnet, an antiferromagnet or a ferrimagnet.

In a ferromagnet—for instance, solid iron—the magnetic moment on each atom is aligned in the same direction (within a magnetic domain). If the domains are also aligned, the solid is a permanent magnet, which is magnetic even in the absence of an external magnetic field. The magnetization disappears when the magnet is heated to the Curie point, which for iron is 768 °C.

An antiferromagnet has two networks of equal and opposite magnetic moments, which cancel each other out so that the net magnetization is zero. For example, in nickel(II) oxide (NiO), half the nickel atoms have moments aligned in one direction and half in the opposite direction.

In a ferrimagnet, the two networks of magnetic moments are opposite but unequal, so that cancellation is incomplete and there is a non-zero net magnetization. An example is magnetite (Fe_3O_4), which contains Fe^{2+} and Fe^{3+} ions with different magnetic moments.

Notes and references

1. ^ It is often stated that more than 99% of the material in the visible universe is plasma. See, for example, D. A. Gurnett, A. Bhattacharjee (2005). *Introduction to Plasma Physics: With Space and Laboratory Applications*. Cambridge, UK: Cambridge University Press. p. 2. ISBN 0-521-36483-3. and K Scherer, H Fichtner, B Heber (2005).*Space Weather: The Physics Behind a Slogan*. Berlin: Springer. p. 138. ISBN 3-540-22907-8.. Essentially, all of the visible light from space comes from stars, which are plasmas with a temperature such that they radiate strongly at visible wavelengths. Most of the ordinary (or baryonic) matter in the universe, however, is found in the intergalactic medium, which is also a plasma, but much hotter, so that it radiates primarily as X-rays. The current scientific consensus is that about 96% of the total energy density in the universe is not plasma or any other form of ordinary matter, but a combination of cold dark matter and dark energy.
2. ^ M.A. Wahab (2005). *Solid State Physics: Structure and Properties of Materials*. Alpha Science. pp. 1–3. ISBN 1-84265-218-4.
3. ^ F. White (2003). *Fluid Mechanics*. McGraw-Hill. p. 4. ISBN 0-07-240217-2.
4. ^ G. Turrell (1997). *Gas Dynamics: Theory and Applications*. John Wiley & Sons. pp. 3–5. ISBN 0-471-97573-7.
5. ^ M. Chaplin (20 August 2009). "Water phase Diagram".*Water Structure and Science*. Retrieved 23 February 2010.
6. ^ D.L. Goodstein (1985). *States of Matter*. Dover Phoenix. ISBN 978-0-486-49506-4.
7. ^ A.P. Sutton (1993). *Electronic Structure of Materials*. Oxford Science Publications. pp. 10–12. ISBN 978-0-19-851754-2.
8. ^ Shao, Y.; Zerda, T. W. (1998). "Phase Transitions of Liquid Crystal PAA in Confined Geometries". *Journal of Physical Chemistry B* **102** (18): 3387–3394. doi:10.1021/jp9734437.

9.

External links

- 2005-06-22, MIT News: MIT physicists create new form of matter Citat: "… They have become the first to create a new type of matter, a gas of atoms that shows high-temperature superfluidity."
- 2003-10-10, Science Daily: Metallic Phase For Bosons Implies New State Of Matter
- 2004-01-15, ScienceDaily: Probable Discovery Of A New, Supersolid, Phase Of Matter Citat: "…We apparently have observed, for the first time, a solid material with the characteristics of a superfluid…but because all its particles are in the identical quantum state, it remains a solid even though its component particles are continually flowing…"
- 2004-01-29, ScienceDaily: NIST/University Of Colorado Scientists Create New Form Of Matter: A Fermionic Condensate
- Short videos demonstrating of States of Matter, solids, liquids and gases by Prof. J M Murrell, University of Sussex

---

This page titled 2.3: States of Matter is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.4: Mineral Identification

Lesson Objectives

- Explain how minerals are identified.
- Describe how color, luster, and streak are used to identify minerals.
- Explain how the hardness of a mineral is measured.
- Identify additional properties that can be used to identify some minerals.

Vocabulary

- cleavage
- fracture
- hardness
- luster
- mineralogist
- streak

Introduction

Minerals can be identified by their physical characteristics. The physical properties of minerals are related to their chemical composition and bonding. Some characteristics, such as a mineral's hardness, are more useful for mineral identification. Color is readily observable and certainly obvious, but it is usually less reliable than other physical properties.

How are Minerals Identified?

Mineralogists are scientists who study minerals. One of the things mineralogists must do is identify and categorize minerals. While a mineralogist might use a high-powered microscope to identify some minerals, most are recognizable using physical properties.

Check out the mineral in **Figure** below. What is the mineral's color? What is its shape? Are the individual crystals shiny or dull? Are there lines (striations) running across the minerals? In this lesson, the properties used to identify minerals are described in more detail.



This mineral has shiny, gold, cubic crystals with striations, so it is pyrite.

Color, Streak, and Luster

Diamonds are popular gemstones because the way they reflect light makes them very sparkly. Turquoise is prized for its striking greenish-blue color. Notice that specific terms are being used to describe the appearance of minerals.

Color

Color is rarely very useful for identifying a mineral. Different minerals may be the same color. Real gold, as seen in **Figure** below, is very similar in color to the pyrite in **Figure** above.



This mineral is shiny, very soft, heavy, and gold in color, and is actually gold.

The same mineral may also be found in different colors. **Figure** below shows one sample of quartz that is colorless and another quartz that is purple. A tiny amount of iron makes the quartz purple. Many minerals are colored by chemical impurities.



Purple quartz, known as amethyst, and clear quartz are the same mineral despite the different colors.

Streak

Streak is the color of a mineral's powder. Streak is a more reliable property than color because streak does not vary. Minerals that are the same color may have a different colored streak. Many minerals, such as the quartz in the **Figure** above, do not have streak.

To check streak, scrape the mineral across an unglazed porcelain plate (**Figure** below). Yellow-gold pyrite has a blackish streak, another indicator that pyrite is not gold, which has a golden yellow streak.



The streak of hematite across an unglazed porcelain plate is red-brown.

Luster

Luster describes the reflection of light off a mineral's surface. Mineralogists have special terms to describe luster. One simple way to classify luster is based on whether the mineral is metallic or non-metallic. Minerals that are opaque and shiny, such as pyrite, have a metallic luster. Minerals such as quartz have a non-metallic luster. Different types of non-metallic luster are described in **Table** below.

Six types of non-metallic luster.

Luster	Appearance
Adamantine	Sparkly
Earthy	Dull, clay-like
Pearly	Pearl-like
Resinous	Like resins, such as tree sap
Silky	Soft-looking with long fibers
Vitreous	Glassy

Can you match the minerals in **Figure** below with the correct luster from **Table** above?



(a) Diamond has an adamantine luster. (b) Quartz is not sparkly and has a vitreous, or glassy, luster. (b) Sulfur reflects less light than quartz, so it has a resinous luster.

Hardness

Hardness is a measure of whether a mineral will scratch or be scratched. Mohs Hardness Scale, shown in **Table** below, is a reference for mineral hardness.

Mohs Hardness Scale: 1 (softest) to 10 (hardest).


Hardness	Mineral
1	Talc
2	Gypsum
3	Calcite
4	Fluorite
5	Apatite
6	Feldspar
7	Quartz
8	Topaz
9	Corundum
10	Diamond

With a Mohs scale, anyone can test an unknown mineral for its hardness. Imagine you have an unknown mineral. You find that it can scratch fluorite or even apatite, but feldspar scratches it. You know then that the mineral's hardness is between 5 and 6. Note that no other mineral can scratch diamond.

Cleavage and Fracture

Breaking a mineral breaks its chemical bonds. Since some bonds are weaker than other bonds, each type of mineral is likely to break where the bonds between the atoms are weaker. For that reason, minerals break apart in characteristic ways.

Cleavage is the tendency of a mineral to break along certain planes to make smooth surfaces. Halite breaks between layers of sodium and chlorine to form cubes with smooth surfaces (**Figure** below).



A close-up view of sodium chloride in a water bubble aboard the International Space Station.

Mica has cleavage in one direction and forms sheets (**Figure** below).



Sheets of mica.

Minerals can cleave into polygons. Fluorite forms octahedrons (**Figure** below).




This rough diamond shows its octahedral cleavage.

One reason gemstones are beautiful is that the cleavage planes make an attractive crystal shape with smooth faces.

Fracture is a break in a mineral that is not along a cleavage plane. Fracture is not always the same in the same mineral because fracture is not determined by the structure of the mineral.

Minerals may have characteristic fractures (**Figure** below). Metals usually fracture into jagged edges. If a mineral splinters like wood, it may be fibrous. Some minerals, such as quartz, form smooth curved surfaces when they fracture.



Chrysotile has splintery fracture.

Other Identifying Characteristics

Some minerals have other unique properties, some of which are listed in **Table** below. Can you name a unique property that would allow you to instantly identify a mineral that's been described quite a bit in this chapter? (Hint: It is most likely found on your dinner table.)

Some minerals have unusual properties that can be used for identification.

Property	Description	Example of Mineral
Fluorescence	Mineral glows under ultraviolet light	Fluorite
Magnetism	Mineral is attracted to a magnet	Magnetite
Radioactivity	Mineral gives off radiation that can be measured with Geiger counter	Uraninite
Reactivity	Bubbles form when mineral is exposed to a weak acid	Calcite
Smell	Some minerals have a distinctive smell	Sulfur (smells like rotten eggs)
Taste	Some minerals taste salty	Halite

A simple lesson on how to identify minerals is seen in this video: http://www.youtube.com/watch?v=JeFVwqBuYl4.

Lesson Summary

- Minerals have distinctive properties that can be used to help identify them.
- Color and luster describe the mineral's outer appearance. Streak is the color of the powder.
- Mohs Hardness Scale is used to compare the hardness of minerals.
- Cleavage or the characteristic way a mineral breaks depends on the crystal structure of the mineral.
- Some minerals have special properties that can be used to help identify them.

Review Questions

1. Which properties of a mineral describe the way it breaks apart?
2. A mineral looks dry and chalky. What sort of luster does it have?
3. What causes a mineral to have the properties that it has?
4. Apatite scratches the surface of an unknown mineral. Which mineral would you use next to test the mineral's hardness — fluorite or feldspar? Explain your reasoning.
5. Why is streak more reliable than color when identifying a mineral?

6. Mineral A has a density of 5 g/cm^3. Mineral B is twice as dense as Mineral A. What is the density of Mineral B?
7. Why do some minerals cleave along certain planes?

Further Reading / Supplemental Links

- Mineral Color: http://geology.csupomona.edu/alert/mineral/color.htm/
- Physical Characteristics of Minerals: http://www.galleries.com/minerals/physical.htm.
- Mineral Identification: http://geology.csupomona.edu/alert/mineral/minerals.htm.

Points to Consider

- If a mineral is magnetic, do you know for certain what mineral it is?
- Some minerals are colored because they contain chemical impurities. How did the impurities get into the mineral?
- What two properties of a mineral sample would you have to measure to calculate its density?
- How much do minerals reflect the environment in which they formed?

This page titled 2.4: Mineral Identification is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.5: Minerals and Mineral Groups

Lesson Objectives

- Describe the characteristics that all minerals share.
- Identify the groups in which minerals are classified and their characteristics.

Vocabulary

- chemical compound
- crystal
- inorganic
- mineral
- silicates

Introduction

Minerals are categorized based on their chemical composition. Owing to similarities in composition, minerals within a same group may have similar characteristics.

What is a Mineral?

Minerals are everywhere! **Figure 2.1** below shows some common household items and the minerals used to make them. The salt you sprinkle on food is the mineral halite. Silver in jewelry is also a mineral. Baseball bats and bicycle frames both contain minerals. Although glass is not a mineral, it is produced from the mineral quartz. Scientists have identified more than 4,000 minerals in Earth's crust. A few are common, but many are uncommon.



Figure 2.1: Silver and halite are minerals; the mineral quartz is used to make glass.

Geologists have a very specific definition for minerals. A material is characterized as a mineral if it meets all of the following traits. **A mineral is an inorganic, crystalline solid. A mineral is formed through natural processes and has a definite chemical composition.** Minerals can be identified by their characteristic physical properties such as crystalline structure, hardness, streak, and cleavage.

Crystalline Solid

Minerals are crystalline solids. A **crystal** is a solid in which the atoms are arranged in a regular, repeating pattern (**Figure 2.2** below). The pattern of atoms in different samples of the same mineral is the same. Is glass a mineral? Without a crystalline structure, even natural glass is not a mineral.




Figure 2.2 Sodium ions (purple balls) bond with chloride ions (green balls) to make table salt (halite). All of the grains of salt that are in a salt shaker have this crystalline structure.

Inorganic Substances

Organic substances are the carbon-based compounds made by living creatures and include proteins, carbohydrates, and oils. **Inorganic** substances have a structure that is not characteristic of living bodies. Coal is made of plant and animal remains. Is it a mineral? Coal is a classified as a sedimentary rock but is not a mineral.

Natural Processes


Minerals are made by natural processes, those that occur in or on Earth. A diamond created deep in Earth's crust is a mineral. Is a diamond created in a laboratory by placing carbon under high pressures a mineral? *No*. Do not buy a laboratory-made "diamond" for jewelry without realizing it is not technically a mineral.

Chemical Composition

Nearly all (98.5%) of Earth's crust is made up of only **eight elements** – **oxygen, silicon, aluminum, iron, calcium, sodium, potassium, and magnesium** – and these are the elements that make up most minerals.

All minerals have a specific chemical composition. The mineral silver is made up of only silver atoms and diamond is made only of carbon atoms, but most minerals are made up of **chemical compounds**. Each mineral has its own chemical formula. Halite, pictured in the **Figure 2.2** above, is NaCl (sodium chloride). Quartz is always made of two oxygen atoms bonded to a silicon atom, SiO_2. If a mineral contains any other elements in its crystal structure, it's not quartz.

A hard mineral containing covalently bonded carbon is diamond, but a softer mineral that also contains calcium and oxygen along with carbon is calcite (**Figure** below).



The structure of calcite shows the relationship of calcium (Ca), carbon (C), and oxygen (O).

Some minerals have a range of chemical composition. Olivine always has silicon and oxygen as well as iron or magnesium or both, $(Mg, Fe)_2SiO_4$.

Physical Properties

The physical properties of minerals include:

- Color: the color of the mineral.
- Streak: the color of the mineral's powder.
- Luster: the way light reflects off the mineral's surface.
- Specific gravity: how heavy the mineral is relative to the same volume of water.
- Cleavage: the mineral's tendency to break along flat surfaces.
- Fracture: the pattern in which a mineral breaks.
- Hardness: what minerals it can scratch and what minerals can scratch it.

How physical properties are used to identify minerals is described in the lesson on *Mineral Formation*.


Mineral Groups

Minerals are divided into groups based on chemical composition. Most minerals fit into one of eight mineral groups.

Silicate Minerals

The roughly 1,000 silicate minerals make up over 90% of Earth's crust. **Silicates** are by far the largest mineral group. Feldspar and quartz are the two most common silicate minerals. Both are extremely common rock-forming minerals.

The basic building block for all silicate minerals is the silica tetrahedron, which is illustrated in **Figure** below. To create the wide variety of silicate minerals, this pyramid-shaped structure is often bound to other elements, such as calcium, iron, and magnesium.



One silicon atom bonds to four oxygen atoms to form a silica tetrahedron.

Silica tetrahedrons combine together in six different ways to create different types of silicates (**Figure** below). Tetrahedrons can stand alone, form connected circles called rings, link into single and double chains, form large flat sheets of pyramids, or join in three dimensions.




The different ways that silica tetrahedrons can join together cause these two minerals to look very different.

Native Elements

Native elements contain atoms of only one type of element. Only a small number of minerals are found in this category. Some of the minerals in this group are rare and valuable. Gold, silver, sulfur, and diamond are examples of native elements.

Carbonates

The basic carbonate structure is one carbon atom bonded to three oxygen atoms. Carbonates include other elements, such as calcium, iron, and copper. Calcite ($CaCO_3$) is the most common carbonate mineral (**Figure** below).



The carbonate ion is one carbon atom bonded to three oxygen atoms.

Calcite crystals.

Calcite is the most common carbonate mineral.


Azurite and malachite, shown in the **Figure** below, are carbonates that contain copper instead of calcium.



Two carbonate minerals: (a) deep blue azurite and (b) opaque green malachite.

Halides


Halide minerals are salts that form when salt water evaporates. Halite is a halide mineral, but table salt is not the only halide. The chemical elements known as the halogens (fluorine, chlorine, bromine, or iodine) bond with various metallic atoms to make halide minerals (see**Figure** below).



Fluorite is a halide containing calcium and fluorine.

Oxides

Oxides contain one or two metal elements combined with oxygen. Many important metals are found as oxides. Hematite (Fe_2O_3), with two iron atoms to three oxygen atoms, and magnetite (Fe_3O_4) (**Figure** below), with three iron atoms to four oxygen atoms, are both iron oxides.



Magnetite is the most magnetic mineral. Magnetite attracts or repels other magnets.

Phosphates

Phosphate minerals are similar in atomic structure to the silicate minerals. In the phosphates, phosphorus, arsenic, or vanadium bond to oxygen to form a tetrahedra. There are many different minerals in the phosphate group, but most are rare (**Figure** below).



Turquoise is a phosphate mineral containing copper, aluminum, and phosphorus.

Sulfates

Sulfate minerals contain sulfur atoms bonded to oxygen atoms. Like halides, they form where salt water evaporates. The sulfate group contains many different minerals, but only a few are common.

Gypsum is a common sulfate with a variety of appearances (**Figure** below). Some gigantic 11-meter gypsum crystals have been found. That is about as long as a school bus!



Although the orange crystals on the left looks nothing like the white sands on the right, both the crystals and sands are gypsum.

Sulfides

Sulfides are formed when metallic elements combine with sulfur. Unlike sulfates, sulfides do not contain oxygen. Pyrite, or iron sulfide, is a common sulfide mineral known as *fool's gold*. People may mistake pyrite for gold because the two minerals are shiny, metallic, and yellow in color.

Lesson Summary

- For a substance to be a mineral, it must be a naturally occurring, inorganic, crystalline solid that has a characteristic chemical composition and crystal structure.
- The atoms in minerals are arranged in regular, repeating patterns that can be used to identify that mineral.
- Minerals are divided into groups based on their chemical composition.
- The chemical feature of each groups is: native elements – only one element; silicates – silica tetrahedron; phosphates – phosphate tetrahedron; carbonates – one carbon atom with three oxygen atoms; halides – a halogen bonded with a metallic atom; oxides – a metal combined with oxygen; sulfates – sulfur and oxygen; sulfides – metal with sulfur, no oxygen.

Review Questions

1. What is a crystal?
2. Which elements do all silicate minerals contain?
3. Obsidian is a glass that forms when lava cools so quickly that the atoms do not have a chance to arrange themselves in crystals. Is obsidian a crystal? Explain your reasoning.
4. What are the eight major mineral groups?
5. What is the same about all minerals in the silicate group? What is different about them?
6. One sample has a chemical composition with a ratio of two iron atoms to three oxygen atoms. Another sample has a chemical composition with a ratio of three iron atoms to four oxygen atoms. They contain the same elements: Are they the same mineral?
7. How does the native elements mineral group differ from all of the other mineral groups?
8. On a trip to the natural history museum you find two minerals that are similar in color. You can see from their chemical formulas that one mineral contains the elements zinc, carbon, and oxygen. The other mineral contains the elements zinc, silicon, oxygen, and hydrogen. Your friend tells you that the minerals are in the same mineral group. Do you agree? Explain your reasoning.

Further Reading / Supplemental Links

- The Definition of a Mineral: http://www.minsocam.org/msa/ima/ima98(04).pdf.
- Mineral Identification: http://geology.csupomona.edu/alert/mineral/minerals.htm.
- Dana Classification of Minerals: http://webmineral.com/danaclass.shtml.
- A Lot of Different Minerals: http://hyperphysics.phy-astr.gsu.edu/hbase/geophys/mineral.html#c1.
- Mineral Groups: http://mineral.galleries.com/minerals/silicate/class.htm.
- Giant Crystal Cave, National Geographic: http://news.nationalgeographic.com/news/2007/04/photogalleries/giant-crystals-cave/photo3.html.

Points to Consider

- Why is obsidian, a natural glass that forms from cooling lava, not a mineral?
- Why are diamonds made in a laboratory not minerals?
- Is coal, formed mostly from decayed plants, a mineral? Is it a rock?
- Artists used to grind up the mineral azurite to make colorful pigments for paints. Is the powdered azurite still crystalline?

This page titled 2.5: Minerals and Mineral Groups is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.6: Mineral Formation

Lesson Objectives

- Describe how melted rock produces minerals.
- Describe how hot rock produces different minerals.
- Explain how minerals form from solutions.

Vocabulary


- lava
- magma
- rock
- vein

Introduction

Minerals form under an enormous range of geologic conditions. There are probably more ways to form minerals than there are types of minerals themselves. Minerals can form from volcanic gases, sediment formation, oxidation, crystallization from magma, or deposition from a saline fluid, to list a few. Some of these methods of mineral formation will be discussed below.

Formation from Hot Material

A **rock** is a collection of minerals. Imagine a rock that becomes so hot it melts. Many minerals start out in liquids that are hot enough to melt rocks. **Magma** is melted rock inside Earth, a molten mixture of substances that can be hotter than 1,000°C. Magma cools slowly inside Earth, which gives mineral crystals time to grow large enough to be seen clearly (**Figure** below).



Granite is rock that forms from slowly cooled magma, containing the minerals quartz (clear), plagioclase feldspar (shiny white), potassium feldspar (pink), and biotite (black).

When magma erupts onto Earth's surface, it is called **lava**. Lava cools much more rapidly than magma when it is below the surface. In a cooling lava, mineral crystals do not have time to form and are very small. The chemical composition will be the same as if the magma cooled slowly.


Existing rocks may be heated enough so that the molecules are released from their structure and can move around. The molecules may match up with different molecules to form new minerals as the rock cools. This occurs during metamorphism, which will be discussed in the chapter "Rocks."

Formation from Solutions

Water on Earth, such as the water in the oceans, contains chemical elements mixed into a solution. Various processes can cause these elements to combine to form solid mineral deposits.

Minerals from Salt Water

When water evaporates, it leaves behind a solid precipitate of minerals, as shown in **Figure** below.



When the water in glass A evaporates, the dissolved mineral particles are left behind.

Water can only hold a certain amount of dissolved minerals and salts. When the amount is too great to stay dissolved in the water, the particles come together to form mineral solids, which sink. Halite easily precipitates out of water, as does calcite. Some lakes, such as Mono Lake in California (**Figure** below) or The Great Salt Lake in Utah, contain many mineral precipitates.



Tufa towers form when calcium-rich spring water at the bottom of Mono Lake bubbles up into the alkaline lake. The tufa towers appear when lake level drops.

Minerals from Hot Underground Water

Magma heats nearby underground water, which reacts with the rocks around it to pick up dissolved particles. As the water flows through open spaces in the rock and cools, it deposits solid minerals. The mineral deposits that form when a mineral fills cracks in rocks are called**veins** (**Figure** below).



Quartz veins formed in this rock.

When minerals are deposited in open spaces, large crystals form (**Figure** below).



Amethyst formed when large crystals grew in open spaces inside the rock. These special rocks are called geodes.

Lesson Summary

- Mineral crystals that form when magma cools slowly are larger than crystals that form when lava cools rapidly.
- Minerals form when rocks are heated enough that atoms of different elements can move around and join into different molecules.
- Minerals are deposited from salty water solutions on Earth's surface and underground.

Review Questions

1. What is the difference between magma and lava?
2. Under what circumstances do large crystals form from a cooling magma?
3. Under what circumstances do small crystals form from a cooling magma?
4. What happens to the mineral particles in salt water when the water evaporates?
5. Explain how mineral veins form.

Further Reading / Supplemental Links

- Gems and Where They're Found: http://socrates.berkeley.edu/~eps2/wisc/Lect3.html.
- How to Grow Your Own Crystals: http://www.sdnhm.org/kids/minerals/grow-crystal.html.

Points to Consider

- Is a mineral a static thing or does it change? If it changes, on what time frame?
- When most minerals form, they combine with other minerals to form rocks. How can these minerals be used?
- The same mineral can be formed by different processes. How can the way a mineral forms affect how the mineral is used?

This page titled 2.6: Mineral Formation is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

2.7: Mining and Mineral Use

Lesson Objectives

- Explain how minerals are mined.
- Describe how metals are made from mineral ores.
- Summarize the ways in which gemstones are used.
- Identify some useful minerals.

Vocabulary

- gemstone
- ore
- ore deposit
- placer
- reclamation

Introduction

Some minerals are very useful. An **ore** is a rock that contains minerals with useful elements. Aluminum in bauxite ore (**Figure** below) is extracted from the ground and refined to be used in aluminum foil and many other products. The cost of creating a product from a mineral depends on how abundant the mineral is and how much the extraction and refining processes cost. Environmental damage from these processes is often not figured into a product's cost. It is important to use mineral resources wisely.



Aluminum is made from the aluminum-bearing minerals in bauxite.

Finding and Mining Minerals

Geologic processes create and concentrate minerals that are valuable natural resources. Geologists study geological formations and then test the physical and chemical properties of soil and rocks to locate possible ores and determine their size and concentration.

A mineral deposit will only be mined if it is profitable. A concentration of minerals is only called an **ore deposit** if it is profitable to mine. There are many ways to mine ores.

Surface Mining

Surface mining allows extraction of ores that are close to Earth's surface. Overlying rock is blasted and the rock that contains the valuable minerals is placed in a truck and taken to a refinery. As pictured in **Figure** below, surface mining includes open-pit mining and mountaintop removal. Other methods of surface mining include strip mining, placer mining, and dredging. Strip mining is like open pit mining but with material removed along a strip.



These different forms of surface mining are methods of extracting ores close to Earth's surface.

Placers are valuable minerals found in stream gravels. California's nickname, the Golden State, can be traced back to the discovery of placer deposits of gold in 1848. The gold weathered out of hard metamorphic rock in the western Sierra Nevada, which also contains deposits of copper, lead, zinc, silver, chromite, and other valuable minerals. The gold traveled down rivers and then settled in gravel deposits. Currently, California has active mines for gold and silver and for non-metal minerals such as sand and gravel, which are used for construction.

Underground Mining

Underground mining is used to recover ores that are deeper into Earth's surface. Miners blast and tunnel into rock to gain access to the ores. How underground mining is approached – from above, below, or sideways – depends on the placement of the ore body, its depth, concentration of ore, and the strength of the surrounding rock.

Underground mining is very expensive and dangerous. Fresh air and lights must also be brought into the tunnels for the miners, and accidents are far too common.

Ore Extraction

The ore's journey to becoming a useable material is only just beginning when the ore leaves the mine (**Figure** below). Rocks are crushed so that the valuable minerals can be separated from the waste rock. Then the minerals are separated out of the ore. A few methods for extracting ore are:

- heap leaching: the addition of chemicals, such as cyanide or acid, to remove ore.
- flotation: the addition of a compound that attaches to the valuable mineral and floats.
- smelting: roasting rock, causing it to segregate into layers so the mineral can be extracted.



The de Young Museum in San Francisco is covered in copper panels. Copper is mined and extracted from copper ores.

To extract the metal from the ore, the rock is melted at a temperature greater than 900°C, which requires a lot of energy. Extracting metal from rock is so energy intensive that if you recycle just 40 aluminum cans, you will save the energy equivalent of one gallon of gasoline.

Mining and the Environment

Although mining provides people with many needed resources, the environmental costs can be high. Surface mining clears the landscape of trees and soil, and nearby streams and lakes are inundated with sediment. Pollutants from the mined rock, such as heavy metals, enter the sediment and water system. Acids flow from some mine sites, changing the composition of nearby waterways (**Figure** below).



Acid drainage from a surface coal mine in Missouri.

U.S. law has changed so that in recent decades a mine region must be restored to its natural state, a process called **reclamation**. This is not true of older mines. Pits may be refilled or reshaped and vegetation planted. Pits may be allowed to fill with water and become lakes or may be turned into landfills. Underground mines may be sealed off or left open as homes for bats.

Valuable Minerals

Some minerals are valuable because they are beautiful. Jade has been used for thousands of years in China. Diamonds sparkle on many engagement rings. Minerals like jade, turquoise, diamonds, and emeralds are gemstones. A **gemstone**, or gem, is a material that is cut and polished for jewelry. Many gemstones, including many in **Figure** below, are minerals.



Gemstones come in many colors.

Gemstones are usually rare and do not break or scratch easily. Most are cut along cleavage faces and then polished so that light bounces back off the cleavage planes (**Figure** below). Light does not pass through gemstones that are opaque, such as turquoise.



Uncut (left) and cut (right) ruby. The cut and polished ruby sparkles.

Gemstones are not just used in jewelry. Diamonds are used to cut and polish other materials, such as glass and metals, because they are so hard. The mineral corundum, of which ruby and sapphire are varieties, is used in products such as sandpaper.

Minerals are used in much less obvious places. The mineral gypsum is used for the sheetrock in homes. Window glass is made from sand, which is mostly quartz. Halite is mined for rock salt. Copper is used in electrical wiring, and bauxite is the source for the aluminum used in soda cans.

Lesson Summary

- Geologists use many methods to find mineral deposits that will be profitable to mine.
- Ore deposits can be mined by surface or underground mining methods.
- Mining provides important resources but has environmental costs.
- By U.S. law, currently mined land must undergo reclamation. This is not true for old mines.
- Metal ores must be melted to make metals.
- Many gems are cut and polished to increase their beauty.
- Minerals are used in a variety of ways.

Review Questions

1. What category of mining would be used to extract ore that is close to the surface? Why?
2. Describe some surface mining methods.
3. What are some disadvantages of underground mining?

4. What are some ways an area can undergo reclamation after being mined?
5. What steps are taken to extract a pure metal from an ore?
6. What makes a gemstone valuable?

Further Reading / Supplemental Links

- Aluminum and Bauxite: http://www.mii.org/Minerals/photoal.html.
- Mining Diamonds: http://www.amnh.org/exhibitions/diamonds/mining.html.
- Gems: http://www.amnh.org/exhibitions/diamonds/mining.html and http://socrates.berkeley.edu/~eps2/wisc/Lect2.html.

Points to Consider

- Are all mineral deposits ores?
- Why might an open pit mine be turned into an underground mine?
- How well does reclaimed land resemble the land before mining began?
- Diamonds are not necessarily the rarest gem so why do people value them more than most other gems?
- Under what circumstances might a mineral deposit be an ore one day and not the next?

This page titled 2.7: Mining and Mineral Use is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

3: Rocks

3.1: Igneous Rocks

3.2: Sedimentary Rocks

3.3: Metamorphic Rocks

This page titled 3: Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

3.1: Igneous Rocks

This lesson will discuss igneous rocks, how they form, how they are classified, and some of their common uses. Igneous rocks may or may not be found naturally where you live, but chances are that you have seen materials made from igneous rocks. One of the most common igneous rocks is granite (Figure 4.1). Granite is used extensively in building materials and making statues. Perhaps you have used a pumice stone to smooth your skin or to do jobs around the house. Pumice is another example of an igneous rock (Figure 4.2). Pumice is used to make stone-washed denim jeans! Pumice stones are put into giant washing machines with newly-manufactured jeans and tumbled around to give jeans that distinctive “stonewashed” look. You also probably use igneous rock when you brush your teeth every morning. Ground up pumice stone is sometimes added to toothpaste to act as an abrasive material that scrubs your teeth clean.



Figure 4.1: Granite is an igneous rock used commonly in statues and building materials.



Figure 4.2: Pumice is a light igneous rock used for abrasive materials.

Lesson Objectives

- Describe how igneous rocks are formed.
- Describe the properties of some common types of igneous rocks.
- Relate some common uses of igneous rocks.

Crystallization

Igneous rocks form when molten material cools and hardens. They may form either below or above Earth’s surface. They make up most of the rocks on Earth. Most igneous rock is buried below the surface and covered with sedimentary rock, and so we do not often see just how much igneous rock there is on Earth. In some places, however, large areas of igneous rocks can be seen at Earth’s surface. Figure 4.3 shows a landscape in California’s Sierra Nevada that consists entirely of granite, an igneous rock.



Figure 4.3: This landscape high in California's Sierra Nevada is completely made up of granite exposed at Earth's surface.

Igneous rocks are called intrusive or **plutonic** when they **cool and solidify beneath the surface**. Because they *form within* the Earth, *cooling occurs slowly*. Such slow cooling allows time for large crystals to form, therefore, intrusive or **plutonic** igneous rocks have relatively large mineral crystals that are easy to see. Granite is the most common intrusive igneous rock (Figure 4.4).



Figure 4.4: A close-up of a granite sample. Notice the black and white portions. Each color represents a different mineral in the rock. You can easily see the mineral crystals that make up this plutonic igneous rock.

Igneous rocks are called extrusive or **volcanic** when they **form above the surface**. They solidify after molten material pours out onto the surface through an opening such as a volcano (Figure 4.5). Extrusive or **volcanic** igneous rocks *cool much more rapidly* and therefore have smaller crystals. Since the rapid cooling time does not allow time for large crystals to form, minerals are not easy to see within the rock. Some volcanic igneous rocks cool so rapidly that crystals do not develop at all. These form a glass, such as **obsidian.** Others, such as pumice, contain holes where gas bubbles were trapped when the material was still hot and molten. The holes make pumice so light that it actually floats in water. The most common extrusive igneous rock is basalt, a rock that is especially common below the oceans (Figure 4.6).



Figure 4.5: Extrusive or volcanic igneous rocks form after lava cools above the surface.



Figure 4.6: These are examples of basalt below the South Pacific Ocean. Basalt is an extrusive igneous rock common below the oceans, though it is also found in some places on continents.

Composition

Igneous rocks are classified according to how and where they formed (in other words, if they're plutonic or volcanic) and their mineral composition (describing the minerals they contain). The mineral compositions of igneous rocks are usually described as being **felsic, intermediate, mafic, or ultramafic** (as examples, see Figure 4.7 and Figure 4.8). **Felsic** rocks are made of light-colored, low-density minerals such as quartz and feldspar. **Mafic** rocks are made of dark-colored, higher-density minerals such as olivine and pyroxene.**Intermediate** rocks have compositions between felsic and mafic. **Ultramafic** rocks contain more than 90% mafic minerals and have very few light, felsic minerals in them. Table 4.1 shows some common igneous rocks classified by mode of occurrence and mineral composition.

Table 4.1: Common Igneous Rocks

Mode of Occurrence	Mineral Composition			
	Felsic	**Intermediate**	**Mafic**	**Ultramafic**
Volcanic/Extrusive	Rhyolite	Andesite	Basalt	Komatiite
Plutonic/Intrusive	Granite	Diorite	Gabbro	Peridotite

The rocks listed in the table above are the most common igneous rocks, but there are actually more than 700 different types of igneous rocks. Granite is perhaps the most useful one to humans. We use granite in many building materials and in art. As discussed in the introduction to this lesson, pumice is commonly used for abrasives. Peridotite is sometimes mined for peridot, a type of gemstone used in jewelry. Diorite is extremely hard and is commonly used for art. It was used extensively by ancient civilizations for vases and other decorative art work



Figure 4.7: Rhyolite is an example of an extrusive or volcanic felsic rock. Notice its light color and small crystals.




Figure 4.8: This is a close-up photograph of peridotite, an extrusive or volcanic ultramafic igneous rock. The green mineral is olivine.

Lesson Summary

- Igneous rocks form either when they cool very slowly deep within the Earth or when magma cools rapidly at the Earth's surface.
- Composition of the magma will determine the minerals that will crystallize forming different types of igneous rocks.

Review Questions

1. What is the difference between a plutonic and a volcanic igneous rock?
2. Why do volcanic igneous rocks usually have smaller crystals than plutonic igneous rocks?
3. How are igneous rocks classified?
4. Describe two ways granite is different from basalt.
5. Occasionally, igneous rocks will contain both large crystals and tiny mineral crystals. Propose a way that both these size crystals might have formed in the rock.

Vocabulary

extrusive or volcanic

A type of igneous rock that forms above Earth's surface.

felsic

A type of igneous rock composition that is made mostly of light minerals such as quartz and feldspar.

intermediate

A type of igneous rock composition that is in between felsic and mafic.

intrusive or plutonic

A type of igneous rock that forms inside the Earth.

mafic

A type of igneous rock composition that is made mostly of dense, dark minerals like olivine and pyroxene.

solidify

To harden after cooling.

ultramafic

A type of igneous rock that contains more than 90% mafic minerals.

Points to Consider

- Do you think igneous rocks could form where you live?
- Would all igneous rocks with the same composition have the same name? Explain why they might not.
- Could an igneous rock cool at two different rates? What would the crystals in such a rock look like?

Lumen Learning authored content

- **License**: *CC BY: Attribution*

This page titled 3.1: Igneous Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

3.2: Sedimentary Rocks

You probably recognize the Figure 4.10 as the White House, the official home and workplace of the President of the United States of America. Do you know why the White House is white? Its color has a lot to do with the stone materials that were used to construct it.

Construction for the White House began in 1792, and most of the work was carried out by people who had only recently come to the newly formed country of America. Its outside walls are made of a type of sedimentary rock called sandstone. The sandstone that was used to construct the White House is very porous, which means that rainwater can easily penetrate the sandstone. This made the White House susceptible to water damage in its early days of construction. To stop the water damage, workers had to cover the sandstone in a mixture of salt, rice, and glue, giving the White House its distinct white color.



Figure 4.10 The White House of the United States of America is made of a sedimentary rock called sandstone.

Lesson Objectives

- Describe how sedimentary rocks are formed.
- Describe the properties of some common sedimentary rocks.
- Relate some common uses of sedimentary rocks.

Sediments

In this lesson, you will learn about sedimentary rocks like sandstone, how they form, how they are classified, and how people often use sedimentary rocks.

Sedimentary rocks are formed by the compaction of sediments. **Sediments** may include:

- fragments of other rocks that have been worn down into small pieces, like sand,
- **organic** materials, or in other words, the remains of once-living organisms,
- or chemical **precipitates**, which are materials that get left behind after the water evaporates from a solution.

Most sediments settle out of water (Figure 4.11). For example, running water in rivers carries huge amounts of sediments. The river dumps these sediments along its banks and at the end of its course. When sediments settle out of water, they form horizontal layers. One layer at a time is put down. Each new layer forms on top of the layers that were already there. Thus, each layer in a sedimentary rock is younger than the layer under it and older than the layer over it. When the sediments harden, the layers are preserved. In large **outcrops** of sedimentary rocks, you can often see layers that show the position and order in which the original sediment layers were deposited. Scientists can figure out the relative ages of layers by knowing that older ones are on the bottom and younger ones are on top.



Figure 4.11: Most sediments settle out of running water, such as in this river.

There are many different types of environments where sedimentary rocks form. Some places where you can see large deposits of sediments today include a beach and a desert. Sediments are also continuously depositing at the bottom of the ocean and in lakes, ponds, rivers, marshes and swamps. Avalanches produce large unsorted piles of sediment. The environment where the sediments are deposited determines the type of sedimentary rock that will form there.


Sedimentary Rock Formation

Sediments accumulate and over time may be hardened into rock. **Lithification** is the hardening of layers of loose sediment into rock (Figure 4.12). Lithification is made up of two processes: cementation and compaction. **Cementation** occurs when substances crystallize or fill in the spaces between the loose particles of sediment. These cementing substances come from the water that moves through the sediments. Sediments may also be hardened into rocks through **compaction**. This occurs when sediments are squeezed together by the weight of layers on top of them. Sedimentary rocks made of cemented, non-organic sediments are called clastic rocks. Those that form from organic remains are called bioclastic rocks, and sedimentary rocks formed by the hardening of chemical precipitates are called chemical sedimentary rocks. Table 4.2 shows some common types of sedimentary rocks and the types of sediments that make them up.



Figure 4.12: This cliff is made of a sedimentary rock called sandstone. The bands of white and red represent different layers of sediment. The layers of sediments were preserved during lithification.

Table 4.2: Common Sedimentary Rocks

Picture	Rock Name	Type of Sedimentary Rock
	Conglomerate	Clastic (sediments are rounded in shape)
	Breccia	Clastic (sediments are angular in shape)
	Sandstone	Clastic
	Siltstone	Clastic
	Shale	Clastic
	Rock Salt	Chemical precipitate
	Rock Gypsum	Chemical precipitate

	Dolostone	Chemical precipitate
	Limestone	Bioclastic (sediments from organic materials, or plant or animal remains)
	Coal	Organic

Note from the pictures in the table that clastic sedimentary rocks vary in the size of their sediments. Both conglomerate and breccia are made of individual stones that have been cemented together. In conglomerate, the stones are rounded; in breccia, the stones are angular around the edges. Sandstone is made of smaller, mostly sand-sized particles cemented together. Siltstone is made mostly of silt, particles that are smaller than sand but larger than clay. Shales have the smallest grain size, being made mostly of clay-sized particles and hardened mud.

Lesson Summary

- Weathering and erosion produce sediments. Once these sediments are deposited, they can become sedimentary rocks.
- Sediments must be compacted and cemented to make sedimentary rock. This process is called lithification.

Review Questions

1. What are three things that the sediments in sedimentary rocks may be made of?
2. If you see a sedimentary rock outcrop and red layers of sand are on top of pale layers of sand, what do you know for sure about the ages of the two layers?
3. Why do sedimentary rocks have layers of different colors sometimes?
4. Describe the two processes necessary for sediments to harden into rock.
5. What type of sedimentary rock is coal?
6. Think back to the story at the start of the lesson about why the White House originally was white. Why do you think sandstone would be a particularly porous rock?

Vocabulary

cementation

Occurs when substances harden crystallize in the spaces between loose sediments.

lithification

The hardening of layers of loose sediment into rock

organic

Made from materials that were once living things.

outcrop

Large rock formation at the surface of the Earth.

precipitate

The solid materials left behind after a liquid evaporates.

compaction

Occurs when sediments are hardened by being squeezed together by the weight of layers on top of them.

CC licensed content, Shared previously

- Earth Science for High School. **Provided by**: CK-12. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/?noindex=None. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 3.2: Sedimentary Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

3.3: Metamorphic Rocks

In this lesson you will learn about metamorphic rocks, how they form, and some of their common uses. Figure 4.13 shows a large outcrop of metamorphic rocks. Notice the platy layers that run from left to right within the rock. It looks as though you could easily break off layers from the front surface of the outcrop. This layering is a result of the process of metamorphism. Metamorphism is the changing of rocks by heat and pressure. During this process, rocks change either physically and/or chemically. They change so much that they become an entirely new rock.



Figure 4.13: The platy layers in this large outcrop of metamorphic rock show the effects of pressure on rocks during metamorphism.

Lesson Objectives

- Describe how metamorphic rocks are formed.
- Describe the properties of some common metamorphic rocks.
- Relate some common uses of metamorphic rocks.

Metamorphism

Metamorphic rocks start off as igneous, sedimentary, or other metamorphic rocks. These rocks are changed when heat or pressure alters the existing rock's physical or chemical make up. One ways rocks may change during metamorphism is by rearrangement of their mineral crystals. When heat and pressure change the environment of a rock, the crystals may respond by rearranging their structure. They will form new minerals that are more **stable** in the new environment. Extreme pressure may also lead to the formation of **foliation**, or flat layers in rocks that form as the rocks are squeezed by pressure. Foliation normally forms when pressure was exerted on a rock from one direction. If pressure is exerted from all directions, then the rock usually does not show foliation.

There are two main types of metamorphism:

1. Contact metamorphism—occurs when magma contacts a rock, changing it by extreme heat (Figure 4.14).
2. Regional metamorphism—occurs when great masses of rock change over a wide area due to pressure exerted on rocks at plate boundaries.




Figure 4.14: This diagram shows hot magma within the earth contacting various rock layers. This is an example of contact metamorphism.

It is important to note that metamorphism does not cause complete melting of the initial rock. It only causes changes to a rock by heat or pressure. The rearrangement of the mineral crystals is the most common way that we notice these changes. Table 4.3 shows some common metamorphic rocks and the original rocks that they come from.

Table 4.3: Common Metamorphic Rocks

Picture	Rock Name	Type	Comments
	Slate	Foliated	Metamorphism of shale
	Phyllite	Foliated	Metamorphism of slate, but under greater heat and pressure than slate
	Schist	Foliated	Often derived from metamorphism of claystone or shale; metamorphosed under more heat and pressure than phyllite
	Gneiss	Foliated	Metamorphism of various different rocks, under extreme conditions of heat and pressure
	Hornfels	Non-foliated	Contact metamorphism of various different rock types
	Quartzite	Non-foliated	Metamorphism of sandstone
	Marble	Non-foliated	Metamorphism of limestone



Figure 4.15: Marble is used for decorative items and in art.

Hornfels, with its alternating bands of dark and light crystals is a good example of how minerals rearrange themselves during metamorphism. In this case, the minerals separated by density and became banded. Gneiss forms by regional metamorphism from both high temperature and pressure.

Quartzite and marble are the most commonly used metamorphic rocks. They are frequently chosen for building materials and artwork. Marble is used for statues and decorative items like vases (Figure 4.15). Ground up marble is also a component of toothpaste, plastics, and paper. Quartzite is very hard and is often crushed and used in building railroad tracks (Figure 4.16). Schist and slate are sometimes used as building and landscape materials.



Figure 4.16: Crushed quartzite is sometimes placed under railroad tracks because it is very hard and durable.

Lesson Summary

- Metamorphic rocks form when heat and pressure transform an existing rock into a new rock.
- Contact metamorphism occurs when hot magma transforms rock that it contacts.
- Regional metamorphism transforms large areas of existing rocks under the tremendous heat and pressure created by tectonic forces.

Review Questions

1. Why do the minerals in a rock sometimes rearrange themselves when exposed to heat or pressure?
2. What is foliation in metamorphic rocks?
3. Describe the different conditions that lead to foliated versus non-foliated metamorphic rocks.
4. List and describe the two main types of metamorphism.
5. How can metamorphic rocks be a clue to how they were formed?
6. Suppose a phyllite sample was metamorphosed again. How might it look different after this second round of metamorphism.

Vocabulary

contact metamorphism

Results from temperature increases when a body of magma contacts a cooler existing rock.

foliation

Property of some metamorphic rocks in which flat layers are formed; seen as evidence of squeezing by pressure.

regional metamorphism

Occurs when great masses of rock change over a wide area due to pressure.

stable

Steady and not likely to change significantly any more.

Points to Consider

- What type of plate boundary would produce the most intense metamorphism of rock?
- Do you think new minerals could form when an existing rock is metamorphosed?

CC licensed content, Shared previously

- Earth Science for High School. **Provided by**: CK-12. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/?noindex=None. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 3.3: Metamorphic Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

4: Earth’s Interior Processes

4.1: Earth’s Interior

4.2: Plate Tectonics

4.3: Volcanoes

4.4: Earthquakes


This page titled 4: Earth’s Interior Processes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

4.1: Earth's Interior

Understanding Earth's Interior

Before you can learn about plate tectonics, you need to know something about the layers that are found inside Earth. These layers are divided by composition into core, mantle, and crust or by mechanical properties into lithosphere and asthenosphere. Scientists use information from earthquakes and computer modeling to learn about Earth's interior.

Humans have never drilled past Earth's crust, and yet we know a lot about the composition of the earth's interior. Rocks yield some clues, but they only reveal information about the outer crust. In rare instances, a mineral, such as a diamond, comes to the surface from deeper down in the crust or the mantle. To learn about Earth's interior, scientists use energy, recorded by seismographs, to "see" the different layers of the Earth, just like doctors can use an MRI, CT scan, or x-ray to see inside our bodies.



SEISMIC WAVES

One ingenious way scientists learn about Earth's interior is by looking at how energy travels from the point of an earthquake, called **seismic waves**. Seismic waves travel outward in all directions from where the ground breaks at an earthquake. Seismograph stations measure the energy released by these earthquakes, but there are two that scientists are most interested with in regards to understanding the interior of the earth.**Primary waves** (also called **P-waves**) are fastest, traveling at about 6 to 7 kilometers (about 4 miles) per second, so they arrive first at the seismometer. P-waves move in a compression/expansion type motion, squeezing and unsqueezing earth materials as they travel. P-waves bend slightly when they travel from one layer into another. Seismic waves move faster through denser or more rigid material. As P-waves encounter the liquid outer core, which is less rigid than the mantle, they slow down. This makes the P-waves arrive later and further away than would be expected. The result is a P-wave shadow zone. No P-waves are picked up at seismographs 104o to 140o from the earthquakes focus.

Secondary waves (also called **S-waves**) are about half as fast as P-waves, traveling at about 3.5 km (2 miles) per second, and arrive second at seismographs. S-waves move in an up and down motion perpendicular to the direction of wave travel. This produces a change in shape for the earth materials they move through. Only solids resist a change in shape, so S-waves are only able to propagate through solids. S-waves cannot travel through liquid. By tracking seismic waves, scientists have learned what makes up the planet's interior. P-waves slow down at the mantle core boundary, so we know the outer core is less rigid than the mantle. S-waves disappear at the mantle core boundary, so the outer core is liquid.


Other clues to Earth's interior includes the fact that we know that Earth's overall density is higher than the density of crustal rocks, so the core must be made of something dense, like metal. Also, since Earth has a magnetic field, there must be metal within the planet. Iron and nickel are both magnetic. Finally, meteorites are the remains of the material that formed the early solar system and are thought to be similar to material in Earth's interior.

The Composition and Structure of Earth

Core, mantle, and crust are divisions based on composition. The crust makes up less than 1 percent of Earth by mass, consisting of oceanic crust and continental crust is often more felsic rock. The mantle is hot and represents about 68 percent of Earth's mass.

Finally, the core is mostly iron metal. The core makes up about 31% of the Earth.

Lithosphere and asthenosphere are divisions based on mechanical properties. The **lithosphere** is composed of both the crust and the portion of the upper mantle that behaves as a brittle, rigid solid. The **asthenosphere** is partially molten upper mantle material that behaves plastically and can flow. This animation by Earthquide shows the layers by composition and by mechanical properties.




CRUST AND LITHOSPHERE

Earth's outer surface is its crust; a cold, thin, brittle outer shell made of rock. The crust is very thin, relative to the radius of the planet. There are two very different types of crust, each with its own distinctive physical and chemical properties.**Oceanic crust** is composed of magma that erupts on the seafloor to create basalt lava flows or cools deeper down to create the intrusive igneous rock gabbro. Sediments, primarily muds and the shells of tiny sea creatures, coat the seafloor. Sediment is thickest near the shore where it comes off the continents in rivers and on wind currents.

Continental crust is made up of many different types of igneous, metamorphic, and sedimentary rocks. The average composition is granite, which is much less dense than the mafic igneous rocks of the oceanic crust. Because it is thick and has relatively low density, continental crust rises higher on the mantle than oceanic crust, which sinks into the mantle to form basins. When filled with water, these basins form the planet's oceans.

The lithosphere is the outermost mechanical layer, which behaves as a brittle, rigid solid. The lithosphere is about 100 kilometers thick. The definition of the lithosphere is based on how earth materials behave, so it includes the crust and the uppermost mantle, which are both brittle. Since it is rigid and brittle, when stresses act on the lithosphere, it breaks. This is what we experience as an earthquake.



MANTLE

The two most important things about the mantle are: (1) it is made of solid rock, and (2) it is hot. Scientists know that the mantle is made of rock based on evidence from seismic waves, heat flow, and meteorites. The properties fit the ultramafic rock peridotite, which is made of the iron- and magnesium-rich silicate minerals. Peridotite is rarely found at Earth's surface.Scientists know that the mantle is extremely hot because of the heat flowing outward from it and because of its physical properties. Heat flows in two different ways within the Earth: conduction and convection. Conduction is defined as the heat transfer that occurs through rapid collisions of atoms, which can only happen if the material is solid. Heat flows from warmer to cooler places until all are the same temperature. The mantle is hot mostly because of heat conducted from the core. Convection is the process of a material that can move and flow may develop convection currents.

Convection in the mantle is the same as convection in a pot of water on a stove. Convection currents within Earth's mantle form as material near the core heats up. As the core heats the bottom layer of mantle material, particles move more rapidly, decreasing its density and causing it to rise. The rising material begins the convection current. When the warm material reaches the surface, it spreads horizontally. The material cools because it is no longer near the core. It eventually becomes cool and dense enough to sink back down into the mantle. At the bottom of the mantle, the material travels horizontally and is heated by the core. It reaches the location where warm mantle material rises, and the mantle convection cell is complete.

Convection in the mantle is the same as convection in a pot of water on a stove. Convection currents within Earth's mantle form as material near the core heats up. As the core heats the bottom layer of mantle material, particles move more rapidly, decreasing its density and causing it to rise. The rising material begins the convection current. When the warm material reaches the surface, it spreads horizontally. The material cools because it is no longer near the core. It eventually becomes cool and dense enough to sink back down into the mantle. At the bottom of the mantle, the material travels horizontally and is heated by the core. It reaches the location where warm mantle material rises, and the mantle convection cell is complete.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=52

CORE

At the planet's center lies a dense metallic core. Scientists know that the core is metal for a few reasons. The density of Earth's surface layers is much less than the overall density of the planet, as calculated from the planet's rotation. If the surface layers are less dense than average, then the interior must be denser than average. Calculations indicate that the core is about 85 percent iron metal with nickel metal making up much of the remaining 15 percent. Also, metallic meteorites are thought to be representative of the core.If Earth's core were not metal, the planet would not have a magnetic field. Metals such as iron are magnetic, but rock, which makes up the mantle and crust, is not. Scientists know that the outer core is liquid and the inner core is solid because S-waves stop at the inner core. The strong magnetic field is caused by convection in the liquid outer core. Convection currents in the outer core are due to heat from the even hotter inner core. The heat that keeps the outer core from solidifying is produced by the breakdown of radioactive elements in the inner core.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-4-plate-tectonics.html. **License**: *CC BY: Attribution*

This page titled 4.1: Earth's Interior is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

4.2: Plate Tectonics

Plate Tectonics

LEARNING OUTCOMES


- **ESSENTIAL QUESTIONS**
  - What is the driving force of plate tectonics and how does this impact earthquakes and volcanoes around the world?
  - How does the theory of plate tectonics help explain the different types of earthquakes and volcanoes around the planet?

Theory of Continental Drift

The continental drift hypothesis was developed in the early part of the 20th century, mostly by Alfred Wegener. Wegener said that continents move around on Earth's surface and that they were once joined together as a single supercontinent. While Wegener was alive, scientists did not believe that the continents could move.

Find a map of the continents and cut each one out. Better yet, use a map where the edges of the continents show the continental shelf. That's the true size and shape of a continent and many can be pieced together like a puzzle. The easiest link is between the eastern Americas and western Africa and Europe, but the rest can fit together too.

Alfred Wegener proposed that the continents were once united into a single supercontinent named Pangaea, meaning all earth in ancient Greek. He suggested that Pangaea broke up long ago and that the continents then moved to their current positions. He called his hypothesis continental drift.




EVIDENCE FOR CONTINENTAL DRIFT

Besides the way the continents fit together, Wegener and his supporters collected a great deal of evidence for the continental drift hypothesis. For one, identical rocks of the same type and age are found on both sides of the Atlantic Ocean. Wegener said the rocks had formed side-by-side and that the land had since moved apart.Mountain ranges with the same rock types, structures, and ages are now on opposite sides of the Atlantic Ocean. The Appalachians of the eastern United States and Canada, for example, are just like mountain ranges in eastern Greenland, Ireland, Great Britain, and Norway. Wegener concluded that they formed as a single mountain range that was separated as the continents drifted.Ancient fossils of the same species of extinct plants and animals are found in rocks of the same age but are on continents that are now widely separated. Wegener proposed that the organisms had lived side by side, but that the lands had moved apart after they were dead and fossilized. He suggested that the organisms would not have been able to travel across the oceans. For example, the fossils of the seed fern *Glossopteris* were too heavy to be carried so far by wind. The reptile Mesosaurus could only swim in fresh water. was a swimming reptile but could only swim in fresh water.*Cynognathus* and *Lystrosaurus* were land reptiles and were unable to swim.Grooves and rock deposits left by ancient glaciers are found today on different continents very close to the equator. This would indicate that the glaciers either formed in the middle of the ocean and/or covered most of the Earth. Today glaciers only form on land and nearer the poles. Wegener thought that the glaciers were centered over the southern land mass close to the South Pole and the continents moved to their present positions later on.Coral reefs and coal-forming swamps are found in tropical and subtropical environments, but ancient coal seams and coral reefs are found in locations where it is much too cold today. Wegener suggested that these creatures were alive in warm climate zones and that the fossils and coal later had drifted to new locations on the continents.Although Wegener's evidence was sound, most geologists at the time rejected his hypothesis of continental drift. *Scientists argued that there was no way to explain how solid*


continents could move through solid oceanic crust. Wegener's idea was nearly forgotten until technological advances presented even more evidence that the continents moved and gave scientists the tools to develop a mechanism for Wegener's drifting continents.

Sea Floor Spreading



SEAFLOOR BATHYMETRY

World War II gave scientists the tools to find the mechanism for continental drift that had eluded Wegener. Maps and other data gathered during the war allowed scientists to develop the **seafloor spreading hypothesis**. This hypothesis traces oceanic crust from its origin at a mid-ocean ridge to its destruction at a deep sea trench and is the mechanism for continental drift.During World War II, battleships and submarines carried echo sounders to locate enemy submarines. Echo sounders produce sound waves that travel outward in all directions, bounce off the nearest object, and then return to the ship. By knowing the speed of sound in seawater, scientists calculate the distance to the object based on the time it takes for the wave to make a round-trip. During the war, most of the sound waves ricocheted off the ocean bottom. This animation shows how sound waves are used to create pictures of the seafloor and ocean crust.After the war, scientists pieced together the ocean depths to produce bathymetric maps, which reveal the features of the ocean floor as if the water were taken away. *Even scientist were amazed that the seafloor was not completely flat*. What was discovered was a large chain of mountains along the deep seafloor, called mid-ocean ridges. Scientists also discovered deep sea trenches along the edges of continents or in the sea near chains of active volcanoes. Finally, large, flat areas called abyssal plains we found. When they first observed these bathymetric maps, scientists wondered what had formed these features.




SEAFLOOR MAGNETISM

Sometimes, for reasons unknown, the magnetic poles switch positions. North becomes south and south becomes north. During normal polarity, the north and south poles are aligned as they are now. With reversed polarity, the north and south poles are in the opposite position. During WWII, magnetometers attached to ships to search for submarines located an astonishing feature; the normal and reversed magnetic polarity of seafloor basalts creates a pattern. Stripes of normal polarity and reversed polarity alternate across the ocean bottom. These stripes also forms a mirror image of itself on either side of the mid-ocean ridges. But the stripes end abruptly at the edges of continents, sometimes at a deep sea trench. The characteristics of the rocks and sediments change with distance from the ridge axis as seen in the **Table below**.

At ridge axis Distance from axis	**Rock Ages** Youngest Becomes older	**Sediment Thickness** None Becomes thicker	**Crust Thickness** Thinnest Becomes thicker	**Heat Flow** Hottest Becomes cooler


A map of sediment thickness is found here. The oldest seafloor is near the edges of continents or deep sea trenches and is less than 180 million years old. Since the oldest ocean crust is so much younger than the oldest continental crust, scientists realized that seafloor was being destroyed in a relatively short time.



SEAFLOOR SPREADING HYPOTHESIS

Scientists brought these observations together in the early 1960s to create the **seafloor spreading hypothesis**. In this hypothesis, hot buoyant mantle rises up a mid-ocean ridge, causing the ridge to rise upward. The hot magma at the ridge erupts as lava that forms new seafloor. When the lava cools, the magnetite crystals take on the current magnetic polarity and as more lava erupts, it pushes the seafloor horizontally away from ridge axis.The magnetic stripes continue across the seafloor. As oceanic crust forms and spreads, moving away from the ridge crest, it pushes the continent away from the ridge axis. If the oceanic crust reaches a deep sea trench, it sinks into the trench and is lost into the mantle. Scientists now know that the oldest crust is coldest and lies deepest in the ocean because it is less buoyant than the hot new crust.*Seafloor spreading is the mechanism for Wegener's drifting continents.* Convection currents within the mantle take the continents on a conveyor-belt ride of oceanic crust that over millions of years takes them around the planet's surface.

Earth's Tectonic Plates



When the concept of seafloor spreading came along, scientists recognized that it was the mechanism to explain how continents could move around Earth's surface. *Scientific data and observation now allows us to merge the ideas of continental drift and seafloor spreading into the* ***theory of plate tectonics****.*

Seafloor and continents move around on Earth's surface, but what is actually moving? What portion of the Earth makes up the "plates" in plate tectonics? This question was also answered because of technology developed during the Cold War. The **plates** are made up of the lithosphere. During the 1950s and early 1960s, scientists set up seismograph networks to see if enemy nations were testing atomic bombs. These seismographs also recorded all of the earthquakes around the planet. The seismic records could be used to locate an earthquake's epicenter, the point on Earth's surface directly above the place where the earthquake occurs. Earthquake epicenters outline these tectonic plates. Mid-ocean ridges, trenches, and large faults mark the edges of these plates along with where earthquakes occur.

The lithosphere is divided into a dozen major and several minor plates. The plates' edges can be drawn by connecting the dots that mark earthquakes' epicenters. *A single plate can be made of all oceanic lithosphere or all continental lithosphere, but nearly all plates are made of a combination of both.* Movement of the plates over Earth's surface is termed **plate tectonics**. Plates move at a rate of a few centimeters a year, about the same rate fingernails grow.

HOW PLATES MOVE


If seafloor spreading drives the plates, what drives seafloor spreading? Picture two convection cells side-by-side in the mantle. Hot mantle from the two adjacent cells rises at the ridge axis, creating new ocean crust. The top limb of the convection cell moves horizontally away from the ridge crest, as does the new seafloor. The outer limbs of the convection cells plunge down into the deeper mantle, dragging oceanic crust as well. This takes place at the deep sea trenches. The material sinks to the core and moves horizontally. The material heats up and reaches the zone where it rises again.

Tectonic Plate Boundaries

Plate boundaries are the edges where two plates meet. Most geologic activities, including volcanoes, earthquakes, and mountain building, take place at plate boundaries. How can two plates move relative to each other?

- Divergent plate boundaries: the two plates move away from each other.
- Convergent plate boundaries: the two plates move towards each other.
- Transform plate boundaries: the two plates slip past each other.

The type of plate boundary and the type of crust found on each side of the boundary determines what sort of geologic activity will be found there.



DIVERGENT PLATE BOUNDARIES

Plates move apart at mid-ocean ridges where new seafloor forms. Between the two plates is a rift valley. Lava flows at the surface cool rapidly to become basalt, but deeper in the crust, magma cools more slowly to form gabbro. So the entire ridge system is made up of igneous rock that is either extrusive or intrusive. Earthquakes are common at mid-ocean ridges since the movement of magma and oceanic crust results in crustal shaking. The vast majority of mid-ocean ridges are located deep below the sea. Here is an animation from the United States Geologic Survey (USGS) of divergent plate boundary at mid-ocean ridge and another animation by IRIS.As divergence occurs, shallow earthquakes can occur along with volcanoes along the rift areas. When the process begins, a valley will develop such as the Great Rift Valley in Africa. Over time that valley can fill up with water creating **linear lakes** (see Figure A below). If divergence continues, a sea can form like the Red Sea (Figure B). and finally an ocean like the Atlantic Ocean. Check out the eastern half of Africa and notice the lakes that look linear. Eastern Africa is tearing apart from these linear lakes, to the Great Rift Valley, and up to the Red Sea. The ultimate divergent boundary is the Mid-Atlantic Ridge which began forming when Pangea broke apart. The spreading along this ridge formed the Atlantic Ocean.



Figure A: Linear lakes within East African Rift Zone



Figure B: Continued spreading will form a sea, here the Red Sea.

CONVERGENT PLATE BOUNDARIES

When two plates converge, the result depends on the type of lithosphere the plates are made of. There are three types of convergent boundaries. **I. Ocean-to-continent convergence** occurs when oceanic crust converges with continental crust, forcing the denser oceanic plate to plunge beneath the continental plate. This process called **subduction**, occurs along oceanic trenches called **subduction zones** where lots of intense earthquakes and volcanic eruptions can occur. The denser, subducting plate begins to heat up under extreme pressure near the mantle and melts to create causes melting in the volcanoes. These coastal volcanic mountains are found in a line above the subducting plate. The volcanoes are known as a continental arc. The movement of crust and magma causes earthquakes. Click here for a map of earthquake epicenters at subduction zones.The volcanoes of northeastern California—Lassen Peak, Mount Shasta, and Medicine Lake volcano—along with the rest of the Cascade Mountains of the Pacific Northwest are the result of subduction of the Juan de Fuca plate beneath the North American plate. The Juan de Fuca plate is created by seafloor spreading just offshore at the Juan de Fuca ridge. Figure C below is of the west coast of South America. The Andes Mountains are created here where the Nazca Plate, composed of oceanic lithosphere, is subducted under the continental lithosphere of the South American plate.



Figure C: South America's western coast and Andes Mountains




Figure D: Oceanic plate convergence along Pacific Ocean (Ring of Fire)

II. Oceanic-to-oceanic plate boundary occurs when two oceanic plates converge, causing the older, denser plate will subduct into the mantle. An ocean trench marks the location where the plate is pushed down into the mantle. The line of volcanoes that grows on the upper oceanic plate is an island arc.

The Ring of Fire, as seen in Figure D above, is a ring around the Pacific Ocean of subduction zones, which most are oceanic-to-oceanic convergence. Here is an animation of an ocean continent plate boundary. Along these subduction zones, volcanic islands (also called volcanic arcs) form. Examples of these regions include Japan, Indonesia, and the Aleutian Islands.






Figure E: India's collision with Asia forming the Himalayan Mountains.

III. Continental-to-continental convergence occurs when two continental plates collide, instead of subduction, the two similar tectonic plates will buckle up to create large mountain ranges like a massive car pile-up. This **continental-to-continental convergence** creates intense folding and faulting rather than volcanic activity.

Examples of mountain ranges created by this process are the Himalayan mountains (see Figure E above) as India is colliding with Asia, the Alps in Europe, and the Appalachian mountains in the United States as the North American plate collided with the African plate when Pangea was forming. The Kashmir India earthquake of 2005 that killed over 80,000 people occurred because of this process. And most recently, the 2008 earthquake in China which killed nearly 85,000 people before the Summer Olympics was because of this tectonic force. The Appalachian Mountains are the remnants of a large mountain range that was created when North America rammed into Eurasia about 250 million years ago.

Transform plate boundaries occur when two tectonic plates slide (or grind) past parallel to each other. The most famous **transform boundary** is the **San Andreas Fault** where the Pacific plate that Los Angeles and Hawaii are on is grinding past the North American plate that San Francisco and the rest of the United States is on at the rate of 3 inches a year. Recently, geologists have stated that San Francisco should expect another disastrous earthquake in the next 30 years. Another important transform boundary is the **North Anatolian Fault** in Turkey. This powerful fault last ruptured in 1999 in Izmit, Turkey which killed 17,000 people in 48 seconds.





INTRAPLATE BOUNDARIES

A small amount of geologic activity, known as **intraplate activity**, does not take place at plate boundaries but within a plate instead. Mantle plumes are pipes of hot rock that rise through the mantle. The release of pressure causes melting near the surface to

form a **hotspot**. Eruptions at the hotspot create a volcano. Hotspot volcanoes are found in a line. Can you figure out why? Hint: The youngest volcano sits above the hotspot and volcanoes become older with distance from the hotspot.An animation of the creation of a hotspot chain is seen here:http://earthguide.ucsd.edu/eoc/teachers/t_tectonics/p_hawaii.html.Geologists use some hotspot chains to tell the direction and the speed a plate is moving.Hotspot magmas rarely penetrate through thick continental crust. One exception is the Yellowstone hotspot.

Earth's Changing Surface

Geologists know that Wegener was right because the movements of continents explain so much about the geology we see. Most of the geologic activity that we see on the planet today is because of the interactions of the moving plates.

In the map of North America, where are the mountain ranges located? Using what you have learned about plate tectonics, try to answer the following questions:

1. What is the geologic origin of the Cascades Range? The Cascades are a chain of volcanoes in the Pacific Northwest. They are not labelled on the diagram but they lie between the Sierra Nevada and the Coastal Range.
2. What is the geologic origin of the Sierra Nevada? (Hint: These mountains are made of granitic intrusions.)
3. What is the geologic origin of the Appalachian Mountains along the Eastern US?

Remember that Wegener used the similarity of the mountains on the west and east sides of the Atlantic as evidence for his continental drift hypothesis. The Appalachian mountains formed at a convergent plate boundary as Pangaea came together.

Before Pangaea came together, the continents were separated by an ocean where the Atlantic is now. The proto-Atlantic ocean shrank as the Pacific ocean grew. Currently, the Pacific is shrinking as the Atlantic is growing. This supercontinent cycle is responsible for most of the geologic features that we see and many more that are long gone (Figure below).

This animation shows the movement of continents over the past 600 million years beginning with the breakup of Rodinia:http://earthguide.ucsd.edu/eoc/teachers/t_tectonics/p_plate_reconstruction_blakey.html


CC licensed content, Shared previously

- Plate Tectonics. **Authored by**: Open Geography. **Located at**: http://www.opengeography.org/ch-5-plate-tectonics.html. **License**: *CC BY: Attribution*

This page titled 4.2: Plate Tectonics is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

4.3: Volcanoes


Where Volcanoes Are Located



Google Earth image of Japan and the oceanic-to-oceanic subduction zones.

VOLCANOES ALONG CONVERGENT PLATE BOUNDARIES


Volcanoes are a vibrant manifestation of plate tectonics processes. Volcanoes are common along convergent and divergent plate boundaries, but are also found within lithospheric plates away from plate boundaries. Wherever mantle is able to melt, volcanoes may be the result.Volcanoes erupt because mantle rock melts. The first stage in creating a volcano is when mantle rock begins to melt because of extremely high temperatures, lithospheric pressure lowers, or water is added.Along subducting plate boundaries, the crust heats up as it sinks into the mantle. Also, ocean water is mixed in with the sediments lying on top of the subducting plate. This water lowers the melting point of the mantle material, which increases melting. Volcanoes at convergent plate boundaries are found all along the Pacific Ocean basin, primarily at the edges of the Pacific, Cocos, and Nazca plates.



Large earthquakes are extremely common along convergent plate boundaries. Since the Pacific Ocean is rimmed by convergent and transform boundaries, about 80% of all earthquakes strike around the Pacific Ocean basin and is why the region is called the Ring of Fire. A description of the Pacific Ring of Fire along western North America is below:

- Subduction at the Middle American Trench creates volcanoes in Central America.
- The San Andreas Fault is a transform boundary.
- Subduction of the Juan de Fuca plate beneath the North American plate creates the Cascade volcanoes like Mount St. Helens, Mount Rainer, Mount Hood and more.
- Subduction of the Pacific plate beneath the North American plate in the north creates the long chain of the Aleutian Islands volcanoes near Alaska.

This incredible explosive eruption of Mount Vesuvius in Italy in A.D. 79 is an example of a composite volcano that forms as the result of a convergent plate boundary.






VOLCANOES ALONG DIVERGENT PLATE BOUNDARIES


Why does melting occur at divergent plate boundaries? Hot mantle rock rises where the plates are moving apart. This releases pressure on the mantle, which lowers its melting temperature allowing lava to erupt through long cracks or fissures. Scientists have captured incredible footage of "Deepest Ocean Eruption Ever Filmed" and other undersea volcanoes. Volcanoes erupt at mid-ocean ridges, such as the Mid-Atlantic ridge, where seafloor spreading creates new seafloor in the rift valleys. Where a hotspot is located along the ridge, such as at Iceland, volcanoes grow high enough to create islands. Eruptions are found at divergent plate boundaries as continents break apart such as the East African Rift between the African and Arabian plates and the Great Basin and Range in the western United States. But those volcanoes of the ladder are now extinct.For some dramatic time-lapsed photography of a volcanic eruption on Iceland, click here. Iceland (image on the right) is a hot spot volcano and the surface manifestation of the mid-Atlantic ridge.

VOLCANIC HOTSPOTS


Although most volcanoes are found at convergent or divergent plate boundaries, intraplate volcanoes are found in the middle of a tectonic plate. The Hawaiian Islands are the exposed peaks of a great chain of volcanoes that lie on the Pacific plate. These islands are in the middle of the Pacific plate. The youngest island sits directly above a column of hot rock called a mantle plume. As the plume rises through the mantle, pressure is released and mantle melts to create a **hotspot**. Earth is home to about 50 known hot spots. Most of these are in the oceans because they are better able to penetrate oceanic lithosphere to create volcanoes. The hotspots that are known beneath continents are extremely large, such as Yellowstone. The video on the right is of the hot spot beneath Hawaii, the origin of the voluminous lava produced by the shield volcano Kilauea. There are several key indicators to determine a

hot spot from island arc volcanoes. At island arcs, the volcanoes are all about the same age. By contrast, at hotspots the volcanoes are youngest at one end of the chain and oldest at the other.





Magma Composition



In 1980, Mount St. Helens blew up in the costliest and deadliest volcanic eruption in United States history. The eruption killed 57 people, destroyed 250 homes and swept away 47 bridges. Mount St. Helens today still has minor earthquakes and eruptions, and now has a horseshoe-shaped crater with a lava dome inside. The dome is formed of viscous lava that oozes into place.

It should first be noted that **magma** is molten material inside the earth, whereas **lava** is molten material on the surface of the earth. The reason for the distinction is because lava can cool quickly from the air and solidify into rock rapidly, whereas magma may never reach the earth's surface. Volcanoes do not always erupt in the same way. Each volcanic **eruption** is unique, differing in size, style, and composition of erupted material. One key to what makes the eruption unique is the chemical composition of the magma that feeds a volcano, which determines (1) the eruption style, (2) the type of volcanic cone that forms, and (3) the composition of rocks that are found at the volcano.

Different minerals within a rocks melt at different temperatures and the amount of partial melting and the composition of the original rock determine the composition of the magma. Magma collects in **magma chambers** in the crust at 160 kilometers (100 miles) beneath the surface of a volcano.

The words that describe composition of igneous rocks also describe magma composition. Mafic magmas are low in silica and contain more dark, magnesium and iron rich mafic minerals, such as olivine and pyroxene. Felsic magmas are higher in silica and contain lighter colored minerals such as quartz and orthoclase feldspar. The higher the amount of silica in the magma, the higher is its **viscosity**. Viscosity is a liquid's resistance to flow.

Viscosity determines what the magma will do. *Mafic magma is not viscous and will flow easily to the surface. Felsic magma is viscous and does not flow easily.* Most felsic magma will stay deeper in the crust and will cool to form igneous intrusive rocks such as granite and granodiorite. If felsic magma rises into a magma chamber, it may be too viscous to move and so it gets stuck. Dissolved gases become trapped by thick magma and the magma chamber begins to build pressure.



EXPLOSIVE ERUPTIONS

The type of magma in the chamber determines the type of volcanic eruption. A **large explosive eruption** creates even more devastation than the force of the atom bomb dropped on Nagasaki at the end of World War II in which more than 40,000 people died. A large explosive volcanic eruption is 10,000 times as powerful. Felsic magmas erupt explosively because of hot, gas-rich magma churning within its chamber. The pressure becomes so great that the magma eventually breaks the seal and explodes, just like when a cork is released from a bottle of champagne. Magma, rock, and ash burst upward in an enormous explosion creating volcanic ash called **tephra**. It should be noted that when looked under a microscope, the volcanic "ash" is actual microscopic shards of glass. That is why it is so dangerous to inhale the air following an eruption.Scorching hot tephra, ash, and gas may speed down the volcano's slopes at 700 km/h (450 mph) as a **pyroclastic flow**. Pyroclastic flows knock down everything in their path. The temperature inside a pyroclastic flow may be as high as 1,000^{o}C (1,800 degrees F).

Prior to the Mount St. Helens eruption in 1980, the Lassen Peak eruption on May 22, 1915, was the most recent Cascades eruption. A column of ash and gas shot 30,000 feet into the air. This triggered a high-speed pyroclastic flow, which melted snow and created a **volcanic mudflow** known as a **lahar**. Lassen Peak currently has geothermal activity and could erupt explosively again. Mt. Shasta, the other active volcano in California, erupts every 600 to 800 years. An eruption would most likely create a large pyroclastic flow, and probably a lahar. Of course, Mt. Shasta could explode and collapse like Mt. Mazama in Oregon.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56

Volcanic gases can form poisonous and invisible clouds in the atmosphere that could contribute to environmental problems such as acid rain and ozone destruction. Particles of dust and ash may stay in the atmosphere for years, disrupting weather patterns and blocking sunlight.

EFFUSIVE ERUPTIONS

Mafic magma creates gentler **effusive eruptions**. Although the pressure builds enough for the magma to erupt, it does not erupt with the same explosive force as felsic magma. People can usually be evacuated before an effusive eruption, so they are much less deadly. Magma pushes toward the surface through fissures and reaches the surface through volcanic vents. Click here to view a lava stream within the vent of a Hawaiian volcano using a thermal camera.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56

Low-viscosity lava flows down mountainsides. Differences in composition and where the lavas erupt result in lava types like a ropy form **pahoehoe** and a chunky form called **aa**. Although effusive eruptions rarely kill anyone, they can be destructive. Even when people know that a lava flow is approaching, there is not much anyone can do to stop it from destroying a building, road, or infrastructure.



Pahoehoe lava



Aa lava

Predicting Volcanic Eruptions

Volcanologists attempt to forecast volcanic eruptions, but this has proven to be nearly as difficult as predicting an earthquake. Many pieces of evidence can mean that a volcano is about to erupt, but the time and magnitude of the eruption are difficult to pin down. This evidence includes the history of previous volcanic activity, earthquakes, slope deformation, and gas emissions.



HISTORY OF VOLCANIC ACTIVITY

A volcano's history, how long since its last eruption and the time span between its previous eruptions, is a good first step to predicting eruptions. If the volcano is considered **active**, it is currently erupting or shows signs of erupting soon. A **dormant** volcano means there is no current activity, but it has erupted recently. Finally, an **extinct** volcano means their is no activity and will probably not erupt again. Active and dormant volcanoes are heavily monitored, especially in populated areas.

EARTHQUAKES

Moving magma shakes the ground, so the number and size of earthquakesincreases before an eruption. A volcano that is about to erupt may produce a sequence of earthquakes. Scientists use seismographs that record the length and strength of each earthquake to try to determine if an eruption is imminent.Magma and gas can push the volcano's slope upward. Most ground deformation is subtle and can only be detected by tiltmeters, which are instruments that measure the angle of the slope of a volcano. But ground swelling may sometimes create huge changes in the shape of a volcano. Mount St. Helens grew a bulge on its north side before its 1980 eruption. Ground swelling may also increase rockfalls and landslides.

GAS EMISSIONS

Gases may be able to escape a volcano before magma reaches the surface. Scientists measure gas emissions in vents on or around the volcano. Gases, such as sulfur dioxide (SO_2), carbon dioxide (CO_2), hydrochloric acid (HCl), and even water vapor can be measured at the site or, in some cases, from a distance using satellites. The amounts of gases and their ratios are calculated to help predict eruptions.

REMOTE MONITORING

Some gases can be monitored using satellite technology. Satellites also monitor temperature readings and deformation. As technology improves, scientists are better able to detect changes in a volcano accurately and safely.

Since volcanologists are usually uncertain about an eruption, officials may not know whether to require an evacuation. If people are evacuated and the eruption doesn't happen, the people will be displeased and less likely to evacuate the next time there is a threat of an eruption. The costs of disrupting business are great. However, scientists continue to work to improve the accuracy of their predictions.

Types of Volcanoes

A volcano is a vent through which molten rock and gas escape from a magma chamber and they can differ in height, shape, and slope steepness. Some volcanoes are tall cones and others are just cracks in the ground. As you might expect, the shape of a volcano is related to the composition of its magma.



COMPOSITE VOLCANOES

Composite volcanoes are some of the most dangerous volcanoes on the planet. They tend to occur along oceanic-to-oceanic or oceanic-to-continental boundaries because of subduction zones. They tend to be made of felsic to intermediate rock and the viscosity of the lava means that eruptions tend to be explosive. The viscous lava cannot travel far down the sides of the volcano before it solidifies, which creates the steep slopes of a composite volcano. Viscosity also causes some eruptions to explode as ash and small rocks. The volcano is constructed layer by layer, as ash and lava solidify, one upon the other and are sometimes called stratovolcanoes or andesite volcanoes. The result is the classic cone shape of composite volcanoes. Examples of composite volcanoes include Mount St. Helens, Mount Rainer, Mount Shasta, Mount Hood, and Mount Pinatubo. Here's a great time-lapse of Mount St. Helens from NASA's Earth Observatory from 1979 to 2013.Sometimes composite volcanoes and other violent volcanoes can erupt so violently that they sometimes collapse in on themselves or actually blow themselves up to produce calderas. One of the most powerful volcanoes in the world – Yellowstone – is a massive caldera that has collapsed several times. Sometimes these calderas can fill up with water to produce beautiful lakes such as Mount Mazama (Crater Lake), in Oregon.




SHIELD VOLCANOES

Shield volcanoes get their name from their shape. Although shield volcanoes are not steep, they may be very large. In fact, Mauna Loa, Hawaii is the tallest mountain in the world. From sea level, Mount Everest is the tallest, but when you consider from the ocean floor to the top of the island, Mauna Loa wins. Shield volcanoes are common at spreading centers or intraplate hot spots.The lava that creates shield volcanoes is fluid and flows easily and creates the shield shape. Shield volcanoes are built by many layers over time and the layers are usually of very similar composition. The low viscosity also means that shield eruptions are non-explosive. Eruptions tend to be mild in comparison to other volcanoes, but lava flows can destroy property and vegetation. The low viscosity

magma can flow not only on the surface as lava, but also underground in lava tubes. The most well known shield volcano is Hawaii. There are two types of lava flows, pahoehoe which is a ropy type of lava that flows easily (low viscosity). The other type is called aa and is a blocky type of lava and has a higher viscosity and does not like to flow well. The following is a short video on Hawaii, an example of a shield volcano.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56



CINDER CONES

Cinder cones are the most common type of volcano. A cinder cone has a cone shape, but is much smaller than a composite volcano. Cinder cones rarely reach 300 meters in height but they have steep sides. Cinder cones grow rapidly, usually from a single eruption cycle. Cinder cones are composed of small fragments of rock, such as pumice, piled on top of one another. The rock shoots up in the air and doesn't fall far from the vent. The exact composition of a cinder cone depends on the composition of the lava ejected from the volcano. Cinder cones usually have a crater at the summit. Cinder cones are often found near larger volcanoes.



SUPERVOLCANOES

Supervolcano eruptions are extremely rare in Earth history. It's a good thing because they are unimaginably large. A supervolcano must erupt more than 1,000 cubic km (240 cubic miles) of material, compared with 1.2 km3 for Mount St. Helens or 25 km3 for Mount Pinatubo, a large eruption in the Philippines in 1991. Not surprisingly, supervolcanoes are the most dangerous type of volcano.Supervolcanoes are a fairly new idea in volcanology. The exact cause of supervolcano eruptions is still debated, however, scientists think that a very large magma chamber erupts entirely in one catastrophic explosion. This creates a huge hole or **caldera** into which the surface collapses.

The largest supervolcano in North America is beneath Yellowstone National Park in Wyoming. Yellowstone sits above a hotspot that has erupted catastrophically three times: 2.1 million, 1.3 million, and 640,000 years ago. Yellowstone has produced many smaller (but still enormous) eruptions more recently. Fortunately, current activity at Yellowstone is limited to the region's famous geysers.
Long Valley Caldera, south of Mono Lake in California, is the second largest supervolcano in North America. Long Valley had an extremely hot and explosive rhyolite explosion about 700,000 years ago. An earthquake swarm in 1980 alerted geologists to the possibility of a future eruption, but the quakes have since calmed down.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56

A supervolcano could change life on Earth as we know it. Ash could block sunlight so much that photosynthesis would be reduced and global temperatures would plummet. Volcanic eruptions could have contributed to some of the mass extinctions in our planet's history. No one knows when the next super eruption will be.

Volcanic Landforms and Geothermal Activity



VOLCANIC LANDFORMS AND VENTS

Volcanoes are associated with many types of landforms. The landforms vary with the composition of the magma that created them. Hot springs and geysers are also examples of surface features related to volcanic activity.The most obvious landforms created by lava are volcanoes, most commonly as cinder cones, composite volcanoes, and shield volcanoes or eruptions that take place through fissures. The eruptions that created the entire ocean floor are essentially fissure eruptions. Magma intrusions ALSO can create landforms. The image on the right is of Shiprock in New Mexico, which is the neck of an old volcano that has eroded away



Lava dome inside Mount St. Helen's crater.

LAVA DOMES

When lava is viscous, it flows slowly. If there is not enough magma or enough pressure to create an explosive eruption, the magma may form a lava dome. But because the viscosity of the magma is so thick, the lava does not flow far from the vent. Lava flows often make mounds right in the middle of craters at the top of volcanoes.LAVA PLATEAUS AND LAND

A **lava plateau** forms when large amounts of fluid lava flows over an extensive area. When the lava solidifies, it creates a large, flat surface of igneous rock. Lava creates new land as it solidifies on the coast or emerges from beneath the water. Over time the


eruptions can create whole islands. The Hawaiian Islands are formed from shield volcano eruptions that have grown over the last 5 million years.



HOT SPRINGS AND GEYSERS

Water sometimes comes into contact with hot rock. The water may emerge at the surface as either a hot spring or a geyser. Water heated below ground that rises through a crack to the surface creates a **hot spring**. The water in hot springs may reach temperatures in the hundreds of degrees Celsius beneath the surface, although most hot springs are much cooler.

Hazards and Benefits of Volcanic Activity



There are several hazards that volcanic activity can produce.

- Eruption clouds occur when massive quantities of ash is ejected into the atmosphere where it can reach heights of 50,000 feet. Eruption clouds have proven to be very dangerous for aviation jets because the ash can shut down the engines. The ash cloud can also be very hazardous in terms of air pollution.
- Lahars are volcanic mudflows. Lahars are very dangerous because they do not require a volcanic eruption yet can travel hundreds of miles. All that is required is loose pyroclastic material on the volcano that mixes with precipitation or melting snow.
- Lava flows are layers of molten rock that flow over the surface, later cooling and solidifying.
- Lava bombs are large chunks of pyroclastic material ejected from a volcano. Larger pyroclastic material is called blocks.

- Pyroclastic flows are some of the most dangerous hazards caused by composite volcanoes. Pyroclastic flows are superheated clouds of pyroclastic material (e.g. hot rock and tephra) ranging in size from small rocks to the size of houses that are over 1,000 degrees F traveling down a mountain at speeds up to 100 mph.
- Tephra (or volcanic ash) is fine particles of pyroclastic material that can be carried thousands of miles away by prevailing winds. Regions hundreds of miles away could suffer collapsed buildings is the falling ash accumulates enough. Tephra can also cool the entire planet if enough is ejected into the atmosphere.

- Poisonous gases such as carbon dioxide, carbon monoxide, and sulfur dioxide can travel down a volcano and asphyxiate (suffocating) wildlife and humans. In 1986, an invisible cloud of carbon dioxide traveled down a volcano in Africa asphyxiating 1,742 people and 3,000 cattle.

There are actually many benefits to volcanic activity. One of the major benefits is the fact that volcanic activity can create very fertile soil for agriculture. The problem is that many civilizations developed near volcanoes for this reason – with sometimes deadly effects. Volcanic activity can also create many mineral resources such as gold, sliver, nickel, copper, and lead. Volcanic rock is often used for landscaping, tile, and cement.

Some of the most amazing landscapes are near volcanoes. This is because volcanic activity builds land creating breathtaking scenery. So volcanoes are economically vital for many regions because of the recreational activity and tourism they bring.

Finally, a new but important trend is **geothermal power**. The heat generated by volcanoes can create electricity to power civilization. Geothermal power is a completely **renewable resource** free of pollution and energy dependency on fossil fuels. Iceland – the surface manifestation of the mid-Atlantic ridge – has a goal of powering the entire nation on geothermal energy. Geothermal energy is also being used in California, Kilauea, Hawaii, and now Utah.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=56

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-5-tectonic-forces.html. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 4.3: Volcanoes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

4.4: Earthquakes

LEARNING OUTCOMES

- **ESSENTIAL QUESTIONS**
  - Could humans live in places along the Wasatch Front without the Wasatch fault and the earthquakes it causes?
  - How does plate tectonics explain where and why earthquakes and volcanoes occur where they do?
  - What role do earthquakes and volcanoes have on civilization? What benefits do they bring us?
  - How could understanding Earth's tectonic forces help save lives?
  - What beneficial roles do tectonic forces have ecologically and for humanity?

Causes and Types of Tectonic Stress



Enormous slabs of lithosphere move unevenly over the planet's spherical surface, resulting in earthquakes. This chapter deals with two types of geological activity that occur because of plate tectonics: mountain building and earthquakes. First, we will consider what can happen to rocks when they are exposed to stress. **Stress** is the force applied to an object. In geosciences, stress is the force per unit area that is placed on a rock. Four types of stresses act on materials.

- A deeply buried rock is pushed down by the weight of all the material above it. Since the rock cannot move, it cannot deform called **confining stress**.
- **Compression** squeezes rocks together, causing rocks to fold or fracture. Compression is the most common stress at convergent plate boundaries.
- Rocks that are pulled apart are under **tension**. Rocks under tension lengthen or break apart. Tension is the major type of stress at divergent plate boundaries.
- When forces are parallel but moving in opposite directions, the stress is called shear. **Shear** stress is the most common stress at transform plate boundaries.

When stress causes a material to change shape, it has undergone strain or **deformation**. Deformed rocks are common in geologically active areas. A rock's response to stress depends on the rock type, the surrounding temperature, and pressure conditions the rock is under, the length of time the rock is under stress, and the type of stress. The rocks then have three possible responses to increasing stress: elastic deformation, plastic deformation, or fracturing. **Elastic deformation** occurs when the rock returns to its original shape when the stress is removed. When rocks under stress do not return to its original shape when the stress is removed, it is called **plastic deformation**. Finally, when a rock under stress breaks, it's called a **fracture.** Fractures can be either **joints** or **faults.**

Under what conditions do you think a rock is more likely to fracture? Is it more likely to break deep within Earth's crust or at the surface? What if the stress applied is sharp rather than gradual? At the Earth's surface, rocks usually break quite quickly, but deeper in the crust, where temperatures and pressures are higher, rocks are more likely to deform plastically. Sudden stress, such as a hit with a hammer, is more likely to make a rock break. Stress applied over time often leads to plastic def

Geologic Structures: Joints and Faults



FAULTS

A rock under enough stress will eventually fracture. If there is no movement on either side of a fracture, the fracture is called a **joint**. But if the blocks of rock on one or both sides of a fracture move, the fracture is called a **fault**. Sudden motions along faults cause rocks to break and move suddenly, releasing the stored up stress energy to create an earthquake.A **slip** is the distance rocks move along a fault and can be up or down the fault plane. Slip is relative, because there is usually no way to know whether both sides moved or only one. Faults lie at an angle to the horizontal surface of the Earth. That angle is called the fault's **dip**. The dip defines which of two basic types a fault is. If the fault's dip is inclined relative to the horizontal, the fault is a dip-slip fault. There are two types of **dip-slip faults**. In **normal faults**, the hanging wall drops down relative to the footwall. Normal faults can be huge and are often times responsible for uplifting mountain ranges in regions experiencing tensional stress.With **reverse faults**, the footwall drops down relative to the hanging wall. A type of reverse fault is a thrust fault, in which the fault plane angle is nearly horizontal. Rocks can slip many miles along thrust faults.A **strike-slip fault** is a dip-slip fault in which the dip of the fault plane is vertical and result from shear stresses. California's San Andreas Fault is the world's most famous strike-slip fault. It is a right-lateral strike slip fault.

Causes of Earthquakes


An **earthquake** is sudden ground movement caused by the sudden release of energy stored in rocks, called the **elastic rebound theory**. Earthquakes happen when so much stress builds up in the rocks that the rocks rupture. The energy is transmitted by seismic waves. Each year there are more than 150,000 earthquakes strong enough to be felt by people and 900,000 recorded by seismometers!

In an earthquake, the initial point where the rocks rupture in the crust is called the **focus** (sometimes called the **hypocenter**). The **epicenter** is the point on the land surface that is directly above the focus. In about 75% of earthquakes, the focus is in the top 10 to 15 kilometers (6 to 9 miles) of the crust. **Shallow earthquakes** cause the most damage because the focus is near where people live. However, it is the epicenter of an earthquake that is reported by scientists and the media.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=58	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=58

Earthquake Zones


Nearly 95% of all earthquakes take place along one of the three types of tectonic plate boundaries, but earthquakes do occur along all three types of plate boundaries. About 80% of all earthquakes strike around the Pacific Ocean basin because it is lined with convergent and transform boundaries. Called the **Ring of Fire**, this is also the location of most volcanoes around the planet. About 15% take place in the Mediterranean-Asiatic Belt, where convergence is causing the Indian Plate to run into the Eurasian Plate creating the largest mountain ranges in the world. The remaining 5% are scattered around other plate boundaries or are **intraplate earthquakes**.



Strike-slip fault

TRANSFORM PLATE BOUNDARIES

Recall that transform plate boundaries occur where two tectonic plates are grinding parallel to each other rather than colliding or subducting. Deadly earthquakes occur at transform plate boundaries, creating **strike-slip faults**, because they tend to have shallow focuses where the rupture occurs. The faults along the San Andreas Fault zone produce around 10,000 earthquakes a year. Most are tiny, but occasionally one is massive. In the San Francisco Bay Area, the Hayward Fault was the site of a magnitude 7.0 earthquake in 1868. The 1906 quake on the San Andreas Fault had a magnitude estimated at about 7.9.During the 1989 World Series, a magnitude 7.1 earthquake struck Loma Prieta, near Santa Cruz, California, killing 63 people, injuring 3,756, and cost $6 billion. A few years later in Northridge, California, a magnitude 6.7 earthquake killed 72 people, injured 12,000 people, and caused $12.5 billion in damage. This earthquake occurred on an unknown fault, because it was a blind thrust fault near Los Angeles, California.Although California is prone to many natural hazards, including volcanic eruptions at Mt. Shasta or Mt. Lassen, and landslides on coastal cliffs, the natural hazard the state is linked with is earthquakes. New Zealand also has strike-slip earthquakes, about 20,000 a year, but only a small percentage of those are large enough to be felt. A 6.3 quake in Christchurch in February 2011 killed about 180 people.




Reverse fault

CONVERGENT PLATE BOUNDARIES

Earthquakes at convergent plate boundaries mark the motions of subducting lithosphere as it plunges through the mantle, creating **reverse** and **thrust faults**. Convergent plate boundaries produce earthquakes all around the Pacific Ocean basin. The Philippine Plate and the Pacific Plate subduct beneath Japan, creating a chain of volcanoes and produces as many as 1,500 earthquakes annually.In March 2011 an enormous 9.0 earthquake struck off of Sendai in northeastern Japan. This quake, called the 2011 Tōhoku earthquake, was the most powerful ever to strike Japan and one of the top five known in the world. Damage from the earthquake was nearly overshadowed by the tsunami it generated, which wiped out coastal cities and towns. Two months after the earthquake, about 25,000 people were dead or missing, and 125,000 buildings had been damaged or destroyed. Aftershocks, some as large as major earthquakes, have continued to rock the region. A map of aftershocks is seen here. Recently, the New York Times created an interactive website of the Japan earthquake and tsunami.The Pacific Northwest of the United States is at risk from a potentially massive earthquake that could strike any time. Subduction of the Juan de Fuca plate beneath North America produces active volcanoes, but large earthquakes only hit every 300 to 600 years. The last was in 1700, with an estimated magnitude of around 9.0. The elastic rebound theory as applied to subduction zones can be viewed here.

Massive earthquakes are the hallmark of the thrust faulting and folding when two continental plates converge. The 2001 Gujarat earthquake in India was responsible for about 20,000 deaths, and many more people became injured or homeless.
In Understanding Earthquakes: From Research to Resilience, scientists try to understand the mechanisms that cause earthquakes and tsunamis and the ways that society can deal with them.


An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=58



Normal fault

DIVERGENT PLATE BOUNDARIES


Earthquakes at mid-ocean ridges are small and shallow because the plates are young, thin, and hot. On land where continents split apart, earthquakes are larger and stronger. A classic example of **normal faulting** along divergent boundaries is the Wasatch Front in Utah and the entire Basin and Range through Nevada.



INTRAPLATE BOUNDARIES


Intraplate earthquakes are the result of stresses caused by plate motions acting in solid slabs of lithosphere. In 1812, a magnitude 7.5 earthquake struck near New Madrid, Missouri. The earthquake was strongly felt over approximately 50,000 square miles and altered the course of the Mississippi River. Because very few people lived there at the time, only 20 people died. Many more people live there today. A similar earthquake today would undoubtedly kill many people and cause a great deal of property damage.

Seismic Waves




Physics shows us that energy is always transmitted in waves. Every wave has a high point called a **crest** and a low point called a **trough**. The height of a wave from the center line to its crest is its **amplitude**. The distance between waves from crest to crest (or trough to trough) is its **wavelength**.

The energy from earthquakes travels in seismic waves, which were discussed in the chapter "Plate Tectonics." The study of seismic waves is known as **seismology**. Seismologists use seismic waves to learn about earthquakes and also to learn about the Earth's interior. The two types of seismic waves described in "Plate Tectonics," P-waves and S-waves, are known as **body waves** because they move through the solid body of the Earth. P-waves travel through solids, liquids, and gases. S-waves only move through solids.**Surface waves** travel along the ground, outward from an earthquake's epicenter. Surface waves are the slowest of all seismic waves, traveling at 2.5 km (1.5 miles) per second. In an earthquake, body waves produce sharp jolts, while rolling motions of surface waves do most of the damage in an earthquake.



MEASURING MAGNITUDE


Seismograms record seismic waves. Over the past century, scientists have developed several ways of measuring earthquake intensity. The currently accepted method is the moment magnitude scale, which measures the total amount of energy released by the earthquake. At this time, seismologists have not found a reliable method for predicting earthquakes.A **seismograph** produces a graph-like representation of the seismic waves it receives and records them onto a **seismogram**. Seismograms contain information that can be used to determine how strong an earthquake was, how long it lasted, and how far away it was. Modern **seismometers** record ground motions using electronic motion detectors. The data are then kept digitally on a computer.



If a seismogram records P-waves and surface waves but not S-waves, the seismograph was on the other side of the Earth from the earthquake because those waves cannot travel through the liquid core of the earth. The amplitude of the waves can be used to determine the magnitude of the earthquake, which will be discussed in a later section.

In order to locate an earthquake epicenter, scientists must first determine the epicenter distance from three different seismographs. The longer the time between the arrival of the P-wave and S-wave, the farther away is the epicenter. So the difference in the P and S wave arrival times determines the distance between the epicenter and a seismometer. This animation shows how distance is determined using P, S, and surface waves.


The scientist then draws a circle with a radius equal to the distance from the epicenter for that seismograph. The epicenter is somewhere along that circle. This is done for three locations. Using data from two seismographs, the two circles will intercept at two points. A third circle will intercept the other two circles at a single point. This point is the earthquake epicenter. Although useful for decades, this technique has been replaced by digital calculations. Seismic stations record ten earthquakes in this animation.



MEASURING EARTHQUAKES

People have always tried to quantify the size of and damage done by earthquakes. Since early in the 20th century, there have been three methods The oldest of the scales is called the **Mercalli Intensity scale**. Earthquakes are described in terms of what nearby residents felt and the damage that was done to nearby structures. This scale is more qualitative in information because it's based on visual damage and not the actual energy released by the earthquake. Today these maps are still important and various seismological stations will create **shake maps** of the surface damage.With the invention of the seismograph station, the **Richter magnitude scale** was created. Developed in 1935 by Charles Richter, this scale uses a seismometer to measure the magnitude of the largest jolt of energy released by an earthquake. Today, the moment magnitude scale has replaced the Richter scale. The **moment magnitude scale** measures the total energy released by an earthquake. Moment magnitude is calculated from the area of the fault that is ruptured and the distance the ground moved along the fault.The Richter scale and the moment magnitude scale are logarithmic. The amplitude of the largest wave increases ten times from one integer to the next. An increase in one integer means that thirty times more energy was released. These two scales often give very similar measurements.How does the amplitude of the largest seismic wave of a magnitude 5 earthquake compare with the largest wave of a magnitude 4 earthquake? How does it compare with a magnitude 3 quake? The amplitude of the largest seismic wave of a magnitude 5 quake is 10 times that of a magnitude 4 quake and 100 times that of a magnitude 3 quake.Each scale has their benefits. As mentioned above, the Mercalli Intensity scale is based on how much damage someone would see. This is relative though because some places have strong building codes, and the rock material underneath will impact ground shaking without changing the energy released at the focus. With the Richter scale, a single sharp jolt measures higher than a very long intense earthquake that releases more energy. The moment magnitude scale more accurately reflects the energy released and the damage caused. Today, most seismologists now use the moment magnitude scale.

Annual Earthquakes



In a single year, on average, more than 900,000 earthquakes are recorded and 150,000 of them are strong enough to be felt. Each year about 18 earthquakes are major with a Richter magnitude of 7.0 to 7.9, and on average one earthquake has a magnitude of 8 to 8.9.

Magnitude 9 earthquakes are rare. The United States Geological Survey lists five since 1900 (see table below). All but the Great Indian Ocean Earthquake of 2004 occurred somewhere around the Pacific Ocean basin.

Location	Year	Magnitude
Valdivia, Chile	1960	9.5
Prince Williams Sound, Alaska	1964	9.2
Great Indian Ocean Earthquake	2004	9.1
Kamchatka, Alaska	1952	9.0
Tohoku, Japan	2011	9.0

Earthquake Prediction

Scientists are a long way from being able to predict earthquakes. A good prediction must be accurate as to where an earthquake will occur, when it will occur, and at what magnitude it will be so that people can evacuate. An unnecessary evacuation is expensive and causes people not to believe authorities the next time an evacuation is ordered.

Where an earthquake will occur is the easiest feature to predict. Scientists know that earthquakes take place at plate boundaries and tend to happen where they've occurred before. Earthquake-prone communities should always be prepared for an earthquake. These communities can implement building codes to make structures earthquake safe.

When an earthquake will occur is much more difficult to predict. Since stress on a fault builds up at the same rate over time, earthquakes should occur at regular intervals. But so far scientists cannot predict when quakes will occur even to within a few years. Signs sometimes come before a large earthquake. Small quakes, called **foreshocks**, sometimes occur a few seconds to a few weeks before a major quake. However, many earthquakes do not have foreshocks and small earthquakes are not necessarily followed by a large earthquake. Often, the rocks around a fault will dilate as **microfractures** form.

Ground tilting, caused by the buildup of stress in the rocks, may precede a large earthquake, but not always. Water levels in wells fluctuate as water moves into or out of fractures before an earthquake. This is also an uncertain predictor of large earthquakes. The relative arrival times of P-waves and S-waves also decreases just before an earthquake occurs.


Folklore tells of animals behaving erratically just before an earthquake. Mostly these anecdotes are told after the earthquake. If indeed animals sense danger from earthquakes or tsunami, scientists do not know what it is they could be sensing, but they would like to find out.

The geology of California underlies the state's wealth of natural resources as well as its natural hazards. This video explores the enormous diversity of California's geology: http://www.youtube.com/watch?v=QzdBx9zL0ZY(57:50).




DAMAGE FROM EARTHQUAKES

Earthquakes are natural disasters that cause enormous amounts of damage, second only to hurricanes. Earthquake-safe construction techniques, securing heavy objects, and preparing an emergency kit are among the precautions people can take to minimize damage.Earthquakes kill people and cause property damage. However, the ground shaking almost never kills people, and the ground does not swallow someone up. The damage depends somewhat on the earthquake size but mostly on the quality of structures. Structures falling on people injure and kill them. More damage is done and more people are killed by the fires that follow an earthquake than the earthquake itself.



There are a variety of ways earthquakes become deadly. Probably the greatest influence is **population density**. The magnitude 9.2 Great Alaska Earthquake, near Anchorage, of 1964 resulted in only 131 deaths. At the time few people lived in the area. Oddly enough, size doesn't matter. Only about 2,000 people died in the 1960 Great Chilean earthquake, the largest earthquake ever recorded. The Indian Ocean earthquake of 2004 was one of the largest ever, but most of the 230,000 fatalities were caused by the tsunami, not the earthquake itself.

Ground type is an important factor in earthquake damage. Solid bedrock vibrates less than soft sediments so there is less damage on bedrock. Sediments that are saturated with water under go **liquefaction** and become like quicksand. This will have dangerous implications for Salt Lake City and the Wasatch Front in Utah. Much of the Wasatch Front is loose soil, left over from the remnants of Lake Bonneville. Along the middle of the two mountain ranges, between the Wasatch Mountains and the Oquirrh Mountains, is the Jordan River which flows from Utah Lake northward to the Great Salt Lake. Near the river and surrounding floodplain, the water table is near the surface. When the ground shakes, the water near the surface shifts upward and over-saturates the soil making it extremely unstable. Earthquake effects on buildings are seen in this animation.



EARTHQUAKE-SAFE STRUCTURES

Construction is a large factor in what happens during an earthquake. For example, many more people died in the 1988 Armenia earthquake where people live in mud houses than in the 1989 earthquake in Loma Prieta. Most buildings in California's earthquake country are designed to be earthquake safe.The key to earthquake safety are the structures we live in. It is often said that earthquakes don't kill people, it's the buildings that kill people. But the reason why not all buildings are not built to withstand earthquakes is cost. More sturdy structures are much more expensive to build, so communities must weigh how great the hazard is, what different building strategies cost, and make an informed decision. But this is the crucial factor in earthquake safety. Let's compare Salt Lake City, UT and Port-au-Prince, Haiti. When Salt Lake City has it's expected 7.0 magnitude earthquake, worst case scenario is that 3,000-4,000 will die. But on January 12, 2010, Haiti experienced a 7.0 magnitude earthquake that killed nearly 300,000 people. The difference – building codes.Skyscrapers and other large structures built on soft ground must be anchored to bedrock, even if it lies hundreds of meters below the ground surface. Larger buildings must sway, but not so much that they touch nearby buildings. Counterweights and diagonal steel beams are used to hold down sway. Large buildings can also be placed on rollers so that they move with the ground.Earthquake prone areas should have building codes that require the use of correct building materials. Houses should also be built with wood and steel rather than brick and stone because they need to be able to bend and sway. New buildings should be built on layers of steel and rubber to absorb the shock of the waves. Connections, such as where the walls meet the foundation, must be made strong to withstand the shaking. In a multi-story building, the first story must be well supported.To make older buildings more earthquake safe, retrofitting with steel or wood can reinforce a building's structure and its connections. Elevated freeways and bridges can also be retrofitted so that they do not collapse.

Fires often cause more damage than the earthquake. Fires start because seismic waves rupture gas and electrical lines, and breaks in water mains make it difficult to fight the fires. Builders zigzag pipes so that they bend and flex when the ground shakes. In San Francisco, water and gas pipelines are separated by valves so that areas can be isolated if one segment breaks.

HUMAN-INDUCED EARTHQUAKES

Can humans create earthquakes? Maybe not intentionally, but the answer is yes and here is why. If a water reservoir is built on top of an active fault line, the water may actually lubricate the fault and weaken the stress built up within it. This may either create a series of small earthquakes or potentially create a large earthquake. Also the shear weight of the reservoir's water can weaken the bedrock causing it to fracture. Then the obvious concern is if the dam fails. Earthquakes can also be generated if humans inject other fluids into a fault such as sewage or chemical waste. Finally, nuclear explosions can trigger earthquakes. In fact, one way to determine if a nation has tested a nuclear bomb is by monitoring the earthquakes and energy released by the explosion.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=58

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-5-tectonic-forces.html. **License**: *CC BY: Attribution*

This page titled 4.4: Earthquakes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

5: Volcanoes

5.1: Volcanic Activity

5.2: Types of Volcanoes

5.3: Volcanic Landforms and Geothermal Activity

5.4: Volcanic Eruptions

This page titled 5: Volcanoes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

5.1: Volcanic Activity

Everybody has heard of volcanoes—the angry Earth spewing up its wrath, the sudden explosion of a distant mountain top, and lava that slithers ominously toward villages. Volcanoes are fantastic displays of the power of the Earth. But what actually is a volcano? How and where are they formed? Why do some places have a history of volcanic activity? Volcanoes explain a key piece of the Earth's geologic puzzle (Figure 8.1).



Figure 1: Mount Merapi, Indonesia

Lesson Objectives

- Describe places where volcanoes occur.
- Describe what volcanic hot spots are and where they occur.

Where Volcanoes Occur

Because volcanoes are vents for magma, it makes sense that volcanoes would be formed above underground magma chambers. If you recall, magma is molten rock that has been heated because of high temperatures and pressures beneath the Earth's crust. This pressure mostly occurs where the tectonic plates meet and subduct. Look at the map of tectonic plates in Figure 8.2.




Figure 8.2: Tectonic plates. Some of the largest tectonic plates meet along the coasts of Asia and North America. Compare this map to the map in Figure 8.3.

So, volcanic activity tends to occur along subduction plate boundaries, where one plate slides underneath another. The edges of the Pacific Plate make up a long subduction boundary. There are a huge number of earthquakes along these boundaries, because these are regions where the plates are colliding. For the same reason, the majority of the volcanic activity on the Earth also occurs along these convergent boundaries. This is called the ***Pacific Ring of Fire*** where over 75% of the world's volcanoes are found. The Cascade Range of volcanoes runs through southwestern Canada and the Pacific Northwest of the United States. These volcanoes are the result of subduction of the Juan de Fuca plate beneath the North American plate. (Figure 8.3).




Figure 8.3: The Pacific Ring of Fire is where the majority of the volcanic activity on the Earth occurs.



Figure 8.4: The Mid-Atlantic Ridge is formed by divergent tectonic plates.

Of course, this is not the only area where volcanoes occur. Beneath the ocean, there are also divergent boundaries, where tectonic plates are pulling away from each other. As the plates pull away from each other, they create a deep canyon or fissure on the sea floor through which molten rock escapes. Mid-ocean ridges, like the Mid-Atlantic ridge, form here as lava flows out through the fissure (Figure 8.4). Submarine volcanoes can also form on the ocean floor. At times these volcanoes can grow to create islands above the water's surface. This explains how Surtsey, a small island of Iceland formed (Figure 8.5).



Figure 8.5: A volcanic eruption in Iceland.

Volcanic Hot Spots

Although most volcanic activity on Earth occurs at plate boundaries, there are some volcanically active spots that are in the middle of a tectonic plate. These areas are called **hot spots**. The islands of Hawaii formed over a hot spot and are not located on the Pacific

Ring of Fire (Figure 8.6). The Hawaiian islands are the exposed peaks of a great chain of volcanoes that were formed over millions of years. The islands are thought to lie directly above a column of hot rock called a **mantle plume**. Mantle plumes are more or less fixed in place and continuously bring magma up form the mantle towards the crust. As the tectonic plates move above them, they leave a trail of volcanic activity, which forms island chains like Hawaii. Scientists believe there are about 50 hot spots on the Earth. Other hot spots include Yellowstone and the Galapagos Islands.




Figure 8.6: Hawaii is a volcanic hot spot. It was formed by volcanic activity fed by mantle plumes.

Don't confuse hot spot volcanoes with islands that are formed by plate tectonics like the Aleutian island chain in Alaska. These long lines of volcanoes form as the edge of a subducted plate melts, producing magma which rises to the surface along the edge of the plate. These volcanic mountains will all be about the same age. When islands form over a hot spot, the youngest island is over the hot spot. As you move along the island chain, each island further from the hot spot will be older than the one before it.

Lesson Summary

- Volcanoes occur most often along plate boundaries.
- Convergent plate boundaries where oceanic crust is subducted form many of the volcanoes found on Earth.
- Divergent plate boundaries produce huge mountain ranges under water in every ocean basin.
- Volcanoes like those that form the islands of Hawaii, form over areas called hot spots.

Review Questions

1. Why are there volcanoes along the west coast of the United States?
2. Why do volcanoes occur where tectonic plates pull apart or diverge?
3. Explain how the Pacific Ring of Fire got its name.
4. What is a mantle plume?
5. Suppose a new volcano suddenly formed in the middle of the United States. How might you explain what caused this volcano?
6. Volcanoes have been found on Venus, Mars, and even Jupiter's moon Io. What do you think this indicates to planetary geologists?

Vocabulary

fissure

A long crack into the Earth's surface, from which lava erupts onto the surface.

hot spot

A fixed region of hot magma that rises through the mantle and creates volcanoes on the Earth's surface. A hot spot is above a mantle plume.

lava

Molten rock that has been erupted onto the Earth's surface. Magma becomes lava once it emerges on the surface.

magma

Molten rock found under the Earth's surface.

mantle plume

A column of very hot rock that rises up through the mantle. Mantle plumes will form hot spots if they make it all the way up to the surface of the Earth's crust.

Points to Consider

- When you look at the map of tectonic plates (Figure 8.3), what areas besides the Pacific Ring of Fire would you expect to have volcanic activity?
- Some volcanoes are extinct; they are no longer active and will probably never be again. What do you think causes a volcano to become extinct?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Volcanic_Activity. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 5.1: Volcanic Activity is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

5.2: Types of Volcanoes

When most people think of volcanoes, they think of a tall mountain with a crater on the top, maybe a little snow at the summit and some trees scattered around the base. There are many volcanoes like this, but volcanoes exist in many other forms as well. Each type of volcano has characteristic features that distinguish it from other types. Volcanoes differ in appearance because of the composition of their magma and the processes that originally created them.

The tall cone shape you usually think of when you think of a volcano describes a composite volcano, one common form of volcanoes. Other types of volcanoes include the shield volcano, the cinder cone, and the supervolcano.

Lesson Objectives

- Describe the basic shapes of volcanoes.
- Compare the features of volcanoes.
- Describe the stages in the formation of volcanoes.

Strato-volcano or Composite Volcano

The picture below shows Mt. Fuji, a classic example of the **strato-volcano** also known as a **composite volcano** (Figure 8.21). This is the type of volcano many people think of when they imagine volcanoes. These volcanoes have broad bases and sides that get steeper and steeper as you get closer to the top. They frequently have a large crater at the top created during its last eruption.



Figure 8.21: Mt. Fuji is a dormant composite volcano that is the highest mountain in Japan.



Figure 8.22: A composite or stratovolcano is created by many levels of alternating materials.

Composite volcanoes are also called stratovolcanoes because of the alternating layers, or **strata**, of which they are made (Figure 8.22). The *magma that creates stratovolcanoes tends to be more viscous, or thick*. Viscous lava creates greater pressure which, in turn, tends to create explosive eruptions. In addition, the viscous lava cannot travel far down the sides of the volcano before it solidifies. This viscous lava thus creates steep sides on stratovolcanoes.

When a stratovolcano erupts, it ejects a great deal of pyroclastic material into the air, which then settles back down on the Earth. After an initial explosion, lava then flows from the volcano creating a second layer of material. As these layers solidify, they create alternating levels, or strata, of material. Ash from the volcanic eruption is also present between the lava layers along the edge of the

volcano. Composite volcanoes are common along the Pacific Ring of Fire and other major tectonic plate boundaries where the presence of water in the magma chamber creates explosive eruptions.

Shield Volcanoes

Shield volcanoes get their name from their shape—a huge shield laid on its side. Figure 8.23 shows the Mauna Loa Volcano. You can see that shield volcanoes do not have the steep mountainous sides of composite volcanoes. They have a very wide base and are much flatter on the top than composite volcano. Although they are not steep, they may be very large. The Mauna Loa Volcano has a diameter of over 112 kilometers (70 miles) and forms a significant part of the island of Hawaii. The Mauna Kea Volcano, also in Hawaii, is another shield volcano that is over ten kilometers (6 miles) high from its base below sea level to its peak.



Figure 8.23: The Mauna Loa Volcano in Hawaii is the largest shield volcano on Earth.




Figure 8.24: A shield volcano is built by layers of more fluid lava that spreads out over broad areas.

Shield volcanoes are more common at spreading centers or volcanic hot spots in the middle of tectonic plates (Figure 8.24). The magma that creates shield volcanoes is less viscous, so it flows much more easily. For this reason, the eruptions of shield volcanoes are non-explosive. In addition, the less viscous lava spreads out more, which makes shield volcanoes much larger and flatter than stratovolcanoes. Although shield volcanoes are built by many layers over time, the composition of the layers do not alternate between ash and lava, as they do in stratovolcanoes.

Cinder Cones



Figure 8.25: Paricutín erupting in 1943, when it first formed. Cinder cones like this one rarely reach even 300 meters high.

Cinder cones are both the most common type of volcano and also the smallest. The cinder cone resembles a composite volcano but on a much smaller scale. They rarely reach even 300 meters in height but have even steeper sides than a composite volcano. They usually have a crater at the summit. Cinder cones are composed of small fragments of rock piled on top of one another. These volcanoes usually do not produce streams of lava.

In 1943, a farmer in Mexico witnessed the first eruption of a cinder cone in his field (Figure 8.25). Within a year, the cinder cone Paricutín grew to 336 meters high. By 1952, it grew to a peak of 424 meters tall, and then stopped erupting. This rapid growth and single eruption cycle is characteristic of cinder cones. For this reason, cinder cones do not reach the sizes of stratovolcanoes or shield volcanoes. Often, cinder cones appear near larger volcanoes, but they also may be found away from all other volcanoes, as was the case with Paricutín. The exact composition of a cinder cone depends on the composition of the lava ejected from the Earth.

Supervolcanoes



Figure 8.26: The caldera at Santorini in Greece is so large that its circular shape can only be seen by satellite.

In certain areas of the world, huge **calderas** have been found to be the remains of volcanic eruptions of enormous scale (Figure 8.26). These calderas are volcanic features that are formed by the collapse of a huge amount of land due to the powerful eruptions. Caldera comes from Latin word, meaning cauldron. Calderas are generally circular shaped geographic formations like the picture in figure 6. These are not singular mountains but entire geographical areas. Yellowstone National Park in Wyoming is another caldera that has blown about a hundred times in the last 16 million years.

Supervolcanoes represent the most dangerous type of volcano. An eruption from a supervolcano could change life on Earth as we know it for many years. Supervolcanoes were not even accepted in volcanology until this millennium. Many supervolcano eruptions are thought to have occurred, the most recent in New Zealand less than 2000 years ago. That explosion was thought to have ejected about 100 cubic kilometers of material. A supervolcano eruption near what is now Colorado was thought to have let loose over 5,000 cubic kilometers of material millions of years ago. In comparison, the Mt. Saint Helens eruption ejected about 1 cubic kilometer of material.

The eruptions from supervolcanoes can be so large that the ash ejected into the air blocks the Sun and lowers the temperature on the entire planet. The lowered temperatures caused by these eruptions is called a volcanic winter. A supervolcano eruption at Lake Toba in northern Sumatra may have annihilated about 60% of the world's human population about 75,000 years ago. One can only imagine how such a huge eruption would change the world in modern times.

The largest supervolcano in North America is the Yellowstone caldera, which had three super eruptions at 2.1 million, 1.3 million and 640,000 years ago, and much more recent smaller (but still enormous) eruptions. Long Valley caldera, south of Mono Lake in California, is the second largest supervolcano in North America, erupting extremely hot and explosive rhyolite around 700,000 years ago. An earthquake swarm in 1980 alerted geologists to the possibility of another eruption in the future, but the timing of such an event is unknown.

Supervolcanoes are a fairly new idea so the exact cause of supervolcano eruptions is still debated. However, scientists believe that an entire and very large magma chamber erupts in a catastrophic explosion. This enormous eruption creates a huge hole or caldera where the surface area collapses.

Lesson Summary

- Composite cones, shield volcanoes, cinder cones and supervolcanoes are some of the types of volcanoes formed.
- Composite cones are tall, cone shaped volcanoes that produce explosive eruptions.
- Shield volcanoes form very large, gently sloped volcanoes with a wide base.
- Cinder cones are the smallest volcanic landform. They are formed from accumulation of many small fragments of ejected material.
- A caldera forms when an explosive eruption leaves a large crater when the mountain blows apart.
- Supervolcanoes are tremendously devastating types of volcanoes that could destroy large areas when they erupt.

Review Questions

1. Rank the four types of volcanoes in order from smallest to largest in diameter.
2. What factor is most important in determining the type of volcano formed in a given area?
3. Which type of volcano is most common?
4. Why do you think that cinder cones are short-lived?
5. If supervolcanoes are so big, why do you think it took so long for scientists to discover them?

Vocabulary

caldera

Circular-shaped geographic features formed from a massive eruption of an ancient volcano, and the subsequent collapse of the volcano back into the ground.

cinder cone

A smaller volcano that grows rapidly but only erupts over a short period of the time. Cinder cones are composed of small rock fragments piled on top of one another. They rarely are more than 300 m in height.

composite volcano

A volcano with a broad base, steep sides, and often a crater at the top. The volcano is composed of alternating layers of ash and lava flows. Also called a stratovolcano.

strata

Layers of rock that are similar in composition to one another.

supervolcano

A massive volcanic eruption that is rare but incredibly powerful. Thousands of cubic kilometers of matter can be ejected, and the dust and ash from their eruption can cool the world's climate for years.

Points to Consider

- Composite volcanoes and volcanic cones usually have craters on the top. Why are the craters not always circular, but sometimes "U" or horseshoe-shaped?
- A shield volcano is relatively flat, and a composite volcano is relatively steep because of the type of magma that creates them. What process might create a volcano that is more steep than a shield volcano but not as steep as a composite volcano?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Types_of_Volcanoes. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 5.2: Types of Volcanoes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

5.3: Volcanic Landforms and Geothermal Activity

As you know, magma is molten rock found beneath the Earth's surface. Sometimes, it appears at the surface of the Earth as lava after moving through a volcano. At other times, magma does not come to the surface, but stays underground. In both cases, the magma eventually solidifies and the resulting rocks and formations are igneous. The rocks that solidify beneath the ground are called **intrusive or plutonic** rocks, while those that solidify above the surface are called **extrusive or volcanic** rocks. Extrusive rocks are sometimes small rocks that you can hold in your hand. At other times, entire landforms are created when lava flows onto the surface. Intrusive rocks do not always remain hidden below the surface. They can appear on the surface when rocks that once covered them are eroded away, exposing the intrusive igneous rock. Hot springs and geysers are some more examples of surface features related to igneous rock.

Lesson Objectives

- List and describe landforms created by lava.
- Explain how magma creates different landforms.
- Describe the processes that create hot springs and geysers.

Landforms from Lava

The most obvious landforms created by lava are volcanoes. Volcanoes, of course, are the places where lava comes to the surface. As already discussed, volcanoes come in many forms, most commonly as cinder cones, stratovolcanoes, and shield volcanoes. However, lava can create other notable landforms, as described below.

Lava Domes

When lava is fairly viscous, it is thick and flows slowly. Although it might not be so viscous that it causes an explosive eruption, it can create a large sort of round "blob" or a lava dome. Because it is so thick, it does not flow far from the vent from which it came. In fact, lava flows often make mounds right in the middle of craters at the top of volcanoes (Figure 8.27 and Figure 8.28).



Figure 8.27: Lava domes are large, round landforms created by thick lava that does not travel far from the vent.



Figure 8.28: Sometimes lava domes are formed in the crater of composite volcanoes. This lava dome is forming in the crater of Mt. St. Helens in Washington State.

Plateaus

A **lava plateau** is caused by a large amount of less viscous lava that flows over a large area. When it solidifies, it creates a large, flat surface of igneous rock. Some plateaus are huge, like the Columbia Plateau in Washington, Oregon, and Idaho that covers over 161,000 square kilometers (63,000 square miles).

Landforms from Magma

Of course, not all magma reaches the surface. Sometimes magma stays beneath the ground where it solidifies. These formations are called **intrusions** (Figure 8.30). Because they form underground, they only become land formations if they arrive at the surface of the Earth. This occurs because of weathering, erosion and plate tectonics. In other words, when tectonic plates collide, they create mountain ranges where one plate is lifted in a subduction zone. This lifting (which occurs at a rate of centimeters per year) and subsequent erosion can eventually uncover intrusive rock formations. As erosion removes the topmost rocks and soil, solidified magma is exposed in the same form in which it cooled and solidified many years before.



Figure 8.30: Devil's Tower in Wyoming is a huge rock formation that was once magma that cooled within a volcano. It rises to nearly 400 meters from its base.

Hot Springs and Geysers

Beneath the surface of the Earth, water works its way through porous rocks or soil. Most caves, for example, are results of water's erosion of the ground. At times, that water crosses paths with volcanic activity. The same heat that melts rock into lava heats the water beneath the surface. If the water makes its way to the surface, it may emerge as either a hot spring or a geyser.

Hot Springs

When that water comes to the surface under regular pressure, it creates a **hot spring** (Figure 8.31). A hot spring is a crack in the Earth through which water reaches the surface, after being heated below the ground. Many people disagree on the exact definition of a hot spring. However, everyone agrees that the water's temperature is higher than normal. The water in hot springs can even reach temperatures in the hundreds of degrees Celsius beneath the surface. Most hot springs do not reach those great temperatures. In fact, many hot springs are used by people as natural hot tubs. Many people believe that hot springs hold curative properties. Hot springs are found all over the world, even in Antarctica!



Figure 8.31: Even some animals enjoy relaxing in nature's hot tubs.

Geysers



Figure 8.32: Old Faithful Geyser in Yellowstone National Park during an eruption.

Like hot springs, **geysers** are created by water that is heated beneath the Earth's surface. When water is both superheated by magma and flows through a narrow passageway underground, the environment becomes ideal for a geyser. The narrow passageway traps the heated water underground, where heat and pressure continue to build. Sooner or later, the pressure grows so great that the superheated water bursts out onto the surface. This explosion is called a geyser. There are only a few areas in the world where the conditions are right for the formation of geysers. About 1,000 geysers exist worldwide and about half of those are found in the United States of America. Perhaps the most famous geyser is Old Faithful, which erupts every 60 to 70 minutes in a plume of hot water nearly 60 meters in the air. It is rare for a geyser to erupt regularly, which contributes to Old Faithful's fame (Figure 8.32).

Lesson Summary

- Very thick lava that doesn't travel very far can produce lava domes at or near the Earth's surface or even within a volcano.
- Lava plateaus form from large lava flows that spread out over large areas.
- Many islands are formed from volcanoes.
- Magma can also cool and crystallize below the Earth's surface forming igneous intrusions.
- When magma heats groundwater, it can form hot springs and geysers.

Review Questions

1. What are three different landforms created by lava?
2. What is the major difference between hot springs and geysers?
3. The geyser called Old Faithful has been erupting for perhaps hundreds of years. One day, it could stop. Why might geysers completely stop erupting?
4. After earthquakes, hot springs sometimes stop bubbling, and new hot springs form. Why might this be?

Vocabulary

lava dome

A dome-shaped plug of viscous lava that cools near the vent of a volcano.

intrusion

A rock mass formed by magma solidifying underground.

hot spring

A stream of hot water that flows out of the ground continuously.

geyser

A fountain of hot water and steam that shoots into the air at either regular or random intervals. The water in a geyser encounters some sort of constriction underground, which forces pressure to build up in the water, eventually causing the geyser to erupt.

Points to Consider

- What might the Earth look like if there were no tectonic plates? Can you think of any planets or satellites (moons) that may not have tectonic plates? How is their surface different from that of the Earth?
- What kind of land formations have you seen that may have been created by volcanic activity? Were the formations made of extrusive or intrusive rock?
- Water is not the only material that can be ejected from geysers and hot springs. Consider the composition of the Earth's crust. What other materials might be ejected from geysers and hot springs?

This page titled 5.3: Volcanic Landforms and Geothermal Activity is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

5.4: Volcanic Eruptions

In 1980, Mount St. Helens erupted in one of the most deadly and costly volcanic eruptions in the United States ever. The eruption was particularly deadly since Mount St. Helens, one of the Cascade Range, is in a populated area between Portland, Oregon and Seattle, Washington. The eruption killed 57 people, destroyed 250 homes, and swept away 47 bridges. The elevation of the volcano dropped by over 400 meters (1,300 feet) because of the immense explosion created by the eruption. Today Mt. St. Helens is still active (Figure 8.8). The volcano now has a horseshoe-shaped crater that is 1.5 km (nearly one mile) across. Within the crater, a new lava dome has formed. How did this eruption occur? Why aren't all volcanoes explosive like Mt. St. Helens? Why did so many people perish if we knew that it was going to erupt? The study of volcanoes has many questions still unanswered. However, scientists have studied volcanoes for many years and are piecing together evidence that explains these powerful geologic phenomena.



Figure 8.8: Mount St. Helens, Washington, two years after its eruption.

Lesson Objectives

- Explain how volcanoes erupt.
- Describe and compare the types of volcanic eruptions.
- Distinguish between different types of lava and understand the difference between magma and lava.
- Describe a method for predicting volcanic eruptions.

How Volcanoes Erupt

All volcanoes share the same basic features. The magma collects in magma chambers that can be 160 kilometers (100 miles) beneath the surface. As the rock heats, it expands, which creates even more pressure. As a result, the magma seeks a way out pushing toward the surface, the magma seeps through cracks in the Earth's crust called vents. Eventually, the magma reaches the surface; when it comes out, we call it an eruption. The word **eruption** is used in other contexts, as well. An eruption can be an outburst or explosion, a violent and sudden occurrence, like when a crowd erupts in anger. But an eruption can also be a spreading of something like a rash on your skin, gradual and relatively calm. These two definitions are similar to the two kinds of eruptions that we see in volcanoes.

Types of Eruptions

Every geological formation is unique. Their composition and construction depend on so many factors, that it would be impossible for two formations to be exactly alike. In the same way, each volcano and its eruptions are unique. However, we tend to see two major kinds of eruptions. We talked about eruption to mean both a violent explosion or a sort of silent spreading. These are the two types of volcanic eruptions that we see–explosive and non-explosive eruptions. When we think of volcanic eruptions, we often think of huge clouds of volcanic ash ejected high into the atmosphere and then thick rivers of red lava snaking down the mountainside. In reality, these two phenomena rarely occur in the same volcano. Volcanic eruptions tend to be one or the other.

Explosive Eruptions

Imagine the devastation and force caused by the atom bomb dropped on Nagasaki at the end of World War II in which over 40,000 people died. Now imagine an explosion 10,000 times as powerful. Explosive volcanic eruptions can be that powerful (Figure 8.9). As hot magma beneath the surface interacts with water, gases accumulate and the magma pressure builds up. This pressure grows and grows until these dissolved gases cause it to burst in an enormous explosion.



Figure 8.9: An explosive eruption from the Mayon Volcano in the Philippines in 1984.

This great explosion takes with it the magma and volcanic gases, which can shoot many kilometers into the sky and forms a mushroom cloud, similar to that formed by a nuclear explosion (Figure 8.10). The debris travels up into the air at very high speeds and cools in the atmosphere to form solid particles called **pyroclasts**. Some of these particles can stay in the atmosphere for years, which can disrupt weather patterns and affect the temperature of the Earth. The rest of the debris comes falling back to Earth where it rains down for kilometers and kilometers around.




Figure 8.10: Explosive eruption of Mt. Redoubt in Alaska, 1989. This huge mushroom cloud reached 45,000 feet and caught a Boeing 747 in its plume.

Sometimes secondary explosions occur that are even greater than the first. Additionally, volcanic gases like water vapor, carbon dioxide, sulfur dioxide, hydrogen sulfide, and hydrogen chloride can form poisonous and invisible clouds that roam about the atmosphere. These gases contribute to environmental problems like acid rain and ozone destruction, and can actually cool the Earth's atmosphere.

In the Cascade Range, the explosive eruption of Mount St. Helens was preceded by the eruption of Lassen Peak, one of the three Cascade Volcanoes in northern California. On May 22, 1915, an explosive eruption sent a column of ash and gas 30,000 feet into the air and triggered a high-speed pyroclastic flow, which melted snow and created a lahar. Lassen continues to have geothermal activity and could erupt explosively again. Mt. Shasta erupts every 600 to 800 years. An eruption would most likely to create a large pyroclastic flow, and perhaps a lahar. However, the volcano could explode like Mt. Mazama, which blew itself in an eruption about 42 times more powerful than Mount St. Helens in 1980, to create Crater Lake.

Non-explosive Eruptions

A second type of volcanic eruption is a non-explosive or **effusive eruption** (Figure 8.11). Because the composition of magma is different in different volcanoes, the properties of the lava are different. In effusive eruptions, lava flows are relatively calm and do not explode out of the volcano. As a result, people generally have a great deal of warning before lava reaches them, so non-explosive eruptions are much less deadly. That does not keep them from being destructive, however. Even when we know that a lava flow is approaching, there are few ways of stopping it, given the huge quantity and temperature of lava.



Magma and Lava

Volcanoes wouldn't be nearly as interesting without the great explosions they create and the glowing red rivers of lava. All igneous rock comes from magma or lava. The next time you go hiking near a volcanic zone, you might try to identify the types of lava that the volcano erupted, based on the types of igneous rocks you find.



Figure 8.12: When lava flows readily, pressure does not build up so great explosions do not occur.

Magma

Deep beneath the Earth, magma forms as the first stage in creating a volcano. This occurs because rock below the surface is subjected to great amounts of pressure from gravity. The decay of radioactive materials generates additional heat. The substantial heat and pressure melt the rock below the surface to form a taffy-like substance. You may have seen a candle that has been left out in the hot sun too long. It becomes softer and more like a liquid. As the molecules absorb heat, they begin to slide past one another becoming more fluid. A similar process occurs with magma. However, different substances melt at different temperatures. For that reason, the temperature at which rocks melt depends on the specific types of rocks. The Earth's crust and mantle are made of many substances so the temperature required to create magma varies. Most magmas are formed between 600°C and 1300°C (Figure 8.13).




Figure 8.13: Cutaway of the Earth. The melting of rock in the crust and upper mantle create magma.

Melted rock or magma can be found in **magma chambers** beneath the Earth. Since the magma chambers are so far beneath the Earth's surface, it is difficult for scientists to study them. Scientists know that magma chambers are created where the heat and pressure are greatest. When tectonic plates collide and rub against each other, magma is formed there. That is how the Pacific Ring of Fire was created. We also know there are volcanoes far away from plate boundaries, so we know there are magma chambers in these areas as well. Magma chambers can be found where there are mantle plumes or hot spots.

Just how or why these hot spots are created isn't exactly known. However, because different substances melt at different temperatures, the creation of magma depends on what substances make it up—its composition. Just like the flavor of a cake depends on the ingredients that you put in it, the behavior of magma and lava depends on its composition. Certain melted rocks act in certain ways. So when the magma becomes lava, not all lava acts the same.

Lava



Figure 8.14: Honey flows slowly; it is more viscous than water.

Once magma reaches the surface, it becomes lava. Consider different liquids that you might see in your house—honey and a bottle of cola, for example. You might agree that the two liquids are different in many regards. They taste different, have different colors, have different gases in them, and they flow differently. In fact, honey is a liquid that resists flowing, whereas cola flows easily. Honey has a higher **viscosity** than the cola; it resists flowing (Figure 8.14). Cola has a low viscosity because it flows easily. One of the major differences in different types of lava is their viscosity.

A highly viscous lava is one that doesn't tend to flow easily. It tends to stay in place. Lavas with high silica contents tend to be more viscous. Since it is so resistant to moving, it clogs the vents in a volcano. The pressure becomes greater and greater until the volcano finally explodes. This type of lava is found in explosive eruptions. It also tends to trap a lot of gas. When the gas is released, it makes the eruption more explosive. Most of this lava is shot up into the air where it hardens and becomes solid rock. This molten rock that solidifies in the air is known as **pyroclastic material**. In an igneous rock like pumice, small holes in the solid rock show where gas bubbles were when the rock was still liquid lava.

Low-viscosity lava slides or flows down mountainsides. There is more than one type of low-viscosity lava. The differences between them come from the lavas' different composition and different spots where they come to the surface. The type of igneous formations formed depends on which type of lava it is. The three major categories are a'a, pahoehoe, and pillow lava.

A'a Lava

A'a lava is the more viscous of the non-explosive lavas (Figure 8.15). This lava forms a thick and brittle crust which is torn into rough and jagged pieces. The solidified surface is jagged and sharp. It can spread over large areas as the lava continues to flow underneath.



Figure 8.15: A'a lava flow.

Pāhoehoe Lava

Pāhoehoe lava is less viscous than a'a lava, and flows more readily. Its surface looks more wrinkly and smooth than the jagged a'a lava. Pāhoehoe lava flows in a series of lobes or rounded areas that form strange twisted shapes and natural rock sculptures (Figure 8.16). Pāhoehoe lava can also form lava tubes beneath the ground (Figure 8.17).



Figure 8.16: Pāhoehoe lava.



Figure 8.17: The Thurston Lava Tube in Hawaii Volcanoes National Park.

Pillow Lava

Pillow lava is lava that comes out from volcanic vents underwater (Figure 8.18). When it comes out underwater, it cools down very quickly and forms roughly spherical rocks that resemble pillows, from which more lava leaks and creates more pillows. Pillow lava is particularly common along underwater spreading centers.



Figure 8.18: Pillow lava.

Predicting Volcanic Eruptions

Volcanic eruptions can be devastating, particularly to the people who are closer to volcanoes. As meteorologists attempt to predict, or forecast, hurricanes and tornadoes, so too do vulcanologist attempt to forecast volcanic eruptions. Although predicting volcanic eruptions is far from perfect, many pieces of evidence can indicate that a volcano is about to erupt. Some of those factors are hard to measure, contributing to the difficulty in predicting eruptions.

History of Volcanic Activities

One important factor in predicting eruptions is a volcano's history. That is, we consider how long since it has erupted and the time span between its previous eruptions. Volcanoes are categorized into three subdivisions—active, dormant, and extinct. An **active volcano** is one that is currently erupting or shows signs of erupting in the near future. A **dormant volcano** no longer shows signs of activity, but has erupted in recent history (Figure 8.19). Finally, an extinct volcano is one that has not erupted in recent history and will probably not erupt again in the future. Both active and dormant volcanoes are heavily monitored because even dormant volcanoes could suddenly show signs of activity.



Figure 8.19: Vesuvius is a dormant volcano near the city of Naples. Although it shows no current signs of eruption, it could one day become active.

Earthquakes

As magma beneath a volcano pushes upward, it shakes the ground and causes earthquakes. Although earthquakes probably occur every day near a volcano, the quantity and size of the earthquakes increases before an eruption. In fact, a volcano that is about to erupt may produce a continuous string of earthquakes, as magma moving underground creates stress on the neighboring rocks. In order to measure these earthquakes, scientists use seismographs that record the length and strength of each earthquake.

Slope Deformation

All that magma and gas pushing upwards can make the ground or the volcano's slope begin to swell. Sometimes, ground swelling reveals huge changes in the shape of a volcano. Most cases of ground deformation are subtle, though, and can only be detected by tiltmeters, which are instruments that measure the angle of the slope of a volcano. Additionally, ground swelling may cause increased rock falls and landslides.

Gas Emissions

Oftentimes, gases are able to escape a volcano before magma reaches the surface in an eruption. So, scientists can measure gas output, or gas emissions, in vents on or around the volcano. Gases, like sulfur dioxide (SO_2), carbon dioxide (CO_2), hydrochloric acid (HCl) and even water vapor can be measured at the site or, in some cases, at a distance with satellites. The amounts of gases and their ratios are calculated to help predict eruptions.

Remote Monitoring

As mentioned, some gases can be monitored using satellite technology (Figure 8.20). Satellites are able to measure other factors, too, like temperature readings of particularly warm spots at a volcano site or areas where the volcano surface is changing. As our technology continues to improve, scientists are better able to detect changes accurately and safely.



Figure 8.20: An Earth-observation satellite before launch.

Although monitoring methods are getting better and better, it is still difficult to predict a volcanic eruption with certainty. No scientist or government agency wants to be considered alarmist by announcing that an eruption is going to occur and then it really doesn't. The cost and disruption to society of a large-scale evacuation would leave many people displeased and the scientists embarrassed. However, the possibility of saving lives and property most certainly makes the pursuit of eruption prediction a worthy cause.

Lesson Summary

- Volcanoes are produced when magma rises towards the Earth's surface because it is less dense than the surrounding rock.
- Volcanic eruptions can be non-explosive or explosive depending on the viscosity of the magma.
- Explosive type eruptions happen along the edges of continents and produce tremendous amounts of material ejected into the air.
- Non-explosive type eruptions mostly produce various types of lava, such as a'a, pāhoehoe and pillow lavas.
- Some signs that a volcano may soon erupt include earthquakes, surface bulging, gases emitted as well as other changes that can be monitored by scientists.

Review Questions

1. What are the two basic types of volcanic eruptions?
2. Several hundred years ago, a volcano erupted near the city of Pompeii. Archaeologists have found the remains of people embracing each other, suffocated by ash and rock that covered everything. What type of eruption must have this been?
3. What is pyroclastic material?
4. Name three liquids that have low viscosity and three that have high viscosity.
5. What is the difference between a magma chamber and a mantle plume?
6. The boiling point of water is 100°C. Why might water make an eruption more explosive?
7. What are three names for non-explosive lava?
8. What factors are considered in predicting volcanic eruptions?
9. Why is predicting volcanoes so important?
10. Given that astronomers are far away from the subjects they study, what evidence might they look for to determine the composition of a planet on which a volcano is found?

Vocabulary

active volcano

A volcano that is currently erupting or just about to erupt.

dormant volcano

A volcano that is not currently erupting, but that has erupted in the recorded past.

effusive eruption

A relatively gentle, non-explosive volcanic eruption.

eruption

The release of magma onto the Earth's surface. Usually an eruption is accompanied by the release of gases as well.

explosive eruption

A volcanic eruption that releases large amounts of gas, so that magma is violently thrown up into the air.

extinct volcano

A volcano that has not erupted in recorded history, and is considered unlikely to erupt again.

magma chamber

A region within Earth surrounded by solid rock and containing magma.

pyroclast

A rock made up of fragments of volcanic rock thrown into the air by volcanic eruptions.

viscosity

The "thickness" or "stickiness" of a liquid. The more viscous a liquid is, the harder it will be for the liquid to flow.

Points to Consider

- What types of evidence do you think would tell scientists whether an ancient volcanic eruption was explosive or non-explosive?
- Are all volcanoes shaped like tall mountains with a crater on the peak?
- What do you think is the origin of the names A'a and Pāhoehoe?
- Earthquakes do not always indicate that a volcano is going to erupt. What factors about an earthquake might indicate a relationship to a volcanic eruption?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Volcanic_Eruptions. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 5.4: Volcanic Eruptions is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

6: Surface Water

6.1: Earth’s Fresh Water
6.2: Streams and Rivers
6.3: Groundwater
6.4: Water Resources
6.5: Floods
6.6: Wetlands

This page titled 6: Surface Water is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.1: Earth's Fresh Water


LEARNING OUTCOMES

- **ESSENTIAL QUESTIONS**
  - How does precipitation affect the topology of the earth?
  - What natural disasters are caused by the water cycle?
  - Is water from a river or from a well more likely to be clean to drink?
  - Why is overuse of groundwater a big concern?
  - What policies might people put in place to conserve water levels in lakes and aquifers?

Introduction

Water is simply two atoms of hydrogen and one atom of oxygen bonded together. Despite its simplicity, water has remarkable properties. Water expands when it freezes, has high surface tension (because of the polar nature of the molecules, they tend to stick together), and others. Without water, life might not be able to exist on Earth and it certainly would not have the tremendous complexity and diversity that we see.

Distribution of Earth's Water




Earth's oceans contain 97% of the planet's water, so just 3% is fresh water, water with low concentrations of salts. Most fresh water is trapped as ice in the vast glaciers and ice sheets of Greenland. A storage location for water such as an ocean, glacier, pond, or even the atmosphere is known as a **reservoir**. A water molecule may pass through a reservoir very quickly or may remain for much longer. The amount of time a molecule stays in a reservoir is known as its residence time.

Earth's oceans contain 97% of the planet's water, so just 3% is fresh water, water with low concentrations of salts. Most fresh water is trapped as ice in the vast glaciers and ice sheets of Greenland. A storage location for water such as an ocean, glacier, pond, or even the atmosphere is known as a reservoir. A water molecule may pass through a reservoir very quickly or may remain for much longer. The amount of time a molecule stays in a reservoir is known as its residence time.

THREE STATES OF WATER

Because of the unique properties of water, water molecules can cycle through almost anywhere on Earth. The water molecule found in your glass of water today could have erupted from a volcano early in Earth history. In the intervening billions of years, the molecule probably spent time in a glacier or far below the ground. The molecule surely was high up in the atmosphere and maybe deep in the belly of a dinosaur. Where will that water molecule go next? Water is the only substance on Earth that is present in all three states of matter – as a solid, liquid or gas. Along with that, Earth is the only planet where water is present in all three states. Because of the ranges in temperature in specific locations around the planet, all three phases may be present in a single location or in a region. The three phases are solid (ice or snow), liquid (water), and gas (water vapor).



THE WATER CYCLE

Because Earth's water is present in all three states, it can get into a variety of environments around the planet. The movement of water around Earth's surface is the **hydrologic (water) cycle**.The *Sun*, many millions of kilometers away, *provides the energy that drives the water cycle*. Our nearest star directly impacts the water cycle by supplying the energy needed for evaporation.Most of Earth's water is stored in the oceans where it can remain for hundreds or thousands of years. The oceans are discussed in detail in the chapter Earth's Oceans.

Water changes from a liquid to a gas by evaporation to become water vapor. The Sun's energy can evaporate water from the ocean surface or from lakes, streams, or puddles on land. Only the water molecules evaporate; the salts remain in the ocean or a freshwater reservoir. The water vapor remains in the atmosphere until it undergoes condensation to become tiny droplets of liquid. The droplets gather in clouds, which are blown about the globe by wind. As the water droplets in the clouds collide and grow, they fall from the sky as precipitation. Precipitation can be rain, sleet, hail, or snow. Sometimes precipitation falls back into the ocean and sometimes it falls onto the land surface. This animation shows the annual cycle of monthly mean precipitation around the world.

When water falls from the sky as rain it may enter streams and rivers that flow downward to oceans and lakes. Water that falls as snow may sit on a mountain for several months. Snow may become part of the ice in a glacier, where it may remain for hundreds or thousands of years. Snow and ice may go directly back into the air by sublimation, the process in which a solid changes directly into a gas without first becoming a liquid. Although you probably have not seen water vapor sublimating from a glacier, you may have seen dry ice sublimate in air.

Snow and ice slowly melt over time to become liquid water, which provides a steady flow of fresh water to streams, rivers, and lakes below. A water droplet falling as rain could also become part of a stream or a lake. At the surface, the water may eventually evaporate and reenter the atmosphere.

A significant amount of water infiltrates into the ground. **Soil moisture** is an important reservoir for water. Water trapped in soil is important for plants to grow. Water may seep through dirt and rock below the soil through pores infiltrating the ground to go into Earth's groundwater system. Groundwater enters aquifers that may store fresh water for centuries. Alternatively, the water may come to the surface through springs or find its way back to the oceans. Plants and animals depend on water to live and they also play a role in the water cycle. Plants take up water from the soil and release large amounts of water vapor into the air through their leaves, a process known as transpiration. NASA has an excellent online guide of the hydrologic cycle.

People also depend on water as a natural resource. Not content to get water directly from streams or ponds, humans create canals, aqueducts, dams, and wells to collect water and direct it to where they want it.

USE	UNITED STATES	GLOBAL
Agriculture	34 percent	70 percent
Domestic (drinking, bathing)	12 percent	10 percent
Industry	5 percent	20 percent
Power plant cooling	49 percent	small

The table above displays water use in the United States and globally (Estimated Use of Water in the United States in 2005, USGS). It is important to note that water molecules cycle around. If climate cools and glaciers and ice caps grow, there is less water for the oceans and sea level will fall. The reverse can also happen.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-8-fresh-water.html. **License**: *CC BY: Attribution*

This page titled 6.1: Earth's Fresh Water is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.2: Streams and Rivers



Fresh water in streams, ponds, and lakes is an extremely important part of the water cycle if only because of its importance to living creatures. Along with wetlands, these fresh water regions contain a tremendous variety of organisms.

Streams are bodies of water that have a current; they are in constant motion. Geologists recognize many categories of streams depending on their size, depth, speed, and location. Creeks, brooks, tributaries, bayous, and rivers might all be lumped together as streams. In streams, water always flows downhill, but the form that downhill movement takes varies with rock type, topography, and many other factors. Stream erosion and deposition are extremely important creators and destroyers of landforms and are described in the Erosion and Deposition chapter.

RIVERS

Rivers are the largest types of stream, moving large amounts of water from higher to lower elevations. The Amazon River, the world's river with the greatest flow, has a flow rate of nearly 220,000 cubic meters per second! People have used rivers since the beginning of civilization as a source of water, food, transportation, defense, power, recreation, and waste disposal.

Rivers and streams complete the hydrologic cycle by returning precipitation that falls on land to the oceans (Figure 10.1). Ultimately, gravity is the driving force, as water moves from mountainous regions to sea level. Some of this water moves over the surface and some moves through the ground as **groundwater** . As this water flows it does the work of both erosion and deposition. You will learn about the erosional effects and the deposits that form as a result of this moving water.



Figure 10.1: As rivers and streams move towards the ocean, they carry weathered materials.


Lesson Objectives

- Describe how surface rivers and streams produce erosion.
- Describe the types of deposits left behind by rivers and streams.


Stream Source

PARTS OF A STREAM/RIVER

Oddly enough, there are a variety of different types streams. A stream originates at its source. The **source** is likely to be in the high mountains where snows collect in winter and melt in summer, or a source might be a spring. The source is known as the **headwaters** or the **head** of the stream. A stream may have more than one sources and when two streams come together it's called a **confluence**. The smaller of the two streams is a **tributary** of the larger stream. A stream may create a pool where water slows and becomes deeper.The point at which a stream comes into a large body of water, like an ocean or a lake is called the **mouth**.



A **divide** is a topographically high area that separates a landscape into different water basins. Rain that falls on the north side of a ridge flows into the northern drainage basin and rain that falls on the south side flows into the southern drainage basin. On a much grander scale, entire continents have divides, known as continental divides.

Erosion and Deposition by Rivers and Streams

Erosion from Runoff

As streams move over the ground, they transport weathered materials. Streams continually erode material away from their banks, especially along the outside curves of meanders. Some of these materials are carried in solution. Many minerals are ionic compounds that dissolve easily in water, so water moves these elements to the sea as part of the **dissolved load** that the stream carries. As groundwater leaches through layers of soil and rock, minerals dissolve and are carried away. Groundwater contributes most of the dissolved components that streams carry. Once an element has completely dissolved, it will likely be carried to the ocean, regardless of the velocity of the stream. In some circumstances, the stream water could become saturated with dissolved materials, in which case elements of those minerals might precipitate out of the water before they reach the ocean.

Another way that rivers and streams move weathered materials is as the **suspended load**. These are pieces of rock that are carried as solids as the river flows. Unlike dissolved load, the size of the particle that can be carried as suspended load is determined by the velocity of the stream. As a stream flows faster, it can carry larger and larger particles. The larger the size particle that can be carried by a stream, the greater the stream's **competence**. If a stream has a steep slope or **gradient**, it will have a faster velocity, which means it will be able to carry larger materials in suspension. At flood stage, rivers flow much faster and do more erosion because the added water increases the stream's velocity. Sand, silt and clay size particles generally make up the suspended load for a stream (Figure 10.2). As a stream slows down, either because the stream's slope decreases or because the stream overflows its banks and broadens its channel, the stream will deposit the largest particles it has been carrying first.



Figure 10.2: Rivers carry sand, silt and clay as suspended load. During flood stage, the suspended load greatly increases as stream velocity increases.

The last way that rivers and streams move weathered materials is as **bed load**. This means that although the water in the stream is capable of bumping and pushing these particles along, it is not able to pick them up and carry them continuously. Bed load is named for the fact that these particles get nudged and rolled along the stream bed as the water flows. Occasionally a larger size particle will get knocked into the main part of the stream flow, but then it settles back down to the stream bed because it is too heavy to remain suspended in the water. This is called **saltation**, which we will learn about later in this chapter with transport of particles by wind. Streams with high velocities and steep gradients do a great deal of downcutting into the stream bed, which is primarily accomplished by movement of particles that make up the bed load. Particles that move along as the bed load of a stream do not move continuously along, but rather in small steps or jumps with periods of remaining stationary in between.

Stream and River Erosion

As a stream moves water from high elevations, like mountains, towards low elevations, like the ocean, which is at sea level, the work of the stream changes. At high elevations, streams are just beginning streams that have small channels and steep gradients. This means that the stream will have a high velocity and will do lots of work eroding its stream bed. The higher the elevation, the farther the stream is from where it eventually meets the sea. **Base level** is the term for where a stream meets sea level or standing water, like a lake or the ocean. Streams will work to downcut their stream beds until they reach base level.

As a stream moves out of high mountainous areas into lower areas closer to sea level, the stream is closer to its base level and does more work eroding the edges of its banks than downcutting into its stream bed. At some point in most streams, there are curves or bends in the stream channel called **meanders** (Figure 10.3). The stream erodes material along its outer banks and deposits material along the inside curves of a meander as it flows to the ocean (Figure 10.4). This causes these meanders to migrate laterally over time. The erosion of the outside edge of the stream's banks begins the work of carving a floodplain, which is a flat level area surrounding the stream channel.



Figure 10.3: Here a stream can be seen actively eroding its outer banks along a meander.



Figure 10.4: This stream has deposited larger materials like gravel and pebbles along the inside curve of a meander.

Stream and River Deposition




Figure 10.5: After many floods, a stream builds natural levees along its banks.

Once a stream nears the ocean, it is very close to its base level and now deposits more materials than it erodes. As you just learned, one place where a river deposits material is along the inside edges of meanders. If you ever decide to pan for gold or look for artifacts from an older town or civilization, you will sift through these deposits. Gold is one of the densest elements on Earth. Streams are lazy and never want to carry more materials than absolutely necessary. It will drop off the heaviest and largest particles first, that is why you might find gold in a stream deposit. Imagine that you had to carry all that you would need for a week as you walked many kilometers. At first you might not mind the weight of what you are carrying at all, but as you get tired, you will look to drop off the heaviest things you are carrying first!

When a river floods or overflows its channel, the area where the stream flows is suddenly much broader and shallower than it was when it was in its channel. This slows down the velocity of the stream's flow and causes the stream to drop off much of its load. The farmers who use the floodplain areas around the Nile River rely on these deposits to supply nutrients to their fields each year as the river floods its banks. At flood stage, a river will also build natural levees as the largest size particles build a higher area around the edges of the stream channel (Figure 10.5).

When a river meets either standing water or nearly flat lying ground, it will deposit its load. If this happens in water, a river may form a **delta**. From its headwaters in the mountains, along a journey of many kilometers, rivers carry the eroded materials that form their stream load. Suddenly the river slows down tremendously in velocity, and drops the tremendous load of sediments it has been carrying. Deltas are relatively flat topped, often triangular shaped deposits of sediments that form where a large river meets the ocean. The name delta comes from the capital Greek letter delta, which is a triangle, even though not all deltas have this shape. A triangular shaped delta forms as the main stream channel splits into many smaller **distributaries**. As the channel shifts back and forth dropping off sediments and moving to a new channel location a wide triangular deposit forms.

There are three types of beds that make up a delta (Figure 10.6). The first particles to be dropped off are the coarsest sediments and these form sloped layers called foreset beds that make up the front edge of the delta. Further out into calmer water, lighter, more

fine grained sediments form thin, horizontal layers. These are called bottomset beds. During floodstage, the whole delta can be covered by finer sediments which will overlie the existing delta. These are called topset beds. These form last and lie on top of the rest of the delta.




Figure 10.6: The three types of beds that form the layers of a delta.

Not all large rivers form deltas as they meet the ocean. Whether a delta forms depends on the action of waves and tides. If the water is quiet water such as a gulf or shallow sea, a delta may form. If the sediments are carried away, then no delta will form. Sediments brought to the shore and distributed along coastlines by longshore transport form our beaches and barrier islands.

If a river or stream suddenly reaches nearly flat ground, like a broad flat valley or plain, an **alluvial fan** develops at the base of the slope (Figure 10.7). An alluvial fan is a curved top, fan shaped deposit of coarse sediments that drop off as the stream suddenly loses velocity. The fan spreads out in a curve in the direction of the flat land as many stream channels move across the curved surface of the alluvial fan, forming and unforming many channels as sediments are deposited. Alluvial fans generally form in more arid regions.



Figure 10.7: This satellite photo of an alluvial fan in Iran shows the typical fan shape of these deposits. The stream forming the alluvial fan runs from the mountains in the southwest (lower left) corner of the photograph toward the flatter land to the northeast (upper right). The green rectangles are farm fields which utilize the distributed water.

Lesson Summary

- Rivers and streams erode the land as they move from higher elevations to the sea.
- Eroded materials can be carried in a river as dissolved load, suspended load, or bed load.
- A river will deeply erode the land when it is far from its base level, the elevation where it enters standing water like the ocean.
- As a river develops bends, called meanders, it forms a broad, flat area known as a floodplain.
- At the end of a stream, a delta or an alluvial fan might form where the river drops off much of the load of sediments it carries.

Review Questions

1. What are the three kinds of load that make up the particles a stream carries. Name and define each type.
2. What is a stream’s gradient? What effect does it have on the work of a stream?
3. Describe several erosional areas produced by streams. Explain why erosion occurs here.
4. What type of gradient or slope would a river have when it is actively eroding its stream bed? Explain.
5. When would a river form an alluvial fan and when will it form a delta? Describe the characteristics of each type of deposit.

CC licensed content, Shared previously

- Earth Science for High School. **Provided by**: CK-12. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/?noindex=None. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 6.2: Streams and Rivers is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.3: Groundwater

Lesson Objectives

- Define groundwater.
- Explain the location, use, and importance of aquifers.
- Define springs.
- Describe how wells work and why they are important.

Vocabulary

- aquifer
- capillary action
- impermeable
- permeability
- porosity
- spring
- subsidence
- water table
- well

Introduction

Although this may seem surprising, water beneath the ground is commonplace. Usually groundwater travels slowly and silently beneath the surface, but in some locations it bubbles to the surface at springs. The products of erosion and deposition by groundwater were described in the Erosion and Deposition chapter.

Groundwater

Groundwater is the largest reservoir of liquid fresh water on Earth and is found in **aquifers**, porous rock and sediment with water in between. Water is attracted to the soil particles and **capillary action**, which describes how water moves through a porous media, moves water from wet soil to dry areas.

Aquifers are found at different depths. Some are just below the surface and some are found much deeper below the land surface. A region may have more than one aquifer beneath it and even most deserts are above aquifers. The source region for an aquifer beneath a desert is likely to be far from where the aquifer is located; for example, it may be in a mountain area.

The amount of water that is available to enter groundwater in a region is influenced by the local climate, the slope of the land, the type of rock found at the surface, the vegetation cover, land use in the area, and water retention, which is the amount of water that remains in the ground. More water goes into the ground where there is a lot of rain, flat land, porous rock, exposed soil, and where water is not already filling the soil and rock.

The residence time of water in a groundwater aquifer can be from minutes to thousands of years. Groundwater is often called “fossil water” because it has remained in the ground for so long, often since the end of the ice ages.

Aquifers


Features of an Aquifer

To be a good aquifer, the rock in the aquifer must have good:

- **porosity**: small spaces between grains
- **permeability**: connections between pores

This animation shows porosity and permeability. The water droplets are found in the pores between the sediment grains, which is porosity. When the water can travel between ores, that’s permeability. http://www.nature.nps.gov/GEOLOGY/usgsnps/animate/POROS_3.MPG

To reach an aquifer, surface water infiltrates downward into the ground through tiny spaces or pores in the rock. The water travels down through the permeable rock until it reaches a layer that does not have pores; this rock is **impermeable** (**Figure** below). This impermeable rock layer forms the base of the aquifer. The upper surface where the groundwater reaches is the **water table**.




Groundwater is found beneath the solid surface. Notice that the water table roughly mirrors the slope of the land's surface. A well penetrates the water table.

The Water Table

For a groundwater aquifer to contain the same amount of water, the amount of recharge must equal the amount of discharge. What are the likely sources of recharge? What are the likely sources of discharge?


In wet regions, streams are fed by groundwater; the surface of the stream is the top of the water table (**Figure** below). In dry regions, water seeps down from the stream into the aquifer. These streams are often dry much of the year. Water leaves a groundwater reservoir in streams or springs. People take water from aquifers, too.

What happens to the water table when there is a lot of rainfall? What happens when there is a drought? Although groundwater levels do not rise and fall as rapidly as at the surface, over time the water table will rise during wet periods and fall during droughts.



The top of the stream is the top of the water table. The stream feeds the aquifer.

One of the most interesting, but extremely atypical types of aquifers is found in Florida. Although aquifers are very rarely underground rivers, in Florida water has dissolved the limestone so that streams travel underground and above ground (**Figure** below).



In Florida, groundwater is sometimes not underground.


Groundwater Use

Groundwater is an extremely important water source for people. Groundwater is a renewable resource and its use is sustainable when the water pumped from the aquifer is replenished. It is important for anyone who intends to dig a well to know how deep beneath the surface the water table is. Because groundwater involves interaction between the Earth and the water, the study of groundwater is called hydrogeology.

Some aquifers are overused; people pump out more water than is replaced. As the water is pumped out, the water table slowly falls, requiring wells to be dug deeper, which takes more money and energy. Wells may go completely dry if they are not deep enough to reach into the lowered water table.

The Ogallala Aquifer supplies about one-third of the irrigation water in the United States (**Figure** below). The aquifer is found from 30 to 100 meters deep over about 440,000 square kilometers! The water in the aquifer is mostly from the last ice age.

The Ogallala Aquifer is widely used by people for municipal and agricultural needs.



The Ogallala Aquifer is found beneath eight states and is heavily used.

About eight times more water is taken from the Ogallala Aquifer each year than is replenished. Much of the water is used for irrigation (**Figure** below).



Farms in Kansas use central pivot irrigation, which is more efficient since water falls directly on the crops instead of being shot in the air. These fields are between 800 and 1600 meters (0.5 and 1 mile) in diameter.

Lowering the water table may cause the ground surface to sink. **Subsidence** may occur beneath houses and other structures (**Figure** below).




The San Joaquin Valley of California is one of the world's major agricultural areas. So much groundwater has been pumped that the land has subsided many tens of feet.

When coastal aquifers are overused, salt water from the ocean may enter the aquifer, contaminating the aquifer and making it less useful for drinking and irrigation. Salt water incursion is a problem in developed coastal regions, such as on Hawaii.

Springs

Groundwater meets the surface in a stream, as shown above, or a **spring** (**Figure** below). A spring may be constant, or may only flow at certain times of year.



(a) Big Spring in Missouri lets out 12,000 liters of water per second. (b) Other springs are just tiny outlets like this one.

Towns in many locations depend on water from springs. Springs can be an extremely important source of water in locations where surface water is scarce (**Figure** below).



In the dry Arizona desert, Oak Creek and many other streams are spring fed.

Wells

A **well** is created by digging or drilling to reach groundwater. When the water table is close to the surface, wells are a convenient method for extracting water. When the water table is far below the surface, specialized equipment must be used to dig a well. Most wells use motorized pumps to bring water to the surface, but some still require people to use a bucket to draw water up (**Figure** below).



An old-fashioned well that uses a bucket drawn up by hand.

Lesson Summary

- Groundwater is the largest reservoir of fresh water.
- The water table is the top of an aquifer below which is water and above is rock or soil mixed with air.
- Aquifers are underground areas of sediment or rock that hold groundwater.
- An aquifer needs good porosity and permeability.
- Where groundwater intersects the ground surface, a spring can form.
- People dig or drill wells to access groundwater.

Review Questions

1. What is groundwater?

2. What is the water table?

3. What are aquifers and why are they so important?

4. Replenishing an aquifer is important because it makes the aquifer a resource that can last a long time. What do you think are ways to keep the amount of water used and the amount of water replenished the same?

5. How does a well work?

6. Since groundwater is largely unseen from the surface, how might you monitor how humans are affecting the amount of groundwater in an aquifer?

Further Reading / Supplemental Links

Earth's water distribution video, University of Waikato, New Zealandhttp://www.sciencelearn.org.nz/contexts/h2o_on_the_go/sci_media/video/earth_s_water_distribution

Points to Consider

- Is water from a river or from a well more likely to be clean to drink?
- Why is overuse of groundwater a big concern?
- What policies might people put in place to conserve water levels in lakes and aquifers?

CC licensed content, Shared previously

- Groundwater. **Authored by**: Joanne Adair. **Provided by**: CK-12. **Located at**: http://www.ck12.org/user:amFkYWlyQGJyZXdlcmVkdS5vcmc./book/General-Environmental-Science/r5/section/7.3/. **License**: *CC BY-NC-SA: Attribution-NonCommercial-ShareAlike*

This page titled 6.3: Groundwater is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.4: Water Resources

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=78

CC licensed content, Original

- **Authored by**: William Hopper. **Provided by**: Florida Memorial University. **Located at**: https://www.youtube.com/watch?v=QuF6SvxfCfc. **License**: *CC BY: Attribution*

This page titled 6.4: Water Resources is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.5: Floods

Floods



Floods are a natural part of the water cycle, but they can be terrifying forces of destruction. Put most simply, a flood is an overflow of water in one place. Floods can occur for a variety of reasons, and their effects can be minimized in several different ways. Perhaps unsurprisingly, floods tend to affect low-lying areas most severely.

Floods usually occur when precipitation falls more quickly than that water can be absorbed into the ground or carried away by rivers or streams. Waters may build up gradually over a period of weeks, when a long period of rainfall or snow-melt fills the ground with water and raises stream levels.

Flash floods are sudden and unexpected, taking place when very intense rains fall over a very brief period. A flash flood may do its damage miles from where the rain actually falls if the water travels far down a dry streambed so that the flash flood occurs far from the location of the original storm.

Heavily vegetated lands are less likely to experience flooding. Plants slow down water as it runs over the land, giving it time to enter the ground. Even if the ground is too wet to absorb more water, plants still slow the water's passage and increase the time between rainfall and the water's arrival in a stream; this could keep all the water falling over a region to hit the stream at once. Wetlands act as a buffer between land and high water levels and play a key role in minimizing the impacts of floods. Flooding is often more severe in areas that have been recently logged.

When a dam breaks along a reservoir, flooding can be catastrophic. High water levels have also caused small dams to break, wreaking havoc downstream. People try to protect areas that might flood with dams, and dams are usually very effective. People may also line a river bank with levees, high walls that keep the stream within its banks during floods. A levee in one location may just force the high water up or downstream and cause flooding there. The New Madrid Overflow in the image above was created with the recognition that the Mississippi River sometimes simply cannot be contained by levees and must be allowed to flood.

Not all the consequences of flooding are negative. Rivers deposit new nutrient-rich sediments when they flood and so floodplains have traditionally been good for farming. Flooding as a source of nutrients was important to Egyptians along the Nile River until the Aswan Dam was built in the 1960s. Although the dam protects crops and settlements from the annual floods, farmers must now use fertilizers to feed their crops.

Floods are also responsible for moving large amounts of sediments about within streams. These sediments provide habitats for animals, and the periodic movement of sediment is crucial to the lives of several types of organisms. Plants and fish along the Colorado River, for example, depend on seasonal flooding to rearrange sand bars.

This page titled 6.5: Floods is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

6.6: Wetlands

A **wetland** is a land area that is saturated with water, either permanently or seasonally, such that it takes on the characteristics of a distinct ecosystem.[2] Primarily, the factor that distinguishes wetlands from other land forms or water bodies is the characteristic vegetation that is adapted to its unique soil conditions. Wetlands consist primarily of hydric soil, which supports aquatic plants.[3][4]

The water found in wetlands can be saltwater, freshwater, or brackish.[4] Main wetland types include swamps, marshes, bogs and fens.[5] Sub-types include mangrove, carr, pocosin, and varzea.

Wetlands play a number of roles in the environment, principally water purification, flood control, and shoreline stability. Wetlands are also considered the most biologically diverse of all ecosystems, serving as home to a wide range of plant and animal life.[6]

Wetlands occur naturally on every continent except Antarctica.[7] They can also be constructed artificially as a water management tool, which may play a role in the developing field of water-sensitive urban design.

The largest wetlands in the world include the Amazon River basin and the West Siberian Plain.[8] Another large wetland is the Pantanal, which straddles Brazil, Bolivia, and Paraguay in South America.[9]

The UN Millennium Ecosystem Assessment determined that environmental degradation is more prominent within wetland systems than any other ecosystem on Earth. International conservation efforts are being used in conjunction with the development of rapid assessment tools to inform people about wetland issues.



Laguna de Rocha, the largest wetland in the urban area in Esteban Echeverría Partido, Argentina



Mangrove swamps are coastal wetlands. This swamp is in the Florida Everglades.[1]



Peat bogs are freshwater wetlands that develop in areas with standing water and low soil fertility.



Marshes develop along the edges of rivers and lakes.



Many species of frogs live in wetlands, while others visit them each year to lay eggs.



Snapping turtles are one of the many kinds of turtles found in wetlands

Definitions

A patch of land that develops pools of water after a rain storm would not be considered a "wetland," even though the land is wet. Wetlands have unique characteristics: they are generally distinguished from other water bodies or landforms based on their water level and on the types of plants that live within them. Specifically, wetlands are characterized as having a water table that stands at or near the land surface for a long enough period each year to support aquatic plants.[10][11]

A more concise definition is a community composed of hydric soil and hydrophytes.[7]

Wetlands have also been described as ecotones, providing a transition between dry land and water bodies.[12] Mitsch and Gosselink write that wetlands exist "…at the interface between truly terrestrial ecosystems and aquatic systems, making them inherently different from each other, yet highly dependent on both."[13]

In environmental decision-making, there are subsets of definitions that are agreed upon to make regulatory and policy decisions.

Technical definitions

A wetland is "an ecosystem that arises when inundation by water produces soils dominated by anaerobic processes, which, in turn, forces the biota, particularly rooted plants, to adapt to flooding."[14] There are four main kinds of wetlands — marsh, swamp, bog and fen (bogs and fens both being types of mires). Some experts also recognize wet meadows and aquatic ecosystems as additional wetland types.[5] The largest wetlands in the world include the swamp forests of the Amazon and the peatlands of Siberia.[8]

Ramsar Convention definition

Under the Ramsar international wetland conservation treaty, wetlands are defined as follows:[15]

- Article 1.1: "…wetlands are areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six metres."
- Article 2.1: "[Wetlands] may incorporate riparian and coastal zones adjacent to the wetlands, and islands or bodies of marine water deeper than six metres at low tide lying within the wetlands."

Regional definitions

Although the general definition given above applies around the world, each county and region tends to have its own definition for legal purposes. For example, in the United States, wetlands are defined as "those areas that are inundated or saturated by surface or groundwater at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs and similar

areas".[16] This definition has been used in the enforcement of the Clean Water Act. Some US states, such as Massachusetts and New York, have separate definitions that may differ from the federal government's.

In the United States Code (16 U.S.C., Section 3801(a)(18)), the term wetland is defined "as land that (A) has a predominance of hydric soils, (B) is inundated or saturated by surface or groundwater at a frequency and duration sufficient to support a prevalence of hydrophytic vegetation typically adapted for life in saturated soil conditions and C) under normal circumstances supports a prevalence of such vegetation." Related to this legal definitions, the term "normal circumstances" are conditions expected to occur during the wet portion of the growing season under normal climatic conditions (not unusually dry or unusually wet), and in the absence of significant disturbance. It is not uncommon for a wetland to be dry for long portions of the growing season. Wetlands can be dry during the dry season and abnormally dry periods during the wet season, but under normal environmental conditions the soils in a wetland will be saturated to the surface or inundated such that the soils become anaerobic, and those conditions will persist through the wet portion of the growing season.

Ecology

The most important factor producing wetlands is flooding. The duration of flooding determines whether the resulting wetland has aquatic, marsh or swamp vegetation. Other important factors include fertility, natural disturbance, competition, herbivory, burial and salinity.[5] When peat accumulates, bogs and swamps arise.

Characteristics

Wetlands vary widely due to local and regional differences in topography, hydrology, vegetation, and other factors, including human involvement. Wetlands can be divided into two main classes: tidal and non-tidal areas.[7]

Hydrology

Wetland hydrology is associated with the spatial and temporal dispersion, flow, and physiochemical attributes of surface and ground water in its reservoirs. Based on hydrology, wetlands can be categorized as riverine (associated with streams), lacustrine (associated with lakes and reservoirs), and palustrine (isolated). Sources of hydrological flows into wetlands are predominately precipitation, surface water, and ground water. Water flows out of wetlands by evapotranspiration, surface runoff, and sub-surface water outflow. Hydrodynamics (the movement of water through and from a wetland) affects hydroperiods (temporal fluctuations in water levels) by controlling the water balance and water storage within a wetland.[17]

Landscape characteristics control wetland hydrology and hydrochemistry. The O_2 and CO_2 concentrations of water depend on temperature and atmospheric pressure. Hydrochemistry within wetlands is determined by the pH, salinity, nutrients, conductivity, soil composition, hardness, and the sources of water. Water chemistry of wetlands varies across landscapes and climatic regions. Wetlands are generally minerotrophic with the exception of bogs.

Bogs receive their water from the atmosphere and therefore their water has low mineral ionic composition because ground water has a higher concentration of dissolved nutrients and minerals in comparison to precipitation.

The water chemistry of fens ranges from low pH and low minerals to alkaline with high accumulation of calcium and magnesium because they acquire their water from precipitation as well as ground water.[18]

Role of salinity

Salinity has a strong influence on wetland water chemistry, particularly in wetlands along the coast.[5][19] In non-riverine wetlands, natural salinity is regulated by interactions between ground and surface water, which may be influenced by human activity.[20]

Soil

Carbon is the major nutrient cycled within wetlands. Most nutrients, such as sulfur, phosphorus, carbon, and nitrogen are found within the soil of wetlands. Anaerobic and aerobic respiration in the soil influences the nutrient cycling of carbon, hydrogen, oxygen, and nitrogen,[21] and the solubility of phosphorus[22] thus contributing to the chemical variations in its water. Wetlands with low pH and saline conductivity may reflect the presence of acid sulfates[23] and wetlands with average salinity levels can be heavily influenced by calcium or magnesium. Biogeochemical processes in wetlands are determined by soils with low redox potential.[24]

Biota

The biota of a wetland system includes its vegetation zones and structure as well as animal populations. The most important factor affecting the biota is the duration of flooding.[5] Other important factors include fertility and salinity. In fens, species are highly dependent on water chemistry. The chemistry of water flowing into wetlands depends on the source of water and the geological material in which it flows through[25] as well as the nutrients discharged from organic matter in the soils and plants at higher elevations in slope wetlands.[26] Biota may vary within a wetland due to season or recent flood regimes.

Flora

There are four main groups of hydrophytes that found in wetland systems throughout the world.[27]

Submerged water plants. This type of vegetation is found completely underwater. Submerged wetland vegetation can grow in saline and fresh-water conditions. Some species have underwater flowers, while others have long stems to allow the flowers to reach the surface.[28] Submerged species provide a food source for native fauna, habitat for invertrebrates, and also possess filtration capabilities. Examples include seagrasses and eelgrass.

Floating water plants. Floating vegetation is usually small although it may take up a large surface area in a wetland system. These hydrophytes have small roots and are only found in slow-moving water with rich-nutrient level water Floating aquatic plants are a food resource for avian species. Examples include water lilies lily pad and duckweed.

Emergent water plants. Emergent water plants can be seen above the surface of the water but whose roots are completely submerged. Many have aerenchyma to transmit oxygen from the atmosphere to their roots.[29] Extensive areas of emergent plants are usually termed marsh. Examples include cattails (Typha) and arrow arum (Peltandra virginica).

Surrounding trees and shrubs. Forested wetlands are generally known as swamps.[5] The upper level of these swamps is determined by high water levels, which are negatively affected by dams.[30] Some swamps can be dominated by a single species, such as silver maple swamps around the Great Lakes.[31] Others, like those of the Amazon Basin, have large numbers of different tree species.[32] Examples include cypress (Taxodium) and mangrove.

Fauna

Fish: Fish are more dependent on wetland ecosystems than any other type of habitat. 75% of the United States' commercial fish and shellfish stocks depend solely on estuaries to survive.[6] Tropical fish species need mangroves for critical hatchery and nursery grounds and the coral reef system for food.

Amphibians: Frogs are the most crucial amphibian species in wetland systems. Frogs need both terrestrial and aquatic habitats in which to reproduce and feed. While tadpoles control algal populations, adult frogs forage on insects. Frogs are used as an indicator of ecosystem health due to their thin skin which absorbs both nutrient and toxins from the surrounding environment resulting in an above average extinction rate in unfavorable and polluted environmental conditions.[33]

Reptiles: Alligators and crocodiles are two common reptilian species. Alligators are found in fresh water along with the fresh water species of the crocodile. The saltwater crocodile is found in estuaries and mangroves and can be seen in the coastline bordering the Great Barrier Reef in Australia. The Florida Everglades is the only place in the world where both crocodiles and alligators co-exist.[1]Snakes, lizards, goannas, and turtles also can be seen throughout wetlands. Snapping turtles are one of the many kinds of turtles found in wetlands.

Mammals: Multiple small mammals as well as large herbivore and apex species such as the Florida Panther live within and around wetlands. The wetland ecosystem attracts mammals due to its prominent seed sources, invertebrate populations, and numbers of small reptiles and amphibians.

Monotremes: The platypus (*Ornithorhynchus anatinus*) is found in eastern Australia living in freshwater rivers or lakes, and much like the beaver creates dams, create burrows for shelter and protection. The platypus swims through the use of webbed feet. Platypuses feed on insect larvae, worms, or other freshwater insects hunting mainly by night by the use of their bill. They turn up mud on the bottom of the lake or river, and with the help of the electroreceptors located on the bill, unearth insects and freshwater insects. The platypus stores their findings in special pouches behind their bill and consumes its prey upon returning to the surface.[34]

Insects and invertebrates: These species total more than half of the 100,000 known animal species in wetlands. Insects and invertebrates can be submerged in the water or soil, on the surface, and in the atmosphere.

Algae

Algae are diverse water plants that can vary in size, color, and shape. Algae occur naturally in habitats such as inland lakes, inter-tidal zones, and damp soil and provide a dedicated food source for animals, fish, and invertebrates. There are three main groups of algae:

Plankton are algae which are microscopic, free-floating algae. This algae is so tiny that on average, if fifty of these microscopic algae were lined up end-to-end, it would only measure one millimetre. Plankton are the basis of the food web and are responsible for primary production in the ocean using photosynthesis to make food. Filamentous algae are long strands of algae cells that form floating mats. Chara and Nitella algae are upright algae that look like a submerged plant with roots.[35]

Climates

Temperature



Wetlands contrast the hot, arid landscape around Middle Spring, Fish Springs National Wildlife Refuge, Utah.

Temperatures vary greatly depending on the location of the wetland. Many of the world's wetlands are in temperate zones (midway between the North or South Pole and the equator). In these zones, summers are warm and winters are cold, but temperatures are not extreme. However, wetlands found in the tropic zone, around the equator, are warm year round. Wetlands on the Arabian Peninsula, for example, can reach 50 °C (122 °F) and would therefore be subject to rapid evaporation. In northeastern Siberia, which has a polar climate, wetland temperatures can be as low as −50 °C (−58 °F). In a moderate zone, such as the Gulf of Mexico, a typical temperature might be 11 °C (51 °F). Wetlands are also located in every climatic zone.

Rainfall

The amount of rainfall a wetland receives varies widely according to its area. Wetlands in Wales, Scotland, and Western Ireland typically receive about 1500 mm (or 60 in) per year. In some places in Southeast Asia, where heavy rains occur, they can receive up to 10,000 mm (about 200 in). In the northern areas of North America, wetlands exist where as little as 180 mm (7 inches) of rain fall each year.

Temporal variation: [36]

- Perennial systems
- Seasonal systems
- Episodic (periodic or intermittent) system of the down
- Surface flow may occur in some segments, with subsurface flow in other segments
- Ephemeral (short-lived) systems
- Migratory species

Human-disturbance: [37]

- Encroachment
  - Drainage
  - Development
  - Over-grazing
  - Mining
  - Unsustainable water use

- Ecosystem Stress
  - Water Scarcity
  - Endangered species
  - Disruption of breeding grounds
  - Imbalance in sediment load and nutrient filtration

Peatswamps of South-east Asia

In Southeast Asia, peatswamp forests and soils are being drained, burnt, mined, and overgrazed, contributing severely to climate change.[37] As a result of peat drainage, the organic carbon that was built up over thousands of years and is normally under water is suddenly exposed to the air. It decomposes and turns into carbon dioxide (CO_2), which is released into the atmosphere. Peat fires cause the same process to occur and in addition create enormous clouds of smoke that cross international borders, such as happens every year in Southeast Asia. While peatlands constitute only 3% of the world's land area, their degradation produces 7% of all fossil fuel CO_2 emissions.

Through the building of dams, Wetlands International is halting the drainage of peatlands in Southeast Asia, hoping to mitigate CO_2 emissions. Concurrent wetland restoration techniques include reforestation with native tree species as well as the formation of community fire brigades. This sustainable approach can be seen in Central Kalimantan and Sumatra, Indonesia.

Aquaculture

Concerns are developing over certain aspects of farm fishing, which uses natural waterways to harvest fish for human consumption and pharmaceuticals. This practice has become especially popular in Asia and the South Pacific. Its impact upon much larger waterways downstream has negatively affected many small island developing states.[38]

The function of natural wetlands can be classified by their ecosystem benefits. United Nations Millennium Ecosystem Assessment and Ramsar Convention found wetlands to be of biosphere significance and societal importance in the following areas:

- Flood control
- Groundwater replenishment
- Shoreline stabilisation and storm protection
- Water purification
- Reservoirs of biodiversity
- Wetland products
- Cultural values
- Recreation and tourism
- Climate change mitigation and adaptation

The economic worth of the ecosystem services provided to society by intact, naturally functioning wetlands is frequently much greater than the perceived benefits of converting them to 'more valuable' intensive land use – particularly as the profits from unsustainable use often go to relatively few individuals or corporations, rather than being shared by society as a whole.*-Ramsar convention*

Unless otherwise cited, Ecosystem services is based on the following series of references. [6]

Flood control

Major wetland type: floodplain

Storage Reservoirs and Flood Protection. The wetland system of floodplains is formed from major rivers downstream from their headwaters. Notable river systems that produce large spans of floodplain include the Nile River, the Niger river inland delta, [the Zambezi River flood plain], [the Okavango River inland delta],[the Kafue River flood plain][the Lake Bangweulu flood plain] (Africa), Mississippi River (USA), Amazon River (South America), Yangtze River (China), Danube River (Central Europe) and Murray-Darling River (Australia). "The floodplains of major rivers act as natural storage reservoirs, enabling excess water to spread out over a wide area, which reduces its depth and speed. Wetlands close to the headwaters of streams and rivers can slow down rainwater runoff and spring snowmelt so that it doesn't run straight off the land into water courses. This can help prevent sudden, damaging floods downstream."[6]

Human-Impact. Converting wetlands through drainage and development have contributed to the issue of irregular flood control through forced adaption of water channels to narrower corridors due to loss of wetland area. These new channels must manage the same amount of precipitation causing flood peaks to be [higher or deeper] and floodwaters to travel faster.

Water management engineering developments in the past century have degraded these wetlands through the construction on artificial embankments. These constructions may be classified as dykes, bunds, levees, weirs, barrages and dams but serve the single purpose of concentrating water into a select source or area. Wetland water sources that were once spread slowly over a large, shallow area are pooled into deep, concentrated locations. Loss of wetland floodplains results in more severe and damaging flooding. Catastrophic human impact in the Mississippi River floodplains was seen in death of several hundred individuals during a levee breach in New Orleans caused by Hurricane Katrina. Ecological catastrophic events from human-made embankments have been noticed along the Yangtze River floodplains after the where the middle of the river has become prone to more frequent and damaging flooding including the loss of riparian vegetation, a 30% loss of the vegetation cover throughout the river's basin, a doubling of the percentage of the land affected by soil erosion, and a reduction in reservoir capacity through siltation build-up in floodplain lakes.[6]

Groundwater replenishment

Major wetland type: marsh, swamp, & subterranean karst and cave hydrological systems

The surface water which is the water visibly seen in wetland systems only represents a portion of the overall water cycle which also includes atmospheric water and groundwater. Wetland systems are directly linked to groundwater and a crucial regulator of both the quantity and quality of water found below the ground. Wetland systems that are made of permeable sediments like limestone or occur in areas with highly variable and fluctuating water tables especially have a role in groundwater replenishment or water recharge. Sediments that are porous allow water to filter down through the soil and overlying rock into aquifers which are the source of 95% of the world's drinking water. Wetlands can also act as recharge areas when the surrounding water table is low and as a discharge zone when it is too high. Karst (cave) systems are a unique example of this system and are a connection of underground rivers influenced by rain and other forms of precipitation. These wetland systems are capable of regulating changes in the water table on upwards of 130 metres (430 ft).

Human-Impact. Groundwater is an important source of water for drinking and irrigation of crops. Over 1 billion people in Asia and 65% of the public water sources in Europe source 100% of their water from groundwater. Irrigation is a massive use of groundwater with 80% of the world's groundwater used for agricultural production.[6]

Unsustainable abstraction of groundwater has become a major concern. In the Commonwealth of Australia, water licensing is being implemented to control use of the water in major agricultural regions. On a global scale, groundwater deficits and water scarcity is one of the most pressing concerns facing the 21st century.[6]

Shoreline stabilisation and storm protection

Wetland type: Mangroves, Coral Reefs, Saltmarsh

Main article: Integrated Coastal Zone Management

Tidal and inter-tidal wetland systems protect and stabilize coastal zones. Coral reefs provide a protective barrier to coastal shoreline. Mangroves stabilize the coastal zone from the interior and will migrate with the shoreline to remain adjacent to the boundary of the water. The main conservation benefit these systems have against storms and tidal waves is the ability to reduce the speed and height of waves and floodwaters.

Human-Impact. The sheer number of people who live and work near the coast is expected to grow immensely over the next 50 years. From an estimated 200 million people that currently live in low-lying coastal regions, the development of urban coastal centers is projected to increase the population by 5 fold within 50 years.[39] The United Kingdom has begun the concept of managed coastal realignment. This management technique provides shoreline protection through restoration of natural wetlands rather than through applied engineering.

Water purification

Wetland Type: Floodplain, Mudflat, Saltmarsh, Mangroves

Nutrient Retention. Wetlands cycle both sediments and nutrients balancing terrestrial and aquatic ecosystems. A natural function of wetland vegetation is the up-take and storage of nutrients found in the surrounding soil and water. These nutrients are retained in the system until the plant dies or is harvested by animals or humans. Wetland vegetation productivity is linked to the climate, wetland type, and nutrient availability. The grasses of fertile floodplains such as the Nile produce the highest yield including plants such as *Arundo donax*(giant reed), *Cyperus papyrus*(papyrus), *Phragmites* (reed) and *Typha* (cattail, bulrush).

Sediment Traps. Rainfall run-off is responsible for moving sediment through waterways. These sediments move towards larger and more sizable waterways through a natural process that moves water towards oceans. All types of sediments which may be composed of clay, sand, silt, and rock can be carried into wetland systems through this process. Reedbeds or forests located in wetlands act as physical barriers to slow waterflow and trap sediment.

Water purification. Many wetland systems possess biofilters, hydrophytes, and organisms that in addition to nutrient up-take abilities have the capacity to remove toxic substances that have come from pesticides, industrial discharges, and mining activities. The up-take occurs through most parts of the plant including the stems, roots, and leaves . Floating plants can absorb and filter heavy metals. *Eichhornia crassipes* (water hyacinth), *Lemna* (duckweed) and *Azolla* (water fern) store iron and copper commonly found in wastewater. Many fast-growing plants rooted in the soils of wetlands such as *Typha* (cattail) and *Phragmites* (reed) also aid in the role of heavy metal up-take. Animals such as the oyster can filter more than 200 liters (53 gallons) of water per day while grazing for food, removing nutrients, suspended sediments, and chemical contaminants in the process.

Capacity. The ability of wetland systems to store nutrients and trap sediment is highly efficient and effective but each system has a threshold. An overabundance of nutrient input from fertilizer run-off, sewage effluent, or non-point pollution will cause eutrophication. Upstream erosion from deforestation can overwhelm wetlands making them shrink in size and see dramatic biodiversity loss through excessive sedimentation load. The capacity of wetland vegetation to store heavy metals is affected by waterflow, number of hectares (acres), climate, and type of plant.

Human-Impact. Introduced hydrophytes in different wetland systems can have devastating results. The introduction of water hyacinth, a native plant of South America into Lake Victoria in East Africa as well as duckweed into non-native areas of Queensland, Australia, have overtaken entire wetland systems suffocating the ecosystem due to their phenomenal growth rate and ability to float and grow on the surface of the water.

Constructed wetlands

Main article: Constructed wetland

The function of most natural wetland systems is not to manage to wastewater, however, their high potential for the filtering and the treatment of pollutants has been recognized by environmental engineers that specialize in the area of wastewater treatment. These constructed artificial wetland systems are highly controlled environments that intend to mimic the occurrences of soil, flora, and microorganisms in natural wetlands to aid in treating wastewater effluent. Artificial wetlands provide the ability to experiment with flow regimes, micro-biotic composition, and flora in order to produce the most efficient treatment process. Other advantages are the control of retention times and hydraulic channels.[40] The most important factors of constructed wetlands are the water flow processes combined with plant growth. Constructed wetland systems can be surface flow systems with only free-floating macrophytes, floating-leaved macrophytes, or submerged macrophytes; however, typical free water surface systems are usually constructed with emergent macrophytes.[41] Constructed wetlands can be adapted to treat raw sewage, secondary domestic sludge, enhance water quality of oxidation ponds' discharge, storm waters, mining waste, and industrial and agricultural waste effluents. The Urrbrae Wetland in Australia was constructed for urban flood control and environmental education. International wastewater management programs can be seen from Kolkata (Calcutta), India to Arcata, California, USA.[42]

Kolkata's constructed wetland. Kolkata is an example of how constructed wetlands are being utilized in developing countries.

Using the purification capacity of wetlands, the Indian city of Kolkata (Calcutta) has pioneered a system of sewage disposal that is both efficient and environmentally friendly. Built to house one million people, Kolkata is now home to over 10 million, many living in slums. But the 8,000-hectare East Kolkata Wetlands Ramsar Site, a patchwork of tree-fringed canals, vegetable plots, rice paddies and fish ponds – and the 20,000 people that work in them – daily transform one-third of the city's sewage and most of its domestic refuse into a rich harvest of fish and fresh vegetables. For example, the Mudially Fishermen's Cooperative Society is a collective of 300 families that lease 70 hectares into which wastewater from the city is released. Through a series of natural

treatment processes – including the use of Eichhornia crassipes and other plants for absorbing oil, grease and heavy metals – the Cooperative has turned the area into a thriving fish farm and nature park. In 2005/06, the Cooperative sold fish worth over US$135,000 and shared income of more than US$55,000 among its members.[6][*not in citation given*]

Reservoirs of biodiversity

Wetland systems' rich biodiversity is becoming a focal point at International Treaty Conventions and within the World Wildlife Fund organization due to the high number of species present in wetlands, the small global geographic area of wetlands, the number of species which are endemic to wetlands, and the high productivity of wetland systems. Hundred of thousands of animal species, 20,000 of them vertebrates, are living in wetland systems. The discovery rate of fresh water fish is at 200 new species per year.

Biodiverse river basins. The Amazon holds 3,000 species of fresh water fish species within the boundaries of its basin whose function it is to disperse the seeds of trees. One of its key species, the Piramutaba catfish, *Brachyplatystoma vaillantii*, migrates more than 3,300 km (2,051 miles) from its nursery grounds near the mouth of the Amazon River to its spawning grounds in Andean tributaries (400 m or 437 yards above sea level) distributing plants seed along the route.

Productive intertidal zones. Intertidal mudflats have a similar productivity even while possessing a low number of species. The abundance of invertebrates found within the mud are a food source for migratory waterfowl.

Critical life-stage habitat. Mudflats, saltmarshes, mangroves, and seagrass beds contain bother species richness and productivity, and are home to important nursery areas for many commercial fish stocks.

Genetic Diversity. Many species in wetland systems are unique due to the long period of time that the ecosystem has been physically isolated from other aquatic sources. The number of endemic species in Lake Baikal in Russia classifies it as a hotspot for biodiversity and one of the most biodiverse wetlands in the entire world.

Lake Baikal. Evidence from a research study by Mazepova *et al.* suggest that the number of crustacean species endemic to Baikal Lake (>690 species and subspecies) exceeds the number of the same groups of animals inhabiting all the fresh water bodies of Eurasia together. Its 150 species of free-living Platyhelminthes alone is analogous to the entire number in all of Eastern Siberia. The 34 species and subspecies number of Baikal sculpins is more than twice the number of the analogous fauna that inhabits Eurasia. One of most exciting discoveries was made by A.V. Shoshin who registered about 300 species of free-living nematodes using only 6 near-shore sampling localities in the Southern Baikal. "If we will take into consideration, that about 60 % of the animals can be found nowhere else except Baikal, it may be assumed that the lake may be the biodiversity center of the Eurasian continent."[43]

Human Impact. Biodiversity loss occurs in wetland systems through land use changes, habitat destruction, pollution, exploitation of resources, and invasive species. Vulnerable, threatened, and endangered species number at 17% of waterfowl, 38% of fresh-water dependent mammals, 33% of fresh water fish, 26% of fresh water amphibians, 72% of fresh water turtles, 86% of marine turtles, 43% of crocodilians and 27% of coral reef-building species.

The impact of maintaining biodiversity is seen at the local level through job creation, sustainability, and community productivity. A good example is the Lower Mekong basin which runs through Cambodia, Laos, and Vietnam. Supporting over 55 million people, the sustainability of the region is enhanced through wildlife tours. The US state of Florida has estimated that US$ 1.6 billion was generated in state revenue from recreational activities associated with wildlife. Sustainable harvesting for medicinal remedies found in native wetlands plants in the Caribbean and Australia include the Red Mangrove *Rhizophora mangle* which possesses antibacterial, wound-healing, anti-ulcer effects, and antioxidant properties.[44]

Wetland products

Wetland systems naturally produce an array of vegetation and other ecological products that can harvested for personal and commercial use. The most significant of these is fish which have all or part of their life-cycle occur within a wetland system. Fresh and saltwater fish are the main source of protein for one billion people and comprise 15% of an additional two billion people's diets. In addition, fish generate a fishing industry that provides 80% of the income and employment to residents in developing countries. Another food staple found in wetland systems is rice, a popular grain that is consumed at the rate of 1/5 of the total global calorie count. In Bangladesh, Cambodia and Vietnam, where rice paddies are predominant on the landscape, rice consumption reach 70%.[44]

Food converted to sweeteners and carbohydrates include the sago palm of Asia and Africa (cooking oil), the nipa palm of Asia (sugar, vinegar, alcohol, and fodder) and honey collection from mangroves. More than supplemental dietary intake, this produce sustains entire villages. Coastal Thailand villages earn the key portion of their income from sugar production while the country of Cuba relocates more than 30,000 hives each year to track the seasonal flowering of the mangrove *Avicennia*.

Other mangrove-derived products:

- Fuelwood
- Salt (produced by evaporating seawater)
- Animal fodder
- Traditional medicines (e.g. from mangrove bark)
- Fibers for textiles
- Dyes and tannins

Human Impact. Over-fishing is the major problem for sustainable use of wetlands. The field of aquaculture within the fisheries industries is eliminating mass areas of wetland systems through practices seen such as in the shrimp farming industry's destruction of mangroves. Aquaculture is continuing to develop rapidly throughout the Asia-Pacific region specifically in China with world holdings in Asia equal to 90% of the total number of aquaculture farms and 80% of its global value.[44] Threats to rice fields mainly stem from inappropriate water management, introduction of invasive alien species, agricultural fertilizers, pesticides, and land use changes. Industrial-scale production of palm oil threatens the biodiversity of wetland ecosystems in parts of south-east Asia, Africa, and other developing countries. Exploitation can occur at the community level as is sometimes seen throughout coastal villages of Southern Thailand where each resident may obtain for themselves every consumable of the mangrove forest (fuelwood, timber, honey, resins, crab, and shellfish) which then becomes threatened through increasing population and continual harvest. Other issues that occur on a global level include an uneven contribution to climate change, point and non-point pollution, and air and water quality issues due to destructive wetland practices.

Wetlands and climate change

Wetlands perform two important functions in relation to climate change. They have mitigation effects through their ability to sink carbon, and adaptation effects through their ability to store and regulate water.[45] Wetlands store approximately 44.6 Tg C y^{-1} globally.[46] In salt marshes and mangrove swamps in particular, the average carbon sequestration rate is 201 g CO_2 m^{-2} y^{-1} while peatlands sequester approximately 20-30g CO_2 m^{-2} y^{-1}.[46][47]

Carbon sequestration has sponsored blue carbon initiatives.More on blue carbon and carbon sequestration

They also are emitters of nitrous oxide, a major greenhouse gas.

Scientific projections

"Low water and occasional drying of the wetland bottom during droughts (dry marsh phase) stimulate plant recruitment from a diverse seed bank and increase productivity by mobilizing nutrients. In contrast, high water during deluges (lake marsh phase) causes turnover in plant populations and creates greater interspersion of element cover and open water, but lowers overall productivity. During a cover cycle that ranges from open water to complete vegetation cover, annual net primary productivity may vary 20-fold."[48]

Nitrous oxide production from wetland soils

Coastal wetlands, such as tropical mangroves and temperate salt marshes are known to be sinks of carbon, therefore mitigating climate change, however they are also emitters of nitrous oxide (N_2O),[49] which is a greenhouse gas with a global warming potential 300 times that of carbon dioxide and the dominant ozone depleting substance emitted in the 21st century.[50] Anthropogenic greenhouse gas (GHG) emissions have rapidly increased to the atmosphere due the combustion of fossil fuels and deforestation practices, and these gases are major contributors to global climate change.

Although wetlands act as natural buffers towards nutrients expelled from surrounding watersheds, excess nutrients mainly through anthropogenic sources have been shown to significantly increase the N_2O fluxes fluxes from their soils through denitrification and nitrification processes (Table 1).[51][52] Anthropogenic sources of nutrients into waterways have increased substantially causing eutrophication especially in coastal systems.[53][54]The main sources of coastal eutrophication are industrially made nitrogen, which

is used as fertilizer in agricultural practices, as well as septic waste runoff.[55] Nitrogen (N) is the limiting nutrient for photosynthetic processes in saline systems, however in excess, it can lead to an overproduction of organic matter that then leads to hypoxic and anoxic zones within the water column.[56] Without oxygen, other organisms cannot survive, including economically important finfish and shellfish species. A study in the intertidal region of a New England salt marsh showed that excess levels of nutrients might increase N_2O emissions rather than sequester them.[51]

Table 1. Nitrous oxide fluxes from different wetland soils. Table adapted from Moseman-Valtierra (2012)[57] and Chen et al. (2010)[58]

Wetland type	Location	N_2O flux (μmol N_2O m^{-2} h^{-1})
Mangrove	Shenzhen and Hong Kong	0.14-23.83 [58]
Mangrove	Muthupet, South India	0.41-0.77 [59]
Mangrove	Bhitarkanika, East India	0.20-4.73 [60]
Mangrove	Pichavaram, South India	0.89-1.89 [60]
Mangrove	Queensland, Australia	-0.045-0.32 [61]
Mangrove	South East Queensland, Australia	0.091-1.48 [62]
Mangrove	Southwest coast, Puerto Rico	0.12-7.8 [63]
Mangrove	Magueyes Island, Puerto Rico	0.05-1.4 [63]
Salt marsh	Chesapeake Bay, USA	0.005-0.12 [64]
Salt marsh	Maryland, USA	0.1 [65]
Salt marsh	NE, China	0.1-0.16 [66]
Salt marsh	Biebrza, Poland	-0.07-0.06 [67]
Salt marsh	Netherlands	0.82-1.64 [68]
Salt marsh	Baltic sea	-0.13 [69]
Salt marsh	Massachusetts, USA	-2.14-1.27 [70]

Nitrous oxide fluxes from wetlands in the southern hemisphere is lacking as are ecosystem based studies including the role of dominant organisms that alter sediment biogeochemistry. Aquatic invertebrates produce ecologically relevant nitrous oxide emissions emissions due to ingestion of denitrifying bacteria that live within the subtidal sediment and water column [71] and thus may also be influencing nitrous oxide production within wetlands.

Conservation

Main article: Wetland conservation

Wetlands have historically been the victim of large draining efforts for real estate development, or flooding for use as recreational lakes. Since the 1970s, more focus has been put on preserving wetlands for their natural function yet by 1993 half the world's wetlands had been drained.[72]Wetlands provide a valuable flood control function. Wetlands are very effective at filtering and cleaning water pollution,[73] (often from agricultural runoff from the farms that replaced the wetlands in the first place). To replace these wetland ecosystem services enormous amounts of money had to be spent on water purification plants, along with the remediation measures for controlling floods: dam and levee construction.

In order to produce sustainable wetlands, short-term, private-sector profits need to come secondary to global equity. Decision-makers must valuate wetland type, provided ecosystem service, long-term benefit, and current subsidies inflating valuation on either the private or public sector side. Analysis using the impact of hurricanes versus storm protection features projected wetland valuation at US$33,000/hectare/year.[74]

Balancing wetland conservation with the needs of people

Wetlands are vital ecosystems that provide livelihoods for the millions of people who live in and around them. The Millennium Development Goals (MDGs) called for different sectors to join forces to secure wetland environments in the context of sustainable development and improving human wellbeing. A three-year project carried out by Wetlands International in partnership with the International Water Management Institute found that it is possible to conserve wetlands while improving the livelihoods of people living among them. Case studies conducted in Malawi and Zambia looked at how dambos – wet, grassy valleys or depressions where water seeps to the surface – can be farmed sustainably to improve livelihoods. Mismanaged or overused dambos often become degraded, however, using a knowledge exchange between local farmers and environmental managers, a protocol was developed using soil and water management practices. Project outcomes included a high yield of crops, development of sustainable farming techniques, and adequate water management generating enough water for use as irrigation. Before the project, there were cases where people had died from starvation due to food shortages. By the end of it, many more people had access to enough water to grow vegetables. A key achievement was that villagers had secure food supplies during long, dry months. They also benefited in other ways: nutrition was improved by growing a wider range of crops, and villagers could also invest in health and education by selling produce and saving money.[75]

Ramsar Convention

Main article: Ramsar Convention

The Convention on Wetlands of International Importance, especially as Waterfowl Habitat, or Ramsar Convention, is an international treaty designed to address global concerns regarding wetland loss and degradation. The primary purposes of the treaty are to list wetlands of international importance and to promote their wise use, with the ultimate goal of preserving the world's wetlands. Methods include restricting access to the majority portion of wetland areas, as well as educating the public to combat the misconception that wetlands are wastelands. The Convention works closely with five International Organisation Partners. These are: Birdlife International, IUCN, International Water Management Institute, Wetlands International and World Wide Fund for Nature. The partners provide technical expertise, help conduct or facilitate field studies and provide financial support. The IOPs also participate regularly as observers in all meetings of the Conference of the Parties and the Standing Committee and as full members of the Scientific and Technical Review Panel.

Valuation

The value of a wetland system to the earth and to humankind is one of the most important valuations that can be computed for sustainable development. A guideline involving assessing a wetland, keeping inventories of known wetlands, and monitoring the same wetlands over time is the current process that is used to educate environmental decision-makers such as governments on the importance of wetland protection and conservation.

Important considerations

- Constructed Wetlands take 10–100 years to fully resemble the vegetative composition of a natural wetland.
- Artificial wetlands do not have hydric soil. The soil has very low levels of organic carbon and total nitrogen compared to natural wetland systems.
- Organic matter can be added to degraded natural wetlands to help restore their productivity before the wetland is destroyed.[76]

Assessment

Rapid assessment.

Five steps to assessing a wetland

1. Collect general biodiversity data in order to inventory and prioritize wetland species, communities and ecosystems. Obtain baseline biodiversity information for a given area.
2. Gather information on the status of a focus or target species such as threatened species. Collect data pertaining to the conservation of a specific species.
3. Gain information on the effects of human or natural disturbance (changes) on a given area or species.
4. Gather information that is indicative of the general ecosystem health or condition of a specific wetland ecosystem.
5. Determine the potential for sustainable use of biological resources in a particular wetland ecosystem.

Inventory

Developing a global inventory of wetlands has proven to be a large and difficult undertaking. Current efforts are based on available data, but both classification and spatial resolution have proven to be inadequate for regional or site-specific environmental management decision-making. It is difficult to identify small, long, and narrow wetlands within the landscape. Many of today's remote sensing satellites do not have sufficient spatial and spectral resolution to monitor wetland conditions, although multispectral IKONOS and QuickBird data may offer improved spatial resolutions once it is 4 m or higher. Majority of the pixels are just mixtures of several plant species or vegetation types and are difficult to isolate which translates into an inability to classify the vegetation that defines the wetland. Improved remote sensing information, coupled with good knowledge domain on wetlands will facilitate expanded efforts in wetland monitoring and mapping. This will also be extremely important because we expect to see major shifts in species composition due to both anthropogenic land use and natural changes in the environment caused by climate change.

Monitoring

Mapping

A wetland system needs to be monitored over time to in order to assess whether it is functioning at an ecologically sustainable level or whether it is becoming degraded. Degraded wetlands will suffer a loss in water quality, a high number of threatened and endangered species, and poor soil conditions.

Due to the large size of wetlands, mapping is an effective tool to monitor wetlands. There are many remote sensing methods that can be used to map wetlands. Remote-sensing technology permits the acquisition of timely digital data on a repetitive basis. This repeat coverage allows wetlands, as well as the adjacent land-cover and land-use types, to be monitored seasonally and/or annually. Using digital data provides a standardized data-collection procedure and an opportunity for data integration within a geographic information system. Traditionally, Landsat 5 Thematic Mapper (TM), Landsat 7 Enhanced Thematic Mapper Plus (ETM +), and the SPOT 4 and 5 satellite systems have been used for this purpose. More recently, however, multispectral IKONOS and QuickBird data, with spatial resolutions of 4 m by 4 m and 2.44 m by 2.44 m, respectively, have been shown to be excellent sources of data when mapping and monitoring smaller wetland habitats and vegetation communities.

For example, Detroit Lakes Wetland Management District assessed area wetlands in Michigan, USA using remote sensing. Through using this technology, satellite images were taken over a large geographic area and extended period. In addition, using this technique was less costly and time-consuming compared to the older method using visual interpretation of aerial photographs. In comparison, most aerial photographs also require experienced interpreters to extract information based on structure and texture while the interpretation of remote sensing data only requires analysis of one characteristic (spectral).

However, there are a number of limitations associated with this type of image acquisition. Analysis of wetlands has proved difficult because to obtain the data it is often linked to other purposes such as the analysis of land cover or land use. Practically, many natural wetlands are difficult to monitor as these areas are quite often difficult to access and require exposure to native wildlife and potential endemic disease.

Further improvements

Methods to develop a classification system for specific biota of interest could assist with technological advances that will allow for identification at a very high accuracy rate. The issue of the cost and expertise involved in remote sensing technology is still a factor hindering further advancements in image acquisition and data processing. Future improvements in current wetland vegetation mapping could include the use of more recent and better geospatial data when it is available.

List of wetland types

Wetland types:[77]

- **A—Marine and Coastal Zone wetlands**

1. Marine waters—permanent shallow waters less than six metres deep at low tide; includes sea bays, straits
2. Subtidal aquatic beds; includes kelp beds, seagrasses, tropical marine meadows
3. Coral reefs
4. Rocky marine shores; includes rocky offshore islands, sea cliffs

5. Sand, shingle or pebble beaches; includes sand bars, spits, sandy islets
6. Intertidal mud, sand or salt flats
7. Intertidal marshes; includes saltmarshes, salt meadows, saltings, raised salt marshes, tidal brackish and freshwater marshes
8. Intertidal forested wetlands; includes mangrove swamps, nipa swamps, tidal freshwater swamp forests
9. Brackish to saline lagoons and marshes with one or more relatively narrow connections with the sea
10. Freshwater lagoons and marshes in the coastal zone
11. Non-tidal freshwater forested wetlands

- **B—Inland wetlands**

1. Permanent rivers and streams; includes waterfalls
2. Seasonal and irregular rivers and streams
3. Inland deltas (permanent)
4. Riverine floodplains; includes river flats, flooded river basins, seasonally flooded grassland, savanna and palm savanna
5. Permanent freshwater lakes (> 8 ha); includes large oxbow lakes
6. Seasonal/intermittent freshwater lakes (> 8 ha), floodplain lakes
7. Permanent saline/brackish lakes
8. Seasonal/intermittent saline lakes
9. Permanent freshwater ponds (< 8 ha), marshes and swamps on inorganic soils; with emergent vegetation waterlogged for at least most of the growing season
10. Seasonal/intermittent freshwater ponds and marshes on inorganic soils; includes sloughs, potholes; seasonally flooded meadows, sedge marshes
11. Permanent saline/brackish marshes
12. Seasonal saline marshes
13. Shrub swamps; shrub-dominated freshwater marsh, shrub carr, alder thicket on inorganic soils
14. Freshwater swamp forest; seasonally flooded forest, wooded swamps; on inorganic soils
15. Peatlands; forest, shrub or open bogs
16. Alpine and tundra wetlands; includes alpine meadows, tundra pools, temporary waters from snow melt
17. Freshwater springs, oases and rock pools
18. Geothermal wetlands
19. Inland, subterranean karst wetlands

- **C—Human-made wetlands**

1. Water storage areas; reservoirs, barrages, hydro-electric dams, impoundments (generally > 8 ha)
2. Ponds, including farm ponds, stock ponds, small tanks (generally < 8 ha)
3. Aquaculture ponds; fish ponds, shrimp ponds
4. Salt exploitation; salt pans, salines
5. Excavations; gravel pits, borrow pits, mining pools
6. Wastewater treatment; sewage farms, settling ponds, oxidation basins
7. Irrigated land and irrigation channels; rice fields, canals, ditches
8. Seasonally flooded arable land, farm land

Wetland names

Variations of names for wetland systems:

• Swamp ◦ Freshwater swamp forest ◦ Coniferous swamp ◦ Peat swamp forest	• Flooded grasslands and savannas • Constructed wetland • Riparian zone • Mire ◦ Fen ◦ Bog • Vernal pool	

	Environment portal
	Ecology portal

References

1. ^ [a] [b] "History of the Everglades | Everglades Forever | Florida DEP". Dep.state.fl.us. 2009-02-11. Retrieved 2012-05-23.
2. ^ "Department of Environmental Protection State of Florida Glossary". State of Florida. Retrieved 2011-09-25.
3. ^ Butler S., ed. (2010). *Macquarie Concise Dictionary* (5th ed.). Sydney Australia: Macquarie Dictionary Publishers Ltd. ISBN 978-1-876429-85-0.
4. ^ [a] [b] "Official page of the Ramsar Convention". Retrieved 2011-09-25.
5. ^ [a] [b] [c] [d] [e] [f] Keddy, Paul A. (2010). *Wetland ecology : principles and conservation* (2nd ed.). New York: Cambridge University Press. p. 497. ISBN 978-0521519403.
6. ^ [a] [b] [c] [d] [e] [f] [g] [h] "Ramsar Convention Ecosystem Services Benefit Factsheets". Retrieved 2011-09-25.
7. ^ [a] [b] [c] "US EPA". Retrieved 2011-09-25.
8. ^ [a] [b] Fraser, L; Keddy, PA, ed. (2005). *The world's largest wetlands : their ecology and conservation*. Cambridge, UK [u.a.]: Cambridge Univ. Press. p. 488. ISBN 978-0521834049.
9. ^ "WWF Pantanal Programme". Retrieved 2011-09-25.
10. ^ "Carpinteria Water District: Glossary of Terms". Cvwd.net. Retrieved 2012-05-23.[*dead link*]
11. ^ "Glossary". Mapping2.orr.noaa.gov. Retrieved 2012-05-23.
12. ^ "Glossary". Alabama Power. Retrieved 2012-05-23.
13. ^ Mitsch, William J.; James G. Gosselink (2007-08-24). *Wetlands* (4th ed.). New York: John Wiley & Sons. ISBN 978-0-471-69967-5.
14. ^ Keddy (2010), p. 2.
15. ^ "The Ramsar 40th Anniversary Message for November". Ramsar. Retrieved 2011-10-10.
16. ^ EPA Regulations listed at 40 CFR 230.3(t). Retrieved 2014-02-18.
17. ^ Richardson, JL, Arndt, JL & Montgomery, JA 2001, 'Hydrology of wetland and related soils' in JL Richardson & MJ Vepraskas (eds), Wetland Soils, Lewis Publishers, Boca Raton
18. ^ Vitt, DH & Chee, W 1990, 'The relationships of vegetation to surface water chemistry and peat chemistry in fens of Alberta, Canada', Plant Ecology, vol. 89, no. 2, pp. 87-106.
19. ^ B.R. Silliman, E.D. Grosholz, and M.D. Bertness (eds.) 2009. Human Impacts on Salt Marshes. A Global Perspective. University of California Press, Berkeley, CA.
20. ^ Smith, MJ, Schreiber, ESG, Kohout, M, Ough, K, Lennie, R, Turnbull, D, Jin, C & Clancy, T 2007 'Wetlands as landscape units: spatial patterns in salinity and water chemistry, Wetlands, Ecology & Management, vol. 15, no. 2, pp. 95-103.
21. ^ Ponnamperuma, FN 1972, 'The chemistry of submerged soils', Advances in Agronomy, vol. 24, pp. 29–96.
22. ^ Moore Jr., PA & Reddy, KR 1994, 'Role of Eh and pH on phosphorus geochemistry in sediments of Lake Okeechobee, Florida' Journal of Environmental Quality, vol. 23, pp. 955–964.
23. ^ Minh LQ, Tuong TP, van Mensvoort MEF, Bouma J 1998 'Soil and water table management effects on aluminum dynamics in an acid sulphate soil in Vietnam', Agriculture, Ecosystems & Environment, vol. 68, no. 3, pp. 255–262.
24. ^ Schlesinger, WA 1997, Biogeochemistry: An analysis of global change, 2nd edn, Academic Press, San Diego

25. ^ Bedford, BL 1996, 'The need to define hydrologic equivalence at the landscape scale for freshwater wetland mitigation', Ecological Applications, vol. 6, no. 1, pp. 57-68.
26. ^ Nelson, ML, Rhoades, CC & Dwire, KA 2011, 'Influences of Bedrock Geology on Water Chemistry of Slope Wetlands and Headwaters Streams in the Southern Rocky Mountains', Wetlands, vol. 31, pp. 251-261.
27. ^ "Blacktown Council wetlands". Retrieved 2011-09-25.[*dead link*]
28. ^ Hutchinson, G. E. 1975. A Treatise on Limnology, Vol. 3, Limnological Botany. New York: John Wiley
29. ^ Hutchinson, G. E. 1975. A Treatise on Limnology, Vol. 3, Limnological Botany. New York: John Wiley.
30. ^ Hughes, F.M.R. (ed.). 2003. The Flooded Forest: Guidance for policy makers and river managers in Europe on the restoration of floodplain forests. FLOBAR2, Department of Geography, University of Cambridge, Cambridge, UK. 96 p
31. ^ Wilcox, D.A, Thompson, T.A., Booth, R.K. and Nicholas, J.R. 2007. Lake-level variability and water availability in the Great Lakes. USGS Circular 1311. 25 p.
32. ^ Goulding, M. 1980. The Fishes and the Forest:Explorations in Amazonian Natural History. Berkeley, CA: University of California Press
33. ^ "Frogs | Bioindicators". Savethefrogs.com. 2011-04-29. Retrieved 2014-01-21.
34. ^ "Taken from Australian fauna". Australian fauna. Retrieved 2012-05-23.
35. ^ "Taken from Blacktown Council Wetland Inventory". Blacktown.nsw.gov.au. Retrieved 2012-05-23.[*dead link*]
36. ^ "Ramsar Convention Technical Reports".
37. ^ *a b* "Wetlands International works to sustain and restore wetlands for people and biodiversity". Wetlands.org. Retrieved 2014-01-21.
38. ^ [1][*dead link*]
39. ^ "United Nations Environment Programme (UNEP) – Home page". Retrieved 2011-12-11.
40. ^ Brix, H 1993, 'Wastewater treatment in constructed wetlands: system design, removal processes, and treatment performance' in AG Moshiri (ed), Constructed Wetlands for Water Quality Improvement, CRC Press, Boca Raton, Florida.
41. ^ Vymazal, J & Kröpfleova, L 2008, 'Wastewater treatment in constructed wetlands with horizontal sub-surface flow', Environmental Pollution, vol. 14.
42. ^ "Arcata, California Constructed Wetland: A Cost-Effective Alternative for Wastewater Treatment". Ecotippingpoints.org. Retrieved 2012-05-23.
43. ^ Timoshkin O.A, ed. (2004). "Index of animal species inhabiting Lake Baikal and its catchment area. Guides and Keys to Identification of Fauna and Flora of Lake Baikal". *2* **1** (1st ed.) (Novosibirsk, Nauka: John Wiley & Sons). ISBN 5-02-031736-5.
44. ^ *a b c* "The Ramsar Information Sheet on Wetlands of International Importance". September 18, 2009. Retrieved November 19, 2011.
45. ^ U.S. EPA, 2009, Synthesis of Adaptation Options for Coastal Areas, Washington DC US, Environmental Protection Agency, Climate Ready Estuaries Program, EPA 430-F-08-024
46. ^ *a b* Chmura, G. L. (2003). Global carbon sequestration in tidal, saline wetland soils. Global Biogeochemical Cycles, 17(4)
47. ^ Roulet, N. T. (2000). Peatlands, Carbon Storage, Greenhouse Gases, And The Kyoto Protocol: Prospects And Significance For Canada. Wetlands, 20(4), 605-615.
48. ^ Johnson W C, Millett B V, Gilmanov T, Voldseth R A, Guntenspergen G R & Naugle D E 2005 Vulnerability of Northern Prairie Wetlands to Climate Change Bio Science 10: 863-872
49. ^ Thompson, A. J., Giannopoulos, G., Pretty, J., Baggs, E. M., & Richardson, D. J. (2012). Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. Philosophical Transactions of the Royal Society B, 367, 1157-1168
50. ^ Nitrous Oxide (N2O): The Dominant Ozone-Depleting Substance Emitted in the 21st Century. A. R. Ravishankara, John S. Daniel, and Robert W. Portmann Science 2 October 2009: 326 (5949), 123-125.Published online 27 August 2009
51. ^ *a b* Moseman -Valtierra S, et al. (2011). Short-term nitrogen additions can shift a coastal wetland from a sink to a source of N2O. Atmospheric Environment 45: 4390–4397.
52. ^ Bange, H. W., Nitrous oxide and methane in European coastal waters. Estuarine Coastal and Shelf Science 2006, 70 (3), 361-374.
53. ^ Valiela, I.; Collins, G.; Kremer, J.; Lajtha, K.; Geist, M.; Seely, B.; Brawley, J.; Sham, C. H., Nitrogen loading from coastal watersheds to receiving estuaries: New method and application. Ecological Applications 1997, 7 (2), 358-380.

54. ^ Nixon, S. W., Nutrients and the productivity of estuarine and coastal marine ecosystems. Journal of the Limnological Society of South Africa, 1986, 12:43–71.
55. ^ Galloway J. The Nitrogen Cascade. Bioscience. 2003;53:341-356.
56. ^ Diaz, R. J., & Rosenberg, R. (2008). Spreading Dead Zones and Consequences for Marine Ecosystems. Science, 321(5891), 926-929
57. ^ Moseman-Valtierra, S. (2012). Chapter 1: Reconcidering the climactic roles of marshes: Are they sinks or sources of greenhouse gases? In D. C. Abreu & S. L. Borbón (Authors), Marshes: Ecology, management and conservation. New York: Nova Science.
58. ^ *a b* Chen, G., Tam, N., & Ye, Y. (2010). Summer fluxes of atmospheric greenhouse gases N2O, CH4 and CO2 from mangrove soil in South China. Science of The Total Environment, 408(13), 2761-2767.
59. ^ Krithika K, Purvaja R, Ramesh R. Fluxes of methane and nitrous oxide from an Indian mangrove. Curr Sci 2008;94:218–24.
60. ^ *a b* Chauhan R, Ramanathan AL, Adhya TK. Assessment of methane and nitrous oxide flux from mangroves along Eastern coast of India. Geofluids 2008;8:321–32.
61. ^ Kreuzwieser J, Buchholz J, Rennenberg H. Emission of methane and nitrous oxide by Australian mangrove ecosystems. Plant Biol 2003;5:423–31.
62. ^ Allen DE, Dalal RC, Rennenberg H, Louise Meyer R, Reeves S, Schmidt S. Spatial and temporal variation of nitrous oxide and methane flux between subtropical mangrove soils and the atmosphere. Soil Biol Biochem 2007;39:622–31.
63. ^ *a b* Sotomayor D, Corredor JE, Morell JM. Methane flux from mangrove soils along the southwestern coast of Puerto Rico. Estuaries 1994;17:140–7.
64. ^ Jordan, T. E., Andrews, M. P., Szuch, R. P., Whigham, D. F., Weller, D. E., & Jacobs, A. D. (2007). Comparing Functional Assessments Of Wetlands To Measurements Of Soil Characteristics And Nitrogen Processing. Wetlands, 27(3), 479-497.
65. ^ Weller, D. E., Cornell, D. L., & Jordan, T. E. (1994). Denitrification in riparian forests receiving agricultural discharges. Global Wetlands: Old World and New, 117-131.
66. ^ Yu, J., Liu, J., Wang, J., Sun, W., Patrick, W. H., & Meixner, F. X. (2007). Nitrous Oxide Emission from Deyeuxia angustifolia Freshwater Marsh in Northeast China. Environmental Management, 40(4), 613-622.
67. ^ Roobroeck, D., Butterbach-Bahl, K., Brüggemann, N., & Boeckx, P. (2010). Dinitrogen and nitrous oxide exchanges from an undrained monolith fen: Short-term responses following nitrate addition. European Journal of Soil Science, 61(5), 662-670.
68. ^ Hefting, M. M., Bobbink, R., & De Caluwe, H. (2003). Nitrous Oxide Emission and Denitrification in Chronically Nitrate-Loaded Riparian Buffer Zones. Journal of Environment Quality, 32(4), 1194.
69. ^ Liikanen, A. (2009). Methane and nitrous oxide fluxes in two coastal wetlands in the northeastern Gulf of Bothnia, Baltic Sea. Boreal environment research, 14(3), 351-368.
70. ^ Moseman-Valtierra S, et al. (2011). Short-term nitrogen additions can shift a coastal wetland from a sink to a source of N2O. Atmospheric Environment 45: 4390–4397.
71. ^ Stief, P., Poulsen, M., Nielsen, et al. (2009). Nitrous oxide emission by aquatic macrofauna. Proceedings of the National Academy of Sciences, 106(11), 4296-4300
72. ^ "unknown title". *New Scientist* (1894): 46. 1993-10-09.
73. ^ "Letting Nature Do the Job". Wild.org. 2008-08-01. Retrieved 2012-05-23.
74. ^ "FAO". Retrieved 2011-09-25.[*dead link*]
75. ^ "Good practices and lessons learned in integrating ecosystem conservation and poverty reduction objectives in wetlands". The Ramsar Convention on Wetlands. December 1, 2008. Retrieved November 19, 2011.
76. ^ Hart T M & Davis S E, 2011, Wetland development in a previously mined landscape of East Texas, USA Wetlands Ecological Management, 19: 317-329
77. ^ "A Directory of Important Wetlands in Australia". Environment.gov.au. 2009-07-27. Retrieved 2012-05-23.

Further reading

- Mitsch, W.J. and J.G. Gosselink. (1986, 1993, 2000, 2007). Wetlands, 1st, 2nd, 3rd and 4th editions. Van Nostrand Reinhold, New York, New York and John Wiley & Sons, Inc., Hoboken, New Jersey.
- Evangelisti, Domenico D'Amelia, Gustavo Di Lallo, Maria Cristina Thaller, Luciana Migliore. 2013. Relationship between Salinity and Bacterioplankton in Three Relic Coastal Ponds (Macchiatonda Wetland, Italy). Vol.5 No.9 Journal of Water Resource and Protection *[2]

- Brinson, M. (1993) A Hydrogeomorphic Classification of Wetlands
- 1987 U.S. Army Corps of Engineers Wetland delineation manual
- Dugan, Patrick (editor) (1993) *Wetlands in Danger*, World Conservation Atlas Series.
- Terra Nuova East Africa. Wetlands in drylands.
- Fredrikson, Leigh H. (1983) "Wetlands: A Vanishing Resource" – Yearbook of Agriculture
- Fraser, L.H. and P.A. Keddy (eds.). 2005. The World's Largest Wetlands: Ecology and Conservation. Cambridge University Press, Cambridge, UK. 488 p.
- Ghabo, A. A. (2007) Wetlands Characterization; Use by Local Communities and Role in Supporting Biodiversity in the Semiarid Ijara District, Kenya.
- Keddy, P.A. 2010. Wetland Ecology: Principles and Conservation (2nd edition). Cambridge University Press, Cambridge, UK. 497 pp.
- MacKenzie W.H. and J.R. Moran (2004) "Wetlands of British Columbia: A Guide to Identification". Ministry of Forests, Land Management Handbook 52.
- Maltby E. and Barker T. (eds) (2009) The Wetlands Handbook. Wiley-Blackwell, Oxford. 1058 pp.
- Mitsch, W.J., J.G. Gosselink, C.J. Anderson, and L. Zhang. (2009) "Wetland Ecosystems". John Wiley & Sons, Inc., Hoboken, 295 pp.
- Romanowski, N 2013,"*Living Waters*", CSIRO Publishing, Melbourne, ISBN 9780643107564

External links

- Short public tv program on Florida Everglades

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Wetlands. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 6.6: Wetlands is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

7: The Ocean

7.1: Introduction to the Oceans
7.2: Composition of Ocean Water
7.3: The Seafloor
7.4: Ocean Currents
7.5: Oceans and Coastal Environments
7.6: Tides
7.7: Currents

This page titled 7: The Ocean is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.1: Introduction to the Oceans

Have you ever heard the Earth called the “Blue Planet”? This term makes sense, because over 70% of the surface of the Earth is covered with water. The vast majority of that water (97.2%) is in the oceans. Without all that water, our world would be a different place. The oceans are an important part of Earth: they help to determine the make-up of the air, they help determine the weather and temperature, and they support great amounts of life. The composition of ocean water is unique to its location and depth. Just as Earth’s interior is divided into layers, the ocean separated into different layers, called the water column.

Lesson Objectives

- Describe how the oceans formed.
- Explain the significance of the oceans.
- Describe the composition of ocean water.
- Define the parts of the water column and oceanic divisions.

How the Oceans Formed

Scientists have developed a number of hypotheses about how the oceans formed. Though these hypotheses have changed over time, one idea now has the wide support of Earth scientists, called the volcanic outgassing theory. This means that water vapor given off by volcanoes erupting over millions or billions of years, cooled and condensed to form Earth’s oceans.

Creation and Collection of Water

When the Earth was formed 4.6 billion years ago, it would never have been called the Blue Planet. There were no oceans, there was no oxygen in the atmosphere, and no life. But there were violent collisions, explosions, and eruptions. In fact, the Earth in its earliest stage was molten. This allowed elements to separate into layers within the Earth — gravity pulled denser elements toward the Earth’s center, while less dense materials accumulated near the surface. This process of separation created the layers of the Earth as we know them.

As temperatures cooled, the surface solidified and an atmosphere was created. Volcanic eruptions released water vapor from the Earth’s crust, while more water came from asteroids and comets that collided with the Earth (Figure 14.1). About 4 billion years ago, temperatures cooled enough for oceans to begin forming.



Figure 14.1: Volcanic activity was common in Earth’s early stages, when the oceans had not yet begun to form.

Present Ocean Formation

As you know, the continents were not always in the same shape or position as they are today. Because of tectonic plate movements, land masses have moved about the Earth since they were created. About 250 million years ago, all of the continents were arranged in one huge mass of land called Pangea (Figure 14.2). This meant that most of Earth’s water was collected in a huge ocean called Panthalassa. By about 180 million years ago, Pangea had begun to break apart because of continental drift. This then separated the Panthalassa Ocean into separate but connected oceans that are the ocean basins we see today on Earth.




Figure 14.2: Pangea was the sole landform 250 million years ago, leaving a huge ocean called Panthalassa, along with a few smaller oceans.

Significance of the Oceans

The Earth's oceans play an important role in maintaining the world as we know it. Indeed, the ocean is largely responsible for keeping the temperatures on Earth fairly steady. It may get pretty cold where you live in the wintertime. Some places on Earth get as cold as -70°C. Some places get as hot as 55°C. This is a range of 125°C. But compare that to the surface temperature on Mercury: it ranges from -180°C to 430°C, a range of 610°C. Mercury has neither an atmosphere nor an ocean to buffer temperature changes so it gets both extremely hot and very cold.



Figure 14.3: Coral reefs are amongst the most densely inhabited and diverse areas on the globe.

On Earth, the oceans absorb heat energy from the Sun. Then the ocean currents move the energy from areas of hot water to areas of cold water, and vice versa. Not only does ocean circulation keep the water temperature moderate, but it also affects the temperature of the air. If you examine land temperatures on the Earth, you will notice that the more extreme temperatures occur in the middle of continents, whereas temperatures near the water tend to be more moderate. This is because water retains heat longer than land. Summer temperatures will therefore not be as hot, and winter temperatures won't be as cold, because the water takes a long time to heat up or cool down. If we didn't have the oceans, the temperature range would be much greater, and humans could not live in those harsh conditions.

The ocean is home to an enormous amount of life. This includes many kinds of microscopic life, plants and algae, invertebrates like sea stars and jellyfish, fish, reptiles, and marine mammals. The many different creatures of the ocean form a vast and complicated food web, that actually makes up the majority of all biomass on Earth. (**Biomass** is the total weight of living organisms in a particular area.) We depend on the ocean as a source of food and even the oxygen created by marine plants. Scientists are still discovering new creatures and features of the oceans, as well as learning more about marine ecosystems (Figure 14.3).

Finally, the ocean provides the starting point for the Earth's water cycle. Most of the water that evaporates into the atmosphere initially comes from the ocean. This water, in turn, falls on land in the form of precipitation. It creates snow and ice, streams and ponds, without which people would have little fresh water. A world without oceans would be a world without you and me.

Composition of Ocean Water

Water has oftentimes been referred to as the "universal solvent", because many things can dissolve in water (Figure 14.4). Many things like salts, sugars, acids, bases, and other organic molecules can be dissolved in water. Pollution of ocean water is a major problem in some areas because many toxic substances easily mix with water.




Figure 14.4: Ocean water is composed of many substances. The salts include sodium chloride, magnesium chloride, and calcium chloride.

Perhaps the most important substance dissolved in the ocean is salt. Everyone knows that ocean water tastes salty. That salt comes from mineral deposits that find their way to the ocean through the water cycle. Salts comprise about 3.5% of the mass of ocean water. Depending on specific location, the salt content or **salinity** can vary. Where ocean water mixes with fresh water, like at the mouth of a river, the salinity will be lower. But where there is lots of evaporation and little circulation of water, salinity can be much higher. The Dead Sea, for example, has 30% salinity—nearly nine times the average salinity of ocean water. It is called the Dead Sea because so few organisms can live in its super salty water.

The density (mass per volume) of seawater is greater than that of fresh water because it has so many dissolved substances in it. When water is more dense, it sinks down to the bottom. Surface waters are usually lower in density and less saline. Temperature affects density too. Warm water is less dense and colder waters are more dense. These differences in density create movement of water or deep ocean **currents** that transport water from the surface to greater depths.

The Water Column

In 1960, one of the deepest parts of the ocean (10,910 meters) was reached by two men in a specially designed submarine called the Trieste (Figure 14.5). This part of the ocean has been named the Challenger Deep. In contrast, the average depth of the ocean is 3,790 meters—still an incredible depth for sea creatures to live at and for humans to travel. What makes it so hard to live at the bottom of the ocean? There are three major factors—the absence of light, low temperature, and extremely high pressure. In order to better understand regions of the ocean, the scientists define different regions by depth (Figure 14.6).




Figure 14.5: The Trieste made a record dive to the Challenger Deep in 1960. James Cameron repeated this feat in 2012.




Figure 14.6: The ocean environment is divided into many regions based on factors like availability of light and nutrients. Organisms adapt to the conditions and resources in the regions in which they live.

Sunlight only penetrates water to a depth of about 200 meters, a region called the **photic zone** (photic means light). Since organisms that photosynthesize depend on sunlight, they can only live in the top 200 meters of water. Such photosynthetic organisms supply almost all the energy and nutrients to the rest of the marine food web. Animals that live deeper than 200 meters mostly feed on whatever drops down from the photic zone.

Beneath the photic zone is the **aphotic zone**, where there is not enough light for photosynthesis. The aphotic zone makes up the majority of the ocean but a minority of its life forms. Descending to the ocean floor, the water temperature decreases while pressure increases tremendously. Each region is progressively deeper and colder, with the very deepest areas in ocean trenches.

The ocean can also be divided by horizontal distance from the shore. Nearest to the shore lies the **intertidal zone**. In this region, you might find waves, changes in tide, and constant motion in the water that exposes the water to large amounts of air. Organisms that live in this zone are adapted to withstand waves and exposure to air in low tides, by having strong attachments and hard shells. The **neritic zone** includes the intertidal zone and the part of the ocean floor that very gradually slopes downward, the continental shelf. Lots of oceanic plants live in this zone, since some sunlight still penetrates to the bottom of the ocean floor in the neritic zone. Beyond the neritic zone is the **oceanic zone**, where the sloping sea floor takes a much even steeper dive and sunlight does not reach. Animals such as sharks, fish, and whales can be found in this zone. They feed on materials that sink from upper levels, or consume one another. At hydrothermal vents, areas of extremely hot water with lots of dissolved materials allow rare and unusual producers to thrive.

Lesson Summary

- Our oceans originally formed as a water vapor released by volcanic outgassing cooled and condensed.
- The oceans serve the very important role of helping to moderate Earth's temperatures.
- The oceans are home to a tremendous diversity of life, and algae which are all photosynthetic organisms.
- The main elements dissolved in seawater are chlorine, sodium, magnesium, sulfate and calcium.
- Usual salinity for the oceans is about 3.5% or 35 parts per thousand.
- Some regions in areas of high evaporation, like the Dead Sea, have exceptionally high salinities.
- The photic zone is the surface layer of the oceans, down to about 200m, where there is enough available light for photosynthesis.
- Below the photic zone, the vast majority of the oceans lies within the aphotic zone, where there is not enough light for photosynthesis.
- On average, the ocean floor is about 3,790m but there are ocean trenches as deep as 10,910m.
- The ocean has many biological zones determined by availability of different abiotic factors.
- Neritic zones are nearshore areas, including the intertidal zone. Oceanic zones are offshore regions of the ocean.

Review Questions

1. What was the name of the single continent that separated to form today's continents?
2. From what three sources did water originate on Earth?
3. What percent of the Earth's surface is covered by water?
4. How do the oceans help to moderate Earth's temperatures?
5. Over time, the Earth's oceans have become more and more salty. Why?
6. What is the most common substance that is dissolved in ocean water?
7. What is density?

Vocabulary

aphotic zone

The zone in the water column deeper than 200 m. Sunlight does not reach this region of the ocean.

biomass

The total mass of living organisms in a certain region.

current

The movement of water in a stream, lake, or ocean.

density

Mass per volume. The units for density are usually $^{g}/_{cm^3}$ or $^{g}/_{mL}$.

intertidal zone

The part of the ocean closest to the shore, between low and high tide.

neritic zone

The part of the ocean where the continental shelf gradually slopes outward from the edge of the continent. Some sunlight can penetrate this region of the ocean.

oceanic zone

The open ocean, where the seafloor is deep. No sunlight reaches the floor of the ocean here.

Pangea

The supercontinent that tectonically broke apart about 200 million years ago, forming the continents and oceans that we see today on Earth.

photic zone

The topmost region of the water column, extending from the surface down to about 200 m in depth. Sunlight easily penetrates this region of the water column.

salinity

A measure of the amount of dissolved salt in water.

water column

A vertical column of ocean water, which is divided into different zones according to their depth.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Introduction_to_the_Oceans. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 7.1: Introduction to the Oceans is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.2: Composition of Ocean Water

Composition of Ocean Water

Water has oftentimes been referred to as the “universal solvent”, because many things can dissolve in water (Figure 14.4). Many things like salts, sugars, acids, bases, and other organic molecules can be dissolved in water. Pollution of ocean water is a major problem in some areas because many toxic substances easily mix with water.




Figure 14.4: Ocean water is composed of many substances. The salts include sodium chloride, magnesium chloride, and calcium chloride.

Perhaps the most important substance dissolved in the ocean is salt. Everyone knows that ocean water tastes salty. That salt comes from mineral deposits that find their way to the ocean through the water cycle. Salts comprise about 3.5% of the mass of ocean water. Depending on specific location, the salt content or **salinity** can vary. Where ocean water mixes with fresh water, like at the mouth of a river, the salinity will be lower. But where there is lots of evaporation and little circulation of water, salinity can be much higher. The Dead Sea, for example, has 30% salinity—nearly nine times the average salinity of ocean water. It is called the Dead Sea because so few organisms can live in its super salty water.

The density (mass per volume) of seawater is greater than that of fresh water because it has so many dissolved substances in it. When water is more dense, it sinks down to the bottom. Surface waters are usually lower in density and less saline. Temperature affects density too. Warm water is less dense and colder waters are more dense. These differences in density create movement of water or deep ocean **currents** that transport water from the surface to greater depths.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Introduction_to_the_Oceans#Composition_of_Ocean_Water. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 7.2: Composition of Ocean Water is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.3: The Seafloor

The ocean surface is vast and hides an entire world underneath it. The ocean floor is sometimes called the final frontier of the modern era. Though people have traveled on the ocean for millennia, people have explored only a tiny fraction of the ocean floor. We know very little about the vast expanse of our oceans. Today's technology has allowed us to learn more about the seafloor, including both its physical properties and its effects on living organisms.

Lesson Objectives

- Describe the obstacles to studying the seafloor and methods for doing so.
- Describe the features of the seafloor.

Ancient myth says that Atlantis was a powerful undersea city whose warriors conquered many parts of Europe. There is little proof that such a city existed, but human fascination with the world under the oceans certainly has existed for centuries. Not much was known about the aphotic zone of the ocean until scientists developed a system modeled after the way that bats and dolphins use echolocation to navigate in the dark (Figure 14.19). Prompted by the need to find submarines during World War II, scientists learned to bounce sound waves through the ocean to detect underwater objects. The sound waves bounce back like an echo off of whatever object may be in the ocean. The distance of the object can be calculated based on the time that it takes for the sound waves to return. Finally, scientists were able to map the ocean floor.



Figure 14.19: Dolphins and whales use echolocation, a natural sonar system, to navigate the ocean.

Three main obstacles have kept us from studying the depths of the ocean: absence of light, very cold temperatures, and high pressure. As you know, light only penetrates the top 200 meters of the ocean; the depths of the ocean can be as much as 11,000 meters deep. Most places in the ocean are completely dark, which makes it impossible for humans to explore without bringing a source of light with them. Secondly, the ocean is very cold; colder than 0°C (32°F) in many places. Such cold temperatures pose significant obstacles to human exploration of the oceans. Finally, the pressure in the ocean increases tremendously as you go deeper. Scuba divers can rarely go deeper than 40 meters due to the pressure. The pressure on a diver at 40 meters would be 4 kilograms/square centimeter (60 lbs/sq in). Even though we don't think about it, the air in our atmosphere has weight. It presses down on us with a force of about 1 kilogram per square centimeter (14.7 lbs/ sq in). In the ocean, for every 10 meters of depth, the pressure increases by nearly 1 atmosphere! Imagine the pressure at 10,000 meters; that would be 1,000 kilograms per square centimeter (14,700 lbs/sq in). Today's submarines usually dive to only about 500 meters; to go deeper than this they must be specially designed for greater depth (Figure 14.20).



Figure 14.20: Submarines are built to withstand great pressure under the sea, up to 680 atmospheres of pressure (10,000 pounds per square inch). They still rarely dive below 400 meters.




Figure 14.21: Alvin allows for a nine hour dive for up to two people and a pilot. It was commissioned in the 1960s.

In the 19th century, explorers mapped ocean floors by painstakingly dropping a line over the side of a ship to measure ocean depths, one tiny spot at a time. SONAR, which stands for *So*und *N*avigation *A*nd *R*anging, has enabled modern researchers to map the ocean floor much more quickly and easily. Researchers send a pulse of sound down to the ocean floor and calculate the depth based on how long it takes the sound to return. Of course, some scientific research requires actually traveling to the bottom of the ocean to collect samples or directly observe the ocean floor, but this is more expensive and can be dangerous.

In the late 1950s, the bathyscaphe (deep boat) *Trieste* was the first manned vehicle to venture to the deepest parts of the ocean, a region of the Marianas Trench named the Challenger Deep. It was built to withstand 1.2 metric tons per square centimeter and plunged to a depth of 10,900 meters. No vehicle has carried humans again to that depth, though robotic submarines have returned to collect sediment samples from the Challenger Deep. *Alvin* is a submersible used by the United States for a great number of studies; it can dive up to 4,500 meters beneath the ocean surface (Figure 14.21).

In order to avoid the expense, dangers and limitations of human missions under the sea, remotely operated vehicles or ROVs, allow scientists to study the ocean's depths by sending vehicles carrying cameras and special measuring devices. Scientists control them electronically with sophisticated operating systems (Figure 14.22).



Figure 14.22: Remotely-operated vehicles like this one allow scientists to study the seafloor.

Features of the Seafloor

Before scientists invented sonar, many people believed the ocean floor was a completely flat surface. Now we know that the seafloor is far from flat. In fact, the tallest mountains and deepest canyons are found on the ocean floor; far taller and deeper than any landforms found on the continents. The same tectonic forces that create geographical features like volcanoes and mountains on land create similar features at the bottom of the oceans.

Look at Figure 14.23. If you follow the ocean floor out from the beach at the top left, the seafloor gently slopes along the **continental shelf**. The sea floor then drops off steeply along the **continental slope**, the true edge of the continent. The smooth, flat regions that make up 40% of the ocean floor are the **abyssal plain**. Running through all the world's oceans is a continuous mountain range, called the **mid-ocean ridge**("submarine ridge" in Figure 14.23). The mid-ocean ridge is formed where tectonic plates are moving apart from each other, allowing magma to seep out in the space where the plates pulled apart. The mid-ocean ridge system is 80,000 kilometers in total length and mostly underwater except for a few places like Iceland. Other underwater mountains include undersea volcanoes (called **seamounts**), which may rise more than 1,000 meters above the ocean floor. Those that reach the surface become volcanic islands, such as the Hawaiian Islands. Deep oceanic **trenches**are created where a tectonic plate dives beneath (subducts) another plate.




Figure 14.23: The seafloor is as varied a landscape as the continents.

Lesson Summary

- Until the development of sonar, we knew very little about the ocean floor.
- The deep ocean is dark, very cold and has tremendous pressure from the overlying water.
- Scuba divers can explore only to about 40 meters, while most submarines dive only to about 500 meters. Scientific research submersibles have explored the ocean's deepest trenches, but most are designed to reach only the ocean floor.
- Today much of our exploration of the oceans happens using sonar and remotely operated vehicles.
- Features of the ocean include the continental shelf, slope, and rise. The ocean floor is called the abyssal plain. Below the ocean floor, there are a few small deeper areas called ocean trenches. Features rising up from the ocean floor include seamounts, volcanic islands and the mid-oceanic ridges and rises.

Review Questions

1. What are three obstacles to studying the seafloor?
2. The atmospheric pressure is about 1 kilogram per centimeter squared (14.7 pounds per square inch or 1 atmosphere) at sea level. About what is the pressure if you are 100 meters deep in the ocean?
3. What invention gave people the ability to map the ocean floor?
4. Which parts of the ocean floor would you expect there to be the greatest amount of living organisms?
5. How much deeper did the *Trieste* submerge than *Alvin*?
6. Compare and contrast the continental shelf and the abyssal plain.
7. Why do you think mapping the seafloor is important to the Navy? Explain.

8. If the mid-ocean ridge is created where the tectonic plates separate, why is a mountain range formed there?

Vocabulary

abyssal plain
The flat bottom of the ocean floor; the deep ocean floor.

continental shelf
The shallow, gradually sloping seabed around the edge of a continent. Usually less than 200 meters in depth. The continental shelf can be thought of as the submerged edge of a continent.

continental slope
The sloped bottom of the ocean that extends from the continental shelf down to the deep ocean bottom.

mid ocean ridge
Mountain range on the ocean floor where magma upwells and new ocean floor is formed.

seamount
A mountain rising from the seafloor that does not reach above the surface of the water. Usually formed from volcanoes.

trench
Deepest areas of the ocean; found where subduction takes place.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/The_Seafloor. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 7.3: The Seafloor is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.4: Ocean Currents

Ocean water is constantly in motion (Figure 14.7). From north to south, east to west, and up and down the shore, ocean water moves all over the place. These movements can be explained as the result of many separate forces, including local conditions of wind, water, the position of the moon and Sun, the rotation of the Earth, and the position of land formations.



Figure 14.7: Ocean waves transfer energy through the water over great distances.

Lesson Objectives

- Describe how surface currents form and how they affect the world's climate.
- Describe the causes of deep currents.
- Relate upwelling areas to their impact on the food chain.

Surface Currents

Wind that blows over the ocean water creates waves. It also creates **surface currents**, which are horizontal streams of water that can flow for thousands of kilometers and can reach depths of hundreds of meters. Surface currents are an important factor in the ocean because they are a major factor in determining climate around the globe.

Causes of Surface Currents




Figure 14.14: The Coriolis Effect causes winds and currents to form circular patterns. The direction that they spin depend on the hemisphere that they are in.

Currents on the surface are determined by three major factors: the major overall global wind patterns, the rotation of the Earth, and the shape of ocean basins.

When you blow across a cup of hot chocolate, you create tiny ripples on its surface that continue to move after you've stopped blowing. The ripples in the cup are tiny waves, just like the waves that wind forms on the ocean surface. The movement of hot chocolate throughout the cup forms a stream or current, just as oceanic water moves when wind blows across it.

But what makes the wind start to blow? When sunshine heats up air, the air expands, which means the density of the air decreases and it becomes lighter. Like a balloon, the light warm air floats upward, leaving a slight vacuum below, which pulls in cooler,

denser air from the sides. The cooler air coming into the space left by the warm air is wind.

Because the Earth's equator is warmed by the most direct rays of the Sun, air at the equator is hotter than air further north or south. This hotter air rises up at the equator and as colder air moves in to take its place, winds begin to blow and push the ocean into waves and currents.

Wind is not the only factor that affects ocean currents. The 'Coriolis Effect' describes how Earth's rotation steers winds and surface currents (Figure 14.14). The Earth is a sphere that spins on its axis in a counterclockwise direction when seen from the North Pole. The further towards one of the poles you move from the equator, the shorter the distance around the Earth. This means that objects on the equator move faster than objects further from the equator. While wind or an ocean current moves, the Earth is spinning underneath it. As a result, an object moving north or south along the Earth will appear to move in a curve, instead of in a straight line. Wind or water that travels toward the poles from the equator is deflected to the east, while wind or water that travels toward the equator from the poles gets bent to the west. The Coriolis Effect bends the direction of surface currents.

The third major factor that determines the direction of surface currents is the shape of ocean basins (Figure 14.15). When a surface current collides with land, it changes the direction of the currents. Imagine pushing the water in a bathtub towards the end of the tub. When the water reaches the edge, it has to change direction.




Figure 14.15: This map shows the major surface currents at sea. Currents are created by wind, and their directions are determined by the Coriolis effect and the shape of ocean basins.

Effect on Global Climate

Surface currents play a large role in determining climate. These currents bring warm water from the equator to cooler parts of the ocean; they transfer heat energy. Let's take the Gulf Stream as an example; you can find the Gulf Stream in the North Atlantic Ocean in Figure 14.15. The Gulf Stream is an ocean current that transports warm water from the equator past the east coast of North America and across the Atlantic to Europe. The volume of water it transports is more than 25 times that of all of the rivers in the world combined, and the energy it transfers is more than 100 times the world's energy demand. It is about 160 kilometers wide and about a kilometer deep. The Gulf Stream's warm waters give Europe a much warmer climate than other places at the same latitude. If the Gulf Stream were severely disrupted, temperatures would plunge in Europe.

Deep Currents

Surface currents occur close to the surface of the ocean and mostly affect the photic zone. Deep within the ocean, equally important currents exist that are called **deep currents**. These currents are not created by wind, but instead by differences in density of masses of water. Density is the amount of mass in a given volume. For example, if you take two full one liter bottles of liquid, one might weigh more, that is it would have greater mass than the other. Because the bottles are both of equal volume, the liquid in the heavier bottle is denser. If you put the two liquids together, the one with greater density would sink and the one with lower density would rise.

Two major factors determine the density of ocean water: salinity (the amount of salt dissolved in the water) and temperature (Figure 14.16). The more salt that is dissolved in the water, the greater its density will be. Temperature also affects density: the colder the temperature, the greater the density. This is because temperature affects volume but not mass. Colder water takes up less space than warmer water (except when it freezes). So, cold water has greater density than warm water.




Figure 14.16: Thermohaline currents are created by differences in density due to temperature (thermo) and salinity (haline). The blue arrows are deep currents and the red ones are surface currents.




Figure 14.17: Surface and deep currents together form convection currents that circulate water from one place to another and back again. A water particle in the convection cycle can take 1600 years to complete the cycle.

More dense water masses will sink towards the ocean floor. Just like convection in air, when denser water sinks, its space is filled by less dense water moving in. This creates convection currents that move enormous amounts of water in the depths of the ocean. Why is the water temperature cooler in some places? Water cools as it moves from the equator to the poles via surface currents. Cooler water is more dense so it begins to sink. As a result, the surface currents and the deep currents are linked. Wind causes surface currents to transport water around the oceans, while density differences cause deep currents to return that water back around the globe (Figure 14.17).

Upwelling

As you have seen, water that has greater density usually sinks to the bottom. However, in the right conditions, this process can be reversed. Denser water from the deep ocean can come up to the surface in an upwelling (Figure 14.18). Generally, an upwelling occurs along the coast when wind blows water strongly away from the shore. As the surface water is blown away from the shore, colder water from below comes up to take its place. This is an important process in places like California, South America, South Africa, and the Arabian Sea because the nutrients brought up from the deep ocean water support the growth of plankton which, in turn, supports other members in the ecosystem. Upwelling also takes place along the equator between the North and South Equatorial Currents.




Figure 14.18: An upwelling forces denser water from below to take the place of less dense water at the surface that is pushed away by the wind.

Lesson Summary

- Ocean waves are energy traveling through the water.
- The highest portion of a wave is the crest and the lowest is the trough.
- The horizontal distance between two wave crests is the wave's length.
- Most waves in the ocean are wind generated waves.
- Ocean surface currents are produced by major overall patterns of atmospheric circulation, the Coriolis Effect and the shape of each ocean basin.
- Ocean surface circulation brings warm equatorial waters towards the poles and cooler polar water towards the equator.
- Deep ocean circulation is density driven circulation produced by differences in salinity and temperature of water masses.
- Upwelling areas are biologically important areas that form as ocean surface waters are blown away from a shore, causing cold, nutrient rich waters to rise to the surface.

Review Questions

1. What factors of wind determine the size of a wave?
2. Define the *crest* and *trough* of a wave.
3. What is the most significant cause of the surface currents in the ocean?
4. How do ocean surface currents affect climate?
5. What is the Coriolis Effect?
6. Some scientists have hypothesized that if enough ice in Greenland melts, the Gulf Stream might be shut down. Without the Gulf Stream to bring warm water northward, Europe would become much colder. Explain why melting ice in Greenland might affect the Gulf Stream.
7. What process can make denser water rise to the top?
8. Why are upwelling areas important to marine life?

Vocabulary

amplitude

The vertical height of a wave, measured from trough to crest.

Coriolis effect

The apparent deflection of a moving object like water or air caused by Earth's rotation.

crest

The highest point in a wave.

deep current

A current deep within the ocean, which moves because of density differences (caused by differences in water temperature and salinity).

rip current

A strong surface current of water that is returning to the ocean from the shore.

surface current

A horizontal movement of ocean water, caused by surface winds.

trough

The lowest point in a wave.

tsunami

A seismic sea wave generated by vertical movement of the ocean floor underwater earthquake, underwater volcanic eruption or landslide or meteorite impact.

upwelling

Cold, nutrient-rich water that rises from oceanic depths usually near the continents, when wind blows the overlying surface away or along the equator.

wave

A change in the shape of water caused by energy moving through the water.

wavelength

The horizontal distance between two troughs, or two crests in a wave.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Ocean_Movements. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 7.4: Ocean Currents is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.


7.5: Oceans and Coastal Environments

Significance of the Oceans

The oceans make up 70 percent of the planet and contains 97 percent of all the water on Earth. The vast amount of water stores the majority of the planet's moisture, terrestrial energy, and heat from the Sun. This energy is transferred between the equator and the two poles by larger surface currents by winds and deep ocean currents driven by differences in ocean density. It also provides the moisture and energy for storm systems and ultimately global climates.


Phytoplankton, microscopic plants and animals in the oceans provide the foundation of the global food web of species. The earth's oceans are so vital for life that over 40 percent of the world's population live 7.4 billion people near coastal areas.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94



MODERATE CLIMATES

As terrestrial creatures, humans think of the importance of the planet's land surfaces, yet Earth is a planet consisting of 70 percent water. From space, the dominance of water is obvious because most of it is stored in Earth's oceans.Earth would not be the same planet without its oceans. The oceans, along with the atmosphere, keep Earth's surface temperatures fairly constant worldwide. While some places on Earth get as cold as -7 degrees Celsius and others as hot as 55 degrees Celsius, the range is only 125 degrees Celsius. On Mercury temperatures go from -180 degrees Celsius to 430 degrees Celsius, a range of 610 degrees Celsius.The oceans, along with the atmosphere, distribute heat around the planet. The oceans absorb heat near the equator and then transport that solar energy to polar regions. The oceans also moderate climate within a region. At the same latitude, the temperature range is smaller along coastal areas compared to areas farther inland. Along coastal areas, summer temperatures are not as hot, and winter temperatures are not as cold, because water takes a long time to heat up or cool down.



BIOLOGICALLY RICH


The oceans are an essential part of Earth's water cycle. Since they cover so much of the planet, most evaporation comes from the ocean and most precipitation falls on the oceans.The oceans are also home to an enormous amount of life. That is, they have

tremendous biodiversity. Tiny ocean plants create the base of a food web that supports all sorts of life forms. Marine life makes up the majority of all biomass on Earth. (**Biomass** is the total mass of living organisms in a given area.) These organisms supply us with food and even the oxygen created by marine plants.



CONTINENTAL MARGIN

Recall from the chapter on Plate Tectonics that the ocean floor is not flat: mid-ocean ridges, deep sea trenches, and other features all rise sharply above or plunge deeply below the abyssal plains. In fact, Earth's tallest mountain is Mauna Kea volcano, which rises 10,203 m (33,476 ft.)meters) from the Pacific Ocean floor to become one of the volcanic mountains of Hawaii. The deepest canyon is also on the ocean floor, the Challenger Deep in the Marianas Trench, 10,916 m (35,814 ft). The mapping of the ocean floor and coastal margins is called **bathymetry**.The continental margin is the transition from the land to the deep sea or, geologically speaking, from continental crust to oceanic crust. More than one-quarter of the ocean basin is continental margin.




Average composition of the ocean.

COMPOSITION OF OCEAN WATER

Water is a polar molecule so it can dissolve many substances such as salts, sugars, acids, bases, and organic molecules. Where does the salt in seawater come from? As water moves through rock and soil on land it picks up ions. This is the flip side of weathering. Salts comprise about 3.5% of the mass of ocean water, but the salt content or **salinity** is different in different locations.In places like estuaries, seawater mixes with fresh water, causing salinity to be much lower than average. Where there is lots of evaporation but little circulation of water, salinity can be much higher. The Dead Sea has 30% salinity—nearly nine times the average salinity of ocean water. It is called the Dead Sea because nearly nothing can survive within it because of its salinity. Earthquide has aninteractive ocean maps can show salinity, temperature, nutrients, and other characteristics.Differences in water density are responsible for deep ocean currents. With so many dissolved substances mixed in seawater, what is the density (mass per volume) of seawater relative to fresh water? Water density increases as: salinity increases; temperature decreases; pressure increases

Layers of the Ocean



In 1960, two men in a specially designed submarine called the **Trieste**descended into a submarine trench called the Challenger Deep (10,910 meters). The average depth of the ocean is 3,790 m, a lot more shallow than the deep trenches but still an incredible depth for sea creatures to live in. There are three major factors that make the deep ocean hard to inhabit: the absence of light, low temperature, and extremely high pressure.

VERTICAL DIVISIONS

To better understand regions of the ocean, scientists define the **water column** by depth. They divide the entire ocean into two zones vertically, based on light level. Large lakes are divided into similar regions. Sunlight only penetrates the sea surface to a depth of about 200 m, creating the **photic zone** (consisting of the Sunlight Zone and Twilight Zone). Organisms that photosynthesize depend on sunlight for food and so are restricted to the photic zone. Since tiny photosynthetic organisms, known as **phytoplankton**, supply nearly all of the energy and nutrients to the rest of the marine food web, most other marine organisms live in or at least visit the photic zone. In the **aphotic zone** (consisting of the Midnight Zone and the Abyss) there is not enough light for photosynthesis. The aphotic zone makes up the majority of the ocean, but has a relatively small amount of its life, both in diversity of type and in numbers.

HORIZONTAL DIVISIONS

The seabed is also divided into the zones described above, but the ocean itself is also divided horizontally by distance from the shore. Nearest to the shore lies the **intertidal** (littoral) **zone**, the region between the high and low tidal marks. This hallmark of the intertidal is change: water is in constant motions in waves, tides, and currents. The land is sometimes under water and sometimes is exposed. The **neritic zone** is from low tide mark and slopes gradually downward to the edge of the seaward side of the continental shelf. Some sunlight penetrates to the seabed here. The oceanic zone is the entire rest of the ocean from the bottom edge of the neritic zone, where sunlight does not reach the bottom. The sea bed and water column are subdivided further, as seen in the figure above.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94

TSUNAMIS

Tsunami are described in the Earthquakes chapter as damaging waves that result from the sharp jolt to the water from an undersea earthquake. Landslides, meteorite impacts, or any other jolt to ocean water may form a tsunami. Tsunami can travel at speeds of

800 kilometers per hour (500 miles per hour). Tsunami have small wave heights and long wavelengths so they are usually unnoticed at sea. As the wave rides up the continental shelf the wave height increases.

The wave speed of a tsunami is also slowed by friction with the shallower ocean floor, which causes the wavelength to decrease, creating a much taller wave. Many people caught in a tsunami have no warning of its approach. Since the wavelength is long, a long time can pass between crests or troughs onshore. In 1755 in Lisbon, an offshore earthquake caused a great deal of damage on land. People rushed to the open space of the shore and discovered that the water was flowing seaward fast. The trough of the tsunami wave reached shore first. People who went out onto the open beach were drowned when the crest of the wave reached shore.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94

Tidal Waves



Tides are the daily rise and fall of sea level at any given place. The pull of the Moon's gravity on Earth is the primarily cause of tides and the pull of the Sun's gravity on Earth is the secondary cause. The Moon has a greater effect because, although it is much smaller than the Sun, it is much closer. The Moon's pull is about twice that of the Sun's.

DAILY TIDE PATTERNS

To understand the tides it is easiest to start with the effect of the Moon on Earth. As the Moon revolves around our planet, its gravity pulls Earth toward it. The lithosphere is unable to move much but the water is pulled by the gravity and a bulge is created. This bulge is the high tide beneath the Moon. The Moon's gravity then pulls the Earth toward it, leaving the water on the opposite side of the planet behind. This creates a second high tide bulge on the opposite side of Earth from the Moon. These two water bulges on opposite sides of the Earth aligned with the Moon are the **high tides**. Click here to learn more.

Since so much water is pulled into the two high tides, **low tides** form between the two high tides. As the Earth rotates beneath the Moon, a single spot will experience two high tides and two low tides every day.

The **tidal range** is the difference between the ocean level at high tide and the ocean at low tide. The tidal range in a location depends on a number of factors, including the slope of the seafloor. Water appears to move a greater distance on a gentle slope than on a steep slope.

MONTHLY TIDAL PATTERNS

If you look at the diagram of high and low tides on a circular Earth above, you'll see that tides are waves. So when the Sun and Moon are aligned, what do you expect the tides to look like?


Waves are additive so when the gravitational pull of both bodies is in the same direction the high tides add and the low tides add. Highs are higher and lows are lower than at other times through the month. These more extreme tides, with a greater tidal range, are called **spring tides**. Spring tides don't just occur in the spring; they occur whenever the Moon is in a new-moon or full-moon phase, about every 14 days. The National Oceanic and Atmospheric Administration (NOAA) has a simple diagram of this process.

Neap tides are tides that have the smallest tidal range, and they occur when the Earth, the Moon, and the Sun form a 90o angle. They occur exactly halfway between the spring tides, when the Moon is at first or last quarter. How do the tides add up to create neap tides? The Moon's high tide occurs in the same place as the Sun's low tide and the Moon's low tide in the same place as the Sun's high tide. At neap tides, the tidal range relatively small. Here is another simple animation of spring and neap tides from NOAA.

High tides occur about twice a day, about every 12 hours and 25 minutes. The reason is that the Moon takes 24 hours and 50 minutes to rotate once around the Earth so the Moon is over the same location 24 hours and 50 minutes later. Since high tides occur twice a day, one arrives each 12 hours and 25 minutes. What is the time between a high tide and the next low tide? This animation shows the effect of the Moon and Sun on the tides.

Some coastal areas do not follow this pattern at all. These coastal areas may have one high and one low tide per day or a different amount of time between two high tides. These differences are often because of local conditions, such as the shape of the coastline that the tide is entering.

Surface Currents



Ocean water moves in predictable ways along the ocean surface.**Surface currents** can flow for thousands of kilometers and can reach depths of hundreds of meters. These surface currents do not depend on weather; they remain unchanged even in large storms because they depend on factors that do not change.

Surface currents are created by three things: *global wind patterns*, the *rotation of the Earth*, and the *shape of the ocean basins*.

Surface currents are extremely important because they distribute heat around the planet and are a major factor influencing climate around the globe.

GLOBAL WIND CURRENTS

Winds on Earth are either global or local. Global winds blow in the same directions all the time and are related to the unequal heating of Earth by the Sun — that is, more solar radiation strikes the equator than the polar regions –- and the rotation of the Earth — that is, the **Coriolis effect**. The causes of the global wind patterns will be described in detail later when we look at the atmosphere. Water in the surface currents is pushed in the direction of the major wind belts:

- **trade winds**: east to west between the equator and 30 degrees North and 30 degrees South
- **westerlies**: west to east in the middle latitudes
- **polar easterlies**: east to west between 50 degrees and 60 degrees north and south of the equator and the north and south pole

ROTATION OF THE EARTH

Wind is not the only factor that affects ocean currents. The Coriolis effect describes how Earth's rotation steers winds and surface ocean currents. Coriolis causes freely moving objects to appear to move to the right in the Northern Hemisphere and to the left in the Southern Hemisphere. The objects themselves are actually moving straight, but the Earth is rotating beneath them, so they seem to bend or curve.

An example might make the Coriolis effect easier to visualize. If an airplane flies 500 miles due north, it will not arrive at the city that was due north of it when it began its journey. Over the time it takes for the airplane to fly 500 miles, that city moved, along with the Earth it sits on. The airplane will therefore arrive at a city to the west of the original city (in the Northern Hemisphere),


unless the pilot has compensated for the change. So to reach his intended destination, the pilot must also veer right while flying north.

As wind or an ocean current moves, the Earth spins underneath it. As a result, an object moving north or south along the Earth will appear to move in a curve, instead of in a straight line. Wind or water that travels toward the poles from the equator is deflected to the east, while wind or water that travels toward the equator from the poles gets bent to the west. The Coriolis effect bends the direction of surface currents to the right in the Northern Hemisphere and left in the Southern Hemisphere. Coriolis effect is demonstrated using a metal ball and a rotating plate in this video. The ball moves in a circular path just like a freely moving particle of gas or liquid moves on the rotating Earth.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94


Shape of the Ocean Basin

When a surface current collides with land, the current must change direction. In the figure below, the Atlantic South Equatorial Current travels westward along the equator until it reaches South America. At Brazil, some of it goes north and some goes south. Because of Coriolis effect, the water goes right in the Northern Hemisphere and left in the Southern Hemisphere.

You can see on the map below the major surface ocean currents that the surface ocean currents create loops called **gyres**. The Antarctic Circumpolar Current is unique because it travels uninhibited around the globe. Why is it the only current to go all the way around?

	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94

Local and Surface Currents




The surface currents described above are all large and unchanging. Local surface currents are also found along shorelines. Two are *long shore currents*and *rip currents*. **Rip currents** are potentially dangerous currents that carry large amounts of water offshore quickly. Look at the rip-current animation to determine what to do if you are caught in a rip current. Each summer in the United States at least a few people die when they are caught in rip currents.

Surface currents play an enormous role in Earth's climate. Even though the equator and poles have very different climates, these regions would have more extremely different climates if ocean currents did not transfer heat from the equatorial regions to the higher latitudes.

The Gulf Stream is a river of warm water in the Atlantic Ocean, about 160 kilometers wide and about a kilometer deep. Water that enters the Gulf Stream is heated as it travels along the equator. The warm water then flows up the east coast of North America and across the Atlantic Ocean to Europe. The energy the Gulf Stream transfers is enormous: more than 100 times the world's energy demand.

The Gulf Stream's warm waters raise temperatures in the North Sea, which raises the air temperatures over land between 3 to 6 degrees C (5 to 11 degrees F). London, U.K., for example, is at the same latitude as Quebec, Canada. However, London's average January temperature is 3.8 degrees C (38 degrees F), while Quebec's is only -12 degrees C (10 degrees F). Because air traveling over the warm water in the Gulf Stream picks up a lot of water, London gets a lot of rain. In contrast, Quebec is much drier and receives its precipitation as snow.

Deep Currents



Red arrows indicate surface water currents and blue arrows indicate deep water currents.

Thermohaline circulation drives deep ocean circulation. Thermo means heat and haline refers to salinity. Differences in temperature and in salinity change the density of seawater. So thermohaline circulation is the result of density differences in water masses because of their different temperature and salinity.

What is the temperature and salinity of very dense water? Lower temperature and higher salinity yield the densest water. When a volume of water is cooled, the molecules move less vigorously so same number of molecules takes up less space and the water is denser. If salt is added to a volume of water, there are more molecules in the same volume so the water is denser.


Changes in temperature and salinity of seawater take place at the surface. Water becomes dense near the poles. Cold polar air cools the water and lowers its temperature, increasing its salinity. Fresh water freezes out of seawater to become sea ice, which also increases the salinity of the remaining water. This very cold, very saline water is very dense and sinks. This sinking is called **downwelling**.

Two things then happen. The dense water pushes deeper water out of its way and that water moves along the bottom of the ocean. This deep water mixes with less dense water as it flows. Surface currents move water into the space vacated at the surface where the dense water sank. Water also sinks into the deep ocean off of Antarctica. Since unlimited amounts of water cannot sink to the bottom of the ocean, water must rise from the deep ocean to the surface somewhere. This process is called **upwelling**.

Generally, upwelling occurs along the coast when wind blows water strongly away from the shore. This leaves a void that is filled by deep water that rises to the surface. Upwelling is extremely important where it occurs. During its time on the bottom, the cold deep water has collected nutrients that have fallen down through the water column. Upwelling brings those nutrients to the surface. Those nutrient support the growth of plankton and form the base of a rich ecosystem. California, South America, South Africa, and the Arabian Sea all benefit from offshore upwelling.

Upwelling also takes place along the equator between the North and South Equatorial Currents. Winds blow the surface water north and south of the equator so deep water undergoes upwelling. The nutrients rise to the surface and support a great deal of life in the equatorial oceans.

Studying the Seafloor



Oceanographers like to say that we know more about the dark side of the Moon than we do about the oceans. That statement is doubly true of the seafloor. Although modern technology has allowed us to learn more about the seafloor, vast regions remain unexplored. Scuba divers can only dive to about 40 meters and they cannot stay down there for very long. Although this is good for researching the organisms and ecosystems very near a coast, most oceanic research requires accessing greater depths.

MAPPING

In the Plate Tectonics chapter you learned that echo sounders designed to locate enemy submarines allowed scientists to create bathymetric maps of the seafloor. Prior to this advance, explorers mapped a small amount of the seafloor by painstakingly dropping a line over the side of a ship to measure the depth at one tiny spot at a time.

SAMPLING REMOTELY

Samples of seawater from different depths in the water column are needed to understand ocean chemistry. To do this bottles are placed along a cable at regular depths and closed as a weight is dropped down the cable. The water trapped in the bottle can be analyzed later in a laboratory.

Scientists are also interested in collecting rock and sediment samples from the seafloor. A dredge is a giant rectangular bucket that is dragged along behind a ship to collect loose rocks. Gravity corers are metal tubes that fall to the seafloor and slice into the sediments to collect a sample. The research vessel, the Joides Resolution, drills deep into the seafloor to collect samples of the sediment and ocean crust. Scientists analyze the samples for chemistry and paleomagnetism.



SUBMERSIBLES

Samples of seawater and rocks can be collected directly by scientists in a **submersible**. These subs can take scientists down to make observations and the subs have arms for collecting samples. The submersible Alvin is an HOV, a human operated vehicle. Alvin can dive up to 4,500 m beneath the ocean surface and has made more than 4000 dives since 1964. Other submersibles can dive deeper. View a slide show of DSV Alvin and its history from the Woods Hole Oceanographic Institution

REMOTELY OPERATED VEHICLES

To avoid the expense, dangers, and limitations of human missions under the sea, remotely operated vehicles or ROV's, allow scientists to study the ocean's depths by using small vehicles carrying cameras and scientific instruments. ROVs were used to study the Titanic, which would have been far too dangerous for a manned sub to enter. Scientists control ROVs electronically with sophisticated operating systems.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=94

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-9-oceans-and-coastal-environments.html. **License**: *CC BY: Attribution*

This page titled 7.5: Oceans and Coastal Environments is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.6: Tides

Tides

Wind is the primary force that causes ocean surface waves, but it does not cause the tides. Tides are the daily changes in the level of the ocean water at any given place. The main factors that causes tides are the gravitational pull of the Moon and the Sun (Figure 14.10).




Figure 14.10: High tide (left) and low tide (right) at Bay of Fundy on the Gulf of Maine in North America. The Bay of Fundy has one of the greatest tidal ranges on Earth.




Figure 14.11: High tide is created by the gravitational pull of the moon which pulls water toward it. Water on the opposite side of the Earth is pulled least by the moon so the water bulges away from the moon. High tide occurs where the water is bulging. Low tide occurs where it is not.

How does the Moon affect the oceans? Since the Moon is a relatively large object in space that is very close to the Earth, its gravity actually pulls Earth's water towards it. Wherever the moon is, as it orbits the Earth, there is a high tide 'bulge' that stays lined up with the Moon. The side of the Earth that is furthest from the Moon also has a high tide 'bulge'. This is because the Earth is closer to the moon than the water on its far side. The Moon's gravity pulls more on the planet than the water on the opposite side. These two water bulges on opposite sides of the Earth aligned with the Moon are the **high tides**. Since ocean water is pulled higher in the areas of the two high tides, there is less water in between the two high tides. These areas are the **low tides** (Figure 14.11).

The **tidal range** is the difference between the ocean level at high tide and the ocean at low tide (Figure 14.12). Some places have a greater tidal range than others. High tides occur about twice a day, about every 12 hours and 24 minutes.




Figure 14.12: The tidal range is the difference between the ocean level at high tide and low tide.




Figure 14.13: Spring tides occur when the Earth, the Sun, and the Moon are aligned, increasing the gravitational pull on the oceans. Sometimes, the Sun and Moon are on opposite sides of the Earth while at other times, they are on the same side. Neap tides occur when the Earth, the Sun, and the Moon form a right angle, at first and last quarters for the Moon.

The Moon's gravity is mostly responsible for our tides, but the Sun also plays a role (Figure 14.13). The Sun is much larger than our Moon. It has a mass about 27,500,000 times greater than the Moon. A very large object like the Sun would produce tremendous tides if it were as near to Earth as the Moon. However it is so far from the Earth that its effect on the tides is only about half as strong as the Moon's. When both the Sun and Moon are aligned, the effect of each is added together, producing higher than normal tides called **spring tides**.

Spring tides are tides with the greatest tidal range. Despite their name, spring tides don't just occur in the spring; they occur throughout the year whenever the Moon is in a new-moon or full-moon phase, or about every 14 days.

Neap tides are tides that have the smallest tidal range, and occur when the Earth, the Moon, and the Sun form a 90° angle. They occur exactly halfway between the spring tides, when the Moon is at first or last quarter. This happens because the Moon's high tide occurs in the same place as the Sun's low tide and the Moon's low tide is added to by the Sun's high tide.

The material that follows on tides was cut from Ocean Movements section under Coastal Processes

Tides are the daily rise and fall of sea level at any given place. The pull of the Moon's gravity on Earth is the primarily cause of tides and the pull of the Sun's gravity on Earth is the secondary cause (**Figure** below). The Moon has a greater effect because, although it is much smaller than the Sun, it is much closer. The Moon's pull is about twice that of the Sun's.




High tide (left) and low tide (right) at Bay of Fundy on the Gulf of Maine. The Bay of Fundy has the greatest tidal ranges on Earth at 38.4 feet.

Daily Tide Patterns

To understand the tides it is easiest to start with the effect of the Moon on Earth. As the Moon revolves around our planet, its gravity pulls Earth toward it. The lithosphere is unable to move much but the water is pulled by the gravity and a bulge is created. This bulge is the high tide beneath the Moon. The Moon's gravity then pulls the Earth toward it, leaving the water on the opposite side of the planet behind. This creates a second high tide bulge on the opposite side of Earth from the Moon. These two water bulges on opposite sides of the Earth aligned with the Moon are the**high tides**.


Since so much water is pulled into the two high tides, **low tides** form between the two high tides (**Figure** below). As the Earth rotates beneath the Moon, a single spot will experience two high tides and two low tides every day.



The gravitational attraction of the Moon to ocean water creates the high and low tides.

A detailed animation of lunar tides is shown here: http://www.pbs.org/wgbh/nova/venice/tides.html.

The **tidal range** is the difference between the ocean level at high tide and the ocean at low tide (**Figure** below). The tidal range in a location depends on a number of factors, including the slope of the seafloor. Water appears to move a greater distance on a gentle slope than on a steep slope.




The tidal range is the difference between the ocean level at high tide and low tide.

Monthly Tide Patterns

If you look at the diagram of high and low tides in the **Figure** above, you'll see that tides are waves. So when the Sun and Moon are aligned, what do you expect the tides to look like?


Waves are additive so when the gravitational pull of both bodies is in the same direction the high tides add and the low tides add (**Figure** below). Highs are higher and lows are lower than at other times through the month. These more extreme tides, with a greater tidal range, are called **spring tides**. Spring tides don't just occur in the spring; they occur whenever the Moon is in a new-moon or full-moon phase, about every 14 days.



Spring tides occur when the tidal bulges from the Moon and Sun are aligned. The Moon is full in this image; in the bottom image the Moon would appear as a new Moon.

Here is a link to see these tides in motion: http://oceanservice.noaa.gov/education/kits/tides/media/tide06a_450.gif.

Neap tides are tides that have the smallest tidal range, and they occur when the Earth, the Moon, and the Sun form a 90° angle (**Figure** below). They occur exactly halfway between the spring tides, when the Moon is at first or last quarter. How do the tides add up to create neap tides? The Moon's high tide occurs in the same place as the Sun's low tide and the Moon's low tide in the same place as the Sun's high tide. At neap tides, the tidal range relatively small.



Neap tides occur when the Earth, the Sun, and the Moon form a right angle; the Moon is in its first or third quarter.

A simple animation of spring and neap tides is found here: http://oceanservice.noaa.gov/education/kits/tides/media/supp_tide06a.html.

High tides occur about twice a day, about every 12 hours and 25 minutes. The reason is that the Moon takes 24 hours and 50 minutes to rotate once around the Earth so the Moon is over the same location 24 hours and 50 minutes later. Since high tides occur twice a day, one arrives each 12 hours and 25 minutes. What is the time between a high tide and the next low tide?

This animation shows the effect of the Moon and Sun on the tides: http://www.onr.navy.mil/focus/ocean/motion/tides1.htm.

Some coastal areas do not follow this pattern at all. These coastal areas may have one high and one low tide per day or a different amount of time between two high tides. These differences are often because of local conditions, such as the shape of the coastline that the tide is entering.

Lumen Learning authored content

- **License**: *Public Domain: No Known Copyright*

This page titled 7.6: Tides is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

7.7: Currents

Surface Currents

Ocean water moves in predictable ways along the ocean surface. **Surface currents** can flow for thousands of kilometers and can reach depths of hundreds of meters. These surface currents do not depend on weather; they remain unchanged even in large storms because they depend on factors that do not change.

Surface currents are created by three things:

- global wind patterns
- the rotation of the Earth
- the shape of the ocean basins

Surface currents are extremely important because they distribute heat around the planet and are a major factor influencing climate around the globe.

Global Wind Patterns

Winds on Earth are either global or local. Global winds blow in the same directions all the time and are related to the unequal heating of Earth by the Sun — that is, more solar radiation strikes the equator than the polar regions –- and the rotation of the Earth — that is, the **Coriolis effect**. The causes of the global wind patterns will be described in detail in the Earth's Atmosphere chapter.

Water in the surface currents is pushed in the direction of the major wind belts:


- trade winds: east to west between the equator and 30°N and 30°S
- westerlies: west to east in the middle latitudes
- polar easterlies: east to west between 50° and 60° north and south of the equator and the north and south pole

Earth's Rotation

Wind is not the only factor that affects ocean currents. The Coriolis effect describes how Earth's rotation steers winds and surface ocean currents (**Figure** below). Coriolis causes freely moving objects to appear to move to the right in the Northern Hemisphere and to the left in the Southern Hemisphere. The objects themselves are actually moving straight, but the Earth is rotating beneath them, so they seem to bend or curve.

An example might make the Coriolis effect easier to visualize. If an airplane flies 500 miles due north, it will not arrive at the city that was due north of it when it began its journey. Over the time it takes for the airplane to fly 500 miles, that city moved, along with the Earth it sits on. The airplane will therefore arrive at a city to the west of the original city (in the Northern Hemisphere), unless the pilot has compensated for the change. So to reach his intended destination, the pilot must also veer right while flying north.

As wind or an ocean current moves, the Earth spins underneath it. As a result, an object moving north or south along the Earth will appear to move in a curve, instead of in a straight line. Wind or water that travels toward the poles from the equator is deflected to the east, while wind or water that travels toward the equator from the poles gets bent to the west. The Coriolis effect bends the direction of surface currents to the right in the Northern Hemisphere and left in the Southern Hemisphere.




The Coriolis effect causes winds and currents to form circular patterns. The direction that they spin depends on the hemisphere that they are in.

Coriolis effect is demonstrated using a metal ball and a rotating plate in this video. The ball moves in a circular path just like a freely moving particle of gas or liquid moves on the rotating Earth **(5b)**: http://www.youtube.com/watch?v=Wda7azMvabE (2:04).


Shape of the Ocean Basins

When a surface current collides with land, the current must change direction. In the **Figure** below, the Atlantic South Equatorial Current travels westward along the equator until it reaches South America. At Brazil, some of it goes north and some goes south. Because of Coriolis effect, the water goes right in the Northern Hemisphere and left in the Southern Hemisphere.



The major surface ocean currents.

You can see on the map of the major surface ocean currents that the surface ocean currents create loops called **gyres** (**Figure** below). The Antarctic Circumpolar Current is unique because it travels uninhibited around the globe. Why is it the only current to go all the way around?



The ocean gyres. Why do the Northern Hemisphere gyres rotate clockwise and the Southern Hemisphere gyres rotate counterclockwise?

This video shows the surface ocean currents set by global wind belts **(5a)**: http://www.youtube.com/watch?v=Hu_Ga0JYFNg (1:20).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=98

Effect on Global Climate

Surface currents play an enormous role in Earth's climate. Even though the equator and poles have very different climates, these regions would have more extremely different climates if ocean currents did not transfer heat from the equatorial regions to the higher latitudes.

The Gulf Stream is a river of warm water in the Atlantic Ocean, about 160 kilometers wide and about a kilometer deep. Water that enters the Gulf Stream is heated as it travels along the equator. The warm water then flows up the east coast of North America and across the Atlantic Ocean to Europe (**Figure** below). The energy the Gulf Stream transfers is enormous: more than 100 times the world's energy demand.

The Gulf Stream's warm waters raise temperatures in the North Sea, which raises the air temperatures over land between 3 to 6oC (5 to 11oF). London, U.K., for example, is at the same latitude as Quebec, Canada. However, London's average January temperature is 3.8oC (38oF), while Quebec's is only -12oC (10oF). Because air traveling over the warm water in the Gulf Stream picks up a lot of water, London gets a lot of rain. In contrast, Quebec is much drier and receives its precipitation as snow.

In a satellite image of water temperature in the western Atlantic it is easy to pick out the Gulf Stream, which brings warmer waters from the equator up eastern North America.

Deep Currents**Thermohaline circulation** drives deep ocean circulation. Thermo means heat and haline refers to salinity. Differences in temperature and in salinity change the density of seawater. So thermohaline circulation is the result of density differences in water masses because of their different temperature and salinity.

What is the temperature and salinity of very dense water? Lower temperature and higher salinity yield the densest water. When a volume of water is cooled, the molecules move less vigorously so same number of molecules takes up less space and the water is denser. If salt is added to a volume of water, there are more molecules in the same volume so the water is denser.

Changes in temperature and salinity of seawater take place at the surface. Water becomes dense near the poles. Cold polar air cools the water and lowers its temperature, increasing its salinity. Fresh water freezes out of seawater to become sea ice, which also increases the salinity of the remaining water. This very cold, very saline water is very dense and sinks. This sinking is called **downwelling**.

This video lecture discusses the vertical distribution of life in the oceans. Seawater density creates currents, which provide different habitats for different creatures **(5d)**: http://www.youtube.com/watch?v=LA1jxeXDsdA (6:12).

Two things then happen. The dense water pushes deeper water out of its way and that water moves along the bottom of the ocean. This deep water mixes with less dense water as it flows. Surface currents move water into the space vacated at the surface where the dense water sank (**Figure** below). Water also sinks into the deep ocean off of Antarctica.

Cold water (blue lines) sinks in the North Atlantic, flows along the bottom of the ocean and upwells in the Pacific or Indian. The water then travels in surface currents (red lines) back to the North Atlantic. Deep water also forms off of Antarctica.

Since unlimited amounts of water cannot sink to the bottom of the ocean, water must rise from the deep ocean to the surface somewhere. This process is called **upwelling** (**Figure** below).

Upwelling forces denser water from below to take the place of less dense water at the surface that is pushed away by the wind.

Generally, upwelling occurs along the coast when wind blows water strongly away from the shore. This leaves a void that is filled by deep water that rises to the surface.

Upwelling is extremely important where it occurs. During its time on the bottom, the cold deep water has collected nutrients that have fallen down through the water column. Upwelling brings those nutrients to the surface. Those nutrient support the growth of plankton and form the base of a rich ecosystem. California, South America, South Africa, and the Arabian Sea all benefit from offshore upwelling.

An animation of upwelling is seen here: http://oceanservice.noaa.gov/education/kits/currents/03coastal4.html.

Upwelling also takes place along the equator between the North and South Equatorial Currents. Winds blow the surface water north and south of the equator so deep water undergoes upwelling. The nutrients rise to the surface and support a great deal of life in the equatorial oceans.

Lesson Summary

- Ocean surface currents are produced by global winds, the Coriolis effect and the shape of each ocean basin.
- The Pacific and Atlantic Oceans have a circular pattern of surface currents called gyres that circle clockwise in the Northern Hemisphere and counterclockwise in the Southern. The Indian Ocean only has a counterclockwise gyre.
- Surface ocean circulation brings warm equatorial waters towards the poles and cooler polar water towards the equator.
- Thermohaline circulation drives deep ocean currents.
- Upwelling of cold, nutrient-rich waters creates biologically rich areas where surface waters are blown away from a shore, or where equatorial waters are blow outward.

Review Questions

6. What causes the patterns of surface currents in the ocean?
7. How do ocean surface currents affect climate?
8. What is the Coriolis effect?
9. What process can make deep, dense water rise to the surface?
10. Why are upwelling areas important to marine life?

Further Reading / Supplemental Links

- Learn About Ocean Currents, 5 min. Life Videopedia http://www.5min.com/Video/Learn-about-Ocean-Currents-117529352
- NOAA's Ocean Explorer program http://oceanexplorer.noaa.gov/edu/welcome.html
- A tutorial for grades 6 to 12 on currents from NOAA: http://oceanservice.noaa.gov/education/tutorial_currents/welcome.html
- Cool jobs: Oceanographer http://news.discovery.com/videos/cool-jobs-cool-jobs-oceanographer.html

Points to Consider

- Some scientists have hypothesized that if enough ice in Greenland melts, the Gulf Stream might be shut down. Why might this happen?
- If the Gulf Stream shuts down, what would be the result on climate in Europe?
- How do the movements of ocean water contribute to the ocean's life?

CC licensed content, Shared previously

- **Provided by**: CK12. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/14.2/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 7.7: Currents is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

8: Coastal Processes

8.1: Waves and Coastal Features

This page titled 8: Coastal Processes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

8.1: Waves and Coastal Features

Introduction

Ocean water is constantly in motion: north-south, east-west, alongshore, and vertically. Seawater motions are the result of waves, tides, and currents (**Figure** below). Ocean movements are the consequence of many separate factors: wind, tides, Coriolis effect, water density differences, and the shape of the ocean basins. Water movements and their causes will be discussed in this lesson.



Ocean waves transfer energy through the water over great distances.

Waves

Waves have been discussed in previous chapters in several contexts: seismic waves traveling through the planet, sound waves traveling through seawater, and ocean waves eroding beaches. Waves transfer energy and the size of a wave and the distance it travels depends on the amount of energy that it carries.

Wind Waves

This lesson studies the most familiar waves, those on the ocean's surface. Ocean waves originate from wind blowing – steady winds or high storm winds – over the water. Sometimes these winds are far from where the ocean waves are seen. What factors create the largest ocean waves?

The largest wind waves form when the wind

- is very strong
- blows steadily for a long time
- blows over a long distance

The wind could be strong, but if it gusts for just a short time, large waves won't form.


Wind blowing across the water transfers energy to that water. The energy first creates tiny ripples that create an uneven surface for the wind to catch so that it may create larger waves. These waves travel across the ocean out of the area where the wind is blowing.

Remember that a wave is a transfer of energy. Do you think the same molecules of water that starts out in a wave in the middle of the ocean later arrive at the shore?

Water molecules in waves make circles or ellipses (**Figure** below). Energy transfers between molecules but the molecules themselves mostly bob up and down in place.

In this animation, a water bottle bobs in place like a water molecule: http://www.onr.navy.mil/focus/ocean/motion/waves1.htm

An animation of motion in wind waves from the Scripps Institution of Oceanography:http://earthguide.ucsd.edu/earthguide/diagrams/waves/swf/wave_wind.html



The circles show the motion of a water molecule in a wind wave. Wave energy is greatest at the surface and decreases with depth. A shows that a water molecule travels in a circular motion in deep water. B shows that molecules in shallow water travel in an elliptical path because of the ocean bottom.


An animation of a deep water wave is seen here: http://en.Wikipedia.org/wiki/File:Deep_water_wave.gif.

An animation of a shallow water wave is seen here: http://commons.wikimedia.org/wiki/File:Shallow_water_wave.gif.

When does a wave break? Do waves only break when they reach shore? Waves break when they become too tall to be supported by their base. This can happen at sea but happens predictably as a wave moves up a shore. The energy at the bottom of the wave is lost by friction with the ground so that the bottom of the wave slows down but the top of the wave continues at the same speed. The crest falls over and crashes down.

Local Surface Currents

The surface currents described above are all large and unchanging. Local surface currents are also found along shorelines (**Figure** below). Two are **longshore currents** and **rip currents**.



Longshore currents move water and sediment parallel to the shore in the direction of the prevailing local winds.

Rip currents are potentially dangerous currents that carry large amounts of water offshore quickly. Look at the rip-current animation to determine what to do if you are caught in a rip current: http://www.onr.navy.mil/focus/ocean/motion/currents2.htm. Each summer in the United States at least a few people die when they are caught in rip currents.

This animation shows the surface currents in the Caribbean, the Gulf of Mexico, and the Atlantic Ocean off of the southeastern United States:http://polar.ncep.noaa.gov/ofs/viewer.shtml?-gulfmex-cur-0-large-rundate=latest.

Wave Action and Erosion

Have you ever been to visit a beach? Some beaches have large, strong rolling waves that rise up and collapse as they crash into the shore. All waves are *energy* traveling through some type of material (Figure 10.13). The waves that we are most familiar with travel

through water. Most of these waves form from wind blowing over the water; sometimes steady winds that blow and sometimes from a storm that forms over the water. The energy of waves does the work of erosion when a wave reaches the shore. When you find a piece of frosted glass along a beach, you have found some evidence of the work of waves. What other evidence might you find?



Figure 10.13: Ocean waves are energy traveling through water.

As wind blows over the surface of the water, it disturbs the water, producing the familiar shape of a wave. You can see this shape in Figure 10.14. The highest part of a wave is called the **wave crest**. The lowest part is called the **wave trough**. The vertical distance from the highest part of a wave to the lowest is called the **wave height**. The horizontal distance between one wave crest and the next crest, is called the **wavelength**. Three things influence how big a wave might get. If the wind is very strong, and it blows steadily for a long time over a long distance, the very largest waves will form. The wind could be strong, but if it gusts for just a short time, large waves won't form. Bigger waves do more work of erosion which changes our shorelines. Each day that waves break along the shore, they steadily erode away a minute bit of the shoreline. When one day, a really big storm like a hurricane arrives, it will do a lot of damage in just a very short time.




Figure 10.14: Each wave form has a wave crest, trough, height, and wavelength.

As waves come into shore, they usually reach the shore at some angle. This means one part of the wave reaches shallow water sooner than the parts of the wave that are further out. As a wave comes into shore, the water 'feels' the bottom which slows down the wave. So the shallower parts of the wave slow down more than the parts that are further from the shore. This makes the wave 'bend', which is called **refraction**. The way that waves bend as they come into shore either concentrates wave energy or disperses it. In quiet water areas like bays, wave energy is dispersed and sand gets deposited. Areas like cliffs that stick out into the water, are eroded away by the strong wave energy that concentrates its power on the cliff (Figure 10.15).



Figure 10.15: Cliffs are eroded by wave action that concentrates energy in these areas.

Wave-cut cliffs form where waves cut into the bottom part of the cliff, eroding away the soil and rocks there. First the waves cut a notch into the base of the cliff. If enough material is cut away, the cliff above can collapse into the water. Many years of this type of erosion can form a **wave-cut platform** (Figure 10.16).



Figure 10.16: This large wave cut platform was formed by the cutting action of waves on the cliffs to the left.



Figure 10.18: A sea stack forms if the upper layers of rock collapse, leaving an isolated pinnacle.

If waves erode a cliff from two sides, the erosion produced can form an open area in the cliff called an **arch** (Figure 10.17). If the material above the arch eventually erodes away, a piece of tall rock can remain in the water, which is called a **sea stack** (Figure 10.18).



Figure 10.17: A sea arch can form if waves erode from opposite sides of a cliff.

Wave Deposition

Rivers carry the sand that comes from erosion of mountains and land areas of the continents to the shore. Soil and rock are also eroded from cliffs and shorelines by waves. That material is transported by waves and deposited in quieter water areas. As the waves come onto shore and break, water and particles move along the shore. When lots of sand accumulates in one place, it forms a beach. Beaches can be made of mineral grains, like quartz, but beaches can also be made of pieces of shell or coral or even bits of broken hardened lava (Figure 10.19).



Figure 10.19: Quartz, rock fragments, and shell make up the sand along a beach.

Waves continually move sand grains along the shore. Smaller particles like silt and clay don't get deposited at the shore because the water here is too turbulent. The work of waves moves sand from the beaches on shore to bars of sand offshore as the seasons change. In the summer time, waves of lower energy bring sand up onto the beach and leave it there. That is good for the many people who enjoy sitting on soft sand when they visit the beach (Figure 10.20). In the wintertime, waves and storms of higher energy bring the sand from the beach back offshore. If you visit your favorite beach in the wintertime, you will find a steeper, rockier beach than the flat, sandy beach of summer. Some communities truck in sand to resupply sand to beaches. It is very important to study the energy of the waves and understand the types of sand particles that normally make up the beach before spending lots of money to do this. If the sand that is trucked in has pieces that are small enough to be carried away by the waves on that beach, the sand will be gone in a very short time.



Figure 10.20: Sand deposits in quiet areas along a shoreline to form a beach.



Figure 10.21: These sand dunes are part of Padre Island off the coast of Texas, which is a barrier island.

Sand transported by the work of waves breaking along the shore can form sand bars that stretch across a bay or ridges of sand that extend away from the shore, called **spits**. Sometimes the end of a spit hooks around towards the quieter waters of the bay as waves refract, causing the sand to curve around in the shape of a hook.

When the land that forms the shore is relatively flat and gently sloping, the shoreline may be lined with long narrow islands called **barrier islands** (Figure 10.21). Most barrier islands are just a few kilometers wide and tens of kilometers long. Many famous beaches, like Miami Beach, are barrier islands. In its natural state, a barrier island acts as the first line of defense against storms like hurricanes.

Instead of keeping barrier islands natural, these areas end up being some of the most built up, urbanized areas of our coastlines. That means storms, like hurricanes, damage houses and businesses rather than hitting soft, vegetated sandy areas. Some hurricanes have hit barrier islands so hard that they break right through the island, removing sand, houses and anything in the way.

Protecting Shorelines

Humans build several different types of structures to try to slow down the regular work of erosion that waves produce and to help prevent damage to homes from large storms. One structure that people build, called a **groyne (or groin)**, is a long narrow pile of rocks that extends out into the water, at right angles to the shoreline (Figure 10.22). The groyne traps sand on one side of the groyne, keeping the sand there, rather than allowing it to move along the coastline. This works well for the person who is on the upcurrent side of the groyne, but it causes problems for the people on the opposite, downcurrent side. Those people no longer have sand reaching the areas in front of their homes. What happens as a result is that people must build another groyne to trap sand there. This means lots of people build groynes, but it is not a very good answer to the problem of wave erosion.




Figure 10.22: Groynes are built perpendicular to the shoreline to help keep sand from moving off the beach.

Some other structures that people build include **breakwaters** and **sea walls** (Figure 10.23). Both of these are built parallel to the shoreline. A breakwater is built out away from the shore in the water while a sea wall is built right along the shore. Breakwaters are built in bay areas to help keep boats safe from the energy of breaking waves. Sometimes enough sand deposits in these quiet water areas that people then need to work to remove the sand. Sea walls are built to protect beach houses from waves during severe storms. If the waves in a storm are very large, sometimes they erode away the whole sea wall, leaving the area unprotected entirely. People do not always want to choose safe building practices, and instead choose to build a beach house right on the beach. If you want your beach house to stay in good shape for many years, it is smarter to build your house away from the shore.



Figure 10.23: Breakwaters are visible in this satellite image, built parallel to the shoreline to protect areas from strong waves.

Lesson Summary

- Waves in the ocean are what we see as energy travels through the water.
- The energy of waves produces erosional formations like cliffs, wave cut platforms, sea arches, and sea stacks.
- When waves reach the shore, deposits like beaches, spits, and barrier islands form in certain areas.
- Groynes, jetties, breakwaters, and sea walls are structures humans build to protect the shore from the erosion of breaking waves.

Review Questions

1. Name three structures that people build to try to prevent wave erosion.
2. Name three natural landforms that are produced by wave erosion.
3. What are the names of the parts of a waveform?
4. Describe the process that produces wave refraction.
5. If you were to visit a beach in a tropical area with coral reefs, what would the beach there be made of?

Vocabulary

arch

An erosional landform that is produced when waves erode through a cliff.

barrier island

Long, narrow island, usually composed of sand that serves as nature's first line of defense against storms.

breakwater

Structure built in the water, parallel to the shore to protect boats or harbor areas from strong waves.

groyne

Long, narrow piles of stone or timbers built perpendicular to the shore to trap sand.

sea stack

Isolated tower of rock that forms when a sea arch collapses.

sea wall

Structure built along the shore, parallel to the shore, to protect against strong waves.

spit

Long, narrow bar of sand that forms as waves transport sand along shore.

wave-cut platform

Flat, level area formed by wave erosion as waves undercut cliffs.

wave crest

The highest part of a wave form.

wave height

The vertical distance from wave crest to wave trough.

wavelength

The horizontal distance from wave crest to wave crest.

wave trough

The lowest part of a wave form.

Points to Consider

- What situations would increase the rate of erosion by waves?
- If barrier islands are nature's first line of defense against ocean storms, why do people build on them?
- Could a sea wall ever increase the amount of damage done by waves?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/14.2/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 8.1: Waves and Coastal Features is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

9: Mountain Building

9.1: Tectonic Stress and Geologic Structures
9.2: Mountain Formation

This page titled 9: Mountain Building is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

9.1: Tectonic Stress and Geologic Structures

Causes and Types of Tectonic Stress



Enormous slabs of lithosphere move unevenly over the planet's spherical surface, resulting in earthquakes. This chapter deals with two types of geological activity that occur because of plate tectonics: mountain building and earthquakes. First, we will consider what can happen to rocks when they are exposed to stress. **Stress** is the force applied to an object. In geosciences, stress is the force per unit area that is placed on a rock. Four types of stresses act on materials.

- When stress causes a material to change shape, it has undergone strain or **deformation**. Deformed rocks are common in geologically active areas. A rock's response to stress depends on the rock type, the surrounding temperature, and pressure conditions the rock is under, the length of time the rock is under stress, and the type of stress. The rocks then have three possible responses to increasing stress: elastic deformation, plastic deformation, or fracturing. **Elastic deformation** occurs when the rock returns to its original shape when the stress is removed. When rocks under stress do not return to its original shape when the stress is removed, it is called **plastic deformation**. Finally, when a rock under stress breaks, it's called a **fracture**.

  Under what conditions do you think a rock is more likely to fracture? Is it more likely to break deep within Earth's crust or at the surface? What if the stress applied is sharp rather than gradual? At the Earth's surface, rocks usually break quite quickly, but deeper in the crust, where temperatures and pressures are higher, rocks are more likely to deform plastically. Sudden stress, such as a hit with a hammer, is more likely to make a rock break. Stress applied over time often leads to plastic def

Geologic Structures


Sedimentary rocks are important for deciphering the geologic history of a region because they follow certain rules. First, sedimentary rocks are formed with the oldest layers on the bottom and the youngest on top. Second, sediments are deposited horizontally, so sedimentary rock layers are originally horizontal, as are some volcanic rocks, such as ash falls. Finally, sedimentary rock layers that are not horizontal are deformed in some manner. Often times looking like they are tiling into the earth.

You can trace the deformation a rock has experienced by seeing how it differs from its original horizontal, oldest-on-bottom position. This deformation produces geologic structures such as folds, joints, and faults that are caused by stresses.


FOLDS



Rocks deforming plastically under compressive stresses crumple into folds. They do not return to their original shape. If the rocks experience more stress, they may undergo more folding or even fracture. There are three major types of rock folding: monoclines, synclines, and anticlines. A **monocline** is a simple bend in the rock layers so that they are no longer horizontal. **Anticlines** are folded rocks that arch upward and dip away from the center of the fold. The oldest rocks are at the center of an anticline and the youngest are draped over them. When rocks arch upward to form a circular structure, that structure is called an **adome**. A **syncline** is a fold that bends downward, causing the youngest rocks are to be at the center and the oldest are on the outside. When rocks bend downward in a circular structure, that structure is called **abasin**. If the rocks are exposed at the surface, where are the oldest rocks located?



Monocline diagram





FAULTS

A rock under enough stress will eventually fracture. If there is no movement on either side of a fracture, the fracture is called a **joint**. But if the blocks of rock on one or both sides of a fracture move, the fracture is called a fault. Sudden motions along faults cause rocks to break and move suddenly, releasing the stored up stress energy to create an earthquake.

A **slip** is the distance rocks move along a fault and can be up or down the fault plane. Slip is relative, because there is usually no way to know whether both sides moved or only one. Faults lie at an angle to the horizontal surface of the Earth. That angle is called the fault's **dip**. The dip defines which of two basic types a fault is. If the fault's dip is inclined relative to the horizontal, the fault is a dip-slip fault. There are two types of **dip-slip faults**. In **normal faults**, the hanging wall drops down relative to the footwall. Normal faults can be huge and are often times responsible for uplifting mountain ranges in regions experiencing tensional stress.

With **reverse faults**, the footwall drops down relative to the hanging wall. A type of reverse fault is a **thrust fault**, in which the fault plane angle is nearly horizontal. Rocks can slip many miles along thrust faults.

A **strike-slip fault** is a dip-slip fault in which the dip of the fault plane is vertical and result from shear stresses. California's San Andreas Fault is the world's most famous strike-slip fault. It is a right-lateral strike slip fault.

STRESS AND MOUNTAIN BUILDING

It is the shear power and strength of two or more converging continental plates smash upwards that create mountain ranges. Stresses from this uplift cause folds, reverse faults, and thrust faults, which allow the crust to rise upwards. Subduction of oceanic lithosphere at convergent plate boundaries also builds mountain ranges.

When tensional stresses pull crust apart, it breaks into blocks that slide up and drop down along normal faults. The result is alternating mountains and valleys, known as a basin-and-range.


CC licensed content, Shared previously

- Tectonic Forces. **Provided by**: Open Geography Education. **Located at**: http://www.opengeography.org/ch-5-tectonic-forces.html. **Project**: Dynamic Earth: Introduction to Physical Geography. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 9.1: Tectonic Stress and Geologic Structures is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

9.2: Mountain Formation

Mountain formation refers to the geological processes that underlie the formation of mountains. These processes are associated with large-scale movements of the earth's crust (plate tectonics).[1] Mountain formation is related to plate tectonics. Folding, faulting, volcanic activity, igneous intrusion and metamorphism are all parts of the orogenic process of mountain building.[2] The understanding of specific landscape features in terms of the underlying tectonic processes is called *tectonic geomorphology*, and the study of geologically young or ongoing processes is called *neotectonics*.[3]



Thrust and reverse fault movement are an important component of mountain formation.

Types of mountains

See also: List of mountain types

There are three main types of mountains: volcanic, fold, and block.[4] A more detailed classification useful on a local scale predates plate tectonics and adds to the above categories.[5]

Volcanic mountains

See also: Stratovolcano, Shield volcano and Guyot

Movements of tectonic plates create volcanoes along the plate boundaries, which erupt and form mountains. A *volcanic arc system* is a series of volcanoes that form near a subduction zone where the crust of a sinking oceanic plate melts.[7]




Mount Baker, a stratovolcano in northwestern Washington

Most volcanoes occur in a band encircling the Pacific Ocean (the Pacific Ring of Fire), and in another that extends from the Mediterranean across Asia to join the Pacific band in the Indonesian Archipelago. The most important types of volcanic mountain are *composite cones* or *stratovolcanoes* (Vesuvius, Kilimanjaro and Mount Fuji are examples) and *shield volcanoes* (such as Mauna Loa on Hawaii, a hotspot volcano).[8][9]




Mauna Loa lava flow; a shield volcano.

A shield volcano has a gently sloping cone due to the low viscosity of the emitted material, primarily basalt. Mauna Loa is the classic example, with a slope of 4°-6°. (The relation between slope and viscosity falls under the topic of angle of repose.[10]) The composite volcano or stratovolcano has a more steeply rising cone (33°-40°),[11] due to the higher viscosity of the emitted material, and eruptions are more violent and less frequent than for shield volcanoes. Besides the examples already mentioned are Mount Shasta, Mount Hood and Mount Rainier.[12]



Stratovolcanoes associated with a subduction (left) and a spreading ridge volcano (right). A hotspot volcano is center.[6]



Elements of volcanoes in a *volcanic arc system*.


Fold mountains

See also: Fold (geology), Fold and thrust belt and Fold mountain

When plates collide or undergo subduction (that is – ride one over another), the plates tend to buckle and fold, forming mountains. Most of the major continental mountain ranges are associated with thrusting and folding or orogenesis. Examples are the Jura and the Zagros mountains.[14]



Zard-Kuh, a fold mountain in the central Zagros range of Iran.




Two processes creating mountains: Top:delamination by intrusion of hot asthenosphere; Bottom: Subduction of ocean crust.[13]

Block mountains

See also: Fault-block mountain

When a fault block is raised or tilted, block mountains can result.[16] Higher blocks are called *horsts* and troughs are called *grabens*. A spreading apart of the surface causes tensional forces. When the tensional forces are strong enough to cause a plate to split apart, it does so such that a center block drops down relative to its flanking blocks.



An example is the Sierra Nevada Range, where delamination created a block 650 km long and 80 km wide that consists of many individual portions tipped gently west, with east facing slips rising abruptly to produce the highest mountain front in the continental United States.[17][18]



Sierra Nevada Mountains (formed by delamination) as seen from the International Space Station.

Models

See also: Extensional tectonics, Rift valley and Rift
See also: Prediction of volcanic activity
See also: Geomorphology

Hotspot volcanoes

Hotspots are supplied by a magma source in the Earth's mantle called a mantle plume. Although originally attributed to a melting of subducted oceanic crust, recent evidence belies this connection.[19] The mechanism for plume formation remains a research topic.

Fault blocks

Several movements of the earth's crust that lead to mountains are associated with faults. These movements actually are amenable to analysis that can predict, for example, the height of a raised block and the width of an intervening rift between blocks using the rheology of the layers and the forces of isostasy. Early bent plate models predicting fractures and fault movements have evolved into today's kinematic and flexural models.[20][21]

Notes

1. ^ Steven M. Stanley (2004). "Mountain building". *Earth system history* (2nd ed.). Macmillan. p. 207. ISBN 0-7167-3907-0.
2. ^ Robert J. Twiss, Eldridge M. Moores (1992). "Plate tectonic models of orogenic core zones". *Structural Geology* (2nd ed.). Macmillan. p. 493. ISBN 0-7167-2252-6.
3. ^ Kurt Stüwe (2007). "§4.5 Geomorphology". *Geodynamics of the lithosphere: an introduction* (2nd ed.). Springer. p. 178. ISBN 3-540-71236-4.
4. ^ "Chapter 6: Mountain building". *Science matters: earth and beyond; module 4*. Pearson South Africa. 2002. p. 75. ISBN 0-7986-6059-7.
5. ^ Andrew Goudie (2004). *Encyclopedia of geomorphology; Volume 2*. Routledge. p. 701. ISBN 0-415-32738-5.
6. ^ Victor Schmidt, William Harbert (2003). *Planet Earth and the New Geoscience* (4th ed.). Kendall Hunt. pp. 46–47. ISBN 0-7872-9355-5.
7. ^ Stephen D Butz (2004). "Chapter 8: Plate tectonics". *Science of Earth Systems*. Thompson/Delmar Learning. p. 136. ISBN 0-7668-3391-7.
8. ^ John Gerrard (1990). "Types of volcano". *Mountain environments: an examination of the physical geography of mountains*. MIT Press. p. 194. ISBN 0-262-07128-2.
9. ^ Robert Wayne Decker, Barbara Decker (2005). "Chapter 8: Hot spots". *Volcanoes* (4th ed.). Macmillan. p. 113 *ff*. ISBN 0-7167-8929-9.
10. ^ Arthur Holmes, Donald Duff (2004). *Holmes Principles of Physical Geology* (4th ed.). Taylor & Francis. p. 209. ISBN 0-7487-4381-2.
11. ^ *Transactions of the American Society of Civil Engineers, Volume 39*. American Society of Civil Engineers. 1898. p. 62.
12. ^ James Shipman, Jerry D. Wilson, Aaron Todd (2007). "Minerals, rocks and volcanoes". *An Introduction to Physical Science* (12th ed.). Cengage Learning. p. 650. ISBN 0-618-93596-7.
13. ^ N. H. Woodcock, Robin A. Strachan (2000). "Chapter 12: The Caledonian Orogeny: a multiple plate collision". *Geological history of Britain and Ireland*. Wiley-Blackwell. p. 202, Figure 12.11. ISBN 0-632-03656-7.
14. ^ Michael P Searle (2007). "Diagnostic features and processes in the construction and evolution of Oman-, Zagros-, Himalyan-, Karakoram-, and Tibetan type orogenic belts". In Robert D. Hatcher Jr., MP Carlson, JH McBride & JR Martinez Catalán. *4-D framework of continental crust*. Geological Society of America. p. 41 *ff*. ISBN 0-8137-1200-9.
15. ^ Chris C. Park (2001). "Figure 6.11". *The environment: principles and applications* (2nd ed.). Routledge. p. 160. ISBN 0-415-21770-9.
16. ^ Scott Ryan (2006). "Figure 13-1". *CliffsQuickReview Earth Science*. Wiley. ISBN 0-471-78937-2.
17. ^ John Gerrard (1990-04-12). *Reference cited*. p. 9. ISBN 0-262-07128-2.
18. ^ Lee, C.-T.; Yin, Q; Rudnick, RL; Chesley, JT; Jacobsen, SB (2000). "Osmium Isotopic Evidence for Mesozoic Removal of Lithospheric Mantle Beneath the Sierra Nevada, California". *Science* **289** (5486): 1912–6. Bibcode:2000Sci…289.1912L. doi:10.1126/science.289.5486.1912. PMID 10988067.
19. ^ Y Niu & MJ O'Hara (2004). "Chapter 7: Mantle plumes are NOT from ancient oceanic crust". In Roger Hékinian, Peter Stoffers, Jean-Louis Cheminée. *Oceanic hotspots: intraplate submarine magmatism and tectonism*. Springer. p. 239 *ff*. ISBN 3-540-40859-2.
20. ^ AB Watts (2001). "§7.2 Extensional tectonics and rifting". *Isostasy and flexure of the lithosphere*. Cambridge University Press. p. 295. ISBN 0-521-00600-7.
21. ^ GD Karner & NW Driscoll (1999). "Style, timing and distribution of tectonic deformation across the Exmouth Plateau, northwest Australia, determined from stratal architecture and quantitative basin modelling". In Conall Mac Niocaill, Paul Desmond Ryan. *Continental tectonics*. Geological society. p. 280. ISBN 1-86239-051-7.

External links

CC licensed content, Shared previously

- Mountain formation. **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Mountain_formation. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 9.2: Mountain Formation is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

10: Geologic History

10.1: Relative Ages of Rocks
10.2: Absolute Ages of Rocks
10.3: Fossils
10.4: History of Earth’s Life Forms
10.5: Early Earth
10.6: Geologic Time Scale

This page titled 10: Geologic History is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.1: Relative Ages of Rocks

In 1666, a young doctor named Nicholas Steno was invited to dissect the head of an enormous great white shark that had been caught by local fisherman near Florence, Italy. Steno was struck by the resemblance of the shark's teeth to fossils, known as "tongue stones", recovered from inland mountains and hills (Figure 11.9).






Figure 11.9
: Fossil shark tooth (left) and modern shark teeth (right).



Figure 11.10: Tyrannosaurus rex fossil resembling a living organism.

While it may seem obvious today, most people at the time did not believe that fossils were once part of living creatures. The reason was that the fossils of clams, snails, and other marine animals were found in tall mountains, miles from any ocean. Two schools of thought explained these fossils. Some religious writers believed that the shells were washed up during the Biblical flood. But this explanation could not account for the fact that fossils were not only found on mountains, but also within mountains, in rocks that had been quarried from deep below Earth's surface. Seeking an alternate explanation, other writers proposed that the fossils had formed *within* the rocks as a result of mysterious forces. In other words, fossil shells, bones, and teeth were never a part of a living creature!

Steno had other ideas. For Steno, the close resemblance between fossils and modern organisms was impossible to ignore. Instead of invoking supernatural forces to explain fossils, Steno concluded that fossils were once parts of living creatures. He then sought to explain how fossil seashells could be found in rocks far from any ocean. As in the Tyrannosaurus rex Figure 11.10, fossils resemble living organisms.

Superposition of Rock Layers

Steno first proposed that if a rock contained the fossils of marine animals, the rock was formed from sediments that were deposited on the seafloor. These rocks were then uplifted to become mountains. Based on those assumptions, Steno made a remarkable series of conjectures that are now known as **Steno's Laws**.

Original Horizontality

Because sediments are deposited under water, they will form flat, horizontal layers (Figure 11.11). If a sedimentary rock is found tilted, the layer was tilted after it was formed.




Figure 11.11: Sedimentary layers that have been deposited horizontally.

Lateral Continuity

Sediments were deposited in continuous sheets that spanned the body of water that they were deposited in. When a valley cuts through sedimentary layers, it can be assumed that the rocks on either side of the valley were originally continuous.

Superposition

Sedimentary rocks are deposited one on top of another. Therefore, the youngest layers are found at the top, and the oldest layers are found at the bottom of the sequence.

Cross-Cutting Relationships



Figure 11.12: Cross cutting relationships: older banded gneiss with a white granite intrusion. The granite must be younger than the gneiss, because it cuts across the existing gneiss.

A rock formation or surface that cuts across other rock layers is younger than the rock layers it disturbs. For example, if an igneous intrusion goes through a series of metamorphic rocks, the intrusion must be younger than the metamorphic rocks that it cuts through (Figure 11.12).

The Grand Canyon provides an excellent illustration of Steno's laws. Figure 11.13 shows the many horizontal layers of sedimentary rock that make up the canyon. This nicely illustrates the principle of original horizontality. The youngest rock layers are at the top of the canyon, while the oldest are at the bottom, which is described by the law of superposition. Distinctive rock layers, such as the Kaibab Limestone, can be matched across the broad expanse of the canyon. We know these rock layers were once connected, which is described in the rule of lateral continuity. Finally, the Colorado River cuts through all the layers of sedimentary rock to form the canyon. Based on the principle of cross-cutting relationships, the river must be younger than all of the rock layers that it cuts through.



Figure 11.13: Grand Canyon, with the Kaibab Limestone visible at the top of the canyon.

Determining the Relative Ages of Rocks

The **relative age** of a rock is its age in comparison with other rocks. If you know the relative ages of two rock layers, you know which is older and which is younger, but you do not know how old the layers are in years. In some cases, it is very tricky to determine the sequence of events that leads to a certain formation. Take the example, Figure 11.14:




Figure 11.14: Cross-section of sedimentary layers: (A-C) igneous intrusion, (D) cross-section, (E) fault.

The principle of cross-cutting relationships states that a fault or intrusion is younger than the rocks that it cuts through. The fault labeled 'E' cuts through all three sedimentary rock layers (A, B, and C) and also cuts through the intrusion (D). So the fault must be the youngest formation that is seen. The intrusion (D) cuts through the three sedimentary rock layers, so it must be younger than those layers.

The principle of superposition states that the oldest sedimentary rock units are at the bottom, and the youngest are at the top. Based on this, layer C is oldest, followed by B and A. So the full sequence of events is as follows:

1. Layer C formed.
2. Layer B formed.
3. Layer A formed.
4. When layers A-B-C were present, intrusion D formed.
5. Intrusion D cut through layers A-C.
6. Fault E formed, shifting rocks A through C and intrusion D.
7. Weathering and erosion occurred, forming a layer of soil on top of layer A.

Unconformities in Rock Layers

Steno discovered the rules for determining the relative age of rock beds, but he did not have a good understanding of how long it would take for these rock formations to form. At the time, most Europeans believed that the Earth was around 6,000 years old, a figure that was based on the amount of time estimated for the events described in the Bible. One of the first to question this time scale was a Scottish geologist named James Hutton (1726-1797). Often described as the founder of modern geology, Hutton formulated a philosophy called **uniformitarianism**: *The present is the key to the past*. According to uniformitarianism, the same processes we see around us today operated in the past as well. For example, if erosion and deposition occur slowly now, they probably have always occurred slowly.

Hutton discovered places where sedimentary rock beds lie on an eroded surface. Such a formation is called an **unconformity**, or a gap in rock layers, where some rocks were eroded away. Hutton reconstructed the sequence of events that led to this formation. For example, consider the famous unconformity at Siccar Point, on the coast of Scotland (Figure 11.15).



Figure 11.15: Hutton's Unconformity on the coast of Scotland.

Based on figure 15, at least nine geological events can be inferred:

1. A series of sedimentary beds is deposited on an ocean floor.
2. The sediments harden into sedimentary rock.
3. The sedimentary rocks are uplifted and tilted, exposing them above the ocean surface.
4. The tilted beds are eroded by rain, ice, and wind to form an irregular surface.
5. A sea covers the eroded sedimentary rock layers.
6. New sedimentary layers are deposited.
7. The new layers harden into sedimentary rock.
8. These layers are tilted.
9. Uplift occurs, exposing the new sedimentary rocks above the ocean surface.

Hutton realized that an enormous period of time was needed to account for the repeated episodes of deposition, rock formation, uplift, and erosion that led to the formation of an unconformity, like the one at Siccar Point. Hutton realized that the age of Earth should not be measured in thousands of years, but millions of years.

Matching Rock Layers

Superposition and cross-cutting are helpful when rocks are touching one another, but are useless when rocks are kilometers or even continents apart. Three kinds of clues help geologists match rock layers across great distances. The first is the fact that some sedimentary rock formations span vast distances, recognizable across large regions. For example, the Pierre Shale formation can be recognized across the Great Plains, from New Mexico to North Dakota. The famous White Cliffs of Dover in southwest England can be matched to similar white cliffs in Denmark and Germany.

A second clue could be the presence of a **key bed**, or a particularly distinctive layer of rock that can be recognized across a large area. Volcanic ash flows are often useful as key beds because they are widespread and easy to identify. Probably the most famous

example of a key bed is a layer of clay found at the boundary between the Cretaceous Period and the Tertiary Period, the time that the dinosaurs went extinct (Figure 11.16). This thin layer of sediment, only a few centimeters thick, contains a high concentration of the element iridium. Iridium is rare on Earth but common in asteroids. In 1980, a team of scientists led by Luis Alvarez and his son Walter proposed that a huge asteroid struck Earth about 66 million years ago, causing forest fires, acid rain, and climate change that wiped out the dinosaurs.



Figure 11.16

: White layer of clay that marks the Cretaceous-Tertiary Boundary.




Figure 11.17: Geologic time scale.

A third type of clue that helps scientists compare different rock layers is index fossils. Recall that index fossils are the remains of organisms that were widespread but only existed for a relatively short period of time. If two rock units both contain the same type of index fossil, their age is probably very similar.

As scientists collected fossils from all over the world, they recognized that rocks of different ages contain distinctive types of fossils. This pattern led to the creation of the **geologic time scale** and helped to inspire Darwin's theory of evolution (Figure 11.17).

Each era, period, and epoch of the geologic time scale is defined by the fossils that appeared at that time. For example, Paleozoic rocks typically contain trilobites, brachiopods, and crinoid fossils. The presence of dinosaur bones indicate that a rock is from the Mesozoic era, and the particular type of dinosaur will allow the rock to be identified as Triassic, Jurassic, or Cretaceous. The Cenozoic Era is also known as the Age of Mammals, and the Quaternary Period represents the time when the first humans spread across Earth.

Lesson Summary

Nicholas Steno first formulated the principles that allow scientists to determine the relative ages of rocks in the 17th century. Steno stated that sedimentary rocks are formed in continuous, horizontal layers, with younger layers on top of older layers. A century later, James Hutton discovered the law of cross-cutting relationships: a fault or igneous intrusion is younger than the rocks that it cuts through. Hutton also was the first to realize the vast amounts of time that would be needed to create an unconformity, a place where sedimentary rocks lie above an eroded surface.

Other methods come into play when comparing rock layers that are separated by a large distance. Many sedimentary rock formations are large and can be recognized across a region. Distinctive rock layers, called key beds, are also useful for correlating rock units. Fossils, especially index fossils, are the most useful way to compare different rock layers. Changes of fossils over time led to the development of the geologic time scale.

Review Questions

1. In the 15th century, a farmer finds a rock that looks exactly like a clamshell. What did the farmer probably conclude about how the fossil got there?
2. Which of Steno's Laws is illustrated by each of the following images in Figure 11.18?
3. What is the sequence of rock units in Figure 11.19, from oldest to youngest?
4. What kind of geological formation is shown in the outcrop in Figure 11.20, and what sequence of events does it represent?
5. The three outcrops in Figure 11.21 are very far apart. Based on what you see, which fossil is an index fossil, and why?




Figure 11.18: Illustration of Steno's Laws.




Figure 11.19: Sequence of rock units.



Figure 11.20: Outcrop.




Figure 11.21: Fossils.

Vocabulary

cross-cutting relationships
One of Steno's principles that states that an intrusion or fault is younger than the rocks that it cuts through.

geologic time scale
A division of Earth's history into blocks of time distinguished by geologic and evolutionary events.

key bed
A distinctive, widespread rock layer that formed at a single time.

lateral continuity
One of Steno's principles that states that a sedimentary rock layer extends sideways as wide as the basin in which it forms.

original horizontality
One of Steno's principles that states that sedimentary layers were horizontal or flat lying at the time they were deposited.

relative age
The age of an object in comparison with the age of other objects.

superposition
One of Steno's principles that states that in a sequence of sedimentary rock layers, the oldest layer is at the bottom and the youngest layer is at the top.

unconformity
A boundary between rocks of very different ages. Unconformities are often marked by an erosional surface.

uniformitarianism

The idea that the geologic processes that shape the land today have acted in basically the same way throughout Earth's history.

Points to Consider

- In Nicholas Steno's time, why didn't most people believe the fossils were the remains of ancient organisms?
- How did Steno explain the presence of marine fossils in high mountains?
- What was the significance of unconformities to James Hutton?
- How can you determine the relative age of two rock layers that are very far apart?

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Relative_Ages_of_Rocks. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.1: Relative Ages of Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.2: Absolute Ages of Rocks

As we learned in the previous lesson, index fossils and superposition are effective methods of determining the relative age of objects. In other words, you can use superposition to tell you that one rock layer is older than another. But determining the **absolute age** of a substance (its age in years) is a much greater challenge. To accomplish this, scientists use a variety of evidence, from tree rings to the amounts of radioactive materials in a rock.

Lesson Objectives

- Define the difference between absolute age and relative age.
- Describe four methods of absolute dating.
- Explain what radioactivity is and give examples of radioactive decay.
- Explain how the decay of radioactive materials helps to establish the age of an object.
- Estimate the age of an object, given the half-life and the amounts of radioactive and daughter materials.
- Give four examples of radioactive materials that are used to date objects, and explain how each is used.

Tree Rings

In regions outside the tropics, trees grow more quickly during the warm summer months than during the cooler winter. This pattern of growth results in alternating bands of light-colored, low density “early wood” and dark, high density “late wood”. Each dark band represents a winter; by counting rings it is possible to find the age of the tree (Figure 11.22). The width of a series of growth rings can give clues to past climates and various disruptions such as forest fires. Droughts and other variations in the climate make the tree grow slower or faster than normal, which shows up in the widths of the tree rings. These tree ring variations will appear in all trees growing in a certain region, so scientists can match up the growth rings of living and dead trees. Using logs recovered from old buildings and ancient ruins, scientists have been able to compare tree rings to create a continuous record of tree rings over the past 2,000 years. This tree ring record has proven extremely useful in creating a record of climate change, and in finding the age of ancient structures.



Figure 11.22: Cross-section showing growth rings. The thick, light-colored part of each ring represents rapid spring and summer growth. The thin, dark part of each ring represents slow autumn and winter growth.

Ice Cores and Varves

Several other processes result in the accumulation of distinct yearly layers that can be used for dating. For example, layers form within glaciers because there tends to be less snowfall in the summertime, allowing a dark layer of dust to accumulate on top of the winter snow (Figure 11.23). To study these patterns, scientists drill deep into ice sheets, producing cores hundreds of meters long. Scientists analyze these ice cores to determine how the climate has changed over time, as well as to measure concentrations of atmospheric gases. The longest cores have helped to form a record of polar climate stretching hundreds of thousands of years back.




Figure 11.23: Ice core section showing annual layers.

Another example of yearly layers is the deposition of sediments in lakes, especially the lakes that are located at the end of glaciers. Rapid melting of the glacier in the summer results in a thick, sandy deposit of sediment. These thick layers alternate with thin, clay-rich layers deposited during the winter. The resulting layers, called **varves**, give scientists clues about past climate conditions. For example, an especially warm summer might result in a very thick layer of sediment deposited from the melting glacier. Thinner varves can indicate colder summers, because the glacier doesn't melt as much and carry as much sediment into the lake.

Age of Earth




Figure 11.24: Lord Kelvin.

While tree rings and other annual layers are useful for dating relatively recent events, they are not of much use on the vast scale of geologic time. During the 18th and 19th centuries, geologists tried to estimate the age of Earth with indirect techniques. For example, geologists measured how fast streams deposited sediment, in order to try to calculate how long the stream had been in existence. Not surprisingly, these methods resulted in wildly different estimates, from a few million years to "quadrillions of years". Probably the most reliable of these estimates was produced by the British geologist Charles Lyell, who estimated that 240 million years have passed since the appearance of the first animals with shells. Today scientists know his estimate was too young; we know that this occurred about 530 million years ago.

In 1892, William Thomson (later known as Lord Kelvin) calculated the age of Earth in a systematic fashion (Figure 11.24). He assumed that the Earth began as a ball of molten rock, which has steadily cooled over time. From these assumptions, he calculated that the Earth was 100 million years old. This estimate was a blow to geologists and supporters of Charles Darwin's theory of evolution, which required an older Earth to provide time for evolution to take place.

Thomson's calculations, however, were soon shown to be flawed when radioactivity was discovered in 1896. **Radioactivity** is the tendency of certain atoms to decay into lighter atoms, emitting energy in the process. Radioactive materials in Earth's interior provide a steady source of heat. Calculations of Earth's age using radioactive decay showed that Earth is actually much older than Thomson calculated.

Radioactive Decay

The discovery of radioactive materials did more than disprove Thomson's estimate of Earth's age. It provided a way to find the absolute age of a rock. To understand how this is done, it is necessary to review some facts about atoms.

Atoms contain three particles: protons, neutrons, and electrons. Protons and neutrons are located in the nucleus, while electrons orbit around the nucleus. The number of protons determines which element you're examining. For example, all atoms of carbon have six protons, all atoms of oxygen have eight protons, and all atoms of gold have 79 protons. The number of neutrons, however, is variable. An atom of an element with a different number of neutrons is an **isotope** of that element. For example, the isotope carbon-12 contains 6 neutrons in its nucleus, while the isotope carbon-13 has 7 neutrons.

Some isotopes are **radioactive**, which means they are unstable and likely to decay. This means the atom will spontaneously change from an unstable form to a stable form. There are two forms of nuclear decay that are relevant in how geologists can date rocks (Table (11.1):

Particle	Composition	Effect on Nucleus
Alpha	2 protons, 2 neutrons	The nucleus contains two fewer protons and two fewer neutrons.
Beta	1 electron	One neutron decays to form a proton and an electron, which is emitted.

If an element decays by losing an alpha particle, it will lose 2 protons and 2 neutrons. If an atom decays by losing a beta particle, it loses just one electron.

So what does this have to do with the age of Earth? Radioactive decay eventually results in the formation of stable **daughter products**. Radioactive materials decay at known rates. As time passes, the proportion of radioactive isotopes will decrease and the proportion of daughter isotopes will increase. A rock with a relatively high proportion of radioactive isotopes is probably very young, while a rock with a high proportion of daughter products is probably very old.

Scientists measure the rate of radioactive decay with a unit called **half-life**. The half-life of a radioactive substance is the amount of time, on average, it takes for half of the atoms to decay. For example, imagine a radioactive substance with a half-life of one year. When a rock is formed, it contains a certain number of radioactive atoms. After one year (one half-life), half of the radioactive atoms have decayed to form stable daughter products, and 50% of the radioactive atoms remain. After another year (two half-lives), half of the remaining radioactive atoms have decayed, and 25% of the radioactive atoms remain. After the third year (three half-lives), 12.5% of the radioactive atoms remain. After four years (four half-lives), 6.25% of the radioactive atoms remain, and after 5 years (five half-lives), only 3.125% of the radioactive atoms remain.

If you find a rock whose radioactive material has a half life of one year and measure 3.125% radioactive atoms and 96.875% daughter atoms, you can assume that the substance is 5 years old. The decay of radioactive materials can be shown with a graph (Figure 11.25). If you find a rock with 75% of the radioactive atoms remaining, about how old is it?




Figure 11.25: Decay of an imaginary radioactive substance with a half-life of one year.

Radiometric Dating of Rocks

In the process of **radiometric dating**, several isotopes are used to date rocks and other materials. Using several different isotopes helps scientists to check the accuracy of the ages that they calculate.

Carbon Dating

Earth's atmosphere contains three isotopes of carbon. Carbon-12 is stable and accounts for 98.9% of atmospheric carbon. Carbon-13 is also stable and accounts for 1.1% of atmospheric carbon. Carbon-14 is radioactive and is found in tiny amounts. Carbon-14 is produced naturally in the atmosphere when cosmic rays interact with nitrogen atoms. The amount of carbon-14 produced in the atmosphere at any particular time has been relatively stable through time.

Radioactive carbon-14 decays to stable nitrogen-14 by releasing a beta particle. The nitrogen atoms are lost to the atmosphere, but the amount of carbon-14 decay can be estimated by measuring the proportion of radioactive carbon-14 to stable carbon-12. As a substance ages, the relative amount of carbon-14 decreases.

Carbon is removed from the atmosphere by plants during the process of photosynthesis. Animals consume this carbon when they eat plants or other animals that have eaten plants. Therefore carbon-14 dating can be used to date plant and animal remains. Examples include timbers from an old building, bones, or ashes from a fire pit. Carbon dating can be effectively used to find the age of materials between 100 and 50,000 years old.

Potassium-Argon Dating

Potassium-40 decays to argon-40 with a half-life of 1.26 billion years. Because argon is a gas, it can escape from molten magma or lava. Therefore any argon that is found in a crystal probably formed as a result of the decay of potassium-40. Measuring the ratio of potassium-40 to argon-40 will yield a good estimate of the age of the sample.

Potassium is a common element found in many minerals such as feldspar, mica, and amphibole. The technique can be used to date igneous rocks from 100,000 years to over a billion years old. Because it can be used to date geologically young materials, the technique has been useful in estimating the age of deposits containing the bones of human ancestors.

Uranium-Lead Dating

Two isotopes of uranium are used for radiometric dating. Uranium-238 decays to form lead-206 with a half-life of 4.47 billion years. Uranium-235 decays to form lead-207 with a half-life of 704 million years.

Uranium-lead dating is usually performed on crystals of the mineral zircon (Figure 11.26). When zircon forms in an igneous rock, the crystals readily accept atoms of uranium but reject atoms of lead. Therefore, if any lead is found in a zircon crystal, it can be assumed that it was produced from the decay of uranium.



Figure 11.26: Zircon crystal.

Uranium-lead dating can be used to date igneous rocks from 1 million years to around 4.5 billion years old. Some of the oldest rocks on Earth have been dated using this method, including zircon crystals from Australia that are 4.4 billion years old.

Limitations of Radiometric Dating

Radiometric dating can only be used on materials that contain measurable amounts of radioactive materials and their daughter products. This includes organic remains (which compared to rocks are relatively young, less than 100,000 years old) and older rocks. Ideally, several different radiometric techniques will be used to date the same rock. Agreement between these values indicates that the calculated age is accurate.

In general, radiometric dating works best for igneous rocks and is not very useful for determining the age of sedimentary rocks. To estimate the age of a sedimentary rock deposit, geologists search for nearby or interlayered igneous rocks that can be dated. For

example, if a sedimentary rock layer is sandwiched between two layers of volcanic ash, its age is between the ages of the two ash layers.

Using a combination of radiometric dating, index fossils, and superposition, geologists have constructed a well-defined timeline of Earth history. For example, an overlying lava flow can give a reliable estimate of the age of a sedimentary rock formation in one location. Index fossils contained in this formation can then be matched to fossils in a different location, providing a good age measurement for that new rock formation as well. As this process has been repeated all over the world, our estimates of rock and fossil ages has become more and more accurate.

Lesson Summary

Techniques such as superposition and index fossils can tell you the relative age of objects, which objects are older and which are younger. Other types of evidence are needed to establish the absolute age of objects in years. Geologists use a variety of techniques to establish absolute age, including radiometric dating, tree rings, ice cores, and annual sedimentary deposits called varves.

Radiometric dating is the most useful of these techniques—it is the only technique that can establish the age of objects older than a few thousand years. The concentrations of several radioactive isotopes (carbon-14, potassium-40, uranium-235 and -238) and their daughter products are used to determine the age of rocks and organic remains.

Review Questions

1. What four techniques are used to determine the absolute age of an object or event?
2. A radioactive substance has a half-life of 5 million years. What is the age of a rock in which 25% of the original radioactive atoms remain?
3. **Figure 11.27**: Radioactive decay of Carbon-14
4. Which radioactive isotope or isotopes would you use to date each of the following objects? Explain each of your choices.
    - A 4 billion year old piece of granite.
    - A one million year old bed of volcanic ash that contains the footprints of hominids (human ancestors).
    - The fur of a woolly mammoth that was recently recovered frozen in a glacier.
    - A fossilized trilobite recovered from a bed of sandstone that is about 500 million years old.
5. The principle of uniformitarionism states that the present is the key to the past. In other words, the processes that we see happening today probably worked in a similar way in the past. Why is it important to assume that the rate of radioactive decay has remained constant over time?

Vocabulary

absolute age
: The age of an object in years.

alpha particle
: Particle consisting of two protons and two neutrons that is ejected from the nucleus during radioactive decay.

beta particle
: Particle consisting of a single electron that is ejected from the nucleus during radioactive decay. A beta particle is created when a neutron decays to form a proton and the emitted electron.

daughter product
: Stable substance that is produced by the decay of a radioactive substance. For example, uranium-238 decays to produce lead-207.

half-life
: Amount of time required for half of the atoms of a radioactive substance to decay and form daughter products.

ice core
: Cylinder of ice extracted from a glacier or ice sheet.

radioactive
: Substance that is unstable and likely to emit energetic particles and radiation.

radioactivity

Emission of high-energy particles and/or radiation by certain unstable atoms.

radiometric dating

Process of using the concentrations of radioactive substances and daughter products to estimate the age of a material. As substances age, the amounts of radioactive atoms decrease while the amounts of daughter materials increase.

tree ring

Layer of wood in a tree that forms in one year. You can determine the age of a tree by counting its rings.

varve

Thin layer of sediment deposited on a lakebed over the course of one year usually found at the bottom of glacial lakes.

Points to Consider

- Why are techniques like tree rings, ice cores, and varves only useful for events that occurred in the last few thousand years?
- Why was it so important for Darwin and his followers to prove that the Earth was very old?
- Why is it important to use more than one method to find the age of a rock or other object?

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Absolute_Ages_of_Rocks. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.2: Absolute Ages of Rocks is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.3: Fossils

Throughout human history, people have discovered fossils and wondered about the creatures that lived long ago. In ancient times, fossils inspired legends of monsters and other strange creatures. The Chinese writer Chang Qu reports the discovery of “dragon bones”, which were probably dinosaur fossils in China 2,000 years ago. The griffin, a mythical creature with a lion’s body and an eagle’s head and wings, was probably based on skeletons of *Protoceratops* that were discovered by nomads in Central Asia (Figure 11.1).





Figure 11.1: Griffin (left) and *Protoceratops* (right).

Another fossil reminded the Greeks of the coiled horns of a ram. The Greeks named them ammonites after the ram god *Ammon*. Similarly, legends of the Cyclops may be based on fossilized elephant skulls found in Crete and other Mediterranean islands. Can you see why (Figure 11.2)?





Figure 11.2: Ammonite (left) and elephant skull (right).

Many of the real creatures whose bones became fossilized were no less marvelous than the mythical creatures they inspired (Figure 11.3). The giant pterosaur *Quetzalcoatlus* had a wingspan of up to 12 meters (39 feet). The dinosaur *Argentinosaurus* had an

estimated weight of 80,000 kg, equal to the weight of seven elephants! Other fossils, such as the trilobite and ammonite, impress us with their bizarre forms and delicate beauty.



Figure 11.3: Kolihapeltis sp (left) and Ammonite (right).

Lesson Objectives

- Explain why it is rare for an organism to be preserved as a fossil.
- Distinguish between body fossils and trace fossils.
- Describe five types of fossilization.
- Explain the importance of index fossils, and give several examples.
- Describe what a living fossil is.

How Fossils Form

A **fossil** is any remains or trace of an ancient organism. Fossils include **body fossils**, left behind when the soft parts have decayed away, as well as **trace fossils**, such as burrows, tracks, or fossilized waste (feces) (Figure 11.4).



Figure 11.4: Coprolite (fossilized waste or feces) from a meat-eating dinosaur.

The process of a once living organism becoming a fossil is called **fossilization**. Fossilization is a very rare process: of all the organisms that have lived on Earth, only a tiny percentage of them ever become fossils. To see why, imagine an antelope that dies on the African plain. Most of its body is quickly eaten by scavengers, and the remaining flesh is soon eaten by insects and bacteria, leaving behind only scattered bones. As the years go by, the bones are scattered and fragmented into small pieces, eventually turning into dust and returning their nutrients to the soil. It would be rare for any of the antelope's remains to actually be preserved as a fossil.




Figure 11.5: Fossil shell that has been attacked by a boring sponge.

On the ocean floor, a similar process occurs when clams, oysters, and other shellfish die. The soft parts quickly decay, and the shells are scattered over the sea floor. If the shells are in shallow water, wave action soon grinds them into sand-sized pieces. Even if they are not in shallow water, the shells are attacked by worms, sponges, and other animals (Figure 11.5).

For animals that lack hard shells or bones, fossilization is even more rare. As a result, the fossil record contains many animals with shells, bones, or other hard parts, and few softbodied organisms. There is virtually no fossil record of jellyfish, worms, or slugs. Insects, which are by far the most common land animals, are only rarely found as fossils. Because mammal teeth are much more resistant than other bones, a large portion of the mammal fossil record consists of teeth. This means the fossil record will show many organisms that had shells, bones or other hard parts and will almost always miss the many soft-bodied organisms that lived at the same time.

Because most decay and fragmentation occurs at the surface, the main factor that contributes to fossilization is quick burial. Marine animals that die near a river delta may be buried by sediment carried by the river. A storm at sea may shift sediment on the ocean floor, covering and helping to preserve skeletal remains.

On land, burial is rare, so consequently fossils of land animals and plants are less common than marine fossils. Land organisms can be buried by mudslides or ash from a volcanic eruption, or covered by sand in a sandstorm. Skeletons can be covered by mud in lakes, swamps, or bogs as well. Some of the best-preserved skeletons of land animals are found in the La Brea Tar Pits of Los Angeles, California. Although the animals trapped in the pits probably suffered a slow, miserable death, their bones were preserved perfectly by the sticky tar.

In spite of the difficulties of preservation, billions of fossils have been discovered, examined, and identified by thousands of scientists. The fossil record is our best clue to the history of life on Earth, and an important indicator of past climates and geological conditions as well. The fossil record also plays a key role in our lives. Fossil fuels such as coal, gas, and oil formed from the decayed remains of plants and animals that lived millions of years ago.

Types of Fossils

Fossilization can occur in many ways. Most fossils are preserved in one of five processes (Figure 11.6): preserved remains, permineralization, molds and casts, replacement, and compression.

Preserved Remains

The rarest form of fossilization is the preservation of original skeletal material and even soft tissue. For example, insects have been preserved perfectly in **amber**, which is ancient tree sap. Several mammoths and even a Neanderthal hunter have been discovered frozen in glaciers. These preserved remains allow scientists the rare opportunity to examine the skin, hair, and organs of ancient creatures. Scientists have collected DNA from these remains and compared the DNA sequences to those of modern creatures.

Permineralization

The most common method of fossilization is **permineralization**. After a bone, wood fragment, or shell is buried in sediment, it may be exposed to mineral-rich water that moves through the sediment. This water will deposit minerals into empty spaces, producing a fossil. Fossil dinosaur bones, petrified wood, and many marine fossils were formed by permineralization.

Molds and Casts

In some cases, the original bone or shell dissolves away, leaving behind an empty space in the shape of the shell or bone. This depression is called a **mold**. Later the space may be filled with other sediments to form a matching **cast** in the shape of the original organism. Many mollusks (clams, snails, octopi and squid) are commonly found as molds and casts because their shells dissolve easily.

Replacement

In some cases, the original shell or bone dissolves away and is replaced by a different mineral. For example, shells that were originally calcite may be replaced by dolomite, quartz, or pyrite. If quartz fossils are surrounded by a calcite matrix, the calcite can be dissolved away by acid, leaving behind an exquisitely preserved quartz fossil.

Compression

Some fossils form when their remains are compressed by high pressure. This can leave behind a dark imprint of the fossil. Compression is most common for fossils of leaves and ferns, but can occur with other organisms, as well.




Figure 11.6: Five types of fossils: insect preserved in amber, petrified wood, cast and mold of a clam shell, compression fossil of a fern and pyritized ammonite.

Exceptional Preservation

Some rock beds have produced exceptional fossils. Fossils from these beds may show evidence of soft body parts that are not normally preserved. Two of the most famous examples of soft organism preservation are the Burgess Shale in Canada and the Solnhofen Limestone in Germany. The Burgess Shale is 505 million years old and records the first explosion of shelled organisms in Earth's oceans. Many of the Burgess Shale fossils are bizarre animals that seem unrelated to any other animal group. The Solnhofen Limestone is 145 million years old and contains fossils of many soft-bodied organisms that are not normally preserved,

such as jellyfish. The most famous Solnhofen fossil is *Archaeopteryx*, one of the earliest birds. Although it resembles a dinosaur fossil, impressions of feathers can clearly be seen (Figure 11.7).




Figure 11.7: Fossils from Lagerstätten: *Archaeopteryx* (left) and *Anomalocaris* (right). Archaeopteryx was an early bird. *Anomalocaris* was an enormous predator (one meter long) that lived 500 million years ago.

Index Fossils and Living Fossils

The fossil record shows clearly that over time, life on Earth has changed. Fossils in relatively young rocks tend to resemble animals and plants that are living today. In older rocks, fossils are less similar to modern organisms.

As scientists collected fossils from different rock layers and formations, they discovered that they could often recognize the rock layer by the assemblage of fossils it contained. Some fossils proved particularly useful in matching up rock layers from different regions. These fossils, called**index fossils**, are widespread but only existed for a relatively brief period of time. When a particular index fossil is found, the relative age of the bed is immediately known.

Many fossils may qualify as index fossils. Ammonites, trilobites, and graptolites are often used as index fossils, as are various **microfossils**, or fossils of microscopic organisms. Fossils of animals that drifted in the upper layers of the ocean are particularly useful as index fossils, as they may be distributed all over the world.

In contrast to index fossils, **living fossils** are organisms that have existed for a tremendously long period of time without changing very much at all. For example, the Lingulata brachiopods have existed from the Cambrian period to the present, a time span of over 500 million years! Modern specimens of Lingulata are almost indistinguishable from their fossil counterparts (Figure 11.8).




Figure 11.8: Fossil *Lingula* (left) and modern *Lingula* (right).

Clues from Fossils

Fossils are our best form of evidence about the history of life on Earth. In addition, fossils can give us clues about past climates, the motions of plates, and other major geological events.

The first clue that fossils can give is whether an environment was **marine** (underwater) or **terrestrial** (on land). Along with the rock characteristics, fossils can indicate whether the water was shallow or deep, and whether the rate of sedimentation was slow or rapid. The amount of wear and fragmentation of a fossil can allow scientists to estimate the amount of wave action or the frequency of storms.

Often fossils of marine organisms are found on or near tall mountains. For example, the Himalayas, the tallest mountains in the world, contain trilobites, brachiopods, and other marine fossils. This indicates that rocks on the seabed have been uplifted to form

huge mountains. In the case of the Himalayas, this happened when the Indian Subcontinent began to ram into Asia about 40 million years ago.

Fossils can also reveal clues about past climate. For example, fossils of plants and coal beds have been found in Antarctica. Although Antarctica is frozen today, in the past it must have been much warmer. This happened both because Earth's climate has changed and because Antarctica has not always been located at the South Pole.

One of the most fascinating patterns revealed by the fossil record is a number of **mass extinctions**, times when many species died off. Although the mass extinction that killed the dinosaurs is most famous, the largest mass extinction in Earth history occurred at the end of the Permian period, about 250 million years ago. In this catastrophe, it is estimated that over 95% of species on Earth went extinct! The cause of these mass extinctions is not definitely known, but most scientists believe that collisions with comets or asteroids were the cause of at least a few of these disasters.

Lesson Summary

- A fossil is any remains of ancient life. Fossils can be body fossils, which are remains of the organism itself or trace fossils, such as burrows, tracks, or other evidence of activity.
- Preservation as a fossil is a relatively rare process. The chances of becoming a fossil are enhanced by quick burial and the presence of preservable hard parts, such as bones or shells.
- Fossils form in five ways: preservation of original remains, permineralization, molds and casts, replacement, and compression.
- Rock formations with exceptional fossils are called very important for scientists to study. They allow us to see information about organisms that we may not otherwise ever know.
- Index fossils are fossils that are widespread but only existed for a short period of time. Index fossils help scientists to find the relative age of a rock layer and match it up with other rock layers.
- Living fossils are organisms that haven't changed much in millions of years and are still alive today.
- Fossils give clues about the history of life on Earth, environments, climate, movement of plates, and other events.

Review Questions

1. What factors make it more likely that an animal will be preserved as a fossil?
2. What are the five main processes of fossilization?
3. A scientist wants to determine the age of a rock. The rock contains an index fossil and an ancient relative of a living fossil. Which fossil will be more useful for dating the rock, and why?
4. The island of Spitzbergen is in the Arctic Ocean north of Norway, near the North Pole. Fossils of tropical fruits have been found in coal deposits in Spitzbergen. What does this indicate?

Vocabulary

amber

Fossilized tree sap.

body fossil

The remains of an ancient organism. Examples include shells, bones, teeth, and leaves.

cast

A structure that forms when sediments fill a mold and harden, forming a replica of the original structure.

fossil

Any remains or trace of an ancient organism.

fossil fuel

A fuel that was formed from the remains of ancient organisms. Examples include coal, oil, and natural gas.

fossilization

The process of becoming a fossil.

index fossil

A fossil that identifies and shows the relative age of the rocks in which it is found. Index fossils come from species that were widespread but existed for a relatively brief period of time.

living fossil

A modern species or genus that has existed on Earth for millions of years without changing very much.

marine

Of or belonging to the sea.

mass extinction

A period of time when an unusually high number of species became extinct.

microfossil

A fossil that must be studied with the aid of a microscope.

mold

An impression made in sediments by the hard parts of an organism.

permineralization

A type of fossilization in which minerals are deposited into the pores of the original hard parts of an organism.

terrestrial

Of or belonging to the land.

trace fossil

Evidence of the activity of an ancient organism. Examples include tracks, trails, burrows, tubes, boreholes, and bite marks.

Points to Consider

- What are some other examples of mythical creatures that may be based on fossils?
- Why is it so rare for an animal to be preserved as a fossil?
- Some organisms are more easily preserved than others. Why is this a problem for scientists who are studying ancient ecosystems?
- Why are examples of amazing fossil preservation so valuable for scientists?
- Many fossils of marine organisms have been found in the middle of continents, far from any ocean. What conclusion can you draw from this?

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Fossils. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.3: Fossils is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.4: History of Earth's Life Forms



Figure 12.15: Charles Doolittle Walcott.

In the summer of 1909, an American scientist named Charles Doolittle Walcott (Figure 12.15) was in the Rocky Mountains of British Columbia, Canada. He was a paleontologist, which is a scientist who studies past life on Earth. He was searching for fossils. Riding on horseback, he was making his way down a mountain trail when he noticed something on the ground. He stopped to pick it up. It was a fossil! He began to dig around the area and found even more fossils. The fossils that Walcott found were of some of the most bizarre organisms anyone had ever seen. One of the organisms preserved in the fossils had a soft body like a worm, five eyes, and a long nose like a vacuum cleaner hose (Figure 12.16). Most of the fossils were the remains of animals that do not live today. They are now extinct, which means that nothing of their kind lives and that they are gone forever.



Figure 12.16: This bizarre animal with five eyes lived during the Cambrian. Fossils of it were discovered by Charles Walcott.

The organisms in Walcott's fossils lived during a time of geologic history known as the Cambrian. The Cambrian period began about 540 million years ago. It marked the beginning of the Phanerozoic Eon. It also marked the beginning of many new and complex life forms appearing on Earth. In fact, the term Phanerozoic means "time of well-displayed life". We still live today in the Phanerozoic Eon. However, life on Earth is very different today than it was 540 million years ago. This lesson covers some of the history of life on Earth. It will show you how living things have developed and changed over the last 540 million years of the Phanerozoic Eon. You will learn about how species adapt and evolve over time.

Lesson Objectives

- Describe how adaptations develop.
- Explain how the fossil record shows us that species evolve over time.
- Describe the general development of Earth's life forms over the last 540 million years.

Evaluating Prior Knowledge

Be sure that you can answer the following questions before you begin this lesson.

- What is a fossil?

- How is geologic time divided?
- How do organisms depend on their environment to live?

Earth's Diversity



Figure 12.17: There is an amazing diversity of organisms on Earth.

There are over 1 million species of plants and animals known to be currently alive on Earth (Figure 12.17). Scientists believe there are millions more that have not been discovered yet. Look around you and you notice that the organisms on this planet have incredible **variation**. One of the most remarkable features of Earth's organisms is their ability to survive in their specific environments. For example, polar bears have thick fur coats that help them stay warm in the icy waters that they hunt in (Figure 12.18). Reindeer have sponge-like hoofs that help them walk on snowy ground without slipping and falling. Plants that live in dry desert environments have special stems and leaves that help them conserve water.



Figure 12.18: Many animals, like this polar bear, have special body features that help them live in a certain environment.

Other organisms have special features that help them hunt for food or avoid being the food of another organism. For example, when zebras in a herd run away from lions, the zebras' dark stripes confuse the lions and make it hard for them to focus on just one zebra during the chase. Hummingbirds have long thin beaks that help them drink nectar from flowers. Some plants have poisonous or foul-tasting substances in them that keep animals from eating them.

Adaptations and Evolution

The characteristics of an organism that help it survive in a given environment are called **adaptations**. Adaptations develop when certain variations in a population help some members survive better than others (Figure 12.19). Often the variation comes from a mutation, or a random change in an organism's genes. The ones that survive pass favorable traits on to their **offspring**.




Figure 12.19: An explanation of how adaptations develop.

To help you understand adaptation, think about a population of oak trees. Imagine that most of the trees are easily killed by a certain fungus but that every now and then, there is one tree that has a natural ability to survive the fungus. That one tree displays a variation that gives it a better chance of surviving its environment. It also has a better chance of living to produce seeds and have offspring. It will reproduce and carry on the species, while the other trees will die off. The tree with the natural ability to resist the fungus will pass that trait on to its offspring. The other trees will not live and have offspring. Eventually the population will change so that most of the individual trees have the trait to survive the fungus. This is an adaptation. Adaptations are inherited traits that an organism gets from its parents. Over time traits that help an organism survive become more common. Traits that hinder survival eventually disappear.

Changes and adaptations in a species accumulate over time. Eventually the descendants are very different from their ancestors and may become a whole new species. Changes in a species over time are called evolution. We learn about evolution from the fossil record. It shows us that many of the life forms that live today developed from earlier, different life forms. For example, horse fossils show us that about 60 million years ago horses were much smaller than they are today (Figure 12.20). Fossils also show us that horses' teeth and hooves have changed several times as horses have adapted to changes in the environment.



Figure 12.20: The horse has evolved over the last 60 million years. Horses today are much larger than earlier horses.

Studying the Fossil Record

Like the organisms that were represented in Walcott's fossils, many of the organisms that once lived on Earth are now extinct. Earth's overall environmental conditions have changed many times since the Cambrian, and many organisms did not have the traits to survive the changes. Those that did survive the changes passed traits on to their offspring. They gave rise to the species that live today.

We study fossils to learn about how species responded to change over the Earth's long history. Fossils show us that simple organisms dominated life on Earth for its first 3 billion years. Then, between 1 and 2 billion years ago, the first multi-cellular organisms appeared on Earth. Life forms gradually evolved and became more complex. During the Cambrian period, animals became more **diverse** and complex. We sometimes refer to this part of the Phanerozoic Eon as the Cambrian Explosion—meaning a time when the Earth "exploded" with incredible numbers of new complex life forms.

Phanerozoic Eon

The Phanerozoic Eon is divided into three chunks of time called eras—the Paleozoic, the Mesozoic, and the Cenozoic (Table (12.1). They span from about 540 million years ago to the present. We live now in the Cenozoic Era. The table below shows how life has changed during the long span of the Phanerozoic Eon. Notice that different types of organisms developed at different times. However, all organisms evolved from a common ancestor. Life gradually became more diverse and new species branched out from that common ancestor. Most modern organisms evolved from species that are now extinct. To get an idea of how an organism can change from a single common ancestor to many different types, think about all the different types of dogs. All dogs evolved from a common wolf ancestor. Today there are hundreds of varieties of dogs that all look very different.

Table 12.1: Development of Life During the Phanerozoic Eon

Era	Millions of Years Ago	Major Forms of Life
Cenozoic	0.2 (200,000 years ago)	First humans
	35	First grasses; grasslands begin to dominate the land
Mesozoic	130	First plants with flowers
	150	First birds on Earth
	200	First mammals on Earth
	251	Age of dinosaurs begins
Paleozoic	300	First reptiles on Earth
	360	First amphibians on Earth
	400	First insects on Earth
	475	First plants and fungi begin growing on land
	500	First fish on Earth

The eras of the Phanerozoic Eon are separated by events called mass extinctions. A mass extinction occurs when large numbers of organisms become extinct in a short amount of time. Between the Paleozoic and the Mesozoic, nearly 95% of all species on Earth died off. The cause or causes of this extinction are still being debated.

Between the Mesozoic and the Cenozoic, about 50% of all animal species on Earth died off. This mass extinction, 65 million years ago, is the one in which the dinosaurs became extinct. Although there are other hypotheses, most scientists think that this mass extinction took place when a giant meteorite struck Earth with the energy of the most powerful nuclear weapon. The impact kicked up a massive dust cloud. When the particles rained back onto the surface they heated the atmosphere until it became as hot as a kitchen oven, roasting animals. Dust that remained in the atmosphere blocked sunlight for a year or more, causing a deep freeze and ending photosynthesis. Sulfur from the impact mixed with water in the atmosphere to form acid rain, which dissolved the shells of the tiny marine plankton that form the base of the food chain. With little food being produced by land plants and plankton, animals starved. Carbon dioxide was also released from the impact and eventually caused global warming. Life forms could not survive the dramatic temperature swings.

Earth's climate changed numerous times during the Phanerozoic Eon. Just before the beginning of the Phanerozoic, much of the Earth was cold and covered with **glaciers** (Figure 12.21). As the Phanerozoic began, however, the climate was changing to a warm and **tropical** one (Figure 12.22). The glaciers were replaced with tropical seas. This allowed the Cambrian Explosion of many new life forms on Earth. During the Phanerozoic, Earth's climate has gone through at least 4 major cycles between times of cold glaciers and times of warm tropical seas. Some organisms survived environmental changes in the climate; others became extinct when the climate changed beyond their capacity to cope with it.



Figure 12.21: Just before the Phanerozoic, many parts of Earth were covered with glaciers. After the Earth began to warm and many of the glaciers melted, there was an explosion of new life on Earth. The glaciers in this picture are from the present. However, glaciers are much less common on Earth today than at other times in Earth's history.



Figure 12.22: The Phanerozoic Eon was often characterized by times of warm tropical climates. The age of the dinosaurs was especially mild for most of the Earth. This allowed plants and animals to spread over large areas of land. This picture shows plants in a modern rainforest. Plants of the Phanerozoic may have looked similar.

Lesson Summary

- Adaptations are favorable traits that organisms inherit. Adaptations develop from variations within a population and help organisms to survive in their given environment.
- Changes in populations accumulate over time; this is called evolution.
- The fossil record shows us that present day life forms evolved from earlier different life forms. It shows us that the first organisms on Earth were simple bacteria that dominated the Earth for several billion years.
- Beginning about 540 million years ago more complex organisms developed on Earth. During the Phanerozoic Eon all of the plant and animal types we know today have evolved.
- Many types of organisms that once lived are now extinct. Eart's overall environment, especially the climate, has changed many times, and organisms change too over time.

Review Questions

1. Describe what is meant by adaptation.
2. The first animals on Earth had soft bodies. Gradually many animal species evolved that had hard outer parts called exoskeletons covering their bodies. How might an exoskeleton be a favorable adaptation?
3. Explain why unfavorable traits do not usually get passed to offspring.
4. List the order in which the major types of animals appeared on Earth.
5. How might climate have affected the ability of plants to grow over large areas during a given time?
6. One cause of mass extinctions is meteorite or comet impacts. What might be some additional causes of mass extinctions?

Vocabulary

adaptation

A trait that an organism inherits that helps it survive in its natural environment.

evolution

The change in an organism's traits over time such that a new species is often the result.

glaciers

Large sheets of flowing ice.

paleontologist

A scientist who studies Earth's past life forms.

tropical

A climate that is warm and humid.

variation

Having many differences.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/History_of_Earth%27s_Life_Forms. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.4: History of Earth's Life Forms is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.5: Early Earth

Imagine that you had a movie that shows the history of Earth from its beginning to the present day—as if a giant camera in space had recorded pictures of Earth over the last $4\,{}^{1}\!/_{2}$ billion years. How do you think the Earth would look in that movie at different times in history? How do you think it has changed?

If you put the movie in fast-forward, you would see lots of action and lots of change! You would see that our planet has undergone remarkable changes over billions of years (Figure 12.3). Huge mountains have formed, been destroyed, and replaced with new mountains. The oceans have opened up and moved around the globe. The continents have moved around, split apart from each other, and collided with each other, until finally reaching their present locations. Life on Earth has also changed tremendously. At first, the Earth was not even able to support life. There was no oxygen in the atmosphere, and Earth's surface was extremely hot. Slowly, over millions of years, the Earth changed so that plants and animals could begin to grow. Living things then changed the Earth even more.



Figure 12.3: The Earth from space. The Earth looks very different today than it did when it first formed over 4 billion years ago.

We often enjoy using our imagination to think about what the Earth was like when dinosaurs roamed around (Figure 12.4). What images come to your mind when you think about the dinosaurs? Now imagine a time on Earth before even the dinosaurs. Imagine the time before any living thing was on Earth. What images come to mind now? How do you think the Earth looked when it was first formed? This lesson will help you understand how the Earth formed, what it looked like during its earliest years, and how life first developed on Earth.



Figure 12.4: The Earth and its dominant life forms have changed throughout the Earth's long history.

Evaluating Prior Knowledge

The following questions are addressed in other chapters and will help you work through this lesson. Research these before you move on.

- What are chemical elements?
- What conditions do plants and animals require to live?
- What is the atmosphere and what is it made of?
- How do weathering and erosion affect the Earth?

Formation of Earth and Our Solar System

We can construct the formation history of our solar system by looking at regions where other stars are forming now. Star formation begins when a giant cloud of gas and dust collapses under its own gravity. As the cloud contracts, it begins to spin faster and settles into a disk-shaped structure. We see these disc-shaped objects (called proplyds) in the Orion Nebula (Figure 12.13), where the new stars are forming today. Most of the dusty disk material drains toward the center where the density gradually increases until the enormous central pressure triggers nuclear fusion reactions and the star is born.




Figure 12.5: Orion Nebula.

However, a relatively small fraction of the disk material is left behind in the form of ice-coated dust grains. The icy mantles of the grains begin sticking together and eventually grow to meter-sized rocky boulders called planetesimals. The planetesimals collide and accrete into larger bodies that are tens of kilometers in diameter called protoplanets. Once the protoplanets clear a gap in the disk, they become bonafide planets and their orbits begin to stabilize (Figure 12.6).




Figure 12.6: An artist’s rendition of a baby star still surrounded by a protoplanetary disc in which planets are forming.

The process of planet formation is messy. Not all of the planetesimals are accreted into planets. Millions of planetesimals remain as the leftover debris and are now the asteroids and ice-coated comets in our solar system. In the first hundred million years after the formation of the Sun, collisions between the leftover planetesimals and the planets were common. We see evidence for heavy bombardment by planetesimals on the surfaces of the moon and Mercury (Figure 12.7 and Figure 12.8).



Figure 12.7: The surface of the moon is scarred by collisions with debris that was meters to kilometers in diameter. Most of the planetesimals were accreted into planets or moons, but some of these objects remain as meteors, asteroids, and comets in our solar system today.




Figure 12.8: The surface of Mercury shows similar collisional cratering. Most of the planetesimals were accreted into planets or moons, but some of these objects remain as meteors, asteroids, and comets in our solar system today.

The same types of collisions would have occurred on the surface of the Earth, however erosive processes have erased all except the most recent of these collisions. Pictured in Figure 12.9 is a Meteor Crater in Arizona.



Figure 12.9: Meteor crater in Arizona was formed about 40,000 years ago by the impact of a meteorite that was about 50 meters in diameter. Such collisions are rare today.

About 100 million years after the formation of the Sun, the gravity of the planets and moons in our solar system had swept up most of the planetesimals. However, millions of these objects still remain in gravitationally stable orbits in the main asteroid belt of the solar system, in the Trojan asteroid belt, or out beyond Neptune and Pluto in the Kuiper belt. Illustrated in the sketch below is the location of the largest reservoir of asteroids in our solar system today (Figure 12.10).




Figure 12.10: This sketch shows the largest reservoir of asteroids in our solar system today.

Earth is the only object in our solar system known to support life (Figure 12.11). Today there are over 1 million known species of plants and animals on Earth.




Figure 12.11: The Earth formed at the same time as the other planets in our solar system about 4 $\frac{1}{2}$ billion years ago.

The materials that came together to form the Earth were made of several different chemical elements. Each element has a different **density**, defined as mass per volume. Density describes how heavy an object is compared to how much space the object takes up. After Earth's early formation, the denser elements sank to the center. The lighter elements rose to the surface. You have probably seen something like this happen if you have ever mixed oil and water in a bottle. The water is denser than oil. If you put both in a bottle, shake it up, and then let it sit for a while, the water settles to the bottom and the oil rises up over the top of the water.

Today, the Earth consists of layers that represent different densities (Figure 12.12). Earth's center is called its core. The core is made of very dense metal elements called iron and nickel. The outermost layer of the Earth is its crust. The crust is made mostly of light elements such as silicon, oxygen, and aluminum. More information on the different layers of the Earth is presented in the lesson on plate tectonics.




Figure 12.12: The Earth is made of several layers that vary in density. The center of the Earth is the core, which is the densest. The outermost layer is the crust, which is the least dense. The middle layers make up the mantle.

Formation of the Earth's Atmosphere

The early Earth was very different from our Earth today. The early Earth experienced frequent impacts from asteroids and meteorites and had much more frequent volcanic eruptions. There was no life on Earth for the first billion years because the **atmosphere** was not suitable for life. Earth's first atmosphere had lots of **water vapor** but had almost no oxygen. Later, frequent volcanic eruptions put several different gases into the air (Figure 12.13). These gases created a new type of atmosphere for Earth. The volcanic eruptions spewed gases such as nitrogen, carbon dioxide, hydrogen, and water vapor into the atmosphere—but no free oxygen. Without oxygen, there was still very little that could live on Earth.



Figure 12.13: Volcanic eruptions occurred almost constantly on the early Earth. Eruptions put water vapor, carbon dioxide and other gases into the air that helped create Earth's early atmosphere.

Slowly, two processes changed Earth's atmosphere to one that is more oxygen-rich—like the one we have today. First, radiation from the Sun caused water vapor **molecules** to split apart. Remember that a molecule of water is made of the elements hydrogen and oxygen, or H2O. Radiation from the Sun split some of the water molecules into hydrogen and oxygen. The hydrogen escaped back to outer space. The oxygen accumulated in the atmosphere. The second process that changed Earth's early atmosphere was photosynthesis (Figure 12.14). About 2.4 billion years ago, a type of organism called cyanobacteria evolved on the early Earth and began carrying out photosynthesis. Photosynthesis uses carbon dioxide and energy from the Sun to produce sugar and oxygen. The cyanobacteria were very simple organisms but performed an important role in changing Earth's early atmosphere. They carried out photosynthesis to produce the materials they needed to grow. They gave off oxygen to the atmosphere as they did this.




Figure 12.14: Bacteria capable of photosynthesis first appeared on Earth about 2.4 billion years ago. Photosynthesis takes sunlight, carbon dioxide, and water and produces sugar and oxygen. Photosynthesis contributed oxygen to Earth's early atmosphere and helped change it from one rich in carbon dioxide to one rich in oxygen.

Oxygen in the atmosphere is important for life for two main reasons. First, oxygen makes up the ozone layer. The ozone layer is in the upper part of the atmosphere, and is made of O_3 molecules—a particular type of oxygen molecule. It blocks harmful **radiation** from the sun and keeps it from reaching Earth's surface. Without an ozone layer, intense radiation from the sun reached the early Earth's surface, making life almost impossible. Secondly, oxygen in the atmosphere is necessary for animals, including humans, to breathe. No animals would have been able to breathe in Earth's early atmosphere. However, there were probably several types of bacteria that lived on Earth during this early time. They would have been anaerobic, meaning that they did not need oxygen to live.

Very simple cells lived on Earth for the first few billion years of Earth's history. Some of the oldest fossils of more complex organisms are from about 2 billion years ago. They are found in Australia.

Besides changes in life and the atmosphere, other changes have also happened since the Earth was first formed. Early volcanic eruptions on Earth released large amounts of water vapor into the atmosphere. The water vapor slowly **condensed** and returned to Earth's surface in rainfall. This formed the oceans. Water began to cycle on Earth, and events like rainfall and storms next began to change the Earth's surface through weathering and erosion. The Earth's Fresh Water chapter gives more detail on how water cycles on Earth.

The continents were in very different locations than they are now. Scientists do not know how Earth's land looked exactly after the planet's first formation. They do know that North America and Greenland formed one giant landmass called Laurentia about 1.8 billion years ago. By about 1 billion years ago, Antarctica may have been close to the equator, even though it now sits at Earth's South Pole. Today, Earth's continents continue to slowly shift around the globe.

Lesson Summary

- The Earth formed more than 4 billion years ago along with the other planets in our solar system.
- The early Earth had no ozone layer and was probably very hot. The early Earth also had no free oxygen.
- Without an oxygen atmosphere very few things could live on the early Earth. Anaerobic bacteria were probably the first living things on Earth.
- The early Earth had no oceans and was frequently hit with meteorites and asteroids. There were also frequent volcanic eruptions. Volcanic eruptions released water vapor that eventually cooled to form the oceans.

- The atmosphere slowly became more oxygen-rich as solar radiation split water molecules and cyanobacteria began the process of photosynthesis. Eventually the atmosphere became like it is today and rich in oxygen.
- The first complex organisms on Earth first developed about 2 billion years ago.

Review Questions

1. Describe how the different layers of the Earth vary by density. When did the materials that make the Earth separate out by density?
2. Explain two reasons why having an oxygen-rich atmosphere is important for life on Earth.
3. Scientists believe that Earth's ozone layer is shrinking because of human activities and air pollution. What affect might this have on Earth's life forms?
4. Describe the role of cyanobacteria in changing Earth's early atmosphere.
5. List three ways the Earth was different today from when it was first formed.
6. Suppose that the Earth had been much cooler when it first formed. How would the Earth's interior be different than it is today?

Vocabulary

atmosphere
: The mixture of gases that surrounds the Earth and contains the air we breathe.

condensed
: Cooled and changed from water vapor to liquid water.

density
: Density means mass per unit area.

molecules
: The smallest possible amounts of a chemical substance.

radiation
: Energy given off by the Sun.

species
: A group of living things that have similar characteristics.

water
: Vapor water in a gas form.

Points to Consider

- How did life on Earth develop from simple bacteria to more complex organisms?
- When did complex organisms like fish, reptiles, and mammals appear on Earth?
- When did the major features of the Earth that we know today first form?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Early_Earth. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.5: Early Earth is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

10.6: Geologic Time Scale

How many years is a “long time”? We often express time in hours or days, and 10 or 20 years certainly feels like a long time. Imagine if you needed to think about one million, 100 million, or even several billion years. These exceptional lengths of time seem unbelievable, but they are exactly the spans of times that scientists use to describe the Earth.

The Earth is 4 $\frac{1}{2}$ billion years old. That’s 4,500,000,000 years! Have places like the Grand Canyon and the Mississippi River been around for all of those years, or were they formed more recently? When did the giant Rocky Mountains form and when did dinosaurs walk the Earth? To answer these questions, you have to think about times that were millions or billions of years ago.

Historical geologists are scientists who study the Earth’s past. They study clues left on the Earth to learn two main things: the *order* in which events happened on Earth, and *how long* it took for those events to happen. For example, they have learned that the Mississippi River formed many millions of years after the Grand Canyon began forming. They have also concluded that dinosaurs lived on the Earth for about 200 million years.

Scientists have put together the **geologic time scale** to describe the order and duration of major events on Earth for the last 4 $\frac{1}{2}$ billion years. Some examples of events listed on the geologic time scale include the first appearance of plant life on Earth, the first appearance of animals on Earth, the formation of Earth’s mountains, and the **extinction** of the dinosaurs.

You will learn about some of the scientific principles that historical geologists use to describe Earth’s past. You will also learn some of the clues that scientists use to learn about the past and shows you what the geologic time scale looks like.

Evaluating Prior Knowledge

Before you work through this lesson, think about the following questions. Be sure that you can answer each one. They will help you better understand this lesson.

- What is a fossil and how does a fossil form?
- How does a sedimentary rock form?
- In what types of locations do sedimentary rocks form?
- How do you determine the relative and absolute ages of rock layers?

Geologic Time

The first principle you need to understand about geologic time is that the laws of nature are always the same. This means that the laws describing how things work are the same today as they were billions of years ago. For example, water freezes at 0°C. This law has always been true and always will be true. Knowing the natural laws helps you think about Earth’s past, because it gives you clues about how things happened very long ago. It means that we can use present-day processes to interpret the past. Imagine you find **fossils** of sea animals in a rock. The laws of nature say that sea animals must live in the sea. That law has never changed, so the rock must have formed near the sea. The rock may be millions of years old, but the fossils in it are a clue for us today about how it formed.

Now imagine that you find that same rock with fossils of a sea animal in a place that is very dry and nowhere near the sea. How could that be? Remember that the laws of nature never change. Therefore, the fossil means that the rock definitely formed by the sea. This tells you that even though the area is now dry, it must have once been underwater. Clues like this have helped scientists learn that Earth’s surface features have changed many times. Spots that were once covered by warm seas may now be cool and dry. Places that now have tall mountains may have once been low, flat ground. These kinds of changes take place over many millions of years, but they are still slowly going on today. The place where you live right now may look very different in the far future.

Relative and Absolute Age Dating of Rocks

The clues in rocks help scientists put together a picture of how places on Earth have changed. Scientists noticed in the 1700s and 1800s that similar layers of sedimentary rocks all over the world contain similar fossils. They used **relative dating** to order the rock layers from oldest to youngest. In the process of relative dating, scientists do not determine the exact age of a fossil but do learn which ones are older or younger than others. They saw that the fossils in older rocks are different from the fossils in younger rocks. For example, older rock layers contain only reptile fossils, but younger rock layers may also contain mammal fossils.

Scientists divided Earth's history into several chunks of time when the fossils showed similar things living on the Earth. They gave each chunk of time a name to help them keep track of how Earth has changed. For example, one chunk of time when many dinosaurs lived is called the Jurassic. We find fossils of Earth's first green plants from the chunk of time named the Ordovician. Many of the scientists who first assigned names to times in Earth's history were from Europe. As a result, many of the names they used came from towns or other local places where they studied in Europe.

Ordering rock layers from oldest to youngest was a first step in creating the geologic time scale. It showed the order in which life on Earth changed. It also showed us how certain areas changed over time in regard to climate or type of environment. However, the early geologic time scale only showed the order of events. It did not show the actual years that events happened. With the discovery of radioactivity in the late 1800s, scientists were able to measure the exact age in years of different rocks. Measuring the amounts of radioactive elements in rocks let scientists use**absolute dating** to give ages to each chunk of time on the geologic time scale. For example, they are now able to state that the Jurassic began about 200 million years ago and that it lasted for about 55 million years.

Geologic Time Scale

Today, the geologic time scale is divided into major chunks of time called **eons**. Eons may be further divided into smaller chunks called eras, and each era is divided into **periods**. Figure 12.1 shows you what the geologic time scale looks like. We now live in the Phanerozoic eon, the Cenozoic era, and the Quarternary period. Sometimes, periods are further divided into epochs, but they are usually just named "early" or "late", for example, "late Jurassic", or "early Cretaceous". Note that chunks of geologic time are not divided into equal numbers of years. Instead, they are divided into blocks of time when the fossil record shows that there were similar organisms on Earth.

EON	ERA	PERIOD	MILLIONS OF YEARS AGO
Phanerozoic	Cenozoic	Quaternary	1.6
		Tertiary	66
	Mesozoic	Cretaceous	138
		Jurassic	205
		Triassic	240
	Paleozoic	Permian	290
		Pennsylvanian	330
		Mississippian	360
		Devonian	410
		Silurian	435
		Ordovician	500
		Cambrian	570
Proterozoic	Late Proterozoic Middle Proterozoic Early Proterozoic		2500
Archean	Late Archean Middle Archean Early Archean		3800?
Pre-Archean			

Figure 12.1: The geologic time scale.

One of the first scientists to understand geologic time was James Hutton. In the late 1700s, he traveled around Great Britain and studied sedimentary rocks and their fossils. He believed that the same processes that work on Earth today formed the rocks and fossils from the past. He knew that these processes take a very long time, so the rocks must have formed over millions of years. Before Hutton, most people believed the Earth was only several thousand years old. His work helped us understand that the laws of nature never change and that the Earth is very old. He is sometimes called the "father of geology".

The geologic time scale is often shown with illustrations of how life on Earth has changed. It sometimes includes major events on Earth, too, such as the formation of the major mountains or the extinction of the dinosaurs. Figure 12.2 shows you a different way of looking at the geologic time scale. It shows how Earth's environment and life forms have changed.




Figure 12.2: A different way of looking at the geologic time scale.

Lesson Summary

- The Earth is very old, and the study of Earth's past requires us to think about times that were millions or even billions of years ago. Scientists use the geologic time scale to illustrate the order in which events on Earth have happened.
- The geologic time scale was developed after scientists observed changes in the fossils going from oldest to youngest sedimentary rocks. They used relative dating to divide Earth's past in several chunks of time when similar organisms were on Earth.
- Later, scientists used absolute dating to determine the actual number of years ago that events happened. The geologic time scale is divided into eons, eras, periods, and epochs.

Review Questions

1. How old is the Earth?
2. Why did early geologic time scales not include the number of years ago that events happened?
3. Dinosaurs went extinct about 66 million years ago. Which period of geologic time was the last in which dinosaurs lived?
4. Can scientists use the same principles they use to study Earth's history to also study the history of other planets?
5. Suppose you are hiking in the mountains of Utah and find a fossil of an animal that lived on the ocean floor. You learn that the rock that holds the fossil is from the Mississippian period. What was the environment like during the Mississippian in Utah?
6. Why are sedimentary rocks more useful than metamorphic or igneous rocks in establishing the relative ages of rock?
7. Which is likely to be more frequently found in rocks: fossils of very old sea creatures or very old land creatures?

Vocabulary

absolute dating
: Methods used to determine how long ago something happened.

extinction
: When an organism completely dies out.

fossils
: The remains of past life, such as bones, shells, or other hard parts; may also include evidence of past life such as footprints or leaf impressions.

geologic time scale
: A timeline that illustrates Earth's past.

relative dating methods
: Used to determine the order of geologic events in Earth's history.

Points to Consider

- How did life on Earth change from one period of geologic time to the next?
- When did life first appear on Earth?
- What conditions were necessary on Earth for living things to survive?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/Geologic_Time_Scale. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 10.6: Geologic Time Scale is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

11: Ocean Water and Life

11.1: Thermohaline circulation

11.2: Thermocline

11.3: Tsunamis


11.4: Ocean Organisms

This page titled 11: Ocean Water and Life is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.


11.1: Thermohaline circulation

Thermohaline circulation (**THC**) is a part of the large-scale ocean circulation that is driven by global density gradients created by surface heat and freshwater fluxes.[1][2] The adjective thermohaline derives from *thermo-* referring to temperature and *-haline* referring to salt content, factors which together determine the density of sea water. Wind-driven surface currents (such as the Gulf Stream) travel polewards from the equatorial Atlantic Ocean, cooling en route, and eventually sinking at high latitudes (forming North Atlantic Deep Water). This dense water then flows into the ocean basins. While the bulk of it upwells in the Southern Ocean, the oldest waters (with a transit time of around 1000 years)[3] upwell in the North Pacific.[4] Extensive mixing therefore takes place between the ocean basins, reducing differences between them and making the Earth's oceans a global system. On their journey, the water masses transport both energy (in the form of heat) and matter (solids, dissolved substances and gases) around the globe. As such, the state of the circulation has a large impact on the climate of the Earth.


The thermohaline circulation is sometimes called the **ocean conveyor belt**, the **great ocean conveyor**, or the **global conveyor belt**. On occasion, it is used to refer to the **meridional overturning circulation** (often abbreviated as **MOC**). The term MOC, indeed, is more accurate and well defined, as it is difficult to separate the part of the circulation which is actually driven by temperature and salinity alone as opposed to other factors such as the wind and tidal forces.[5] Moreover, temperature and salinity gradients can also lead to circulation effects that are not included in the MOC itself.



A summary of the path of the thermohaline circulation. Blue paths represent deep-water currents, while red paths represent surface currents.



Thermohaline circulation



Global map of average Sea Surface Density

Overview



The global conveyor belt on a continuous-ocean map

The movement of surface currents pushed by the wind is fairly intuitive. For example, the wind easily produces ripples on the surface of a pond. Thus the deep ocean — devoid of wind — was assumed to be perfectly static by early oceanographers. However, modern instrumentation shows that current velocities in deep water masses can be significant (although much less than surface speeds).

In the deep ocean, the predominant driving force is differences in density, caused by salinity and temperature variations (increasing salinity and lowering the temperature of a fluid both increase its density). There is often confusion over the components of the circulation that are wind and density driven.[6][7] Note that ocean currents due to tides are also significant in many places; most prominent in relatively shallow coastal areas, tidal currents can also be significant in the deep ocean.

The density of ocean water is not globally homogeneous, but varies significantly and discretely. Sharply defined boundaries exist between water masses which form at the surface, and subsequently maintain their own identity within the ocean. They position themselves one above or below each other according to their density, which depends on both temperature and salinity.

Warm seawater expands and is thus less dense than cooler seawater. Saltier water is denser than fresher water because the dissolved salts fill interstices between water molecules, resulting in more mass per unit volume. Lighter water masses float over denser ones (just as a piece of wood or ice will float on water, see buoyancy). This is known as “stable stratification”. When dense water masses are first formed, they are not stably stratified. In order to take up their most stable positions, water masses of different densities must flow, providing a driving force for deep currents.

The thermohaline circulation is mainly triggered by the formation of deep water masses in the North Atlantic and the Southern Ocean caused by differences in temperature and salinity of the water.

The great quantities of dense water sinking at polar ocean basin edges must be offset by equal quantities of water rising elsewhere. Note that cold water in polar zones sink relatively rapidly over a small area, while warm water in temperate and tropical zones rise more gradually across a much larger area. It then slowly returns poleward near the surface to repeat the cycle. The continual diffuse upwelling of deep water maintains the existence of the permanent thermocline found everywhere at low and mid-latitudes. This slow upward movement is estimated to be about 1 centimeter (0.5 inch) per day over most of the ocean. If this rise were to stop, downward movement of heat would cause the thermocline to descend and would reduce its steepness.

Formation of deep water masses

The dense water masses that sink into the deep basins are formed in quite specific areas of the North Atlantic and the Southern Ocean. In the North Atlantic, seawater at the surface of the ocean is intensely cooled by the wind. Wind moving over the water also produces a great deal of evaporation, leading to a decrease in temperature, called evaporative cooling. Evaporation removes only water molecules, resulting in an increase in the salinity of the seawater left behind, and thus an increase in the density of the water mass. In the Norwegian Sea evaporative cooling is predominant, and the sinking water mass, the North Atlantic Deep Water (NADW), fills the basin and spills southwards through crevasses in the submarine sills that connect Greenland, Iceland and Great Britain. It then flows very slowly into the deep abyssal plains of the Atlantic, always in a southerly direction. Flow from the Arctic Ocean Basin into the Pacific, however, is blocked by the narrow shallows of the Bering Strait.

In the Southern Ocean, strong katabatic winds blowing from the Antarctic continent onto the ice shelves will blow the newly formed sea ice away, opening polynyas along the coast. The ocean, no longer protected by sea ice, suffers a brutal and strong

cooling. Meanwhile, sea ice starts reforming, so the surface waters also get saltier, hence very dense. In fact, the formation of sea ice contributes to an increase in surface seawater salinity; saltier brine is left behind as the sea ice forms around it (pure water preferentially being frozen). Increasing salinity lowers the freezing point of seawater, so cold liquid brine is formed in inclusions within a honeycomb of ice. The brine progressively melts the ice just beneath it, eventually dripping out of the ice matrix and sinking. This process is known as **brine rejection**.

The resulting Antarctic Bottom Water (AABW) sinks and flows north and east, but is so dense it actually underflows the NADW. AABW formed in the Weddell Sea will mainly fill the Atlantic and Indian Basins, whereas the AABW formed in the Ross Sea will flow towards the Pacific Ocean.

The dense water masses formed by these processes flow downhill at the bottom of the ocean, like a stream within the surrounding less dense fluid, and fill up the basins of the polar seas. Just as river valleys direct streams and rivers on the continents, the bottom topography steers the deep and bottom water masses.

Note that, unlike fresh water, seawater does not have a density maximum at 4°C but gets denser as it cools all the way to its freezing point of approximately −1.8°C.

Movement of thermohaline circulation

Formation and movement of the deep water masses at the North Atlantic Ocean, creates sinking water masses that fill the basin and flows very slowly into the deep abyssal plains of the Atlantic. This high latitude cooling and the low latitude heating drives the movement of the deep water in a polar southward flow. The deep water flows through the Antarctic Ocean Basin around South Africa where it is split into two routes: one into the Indian Ocean and one past Australia into the Pacific.

At the Indian Ocean, some of the cold and salty water from the Atlantic — drawn by the flow of warmer and fresher upper ocean water from the tropical Pacific — causes a vertical exchange of dense, sinking water with lighter water above. It is known as **overturning**. In the Pacific Ocean, the rest of the cold and salty water from the Atlantic undergoes haline forcing, and becomes warmer and fresher more quickly.

The out-flowing undersea of cold and salty water makes the sea level of the Atlantic slightly lower than the Pacific and salinity or halinity of water at the Atlantic higher than the Pacific. This generates a large but slow flow of warmer and fresher upper ocean water from the tropical Pacific to the Indian Ocean through the Indonesian Archipelago to replace the cold and salty Antarctic Bottom Water. This is also known as ‘haline forcing’ (net high latitude freshwater gain and low latitude evaporation). This warmer, fresher water from the Pacific flows up through the South Atlantic to Greenland, where it cools off and undergoes evaporative cooling and sinks to the ocean floor, providing a continuous thermohaline circulation.[8]

Hence, a recent and popular name for the thermohaline circulation, emphasizing the vertical nature and pole-to-pole character of this kind of ocean circulation, is the **meridional overturning circulation**.

Quantitative estimation

The deep water masses that participate in the MOC have chemical, temperature and isotopic ratio signatures and can be traced, their flow rate calculated, and their age determined. These include ^{231}Pa / ^{230}Th ratios.

Gulf Stream



Benjamin Franklin’s map of the Gulf Stream

Main article: Gulf Stream

The Gulf Stream, together with its northern extension towards Europe, the North Atlantic Drift, is a powerful, warm, and swift Atlantic ocean current that originates at the tip of Florida, and follows the eastern coastlines of the United States and Newfoundland before crossing the Atlantic Ocean. The process of western intensification causes the Gulf Stream to be a northward accelerating current off the east coast of North America.[9] At about 40°0′N 30°0′W / 40.000°N 30.000°W, it splits in two, with the northern stream crossing to northern Europe and the southern stream recirculating off West Africa. The Gulf Stream influences the climate of the east coast of North America from Florida to Newfoundland, and the west coast of Europe. Although there has been recent debate, there is consensus that the climate of Western Europe and Northern Europe is warmer than it would otherwise be due to the North Atlantic drift,[10][11] one of the branches from the tail of the Gulf Stream. It is part of the North Atlantic Gyre. Its presence has led to the development of strong cyclones of all types, both within the atmosphere and within the ocean. The Gulf Stream is also a significant potential source of renewable power generation.[12][13]

Upwelling

Main article: Upwelling

All these dense water masses sinking into the ocean basins displace the older deep water masses which were made less dense by ocean mixing. To maintain a balance water must be rising elsewhere. However, because this thermohaline upwelling is so widespread and diffuse, its speeds are very slow even compared to the movement of the bottom water masses. It is therefore difficult to measure where upwelling occurs using current speeds, given all the other wind-driven processes going on in the surface ocean. Deep waters have their own chemical signature, formed from the breakdown of particulate matter falling into them over the course of their long journey at depth. A number of scientists have tried to use these tracers to infer where the upwelling occurs.

Wallace Broecker, using box models, has asserted that the bulk of deep upwelling occurs in the North Pacific, using as evidence the high values of silicon found in these waters. Other investigators have not found such clear evidence. Computer models of ocean circulation increasingly place most of the deep upwelling in the Southern Ocean,[14] associated with the strong winds in the open latitudes between South America and Antarctica. While this picture is consistent with the global observational synthesis of William Schmitz at Woods Hole and with low observed values of diffusion, not all observational syntheses agree. Recent papers by Lynne Talley at the Scripps Institution of Oceanography and Bernadette Sloyan and Stephen Rintoul in Australia suggest that a significant amount of dense deep water must be transformed to light water somewhere north of the Southern Ocean.

Effects on global climate

The thermohaline circulation plays an important role in supplying heat to the polar regions, and thus in regulating the amount of sea ice in these regions, although poleward heat transport outside the tropics is considerably larger in the atmosphere than in the ocean.[15] Changes in the thermohaline circulation are thought to have significant impacts on the Earth's radiation budget. Insofar as the thermohaline circulation governs the rate at which deep waters are exposed to the surface, it may also play an important role in determining the concentration of carbon dioxide in the atmosphere. While it is often stated that the thermohaline circulation is the primary reason that Western Europe is so temperate, it has been suggested that this is largely incorrect, and that Europe is warm mostly because it lies downwind of an ocean basin, and because of the effect of atmospheric waves bringing warm air north from the subtropics.[16] However, the underlying assumptions of this particular analysis have likewise been challenged.[17]

Large influxes of low-density meltwater from Lake Agassiz and deglaciation in North America are thought to have led to a disruption of deep water formation and subsidence in the extreme North Atlantic and caused the climate period in Europe known as the Younger Dryas.[18]

In 2005, British researchers noticed that the net flow of the northern Gulf Stream had decreased by about 30% since 1957. Coincidentally, scientists at Woods Hole had been measuring the freshening of the North Atlantic as Earth becomes warmer. Their findings suggested that precipitation increases in the high northern latitudes, and polar ice melts as a consequence. By flooding the northern seas with lots of extra fresh water, global warming could, in theory, divert the Gulf Stream waters that usually flow northward past England and Norway, and cause them to instead circulate toward the equator. If this were to happen, Europe's climate would be seriously impacted.

References

1. ^ Rahmstorf, S (2003). "The concept of the thermohaline circulation". *Nature* **421** (6924): 699. Bibcode:2003Natur.421..699R. doi:10.1038/421699a. PMID 12610602.
2. ^ Lappo, SS (1984). "On reason of the northward heat advection across the Equator in the Atlantic ocean". *Study of Ocean and Atmosphere Interaction Processes* (Moscow Department of Gidrometeoizdat (in Russian)): 125–9.
3. ^ The global ocean conveyor belt is a constantly moving system of deep-ocean circulation driven by temperature and salinity;http://oceanservice.noaa.gov/facts/conveyor.html
4. ^ Primeau, F (2005). "Characterizing transport between the surface mixed layer and the ocean interior with a forward and adjoint global ocean transport model". *Journal of Physical Oceanography* **35** (4): 545–64. Bibcode:2005JPO….35..545P. doi:10.1175/JPO2699.1.
5. ^ Wunsch, C (2002). "What is the thermohaline circulation?".*Science* **298** (5596): 1179–81. doi:10.1126/science.1079329. PMID 12424356.
6. ^ Wyrtki, K (1961). "The thermohaline circulation in relation to the general circulation in the oceans". *Deep Sea Research* **8** (1): 39–64. Bibcode:1961DSR…..8…39W. doi:10.1016/0146-6313(61)90014-4.
7. ^ Schmidt, G., 2005, Gulf Stream slowdown?, RealClimate
8. ^ United Nations Environment Programme / GRID-Arendal, 2006, [1]. *Potential Impact of Climate Change*
9. ^ National Environmental Satellite, Data, and Information Service (2009). Investigating the Gulf Stream. North Carolina State University. Retrieved 2009-05-06.
10. ^ Hennessy (1858). *Report of the Annual Meeting: On the Influence of the Gulf-stream on the Climate of Ireland*. Richard Taylor and William Francis. Retrieved 2009-01-06.
11. ^ "Satellites Record Weakening North Atlantic Current Impact". NASA. Retrieved 2008-09-10.
12. ^ The Institute for Environmental Research & Eductation.Tidal.pdf Retrieved on 2010-07-28.
13. ^ Jeremy Elton Jacquot. Gulf Stream's Tidal Energy Could Provide Up to a Third of Florida's Power. Retrieved 2008-09-21.
14. ^ Marshall, John; Speer, Kevin (26 February 2012). "Closure of the meridional overturning circulation through Southern Ocean upwelling". *Nature Geoscience* **5** (3): 171–180. doi:10.1038/ngeo1391.
15. ^ Trenberth, K; Caron, J (2001). "Estimates of Meridional Atmosphere and Ocean Heat Transports". *Journal of Climate***14** (16): 3433–43. Bibcode:2001JCli…14.3433T. doi:10.1175/1520-0442(2001)014<3433:EOMAAO>2.0.CO;2.
16. ^ Seager,R., 2006, The Source of Europe's Mild Climate,*American Scientist*
17. ^ Rhines and Hakkinen, 2003, Is the Oceanic Heat Transport in the North Atlantic Irrelevant to the Climate in Europe? *ASOF Newsletter*
18. ^ Broecker, WSyear=2006 (2006). "Was the Younger Dryas Triggered by a Flood?". *Science* **312** (5777): 1146–8. doi:10.1126/science.1123253. PMID 16728622.

Other Sources

- Apel, JR (1987). *Principles of Ocean Physics*. Academic Press. ISBN 0-12-058866-8.
- Gnanadesikan, A., R. D. Slater, P. S. Swathi, and G. K. Vallis (2005). "The energetics of ocean heat transport". *Journal of Climate* **18**(14): 2604–16. Bibcode:2005JCli…18.2604G. doi:10.1175/JCLI3436.1.
- Knauss, JA (1996). *Introduction to Physical Oceanography*. Prentice Hall. ISBN 0-13-238155-9.
- United Nations Environment Programme / GRID-Arendal, 2006, [2]. *Potential Impact of Climate Change*
- Rahmstorf, S (2006). "Thermohaline Ocean Circulation". In Elias, S. A. *Encyclopedia of Quaternary Sciences*. Elsevier Science. ISBN 0-444-52747-8.

External links

- Ocean Conveyor Belt
- THOR FP7 projects http://www.eu-thor.eu investigates on the topic "Thermohaline overturning- at risk?" and the predictability of changes of the THC. THOR is financed by the 7th Framework Programme of the European Commission.
- Google Map of the Geography of the Thermohaline Circulation of our Oceans

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Thermohaline_circulation. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 11.1: Thermohaline circulation is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.


11.2: Thermocline

A **thermocline** (sometimes **metalimnion** in lakes) is a thin but distinct layer in a large body of fluid (e.g. water, such as an ocean or lake, or air, such as an atmosphere) in which temperature changes more rapidly with depth than it does in the layers above or below. In the ocean, the thermocline may be thought of as an invisible blanket which separates the upper mixed layer from the calm deep water below. Depending largely on season, latitude and turbulent mixing by wind, thermoclines may be a semi-permanent feature of the body of water in which they occur or they may form temporarily in response to phenomena such as the radiative heating/cooling of surface water during the day/night. Factors that affect the depth and thickness of a thermocline include seasonal weather variations, latitude and local environmental conditions, such as tides and currents.

Aquatic layers	
	Pelagic
	Photic
	Epipelagic
	Aphotic
	Mesopelagic
	Bathyalpelagic
	Abyssopelagic
	Hadopelagic
	Demersal
Stratification	
	Pycnocline
	Isopycnal
	Chemocline
	Halocline
	Thermocline
	Thermohaline
Marine habitats	
Lake stratification	
Aquatic ecosystems	
Wild fisheries	

Oceans

For the use of ocean thermoclines to generate energy, see Ocean thermal energy conversion.



Graph showing a tropical ocean thermocline (depth vs. temperature). Note the rapid change between 100 and 1000 meters. The temperature is nearly constant after 1500 meters depth.

Most of the heat energy of sunlight is absorbed in the first few centimeters at the ocean's surface, which heats during the day and cools at night as heat energy is lost to space by radiation. Waves mix the water near the surface layer and distribute heat to deeper water such that the temperature may be relatively uniform in the upper 100 m (300 ft), depending on wave strength and the existence of surface turbulence caused by currents. Below this mixed layer, the temperature remains relatively stable over day/night cycles. The temperature of the deep ocean drops gradually with depth. As saline water does not freeze until it reaches −2.3 °C (colder as depth and pressure increase) the temperature well below the surface is usually not far from zero degrees. [1]

The thermocline varies in depth. It is semi-permanent in the tropics, variable in temperate regions (often deepest during the summer) and shallow to nonexistent in the polar regions, where the water column is cold from the surface to the bottom. A layer of sea ice will act as an insulation blanket.

In the open ocean, the thermocline is characterized by a negative sound speed gradient, making the thermocline important in submarine warfare because it can reflect active sonar and other acoustic signals. Technically, this effect stems from a discontinuity in the acoustic impedance of water created by the sudden change in density.

When scuba diving, a thermocline where water drops in temperature by a few degrees Celsius quite suddenly can sometimes be observed between two bodies of water, for example where colder upwelling water runs into a surface layer of warmer water. It gives the water an appearance of wrinkled glass that is often used to obscure bathroom windows and is caused by the altered refractive index of the cold or warm water column. These same schlieren can be observed when hot air rises off the tarmac at airports or desert roads and is the cause of mirages.

Other water bodies

Thermoclines can also be observed in lakes. In colder climates, this leads to a phenomenon called stratification. During the summer, warm water, which is less dense, will sit on top of colder, denser, deeper water with a thermocline separating them. The warm layer is called the epilimnion and the cold layer is called the hypolimnion. Because the warm water is exposed to the sun during the day, a stable system exists and very little mixing of warm water and cold water occurs, particularly in calm weather.

One result of this stability is that as the summer wears on, there is less and less oxygen below the thermocline as the water below the thermocline never circulates to the surface and organisms in the water deplete the available oxygen. As winter approaches, the temperature of the surface water will drop as nighttime cooling dominates heat transfer. A point is reached where the density of the cooling surface water becomes greater than the density of the deep water and overturning begins as the dense surface water moves down under the influence of gravity. This process is aided by wind or any other process (currents for example) that agitates the water. This effect also occurs in Arctic and Antarctic waters, bringing water to the surface which, although low in oxygen, is higher in nutrients than the original surface water. This enriching of surface nutrients may produce blooms of phytoplankton, making these areas productive.

As the temperature continues to drop, the water on the surface may get cold enough to freeze and the lake/ocean begins to ice over. A new thermocline develops where the densest water (4 °C) sinks to the bottom, and the less dense water (water that is approaching the freezing point) rises to the top. Once this new stratification establishes itself, it lasts until the water warms enough for the

‘spring turnover,’ which occurs after the ice melts and the surface water temperature rises to 4 °C. During this transition, a thermal bar may develop.

Waves can occur on the thermocline, causing the depth of the thermocline as measured at a single location to oscillate (usually as a form of seiche). Alternately, the waves may be induced by flow over a raised bottom, producing a thermocline wave which does not change with time, but varies in depth as one moves into or against the flow.

Atmosphere

The lower atmosphere also typically contains a boundary between two distinct regions (the troposphere and stratosphere), but that boundary (the tropopause) displays quite different behavior. However, atmospheric thermoclines, or *inversions*, can occur, e.g. as nighttime cooling of the Earth’s surface produces cold, dense, often calm air adjacent to the ground. The coldest air is next to the ground, with air temperature increasing with height. At the top of this nighttime boundary layer (which may be only a hundred meters) the normal adiabatic temperature profile of the troposphere (i.e. temperature decreasing with altitude) is again observed. The thermocline or inversion layer occurs where the temperature profile changes from positive to negative with increasing height. The stability of the night time inversion is usually destroyed soon after sunrise as the sun’s energy warms the ground, which warms the air in the inversion layer. The warmer, less dense air then rises, destroying the stability that characterizes the nightly inversion.

This phenomenon was first applied to the field of noise pollution study in the 1960s, contributing to the design of urban highways and noise barriers.[2]

References

1. ^ “Temperature of Ocean Water”. UCAR. Retrieved 2008-05-09.
2. ^ Hogan, C. Michael (September 1973). “Analysis of highway noise”. *Water, Air, & Soil Pollution* **2** (3): 387–392. doi:10.1007/BF00159677.

External links

- “Temperature of Ocean Water”, University Corporation for Atmospheric Research, University of Michigan.

Retrieved from “http://en.Wikipedia.org/w/index.php?title=Thermocline&oldid=612864124“

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Thermocline. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 11.2: Thermocline is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

11.3: Tsunamis

Tsunami

Tsunami are described in the Earthquakes chapter as damaging waves that result from the sharp jolt to the water from an undersea earthquake. Landslides, meteorite impacts, or any other jolt to ocean water may form a tsunami (**Figure** below). Tsunami can travel at speeds of 800 kilometers per hour (500 miles per hour).

Tsunami have small wave heights and long wavelengths so they are usually unnoticed at sea. As the wave rides up the continental shelf the wave height increases.

A video explanation of tsunami is here: http://www.youtube.com/watch?v=StdqGoezNrY

The wave speed of a tsunami is also slowed by friction with the shallower ocean floor, which causes the wavelength to decrease, creating a much taller wave.

Many people caught in a tsunami have no warning of its approach. Since the wavelength is long, a long time can pass between crests or troughs onshore. In 1755 in Lisbon, an offshore earthquake caused a great deal of damage on land. People rushed to the open space of the shore and discovered that the water was flowing seaward fast. The trough of the tsunami wave reached shore first. People who went out onto the open beach were drowned when the crest of the wave reached shore.



A wave from the 2004 Boxing Day Tsunami hits the Maldives in the Indian Ocean.

Large tsunami in the Indian Ocean and more recently Japan have killed hundreds of thousands of people in recent years. The west coast is vulnerable to tsunami since it sits on the Pacific Ring of Fire. Scientists are trying to learn everything they can about predicting tsunamis before a massive one strikes a little closer to home. Learn more at: http://science.kqed.org/quest/video/scary-tsunamis/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=130

Lumen Learning authored content

- **License**: *Public Domain: No Known Copyright*

This page titled 11.3: Tsunamis is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

11.4: Ocean Organisms

Ocean Resources



LIVING RESOURCES

The ocean provides important living and nonliving resources. To be maintained for future use, these resources must be managed sustainably. Most fish are caught by lines or nets as they swim in the open waters of the ocean. Some species of fish are being over-harvested, which means their rate of reproduction cannot keep up with the rate at which people consume them.**Bottom trawling** is a method of fishing that involves towing a weighted net across the seafloor to harvest fish. In many areas where bottom trawling is done, ecosystems are severely disturbed by the large nets. For this reason, in a few areas in the world, laws limit bottom trawling to waters not more than 1,000 m deep or waters far from protected and sensitive areas. Still these actions protect some of the seafloor. Besides food, ocean organisms have other uses. Some provide us with medications.

NONLIVING RESOURCES

Oil and natural gas are the most valuable non-living resources taken from the ocean. Extracting these resources requires drilling into the seafloor. Oil platforms have dozens of oil wells that are drilled in places where the ocean is sometimes 2,000 m deep. A description of the Deepwater Horizon oil spill affecting the Gulf of Mexico is located in the Human Actions and Earth's Waters chapter.

The seafloor has some valuable minerals. **Manganese nodules** containing manganese, iron, copper, nickel, phosphate, and cobalt may be as small as a pea or as large as a basketball. Estimates are that there may be as much as 500 billion tons of nodules on the seafloor. The minerals in manganese nodules have many uses in the industrial world, but currently they are not being mined. Think back to the discussion of ore deposits in the Earth's Minerals chapter. Why do you think these seafloor resources are not being mined?

Types of Ocean Organisms

Oceans are a harsh placed to live. In the intertidal zone, conditions change rapidly as water covers and uncovers the region and waves pound on the rocks. Most of the environments at sea are cold and at just about any depth below the surface the pressure is very high. Beyond the photic zone, the ocean is entirely black. Organisms have adapted to these conditions in many interesting and effective ways. The size and variety of different habitats means that the oceans are home to a large portion of all life on Earth. It's interesting to note that the smallest and largest animals on Earth live in the oceans.

Marine animals breathe air or extract oxygen from the water. Some float on the surface and others dive into the ocean's depths. There are animals that eat other animals, and plants generate food from sunlight. A few bizarre creatures break down chemicals to make food! The following section divides ocean life into seven basic groups.



PLANKTON

Plankton are organisms that cannot swim but that float along with the current. The word “plankton” comes from the Greek for wanderer. Most plankton are microscopic, but some are visible to the naked eye.**Phytoplankton** are tiny plants that make food by photosynthesis. Because they need sunlight, phytoplankton live in the photic zone. Phytoplankton are responsible for about half of the total primary productivity (food energy) on Earth. Like other plants, phytoplankton release oxygen as a waste product.

Zooplankton, or animal plankton, eat phytoplankton as their source of food. Some zooplankton live as plankton all their lives and others are juvenile forms of animals that will attach to the bottom as adults. Some small invertebrates live as zooplankton.



PLANTS AND ALGAE

The few true plants found in the oceans include salt marsh grasses and mangrove trees. Although they are not true plants, large algae, which are called seaweed, also use photosynthesis to make food. Plants and seaweeds are found in the neritic zone where the light they need penetrates so that they can photosynthesize.

MARINE INVERTEBRATES

The variety and number of **invertebrates**, animals without a backbone, is truly remarkable. Marine invertebrates include sea slugs, sea anemones, starfish, octopi, clams, sponges, sea worms, crabs, and lobsters. Most of these animals are found close to the shore, but they can be found throughout the ocean.



FISH

Fish are **vertebrates**; they have a backbone. What are some of the features fish have that allows them to live in the oceans? All fish have most or all of these traits.

- Included among the fish are sardines, salmon, and eels, as well as the sharks and rays (which lack swim bladders).



REPTILES

Only a few types of reptiles live in the oceans and they live in warm water. Why are reptiles so restricted in their ability to live in the sea? Sea turtles, sea snakes, saltwater crocodiles, and marine iguana that are found only at the Galapagos Islands sum up the marine reptile groups. Sea snakes bear live young in the ocean, but turtles, crocodiles, and marine iguanas all lay their eggs on land.



SEABIRDS AND MARINE MAMMALS

Many types of birds are adapted to living in the sea or on the shore. What are the common traits of mammals? Mammals are endothermic (warm-blooded) vertebrates that give birth to live young; feed them with milk; and have hair, ears, and a jaw bone with teeth. What traits might mammals have to be adapted to life in the ocean?

- For swimming: streamlined bodies, slippery skin or hair, fins.
- For warmth: Fur, fat, high metabolic rate, small surface area to volume, specialized blood system.
- For salinity: kidneys that excrete salt, impervious skin.

Interactions Among Ocean Organisms



The previous section briefly discussed the adaptations different types of organisms have to live in the ocean. A look at a few of the different habitats organisms live in can focus even more on these important adaptations.

INTERTIDAL

A great abundance of life is found in the intertidal zone. High energy waves pound the organisms that live in this zone and so they must be adapted to pounding waves and exposure to air during low tides. Hard shells protect from pounding waves and also protect against drying out when the animal is above water. Strong attachments keep the animals anchored to the rock.

In a tide pool, as in the photo, what organisms are found where and what specific adaptations do they have to that zone? The mussels on the top left have hard shells for protection and to prevent drying because they are often not covered by water. The sea anemones in the lower right are more often submerged and have strong attachments but can close during low tides. Many young organisms get their start in estuaries and so they must be adapted to rapid shifts in salinity.

REEFS

Corals and other animals deposit calcium carbonate to create rock **reefs** near the shore. Coral reefs are the "rainforests of the oceans" with a tremendous amount of species diversity.

Reefs can form interesting shapes in the oceans. Remember that hot spots create volcanoes on the seafloor. If these volcanoes rise above sea level to become islands, and if they occur in tropical waters, coral reefs will form on them. Since the volcanoes are cones, the reef forms in a circle around the volcano. As the volcano comes off the hot spot, the crust cools. The volcano subsides and then begins to erode away. Eventually, all that is left is a reef island called an **atoll**. A **lagoon** is found inside the reef.

Coral reef are near shore and so are subject to pollution from land. The coral animals are very sensitive to temperature and reefs around the world are stressed from rising ocean temperatures.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=132	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=132

Oceanic Zone

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=132	The open ocean is a vast area. Food either washes down from the land or is created by photosynthesizing plankton. Zooplankton and larger animals feed on the phytoplankton and on each other. Larger animals such as whales and giant groupers may live their entire lives in the open water. The few species that live in the greatest depths are very specialized. Since it's rare to find a meal, the fish use very little energy; they move very little, breathe slowly, have minimal bone structure and a slow metabolism. These fish are very small. To maximize the chance of getting a meal, some species may have jaws that unhinge to accept a larger fish or backward-folding teeth to keep prey from escaping.

Hydrothermal Vents

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=132	At mid-ocean ridges at **hydrothermal vents**, bacteria that use chemosynthesis for food energy are the base of a unique ecosystem. This ecosystem is entirely separate from the photosynthesis at the surface. Shrimp, clams, fish, and giant tube worms have been found in these extreme places.

This page titled 11.4: Ocean Organisms is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

12: The Atmosphere

12.1: The Atmosphere

This page titled 12: The Atmosphere is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

12.1: The Atmosphere



Earth's atmosphere is a thin blanket of gases and tiny particles — together called air. We are most aware of air when it moves and creates wind. All living things need some of the gases in air for life support. Without an atmosphere, Earth would likely be just another lifeless rock.

Earth's atmosphere, along with the abundant liquid water at Earth's surface, are the keys to our planet's unique place in the solar system. Much of what makes Earth exceptional depends on the atmosphere. Let's consider some of the reasons we are lucky to have an atmosphere.

INDISPENSABLE FOR LIFE ON EARTH


Without the atmosphere, Earth would look a lot more like the Moon. Atmospheric gases, especially carbon dioxide (CO_2) and oxygen (O_2), are extremely important for living organisms. How does the atmosphere make life possible? How does life alter the atmosphere?

In **photosynthesis** plants use CO_2 and create O_2. Photosynthesis is responsible for nearly all of the oxygen currently found in the atmosphere. By creating oxygen and food, plants have made an environment that is favorable for animals. In respiration, animals use oxygen to convert sugar into food energy they can use. Plants also go through respiration and consume some of the sugars they produce.



CRUCIAL PART OF THE WATER CYCLE

As part of the hydrologic cycle, which was detailed in the Earth's Fresh Water chapter, water spends a lot of time in the atmosphere, mostly as water vapor.All **weather** takes place in the atmosphere, virtually all of it in the lower atmosphere. Weather describes what the atmosphere is like at a specific time and place, and may include temperature, wind, and precipitation. Weather is the change we experience from day to day. Climate is the long-term average of weather in a particular spot. Although the weather for a particular winter day in Tucson, Arizona, may include snow, the climate of Tucson is generally warm and dry.




OZONE LAYER MAKES LIFE POSSIBLE

Ozone is a molecule composed of three oxygen atoms, (O_3). Ozone in the upper atmosphere absorbs high-energy **ultraviolet (UV) radiation** coming from the Sun. This protects living things on Earth's surface from the Sun's most harmful rays. Without ozone for protection, only the simplest life forms would be able to live on Earth.MODERATES EARTH'S TEMPERATURE

Along with the oceans, the **atmosphere** keeps Earth's temperatures within an acceptable range.**Greenhouse gases** trap heat in the atmosphere so they help to moderate global temperatures. Without an atmosphere with greenhouse gases, Earth's temperatures would be frigid at night and scorching during the day. Important greenhouse gases include carbon dioxide, methane, water vapor, and ozone.


Atmospheric Gasses



COMPOSITION OF THE ATMOSPHERE

Nitrogen and oxygen together make up 99 percent of the planet's atmosphere. The rest of the gases are minor components but sometimes are very important. **Humidity** is the amount of water vapor in the air. Humidity varies from place to place and season to season. This fact is obvious if you compare a summer day in Atlanta, Georgia, where humidity is high, with a winter day in Phoenix, Arizona, where humidity is low. When the air is very humid, it feels heavy or sticky. Dry air usually feels more comfortable.Where around the globe is mean atmospheric water vapor higher and where is it lower and why? Higher humidity is found around the equatorial regions because air temperatures are higher and warm air can hold more moisture than cooler air. Of course, humidity is lower near the polar regions because air temperature is lower.

Some of what is in the atmosphere is not gas. Particles of dust, soil, fecal matter, metals, salt, smoke, ash, and other solids make up a small percentage of the atmosphere. Particles provide starting points (or nuclei) for water vapor to condense on and form raindrops. Some particles are pollutants, which are discussed in the Human Actions and the Atmosphere chapter.




ATMOSPHERIC PRESSURE AND DENSITY

The atmosphere has different properties at different elevations above sea level, or **altitudes**. The air density (the number of molecules in a given volume) decreases with increasing altitude. This is why people who climb tall mountains, such as Mt. Everest, have to set up camp at different elevations to let their bodies get used to the decreased air.Why does air density decrease with altitude? Gravity pulls the gas molecules towards Earth's center. The pull of gravity is stronger closer to the center at sea level. Air is denser at sea level where the gravitational pull is greater.Gases at sea level are also compressed by the weight of the atmosphere above them. The force of the air weighing down over a unit of area is known as its**atmospheric pressure**. The reason why we are not crushed by this weight is because the molecules inside our bodies are pushing outward to compensate. Atmospheric pressure is felt from all directions, not just from above.

At higher altitudes the atmospheric pressure is lower and the air is less dense than at higher altitudes. If your ears have ever "popped", you have experienced a change in air pressure. Gas molecules are found inside and outside your ears. When you change altitude quickly, like when an airplane is descending, your inner ear keeps the density of molecules at the original altitude. Eventually the air molecules inside your ear suddenly move through a small tube in your ear to equalize the pressure. This sudden rush of air is felt as a popping sensation.

Although the density of the atmosphere changes with altitude, the composition stays the same with altitude, with one exception. In the ozone layer, at about 20 km to 40 km above the surface, there is a greater concentration of ozone molecules than in other portions of the atmosphere.

Layers of the Atmosphere



The atmosphere is layered, corresponding with how the atmosphere's temperature changes with altitude. By understanding the way temperature changes with altitude, we can learn a lot about how the atmosphere works. While weather takes place in the lower atmosphere, interesting things, such as the beautiful aurora, happen higher in the atmosphere.

Why does warm air rise? Gas molecules are able to move freely and if they are uncontained, as they are in the atmosphere, they can take up more or less space.

- When gas molecules are cool, they are sluggish and do not take up as much space. With the same number of molecules in less space, both air density and air pressure are higher.
- When gas molecules are warm, they move vigorously and take up more space. Air density and air pressure are lower.

Warmer, lighter air is more buoyant than the cooler air above it, so it rises. The cooler air then sinks down, because it is denser than the air beneath it. This is convection, which was described in the Plate Tectonics chapter.

The property that changes most strikingly with altitude is air temperature. Unlike the change in pressure and density, which decrease with altitude, changes in air temperature are not regular. A change in temperature with distance is called a **temperature gradient**.

The atmosphere is divided into layers based on how the temperature in that layer changes with altitude, the layer's temperature gradient. The temperature gradient of each layer is different. In some layers, temperature increases with altitude and in others it decreases. The temperature gradient in each layer is determined by the heat source of the layer. Most of the important processes of the atmosphere take place in the lowest two layers: the troposphere and the stratosphere.



TROPOSPHERE


The temperature of the **troposphere** is highest near the surface of the Earth and decreases with altitude. On average, the temperature gradient of the troposphere is 6.5°C per 1,000 m (3.6°F per 1,000 ft.) of altitude. What is the source of heat for the troposphere? Earth's surface is a major source of heat for the troposphere, although nearly all of that heat comes from the Sun. Rock, soil, and water on Earth absorb the Sun's light and radiate it back into the atmosphere as heat. The temperature is also higher near the surface because of the greater density of gases.

Notice that in the troposphere warmer air is beneath cooler air. What do you think the consequence of this is? This condition is unstable. The warm air near the surface rises and cool air higher in the troposphere sinks. So air in the troposphere does a lot of mixing. This mixing causes the temperature gradient to vary with time and place. The rising and sinking of air in the troposphere means that all of the planet's weather takes place in the troposphere.

Sometimes there is a temperature **inversion**, air temperature in the troposphere increases with altitude and warm air sits over cold air. Inversions are very stable and may last for several days or even weeks. They form:

- Over land at night or in winter when the ground is cold. The cold ground cools the air that sits above it, making this low layer of air denser than the air above it.
- Near the coast where cold seawater cools the air above it. When that denser air moves inland, it slides beneath the warmer air over the land.

Since temperature inversions are stable, they often trap pollutants and produce unhealthy air conditions in cities. At the top of the troposphere is a thin layer in which the temperature does not change with height. This means that the cooler, denser air of the troposphere is trapped beneath the warmer, less dense air of the stratosphere. Air from the troposphere and stratosphere rarely mix.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=136



STRATOSPHERE


Ash and gas from a large volcanic eruption may burst into the **stratosphere**, the layer above the troposphere. Once in the stratosphere, it remains suspended there for many years because there is so little mixing between the two layers. Pilots like to fly in the lower portions of the stratosphere because there is little air turbulence.In the stratosphere, temperature increases with altitude. What is the heat source for the stratosphere? The direct heat source for the stratosphere is the Sun. Air in the stratosphere is stable because warmer, less dense air sits over cooler, denser air. As a result, there is little mixing of air within the layer.The **ozone layer** is found within the stratosphere between 15 to 30 km (9 to 19 miles) altitude. The thickness of the ozone layer varies by the season and also by latitude.

The ozone layer is extremely important because ozone gas in the stratosphere absorbs most of the Sun's harmful ultraviolet (UV) radiation. Because of this, the ozone layer protects life on Earth. High-energy UV light penetrates cells and damages DNA, leading to cell death (which we know as a bad sunburn). Organisms on Earth are not adapted to heavy UV exposure, which kills or damages them. Without the ozone layer to reflect UVC and UVB radiation, most complex life on Earth would not survive long.

MESOSPHERE

Temperatures in the **mesosphere** decrease with altitude. Because there are few gas molecules in the mesosphere to absorb the Sun's radiation, the heat source is the stratosphere below. The mesosphere is extremely cold, especially at its top, about -90 degrees C (-130 degrees F).

The air in the mesosphere has extremely low density: 99.9 percent of the mass of the atmosphere is below the mesosphere. As a result, air pressure is very low. A person traveling through the mesosphere would experience severe burns from ultraviolet light since the ozone layer which provides UV protection is in the stratosphere below. There would be almost no oxygen for breathing. Stranger yet, an unprotected traveler's blood would boil at normal body temperature because the pressure is so low.



THERMOSPHERE


The density of molecules is so low in the **thermosphere** that one gas molecule can go about 1 km before it collides with another molecule. Since so little energy is transferred, the air feels very cold. Within the thermosphere is the **ionosphere**. The ionosphere gets its name from the solar radiation that ionizes gas molecules to create a positively charged ion and one or more negatively charged electrons. The freed electrons travel within the ionosphere as electric currents. Because of the free ions, the ionosphere has many interesting characteristics. At night, radio waves bounce off the ionosphere and back to Earth. This is why you can often pick up an AM radio station far from its source at night.The Van Allen radiation belts are two doughnut-shaped zones of highly charged particles that are located beyond the atmosphere in the **magnetosphere**. The particles originate in solar flares and fly to Earth on the solar wind. Once trapped by Earth's magnetic field, they follow along the field's magnetic lines of force. These lines extend from above the equator to the North Pole and also to the South Pole then return to the equator.

When massive solar storms cause the Van Allen belts to become overloaded with particles, the result is the most spectacular feature of the ionosphere — the **aurora**. The particles spiral along magnetic field lines toward the poles. The charged particles energize oxygen and nitrogen gas molecules, causing them to light up. Each gas emits a particular color of light.

There is no real outer limit to the **exosphere**, the outermost layer of the atmosphere; the gas molecules finally become so scarce that at some point there are no more. Beyond the atmosphere is the solar wind. The solar wind is made of high-speed particles, mostly protons and electrons, traveling rapidly outward from the Sun.

There is no real outer limit to the **exosphere**, the outermost layer of the atmosphere; the gas molecules finally become so scarce that at some point there are no more. Beyond the atmosphere is the solar wind. The solar wind is made of high-speed particles, mostly protons and electrons, traveling rapidly outward from the Sun. This video is very thorough in its discussion of the layers of the atmosphere.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=136

Atmospheric Energy, Temperature, and Heat



ENERGY

Energy travels through space or material. This is obvious when you stand near a fire and feel its warmth or when you pick up the handle of a metal pot even though the handle is not sitting directly on the hot stove. Invisible energy waves can travel through air, glass, and even the vacuum of outer space. These waves have electrical and magnetic properties, so they are called electromagnetic waves. The transfer of energy from one object to another through electromagnetic waves is known as radiation. Different wavelengths of energy create different types of electromagnetic waves.

- Can you think of some objects that appear to radiate visible light, but actually do not? The moon and the planets do not emit light of their own; they reflect the light of the Sun. **Reflection** is when light (or another wave) bounces back from a surface. **Albedo** is a measure of how well a surface reflects light. A surface with high albedo reflects a large percentage of light. A snow field has high albedo.

  One important fact to remember is that energy cannot be created or destroyed — it can only be changed from one form to another. This is such a fundamental fact of nature that it is a law: the law of conservation of energy.

  ### TEMPERATURE

  Temperature is a measure of how fast the atoms in a material are vibrating. High temperature particles vibrate faster than low temperature particles. Rapidly vibrating atoms smash together, which generates heat. As a material cools down, the atoms vibrate more slowly and collide less frequently. As a result, they emit less heat. What is the difference between heat and temperature?

  - Which has higher heat and which has higher temperature: a candle flame or a bathtub full of hot water?

    - #### HEAT

      Heat is taken in or released when an object changes state, or changes from a gas to a liquid, or a liquid to a solid. This heat is called **latent heat**. When a substance changes state, latent heat is released or absorbed. A substance that is changing its state of matter does not change temperature. All of the energy that is released or absorbed goes toward changing the material's state.

      For example, imagine a pot of boiling water on a stove burner: that water is at 100 degrees C (212 degrees F). If you increase the temperature of the burner, more heat enters the water. The water remains at its boiling temperature, but the

additional energy goes into changing the water from liquid to gas. With more heat the water evaporates more rapidly. When water changes from a liquid to a gas it takes in heat. Since evaporation takes in heat, this is called evaporative cooling. Evaporative cooling is an inexpensive way to cool homes in hot, dry areas.


Energy from the Sun

The earth constantly tries to maintain an energy balance with the atmosphere. Most of the energy that reaches the Earth's surface comes from the Sun. About 44% of solar radiation is in the visible light wavelengths, but the Sun also emits infrared, ultraviolet, and other wavelengths. When viewed together, all of the wavelengths of visible light appear white. But a prism or water droplets can break the white light into different wavelengths so that separate colors appear.

Of the solar energy that reaches the outer atmosphere, UV wavelengths have the greatest energy. Only about 7 percent of solar radiation is in the UV wavelengths. The three types are:

- UVC: the highest energy ultraviolet, does not reach the planet's surface at all.
- UVB: the second highest energy, is also mostly stopped in the atmosphere.
- UVA: the lowest energy, travels through the atmosphere to the ground.

The remaining solar radiation is the longest wavelength, infrared. Most objects radiate infrared energy, which we feel as heat. Some of the wavelengths of solar radiation traveling through the atmosphere may be lost because they are absorbed by various gases. Ozone completely removes UVC, most UVB and some UVA from incoming sunlight. Oxygen, carbon dioxide, and water vapor also filter out some wavelengths.



Heat Transfer in the Atmosphere

Heat moves in the atmosphere the same way it moves through the solid Earth (Plate Tectonics chapter) or another medium. What follows is a review of the way heat flows and is transferred, but applied to the atmosphere.

Radiation is the transfer of energy between two objects by electromagnetic waves. Heat radiates from the ground into the lower atmosphere.

In **conduction**, heat moves from areas of more heat to areas of less heat by direct contact. Warmer molecules vibrate rapidly and collide with other nearby molecules, transferring their energy. In the atmosphere, conduction is more effective at lower altitudes where air density is higher; transfers heat upward to where the molecules are spread further apart or transfers heat laterally from a warmer to a cooler spot, where the molecules are moving less vigorously.

Heat transfer by movement of heated materials is called **convection**. Heat that radiates from the ground initiates convection cells in the atmosphere.

HEAT AT EARTH'S SURFACE


About half of the solar radiation that strikes the top of the atmosphere is filtered out before it reaches the ground. This energy can be absorbed by atmospheric gases, reflected by clouds, or scattered. Scattering occurs when a light wave strikes a particle and bounces off in some other direction.

About 3% of the energy that strikes the ground is reflected back into the atmosphere. The rest is absorbed by rocks, soil, and water and then radiated back into the air as heat. These infrared wavelengths can only be seen by infrared sensors. Because solar energy continually enters Earth's atmosphere and ground surface, is the planet getting hotter? The answer

is no (although the next section contains an exception) because energy from Earth escapes into space through the top of the atmosphere. If the amount that exits is equal to the amount that comes in, then average global temperature stays the same. This means that the planet's heat budget is in balance. What happens if more energy comes in than goes out? If more energy goes out than comes in?

To say that the Earth's heat budget is balanced ignores an important point. The amount of incoming solar energy is different at different latitudes). Where do you think the most solar energy ends up and why? Where does the least solar energy end up and why? The difference in solar energy received at different latitudes drives atmospheric circulation.

	Day Length	**Sun Angle**	**Solar Radiation**	**Albedo**
Equatorial Regions	Nearly same all year	High	High	Low
Polar Regions	Night 6 months	Low	Low	High



THE GREENHOUSE EFFECT

The exception to Earth's temperature being in balance is caused by greenhouse gases. But first the role of greenhouse gases in the atmosphere must be explained. Greenhouse gases warm the atmosphere by trapping heat. Some of the heat radiation out from the ground is trapped by greenhouse gases in the troposphere. Like a blanket on a sleeping person, greenhouse gases act as insulation for the planet. The warming of the atmosphere because of insulation by greenhouse gases is called the **greenhouse effect**. Greenhouse gases are the component of the atmosphere that moderate Earth's temperatures.Greenhouse gases include CO2, H2O, methane, O3, nitrous oxides (NO and NO2), and chlorofluorocarbons (CFCs). All are a normal part of the atmosphere except CFCs. The table below shows how each greenhouse gas naturally enters the atmosphere.

Greenhouse Gas	**Where It Comes From**
Carbon dioxide (CO_2)	Respiration, volcanic eruptions, decomposition of plant material; burning of fossil fuels
Methane	Decomposition of plant material under some conditions, biochemical reactions in stomachs
Nitrous oxide	Produced by bacteria; burning fossil fuels
Ozone	Atmospheric processes, chemical reactions resulting from burning fossil fuels
Chlorofluorocarbons (CFC)	Not naturally occurring; made by humans

Different greenhouse gases have different abilities to trap heat. For example, one methane molecule traps 30 times as much heat as one CO_2 molecule. One CFC-12 molecule (a type of CFC) traps 10,600 times as much heat as one CO_2. Still, CO_2 is a very important greenhouse gas because it is much more abundant in the atmosphere.

Human activity has significantly raised the levels of many of greenhouse gases in the atmosphere. Methane levels are about 2 1/2 times higher as a result of human activity. Carbon dioxide has increased more than 35%. CFCs have only recently existed.

What do you think happens as atmospheric greenhouse gas levels increase? More greenhouse gases trap more heat and warm the atmosphere. The increase or decrease of greenhouse gases in the atmosphere affect climate and weather the

world over.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-10-the-atmosphere.html. **License**: *CC BY: Attribution*

This page titled 12.1: The Atmosphere is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

13: Weather

13.1: Weather Processes and Systems

This page titled 13: Weather is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

13.1: Weather Processes and Systems

LEARNING OUTCOMES

- Describe the various aspects and elements of weather and atmospheric water.
- Explain how air masses and weather fronts together form mid-latitude cyclones.
- Describe the three phases a thunderstorm goes through in it's life cycle.
- Determine the causes and geographic location of most tornadoes around the world.
- Describe the genesis of hurricanes formation
- Compare the difference between blizzards lake effects, and heat waves.

Weather and Atmospheric Water

ELEMENTS OF WEATHER

If someone across country asks you what the weather is like today, you need to consider several factors. Air temperature, humidity, wind speed, the amount and types of clouds, and precipitation are all part of a thorough weather report. In this chapter, you will learn about many of these features in more detail.

Weather is what is going on in the atmosphere at a particular place at a particular time. Weather can change rapidly. A location's weather depends on air temperature; air pressure; fog; humidity; cloud cover; precipitation; wind speed and direction. All of these are directly related to the amount of energy that is in the system and where that energy is. The ultimate source of this energy is the sun.

Climate is the average of a region's weather over time. The climate for a particular place is steady, and changes only very slowly. Climate is determined by many factors, including the angle of the Sun, the likelihood of cloud cover, and the air pressure. All of these factors are related to the amount of energy that is found in that location over time.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140



HUMIDITY

Humidity is the amount of water vapor in the air in a particular spot. We usually use the term to mean **relative humidity**, the percentage of water vapor a certain volume of air is holding relative to the maximum amount it can contain. If the humidity today is 80%, it means that the air contains 80% of the total amount of water it can hold at that temperature. What will happen if the humidity increases to more than 100%? The excess water condenses and forms precipitation.

Since warm air can hold more water vapor than cool air, raising or lowering temperature can change air's relative humidity. The temperature at which air becomes saturated with water is called the air's dew point. This term makes sense, because water condenses from the air as dew, if the air cools down overnight and reaches 100% humidity.



CLOUDS

Clouds have a big influence on weather by preventing solar radiation from reaching the ground; absorbing warmth that is re-emitted from the ground; and as the source of precipitation.

When there are no clouds, there is less insulation. As a result, cloudless days can be extremely hot, and cloudless nights can be very cold. For this reason, cloudy days tend to have a lower range of temperatures than clear days.

There are a variety of conditions needed for clouds to form. First, clouds form when air reaches its dew point. This can happen in two ways: (1) Air temperature stays the same but humidity increases. This is common in locations that are warm and humid. (2) Humidity can remain the same, but temperature decreases. When the air cools enough to reach 100% humidity, water droplets form. Air cools when it comes into contact with a cold surface or when it rises.

Rising air creates clouds when it has been warmed at or near the ground level and then is pushed up over a mountain or mountain range or is thrust over a mass of cold, dense air. Water vapor is not visible unless it condenses to become a cloud. Water vapor condenses around a nucleus, such as dust, smoke, or a salt crystal. This forms a tiny liquid droplet. Billions of these water droplets together make a cloud.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

Clouds are classified in several ways. The most common classification used today divides clouds into four separate cloud groups, which are determined by their altitude and if precipitation is occurring or not.

High-level clouds form from ice crystals where the air is extremely cold and can hold little water vapor. **Cirrus**, **cirrostratus**, and **cirrocumulus**are all names of high clouds. Cirrocumulus clouds are small, white puffs that ripple across the sky, often in rows. Cirrus clouds may indicate that a storm is coming.

Middle-level clouds, including **altocumulus** and **altostratus** clouds, may be made of water droplets, ice crystals or both, depending on the air temperatures. Thick and broad altostratus clouds are gray or blue-gray. They often cover the entire sky and usually mean a large storm, bearing a lot of precipitation, is coming.

Low-level clouds are nearly all water droplets. **Stratus**, **stratocumulus** and **nimbostratus** clouds are common low clouds. Nimbostratus clouds are thick and dark that produce precipitation. Clouds with the prefix ‘cumulo-‘ grow vertically instead of horizontally and have their bases at low altitude and their tops at high or middle altitude. Clouds grow vertically when strong unstable air currents are rising upward. Common clouds include **cumulus humilis**, **cumulus mediocris**, **cumulus congestus**, and **cumulonimbus**.

To learn more about the various types of cloud formations from the National Oceanic and Atmospheric Administration (NOAA), click here.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140



FOG

Fog is a cloud located at or near the ground . When humid air near the ground cools below its dew point, fog is formed. The several types of fog that each form in a different way.

Radiation fog forms at night when skies are clear and the relative humidity is high. As the ground cools, the bottom layer of air cools below its dew point. Tule fog is an extreme form of radiation fog found in some regions. San Francisco, California, is famous for its summertime **advection fog**. Warm, moist Pacific Ocean air blows over the cold California current and cools below its dew point. Sea breezes bring the fog onshore. **Steam fog**appears in autumn when cool air moves over a warm lake. Water evaporates from the lake surface and condenses as it cools, appearing like steam. Warm humid air travels up a hillside and cools below its dew point to create **upslope fog**.




PRECIPITATION

Precipitation is an extremely important part of weather. Some precipitation forms in place. The most common precipitation comes from clouds. Rain or snow droplets grow as they ride air currents in a cloud and collect other droplets. They fall when they become heavy enough to escape from the rising air currents that hold them up in the cloud. One million cloud droplets will combine to make only one rain drop ! If temperatures are cold, the droplet will hit the ground as a snowflake.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

Air Masses

An **air mass** is a large mass of air that has similar characteristics of*temperature* and *humidity* within it. An air mass acquires these characteristics above an area of land or water known as its source region. When the air mass sits over a region for several days, or longer, it picks up the distinct temperature and humidity characteristics of that region.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140



AIR MASS FORMATION

Where an air mass receives it's characteristics of temperature and humidity is called the **source region**. Air masses are slowly pushed along by high-level winds, when an air mass moves over a new region, it shares its temperature and humidity with that region. So the temperature and humidity of a particular location depends partly on the characteristics of the air mass that sits over it.

Storms arise if the air mass and the region it moves over have different characteristics. For example, when a colder air mass moves over warmer ground, the bottom layer of air is heated. That air rises, forming clouds, rain, and sometimes thunderstorms. How would a moving air mass form an inversion? When a warmer air mass travels over colder ground, the bottom layer of air cools and, because of its high density, is trapped near the ground.

In general, cold air masses tend to flow toward the equator and warm air masses tend to flow toward the poles. This brings heat to cold areas and cools down areas that are warm. It is one of the many processes that act towards balancing out the planet's temperatures. Air masses are slowly pushed along by high-level winds. When an air mass moves over a new region, it shares its temperature and humidity with that region. So the temperature and humidity of a particular location depends partly on the characteristics of the air mass that sits over it. Air masses are classified based on their temperature and humidity characteristics. Below are examples of how air masses are classified over North America.

- Storms arise if the air mass and the region it moves over have different characteristics. For example, when a colder air mass moves over warmer ground, the bottom layer of air is heated. That air rises, forming clouds, rain, and sometimes thunderstorms. How would a moving air mass form an inversion? When a warmer air mass travels over colder ground, the bottom layer of air cools and, because of its high density, is trapped near the ground.

  In general, cold air masses tend to flow toward the equator and warm air masses tend to flow toward the poles. This brings heat to cold areas and cools down areas that are warm. It is one of the many processes that act towards balancing out the planet's temperatures.

Weather Front

The rest of this section will be devoted to four types of fronts. Three of these fronts move and one is stationary. With cold fronts and warm fronts, the air mass at the leading edge of the front gives the front its name. In other words, a cold front is right at the leading edge of moving cold air and a warm front marks the leading edge of moving warm air.

When two air masses meet together, the boundary between the two is called a **weather front**. At a front, the two air masses have different densities, based on temperature, and do not easily mix. One air mass is lifted above the other, creating a low pressure zone. If the lifted air is moist, there will be condensation and precipitation. Winds are common at a front. The greater the temperature difference between the two air masses, the stronger the winds will be. Fronts are the main cause of stormy weather.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140



The rest of this section will be devoted to four types of fronts. Three of these fronts move and one is stationary. With cold fronts and warm fronts, the air mass at the leading edge of the front gives the front its name. In other words, a cold front is right at the leading edge of moving cold air and a warm front marks the leading edge of moving warm air.

STATIONARY FRONTS

At a **stationary front** the air masses do not move. A front may become stationary if an air mass is stopped by a barrier, such as a mountain range.


A stationary front may bring days of rain, drizzle, and fog. Winds usually blow parallel to the front, but in opposite directions. After several days, the front will likely break apart. When a cold air mass takes the place of a warm air mass, there is a cold front.



COLD FRONTS

Imagine that you are standing in one spot as a **cold front** approaches. Along the cold front, the denser, cold air pushes up the warm air, causing the air pressure to decrease. If the humidity is high enough, some types of cumulus clouds will grow. High in the atmosphere, winds blow ice crystals from the tops of these clouds to create cirrostratus and cirrus clouds. At the front, there will be a line of rain showers, snow showers, or thunderstorms with blustery winds. A **squall line** is a line of severe thunderstorms that forms along a cold front. Behind the front is the cold air mass. This mass is drier so precipitation stops. The weather may be cold and clear or only partly cloudy. Winds may continue to blow into the low pressure zone at the front. The weather at a cold front varies with the season.

- spring and summer: The air is unstable so thunderstorms or tornadoes may form.
- spring: If the temperature gradient is high, strong winds blow.
- autumn: Strong rains fall over a large area.
- winter: The cold air mass is likely to have formed in the frigid arctic so there are frigid temperatures and heavy snows.



WARM FRONTS

Along a **warm front**, a warm air mass slides over a cold air mass. When warm, less dense air moves over the colder, denser air, the atmosphere is relatively stable.

Imagine that you are on the ground in the wintertime under a cold winter air mass with a warm front approaching. The transition from cold air to warm air takes place over a long distance so the first signs of changing weather appear long before the front is actually over you. Initially, the air is cold: the cold air mass is above you and the warm air mass is above it. High cirrus clouds mark the transition from one air mass to the other.


Over time, cirrus clouds become thicker and cirrostratus clouds form. As the front approaches, altocumulus and altostratus clouds appear and the sky turns gray. Since it is winter, snowflakes fall. The clouds thicken and nimbostratus clouds form. Snowfall increases. Winds grow stronger as the low pressure approaches. As the front gets closer, the cold air mass is just above you but the warm air mass is not too far above that. The weather worsens. As the warm air mass approaches, temperatures rise and snow turns to sleet and freezing rain. Warm and cold air mix at the front, leading to the formation of stratus clouds and fog.

OCCLUDED FRONTS

An **occluded front** usually forms around a low pressure system. The occlusion starts when a cold front catches up to a warm front. The air masses, in order from front to back, are cold, warm, and then cold again.

Coriolis Effect curves the boundary where the two fronts meet towards the pole. If the air mass that arrives third is colder than either of the first two air masses, that air mass slip beneath them both. This is called a cold occlusion. If the air mass that arrives third is warm, that air mass rides over the other air mass. This is called a warm occlusion.

The weather at an occluded front is especially fierce right at the occlusion. Precipitation and shifting winds are typical. The Pacific Coast has frequent occluded fronts.



Remember, a weather front is basically the boundary between two air masses of different densities. At the center of each air mass is typically a high pressure. This means that weather is typically sunny within air masses, but their temperatures could vary with the season and humidity could vary based on the source region of the air mass.

Now more often than not, these weather fronts are not isolated events. Often times, they are part of a larger rotating system called a mid-latitude cyclone. This type of cyclone will be discussed later in this chapter, but as an introduction it is a low pressure system that is usually mixing warmer air from the south (in the Northern Hemisphere) and colder air from the north.

Thunderstorms



Weather happens every day, but only some days have storms. Storms vary immensely depending on whether they're warm or cold, coming off the ocean or off a continent, occurring in summer or winter, and many other factors. The effects of storms also vary depending on whether they strike a populated area or a natural landscape. Hurricane Katrina is a good example, since the

flooding after the storm severely damaged New Orleans, while a similar storm in an unpopulated area would have done little damage.


Thunderstorms are extremely common: Worldwide there are 14 million per year; that's 40,000 per day! Most drop a lot of rain on a small area quickly, but some are severe and highly damaging. They form when ground temperatures are high, ordinarily in the late afternoon or early evening in spring and summer. The two figures below show two stages of thunderstorm buildup.

THUNDERSTORM GENESIS

All thunderstorms go through a three-stage life cycle. The first stage is called the **cumulus stage**, where an air parcel is forced to rise, cool, and condense, called the lower condensation level, to develop into a cumulus cloud. The process of water vapor condensing into liquid water releases large quantities of latent heat, which makes the air within the cloud warmer, and unstable causing the cloud continues to grow upward like a hot air balloon. These rising air parcels, called updrafts, prevent precipitation from falling from the cloud. But once the precipitation becomes too heavy for the updrafts to hold up, the moisture begins to fall creating downdrafts within the cloud. The downdrafts also begin to pull cold, dry air from outside the cloud toward the ground in a process called**entrainment**.

Once the precipitation begins to fall from the cloud, the storm has reached the **mature stage**. During this stage, updrafts and downdrafts exist side-by-side and the cumulonimbus is called a **cell**. If the updrafts reach the top of the troposphere, the cumulus cloud will begin to spread outward creating a defined **anvil**. At the same time, the downdrafts spread within the cloud and at first make the cloud become wider, but eventually overtaking the updrafts. Cool downdrafts form when precipitation and the cool air from entrainment are dragged down to the lower regions of a thunderstorm. It is also during the mature stage when the storm is most intense producing strong, gusting winds, heavy precipitation, lightning, and possibly small hail.

Once the downdrafts overtake the updrafts, which also prevents the release of latent heat energy, the thunderstorm will begin to weaken into the third and final stage, called the **dissipating stage**. During this stage, light precipitation and downdrafts become the dominate feature within the cloud as it weakens. In all, only twenty percent of the moisture within the cloud fell as precipitation whereas the other eighty percent evaporates back into the atmosphere.





With severe thunderstorms, the downdrafts are so intense that when they hit the ground it sends warm air from the ground upward into the storm. The warm air gives the convection cells more energy. Rain and hail grow huge before gravity pulls them to Earth. Severe thunderstorms can last for hours and can cause a lot of damage because of high winds, flooding, intense hail, and tornadoes.

Thunderstorms can form individually or in squall lines along a cold front. In the United States, squall lines form in spring and early summer in the Midwest where the maritime tropical (mT) air mass from the Gulf of Mexico meets the continental polar

(cP) air mass from Canada.

So much energy collects in cumulonimbus clouds that a huge release of electricity, called **lightning**, may result. The electrical discharge may be between one part of the cloud and another, two clouds, or a cloud and the ground.

Lightning heats the air so that it expands explosively. The loud clap is thunder. Light waves travel so rapidly that lightning is seen instantly. Sound waves travel much more slowly, so a thunderclap may come many seconds after the lightning is spotted. Thunderstorms kill approximately 200 people in the United States and injure about 550 Americans per year, mostly from lightning strikes. Have you heard the common misconception that lightning doesn't strike the same place twice? In fact, lightning strikes the New York City's Empire State Building about 100 times per year.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

Tornadoes



Tornadoes, also called twisters, are fierce products of severe thunderstorms. As air in a thunderstorm rises, the surrounding air races in to fill the gap, forming a funnel. A tornado lasts from a few seconds to several hours. The average wind speed is about 177 kph (110 mph), but some winds are much faster. A tornado travels over the ground at about 45 km per hour (28 miles per hour) and goes about 25 km (16 miles) before losing energy and disappearing.

An individual tornado strikes a small area, but it can destroy everything in its path. Most injuries and deaths from tornadoes are caused by flying debris. In the United States an average of 90 people are killed by tornadoes each year. The most violent two percent of tornadoes account for 70% of the deaths by tornadoes.

Tornadoes form at the front of severe thunderstorms. Lines of these thunderstorms form in the spring where where maritime tropical (mT) and continental polar (cP) air masses meet. Although there is an average of 770 tornadoes annually, the number of tornadoes each year varies greatly.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

In late April 2011, the situation was ripe for the deadliest set of tornadoes in 25 years. In addition to the meeting of cP and mT mentioned above, the jet stream was blowing strongly in from the west. The result was more than 150 tornadoes reported throughout the day.

The entire region was alerted to the possibility of tornadoes in those late April days. But meteorologists can only predict tornado danger over a very wide region. No one can tell exactly where and when a tornado will touch down. Once a tornado is sighted

on radar, its path is predicted and a warning is issued to people in that area. The exact path is unknown because tornado movement is not very predictable.

The intensity of tornadoes is measured on the Fujita Scale, which assigns a value based on wind speed and damage.

F-Scale	**3 Second Gust (mph)**	**Damage**
EF0 (Gale)	65-85	Light – tree branches fall and chimneys may collapse
EF1 (Weak)	86-110	Moderate – mobile homes, automobiles pushed aside
EF2 (Strong)	111-135	Considerate – roofs torn off houses, large trees uprooted
EF3 (Severe)	136-165	Severe – houses torn apart, trees uprooted, cars lifted
EF4 (Devastating)	166-200	Devastating – houses leveled, cars thrown
EF5 (Incredible)	Over 200	Incredible – structures fly, cars become missiles

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140


Cyclones



Cyclones can be the most intense storms on Earth. A **cyclone** is a system of winds rotating counterclockwise in the Northern Hemisphere around a low pressure center. The swirling air rises and cools, creating clouds and precipitation.

There are two types of cyclones: middle latitude (mid-latitude) cyclones and tropical cyclones. Mid-latitude cyclones are the main cause of winter storms in the middle latitudes. Tropical cyclones are also known as hurricanes.

An **anticyclone** is the opposite of a cyclone. An anticyclone's winds rotate clockwise in the Northern Hemisphere around a center of high pressure. Air comes in from above and sinks to the ground. High pressure centers generally have fair weather.




MID-LATITUDE CYCLONES

Mid-latitude cyclones, sometimes called **extratropical cyclones**, form at the polar front when the temperature difference between two air masses is large. These air masses blow past each other in opposite directions. Coriolis Effect deflects winds to the right in the Northern Hemisphere, causing the winds to strike the polar front at an angle. Warm and cold fronts form next to each other. Most winter storms in the middle latitudes, including most of the United States and Europe, are caused by mid-latitude cyclones.

The warm air at the cold front rises and creates a low pressure cell. Winds rush into the low pressure and create a rising column of air. The air twists, rotating counterclockwise in the Northern Hemisphere and clockwise in the Southern Hemisphere. Since the rising air is moist, rain or snow falls.

Mid-latitude cyclones form in winter in the mid-latitudes and move eastward with the westerly winds. These two- to five-day storms can reach 1,000 to 2,500 km (625 to 1,600 miles) in diameter and produce winds up to 125 km (75 miles) per hour. Like tropical cyclones, they can cause extensive beach erosion and flooding.

Mid-latitude cyclones are especially fierce in the mid-Atlantic and New England states where they are called **nor'easters**, because they come from the northeast. About 30 nor'easters strike the region each year.



HURRICANES

Tropical cyclones have many names. They are called hurricanes in the North Atlantic and eastern Pacific oceans, typhoons in the western Pacific Ocean, tropical cyclones in the Indian Ocean, and willi-willi's in the waters near Australia. By any name, they are the most damaging storms on Earth.

Hurricanes arise in the tropical latitudes (between 10 degrees and 25 degrees N) in summer and autumn when sea surface temperature are 28 degrees C (82 degrees F) or higher. The warm seas create a large humid air mass. The warm air rises and forms a low pressure cell, known as a tropical depression. Thunderstorms materialize around the tropical depression.

If the temperature reaches or exceeds 28 degrees C (82 degrees F) the air begins to rotate around the low pressure (counterclockwise in the Northern Hemisphere and clockwise in the Southern Hemisphere). As the air rises, water vapor condenses, releasing energy from latent heat. If wind shear is low, the storm builds into a hurricane within two to three days.

Hurricanes are huge with high winds. The exception is the relatively calm eye of the storm where air is rising upward. Rainfall can be as high as 2.5 cm (1″) per hour, resulting in about 20 billion metric tons of water released daily in a hurricane. The release of latent heat generates enormous amounts of energy, nearly the total annual electrical power consumption of the United States from one storm. Hurricanes can also generate tornadoes.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

Hurricanes are huge with high winds. The exception is the relatively calm eye of the storm where air is rising upward. Rainfall can be as high as 2.5 cm (1″) per hour, resulting in about 20 billion metric tons of water released daily in a hurricane. The release of latent heat generates enormous amounts of energy, nearly the total annual electrical power consumption of the United States from one storm. Hurricanes can also generate tornadoes.


Hurricanes are strange creatures because they are deadly monsters, yet have a gentle, but cold heart. The anatomy of a hurricane is fairly simple, though the processes involved are quite complex. As a low pressure disturbance forms, the warm, moist air rushes towards the low pressure in order to rise upward to form towering thunderstorms. Around the low pressure disturbance is a wall of clouds called an **eye wall**. Within the eye wall, the wind speeds are greatest, the clouds are the tallest, atmospheric pressure is at its lowest, and precipitation is most intense.

At the center or heart of the hurricane is called the **eye**. Within the eye of a hurricane, winds are light, precipitation is minimal, and occasionally the skies above are clear. It is the calm region of the tropical storm, but that is what makes it so dangerous. Many people tend to go outside as the eye moves overhead because they believe the storm is over. But what some don't realize is that "round two" is coming from behind.

Moving away from the eye wall are organized, intense thunderstorms, called **spiral rain bands**, that rotate around and toward the storm's eye wall. These rain bands are the first

Hurricanes are assigned to categories based on their wind speed. The categories are listed on the **Saffir-Simpson Scale**.

Category	**MPH**	**Estimated Damage**
1 (Weak)	74-95	Above normal; no read damage to structures
2 (Moderate)	96-110	Some property damage; considerable damage to vegetation
3 (Strong)	111-130	Some buildings damaged; mobile homes destroyed
4 (Very strong)	131-156	Complete roof failure on small residences; major beach erosion of beach areas
5 (Devastating)	Over 156	Complete roof and some building failure on most residential and industrial buildings



Hurricanes move with the prevailing winds. In the Northern Hemisphere, theyoriginate in the trade winds and move to the west. When they reach the latitude of the westerlies, they switch direction and travel toward the north or northeast. Hurricanes may cover 800 km (500 miles) in one day.

Damage from hurricanes comes from the high winds, rainfall, and storm surge.**Storm surge** occurs as the storm's low pressure center comes onto land, causing the sea level to rise unusually high. A storm surge is often made worse by the hurricane's high winds blowing seawater across the ocean onto the shoreline. Flooding can be devastating, especially along low-lying coastlines such as the Atlantic and Gulf Coasts. Hurricane Camille in 1969 had a 7.3 m (24 foot) storm surge that traveled 125 miles (200 km) inland.

Hurricanes typically last for 5 to 10 days. Over cooler water or land, the hurricane's latent heat source shut downs and the storm weakens. When a hurricane disintegrates, it is replaced with intense rains and tornadoes.

There are about 100 hurricanes around the world each year, plus many smaller tropical storms and tropical depressions. As people develop coastal regions, property damage from storms continues to rise. However, scientists are becoming better at predicting the paths of these storms and fatalities are decreasing. There is, however, one major exception to the previous statement: Hurricane Katrina.

For over a decade, the conversation in the United States over hurricanes was directed toward Hurricane Katrina. But in late October 2012, a Category 3 hurricane called Hurricane Sandy, sometimes called Superstorm Sandy, struck the eastern coast of the United States. It is estimaged that the storm system caused $68 billion in damage and killed nearly 300 people. What made Hurricane Sandy different from most tropical storms was how powerful it was so late in the hurricane season. The concern is that as the planet warms from anthropogenic (human-enhanced) climate change, the oceans will likely warm up allow hurricanes to become stronger, travel farther toward the poles, and the hurricane season last longer.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

To view an interactive map on hurricanes created by the National Oceanic and Atmospheric Administration, click here.



Lake-effect Snow off the Great Lakes.

BLIZZARDS AND LAKE EFFECTS

A **blizzard** is distinguished by certain conditions:

- Blizzards happen across the middle latitudes and toward the poles, usually as part of a mid-latitude cyclone. Blizzards are most common in winter, when the jet stream has traveled south and a cold, northern air mass comes into contact with a warmer, semitropical air mass. The very strong winds develop because of the pressure gradient between the low pressure storm and the higher pressure west of the storm. Snow produced by the storm gets caught in the winds and blows nearly horizontally. Blizzards can also produce **sleet** or **freezing rain**.

  In winter, a continental polar air mass travels down from Canada. As the frigid air travels across one of the Great Lakes, it warms and absorbs moisture. When the air mass reaches the leeward side of the lake, it is very unstable and it drops tremendous amounts of snow. This **lake-effect snow** falls on the snowiest, metropolitan areas in the United States: Buffalo and Rochester, New York.



High pressure in the middle layers of the atmosphere acts as a dome or cap allowing heat to build up at the earth's surface.

HEAT WAVES

Even more insidious are the deadliest weather phenomena, a **heat wave**. A heat wave is different for different locations; it is a long period of hot weather, at least 86 degrees F (30 degrees C) for at least three days in cooler locations but much more in hotter locations. Heat waves have increased in frequency and duration in recent years.

What do you think caused the heat wave in the image below? A high pressure zone kept the jet stream further north than normal for August.

Collecting Weather Data

Weather forecasts are better than they ever have been. According to the World Meteorological Organization (WMO), a 5-day weather forecast today is as reliable as a 2-day forecast was 20 years ago! This is because forecasters now use advanced technologies to gather weather data, along with the world's most powerful computers. Together, the data and computers produce complex models that more accurately represent the conditions of the atmosphere. These models can be programmed to predict how the atmosphere and the weather will change. Despite these advances, weather forecasts are still often incorrect. Weather is extremely difficult to predict because it is a complex and chaotic system.

To make a weather forecast, the conditions of the atmosphere must be known for that location and for the surrounding area. Temperature, air pressure, and other characteristics of the atmosphere must be measured and the data collected.



THERMOMETER

Thermometers measure temperature. In an old-style mercury thermometer, mercury is placed in a long, very narrow tube with a bulb. Because mercury is temperature sensitive, it expands when temperatures are high and contracts when they are low. A scale on the outside of the thermometer matches up with the air temperature.

Some modern thermometers use a coiled strip composed of two kinds of metal, each of which conducts heat differently. As the temperature rises and falls, the coil unfolds or curls up tighter. Other modern thermometers measure infrared radiation or electrical resistance. Modern thermometers usually produce digital data that can be fed directly into a computer.

BAROMETER

Meteorologists use **barometers** to measure air pressure. A barometer may contain water, air, or mercury, but like thermometers, barometers are now mostly digital. A change in barometric pressure indicates that a change in weather is coming. If air pressure rises, a high pressure cell is on the way and clear skies can be expected. If pressure falls, a low pressure cell is coming and will likely bring storm clouds. Barometric pressure data over a larger area can be used to identify pressure systems, fronts, and other weather systems.




WEATHER STATIONS

Weather stations contain some type of thermometer and barometer. Other instruments measure different characteristics of the atmosphere such as wind speed, wind direction, humidity, and amount of precipitation. These instruments are placed in various locations so that they can check the atmospheric characteristics of that location. According to the WMO, weather information is collected from 15 satellites, 100 stationary buoys, 600 drifting buoys, 3,000 aircraft, 7,300 ships, and some 10,000 land-based stations. The official weather stations used by the National Weather Service is called the Automated Surface Observing System (ASOS).

RADIOSONDES

Radiosondes is a balloon that measures atmospheric characteristics, such as temperature, pressure, and humidity as they move through the air. Radiosondes in flight can be tracked to obtain wind speed and direction. Radiosondes use a radio to communicate the data they collect to a computer. Radiosondes are launched from about 800 sites around the globe twice daily to provide a profile of the atmosphere. Radiosondes can be dropped from a balloon or airplane to make measurements as they fall. This is done to monitor storms, for example, since they are dangerous places for airplanes to fly.



RADAR


Radar stands for Radio Detection and Ranging. A transmitter sends out radio waves that bounce off the nearest object and then return to a receiver. Weather radar can sense many characteristics of precipitation: its location, motion, intensity, and the likelihood of future precipitation. Doppler radar can also track how fast the precipitation falls. Radar can outline the structure of a storm and can be used to estimate its possible effects.



SATELLITES


Weather satellites have been increasingly important sources of weather data since the first one was launched in 1952 and are the best way to monitor large scale systems, such as storms. Satellites are able to record long-term changes, such as the amount of ice cover over the Arctic Ocean in September each year. They also observe all energy from all wavelengths in the electromagnetic spectrum.

The flagship of the National Weather Service is the Geostationary Operational Environmental Satellites (GOES). These satellites are the ones you see on the nightly news where it looks like the clouds are moving, but not the planet. That is because these satellites are "geo-fixed" on a particular location over Earth rotating around the planet as fast as Earth's rotation at a distance of over 23,000 miles above the planet. There are basically three different types of GOES: visible, infrared, and water vapor. Visible light images record storms, clouds, fires, and smog. Infrared images record clouds, water and land temperatures, and features of the ocean, such as ocean currents**.** The final type of GOES imagery is water vapor. This type of imagery looks at the moisture content in the upper-half of the atmosphere. This is important for determining if clouds can grow to great heights like cumulonimbus thunderstorms.



The other type of satellite commonly used in weather forecasting is called a Polar Orbiting Environmental Satellites (POES). These types of satellites fly much lower to the earth, only about 530 miles, and orbit the planet pole-to-pole. You've probably seen these satellites at night when you see one crossing the sky. Look for their direction and odds are they are moving northward or southward toward each pole.

Just like the weather satellites on the news, you've seen these images often when you are looking at natural disasters like hurricanes or volcanic eruptions, wars like have occurred in Afghanistan, Iraq, or recently in Syria. Even the Malaysian flight that "disappeared" in the Indian Ocean for weeks was ultimately discovered using polar orbiting satellites. Common types of these satellites include: Landsat,MODIS, and the Tropical Rainfall Measuring Mission (TRMM).



NUMERICAL WEATHER PREDICTION

The most accurate weather forecasts are made by advanced computers, with analysis and interpretation added by experienced meteorologists. These computers have up-to-date mathematical models that can use much more data and make many more calculations than would ever be possible by scientists working with just maps and calculators. Meteorologists can use these results to give much more accurate weather forecasts and climate predictions.

In Numerical Weather Prediction (NWP), atmospheric data from many sources are plugged into supercomputers running complex mathematical models. The models then calculate what will happen over time at various altitudes for a grid of evenly spaced locations. The grid points are usually between 10 and 200 kilometers apart. Using the results calculated by the model, the program projects weather further into the future. It then uses these results to project the weather still further into the future, as far as the meteorologists want to go. Once a forecast is made, it is broadcast by satellites to more than 1,000 sites around the world.

NWP produces the most accurate weather forecasts, but as anyone knows, even the best forecasts are not always right. Weather prediction is extremely valuable for reducing property damage and even fatalities. If the proposed track of a hurricane can be predicted, people can try to secure their property and then evacuate.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=140

WEATHER MAPS

Weather maps, also called **synoptic maps**, simply and graphically depict meteorological conditions in the atmosphere from a spatial perspective. Weather maps may display only one feature of the atmosphere or multiple features. They can depict information from computer models or from human observations.

On a weather map, important meteorological conditions are plotted for each weather station. Metorologists use many different symbols as a quick and easy way to display information on the map.



Once conditions have been plotted, points of equal value can be connected by isolines. Weather maps can have many types of connecting lines. For example:

- Surface weather analysis maps are weather maps that only show conditions on the ground.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-11-weather.html. **License**: *CC BY: Attribution*

This page titled 13.1: Weather Processes and Systems is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

14: Astronomy

14.1: The Universe

This page titled 14: Astronomy is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

14.1: The Universe



INTRODUCTION

The Whirlpool Galaxy, also known as M51, is a spiral galaxy about 23 million light-years from Earth. Its interactions with the yellowish dwarf galaxy NGC 5195 are of interest to astronomers because the galaxies are near enough to Earth to be well-studied.Decades ago astronomers could not tell if these two galaxies were just passing each other but radio astronomy has supplied astronomers with important data outlining their interactions. Using this data, astronomers have simulated the interaction. NGC 5195 came from behind and then passed through the main disk of M51 about 500 to 600 million years ago. The dwarf galaxy crossed the disk again between 50 and 100 million years ago and is now slightly behind M51. These interactions appear to have intensified the spiral arms that are the dominant characteristic of the Whirlpool Galaxy.Astronomers are able to learn about objects unimaginably far away from Earth using telescopes that sense all wavelengths on the electromagnetic spectrum. Imagine what Galileo would do if he could see the images and data astronomers have available to them now. The study of the universe is called **cosmology**. **Cosmologists** study the structure and changes in the present universe. The universe contains all of the star systems, galaxies, gas and dust, plus all the matter and energy that exists now, that existed in the past, and that will exist in the future. The universe includes all of space and time.LEARNING OUTCOMES

- Understand the scientific ideas of how the universe formed and is expanding
- Compare and contrast the difference and similarities between dark matter and dark energy
- Describe star systems and the various types of galaxies

Expanding Universe



What did the ancient Greeks recognize as the universe? In their model, the universe contained Earth at the center, the Sun, the Moon, five planets, and a sphere to which all the stars were attached. This idea held for many centuries until Galileo's telescope helped allow people to realize that Earth is not the center of the universe. They also found out that there are many more stars than were visible to the naked eye. All of those stars were in the Milky Way Galaxy.


In the early 20th century, an astronomer named Edwin Hubble Figure below discovered that what scientists called the Andromeda Nebula was actually over 2 million light years away, many times farther than the farthest distances that had ever been measured. Hubble realized that many of the objects that astronomers called nebulas were not actually clouds of gas, but were collections of millions or billions of stars that we now call **galaxies**.

Hubble showed that the universe was much larger than our own galaxy. Today, we know that the universe contains about a hundred billion galaxies, about the same number of galaxies as there are stars in the Milky Way Galaxy. After discovering that there are galaxies beyond the Milky Way, Edwin Hubble went on to measure the distance to hundreds of other galaxies. His data would eventually show how the universe is changing, and would even yield clues as to how the universe formed. Today we know know that the universe in nearly 14 billion years old.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144

REDSHIFT

If you look at a star through a prism, you will see a spectrum, or a range of colors through the rainbow. The spectrum will have specific dark bands where elements in the star absorb light of certain energies. By examining the arrangement of these dark absorption lines, astronomers can determine the composition of elements that make up a distant star. In fact, the element helium was first discovered in our Sun, not on Earth, by analyzing the absorption lines in the spectrum of the Sun.While studying the spectrum of light from distant galaxies, astronomers noticed something strange. The dark lines in the spectrum were in the patterns they expected, but they were shifted toward the red end of the spectrum, as shown in Figure below. This shift of absorption bands toward the red end of the spectrum is known as redshift.**Redshift** occurs when the light source is moving away from the observer or when the space between the observer and the source is stretched. What does it mean that stars and galaxies are redshifted? When astronomers see redshift in the light from a galaxy, they know that the galaxy is moving away from Earth. What astronomers are noticing is that all the galaxies have a redshift, strongly indicating that all galaxies are moving away from each other causing the Universe to expand.Redshift can occur with other types of waves too, called the **Doppler Effect**. An analogy to redshift is the noise a siren makes as it passes you. You may have noticed that an ambulance seems to lower the pitch of its siren after it passes you. The sound waves shift towards a lower pitch when the ambulance speeds away from you. Though redshift involves light instead of sound, a similar principle operates in both situations. Click here to view an animation of Doppler Effect.






THE EXPANDING UNIVERSE

Edwin Hubble combined his measurements of the distances to galaxies with other astronomers' measurements of redshift. From this data, he noticed a relationship, which is now called **Hubble's Law**. The law states that the farther away a galaxy is, the faster it is moving away from us. What this leads to is the hypothesis that the universe is expanding. For an animation demonstrating the expanding universe, click here.The figure here by NASA shows a simplified diagram of the expansion of the universe. If you look closely at the diagram, you that on the left was the formation of the universe and the energy is quite high. Over the course of the 13.7 billion years, the energy begins to cool enough to create trillions of stars and over time develop into galaxies. Over time, the galaxies continue to cool and expand farther apart from each other.n

Formation of the Universe



Before Hubble, most astronomers thought that the universe didn't change. But if the universe is expanding, what does that say about where it was in the past? If the universe is expanding, the next logical thought is that in the past it had to have been smaller.


THE BIG BANG THEORY

The Big Bang theory is the most widely accepted cosmological explanation of how the universe formed. If we start at the present and go back into the past, the universe is contracting, getting smaller and smaller. What is the end result of a contracting universe?

According to the Big Bang theory, the universe began about 13.7 billion years ago. Everything that is now in the universe was squeezed into a very small volume. Imagine all of the known universe in a single, hot, chaotic mass. An enormous explosion — a big bang — caused the universe to start expanding rapidly. All the matter and energy in the universe, and even space itself, came out of this explosion. What came before the Big Bang? There is no way for scientists to know since there is no remaining evidence.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144




AFTER THE BIG BANG

In the first few moments after the Big Bang, the universe was unimaginably hot and dense. As the universe expanded, it became less dense and began to cool. After only a few seconds, protons, neutrons, and electrons could form. After a few minutes, those subatomic particles came together to create hydrogen. Energy in the universe was great enough to initiate nuclear fusion and hydrogen nuclei were fused into helium nuclei. The first neutral atoms that included electrons did not form until about 380,000 years later.The matter in the early universe was not smoothly distributed across space. Dense clumps of matter held close together by gravity were spread around. Eventually, these clumps formed countless trillions of stars, billions of galaxies, and other structures that now form most of the visible mass of the universe. If you look at an image of galaxies at the far edge of what we can see, you are looking at great distances. But you are also looking across a different type of distance. What do those far away galaxies represent? Because it takes so long for light from so far away to reach us, you are also looking back in time.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144

Dark Matter and Dark Energy



The Big Bang theory is still the best scientific model we have for explaining the formation of the universe and many lines of evidence support it. However, recent discoveries continue to shake up our understanding of the universe. Astronomers and other scientists are now wrestling with some unanswered questions about what the universe is made of and why it is expanding. A lot of what cosmologists do is create mathematical models and computer simulations to account for these unknown phenomena, such as dark energy and dark matter.

DARK MATTER

The things we observe in space are objects that emit some type of electromagnetic radiation. However, scientists think that matter that emits light makes up only a small part of the matter in the universe. The rest of the matter, about 80 percent, is dark matter.Dark matter emits no electromagnetic radiation so we can't observe it directly. However, astronomers know that dark matter exists because its gravity affects the motion of objects around it. When astronomers measure how spiral galaxies rotate, they find that the outside edges of a galaxy rotate at the same speed as parts closer to the center. This can only be explained if there is a lot more matter in the galaxy than they can see.

Gravitational lensing occurs when light is bent from a very distant bright source around a super-massive object. To explain strong gravitational lensing, more matter than is observed must be present. With so little to go on, astronomers don't really know much about the nature of dark matter. One possibility is that it could just be ordinary matter that does not emit radiation in objects such as black holes, neutron stars, and brown dwarfs, objects larger than Jupiter but smaller than the smallest stars. But astronomers cannot find enough of these types of objects, which they have named MACHOS (massive astrophysical compact halo object), to account for all the dark matter, so they are thought to be only a small part of the total.

Another possibility is that the dark matter is thought to be much different from the ordinary matter we see. Some appear to be particles that have gravity, but don't otherwise appear to interact with other particles. Scientists call these theoretical particles WIMPs, which stands for Weakly Interactive Massive Particles.

Most scientists who study dark matter think that the dark matter in the universe is a combination of MACHOS and some type of exotic matter such as WIMPs. Researching dark matter is an active area of scientific research, and astronomers' knowledge about dark matter is changing rapidly.

DARK ENERGY

Astronomers who study the expansion of the universe are interested in knowing the rate of that expansion. Is the rate fast enough to overcome the attractive pull of gravity?

- If yes, then the universe will expand forever, although the expansion will slow down over time.
- If no, then the universe would someday start to contract, and eventually get squeezed together in a big crunch, the opposite of the Big Bang.

Recently astronomers have made a discovery that answers that question: the rate at which the universe is expanding is actually increasing. In other words, the universe is expanding faster now than ever before, and in the future it will expand even faster. So now astronomers think that the universe will keep expanding forever. But it also proposes a perplexing new question: What is causing the expansion of the universe to accelerate? One possible hypothesis involves a new, hypothetical form of energy called dark energy. Some scientists think that dark energy makes up as much as 72% of the total energy content of the universe.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=144

Star Systems



Although constellations have stars that usually only appear to be close together, stars may be found in the same portion of space. Stars that are grouped closely together are called **star systems**. Larger groups of hundreds or thousands of stars are called **star clusters**. The image shown here is a famous star cluster classed M45, also known as Pleides, which can be seen with the naked autumn sky.

Although the star humans know best is a single star, many stars—in fact, more than half of the bright stars in our galaxy—are star systems. A system of two stars orbiting each other is a binary star. A system with more than two stars orbiting each other is a multiple star system. The stars in a binary or multiple star system are often so close together that they appear as one and only through a telescope can the pair be distinguished.

Star clusters are divided into two main types, open clusters and globular clusters. **Open clusters** are groups of up to a few thousand stars that are loosely held together by gravity. Pleiades is an open cluster that is also called the Seven Sisters. Open clusters tend to be blue in color and often contain glowing gas and dust. Why do you think that open clusters have these features? Open clusters are made of young stars that formed from the same nebula. The stars may eventually be pulled apart by gravitational attraction to other objects.



Globular clusters are groups of tens to hundreds of thousands of stars held tightly together by gravity. Figure below shows an example of a globular cluster. Globular clusters have a definite, spherical shape and contain mostly reddish stars. The stars are closer together, closer to the center of the cluster. Globular clusters don't have much dust in them — the dust has already formed into stars.

To view more amazing images taken by the Hubble telescope, click here. If you would like to download free eBooks of Hubble imagery, click here.

Types of Galaxies

Galaxies are the biggest groups of stars and can contain anywhere from a few million stars to many billions of stars. Every star that is visible in the night sky is part of the Milky Way Galaxy. To the naked eye the closest major galaxy, the Andromeda Galaxy, looks like only a dim, fuzzy spot but that fuzzy spot contains one trillion stars.



SPIRAL GALAXIES

Spiral galaxies spin, so they appear as a rotating disk of stars and dust, with a bulge in the middle, like the Sombrero Galaxy. Several arms spiral outward in the Pinwheel Galaxy and are appropriately called **spiral arms**. Spiral galaxies have lots of gas and dust and lots of young stars.Figure below shows a typical egg-shaped **elliptical galaxy.** The smallest elliptical galaxies are as small as some globular clusters. Giant elliptical galaxies, on the other hand, can contain over a trillion stars. Elliptical galaxies are reddish to yellowish in color because they contain mostly old stars.Most elliptical galaxies contain very little gas and dust because they had already formed. However, some elliptical galaxies contain lots of dust. Why might some elliptical galaxies contain dust?



IRREGULAR AND DWARF GALAXIES

Galaxies that are not clearly elliptical galaxies or spiral galaxies are **irregular galaxies**. Most irregular galaxies were once spiral or elliptical galaxies that were then deformed either by gravitational attraction to a larger galaxy or by a collision with another galaxy.**Dwarf galaxies** are small galaxies containing only a few million to a few billion stars. Dwarf galaxies are the most common type in the universe. However, because they are relatively small and dim, we don't see as many dwarf galaxies from Earth. Most dwarf galaxies are irregular in shape. However, there are also dwarf elliptical galaxies and dwarf spiral galaxiesLook back at the picture of the spiral galaxy, Andromeda. Next to our closest galaxy neighbor are two dwarf elliptical galaxies that are companions to the Andromeda Galaxy. One is a bright sphere to the left of center, and the other is a long ellipse below and to the right of center. Dwarf galaxies are often found near larger galaxies. They sometimes collide with and merge into their larger neighbors.


Milky Way Galaxy



On a dark, clear night, you will see a milky band of light stretching across the sky. This band is the disk of a galaxy, the Milky Way Galaxy, is our galaxy and is made of millions of stars along with a lot of gas and dust.

SHAPE and SIZE


Although it is difficult to know what the shape of the Milky Way Galaxy is because we are inside of it, astronomers have identified it as a typical spiral galaxy containing about 100 billion to 400 billion stars.



Like other spiral galaxies, our galaxy has a disk, a central bulge, and spiral arms. The disk is about 100,000 light-years across and 3,000 light-years thick. Most of the Galaxy's gas, dust, young stars, and open clusters are in the disk. What data and evidence do astronomers find that lets them know that the Milky Way is a spiral galaxy?

1. The shape of the galaxy as we see it.
2. The velocities of stars and gas in the galaxy show a rotational motion.
3. The gases, color, and dust are typical of spiral galaxies.

The central bulge is about 12,000 to 16,000 light-years wide and 6,000 to 10,000 light-years thick. The central bulge contains mostly older stars and globular clusters. Some recent evidence suggests the bulge might not be spherical, but is instead shaped like a bar. The bar might be as long as 27,000 light-years long. The disk and bulge are surrounded by a faint, spherical halo, which also includes old stars and globular clusters. Astronomers have discovered that there is a gigantic black hole at the center of the galaxy.



The Milky Way Galaxy is a significant place. Our solar system, including the Sun, Earth, and all the other planets, is within one of the spiral arms in the disk of the Milky Way Galaxy. Most of the stars we see in the sky are relatively nearby stars that are also in this spiral arm. Earth is about 26,000 light-years from the center of the galaxy, a little more than halfway out from the center of the galaxy to the edge.

Just as Earth orbits the Sun, the Sun and solar system orbit the center of the Galaxy. One orbit of the solar system takes about 225 to 250 million years. The solar system has orbited 20 to 25 times since it formed 4.6 billion years ago. Astronomers have recently found that at the center of the Milky Way, and most other galaxies, is a supermassive black hole, though a black hole cannot be seen.

Banner Image: Sombrero Galaxy (M104) from the Hubble Telescope.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-2-the-universe.html. **License**: *CC BY: Attribution*

This page titled 14.1: The Universe is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

15: Stellar Evolution and Our Sun

15.1: Stars

15.2: Stellar Life Cycle

15.3: The Sun

15.4: The Sun and the Earth-Moon System

15.5: Sun – Earth Relationship- The Seasons

This page titled 15: Stellar Evolution and Our Sun is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

15.1: Stars

Lesson Objectives

- Define constellation.
- Describe the flow of energy in a star.
- Classify stars based on their properties.

Vocabulary

- asterism
- nuclear fusion reaction
- parallax
- star

Introduction

When you look at the sky on a clear night, you can see dozens, perhaps even hundreds, of tiny points of light. Almost every one of these points of light is a **star**, a giant ball of glowing gas at a very, very high temperature. Stars differ in size, temperature, and age, but they all appear to be made up of the same elements and to behave according to the same principles

Constellations

People of many different cultures, including the Greeks, identified patterns of stars in the sky. We call these patterns constellations. **Figure** below shows one of the most easily recognized constellations.



The ancient Greeks thought this group of stars looked like a hunter, so they named it Orion after their mythical hunter. The line of three stars at the center is “Orion’s Belt”.

Why do the patterns in constellations and in groups or clusters of stars, called **asterisms**, stay the same night after night? Although the stars move across the sky, they stay in the same patterns. This is because the apparent nightly motion of the stars is actually caused by the rotation of Earth on its axis. The patterns also shift in the sky with the seasons as Earth revolves around the Sun. As a result, people in a particular location can see different constellations in the winter than in the summer. For example, in the Northern Hemisphere Orion is a prominent constellation in the winter sky, but not in the summer sky. This is the annual traverse of the constellations.

Apparent Versus Real Distances


Although the stars in a constellation appear close together as we see them in our night sky, they are not at all close together out in space. In the constellation Orion, the stars visible to the naked eye are at distances ranging from just 26 light-years (which is relatively close to Earth) to several thousand light-years away.

How Stars Are Classified

The many different colors of stars reflect the star's temperature. In Orion (as shown in the **Figure** above) the bright, red star in the upper left named Betelgeuse (pronounced BET-ul-juice) is not as hot than the blue star in the lower right named Rigel.

Color and Temperature

Think about how the color of a piece of metal changes with temperature. A coil of an electric stove will start out black but with added heat will start to glow a dull red. With more heat the coil turns a brighter red, then orange. At extremely high temperatures the coil will turn yellow-white, or even blue-white (it's hard to imagine a stove coil getting that hot). A star's color is also determined by the temperature of the star's surface. Relatively cool stars are red, warmer stars are orange or yellow, and extremely hot stars are blue or blue-white (**Figure** below).



A Hertzsprung-Russell diagram shows the brightness and color of main sequence stars. The brightness is indicated by luminosity and is higher up the y-axis. The temperature is given in degrees Kelvin and is higher on the left side of the x-axis. How does our Sun fare in terms of brightness and color compared with other stars?

Classifying Stars by Color

Color is the most common way to classify stars. **Table** below shows the classification system. The class of a star is given by a letter. Each letter corresponds to a color, and also to a range of temperatures. Note that these letters don't match the color names; they are left over from an older system that is no longer used.

Classification of Stars By Color and Temperature

Class	Color	Temperature Range	Sample Star
O	Blue	30,000 K or more	Zeta Ophiuchi
B	Blue-white	10,000–30,000 K	Rigel
A	White	7,500–10,000 K	Altair
F	Yellowish-white	6,000–7,500 K	Procyon A
G	Yellow	5,500–6,000 K	Sun
K	Orange	3,500–5,000 K	Epsilon Indi
M	Red	2,000–3,500 K	Betelgeuse, Proxima Centauri

For most stars, surface temperature is also related to size. Bigger stars produce more energy, so their surfaces are hotter. These stars tend toward bluish white. Smaller stars produce less energy. Their surfaces are less hot and so they tend to be yellowish.

Measuring Star Distances

How can you measure the distance of an object that is too far away to measure? Now what if you don't know the size of the object or the size or distance of any other objects like it? That would be very difficult, but that is the problem facing astronomers when they try to measure the distances to stars.


Parallax

Distances to stars that are relatively close to us can be measured using **parallax**. Parallax is an apparent shift in position that takes place when the position of the observer changes.

To see an example of parallax, try holding your finger about 1 foot (30 cm) in front of your eyes. Now, while focusing on your finger, close one eye and then the other. Alternate back and forth between eyes, and pay attention to how your finger appears to move. The shift in position of your finger is an example of parallax. Now try moving your finger closer to your eyes, and repeat the experiment. Do you notice any difference? The closer your finger is to your eyes, the greater the position changes because of parallax.

As **Figure** below shows, astronomers use this same principle to measure the distance to stars. Instead of a finger, they focus on a star, and instead of switching back and forth between eyes, they switch between the biggest possible differences in observing position. To do this, an astronomer first looks at the star from one position and notes where the star is relative to more distant stars. Now where will the astronomer go to make an observation the greatest possible distance from the first observation? In six months, after Earth moves from one side of its orbit around the Sun to the other side, the astronomer looks at the star again. This time parallax causes the star to appear in a different position relative to more distant stars. From the size of this shift, astronomers can calculate the distance to the star.

For more about parallax, visit http://starchild.gsfc.nasa.gov/docs/StarChild/questions/parallax.html.



Parallax is used to measure the distance to stars that are relatively nearby.

A parallax exercise is seen here: http://www.astro.ubc.ca/~scharein/a311/Sim/new-parallax/Parallax.html.

Other Methods

Even with the most precise instruments available, parallax is too small to measure the distance to stars that are more than a few hundred light years away. For these more distant stars, astronomers must use more indirect methods of determining distance. Most of these methods involve determining how bright the star they are looking at really is. For example, if the star has properties similar to the Sun, then it should be about as bright as the Sun. The astronomer compares the observed brightness to the expected brightness.

Lesson Summary

- Constellations and asterisms are apparent patterns of stars in the sky.
- Stars in the same constellation are often not close to each other in space.
- A star generates energy by nuclear fusion reactions in its core.
- The color of a star is determined by its surface temperature.
- Stars are classified by color and temperature: O (blue), B (bluish white), A (white), F (yellowish white), G (yellow), K (orange), and M (red), from hottest to coolest.
- Parallax is an apparent shift in an object's position when the position of the observer changes. Astronomers use parallax to measure the distance to relatively nearby stars.

Review Questions

1. What distinguishes a nebula and a star?
2. What kind of reactions provide a star with energy?
3. Stars are extremely massive. Why don't they collapse under the weight of their own gravity?
4. Of what importance are particle accelerators to scientists?
5. Which has a higher surface temperature: a blue star or a red star?
6. What is a light year?
7. Why don't astronomers use parallax to measure the distance to stars that are very far away?

Further Reading / Supplemental Links

- Myths and history of constellations: http://www.ianridpath.com/startales/contents.htm
- NASA World Book, Stars: http://www.nasa.gov/worldbook/star_worldbook.html
- NASA, parts of a star: http://imagine.gsfc.nasa.gov/docs/science/know_l1/stars.html

Points to Consider

- Although stars may appear to be close together in constellations, they are usually not close together out in space. Can you think of any groups of astronomical objects that are relatively close together in space?
- Most nebulas contain more mass than a single star. If a large nebula collapsed into several different stars, what would the result be like?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/26.1/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 15.1: Stars is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

15.2: Stellar Life Cycle


Stellar evolution is the process by which a star undergoes a sequence of radical changes during its lifetime. Depending on the mass of the star, this lifetime ranges from only a few million years for the most massive to trillions of years for the least massive, which is considerably longer than the age of the universe. The table shows the lifetimes of stars as a function of their masses. [1] **All stars are born from collapsing clouds of gas and dust, often called nebulae or molecular clouds.** Over the course of millions of years, these **protostars** settle down into a state of equilibrium, becoming what is known as a **main-sequence star**.

Nuclear fusion powers a star for most of its life. Initially the energy is generated by the fusion of hydrogen atoms at the core of the main-sequence star. Later, as the preponderance of atoms at the core becomes helium, stars like the Sun begin to fuse hydrogen along a spherical shell surrounding the core. This process causes the star to gradually grow in size, passing through the subgiant stage until it reaches the red giant phase. Stars with at least half the mass of the Sun can also begin to generate energy through the fusion of helium at their core, whereas more massive stars can fuse heavier elements along a series of concentric shells. Once a star like the Sun has exhausted its nuclear fuel, its core collapses into a dense white dwarf and the outer layers are expelled as a planetary nebula. Stars with around ten or more times the mass of the Sun can explode in a supernova as their inert iron cores collapse into an extremely dense neutron star or black hole. Although the universe is not old enough for any of the smallest red dwarfs to have reached the end of their lives, stellar models suggest they will slowly become brighter and hotter before running out of hydrogen fuel and becoming low-mass white dwarfs.[2]


Stellar evolution is not studied by observing the life of a single star, as most stellar changes occur too slowly to be detected, even over many centuries. Instead, astrophysicists come to understand how stars evolve by observing numerous stars at various points in their lifetime, and by simulating stellar structure using computer models.

Mass (solar masses)	Time (years)	Spectral type
60	3 million	O3
30	11 million	O7
10	32 million	B4
3	370 million	A5
1.5	3 billion	F5
1	10 billion	G2 (Sun)
0.1	1000s billions	M7

Representative lifetimes of stars as a function of their masses




The life cycle of a Sun-like star.



Artist's depiction of the life cycle of a Sun-like star, starting as a main-sequence star at lower left then expanding through the subgiant and giant phases, until its outer envelope is expelled to form a planetary nebula at upper right.

Birth of a star



Schematic of stellar evolution.

Protostar

Stellar evolution begins with the gravitational collapse of a giant molecular cloud. Typical giant molecular clouds are roughly 100 light-years (9.5×10^{14} km) across and contain up to 6,000,000 solar masses (1.2×10^{37} kg). As it collapses, a giant molecular cloud breaks into smaller and smaller pieces. In each of these fragments, the collapsing gas releases gravitational potential energy as heat. As its temperature and pressure increase, a fragment condenses into a rotating sphere of superhot gas known as a protostar.[3]


The further development heavily depends on the mass of the evolving protostar; in the following, The protostar mass is compared to the mass of the Sun: 1.0 $M_\odot$ (2.0×10^{30} kg) means 1 solar mass.

Protostars are encompassed in dust, and are thus more readily visible at infrared wavelengths. Observations from the Wide-field Infrared Survey Explorer (WISE) have been especially important for unveiling numerous Galactic protostars and their parent star clusters.[4][5]

Brown dwarfs and sub-stellar objects

Protostars with masses less than roughly 0.08 $M_\odot$ (1.6×10^{29} kg) never reach temperatures high enough for nuclear fusion of hydrogen to begin. These are known as brown dwarfs. The International Astronomical Union defines brown dwarfs as stars massive enough to fuse deuterium at some point in their lives (13 Jupiter masses, 2.5×10^{28} kg, or 0.0125 solar masses). Objects smaller than 13 Jupiter masses are classified as sub-brown dwarfs (but if they orbit around another stellar object they are classified as planets).[6] Both types, deuterium-burning and not, shine dimly and die away slowly, cooling gradually over hundreds of millions of years.


Hydrogen fusion



A dense starfield in Sagittarius

For a more massive protostar, the core temperature will eventually reach 10 million kelvin, initiating the proton-proton chain reaction and allowing hydrogen to fuse, first to deuterium and then to helium. In stars of slightly over 1 $M_\odot$ (2.0×10^{30} kg), the carbon–nitrogen–oxygen fusion reaction (CNO cycle) contributes a large portion of the energy generation. The onset of nuclear fusion leads relatively quickly to a hydrostatic equilibrium in which energy released by the core exerts a “radiation pressure” balancing the weight of the star’s matter, preventing further gravitational collapse. The star thus evolves rapidly to a stable state, beginning the main-sequence phase of its evolution.

A new star will sit at a specific point on the main sequence of the Hertzsprung–Russell diagram, with the main-sequence spectral type depending upon the mass of the star. Small, relatively cold, low-mass red dwarfs fuse hydrogen slowly and will remain on the main sequence for hundreds of billions of years or longer, whereas massive, hot supergiants will leave the main sequence after just a few million years. A mid-sized star like the Sun will remain on the main sequence for about 10 billion years. The Sun is thought to be in the middle of its lifespan; thus, it is currently on the main sequence.



The graph above illustrates the evolutionary tracks of stars with different initial masses on the Hertzsprung–Russell diagram. The tracks start once the star has evolved to the main sequence and stop when fusion stops.
A yellow track is shown for the Sun, which will become a red giant after its main-sequence phase ends before expanding further along the asymptotic giant branch, which will be the last phase in which the Sun undergoes fusion.

Mature stars

Eventually the core exhausts its supply of hydrogen and the star begins to evolve off of the main sequence. Without the outward pressure generated by the fusion of hydrogen to counteract the force of gravity the core contracts until either electron degeneracy becomes sufficient to oppose gravity or the core becomes hot enough (around 100 MK) for helium fusion to begin. Which of these happens first depends upon the star's mass.


Low-mass stars

What happens after a low-mass star ceases to produce energy through fusion has not been directly observed; the universe is thought to be around 13.8 billion years old, which is less time (by several orders of magnitude, in some cases) than it takes for fusion to cease in such stars.

Recent astrophysical models suggest that red dwarfs of 0.1 solar mass may stay on the main sequence for some six to twelve trillion years, gradually increasing in both temperature and luminosity, and take several hundred billion more to slowly collapse into a white dwarf.[7][8]Such stars are fully convective and will not develop a degenerate helium core with hydrogen burning shells, or at least not until almost the whole star is helium, so they don't ever expand into a red giant.

Slightly more massive stars do expand into red giants, but their helium cores are not massive enough to ever reach the temperatures required for helium fusion so they never reach the tip of the red giant branch. When hydrogen shell burning finishes, these stars move directly off the red giant branch like a post AGB star, but at lower luminosity, to become a white dwarf.[2] A star of about 0.5 solar mass will be able to reach temperatures high enough to fuse helium, and these "mid-sized" stars go on to further stages of evolution beyond the red giant branch.

Mid-sized stars



Internal structures of main-sequence stars, convection zones with arrowed cycles and radiative zones with red flashes. To the left a **low-mass** red dwarf, in the center a **mid-sized** yellow dwarf and at the right a **massive** blue-white main-sequence star.



The Cat's Eye Nebula, a planetary nebula formed by the death of a star with about the same mass as the Sun


Stars of roughly 0.5–10 solar masses become red giants, which are large non-main-sequence stars of stellar classification K or M. Red giants lie along the right edge of the Hertzsprung–Russell diagram due to their red color and large luminosity. Examples include Aldebaran in the constellation Taurus and Arcturus in the constellation of Boötes. Red giants all have inert cores with hydrogen-burning shells: concentric layers atop the core that are still fusing hydrogen into helium.

Mid-sized stars are red giants during two different phases of their post-main-sequence evolution: red-giant-branch stars, whose inert cores are made of helium, and asymptotic-giant-branch stars, whose inert cores are made of carbon. Asymptotic-giant-branch stars have helium-burning shells inside the hydrogen-burning shells, whereas red-giant-branch stars have hydrogen-burning shells only.[9] In either case, the accelerated fusion in the hydrogen-containing layer immediately over the core causes the star to expand. This lifts the outer layers away from the core, reducing the gravitational pull on them, and they expand faster than the energy production increases. This causes the outer layers of the star to cool, which causes the star to become redder than it was on the main sequence.

Red-giant-branch phase

The red-giant-branch phase of a star's life follows the main sequence. Initially, the cores of red-giant-branch stars collapse, as the internal pressure of the core is insufficient to balance gravity. This gravitational collapse releases energy, heating concentric shells immediately outside the inert helium core so that hydrogen fusion continues in these shells. The core of a red-giant-branch star of up to a few solar masses stops collapsing when it is dense enough to be supported by electron degeneracy pressure. Once this occurs, the core reaches hydrostatic equilibrium: the electron degeneracy pressure is sufficient to balance gravitational pressure.[10] The core's gravity compresses the hydrogen in the layer immediately above it, causing it to fuse faster than hydrogen would fuse in a main-sequence star of the same mass. This in turn causes the star to become more luminous (from 1,000–10,000 times brighter) and expand; the degree of expansion outstrips the increase in luminosity, causing the effective temperature to decrease.

The expanding outer layers of the star are convective, with the material being mixed by turbulence from near the fusing regions up to the surface of the star. For all but the lowest-mass stars, the fused material has remained deep in the stellar interior prior to this point, so the convecting envelope makes fusion products visible at the star's surface for the first time. At this stage of evolution, the results are subtle, with the largest effects, alterations to the isotopes of hydrogen and helium, being unobservable. The effects of the CNO cycle appear at the surface, with lower $^{12}C/^{13}C$ ratios and altered proportions of carbon and nitrogen. These are detectable with spectroscopy and have been measured for many evolved stars.



The Evolution of a Sun-like star

As the hydrogen around the core is consumed, the core absorbs the resulting helium, causing it to contract further, which in turn causes the remaining hydrogen to fuse even faster. This eventually leads to ignition of helium fusion (which includes the triple-alpha process) in the core. In stars of more than approximately solar mass, it can take a billion years or more for the core to reach helium ignition temperatures.

When the temperature and pressure in the core become sufficient to ignite helium fusion, a helium flash will occur if the core is largely supported by electron degeneracy pressure (stars under 1.4 solar mass). In more massive stars, the ignition of helium fusion occurs relatively quietly. Even if a helium flash does occur, the time of very rapid energy release (on the order of 10^8 Suns) is brief, so that the visible outer layers of the star are relatively undisturbed.[11] The energy released by helium fusion causes the core to expand, so that hydrogen fusion in the overlying layers slows and total energy generation decreases. The star contracts, although not all the way to the main sequence, and it migrates to the horizontal branch on the Hertzsprung–Russell diagram, gradually shrinking in radius and increasing its surface temperature. Core helium flash stars evolve to the red end of the horizontal branch but do not migrate to higher temperatures before they gain a degenerate carbon-oxygen core and start helium shell burning. These stars are often observed as a red clump of stars in the colour-magnitude diagram of a cluster, hotter and less luminous than the red giants. Higher-mass stars with larger helium cores move along the horizontal branch to higher temperatures, some becoming unstable pulsating stars in the yellow instability strip (RR Lyrae variables), whereas some become even hotter and can form a blue tail or blue hook to the horizontal branch. The exact morphology of the horizontal branch depends on parameters such as metallicity, age, and helium content, but the exact details are still being modelled.[12]

Asymptotic-giant-branch phase

After a star has consumed the helium at the core, fusion continues in a shell around a hot core of carbon and oxygen. The star follows the asymptotic giant branch on the Hertzsprung–Russell diagram, paralleling the original red giant evolution, but with even faster energy generation (which lasts for a shorter time).[13] Although helium is being burnt in a shell, the majority of the energy is produced by hydrogen burning in a shell closer to the surface of the star. Helium from these hydrogen burning shells drops towards the center of the star and periodically the energy output from the helium shell increases dramatically. This is known as a thermal pulse and they occur towards the end of the asymptotic-giant-branch phase, sometimes even into the post-asymptotic-giant-branch phase. Depending on mass and composition, there may be several to hundreds of thermal pulses.

There is a phase on the ascent of the asymptotic-giant-branch where a deep convective zone forms and can bring carbon from the core to the surface, This is known as the second dredge up, and in some stars there may even be a third dredge up. In this way a carbon star is formed, very cool and strongly reddened stars showing strong carbon lines in their spectra. A process known as hot bottom burning may convert carbon into oxygen and nitrogen before it can be dredged to the surface, and the interaction between these processes determines the observed luminosities and spectra of carbon stars in particular clusters.[14]

Another well known class of asymptotic-giant-branch stars are the Mira variables, which pulsate with well-defined periods of tens to hundreds of days and large amplitudes up to about 10 magnitudes (in the visual, total luminosity changes by a much smaller

amount). In more massive stars the stars become more luminous and the pulsation period is longer, leading to enhanced mass loss, and the stars become heavily obscured at visual wavelengths. These stars can be observed as OH/IR stars, pulsating in the infra-red and showing OH maser activity. These stars are clearly oxygen rich, in contrast to the carbon stars, but both must be produced by dredge ups.

These mid-range stars ultimately reach the tip of the asymptotic-giant-branch and run out of fuel for shell burning. They are not sufficiently massive to start full-scale carbon fusion, so they contract again, going through a period of post-asymptotic-giant-branch superwind to produce a planetary nebula with an extremely hot central star. The central star then cools to a white dwarf. The expelled gas is relatively rich in heavy elements created within the star and may be particularly oxygen or carbon enriched, depending on the type of the star. The gas builds up in an expanding shell called a circumstellar envelope and cools as it moves away from the star, allowing dust particles and molecules to form. With the high infrared energy input from the central star, ideal conditions are formed in these circumstellar envelopes for maser excitation.

It is possible for thermal pulses to be produced once post-asymptotic-giant-branch evolution has begun, producing a variety of unusual and poorly understood stars known as born-again asymptotic-giant-branch stars.[15] These may result in extreme horizontal-branch stars (subdwarf B stars), hydrogen deficient post-asymptotic-giant-branch stars, variable planetary nebula central stars, and R Coronae Borealis variables.

Massive stars




The Crab Nebula, the shattered remnants of a star which exploded as a supernova, the light of which reached Earth in 1054 AD

In massive stars, the core is already large enough at the onset of the hydrogen burning shell that helium ignition will occur before electron degeneracy pressure has a chance to become prevalent. Thus, when these stars expand and cool, they do not brighten as much as lower-mass stars; however, they were much brighter than lower-mass stars to begin with, and are thus still brighter than the red giants formed from less massive stars. These stars are unlikely to survive as red supergiants; instead they will destroy themselves as type II supernovas.

Extremely massive stars (more than approximately 40 solar masses), which are very luminous and thus have very rapid stellar winds, lose mass so rapidly due to radiation pressure that they tend to strip off their own envelopes before they can expand to become red supergiants, and thus retain extremely high surface temperatures (and blue-white color) from their main-sequence time onwards. The largest stars of the current generation are about 100-150 solar masses because the outer layers would be expelled by the extreme radiation. Although lower-mass stars normally do not burn off their outer layers so rapidly, they can likewise avoid becoming red giants or red supergiants if they are in binary systems close enough so that the companion star strips off the envelope as it expands, or if they rotate rapidly enough so that convection extends all the way from the core to the surface, resulting in the absence of a separate core and envelope due to thorough mixing.[16]

The core grows hotter and denser as it gains material from fusion of hydrogen at the base of the envelope. In all massive stars, electron degeneracy pressure is insufficient to halt collapse by itself, so as each major element is consumed in the center, progressively heavier elements ignite, temporarily halting collapse. If the core of the star is not too massive (less than approximately 1.4 solar mass, taking into account mass loss that has occurred by this time), it may then form a white dwarf

(possibly surrounded by a planetary nebula) as described above for less massive stars, with the difference that the white dwarf is composed chiefly of oxygen, neon, and magnesium.



The onion-like layers of a massive, evolved star just before core collapse. (Not to scale.)

Above a certain mass (estimated at approximately 2.5 solar masses and whose star's progenitor was around 10 solar masses), the core will reach the temperature (approximately 1.1 gigakelvins) at which neon partially breaks down to form oxygen and helium, the latter of which immediately fuses with some of the remaining neon to form magnesium; then oxygen fuses to form sulfur, silicon, and smaller amounts of other elements. Finally, the temperature gets high enough that any nucleus can be partially broken down, most commonly releasing an alpha particle (helium nucleus) which immediately fuses with another nucleus, so that several nuclei are effectively rearranged into a smaller number of heavier nuclei, with net release of energy because the addition of fragments to nuclei exceeds the energy required to break them off the parent nuclei.

A star with a core mass too great to form a white dwarf but insufficient to achieve sustained conversion of neon to oxygen and magnesium, will undergo core collapse (due to electron capture) before achieving fusion of the heavier elements.[17] Both heating and cooling caused by electron capture onto minor constituent elements (such as aluminum and sodium) prior to collapse may have a significant impact on total energy generation within the star shortly before collapse.[18] This may produce a noticeable effect on the abundance of elements and isotopes ejected in the subsequent supernova.

Supernova

Once the nucleosynthesis process arrives at iron-56, the continuation of this process consumes energy (the addition of fragments to nuclei releases less energy than required to break them off the parent nuclei). If the mass of the core exceeds the Chandrasekhar limit, electron degeneracy pressure will be unable to support its weight against the force of gravity, and the core will undergo sudden, catastrophic collapse to form a neutron star or (in the case of cores that exceed the Tolman-Oppenheimer-Volkoff limit), a black hole. Through a process that is not completely understood, some of the gravitational potential energy released by this core collapse is converted into a Type Ib, Type Ic, or Type II supernova. It is known that the core collapse produces a massive surge of neutrinos, as observed with supernova SN 1987A. The extremely energetic neutrinos fragment some nuclei; some of their energy is consumed in releasing nucleons, including neutrons, and some of their energy is transformed into heat and kinetic energy, thus augmenting the shock wave started by rebound of some of the infalling material from the collapse of the core. Electron capture in very dense parts of the infalling matter may produce additional neutrons. Because some of the rebounding matter is bombarded by the neutrons, some of its nuclei capture them, creating a spectrum of heavier-than-iron material including the radioactive elements up to (and likely beyond) uranium.[19] Although non-exploding red giants can produce significant quantities of elements heavier than iron using neutrons released in side reactions of earlier nuclear reactions, the abundance of elements heavier than iron (and in particular, of certain isotopes of elements that have multiple stable or long-lived isotopes) produced in such reactions is quite different from that produced in a supernova. Neither abundance alone matches that found in the Solar System, so both supernovae and ejection of elements from red giants are required to explain the observed abundance of heavy elements and isotopes there of.

The energy transferred from collapse of the core to rebounding material not only generates heavy elements, but provides for their acceleration well beyond escape velocity, thus causing a Type Ib, Type Ic, or Type II supernova. Note that current understanding of this energy transfer is still not satisfactory; although current computer models of Type Ib, Type Ic, and Type II supernovae account for part of the energy transfer, they are not able to account for enough energy transfer to produce the observed ejection of material.

[20] Some evidence gained from analysis of the mass and orbital parameters of binary neutron stars (which require two such supernovae) hints that the collapse of an oxygen-neon-magnesium core may produce a supernova that differs observably (in ways other than size) from a supernova produced by the collapse of an iron core.[21]

The most massive stars that exist today may be completely destroyed by a supernova with an energy greatly exceeding its gravitational binding energy. This rare event, caused by pair-instability, leaves behind no black hole remnant.[22] In the past history of the universe, some stars were even larger than the largest that exists today, and they would immediately collapse into a black hole at the end of their lives, due to photodisintegration.

Stellar remnants

After a star has burned out its fuel supply, its remnants can take one of three forms, depending on the mass during its lifetime.

White and black dwarfs

For a star of 1 solar mass, the resulting white dwarf is of about 0.6 solar mass, compressed into approximately the volume of the Earth. White dwarfs are stable because the inward pull of gravity is balanced by the degeneracy pressure of the star's electrons, a consequence of the Pauli exclusion principle. Electron degeneracy pressure provides a rather soft limit against further compression; therefore, for a given chemical composition, white dwarfs of higher mass have a smaller volume. With no fuel left to burn, the star radiates its remaining heat into space for billions of years.

A white dwarf is very hot when it first forms, more than 100,000 K at the surface and even hotter in its interior. It is so hot that a lot of its energy is lost in the form of neutrinos for the first 10 million years of its existence, but will have lost most of its energy after a billion years.[23]

The chemical composition of the white dwarf depends upon its mass. A star of a few solar masses will ignite carbon fusion to form magnesium, neon, and smaller amounts of other elements, resulting in a white dwarf composed chiefly of oxygen, neon, and magnesium, provided that it can lose enough mass to get below the Chandrasekhar limit (see below), and provided that the ignition of carbon is not so violent as to blow the star apart in a supernova.[24] A star of mass on the order of magnitude of the Sun will be unable to ignite carbon fusion, and will produce a white dwarf composed chiefly of carbon and oxygen, and of mass too low to collapse unless matter is added to it later (see below). A star of less than about half the mass of the Sun will be unable to ignite helium fusion (as noted earlier), and will produce a white dwarf composed chiefly of helium.

In the end, all that remains is a cold dark mass sometimes called a black dwarf. However, the universe is not old enough for any black dwarfs to exist yet.

If the white dwarf's mass increases above the Chandrasekhar limit, which is 1.4 solar mass for a white dwarf composed chiefly of carbon, oxygen, neon, and/or magnesium, then electron degeneracy pressure fails due to electron capture and the star collapses. Depending upon the chemical composition and pre-collapse temperature in the center, this will lead either to collapse into a neutron star or runaway ignition of carbon and oxygen. Heavier elements favor continued core collapse, because they require a higher temperature to ignite, because electron capture onto these elements and their fusion products is easier; higher core temperatures favor runaway nuclear reaction, which halts core collapse and leads to a Type Ia supernova.[25] These supernovae may be many times brighter than the Type II supernova marking the death of a massive star, even though the latter has the greater total energy release. This inability to collapse means that no white dwarf more massive than approximately 1.4 solar mass can exist (with a possible minor exception for very rapidly spinning white dwarfs, whose centrifugal force due to rotation partially counteracts the weight of their matter). Mass transfer in a binary system may cause an initially stable white dwarf to surpass the Chandrasekhar limit.

If a white dwarf forms a close binary system with another star, hydrogen from the larger companion may accrete around and onto a white dwarf until it gets hot enough to fuse in a runaway reaction at its surface, although the white dwarf remains below the Chandrasekhar limit. Such an explosion is termed a nova.

Neutron stars



Bubble-like shock wave still expanding from a supernova explosion 15,000 years ago.

When a stellar core collapses, the pressure causes electron capture, thus converting the great majority of the protons into neutrons. The electromagnetic forces keeping separate nuclei apart are gone (proportionally, if nuclei were the size of dust mites, atoms would be as large as football stadiums), and most of the core of the star becomes a dense ball of contiguous neutrons (in some ways like a giant atomic nucleus), with a thin overlying layer of degenerate matter (chiefly iron unless matter of different composition is added later). The neutrons resist further compression by the Pauli Exclusion Principle, in a way analogous to electron degeneracy pressure, but stronger.

These stars, known as neutron stars, are extremely small—on the order of radius 10 km, no bigger than the size of a large city—and are phenomenally dense. Their period of rotation shortens dramatically as the stars shrink (due to conservation of angular momentum); observed rotational periods of neutron stars range from about 1.5 milliseconds (over 600 revolutions per second) to several seconds.[26] When these rapidly rotating stars' magnetic poles are aligned with the Earth, we detect a pulse of radiation each revolution. Such neutron stars are called pulsars, and were the first neutron stars to be discovered. Though electromagnetic radiation detected from pulsars is most often in the form of radio waves, pulsars have also been detected at visible, X-ray, and gamma ray wavelengths.[27]

Black holes

If the mass of the stellar remnant is high enough, the neutron degeneracy pressure will be insufficient to prevent collapse below the Schwarzschild radius. The stellar remnant thus becomes a black hole. The mass at which this occurs is not known with certainty, but is currently estimated at between 2 and 3 solar masses.

Black holes are predicted by the theory of general relativity. According to classical general relativity, no matter or information can flow from the interior of a black hole to an outside observer, although quantum effects may allow deviations from this strict rule. The existence of black holes in the universe is well supported, both theoretically and by astronomical observation.

Because the core-collapse supernova mechanism itself is imperfectly understood, it is still not known whether it is possible for a star to collapse directly to a black hole without producing a visible supernova, or whether some supernovae initially form unstable neutron stars which then collapse into black holes; the exact relation between the initial mass of the star and the final remnant is also not completely certain. Resolution of these uncertainties requires the analysis of more supernovae and supernova remnants.

Models

A stellar evolutionary model is a mathematical model that can be used to compute the evolutionary phases of a star from its formation until it becomes a remnant. The mass and chemical composition of the star are used as the inputs, and the luminosity and surface temperature are the only constraints. The model formulae are based upon the physical understanding of the star, usually under the assumption of hydrostatic equilibrium. Extensive computer calculations are then run to determine the changing state of the star over time, yielding a table of data that can be used to determine the evolutionary track of the star across the Hertzsprung–Russell diagram, along with other evolving properties.[28] Accurate models can be used to estimate the current age of a star by comparing its physical properties with those of stars along a matching evolutionary track.[29]

Further reading

- Astronomy 606 (Stellar Structure and Evolution) lecture notes, Cole Miller, Department of Astronomy, University of Maryland
- Astronomy 162, Unit 2 (The Structure & Evolution of Stars) lecture notes, Richard W. Pogge, Department of Astronomy, Ohio State University
- *Hansen, Carl J.; Kawaler, Steven D.; Trimble, Virginia (2004). Stellar interiors: physical principles, structure, and evolution (2nd ed.). Springer-Verlag. ISBN 0-387-20089-4*

External links

- Stellar evolution simulator

References

1. ^ Bertulani, Carlos A. (2013). *Nuclei in the Cosmos*. World Scientific. ISBN 978-981-4417-66-2.
2. ^ *a b* The End of the Main Sequence, Gregory Laughlin, Peter Bodenheimer, and Fred C. Adams, *The Astrophysical Journal*,**482** (June 10, 1997), pp. 420–432. Bibcode: 1997ApJ…482..420L. doi:10.1086/304125.
3. ^ Dina Prialnik (2000). *An Introduction to the Theory of Stellar Structure and Evolution*. Cambridge University Press. chapter 10. ISBN 0-521-65065-8.
4. ^ "Wide-field Infrared Survey Explorer Mission". NASA.
5. ^ Majaess, D. (2013). *Discovering protostars and their host clusters via WISE*, ApSS, 344, 1 (*VizieR catalog*)
6. ^ "Working Group on Extrasolar Planets: Definition of a "Planet"". *IAU position statement*. 2003-02-28. Retrieved 2012-05-30. [*dead link*]
7. ^ "Why the Smallest Stars Stay Small". *Sky & Telescope* (22). November 1997.
8. ^ Adams, F. C.; P. Bodenheimer, G. Laughlin (2005). "M dwarfs: planet formation and long term evolution".*Astronomische Nachrichten* **326** (10): 913–919. Bibcode:2005AN….326..913A. doi:10.1002/asna.200510440.
9. ^ Hansen, Kawaler & Trimble (2004, pp. 55–56)
10. ^ Hansen, Kawaler & Trimble (2004, pp. 62–63)
11. ^ Alan C. Edwards (1969). "The hydrodynamics of the helium flash". *Monthly Notices of the Royal Astronomical Society***146**: 445–472. Bibcode:1969MNRAS.146..445E.
12. ^ Gratton, R. G.; Carretta, E.; Bragaglia, A.; Lucatello, S.; d'Orazi, V. (2010). "The second and third parameters of the horizontal branch in globular clusters". *Astronomy and Astrophysics* **517**: A81. arXiv:1004.3862. Bibcode:2010A&A…517A..81G. doi:10.1051/0004-6361/200912572. edit
13. ^ Sackmann, I. -J.; Boothroyd, A. I.; Kraemer, K. E. (1993). "Our Sun. III. Present and Future". *The Astrophysical Journal***418**: 457. Bibcode:1993ApJ…418..457S. doi:10.1086/173407.edit
14. ^ van Loon; Zijlstra; Whitelock; Peter te Lintel Hekkert; Chapman; Cecile Loup; Groenewegen; Waters; Trams (1998). "Obscured Asymptotic Giant Branch stars in the Magellanic Clouds IV. Carbon stars and OH/IR stars". *Astronomy and Astrophysics* **329** (169): 32. arXiv:astro-ph/9709119v1. Bibcode:1996MNRAS.279…32Z.
15. ^ Bibcode: 1991IAUS..145..363H
16. ^ D. Vanbeveren; De Loore, C.; Van Rensbergen, W. (1998)."Massive stars". *The Astronomy and Astrophysics Review* **9**(1–2): 63–152. Bibcode:1998A&ARv…9…63V. doi:10.1007/s001590050015. Archived from the original on 2009-03-27.
17. ^ Ken'ichi Nomoto (1987). "Evolution of 8–10 solar mass stars toward electron capture supernovae. II – Collapse of an O + Ne + Mg core". *Astrophysical Journal*. 322. Part 1: 206–214. Bibcode:1987ApJ…322..206N. doi:10.1086/165716.
18. ^ Claudio Ritossa et al. (1999). "On the Evolution of Stars that Form Electron-degenerate Cores Processed by Carbon Burning. V. Shell Convection Sustained by Helium Burning, Transient Neon Burning, Dredge-out, URCA Cooling, and Other Properties of an 11 M_solar Population I Model Star".*The Astrophysical Journal* **515** (1): 381–397. Bibcode:1999ApJ…515..381R. doi:10.1086/307017.
19. ^ How do Massive Stars Explode?
20. ^ Robert Buras, et al. (June 2003). "Supernova Simulations Still Defy Explosions". *Research Highlights*. Max-Planck-Institut für Astrophysik.
21. ^ E. P. J. van den Heuvel (2004). "X-Ray Binaries and Their Descendants: Binary Radio Pulsars; Evidence for Three Classes of Neutron Stars?". *Proceedings of the 5th INTEGRAL Workshop on the INTEGRAL Universe (ESA SP-552)* **552**: 185–194. arXiv:astro-ph/0407451. Bibcode:2004inun.conf..185V.

22. ^ Pair Instability Supernovae and Hypernovae.[*dead link*], Nicolay J. Hammer, (2003), accessed May 7, 2007.
23. ^ Fossil Stars (1): White Dwarfs
24. ^ Ken'ichi Nomoto (1984). "Evolution of 8–10 solar mass stars toward electron capture supernovae. I – Formation of electron-degenerate O + Ne + Mg cores". *Astrophysical Journal*. Part 1 **277**: 791–805. Bibcode:1984ApJ…277..791N. doi:10.1086/161749.
25. ^ Ken'ichi Nomoto and Yoji Kondo (1991). "Conditions for accretion-induced collapse of white dwarfs". *Astrophysical Journal*. 367. Part 2: L19–L22. Bibcode:1991ApJ…367L..19N. doi:10.1086/185922.
26. ^ D'Amico, N.; Stappers, B. W.; Bailes, M.; Martin, C. E.; Bell, J. F.; Lyne, A. G.; Manchester, R. N (1998). "The Parkes Southern Pulsar Survey – III. Timing of long-period pulsars".*Monthly Notices of the Royal Astronomical Society* **297**: 28. Bibcode:1998MNRAS.297…28D. doi:10.1046/j.1365-8711.1998.01397.x.
27. ^ Courtland, Rachel (17 October 2008). "Pulsar Detected by Gamma Waves Only". *New Scientist*.[*dead link*]
28. ^ Demarque, P.; Guenther, D. B.; Li, L. H.; Mazumdar, A.; Straka, C. W. (August 2008). "YREC: the Yale rotating stellar evolution code". *Astrophysics and Space Science* **316** (1–4): 31–41. arXiv:0710.4003. Bibcode:2008Ap&SS.316…31D. doi:10.1007/s10509-007-9698-y. ISBN 9781402094408.
29. ^ Ryan, Seán; Norton, Andrew J. (2010). "Assigning ages from hydrogen-burning timescales". *Stellar Evolution and Nucleosynthesis*. Cambridge University Press. p. 79. ISBN 0-521-13320-3.

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.Wikipedia.org/wiki/Stellar_life_cycle. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 15.2: Stellar Life Cycle is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

15.3: The Sun

Consider the Earth, the Moon, and all the other planets in our solar system. Think about the mass that all those objects must have when they are all added together. Added all together, however, they account for only 0.2% of the total mass of the solar system. The Sun makes up the remaining 99.8% of all the mass in the solar system (Figure 24.17)! The Sun is the center of the solar system and the largest object in the solar system. Our Sun is a star that provides light and heat and supports almost all life on Earth.

In this lesson you will learn about the features of the Sun. We will discuss the composition of the Sun, its atmosphere, and some of its surface features.



Figure 24.17: The Sun.

Layers of the Sun

The Sun is a sphere, but unlike the Earth and the Moon, is not solid. Most atoms in the Sun exist as **plasma**, or a fourth state of matter made up of superheated gas with an electrical charge. Our Sun consists almost entirely of the elements hydrogen and helium. Because the Sun is not solid, it does not have a defined outer boundary. It does, however, have a definite internal structure. There are several identifiable layers of the Sun:

The **core** is the innermost or central layer of the Sun. The core is plasma, but moves similarly to a gas. Its temperature is around 27 million degrees Celsius. In the core, nuclear reactions combine hydrogen atoms to form helium, releasing vast amounts of energy in the process. The energy released then begins to move outward, towards the outer layers of the Sun.

The **radiative zone** is just outside the core, which has a temperature of about 7 million degrees Celsius. The energy released in the core travels extremely slowly through the radiative zone. Particles of light called photons can only travel a few millimeters before they hit another particle in the Sun, are absorbed and then released again. It can take a photon as long as 50 million years to travel all the way through the radiative zone.

The **convection zone** surrounds the radiative zone. In the convection zone, hot material from near the Sun's center rises, cools at the surface, and then plunges back downward to receive more heat from the radiative zone. This movement helps to create solar flares and sunspots, which we'll learn more about in a bit. These first three layers make up what we would actually call "the Sun". The next three layers make up the Sun's atmosphere. Of course, there are no solid layers to any part of the Sun, so these boundaries are fuzzy and indistinct.

The Sun's "Atmosphere"

The **photosphere** is the visible surface of the Sun (Figure 24.18). This is the region of the Sun that emits sunlight. It's also one of the coolest layers of the Sun—only about 6700°C. Looking at a photograph of the Sun's surface, you can see that it has several different colors; oranges, yellow and reds, giving it a grainy appearance. We cannot see this when we glance quickly at the Sun.

Our eyes can't focus that quickly and the Sun is too bright for us to look at for more than a brief moment. Looking at the Sun for any length of time can cause blindness, so don't try it! Sunlight is emitted from the Sun's photosphere. A fraction of the light that travels from the Sun reaches Earth. It travels as light in a range of wavelengths, including visible light, ultraviolet, and infrared **radiation**. Visible light is all the light we can see with our eyes. We can't see ultraviolet and infrared radiation, but their effects can still be detected. For example, a sunburn is caused by ultraviolet radiation when you spend too much time in the Sun.




Figure 24.18: The layers of the Sun.

The **chromosphere** is the zone about 2,000 kilometers thick that lies directly above the photosphere. The chromosphere is a thin region of the Sun's atmosphere that glows red as it is heated by energy from the photosphere. Temperatures in the chromosphere range from about 4000°C to about 10,000°C. Jets of gas fire up through the chromosphere at speeds up to 72,000 kilometers per hour, reaching heights as high as 10,000 kilometers.

The **corona** is the outermost layer of the Sun and is the outermost part of its atmosphere. It is the Sun's halo or "crown". It has a temperature of 2 to 5 million degrees Celsius and is much hotter than the visible surface of the Sun, or photosphere. The corona extends millions of kilometers into space. If you ever have the chance to see a total solar eclipse, you will be able to see the Sun's corona, shining out into space.

In the Sun's core, **nuclear fusion** reactions generate energy by converting hydrogen to helium. Fusion is a process where the nuclei of atoms join together to form a heavier chemical element. Fusion reactions in the Sun's core produce energy, which we experience as heat and light. The rest of the Sun is heated by movement of heat energy outward from the core. Light energy from the Sun is emitted from the photosphere. It travels through space, and some of it reaches the Earth. The Sun is the source of almost all the energy on Earth and sunlight powers **photosynthesis**, as well as warming and illuminating our Earth.

Surface Features of the Sun

The most noticeable surface feature of the Sun is the presence of sunspots, which are cooler, darker areas on the Sun's surface. Sunspots are only visible with special light-filtering lenses. They exhibit intense magnetic activity. These areas are cooler and darker because loops of the Sun's magnetic field break through the surface and disrupt the smooth transfer of heat from lower layers. Sunspots usually occur in pairs. When a loop of the Sun's magnetic field breaks through the surface, it usually creates a sunspot both where it comes out and one where it goes back in again. Sunspots usually occur in 11 year cycles, beginning when the number of sunspots is at a minimum, increasing to a maximum number of sunspots and then gradually decreasing to a minimum number of sunspots again.

If a loop of the sun's magnetic field snaps and breaks, it creates solar flares, which are violent explosions that release huge amounts of energy (Figure 24.19). They release streams of highly energetic particles that make up the solar wind. The solar wind can be dangerous to spacecraft and astronauts. It sends out large amounts of radiation, which can harm the human body. Solar flares have knocked out entire power grids and can disturb radio, satellite, and cell phone communications.




Figure 24.19: A solar flare.

Another highly visible feature on the Sun are solar prominences. If plasma flows along a loop of the Sun's magnetic field from sunspot to sunspot, it forms a glowing arch that reaches thousands of kilometers into the Sun's atmosphere. Prominences can last for a day to several months. Prominences are also visible during a total solar eclipse.

A beautiful and mysterious effect of the Sun's electrically charged particles are auroras, which form around the polar regions high in Earth's atmosphere. Gases in Earth's atmosphere are excited by the electrically charged particles of the solar wind and glow producing curtains of light, which bend and change as you watch.

Lesson Summary

- The mass of the Sun is tremendous. It makes up 99.8% of the mass of our solar system.
- The Sun is mostly made of hydrogen with smaller amounts of helium in the form of plasma.
- The main part of the Sun has three layers: the core, the radiative zone, and the convection zone.
- The Sun's atmosphere also has three layers: the photosphere, the chromosphere, and the corona.
- Nuclear fusion of hydrogen in the core of the Sun produces tremendous amounts of energy that radiate out from the Sun.
- Some features of the Sun's surface include sunspots, solar flares, and prominences.

Review Questions

1. In what way does the Sun support all life on Earth?
2. Which two elements make up the Sun almost in entirety?
3. Which process is the source of heat in the Sun and where does it take place?
4. Some scientists would like to plan a trip to take humans to Mars. One of the things standing in the way of our ability to do this is solar wind. Why will we have to be concerned with solar wind?
5. Describe how movements in the convection zone contribute to solar flares.
6. Do you think fusion reactions in the Sun's core will continue forever and go on with no end? Explain your answer.

Vocabulary

chromosphere

Thin layer of the Sun's atmosphere that lies directly above the photosphere; glows red.

convection zone

Layer of the Sun that surrounds the Radiative Zone; energy moves as flowing cells of gas.

core

Innermost or central layer of the Sun.

corona

Outermost layer of the Sun; a plasma that extends millions of kilometers into space.

nuclear fusion

The merging together of the nuclei of atoms to form new, heavier chemical elements; huge amounts of nuclear energy are released in the process.

photosphere

Layer of the Sun that we see; the visible surface of the Sun.

photosynthesis

The process that green plants use to convert sunlight to energy.

plasma

A high energy, high temperature form of matter. Electrons are removed from atoms, leaving each atom with an electrical charge.

radiation

Electromagnetic energy; photons.

radiative zone

Layer of the Sun immediately surrounding the core; energy moves atom to atom as electromagnetic waves.

solar flare

A violent explosion on the Sun's surface.

solar wind

A stream of radiation emitted by a solar flare. The solar wind extends millions of kilometers out into space and can even reach the Earth.

sunspots

Cooler, darker areas on the Sun's surface that have lower temperatures than surrounding areas; sunspots usually occur in pairs.

Points to Consider

- If something were to suddenly cause nuclear fusion to stop in the Sun, how would we know?
- Are there any types of dangerous energy from the Sun? What might be affected by them?
- If the Sun is all made of gases like hydrogen and helium, how can it have layers?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/The_Sun. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 15.3: The Sun is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

15.4: The Sun and the Earth-Moon System

The solar system is made up of the Sun, the planets that orbit the Sun, their satellites, dwarf planets and many, many small objects, like asteroids and comets. All of these objects move and we can see these movements. We notice the Sun rises in the eastern sky in the morning and sets in the western sky in the evening. We observe different stars in the sky at different times of the year. When ancient people made these observations, they imagined that the sky was actually moving while the Earth stood still. In 1543, Nicolaus Copernicus (Figure 24.21) proposed a radically different idea: the Earth and the other planets make regular revolutions around the Sun. He also suggested that the Earth rotates once a day on its axis. Copernicus' idea slowly gained acceptance and today we base our view of motions in the solar system on his work. We also now know that everything in the universe is moving.




Figure 24.21: Nicholas Copernicus.

In this lesson you will learn about how the movements of the Earth, Moon, and Sun affect different phenomena on Earth, including day and night, the seasons, tides, and phases of the Moon.

Lesson Objectives

- Describe how Earth's movements affect seasons and cause day and night.
- Explain solar and lunar eclipses.
- Describe the phases of the Moon and explain why they occur.
- Explain how movements of the Earth and Moon affect Earth's tides.

Positions and Movements

Earlier we discussed Earth's rotation and revolution. The Earth rotates once on its axis about every 24 hours. If you were to look at Earth from the North Pole, it would be spinning counterclockwise. As the Earth rotates, observers on Earth see the Sun moving across the sky from east to west with the beginning of each new day. We often say that the Sun is "rising" or "setting", but actually it is the Earth's rotation that gives us the perception of the Sun rising up or setting over the horizon. When we look at the Moon or the stars at night, they also seem to rise in the east and set in the west. Earth's rotation is also responsible for this. As Earth turns, the Moon and stars change position in our sky.

Earth's Day and Night

Another effect of Earth's rotation is that we have a cycle of daylight and darkness approximately every 24 hours. This is called a day. As Earth rotates, the side of Earth facing the Sun experiences daylight, and the opposite side (facing away from the Sun) experiences darkness or nighttime. Since the Earth completes one rotation in about 24 hours, this is the time it takes to complete one day-night cycle. As the Earth rotates, different places on Earth experience sunset and sunrise at a different time. As you move towards the poles, summer and winter days have different amounts of daylight hours in a day. For example, in the Northern hemisphere, we begin summer on June 21. At this point, the Earth's North Pole is pointed directly toward the Sun. Therefore, areas north of the equator experience longer days and shorter nights because the northern half of the Earth is pointed toward the Sun. Since the southern half of the Earth is pointed away from the Sun at that point, they have the opposite effect—longer nights and shorter days.

For people in the Northern hemisphere, winter begins on December 21. At this point, it is Earth's South Pole that is tilted toward the Sun, and so there are shorter days and longer nights for those who are north of the equator.

Earth's Seasons

It is a common misconception that summer is warm and winter is cold because the Sun is closer to Earth in the summer and farther away from it during the winter. Remember that seasons are caused by the 23.5 degree tilt of Earth's axis of rotation and Earth's yearly revolution around the Sun (Figure 24.22). This results in one part of the Earth being more directly exposed to rays from the Sun than the other part. The part tilted away from the Sun experiences a cool season, while the part tilted toward the Sun experiences a warm season. Seasons change as the Earth continues its revolution, causing the hemisphere tilted away from or towards the Sun to change accordingly. When it is winter in the Northern hemisphere, it is summer in the Southern hemisphere, and vice versa.





Figure 24.22: The Earth's tilt on its axis leads to one hemisphere facing the Sun more than the other hemisphere and gives rise to seasons.

NORTHERN HEMISPHERE SUMMER

The North Pole is tilted towards the Sun and the Sun's rays strike the Northern Hemisphere more directly in summer. At the **summer solstice**, June 21 or 22, the Sun's rays hit the Earth most directly along the Tropic of Cancer (23.5 degrees N); that is, the angle of incidence of the sun's rays there is zero (the angle of incidence is the deviation in the angle of an incoming ray from straight on). When it is summer solstice in the Northern Hemisphere, it is winter solstice in the Southern Hemisphere.

NORTHERN HEMISPHERE WINTER


Winter solstice for the Northern Hemisphere happens on December 21 or 22. The tilt of Earth's axis points away from the Sun. Light from the Sun is spread out over a larger area, so that area isn't heated as much. With fewer daylight hours in winter, there is also less time for the Sun to warm the area. When it is winter in the Northern Hemisphere, it is summer in the Southern Hemisphere.



EQUINOX

Halfway between the two solstices, the Sun's rays shine most directly at the equator, called an "**equinox**." The daylight and nighttime hours are exactly equal on an equinox. The autumnal equinox happens on September 22 or 23 and the vernal or spring equinox happens March 21 or 22 in the Northern Hemisphere.

Solar Eclipses



A **solar eclipse** occurs when the new moon passes directly between the Earth and the Sun (Figure 24.23). This casts a shadow on the Earth and blocks our view of the Sun. A total solar eclipse occurs when the Moon's shadow completely blocks the Sun (Figure 24.24). When only a portion of the Sun is out of view, it is called a partial solar eclipse. Solar eclipses are rare events that usually only last a few minutes. That is because the Moon's shadow only covers a very small area on Earth and Earth is turning very rapidly. As the Sun is covered by the moon's shadow, it will actually get cooler outside. Birds may begin to sing, and stars will become visible in the sky. During a solar eclipse, the corona and solar prominences can be seen.

Figure 24.23: A solar eclipse.




Figure 24.24: Photo of a total solar eclipse. The corona is the white region surrounding the Sun.

A Lunar Eclipse

A **lunar eclipse** occurs when the full moon moves through the shadow of the Earth (Figure 24.25). This can only happen when the Earth is between the Moon and the Sun and all three are lined up in the same plane, called the ecliptic. The ecliptic is the plane of Earth's orbit around the Sun. The Earth's shadow has two distinct parts: the umbra and the penumbra. The **umbra** is the inner, cone shaped part of the shadow, in which all of the light has been blocked. The outer part of Earth's shadow is the **penumbra** where only part of the light is blocked. In the penumbra, the light is dimmed but not totally absent. A total lunar eclipse occurs when the Moon travels completely in Earth's umbra. During a partial lunar eclipse, only a portion of the Moon enters Earth's umbra. A penumbral eclipse happens when the Moon passes through Earth's penumbra. The Earth's shadow is quite large, so a lunar eclipse lasts for hours and can be seen by anyone with a view of the Moon at the time of the eclipse.




Figure 24.25: The formation of a lunar eclipse.

Partial lunar eclipses occur at least twice a year, but total lunar eclipses are less common. The next total lunar eclipse will occur December 21, 2010. The moon glows with a dull red coloring during a total lunar eclipse.

The Phases of the Moon

The Moon does not produce any light of its own—it only reflects light from the Sun. As the Moon moves around the Earth, we see different parts of the near side of the Moon illuminated by the Sun. This causes the changes in the shape of the Moon that we notice on a regular basis, called the phases of the Moon. As the Moon revolves around Earth, the illuminated portion of the near side of the Moon will change from fully lit to completely dark and back again.

A full moon is the lunar phase seen when the whole of the Moon's lit side is facing Earth. This phase happens when Earth is between the Moon and the Sun. About one week later, the Moon enters the quarter-moon phase. At this point, the Moon appears as a half-circle, since only half of the Moon's lit surface is visible from Earth. When the Moon moves between Earth and the Sun, the side facing Earth is completely dark. This is called the new moon phase, and we do not usually see the Moon at this point. Sometimes you can just barely make out the outline of the new moon in the sky. This is because some sunlight reflects off the Earth and hits the moon. Before and after the quarter-moon phases are the **gibbous** and **crescent** phases. During the gibbous moon phase, the moon is more than half lit but not full. During the crescent moon phase, the moon is less than half lit and is seen as only a sliver or crescent shape. It takes about 29.5 days for the Moon to revolve around Earth and go through all the phases (Figure 24.26).




Figure 24.26: The phases of the Moon. Note that the Sun would be above the top of this picture, and thus, the Sun's rays would be directed downward.

The Tides

Tides are the regular rising and falling of Earth's surface water in response to gravitational attraction from the Moon and Sun. The Moon's gravity causes the oceans to bulge out in the direction of the Moon. In other words, the Moon's gravity is pulling upwards on Earth's water, producing a high tide. On the other side of the Earth, there is another high tide area, produced where the Moon's pull is weakest. As the Earth rotates on its axis, the areas directly in line with the Moon will experience high tides. Each place on

Earth experiences changes in the height of the water throughout the day as it changes from high tide to low tide. There are two high tides and two low tides each tidal day. Figure 24.27 and Figure 24.28 will help you better understand how tides work.




Figure 24.28: A neap tide.

The first picture shows what is called a **spring tide**. Confusingly, this tide has nothing to do with the season "Spring", but means that the tide waters seem to spring forth. During a spring tide, the Sun and Moon are in line. This happens at both the new moon and the full moon. The Sun's gravity pulls on Earth's water, while the Moon's gravity pulls on the water in the same places. The high tide produced by Sun adds to the high tide produced by the Moon. So spring tides have higher than normal high tides. This water is shown on the picture as the gray bulges on opposite sides of the Earth. Notice that perpendicular to the gray areas, the water is at a relatively low level. The places where the water is being pulled out experience high tides, while the areas perpendicular to them experience low tides. Since the Earth is rotating on its axis, the high-low tide cycle moves around the globe in a 24-hour period.

The second picture shows a **neap tide**. A neap tide occurs when the Earth and Sun are in line but the Moon is perpendicular to the Earth. This happens when the moon is at first or last quarter moon phase. In this case, the pull of gravity from the Sun partially cancels out the pull of gravity from the Moon, and the tides are less pronounced. Neap tides produce less extreme tides than the normal tides. This is because the high tide produced by the Sun adds to the low tide area of the Moon and vice versa. So high tide is not as high and low tide is not as low as it usually might be.

Lesson Summary

- As the Earth rotates on its axis and revolves around the Sun, several different effects are produced.
- When the new moon comes between the Earth and the Sun along the ecliptic, a solar eclipse is produced.
- When the Earth comes between the full moon and the Sun along the ecliptic, a lunar eclipse occurs.
- Observing the Moon from Earth, we see a sequence of phases as the side facing us goes from completely darkened to completely illuminated and back again once every 29.5 days.
- Also as the Moon orbits Earth, it produces tides aligned with the gravitational pull of the Moon.
- The Sun also produces a smaller solar tide. When the solar and lunar tide align, at new and full moons, we experience higher than normal tidal ranges, called spring tides.
- At first and last quarter moons, the solar tide and lunar tide interfere with each other, producing lower than normal tidal ranges called neap tides.

Review Questions

1. The globe is divided into time zones, so that any given hour of the day in one time zone occurs at a different time in other time zones. For example, New York City is in one time zone and Los Angeles is in another time zone. When it is 8 am in New York City, it is only 5 am in Los Angeles. Explain how Earth's motions cause this difference in times.
2. Explain how Earth's tilt on its axis accounts for seasons on Earth.
3. Explain how the positions of the Earth, Moon, and Sun vary during a solar eclipse and a lunar eclipse.
4. Draw a picture that shows how the Earth, Moon, and Sun are lined up during the new moon phase.
5. Why are neap tides less extreme than spring tides?

Vocabulary

crescent
Phase of the moon when it is less than half full but still slightly lit.

gibbous
Phase of the moon when it is more than half lit but not completely full.

lunar eclipse
An eclipse that occurs when the Moon moves through the shadow of the Earth and is blocked from view.

neap tide
Type of tide event when the Sun and Earth are in line and the Moon is perpendicular to the Earth.

penumbra
Outer part of shadow that remains partially lit during an eclipse.

solar eclipse
Occurs when moon passes directly between the Earth and Sun; the Moon's shadow blocks the Sun from view.

spring tide
An extreme tide event that happens when the Earth, Moon, and the Sun are aligned; happens at full and new moon phases.

tide
The regular rising and falling of Earth's surface waters twice a tidal day as a result of the Moon's and Sun's gravitational attraction.

umbra
Inner cone shaped part of a shadow when all light is blocked during an eclipse.

Points to Consider

- Why don't eclipses occur every single month at the full and new moons?
- The planet Mars has a tilt that is very similar to Earth's. What does this produce on Mars?
- Venus comes between the Earth and the Sun. Why don't we see an eclipse when this happens?

CC licensed content, Shared previously

- **Provided by**: Wikibooks. **Located at**: http://en.wikibooks.org/wiki/High_School_Earth_Science/The_Sun_and_the_Earth-Moon_System. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-10-the-atmosphere.html%20. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 15.4: The Sun and the Earth-Moon System is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

15.5: Sun – Earth Relationship- The Seasons

SOLAR RADIATION ON EARTH

Different parts of the Earth receive different amounts of solar radiation. Which part of the planet receives the most insolation? The Sun's rays strike the surface most directly at the equator.Different areas also receive different amounts of sunlight in different seasons. What causes the seasons? The seasons are caused by the direction Earth's axis is pointing relative to the Sun.

The Earth revolves around the Sun once each year and spins on its axis of rotation once each day. This axis of rotation is tilted 23.5 degrees relative to its plane of orbit around the Sun. The axis of rotation is pointed toward Polaris, the North Star. As the Earth orbits the Sun, the tilt of Earth's axis stays lined up with the North Star.



NORTHERN HEMISPHERE SUMMER

The North Pole is tilted towards the Sun and the Sun's rays strike the Northern Hemisphere more directly in summer. At the summer solstice, June 21 or 22, the Sun's rays hit the Earth most directly along the Tropic of Cancer (23.5 degrees N); that is, the angle of incidence of the sun's rays there is zero (the angle of incidence is the deviation in the angle of an incoming ray from straight on). When it is summer solstice in the Northern Hemisphere, it is winter solstice in the Southern Hemisphere.

NORTHERN HEMISPHERE WINTER

Winter solstice for the Northern Hemisphere happens on December 21 or 22. The tilt of Earth's axis points away from the Sun. Light from the Sun is spread out over a larger area, so that area isn't heated as much. With fewer daylight hours in winter, there is also less time for the Sun to warm the area. When it is winter in the Northern Hemisphere, it is summer in the Southern Hemisphere.

EQUINOX

Halfway between the two solstices, the Sun's rays shine most directly at the equator, called an "equinox." The daylight and nighttime hours are exactly equal on an equinox. The autumnal equinox happens on September 22 or 23 and the vernal or spring equinox happens March 21 or 22 in the Northern Hemisphere.

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-10-the-atmosphere.html%20. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 15.5: Sun – Earth Relationship- The Seasons is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

16: Light and the Sun

16.1: Electromagnetic Spectrum

16.2: Doppler Effect

16.3: Telescopes

16.4: Nuclear Fusion

This page titled 16: Light and the Sun is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

16.1: Electromagnetic Spectrum

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=160

CC licensed content, Original

- **Authored by**: William Hopper. **Provided by**: Florida Memorial University. **License**: *CC BY: Attribution*

This page titled 16.1: Electromagnetic Spectrum is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.


16.2: Doppler Effect

The **Doppler effect** (or **Doppler shift**), named after the Austrian physicist Christian Doppler, who proposed it in 1842 in Prague, is the change in frequency of a wave (or other periodic event) for an observer moving relative to its source. It is commonly heard when a vehicle sounding a siren or horn approaches, passes, and recedes from an observer. Compared to the emitted frequency, the received frequency is higher during the approach, identical at the instant of passing by, and lower during the recession.

When the source of the waves is moving toward the observer, each successive wave crest is emitted from a position closer to the observer than the previous wave. Therefore each wave takes slightly less time to reach the observer than the previous wave. Therefore the time between the arrival of successive wave crests at the observer is reduced, causing an increase in the frequency. While they are travelling, the distance between successive wave fronts is reduced; so the waves “bunch together”. Conversely, if the source of waves is moving away from the observer, each wave is emitted from a position farther from the observer than the previous wave, so the arrival time between successive waves is increased, reducing the frequency. The distance between successive wave fronts is increased, so the waves “spread out”.


Astronomy

The Doppler effect for electromagnetic waves such as light is of great use in astronomy and results in either a so-called redshift or blueshift. It has been used to measure the speed at which stars and galaxies are approaching or receding from us, that is, the radial velocity. This is used to detect if an apparently single star is, in reality, a close binary and even to measure the rotational speed of stars and galaxies.




Redshift of spectral lines in the optical spectrum of a supercluster of distant galaxies (right), as compared to that of the Sun (left)

The use of the Doppler effect for light in astronomy depends on our knowledge that the spectra of stars are not homogeneous. They exhibit absorption lines at well defined frequencies that are correlated with the energies required to excite electrons in various elements from one level to another. The Doppler effect is recognizable in the fact that the absorption lines are not always at the frequencies that are obtained from the spectrum of a stationary light source. Since blue light has a higher frequency than red light, the spectral lines of an approaching astronomical light source exhibit a blueshift and those of a receding astronomical light source exhibit a redshift.



Redshift and blueshift

Example



Doppler effect, yellow (~575 nm wavelength) ball appears greenish (blueshift to ~565 nm wavelength) approaching observer, turns orange (redshift to ~585 nm wavelength) as it passes, and returns to yellow when motion stops. To observe such a change in color, the object would have to be traveling at approximately 5200 km/s, or about 75 times faster than the speed record for the fastest manmade space probe.

Among the nearby stars, the largest radial velocities with respect to the Sun are +308 km/s (BD-15°4041, also known as LHS 52, 81.7 light-years away) and -260 km/s (Woolley 9722, also known as Wolf 1106 and LHS 64, 78.2 light-years away). Positive radial velocity means the star is receding from the Sun, negative that it is approaching.

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Doppler_effect. **License**: *CC BY-SA: Attribution-ShareAlike*
- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Redshift. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 16.2: Doppler Effect is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

16.3: Telescopes

Lesson Objectives

- Explain how astronomers use the whole electromagnetic spectrum to study the universe beyond Earth.
- Identify different types of telescopes.
- Describe historical and modern observations made with telescopes.

Vocabulary

- astronomer
- catadioptric telescope
- constellation
- electromagnetic (EM) radiation
- electromagnetic spectrum
- frequency
- gamma ray
- infrared light
- light-year
- microwave
- planet
- radio telescope
- radio wave
- reflecting telescope
- refracting telescope
- space telescope
- spectrometer
- ultraviolet (UV)
- visible light
- wavelength
- X-ray

Introduction

Many scientists interact directly with what they are studying. Biologists can collect cells, seeds, or sea urchins and put them in a controlled laboratory environment. Physicists can subject metals to stress or smash atoms into each other. Geologists can chip away at rocks to see what is inside. But **astronomers**, scientists who study the universe beyond Earth, rarely have a chance for direct contact with their subject. Astronomers observe their subjects at a distance, usually a very large distance!

Electromagnetic Radiation

Earth is separated from the rest of the universe by very large expanses of space. Very rarely matter from outside Earth's environment reaches us, such as when a meteorite makes it through the atmosphere from elsewhere in the solar system. But for the most part, astronomers have one main source for their data — light. Light can travel across empty space, and as it does, so it carries both energy and information. Light is one type of**electromagnetic (EM) radiation**, energy that is transmitted through space as a wave.

These videos discuss infrared, ultraviolet, and radio telescopes, as well as telescopes that detect visible light, and reveal tremendous features of the stars and galaxies around the Universe **(2d)**: http://www.youtube.com/watch?v=AK-gtuAJ-B4 (2:19), http://www.youtube.com/watch?v=aQJQH7lS27s (5:30).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=164

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=164

The Speed of Light

Light travels faster than anything else in the universe. In the almost completely empty vacuum of space, light travels at a speed of approximately 300,000,000 meters per second (670,000,000 miles per hour). To give you an idea of how fast that is, a beam of light could travel from New York to Los Angeles and back again nearly 40 times in just one second. Even though light travels extremely fast, objects in space are so far away that it takes a significant amount of time for light from those objects to reach us. For example, light from the Sun takes about 8 minutes to reach Earth.

Light-Years

Since astronomical distances are so large, it helps to have a unit of measurement that is good for expressing those large distances. A **light-year** is a unit of distance that is defined as the distance that light travels in one year. One light-year is approximately equal to 9,500,000,000,000 (9.5 trillion) kilometers, or 5,900,000,000,000 (5.9 trillion) miles (**Figure** below). That's a long way! But by astronomical standards, it's actually a pretty short distance.

Proxima Centauri, the closest star to us after the Sun, is 4.22 light-years away. That means the light from Proxima Centauri takes 4.22 years to reach us. The galaxy we live in, the Milky Way Galaxy, is about 100,000 light-years across. How long does it take light to travel from one side of the galaxy to the other? 100,000 years! If an astronomer looks through a telescope at a star that is 1,000 light years away, is she seeing the star as it is now?



The smallest red-shifted galaxies are among the most distant galaxies we have detected so far at 13 billion light-years away. That's over a hundred-billion-trillion (100,000,000,000,000,000,000,000) kilometers!

Looking Back in Time

When we look at astronomical objects such as stars and galaxies, we are not just seeing over great distances—we are also seeing back in time. Because light takes time to travel, the image we see of a distant galaxy is an image of how the galaxy used to look. For example, the Andromeda Galaxy, shown in (**Figure** below), is about 2.5 million light years from Earth. If you look at the Andromeda Galaxy through a telescope, what are you seeing? You are seeing the galaxy as it was 2.5 million years ago. If the galaxy ceased to exist 1 million years ago, when would you know that? If you want to see the galaxy as it is now, you will have to wait and look again 2.5 million years into the future.




This recent picture of the Andromeda Galaxy actually shows the galaxy as it was about 2.5 million years ago.

Electromagnetic Waves

Light is one type of EM radiation; light is energy that travels in the form of an *electromagnetic wave*. (**Figure** below) shows a diagram of an electromagnetic wave. An EM wave has two components: an electric field and a magnetic field. Each of these components oscillates between positive and negative values, which is what makes the "wavy" shape in the diagram.

The distance between two adjacent oscillations is called **wavelength**. A related value is **frequency**, which measures the number of wavelengths that pass a given point every second. Wavelength and frequency are reciprocal, which means that as one increases, the other decreases.



An electromagnetic wave consists of oscillating electric and magnetic fields.

The Electromagnetic Spectrum

Visible light — the light that human eyes can see — comes in a variety of colors. The color of visible light is determined by its wavelength. Visible light ranges from wavelengths of 400 nm to 700 nm, corresponding to the colors violet through red. EM radiation with wavelengths shorter than 400 nm or longer than 700 nm exists all around you — you just can't see it. The full range of electromagnetic radiation, or the**electromagnetic spectrum**, is shown in **Figure** below.



(a) Visible light is part of the electromagnetic spectrum, which ranges from gamma rays with very short wavelengths, to radio waves with very long wavelengths. (b) These are images of the same scene. In the top, only the wavelengths of visible light show. In the bottom, a layer of thick clouds appears in the infrared wavelengths.

Like our Sun, every star emits light at a wide range of wavelengths, all across the visible spectrum and even outside the visible spectrum. Astronomers can learn a lot from studying the details of the spectrum of light from a star.

Some very hot stars emit light primarily at **ultraviolet (UV)** wavelengths, while some very cool stars emit mostly in the **infrared**. There are extremely hot objects that emit **X-rays** and even **gamma rays**. Light from some of the faintest, most distant objects is in the form of **radio waves**. In fact, a lot of the objects most interesting to astronomers today can't even be seen with the naked eye. Astronomers use telescopes to detect the faint light from distant objects and to see objects at wavelengths all across the electromagnetic spectrum.


To learn more about star's spectra, check out http://www.colorado.edu/physics/PhysicsInitiative/Physics2000/quantumzone/.

Types of Telescopes

Optical Telescopes

People have been making and using lenses for magnification for thousands of years. However, the first true telescopes were made in Europe in the late 16th century. These telescopes used a combination of two lenses to make distant objects appear both nearer and larger. The term *telescope*was coined by the Italian scientist and mathematician Galileo Galilei (1564–1642). Galileo built his first telescope in 1608 and subsequently made many improvements to telescope design.

Telescopes that rely on the refraction, or bending, of light by lenses are called **refracting telescopes**, or simply *refractors*. The earliest telescopes, including Galileo's, were all refractors. Many of the small telescopes used by amateur astronomers today are refractors. Refractors are particularly good for viewing details within our solar system, such as the surface of Earth's moon or the rings around Saturn (**Figure** below).



The largest refracting telescope in the world is at the University of Chicago's Yerkes Observatory in Wisconsin and was built in 1897. Its largest lens has a diameter of 102 cm.

Around 1670, another famous scientist and mathematician — Sir Isaac Newton (1643–1727) — built a different kind of telescope. Newton used curved mirrors to focus light and so created the first **reflecting telescopes**, or *reflectors* (**Figure** below). The mirrors in a reflecting telescope are much lighter than the heavy glass lenses in a refractor. This is significant, because:

- To support the thick glass lenses a refractor must be strong and heavy.
- Mirrors are easier to make precisely than it is to make glass lenses.
- Because they do not need to be as heavy to support the same size lens, reflectors can be made larger than refractors.

Larger telescopes can collect more light and so they can study dimmer or more distant objects. The largest optical telescopes in the world today are reflectors.



(a) Reflecting telescopes used by amateur astronomers today are similar to the one designed by Isaac Newton in the 17th century. (b) The South African Large Telescope (SALT) is one of the largest reflecting telescopes on Earth. SALT's primary mirror consists of 91 smaller hexagonal mirrors, each with sides 1 m long. (c) Many amateur astronomers today use catadioptric telescopes.

Catadioptric telescopes have a combination of mirrors and lenses to focus light. Catadioptric telescopes have large mirrors to collect a lot of light, but short tubes for portability.

KQED: Amateur Astronomers

Amateur astronomers enjoy observing and studying stars and other celestial objects. Both professional and amateur astronomers use telescopes. A telescope is an instrument that makes faraway objects look closer. Learn more at: http://science.kqed.org/quest/video/amateur-astronomers/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=164

Radio Telescopes

Notice it says above that the largest *optical* telescopes in the world are reflectors. Optical telescopes collect visible light. Even larger telescopes are built to collect light at longer wavelengths — radio waves. What do you think these telescopes are called? **Radio telescopes** look a lot like satellite dishes because both are designed to do the same thing — to collect and focus radio waves or **microwaves** (the shortest wavelength waves) from space.

The largest single telescope in the world is at the Arecibo Observatory in Puerto Rico (**Figure** below). This telescope is located in a naturally occurring sinkhole that formed when water flowing underground dissolved the limestone rock. If this telescope were not supported by the ground, it would collapse under its own weight. Since the telescope is set into the ground it cannot be aimed to different parts of the sky and so can only observe the part of the sky that happens to be overhead at a given time.



The radio telescope at the Arecibo Observatory has a diameter of 305 m.

A group of radio telescopes can be linked together with a computer so that they are all observing the same object (**Figure** below). The computer combines the data, making the group function like one single telescope.

For more on radio telescopes and radio astronomy in general, go to http://www.nrao.edu/whatisra/index.shtml.



The Very Large Array in New Mexico has 27 radio dishes, each 25 m in diameter. When all the dishes are pointed at the same object, they are like a single telescope with a diameter of 22.3 mi.

KQED: SETI: The New Search for ET

Scientists have upped their search for extraterrestrial intelligence with the Allen Telescope Array, a string of 350 radio telescopes, located 300 miles north of San Francisco. Find out why SETI scientists now say we might be hearing from ET sooner than you think. Learn more at:http://science.kqed.org/quest/video/seti-the-new-search-for-et/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=164

KQED: Interview with Astronomer Jill Tartar

SETI listens for signs of other civilization's technology. Dr. Jill Tartar explains the program: What it's looking for; what the problems are; what the potential benefits are. Learn more at: http://www.youtube.com/watch?v=QwEm3WHvNHI.


An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=164

Space Telescopes

Telescopes on Earth all have one significant limitation: the electromagnetic radiation they gather must pass through Earth's atmosphere. The atmosphere blocks some radiation in the infrared part of the spectrum and almost all radiation in the ultraviolet and higher frequency ranges. Furthermore, motion in the atmosphere distorts light. That distortion is why stars twinkle in the night sky. To minimize these problems, many observatories are built on high mountains, where there is less atmosphere above the telescope. Even better, **space telescopes** avoid such problems completely because they orbit outside Earth's atmosphere in space. Space telescopes can carry instruments to observe objects emitting various types of electromagnetic radiation such as visible, infrared or ultraviolet light; gamma rays; or x-rays. X-ray telescopes, such as the Chandra X-ray Observatory, use X-ray optics to observe remote objects in the X-ray spectrum.

The Hubble Space Telescope (HST), shown in (**Figure** below), is perhaps the best known space telescope. The Hubble was put into orbit by the Space Shuttle Atlantis in 1990. Once it was in orbit, scientists discovered that there was a flaw in the shape of the mirror. A servicing mission to the Hubble by the Space Shuttle Endeavor in 1994 corrected the problem. Since that time, the Hubble has provided huge amounts of data that have helped to answer many of the biggest questions in astronomy.


Find out more by visiting the Hubble Space Telescope website at http://hubblesite.org.



(a) The Hubble Space Telescope orbits Earth at an altitude of 589 km (366 mi). It collects data in visible, infrared, and ultraviolet wavelengths. (b) This starburst cluster is one of the many fantastic images taken by the HST over the past two decades.

The National Aeronautics and Space Administration (NASA) has placed three other major space telescopes in orbit, comprising what NASA calls the 'Great Observatories'. Each of these telescopes specializes in a different part of the electromagnetic spectrum (**Figure** below). NASA is planning for another telescope, the James Webb Space Telescope, to serve as a replacement for the aging Hubble. The James Webb is scheduled to launch in 2018.

To learn more about NASA's great observatories, check outhttp://www.nasa.gov/audience/forstudents/postsecondary/features/F_NASA_Great_Observatories_PS.html.



NASA's four space-based Great Observatories were designed to view the universe in different ranges of the electromagnetic spectrum. A. Hubble Space Telescope: visible, infrared and ultraviolet light; B. Compton Gamma Ray Observatory (inactive): gamma ray; C. Spitzer Space Telescope: infrared; D. Chandra X-ray Observatory: X-ray.

Observations with Telescopes

Ancient Astronomers

Humans have been studying the night sky for thousands of years. Observing the patterns and motions in the sky helped ancient peoples keep track of time (**Figure** below). By understanding annual rhythms, people could know when to plant crops. They also timed many of their religious ceremonies to coincide with events in the heavens.



Many archaeologists think that Stonehenge was used to observe the movement of the moon and the sun.

The ancient Greeks made careful observations of the locations of stars in the sky. They noticed that some of the 'stars' moved against the background of other stars. They called these bright and odd bodies in the sky **planets**, which in Greek means "wanderers." Today we know that the planets are not stars, but members of our solar system that orbit the Sun. Greeks also identified **constellations**, patterns of stars in the sky (**Figure** below). They associated the constellations with stories and myths from their culture.




Stars in the constellation Orion. Constellations help astronomers today identify different regions of the night sky.

Galileo's Observations

Ancient astronomers knew a lot about the patterns of stars and the movement of objects in the sky, but they did not know what these objects actually were. That understanding began in the year 1610, when Galileo turned a telescope toward the heavens. With his telescope, Galileo made the following discoveries (among others):

- There are more stars in the night sky than the naked eye can see.
- The band of stars called the Milky Way consists of many stars.
- The Moon has craters (**Figure** below).
- Venus has phases like the Moon.
- Jupiter has orbiting moons.
- There are dark spots that move across the surface of the Sun.



Galileo was the first person known to look at the Moon through a telescope. Galileo made the drawing on the left in 1610; on the right is a modern photograph of the Moon.


Galileo's observations challenged people to think in new ways about the universe and Earth's place in it. About 100 years before Galileo, Nicolaus Copernicus had proposed a controversial new model of the universe in which Earth and the other planets revolve around the Sun. In Galileo's time, most people still believed that the Sun and planets revolved around Earth. Galileo's observations provided direct evidence to support Copernicus' model.

Observations with Modern Telescopes

Equipped with no more than a good pair of binoculars, you can see all of the things Galileo saw, and more. You can even see sunspots, but be sure to use special filters on the lenses to protect your eyes. With a basic telescope like those used by many

amateur astronomers, you can see more than Galileo saw, such as polar caps on Mars, the rings of Saturn, and bands in the atmosphere of Jupiter.

All of the objects mentioned above are within our solar system. With a telescope you can also see many times more stars than without a telescope. However, because they are so far away, the stars will appear as points of light. This is true even of the most powerful professional telescopes, with one rare exception (**Figure** below).



This ultraviolet image of the red supergiant star Betelgeuse taken with the Hubble Space Telescope in 1996 was the first direct image taken of the disk of a star other than the Sun.

Very few professional astronomers today look directly through the eyepiece of a telescope. Instead, they attach to the telescopes sophisticated instruments that capture and process the light. The astronomers then look at the images or data shown on these instruments. Most often the instruments then pass the data on to a computer where the data can be stored for later use. An astronomer may take weeks or months to analyze all the data collected from just a single night.

Astronomers use **spectrometers** to study the light from a telescope. A spectrometer uses a prism or other device to break light down into its component colors. The spectrum produced can be observed directly, captured on film, or stored digitally on a computer (**Figure** below).



A simplified example of starlight after it passes through a spectrometer. The dark lines in a star's spectrum are caused by gases absorbing light in the star's outer atmosphere.

From a single spectrum of a star, an astronomer can tell:

- How hot the star is (by the relative brightness of different colors).
- What elements the star contains (by the pattern of dark lines).
- Whether and how fast the star is moving toward or away from Earth (by how far the dark lines are shifted from their normal positions).

Using telescopes, astronomers can also learn how stars evolve, what kind of matter is found throughout the universe, how that matter is distributed, and even how the universe might have formed.

Lesson Summary

- Astronomers study light from distant objects.
- Light travels at 300,000,000 meters per second — faster than anything else in the universe.
- A light-year is equal to the distance light travels in one year, 9.5 trillion kilometers.
- When we see distant objects, we see them as they were in the past, because their light has been traveling to us for many years.
- Visible light is part of the electromagnetic spectrum.
- Telescopes make distant objects appear both nearer and larger.
- Optical telescopes collect visible light. The three main types are reflecting telescopes, refracting telescopes, and catadioptric telescopes.
- Radio telescopes collect and focus radio waves from distant objects.
- Space telescopes orbit Earth, collecting wavelengths of light that are normally blocked by the atmosphere.
- Galileo was the first person known to use a telescope to study the sky. His discoveries helped change the way humans think about the universe.

- Modern telescopes collect data that can be stored on a computer.
- Astronomers can learn a lot about a star by studying its spectrum.

Review Questions

1. Betelgeuse is around 640 light-years from Earth. Light travels 9.5 trillion kilometers in one year. How far away is Betelgeuse in kilometers?
2. Identify four regions of the electromagnetic spectrum that astronomers use when observing objects in space.
3. List the three main types of optical telescopes, and describe their differences.
4. Explain the advantages of putting a telescope into orbit around Earth.
5. Describe two observations that Galileo was the first to make with his telescope.
6. List three things that an astronomer can learn about a star by studying its spectrum.

Further Reading / Supplemental Links

- Lots of news and information from NASA at http://www.nasa.gov/
- The first stars in the Universe: http://science.nasa.gov/headlines/y2002/08feb_gravlens.htm
- How many stars can you observe? http://www.stargazing.net/David/constel/howmanystars.html
- Archeastronomy: http://www.astronomy.pomona.edu/archeo/
- An activity found online with directions for how to determine if that bright point you see in the sky is a planet or a star:http://cse.ssl.berkeley.edu/SegwayEd/lessons/findplanets/Find-hmpg2.html

Points to Consider

- Radio waves are used for communicating with spacecraft. A round-trip communication from Earth to Mars takes anywhere from 6 to 42 minutes. What challenges does this present for sending unmanned spacecraft and probes to Mars?
- The Hubble Space Telescope is a very important source of data for astronomers. The fascinating and beautiful images from the Hubble also help to maintain public support for science. However, the Hubble is growing old. Missions to service and maintain the telescope are extremely expensive and put the lives of astronauts at risk. Do you think there should be another servicing mission to the Hubble?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/23.1/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 16.3: Telescopes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.


16.4: Nuclear Fusion

Star Power

The Sun is Earth's major source of energy, yet the planet only receives a small portion of its energy and the Sun is just an ordinary star. Many stars produce much more energy than the Sun. The energy source for all stars is nuclear fusion.


Nuclear Fusion

Stars are made mostly of hydrogen and helium, which are packed so densely in a star that in the star's center the pressure is great enough to initiate nuclear fusion reactions. In a **nuclear fusion reaction**, the nuclei of two atoms combine to create a new atom. Most commonly, in the core of a star, two hydrogen atoms fuse to become a helium atom. Although nuclear fusion reactions require a lot of energy to get started, once they are going they produce enormous amounts of energy (**Figure** below).



A thermonuclear bomb is an uncontrolled fusion reaction in which enormous amounts of energy are released.

In a star, the energy from fusion reactions in the core pushes outward to balance the inward pull of gravity. This energy moves outward through the layers of the star until it finally reaches the star's outer surface. The outer layer of the star glows brightly, sending the energy out into space as electromagnetic radiation, including visible light, heat, ultraviolet light, and radio waves (**Figure** below).



A diagram of a star like the Sun.

In particle accelerators, subatomic particles are propelled until they have attained almost the same amount of energy as found in the core of a star (**Figure** below). When these particles collide head-on, new particles are created. This process simulates the nuclear fusion that takes place in the cores of stars. The process also simulates the conditions that allowed for the first helium atom to be produced from the collision of two hydrogen atoms in the first few minutes of the universe.



The SLAC National Accelerator Lab in California can propel particles a straight 2 mi (3.2 km).

The CERN Particle Accelerator presented in this video is the world's largest and most powerful particle accelerator and can boost subatomic particles to energy levels that simulate conditions in the stars and in the early history of the universe before stars formed **(2e)**:http://www.youtube.com/watch?v=sxAxV7g3yf8 (6:16).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=166

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/26.1/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 16.4: Nuclear Fusion is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

17: The Solar System

17.1: Introduction to the Solar System

17.2: Inner Planets

17.3: Outer Planets

17.4: Other Objects in the Solar System

This page titled 17: The Solar System is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

17.1: Introduction to the Solar System

Lesson Objectives

- Describe historical views of the solar system.
- Name the planets, and describe their motion around the sun.
- Explain how the solar system formed.

Vocabulary

- geocentric model
- heliocentric model
- moon
- nebula
- nebular hypothesis
- solar system

Changing Views of the Solar System

Humans' view of the **solar system** has evolved as technology and scientific knowledge have increased. The ancient Greeks identified five of the planets and for many centuries they were the only planets known. Since then, scientists have discovered two more planets, many other solar-system objects and even planets found outside our solar system.

The Geocentric Universe

The ancient Greeks believed that Earth was at the center of the universe, as shown in **Figure** below. This view is called the **geocentric model** of the universe. Geocentric means "Earth-centered." In the geocentric model, the sky, or heavens, are a set of spheres layered on top of one another. Each object in the sky is attached to a sphere and moves around Earth as that sphere rotates. From Earth outward, these spheres contain the Moon, Mercury, Venus, the Sun, Mars, Jupiter, and Saturn. An outer sphere holds all the stars. Since the planets appear to move much faster than the stars, the Greeks placed them closer to Earth.



Model of a geocentric universe. This diagram of the universe from the Middle Ages shows Earth at the center, with the Moon, the Sun, and the planets orbiting Earth.


The geocentric model worked well, by explaining why all the stars appear to rotate around Earth once per day. The model also explained why the planets move differently from the stars and from each other.

One problem with the geocentric model is that some planets seem to move backwards (in retrograde) instead of in their usual forward motion around Earth.

A demonstration animation of retrograde motion of Mars as it appears to Earth can be found here:

http://projects.astro.illinois.edu/data/Retrograde/index.html

Around 150 A.D. the astronomer Ptolemy resolved this problem by using a system of circles to describe the motion of planets (**Figure** below). In Ptolemy's system, a planet moves in a small circle, called an epicycle. This circle moves around Earth in a larger circle, called a deferent. Ptolemy's version of the geocentric model worked so well that it remained the accepted model of the universe for more than a thousand years.




According to Ptolemy, a planet moves on a small circle (epicycle) that in turn moves on a larger circle (deferent) around Earth.

An animation of Ptolemy's system is seen here:

http://www.youtube.com/watch?v=FHSWVLwbbNw

The Heliocentric Universe


Ptolemy's geocentric model worked but it was not only complicated, it occasionally made errors in predicting the movement of planets. At the beginning of the 16th century A.D., Nicolaus Copernicus proposed that Earth and all the other planets orbit the Sun. With the Sun at the center, this model is called the **heliocentric model** or "sun-centered" model of the universe (**Figure** below). Copernicus' model explained the motion of the planets as well as Ptolemy's model did, but it did not require complicated additions like epicycles and deferents.



Unlike the geocentric model, the heliocentric model had the Sun at the center and did not require epicycles.

Although Copernicus' model worked more simply than Ptolemy's, it still did not perfectly describe the motion of the planets because, like Ptolemy, Copernicus thought planets moved in perfect circles. Not long after Copernicus, Johannes Kepler refined the heliocentric model so that the planets moved around the Sun in ellipses (ovals), not circles (**Figure** below). Kepler's model matched observations perfectly.

Animation of Kepler's Laws of Planetary Motion: http://projects.astro.illinois.edu/data/KeplersLaws/index.html




Kepler's model showed the planets moving around the sun in ellipses. The elliptical orbits are exaggerated in this image.

Because people were so used to thinking of Earth at the center of the universe, the heliocentric model was not widely accepted at first. However, when Galileo Galilei first turned a telescope to the heavens in 1610, he made several striking discoveries. Galileo discovered that the planet Jupiter has **moons** orbiting around it. This provided the first evidence that objects could orbit something besides Earth.

An animation of three of Jupiter's moons orbiting the planet is seen here:http://upload.wikimedia.org/Wikipedia/commons/e/e7/Galilean_moon_Laplace_resonance_animation_de.gif.

Galileo also discovered that Venus has phases like the Moon (**Figure** below), which provides direct evidence that Venus orbits the Sun.



The phases of Venus.

Galileo's discoveries caused many more people to accept the heliocentric model of the universe, although Galileo himself was found guilty of heresy for his ideas. The shift from an Earth-centered view to a Sun-centered view of the universe is referred to as the Copernican Revolution.

Watch this animation of the Ptolemaic and Copernican models of the solar system. Ptolemy made the best model he could with the assumption that Earth was the center of the universe, but by letting that assumption go, Copernicus came up with a much simpler model. Before people would accept that Copernicus was right, they needed to accept that the Sun was the center of the solar system **(1n – I&E Stand.)**:http://www.youtube.com/watch?v=VyQ8Tb85HrU (0:47).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170


The Modern Solar System

Today, we know that our solar system is just one tiny part of the universe as a whole. Neither Earth nor the Sun are at the center of the universe. However, the heliocentric model accurately describes the solar system. In our modern view of the solar system, the

Sun is at the center, with the planets moving in elliptical orbits around the Sun. The planets do not emit their own light, but instead reflect light from the Sun.

Extrasolar Planets or Exoplanets

Since the early 1990s, astronomers have discovered other solar systems, with planets orbiting stars other than our own Sun (called "extrasolar planets" or simply "exoplanets") (**Figure** below).



The extrasolar planet Fomalhaut is surrounded by a large disk of gas. The disk is not centered on the planet, suggesting that another planet may be pulling on the gas as well.

Some extrasolar planets have been directly imaged, but most have been discovered by indirect methods. One technique involves detecting the very slight motion of a star periodically moving toward and away from us along our line-of-sight (also known as a star's "radial velocity"). This periodic motion can be attributed to the gravitational pull of a planet or, sometimes, another star orbiting the star.

This is in line with the plane of the system: http://en.Wikipedia.org/wiki/File:Dopspec-inline.gif.

A planet may also be identified by measuring a star's brightness over time. A temporary, periodic decrease in light emitted from a star can occur when a planet crosses in front of (or "transits") the star it is orbiting, momentarily blocking out some of the starlight.

More than 3,600 extrasolar planets have been identified and the rate of discovery is increasing rapidly.

Extrasolar Planet from the ESA discusses extrasolar planets and particularly a planetary system very similar to our solar system **(1g)**:http://www.youtube.com/watch?v=ouJahDONTWc (3:29).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170

An introduction to extrasolar planets from NASA is available at **(1g)**: http://www.youtube.com/watch?v=oeeZCHDNTvQ (3:14).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170

KQED: The Planet Hunters


Hundreds of exoplanets have now been discovered. To learn something about how planet hunters find these balls of rock they usually can't even see, watch this QUEST video. Learn more at: http://science.kqed.org/quest/video/the-planet-hunters/ andhttp://science.kqed.org/quest/audio/exoplanets/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170

Planets and Their Motions

Since the time of Copernicus, Kepler, and Galileo, we have learned a lot more about our solar system. Astronomers have discovered two more planets (Uranus and Neptune), four dwarf planets (Ceres, Pluto, Makemake, Haumea, and Eris), more than 150 moons, and many, many asteroids and other small objects.

(**Figure** below) shows the Sun and the major objects that orbit the Sun. There are eight planets (Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune) and the five known dwarf planets and the five known dwarf planets (Ceres, Pluto, Makemake, Haumea, and Eris).



Relative sizes of the Sun, planets and dwarf planets. The relative sizes are correct and their position relative to each other is correct, but the relative distances are not correct.


Although the Sun is just an average star compared to other stars, it is by far the largest object in the solar system. The Sun is more than 500 times the mass of everything else in the solar system combined! Table below gives data on the sizes of the Sun and planets relative to Earth.

Sizes of Solar System Objects Relative to Earth


Object	Mass (Relative to Earth)	Diameter of Planet (Relative to Earth)
Sun	333,000 Earth's mass	109.2 Earth's diameter
Mercury	0.06 Earth's mass	0.39 Earth's diameter
Venus	0.82 Earth's mass	0.95 Earth's diameter
Earth	1.00 Earth's mass	1.00 Earth's diameter
Mars	0.11 Earth's mass	0.53 Earth's diameter
Jupiter	317.8 Earth's mass	11.21 Earth's diameter
Saturn	95.2 Earth's mass	9.41 Earth's diameter
Uranus	14.6 Earth's mass	3.98 Earth's diameter
Neptune	17.2 Earth's mass	3.81 Earth's diameter

The Size and Shape of Orbits

Figure below shows the relative sizes of the orbits of the planets, asteroid belt, and Kuiper belt. In general, the farther away from the Sun, the greater the distance from one planet's orbit to the next. The orbits of the planets are not circular but slightly elliptical with the Sun located at one of the foci (**Figure** below).



The relative sizes of the orbits of planets in the solar system. The inner solar system and asteroid belt is on the upper left. The upper right shows the outer planets and the Kuiper belt.



The planets orbit the Sun in regular paths.

While studying the solar system, Johannes Kepler discovered the relationship between the time it takes a planet to make one complete orbit around the Sun, its “orbital period,” and the distance from the Sun to the planet. If the orbital period of a planet is known, then it is possible to determine the planet’s distance from the Sun. This is how astronomers without modern telescopes could determine the distances to other planets within the solar system.

Distances in the solar system are often measured in **astronomical units** (AU). One astronomical unit is defined as the distance from Earth to the Sun. 1 AU equals about 150 million km, or 93 million mi. **Table** below shows the distances to the planets (the average radius of orbits) in AU. The table also shows how long it takes each planet to spin on its axis (the length of a day) and how long it takes each planet to complete an orbit (the length of a year); in particular, notice how slowly Venus rotates relative to Earth.

Distances to the Planets and Properties of Orbits Relative to Earth’s Orbit

Planet	Average Distance from Sun (AU)	Length of Day (In Earth Days)	Length of Year (In Earth Years)
Mercury	0.39 AU	56.84 days	0.24 years
Venus	0.72	243.02	0.62
Earth	1.00	1.00	1.00
Mars	1.52	1.03	1.88
Jupiter	5.20	0.41	11.86
Saturn	9.54	0.43	29.46
Uranus	19.22	0.72	84.01
Neptune	30.06	0.67	164.8

How old are you on Earth? How old would you be if you lived on Jupiter? How many days is it until your birthday on Earth? How many days until your birthday if you lived on Saturn?


Scaling the solar system creates a scale to measure all objects in solar system **(1i – I&E Stand.)**: http://www.youtube.com/watch?v=-6szEDHMxP4 (4:44).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170

The Role of Gravity

Isaac Newton was one of the first scientists to explore gravity. He understood that the Moon circles the Earth because a force is pulling the Moon toward Earth's center. Without that force, the Moon would continue moving in a straight line off into space. Newton also came to understand that the same force that keeps the Moon in its orbit is the same force that causes objects on Earth to fall to the ground.

Newton defined the Universal Law of Gravitation, which states that a force of attraction, called gravity, exists between all objects in the universe (**Figure** below). The strength of the gravitational force depends on how much mass the objects have and how far apart they are from each other. The greater the objects' mass, the greater the force of attraction; in addition, the greater the distance between the objects, the smaller the force of attraction.



The force of gravity exists between all objects in the universe; the strength of the force depends on the mass of the objects and the distance between them.

The distance between the Sun and each of its planets is very large, but the Sun and each of the planets are also very large. Gravity keeps each planet orbiting the Sun because the star and its planets are very large objects. The force of gravity also holds moons in orbit around planets.

Formation of the Solar System

There are two additional key features of the solar system:

1. All the planets lie in nearly the same plane, or flat disk like region.

2. All the planets orbit in the same direction around the Sun.


These two features are clues to how the solar system formed.

A Giant Nebula

The most widely accepted explanation of how the solar system formed is called the **nebular hypothesis**. According to this hypothesis, the Sun and the planets of our solar system formed about 4.6 billion years ago from the collapse of a giant cloud of gas and dust, called a **nebula**.

The nebula was drawn together by gravity, which released gravitational potential energy. As small particles of dust and gas smashed together to create larger ones, they released kinetic energy. As the nebula collapsed, the gravity at the center increased and the cloud started to spin because of its angular momentum. As it collapsed further, the spinning got faster, much as an ice skater spins faster when he pulls his arms to his sides during a spin.

Much of the cloud's mass migrated to its center but the rest of the material flattened out in an enormous disk, as shown in **Figure** below. The disk contained hydrogen and helium, along with heavier elements and even simple organic molecules.



An artist's painting of a protoplanetary disk.

Formation of the Sun and Planets

As gravity pulled matter into the center of the disk, the density and pressure at the center became intense. When the pressure in the center of the disk was high enough, nuclear fusion began. A star was born—the Sun. The burning star stopped the disk from collapsing further.

Meanwhile, the outer parts of the disk were cooling off. Matter condensed from the cloud and small pieces of dust started clumping together. These clumps collided and combined with other clumps. Larger clumps, called planetesimals, attracted smaller clumps with their gravity. Gravity at the center of the disk attracted heavier particles, such as rock and metal and lighter particles remained further out in the disk. Eventually, the planetesimals formed protoplanets, which grew to become the planets and moons that we find in our solar system today.

Because of the gravitational sorting of material, the inner planets — Mercury, Venus, Earth, and Mars — formed from dense rock and metal. The outer planets — Jupiter, Saturn, Uranus and Neptune — condensed farther from the Sun from lighter materials such as hydrogen, helium, water, ammonia, and methane. Out by Jupiter and beyond, where it's very cold, these materials form solid particles.

The nebular hypothesis was designed to explain some of the basic features of the solar system:

- The orbits of the planets lie in nearly the same plane with the Sun at the center
- The planets revolve in the same direction
- The planets mostly rotate in the same direction
- The axes of rotation of the planets are mostly nearly perpendicular to the orbital plane

- The oldest moon rocks are 4.5 billion years

This video, from the ESA, discusses the Sun, planets, and other bodies in the Solar System and how they formed **(1a, 1d)**. The first part of the video explores the evolution of our view of the solar system starting with the early Greeks who reasoned that since some points of light – which they called planets – moved faster than the stars, they must be closer: http://www.youtube.com/watch?v=-NxfBOhQ1CY (8:34).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=170

Lesson Summary

- The solar system is the Sun and all the objects that are bound to the Sun by gravity.
- The solar system has eight planets: Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune. Ceres, Makemake, Pluto and Eris are dwarf planets.
- The ancient Greeks and people for centuries afterwards believed in a geocentric model of the universe, with Earth at the center and everything else orbiting our planet.
- Copernicus, Kepler, and Galileo promoted a heliocentric model of the universe, with the Sun at the center and Earth and the other planets orbiting the Sun.
- Gravity holds planets in elliptical orbits around the Sun.
- The nebular hypothesis describes how the solar system formed from a giant cloud of gas and dust about 4.6 billion years ago.

Review Questions

1. What does geocentric mean?

2. Describe the geocentric model and heliocentric model of the universe.

3. How was Kepler's version of the heliocentric model different from Copernicus'?

4. Name the eight planets in order from the Sun outward. Which are the inner planets and which are the outer planets?

5. Compare and contrast the inner planets and the outer planets.

6. What object used to be considered a planet, but is now considered a dwarf planet? What are the other dwarf planets?

7. What keeps planets and moons in their orbits?

8. How old is the solar system? How old is Earth?

9. Use the nebular hypothesis to explain why the planets all orbit the Sun in the same direction.

Further Reading / Supplemental Links

- More information about the solar system from NASA: http://sse.jpl.nasa.gov/planets/index.cfm
- Lots of information about the solar system from the BBC: BBC Explore the solar system http://www.bbc.co.uk/solarsystem/
- Information about solar system objects: http://www.solarviews.com/eng/homepage.htm
- A multimedia tour of the solar system: http://www.nineplanets.org/
- Windows to the Universe: http://www.windows.ucar.edu/tour/link=/our_solar_system/formation.html
- Space news: http://www.space.com/

Points to Consider

- Would you expect all the planets in the solar system to be made of similar materials? Why or why not?
- The planets are often divided into two groups: the inner planets and the outer planets. Which planets do you think are in each of these two groups? What do members of each group have in common?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/25.1/. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 17.1: Introduction to the Solar System is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

17.2: Inner Planets

Lesson Objectives

- Describe key features of each of the inner planets.
- Compare each of the inner planets to Earth and to one another.

Vocabulary


- day
- inner planets
- terrestrial planets
- year

Introduction

What evidence do planetary geologists have to go on to determine the geology of the inner planets? On Earth, scientists can collect and analyze the chemistry of samples, do radiometric dating to determine their ages, and look at satellite images to see large-scale features. Rovers have landed on Mars and sent back enormous amounts of information but much of the rest of what is known about the inner planets is from satellite images.

The Inner Planets

The **inner planets**, or **terrestrial planets**, are the four planets closest to the Sun: Mercury, Venus, Earth, and Mars. **Figure** below shows the relative sizes of these four inner planets.



This composite shows the relative sizes of the four inner planets. From left to right, they are Mercury, Venus, Earth, and Mars.

Unlike the outer planets, which have many of satellites, Mercury and Venus do not have moons, Earth has one, and Mars has two. Of course, the inner planets have shorter orbits around the Sun, and they all spin more slowly. Geologically, the inner planets are all made of cooled igneous rock with iron cores, and all have been geologically active, at least early in their history. None of the inner planets has rings.

Earth

Although Earth is the third planet out from the Sun this lesson will start here. We know a lot more about Earth, so what we know can be used for comparison with the other planets.



What are Earth's most distinctive features? This famous image of Earth was taken during the Apollo 17 mission to the moon. Can you find a hurricane? A storm spinning in the opposite direction from the hurricane?

Earth's Surface and Life

As you can see in (**Figure** above), Earth has vast oceans of liquid water, large masses of exposed land, and a dynamic atmosphere with clouds of water vapor. Earth also has ice covering its polar regions. Earth's average surface temperature is 14°C (57°F). Water is a liquid at this temperature, but the planet also has water in its other two states, solid and gas. The oceans and the atmosphere help keep Earth's surface temperatures fairly steady.

As yet Earth is the only planet known to have life. The presence of liquid water, the ability of the atmosphere to filter out harmful radiation, and many other features make the planet uniquely suited to harbor life. Life and Earth now affect each other; for example, the evolution of plants allowed oxygen to enter the atmosphere in large enough quantities for animals to evolve. Although life has not been found elsewhere in the solar system, other planets or satellites may harbor primitive life forms. Life may also be found elsewhere in the universe.

Structure and Plate Tectonics


The heat that remained from the planet's accretion, gravitational compression, and radioactive decay allowed the Earth to melt, probably more than once. As it subsequently cooled, gravity pulled metal into the center to create the core. Heavier rocks formed the mantle and lighter rocks formed the crust.

Earth's crust is divided into tectonic plates, which move around on the surface because of the convecting mantle below. Movement of the plates causes other geological activity, such as earthquakes, volcanoes, and the formation of mountains. The locations of these features are mostly related to current or former plate boundaries. Earth is the only planet known to have plate tectonics.

Earth's Motions and Satellites

Earth rotates on its axis once per **day**, by definition. Earth orbits the Sun once every 365.24 days, which is defined as a **year**. Earth has one large moon, which orbits Earth once every 29.5 days, a period known as a month.

Earth's moon is the only large moon orbiting a terrestrial planet in the solar system. The Moon is covered with craters; it also has large plains of lava. The huge number of craters suggests that Moon's surface is ancient. There is evidence that the Moon formed when a large object — perhaps as large as the planet Mars — struck Earth in the distant past (**Figure** below).



Besides its Moon, Earth is orbited by a great deal of space debris, the remains of satellites, and rocket stages.


Mercury

The smallest planet, Mercury, is the planet closest to the Sun. Because Mercury is so close to the Sun, it is difficult to observe from Earth, even with a telescope. However, the Mariner 10 spacecraft, shown in **Figure** below, visited Mercury from 1974 to 1975. The MESSENGER spacecraft has been studying Mercury in detail since 2005. The craft is currently in orbit around the planet, where it is creating detailed maps. MESSENGER stands for Mercury Surface, Space Environment, Geochemistry and Ranging.



(a) Mariner 10 made three flybys of Mercury in 1974 and 1975. (b) A 2008 image of compiled from a flyby by MESSENGER.

As **Figure** below shows, the surface of Mercury is covered with craters, like Earth's moon. Ancient impact craters means that for billions of years Mercury hasn't changed much geologically. Also, with very little atmosphere, the processes of weathering and erosion do not wear down structures on the planet.



Mercury is covered with craters, like Earth's moon. MESSENGER has taken extremely detailed pictures of the planet's surface.

There are many images, movies and activities on the MESSENGER site: http://messenger.jhuapl.edu/index.php.

Short Year, Long Days

Mercury is named for the Roman messenger god, who could run extremely quickly, just as the planet moves very quickly in its orbit around the Sun. A year on Mercury — the length of time it takes to orbit the Sun — is just 88 Earth days.


Despite its very short years, Mercury has very long days. A day is defined as the time it takes a planet to turn on its axis. Mercury rotates slowly on its axis, turning exactly three times for every two times it orbits the Sun. Therefore, each day on Mercury is 57 Earth days long. In other words, on Mercury, a year is only a Mercury day and a half long!

Extreme Temperatures

Mercury is close to the Sun, so it can get very hot. However, Mercury has virtually no atmosphere, no water to insulate the surface, and it rotates very slowly. For these reasons, temperatures on the surface of Mercury vary widely. In direct sunlight, the surface can be as hot as 427°C (801°F). On the dark side, or in the shadows inside craters, the surface can be as cold as -183°C (-297°F)! Although most of Mercury is extremely dry, scientists think there may be a small amount of water in the form of ice at the poles of Mercury, in areas that never receive direct sunlight.

A Liquid Metal Core

Figure below shows a diagram of Mercury's interior. Mercury is one of the densest planets. It's relatively large, liquid core, made mostly of melted iron, takes up about 42% of the planet's volume.




Mercury contains a thin crust, a mantle, and a large, liquid core that is rich in iron.

Venus

Named after the Roman goddess of love, Venus is the only planet named after a female. Venus' thick clouds reflect sunlight well so Venus is very bright. When it is visible, Venus is the brightest object in the sky besides the Sun and the Moon. Because the orbit of Venus is inside Earth's orbit, Venus always appears close to the Sun. When Venus rises just before the Sun rises, the bright object is called the morning star. When it sets just after the Sun sets, it is the evening star.

Of the planets, Venus is most similar to Earth in size and density. Venus is also our nearest neighbor. The planet's interior structure is similar to Earth's with a large iron core and a silicate mantle (**Figure** below). But the resemblance between the two inner planets ends there.

Find out more about Venus at the following link: http://www.nasa.gov/worldbook/venus_worldbook.html.



Venus's interior is similar to Earth's.

Motion

Venus rotates in a direction opposite the other planets and opposite to the direction it orbits the Sun. This rotation is extremely slow, only one turn every 243 days. This is longer than a year on Venus—it takes Venus only 224 days to orbit the Sun.

Extreme Atmosphere

Venus is covered by a thick layer of clouds, as shown in pictures of Venus taken at ultraviolet wavelengths (**Figure** below).



This ultraviolet image from the Pioneer Venus Orbiter shows thick layers of clouds in the atmosphere of Venus.

Venus' clouds are not made of water vapor like Earth's clouds. Clouds on Venus are made mostly of carbon dioxide with a bit of sulfur dioxide — and they also contain corrosive sulfuric acid. Because carbon dioxide is greenhouse gas, the atmosphere traps heat from the Sun and creates a powerful greenhouse effect. Even though Venus is further from the Sun than Mercury, the greenhouse effect makes Venus the hottest planet. Temperatures at the surface reach 465°C (860°F). That's hot enough to melt lead.


The atmosphere of Venus is full of acid, its pressure is crushing, and the enormous amount of carbon dioxide causes runaway greenhouse effect**(4d)**: http://www.youtube.com/watch?v=HqFVxWfVtoo (2:05).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=172

The atmosphere of Venus is so thick that the atmospheric pressure on the planet's surface is 90 times greater than the atmospheric pressure on Earth's surface. The dense atmosphere totally obscures the surface of Venus, even from spacecraft orbiting the planet.

Venus's Surface


Since spacecraft cannot see through the thick atmosphere, radar is used to map Venus' surface. Many features found on the surface are similar to Earth and yet are very different. **Figure** below shows a topographical map of Venus produced by the Magellan probe using radar.



This false color image of Venus was made from radar data collected by the Magellan probe between 1990 and 1994. What features can you identify?

Orbiting spacecraft have used radar to reveal mountains, valleys, and canyons. Most of the surface has large areas of volcanoes surrounded by plains of lava. In fact, Venus has many more volcanoes than any other planet in the solar system and some of those volcanoes are very large.

Most of the volcanoes are no longer active, but scientists have found evidence that there is some active volcanism (**Figure** below). Think about what you know about the geology of Earth and what produces volcanoes. What does the presence of volcanoes suggest about the geology of Venus? What evidence would you look for to find the causes of volcanism on Venus?



This image of the Maat Mons volcano with lava beds in the foreground was generated by a computer from radar data. The reddish-orange color is close to what scientists think the color of sunlight would look like on the surface of Venus.

Venus also has very few impact craters compared with Mercury and the Moon. What is the significance of this? Earth has fewer impact craters than Mercury and the Moon too. Is this for the same reason that Venus has fewer impact craters?

It's difficult for scientists to figure out the geological history of Venus. The environment is too harsh for a rover to go there. It is even more difficult for students to figure out the geological history of a distant planet based on the information given here. Still we can piece together a few things.

On Earth, volcanism is generated because the planet's interior is hot. Much of the volcanic activity is caused by plate tectonic activity. But on Venus, there is no evidence of plate boundaries and volcanic features do not line up the way they do at plate boundaries.


Because the density of impact craters can be used to determine how old a planet's surface is, the small number of impact craters means that Venus' surface is young. Scientists think that there is frequent, planet-wide resurfacing of Venus with volcanism taking place in many locations. The cause is heat that builds up below the surface that has no escape until finally it destroys the crust and results in volcanoes.

Mars

Mars is the fourth planet from the Sun, and the first planet beyond Earth's orbit (**Figure** below). Mars is a quite different from Earth and yet more similar than any other planet. Mars is smaller, colder, drier, and appears to have no life, but volcanoes are common to both planets and Mars has many.

Mars is easy to observe so Mars has been studied more thoroughly than any other extraterrestrial planet. Space probes, rovers, and orbiting satellites have all yielded information to planetary geologists. Although no humans have ever set foot on Mars, both NASA and the European Space Agency have set goals of sending people to Mars sometime between 2030 and 2040.

Find out all you want to know about Mars at http://mars.jpl.nasa.gov/extreme/.



This image of Mars, taken by the Hubble Space Telescope in October, 2005, shows the planet's red color, a small ice cap on the south pole, and a dust storm.


A Red Planet

Viewed from Earth, Mars is reddish in color. The ancient Greeks and Romans named the planet after the god of war. But the surface is not red from blood but from large amounts of iron oxide in the soil.

The Martian atmosphere is very thin relative to Earth's and has much lower atmospheric pressure. Although the atmosphere is made up mostly of carbon dioxide, the planet has only a weak greenhouse effect so temperatures are only slightly higher than if the planet had no atmosphere.

Surface Features

Mars has mountains, canyons, and other features similar to Earth. Some of these surface features are amazing for their size! Olympus Mons is a shield volcano, similar to the volcanoes that make up the Hawaiian Islands. But Olympus Mons is also the largest mountain in the solar system (**Figure** below).



Olympus Mons is about 27 km (16.7 miles/88,580 ft) above the Martian surface, more than three times taller than Mount Everest. The volcano's base is about the size of the state of Arizona.

Mars also has the largest canyon in the solar system, Valles Marineris (**Figure** below).




Valles Marineris is 4,000 km (2,500 mi) long, as long as Europe is wide, and one-fifth the circumference of Mars. The canyon is 7 km (4.3 mi) deep. By comparison, the Grand Canyon on Earth is only 446 km (277 mi) long and about 2 km (1.2 mi) deep.

Mars has more impact craters than Earth, though fewer than the Moon. A video comparing geologic features on Mars and Earth is seen here: Mars tectonics video http://news.discovery.com/videos/space-3-questions-mars-tectonics.html

Is There Water on Mars?

Water cannot stay in liquid form on Mars because the atmospheric pressure is too low. However, there is a lot of water in the form of ice and even prominent ice caps (**Figure** below). Scientists also think that there is a lot of water ice present just under the Martian surface. This ice can melt when volcanoes erupt, and water can flow across the surface temporarily.



The north polar ice cap on Mars.

Scientists think that water once flowed over the Martian surface because there are surface features that look like water-eroded canyons (**Figure** below). The presence of water on Mars, even though it is now frozen as ice, suggests that it might have been possible for life to exist on Mars in the past.




The Mars rover collected rounded clumps of crystals that, on Earth, are known to form in water.

A video of the top five Phoenix Lander sites on Mars is seen here: http://news.discovery.com/videos/space-top-5-mars-phoenix-lander-images.html.

Two Martian Moons

Mars has two very small moons that are irregular rocky bodies (**Figure** below). Phobos and Deimos are named after characters in Greek mythology — the two sons of Ares, who followed their father into war. Ares is equivalent to the Roman god Mars.



Mars has two small moons, Phobos (left) and Deimos (right). Both were discovered in 1877 and are thought to be captured asteroids.

An animation of the moons orbiting Mars is seen here: http://commons.wikimedia.org/wiki/File:Orbits_of_Phobos_and_Deimos.gif.

KQED: Searching for Life on Mars

The Mars Science Laboratory was launched on November 26, 2011 and will search for any evidence that the Red Planet was once capable of supporting life. Curiosity is a car-sized rover that is scouring the red planet for clues; it landed in August 2012. Learn more at:http://science.kqed.org/quest/video/searching-for-life-on-mars/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=172

Lesson Summary

- The four inner planets, or terrestrial planets, have solid, rocky surfaces.
- Earth, the third planet from the Sun, is the only planet with large amounts of liquid water, and the only planet known to support life. Earth has a large round moon.
- Mercury is the smallest planet and is the closest to the Sun. With its extremely thin atmosphere, Mercury has a large temperature range. Like the Moon, it is covered with craters.
- Venus is the second planet from the Sun and the closest planet to Earth, in distance and in size. With its thick, corrosive atmosphere, the surface temperature is extremely high.
- Venus has mountainous areas, as well as volcanoes surrounded by plains of lava.
- Mars is the fourth planet from the Sun. Mars is reddish in color and has the largest mountain and the largest canyon in the solar system. It has two small moons.
- Water ice is found in the polar ice caps and under the surface of Mars.

Review Questions

1. Name the inner planets from the Sun outward. Then name them from smallest to largest.

2. Why do the temperatures on some planets vary widely? Why are some temperatures much less variable?

3. Why does Venus have higher temperatures than Mercury?

4. How are maps of Venus made?

5. Name two major ways in which Earth is unlike any other planet.

6. Why is Mars red?

7. Suppose you are planning a mission to Mars. Identify two places where you might be able to get water on the planet. Why is this important?

Further Reading / Supplemental Links

- The Jet Propulsion Lab home page has all the current and past missions with media and activities and great images:http://www.jpl.nasa.gov/.

- NASA Solar System Explorer, Mercury: http://solarsystem.nasa.gov/planetselector.cfm?Object=Mercury;http://solarsystem.jpl.nasa.gov/planets/profile.cfm?Object=Mercury&Display=Kids
- Google maps has Mars! http://www.google.com/mars/
- Home page of the Mars Exploration Rover Mission: http://marsrovers.jpl.nasa.gov/home/index.html
- A short video about Mercury: http://www.youtube.com/watch?v=U8-DTJpygyk
- A short video about Venus: http://www.youtube.com/watch?v=HqFVxWfVtoo
- A short video about Mars: http://www.youtube.com/watch?v=M-KfYEQUg2s

Points to Consider

- The first humans may reach Mars sometime in the next few decades. What conditions will they face? Why do you think we are going to Mars instead of Mercury or Venus?
- Why are the four inner planets called terrestrial planets? What might a planet be like if it weren't a terrestrial planet?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/25.2. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 17.2: Inner Planets is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

17.3: Outer Planets

Lesson Objectives

- Describe key features of the outer planets and their moons.
- Compare the outer planets to each other and to Earth.

Vocabulary


- Galilean moons
- gas giants
- Great Red Spot
- outer planets
- planetary rings

Introduction

The four outer planets are farther from the Sun as well as farther from Earth. They are much more difficult to learn about since they are very different from our home planet.

The Outer Planets

The four planets farthest from the Sun are the **outer planets**. **Figure** below shows the relative sizes of the outer planets and the Sun. These planets are much larger than the inner planets and are made primarily of gases and liquids, so they are also called **gas giants**.



This image shows the four outer planets and the Sun, with sizes to scale. From left to right, the outer planets are Jupiter, Saturn, Uranus, and Neptune.

The gas giants are made up primarily of hydrogen and helium, the same elements that make up most of the Sun. Astronomers think that hydrogen and helium gases comprised much of the solar system when it first formed. Since the inner planets didn't have enough mass to hold on to these light gases, their hydrogen and helium floated away into space. The Sun and the massive outer planets had enough gravity to keep hydrogen and helium from drifting away.

All of the outer planets have numerous moons. They all also have **planetary rings**, composed of dust and other small particles that encircle the planet in a thin plane.

Jupiter

Because Jupiter is so large, it reflects a lot of sunlight. Jupiter is extremely bright in the night sky; only the Moon and Venus are brighter (**Figure** below). This brightness is all the more impressive because Jupiter is quite far from the Earth — 5.20 AUs away. It takes Jupiter about 12 Earth years to orbit once around the Sun.




This image of Jupiter was taken by Voyager 2 in 1979. The colors were later enhanced to bring out more details.

Jupiter is named for the king of the gods in Roman mythology. The planet is enormous, the largest object in the solar system besides the Sun. Although Jupiter is over 1,300 times Earth's volume, it has only 318 times the mass of Earth. Like the other gas giants, it is much less dense than Earth.

Check out NASA's world book to learn more about Jupiter: http://www.nasa.gov/worldbook/jupiter_worldbook.html.

A Ball of Gas and Liquid

Astronauts trying to land a spaceship on the surface of Jupiter would find that there is no solid surface at all! Jupiter is made mostly of hydrogen, with some helium, and small amounts of other elements (**Figure** below).



Jupiter's atmosphere is composed of hydrogen and helium. Deeper within the planet, pressure compresses the gases into a liquid. Some evidence suggests that Jupiter may have a small rocky core of heavier elements at its center.

A Stormy Atmosphere

The upper layer of Jupiter's atmosphere contains clouds of ammonia (NH_3) in bands of different colors. These bands rotate around the planet, but also swirl around in turbulent storms. The **Great Red Spot** (**Figure** below) is an enormous, oval-shaped storm found south of Jupiter's equator. This storm is more than three times as wide as the entire Earth. Clouds in the storm rotate in a counterclockwise direction, making one complete turn every six days or so. The Great Red Spot has been on Jupiter for at least 300 years, since astronomers could first see the storm through telescopes. Do you think the Great Red Spot is a permanent feature on Jupiter? How could you know?



This image of Jupiter's Great Red Spot (upper right of image) was taken by the Voyager 1 spacecraft. The white storm just below the Great Red Spot is about the same diameter as Earth.


Jupiter's Moons and Rings

Jupiter has a very large number of moons — 63 have been discovered so far. Four are big enough and bright enough to be seen from Earth, using no more than a pair of binoculars. These moons — Io, Europa, Ganymede, and Callisto — were first discovered by Galileo in 1610, so they are sometimes referred to as the **Galilean moons** (**Figure** below). The Galilean moons are larger than the dwarf planets Pluto, Ceres, and Eris. Ganymede is not only the biggest moon in the solar system it is even larger than the planet Mercury!



This composite image shows the four Galilean moons and their sizes relative to the Great Red Spot. From top to bottom, the moons are Io, Europa, Ganymede, and Callisto. Jupiter's Great Red Spot is in the background. Sizes are to scale.

Scientists are particularly interested in Europa because it may be a place to find extraterrestrial life. What features might make a satellite so far from the Sun a candidate for life? Although the surface of Europa is a smooth layer of ice, there is evidence that there is an ocean of liquid water underneath (**Figure** below). Europa also has a continual source of energy — it is heated as it is stretched and squashed by tidal forces from Jupiter. Numerous missions have been planned to explore Europa, including plans to drill through the ice and send a probe into the ocean. However, no such mission has yet been attempted.




An enhanced color image of a portion of Europa's icy surface. The surface ice may have motions similar to plate tectonics on Earth.

In 1979, two spacecrafts — Voyager 1 and Voyager 2 — visited Jupiter and its moons. Photos from the Voyager missions showed that Jupiter has a ring system. This ring system is very faint, so it is difficult to observe from Earth.

Saturn

Saturn, shown in **Figure** below, is famous for its beautiful rings. Although all the gas giants have rings, only Saturn's can be easily seen from Earth. In Roman mythology, Saturn was the father of Jupiter.

Saturn's mass is about 95 times the mass of Earth, and its volume is 755 times Earth's volume, making it the second largest planet in the solar system. Saturn is also the least dense planet in the solar system. It is less dense than water. What would happen if you had a large enough bathtub to put Saturn in? Saturn would float! Saturn orbits the Sun once about every 30 Earth years.



This image of Saturn and its rings is a composite of pictures taken by the Cassini orbiter in 2008

Like Jupiter, Saturn is made mostly of hydrogen and helium gases in the outer layers and liquids at greater depths. The upper atmosphere has clouds in bands of different colors. These rotate rapidly around the planet, but there seems to be less turbulence and fewer storms on Saturn than on Jupiter. One interesting phenomena that has been observed in the storms on Saturn is the presence of thunder and lightning (see video, below). The planet likely has a small rocky and metallic core.

Cassini scientists waited years for the right conditions to produce the first movie that shows lightning on another planet — Saturn.http://saturn.jpl.nasa.gov/video/videodetails/?videoID=210 Lots more videos from the Cassini mission are indexed here:http://saturn.jpl.nasa.gov/video/

Saturn's Rings


In 1610 Galileo first observed Saturn's rings with his telescope, but he thought they might be two large moons, one on either side of the planet. In 1659, the Dutch astronomer Christian Huygens realized that the features were rings (**Figure** below).

Saturn's rings circle the planet's equator and appear tilted because Saturn itself is tilted about 27 degrees. The rings do not touch the planet.



A color-exaggerated mosaic of Saturn and its rings taken by Cassini as Saturn eclipses the Sun.

The Voyager 1 and 2 spacecraft in 1980 and 1981 sent back detailed pictures of Saturn, its rings, and some of its moons. Saturn's rings are made of particles of water and ice, with some dust and rocks (**Figure** below). There are several gaps in the rings that scientists think have originated because (1) the material was cleared out by the gravitational pull within the rings or (2) by the gravitational forces of Saturn and of moons outside the rings.



A close-up of Saturn's outer C ring showing areas with higher particle concentration and gaps.

The rings were likely formed by the breakup of one of Saturn's moons or from material that never accreted into the planet when Saturn originally formed.

An animation of dark spokes in Saturn's rings is seen here:http://en.Wikipedia.org/wiki/File:Saturn_ring_spokes_PIA11144_300px_secs15.5to23_20080926.ogv. The spokes appear seasonally and their origin is as yet unknown.

Saturn's Moons

Most of Saturn's moons are very small and only seven are large enough for gravity to have made them spherical. Only Titan is larger than Earth's Moon at about 1.5 times its size. Titan is even larger than the planet Mercury.

Scientists are interested in Titan because its atmosphere is similar to what Earth's was like before life developed. Nitrogen is dominant and methane is the second most abundant gas. Titan may have a layer of liquid water and ammonia under a layer of surface ice. Lakes of liquid methane (CH_4) and ethane (C_2H_6) are found on Titan's surface. Although conditions are similar enough to those of early Earth for scientists to speculate that extremely primitive life may exist on Titan, the extreme cold and lack of carbon dioxide make it unlikely (**Figure** below).



This composite image compares Saturn's largest moon, Titan (right) to Earth (left).

Uranus

Uranus (YOOR-uh-nuhs) is named for the Greek god of the sky (**Figure** below). From Earth, Uranus is so faint that it was unnoticed by ancient observers. William Herschel first discovered the planet in 1781.



Uranus is an icy blue-green ball.

Although Uranus is very large, it is extremely far away, about 2.8 billion km (1.8 billion mi) from the Sun. Light from the Sun takes about 2 hours and 40 minutes to reach Uranus. Uranus orbits the Sun once about every 84 Earth years.

Uranus has a mass about 14 times the mass of Earth, but it is much less dense than Earth. Gravity at the surface of Uranus is weaker than on Earth's surface so if you were at the top of the clouds on Uranus, you would weigh about 10% less than what you weigh on Earth.

An Icy Blue-Green Ball

Like Jupiter and Saturn, Uranus is composed mainly of hydrogen and helium, with an outer gas layer that gives way to liquid on the inside. Uranus has a higher percentage of icy materials, such as water, ammonia (NH_3), and methane (CH_4), than Jupiter and Saturn.


When sunlight reflects off Uranus, clouds of methane filter out red light, giving the planet a blue-green color. There are bands of clouds in the atmosphere of Uranus, but they are hard to see in normal light, so the planet looks like a plain blue ball.

The Sideways Planet

Most of the planets in the solar system rotate on their axes in the same direction that they move around the Sun. Uranus, though, is tilted on its side so its axis is almost parallel to its orbit. In other words, it rotates like a top that was turned so that it was spinning parallel to the floor. Scientists think that Uranus was probably knocked over by a collision with another planet-sized object billions of years ago.

Rings and Moons of Uranus

Uranus has a faint system of rings (**Figure** below). The rings circle the planet's equator, but because Uranus is tilted on its side, the rings are almost perpendicular to the planet's orbit.



This image from the Hubble Space Telescope shows the faint rings of Uranus. The planet is tilted on its side, so the rings are nearly vertical.

Uranus has 27 known moons and all but a few of them are named for characters from the plays of William Shakespeare. The five biggest moons of Uranus — Miranda, Ariel, Umbriel, Titania, and Oberon — are shown in **Figure** below.




These Voyager 2 photos have been resized to show the relative sizes of the five main moons of Uranus.

Neptune


Neptune, shown in **Figure** below, is the only major planet that can't be seen from Earth without a telescope. Scientists predicted the existence of Neptune before it was discovered because Uranus did not always appear exactly where it should appear. They knew that the gravitational pull of another planet beyond Uranus must be affecting Uranus' orbit.

Neptune was discovered in 1846, in the position that had been predicted, and it was named Neptune for the Roman god of the sea because of its bluish color.



This image of Neptune was taken by Voyager 2 in 1989. The Great Dark Spot seen on the left center in the picture has since disappeared, but a similar dark spot has appeared on another part of the planet.

In many respects, Neptune is similar to Uranus (**Figure** below). Neptune has slightly more mass than Uranus, but it is slightly smaller in size. Neptune is much farther from the Sun at nearly 4.5 billion km (2.8 billion mi). The planet's slow orbit means that it takes 165 Earth years to go once around the Sun.



Neptune's composition is that of a gas giant: (1) upper atmosphere, (2) atmosphere composed of hydrogen, helium and methane gas, (3) mantle of water, ammonia and methane ice, (4) core of rock and ice.

Extremes of Cold and Wind


Neptune's blue color is mostly because of frozen methane (CH_4). When Voyager 2 visited Neptune in 1986, there was a large dark-blue spot that scientists named the Great Dark Spot, south of the equator. When the Hubble Space Telescope took pictures of Neptune in 1994, the Great Dark Spot had disappeared but another dark spot had appeared north of the equator. Astronomers think that both of these spots represent gaps in the methane clouds on Neptune.

The changing appearance of Neptune is caused by its turbulent atmosphere. The winds on Neptune are stronger than on any other planet in the solar system, reaching speeds of 1,100 km/h (700 mi/h), close to the speed of sound. This extreme weather surprised astronomers, since the planet receives little energy from the Sun to power weather systems. Neptune is also one of the coldest places in the solar system. Temperatures at the top of the clouds are about -218°C (-360°F).

Neptune's Rings and Moons

Neptune has faint rings of ice and dust that may change or disappear in fairly short time frames.

Neptune has 13 known moons. Triton, shown in **Figure** below, is the only one of them that has enough mass to be spherical in shape. Triton orbits in the direction opposite to the orbit of Neptune. Scientists think Triton did not form around Neptune, but instead was captured by Neptune's gravity as it passed by.



This image Triton, Neptune's largest moon, was taken by Voyager 2 in 1989.

Fly by Neptune's moon Triton by watching this video: http://www.space.com/common/media/video/player.php?videoRef=mm32_SunDeath#playerTop

Lesson Summary

- The four outer planets are all gas giants made primarily of hydrogen and helium. They have thick gaseous outer layers and liquid interiors.
- The outer planets have numerous moons, as well as planetary rings.
- Jupiter, by far the largest planet in the solar system, has bands of different colored clouds, and a long-lasting storm called the Great Red Spot.
- Jupiter has more than 60 moons including the four largest, the Galilean moons.
- Europa has an ocean of liquid water under a layer of ice where life may have formed.

- Saturn is smaller than Jupiter but has a large system of beautiful rings.
- Titan's atmosphere is similar to early Earth's and the moon could harbor primitive life.
- Uranus and Neptune were discovered relatively recently since they are so far away.
- Uranus is tilted on its side, probably because of a past collision with a large object.
- Neptune is very cold and has strong winds. Its dark spots are storms in Neptune's atmosphere.

Review Questions

1. Name the outer planets a) in order from the Sun outward, b) from largest to smallest by mass, and c) from largest to smallest by size.

2. Why are the outer planets called gas giants?

3. How do the Great Red Spot and Great Dark Spot differ?

4. Name the Galilean moons, and explain why they have that name.

5. Why might Europa be a likely place to find extraterrestrial life?

6. What causes gaps in Saturn's rings?

7. Why are scientists interested in the atmosphere of Saturn's moon Titan?

8. What liquid is found on the surface of Titan?

9. Why is Uranus blue-green in color?

10. What is the name of Neptune's largest moon?

Further Reading / Supplemental Links

- Cool jobs! Planetary meteorologist: http://news.discovery.com/videos/discovery-news-2009-planetary-meteorologist.html
- About the Cassini Mission to Saturn: http://saturn.jpl.nasa.gov/
- NASA's world book, Jupiter: http://www.nasa.gov/worldbook/jupiter_worldbook.html
- NASA's planet selector: http://solarsystem.nasa.gov/planetselector.cfm
- Short videos of the planet Jupiter: http://www.youtube.com/watch?v=5iVw72sX3Bg;
- Video of Saturn: http://www.youtube.com/watch?v=iLXeUVCNoX8
- From the BBC Documentary, The Planets, Neptune: http://www.youtube.com/watch?v=29wfzotaBIg

Points to Consider

- The inner planets are small and rocky, while the outer planets are large and gaseous. Why might the planets have formed into these two groups?
- We have discussed the Sun, the planets, and the moons of the planets. What other objects can you think of that can be found in our solar system?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/25.3. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 17.3: Outer Planets is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

17.4: Other Objects in the Solar System

Lesson Objectives

- Locate and describe the asteroid belt.
- Explain where comets come from and what causes their tails.
- Differentiate between meteors, meteoroids, and meteorites.

Vocabulary

- asteroid
- asteroid belt
- comet
- dwarf planet
- Kuiper belt
- meteor
- meteor shower
- meteoroid

Introduction

When the solar system formed, most of the matter ended up in the Sun. Material spinning in a disk around the Sun clumped together into larger and larger pieces to form the eight planets. But some of the smaller pieces of matter never joined one of these larger bodies and are still out there in space.

Asteroids

Asteroids are very small, rocky bodies that orbit the Sun. “Asteroid” means “star-like,” and in a telescope, asteroids look like points of light, just like stars. Asteroids are irregularly shaped because they do not have enough gravity to become round. They are also too small to maintain an atmosphere and without internal heat they are not geologically active (**Figure** below). Collisions with other bodies may break up the asteroid or create craters on its surface.




In 1991, Asteroid 951 Gaspra was the first asteroid photographed at close range. Gaspra is a medium-sized asteroid, measuring about 19 by 12 by 11 km (12 by 7.5 by 7 mi).

Asteroid impacts have had dramatic impacts on the shaping of the planets, including Earth. Early impacts caused the planets to grow as they cleared their portions of space. An impact with an asteroid about the size of Mars caused fragments of Earth to fly into space and ultimately create the Moon. Asteroid impacts are linked to mass extinctions throughout Earth history.

The Asteroid Belt

Hundreds of thousands of asteroids have been discovered in our solar system. They are still being discovered at a rate of about 5,000 new asteroids per month. The majority of the asteroids are found in between the orbits of Mars and Jupiter, in a region called the **asteroid belt**, as shown in **Figure** below. Although there are many thousands of asteroids in the asteroid belt, their total mass adds up to only about 4% of Earth's moon.



The white dots in the figure are asteroids in the main asteroid belt. Other groups of asteroids closer to Jupiter are called the Hildas (orange), the Trojans (green), and the Greeks (also green).

Scientists think that the bodies in the asteroid belt formed during the formation of the solar system. The asteroids might have come together to make a single planet, but they were pulled apart by the intense gravity of Jupiter.

Near-Earth Asteroids

More than 4,500 asteroids cross Earth's orbit; they are near-Earth asteroids. Between 500 and 1,000 of these are over 1 km in diameter.

Any object whose orbit crosses Earth's can collide with Earth and many asteroids do. On average, each year a rock about 5–10 m in diameter hits Earth (**Figure** below). Since past asteroid impacts have been implicated in mass extinctions, astronomers are always on the lookout for new asteroids, and follow the known near-Earth asteroids closely, so they can predict a possible collision as early as possible.




A painting of what an asteroid a few kilometers across might look like as it strikes Earth.

Asteroid Missions

Scientists are interested in asteroids because they are representatives of the earliest solar system (**Figure** below). Eventually asteroids could be mined for rare minerals or for construction projects in space. A few missions have studied asteroids directly.

NASA's DAWN mission orbited asteroid Vesta from July 2011 to September 2012 and is on its way to meet dwarf planet Ceres in 2015.



The NEAR Shoemaker probe took this photo as it was about to land on 433 Eros in 2001.

KQED: Asteroid Hunters

Thousands of objects, including comets and asteroids, are zooming around our solar system; some could be on a collision course with Earth. QUEST explores how these Near Earth Objects are being tracked and what scientists are saying should be done to prevent a deadly impact. Learn more at: http://science.kqed.org/quest/video/asteroid-hunters/.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=176

Meteors

A **meteor**, such as in **Figure** below, is a streak of light across the sky. People call them shooting stars but they are actually small pieces of matter burning up as they enter Earth's atmosphere from space.



A meteor streaks across the sky.

Meteors are called **meteoroids** before they reach Earth's atmosphere. Meteoroids are smaller than asteroids and range from the size of boulders down to the size of tiny sand grains. Still smaller objects are called interplanetary dust. When Earth passes through a cluster of meteoroids, there is a **meteor shower**. These clusters are often remnants left behind by comet tails.

Meteorites


Although most meteors burn up in the atmosphere, larger meteoroids may strike the Earth's surface to create a meteorite. Meteorites are valuable to scientists because they provide clues about our solar system. Many meteorites are from asteroids that formed when the solar system formed (**Figure** below). A few meteorites are made of rocky material that is thought to have come from Mars when an asteroid impact shot material off the Martian surface and into space.



A lunar meteorite originates on the Moon and strikes Earth.

Comets

Comets are small, icy objects that have very elliptical orbits around the Sun. Their orbits carry them from the outer solar system to the inner solar system, close to the Sun (**Figure** below). Early in Earth's history, comets may have brought water and other substances to Earth during collisions.



The highly elliptical orbit of Kohoutek (red) relative to Earth's more circular orbit (blue) and the position of the Sun.

Comet tails form the outer layers of ice melt and evaporate as the comet flies close to the Sun. The ice from the comet vaporizes and forms a glowing coma, which reflects light from the Sun. Radiation and particles streaming from the Sun push this gas and dust into a long tail that always points away from the Sun (**Figure** below). Comets appear for only a short time when they are near the Sun, then seem to disappear again as they move back to the outer solar system.



Comet Hale-Bopp, also called the Great Comet of 1997, shone brightly for several months in 1997. The comet has two visible tails: a bright, curved dust tail and a fainter, straight tail of ions (charged atoms) pointing directly away from the Sun.

The time between one appearance of a comet and the next is called the comet's period. Halley's comet, with a period of 75 years, will next be seen in 2061. The first mention of the comet in historical records may go back as much as two millennia.

Where Comets Come From

Short-period comets, with periods of about 200 years or less, come from a region beyond the orbit of Neptune. The **Kuiper belt** (pronounced "KI-per") contains not only comets, but asteroids, and at least two dwarf planets.


Comets with periods as long as thousands or even millions of years come from a very distant region of the solar system called the Oort cloud, about 50,000–100,000 AU from the Sun (50,000–100,000 times the distance from the Sun to Earth).

Dwarf Planets

The **dwarf planets** of our solar system are exciting proof of how much we are learning about our solar system. With the discovery of many new objects in our solar system, in 2006, astronomers refined the definition of a planet. Their subsequent reclassification of Pluto to the new category dwarf planet stirred up a great deal of controversy. How the classification of Pluto has evolved is an interesting story in science. The question is: What is and is not a planet?


Pluto

From the time it was discovered in 1930 until the early 2000s Pluto was considered the ninth planet. When astronomers first located Pluto, the telescopes were not as good so Pluto and its moon, Charon, were seen as one much larger object (**Figure** below). With better telescopes, astronomers realized that Pluto was much smaller than they had thought.



Pluto and its moon Charon are actually two objects.

Better technology also allowed astronomers to discover many smaller objects like Pluto that orbit the Sun. One of them, Eris, discovered in 2005, is even larger than Pluto (**Figure** below).



Eris and its moon Dysnomia.

Even when it was considered a planet, Pluto was an oddball. Unlike the other outer planets in the solar system, which are all gas giants, it is small, icy, and rocky. With a diameter of about 2,400 km, it is only about one-fifth the mass of Earth's Moon. Pluto's orbit is tilted relative to the other planets and is shaped like a long, narrow ellipse. Pluto's orbit sometimes even passes inside Neptune's orbit.

In 1992 Pluto's orbit was recognized to be part of the Kuiper belt. With more than 200 million Kuiper belt objects, Pluto has failed the test of clearing other bodies out its orbit.

From what you've read above, do you think Pluto should be called a planet? Why are people hesitant to take away Pluto's planetary status?

In 2006, the International Astronomical Union decided that there were too many questions surrounding what could be called a planet and so refined the definition of a planet.

According to the new definition, a planet must:

- Orbit a star.
- Be big enough that its own gravity causes it to be shaped as a sphere.
- Be small enough that it isn't a star itself.
- Have cleared the area of its orbit of smaller objects.

A dwarf planet is an object that meets items the first three items in the list above, but not but not the fourth. Pluto is now called a dwarf planet, along with the objects Ceres, Makemake, and Eris.

According to the IAU, a dwarf planet must:

- Orbit a star.
- Have enough mass to be nearly spherical.
- Not have cleared the area around its orbit of smaller objects.
- Not be a moon.

A video showing why Pluto isn't a planet any more: http://www.youtube.com/watch?v=FqX2YdnwtRc.

Pluto has three moons of its own. The largest, Charon, is big enough that the Pluto-Charon system is sometimes considered to be a double dwarf planet (**Figure** below). Two smaller moons, Nix and Hydra, were discovered in 2005. But having moons is not enough to make an object a planet.

Ceres

Ceres is the largest object in the asteroid belt (**Figure** below). Before 2006, Ceres was considered the largest of the asteroids, with only about 1.3% of the mass of the Earth's Moon. But unlike the asteroids, Ceres has enough mass that its gravity causes it to be shaped like a sphere. Like Pluto, Ceres is rocky.


Is Ceres a planet? How does it match the criteria above? Ceres orbits the Sun, is round, and is not a moon. As part of the asteroid belt, its orbit is full of other smaller bodies, so Ceres fails the fourth criterion for being a planet.



This composite image compares the size of the dwarf planet Ceres to Earth and the Moon.

Makemake

Makemake is the third largest and second brightest dwarf planet we have discovered so far (**Figure** below). With a diameter estimated to be between 1,300 and 1,900 km, it is about three-quarters the size of Pluto. Makemake orbits the Sun in 310 years at a distance between 38.5 to 53 AU. It is thought to be made of methane, ethane, and nitrogen ices.



Largest Known Trans-Neptunian Objects. Makemake is named after the deity that created humanity in the mythology of the people of Easter Island.

Eris

Eris is the largest known dwarf planet in the solar system — about 27% more massive than Pluto. The object was not discovered until 2003 because it is about three times farther from the Sun than Pluto, and almost 100 times farther from the Sun than Earth is. For a short time Eris was considered the “tenth planet” in the solar system, but its discovery helped to prompt astronomers to better define planets and dwarf planets in 2006. Eris also has a small moon, Dysnomia that orbits it once about every 16 days.

Astronomers know there may be other dwarf planets in the outer reaches of the solar system. Haumea was made a dwarf planet in 2008 and so now the total is five. Quaoar, Varuna and Orcus may be added to the list of dwarf planets in the future. We still have a lot to discover and explore.

Lesson Summary

- Asteroids are irregularly shaped, rocky bodies that orbit the Sun. Most are found in the asteroid belt, between the orbits of Mars and Jupiter.

- Meteoroids are smaller than asteroids, ranging from the size of boulders to the size of sand grains. When meteoroids enter Earth's atmosphere, they vaporize, creating a trail of glowing gas called a meteor. If any of the meteoroid reaches Earth, it is a meteorite.
- Comets are small, icy objects that have very elliptical orbits. When they are close to the Sun, they form comas and tails, which glow and make the comet more visible.
- Short-period comets come from the Kuiper belt, beyond Neptune. Long-period comets come from the very distant Oort cloud.
- Dwarf planets are spherical bodies that orbit the Sun, but that have not cleared their orbit of smaller bodies. Ceres is a dwarf planet in the asteroid belt. Pluto, Makemake and Eris are dwarf planets in the Kuiper belt.

Review Questions

1. Arrange the following from smallest to largest: asteroid, star, meteoroid, planet, dwarf planet.

2. Where are most asteroids found?

3. What is the difference between a meteor, a meteoroid, and a meteorite?

4. Why are meteorites extremely valuable to scientists?

5. What objects would scientists study to learn about the composition of the Oort cloud?

6. Why is Pluto no longer considered a planet?

7. Name the four known dwarf planets in our solar system.

Further Reading / Supplemental Links

- NASA worldbook on asteroids: http://www.nasa.gov/worldbook/asteroid_worldbook.html

Points to Consider

- In 2006, astronomers changed the definition of a planet and created a new category of dwarf planets. Do you think planets, dwarf planets, moons, asteroids, and meteoroids are clearly separate groups?
- What defines each of these groups, and what do objects in these different groups have in common? Could an object change from being in one group to another? How?
- We have learned about many different kinds of objects that are found within our solar system. What objects or systems of objects can you think of that are found outside our solar system?

CC licensed content, Shared previously

- **Provided by**: CK12.org. **Located at**: http://www.ck12.org/book/CK-12-Earth-Science-For-High-School/section/25.4. **License**: *CC BY-NC: Attribution-NonCommercial*

This page titled 17.4: Other Objects in the Solar System is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

18: Climate and Climate Change

18.1: Climate Systems and Change

18.2: Climate and Weather

18.3: Global Climate Change

This page titled 18: Climate and Climate Change is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

18.1: Climate Systems and Change

LEARNING OUTCOMES

- **ESSENTIAL QUESTIONS**
  - In what ways does climate influence regional culture?
  - What impacts could a changing climate have on local, regional, and international social, economic, and geopolitical stability?
  - Should global environmental issues like climate change be resolved by individuals, nations, or international institutions like the United Nations?
  - What is your worldview on global environmental issues and how they should be addressed?

Controls of Climate




Although almost anything can happen with the weather, climate is more predictable. The weather on a particular winter day in San Diego may be colder than on the same day in Lake Tahoe, but, on average, Tahoe's winter climate is significantly colder than San Diego's. Climate then is the long-term average of weather. Good climate is why we choose to vacation in Hawaii in February, even though the weather is not guaranteed to be good!

Climate is the average of weather in that location over a long period of time, usually for at least 30 years. A location's climate can be described by its air temperature, humidity, wind speed and direction, and the type, quantity, and frequency of precipitation. Climate can change, but only over long periods of time. The climate of a region depends on its position relative to many things. These factors are described in the next sections.

LATITUDE

The main factor influencing the climate of a region is latitude because different latitudes receive different amounts of solar radiation. To review from the Earth's Atmosphere chapter:

- The equator receives the most solar radiation. Days are equally long year-round and the sun is just about directly overhead at midday.
- The polar regions receive the least solar radiation. The night lasts six months during the winter. Even in summer, the sun never rises very high in the sky. Sunlight filters through a thick wedge of atmosphere, making the sunlight much less intense. The high albedo, because of ice and snow, reflects a good portion of the sun's light.



ATMOSPHERIC CIRCULATION

Recall from the Earth's Atmosphere chapter the circulation cells and global wind belts**.** The position of a region relative to the circulation cells and wind belts has a great effect on its climate. In an area where the air is mostly rising or sinking, there is not much wind.

INTERTROPICAL CONVERGENCE ZONE

The **Intertropical Convergence Zone (ITCZ)** is the low pressure area near the equator in the boundary between the two Hadley Cells. The air rises so that it cools and condenses to create clouds and rain. Climate along the ITCZ is therefore warm and wet. Early mariners called this region the doldrums because their ships were often unable to sail because there were no steady winds.

The ITCZ migrates slightly with the season. Land areas heat more quickly than the oceans. Because there are more land areas in the Northern Hemisphere, the ITCZ is influenced by the heating effect of the land. In Northern Hemisphere summer, it is approximately 5o north of the equator while in the winter it shifts back and is approximately at the equator. As the ITCZ shifts, the major wind belts also shift slightly north in summer and south in winter, which causes the wet and dry seasons in this area

.

HADLEY CELL AND FERRELL CELL BOUNDARIES

At about 30 degrees N and 30 degrees S, the air is fairly warm and dry because much of it came from the equator where it lost most of its moisture at the ITCZ. At this location the air is descending, and sinking air warms and causes evaporationMariners named this region the horse latitudes. Sailing ships were sometimes delayed for so long by the lack of wind that they would run out of water and food for their livestock. Sailors tossed horses and other animals over the side after they died. Sailors sometimes didn't make it either.

PREVAILING WINDS

The prevailing winds are the bases of the Hadley, Ferrell, and Polar Cells. These winds greatly influence the climate of a region because they bring the weather from the locations they come from. For example, in California, the predominant winds are the westerlies blowing in from the Pacific Ocean, which bring in relatively cool air in summer and relatively warm air in winter. Local winds also influence local climate. For example, land breezes and sea breezes moderate coastal temperatures.

CONTINENTAL POSITION

When a particular location is near an ocean or large lake, the body of water plays an extremely important role in affecting the region's climate.

- A **maritime climate** is strongly influenced by the nearby sea. Temperatures vary a relatively small amount seasonally and daily. For a location to have a true maritime climate, the winds must most frequently come off the sea.
- A **continental climate** is more extreme, with greater temperature differences between day and night and between summer and winter.


The ocean's influence in moderating climate can be seen in the following temperature comparisons. Each of these cities is located at 37oN latitude, within the westerly winds**.**



Maritime climate



Continental climate



OCEAN CURRENTS

The temperature of the water offshore influences the temperature of a coastal location, particularly if the winds come off the sea. The cool waters of along the western United States is caused by a clockwise rotating ocean current that is bringing cold water from the arctic toward the equator. The climatic effect is that coastal regions of California, Oregon, and Washington are are cool. Coastal upwelling also brings cold, deep water up to the ocean surface off of California, which contributes to the cool coastal temperatures. But that same ocean current brings warm, tropical water to eastern Japan.In the Atlantic Ocean, the northern ocean current, called the **Gulf Stream**, brings warm water from the tropics to the southern states. This is a major reason why the southern states experience humid conditions in the summer and tornadoes because of all this warm moisture. The Gulf Stream also impacts Europe by bringing warm water northward, making this region that is rather northward warmer than expected.



ALTITUDE AND MOUNTAIN RANGES

Air pressure and air temperature decreases with altitude. The closer molecules are packed together, the more likely they are to collide. Collisions between molecules give off heat, which warms the air. At higher altitudes, the air is less dense and air molecules are more spread out and less likely to collide. A location in the mountains has lower average temperatures than one at the base of the mountains. In Colorado, for example, Lakewood (5,640 feet) average annual temperature is 62 degrees F (17 degrees C), while Climax Lake (11,300 feet) is 42 degrees F (5.4 degrees C). Mountain ranges have two effects on the climate of the surrounding region. The first is something called the **rainshadow effect**, which brings warm dry climate to the leeward size of a mountain range, was described in the Earth's Atmosphere chapter**.**The second effect mountains have on climate systems is the ability to separate coastal regions from the rest of the continent. Since a maritime air mass may have trouble rising over a mountain range, the coastal area will have a maritime climate but the inland area on the leeward side will have a continental climate.

Climate Zones and Biomes


A **climate zone** results from the climate conditions of an area: its temperature, humidity, amount and type of precipitation, and the season. A climate zone is reflected in a region's natural vegetation. Perceptive travelers can figure out which climate zone they are in by looking at the vegetation, even if the weather is unusual for the climate on that day.

The major factors that influence climate determine the different climate zones. In general, the same type of climate zone will be found at similar latitudes and in similar positions on nearly all continents, both in the Northern and Southern Hemispheres. The one exception to this pattern is the climate zones called the continental climates, which are not found at higher latitudes in the Southern Hemisphere. This is because the Southern Hemisphere land masses are not wide enough to produce a continental climate.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

The most common system used to classify climatic zones is the **Köppen classification system**. This system is based on the *temperature*, the amount of *precipitation*, and the times of year when precipitation occurs. Since climate determines the type of vegetation that grows in an area, vegetation is used as an indicator of climate type.

A climate type and its plants and animals make up a **biome**. The organisms of a biome share certain characteristics around the world, because their environment has similar advantages and challenges. The organisms have adapted to that environment in similar ways over time. For example, different species of cactus live on different continents, but they have adapted to the harsh desert in similar ways**.**



An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

The Köppen classification system recognizes five major climate groups, each with a distinct capital letter A through E. Each lettered group is divided into subcategories. Some of these subcategories are forest (f), monsoon (m), and wet/dry (w) types, based on the amount of precipitation and season when that precipitation occurs**.**

Tropical Moist Climates (Group A)



Tropical Moist (Group A) climates are found in a band about 15o to 25o N and S of the equator. What climate characteristics is the tropical moist climate group likely to have?

- Temperature: Intense sunshine; each month has an average temperature of at least 18oC (64oF).
- Rainfall: Abundant, at least 150 cm (59 inches) per year.

The wet tropics have almost no annual temperature variation and tremendous amounts of rain fall year round, between 175 and 250 cm (65 and 100 inches). These conditions support the tropical rainforest biome. **Tropical rainforests** are dominated by densely packed, broadleaf evergreen trees. These rainforests have the highest number of species or **biodiversity** of any ecosystem.

TROPICAL MONSOON (Am)

The tropical monsoon climate has very low precipitation for one to two months each year. Rainforests grow here because the dry period is short, and the trees survive off of soil moisture. This climate is found where the monsoon winds blow, primarily in southern Asia, western Africa, and northeastern South America**.**

TROPICAL WET AND DRY

The tropical wet and dry climate lies between about 5o and 20o latitude, around the location of the ITCZ. In the summer, when the ITCZ drifts northward, the zone is wet. In the winter, when the ITCZ moves toward the equator, the region is dry. This climate exists where strong monsoon winds blow from land to sea, such as in India.

Rainforests cannot survive the months of low rainfall, so the typical vegetation is **savanna.** This biome consists mostly of grasses, with widely scattered deciduous trees and rare areas of denser forests.

Dry Climates (Group B)



The Dry Climates (Group B) have less precipitation than evaporation. Dry climate zones cover about 26% of the world's land area**.** What climate characteristics is the dry climate group likely to have?

- Temperature: Abundant sunshine. Summer temperatures are high; winters are cooler and longer than Tropical Moist climates
- Rainfall: Irregular; several years of drought are often followed by a single year of abundant rainfall

ARID DESERT (Bw)

Low-latitude, arid deserts are found between 15o to 30o N and S latitudes. This is where warm dry air sinks at high pressure zones. True deserts make up around 12% of the world's lands**.**

In the Sonoran Desert of the southwestern United States and northern Mexico, skies are clear. The typical weather is extremely hot summer days and cold winter nights. Although annual rainfall is less than 25 cm (10 inches), rain falls during two seasons. Pacific storms bring winter rains and monsoons bring summer rains. Since organisms do not have to go too many months without some rain, a unique group of plants and animals can survive in the Sonoran desert.



SEMI-ARID OR STEPPE (Bs)

Higher latitude semi-arid deserts, also called **steppe**, are found in continental interiors or in rainshadows. Semi-arid deserts receive between 20 and 40 cm (8 to 16 inches) of rain annually. The annual temperature range is large. In the United States, the Great Plains, portions of the southern California coast, and the Great Basin are semi-arid deserts.

Moist Subtropical Mid-latitude (Group C)



What climate characteristics is the moist subtropical group likely to have?

- Temperature: The coldest month ranges from just below freezing to almost balmy, between -3oC and 18oC (27o to 64oF). Summers are mild with average temperatures above 10oC (50oF). Seasons are distinct.
- Rainfall: There is plentiful annual rainfall.

DRY SUMMER SUBTROPICAL OR MEDITERRANEAN CLIMATES (Cs)

The Dry Summer Subtropical climate is found on the western sides of continents between 30o and 45o latitude. Annual rainfall is 30 to 90 cm (14 to 35 inches), most of which comes in the winter.

The climate is typical of coastal California, which sits beneath a summertime high pressure for about five months each year. Land and sea breezes make winters moderate and summers cool. Vegetation must survive long summer droughts. The scrubby, woody vegetation that thrives in this climate is called **chaparral**.



HUMID SUBTROPICAL (Cfa)

The Humid Subtropical climate zone is found mostly on the eastern sides of continents**.** Rain falls throughout the year with annual averages between 80 and 165 cm (31 and 65 inches). Summer days are humid and hot, from the lower 30's up to 40oC (mid-80's up to 104oF). Afternoon and evening thunderstorms are common. These conditions are caused by warm tropical air passing over the hot continent. Winters are mild, but middle-latitude storms called cyclones may bring snow and rain. The southeastern United States, with its hot humid summers and mild, but frosty winters, is typical of this climate zone.

MARINE WEST COAST CLIMATE (Cfb)

This climate lines western North America between 40o and 65o latitude, an area known as the Pacific Northwest**.** Ocean winds bring mild winters and cool summers. The temperature range, both daily and annually, is fairly small. Rain falls year round, although summers are drier as the jet stream moves northward. Low clouds, fog, and drizzle are typical. In Western Europe the climate covers a larger region since no high mountains are near the coast to block wind blowing off the Atlantic.

Humid Continental (Group D)

Continental (Group D) climates are found in most of the North American interior from about 40°N to 70°N. What climate characteristics is the continental group most likely to have?

- Temperature: The average temperature of the warmest month is higher than 10°C (50°F) and the coldest month is below -3°C (-27°F).
- Precipitation: Winters are cold and stormy (look at the latitude of this zone and see if you can figure out why). Snowfall is common and snow stays on the ground for long periods of time.

Trees grow in continental climates, even though winters are extremely cold, because the average annual temperature is fairly mild. Continental climates are not found in the Southern Hemisphere because of the absence of a continent large enough to generate this effect.

HUMID CONTINENTAL (Dfa, Dfb)

The humid continental climates are found around the polar front in North America and Europe (**Figure** below). In the winter, middle-latitude cyclones bring chilly temperatures and snow. In the summer, westerly winds bring continental weather and warm temperatures. The average July temperature is often above 20oC (70oF). The region is typified by deciduous trees, which protect themselves in winter by losing their leaves.

The two variations of this climate are based on summer temperatures.

- Dfa, long, hot summers: summer days may be over 38oC (100oF), nights are warm and the temperature range is large, perhaps as great as 31oC (56oF). The long summers and high humidity foster plant growth**.**
- Dfb, long, cool summers: summer temperatures and humidity are lower. Winter temperatures are below -18oC (0oF) for long periods.



SUBPOLAR (Dfc)

The subpolar climate is dominated by the continental polar air that masses over the frigid continent. Snowfall is light, but cold temperatures keep snow on the ground for months. Most of the approximately 50 cm (20 inches) of annual precipitation falls during summer cyclonic storms. The angle of the Sun's rays is low but the Sun is visible in the sky for most or all of the day during

the summer, so temperatures may get warm, but are rarely hot. These continental regions have extreme annual temperature ranges. The boreal, coniferous forests found in the subpolar climate are called **taiga** and have small, hardy, and widely spaced trees. Taiga vast forests stretch across Eurasia and North America.

Polar Climates (Group E)



Polar climates are found across the continents that border the Arctic Ocean, Greenland, and Antarctica. What climate characteristics is the polar climate group most likely to have?

- Temperature: Winters are entirely dark and bitterly cold. Summer days are long, but the sun is low on the horizon so summers are cool. The average temperature of the warmest month at less than 10oC (50oF). The annual temperature range is large.
- Precipitation: The region is dry with less than 25 cm (10 inches) of precipitation annually; most precipitation occurs during the summer.

POLAR TUNDRA (ET)

The polar tundra climate is continental, with severe winters. Temperatures are so cold that a layer of permanently frozen ground, called **permafrost** forms below the surface. This frozen layer can extend hundreds of meters deep. The average temperature of the warmest months is above freezing, so summer temperatures defrost the uppermost portion of the permafrost. In winter, the permafrost prevents water from draining downward. In summer, the ground is swampy. Although the precipitation is low enough in many places to qualify as a desert, evaporation rates are also low, so the landscape receives more usable water than a desert.

Because of the lack of ice-free land near the South Pole, there is very little tundra in the Southern Hemisphere (**Figure** below). The only plants that can survive the harsh winters and soggy summers are small ground-hugging plants like mosses, lichens, small shrubs, and scattered small trees that make up the **tundra**.



ICE CAP

Ice caps are found mostly on Greenland and Antarctica, about 9% of the Earth's land area**.** Ice caps may be thousands of meters thick. Ice cap areas have extremely low average annual temperatures, e.g. -29oC (-20oF) at Eismitte, Greenland. Precipitation is low because the air is too cold to hold much moisture. Snow occasionally falls in the summer.

Hightland Climates (Group H)



When climate conditions in a small area are different from those of the surroundings, the climate of the small area is called a **microclimate**. The microclimate of a valley may be cool relative to its surroundings since cold air sinks. The ground surface may be hotter or colder than the air a few feet above it, because rock and soil gain and lose heat readily. Different sides of a mountain will have different microclimates. In the Northern Hemisphere, a south-facing slope receives more solar energy than a north-facing slope, so each side supports different amounts and types of vegetation.

Altitude mimics latitude in climate zones**.** Climates and biomes typical of higher latitudes may be found in other areas of the world at high altitudes.

Climate Change in Earth History

For the past two centuries, climate has been relatively stable. People placed their farms and cities in locations that were in a favorable climate without thinking that the climate could change. But climate has changed throughout Earth history, and a stable climate is not the norm. In recent years, Earth's climate has begun to change again. Most of this change is warming because of human activities that release greenhouse gases into the atmosphere. The effects of warming are already being seen and will become more extreme as temperature rise.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180



Climate has changed throughout Earth history. Much of the time Earth's climate was hotter and more humid than it is today, but climate has also been colder, as when glaciers covered much more of the planet. The most recent ice ages were in the Pleistocene Epoch, between 1.8 million and 10,000 years ago. Glaciers advanced and retreated in cycles, known as glacial and interglacial periods. With so much of the world's water bound into the ice, sea level was about 125 meters (395 feet) lower than it is today. Many scientists think that we are now in a warm, interglacial period that has lasted about 10,000 years.

For the past 2,000 years, climate has been relatively mild and stable when compared with much of Earth's history. Why has climate stability been beneficial for human civilization? Stability has allowed the expansion of agriculture and the development of towns and cities.


Fairly small temperature changes can have major effects on global climate. The average global temperature during glacial periods was only about 5.5 degrees C (10 degrees F) less than Earth's current average temperature. Temperatures during the interglacial

periods were about 1.1 degrees C (2.0 degrees F) higher than today.

Since the end of the Pleistocene, the global average temperature has risen about 4 degrees C (7 degrees F). Glaciers are retreating and sea level is rising. While climate is getting steadily warmer, there have been a few more extreme warm and cool times in the last 10,000 years. Changes in climate have had effects on human civilization.

- The Medieval Warm Period from 900 to 1300 A.D. allowed Vikings to colonize Greenland and Great Britain to grow wine grapes.
- The Little Ice Age, from the 14th to 19th centuries, the Vikings were forced out of Greenland and humans had to plant crops further south.

Short-term Climate Changes



Short-term changes in climate are common. The largest and most important of these is the oscillation between El Niño and La Niña conditions. This cycle is called the ENSO (El Niño southern oscillation). The ENSO drives changes in climate that are felt around the world about every two to seven years.

In a normal year, the trade winds blow across the Pacific Ocean near the equator from east to west (toward Asia). A low pressure cell rises above the western equatorial Pacific. Warm water in the western Pacific Ocean and raises sea levels by one-half meter. Along the western coast of South America, the Peru Current carries cold water northward, and then westward along the equator with the trade winds. Upwelling brings cold, nutrient-rich waters from the deep sea.

In an **El Niño** year, when water temperature reaches around 28 degrees C (82 degrees F), the trade winds weaken or reverse direction and blow east (toward South America). Warm water is dragged back across the Pacific Ocean and piles up off the west coast of South America. With warm, low-density water at the surface, upwelling stops. Without upwelling, nutrients are scarce and plankton populations decline. Since plankton form the base of the food web, fish cannot find food, and fish numbers decrease as well. All the animals that eat fish, including birds and humans, are affected by the decline in fish.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

By altering atmospheric and oceanic circulation, El Niño events change global climate patterns.

- Some regions receive more than average rainfall, including the west coast of North and South America, the southern United States, and Western Europe.
- Drought occurs in other parts of South America, the western Pacific, southern and northern Africa, and southern Europe.

An El Niño cycle lasts one to two years. Often normal circulation patterns resume. Sometimes circulation patterns bounce back quickly and extremely, called **La Niña**.

In a La Niña year, as in a normal year, trade winds moves from east to west and warm water piles up in the western Pacific Ocean. Ocean temperatures along coastal South America are colder than normal (instead of warmer, as in El Niño). Cold water reaches farther into the western Pacific than normal.

Other important oscillations are smaller and have a local, rather than global, effect. The North Atlantic Oscillation mostly alters climate in Europe. The Mediterranean also goes through cycles, varying between being dry at some times, and warm and wet at others.

Causes of Long-term Climate Change

Many processes can cause climate to change. These include changes in the amount of energy the Sun produces over years; the positions of the continents over millions of years; in the tilt of Earth's axis; orbit over thousands of years; that are sudden and dramatic because of random catastrophic events, such as a large asteroid impact; in greenhouse gases in the atmosphere, caused naturally or by human activities.




PLATE TECTONICS

Plate tectonic movements can alter climate. Over millions of years as seas open and close, ocean currents may distribute heat differently. For example, when all the continents are joined into one supercontinent (such as Pangaea), nearly all locations experience a continental climate. When the continents separate, heat is more evenly distributed.Plate tectonic movements may help start an ice age. When continents are located near the poles, ice can accumulate, which may increase albedo and lower global temperature. Low enough temperatures may start a global ice age.

Plate motions trigger volcanic eruptions, which release dust and CO2 into the atmosphere. Ordinary eruptions, even large ones, have only a short-term effect on weather. Massive eruptions of the fluid lavas that create lava plateaus release much more gas and dust, and can change climate for many years. This type of eruption is exceedingly rare; none has occurred since humans have lived on Earth.

MILANKOVITCH CYCLES


The most extreme climate of recent Earth history was the Pleistocene. Scientists attribute a series of ice ages to variation in the Earth's position relative to the Sun, known as **Milankovitch cycles**. The Earth goes through regular variations in its position relative to the Sun:

The shape of the Earth's orbit changes slightly as it goes around the Sun. The orbit varies from more circular to more elliptical in a cycle lasting between 90,000 and 100,000 years. When the orbit is more elliptical, there is a greater difference in solar radiation between winter and summer.The planet wobbles on its axis of rotation. At one extreme of this 27,000 year cycle, the Northern Hemisphere points toward the Sun when the Earth is closest to the Sun. Summers are much warmer and winters are much colder than now. At the opposite extreme, the Northern Hemisphere points toward the Sun when it is farthest from the Sun. This results in chilly summers and warmer winters.


The planet's tilt on its axis varies between 22.1 degrees and 24.5 degrees. Seasons are caused by the tilt of Earth's axis of rotation, which is at a 23.5o angle now. When the tilt angle is smaller, summers and winters differ less in temperature. This cycle lasts 41,000 years.

When these three variations are charted out, a climate pattern of about 100,000 years emerges. Ice ages correspond closely with Milankovitch cycles. Since glaciers can form only over land, ice ages only occur when landmasses cover the polar regions.


Therefore, Milankovitch cycles are also connected to plate tectonics.




Eccentricity – circular




Eccentricity – elliptical



Precession




Obliquity



SUN VARIATION

The amount of energy the Sun radiates is variable. **Sunspots** are magnetic storms on the Sun's surface that increase and decrease over an 11-year cycle. When the number of sunspots is high, solar radiation is also relatively high. But the entire variation in solar radiation is tiny relative to the total amount of solar radiation that there is and there is no known 11-year cycle in climate variability. The Little Ice Age corresponded to a time when there were no sunspots on the Sun.



CHANGES IN ATMOSPHERIC GREENHOUSE GAS LEVELS

Since greenhouse gases trap the heat that radiates off the planet's surfaces what would happen to global temperatures if atmospheric greenhouse gas levels decreased? What if greenhouse gases increased? A decrease in greenhouse gas levels decreases global temperature and an increase raises air temperature.Greenhouse gas levels have varied throughout Earth history. For example, CO2 has been present at concentrations less than 200 parts per million (ppm) and more than 5,000 ppm. But for at least 650,000 years, CO2 has never risen above 300 ppm, during either glacial or interglacial periods. Natural processes add (volcanic eruptions and the decay or burning of organic matter) and remove absorption by plants, animal tissue, and the ocean) CO2 from the atmosphere. When plants are turned into fossil fuels the CO2 in their tissue is stored with them. So CO2 is removed from the atmosphere. What does this do to Earth's average temperature?

Fossil fuel use has skyrocketed in the past few decades more people want more cars and industrial products. This has released CO2 into the atmosphere. Burning tropical rainforests, to clear land for agriculture, a practice called **slash-and-burn agriculture**, also increases atmospheric CO2. By cutting down trees, they can no longer remove CO2 from the atmosphere. Burning the trees releases all the CO2 stored in the trees into the atmosphere.

There is now nearly 40% more CO2 in the atmosphere than there was 200 years ago, before the Industrial Revolution. About 65% of that increase has occurred since the first CO2 measurements were made on Mauna Loa Volcano, Hawaii, in 1958. CO2 is the most important greenhouse gas that human activities affect because it is so abundant. But other greenhouse gases are increasing as well. A few are:

- Methane: released from raising livestock, rice production, and the incomplete burning of rainforest plants.
- Chlorofluorocarbons (CFCs): human-made chemicals that were invented and used widely in the 20th century.
- Tropospheric ozone: from vehicle exhaust, it has more than doubled since 1976.

Anthropogenic Climate Change

TEMPERATURES ARE RISING

With more greenhouse gases trapping heat, average annual global temperatures are rising. This is known as **global warming**. While temperatures have risen since the end of the Pleistocene, 10,000 years ago, this rate of increase has been more rapid in the past century, and has risen even faster since 1990. The nine warmest years on record have all occurred since 1998, and the 10 of the 11 warmest years have occurred since 2001 (through 2012). The 2000s were the warmest decade yet. Annual variations aside, the average global temperature increased about 0.8 degrees C (1.5 degrees F) between 1880 and 2010, according to the Goddard Institute for Space Studies, NOAA.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

The United States has long been the largest emitter of greenhouse gases, with about 20% of total emissions in 2004. As a result of China's rapid economic growth, its emissions surpassed those of the United States in 2008. However, it's also important to keep in mind that the United States has only about one-fifth the population of China. What's the significance of this? The average United States citizen produces far more greenhouse gases than the average Chinese person.

If nothing is done to decrease the rate of CO2 emissions, by 2030, CO2 emissions are projected to be 63% greater than they were in 2002.

FUTURE WARMING

The amount CO2 levels will rise in the next decades is unknown. What will this number depend on in the developed nations? What will it depend on in the developing nations? In the developed nations it will depend on technological advances or lifestyle changes that decrease emissions. In the developing nations, it will depend on how much their lifestyles improve and how these improvements are made. Computer models are used to predict the effects of greenhouse gas increases on climate for the planet as a whole and also for specific regions.

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

If nothing is done to control greenhouse gas emissions and they continue to increase at current rates, the surface temperature of the Earth can be expected to increase between 0.5 degrees C and 2.0 degrees C (0.9 degrees F and 3.6 degrees F) by 2050 and between

2 degrees and 4.5 degrees C (3.5 degrees and 8 degrees F) by 2100, with CO2 levels over 800 parts per million (ppm). On the other hand, if severe limits on CO2 emissions begin soon, temperatures could rise less than 1.1 degrees C (2 degrees F) by 2100.

Whatever the temperature increase, it will not be uniform around the globe. A rise of 2.8 degrees C (5 degrees F) would result in 0.6 degrees to 1.2 degrees C (1 degree to 2 degrees F) at the equator, but up to 6.7 degrees C (12 degrees F) at the poles. So far, global warming has affected the North Pole more than the South Pole, but temperatures are still increasing at Antarctica. The timing of events for species is changing. Mating and migrations take place earlier in the spring months. Species that can are moving their ranges uphill. Some regions that were already marginal for agriculture are no longer farmable because they have become too warm or dry.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180



Glaciers are melting and vegetation zones are moving uphill. If fossil fuel use exploded in the 1950s, why do these changes begin early in the animation? Does this mean that the climate change we are seeing is caused by natural processes and not by fossil fuel use?

As greenhouse gases increase, changes will be more extreme. Oceans will become slightly more acidic, making it more difficult for creatures with carbonate shells to grow, and that includes coral reefs. A study monitoring ocean acidity in the Pacific Northwest found ocean acidity increasing ten times faster than expected and 10% to 20% of shellfish (mussels) being replaced by acid tolerant algae.

Plant and animal species seeking cooler temperatures will need to move poleward 100 to 150 km (60 to 90 miles) or upward 150 m (500 feet) for each 1.0 degrees C (8 degrees F) rise in global temperature. There will be a tremendous loss of biodiversity because forest species can’t migrate that rapidly. Biologists have already documented the extinction of high-altitude species that have nowhere higher to go.

Decreased snow packs, shrinking glaciers, and the earlier arrival of spring will all lessen the amount of water available in some regions of the world, including the western United States and much of Asia. Ice will continue to melt and sea level is predicted to rise 18 to 97 cm (7 to 38 inches) by 2100. An increase this large will gradually flood coastal regions where about one-third of the world’s population lives, forcing billions of people to move inland.

Weather will become more extreme with heat waves and droughts. Some modelers predict that the Midwestern United States will become too dry to support agriculture and that Canada will become the new breadbasket. In all, about 10% to 50% of current cropland worldwide may become unusable if CO2 doubles. There are global monitoring systems to help monitor potential droughts that could turn into famines if they occur in politically and socially unstable regions of the world and if appropriate action isn't taken in time. One example isFEWS NET, which is a network of social and environmental scientists using geospatial technology to monitor these situations. But even with proper monitoring, if nations don't act, catastrophes can occur like in Somalia from 2010-2012.	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

Although scientists do not all agree, hurricanes are likely to become more severe and possibly more frequent. Tropical and subtropical insects will expand their ranges, resulting in the spread of tropical diseases such as malaria, encephalitis, yellow fever, and dengue fever.

You may notice that the numerical predictions above contain wide ranges. Sea level, for example, is expected to rise somewhere between 18 and 97 cm — quite a wide range. What is the reason for this uncertainty? It is partly because scientists cannot predict exactly how the Earth will respond to increased levels of greenhouses gases. How quickly greenhouse gases continue to build up in the atmosphere depends in part on the choices we make.

An important question people ask is this: Are the increases in global temperature natural? In other words, can natural variations in temperature account for the increase in temperature that we see? The scientific data shows no, natural variations can not explain the dramatic increase in global temperatures. Changes in the Sun's irradiance, El Niño and La Niña cycles, natural changes in greenhouse gas, plate tectonics, and the Milankovitch Cycles cannot account for the increase in temperature that has already happened in the past decades.

In December 2013 and April 2014, the Intergovernmental Panel on Climate Change (IPCC) released a series of damaging reports on not only the current scientific knowledge of climate change, but also on the vulnerability and impacts to humans and ecosystems. Below are two videos detailing the physical science of climate change (Working Group I), and the risks and impacts to the planet (Working Group II). But it is important to get a strong, data driven understanding of climate change. Along with the IPCC, other organizations like the United Nations Environmental Programme (UNEP), World Health Organizations, World Meteorological Organization (WMO), NASA, the National Oceanic and Atmospheric Administration (NOAA), and the U.S. Environmental Protection Agency (EPA).

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180	An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=180

CC licensed content, Shared previously

- **Provided by**: Open Geography. **Located at**: http://www.opengeography.org/ch-12-climate-systems-and-change.html. **License**: *CC BY: Attribution*

This page titled 18.1: Climate Systems and Change is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

18.2: Climate and Weather

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=182

CC licensed content, Original

- **Authored by**: William Hopper. **Provided by**: Florida Memorial University. **Located at**: https://www.youtube.com/watch?v=C2xWEuS0Gpk. **License**: *CC BY: Attribution*

This page titled 18.2: Climate and Weather is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

18.3: Global Climate Change

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=184

An interactive or media element has been excluded from this version of the text. You can view it online here: http://pb.libretexts.org/es/?p=184

CC licensed content, Original

- **Authored by**: William Hopper. **Provided by**: Florida Memorial University. **License**: *CC BY: Attribution*

This page titled 18.3: Global Climate Change is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

CHAPTER OVERVIEW

19: Deserts

19.1: Deserts

19.2: Desertification

19.3: Dunes

19.4: Video- Desertification

This page titled 19: Deserts is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

19.1: Deserts



Sand dunes in the Rub' al Khali ("Empty quarter") of Saudi Arabia

A **desert** is a barren area of land where little precipitation occurs and consequently living conditions are hostile for plant and animal life. The lack of vegetation exposes the unprotected surface of the ground to the processes of denudation. About one third of the land surface of the world is arid or semi-arid. This includes much of the polar regions where little precipitation occurs and which are sometimes called "cold deserts". Deserts can be classified by the amount of precipitation that falls, by the temperature that prevails, by the causes of desertification or by their geographical location.

Deserts are formed by weathering processes as large variations in temperature between day and night put strains on the rocks which consequently break in pieces. Although rain seldom occurs in deserts, there are occasional downpours that can result in flash floods. Rain falling on hot rocks can cause them to shatter and the resulting fragments and rubble strewn over the desert floor is further eroded by the wind. This picks up particles of sand and dust and wafts them aloft in sand or dust storms. Wind-blown sand grains striking any solid object in their path can abrade the surface. Rocks are smoothed down, and the wind sorts sand into uniform deposits. The grains end up as level sheets of sand or are piled high in billowing sand dunes. Other deserts are flat, stony plains where all the fine material has been blown away and the surface consists of a mosaic of smooth stones. These areas are known as desert pavements and little further erosion takes place. Other desert features include rock outcrops, exposed bedrock and clays once deposited by flowing water. Temporary lakes may form and salt pans may be left when waters evaporate. There may be underground sources of water in the form of springs and seepages from aquifers. Where these are found, oases can occur.

Plants and animals living in the desert need special adaptations to survive in the harsh environment. Plants tend to be tough and wiry with small or no leaves, water-resistant cuticles and often spines to deter herbivory. Some annual plants germinate, bloom and die in the course of a few weeks after rainfall while other long-lived plants survive for years and have deep root systems able to tap underground moisture. Animals need to keep cool and find enough food and water to survive. Many are nocturnal and stay in the shade or underground during the heat of the day. They tend to be efficient at conserving water, extracting most of their needs from their food and concentrating their urine. Some animals remain in a state of dormancy for long periods, ready to become active again when the rare rains fall. They then reproduce rapidly while conditions are favorable before returning to dormancy.

People have struggled to live in deserts and the surrounding semi-arid lands for millennia. Nomads have moved their flocks and herds to wherever grazing is available and oases have provided opportunities for a more settled way of life. The cultivation of semi-arid regions encourages erosion of soil and is one of the causes of increased desertification. Desert farming is possible with the aid of irrigation and the Imperial Valley in California provides an example of how previously barren land can be made productive by the import of water from an outside source. Many trade routes have been forged across deserts, especially across the Sahara, and traditionally were used by caravans of camels carrying salt, gold, ivory and other goods. Large numbers of slaves were also taken northwards across the Sahara. Some mineral extraction also takes place in deserts and the uninterrupted sunlight gives potential for the capture of large quantities of solar energy.

Etymology

English *desert* and its Romance cognates (including Italian and Portuguese *deserto*, French *désert* and Spanish *desierto*) all come from the ecclesiastical Latin *dēsertum* (originally "an abandoned place"), a participle of *dēserere*, "to abandon".[1] The correlation between aridity and sparse population is complex and dynamic, varying by culture, era, and technologies; thus the use of the word *desert* can cause confusion. In English before the 20th century, *desert* was often used in the sense of "unpopulated area", without specific reference to aridity;[1] but today the word is most often used in its climate-science sense (an area of low precipitation).[2]

Phrases such as “desert island”[3] and “Great American Desert” in previous centuries did not necessarily imply sand or aridity; their focus was the sparse population.[4]

The transliteration of the Ancient Egyptian term for the *Red land* (i.e. the deserts on either side of the fertile Black land irrigated by the Nile) is*dšrt* (conventionally pronounced *deshret*); it has been stated that “It is not impossible that the very word *deserta* entered the Latin language by way of Egyptian”.[5]

Physical geography

A desert is a region of land that is very dry because it receives low amounts of precipitation (usually in the form of rain but may be snow, mist or fog), often has little coverage by plants, and in which streams dry up unless they are supplied by water from outside the area.[6] Deserts can also be described as areas where more water is lost by evapotranspiration than falls as precipitation.[7] Deserts generally receive less than 250 mm (10 in) of precipitation each year.[6]Semideserts are regions which receive between 250 and 500 mm (10 and 20 in) and when clad in grass, these are known as steppes.[7][8]

Classification

Deserts have been defined and classified in a number of ways, generally combining total precipitation, number of days on which this falls, temperature, and humidity, and sometimes additional factors.[8] For example, Phoenix, Arizona receives less than 250 mm (9.8 in) of precipitation per year, and is immediately recognized as being located in a desert because of its arid-adapted plants. The North Slope of Alaska’s Brooks Range also receives less than 250 mm (9.8 in) of precipitation per year and is often classified as a cold desert.[9] Other regions of the world have cold deserts, including areas of the Himalayas[10] and other high altitude areas in other parts of the world.[11] Polar deserts cover much of the ice-free areas of the Arctic and Antarctic.[12][13] A non-technical definition is that deserts are those parts of the Earth’s surface that have insufficient vegetation cover to support a human population.[14]

Potential evapotranspiration supplements the measurement of precipitation in providing a scientific measurement-based definition of a desert. The water budget of an area can be calculated using the formula $P - PE \pm S$, wherein P is precipitation, PE is potential evapotranspiration rates and $S$is amount of surface storage of water. Evapotranspiration is the combination of water loss through atmospheric evaporation and through the life processes of plants. Potential evapotranspiration, then, is the amount of water that *could* evaporate in any given region. As an example, Tucson, Arizona receives about 300 mm (12 in) of rain per year, however about 2,500 mm (98 in) of water could evaporate over the course of a year.[15]In other words, about eight times more water could evaporate from the region than actually falls as rain. Rates of evapotranspiration in cold regions such as Alaska are much lower because of the lack of heat to aid in the evaporation process.[16]

Deserts are sometimes classified as “hot” or “cold”, “semiarid” or “coastal”.[14] The characteristics of hot deserts include high temperatures in summer; greater evaporation than precipitation usually exacerbated by high temperatures, strong winds and lack of cloud cover; considerable variation in the occurrence of precipitation, its intensity and distribution; and low humidity. Winter temperatures vary considerably between different deserts and are often related to the location of the desert on the continental landmass and the latitude. Daily variations in temperature can be as great as 22 °C (40 °F) or more, with heat loss by radiation at night being increased by the clear skies.[17]



Cold desert: snow surface at Dome C Station, Antarctica

Cold deserts, sometimes known as temperate deserts, occur at higher latitudes than hot deserts, and the aridity is caused by the dryness of the air. Some cold deserts are far from the ocean and others are separated by mountain ranges from the sea and in both cases there is insufficient moisture in the air to cause much precipitation. The largest of these deserts are found in Central Asia. Others occur on the eastern side of the Rocky Mountains, the eastern side of the southern Andes and in southern Australia.[7] Polar deserts are a particular class of cold desert. The air is very cold and carries little moisture so little precipitation occurs and what

does fall, usually as snow, is carried along in the often strong wind and may form blizzards, drifts and dunes similar to those caused by dust and sand in other desert regions. In Antarctica, for example, the annual precipitation is about 50 mm (2 in) on the central plateau and some ten times that amount on some major peninsulas.[17]

Based on precipitation alone, hyperarid deserts receive less than 25 mm (1 in) of rainfall a year; they have no annual seasonal cycle of precipitation and experience twelve-month periods with no rainfall at all.[17][18] Arid deserts receive between 25 and 200 mm (1 and 8 in) in a year and semiarid deserts between 200 and 500 mm (8 and 20 in). However, such factors as the temperature, humidity, rate of evaporation and evapotranspiration, and the moisture storage capacity of the ground have a marked effect on the degree of aridity and the plant and animal life that can be sustained. Rain falling in the cold season may be more effective at promoting plant growth, and defining the boundaries of deserts and the semiarid regions that surround them on the grounds of precipitation alone is problematic.[17]

Coastal deserts are mostly found on the western edges of continental land masses in regions where cold currents approach the land or cold water upwellings rise from the ocean depths. The cool winds crossing this water pick up little moisture and the coastal regions have low temperatures and very low rainfall, the main precipitation being in the form of fog and dew. The range of temperatures on a daily and annual scale is relatively low, being 11 °C (20 °F) and 5 °C (9 °F) respectively in the Atacama Desert. Deserts of this type are often long and narrow and bounded to the east by mountain ranges. They occur in south-west Africa, Chile, southern California and Baja California. Other coastal deserts influenced by cold currents are found in Western Australia, the Arabian Peninsula and Horn of Africa, and the western fringes of the Sahara.[17]

In 1961, Peveril Meigs divided desert regions on Earth into three categories according to the amount of precipitation they received. In this now widely accepted system, extremely arid lands have at least twelve consecutive months without precipitation, arid lands have less than 250 mm (10 in) of annual precipitation, and semiarid lands have a mean annual precipitation of between 250 and 500 mm (10–20 in). Both extremely arid and arid lands are considered to be deserts while semiarid lands are generally referred to as steppes when they are grasslands.[8]



The Agasthiyamalai hills cut off Tirunelveli in India from the monsoons, creating a rainshadow region.

Deserts are also classified, according to their geographical location and dominant weather pattern, as trade wind, mid-latitude, rain shadow, coastal, monsoon, or polar deserts.[19] Trade wind deserts occur either side of the horse latitudes at 30° to 35° North and South. These belts are associated with the subtropical anticyclone and the large-scale descent of dry air moving from high-altitudes toward the poles. The Sahara Desert is of this type.[20] Mid-latitude deserts occur between 30° and 50° North and South. They are mostly in areas remote from the sea where most of the moisture has already precipitated from the prevailing winds. They include the Tengger and Sonoran Deserts.[19] Monsoon deserts are similar. They occur in regions where large temperature differences occur between sea and land. Moist warm air rises over the land, deposits its water content and circulates back to sea. Further inland, areas receive very little precipitation. The Thar Desert near the India/Pakistan border is of this type.[19]


In some parts of the world, deserts are created by a rain shadow effect. Orographic lift occurs as air masses rise to pass over high ground. In the process they cool and lose much of their moisture by precipitation on the windward slope of the mountain range. When they descend on the leeward side, they warm and their capacity to hold moisture increases so an area with relatively little precipitation occurs.[21] The Taklamakan Desert is an example, lying in the rain shadow of the Himalayas and receiving less than 38 mm (1.5 in) precipitation annually.[22] Other areas are arid by virtue of being a very long way from the nearest available sources of moisture.[23]

Montane deserts are arid places with a very high altitude; the most prominent example is found north of the Himalayas, in the Kunlun Mountains and the Tibetan Plateau. Many locations within this category have elevations exceeding 3,000 m (9,800 ft) and the thermal regime can be hemiboreal. These places owe their profound aridity (the average annual precipitation is often less than 40 mm or 1.5 in) to being very far from the nearest available sources of moisture and are often in the lee of mountain ranges.

Montane deserts are normally cold, or may be scorchingly hot by day and very cold by night as is true of the northeastern slopes of Mount Kilimanjaro.[24]

Polar deserts such as McMurdo Dry Valleys remain ice-free because of the dry katabatic winds that flow downhill from the surrounding mountains.[25] Former desert areas presently in non-arid environments, such as the Sandhills in Nebraska, are known as paleodeserts.[19] In the Köppen climate classification system, deserts are classed as *BWh* (hot desert) or *BWk* (temperate desert). In the Thornthwaite climate classification system, deserts would be classified as arid megathermal climates.[26][27]

Weathering processes



Exfoliation of weathering rocks in Texas

Deserts usually have a large diurnal and seasonal temperature range, with high daytime temperatures falling sharply at night. The diurnal range may be as much as 20 to 30 °C (36 to 54 °F) and the rock surface experiences even greater temperature differentials.[28] During the day the sky is usually clear and most of the sun's radiation reaches the ground, but as soon as the sun sets, the desert cools quickly by radiating heat into space. In hot deserts, the temperature during daytime can exceed 45 °C (113 °F) in summer and plunge below freezing point at night during winter.[29]



One square centimeter
(0.16 sq in) of windblown sand from the Gobi Desert

Such large temperature variations have a destructive effect on the exposed rocky surfaces. The repeated fluctuations put a strain on exposed rock and the flanks of mountains crack and shatter. Fragmented strata slide down into the valleys where they continue to break into pieces due to the remorseless sun by day and chill by night. Successive strata are exposed to further weathering. The relief of the internal pressure that has built up in rocks that have been underground for aeons can cause them to shatter.[30]Exfoliation also occurs when the outer surfaces of rocks split off in flat flakes. This is believed to be caused by the stresses put on the rock by repeated expansions and contractions which induces fracturing parallel to the original surface.[28] Chemical weathering processes probably play a more important role in deserts than was previously thought. The necessary moisture may be present in the form of dew or mist. Ground water may be drawn to the surface by evaporation and the formation of salt crystals may dislodge rock particles as sand or disintegrate rocks by exfoliation. Shallow caves are sometimes formed at the base of cliffs by this means.[28]

As the desert mountains decay, large areas of shattered rock and rubble occur. The process continues and the end products are either dust or sand. Dust is formed from solidified clay or volcanic deposits whereas sand results from the fragmentation of harder granites, limestone and sandstone.[31] There is a certain critical size (about 500μ) below which further temperature-induced weathering of rocks does not occur and this provides a minimum size for sand grains.[32]


As the mountains are eroded, more and more sand is created. At high wind speeds, sand grains are picked up off the surface and blown along, a process known as saltation. The whirling airborne grains act as a sand blasting mechanism which grinds away solid objects in its path as the kinetic energy of the wind is transferred to the ground.[33] The sand eventually ends up deposited in level areas known as sand-fields or sand-seas, or piled up in dunes.[34]

Dust storms and sandstorms

Main article: Dust storm 

Dust storm about to engulf a military camp in Iraq, 2005

Sand and dust storms are natural events that occur in arid regions where the land is not protected by a covering of vegetation. Dust storms usually start in desert margins rather than the deserts themselves where the finer materials have already been blown away. As a steady wind begins to blow, fine particles lying on the exposed ground begin to vibrate. At greater wind speeds, some particles are lifted into the air stream. When they land, they strike other particles which may be jerked into the air in their turn, starting a chain reaction. Once ejected, these particles move in one of three possible ways, depending on their size, shape and density; suspension, saltation or creep. Suspension is only possible for particles less than 0.1 mm (0.004 in) in diameter. In a dust storm, these fine particles are lifted up and wafted aloft to heights of up to 6 km (3.7 mi). They reduce visibility and can remain in the atmosphere for days on end, conveyed by the trade winds for distances of up to 6,000 km (3,700 mi).[35]Denser clouds of dust can be formed in stronger winds, moving across the land with a billowing leading edge. The sunlight can be obliterated and it may become as dark as night at ground level.[36] In a study of a dust storm in China in 2001, it was estimated that 6.5 million tons of dust were involved, covering an area of 134,000,000 km^2 (52,000,000 sq mi). The mean particle size was 1.44 μm.[37] A much smaller scale, short-lived phenomenon can occur in calm conditions when hot air near the ground rises quickly through a small pocket of cooler, low-pressure air above forming a whirling column of particles, a dust devil.[38]



Wind-blown particles: 1=Creep 2=Saltation 3=Suspension 4=Wind current

Sandstorms occur with much less frequency than dust storms. They are often preceded by severe dust storms and occur when the wind velocity increases to a point where it can lift heavier particles. These grains of sand, up to about 0.5 mm (0.020 in) in diameter are jerked into the air but soon fall back to earth, ejecting other particles in the process. Their weight prevents them from being airborne for long and most only travel a distance of a few meters (yards). The sand streams along above the surface of the ground like a fluid, often rising to heights of about 30 cm (12 in).[35] In a really severe steady blow, 2 m (6 ft 7 in) is about as high as the sand stream can rise as the largest sand grains do not become airborne at all. They are transported by creep, being rolled along the desert floor or performing short jumps.[36]

During a sandstorm, the wind-blown sand particles become electrically charged. Such electric fields, which range in size up to 80 kV/m, can produce sparks and cause interference with telecommunications equipment. They are also unpleasant for humans and can cause headaches and nausea.[36] The electric fields are caused by collision between airborne particles and by the impacts of saltating sand grains landing on the ground. The mechanism is little understood but the particles usually have a negative charge when their diameter is under 250 μm and a positive one when they are over 500 μm.[39][40]

Major deserts

See also: List of deserts by area 

The world's largest non-polar deserts


Deserts take up about one third of the Earth's land surface.[8] Bottomlands may be salt-covered flats. Eolian processes are major factors in shaping desert landscapes. Polar deserts (also seen as "cold deserts") have similar features, except the main form of precipitation is snow rather than rain. Antarctica is the world's largest cold desert (composed of about 98% thick continental ice sheet and 2% barren rock). Some of the barren rock is to be found in the so-called Dry Valleys of Antarctica that almost never get snow, which can have ice-encrusted saline lakes that suggest evaporation far greater than the rare snowfall due to the strong katabatic winds that even evaporate ice.

The ten largest deserts[41]

Rank	Desert	Area (km²)	Area (mi²)
1	**Antarctic Desert** (Antarctica)	14,200,000	5,500,000
2	**Arctic Desert** (Arctic)	13,900,000	5,400,000
3	**Sahara Desert** (Africa)	9,100,000	3,500,000
4	**Arabian Desert** (Middle East)	2,600,000	1,000,000
5	**Gobi Desert** (Asia)	1,300,000	500,000
6	**Patagonian Desert** (South America)	670,000	260,000
7	**Great Victoria Desert** (Australia)	647,000	250,000
8	**Kalahari Desert** (Africa)	570,000	220,000
9	**Great Basin Desert** (North America)	490,000	190,000
10	**Syrian Desert** (Middle East)	490,000	190,000

Deserts, both hot and cold, play a part in moderating the Earth's temperature. This is because they reflect more of the incoming light and their albedo is higher than that of forests or the sea.[42]

Features



Aerial view of Makhtesh Ramon, an erosion cirque of a type unique to the Negev

Many people think of deserts as consisting of extensive areas of billowing sand dunes because that is the way they are often depicted on TV and in films,[43] but deserts do not always look like this.[44] Across the world, around 20% of desert is sand, varying from only 2% in North America to 30% in Australia and over 45% in Central Asia.[45] Where sand does occur, it is usually in large quantities in the form of sand sheets or extensive areas of dunes.[45]

A sand sheet is a near-level, firm expanse of partially consolidated particles in a layer that varies from a few centimeters to a few meters thick. The structure of the sheet consists of thin horizontal layers of coarse silt and very fine to medium grain sand, separated by layers of coarse sand and pea-gravel which are a single grain thick. These larger particles anchor the other particles in place and may also be packed together on the surface so as to form a miniature desert pavement.[46] Small ripples form on the sand sheet when the wind exceeds 24 kph (15 mph). They form perpendicular to the wind direction and gradually move across the surface as the wind continues to blow. The distance between their crests corresponds to the average length of jumps made by particles during saltation. The ripples are ephemeral and a change in wind direction causes them to reorganise.[47]




Diagram showing barchan dune formation, with the wind blowing from the left

Sand dunes are accumulations of windblown sand piled up in mounds or ridges. They form downwind of copious sources of dry, loose sand and occur when topographic and climatic conditions cause airborne particles to settle. As the wind blows, saltation and creep take place on the windward side of the dune and individual grains of sand move uphill. When they reach the crest, they cascade down the far side. The upwind slope typically has a gradient of 10° to 20° while the lee slope is around 32°, the angle at which loose dry sand will slip. As this wind-induced movement of sand grains takes place, the dune moves slowly across the surface of the ground.[48] Dunes are sometimes solitary, but they are more often grouped together in dune fields. When these are extensive, they are known as sand seas or ergs.[49]

The shape of the dune depends on the characteristics of the prevailing wind. Barchan dunes are produced by strong winds blowing across a level surface, and are crescent-shaped with the concave side away from the wind. When there are two directions from which winds regularly blow, a series of long, linear dunes known as seif dunes may form. These also occur parallel to a strong wind that blows in one general direction. Transverse dunes run at a right angle to the prevailing wind direction. Star dunes are formed by variable winds, and have several ridges and slip faces radiating from a central point. They tend to grow vertically; they can reach a height of 500 m (1,600 ft), making them the tallest type of dune. Rounded mounds of sand without a slip face are the rare dome dunes, found on the upwind edges of sand seas.[49]



Windswept desert pavement of small, smooth, closely packed stones in the Mojave desert

A large part of the surface area of the world's deserts consists of flat, stone-covered plains dominated by wind erosion. In "eolian deflation", the wind continually removes fine-grained material, which becomes wind-blown sand. This exposes coarser-grained material, mainly pebbles with some larger stones or cobbles,[34][45] leaving a desert pavement, an area of land overlaid by closely packed smooth stones forming a tessellated mosaic. Different theories exist as to how exactly the pavement is formed. It may be that after the sand and dust is blown away by the wind the stones jiggle themselves into place; alternatively, stones previously below ground may in some way work themselves to the surface. Very little further erosion takes place after the formation of a pavement, and the ground becomes stable. Evaporation brings moisture to the surface by capillary action and calcium salts may be precipitated, binding particles together to form a desert conglomerate.[50] In time, bacteria that live on the surface of the stones accumulate a film of minerals and clay particles, forming a shiny brown coating known as desert varnish.[51]

Other non-sandy deserts consist of exposed outcrops of bedrock, dry soils or aridisols, and a variety of landforms affected by flowing water, such as alluvial fans, sinks or playas, temporary or permanent lakes, and oases.[45] A hamada is a type of desert landscape consisting of a high rocky plateau where the sand has been removed by aeolian processes. Other landforms include plains largely covered by gravels and angular boulders, from which the finer particles have been stripped by the wind. These are called "reg" in the western Sahara, "serir" in the eastern Sahara, "gibber plains" in Australia and "saï" in central Asia.[52] The Tassili Plateau in Algeria is an impressive jumble of eroded sandstone outcrops, canyons, blocks, pinnacles, fissures, slabs and ravines. In some places the wind has carved holes or arches and in others it has created mushroom-like pillars narrower at the base than the top.[53] In the Colorado Plateau it is water that has been the eroding force. Here the Colorado River has cut its way over the millennia through the high desert floor creating a canyon that is over a mile (6,000 feet or 1,800 meters) deep in places, exposing strata that are over two billion year old.[54]

Water



Atacama, the world's driest desert

One of the driest places on Earth is the Atacama Desert.[55] It is virtually devoid of life because it is blocked from receiving precipitation by the Andes mountains to the east and the Chilean Coast Range to the west. The cold Humboldt Current and the anticyclone of the Pacific are essential to keep the dry climate of the Atacama. The average precipitation in the Chilean region of Antofagasta is just 1 mm (0.039 in) per year. Some weather stations in the Atacama have never received rain. Evidence suggests that the Atacama may not have had any significant rainfall from 1570 to 1971. It is so arid that mountains that reach as high as 6,885 m (22,589 ft) are completely free of glaciers and, in the southern part from 25°S to 27°S, may have been glacier-free throughout the Quaternary, though permafrost extends down to an altitude of 4,400 m (14,400 ft) and is continuous above 5,600 m (18,400 ft).[56][57] Nevertheless, there is some plant life in the Atacama, in the form of specialist plants that obtain moisture from dew and the fogs that blow in from the Pacific.[55]



Flash flood in the Gobi

When rain falls in deserts, as it occasionally does, it is often with great violence. The desert surface is evidence of this with dry stream channels known as arroyos or wadis meandering across its surface. These can experience flash floods, becoming raging torrents with surprising rapidity after a storm that may be many kilometers away. Most deserts are in basins with no drainage to the sea but some are crossed by exotic rivers sourced in mountain ranges or other high rainfall areas beyond their borders. The River Nile, the Colorado River and the Yellow River do this, losing much of their water through evaporation as they pass through the desert and raising groundwater levels nearby. There may also be underground sources of water in deserts in the form of springs, aquifers, underground rivers or lakes. Where these lie close to the surface, wells can be dug and oases may form where plant and animal life can flourish.[45] The Nubian Sandstone Aquifer System under the Sahara Desert is the largest known accumulation of fossil water. The Great Man-Made River is a scheme launched by Libya's Colonel Gadaffi to tap this aquifer and supply water to coastal cities.[58]Kharga Oasis in Egypt is 150 km (93 mi) long and is the largest oasis in the Libyan Desert. A lake occupied this depression in ancient times and thick deposits of sandy-clay resulted. Wells are dug to extract water from the porous sandstone that lies underneath.[59] Seepages may occur in the walls of canyons and pools may survive in deep shade near the dried up watercourse below.[60]

Lakes may form in basins where there is sufficient precipitation or meltwater from glaciers above. They are usually shallow and saline, and wind blowing over their surface can cause stress, moving the water over nearby low-lying areas. When the lakes dry up, they leave a crust or hardpan behind. This area of deposited clay, silt or sand is known as a playa. The deserts of North America have more than one hundred playas, many of them relics of Lake Bonneville which covered parts of Utah, Nevada and Idaho during the last ice age when the climate was colder and wetter.[61] These include the Great Salt Lake, Utah Lake, Sevier Lake and many dry lake beds. The smooth flat surfaces of playas have been used for attempted vehicle speed records at Black Rock Desert and Bonneville Speedway and the United States Air Force uses Rogers Dry Lake in the Mojave Desert as runways for aircraft and the space shuttle.[45]

Flora



Xerophytes: Cardón cacti in the Baja California Desert, Cataviña region, Mexico

Plants face severe challenges in arid environments. Problems they need to solve include how to obtain enough water, how to avoid being eaten and how to reproduce. Photosynthesis is the key to plant growth. It can only takes place during the day as energy from the sun is required, but during the day, many deserts become very hot. Opening stomata to allow in the carbon dioxide necessary for the process causes evapotranspiration, and conservation of water is a top priority for desert vegetation. Some plants have resolved this problem by adopting crassulacean acid metabolism, allowing them to open their stomata during the night to allow CO_2 to enter, and close them during the day,[62] or by using C4 carbon fixation.[63]

Many desert plants have reduced the size of their leaves or abandoned them altogether. Cacti are desert specialists and in most species the leaves have been dispensed with and the chlorophyll displaced into the trunks, the cellular structure of which has been modified to allow them to store water. When rain falls, the water is rapidly absorbed by the shallow roots and retained to allow them to survive until the next downpour, which may be months or years away.[64] The giant saguaro cacti of the Sonoran Desert form “forests”, providing shade for other plants and nesting places for desert birds. Saguaro grow slowly but may live for up to two hundred years. The surface of the trunk is folded like a concertina, allowing it to expand, and a large specimen can hold eight tons of water after a good downpour.[64]

Cacti are restricted to the New World but other xerophytic plants have developed similar strategies by a process known as convergent evolution.[65] They limit water loss by reducing the size and number of stomata, by having waxy coatings and hairy or tiny leaves. Some are deciduous, shedding their leaves in the driest season, and others curl their leaves up to reduce transpiration. Others store water in succulent leaves or stems or in fleshy tubers. Desert plants maximize water uptake by having shallow roots that spread widely, or by developing long taproots that reach down to deep rock strata for ground water.[66] The saltbush in Australia has succulent leaves and secretes salt crystals, enabling it to live in saline areas.[66][67] In common with cacti, many have developed spines to ward off browsing animals.[64]



The Camel Thorn Tree (Acacia erioloba) in the Namib Desert is nearly leafless in dry periods.

Some desert plants produce seed which lies dormant in the soil until sparked into growth by rainfall. When annuals, such plants grow with great rapidity and may flower and set seed within weeks, aiming to complete their development before the last vestige of water dries up. For perennial plants, reproduction is more likely to be successful if the seed germinates in a shaded position, but not so close to the parent plant as to be in competition with it. Some seed will not germinate until it has been blown about on the desert floor to scarify the seed coat. The seed of the mesquite tree which grows in deserts in American is hard and fails to sprout even when planted carefully. When it has passed through the gut of a pronghorn it germinates readily, and the little pile of moist dung provides an excellent start to life well away from the parent tree.[64] The stems and leaves of some plants lower the surface velocity of sand-carrying winds and protect the ground from erosion. Even small fungi and microscopic plant organisms found on the soil surface (so-called *cryptobiotic soil*) can be a vital link in preventing erosion and providing support for other living organisms. Cold deserts often have high concentrations of salt in the soil. Grasses and low shrubs are the dominant vegetation here and the ground may be covered with lichens. Most shrubs have spiny leaves and shed them in the coldest part of the year.[68]

Fauna

Main article: Xerocole

Animals adapted to live in deserts are called xerocoles. There is no evidence that body temperature of mammals and birds is adaptive to the different climates, either of great heat or cold. In fact, with a very few exceptions, their basal metabolic rate is determined by body size, irrespective of the climate in which they live.[69] Many desert animals (and plants) show especially clear evolutionary adaptations for water conservation or heat tolerance and so are often studied in comparative physiology, ecophysiology, and evolutionary physiology. One well-studied example is the specializations of mammalian kidneys shown by desert-inhabiting species.[70] Many examples of convergent evolution have been identified in desert organisms, including between cacti and Euphorbia, kangaroo rats and jerboas, *Phrynosoma* and *Moloch* lizards.[71]



The cream-colored courser,*Cursorius cursor*, is a well-camouflaged desert resident with its dusty coloration, countershading, and disruptive head markings.

Deserts present a very challenging environment for animals. Not only do they require food and water but they also need to keep their body temperature at a tolerable level. In many ways birds are the most able to do this of the higher animals. They can move to areas of greater food availability as the desert blooms after local rainfall and can fly to faraway waterholes. In hot deserts, gliding birds can remove themselves from the over-heated desert floor by using thermals to soar in the cooler air at great heights. In order to conserve energy, other desert birds run rather than fly. The cream-colored courser flits gracefully across the ground on its long legs, stopping periodically to snatch up insects. Like other desert birds it is well-camouflaged by its coloring and can merge into the landscape when stationary. The sandgrouse is an expert at this and nests on the open desert floor dozens of kilometers (miles) away from the waterhole it needs to visit daily. Some small diurnal birds are found in very restricted localities where their plumage matches the color of the underlying surface. The desert lark takes frequent dust baths which ensures that it matches its environment.[72]

Water and carbon dioxide are metabolic end products of oxidation of fats, proteins, and carbohydrates.[73] Oxidising a gram of carbohydrate produces 0.60 grams of water; a gram of protein produces 0.41 grams of water; and a gram of fat produces 1.07 grams of water,[74] making it possible for xerocoles to live with little or no access to drinking water.[75] The kangaroo rat for example makes use of this water of metabolism and conserves water both by having a low basal metabolic rate and by remaining underground during the heat of the day,[76] reducing loss of water through its skin and respiratory system when at rest.[75][77]Herbivorous mammals obtain moisture from the plants they eat. Species such as the addax antelope,[78]dik-dik, Grant's gazelle and oryx are so efficient at doing this that they apparently never need to drink.[79] The camel is a superb example of a mammal adapted to desert life. It minimizes its water loss by producing concentrated urine and dry dung, and is able to lose 40% of its body weight through water loss without dying of dehydration.[80]Carnivores can obtain much of their water needs from the body fluids of their prey.[81] Many other hot desert animals are nocturnal, seeking out shade during the day or dwelling underground in burrows. At depths of more than 50 cm (20 in), these remain at between 30 and 32 °C (86 and 90 °F) regardless of the external temperature.[81]Jerboas, desert rats, kangaroo rats and other small rodents emerge from their burrows at night and so do the foxes, coyotes, jackals and snakes that prey on them. Kangaroos keep cool by increasing their respiration rate, panting, sweating and moistening the skin of their forelegs with saliva.[82] Mammals living in cold deserts have developed greater insulation through warmer body fur and insulating layers of fat beneath the skin. The arctic weasel has a metabolic rate that is two or three times as high as would be expected for an animal of its size. Birds have avoided the problem of losing heat through their feet by not attempting to maintain them at the same temperature as the rest of their bodies, a form of adaptive insulation.[69] The emperor penguin has dense plumage, a downy under layer, an air insulation layer next the skin and various thermoregulatory strategies to maintain its body temperature in one of the harshest environments on Earth.[83]

Being ectotherms, reptiles are unable to live in cold deserts but are well-suited to hot ones. In the heat of the day in the Sahara, the temperature can rise to 50 °C (122 °F). Reptiles cannot survive at this temperature and lizards will be prostrated by heat at 45 °C (113 °F). They have few adaptations to desert life and are unable to cool themselves by sweating so they shelter during the heat of the day. In the first part of the night, as the ground radiates the heat absorbed during the day, they emerge and search for prey. Lizards and snakes are the most numerous in arid regions and certain snakes have developed a novel method of locomotion that enables them to move sidewards and navigate high sand-dunes. These include the horned viper of Africa and the sidewinder of North America, evolutionarily distinct but with similar behavioural patterns because of convergent evolution. Many desert reptiles are ambush predators and often bury themselves in the sand, waiting for prey to come within range.[84]

Amphibians might seem unlikely desert-dwellers, because of their need to keep their skins moist and their dependence on water for reproductive purposes. In fact, the few species that are found in this habitat have made some remarkable adaptations. Most of them are fossorial, spending the hot dry months aestivating in deep burrows. While there they shed their skins a number of times and retain the remnants around them as a waterproof cocoon to retain moisture. In the Sonoran Desert, Couch's spadefoot toad spends most of the year dormant in its burrow. Heavy rain is the trigger for emergence and the first male to find a suitable pool calls to attract others. Eggs are laid and the tadpoles grow rapidly as they must reach metamorphosis before the water evaporates. As the desert dries out, the adult toads rebury themselves. The juveniles stay on the surface for a while, feeding and growing, but soon dig themselves burrows. Few make it to adulthood.[85] The water holding frog in Australia has a similar life cycle and may aestivate for as long as five years if no rain falls.[86] The Desert rain frog of Namibia is nocturnal and survives because of the damp sea fogs that roll in from the Atlantic.[87]



Tadpole shrimp survive dry periods as eggs, which rapidly hatch and develop after rain.

Invertebrates, particularly arthropods, have successfully made their homes in the desert. Flies, beetles, ants, termites, locusts, millipedes, scorpions and spiders[88] have hard cuticles which are impervious to water and many of them lay their eggs underground and their young develop away from the temperature extremes at the surface.[89] The Saharan silver ant (*Cataglyphis bombycina*) uses a heat shock protein in a novel way and forages in the open during brief forays in the heat of the day.[90] The long-legged darkling beetle in Namibia stands on its front legs and raises its carapace to catch the morning mist as condensate, funnelling the water into its mouth.[91] Some arthropods make use of the ephemeral pools that form after rain and complete their life cycle in a matter of days. The desert shrimp does this, appearing "miraculously" in new-formed puddles as the dormant eggs hatch. Others, such as brine shrimps, fairy shrimps and tadpole shrimps, are cryptobiotic and can lose up to 92% of their bodyweight, rehydrating as soon as it rains and their temporary pools reappear.[92]

Human relations

Humans have long made use of deserts as places to live,[93] and more recently have started to exploit them for minerals[94] and energy capture.[95] Deserts play a significant role in human culture with an extensive literature.[96]

History



Shepherd near Marrakech leading his flock to new pasture

People have been living in deserts for millennia. Many, such as the Bushmen in the Kalahari, the Aborigines in Australia and various tribes of North American Indians, were originally hunter-gatherers. They developed skills in the manufacture and use of weapons, animal tracking, finding water, foraging for edible plants and using the things they found in their natural environment to supply their everyday needs. Their self-sufficient skills and knowledge were passed down through the generations by word of

mouth.[93] Other cultures developed a nomadic way of life as herders of sheep, goats, cattle, camels, yaks, lamas or reindeer. They travelled over large areas with their herds, moving to new pastures as seasonal and erratic rainfall encouraged new plant growth. They took with them their tents made of cloth or skins draped over poles and their diet included milk, blood and sometimes meat.[97]



Salt caravan travelling between Agadez and the Bilma salt mines

The desert nomads were also traders. The Sahara is a very large expanse of land stretching from the Atlantic rim to Egypt. Trade routes were developed linking the Sahel in the south with the fertile Mediterranean region to the north and large numbers of camels were used to carry valuable goods across the desert interior. The Tuareg were traders and the goods transported traditionally included slaves, ivory and gold going northwards and salt going southwards. Berbers with knowledge of the region were employed to guide the caravans between the various oases and wells.[98] Several million slaves may have been taken northwards across the Sahara between the 8th and 18th centuries.[99] Traditional means of overland transport declined with the advent of motor vehicles, shipping and air freight, but caravans still travel along routes between Agadez and Bilma and between Timbuktu and Taoudenni carrying salt from the interior to desert-edge communities.[100]

Round the rims of deserts, where more precipitation occurred and conditions were more suitable, some groups took to cultivating crops. This may have happened when drought caused the death of herd animals, forcing herdsmen to turn to cultivation. With few inputs, they were at the mercy of the weather and may have lived at bare subsistence level. The land they cultivated reduced the area available to nomadic herders, causing disputes over land. The semi-arid fringes of the desert have fragile soils which are at risk of erosion when exposed, as happened in the American Dust Bowl in the 1930s. The grasses that held the soil in place were ploughed under, and a series of dry years caused crop failures, while enormous dust storms blew the topsoil away. Half a million Americans were forced to leave their land in this catastrophe.[101]

Similar damage is being done today to the semi-arid areas that rim deserts and about twelve million hectares of land are being turned to desert each year.[102]Desertification is caused by such factors as drought, climatic shifts, tillage for agriculture, overgrazing and deforestation. Vegetation plays a major role in determining the composition of the soil. In many environments, the rate of erosion and run off increases dramatically with reduced vegetation cover.[103] Unprotected dry surfaces tend to be blown away by the wind or be washed away by flash floods, leaving infertile soil layers that bake in the sun and become unproductive hardpan. Although overgrazing has historically been considered to be a cause of desertification, there is some evidence that wild and domesticated animals actually improve fertility and vegetation cover, and that their removal encourages erosive processes.[104]

Natural resource extraction



A mining plant near Jodhpur, India

Deserts contain substantial mineral resources, sometimes over their entire surface, giving them their characteristic colors. For example, the red of many sand deserts comes from laterite minerals.[105] Geological processes in a desert climate can concentrate minerals into valuable deposits. Leaching by ground water can extract ore minerals and redeposit them, according to the water table, in concentrated form.[94] Similarly, evaporation tends to concentrate minerals in desert lakes, creating dry lake beds or playas rich in minerals. Evaporation can concentrate minerals as a variety of evaporite deposits, including gypsum, sodium nitrate, sodium chloride and borates.[94] Evaporites are found in the USA's Great Basin Desert, historically exploited by the "20-mule teams" pulling carts of borax from Death Valley to the nearest railway.[94] A desert especially rich in mineral salts is the Atacama Desert, Chile, where sodium nitrate has been mined for explosives and fertilizer since around 1850.[94] Other desert minerals are copper

from Chile, Peru, and Iran, and iron and uranium in Australia. Many other metals, salts and commercially valuable types of rock such as pumice are extracted from deserts around the world.[94]

Oil and gas form on the bottom of shallow seas when micro-organisms decompose under anoxic conditions and later become covered with sediment. Many deserts were at one time the sites of shallow seas and others have had underlying hydrocarbon deposits transported to them by the movement of tectonic plates.[106] Some major oilfields such as Ghawar are found under the sands of Saudi Arabia.[94] Geologists believe that other oil deposits were formed by aeolian processes in ancient deserts as may be the case with some of the major American oil fields.[94]

Farming

Main article: Desert farming

Traditional desert farming systems have long been established in North Africa, irrigation being the key to success in an area where water stress is a limiting factor to growth. Techniques that can be used include drip irrigation, the use of organic residues or animal manures as fertilisers and other traditional agricultural management practises. Once fertility has been built up, further crop production preserves the soil from destruction by wind and other forms of erosion.[107] It has been found that plant growth-promoting bacteria play a role in increasing the resistance of plants to stress conditions and these rhizobacterial suspensions could be inoculated into the soil in the vicinity of the plants. A study of these microbes found that desert farming hampers desertification by establishing islands of fertility allowing farmers to achieve increased yields despite the adverse environmental conditions.[107] A field trial in the Sonoran Desert which exposed the roots of different species of tree to rhizobacteria and the nitrogen fixing bacterium *Azospirillum brasilense* with the aim of restoring degraded lands was only partially successful.[107]

The Judean Desert was farmed in the 7th century BC during the Iron Age to supply food for desert forts.[108] Native Americans in the south western United States became agriculturalists around 600 AD when seeds and technologies became available from Mexico. They used terracing techniques and grew gardens beside seeps, in moist areas at the foot of dunes, near streams providing flood irrigation and in areas irrigated by extensive specially built canals. The Hohokam tribe constructed over 500 miles (800 km) of large canals and maintained them for centuries, an impressive feat of engineering. They grew maize, beans, squash and peppers.[109]



Mosaic of fields in Imperial Valley

A modern example of desert farming is the Imperial Valley in California, which has high temperatures and average rainfall of just 3 in (76 mm) per year.[110] The economy is heavily based on agriculture and the land is irrigated through a network of canals and pipelines sourced entirely from the Colorado River via the All-American Canal. The soil is deep and fertile, being part of the river's flood plains, and what would otherwise have been desert has been transformed into one of the most productive farming regions in California. Other water from the river is piped to urban communities but all this has been at the expense of the river, which below the extraction sites no longer has any above-ground flow during most of the year. Another problem of growing crops in this way is the build-up of salinity in the soil caused by evaporation of river water.[111] The greening of the desert remains an aspiration and was at one time viewed as a future means for increasing food production for the world's growing population. This prospect has proved false as it disregarded the environmental damage caused elsewhere by the diversion of water for desert project irrigation.[112]

Solar energy capture




Desertec proposes using the Saharan and Arabian deserts to produce solar energy to power Europe and the Middle East.

Deserts are increasingly seen as sources for solar energy, partly due to low amounts of cloud cover. Many successful solar power plants have been built in the Mojave Desert. These plants have a combined capacity of 354 megawatts (MW) making them the largest solar power installation in the world.[113] Large swaths of this desert are covered in mirrors,[114] including nine fields of solar collectors. The Mojave Solar Park is currently under construction and will produce 280MW when completed.[115]

The potential for generating solar energy from the Sahara desert is immense. Professor David Faiman of Ben-Gurion University has stated that the technology now exists to supply all of the world's electricity needs from 10% of the Sahara Desert.[116]Desertec Industrial Initiative is a consortium seeking $560 billion to invest in North African solar and wind installations over the next forty years to supply electricity to Europe via cable lines running under the Mediterranean Sea. European interest in the Sahara Desert stems from its two aspects: the almost continual daytime sunshine and plenty of unused land. The Sahara receives more sunshine per acre than any part of Europe. The Sahara Desert also has the empty space totalling hundreds of square miles required to house fields of mirrors for solar plants.[117]

The Negev Desert, Israel, and the surrounding area, including the Arava Valley, receive plenty of sunshine and are generally not arable. This has resulted in the construction of many solar plants.[95] David Faiman has proposed that "giant" solar plants in the Negev could supply all of Israel's needs for electricity.[116]



Deployment of forces on the eve of the Second Battle of El Alamein in 1942

Warfare

Main article: Desert warfare

Both world wars saw fighting in the desert. In the First World War, the Ottoman Turks were engaged with the British regular army in a campaign that spanned the Arabian peninsula. The Turks were defeated by the British, who had the backing of irregular Arab forces that were seeking to revolt against the Turks in the Hejaz, made famous in T. E. Lawrence's book *Seven Pillars of Wisdom*.[118][119]

In the Second World War, the Western Desert Campaign began in Italian Libya. Warfare in the desert offered great scope for tacticians to use the large open spaces without the distractions of casualties among civilian populations. Tanks and armoured vehicles were able to travel large distances unimpeded and land mines were laid in large numbers. However the size and harshness of the terrain meant that all supplies needed to be brought in from great distances. The victors in a battle would advance and their supply chain would necessarily become longer, while the defeated army could retreat, regroup and resupply. For these reasons, the front line moved back and forth through hundreds of kilometers as each side lost and regained momentum.[120] Its most easterly point was at El Alamein in Egypt, where the Allies decisively defeated the Axis forces in 1942.[121]

In culture



Marco Polo arriving in a desert land with camels. 14th century miniature from *Il milione*.

The desert is generally thought of as a barren and empty landscape. It has been portrayed by writers, film-makers, philosophers, artists and critics as a place of extremes, a metaphor for anything from death, war or religion to the primitive past or the desolate future.[122]

There is an extensive literature on the subject of deserts.[96] An early historical account is that of Marco Polo (c. 1254–1324), who travelled through Central Asia to China, crossing a number of deserts in his twenty four year trek.[123] Some accounts give vivid descriptions of desert conditions, though often accounts of journeys across deserts are interwoven with reflection, as is the case in Charles Montagu Doughty's major work, *Travels in Arabia Deserta* (1888).[124]Antoine de Saint-Exupéry described both his flying and the desert in *Wind, Sand and Stars*[125]and Gertrude Bell travelled extensively in the Arabian desert in the early part of the 20th century, becoming an expert on the subject, writing books and advising the British government on dealing with the Arabs.[126] Another woman explorer was Freya Stark who travelled alone in the Middle East, visiting Turkey, Arabia, Yemen, Syria, Persia and Afghanistan, writing over twenty books on her experiences.[127] The German naturalist Uwe George spent several years living in deserts, recording his experiences and research in his book, *In the Deserts of this Earth*.[128]

The American poet Robert Frost expressed his bleak thoughts in his poem, *Desert Places*, which ends with the stanza "They cannot scare me with their empty spaces / Between stars – on stars where no human race is. / I have it in me so much nearer home / To scare myself with my own desert places."[129]

Deserts on other planets



View of the Martian desert seen by the probe *Spirit* in 2004.

Mars is the only planet in the Solar System on which deserts have been identified. Despite its low surface atmospheric pressure (only 1/100 of that of the Earth), the patterns of atmospheric circulation on Mars have formed a sea of circumpolar sand more than 5 million km² (1.9 million sq mi) in area, much larger than deserts on Earth. The Martian deserts principally consist of dunes in the form of half-moons in flat areas near the permanent polar ice caps in the north of the planet. The smaller dune fields occupy the bottom of many of the craters situated in the Martian polar regions.[130] Examination of the surface of rocks by laser beamed from the Mars Exploration Rover have shown a surface film that resembles the desert varnish found on Earth although it might just be surface dust.[131] The surface of Titan, a moon of Saturn, also has a desert-like surface with dune seas.[132]

- Aridification
- Arid Lands Information Network
- Desert greening
- Desertification
- Deserts of Australia
- International Center for Agricultural Research in the Dry Areas
- List of deserts
- List of deserts by area
- List of North American deserts
- Sediment precipitation

- Semi-arid climate

References

1. ^ [a] [b] Harper, Douglas (2012). "Desert". *Online Etymology Dictionary*. Retrieved 2013-05-12.
2. ^ "Desert". *The Free Dictionary*. Farlex. Retrieved 2013-05-12.
3. ^ "Desert Island". *The Free Dictionary*. Farlex. Retrieved 2013-05-12.
4. ^ Meinig, Donald W. (1993). *The Shaping of America: A Geographical Perspective on 500 Years of History, Volume 2: Continental America, 1800–1867*. Yale University Press. p. 76. ISBN 978-0-300-05658-7.
5. ^ Wilson, Hilary (1995). *Understanding Hieroglyphs: A Complete Introductory Guide*. Michael O'Mara Books. p. 69. ISBN 978-1-86019-905-9.
6. ^ [a] [b] Marshak (2009). *Essentials of Geology, 3rd ed.*. W. W. Norton & Co. p. 452. ISBN 978-0-393-19656-6.
7. ^ [a] [b] [c] Smith, Jeremy M. B. "Desert". Encyclopædia Britannica online. Retrieved 2013-09-24.
8. ^ [a] [b] [c] [d] "What is a desert?". United States Geological Survey. Retrieved 2013-05-23.
9. ^ Walter, Heinrich; Breckle, Siegmar-W. (2002). *Walter's Vegetation of the Earth: The Ecological Systems of the Geo-biosphere*. Springer. p. 457. ISBN 978-3-540-43315-6.
10. ^ Negi, S.S. (2002). *Cold Deserts of India*. Indus Publishing. p. 9. ISBN 978-81-7387-127-6.
11. ^ Rohli, Robert V.; Vega, Anthony J. (2008). *Climatology*. Jones & Bartlett Learning. p. 207. ISBN 978-0-7637-3828-0.
12. ^ Thomas, David Neville et al. (2008). *The biology of polar regions*. Oxford University Press. p. 64. ISBN 978-0-19-929813-6.
13. ^ Lyons, W. Berry; Howard-Williams, C. and Hawes, Ian (1997). *Ecosystem processes in Antarctic ice-free landscapes: proceedings of an International Workshop on Polar Desert Ecosystems : Christchurch, New Zealand, 1–4 July 1996*. Taylor & Francis. pp. 3–10. ISBN 978-90-5410-925-9.
14. ^ [a] [b] "Desert". *1911 Encyclopædia Britannica, Volume 8*. 1911. p. 93. Retrieved 2013-09-24.
15. ^ Buel, S. W. (1964). "Calculated actual and potential evapotranspiration in Arizona". *Tucson, Arizona University Agricultural Experiment Station Technical Bulletin* **162**: 48.
16. ^ Mendez, J.; Hinzman, L. D.; Kane, D. L. (1998). "Evapotranspiration from a wetland complex on the Arctic coastal plain of Alaska".*Nordic Hydrology* **29** (4–5): 303–330. ISSN 0029-1277.
17. ^ [a] [b] [c] [d] [e] Laity, Julie J. (2009). *Deserts and Desert Environments: Volume 3 of Environmental Systems and Global Change Series*. John Wiley & Sons. pp. 2–7, 49. ISBN 9781444300741.
18. ^ John E. Oliver (1 January 2005). *The Encyclopedia of World Climatology*. Springer. p. 86. ISBN 978-1-4020-3264-6.
19. ^ [a] [b] [c] [d] "Types of deserts". *Deserts: Geology and Resources*. United States Geological Survey. Retrieved 2013-05-11.
20. ^ "The formation of deserts". *Desert*. Oracle ThinkQuest Education Foundation. Retrieved 2013-05-11.
21. ^ Brinch, Brian (2007-11-01). "How mountains influence rainfall patterns". *USA Today*. Retrieved 2013-05-08.
22. ^ "Taklamakan Desert". *Encyclopedia Britannica online*. Retrieved 2007-08-11.
23. ^ Pidwirny, Michael (2008). "CHAPTER 8: Introduction to the Hydrosphere (e) Cloud Formation Processes". Physical Geography. Retrieved 2009-01-01.
24. ^ Mares, Michael (ed.) (1999). *Encyclopedia of Deserts: Deserts, Montane*. University of Oklahoma Press. p. 172. ISBN 978-0-8061-3146-7.
25. ^ Bockheim, J. G. (2002). "Landform and soil development in the McMurdo Dry Valleys, Antarctica: a regional synthesis". *Arctic, Antarctic, and Alpine Research* **34** (3): 308–17. doi:10.2307/1552489.
26. ^ Fredlund, D.G.; Rahardjo, H. (1993). *Soil Mechanics for Unsaturated Soils* (PDF). Wiley-Interscience. ISBN 978-0-471-85008-3. Retrieved 2008-05-21.
27. ^ Allaby, Michael (2004). "Thornthwaite climate classification". *A Dictionary of Ecology*. Encyclopedia.com. Retrieved 2013-09-23.
28. ^ [a] [b] [c] Briggs, Kenneth (1985). *Physical Geography: Process and System*. Hodder & Stoughton. pp. 8, 59–62. ISBN 978-0-340-35951-8.
29. ^ George, 1978. p. 11
30. ^ George, 1978. p. 21
31. ^ George, 1978. p. 22
32. ^ Smalley, I. J.; Vita-Finzi, C. (1968). "The formation of fine particles in sandy deserts and the nature of 'desert' loess". *Journal of Sedimentary Petrology* **38** (3): 766–774.

33. ^ Pye & Tsoar, 2009. p. 4
34. ^ [a] [b] Pye & Tsoar, 2009. p. 141
35. ^ [a] [b] Yang, Youlin; Squires, Victor; Lu, Qi, ed. (2001). "Physics, Mechanics and Processes of Dust and Sandstorms" (PDF). *Global Alarm: Dust and Sandstorms from the World's Drylands*. United Nations Convention to Combat Desertification. p. 17.
36. ^ [a] [b] [c] George, 1978. pp. 17–20
37. ^ Gu, Yingxin; Rose, William I.; Bluth, Gregg J. S. (2003). "Retrieval of mass and sizes of particles in sandstorms using two MODIS IR bands: A case study of April 7, 2001 sandstorm in China". *Geophysical Research Letters* **30** (15). Bibcode:2003GeoRL..30.1805G. doi:10.1029/2003GL017405.
38. ^ Sinclair, Peter C. (1969). "General characteristics of dust devils". *Journal of Applied Meteorology* **8** (1): 32–45. doi:10.1175/1520-0450(1969)008<0032:GCODD>2.0.CO;2.
39. ^ Zheng, Xiao Jing; Huang, Ning; Zhou, You-He (2003). "Laboratory measurement of electrification of wind-blown sands and simulation of its effect on sand saltation movement". *Journal of Geophysical Research: Atmospheres* **108** (D10). Bibcode:2003JGRD..108.4322Z. doi:10.1029/2002JD002572.
40. ^ Latham, J. (1964). "The electrification of snowstorms and sandstorms". *Quarterly Journal of the Royal Meteorological Society* **90**(383): 91–95. doi:10.1002/qj.49709038310.
41. ^ "The World's Largest Deserts". Geology.com. Retrieved 2013-05-12.
42. ^ Coakley, J. A.; Holton, J. R.; Curry, J. A. (eds.) (2002). *Reflectance and albedo, surface in "Encyclopedia of the Atmosphere"*. Academic Press. pp. 1914–1923.
43. ^ "Misconceptions surround desert terrain, vegetation". *Ask a Scientist*. Cornell Center for Materials Research. 2001-07-11. Retrieved 2013-09-24.
44. ^ "Habitats: Desert". BBC Nature. 2013. Retrieved 2013-05-23.
45. ^ [a] [b] [c] [d] [e] [f] "Desert Features". United States Geological Survey. 1997-10-29. Retrieved 2013-05-23.
46. ^ "Sand Plains/Sand Sheets". *Desert Guide*. US Army Corps of Engineers. Retrieved 2013-05-23.
47. ^ "Ripples, Sand". *Desert Guide*. US Army Corps of Engineers. Retrieved 2013-05-23.
48. ^ "Dunes, General". *Desert Guide*. US Army Corps of Engineers. Retrieved 2013-05-23.
49. ^ [a] [b] "Types of Dunes". United States Geological Survey. 1997-10-29. Retrieved 2013-05-23.
50. ^ "Desert pavement". *Encyclopædia Britannica online*. Retrieved 2013-05-23.
51. ^ Perry, R. S.; Adams, J. B. (1978). "Desert varnish: evidence for cyclic deposition of manganese". *Nature* **276** (5687): 489–491. doi:10.1038/276489a0.
52. ^ "Hamada, Reg, Serir, Gibber, Saï". Springer Reference. 2013. Retrieved 2013-05-23.
53. ^ George, 1978. pp. 29–30
54. ^ Foos, Annabelle. "Geology of Grand Canyon National Park, North Rim". Retrieved 2013-09-24.
55. ^ [a] [b] Westbeld, A.; Klemm, O.; Grießbaum, F.; Strater, E.; Larrain, H.; Osses, P.; Cereceda, P. (2009). "Fog deposition to a *Tillandsia*carpet in the Atacama Desert". *Annales Geophysicae* **27**: 3571–3576. doi:10.5194/angeo-27-3571-2009.
56. ^ McKay, C. P. (May–June 2002). "Too dry for life: The Atacama Desert and Mars" (PDF). *Ad Astra* (NASA): 30. Retrieved 2010-10-16.
57. ^ Boehm, Richard G.; Editors and writers of *The World and Its People* (2006). *The World and Its People* (2005 ed.). Glencoe. p. 276. ISBN 978-0-07-860977-0.
58. ^ Preston, Benjamin (2011-04-01). "Colonel Qaddafi and the Great Man-made River". *State of the Planet*. Earth Institute: Columbia University. Retrieved 2013-10-02.
59. ^ Bayfield, Su (2011). "Introduction to Kharga Oasis". *Egyptian monuments*. Retrieved 2013-10-02.
60. ^ "Desert Survival". Public Broadcasting Service. Retrieved 2010-10-16.
61. ^ "Lake Bonneville". Utah Geological Survey. Retrieved 2013-05-24.
62. ^ Eduardo Zeiger (1987). *Stomatal function*. Stanford, Calif.: Stanford University Press. ISBN 9780804713474. Retrieved 22 December 2013.
63. ^ Osborne, Colin P.; Beerling, David J. (2006). "Nature's green revolution: the remarkable evolutionary rise of C4 plants". *Philosophical Transactions of the Royal Society B* **361** (1465): 173–194. doi:10.1098/rstb.2005.1737. ISSN 1471-2970. PMC 1626541. PMID 16553316.
64. ^ [a] [b] [c] [d] George, 1978. pp. 122–123
65. ^ Council-Garcia, Cara Lea (2002). "Plant adaptations". University of New Mexico. Retrieved 2013-09-24.

66. ^ [a] [b] "Desert Flora". Australian Department of the Environment and Heritage. Retrieved 2013-05-13.
67. ^ Dimmitt, Mark A. (1997). "How Plants Cope with the Desert Climate". Arizona-Sonora Desert Museum. Retrieved 2013-05-13.
68. ^ "Cold Deserts". *The desert biome*. University of California Museum of Paleontology. 1996. Retrieved 2013-09-23.
69. ^ [a] [b] Scholander, P. F.; Hock, Raymond; Walters, Vladimir; Irving, Laurence (1950). "Adaptation to cold in arctic and tropical mammals and birds in relation to body temperature, insulation, and basal metabolic rate". *Biological Bulletin* **50** (2): 269.
70. ^ Al-kahtani, M.A.; C. Zuleta, E. Caviedes-Vidal, and T. Garland, Jr. (2004). "Kidney mass and relative medullary thickness of rodents in relation to habitat, body size, and phylogeny". *Physiological and Biochemical Zoology* **77** (3): 346–365. doi:10.1086/420941. PMID 15286910.
71. ^ Pianka, Eric R. "Convergent Evolution". *Biology Reference*. Retrieved 2013-05-28.
72. ^ George, 1978. p. 141
73. ^ Campbell, Mary K; Farrell, Shawn O (2006). *Biochemistry (fifth edition)*. USA: Thomson Brooks/Cole. p. 511. ISBN 0-453-40521-5.
74. ^ Morrison, S. D. (1953). "A method for the calculation of metabolic water". *J. Physiology* **122** (2): 399–402. Morrison cites Brody, S.*Bioenergetics and Growth*. Reinhold, 1945. p. 36 for the figures.
75. ^ [a] [b] Mellanby, Kenneth (1942). "Metabolic water and desiccation". *Nature* **150**: 21. doi:10.1038/150021a0. ISSN 0028-0836.
76. ^ Best, T. L., et al. (1989) Dipodomys deserti. Mammalian Species 339:1-8 [1]
77. ^ Lidicker, W. Z. (1960). *An Analysis of Intraspecific Variation in the Kangaroo Rat* Dipodomus merriami. University of California Press.
78. ^ Lacher, Jr., Thomas E. *Encyclopedia of Deserts: Addax*. University of Oklahoma Press. p. 7. ISBN 978-0-8061-3146-7.
79. ^ Maloiy, G. M. O. (November 1973). "The water metabolism of a small East African antelope: the dik-dik". *Proceedings of the Royal Society B* **184** (1075): 167–178. doi:10.1098/rspb.1973.0041.
80. ^ Vann Jones, Kerstin. "What secrets lie within the camel's hump?". Lund University. Retrieved 2013-05-21.
81. ^ [a] [b] Silverstein, Alvin; Silverstein, Virginia B.; Silverstein, Virginia; Silverstein Nunn, Laura (2008). *Adaptation*. Twenty-First Century Books. pp. 42–43. ISBN 978-0-8225-3434-1.
82. ^ Monroe, M. H. "The Red Kangaroo". *Australia: The Land Where Time Began*. Retrieved 2013-10-03.
83. ^ Hile, J. (2004-03-29). "Emperor Penguins: Uniquely Armed for Antarctica". *National Geographic*. Retrieved 2013-10-02.
84. ^ Lacher, Jr., Thomas E. *Encyclopedia of Deserts:* Cerastes. p. 108. ISBN 978-0-8061-3146-7.
85. ^ "Couch's spadefoot (*Scaphiopus couchi*)". Arizona-Sonora Desert Museum. Retrieved 2013-05-21.
86. ^ Withers, P. C. (1993). "Metabolic Depression During Estivation in the Australian Frogs, *Neobatrachus* and *Cyclorana*". *Australian Journal of Zoology* **41** (5): 467–473. doi:10.1071/ZO9930467.
87. ^ Castillo, Nery (2011-06-23). "*Breviceps macrops*". AmphibiaWeb. Retrieved 2012-10-20.
88. ^ "Invertebrates: A Vertebrate Looks at Arthropods". Arizona-Sonora Desert Museum. Retrieved 2013-05-21.
89. ^ "Invertebrates in the Desert". *ThinkQuest*. Oracle. Retrieved 2013-05-22.
90. ^ Moseley, Pope L. (1997). "Heat shock proteins and heat adaptation of the whole organism". *Journal of Applied Physiology* **83** (5): 1413–1417.
91. ^ Picker, Mike; Griffiths, Charles; Weaving, Alan (2004). *Field Guide to the Insects of South Africa*. Struik. p. 232. ISBN 978-1-77007-061-5.
92. ^ "Ephemeral Pools". *Arches National Park, Utah*. National Park Service. Retrieved 2013-05-22.
93. ^ [a] [b] Fagan, Brian M. (2004). *People of the Earth*. Pearson Prentice Hall. pp. 169–181. ISBN 978-0-205-73567-9.
94. ^ [a] [b] [c] [d] [e] [f] [g] [h] "Mineral Resources in Deserts". US Geological Survey. 1997-10-29. Retrieved 2013-05-24.
95. ^ [a] [b] Waldoks, Ehud Zion (2008-03-18). "Head of Kibbutz Movement: We will not be discriminated against by the government". *The Jerusalem Post*. Retrieved 2013-09-22.
96. ^ [a] [b] Bancroft, John (ed.) (1994). "The Deserts in Literature". *The Arid Lands Newsletter* (35). ISSN 1092-5481.
97. ^ Dyson-Hudson, Rada; Dyson-Hudson, Neville (1980). "Nomadic pastoralism". *Annual Review of Anthropology* **9**: 15–61. doi:10.1146/annurev.an.09.100180.000311. JSTOR 2155728.
98. ^ Masonen, Pekka (1995). "Trans-Saharan trade and the West African discovery of the Mediterranean". *Nordic Research on the Middle East* **3**: 116–142.
99. ^ Wright, John (2007). *The Trans-Saharan Slave Trade*. Routledge. p. 22. ISBN 978-0-203-96281-7.
100. ^ "Sahara salt trade camel caravans". *National Geographic News*. 2010-10-28. Retrieved 2013-09-22.

101. ^ "First Measured Century: Interview:James Gregory". Public Broadcasting Service. Retrieved 2013-05-25.
102. ^ "Desertification: Facts and figures". United Nations. Retrieved 2013-05-26.
103. ^ Geeson, Nicola; Brandt, C. J.; Thornes, J. B. (2003). *Mediterranean Desertification: A Mosaic of Processes and Responses*. Wiley. p. 58. ISBN 978-0-470-85686-4.
104. ^ Savory, Allen (2013-03-04). "How to green the world's deserts and reverse climate change".
105. ^ van Dalen, Dorrit (2009) *Landenreeks Tsjaad*, KIT Publishers, ISBN 978-90-6832-690-1.
106. ^ Anderson, Roger N. (2006-01-16). "Why is oil usually found in deserts and Arctic areas?". *Scientific American*. Retrieved 2013-05-26.
107. ^ [a] [b] [c] Marasco, Ramona; Rolli, Eleonora; Ettoumi, Besma; Vigani, Gianpiero; Mapelli, Francesca; Borin, Sara; Abou-Hadid, Ayman F.; El-Behairy, Usama A.; Sorlini, Claudia; Cherif, Ameur; Zocchi, Graziano; Daffonchio, Daniele (2012). "A drought resistance-promoting microbiome is selected by root system under desert farming". *PLOSone*.
108. ^ Stager, Lawrence E. (1976). "Farming in the Judean Desert during the Iron Age". *Bulletin of the American Schools of Oriental Research* **221**: 145–158. JSTOR 1356097.
109. ^ Smith, Chuck (2002-10-14). "Agricultural Societies In Pre-European Times: Southwestern U.S. and Northwestern Mexico". *Native Peoples of North America*. Retrieved 2013-09-28.
110. ^ "Imperial County Agriculture". University of California Cooperative Extension. 2012-02-15. Retrieved 2013-09-28.
111. ^ Meadows, Robin (2012). "Research news: UC Desert Research and Extension Center celebrates 100 years". *California Agriculture***66** (4): 122–126. doi:10.3733/ca.v066n04p122.
112. ^ Clemings, R. (1996). *Mirage: the false promise of desert agriculture*. Sierra Club Books. pp. 1–247. ISBN 0-87156-416-5.
113. ^ "Solar Trough Systems". National Renewable Energy Laboratory. Retrieved 2013-09-22.
114. ^ Parry, Tom (2007-08-15). "Looking to the sun". Canadian Broadcasting Corporation. Retrieved 2013-09-22.
115. ^ Sandler, Neal (2006-02-14). "Israeli Solar Startup Shines". Businessweek.com. Retrieved 2010-10-16.
116. ^ [a] [b] Lettice, John (2008-01-25). "Giant solar plants in Negev could power Israel's future". *The Register*. Retrieved 2013-09-22.
117. ^ Matlack, Carol (2010-12-16). "Sahara Solar Energy Could Power Europe". *BloombergBusinessweek*. Bloomberg. Retrieved 2013-09-22.
118. ^ Lawrence, T. E. (1922). *Seven Pillars of Wisdom*. Private edition.
119. ^ Murphy, David (2008). *The Arab Revolt 1916–18: Lawrence Sets Arabia Ablaze*. Osprey. ISBN 978-1-84603-339-1.
120. ^ Woolley, Jo (2008). "Desert warfare". *History Today* **52** (10).
121. ^ Latimer, Jon (2002). *Alamein*. John Murray. ISBN 978-0-7195-6203-7.
122. ^ "Writing the Void: Desert Literature". University of Strathclyde. Retrieved 2013-09-24.
123. ^ Bergreen, Laurence (2007). *Marco Polo: From Venice to Xanadu*. Quercus. pp. 1–415. ISBN 978-1-84724-345-4.
124. ^ Taylor, Andrew (1999). *God's Fugitive*. Harper Collins. pp. 295–298. ISBN 978-0-00-255815-0.
125. ^ Hutchinson, Charles F. (1994). "Wind, Sand and Stars Revisited". Retrieved 2013-05-26.
126. ^ Howell, Georgina (2007). *Gertrude Bell: Queen of the Desert, Shaper of Nations*. Farrar, Straus and Giroux. ISBN 978-0-374-16162-0.
127. ^ Moorehead, C. (1985). *Freya Stark*. Penguin. ISBN 978-0-14-008108-4.
128. ^ George, 1978.
129. ^ Frost, Robert. "Desert Places". Poemhunter. Retrieved 2013-05-26.
130. ^ "Strange Land Formations on Mars". *The Blue Bird Files*. 2007-04-11. Retrieved 2013-09-27.
131. ^ "Do Mars Rocks Have Desert Varnish?". *Astrobiology*. 2013-03-23. Retrieved 2013-09-27.
132. ^ Arnold, K.; Radebaugh, J.; Savage, C. J.; Turtle, E. P.; Lorenz, R. D.; Stofan, E. R.; Le-Gall, A. "Areas of Sand Seas on Titan from Cassini Radar and ISS: Fensal and Aztlan". *42nd Lunar and Planetary Science Conference, March 7–11, 2011 at The Woodlands, Texas. LPI Contribution No. 1608*. Lunar and Planetary Institute.

Bibliography

Further reading

- Bagnold, Ralph A. (1941). *The physics of blown sand and desert dunes*. Methuen.
- Macmahon, James (1988). *Deserts*. National Audubon Society nature guides. Random House / Chanticleer Press. ISBN 978-0-394-73139-1.

External links

- "Global Deserts Outlook". United Nations Environment Programme (UNEP). 2006., a report in the Global Environment Outlook (GEO) series.
  - Global Deserts Outlook in PDF at the Wayback Machine (archived June 10, 2006)
  - Map with biodiversity scenarios for desert areas, from the Global Deserts Outlook.


CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Deserts. **License**: *CC BY-SA: Attribution-ShareAlike*


This page titled 19.1: Deserts is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

19.2: Desertification

Desertification is a type of land degradation in which a relatively dry land region becomes increasingly arid, typically losing its bodies of water as well as vegetation and wildlife.[2] It is caused by a variety of factors, such as climate change and human activities. Desertification is a significant global ecological and environmental problem.[3]



Global Desertification Vulnerability Map



Lake Chad in a 2001 satellite image, with the actual lake in blue. The lake has shrunk by 94% since the 1960s.[1]

Definitions

Considerable controversy exists over the proper definition of the term “desertification” for which Helmut Geist (2005) has identified more than 100 formal definitions.The most widely accepted[2] of these is that of the Princeton University Dictionary which defines it as “the process of fertile land transforming into desert typically as a result of deforestation, drought or improper/inappropriate agriculture”[4]

The earliest known discussion of the topic arose soon after the French colonization of West Africa, when the Comité d’Etudes commissioned a study on *desséchement progressif* to explore the prehistoric expansion of the Sahara Desert.[5]

History

The world’s most noted deserts have been formed by natural processes interacting over long intervals of time. During most of these times, deserts have grown and shrunk independent of human activities. Paleodeserts are large sand seas now inactive because they are stabilized by vegetation, some extending beyond the present margins of core deserts, such as the Sahara, the largest hot desert. [6]

Desertification has played a significant role in human history, contributing to the collapse of several large empires, such as Carthage, Greece, and the Roman Empire, as well as causing displacement of local populations.[7][3][8][9][10]Historical evidence shows that the serious and extensive land deterioration occurring several centuries ago in arid regions had three epicenters: the Mediterranean, the Mesopotamian Valley, and the loessial plateau of China, where population was dense.[11][12]

Areas affected



Sun, Moon and Telescopes above the Desert.[13]

Drylands occupy approximately 40–41% of Earth's land area[14][15] and are home to more than 2 billion people.[15] It has been estimated that some 10–20% of drylands are already degraded, the total area affected by desertification being between 6 and 12 million square kilometres, that about 1–6% of the inhabitants of drylands live in desertified areas, and that a billion people are under threat from further desertification.[16][17]

As of 1998, the then-current degree of southward expansion of the Sahara was not well known, due to a lack of recent, measurable expansion of the desert into the Sahel at the time.[18]

Vegetation patterning

As the desertification takes place, the landscape may progress through different stages and continuously transform in appearance. On gradually sloped terrain, desertification can create increasingly larger empty spaces over a large strip of land, a phenomenon known as "Brousse tigrée". A mathematical model of this phenomenon proposed by C. Klausmeier attributes this patterning to dynamics in plant-water interaction.[19] One outcome of this observation suggests an optimal planting strategy for agriculture in arid environments.[20]

Causes



A herd of goats in Norte Chico, Chile. Overgrazing of drylands is one of the primary causes of desertification.



A shepherd guiding his sheep through the high desert outside of Marrakech, Morocco.

The immediate cause is the removal of most vegetation. This is driven by a number of factors, alone or in combination, such as drought, climatic shifts, tillage for agriculture, overgrazing and deforestation for fuel or construction materials. Vegetation plays a major role in determining the biological composition of the soil. Studies have shown that, in many environments, the rate of erosion and runoff decreases exponentially with increased vegetation cover.[21] Unprotected, dry soil surfaces blow away with the wind or are washed away by flash floods, leaving infertile lower soil layers that bake in the sun and become an unproductive hardpan. Alternatively, recent research has suggested that the movement and migration of large herds of livestock and wildlife has an integral role in the preservation of vegetation and soil fertilization, and that the removal of livestock and wildlife (largely by human influence) has been the main driver of increasing desertification.[22][23][24][25][26]

Poverty

At least 90% of the inhabitants of drylands live in developing nations, where they also suffer from poor economic and social conditions.[16] This situation is exacerbated by land degradation because of the reduction in productivity, the precariousness of

living conditions and the difficulty of access to resources and opportunities.[27]

A downward spiral is created in many underdeveloped countries by overgrazing, land exhaustion and overdrafting of groundwater in many of the marginally productive world regions due to overpopulation pressures to exploit marginal drylands for farming. Decision-makers are understandably averse to invest in arid zones with low potential. This absence of investment contributes to the marginalisation of these zones. When unfavourable agro-climatic conditions are combined with an absence of infrastructure and access to markets, as well as poorly adapted production techniques and an underfed and undereducated population, most such zones are excluded from development.[28]

Desertification often causes rural lands to become unable to support the same sized populations that previously lived there. This results in mass migrations out of rural areas and into urban areas, particularly in Africa. These migrations into the cities often cause large numbers of unemployed people, who end up living in slums.[29][30]

Countermeasures and prevention



Anti-sand shields in north Sahara, Tunisia.



Jojoba plantations, such as those shown, have played a role in combating edge effects of desertification in the Thar Desert, India. [*citation needed*]

Techniques exist for mitigating or reversing the effects of desertification, however there are numerous barriers to their implementation. One of these is that the costs of adopting sustainable agricultural practices sometimes exceed the benefits for individual farmers, even while they are socially and environmentally beneficial.[*citation needed*] Another issue is a lack of political will, and lack of funding to support land reclamation and anti-desertification programs.[31]

Desertification is recognized as a major threat to biodiversity. Some countries have developed Biodiversity Action Plans to counter its effects, particularly in relation to the protection of endangered flora and fauna.[32][33]

Reforestation gets at one of the root causes of desertification and isn't just a treatment of the symptoms. Environmental organizations[34] work in places where deforestation and desertification are contributing to extreme poverty. There they focus primarily on educating the local population about the dangers of deforestation and sometimes employ them to grow seedlings, which they transfer to severely deforested areas during the rainy season.[35]

Techniques focus on two aspects: provisioning of water, and fixation and hyper-fertilizing soil.

Fixating the soil is often done through the use of shelter belts, woodlots and windbreaks. Windbreaks are made from trees and bushes and are used to reduce soil erosion and evapotranspiration. They were widely encouraged by development agencies from the middle of the 1980s in the Sahel area of Africa.

Some soils (for example, clay), due to lack of water can become consolidated rather than porous (as in the case of sandy soils). Some techniques as zaï or tillage are then used to still allow the planting of crops.[36]

Another technique that is useful is contour trenching. This involves the digging of 150m long, 1m deep trenches in the soil. The trenches are made parallel to the height lines of the landscape, preventing the water of flowing within the trenches and causing

erosion. Stone walls are placed around the trenches to prevent the trenches of closing up again. The method was invented by Peter Westerveld.[37]

Enriching of the soil and restoration of its fertility is often done by plants. Of these, the Leguminous plants which extract nitrogen from the air and fixes it in the soil, and food crops/trees as grains, barley, beans and dates are the most important. Sand fences can also be used to control drifting of soil and sand erosion.[38]

As there are many different types of deserts, there are also different types of desert reclamation methodologies. An example for this is the sa ">[39]

Farmer-managed natural regeneration (FMNR) is another technique that has produced successful results for desert reclamation. Since 1980, this method to reforest degraded landscape has been applied with some success in Niger. This simple and low-cost method has enabled farmers to regenerate some 30,000 square kilometers in Niger. The process involves enabling native sprouting tree growth through selective pruning of shrub shoots. The residue from pruned trees can be used to provide mulching for fields thus increasing soil water retention and reducing evaporation. Additionally, properly spaced and pruned trees can increase crop yields. The Humbo Assisted Regeneration Project which uses FMNR techniques in Ethiopia has received money from The World Bank's BioCarbon Fund, which supports projects that sequester or conserve carbon in forests or agricultural ecosystems.[40]



Managed grazing methods are argued to be able to restore grasslands, thereby decreasing atmospheric CO_2 levels.[41]

Managed grazing

Restoring grasslands store CO_2 from the air into plant material. Grazing livestock, usually not left to wander, would eat the grass and would minimize any grass growth while grass left alone would eventually grow to cover its own growing buds, preventing them from photosynthesizing and killing the plant.[42] A method proposed to restore grasslands uses fences with many small paddocks and moving herds from one paddock to another after a day a two in order to mimick natural grazers and allowing the grass to grow optimally.[42][43][44] It is estimated that increasing the carbon content of the soils in the world's 3.5 billion hectares of agricultural grassland would offset nearly 12 years of CO_2 emissions.[42]Allan Savory, as part of holistic management, claims that while large herds are often blamed for desertification, prehistoric lands used to support large or larger herds and areas where herds were removed in the United States are still desertifying.[41]



Visible soil erosion outside of Leuchars.

Mitigation:

- Arid Lands Information Network—Kenya
- Biochar Fertilisation using carbon
- United Nations Convention to Combat Desertification
- Desert greening
- Ecological engineering
- Green Wall of China

- Holistic management

References

1. ^ Mayell, Hillary (April 26, 2001). “Shrinking African Lake Offers Lesson on Finite Resources”. *National Geographic News*. Retrieved 20 June 2011.
2. ^ [a] [b] Geist (2005), p. 2
3. ^ [a] [b] Geist (2005), p. 4
4. ^ Google Dictionary (2012)
5. ^ Mortimore, Michael (1989). *Adapting to drought: farmers, famines, and desertification in west Africa*. Cambridge University Press. p. 12. ISBN 978-0-521-32312-3.
6. ^ United States Geological Survey, “Desertification”, 1997
7. ^ LOWDERMILK, W C. “CONQUEST OF THE LAND THROUGH SEVEN THOUSAND YEARS”. *Soil Conservation Service*. United States Department of Agriculture. Retrieved 9 April 2014.
8. ^ Whitford, Walter G. (2002). *Ecology of desert systems*. Academic Press. p. 277. ISBN 978-0-12-747261-4.
9. ^ Bogumil Terminski (2011), Towards Recognition and Protection of Forced Environmental Migrants in the Public International Law: Refugee or IDPs Umbrella, Policy Studies Organization (PSO), Washington.
10. ^ Geist, Helmut. “The causes and progression of desertification”. *Antony Rowe Ltd*. Ashgate publishing limited. Retrieved 6 July 2013.
11. ^ LOWDERMILK, W C. “CONQUEST OF THE LAND THROUGH SEVEN THOUSAND YEARS”. *Soil Conservation Service*. United States Department of Agriculture. Retrieved 9 April 2014.
12. ^ Dregne, H.E. “Desertification of Arid Lands”. Columbia University. Retrieved 3 December 2013.
13. ^ “Sun, Moon and Telescopes above the Desert”. *ESO Picture of the Week*. Retrieved 30 April 2012.
14. ^ Bauer (2007), p. 78
15. ^ [a] [b] Johnson et al (2006), p. 1
16. ^ [a] [b] Holtz (2007)
17. ^ World Bank (2009). *Gender in agriculture sourcebook*. World Bank Publications. p. 454. ISBN 978-0-8213-7587-7.
18. ^ [1]
19. ^ Klausmeier, Christopher (1999). “Regular and irregular patterns in semiarid vegetation”. *Science* **284** (5421): 1826–1828. doi:10.1126/science.284.5421.1826.
20. ^ Straw squares method: planting with empty spaces in between plants to combat desertification
21. ^ Geeson, Nichola et al (2002). *Mediterranean desertification: a mosaic of processes and responses*. John Wiley & Sons. p. 58. ISBN 978-0-470-84448-9.
22. ^ Savory, Allan. “Allan Savory: How to green the world’s deserts and reverse climate change”.
23. ^ Savory, Allan. “Holistic resource management: a conceptual framework for ecologically sound economic modelling”. *Ecological Economics*. Elsevier Science Publishers. Retrieved 10 March 2013.
24. ^ Butterfield, Jody (2006). *Holistic Management Handbook: Healthy Land, Healthy Profits, Second Edition*. Island Press. ISBN 1559638850.
25. ^ Savory, Allan. “Response to request for information on the “science” and “methodology” underpinning Holistic Management and holistic planned grazing”. *Savory Institute*. Retrieved 10 March 2013.
26. ^ Drury, Steve. “Large-animal extinction in Australia linked to human hunters”. Earth-Pages. Retrieved 9 June 2014.
27. ^ Dobie, Ph. 2001. “Poverty and the drylands”, in Global Drylands Imperative, Challenge paper, Undp, Nairobi (Kenya) 16 p.
28. ^ Cornet A., 2002. Desertification and its relationship to the environment and development: a problem that affects us all. In: Ministère des Affaires étrangères/adpf, Johannesburg. World Summit on Sustainable Development. 2002. What is at stake? The contribution of scientists to the debate: 91-125..
29. ^ Pasternak, Dov & Schlissel, Arnold (2001). *Combating desertification with plants*. Springer. p. 20. ISBN 978-0-306-46632-8.
30. ^ Briassoulis, Helen (2005). *Policy integration for complex environmental problems: the example of Mediterranean desertification*. Ashgate Publishing. p. 161. ISBN 978-0-7546-4243-5.
31. ^ Briassoulis, Helen (2005). *Policy integration for complex environmental problems: the example of Mediterranean desertification*. Ashgate Publishing. p. 237. ISBN 978-0-7546-4243-5.
32. ^ Techniques for Desert Reclamation by Andrew S. Goudie

33. ^ Desert reclamation projects
34. ^ For example, Eden Reforestation Projects.
35. ^ This article incorporates public domain material from the United States Government document "http://pubs.usgs.gov/gip/deserts/desertification/".
36. ^ Arid sandy soils becoming consolidated; zai-system
37. ^ Westerveld's Naga Foundation
38. ^ List of plants to halt desertification; some of which may be soil-fixating
39. ^ Rethinking landscapes, Nicol-André Berdellé July 2011 H2O magazine
40. ^ "Sprouting Trees From the Underground Forest — A Simple Way to Fight Desertification and Climate Change – Water Matters – State of the Planet". Blogs.ei.columbia.edu. 2011-10-18. Retrieved 2012-08-11.
41. ^ [a] [b] "How cows could repair the world". nationalgeographic.com. March 6, 2013. Retrieved May 5, 2013.
42. ^ [a] [b] [c] "How fences could save the planet". newstatesman.com. January 13, 2011. Retrieved May 5, 2013.
43. ^ "Restoring soil carbon can reverse global warming, desertification and biodiversity". mongabay.com. February 21, 2008. Retrieved May 5, 2013.
44. ^ "How eating grass-fed beef could help fight climate change". time.com. January 25, 2010. Retrieved May 11, 2013.

Bibliography

- Arnalds, Ólafur & Archer, Steve (2000). *Rangeland Desertification*. Springer. ISBN 978-0-7923-6071-1.
- Barbault R., Cornet A., Jouzel J., Mégie G., Sachs I., Weber J. (2002). Johannesburg. World Summit on Sustainable Development. 2002. What is at stake? The contribution of scientists to the debate. Ministère des Affaires étrangères/adpf.
- Bauer, Steffan (2007). "Desertification". In Thai, Khi V. et al. *Handbook of globalization and the environment*. CRC Press. ISBN 978-1-57444-553-4.
- Batterbury, S.P.J. & A.Warren (2001) Desertification. in N. Smelser & P. Baltes (eds.) International Encyclopædia of the Social and Behavioral Sciences. Elsevier Press. pp. 3526–3529
- Geist, Helmut (2005). *The causes and progression of desertification*. Ashgate Publishing. ISBN 978-0-7546-4323-4.
- Hartman, Ingrid (2008). "Desertification". In Philander, S. George. *Encyclopedia of global warming and climate change, Volume 1*. SAGE. ISBN 978-1-4129-5878-3.
- Hinman, C. Wiley & Hinman, Jack W. (1992). *The plight and promise of arid land agriculture*. Columbia University Press. ISBN 978-0-231-06612-9.
- Holtz, Uwe (2007). *Implementing the United Nations Convention to Combat Desertification from a parliamentary point of view – Critical assessment and challenges ahead*. Online at [2]
- Holtz, Uwe (2013). *Role of parliamentarians in the implementation process of the UN Convention to Combat Desertification. A guide to Parliamentary Action*, ed. Secretariat of the UNCCD, Bonn ISBN 978-92-95043-69-5. Online at [3]
- Johnson, Pierre Marc et al., ed. (2006). *Governing global desertification: linking environmental degradation, poverty and participation*. Ashgate Publishing. ISBN 978-0-7546-4359-3.
- Lucke, Bernhard (2007): Demise of the Decapolis. Past and Present Desertification in the Context of Soil Development, Land Use, and Climate. Online at [4]
- Mensah, Joseph (2006). "Desertification". In Leonard, Thomas M. *Encyclopedia of the developing world, Volume 1*. Taylor & Francis. ISBN 978-0-415-97662-6.
- Millennium Ecosystem Assessment (2005) *Desertification Synthesis Report*
- Moseley, W.G. and E. Jerme 2010. "Desertification." In: Warf, B. (ed). Encyclopedia of Geography. Sage Publications. Volume 2, pp. 715–719.
- Oliver, John E., ed. (2005). "Desertification". *Encyclopedia of world climatology*. Springer. ISBN 978-1-4020-3264-6.
- Parrillo, Vincent N., ed. (2008). "Desertification". *Encyclopedia of social problems, Volume 2*. SAGE. ISBN 978-1-4129-4165-5.
- Reynolds, James F., and D. Mark Stafford Smith (ed.) (2002) *Global Desertification – Do Humans Cause Deserts?* Dahlem Workshop Report 88, Berlin: Dahlem University Press
- Stelt, Sjors van der (2012) *Rise and Fall of Periodic Patterns for a Generalized Klausmeier-Gray-Scott Model*, PhD Thesis University of Amsterdam
- UNCCD (1994) United Nations Convention to Combat Desertification

- The End of Eden a 90-minute documentary by South African filmmaker Rick Lomba in 1984 on African desertification

Attribution

- This article incorporates public domain material from the United States Government document "http://pubs.usgs.gov/gip/deserts/desertification/".

External links

- Beyerlin, Ulrich. Desertification, Max Planck Encyclopedia of Public International Law
- Bell, Trudy; Phillips, Tony, Dr. (December 6, 2002). "City-swallowing Sand Dunes". NASA. Retrieved 2006-04-28.
- Desert Research Institute in Nevada, United States
- Environmental Issues – Desertification in Africa, The Environmental Blog
- Eden Foundation article on desertification
- FAO Information Portal – Properties and Management of Drylands
- UNEP (2006): *Global Deserts Outlook*
- UNEP *Programme on Success Stories in Land Degradation/ Desertification Control*
- United Nations Convention to Combat Desertification – Secretariat
- Procedural history and related documents on the *United Nations Convention to Combat Desertification in those Countries Experiencing Serious Drought and/or Dersertification, Particularly in Africa* in the Historic Archives of the United Nations Audiovisual Library of International Law
- A guide for desert and dryland restoration by David A. Bainbridge
- French Scientific Committee on Desertification (CSFD)
- Olive Trees May Be The Answer To Desertification
- The End of Eden on Youtube

News

- Fighting Desertification Through Conservation Report on a project to stop the advance of the Sahara in Algeria – IPS, 27 February 2007

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Desertification. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 19.2: Desertification is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

19.3: Dunes

In physical geography, a **dune** is a hill of sand built by either wind or water flow. Dunes occur in different forms and sizes, formed by interaction with the flow of air or water. Most kinds of dunes are longer on the windward side where the sand is pushed up the dune and have a shorter “slip face” in the lee of the wind. The valley or trough between dunes is called a *slack*. A “dune field” is an area covered by extensive sand dunes. Large dune fields are known as ergs.

Some coastal areas have one or more sets of dunes running parallel to the shoreline directly inland from the beach. In most cases the dunes are important in protecting the land against potential ravages by storm waves from the sea. Although the most widely distributed dunes are those associated with coastal regions, the largest complexes of dunes are found inland in dry regions and associated with ancient lake or sea beds.

Dunes also form under the action of water flow (fluvial processes), and on sand or gravel beds of rivers, estuaries and the sea-bed.


The modern word “dune” came into English from French circa 1790. In ancient times, words cognate to “dune” probably had the meaning of a bui ">[1]



Erg Chebbi, Morocco

Conservation

Dune habitats provide niches for highly specialised plants and animals, including numerous rare species and some endangered species. Due to widespread human population expansion, dunes face destruction through land development and recreational usages, as well as alteration to prevent the encroachment of sand onto inhabited areas. Some countries, notably the United States, Australia, Canada, New Zealand, the United Kingdom, and Netherlands, have developed significant programs of dune protection through the use of sand dune stabilization. In the U.K., a Biodiversity Action Plan has been developed to assess dunes loss and to prevent future dunes destruction.





Sand blowing off a crest in the Kelso Dunes of the Mojave Desert, California.

Aeolian dune shapes

Crescentic

“Crescent-shaped mounds are generally wider than they are long. The slipfaces are on the concave sides of the dunes. These dunes form under winds that blow consistently from one direction, and they also are known as barchans, or transverse dunes. Some types of crescentic dunes move more quickly over desert surfaces than any other type of dune. A group of dunes moved more than 100 metres per year between 1954 and 1959 in the China’s Ningxia Province, and similar speeds have been recorded in the Western

Desert of Egypt. The largest crescentic dunes on Earth, with mean crest-to-crest widths of more than 3 kilometres, are in China's Taklamakan Desert."[2]

Linear

"Straight or slightly sinuous sand ridges typically much longer than they are wide are known as linear dunes. They may be more than 160 kilometres (100 miles) long. Some linear dunes merge to form Y-shaped compound dunes. Many form in bidirectional wind regimes. The long axes of these dunes extend in the resultant direction of sand movement."[3]

Linear loess hills known as pahas are superficially similar. These hills appear to have been formed during the last ice age under permafrost conditions dominated by sparse tundra vegetation.

Star

Radially symmetrical, star dunes are pyramidal sand mounds with slipfaces on three or more arms that radiate from the high center of the mound. They tend to accumulate in areas with multidirectional wind regimes. Star dunes grow upward rather than laterally. They dominate the Grand Erg Oriental of the Sahara. In other deserts, they occur around the margins of the sand seas, particularly near topographic barriers. In the southeast Badain Jaran Desert of China, the star dunes are up to 500 metres tall and may be the tallest dunes on Earth.

Dome

Oval or circular mounds that generally lack a slipface. Dome dunes are rare and occur at the far upwind margins of sand seas.

Parabolic

U-shaped mounds of sand with convex noses trailed by elongated arms are parabolic dunes. These dunes are formed from blowout dunes where the erosion of vegetated sand leads to a U-shaped depression. The elongated arms are held in place by vegetation; the largest arm known on Earth reaches 12 km. Sometimes these dunes are called U-shaped, blowout, or hairpin dunes, and they are well known in coastal deserts. Unlike crescent shaped dunes, their crests point upwind. The bulk of the sand in the dune migrates forward.

In plan view, these are U-shaped or V-shaped mounds of well-sorted, very fine to medium sand with elongated arms that extend upwind behind the central part of the dune. There are slip faces that often occur on the outer side of the nose and on the outer slopes of the arms.

These dunes often occur in semiarid areas where the precipitation is retained in the lower parts of the dune and underlying soils. The stability of the dunes was once attributed to the vegetative cover but recent research has pointed to water as the main source of parabolic dune stability. The vegetation that covers them-grasses,shrubs,and trees, help anchor the trailing arms. In inland deserts, parabolic dunes commonly originate and extend downwind from blowouts in sand sheets only partly anchored by vegetation. They can also originate from beach sands and extend inland into vegetated areas in coastal zones and on shores of large lakes.

Most parabolic dunes do not reach heights higher than a few tens of metres except at their nose, where vegetation stops or slows the advance of accumulating sand.

Simple parabolic dunes have only one set of arms that trail upwind, behind the leading nose. Compound parabolic dunes are coalesced features with several sets of trailing arms. Complex parabolic dunes include subsidiary superposed or coalesced forms, usually of barchanoid or linear shapes.

Parabolic dunes, like crescent dunes, occur in areas where very strong winds are mostly unidirectional. Although these dunes are found in areas now characterized by variable wind speeds, the effective winds associated with the growth and migration of both the parabolic and crescent dunes probably are the most consistent in wind direction.

The grain size for these well-sorted, very fine to medium sands is about 0.06 to 0.5 mm. Parabolic dunes have loose sand and steep slopes only on their outer flanks. The inner slopes are mostly well packed and anchored by vegetation, as are the corridors between individual dunes. Because all dune arms are oriented in the same direction, and, the inter-dune corridors are generally swept clear of loose sand, the corridors can usually be traversed in between the trailing arms of the dune. However to cross straight over the dune by going over the trailing arms, can be very difficult. Also, traversing the nose is very difficult as well because the nose is usually made up of loose sand without much if any vegetation.

A type of extensive parabolic dune that lacks discernible slip faces and has mostly coarse grained sand is known as a Zibar.[4] The term zibar comes from the Arabic word to describe “rolling transverse ridges…with a hard surface.”[5] The dunes are small, have low relief, and can be found in many places across the planet from Wyoming to Saudi Arabia to Australia. Spacing between zibars ranges from 50 to 400 meters and they don’t become more than 10 meters high.[6] The dunes form at about ninety degrees to the prevailing wind which blows away the small, fine-grained sand leaving behind the coarser grained sand to form the crest.[7]

Longitudinal (Seif) dunes

Longitudinal dunes (also called Seif dunes, after the Arabic word for “sword”), elongate parallel to the prevailing wind, possibly caused by a larger dune having its smaller sides blown away. Seif dunes are sharp-crested and are common in the Sahara. They range up to 300 m (980 ft) in height and 300 km (190 mi) in length. In the southern third of the Arabian Peninsula, a vast erg called the Rub’ al Khali or the Empty Quarter, contains seif dunes that stretch for almost 200 km and reach heights of over 300 m.

Seif dunes are thought to develop from barchans if a change of the usual wind direction occurs. The new wind direction will lead to the development of a new wing and the over development of one of the original wings. If the prevailing wind then becomes dominant for a lengthy period of time the dune will revert to its barchan form, with one exaggerated wing. Should the strong wind then return the exaggerated wing will further extend so that eventually it will be supplied with sand when the prevailing wind returns. The wing will continue to grow under both wind conditions, thus producing a seif dune. On a seif dune the slip face develops on the side facing away from the strong wind, while the slip face of a barchan faces the direction of movement. In the sheltered troughs between highly developed seif dunes barchans may be formed because the wind is unidirectional.

A transverse dune is perpendicular to the prevailing wind, probably caused by a steady build-up of sand on an already existing minuscule mound.

Reversing dunes

Occurring wherever winds periodically reverse direction, reversing dunes are varieties of any of the above shapes. These dunes typically have major and minor slipfaces oriented in opposite directions.

All these dune shapes may occur in three forms: simple, compound, and complex. Simple dunes are basic forms with a minimum number of slipfaces that define the geometric type. Compound dunes are large dunes on which smaller dunes of similar type and slipface orientation are superimposed, and complex dunes are combinations of two or more dune types. A crescentic dune with a star dune superimposed on its crest is the most common complex dune. Simple dunes represent a wind regime that has not changed in intensity or direction since the formation of the dune, while compound and complex dunes suggest that the intensity and direction of the wind has changed.

Sub-aqueous dunes

Sub-aqueous (underwater) dunes form on a bed of sand or gravel under the actions of water flow. They are ubiquitous in natural channels such as rivers and estuaries, and also form in engineered canals and pipelines. Dunes move downstream as the upstream slope is eroded and the sediment deposited on the downstream or lee slope in typical bedform construction.[8]

These dunes most often form as a continuous ‘train’ of dunes, showing remarkable similarity in wavelength and height.

Dunes on the bed of a channel significantly increase flow resistance, their presence and growth playing a major part in river flooding.

Lithified dunes

A lithified (consolidated) sand dune is a type of sandstone that is formed when a marine or aeolian sand dune becomes compacted and hardened. Once in this form, water passing through the rock can carry and deposit minerals, which can alter the color of the rock. Cross-bedded layers of stacks of lithified dunes can produce the cross-hatching patterns, such as those seen in the Zion National Park in the western United States.

A slang term that is used in the Southwestern States (of the U.S.A.) for those consolidated and hardened sand dunes is “slickrock”, a name that was introduced by pioneers of the Old West because their steel-rimmed wagon wheels could not gain traction on the rock.

Coastal dunes

Dunes form where **constructive waves** encourage the accumulation of sand, and where prevailing **onshore winds** blow this sand inland. There need to be obstacles—for example, vegetation, pebbles and so on—to trap the moving sand grains. As the sand grains get trapped they start to accumulate, starting dune formation. The wind then starts to affect the mound of sand by eroding sand particles from the windward side and depositing them on the leeward side. Gradually this action causes the dune to "migrate" inland, as it does so it accumulates more and more sand. Dunes provide privacy and shelter from the wind.

Ecological succession on coastal dunes

As a dune forms, plant succession occurs. The conditions on an *embryo dune* are harsh, with salt spray from the sea carried on strong winds. The dune is well drained and often dry, and composed of calcium carbonate from seashells. Rotting seaweed, brought in by storm waves adds nutrients to allow pioneer species to colonize the dune. These pioneer species are marram grass, sea wort grass and other sea grasses in the United Kingdom. These plants are well adapted to the harsh conditions of the foredune typically having deep roots which reach the water table, root nodules that produce nitrogen compounds, and protected stoma, reducing transpiration. Also, the deep roots bind the sand together, and the dune grows into a foredune as more sand is blown over the grasses. The grasses add nitrogen to the soil, meaning other, less hardy plants can then colonize the dunes. Typically these are heather, heaths and gorses. These too are adapted to the low soil water content and have small, prickly leaves which reduce transpiration. Heather adds humus to the soil and is usually replaced by coniferous trees, which can tolerate low soil pH, caused by the accumulation and decomposition of organic matter with nitrate leaching.[9] Coniferous forests and heathland are common climax communities for sand dune systems.

Young dunes are called yellow dunes and dunes which have high humus content are called grey dunes. Leaching occurs on the dunes, washing humus into the slacks, and the slacks may be much more developed than the exposed tops of the dunes. It is usually in the slacks that more rare species are developed and there is a tendency for the dune slacks soil to be waterlogged and where only marsh plants can survive. These plants would include: creeping willow, cotton grass, yellow ins, reeds, and rushes. As for the species, there is a tendency for natterjack toads to breed here.

Desertification

Main article: Desertification

Sand dunes can have a negative impact on humans when they encroach on human habitats. Sand dunes move via a few different means, all of them helped along by wind. One way that dunes can move is by saltation, where sand particles skip along the ground like a bouncing ball. When these skipping particles land, they may knock into other particles and cause them to move as well, in a process known as creep. With slightly stronger winds, particles collide in mid-air, causing sheet flows. In a major dust storm, dunes may move tens of metres through such sheet flows. Also as in the case of snow, sand avalanches, falling down the slipface of the dunes – that face away from the winds – also move the dunes forward.

Sand threatens buildings and crops in Africa, the Middle East, and China. Drenching sand dunes with oil stops their migration, but this approach is quite destructive to the dunes' animal habitats and uses a valuable resource. Sand fences might also slow their movement to a crawl, but geologists are still analyzing results for the optimum fence designs.[*citation needed*] Preventing sand dunes from overwhelming towns, villages, and agricultural areas has become a priority for the United Nations Environment Programme. Planting dunes with vegetation also helps to stabilise them.

Examples

Africa

- Alexandria Coastal Dunefields, in the Eastern Cape, South Africa[10]
- Western Sahara
- Sossusvlei, Namibia[11]



Wind ripples on crescent-shaped sand dunes (Barchans) in Southwest Afghanistan (Sistan).

Asia

- The dunes in the Thar desert in Rajasthan, India
- Tottori Sand Dunes, Tottori Prefecture, Japan
- Rig-e Jenn in the Central Desert of Iran.
- Rig-e Lut in the Southeast of Iran.
- The Ilocos Norte Sand Dunes in the Philippines, particularly Paoay Sand Dunes.
- Mer'eb Dune (also written as Merheb) in United Arab Emirates, used as an arena for motor sports and skiing.
- Gumuk Pasir Parangkusumo near Parangtritis beach in Yogyakarta, Indonesia.
- Mui Ne, Vietnam.
- Wahiba Sands, Oman[12]

Europe

- The Dune of Pyla, not far from Bordeaux, France, is the largest known sand dune in Europe
- The Dunes of Piscinas, in the south west of Sardinia island.
- Sands of Forvie within the Ythan Estuary complex, Aberdeenshire, Scotland.
- Oxwich Dunes, near Swansea, is on the Gower Peninsula in Wales.
- Winterton Dunes – Norfolk, England
- Słowiński National Park, Poland
- Akrotiri Sand Dune, Lemesos, Cyprus
- Råbjerg mile, Northern Jutland, Denmark
- Thy National Park, North Denmark Region, Denmark

North America



Guadalupe-Nipomo Dunes



Oregon Dunes National Recreation Area

- Great Kobuk Sand Dunes, Kobuk Valley National Park, Alaska[13]
- The Athabasca Sand Dunes, the northmost sand dunes in the world, located in the Athabasca Sand Dunes Provincial Park, Saskatchewan.
- The Kelso Dunes in the Mojave Desert of California.
- Great Sand Dunes National Park, Colorado.
- White Sands National Monument, New Mexico.
- Little Sahara Recreation Area, Utah.
- Indiana Dunes National Lakeshore, Indiana, on the south shore of Lake Michigan.[14][15]
- Warren Dunes State Park and Sleeping Bear Dunes National Lakeshore, Michigan, on the east shore of Lake Michigan.
- Grand Sable Dunes, in the Pictured Rocks National Lakeshore, Michigan.
- Samalayuca Dunes, in the state of Chihuahua, Mexico
- Algodones Dunes near Brawley, California.
- Guadalupe-Nipomo Dunes, on the central coast of California.
- Monahans Sandhills State Park near Odessa, Texas.
- Beaver Dunes State Park near Beaver, Oklahoma.
- The Killpecker sand dunes of the Red Desert in southwestern Wyoming.
- Jockey's Ridge State Park – on the Outer Banks, North Carolina.
- Oregon Dunes National Recreation Area near Florence, Oregon, on the Pacific Coast.
- Bruneau Dunes State Park – Owyhee Desert, Idaho
- Hoffmaster State Park – Muskegon, Michigan

South America

- Lençóis Maranhenses National Park in the state of Maranhão, Brazil
- Jericoacoara National Park, in the state of Ceará, Brazil
- Genipabu in Natal, Brazil
- Medanos de Coro National Park near the town of Coro, in Falcón State, Venezuela
- Duna Federico Kirbus in Catamarca Province, Argentina
- Villa Gesell in Buenos Aires Province, Argentina
- Cerro Blanco in Nazca Province, Peru
- Cerro Medanoso in Atacama Region, Chile
- Colún Beach, Valdivian Coastal Reserve in Chile

Oceania

- Simpson Desert sand dunes, Queensland and Northern Territory, Australia
- Fraser Island in Queensland, Australia
- Cronulla sand dunes in New South Wales, Australia
- Stockton Beach in New South Wales, Australia

World's highest dunes

Note: This table is partially based on estimates and incomplete information.

Dune	Height from Base feet/metres	Height from Sea Level feet/metres	Location	Notes
Duna Federico Kirbus	~4,035/1,230	~9,334/2,845	Bolsón de Fiambalá, Fiambalá, Catamarca Province, Argentina	Highest in the world[16]
Cerro Blanco	~3,860/1,176	~6,791/2,080	Nasca Valley, Department of Ica, Peru14°52′05″S 74°50′17″W / 14.868°S 74.838°W	Highest in Peru, second highest in the world
Badain Jaran Dunes	~1,640/500	~6,640/2,020	Badain Jaran Desert, Alashan Plain, Inner Mongolia, Gobi Desert, China	World's tallest stationary dunes and highest in Asia[17]
Rig-e Yalan Dune	~1,542/470	~3,117/950	Lut Desert, Kerman, Iran	Near World the hottest place (Gandom Beryan)
Average Highest Area Dunes	1,410/430?	~6,500/~1,980?	Isaouane-n-Tifernine Sand Sea, Algerian Sahara	Highest in Africa
Big Daddy/Dune 7 (Big Mama?)[18]	1,256/383	~1,870/570	Sossusvlei Dunes, Namib Desert, Namibia / Near Walvis Bay Namib Desert, Namibia	according to the Namibian Ministry of Environment & Tourism the highest dune in the world
Mount Tempest	~920/280	~920/280	Moreton Island, Brisbane, Australia	Highest in Australia
Star Dune	>750/230	~8,950/2,730	Great Sand Dunes National Park and Preserve, Colorado, USA	Highest in North America
Dune of Pyla	~345/105	~699/130	Bay of Arcachon, Aquitaine, France	Highest in Europe
Ming-Sha Dunes	?	5,660/1,725	Dunhuang Oasis, Taklamakan Desert, Gansu, China	
Medanoso Dune	~1805/550	~5446/1,660	Atacama Desert, Chile	Highest in Chile


Sand dune systems

(coastal dunes featuring succession)

- Athabasca Sand Dunes Provincial Park, Alberta and Saskatchewan
- Ashdod Sand Dune, Israel
- Bamburgh Dunes, Northumberland, England
- Bradley Beach, New Jersey
- Circeo National Park, a Mediterranean dune area on the southwest coast of the Lazio region of Italy
- Cronulla sand dunes, NSW, Australia

- Crymlyn Burrows, Wales
- Fraser Island, Queensland Australia, largest sand island in the world
- Kenfig Burrows, Wales
- Margam burrows, Wales
- Murlough Sand Dunes, Newcastle, Co Down, Northern Ireland
- Morfa Harlech sand dunes, Gwynedd, Wales
- Newborough Warren, North Wales
- Oregon Dunes National Recreation Area, near North Bend, Oregon
- Sleeping Bear Dunes National Lakeshore, Michigan
- Studland, Dorsetshire, England
- Thy National Park, North Denmark Region, Denmark
- Winterton, Norfolk, England
- Ynyslas Sand Dunes, Wales

Extraterrestrial dunes



Sand dune on Mars

See also: List of extraterrestrial dune fields

Dunes can likely be found in any environment where there is a substantial atmosphere, winds, and dust to be blown. Dunes are common on Mars and in the equatorial regions of Titan.

Titan's dunes include large expanses with modal lengths of about 20–30 km. The regions are not topographically confined, resembling sand seas. These dunes are interpreted to be longitudinal dunes whose crests are oriented parallel to the dominant wind direction, which generally indicates west-to-east wind flow. The sand is likely composed of hydrocarbon particles, possibly with some water ice mixed in.[19]

Notes

1. ^ dune – Definitions from Dictionary.com
2. ^ "Types of Dunes". Retrieved 8 March 2012.
3. ^ "Types of Dunes". Retrieved 23 May 2014.
4. ^ Goudie, Ron Cooke; Andrew Warren; Andrew (1996).*Desert geomorphology* (2. impr. ed.). London: UCL Press. pp. 395–396. ISBN 1-85728-017-2.
5. ^ Goudie, Ron Cooke; Andrew Warren; Andrew (1996).*Desert geomorphology* (2. impr. ed.). London: UCL Press. p. 395. ISBN 1-85728-017-2.
6. ^ "USGS Landform Glossary". *United States Geological Survey*. Retrieved 3 October 2013.
7. ^ Nielsen, Ole. "Zibar Dunes". *My Opera*. Retrieved 3 October 2013.
8. ^ Prothero, D. R. and Schwab, F., 1996, Sedimentary Geology, pg. 45-49, ISBN 0-7167-2726-9
9. ^ Miles, J. (1985). "The pedogenic effects of different species and vegetation types and the implications of succession".*European Journal of Soil Science* **36** (4): 571. doi:10.1111/j.1365-2389.1985.tb00359.x.
10. ^ "Alexandria Coastal Dunefields". UNESCO World Heritage. Retrieved 11 January 2010.
11. ^ Sossusvlei
12. ^ Wahiba Sands
13. ^ Mann, D.H.; Heiser P.A.; Finney B.P (2002). "Holocene history of the Great Kobuk Sand Dunes, Northwestern Alaska".*Quaternary Science Reviews* (Elsevier) **21** (4): 709–731. doi:10.1016/S0277-3791(01)00120-2.
14. ^ Smith, S. & Mark, S. (2006). Alice Gray, Dorothy Buell, and Naomi Svihla: Preservationists of Ogden Dunes. The South Shore Journal, 1.http://www.southshorejournal.org/index.php/issues/volume-1-2006/78-journals/vol-1-2006/117-alice-gray-dorothy-buell-and-naomi-svihla-preservationists-of-ogden-dunes

15. ^ Smith, S. & Mark, S. (2009). The Historical Roots of the Nature Conservancy in the Northwest Indiana/Chicagoland Region: From Science to Preservation. The South Shore Journal, 3. http://www.southshorejournal.org/index.php/issues/volume-3-2009/83-journals/vol-3-2009/75-the-historical-roots-of-the-nature-conservancy-in-the-northwest-indianachicagoland-region-from-science-to-preservation
16. ^ Sandboard Magazine
17. ^ Mystery of world's tallest sand dunes solved – 24 November 2004 – New Scientist
18. ^ Big Mama highest dune
19. ^ Peeking Through the Haze: Titan's Surface, part II – The Planetary Society Blog | The Planetary Society

References

- Bagnold, Ralph ([1941] 2005). *The Physics of Blown Sand and Desert Dunes*. Dover Publications. ISBN 978-0-486-43931-0.
- Anthony J. Parsons, A. D. Abrahams, ed. (2009). *Geomorphology of Desert Environments*. Springer. ISBN 978-1-4020-5718-2.
- Pye, Kenneth; Tsoar, Haim (2009). *Aeolian Sand and Sand Dunes*. Springer. ISBN 978-3-540-85909-3.
- "Nouakchott, Mauritania". NASA Earth Observatory. Retrieved 2006-04-28.
- V. Badescu, R. B. Cathcart and A. A. Bolonkin, "Sand Dune Fixation: a solar-powered Sahara seawater pipeline macroproject", in: *Land Degradation & Development*; 19, (2008): doi:10.1002/ldr.864.
- "Types of Dunes". USGS. Retrieved 2010-10-06.
- "Summary: Dunes, Parabolic". Desert Processes Working Group; Knowledge Sciences, Inc. Retrieved 2010-10-06.
- "Fighting wind erosion. One aspect of the combat against desertification". Les dossiers thématiques du CSFD. Retrieved 2011-01-04.

External links

- Coastal Sand Dunes
- Dune pattern identification, U.S. Army
- The Bibliography of Aeolian Research
- Gold Coast Dune Management Policy
- USGS page on dune types
- Great Sand Dunes National Park

CC licensed content, Shared previously

- **Provided by**: Wikipedia. **Located at**: http://en.Wikipedia.org/wiki/Dune. **License**: *CC BY-SA: Attribution-ShareAlike*

This page titled 19.3: Dunes is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

Welcome to the Geosciences Library. This Living Library is a principal hub of the LibreTexts project, which is a multi-institutional collaborative venture to develop the next generation of open-access texts to improve postsecondary education at all levels of higher learning. The LibreTexts approach is highly collaborative where an Open Access textbook environment is under constant revision by students, faculty, and outside experts to supplant conventional paper-based books.

CHAPTER OVERVIEW

20: Glaciers

20.1: Glacial Erosion and Deposition
20.2: Glaciers

This page titled 20: Glaciers is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

20.1: Glacial Erosion and Deposition

Lesson Objectives

- Discuss the different erosional features formed by alpine glaciers.
- Describe the processes by which glaciers change the underlying rocks.
- Discuss the particles deposited by glaciers as they advance and recede.
- Describe the landforms created by glacial deposits.

Vocabulary

- alpine (valley) glacier
- continental glacier
- end moraine
- glacial erratic
- glacial striations
- glacial till
- glaciers
- ground moraine
- hanging valley
- lateral moraine
- medial moraine
- moraine
- plucking
- terminal moraine
- varve


Introduction

Glaciers cover about 10% of the land surface near Earth's poles and they are also found in high mountains. During the Ice Ages, glaciers covered as much as 30% of Earth. Around 600 to 800 million years ago, geologists think that almost all of the Earth was covered in snow and ice. Scientists use the evidence of erosion and deposition left by glaciers to do a kind of detective work to figure out where the ice once was.

Formation and Movement of Glaciers

Glaciers are solid ice that move extremely slowly along the land surface (**Figure** below). Glacial ice erodes and shapes the underlying rocks. Glaciers also deposit sediments in characteristic landforms. The two types of glaciers are:

- **Continental glaciers** are large ice sheets that cover relatively flat ground. These glaciers flow outward from where the greatest amount of snow and ice accumulate.
- **Alpine** or **valley glaciers** flow downhill through mountains along existing valleys.



A satellite image of glaciers in the Himalaya with some features labeled.


Glacial Erosion

Glaciers erode the underlying rock by abrasion and **plucking.** Glacial meltwater seeps into cracks of the underlying rock, the water freezes and pushes pieces of rock outward. The rock is then plucked out and carried away by the flowing ice of the moving glacier (**Figure** below). With the weight of the ice over them, these rocks can scratch deeply into the underlying bedrock making long, parallel grooves in the bedrock, called **glacial striations.**



Glacial striations point the direction a glacier has gone.

Mountain glaciers leave behind unique erosional features. When a glacier cuts through a 'V' shaped river valley, the glacier pucks rocks from the sides and bottom. This widens the valley and steepens the walls, making a 'U' shaped valley (**Figure** below).



A U shaped valley in Glacier National Park.

Smaller tributary glaciers, like tributary streams, flow into the main glacier in their own shallower 'U' shaped valleys. A **hanging valley**forms where the main glacier cuts off a tributary glacier and creates a cliff. Streams plunge over the cliff to create waterfalls

(**Figure** below).




Yosemite Valley is known for waterfalls that plunge from hanging valleys.

Up high on a mountain, where a glacier originates, rocks are pulled away from valley walls. Some of the resulting erosional features are shown: **Figure** below and **Figure** below.



(a) A bowl-shaped cirque in Glacier National Park was carved by glaciers. (b) A high altitude lake, called a tarn, forms from meltwater trapped in the cirque. (c) Several cirques from glaciers flowing in different directions from a mountain peak, leave behind a sharp sided horn, like the Matterhorn in Switzerland. (d) When glaciers move down opposite sides of a mountain, a sharp edged ridge, called an arête, forms between them.



A roche moutonée forms where a glacier smooths the uphill side of the bedrock and plucks away rock from the downslope side.

Depositional Features of Glaciers

As glaciers flow, mechanical weathering loosens rock on the valley walls, which falls as debris on the glacier. Glaciers can carry rock of any size, from giant boulders to silt (**Figure** below). These rocks can be carried for many kilometers for many years. These rocks with a different rock type or origin from the surrounding bedrock are **glacial erratics**. Melting glaciers deposit all the big and small bits of rocky material they are carrying in a pile. These unsorted deposits of rock are called **glacial till**.



A large boulder dropped by a glacier is a glacial erratic.

Glacial till is found in different types of deposits. Linear rock deposits are called **moraines.** Geologists study moraines to figure out how far glaciers extended and how long it took them to melt away. Moraines are named by their location relative to the glacier:

- **Lateral moraines** form at the edges of the glacier as material drops onto the glacier from erosion of the valley walls.
- **Medial moraines** form where the lateral moraines of two tributary glaciers join together in the middle of a larger glacier (**Figure** below).




The long, dark lines are medial and lateral moraines.

- Sediment from underneath the glacier becomes a **ground moraine** after the glacier melts. Ground moraine contributes to the fertile transported soils in many regions.
- **Terminal moraines** are long ridges of till left at the furthest point the glacier reached.
- **End moraines** are deposited where the glacier stopped for a long enough period to create a rocky ridge as it retreated. Long Island in New York is formed by two end moraines.



(a) An esker is a winding ridge of sand and gravel deposited under a glacier by a stream of meltwater. (b) A drumlin is an asymmetrical hill made of sediments that points in the direction the ice moved. Usually drumlins are found in groups called drumlin fields.

While glaciers dump unsorted sediments, glacial meltwater can sort and re-transport the sediments (**Figure** below).



(a) A sorted deposit of sand and smaller particles is stratified drift. A broad area of stratified drift from meltwater over broad region is an outwash plain. (b) Kettle lakes form as blocks of ice in glacial till melt.

- Try to pick out some of the glacial features seen in this Glacier National Park video:http://www.visitmt.com/national_parks/glacier/video_series/part_3.htm.

Several types of stratified deposits form in glacial regions but are not formed directly by the ice. **Varves** form where lakes are covered by ice in the winter. Dark, fine-grained clays sink to the bottom in winter but melting ice in spring brings running water that deposits lighter colored sands. Each alternating dark/light layer represents one year of deposits. If during a year, a glacier accumulates more ice than melts away, the glacier advances downhill. If a glacier melts more than it accumulates over a year, it is retreating (**Figure** below).



Grinnell Glacier in Glacier National Park has been retreating over the past 70 years.

Lesson Summary

- The movement of ice in the form of glaciers has transformed our mountainous land surfaces with its tremendous power of erosion.
- U-shaped valleys, hanging valleys, cirques, horns, and aretes are features sculpted by ice.
- The eroded material is later deposited as large glacial erratics, in moraines, stratified drift, outwash plains, and drumlins.
- Varves are a very useful yearly deposit that forms in glacial lakes.

Review Questions

1. How much of the Earth's land surface is covered by glaciers today? Where are they found?
2. What are the two types of glaciers and how are they different from each other?
3. What is the shape of a valley that has been eroded by rivers? How does a glacier change that shape and what does it become?
4. What two different features form as smaller side glaciers join the central main glacier?
5. How do glaciers erode the surrounding rocks?
6. Name the erosional features that are formed by glaciers high in the mountains and describe how they form.
7. Describe the different types of moraines formed by glaciers.
8. Describe the difference between glacial till and stratified drift. Give an example of how each type of deposit forms.
9. Name and describe the two asymmetrical hill shaped landforms created by glaciers.

Further Reading / Supplemental Links

- Glacial landforms illustrated: http://www.uwsp.edu/geo/faculty/lemke/alpine_glacial_glossary/glossary.html

Points to Consider

- What features would you look for to determine if glaciers had ever been present?
- If glaciers had never formed, how would soil in Midwestern North America be different?
- Can the process of erosion produce landforms that are beautiful?

This page titled 20.1: Glacial Erosion and Deposition is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

20.2: Glaciers

A **glacier** (US /ˈɡleɪʃər/ or UK /ˈɡlæsiə/) is a persistent body of dense ice that is constantly moving under its own weight; it forms where the accumulation of snow exceeds its ablation (melting and sublimation) over many years, often centuries. Glaciers slowly deform and flow due to stresses induced by their weight, creating crevasses, seracs, and other distinguishing features. They also abrade rock and debris from their substrate to create landforms such as cirques and moraines. Glaciers form only on land and are distinct from the much thinner sea ice and lake ice that form on the surface of bodies of water.



The Baltoro Glacier in the Karakoram, Baltistan, Northern Pakistan. At 62 kilometres (39 mi) in length, it is one of the longest alpine glaciers on earth.



Ice calving from the terminus of the Perito Moreno Glacier in western Patagonia, Argentina



The Aletsch Glacier, the largest glacier of the Alps, in Switzerland



The Quelccaya Ice Cap is the largest glaciated area in the tropics, in Peru

On Earth, 99% of glacial ice is contained within vast ice sheets in the polar regions, but glaciers may be found in mountain ranges on every continent except Australia, and on a few high-latitude oceanic islands. Between 35°N and 35°S, glaciers occur only in the Himalayas, Andes, Rocky Mountains, a few high mountains in East Africa, Mexico, New Guinea and on Zard Kuh in Iran. [1]

Glacial ice is the largest reservoir of freshwater on Earth.[2] Many glaciers from temperate, alpine and seasonal polar climates store water as ice during the colder seasons and release it later in the form of meltwater as warmer summer temperatures cause the glacier to melt, creating a water source that is especially important for plants, animals and human uses when other sources may be scant. Within high altitude and Antarctic environments, the seasonal temperature difference is often not sufficient to release meltwater.

Because glacial mass is affected by long-term climate changes, e.g., precipitation, mean temperature, and cloud cover, glacial mass changes are considered among the most sensitive indicators of climate change and are a major source of variations in sea level.

Etymology and related terms

The word *glacier* comes from French. It is derived from the Vulgar Latin *glacia* and ultimately from Latin *glacies* meaning “ice”.[3] The processes and features caused by glaciers and related to them are referred to as **glacial**. The process of glacier establishment, growth and flow is called **glaciation**. The corresponding area of study is called glaciology. Glaciers are important components of the global cryosphere.

Types

Main article: Glacier morphology



Mouth of the Schlatenkees Glacier near Innergschlöß, Austria

Glaciers are categorized by their morphology, thermal characteristics, and behavior. *Alpine glaciers*, also known as *mountain glaciers*or *cirque glaciers*, form on the crests and slopes of mountains. An alpine glacier that fills a valley is sometimes called a *valley glacier*. A large body of glacial ice astride a mountain, mountain range, or volcano is termed an ice cap or ice field.[4] Ice caps have an area less than 50,000 km² (20,000 mile²) by definition.

Glacial bodies larger than 50,000 km² are called ice sheets or continental glaciers.[5] Several kilometers deep, they obscure the underlying topography. Only *nunataks* protrude from their surfaces. The only extant ice sheets are the two that cover most of Antarctica and Greenland. They contain vast quantities of fresh water, enough that if both melted, global sea levels would rise by over 70 meters.[6]Portions of an ice sheet or cap that extend into water are called *ice shelves;* they tend to be thin with limited slopes and reduced velocities.[7] Narrow, fast-moving sections of an ice sheet are called *ice streams.*[8][9] In Antarctica, many ice streams drain into large ice shelves. Some drain directly into the sea, often with an ice tongue, like Mertz Glacier.



Sightseeing boat in front of a tidewater glacier, Kenai Fjords National Park, Alaska

Tidewater glaciers are glaciers that terminate in the sea, including most glaciers flowing from Greenland, Antarctica, Baffin and Ellesmere Islands in Canada, Southeast Alaska, and the Northern and Southern Patagonian Ice Fields. As the ice reaches the sea, pieces break off, or *calve*, forming icebergs. Most tidewater glaciers calve above sea level, which often results in a tremendous impact as the iceberg strikes the water. Tidewater glaciers undergo centuries-long cycles of advance and retreat that are much less affected by the climate change than those of other glaciers.

Thermally, a *temperate glacier* is at melting point throughout the year, from its surface to its base. The ice of a *polar glacier* is always below freezing point from the surface to its base, although the surface snowpack may experience seasonal melting. A *sub-polar glacier*includes both temperate and polar ice, depending on depth beneath the surface and position along the length of the glacier. In a similar way, the thermal regime of a glacier is often described by the temperature at its base alone. A *cold-based* glacier is below freezing at the ice-ground interface, and is thus frozen to the underlying substrate. A *warm-based* glacier is above or at freezing at the interface, and is able to slide at this contact.[10] This contrast is thought to a large extent to govern the ability of a glacier to effectively erode its bed, as sliding ice promotes plucking at rock from the surface below.[11] Glaciers which are partly cold-based and partly warm-based are known as *polythermal*.[10]

Formation



Gorner Glacier in Switzerland

Glaciers form where the accumulation of snow and ice exceeds ablation. The area in which a glacier forms is called a cirque (corrie or cwm) – a typically armchair-shaped geological feature (such as a depression between mountains enclosed by arêtes) – which collects and compresses through gravity the snow which falls into it. This snow collects and is compacted by the weight of the snow falling above it forming névé. Further crushing of the individual snowflakes and squeezing the air from the snow turns it into extremely dense 'glacial ice'. This glacial ice will fill the cirque until it 'overflows' through a geological weakness or vacancy, such as the gap between two mountains. When the mass of snow and ice is sufficiently thick, it begins to move due to a combination of surface slope, gravity and pressure. On steeper slopes, this can occur with as little as 15 m (50 ft) of snow-ice.

In temperate glaciers, snow repeatedly freezes and thaws, changing into granular ice called firn. Under the pressure of the layers of ice and snow above it, this granular ice fuses into denser and denser firn. Over a period of years, layers of firn undergo further compaction and become glacial ice. Glacier ice is slightly less dense than ice formed from frozen water because it contains tiny trapped air bubbles.

Glacial ice has a distinctive blue tint because it absorbs some red light due to an overtone of the infrared OH stretching mode of the water molecule. Liquid water is blue for the same reason. The blue of glacier ice is sometimes misattributed to Rayleigh scattering due to bubbles in the ice.[12]



A glacier cave located on the Perito Moreno Glacier in Argentina.

Structure

A glacier originates at a location called its *glacier head* and terminates at its *glacier foot*, or terminus.

Glaciers are broken into zones based on surface snowpack and melt conditions.[13] The *ablation zone* is the region where there is a net loss in glacier mass. The *equilibrium line* separates the ablation zone and the accumulation zone; it is the altitude where the amount of new snow gained by accumulation is equal to the amount of ice lost through ablation. The upper part of a glacier, where accumulation exceeds ablation, is called the *accumulation zone*. In general, the accumulation zone accounts for 60–70% of the glacier's surface area, more if the glacier calves icebergs. Ice in the accumulation zone is deep enough to exert a downward force that erodes underlying rock. After a glacier melts, it often leaves behind a bowl- or amphitheater-shaped depression that ranges in size from large basins like the Great Lakes to smaller mountain depressions known as cirques.

The accumulation zone can be subdivided based on its melt conditions.

1. The *dry snow zone* is a region where no melt occurs, even in the summer, and the snowpack remains dry.
2. The *percolation zone* is an area with some surface melt, causing meltwater to percolate into the snowpack. This zone is often marked by refrozen ice lenses, glands, and layers. The snowpack also never reaches melting point.
3. Near the equilibrium line on some glaciers, a *superimposed ice zone* develops. This zone is where meltwater refreezes as a cold layer in the glacier, forming a continuous mass of ice.
4. The *wet snow zone* is the region where all of the snow deposited since the end of the previous summer has been raised to 0 °C.

The "health" of a glacier is usually assessed by determining the glacier mass balance or observing terminus behavior. Healthy glaciers have large accumulation zones, more than 60% of their area snowcovered at the end of the melt season, and a terminus with vigorous flow.

Following the Little Ice Age's end around 1850, glaciers around the Earth have retreated substantially. A slight cooling led to the advance of many alpine glaciers between 1950–1985, but since 1985 glacier retreat and mass loss has become larger and increasingly ubiquitous.[14][15][16]

Motion



Shear or herring-bone crevasses on Emmons Glacier (Mount Rainier); such crevasses often form near the edge of a glacier where interactions with underlying or marginal rock impede flow. In this case, the impediment appears to be some distance from the near margin of the glacier.

Main article: Ice sheet dynamics

Glaciers move, or flow, downhill due to gravity and the internal deformation of ice.[17] Ice behaves like a brittle solid until its thickness exceeds about 50 m (160 ft). The pressure on ice deeper than 50 m causes plastic flow. At the molecular level, ice consists of stacked layers of molecules with relatively weak bonds between layers. When the stress on the layer above exceeds the inter-layer binding strength, it moves faster than the layer below.[18]

Glaciers also move through basal sliding. In this process, a glacier slides over the terrain on which it sits, lubricated by the presence of liquid water. The water is created from ice that melts under high pressure from frictional heating. Basal sliding is dominant in temperate, or warm-based glaciers.

Fracture zone and cracks



Ice cracks in the Titlis Glacier

The top 50 metres (160 ft) of a glacier are rigid because they are under low pressure. This upper section is known as the *fracture zone*; it mostly moves as a single unit over the plastically flowing lower section. When a glacier moves through irregular terrain, cracks called crevasses develop in the fracture zone. Crevasses form due to differences in glacier velocity. If two rigid sections of a glacier move at different speeds and directions, shear forces cause them to break apart, opening a crevasse. Crevasses are seldom more than 150 feet (46 m) deep but in some cases can be 1,000 feet (300 m) or even deeper. Beneath this point, the plasticity of the ice is too great for cracks to form. Intersecting crevasses can create isolated peaks in the ice, called seracs.

Crevasses can form in several different ways. Transverse crevasses are transverse to flow and form where steeper slopes cause a glacier to accelerate. Longitudinal crevasses form semi-parallel to flow where a glacier expands laterally. Marginal crevasses form from the edge of the glacier, due to the reduction in speed caused by friction of the valley walls. Marginal crevasses are usually largely transverse to flow. Moving glacier ice can sometimes separate from stagnant ice above, forming a *bergschrund*. Bergschrunds resemble crevasses but are singular features at a glacier's margins.

Crevasses make travel over glaciers hazardous, especially when they are hidden by fragile snow bridges.



Crossing a crevasse on the Easton Glacier, Mount Baker, in the North Cascades, United States

Below the equilibrium line, glacial meltwater is concentrated in stream channels. Meltwater can pool in proglacial lakes on top of a glacier or descend into the depths of a glacier via moulins. Streams within or beneath a glacier flow in englacial or sub-glacial tunnels. These tunnels sometimes reemerge at the glacier's surface.[19]

Speed

The speed of glacial displacement is partly determined by friction. Friction makes the ice at the bottom of the glacier move more slowly than ice at the top. In alpine glaciers, friction is also generated at the valley's side walls, which slows the edges relative to the center.

Mean speeds vary greatly, but is typically around 1 meter per day.[20] There may be no motion in stagnant areas; for example, in parts of Alaska, trees can establish themselves on surface sediment deposits. In other cases, glaciers can move as fast as 20–30 m per day, such as in Greenland's Jakobshavn Isbræ (Greenlandic: *Sermeq Kujalleq*). Velocity increases with increasing slope, increasing thickness, increasing snowfall, increasing longitudinal confinement, increasing basal temperature, increasing meltwater production and reduced bed hardness.

A few glaciers have periods of very rapid advancement called surges. These glaciers exhibit normal movement until suddenly they accelerate, then return to their previous state. During these surges, the glacier may reach velocities far greater than normal speed.[21] These surges may be caused by failure of the underlying bedrock, the pooling of meltwater at the base of the glacier[22] — perhaps delivered from a supraglacial lake — or the simple accumulation of mass beyond a critical "tipping point".[23]


In glaciated areas where the glacier moves faster than one km per year, glacial earthquakes occur. These are large scale tremblors that have seismic magnitudes as high as 6.1.[24][25] The number of glacial earthquakes in Greenland peaks every year in July, August and September and is increasing over time. In a study using data from January 1993 through October 2005, more events were detected every year since 2002, and twice as many events were recorded in 2005 as there were in any other year. This increase in the numbers of glacial earthquakes in Greenland may be a response to global warming.[24][25]

Ogives

Ogives are alternating wave crests and valleys that appear as dark and light bands of ice on glacier surfaces. They are linked to seasonal motion of glaciers; the width of one dark and one light band generally equals the annual movement of the glacier. Ogives are formed when ice from an icefall is severely broken up, increasing ablation surface area during summer. This creates a swale and space for snow accumulation in the winter, which in turn creates a ridge.[26] Sometimes ogives consist only of undulations or color bands and are described as wave ogives or band ogives.[27]

Geography

For more details on this topic, see List of glaciers, and Retreat of glaciers since 1850.



Black ice glacier near Aconcagua, Argentina

Glaciers are present on every continent and approximately fifty countries, excluding those (Australia, South Africa) that have glaciers only on distant subantarctic island territories. Extensive glaciers are found in Antarctica, Chile, Canada, Alaska, Greenland and Iceland. Mountain glaciers are widespread, especially in the Andes, the Himalayas, the Rocky Mountains, the Caucasus, and the Alps. Mainland Australia currently contains no glaciers, although a small glacier on Mount Kosciuszko was present in the last glacial period.[28] In New Guinea, small, rapidly diminishing, glaciers are located on its highest summit massif of Puncak Jaya.[29] Africa has glaciers on Mount Kilimanjaro in Tanzania, on Mount Kenya and in the Rwenzori Mountains. Oceanic islands with glaciers occur on Iceland, Svalbard, New Zealand, Jan Mayen and the subantarctic islands of Marion, Heard, Grande Terre (Kerguelen) and Bouvet. During glacial periods of the Quaternary, Taiwan, Hawaii on Mauna Kea[30] and Tenerife also had large alpine glaciers, while the Faroe and Crozet Islands[31] were completely glaciated.

The permanent snow cover necessary for glacier formation is affected by factors such as the degree of slope on the land, amount of snowfall and the winds. Glaciers can be found in all latitudes except from 20° to 27° north and south of the equator where the presence of the descending limb of the Hadley circulation lowers precipitation so much that with high insolation snow lines reach above 6,500 metres (21,330 ft). Between 19˚N and 19˚S, however, precipitation is higher and the mountains above 5,000 metres (16,400 ft) usually have permanent snow.

Even at high latitudes, glacier formation is not inevitable. Areas of the Arctic, such as Banks Island, and the McMurdo Dry Valleys in Antarctica are considered polar deserts where glaciers cannot form because they receive little snowfall despite the bitter cold. Cold air, unlike warm air, is unable to transport much water vapor. Even during glacial periods of the Quaternary, Manchuria, lowland Siberia,[32]and central and northern Alaska,[33] though extraordinarily cold, had such light snowfall that glaciers could not form.[34][35]

In addition to the dry, unglaciated polar regions, some mountains and volcanoes in Bolivia, Chile and Argentina are high (4,500 metres (14,800 ft) – 6,900 m (22,600 ft)) and cold, but the relative lack of precipitation prevents snow from accumulating into glaciers. This is because these peaks are located near or in the hyperarid Atacama Desert.

Glacial geology




Diagram of glacial plucking and abrasion



Glacially plucked granitic bedrock near Mariehamn, Åland Islands

Glaciers erode terrain through two principal processes: **abrasion** and **plucking.**

As glaciers flow over bedrock, they soften and lift blocks of rock into the ice. This process, called plucking, is caused by subglacial water that penetrates fractures in the bedrock and subsequently freezes and expands. This expansion causes the ice to act as a lever that loosens the rock by lifting it. Thus, sediments of all sizes become part of the glacier's load. If a retreating glacier gains enough debris, it may become a rock glacier, like the Timpanogos Glacier in Utah.

Abrasion occurs when the ice and its load of rock fragments slide over bedrock and function as sandpaper, smoothing and polishing the bedrock below. The pulverized rock this process produces is called rock flour and is made up of rock grains between 0.002 and 0.00625 mm in size. Abrasion leads to steeper valley walls and mountain slopes in alpine settings, which can cause avalanches and rock slides. These add even more material to the glacier.

Glacial abrasion is commonly characterized by glacial striations. Glaciers produce these when they contain large boulders that carve long scratches in the bedrock. By mapping the direction of the striations, researchers can determine the direction of the glacier's movement. Similar to striations are chatter marks, lines of crescent-shape depressions in the rock underlying a glacier. They are formed by abrasion when boulders in the glacier are repeatedly caught and released as they are dragged along the bedrock.

The rate of glacier erosion is variable. Six factors control erosion rate:

- Velocity of glacial movement
- Thickness of the ice
- Shape, abundance and hardness of rock fragments contained in the ice at the bottom of the glacier
- Relative ease of erosion of the surface under the glacier
- Thermal conditions at the glacier base
- Permeability and water pressure at the glacier base

Material that becomes incorporated in a glacier is typically carried as far as the zone of ablation before being deposited. Glacial deposits are of two distinct types:

- *Glacial till*: material directly deposited from glacial ice. Till includes a mixture of undifferentiated material ranging from clay size to boulders, the usual composition of a moraine.
- *Fluvial and outwash sediments*: sediments deposited by water. These deposits are stratified by size.

Larger pieces of rock that are encrusted in till or deposited on the surface are called "glacial erratics". They range in size from pebbles to boulders, but as they are often moved great distances, they may be drastically different from the material upon which they are found. Patterns of glacial erratics hint at past glacial motions.

Moraines




Glacial moraines above Lake Louise, Alberta, Canada

Glacial moraines are formed by the deposition of material from a glacier and are exposed after the glacier has retreated. They usually appear as linear mounds of till, a non-sorted mixture of rock, gravel and boulders within a matrix of a fine powdery material. Terminal or end moraines are formed at the foot or terminal end of a glacier. Lateral moraines are formed on the sides of the glacier. Medial moraines are formed when two different glaciers merge and the lateral moraines of each coalesce to form a moraine in the middle of the combined glacier. Less apparent are ground moraines, also called *glacial drift*, which often blankets the surface underneath the glacier downslope from the equilibrium line.

The term *moraine* is of French origin. It was coined by peasants to describe alluvial embankments and rims found near the margins of glaciers in the French Alps. In modern geology, the term is used more broadly, and is applied to a series of formations, all of which are composed of till. Moraines can also create moraine dammed lakes.

Drumlins




A drumlin field forms after a glacier has modified the landscape. The teardrop-shaped formations denote the direction of the ice flow.

Drumlins are asymmetrical, canoe shaped hills made mainly of till. Their heights vary from 15 to 50 meters and they can reach a kilometer in length. The tilted side of the hill faces the direction from which the ice advanced (*stoss*), while the longer slope follows the ice's direction of movement (*lee*).

Drumlins are found in groups called *drumlin fields* or *drumlin camps*. One of these fields is found east of Rochester, New York; it is estimated to contain about 10,000 drumlins.

Although the process that forms drumlins is not fully understood, their shape implies that they are products of the plastic deformation zone of ancient glaciers. It is believed that many drumlins were formed when glaciers advanced over and altered the deposits of earlier glaciers.

Glacial valleys, cirques, arêtes, and pyramidal peaks



Features of a glacial landscape

Before glaciation, mountain valleys have a characteristic "V" shape, produced by eroding water. During glaciation, these valleys are widened, deepened, and smoothed, forming a "U"-shaped glacial valley. The erosion that creates glacial valleys eliminates the spurs of earth that extend across mountain valleys, creating triangular cliffs called truncated spurs. Within glacial valleys, depressions created by plucking and abrasion can be filled by lakes, called paternoster lakes. If a glacial valley runs into a large body of water, it forms a fjord.

Many glaciers deepen their valleys more than their smaller tributaries. Therefore, when glaciers recede, the valleys of the tributary glaciers remain above the main glacier's depression and are called hanging valleys.

At the start of a classic valley glacier is a bowl-shaped cirque, which has escarped walls on three sides but is open on the side that descends into the valley. Cirques are where ice begins to accumulate in a glacier. Two glacial cirques may form back to back and erode their backwalls until only a narrow ridge, called an arête is left. This structure may result in a mountain pass. If multiple cirques encircle a single mountain, they create pointed pyramidal peaks; particularly steep examples are called horns.

Roche moutonnée


Some rock formations in the path of a glacier are sculpted into small hills called *roche moutonnée*, or "sheepback" rock. Roche moutonnée are elongated, rounded, and asymmetrical bedrock knobs can be produced by glacier erosion. They range in length from less than a meter to several hundred meters long.[36] Roche moutonnée have a gentle slope on their up-glacier sides and a steep to vertical face on their down-glacier sides. The glacier abrades the smooth slope on the upstream side as it flows along, but tears loose and carries away rock from the downstream side via plucking.

Alluvial stratification

As the water that rises from the ablation zone moves away from the glacier, it carries fine eroded sediments with it. As the speed of the water decreases, so does its capacity to carry objects in suspension. The water thus gradually deposits the sediment as it runs, creating an alluvial plain. When this phenomenon occurs in a valley, it is called a *valley train*. When the deposition is in an estuary, the sediments are known as bay mud.

Outwash plains and valley trains are usually accompanied by basins known as "kettles". These are small lakes formed when large ice blocks that are trapped in alluvium melt and produce water-filled depressions. Kettle diameters range from 5 m to 13 km, with depths of up to 45 meters. Most are circular in shape because the blocks of ice that formed them were rounded as they melted.[37]

Glacial deposits



Landscape produced by a receding glacier

When a glacier's size shrinks below a critical point, its flow stops and it becomes stationary. Meanwhile, meltwater within and beneath the ice leaves stratified alluvial deposits. These deposits, in the forms of columns, terraces and clusters, remain after the glacier melts and are known as "glacial deposits".

Glacial deposits that take the shape of hills or mounds are called *kames*. Some kames form when meltwater deposits sediments through openings in the interior of the ice. Others are produced by fans or deltas created by meltwater. When the glacial ice occupies a valley, it can form terraces or kames along the sides of the valley.


Long, sinuous glacial deposits are called *eskers*. Eskers are composed of sand and gravel that was deposited by meltwater streams that flowed through ice tunnels within or beneath a glacier. They remain after the ice melts, with heights exceeding 100 meters and lengths of as long as 100 km.

Loess deposits

Very fine glacial sediments or rock flour is often picked up by wind blowing over the bare surface and may be deposited great distances from the original fluvial deposition site. These eolian loess deposits may be very deep, even hundreds of meters, as in areas of China and the Midwestern United States of America. Katabatic winds can be important in this process.

Isostatic rebound

Main article: Isostatic rebound




Isostatic pressure by a glacier on the Earth's crust

Large masses, such as ice sheets or glaciers, can depress the crust of the Earth into the mantle. The depression usually totals a third of the ice sheet or glacier's thickness. After the ice sheet or glacier melts, the mantle begins to flow back to its original position, pushing the crust back up. This post-glacial rebound, which proceeds very slowly after the melting of the ice sheet or glacier, is currently occurring in measurable amounts in Scandinavia and the Great Lakes region of North America.

A geomorphological feature created by the same process on a smaller scale is known as *dilation-faulting*. It occurs where previously compressed rock is allowed to return to its original shape more rapidly than can be maintained without faulting. This leads to an effect similar to what would be seen if the rock were hit by a large hammer. Dilation faulting can be observed in recently de-glaciated parts of Iceland and Cumbria.

On Mars



Northern polar ice cap on Mars

Main article: Glaciers on Mars

The polar ice caps of Mars show geologic evidence of glacial deposits. The south polar cap is especially comparable to glaciers on Earth.[38] Topographical features and computer models indicate the existence of more glaciers in Mars' past.[39]

At mid-latitudes, between 35° and 65° north or south, Martian glaciers are affected by the thin Martian atmosphere. Because of the low atmospheric pressure, ablation near the surface is solely due to sublimation, not melting. As on Earth, many glaciers are covered with a layer of rocks which insulates the ice. A radar instrument on board the Mars Reconnaissance Orbiter found ice under a thin layer of rocks in formations called Lobate Debris Aprons (LDAs).[40][41][42][43][44]

Notes

1. ^ Post, Austin; LaChapelle, Edward R (2000). *Glacier ice*. Seattle, Washington: University of Washington Press. ISBN 0-295-97910-0.
2. ^ Brown, Molly Elizabeth; Ouyang, Hua; Habib, Shahid; Shrestha, Basanta; Shrestha, Mandira; Panday, Prajjwal; Tzortziou, Maria; Policelli, Frederick; Artan, Guleid; Giriraj, Amarnath; Bajracharya, Sagar R.; Racoviteanu, Adina. "HIMALA: Climate Impacts on Glaciers, Snow, and Hydrology in the Himalayan Region". *Mountain Research and Development*. International Mountain Society. Retrieved 16 September 2011.
3. ^ Simpson, D.P. (1979). *Cassell's Latin Dictionary* (5 ed.). London: Cassell Ltd. p. 883. ISBN 0-304-52257-0.
4. ^ "Retreat of Alaskan glacier Juneau icefield". Nichols.edu. Retrieved 2009-01-05.
5. ^ "American Meteorological Society, Glossary of Meteorology". Amsglossary.allenpress.com. Retrieved 2013-01-04.
6. ^ "Sea Level and Climate". *USGS FS 002-00*. USGS. 2000-01-31. Retrieved 2009-01-05.
7. ^ * National Snow and Ice Data Center. "Types of Glacier".
8. ^ Bindschadler, R.A. and T.A. Scambos. Satellite-image-derived velocity field of an Antarctic ice stream. *Science*, 252(5003), 242-246, 1991
9. ^ British Antarctic Survey. "Description of Ice Streams". Retrieved 2009-01-26.
10. ^ [a] [b] http://link.springer.com/referenceworkentry/10.1007%2F978-90-481-2642-2_72/fulltext.html
11. ^ Boulton, G.S. [1974] "Processes and patterns of glacial erosion", (In Coates, D.R. ed., *Glacial Geomorphology*. A Proceedings Volume of the fifth Annual Geomorphology Symposia series, held at Binghamton, New York, September 26–28, 1974. Binghamton, N.Y., State University of New York, p. 41-87. (Publications in Geomorphology))
12. ^ "What causes the blue color that sometimes appears in snow and ice ?". Webexhibits.org. Retrieved 2013-01-04.
13. ^ Benson, C.S., 1961, "Stratigraphic studies in the snow and firn of the Greenland Ice Sheet", *Res. Rep. 70*, U.S. Army Snow, Ice and Permafrost Res Establ., Corps of Eng., 120 pp
14. ^ "Glacier change and related hazards in Switzerland". UNEP. Retrieved 2009-01-05.
15. ^ "Frank Paul, et al., 2004, Rapid disintegration of Alpine glaciers observed with satellite data, GEOPHYSICAL RESEARCH LETTERS, VOL. 31, L21402, " (PDF). 2004.
16. ^ "Recent Global Glacier Retreat Overview" (PDF). Retrieved 2013-01-04.
17. ^ Greve, R.; Blatter, H. (2009). *Dynamics of Ice Sheets and Glaciers*. Springer. doi:10.1007/978-3-642-03415-2. ISBN 978-3-642-03414-5.
18. ^ W.S.B. Paterson, Physics of ice
19. ^ "Moulin 'Blanc': NASA Expedition Probes Deep Within a Greenland Glacier". NASA. 2006-12-11. Retrieved 2009-01-05.
20. ^ Glacier properties Hunter College CUNY lectures
21. ^ T. Strozzi et al.: *The Evolution of a Glacier Surge Observed with the ERS Satellites* (pdf, 1.3 Mb)
22. ^ "The Brúarjökull Project: Sedimentary environments of a surging glacier. The Brúarjökull Project research idea". Hi.is. Retrieved 2013-01-04.
23. ^ Meier & Post (1969)
24. ^ [a] [b] http://people.deas.harvard.edu/~vtsai/files/EkstromNettlesTsai_Science2006.pdf Ekström, G., M. Nettles, and V. C. Tsai (2006)"Seasonality and Increasing Frequency of Greenland Glacial Earthquakes", *Science*, 311, 5768, 1756-1758, doi:10.1126/science.1122112
25. ^ [a] [b] http://people.deas.harvard.edu/~vtsai/files/TsaiEkstrom_JGR2007.pdf Tsai, V. C. and G. Ekström (2007). "Analysis of Glacial Earthquakes", J. Geophys. Res., 112, F03S22, doi:10.1029/2006JF000596

26. ^ Easterbrook, D.J. (1999). *Surface Processes and Landforms* (2 ed.). New Jersey: Prentice-Hall, Inc. p. 546. ISBN 0-13-860958-6.
27. ^ "Glossary of Glacier Terminology". Pubs.usgs.gov. 2012-06-20. Retrieved 2013-01-04.
28. ^ "C.D. Ollier: "Australian Landforms and their History", National Mapping Fab, Geoscience Australia". Ga.gov.au. 2010-11-18. Retrieved 2013-01-04.
29. ^ KINCAID, JONI L.; KLEIN, ANDREW G. (2004). "Retreat of the Irian Jaya Glaciers from 2000 to 2002 as Measured from IKONOS Satellite Images". Portland, Maine, USA. pp. 147–157. Retrieved 2009-01-05.
30. ^ "Hawaiian Glaciers Reveal Clues to Global Climate Change". Geology.com. 2007-01-26. Retrieved 2013-01-04.
31. ^ "French Colonies – Crozet Archipelago". Discoverfrance.net. 2010-12-09. Retrieved 2013-01-04.
32. ^ Collins, Henry Hill; **Europe and the USSR**; p. 263. ISBN 1-256-35000-3
33. ^ "Yukon Beringia Interpretive Center". Beringia.com. 1999-04-12. Retrieved 2013-01-04.
34. ^ Earth History 2001 (page 15)
35. ^ "On the Zoogeography of the Holarctic Region". Wku.edu. Retrieved 2013-01-04.
36. ^ 'Glaciers & Glaciation' (Arnold, London 1998) Douglas Benn and David Evans, pp324-326
37. ^ "Kettle geology". Britannica Online. Retrieved 2009-03-12.
38. ^ "Kargel, J.S. et al.:"Martian Polar Ice Sheets and Mid-Latitude Debris-Rich Glaciers, and Terrestrial Analogs", Third International Conference on Mars Polar Science and Exploration, Alberta, Canada, October 13-17, 2003 (pdf 970 Kb)" (PDF). Retrieved 2013-01-04.
39. ^ "Martian glaciers: did they originate from the atmosphere? ESA Mars Express, 20 January 2006". Esa.int. 2006-01-20. Retrieved 2013-01-04.
40. ^ Head, J. et al. 2005. Tropical to mid-latitude snow and ice accumulation, flow and glaciation on Mars. Nature: 434. 346-350
41. ^ Source: Brown University Posted Monday, October 17, 2005 (2005-10-17). "Mars' climate in flux: Mid-latitude glaciers | SpaceRef – Your Space Reference". Marstoday.com. Retrieved 2013-01-04.
42. ^ Richard Lewis (2008-04-23). "Glaciers Reveal Martian Climate Has Been Recently Active | Brown University News and Events". News.brown.edu. Retrieved 2013-01-04.
43. ^ Plaut, J. et al. 2008. Radar Evidence for Ice in Lobate Debris Aprons in the Mid-Northern Latitudes of Mars. Lunar and Planetary Science XXXIX. 2290.pdf
44. ^ Holt, J. et al. 2008. Radar Sounding Evidence for Ice within Lobate Debris Aprons near Hellas Basin, Mid-Southern Latitudes of Mars. Lunar and Planetary Science XXXIX. 2441.pdf

References

- *This article draws heavily on the corresponding article in the Spanish-language Wikipedia, which was accessed in the version of 24 July 2005.*
- Hambrey, Michael; Alean, Jürg (2004). *Glaciers* (2nd ed.). Cambridge University Press. ISBN 0-521-82808-2. OCLC 54371738. An excellent less-technical treatment of all aspects, with superb photographs and firsthand accounts of glaciologists' experiences. All images of this book can be found online (see Weblinks: Glaciers-online)
- Benn, Douglas I.; Evans, David J. A. (1999). *Glaciers and Glaciation*. Arnold. ISBN 0-470-23651-5. OCLC 38329570.
- Bennett, M. R.; Glasser, N. F. (1996). *Glacial Geology: Ice Sheets and Landforms*. John Wiley & Sons. ISBN 0-471-96344-5. OCLC 33359888 37536152.
- Hambrey, Michael (1994). *Glacial Environments*. University of British Columbia Press, UCL Press. ISBN 0-7748-0510-2. OCLC 30512475. An undergraduate-level textbook.
- Knight, Peter G (1999). *Glaciers*. Cheltenham: Nelson Thornes. ISBN 0-7487-4000-7. OCLC 42656957 63064183 77294832. A textbook for undergraduates avoiding mathematical complexities
- Walley, Robert (1992). *Introduction to Physical Geography*. Wm. C. Brown Publishers. A textbook devoted to explaining the geography of our planet.
- W. S. B. Paterson (1994). *Physics of Glaciers* (3rd ed.). Pergamon Press. ISBN 0-08-013972-8. OCLC 26188. A comprehensive reference on the physical principles underlying formation and behavior.

External links

- Glacial structures – photo atlas
- Glaciers of the Pyrenees
- NOW on PBS “On Thin Ice”
- Photo project tracks changes in Himalayan glaciers since 1921
- Short radio episode ***California Glaciers*** from *The Mountains of California* by John Muir, 1894. California Legacy Project
- Dyanamics of Glaciers

This page titled 20.2: Glaciers is shared under a not declared license and was authored, remixed, and/or curated by Lumen Learning via source content that was edited to the style and standards of the LibreTexts platform.

Index

A

Absolute age
10.2: Absolute Ages of Rocks
absolute ages
10: Geologic History
absolute dating
10.6: Geologic Time Scale
abyssal plain
7.3: The Seafloor
accessible education
InfoPage
adaptation development
10.4: History of Earth's Life Forms
adaptations
11.4: Ocean Organisms
19.1: Deserts
adaptive homework
InfoPage
age
10.2: Absolute Ages of Rocks
agriculture
18: Climate and Climate Change
18.1: Climate Systems and Change
air
12.1: The Atmosphere
air masses
13: Weather
air pressure
13.1: Weather Processes and Systems
alpine glaciers
20: Glaciers
20.1: Glacial Erosion and Deposition
amateur astronomers
16.3: Telescopes
ancient environments
10.6: Geologic Time Scale
ancient organisms
10.3: Fossils
angle of incidence
15.5: Sun – Earth Relationship- The Seasons
applications
3: Rocks
aquatic
11.2: Thermocline
aquatic ecosystems
11: Ocean Water and Life
aquifer
6.3: Groundwater
aquifers
6: Surface Water
Arecibo Observatory
16.3: Telescopes
asteroid
17.4: Other Objects in the Solar System
asteroid impact
17.4: Other Objects in the Solar System
asteroids
17: The Solar System
astronomical discoveries
17.1: Introduction to the Solar System
astronomical observations
16: Light and the Sun
astronomy
1: Introduction to Earth Science
14.1: The Universe
15.1: Stars
16.2: Doppler Effect
17.4: Other Objects in the Solar System
astronomy education
16.3: Telescopes
astrophysics
15.1: Stars
15.2: Stellar Life Cycle
16.3: Telescopes
atmosphere
10.5: Early Earth
12: The Atmosphere
12.1: The Atmosphere
atmospheric pressure
12.1: The Atmosphere
atomic mass
2.2: Atoms to Molecules
atoms
2: Minerals
2.2: Atoms to Molecules
attribution
16.1: Electromagnetic Spectrum

B

basalt
3.1: Igneous Rocks
beach
8.1: Waves and Coastal Features
beach formation
8: Coastal Processes
big bang theory
14: Astronomy
biodiversity
6.6: Wetlands
7: The Ocean
7.5: Oceans and Coastal Environments
10.4: History of Earth's Life Forms
10.5: Early Earth
11: Ocean Water and Life
11.4: Ocean Organisms
18: Climate and Climate Change
18.1: Climate Systems and Change
19.1: Deserts
19.2: Desertification
19.3: Dunes
black hole
15.2: Stellar Life Cycle
black holes
15: Stellar Evolution and Our Sun
blizzards
13.1: Weather Processes and Systems
blue planet
7.1: Introduction to the Oceans
blueshift
16.2: Doppler Effect
body fossils
10.3: Fossils
brightness
15.1: Stars

C

Cambrian period
10.4: History of Earth's Life Forms
carbon dating
10.2: Absolute Ages of Rocks
CC
16.1: Electromagnetic Spectrum
celestial objects
17.4: Other Objects in the Solar System
cementation
3.2: Sedimentary Rocks
characteristics
2.5: Minerals and Mineral Groups
chemical bonding
2: Minerals
chemical composition
2.5: Minerals and Mineral Groups
2.6: Mineral Formation
chemical elements
2.2: Atoms to Molecules
10.5: Early Earth
chemistry
2.2: Atoms to Molecules
cinder cone
5.2: Types of Volcanoes
cirques
20: Glaciers
classification
2.4: Mineral Identification
2.5: Minerals and Mineral Groups
3.1: Igneous Rocks
cleavage
2.4: Mineral Identification
climate
7.4: Ocean Currents
7.5: Oceans and Coastal Environments
10.5: Early Earth
12.1: The Atmosphere
13.1: Weather Processes and Systems
18: Climate and Climate Change
18.1: Climate Systems and Change
19.1: Deserts
climate change
1: Introduction to Earth Science
1.1: Principles of Science
6: Surface Water
6.1: Earth's Fresh Water
6.6: Wetlands
11.1: Thermohaline circulation
18: Climate and Climate Change
18.1: Climate Systems and Change
19: Deserts
19.2: Desertification
20.2: Glaciers
climate impact
11.1: Thermohaline circulation

climate influence
7.7: Currents
11: Ocean Water and Life
climate regulation
7: The Ocean
climatology
1.2: Earth Science and Its Branches
coastal features
8: Coastal Processes
coastal management
8: Coastal Processes
coastal regions
19.3: Dunes
coastline
11.3: Tsunamis
coding
Back Matter
collaborative platform
Front Matter
InfoPage
color
2.4: Mineral Identification
comet
17.4: Other Objects in the Solar System
compaction
3.2: Sedimentary Rocks
composite volcano
5.2: Types of Volcanoes
composite volcanoes
5: Volcanoes
compounds
2.2: Atoms to Molecules
condensation
6.1: Earth's Fresh Water
conservation
19.3: Dunes
conservation efforts
6.6: Wetlands
constellation
15.1: Stars
constellations
15: Stellar Evolution and Our Sun
construction
3: Rocks
contact metamorphism
3.3: Metamorphic Rocks
contamination
6: Surface Water
content
16.1: Electromagnetic Spectrum
continental drift
4.2: Plate Tectonics
continental glaciers
20: Glaciers
20.1: Glacial Erosion and Deposition
continental shelf
7.3: The Seafloor
convection currents
4.1: Earth's Interior
Copernican Revolution
17.1: Introduction to the Solar System
Copernicus
17: The Solar System
core
4: Earth's Interior Processes
core collapse
15.2: Stellar Life Cycle
core mantle crust
4.1: Earth's Interior
Coriolis
7.4: Ocean Currents
Coriolis effect
7.7: Currents
cosmic events
14: Astronomy
cosmology
14.1: The Universe
crust
2.1: Materials of Earth's Crust
4: Earth's Interior Processes
crystal structure
2.5: Minerals and Mineral Groups
crystalline solids
2.3: States of Matter
currents
7.4: Ocean Currents
8: Coastal Processes
customization
Front Matter

D

damage
11.3: Tsunamis
dams
6.5: Floods
dark energy
14: Astronomy
14.1: The Universe
dark matter
14: Astronomy
14.1: The Universe
data
1.1: Principles of Science
Dating Methods
10.2: Absolute Ages of Rocks
day and night
15.4: The Sun and the Earth-Moon System
deep ocean
7.3: The Seafloor
deep ocean currents
7.2: Composition of Ocean Water
7.7: Currents
deep water formation
11: Ocean Water and Life
definitions
Back Matter
Deforestation
19.2: Desertification
deformation
9.1: Tectonic Stress and Geologic Structures
density
7.2: Composition of Ocean Water
7.4: Ocean Currents
density gradients
11.1: Thermohaline circulation
Department of Education
Front Matter
deposition
20: Glaciers
depth
11.2: Thermocline
desert
19.1: Deserts
19.3: Dunes
desert features
19.1: Deserts
desert reclamation
19.2: Desertification
desertification
19: Deserts
19.1: Deserts
19.2: Desertification
deserts
19: Deserts
dinosaur extinction
10.6: Geologic Time Scale
dinosaur fossils
10.3: Fossils
discharge
6.3: Groundwater
divergent boundaries
5.1: Volcanic Activity
Doppler effect
16: Light and the Sun
16.2: Doppler Effect
dune ecosystems
19.3: Dunes
dune habitats
19.3: Dunes
dwarf planet
17.4: Other Objects in the Solar System
dwarf planets
17: The Solar System

E

Earth
2.1: Materials of Earth's Crust
10.5: Early Earth
12: The Atmosphere
15.1: Stars
15.4: The Sun and the Earth-Moon System
17.2: Inner Planets
earth density
4.1: Earth's Interior
Earth Fresh Water
6: Surface Water
Earth history
10.6: Geologic Time Scale
Earth science
1: Introduction to Earth Science
1.1: Principles of Science
1.2: Earth Science and Its Branches
2.6: Mineral Formation
3.1: Igneous Rocks
3.3: Metamorphic Rocks
20.1: Glacial Erosion and Deposition

Earth's axis
15.5: Sun – Earth Relationship- The Seasons
Earth's Crust
2: Minerals
earth's interior
4: Earth's Interior Processes
earthquake causes
4.4: Earthquakes
earthquake energy
4.1: Earth's Interior
earthquake prediction
4.4: Earthquakes
earthquakes
4: Earth's Interior Processes
4.2: Plate Tectonics
4.4: Earthquakes
9: Mountain Building
9.1: Tectonic Stress and Geologic Structures
11.3: Tsunamis
ecological assessment
6.6: Wetlands
ecological importance
6.2: Streams and Rivers
ecology
1: Introduction to Earth Science
19: Deserts
19.1: Deserts
19.3: Dunes
ecosystem
6.6: Wetlands
ecosystem services
6.6: Wetlands
ecosystems
6: Surface Water
7.5: Oceans and Coastal Environments
11.2: Thermocline
11.4: Ocean Organisms
18: Climate and Climate Change
18.1: Climate Systems and Change
19.1: Deserts
edwin hubble
14: Astronomy
elastic deformation
9: Mountain Building
9.1: Tectonic Stress and Geologic Structures
electromagnetic radiation
16.4: Nuclear Fusion
electromagnetic spectrum
16: Light and the Sun
electromagnetic waves
16.2: Doppler Effect
electrons
2.2: Atoms to Molecules
energy
15.1: Stars
15.3: The Sun
16.4: Nuclear Fusion
energy transfer
8.1: Waves and Coastal Features
environmental changes
10.4: History of Earth's Life Forms
environmental impact
2.7: Mining and Mineral Use
18: Climate and Climate Change
Environmental Issues
18.1: Climate Systems and Change
environmental science
1: Introduction to Earth Science
1.2: Earth Science and Its Branches
19.1: Deserts
equilibrium
2.3: States of Matter
equinox
15.5: Sun – Earth Relationship- The Seasons
erosion
2.1: Materials of Earth's Crust
3.2: Sedimentary Rocks
8: Coastal Processes
8.1: Waves and Coastal Features
19: Deserts
19.3: Dunes
erosion process
6.2: Streams and Rivers
erosional features
20: Glaciers
eruption
5.4: Volcanic Eruptions
eruptions
4.3: Volcanoes
5.3: Volcanic Landforms and Geothermal Activity
essential questions
4.4: Earthquakes
6.1: Earth's Fresh Water
evaporation
6.1: Earth's Fresh Water
evolution
1.1: Principles of Science
10: Geologic History
10.5: Early Earth
10.6: Geologic Time Scale
examples
Back Matter
explosive eruptions
4.3: Volcanoes
5.4: Volcanic Eruptions
extinction
10.4: History of Earth's Life Forms
extraterrestrial life
17.3: Outer Planets
extreme weather
18: Climate and Climate Change
extrusive
5.3: Volcanic Landforms and Geothermal Activity

F

faulting
9.2: Mountain Formation
faults
9: Mountain Building
fauna
19: Deserts
financial burden
InfoPage
fishing methods
11.4: Ocean Organisms
flash floods
6.5: Floods
flood control
6.6: Wetlands
floods
6.5: Floods
flora
19: Deserts
Florida Memorial University
16.1: Electromagnetic Spectrum
folding
9.2: Mountain Formation
foliation
3.3: Metamorphic Rocks
formation
3: Rocks
3.1: Igneous Rocks
10.5: Early Earth
formation of rocks
3.3: Metamorphic Rocks
fossil record
10.3: Fossils
fossilization
10.3: Fossils
fossils
10: Geologic History
10.1: Relative Ages of Rocks
10.2: Absolute Ages of Rocks
10.3: Fossils
10.4: History of Earth's Life Forms
10.6: Geologic Time Scale
fracture
2.4: Mineral Identification
frequency
16: Light and the Sun
frequency change
16.2: Doppler Effect
fresh water
6.2: Streams and Rivers
7.2: Composition of Ocean Water
functionality
Back Matter

G

galaxies
14.1: The Universe
galaxy movement
14: Astronomy
Galilean moons
17.3: Outer Planets
Galileo
17: The Solar System
gas
2.3: States of Matter
gas giants
17.3: Outer Planets
gases
12: The Atmosphere
geocentric model
17: The Solar System
17.1: Introduction to the Solar System
geologic structures
9: Mountain Building

geologic time
10.4: History of Earth's Life Forms
10.6: Geologic Time Scale
geologic time scale
10.6: Geologic Time Scale
geological
2.5: Minerals and Mineral Groups
geological activity
4.4: Earthquakes
9.1: Tectonic Stress and Geologic Structures
17.2: Inner Planets
geological evidence
4.2: Plate Tectonics
geological features
4.2: Plate Tectonics
geological formations
5.3: Volcanic Landforms and Geothermal Activity
geological monitoring
5.4: Volcanic Eruptions
geological processes
3.2: Sedimentary Rocks
9.2: Mountain Formation
geological time
10: Geologic History
10.1: Relative Ages of Rocks
10.2: Absolute Ages of Rocks
geology
1: Introduction to Earth Science
1.2: Earth Science and Its Branches
2.6: Mineral Formation
2.7: Mining and Mineral Use
3.1: Igneous Rocks
3.3: Metamorphic Rocks
4.1: Earth's Interior
5.2: Types of Volcanoes
6.2: Streams and Rivers
10: Geologic History
10.1: Relative Ages of Rocks
10.5: Early Earth
geoscience
1.2: Earth Science and Its Branches
geothermal
4.3: Volcanoes
glacial deposits
20.1: Glacial Erosion and Deposition
glacial erosion
20: Glaciers
20.1: Glacial Erosion and Deposition
glacial features
20.1: Glacial Erosion and Deposition
glaciation
20.2: Glaciers
glacier dynamics
20.2: Glaciers
glacier formation
20.2: Glaciers
glacier mass balance
20.2: Glaciers
glacier retreat
20.2: Glaciers
glaciers
20: Glaciers
20.1: Glacial Erosion and Deposition
glaciology
20.2: Glaciers
global conveyor belt
11: Ocean Water and Life
11.1: Thermohaline circulation
global warming
18: Climate and Climate Change
20.2: Glaciers
global winds
7.7: Currents
glossary
Back Matter
granite
3.1: Igneous Rocks
gravitational collapse
15.2: Stellar Life Cycle
gravitational pull
7.6: Tides
gravity
15.1: Stars
16.4: Nuclear Fusion
17.1: Introduction to the Solar System
greenhouse gases
12.1: The Atmosphere
18.1: Climate Systems and Change
groundwater
6.1: Earth's Fresh Water
6.2: Streams and Rivers
6.3: Groundwater
groundwater overuse
6: Surface Water
guidelines
Back Matter
Gulf Stream
7: The Ocean
7.7: Currents
11: Ocean Water and Life

H

habitats
11.2: Thermocline
hardness
2.4: Mineral Identification
hazards
4.3: Volcanoes
heat
3: Rocks
heliocentric model
17: The Solar System
17.1: Introduction to the Solar System
helium
15.3: The Sun
16.4: Nuclear Fusion
high tide
7.6: Tides
historical views
17.1: Introduction to the Solar System
history
10.5: Early Earth
11.3: Tsunamis
hot spots
5: Volcanoes
5.1: Volcanic Activity
Hubble Space Telescope
16.3: Telescopes
Hubble's Law
14.1: The Universe
human activities
19.2: Desertification
human interaction
19.1: Deserts
humidity
12.1: The Atmosphere
13: Weather
13.1: Weather Processes and Systems
hurricanes
13: Weather
13.1: Weather Processes and Systems
hydrogen
15.3: The Sun
16.4: Nuclear Fusion
hydrologic cycle
6: Surface Water
6.1: Earth's Fresh Water
hypothesis
1.1: Principles of Science

I

ice ages
20.1: Glacial Erosion and Deposition
ice flow
20.2: Glaciers
ice sheet
20.2: Glaciers
identification
2.4: Mineral Identification
igneous
2.1: Materials of Earth's Crust
5.3: Volcanic Landforms and Geothermal Activity
igneous rocks
2: Minerals
3: Rocks
3.1: Igneous Rocks
3.3: Metamorphic Rocks
index fossils
10.3: Fossils
inner planets
17.2: Inner Planets
insolation
15.5: Sun – Earth Relationship- The Seasons
interactive textbooks
InfoPage
intrusive
5.3: Volcanic Landforms and Geothermal Activity

J

James Webb Space Telescope
16.3: Telescopes
Jupiter
17.3: Outer Planets

K

Kepler
17: The Solar System
Kuiper belt
17.4: Other Objects in the Solar System

L

lake
- 11.2: Thermocline

land degradation
- 19.2: Desertification

landforms
- 5.3: Volcanic Landforms and Geothermal Activity
- 20: Glaciers

lava
- 2.6: Mineral Formation
- 4.3: Volcanoes
- 5.3: Volcanic Landforms and Geothermal Activity

lava flow
- 5.4: Volcanic Eruptions

lava viscosity
- 5: Volcanoes

layers of earth
- 4.1: Earth's Interior

learning outcomes
- 4.4: Earthquakes
- 6.1: Earth's Fresh Water

levees
- 6.5: Floods

LibreTexts
- Front Matter
- InfoPage

license
- 16.1: Electromagnetic Spectrum

licensing
- Front Matter

life
- 12: The Atmosphere

life cycle
- 15: Stellar Evolution and Our Sun

liquid
- 2.3: States of Matter

lithification
- 3.2: Sedimentary Rocks

lithosphere
- 4: Earth's Interior Processes
- 4.2: Plate Tectonics

lithosphere asthenosphere
- 4.1: Earth's Interior

living fossils
- 10.3: Fossils

longshore currents
- 8.1: Waves and Coastal Features

low tide
- 7.6: Tides

luster
- 2.4: Mineral Identification

M

m51
- 14.1: The Universe

magma
- 2.6: Mineral Formation
- 4.3: Volcanoes
- 5.1: Volcanic Activity
- 5.2: Types of Volcanoes
- 5.3: Volcanic Landforms and Geothermal Activity
- 5.4: Volcanic Eruptions

mantle
- 4: Earth's Interior Processes

marine
- 11.2: Thermocline

marine ecosystems
- 7: The Ocean
- 7.1: Introduction to the Oceans

marine life
- 7.1: Introduction to the Oceans
- 7.5: Oceans and Coastal Environments
- 11.4: Ocean Organisms

marine science
- 7.2: Composition of Ocean Water

Mars
- 17.2: Inner Planets

Mars exploration
- 17.2: Inner Planets

mass
- 15.2: Stellar Life Cycle

mass extinction
- 10.4: History of Earth's Life Forms

mass extinctions
- 10: Geologic History

matter
- 2.2: Atoms to Molecules

media
- 16.1: Electromagnetic Spectrum

melting
- 2.3: States of Matter

meltwater
- 20: Glaciers

mercury
- 17.2: Inner Planets

Meridional Overturning Circulation
- 11.1: Thermohaline circulation

metal production
- 2.7: Mining and Mineral Use

metamorphic
- 2.1: Materials of Earth's Crust

metamorphic rocks
- 2: Minerals
- 3: Rocks
- 3.3: Metamorphic Rocks

metamorphism
- 2.6: Mineral Formation
- 3: Rocks
- 3.3: Metamorphic Rocks
- 9.2: Mountain Formation

meteor
- 17.4: Other Objects in the Solar System

meteorology
- 1: Introduction to Earth Science
- 1.2: Earth Science and Its Branches
- 13.1: Weather Processes and Systems

mineral composition
- 3: Rocks
- 3.1: Igneous Rocks

mineral extraction
- 2.7: Mining and Mineral Use

mineral formation
- 2.6: Mineral Formation

mineral groups
- 2.5: Minerals and Mineral Groups

mineral identification
- 2: Minerals
- 2.5: Minerals and Mineral Groups

minerals
- 2.1: Materials of Earth's Crust
- 2.4: Mineral Identification
- 2.5: Minerals and Mineral Groups
- 2.6: Mineral Formation
- 2.7: Mining and Mineral Use

mining
- 2.7: Mining and Mineral Use

molecules
- 2: Minerals
- 2.2: Atoms to Molecules

moon
- 7.6: Tides
- 15.4: The Sun and the Earth-Moon System

moons
- 17: The Solar System

moraines
- 20.1: Glacial Erosion and Deposition

motion
- 17.1: Introduction to the Solar System

Mount St. Helens
- 5.4: Volcanic Eruptions

mountain building
- 4.4: Earthquakes
- 9: Mountain Building
- 9.1: Tectonic Stress and Geologic Structures

N

NASA
- 16.3: Telescopes

National Science Foundation
- Front Matter

natural disasters
- 6.5: Floods
- 11.3: Tsunamis

natural hazards
- 4.4: Earthquakes

neap tides
- 7.6: Tides

Neptune
- 17.3: Outer Planets

neutron stars
- 15.2: Stellar Life Cycle

neutrons
- 2.2: Atoms to Molecules

ngc 5195
- 14: Astronomy

northern hemisphere
- 15.5: Sun – Earth Relationship- The Seasons

nuclear fusion
- 15: Stellar Evolution and Our Sun
- 15.3: The Sun
- 16: Light and the Sun
- 16.4: Nuclear Fusion

nutrient transport
- 7: The Ocean

O

observation
- 1.1: Principles of Science

observer
16: Light and the Sun
16.2: Doppler Effect
ocean
7.3: The Seafloor
7.4: Ocean Currents
7.6: Tides
8.1: Waves and Coastal Features
11.2: Thermocline
11.3: Tsunamis
11.4: Ocean Organisms
ocean basins
7.7: Currents
ocean circulation
11.1: Thermohaline circulation
ocean currents
7: The Ocean
7.1: Introduction to the Oceans
7.5: Oceans and Coastal Environments
11.1: Thermohaline circulation
ocean dynamics
8: Coastal Processes
ocean movement
7.7: Currents
ocean significance
7.1: Introduction to the Oceans
ocean trenches
7.1: Introduction to the Oceans
ocean water
7: The Ocean
7.2: Composition of Ocean Water
ocean zones
7.5: Oceans and Coastal Environments
oceanography
1: Introduction to Earth Science
1.2: Earth Science and Its Branches
7: The Ocean
7.3: The Seafloor
7.5: Oceans and Coastal Environments
oceans
7: The Ocean
7.5: Oceans and Coastal Environments
oceans formation
7.1: Introduction to the Oceans
OER
InfoPage
ore
2.7: Mining and Mineral Use
organisms
6.2: Streams and Rivers
10.4: History of Earth's Life Forms
orogenic process
9.2: Mountain Formation
outer planets
17.3: Outer Planets
overgrazing
19.2: Desertification
ozone layer
12: The Atmosphere

P

Pacific Ring of Fire
5: Volcanoes
5.1: Volcanic Activity
paleontology
1.2: Earth Science and Its Branches
10.1: Relative Ages of Rocks
10.4: History of Earth's Life Forms
parallax
15.1: Stars
pelagic
11.2: Thermocline
permeability
6.3: Groundwater
phase transitions
2.3: States of Matter
physical characteristics
2.4: Mineral Identification
physical geography
19.3: Dunes
physical properties
2.5: Minerals and Mineral Groups
planetary geology
17.2: Inner Planets
planetary motion
17.1: Introduction to the Solar System
planetary rings
17.3: Outer Planets
planets
15.4: The Sun and the Earth-Moon System
17: The Solar System
17.1: Introduction to the Solar System
plasma
2.3: States of Matter
plastic deformation
9: Mountain Building
9.1: Tectonic Stress and Geologic Structures
plate tectonics
1.1: Principles of Science
4: Earth's Interior Processes
4.1: Earth's Interior
4.2: Plate Tectonics
9: Mountain Building
9.1: Tectonic Stress and Geologic Structures
9.2: Mountain Formation
17.2: Inner Planets
pollution
7.2: Composition of Ocean Water
11: Ocean Water and Life
12.1: The Atmosphere
porosity
6.3: Groundwater
postsecondary education
Front Matter
precipitation
6.5: Floods
13: Weather
13.1: Weather Processes and Systems
18.1: Climate Systems and Change
19: Deserts
prediction
11.3: Tsunamis
prehistoric creatures
10.3: Fossils
Pressure
3: Rocks
properties
2.4: Mineral Identification
protons
2.2: Atoms to Molecules
pumice
3.1: Igneous Rocks

R

radiation
15.3: The Sun
radio astronomy
14: Astronomy
radioactivity
10.2: Absolute Ages of Rocks
radiometric dating
10: Geologic History
10.2: Absolute Ages of Rocks
recharge
6.3: Groundwater
reclamation
2.7: Mining and Mineral Use
red giants
15: Stellar Evolution and Our Sun
redshift
16.2: Doppler Effect
reflecting telescopes
16.3: Telescopes
regional metamorphism
3.3: Metamorphic Rocks
Relative age
10.2: Absolute Ages of Rocks
relative ages
10: Geologic History
relative dating
10.1: Relative Ages of Rocks
10.6: Geologic Time Scale
reservoirs
6.5: Floods
reverse fault
9.2: Mountain Formation
revolution
15.4: The Sun and the Earth-Moon System
Ring of Fire
4.3: Volcanoes
5.4: Volcanic Eruptions
rip currents
8.1: Waves and Coastal Features
rivers
6.2: Streams and Rivers
rock
2.6: Mineral Formation
rock cycle
2.1: Materials of Earth's Crust
rock formation
10: Geologic History
rock fracturing
9: Mountain Building
rock stresses
9: Mountain Building
rocks
2.1: Materials of Earth's Crust
3.1: Igneous Rocks
9.1: Tectonic Stress and Geologic Structures

rotation
15.4: The Sun and the Earth-Moon System

S

salinity
7: The Ocean
7.2: Composition of Ocean Water
7.4: Ocean Currents
7.5: Oceans and Coastal Environments
11.1: Thermohaline circulation
salinity gradients
11: Ocean Water and Life
sand movement
19.3: Dunes
sandstone
3.2: Sedimentary Rocks
Saturn
17.3: Outer Planets
science
1: Introduction to Earth Science
1.1: Principles of Science
scientific method
1: Introduction to Earth Science
1.1: Principles of Science
seafloor
7.3: The Seafloor
seamount
7.3: The Seafloor
seasons
15.4: The Sun and the Earth-Moon System
sediment
2.6: Mineral Formation
sediment deposition
3.2: Sedimentary Rocks
6.2: Streams and Rivers
sedimentary
2.1: Materials of Earth's Crust
sedimentary rock types
3.2: Sedimentary Rocks
sedimentary rocks
2: Minerals
3: Rocks
3.2: Sedimentary Rocks
3.3: Metamorphic Rocks
10.1: Relative Ages of Rocks
10.6: Geologic Time Scale
seismic activity
5.1: Volcanic Activity
seismic waves
4: Earth's Interior Processes
4.1: Earth's Interior
4.4: Earthquakes
seismology
1.2: Earth Science and Its Branches
severe weather
13: Weather
shield volcano
5.2: Types of Volcanoes
shield volcanoes
5: Volcanoes
shoreline preservation
8: Coastal Processes
silicate minerals
2: Minerals
silicates
2.5: Minerals and Mineral Groups
sky
15.1: Stars
soil erosion
19.2: Desertification
solar flare
15.3: The Sun
solar prominence
15.3: The Sun
solar radiation
12.1: The Atmosphere
15.5: Sun – Earth Relationship- The Seasons
solar system
10.5: Early Earth
15: Stellar Evolution and Our Sun
15.3: The Sun
15.4: The Sun and the Earth-Moon System
17: The Solar System
17.1: Introduction to the Solar System
17.3: Outer Planets
17.4: Other Objects in the Solar System
solid
2.3: States of Matter
sonar
7.3: The Seafloor
sound
16: Light and the Sun
space exploration
16.3: Telescopes
spectral lines
16.2: Doppler Effect
spiral galaxy
14.1: The Universe
spring tides
7.6: Tides
springs
6.3: Groundwater
star power
16.4: Nuclear Fusion
stars
15: Stellar Evolution and Our Sun
15.1: Stars
states of matter
2.3: States of Matter
stellar evolution
15: Stellar Evolution and Our Sun
15.2: Stellar Life Cycle
stellar remnants
15.2: Stellar Life Cycle
strain
9.1: Tectonic Stress and Geologic Structures
stratosphere
12: The Atmosphere
stratovolcano
5.2: Types of Volcanoes
stream erosion
6.2: Streams and Rivers
structured approach
Back Matter
subduction
5.1: Volcanic Activity
subduction zones
4.2: Plate Tectonics
subsidence
6.3: Groundwater
summer solstice
15.5: Sun – Earth Relationship- The Seasons
sun
7.6: Tides
15: Stellar Evolution and Our Sun
15.3: The Sun
15.4: The Sun and the Earth-Moon System
16.4: Nuclear Fusion
sunspots
15.3: The Sun
supernova
15.2: Stellar Life Cycle
supernovae
15: Stellar Evolution and Our Sun
superposition
10.1: Relative Ages of Rocks
10.2: Absolute Ages of Rocks
surface currents
7.7: Currents
8.1: Waves and Coastal Features
surface mining
2.7: Mining and Mineral Use
sustainability
6: Surface Water
sustainable management
11.4: Ocean Organisms
sustainable practices
19: Deserts
19.2: Desertification
synonyms
Back Matter

T

table
Back Matter
tectonic geomorphology
9.2: Mountain Formation
tectonic plate boundaries
4: Earth's Interior Processes
5: Volcanoes
tectonic plates
4.2: Plate Tectonics
5.1: Volcanic Activity
5.2: Types of Volcanoes
5.3: Volcanic Landforms and Geothermal Activity
tectonic stress
4.4: Earthquakes
9: Mountain Building
9.1: Tectonic Stress and Geologic Structures
tectonics
4.3: Volcanoes
telescopes
16: Light and the Sun
16.3: Telescopes

temperature
7.2: Composition of Ocean Water
7.4: Ocean Currents
11.1: Thermohaline circulation
11.2: Thermocline
12: The Atmosphere
12.1: The Atmosphere
13: Weather
18: Climate and Climate Change
18.1: Climate Systems and Change
19: Deserts
temperature gradients
11: Ocean Water and Life
temperature regulation
7.1: Introduction to the Oceans
temperature variations
12: The Atmosphere
terms
Back Matter
terrestrial planets
17.2: Inner Planets
textbook costs
Front Matter
theory
1.1: Principles of Science
thermal equilibrium
2.3: States of Matter
thermocline
11.2: Thermocline
thermohaline circulation
7.7: Currents
11: Ocean Water and Life
11.1: Thermohaline circulation
thrust
9.2: Mountain Formation
thunderstorms
13: Weather
13.1: Weather Processes and Systems
tidal range
7.6: Tides
tides
7.6: Tides
8: Coastal Processes
15.4: The Sun and the Earth-Moon System
tilt of axis
15.5: Sun – Earth Relationship- The Seasons
tornadoes
13: Weather
13.1: Weather Processes and Systems
trace fossils
10.3: Fossils
traditional textbooks
InfoPage
trench
7.3: The Seafloor
Tropic of Cancer
15.5: Sun – Earth Relationship- The Seasons
troposphere
12: The Atmosphere
tsunami
11.3: Tsunamis

U

ultraviolet light
16.4: Nuclear Fusion
unconformity
10.1: Relative Ages of Rocks
underground mining
2.7: Mining and Mineral Use
understanding earth interior
4.1: Earth's Interior
uniformitarianism
10: Geologic History
10.1: Relative Ages of Rocks
universal solvent
7.2: Composition of Ocean Water
universe expansion
14: Astronomy
universe formation
14.1: The Universe
upwelling
7.4: Ocean Currents
7.7: Currents
Uranus
17.3: Outer Planets
UV radiation
12: The Atmosphere

V

vegetation
6.5: Floods
Venus
17.2: Inner Planets
visible light
16.4: Nuclear Fusion
volcanic
5.3: Volcanic Landforms and Geothermal Activity
volcanic activity
5: Volcanoes
5.1: Volcanic Activity
5.4: Volcanic Eruptions
9.2: Mountain Formation
volcanic eruptions
4.3: Volcanoes
5: Volcanoes
5.1: Volcanic Activity
5.2: Types of Volcanoes
volcanic hazards
5.4: Volcanic Eruptions
volcanic outgassing theory
7.1: Introduction to the Oceans
volcanic types
5: Volcanoes
volcano
4.3: Volcanoes
volcanoes
4: Earth's Interior Processes
4.2: Plate Tectonics
5: Volcanoes
5.1: Volcanic Activity
5.2: Types of Volcanoes
5.4: Volcanic Eruptions
volcanology
1.2: Earth Science and Its Branches
5.2: Types of Volcanoes

W

warning
11.3: Tsunamis
water
7.4: Ocean Currents
water composition
7.1: Introduction to the Oceans
water conservation
6: Surface Water
6.1: Earth's Fresh Water
water cycle
6.1: Earth's Fresh Water
6.2: Streams and Rivers
6.5: Floods
7.2: Composition of Ocean Water
7.5: Oceans and Coastal Environments
water distribution
6.1: Earth's Fresh Water
water motion
8.1: Waves and Coastal Features
water movement
8: Coastal Processes
water properties
6: Surface Water
water purification
6.6: Wetlands
water table
6.3: Groundwater
wave
16.2: Doppler Effect
waves
7.4: Ocean Currents
8: Coastal Processes
8.1: Waves and Coastal Features
11.3: Tsunamis
16: Light and the Sun
weather
12.1: The Atmosphere
13: Weather
13.1: Weather Processes and Systems
18: Climate and Climate Change
18.1: Climate Systems and Change
weather fronts
13: Weather
weathering
2.1: Materials of Earth's Crust
3.2: Sedimentary Rocks
wells
6.3: Groundwater
wetland
6.6: Wetlands
wetlands
6.5: Floods
6.6: Wetlands
whirlpool galaxy
14: Astronomy
14.1: The Universe
white dwarfs
15.2: Stellar Life Cycle
William Hopper
16.1: Electromagnetic Spectrum
wind waves
8.1: Waves and Coastal Features

winter solstice

15.5: Sun – Earth Relationship- The Seasons

Glossary

Sample Word 1 | Sample Definition 1

Detailed Licensing

Overview

Title: Earth Science (Lumen)

Webpages: 101

All licenses found:

- Undeclared: 100% (101 pages)

By Page

- Earth Science (Lumen) - *Undeclared*
  - Front Matter - *Undeclared*
    - TitlePage - *Undeclared*
    - InfoPage - *Undeclared*
    - Table of Contents - *Undeclared*
    - Licensing - *Undeclared*
  - 1: Introduction to Earth Science - *Undeclared*
    - 1.1: Principles of Science - *Undeclared*
    - 1.2: Earth Science and Its Branches - *Undeclared*
  - 2: Minerals - *Undeclared*
    - 2.1: Materials of Earth's Crust - *Undeclared*
    - 2.2: Atoms to Molecules - *Undeclared*
    - 2.3: States of Matter - *Undeclared*
    - 2.4: Mineral Identification - *Undeclared*
    - 2.5: Minerals and Mineral Groups - *Undeclared*
    - 2.6: Mineral Formation - *Undeclared*
    - 2.7: Mining and Mineral Use - *Undeclared*
  - 3: Rocks - *Undeclared*
    - 3.1: Igneous Rocks - *Undeclared*
    - 3.2: Sedimentary Rocks - *Undeclared*
    - 3.3: Metamorphic Rocks - *Undeclared*
  - 4: Earth's Interior Processes - *Undeclared*
    - 4.1: Earth's Interior - *Undeclared*
    - 4.2: Plate Tectonics - *Undeclared*
    - 4.3: Volcanoes - *Undeclared*
    - 4.4: Earthquakes - *Undeclared*
  - 5: Volcanoes - *Undeclared*
    - 5.1: Volcanic Activity - *Undeclared*
    - 5.2: Types of Volcanoes - *Undeclared*
    - 5.3: Volcanic Landforms and Geothermal Activity - *Undeclared*
    - 5.4: Volcanic Eruptions - *Undeclared*
  - 6: Surface Water - *Undeclared*
    - 6.1: Earth's Fresh Water - *Undeclared*
    - 6.2: Streams and Rivers - *Undeclared*
    - 6.3: Groundwater - *Undeclared*
    - 6.4: Water Resources - *Undeclared*
    - 6.5: Floods - *Undeclared*
    - 6.6: Wetlands - *Undeclared*
  - 7: The Ocean - *Undeclared*
    - 7.1: Introduction to the Oceans - *Undeclared*
    - 7.2: Composition of Ocean Water - *Undeclared*
    - 7.3: The Seafloor - *Undeclared*
    - 7.4: Ocean Currents - *Undeclared*
    - 7.5: Oceans and Coastal Environments - *Undeclared*
    - 7.6: Tides - *Undeclared*
    - 7.7: Currents - *Undeclared*
  - 8: Coastal Processes - *Undeclared*
    - 8.1: Waves and Coastal Features - *Undeclared*
  - 9: Mountain Building - *Undeclared*
    - 9.1: Tectonic Stress and Geologic Structures - *Undeclared*
    - 9.2: Mountain Formation - *Undeclared*
  - 10: Geologic History - *Undeclared*
    - 10.1: Relative Ages of Rocks - *Undeclared*
    - 10.2: Absolute Ages of Rocks - *Undeclared*
    - 10.3: Fossils - *Undeclared*
    - 10.4: History of Earth's Life Forms - *Undeclared*
    - 10.5: Early Earth - *Undeclared*
    - 10.6: Geologic Time Scale - *Undeclared*
  - 11: Ocean Water and Life - *Undeclared*
    - 11.1: Thermohaline circulation - *Undeclared*
    - 11.2: Thermocline - *Undeclared*
    - 11.3: Tsunamis - *Undeclared*
    - 11.4: Ocean Organisms - *Undeclared*
  - 12: The Atmosphere - *Undeclared*
    - 12.1: The Atmosphere - *Undeclared*
  - 13: Weather - *Undeclared*
    - 13.1: Weather Processes and Systems - *Undeclared*
  - 14: Astronomy - *Undeclared*
    - 14.1: The Universe - *Undeclared*
  - 15: Stellar Evolution and Our Sun - *Undeclared*

- 15.1: Stars - *Undeclared*
- 15.2: Stellar Life Cycle - *Undeclared*
- 15.3: The Sun - *Undeclared*
- 15.4: The Sun and the Earth-Moon System - *Undeclared*
- 15.5: Sun – Earth Relationship- The Seasons - *Undeclared*

16: Light and the Sun - *Undeclared*
- 16.1: Electromagnetic Spectrum - *Undeclared*
- 16.2: Doppler Effect - *Undeclared*
- 16.3: Telescopes - *Undeclared*
- 16.4: Nuclear Fusion - *Undeclared*

17: The Solar System - *Undeclared*
- 17.1: Introduction to the Solar System - *Undeclared*
- 17.2: Inner Planets - *Undeclared*
- 17.3: Outer Planets - *Undeclared*
- 17.4: Other Objects in the Solar System - *Undeclared*

18: Climate and Climate Change - *Undeclared*
- 18.1: Climate Systems and Change - *Undeclared*
- 18.2: Climate and Weather - *Undeclared*
- 18.3: Global Climate Change - *Undeclared*

19: Deserts - *Undeclared*
- 19.1: Deserts - *Undeclared*
- 19.2: Desertification - *Undeclared*
- 19.3: Dunes - *Undeclared*
- 19.4: Video- Desertification - *Undeclared*

20: Glaciers - *Undeclared*
- 20.1: Glacial Erosion and Deposition - *Undeclared*
- 20.2: Glaciers - *Undeclared*

Back Matter - *Undeclared*
- Index - *Undeclared*
- Glossary - *Undeclared*
- Detailed Licensing - *Undeclared*