MTH 098 ELEMENTARY ALGEBRA



LibreTexts™

MTH 098 Elementary Algebra

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (https://LibreTexts.org) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide and accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden born from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by NICE CXone Expert and was supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptions contact info@LibreTexts.org or visit our main website at https://LibreTexts.org.

This text was compiled on 06/09/2025

TABLE OF CONTENTS

1: Foundations

- 1.1: Prelude to Foundations of Algebra
- 1.2: Introduction to Whole Numbers
  - 1.2E: Exercises
- 1.3: The Language of Algebra and the Order of Operations
  - 1.3E: Exercises
- 1.4: Add and Subtract Integers
  - 1.4E: Exercises
- 1.5: Multiply and Divide Integers
  - 1.5E: Exercises
- 1.6: Visualize Fractions
  - 1.6E: Exercises
- 1.7: Add and Subtract Fractions
  - 1.7E: Exercises
- 1.8: Decimals
  - 1.8E: Exercises
- 1.9: The Real Numbers
  - 1.9E: Exercises
- 1.10: Properties of Real Numbers
  - 1.10E: Exercises
- 1.11: Systems of Measurement
  - 1.11E: Exercises
- Chapter 1 Review Exercises

2: Solving Linear Equations and Inequalities

- 2.1: Solve Equations Using the Subtraction and Addition Properties of Equality
  - 2.1E: Exercises
- 2.2: Solve Equations using the Division and Multiplication Properties of Equality
  - 2.2E: Exercises
- 2.3: Solve Equations with Variables and Constants on Both Sides
  - 2.3E: Exercises
- 2.4: Use a General Strategy to Solve Linear Equations
  - 2.4E: Exercises
- 2.5: Solve Equations with Fractions or Decimals
  - 2.5E: Exercises
- 2.6: Solve Proportions and Similar Figure Applications
  - 2.6E: Exercises
- 2.7: Solve Linear Inequalities
  - 2.7E: Exercises for Section 2.7
  - 2.7E2: Exercises
- 2.8: Solve Compound Inequalities

- 2.8E: Exercises for Section 2.8
- 2.8b: Solve Compound Inequalities - additional explanations
  - 2.8bE: Exercises
- Chapter 2 Review Exercises

3: Math Models

- 3.1: Solve a Formula for a Specific Variable
  - 3.1E: Exercises
- 3.2: Use a Problem-Solving Strategy
  - 3.2E: Exercises
- 3.3: Solve Percent Applications
  - 3.3E: Exercises
- 3.4: Solve Mixture Applications
  - 3.4E: Exercises
- 3.5: Triangles, Rectangles, and the Pythagorean Theorem
  - 3.5E: Exercises
- 3.6: Solve Uniform Motion Applications
  - 3.6E: Exercises
- 3.7: Solve Applications with Linear Inequalities
  - 3.7E: Exercises
- Chapter 3 Review Exercises

4: Graphs

- 4.1: Use the Rectangular Coordinate System
  - 4.1E: Exercises
- 4.2: Graph Linear Equations in Two Variables
  - 4.2E: Exercises
- 4.3: Graph with Intercepts
  - 4.3E: Exercises
- 4.4: Understanding the Slope of a Line
  - 4.4E: Exercises
- 4.5: Use the Slope–Intercept Form of an Equation of a Line
  - 4.5E: Exercises
- 4.6: Find the Equation of a Line
  - 4.6E: Exercises
- 4.7: Graphs of Linear Inequalities
  - 4.7E: Exercises
- Chapter 4 Review Exercises

5: Systems of Linear Equations

- 5.1: Solve Systems of Equations by Graphing
  - 5.1E: Exercises
- 5.2: Solve Systems of Equations by Substitution
  - 5.2E: Exercises

- 5.3: Solve Systems of Equations by Elimination
  - 5.3E: Exercises
- 5.4: Solve Applications with Systems of Equations
  - 5.4E: Exercises
- 5.5: Solve Mixture Applications with Systems of Equations
  - 5.5E: Exercises
- 5.6: Graphing Systems of Linear Inequalities
  - 5.6E: Exercises
- Chapter 5 Review Exercises

6: Polynomials

- 6.1: Add and Subtract Polynomials
  - 6.1E: Exercises
- 6.2: Use Multiplication Properties of Exponents
  - 6.2E: Exercises
- 6.3: Multiply Polynomials
  - 6.3E: Exercises
- 6.4: Special Products
  - 6.4E: Exercises
- 6.5: Divide Monomials
  - 6.5E: Exercises
- 6.6: Divide Polynomials
  - 6.6E: Exercises
- 6.7: Integer Exponents and Scientific Notation
  - 6.7E: Exercises
- Chapter 6 Review Exercises

7: Factoring

- 7.1: Greatest Common Factor and Factor by Grouping
  - 7.1E: Exercises
- 7.2: Factor Quadratic Trinomials with Leading Coefficient 1
  - 7.2E: Exercises
- 7.3: Factor Quadratic Trinomials with Leading Coefficient Other than 1
  - 7.3E: Exercises
- 7.4: Factor Special Products
  - 7.4E: Exercises
- 7.5: General Strategy for Factoring Polynomials
  - 7.5E: Exercises
- 7.6: Quadratic Equations
  - 7.6E: Exercises
- Chapter 7 Review Exercises

CHAPTER OVERVIEW

1: Foundations

1.1: Prelude to Foundations of Algebra
1.2: Introduction to Whole Numbers
1.2E: Exercises
1.3: The Language of Algebra and the Order of Operations
1.3E: Exercises
1.4: Add and Subtract Integers
1.4E: Exercises
1.5: Multiply and Divide Integers
1.5E: Exercises
1.6: Visualize Fractions
1.6E: Exercises
1.7: Add and Subtract Fractions
1.7E: Exercises
1.8: Decimals
1.8E: Exercises
1.9: The Real Numbers
1.9E: Exercises
1.10: Properties of Real Numbers
1.10E: Exercises
1.11: Systems of Measurement
1.11E: Exercises
Chapter 1 Review Exercises

This page titled 1: Foundations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

1.1: Prelude to Foundations of Algebra

Just like a building needs a firm foundation to support it, your study of algebra needs to have a firm foundation. To ensure this, we begin this book with a review of arithmetic operations with whole numbers, integers, fractions, and decimals, so that you have a solid base that will support your study of algebra.



Figure 1.1.1: In order to be structurally sound, the foundation of a building must be carefully constructed.

Contributors and Attributions

- Lynn Marecek (Santa Ana College) and MaryAnne Anthony-Smith (formerly of Santa Ana College). This content produced by OpenStax and is licensed under a Creative Commons Attribution License 4.0 license.

This page titled 1.1: Prelude to Foundations of Algebra is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.1: Prelude to Foundations of Algebra** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.2: Introduction to Whole Numbers

Learning Objectives

- By the end of this section, you will be able to:
- Use place value with whole numbers
- Identify multiples and and apply divisibility tests
- Find prime factorizations and least common multiples

As we begin our study of elementary algebra, we need to refresh some of our skills and vocabulary. This chapter will focus on whole numbers, integers, fractions, decimals, and real numbers. We will also begin our use of algebraic notation and vocabulary.

Use Place Value with Whole Numbers

The most basic numbers used in algebra are the numbers we use to count objects in our world: $1, 2, 3, 4$, and so on. These are called the **counting numbers**. Counting numbers are also called *natural numbers*. If we add zero to the counting numbers, we get the set of **whole numbers**.

- Counting Numbers: $1, 2, 3, \ldots$
- Whole Numbers: $0, 1, 2, 3, \ldots$

The notation "$\ldots$" is called ellipsis and means "and so on," or that the pattern continues endlessly.

We can visualize counting numbers and whole numbers on a **number line** (see Figure $1.2.1$).




Figure 1.2.1: The numbers on the number line get larger as they go from left to right, and smaller as they go from right to left. While this number line shows only the whole numbers 0 through 6, the numbers keep going without end.

Doing the Manipulative Mathematics activity "Number Line-Part 1" will help you develop a better understanding of the counting numbers and the whole numbers.

Our **number system** is called a place value system, because the value of a digit depends on its position in a number. Figure $1.2.2$ shows the **place values**. The place values are separated into groups of three, which are called periods. The periods are *ones*, *thousands*, *millions*, *billions*, *trillions*, and so on. In a written number, commas separate the periods.

Place Value														
Trillions			Billions			Millions			Thousands			Ones		
Hundred trillions	Ten trillions	Trillions	Hundred billions	Ten billions	Billions	Hundred millions	Ten millions	Millions	Hundred thousands	Ten thousands	Thousands	Hundreds	Tens	Ones
								5	2	7	8	1	9	4

Figure 1.2.2: The number 5278194 is shown in the chart. The digit 5 is in the millions place. The digit 2 is in the hundred-thousands place. The digit 7 is in the ten-thousands place. The digit 8 is in the thousands place. The digit 1 is in the hundreds place. The digit 9 is in the tens place. The digit 4 is in the ones place.

Exercise 1.2.1

In the number 63407218, find the place value of each digit:

1. 7

2. 0
3. 1
4. 6
5. 3

Answer

Place the number in the place value chart:

Trillions			Billions			Millions			Thousands			Ones		
Hundred trillions	Ten trillions	Trillions	Hundred billions	Ten billions	Billions	Hundred millions	Ten millions	Millions	Hundred thousands	Ten thousands	Thousands	Hundreds	Tens	Ones
							6	3	4	0	7	2	1	8

1. The 7 is in the thousands place.
2. The 0 is in the ten thousands place.
3. The 1 is in the tens place.
4. The 6 is in the ten-millions place.
5. The 3 is in the millions place.

Exercise 1.2.2

For the number 27493615 find the place value of each digit:

1. 2
2. 1
3. 4
4. 7
5. 5

Answer

1. ten millions
2. tens
3. hundred thousands
4. millions
5. ones

Exercise 1.2.3

For the number 519711641328 find the place value of each digit:

1. 9
2. 4
3. 2
4. 6
5. 7

Answer

1. billions
2. ten thousands

3. tens
4. hundred thousands
5. hundred millions

When you write a check, you write out the number in words as well as in digits. To write a number in words, write the number in each period, followed by the name of the period, without the *s* at the end. Start at the left, where the periods have the largest value. The ones period is not named. The commas separate the periods, so wherever there is a comma in the number, put a comma between the words (see Figure 1.2.3). The number 74218369is written as seventy-four million, two hundred eighteen thousand, three hundred sixty-nine.




Figure 1.2.3

NAME A WHOLE NUMBER IN WORDS.


1. Start at the left and name the number in each period, followed by the period name.
2. Put commas in the number to separate the periods.
3. Do not name the ones period.

Exercise 1.2.4

Name the number 8165432098710using words.

Answer

Name the number in each period, followed by the period name.



Put the commas in to separate the periods.

So, 8165432098710is named as eight trillion, one hundred sixty-five billion, four hundred thirty-two million, ninety-eight thousand, seven hundred ten.

Exercise 1.2.5

Name the number 9,258,137,904,0619,258,137,904,061 using words.

Answer

nine trillion, two hundred fifty-eight billion, one hundred thirty-seven million, nine hundred four thousand, sixty-one

Exercise 1.2.6

Name the number 17,864,325,619,00417,864,325,619,004 using words.

Answer

seventeen trillion, eight hundred sixty-four billion, three hundred twenty-five million, six hundred nineteen thousand four

We are now going to reverse the process by writing the digits from the name of the number. To write the number in digits, we first look for the clue words that indicate the periods. It is helpful to draw three blanks for the needed periods and then fill in the blanks with the numbers, separating the periods with commas.

WRITE A WHOLE NUMBER USING DIGITS.


1. Identify the words that indicate periods. (Remember, the ones period is never named.)
2. Draw three blanks to indicate the number of places needed in each period. Separate the periods by commas.
3. Name the number in each period and place the digits in the correct place value position.

Exercise 1.2.7

Write *nine billion, two hundred forty-six million, seventy-three thousand, one hundred eighty-nine* as a whole number using digits.

Answer

Identify the words that indicate periods.
Except for the first period, all other periods must have three places. Draw three blanks to indicate the number of places needed in each period. Separate the periods by commas.
Then write the digits in each period.



The number is 9,246,073,189.

Exercise 1.2.8

Write the number two billion, four hundred sixty-six million, seven hundred fourteen thousand, fifty-one as a whole number using digits.

Answer

2,466,714,051

Exercise 1.2.9

Write the number eleven billion, nine hundred twenty-one million, eight hundred thirty thousand, one hundred six as a whole number using digits.

Answer

11,921,830,106

In 2013, the U.S. Census Bureau estimated the population of the state of New York as 19,651,127. We could say the population of New York was approximately 20 million. In many cases, you don't need the exact value; an approximate number is good enough.

The process of approximating a number is called **rounding**. Numbers are rounded to a specific place value, depending on how much accuracy is needed. Saying that the population of New York is approximately 20 million means that we rounded to the

millions place.

Exercise 1.2.10 How to Round Whole Numbers

Round 23,658 to the nearest hundred.

Answer

Step 1. Locate the given place value with an arrow. All digits to the left do not change.	Locate the hundreds place in 23,658.	hundredths place ↓ 23,658
Step 2. Underline the digit to the right of the given place value.	Underline the 5, which is to the right of the hundreds place.	hundredths place ↓ 23,658
Step 3. Is this digit greater than or equal to 5? Yes–add 1 to the digit in the given place value. No–do not change the digit in the given place value.	Add 1 to the 6 in the hundreds place, since 5 is greater than or equal to 5.	23,658 add 1 ↗
Step 4. Replace all digits to the right of the given place value with zeros.	Replace all digits to the right of the hundreds place with zeros.	23,700 add 1 ↗ ↖ replace with 0s So, 23,700 is rounded to the nearest hundred.

Exercise 1.2.11

Round to the nearest hundred: 17,852.

Answer

17,900

Exercise 1.2.12

Round to the nearest hundred: 468,751.

Answer

468,800

ROUND WHOLE NUMBERS.

1. Locate the given place value and mark it with an arrow. All digits to the left of the arrow do not change.
2. Underline the digit to the right of the given place value.
3. Is this digit greater than or equal to 5?
   - Yes–add 11 to the digit in the given place value.
   - No–do not change the digit in the given place value.
4. Replace all digits to the right of the given place value with zeros.


Exercise 1.2.13

Round 103,978103,978 to the nearest:


1. hundred
2. thousand
3. ten thousand

Answer

1.

Locate the hundreds place in 103,978.	
Underline the digit to the right of the hundreds place.	
Since 7 is greater than or equal to 5, add 1 to the 9. Replace all digits to the right of the hundreds place with zeros.	
	So, 104,000 is 103,978 rounded to the nearest hundred.

2.

Locate the thousands place and underline the digit to the right of the thousands place.	
Since 9 is greater than or equal to 5, add 1 to the 3. Replace all digits to the right of the hundreds place with zeros.	
	So, 104,000 is 103,978 rounded to the nearest thousand.

3.

Locate the ten thousands place and underline the digit to the right of the ten thousands place.	
Since 3 is less than 5, we leave the 0 as is, and then replace the digits to the right with zeros.	
	So, 100,000 is 103,978 rounded to the nearest ten thousand.

Exercise 1.2.14

Round 206,981 to the nearest: 1. hundred 2. thousand 3. ten thousand.

Answer

1. 207,000
2. 207,000

3. 210,000

Exercise 1.2.15

Round 784,951 to the nearest: 1. hundred 2. thousand 3. ten thousand.

Answer

1. 785,000
2. 785,000
3. 780,000

Identify Multiples and Apply Divisibility Tests

The numbers 2, 4, 6, 8, 10, and 12 are called **multiples** of 2. A multiple of 2 can be written as the **product** of a counting number and 2.

$$\underset{2\cdot 1,}{2,}\quad \underset{2\cdot 2,}{4,}\quad \underset{2\cdot 3,}{6,}\quad \underset{2\cdot 4,}{8,}\quad \underset{2\cdot 5,}{10,}\quad \underset{2\cdot 6}{12,\ldots}$$

Figure 1.2.4

Similarly, a multiple of 3 would be the product of a counting number and 3.

$$\underset{3\cdot 1,}{3,}\quad \underset{3\cdot 2,}{6,}\quad \underset{3\cdot 3,}{9,}\quad \underset{3\cdot 4,}{12,}\quad \underset{3\cdot 5,}{15,}\quad \underset{3\cdot 6}{18,\ldots}$$

Figure 1.2.5

We could find the multiples of any number by continuing this process.

Note

Doing the Manipulative Mathematics activity “Multiples” will help you develop a better understanding of multiples.

Table 1.2.1 shows the multiples of 2 through 9 for the first 12 counting numbers.

Table 1.2.1

Counting Number	1	2	3	4	5	6	7	8	9	10	11	12
Multiples of 2	2	4	6	8	10	12	14	16	18	20	22	24
Multiples of 3	3	6	9	12	15	18	21	24	27	30	33	36
Multiples of 4	4	8	12	16	20	24	28	32	36	40	44	48
Multiples of 5	5	10	15	20	25	30	35	40	45	50	55	60
Multiples of 6	6	12	18	24	30	36	42	48	54	60	66	72
Multiples of 7	7	14	21	28	35	42	49	56	63	70	77	84
Multiples of 8	8	16	24	32	40	48	56	64	72	80	88	96

Counting Number	1	2	3	4	5	6	7	8	9	10	11	12
Multiples of 9	9	18	27	36	45	54	63	72	81	90	99	108
Multiples of 10	10	20	30	40	50	60	70	80	90	100	110	120

MULTIPLE OF A NUMBER

A number is a **multiple** of n if it is the product of a counting number and n.

Another way to say that 15 is a multiple of 3 is to say that 15 is **divisible** by 3. That means that when we divide 3 into 15, we get a counting number. In fact, $15 \div 3$ is 5, so 15 is $5 \cdot 3$.

DIVISIBLE BY A NUMBER

If a number m is a multiple of n, then m is **divisible** by n

Look at the multiples of 5 in Table 1.2.1. They all end in 5 or 0. Numbers with last digit of 5 or 0 are divisible by 5. Looking for other patterns in Table 1.2.1 that shows multiples of the numbers 2 through 9, we can discover the following divisibility tests:

DIVISIBILITY TESTS

A number is divisible by:

- 2 if the last digit is 0, 2, 4, 6, or 8.
- 3 if the sum of the digits is divisible by 3.
- 5 if the last digit is 5 or 0.
- 6 if it is divisible by both 2 and 3.
- 10 if it ends with 0.

Exercise 1.2.16

Is 5625 divisible by 2? By 3? By 5? By 6? By 10?

Answer

Is 5625 divisible by 2?
Does it end in 0, 2, 4, 6, or 8? No. 5625 is not divisible by 2. (1.2.1)

Is 5625 divisible by 3?
What is the sum of the digits? $5+6+2+5=18$
Is the sum divisible by 3? Yes, 5625 is divisible by 3. (1.2.2)

Is 5625 divisible by 5 or 10?
What is the last digit? It is 5. 5625 is divisible by 5 but not by 10. (1.2.3)

Is 5625 divisible by 6?
Is it divisible by both 2 and 3? No, 5625 is not divisible by 2, so 5625 is not divisible by 6. (1.2.4)

Exercise 1.2.17

Determine whether 4,962 is divisible by 2, by 3, by 5, by 6, and by 10.

Answer

by 2, 3, and 6

Exercise 1.2.18

Determine whether 3,765 is divisible by 2, by 3, by 5, by 6, and by 10.

Answer

by 3 and 5

Find Prime Factorizations and Least Common Multiples

In mathematics, there are often several ways to talk about the same ideas. So far, we've seen that if m is a multiple of n, we can say that m is divisible by n. For example, since 72 is a multiple of 8, we say 72 is divisible by 8. Since 72 is a multiple of 9, we say 72 is divisible by 9. We can express this still another way.

Since $8 \cdot 9 = 72$, we say that 8 and 9 are **factors** of 72. When we write $72 = 8 \cdot 9$, we say we have factored 72.



Figure 1.2.6

Other ways to factor 72 are $1 \cdot 72$, $2 \cdot 36$, $3 \cdot 24$, $4 \cdot 18$ and $6 \cdot 12$. Seventy-two has many factors: 1, 2, 3, 4, 6, 8, 9, 12, 18, 36, and 72.

FACTORS

If $a \cdot b = m$, then a and b are **factors** of m.

Some numbers, like 72, have many factors. Other numbers have only two factors.

Note

Doing the Manipulative Mathematics activity "Model Multiplication and Factoring" will help you develop a better understanding of multiplication and factoring.

PRIME NUMBER AND COMPOSITE NUMBER

A **prime number** is a counting number greater than 1, whose only factors are 1 and itself.

A **composite number** is a counting number that is not prime. A composite number has factors other than 1 and itself.

Note

Doing the Manipulative Mathematics activity "Prime Numbers" will help you develop a better understanding of prime numbers.

The counting numbers from 2 to 19 are listed in Figure 1.2.7, with their factors. Make sure to agree with the "prime" or "composite" label for each!

Number	Factors	Prime or Composite?
2	1,2	Prime
3	1,3	Prime
4	1,2,4	Composite
5	1,5	Prime
6	1,2,3,6	Composite
7	1,7	Prime
8	1,2,4,8	Composite
9	1,3,9	Composite
10	1,2,5,10	Composite

Number	Factors	Prime or Composite?
11	1,11	Prime
12	1,2,3,4,6,12	Composite
13	1,13	Prime
14	1,2,7,14	Composite
15	1,3,5,15	Composite
16	1,2,4,8,16	Composite
17	1,17	Prime
18	1,2,3,6,9,18	Composite
19	1,19	Prime

Figure 1.2.7

The **prime numbers** less than 20 are 2, 3, 5, 7, 11, 13, 17, and 19. Notice that the only even prime number is 2.

A composite number can be written as a unique **product** of primes. This is called the **prime factorization** of the number. Finding the prime factorization of a composite number will be useful later in this course.

PRIME FACTORIZATION

The **prime factorization** of a number is the product of prime numbers that equals the number.

To find the prime factorization of a composite number, find any two factors of the number and use them to create two branches. If a factor is prime, that branch is complete. Circle that prime!



If the factor is not prime, find two factors of the number and continue the process. Once all the branches have circled primes at the end, the factorization is complete. The composite number can now be written as a product of prime numbers.

Exercise 1.2.19

Factor 48.

Answer

Step 1. Find two factors whose product is the given number. Use these numbers to create two branches.	$48 = 2 \cdot 24$	48 2 24
Step 2. If a factor is prime, that branch is complete. Circle the prime.	2 is prime. Circle the prime.	48 (2) 24

Step 3. If a factor is not prime, write it as the product of two factors and continue the process.	24 is not prime. Break it into 2 more factors.	
	4 and 6 are not prime. Break them each into two factors.	
	2 and 3 are prime, so circle them.	
Step 4. Write the composite number as the product of all the circled primes.		$48 = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 3$

We say $2 \cdot 2 \cdot 2 \cdot 2 \cdot 3$ is the prime factorization of 48. We generally write the primes in ascending order. Be sure to multiply the factors to verify your answer!

If we first factored 48 in a different way, for example as $6 \cdot 8$, the result would still be the same. Finish the prime factorization and verify this for yourself.

Exercise 1.2.20

Find the prime factorization of 80.

Answer

$2 \cdot 2 \cdot 2 \cdot 2 \cdot 5$

Exercise 1.2.21

Find the prime factorization of 60.

Answer

$2 \cdot 2 \cdot 3 \cdot 5$


FIND THE PRIME FACTORIZATION OF A COMPOSITE NUMBER.

1. Find two factors whose product is the given number, and use these numbers to create two branches.
2. If a factor is prime, that branch is complete. Circle the prime, like a bud on the tree.
3. If a factor is not prime, write it as the product of two factors and continue the process.
4. Write the composite number as the product of all the circled primes.

Exercise 1.2.22

Find the prime factorization of 252.

Answer

Step 1. Find two factors whose product is 252. 12 and 21 are not prime. Break 12 and 21 into two more factors. Continue until all primes are factored.	
Step 2. Write 252 as the product of all the circled primes.	$252 = 2 \cdot 2 \cdot 3 \cdot 3 \cdot 7$

? Exercise 1.2.23

Find the prime factorization of 126.

Answer

$2 \cdot 3 \cdot 3 \cdot 7$

? Exercise 1.2.24

Find the prime factorization of 294.

Answer

$2 \cdot 3 \cdot 7 \cdot 7$

One of the reasons we look at multiples and primes is to use these techniques to find the **least common multiple** of two numbers. This will be useful when we add and subtract fractions with different **denominator**s. Two methods are used most often to find the least common multiple and we will look at both of them.

The first method is the Listing Multiples Method. To find the least common multiple of 12 and 18, we list the first few multiples of 12 and 18:

12: 12, 24, **36**, 48, 60, **72**, 84, 96, **108**...

18: 18, **36**, 54, **72**, 90, **108**...

Common Multiples: 36, 72, 108...

Least Common Multiple: 36

Figure 1.2.8

Notice that some numbers appear in both lists. They are the **common multiples** of 12 and 18.

We see that the first few common multiples of 12 and 18 are 36, 72, and 108. Since 36 is the smallest of the common multiples, we call it the *least common multiple*. We often use the abbreviation LCM.

LEAST COMMON MULTIPLE

The **least common multiple** (LCM) of two numbers is the smallest number that is a multiple of both numbers.

The procedure box lists the steps to take to find the LCM using the prime factors method we used above for 12 and 18.

FIND THE LEAST COMMON MULTIPLE BY LISTING MULTIPLES.

1. List several multiples of each number.
2. Look for the smallest number that appears on both lists.
3. This number is the LCM.

Exercise 1.2.25

Find the least common multiple of 15 and 20 by listing multiples.

Answer

Make lists of the first few multiples of 15 and of 20, and use them to find the least common multiple.	**15:** 15, 30, 45, **60**, 75, 90, 105, 120 **20:** 20, 40, **60**, 80, 100, 120, 140, 160
Look for the smallest number that appears in both lists.	The first number to appear on both lists is 60, so 60 is the least common multiple of 15 and 20.

Notice that 120 is in both lists, too. It is a common multiple, but it is not the *least* common multiple.

Exercise 1.2.26

Find the least common multiple by listing multiples: 9 and 12.

Answer

36

Exercise 1.2.27

Find the least common multiple by listing multiples: 18 and 24.

Answer


72

Our second method to find the least common multiple of two numbers is to use The Prime Factors Method. Let's find the LCM of 12 and 18 again, this time using their prime factors.

Exercise 1.2.28

Find the Least Common Multiple (LCM) of 12 and 18 using the prime factors method.

Answer

Step 1. Write each number as a product of primes.		
Step 2. List the primes of each number. Match primes vertically when possible.	List the primes of 12. List the primes of 18. Line up with the primes of 12 when possible. If not create a new column.	$12 = 2 \cdot 2 \cdot 3$ $18 = 2 \cdot \quad 3 \cdot 3$
Step 3. Bring down the number from each column.		$12 = 2 \cdot 2 \cdot 3$ $18 = 2 \cdot \quad 3 \cdot 3$ $\text{LCM} = 2 \cdot 2 \cdot 3 \cdot 3$
Step 4. Multiply the factors.		$\text{LCM} = 36$

Notice that the prime factors of $12(2 \cdot 2 \cdot 3)$ and the prime factors of $18(2 \cdot 3 \cdot 3)$ are included in the LCM $(2 \cdot 2 \cdot 3 \cdot 3)$. So 36 is the least common multiple of 12 and 18.

By matching up the common primes, each common prime factor is used only once. This way you are sure that 36 is the *least* common multiple.

Exercise 1.2.29

Find the LCM using the prime factors method: 9 and 12.

Answer

36

Exercise 1.2.30

Find the LCM using the prime factors method: 18 and 24.

Answer

72

FIND THE LEAST COMMON MULTIPLE USING THE PRIME FACTORS METHOD.

1. Write each number as a product of primes.
2. List the primes of each number. Match primes vertically when possible.
3. Bring down the columns.
4. Multiply the factors.

Exercise 1.2.31

Find the Least Common Multiple (LCM) of 24 and 36 using the prime factors method.

Answer

Find the primes of 24 and 36. Match primes vertically when possible. Bring down all columns.	$24 = 2 \cdot 2 \cdot 2 \cdot 3$ $36 = 2 \cdot 2 \cdot \quad 3 \cdot 3$ $\text{LCM} = 2 \cdot 2 \cdot 2 \cdot 3 \cdot 3$
Multiply the factors.	$\text{LCM} = 72$
	The LCM of 24 and 36 is 72.

Exercise 1.2.32

Find the LCM using the prime factors method: 21 and 28.

Answer

84

Exercise 1.2.33

Find the LCM using the prime factors method: 24 and 32.

Answer

96

Note

Access this online resource for additional instruction and practice with using whole numbers. You will need to enable Java in your web browser to use the application.

- Sieve of Eratosthenes

Key Concepts

- **Place Value** as in Figure.
- **Name a Whole Number in Words**
  1. Start at the left and name the number in each period, followed by the period name.
  2. Put commas in the number to separate the periods.
  3. Do not name the ones period.
- **Write a Whole Number Using Digits**
  1. Identify the words that indicate periods. (Remember the ones period is never named.)
  2. Draw 3 blanks to indicate the number of places needed in each period. Separate the periods by commas.
  3. Name the number in each period and place the digits in the correct place value position.
- **Round Whole Numbers**
  1. Locate the given place value and mark it with an arrow. All digits to the left of the arrow do not change.
  2. Underline the digit to the right of the given place value.
  3. Is this digit greater than or equal to 5?
     - Yes—add 1 to the digit in the given place value.
     - No—do not change the digit in the given place value.
  4. Replace all digits to the right of the given place value with zeros.
- **Divisibility Tests:** A number is divisible by:
  - 2 if the last digit is 0, 2, 4, 6, or 8.
  - 3 if the sum of the digits is divisible by 3.
  - 5 if the last digit is 5 or 0.
  - 6 if it is divisible by both 2 and 3.
  - 10 if it ends with 0.
- **Find the Prime Factorization of a Composite Number**
  1. Find two factors whose product is the given number, and use these numbers to create two branches.
  2. If a factor is prime, that branch is complete. Circle the prime, like a bud on the tree.
  3. If a factor is not prime, write it as the product of two factors and continue the process.
  4. Write the composite number as the product of all the circled primes.
- **Find the Least Common Multiple by Listing Multiples**
  1. List several multiples of each number.
  2. Look for the smallest number that appears on both lists.
  3. This number is the LCM.
- **Find the Least Common Multiple Using the Prime Factors Method**
  1. Write each number as a product of primes.
  2. List the primes of each number. Match primes vertically when possible.
  3. Bring down the columns.

4. Multiply the factors.

Glossary

composite number

A composite number is a counting number that is not prime. A composite number has factors other than 1 and itself.

counting numbers

The counting numbers are the numbers 1, 2, 3, …

divisible by a number

If a number m is a multiple of n, then m is divisible by n. (If 6 is a multiple of 3, then 6 is divisible by 3.)

factors

If $a \cdot b = m$, then a and b are factors of m. Since $3 \cdot 4 = 12$, then 3 and 4 are factors of 12.

least common multiple

The least common multiple of two numbers is the smallest number that is a multiple of both numbers.

multiple of a number

A number is a multiple of n if it is the product of a counting number and n.

number line

A number line is used to visualize numbers. The numbers on the number line get larger as they go from left to right, and smaller as they go from right to left.

origin

The origin is the point labeled 0 on a number line.

prime factorization

The prime factorization of a number is the product of prime numbers that equals the number.

prime number

A prime number is a counting number greater than 1, whose only factors are 1 and itself.

whole numbers

The whole numbers are the numbers 0, 1, 2, 3, ...

This page titled 1.2: Introduction to Whole Numbers is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.2: Introduction to Whole Numbers** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.2E: Exercises

Practice Makes Perfect

Use Place Value with Whole Numbers

In the following exercises, find the place value of each digit in the given numbers.

Exercise 1.2E.34

51,493

1. 1
2. 4
3. 9
4. 5
5. 3

Answer

1. thousands
2. hundreds
3. tens
4. ten thousands
5. ones

Exercise 1.2E.35

87,210

1. 2
2. 8
3. 0
4. 7
5. 1

Exercise 1.2E.36

164,285

1. 5
2. 6
3. 1
4. 8
5. 2

Answer

1. ones
2. ten thousands
3. hundred thousands
4. tens
5. hundreds

Exercise 1.2E.37

395,076

1. 5
2. 3
3. 7
4. 0
5. 9

Exercise 1.2E.38

93,285,170

1. 9
2. 8
3. 7
4. 5
5. 3

Answer

1. ten millions
2. ten thousands
3. tens
4. thousands
5. millions

Exercise 1.2E.39

36,084,215

1. 8
2. 6
3. 5
4. 4
5. 3

Exercise 1.2E.40

7,284,915,860,132

1. 7
2. 4
3. 5
4. 3
5. 0

Answer

1. trillions
2. billions
3. millions
4. tens
5. thousands

Exercise 1.2E.41

2,850,361,159,433

1. 9
2. 8
3. 6
4. 4
5. 2

In the following exercises, name each number using words.

Exercise 1.2E.42

1,078

Answer

one thousand, seventy-eight

Exercise 1.2E.43

5,902

Exercise 1.2E.44

364,510

Answer

three hundred sixty-four thousand, five hundred ten

Exercise 1.2E.45

146,023

Exercise 1.2E.46

5,846,103

Answer

five million, eight hundred forty-six thousand, one hundred three

Exercise 1.2E.47

1,458,398

Exercise 1.2E.48

37,889,005

Answer

thirty-seven million, eight hundred eighty-nine thousand, five

Exercise 1.2E.49

62,008,465

In the following exercises, write each number as a whole number using digits.

Exercise 1.2E.50

four hundred twelve

Answer

412

Exercise 1.2E.51

two hundred fifty-three

Exercise 1.2E.52

thirty-five thousand, nine hundred seventy-five

Answer

35,975

Exercise 1.2E.53

sixty-one thousand, four hundred fifteen

Exercise 1.2E.54

eleven million, forty-four thousand, one hundred sixty-seven

Answer

11,044,167

Exercise 1.2E.55

eighteen million, one hundred two thousand, seven hundred eighty-three

Exercise 1.2E.56

three billion, two hundred twenty-six million, five hundred twelve thousand, seventeen

Answer

3,226,512,017

Exercise 1.2E.57

eleven billion, four hundred seventy-one million, thirty-six thousand, one hundred six

In the following, round to the indicated place value.

Exercise 1.2E.58

Round to the nearest ten.

1. 386
2. 2,931

Answer

1. 390
2. 2,930

Exercise 1.2E.59

Round to the nearest ten.

1. 792
2. 5,647

Exercise 1.2E.60

Round to the nearest hundred.

1. 13,748
2. 391,794

Answer

1. 13,700
2. 391,800

Exercise 1.2E.61

Round to the nearest hundred.

1. 28,166
2. 481,628

Exercise 1.2E.62

Round to the nearest ten.

1. 1,492
2. 1,497

Answer

1. 1,490
2. 1,500

Exercise 1.2E.63

Round to the nearest ten.

1. 2,791
2. 2,795

Exercise 1.2E.64

Round to the nearest hundred.

1. 63,994
2. 63,940

Answer

1. 64,000
2. 63,900

Exercise 1.2E.65

Round to the nearest hundred.

1. 49,584
2. 49,548

In the following exercises, round each number to the nearest ⓐ hundred, ⓑ thousand, ⓒ ten thousand.

Exercise 1.2E.66

392,546

Answer

1. 392,500
2. 393,000
3. 390,000

Exercise 1.2E.67

619,348

Exercise 1.2E.68

2,586,991

Answer

1. 2,587,000
2. 2,587,000
3. 2,590,000

Exercise 1.2E.69

4,287,965

Identify Multiples and Factors

In the following exercises, use the divisibility tests to determine whether each number is divisible by 2, 3, 5, 6, and 10.

Exercise 1.2E.70

84

Answer

divisible by 2, 3, and 6

Exercise 1.2E.71

9,696

Exercise 1.2E.72

75

Answer

divisible by 3 and 5

Exercise 1.2E.73

78

Exercise 1.2E.74

900

Answer

divisible by 2, 3, 5, 6, and 10

Exercise 1.2E.75

800

Exercise 1.2E.76

986

Answer

divisible by 2

Exercise 1.2E.77

942

Exercise 1.2E.78

350

Answer

divisible by 2, 5, and 10

Exercise 1.2E.79

550

Exercise 1.2E.80

22,335

Answer

divisible by 3 and 5

Exercise 1.2E.81

39,075

Find Prime Factorizations and Least Common Multiples

In the following exercises, find the prime factorization.

Exercise 1.2E.82

86

Answer

$2 \cdot 43$

Exercise 1.2E.83

78

Exercise 1.2E.84

132

Answer

$2 \cdot 2 \cdot 3 \cdot 11$

Exercise 1.2E.85

455

Exercise 1.2E.86

693

Answer

$3 \cdot 3 \cdot 7 \cdot 11$

Exercise 1.2E.87

400

Exercise 1.2E.88

432

Answer

$2 \cdot 2 \cdot 2 \cdot 2 \cdot 3 \cdot 3 \cdot 3$

Exercise 1.2E.89

627

Exercise 1.2E.90

2,160

Answer

$2 \cdot 2 \cdot 2 \cdot 2 \cdot 3 \cdot 3 \cdot 3 \cdot 5$

Exercise 1.2E.91

2,520

In the following exercises, find the least common multiple of the each pair of numbers using the multiples method.

Exercise 1.2E.92

8, 12

Answer

24

Exercise 1.2E.93

4, 3

Exercise 1.2E.94

12, 16

Answer

48

Exercise 1.2E.95

30, 40

Exercise 1.2E.96

20, 30

Answer

60

Exercise 1.2E.97

44, 55

In the following exercises, find the least common multiple of each pair of numbers using the prime factors method.

Exercise 1.2E.98

8, 12

Answer

24

Exercise 1.2E.99

12, 16

Exercise 1.2E.100

28, 40

Answer

280

Exercise 1.2E.101

84, 90

Exercise 1.2E.102

55, 88

Answer

440

Exercise 1.2E.103

60, 72

Everyday Math

Exercise 1.2E.104

Writing a Check Jorge bought a car for $24,493. He paid for the car with a check. Write the purchase price in words.

Answer

twenty-four thousand, four hundred ninety-three dollars

Exercise 1.2E.105

Writing a Check Marissa's kitchen remodeling cost $18,549. She wrote a check to the contractor. Write the amount paid in words.

Exercise 1.2E.106

Buying a Car Jorge bought a car for $24,493. Round the price to the nearest

1. ten
2. hundred
3. thousand; and
4. ten-thousand.

Answer

1. $24,490
2. $24,500
3. $24,000
4. $20,000

Exercise 1.2E.107

Remodeling a Kitchen Marissa's kitchen remodeling cost $18,549, Round the cost to the nearest

1. ten
2. hundred
3. thousand and
4. ten-thousand.

Exercise 1.2E.108

Population The population of China was 1,339,724,852 on November 1, 2010. Round the population to the nearest

1. billion
2. hundred-million; and
3. million.

Answer

1. 1,000,000,000
2. 1,300,000,000
3. 1,340,000,000

Exercise 1.2E.109

Astronomy The average distance between Earth and the sun is 149,597,888 kilometers. Round the distance to the nearest

1. hundred-million
2. ten-million; and
3. million.

Exercise 1.2E.110

Grocery Shopping Hot dogs are sold in packages of 10, but hot dog buns come in packs of eight. What is the smallest number that makes the hot dogs and buns come out even?

Answer

40

Exercise 1.2E.111

Grocery Shopping Paper plates are sold in packages of 12 and party cups come in packs of eight. What is the smallest number that makes the plates and cups come out even?

Writing Exercises

Exercise 1.2E.112

Give an everyday example where it helps to round numbers.

Exercise 1.2E.113

If a number is divisible by 2 and by 3 why is it also divisible by 6?

Exercise 1.2E.114

What is the difference between prime numbers and composite numbers?

Answer

Answers may vary.

Exercise 1.2E.115

Explain in your own words how to find the prime factorization of a composite number, using any method you prefer.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
use place value with whole numbers.			
identify multiples and apply divisibility tests.			
find prime factorizations and least common multiples.			

ⓑ If most of your checks were:

…confidently. Congratulations! You have achieved the objectives in this section. Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific.

…with some help. This must be addressed quickly because topics you do not master become potholes in your road to success. In math, every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no—I don't get it! This is a warning sign and you must not ignore it. You should get help right away or you will quickly be overwhelmed. See your instructor as soon as you can to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 1.2E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.2E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.3: The Language of Algebra and the Order of Operations

Learning Objectives

By the end of this section, you will be able to:

- Use variables and algebraic symbols
- Simplify expressions using the order of operations
- Evaluate an expression
- Identify and combine like terms
- Translate an English phrase to an algebraic expression

Use Variables and Algebraic Symbols

Suppose this year Greg is 20 years old and Alex is 23. You know that Alex is 3 years older than Greg. When Greg was 12, Alex was 15. When Greg is 35, Alex will be 38. No matter what Greg's age is, Alex's age will always be 3 years more, right? In the language of algebra, we say that Greg's age and Alex's age are variables and the 3 is a constant. The ages change ("vary") but the 3 years between them always stays the same ("constant"). Since Greg's age and Alex's age will always differ by 3 years, 3 is the constant. In algebra, we use letters of the alphabet to represent variables. So if we call Greg's age g, then we could use $g+3g+3$ to represent Alex's age. See Table 1.3.1.

Table 1.3.1

Greg's age	Alex's age
12	15
20	23
35	38
g	$g+3$

The letters used to represent these changing ages are called variables. The letters most commonly used for variables are x, y, a, b, and c.

Definition: VARIABLE

A **variable** is a letter that represents a number whose value may change.

Definition: CONSTANT

A **constant** is a number whose value always stays the same.

To write algebraically, we need some operation symbols as well as numbers and variables. There are several types of symbols we will be using.

There are four basic arithmetic **operations: addition, subtraction, multiplication,** and **division**. We'll list the symbols used to indicate these operations below (Table 1.3.2). You'll probably recognize some of them.

Table 1.3.2

Operation	Notation	Say:	The result is...
Addition	$a+b$	a plus b	the sum of a and b
Subtraction	$a-b$	a minus b	the difference of a and b
Multiplication	$a \cdot b, ab, (a)(b), (a)b, a(b)$	a times b	the product of a and b

Operation	Notation	Say:	The result is...
Division	$a \div b, a/b, \frac{a}{b}, b\overline{)a}$	a divided by b	the quotient of a and b, a is called the dividend, and b is called the divisor

We perform these operations on two numbers. When translating from symbolic form to English, or from English to symbolic form, pay attention to the words "of" and "and."

- The *difference* of 9 and 2 means subtract 9 and 2, in other words, 9 minus 2, which we write symbolically as $9-2$.
- The *product* of 4 and 8 means multiply 4 and 8, in other words 4 times 8, which we write symbolically as $4 \cdot 8$.

In algebra, the cross symbol, $\times$, is not used to show multiplication because that symbol may cause confusion. Does $3xy$ mean $3 \times y$ ('three times y') or $3 \cdot x \cdot y$ (three times x times y)? To make it clear, use $\cdot$ or parentheses for multiplication.

When two quantities have the same value, we say they are equal and connect them with an **equal sign**.

EQUALITY SYMBOL

$a = b$ is read "a is equal to b"

The symbol "$=$" is called the **equal sign**.

On the number line, the numbers get larger as they go from left to right. The number line can be used to explain the symbols "$<$" and "$>$".

INEQUALITY

$a < b$ is read "a is less than b"

a is to the left of b on the number line




Figure 1.3.1

$a > b$ is read "a is greater than b"

a is to the right of b on the number line




Figure 1.3.2

The expressions $a < b$ or $a > b$ can be read from left to right or right to left, though in English we usually read from left to right Table 1.3.3. In general, $a < b$ is equivalent to $b > a$. For example $7 < 11$ is equivalent to $11 > 7$. And $a > b$ is equivalent to $b < a$. For example $17 > 4$ is equivalent to $4 < 17$.

Table 1.3.3

Inequality Symbols	Words
$a \neq b$	a is not equal to b
$a < b$	a is less than b
$a \leq b$	a is less than or equal to b
$a > b$	a is greater than b
$a \geq b$	a is greater than or not equal to b

Example 1.3.1

Translate from algebra into English:

1. $17 \leq 26$
2. $8 \neq 17 - 3$
3. $12 > 27 \div 3$
4. $y + 7 < 19$

Solution

1. $17 \leq 26$, 17 is less than or equal to 26
2. $8 \neq 17 - 3$, 8 is not equal to 17 minus 3
3. $12 > 27 \div 3$, 12 is greater than 27 divided by 3
4. $y + 7 < 19$, y plus 7 is less than 19

Try It 1.3.2

Translate from algebra into English:

1. $14 \leq 27$
2. $19 - 2 \neq 8$
3. $12 > 4 \div 2$
4. $x - 7 < 1$

Answer

1. 14 is less than or equal to 27
2. 19 minus 2 is not equal to 8
3. 12 is greater than 4 divided by 2
4. x minus 7 is less than 1

Try It 1.3.3

Translate from algebra into English:

1. $19 \leq 15$
2. $7 = 12 - 5$
3. $15 \div 3 < 8$
4. $y + 3 < 6$

Answer

1. 19 is greater then or equal to 15
2. 7 is equal to 12 minus 5
3. 15 divided by 3 is less than 8
4. y plus 3 is greater than 6

Grouping symbols in algebra are much like the commas, colons, and other punctuation marks in English. They help to make clear which expressions are to be kept together and separate from other expressions. We will introduce three types now.

GROUPING SYMBOLS

Parentheses	()
Brackets	[]
Braces	{}

Here are some examples of expressions that include **grouping symbols**. We will simplify expressions like these later in this section.

$$8(14-8) \qquad 21-3[2+4(9-8)] \qquad 24\div\{13-2[1(6-5)+4]\}$$

What is the difference in English between a phrase and a sentence? A phrase expresses a single thought that is incomplete by itself, but a sentence makes a complete statement. “Running very fast” is a phrase, but “The football player was running very fast” is a sentence. A sentence has a subject and a verb. In algebra, we have expressions and *equations*.

EXPRESSION

An **expression** is a number, a variable, or a combination of numbers and variables using operation symbols.

An **expression** is like an English phrase. Here are some examples of expressions:

Table 1.3.4

Expression	Words	English Phrase
$3+5$	3 plus 5	the sum of three and five
$n-1$	n minus one	the difference of n and one
$6\cdot 7$	6 times 7	the product of six and seven
$\frac{x}{y}$	x divided by y	the quotient of x and y

Notice that the English phrases do not form a complete sentence because the phrase does not have a verb. An **equation** is two expressions linked with an **equal sign**. When you read the words the symbols represent in an equation, you have a complete sentence in English. The equal sign gives the verb.

Definition: EQUATION

An **equation** is two expressions connected by an equal sign.

Here are some examples of equations.

Table 1.3.5

Equation	English Sentence
$3+5=8$	the sum of three and five is equal to eight
$n-1=14$	n minus one equals fourteen
$6\cdot 7=42$	The product of six ans seven is equal to forty-two
$x=53$	x is equal to fifty-three
$y+9=2y-3$	y plus nine is equal to two y minus three

Example 1.3.4

Determine if each is an expression or an equation:

1. $2(x+3)=10$
2. $4(y-1)+1$
3. $x\div 25$
4. $y+8=40$

Solution

1. $2(x+3)=10$. This is an equation — two expressions are connected with an equal sign.

2. $4(y-1)+1$. This is an expression — no equal sign.
3. $x \div 25$. This is an expression — no equal sign.
4. $y+8=40$. This is an equation — two expressions are connected with an equal sign.

Try It 1.3.5

Determine if each is an expression or an equation:

1. $3(x-7)=27$
2. $5(4y-2)-7$

Answer

1. equation
2. expression

Try It 1.3.6

Determine if each is an expression or an equation:

1. $y^3 \div 14$
2. $4x-6=22$

Answer

1. expression
2. equation

Suppose we need to multiply nine factors of 2. We could write this as $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$. This is tedious and it can be hard to keep track of all those 2s, so we use exponents. We write $2 \cdot 2 \cdot 2$ as 2^3 and $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2$ as 2^9. In expressions such as 2^3, the 2 is called the base and the 3 is called the *exponent*. The exponent tells us how many times we need to multiply the base.




Figure 1.3.3

We read 2^3 as "two to the third power" or "two cubed."

We say 2^3 is in exponential notation and $2 \cdot 2 \cdot 2$ is in expanded notation.

EXPONENTIAL NOTATION

a^n means the product of n factors of a.




Figure 1.3.4

The expression a^n is read a to the nth power.

While we read a^n as "a to the nth power," we usually read:

- a^2 "a squared"
- a^3 "a cubed"

We'll see later why a^2 and a^3 have special names.

Table 1.3.6 shows how we read some expressions with exponents.

Table 1.3.6

Expression	In Words

Expression	In Words
7^2	7 to the second power or 7 squared
5^3	5 to the third power or 5 cubed
9^4	9 to the fourth power
12^5	12 to the fifth power

Example 1.3.7

Simplify: 3^4

Solution

$$3^4$$

Expand the expression	$3 \cdot 3 \cdot 3 \cdot 3$
Multiply left to right	$9 \cdot 3 \cdot 3$
Multiply	$27 \cdot 3$
Multiply	81

Try It 1.3.8

Simplify:

1. 5^3
2. 1^7

Answer

1. 125
2. 1

Try It 1.3.9

1. 7^2
2. 0^5

Answer

1. 49
2. 0

Simplify Expressions Using the Order of Operations

To **simplify an expression** means to do all the math possible. For example, to simplify $4 \cdot 2 + 1$ we'd first multiply $4 \cdot 2$ to get 8 and then add the 1 to get 9. A good habit to develop is to work down the page, writing each step of the process below the previous step. The example just described would look like this:

$$4 \cdot 2 + 1$$

$$8 + 1$$

$$9$$

By not using an **equal sign** when you simplify an expression, you may avoid confusing expressions with equations.

SIMPLIFY AN EXPRESSION

To **simplify an expression**, do all operations in the expression.

We've introduced most of the symbols and notation used in algebra, but now we need to clarify the order of operations. Otherwise, expressions may have different meanings, and they may result in different values. For example, consider the expression:

$$4+3\cdot 7$$

If you simplify this expression, what do you get?

Some students say 49,

$$4+3\cdot 7$$

Since $4+3$ gives 7.

$$7\cdot 7$$

And $7\cdot 7$ is 49

$$49$$

Others say 25,

$$4+3\cdot 7$$

Since $3\cdot 7$ is 21.

$$4+21$$

And $21+4$ makes 25.

$$25$$

Imagine the confusion in our banking system if every problem had several different correct answers!

The same expression should give the same result. So mathematicians early on established some guidelines that are called the **Order of Operations**.

PERFORM THE ORDER OF OPERATIONS.

1. Parentheses and Other Grouping Symbols
   - Simplify all expressions inside the parentheses or other grouping symbols, working on the innermost parentheses first.
2. Exponents
   - Simplify all expressions with exponents.
3. Multiplication and Division
   - Perform all multiplication and division in order from left to right. These operations have equal priority.
4. Addition and Subtraction
   - Perform all addition and subtraction in order from left to right. These operations have equal priority.

Note

Doing the Manipulative Mathematics activity "Game of 24" will give you practice using the order of operations.

Students often ask, "How will I remember the order?" Here is a way to help you remember: Take the first letter of each key word and substitute the silly phrase: "Please Excuse My Dear Aunt Sally."

Parentheses	**P**lease
Exponents	**E**xcuse
Multiplication **D**ivision	**M**y **D**ear
Addition **S**ubtraction	**A**unt **S**ally

It's good that "**M**y **D**ear" goes together, as this reminds us that **m**ultiplication and **d**ivision have equal priority. We do not always do multiplication before division or always do division before multiplication. We do them in order from left to right.

Similarly, "**A**unt **S**ally" goes together and so reminds us that **a**ddition and **s**ubtraction also have equal priority and we do them in order from left to right.

Let's try an example.

Example 1.3.10

Simplify:

1. $4+3\cdot 7$
2. $(4+3)\cdot 7$

Solution

1.

	$4+3\cdot 7$
Are there any **p**arentheses? No.	
Are there any **e**xponents? No.	
Is there any **m**ultiplication or **d**ivision? Yes.	
Multiply first.	$4+3\cdot 7$
Add.	$4+21$
	25

2.

	$(4+3)\cdot 7$
Are there any **p**arentheses? Yes.	$(4+3)\cdot 7$
Simplify inside the parentheses.	$(7)7$
Are there any **e**xponents? No.	
Is there any **m**ultiplication or **d**ivision? Yes.	
Multiply.	49

Try It 1.3.11

Simplify:

1. $12-5\cdot 2$
2. $(12-5)\cdot 2$

Answer

1. 2
2. 14

Try It 1.3.12

Simplify:

1. $8+3\cdot 9$
2. $(8+3)\cdot 9$

Answer

1. 35
2. 99

Example 1.3.13

Simplify: $18\div 6+4(5-2)$

Solution

Parentheses? Yes, subtract first.	$18\div 6+4(5-2)$ $18\div 6+4(3)$
Exponents? No.	
Multiplication or division? Yes.	$18\div 6+4(3)$
Divide first because we multiply and divide left to right.	$3+4(3)$
Any other multiplication or division? Yes.	
Multiply.	$3+12$
Any other multiplication or division? No.	
Any addition or subtraction? Yes.	15

Try It 1.3.14

Simplify: $30\div 5+10(3-2)$

Answer

16

Try It 1.3.15

Simplify: $70\div 10+4(6-2)$

Answer

23

When there are multiple grouping symbols, we simplify the innermost parentheses first and work outward.

Example 1.3.16

Simplify: $5+2^3+3[6-3(4-2)]$.

Solution

	$5+2^3+3[6-3(4-2)]$

Are there any parentheses (or other grouping symbol)? Yes.	
Focus on the parentheses that are inside the brackets.	$5+2^3+3[6-3(4-2)]$
Subtract.	$5+2^3+3[6-3(2)]$
Continue inside the brackets and multiply.	$5+2^3+3[6-6]$
Continue inside the brackets and subtract.	$5+2^3+3[0]$
The expression inside the brackets requires no further simplification.	
Are there any exponents? Yes.	$5+2^3+3[0]$
Simplify exponents.	$5+8+3[0]$
Is there any multiplication or division? Yes.	
Multiply.	$5+8+0$
Is there any addition or subtraction? Yes.	
Add.	$13+0$
Add.	13

Try It 1.3.17

Simplify: $9+5^3-[4(9+3)]$.

Answer

86

Try It 1.3.18

Simplify: $7^2-2[4(5+1)]$.

Answer

1

Evaluate an Expression

In the last few examples, we simplified expressions using the order of operations. Now we'll evaluate some expressions—again following the order of operations. To **evaluate an expression** means to find the value of the expression when the variable is replaced by a given number.

EVALUATE AN EXPRESSION

To **evaluate an expression** means to find the value of the expression when the variable is replaced by a given number.

To evaluate an expression, substitute that number for the variable in the expression and then simplify the expression.

Example 1.3.19

Evaluate $7x-4$, when

1. $x=5$
2. $x=1$

Solution

1.

when $x = 5$	$7x - 4$
	$7(5) - 4$
Multiply.	$35 - 4$
Subtract.	31

2.

when $x = 1$	$7x - 4$
	$7(1) - 4$
Multiply.	$7 - 4$
Subtract.	3

Try It 1.3.20

Evaluate $8x - 3$, when

1. $x = 2$
2. $x = 1$

Answer

1. 13
2. 5

Try It 1.3.21

Evaluate $4y - 4$, when

1. $y = 3$
2. $y = 5$

Answer

1. 8
2. 16

Example 1.3.22

Evaluate $x = 4$, when

1. x^2
2. 3^x

Solution

1.

	x^2
Replace x with 4.	$(4)^2$
Use definition of exponent.	$4 \cdot 4$

Simplify.	16

2.

	3^x
Replace x with 4.	3^4
Use definition of exponent.	$3 \cdot 3 \cdot 3 \cdot 3$
Simplify.	81

Try It 1.3.23

Evaluate $x = 3$, when

1. x^2
2. 4^x

Answer

1. 9
2. 64

Try It 1.3.24

Evaluate $x = 6$, when

1. x^3
2. 2^x

Answer

1. 216
2. 64

Example 1.3.25

Evaluate $2x^2 + 3x + 8$ when $x = 4$.

Solution

	$2x^2 + 3x + 8$
Substitute $x = 4$.	$2x^2 + 3x + 8$ $2(4)^2 + 3(4) + 8$
Follow the order of operations.	$2(16) + 3(4) + 8$
	$32 + 12 + 8$
	52

Try It 1.3.26

Evaluate $3x^2 + 4x + 1$ when $x = 3$.

Answer

40

Try It 1.3.27

Evaluate $6x^2 - 4x - 7$ when $x = 2$.

Answer

9

Identify and Combine Like Terms

Algebraic expressions are made up of terms. A **term** is a constant, or the product of a constant and one or more variables.

TERM

A **term** is a constant, or the product of a constant and one or more variables.

Examples of terms are $7, y, 5x^2, 9a$, and b^5.

The constant that multiplies the variable is called the **coefficient**.

COEFFICIENT

The **coefficient** of a term is the constant that multiplies the variable in a term.

Think of the coefficient as the number in front of the variable. The coefficient of the term $3x$ is 3. When we write x, the coefficient is 1, since $x = 1 \cdot x$.

Example 1.3.28

Identify the coefficient of each term:

1. $14y$
2. $15x^2$
3. a

Solution

1. The coefficient of $14y$ is 14
2. The coefficient of $15x^2$ is 15
3. The coefficient of a is 1 since $a = 1a$.

Try It 1.3.29

Identify the coefficient of each term:

1. $17x$
2. $41b^2$
3. z

Answer

1. 14
2. 41
3. 1

Try It 1.3.30

Identify the coefficient of each term:

1. $9p$
2. $13a^2$
3. y^3

Answer

1. 9
2. 13
3. 1

Some terms share common traits. Look at the following 6 terms. Which ones seem to have traits in common?

$$5x \quad 7 \quad n^2 \quad 4 \quad 3x \quad 9n^2$$

The 7 and the 4 are both constant terms.

The $5x$ and the $3x$ are both terms with x.

The n^2 and the $9n^2$ are both terms with n^2.

When two terms are constants or have the same variable and exponent, we say they are **like terms**.

- 7 and 4 are like terms.
- $5x$ and $3x$ are like terms.
- x^2 and $9x^2$ are like terms.

LIKE TERMS

Terms that are either constants or have the same variables raised to the same powers are called **like terms**.

Example 1.3.31

Identify the like terms: $y^3, 7x^2, 14, 23, 4y^3, 9x, 5x^2$

Solution

y^3 and $4y^3$ are like terms because both have y^3; the variable and the exponent match.

$7x^2$ and $5x^2$ are like terms because both have x^2; the variable and the exponent match.

14 and 23 are like terms because both are constants.

There is no other term like $9x$.

Try It 1.3.32

Identify the like terms: $9, 2x^3, y^2, 8x^3, 15, 9y, 11y^2$

Answer

9 and 15, y^2 and $11y^2$, $2x^3$ and $8x^3$

Try It 1.3.33

Identify the like terms: $4x^3, 8x^2, 19, 3x^3, 24, 6x^3$

Answer

19 and 24, $8x^2$ and $3x^2$, $4x^3$ and $6x^3$

Adding or subtracting terms forms an expression. In the expression $2x^2+3x+8$, from Example, the three terms are $2x^2$, $3x$, and 8.

Example 1.3.34

Identify the terms in each expression.

1. $9x^2+7x+12$
2. $8x+3y$

Solution

1. The terms of $9x^2+7x+12$ are $9x^2$, $7x$, and 12.
2. The terms of $8x+3y$ are $8x$ and $3y$.

Try It 1.3.35

Identify the terms in the expression $4x^2+5x+17$.

Answer

$4x^2, 5x, 17$

Try It 1.3.36

Identify the terms in the expression $5x+2y$.

Answer

$5x, 2y$

If there are like terms in an expression, you can simplify the expression by combining the like terms. What do you think $4x+7x+x$ would simplify to? If you thought $12x$, you would be right!

$$\begin{array}{c} 4x+7x+x \\ x+x+x+x \quad +x+x+x+x+x+x+x \quad +x \\ 12x \end{array} \tag{1.3.1}$$

Add the coefficients and keep the same variable. It doesn't matter what x is—if you have 4 of something and add 7 more of the same thing and then add 1 more, the result is 12 of them. For example, 4 oranges plus 7 oranges plus 1 orange is 12 oranges. We will discuss the mathematical properties behind this later.

Simplify: $4x+7x+x$

Add the coefficients. $12x$

Example 1.3.37: How To Combine Like Terms

Simplify: $2x^2+3x+7+x^2+4x+5$

Solution

Step 1. Identify the like terms.	$2x^2+3x+7+x^2+4x+5$ $2x^2+3x+7+x^2+4x+5$
Step 2. Rearrange the expression so the like terms are together.	$2x^2+x^2+3x+4x+7+5$

Step 3. Combine like terms.	$3x^2 + 7x + 12$

Try It 1.3.38

Simplify: $3x^2+7x+9+7x^2+9x+8$.

Answer

$10x^2+16x+17$

Try It 1.3.39

Simplify: $4y^2+5y+2+8y^2+4y+5$.

Answer

$12y^2+9y+7$

COMBINE LIKE TERMS.

1. Identify like terms.
2. Rearrange the expression so like terms are together.
3. Add or subtract the coefficients and keep the same variable for each group of like terms.

Translate an English Phrase to an Algebraic Expression

In the last section, we listed many operation symbols that are used in algebra, then we translated expressions and equations into English phrases and sentences. Now we'll reverse the process. We'll translate English phrases into algebraic expressions. The symbols and variables we've talked about will help us do that. Table 1.3.7 summarizes them.

Table 1.3.7

Operation	Phrase	Expression	
Addition	a plus b the sum of a and b a increased by b b more than a the total of a and b b added to a	$a+b$	(1.3.2)
Subtraction	a minus b the difference of a and b a decreased by b b less than a b subtracted from a	$a-b$	(1.3.3)
Multiplication	a times b the product of a and b twice a	$a \cdot b, ab, a(b), (a)(b)$ $2a$	(1.3.4) (1.3.5)
Division	a divided by b the quotient of a and b the ratio of a and b b divided into a	$a \div b, a/b, \frac{a}{b}, b\overline{)a}$	(1.3.6)

Look closely at these phrases using the four operations:

the **sum** *of a and b*

the **difference** *of a and b*

the **product** *of a and b*

the **quotient** *of a and b*

Figure 1.3.5

Each phrase tells us to operate on two numbers. Look for the words *of* and *and* to find the numbers.

Example 1.3.40

Translate each English phrase into an algebraic expression:

1. the difference of $17x$ and 5
2. the quotient of $10x^2$ and 7.

Solution

1. The key word is *difference*, which tells us the operation is subtraction. Look for the words *of* and *and* to find the numbers to subtract.

   the *difference of* $17x$ *and* 5

   $17x$ minus 5

   $17x - 5$

2. The key word is "quotient," which tells us the operation is division.

   the *quotient of* $10x^2$ *and* 7

   divide $10x^2$ by 7

   $10x^2 \div 7$

This can also be written $10x^2/7$ or $\frac{10x^2}{7}$.

Try It 1.3.41

Translate each English phrase into an algebraic expression:

1. the difference of $14x^2$ and 13
2. the quotient of $12x$ and 2.

Answer

1. $14x^2 - 13$
2. $12x \div 2$

Try It 1.3.42

Translate each English phrase into an algebraic expression:

1. the sum of $17y^2$ and 19
2. the product of 7 and y.

Answer

1. $17y^2 + 19$
2. $7y$

How old will you be in eight years? What age is eight more years than your age now? Did you add 8 to your present age? Eight "more than" means 8 added to your present age. How old were you seven years ago? This is 7 years less than your age now. You subtract 7 from your present age. Seven "less than" means 7 subtracted from your present age.

Example 1.3.43

Translate the English phrase into an algebraic expression:

1. Seventeen more than y
2. Nine less than $9x^2$.

Solution

1. The key words are *more than.* They tell us the operation is addition. *More than* means "added to."

$$\begin{aligned} &\text{Seventeen more than } y \\ &\text{Seventeen added to } y \\ &y+17 \end{aligned}$$

2. The key words are *less than.* They tell us to subtract. *Less than* means "subtracted from."

$$\begin{aligned} &\text{Nine less than } 9x^2 \\ &\text{Nine subtracted from } 9x^2 \\ &9x^2-9 \end{aligned}$$

Try It 1.3.44

Translate the English phrase into an algebraic expression:

1. Eleven more than x
2. Fourteen less than $11a$.

Answer

1. $x+11$
2. $11a-14$

Try It 1.3.45

Translate the English phrase into an algebraic expression:

1. 13 more than z
2. 18 less than $8x$.

Answer

1. $z+13$
2. $8x-18$

Example 1.3.46

Translate the English phrase into an algebraic expression:

1. five times the sum of m and n
2. the sum of five times m and n.

Solution

There are two operation words—*times* tells us to multiply and *sum* tells us to add.
1. Because we are multiplying 5 times the sum we need parentheses around the sum of m and n, $(m+n)$. This forces us to determine the sum first. (Remember the order of operations.)

$$\begin{aligned} &\text{five times the sum of } m \text{ and } n \\ &5(m+n) \end{aligned} \tag{1.3.7}$$

2. To take a sum, we look for the words “of” and “and” to see what is being added. Here we are taking the sum *of* five times m and \(n\.)

$$\begin{array}{c} \text{the sum of five times } m \text{ and } n \\ 5m+n \end{array} \tag{1.3.8}$$

Try It 1.3.47

Translate the English phrase into an algebraic expression:

1. four times the sum of p and q
2. the sum of four times p and q.

Answer

1. $4(p+q)$
2. $4p+q$

Try It 1.3.48

Translate the English phrase into an algebraic expression:

1. the difference of two times x and 8,
2. two times the difference of x and 8.

Answer

1. $2x-8$
2. $2(x-8)$

Later in this course, we’ll apply our skills in algebra to solving applications. The first step will be to translate an English phrase to an algebraic expression. We’ll see how to do this in the next two examples.

Example 1.3.49

The length of a rectangle is 6 less than the width. Let w represent the width of the rectangle. Write an expression for the length of the rectangle.

Solution

$$\begin{array}{ll} \text{Write a phrase about the length of the rectangle.} & \text{6 less than the width} \\ \text{Substitute } w \text{ for "the width."} & \text{6 less then w} \\ \text{Rewrite "less than" as "subtracted from."} & \text{6 subtracted from w} \\ \text{Translate the phrase into algebra.} & w-6 \end{array} \tag{1.3.9}$$

Try It 1.3.50

The length of a rectangle is 7 less than the width. Let w represent the width of the rectangle. Write an expression for the length of the rectangle.

Answer

$w-7$

Try It 1.3.51

The width of a rectangle is 6 less than the length. Let l represent the length of the rectangle. Write an expression for the width of the rectangle.

Answer

$l-6$

Example 1.3.52

June has dimes and quarters in her purse. The number of dimes is three less than four times the number of quarters. Let q represent the number of quarters. Write an expression for the number of dimes.

Solution

Write the phrase about the number of dimes.	three less than four times the number of quarters
Substitute q for the number of quarters.	3 less than 4 times q
Translate "4 times q ."	3 less than 4q
Translate the phrase into algebra.	4q - 3

(1.3.10)

Try It 1.3.53

Geoffrey has dimes and quarters in his pocket. The number of dimes is eight less than four times the number of quarters. Let q represent the number of quarters. Write an expression for the number of dimes.

Answer

$4q-8$

Try It 1.3.54

Lauren has dimes and nickels in her purse. The number of dimes is three more than seven times the number of nickels. Let n represent the number of nickels. Write an expression for the number of dimes.

Answer

$7n+3$

Key Concepts

- **Notation** — **The result is…**
  - $a+b$ — the sum of a and b
  - $a-b$ — the difference of a and b
  - $a \cdot b, ab, (a)(b)(a)b, a(b)$ — the product of a and b
  - $a \div b, a/b, \frac{a}{b}, b\overline{)a}$ — the quotient of a and b
- **Inequality**
  - $a < b$ is read "a is less than b" — a is to the left of b on the number line
  - $a > b$ is read "a is greater than b" — a is to the right of b on the number line
- **Inequality Symbols** — **Words**
  - $a \neq b$ — a is not equal to b
  - $a < b$ — a is less than b
  - $a \leq b$ — a is less than or equal to b
  - $a > b$ — a is greater than b
  - $a \geq b$ — a is greater than or equal to b

- **Grouping Symbols**
  - Parentheses ()
  - Brackets []
  - Braces {}
- **Exponential Notation**
  - a^n means the product of n factors of a. The expression a^n is read a to the nth power.
- **Order of Operations:** When simplifying mathematical expressions perform the operations in the following order:
  1. Parentheses and other Grouping Symbols: Simplify all expressions inside the parentheses or other grouping symbols, working on the innermost parentheses first.
  2. Exponents: Simplify all expressions with exponents.
  3. Multiplication and Division: Perform all multiplication and division in order from left to right. These operations have equal priority.
  4. Addition and Subtraction: Perform all addition and subtraction in order from left to right. These operations have equal priority.
- **Combine Like Terms**
  1. Identify like terms.
  2. Rearrange the expression so like terms are together.
  3. Add or subtract the coefficients and keep the same variable for each group of like terms.

Glossary

coefficient

The coefficient of a term is the constant that multiplies the variable in a term.

constant

A constant is a number whose value always stays the same.

equality symbol

The symbol "$=$" is called the equal sign. We read $a = b$ as "a is equal to b."

equation

An equation is two expressions connected by an equal sign.

evaluate an expression

To evaluate an expression means to find the value of the expression when the variable is replaced by a given number.

expression

An expression is a number, a variable, or a combination of numbers and variables using operation symbols.

like terms

Terms that are either constants or have the same variables raised to the same powers are called like terms.

simplify an expression

To simplify an expression, do all operations in the expression.

term

A term is a constant or the product of a constant and one or more variables.

variable

A variable is a letter that represents a number whose value may change.

This page titled 1.3: The Language of Algebra and the Order of Operations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.3: Use the Language of Algebra** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.3E: Exercises

Practice Makes Perfect

Use Variables and Algebraic Symbols

In the following exercises, translate from algebra to English.

Exercise 1.3E.55

$16 - 9$

Answer

16 minus 9, the difference of sixteen and nine

Exercise 1.3E.56

$3 \cdot 9$

Exercise 1.3E.57

$28 \div 4$

Answer

28 divided by 4, the quotient of twenty-eight and four

Exercise 1.3E.58

$x + 11$

Exercise 1.3E.59

$(2)(7)$

Answer

2 times 7, the product of two and seven

Exercise 1.3E.60

$(4)(8)$

Exercise 1.3E.61

$14 < 21$

Answer

fourteen is less than twenty-one

Exercise 1.3E.62

$17 < 35$

Exercise 1.3E.63

$36 \geq 19$

Answer

thirty-six is greater than or equal to nineteen

Exercise 1.3E.64

$6n = 36$

Exercise 1.3E.65

$y - 1 > 6$

Answer

y minus 1 is greater than 6, the difference of y and one is greater than six

Exercise 1.3E.66

$y - 4 > 8$

Exercise 1.3E.67

$2 \leq 18 \div 6$

Answer

2 is less than or equal to 18 divided by 6; 2 is less than or equal to the quotient of eighteen and six

Exercise 1.3E.68

$a \neq 1 \cdot 12$

In the following exercises, determine if each is an expression or an equation.

Exercise 1.3E.69

$9 \cdot 6 = 54$

Answer

equation

Exercise 1.3E.70

$7 \cdot 9 = 63$

Exercise 1.3E.71

$5 \cdot 4 + 3$

Answer

expression

Exercise 1.3E.72

$x+7$

Exercise 1.3E.73

$x+9$

Answer

expression

Exercise 1.3E.74

$y-5=25$

Simplify Expressions Using the Order of Operations

In the following exercises, simplify each expression.

Exercise 1.3E.75

5^3

Answer

125

Exercise 1.3E.76

8^3

Exercise 1.3E.77

2^8

Answer

256

Exercise 1.3E.78

10^5

In the following exercises, simplify using the order of operations.

Exercise 1.3E.79

1. $3+8\cdot 5$
2. $(3+8)\cdot 5$

Answer

1. 43
2. 55

Exercise 1.3E.80

1. $2+6\cdot 3$
2. $(2+6)\cdot 3$

Exercise 1.3E.81

$2^3-12\div(9-5)$

Answer

5

Exercise 1.3E.82

$3^2-18\div(11-5)$

Exercise 1.3E.83

$3\cdot 8+5\cdot 2$

Answer

34

Exercise 1.3E.84

$4\cdot 7+3\cdot 5$

Exercise 1.3E.85

$2+8(6+1)$

Answer

58

Exercise 1.3E.86

$4+6(3+6)$

Exercise 1.3E.87

$4\cdot 12/8$

Answer

6

Exercise 1.3E.88

$2\cdot 36/6$

Exercise 1.3E.89

$(6+10)\div(2+2)$

Answer

4

Exercise 1.3E.90

$(9+12)\div(3+4)$

Exercise 1.3E.91

$20\div4+6\cdot5$

Answer

35

Exercise 1.3E.92

$33\div3+8\cdot2$

Exercise 1.3E.93

3^2+7^2

Answer

58

Exercise 1.3E.94

$(3+7)^2$

Exercise 1.3E.95

$3(1+9\cdot6)-4^2$

Answer

149

Exercise 1.3E.96

$5(2+8\cdot4)-7^2$

Exercise 1.3E.97

$2[1+3(10-2)]$

Answer

50

Exercise 1.3E.98

$5[2+4(3-2)]$

Evaluate an Expression

In the following exercises, evaluate the following expressions.

Exercise 1.3E.99

$7x+8$ when $x=2$

Answer

22

Exercise 1.3E.100

$8x-6$ when $x=7$

Exercise 1.3E.101

x^2 when $x=12$

Answer

144

Exercise 1.3E.102

x^3 when $x=5$

Exercise 1.3E.103

x^5 when $x=2$

Answer

32

Exercise 1.3E.104

4^x when $x=2$

Exercise 1.3E.105

x^2+3x-7 when $x=4$

Answer

21

Exercise 1.3E.106

$6x+3y-9$ when $x=10, y=7$

Answer

9

Exercise 1.3E.107

$(x+y)^2$ when $x=6, y=9$

Exercise 1.3E.108

a^2+b^2 when $a=3, b=8$

Answer

73

Exercise 1.3E.109

r^2-s^2 when $r=12, s=5$

Exercise 1.3E.110

$2l+2w$ when $l=15, w=12$

Answer

54

Exercise 1.3E.111

$2l+2w$ when $l=18, w=14$

Simplify Expressions by Combining Like Terms

In the following exercises, identify the coefficient of each term.

Exercise 1.3E.112

$8a$

Answer

8

Exercise 1.3E.113

$13m$

Exercise 1.3E.114

$5r^2$

Answer

5

Exercise 1.3E.115

$6x^3$

In the following exercises, identify the like terms.

Exercise 1.3E.116

$x^3, 8x, 14, 8y, 5, 8x^3$

Answer

x^3 and $8x^3$, 14 and 5

Exercise 1.3E.117

$6z, 3w^2, 1, 6z^2, 4z, w^2$

Exercise 1.3E.118

$9a, a^2, 16, 16b^2, 4, 9b^2$

Answer

16 and 4, $16b^2$ and $9b^2$

Exercise 1.3E.119

$3, 25r^2, 10s, 10r, 4r^2, 3s$

In the following exercises, identify the terms in each expression.

Exercise 1.3E.120

$15x^2 + 6x + 2$

Answer

$15x^2, 6x, 2$

Exercise 1.3E.121

$11x^2 + 8x + 5$

Exercise 1.3E.122

$10y^3 + y + 2$

Answer

$10y^3, y, 2$

Exercise 1.3E.123

$9y^3 + y + 5$

In the following exercises, simplify the following expressions by combining like terms.

Exercise 1.3E.124

$10x + 3x$

Answer

$13x$

Exercise 1.3E.125

$15x + 4x$

Exercise 1.3E.126

$4c + 2c + c$

Answer

$7c$

Exercise 1.3E.127

$6y + 4y + y$

Exercise 1.3E.128

$7u + 2 + 3u + 1$

Answer

$10u + 3$

Exercise 1.3E.129

$8d + 6 + 2d + 5$

Exercise 1.3E.130

$10a + 7 + 5a - 2 + 7a - 4$

Answer

$22a + 1$

Exercise 1.3E.131

$7c + 4 + 6c - 3 + 9c - 1$

Exercise 1.3E.132

$3x^2 + 12x + 11 + 14x^2 + 8x + 5$

Answer

$17x^2 + 20x + 16$

Exercise 1.3E.133

$5b^2+9b+10+2b^2+3b-4$

Translate an English Phrase to an Algebraic Expression

In the following exercises, translate the phrases into algebraic expressions.

Exercise 1.3E.134

the difference of 14 and 9

Answer

$14-9$

Exercise 1.3E.135

the difference of 19 and 8

Exercise 1.3E.136

the product of 9 and 7

Answer

$9 \cdot 7$

Exercise 1.3E.137

the product of 8 and 7

Exercise 1.3E.138

the quotient of 36 and 9

Answer

$36 \div 9$

Exercise 1.3E.139

the quotient of 42 and 7

Exercise 1.3E.140

the sum of $8x$ and $3x$

Answer

$8x+3x$

Exercise 1.3E.141

the sum of $13x$ and $3x$

Exercise 1.3E.142

the quotient of y and 3

Answer

$\frac{y}{3}$

Exercise 1.3E.143

the quotient of y and 8

Exercise 1.3E.144

eight times the difference of y and nine

Answer

$8(y-9)$

Exercise 1.3E.145

seven times the difference of y and one

Exercise 1.3E.146

Eric has rock and classical CDs in his car. The number of rock CDs is 3 more than the number of classical CDs. Let c represent the number of classical CDs. Write an expression for the number of rock CDs.

Answer

$c+3$

Exercise 1.3E.147

The number of girls in a second-grade class is 4 less than the number of boys. Let b represent the number of boys. Write an expression for the number of girls.

Exercise 1.3E.148

Greg has nickels and pennies in his pocket. The number of pennies is seven less than twice the number of nickels. Let n represent the number of nickels. Write an expression for the number of pennies.

Answer

$2n-7$

Exercise 1.3E.149

Jeannette has $5 and $10 bills in her wallet. The number of fives is three more than six times the number of tens. Let t represent the number of tens. Write an expression for the number of fives.

Everyday Math

Exercise 1.3E.150

Car insurance Justin's car insurance has a $750 deductible per incident. This means that he pays $750 and his insurance company will pay all costs beyond $750. If Justin files a claim for $2,100.

1. how much will he pay?
2. how much will his insurance company pay?

Answer

1. $750
2. $1,350

Exercise 1.3E.151

Home insurance Armando's home insurance has a $2,500 deductible per incident. This means that he pays $2,500 and the insurance company will pay all costs beyond $2,500. If Armando files a claim for $19,400.

1. how much will he pay?
2. how much will the insurance company pay?

Writing Exercises

Exercise 1.3E.152

Explain the difference between an expression and an equation.

Answer

Answers may vary

Exercise 1.3E.153

Why is it important to use the order of operations to simplify an expression?

Exercise 1.3E.154

Explain how you identify the like terms in the expression $8a^2+4a+9-a^2-1$

Answer

Answers may vary

Exercise 1.3E.155

Explain the difference between the phrases "4 times the sum of x and y" and "the sum of 4 times x and y."

Self Check

ⓐ Use this checklist to evaluate your mastery of the objectives of this section.

the *quotient of* $10x^2$ *and* 7
divide $10x^2$ by 7
$10x^2 \div 7$

ⓑ After reviewing this checklist, what will you do to become confident for all objectives?

More Exponent Practice

Exercise $1.3E.156$

compute the exact value of the given exponential expression

1. 0^1	5. 4^1	9. -3^2	13. $(-5)^2$	17. $(-6)^2$	21. $-(-5)^2$
2. 1^3	6. 5^2	10. -3^3	14. $(-2)^4$	18. $(-4)^2$	22. $-(-3)^4$
3. 3^2	7. 4^3	11. -4^2	15. $(-3)^3$	19. $(-5)^3$	23. $-(-4)^3$
4. 3^4	8. 4^2	12. -2^4	16. $(-2)^5$	20. $(-4)^3$	24. $-(-2)^5$

Answers to odd problems for E.156

1. 0	5. 4	9. -9	13. 25	17. 36	21. -25
3. 9	7. 64	11. -64	15. -27	19. -125	23. 64

More Order of Operations Practice

Exercise $1.3E.157$

Compute the exact value of the given expression.

1. $9 - 1(-7)$	7. $10 + 12(-5)$	13. $15 \div 1 \cdot 3$	19. $10 - 72 \div 6 \cdot 3+8$
2. $85 - 8(9)$	8. $4 + 12(4)$	14. $18 \div 6 \cdot 1$	20. $8 - 120 \div 5 \cdot 6+7$
3. $3 + 9(4)$	9. $30 \cdot 5 \div 3$	15. $40 \div 5 \cdot 4$	21. $3 - 24 \div 4 \cdot 3+4$
4. $6 + 7(-1)$	10. $72 \cdot 6 \div 4$	16. $30 \div 6 \cdot 5$	22. $4 - 40 \div 5 \cdot 4+9$
5. $-2 - 3(-5)$	11. $32 \div 4 \cdot 4$		23. $2+6 \div 1 \cdot 6 - 1$
6. $64 - 7(7)$	12. $64 \div 4 \cdot 4$		24. $1 + 12 \div 2 \cdot 2 - 6$

Answers to odd problems for E.157

1. 16	7. -50	13. 45	19. -18
3. 39	9. 50	15. -216	21. -11
5. 13	11. 32		23. 37

More Practice Changing English Expressions into Algebraic Expressions

Exercise $1.3E.158$

Translate the following phrases into algebraic expressions.

1. a. the difference of $5x^2$ and $6xy$
   b. the quotient of $6y^2$ and $5x$
   c. Twenty-one more than y^2
   d. $6x$ less than $81x^2$
2. a. the difference of $17x^2$ and $17x^2$ and $5xy$
   b. the quotient of $8y^3$ and $3x$
   c. Eighteen more than a^2;
   d. $11b$ less than $100b^2$
3. a. the sum of $4ab^2$ and $3a^2b$
   b. the product of $4y^2$ and $5x$
   c. Fifteen more than m
   d. $9x$ less than $121x^2$
4. a. the sum of $3x^2y$ and $7xy^2$
   b. The product of $6xy^2$ and $4z$
   c. Twelve more than $3x^2$
   d. $7x^2$ less than $63x^3$

Answers to odd problems for E.158 #1-4

1a. $5x^2-6xy$ b. $\frac{6y^2}{5x}$ c. y^2+21 d. $81x^2-6x$ 3a. $4ab^2+3a^2b$ b. $20xy^2$ c. $m+15$ d. $121x^2-9x$.

Translate the following phrases into algebraic expressions.

5. a. four times the difference of y and six
   b. the difference of four times y and 6
6. a. five times the difference of y and two
   b. the difference of five times y and 2
7. a. five times the sum of $3x$ and y
   b. the sum of five times $3x$ and y
8. a. eleven times the sum of $4x^2$ and $5x$
   b. the sum of eleven times $4x^2$ and $5x$

Answers to odd problems for E.158 #5-8

5 a. $4(y-6)$ b. $4y-6$ 7 a. $5(3x+y)$ b. $5(3x)+y$.

Translate the following phrases into algebraic expressions.

9. The length of a rectangle is 5 inches less than the width. Let w represent the width of the rectangle. Write an expression for the length of the rectangle.

10. The width of a rectangle is 2 meters greater than the length. Let L represent the length of the rectangle. Write an expression for the width of the rectangle

11. A collection contains jazz and classical CDs. The number of jazz CDs is 3 more than the number of classical CDs. Let c represent the number of classical CDs. Write an expression for the number of jazz CDs.

12. The number of girls in a second-grade class is 4 less than the number of boys. Let b represent the number of boys. Write an expression for the number of girls.

13. A playlist contains rock and country songs. The number of rock songs is 14 more than twice the number of country songs. Let c represent the number of country songs. Write an expression for the number of rock songs.

14. The number of women in a Statistics class is 8 less than twice the number of men. Let m represent the number of men. Write an expression for the number of women.

15. Greg has nickels and pennies in his pocket. The number of pennies is seven less than three times the number of nickels. Let n represent the number of nickels. Write an expression for the number of pennies.

16. Greg has nickels and pennies in his pocket. The number of pennies is four more than twice the number of nickels. Let n represent the number of nickels. Write an expression for the number of pennies.

Answers to odd problems for E.158 #9-16

9. $w-5$ 11. $c+3$ 13. $2c+14$ 15. $3n-7$.

This page titled 1.3E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.3E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **1.2E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/intermediate-algebra-2e.

1.4: Add and Subtract Integers

Learning Objectives

- Use negatives and opposites
- Simplify: expressions with absolute value
- Add integers
- Subtract integers

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapter, **Integers**.

Use Negatives and Opposites

Our work so far has only included the counting numbers and the whole numbers. But if you have ever experienced a temperature below zero or accidentally overdrawn your checking account, you are already familiar with negative numbers. **Negative numbers** are numbers less than 0. The **negative numbers** are to the left of zero on the number line. See Figure 1.4.1.




Figure 1.4.1. The number line shows the location of positive and negative numbers.

The arrows on the ends of the number line indicate that the numbers keep going forever. There is no biggest positive number, and there is no smallest negative number.

Is zero a positive or a negative number? Numbers larger than zero are positive, and numbers smaller than zero are negative. Zero is neither positive nor negative.

Consider how numbers are ordered on the number line. Going from left to right, the numbers increase in value. Going from right to left, the numbers decrease in value. See Figure 1.4.2.




Figure 1.4.2. The numbers on a number line increase in value going from left to right and decrease in value going from right to left.

Doing the Manipulative Mathematics activity “Number Line-part 2” will help you develop a better understanding of integers.

Remember that we use the notation:

$a < b$ (read “a is less than b”) when a is to the left of b on the number line.

$a > b$ (read “a is greater than b”) when a is to the right of b on the number line.

Now we need to extend the number line which showed the whole numbers to include negative numbers, too. The numbers marked by points in Figure 1.4.3. are called the integers. The integers are the numbers $\ldots -3, -2, -1, 0, 1, 2, 3 \ldots$




Figure 1.4.3. All the marked numbers are called *integers*.

Example 1.4.1

Order each of the following pairs of numbers, using $<$ or $>$:

1. 14 ___ 6
2. -1 ___ 9
3. -1 ___ -4
4. 2 ___ -20

Solution:

It may be helpful to refer to the number line shown.




Figure 1.4.4

1. $14 \_\_\_ 6$

   $14 > 6$ 14 is to the right of 6 on the number line.

2. $-1 \_\_\_ 9$

   $-1 < 9$ -1 is to the left of 9 on the number line.

3. $-1 \_\_\_ -4$

   $-1 > -4$ -1 is to the right of -4 on the number line.

4. $2 \_\_\_ -20$

   $2 > -20$ 2 is to the right of -20 on the number line.

Try It 1.4.1

Order each of the following pairs of numbers, using $<$ or $>$:

1. $15 \_\_\_ 7$
2. $-2 \_\_\_ 5$
3. $-3 \_\_\_ -7$
4. $5 \_\_\_ -17$

Answer

1. $<$
2. $>$
3. $<$
4. $>$

Try It 1.4.2

Order each of the following pairs of numbers, using $<$ or $>$:

1. $8 \_\_\_ 13$
2. $3 \_\_\_ -4$
3. $-5 \_\_\_ -2$
4. $9 \_\_\_ -21$

Answer

1. $<$
2. $>$
3. $<$
4. $>$

You may have noticed that, on the **number line**, the negative numbers are a mirror image of the positive numbers, with zero in the middle. Because the numbers 2 and -2 are the same distance from zero, they are called **opposites**. The opposite of 2 is -2, and the opposite of -2 is 2.

OPPOSITE

The **opposite** of a number is the number that is the same distance from zero on the number line but on the opposite side of zero.

Figure 1.4.5 illustrates the definition.



Figure 1.4.5: The opposite of 3 is −3.

Sometimes in algebra the same symbol has different meanings. Just like some words in English, the specific meaning becomes clear by looking at how it is used. You have seen the symbol "−" used in three different ways.

$10-4$ Between two numbers, it indicates the operation of *subtraction*.
We read $10-4$ as "10 minus 4."

-8 In front of a number, it indicates a *negative* number.
We read -8 as "negative eight."

$-x$ In front of a variable, it indicates the *opposite*.
We read $-x$ as "the opposite of x"

$-(-2)$ Here there are two "−" signs.
The one in the parentheses tells us the number is negative 2.
The one outside the parentheses tells us to take the *opposite* of -2.
We read $-(-2)$ as "the opposite of negative two."

OPPOSITE NOTATION

$-a$ means the opposite of the number a.

The notation $-a$ is read as "the opposite of a."

Example 1.4.2

Find:

1. the opposite of 7
2. the opposite of -10
3. the opposite of $-(-6)$

Solution:

1. -7 is the same distance from 0 as 7, but on the opposite side of 0. The opposite of 7 is -7.



2. 10 is the same distance from 0 as -10, but on the opposite side of 0. The opposite of -10 is 10.



3. The opposite of $-(-6)$ is -6.



Try It 1.4.3

Find:

1. the opposite of 4
2. the opposite of -3
3. $-(-1)$

Answer

1. -4
2. 3
3. 1

Try It 1.4.4

Find:

1. the opposite of 8
2. the opposite of -5
3. $-(-5)$

Answer

1. -8
2. 5
3. 5

Our work with opposites gives us a way to define the integers. The whole numbers and their opposites are called the **integers**. The integers are the numbers $\ldots -3, -2, -1, 0, 1, 2, 3 \ldots$

INTEGERS

The whole numbers and their opposites are called the **integers**.

The integers are the numbers

$$\ldots -3, -2, -1, 0, 1, 2, 3 \ldots$$

When evaluating the opposite of a **variable**, we must be very careful. Without knowing whether the variable represents a positive or negative number, we don't know whether −x−x is positive or negative. We can see this in Example 1.4.1.

Example 1.4.3:

Evaluate

1. $-x$, when $x = 8$
2. $-x$, when $x = -8$

Solution:

1.

To evaluate when $x = 8$ means to substitute 8 for x.	
	-x
Substitute 8 for x.	–(8)
Write the opposite of 8.	-8

2.

To evaluate when $x = -8$ means to substitute -8 for $-x$.	
	-x
Substitute -8 for x.	$-(-8)$
Write the opposite of -8.	8

Try It 1.4.5

Evaluate $-n$, when

1. $n = 4$
2. $n = -4$

Answer

1. -4
2. 4

Try It 1.4.6

Evaluate $-m$, when

1. $m = 11$
2. $m = -11$

Answer

1. -11
2. 11

Simplify: Expressions with Absolute Value

We saw that numbers such as 2 and -2 are opposites because they are the same distance from 0 on the number line. They are both two units from 0. The distance between 0 and any number on the number line is called the **absolute value** of that number.

ABSOLUTE VALUE

The **absolute value** of a number is its distance from 0 on the number line.

The absolute value of a number \(n\) is written as $|n|$.

For example,

- -5 is 5 units away from 0, so $|-5| = 5$.
- 5 is 5 units away from 0, so $|5| = 5$.

Figure $1.4.6$ illustrates this idea.




Figure $1.4.6$: The integers 5 and are 5 units away from 0.

The absolute value of a number is never negative (because distance cannot be negative). The only number with absolute value equal to zero is the number zero itself, because the distance from 0 to 0 on the number line is zero units.

PROPERTY OF ABSOLUTE VALUE

$|n| \geq 0$ for all numbers

Absolute values are always greater than or equal to zero!

Mathematicians say it more precisely, “absolute values are always non-negative.” Non-negative means greater than or equal to zero.

Example 1.4.4:

Simplify:

1. $|3|$
2. $|-44|$
3. $|0|$

Solution:

The absolute value of a number is the distance between the number and zero. Distance is never negative, so the absolute value is never negative.

1. $|3| = 3$
2. $|-44| = 44$
3. $|0| = 0$

Try It 1.4.7

Simplify:

1. $|4|$
2. $|-28|$
3. $|0|$

Answer

1. 4
2. 28
3. 0

Try It 1.4.8

Simplify:

1. $|-13|$
2. $|47|$

Answer

1. 13
2. 47

In the next example, we’ll order expressions with absolute values. Remember, positive numbers are always greater than negative numbers!

Example 1.4.5:

Fill in $<, >,$ or $=$ for each of the following pairs of numbers:

1. $|-5|$ ____ $-|-5|$
2. 8 ____ $-|-8|$
3. -9 ____ $-|-9|$
4. $-(-16)$ ____ $-|-16|$

Solution:

1. Simplification $|-5| \quad \_\_\_\_ \quad -|-5|$
   Order. $5 \quad \_\_\_\_ \quad -5$
   $5 \quad > \quad -5$
   $|-5| \quad > \quad -|-5|$
2. Simplification $8 \quad \_\_\_\_ \quad -|-8|$
   Order. $8 \quad \_\_\_\_ \quad -8$
   $8 \quad > \quad -8$
   so
   $|8| \quad > \quad -|-8|$
3. Simplification $-9 \quad \_\_\_\_ \quad -|-9|$
   Order. $-9 \quad \_\_\_\_ \quad -9$
   $-9 \quad = \quad -9$
   so
   $-9 \quad = \quad -|-9|$
4. Simplification $-(-16) \quad \_\_\_\_ \quad -|16|$
   Order. $16 \quad \_\_\_\_ \quad -16$
   $16 \quad > \quad -16$
   so
   $-(-16) \quad > \quad -|-16|$

Try It 1.4.9

Fill in $<, >,$ or $=$ for each of the following pairs of numbers:

1. $|-9|$ ____ $-|-9|$
2. 2 ____ $-|-2|$
3. -8 ____ $-|-8|$
4. $-(-9)$ ____ $-|-9|$

Answer

1. $>$
2. $>$
3. $<$
4. $>$

Try It 1.4.10

Fill in $<, >,$ or $=$ for each of the following pairs of numbers:

1. 7 ____ $-|-7|$
2. $-(-10)$ ____ $-|-10|$
3. $|-4|$ ____ $-|-4|$
4. -1 ____ $|-1|$

Answer

1. $>$
2. $>$
3. $>$
4. $<$

We now add absolute value bars to our list of grouping symbols. When we use the order of operations, first we simplify inside the absolute value bars as much as possible, then we take the **absolute value** of the resulting number.

GROUPING SYMBOLS

Parentheses	$()$	Braces	$\{\}$
Brackets	$[\,]$	Absolute	$\|\,\|$

In the next example, we simplify the expressions inside absolute value bars first, just like we do with parentheses.

Example 1.4.6:

Simplify: $24-|19-3(6-2)|$

Solution:

	$24-\|19-3(6-2)\|$
Work inside parentheses first: subtract 2 from 6	$24-\|19-3(4)\|$
Multiply $3(4)$	$24-\|19-12\|$
Subtract inside the absolute value bars.	$24-\|7\|$
Take the absolute value.	$24-7$
Subtract.	17

Try It 1.4.11

Simplify: $19-|11-4(3-1)|$

Answer

16

Try It 1.4.12

Simplify: $9-|8-4(7-5)|$

Answer

9

Example 1.4.7

Evaluate:

1. $|x|$ when $x = -35$
2. $|y|$ when $y = -20$
3. $-|u|$ when $u = 12$
4. $-|p|$ when $p = -14$

Solution:

1. $|x|$ when $x = -35$

	$\|x\|$
Substitute -35 for x	$\|-35\|$
Take the absolute value.	35

2. $|y|$ when $y = -20$

	$\|-y\|$
Substitute -20 for y	$\|-(-20)\|$
Simplify	$\|20\|$
Take the absolute value.	20

3. $-|u|$ when $u = 12$

	$-\|u\|$
Substitute 12 for u	$\|-12\|$
Take the absolute value.	-12

4. $-|p|$ when $p = -14$

	$-\|p\|$
Substitute -14 for p	$-\|-14\|$
Take the absolute value.	-14

Try It 1.4.13

Evaluate:

1. $|x|$ when $x = -17$
2. $|y|$ when $y = -39$
3. $-|m|$ when $m = 22$
4. $-|p|$ when $p = -11$

Answer

1. 17
2. 39
3. -22
4. -11

Try It 1.4.14

Evaluate:

1. $|y|$ when $y = -23$
2. $|-y|$ when $y = -21$
3. $-|n|$ when $n = 37$
4. $-|q|$ when $q = -49$

Answer

1. 23
2. 21
3. -37
4. -49

Add Integers

Most students are comfortable with the addition and subtraction facts for positive numbers. But doing addition or subtraction with both positive and negative numbers may be more challenging.

Doing the Manipulative Mathematics activity "Addition of Signed Numbers" will help you develop a better understanding of adding integers."

We will use two color counters to model addition and subtraction of negatives so that you can visualize the procedures instead of memorizing the rules.

We let one color (blue) represent positive. The other color (red) will represent the negatives. If we have one positive counter and one negative counter, the value of the pair is zero. They form a neutral pair. The value of this neutral pair is zero.





Figure 1.4.7

We will use the counters to show how to add the four addition facts using the numbers $5, -5$ and $3, -3$.


$$5+3 \quad -5+(-3) \quad -5+3 \quad 5+(-3)$$

To add $5+3$, we realize that $5+3$ means the sum of 5 and 3.

We start with 5 positives.	
And then we add 3 positives.	
We now have 8 positives. The sum of 5 and 3 is 8.	

Now we will add $-5+(-3)$. Watch for similarities to the last example $5+3=8$.

To add $-5+(-3)$, we realize this means the sum of -5 and -3.

We start with 5 negatives.	
And then we add 3 negatives.	
We now have 8 negative. The sum of -5 and -3 is -8.	

In what ways were these first two examples similar?

- The first example adds 5 positives and 3 positives—both positives.
- The second example adds 5 negatives and 3 negatives—both negatives.

In each case we got 8 — either 8 positives or 8 negatives.

When the signs were the same, the counters were all the same color, and so we added them.



Figure 1.4.7

? Example 1.4.8

Add:

1. $1+4$
2. $-1+(-4)$

Solution:

1. 

1 positive plus 4 positives is 5 positives.

2. 

1 negative plus 4 negatives is 5 negatives.

? Try It 1.4.15

Add:

1. $2+4$
2. $-2+(-4)$

Answer

1. 6
2. -6

? Try It 1.4.16

Add:

1. $2+5$
2. $-2+(-5)$



Answer

1. 7
2. -7

So what happens when the signs are different? Let's add $-5+3$. We realize this means the sum of -5 and 3. When the counters were the same color, we put them in a row. When the counters are a different color, we line them up under each other.

Table 1.4.1



	$-5+3$ means the sum of -5 and 3.
We start with 5 negatives.	

And then we add 3 positives.	
We remove any neutral pairs.	
We have 2 negatives left.	 2 negatives
The sum of -5 and 3 is -2.	$-5+3=2$

Notice that there were more negatives than positives, so the result was negative.

Let's now add the last combination, $5+(-3)$.

Table 1.4.2

	$5+(-3)$ means the sum of -5 and -3.
We start with 5 positives.	
And then we add 3 negatives.	
We remove any neutral pairs.	
We have 2 positives left.	 2 positives
The sum of 5 and -3 is 2.	$5+(-3)=2$

When we use counters to model addition of positive and negative integers, it is easy to see whether there are more positive or more negative counters. So we know whether the sum will be positive or negative.




Figure 1.4.8


? Example 1.4.9

Add:

1. $-1+5$
2. $1+(-5)$

Solution:


1. $-1+5$



There are more positives, so the sum is positive.

So, $-1+5=4$.

2. $1+(-5)$



There are more negatives, so the sum is negative.

So, $1+(-5)=-4$

Try It 1.4.17

Add:

1. $-2+4$
2. $2+(-4)$

Answer

1. 2
2. -2

Try It 1.4.18

Add:

1. $-2+5$
2. $2+(-5)$

Answer

1. 3
2. -3

Now that we have added small positive and negative integers with a model, we can visualize the model in our minds to simplify problems with any numbers.

When you need to add numbers such as $37+(-53)$, you really don't want to have to count out 37 blue counters and 53 red counters. With the model in your mind, can you visualize what you would do to solve the problem?

Picture 37 blue counters with 53 red counters lined up underneath. Since there would be more red (negative) counters than blue (positive) counters, the sum would be *negative*. How many more red counters would there be? Because $53-37=16$, there are 16 more red counters.

Therefore, the sum of $37+(-53)$ is -16.

$$37+(-53)=-16$$

Let's try another one. We'll add $-74+(-27)$. Again, imagine 74 red counters and 27 more red counters, so we'd have 101 red counters. This means the sum is -101.

$$-74+(-27)=-101$$

Let's look again at the results of adding the different combinations of $5, -5$ and $3, -3$.

ADDITION OF POSITIVE AND NEGATIVE INTEGERS

$5+3=8$ — both positive, sum positive

$-5+(-3)=-8$ — both negative, sum negative

When the signs are the same, the counters would be all the same color, so add them.

$-5+3=-2$ — different signs, more negatives, sum negative

$5+(-3)=2$ — different signs, more positives, sum positive

When the signs are different, some of the counters would make neutral pairs, so subtract to see how many are left.

Visualize the model as you simplify the expressions in the following examples.

Example 1.4.10

Simplify:

1. $19+(-47)$
2. $-14+(-36)$

Solution:

1. Since the signs are different, we subtract 19 from 47. The answer will be negative because there are more negatives than positives.

Add. $19+(-47)=-28$

2. Since the signs are same, we add. The answer will be negative because there are more negatives than positives.

Add. $-14+(-36)=-50$

Try It 1.4.19

Simplify:

1. $-31+(-19)$
2. $15+(-32)$

Answer

1. -50
2. -17

Try It 1.4.20

Simplify:

1. $-42+(-28)$
2. $25+(-61)$

Answer

1. -70
2. -36

The techniques used up to now extend to more complicated problems, like the ones we've seen before. Remember to follow the order of operations!

Example 1.4.11

Simplify:

$-5+3(-2+7)$

Solution:

	$-5+3(-2+7)$
Simplify inside the parenthesis	$-5+3(5)$
Multiply	$-5+15$
add left to right	10

Try It 1.4.21

Simplify:

$-2+5(-4+7)$

Answer

13

Try It 1.4.22

Simplify:

$-4+2(-3+5)$

Answer

0

Subtract Integers

Note

Doing the Manipulative Mathematics activity "Subtraction of Signed Numbers" will help you develop a better understanding of subtracting integers.

We will continue to use counters to model the subtraction. Remember, the blue counters represent positive numbers and the red counters represent negative numbers.


Perhaps when you were younger, you read "$5-3$" as "5 take away 3." When you use counters, you can think of subtraction the same way!

We will model the four subtraction facts using the numbers 5 and 3.

$$5-3 \quad -5-(-3)) \quad -5-3 \quad 5-(-3)$$

To subtract $5-3$, we restate the problem as "5 take away 3."

Table 1.4.3



We start with 5 positives.	
We 'take away' 3 positives.	

We have 2 positives left.	
The difference of 5 and 3 is 2.	2

Now we will subtract $-5-(-3)$. Watch for similarities to the last example $5-3=2$.

To subtract $-5-(-3)$, we restate this as "-5 take away -3"

Table 1.4.4

We start with 5 negatives.	
We 'take away' 3 negatives.	
We have 2 negatives left.	
The difference of -5 and -3 is -2.	-2

Notice that these two examples are much alike: The first example, we subtract 3 positives from 5 positives and end up with 2 positives.

In the second example, we subtract 3 negatives from 5 negatives and end up with 2 negatives.

Each example used counters of only one color, and the "take away" model of subtraction was easy to apply.




Figure 1.4.9

? Example 1.4.12

Subtract:

1. $7-5$
2. $-7-(-5)$

Solution:

1. $7-5$
   2 Take 5 positives from 7 positives and get 2 positives
2. $-7-(-5)$
   -2 Take 5 negatives from 7 negatives and get 2 negatives

? Try It 1.4.23

Subtract:

1. $6-4$
2. $-6-(-4)$

Answer

1. 2
2. -2

https://math.libretexts.org/@go/page/30456

Try It 1.4.24

Subtract:

1. $7 - 4$
2. $-7 - (-4)$

Answer

1. 3
2. -3


What happens when we have to subtract one positive and one negative number? We'll need to use both white and red counters as well as some **neutral pairs**. Adding a neutral pair does not change the value. It is like changing quarters to nickels—the value is the same, but it looks different.

- To subtract $-5 - 3$, we restate it as -5 take away 3.

We start with 5 negatives. We need to take away 3 positives, but we do not have any positives to take away.


Remember, a neutral pair has value zero. If we add 0 to 5 its value is still 5. We add neutral pairs to the 5 negatives until we get 3 positives to take away.

Table 1.4.5

	$-5 - 3$ means -5 take away 3
We start with 5 negatives.	
We now add the neutrals needed to get 3 positives.	
We remove the 3 positives.	
We are left with 8 negatives.	
The difference of -5 and 3 is -8.	$-5 - 3 = -8$

And now, the fourth case, $5 - (-3)$. We start with 5 positives. We need to take away 3 negatives, but there are no negatives to take away. So we add neutral pairs until we have 3 negatives to take away.

Table 1.4.6

	$5 - (-3)$ means 5 take away -3
We start with 5 positives.	
We now add the needed neutrals pairs.	
We remove the 3 negatives.	

We are left with 8 positives.	 8 positives
The difference of 5 and -3 is 8.	$5-(-3)=8$


Example 1.4.13

Subtract:


1. $-3-1$
2. $3-(-1)$

Solution:

1.

Take 1 positive from the one added neutral pair.		$-3-1$ -4

2.

Take 1 negative from the one added neutral pair.		$3-(-1)$ 4

Try It 1.4.25

Subtract:

1. $-6-4$
2. $6-(-4)$

Answer

1. -10
2. 10

Try It 1.4.26

Subtract:

1. $-7-4$
2. $7-(-4)$

Answer

1. -11
2. 11

Have you noticed that *subtraction of signed numbers can be done by adding the opposite*? In Exercise 1.4.33, $-3-1$ is the same as $-3+(-1)$ and $3-(-1)$ is the same as $3+1$. You will often see this idea, the **subtraction property**, written as follows:

SUBTRACTION PROPERTY

$$a-b=a+(-b)$$

Subtracting a number is the same as adding its opposite.

Look at these two examples.




Figure 1.4.10: $6-4$ gives the same answer as $6+(-4)$.

Of course, when you have a subtraction problem that has only positive numbers, like $6-4$, you just do the subtraction. You already knew how to subtract $6-4$ long ago. But *knowing* that $6-4$ gives the same answer as $6+(-4)$ helps when you are subtracting negative numbers. Make sure that you understand how $6-4$ and $6+(-4)$ give the same results!

Example 1.4.14

Simplify:

1. $13-8$ and $13+(-8)$
2. $-17-9$ and $-17+(-9)$

Solution:

1. Subtract. $13-8$ and $13+(-8)$
   5 5
2. Subtract. $-17-9$ and $-17+(-9)$
   -26 -26

Try It 1.4.27

Simplify:

1. $21-13$ and $21+(-13)$
2. $-11-7$ and $-11+(-7)$

Answer

1. 8
2. -18

Try It 1.4.28

Simplify:

1. $15-7$ and $15+(-7)$
2. $-14-8$ and $-14+(-8)$

Answer

1. 8
2. -22

Look at what happens when we subtract a negative.




Figure 1.4.11: $8-(-5)$ gives the same answer as $8+5$

Subtracting a negative number is like adding a positive!

You will often see this written as $a-(-b)=a+b$.

Does that work for other numbers, too? Let's do the following example and see.

Example 1.4.15

Simplify:

1. $9-(-15)$ and $9+15$
2. $-7-(-4)$ and $-7+4$

Solution:

1.		$9-(-15)$	$9+15$
	Subtract	24	24
2.		$-7-(-4)$	$-7+4$
	Subtract	-3	-3

Try It 1.4.29

Simplify:

1. $6-(-13)$ and $6+13$
2. $-5-(-1)$ and $-5+1$

Answer

1. 19
2. -4

Try It 1.4.30

Simplify:

1. $4-(-19)$ and $4+19$
2. $-4-(-7)$ and $-4+7$

Answer

1. 23
2. 3

Let's look again at the results of subtracting the different combinations of $5, -5$ and $3, -3$.

SUBTRACTION OF INTEGERS

$5-3$	$-5-(-3)$
2	-2
5 positives take away 3 positives	5 negatives take away 3 negatives
2 positives	2 negatives

When there would be enough counters of the color to take away, subtract.

$-5-3$	$5-(-3)$
-8	8
5 negatives, want to take away 3 positives need neutral pairs	5 positives, want to take away 3 negatives need neutral pairs

When there would be not enough counters of the color to take away, add.

What happens when there are more than three integers? We just use the order of operations as usual.

? Example 1.4.16

Simplify:

$7-(-4-3)-9$

Solution:

	$7-(-4-3)-9$
Simplify inside the parenthesis first.	$7-(-7)-9$
Subtract left to right.	$14-9$
Subtract	5

? Try It 1.4.31

Simplify:

$8-(-3-1)-9$

Answer

3

? Try It 1.4.32

Simplify:

$12-(-9-6)-14$

Answer

12

Access these online resources for additional instruction and practice with adding and subtracting integers. You will need to enable Java in your web browser to use the applications.

- Add Colored Chip
- Subtract Colored Chip

Key Concepts

- **Addition of Positive and Negative Integers**

$5+3$	$-5+(-3)$
8	-8
both positive, sum positive	both negative, sum negative

$-5+3$	$5+(-3)$
-2	2
different signs, more negatives sum negative	different signs, more positives sum positive

- **Property of Absolute Value**:

$|n| \geq 0$ for all numbers. Absolute values are always greater than or equal to zero!

- **Subtraction of Integers**

$5-3$	$-5-(-3)$
2	-2
5 positives take away 3 positives 2 positives	5 negatives take away 3 negatives 2 negatives
$-5-3$	$5-(-3)$
-8	8
5 negatives, want to subtract 3 positives need neutral pairs	5 positives, want to subtract 3 negatives need neutral pairs

- **Subtraction Property:** Subtracting a number is the same as adding its opposite.

Glossary

absolute value

The absolute value of a number is its distance from 0 on the number line. The absolute value of a number nn is written as |n|.

integers

The whole numbers and their opposites are called the integers: $\ldots -3, -2, -1, 0, 1, 2, 3 \ldots$

opposite

The opposite of a number is the number that is the same distance from zero on the number line but on the opposite side of zero: $-a$ means the opposite of the number. The notation $-a$ is read “the opposite of a.”

This page titled 1.4: Add and Subtract Integers is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.4: Add and Subtract Integers** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.4E: Exercises

Practice Makes Perfect

Use Negatives and Opposites of Integers

In the following exercises, order each of the following pairs of numbers, using < or >.

Exercise 1.4E.1

1. 9___4
2. −3___6
3. −8___−2
4. 1___−10

Answer

1. >
2. <
3. <
4. >

Exercise 1.4E.2

1. −7___3
2. −10___−5
3. 2___−6
4. 8___9

In the following exercises, find the opposite of each number.

Exercise 1.4E.3

1. 2
2. −6

Answer

1. −2
2. 6

Exercise 1.4E.4

1. 9
2. −4

In the following exercises, simplify.

Exercise 1.4E.5

−(−4)

Answer

4

Exercise 1.4E.6

$-(-8)$

Exercise 1.4E.7

$-(-15)$

Answer

15

Exercise 1.4E.8

$-(-11)$

In the following exercises, evaluate.

Exercise 1.4E.9

$-c$ when

1. $c=12$
2. $c=-12$

Answer

1. -12
2. 12

1. Exercise 1.4E.10

$-d$ when

1. $d=21$
2. $d=-21$

Simplify Expressions with Absolute Value

In the following exercises, simplify.

Exercise 1.4E.11

1. $|-32|$
2. $|0|$
3. $|16|$

Answer

1. 32
2. 0
3. 16

Exercise 1.4E.12

1. $|0|$
2. $|-40|$
3. $|22|$

In the following exercises, fill in <, >, or = for each of the following pairs of numbers.

Exercise 1.4E.13

1. $-6 \_\_\_ |-6|$
2. $-|-3| \_\_\_ -3$

Answer

1. <
2. =

Exercise 1.4E.14

1. $|-5| \_\_\_ -|-5|$
2. $9 \_\_\_ -|-9|$

In the following exercises, simplify.

Exercise 1.4E.15

$-(-5)$ and $-|-5|$

Answer

$5, -5$

Exercise 1.4E.16

$-|-9|$ and $-(-9)$

Exercise 1.4E.17

$8|-7|$

Answer

56

Exercise 1.4E.18

$5|-5|$

Exercise 1.4E.19

$|15-7|-|14-6|$

Answer

0

Exercise 1.4E.20

$|17-8|-|13-4|$

Exercise 1.4E.21

$18-|2(8-3)|$

Answer

8

Exercise 1.4E.22

$18-|3(8-5)|$

In the following exercises, evaluate.

Exercise 1.4E.23

1. $-|p|$ when $p=19$
2. $-|q|$ when $q=-33$

Answer

1. -19
2. -33

Exercise 1.4E.24

1. $-|a|$ when $a=60$
2. $-|b|$ when $b=-12$

Add Integers

In the following exercises, simplify each expression.

Exercise 1.4E.25

$-21+(-59)$

Answer

-80

Exercise 1.4E.26

$-35+(-47)$

Exercise 1.4E.27

$48+(-16)$

Answer

32

Exercise 1.4E.28

$34+(-19)$

Exercise 1.4E.29

$-14+(-12)+4$

Answer

-22

Exercise 1.4E.30

$-17+(-18)+6$

Exercise 1.4E.31

$135+(-110)+83$

Answer

108

Exercise 1.4E.32

$-38+27+(-8)+12$

Exercise 1.4E.33

$19+2(-3+8)$

Answer

29

Exercise 1.4E.34

$24+3(-5+9)$

Subtract Integers

In the following exercises, simplify.

Exercise 1.4E.35

$8-2$

Answer

6

Exercise 1.4E.36

$-6-(-4)$

Exercise 1.4E.37

$-5-4$

Answer

-9

Exercise 1.4E.38

$-7-2$

Exercise 1.4E.39

$8-(-4)$

Answer

12

Exercise 1.4E.40

$7-(-3)$

Exercise 1.4E.41

1. $44-28$
2. $44+(-28)$

Answer

1. 16
2. 16

Exercise 1.4E.42

1. $35-16$
2. $35+(-16)$

Exercise 1.4E.43

1. $27-(-18)$
2. $27+18$

Answer

1. 45
2. 45

Exercise 1.4E.44

1. $46-(-37)$
2. $46+37$

In the following exercises, simplify each expression.

Exercise 1.4E.45

$15-(-12)$

Answer

27

Exercise 1.4E.46

$14-(-11)$

Exercise 1.4E.47

$48-87$

Answer

-39

Exercise 1.4E.48

$45-69$

Exercise 1.4E.49

$-17-42$

Answer

-59

Exercise 1.4E.50

$-19-46$

Exercise 1.4E.51

$-103-(-52)$

Answer

-51

Exercise 1.4E.52

$-105-(-68)$

Exercise 1.4E.53

$-45-(-54)$

Answer

9

Exercise 1.4E.54

$-58-(-67)$

Exercise 1.4E.55

$8-3-7$

Answer

-2

Exercise 1.4E.56

$9-6-5$

Exercise 1.4E.57

$-5-4+7$

Answer

-2

Exercise 1.4E.58

$-3-8+4$

Exercise 1.4E.59

$-14-(-27)+9$

Answer

22

Exercise 1.4E.60

$64+(-17)-9$

Exercise 1.4E.61

$(2-7)-(3-8)$

Answer

0

Exercise 1.4E.62

$(1-8)-(2-9)$

Exercise 1.4E.63

$-(6-8)-(2-4)$

Answer

4

Exercise 1.4E.64

$-(4-5)-(7-8)$

Exercise 1.4E.65

$25-[10-(3-12)]$

Answer

6

Exercise 1.4E.66

$32-[5-(15-20)]$

Exercise 1.4E.67

$6.3-4.3-7.2$

Answer

-5.2

Exercise 1.4E.68

$5.7-8.2-4.9$

Exercise 1.4E.69

5^2-6^2

Answer

-11

Exercise 1.4E.70

6^2-7^2

Everyday Math

Exercise 1.4E.71

Elevation The highest elevation in the United States is Mount McKinley, Alaska, at 20,320 feet above sea level. The lowest elevation is Death Valley, California, at 282 feet below sea level.

Use integers to write the elevation of:

1. Mount McKinley.
2. Death Valley.

Answer

1. 20,329
2. −282

Exercise 1.4E.72

Extreme temperatures The highest recorded temperature on Earth was 58° Celsius, recorded in the Sahara Desert in 1922. The lowest recorded temperature was 90° below 0° Celsius, recorded in Antarctica in 1983.

Use integers to write the:

1. highest recorded temperature.
2. lowest recorded temperature.

Exercise 1.4E.73

State budgets In June, 2011, the state of Pennsylvania estimated it would have a budget surplus of $540 million. That same month, Texas estimated it would have a budget deficit of $27 billion.

Use integers to write the budget of:

1. Pennsylvania.
2. Texas.

Answer

1. $540 million
2. −$27 billion

Exercise 1.4E.74

College enrollments Across the United States, community college enrollment grew by 1,400,000 students from Fall 2007 to Fall 2010. In California, community college enrollment declined by 110,171 students from Fall 2009 to Fall 2010.

Use integers to write the change in enrollment:

1. in the U.S. from Fall 2007 to Fall 2010.
2. in California from Fall 2009 to Fall 2010.

Exercise 1.4E.75

Stock Market The week of September 15, 2008 was one of the most volatile weeks ever for the US stock market. The closing numbers of the Dow Jones Industrial Average each day were:

Monday	−504
Tuesday	+142
Wednesday	−449
Thursday	+410
Friday	+369

What was the overall change for the week? Was it positive or negative?

Answer

-32

Exercise 1.4E.76

Stock Market During the week of June 22, 2009, the closing numbers of the Dow Jones Industrial Average each day were:

Monday	−201
Tuesday	−16
Wednesday	−23
Thursday	+172
Friday	−34

What was the overall change for the week? Was it positive or negative?

Writing Exercises

Exercise 1.4E.77

Give an example of a negative number from your life experience.

Answer

Answers may vary

Exercise 1.4E.78

What are the three uses of the "−" sign in algebra? Explain how they differ.

Exercise 1.4E.79

Explain why the sum of −8 and 2 is negative, but the sum of 8 and −2 is positive.

Answer

Answers may vary

Exercise 1.4E.80

Give an example from your life experience of adding two negative numbers.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
use negatives and opposites of integers.			
simplify expressions with absolute value.			
add integers.			
subtract integers.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

More Inequality Practice

Exercise 1.4E.81

Enter the inequality symbol < or the symbol > in the box in order that the resulting inequality is a true statement.

1. $-4 \square 0$	5. $-3 \square -1$	9. $-3 \square -6$
2. $-4 \square 3$	6. $6 \square 5$	10. $0 \square -3$
3. $-2 \square -1$	7. $3 \square 6$	11. $-1 \square 4$
4. $3 \square 0$	8. $-4 \square -2$	12. $1 \square -4$

Answers to odd problems for E.81

1. −4 <0	3. −2 < −1	5. −3 < −1	7. 3 < 6	9. −3 > −6	11. −1 < 4

More Absolute Value Practice

Exercise 1.4E.82

Simplify each of the following expressions.

1. $\|7\|$.	5. $-\|-11\|$.	9. $-\|-20\|$.
2. $\|1\|$.	6. $-\|-1\|$.	10. $-\|-8\|$.
3. $\|-5\|$.	7. $\|-5\|$.	11. $\|-4\|$.
4. $\|-3\|$.	8. $\|-1\|$.	12. $\|-3\|$.

Answers to odd problems for E.82

1. 7	3. 5	5. -11	7. 5	9. -20	11. 4

More Addition Practice

Exercise 1.4E.83

Find the sum of the given integers.

1. −15 + 1	9. 2+5	17. −11 + (−6)
2. −1 + 18	10. 14 + 1	18. −7+(−8)
3. 18 + (−10)	11. 19 + (−15)	19. 14 + (−9)
4. 2+(−19)	12. 20 + (−17)	20. 5+(−15)
5. −10 + (−12)	13. −2+(−7)	21. 10 + 11
6. −1+(−7)	14. −14 + (−6)	22. 14 + 11
7. 5 + 10	15. −6 + 16	23. −13 + 1
8. 1 + 12	16. −2 + 14	24. −8+2

Answers to odd problems for E.83

1. −14	9. 7	17. −17
3. 8	11. 4	19. 5
5. −22	13. −9	21. 21
7. 15	15. 10	23. −12

.

More Subtraction Practice

Exercise 1.4E. 84

Find the difference of the given integers.

1. 16 − 20	9. −4 − (−9)	17. 13 − (−1)
2. 17 − 2	10. −13 − (−3)	18. 12 − (−13)
3. 10 − 12	11. 8 − (−3)	19. −4 − (−2)
4. 16 − 8	12. 14 − (−20)	20. −6 − (−8)
5. 14 − 11	13. 2 − 11	21. 7 − (−8)
6. 5 − 8	14. 16 − 2	22. 13 − (−14)
7. 7 − (−16)	15. −8 − (−10)	23. −3 − (−10)
8. 20 − (−10)	16. −14 − (−2)	24. −13 − (−9)

Answers to odd problems for E.84

1. −4	9. 5	17. 14
3. −2	11. 11	19. −2
5. 3	13. −9	21. 15
7. 23	15. 2	23. 7

.

This page titled 1.4E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.4E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.5: Multiply and Divide Integers

Learning Objectives

By the end of this section, you will be able to:

- Multiply integers
- Divide integers
- Simplify expressions with integers
- Evaluate variable expressions with integers
- Translate English phrases to algebraic expressions
- Use integers in applications

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapter, **Integers**.

Multiply Integers

Since multiplication is mathematical shorthand for repeated addition, our model can easily be applied to show **multiplication** of **integers**. Let's look at this concrete model to see what patterns we notice. We will use the same examples that we used for addition and subtraction. Here, we will use the model just to help us discover the pattern.

We remember that $a \cdot b$ means add a, b times. Here, we are using the model just to help us discover the pattern.




Figure 1.5.1

The next two examples are more interesting.

What does it mean to multiply 5 by -3? It means subtract $5, 3$ times. Looking at subtraction as "taking away," it means to take away $5, 3$ times. But there is nothing to take away, so we start by adding neutral pairs on the workspace. Then we take away 5 three times.




Figure 1.5.2

In summary:

$$5 \cdot 3 = 15 \qquad -5(3) = -15$$
$$5(-3) = -15 \qquad (-5)(-3) = 15$$

Notice that for multiplication of two signed numbers, when the:

- signs are the *same*, the product is *positive*.
- signs are *different*, the product is *negative*.

We'll put this all together in the chart below.

MULTIPLICATION OF SIGNED NUMBERS

For multiplication of two signed numbers:

Table 1.5.1

Same signs	Product	Example
Two positives	Positive	$7 \cdot 4 = 28$
Two negatives	Positive	$-8(-6) = 48$

Table 1.5.2

Different signs	Product	Example
Positive · negative	Negative	$7(-9) = -63$
Negative · positive	Negative	$-5 \cdot 10 = -50$

Example 1.5.1

Multiply:

1. $-9 \cdot 3$
2. $-2(-5)$
3. $4(-8)$
4. $7 \cdot 6$

Solution

1. $-9 \cdot 3 = -27$ Multiply, noting that the signs are different, so the product is negative.
2. $-2(-5) = 10$ Multiply, noting that the signs are same, so the product is positive.
3. $4(-8) = -32$ Multiply, with different signs.
4. $7 \cdot 6 = 42$ Multiply, with the same signs.

Try It 1.5.2

Multiply:

1. $-6 \cdot 8$
2. $-4(-7)$
3. $9(-7)$
4. $5 \cdot 12$

Answer

1. -48
2. 28
3. -63
4. 60

Try It 1.5.3

Multiply:

1. $-8 \cdot 7$
2. $-6(-9)$
3. $7(-4)$
4. $3 \cdot 13$

Answer

1. -56
2. 54
3. -28
4. 39

When we multiply a number by 1, the result is the same number. What happens when we multiply a number by -1? Let's multiply a positive number and then a negative number by -1 to see what we get.

Multiply.	Multiply.
$-1 \cdot 4 = -4$	$-1(-3) = 3$
Note that -4 is the opposite of 4.	Note that 3 is the opposite of -3

Each time we multiply a number by -1, we get its opposite!

MULTIPLICATION BY −1

$$-1a = -a$$

Multiplying a number by -1 gives its opposite.

Example 1.5.4

Multiply:

1. $-1 \cdot 7$
2. $-1(-11)$

Solution

1.		$-1 \cdot 7$
	Multiply, noting that the signs are different so the product is negative.	-7
		-7 is the opposite of 7.
2.		$-1(-11)$
	Multiply, noting that the signs are the same so the product is positive.	11
		11 is the opposite of -11.

Try It 1.5.5

Multiply:

1. $-1 \cdot 9$
2. $-1 \cdot (-17)$

Answer

1. -9
2. 17

Try It 1.5.6

Multiply:

1. $-1 \cdot 8$
2. $-1 \cdot (-16)$

Answer

1. -8
2. 16

Divide Integers

What about **division**? Division is the inverse operation of multiplication. So, $15 \div 3 = 5$ because $5 \cdot 3 = 15$. In words, this expression says that 15 can be divided into three groups of five each because adding five three times gives 15. Look at some examples of multiplying integers, to figure out the rules for dividing integers.

$$\begin{aligned} 5 \cdot 3 &= 15 \quad \text{so} \quad 15 \div 3 = 5 \\ -5 \cdot 3 &= -15 \quad \text{so} \quad -15 \div 3 = -5 \\ (5)(-3) &= -15 \quad \text{so} \quad -15 \div (-3) = 5 \\ (-5)(-3) &= 15 \quad \text{so} \quad 15 \div (-3) = -5 \end{aligned}$$

Division follows the same rules as multiplication!

For division of two signed numbers, when the:

- signs are the *same*, the quotient is *positive*.
- signs are *different*, the quotient is *negative*.

And remember that we can always check the answer of a division problem by multiplying.

MULTIPLICATION AND DIVISION OF SIGNED NUMBERS

For multiplication and division of two signed numbers:

- If the signs are the same, the result is positive.
- If the signs are different, the result is negative.

Table 1.5.3

Same signs	Result
Two positives	Positive
Two negatives	Positive
If the signs are the same, the result is positive.	

Table 1.5.4

Different signs	Result
Positive and negative	Negative
Negative and positive	Negative
If the signs are different, the result is negative.	

Example 1.5.7

1. $-27 \div 3$
2. $-100 \div (-4)$

Solution

1. $-27 \div 3$
   Divide, with different signs, the quotient is negative. -9
2. $-100 \div (-4)$
   Divide, with signs that are the same the quotient is negative. 25

Try It 1.5.8

Divide:

1. $-42 \div 6$
2. $-117 \div (-3)$

Answer

1. -7
2. 39

Try It 1.5.9

Divide:

1. $-63 \div 7$
2. $-115 \div (-5)$

Answer

1. -9
2. 23

Simplify Expressions with Integers

What happens when there are more than two numbers in an expression? The order of operations still applies when negatives are included. Remember My Dear Aunt Sally?

Let's try some examples. We'll simplify expressions that use all four operations with integers—addition, subtraction, multiplication, and division. Remember to follow the order of operations.

Example 1.5.10

Simplify:

$7(-2)+4(-7)-6$

Solution

	$7(-2)+4(-7)-6$
Multiply first.	$-14+(-28)-6$
Add.	$-42-6$
Subtract	-48

Try It 1.5.11

Simplify:

$8(-3)+5(-7)-4$

Answer

-63

Try It 1.5.12

Simplify:

$9(-3)+7(-8)-1$

Answer

-84

Example 1.5.13

Simplify:

1. $(-2)^4$
2. -2^4

Solution

1. $(-2)^4$

Write in expanded form.	$(-2)(-2)(-2)(-2)$
Multiply	$4(-2)(-2)$
Multiply	$-8(-2)$
Multiply	16

2. -2^4

Write in expanded form. We are asked to find the opposite of 2^4.	$-(2\cdot 2\cdot 2\cdot 2)$
Multiply	$-(4\cdot 2\cdot 2)$
Multiply	$-(8\cdot 2)$
Multiply	-16

Notice the difference in parts (1) and (2). In part (1), the exponent means to raise what is in the parentheses, the (-2) to the 4th power. In part (2), the exponent means to raise just the 2 to the 4th power and then take the opposite.

? Try It 1.5.14

Simplify:

1. $(-3)^4$
2. -3^4

Answer

1. 81
2. -81

? Try It 1.5.15

Simplify:

1. $(-7)^2$
2. -7^2

Answer

1. 49
2. -49

The next example reminds us to simplify inside parentheses first.

✓ Example 1.5.16

Simplify:

$12-3(9-12)$

Solution

$12-3(9-12)$	
$=12-3(-3)$	Subtract in parentheses first.
$=12-(-9)$	Multiply.
$=12+9$	Add the opposite.
$=21$	

Try It 1.5.17

Simplify:

$17-4(8-11)$

Answer

29

Try It 1.5.18

Simplify:

$16-6(7-13)$

Answer

52

Example 1.5.19

Simplify:

$8(-9)\div(-2)^3$

Solution

	$8(-9)\div(-2)^3$
Exponents first	$8(-9)\div(-8)$
Multiply.	$-72\div(-8)$
Divide	9

Try It 1.5.20

Simplify:

$12(-9)\div(-3)^3$

Answer

4

Try It 1.5.21

Simplify:

$18(-4)\div(-2)^3$

Answer

9

Example 1.5.22

Simplify:

$-30\div 2+(-3)(-7)$

Solution

	$-30 \div 2 + (-3)(-7)$
Multiply and divide left to right, so divide first.	$-15 + (-3)(-7)$
Multiply.	$-15 + 21$
Add	6

Try It 1.5.23

Simplify:

$-27 \div 3 + (-5)(-6)$

Answer

21

Try It 1.5.24

Simplify:

$-32 \div 4 + (-2)(-7)$

Answer

6

Evaluate Variable Expressions with Integers

Remember that to evaluate an expression means to substitute a number for the variable in the expression. Now we can use negative numbers as well as positive numbers.

Example 1.5.25

When $n = -5$, evaluate:

1. $n + 1$
2. $-n + 1$.

Solution

1.

	$n + 1$
Substitute -5 for n	$-5 + 1$
Simplify.	-4

2.

	$n + 1$
Substitute -5 for n	$-(-5) + 1$
Simplify.	$5 + 1$
Add.	6

Try It 1.5.26

When $n = -8$, evaluate:

1. $n + 2$
2. $-n + 2$.

Answer

1. -6
2. 10

Try It 1.5.27

When $y = -9$, evaluate:

1. $y + 8$
2. $-y + 8$.

Answer

1. -1
2. 17

Example 1.5.28

Evaluate $(x + y)^2$ when $x = -18$ and $y = 24$.

Solution

	$(x+y)^2$
Substitute -18 for x and 24 for y	$(-18+24)^2$
Add inside parentheses	$(6)^2$
Simplify.	36

Try It 1.5.29

Evaluate $(x + y)^2$ when $x = -15$ and $y = 29$.

Answer

196

Try It 1.5.30

Evaluate $(x + y)^3$ when $x = -8$ and $y = 10$.

Answer

8

Example 1.5.31

Evaluate $20 - z$ when

1. $z = 12$
2. $z = -12$

Solution

1.

	$20 - z$
Substitute 12 for z.	$20 - 12$
Subtract	8

2.

	$20 - z$
Substitute -12 for z.	$20 - (-12)$
Subtract	32

Try It 1.5.32

Evaluate $17 - k$ when

1. $k = 19$
2. $k = -19$

Answer

1. -2
2. 36

Try It 1.5.33

Evaluate $-5 - b$ when

1. $b = 14$
2. $b = -14$

Answer

1. -19
2. 9

Example 1.5.34

Evaluate:

$2x^2 + 3x + 8$ when $x = 4$.

Solution

Substitute 4 for x. Use parentheses to show multiplication.

	$2x^2 + 3x + 8$
Substitute	$2(4)^2 + 3(4) + 8$
Evaluate exponents.	$2(16) + 3(4) + 8$
Multiply.	$32 + 12 + 8$
Add.	52

Try It 1.5.35

Evaluate:

$3x^2 - 2x + 6$ when $x = -3$.

Answer

39

Try It 1.5.36

Evaluate:

$4x^2 - x - 5$ when $x = -2$.

Answer

13

Translate Phrases to Expressions with Integers

Our earlier work translating English to algebra also applies to phrases that include both positive and negative numbers.

Example 1.5.37

Translate and simplify: the sum of 8 and -12, increased by 3.

Solution

	the **sum** of 8 and -12, increased by 3
Translate.	$[8+(-12)]+3$
Simplify. Be careful not to confuse the brackets with an absolute value sign.	$(-4)+3$
Add.	-1

Try It 1.5.38

Translate and simplify: the sum of 9 and -16, increased by 4.

Answer

$(9+(-16))+4-3$

Try It 1.5.39

Translate and simplify: the sum of -8 and -12, increased by 7.

Answer

$(-8+(-12))+7-13$

When we first introduced the operation symbols, we saw that the expression may be read in several ways. They are listed in the chart below.

Table 1.5.5

$a-b$
a minus b
the difference of a and b
b subtracted from a
b less than a

Be careful to get a and b in the right order!

Example 1.5.40

Translate and then simplify

1. the difference of 13 and -21
2. subtract 24 from -19.

Solution

1.

	the **difference** of 13 and -21
Translate.	$13-(-21)$
Simplify.	34

2.

	subtract 24 **from** -19
Translate.	$-19-24$
Remember, subtract b from a means $a-b$	
Simplify.	-43

Try It 1.5.41

Translate and simplify

1. the difference of 14 and -23
2. subtract 21 from -17.

Answer

1. $14-(-23); 37$
2. $-17-21; -38$

Try It 1.5.42

Translate and simplify

1. the difference of 11 and -19
2. subtract 18 from -11.

Answer

1. $11-(-19); 30$
2. $-11-18; -29$

Once again, our prior work translating English to algebra transfers to phrases that include both multiplying and dividing integers. Remember that the key word for **multiplication** is "**product**" and for **division** is "**quotient**."

Example 1.5.43

Translate to an algebraic expression and simplify if possible: the product of -2 and 14.

Solution

	the product of -2 and 14
Translate.	$(-2)(14)$
Simplify.	-28

Try It 1.5.44

Translate to an algebraic expression and simplify if possible: the product of -5 and 12.

Answer

$-5(12); -60$

Try It 1.5.45

Translate to an algebraic expression and simplify if possible: the product of 8 and -13.

Answer

$-8(13); -104$

Example 1.5.46

Translate to an algebraic expression and simplify if possible: the quotient of -56 and -7.

Solution

$$\begin{array}{ll} & \text{the quotient of } -56 \text{ and } -7 \\ \text{Translate.} & -56 \div (-7) \\ \text{Simplify.} & 8 \end{array} \quad (1.5.1)$$

Try It 1.5.47

Translate to an algebraic expression and simplify if possible: the quotient of -63 and -9.

Answer

$-63 \div (-9); 7$

Try It 1.5.48

Translate to an algebraic expression and simplify if possible: the quotient of -72 and -9.

Answer

$-72 \div (-9); 8$

Use Integers in Applications

We'll outline a plan to solve applications. It's hard to find something if we don't know what we're looking for or what to call it! So when we solve an application, we first need to determine what the problem is asking us to find. Then we'll write a phrase that gives the information to find it. We'll translate the phrase into an expression and then simplify the expression to get the answer. Finally, we summarize the answer in a sentence to make sure it makes sense.

How to Apply a Strategy to Solve Applications with Integers

Example 1.5.49

The temperature in Urbana, Illinois one morning was 11 degrees. By mid-afternoon, the temperature had dropped to -9 degrees. What was the difference of the morning and afternoon temperatures?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.	
Step 2. Identify what we are asked to find.	the difference of the morning and afternoon temperatures
Step 3. Write a phrase that gives the information to find it.	the difference of 11 and -9
Step 4. Translate the phrase to an expression.	$11 - (-9)$
Step 5. Simplify the expression.	20
Step 6. Write a complete sentence that answers the question.	The difference in temperatures was 20 degrees.

Try It 1.5.50

The temperature in Anchorage, Alaska one morning was 15 degrees. By mid-afternoon the temperature had dropped to 30 degrees below zero. What was the difference in the morning and afternoon temperatures?

Answer

The difference in temperatures was 45 degrees.

Try It 1.5.51

The temperature in Denver was -6 degrees at lunchtime. By sunset the temperature had dropped to -15 degrees. What was the difference in the lunchtime and sunset temperatures?

Answer

The difference in temperatures was 9 degrees.

APPLY A STRATEGY TO SOLVE APPLICATIONS WITH INTEGERS.

1. Read the problem. Make sure all the words and ideas are understood
2. Identify what we are asked to find.
3. Write a phrase that gives the information to find it.
4. Translate the phrase to an expression.
5. Simplify the expression.
6. Answer the question with a complete sentence.

Example 1.5.52

The Mustangs football team received three penalties in the third quarter. Each penalty gave them a loss of fifteen yards. What is the number of yards lost?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.	
Step 2. Identify what we are asked to find.	the number of yards lost
Step 3. Write a phrase that gives the information to find it.	three times a 15-yard penalty
Step 4. Translate the phrase to an expression.	$3(-15)$
Step 5. Simplify the expression.	-45
Step 6. Write a complete sentence that answers the question.	The team lost 45 yards.

Try It 1.5.53

The Bears played poorly and had seven penalties in the game. Each penalty resulted in a loss of 15 yards. What is the number of yards lost due to penalties?

Answer

The Bears lost 105 yards.

? Try It 1.5.54

Bill uses the ATM on campus because it is convenient. However, each time he uses it he is charged a $2 fee. Last month he used the ATM eight times. How much was his total fee for using the ATM?

Answer

A $16 fee was deducted from his checking account.

Key Concepts

- **Multiplication and Division of Two Signed Numbers**
  - Same signs—Product is positive
  - Different signs—Product is negative
- **Strategy for Applications**
  1. Identify what you are asked to find.
  2. Write a phrase that gives the information to find it.
  3. Translate the phrase to an expression.
  4. Simplify the expression.
  5. Answer the question with a complete sentence.

This page titled 1.5: Multiply and Divide Integers is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.5: Multiply and Divide Integers** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.5E: Exercises

Practice Makes Perfect

Multiply Integers

In the following exercises, multiply.

Exercise 1.5E.55

$-4 \cdot 8$

Answer

-32

Exercise 1.5E.56

$-3 \cdot 9$

Exercise 1.5E.57

$9(-7)$

Answer

-63

Exercise 1.5E.58

$13(-5)$

Exercise 1.5E.59

$-1 \cdot 6$

Answer

-6

Exercise 1.5E.60

$-1 \cdot 3$

Exercise 1.5E.61

$-1(-14)$

Answer

14

Exercise 1.5E.62

$-1(-19)$

Divide Integers

In the following exercises, divide.

Exercise 1.5E.63

$-24 \div 6$

Answer

-4

Exercise 1.5E.64

$35 \div (-7)$

Exercise 1.5E.65

$-52 \div (-4)$

Answer

13

Exercise 1.5E.66

$-84 \div (-6)$

Exercise 1.5E.67

$-180 \div 15$

Answer

-12

Exercise 1.5E.68

$-192 \div 12$

Simplify Expressions with Integers

In the following exercises, simplify each expression.

Exercise 1.5E.69

$5(-6)+7(-2)-3$

Answer

-47

Exercise 1.5E.70

$8(-4)+5(-4)-6$

Exercise 1.5E.71

$(-2)^6$

Answer

64

Exercise 1.5E.72

$(-3)^5$

Exercise 1.5E.73

$(-4)^2$

Answer

-16

Exercise 1.5E.74

$(-6)^2$

Exercise 1.5E.75

$-3(-5)(6)$

Answer

90

Exercise 1.5E.76

$-4(-6)(3)$

Exercise 1.5E.77

$(8-11)(9-12)$

Answer

9

Exercise 1.5E.78

$(6-11)(8-13)$

Exercise 1.5E.79

$26-3(2-7)$

Answer

41

Exercise 1.5E.80

$23-2(4-6)$

Exercise 1.5E.81

$65 \div (-5) + (-28) \div (-7)$

Answer

-9

Exercise 1.5E.82

$52 \div (-4) + (-32) \div (-8)$

Exercise 1.5E.83

$9-2[3-8(-2)]$

Answer

-29

Exercise 1.5E.84

$11-3[7-4(-20)]$

Exercise 1.5E.85

$(-3)^2 - 24 \div (8-2)$

Answer

5

Exercise 1.5E.86

$(-4)^2 - 32 \div (12-4)$

Evaluate Variable Expressions with Integers

In the following exercises, evaluate each expression.

Exercise 1.5E.87

y+(−14) when

1. y=−33
2. y=30

Answer

1. −47
2. 16

Exercise 1.5E.88

x+(−21) when

1. x=−27
2. x=44

Exercise 1.5E.89

1. a+3 when a=−7
2. −a+3 when a=−7

Answer

1. −4
2. 10

Exercise 1.5E.90

1. d+(−9) when d=−8
2. −d+(−9) when d=−8

Exercise 1.5E.91

m+n when
m=−15,n=7

Answer

-8

Exercise 1.5E.92

p+q when
p=−9,q=17

Exercise 1.5E.93

r+s when r=−9,s=−7

Answer

-16

Exercise 1.5E.94

t+u when t=−6,u=−5

Exercise 1.5E.95

$(x+y)^2$ when
x=−3,y=14

Answer

121

Exercise 1.5E.96

$(y+z)^2$ when
y=−3, z=15

Exercise 1.5E.97

−2x+17 when

1. x=8
2. x=−8

Answer

1. 1
2. 33

Exercise 1.5E.98

−5y+14 when

1. y=9
2. y=−9

Exercise 1.5E.99

10−3m when

1. m=5
2. m=−5

Answer

1. −5
2. 25

Exercise 1.5E.100

18−4n when

1. n=3
2. n=−3

Exercise 1.5E.101

$2w^2-3w+7$ when
w=−2

Answer

21

Exercise 1.5E.102

$3u^2-4u+5$

Exercise 1.5E.103

9a−2b−8 when
a=−6 and b=−3

Answer

-56

Exercise 1.5E.104

7m−4n−2 when
m=−4 and n=−9

Translate English Phrases to Algebraic Expressions

In the following exercises, translate to an algebraic expression and simplify if possible.

Exercise 1.5E.105

the sum of 3 and −15, increased by 7

Answer

(3+(−15))+7;−5

Exercise 1.5E.106

the sum of −8 and −9, increased by 23

Exercise 1.5E.107

the difference of 10 and −18

Answer

10−(−18);28

Exercise 1.5E.108

subtract 11 from −25

Exercise 1.5E.109

the difference of −5 and −30

Answer

−5−(−30);25

Exercise 1.5E.110

subtract −6 from −13

Exercise 1.5E.111

the product of −3 and 15

Answer

$-3 \cdot 15; -45$

Exercise 1.5E.112

the product of −4 and 16

Exercise 1.5E.113

the quotient of −60 and −20

Answer

$-60 \div (-20)\,; 3$

Exercise 1.5E.114

the quotient of −40 and −20

Exercise 1.5E.115

the quotient of −6 and the sum of *a* and *b*

Answer

$\frac{-6}{a+b}$

Exercise 1.5E.116

the quotient of −6 and the sum of *a* and *b*

Exercise 1.5E.117

the product of −10 and the difference of p and q

Answer

−10(p−q)

Exercise 1.5E.118

the product of −13 and the difference of c and d

Use Integers in Applications

In the following exercises, solve.

Exercise 1.5E.119

Temperature On January 15, the high temperature in Anaheim, California, was 84°. That same day, the high temperature in Embarrass, Minnesota was −12°. What was the difference between the temperature in Anaheim and the temperature in Embarrass?

Answer

96°

Exercise 1.5E.120

Temperature On January 21, the high temperature in Palm Springs, California, was 89°, and the high temperature in Whitefield, New Hampshire was −31°. What was the difference between the temperature in Palm Springs and the temperature in Whitefield?

Exercise 1.5E.121

Football At the first down, the Chargers had the ball on their 25 yard line. On the next three downs, they lost 6 yards, gained 10 yards, and lost 8 yards. What was the yard line at the end of the fourth down?

Answer

21

Exercise 1.5E.122

Football At the first down, the Steelers had the ball on their 30 yard line. On the next three downs, they gained 9 yards, lost 14 yards, and lost 2 yards. What was the yard line at the end of the fourth down?

Exercise 1.5E.123

Checking Account Mayra has $124 in her checking account. She writes a check for $152. What is the new balance in her checking account?

Answer

−$28

Exercise 1.5E.124

Checking Account Selina has $165 in her checking account. She writes a check for $207. What is the new balance in her checking account?

Exercise 1.5E.125

Checking Account Diontre has a balance of −$38 in his checking account. He deposits $225 to the account. What is the new balance?

Answer

$187

Exercise 1.5E.126

Checking Account Reymonte has a balance of −$49 in his checking account. He deposits $281 to the account. What is the new balance?

Everyday Math

Exercise 1.5E.127

Stock market Javier owns 300 shares of stock in one company. On Tuesday, the stock price dropped $12 per share. What was the total effect on Javier's portfolio?

Answer

Weight loss In the first week of a diet program, eight women lost an average of 3 pounds each. What was the total weight change for the eight women?

Exercise 1.5E.128

Weight loss In the first week of a diet program, eight women lost an average of 3 pounds each. What was the total weight change for the eight women?

Writing Exercises

Exercise 1.5E.129

In your own words, state the rules for multiplying integers.

Answer

Answers may vary

Exercise 1.5E.130

In your own words, state the rules for dividing integers.

Exercise 1.5E.131

Why is $-2^4 \neq (-2)^4$?

Answer

Answers may vary

Exercise 1.5E.132

Why is $-4^3 \neq (-4)^3$?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
multiply integers.			
divide integers.			
simplify expressions with integers.			
evaluate variable expressions with integers.			
translate English phrases to algebraic expressions.			
use integers in applications.			

ⓑ On a scale of 1–10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 1.5E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.5E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.6: Visualize Fractions

Learning Objectives

By the end of this section, you will be able to:

- Find equivalent fractions
- Simplify fractions
- Multiply fractions
- Divide fractions
- Simplify expressions written with a fraction bar
- Translate phrases to expressions with fractions

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapter, **Fractions**.

Find Equivalent Fractions

Fractions are a way to represent parts of a whole. The fraction $\frac{1}{3}$ means that one whole has been divided into 3 equal parts and each part is one of the three equal parts. See Figure 1.6.1. The fraction $\frac{2}{3}$ represents two of three equal parts. In the fraction $\frac{2}{3}$, the 2 is called the **numerator** and the 3 is called the **denominator**.




Figure 1.6.1: The circle on the left has been divided into 3 equal parts. Each part is $\frac{1}{3}$ of the 3 equal parts. In the circle on the right, $\frac{2}{3}$ of the circle is shaded (2 of the 3 equal parts).

Doing the Manipulative Mathematics activity "Model Fractions" will help you develop a better understanding of fractions, their numerators and denominators.

FRACTION

A **fraction** is written $\frac{a}{b}$, where $b \neq 0$ and

- a is the **numerator** and b is the **denominator**.

A fraction represents parts of a whole. The denominator b is the number of equal parts the whole has been divided into, and the numerator a indicates how many parts are included.

If a whole pie has been cut into 6 pieces and we eat all 6 pieces, we ate $\frac{6}{6}$ pieces, or, in other words, one whole pie.



Figure 1.6.2

So $\frac{6}{6} = 1$. This leads us to the property of one that tells us that any number, except zero, divided by itself is 1.

PROPERTY OF ONE

$$\frac{a}{a} = 1 \quad (a \neq 0) \tag{1.6.1}$$

Any number, except zero, divided by itself is one.

Note

Doing the Manipulative Mathematics activity "Fractions Equivalent to One" will help you develop a better understanding of fractions that are equivalent to one.

If a pie was cut in 6 pieces and we ate all 6, we ate $\frac{6}{6}$ pieces, or, in other words, one whole pie. If the pie was cut into 8 pieces and we ate all 8, we ate $\frac{8}{8}$ pieces, or one whole pie. We ate the same amount—one whole pie.

The fractions $\frac{6}{6}$ and $\frac{8}{8}$ have the same value, 1, and so they are called equivalent fractions. **Equivalent fractions** are fractions that have the same value.

Let's think of pizzas this time. Figure 1.6.3 shows two images: a single pizza on the left, cut into two equal pieces, and a second pizza of the same size, cut into eight pieces on the right. This is a way to show that $\frac{1}{2}$ is equivalent to $\frac{4}{8}$. In other words, they are **equivalent fractions**.




Figure 1.6.3: Since the same amount is of each pizza is shaded, we see that $\frac{1}{2}$ is equivalent to $\frac{4}{8}$. They are equivalent fractions.

EQUIVALENT FRACTIONS

Equivalent fractions are fractions that have the same value.

How can we use mathematics to change $\frac{1}{2}$ into $\frac{4}{8}$? How could we take a pizza that is cut into 2 pieces and cut it into 8 pieces? We could cut each of the 2 larger pieces into 4 smaller pieces! The whole pizza would then be cut into 88 pieces instead of just 2. Mathematically, what we've described could be written like this as $\frac{1 \cdot 4}{2 \cdot 4} = \frac{4}{8}$. See Figure 1.6.4.




Figure 1.6.4: Cutting each half of the pizza into 4 pieces, gives us pizza cut into 8 pieces: $\frac{1 \cdot 4}{2 \cdot 4} = \frac{4}{8}$

This model leads to the following property:

EQUIVALENT FRACTIONS PROPERTY

If a, b, c are numbers where $b \neq 0, c \neq 0$, then

$$\frac{a}{b} = \frac{a \cdot c}{b \cdot c} \tag{1.6.2}$$

If we had cut the pizza differently, we could get

$$\frac{1\cdot 2}{2\cdot 2}=\frac{2}{4} \text{ so } \frac{1}{2}=\frac{2}{4}$$

$$\frac{1\cdot 3}{2\cdot 3}=\frac{3}{6} \text{ so } \frac{1}{2}=\frac{3}{6}$$

$$\frac{1\cdot 10}{2\cdot 10}=\frac{10}{20} \text{ so } \frac{1}{2}=\frac{10}{20}$$

Figure 1.6.5

So, we say $\frac{1}{2}$, $\frac{2}{4}$, $\frac{3}{6}$, and $\frac{10}{20}$ are equivalent fractions.

Note

Doing the Manipulative Mathematics activity "Equivalent Fractions" will help you develop a better understanding of what it means when two fractions are equivalent.

Example 1.6.1

Find three fractions equivalent to $\frac{2}{5}$.

Solution

To find a fraction equivalent to $\frac{2}{5}$, we multiply the numerator and denominator by the same number. We can choose any number, except for zero. Let's multiply them by 2, 3, and then 5.

$$\frac{2\cdot 2}{5\cdot 2}=\frac{4}{10} \quad \frac{2\cdot 3}{5\cdot 3}=\frac{6}{15} \quad \frac{2\cdot 5}{5\cdot 5}=\frac{10}{25}$$

So, $\frac{4}{10}$, $\frac{6}{15}$, and $\frac{10}{25}$ are equivalent to $\frac{2}{5}$.

Try It 1.6.2

Find three fractions equivalent to $\frac{3}{5}$.

Answer

$\frac{6}{10}$, $\frac{9}{15}$, $\frac{12}{20}$; answers may vary

Try It 1.6.3

Find three fractions equivalent to $\frac{4}{5}$.

Answer

$\frac{8}{10}$, $\frac{12}{15}$, $\frac{16}{20}$; answers may vary

Simplify Fractions

A fraction is considered **simplified** if there are no common factors, other than 1, in its numerator and denominator.

For example,

- $\frac{2}{3}$ is simplified because there are no common factors of 2 and 3.
- $\frac{10}{15}$ is not simplified because 5 is a common factor of 10 and 15.

SIMPLIFIED FRACTION

A fraction is considered **simplified** if there are no common factors in its numerator and denominator.

The phrase *reduce a fraction* means to simplify the fraction. We simplify, or reduce, a fraction by removing the common factors of the numerator and denominator. A fraction is not simplified until all common factors have been removed. If an expression has fractions, it is not completely simplified until the fractions are simplified.

In Exercise 1.6.4, we used the equivalent fractions property to find equivalent fractions. Now we'll use the equivalent fractions property in reverse to simplify fractions. We can rewrite the property to show both forms together.

EQUIVALENT FRACTIONS PROPERTY

If a, b, c are numbers where $b \neq 0, c \neq 0$,

$$\text{then } \frac{a}{b} = \frac{a \cdot c}{b \cdot c} \text{ and } \frac{a \cdot c}{b \cdot c} = \frac{a}{b} \tag{1.6.3}$$

Example 1.6.4

Simplify: $-\frac{32}{56}$

Solution

	$-\frac{32}{56}$
Rewrite the numerator and denominator showing the common factors.	$-\frac{4 \cdot 8}{7 \cdot 8}$
Simplify using the equivalent fractions property.	$-\frac{4}{7}$

Notice that the fraction $-\frac{4}{7}$ is simplified because there are no more common factors.

p

Try It 1.6.5

Simplify: $-\frac{42}{54}$

Answer

$-\frac{7}{9}$

Try It 1.6.6

Simplify: $-\frac{30}{54}$

Answer

$-\frac{5}{9}$

Sometimes it may not be easy to find common factors of the numerator and denominator. When this happens, a good idea is to factor the numerator and the denominator into **prime numbers**. Then divide out the common factors using the equivalent fractions property.

Example 1.6.7

Simplify: $-\frac{210}{385}$

Solution

Step 1. Rewrite the numerator and denominator to show the common factors. If needed, factor the numerator and denominator into prime numbers first.	Rewrite 210 and 385 as the product of the primes.	$-\frac{210}{385}$ $-\frac{2\cdot 3\cdot 5\cdot 7}{5\cdot 7\cdot 11}$
Step 2. Simplify using the equivalent fractions property by dividing out common factors.	Mark the common factors 5 and 7. Divide out the common factors.	$-\frac{2\cdot 3\cdot \cancel{5}\cdot \cancel{7}}{\cancel{5}\cdot \cancel{7}\cdot 11}$ $-\frac{2\cdot 3}{11}$
Step 3. Multiplify the remaining factors, if necessary.		$-\frac{6}{11}$

Try It 1.6.8

Simplify: $-\frac{69}{120}$

Answer

$-\frac{23}{40}$

Try It 1.6.9

Simplify: $-\frac{120}{192}$

Answer

$-\frac{5}{8}$

We now summarize the steps you should follow to simplify fractions.

SIMPLIFY A FRACTION.

1. Rewrite the numerator and denominator to show the common factors.
   If needed, factor the numerator and denominator into prime numbers first.
2. Simplify using the equivalent fractions property by dividing out common factors.
3. Multiply any remaining factors, if needed.

Example 1.6.10

Simplify: $\frac{5x}{5y}$

Solution

	$\frac{5x}{5y}$

Rewrite showing the common factors, then divide out the common factors.	Rewrite showing the common factors, then divide out the common factors. $\frac{5x}{5y}$ $\frac{5 \cdot x}{5 \cdot y}$ Simplify. $\frac{x}{y}$
Simplify.	$\frac{x}{y}$

Try It 1.6.11

Simplify: $\frac{7x}{7y}$

Answer

$\frac{x}{y}$

Try It 1.6.12

Simplify: $\frac{3a}{3b}$

Answer

$\frac{a}{b}$

Multiply Fractions

Many people find multiplying and dividing fractions easier than adding and subtracting fractions. So we will start with fraction multiplication.

Doing the Manipulative Mathematics activity "Model Fraction Multiplication" will help you develop a better understanding of multiplying fractions.

We'll use a model to show you how to multiply two fractions and to help you remember the procedure. Let's start with $\frac{3}{4}$.



Figure 1.6.6

Now we'll take $\frac{1}{2}$ of $\frac{3}{4}$.



Figure 1.6.6

Notice that now, the whole is divided into 8 equal parts. So $\frac{1}{2} \cdot \frac{3}{4} = \frac{3}{8}$.

To multiply fractions, we multiply the numerators and multiply the denominators.

FRACTION MULTIPLICATION

If a, b, c and d are numbers where $b \neq 0$ and $d \neq 0$, then

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd} \tag{1.6.4}$$

To multiply fractions, multiply the numerators and multiply the denominators.

When **multiplying fractions**, the properties of positive and negative numbers still apply, of course. It is a good idea to determine the sign of the **product** as the first step. In Exercise 1.6.13, we will multiply negative and a positive, so the product will be negative.

Example 1.6.13

Multiply: $-\frac{11}{12} \cdot \frac{5}{7}$

Solution

The first step is to find the sign of the product. Since the signs are the different, the product is negative.

$$\begin{aligned} & -\frac{11}{12} \cdot \frac{5}{7} \\ \text{Determine the sign of the product; multiply.} \quad & -\frac{11 \cdot 5}{12 \cdot 7} \\ \text{Are there any common factors in the numerator} & \\ \text{and the denominator? No} \quad & -\frac{55}{84} \end{aligned} \tag{1.6.5}$$

Try It 1.6.14

Multiply: $-\frac{10}{28} \cdot \frac{8}{15}$

Answer

$-\frac{4}{21}$

Try It 1.6.15

Multiply: $-\frac{9}{20} \cdot \frac{5}{12}$

Answer

$-\frac{3}{16}$

When multiplying a fraction by an integer, it may be helpful to write the integer as a fraction. Any integer, a, can be written as $\frac{a}{1}$. So, for example, $3 = \frac{3}{1}$.

Example 1.6.16

Multiply: $-\frac{12}{5}(-20x)$

Solution

Determine the sign of the product. The signs are the same, so the product is positive.

	$-\frac{12}{5}(-20x)$
Write $20x$ as a fraction.	$\frac{12}{5}(\frac{20x}{1})$
Multiply.	
Rewrite 20 to show the common factor 5 and divide it out.	$\frac{12 \cdot 4 \cdot \cancel{5}x}{\cancel{5} \cdot 1}$
Simplify.	$48x$

Try It 1.6.17

Multiply: $\frac{11}{3}(-9a)$

Answer

$-33a$

Try It 1.6.18

Multiply: $\frac{13}{7}(-14b)$

Answer

$-26b$

Divide Fractions

Now that we know how to multiply fractions, we are almost ready to divide. Before we can do that, that we need some vocabulary.

The **reciprocal** of a fraction is found by inverting the fraction, placing the numerator in the denominator and the denominator in the numerator. The reciprocal of $\frac{2}{3}$ is $\frac{3}{2}$.

Notice that $\frac{2}{3} \cdot \frac{3}{2} = 1$. A number and its reciprocal multiply to 1.

To get a **product** of positive 1 when multiplying two numbers, the numbers must have the same sign. So reciprocals must have the same sign.

The reciprocal of $-\frac{10}{7}$ is $-\frac{7}{10}$, since $-\frac{10}{7}(-\frac{7}{10}) = 1$.

RECIPROCAL

The **reciprocal** of $\frac{a}{b}$ is $\frac{b}{a}$.

A number and its reciprocal multiply to one $\frac{a}{b} \cdot \frac{b}{a} = 1$

Note

Doing the Manipulative Mathematics activity “Model Fraction Division” will help you develop a better understanding of dividing fractions.

To divide fractions, we multiply the first fraction by the reciprocal of the second.

FRACTION DIVISION

If a, b, c and d are numbers where $b \neq 0, c \neq 0$ and $d \neq 0$, then

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} \tag{1.6.6}$$

To divide fractions, we multiply the first fraction by the reciprocal of the second.

We need to say $b \neq 0, c \neq 0$ and $d \neq 0$ to be sure we don't divide by zero!

Example 1.6.19

Divide: $-\frac{2}{3} \div \frac{n}{5}$

Solution

	$-\frac{2}{3} \div \frac{n}{5}$
To divide, multiply the first fraction by the reciprocal of the second.	$-\frac{2}{3} \cdot \frac{5}{n}$
Multiply.	$-\frac{10}{3n}$

(1.6.7)

Try It 1.6.20

Divide: $-\frac{3}{5} \div \frac{p}{7}$.

Answer

$-\frac{21}{5p}$

Try It 1.6.21

Divide: $-\frac{5}{8} \div \frac{q}{3}$.

Answer

$-\frac{15}{8q}$

Example 1.6.22

Find the quotient:

$-\frac{7}{18} \div (-\frac{14}{27})$

Solution

	$-\frac{7}{18} \div (-\frac{14}{27})$

To divide, multiply the first fraction by the reciprocal of the second.	$-\frac{7}{18} \cdot -\frac{27}{14}$
Determine the sign of the product, and then multiply..	$\frac{7 \cdot 27}{18 \cdot 14}$
Rewrite showing common factors.	$\frac{7 \cdot 9 \cdot 3}{9 \cdot 2 \cdot 7 \cdot 2}$
Remove common factors.	$\frac{3}{2 \cdot 2}$
Simplify.	$\frac{3}{4}$

Try It 1.6.23

Find the quotient:

$-\frac{7}{8} \div (-\frac{14}{27})$

Answer

$\frac{4}{15}$

Try It 1.6.24

Find the quotient:

$-\frac{7}{8} \div (-\frac{14}{27})$

Answer

$\frac{2}{3}$

There are several ways to remember which steps to take to multiply or divide fractions. One way is to repeat the call outs to yourself. If you do this each time you do an exercise, you will have the steps memorized.

- “To multiply fractions, multiply the numerators and multiply the denominators.”
- “To divide fractions, multiply the first fraction by the reciprocal of the second.”

Another way is to keep two examples in mind:

One fourth of two pizzas is one half of a pizza.



$$2 \cdot \frac{1}{4}$$

$$\frac{2}{1} \cdot \frac{1}{4}$$

$$\frac{2}{4}$$

$$\frac{1}{2}$$

There are eight quarters in $2.00.



$$2 \div \frac{1}{4}$$

$$\frac{2}{1} \div \frac{1}{4}$$

$$\frac{2}{1} \cdot \frac{4}{1}$$

$$8$$

Figure 1.6.7

The numerators or denominators of some fractions contain fractions themselves. A fraction in which the numerator or the denominator is a fraction is called a **complex fraction**.

COMPLEX FRACTION

A **complex fraction** is a fraction in which the numerator or the denominator contains a fraction.

Some examples of complex fractions are:

$$\frac{\frac{6}{7}}{3} \quad \frac{\frac{3}{4}}{\frac{5}{8}} \quad \frac{\frac{x}{2}}{\frac{5}{6}} \tag{1.6.8}$$

To simplify a complex fraction, we remember that the fraction bar means **division**. For example, the complex fraction $\frac{\frac{3}{4}}{\frac{5}{8}}$ means $\frac{3}{4} \div \frac{5}{8}$.

Example 1.6.25

Simplify: $\frac{\frac{3}{4}}{\frac{5}{8}}$

Solution

	$\frac{\frac{3}{4}}{\frac{5}{8}}$
Rewrite as division.	$\frac{3}{4} \div \frac{5}{8}$
Multiply the first fraction by the reciprocal of the second.	$\frac{3}{4} \cdot \frac{8}{5}$
Multiply.	$\frac{3 \cdot 8}{4 \cdot 5}$
Look for common factors.	$\frac{3 \cdot \cancel{4} \cdot 2}{\cancel{4} \cdot 5}$
Divide out common factors and simplify.	$\frac{6}{5}$

Try It 1.6.26

Simplify: $\frac{\frac{2}{3}}{\frac{5}{6}}$

Answer

$\frac{4}{5}$

Try It 1.6.27

Simplify: $\frac{\frac{3}{7}}{\frac{6}{11}}$

Answer

$\frac{11}{14}$

Example 1.6.28

Simplify: $\frac{\frac{x}{2}}{\frac{xy}{6}}$

Solution

	$\frac{\frac{x}{2}}{\frac{xy}{6}}$
Rewrite as division.	$\frac{x}{2} \div \frac{xy}{6}$
Multiply the first fraction by the reciprocal of the second.	$\frac{x}{2} \cdot \frac{6}{xy}$
Multiply.	$\frac{x \cdot 6}{2 \cdot xy}$
Look for common factors.	$\frac{x \cdot 3 \cdot 2}{2 \cdot x \cdot y}$
Divide out common factors and simplify.	$\frac{3}{y}$

Try It 1.6.29

Simplify: $\frac{\frac{a}{8}}{\frac{ab}{6}}$

Answer

$\frac{3}{4b}$

Try It 1.6.30

Simplify: $\frac{\frac{p}{2}}{\frac{pq}{8}}$

Answer

$\frac{4}{q}$

Simplify Expressions with a Fraction Bar

The line that separates the numerator from the denominator in a fraction is called a fraction bar. A **fraction bar** acts as grouping symbol. The order of operations then tells us to simplify the numerator and then the denominator. Then we divide.

To simplify the expression $\frac{5-3}{7+1}$, we first simplify the numerator and the denominator separately. Then we divide.

$$\begin{array}{l} \frac{5-3}{7+1} \\ \frac{2}{8} \\ \frac{1}{4} \end{array} \tag{1.6.9}$$

SIMPLIFY AN EXPRESSION WITH A FRACTION BAR.

1. Simplify the expression in the numerator. Simplify the expression in the denominator.
2. Simplify the fraction.

Example 1.6.31

Simplify: $\frac{4-2(3)}{2^2+2}$

Solution

$$\begin{array}{ll} & \frac{4-2(3)}{2^2+2} \\ \text{Use the order of operations to simplify the numerator and the denominator.} & \frac{4-6}{4+2} \\ \text{Simplify the numerator and the denominator} & \frac{-2}{6} \\ \text{Simplify. A negative divided by a positive is negative.} & -\frac{1}{3} \end{array} \tag{1.6.10}$$

Try It 1.6.32

Simplify: $\frac{6-3(5)}{3^2+3}$

Answer

$-\frac{3}{4}$

Try It 1.6.33

Simplify: $\dfrac{4-4(6)}{3^2+3}$

Answer

$-\dfrac{5}{3}$

Where does the negative sign go in a fraction? Usually the negative sign is in front of the fraction, but you will sometimes see a fraction with a negative numerator, or sometimes with a negative denominator. Remember that fractions represent division. When the numerator and denominator have different signs, the quotient is negative.

$$\begin{aligned} \tfrac{-1}{3} &= -\tfrac{1}{3} \quad \tfrac{\text{negative}}{\text{positive}} = \text{negative} \\ \tfrac{1}{-3} &= -\tfrac{1}{3} \quad \tfrac{\text{positive}}{\text{negative}} = \text{negative} \end{aligned} \tag{1.6.11}$$

PLACEMENT OF NEGATIVE SIGN IN A FRACTION

For any positive numbers a and b,

$$\frac{-a}{b} = \frac{a}{-b} = -\frac{a}{b} \tag{1.6.12}$$

Example 1.6.34

Simplify: $\frac{4(-3)+6(-2)}{-3(2)-2}$

Solution

The fraction bar acts like a grouping symbol. So completely simplify the numerator and the denominator separately.

$$\begin{aligned} & \tfrac{4(-3)+6(-2)}{-3(2)-2} \\ \text{Multiply.} \quad & \tfrac{-12+(-12)}{-6-2} \\ \text{Simplify.} \quad & \tfrac{-24}{-8} \\ \text{Divide.} \quad & 3 \end{aligned} \tag{1.6.13}$$

Try It 1.6.35

Simplify: $\frac{8(-2)+4(-3)}{-5(2)+3}$

Answer

4

Try It 1.6.36

Simplify: $\frac{7(-1)+9(-3)}{-5(3)-2}$

Answer

2

Translate Phrases to Expressions with Fractions

Now that we have done some work with fractions, we are ready to translate phrases that would result in expressions with fractions.

The English words **quotient** and **ratio** are often used to describe fractions. Remember that "quotient" means **division**. The quotient of aa and bb is the result we get from dividing a by b, or $\frac{a}{b}$.

Example 1.6.37

Translate the English phrase into an algebraic expression: the quotient of the difference of m and n, and p.

Solution

We are looking for the *quotient of* the difference of m and n, and p. This means we want to divide the difference of m and n, and p.

$$\frac{m-n}{p} \tag{1.6.14}$$

Try It 1.6.38

Translate the English phrase into an algebraic expression: the quotient of the difference of a and b, and cd.

Answer

$\frac{a-b}{cd}$

Try It 1.6.39

Translate the English phrase into an algebraic expression: the quotient of the sum of p and q, and r.

Answer

$\frac{p+q}{r}$

Key Concepts

- **Equivalent Fractions Property:** If a, b, c are numbers where $b \neq 0, c \neq 0$, then $\frac{a}{b} = \frac{a \cdot c}{b \cdot c}$ and $\frac{a \cdot c}{b \cdot c} = \frac{a}{b}$
- **Fraction Division:** If a, b, c and d are numbers where $b \neq 0, c \neq 0$ and $d \neq 0$, then $\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c}$. To divide fractions, multiply the first fraction by the reciprocal of the second.
- **Fraction Multiplication:** If a, b, c and d are numbers where $b \neq 0, d \neq 0$, then $\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$. To multiply fractions, multiply the numerators and multiply the denominators.
- **Placement of Negative Sign in a Fraction:** For any positive numbers a and b, $\frac{-a}{a} = \frac{a}{-a} = -\frac{a}{b}$
- **Property of One:** $\frac{a}{a} = 1$; Any number, except zero, divided by itself is one.
- **Simplify a Fraction**
  1. Rewrite the numerator and denominator to show the common factors. If needed, factor the numerator and denominator into prime numbers first.
  2. Simplify using the equivalent fractions property by dividing out common factors.
  3. Multiply any remaining factors.
- **Simplify an Expression with a Fraction Bar**
  1. Simplify the expression in the numerator. Simplify the expression in the denominator.
  2. Simplify the fraction.

Glossary

complex fraction

A complex fraction is a fraction in which the numerator or the denominator contains a fraction.

denominator

The denominator is the value on the bottom part of the fraction that indicates the number of equal parts into which the whole has been divided.

equivalent fractions

Equivalent fractions are fractions that have the same value.

fraction

A fraction is written $\frac{a}{b}$, where $b \neq 0$, a is the numerator and b is the denominator. A fraction represents parts of a whole. The denominator b is the number of equal parts the whole has been divided into, and the numerator aa indicates how many parts are included.

numerator

The numerator is the value on the top part of the fraction that indicates how many parts of the whole are included.

reciprocal

The reciprocal of $\frac{a}{b}$ is $\frac{b}{a}$. A number and its reciprocal multiply to one: $\frac{a}{b} \cdot \frac{b}{a} = 1$.

simplified fraction

A fraction is considered simplified if there are no common factors in its numerator and denominator.

This page titled 1.6: Visualize Fractions is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.6: Visualize Fractions** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.6E: Exercises

Practice Makes Perfect

Find Equivalent Fractions

In the following exercises, find three fractions equivalent to the given fraction. Show your work, using figures or algebra.

Exercise 1.6E.1

$\frac{3}{8}$

Answer

$\frac{6}{16}, \frac{9}{24}, \frac{12}{32}$, answers may vary

Exercise 1.6E.2

$\frac{5}{8}$

Exercise 1.6E.3

$\frac{5}{9}$

Answer

$\frac{10}{18}, \frac{15}{27}, \frac{20}{36}$, answers may vary

Exercise 1.6E.4

$\frac{1}{8}$

Simplify Fractions

In the following exercises, simplify.

Exercise 1.6E.5

$-\frac{40}{88}$

Answer

$-\frac{5}{11}$

Exercise 1.6E.6

$-\frac{63}{99}$

Exercise 1.6E.7

$$-\frac{108}{63}$$

Answer

$$-\frac{12}{7}$$

Exercise 1.6E.8

$$-\frac{104}{48}$$

Exercise 1.6E.9

$$\frac{120}{252}$$

Answer

$$\frac{10}{21}$$

Exercise 1.6E.10

$$\frac{182}{294}$$

Exercise 1.6E.11

$$-\frac{3x}{12y}$$

Answer

$$-\frac{x}{4y}$$

Exercise 1.6E.12

$$-\frac{4x}{32y}$$

Exercise 1.6E.13

$$\frac{14x^2}{21y}$$

Answer

$$\frac{2x^2}{3y}$$

Exercise 1.6E.14

$$\frac{24a}{32b^2}$$

Multiply Fractions

In the following exercises, multiply.

Exercise 1.6E.15

$$\frac{3}{4} \cdot \frac{9}{10}$$

Answer

$$\frac{27}{40}$$

Exercise 1.6E.16

$$\frac{4}{5} \cdot \frac{2}{7}$$

Exercise 1.6E.17

$$-\frac{2}{3} \cdot -\frac{3}{8}$$

Answer

$$\frac{1}{4}$$

Exercise 1.6E.18

$$-\frac{3}{4}\left(-\frac{4}{9}\right)$$

Exercise 1.6E.19

$$-\frac{5}{9} \cdot \frac{3}{10}$$

Answer

$$-\frac{1}{6}$$

Exercise 1.6E.20

$$-\frac{3}{8} \cdot \frac{4}{15}$$

Exercise 1.6E.21

$$\left(-\frac{14}{15}\right)\left(\frac{9}{20}\right)$$

Answer

$$-\frac{21}{50}$$

Exercise 1.6E.22

$$\left(-\frac{9}{10}\right)\left(\frac{25}{33}\right)$$

Exercise 1.6E.23

$$\left(-\frac{63}{84}\right)\left(-\frac{44}{90}\right)$$

Answer

$$\frac{11}{30}$$

Exercise 1.6E.24

$$\left(-\frac{63}{60}\right)\left(-\frac{40}{88}\right)$$

Exercise 1.6E.25

$$4 \cdot \frac{5}{11}$$

Answer

$$\frac{20}{11}$$

Exercise 1.6E.26

$$5 \cdot \frac{8}{3}$$

Exercise 1.6E.27

$$\frac{3}{7} \cdot 21n$$

Answer

9n

Exercise 1.6E.28

$$\frac{5}{6} \cdot 30m$$

Exercise 1.6E.29

$-8 \cdot \frac{17}{4}$

Answer

-34

Exercise 1.6E.30

$(-1)\left(-\frac{6}{7}\right)$

Divide Fractions

In the following exercises, divide.

Exercise 1.6E.31

$\frac{3}{4} \div \frac{2}{3}$

Answer

$\frac{9}{8}$

Exercise 1.6E.32

$\frac{4}{5} \div \frac{3}{4}$

Exercise 1.6E.33

$-\frac{7}{9} \div \left(-\frac{7}{4}\right)$

Answer

$\frac{4}{9}$

Exercise 1.6E.34

$-\frac{5}{6} \div \left(-\frac{5}{6}\right)$

Exercise 1.6E.35

$\frac{3}{4} \div \frac{x}{11}$

Answer

$\frac{33}{4x}$

Exercise 1.6E.36

$\frac{2}{5} \div \frac{y}{9}$

Exercise 1.6E.37

$\frac{5}{18} \div -\frac{15}{24}$

Answer

$-\frac{4}{9}$

Exercise 1.6E.38

$\frac{7}{18} \div \left(-\frac{14}{27}\right)$

Exercise 1.6E.39

$\frac{8u}{15} \div \frac{12v}{25}$

Answer

$\frac{10u}{9v}$

Exercise 1.6E.40

$\frac{12r}{25} \div \frac{18s}{35}$

Exercise 1.6E.41

$-5 \div \frac{1}{2}$

Answer

-10

Exercise 1.6E.42

$-3 \div \frac{1}{4}$

Exercise 1.6E.43

$\frac{3}{4} \div (-12)$

Answer

$\frac{1}{16}$

Exercise 1.6E.44

$-15 \div -\frac{5}{3}$

In the following exercises, simplify.

Exercise 1.6E.45

$$\frac{-\frac{8}{21}}{\frac{12}{35}}$$

Answer

$-\frac{10}{9}$

Exercise 1.6E.46

$$\frac{-\frac{9}{16}}{\frac{33}{40}}$$

Exercise 1.6E.47

$$\frac{-\frac{4}{5}}{2}$$

Answer

$-\frac{2}{5}$

Exercise 1.6E.48

$$\frac{5}{\frac{3}{10}}$$

Exercise 1.6E.49

$$\frac{\frac{m}{3}}{\frac{n}{2}}$$

Answer

$\frac{2m}{3n}$

Exercise 1.6E.50

$$\frac{-\frac{3}{8}}{-\frac{y}{12}}$$

Simplify Expressions Written with a Fraction Bar

In the following exercises, simplify.

Exercise 1.6E.51

$$\frac{22+3}{10}$$

Answer

$$\frac{5}{2}$$

Exercise 1.6E.52

$$\frac{19-4}{6}$$

Exercise 1.6E.53

$$\frac{48}{24-15}$$

Answer

$$\frac{16}{3}$$

Exercise 1.6E.54

$$\frac{46}{4+4}$$

Exercise 1.6E.55

$$\frac{-6+6}{8+4}$$

Answer

$$0$$

Exercise 1.6E.56

$$\frac{-6+3}{17-8}$$

Exercise 1.6E.57

$$\frac{4 \cdot 3}{6 \cdot 6}$$

Answer

$$\frac{1}{3}$$

Exercise 1.6E.58

$$\frac{6 \cdot 6}{9 \cdot 2}$$

Exercise 1.6E.59

$$\frac{4^2 - 1}{25}$$

Answer

$$\frac{3}{5}$$

Exercise 1.6E.60

$$\frac{7^2 + 1}{60}$$

Exercise 1.6E.61

$$\frac{8 \cdot 3 + 2 \cdot 9}{14 + 3}$$

Answer

$$\frac{42}{17}$$

Exercise 1.6E.62

$$\frac{9 \cdot 6 - 4 \cdot 7}{22 + 3}$$

Exercise 1.6E.63

$$\frac{5 \cdot 6 - 3 \cdot 4}{4 \cdot 5 - 2 \cdot 3}$$

Answer

$$\frac{9}{7}$$

Exercise 1.6E.64

$$\frac{8 \cdot 9 - 7 \cdot 6}{5 \cdot 6 - 9 \cdot 2}$$

Exercise 1.6E.65

$$\frac{5^2 - 3^2}{3 - 5}$$

Answer

-8

Exercise 1.6E.66

$$\frac{6^2 - 4^2}{4 - 6}$$

Exercise 1.6E.67

$$\frac{7 \cdot 4 - 2(8 - 5)}{9 \cdot 3 - 3 \cdot 5}$$

Answer

$$\frac{11}{6}$$

Exercise 1.6E.68

$$\frac{9 \cdot 7 - 3(12 - 8)}{8 \cdot 7 - 6 \cdot 6}$$

Exercise 1.6E.69

$$\frac{9(8 - 2) - 3(15 - 7)}{6(7 - 1) - 3(17 - 9)}$$

Answer

$$\frac{5}{2}$$

Exercise 1.6E.70

$$\frac{8(9 - 2) - 4(14 - 9)}{7(8 - 3) - 3(16 - 9)}$$

Translate Phrases to Expressions with Fractions

In the following exercises, translate each English phrase into an algebraic expression.

Exercise 1.6E.71

the quotient of r and the sum of s and 10

Answer

$$\frac{r}{s+10}$$

Exercise 1.6E.72

the quotient of A and the difference of 3 and B

Exercise 1.6E.73

the quotient of the difference of x and y, and -3

Answer

$$\frac{x-y}{-3}$$

Exercise 1.6E.74

the quotient of the sum of m and n, and $4q$

Everyday Math

Exercise 1.6E.75

Baking. A recipe for chocolate chip cookies calls for $\frac{3}{4}$ cup brown sugar. Imelda wants to double the recipe.

1. How much brown sugar will Imelda need? Show your calculation.
2. Measuring cups usually come in sets of $\frac{1}{4}$, $\frac{1}{3}$, $\frac{1}{2}$, and 1 cup. Draw a diagram to show two different ways that Imelda could measure the brown sugar needed to double the cookie recipe.

Answer

1. $1\frac{1}{2}$ cups
2. answers will vary

Exercise 1.6E.76

Baking. Nina is making 4 pans of fudge to serve after a music recital. For each pan, she needs $\frac{2}{3}$ cup of condensed milk.

1. How much condensed milk will Nina need? Show your calculation.
2. Measuring cups usually come in sets of $\frac{1}{4}$, $\frac{1}{3}$, $\frac{1}{2}$, and 1 cup. Draw a diagram to show two different ways that Nina could measure the condensed milk needed for 4 pans of fudge.

Exercise 1.6E.77

Portions Don purchased a bulk package of candy that weighs 5 pounds. He wants to sell the candy in little bags that hold $\frac{1}{4}$ pound. How many little bags of candy can he fill from the bulk package?

Answer

20 bags

Exercise 1.6E.78

Portions Kristen has $\frac{3}{4}$ yards of ribbon that she wants to cut into 6 equal parts to make hair ribbons for her daughter's 6 dolls. How long will each doll's hair ribbon be?

Writing Exercises

Exercise 1.6E.79

Rafael wanted to order half a medium pizza at a restaurant. The waiter told him that a medium pizza could be cut into 6 or 8 slices. Would he prefer 3 out of 6 slices or 4 out of 8 slices? Rafael replied that since he wasn't very hungry, he would prefer 3 out of 6 slices. Explain what is wrong with Rafael's reasoning.

Answer

Answers may vary

Exercise 1.6E.80

Give an example from everyday life that demonstrates how $\frac{1}{2} \cdot \frac{2}{3}$ is $\frac{1}{3}$.

Exercise 1.6E.81

Explain how you find the reciprocal of a fraction.

Answer

Answers may vary

Exercise 1.6E.82

Explain how you find the reciprocal of a negative number.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
find equivalent fractions.			
simplify fractions.			
multiply fractions.			
divide fractions.			
simplify expressions written with a fraction bar.			
translate phrases to expressions with fractions.			

ⓑ After looking at the checklist, do you think you are well prepared for the next section? Why or why not?

This page titled 1.6E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.6E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.7: Add and Subtract Fractions

Learning Objectives

By the end of this section, you will be able to:

- Add or subtract fractions with a common denominator
- Add or subtract fractions with different denominators
- Use the order of operations to simplify complex fractions
- Evaluate variable expressions with fractions

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra*chapter, **Fractions**.

Add or Subtract Fractions with a Common Denominator

When we multiplied fractions, we just multiplied the numerators and multiplied the denominators right straight across. To add or subtract fractions, they must have a common denominator.

FRACTION ADDITION AND SUBTRACTION

If a, b, and c are numbers where $c \neq 0$, then

$$\frac{a}{c}+\frac{b}{c}=\frac{a+b}{c} \quad \text{and} \quad \frac{a}{c}-\frac{b}{c}=\frac{a-b}{c} \tag{1.7.1}$$

To add or subtract fractions, add or subtract the numerators and place the result over the common denominator.

Manipulative Mathematics

Doing the Manipulative Mathematics activities “Model Fraction Addition” and “Model Fraction Subtraction” will help you develop a better understanding of adding and subtracting fractions.

Example 1.7.1

Find the sum: $\frac{x}{3}+\frac{2}{3}$.

Solution

$$\begin{array}{ll} & \frac{x}{3}+\frac{2}{3} \\ \text{Add the numerators and place the sum over the common denominator} & \frac{x+2}{3} \end{array} \tag{1.7.2}$$

Try It 1.7.2

Find the sum: $\frac{x}{4}+\frac{3}{4}$.

Answer

$\frac{x+3}{4}$

Try It 1.7.3

Find the sum: $\frac{y}{8}+\frac{5}{8}$.

Answer

$\frac{y+5}{8}$

Example 1.7.4

Find the difference: $-\frac{23}{24}-\frac{13}{24}$

Solution

	$-\frac{23}{24}-\frac{13}{24}$
Subtract the numerators and place the difference over the common denominator	$\frac{-23-13}{24}$ (1.7.3)
Simplify.	$\frac{-36}{24}$
Simplify. Remember, $-\frac{a}{b}=\frac{-a}{b}$	$-\frac{3}{2}$

Try It 1.7.5

Find the difference: $-\frac{19}{28}-\frac{7}{28}$

Answer

$-\frac{26}{28}$

Try It 1.7.6

Find the difference: $-\frac{27}{32}-\frac{1}{32}$

Answer

$-\frac{7}{8}$

Example 1.7.7

Find the difference: $-\frac{10}{x}-\frac{4}{x}$

Solution

	$-\frac{10}{x}-\frac{4}{x}$	
Subtract the numerators and place the difference over the common denominator	$\frac{-14}{x}$	(1.7.4)
Rewrite with the sign in front of the fraction.	$-\frac{14}{x}$	

Try It 1.7.8

Find the difference: $-\frac{9}{x}-\frac{7}{x}$

Answer

$-\frac{16}{x}$

Try It 1.7.9

Find the difference: $-\frac{17}{a}-\frac{5}{a}$

Answer

$-\frac{22}{a}$

Now we will do an example that has both addition and subtraction.

Example 1.7.10

Simplify: $\frac{3}{8}+(-\frac{5}{8})-\frac{1}{8}$

Solution

Add and Subtract fractions — do they have a common denominator? Yes.	$\frac{3}{8}+(-\frac{5}{8})-\frac{1}{8}$	
Add and subtract the numerators and place the result over the common denominator.	$\frac{3+(-5)-1}{8}$	(1.7.5)
Simplify left to right.	$\frac{-2-1}{8}$	
Simplify.	$-\frac{3}{8}$	

Try It 1.7.11

Simplify: $\frac{2}{9}+(-\frac{4}{9})-\frac{7}{9}$

Answer

-1

Try It 1.7.12

Simplify: $\frac{2}{5}+(-\frac{4}{9})-\frac{7}{9}$

Answer

$-\frac{2}{3}$

Add or Subtract Fractions with Different Denominators

As we have seen, to add or subtract fractions, their denominators must be the same. The **least common denominator** (LCD) of two fractions is the smallest number that can be used as a common **denominator** of the fractions. The LCD of the two fractions is the least common multiple (LCM) of their denominators.

LEAST COMMON DENOMINATOR

The least common denominator (LCD) of two fractions is the least common multiple (LCM) of their denominators.

Note

Doing the Manipulative Mathematics activity "Finding the Least Common Denominator" will help you develop a better understanding of the LCD.

After we find the least common denominator of two fractions, we convert the fractions to equivalent fractions with the LCD. Putting these steps together allows us to add and subtract fractions because their denominators will be the same!

Example 1.7.13

Add: $\frac{7}{12}+\frac{5}{18}$

Solution

Step 1. Do they have a common denominator? No—rewrite each fraction with the LCD (least common denominator).	No. Find the LCD of 12, 18. Change into equivalent fractions with the LCD, 36. Do not simplify the equivalent fractions! If you do, you'll get back to the original fractions and lose the common denominator!	$12=2\cdot2\cdot3$ $18=2\cdot\quad3\cdot3$ $\text{LCD}=2\cdot2\cdot3\cdot3$ $\text{LCD}=36$ $\frac{7}{12}+\frac{5}{18}$ $\frac{7\cdot3}{12\cdot3}+\frac{5\cdot2}{18\cdot2}$ $\frac{21}{36}+\frac{10}{36}$
Step 2. Add or subtract the fractions.	Add.	$\frac{31}{36}$
Step 3. Simplify, if possible.	Because 31 is a prime number, it has no factors in common with 36. The answer is simplified.	

Try It 1.7.14

Add: $\frac{7}{12}+\frac{11}{15}$

Answer

$\frac{79}{60}$

Try It 1.7.15

Add: $\frac{13}{15}+\frac{17}{20}$

Answer

$\frac{103}{60}$

ADD OR SUBTRACT FRACTIONS.

1. Do they have a common denominator?
   - Yes—go to step 2.
   - No—rewrite each fraction with the LCD (least common denominator). Find the LCD. Change each fraction into an equivalent fraction with the LCD as its denominator.
2. Add or subtract the fractions.
3. Simplify, if possible.

When finding the equivalent fractions needed to create the common denominators, there is a quick way to find the number we need to multiply both the numerator and denominator. This method works if we found the LCD by factoring into primes.

Look at the factors of the LCD and then at each column above those factors. The "missing" factors of each denominator are the numbers we need.




Figure: 1.7.1

In Exercise 1.7.13, the LCD, 36, has two factors of 2 and two factors of 3.

The numerator 12 has two factors of 2 but only one of 3—so it is "missing" one 3—we multiply the numerator and denominator by 3.

The numerator 18 is missing one factor of 2—so we multiply the numerator and denominator by 2.

We will apply this method as we subtract the fractions in Exercise 1.7.16.

Example 1.7.16

Subtract: $\frac{7}{15}-\frac{19}{24}$

Solution

Do the fractions have a common denominator? No, so we need to find the LCD.

Find the LCD. $\frac{7}{15}-\frac{19}{24}$ $15=3\cdot 5$ $24=2\cdot 2\cdot 2\cdot 3$ $LCD=2\cdot 2\cdot 2\cdot 3\cdot 5$ $LCD=120$	
Notice, 15 is “missing” three factors of 2 and 24 is “missing” the 5 from the factors of the LCD. So we multiply 8 in the first fraction and 5 in the second fraction to get the LCD.	
Rewrite as equivalent fractions with the LCD.	$\frac{7\cdot 8}{15\cdot 8}-\frac{19\cdot 5}{24\cdot 5}$
Simplify.	$\frac{56}{120}-\frac{95}{120}$
Subtract.	$-\frac{39}{120}$
Check to see if the answer can be simplified.	$-\frac{13\cdot 3}{40\cdot 3}$
Both 39 and 120 have a factor of 3.	
Simplify.	$-\frac{13}{40}$

Do not simplify the equivalent fractions! If you do, you’ll get back to the original fractions and lose the common denominator!

Try It 1.7.17

Subtract: $\frac{13}{24}-\frac{17}{32}$

Answer

$\frac{1}{96}$

Try It 1.7.18

Subtract: $\frac{7}{15}-\frac{19}{24}$

Answer

$\frac{75}{224}$

In the next example, one of the fractions has a variable in its numerator. Notice that we do the same steps as when both numerators are numbers.

Example 1.7.19

Add: $\frac{3}{5}+\frac{x}{8}$

Solution

The fractions have different denominators.

	$\frac{3}{5}+\frac{x}{8}$

Find the LCD. $\begin{array}{l} 5= \quad 5 \\ 8=2\cdot 2\cdot 2 \\ \text{LCD}=2\cdot 2\cdot 2\cdot 5 \\ \text{LCD}=40 \end{array}$	
Rewrite as equivalent fractions with the LCD.	$\frac{3\cdot 8}{5\cdot 8}+\frac{x\cdot 5}{8\cdot 5}$
Simplify.	$\frac{24}{40}+\frac{5x}{40}$
Add.	$\frac{24+5x}{40}$

Remember, we can only add like terms: 24 and $5x$ are not like terms.

Try It 1.7.20

Add: $\frac{y}{6}+\frac{7}{9}$

Answer

$\frac{3y+14}{18}$

Try It 1.7.21

Add: $\frac{x}{6}+\frac{7}{15}$

Answer

$\frac{15x+42}{153}$

We now have all four operations for fractions. Table $1.7.1$ summarizes **fraction operations**.

Table 1.7.1

Fraction Multiplication	**Fraction Division**
$\frac{a}{b}\cdot\frac{c}{d}=\frac{ac}{bd}$ Multiply the numerators and multiply the denominators	$\frac{a}{b}\div\frac{c}{d}=\frac{a}{b}\cdot\frac{d}{c}$ Multiply the first fraction by the reciprocal of the second.
Fraction Addition	**Fraction Subtraction**
$\frac{a}{c}+\frac{b}{c}=\frac{a+b}{c}$ Add the numerators and place the sum over the common denominator.	$\frac{a}{c}-\frac{b}{c}=\frac{a-b}{c}$ Subtract the numerators and place the difference over the common denominator.
To multiply or divide fractions, and LCD is NOT needed. To add or subtract fractions, an LCD is needed.	

Example 1.7.22

Simplify:

1. $\frac{5x}{6}-\frac{3}{10}$
2. $\frac{5x}{6}\cdot\frac{3}{10}$.

Solution

First ask, “What is the operation?” Once we identify the operation that will determine whether we need a common denominator. Remember, we need a common denominator to add or subtract, but not to multiply or divide.

1. What is the operation? The operation is subtraction.

$$
\begin{array}{ll}
\text{Do the fractions have a common denominator? No.} & \frac{5x}{6}-\frac{3}{10} \\
\text{Rewrite each fractions as an equivalent fraction with the LCD.} & \frac{5x\cdot 5}{6\cdot 5}-\frac{3\cdot 3}{10\cdot 3} \\
 & \frac{25x}{30}-\frac{9}{30} \\
\text{Subtract the numerators and place the difference over the common denominators.} & \frac{25x-9}{30} \\
\text{Simplify, if possible. There are no common factors.} & \\
\text{The fraction is simplified.} &
\end{array}
\tag{1.7.6}
$$

2. What is the operation? Multiplication.

$$
\begin{array}{ll}
 & \frac{5x}{6}\cdot\frac{3}{10} \\
\text{To multiply fractions, multiply the numerators and multiply the denominators} & \frac{5x\cdot 3}{6\cdot 10} \\
\text{Rewrite, showing common factors.} & \frac{5x\cdot 3}{2\cdot 3\cdot 2\cdot 5} \\
\text{common denominators.} & \\
\text{Simplify.} & \frac{x}{4}
\end{array}
\tag{1.7.7}
$$

? Try It 1.7.23

Simplify:

1. $\frac{3a}{4}-\frac{8}{9}$
2. $\frac{3a}{4}\cdot\frac{8}{9}$

Answer

1. $\frac{27a-32}{36}$
2. $\frac{2a}{3}$

? Try It 1.7.24

Simplify:

1. $\frac{4k}{5}-\frac{1}{6}$
2. $\frac{4k}{5}\cdot\frac{1}{6}$

Answer

1. $\frac{24k-5}{30}$
2. $\frac{2k}{15}$

Use the Order of Operations to Simplify Complex Fractions

We have seen that a complex fraction is a fraction in which the numerat or or denominator contains a fraction. The fraction bar indicates **division**. We simplified the complex fraction $\frac{\frac{3}{4}}{\frac{5}{8}}$ by dividing $\frac{3}{4}$ by $\frac{5}{8}$.

Now we'll look at complex fractions where the numerator or denominator contains an expression that can be simplified. So we first must completely simplify the numerator and denominator separately using the order of operations. Then we divide the numerator by the denominator.

Example 1.7.25: How to simplify complex fractions

Simplify: $\frac{(\frac{1}{2})^2}{4+3^2}$

Solution

Step 1. Simplify the numerator. * Remember, $\left(\frac{1}{2}\right)^2$ means $\frac{1}{2}\cdot\frac{1}{2}$.	$\frac{\left(\frac{1}{2}\right)^2}{4+3^2}$ $\frac{\frac{1}{4}}{4+3^2}$
Step 2. Simplify the denominator.	$\frac{\frac{1}{4}}{4+9}$ $\frac{\frac{1}{4}}{13}$
Step 3. Divide the numerator by the denominator. Simplify if possible. * Remember, $13=\frac{13}{1}$	$\frac{1}{4}\div 13$ $\frac{1}{4}\cdot\frac{1}{13}$ $\frac{1}{52}$

Try It 1.7.26

Simplify: $\frac{(\frac{1}{3})^2}{2^3+2}$

Answer

$\frac{1}{90}$

Try It 1.7.27

Simplify: $\frac{1+4^2}{(\frac{1}{4})^2}$

Answer

272

SIMPLIFY COMPLEX FRACTIONS.

1. Simplify the numerator.
2. Simplify the denominator.
3. Divide the numerator by the denominator. Simplify if possible.

Example 1.7.28

Simplify: $\dfrac{\frac{1}{2}+\frac{2}{3}}{\frac{3}{4}-\frac{1}{6}}$

Solution

	$\dfrac{(\frac{1}{2}+\frac{2}{3})}{(\frac{3}{4}-\frac{1}{6})}$
Simplify the numerator (LCD $= 6$) and simplify the denominator (LCD $= 12$).	$\dfrac{(\frac{3}{6}+\frac{4}{6})}{(\frac{9}{12}-\frac{2}{12})}$
Simplify.	$\dfrac{(\frac{7}{6})}{(\frac{7}{12})}$ (1.7.8)
Divide the numerator by the denominator.	$\frac{7}{6} \div \frac{7}{12}$
Simplify.	$\frac{7}{6} \cdot \frac{12}{7}$
Divide out common factors.	$\frac{7 \cdot 6 \cdot 2}{6 \cdot 7}$
Simplify.	2

Try It 1.7.29

Simplify: $\dfrac{\frac{1}{3}+\frac{1}{2}}{\frac{3}{4}-\frac{1}{3}}$

Answer

2

Try It 1.7.30

Simplify: $\dfrac{\frac{2}{3}-\frac{1}{2}}{\frac{1}{4}+\frac{1}{3}}$

Answer

$\frac{2}{7}$

Evaluate Variable Expressions with Fractions

We have evaluated expressions before, but now we can evaluate expressions with fractions. Remember, to evaluate an expression, we substitute the value of the variable into the expression and then simplify.

Example 1.7.31

Evaluate $x+\dfrac{1}{3}$ when

1. $x=-\dfrac{1}{3}$

2. $x = -\frac{3}{4}$

Solution

1. To evaluate $x + \frac{1}{3}$ when $x = -\frac{1}{3}$, substitute $-\frac{1}{3}$ for x in the expression.

	$x+\frac{1}{3}$
Substitute $-\frac{1}{3}$ for x.	$-\frac{1}{3}+\frac{1}{3}$
Simplify.	0

2. To evaluate $x + \frac{1}{3}$ when $x = -\frac{3}{4}$, substitute $-\frac{3}{4}$ for x in the expression.

	$x+\frac{1}{3}$
Substitute $-\frac{3}{4}$ for x.	$-\frac{3}{4}+\frac{1}{3}$
Rewrite as equivalent fractions with the LCD, 12.	$-\frac{3\cdot3}{4\cdot3}+\frac{1\cdot4}{3\cdot4}$
Simplify.	$-\frac{9}{12}+\frac{4}{12}$
Add.	$-\frac{5}{12}$

? Try It 1.7.32

Evaluate $x + \frac{3}{4}$ when

1. $x = -\frac{7}{4}$
2. $x = -\frac{5}{4}$

Answer

1. -1
2. $-\frac{1}{2}$

? Try It 1.7.33

Evaluate $y + \frac{1}{2}$ when

1. $y = \frac{2}{3}$
2. $y = -\frac{3}{4}$

Answer

1. $\frac{7}{6}$
2. $-\frac{1}{12}$

Example 1.7.34

Evaluate $-\frac{5}{6}-y$ when $y=-\frac{2}{3}$

Solution

	$-\frac{5}{6}-y$
Substitute $-\frac{2}{3}$ for y.	$-\frac{5}{6}-\left(-\frac{2}{3}\right)$
Rewrite as equivalent fractions with the LCD, 6.	$-\frac{5}{6}-\left(-\frac{4}{6}\right)$
Subtract.	$\frac{-5-(-4)}{6}$
Simplify.	$-\frac{1}{6}$

Try It 1.7.35

Evaluate $y+\frac{1}{2}$ when $y=\frac{2}{3}$

Answer

$-\frac{1}{4}$

Try It 1.7.36

Evaluate $y+\frac{1}{2}$ when $y=\frac{2}{3}$

Answer

$-\frac{17}{8}$

Example 1.7.37

Evaluate $2x^2y$ when $x=\frac{1}{4}$ and $y=-\frac{2}{3}$.

Solution

Substitute the values into the expression.

	$2x^2y$
Substitute $\frac{1}{4}$ for x and $-\frac{2}{3}$ for y.	$2\left(\frac{1}{4}\right)^2\left(-\frac{2}{3}\right)$
Simplify exponents first.	$2\left(\frac{1}{16}\right)\left(-\frac{2}{3}\right)$
Multiply. Divide out the common factors. Notice we write 16 as $2\cdot 2\cdot 4$ to make it easy to remove	$-\frac{\cancel{2}\cdot 1\cdot\cancel{2}}{\cancel{2}\cdot\cancel{2}\cdot 4\cdot 3}$
Simplify.	$-\frac{1}{12}$

Try It 1.7.38

Evaluate $3ab^2$ when $a=-\frac{2}{3}$ and $b=-\frac{1}{2}$.

Answer

$-\frac{1}{2}$

Try It 1.7.39

Evaluate $4c^3d$ when $c=-\frac{1}{2}$ and $d=-\frac{4}{3}$.

Answer

$\frac{2}{3}$

The next example will have only variables, no constants.

Example 1.7.40

Evaluate $\frac{p+q}{r}$ when $p=-4, q=-2$, and $r=8$.

Solution

To evaluate $\frac{p+q}{r}$ when $p=-4, q=-2$, and $r=8$, we substitute the values into the expression.

	$\frac{p+q}{r}$
Substitute –4 for p, –2 for q and 8 for r.	$\frac{-4+(-2)}{8}$
Add in the numerator first.	$\frac{-6}{8}$
Simplify.	$-\frac{3}{4}$

Try It 1.7.41

Evaluate $\frac{a+b}{c}$ when $a=-8, b=-7$, and $c=6$.

Answer

$-\frac{5}{2}$

Try It 1.7.42

Evaluate $\frac{x+y}{z}$ when $x=9, y=-18$, and $z=-6$.

Answer

$\frac{3}{2}$

Key Concepts

- **Fraction Addition and Subtraction:** If a, b, and c are numbers where $c \neq 0$, then
  $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$ and $\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$
  To add or subtract fractions, add or subtract the numerators and place the result over the common denominator.
- **Strategy for Adding or Subtracting Fractions**
  1. Do they have a common denominator?
     Yes—go to step 2.
     No—Rewrite each fraction with the LCD (Least Common Denominator). Find the LCD. Change each fraction into an equivalent fraction with the LCD as its denominator.
  2. Add or subtract the fractions.
  3. Simplify, if possible. To multiply or divide fractions, an LCD IS NOT needed. To add or subtract fractions, an LCD IS needed.
- **Simplify Complex Fractions**
  1. Simplify the numerator.
  2. Simplify the denominator.
  3. Divide the numerator by the denominator. Simplify if possible.

This page titled 1.7: Add and Subtract Fractions is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.7: Add and Subtract Fractions** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.7E: Exercises

Practice Makes Perfect

Add and Subtract Fractions with a Common Denominator

In the following exercises, add.

Exercise 1.7E.1

$\frac{6}{13}+\frac{5}{13}$

Answer

$\frac{11}{13}$

Exercise 1.7E.2

$\frac{4}{15}+\frac{7}{15}$

Exercise 1.7E.3

$\frac{x}{4}+\frac{3}{4}$

Answer

$\frac{x+3}{4}$

Exercise 1.7E.4

$\frac{8}{q}+\frac{6}{q}$

Exercise 1.7E.5

$-\frac{3}{16}+\left(-\frac{7}{16}\right)$

Answer

$-\frac{5}{8}$

Exercise 1.7E.6

$-\frac{5}{16}+\left(-\frac{9}{16}\right)$

Exercise 1.7E.7

$-\frac{8}{17}+\frac{15}{17}$

Answer

$\frac{7}{17}$

Exercise 1.7E.8

$$-\frac{9}{19}+\frac{17}{19}$$

Exercise 1.7E.9

$$\frac{6}{13}+\left(-\frac{10}{13}\right)+\left(-\frac{12}{13}\right)$$

Answer

$-\frac{16}{13}$

Exercise 1.7E.10

$$\frac{5}{12}+\left(-\frac{7}{12}\right)+\left(-\frac{11}{12}\right)$$

In the following exercises, subtract.

Exercise 1.7E.11

$$\frac{11}{15}-\frac{7}{15}$$

Answer

$\frac{4}{15}$

Exercise 1.7E.12

$$\frac{9}{13}-\frac{4}{13}$$

Exercise 1.7E.13

$$\frac{11}{12}-\frac{5}{12}$$

Answer

$\frac{1}{2}$

Exercise 1.7E.14

$$\frac{7}{12}-\frac{5}{12}$$

Exercise 1.7E.15

$$\frac{19}{21}-\frac{4}{21}$$

Answer

$\frac{5}{7}$

Exercise 1.7E.16

$$\frac{17}{21} - \frac{8}{21}$$

Exercise 1.7E.17

$$\frac{5y}{8} - \frac{7}{8}$$

Answer

$\frac{5y-7}{8}$

Exercise 1.7E.18

$$\frac{11z}{13} - \frac{8}{13}$$

Exercise 1.7E.19

$$-\frac{23}{u} - \frac{15}{u}$$

Answer

$-\frac{38}{u}$

Exercise 1.7E.20

$$-\frac{29}{v} - \frac{26}{v}$$

Exercise 1.7E.21

$$-\frac{3}{5} - \left(-\frac{4}{5}\right)$$

Answer

$\frac{1}{5}$

Exercise 1.7E.22

$$-\frac{3}{7} - \left(-\frac{5}{7}\right)$$

Exercise 1.7E.23

$$-\frac{7}{9} - \left(-\frac{5}{9}\right)$$

Answer

$-\frac{2}{9}$

Exercise 1.7E.24

$$-\frac{8}{11}-\left(-\frac{5}{11}\right)$$

Mixed Practice

In the following exercises, simplify.

Exercise 1.7E.25

$$-\frac{5}{18}\cdot\frac{9}{10}$$

Answer

$-\frac{1}{4}$

Exercise 1.7E.26

$$-\frac{3}{14}\cdot\frac{7}{12}$$

Exercise 1.7E.27

$$\frac{n}{5}-\frac{4}{5}$$

Answer

$\frac{n-4}{5}$

Exercise 1.7E.28

$$\frac{6}{11}-\frac{s}{11}$$

Exercise 1.7E.29

$$-\frac{7}{24}+\frac{2}{24}$$

Answer

$-frac524$

Exercise 1.7E.30

$$-\frac{5}{18}+\frac{1}{18}$$

Exercise 1.7E.31

$$\frac{8}{15}\div\frac{12}{5}$$

Answer

$\frac{2}{9}$

Exercise 1.7E.32

$\frac{7}{12} \div \frac{9}{28}$

Add or Subtract Fractions with Different Denominators

In the following exercises, add or subtract.

Exercise 1.7E.33

$\frac{1}{2} + \frac{1}{7}$

Answer

$\frac{9}{14}$

Exercise 1.7E.34

$\frac{1}{3} + \frac{1}{8}$

Exercise 1.7E.35

$\frac{1}{3} - \left(-\frac{1}{9}\right)$

Answer

$\frac{4}{9}$

Exercise 1.7E.36

$\frac{1}{4} - \left(-\frac{1}{8}\right)$

Exercise 1.7E.37

$\frac{7}{12} + \frac{5}{12}$

Answer

$\frac{29}{24}$

Exercise 1.7E.38

$\frac{5}{12} + \frac{3}{8}$

Exercise 1.7E.39

$\frac{7}{12} - \frac{9}{16}$

Answer

$\frac{1}{48}$

Exercise 1.7E.40

$\frac{7}{16} - \frac{5}{12}$

Exercise 1.7E.41

$\frac{2}{3} - \frac{3}{8}$

Answer

$\frac{7}{24}$

Exercise 1.7E.42

$\frac{5}{6} - \frac{3}{4}$

Exercise 1.7E.43

$-\frac{11}{30} + \frac{27}{40}$

Answer

$\frac{37}{120}$

Exercise 1.7E.44

$-\frac{9}{20} + \frac{17}{30}$

Exercise 1.7E.45

$-\frac{13}{30} + \frac{25}{42}$

Answer

$\frac{17}{105}$

Exercise 1.7E.46

$-\frac{23}{30} + \frac{5}{48}$

Exercise 1.7E.47

$-\frac{39}{56} - \frac{22}{35}$

Answer

$-\frac{53}{40}$

Exercise 1.7E.48

$-\frac{33}{49} - \frac{18}{35}$

Exercise 1.7E.49

$-\frac{2}{3}-\left(-\frac{3}{4}\right)$

Answer

$\frac{1}{12}$

Exercise 1.7E.50

$-\frac{3}{4}-\left(-\frac{4}{5}\right)$

Exercise 1.7E.51

$1+\frac{7}{8}$

Answer

$\frac{15}{8}$

Exercise 1.7E.52

$1-\frac{3}{10}$

Exercise 1.7E.53

$\frac{x}{3}+\frac{1}{4}$

Answer

$\frac{4x+3}{12}$

Exercise 1.7E.54

$\frac{y}{2}+\frac{2}{3}$

Exercise 1.7E.55

$\frac{y}{4}-\frac{3}{5}$

Answer

$\frac{5y-12}{20}$

Exercise 1.7E.56

$\frac{x}{5}-\frac{1}{4}$

Mixed Practice

In the following exercises, simplify.

Exercise 1.7E.57

1. $\frac{2}{3}+\frac{1}{6}$
2. $\frac{2}{3} \div \frac{1}{6}$

Answer

1. $\frac{5}{6}$
2. 4

Exercise 1.7E.58

1. $-\frac{2}{5}-\frac{1}{8}$
2. $-\frac{2}{5} \cdot \frac{1}{8}$

Exercise 1.7E.59

1. $\frac{5n}{6} \div \frac{8}{15}$
2. $\frac{5n}{6}-\frac{8}{15}$

Answer

1. $\frac{25n}{16}$
2. $\frac{25n-16}{30}$

Exercise 1.7E.60

1. $\frac{3a}{8} \div \frac{7}{12}$
2. $\frac{3a}{8}-\frac{7}{12}$

Exercise 1.7E.61

$-\frac{3}{8} \div \left(-\frac{3}{10}\right)$

Answer

$\frac{5}{4}$

Exercise 1.7E.62

$-\frac{5}{12} \div \left(-\frac{5}{9}\right)$

Exercise 1.7E.63

$-\frac{3}{8}+\frac{5}{12}$

Answer

$\frac{1}{24}$

Exercise 1.7E.64

$-\frac{1}{8}+\frac{7}{12}$

Exercise 1.7E.65

$\frac{5}{6} - \frac{1}{9}$

Answer

$\frac{13}{18}$

Exercise 1.7E.66

$\frac{5}{9} - \frac{1}{6}$

Exercise 1.7E.67

$-\frac{7}{15} - \frac{y}{4}$

Answer

$\frac{-28-15y}{60}$

Exercise 1.7E.68

$-\frac{3}{8} - \frac{x}{11}$

Exercise 1.7E.69

$\frac{11}{12a} \cdot \frac{9a}{16}$

Answer

$\frac{33}{64}$

Exercise 1.7E.70

$\frac{10y}{13} \cdot \frac{8}{15y}$

Use the Order of Operations to Simplify Complex Fractions

In the following exercises, simplify.

Exercise 1.7E.71

$\frac{2^3+4^2}{\left(\frac{2}{3}\right)^2}$

Answer

54

Exercise 1.7E.72

$\frac{3^3-3^2}{\left(\frac{3}{4}\right)^2}$

Exercise $1.7E.73$

$$\frac{\left(\frac{3}{5}\right)^2}{\left(\frac{3}{7}\right)^2}$$

Answer

$\frac{49}{25}$

Exercise $1.7E.74$

$$\frac{\left(\frac{3}{4}\right)^2}{\left(\frac{5}{8}\right)^2}$$

Exercise $1.7E.75$

$$\frac{2}{\frac{1}{3}+\frac{1}{5}}$$

Answer

$\frac{15}{4}$

Exercise $1.7E.76$

$$\frac{5}{\frac{1}{4}+\frac{1}{3}}$$

Exercise $1.7E.77$

$$\frac{\frac{7}{8}-\frac{2}{3}}{\frac{1}{2}+\frac{3}{8}}$$

Answer

$\frac{5}{21}$

Exercise $1.7E.78$

$$\frac{\frac{3}{4}-\frac{3}{5}}{\frac{1}{4}+\frac{2}{5}}$$

Exercise $1.7E.79$

$\frac{1}{2}+\frac{2}{3}\cdot\frac{5}{12}$

Answer

$\frac{7}{9}$

Exercise $1.7E.80$

$\frac{1}{3}+\frac{2}{5}\cdot\frac{3}{4}$

Exercise 1.7E.81

$1-\frac{3}{5} \div \frac{1}{10}$

Answer

-5

Exercise 1.7E.82

$1-\frac{5}{6} \div \frac{1}{12}$

Exercise 1.7E.83

$\frac{2}{3}+\frac{1}{6}+\frac{3}{4}$

Answer

$\frac{19}{12}$

Exercise 1.7E.84

$\frac{2}{3}+\frac{1}{4}+\frac{3}{5}$

Exercise 1.7E.85

$\frac{3}{8}-\frac{1}{6}+\frac{3}{4}$

Answer

$\frac{23}{24}$

Exercise 1.7E.86

$\frac{2}{5}+\frac{5}{8}-\frac{3}{4}$

Exercise 1.7E.87

$12\left(\frac{9}{20}-\frac{4}{15}\right)$

Answer

$\frac{11}{5}$

Exercise 1.7E.88

$8\left(\frac{15}{16}-\frac{5}{6}\right)$

Exercise 1.7E.89

$\frac{\frac{5}{8}+\frac{1}{6}}{\frac{19}{24}}$

Answer

1

Exercise 1.7E.90

$\frac{\frac{1}{6}+\frac{3}{10}}{\frac{14}{30}}$

Exercise 1.7E.91

$\left(\frac{5}{9}+\frac{1}{6}\right) \div \left(\frac{2}{3}-\frac{1}{2}\right)$

Answer

$\frac{13}{3}$

Exercise 1.7E.92

$\left(\frac{3}{4}+\frac{1}{6}\right) \div \left(\frac{5}{8}-\frac{1}{3}\right)$

Evaluate Variable Expressions with Fractions

In the following exercises, evaluate.

Exercise 1.7E.93

$x+\left(-\frac{5}{6}\right)$ when

1. $x=\frac{1}{3}$
2. $x=-\frac{1}{6}$

Answer

1. $-\frac{1}{2}$
2. -1

Exercise 1.7E.94

$x+\left(-\frac{11}{12}\right)$ when

1. $x=\frac{11}{12}$
2. $x=-\frac{3}{4}$

Exercise 1.7E.95

$x-\frac{2}{5}$ when

1. $x=\frac{3}{5}$
2. $x=-\frac{3}{5}$

Answer

1. $\frac{1}{5}$
2. -1

Exercise 1.7E.96

$x - \frac{1}{3}$ when

1. $x = \frac{2}{3}$
2. $x = -\frac{2}{3}$

Exercise 1.7E.97

$\frac{7}{10} - w$ when

1. $w = \frac{1}{2}$
2. $w = -\frac{1}{2}$

Answer

1. $\frac{1}{5}$
2. $\frac{6}{5}$

Exercise 1.7E.98

$\frac{5}{12} - w$ when

1. $w = \frac{1}{4}$
2. $w = -\frac{1}{4}$

Exercise 1.7E.99

$2x^2y^3$ when $x = -\frac{2}{3}$ and $y = -\frac{1}{2}$

Answer

$-\frac{1}{9}$

Exercise 1.7E.100

$8u^2v^3$ when $u = -\frac{3}{4}$ and $v = -\frac{1}{2}$

Exercise 1.7E.101

$\frac{a+b}{a-b}$ when $a = -3, b = 8$

Answer

$-\frac{5}{11}$

Exercise 1.7E.102

$\frac{r-s}{r+s}$ when $r = 10, s = -5$

Everyday Math

Exercise 1.7E.103

Decorating Laronda is making covers for the throw pillows on her sofa. For each pillow cover, she needs $\frac{1}{2}$ yard of print fabric and \frac{3}{8}\) yard of solid fabric. What is the total amount of fabric Laronda needs for each pillow cover?

Answer

$\frac{7}{8}$ yard

Exercise 1.7E.104

Baking Vanessa is baking chocolate chip cookies and oatmeal cookies. She needs $\frac{1}{2}$ cup of sugar for the chocolate chip cookies and $\frac{1}{4}$ of sugar for the oatmeal cookies. How much sugar does she need altogether?

Writing Exercises

Exercise 1.7E.105

Why do you need a common denominator to add or subtract fractions? Explain.

Answer

Answers may vary

Exercise 1.7E.106

How do you find the LCD of 2 fractions?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
add and subtract fractions with different denominators.			
identify and use fraction operations.			
use the order of operations to simplify complex fractions.			
evaluate variable expressions with fractions.			

ⓑ After looking at the checklist, do you think you are well-prepared for the next chapter? Why or why not?

1.7E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **1.7E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.8: Decimals

Learning Objectives

By the end of this section, you will be able to:

- Name and write decimals
- Round decimals
- Add and subtract decimals
- Multiply and divide decimals
- Convert decimals, fractions, and percents

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra*chapter, **Decimals**.

Name and Write Decimals

Decimals are another way of writing **fraction**s whose denominators are powers of 10.

$$\begin{aligned} 0.1 &= \tfrac{1}{10} && 0.1 \text{ is "one tenth"} \\ 0.01 &= \tfrac{1}{100} && 0.01 \text{ is "one hundredth} \\ 0.001 &= \tfrac{1}{1{,}000} && 0.001 \text{ is "one thousandth} \\ 0.0001 &= \tfrac{1}{10{,}000} && 0.0001 \text{ is "one ten-thousandth"} \end{aligned} \tag{1.8.1}$$

Notice that "ten thousand" is a number larger than one, but "one ten-thousand**th**" is a number smaller than one. The "th" at the end of the name tells you that the number is smaller than one.

When we name a whole number, the name corresponds to the place value based on the powers of ten. We read 10,000 as "ten thousand" and 10,000,000 as "ten million." Likewise, the names of the **decimal** places correspond to their fraction values. Figure 1.8.1 shows the names of the **place values** to the left and right of the decimal point.




Figure 1.8.1: Place value of decimal numbers are shown to the left and right of the decimal point.

Exercise 1.8.1

Name the decimal 4.3.

Answer

Step 1. Name the number to the left of the decimal point.	4 is to the left of the decimal point.	four	4.3	**Step 2.** Write 'and' for the decimal point.	four and
Step 3. Name the 'number' part to the right of the decimal point as if it were a whole number.	3 is to the right of the decimal point.	four and three		**Step 4.** Name the decimal place.	four and three tenths

Exercise 1.8.2

Name the decimal 6.7.

Answer

six and seven tenths

Exercise 1.8.3

Name the decimal 5.8.

Answer

five and eight tenths

We summarize the steps needed to name a **decimal** below.

NAME A DECIMAL.

1. Name the number to the left of the decimal point.
2. Write "and" for the decimal point.
3. Name the "number" part to the right of the decimal point as if it were a whole number.
4. Name the decimal place of the last digit.

Exercise 1.8.4

Name the decimal: -15.571.

Answer

	-15.571
Name the number to the left of the decimal point.	negative fifteen _____
Write “and” for the decimal point.	negative fifteen and _____
Name the number to the right of the decimal point.	negative fifteen and five hundred seventy-one _____
The 1 is in the thousandths place.	negative fifteen and five hundred seventy-one thousandths

Exercise 1.8.5

Name the decimal: -13.461.

Answer

negative thirteen and four hundred sixty-one thousandths

Exercise 1.8.6

Name the decimal: -2.053.

Answer

negative two and fifty-three thousandths

When we write a check we write both the numerals and the name of the number. Let’s see how to write the decimal from the name.

Exercise 1.8.7: How to Write decimals

Write “fourteen and twenty-four thousandths” as a decimal.

Answer

Step 1. Look for the word ‘and’; it locates the decimal point. Place a decimal point under the word ‘and’. Translate the words before ‘and’ into the whole number and place to the left of the decimal point.		fourteen and twenty-four thousandths fourteen and twenty-four thousandths ____ . _____ 14. _____
Step 2. Mark the number of decimal places needed to the right of the decimal point by noting the place value indicated by the last word.	The last word is ‘thousandths’.	14. ____ ____ ____ tenths hundredths thousandths
Step 3. Translate the words after ‘and’ into the number to the right of the decimal point. Write the number in the spaces – putting the final digit in the last place.		14. ____ 2 4
Step 4. Fill in zeros for empty place holders as needed.	Zeros are needed in the tenths place.	14. 0 2 4 Fourteen and twenty-four thousandths is written 14.024.

Exercise 1.8.8

Write as a decimal: thirteen and sixty-eight thousandths.

Answer

13.068

Exercise 1.8.9

Write as a decimal: five and ninety-four thousandths.

Answer

5.094

We summarize the steps to writing a **decimal**.

WRITE A DECIMAL.

1. Look for the word “and”—it locates the decimal point.
   - Place a decimal point under the word “and.” Translate the words before “and” into the whole number and place it to the left of the decimal point.
   - If there is no “and,” write a “0” with a decimal point to its right.
2. Mark the number of decimal places needed to the right of the decimal point by noting the place value indicated by the last word.
3. Translate the words after “and” into the number to the right of the decimal point. Write the number in the spaces—putting the final digit in the last place.
4. Fill in zeros for place holders as needed.

Round Decimals

Rounding decimals is very much like rounding whole numbers. We will round decimals with a method based on the one we used to round whole numbers.

Exercise 1.8.10

Round 18.379 to the nearest hundredth.

Answer

Step 1. Locate the given place value and mark it with an arrow.		hundredths place ↓ 18.379	**Step 2.** Underline the digit to the right of the given place value.		hundredths place ↓ 18.379
Step 3. Is this digit greater than or equal to 5? **Yes:** Add 1 to the digit in the given place value. **No:** Do not change the digit in the given place value.	Because 9 is greater than or equal to 5, add 1 to the 7.	18.37 9 add 1 delete	**Step 4.** Rewrite the number, removing all digits to the right of the rounding digit.		18.38 18.38 is 18.379 rounded to the nearest hundredth.

Exercise 1.8.11

Round to the nearest hundredth: 1.047.

Answer

1.05

Exercise 1.8.12

Round to the nearest hundredth: 9.173.

Answer

9.17

We summarize the steps for rounding a **decimal** here.

ROUND DECIMALS.

1. Locate the given place value and mark it with an arrow.
2. Underline the digit to the right of the place value.
3. Is this digit greater than or equal to 5?
   - Yes—add 1 to the digit in the given place value.
   - No—do not change the digit in the given place value.
4. Rewrite the number, deleting all digits to the right of the rounding digit.

Exercise 1.8.13

Round 18.379 to the nearest

1. tenth
2. whole number.

Answer

Round 18.379

1. to the nearest tenth

Locate the tenths place with an arrow.	tenths place ↓ 18.379
Underline the digit to the right of the given place value.	tenths place ↓ 18.379
Because 7 is greater than or equal to 5, add 1 to the 3.	18.379 add 1 delete
Rewrite the number, deleting all digits to the right of the rounding digit.	18.4
Notice that the deleted digits were NOT replaced with zeros.	So, 18.379 rounded to the nearest tenth is 18.4.

2. to the nearest whole number

Locate the ones place with an arrow.	ones place ↓ 18.379
Underline the digit to the right of the given place value.	ones place ↓ 18.379

Since 3 is not greater than or equal to 5, do not add 1 to the 8.	18.379 do not add 1 delete
Rewrite the number, deleting all digits to the right of the rounding digit.	18
	So, 18.379 rounded to the nearest whole number is 18.

? Exercise 1.8.14

Round 6.582 to the nearest

1. hundredth
2. tenth
3. whole number.

Answer

1. 6.58
2. 6.6
3. 7

? Exercise 1.8.15

Round 15.2175 to the nearest

1. thousandth
2. hundredth
3. tenth.

Answer

1. 15.218
2. 15.22
3. 15.2

Add and Subtract Decimals

To add or subtract decimals, we line up the decimal points. By lining up the decimal points this way, we can add or subtract the corresponding **place values**. We then add or subtract the numbers as if they were whole numbers and then place the decimal point in the sum.

ADD OR SUBTRACT DECIMALS.

1. Write the numbers so the decimal points line up vertically.
2. Use zeros as place holders, as needed.
3. Add or subtract the numbers as if they were whole numbers. Then place the decimal point in the answer under the decimal points in the given numbers.

? Exercise 1.8.16

Add: $23.5+41.38$.

Answer

Write the numbers so that the decimal points line up vertically.

$$\begin{array}{r} 23.50 \\ +41.38 \\ \hline \end{array} \tag{1.8.2}$$

Put 0 as a placeholder after the 5 in 23.5. Remember, $\frac{5}{10}=\frac{50}{100}$, so $0.5=0.50$

$$\begin{array}{r} 23.50 \\ +41.38 \\ \hline \end{array} \tag{1.8.3}$$

Add the numbers as if they were whole numbers . Then place the decimal point in the sum.

$$\begin{array}{r} 23.50 \\ +41.38 \\ \hline 64.88 \end{array} \tag{1.8.4}$$

? Exercise 1.8.17

Add: $4.8+11.69$.

Answer

16.49

? Exercise 1.8.18

Add: $5.123+18.47$.

Answer

23.593

Exercise 1.8.19

Subtract: $20 - 14.65$.

Answer

Write the numbers so that the decimal points line up vertically.

$$\begin{array}{r} 20 - 14.65 \\ 20. \\ -14.65 \\ \hline \end{array}$$

Remember, 20 is a whole number, so place the decimal point after the 0.

Put zeros to the right as placeholders.

$$\begin{array}{rr} 20.00 & (1.8.8) \\ -14.65 & (1.8.9) \\ \hline \end{array}$$

Write the numbers so that the decimal points line up vertically.

$$\begin{array}{r} {}^{9}\quad {}^{9} \\ 1\,\not{1}\not{0}\,\not{1}\not{0}\,\not{1}\not{0} \\ \not{2}\,\not{0}.\,\not{0}\,\not{0} \\ -14.65 \\ \hline 5.35 \end{array}$$

Exercise 1.8.20

Subtract: $10 - 9.58$.

Answer

0.42

Exercise 1.8.21

Subtract: $50 - 37.42$.

Answer

12.58

Multiply and Divide Decimals

Multiplying decimals is very much like multiplying whole numbers—we just have to determine where to place the decimal point. The procedure for multiplying decimals will make sense if we first convert them to fractions and then multiply.

So let's see what we would get as the **product** of decimals by converting them to fractions first. We will do two examples side-by-side. Look for a pattern!

Table 1.8.1

	(0.3) (0.7)	(0.2) (0.46)
Convert to fractions.	$\frac{3}{10} \cdot \frac{7}{10}$	$\frac{2}{10} \cdot \frac{46}{100}$
Multiply.	$\frac{21}{100}$	$\frac{92}{1000}$
Convert to decimals.	0.21	0.092

Notice, in the first example, we multiplied two numbers that each had one digit after the decimal point and the product had two decimal places. In the second example, we multiplied a number with one decimal place by a number with two decimal places and the product had three decimal places.

We multiply the numbers just as we do whole numbers, temporarily ignoring the decimal point. We then count the number of decimal points in the factors and that sum tells us the number of decimal places in the product.

The rules for multiplying positive and negative numbers apply to decimals, too, of course!

When *multiplying* two numbers,

- if their signs are the *same* the product is *positive*.
- if their signs are *different* the product is *negative*.

When we multiply signed decimals, first we determine the sign of the product and then multiply as if the numbers were both positive. Finally, we write the product with the appropriate sign.

MULTIPLY DECIMALS.

1. Determine the sign of the product.
2. Write in vertical format, lining up the numbers on the right. Multiply the numbers as if they were whole numbers, temporarily ignoring the decimal points.
3. Place the decimal point. The number of decimal places in the product is the sum of the number of decimal places in the factors.
4. Write the product with the appropriate sign.

Exercise 1.8.22

Multiply: $(-3.9)(4.075)$.

Answer

	$(-3.9)(4.075)$
The signs are different. The product will be negative.	
Write in vertical format, lining up the numbers on the right.	4.075 × 3.9
Multiply.	4.075 × 3.9 36675 12225 158925
Add the number of decimal places in the factors $(1+3)$. (–3.9) (4.075) 1 place 3 places Place the decimal point 4 places from the right.	4.075 × 3.9 36675 12225 15.8925 4 places
The signs are different, so the product is negative.	$(-3.9)(4.075) = -15.8925$

Exercise 1.8.23

Multiply: $-4.5(6.107)$

Answer

-27.4815

Exercise 1.8.24

Multiply: $-10.79(8.12)$.

Answer

-87.6148

In many of your other classes, especially in the sciences, you will multiply decimals by powers of 10 (10, 100, 1000, etc.). If you multiply a few products on paper, you may notice a pattern relating the number of zeros in the power of 10 to number of decimal places we move the decimal point to the right to get the product.

MULTIPLY A DECIMAL BY A POWER OF TEN.

1. Move the decimal point to the right the same number of places as the number of zeros in the power of 10.
2. Add zeros at the end of the number as needed.

Exercise 1.8.25

Multiply 5.63

1. by 10
2. by 100
3. by 1,000.

Answer

By looking at the number of zeros in the multiple of ten, we see the number of places we need to move the decimal to the right.

ⓐ

	$5.63(10)$
There is 1 zero in 10, so move the decimal point 1 place to the right.	5.63 56.3

ⓑ

	$5.63(100)$
There are 2 zeros in 100, so move the decimal point 2 places to the right.	There are 2 zeros in 100, so move the decimal point 2 places to the right. 5.63 (100) 5,63 563

ⓒ

There are 3 zeros in 1,000, so move the decimal point 3 places to the right.	5.63
A zero must be added at the end.	5,630

Exercise 1.8.26

Multiply 2.58

1. by 10
2. by 100
3. by 1,000.

Answer

1. 25.8
2. 258
3. 2,580

Exercise 1.8.27

Multiply 14.2

1. by 10
2. by 100
3. by 1,000.

Answer

1. 142
2. 1,420
3. 14,200

Just as with multiplication, **division** of decimals is very much like dividing whole numbers. We just have to figure out where the decimal point must be placed.

To divide decimals, determine what power of 10 to multiply the denominator by to make it a whole number. Then multiply the numerator by that same power of 10. Because of the equivalent fractions property, we haven't changed the value of the fraction! The effect is to move the decimal points in the numerator and denominator the same number of places to the right. For example:

$$\begin{array}{c} \frac{0.8}{0.4} \\ \frac{0.8(10)}{0.4(10)} \\ \frac{8}{4} \end{array} \tag{1.8.16}$$

We use the rules for dividing positive and negative numbers with decimals, too. When dividing signed decimals, first determine the sign of the quotient and then divide as if the numbers were both positive. Finally, write the quotient with the appropriate sign.

We review the notation and vocabulary for division:

$$\underset{\text{dividend}}{a} \div \underset{\text{divisor}}{b} = \underset{\text{quotient}}{c} \qquad \begin{array}{r} \overset{\text{quotient}}{c} \\ \underset{\text{divisor}}{b}\,\overline{)\,\underset{\text{dividend}}{a}} \end{array} \tag{1.8.17}$$

We'll write the steps to take when dividing decimals, for easy reference.

DIVIDE DECIMALS.

1. Determine the sign of the quotient.
2. Make the divisor a whole number by "moving" the decimal point all the way to the right. "Move" the decimal point in the dividend the same number of places—adding zeros as needed.
3. Divide. Place the decimal point in the quotient above the decimal point in the dividend.
4. Write the quotient with the appropriate sign.

Exercise 1.8.28

Divide: $-25.65 \div (-0.06)$.

Answer

Remember, you can "move" the decimals in the divisor and dividend because of the Equivalent Fractions Property.

	$-25.65 \div (-0.06)$
The signs are the same.	The quotient is positive.
Make the divisor a whole number by "moving" the decimal point all the way to the right.	
"Move" the decimal point in the dividend the same number of places.	$0.06\overline{)25.65}$
Divide. Place the decimal point in the quotient above the decimal point in the dividend.	427.5 006.)2565.0 −24 16 −12 45 −42 30 30
Write the quotient with the appropriate sign.	$-25.65 \div (-0.06) = 427.5$

Exercise 1.8.29

Divide: $-23.492 \div (-0.04)$.

Answer

687.3

Exercise 1.8.30

Divide: $-4.11 \div (-0.12)$.

Answer

34.25

A common application of dividing whole numbers into decimals is when we want to find the price of one item that is sold as part of a multi-pack. For example, suppose a case of 24 water bottles costs $3.99. To find the price of one water bottle, we would divide $3.99 by 24. We show this division in Exercise 1.8.31. In calculations with money, we will round the answer to the nearest cent (hundredth).

Exercise 1.8.31

Divide: $3.99 \div 24$.

Answer

.99 divided by 24 is given. A long division problem is set up with 24 dividing 3.99. A table is given with directions on the left and the mathematical steps on the right. The first step reads “Place the decimal point in the quotient above the decimal point in the dividend. Divide as usual. When do we stop? Since this division involves money, we round it to the nearest cent (hundredth). To do this, we must carry the division to the thousandths place.” To the right of this, we have a long division problem set up with 24 dividing 3.990. The quotient is given as 0.166. To show the work, below 3.990 it reads 24, solid horizontal line, 159, 144, solid horizontal line, 150, 144, solid horizontal line, and finally 6. The fifth step reads “Round to the nearest cent.” To the right of this, we have $0.166 is approximately equal to $0.17 and hence >.99 divided by 24 is $0.17.">

	$3.99 \div 24$
Place the decimal point in the quotient above the decimal point in the dividend.	
Divide as usual. When do we stop? Since this division involves money, we round it to the nearest cent (hundredth.) To do this, we must carry the division to the thousandths place.	$$\begin{array}{r} 0.166 \\ 24\overline{)3.990} \\ \underline{24} \\ 159 \\ \underline{144} \\ 150 \\ \underline{144} \\ 6 \end{array}$$
Round to the nearest cent.	$0.166 \approx $0.17 $3.99 \div 2 \approx $0.17

Exercise 1.8.32

Divide: $6.99 \div 36$.

Answer

$0.19

Exercise 1.8.33

Divide: $4.99 \div 12$.

Answer

$0.42

Convert Decimals, Fractions, and Percents

We convert decimals into fractions by identifying the place value of the last (farthest right) digit. In the decimal 0.03 the 3 is in the hundredths place, so 100 is the denominator of the fraction equivalent to 0.03.

$$00.03 = \frac{3}{100} \tag{1.8.18}$$

Notice, when the number to the left of the decimal is zero, we get a fraction whose numerator is less than its denominator. Fractions like this are called **proper fractions**.

The steps to take to convert a decimal to a fraction are summarized in the procedure box.

CONVERT A DECIMAL TO A PROPER FRACTION.

1. Determine the place value of the final digit.
2. Write the fraction.
   - numerator—the “numbers” to the right of the decimal point
   - denominator—the place value corresponding to the final digit

Exercise 1.8.34

Write 0.374 as a fraction.

Answer

	0.374
Determine the place value of the final digit.	0.3 (tenths) 7 (hundredths) 4 (thousandths)
Write the fraction for 0.374: • The numerator is 374. • The denominator is 1,000.	$\frac{374}{1000}$
Simplify the fraction.	$\frac{2 \cdot 187}{2 \cdot 500}$
Divide out the common factors.	$\frac{187}{500}$ so, $0.374 = \frac{187}{500}$

Did you notice that the number of zeros in the denominator of $\frac{374}{1000}$ is the same as the number of decimal places in 0.374?

Exercise 1.8.35

Write 0.234 as a fraction.

Answer

$\frac{117}{500}$

Exercise 1.8.36

Write 0.024 as a fraction.

Answer

$\frac{3}{125}$

We've learned to convert decimals to fractions. Now we will do the reverse—convert fractions to decimals. Remember that the fraction bar means division. So $\frac{4}{5}$ can be written $4 \div 5$ or $5\overline{)4}$. This leads to the following method for converting a fraction to a decimal.

CONVERT A FRACTION TO A DECIMAL.

To convert a fraction to a decimal, divide the numerator of the fraction by the denominator of the fraction.

Exercise 1.8.37

Write $-\frac{5}{8}$ as a decimal.

Answer

Since a fraction bar means division, we begin by writing $\frac{5}{8}$ as $8\overline{)5}$. Now divide.

$$\begin{array}{r} 0.625 \\ 8\overline{)5.000} \\ \underline{48} \\ 20 \\ \underline{16} \\ 40 \\ \underline{40} \\ 0 \end{array}$$

so, $-\frac{5}{8} = -0.625$

Exercise 1.8.38

Write $-\frac{7}{8}$ as a decimal.

Answer

−0.875

Exercise 1.8.39

Write $-\frac{3}{8}$ as a decimal.

Answer

−0.375

When we divide, we will not always get a zero remainder. Sometimes the quotient ends up with a decimal that repeats. A **repeating decimal** is a decimal in which the last digit or group of digits repeats endlessly. A bar is placed over the repeating block of digits to indicate it repeats.

REPEATING DECIMAL


A **repeating decimal** is a decimal in which the last digit or group of digits repeats endlessly.

A bar is placed over the repeating block of digits to indicate it repeats.

Exercise 1.8.40

Write $\frac{43}{22}$ as a decimal.

Answer



Exercise 1.8.41

Write $\frac{27}{11}$ as a decimal.

Answer

$2.\overline{45}$

Exercise 1.8.42

Write $\frac{51}{22}$ as a decimal.

Answer

$2.3\overline{18}$

Sometimes we may have to simplify expressions with fractions and decimals together.

Exercise 1.8.43

Simplify: $\frac{7}{8}+6.4$.

Answer

First we must change one number so both numbers are in the same form. We can change the fraction to a decimal, or change the decimal to a fraction. Usually it is easier to change the fraction to a decimal.

		$\frac{7}{8}+6.4$
Change $\frac{7}{8}$ to a decimal.	$8\overline{)7.000}$ 64 60 56 40 40 0	
Add.		$0.875+6.4$
		7.275
		So, $\frac{7}{8}+6.4=7.275$

Exercise 1.8.44

Simplify: $\frac{3}{8}+4.9$.

Answer

5.275

Exercise 1.8.45

Simplify: $5.7+\frac{13}{20}$.

Answer

6.35

A **percent** is a ratio whose denominator is 100. Percent means per hundred. We use the percent symbol, %, to show percent.

PERCENT

A **percent** is a ratio whose denominator is 100.

Since a percent is a ratio, it can easily be expressed as a fraction. Percent means per 100, so the denominator of the fraction is 100. We then change the fraction to a decimal by dividing the numerator by the denominator.

	6%	78%	135%
Write as a ratio with denominator 100.	$\frac{6}{100}$	$\frac{78}{100}$	$\frac{135}{100}$
Change the fraction to a decimal by dividing the numerator by the denominator.	0.06	0.78	1.35

(1.8.19)

Do you see the pattern? *To convert a percent number to a decimal number, we move the decimal point two places to the left.*



Figure 1.8.2

Exercise 1.8.46

Convert each percent to a decimal:

1. 62%
2. 135%
3. 35.7%.

Answer

1.	
	62%
Move the decimal point two places to the left.	0.62
2.	
	135%
Move the decimal point two places to the left.	1.35
3.	
	5.7%
Move the decimal point two places to the left.	0.057

Exercise 1.8.47

Convert each percent to a decimal:

1. 9%
2. 87%
3. 3.9%.

Answer

1. 0.09
2. 0.87
3. 0.039

Exercise 1.8.48

Convert each percent to a decimal:

1. 3%
2. 91%
3. 8.3%.

Answer

1. 0.03
2. 0.91
3. 0.083

Converting a decimal to a percent makes sense if we remember the definition of percent and keep place value in mind.

To convert a decimal to a percent, remember that percent means per hundred. If we change the decimal to a fraction whose denominator is 100, it is easy to change that fraction to a percent.

	0.83	1.05	0.075
Write as a fraction	$\frac{83}{100}$	$1\frac{5}{100}$	$\frac{75}{1000}$
The denominator is 100.		$\frac{105}{100}$	$\frac{7.5}{100}$
Write the ratio as a percent.	83%	105%	7.5%

(1.8.20)

Recognize the pattern? *To convert a decimal to a percent, we move the decimal point two places to the right and then add the percent sign.*




Figure 1.8.3

https://math.libretexts.org/@go/page/30463

Exercise 1.8.49

Convert each decimal to a percent:

1. 0.51
2. 1.25
3. 0.093.

Answer

1.	
	0.51
Move the decimal point two places to the right.	51

2.	
	1.25
Move the decimal point two places to the right.	125

3.	
	0.093
Move the decimal point two places to the right.	9.3

Exercise 1.8.50

Convert each decimal to a percent:

1. 0.17
2. 1.75
3. 0.0825

Answer

1. 17%
2. 175%
3. 8.25%

Exercise 1.8.51

Convert each decimal to a percent:

1. 0.41
2. 2.25
3. 0.0925.

Answer

1. 41%
2. 225%
3. 9.25%

Key Concepts

- **Name a Decimal**
  1. Name the number to the left of the decimal point.
  2. Write ”and” for the decimal point.
  3. Name the “number” part to the right of the decimal point as if it were a whole number.
  4. Name the decimal place of the last digit.
- **Write a Decimal**
  1. Look for the word ‘and’—it locates the decimal point. Place a decimal point under the word ‘and.’ Translate the words before ‘and’ into the whole number and place it to the left of the decimal point. If there is no “and,” write a “0” with a decimal point to its right.
  2. Mark the number of decimal places needed to the right of the decimal point by noting the place value indicated by the last word.
  3. Translate the words after ‘and’ into the number to the right of the decimal point. Write the number in the spaces—putting the final digit in the last place.
  4. Fill in zeros for place holders as needed.
- **Round a Decimal**
  1. Locate the given place value and mark it with an arrow.
  2. Underline the digit to the right of the place value.
  3. Is this digit greater than or equal to 5? Yes—add 1 to the digit in the given place value. No—do not change the digit in the given place value.
  4. Rewrite the number, deleting all digits to the right of the rounding digit.
- **Add or Subtract Decimals**
  1. Write the numbers so the decimal points line up vertically.
  2. Use zeros as place holders, as needed.
  3. Add or subtract the numbers as if they were whole numbers. Then place the decimal in the answer under the decimal points in the given numbers.
- **Multiply Decimals**
  1. Determine the sign of the product.

2. Write in vertical format, lining up the numbers on the right. Multiply the numbers as if they were whole numbers, temporarily ignoring the decimal points.
3. Place the decimal point. The number of decimal places in the product is the sum of the decimal places in the factors.
4. Write the product with the appropriate sign.

- **Multiply a Decimal by a Power of Ten**
  1. Move the decimal point to the right the same number of places as the number of zeros in the power of 10.
  2. Add zeros at the end of the number as needed.
- **Divide Decimals**
  1. Determine the sign of the quotient.
  2. Make the divisor a whole number by “moving” the decimal point all the way to the right. “Move” the decimal point in the dividend the same number of places - adding zeros as needed.
  3. Divide. Place the decimal point in the quotient above the decimal point in the dividend.
  4. Write the quotient with the appropriate sign.
- **Convert a Decimal to a Proper Fraction**
  1. Determine the place value of the final digit.
  2. Write the fraction: numerator—the ‘numbers’ to the right of the decimal point; denominator—the place value corresponding to the final digit.
- **Convert a Fraction to a Decimal** Divide the numerator of the fraction by the denominator.

This page titled 1.8: Decimals is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.8: Decimals** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.8E: Exercises

Practice Makes Perfect

Needs to be integrated from https://cnx.org/contents/CImQfPDv@8.46:NciJ1qa0@19/1-7-Decimals

Name and Write Decimals

In the following exercises, write as a decimal.

531.

Twenty-nine and eighty-one hundredths

532.

Sixty-one and seventy-four hundredths

533.

Seven tenths

534.

Six tenths

535.

Twenty-nine thousandth

536.

Thirty-five thousandths

537.

Negative eleven and nine ten-thousandths

538.

Negative fifty-nine and two ten-thousandths

In the following exercises, name each decimal.

539.

5.5

540.

14.02

541.

8.71

542.

2.64

543.

0.002

544.

0.479

545.

−17.9−17.9

546.

−31.4−31.4

Round Decimals

In the following exercises, round each number to the nearest tenth.

547.

0.67

548.

0.49

549.

2.84

550.

4.63

In the following exercises, round each number to the nearest hundredth.

551.

0.845

552.

0.761

553.

0.299

554.

0.697

555.

4.098

556.

7.096

In the following exercises, round each number to the nearest ⓐ hundredth ⓑ tenth ⓒ whole number.

557.

5.781

558.

1.6381

559.

63.479

560.

84.28184.281

Add and Subtract Decimals

In the following exercises, add or subtract.

561.

16.92+7.5616.92+7.56

562.

248.25−91.29248.25−91.29

563.

21.76−30.9921.76−30.99

564.

38.6+13.6738.6+13.67

565.

−16.53−24.38−16.53−24.38

566.

−19.47−32.58−19.47−32.58

567.

−38.69+31.47−38.69+31.47

568.

29.83+19.7629.83+19.76

569.

72.5−10072.5−100

570.

86.2−10086.2−100

571.

15+0.7315+0.73

572.

27+0.8727+0.87

573.

91.95−(−10.462)91.95−(−10.462)

574.

94.69−(−12.678)94.69−(−12.678)

575.

55.01−3.755.01−3.7

576.

59.08−4.659.08−4.6

577.

2.51−7.42.51−7.4

578.

3.84−6.13.84−6.1

Multiply and Divide Decimals

In the following exercises, multiply.

579.

(0.24)(0.6)(0.24)(0.6)

580.

(0.81)(0.3)(0.81)(0.3)

581.

(5.9)(7.12)(5.9)(7.12)

582.

(2.3)(9.41)(2.3)(9.41)

583.

(−4.3)(2.71)(−4.3)(2.71)

584.

(−8.5)(1.69)(−8.5)(1.69)

585.

(−5.18)(−65.23)(−5.18)(−65.23)

586.

(−9.16)(−68.34)(−9.16)(−68.34)

587.

(0.06)(21.75)(0.06)(21.75)

588.

(0.08)(52.45)(0.08)(52.45)

589.

(9.24)(10)(9.24)(10)

590.

(6.531)(10)(6.531)(10)

591.

(55.2)(1000)(55.2)(1000)

592.

(99.4)(1000)(99.4)(1000)

In the following exercises, divide.

593.

4.75÷254.75÷25

594.

12.04÷4312.04÷43

595.

$117.25÷48$117.25÷48

596.

$109.24÷36$109.24÷36

597.

0.6÷0.20.6÷0.2

598.

0.8÷0.40.8÷0.4

599.

1.44÷(−0.3)1.44÷(−0.3)

600.

1.25÷(−0.5)1.25÷(−0.5)

601.

−1.75÷(−0.05)−1.75÷(−0.05)

602.

−1.15÷(−0.05)−1.15÷(−0.05)

603.

5.2÷2.55.2÷2.5

604.

6.5÷3.256.5÷3.25

605.

11÷0.5511÷0.55

606.

14÷0.3514÷0.35

Convert Decimals, Fractions and Percents

In the following exercises, write each decimal as a fraction.

607.

0.04

608.

0.19

609.

0.52

610.

0.78

611.

1.25

612.

1.35

613.

0.375

614.

0.464

615.

0.095

616.

0.085

In the following exercises, convert each fraction to a decimal.

617.

17201720

618.

13201320

619.

114114

620.

174174

621.

−31025−31025

622.

−28425−28425

623.

15111511

624.

18111811

625.

1511115111

626.

2511125111

627.

2.4+582.4+58

628.

3.9+9203.9+920

In the following exercises, convert each percent to a decimal.

629.

1%

630.

2%

631.

63%

632.

71%

633.

150%

634.

250%

635.

21.4%

636.

39.3%

637.

7.8%

638.

6.4%

In the following exercises, convert each decimal to a percent.

639.

0.01

640.

0.03

641.

1.35

642.

1.56

643.

3

644.

4

645.

0.0875

646.

0.0625

647.

2.254

648.

2.317

Everyday Math

649.

Salary Increase Danny got a raise and now makes $58,965.95 a year. Round this number to the nearest
ⓐ dollar
ⓑ thousand dollars
ⓒ ten thousand dollars.

650.

New Car Purchase Selena's new car cost $23,795.95. Round this number to the nearest
ⓐ dollar
ⓑ thousand dollars
ⓒ ten thousand dollars.

651.

Sales Tax Hyo Jin lives in San Diego. She bought a refrigerator for $1,624.99 and when the clerk calculated the sales tax it came out to exactly $142.186625. Round the sales tax to the nearest
ⓐ penny and
ⓑ dollar.

652.

Sales Tax Jennifer bought a $1,038.99 dining room set for her home in Cincinnati. She calculated the sales tax to be exactly $67.53435. Round the sales tax to the nearest
ⓐ penny and
ⓑ dollar.

653.

Paycheck Annie has two jobs. She gets paid $14.04 per hour for tutoring at City College and $8.75 per hour at a coffee shop. Last week she tutored for 8 hours and worked at the coffee shop for 15 hours.
ⓐ How much did she earn?
ⓑ If she had worked all 23 hours as a tutor instead of working both jobs, how much more would she have earned?

654.

Paycheck Jake has two jobs. He gets paid $7.95 per hour at the college cafeteria and $20.25 at the art gallery. Last week he worked 12 hours at the cafeteria and 5 hours at the art gallery.

ⓐ How much did he earn?

ⓑ If he had worked all 17 hours at the art gallery instead of working both jobs, how much more would he have earned?

Writing Exercises

655.

How does knowing about US money help you learn about decimals?

656.

Explain how you write “three and nine hundredths” as a decimal.

657.

Without solving the problem “44 is 80% of what number” think about what the solution might be. Should it be a number that is greater than 44 or less than 44? Explain your reasoning.

658.

When the Szetos sold their home, the selling price was 500% of what they had paid for the house 30 years ago. Explain what 500% means in this context.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
name and write decimals.			
round decimals.			
add and subtract decimals.			
multiply and divide decimals.			
convert decimals, fractions, and percents.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 1.8E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.8E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.9: The Real Numbers

Learning Objectives

By the end of this section, you will be able to:

- Simplify expressions with square roots
- Identify integers, rational numbers, irrational numbers, and real numbers
- Locate fractions on the number line
- Locate decimals on the number line

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapters, **Decimals** and **Properties of Real Numbers**.

Simplify Expressions with Square Roots

Remember that when a number n is multiplied by itself, we write n^2 and read it "n squared." The result is called the **square** of n. For example,

$$\begin{array}{ll} 8^2 & \text{read '8 squared'} \\ 64 & 64 \text{ is called the square of } 8\text{ .} \end{array}$$

Similarly, 121 is the square of 11, because 11^2 is 121.

SQUARE OF A NUMBER

If $n^2 = m$, then m is the **square** of n.

Note

Doing the Manipulative Mathematics activity "Square Numbers" will help you develop a better understanding of perfect square numbers.

Complete the following table to show the squares of the counting numbers 1 through 15.

Number	n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Square	n^2								64			121				

Figure 1.9.1

The numbers in the second row are called perfect square numbers. It will be helpful to learn to recognize the perfect square numbers.

The squares of the counting numbers are positive numbers. What about the squares of negative numbers? We know that when the signs of two numbers are the same, their product is positive. So the square of any negative number is also positive.

$$(-3)^2 = 9 \quad (-8)^2 = 64 \quad (-11)^2 = 121 \quad (-15)^2 = 225$$

Did you notice that these squares are the same as the squares of the positive numbers?

Sometimes we will need to look at the relationship between numbers and their squares in reverse. Because $10^2 = 100$, we say 100 is the square of 10. We also say that 10 is a *square root* of 100. A number whose square is mm is called a **square root** of m.

SQUARE ROOT OF A NUMBER

If $n^2 = m$, then n is a **square root** of m.

Notice $(-10)^2 = 100$ also, so -10 is also a square root of 100. Therefore, both 10 and -10 are square roots of 100.

So, every positive number has two square roots—one positive and one negative. What if we only wanted the positive square root of a positive number? The **radical sign**, $\sqrt{m}$, denotes the positive square root. The positive square root is called the **principal square root**. When we use the radical sign that always means we want the principal square root.

We also use the radical sign for the square root of zero. Because $0^2 = 0$, $\sqrt{0} = 0$. Notice that zero has only one square root.

SQUARE ROOT NOTATION

$\sqrt{m}$ is read "the square root of m"

radical sign $\rightarrow \sqrt{m} \leftarrow$ radicand

If $m = n^2$, then $\sqrt{m} = n$, for $n \geq 0$.

The square root of m, $\sqrt{m}$, is the positive number whose square is m.

Since 10 is the principal square root of 100, we write $\sqrt{100} = 10$. You may want to complete the following table to help you recognize square roots.

$\sqrt{1}$	$\sqrt{4}$	$\sqrt{9}$	$\sqrt{16}$	$\sqrt{25}$	$\sqrt{36}$	$\sqrt{49}$	$\sqrt{64}$	$\sqrt{81}$	$\sqrt{100}$	$\sqrt{121}$	$\sqrt{144}$	$\sqrt{169}$	$\sqrt{196}$	$\sqrt{225}$
									10					

Figure 1.9.2

Exercise 1.9.1

Simplify:

1. $\sqrt{25}$
2. $\sqrt{121}$

Answer

1. $\sqrt{25} = 5$, since $5^2 = 25$.
2. $\sqrt{121} = 11$, since $11^2 = 121$.

Exercise 1.9.2

Simplify:

1. $\sqrt{36}$
2. $\sqrt{169}$

Answer

1. $\sqrt{36} = 6$
2. $\sqrt{169} = 13$

Exercise 1.9.3

Simplify:

1. $\sqrt{16}$
2. $\sqrt{196}$

Answer

1. $\sqrt{16} = 4$
2. $\sqrt{196} = 14$

We know that every positive number has two square roots and the radical sign indicates the positive one. We write $\sqrt{100}=10$. If we want to find the negative square root of a number, we place a negative in front of the radical sign. For example, $-\sqrt{100}=-10$. We read $-\sqrt{100}$ as "the opposite of the square root of 10."

Exercise 1.9.4

Simplify:

1. $-\sqrt{9}$
2. $-\sqrt{144}$

Answer

1. $-\sqrt{9}=-3$, since the negative is in front of the radical sign.
2. $-\sqrt{144}=-12$, since the negative is in front of the radical sign.

Exercise 1.9.5

Simplify:

1. $-\sqrt{4}$
2. $-\sqrt{225}$

Answer

1. $-\sqrt{4}=-2$
2. $-\sqrt{225}=-15$

Exercise 1.9.6

Simplify:

1. $-\sqrt{81}$
2. $-\sqrt{100}$

Answer

1. $-\sqrt{81}=-9$
2. $-\sqrt{100}=-10$

Identify Integers, Rational Numbers, Irrational Numbers, and Real Numbers

We have already described numbers as ***counting numbers***, ***whole numbers***, and ***integers***. What is the difference between these types of numbers?

Counting numbers	$1, 2, 3, 4, \ldots$
Whole numbers	$0, 1, 2, 3, 4, \ldots$
Integers	$\cdots -3, -2, -1, 0, 1, 2, 3, \ldots$

What type of numbers would we get if we started with all the integers and then included all the fractions? The numbers we would have form the set of rational numbers. A **rational number** is a number that can be written as a ratio of two integers.

RATIONAL NUMBER

A **rational number** is a number of the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$

A rational number can be written as the ratio of two integers.

All signed fractions, such as $\frac{4}{5}, -\frac{7}{8}, \frac{13}{4}, -\frac{20}{3}$ are rational numbers. Each numerator and each denominator is an integer.

Are integers rational numbers? To decide if an integer is a rational number, we try to write it as a ratio of two integers. Each integer can be written as a ratio of integers in many ways. For example, 3 is equivalent to $\frac{3}{1}, -\frac{6}{2}, \frac{9}{3}, \frac{12}{4}, -\frac{15}{5} \ldots$

An easy way to write an integer as a ratio of integers is to write it as a fraction with denominator one.

$$3 = \frac{3}{1} \quad -8 = -\frac{8}{1} \quad 0 = \frac{0}{1}$$

Since any integer can be written as the ratio of two integers, *all integers are rational numbers*! Remember that the counting numbers and the whole numbers are also integers, and so they, too, are rational.

What about decimals? Are they rational? Let's look at a few to see if we can write each of them as the ratio of two integers.

We've already seen that integers are rational numbers. The integer -8 could be written as the decimal -8.0. So, clearly, some decimals are rational.

Think about the decimal 7.3. Can we write it as a ratio of two integers? Because 7.3 means $7\frac{3}{10}$, we can write it as an improper fraction, $\frac{73}{10}$. So 7.3 is the ratio of the integers 73 and 10. It is a rational number.

In general, any decimal that ends after a number of digits (such as 7.3 or -1.2684) is a rational number. We can use the place value of the last digit as the denominator when writing the decimal as a fraction.

? Exercise 1.9.7

Write as the ratio of two integers:

1. -27
2. 7.31

Answer

1. Writing -27 as a fraction with denominator 1, we get $\frac{-27}{1}$.
2. For 7.31, we have two ways to think of it. We could see that it is equal to 731 hundredths or $\frac{731}{100}$,

   or we could start by rewriting 7.31 as a mixed number. That is, $7.31 = 7\frac{31}{100}$. Then we can convert this mixed number to an improper fraction to get $\frac{731}{100}$.

Since both numbers can be written as the ratio of two integers, we see that -27 and 7.31 are both rational numbers.

? Exercise 1.9.8

Write as the ratio of two integers:

1. -24
2. 3.57

Answer

1. $-24 = \frac{-24}{1}$
2. $3.57 = \frac{357}{100}$

Exercise 1.9.9

Write as the ratio of two integers:

1. -19
2. 8.41

Answer

1. $-19 = \dfrac{-19}{1}$
2. $8.41 = \dfrac{841}{100}$

Let's look at the decimal form of the numbers we know are rational.

We have seen that *every integer is a rational number*, since $a = \dfrac{a}{1}$ for any integer, a. We can also change any integer to a decimal by adding a decimal point and a zero.

Integer	-2	-1	0	1	2	3
Decimal form	-2.0	-1.0	0.0	1.0	2.0	3.0

These decimal numbers stop.

We have also seen that *every fraction is a rational number*. Look at the decimal form of the fractions we considered above.

Ratio of integers	$\frac{4}{5}$	$-\frac{7}{8}$	$\frac{13}{4}$	$-\frac{20}{3}$
The decimal form	0.8	-0.875	3.25	$-6.666\ldots$
			$-6.\overline{6}$	

These decimal numbers either stop or repeat.

What do these examples tell us?

Every rational number can be written both as a ratio of integers, ($\dfrac{p}{q}$, where p and q are integers and $q \neq 0$), and as a decimal that either stops or repeats.

Here are the numbers we looked at above expressed as a ratio of integers and as a decimal:

Table 1.9.1

Fractions					Integers					
Number	$\frac{4}{5}$	$\frac{7}{8}$	$\frac{13}{4}$	$-\frac{20}{3}$	-2	-1	0	1	2	3
Ratio of Integers	$\frac{4}{5}$	$\frac{7}{8}$	$\frac{13}{4}$	$-\frac{20}{3}$	$-\frac{2}{1}$	$-\frac{1}{1}$	$\frac{0}{1}$	$\frac{1}{1}$	$\frac{2}{1}$	$\frac{3}{1}$
Decimal Form	0.8	−0.875	3.25	$-6.\overline{6}$	−2.0	−1.0	0.0	1.0	2.0	3.0

RATIONAL NUMBER

A **rational number** is a number of the form $\frac{p}{q}$, where p and q are integers and $q \neq 0$

Its decimal form stops or repeats.

Are there any decimals that do not stop or repeat? Yes!

The number π (the Greek letter *pi*, pronounced "pie"), which is very important in describing circles, has a decimal form that does not stop or repeat.

$$\pi = 3.141592654\ldots \tag{1.9.1}$$

We can even create a decimal pattern that does not stop or repeat, such as

$$2.01001000100001\ldots \tag{1.9.2}$$

Numbers whose decimal form does not stop or repeat cannot be written as a fraction of integers. We call these numbers irrational.

IRRATIONAL NUMBER

An **irrational number** is a number that cannot be written as the ratio of two integers.

Its decimal form does not stop and does not repeat.

Let's summarize a method we can use to determine whether a number is rational or irrational.

RATIONAL OR IRRATIONAL?

If the decimal form of a number

- *repeats or stops*, the number is **rational**.
- *does not repeat and does not stop*, the number is **irrational**.

Exercise 1.9.10

Given the numbers $0.58\overline{3}, 0.47, 3.605551275\ldots$ list the

1. rational numbers
2. irrational numbers.

Answer

1.	Look for decimals that repeat or stop	The 3 repeats in $0.58\overline{3}$. The decimal 0.47 stops after the 7. So $0.58\overline{3}$ and $0.47$are rational
2.	Look for decimals that repeat or stop	$3.605551275\ldots$ has no repeating block of digits and it does not stop. So $3.605551275\ldots$ is irrational.

Exercise 1.9.11

For the given numbers list the

1. rational numbers
2. irrational numbers: $0.29, 0.81\overline{6}, 2.515115111\ldots.$

Answer

1. $0.29, 0.81\overline{6}$
2. $2.515115111\ldots.$

Exercise 1.9.12

For the given numbers list the

1. rational numbers
2. irrational numbers: $2.6\overline{3}, 0.125, 0.418302\ldots$

Answer

1. $2.6\overline{3}, 0.125$

2. $0.418302\ldots$

Exercise 1.9.13

For each number given, identify whether it is rational or irrational:

1. $\sqrt{36}$
2. $\sqrt{44}$

Answer

1. Recognize that 36 is a perfect square, since $6^2 = 36$. So $\sqrt{36} = 6$, therefore $\sqrt{36}$ is rational.
2. Remember that $6^2 = 36$ and $7^2 = 49$, so 44 is not a perfect square. Therefore, the decimal form of $\sqrt{44}$ will never repeat and never stop, so $\sqrt{44}$ is irrational.

Exercise 1.9.14

For each number given, identify whether it is rational or irrational:

1. $\sqrt{81}$
2. $\sqrt{17}$

Answer

1. rational
2. irrational

Exercise 1.9.15

For each number given, identify whether it is rational or irrational:

1. $\sqrt{116}$
2. $\sqrt{121}$

Answer

1. irrational
2. rational

We have seen that all counting numbers are whole numbers, all whole numbers are integers, and all integers are rational numbers. The irrational numbers are numbers whose decimal form does not stop and does not repeat. When we put together the rational numbers and the irrational numbers, we get the set of **real numbers**.

REAL NUMBER

A **real number** is a number that is either rational or irrational.

All the numbers we use in elementary algebra are real numbers. Figure 1.9.3 illustrates how the number sets we've discussed in this section fit together.




Figure 1.9.3: This chart shows the number sets that make up the set of real numbers. Does the term “real numbers” seem strange to you? Are there any numbers that are not “real,” and, if so, what could they be?

Can we simplify $\sqrt{-25}$? Is there a number whose square is -25?

$$(\quad)^2 = -25?$$

None of the numbers that we have dealt with so far has a square that is -25. Why? Any positive number squared is positive. Any negative number squared is positive. So we say there is no real number equal to $\sqrt{-25}$.

The square root of a negative number is not a real number.

Exercise 1.9.16

For each number given, identify whether it is a real number or not a real number:

1. $\sqrt{-169}$
2. $-\sqrt{64}$

Answer

1. There is no real number whose square is -169. Therefore, $\sqrt{-169}$ is not a real number.
2. Since the negative is in front of the radical, $-\sqrt{64}$ is -8, Since -8 is a real number, $-\sqrt{64}$ is a real number.

Exercise 1.9.17

For each number given, identify whether it is a real number or not a real number:

1. $\sqrt{-196}$
2. $-\sqrt{81}$

Answer

1. not a real number
2. real number

Exercise 1.9.18

For each number given, identify whether it is a real number or not a real number:

1. $-\sqrt{49}$
2. $\sqrt{-121}$

Answer

1. real number

2. not a real number

Exercise 1.9.19

Given the numbers $-7, \frac{14}{5}, 8, \sqrt{5}, 5.9, \sqrt{64}$ list the

1. whole numbers
2. integers
3. rational numbers
4. irrational numbers
5. real numbers

Answer

1. Remember, the whole numbers are 0, 1, 2, 3, … and 8 is the only whole number given.
2. The integers are the whole numbers, their opposites, and 0. So the whole number 8 is an integer, and -7 is the opposite of a whole number so it is an integer, too. Also, notice that 64 is the square of 8 so $-\sqrt{64} = -8$. So the integers are $-7, 8, \sqrt{64}$.
3. Since all integers are rational, then $-7, 8, -\sqrt{64}$ are rational. Rational numbers also include fractions and decimals that repeat or stop, so $\frac{14}{5}$ and 5.9 are rational. So the list of rational numbers is $-7, \frac{14}{5}, 8, 5.9, \sqrt{64}$
4. Remember that 5 is not a perfect square, so $\sqrt{5}$ is irrational.
5. All the numbers listed are real numbers.

Exercise 1.9.20

For the given numbers, list the

1. whole numbers
2. integers
3. rational numbers
4. irrational numbers
5. real numbers: $-3, -\sqrt{2}, 0.\overline{3}, \frac{9}{5}, 4, \sqrt{49}$

Answer

1. $4, \sqrt{49}$.
2. $-3, 4, \sqrt{49}$
3. $-3, 0.\overline{3}, \frac{9}{5}, 4, \sqrt{49}$
4. $-\sqrt{2}$
5. $-3, \sqrt{2}, 0.\overline{3}, \frac{9}{5}, 4, \sqrt{49}$

Exercise 1.9.21

For the given numbers, list the

1. whole numbers
2. integers
3. rational numbers
4. irrational numbers
5. real numbers: $-\sqrt{25}, -\frac{3}{8}, -1, 6, \sqrt{121}, 2.041975\ldots$

Answer

1. $6, \sqrt{121}$.
2. $-\sqrt{25}, -1, 6, \sqrt{121}$
3. $-\sqrt{25}, -\frac{3}{8}, -1, 6, \sqrt{121}$

4. $2.041975\ldots$
5. $-\sqrt{25}, -\frac{3}{8}, -1, 6, \sqrt{121}, 2.041975\ldots$

Locate Fractions on the Number Line

The last time we looked at the **number line**, it only had positive and negative integers on it. We now want to include **fraction**s and decimals on it.

Note

Doing the Manipulative Mathematics activity "Number Line Part 3" will help you develop a better understanding of the location of fractions on the number line.

Let's start with fractions and locate $\frac{1}{5}, -\frac{4}{5}, 3, \frac{7}{4}, -\frac{9}{2}, -5$ and $\frac{8}{3}$ on the number line.

We'll start with the whole numbers 3 and −5. because they are the easiest to plot. See Figure 1.9.4.

The **proper fractions** listed are $\frac{1}{5}$ and $-\frac{4}{5}$. We know the proper fraction $\frac{1}{5}$ has value less than one and so would be located between 0 and 1. The denominator is 5, so we divide the unit from 0 to 1 into 5 equal parts $\frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}$. We plot $\frac{1}{5}$. See Figure 1.9.4.

Similarly, $-\frac{4}{5}$ is between 0 and −1. After dividing the unit into 5 equal parts we plot $-\frac{4}{5}$. See Figure 1.9.4.

Finally, look at the **improper fractions** $\frac{7}{4}, -\frac{9}{2}, \frac{8}{3}$. These are fractions in which the numerator is greater than the denominator. Locating these points may be easier if you change each of them to a mixed number. See Figure 1.9.4.

$$\frac{7}{4} = 1\frac{3}{4} \quad -\frac{9}{2} = -4\frac{1}{2} \quad \frac{8}{3} = 2\frac{2}{3} \tag{1.9.3}$$

Figure 1.9.4 shows the number line with all the points plotted.




Figure 1.9.4

Exercise 1.9.22


Locate and label the following on a number line: $4, \frac{3}{4}, -\frac{1}{4}, -3, \frac{6}{5}, -\frac{5}{2}$ and $\frac{7}{3}$.

Answer

Locate and plot the integers, 4,−3.

Locate the proper fraction $\frac{3}{4}$ first. The fraction $\frac{3}{4}$ is between 0 and 1. Divide the distance between 0 and 1 into four equal parts then, we plot $\frac{3}{4}$. Similarly plot $-\frac{1}{4}$.


Now locate the improper fractions $\frac{6}{5}, -\frac{5}{2}, \frac{7}{3}$. It is easier to plot them if we convert them to mixed numbers and then plot them as described above: $\frac{6}{5} = 1\frac{1}{5}, -\frac{5}{2} = -2\frac{1}{2}, \frac{7}{3} = 2\frac{1}{3}$.



Exercise 1.9.23

Locate and label the following on a number line: $-1, \frac{1}{3}, \frac{6}{5}, -\frac{7}{4}, \frac{9}{2}, 5$ and $-\frac{8}{3}$.


Answer



Exercise 1.9.24

Locate and label the following on a number line: $\frac{1}{5}, -\frac{4}{5}, 3, \frac{7}{4}, -\frac{9}{2}, -5$ and $\frac{8}{3}$.

Answer



In Exercise 1.9.25, we'll use the inequality symbols to order fractions. In previous chapters we used the number line to order numbers.

- $a < b$ "a is less than b" when a is to the left of b on the number line.
- $a < b$ " a is greater than b" when a is to the right of b on the number line.

As we move from left to right on a number line, the values increase.

Exercise 1.9.25

Order each of the following pairs of numbers, using $<$ or $>$. It may be helpful to refer Figure 1.9.5.

1. $-\frac{2}{3}$___ -1
2. $-3\frac{1}{2}$___ -3
3. $-\frac{3}{4}$___ $-\frac{1}{4}$
4. -2___ $-\frac{8}{3}$




Figure 1.9.5

Answer

Be careful when ordering negative numbers.

1. Since $-\frac{2}{3}$ is to the right of -1 on the number line, we know that $-\frac{2}{3} > -1$.
2. Since $-3\frac{1}{2}$ is to the left of -3 on the number line, we know that $-3\frac{1}{2} < -3$.
3. Since $-\frac{3}{4}$ is to the left of $-\frac{1}{4}$ on the number line, we know that $-\frac{3}{4} < -\frac{1}{4}$.
4. Since -2 is to the right of $-\frac{8}{3}$ on the number line, we know that $-2 > -\frac{8}{3}$.

Exercise 1.9.26

Order each of the following pairs of numbers, using $<$ or $>$.

1. $-\frac{1}{3}$___ -1
2. $-1\frac{1}{2}$___ -2
3. $-\frac{2}{3}$___ $-\frac{1}{3}$
4. -3___ $-\frac{7}{3}$

Answer

1. $-\frac{1}{3} > -1$
2. $-1\frac{1}{2} > -2$

3. $-\frac{2}{3} < -\frac{1}{3}$
4. $-3 < -\frac{7}{3}$

Exercise 1.9.27

Order each of the following pairs of numbers, using $<$ or $>$.

1. $-1 \_\_\_ -\frac{2}{3}$
2. $-2\frac{1}{4} \_\_\_ -2$
3. $-\frac{3}{5} \_\_\_ -\frac{4}{5}$
4. $-4 \_\_\_ -\frac{10}{3}$

Answer

1. $-1 < -\frac{2}{3}$
2. $-2\frac{1}{4} < -2$
3. $-\frac{3}{5} > -\frac{4}{5}$
4. $-4 < -\frac{10}{3}$

Locate Decimals on the Number Line

Since decimals are forms of fractions, locating decimals on the number line is similar to locating fractions on the number line.

Exercise 1.9.28

Locate 0.4 on the number line.

Answer

A proper fraction has value less than one. The decimal number 0.4 is equivalent to $\frac{4}{10}$, a proper fraction, so 0.4 is located between 0 and 1. On a number line, divide the interval between 0 and 1 into 10 equal parts. Now label the parts $0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0$. We write 0 as 0.0 and 1 and 1.0, so that the numbers are consistently in tenths. Finally, mark 0.4 on the number line. See Figure 1.9.6




Figure 1.9.6

Exercise 1.9.29

Locate on the number line: 0.6.

Answer



Exercise 1.9.30

Locate on the number line: 0.9.

Answer



Exercise 1.9.31

Locate -0.74 on the number line.

Answer

The decimal (−0.74\) is equivalent to $-\frac{74}{100}$, so it is located between 0 and −1. On a number line, mark off and label the hundredths in the interval between 0 and −1. See Figure 1.9.7.





Figure 1.9.7

Exercise 1.9.32

Locate on the number line: −0.6.


Answer



Exercise 1.9.33

Locate on the number line: −0.7.

Answer



Which is larger, 0.04 or 0.40? If you think of this as money, you know that \$0.40 (forty cents) is greater than \$0.04 (four cents). So, $0.40 > 0.04$

Again, we can use the number line to order numbers.

- $a < b$ "a is less than b" when a is to the left of b on the number line.
- $a < b$ " a is greater than b" when a is to the right of b on the number line.

Where are 0.04 and 0.40 located on the number line? See Figure 1.9.8.




Figure 1.9.8

We see that 0.40 is to the right of 0.04 on the number line. This is another way to demonstrate that $0.40 > 0.04$.

How does 0.31 compare to 0.308? This doesn't translate into money to make it easy to compare. But if we convert 0.31 and 0.308 into fractions, we can tell which is larger.

Table 1.9.2

	0.31	0.308
Convert to fractions.	$\frac{31}{100}$	$\frac{308}{1000}$
We need a common denominator to compare them.	$\frac{31 \cdot 10}{100 \cdot 10}$	$\frac{308}{1000}$
	$\frac{310}{1000}$	$\frac{308}{1000}$

Because $310 > 308$, we know that $\frac{310}{1000} > \frac{308}{1000}$. Therefore, $0.31 > 0.308$.

Notice what we did in converting 0.31 to a fraction—we started with the fraction $\frac{31}{100}$ and ended with the equivalent fraction $\frac{310}{1000}$.Converting $\frac{310}{1000}$ back to a decimal gives 0.310. So 0.31 is equivalent to 0.310. Writing zeros at the end of a decimal does not change its value!

$$\frac{31}{100} = \frac{310}{1000} \quad \text{and} \quad 0.31 = 0.310$$

We say 0.31 and 0.310 are **equivalent decimals**.

EQUIVALENT DECIMALS

Two decimals are equivalent if they convert to equivalent fractions.

We use equivalent decimals when we order decimals.

The steps we take to order decimals are summarized here.

ORDER DECIMALS.

1. Write the numbers one under the other, lining up the decimal points.
2. Check to see if both numbers have the same number of digits. If not, write zeros at the end of the one with fewer digits to make them match.
3. Compare the numbers as if they were whole numbers.
4. Order the numbers using the appropriate inequality sign.

Exercise 1.9.34

Order 0.64 ___ 0.6using $<$ or $>$.

Answer

Write the numbers one under the other, lining up the decimal points.	0.64 0.6
Add a zero to 0.6 to make it a decimal with 2 decimal places. Now they are both hundredths.	0.64 0.60
64 is greater than 60.	$64 > 60$
64 hundredths is greater than 60 hundredths.	$0.64 > 0.60$
	$0.64 > 0.6$

Exercise 1.9.35

Order each of the following pairs of numbers, using $<$ or $>$: 0.42 ___ 0.4

Answer

$0.42 > 0.4$

https://math.libretexts.org/@go/page/30465

Exercise 1.9.36

Order each of the following pairs of numbers, using $<$ or $>$: 0.18 ___ 0.1.

Answer

$0.18 > 0.1$

Exercise 1.9.37

Order 0.83 ___ 0.803using $<$ or $>$.

Answer

	0.83 ___ 0.803
Write the numbers one under the other, lining up the decimal points.	0.83 0.803
They do not have the same number of digits. Write one zero at the end of 0.83.	0.830 0.803
Since $830 > 803$, 830 hundredths is greater than 803 thousandths.	$0.830 > 0.803$
	$0.83 > 0.803$

Exercise 1.9.38

Order each of the following pairs of numbers, using $<$ or $>$: 0.76 ___ 0.706

Answer

$0.76 > 0.706$

Exercise 1.9.39

Order each of the following pairs of numbers, using $<$ or $>$: 0.305 ___ 0.35

Answer

$0.305 < 0.35$

When we order negative decimals, it is important to remember how to order negative integers. Recall that larger numbers are to the right on the number line. For example, because -2 lies to the right of -3 on the number line, we know that $-2 > -3$. Similarly, smaller numbers lie to the left on the number line. For example, because -9 lies to the left of -6 on the number line, we know that $-9 < -6$. See Figure 1.9.9.




Figure 1.9.9

If we zoomed in on the interval between 0 and -1, as shown in Exercise 1.9.40, we would see in the same way that $-0.2 > -0.3$ and $-0.9 < -0.6$.

https://math.libretexts.org/@go/page/30465

? Exercise 1.9.40

Use $<$ or $>$ to order -0.1 ___ -0.8.

Answer

	-0.1 ___ -0.8
Write the numbers one under the other, lining up the decimal points. They have the same number of digits.	-0.1 -0.8
Since $-1 > -8$, we know that -1 tenth is greater than -8 tenths. So, we know that	$-0.1 > -0.8$

? Exercise 1.9.41

Order the following pair of numbers, using $<$ or $>$: -0.3 ___ -0.5.

Answer

$-0.3 > -0.5$

? Exercise 1.9.42

Order the following pair of numbers, using $<$ or $>$: -0.6 ___ -0.7.

Answer

$-0.6 > -0.7$

Key Concepts

- **Square Root Notation**
  $\sqrt{m}$ is read 'the square root of m.' If $m = n^2$, then $\sqrt{m} = n$, for $n \geq 0$.
- **Order Decimals**
  1. Write the numbers one under the other, lining up the decimal points.
  2. Check to see if both numbers have the same number of digits. If not, write zeros at the end of the one with fewer digits to make them match.
  3. Compare the numbers as if they were whole numbers.
  4. Order the numbers using the appropriate inequality sign.

Practice Makes Perfect

Simplify Expressions with Square Roots

In the following exercises, simplify.

This page titled 1.9: The Real Numbers is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.9: The Real Numbers** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.9E: Exercises

Practice Makes Perfect

Simplify Expressions with Square Roots

In the following exercises, simplify.

1. $\sqrt{36}$

Answer

$\sqrt{36} = 6$

2. $\sqrt{4}$
3. $\sqrt{64}$

Answer

$\sqrt{64} = 8$

4. $\sqrt{169}$
5. $\sqrt{9}$

Answer

$\sqrt{9} = 3$

6. $\sqrt{16}$
7. $\sqrt{100}$

Answer

$\sqrt{100} = 10$

8. $\sqrt{144}$
9. $\sqrt{-4}$

Answer

$\sqrt{-4}$ is not a real number.

10. $\sqrt{-100}$
11. $\sqrt{-1}$

Answer

$\sqrt{-1}$ is not a real number.

12. $\sqrt{-121}$

Identify Integers, Rational Numbers, Irrational Numbers, and Real Numbers

In the following exercises, write each rational number as the ratio of two integers.

13. ⓐ 5 ⓑ 3.19

Answer

a. $5 = \frac{5}{1}$ b. $3.19 = \frac{319}{100}$

14. ⓐ 8 ⓑ 1.61

15. ⓐ -12 ⓑ 9.279

Answer

a. $-12 = \frac{-12}{1}$ b. $9.279 = \frac{9279}{1000}$

16. ⓐ -16 ⓑ 4.399

In the following exercises, list the ⓐ rational numbers, ⓑ irrational numbers

17. $\{0.75, 0.22\overline{3}, 1.39174\}$

Answer

All of these numbers are rational.

18. $\{0.36, 0.94729\ldots, 2.52\overline{8}\}$

19. $\{0.4\overline{5}, 1.919293\ldots, 3.59\}$

Answer

a. $0.4\overline{5}$and 3.59 are rational numbers. b. 1.919293… is irrational, since it does not terminate or repeat.

20. $\{0.1\overline{3}, 0.42982\ldots, 1.875\}$

In the following exercises, identify whether each number is rational or irrational.

21. ⓐ $\sqrt{25}$ ⓑ $\sqrt{30}$

Answer

a. $\sqrt{25} = 5$ is rational. b. $\sqrt{30}$ is irrational.

22. ⓐ $\sqrt{44}$ ⓑ $\sqrt{49}$

23. ⓐ $\sqrt{164}$ ⓑ $\sqrt{169}$

Answer

a. $\sqrt{164}$ is irrational. b. $\sqrt{169} = 13$ is rational.

24. ⓐ $\sqrt{225}$ ⓑ $\sqrt{216}$

In the following exercises, identify whether each number is a real number or not a real number.

25. ⓐ $-\sqrt{81}$ ⓑ $\sqrt{-121}$

Answer

a. $-\sqrt{81} = -9$ is a real number. b. $\sqrt{-121}$ is not a real number.

26. ⓐ $-\sqrt{64}$ ⓑ $\sqrt{-9}$

27. ⓐ $\sqrt{-36}$ ⓑ $-\sqrt{144}$

Answer

a. $\sqrt{-36}$ is not a real number. b. $-\sqrt{144} = -12$ is a real number.

28. ⓐ $\sqrt{-49}$ ⓑ $-\sqrt{144}$

In the following exercises, for each set of numbers, list the ⓐ whole numbers, ⓑ integers, ⓒ rational numbers, ⓓ irrational numbers, ⓔ real numbers.

29. $\{-8, 0, 1.95286\ldots, \frac{12}{5}, \sqrt{36}, 9\}$

Answer

a. whole numbers: 0, $\sqrt{36}=6$, 9
b. integers: -8, 0, $\sqrt{36}=6$, 9
c. rational numbers: -8, 0, $\sqrt{36}=6$, 9, $\frac{12}{5}$
d. irrational numbers: 1.95286…
e. real numbers: All of these numbers are real numbers.

30. $\{-9, -3\frac{4}{9}, -\sqrt{9}, 0.40\overline{9}, 116, 7\}$

31. $\{-\sqrt{100}, -7, -\frac{8}{3}, -1, 0.77, 3\frac{1}{4}\}$

Answer

a. whole numbers: None of these numbers are whole numbers.
b. integers: $-\sqrt{100}=-10$, −7, -1
c. rational numbers: All of these numbers are rational.
d. irrational numbers: None of these numbers is irrational.
e. real numbers: All of these numbers are real numbers.


32. $\{-6, -\frac{5}{2}, 0, 0.\overline{714285}, 2\frac{1}{5}, \sqrt{14}\}$

Locate Fractions on the Number Line

In the following exercises, locate the numbers on a number line.

33. $\frac{3}{4}, \frac{8}{5}, \frac{10}{3}$


Answer



34. $\frac{1}{4}, \frac{9}{5}, \frac{11}{3}$

35. $\frac{3}{10}, \frac{7}{2}, \frac{11}{6}, 4$


Answer



36. $\frac{7}{10}, \frac{5}{2}, \frac{13}{8}, 3$

37. $\frac{2}{5}, -\frac{2}{5}$


Answer



38. $\frac{3}{4}, -\frac{3}{4}$

39. $\frac{3}{4}, -\frac{3}{4}, 1\frac{2}{3}, -1\frac{2}{3}, \frac{5}{2}, -\frac{5}{2}$

Answer



40. $\frac{1}{5}, -\frac{2}{5}, 1\frac{3}{4}, -1\frac{3}{4}, \frac{8}{3}, -\frac{8}{3}$

In the following exercises, order each of the pairs of numbers, using < or >.

41. -1 ___ $-\frac{1}{4}$

Answer

$-1 < -\frac{1}{4}$

42. -1 ___ $-\frac{1}{3}$

43. $-2\frac{1}{2}$ ___ -3

Answer

$-2\frac{1}{2} > -3$

44. $-1\frac{3}{4}$ ___ -2

45. $-\frac{5}{12}$ ___ $-\frac{7}{12}$

Answer

$-\frac{5}{12} > -\frac{7}{12}$

46. $-\frac{9}{10}$ ___ $-\frac{3}{10}$

47. -3 ___ $-\frac{13}{5}$

Answer


$-3 < -\frac{13}{5}$

48. -4 ___ $-\frac{23}{6}$

Locating Decimals on the Number Line In the following exercises, locate the number on the number line.

49. 0.8

Answer



50. −0.9

51. −1.6

Answer



52. 3.1

In the following exercises, order each pair of numbers, using < or >.

53. 0.37 ___ 0.63

Answer

$0.37 < 0.63$

54. 0.86 ___ 0.69

55. 0.91 ___ 0.901

Answer

$0.91 > 0.901$

56. 0.415 ___ 0.41

57. -0.5 ___ -0.3

Answer

$-0.5 < -0.3$

58. -0.1 ___ -0.4

59. -0.62 ___ -0.619

Answer

$-0.62 < -0.619$

60. -7.31 ___ -7.3

Everyday Math

61. **Field trip** All the 5th graders at Lincoln Elementary School will go on a field trip to the science museum. Counting all the children, teachers, and chaperones, there will be 147 people. Each bus holds 44 people.

ⓐ How many busses will be needed?
ⓑ Why must the answer be a whole number?
ⓒ Why shouldn't you round the answer the usual way, by choosing the whole number closest to the exact answer?

62. **Child care** Serena wants to open a licensed child care center. Her state requires there be no more than 12 children for each teacher. She would like her child care center to serve 40 children.

ⓐ How many teachers will be needed?
ⓑ Why must the answer be a whole number?
ⓒ Why shouldn't you round the answer the usual way, by choosing the whole number closest to the exact answer?

Writing Exercises

63. In your own words, explain the difference between a rational number and an irrational number.

64. Explain how the sets of numbers (counting, whole, integer, rational, irrationals, reals) are related to each other.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objective of this section.

I can...	Confidently	With some help	No-I don't get it!
simplify expressions with square roots.			
identify integers, rational numbers, irrational numbers, and real numbers.			
locate fractions on the number line.			
locate decimals on the number line.			

ⓑ On a scale of 1−10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 1.9E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.9E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.10: Properties of Real Numbers

Learning Objectives

By the end of this section, you will be able to:

- Use the commutative and associative properties
- Use the identity and inverse properties of addition and multiplication
- Use the properties of zero
- Simplify expressions using the distributive property

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapter, **The Properties of Real Numbers**.

Use the Commutative and Associative Properties

Think about adding two numbers, say 5 and 3. The order we add them doesn't affect the result, does it?

$$\begin{array}{ll} 5+3 & 3+5 \\ 8 & 8 \\ 5+3= & 3+5 \end{array}$$

The results are the same.

As we can see, the order in which we add does not matter!

What about multiplying 5 and 3?

$$\begin{array}{ll} 5\cdot 3 & 3\cdot 5 \\ 15 & 15 \\ 5\cdot 3= & 3\cdot 5 \end{array}$$

Again, the results are the same!

The order in which we multiply does not matter!

These examples illustrate the **commutative property**. When adding or multiplying, changing the *order* gives the same result.

COMMUTATIVE PROPERTY

of Addition	If a, b are real numbers, then	$a+b=b+a$
of Multiplication	If a, b are real numbers, then	$a\cdot b=b\cdot a$

When adding or multiplying, changing the *order* gives the same result.

The commutative property has to do with order. If you change the order of the numbers when adding or multiplying, the result is the same.

What about subtraction? Does order matter when we subtract numbers? Does $7-3$ give the same result as $3-7$?

$$\begin{array}{ll} 7-3 & 3-7 \\ 4 & -4 \end{array}$$

$$4 \neq -4$$

$$7-3 \neq 3-7$$

The results are not the same.

Since changing the order of the subtraction did not give the same result, we know that *subtraction is not commutative*.

Let's see what happens when we divide two numbers. Is **division** commutative?

$$\begin{array}{ll} 12 \div 4 & 4 \div 12 \\ \frac{12}{4} & \frac{4}{12} \\ 3 & \frac{1}{3} \end{array}$$

$$3 \neq \frac{1}{3}$$
$$12 \div 4 \neq 4 \div 12$$

The results are not the same.

Since changing the order of the division did not give the same result, *division is not commutative*. The commutative properties only apply to addition and multiplication!

- Addition and multiplication *are* commutative.
- Subtraction and Division *are not* commutative.

If you were asked to simplify this expression, how would you do it and what would your answer be?

$$7+8+2$$

Some people would think $7+8$ is 15 and then $15+2$ is 17. Others might start with $8+2$ makes 10 and then $7+10$ makes 17.

Either way gives the same result. Remember, we use parentheses as grouping symbols to indicate which operation should be done first.

Add $7+8$.	$(7+8)+2$
Add.	$15+2$
Add.	17

	$7+(8+2)$
Add $8+2$.	$7+10$
Add.	77

$$(7+8)+2=7+(8+2)$$

When adding three numbers, changing the grouping of the numbers gives the same result.

This is true for multiplication, too.

	$(5 \cdot \frac{1}{3}) \cdot 3$
Multiply. $5 \cdot \frac{1}{3}$	$\frac{5}{3} \cdot 3$
Multiply.	5

	$5 \cdot (\frac{1}{3} \cdot 3)$
Multiply. $\frac{1}{3} \cdot 3$	$5 \cdot 1$
Multiply.	5

$$(5 \cdot \tfrac{1}{3}) \cdot 3 = 5 \cdot (\tfrac{1}{3} \cdot 3)$$

When multiplying three numbers, changing the grouping of the numbers gives the same result.

You probably know this, but the terminology may be new to you. These examples illustrate the **associative property**.

ASSOCIATIVE PROPERTY

of Addition	If a, b, c are real numbers, then $(a+b)+c=a+(b+c)$
of Multiplication	If a, b, c are real numbers, then $(a \cdot b) \cdot c=a \cdot(b \cdot c)$

When adding or multiplying, changing the *grouping* gives the same result.

Let's think again about multiplying $5 \cdot \frac{1}{3} \cdot 3$. We got the same result both ways, but which way was easier? Multiplying $\frac{1}{3}$ and 3 first, as shown above on the right side, eliminates the fraction in the first step. Using the associative property can make the math easier!

The **associative property** has to do with grouping. If we change how the numbers are grouped, the result will be the same. Notice it is the same three numbers in the same order—the only difference is the grouping.

We saw that subtraction and division were not commutative. They are not associative either.

When simplifying an expression, it is always a good idea to plan what the steps will be. In order to combine like terms in the next example, we will use the commutative property of addition to write the like terms together.

Example 1.10.1

Simplify: $18p+6q+15p+5q$.

Solution

	$18p+6q+15p+5q$
Use the commutative property of addition to re-order so that like terms are together.	$18p+15p+6q+5q$
Add like terms.	$33p+11q$

Try It 1.10.2

Simplify: $23r+14s+9r+15s$.

Answer

$32r+29s$

Try It 1.10.3

Simplify: $37m+21n+4m-15n$.

Answer

$41m+6n$

When we have to simplify algebraic **expressions**, we can often make the work easier by applying the commutative or associative property first, instead of automatically following the order of operations. When adding or subtracting fractions, combine those with a common denominator first.

Example 1.10.4

Simplify: $\left(\frac{5}{13}+\frac{3}{4}\right)+\frac{1}{4}$

Solution

	$(\frac{5}{13}+\frac{3}{4})+\frac{1}{4}$
Notice that the last 2 terms have a common denominator, so change the grouping.	$\frac{5}{13}+(\frac{3}{4}+\frac{1}{4})$
Add in parentheses first.	$\frac{5}{13}+(\frac{4}{4})$
Simplify the fraction.	$\frac{5}{13}+1$
Add.	$1\frac{5}{13}$
Convert to an improper fraction.	$\frac{18}{13}$

Try It 1.10.5

Simplify: $(\frac{7}{15}+\frac{5}{8})+\frac{3}{8}$

Answer

$1\frac{7}{15}$

Try It 1.10.6

Simplify: $(\frac{2}{9}+\frac{7}{12})+\frac{5}{12}$

Answer

$1\frac{2}{9}$

Example 1.10.7

Use the associative property to simplify $6(3x)$.

Solution

Use the associative property of multiplication, $(a\cdot b)\cdot c=a\cdot(b\cdot c)$, to change the grouping.

	$6(3x)$
Change the grouping.	$(6\cdot 3)x$
Multiply in the parentheses.	18

Notice that we can multiply $6\cdot 3$ but we could not multiply $3x$ without having a value for x.

Try It 1.10.8

Use the associative property to simplify $8(4x)$.

Answer

$32x$

Try It 1.10.9

Use the associative property to simplify $-9(7y)$.

Answer

$-63y$

Use the Identity and Inverse Properties of Addition and Multiplication

What happens when we add 0 to any number? Adding 0 doesn't change the value. For this reason, we call 0 the **additive identity**.

For example,

$$\begin{array}{ccc} 13+0 & -14+0 & 0+(-8) \\ 13 & -14 & -8 \end{array}$$

These examples illustrate the **Identity Property of Addition** that states that for any real number a, $a+0=a$ and $0+a=a$.

What happens when we multiply any number by one? Multiplying by 1 doesn't change the value. So we call 1 the **multiplicative identity**.

For example,

$$\begin{array}{ccc} 43\cdot 1 & -27\cdot 1 & 1\cdot\frac{3}{5} \\ 43 & -27 & \frac{3}{5} \end{array}$$

These examples illustrate the **Identity Property of Multiplication** that states that for any real number a, $a\cdot 1=a$ and $1\cdot a=a$.

We summarize the Identity Properties below.

IDENTITY PROPERTY

of addition For any real number a: $a+0=a \quad 0+a=a$
0 is the **additive identity**
of multiplication For any real number a: $a\cdot 1=a \quad 1\cdot a=a$
1 is the **multiplicative identity**

What number added to 5 gives the additive identity, 0?
$5+\_\_\_\_\_=0$ We know $5+(-5)=0$
What number added to -6 gives the additive identity, 0?
$-6+\_\_\_\_\_=0$ We know $-6+6=0$

Figure 1.10.1

Notice that in each case, the missing number was the opposite of the number!

We call $-a$. the **additive inverse** of a. *The opposite of a number is its additive inverse.* A number and its opposite add to zero, which is the additive identity. This leads to the **Inverse Property of Addition** that states for any real number $a, a+(-a)=0$. Remember, a number and its opposite add to zero.

What number multiplied by $\frac{2}{3}$ gives the multiplicative identity, 1? In other words, $\frac{2}{3}$ times what results in 1?

$\frac{2}{3}\cdot\_\_\_=1$ We know $\frac{2}{3}\cdot\frac{3}{2}=1$

Figure 1.10.2

What number multiplied by 2 gives the multiplicative identity, 1? In other words 2 times what results in 1?

$2\cdot\_\_\_=1$ We know $2\cdot\frac{1}{2}=1$

Figure 1.10.3

Notice that in each case, the missing number was the **reciprocal** of the number!

We call $\frac{1}{a}$ the **multiplicative inverse** of a. *The reciprocal of aa number is its multiplicative inverse.* A number and its reciprocal multiply to one, which is the multiplicative identity. This leads to the **Inverse Property of Multiplication** that states that for any real number $a, a \neq 0, a \cdot \frac{1}{a} = 1$.

We'll formally state the inverse properties here:

INVERSE PROPERTY

of addition	For any real number a, $-a$. is the **additive inverse** of a A number and its opposite add to zero.	$a + (-a) = 0$
of multiplication	For any real number $a, a \neq 0$ $\frac{1}{a}$. is the **multiplicative inverse** of a A number and its reciprocal multiply to one.	$a \cdot \frac{1}{a} = 1$

Example 1.10.10

Find the additive inverse of

1. $\frac{5}{8}$
2. 0.6
3. -8
4. $-\frac{4}{3}$

Solution

To find the additive inverse, we find the opposite.

1. The additive inverse of $\frac{5}{8}$ is the opposite of $\frac{5}{8}$. The additive inverse of $\frac{5}{8}$ is $-\frac{5}{8}$
2. The additive inverse of 0.6 is the opposite of 0.6. The additive inverse of 0.6 is -0.6.
3. The additive inverse of -8 is the opposite of -8. We write the opposite of -8 as $-(-8)$, and then simplify it to 8. Therefore, the additive inverse of -8 is 8.
4. The additive inverse of $-\frac{4}{3}$ is the opposite of $-\frac{4}{3}$. We write this as $-(-\frac{4}{3})$, and then simplify to $\frac{4}{3}$. Thus, the additive inverse of $-\frac{4}{3}$ is $\frac{4}{3}$.

Try It 1.10.11

Find the additive inverse of

1. $\frac{7}{9}$
2. 1.2
3. -14
4. $-\frac{9}{4}$

Answer

1. $-\frac{7}{9}$
2. -1.2
3. 14
4. $\frac{9}{4}$

Try It 1.10.12

Find the additive inverse of

1. $\frac{7}{13}$
2. 8.4
3. -46
4. $-\frac{5}{2}$

Answer

1. $-\frac{7}{13}$
2. -8.4
3. 46
4. $\frac{5}{2}$

Example 1.10.13

Find the multiplicative inverse of

1. 9
2. $-\frac{1}{9}$
3. 0.9

Solution

To find the multiplicative inverse, we find the reciprocal.

1. The multiplicative inverse of 9 is the reciprocal of 9, which is $\frac{1}{9}$. Therefore, the multiplicative inverse of 9 is $\frac{1}{9}$.
2. The multiplicative inverse of $-\frac{1}{9}$ is the reciprocal of $-\frac{1}{9}$, which is -9. Thus, the multiplicative inverse of $-\frac{1}{9}$ is -9.
3. To find the multiplicative inverse of 0.9, we first convert 0.9 to a fraction, $\frac{9}{10}$. Then we find the reciprocal of the fraction. The reciprocal of $\frac{9}{10}$ is $\frac{10}{9}$. So the multiplicative inverse of 0.9 is $\frac{10}{9}$.

Try It 1.10.14

Find the multiplicative inverse of

1. 4
2. $-\frac{1}{7}$
3. 0.3

Answer

1. $\frac{1}{4}$
2. -7
3. $\frac{10}{3}$

Try It 1.10.15

Find the multiplicative inverse of

1. 18
2. $-\frac{4}{5}$
3. 0.6

Answer

1. $\frac{1}{18}$

2. $-\frac{5}{4}$
3. $\frac{5}{3}$

Use the Properties of Zero

The identity property of addition says that when we add 0 to any number, the result is that same number. What happens when we multiply a number by 0? Multiplying by 0 makes the **product** equal zero.

MULTIPLICATION BY ZERO

For any real number a.

$$a \cdot 0 = 0 \quad 0 \cdot a = 0$$

The product of any real number and 0 is 0.

What about **division** involving zero? What is $0 \div 3$? Think about a real example: If there are no cookies in the cookie jar and 3 people are to share them, how many cookies does each person get? There are no cookies to share, so each person gets 0 cookies. So,

$$0 \div 3 = 0$$

We can check division with the related multiplication fact.

$$12 \div 6 = 2 \text{ because } 2 \cdot 6 = 12$$

So we know $0 \div 3 = 0$ because $0 \cdot 3 = 0$.

DIVISION OF ZERO

For any real number a, except 0, $\frac{0}{a} = 0$ and $0 \div a = 0$.

Zero divided by any real number except zero is zero.

Now think about dividing *by* zero. What is the result of dividing 4 by 0? Think about the related multiplication fact: $4 \div 0 = ?$ means $? \cdot 0 = 4$. Is there a number that multiplied by 0 gives 4? Since any real number multiplied by 0 gives 0, there is no real number that can be multiplied by 0 to obtain 4.

We conclude that there is no answer to $4 \div 0$ and so we say that division by 0 is undefined.

DIVISION BY ZERO

For any real number a, $\frac{a}{0}$ and $a \div 0$ are undefined.

Division by zero is undefined.

We summarize the properties of zero below.

PROPERTIES OF ZERO

Multiplication by Zero: For any real number a,

$$a \cdot 0 = 0 \quad 0 \cdot a = 0 \quad \text{The product of any number and 0 is 0}$$

Division of Zero, Division by Zero: For any real number $a, a \neq 0$

$\frac{0}{a} = 0$	Zero divided by any real number, except itself is zero.
$\frac{a}{0}$ is undefined	Division by zero is undefined.

Example 1.10.16

Simplify:

1. $-8 \cdot 0$
2. $\frac{0}{-2}$
3. $\frac{-32}{0}$

Solution

1. $-8 \cdot 0$

   The product of any real number and 0 is 0 — 0

2. $\frac{0}{-2}$

   Zero divided by any real number, except itself, is 0 — 0

3. $\frac{-32}{0}$

   Division by 0 is undefined. — undefined

Try It 1.10.17

Simplify:

1. $-14 \cdot 0$
2. $\frac{0}{-6}$
3. $\frac{-2}{0}$

Answer

1. 0
2. 0
3. undefined

Try It 1.10.18

Simplify:

1. $0(-17)$
2. $\frac{0}{-10}$
3. $\frac{-5}{0}$

Answer

1. 0
2. 0
3. undefined

We will now practice using the properties of identities, inverses, and zero to simplify expressions.

Example 1.10.19

Simplify:

1. $\frac{0}{n+5}$, where $n \neq -5$
2. $\frac{10-3p}{0}$ where $10-3p \neq 0$

Solution

1. $\frac{0}{n+5}$

   Zero divided by any real number except itself is 0. 0

2. $\frac{10-3p}{0}$

   Division by 0 is undefined undefined

Try It 1.10.26

Simplify:

1. $\frac{0}{m+7}$, where $m \neq -7$
2. $\frac{18-6c}{0}$, where $18-6c \neq 0$

Answer

1. 0
2. undefined

Try It 1.10.27

Simplify:

1. $\frac{0}{d-4}$, where $d \neq 4$
2. $\frac{15-4q}{0}$, where $15-4q \neq 0$

Answer

1. 0
2. undefined

Example 1.10.20

Simplify: $-84n+(-73n)+84n$.

Solution

	$-84n+(-73n)+84n$
Notice that the first and third terms are opposites; use the commutative property of addition to re-order the terms.	$-84n+84n+(-73n)$
Add left to right.	$0+(-73)$
Add.	$-73n$

Try It 1.10.21

Simplify: $-27a+(-48a)+27a$.

Answer

$-48a$

https://math.libretexts.org/@go/page/30467

Try It 1.10.22

Simplify: $39x + (-92x) + (-39x)$.

Answer

$-92x$

Now we will see how recognizing reciprocals is helpful. Before multiplying left to right, look for reciprocals—their product is 1.

Example 1.10.23

Simplify: $\frac{7}{15} \cdot \frac{8}{23} \cdot \frac{15}{7}$

Solution

	$\frac{7}{15} \cdot \frac{8}{23} \cdot \frac{15}{7}$
Notice that the first and third terms are reciprocals, so use the commutative property of multiplication to re-order the factors.	$\frac{7}{15} \cdot \frac{15}{7} \cdot \frac{8}{23}$
Multiply left to right.	$1 \cdot \frac{8}{23}$
Multiply.	$\frac{8}{23}$

Exercise 1.10.24

Simplify: $\frac{9}{16} \cdot \frac{5}{49} \cdot \frac{16}{9}$

Answer

$\frac{5}{49}$

Try It 1.10.25

Simplify: $\frac{6}{17} \cdot \frac{11}{25} \cdot \frac{17}{6}$

Answer

$\frac{11}{25}$

Example 1.10.28

Simplify: $\frac{3}{4} \cdot \frac{4}{3}(6x + 12)$

Solution

	$\frac{3}{4} \cdot \frac{4}{3}(6x+12)$
There is nothing to do in the parentheses, so multiply the two fractions first—notice, they are reciprocals.	$1(6x+12)$
Simplify by recognizing the multiplicative identity.	$6x+12$

? Try It 1.10.29

Simplify: $\frac{2}{5} \cdot \frac{5}{2}(20y+50)$

Answer

$20y+50$

? Try It 1.10.30

Simplify: $\frac{3}{8} \cdot \frac{8}{3}(12z+16)$

Answer

$12z+16$

Simplify Expressions Using the Distributive Property

Suppose that three friends are going to the movies. They each need $9.25—that's 9 dollars and 1 quarter—to pay for their tickets. How much money do they need all together?

You can think about the dollars separately from the quarters. They need 3 times $9 so $27, and 3 times 1 quarter, so 75 cents. In total, they need $27.75. If you think about doing the math in this way, you are using the **distributive property**.

DISTRIBUTIVE PROPERTY

If a, b, c are real numbers, then	$a(b+c) = ab+ac$
Also,	$(b+c)a = ba+ca$
	$a(b-c) = ab-ac$
	$(b-c)a = ba-ca$

Back to our friends at the movies, we could find the total amount of money they need like this:

$$3(9.25)$$
$$3(9+0.25)$$
$$3(9)+3(0.25)$$
$$27+0.75$$
$$27.75$$

In algebra, we use the **distributive property** to remove parentheses as we simplify expressions.

For example, if we are asked to simplify the expression $3(x+4)$, the order of operations says to work in the parentheses first. But we cannot add x and 4, since they are not like terms. So we use the distributive property, as shown in Exercise 1.10.31

Example 1.10.31

Simplify: $3(x+4)$.

Solution

	$3(x+4)$
Distribute.	$3\cdot x+3\cdot 4$
Multiply.	$3x+12$

Try It 1.10.32

Simplify: $4(x+2)$.

Answer

$4x+8$

Try It 1.10.33

Simplify: $6(x+7)$.

Answer

$6x+42$

Some students find it helpful to draw in arrows to remind them how to use the distributive property. Then the first step in Exercise 1.10.31would look like this:



Example 1.10.34

Simplify: $8(\frac{3}{8}x+\frac{1}{4})$.

Solution

	$8\left(\frac{3}{8}x+\frac{1}{4}\right)$
Distribute.	$8\cdot\frac{3}{8}x+8\cdot\frac{1}{4}$
Multiply.	$3x+2$

Try It 1.10.35

Simplify: $6(\frac{5}{6}y+\frac{1}{2})$.

Answer

$5y+3$

Try It 1.10.36

Simplify: $12(\frac{1}{3}n+\frac{3}{4})$.

Answer

$4n+9$

Using the distributive property as shown in Exercise 1.10.37 will be very useful when we solve money applications in later chapters.

Example 1.10.37

Simplify: $100(0.3+0.25q)$.

Solution

	$100(0.3+0.25q)$
Distribute.	$100(0.3)+100(0.25q)$
Multiply.	$30+25q$

Try It 1.10.38

Simplify: $100(0.7+0.15p)$.

Answer

$70+15p$

Try It 1.10.39

Simplify: $100(0.04+0.35d)$.

Answer

$4+35d$

When we distribute a negative number, we need to be extra careful to get the signs correct!

Example 1.10.40

Simplify: $-2(4y+1)$.

Solution

	$-2(4y+1)$
Distribute.	$-2 \cdot 4y+(-2) \cdot 1$
Multiply.	$-8y-2$

Try It 1.10.41

Simplify: $-3(6m+5)$.

Answer

$-18m-15)$

Try It 1.10.42

Simplify: $-6(8n+11)$.

Answer

$-48n-66)$

Example 1.10.43

Simplify: $-11(4-3a)$.

Solution

Distribute.	$-11(4-3a)$
Multiply.	$-11 \cdot 4-(-11) \cdot 3a$ $-44-(-33a)$
Simplify.	$-44+33a$

Notice that you could also write the result as $33a-44$. Do you know why?

Try It 1.10.44

Simplify: $-5(2-3a)$.

Answer

$10+15a$

Try It 1.10.45

Simplify: $-7(8-15y)$.

Answer

$-56+105y$

Exercise 1.10.46will show how to use the distributive property to find the opposite of an expression.

Example 1.10.46

Simplify: $-(y+5)$.

Solution

	$-(y+5)$
Multiplying by -1 results in the opposite.	$-1(y+5)$
Distribute.	$-1 \cdot y+(-1) \cdot 5$
Simplify.	$-y+(-5)$
	$-y-5$

Try It 1.10.47

Simplify: $-(z-11)$.

Answer

$-z+11$

Try It 1.10.48

Simplify: $-(x-4)$.

Answer

$-x+4$

There will be times when we'll need to use the **distributive property** as part of the order of operations. Start by looking at the parentheses. If the expression inside the parentheses cannot be simplified, the next step would be multiply using the distributive property, which removes the parentheses. The next two examples will illustrate this.

Example 1.10.49

Simplify: $8-2(x+3)$.

Be sure to follow the order of operations. Multiplication comes before subtraction, so we will distribute the 2 first and then subtract.

Solution

	$8-2(x+3)$
Distribute.	$8-2\cdot x-2\cdot 3$
Multiply.	$8-2x-6$
Combine like terms.	$-2x+2$

Try It 1.10.50

Simplify: $9-3(x+2)$.

Answer

$3-3x$

Try It 1.10.51

Simplify: $7x-5(x+4)$.

Answer

$2x-20$

Example 1.10.52

Simplify: $4(x-8)-(x+3)$.

Solution

	$4(x-8)-(x+3)$
Distribute.	$4x-32-x-3$
Combine like terms.	$3x-35$

Try It 1.10.53

Simplify: $6(x-9)-(x+12)$.

Answer

$5x-66$

Try It 1.10.54

Simplify: $8(x-1)-(x+5)$.

Answer

$7x-13$

All the properties of real numbers we have used in this chapter are summarized in Table 1.10.1.

Table 1.10.1

Commutative Property	
of addition If a,b are real numbers, then	$a+b=b+a$
of multiplication If a,b are real numbers, then	$a\cdot b=b\cdot a$
Associative Property	
of addition If a,b,c are real numbers, then	$(a+b)+c=a+(b+c)$
of multiplication If a,b,c are real numbers, then	$(a\cdot b)\cdot c=a\cdot(b\cdot c)$
Distributive Property	
If a,b,c are real numbers, then	$a(b+c)=ab+ac$
Identity Property	
of addition For any real number a: 0 is the **additive identity**	$a+0=a$ $0+a=a$
of multiplication For any real number a: **1** is the **multiplicative identity**	$a\cdot 1=a$ $1\cdot a=a$
Inverse Property	

of addition For any real number a, $-a$ is the **additive inverse** of a	$a+(-a)=0$
of multiplication For any real number $a, a \neq 0$ $\frac{1}{a}$ is the **multiplicative inverse** of a	$a \cdot \frac{1}{a}=1$
Properties of Zero	
For any real number a,	$a \cdot 0=0$
For any real number $a, a \neq 0$	$0 \cdot a=0$
	$\frac{0}{a}=0$
For any real number $a, a \neq 0$	$\frac{a}{0}$ is undefined

Key Concepts

- **Commutative Property of**
  - **Addition:** If a,b are real numbers, then $a+b=b+a$.
  - **Multiplication:** If a,b are real numbers, then $a \cdot b=b \cdot a$. When adding or multiplying, changing the *order* gives the same result.
- **Associative Property of**
  - **Addition:** If a,b,c are real numbers, then $(a+b)+c=a+(b+c)$.
  - **Multiplication:** If a,b,c are real numbers, then $(a \cdot b) \cdot c=a \cdot(b \cdot c)$. When adding or multiplying, changing the *grouping* gives the same result.
- **Distributive Property:** If a,b,c are real numbers, then
  - $a(b+c)=ab+ac$
  - $(b+c)a=ba+ca$
  - $a(b-c)=ab-ac$
  - $(b+c)a=ba+ca$
- **Identity Property**
  - **of Addition:** For any real number a: $a+0=a$ **0** is the **additive identity**
  - **of Multiplication:** For any real number a: $a \cdot 1=a \quad 1 \cdot a=a$ **1**1 is the **multiplicative identity**
- **Inverse Property**
  - **of Addition:** For any real number $a, a+(-a)=0$. A number and its *opposite* add to zero. $-a$ is the **additive inverse** of a.
  - **of Multiplication:** For any real number $a,(a \neq 0) a \cdot \frac{1}{a}=1$. A number and its *reciprocal* multiply to one. $\frac{1}{a}$ is the **multiplicative inverse** of a.
- **Properties of Zero**
  - For any real number a, $a \cdot 0=0 \quad 0 \cdot a=0$ – The product of any real number and 0 is 0.
  - $\frac{0}{a}=0$ for $a \neq 0$ – Zero divided by any real number except zero is zero.
  - $\frac{a}{0}$ is undefined – Division by zero is undefined.

This page titled 1.10: Properties of Real Numbers is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.10: Properties of Real Numbers** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.10E: Exercises

Practice Makes Perfect

Use the Commutative and Associative Properties

In the following exercises, use the associative property to simplify.

Exercise 1

$3(4x)$

Answer

$12x$

Exercise 2

$4(7m)$

Exercise 3

$(y+12)+28$

Answer

$y+40$

Exercise 4

$(n+17)+33$

In the following exercises, simplify.

Exercise 5

$\frac{1}{2}+\frac{7}{8}+\left(-\frac{1}{2}\right)$

Answer

$\frac{7}{8}$

Exercise 6

$\frac{2}{5}+\frac{5}{12}+\left(-\frac{2}{5}\right)$

Exercise 7

$\frac{3}{20}\cdot\frac{49}{11}\cdot\frac{20}{3}$

Answer

$\frac{49}{11}$

Exercise 8

$\frac{13}{18} \cdot \frac{25}{7} \cdot \frac{18}{13}$

Exercise 9

$-24 \cdot 7 \cdot \frac{3}{8}$

Answer

-63

Exercise 10

$-36 \cdot 11 \cdot \frac{4}{9}$

Exercise 11

$\left(\frac{5}{6} + \frac{8}{15}\right) + \frac{7}{15}$

Answer

$1\frac{5}{6}$

Exercise 12

$\left(\frac{11}{12} + \frac{4}{9}\right) + \frac{5}{9}$

Exercise 13

$17(0.25)(4)$

Answer

17

Exercise 14

$36(0.2)(5)$

Exercise 15

$[2.48(12)](0.5)$

Answer

14.88

Exercise 16

$[9.731(4)](0.75)$

Exercise 17

$7(4a)$

Answer

$28a$

Exercise 18

$9(8w)$

Exercise 19

$-15(5m)$

Answer

$-75m$

Exercise 20

$-23(2n)$

Exercise 21

$12(\frac{5}{6}p)$

Answer

$10p$

Exercise 22

$20(\frac{3}{5}q)$

Exercise 23

$43m+(-12n)+(-16m)+(-9n)$

Answer

$27m+(-21n)$

Exercise 24

$-22p+17q+(-35p)+(-27q)$

Exercise 25

$\frac{3}{8}g+\frac{1}{12}h+\frac{7}{8}g+\frac{5}{12}h$

Answer

$\frac{5}{4}g+\frac{1}{2}h$

Exercise 26

$\frac{5}{6}a+\frac{3}{10}b+\frac{1}{6}a+\frac{9}{10}b$

Exercise 27

$6.8p+9.14q+(-4.37p)+(-0.88q)$

Answer

$2.43p+8.26q$

Exercise 28

$9.6m+7.22n+(-2.19m)+(-0.65n)$

Use the Identity and Inverse Properties of Addition and Multiplication

In the following exercises, find the additive inverse of each number

Exercise 29

1. $\frac{2}{5}$
2. 4.3
3. -8
4. $-\frac{10}{3}$

Answer

1. $-\frac{2}{5}$
2. -4.3
3. 8
4. $\frac{10}{3}$

Exercise 30

1. $\frac{5}{9}$
2. 2.1
3. -3
4. $-\frac{9}{5}$

Exercise 31

1. $-\frac{7}{6}$
2. -0.075
3. 23
4. $\frac{1}{4}$

Answer

1. $\frac{7}{6}$
2. 0.075
3. -23
4. $-\frac{1}{4}$

Exercise 32

1. $-\frac{8}{3}$
2. -0.019
3. 52
4. $\frac{5}{6}$

In the following exercises, find the multiplicative inverse of each number.

Exercise 33

1. 6
2. $-\frac{3}{4}$
3. 0.7

Answer

1. $\frac{1}{6}$
2. $-\frac{4}{3}$
3. $\frac{10}{7}$

Exercise 34

1. 12
2. $-\frac{9}{2}$
3. 0.13

Exercise 35

1. $\frac{11}{12}$
2. -1.1
3. -4

Answer

1. $\frac{12}{11}$
2. $-\frac{10}{11}$
3. $-\frac{1}{4}$

Exercise 36

1. $\frac{17}{20}$
2. -1.5
3. -3

Use the Properties of Zero

In the following exercises, simplify.

Exercise 37

$\frac{0}{6}$

Answer

0

Exercise 38

$\frac{3}{0}$

Exercise 39

$0 \div \frac{11}{12}$

Answer

0

Exercise 40

$\frac{0}{3}$

Exercise 41

$\frac{6}{0}$

Answer

undefined

Exercise 42

$0 \cdot \frac{8}{15}$

Exercise 43

$(-3.14)(0)$

Answer

0

Exercise 44

$\frac{\frac{1}{10}}{0}$

Mixed Practice

In the following exercises, simplify.

Exercise 45

$19a + 44 - 19a$

Answer

44

Exercise 46

$27c + 16 - 27c$

Exercise 47

$10(0.1d)$

Answer

$1d$

Exercise 48

$100(0.01p)$

Exercise 49

$\frac{0}{u-4.99}$, where $u \neq 4.99$

Answer

0

Exercise 50

$\frac{0}{v-65.1}$, where $v \neq 65.1$

Exercise 51

$0 \div \left(x - \frac{1}{2}\right)$, where $x \neq \frac{1}{2}$

Answer

0

Exercise 52

$0 \div \left(y - \frac{1}{6}\right)$, where $y \neq \frac{1}{6}$

Exercise 53

$\frac{32-5a}{0}$, where $32 - 5a \neq 0$

Answer

undefined

Exercise 54

$\frac{28-9b}{0}$, where $28 - 9b \neq 0$

? Exercise 55

$\left(\frac{3}{4}+\frac{9}{10}m\right)\div 0$ where $\frac{3}{4}+\frac{9}{10}m\neq 0$

Answer

undefined

? Exercise 56

$\left(\frac{5}{16}n-\frac{3}{7}\right)\div 0$ where $\frac{5}{16}n-\frac{3}{7}\neq 0$

? Exercise 57

$15\cdot\frac{3}{5}(4d+10)$

Answer

$36d+90$

? Exercise 58

$18\cdot\frac{5}{6}(15h+24)$

Simplify Expressions Using the Distributive Property

In the following exercises, simplify using the distributive property.

? Exercise 59

$8(4y+9)$

Answer

$32y+72$

? Exercise 60

$9(3w+7)$

? Exercise 61

$6(c-13)$

Answer

$6c-78$

? Exercise 62

$7(y-13)$

? Exercise 63

$\frac{1}{4}(3q+12)$

Answer

$\frac{3}{4}q+3$

Exercise 64

$\frac{1}{5}(4m+20)$

Exercise 65

$9\left(\frac{5}{9}y-\frac{1}{3}\right)$

Answer

$5y-3$

Exercise 66

$10\left(\frac{3}{10}x-\frac{2}{5}\right)$

Exercise 67

$12\left(\frac{1}{4}+\frac{2}{3}r\right)$

Answer

$3+8r$

Exercise 68

$12\left(\frac{1}{6}+\frac{3}{4}s\right)$

Exercise 69

$r(s-18)$

Answer

$rs-18r$

Exercise 70

$u(v-10)$

Exercise 71

$(y+4)p$

Answer

$yp+4p$

Exercise 72

$(a+7)x$

Exercise 73

$-7(4p+1)$

Answer

$-28p-7$

Exercise 74

$-9(9a+4)$

Exercise 75

$-3(x-6)$

Answer

$-3x+18$

Exercise 76

$-4(q-7)$

Exercise 77

$-(3x-7)$

Answer

$-3x+7$

Exercise 78

$-(5p-4)$

Exercise 79

$16-3(y+8)$

Answer

$-3y-8$

Exercise 80

$18-4(x+2)$

Exercise 81

$4-11(3c-2)$

Answer

$-33c+26$

Exercise 82

$9-6(7n-5)$

Exercise 83

$22-(a+3)$

Answer

$-a+19$

Exercise 84

$8-(r-7)$

Exercise 85

$(5m-3)-(m+7)$

Answer

$4m-10$

Exercise 86

$(4y-1)-(y-2)$

Exercise 87

$5(2n+9)+12(n-3)$

Answer

$22n+9$

Exercise 88

$9(5u+8)+2(u-6)$

Exercise 89

$9(8x-3)-(-2)$

Answer

$72x-25$

Exercise 90

$4(6x-1)-(-8)$

Exercise 91

$14(c-1)-8(c-6)$

Answer

$6c+34$

Exercise 92

$11(n-7)-5(n-1)$

Exercise 93

$6(7y+8)-(30y-15)$

Answer

$12y+63$

Exercise 94

$7(3n+9)-(4n-13)$

Everyday Math

Exercise 95

Insurance copayment Carrie had to have 5 fillings done. Each filling cost $80. Her dental insurance required her to pay 20% of the cost as a copay. Calculate Carrie's copay:

1. First, by multiplying 0.20 by 80 to find her copay for each filling and then multiplying your answer by 5 to find her total copay for 5 fillings.
2. Next, by multiplying [5(0.20)](80)
3. Which of the properties of real numbers says that your answers to parts (a), where you multiplied 5[(0.20)(80)] and (b), where you multiplied [5(0.20)](80), should be equal?

Answer

1. $80
2. $80
3. answers will vary

Exercise 96

Cooking time Helen bought a 24-pound turkey for her family's Thanksgiving dinner and wants to know what time to put the turkey in to the oven. She wants to allow 20 minutes per pound cooking time. Calculate the length of time needed to roast the turkey:

1. First, by multiplying 24·20 to find the total number of minutes and then multiplying the answer by $\frac{1}{60}$ to convert minutes into hours.
2. Next, by multiplying $24(20\cdot\frac{1}{60})$.
3. Which of the properties of real numbers says that your answers to parts (a), where you multiplied $(24\cdot 20)\frac{1}{60}$, and (b), where you multiplied $24(20\cdot\frac{1}{60})$, should be equal?

Exercise 97

Buying by the case Trader Joe's grocery stores sold a bottle of wine they called "Two Buck Chuck" for $1.99. They sold a case of 12 bottles for $23.88. To find the cost of 12 bottles at $1.99, notice that 1.99 is $2-0.01$.

1. Multiply $12(1.99)$ by using the distributive property to multiply $12(2-0.01)$.
2. Was it a bargain to buy "Two Buck Chuck" by the case?

Answer

1. $23.88
2. no, the price is the same

Exercise 98

Multi-pack purchase Adele's shampoo sells for $3.99 per bottle at the grocery store. At the warehouse store, the same shampoo is sold as a 3 pack for $10.49. To find the cost of 3 bottles at $3.99, notice that 3.99 is $4-0.01$.

1. Multiply $3(3.99)$ by using the distributive property to multiply $3(4-0.01)$.
2. How much would Adele save by buying 3 bottles at the warehouse store instead of at the grocery store?

Writing Exercises

Exercise 99

In your own words, state the commutative property of addition.

Answer

$Answers may vary$

Exercise 100

What is the difference between the additive inverse and the multiplicative inverse of a number?

Exercise 101

Simplify $8\left(x-\frac{1}{4}\right)$ using the distributive property and explain each step.

Answer

$Answers may vary$

Exercise 102

Explain how you can multiply $4(\$5.97)$ without paper or calculator by thinking of $5.97 as $6-0.03$ and then using the distributive property.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
use the commutative and associative properties.			
use the identity and inverse properties of addition and multiplication.			
use the properties of zero.			
simplify expressions using the distributive property.			

ⓑ After reviewing this checklist, what will you do to become confident for all objectives?

This page titled 1.10E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.10E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.11: Systems of Measurement

Learning Objectives

By the end of this section, you will be able to:

- Make unit conversions in the US system
- Use mixed units of measurement in the US system
- Make unit conversions in the metric system
- Use mixed units of measurement in the metric system
- Convert between the US and the metric systems of measurement
- Convert between Fahrenheit and Celsius temperatures

Note

A more thorough introduction to the topics covered in this section can be found in the *Prealgebra* chapter, **The Properties of Real Numbers**.

Make Unit Conversions in the U.S. System

There are two systems of measurement commonly used around the world. Most countries use the metric system. The U.S. uses a different system of measurement, usually called the **U.S. system**. We will look at the U.S. system first.

The U.S. system of measurement uses units of inch, foot, yard, and mile to measure length and pound and ton to measure weight. For capacity, the units used are cup, pint, quart, and gallons. Both the U.S. system and the metric system measure time in seconds, minutes, and hours.

The equivalencies of measurements are shown in Table 1.11.1. The table also shows, in parentheses, the common abbreviations for each measurement.

Table 1.11.1

U.S. System of Measurement			
Length	1 foot (ft.)	=	12 inches (in.)
	1 yard (yd.)	=	3 feet (ft.)
	1 mile (mi.)	=	5280 feet (ft.)
Volume	3 teaspoons (t)	=	1 tablespoon (T)
	16 tablespoons (T)	=	1 cup (C)
	1 cup (C)	=	8 fluid ounces (fl.oz.)
	1 pint (pt.)	=	2 cups (C)
	1 quart (qt.)	=	2 pints (ft.)
	1 gallon (gal)	=	4 quarts (qt.)
Weight	1 pound (lb.)	=	16 ounces (oz.)
	1 ton	=	2000 pounds (lb.)
Time	1 minute (min)	=	60 seconds (sec)
	1 hour (hr)	=	60 minutes (min)
	1 day	=	24 hours (hr)
	1 week (wk)	=	7 days
	1 year (yr)	=	365 days

In many real-life applications, we need to convert between units of measurement, such as feet and yards, minutes and seconds, quarts and gallons, etc. We will use the identity property of multiplication to do these conversions. We'll restate the identity property of multiplication here for easy reference.

IDENTITY PROPERTY OF MULTIPLICATION

For any real number a: $a \cdot 1 = a \quad 1 \cdot a = a$

1 is the **multiplicative identity**

To use the identity property of multiplication, we write 1 in a form that will help us convert the units. For example, suppose we want to change inches to feet. We know that 1 foot is equal to 12 inches, so we will write 1 as the fraction $\frac{1 \text{ foot}}{12 \text{ inches}}$. When we multiply by this fraction we do not change the value, but just change the units.

But $\frac{12 \text{ inches}}{1 \text{ foot}}$ also equals 1. How do we decide whether to multiply by $\frac{1 \text{ foot}}{12 \text{ inches}}$ or $\frac{12 \text{ inches}}{1 \text{ foot}}$? We choose the fraction that will make the units we want to convert *from* divide out. Treat the unit words like **factors** and "divide out" common units like we do common factors. If we want to convert 6666 inches to feet, which multiplication will eliminate the inches?

$$66 \text{ inches} \cdot \frac{1 \text{ foot}}{12 \text{ inches}} \quad \text{or} \quad 66 \text{ inches} \cdot \frac{12 \text{ inches}}{1 \text{ foot}}$$

The first form works since $66 \text{ inches} \cdot \frac{1 \text{ foot}}{12 \text{ inches}}$.

Figure 1.11.1

The inches divide out and leave only feet. The second form does not have any units that will divide out and so will not help us.

Example 1.11.1

Mary Anne is 66 inches tall. Convert her height into feet.

Solution

Step 1. Multiply the measurement to be converted by 1; write 1 as a fraction relating the units given and the units needed.	Multiply 66 inches by 1, writing 1 as a fraction relating inches and feet. We need inches in the denominator so that the inches will divide out!	$66 \text{ inches} \cdot 1$ $66 \text{ inches} \cdot \frac{1 \text{ foot}}{12 \text{ inches}}$
Step 2. Multiply.	Think of 66 inches as $\frac{66 \text{ inches}}{1}$.	$\frac{66 \text{ inches} \cdot 1 \text{ foot}}{12 \text{ inches}}$
Step 3. Simplify the fraction.	Notice: inches divide out.	$\frac{66 \text{ inches} \cdot 1 \text{ foot}}{12 \text{ inches}}$ $\frac{66 \text{ feet}}{12}$
Step 4. Simplify.	Divide 66 by 12.	5.5 feet

Try It 1.11.2

Lexie is 30 inches tall. Convert her height to feet.

Answer

2.5 feet

Try It 1.11.3

Rene bought a hose that is 18 yards long. Convert the length to feet.

Answer

54 feet

MAKE UNIT CONVERSIONS.

1. Multiply the measurement to be converted by 1; write 1 as a fraction relating the units given and the units needed.
2. Multiply.
3. Simplify the fraction.
4. Simplify.

When we use the identity property of multiplication to convert units, we need to make sure the units we want to change from will divide out. Usually this means we want the conversion fraction to have those units in the denominator.

Example 1.11.4

Ndula, an elephant at the San Diego Safari Park, weighs almost 3.2 tons. Convert her weight to pounds.

Solution

We will convert 3.2 tons into pounds. We will use the identity property of multiplication, writing 1 as the fraction $\frac{2000 \text{ pounds}}{1 \text{ ton}}$.

	3.2 tons
Multiply the measurement to be converted, by 1.	$3.2 \text{ tons} \cdot 1$
Write 1 as a fraction relating tons and pounds.	$3.2 \text{ tons} \cdot \frac{2000 \text{ pounds}}{1 \text{ ton}}$
Simplify.	$\frac{3.2 \text{ tons} \cdot 2{,}000 \text{ pounds}}{1 \text{ ton}}$
Multiply.	6400 pounds
	Ndula weighs almost 6400 pounds.

Try It 1.11.5

Arnold's SUV weighs about 4.3 tons. Convert the weight to pounds.

Answer

8600 pounds

Try It 1.11.6

The Carnival *Destiny* cruise ship weighs 51000 tons. Convert the weight to pounds.

Answer

102000000 pounds

Sometimes, to convert from one unit to another, we may need to use several other units in between, so we will need to multiply several fractions.

Example 1.11.7

Juliet is going with her family to their summer home. She will be away from her boyfriend for 9 weeks. Convert the time to minutes.

Solution

To convert weeks into minutes we will convert weeks into days, days into hours, and then hours into minutes. To do this we will multiply by conversion factors of 1.

	9 weeks
Write 1 as $\frac{7\text{ days}}{1\text{ week}}$, and $\frac{60\text{ minutes}}{1\text{ hour}}$.	$\frac{9\text{ wk}}{1}\cdot\frac{7\text{ days}}{1\text{ wk}}\cdot\frac{24\text{ hr}}{1\text{ day}}\cdot\frac{60\text{ min}}{1\text{ hr}}$
Divide out the common units.	9 weeks Write 1 as $\frac{7\text{ days}}{1\text{ week}}$, $\frac{24\text{ hours}}{1\text{ day}}$, and $\frac{60\text{ minutes}}{1\text{ hour}}$. $\frac{9\text{ wk}}{1}\cdot\frac{7\text{ days}}{1\text{ wk}}\cdot\frac{24\text{ hr}}{1\text{ day}}\cdot\frac{60\text{ min}}{1\text{ hr}}$ Divide out the common units. $\frac{9\,\cancel{\text{wk}}}{1}\cdot\frac{7\,\cancel{\text{days}}}{1\,\cancel{\text{wk}}}\cdot\frac{24\,\cancel{\text{hr}}}{1\,\cancel{\text{day}}}\cdot\frac{60\text{ min}}{1\,\cancel{\text{hr}}}$ Multiply. $\frac{9\cdot7\cdot24\cdot60\text{ min}}{1\cdot1\cdot1\cdot1}$ Multiply. 90,720 min
Multiply.	$\frac{9\cdot7\cdot24\cdot60\text{ min}}{1\cdot1\cdot1\cdot1}$
Multiply.	90,720 min

Juliet and her boyfriend will be apart for 90,720 minutes (although it may seem like an eternity!).

Try It 1.11.8

The distance between the earth and the moon is about 250,000 miles. Convert this length to yards.

Answer

440,000,000 yards

Try It 1.11.9

The astronauts of Expedition 28 on the International Space Station spend 15 weeks in space. Convert the time to minutes.

Answer

151,200 minutes

Example 1.11.10

How many ounces are in 1 gallon?

Solution

We will convert gallons to ounces by multiplying by several conversion factors. Refer to Table 1.11.1.

	1 gallon
Multiply the measurement to be converted by 1.	$\frac{1\text{ gallon}}{1}\cdot\frac{4\text{ quarts}}{1\text{ gallon}}\cdot\frac{2\text{ pints}}{1\text{ quart}}\cdot\frac{2\text{ cups}}{1\text{ pint}}\cdot\frac{8\text{ ounces}}{1\text{ cup}}$
Use conversion factors to get to the right unit. Simplify.	1 gallon Multiply the measurement to be converted by 1. $\frac{1\text{ gallon}}{1}\cdot\frac{4\text{ quarts}}{1\text{ gallon}}\cdot\frac{2\text{ pints}}{1\text{ quart}}\cdot\frac{2\text{ cups}}{1\text{ pint}}\cdot\frac{8\text{ ounces}}{1\text{ cup}}$ Use conversion factors to get to the right unit. Simplify. $\frac{1\text{ gallon}}{1}\cdot\frac{4\,\cancel{\text{quarts}}}{1\,\cancel{\text{gallon}}}\cdot\frac{2\,\cancel{\text{pints}}}{1\,\cancel{\text{quart}}}\cdot\frac{2\,\cancel{\text{cups}}}{1\,\cancel{\text{pint}}}\cdot\frac{8\text{ ounces}}{1\,\cancel{\text{cup}}}$ Multiply. $\frac{1\cdot4\cdot2\cdot2\cdot8\text{ ounces}}{1\cdot1\cdot1\cdot1\cdot1}$ Simplify. 128 ounces There are 128 ounces in a gallon.
Multiply.	$\frac{1\cdot4\cdot2\cdot2\cdot8\text{ ounces}}{1\cdot1\cdot1\cdot1\cdot1}$
Simplify.	128 ounces

Try It 1.11.11

How many cups are in 1 gallon?

Answer

16 cups

Try It 1.11.12

How many teaspoons are in 1 cup?

Answer

48 teaspoons

Use Mixed Units of Measurement in the U.S. System

We often use mixed units of measurement in everyday situations. Suppose Joe is 5 feet 10 inches tall, stays at work for 7 hours and 45 minutes, and then eats a 1 pound 2 ounce steak for dinner—all these measurements have mixed units.

Performing arithmetic operations on measurements with mixed units of measures requires care. Be sure to add or subtract like units!

Example 1.11.13

Seymour bought three steaks for a barbecue. Their weights were 14 ounces, 1 pound 2 ounces and 1 pound 6 ounces. How many total pounds of steak did he buy?

Solution

We will add the weights of the steaks to find the total weight of the steaks.

Add the ounces. Then add the pounds.	14 ounces 1 pound 2 ounces + 1 pound 6 ounces 2 pounds 22 ounces
Convert 22 ounces to pounds and ounces.	2 pounds + 1 pound, 6 ounces 3 pounds, 6 ounces
Add the pounds.	Seymour bought 3 pounds 6 ounces of steak.

Add the ounces. Then add the pounds.	
Convert 22 ounces to pounds and ounces.	2 pounds + 1 pound, 6 ounces
Add the pounds.	3 pounds, 6 ounces
	Seymour bought 3 pounds 6 ounces of steak.

Try It 1.11.14

Laura gave birth to triplets weighing 3 pounds 3 ounces, 3 pounds 3 ounces, and 2 pounds 9 ounces. What was the total birth weight of the three babies?

Answer

8 lbs. 15 oz

Try It 1.11.15

Stan cut two pieces of crown molding for his family room that were 8 feet 7 inches and 12 feet 11 inches. What was the total length of the molding?

Answer

21 ft. 6 in.

Example 1.11.16

Anthony bought four planks of wood that were each 6 feet 4 inches long. What is the total length of the wood he purchased?

Solution

We will multiply the length of one plank to find the total length.

Multiply the inches and then the feet.	6 feet 4 inches × 4 24 feet 16 inches
Convert the 16 inches to feet. Add the feet.	24 feet + 1 foot 4 inches 25 feet 4 inches
	Anthony bought 25 feet and 4 inches of wood.

Try It 1.11.17

Henri wants to triple his spaghetti sauce recipe that uses 1 pound 8 ounces of ground turkey. How many pounds of ground turkey will he need?

Answer

4 lbs. 8 oz.

Try It 1.11.18

Joellen wants to double a solution of 5 gallons 3 quarts. How many gallons of solution will she have in all?

Answer

11 gallons 2 qt.

Make Unit Conversions in the Metric System

In the **metric system**, units are related by powers of 10. The roots words of their names reflect this relation. For example, the basic unit for measuring length is a meter. One kilometer is 1,000 meters; the prefix *kilo* means *thousand*. One centimeter is $\frac{1}{100}$ of a meter, just like one cent is $\frac{1}{100}$ of one dollar.

The equivalencies of measurements in the metric system are shown in Table 1.11.2 The common abbreviations for each measurement are given in parentheses.

Table 1.11.2

Metric System of Measurement		
Length	Mass	Capacity

Metric System of Measurement		
Length	Mass	Capacity
1 kilometer (km) = 1,000 m	1 kilogram (kg) = 1,000 g	1 kiloliter (kL) = 1,000 L
1 hectometer (hm) = 100 m	1 hectogram (hg) = 100 g	1 hectoliter (hL) = 100 L
1 dekameter (dam) = 10 m	1 dekagram (dag) = 10 g	1 dekaliter (daL) = 10 L
1 meter (m) = 1 m	1 gram (g) = 1 g	1 liter (L) = 1 L
1 decimeter (dm) = 0.1 m	1 decigram (dg) = 0.1 g	1 deciliter (dL) = 0.1 L
1 centimeter (cm) = 0.01 m	1 centigram (cg) = 0.01 g	1 centiliter (cL) = 0.01 L
1 millimeter (mm) = 0.001 m	1 milligram (mg) = 0.001 g	1 milliliter (mL) = 0.001 L
1 meter = 100 centimeters	1 gram = 100 centigrams	1 liter = 100 centiliters
1 meter = 1,000 millimeters	1 gram = 1,000 milligrams	1 liter = 1,000 milliliters

To make conversions in the metric system, we will use the same technique we did in the US system. Using the identity property of multiplication, we will multiply by a conversion factor of one to get to the correct units.

Have you ever run a 5K or 10K race? The length of those races are measured in kilometers. The metric system is commonly used in the United States when talking about the length of a race.

Example 1.11.19

Nick ran a 10K race. How many meters did he run?

Solution

We will convert kilometers to meters using the identity property of multiplication.

	10 kilometers
Multiply the measurement to be converted by 1.	$10 \text{ kilometers} \cdot 1$
Write 1 as a fraction relating kilometers and meters.	$10 \text{ kilometers} \cdot \frac{1{,}000 \text{ meters}}{1 \text{ kilometers}}$
Simplify.	$\frac{10 \text{ kilometers} \cdot 1{,}000 \text{ m}}{1 \text{ kilometers}}$
Multiply.	10,000 meters
	Nick ran 10,000 meters.

Try It 1.11.20

Sandy completed her first 5K race! How many meters did she run?

Answer

5,000 meters

Try It 1.11.21

Herman bought a rug 2.5 meters in length. How many centimeters is the length?

Answer

250 centimeters

Example 1.11.22

Eleanor's newborn baby weighed 3,200 grams. How many kilograms did the baby weigh?

Solution

We will convert grams into kilograms.

	$3{,}200\ \text{grams}$
Multiply the measurement to be converted by 1.	$3{,}200\ \text{grams} \cdot 1$
Write 1 as a function relating kilograms and grams.	$3{,}200\ \text{grams} \cdot \frac{1\ \text{kg}}{1{,}000\ \text{grams}}$
Simplify.	$3{,}200\ \text{grams} \cdot \frac{1\ \text{kg}}{1{,}000\ \text{grams}}$
Multiply.	$\frac{3{,}200\ \text{kilograms}}{1{,}000}$
Divide.	3.2 kilograms The baby weighed 3.2 kilograms.

Try It 1.11.23

Kari's newborn baby weighed 2,800 grams. How many kilograms did the baby weigh?

Answer

2.8 kilograms

Try It 1.11.24

Anderson received a package that was marked 4,500 grams. How many kilograms did this package weigh?

Answer

4.5 kilograms

As you become familiar with the metric system you may see a pattern. Since the system is based on multiples of ten, the calculations involve multiplying by multiples of ten. We have learned how to simplify these calculations by just moving the decimal.

To multiply by 10, 100, or 1,000, we move the decimal to the right one, two, or three places, respectively. To multiply by 0.1, 0.01, or 0.001, we move the decimal to the left one, two, or three places, respectively.

We can apply this pattern when we make measurement conversions in the metric system. In Exercise 1.11.25 we changed 3,200 grams to kilograms by multiplying by $\frac{1}{1000}$ (or 0.001). This is the same as moving the decimal three places to the left.

$3{,}200 \cdot \frac{1}{1{,}000}$ $\qquad$ $3{,}200.$

3.2 $\qquad$ 3.2

Figure 1.11.2

Example 1.11.25

Convert

1. 350 L to kiloliters
2. 4.1 L to milliliters.

Solution

1. We will convert liters to kiloliters. In Table 1.11.2, we see that 1 kiloliter=1,000 liters.1kiloliter=1,000 liters.

	350 L
Multiply by 1, writing 1 as a fraction relating liters to kiloliters.	$350\ \text{L}\ \frac{1\ \text{kL}}{1000\text{L}}$
Simplify.	$350\ \cancel{\text{L}}\ \frac{1\ \text{kL}}{1000\ \cancel{\text{L}}}$
Multiply by 1, writing 1 as a fraction relating liters to kiloliters. Simplify. Move the decimal 3 units to the left. (350.)	350 L $350\ \cancel{\text{L}} \cdot \frac{1\ \text{kL}}{1{,}000\ \cancel{\text{L}}}$ 0.35 kL
	0.35 kL

2. We will convert liters to milliliters. From Table 1.11.2we see that 1 liter=1,000 milliliters.1 liter=1,000 milliliters.

	4.1 L
Multiply by 1, writing 1 as a fraction relating liters to milliliters.	$4.1\ \text{L} \cdot \frac{1{,}000\ \text{mL}}{1\ \text{L}}$
Simplify.	$4.1\ \cancel{\text{L}} \cdot \frac{1{,}000\ \text{mL}}{1\ \cancel{\text{L}}}$
Move the decimal 3 units to the right.	4.100 mL
	4,100 mL

Try It 1.11.26

Convert:

1. 725 L to kiloliters
2. 6.3 L to milliliters

Answer

1. 7,250 kiloliters
2. 6,300 milliliters

Try It 1.11.27

Convert:

1. 350 hL to liters
2. 4.1 L to centiliters

Answer

1. 35,000 liters
2. 410 centiliters

Use Mixed Units of Measurement in the Metric System

Performing arithmetic operations on measurements with mixed units of measures in the metric system requires the same care we used in the US system. But it may be easier because of the relation of the units to the powers of 10. Make sure to add or subtract like units.

Example 1.11.28

Ryland is 1.6 meters tall. His younger brother is 85 centimeters tall. How much taller is Ryland than his younger brother?

Solution

We can convert both measurements to either centimeters or meters. Since meters is the larger unit, we will subtract the lengths in meters. We convert 85 centimeters to meters by moving the decimal 2 places to the left.

$$\begin{array}{ll} \text{Write the 85 centimeters as meters.} & 1.60\text{m} \\ & -0.85\text{m} \\ & \hline 0.75\text{m} \end{array} \qquad (1.11.1)$$

Ryland is 0.75 m0.75 m taller than his brother.

Try It 1.11.29

Mariella is 1.58 meters tall. Her daughter is 75 centimeters tall. How much taller is Mariella than her daughter? Write the answer in centimeters.

Answer

83 centimeters

Try It 1.11.30

The fence around Hank's yard is 2 meters high. Hank is 96 centimeters tall. How much shorter than the fence is Hank? Write the answer in meters.

Answer

1.04 meters

Example 1.11.31

Dena's recipe for lentil soup calls for 150 milliliters of olive oil. Dena wants to triple the recipe. How many liters of olive oil will she need?

Solution

We will find the amount of olive oil in milliliters then convert to liters.

	Triple 150 mL
Translate to algebra.	$3 \cdot 150$ mL
Multiply.	450 mL
Convert to liters.	$450 \cdot \frac{0.001\text{ L}}{1\text{ ml}}$
Simplify.	0.45 L
	Dena needs 0.45 liters of olive oil.

Try It 1.11.32

A recipe for Alfredo sauce calls for 250 milliliters of milk. Renata is making pasta with Alfredo sauce for a big party and needs to multiply the recipe amounts by 8. How many liters of milk will she need?

Answer

2 liters

Try It 1.11.33

To make one pan of baklava, Dorothea needs 400 grams of filo pastry. If Dorothea plans to make 6 pans of baklava, how many kilograms of filo pastry will she need?

Answer

2.4 kilograms

Convert Between the U.S. and the Metric Systems of Measurement

Many measurements in the United States are made in metric units. Our soda may come in 2-liter bottles, our calcium may come in 500-mg capsules, and we may run a 5K race. To work easily in both systems, we need to be able to convert between the two systems.

Table 1.11.3 shows some of the most common conversions.

Table 1.11.3

Conversion Factors Between U.S. and Metric Systems		
Length	Mass	Capacity
1 in. = 2.54 cm		
1 ft. = 0.305 m	1 lb. = 0.45 kg	1 qt. = 0.95 L
1 yd. = 0.914 m	1 oz. = 28 g	1 fl. oz. = 30 ml
1 mi. = 1.61 km	1 kg = 2.2 lb	1 L = 1.06 lb
1 m = 3.28 ft		

Figure 1.11.3 shows how inches and centimeters are related on a ruler.




Figure 1.11.3: This ruler shows inches and centimeters.

Figure 1.11.4 shows the ounce and milliliter markings on a measuring cup.




Figure 1.11.4: This measuring cup shows ounces and cups.

Figure 1.11.5 shows how pounds and kilograms marked on a bathroom scale.



Figure 1.11.5: This scale shows pounds and kilograms.

We make conversions between the systems just as we do within the systems—by multiplying by unit conversion factors.

Example 1.11.34

Lee's water bottle holds 500 mL of water. How many ounces are in the bottle? Round to the nearest tenth of an ounce.

Solution

	500 mL
Multiplying by a unit conversion factor relating mL and ounces	$500 \text{ milliliters} \cdot \frac{1 \text{ ounce}}{30 \text{ milliliters}}$
Simplify.	$\frac{50 \text{ ounce}}{30}$
Divide.	16.7 ounces
	The water bottle has 16.7 ounces

Try It 1.11.35

How many quarts of soda are in a 2-L bottle?

Answer

2.12 quarts

Try It 1.11.36

How many liters are in 4 quarts of milk?

Answer

3.8 liters

Example 1.11.37

Soleil was on a road trip and saw a sign that said the next rest stop was in 100 kilometers. How many miles until the next rest stop?

Solution

	100 kilometers
Multiplying by a unit conversion factor relating km and mi.	$100 \text{ kilometers} \cdot \frac{1 \text{ mile}}{1.61 \text{ kilometers}}$
Simplify.	$\frac{100 \text{ miles}}{1.61}$
Divide.	62 miles
	Soleil will travel 62 miles.

Try It 1.11.38

The height of Mount Kilimanjaro is 5,895 meters. Convert the height to feet.

Answer

19,335.6 feet

Try It 1.11.39

The flight distance from New York City to London is 5,586 kilometers. Convert the distance to miles.

Answer

3,469.57 miles

Convert between Fahrenheit and Celsius Temperatures

Have you ever been in a foreign country and heard the weather forecast? If the forecast is for 22°C, what does that mean?

The U.S. and metric systems use different scales to measure temperature. The U.S. system uses degrees Fahrenheit, written °F. The metric system uses degrees Celsius, written °C. Figure 1.11.6 shows the relationship between the two systems.




Figure 1.11.6: The diagram shows normal body temperature, along with the freezing and boiling temperatures of water in degrees Fahrenheit and degrees Celsius.

TEMPERATURE CONVERSION

To convert from Fahrenheit temperature, F, to Celsius temperature, C, use the formula

$$C = \frac{5}{9}(F - 32) \tag{1.11.2}$$

To convert from Celsius temperature, C, to Fahrenheit temperature, F, use the formula

$$F = \frac{9}{5}C + 32 \tag{1.11.3}$$

Example 1.11.40

Convert 50° Fahrenheit into degrees Celsius.

Solution

We will substitute 50°F into the formula to find C.

	$C = \frac{5}{9}(F - 32)$
Substitute 50 for F.	$C = \frac{5}{9}(50 - 32)$
Simplify in parentheses.	$C = \frac{5}{9}(18)$
Multiply.	$C = 10$
	So we found that 50°F is equivalent to 10°C.

Try It 1.11.41

Convert the Fahrenheit temperature to degrees Celsius: 59° Fahrenheit.

Answer

15°C

Try It 1.11.42

Convert the Fahrenheit temperature to degrees Celsius: 41° Fahrenheit.

Answer

5°C

Example 1.11.43

While visiting Paris, Woody saw the temperature was 20° Celsius. Convert the temperature into degrees Fahrenheit.

Solution

We will substitute 20°C into the formula to find F.

	$F = \frac{9}{5}C + 32$
Substitute 20 for C.	$F = \frac{9}{5}(20) + 32$
Multiply.	$F = 36 + 32$
Add.	$F = 68$
	So we found that 20°C is equivalent to 68°F.

Try It 1.11.44

Convert the Celsius temperature to degrees Fahrenheit: the temperature in Helsinki, Finland, was 15° Celsius.

Answer

59°F

Try It 1.11.45

Convert the Celsius temperature to degrees Fahrenheit: the temperature in Sydney, Australia, was 10° Celsius.

Answer

50°F

Key Concepts

- **Metric System of Measurement**
  - **Length**

    1 kilometer (km) = 1,000 m

    1 hectometer (hm) = 100 m

    1 dekameter (dam) = 10 m

    1 meter (m) = 1 m

    1 decimeter (dm) = 0.1 m

    1 centimeter (cm) = 0.01 m

    1 millimeter (mm) = 0.001 m

    1 meter = 100 centimeters

    1 meter = 1,000 millimeters

- **Mass**

  1 kilogram (kg) = 1,000 g

  1 hectogram (hg) = 100 g

  1 dekagram (dag) = 10 g

  1 gram (g) = 1 g

  1 decigram (dg) = 0.1 g

  1 centigram (cg) = 0.01 g

  1 milligram (mg) = 0.001 g

  1 gram = 100 centigrams

  1 gram = 1,000 milligrams

- **Capacity**

  1 kiloliter (kL) = 1,000 L

1 hectoliter (hL) = 100 L

1 dekaliter (daL) = 10 L

1 liter (L) = 1 L

1 deciliter (dL) = 0.1 L

1 centiliter (cL) = 0.01 L

1 milliliter (mL) = 0.001 L

1 liter = 100 centiliters

1 liter = 1,000 milliliters

- **Temperature Conversion**
  - To convert from Fahrenheit temperature, F, to Celsius temperature, C, use the formula $C = \frac{5}{9}(F - 32)$
  - To convert from Celsius temperature, C, to Fahrenheit temperature, F, use the formula $F = \frac{9}{5}C + 32$

This page titled 1.11: Systems of Measurement is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **1.11: Systems of Measurement** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

1.11E: Exercises

Practice Makes Perfect

Make Unit Conversions in the U.S. System

In the following exercises, convert the units.

Exercise 1.11E.1

A park bench is 6 feet long. Convert the length to inches.

Answer

72 inches

Exercise 1.11E.2

A floor tile is 2 feet wide. Convert the width to inches.

Exercise 1.11E.3

A ribbon is 18 inches long. Convert the length to feet.

Answer

1.5 feet

Exercise 1.11E.4

Carson is 45 inches tall. Convert his height to feet.

Exercise 1.11E.5

A football field is 160 feet wide. Convert the width to yards.

Answer

$53\frac{1}{3}$ yards

Exercise 1.11E.6

On a baseball diamond, the distance from home plate to first base is 30 yards. Convert the distance to feet.

Exercise 1.11E.7

On a baseball diamond, the distance from home plate to first base is 30 yards. Convert the distance to feet.

Answer

7,920 feet

Exercise 1.11E.8

Denver, Colorado, is 5,183 feet above sea level. Convert the height to miles.

Exercise 1.11E. 9

A killer whale weighs 4.6 tons. Convert the weight to pounds.

Answer

9,200 pounds

Exercise 1.11E. 10

Blue whales can weigh as much as 150 tons. Convert the weight to pounds.

Exercise 1.11E. 11

An empty bus weighs 35,000 pounds. Convert the weight to tons.

Answer

$17\frac{1}{2}$ tons

Exercise 1.11E. 12

At take-off, an airplane weighs 220,000 pounds. Convert the weight to tons.

Exercise 1.11E. 13

Rocco waited $1\frac{1}{2}$ hours for his appointment. Convert the time to seconds.

Answer

5,400 s

Exercise 1.11E. 14

Misty's surgery lasted $2\frac{1}{4}$ hours. Convert the time to seconds.

Exercise 1.11E. 15

How many teaspoons are in a pint?

Answer

96 teaspoons

Exercise 1.11E. 16

How many tablespoons are in a gallon?

Exercise 1.11E. 17

JJ's cat, Posy, weighs 14 pounds. Convert her weight to ounces.

Answer

224 ounces

Exercise 1.11E.18

April's dog, Beans, weighs 8 pounds. Convert his weight to ounces.

Exercise 1.11E.19

Crista will serve 20 cups of juice at her son's party. Convert the volume to gallons.

Answer

$1\frac{1}{4}$ gallons

Exercise 1.11E.20

Lance needs 50 cups of water for the runners in a race. Convert the volume to gallons.

Exercise 1.11E.21

Jon is 6 feet 4 inches tall. Convert his height to inches.

Answer

26 in.

Exercise 1.11E.22

Faye is 4 feet 10 inches tall. Convert her height to inches.

Exercise 1.11E.23

The voyage of the *Mayflower* took 2 months and 5 days. Convert the time to days.

Answer

65 days

Exercise 1.11E.24

Lynn's cruise lasted 6 days and 18 hours. Convert the time to hours.

Exercise 1.11E.25

Baby Preston weighed 7 pounds 3 ounces at birth. Convert his weight to ounces.

Answer

115 ounces

Exercise 1.11E.26

Baby Audrey weighted 6 pounds 15 ounces at birth. Convert her weight to ounces.

Use Mixed Units of Measurement in the U.S. System

In the following exercises, solve.

Exercise 1.11E.27

Eli caught three fish. The weights of the fish were 2 pounds 4 ounces, 1 pound 11 ounces, and 4 pounds 14 ounces. What was the total weight of the three fish?

Answer

8 lbs. 13 oz.

Exercise 1.11E.28

Judy bought 1 pound 6 ounces of almonds, 2 pounds 3 ounces of walnuts, and 8 ounces of cashews. How many pounds of nuts did Judy buy?

Exercise 1.11E.29

One day Anya kept track of the number of minutes she spent driving. She recorded 45, 10, 8, 65, 20, and 35. How many hours did Anya spend driving?

Answer

3.05 hours

Exercise 1.11E.30

Last year Eric went on 6 business trips. The number of days of each was 5, 2, 8, 12, 6, and 3. How many weeks did Eric spend on business trips last year?

Exercise 1.11E.31

Renee attached a 6 feet 6 inch extension cord to her computer's 3 feet 8 inch power cord. What was the total length of the cords?

Answer

10 ft. 2 in.

Exercise 1.11E.32

Fawzi's SUV is 6 feet 4 inches tall. If he puts a 2 feet 10 inch box on top of his SUV, what is the total height of the SUV and the box?

Exercise 1.11E.33

Leilani wants to make 8 placemats. For each placemat she needs 18 inches of fabric. How many yards of fabric will she need for the 8 placemats?

Answer

4 yards

Exercise 1.11E.34

Mireille needs to cut 24 inches of ribbon for each of the 12 girls in her dance class. How many yards of ribbon will she need altogether?

Make Unit Conversions in the Metric System

In the following exercises, convert the units.

Exercise 1.11E.35

Ghalib ran 5 kilometers. Convert the length to meters.

Answer

5,000 meters

Exercise 1.11E.36

Kitaka hiked 8 kilometers. Convert the length to meters.

Exercise 1.11E.37

Estrella is 1.55 meters tall. Convert her height to centimeters.

Answer

155 centimeters

Exercise 1.11E.38

The width of the wading pool is 2.45 meters. Convert the width to centimeters.

Exercise 1.11E.39

Mount Whitney is 3,072 meters tall. Convert the height to kilometers.

Answer

3.072 kilometers

Exercise 1.11E.40

The depth of the Mariana Trench is 10,911 meters. Convert the depth to kilometers.

Exercise 1.11E.41

June’s multivitamin contains 1,500 milligrams of calcium. Convert this to grams.

Answer

1.5 grams

Exercise 1.11E.42

A typical ruby-throated hummingbird weights 3 grams. Convert this to milligrams.

Exercise 1.11E.43

One stick of butter contains 91.6 grams of fat. Convert this to milligrams.

Answer

91,600 milligrams

Exercise 1.11E.44

One serving of gourmet ice cream has 25 grams of fat. Convert this to milligrams.

Exercise 1.11E.45

The maximum mass of an airmail letter is 2 kilograms. Convert this to grams.

Answer

2,000 grams

Exercise 1.11E.46

Dimitri's daughter weighed 3.8 kilograms at birth. Convert this to grams.

Exercise 1.11E.47

A bottle of wine contained 750 milliliters. Convert this to liters.

Answer

0.75 liters

Exercise 1.11E.48

A bottle of medicine contained 300 milliliters. Convert this to liters.

Use Mixed Units of Measurement in the Metric System

In the following exercises, solve.

Exercise 1.11E.49

Matthias is 1.8 meters tall. His son is 89 centimeters tall. How much taller is Matthias than his son?

Answer

91 centimeters

Exercise 1.11E.50

Stavros is 1.6 meters tall. His sister is 95 centimeters tall. How much taller is Stavros than his sister?

Exercise 1.11E.51

A typical dove weighs 345 grams. A typical duck weighs 1.2 kilograms. What is the difference, in grams, of the weights of a duck and a dove?

Answer

855 grams

Exercise 1.11E.52

Concetta had a 2-kilogram bag of flour. She used 180 grams of flour to make biscotti. How many kilograms of flour are left in the bag?

Exercise 1.11E.53

Harry mailed 5 packages that weighed 420 grams each. What was the total weight of the packages in kilograms?

Answer

2.1 kilograms

Exercise 1.11E.54

One glass of orange juice provides 560 milligrams of potassium. Linda drinks one glass of orange juice every morning. How many grams of potassium does Linda get from her orange juice in 30 days?

Exercise 1.11E.55

Jonas drinks 200 milliliters of water 8 times a day. How many liters of water does Jonas drink in a day?

Answer

1.6 liters

Exercise 1.11E.56

One serving of whole grain sandwich bread provides 6 grams of protein. How many milligrams of protein are provided by 7 servings of whole grain sandwich bread?

Convert Between the U.S. and the Metric Systems of Measurement

In the following exercises, make the unit conversions. Round to the nearest tenth.

Exercise 1.11E.57

Bill is 75 inches tall. Convert his height to centimeters.

Answer

190.5 centimeters

Exercise 1.11E.58

Frankie is 42 inches tall. Convert his height to centimeters.

Exercise 1.11E.59

Marcus passed a football 24 yards. Convert the pass length to meters

Answer

21.9 meters

Exercise 1.11E.60

Connie bought 9 yards of fabric to make drapes. Convert the fabric length to meters.

Exercise 1.11E.61

Each American throws out an average of 1,650 pounds of garbage per year. Convert this weight to kilograms.

Answer

742.5 kilograms

Exercise 1.11E.62

An average American will throw away 90,000 pounds of trash over his or her lifetime. Convert this weight to kilograms.

Exercise 1.11E.63

A 5K run is 5 kilometers long. Convert this length to miles.

Answer

3.1 miles

Exercise 1.11E.64

Kathryn is 1.6 meters tall. Convert her height to feet.

Exercise 1.11E.65

Dawn's suitcase weighed 20 kilograms. Convert the weight to pounds.

Answer

44 pounds

Exercise 1.11E.66

Jackson's backpack weighed 15 kilograms. Convert the weight to pounds.

Exercise 1.11E.67

Ozzie put 14 gallons of gas in his truck. Convert the volume to liters.

Answer

53.2 liters

Exercise 1.11E.68

Bernard bought 8 gallons of paint. Convert the volume to liters.

Convert between Fahrenheit and Celsius Temperatures

In the following exercises, convert the Fahrenheit temperatures to degrees Celsius. Round to the nearest tenth.

Exercise 1.11E.69

86° Fahrenheit

Answer

30°C

Exercise 1.11E.70

77° Fahrenheit

Exercise 1.11E.71

104° Fahrenheit

Answer

40°C

Exercise 1.11E.72

14° Fahrenheit

Exercise 1.11E.73

72° Fahrenheit

Answer

22.2°C

Exercise 1.11E.74

4° Fahrenheit

Exercise 1.11E.75

0° Fahrenheit

Answer

−17.8°C

Exercise 1.11E.76

120° Fahrenheit

In the following exercises, convert the Celsius temperatures to degrees Fahrenheit. Round to the nearest tenth.

Exercise 1.11E.77

5° Celsius

Answer

41°F

Exercise 1.11E.78

25° Celsius

Exercise 1.11E.79

−10° Celsius

Answer

14°F

Exercise 1.11E.80

−15° Celsius

Exercise 1.11E.81

22° Celsius

Answer

71.6°F

Exercise 1.11E.82

8° Celsius

Exercise 1.11E.83

43° Celsius

Answer

109.4°F

Exercise 1.11E.84

16° Celsius

Everyday Math

Exercise 1.11E.85

Nutrition Julian drinks one can of soda every day. Each can of soda contains 40 grams of sugar. How many kilograms of sugar does Julian get from soda in 1 year?

Answer

14.6 kilograms

Exercise 1.11E.86

Reflectors The reflectors in each lane-marking stripe on a highway are spaced 16 yards apart. How many reflectors are needed for a one mile long lane-marking stripe?

Writing Exercises

Exercise 1.11E. 87

Some people think that 65° to 75° Fahrenheit is the ideal temperature range.

1. What is your ideal temperature range? Why do you think so?
2. Convert your ideal temperatures from Fahrenheit to Celsius.

Answer

Answers may vary.

Exercise 1.11E. 88

1. Did you grow up using the U.S. or the metric system of measurement?
2. Describe two examples in your life when you had to convert between the two systems of measurement.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
define U.S. units of measurement and convert from one unit to another.			
use U.S units of measurement.			
define metric units of measurement and convert from one unit to another.			
use metric units of measurement.			
convert between the U.S. and the metric system of measurement.			
convert between Fahrenheit and Celsius temperatures.			

ⓑ Overall, after looking at the checklist, do you think you are well-prepared for the next Chapter? Why or why not?

This page titled 1.11E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **1.11E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 1 Review Exercises

Chapter Review Exercises

Introduction to Whole Numbers

Use Place Value with Whole Number

In the following exercises find the place value of each digit.

Exercise 1

26,915

1. 1
2. 2
3. 9
4. 5
5. 6

Answer

1. tens
2. ten thousands
3. hundreds
4. ones
5. thousands

Exercise 2

359,417

1. 9
2. 3
3. 4
4. 7
5. 1

Exercise 3

58,129,304

1. 5
2. 0
3. 1
4. 8
5. 2

Answer

1. ten millions
2. tens
3. hundred thousands
4. millions
5. ten thousands

Exercise 4

9,430,286,157

1. 6
2. 4
3. 9
4. 0
5. 5

In the following exercises, name each number.

Exercise 5

6,104

Answer

six thousand, one hundred four

Exercise 6

493,068

Exercise 7

3,975,284

Answer

three million, nine hundred seventy-five thousand, two hundred eighty-four

Exercise 8

85,620,435

In the following exercises, write each number as a whole number using digits.

Exercise 9

three hundred fifteen

Answer

315

Exercise 10

sixty-five thousand, nine hundred twelve

Exercise 11

ninety million, four hundred twenty-five thousand, sixteen

Answer

90,425,016

Exercise 12

one billion, forty-three million, nine hundred twenty-two thousand, three hundred eleven

In the following exercises, round to the indicated place value.

Exercise 13

Round to the nearest ten.

1. 407
2. 8,564

Answer

1. 410
2. 8,560

Exercise 14

Round to the nearest hundred.

1. 25,846
2. 25,864

In the following exercises, round each number to the nearest 1. hundred 2. thousand 3. ten thousand.

Exercise 15

864,951

Answer

1. 865,000865,000
2. 865,000865,000
3. 860,000

Exercise 16

3,972,849

Identify Multiples and Factors

In the following exercises, use the divisibility tests to determine whether each number is divisible by 2, by 3, by 5, by 6, and by 10.

Exercise 17

168

Answer

by 2,3,6

Exercise 18

264

Exercise 19

375

Answer

by 3,5

Exercise 20

750

Exercise 21

1430

Answer

by 2,5,10

Exercise 22

1080

Find Prime Factorizations and Least Common Multiples

In the following exercises, find the prime factorization.

Exercise 23

420

Answer

$2 \cdot 2 \cdot 3 \cdot 5 \cdot 7$

Exercise 24

115

Exercise 25

225

Answer

$3 \cdot 3 \cdot 5 \cdot 5$

Exercise 26

2475

Exercise 27

1560

Answer

$2 \cdot 2 \cdot 2 \cdot 3 \cdot 5 \cdot 13$

Exercise 28

56

Exercise 29

72

Answer

$2 \cdot 2 \cdot 2 \cdot 3 \cdot 3$

Exercise 30

168

Exercise 31

252

Answer

$2 \cdot 2 \cdot 3 \cdot 3 \cdot 7$

Exercise 32

391

In the following exercises, find the least common multiple of the following numbers using the multiples method.

Exercise 33

6,15

Answer

30

Exercise 34

60, 75

In the following exercises, find the least common multiple of the following numbers using the prime factors method.

Exercise 35

24, 30

Answer

120

Exercise 36

70, 84

Use the Language of Algebra

Use Variables and Algebraic Symbols

In the following exercises, translate the following from algebra to English.

Exercise 37

$25-7$

Answer

25 minus 7, the difference of twenty-five and seven

Exercise 38

$5 \cdot 6$

Exercise 39

$45 \div 5$

Answer

45 divided by 5, the quotient of forty-five and five

Exercise 40

$x+8$

Exercise 41

$42 \geq 27$

Answer

forty-two is greater than or equal to twenty-seven

Exercise 42

$3n=24$

Exercise 43

$3 \leq 20 \div 4$

Answer

3 is less than or equal to 20 divided by 4, three is less than or equal to the quotient of twenty and four

Exercise 44

$a \neq 7 \cdot 4$

In the following exercises, determine if each is an expression or an equation.

Exercise 45

$6 \cdot 3 + 5$

Answer

expression

Exercise 46

y−8=32

Simplify Expressions Using the Order of Operations

In the following exercises, simplify each expression.

Exercise 47

3^5

Answer

243

Exercise 48

10^8

In the following exercises, simplify

Exercise 49

6+10/2+2

Answer

13

Exercise 50

9+12/3+4

Exercise 51

$20 \div (4 + 6) \cdot 5$

Answer

10

Exercise 52

$33 \div (3+8) \cdot 2$

Exercise 53

$4^2 + 5^2$

Answer

41

Exercise 54

$(4+5)^2$

Evaluate an Expression

In the following exercises, evaluate the following expressions.

Exercise 55

9x+7 when x=3

Answer

34

Exercise 56

5x−4 when x=6

Exercise 57

x^4 when $x = 3$

Answer

81

Exercise 58

3^x when $x = 3$

Exercise 59

$x^2 + 5x - 8$ when $x = 6$

Answer

58

Exercise 60

$2x + 4y - 5$ when
$x = 7, y = 8$

Simplify Expressions by Combining Like Terms

In the following exercises, identify the coefficient of each term.

Exercise 61

12n

Answer

12

Exercise 62

$9x^2$

In the following exercises, identify the like terms.

Exercise 63

$3n, n^2, 12, 12p^2, 3, 3n^2$

Answer

12 and 3, n^2 and $3n^2$

Exercise 64

$5, 18r^2, 9s, 9r, 5r^2, 5s$

In the following exercises, identify the terms in each expression.

Exercise 65

$11x^2 + 3x + 6$

Answer

$11x^2, 3x, 6$

Exercise 66

$22y^3 + y + 15$

In the following exercises, simplify the following expressions by combining like terms.

Exercise 67

17a+9a

Answer

26a

Exercise 68

18z+9z

Exercise 69

9x+3x+8

Answer

12x+8

Exercise 70

8a+5a+9

Exercise 71

7p+6+5p−4

Answer

12p+2

Exercise 72

8x+7+4x−5

Translate an English Phrase to an Algebraic Expression

In the following exercises, translate the following phrases into algebraic expressions.

Exercise 73

the sum of 8 and 12

Answer

8+12

Exercise 74

the sum of 9 and 1

Exercise 75

the difference of x and 4

Answer

x−4

Exercise 76

the difference of x and 3

Exercise 77

the product of 6 and y

Answer

6y

Exercise 78

the product of 9 and y

Exercise 79

Adele bought a skirt and a blouse. The skirt cost $15 more than the blouse. Let bb represent the cost of the blouse. Write an expression for the cost of the skirt.

Answer

b+15

Exercise 80

Marcella has 6 fewer boy cousins than girl cousins. Let g represent the number of girl cousins. Write an expression for the number of boy cousins.

Add and Subtract Integers

Use Negatives and Opposites of Integers

In the following exercises, order each of the following pairs of numbers, using < or >.

Exercise 81

1. 6___2
2. −7___4
3. −9___−1
4. 9___−3

Answer

1. >
2. <
3. <
4. >

Exercise 82

1. −5___1
2. −4___−9
3. 6___10
4. 3___−8

In the following exercises,, find the opposite of each number.

Exercise 83

1. −8
2. 1

Answer

1. 8
2. −1

Exercise 84

1. −2
2. 6

In the following exercises, simplify.

Exercise 85

−(−19)

Answer

19

Exercise 86

−(−53)

In the following exercises, simplify.

Exercise 87

−m when

1. m=3
2. m=−3

Answer

1. −3
2. 3

Exercise 88

−p when

1. p=6
2. p=−6

Simplify Expressions with Absolute Value

In the following exercises,, simplify.

Exercise 89

1. |7|
2. |−25|
3. |0|

Answer

1. 7
2. 25
3. 0

Exercise 90

1. |5|
2. |0|
3. |−19|

In the following exercises, fill in <, >, or = for each of the following pairs of numbers.

Exercise 91

1. −8___|−8|
2. −|−2|___−2

Answer

1. <
2. =

Exercise 92

1. |−3|___−|−3|
2. 4___−|−4|

In the following exercises, simplify.

Exercise 93

|8−4|

Answer

4

Exercise 94

|9−6|

Exercise 95

8(14−2|−2|)

Answer

80

Exercise 96

6(13−4|−2|)

In the following exercises, evaluate.

Exercise 97

1. |x| when x=−28

Answer

1. 28
2. 15

Exercise 98

1. |y| when y=−37
2. |−z| when z=−24

Add Integers

In the following exercises, simplify each expression.

Exercise 99

−200+65

Answer

−135

Exercise 100

−150+45

Exercise 101

2+(−8)+6

Answer

0

Exercise 102

4+(−9)+7

Exercise 103

140+(−75)+67

Answer

132

Exercise 104

−32+24+(−6)+10

Subtract Integers

In the following exercises, simplify.

Exercise 105

$9-3$

Answer

6

Exercise 106

$-5-(-1)$

Exercise 107

1. $15-6$
2. $15+(-6)$

Answer

1. 9
2. 9

Exercise 108

1. $12-9$
2. $12+(-9)$

Exercise 109

1. $8-(-9)$
2. $8+9$

Answer

1. 17
2. 17

Exercise 110

1. $4-(-4)$
2. $4+4$

In the following exercises, simplify each expression.

Exercise 111

$10-(-19)$

Answer

29

Exercise 112

$11-(-18)$

Exercise 113

$31-79$

Answer

-48

Exercise 114

$39-81$

Exercise 115

$-31-11$

Answer

-42

Exercise 116

$-32-18$

Exercise 117

$-15-(-28)+5$

Answer

18

Exercise 118

$71+(-10)-8$

Exercise 119

$-16-(-4+1)-7$

Answer

-20

Exercise 120

$-15-(-6+4)-3$

Multiply Integers

In the following exercises, multiply.

Exercise 121

$-5(7)$

Answer

−35

Exercise 122

$-8(6)$

Exercise 123

$-18(-2)$

Answer

36

Exercise 124

$-10(-6)$

Divide Integers

In the following exercises, divide.

Exercise 125

$-28 \div 7$

Answer

-4

Exercise 126

$56 \div (-7)$

Exercise 127

$-120 \div (-20)$

Answer

6

Exercise 128

$-200 \div 25$

Simplify Expressions with Integers

In the following exercises, simplify each expression.

Exercise 129

$-8(-2)-3(-9)$

Answer

43

Exercise 130

$-7(-4)-5(-3)$

Exercise 131

$(-5)^3$

Answer

-125

Exercise 132

$(-4)^3$

Exercise 133

$-4 \cdot 2 \cdot 11$

Answer

-88

Exercise 134

$-5 \cdot 3 \cdot 10$

Exercise 135

$-10(-4) \div (-8)$

Answer

-5

Exercise 136

$-8(-6) \div (-4)$

Exercise 137

$31-4(3-9)$

Answer

55

Exercise 138

$24-3(2-10)$

Evaluate Variable Expressions with Integers

In the following exercises, evaluate each expression.

Exercise 139

x+8 when

1. x=−26
2. x=−95

Answer

1. −18
2. −87

Exercise 140

y+9 when

1. y=−29
2. y=−84

Exercise 141

When b=−11, evaluate:

1. b+6
2. −b+6

Answer

1. −5
2. 17

Exercise 142

When c=−9, evaluate:

1. c+(−4)c+(−4)
2. −c+(−4)

Exercise 143

$p^2 - 5p + 2$ when
$p = -1$

Answer

8

Exercise 144

$q^2 - 2q + 9$ when $q = -2$

Exercise 145

$6x - 5y + 15$ when $x = 3$ and $y = -1$

Answer

38

Exercise 146

$3p-2q+9$ when $p=8$ and $q=-2$

Translate English Phrases to Algebraic Expressions

In the following exercises, translate to an algebraic expression and simplify if possible.

Exercise 147

the sum of −4 and −17, increased by 32

Answer

(−4+(−17))+32;11

Exercise 148

1. the difference of 15 and −7
2. subtract 15 from −7

Exercise 149

the quotient of −45 and −9

Answer

$\frac{-45}{-9}; 5$

Exercise 150

the product of −12 and the difference of c and d

Use Integers in Applications

In the following exercises, solve.

Exercise 151

Temperature The high temperature one day in Miami Beach, Florida, was 76°. That same day, the high temperature in Buffalo, New York was −8°. What was the difference between the temperature in Miami Beach and the temperature in Buffalo?

Answer

84 degrees

Exercise 152

Checking Account Adrianne has a balance of −$22 in her checking account. She deposits $301 to the account. What is the new balance?

Visualize Fractions

Find Equivalent Fractions

In the following exercises, find three fractions equivalent to the given fraction. Show your work, using figures or algebra.

Exercise 153

$\frac{1}{4}$

Answer

$\frac{2}{8}, \frac{3}{12}, \frac{4}{16}$ answers may vary

Exercise 154

$\frac{1}{3}$

Exercise 155

$\frac{5}{6}$

Answer

$\frac{10}{12}, \frac{15}{18}, \frac{20}{24}$ answers may vary

Exercise 156

$\frac{2}{7}$

Simplify Fractions

In the following exercises, simplify.

Exercise 157

$\frac{7}{21}$

Answer

$\frac{1}{3}$

Exercise 158

$\frac{8}{24}$

Exercise 159

$\frac{15}{20}$

Answer

$\frac{3}{4}$

Exercise 160

$\frac{12}{18}$

Exercise 161

$-\frac{168}{192}$

Answer

$-\frac{7}{8}$

Exercise 162

$-\frac{140}{224}$

Exercise 163

$\frac{11x}{11y}$

Answer

$\frac{x}{y}$

Exercise 164

$\frac{15a}{15b}$

Multiply Fractions

In the following exercises, multiply.

Exercise 165

$\frac{2}{5} \cdot \frac{1}{3}$

Answer

$\frac{2}{15}$

Exercise 166

$\frac{1}{2} \cdot \frac{3}{8}$

Exercise 167

$\frac{7}{12}\left(-\frac{8}{21}\right)$

Answer

$-\frac{2}{9}$

Exercise 168

$\frac{5}{12}\left(-\frac{8}{15}\right)$

Exercise 169

$-28p\left(-\frac{1}{4}\right)$

Answer

7p

Exercise 170

$-51q\left(-\frac{1}{3}\right)$

Exercise 172

$\frac{14}{5}(-15)$

Answer

-42

Exercise 173

$-1\left(-\frac{3}{8}\right)$

Divide Fractions

In the following exercises, divide

Exercise 174

$\frac{1}{2} \div \frac{1}{4}$

Answer

2

Exercise 175

$\frac{1}{2} \div \frac{1}{8}$

Exercise 176

$-\frac{4}{5} \div \frac{4}{7}$

Answer

$-\frac{7}{5}$

Exercise 177

$-\frac{3}{4} \div \frac{3}{5}$

Exercise 178

$\frac{5}{8} \div \frac{a}{10}$

Answer

$\frac{25}{4a}$

Exercise 179

$\frac{5}{6} \div \frac{c}{15}$

Exercise 180

$\frac{7p}{12} \div \frac{21p}{8}$

Answer

$\frac{2}{9}$

Exercise 181

$\frac{5q}{12} \div \frac{15q}{8}$

Exercise 182

$\frac{2}{5} \div (-10)$

Answer

$-\frac{1}{25}$

Exercise 183

$-18 \div -\left(\frac{9}{2}\right)$

In the following exercises, simplify.

Exercise 184

$\frac{\frac{2}{3}}{\frac{8}{9}}$

Answer

$\frac{3}{4}$

Exercise 185

$\frac{\frac{4}{5}}{\frac{8}{15}}$

https://math.libretexts.org/@go/page/30471

Exercise 186

$\frac{-\frac{9}{10}}{3}$

Answer

$-\frac{3}{10}$

Exercise 187

$\frac{2}{\frac{5}{8}}$

Exercise 188

$\frac{\frac{r}{5}}{\frac{s}{3}}$

Answer

$\frac{3r}{5s}$

Exercise 189

$\frac{-\frac{x}{6}}{-\frac{8}{9}}$

Simplify Expressions Written with a Fraction Bar

In the following exercises, simplify.

Exercise 190

$\frac{4+11}{8}$

Answer

$\frac{15}{8}$

Exercise 191

$\frac{9+3}{7}$

Exercise 192

$\frac{30}{7-12}$

Answer

-6

Exercise 193

$\frac{15}{4-9}$

Exercise 194

$\frac{22-14}{19-13}$

Answer

$\frac{4}{3}$

Exercise 195

$\frac{15+9}{18+12}$

Exercise 196

$\frac{5 \cdot 8}{-10}$

Answer

-4

Exercise 197

$\frac{3 \cdot 4}{-24}$

Exercise 198

$\frac{15 \cdot 5-5^2}{2 \cdot 10}$

Answer

$\frac{5}{2}$

Exercise 199

$\frac{12 \cdot 9-3^2}{3 \cdot 18}$

Exercise 200

$\frac{2+4(3)}{-3-2^2}$

Answer

-2

Exercise 201

$\frac{7+3(5)}{-2-3^2}$

Translate Phrases to Expressions with Fractions

In the following exercises, translate each English phrase into an algebraic expression.

Exercise 202

the quotient of c and the sum of d and 9.

Answer

$\frac{c}{d+9}$

Exercise 203

the quotient of the difference of h and k, and -5.

Add and Subtract Fractions

Add and Subtract Fractions with a Common Denominator

In the following exercises, add.

Exercise 204

$\frac{4}{9}+\frac{1}{9}$

Answer

$\frac{5}{9}$

Exercise 205

$\frac{2}{9}+\frac{5}{9}$

Exercise 206

$\frac{y}{3}+\frac{2}{3}$

Answer

$\frac{y+2}{3}$

Exercise 207

$\frac{7}{p}+\frac{9}{p}$

Exercise 208

$-\frac{1}{8}+\left(-\frac{3}{8}\right)$

Answer

$-\frac{1}{2}$

Exercise 209

$-\frac{1}{8}+\left(-\frac{5}{8}\right)$

In the following exercises, subtract.

Exercise 210

$\frac{4}{5} - \frac{1}{5}$

Answer

$\frac{3}{5}$

Exercise 211

$\frac{4}{5} - \frac{3}{5}$

Exercise 212

$\frac{y}{17} - \frac{9}{17}$

Answer

$\frac{y-9}{17}$

Exercise 213

$\frac{x}{19} - \frac{8}{19}$

Exercise 214

$-\frac{8}{d} - \frac{3}{d}$

Answer

$-\frac{11}{d}$

Exercise 215

$-\frac{7}{c} - \frac{7}{c}$

Add or Subtract Fractions with Different Denominators

In the following exercises, add or subtract.

Exercise 216

$\frac{1}{3} + \frac{1}{5}$

Answer

$\frac{8}{15}$

Exercise 217

$\frac{1}{4} + \frac{1}{5}$

Exercise 218

$\frac{1}{5} - \left(-\frac{1}{10}\right)$

Answer

$\frac{3}{10}$

Exercise 219

$\frac{1}{2} - \left(-\frac{1}{6}\right)$

Exercise 220

$\frac{2}{3} + \frac{3}{4}$

Answer

$\frac{17}{12}$

Exercise 221

$\frac{3}{4} + \frac{2}{5}$

Exercise 222

$\frac{11}{12} - \frac{3}{8}$

Answer

$\frac{13}{24}$

Exercise 223

$\frac{5}{8} - \frac{7}{12}$

Exercise 224

$-\frac{9}{16} - \left(-\frac{4}{5}\right)$

Answer

$\frac{19}{80}$

Exercise 225

$-\frac{7}{20} - \left(-\frac{5}{8}\right)$

Exercise 226

$1 + \frac{5}{6}$

Answer

$\frac{11}{6}$

Exercise 227

$1-\frac{5}{9}$

Use the Order of Operations to Simplify Complex Fractions

In the following exercises, simplify.

Exercise 228

$\frac{\left(\frac{1}{5}\right)^2}{2+3^2}$

Answer

$\frac{1}{275}$

Exercise 229

$\frac{\left(\frac{1}{3}\right)^2}{5+2^2}$

Exercise 230

$\frac{\frac{2}{3}+\frac{1}{2}}{\frac{3}{4}-\frac{2}{3}}$

Answer

14

Exercise 231

$\frac{\frac{3}{4}+\frac{1}{2}}{\frac{5}{6}-\frac{2}{3}}$

Evaluate Variable Expressions with Fractions

In the following exercises, evaluate.

Exercise 232

$x+\frac{1}{2}$ when

1. $x=-\frac{1}{8}$
2. $x=-\frac{1}{2}$

Answer

1. $\frac{3}{8}$
2. 0

Exercise 233

$x+\frac{2}{3}$ when

1. $x=-\frac{1}{6}$

2. $x = -\frac{5}{3}$

Exercise 234

$4p^2q$ when $p = -\frac{1}{2}$ and $q = \frac{5}{9}$

Answer

$\frac{5}{9}$

Exercise 235

$5m^2n$ when $m = -\frac{2}{5}$ and $n = \frac{1}{3}$

Exercise 236

$\frac{u+v}{w}$ when
$u = -4, v = -8, w = 2$

Answer

-6

Exercise 237

$\frac{m+n}{p}$ when
$m = -6, n = -2, p = 4$

Decimals

Name and Write Decimals

In the following exercises, write as a decimal.

Exercise 238

Eight and three hundredths

Answer

8.03

Exercise 239

Nine and seven hundredths

Exercise 240

One thousandth

Answer

0.001

Exercise 241

Nine thousandths

In the following exercises, name each decimal.

Exercise 242

7.8

Answer

seven and eight tenths

Exercise 243

5.01

Exercise 244

0.005

Answer

five thousandths

Exercise 245

0.381

Round Decimals

In the following exercises, round each number to the nearest

1. hundredth
2. tenth
3. whole number.

Exercise 246

5.7932

Answer

1. 5.79
2. 5.8
3. 6

Exercise 247

3.6284

Exercise 248

12.4768

Answer

1. 12.48
2. 12.5
3. 12

Exercise 249

25.8449

Add and Subtract Decimals

In the following exercises, add or subtract.

Exercise 250

18.37+9.36

Answer

27.73

Exercise 251

256.37−85.49

Exercise 252

15.35−20.88

Answer

−5.53

Exercise 253

37.5+12.23

Exercise 254

−4.2+(−9.3)

Answer

−13.5

Exercise 255

−8.6+(−8.6)

Exercise 256

100−64.2

Answer

35.8

Exercise 257

100−65.83

Exercise 258

2.51+40

Answer

42.51

Exercise 259

9.38+60

Multiply and Divide Decimals

In the following exercises, multiply.

Exercise 260

(0.3)(0.4)

Answer

0.12

Exercise 261

(0.6)(0.7)

Exercise 262

(8.52)(3.14)

Answer

26.7528

Exercise 263

(5.32)(4.86)

Exercise 264

(0.09)(24.78)

Answer

2.2302

Exercise 265

(0.04)(36.89)

In the following exercises, divide.

Exercise 266

$0.15 \div 5$

Answer

0.03

Exercise 267

$0.27 \div 3$

Exercise 268

$\$8.49 \div 12$

Answer

$0.71

Exercise 269

$\$16.99 \div 9$

Exercise 270

$12 \div 0.08$

Answer

150

Exercise 271

$5 \div 0.04$

Convert Decimals, Fractions, and Percents

In the following exercises, write each decimal as a fraction.

Exercise 272

0.08

Answer

$\frac{2}{25}$

Exercise 273

0.17

Exercise 274

0.425

Answer

$\frac{17}{40}$

Exercise 275

0.184

Exercise 276

1.75

Answer

$\frac{7}{4}$

Exercise 277

0.035

In the following exercises, convert each fraction to a decimal.

Exercise 278

$\frac{2}{5}$

Answer

0.4

Exercise 279

$\frac{4}{5}$

Exercise 280

$-\frac{3}{8}$

Answer

-0.375

Exercise 281

$-\frac{5}{8}$

Exercise 282

$\frac{5}{9}$

Answer

$0.\overline{5}$

Exercise 283

$\frac{2}{9}$

Exercise 284

$\frac{1}{2} + 6.5$

Answer

7

Exercise 285

$\frac{1}{4} + 10.75$

In the following exercises, convert each percent to a decimal.

Exercise 286

5%

Answer

0.05

Exercise 287

9%

Exercise 288

40%

Answer

0.4

Exercise 289

50%

Exercise 290

115%

Answer

1.15

Exercise 291

125%

In the following exercises, convert each decimal to a percent.

Exercise 292

0.18

Answer

18%

Exercise 293

0.15

Exercise 294

0.009

Answer

0.9%

Exercise 295

0.008

Exercise 296

1.5

Answer

150%

Exercise 297

2.2

The Real Numbers

Simplify Expressions with Square Roots

In the following exercises, simplify.

Exercise 298

$\sqrt{64}$

Answer

8

Exercise 299

$\sqrt{144}$

Exercise 300

$-\sqrt{25}$

Answer

-5

Exercise 301

$-\sqrt{81}$

Identify Integers, Rational Numbers, Irrational Numbers, and Real Numbers

In the following exercises, write as the ratio of two integers.

Exercise 302

1. 9
2. 8.47

Answer

1. $\frac{9}{1}$
2. $\frac{847}{100}$

Exercise 303

1. -15
2. 3.591

In the following exercises, list the

1. rational numbers,
2. irrational numbers.

Exercise 304

$0.84, 0.79132\ldots, 1.\overline{3}$

Answer

1. $0.84, 1.3$
2. $0.79132\ldots$

Exercise 305

$2.3\overline{8}, 0.572, 4.93814\ldots$

In the following exercises, identify whether each number is rational or irrational.

Exercise 306

1. $\sqrt{121}$
2. $\sqrt{48}$

Answer

1. rational

2. irrational

Exercise 307

1. $\sqrt{56}$
2. $\sqrt{16}$

In the following exercises, identify whether each number is a real number or not a real number.

Exercise 308

1. $\sqrt{-9}$
2. $-\sqrt{169}$

Answer

1. not a real number
2. real number

Exercise 309

1. $\sqrt{-64}$
2. $-\sqrt{81}$

In the following exercises, list the

1. whole numbers,
2. integers,
3. rational numbers,
4. irrational numbers,
5. real numbers for each set of numbers.

Exercise 310

$-4, 0, \frac{5}{6}, \sqrt{16}, \sqrt{18}, 5.2537\ldots$

Answer

1. $0, \sqrt{16}$
2. $-4, 0, \sqrt{16}$
3. $-4, 0, \frac{5}{6}, \sqrt{16}$
4. $\sqrt{18}, 5.2537\ldots$
5. $-4, 0, \frac{5}{6}, \sqrt{16}, \sqrt{18}, 5.2537\ldots$

Exercise 311

$-\sqrt{4}, 0.\overline{36}, \frac{13}{3}, 6.9152\ldots, \sqrt{48}, 10\frac{1}{2}$


Locate Fractions on the Number Line

In the following exercises, locate the numbers on a number line.

Exercise 312

$\frac{2}{3}, \frac{5}{4}, \frac{12}{5}$

Answer




Exercise 313

$\frac{1}{3}, \frac{7}{4}, \frac{13}{5}$

Exercise 314

$2\frac{1}{3}, -2\frac{1}{3}$

Answer



Exercise 315

$1\frac{3}{5}, -1\frac{3}{5}$

In the following exercises, order each of the following pairs of numbers, using < or >.

Exercise 316

$-1 \_\_\_ -\frac{1}{8}$

Answer

<

Exercise 317

$-3\frac{1}{4} \_\_\_ -4$

Exercise 318

$-\frac{7}{9} \_\_\_ \frac{4}{9}$

Answer

>

Exercise 319

$-2 \_\_\_ \frac{19}{8}$


Locate Decimals on the Number Line

In the following exercises, locate on the number line.

Exercise 320

0.3

Answer




Exercise 321

−0.2

Exercise 322

−2.5

Answer



Exercise 323

2.7

In the following exercises, order each of the following pairs of numbers, using < or >.

Exercise 324

0.9___0.6

Answer

>

Exercise 325

0.7___0.8

Exercise 326

−0.6___−0.59

Answer

>

Exercise 327

−0.27___−0.3

Properties of Real Numbers

Use the Commutative and Associative Properties

In the following exercises, use the Associative Property to simplify.

Exercise 328

$-12(4m)$

Answer

$-48m$

Exercise 329

$30\left(\frac{5}{6}q\right)$

Exercise 330

(a+16)+31

Answer

a+47

Exercise 331

(c+0.2)+0.7

In the following exercises, simplify.

Exercise 332

6y+37+(−6y)

Answer

37

Exercise 333

$\frac{1}{4}+\frac{11}{15}+\left(-\frac{1}{4}\right)$

Exercise 334

$\frac{14}{11}\cdot\frac{35}{9}\cdot\frac{14}{11}$

Answer

$\frac{35}{9}$

Exercise 335

$-18\cdot 15\cdot\frac{2}{9}$

Exercise 336

$\left(\frac{7}{12}+\frac{4}{5}\right)+\frac{1}{5}$

Answer

$1\frac{7}{12}$

Exercise 337

(3.98d+0.75d)+1.25d

Exercise 338

11x+8y+16x+15y

Answer

27x+23y

Exercise 339

52m+(−20n)+(−18m)+(−5n)

Use the Identity and Inverse Properties of Addition and Multiplication

In the following exercises, find the additive inverse of each number.

Exercise 340

1. $\frac{1}{3}$
2. 5.1
3. -14
4. $-\frac{8}{5}$

Answer

1. $-\frac{1}{3}$
2. -5.1
3. -14
4. $-\frac{8}{5}$

Exercise 341

1. $-\frac{7}{8}$
2. -0.03
3. 17
4. $\frac{12}{5}$

In the following exercises, find the multiplicative inverse of each number.

Exercise 342

1. 10
2. $-\frac{4}{9}$
3. 0.6

Answer

1. $\frac{1}{10}$
2. $-\frac{9}{4}$
3. $\frac{5}{3}$

Exercise 343

1. $-\frac{9}{2}$
2. -7
3. 2.1

Use the Properties of Zero

In the following exercises, simplify.

Exercise 344

$83 \cdot 0$

Answer

0

Exercise 345

$\frac{0}{9}$

Exercise 346

$\frac{5}{0}$

Answer

undefined

Exercise 347

$0 \div \frac{2}{3}$

In the following exercises, simplify.

Exercise 348

$43+39+(-43)$

Answer

39

Exercise 349

$(n+6.75)+0.25$

Exercise 350

$\frac{5}{13} \cdot 57 \cdot \frac{13}{5}$

Answer

57

Exercise 351

$\frac{1}{6} \cdot 17 \cdot 12$

Exercise 352

$\frac{2}{3} \cdot 28 \cdot \frac{3}{7}$

Answer

8

Exercise 353

$9(6x - 11) + 15$

Simplify Expressions Using the Distributive Property

In the following exercises, simplify using the Distributive Property.

Exercise 354

7(x+9)

Answer

7x+63

Exercise 355

9(u−4)

Exercise 356

−3(6m−1)

Answer

−18m+3

Exercise 357

−8(−7a−12)

Exercise 358

$\frac{1}{3}(15n - 6)$

Answer

5n−2

Exercise 359

$(y + 10) \cdot p$

Exercise 360

$(a-4)-(6a+9)$

Answer

$-5a-13$

Exercise 361

$4(x+3)-8(x-7)$

Systems of Measurement

1.1 Define U.S. Units of Measurement and Convert from One Unit to Another

In the following exercises, convert the units. Round to the nearest tenth.

Exercise 362

A floral arbor is 7 feet tall. Convert the height to inches.

Answer

84 inches

Exercise 363

A picture frame is 42 inches wide. Convert the width to feet.

Exercise 364

Kelly is 5 feet 4 inches tall. Convert her height to inches.

Answer

64 inches

Exercise 365

A playground is 45 feet wide. Convert the width to yards.

Exercise 366

The height of Mount Shasta is 14,179 feet. Convert the height to miles.

Answer

2.7 miles

Exercise 367

Shamu weights 4.5 tons. Convert the weight to pounds.

Exercise 368

The play lasted $1\frac{3}{4}$ hours. Convert the time to minutes.

Answer

105 minutes

Exercise 369

How many tablespoons are in a quart?

Exercise 370

Naomi's baby weighed 5 pounds 14 ounces at birth. Convert the weight to ounces.

Answer

94 ounces

Exercise 371

Trinh needs 30 cups of paint for her class art project. Convert the volume to gallons.

Use Mixed Units of Measurement in the U.S. System.

In the following exercises, solve.

Exercise 372

John caught 4 lobsters. The weights of the lobsters were 1 pound 9 ounces, 1 pound 12 ounces, 4 pounds 2 ounces, and 2 pounds 15 ounces. What was the total weight of the lobsters?

Answer

10 lbs. 6 oz.

Exercise 373

Every day last week Pedro recorded the number of minutes he spent reading. The number of minutes were 50, 25, 83, 45, 32, 60, 135. How many hours did Pedro spend reading?

Exercise 374

Fouad is 6 feet 2 inches tall. If he stands on a rung of a ladder 8 feet 10 inches high, how high off the ground is the top of Fouad's head?

Answer

15 feet

Exercise 375

Dalila wants to make throw pillow covers. Each cover takes 30 inches of fabric. How many yards of fabric does she need for 4 covers?

Make Unit Conversions in the Metric System

In the following exercises, convert the units.

Exercise 376

Donna is 1.7 meters tall. Convert her height to centimeters.

Answer

170 centimeters

Exercise 377

Mount Everest is 8,850 meters tall. Convert the height to kilometers.

Exercise 378

One cup of yogurt contains 488 milligrams of calcium. Convert this to grams.

Answer

0.488 grams

Exercise 379

One cup of yogurt contains 13 grams of protein. Convert this to milligrams.

Exercise 380

Sergio weighed 2.9 kilograms at birth. Convert this to grams.

Answer

2,900 grams

Exercise 381

A bottle of water contained 650 milliliters. Convert this to liters.

Use Mixed Units of Measurement in the Metric System

In the following exerices, solve.

Exercise 382

Minh is 2 meters tall. His daughter is 88 centimeters tall. How much taller is Minh than his daughter?

Answer

1.12 meter

Exercise 383

Selma had a 1 liter bottle of water. If she drank 145 milliliters, how much water was left in the bottle?

Exercise 384

One serving of cranberry juice contains 30 grams of sugar. How many kilograms of sugar are in 30 servings of cranberry juice?

Answer

0.9 kilograms

Exercise 385

One ounce of tofu provided 2 grams of protein. How many milligrams of protein are provided by 5 ounces of tofu?

Convert between the U.S. and the Metric Systems of Measurement

In the following exercises, make the unit conversions. Round to the nearest tenth.

Exercise 386

Majid is 69 inches tall. Convert his height to centimeters.

Answer

175.3 centimeters

Exercise 387

A college basketball court is 84 feet long. Convert this length to meters.

Exercise 388

Caroline walked 2.5 kilometers. Convert this length to miles.

Answer

1.6 miles

Exercise 389

Lucas weighs 78 kilograms. Convert his weight to pounds.

Exercise 390

Steve’s car holds 55 liters of gas. Convert this to gallons.

Answer

14.6 gallons

Exercise 391

A box of books weighs 25 pounds. Convert the weight to kilograms.

Convert between Fahrenheit and Celsius Temperatures

In the following exercises, convert the Fahrenheit temperatures to degrees Celsius. Round to the nearest tenth.

Exercise 392

95° Fahrenheit

Answer

35° C

Exercise 393

23° Fahrenheit

Exercise 394

20° Fahrenheit

Answer

–6.7° C

Exercise 395

64° Fahrenheit

In the following exercises, convert the Celsius temperatures to degrees Fahrenheit. Round to the nearest tenth.

Exercise 396

30° Celsius

Answer

86° F

Exercise 397

–5° Celsius

Exercise 398

–12° Celsius

Answer

10.4° F

Exercise 399

24° Celsius

Chapter Practice Test

Exercise 1

Write as a whole number using digits: two hundred five thousand, six hundred seventeen.

Answer

205,617

Exercise 2

Find the prime factorization of 504.

Exercise 3

Find the Least Common Multiple of 18 and 24.

Answer

72

Exercise 4

Combine like terms: 5n+8+2n−1.

In the following exercises, evaluate.

Exercise 5

$-|x|$ when $x = -2$

Answer

−2

Exercise 6

11−a when a=−3

Exercise 7

Translate to an algebraic expression and simplify: twenty less than negative 7.

Answer

−7−20;−27

Exercise 8

Monique has a balance of −$18 in her checking account. She deposits $152 to the account. What is the new balance?

Exercise 9

Round 677.1348 to the nearest hundredth.

Answer

677.13

Exercise 10

Convert $\frac{4}{5}$ to a decimal.

Exercise 11

Convert 1.85 to a percent.

Answer

185%

Exercise 12

Locate $\frac{2}{3}, -1.5,$ and $\frac{9}{4}$ on a number line.

In the following exercises, simplify each expression.

Exercise 13

$4+10(3+9)-5^2$

Answer

99

Exercise 14

$-85+42$

Exercise 15

$-19-25$

Answer

-44

Exercise 16

$(-2)^4$

Exercise 17

$-5(-9)\div 15$

Answer

3

Exercise 18

$\frac{3}{8}\cdot\frac{11}{12}$

Exercise 19

$\frac{4}{5} \div \frac{9}{20}$

Answer

$\frac{16}{9}$

Exercise 20

$\frac{12+3 \cdot 5}{15-6}$

Exercise 21

$\frac{m}{7} + \frac{10}{7}$

Answer

$\frac{m+10}{7}$

Exercise 22

$\frac{7}{12} - \frac{3}{8}$

Exercise 23

$-5.8 + (-4.7)$

Answer

−10.5

Exercise 24

100−64.25

Exercise 25

(0.07)(31.95)

Answer

2.2365

Exercise 26

$9 \div 0.05$

Exercise 27

$-14\left(\frac{5}{7}p\right)$

Answer

−10p

Exercise 28

$(u+8)-9$

Exercise 29

$6x+(-4y)+9x+8y$

Answer

$15x+4y$

Exercise 30

$\frac{0}{23}$

Exercise 31

$\frac{75}{0}$

Answer

undefined

Exercise 32

$-2(13q-5)$

Exercise 33

A movie lasted $1\frac{2}{3}$ hours. How many minutes did it last? (1 hour $=$ 60 minutes)

Answer

100 minutes

Exercise 34

Mike's SUV is 5 feet 11 inches tall. He wants to put a rooftop cargo bag on the the SUV. The cargo bag is 1 foot 6 inches tall. What will the total height be of the SUV with the cargo bag on the roof? *(1 foot = 12 inches)*

Exercise 35

Jennifer ran 2.8 miles. Convert this length to kilometers. *(1 mile = 1.61 kilometers)*

Answer

4.508 km

This page titled Chapter 1 Review Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **Chapter 1 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

CHAPTER OVERVIEW

2: Solving Linear Equations and Inequalities

2.1: Solve Equations Using the Subtraction and Addition Properties of Equality

2.1E: Exercises

2.2: Solve Equations using the Division and Multiplication Properties of Equality

2.2E: Exercises

2.3: Solve Equations with Variables and Constants on Both Sides

2.3E: Exercises

2.4: Use a General Strategy to Solve Linear Equations

2.4E: Exercises

2.5: Solve Equations with Fractions or Decimals

2.5E: Exercises

2.6: Solve Proportions and Similar Figure Applications

2.6E: Exercises

2.7: Solve Linear Inequalities

2.7E: Exercises for Section 2.7

2.7E2: Exercises

2.8: Solve Compound Inequalities

2.8E: Exercises for Section 2.8

Chapter 2 Review Exercises

This page titled 2: Solving Linear Equations and Inequalities is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

2.1: Solve Equations Using the Subtraction and Addition Properties of Equality

Learning Objectives

By the end of this section, you will be able to:

- Verify a solution of an equation
- Solve equations using the Subtraction and Addition Properties of Equality
- Solve equations that require simplification
- Translate to an equation and solve
- Translate and solve applications

Quiz

Before you get started, take this readiness quiz.

1. Evaluate $x+4$ when $x=-3$.
   If you missed this problem, review Exercise 1.5.25.
2. Evaluate $15-y$ when $y=-5$.
   If you missed this problem, review Exercise 1.5.31.
3. Simplify $4(4n+1)-15n$.
   If you missed this problem, review Exercise 1.10.49.
4. Translate into algebra "5 is less than x."
   If you missed this problem, review Exercise 1.3.43.

Verify a Solution of an Equation

Solving an equation is like discovering the answer to a puzzle. The purpose in solving an equation is to find the value or values of the variable that make each side of the equation the same – so that we end up with a true statement. Any value of the variable that makes the equation true is called a solution to the equation. It is the answer to the puzzle!

Definition: Solution of an Equation

A **solution of an equation** is a value of a variable that makes a true statement when substituted into the equation.

TO DETERMINE WHETHER A NUMBER IS A Solution TO AN EQUATION

1. Substitute the number in for the variable in the equation.
2. Simplify the expressions on both sides of the equation.
3. Determine whether the resulting equation is true (the left side is equal to the right side)
   - If it is true, the number is a solution.
   - If it is not true, the number is not a solution.

Example 2.1.1

Determine whether $x=\frac{3}{2}$ is a solution of $4x-2=2x+1$.

Solution

Since a solution to an equation is a value of the variable that makes the equation true, begin by substituting the value of the solution for the variable.

$$\begin{array}{l|l|l} 4x-2=2x+1 & & \\ 4\left(\frac{3}{2}\right)-2 & 2\left(\frac{3}{2}\right)+1 & \text{Substitute } \frac{3}{2} \text{ for } x. \\ =6-2 & =3+1 & \text{Multiply.} \\ =4 & =4 & \text{Subtract.} \end{array}$$

$$= \checkmark$$

Since $x = \frac{3}{2}$ results in both sides of the equation being equal (4 is in fact equal to 4), $\frac{3}{2}$ is a solution to the equation $4x - 2 = 2x + 1$.

Try It 2.1.2

Is $y = \frac{4}{3}$ a solution of $9y + 2 = 6y + 3$?

Answer

no

Try It 2.1.3

Is $y = \frac{7}{5}$ a solution of $5y + 3 = 10y - 4$?

Answer

yes

Solve Equations Using the Subtraction and Addition Properties of Equality

We are going to use a model to clarify the process of solving an equation. An envelope represents the variable – since its contents are unknown – and each counter represents one. We will set out one envelope and some counters on our workspace, as shown in Figure 2.1.1. Both sides of the workspace have the same number of counters, but some counters are "hidden" in the envelope. Can you tell how many counters are in the envelope?




Figure 2.1.1: The illustration shows a model of an equation with one variable. On the left side of the workspace is an unknown (envelope) and three counters, while on the right side of the workspace are eight counters.

What are you thinking? What steps are you taking in your mind to figure out how many counters are in the envelope?

Perhaps you are thinking: "I need to remove the 3 counters at the bottom left to get the envelope by itself. The 3 counters on the left can be matched with 3 on the right and so I can take them away from both sides. That leaves five on the right—so there must be 5 counters in the envelope." See Figure 2.1.2 for an illustration of this process.




Figure 2.1.2: The illustration shows a model for solving an equation with one variable. On both sides of the workspace remove three counters, leaving only the unknown (envelope) and five counters on the right side. The unknown is equal to five counters.

What algebraic equation would match this situation? In Figure 2.1.3 each side of the workspace represents an expression and the center line takes the place of the equal sign. We will call the contents of the envelope x.




Figure 2.1.3: The illustration shows a model for the equation $x + 3 = 8$.

Let's write algebraically the steps we took to discover how many counters were in the envelope:

Table 2.1.1

	$x + 3 = 8$
First, we took away three from each side.	$x + 3 = 8$ $-3 \quad -3$

Then we were left with five.	$x = 5$

Check:

Five in the envelope plus three more does equal eight!

$$5 + 3 = 8$$

Our model has given us an idea of what we need to do to solve one kind of equation. The goal is to isolate the variable by itself on one side of the equation. To solve equations such as these mathematically, we use the **Subtraction Property of Equality**.

SUBTRACTION PROPERTY OF EQUALITY

For any numbers a, b, and c,

$$\begin{aligned} &\text{If} \quad a = b \\ &\text{then} \quad a - c = b - c \end{aligned}$$

When you subtract the same quantity from both sides of an equation, you still have equality.

Note

Doing the Manipulative Mathematics activity "Subtraction Property of Equality" will help you develop a better understanding of how to solve equations by using the **Subtraction Property of Equality**.

Let's see how to use this property to solve an equation. Remember, the goal is to isolate the variable on one side of the equation. And we check our solutions by substituting the value into the equation to make sure we have a true statement.

Example 2.1.4

Solve: $y + 37 = -13$.

Solution

To get y by itself, we will undo the addition of 37 by using the Subtraction Property of Equality.

	$y + 37 = -13$
Subtract 37 from each side to 'undo' the addition.	$\begin{array}{rr} y + 37 = -13 \\ -37 \quad -37 \end{array}$
Simplify.	$y = -50$

To check, we substitute the value -50 in for the variable y on the left side of the equation and simplify. Note that since the variable is only present on the left side of the equation, we don't need to show both sides of the equation until we evaluate the left side to see if it gives us the right number.

Check:

$$\begin{aligned} y + 37 &= -13 \\ (-50) + 37 &= -13 \checkmark \end{aligned}$$

Since y=−50 makes y+37=−13 a true statement, we have the solution to this equation.

Note that if we had incorrectly solved for y, and we were checking whether $y = 24$ is a solution, we would have done the following work to check it in the equation.

Check:

$$\begin{aligned} y + 37 &= -13 \\ (24) + 37 &= 61 \neq -13 \end{aligned}$$

Since this value of y does not make this equation true (it does **not** give us -13 as the result), we know that $y = 24$ is **not** a solution.

Try It 2.1.5

Solve: $x+19=-27$.

Answer

$x=-46$

Try It 2.1.6

Solve: $x+16=-34$.

Answer

$x=-50$

What happens when an equation has a number subtracted from the variable, as in the equation $x-5=8$? We use another property of equations to solve equations where a number is subtracted from the variable. We want to isolate the variable, so to 'undo' the subtraction we will add the number to both sides. We use the **Addition Property of Equality**.

ADDITION PROPERTY OF EQUALITY

For any numbers a, b, and c,

$$\begin{aligned} &\text{If} \quad a=b \\ &\text{then} \quad a+c=b+c \end{aligned}$$

When you add the same quantity from both sides of an equation, you still have equality.

In Exercise 2.1.4, 37 was added to the y and so we subtracted 37 to 'undo' the addition. In Exercise 2.1.7, we will need to 'undo' subtraction by using the **Addition Property of Equality**.

Example 2.1.7

Solve: $a-28=-37$.

Solution

	$a-28=-37$
Add 28 to each side to 'undo' the subtraction.	$a-28+28=-37+28$
Simplify.	$a=-9$
Check:	$a-28=-37$
Substitute $a=-9$	$-9-28=-37$
	$-37\stackrel{?}{=}-37$ ✓
	The solution to $a-28=-37$ is $a=-9$.

Try It 2.1.8

Solve: $n-61=-75$.

Answer

$n=-14$

Try It 2.1.9

Solve: $p-41=-73$.

Answer

$p=-32$

Example 2.1.10

Solve: $x-\frac{5}{8}=\frac{3}{4}$

Solution

	$x-\frac{5}{8}=\frac{3}{4}$
Use the Addition Property of Equality.	$x-\frac{5}{8}+\frac{5}{8}=\frac{3}{4}+\frac{5}{8}$
Find the LCD to add the fractions on the right.	$x=\frac{6}{8}+\frac{5}{8}$
Simplify.	$x=\frac{11}{8}$
Check:	$x-\frac{5}{8}=\frac{3}{4}$
Substitute $x=\frac{11}{8}$	$\frac{11}{8}-\frac{5}{8}\stackrel{?}{=}\frac{3}{4}$
Subtract.	$\frac{6}{8}\stackrel{?}{=}\frac{3}{4}$
Simplify.	$\frac{3}{4}=\frac{3}{4}\checkmark$
	The solution to $x-\frac{5}{8}=\frac{3}{4}$ is $x=\frac{11}{8}$.

Try It 2.1.11

Solve: $p-\frac{2}{3}=\frac{5}{6}$.

Answer

$p=\frac{9}{6}p=\frac{3}{2}$

Try It 2.1.12

Solve: $q-\frac{1}{2}=\frac{5}{6}$.

Answer

$q=\frac{4}{3}$

The next example will be an equation with decimals.

Example 2.1.13

Solve: $n-0.63=-4.2$.

Solution

	$n-0.63=-4.2$
Use the Addition Property of Equality.	$n-0.63+0.63=-4.2+0.63$
Add.	$n=-3.57$
Check:	$n=-3.57$

Let $n = -3.57$.	$-3.57 - 0.63 \stackrel{?}{=} -4.2$	
	$-4.2 = -4.2$ ✓	

Try It 2.1.14

Solve: $b - 0.47 = -2.1$.

Answer

$b = -1.63$

Try It 2.1.15

Solve: $c - 0.93 = -4.6$.

Answer

$c = -3.67$

Solve Equations That Require Simplification

In the previous examples, we were able to isolate the variable with just one operation. Most of the equations we encounter in algebra will take more steps to solve. Usually, we will need to simplify one or both sides of an equation before using the Subtraction or Addition Properties of Equality.

You should always simplify as much as possible before you try to isolate the variable. Remember that to simplify an expression means to do all the operations in the expression. Simplify one side of the equation at a time. Note that **simplification** is different from the process used to solve an equation in which we apply an operation to both sides.

Example 2.1.16: How to Solve Equations That Require Simplification

Solve: $9x - 5 - 8x - 6 = 7$.

Solution

Step 1. Simplify the expressions on each side as much as possible.	Rearrange the terms, using the Commutative Property of Addition. Combine like terms. Notice that each side is now simplified as much as possible.	$9x - 5 - 8x - 6 = 7$ $9x - 8x - 5 - 6 = 7$ $x - 11 = 7$
Step 2. Isolate the variable.	Now isolate x. Undo subtraction by adding 11 to both sides.	$x - 11 + 11 = 7 + 11$
Step 3. Simplify the expressions on both sides of the equation.		$x = 18$
Step 4. Check the solution.		**Check:** Substitute $x = 18$. $9x - 5 - 8x - 6 = 7$ $9(18) - 5 - 8(18) - 6 \stackrel{?}{=} 7$ $162 - 5 - 144 - 6 \stackrel{?}{=} 7$ $157 - 144 - 6 \stackrel{?}{=} 7$ $13 - 6 \stackrel{?}{=} 7$ $7 = 7$ ✓ The solution to $9x - 5 - 8x - 6 = 7$ is $x = 18$.

Try It 2.1.17

Solve: $8y-4-7y-7=4$.

Answer

$y=15$

Try It 2.1.18

Solve: $6z+5-5z-4=3$.

Answer

$z=2$

Example 2.1.19

Solve: $5(n-4)-4n=-8$.

Solution

We simplify both sides of the equation as much as possible before we try to isolate the variable.

	$5(n-4)-4n=-8$
Distribute on the left.	$5n-20-4n=-8$
Use the Commutative Property to rearrange terms.	$5n-4n-20=-8$
Combine like terms.	$n-20=-8$
Each side is as simplified as possible. Next, isolate n.	
Undo subtraction by using the Addition Property of Equality.	$n-20+20=-8+20$
Add.	$n=12$
Check. Substitute $n=12$. $5(n-4)-4n=-8$ $5(12-4)-4(12)\stackrel{?}{=}-8z$ $5(8)-48\stackrel{?}{=}-8$ $40-48\stackrel{?}{=}-8$ $-8=-8$ ✓	
	The solution to $5(n-4)-4n=-8$ is $n=12$.

Try It 2.1.20

Solve: $5(p-3)-4p=-10$.

Answer

$p=5$

Try It 2.1.21

Solve: $4(q+2)-3q=-8$.

Answer

$q=-16$

Example 2.1.22

Solve: $3(2y-1)-5y=2(y+1)-2(y+3)$.

Solution

We simplify both sides of the equation before we isolate the variable.

	$3(2y-1)-5y=2(y+1)-2(y+3)$
Distribute on both sides.	$6y-3-5y=2y+2-2y-6$
Use the Commutative Property of Addition.	$6y-5y-3=2y-2y+2-6$
Combine like terms.	$y-3=-4$
Each side is as simplified as possible. Next, isolate y.	
Undo subtraction by using the Addition Property of Equality.	$y-3+3=-4+3$
Add.	$y=-1$
Check. Let y=−1. $3(2y-1)-5y=2(y+1)-2(y+3)$ $3(2(-1)-1)-5(-1)\stackrel{?}{=}2(-1+1)-2(-1+3)$ $3(-2-1)+5\stackrel{?}{=}2(0)-2(2)$ $3(-3)+5\stackrel{?}{=}-4$ $-9+5\stackrel{?}{=}-4$ $-4=-4\checkmark$	
	The solution to $3(2y-1)-5y=2(y+1)-2(y+3)3(2y-1)-5y=2(y+1)-2(y+3$ is $y=-1$.

Try It 2.1.23

Solve: $4(2h-3)-7h=6(h-2)-6(h-1)$.

Answer

$h=6$

Try It 2.1.24

Solve: $2(5x+2)-9x=3(x-2)-3(x-4)$.

Answer

$x=2$

Translate to an Equation and Solve

To solve applications algebraically, we will begin by translating from English sentences into equations. Our first step is to look for the word (or words) that would translate to the **equals sign**. Here are some of the words that are commonly used.

Equals =

- is
- is equal to
- is the same as
- the result is
- gives

- was
- will be

The steps we use to translate a sentence into an equation are listed below.

TRANSLATE AN ENGLISH SENTENCE TO AN ALGEBRAIC EQUATION

1. Locate the “equals” word(s). Translate to an equals sign (=).
2. Translate the words to the left of the “equals” word(s) into an algebraic expression.
3. Translate the words to the right of the “equals” word(s) into an algebraic expression.

Example 2.1.25

Translate and solve: Eleven more than x is equal to 54.

Solution

Translate.	Eleven more than x is equal to 54 $x+11 = 54$
Subtract 11 from both sides.	$x+11-11 = 54-11$
Simplify.	$x = 43$
Check: Is 54 eleven more than 43? $43+11 = 54\checkmark$	

Try It 2.1.26

Translate and solve: Ten more than x is equal to 41.

Answer

$x+10=41;\quad x=31$

Try It 2.1.27

Translate and solve: Twelve less than x is equal to 51.

Answer

$y-12=51;\quad y=63$

Example 2.1.28

Translate and solve: The difference of $12t$ and $11t$ is -14.

Solution

Translate.	The difference of $12t$ and $11t$ is -14 $12t-11t = -14$
Simplify.	$t = -14$
Check: $12(-14)-11(-14) \stackrel{?}{=} -14$ $-168+154 \stackrel{?}{=} -14$ $-14 = -14\checkmark$	

Try It 2.1.29

Translate and solve: The difference of $4x$ and $3x$ is 14.

Answer

$4x - 3x = 14; \quad x = 14$

Try It 2.1.30

Translate and solve: The difference of $7a$ and $6a$ is -8.

Answer

$7a - 6a = -8; \quad a = -8$

Translate and Solve Applications

Most of the time a question that requires an algebraic solution comes out of a real life question. To begin with that question is asked in English (or the language of the person asking) and not in math symbols. Because of this, it is an important skill to be able to translate an everyday situation into algebraic language.

We will start by restating the problem in just one sentence, assign a variable, and then translate the sentence into an equation to solve. When assigning a variable, choose a letter that reminds you of what you are looking for. For example, you might use q for the number of quarters if you were solving a problem about coins.

Example 2.1.31: How to Solve Translate and Solve Applications

The MacIntyre family recycled newspapers for two months. The two months of newspapers weighed a total of 57 pounds. The second month, the newspapers weighed 28 pounds. How much did the newspapers weigh the first month?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.	The problem is about the weight of newspapers.	
Step 2. Identify what we are asked to find.	What are we asked to find?	"How much did the newspapers weigh the 2nd month?"
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Choose a variable.	Let w = weight of the newspapers the 1st month
Step 4. Translate into an equation. It may be helpful to restate the problem in one sentence with the important information.	Restate the problem. We know the weight of the newspapers the second month is 28 pounds. Translate into an equation, using the variable w.	Weight of newspapers the 1st month plus the weight of the newspapers the 2nd month equals 57 pounds. Weight from 1st month plus 28 equals 57. $w + 28 = 57$
Step 5. Solve the equation using good algebra techniques.	Solve.	$w + 28 - 28 = 57 - 28$ $w = 29$

Step 6. Check the answer in the problem and make sure it makes sense.	Does 1st month's weight plus 2nd month's weight equal 57 pounds?	**Check:** Does 1st month's weight plus 2nd month's weight equal 57 pounds? $29 + 28 \stackrel{?}{=} 57$ $57 = 57$ ✓
Step 7. Answer the question with a complete sentence.	Write a sentence to answer "How much did the newspapers weigh the 2nd month?"	The 2nd month the newspapers weighed 29 pounds.

? Try It 2.1.32

Translate into an algebraic equation and solve:

The Pappas family has two cats, Zeus and Athena. Together, they weigh 23 pounds. Zeus weighs 16 pounds. How much does Athena weigh?

Answer

7 pounds

? Try It 2.1.33

Translate into an algebraic equation and solve:

Sam and Henry are roommates. Together, they have 68 books. Sam has 26 books. How many books does Henry have?

Answer

42 books

SOLVE AN APPLICATION.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with the important information.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

✓ Example 2.1.34

Randell paid $28,675 for his new car. This was $875 less than the sticker price. What was the sticker price of the car?

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for.	"What was the sticker price of the car?"
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let s = the sticker price of the car.
Step 4. Translate into an equation. Restate the problem in one sentence.	$28675 is $875 less than the sticker price
Step 5. Solve the equation.	$28675 is $875 less than s $28675 = s - 875$ $28675 + 875 = s - 875 + 875$ $29550 = s$
Step 6. Check the answer. Is $875 less than $29550 equal to $28675? $29550 - 875 \stackrel{?}{=} 28675$ $28675 = 28675$✓	
Step 7. Answer the question with a complete sentence.	The sticker price of the car was $29550.

? Try It 2.1.35

Translate into an algebraic equation and solve:

Eddie paid $19875 for his new car. This was $1025 less than the sticker price. What was the sticker price of the car?

Answer

$20900

? Try It 2.1.36

Translate into an algebraic equation and solve:

The admission price for the movies during the day is $7.75. This is $3.25 less the price at night. How much does the movie cost at night?

Answer

$11.00

Key Concepts

- **To Determine Whether a Number is a Solution to an Equation**
  1. **Substitute the number in for the variable in the equation.**
  2. **Simplify the expressions on both sides of the equation.**
  3. **Determine whether the resulting statement is true.**
     - If it is true, the number is a solution.
     - If it is not true, the number is not a solution.
- **Addition Property of Equality**
  - For any numbers a, b and c, if $a = b$, then $a + c = b + c$.
- **Subtraction Property of Equality**
  - For any numbers a, b and c, if $a = b$, then $a - c = b - c$.
- **To Translate a Sentence to an Equation**

1. Locate the “equals” word(s). Translate to an equal sign (=).
2. Translate the words to the left of the “equals” word(s) into an algebraic expression.
3. Translate the words to the right of the “equals” word(s) into an algebraic expression.

- **To Solve an Application**
  1. Read the problem. Make sure all the words and ideas are understood.
  2. Identify what we are looking for.
  3. Name what we are looking for. Choose a variable to represent that quantity.
  4. Translate into an equation. It may be helpful to restate the problem in one sentence with the important information.
  5. Solve the equation using good algebra techniques.
  6. Check the answer in the problem and make sure it makes sense.
  7. Answer the question with a complete sentence.

Glossary

solution of an equation

A solution of an equation is a value of a variable that makes a true statement when substituted into the equation.

This page titled 2.1: Solve Equations Using the Subtraction and Addition Properties of Equality is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.1: Solve Equations Using the Subtraction and Addition Properties of Equality** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.1E: Exercises

Practice Makes Perfect

Verify a Solution of an Equation

In the following exercises, determine whether the given value is a solution to the equation.

Exercise 2.1E.1

Is $y = \frac{5}{3}$ a solution of
$6y + 10 = 12y$?

Answer

Yes

Exercise 2.1E.2

Is $x = \frac{9}{4}$ a solution of
$4x + 9 = 8x$?

Exercise 2.1E.3

Is $u = -\frac{1}{2}$ a solution of
$8u - 1 = 6u$?

Answer

No

Exercise 2.1E.4

Is $v = -\frac{1}{3}$ a solution of
$9v - 2 = 3v$?

Solve Equations using the Subtraction and Addition Properties of Equality

In the following exercises, solve each equation using the Subtraction and Addition Properties of Equality.

Exercise 2.1E.5

$x + 24 = 35$

Answer

x = 11

Exercise 2.1E.6

$x + 17 = 22$

Exercise 2.1E.7

$y + 45 = -66$

Answer

y = -111

Exercise 2.1E.8

$y + 39 = -83$

Exercise 2.1E.9

$b + \frac{1}{4} = \frac{3}{4}$

Answer

$b = \frac{1}{2}$

Exercise 2.1E.10

$a + \frac{2}{5} = \frac{4}{5}$

Exercise 2.1E.11

$p + 2.4 = -9.3$

Answer

p = -11.7

Exercise 2.1E.12

$m + 7.9 = 11.6$

Exercise 2.1E.13

$a - 45 = 76$

Answer

a = 121

Exercise 2.1E.14

$a - 30 = 57$

Exercise 2.1E.15

$m - 18 = -200$

Answer

m = -182

Exercise 2.1E.16

$m - 12 = -12$

Exercise 2.1E.17

$x - \frac{1}{3} = 2$

Answer

$x = \frac{7}{3}$

Exercise 2.1E.18

$x - \frac{1}{5} = 4$

Exercise 2.1E.19

$y - 3.8 = 10$

Answer

y = 10.8

Exercise 2.1E.20

$y - 7.2 = 5$

Exercise 2.1E.21

$x - 165 = -420$

Answer

$x = -255$

Exercise 2.1E.22

$z - 101 = -314$

Exercise 2.1E.23

$z + 0.52 = -8.5$

Answer

$z = -9.02$

Exercise 2.1E.24

$x + 0.93 = -4.1$

Exercise 2.1E.25

$q + \frac{3}{4} = \frac{1}{2}$

Answer

$q = -\frac{1}{4}$

Exercise 2.1E.26

$p+\frac{1}{3}=\frac{5}{6}$

Exercise 2.1E.27

$p-\frac{2}{5}=\frac{2}{3}$

Answer

$p=\frac{16}{15}$

Exercise 2.1E.28

$y-\frac{3}{4}=\frac{3}{5}$

Solve Equations that Require Simplification

In the following exercises, solve each equation.

Exercise 2.1E.29

$c+31-10=46$

Answer

c = 25

Exercise 2.1E.30

$m+16-28=5$

Exercise 2.1E.31

$9x+5-8x+14=20$

Answer

x = 1

Exercise 2.1E.32

$6x+8-5x+16=32$

Exercise 2.1E.33

$-6x-11+7x-5=-16$

Answer

x = 0

Exercise 2.1E.34

$-8n-17+9n-4=-41$

Exercise 2.1E.35

$5(y-6)-4y=-6$

Answer

$y=24$

Exercise 2.1E.36

$9(y-2)-8y=-16$

Exercise 2.1E.37

$8(u+1.5)-7u=4.9$

Answer

$u=-7.1$

Exercise 2.1E.38

$5(w+2.2)-4w=9.3$

Exercise 2.1E.39

$6a-5(a-2)+9=-11$

Answer

$a=-30$

Exercise 2.1E.40

$8c-7(c-3)+4=-16$

Exercise 2.1E.41

$6(y-2)-5y=4(y+3)-4(y-1)$

Answer

y =28

Exercise 2.1E.42

$9(x-1)-8x=-3(x+5)+3(x-5)$

Exercise 2.1E.43

$3(5n-1)-14n+9=10(n-4)-6n-4(n+1)$

Answer

n = -50

Exercise 2.1E.44

$2(8m+3)-15m-4=9(m+6)-2(m-1)-7m$

Exercise 2.1E.45

$-(j+2)+2j-1=5$

Answer

j = 8

Exercise 2.1E.46

$-(k+7)+2k+8=7$

Exercise 2.1E.47

$-\left(\frac{1}{4}a-\frac{3}{4}\right)+\frac{5}{4}a=-2$

Answer

$a=-\frac{11}{4}$

Exercise 2.1E.48

$-\left(\frac{2}{3}d-\frac{1}{3}\right)+\frac{5}{3}d=-4$

Exercise 2.1E.49

$8(4x+5)-5(6x)-x$
$=53-6(x+1)+3(2x+2)$

Answer

x=13

Exercise 2.1E.50

$6(9y-1)-10(5y)-3y$
$=22-4(2y-12)+8(y-6)$

Translate to an Equation and Solve

In the following exercises, translate to an equation and then solve it.

Exercise 2.1E.51

Nine more than x is equal to 52.

Answer

$x+9=52; x=43$

Exercise 2.1E.52

The sum of x and -15 is 23.

Exercise 2.1E.53

Ten less than m is -14.

Answer

$m-10=-14; m=-4$

Exercise 2.1E.54

Three less than y is -19.

Exercise 2.1E.55

The sum of y and -30 is 40.

Answer

$y+(-30)=40; y=70$

Exercise 2.1E.56

Twelve more than p is equal to 67.

Exercise 2.1E.57

The difference of $9x$ and $8x$ is 107.

Answer

$9x-8x=107; 107$

Exercise 2.1E.58

The difference of $5c$ and $4c$ is 602.

Exercise 2.1E.59

The difference of $5c$ and $4c$ is 602

Answer

$n-\frac{1}{6}=\frac{1}{2}; \frac{2}{3}$

Exercise 2.1E.60

The difference of f and $\frac{1}{3}$ is $\frac{1}{12}$.

Exercise 2.1E.61

The sum of $-4n$ and $5n$ is -82

Answer

$-4n+5n=-82;-82$

Exercise 2.1E.62

The sum of $-9m$ and $10m$ is -95

Translate and Solve Applications

In the following exercises, translate into an equation and solve.

Exercise 2.1E.63

Distance Avril rode her bike a total of 18 miles, from home to the library and then to the beach. The distance from Avril's house to the library is 7 miles. What is the distance from the library to the beach?

Answer

11 miles

Exercise 2.1E.64

Reading Jeff read a total of 54 pages in his History and Sociology textbooks. He read 41 pages in his History textbook. How many pages did he read in his Sociology textbook?

Exercise 2.1E.65

Age Eva's daughter is 15 years younger than her son. Eva's son is 22 years old. How old is her daughter?

Answer

7 years old

Exercise 2.1E.66

Age Pablo's father is 3 years older than his mother. Pablo's mother is 42 years old. How old is his father?

Exercise 2.1E.67

Groceries For a family birthday dinner, Celeste bought a turkey that weighed 5 pounds less than the one she bought for Thanksgiving. The birthday turkey weighed 16 pounds. How much did the Thanksgiving turkey weigh?

Answer

21 pounds

Exercise 2.1E.68

Weight Allie weighs 8 pounds less than her twin sister Lorrie. Allie weighs 124 pounds. How much does Lorrie weigh?

Exercise 2.1E. 69

Health Connor's temperature was 0.7 degrees higher this morning than it had been last night. His temperature this morning was 101.2 degrees. What was his temperature last night?

Answer

100.5 degrees

Exercise 2.1E. 70

Health The nurse reported that Tricia's daughter had gained 4.2 pounds since her last checkup and now weighs 31.6 pounds. How much did Tricia's daughter weigh at her last checkup?

Exercise 2.1E. 71

Salary Ron's paycheck this week was $17.43 less than his paycheck last week. His paycheck this week was $103.76. How much was Ron's paycheck last week?

Answer

$121.19

Exercise 2.1E. 72

Textbooks Melissa's math book cost $22.85 less than her art book cost. Her math book cost $93.75. How much did her art book cost?

Everyday Math

Exercise 2.1E. 73

Construction Miguel wants to drill a hole for a $\frac{5}{8}$ inch screw. The hole should be $\frac{1}{12}$ inch smaller than the screw. Let d equal the size of the hole he should drill. Solve the equation $d = \frac{5}{8} - \frac{1}{12}$ to see what size the hole should be.

Answer

$d = \frac{13}{24}$ inch

Exercise 2.1E. 74

Baking Kelsey needs $\frac{2}{3}$ cup of sugar for the cookie recipe she wants to make. She only has $\frac{3}{8}$ cup of sugar and will borrow the rest from her ner neighbor. Let s equal the amount of sugar she will borrow. Solve the equation $\frac{3}{8} + s = \frac{2}{3}$ to find the amount of sugar she should ask to borrow.

Writing Exercises

Exercise 2.1E. 75

Is -8 a solution to the equation $3x = 16 - 5x$? How do you know?

Answer

No. Justifications will vary.

? Exercise $2.1E.76$

What is the first step in your solution to the equation $10x + 2 = 4x + 26$?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
verify a solution of an equation.			
solve equations using the subtraction and addition properties of equality.			
solve equations that require simplification.			
translate to an equation and solve.			
translate and solve applications.			

ⓑ If most of your checks were:

…confidently. Congratulations! You have achieved your goals in this section! Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific!

…with some help. This must be addressed quickly as topics you do not master become potholes in your road to success. Math is sequential - every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no - I don't get it! This is critical and you must not ignore it. You need to get help immediately or you will quickly be overwhelmed. See your instructor as soon as possible to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 2.1E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.1E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.2: Solve Equations using the Division and Multiplication Properties of Equality

Learning Objectives

By the end of this section, you will be able to:

- Solve equations using the Division and Multiplication Properties of Equality
- Solve equations that require simplification
- Translate to an equation and solve
- Translate and solve applications

Note

Before you get started, take this readiness quiz.

1. Simplify: $-7\left(\frac{1}{-7}\right)$.
   If you missed this problem, review Exercise 1.6.13.
2. Evaluate $9x+2$ when $x=-3$.
   If you missed this problem, review Exercise 1.5.34.

Solve Equations Using the Division and Multiplication Properties of Equality

You may have noticed that all of the equations we have solved so far have been of the form $x+a=b$ or $x-a=b$. We were able to isolate the variable by adding or subtracting the constant term on the side of the equation with the variable. Now we will see how to solve equations that have a variable multiplied by a constant and so will require division to isolate the variable.

Let's look at our puzzle again with the envelopes and counters in Figure 2.2.1.




Figure 2.2.1: The illustration shows a model of an equation with one variable multiplied by a constant. On the left side of the workspace are two instances of the unknown (envelope), while on the right side of the workspace are six counters.

In the illustration there are two identical envelopes that contain the same number of counters. Remember, the left side of the workspace must equal the right side, but the counters on the left side are "hidden" in the envelopes. So how many counters are in each envelope?

How do we determine the number? We have to separate the counters on the right side into two groups of the same size to correspond with the two envelopes on the left side. The 6 counters divided into 2 equal groups gives 3 counters in each group (since $6\div 2=3$).

What equation models the situation shown in Figure 2.2.2? There are two envelopes, and each contains xx counters. Together, the two envelopes must contain a total of 6 counters.




Figure 2.2.2: The illustration shows a model of the equation $2x=6$.

	$2x=6$

If we divide both sides of the equation by 2, as we did with the envelopes and counters,	$\frac{2x}{2}=\frac{6}{2}$
we get:	$x=3$

We found that each envelope contains 3 counters. Does this check? We know $2 \cdot 3 = 6$, so it works! Three counters in each of two envelopes does equal six!

This example leads to the **Division Property of Equality**.

THE DIVISION PROPERTY OF EQUALITY

For any numbers *a*, *b*, and *c*, and $c \neq 0$,

$$\begin{array}{rcl} \text{If} & a & = & b \\ \text{then} & \frac{a}{c} & = & \frac{b}{c} \end{array} \qquad (2.2.1)$$

When you divide both sides of an equation by any non-zero number, you still have equality.

Note

Doing the Manipulative Mathematics activity "**Division Property of Equality**" will help you develop a better understanding of how to solve equations by using the Division Property of Equality.

The goal in solving an equation is to 'undo' the operation on the variable. In the next example, the variable is multiplied by 5, so we will divide both sides by 5 to 'undo' the multiplication.

Example 2.2.1

Solve: $5x = -27$.

Solution

To isolate x, "undo" the multiplication by 5.		$5x=-27$
Divide to 'undo' the multiplication.		$\frac{5x}{5}=\frac{-27}{5}$
Simplify.		$x=-\frac{27}{5}$
Check:	$5x=-27$	
Substitute $-\frac{27}{5}$ for x.	$5\left(-\frac{27}{5}\right)\stackrel{?}{=}-27$	
	$-27=-27$ ✓	
		Since this is a true statement, $x=-\frac{27}{5}$ is the solution to $5x=-27$.

Try It 2.2.2

Solve: $3y = -41$.

Answer

$y=-\frac{41}{3}$

Try It 2.2.3

Solve: $4z = -55$.

Answer

$y = -\frac{55}{4}$

Consider the equation $\frac{x}{4} = 3$. We want to know what number divided by 4 gives 3. So to "undo" the division, we will need to multiply by 4. The **Multiplication Property of Equality** will allow us to do this. This property says that if we start with two equal quantities and multiply both by the same number, the results are equal.

THE MULTIPLICATION PROPERTY OF EQUALITY

For any numbers *a*, *b*, and *c*,

$$\begin{array}{lll} \text{If} & a & = & b \\ \text{then} & ac & = & bc \end{array} \qquad (2.2.2)$$

If you multiply both sides of an equation by the same number, you still have equality.

Example 2.2.4

Solve: $\frac{y}{-7} = -14$

Solution

Here y is divided by −7. We must multiply by −7 to isolate y.

	$\frac{y}{-7} = -14$
Multiply both sides by −7.	$-7\left(\frac{y}{-7}\right) = -7(-14)$
Multiply.	$\frac{-7y}{7} = 98$
Simplify.	$y = 98$
Check: $\frac{y}{-7} = -14$	
Substitute y=98.	$\frac{98}{-7} \stackrel{?}{=} -14$
Divide.	$-14 = -14$ ✓

Try It 2.2.5

Solve: $\frac{a}{-7} = -42$

Answer

$a = 294$

Try It 2.2.6

Solve: $\frac{b}{-6} = -24$

Answer

$b = 144$

Example 2.2.7

Solve: $-n = 9$

Solution

	$-n = 9$
Remember −n is equivalent to −1n.	$-1n = 9$
Divide both sides by −1.	$\frac{-1n}{-1} = \frac{9}{-1}$
Divide.	$n = -9$
Notice that there are two other ways to solve −n=9. We can also solve this equation by multiplying both sides by −1 and also by taking the opposite of both sides.	
Check:	$-n = 9$
Substitute n=−9.	$-(-9) \stackrel{?}{=} 9$
Simplify.	$9 = 9$ ✓

Try It 2.2.8

Solve: $-k = 8$.

Answer

$k = -8$

Try It 2.2.9

Solve: $-g = 3$.

Answer

$g = -3$

Example 2.2.10

Solve: $\frac{3}{4}x = 12$

Solution

Since the product of a number and its reciprocal is 1, our strategy will be to isolate x by multiplying by the reciprocal of $\frac{3}{4}$.

	$\frac{3}{4}x = 12$
Multiply by the reciprocal of $\frac{3}{4}$.	$\frac{4}{3} \cdot \frac{3}{4}x = \frac{4}{3} \cdot 12$
Reciprocals multiply to 1.	$1x = \frac{4}{3} \cdot \frac{12}{1}$
Multiply.	$x = 16$
Notice that we could have divided both sides of the equation $\frac{3}{4}x = 12$ by $\frac{3}{4}$ to isolate x. While this would work, most people would find multiplying by the reciprocal easier.	
Check:	$\frac{3}{4}x = 12$

Substitute $x = 16$.	$\frac{3}{4} \cdot 16 \stackrel{?}{=} 12$
	$12 = 12$ ✓

Try It 2.2.11

Solve: $\frac{2}{5}n = 14$.

Answer

$n = 35$

Try It 2.2.12

Solve: $\frac{5}{6}y = 15$.

Answer

$y = 18$

In the next example, all the variable terms are on the right side of the equation. As always, our goal in solving the equation is to isolate the variable.

Example 2.2.13

Solve: $\frac{8}{15} = -\frac{4}{5}x$

Solution

	$\frac{8}{15} = -\frac{4}{5}x$
Multiply by the reciprocal of $-\frac{4}{5}$.	$\left(-\frac{5}{4}\right)\left(\frac{8}{15}\right) = \left(-\frac{5}{4}\right)\left(-\frac{4}{5}x\right)$
Reciprocals multiply to 1.	$-\frac{\cancel{5} \cdot \cancel{4} \cdot 2}{\cancel{4} \cdot 3 \cdot \cancel{5}} = 1x$
Multiply.	$-\frac{2}{3} = x$
Check:	$\frac{8}{15} = -\frac{4}{5}x$
Let $x = -\frac{2}{3}$.	$\frac{8}{15} = -\frac{4}{5}\left(-\frac{2}{3}\right)$
	$\frac{8}{15} = \frac{8}{15}$ ✓

Try It 2.2.14

Solve: $\frac{9}{25} = -\frac{4}{5}z$

Answer

$z = -\frac{9}{5}$

Try It 2.2.15

$\frac{5}{6} = -\frac{8}{3}r$

Answer

$r = -\frac{5}{16}$

Solve Equations That Require Simplification

Many equations start out more complicated than the ones we have been working with.

With these more complicated equations the first step is to simplify both sides of the equation as much as possible. This usually involves combining like terms or using the distributive property.

Example 2.2.16

Solve: $14 - 23 = 12y - 4y - 5y$.

Solution

Begin by simplifying each side of the equation.

	$14 - 23 \stackrel{?}{=} -36 + 12 + 15$
Simplify each side.	$-9 = -9$ ✓
Divide both sides by 3 to isolate y.	$14 - 23 = 12y - 4y - 5y$
Divide.	$-9 = 3y$
Check:	$14 - 23 = 12y - 4y - 5y$
Substitute $y = -3$.	$14 - 23 \stackrel{?}{=} 12(-3) - 4(-3) - 5(-3)$
	$14 - 23 \stackrel{?}{=} -36 + 12 + 15$
	$-9 = -9$ ✓

Try It 2.2.17

Solve: $18 - 27 = 15c - 9c - 3c$.

Answer

$c = -3$

Try It 2.2.18

Solve: $18 - 22 = 12x - x - 4x$.

Answer

$x = -\frac{4}{7}$

Example 2.2.19

Solve: $-4(a - 3) - 7 = 25$.

Solution

Here we will simplify each side of the equation by using the distributive property first.

	$-4(a-3)-7=25$
Distribute.	$-4a+12-7=25$
Simplify.	$-4a+5=25$
Simplify.	$-4a=20$
Divide both sides by -4 to isolate a.	$\frac{-4a}{-4}=\frac{20}{-4}$
Divide.	$a=-5$
Check:	$-4(a-3)-7=25$
Substitute $a=-5$	$-4(-5-3)-7\stackrel{?}{=}25$
	$-4(-8)-7\stackrel{?}{=}25$
	$32-7\stackrel{?}{=}25$
	$25=25$ ✓

Try It 2.2.20

Solve: $-4(q-2)-8=24$.

Answer

$q=-6$

Try It 2.2.21

Solve: $-6(r-2)-12=30$.

Answer

$r=-5$

Now we have covered all four properties of equality—subtraction, addition, division, and multiplication. We'll list them all together here for easy reference.

PROPERTIES OF EQUALITY

When you add, subtract, multiply, or divide the same quantity from both sides of an equation, you still have equality.

$$\begin{array}{ll} \textbf{Subtraction Property of Equality} & \textbf{Addition Property of Equality} \\ \text{For any real numbers a, b, and c,} & \text{For any real numbers a, b, and c,} \\ \text{if } a=b, & \text{if } a=b, \\ \text{then } a-c=b-c & \text{then } a+c=b+c \\ \textbf{Divition Property of Equality} & \textbf{Multiplication Property of Equality} \\ \text{For any real numbers a, b, and c,} & \text{For any real numbers a, b, and c,} \\ \text{if } a=b, & \text{if } a=b, \\ \text{then } a-c=b-c & \text{then } a+c=b+c \end{array} \tag{2.2.3}$$

When you add, subtract, multiply, or divide the same quantity from both sides of an equation, you still have equality.

Translate to an Equation and Solve

In the next few examples, we will translate sentences into equations and then solve the equations. You might want to review the translation table in the previous chapter.

Example 2.2.22

Translate and solve: The number 143 is the product of −11 and *y*.

Solution

Translate.	The number 143 is the product of –11 and y. $143 = -11y$
Divide by −11.	$\frac{143}{-11} = \frac{-11y}{-11}$
Simplify.	$-13 = y$
Check: $143 = -11y$ $143 \stackrel{?}{=} -11(-13)$ (2.2.4) $143 = 143\checkmark$	

Try It 2.2.23

Translate and solve: The number 132 is the product of *−12* and *y*.

Answer

132=−12y;y=−11

Try It 2.2.24

Translate and solve: The number 117 is the product of *−13* and *z*.

Answer

117=−13z;z=−9

Example 2.2.25

Translate and solve: *n* divided by 8 is −32.

Solution

Begin by translating the sentence into an equation. Translate.		*n* divided by 8 is –32. $\frac{n}{8} = -32$
Multiple both sides by 8.		$8 \cdot \frac{n}{8} = 8(-32)$
Simplify.		$n = -256$
Check:	Is nn divided by 8 equal to −32?	
Let $n = -256$.	Is −256 divided by 88 equal to −32?	
Translate.	$\frac{-256}{8} \stackrel{?}{=} -32$	

Simplify.	$-32 = -32$✓

? Try It 2.2.26

Translate and solve: nn divided by 7 is equal to −21.

Answer

$\frac{n}{7} = -21; n = -147$

? Try It 2.2.27

Translate and solve: n divided by 8 is equal to −56.

Answer

$\frac{n}{8} = -56; n = -448$

✓ Example 2.2.28

Translate and solve: The quotient of yy and −4 is 68.

Solution

Begin by translating the sentence into an equation.

Translate.	The quotient of y and -4 is 68. $\frac{y}{-4} = 68$
Multiply both sides by -4.	$-4\left(\frac{y}{-4}\right) = -4(68)$
Simplify.	$y = -272$
Check:	Is the quotient of y and −4 equal to 68?
Let y=−272.	Is the quotient of −272 and −4 equal to 68?
Translate.	$\frac{-272}{-4} \stackrel{?}{=} 68$
Simplify.	$68 = 68$✓

? Try It 2.2.29

Translate and solve: The quotient of q and −8 is 72.

Answer

$\frac{q}{-8} = 72; q = -576$

? Try It 2.2.30

Translate and solve: The quotient of p and −9 is 81.

Answer

$\frac{p}{-9} = 81; p = -729$

Example 2.2.31

Translate and solve: Three-fourths of p is 18.

Solution

Begin by translating the sentence into an equation. Remember, “of” translates into multiplication.

Translate.	Three-fourths of p is 18. $\frac{3}{4}p = 18$
Multiply both sides by $\frac{4}{3}$.	$\frac{4}{3} \cdot \frac{3}{4}p = \frac{4}{3} \cdot 18$
Simplify.	$p = 24$
Check:	Is three-fourths of p equal to 18?
Let p = 24.	Is three-fourths of 24 equal to 18?
Translate.	$\frac{3}{4} \cdot 24 \stackrel{?}{=} 18$
Simplify.	$18 = 18$✓

Try It 2.2.32

Translate and solve: Two-fifths of f is 16.

Answer

$\frac{2}{5}f = 16; f = 40$

Try It 2.2.33

Translate and solve: Three-fourths of f is 21.

Answer

$\frac{3}{4}f = 21; f = 28$

Example 2.2.34

Translate and solve: The sum of three-eighths and x is one-half.

Solution

Begin by translating the sentence into an equation.

Translate.	The sum of three – eighths and x is $\frac{1}{2}$ $\frac{3}{8} + x = \frac{1}{2}$
Subtract $\frac{3}{8}$ from each side.	$\frac{3}{8} - \frac{3}{8} + x = \frac{1}{2} - \frac{3}{8}$
Simplify and rewrite fractions with common denominators.	$x = \frac{4}{8} - \frac{3}{8}$
Simplify.	$x = \frac{1}{8}$

Check:	Is the sum of three-eighths and x equal to one-half?
Let $x = \frac{1}{8}$.	Is the sum of three-eighths and one-eighth equal to one-half?
Translate.	$\frac{3}{8} + \frac{1}{8} \stackrel{?}{=} \frac{1}{2}$
Simplify.	$\frac{4}{8} \stackrel{?}{=} \frac{1}{2}$
Simplify.	$\frac{1}{2} = \frac{1}{2}$ ✓

Try It 2.2.35

Translate and solve: The sum of five-eighths and *x* is one-fourth.

Answer

$\frac{5}{8} + x = \frac{1}{4}; x = -\frac{3}{8}$

Try It 2.2.36

Translate and solve: The sum of three-fourths and *x* is five-sixths.

Answer

$\frac{3}{4} + x = \frac{5}{6}; x = \frac{1}{12}$

Translate and Solve Applications

To solve applications using the Division and Multiplication Properties of Equality, we will follow the same steps we used in the last section. We will restate the problem in just one sentence, assign a variable, and then translate the sentence into an equation to solve.

Example 2.2.37

Denae bought 6 pounds of grapes for $10.74. What was the cost of one pound of grapes?

Solution

What are you asked to find?	The cost of 1 pound of grapes	
Assign a variable.	Let c = the cost of one pound.	
Write a sentence that gives the information to find it.	The cost of 6 pounds is $10.74	
Translate into an equation.	$6c = 10.74$	
Solve.	$\frac{6c}{c} = \frac{10.74}{6}$	(2.2.5)
	$c = 1.79$	
	The grapes cost $1.79 per pound.	
Check: If one pound costs $1.79, *do* 6 pounds cost $10.74?		
$6(1.79) \stackrel{?}{=} 10.74$		
$10.74 = 10.74$✓		

? Try It 2.2.38

Translate and solve:

Arianna bought a 24-pack of water bottles for $9.36. What was the cost of one water bottle?

Answer

$0.39

? Try It 2.2.39

Translate and solve:

At JB's Bowling Alley, 6 people can play on one lane for $34.98. What is the cost for each person?

Answer

$5.83

✓ Example 2.2.40

Andreas bought a used car for $12,000. Because the car was 4-years old, its price was $\frac{3}{4}$ of the original price, when the car was new. What was the original price of the car?

Solution

What are you asked to find?	The original price of the car
Assign a variable.	Let p = the original price.
Write a sentence that gives the information to find it.	\$12000 is $\frac{3}{4}$ of the original price.
Translate into an equation.	$12000 = \frac{3}{4}p$
Solve.	$\frac{3}{4}(12000) = \frac{4}{3} \cdot \frac{3}{4}p$ $16000 = p$
	The original cost of the car was \$16000.

(2.2.6)

Check: Is $\frac{3}{4}$ of \$16000 equal to \$12000

$\frac{3}{4} \cdot 16000 \stackrel{?}{=} 12000$
$12000 = 12000$✓

? Try It 2.2.41

Translate and solve:

The annual property tax on the Mehta's house is \$1,800, calculated as $\frac{15}{1000}$ of the assessed value of the house. What is the assessed value of the Mehta's house?

Answer

\$120000

? Try It 2.2.42

Translate and solve:

Stella planted 14 flats of flowers in $\frac{2}{3}$ of her garden. How many flats of flowers would she need to fill the whole garden?

Answer

21 flats

Key Concepts

- **The Division Property of Equality**—For any numbers a, b, and c, and $c \neq 0$, if $a = b$, then $\frac{a}{c} = \frac{b}{c}$.
  When you divide both sides of an equation by any non-zero number, you still have equality.
- **The Multiplication Property of Equality**—For any numbers a, b, and c, if $a = b$, then $ac = bc$.
  If you multiply both sides of an equation by the same number, you still have equality.

This page titled 2.2: Solve Equations using the Division and Multiplication Properties of Equality is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.2: Solve Equations using the Division and Multiplication Properties of Equality** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.2E: Exercises

Practice Makes Perfect

Solve Equations Using the Division and Multiplication Properties of Equality

In the following exercises, solve each equation using the Division and Multiplication Properties of Equality and check the solution.

Exercise 2.2E.1

$8x = 56$

Answer

$x = 7$

Exercise 2.2E.2

$7p = 63$

Exercise 2.2E.3

$-5c = 55$

Answer

$c = -11$

Exercise 2.2E.4

$-9x = -27$

Exercise 2.2E.5

$-809 = 15y$

Answer

$y = -\frac{809}{15}$

Exercise 2.2E.6

$-731 = 19y$

Exercise 2.2E.7

$-37p = -541$

Answer

$p = -\frac{541}{37}$

Exercise 2.2E.8

$-19m = -586$

Exercise 2.2E.9

$0.25z = 3.25$

Answer

z= 13

Exercise 2.2E.10

$0.75a = 11.25$

Exercise 2.2E.11

$-13x = 0$

Answer

$x = 0$

Exercise 2.2E.12

$24x = 0$

Exercise 2.2E.13

$\frac{x}{4} = 35$

Answer

$x = 140$

Exercise 2.2E.14

$\frac{z}{2} = 54$

Exercise 2.2E.15

$-20 = \frac{q}{-5}$

Answer

$q = 100$

Exercise 2.2E.16

$\frac{c}{-3} = -12$

Exercise 2.2E.17

$\frac{y}{9} = -16$

Answer

$y = -144$

Exercise 2.2E.18

$\frac{q}{6} = -38$

Exercise 2.2E.19

$\frac{m}{-12} = 45$

Answer

$m = -540$

Exercise 2.2E.20

$-24 = \frac{p}{-20}$

Exercise 2.2E.21

$-y = 6$

Answer

$y = -6$

Exercise 2.2E.22

$-u = 15$

Exercise 2.2E.23

$-v = -72$

Answer

$v = 72$

Exercise 2.2E.24

$-x = -39$

Exercise 2.2E.25

$\frac{2}{3}y = 48$

Answer

$y = 72$

Exercise 2.2E.26

$\frac{3}{5}r = 75$

Exercise 2.2E.27

$-\frac{5}{8}w = 40$

Answer

$w = -64$

Exercise 2.2E.28

$24 = -\frac{3}{4}x$

Exercise 2.2E.29

$-\frac{2}{5} = \frac{1}{10}a$

Answer

$a = -4$

Exercise 2.2E.30

$-\frac{1}{3}q = -\frac{5}{6}$

Exercise 2.2E.31

$-\frac{7}{10}x = -\frac{14}{3}$

Answer

$x = \frac{20}{3}$

Exercise 2.2E.32

$\frac{3}{8}y = -\frac{1}{4}$

Exercise 2.2E.33

$\frac{7}{12} = -\frac{3}{4}p$

Answer

$p = -\frac{7}{9}$

Exercise 2.2E.34

$\frac{11}{18} = -\frac{5}{6}q$

Exercise 2.2E.35

$-\frac{5}{18} = -\frac{10}{9}u$

Answer

$u = \frac{1}{4}$

Exercise 2.2E.36

$-\frac{7}{20}=-\frac{7}{4}v$

Solve Equations That Require Simplification

In the following exercises, solve each equation requiring simplification.

Exercise 2.2E.37

$100-16=4p-10p-p$

Answer

$p=-12$

Exercise 2.2E.38

$-18-7=5t-9t-6t$

Exercise 2.2E.39

$\frac{7}{8}n-\frac{3}{4}n=9+2$

Answer

$n=88$

Exercise 2.2E.40

$\frac{5}{12}q+\frac{1}{2}q=25-3$

Exercise 2.2E.41

$0.25d+0.10d=6-0.75$

Answer

d=15

Exercise 2.2E.42

$0.05p-0.01p=2+0.24$

Exercise 2.2E.43

$-10(q-4)-57=93$

Answer

$q=-11$

Exercise 2.2E.44

$-12(d-5)-29=43$

Exercise 2.2E.45

$-10(x+4)-19=85$

Answer

$x=-\frac{72}{5}$

Exercise 2.2E.46

$-15(z+9)-11=75$

Mixed Practice

In the following exercises, solve each equation.

Exercise 2.2E.47

$\frac{9}{10}x=90$

Answer

$x=100$

Exercise 2.2E.48

$\frac{5}{12}y=60$

Exercise 2.2E.49

$y+46=55$

Answer

$y=9$

Exercise 2.2E.50

$x+33=41$

Exercise 2.2E.51

$\frac{w}{-2}=99$

Answer

$w=-198$

Exercise 2.2E.52

$\frac{s}{-3}=-60$

Exercise 2.2E.53

$27 = 6a$

Answer

$a = \frac{9}{2}$

Exercise 2.2E.54

$-a = 7$

Exercise 2.2E.55

$-x = 2$

Answer

$x = -2$

Exercise 2.2E.56

$z - 16 = -59$

Exercise 2.2E.57

$m - 41 = -14$

Answer

$m = 27$

Exercise 2.2E.58

$0.04r = 52.60$

Exercise 2.2E.59

$63.90 = 0.03p$

Answer

$p = 2130$

Exercise 2.2E.60

$-15x = -120$

Exercise 2.2E.61

$84 = -12z$

Answer

$y = -7$

Exercise 2.2E.62

$19.36 = x - 0.2x$

Exercise 2.2E.63

$c - 0.3c = 35.70$

Answer

$c = 51$

Exercise 2.2E.64

$-y = -9$

Exercise 2.2E.65

$-x = -8$

Answer

$x = 8$

Translate to an Equation and Solve

In the following exercises, translate to an equation and then solve.

Exercise 2.2E.66

187 is the product of -17 and m

Exercise 2.2E.67

133 is the product of -19 and n

Answer

$133 = -19n; n = -7$

Exercise 2.2E.68

-184 is the product of 23 and p

Exercise 2.2E.69

-152 is the product of 8 and q

Answer

$-152 = 8q; q = -19$

Exercise 2.2E.70

u divided by 7 is equal to -49

Exercise 2.2E.71

r divided by 12 is equal to -48

Answer

$\frac{r}{12} = -48; r = -576$

Exercise 2.2E.72

h divided by -13 is equal to -65

Exercise 2.2E.73

j divided by -20 is equal to -80

Answer

$\frac{j}{-20} = -80; j = 1,600$

Exercise 2.2E.74

The quotient c and -19 is 38.

Exercise 2.2E.75

The quotient of b and -6 is 18

Answer

$\frac{b}{-6} = 18; b = -108$

Exercise 2.2E.76

The quotient of h and 26 is -52

Exercise 2.2E.77

The quotient k and 22 is -66

Answer

$\frac{k}{22} = -66; k = -1,452$

Exercise 2.2E.78

Five-sixths of y is 15

Exercise 2.2E.79

Three-tenths of x is 15

Answer

$\frac{3}{10}x = 15; x = 50$

Exercise 2.2E.80

Four-thirds of w is 36

Exercise 2.2E.81

Five-halves of v is 50

Answer

$\frac{5}{2}v = 50; v = 20$

Exercise 2.2E.82

The sum of nine-tenths and g is two-thirds.

Exercise 2.2E.83

The sum of two-fifths and f is one-half.

Answer

$\frac{2}{5} + f = \frac{1}{2}; f = \frac{1}{10}$

Exercise 2.2E.84

The difference of p and one-sixth is two-thirds.

Exercise 2.2E.85

The difference of q and one-eighth is three-fourths.

Answer

$q - \frac{1}{8} = \frac{3}{4}; q = \frac{7}{8}$

Translate and Solve Applications

In the following exercises, translate into an equation and solve.

Exercise 2.2E.86

Kindergarten Connie's kindergarten class has 24 children. She wants them to get into 4 equal groups. How many children will she put in each group?

Exercise 2.2E.87

Balloons Ramona bought 18 balloons for a party. She wants to make 3 equal bunches. How many balloons did she use in each bunch?

Answer

6 balloons

Exercise 2.2E.88

Tickets Mollie paid $36.25 for 5 movie tickets. What was the price of each ticket?

Exercise 2.2E.89

Shopping Serena paid $12.96 for a pack of 12 pairs of sport socks. What was the price of pair of sport socks?

Answer

$1.08

Exercise 2.2E.90

Sewing Nancy used 14 yards of fabric to make flags for one-third of the drill team. How much fabric, would Nancy need to make flags for the whole team?

Exercise 2.2E.91

MPG John's SUV gets 18 miles per gallon (mpg). This is half as many mpg as his wife's hybrid car. How many miles per gallon does the hybrid car get?

Answer

36 mpg

Exercise 2.2E.92

Height Aiden is 27 inches tall. He is $\frac{3}{8}$ as tall as his father. How tall is his father?

Exercise 2.2E.92

Real estate Bea earned $\$11,700$ commission for selling a house, calculated as $\frac{6}{100}$ of the selling price. What was the selling price of the house?

Answer

$195,000

Everyday Math

Exercise 2.2E.93

Commissions Every week Perry gets paid $\$150$ plus 12% of his total sales amount. Solve the equation $840 = 150 + 0.12(a - 1250)$ for a to find the total amount Perry must sell in order to be paid $\$840$ one week.

Exercise 2.2E.94

Stamps Travis bought $9.45 worth of 49-cent stamps and 21-cent stamps. The number of 21-cent stamps was 5 less than the number of 49-cent stamps. Solve the equation 0.49s+0.21(s−5)=9.45 for s, to find the number of 49-cent stamps Travis bought.

Answer

15 49-cent stamps

Writing Exercises

Exercise 2.2E.95

Frida started to solve the equation −3x=36 by adding 3 to both sides. Explain why Frida's method will not solve the equation.

Exercise 2.2E.96

Emiliano thinks $x = 40$ is the solution to the equation $\frac{1}{2}x = 80$. Explain why he is wrong.

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve equations using the Division and Multiplication Properties of equality.			
solve equations that require simplification.			
translate to an equation and solve.			
translate and solve applications.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 2.2E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.2E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.3: Solve Equations with Variables and Constants on Both Sides

Learning Objectives

By the end of this section, you will be able to:

- Solve an equation with constants on both sides
- Solve an equation with variables on both sides
- Solve an equation with variables and constants on both sides

Note

Before you get started, take this readiness quiz.

1. Simplify: 4y−9+9.
   If you missed this problem, review Exercise 1.10.20.

Solve Equations with Constants on Both Sides

In all the equations we have solved so far, all the variable terms were on only one side of the equation with the constants on the other side. This does not happen all the time—so now we will learn to solve equations in which the variable terms, or constant terms, or both are on both sides of the equation.

Our strategy will involve choosing one side of the equation to be the "variable side", and the other side of the equation to be the "constant side." Then, we will use the Subtraction and Addition Properties of Equality to get all the variable terms together on one side of the equation and the constant terms together on the other side.

By doing this, we will transform the equation that began with variables and constants on both sides into the form $ax = b$. We already know how to solve equations of this form by using the Division or Multiplication Properties of Equality.

Example 2.3.1

Solve: $7x + 8 = -13$.

Solution

In this equation, the variable is found only on the left side. It makes sense to call the left side the "variable" side. Therefore, the right side will be the "constant" side. We will write the labels above the equation to help us remember what goes where.

$$\overset{\text{variable}}{7x} + 8 = \overset{\text{constant}}{-13}$$

Since the left side is the "xx", or variable side, the 8 is out of place. We must "undo" adding 8 by subtracting 8, and to keep the equality we must subtract 8 from both sides.

	$\overset{\text{variable}}{7x} + 8 = \overset{\text{constant}}{-13}$
Use the Subtraction Property of Equality.	$7x + 8 - 8 = -13 - 8$
Simplify.	$7x = -21$
Now all the variables are on the left and the constant on the right. The equation looks like those you learned to solve earlier.	
Use the Division Property of Equality.	$\frac{7x}{7} = \frac{-21}{7}$
Simplify.	$x = -3$
Check:	$7x + 8 = -13$
Let x=−3.	$7(-3) + 8 \stackrel{?}{=} -13$

	$-21+8\stackrel{?}{=}-13$ $-13=-13\checkmark$	

Try It 2.3.2

Solve: $3x+4=-8$.

Answer

$x=-4$

Try It 2.3.3

Solve: $5a+3=-37$.

Answer

$a=-8$

Example 2.3.4

Solve: $8y-9=31$.

Solution

Notice, the variable is only on the left side of the equation, so we will call this side the "variable" side, and the right side will be the "constant" side. Since the left side is the "variable" side, the 9 is out of place. It is subtracted from the 8y, so to "undo" subtraction, add 9 to both sides. Remember, whatever you do to the left, you must do to the right.

	variable constant $8y-9=31$
Add 9 to both sides.	$8y-9=31$ $+9\quad+9$
Simplify.	$8y=40$
	The variable is now on one side and the constants on the other. We continue from here as we did earlier.
Divide both sides by 8.	$\frac{8y}{8}=\frac{40}{8}$
Simplify.	$y=5$
Check:	$8y-9=31$
Let y=5.	$8\cdot5-9\stackrel{?}{=}31$
	$40-9\stackrel{?}{=}31$
	$31=31\checkmark$

Try It 2.3.5

Solve: $5y-9=16$.

Answer

$y = 5$

Try It 2.3.6

Solve: $3m - 8 = 19$.

Answer

$m = 9$

Solve Equations with Variables and Constants on Both Sides

The next example will be the first to have variables and constants on both sides of the equation. It may take several steps to solve this equation, so we need a clear and organized strategy.

Example 2.3.7

Solve: $9x = 8x - 6$.

Solution

Here the variable is on both sides, but the constants only appear on the right side, so let's make the right side the "constant" side. Then the left side will be the "variable" side.

	$9x = 8x - 6$
We don't want any x's on the right, so subtract the 8x from both sides.	$9x - 8x = 8x - 8x - 6$
Simplify.	$x = -6$
We succeeded in getting the variables on one side and the constants on the other, and have obtained the solution.	
Check:	$9x = 8x - 6$
Let x=−6.	$9(-6) \stackrel{?}{=} 8(-6) - 6$
	$-54 \stackrel{?}{=} -48 - 6$
	$-54 = -54$ ✓

Try It 2.3.8

Solve: $6n = 5n - 10$.

Answer

$n = -10$

Try It 2.3.9

Solve: $-6c = -7c - 1$

Answer

$c = -1$

Example 2.3.10

Solve: $5y - 9 = 8y$

Solution

The only constant is on the left and the y's are on both sides. Let's leave the constant on the left and get the variables to the right.

	constant variable $5y - 9 = 8y$
Subtract 5y from both sides.	$5y - 5y - 9 = 8y - 5y$
Simplify.	$-9 = 3y$
We have the y's on the right and the constants on the left. Divide both sides by 3.	$\frac{-9}{3} = \frac{3y}{3}$
Simplify.	$-3 = y$
Check:	$5y - 9 = 8y$
Let $y = -3$.	$5(-3) - 9 \stackrel{?}{=} 8(-3)$
	$-15 - 9 \stackrel{?}{=} -24$
	$-24 = -24$ ✓

Try It 2.3.11

Solve: $3p - 14 = 5p$.

Answer

$p = -7$

Try It 2.3.12

Solve: $8m + 9 = 5m$

Answer

$m = -3$

Example 2.3.13

Solve: $12x = -x + 26$

Solution

The only constant is on the right, so let the left side be the "variable" side.

	variable constant $12x = -x + 26$
Remove the $-x$ from the right side by adding x to both sides.	$12x + x = -x + x + 26$
Simplify.	$13x = 26$
All the x's are on the left and the constants are on the right. Divide both sides by 13.	$\frac{13x}{13} = \frac{26}{13}$

Simplify.	$x = 2$

Try It 2.3.14

Solve: $12j = -4j + 32$

Answer

$j = 2$

Try It 2.3.15

Solve: $8h = -4h + 12$

Answer

$h = 1$

Solve Equations with Variables and Constants on Both Sides

The next example will be the first to have variables and constants on both sides of the equation. It may take several steps to solve this equation, so we need a clear and organized strategy.

Example 2.3.16: How to Solve Equations with Variables and Constants on Both Sides

Solve: $7x + 5 = 6x + 2$

Solution

Step 1. Choose which side will be the "variable" side—the other side will be the "constant" side.	The variable terms are $7x$ and $6x$. Since 7 is greater than 6, we will make the left side the "x" side. The right side will be the "constant" side.	variable constant $7x + 5 = 6x + 2$
Step 2. Collect the variable terms to the "variable" side of the equation, using the addition or subtraction property of equality.	With the right side as the "constant" side, the $6x$ is out of place, so subtract $6x$ from both sides. Combine like terms. Now, the variable is only on the left side!	$7x - 6x + 5 = 6x - 6x + 2$ $x + 5 = 2$
Step 3. Collect all the constants to the other side of the equation, using the addition or subtraction property of equality.	The right side is the "constant" side, so the 5 is out of place. Subtract 5 from both sides. Simplify.	$x + 5 - 5 = 2 - 5$ $x = -3$

Step 4. Make the coefficient of the variable equal 1, using the multiplication or division property of equality.	The coefficient of x is one. The equation is solved.	
Step 5. Check.	Let $x = -3$ Simplify. Add.	**Check:** $7x + 6 = 6x + 2$ $(-3) + 5 = 6(-3) + 2$ $-21 + 5 = -18 + 2$ $-16 = -16$ ✓

? Try It 2.3.17

Solve: $12x + 8 = 6x + 2$.

Answer

$x = -1$

? Try It 2.3.18

Solve: $9y + 4 = 7y + 12$.

Answer

$y = 4$

We'll list the steps below so you can easily refer to them. But we'll call this the 'Beginning Strategy' because we'll be adding some steps later in this chapter.

BEGINNING STRATEGY FOR SOLVING EQUATIONS WITH VARIABLES AND CONSTANTS ON BOTH SIDES OF THE EQUATION.

1. Choose which side will be the "variable" side—the other side will be the "constant" side.
2. Collect the variable terms to the "variable" side of the equation, using the Addition or Subtraction Property of Equality.
3. Collect all the constants to the other side of the equation, using the Addition or Subtraction Property of Equality.
4. Make the coefficient of the variable equal 1, using the Multiplication or Division Property of Equality.
5. Check the solution by substituting it into the original equation.

In Step 1, a helpful approach is to make the "variable" side the side that has the variable with the larger coefficient. This usually makes the arithmetic easier.

✓ Example 2.3.19

Solve: $8n - 4 = -2n + 6$.

Solution

In the first step, choose the variable side by comparing the coefficients of the variables on each side.

Since $8 > -2$, make the left side the "variable" side.	variable constant $8n - 4 = -2n + 6$
We don't want variable terms on the right side—add 2n to both sides to leave only constants on the right.	$8n + 2n - 4 = -2n + 2n + 6$
Combine like terms.	$10n - 4 = 6$

We don't want any constants on the left side, so add 4 to both sides.	$10n-4+4=6+4$
Simplify.	$10n=10$
The variable term is on the left and the constant term is on the right. To get the coefficient of nn to be one, divide both sides by 10.	$\frac{10n}{10}=\frac{10}{10}$
Simplify.	$n=1$
Check:	$8n-4=-2n+6$
Let n=1.	$8\cdot 1-4\stackrel{?}{=}-2\cdot 1+6$
	$8-4\stackrel{?}{=}-2+6$
	$4=4\checkmark$

Try It 2.3.20

Solve: $8q-5=-4q+7$

Answer

$q=1$

Try It 2.3.21

Solve: $7n-3=n+3$

Answer

$n=1$

Example 2.3.22

Solve: $7a-3=13a+7$

Solution

In the first step, choose the variable side by comparing the coefficients of the variables on each side.

Since 13>7, make the right side the "variable" side and the left side the "constant" side.

	constant variable $7a-3=13a+7$
Subtract 7a from both sides to remove the variable term from the left.	$7a-7a-3=13a-7a+7$
Combine like terms.	$-3=6a+7$
Subtract 7 from both sides to remove the constant from the right.	$-3-7=6a+7-7$
Simplify.	$-10=6a$
Divide both sides by 6 to make 1 the coefficient of a.	$\frac{-10}{6}=\frac{6a}{6}$
Simplify.	$-\frac{5}{3}=a$
Check:	$7a-3=13a+7$
Let $a=-\frac{5}{3}$.	$7\left(-\frac{5}{3}\right)-3\stackrel{?}{=}13\left(-\frac{5}{3}\right)+7$

	$-\frac{35}{3}-\frac{9}{3}\stackrel{?}{=}-\frac{65}{3}+\frac{21}{3}$	
	$-\frac{54}{3}=-\frac{54}{3}\checkmark$	

Try It 2.3.23

Solve: $2a-2=6a+18$

Answer

$a=-5$

Try It 2.3.24

Solve: $4k-1=7k+17$

Answer

$k=-6$

In the last example, we could have made the left side the "variable" side, but it would have led to a negative coefficient on the variable term. (Try it!) While we could work with the negative, there is less chance of errors when working with positives. The strategy outlined above helps avoid the negatives!

To solve an equation with fractions, we just follow the steps of our strategy to get the solution!

Example 2.3.25

Solve: $\frac{4}{5}x+6=\frac{1}{4}x-2$

Solution

Since $\frac{5}{4}>\frac{1}{4}$, make the left side the "variable" side and the right side the "constant" side.

	variable constant $\frac{5}{4}x+6=\frac{1}{4}x-2$
Subtract $\frac{1}{4}x$ from both sides.	$\frac{5}{4}x-\frac{1}{4}x+6=\frac{1}{4}x-\frac{1}{4}x-2$
Combine like terms.	$x+6=-2$
Subtract 6 from both sides.	$x+6-6=-2-6$
Simplify.	$x=-8$
Check: Let $x=-8$	$\frac{5}{4}x+6 = \frac{1}{4}x-2$ $\frac{5}{4}(-8)+6 \stackrel{?}{=} \frac{1}{4}(-8)-2$ $-10+6 \stackrel{?}{=} -2-2$ $-4 = -4\checkmark$

Try It 2.3.26

Solve: $\frac{7}{8}x-12=-\frac{1}{8}x-2$

Answer

$x = 10$

Try It 2.3.27

Solve: $\frac{7}{6}x + 11 = \frac{1}{6}y + 8$

Answer

$y = -3$

We will use the same strategy to find the solution for an equation with decimals.

Example 2.3.28

Solve: $7.8x + 4 = 5.4x - 8$.

Solution

Since $7.8 > 5.4$, make the left side the "variable" side and the right side the "constant" side.

	variable side constant side $7.8x + 4 = 5.4x - 8$
Subtract 5.4x from both sides.	$7.8x - 5.4x + 4 = 5.4x - 5.4x - 8$
Combine like terms.	$2.4x + 4 = -8$
Subtract 4 from both sides.	$2.4x + 4 - 4 = -8 - 4$
Simplify.	$2.4x = -12$
Use the Division Property of Equality.	$\frac{2.4x}{2.4} = \frac{-12}{2.4}$
Simplify.	$x = -5$
Check:	$7.8x + 4 = 5.4x - 5$
Let $x = -5$	$7.8(-5) + 4 = 5.4(-5) - 8$
	$-39 + 4 \stackrel{?}{=} -27 - 8$
	$-35 = -35$ ✓

Try It 2.3.29

Solve: $2.8x + 12 = -1.4x - 9$

Answer

$x = -5$

Try It 2.3.30

Solve: $3.6y + 8 = 1.2y - 4$

Answer

$y = -5$

Key Concepts

- **Beginning Strategy for Solving an Equation with Variables and Constants on Both Sides of the Equation**
  1. Choose which side will be the “variable” side—the other side will be the “constant” side.
  2. Collect the variable terms to the “variable” side of the equation, using the Addition or Subtraction Property of Equality.
  3. Collect all the constants to the other side of the equation, using the Addition or Subtraction Property of Equality.
  4. Make the coefficient of the variable equal 1, using the Multiplication or Division Property of Equality.
  5. Check the solution by substituting it into the original equation.

This page titled 2.3: Solve Equations with Variables and Constants on Both Sides is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.3: Solve Equations with Variables and Constants on Both Sides** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.3E: Exercises

Practice Makes Perfect

Solve Equations with Constants on Both Sides

In the following exercises, solve the following equations with constants on both sides.

Exercise 2.3E.1

$9x - 3 = 60$

Exercise 2.3E.2

$12x - 8 = 64$

Answer

$x = 6$

Exercise 2.3E.3

$14w + 5 = 117$

Exercise 2.3E.4

$15y + 7 = 97$

Answer

$y = 6$

Exercise 2.3E.5

$2a + 8 = -28$

Exercise 2.3E.6

$3m + 9 = -15$

Answer

$m = -8$

Exercise 2.3E.7

$-62 = 8n - 6$

Exercise 2.3E.8

$-77 = 9b - 5$

Answer

$b = -8$

Exercise 2.3E.9

$35 = -13y + 9$

Exercise 2.3E.10

$60 = -21x - 24$

Answer

$x = -4$

Exercise 2.3E.11

$-12p - 9 = 9$

Exercise 2.3E.12

$-14q - 2 = 16$

Answer

$q = -\frac{9}{7}$

Solve Equations with Variables on Both Sides

In the following exercises, solve the following equations with variables on both sides.

Exercise 2.3E.13

$19z = 18z - 7$

Exercise 2.3E.14

$21k = 20k - 11$

Answer

$k = -11$

Exercise 2.3E.15

$9x + 36 = 15x$

Exercise 2.3E.16

$8x + 27 = 11x$

Answer

$x = 9$

Exercise 2.3E.17

$c = -3c - 20$

Exercise 2.3E.18

$b = -4b - 15$

Answer

$b = -3$

Exercise 2.3E.19

$9q = 44 - 2q$

Exercise 2.3E.20

$5z = 39 - 8z$

Answer

$z = 3$

Exercise 2.3E.21

$6y + \frac{1}{2} = 5y$

Exercise 2.3E.22

$4x + \frac{3}{4} = 3x$

Answer

$x = -\frac{3}{4}$

Exercise 2.3E.23

$-18a - 8 = -22a$

Exercise 2.3E.24

$-11r - 8 = -7r$

Answer

$r = -2$

Solve Equations with Variables and Constants on Both Sides

In the following exercises, solve the following equations with variables and constants on both sides.

Exercise 2.3E.25

$8x - 15 = 7x + 3$

Exercise 2.3E.26

$6x - 17 = 5x + 2$

Answer

$x = 19$

Exercise 2.3E.27

$26 + 13d = 14d + 11$

Exercise 2.3E.28

$21 + 18f = 19f + 14$

Answer

$f = 7$

Exercise 2.3E.29

$2p - 1 = 4p - 33$

Exercise 2.3E.30

$12q - 5 = 9q - 20$

Answer

$q = -5$

Exercise 2.3E.31

$4a + 5 = -a - 40$

Exercise 2.3E.32

$8c + 7 = -3c - 37$

Answer

$c = -4$

Exercise 2.3E.33

$5y - 30 = -5y + 30$

Exercise 2.3E.34

$7x - 17 = -8x + 13$

Answer

$x = 2$

Exercise 2.3E.35

$7s + 12 = 5 + 4s$

Exercise 2.3E.36

$9p+14=6+4p$

Answer

$p=-\frac{8}{5}$

Exercise 2.3E.37

$2z-6=23-z$

Exercise 2.3E.38

$3y-4=12-y$

Answer

$y=4$

Exercise 2.3E.39

$\frac{5}{3}c-3=\frac{2}{3}c-16$

Exercise 2.3E.40

$\frac{7}{4}m-7=\frac{3}{4}m-13$

Answer

$m=-6$

Exercise 2.3E.41

$8-\frac{2}{5}q=\frac{3}{5}q+6$

Exercise 2.3E.42

$11-\frac{1}{5}a=\frac{4}{5}a+4$

Answer

$a=7$

Exercise 2.3E.43

$\frac{4}{3}n+9=\frac{1}{3}n-9$

Exercise 2.3E.44

$\frac{5}{4}a+15=\frac{3}{4}a-5$

Answer

$a=-40$

Exercise 2.3E.45

$\frac{1}{4}y+7=\frac{3}{4}y-3$

Exercise 2.3E.46

$\frac{3}{5}p+2=\frac{4}{5}p-1$

Answer

$p=15$

Exercise 2.3E.47

$14n+8.25=9n+19.60$

Exercise 2.3E.48

$13z+6.45=8z+23.75$

Answer

$z=3.46$

Exercise 2.3E.49

$2.4w-100=0.8w+28$

Exercise 2.3E.50

$2.7w-80=1.2w+10$

Answer

$w=60$

Exercise 2.3E.51

$5.6r+13.1=3.5r+57.2$

Exercise 2.3E.52

$6.6x-18.9=3.4x+54.7$

Answer

$x=23$

Everyday Math

Exercise 2.3E.53

Concert tickets At a school concert the total value of tickets sold was $1506. Student tickets sold for $6 and adult tickets sold for $9. The number of adult tickets sold was 5 less than 3 times the number of student tickets. Find the number of student tickets sold, *s*, by solving the equation 6s+27s−45=1506. Add exercises text here.

Exercise $2.3E.54$

Making a fence Jovani has 150 feet of fencing to make a rectangular garden in his backyard. He wants the length to be 15 feet more than the width. Find the width, w, by solving the equation $150 = 2w + 30 + 2w$.

Answer

30 feet

Writing Exercises

Exercise $2.3E.55$

Solve the equation $\frac{6}{5}y - 8 = \frac{1}{5}y + 7$ explaining all the steps of your solution as in the examples in this section.

Exercise $2.3E.56$

Solve the equation $10x + 14 = -2x + 38$ explaining all the steps of your solution as in this section.

Answer

$x = 2$ Justifications will vary.

Exercise $2.3E.57$

When solving an equation with variables on both sides, why is it usually better to choose the side with the larger coefficient of x to be the "variable" side?

Exercise $2.3E.58$

Is $x = -2$ a solution to the equation $5 - 2x = -4x + 1$? How do you know?

Answer

Yes. Justifications will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve an equation with constants on both sides.			
solve an equation with variables on both sides.			
solve an equation with variables and constants on both sides.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 2.3E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.3E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.4: Use a General Strategy to Solve Linear Equations

Learning Objectives

By the end of this section, you will be able to:

- Solve equations using a general strategy
- Classify equations

Note

Before you get started, take this readiness quiz.

1. Simplify: $-(a-4)$.
   If you missed this problem, review Exercise 1.10.46
2. Multiply: $\frac{3}{2}(12x+20)$
   If you missed this problem, review Exercise 1.10.34.
3. Simplify: $5-2(n+1)$
   If you missed this problem, review Exercise 1.10.49.
4. Multiply: $3(7y+9)$
   If you missed this problem, review Exercise 1.10.34.
5. Multiply: $(2.5)(6.4)$
   If you missed this problem, review Exercise 1.8.19.

Solve Equations Using the General Strategy

Until now we have dealt with solving one specific form of a linear equation. It is time now to lay out one overall strategy that can be used to solve any linear equation. Some equations we solve will not require all these steps to solve, but many will.

Beginning by simplifying each side of the equation makes the remaining steps easier.

Example 2.4.1: How to Solve Linear Equations Using the General Strategy

Solve: $-6(x+3)=24$.

Solution

Step 1. Simplify each side of the equation as much as possible.	Use the Distributive Property. Notice that each side of the equation is simplified as much as possible.	$-6(x+3)=24$ $-6x-18=24$
Step 2. Collect all variable terms on one side of the equation.	Nothing to do – all x's are on the left side.	
Step 3. Collect constant terms on the other side of the equation.	To get constants only on the right, add 18 to each side. Simplify.	$-6x-18+18=24+18$ $-6x=42$

Step 4. Make the coefficient of the variable term to equal to 1.	Divide each side by –6.	$\frac{-6x}{-6} = \frac{42}{-6}$
	Simplify.	$x = -7$
Step 5. Check the solution.	Let $x = -7$	**Check:** $-6(x + 3) = 24$
	Simplify.	$-6(-7 + 3) \stackrel{?}{=} 24$
	Multiply.	$-6(-4) \stackrel{?}{=} 24$
		$24 = 24$ ✓

Try It 2.4.2

Solve: $5(x+3) = 35$

Answer

$x = 4$

Try It 2.4.3

Solve: $6(y-4) = -18$

Answer

$y = 1$

GENERAL STRATEGY FOR SOLVING LINEAR EQUATIONS.

1. **Simplify each side of the equation as much as possible.**
   Use the Distributive Property to remove any parentheses.
   Combine like terms.
2. **Collect all the variable terms on one side of the equation.**
   Use the Addition or Subtraction Property of Equality.
3. **Collect all the constant terms on the other side of the equation.**
   Use the Addition or Subtraction Property of Equality.
4. **Make the coefficient of the variable term to equal to 1.**
   Use the Multiplication or Division Property of Equality.
   State the solution to the equation.
5. **Check the solution.** Substitute the solution into the original equation to make sure the result is a true statement.

Example 2.4.4

Solve: $-(y+9) = 8$

Solution

	$-(y + 9) = 8$
Simplify each side of the equation as much as possible by distributing.	$-y - 9 = 8$
The only y term is on the left side, so all variable terms are on the left side of the equation.	
Add 9 to both sides to get all constant terms on the right side of the equation.	$-y - 9 + 9 = 8 + 9$
Simplify.	$-y = 17$

Rewrite $-y$ as $-1y$.		$-1y = 17$
Make the coefficient of the variable term to equal to 1 by dividing both sides by -1.		$\frac{-1y}{-1} = \frac{17}{-1}$
Simplify.		$y = -17$
Check:	$-(y+9) = 8$	
Let y=−17.	$-(-17+9) \stackrel{?}{=} 8$	
	$-(-8) \stackrel{?}{=} 8$	
	$8 = 8$ ✓	

Try It 2.4.5

Solve: $-(y+8) = -2$

Answer

$y = -6$

Try It 2.4.6

Solve: $-(z+4) = -12$

Answer

$z = 8$

Example 2.4.7

Solve: $5(a-3)+5 = -10$

Solution

		$5(a-3)+5 = -10$
Simplify each side of the equation as much as possible.		
Distribute.		$5a-15+5 = -10$
Combine like terms.		$5a-10 = -10$
The only a term is on the left side, so all variable terms are on one side of the equation.		
Add 10 to both sides to get all constant terms on the other side of the equation.		$5a-10+10 = -10+10$
Simplify.		$5a = 0$
Make the coefficient of the variable term to equal to 11 by dividing both sides by 55.		$\frac{5a}{5} = \frac{0}{5}$
Simplify.		$a = 0$
Check:	$5(a-3)+5 = -10$	
Let a=0.	$5(0-3)+5 \stackrel{?}{=} -10$	
	$5(-3)+5 \stackrel{?}{=} -10$	
	$-15+5 \stackrel{?}{=} -10$	

	$-10 = -10$ ✓

? Try It 2.4.8

Solve: $2(m-4)+3=-1$

Answer

$m=2$

? Try It 2.4.9

Solve:$7(n-3)-8=-15$

Answer

$n=2$

✓ Example 2.4.10

Solve: $\frac{2}{3}(6m-3)=8-m$

Solution

	$\frac{2}{3}(6m-3)=8-m$
Distribute.	$4m-2=8-m$
Add m to get the variables only to the left.	$4m+m-2=8-m+m$
Simplify.	$5m-2=8$
Add 2 to get constants only on the right.	$5m-2+2=8+2$
Simplify.	$5m=10$
Divide by 5.	$\frac{5m}{5}=\frac{10}{5}$
Simplify.	$m=2$
Check:	$\frac{2}{3}(6m-3)=8-m$
Let m=2.	$\frac{2}{3}(6\cdot 2-3)\stackrel{?}{=}8-2$
	$\frac{2}{3}(12-3)\stackrel{?}{=}6$
	$\frac{2}{3}(9)\stackrel{?}{=}6$
	$6=6$ ✓

? Try It 2.4.11

Solve: $\frac{1}{3}(6u+3)=7-u$

Answer

$u=2$

Try It 2.4.12

Solve: $\frac{2}{3}(9x-12)=8+2x$

Answer

$x=4$

Example 2.4.13

Solve: $8-2(3y+5)=0$

Solution

	$8-2(3y+5)=0$
Simplify—use the Distributive Property.	$8-6y-10=0$
Combine like terms.	$-6y-2=0$
Add 2 to both sides to collect constants on the right.	$-6y-2+2=0+2$
Simplify.	$-6y=2$
Divide both sides by -6−6.	$\frac{-6y}{-6}=\frac{2}{-6}$
Simplify.	$y=-\frac{1}{3}$
Check: Let y=−13. $8-2(3y+5)=0$ $8-2\left[3\left(-\frac{1}{3}\right)+5\right]=0$ $8-2(-1+5)\stackrel{?}{=}0$ $8-2(4)\stackrel{?}{=}0$ $8-8\stackrel{?}{=}0$ $0=0$ ✓	

Try It 2.4.14

Solve: $12-3(4j+3)=-17$

Answer

$j=\frac{5}{3}$

Try It 2.4.15

Solve: $-6-8(k-2)=-10$

Answer

$k=\frac{5}{2}$

Example 2.4.16

Solve: $4(x-1)-2=5(2x+3)+6$

Solution

	$4(x-1)-2=5(2x+3)+6$
Distribute.	$4x-4-2=10x+15+6$
Combine like terms.	$4x-6=10x+21$
Subtract 4x to get the variables only on the right side since $10>4$.	$4x-4x-6=10x-4x+21$
Simplify.	$-6=6x+21$
Subtract 21 to get the constants on left.	$-6-21=6x+21-21$
Simplify.	$-27=6x$
Divide by 6.	$\frac{-27}{6}=\frac{6x}{6}$
Simplify.	$-\frac{9}{2}=x$
Check:	$4(x-1)-2=5(2x+3)+6$
Let $x=-\frac{9}{2}$.	$4\left(-\frac{9}{2}-1\right)-2\stackrel{?}{=}5\left[2\left(-\frac{9}{2}\right)+3\right]+6$
	$4\left(-\frac{11}{2}\right)-2\stackrel{?}{=}5(-9+3)+6$
	$-22-2\stackrel{?}{=}5(-6)+6$
	$-24\stackrel{?}{=}-30+6$
	$-24=-24$ ✓

Try It 2.4.17

Solve: $6(p-3)-7=5(4p+3)-12$

Answer

$p=-2$

Try It 2.4.18

Solve: $8(q+1)-5=3(2q-4)-1$

Answer

$q=-8$

Example 2.4.19

Solve: $10[3-8(2s-5)]=15(40-5s)$

Solution

	$10[3-8(2s-5)]=15(40-5s)$

Simplify from the innermost parentheses first.		$10[3-16s+40]=15(40-5s)$
Combine like terms in the brackets.		$10[43-16s]=15(40-5s)$
Distribute.		$430-160s=600-75s$
Add 160s to get the s's to the right.		$430-160s+160s=600-75s+160s$
Simplify.		$430=600+85s$
Subtract 600 to get the constants to the left.		$430-600=600+85s-600$
Simplify.		$-170=85s$
Divide.		$\frac{-170}{85}=\frac{85s}{85}$
Simplify.		$-2=s$
Check:	$10[3-8(2s-5)]=15(40-5s)$	
Substitute s=−2.	$10[3-8(2(-2)-5)] \stackrel{?}{=} 15(40-5(-2))$	
	$10[3-8(-4-5)] \stackrel{?}{=} 15(40+10)$	
	$10[3-8(-9)] \stackrel{?}{=} 15(50)$	
	$10[3+72] \stackrel{?}{=} 750$	
	$10[75] \stackrel{?}{=} 750$	
	$750=750$ ✓	

? Try It 2.4.20

Solve: $6[4-2(7y-1)]=8(13-8y)$.

Answer

$y=-\frac{17}{5}$

? Try It 2.4.21

Solve: $12[1-5(4z-1)]=3(24+11z)$.

Answer

$z=0$

✓ Example 2.4.22

Solve: $0.36(100n+5)=0.6(30n+15)$.

Solution

	$0.36(100n+5)=0.6(30n+15)$
Distribute.	$36n+1.8=18n+9$
Subtract 18n to get the variables to the left.	$36n-18n+1.8=18n-18n+9$
Simplify.	$18n+1.8=9$

Subtract 1.8 to get the constants to the right.		$18n + 1.8 - 1.8 = 9 - 1.8$
Simplify.		$18n = 7.2$
Divide.		$\frac{18n}{18} = \frac{7.2}{18}$
Simplify.		$n = 0.4$
Check:	$0.36(100n + 5) = 0.6(30n + 15)$	
Let n=0.4.	$0.36(100(0.4) + 5) \stackrel{?}{=} 0.6(30(0.4) + 15)$	
	$0.36(40 + 5) \stackrel{?}{=} 0.6(12 + 15)$	
	$0.36(45) \stackrel{?}{=} 0.6(27)$	
	$16.2 = 16.2$ ✓	

? Try It 2.4.23

Solve: $0.55(100n+8) = 0.6(85n+14)$.

Answer

$n = 1$

? Try It 2.4.24

Solve: $0.15(40m-120) = 0.5(60m+12)$.

Answer

$m = -1$

Classify Equations

Consider the equation we solved at the start of the last section, 7x+8=−13. The solution we found was x=−3. This means the equation 7x+8=−13 is true when we replace the variable, *x*, with the value −3. We showed this when we checked the solution x=−3 and evaluated 7x+8=−13 for x=−3.

$$7(-3) + 8 \stackrel{?}{=} -13$$
$$-21 + 8 \stackrel{?}{=} -13$$
$$-13 = -13 \checkmark$$

If we evaluate 7x+8 for a different value of *x*, the left side will not be −13.

The equation 7x+8=−13 is true when we replace the variable, *x*, with the value −3, but not true when we replace *x* with any other value. Whether or not the equation 7x+8=−13 is true depends on the value of the variable. Equations like this are called conditional equations.

All the equations we have solved so far are conditional equations.

CONDITIONAL EQUATION

An equation that is true for one or more values of the variable and false for all other values of the variable is a **conditional equation**.

Now let's consider the equation 2y+6=2(y+3). Do you recognize that the left side and the right side are equivalent? Let's see what happens when we solve for *y*.

	$2y+6=2(y+3)$
Distribute.	$2y+6=2y+6$
Subtract 2y to get the y's to one side.	$2y-2y+6=2y-2y+6$
Simplify—the y's are gone!	$6=6$

But 6=6 is true.

This means that the equation 2y+6=2(y+3) is true for any value of *y*. We say the solution to the equation is all of the real numbers. An equation that is true for any value of the variable like this is called an **identity**.

IDENTITY

An equation that is true for any value of the variable is called an **identity**.

The solution of an identity is all real numbers.

What happens when we solve the equation 5z=5z−1?

	$5z=5z-1$
Subtract 5z to get the constant alone on the right.	$5z-5z=5z-5z-1$
Simplify—the z's are gone!	$0\neq -1$

But $0 \neq -1$.

Solving the equation 5z=5z−1 led to the false statement 0=−1. The equation 5z=5z−1 will not be true for any value of z. It has no solution. An equation that has no solution, or that is false for all values of the variable, is called a **contradiction**.

CONTRADICTION

An equation that is false for all values of the variable is called a **contradiction**.

A contradiction has no solution.

Example 2.4.25

Classify the equation as a conditional equation, an identity, or a contradiction. Then state the solution.

$6(2n-1)+3=2n-8+5(2n+1)$

Solution

$6(2n-1)+3=2n-8+5(2n+1)$	
Distribute.	$12n-6+3=2n-8+10n+5$
Combine like terms.	$12n-3=12n-3$
Subtract 12n to get the nn's to one side.	$12n-12n-3=12n-12n-3$
Simplify.	$-3=-3$
This is a true statement.	The equation is an identity. The solution is all real numbers.

Try It 2.4.26

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution:

$4+9(3x-7)=-42x-13+23(3x-2)$

Answer

identity; all real numbers

Try It 2.4.27

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution:

$8(1-3x)+15(2x+7)=2(x+50)+4(x+3)+1$

Answer

identity; all real numbers

Example 2.4.28

Classify as a conditional equation, an identity, or a contradiction. Then state the solution.

$10+4(p-5)=0$

Solution

	$10+4(p-5)=0$
Distribute.	$10+4p-20=0$
Combine like terms.	$4p-10=0$
Add 10 to both sides.	$4p-10+10=0+10$
Simplify.	$4p=10$
Divide.	$\frac{4p}{4}=\frac{10}{4}$
Simplify.	$p=\frac{5}{2}$
The equation is true when $p=frac52$.	This is a conditional equation. The solution is $p=frac52$.

Try It 2.4.29

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution: $11(q+3)-5=19$

Answer

conditional equation; \(q = \frac{9}{11}\

Try It 2.4.30

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution: $6+14(k-8)=95$

Answer

conditional equation; $k = \frac{193}{14}$

Example 2.4.31

Classify the equation as a conditional equation, an identity, or a contradiction. Then state the solution.

$5m + 3(9 + 3m) = 2(7m - 11)$

Solution

	$5m + 3(9 + 3m) = 2(7m - 11)$
Distribute.	$5m + 27 + 9m = 14m - 22$
Combine like terms.	$14m + 27 = 14m - 22$
Subtract 14m from both sides.	$14m + 27 - 14m = 14m - 22 - 14m$
Simplify.	$27 \neq -22$
But $27 \neq -22$.	The equation is a contradiction. It has no solution.

Try It 2.4.32

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution:

$12c + 5(5 + 3c) = 3(9c - 4)$

Answer

contradiction; no solution

Try It 2.4.33

Classify the equation as a conditional equation, an identity, or a contradiction and then state the solution:

$4(7d + 18) = 13(3d - 2) - 11d$

Answer

contradiction; no solution

Type of equation	What happens when you solve it?	Solution
Conditional Equation	True for one or more values of the variables and false for all other values	One or more values
Identity	**True** for any value of the variable	All real numbers
Contradiction	**False** for all values of the variable	No solution

Key Concepts

- **General Strategy for Solving Linear Equations**
  1. Simplify each side of the equation as much as possible.
     Use the Distributive Property to remove any parentheses.
     Combine like terms.

2. Collect all the variable terms on one side of the equation.
   Use the Addition or Subtraction Property of Equality.
3. Collect all the constant terms on the other side of the equation.
   Use the Addition or Subtraction Property of Equality.
4. Make the coefficient of the variable term to equal to 1.
   Use the Multiplication or Division Property of Equality.
   State the solution to the equation.
5. Check the solution.
   Substitute the solution into the original equation.

This page titled 2.4: Use a General Strategy to Solve Linear Equations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.4: Use a General Strategy to Solve Linear Equations** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.4E: Exercises

Practice Makes Perfect

Solve Equations Using the General Strategy for Solving Linear Equations

In the following exercises, solve each linear equation.

Exercise 2.4E.1

$15(y-9)=-60$

Exercise 2.4E.2

$21(y-5)=-42$

Answer

$y=3$

Exercise 2.4E.3

$-9(2n+1)=36$

Exercise 2.4E.4

$-16(3n+4)=32$

Answer

$n=-2$

Exercise 2.4E.5

$8(22+11r)=0$

Exercise 2.4E.6

$5(8+6p)=0$

Answer

$p=-\frac{4}{3}$

Exercise 2.4E.7

$-(w-12)=30$

Exercise 2.4E.8

$-(t-19)=28$

Answer

$t=-9$

Exercise 2.4E.9

$9(6a+8)+9=81$

Exercise 2.4E.10

$8(9b-4)-12=100$

Answer

$b=2$

Exercise 2.4E.11

$32+3(z+4)=41$

Exercise 2.4E.12

$21+2(m-4)=25$

Answer

$m=6$

Exercise 2.4E.13

$51+5(4-q)=56$

Exercise 2.4E.14

$-6+6(5-k)=15$

Answer

$k=\frac{3}{2}$

Exercise 2.4E.15

$2(9s-6)-62=16$

Exercise 2.4E.16

$8(6t-5)-35=-27$

Answer

$t=1$

Exercise 2.4E.17

$3(10-2x)+54=0$

Exercise 2.4E.18

$-2(11-7x)+54=4$

Answer

$x=-2$

Exercise 2.4E.19

$\frac{2}{3}(9c-3)=22$

Exercise 2.4E.20

$\frac{3}{5}(10x-5)=27$

Answer

$x=5$

Exercise 2.4E.21

$\frac{1}{5}(15c+10)=c+7$

Exercise 2.4E.22

$\frac{1}{4}(20d+12)=d+7$

Answer

$d=1$

Exercise 2.4E.23

$18-(9r+7)=-16$

Exercise 2.4E.24

$15-(3r+8)=28$

Answer

$r=-7$

Exercise 2.4E.25

$5-(n-1)=19$

Exercise 2.4E.26

$-3-(m-1)=13$

Answer

$m=-15$

Exercise 2.4E.27

$11-4(y-8)=43$

Exercise 2.4E.28

$18-2(y-3)=32$

Answer

$y=-4$

Exercise 2.4E.29

$24-8(3v+6)=0$

Exercise 2.4E.30

$35-5(2w+8)=-10$

Answer

$w=\frac{1}{2}$

Exercise 2.4E.31

$4(a-12)=3(a+5)$

Exercise 2.4E.32

$-2(a-6)=4(a-3)$

Answer

$a=4$

Exercise 2.4E.33

$2(5-u)=-3(2u+6)$

Exercise 2.4E.34

$5(8-r)=-2(2r-16)$

Answer

$r=8$

Exercise 2.4E.35

$3(4n-1)-2=8n+3$

Exercise 2.4E.36

$9(2m-3)-8=4m+7$

Answer

$m=3$

Exercise 2.4E.37

$12+2(5-3y)=-9(y-1)-2$

Exercise 2.4E.38

$-15+4(2-5y)=-7(y-4)+4$

Answer

$y=-3$

Exercise 2.4E.39

$8(x-4)-7x=14$

Exercise 2.4E.40

$5(x-4)-4x=14$

Answer

$x=34$

Exercise 2.4E.41

$5+6(3s-5)=-3+2(8s-1)$

Exercise 2.4E.42

$-12+8(x-5)=-4+3(5x-2)$

Answer

$x=-6$

Exercise 2.4E.43

$4(u-1)-8=6(3u-2)-7$

Exercise 2.4E.44

$7(2n-5)=8(4n-1)-9$

Answer

$n=-1$

Exercise 2.4E.45

$4(p-4)-(p+7)=5(p-3)$

Exercise 2.4E.46

$3(a-2)-(a+6)=4(a-1)$

Answer

$a=-4$

Exercise 2.4E.47

$-(9y+5)-(3y-7)$
$=16-(4y-2)$

Exercise 2.4E.48

$-(7m+4)-(2m-5)$
$=14-(5m-3)$

Answer

$m=-4$

Exercise 2.4E.49

$4[5-8(4c-3)]$
$=12(1-13c)-8$

Exercise 2.4E.50

$5[9-2(6d-1)]$
$=11(4-10d)-139$

Answer

$d=-3$

Exercise 2.4E.51

$3[-9+8(4h-3)]$
$=2(5-12h)-19$

Exercise 2.4E.52

$3[-14+2(15k-6)]$
$=8(3-5k)-24$

Answer

$k=\frac{3}{5}$

Exercise 2.4E.53

$5[2(m+4)+8(m-7)]$
$=2[3(5+m)-(21-3m)]$

Exercise 2.4E.54

$10[5(n+1)+4(n-1)]$
$=11[7(5+n)-(25-3n)]$

Answer

$n=-5$

Exercise 2.4E.55

$5(1.2u-4.8)=-12$

Exercise 2.4E.56

$4(2.5v-0.6)=7.6$

Answer

$v=1$

Exercise 2.4E.57

$0.25(q-6)=0.1(q+18)$

Exercise 2.4E.58

$0.2(p-6)=0.4(p+14)$

Answer

$p=-34$

Exercise 2.4E.59

$0.2(30n+50)=28$

Exercise 2.4E.60

$0.5(16m+34)=-15$

Answer

$m=-4$

Classify Equations

In the following exercises, classify each equation as a conditional equation, an identity, or a contradiction and then state the solution.

Exercise 2.4E.61

$23z+19=3(5z-9)+8z+46$

Exercise 2.4E.62

$15y+32=2(10y-7)-5y+46$

Answer

identity; all real numbers

Exercise 2.4E.63

$5(b-9)+4(3b+9)=6(4b-5)-7b+21$

Exercise 2.4E.64

$9(a-4)+3(2a+5)=7(3a-4)-6a+7$

Answer

identity; all real numbers

Exercise 2.4E.65

$18(5j-1)+29=47$

Exercise 2.4E.66

$24(3d-4)+100=52$

Answer

conditional equation; $d=\frac{2}{3}$

Exercise 2.4E.67

$22(3m-4)=8(2m+9)$

Exercise 2.4E.68

$30(2n-1)=5(10n+8)$

Answer

conditional equation; $n=7$

Exercise 2.4E.69

$7v+42=11(3v+8)-2(13v-1)$

Exercise 2.4E.70

$18u-51=9(4u+5)-6(3u-10)$

Answer

contradiction; no solution

Exercise 2.4E.71

$3(6q-9)+7(q+4)=5(6q+8)-5(q+1)$

Exercise 2.4E.72

$5(p+4)+8(2p-1)=9(3p-5)-6(p-2)$

Answer

contradiction; no solution

Exercise 2.4E.73

$12(6h-1)=8(8h+5)-4$

Exercise 2.4E.74

$9(4k-7)=11(3k+1)+4$

Answer

conditional equation; $k=26$

Exercise 2.4E.75

$45(3y-2)=9(15y-6)$

Exercise 2.4E.76

$60(2x-1)=15(8x+5)$

Answer

contradiction; no solution

Exercise 2.4E.77

$16(6n+15)=48(2n+5)$

Exercise 2.4E.78

$36(4m+5)=12(12m+15)$

Answer

identity; all real numbers

Exercise 2.4E.79

$9(14d+9)+4d=13(10d+6)+3$

Exercise 2.4E.80

$11(8c+5)-8c=2(40c+25)+5$

Answer

identity; all real numbers

Everyday Math

Exercise 2.4E.81

Fencing Micah has 44 feet of fencing to make a dog run in his yard. He wants the length to be 2.5 feet more than the width. Find the length, L, by solving the equation 2L+2(L−2.5)=44.

Exercise 2.4E.82

Coins Rhonda has $\$1.90$ in nickels and dimes. The number of dimes is one less than twice the number of nickels. Find the number of nickels, n, by solving the equation $0.05n+0.10(2n-1)=1.90$.

Answer

8 nickels

Writing Exercises

Exercise 2.4E.83

Using your own words, list the steps in the general strategy for solving linear equations.

Exercise 2.4E.84

Explain why you should simplify both sides of an equation as much as possible before collecting the variable terms to one side and the constant terms to the other side.

Answer

Answers will vary.

Exercise 2.4E.85

What is the first step you take when solving the equation $3-7(y-4)=38$? Why is this your first step?

Exercise 2.4E.86

Solve the equation $\frac{1}{4}(8x+20)=3x-4$ explaining all the steps of your solution as in the examples in this section.

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objective of this section.

I can...	Confidently	With some help	No-I don't get it!
solve equations using the general strategy for solving linear equations.			
classify equations.			

ⓑ On a scale of 1-10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

Glossary

conditional equation

An equation that is true for one or more values of the variable and false for all other values of the variable is a conditional equation.

contradiction

An equation that is false for all values of the variable is called a contradiction. A contradiction has no solution.

identity

An equation that is true for any value of the variable is called an identity. The solution of an identity is all real numbers.

This page titled 2.4E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.4E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.5: Solve Equations with Fractions or Decimals

Learning Objectives

By the end of this section, you will be able to:

- Solve equations with fraction coefficients
- Solve equations with decimal coefficients

Note

Before you get started, take this readiness quiz.

1. Multiply: $8 \cdot 38$.
   If you missed this problem, review Exercise 1.6.16.
2. Find the LCD of $\frac{5}{6}$ and $\frac{1}{4}$.
   If you missed this problem, review Exercise 1.7.16.
3. Multiply 4.78 by 100.
   If you missed this problem, review Exercise 1.8.22.

Solve Equations with Fraction Coefficients

Let's use the general strategy for solving linear equations introduced earlier to solve the equation, $\frac{1}{8}x + \frac{1}{2} = \frac{1}{4}$.

Table 2.5.1

	$\frac{1}{8}x+\frac{1}{2}=\frac{1}{4}$
To isolate the x term, subtract $\frac{1}{2}$ from both sides.	$\frac{1}{8}x+\frac{1}{2}-\frac{1}{2}=\frac{1}{4}-\frac{1}{2}$
Simplify the left side.	$\frac{1}{8}x=\frac{1}{4}-\frac{1}{2}$
Change the constants to equivalent fractions with the LCD.	$\frac{1}{8}x=\frac{1}{4}-\frac{2}{4}$
Subtract.	$\frac{1}{8}x=-\frac{1}{4}$
Multiply both sides by the reciprocal of $\frac{1}{8}$.	$\frac{8}{1}\cdot\frac{1}{8}x=\frac{8}{1}\left(-\frac{1}{4}\right)$
Simplify.	$x=-2$

This method worked fine, but many students do not feel very confident when they see all those fractions. So, we are going to show an alternate method to solve equations with fractions. This alternate method eliminates the fractions.

We will apply the Multiplication Property of Equality and multiply both sides of an equation by the least common denominator of all the fractions in the equation. The result of this operation will be a new equation, equivalent to the first, but without fractions. This process is called "clearing" the equation of fractions.

Let's solve a similar equation, but this time use the method that eliminates the fractions.

Example 2.5.1: How to Solve Equations with Fraction Coefficients

Solve: $\frac{1}{6}y - \frac{1}{3} = \frac{5}{6}$

Solution

Step 1. Find the least common denominator of *all* the fractions in the equation.	What is the LCD of $\frac{1}{6}$, $\frac{1}{3}$, and $\frac{5}{6}$?	$\frac{1}{6}y-\frac{1}{3}=\frac{5}{6}$ LCD $=6$
Step 2. Multiply both sides of the equation by that LCD. This clears the fractions.	Multiply both sides of the equation by the LCD 6. Use the Distributive Property. Simplify – and notice, no more fractions!	$6\left(\frac{1}{6}y-\frac{1}{3}\right)=6\left(\frac{5}{6}\right)$ $6\cdot\frac{1}{6}y-6\cdot\frac{1}{3}=6\cdot\frac{5}{6}$ $y-2=5$
Step 3. Solve using the General Strategy for Solving Linear Equations.	To isolate the "y" term, add 2. Simplify.	$y-2+2=5+2$ $y=7$

Try It 2.5.2

Solve: $\frac{1}{4}x+\frac{1}{2}=\frac{5}{8}$

Answer

$x=\frac{1}{2}$

Try It 2.5.3

Solve: $\frac{1}{8}x+\frac{1}{2}=\frac{1}{4}$

Answer

$x=-2$

Notice in Exercise 2.5.1, once we cleared the equation of fractions, the equation was like those we solved earlier in this chapter. We changed the problem to one we already knew how to solve! We then used the General Strategy for Solving Linear Equations.

STRATEGY TO SOLVE EQUATIONS WITH FRACTION COEFFICIENTS.

1. Find the least common denominator of *all* the fractions in the equation.
2. Multiply both sides of the equation by that LCD. This clears the fractions.
3. Solve using the General Strategy for Solving Linear Equations.

Example 2.5.4

Solve: $6=\frac{1}{2}v+\frac{2}{5}v-\frac{3}{4}v$

Solution

We want to clear the fractions by multiplying both sides of the equation by the LCD of all the fractions in the equation.

Find the LCD of all fractions in the equation.	$6=\frac{1}{2}v+\frac{2}{5}v-\frac{3}{4}v$
The LCD is 20.	
Multiply both sides of the equation by 20.	$20(6)=20\cdot\left(\frac{1}{2}v+\frac{2}{5}v-\frac{3}{4}v\right)$
Distribute.	$20(6)=20\cdot\frac{1}{2}v+20\cdot\frac{2}{5}v-20\cdot\frac{3}{4}v$

Simplify—notice, no more fractions!		$120 = 10v + 8v - 15v$
Combine like terms.		$120 = 3v$
Divide by 3.		$\frac{120}{3} = \frac{3v}{3}$
Simplify.		$40 = v$
Check:	$6 = \frac{1}{2}v + \frac{2}{5}v - \frac{3}{4}v$	
Let v=40.	$6 \stackrel{?}{=} \frac{1}{2}(40) + \frac{2}{5}(40) - \frac{3}{4}(40)$	
	$6 \stackrel{?}{=} 20 + 16 - 30$	
	$6 = 6$ ✓	

Try It 2.5.5

Solve: $7 = \frac{1}{2}x + \frac{3}{4}x - \frac{2}{3}x$

Answer

$x = 12$

Try It 2.5.6

Solve: $-1 = \frac{1}{2}u + \frac{1}{4}u - \frac{2}{3}u$

Answer

$u = -12$

In the next example, we again have variables on both sides of the equation.

Example 2.5.7

Solve: $a + \frac{3}{4} = \frac{3}{8}a - \frac{1}{2}$

Solution

	$a + \frac{3}{4} = \frac{3}{8}a - \frac{1}{2}$
Find the LCD of all fractions in the equation. The LCD is 8.	
Multiply both sides by the LCD.	$8\left(a + \frac{3}{4}\right) = 8\left(\frac{3}{8}a - \frac{1}{2}\right)$
Distribute.	$8 \cdot a + 8 \cdot \frac{3}{4} = 8 \cdot \frac{3}{8}a - 8 \cdot \frac{1}{2}$
Simplify—no more fractions.	$8a + 6 = 3a - 4$
Subtract 3a3a from both sides.	$8a - 3a + 6 = 3a - 3a - 4$
Simplify.	$5a + 6 = -4$
Subtract 6 from both sides.	$5a + 6 - 6 = -4 - 6$
Simplify.	$5a = -10$

Divide by 5.		$\frac{5a}{5}=\frac{-10}{5}$
Simplify.		$a=-2$
Check:	$a+\frac{3}{4}=\frac{3}{8}a-\frac{1}{2}$	
Let a=−2.	$-2+\frac{3}{4}\stackrel{?}{=}\frac{3}{8}(-2)-\frac{1}{2}$	
	$-\frac{8}{4}+\frac{3}{4}\stackrel{?}{=}-\frac{16}{8}-\frac{4}{8}$	
	$-\frac{5}{4}=-\frac{10}{8}$	
	$-\frac{5}{4}=-\frac{5}{4}\checkmark$	

? Try It 2.5.8

Solve: $x+\frac{1}{3}=\frac{1}{6}x-\frac{1}{2}$

Answer

$x=-1$

? Try It 2.5.9

Solve: $c+\frac{3}{4}=\frac{1}{2}c-\frac{1}{4}$

Answer

$c=-2$

In the next example, we start by using the Distributive Property. This step clears the fractions right away.

✓ Example 2.5.10

Solve: $-5=\frac{1}{4}(8x+4)$

Solution

	$-5=\frac{1}{4}(8x+4)$
Distribute.	$-5=\frac{1}{4}\cdot 8x+\frac{1}{4}\cdot 4$
Simplify. Now there are no fractions.	$-5=2x+1$
Subtract 1 from both sides.	$-5-1=2x+1-1$
Simplify.	$-6=2x$
Divide by 2.	$\frac{-6}{2}=\frac{2x}{2}$
Simplify.	$-3=x$
Check:	$-5=\frac{1}{4}(8x+4)$
Let x=−3.	$-5\stackrel{?}{=}\frac{1}{2}(4(-3)+2)$

	$-5 \stackrel{?}{=} \frac{1}{2}(-12+2)$	
	$-5 \stackrel{?}{=} \frac{1}{2}(-10)$	
	$-5 = -5$ ✓	

Try It 2.5.11

Solve: $-11 = \frac{1}{2}(6p+2)$

Answer

$p = -4$

Try It 2.5.12

Solve: $8 = \frac{1}{3}(9q+6)$

Answer

$q = 2$

In the next example, even after distributing, we still have fractions to clear.

Example 2.5.13

Solve: $\frac{1}{2}(y-5) = \frac{1}{4}(y-1)$

Solution

	$\frac{1}{2}(y-5) = \frac{1}{4}(y-1)$
Distribute.	$\frac{1}{2} \cdot y - \frac{1}{2} \cdot 5 = \frac{1}{4} \cdot y - \frac{1}{4} \cdot 1$
Simplify.	$\frac{1}{2}y - \frac{5}{2} = \frac{1}{4}y - \frac{1}{4}$
Multiply by the LCD, 4.	$4\left(\frac{1}{2}y - \frac{5}{2}\right) = 4\left(\frac{1}{4}y - \frac{1}{4}\right)$
Distribute.	$4 \cdot \frac{1}{2}y - 4 \cdot \frac{5}{2} = 4 \cdot \frac{1}{4}y - 4 \cdot \frac{1}{4}$
Simplify.	$2y - 10 = y - 1$
Collect the variables to the left.	$2y - y - 10 = y - y - 1$
Simplify.	$y - 10 = -1$
Collect the constants to the right.	$y - 10 + 10 = -1 + 10$
Simplify.	$y = 9$
Check:	$\frac{1}{2}(y-5) = \frac{1}{4}(y-1)$
Let y=9.	$\frac{1}{2}(9-5) \stackrel{?}{=} \frac{1}{4}(9-1)$
Finish the check on your own.	

Try It 2.5.14

Solve: $\frac{1}{5}(n+3)=\frac{1}{4}(n+2)$

Answer

$n=2$

Try It 2.5.15

Solve: $\frac{1}{2}(m-3)=\frac{1}{4}(m-7)$

Answer

$m=-1$

Example 2.5.16

Solve: $\frac{5x-3}{4}=\frac{x}{2}$

Solution

	$\frac{5x-3}{4}=\frac{x}{2}$
Multiply by the LCD, 4.	$4\left(\frac{5x-3}{4}\right)=4\left(\frac{x}{2}\right)$
Simplify.	$5x-3=2x$
Collect the variables to the right.	$5x-5x-3=2x-5x$
Simplify.	$-3=-3x$
Divide.	$\frac{-3}{-3}=\frac{-3x}{-3}$
Simplify.	$1=x$
Check:	$\frac{5x-3}{4}=\frac{x}{2}$
Let x=1.	$\frac{5(1)-3}{4}\stackrel{?}{=}\frac{1}{2}$
	$\frac{2}{4}\stackrel{?}{=}\frac{1}{2}$
	$\frac{1}{2}=\frac{1}{2}\checkmark$

Try It 2.5.17

Solve: $\frac{4y-7}{3}=\frac{y}{6}$

Answer

$y=2$

Try It 2.5.18

Solve: $\frac{-2z-5}{4}=\frac{z}{8}$

Answer

$z = -2$

Example 2.5.19

Solve: $\frac{a}{6} + 2 = \frac{a}{4} + 3$

Solution

	$\frac{a}{6} + 2 = \frac{a}{4} + 3$
Multiply by the LCD, 12.	$12\left(\frac{a}{6} + 2\right) = 12\left(\frac{a}{4} + 3\right)$
Distribute.	$12 \cdot \frac{a}{6} + 12 \cdot 2 = 12 \cdot \frac{a}{4} + 12 \cdot 3$
Simplify.	$2a + 24 = 3a + 36$
Collect the variables to the right.	$2a - 2a + 24 = 3a - 2a + 36$
Simplify.	$24 = a + 36$
Collect the constants to the left.	$24 - 36 = a + 36 - 36$
Simplify.	$a = -12$
Check:	$\frac{a}{6} + 2 = \frac{a}{4} + 3$
Let a=−12.	$\frac{-12}{6} + 2 \stackrel{?}{=} \frac{-12}{4} + 3$
	$-2 + 2 \stackrel{?}{=} -3 + 3$
	$0 = 0$ ✓

Try It 2.5.20

Solve: $\frac{b}{10} + 2 = \frac{b}{4} + 5$

Answer

$b = -20$

Try It 2.5.21

Solve: $\frac{c}{6} + 3 = \frac{c}{3} + 4$

Answer

$c = -6$

Example 2.5.22

Solve: $\frac{4q+3}{2} + 6 = \frac{3q+5}{4}$

Solution

	$\frac{4q+3}{2} + 6 = \frac{3q+5}{4}$
Multiply by the LCD, 4.	$4\left(\frac{4q+3}{2} + 6\right) = 4\left(\frac{3q+5}{4}\right)$

Distribute.	$4\left(\frac{4q+3}{2}\right)+4\cdot 6=4\cdot\left(\frac{3q+5}{4}\right)$
Simplify.	$2(4q+3)+24=3q+5$ $8q+6+24=3q+5$ $8q+30=3q+5$
Collect the variables to the left.	$8q-3q+30=3q-3q+5$
Simplify.	$5q+30=5$
Collect the constants to the right.	$5q+30-30=5-30$
Simplify.	$5q=-25$
Divide by 5.	$\frac{5q}{5}=\frac{-25}{5}$
Simplify.	$q=-5$
Check:	$\frac{4q+3}{2}+6=\frac{3q+5}{4}$
Let q=−5.	$\frac{4(-5)+3}{2}+6\stackrel{?}{=}\frac{3(-5)+5}{4}$
Finish the check on your own.	

Try It 2.5.23

Solve: $\frac{3r+5}{6}+1=\frac{4r+3}{3}$

Answer

$r=1$

Try It 2.5.24

Solve: $\frac{2s+3}{2}+1=\frac{3s+2}{4}$

Answer

$s=-8$

Solve Equations with Decimal Coefficients

Some equations have decimals in them. This kind of equation will occur when we solve problems dealing with money or percentages. But decimals can also be expressed as fractions. For example, $0.3=\frac{3}{10}$ and $0.17=\frac{17}{100}$. So, with an equation with decimals, we can use the same method we used to clear fractions—multiply both sides of the equation by the least common denominator.

Example 2.5.25

Solve: $0.06x+0.02=0.25x-1.5$

Solution

Look at the decimals and think of the equivalent fractions.

$0.06=\frac{6}{100}\quad 0.02=\frac{2}{100}\quad 0.25=\frac{25}{100}\quad 1.5=1\frac{5}{10}$

Notice, the LCD is 100.

By multiplying by the LCD, we will clear the decimals from the equation.

	$0.06x + 0.02 = 0.25x - 1.5$
Multiply both side by 100.	$100(0.06x + 0.02) = 100(0.25x - 1.5)$
Distribute.	$100(0.06x) + 100(0.02) = 100(0.25x) - 100(1.5)$
Multiply, and now we have no more decimals.	$6x + 2 = 25x - 150$
Collect the variables to the right.	$6x - 6x + 2 = 25x - 6x - 150$
Simplify.	$2 = 19x - 150$
Collect the variables to the right.	$2 + 150 = 19x - 150 + 150$
Simplify.	$152 = 19x$
Divide by 19.	$\frac{152}{19} = \frac{19x}{19}$
Simplify.	$8 = x$
Check: Let x=8 $0.06(8) + 0.02 \stackrel{?}{=} 0.25(8) - 1.5$ $0.48 + 0.02 \stackrel{?}{=} 2.00 - 1.5$ $0.50 = 0.50$ ✓	

Try It 2.5.26

Solve: $0.14h + 0.12 = 0.35h - 2.4$

Answer

$h = 12$

Try It 2.5.27

Solve: $0.65k - 0.1 = 0.4k - 0.35$

Answer

$k = -1$

The next example uses an equation that is typical of the money applications in the next chapter. Notice that we distribute the decimal before we clear all the decimals.

Example 2.5.28

Solve: $0.25x + 0.05(x + 3) = 2.85$

Solution

	$0.25x + 0.05(x + 3) = 2.85$
Distribute first.	$0.25x + 0.05x + 0.15 = 2.85$
Combine like terms.	$0.30x + 0.15 = 2.85$
To clear decimals, multiply by 100.	$100(0.30x + 0.15) = 100(2.85)$
Distribute.	$30x + 15 = 285$
Subtract 15 from both sides.	$30x + 15 - 15 = 285 - 15$

Simplify.	$30x = 270$
Divide by 30.	$\frac{30x}{30} = \frac{270}{30}$
Simplify.	$x = 9$
Check it yourself by substituting x=9 into the original equation.	

Try It 2.5.29

Solve: $0.25n + 0.05(n + 5) = 2.95$

Answer

$n = 9$

Try It 2.5.30

Solve: $0.10d + 0.05(d - 5) = 2.15$

Answer

$d = 16$

Key Concepts

- **Strategy to Solve an Equation with Fraction Coefficients**
  1. Find the least common denominator of all the fractions in the equation.
  2. Multiply both sides of the equation by that LCD. This clears the fractions.
  3. Solve using the General Strategy for Solving Linear Equations.

This page titled 2.5: Solve Equations with Fractions or Decimals is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.5: Solve Equations with Fractions or Decimals** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.5E: Exercises

Practice Makes Perfect

Solve Equations with Fraction Coefficients

In the following exercises, solve each equation with fraction coefficients.

Exercise 2.5E.1

$\frac{1}{4}x - \frac{1}{2} = -\frac{3}{4}$

Exercise 2.5E.2

$\frac{3}{4}x - \frac{1}{2} = \frac{1}{4}$

Answer

x=1

Exercise 2.5E.3

$\frac{5}{6}y - \frac{2}{3} = -\frac{3}{2}$

Exercise 2.5E.4

$\frac{5}{6}y - \frac{1}{3} = -\frac{7}{6}$

Answer

$y = -1$

Exercise 2.5E.5

$\frac{1}{2}a + \frac{3}{8} = \frac{3}{4}$

Exercise 2.5E.6

$\frac{5}{8}b + \frac{1}{2} = -\frac{3}{4}$

Answer

$b = -2$

Exercise 2.5E.7

$2 = \frac{1}{3}x - \frac{1}{2}x + \frac{2}{3}x$

Exercise 2.5E.8

$2 = \frac{3}{5}x - \frac{1}{3}x + \frac{2}{5}x$

Answer

$x = 3$

Exercise 2.5E. 9

$\frac{1}{4}m - \frac{4}{5}m + \frac{1}{2}m = -1$

Exercise 2.5E. 10

$\frac{5}{6}n - \frac{1}{4}n - \frac{1}{2}n = -2$

Answer

$n = -24$

Exercise 2.5E. 11

$x + \frac{1}{2} = \frac{2}{3}x - \frac{1}{2}$

Exercise 2.5E. 12

$x + \frac{3}{4} = \frac{1}{2}x - \frac{5}{4}$

Answer

$x = -4$

Exercise 2.5E. 13

$\frac{1}{3}w + \frac{5}{4} = w - \frac{1}{4}$

Exercise 2.5E. 14

$\frac{3}{2}z + \frac{1}{3} = z - \frac{2}{3}$

Answer

$z = -2$

Exercise 2.5E. 15

$\frac{1}{2}x - \frac{1}{4} = \frac{1}{12}x + \frac{1}{6}$

Exercise 2.5E. 16

$\frac{1}{2}a - \frac{1}{4} = \frac{1}{6}a + \frac{1}{12}$

Answer

$a = 1$

Exercise 2.5E. 17

$\frac{1}{3}b + \frac{1}{5} = \frac{2}{5}b - \frac{3}{5}$

Exercise 2.5E.18

$\frac{1}{3}x+\frac{2}{5}=\frac{1}{5}x-\frac{2}{5}$

Answer

$x=-6$

Exercise 2.5E.19

$1=\frac{1}{6}(12x-6)$

Exercise 2.5E.20

$1=\frac{1}{5}(15x-10)$

Answer

$x=1$

Exercise 2.5E.21

$\frac{1}{4}(p-7)=\frac{1}{3}(p+5)$

Exercise 2.5E.22

$\frac{1}{5}(q+3)=\frac{1}{2}(q-3)$

Answer

$q=7$

Exercise 2.5E.23

$\frac{1}{2}(x+4)=\frac{3}{4}$

Exercise 2.5E.24

$\frac{1}{3}(x+5)=\frac{5}{6}$

Answer

$x=-\frac{5}{2}$

Exercise 2.5E.25

$\frac{5q-8}{5}=\frac{2q}{10}$

Exercise 2.5E.26

$\frac{4m+2}{6}=\frac{m}{3}$

Answer

$m=-1$

Exercise 2.5E.27

$\frac{4n+8}{4} = \frac{n}{3}$

Exercise 2.5E.28

$\frac{3p+6}{3} = \frac{p}{2}$

Answer

$p = -4$

Exercise 2.5E.29

$\frac{u}{3} - 4 = \frac{u}{2} - 3$

Exercise 2.5E.30

$\frac{v}{10} + 1 = \frac{v}{4} - 2$

Answer

$v = 20$

Exercise 2.5E.31

$\frac{c}{15} + 1 = \frac{c}{10} - 1$

Exercise 2.5E.32

$\frac{d}{6} + 3 = \frac{d}{8} + 2$

Answer

$d = -24$

Exercise 2.5E.33

$\frac{3x+4}{2} + 1 = \frac{5x+10}{8}$

Exercise 2.5E.34

$\frac{10y-2}{3} + 3 = \frac{10y+1}{9}$

Answer

$y = -1$

Exercise 2.5E.35

$\frac{7u-1}{4} - 1 = \frac{4u+8}{5}$

Exercise 2.5E.36

$\frac{3v-6}{2}+5=\frac{11v-4}{5}$

Answer

$v=4$

Solve Equations with Decimal Coefficients

In the following exercises, solve each equation with decimal coefficients.

Exercise 2.5E.37

$0.6y+3=9$

Exercise 2.5E.38

$0.4y-4=2$

Answer

$y=15$

Exercise 2.5E.39

$3.6j-2=5.2$

Exercise 2.5E.40

$2.1k+3=7.2$

Answer

$k=2$

Exercise 2.5E.41

$0.4x+0.6=0.5x-1.2$

Exercise 2.5E.42

$0.7x+0.4=0.6x+2.4$

Answer

$x=20$

Exercise 2.5E.43

$0.23x+1.47=0.37x-1.05$

Exercise 2.5E.44

$0.48x+1.56=0.58x-0.64$

Answer

$x = 22$

Exercise 2.5E.45

$0.9x - 1.25 = 0.75x + 1.75$

Exercise 2.5E.46

$1.2x - 0.91 = 0.8x + 2.29$

Answer

$x = 8$

Exercise 2.5E.47

$0.05n + 0.10(n + 8) = 2.15$

Exercise 2.5E.48

$0.05n + 0.10(n + 7) = 3.55$

Answer

$n = 19$

Exercise 2.5E.49

$0.10d + 0.25(d + 5) = 4.05$

Exercise 2.5E.50

$0.10d + 0.25(d + 7) = 5.25$

Answer

$d = 10$

Exercise 2.5E.51

$0.05(q - 5) + 0.25q = 3.05$

Exercise 2.5E.52

$0.05(q - 8) + 0.25q = 4.10$

Answer

$q = 15$

Everyday Math

Exercise 2.5E.53

Coins Taylor has $200 in dimes and pennies. The number of pennies is 2 more than the number of dimes. Solve the equation $0.10d+0.01(d+2)=2$ for d, the number of dimes.

Exercise 2.5E.54

Stamps Paula bought $22.82 worth of 49-cent stamps and 21-cent stamps. The number of 21-cent stamps was 8 less than the number of 49-cent stamps. Solve the equation $0.49s+0.21(s-8)=22.82$ for s, to find the number of 49-cent stamps Paula bought.

Answer

$s=35$

Writing Exercises

Exercise 2.5E.55

Explain how you find the least common denominator of $\frac{3}{8}$, $\frac{1}{6}$, and $\frac{2}{3}$

Exercise 2.5E.56

If an equation has several fractions, how does multiplying both sides by the LCD make it easier to solve?

Answer

Answers will vary.

Exercise 2.5E.57

If an equation has fractions only on one side, why do you have to multiply both sides of the equation by the LCD?

Exercise 2.5E.58

In the equation $0.35x+2.1=3.85$ what is the LCD? How do you know?

Answer

100. Justifications will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve equations with fraction coefficients.			
solve equations with decimal coefficients.			

ⓑ Overall, after looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

This page titled 2.5E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.5E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.6: Solve Proportions and Similar Figure Applications

Learning Objectives

By the end of this section, you will be able to:

- Solve proportions
- Solve similar figure applications

Note

Before you get started, take this readiness quiz.

If you miss a problem, go back to the section listed and review the material.

1. Solve $\frac{n}{3} = 30$.
   If you missed this problem, review Exercise 2.2.25.
2. The perimeter of a triangular window is 23 feet. The lengths of two sides are ten feet and six feet. How long is the third side?
   If you missed this problem, review Example 3.4.2

Solve Proportions

When two rational expressions are equal, the equation relating them is called a *proportion*.

Definition: PROPORTION

A **proportion** is an equation of the form $\frac{a}{b} = \frac{c}{d}$, where $b \neq 0$, $d \neq 0$.

The proportion is read “a is to b, as c is to d"

The equation $\frac{1}{2} = \frac{4}{8}$ is a proportion because the two fractions are equal.

The proportion $\frac{1}{2} = \frac{4}{8}$ is read “1 is to 2 as 4 is to 8.”

Proportions are used in many applications to ‘scale up’ quantities. We’ll start with a very simple example so you can see how proportions work. Even if you can figure out the answer to the example right away, make sure you also learn to solve it using proportions.

Suppose a school principal wants to have 1 teacher for 20 students. She could use proportions to find the number of teachers for 60 students. We let x be the number of teachers for 60 students and then set up the proportion:

$$\frac{1 \text{ teacher}}{20 \text{ students}} = \frac{x \text{ teachers}}{60 \text{ students}}$$

We are careful to match the units of the numerators and the units of the denominators—teachers in the numerators, students in the denominators.

Since a proportion is an equation with rational expressions, we will solve proportions the same way we solved equations in Solve Rational Equations. We’ll multiply both sides of the equation by the LCD to clear the fractions and then solve the resulting equation.

	$\frac{1}{20} = \frac{x}{60}$
Multiply both sides by the LCD, 60.	$\frac{1}{20} \cdot 60 = \frac{x}{60} \cdot 60$
Simplify.	$3 = x$

	The principal needs 3 teachers for 60 students.

Now we'll do a few examples of solving numerical proportions without any units. Then we will solve applications using proportions.

Example 2.6.1

$\frac{x}{63} = \frac{4}{7}$.

Solution

		$\frac{x}{63} = \frac{4}{7}$
To isolate x, multiply both sides by the LCD, 63.		$63\left(\frac{x}{63}\right) = 63\left(\frac{4}{7}\right)$
Simplify.		$x = \frac{9 \cdot 7 \cdot 4}{7}$
Divide the common factors.		$x = 36$
Check. To check our answer, we substitute into the original proportion.		
	$\frac{x}{63} = \frac{4}{7}$	
Substitute $x = 36$.	$\frac{36}{63} \stackrel{?}{=} \frac{4}{7}$	
Show common factors.	$\frac{4 \cdot 9}{7 \cdot 9} \stackrel{?}{=} \frac{4}{7}$	
Simplify.	$\frac{4}{7} = \frac{4}{7}$ ✓	

Try It 2.6.1

$\frac{n}{84} = \frac{11}{12}$.

Answer

77

Try It 2.6.2

$\frac{y}{96} = \frac{13}{12}$.

Answer

104

Example 2.6.2

$\frac{144}{a} = \frac{9}{4}$.

Solution

		$\frac{144}{a}=\frac{9}{4}$
Multiply both sides by the LCD.		$\frac{144}{a}\cdot 4a=\frac{9}{4}\cdot 4a$
Remove common factors on each side.		$4\cdot 144=a\cdot 9$
Simplify.		$576=9a$
Divide both sides by 9.		$\frac{576}{9}=\frac{9a}{9}$
Simplify.		$64=a$
Check.		
	$\frac{144}{a}=\frac{9}{4}$	
Substitute $a=64$.	$\frac{144}{64}\stackrel{?}{=}\frac{9}{4}$	
Show common factors.	$\frac{9\cdot 16}{4\cdot 16}\stackrel{?}{=}\frac{9}{4}$	
Simplify.	$\frac{9}{4}=\frac{9}{4}$ ✓	

Try It 2.6.3

$\frac{91}{b}=\frac{7}{5}$.

Answer

65

Try It 2.6.4

$\frac{39}{c}=\frac{13}{8}$.

Answer

24

Example 2.6.3

$\frac{n}{n+14}=\frac{5}{7}$.

Solution

		$\frac{n}{n+14}=\frac{5}{7}$
Multiply both sides by the LCD.		$7(n+14)\left(\frac{n}{n+14}\right)=7(n+14)\left(\frac{5}{7}\right)$
Remove common factors on each side.		$7n=5(n+14)$
Simplify.		$7n=5n+70$
Solve for n.		$2n=70$
		$n=35$

Check.		
	$\frac{n}{n+14}=\frac{5}{7}$	
Substitute $n=35$.	$\frac{35}{35+14}\stackrel{?}{=}\frac{5}{7}$	
Simplify.	$\frac{35}{49}\stackrel{?}{=}\frac{5}{7}$	
Show common factors.	$\frac{5\cdot 7}{7\cdot 7}\stackrel{?}{=}\frac{5}{7}$	
Simplify.	$\frac{5}{7}=\frac{5}{7}$ ✓	

Try It 2.6.5

$\frac{y}{y+55}=\frac{3}{8}$.

Answer

33

Try It 2.6.6

$\frac{z}{z-84}=-\frac{1}{5}$.

Answer

14

Example 2.6.4

$\frac{p+12}{9}=\frac{p-12}{6}$.

Solution

		$\frac{p+12}{9}=\frac{p-12}{6}$
Multiply both sides by the LCD, 18.		$18\cdot\frac{p+12}{9}=18\cdot\frac{p+12}{6}$
Simplify.		$2(p+12)=3(p-12)$
Distribute.		$2p+24=3p-36$
Solve fpr p.		$60=p$
Check.		
	$\frac{p+12}{9}=\frac{p-12}{6}$	
Substitute $p=60$.	$\frac{60+12}{9}\stackrel{?}{=}\frac{60-12}{6}$	
Simplify.	$\frac{72}{9}\stackrel{?}{=}\frac{48}{6}$	
Divide.	$8=8$ ✓	

Try It 2.6.7

$\frac{v+30}{8} = \frac{v+66}{12}$.

Answer

42

Try It 2.6.8

$\frac{2x+15}{9} = \frac{7x+3}{15}$.

Answer

6

To solve applications with proportions, we will follow our usual strategy for solving applications. But when we set up the proportion, we must make sure to have the units correct—the units in the numerators must match and the units in the denominators must match.

Example 2.6.5

When pediatricians prescribe acetaminophen to children, they prescribe 5 milliliters (ml) of acetaminophen for every 25 pounds of the child's weight. If Zoe weighs 80 pounds, how many milliliters of acetaminophen will her doctor prescribe?

Solution

Identify what we are asked to find, and choose a variable to represent it.	How many ml of acetaminophen will the doctor prescribe?
	Let a=ml of acetaminophen.
Write a sentence that gives the information to find it.	If 5 ml is prescribed for every 25 pounds, how much will be prescribed for 80 pounds?
Translate into a proportion–be careful of the units. $\frac{5}{25} = \frac{a}{80}$	
Multiply both sides by the LCD, 400.	$400\left(\frac{5}{25}\right) = 400\left(\frac{a}{80}\right)$
Remove common factors on each side.	$25 \cdot 16\left(\frac{5}{25}\right) = 80 \cdot 5\left(\frac{a}{80}\right)$
Simplify, but don't multiply on the left. Notice what the next step will be.	$16 \cdot 5 = 5a$
Solve for a.	$\frac{16 \cdot 5}{5} = \frac{5a}{5}$
Check.	$16 = a$
Is the answer reasonable?	
Yes, since 80 is about 3 times 25, the medicine should be about 3 times 5. So 16 ml makes sense.	

Substitute $a = 16$ in the original proportion. $\frac{5}{25} = \frac{a}{80}$ $\frac{5}{25} \stackrel{?}{=} \frac{16}{80}$ $\frac{1}{5} = \frac{1}{5}$ ✓	
Write a complete sentence.	The pediatrician would prescribe 16 ml of acetaminophen to Zoe.

Try It 2.6.9

Pediatricians prescribe 5 milliliters (ml) of acetaminophen for every 25 pounds of a child's weight. How many milliliters of acetaminophen will the doctor prescribe for Emilia, who weighs 60 pounds?

Answer

12 ml

Try It 2.6.10

For every 1 kilogram (kg) of a child's weight, pediatricians prescribe 15 milligrams (mg) of a fever reducer. If Isabella weighs 12 kg, how many milligrams of the fever reducer will the pediatrician prescribe?

Answer

180 ml

Example 2.6.6

A 16-ounce iced caramel macchiato has 230 calories. How many calories are there in a 24-ounce iced caramel macchiato?

Solution

Identify what we are asked to find, and choose a variable to represent it.	How many calories are in a 24 ounce iced caramel macchiato?
	Let c=calories in 24 ounces.
Write a sentence that gives the information to find it.	If there are 230 calories in 16 ounces, then how many calories are in 24 ounces?
Translate into a proportion–be careful of the units. $\frac{230}{16} = \frac{c}{24}$	
Multiply both sides by the LCD, 48.	$48\left(\frac{230}{16}\right) = 48\left(\frac{c}{24}\right)$
Remove common factors on each side.	$16 \cdot 3\left(\frac{230}{16}\right) = 24 \cdot 2\left(\frac{c}{24}\right)$
Simplify.	$690 = 2c$
Solve for c.	$\frac{690}{2} = \frac{2c}{2}$
	$345 = c$
Check.	

Is the answer reasonable?	
Yes, 345 calories for 24 ounces is more than 290 calories for 16 ounces, but not too much more.	
Substitute $c = 345$ in the original proportion. $\frac{230}{16} = \frac{c}{24}$ $\frac{230}{16} \stackrel{?}{=} \frac{345}{24}$ $\frac{115}{8} = \frac{115}{8}$ ✓	
Write a complete sentence.	There are 345 calories in a 24-ounce iced caramel macchiato.

Try It 2.6.11

At a fast-food restaurant, a 22-ounce chocolate shake has 850 calories. How many calories are in their 12-ounce chocolate shake? Round your answer to nearest whole number.

Answer

464 calories

Try It 2.6.12

Yaneli loves Starburst candies, but wants to keep her snacks to 100 calories. If the candies have 160 calories for 8 pieces, how many pieces can she have in her snack?

Answer

5 pieces

Example 2.6.7

Josiah went to Mexico for spring break and changed $325 dollars into Mexican pesos. At that time, the exchange rate had $1 US is equal to 12.54 Mexican pesos. How many Mexican pesos did he get for his trip?

Solution

What are you asked to find?	How many Mexican pesos did Josiah get?
Assign a variable.	Let p=the number of Mexican pesos.
Write a sentence that gives the information to find it.	If $1 US is equal to 12.54 Mexican pesos, then $325 is how many pesos?
Translate into a proportion–be careful of the units. $\frac{1}{12.54} = \frac{325}{p}$	
Multiply both sides by the LCD, 12.54p.	$12.54p\left(\frac{1}{12.54}\right) = 12.54p\left(\frac{325}{p}\right)$
Remove common factors on each side.	$\cancel{12.54p}\left(\frac{1}{\cancel{12.54}}\right) = 12.54\cancel{p}\left(\frac{325}{\cancel{p}}\right)$
Simplify.	$p = 4075.5$
Check.	
Is the answer reasonable?	

Yes, $100 would be 1,254 pesos. $325 is a little more than 3 times this amount, so our answer of 4075.5 pesos makes sense.	
Substitute $p = 4075.5$ in the original proportion. Use a calculator. $\frac{1}{12.54} = \frac{325}{p}$ $\frac{1}{12.54} \stackrel{?}{=} \frac{325}{4075.5}$ $0.07874... = 0.07874...$ ✓	
Write a complete sentence.	Josiah got 4075.5 pesos for his spring break trip.

? Try It 2.6.13

Yurianna is going to Europe and wants to change $800 dollars into Euros. At the current exchange rate, $1 US is equal to 0.738 Euro. How many Euros will she have for her trip?

Answer

590.4 Euros

? Try It 2.6.14

Corey and Nicole are traveling to Japan and need to exchange $600 into Japanese yen. If each dollar is 94.1 yen, how many yen will they get?

Answer

56,460 yen

In the example above, we related the number of pesos to the number of dollars by using a proportion. We could say the number of pesos *is proportional to* the number of dollars. If two quantities are related by a proportion, we say that they are proportional.


Solve Similar Figure Applications

When you shrink or enlarge a photo on a phone or tablet, figure out a distance on a map, or use a pattern to build a bookcase or sew a dress, you are working with **similar figures**. If two figures have exactly the same shape, but different sizes, they are said to be *similar.* One is a scale model of the other. All their corresponding angles have the same measures and their corresponding sides are in the same ratio.

Definition: SIMILAR FIGURES

Two figures are similar if the measures of their corresponding angles are equal and their corresponding sides are in the same ratio.

For example, the two triangles in the figure below are similar. Each side of ΔABC is 4 times the length of the corresponding side of ΔXYZ.



$m\angle A = m\angle X$

$m\angle B = m\angle Y$


$m\angle C = m\angle Z$

$\frac{16}{4} = \frac{20}{5} = \frac{12}{3}$

This is summed up in the Property of Similar Triangles.

Definition: PROPERTY OF SIMILAR TRIANGLES

If ΔABC is similar to ΔXYZ



$m\angle A = m\angle X$

$m\angle B = m\angle Y$

$m\angle C = m\angle Z$

$\frac{a}{x} = \frac{b}{y} = \frac{c}{z}$


To solve applications with similar figures we will follow the Problem-Solving Strategy for Geometry Applications we used earlier.

Definition: SOLVE GEOMETRY APPLICATIONS.

1. **Read** the problem and make all the words and ideas are understood. Draw the figure and label it with the given information.
2. **Identify** what we are looking for.
3. **Name** what we are looking for by choosing a variable to represent it.
4. **Translate** into an equation by writing the appropriate formula or model for the situation. Substitute in the given information.
5. **Solve the equation** using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Example 2.6.8

ΔABC is similar to ΔXYZ. The lengths of two sides of each triangle are given in the figure.



Find the length of the sides of the similar triangles.


Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	Figure is given.
Step 2. Identify what we are looking for.	the length of the sides of similar triangles.
Step 3. Name the variables.	Let a= length of the third side of ΔABC. y= length of the third side of ΔXYZ.
Step 4. Translate.	Since the triangles are similar, the corresponding sides are proportional.
We need to write an equation that compares the side we are looking for to a known ratio. Since the side $AB = 4$ corresponds to the side $XY = 3$ we know $\frac{AB}{XY} = \frac{4}{3}$. So we write equations with $\frac{AB}{XY}$ to find the sides we are looking for. Be careful to match up corresponding sides correctly.	$\frac{AB}{XY} = \frac{BC}{YZ} = \frac{AC}{XZ}$. To find a: $\frac{AB}{XY} = \frac{BC}{YZ}$ (sides of large triangle → / sides of small triangle →) To find y: $\frac{AB}{XY} = \frac{AC}{XZ}$
Substitute.	$\frac{4}{3} = \frac{a}{4.5}$ $\frac{4}{3} = \frac{3.2}{y}$

Step 5. Solve the equation.	$3a = 4(4.5)$ Figure_08_07_030f_img_new.jpg" />
	$a = 6$ $y = 2.4$
Step 6. Check.	
Step 7. Answer the question.	The third side of ΔABC is 6 and the third side of ΔXYZ is 2.4.

Try It 2.6.15

ΔABC is similar to ΔXYZ. The lengths of two sides of each triangle are given in the figure.



Find the length of side a

Answer

8

Try It 2.6.16

ΔABC is similar to ΔXYZ. The lengths of two sides of each triangle are given in the figure.


Answer

22.5

The next example shows how similar triangles are used with maps.

Example 2.6.9

On a map, San Francisco, Las Vegas, and Los Angeles form a triangle whose sides are shown in the figure below. If the actual distance from Los Angeles to Las Vegas is 270 miles find the distance from Los Angeles to San Francisco.




Solution

Read the problem. Draw the figures and label with the given information.	The figures are shown above.
Identify what we are looking for.	The actual distance from Los Angeles to San Francisco.

Name the variables.	Let x= distance from Los Angeles to San Francisco.
Translate into an equation. Since the triangles are similar, the corresponding sides are proportional. We'll make the numerators "miles" and the denominators "inches."	$\frac{x \text{ miles}}{1.3 \text{ inches}} = \frac{270 \text{ miles}}{1 \text{ inch}}$
	$1.3\left(\frac{x}{1.3}\right) = 1.3\left(\frac{270}{1}\right)$
Solve the equation.	$x = 351$
Check.	
On the map, the distance from Los Angeles to San Francisco is more than the distance from Los Angeles to Las Vegas. Since 351 is more than 270 the answer makes sense.	
Check $x = 351$ in the original proportion. Use a calculator. $\frac{x \text{ miles}}{1.3 \text{ inches}} = \frac{270 \text{ miles}}{1 \text{ inch}}$ $\frac{351 \text{ miles}}{1.3 \text{ inches}} \stackrel{?}{=} \frac{270 \text{ miles}}{1 \text{ inch}}$ $\frac{270 \text{ miles}}{1 \text{ inch}} = \frac{270 \text{ miles}}{1 \text{ inch}}$ ✓	
Answer the question.	The distance from Los Angeles to San Francisco is 351 miles.

Try It 2.6.17

On the map, Seattle, Portland, and Boise form a triangle whose sides are shown in the figure below. If the actual distance from Seattle to Boise is 400 miles, find the distance from Seattle to Portland.



Answer

150 miles

Try It 2.6.18

Using the map above, find the distance from Portland to Boise.

Answer


350 miles

We can use **similar figures** to find heights that we cannot directly measure.

Example 2.6.10

Tyler is 6 feet tall. Late one afternoon, his shadow was 8 feet long. At the same time, the shadow of a tree was 24 feet long. Find the height of the tree.

Solution

Read the problem and draw a figure.	
We are looking for h, the height of the tree.	
We will use similar triangles to write an equation.	
The small triangle is similar to the large triangle.	$\frac{h}{24}=\frac{6}{8}$
Solve the proportion.	$24\left(\frac{6}{8}\right)=24\left(\frac{h}{24}\right)$
Simplify.	$18=h$
Check.	
Tyler's height is less than his shadow's length so it makes sense that the tree's height is less than the length of its shadow.	
Check $h=18$ in the original proportion. $\frac{6}{8}=\frac{h}{24}$ $\frac{6}{8}\stackrel{?}{=}\frac{18}{24}$ $\frac{3}{4}=\frac{3}{4}$ ✓	

Try It 2.6.19

A telephone pole casts a shadow that is 50 feet long. Nearby, an 8 foot tall traffic sign casts a shadow that is 10 feet long. How tall is the telephone pole?

Answer

40 feet

Try It 2.6.20

A pine tree casts a shadow of 80 feet next to a 30-foot tall building which casts a 40 feet shadow. How tall is the pine tree?

Answer

60 feet

Key Concepts

- **Property of Similar Triangles**
  - If ΔABC is similar to ΔXYZ
- **Problem Solving Strategy for Geometry Applications**

1. **Read** the problem and make sure all the words and ideas are understood. Draw the figure and label it with the given information.
2. **Identify** what we are looking for.
3. **Name** what we are looking for by choosing a variable to represent it.
4. **Translate** into an equation by writing the appropriate formula or model for the situation. Substitute in the given information.
5. **Solve the equation** using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Glossary

proportion

A proportion is an equation of the form $\frac{a}{b} = \frac{c}{d}, where \(b \neq 0$, $d \neq 0$. The proportion is read “a is to b c is to d."

similar figures

Two figures are similar if the measures of their corresponding angles are equal and their corresponding sides are in the same ratio.

This page titled 2.6: Solve Proportions and Similar Figure Applications is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **8.7: Solve Proportion and Similar Figure Applications** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.6E: Exercises

Practice Makes Perfect

Solve Proportions

In the following exercises, solve.

Example 2.6E.31

$\frac{x}{56} = \frac{7}{8}$

Answer

49

Example 2.6E.32

$\frac{n}{91} = \frac{8}{13}$

Example 2.6E.32

$\frac{49}{63} = \frac{z}{9}$

Answer

7

Example 2.6E.33

$\frac{56}{72} = \frac{y}{9}$

Example 2.6E.34

$\frac{5}{a} = \frac{65}{11}$

Answer

9

Example 2.6E.35

$\frac{4}{b} = \frac{64}{144}$

Example 2.6E.36

$\frac{98}{154} = -\frac{7}{p}$

Answer

-11

Example 2.6E.37

$$\frac{72}{156} = -\frac{6}{q}$$

Example 2.6E.38

$$\frac{a}{-8} = \frac{-42}{48}$$

Answer

7

Example 2.6E.39

$$\frac{b}{-7} = \frac{-30}{42}$$

Example 2.6E.40

$$\frac{2.7}{j} = \frac{0.9}{0.2}$$

Answer

0.6

Example 2.6E.41

$$\frac{2.8}{k} = \frac{2.1}{1.5}$$

Example 2.6E.42

$$\frac{a}{a+12} = \frac{4}{7}$$

Answer

16

Example 2.6E.43

$$\frac{b}{b-16} = \frac{11}{9}$$

Example 2.6E.44

$$\frac{c}{c-104} = -\frac{5}{8}$$

Answer

$-\frac{5}{8}$

Example 2.6E.45

$$\frac{d}{d-48}=-\frac{13}{3}$$

Example 2.6E.46

$$\frac{m+90}{25}=\frac{m+30}{15}$$

Answer

60

Example 2.6E.47

$$\frac{n+10}{4}=\frac{40-n}{6}$$

Example 2.6E.48

$$\frac{2p+4}{8}=\frac{p+18}{6}$$

Answer

30

Example 2.6E.49

$$\frac{q-2}{2}=\frac{2q-7}{18}$$

Example 2.6E.50

Pediatricians prescribe 5 milliliters (ml) of acetaminophen for every 25 pounds of a child's weight. How many milliliters of acetaminophen will the doctor prescribe for Jocelyn, who weighs 45 pounds?

Answer

9 ml

Example 2.6E.51

Brianna, who weighs 6 kg, just received her shots and needs a pain killer. The pain killer is prescribed for children at 15 milligrams (mg) for every 1 kilogram (kg) of the child's weight. How many milligrams will the doctor prescribe?

Example 2.6E.52

A veterinarian prescribed Sunny, a 65 pound dog, an antibacterial medicine in case an infection emerges after her teeth were cleaned. If the dosage is 5 mg for every pound, how much medicine was Sunny given?

Answer

325 mg

Example 2.6E.53

Belle, a 13 pound cat, is suffering from joint pain. How much medicine should the veterinarian prescribe if the dosage is 1.8 mg per pound?

Example 2.6E.54

A new energy drink advertises 106 calories for 8 ounces. How many calories are in 12 ounces of the drink?

Answer

159 calories

Example 2.6E.55

One 12 ounce can of soda has 150 calories. If Josiah drinks the big 32 ounce size from the local mini-mart, how many calories does he get?

Example 2.6E.56

A new 7 ounce lemon ice drink is advertised for having only 140 calories. How many ounces could Sally drink if she wanted to drink just 100 calories?

Answer

5 oz

Example 2.6E.57

Reese loves to drink healthy green smoothies. A 16 ounce serving of smoothie has 170 calories. Reese drinks 24 ounces of these smoothies in one day. How many calories of smoothie is he consuming in one day?

Example 2.6E.58

Janice is traveling to Canada and will change $250 US dollars into Canadian dollars. At the current exchange rate, $1 US is equal to $1.01 Canadian. How many Canadian dollars will she get for her trip?

Answer

252.5 Canadian dollars

Example 2.6E.59

Todd is traveling to Mexico and needs to exchange $450 into Mexican pesos. If each dollar is worth 12.29 pesos, how many pesos will he get for his trip?

Example 2.6E.60

Steve changed $600 into 480 Euros. How many Euros did he receive for each US dollar?

Answer

0.80 Euros

Example 2.6E.61

Martha changed $350 US into 385 Australian dollars. How many Australian dollars did she receive for each US dollar?

Example 2.6E.62

When traveling to Great Britain, Bethany exchanged her $900 into 570 British pounds. How many pounds did she receive for each American dollar?

Answer

0.63 British pounds

Example 2.6E.63

A missionary commissioned to South Africa had to exchange his $500 for the South African Rand which is worth 12.63 for every dollar. How many Rand did he have after the exchange?

Example 2.6E.64

Ronald needs a morning breakfast drink that will give him at least 390 calories. Orange juice has 130 calories in one cup. How many cups does he need to drink to reach his calorie goal?

Answer

3 cups

Example 2.6E.65

Sarah drinks a 32-ounce energy drink containing 80 calories per 12 ounce. How many calories did she drink?

Example 2.6E.66

Elizabeth is returning to the United States from Canada. She changes the remaining 300 Canadian dollars she has to $230.05 in American dollars. What was $1 worth in Canadian dollars?

Answer

1.30 Canadian dollars

Example 2.6E.67

Ben needs to convert $1000 to the Japanese Yen. One American dollar is worth 123.3 Yen. How much Yen will he have?

Example 2.6E.68

A golden retriever weighing 85 pounds has diarrhea. His medicine is prescribed as 1 teaspoon per 5 pounds. How much medicine should he be given?

Answer

17 tsp

Example 2.6E.69

Five-year-old Lacy was stung by a bee. The dosage for the anti-itch liquid is 150 mg for her weight of 40 pounds. What is the dosage per pound?

Example 2.6E.70

Karen eats $\frac{1}{2}$ cup of oatmeal that counts for 2 points on her weight loss program. Her husband, Joe, can have 3 points of oatmeal for breakfast. How much oatmeal can he have?

Answer


$\frac{3}{4}$ cup

Example 2.6E.71

An oatmeal cookie recipe calls for $\frac{1}{2}$ cup of butter to make 4 dozen cookies. Hilda needs to make 10 dozen cookies for the bake sale. How many cups of butter will she need?

Solve Similar Figure Applications

In the following exercises, ΔABC is similar to ΔXYZ



Example 2.6E.72

side *b*


Answer

12

Example 2.6E.73

side *x*

In the following exercises, ΔDEF is similar to ΔNPQ



Example 2.6E.74

Find the length of side *d*.


Answer

$\frac{77}{18}$

Example 2.6E.75

Find the length of side q.

In the following two exercises, use the map shown. On the map, New York City, Chicago, and Memphis form a triangle whose sides are shown in the figure below. The actual distance from New York to Chicago is 800 miles.



Example 2.6E.76

Find the actual distance from New York to Memphis.


Answer

950 miles

Example 2.6E.77

Find the actual distance from Chicago to Memphis.

In the following two exercises, use the map shown. On the map, Atlanta, Miami, and New Orleans form a triangle whose sides are shown in the figure below. The actual distance from Atlanta to New Orleans is 420 miles.



Example 2.6E.78

Find the actual distance from New Orleans to Miami.

Answer

680 miles

Example 2.6E.79

Find the actual distance from Atlanta to Miami.

Example 2.6E.80

A 2 foot tall dog casts a 3 foot shadow at the same time a cat casts a one foot shadow. How tall is the cat?

Answer

$\frac{2}{3}$ foot (8in)

Example 2.6E.81

Larry and Tom were standing next to each other in the backyard when Tom challenged Larry to guess how tall he was. Larry knew his own height is 6.5 feet and when they measured their shadows, Larry's shadow was 8 feet and Tom's was 7.75 feet long. What is Tom's height?

Example 2.6E.82

The tower portion of a windmill is 212 feet tall. A six foot tall person standing next to the tower casts a seven foot shadow. How long is the windmill's shadow?

Answer

247.3 feet

Example 2.6E.83

The height of the Statue of Liberty is 305 feet. Nicole, who is standing next to the statue, casts a 6 foot shadow and she is 5 feet tall. How long should the shadow of the statue be?

Everyday Math

Example 2.6E.84

Heart Rate At the gym, Carol takes her pulse for 10 seconds and counts 19 beats.

1. How many beats per minute is this?
2. Has Carol met her target heart rate of 140 beats per minute?

Answer

1. 114 beats per minute
2. no

Example 2.6E.85

Heart Rate Kevin wants to keep his heart rate at 160 beats per minute while training. During his workout he counts 27 beats in 10 seconds.

1. How many beats per minute is this?
2. Has Kevin met his target heart rate?

Example 2.6E.86

Cost of a Road Trip Jesse's car gets 30 miles per gallon of gas.

1. If Las Vegas is 285 miles away, how many gallons of gas are needed to get there and then home?
2. If gas is $3.09 per gallon, what is the total cost of the gas for the trip?

Answer

1. 19 gallons
2. $58.71

Example 2.6E.87

Cost of a Road Trip Danny wants to drive to Phoenix to see his grandfather. Phoenix is 370 miles from Danny's home and his car gets 18.5 miles per gallon.

1. How many gallons of gas will Danny need to get to and from Phoenix?
2. If gas is $3.19 per gallon, what is the total cost for the gas to drive to see his grandfather?

Example 2.6E.88

Lawn Fertilizer Phil wants to fertilize his lawn. Each bag of fertilizer covers about 4,000 square feet of lawn. Phil's lawn is approximately 13,500 square feet. How many bags of fertilizer will he have to buy?

Answer

4 bags

Example 2.6E.89

House Paint April wants to paint the exterior of her house. One gallon of paint covers about 350 square feet, and the exterior of the house measures approximately 2000 square feet. How many gallons of paint will she have to buy?

Example 2.6E.90

Cooking Natalia's pasta recipe calls for 2 pounds of pasta for 1 quart of sauce. How many pounds of pasta should Natalia cook if she has 2.5 quarts of sauce?

Answer

5

Example 2.6E.91

Heating Oil A 275 gallon oil tank costs $400 to fill. How much would it cost to fill a 180 gallon oil tank?

Writing Exercises

Example 2.6E.92

Marisol solves the proportion $\frac{144}{a}=\frac{9}{4}$ by 'cross multiplying', so her first step looks like 4·144=9·a. Explain how this differs from the method of solution shown in Example.

Answer

Answers will vary.

Example 2.6E.93

Find a printed map and then write and solve an application problem similar to Example.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve proportions.			
solve similar figure applications.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 2.6E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **8.7E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

2.7: Solve Linear Inequalities

In Chapter 1, we introduced the natural numbers $\mathbb{N} = \{1, 2, 3, \dots\}$, the whole numbers $\mathbb{W} = \{0, 1, 2, 3, \dots\}$, and the integers $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$. Later in the chapter, we introduced the rational numbers, numbers of the form p/q, where p and q are integers. We noted that both terminating and repeating decimals are rational numbers. Each of these numbers has a unique position on the number line (see Figure 2.7.1).




Figure 2.7.1: Positioning numbers on the number line.

The natural numbers, whole numbers, and integers are also rational numbers, because each can be expressed in the form $\frac{p}{q}$, where p and q are integers. For example, $0 = \frac{0}{12}$, $4 = \frac{4}{1}$, and $-3 = \frac{-12}{4}$. Indeed, the rational numbers contain all of the numbers we've studied up to this point in the course. However, not all numbers are rational numbers. For example, consider the decimal number $-3.10110111011110\ldots$, which neither terminates nor repeats. The number $\sqrt{2} = 1.414213562373095\ldots$ also equals a decimal number that never terminates and never repeats. A similar statement can be made about the number $\pi = 3.141592653589793\ldots$ Each of these irrational (not rational) numbers also has a unique position on the number line (see Figure 2.7.2).




Figure 2.7.2: Positioning numbers on the number line.

Two other irrational numbers you may encounter in your mathematical studies are e (Euler's constant), which is approximately equal to $e \approx 2.71828182845904\ldots$, and ϕ (pronounced "phi"), called the golden ratio, which equals $\phi = \frac{1+\sqrt{5}}{2}$. The number e arises in applications involving compound interest, probability, and other areas of mathematics. The number ϕ is used in financial markets and is also arguably the ratio of beauty in art and architecture.

The Real Numbers

If we combine all of the rational and irrational numbers into one collection, then we have a set of numbers that is called the set of real numbers. The set of real numbers is denoted by the symbol $\mathbb{R}$.

Every point on the number line is associated with a unique real number. Conversely, every real number is associated with a unique position on the number line. In lieu of this correspondence, the number line is usually called the real line.

Ordering the Real Numbers

The real numbers are ordered on the real line in a manner identical to how we ordered the integers on the number line in Section 1 of Chapter 1.

Order on the Real Line

Suppose that a and b are real numbers positioned on the real line as shown below.



- Because a lies to the "left of" b, we say that a is "less than" b, or in mathematical symbols, $a < b$. The inequality symbol $<$ is read "less than."
- Alternately, b lies to the "right of" a, so we can also say that b is "greater than" a, or in mathematical symbols, $b > a$. The inequality symbol $>$ is read "greater than."

Here are two more inequality symbols that we will use in this section.

Less than or equal to

If we want to say that a lies to the "left of" b, or shares the same position as b, then we say that a is "less than or equal to" b and write $a \leq b$. The inequality symbol $\leq$ is pronounced "less than or equal to."

Greater than or equal to

If we want to say that b lies to the "right of" a, or shares the same position as a, then we say that b is "greater than or equal to" a and write $b \geq a$. The inequality symbol $\geq$ is pronounced "greater than or equal to."

Set-Builder Notation

Mathematicians use a construct called set-builder notation to describe sets or collections of numbers. The general form of set-builder notation looks as follows:

$$\{x \mid \text{ some statement about } x\}$$

For example, suppose that we want to describe the set of "all real numbers that are less than 2." We could use the following notation:

$$A = \{x \mid x < 2\}$$

This is read aloud as follows: "A equals the set of all x such that x is less than 2." Some prefer to use a colon instead of a vertical bar.

$$A = \{x : x < 2\}$$

In this text we use the vertical bar in set-builder notation, but feel free to use a colon instead. They mean the same thing. One might still object that the notation

$$\{x \mid x < 2\}$$

is a bit vague. One objection could be "What type of numbers x are you referring to? Do you want the integers that are less than two or do you want the real numbers that are less than two?" As you can see, this is a valid objection. One way of addressing this objection is to write:

$$A = \{x \in \mathbb{R} \mid x < 2\} \quad \text{or} \quad A = \{x \in \mathbb{N} \mid x < 2\}$$

The first is read "A is the set of all x in R that are less than two," while the second is read "A is the set of all x in N that are less than two."

Set-builder Assumption

In this text, unless there is a specific reference to the set of numbers desired, we will assume that the notation $\{x \mid x < 2\}$ is asking for the set of all real numbers less than 2.

In Figure 2.7.3, we've shaded the set of real numbers $\{x \mid x < 2\}$. Because




Figure 2.7.3: Shading the numbers less than 2.

"less than" is the same as saying "left of," we've shaded (in red) all points on the real line that lie to the left of the number two. Note that there is an "empty circle" at the number two. The point representing the number two is not shaded because we were only asked to shade the numbers that are strictly less than two.

While the shading in Figure 2.7.3 is perfectly valid, much of the information provided in Figure 2.7.3 is unnecessary (and perhaps distracting). We only need to label the endpoint and shade the real numbers to the left of two, as we've done in constructing Figure 2.7.4.



Figure 2.7.4: You only need to label the endpoint.

For contrast, suppose instead that we're asked to shade the set of real numbers $\{x \mid x \leq 2\}$. This means that we must shade all the real numbers



Figure 2.7.5: You only need to label the endpoint.

that are "less than or equal to 2" or "left of and including 2." The resulting set is shaded in Figure 2.7.5.


Note the difference between Figures 2.7.4 and 2.7.45. In Figures 2.7.4 we're shading the set $\{x \mid x < 2\}$, so the number 2 is left unshaded (an empty dot). In Figures 2.7.5, we're shading the set $\{x \mid x \leq 2\}$, so the number 2 is shaded (a filled-in dot).

Example 2.7.1

Shade the set $\{x \mid x \geq -3\}$ on the real line.

Solution

The notation $\{x \mid x \geq -3\}$ is pronounced "the set of all real numbers x such that x is greater than or equal to -3." Thus, we need to shade the number -3 and all real numbers to the right of -3.



Try It 2.7.1


Shade $\{x \mid x \leq 4\}$ on the real line.

Answer



Example 2.7.2

Use set-builder notation to describe the set of real numbers that are shaded on the number line below.




Solution

The number -1 is not shaded. Only the numbers to the left of -1 are shaded. This is the set of all real numbers x such that x is "less than" -1. Thus, we describe this set with the following set-builder notation:

$$\{x \mid x < -1\}$$

Try It 2.7.2

Use set-builder notation to describe the following set of real numbers:




Answer

$\{x \mid x > -10\}$

Interval Notation


In Examples 2.7.1 and 2.7.2, we used set-builder notation to describe the set of real numbers greater than or equal to -3 and a second set of real numbers less than -1. There is another mathematical symbolism, called interval notation, that can be used to describe these sets of real numbers. Consider the first set of numbers from Example 2.7.1, $\{x \mid x \geq -3\}$.



Sweeping our eyes "from left to right", we use $[-3, \infty)$ to describe this set of real numbers. Some comments are in order:

1. The bracket at the left end means that -3 is included in the set.
2. As you move toward the right end of the real line, the numbers grow without bound. Hence, the ∞ symbol (positive infinity) is used to indicate that we are including all real numbers to the right of -3. However, ∞ is not really a number, so we use a parentheses to indicate we are "not including" this fictional point.

The set of numbers from Example 2.7.1 is $\{x \mid x < -1\}$.



Sweeping our eyes "from left to right", this set of real numbers is described with $(-\infty, -1)$. Again, comments are in order:

1. The number -1 is not included in this set. To indicate that it is not included, we use a parenthesis.
2. As you move toward the left end of the real line, the numbers decrease without bound. Hence, the $-\infty$ symbol (negative infinity) is used to indicate that we are including all real numbers to the left of -1. Again, $-\infty$ is not an actual number, so we use a parenthesis to indicate that we are not including this "fictional" point.

Sweep your eyes from "left to right"

If you would like to insure that you correctly use interval notation, place the numbers in your interval notation in the same order as they are encountered as you sweep your eyes from "left to right" on the real line.

A nice summary of set-builder and interval notation is presented in Table 2.7.1 at the end of the section.

Equivalent Inequalities

Like equations, two inequalities are equivalent if they have the same solution sets.

Adding or Subtracting the Same Quantity from Both Sides of an Inequality

Let a and b be real numbers with $a < b$
If c is any real number, then $a + c < b + c$ and $a - c < b - c$.

That is, adding or subtracting the same amount from both sides of an inequality produces an equivalent inequality (does not change the solution).

Example 2.7.3

Solve for x : $x - 2 \leq 7$. Sketch the solution on the real line, then use set-builder and interval notation to describe your solution.

Solution

To "undo" subtracting 2, we add 2 to both sides of the inequality.

$$\begin{aligned} x - 2 &\leq 7 && \text{Original inequality.} \\ x - 2 + 2 &\leq 7 + 2 && \text{Add 2 to both sides.} \\ x &\leq 9 && \text{Simplify both sides.} \end{aligned}$$

To shade the real numbers less than or equal to 9, we shade the number 9 and all real numbers to the left of 9.



Using set-builder notation, the solution is $\{x \mid x \leq 9\}$. Using interval notation, the solution is $(-\infty, 9]$.

Try It 2.7.3

Use use set-builder and interval notation to describe the solution of: $x - 7 < -8$.

Answer

Using set-builder notation, the solution is $\{x \mid x < -1\}$. Using interval notation, the solution is $(-\infty, -1)$

If we multiply or divide both sides of an inequality by a positive number, we have an equivalent inequality.

Multiplying or Dividing by a Positive Number

Let a and b be real numbers with $a < b$.

If c is a real positive number, then $ac < bc$ and $\frac{a}{c} < \frac{b}{c}$

Example 2.7.4


Solve for x : $3x \leq -9$ Sketch the solution on the real line, then use set-builder and interval notation to describe your solution.

Solution

To "undo" multiplying by 3, divide both sides of the inequality by 3. Because we are dividing both sides by a positive number, we do not reverse the inequality sign.

$$\begin{aligned} 3x &\leq -9 && \text{Original inequality.} \\ \frac{3x}{3} &\leq \frac{-9}{3} && \text{Divide both sides by 3.} \\ x &\leq -3 && \text{Simplify both sides.} \end{aligned}$$

Shade the real numbers less than or equal to -3.



Using set-builder notation, the solution is $\{x \mid x \leq -3\}$. Using interval notation, the solution is $(-\infty, -3]$.

Try It 2.7.4

Use use set-builder and interval notation to describe the solution of:

$$2x > -8$$

Answer

Using set-builder notation, the solution is $\{x \mid x > -4\}$. Using interval notation, the solution is $(-4, \infty)$

Reversing the Inequality Sign

Up to this point, it seems that the technique for solving inequalities is pretty much identical to the technique used to solve equations. However, in this section we are going to encounter one exception.

Suppose we start with the valid inequality $-2 < 5$, then we multiply both sides by 2, 3, and 4.

$$\begin{array}{rrr} -2<5 & -2<5 & -2<5 \\ 2(-2)<2(5) & 3(-2)<3(5) & 4(-2)<4(5) \\ -4<10 & -6<15 & -8<20 \end{array}$$

Note that in each case, the resulting inequality is still valid.

Caution! We're about to make an error!

Start again with $-2<5$, but this time multiply both sides by -2, -3, and -4.

$$\begin{array}{rrr} -2<5 & -2<5 & -2<5 \\ -2(-2)<-2(5) & -3(-2)<-3(5) & -4(-2)<-4(5) \\ 4<-10 & 6<-15 & 8<-20 \end{array}$$

In each of the resulting inequalities, the inequality symbol is pointing the wrong way!

When you multiply both sides of an inequality by a negative number, you must reverse the inequality sign. Starting with $-2<5$, multiply both sides by -2, -3, and -4, but reverse the inequality symbol.

Some readers might prefer a more formal reason as to why we reverse the inequality when we multiply both sides by a negative number. Suppose that $a<b$. Then, subtracting b from both sides gives the result $a-b<0$. This means that $a-b$ is a negative number. Now, if c is a negative number, then the product $(a-b)c$ is positive. Then:

$$\begin{aligned} a &< b \\ a-b &< 0 \\ (a-b)c &> 0 \\ ac-bc &> 0 \\ ac-bc+bc &> 0+bc \\ ac &> bc \end{aligned}$$

Thus, if you start with $a<b$ and $c<0$, then $ac>bc$.

Multiplying or Dividing by a Negative Number

Let a and b be real numbers with $a<b$.

If c is a real negative number, then \[a c>b c \) and $\frac{a}{c}>\frac{b}{c}$

That is, when multiplying or dividing both sides of an inequality by a negative number, you must **reverse** the inequality sign.

Example 2.7.5


Solve for x: $-2x<4$. Sketch the solution on the real line, then use set-builder and interval notation to describe your solution.

Solution

To "undo" multiplying by -2, divide both sides by -2. Because we are dividing both sides by a negative number, we reverse the inequality sign.

$$\begin{array}{rl} -2x<4 & \text{Original inequality.} \\ \frac{-2x}{-2}>\frac{4}{-2} & \text{Divide both sides by } -2 \\ x>-2 & \text{Reverse the inequality sign.} \\ x>-2 & \text{Simplify both sides.} \end{array}$$

Shade the real numbers greater than -2.



Using set-builder notation, the solution is $\{x \mid x > -2\}$. Using interval notation, the solution is $(-2, \infty)$.

Try It 2.7.5

Use use set-builder and interval notation to describe the solution of:

$$-3x \geq -6$$

Answer

Using set-builder notation, the solution is $\{x \mid x \leq 2\}$. Using interval notation, the solution is $(-\infty, 2]$

Multiple Steps

Sometimes you need to perform a sequence of steps to arrive at the solution.

Example 2.7.6

Solve for x: $2x + 5 > -7$. Sketch the solution on the real line, then use set-builder and interval notation to describe your solution.

Solution


To "undo" adding 5, subtract 5 from both sides of the inequality.

$$\begin{aligned} 2x + 5 &> -7 && \text{Original inequality.} \\ 2x + 5 - 5 &> -7 - 5 && \text{Subtract 5 from both sides.} \\ 2x &> -12 && \text{Simplify both sides.} \end{aligned}$$

To "undo" multiplying by 2, divide both sides by 2. Because we are dividing both sides by a positive number, we do not reverse the inequality sign.

$$\begin{aligned} \frac{2x}{2} &> \frac{-12}{2} && \text{Divide both sides by 2} \\ x &> -6 && \text{Simplify both sides.} \end{aligned}$$

Shade the real numbers greater than -6.



Using set-builder notation, the solution is $\{x \mid x > -6\}$. Using interval notation, the solution is $(-6, \infty)$.

Try It 2.7.6

Use use set-builder and interval notation to describe the solution of:

$$3x - 2 \leq 4$$

Answer

Using set-builder notation, the solution is $\{x \mid x \leq 2\}$. Using interval notation, the solution is $(-\infty, 2]$

Example 2.7.7

Solve for x: $3 - 5x \leq 2x + 17$. Sketch the solution on the real line, then use set-builder and interval notation to describe your solution.

Solution

We need to isolate terms containing x on one side of the inequality. Start by subtracting $2x$ from both sides of the inequality.


$$\begin{aligned} 3-5x &\leq 2x+17 && \text{Original inequality.} \\ 3-5x-2x &\leq 2x+17-2x && \text{Subtract } 2x \text{ from both sides.} \\ 3-7x &\leq 17 && \text{Simplify both sides.} \end{aligned}$$

We continue to isolate terms containing x on one side of the inequality. Subtract 3 from both sides.

$$\begin{aligned} 3-7x-3 &\leq 17-3 && \text{Subtract 3 from both sides.} \\ -7x &\leq 14 && \text{Simplify both sides.} \end{aligned}$$

To "undo" multiplying by -7, divide both sides by -7. Because we are dividing both sides by a negative number, we reverse the inequality sign.

$$\begin{aligned} \frac{-7x}{-7} &\geq \frac{14}{-7} && \text{Divide both sides by } -7 \\ x &\geq -2 && \text{Simplify both sides.} \end{aligned}$$



Using set-builder notation, the solution is $\{x \mid x \geq -2\}$. Using interval notation, the solution is $[-2, \infty)$.

? Try It 2.7.7

Use use set-builder and interval notation to describe the solution of:

$$4-x>2x+1$$

Answer

Using set-builder notation, the solution is $\{x \mid x<1\}$. Using interval notation, the solution is $(-\infty, 1)$

We clear fractions from an inequality in the usual manner, by multiplying both sides by the least common denominator.

✓ Example 2.7.8

Solve for x : $\quad \frac{3}{4}-\frac{x}{12}>\frac{1}{3}$.

Solution

First, clear the fractions from the inequality by multiplying both sides by the least common denominator, which in this case is 12.

$$\begin{aligned} \frac{3}{4}-\frac{x}{12} &> \frac{1}{3} && \text{Original inequality.} \\ 12\left[\frac{3}{4}-\frac{x}{12}\right] &> \left[\frac{1}{3}\right]12 && \text{Multiply both sides by 12.} \\ 12\left[\frac{3}{4}\right]-12\left[\frac{x}{12}\right] &> \left[\frac{1}{3}\right]12 && \text{Distribute the 12.} \\ 9-x &> 4 && \text{Cancel and Multiply.} \end{aligned}$$


To "undo" adding 9, subtract 9 from both sides.

$$\begin{aligned} 9-x-9 &> 4-9 && \text{Subtract 9 from both sides.} \\ -x &> -5 && \text{Simplify both sides.} \end{aligned}$$

We could divide both sides by -1, but multiplying both sides by -1 will also do the job. Because we are multiplying both sides by a negative number, we reverse the inequality sign.

$$\begin{aligned} (-1)(-x) &< (-5)(-1) && \text{Multiply both sides by } -1. \text{ Reverse the inequality sign.} \\ x &< 5 && \text{Simplify both sides.} \end{aligned}$$

Shade the real numbers less than 5.



Using set-builder notation, the solution is $\{x \mid x < 5\}$. Using interval notation, the solution is $(-\infty, 5)$.

Try It 2.7.8

Use use set-builder and interval notation to describe the solution of:

$$\frac{2x}{3} - \frac{3}{4} \geq -\frac{3}{2}$$

Answer

Using set-builder notation, the solution is $\{x \mid x \geq -\frac{9}{8}\}$. Using interval notation, the solution is $[-\frac{9}{8}, \infty)$

We clear decimals from an inequality in the usual manner, by multiplying both sides by the appropriate power of ten.

Example 2.7.9

Solve for x : $3.25 - 1.2x > 4.6$.

Solution

First, clear the decimals from the inequality by multiplying both sides by 100, which moves each decimal point two places to the right.

$$\begin{aligned} 3.25 - 1.2x &> 4.6 && \text{Original inequality.} \\ 325 - 120x &> 460 && \text{Multiply both sides by 100.} \\ 325 - 120x - 325 &> 460 - 325 && \text{Subtract 325 from both sides.} \\ -120x &> 135 && \text{Simplify both sides.} \\ \frac{-120x}{-120} &< \frac{135}{-120} && \text{Divide both sides by } -120. \text{ Reverse the inequality sign.} \\ x &< -\frac{9}{8} && \text{Reduce to lowest terms.} \end{aligned}$$

Shade the real numbers less than $-9/8$.



Using set-builder notation, the solution is $\{x \mid x < -9/8\}$. Using interval notation, the solution is $(-\infty, -9/8)$.

Try It 2.7.9

Use use set-builder and interval notation to describe the solution of:

$$2.3x - 5.62 \geq -1.4$$

Answer

Using set-builder notation, the solution is $\{x \mid x \geq 1.83\}$. Using interval notation, the solution is $[\frac{211}{115}, \infty)$

Solve Applications with Linear Inequalities

Many real-life situations require us to solve inequalities. The method we will use to solve applications with linear inequalities is very much like the one we used when we solved applications with equations.

We will read the problem and make sure all the words are understood. Next, we will identify what we are looking for and assign a variable to represent it. We will restate the problem in one sentence to make it easy to translate into an inequality. Then, we will solve the inequality.

Sometimes an application requires the solution to be a whole number, but the algebraic solution to the inequality is not a whole number. In that case, we must round the algebraic solution to a whole number. The context of the application will determine whether we round up or down.

Example 2.7.28

Dawn won a mini-grant of $4,000 to buy tablet computers for her classroom. The tablets she would like to buy cost $254.12 each, including tax and delivery. What is the maximum number of tablets Dawn can buy?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the maximum number of tablets Dawn can buy
Step 3. Name what you are looking for. Choose a variable to represent that quantity.	Let n = the number of tablets.
Step 4. Translate Write a sentence that gives the information to find it.	$254.12 times the number of tablets is no more than $4,000.
Translate into an inequality.	$254.12n \leq 4000$
Step 5. Solve the inequality.	$n \leq 15.74$ but n must be a whole number of tablets, so round to 15. $n \leq 15$ tablets
Step 6. Check the answer in the problem and make sure it makes sense.	Rounding down the price to $250, 15 tablets would cost $3,750, while 16 tablets would be $4,000. So a maximum of 15 tablets at $254.12 seems reasonable.
Step 7. Write a sentence that answers the question.	Dawn can buy a maximum of 15 tablets.

Try It 2.7.29

Angie has $20 to spend on juice boxes for her son's preschool picnic. Each pack of juice boxes costs $2.63. What is the maximum number of packs she can buy?

Answer

Angie can buy 7 packs of juice.

Try It 2.7.30

Daniel wants to surprise his girlfriend with a birthday party at her favorite restaurant. It will cost $42.75 per person for dinner, including tip and tax. His budget for the party is $500. What is the maximum number of people Daniel can have at the party?

Answer

Daniel can have 11 people at the party.

Example 2.7.31

Taleisha's phone plan costs her $28.80 a month plus $0.20 per text message. How many text messages can she send/receive and keep her monthly phone bill no more than $50?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the number of text messages Taleisha can make
Step 3. Name what you are looking for. Choose a variable to represent that quantity.	Let $t =$ the number of text messages.
Step 4. Translate Write a sentence that gives the information to find it.	$28.80 plus $0.20 times the number of text messages is less than or equal to $50.
Translate into an inequality.	$28.80 + 0.20t \leq 50$
Step 5. Solve the inequality.	$0.2t \leq 21.2$ $t \leq 106$ text messages
Step 6. Check the answer in the problem and make sure it makes sense.	Yes, $28.80 + 0.20(106) = 50$.
Step 7. Write a sentence that answers the question.	Taleisha can send/receive no more than 106 text messages to keep her bill no more than $50.

Try It 2.7.32

Sergio and Lizeth have a very tight vacation budget. They plan to rent a car from a company that charges $75 a week plus $0.25 a mile. How many miles can they travel during the week and still keep within their $200 budget?

Answer

Sergio and Lizeth can travel no more than 500 miles.

Try It 2.7.33

Rameen's heating bill is $5.42 per month plus $1.08 per therm. How many therms can Rameen use if he wants his heating bill to be a maximum of $87.50.

Answer

Rameen's heating bill is $5.42 per month plus $1.08 per therm. How many therms can Rameen use if he wants his heating bill to be a maximum of $87.50.

Profit is the money that remains when the costs have been subtracted from the revenue. In the next example, we will find the number of jobs a small businesswoman needs to do every month in order to make a certain amount of profit.

Example 2.7.34

Felicity has a calligraphy business. She charges $2.50 per wedding invitation. Her monthly expenses are $650. How many invitations must she write to earn a profit of at least $2,800 per month?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the number of invitations Felicity needs to write

Step 3. Name what you are looking for. Choose a variable to represent that quantity.	Let $j =$ the number of invitations.
Step 4. Translate Write a sentence that gives the information to find it.	$2.50 times the number of invitations minus $650 is at least $2,800.
Translate into an inequality.	$2.50j - 650 \geq 2,800$
Step 5. Solve the inequality.	$2.5j \geq 3,450$ $j \geq 1,380$ invitations
Step 6. Check the answer in the problem and make sure it makes sense.	If Felicity wrote 1400 invitations, her profit would be $2.50(1400) - 650$, or $2,850. This is more than $2800.
Step 7. Write a sentence that answers the question.	Felicity must write at least 1,380 invitations.

Try It 2.7.35

Caleb has a pet sitting business. He charges $32 per hour. His monthly expenses are $2,272. How many hours must he work in order to earn a profit of at least $800 per month?

Answer

Caleb must work at least 96 hours.

Try It 2.7.36

Elliot has a landscape maintenance business. His monthly expenses are $1,100. If he charges $60 per job, how many jobs must he do to earn a profit of at least $4,000 a month?

Answer

Elliot must work at least 85 jobs.

There are many situations in which several quantities contribute to the total expense. We must make sure to account for all the individual expenses when we solve problems like this.

Example 2.7.37

Malik is planning a six-day summer vacation trip. He has $840 in savings, and he earns $45 per hour for tutoring. The trip will cost him $525 for airfare, $780 for food and sightseeing, and $95 per night for the hotel. How many hours must he tutor to have enough money to pay for the trip?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the number of hours Malik must tutor
Step 3. Name what you are looking for. Choose a variable to represent that quantity.	Let h = the number of hours.
Step 4. Translate Write a sentence that gives the information to find it.	The expenses must be less than or equal to the income. The cost of airfare plus the cost of food and sightseeing and the hotel bill must be less than the savings plus the amount earned tutoring.
Translate into an inequality.	$525 + 780 + 95(6) \leq 840 + 45h$

Step 5. Solve the inequality.	$1,875 \leq 840 + 45h$ $1,035 \leq 45h$ $h \geq 23$ hours
Step 6. Check the answer in the problem and make sure it makes sense.	We substitute 23 into the inequality. $1,875 \leq 840 + 45h$ $1,875 \leq 840 + 45(23)$ $1,875 \leq 1875$
Step 7. Write a sentence that answers the question.	Malik must tutor at least 23 hours.

? Try It 2.7.38

Brenda's best friend is having a destination wedding and the event will last three days. Brenda has $500 in savings and can earn $15 an hour babysitting. She expects to pay $350 airfare, $375 for food and entertainment and $60 a night for her share of a hotel room. How many hours must she babysit to have enough money to pay for the trip?

Answer

Brenda must babysit at least 27 hours.

? Try It 2.7.39

Josue wants to go on a 10-night road trip with friends next spring. It will cost him $180 for gas, $450 for food, and $49 per night to share a motel room. He has $520 in savings and can earn $30 per driveway shoveling snow. How many driveways must he shovel to have enough money to pay for the trip?




Answer

Josue must shovel at least 20 driveways.

Summary Table of Set-Builder and Interval Notation

A summary table of the set-builder and interval notation is presented in Table 2.7.1.

Table 2.7.1: Examples of set-builder and interval notation.

Shading on the real line	Set-builder	Interval
	$\{x \mid x > -5\}$	$(-5, \infty)$
	$\{x \mid x \geq -5\}$	$[-5, \infty)$
	$\{x \mid x < -5\}$	$(-\infty, -5)$
	$\{x \mid x \leq -5\}$	$(-\infty, -5]$

Contributors and Attributions

- David Arnold (Retired Professor (Mathematics) at College of the Redwoods)

This page titled 2.7: Solve Linear Inequalities is shared under a CC BY license and was authored, remixed, and/or curated by David Arnold.


- **2.6: Inequalities** by David Arnold is licensed CC BY-NC-ND 3.0. Original source: https://math.libretexts.org/api/deki/files/80188/ElementaryAlgebra.pdf.

- **8.7: Solve Proportion and Similar Figure Applications** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **2.6: Solve Linear Inequalities** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/intermediate-algebra-2e.

2.7E: Exercises for Section 2.7

1) Draw a number line, then plot the numbers $4, 3, -4, 7/8$, and $-8/3$ on your number line. Label each point with its value. Finally, list the numbers in order, from smallest to largest.


Answer



2) Draw a number line, then plot the numbers $5, 3, -4, 5/7$, and $-4/3$ on your number line. Label each point with its value. Finally, list the numbers in order, from smallest to largest.

3) Draw a number line, then plot the numbers $-5, 5, 4, 2/3$, and $8/3$ on your number line. Label each point with its value. Finally, list the numbers in order, from smallest to largest.

Answer




4) Draw a number line, then plot the numbers $-3, -2, 4, 1/3$, and $5/2$ on your number line. Label each point with its value. Finally, list the numbers in order, from smallest to largest.

In Exercises 5-20, shade each of the following sets on a number line.

5) $\{x \mid x \geq -7\}$


Answer



6) $\{x \mid x \geq -1\}$

7) $\{x \mid x < 2\}$

Answer



8) $\{x \mid x < -6\}$

9) $(-\infty, 2)$

Answer



10) $(-\infty, -9)$

11) $(6, \infty)$

Answer



12) $(5, \infty)$

13) $\{x \mid x > 7\}$

Answer



14) $\{x \mid x > -8\}$

15) $[0, \infty)$


Answer



16) $[7, \infty)$

17) $\{x \mid x \leq -2\}$

Answer



18) $\{x \mid x \leq 7\}$

19) $(-\infty, 3]$

Answer



20) $(-\infty, -1]$

In Exercises 21-28, use set-builder notation to describe the shaded region on the given number line.

21)




Answer

$\{x \mid x \leq 9\}$

22)



23)




Answer

$\{x \mid x < -8\}$

24)




25)




Answer

$\{x \mid x > -2\}$

26)



27)



Answer


$\{x \mid x \geq -3\}$

28)



In Exercises 29-36, use interval notation to describe the shaded region on the given number line.


29)




Answer

$(4, \infty)$

30)




31)



Answer

$(-\infty, -2)$

32)



33)



Answer

$(-\infty, 5]$

34)




35)



Answer

$[1, \infty)$

36)



In Exercises 37-44, solve each of the given inequalities. Sketch the solution on a number line, then use set-builder and interval notation to describe your solution.

37) $x + 10 < 19$

Answer

$(-\infty, 9)$

38) $x + 17 \geq 7$

39) $4x < 8$

Answer

$(-\infty, 2)$

40) $16x \geq -2$

41) $-2x \leq -2$

Answer

$[1, \infty)$

42) $-18x > -20$

43) $x - 18 > -10$

Answer

$(8, \infty)$

44) $x - 8 \leq -18$

In Exercises 45-62, solve each of the given inequalities. Sketch the solution on a number line, then use set-builder and interval notation to describe your solution.

45) $-5x - 6 \geq 4 - 9x$

Answer

$[5/2, \infty)$

46) $2x - 7 \geq -3 - 4x$

47) $16x - 6 \leq 18$

Answer

$(-\infty, 3/2]$

48) $8x - 14 \leq -12$

49) $-14x-6 \geq -10-4x$

Answer

$(-\infty, 2/5]$

50) $-13x-4 \geq -2-5x$

51) $5x+18 < 38$

Answer

$(-\infty, 4)$

52) $9x+16 < 79$

53) $-16x-5 \geq -11-6x$

Answer

$(-\infty, 3/5]$

54) $-11x-7 \geq -15-5x$

55) $2x-9 \geq 5-8x$

Answer

$[7/5, \infty)$

56) $-3x-6 \geq -2-9x$

57) $-10x-4 \leq 18$

Answer

$[-11/5, \infty)$

58) $-6x-14 \leq 1$

59) $-12x+4 < -56$

Answer

$(5, \infty)$

60) $-18x+6 < -12$

61) $15x+5 < 6x+2$

Answer

$(-\infty, -1/3)$

62) $12x+8 < 3x+5$

In Exercises 63-76, solve each of the given inequalities. Sketch the solution on a number line, then use set-builder and interval notation describe your solution.

63) $\frac{3}{2}x > \frac{9}{8}$

Answer

$(3/4, \infty)$

64) $\frac{6}{7}x > \frac{3}{4}$

65) $x+\frac{3}{2}<\frac{9}{5}$

Answer

$(-\infty, 3/10)$

66) $x+\frac{1}{4}<-\frac{1}{5}$

67) $\frac{4}{7}-\frac{1}{6}x \leq \frac{4}{3}x-\frac{1}{2}$

Answer

$[5/7, \infty)$

68) $\frac{5}{3}-\frac{3}{4}x \leq \frac{7}{4}x-\frac{3}{5}$

69) $x-\frac{3}{8} \geq -\frac{9}{7}$

Answer

$[-51/56, \infty)$

70) $x-\frac{7}{2} \geq \frac{1}{5}$

71) $\frac{6}{5}x \leq -\frac{4}{7}$

Answer

$[10/21, \infty)$

72) $\frac{4}{3}x \leq \frac{2}{9}$

73) $-\frac{6}{5}x-\frac{7}{3} \leq \frac{5}{9}-\frac{2}{9}x$

Answer

$[-65/22, \infty)$

74) $-\frac{3}{7}x-\frac{1}{2} \leq \frac{3}{2}-\frac{2}{7}x$

75) $\frac{9}{7}x+\frac{9}{2}>\frac{1}{7}x+\frac{7}{2}$

Answer

$(-7/8, \infty)$

76) $\frac{5}{7}x+\frac{9}{2}>\frac{1}{3}x+\frac{5}{2}$

In Exercises 77-84, solve each of the given inequalities. Sketch the solution on a number line, then use set-builder and interval notation containing fractions in reduced form to describe your solution.

77) $-3.7x-1.98 \leq 3.2$

Answer

$[-7/5, \infty)$

78) $-3.6x - 3.32 \leq 0.8$

79) $-3.4x + 3.5 \geq 0.9 - 2.2x$

Answer

$(-\infty, 13/6]$

80) $-2.6x + 3.1 \geq -2.9 - 1.7x$

81) $-1.3x + 2.9 > -2.6 - 3.3x$

Answer

$(-11/4, \infty)$

82) $2.5x + 2.1 > 1.4 - 3.8x$

83) $2.2x + 1.9 < -2.3$

Answer

$(-\infty, -21/11)$

84) $1.6x + 1.2 < 1.6$

Contributors and Attributions

- David Arnold (Retired Professor (Mathematics) at College of the Redwoods)

2.7E: Exercises for Section 2.7 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **2.E: Solving Linear Equations and Inequalities (Exercises)** by David Arnold is licensed CC BY-NC-ND 3.0. Original source: https://math.libretexts.org/api/deki/files/80188/ElementaryAlgebra.pdf.

2.7E2: Exercises

Practice Makes Perfect

Graph Inequalities on the Number Line

In the following exercises, graph each inequality on the number line


Exercise 2.7E2.1


1. $x \leq 2$
2. x>−1
3. x<0


Exercise 2.7E2.2

1. x>1
2. x<−2
3. $x \geq -3$

Answer

1. 

2. 

3. 


Exercise 2.7E2.3


1. $x \geq -3$
2. x<4
3. $x \leq -2$


Exercise 2.7E2.4

1. $x \leq 0$
2. x>−4
3. $x \geq -1$

Answer

1. 

2. 

3. 

In the following exercises, graph each inequality on the number line and write in interval notation.


Exercise 2.7E2.5


1. x<−2
2. $x \geq -3.5$
3. $x \leq \frac{2}{3}$


Exercise 2.7E2.6

1. $x > 3$
2. $x \leq -0.5$
3. $x \geq \frac{1}{3}$

Answer

1. 

2. 

3. 


Exercise 2.7E2.7


1. $x \geq -4$
2. x<2.5
3. $x > -\frac{3}{2}$


Exercise 2.7E2.8

1. $x \leq 5$
2. $x \geq -1.5x$
3. x<−73

Answer

1. 

2. 

3. 

Solve Inequalities using the Subtraction and Addition Properties of Inequality

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 2.7E2.9

$n - 11 < 33$

Exercise 2.7E2.10

$m - 45 \leq 62$

Answer




Exercise 2.7E2.11

$u+25>21$

Exercise 2.7E2.12

$v+12>3$

Answer




Exercise 2.7E2.13

$a+\frac{3}{4}\geq\frac{7}{10}$

Exercise 2.7E2.14

$b+\frac{7}{8}\geq\frac{1}{6}$

Answer




Exercise 2.7E2.15

$f-\frac{13}{20}<-\frac{5}{12}$

Exercise 2.7E2.16

$g-\frac{11}{12}<-\frac{5}{18}$

Answer



Solve Inequalities using the Division and Multiplication Properties of Inequality

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 2.7E2.17

$8x > 72$

Exercise 2.7E2.18

$6y < 48$

Answer




Exercise 2.7E2.19

$7r \leq 56$

Exercise 2.7E2.20

$9s \geq 81$

Answer




Exercise 2.7E2.21

$-5u \geq 65$

Exercise 2.7E2.22

$-8v \leq 96$

Answer




Exercise 2.7E2.23

$-9c < 126$

Exercise 2.7E2.24

$-7d > 105$

Answer




Exercise 2.7E2.25

$20 > \frac{2}{5}h$

Exercise 2.7E2.26

$40 < \frac{5}{8}k$

Answer




Exercise 2.7E2.27

$\frac{7}{6}j \geq 42$

Exercise 2.7E2.28

$\frac{9}{4}g \leq 36$

Answer




Exercise 2.7E2.29

$\frac{a}{-3} \leq 9$

Exercise 2.7E2.30

$\frac{b}{-10} \geq 30$

Answer




Exercise 2.7E2.31

$-25 < \frac{p}{-5}$

Exercise 2.7E2.32

$-18 > \frac{q}{-6}$

Answer




Exercise 2.7E2.33

$9t \geq -27$

Exercise 2.7E2.34

$7s < -28$

Answer




Exercise 2.7E2.35

$\frac{2}{3}y > -36$

Exercise 2.7E2.36

$\frac{3}{5}x \leq -45$

Answer



Solve Inequalities That Require Simplification

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 2.7E2.37

$4v \geq 9v - 40$

Exercise 2.7E2.38

$5u \leq 8u - 21$

Answer




Exercise 2.7E2.39

$13q < 7q - 29$

Exercise 2.7E2.40

$9p > 14p - 18$

Answer




Exercise 2.7E2.41

$12x + 3(x + 7) > 10x - 24$

Exercise 2.7E2.42

$9y + 5(y + 3) < 4y - 35$

Answer




Exercise 2.7E2.43

$6h - 4(h - 1) \leq 7h - 11$

Exercise 2.7E2.44

$4k - (k - 2) \geq 7k - 26$

Answer




Exercise 2.7E2.45

$8m - 2(14 - m) \geq 7(m - 4) + 3m$

Exercise 2.7E2.46

$6n - 12(3 - n) \leq 9(n - 4) + 9n$

Answer




Exercise 2.7E2.47

$\frac{3}{4}b-\frac{1}{3}b<\frac{5}{12}b-\frac{1}{2}$

Exercise 2.7E2.48

$9u+5(2u-5)\geq 12(u-1)+7u$

Answer




Exercise 2.7E2.49

$\frac{2}{3}g-\frac{1}{2}(g-14)\leq \frac{1}{6}(g+42)$

Exercise 2.7E2.50

$\frac{5}{6}a-\frac{1}{4}a>\frac{7}{12}a+\frac{2}{3}$

Answer




Exercise 2.7E2.51

$\frac{4}{5}h-\frac{2}{3}(h-9)\geq \frac{1}{15}(2h+90)$

Exercise 2.7E2.52

$12v+3(4v-1)\leq 19(v-2)+5v$

Answer



Mixed practice

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 2.7E2.53

$15k\leq -40$

Exercise $2.7E2.54$

$35k \geq -77$

Answer




Exercise $2.7E2.55$

$23p - 2(6 - 5p) > 3(11p - 4)$

Exercise $2.7E2.56$

$18q - 4(10 - 3q) < 5(6q - 8)$

Answer




Exercise $2.7E2.57$

$-\frac{9}{4}x \geq -\frac{5}{12}$

Exercise $2.7E2.58$

$-\frac{21}{8}y \leq -\frac{15}{28}$

Answer




Exercise $2.7E2.59$

$c + 34 < -99$

Exercise $2.7E2.60$

$d + 29 > -61$

Answer




Exercise 2.7E2.61

$\frac{m}{18} \geq -4$

Exercise 2.7E2.62

$\frac{n}{13} \leq -6$

Answer



Translate to an Inequality and Solve

In the following exercises, translate and solve .Then write the solution in interval notation and graph on the number line.


Exercise 2.7E2.63

Fourteen times d is greater than 56.

Exercise 2.7E2.64

Ninety times c is less than 450.

Answer




Exercise 2.7E2.65

Eight times z is smaller than -40

Exercise 2.7E2.66

Ten times y is at most -110

Answer




Exercise 2.7E2.67

Three more than h is no less than 25

Exercise 2.7E2.68

Six more than k exceeds 25

Answer




Exercise 2.7E2.69

Ten less than w is at least 39.

Exercise 2.7E2.70

Twelve less than x is no less than 21

Answer




Exercise 2.7E2.71

Negative five times r is no more than 95.

Exercise 2.7E2.72

Negative two times s is lower than 56

Answer




Exercise 2.7E2.73

Nineteen less than b is at most -22

Exercise 2.7E2.74

Fifteen less than a is at least -7

Answer



Everyday Math

Exercise 2.7E2.75

Safety A child's height, *h*, must be at least 57 inches for the child to safely ride in the front seat of a car. Write this as an inequality.

Exercise 2.7E2.76

Fighter pilots The maximum height, *h*, of a fighter pilot is 77 inches. Write this as an inequality.

Answer

$h \leq 77$

Exercise 2.7E2.77

Elevators The total weight, *w*, of an elevator's passengers can be no more than 1,200 pounds. Write this as an inequality.

Exercise 2.7E2.78

Shopping The number of items, *n*, a shopper can have in the express check-out lane is at most 8. Write this as an inequality.

Answer

$n \leq 8$

Writing Exercises

Exercise 2.7E2.79

Give an example from your life using the phrase 'at least'.

Exercise 2.7E2.80

Give an example from your life using the phrase 'at most'.

Answer

Answers will vary.

Exercise 2.7E2.81

Explain why it is necessary to reverse the inequality when solving $-5x > 10$

Exercise 2.7E2.82

Explain why it is necessary to reverse the inequality when solving $\frac{n}{-3} < 12$

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
graph inequalities on the number line.			
solve inequalities using the Subtraction and Addition Properties of Inequality.			
solve inequalities using the Division and Multiplication Properties of Inequality.			
solve inequalities that require simplification.			
translate to an inequality and solve.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

More Practice: Solve Applications with Linear Inequalities

? Exercise $2.7E2.83$

1. Alan is loading a pallet with boxes that each weighs 45 pounds. The pallet can safely support no more than 900 pounds. How many boxes can he safely load onto the pallet?

2. The elevator in Yehire's apartment building has a sign that says the maximum weight is 2100 pounds. If the average weight of one person is 150 pounds, how many people can safely ride the elevator?

3. Andre is looking at apartments with three of his friends. They want the monthly rent to be no more than $2,360. If the roommates split the rent evenly among the four of them, what is the maximum rent each will pay?

4. Arleen got a $20 gift card for the coffee shop. Her favorite iced drink costs $3.79. What is the maximum number of drinks she can buy with the gift card?

5. Teegan likes to play golf. He has budgeted $60 next month for the driving range. It costs him $10.55 for a bucket of balls each time he goes. What is the maximum number of times he can go to the driving range next month?

6. Ryan charges his neighbors $17.50 to wash their car. How many cars must he wash next summer if his goal is to earn at least $1,500?

7. Keshad gets paid $2,400 per month plus 6% of his sales. His brother earns $3,300 per month. For what amount of total sales will Keshad's monthly pay be higher than his brother's monthly pay?

8. Kimuyen needs to earn $4,150 per month in order to pay all her expenses. Her job pays her $3,475 per month plus 4% of her total sales. What is the minimum Kimuyen's total sales must be in order for her to pay all her expenses?

9. Andre has been offered an entry-level job. The company offered him $48,000 per year plus 3.5% of his total sales. Andre knows that the average pay for this job is $62,000. What would Andre's total sales need to be for his pay to be at least as high as the average pay for this job?

10. Nataly is considering two job offers. The first job would pay her $83,000 per year. The second would pay her $66,500 plus 15% of her total sales. What would her total sales need to be for her salary on the second offer be higher than the first?

11. Jake's water bill is $24.80 per month plus $2.20 per ccf (hundred cubic feet) of water. What is the maximum number of ccf Jake can use if he wants his bill to be no more than $60?

12. Kiyoshi's phone plan costs $17.50 per month plus $0.15 per text message. What is the maximum number of text messages Kiyoshi can use so the phone bill is no more than $56.60?

13. Marlon's TV plan costs $49.99 per month plus $5.49 per first-run movie. How many first-run movies can he watch if he wants to keep his monthly bill to be a maximum of $100?

14. Kellen wants to rent a banquet room in a restaurant for her cousin's baby shower. The restaurant charges $350 for the banquet room plus $32.50 per person for lunch. How many people can Kellen have at the shower if she wants the maximum cost to be $1,500?

15. Moshde runs a hairstyling business from her house. She charges $45 for a haircut and style. Her monthly expenses are $960. She wants to be able to put at least $1,200 per month into her savings account order to open her own salon. How many

"cut & styles" must she do to save at least $1,200 per month?

16. Noe installs and configures software on home computers. He charges $125 per job. His monthly expenses are $1,600. How many jobs must he work in order to make a profit of at least $2,400?

17. Katherine is a personal chef. She charges $115 per four-person meal. Her monthly expenses are $3,150. How many four-person meals must she sell in order to make a profit of at least $1,900?

18. Melissa makes necklaces and sells them online. She charges $88 per necklace. Her monthly expenses are $3,745. How many necklaces must she sell if she wants to make a profit of at least $1,650?

19. Five student government officers want to go to the state convention. It will cost them $110 for registration, $375 for transportation and food, and $42 per person for the hotel. There is $450 budgeted for the convention in the student government savings account. They can earn the rest of the money they need by having a car wash. If they charge $5 per car, how many cars must they wash in order to have enough money to pay for the trip?

20. Cesar is planning a four-day trip to visit his friend at a college in another state. It will cost him $198 for airfare, $56 for local transportation, and $45 per day for food. He has $189 in savings and can earn $35 for each lawn he mows. How many lawns must he mow to have enough money to pay for the trip?

21. Alonzo works as a car detailer. He charges $175 per car. He is planning to move out of his parents' house and rent his first apartment. He will need to pay $120 for application fees, $950 for security deposit, and first and last months' rent at $1,140 per month. He has $1,810 in savings. How many cars must he detail to have enough money to rent the apartment?

22. Eun-Kyung works as a tutor and earns $60 per hour. She has $792 in savings. She is planning an anniversary party for her parents. She would like to invite 40 guests. The party will cost her $1,520 for food and drinks and $150 for the photographer. She will also have a favor for each of the guests, and each favor will cost $7.50. How many hours must she tutor to have enough money for the party?

23. In 2014, a high school stage collapsed in Fullerton, California, when 250 students got on stage for the finale of a musical production. Two dozen students were injured. The stage could support a maximum of 12,750 pounds. If the average weight of a student is assumed to be 140 pounds, what is the maximum number of students who could safely be on the stage?

24. In 2004, a water taxi sank in Baltimore harbor and five people drowned. The water taxi had a maximum capacity of 3,500 pounds (25 people with average weight 140 pounds). The average weight of the 25 people on the water taxi when it sank was 168 pounds per person. What should the maximum number of people of this weight have been?

25. Adele and Walter found the perfect venue for their wedding reception. The cost is $9850 for up to 100 guests, plus $38 for each additional guest. How many guests can attend if Adele and Walter want the total cost to be no more than $12,500?

26. Penny is planning a baby shower for her daughter-in-law. The restaurant charges $950 for up to 25 guests, plus $31.95 for each additional guest. How many guests can attend if Penny wants the total cost to be no more than $1,500?

Answer

2. A maximum of 14 people can safely ride in the elevator. 4. five drinks 6. 86 cars 8. $16,875 10. $110,000

12. 260 messages 14. 35 people 16. 32 jobs 18. 62 necklaces 20. seven lawns 22. 20 hours 24. 20 people 26. 42 guests.

This page titled 2.7E2: Exercises is shared under a not declared license and was authored, remixed, and/or curated by David Arnold.

- **2.7E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **2.6E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/intermediate-algebra-2e.

2.8: Solve Compound Inequalities

In this section, we will explore compound inequalities. Like a compound sentence is the combination of two (or more) sentences into a larger sentence using conjunctions like "and" and "or", compound inequalities are statements formed by linking two (or more) inequalities together, either using the word "and" or "or", like

$$x \leq -1 \text{ or } x > 5,$$

or by creating a single statement (sometimes called a *chain inequality*) that incorporates multiple inequality symbols, e.g.,

$$-4 \leq x < 7.$$

In order to better understand how these compound inequalities work, we first need to define the intersection and union of two sets.

Intersection and Union

The intersection of two sets A and B is defined as follows.

Definition

The **intersection** of the sets A and B is the set of all objects that are in A and in B.

In symbols, we write

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\} \tag{2.8.1}$$

In order to understand this definition, it's absolutely crucial that we understand the meaning of the word "and." The word "and" is a conjunction, used between statements P and Q, as in "It is raining today **and** my best friend is the Lone Ranger." In order to determine the truth or falsehood of this statement, you must first examine the truth or falsehood of the statements P and Q on each side of the word "and."

The only way that the speaker is telling the truth is if both statements P and Q are true. In other words, the statement "It is raining today and my best friend is the Lone Ranger" is true if and only if the statement "It is raining today" is true and the statement "my best friend is the Lone Ranger" is also true. Logicians like to make up a construct called a truth table, like the one shown in Table 2.8.2.

Points in Table 2.8.2 to consider:

Table 2.8.2 Truth table for the conjunction "and."

P	Q	P **and** Q
T	T	T
T	F	F
F	T	F
F	F	F

- In the first row (after the header row) of Table 2.8.2, if statements P and Q are both true (indicated with a T), then the statement "P and Q" is also true.
- In the remaining rows of Table 2.8.2, one or the other of statements P or Q are false (indicated with an F), so the statement "P and Q" is also false.

Therefore, the statement "P and Q" is true if and only if P is true and Q is true.

Example 2.8.1

If $A = \{1, 3, 5, 7, 9\}$ and $B = \{2, 5, 7, 8, 11\}$, find the intersection of A and B.

Solution

As a reminder, the *intersection* of sets A and B is

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\} \tag{2.8.2}$$

Thus, we are looking for the objects that are in A and in B. The only objects that are in A and in B (remember, both statements "in A" and "in B" must be true) are 5 and 7, so we write:

$$A \cap B = \{5, 7\} \tag{2.8.3}$$

Mathematicians and logicians both use a visual aid called a Venn Diagram to represent sets. John Venn was an English mathematician who devised this visualization of logical relationships. Consider the ellipse A in Figure 2.8.5. Everything inside the boundary of this ellipse constitutes the set $A = \{1, 3, 5, 7, 9\}$. That's why you see these numbers inside the boundary of this ellipse.

Consider the ellipse B in Figure 2.8.5. Everything inside the boundary of this ellipse constitutes the set $B = \{2, 5, 7, 8, 11\}$. That's why you see these numbers inside the boundary of this ellipse.




Figure 2.8.5 Venn Diagram

Now, note that only two numbers, 5 and 7, are contained within the boundaries of both A and B. These are the numbers that are in the intersection of the sets A and B.

The shaded region in Figure 2.8.6 is the area that belongs to both of the sets A and B. Note how this shaded region is aptly named "the intersection of the sets A and B." This is the region that is in common to the sets A and B, the region where the sets A and B overlap or "intersect."




Figure 2.8.6 The shaded region is the intersection of the sets A and B. That is, the shaded region is $A \cap B$.

This leads to the following important piece of advice.

Note

When asked to find the intersection of two sets A and B, look to see where the sets intersect or overlap. That is, look to see the elements that are in both sets A and B.

Let's move on to the definition of the union of two sets A and B.

Definition

The **union** of the sets A and B is the set of all objects that are in A or in B.

In symbols, we write

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\} \tag{2.8.4}$$

In order to understand this definition, it's critical that we understand the meaning of the word "or." The word "or" is a disjunction, used between statements P and Q, as in "It is raining today or my best friend is the Lone Ranger." In order to determine the truth

or falsehood of this statement, you must first examine the truth or falsehood of the statements P and Q on each side of the word "or."

The speaker is telling the truth if either statement P is true or statement Q is true. In other words, the statement "It is raining today or my best friend is the Lone Ranger" is true if and only if the statement "It is raining today" is true or the statement "my best friend is the Lone Ranger" is true. Logicians like to make up a construct called a truth table, like the one shown in Table 2.8.3.

Table 2.8.3 Truth table for the conjunction "or."

P	Q	P **or** Q
T	T	T
T	F	T
F	T	T
F	F	F

Points in Table 2.8.3 to consider:

- In the last row of Table 2.8.3, both statements P and Q are false (indicated with an F), so the statement P or Q is also false.
- In the first three rows (after the header row) of Table 2.8.3, either statement P is true or statement Q is true (indicated with a T), so the statement P or Q is also true.

Therefore, the statement "P or Q" is true if and only if either statement, P or Q, is true.

Example 2.8.2

If $A = \{1, 3, 5, 7, 9\}$ and $B = \{2, 5, 7, 8, 11\}$, find the union of A and B.

Solution

As a reminder, the union of A and B is

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\} \tag{2.8.5}$$

Thus, an object is in the union of A and B if and only if it is in either set. The numbers that are in either set are the numbers

$$A \cup B = \{1, 2, 3, 5, 7, 8, 9, 11\} \tag{2.8.6}$$

If we look again at the Venn Diagram in Figure 2.8.5, we see that this union $A \cup B = \{1, 2, 3, 5, 7, 8, 9, 11\}$ lists every number that is in either set in Figure 2.8.5.

Thus, the shaded region in Figure 2.8.7 is the union of sets A and B. Note how this region is well-named, as that's what you're actually doing, taking the "union" of the two sets A and B. That is, the union contains all elements that belong to either A or B. Less formally, the union is a way of combining everything that occurs in either set.




Figure 2.8.7. The shaded region is the union of sets A and B. That is, the shaded region is $A \cup B$.

This leads to the following important piece of advice.

Note

When asked to find the union of two sets A and B, in your answer, include everything from both sets.

Simple Compound Inequalities

Let's apply what we've learned to find the unions and/or intersections of intervals of real numbers. The easiest approach is through a series of examples. Let's begin.

Example 2.8.3

On the real line, sketch the set of real numbers in the set $\{x \mid x < 3 \text{ or } x < 5\}$. Use set-builder notation and interval notation to describe your final answer.

Solution

First, let's sketch two sets, $\{x \mid x < 3\}$ and $\{x \mid x < 5\}$, on separate real lines, one above the other as shown in Figure 2.8.8.




Figure 2.8.8 Sketch each set separately.

Now, to sketch the solution, note the word "or" in the set $\{x \mid x < 3 \text{ or } x < 5\}$. Thus, we need to take the union of the two shaded real lines in Figure 2.8.8. That is, we need to shade everything that is shaded on either of the two number lines. Of course, this would be everything less than 5, as shown in Figure 2.8.9.




Figure 2.8.9 The final solution is the union of the two shaded sets in Figure 2.8.8.

Thus, the final simplified solution is

Set-Builder Notation: $\{x \mid x < 5\}$

Interval notation: $(-\infty, 5)$

Let's look at another example.

Example 2.8.4

On the real line, sketch the set of real numbers in the set $\{x \mid x < 3 \text{ and } x < 5\}$. Use interval notation to describe your final answer.

Solution

In Example 2.8.3, you were asked to shade the set $\{x \mid x < 3 \text{ or } x < 5\}$ on the real line. In this example, we're asked to sketch the set $\{x \mid x < 3 \text{ and } x < 5\}$. Note that the set-builder notations are identical except for one change, the "or" of Example 2.8.3 has been replaced with the word "and."

Our solution will be the numbers that are solutions to *both* inequalities known as the intersection of the two inequalities. Consider the intersection of two streets—the part where the streets overlap—belongs to both streets.



Figure 2.8.10: The intersection of two streets is where they overlap.

To find the solution of an "and" compound inequality, we look at the graphs of each inequality and then find the numbers that belong to both graphs—where the graphs overlap.

Again, sketch two sets, $\{x \mid x < 3\}$ and $\{x \mid x < 5\}$, on separate real lines, one above the other as shown in Figure 2.8.11.




Figure 2.8.11. Sketch each set separately.

Now, to sketch the solution, note the word "and" in the set $\{x \mid x < 3 \text{ and } x < 5\}$. Thus, we need to take the intersection of the two shaded real lines in Figure 2.8.11. That is, we need to shade everything that is common to the two number lines. Here this would be everything less than 3, as shown in Figure 2.8.12.



Figure 2.8.12. The final solution is the intersection of the two shaded sets in Figure 2.8.11.

Thus, the final simplified solution is

Set-Builder Notation: $\{x \mid x < 3\}$

Interval notation: $(-\infty, 3)$.

Note

If you answer "or" when the answer requires "and," or vice-versa, you have not made a minor mistake. Indeed, this is a huge error, as demonstrated in Example 2.8.3 and Example 2.8.4.

Before attempting another example, we pause to define a bit of notation that will be extremely important in our upcoming work.

Definition

The notation

$$a < x < b \tag{2.8.7}$$

is interpreted to mean

$$x > a \text{ and } x < b \tag{2.8.8}$$

Alternatively, we could have said that $a < x < b$ is identical to saying "$a < x$ and $x < b$," but saying "$a < x$" is the same as saying "$x > a$." We prefer to say "$x > a$ and $x < b$," and will use this order throughout our work, but the form "$a < x$ and $x < b$" is equally valid.

The really key point to make here is the fact that the statement $a < x < b$ is an "and" statement. If it is used properly, it's a good way to describe the numbers that lie between a and b.

Let's look at an example.

Example 2.8.5

On the real line, sketch the set of real numbers in the set $\{x \mid x > 3 \text{ and } x < 5\}$. Give both the simplified set-builder notation and interval notation to describe your answer.

Solution

The first step is to sketch the sets $\{x \mid x > 3\}$ and $\{x \mid x < 5\}$ on individual real lines, stacked one above the other, as shown in Figure 2.8.13.



Figure 2.8.13. Sketch each set separately.

Now, to sketch the solution, note the word "and" in the set $\{x \mid x > 3 \text{ and } x < 5\}$. Thus, we need to take the intersection of the two graphs in Figure 2.8.13. That is, we need to shade the numbers on the real line that are in common to the two graphs shown in Figure 2.8.13. The numbers 3 and 5 are not shaded in both sets in Figure 2.8.13, so they will not be shaded in our final solution. However, all real numbers between 3 and 5 are shaded in both sets in Figure 2.8.13, so these numbers will be shaded in the final solution shown in Figure 2.8.14.




Figure 2.8.14. The final solution is the intersection of the two shaded sets in Figure 2.8.13.

Since the numbers that make the original compound inequality true are just those that are between 3 and 5 on the number line, we can simplify the condition in set-builder notation to





Set-Builder Notation: $\{x \mid 3 < x < 5\}$

And the interval notation for the shaded solution in Figure 2.8.14 is

Interval Notation: $(3, 5)$

Similarly, here are the set-builder and interval notations, as well as shading of the sets on the real line, for several situations, including the one just discussed.

Table 2.8.4. Number line graphs, set-builder notation, and interval notation.

Number line graph	Set-builder notation	Interval notation
	$\{x \mid 3 < x < 5\}$	$(3, 5)$
	$\{x \mid 3 \leq x \leq 5\}$	$[3, 5]$
	$\{x \mid 3 \leq x < 5\}$	$[3, 5)$
	$\{x \mid 3 < x \leq 5\}$	$(3, 5]$

There are several points of emphasis regarding the intervals in Table 2.8.4.

1. When we want to emphasize that we are not including a point on the real line, we use an "empty circle." Conversely, a "filled circle" means that we are including the point on the real line. Thus, the interval in the first row of Table 2.8.4 does not include the endpoints at 3 and 5, but the interval in the second row of Table 2.8.4 does include the endpoints at 3 and 5.
2. The use of a parenthesis in interval notation means that we are not including that endpoint in the interval. Thus, the parentheses used in $(3, 5)$ in the first row of Table 2.8.4 means that we are not including the numbers 3 and 5 in that interval.
3. The use of a bracket in interval notation means that we are including the bracketed number in the interval. Thus, the brackets used in $[3, 5]$, as seen in the second row of Table 2.8.4, means that we are including the numbers 3 and 5 in the interval.
4. Finally, note that some of our intervals are "open" on one end but "closed" (filled) on the other end, such as those in rows 3 and 4 of Table 2.8.4.

Definition

Some terminology:

- The interval $(3, 5)$ is open at each end. Therefore, we call the interval $(3, 5)$ an open interval.
- The interval $[3, 5]$ is closed (filled) at each end. Therefore, we call the interval $[3, 5]$ a closed interval.
- The intervals $(3, 5]$ and $[3, 5)$ are neither open nor closed.

Let's look at another example.

Example 2.8.6

On the real line, sketch the set of all real numbers in the set $\{x \mid x > 3 \text{ or } x < 5\}$. Use set-builder notation and interval notation to describe your answer.

Solution

Note that the only difference between this current example and the set shaded in Example 2.8.4 is the fact that we've replaced the word "and" in $\{x \mid x > 3 \text{ and } x < 5\}$ with the word "or" in $\{x \mid x > 3 \text{ or } x < 5\}$. But, as we've seen before, this can make a world of difference.

Thus, the first step is to sketch the sets $\{x \mid x > 3\}$ and $\{x \mid x < 5\}$ on individual real lines, stacked one above the other, as shown in Figure 2.8.15.



Figure 2.8.15. Sketch each set separately

Now, to sketch the solution, note the word "or" in the set $\{x \mid x > 3 \text{ or } x < 5\}$. Thus, we need to take the union of the two graphs in Figure 2.8.15. That is, we need to shade the numbers on the real line that are shaded on either of the two graphs shown in Figure 2.8.15. However, this means that we will have to shade every number on the line, as shown in Figure 2.8.16. You'll note no labels for 3 and 5 on the real line in Figure 2.8.16, as there are no endpoints in this solution. The endpoints, if you will, are at negative and positive infinity.



Figure 2.8.16. The final solution is the union of the two shaded sets in Figure 2.8.15.

Thus, in a most natural way, the interval notation for the shaded solution in Figure 2.8.16 is

Interval Notation: $(-\infty, \infty)$.

The set-builder notation for this solution needs to represent the whole real line or all real numbers, so we write it in the following way:

Set-Builder Notation: $\{x \mid x \in \mathbb{R}\}$.

Let's look at another example.

Example 2.8.7

On the real line, sketch the set of all real numbers in the set $\{x \mid x < -1 \text{ or } x > 3\}$. Use interval notation to describe your answer.

Solution

The first step is to sketch the sets $\{x \mid x < -1\}$ and $\{x \mid x > 3\}$ on separate real lines, stacked one above the other, as shown in Figure 2.8.17.




Figure 2.8.17. Sketch each set separately.

To sketch the solution, note the word "or" in the set $\{x \mid x < -1 \text{ or } x > 3\}$. Thus, we need to take the union of the two shaded real lines in Figure 2.8.17. That is, we need to shade the numbers on the real line that are shaded on either real line in Figure 2.8.17. Thus, every number smaller than -1 is shaded, as well as every number greater than 3. The result is shown in Figure 2.8.18.




Figure 2.8.18. The final solution is the union of the shaded real lines in Figure 2.8.17.

Since the graph of this solution is not a single interval on the numberline, but is instead made up of two disjoint (non-overlapping) intervals, in this one situation, we will need to leave the word "or" in the condition in the set-builder notation for this solution.

Set-Builder Notation: $\{x \mid x < -1 \text{ or } x > 3\}$

Next let's figure out how to write this solution using interval notation.

Here is an important tip.

Note

If you wish to use interval notation correctly, follow one simple rule: Always sweep your eyes from left to right describing what you see shaded on the real line.

If we follow this advice, as we sweep our eyes from left to right across the real line shaded in Figure 2.8.18, we see that numbers are shaded from negative infinity to -1, and from 3 to positive infinity. Thus, in a most natural way, the interval notation for the shaded solution set in Figure 2.8.18 is

Interval Notation: $(-\infty, -1) \cup (3, \infty)$

There are several important points to make here:





Note how we used the union symbol $\cup$ to join the two intervals in $(-\infty, -1) \cup (3, \infty)$ in a natural manner.

The union symbol is used between sets of numbers, while the word “or” is used between statements about numbers. Thus it is incorrect to exchange the roles of the union symbol and the word “or.” Thus, writing $\{x \mid x < -1 \cup x > 3\}$ is incorrect, as it would also be to write $(-\infty, -1)$ or $(3, \infty)$.

Note that we will never use an intersection symbol ($\cap$) in our interval notation. Intersections of intervals can always be simplified to a single interval. Only unions (combined using the word "or") can form solutions made up of disjoint (non-connected) intervals like the one shown in Figure 2.8.18.

Let's reinforce our earlier discussion about the difference between “filled” and “open” circles, brackets, and parentheses using the examples shown below in Table 2.8.5. Here we include several comparisons of interval and set-builder notation, including the current solution to Example 2.8.7.

Table 2.8.5. Number line graphs, set-builder notation, and interval notation.

Number line graph	Set-builder notation	Interval notation
	$\{x \mid x < -1 \text{ or } x > 3\}$	$(-\infty, -1) \cup (3, \infty)$
	$\{x \mid x \leq -1 \text{ or } x \geq 3\}$	$(-\infty, -1] \cup [3, \infty)$
	$\{x \mid x \leq -1 \text{ or } x > 3\}$	$(-\infty, -1] \cup (3, \infty)$
	$\{x \mid x < -1 \text{ or } x \geq 3\}$	$(-\infty, -1) \cup [3, \infty)$

Again, we reinforce the following points.

- Note how sweeping your eyes from left to right, describing what is shaded on the real line, insures that you write the interval notation in the correct order.
- A bracket is equivalent to a filled dot and includes the endpoint, while a parenthesis is equivalent to an open dot and does not include the endpoint.

Let’s do one last example that should forever cement the notion that there is a huge difference between the words “and” and “or.”

Example 2.8.8

On the real line, sketch the set of all real numbers in the set $\{x \mid x < -1 \text{ and } x > 3\}$. Describe your solution.

Solution

First and foremost, note that the only difference between this example and Example 2.8.6 is the fact that we changed the "or" in $\{x \mid x < -1 \text{ or } x > 3\}$ to an "and" in $\{x \mid x < -1 \text{ and } x > 3\}$. The preliminary sketches are identical to those in Figure 2.8.17.




Figure 2.8.19. Sketch $\{x \mid x < -1\}$ and $\{x \mid x > 3\}$ on separate real lines.

Now, note the word "and" in $\{x \mid x < -1 \text{ and } x > 3\}$. Thus, we need to take the intersection of the shaded real lines in Figure 2.8.19. That is, we need to shade on a single real line all of the numbers that are shaded on both real lines in Figure 2.8.19. However, there are no points shaded in common on the real lines in Figure 2.8.19, so the solution set is empty, as shown in Figure 2.8.20.

Figure 2.8.20. The solution is empty so we leave the real line blank.

This compound inequality has **No Solution**, so we would write its solution set as: $\{\}$.

There would be no way to write interval notation or set-builder notation in this case.

Pretty impressive! The last two examples clearly demonstrate that if you interchange the roles of "and" and "or," you have not made a minor mistake. Indeed, you've changed the whole meaning of the problem. So, be careful with your "ands" and "ors."

Solving Compound Inequalities

We now turn our attention to the business of solving compound inequalities.

In this case, the best way of learning is by doing. Let's start with an example.

Example 2.8.9 Solving an "OR" Compound Inequality

Solve the following compound inequality for x.


$$3 - 2x < -1 \quad \text{or} \quad 3 - 2x > 1$$

Solution

First, solve each of the inequalities independently. With the first inequality, add -3 to both sides of the inequality, then divide by -2, reversing the inequality sign.


$$\begin{aligned} 3 - 2x &< -1 \\ -2x &< -4 \\ x &> 2 \end{aligned}$$

Shade the solution on a number line.



The exact same sequence of operations can be used to solve the second inequality

$$\begin{aligned} 3 - 2x &> 1 \\ -2x &> -2 \\ x &< 1 \end{aligned}$$



Although you solve each side of the inequality independently, you will want to arrange your work as follows, stacking the number line solution for the first inequality above that of the second inequality.

$$\begin{array}{rcl} 3-2x & < & -1 \\ -2x & < & -4 \\ x & > & 2 \end{array} \quad \text{or} \quad \begin{array}{rcl} 3-2x & > & 1 \\ -2x & > & -2 \\ x & < & 1 \end{array}$$




Figure 2.8.21. The solution of the compound inequality $3-2x<-1$ or $3-2x>1$

The answer, described in both interval and set-builder notation, is

Interval Notation: $(-\infty,1)\cup(2,\infty)$

Set-Builder Notation: $\{x \mid <1 \text{ or } x>2\}$.

Let's look at another example.

Example 2.8.10 Solving an "AND" Compound Inequality

Solve the following compound inequality for x.

$$3-2x>-1 \quad \text{and} \quad 3-2x<1 \tag{2.8.9}$$

Solution

Solve each inequality independently, arranging your work as follows.

$$\begin{array}{rcl} 3-2x & > & -1 \\ -2x & > & -4 \\ x & < & 2 \end{array} \quad \text{and} \quad \begin{array}{rcl} 3-2x & < & 1 \\ -2x & < & -2 \\ x & > & 1 \end{array} \tag{2.8.10}$$

Shade the solution of each inequality on separate real lines, one above the other




Figure 2.8.22. The solution of the compound inequality $-1<3-2x<1$.

The answer, described in both interval and set-builder notation, is

Interval Notation: $(1,2)$

Set-Builder Notation: $\{x \mid 1<x<2\}$

Note that we used the compact form of the compound inequality in our answer, as this condition is simpler than using an "and" compound inequality.

Alternative approach. You might have noted that in solving the second inequality in 2.8.10, you found yourself repeating the identical operations used to solve the first inequality. That is, you subtracted 3 from both sides of the inequality, then divided both sides of the inequality by -2, reversing the inequality sign.

This repetition is annoying and suggests a possible shortcut in this particular situation. Because the statement $x > a$ and $x < b$ is identical to the statement $a < x < b$, solving the two part "AND" inequality (as in 2.8.9) could be solved more efficiently by first rewriting it in its equivalent "chain" inequality form

$$-1 < 3 - 2x < 1 \tag{2.8.11}$$

Now, here are the rules for working with this form.

Property 2.8.4

When working with a compound inequality having the form

$$a < x < b \tag{2.8.12}$$

you may add (or subtract) the same amount to (from) all three parts of the inequality, as in

$$a + c < x + c < b + c \tag{2.8.13}$$

or

$$a - c < x - c < b - c \tag{2.8.14}$$

You may also multiply all three parts by the same **positive** number $c > 0$, as in

$$ca < cx < cb \tag{2.8.15}$$

However, if you multiply all three parts by the same **negative** number $c < 0$, then don't forget to reverse the inequality signs, as in

$$ca > cx > cb \tag{2.8.16}$$

The rules for division are identical to the multiplication rules. If $c > 0$ (positive), then

$$\frac{a}{c} < \frac{x}{c} < \frac{b}{c} \tag{2.8.17}$$

If $c < 0$ (negative), then reverse the inequality signs when you divide.

$$\frac{a}{c} > \frac{x}{c} > \frac{b}{c} \tag{2.8.18}$$

Each of the tools in Property 2.8.4 always produce equivalent inequalities.

So, let's return to the compound inequality 2.8.9. The following example solves it using its equivalent "Chain" inequality form 2.8.11.

Example 2.8.10*b* Solving a "chain" Compound Inequality

Solve the following compound inequality for x.

$$-1 < 3 - 2x < 1 \tag{2.8.19}$$

Solution

$$\begin{gathered} -1 < 3 - 2x < 1 \\ -1 - 3 < 3 - 2x - 3 < 1 - 3 \\ -4 < -2x < -2 \end{gathered}$$

Next, divide all three members by -2, reversing the inequality signs as you do so.

$$\begin{gathered} -4 < -2x < -2 \\ \frac{-4}{-2} > \frac{-2x}{-2} > \frac{-2}{-2} \\ 2 > x > 1 \end{gathered}$$

It is conventional to change the order of this last inequality. By reading the inequality from right to left, we get

$$1 < x < 2$$

which describes the real numbers that are greater than 1 and less than 2. The solution is drawn on the following real line.



Figure 2.8.23. The solution of the compound inequality $-1 < 3 - 2x < 1$.

Note that this is identical to the solution set on the real line in Figure 2.8.22 Note also that this second alternative method is more efficient, particularly if you do a bit of work in your head. Consider the following sequence where we subtract three from all three members, then divide all three members by -2, reversing the inequality signs, then finally read the inequality in the opposite direction.

$$\begin{aligned} -1 &< 3 - 2x < 1 \\ -4 &< -2x < -2 \\ 2 &> x > 1 \\ 1 &< x < 2 \end{aligned}$$

Let's look at another example.

Example 2.8.11

Solve the following compound inequality for x.

$$-1 < x - \frac{x+1}{2} \leq 2$$

Solution

First, let's multiply all three members by 2, in order to clear the fractions.

$$2(-1) < 2\left(x - \frac{x+1}{2}\right) \leq 2(2)$$

$$-2 < 2(x) - 2\left(\frac{x+1}{2}\right) \leq 4$$

Cancel. Note the use of parentheses, which is crucial when a minus sign is involved.

$$-2 < 2x - \cancel{2}\left(\frac{x+1}{\cancel{2}}\right) \leq 4$$

$$-2 < 2x - (x+1) \leq 4$$

Distribute the minus sign and simplify.

$$\begin{aligned} -2 < \quad & 2x - x - 1 \quad && \leq 4 \\ -2 < \quad & x - 1 && \leq 4 \end{aligned}$$

Add 1 to all three members and simplify.

$$-2 + 1 < x - 1 + 1 \leq 4 + 1$$

$$-1 < x \leq 5$$

This solution describes the real numbers that are greater than -1 and less than 5, including 5. That is, the real numbers that fall between -1 and 5, including 5, shaded on the real line in Figure 2.8.24



Figure 2.8.24. The solution set of $-1 < x - (x+1)/2 \leq 2$.

The answer, described in both interval and set-builder notation, is

Interval Notation: $(-1, 5]$

Set-Builder Notation: $\{x \mid -1 < x \leq 5\}$

Let's look at another example.

Example 2.8.12 Solve a "chain" inequality with the variable in several parts of the inequality

Solve the following compound inequality for x.

$$x \leq 2x - 3 \leq 5$$

Solution

Suppose that we try to isolate x as we did in Example 2.8.11. Perhaps we would try adding $-x$ to all three members.

$$\begin{aligned} x \leq & \quad 2x - 3 & \leq 5 \\ x - x \leq & \quad 2x - 3 - x & \leq 5 - x \\ 0 \leq & \quad x - 3 & \leq 5 - x \end{aligned}$$

Well, that didn't help much, just transferring the problem with x to the other end of the inequality. Similar attempts will not help in isolating x. So, what do we do?

The solution is we split the inequality (with the word "and," of course).

$$x \leq 2x - 3 \quad \text{and} \quad 2x - 3 \leq 5$$

We can solve the first inequality by subtracting $2x$ from both sides of the inequality, then multiplying both sides by -1, reversing the inequality in the process

$$\begin{aligned} x &\leq 2x - 3 \\ -x &\leq -3 \\ x &\geq 3 \end{aligned}$$

To solve the second inequality, add 3 to both sides, then divide both sides by 2:

$$\begin{aligned} 2x - 3 &\leq 5 \\ 2x &\leq 8 \\ x &\leq 4 \end{aligned}$$

Of course, you'll probably want to arrange your work as follows

$$\begin{array}{rlcrl} x & \leq 2x - 3 & \text{and} & 2x - 3 & \leq 5 \\ -x & \leq -3 & & 2x & \leq 8 \\ x & \geq 3 & & x & \leq 4 \end{array}$$

Thus, we need to shade on a number line all real numbers that are greater than or equal to 3 and less than or equal to 4, as shown in Figure 2.8.25.



Figure 2.8.25 When shading the solution of $x \leq 2x - 3 \leq 5$, we "fill-in" the endpoints.

The solution, described in both interval and set-builder notation, is

Interval Notation: $[3, 4]$

Set-Builder Notation: $\{x \mid 3 \leq x \leq 4\}$

Contributors

Adapted and customized by Paul Seeburger (Monroe Community College) based primarily on material by

David Arnold (Retired Professor (Mathematics) at College of the Redwoods)

Figure 2.8.10 and some of the surrounding text comes from Section 2.7 of the OpenStax Intermediate Algebra text by Lynn Marecek.


2.8: Solve Compound Inequalities is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **1.3: Logic** by David Arnold is licensed CC BY-NC-SA 2.5. Original source: https://web.archive.org/web/20200814023923/http://msenux2.redwoods.edu/IntAlgText/.
- **2.7: Solve Compound Inequalities** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/intermediate-algebra-2e.
- **1.4: Compound Inequalities** by David Arnold is licensed CC BY-NC-SA 2.5. Original source: https://web.archive.org/web/20200814023923/http://msenux2.redwoods.edu/IntAlgText/.

2.8E: Exercises for Section 2.8


In Exercises 1 - 8, use both interval and set-builder notation to describe the intersection of the two intervals shown on the graph. Also, sketch the graph of the intersection on the real number line.

Exercise 1



Answer


The intersection is the set of points that are in both intervals (shaded on both graphs). Graph of the intersection:




Set-Builder Notation: $\{x \mid x \geq 1\}$

Interval Notation: $[1, \infty)$

Exercise 2



Exercise 3



Answer


There are no points that are in both intervals (shaded in both), so there is no intersection. Graph of the intersection:

no intersection

Set-Builder Notation: $\{\ \}$

Interval Notation: none

Exercise 4



Exercise 5



Answer

The intersection is the set of points that are in both intervals (shaded in both). Graph of the intersection:




Set-Builder Notation: $\{x \mid -6 \leq x \leq 2\}$

Interval Notation: $[-6, 2]$

Exercise 6



Exercise 7



Answer


The intersection is the set of points that are in both intervals (shaded in both). Graph of the intersection:



Set-Builder Notation: $\{x \mid x \geq 9\}$


Interval Notation: $[9, \infty)$

Exercise 8



In Exercises 9 - 16, use both interval notation and set-builder notation to describe the union of the two intervals shown on the graph. Also, sketch the graph of the union on the real number line.

Exercise 9



Answer

The union is the set of all points that are in one interval or the other (shaded in either graph). Graph of the union:




Set-Builder Notation: $\{x \mid x \leq -8\}$

Interval Notation: $(-\infty, -8]$

Exercise 10



Exercise 11



Answer


The union is the set of all points that are in one interval or the other (shaded in either graph). Graph of the union:




Set-Builder Notation: $\{x \mid x \leq 9 \text{ or } x > 15\}$

Interval Notation: $(-\infty, 9] \cup (15, \infty)$

Exercise 12



Exercise 13



Answer


The union is the set of all points that are in one interval or the other (shaded in either). Graph of the union:




Set-Builder Notation: $\{x \mid x < 3\}$

Interval Notation: $(-\infty, 3)$

Exercise 14



Exercise 15



Answer


The union is the set of all points that are in one interval or the other (shaded in either). Graph of the union:



Set-Builder Notation: $\{x \mid x \geq 9\}$

Interval Notation: $[9, \infty)$

Exercise 16




In Exercises 17 - 32, use interval notation to describe the given set. Also, sketch the graph of the set on the real number line.

Exercise 17

$\{x \mid x \geq -6 \text{ and } x > -5\}$

Answer

This set is the same as $\{x \mid x > -5\}$, which is $(-5, \infty)$ in interval notation. Graph of the set:



Exercise 18

$\{x \mid x \leq 6 \text{ and } x \geq 4\}$

Exercise 19

$\{x \mid x \geq -1 \text{ or } x < 3\}$

Answer

Every real number is in one or the other of the two intervals. Therefore, the set is the set of all real numbers $(-\infty, \infty)$. Graph of the set:

Exercise 20

$\{x \mid x > -7 \text{ and } x > -4\}$

Exercise 21

$\{x \mid x \geq -1 \text{ or } x > 6\}$

Answer

This set is the same as $\{x \mid x \geq -1\}$, which is $[-1, \infty)$ in interval notation. Graph of the set:



Exercise 22


$\{x \mid x \geq 7 \text{ or } x < -2\}$

Exercise 23

$\{x \mid x \geq 6 \text{ or } x > -3\}$

Answer

This set is the same as $\{x \mid x > -3\}$, which is $(-3, \infty)$ in interval notation. Graph of the set:



Exercise 24


$\{x \mid x \leq 1 \text{ or } x > 0\}$

Exercise 25

$\{x \mid x < 2 \text{ and } x < -7\}$

Answer

This set is the same as $\{x \mid x < -7\}$, which is $(-\infty, -7)$ in interval notation. Graph of the set:



Exercise 26


$\{x \mid x \leq -3 \text{ and } x < -5\}$

Exercise 27

$\{x \mid x \leq -3 \text{ or } x \geq 4\}$

Answer

This set is the union of two intervals, $(-\infty, -3] \cup [4, \infty)$. Graph of the set:



Exercise 28

$\{x \mid x < 11 \text{ or } x \leq 8\}$

Exercise 29

$\{x \mid x \geq 5 \text{ and } x \leq 1\}$

Answer

There are no numbers that satisfy both inequalities. Thus, there is no intersection. Graph of the set:

Exercise 30


$\{x \mid x < 5 \text{ or } x < 10\}$

Exercise 31

$\{x \mid x \leq 5 \text{ and } x \geq -1\}$

Answer

This set is the same as $\{x \mid -1 \leq x \leq 5\}$, which is $[-1, 5]$ in interval notation. Graph of the set



Exercise 32

$\{x \mid x > -3 \text{ and } x < -6\}$

In Exercises 33 - 44, solve the inequality. Express your answer in both interval and set-builder notations, and graph the solution on a number line. Answers include worked out solutions.


Exercise 33

$-8x-3 \leq -16x-1$

Answer

$$\begin{aligned} & -8x-3 \leq -16x-1 \\ \Longrightarrow \quad & -8x+16x \leq -1+3 \\ \Longrightarrow \quad & 8x \leq 2 \\ \Longrightarrow \quad & x \leq \frac{1}{4} \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: $\{x \mid x \leq \frac{1}{4}\}$

Interval Notation: $(-\infty, \frac{1}{4}]$

Exercise 34

$6x-6 > 3x+3$


Exercise 35

$-12x+5 \leq -3x-4$

Answer

$$\begin{aligned} & -12x+5 \leq -3x-4 \\ \Longrightarrow \quad & -12x+3x \leq -4-5 \\ \Longrightarrow \quad & -9x \leq -9 \\ \Longrightarrow \quad & x \geq 1 \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: \(\\{x \, | \, x\geq 1\}\)

Interval Notation: $[1, \infty)$

Exercise 36

$7x+3 \leq -2x-8$

Exercise 37

$-11x-9 < -3x+1$

Answer

$$\begin{aligned} & -11x-9 < -3x+1 \\ \Longrightarrow \quad & -11x+3x < 1+9 \\ \Longrightarrow \quad & -8x < 10 \\ \Longrightarrow \quad & x > -\frac{5}{4} \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: $\{x \mid x > -\frac{5}{4}\}$

Interval Notation: $(-\frac{5}{4}, \infty)$

Exercise 38

$4x - 8 \geq -4x - 5$

Exercise 39

$4x - 5 > 5x - 7$

Answer

$$\begin{aligned} & 4x - 5 > 5x - 7 \\ \implies \quad & 4x - 5x > -7 + 5 \\ \implies \quad & -x > -2 \\ \implies \quad & x < 2 \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: $\{x \mid x < 2\}$

Interval Notation: $(-\infty, 2)$

Exercise 40

$-14x + 4 > -6x + 8$


Exercise 41

$2x - 1 > 7x + 2$

Answer

$$\begin{aligned} & 2x - 1 > 7x + 2 \\ \implies \quad & 2x - 7x > 2 + 1 \\ \implies \quad & -5x > 3 \\ \implies \quad & x < -\frac{3}{5} \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: $\{x \mid x < -\frac{3}{5}\}$

Interval Notation: $(-\infty, -\frac{3}{5})$

Exercise 42

$-3x-2>-4x-9$


Exercise 43

$-3x+3<-11x-3$

Answer

$$\begin{aligned} & -3x+3<-11x-3 \\ \Longrightarrow \quad & -3x+11x<-3-3 \\ \Longrightarrow \quad & 8x<-6 \\ \Longrightarrow \quad & x<-\frac{3}{4} \end{aligned}$$

Thus, the solution interval is



Set-Builder Notation: $\{x \mid x<-\frac{3}{4}\}$

Interval Notation: $(-\infty,-\frac{3}{4})$

Exercise 44

$6x+3<8x+8$

In Exercises 45-53, solve the compound inequality. Express your answer in both interval and set-builder notations, and graph the solution on a number line. Worked out solutions are provided in the Answers for #45 and #47. Only the interval notation version of the answer is given for Answers in #49-82.

Exercise 45

$2x-1<4$ or $7x+1\geq -4$

Answer

$$\begin{aligned} & 2x-1<4 \text{ or } 7x+1\geq -4 \\ \Longrightarrow \quad & 2x<5 \quad \text{or} \quad 7x\geq -5 \\ \Longrightarrow \quad & x<\frac{5}{2} \quad \text{or} \quad x\geq -\frac{5}{7} \end{aligned}$$



For the union, shade anything shaded in either graph. The solution is the set of all real numbers. $(-\infty,\infty)$.

Set-Builder Notation: $\{x \mid x\in\mathbb{R}\}$

Interval Notation: $(-\infty, \infty)$

Exercise 46


$-8x+9<-3$ and $-7x+1>3$

Exercise 47


$-6x-4<-4$ and $-3x+7\geq -5$

Answer

$$\begin{aligned} & -6x-4<-4 \text{ and } -3x+7\geq -5 \\ \Longrightarrow \quad & -6x<0 \quad \text{and} \quad -3x\geq -12 \\ \Longrightarrow \quad & x>0 \quad \text{and} \quad x\leq 4 \\ \Longrightarrow \quad & 0<x\leq 4 \end{aligned}$$



The intersection is all points shaded in both graphs, so the solution is



Set-Builder Notation: $\{x \mid 0<x\leq 4\}$

Interval Notation: $(0, 4]$

Exercise 48

$-3x+3\leq 8$ and $-3x-6>-6$

Exercise 49

$8x+5\leq -1$ and $4x-2>-1$

Answer

No Solution
Set-Builder Notation: { }

Exercise 50

$-x-1<7$ and $-6x-9\geq 8$

Exercise 51

$-3x+8\leq -5$ or $-2x-4\geq -3$

Answer

No Solution
Set-Builder Notation: { }

Exercise 52

$-6x-7<-3$ and $-8x\geq 3$

Exercise 53

$9x-9\leq 9$ and $5x>-1$

Answer

Interval Notation: $\left(-\frac{1}{5},2\right]$

Exercise 2.8E.54

$-7x+3<-3$ or $-8x\geq 2$

Exercise 2.8E.55

$3x-5<4$ and $-x+9>3$

Answer

Interval Notation: $(-\infty,3)$

Exercise 2.8E.56

$-8x-6<5$ or $4x-1\geq 3$

Exercise 2.8E.57

$9x+3\leq -5$ or $-2x-4\geq 9$

Answer

Interval Notation: $\left(-\infty,-\frac{8}{9}\right)$

Exercise 2.8E.58

$-7x+6<-4$ or $-7x-5>7$

Exercise 2.8E.59

$4x-2\leq 2$ or $3x-9\geq 3$

Answer

Add texts here. $(-\infty,1]\cup[4,\infty)$

Exercise 2.8E.60

$-5x+5<-4$ or $-5x-5\geq -5$

Exercise 2.8E.61

$5x+1<-6$ and $3x+9>-4$

Answer

Interval Notation: $\left(-\frac{13}{3},-\frac{7}{5}\right)$

Exercise 2.8E.62

$7x+2<-5$ or $6x-9\geq -7$

Exercise 2.8E.63

$-7x-7<-2$ and $3x\geq 3$

Answer

Interval Notation: $[1,\infty)$

Exercise 2.8E.64

$4x+1<0$ or $8x+6>9$

Exercise 2.8E.65

$7x+8<-3$ and $8x+3\geq -9$

Answer

No Solution
Set-Builder Notation: $\{\ \}$

Exercise 2.8E.66

$3x<2$ and $-7x-8\geq 3$

Exercise 2.8E.67

$-5x+2\leq -2$ and $-6x+2\geq 3$

Answer

No Solution
Set-Builder Notation: $\{\ \}$

Exercise 2.8E.68

$4x-1\leq 8$ or $3x-9>0$

Exercise 2.8E.69

$2x-5\leq 1$ and $4x+7>7$

Answer

$(0, 3]$

Exercise 2.8E.70

$3x + 1 < 0$ or $5x + 5 > -8$

Exercise 2.8E.71

$-8x + 7 \leq 9$ or $-5x + 6 > -2$

Answer

$(-\infty, \infty)$

Exercise 2.8E.72

$x - 6 \leq -5$ and $6x - 2 > -3$

Exercise 2.8E.73

$-4x - 8 < 4$ or $-4x + 2 > 3$

Answer

$(-\infty, \infty)$

Exercise 2.8E.74

$9x - 5 < 2$ or $-8x - 5 \geq -6$

Exercise 2.8E.75

$-9x - 5 \leq -3$ or $x + 1 > 3$

Answer

$\left(-\frac{2}{9}, \infty\right)$

Exercise 2.8E.76

$-5x - 3 \leq 6$ and $2x - 1 \geq 6$

Exercise 2.8E.77

$-1 \leq -7x - 3 \leq 2$

Answer

Interval Notation: $\left[-\frac{5}{7}, -\frac{2}{7}\right]$


Exercise 2.8E.78

$0 < 5x - 5 < 9$

Exercise 79

$5 < 9x - 3 \leq 6$

Answer

Graph of the solution: 

Set-Builder Notation: $\{x \mid \frac{8}{9} < x \leq 1\}$

Interval Notation: $(\frac{8}{9}, 1]$

Exercise 2.8E.80

$-6 < 7x + 3 \leq 2$

Exercise 2.8E.81

$-2 < -7x + 6 < 6$

Answer

Interval Notation: $\left(0, \frac{8}{7}\right)$

Exercise 82

$-9 < -2x + 5 \leq 1$

In Exercises 83-94, solve the given inequality for x. Graph the solution set on a number line, then use interval and setbuilder notation to describe the solution set. **Only the interval notation version of the answer is given for Answers in #83-94.**

Exercise 2.8E.83

$-\frac{1}{3} < \frac{x}{2} + \frac{1}{4} < \frac{1}{3}$

Answer

Interval Notation: $\left(-\frac{7}{6}, \frac{1}{6}\right)$

Exercise 2.8E.84

$-\frac{1}{5} < \frac{x}{2} - \frac{1}{4} < \frac{1}{5}$

Exercise 2.8E.85

$-\frac{1}{2} < \frac{1}{3} - \frac{x}{2} < \frac{1}{2}$

Answer

Interval Notation: $\left(-\frac{1}{3}, \frac{5}{3}\right)$

Exercise 86

$-\frac{2}{3} \leq \frac{1}{2} - \frac{x}{5} \leq \frac{2}{3}$

Exercise 2.8E.87

$-1 < x - \frac{x+1}{5} < 2$

Answer

Interval Notation: $\left(-1, \frac{11}{4}\right)$

Exercise 2.8E.88

$-2 < x - \frac{2x-1}{3} < 4$

Exercise 2.8E.89

$-2 < \frac{x+1}{2} - \frac{x+1}{3} \leq 2$

Answer

Interval Notation: $(-13, 11]$

Exercise 2.8E.90

$-3 < \frac{x-1}{3} - \frac{2x-1}{5} \leq 2$

Exercise 2.8E.91

$x < 4 - x < 5$

Answer

Interval Notation: $(-1, 2)$

Exercise 2.8E.92

$-x < 2x + 3 \leq 7$

Exercise 2.8E.93

$-x < x + 5 \leq 11$

Answer

Interval Notation: $\left(-\frac{5}{2}, 6\right]$

Exercise 2.8E.94

$-2x < 3 - x \leq 8$

More Practice with Compound Inequalities

Exercise $2.8E.95$

For each graph below, describe the interval (a) using set-builder notation and (b) using interval notation.


1.


2.


3.


4.


5.


6.


7.


8.


Answer

1. (a) $\{x|\ 3 \leq x \leq 4\}$ (b) $[3, 4]$
3. (a) $\{x|\ -4 < x < 1\}$ (b) $(-4, 1)$
5. (a) $\{x|\ -1 \leq x \leq 5\}$ (b) $[-1, 5]$
7. (a) $\{x|\ -1 < x \leq 5\}$ (b) $(-1, 5]$

2.8E: Exercises for Section 2.8 is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **1.5: Chapter 1 Exercises with Solutions** has no license indicated.

Chapter 2 Review Exercises

Chapter 2 Review Exercises

Solve Equations using the Subtraction and Addition Properties of Equality

Verify a Solution of an Equation

In the following exercises, determine whether each number is a solution to the equation.

Exercise 1

$10x - 1 = 5x; x = \frac{1}{5}$

Exercise 2

$w + 2 = \frac{5}{8}; w = \frac{3}{8}$

Answer

no

Exercise 3

$-12n + 5 = 8n; n = -\frac{5}{4}$

Exercise 4

$6a - 3 = -7a, a = \frac{3}{13}$

Answer

yes

Solve Equations using the Subtraction and Addition Properties of Equality

In the following exercises, solve each equation using the Subtraction Property of Equality.

Exercise 5

$x + 7 = 19$

Exercise 6

$y + 2 = -6$

Answer

$y = -8$

Exercise 7

$a + \frac{1}{3} = \frac{5}{3}$

Exercise 8

$n+3.6=5.1$

Answer

$n=1.5$

In the following exercises, solve each equation using the Addition Property of Equality.

Exercise 9

$u-7=10$

Exercise 10

$x-9=-4$

Answer

$x=5$

Exercise 11

$c-\frac{3}{11}=\frac{9}{11}$

Exercise 12

$p-4.8=14$

Answer

$p=18.8$

In the following exercises, solve each equation.

Exercise 13

$n-12=32$

Exercise 14

$y+16=-9$

Answer

$y=-25$

Exercise 15

$f+\frac{2}{3}=4$

Exercise 16

$d-3.9=8.2$

Answer

$d = 12.1$

Solve Equations That Require Simplification

In the following exercises, solve each equation.

Exercise 17

$y + 8 - 15 = -3$

Exercise 18

$7x + 10 - 6x + 3 = 5$

Answer

$x = -8$

Exercise 19

$6(n - 1) - 5n = -14$

Exercise 20

$8(3p + 5) - 23(p - 1) = 35$

Answer

$p = -28$

Translate to an Equation and Solve

In the following exercises, translate each English sentence into an algebraic equation and then solve it.

Exercise 21

The sum of -6 and m is 25

Exercise 22

Four less than n is 13

Answer

$n - 4 = 13; n = 17$

Translate and Solve Applications

In the following exercises, translate into an algebraic equation and solve.

Exercise 23

Rochelle's daughter is 11 years old. Her son is 3 years younger. How old is her son?

Exercise 24

Tan weighs 146 pounds. Minh weighs 15 pounds more than Tan. How much does Minh weigh?

Answer

161 pounds

Exercise 25

Peter paid $9.75 to go to the movies, which was $46.25 less than he paid to go to a concert. How much did he pay for the concert?

Exercise 26

Elissa earned $\$152.84$this week, which was $\$2..65$more than she earned last week. How much did she earn last week?

Answer

$\$131.19$

Solve Equations using the Division and Multiplication Properties of Equality

Solve Equations Using the Division and Multiplication Properties of Equality

In the following exercises, solve each equation using the division and multiplication properties of equality and check the solution.

Exercise 27

$8x = 72$

Exercise 28

$13a = -65$

Answer

$a = -5$

Exercise 29

$0.25p = 5.25$

Exercise 30

$-y = 4$

Answer

$y = -4$

Exercise 31

$\frac{n}{6} = 18$

Exercise 32

$\frac{y}{-10} = 30$

Answer

$y = -300$

Exercise 33

$36 = \frac{3}{4}x$

Exercise 34

$\frac{5}{8}u = \frac{15}{16}$

Answer

$u = \frac{3}{2}$

Exercise 35

$-18m = -72$

Exercise 36

$\frac{c}{9} = 36$

Answer

$c = 324$

Exercise 37

$0.45x = 6.75$

Exercise 38

$\frac{11}{12} = \frac{2}{3}y$

Answer

$y = \frac{11}{8}$

Solve Equations That Require Simplification

In the following exercises, solve each equation requiring simplification.

Exercise 39

$5r - 3r + 9r = 35 - 2$

Exercise 40

$24x + 8x - 11x = -7 - 14$

Answer

$x = -1$

Exercise 41

$\frac{11}{12}n - \frac{5}{6}n = 9 - 5$

Exercise 42

$-9(d - 2) - 15 = -24$

Answer

$d = 3$

Translate to an Equation and Solve

In the following exercises, translate to an equation and then solve.

Exercise 43

143 is the product of -11 and y

Exercise 44

The quotient of b and and 9 is -27

Answer

$\frac{b}{9} = -27; b = -243$

Exercise 45

The sum of q and one-fourth is one.

Exercise 46

The difference of s and one-twelfth is one fourth.

Answer

$s - \frac{1}{12} = \frac{1}{4}; s = \frac{1}{3}$

Translate and Solve Applications

In the following exercises, translate into an equation and solve.

Exercise 47

Ray paid $21 for 12 tickets at the county fair. What was the price of each ticket?

Exercise 48

Janet gets paid $\$24$ per hour. She heard that this is $\frac{3}{4}$ of what Adam is paid. How much is Adam paid per hour?

Answer

$\$32$

Solve Equations with Variables and Constants on Both Sides

Solve an Equation with Constants on Both Sides

In the following exercises, solve the following equations with constants on both sides.

Exercise 49

$8p+7=47$

Exercise 50

$10w-5=65$

Answer

$w=7$

Exercise 51

$3x+19=-47$

Exercise 52

$32=-4-9n$

Answer

$n=-4$

Solve an Equation with Variables on Both Sides

In the following exercises, solve the following equations with variables on both sides.

Exercise 53

$7y=6y-13$

Exercise 54

$5a+21=2a$

Answer

$a=-7$

Exercise 55

$k=-6k-35$

Exercise 56

$4x - \frac{3}{8} = 3x$

Answer

$x = \frac{3}{8}$

Solve an Equation with Variables and Constants on Both Sides

In the following exercises, solve the following equations with variables and constants on both sides.

Exercise 57

$12x - 9 = 3x + 45$

Exercise 58

$5n - 20 = -7n - 80$

Answer

$n = -5$

Exercise 59

$4u + 16 = -19 - u$

Exercise 60

$\frac{5}{8}c - 4 = \frac{3}{8}c + 4$

Answer

$c = 32$

Use a General Strategy for Solving Linear Equations

Solve Equations Using the General Strategy for Solving Linear Equations

In the following exercises, solve each linear equation.

Exercise 61

$6(x + 6) = 24$

Exercise 62

$9(2p - 5) = 72$

Answer

$p = \frac{13}{2}$

Exercise 63

$-(s+4)=18$

Exercise 64

$8+3(n-9)=17$

Answer

$n=12$

Exercise 65

$23-3(y-7)=8$

Exercise 66

$\frac{1}{3}(6m+21)=m-7$

Answer

$m=-14$

Exercise 67

$4(3.5y+0.25)=365$

Exercise 68

$0.25(q-8)=0.1(q+7)$

Answer

$q=18$

Exercise 69

$8(r-2)=6(r+10)$

Exercise 70

$5+7(2-5x)=2(9x+1)$
$-(13x-57)$

Answer

$x=-1$

Exercise 71

$(9n+5)-(3n-7)$
$=20-(4n-2)$

Exercise 72

$2[-16+5(8k-6)]$
$=8(3-4k)-32$

Answer

$k=\frac{3}{4}$

Classify Equations

In the following exercises, classify each equation as a conditional equation, an identity, or a contradiction and then state the solution.

Exercise 73

$17y-3(4-2y)=11(y-1)$
$+12y-1$

Exercise 74

$9u+32=15(u-4)$
$-3(2u+21)$

Answer

contradiction; no solution

Exercise 75

$-8(7m+4)=-6(8m+9)$

Exercise 76

$21(c-1)-19(c+1)$
$=2(c-20)$

Answer

identity; all real numbers

Solve Equations with Fractions and Decimals

Solve Equations with Fraction Coefficients

In the following exercises, solve each equation with fraction coefficients.

Exercise 77

$\frac{2}{5}n-\frac{1}{10}=\frac{7}{10}$

Exercise 78

$\frac{1}{3}x+\frac{1}{5}x=8$

Answer

$x=15$

Exercise 79

$\frac{3}{4}a-\frac{1}{3}=\frac{1}{2}a-\frac{5}{6}$

Exercise 80

$\frac{1}{2}(k-3)=\frac{1}{3}(k+16)$

Answer

$k=41$

Exercise 81

$\frac{3x-2}{5}=\frac{3x+4}{8}$

Exercise 82

$\frac{5y-1}{3}+4=\frac{-8y+4}{6}$

Answer

$y=-1$

Solve Equations with Decimal Coefficients

In the following exercises, solve each equation with decimal coefficients.

Exercise 83

$0.8x-0.3=0.7x+0.2$

Exercise 84

$0.36u+2.55=0.41u+6.8$

Answer

$u=-85$

Exercise 85

$0.6p-1.9=0.78p+1.7$

Exercise 86

$0.6p-1.9=0.78p+1.7$

Answer

$d=-20$

Solve a Formula for a Specific Variable

Use the Distance, Rate, and Time Formula

In the following exercises, solve.

Exercise 87

Natalie drove for $7\frac{1}{2}$ hours at 60 miles per hour. How much distance did she travel?

Exercise 88

Mallory is taking the bus from St. Louis to Chicago. The distance is 300 miles and the bus travels at a steady rate of 60 miles per hour. How long will the bus ride be?

Answer

5 hours

Exercise 89

Aaron's friend drove him from Buffalo to Cleveland. The distance is 187 miles and the trip took 2.75 hours. How fast was Aaron's friend driving?

Exercise 90

Link rode his bike at a steady rate of 15 miles per hour for $2\frac{1}{2}$ hours. How much distance did he travel?

Answer

37.5 miles

Solve a Formula for a Specific Variable

In the following exercises, solve.

Exercise 91

Use the formula. d=rt to solve for t

1. when d=510 and r=60
2. in general

Exercise 92

Use the formula. d=rt to solve for r

1. when when d=451 and t=5.5
2. in general

Answer

1. r=82mph
2. $r = \frac{D}{t}$

Exercise 93

Use the formula $A = \frac{1}{2}bh$ to solve for b

1. when A=390 and h=26
2. in general

Exercise 94

Use the formula $A = \frac{1}{2}bh$ to solve for b

1. when A=153 and b=18
2. in general

Answer

1. $h = 17$
2. $h = \frac{2A}{b}$

Exercise 95

Use the formula I=Prt to solve for the principal, *P* for

1. I=$2,501,r=4.1%, t=5 years
2. in general

Exercise 96

Solve the formula 4x+3y=6 for *y*

1. when x=−2
2. in general

Answer

ⓐ $y = \frac{14}{3}$ ⓑ $y = \frac{6-4x}{3}$

Exercise 97

Solve $180 = a + b + c$ for c

Exercise 98

Solve the formula $V = LWH$ for H

Answer

$H = \frac{V}{LW}$

Solve Linear Inequalities

Graph Inequalities on the Number Line

In the following exercises, graph each inequality on the number line.


Exercise 99


1. $x \leq 4$
2. x>−2
3. x<1


Exercise 100

1. x>0
2. x<−3
3. $x \geq -1$

Answer

1. 

2. 

3. 

In the following exercises, graph each inequality on the number line and write in interval notation.


Exercise 101


1. $x < -1$
2. $x \geq -2.5$
3. $x \leq \frac{5}{4}$


Exercise 102

1. $x > 2$
2. $x \leq -1.5$
3. $x \geq \frac{5}{3}$

Answer

1. 

2. 

3. 

Solve Inequalities using the Subtraction and Addition Properties of Inequality

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 103

$n - 12 \leq 23$

Exercise 104

$m + 14 \leq 56$

Answer




Exercise 105

$a+\frac{2}{3} \geq \frac{7}{12}$

Exercise 106

$b-\frac{7}{8} \geq -\frac{1}{2}$

Answer



Solve Inequalities using the Division and Multiplication Properties of Inequality

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 107

$9x > 54$

Exercise 108

$-12d \leq 108$

Answer




Exercise 109

$\frac{5}{2}j < -60$

Exercise 110

$\frac{q}{-2} \geq -24$

Answer



Solve Inequalities That Require Simplification

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 111

$6p > 15p - 30$

Exercise 112

$9h - 7(h-1) \leq 4h - 23$

Answer




Exercise 113

$5n - 15(4-n) < 10(n-6) + 10n$

Exercise 114

$\frac{3}{8}a - \frac{1}{12}a > \frac{5}{12}a + \frac{3}{4}$

Answer



Translate to an Inequality and Solve

In the following exercises, translate and solve. Then write the solution in interval notation and graph on the number line.


Exercise 115

Five more than z is at most 19.

Exercise 116

Three less than c is at least 360.

Answer




Exercise 117

Nine times n exceeds 42.

Exercise 118

Negative two times a is no more than 8.

Answer



Everyday Math

Exercise 119

Describe how you have used two topics from this chapter in your life outside of your math class during the past month.

Chapter 2 Practice Test

Exercise 1

Determine whether each number is a solution to the equation $6x - 3 = x + 20$

1. 5
2. $\frac{23}{5}$

Answer

1. no
2. yes

In the following exercises, solve each equation.

Exercise 2

$n - \frac{2}{3} = \frac{1}{4}$

Exercise 3

$\frac{9}{2}c = 144$

Answer

c=32

Exercise 4

$4y - 8 = 16$

Exercise 5

$-8x - 15 + 9x - 1 = -21$

Answer

$x = -5$

Exercise 6

$-15a = 120$

Exercise 7

$\frac{2}{3}x = 6$

Answer

$x = 9$

Exercise 8

$x - 3.8 = 8.2$

Exercise 9

$10y = -5y - 60$

Answer

$y = -4$

Exercise 10

$8n - 2 = 6n - 12$

Exercise 11

$9m - 2 - 4m - m = 42 - 8$

Answer

$m = 9$

Exercise 12

$-5(2x - 1) = 45$

Exercise 13

$-(d - 9) = 23$

Answer

$d = -14$

Exercise 14

$\frac{1}{4}(12m - 28) = 6 - 2(3m - 1)$

Exercise 15

$2(6x - 5) - 8 = -22$

Answer

$x = -\frac{1}{3}$

Exercise 16

$8(3a-5)-7(4a-3)=20-3a$

Exercise 17

$\frac{1}{4}p-\frac{1}{3}=\frac{1}{2}$

Answer

$p=\frac{10}{3}$

Exercise 18

$0.1d+0.25(d+8)=4.1$

Exercise 19

$14n-3(4n+5)=-9+2(n-8)$

Answer

contradiction; no solution

Exercise 20

$9(3u-2)-4[6-8(u-1)]=3(u-2)$

Exercise 21

Solve the formula x−2y=5 for *y*

1. when x=−3
2. in general

Answer

1. y=4
2. $y=\frac{5-x}{2}$

In the following exercises, graph on the number line and write in interval notation.


Exercise 22

$x\geq -3.5$

Exercise 23

$x<\frac{11}{4}$

Answer



https://math.libretexts.org/@go/page/30487

In the following exercises,, solve each inequality, graph the solution on the number line, and write the solution in interval notation.


Exercise 24

$8k \geq 5k - 120$

Exercise 25

$3c - 10(c-2) < 5c + 16$

Answer



In the following exercises, translate to an equation or inequality and solve.

Exercise 26

4 less than twice *x* is 16.

Exercise 27

Fifteen more than *n* is at least 48.

Answer

$n + 15 \geq 48; n \geq 33$

Exercise 28

Samuel paid $25.82 for gas this week, which was $3.47 less than he paid last week. How much had he paid last week?

Exercise 29

Jenna bought a coat on sale for $\$120$, which was $\frac{2}{3}$ of the original price. What was the original price of the coat?

Answer

$120 = \frac{2}{3}p$; The original price was $\$180$

Exercise 30

Sean took the bus from Seattle to Boise, a distance of 506 miles. If the trip took $7\frac{2}{3}$ hours, what was the speed of the bus?

Review for 2.7 Ratio and Proportions and Similar Triangles

Solve Proportions

In the following exercises, solve.

Exercise 74

$\frac{x}{4} = \frac{3}{5}$

Answer

$\frac{12}{5}$

Exercise 75

$\frac{3}{y} = \frac{9}{5}$

Exercise 76

$\frac{s}{s+20} = \frac{3}{7}$

Answer

15

Exercise 77

$\frac{t-3}{5} = \frac{t+2}{9}$

In the following exercises, solve using proportions.

Exercise 78

Rachael had a 21 ounce strawberry shake that has 739 calories. How many calories are there in a 32 ounce shake?

Answer

1161calories

Exercise 79


Leo went to Mexico over Christmas break and changed $\$525$ dollars into Mexican pesos. At that time, the exchange rate had $\$1$ US is equal to 16.25 Mexican pesos. How many Mexican pesos did he get for his trip?

Solve Similar Figure Applications

In the following exercises, solve.

Exercise 80

ΔABC is similar to ΔXYZ. The lengths of two sides of each triangle are given in the figure. Find the lengths of the third sides.




Answer

$b = 9; x = 2\frac{1}{3}$

Exercise 81

On a map of Europe, Paris, Rome, and Vienna form a triangle whose sides are shown in the figure below. If the actual distance from Rome to Vienna is 700 miles, find the distance from

1. a. Paris to Rome
2. b. Paris to Vienna



Exercise 82

Tony is 5.75 feet tall. Late one afternoon, his shadow was 8 feet long. At the same time, the shadow of a nearby tree was 32 feet long. Find the height of the tree.

Answer

23 feet

Exercise 83

The height of a lighthouse in Pensacola, Florida is 150 feet. Standing next to the statue, 5.5 foot tall Natalie cast a 1.1 foot shadow How long would the shadow of the lighthouse be?


Review for 2.9 Compound Inequalities

Solve Compound Inequalities with "and"

In each of the following exercises, solve each inequality, graph the solution, and write the solution in interval notation.

98. $x \leq 5$ and $x > -3$


Answer



99. $4x-2\leq 4$ and $7x-1>-8$

100. $5(3x-2)\leq 5$ and $4(x+2)<3$


Answer



101. $34(x-8)\leq 3$ and $15(x-5)\leq 3$

102. $34x-5\geq -2$ and $-3(x+1)\geq 6$

Answer




103. $-5\leq 4x-1<7$

Solve Compound Inequalities with "or"

In the following exercises, solve each inequality, graph the solution on the number line, and write the solution in interval notation.

104. $5-2x\leq -1$ or $6+3x\leq 4$


Answer



105. $3(2x-3)<-5$ or $4x-1>3$

106. $34x-2>4$ or $4(2-x)>0$


Answer



107. $2(x+3)\geq 0$ or $3(x+4)\leq 6$

108. $12x-3\leq 4$ or $13(x-6)\geq -2$

Answer



Solve Applications with Compound Inequalities

In the following exercises, solve.

109. Liam is playing a number game with his sister Audry. Liam is thinking of a number and wants Audry to guess it. Five more than three times her number is between 2 and 32. Write a compound inequality that shows the range of numbers that Liam might be thinking of.

110. Elouise is creating a rectangular garden in her back yard. The length of the garden is 12 feet. The perimeter of the garden must be at least 36 feet and no more than 48 feet. Use a compound inequality to find the range of values for the width of the garden.

Answer

$6 \leq w \leq 12$

This page titled Chapter 2 Review Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **Chapter 2 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **Chapter 8 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **Chapter 2 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/intermediate-algebra-2e.

CHAPTER OVERVIEW

3: Math Models

- 3.1: Solve a Formula for a Specific Variable
  - 3.1E: Exercises
- 3.2: Use a Problem-Solving Strategy
  - 3.2E: Exercises
- 3.3: Solve Percent Applications
  - 3.3E: Exercises
- 3.4: Solve Mixture Applications
  - 3.4E: Exercises
- 3.5: Triangles, Rectangles, and the Pythagorean Theorem
  - 3.5E: Exercises
- 3.6: Solve Uniform Motion Applications
  - 3.6E: Exercises
- 3.7: Solve Applications with Linear Inequalities
  - 3.7E: Exercises
- Chapter 3 Review Exercises

Thumbnail: https://www.wikihow.com/Make-a-Mathematical-Model

This page titled 3: Math Models is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

3.1: Solve a Formula for a Specific Variable

Learning Objectives

By the end of this section, you will be able to:

- Use the Distance, Rate, and Time formula
- Solve a formula for a specific variable

Quiz

Before you get started, take this readiness quiz.

1. Solve: $15t = 120$.
   If you missed this problem, review Exercise 2.2.1.
2. Solve: $6x + 24 = 96$.
   If you missed this problem, review Exercise 2.3.1.

Use the Distance, Rate, and Time Formula

One formula you will use often in algebra and in everyday life is the formula for distance traveled by an object moving at a constant rate. Rate is an equivalent word for "speed." The basic idea of rate may already familiar to you. Do you know what distance you travel if you drive at a steady rate of 60 miles per hour for 2 hours? (This might happen if you use your car's cruise control while driving on the highway.) If you said 120 miles, you already know how to use this formula!

DISTANCE, RATE, AND TIME

For an object moving at a uniform (constant) rate, the distance traveled, the elapsed time, and the rate are related by the formula:

$$d = rt \quad \text{where} \quad \begin{aligned} d &= \text{distance} \\ r &= \text{rate} \\ t &= \text{time} \end{aligned} \tag{3.1.1}$$

We will use the Strategy for Solving Applications that we used earlier in this chapter. When our problem requires a formula, we change Step 4. In place of writing a sentence, we write the appropriate formula. We write the revised steps here for reference.

SOLVE AN APPLICATION (WITH A FORMULA).

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
4. **Translate** into an equation. Write the appropriate formula for the situation. Substitute in the given information.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

You may want to create a mini-chart to summarize the information in the problem. See the chart in this first example.

Example 3.1.1

Jamal rides his bike at a uniform rate of 12 miles per hour for $3\frac{1}{2}$ hours. What distance has he traveled?

Solution

Step 1. Read the problem.	

Step 2. Identify what you are looking for.	distance traveled
Step 3. Name. Choose a variable to represent it.	Let d = distance.
Step 4. Translate: Write the appropriate formula.	$d = rt$
	$d = ?$ $r = 12$ mph $t = 3\frac{1}{2}$ hours
Substitute in the given information.	$d = 12 \cdot 3\frac{1}{2}$
Step 5. Solve the equation.	$d = 42$ miles
Step 6. Check	
Does 42 miles make sense?	
Jamal rides:	
12 miles in 1 hour, 24 miles in 2 hours, 36 miles in 3 hours, 42 miles in $3\frac{1}{2}$ hours is reasonable 48 miles in 4 hours.	
Step 7. Answer the question with a complete sentence.	Jamal rode 42 miles.

Try It 3.1.2

Lindsay drove for $5\frac{1}{2}$ hours at 60 miles per hour. How much distance did she travel?

Answer

330 miles

Try It 3.1.3

Trinh walked for $2\frac{1}{3}$ hours at 3 miles per hour. How far did she walk?

Answer

7 miles

Example 3.1.4

Rey is planning to drive from his house in San Diego to visit his grandmother in Sacramento, a distance of 520 miles. If he can drive at a steady rate of 65 miles per hour, how many hours will the trip take?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	How many hours (time)
Step 3. Name. Choose a variable to represent it.	Let t = time.
	$d = 520$ miles $r = 65$ mph $t = ?$ hours

Step 4. Translate. Write the appropriate formula.	$d = rt$
Substitute in the given information.	$520 = 65t$
Step 5. Solve the equation.	$t = 8$
Step 6. Check. Substitute the numbers into the formula and make sure the result is a true statement.	
$d = rt$ $520 \stackrel{?}{=} 65 \cdot 8$ $520 = 520 \checkmark$	
Step 7. Answer the question with a complete sentence. Rey's trip will take 8 hours.	

Try It 3.1.5

Lee wants to drive from Phoenix to his brother's apartment in San Francisco, a distance of 770 miles. If he drives at a steady rate of 70 miles per hour, how many hours will the trip take?

Answer

11 hours

Try It 3.1.6

Yesenia is 168 miles from Chicago. If she needs to be in Chicago in 3 hours, at what rate does she need to drive?

Answer

56 mph

Solve a Formula for a Specific Variable

You are probably familiar with some geometry formulas. A formula is a mathematical description of the relationship between variables. Formulas are also used in the sciences, such as chemistry, physics, and biology. In medicine they are used for calculations for dispensing medicine or determining body mass index. Spreadsheet programs rely on formulas to make calculations. It is important to be familiar with formulas and be able to manipulate them easily.

In Exercise 3.1.1 and Exercise 3.1.4, we used the formula $d = rt$. This formula gives the value of d, distance, when you substitute in the values of r and t, the rate and time. But in Exercise 3.1.4, we had to find the value of t. We substituted in values of d and r and then used algebra to solve for tt. If you had to do this often, you might wonder why there is not a formula that gives the value of t when you substitute in the values of d and r. We can make a formula like this by solving the formula $d = rt$ for t.

To solve a formula for a specific variable means to isolate that variable on one side of the equals sign with a coefficient of 1. All other variables and constants are on the other side of the equals sign. To see how to solve a formula for a specific variable, we will start with the distance, rate and time formula.

Example 3.1.7

Solve the formula d=rt for t:

1. when d=520 and r=65
2. in general

Solution

We will write the solutions side-by-side to demonstrate that solving a formula in general uses the same steps as when we have numbers to substitute.

1. when d=520 and r=65		2. in general	
Write the formula.	$d = rt$	Write the formula.	$d = rt$
Substitute.	$520 = 65t$		
Divide, to isolate t.	$\frac{520}{65} = \frac{65t}{65}$	Divide, to isolate tt.	$\frac{d}{r} = \frac{rt}{t}$
Simplify.	$8 = t$	Simplify.	$\frac{d}{r} = t$

We say the formula $t = \frac{d}{r}$ is solved for t.

Try It 3.1.8

Solve the formula $d = rt$ for r:

1. when d=180 and t=4
2. in general

Answer

1. $r = 45$
2. $r = \frac{d}{t}$

Try It 3.1.9

Solve the formula $d = rt$ for r:

1. when d=780 and t=12
2. in general

Answer

1. $r = 65$
2. \(r = \frac{d}{rt}\)

Example 3.1.10

Solve the formula $A = \frac{1}{2}bh$ for h:

1. when $A = 90$ and $b = 15$
2. in general

Solution

1. when $A = 90$ and $b = 15$		2. in general	
Write the formula.	$A = \frac{1}{2}bh$	Write the formula.	$A = \frac{1}{2}bh$
Substitute.	$90 = \frac{1}{2} \cdot 15 \cdot h$		
Clear the fractions.	$2 \cdot 90 = 2 \cdot \frac{1}{2}15h$	Clear the fractions.	$2 \cdot A = 2 \cdot \frac{1}{2}bh$
Simplify.	$180 = 15h$	Simplify.	$2A = bh$
Solve for h.	$12 = h$	Solve for hh.	$\frac{2A}{b} = h$

We can now find the height of a triangle, if we know the area and the base, by using the formula $h = \frac{2A}{b}$

Try It 3.1.11

Solve the formula $A = \frac{1}{2}bh$ for h:

1. when $A = 170$ and $b = 17$
2. in general

Answer

1. $h = 20$
2. $h = \frac{2A}{b}$

Try It 3.1.12

Solve the formula $A = \frac{1}{2}bh$ for h:

1. when $A = 62$ and $h = 31$
2. in general

Answer

1. $b = 4$
2. $b = \frac{2A}{h}$

The formula $I = Prt$ is used to calculate simple interest, *I*, for a principal, *P*, invested at rate, *r*, for *t* years.

Example 3.1.13

Solve the formula I=Prt to find the principal, P:

1. when I=$5,600, r=4% ,t=7years
2. in general

Solution

1. I=$5,600, r=4% ,t=7years		2. in general	
Write the formula.	$I = Prt$	Write the formula.	$I = Prt$
Substitute.	$5600 = P(0.04)(7)$		
Simplify.	$5600 = P(0.28)$	Simplify.	$I = P(rt)$
Divide, to isolate *P*.	$\frac{5600}{0.28} = \frac{P(0.28)}{0.28}$	Divide, to isolate *P*.	$\frac{I}{rt} = \frac{P(rt)}{rt}$
Simplify.	$20,000 = P$	Simplify.	$\frac{I}{rt} = P$
The principal is	$20,000		$P = \frac{I}{rt}$

Try It 3.1.14

Solve the formula I=Prt to find the principal, P:

1. when I=$2160, r=6% ,t=3 years
2. in general

Answer

1. $12000
2. $P = \frac{1}{rt}$

Try It 3.1.15

Solve the formula I=Prt to find the principal, P:

1. when I=$5400, r= 12% ,t=5 years
2. in general

Answer

1. $9000
2. $P = \frac{1}{rt}$

Later in this class, and in future algebra classes, you'll encounter equations that relate two variables, usually *x* and *y*. You might be given an equation that is solved for *y* and need to solve it for *x*, or vice versa. In the following example, we're given an equation with both *x* and *y* on the same side and we'll solve it for *y*.

Example 3.1.16

Solve the formula 3x+2y=18 for *y*:

1. when x=4
2. in general

Solution

1. when x=4		2. in general	
	$3x + 2y = 18$		$3x + 2y = 18$
Substitute.	$3(4) + 2y = 18$		
Subtract to isolate the y-term.	$12 - 12 + 2y = 18 - 12$	Subtract to isolate the y-term.	$3x - 3x + 2y = 18 - 3x$
Divide.	$\frac{2y}{2} = \frac{6}{2}$	Divide.	$\frac{2y}{2} = \frac{18}{2} - \frac{3x}{2}$
Simplify.	$y = 3$	Simplify.	$y = -\frac{3x}{2} + 9$

Try It 3.1.17

Solve the formula 3x+4y=10 for *y*:

1. when $x = \frac{14}{3}$
2. in general

Answer

1. $y = -1$
2. $y = \frac{10-3x}{4}$

Try It 3.1.18

Solve the formula 5x+2y=18 for *y*:

1. when $x = 4$

2. in general

Answer

1. $y = -1$
2. $y = \frac{18-5x}{2}$

In Exercise 3.1.7 through Exercise 3.1.18 we used the numbers in part 1 as a guide to solving in general in part 2. Now we will solve a formula in general without using numbers as a guide.

Example 3.1.19

Solve the formula P=a+b+c for a.

Solution

We will isolate aa on one side of the equation.	$P = a + b + c$
Both b and c are added to a, so we subtract them from both sides of the equation.	$P - b - c = a + b + c - b - c$
Simplify.	$P - b - c = a$ $a = P - b - c$

Try It 3.1.20

Solve the formula P=a+b+c for *b*.

Answer

b=P−a−c

Try It 3.1.21

Solve the formula P=a+b+c for *c*.

Answer

c=P−a−b

Example 3.1.22

Solve the formula 6x+5y=13 for *y*.

Solution

	$6x + 5y = 13$
Subtract 6x from both sides to isolate the term with y.	$6x - 6x + 5y = 13 - 6x$
Simplify.	$5y = 13 - 6x$
Divide by 5 to make the coefficient 1.	$\frac{5y}{5} = \frac{13-6x}{5}$
Simplify.	$y = \frac{13-6x}{5}$

The fraction is simplified. We cannot divide 13−6x by 5.

Try It 3.1.23

Solve the formula 4x+7y=9 for *y*.

Answer

$y = \frac{9-4x}{7}$

Try It 3.1.24

Solve the formula 5x+8y=1 for *y*.

Answer

$y = \frac{1-5x}{8}$

Key Concepts

- **To Solve an Application (with a formula)**
  1. **Read** the problem. Make sure all the words and ideas are understood.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose a variable to represent that quantity.
  4. **Translate** into an equation. Write the appropriate formula for the situation. Substitute in the given information.
  5. **Solve** the equation using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.
- **Distance, Rate and Time**
  For an object moving at a uniform (constant) rate, the distance traveled, the elapsed time, and the rate are related by the formula: d=rt where *d* = distance, *r* = rate, *t* = time.
- **To solve a formula for a specific variable** means to get that variable by itself with a coefficient of 1 on one side of the equation and all other variables and constants on the other side.

This page titled 3.1: Solve a Formula for a Specific Variable is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **2.6: Solve a Formula for a Specific Variable** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.1E: Exercises

Practice Makes Perfect

Use the Distance, Rate, and Time Formula

In the following exercises, solve.

Exercise 3.1E.1

Steve drove for $8\frac{1}{2}$ hours at 72 miles per hour. How much distance did he travel?

Exercise 3.1E.2

Socorro drove for $4\frac{5}{6}$ hours at 60 miles per hour. How much distance did she travel?

Answer

290 miles

Exercise 3.1E.3

Yuki walked for $1\frac{3}{4}$ hours at 4 miles per hour. How far did she walk?

Exercise 3.1E.4

Francie rode her bike for $2\frac{1}{2}$ hours at 12 miles per hour. How far did she ride?

Answer

30 miles

Exercise 3.1E.5

Connor wants to drive from Tucson to the Grand Canyon, a distance of 338 miles. If he drives at a steady rate of 52 miles per hour, how many hours will the trip take?

Exercise 3.1E.6

Megan is taking the bus from New York City to Montreal. The distance is 380 miles and the bus travels at a steady rate of 76 miles per hour. How long will the bus ride be?

Answer

5 hours

Exercise 3.1E.7

Aurelia is driving from Miami to Orlando at a rate of 65 miles per hour. The distance is 235 miles. To the nearest tenth of an hour, how long will the trip take?

Exercise 3.1E.8

Kareem wants to ride his bike from St. Louis to Champaign, Illinois. The distance is 180 miles. If he rides at a steady rate of 16 miles per hour, how many hours will the trip take?

Answer

11.25 hours

Exercise 3.1E. 9

Javier is driving to Bangor, 240 miles away. If he needs to be in Bangor in 4 hours, at what rate does he need to drive?

Exercise 3.1E. 10

Alejandra is driving to Cincinnati, 450 miles away. If she wants to be there in 6 hours, at what rate does she need to drive?

Answer

75 mph

Exercise 3.1E. 11

Aisha took the train from Spokane to Seattle. The distance is 280 miles and the trip took 3.5 hours. What was the speed of the train?

Exercise 3.1E. 12

Philip got a ride with a friend from Denver to Las Vegas, a distance of 750 miles. If the trip took 10 hours, how fast was the friend driving?

Answer

75 mph

Solve a Formula for a Specific Variable

In the following exercises, use the formula $d = rt$.

Exercise 3.1E. 13

Solve for t

1. when $d = 350$ and $r = 70$
2. in general

Exercise 3.1E. 14

Solve for t

1. when $d = 240$ and $r = 60$
2. in general

Answer

1. $t = 4$
2. $t = \frac{d}{r}$

Exercise 3.1E. 15

Solve for t

1. when $d = 510$ and $r = 60$

2. in general

Exercise 3.1E.16

Solve for t

1. when $d = 175$ and $r = 50$
2. in general

Answer

1. $t = 3.5$
2. $t = \frac{d}{r}$

Exercise 3.1E.17

Solve for r

1. when $d = 204$ and $t = 3$
2. in general

Exercise 3.1E.18

Solve for r

1. when $d = 420$ and $t = 6$
2. in general

Answer

1. $r = 70$
2. $r = \frac{d}{t}$

Exercise 3.1E.19

Solve for r

1. when $d = 160$ and $t = 2.5$
2. in general

Exercise 3.1E.20

Solve for r

1. when $d = 180$ and $t = 4.5$
2. in general

Answer

1. $r = 40$
2. $r = \frac{d}{t}$

In the following exercises, use the formula $A = \frac{1}{2}bh$

Exercise 3.1E.21

Solve for b

1. when $A = 126$ and $h = 18$
2. in general

Exercise 3.1E.22

Solve for h

1. when $A = 176$ and $b = 22$
2. in general

Answer

1. $h = 16$
2. $h = \frac{2A}{b}$

Exercise 3.1E.23

Solve for h

1. when $A = 375$ and $b = 25$
2. in general

Exercise 3.1E.24

Solve for b

1. when $A = 65$ and $h = 13$
2. in general

Answer

1. $b = 10$
2. $b = \frac{2A}{h}$

In the following exercises, use the formula $I = Prt$.

Exercise 3.1E.25

Solve for the principal, P for

1. $I = \$5,480$, $r = 4\%$, $t = 7$ years
2. in general

Exercise 3.1E.26

Solve for the principal, P for

1. $I = \$3,950$, $r = 6\%$, $t = 5$ years
2. in general

Answer

1. $P = \$13,166.67$
2. $P = \frac{I}{rt}$

Exercise 3.1E.27

Solve for the time, t for

1. $I = \$2,376$, $P = \$9,000$, $r = 4.4\%$
2. in general

Exercise 3.1E.28

Solve for the time, t for

1. $I = \$624$, $P = \$6,000$, $r = 5.2\%$
2. in general

Answer

1. $t = 2$ years
2. $t = \frac{I}{Pr}$

In the following exercises, solve.

Exercise 3.1E.29

Solve the formula $2x + 3y = 12$ for y

1. when $x = 3$
2. in general

Exercise 3.1E.30

Solve the formula $5x + 2y = 10$ for y

1. when $x = 4$
2. in general

Answer

1. $y = -5$
2. $y = \frac{10-5x}{2}$

Exercise 3.1E.31

Solve the formula $3x - y = 7$ for y

1. when $x = -2$
2. in general

Exercise 3.1E.32

Solve the formula $4x + y = 5$ for y

1. when $x = -3$
2. in general

Answer

1. $y = 17$
2. $y = 5 - 4x$

Exercise 3.1E.33

Solve $a+b=90$ for b.

Exercise 3.1E.34

Solve $a+b=90$ for a

Answer

$a=90-b$

Exercise 3.1E.35

Solve $180=a+b+c$ for a

Exercise 3.1E.36

Solve $180=a+b+c$ for c

Answer

$c=180-a-b$

Exercise 3.1E.37

Solve the formula $8x+y=15$ for y

Exercise 3.1E.38

Solve the formula $9x+y=13$ for y

Answer

$y=13-9x$

Exercise 3.1E.39

Solve the formula $-4x+y=-6$ for y

Exercise 3.1E.40

Solve the formula $-5x+y=-1$ for y

Answer

$y=-1+5x$

Exercise 3.1E.41

Solve the formula $4x+3y=7$ for y

Exercise 3.1E.42

Solve the formula $3x + 2y = 11$ for y

Answer

$y = \frac{11-3x}{2}$

Exercise 3.1E.43

Solve the formula $x - y = -4$ for y

Exercise 3.1E.44

Solve the formula $x - y = -3$ for y

Answer

$y = 3 + x$

Exercise 3.1E.45

Solve the formula $P = 2L + 2W$ for L

Exercise 3.1E.46

Solve the formula $P = 2L + 2W$ for W

Answer

$W = \frac{P-2L}{2}$

Exercise 3.1E.47

Solve the formula $C = \pi d$ for d

Exercise 3.1E.48

Solve the formula $C = \pi d$ for π

Answer

$\pi = \frac{C}{d}$

Exercise 3.1E.49

Solve the formula $V = LWH$ for L

Exercise 3.1E.50

Solve the formula $V = LWH$ for H

Answer

$H = \frac{V}{LW}$

Everyday Math

Exercise 3.1E.51

Converting temperature While on a tour in Greece, Tatyana saw that the temperature was 40° Celsius. Solve for F in the formula $C = \frac{5}{9}(F - 32)$ to find the Fahrenheit temperature.

Exercise 3.1E.52

Converting temperature Yon was visiting the United States and he saw that the temperature in Seattle one day was 50ºFahrenheit. Solve for C in the formula $F = \frac{9}{5}C + 32$ to find the Celsius temperature.

Answer

10°C

Writing Exercises

Exercise 3.1E.53

Solve the equation $2x + 3y = 6$ for y

1. when $x = -3$
2. in general
3. Which solution is easier for you, 1 or 2? Why?

Exercise 3.1E.54

Solve the equation $5x - 2y = 10$ for x

1. when $y = 10$
2. in general
3. Which solution is easier for you, 1 or 2? Why?

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
use the distance, rate, and time formula.			
solve a formula for a specific variable.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 3.1E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **2.6E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.2: Use a Problem-Solving Strategy

Learning Objectives

By the end of this section, you will be able to:

- Approach word problems with a positive attitude
- Use a problem-solving strategy for word problems
- Solve number problems

Note

Before you get started, take this readiness quiz.

1. Translate "6 less than twice x" into an algebraic expression.
   If you missed this problem, review Exercise 1.3.43.
2. Solve: $\frac{2}{3}x = 24$.
   If you missed this problem, review Exercise 2.2.10.
3. Solve: $3x + 8 = 14$.
   If you missed this problem, review Exercise 2.3.1.

Approach Word Problems with a Positive Attitude

"If you think you can… or think you can't… you're right."—Henry Ford

The world is full of word problems! Will my income qualify me to rent that apartment? How much punch do I need to make for the party? What size diamond can I afford to buy my girlfriend? Should I fly or drive to my family reunion? How much money do I need to fill the car with gas? How much tip should I leave at a restaurant? How many socks should I pack for vacation? What size turkey do I need to buy for Thanksgiving dinner, and then what time do I need to put it in the oven? If my sister and I buy our mother a present, how much does each of us pay?

Now that we can solve equations, we are ready to apply our new skills to word problems. Do you know anyone who has had negative experiences in the past with word problems? Have you ever had thoughts like the student below (Figure 3.2.1)?




Figure 3.2.1: Negative thoughts can be barriers to success.

When we feel we have no control, and continue repeating negative thoughts, we set up barriers to success. We need to calm our fears and change our negative feelings.

Start with a fresh slate and begin to think positive thoughts. If we take control and believe we can be successful, we will be able to master word problems! Read the positive thoughts in Figure 3.2.2 and say them out loud.



Figure 3.2.2: Thinking positive thoughts is a first step towards success.

Think of something, outside of school, that you can do now but couldn't do 3 years ago. Is it driving a car? Snowboarding? Cooking a gourmet meal? Speaking a new language? Your past experiences with word problems happened when you were younger—now you're older and ready to succeed!

Use a Problem-Solving Strategy for Word Problems

We have reviewed translating English phrases into algebraic expressions, using some basic mathematical vocabulary and symbols. We have also translated English sentences into algebraic equations and solved some word problems. The word problems applied math to everyday situations. We restated the situation in one sentence, assigned a variable, and then wrote an equation to solve the problem. This method works as long as the situation is familiar and the math is not too complicated.

Now, we'll expand our strategy so we can use it to successfully solve any word problem. We'll list the strategy here, and then we'll use it to solve some problems. We summarize below an effective strategy for problem solving.

USE A PROBLEM-SOLVING STRATEGY TO SOLVE WORD PROBLEMS.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information. Then, translate the English sentence into an algebraic equation.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Example 3.2.1

Pilar bought a purse on sale for $18, which is one-half of the original price. What was the original price of the purse?

Solution

Step 1. Read the problem. Read the problem two or more times if necessary. Look up any unfamiliar words in a dictionary or on the internet.

In this problem, is it clear what is being discussed? Is every word familiar?

Let p = the original price of the purse.

Step 2. Identify what you are looking for. Did you ever go into your bedroom to get something and then forget what you were looking for? It's hard to find something if you are not sure what it is! Read the problem again and look for words that tell you what you are looking for!

In this problem, the words "what was the original price of the purse" tell us what we need to find.

Step 3. Name what we are looking for. Choose a variable to represent that quantity. We can use any letter for the variable, but choose one that makes it easy to remember what it represents.

Step 4. Translate into an equation. It may be helpful to restate the problem in one sentence with all the important information. Translate the English sentence into an algebraic equation.

Reread the problem carefully to see how the given information is related. Often, there is one sentence that gives this information, or it may help to write one sentence with all the important information. Look for clue words to help translate the sentence into algebra. Translate the sentence into an equation.

Restate the problem in one sentence with all the important information.	**18** **is** **one-half the original price.**
Translate into an equation.	$18 = \frac{1}{2} \cdot p$

Step 5. Solve the equation using good algebraic techniques. Even if you know the solution right away, using good algebraic techniques here will better prepare you to solve problems that do not have obvious answers.

Solve the equation.	$18 = \frac{1}{2}p$
Multiply both sides by 2.	$2 \cdot 18 = 2 \cdot \frac{1}{2}p$
Simplify.	$36 = p$

Step 6. Check the answer in the problem to make sure it makes sense. We solved the equation and found that $p = 36$,which means "the original price" was $\$36$.

Does $36 make sense in the problem? Yes, because 18 is one-half of 36, and the purse was on sale at half the original price.

If this were a homework exercise, our work might look like this:

Pilar bought a purse on sale for $\$18$, which is one-half the original price. What was the original price of the purse?

Step 7. Answer the question with a complete sentence. The problem asked "What was the original price of the purse?"

The answer to the question is: "The original price of the purse was $36."

	Let $p =$ the original price.
	18 is one-half the original price.
	$18 = \frac{1}{2}p$
Multiply both sides by 2.	$2 \cdot 18 = 2 \cdot \frac{1}{2}p$
Simplify.	$36 = p$
Check. Is $36 a reasonable price for a purse?	
Yes.	
Is 18 one half of 36?	
$18 \stackrel{?}{=} \frac{1}{2} \cdot 36$	
$18 = 18$✓	
	The original price of the purse was $36.

Try It 3.2.2

Joaquin bought a bookcase on sale for $120, which was two-thirds of the original price. What was the original price of the bookcase?

Answer

$180

Try It 3.2.3

Two-fifths of the songs in Mariel's playlist are country. If there are 16 country songs, what is the total number of songs in the playlist?

Answer

40

Let's try this approach with another example.

Example 3.2.4

Ginny and her classmates formed a study group. The number of girls in the study group was three more than twice the number of boys. There were 11 girls in the study group. How many boys were in the study group?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	How many boys were in the study group?
Step 3. Name. Choose a variable to represent the number of boys.	Let $n =$ the number of boys.
Step 4. Translate. Restate the problem in one sentence with all the important information.	**The number of girls** (11) **was** **three more than twice the number of boys**
Translate into an equation.	$11 \quad = \quad 2b+3$
Step 5. Solve the equation.	$11 = 2b+3$
Subtract 3 from each side.	$11-3 = 2b+3-3$
Simplify.	$8 = 2b$
Divide each side by 2.	$\frac{8}{2} = \frac{2b}{2}$
Simplify.	$4 = b$
Step 6. Check. First, is our answer reasonable? Yes, having 4 boys in a study group seems OK. The problem says the number of girls was 3 more than twice the number of boys. If there are four boys, does that make eleven girls? Twice 4 boys is 8. Three more than 8 is 11.	
Step 7. Answer the question.	There were 4 boys in the study group.

Try It 3.2.5

Guillermo bought textbooks and notebooks at the bookstore. The number of textbooks was 3 more than twice the number of notebooks. He bought 7 textbooks. How many notebooks did he buy?

Answer

2

Try It 3.2.6

Gerry worked Sudoku puzzles and crossword puzzles this week. The number of Sudoku puzzles he completed is eight more than twice the number of crossword puzzles. He completed 22 Sudoku puzzles. How many crossword puzzles did he do?

Answer

7

Solve Number Problems

Now that we have a problem solving strategy, we will use it on several different types of word problems. The first type we will work on is “number problems.” Number problems give some clues about one or more numbers. We use these clues to write an equation. Number problems don’t usually arise on an everyday basis, but they provide a good introduction to practicing the problem solving strategy outlined above.

Example 3.2.7

The difference of a number and six is 13. Find the number.

Solution

Step 1. Read the problem. Are all the words familiar?	
Step 2. Identify what we are looking for.	the number
Step 3. Name. Choose a variable to represent the number.	Let $n =$ the number.
Step 4. Translate. Remember to look for clue words like "difference... of... and..."	
Restate the problem as one sentence.	$\underbrace{\textbf{The difference of the number and 6}}$ $\underbrace{\textbf{is}}$ $\underbrace{\textbf{13}}$
Translate into an equation.	$n-6 \quad = \quad 13$
Step 5. Solve the equation.	$n-6=13$
Simplify.	$n=19$
Step 6. Check.	
The difference of 19 and 6 is 13. It checks!	
Step 7. Answer the question.	The number is 19.

Try It 3.2.8

The difference of a number and eight is 17. Find the number.

Answer

25

Try It 3.2.9

The difference of a number and eleven is -7. Find the number.

Answer

4

Example 3.2.10

The sum of twice a number and seven is 15. Find the number.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the number
Step 3. Name. Choose a variable to represent the number.	Let $n =$ the number.
Step 4. Translate.	
Restate the problem as one sentence.	The sum of twice a number and 7 is 15
Translate into an equation.	$2n + 7 = 15$
Step 5. Solve the equation.	$2n + 7 = 15$
Subtract 7 from each side and simplify.	$2n = 8$
Divide each side by 2 and simplify.	$n = 4$
Step 6. Check.	
Is the sum of twice 4 and 7 equal to 15?	
$2 \cdot 4 + 7 \stackrel{?}{=} 15$ $15 = 15\checkmark$	
Step 7. Answer the question.	The number is 4.

Did you notice that we left out some of the steps as we solved this equation? If you're not yet ready to leave out these steps, write down as many as you need.

Try It 3.2.11

The sum of four times a number and two is 14. Find the number.

Answer

3

Try It 3.2.12

The sum of three times a number and seven is 25. Find the number.

Answer

6

Some number word problems ask us to find two or more numbers. It may be tempting to name them all with different variables, but so far we have only solved equations with one variable. In order to avoid using more than one variable, we will define the numbers in terms of the same variable. Be sure to read the problem carefully to discover how all the numbers relate to each other.

Example 3.2.13

One number is five more than another. The sum of the numbers is 21. Find the numbers.

Solution

Step 1. Read the problem.		
Step 2. Identify what we are looking for.		We are looking for two numbers.
Step 3. Name. We have two numbers to name and need a name for each.		
Choose a variable to represent the first number.		Let $n = 1^{st}$ number.
What do we know about the second number?		One number is five more than another.
		$n + 5 = 2^{nd}$ number
Step 4. Translate. Restate the problem as one sentence with all the important information.		The sum of the 1st number and the 2nd number is 21.
Translate into an equation.		$1^{st}\text{ number} + 2^{nd}\text{ number} = 21$
Substitute the variable expressions.		$n + n + 5 = 21$
Step 5. Solve the equation.		$n + n + 5 = 21$
Combine like terms.		$2n + 5 = 21$
Subtract 5 from both sides and simplify.		$2n = 16$
Divide by 2 and simplify.		$n = 8$ 1st number
Find the second number, too.		$n + 5$ 2nd number
		$8 + 5$
		13
Step 6. Check.		
Do these numbers check in the problem?		
Is one number 5 more than the other?	$13 \stackrel{?}{=} 8 + 5$	
Is thirteen 5 more than 8? Yes.	$13 = 13$✓	
Is the sum of the two numbers 21?	$8 + 13 \stackrel{?}{=} 21$	
	$21 = 21$✓	
Step 7. Answer the question.		The numbers are 8 and 13.

Try It 3.2.14

One number is six more than another. The sum of the numbers is twenty-four. Find the numbers.

Answer

9, 15

Try It 3.2.15

The sum of two numbers is fifty-eight. One number is four more than the other. Find the numbers.

Answer

27, 31

Example 3.2.16

The sum of two numbers is negative fourteen. One number is four less than the other. Find the numbers.

Solution

Step 1. Read the problem.		
Step 2. Identify what we are looking for.		We are looking for two numbers.
Step 3. Name.		
Choose a variable.		Let $n = 1^{st}$ number.
One number is 4 less than the other.		$n - 4 = 2^{nd}$ number
Step 4. Translate.		
Write as one sentence.		The sum of the 2 numbers is negative 14.
Translate into an equation.		1st number + 2nd number is negative fourteen
Step 5. Solve the equation.		$n + n - 4 = -14$
Combine like terms.		$n + n - 4 = -14$
Add 4 to each side and simplify.		$2n - 4 = -14$
Simplify.		$2n = -10$
		$n = -5$ 1st number
		$n - 4$ 2nd number
		$-5 - 4$
		-9
Step 6. Check.		
Is -9 four less than -5?	$-5 - 4 \stackrel{?}{=} -9$	
	$-9 = -9$✓	
Is their sum -14?	$-5 + (-9) \stackrel{?}{=} -14$	
	$-14 = -14$✓	
Step 7. Answer the question.		The numbers are -5 and -9.

Try It 3.2.17

The sum of two numbers is negative twenty-three. One number is seven less than the other. Find the numbers.

Answer

-15, -8

Try It 3.2.18

The sum of two numbers is -18. One number is 40 more than the other. Find the numbers.

Answer

-29, 11

Example 3.2.19

One number is ten more than twice another. Their sum is one. Find the numbers.

Solution

Step 1. Read the problem.		
Step 2. Identify what you are looking for.		We are looking for two numbers.
Step 3. Name.		
Choose a variable.		Let $x = 1^{st}$ number.
One number is 10 more than twice another.		$2x + 10 = 2^{nd}$ number
Step 4. Translate.		
Restate as one sentence.		Their sum is one.
		The sum of the two numbers is 1.
Translate into an equation.		$x + 2x + 10 = 1$
Step 5. Solve the equation.		
Combine like terms.		$x + 2x + 10 = 1$
Subtract 10 from each side.		$3x + 10 = 1$
Divide each side by 3.		$3x = -9$
		$x = -3$ 1st number
		$2x + 10$ 2nd number
		$2(-3) + 10$
		4
Step 6. Check.		
Is ten more than twice −3 equal to 4?	$2(-3) + 10 \stackrel{?}{=} 4$	
	$-6 + 10 \stackrel{?}{=} 4$	
	$4 = 4$✓	

Is their sum 1?	$-3+4\stackrel{?}{=}1$	
	$1=1\checkmark$	
Step 7. Answer the question.		The numbers are −3 and 4.

Try It 3.2.20

One number is eight more than twice another. Their sum is negative four. Find the numbers.

Answer

$-4,\ 0$

Try It 3.2.21

One number is three more than three times another. Their sum is -5. Find the numbers.

Answer

$-3,\ -2$

Some number problems involve consecutive integers. *Consecutive integers* are integers that immediately follow each other. Examples of consecutive integers are:

$$\begin{aligned} &1, 2, 3, 4 \\ &-10, -9, -8, -7 \\ &150, 151, 152, 153 \end{aligned} \tag{3.2.1}$$

Notice that each number is one more than the number preceding it. So if we define the first integer as n, the next consecutive integer is $n+1$. The one after that is one more than $n+1$, so it is $n+1+1$, which is $n+2$.

$$\begin{aligned} &n && 1^{\text{st}} \text{ integer} \\ &n+1 && 2^{\text{nd}} \text{ consecutive integer} \\ &n+2 && 3^{\text{rd}} \text{ consecutive integer } \ldots \text{ etc.} \end{aligned} \tag{3.2.2}$$

Example 3.2.22

The sum of two consecutive integers is 47. Find the numbers.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	two consecutive integers
Step 3. Name each number.	Let $n = 1^{st}$ integer.
	$n+1 =$ next consecutive integer
Step 4. Translate.	
Restate as one sentence.	The sum of the integers is 47.
Translate into an equation.	$n + n + 1 = 47$
Step 5. Solve the equation.	$n + n + 1 = 47$
Combine like terms.	$2n + 1 = 47$
Subtract 1 from each side.	$2n = 46$

Divide each side by 2.	$n = 23$ 1st integer
	$n + 1$ next consecutive integer
	$23 + 1$
	24
Step 6. Check.	
$23 + 24 \stackrel{?}{=} 47$ $47 = 47$✓	
Step 7. Answer the question.	The two consecutive integers are 23 and 24.

Try It 3.2.23

The sum of two consecutive integers is 95. Find the numbers.

Answer

47, 48

Try It 3.2.24

The sum of two consecutive integers is −31. Find the numbers.

Answer

-16, -15

Example 3.2.25

Find three consecutive integers whose sum is −42.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	three consecutive integers
Step 3. Name each of the three numbers.	Let $n = 1^{st}$ integer.
	$n + 1 = 2^{nd}$ consecutive integer
	$n + 2 = 3^{rd}$ consecutive integer
Step 4. Translate.	
Restate as one sentence.	The sum of the three integers is -42.
Translate into an equation.	$n + n + 1 + n + 2 = -42$
Step 5. Solve the equation.	$n + n + 1 + n + 2 = -42$
Combine like terms.	$3n + 3 = -42$
Subtract 3 from each side.	$3n = -45$
Divide each side by 3.	$n = -15$ 1st integer
	$n + 1$ 2nd integer

	$-15+1$
	-14
	$n+2$ 3rd integer
	$-15+2$
	-13
Step 6. Check.	
$-13+(-14)+(-15) \stackrel{?}{=} -42$ $-42 = -42\checkmark$	
Step 7. Answer the question.	The three consecutive integers are −13, −14, and −15.

Try It 3.2.26

Find three consecutive integers whose sum is −96.

Answer

-33, -32, -31

Try It 3.2.27

Find three consecutive integers whose sum is −36.

Answer

-13, -12, -11

Now that we have worked with consecutive integers, we will expand our work to include consecutive even integers and consecutive odd integers. *Consecutive even integers* are even integers that immediately follow one another. Examples of consecutive even integers are:

$$\begin{aligned} &18, 20, 22 \\ &64, 66, 68 \\ &-12, -10, -8 \end{aligned} \tag{3.2.3}$$

Notice each integer is 2 more than the number preceding it. If we call the first one n, then the next one is $n+2$. The next one would be $n+2+2$ or $n+4$.

$$\begin{aligned} n &\quad 1^{\text{st}} \text{ even integer} \\ n+2 &\quad 2^{\text{nd}} \text{ consecutive even integer} \\ n+4 &\quad 3^{\text{rd}} \text{ consecutive even integer } \ldots \text{ etc.} \end{aligned} \tag{3.2.4}$$

Consecutive odd integers are odd integers that immediately follow one another. Consider the consecutive odd integers 77, 79, and 81.

$$\begin{aligned} &77, 79, 81 \\ &n, n+2, n+4 \end{aligned} \tag{3.2.5}$$

$$\begin{aligned} n &\quad 1^{\text{st}} \text{ odd integer} \\ n+2 &\quad 2^{\text{nd}} \text{ consecutive odd integer} \\ n+4 &\quad 3^{\text{rd}} \text{ consecutive odd integer } \ldots \text{ etc.} \end{aligned} \tag{3.2.6}$$

Does it seem strange to add 2 (an even number) to get from one odd integer to the next? Do you get an odd number or an even number when we add 2 to 3? to 11? to 47?

Whether the problem asks for consecutive even numbers or odd numbers, you don't have to do anything different. The pattern is still the same—to get from one odd or one even integer to the next, add 2.

Example 3.2.28

Find three consecutive even integers whose sum is 84.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	three consecutive even integers
Step 3. Name the integers.	Let $n = 1^{st}$ even integers.
	$n+2 = 2^{nd}$ consecutive even integer
	$n+4 = 3^{rd}$ consecutive even integer
Step 4. Translate.	
Restate as one sentence.	The sum of the three even integers is 84.
Translate into an equation.	$n+n+2+n+4=84$
Step 5. Solve the equation.	
Combine like terms.	$n+n+2+n+4=84$
Subtract 6 from each side.	$3n+6=84$
Divide each side by 3.	$3n=78$
	$n=26$ 1st integer
	$n+2$ 2nd integer
	$26+2$
	28
	$n+4$ 3rd integer
	$26+4$
	30
Step 6. Check.	
$26+28+30 \stackrel{?}{=} 84$	
$84=84$✓	
Step 7. Answer the question.	The three consecutive integers are 26, 28, and 30.

(3.2.7)

Try It 3.2.29

Find three consecutive even integers whose sum is 102.

Answer

32, 34, 36

Try It 3.2.30

Find three consecutive even integers whose sum is −24.

Answer

$-10, -8, -6$

Example 3.2.31

A married couple together earns $110,000 a year. The wife earns $16,000 less than twice what her husband earns. What does the husband earn?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	How much does the husband earn?
Step 3. Name.	
Choose a variable to represent the amount the husband earns.	Let $h =$ the amount the husband earns.
The wife earns $\$16,000$ less than twice that.	$2h - 16,000$ the amount the wife earns.
Step 4. Translate.	Together the husband and wife earn $\$110,000$.
Restate the problem in one sentence with all the important information.	The amount the husband earns plus the amount the wife earns is $110,000
Translate into an equation.	$h + 2h - 16,000 = 110,000$
Step 5. Solve the equation.	$h + 2h - 16,000 = 110,000$
Combine like terms.	$3h - 16,000 = 110,000$
Add $16,000$ to both sides and simplify.	$3h = 126,000$
Divide each side by 3.	$h = 42,000$
	$\$42,000$ amount husband earns
	$2h - 16,000$ amount wife earns
	$2(42,000) - 16,000$
	$84,000 - 16,000$
	$68,000$
Step 6. Check.	
If the wife earns $\$68,000$ and the husband earns $\$42,000$ is the total $\$110,000$? Yes!	
Step 7. Answer the question.	The husband earns $\$42,000$ a year.

Try It 3.2.32

According to the National Automobile Dealers Association, the average cost of a car in 2014 was $28,500. This was $1,500 less than 6 times the cost in 1975. What was the average cost of a car in 1975?

Answer

$5000

? Try It 3.2.33

U.S. Census data shows that the median price of new home in the United States in November 2014 was $280,900. This was $10,700 more than 14 times the price in November 1964. What was the median price of a new home in November 1964?

Answer

$19300

Key Concepts

- **Problem-Solving Strategy**
  1. **Read** the problem. Make sure all the words and ideas are understood.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose a variable to represent that quantity.
  4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information. Then, translate the English sentence into an algebra equation.
  5. **Solve** the equation using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.
- **Consecutive Integers**
  Consecutive integers are integers that immediately follow each other.

$$\begin{array}{ll} n & 1^{\text{st}} \text{ integer} \\ n+1 & 2^{\text{nd}} \text{ consecutive integer} \\ n+2 & 3^{\text{rd}} \text{ consecutive integer } \ldots \text{ etc.} \end{array} \tag{3.2.8}$$

  Consecutive even integers are even integers that immediately follow one another.

$$\begin{array}{ll} n & 1^{\text{st}} \text{ integer} \\ n+2 & 2^{\text{nd}} \text{ consecutive even integer} \\ n+4 & 3^{\text{rd}} \text{ consecutive even integer } \ldots \text{ etc.} \end{array} \tag{3.2.9}$$

  Consecutive odd integers are odd integers that immediately follow one another.

$$\begin{array}{ll} n & 1^{\text{st}} \text{ integer} \\ n+2 & 2^{\text{nd}} \text{ consecutive odd integer} \\ n+4 & 3^{\text{rd}} \text{ consecutive odd integer } \ldots \text{ etc.} \end{array} \tag{3.2.10}$$

This page titled 3.2: Use a Problem-Solving Strategy is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.1: Use a Problem-Solving Strategy** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.2E: Exercises

Practice Makes Perfect

Use the Approach Word Problems with a Positive Attitude

In the following exercises, prepare the lists described.

Exercise 3.2E.1

List five positive thoughts you can say to yourself that will help you approach word problems with a positive attitude. You may want to copy them on a sheet of paper and put it in the front of your notebook, where you can read them often.

Answer

Answers will vary

Exercise 3.2E.2

List five negative thoughts that you have said to yourself in the past that will hinder your progress on word problems. You may want to write each one on a small piece of paper and rip it up to symbolically destroy the negative thoughts.

Use a Problem-Solving Strategy for Word Problems

In the following exercises, solve using the problem solving strategy for word problems. Remember to write a complete sentence to answer each question.

Exercise 3.2E.3

Two-thirds of the children in the fourth-grade class are girls. If there are 20 girls, what is the total number of children in the class?

Answer

30

Exercise 3.2E.4

Three-fifths of the members of the school choir are women. If there are 24 women, what is the total number of choir members?

Exercise 3.2E.5

Zachary has 25 country music CDs, which is one-fifth of his CD collection. How many CDs does Zachary have?

Answer

125

Exercise 3.2E.6

One-fourth of the candies in a bag of M&M's are red. If there are 23 red candies, how many candies are in the bag?

Exercise 3.2E.7

There are 16 girls in a school club. The number of girls is four more than twice the number of boys. Find the number of boys.

Answer

6

Exercise 3.2E.8

There are 18 Cub Scouts in Pack 645. The number of scouts is three more than five times the number of adult leaders. Find the number of adult leaders.

Exercise 3.2E.9

Huong is organizing paperback and hardback books for her club's used book sale. The number of paperbacks is 12 less than three times the number of hardbacks. Huong had 162 paperbacks. How many hardback books were there?

Answer

58

Exercise 3.2E.10

Jeff is lining up children's and adult bicycles at the bike shop where he works. The number of children's bicycles is nine less than three times the number of adult bicycles. There are 42 adult bicycles. How many children's bicycles are there?

Exercise 3.2E.11

Philip pays $1,620 in rent every month. This amount is $120 more than twice what his brother Paul pays for rent. How much does Paul pay for rent?

Answer

$750

Exercise 3.2E.12

Marc just bought an SUV for $54,000. This is $7,400 less than twice what his wife paid for her car last year. How much did his wife pay for her car?

Exercise 3.2E.13

Laurie has $46,000 invested in stocks and bonds. The amount invested in stocks is $8,000 less than four times the amount invested in bonds. How much does Laurie have invested in bonds?

Answer

$13,500

Exercise 3.2E.14

Erica earned a total of $50,450 last year from her two jobs. The amount she earned from her job at the store was $1,250 more than three times the amount she earned from her job at the college. How much did she earn from her job at the college?

Solve Number Problems

In the following exercises, solve each number word problem.

Exercise 3.2E.15

The sum of a number and eight is 12. Find the number.

Answer

4

Exercise 3.2E.16

The sum of a number and nine is 17. Find the number.

Exercise 3.2E.17

The difference of a number and 12 is three. Find the number.

Answer

15

Exercise 3.2E.18

The difference of a number and eight is four. Find the number.

Exercise 3.2E.19

The sum of three times a number and eight is 23. Find the number.

Answer

5

Exercise 3.2E.20

The sum of twice a number and six is 14. Find the number.

Exercise 3.2E.21

The difference of twice a number and seven is 17. Find the number.

Answer

12

Exercise 3.2E.22

The difference of four times a number and seven is 21. Find the number.

Exercise 3.2E.23

Three times the sum of a number and nine is 12. Find the number.

Answer

-5

Exercise 3.2E.24

Six times the sum of a number and eight is 30. Find the number.

Exercise 3.2E.25

One number is six more than the other. Their sum is 42. Find the numbers.

Answer

18, 24

Exercise 3.2E.26

One number is five more than the other. Their sum is 33. Find the numbers.

Exercise 3.2E.27

The sum of two numbers is 20. One number is four less than the other. Find the numbers.

Answer

8, 12

Exercise 3.2E.28

The sum of two numbers is 27. One number is seven less than the other. Find the numbers.

Exercise 3.2E.29

The sum of two numbers is −45. One number is nine more than the other. Find the numbers.

Answer

−18,−27

Exercise 3.2E.30

The sum of two numbers is −61. One number is 35 more than the other. Find the numbers.

Exercise 3.2E.31

The sum of two numbers is −316. One number is 94 less than the other. Find the numbers.

Answer

−111,−205

Exercise 3.2E.32

The sum of two numbers is −284. One number is 62 less than the other. Find the numbers.

Exercise 3.2E.33

One number is 14 less than another. If their sum is increased by seven, the result is 85. Find the numbers.

Answer

32, 46

Exercise 3.2E.34

One number is 11 less than another. If their sum is increased by eight, the result is 71. Find the numbers.

Exercise 3.2E.35

One number is five more than another. If their sum is increased by nine, the result is 60. Find the numbers.

Answer

23, 28

Exercise 3.2E.36

One number is eight more than another. If their sum is increased by 17, the result is 95. Find the numbers.

Exercise 3.2E.37

One number is one more than twice another. Their sum is −5. Find the numbers.

Answer

−2,−3

Exercise 3.2E.38

One number is six more than five times another. Their sum is six. Find the numbers.

Exercise 3.2E.39

The sum of two numbers is 14. One number is two less than three times the other. Find the numbers.

Answer

4, 10

Exercise 3.2E.40

The sum of two numbers is zero. One number is nine less than twice the other. Find the numbers.

Exercise 3.2E.41

The sum of two consecutive integers is 77. Find the integers.

Answer

38, 39

Exercise 3.2E.42

The sum of two consecutive integers is 89. Find the integers.

Exercise 3.2E.43

The sum of two consecutive integers is −23. Find the integers.

Answer

−11,−12

Exercise 3.2E.44

The sum of two consecutive integers is −37. Find the integers.

Exercise 3.2E.45

The sum of three consecutive integers is 78. Find the integers.

Answer

25, 26, 27

Exercise 3.2E.46

The sum of three consecutive integers is 60. Find the integers.

Exercise 3.2E.47

Find three consecutive integers whose sum is −36.

Answer

−11,−12,−13

Exercise 3.2E.48

Find three consecutive integers whose sum is −3.

Exercise 3.2E.49

Find three consecutive even integers whose sum is 258.

Answer

84, 86, 88

Exercise 3.2E.50

Find three consecutive even integers whose sum is 222.

Exercise 3.2E.51

Find three consecutive odd integers whose sum is 171.

Answer

55, 57, 59

Exercise 3.2E.52

Find three consecutive odd integers whose sum is 291.

Exercise 3.2E.53

Find three consecutive even integers whose sum is −36.

Answer

−10,−12,−14

Exercise 3.2E.54

Find three consecutive even integers whose sum is −84.

Exercise 3.2E.55

Find three consecutive odd integers whose sum is −213.

Answer

−69,−71,−73

Exercise 3.2E.56

Find three consecutive odd integers whose sum is −267.

Everyday Math

Exercise 3.2E.57

Sale Price Patty paid $35 for a purse on sale for $10 off the original price. What was the original price of the purse?

Answer

$45

Exercise 3.2E.58

Sale Price Travis bought a pair of boots on sale for $25 off the original price. He paid $60 for the boots. What was the original price of the boots?

Exercise 3.2E.59

Buying in Bulk Minh spent $6.25 on five sticker books to give his nephews. Find the cost of each sticker book.

Answer

$1.25

Exercise 3.2E.60

Buying in Bulk Alicia bought a package of eight peaches for $3.20. Find the cost of each peach.

Exercise 3.2E.61

Price before Sales Tax Tom paid $1,166.40 for a new refrigerator, including $86.40 tax. What was the price of the refrigerator?

Answer

$1080

Exercise 3.2E.62

Price before Sales Tax Kenji paid $2,279 for a new living room set, including $129 tax. What was the price of the living room set?

Writing Exercises

Exercise 3.2E.63

What has been your past experience solving word problems?

Answer

answers will vary

Exercise 3.2E.64

When you start to solve a word problem, how do you decide what to let the variable represent?

Exercise 3.2E.65

What are consecutive odd integers? Name three consecutive odd integers between 50 and 60.

Answer

Consecutive odd integers are odd numbers that immediately follow each other. An example of three consecutive odd integers between 50 and 60 would be 51, 53, and 55.

Exercise 3.2E.66

What are consecutive even integers? Name three consecutive even integers between −50 and −40.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
approach word problems with a positive attitude.			
use a problem solving strategy for word problems.			
solve number problems.			

ⓑ If most of your checks were:

…confidently. Congratulations! You have achieved your goals in this section! Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific!

…with some help. This must be addressed quickly as topics you do not master become potholes in your road to success. Math is sequential—every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no—I don't get it! This is critical and you must not ignore it. You need to get help immediately or you will quickly be overwhelmed. See your instructor as soon as possible to discuss your situation. Together you can come up with a plan to get you the help you need.

Practice for fixed and variable cost problems

? Exercise $3.2E.67$

Construct an equation and use it to solve the following problems.

1. A restaurant charges $350 for use of the banquet room plus $32.50 per person. If the total cost was $3470, how many people attended the event?
2. A storage facility charges a fixed cost of $1200 a month for rent and a variable cost of $5 per box that is stored. If the total cost for a month was $1315, how many boxes were stored?
3. A car was rented for two days at $19.95 per day plus $0.10 per mile. How many miles were driven if the total bill was $89.70?
4. A car was rented for two days at $25 a day plus $0.15 per mile. How many miles were driven if the total bill was $102.50?
5. An Internet Provider charges $12.95 a month for 6 hours and $0.25 for each additional hour. How many hours were spent on-line if the Internet bill for the month was $21.20?
6. An Internet Provider charges $10 a month for 8 hours of service plus $0.50 an hour for each hour over the initial 8 hours. How many hours were spent online if the Internet bill for a month was $18.50?
7. If $100 is charged for bar-coding the first 500 pieces of mail, $0.10 is charged for each additional piece of mail, and the total bill for bar coding was $143.20, how many pieces of mail were bar coded?
8. An apartment costs $755 a month for rent, plus $3.60 an hour for cable usage. If the total cost in a month was $826.20, how much time spent using the cable service?
9. A hockey player is paid $175,000 a year plus $200 per goal. How many goals were made if the salary for a year was $199,000?
10. A musician at a club is paid $175 a week plus $1.50 for each paid admission. How many paid admissions did the club have if the musician's paycheck for a week was for $340?

Answer

1. 96 people. 3. 498 miles. 5. $6+33=39$ hours. 7. $500+432=932$ pieces of mail. 9. 120 goals.

This page titled 3.2E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.1E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.3: Solve Percent Applications

Learning Objectives

By the end of this section, you will be able to:

- Translate and solve basic percent equations
- Solve percent applications
- Find percent increase and percent decrease
- Solve simple interest applications
- Solve applications with discount or mark-up

Be prepared

Before you get started, take this readiness quiz.

1. Convert 4.5% to a decimal.
   If you missed this problem, review [link].
2. Convert 0.6 to a percent.
   If you missed this problem, review [link].
3. Round 0.875 to the nearest hundredth.
   If you missed this problem, review [link].
4. Multiply (4.5)(2.38).
   If you missed this problem, review [link].
5. Solve 3.5=0.7n.
   If you missed this problem, review [link].
6. Subtract 50−37.45.
   If you missed this problem, review [link].

Translate and Solve Basic Percent Equations

We will solve percent equations using the methods we used to solve equations with fractions or decimals. Without the tools of algebra, the best method available to solve percent problems was by setting them up as proportions. Now as an algebra student, you can just translate English sentences into algebraic equations and then solve the equations.

We can use any letter you like as a variable, but it is a good idea to choose a letter that will remind us of what you are looking for. We must be sure to change the given percent to a decimal when we put it in the equation.

Example 3.3.1

Translate and solve: What number is 35% of 90?

Solution

	What number is 35% of 90?
Translate into algebra. Let n= the number.	$n = 0.35 \cdot 90$
Remember "of" means multiply, "is" means equals.	
Multiply.	$n = 31.5$
	31.5 is 35% of 90

Try It 3.3.2

Translate and solve:

What number is 45% of 80?

Answer

36

Try It 3.3.3

Translate and solve:

What number is 55% of 60?

Answer

33

We must be very careful when we translate the words in the next example. The unknown quantity will not be isolated at first, like it was in Example. We will again use direct translation to write the equation.

Example 3.3.4

Translate and solve: 6.5% of what number is $1.17?

Solution

	6.5% of what number is $1.17?
Translate. Let n= the number.	$0.065 \cdot n = 1.17$
Multiply.	$0.065n = 1.17$
Divide both sides by 0.065 and simplify.	$n = 18$
	6.5% of $18 is $1.17

Try It 3.3.5

Translate and solve:

7.5% of what number is $1.95?

Answer

$26

Try It 3.3.6

Translate and solve:

8.5% of what number is $3.06?

Answer

$36

In the next example, we are looking for the percent.

Example 3.3.7

Translate and solve: 144 is what percent of 96?

Solution

	144 is what percent of 96?
Translate into algebra. Let p= the percent.	$144 = p \cdot 96$
Multiply.	$144 = 96p$
Divide by 96 and simplify.	$1.5 = p$
Convert to percent.	$150\% = p$
	144 is 150% of 96

Note that we are asked to find percent, so we must have our final result in percent form.

Try It 3.3.8

Translate and solve:

110 is what percent of 88?

Answer

125%

Try It 3.3.9

Translate and solve:

126 is what percent of 72?

Answer

175%

Solve Applications of Percent

Many applications of **percent**—such as tips, sales tax, discounts, and interest—occur in our daily lives. To solve these applications we'll translate to a basic percent equation, just like those we solved in previous examples. Once we translate the sentence into a percent equation, we know how to solve it.

We will restate the problem solving strategy we used earlier for easy reference.

USE A PROBLEM-SOLVING STRATEGY TO SOLVE AN APPLICATION.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information. Then, translate the English sentence into an algebraic equation.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Now that we have the strategy to refer to, and have practiced solving basic percent equations, we are ready to solve percent applications. Be sure to ask yourself if your final answer makes sense—since many of the applications will involve everyday situations, you can rely on your own experience.

Example 3.3.10

Dezohn and his girlfriend enjoyed a nice dinner at a restaurant and his bill was $68.50. He wants to leave an 18% tip. If the tip will be 18% of the total bill, how much tip should he leave?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the amount of tip should Dezohn leave
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let t = amount of tip.
Step 4. Translate into an equation.	The tip is 18% of the total bill.
Write a sentence that gives the information to find it.	The tip is 18% of $68.50
Translate the sentence into an equation.	$t = 0.18 \cdot 68.50$
Step 5. Solve the equation. Multiply.	$t = 12.33$
Step 6. Check. Does this make sense?	
Yes, 20% is close to 18%, and 20% of $70 is $14.	
Step 7. Answer the question with a complete sentence.	Dezohn should leave a tip of $12.33.

Notice that we used t to represent the unknown tip.

Try It 3.3.11

Cierra and her sister enjoyed a dinner in a restaurant and the bill was $81.50. If she wants to leave 18% of the total bill as her tip, how much should she leave?

Answer

$14.67

Try It 3.3.12

Kimngoc had lunch at her favorite restaurant. She wants to leave 15% of the total bill as her tip. If her bill was $14.40, how much will she leave for the tip?

Answer

$2.16

Example 3.3.13

The label on Masao's breakfast cereal said that one serving of cereal provides 85 milligrams (mg) of potassium, which is 2% of the recommended daily amount. What is the total recommended daily amount of potassium?

Solution

Step 1. Read the problem.	

Step 2. Identify what we are looking for.	the total amount of potassium that is recommended
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let a= total amount of potassium.
Step 4. Translate. Write a sentence that gives the information to find it.	85 mg is 2% of the total amount
Translate into an equation.	$85 = 0.02 \cdot a$
Step 5. Solve the equation.	$4{,}250 = a$
Step 6. Check. Does this make sense?	
Yes, 2% is a small percent and 85 is a small part of 4,250.	
Step 7. Answer the question with a complete sentence.	The amount of potassium that is recommended is 4,250 mg.

Try It 3.3.14

One serving of wheat square cereal has seven grams of fiber, which is 28% of the recommended daily amount. What is the total recommended daily amount of fiber?

Answer

25 grams

Try It 3.3.15

One serving of rice cereal has 190 mg of sodium, which is 8% of the recommended daily amount. What is the total recommended daily amount of sodium?

Answer

2,375 mg

Example 3.3.16

Mitzi received some gourmet brownies as a gift. The wrapper said each brownie was 480 calories, and had 240 calories of fat. What percent of the total calories in each brownie comes from fat?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the percent of the total calories from fat
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let p= percent of fat.
Step 4. Translate. Write a sentence that gives the information to find it.	What percent of 480 is 240?
Translate into an equation.	$p \cdot 480 = 240$
Step 5. Solve the equation.	$480\,p = 240$
Divide by 480.	$p = 0.5$
Put in a percent form.	$p = 50\%$

Step 6. Check. Does this make sense?	
Yes, 240 is half of 480, so 50% makes sense.	
Step 7. Answer the question with a complete sentence.	Of the total calories in each brownie, 50% is fat.

Try It 3.3.17

Solve. Round to the nearest whole percent.

Veronica is planning to make muffins from a mix. The package says each muffin will be 230 calories and 60 calories will be from fat. What percent of the total calories is from fat?

Answer

26%

Try It 3.3.18

Solve. Round to the nearest whole percent.

The mix Ricardo plans to use to make brownies says that each brownie will be 190 calories, and 76 calories are from fat. What percent of the total calories are from fat?

Answer

40%

Find Percent Increase and Percent Decrease

People in the media often talk about how much an amount has increased or decreased over a certain period of time. They usually express this increase or decrease as a percent.

To find the **percent increase**, first we find the amount of increase, the difference of the new amount and the original amount. Then we find what percent the amount of increase is of the original amount.

FIND THE PERCENT INCREASE.

1. Find the amount of increase.

   $$\text{new amount} - \text{original amount} = \text{increase}$$
2. Find the percent increase.

   The increase is what percent of the original amount?

Example 3.3.19

In 2011, the California governor proposed raising community college fees from $26 a unit to $36 a unit. Find the percent increase. (Round to the nearest tenth of a percent.)

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the percent increase
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let p = the percent.

Step 4. Translate. Write a sentence that gives the information to find it.	
First find the amount of increase.	new amount − original amount = increase
	$36-26=10$
Find the percent.	Increase is what percent of the original amount?
	10 is what percent of 26?
Translate into an equation.	$10 = p \cdot 26$
Step 5. Solve the equation.	$10 = 26p$
Divide by 26.	$0.3846\ldots = p$
Change to percent form; round to the nearest tenth.	$38.5\% = p$
Step 6. Check. Does this make sense?	
Yes, 38.5% is close to $\frac{1}{3}$, and 10 is close to $\frac{1}{3}$ of 26.	
Step 7. Answer the question with a complete sentence.	The new fees represent a 38.5% increase over the old fees.

Try It 3.3.20

Find the percent increase. (Round to the nearest tenth of a percent.)

In 2011, the IRS increased the deductible mileage cost to 55.5 cents from 51 cents.

Answer

8.8%

Try It 3.3.21

Find the percent increase.

In 1995, the standard bus fare in Chicago was $1.50. In 2008, the standard bus fare was $2.25.

Answer

50%

Finding the **percent decrease** is very similar to finding the percent increase, but now the amount of decrease is the difference of the original amount and the new amount. Then we find what percent the amount of decrease is of the original amount.

FIND THE PERCENT DECREASE.

1. Find the amount of decrease.

   $$\text{original amount} - \text{new amount} = \text{decrease}$$
2. Find the percent decrease.

   The decrease is what percent of the original amount?

Example 3.3.22

The average price of a gallon of gas in one city in June 2014 was $3.71. The average price in that city in July was $3.64. Find the percent decrease.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the percent decrease
Step 3. Name what we are looking for.	
Choose a variable to represent that quantity.	Let p= the percent decrease.
Step 4. Translate. Write a sentence that gives the information to find it.	
First find the amount of decrease.	3.71−3.64=0.07
Find the percent.	Decrease is what percent of the original amount?
	0.07 is what percent of 3.71?
Translate into an equation.	$0.07 = p \cdot 3.71$
Step 5. Solve the equation.	$0.07 = 3.71\,p$
Divide by 3.71.	$0.0188\ldots = p$
Change to percent form; round to the nearest tenth.	$1.9\% = p$
Step 6. Check. Does this make sense?	
Yes, if the original price was $4, a 2% decrease would be 8 cents.	
Step 7. Answer the question with a complete sentence.	The price of gas decreased 1.9%.

Try It 3.3.23

Find the percent decrease. (Round to the nearest tenth of a percent.)

The population of North Dakota was about 672,000 in 2010. The population is projected to be about 630,000 in 2020.

Answer

6.3%

Try It 3.3.24

Find the percent decrease.

Last year, Sheila's salary was $42,000. Because of furlough days, this year, her salary was $37,800.

Answer

10%

Solve Simple Interest Applications

Do you know that banks pay you to keep your money? The money a customer puts in the bank is called the **principal**, P, and the money the bank pays the customer is called the **interest**. The interest is computed as a certain percent of the principal; called the

rate of interest, *r*. We usually express rate of interest as a percent per year, and we calculate it by using the decimal equivalent of the percent. The variable *t*, (for *time*) represents the number of years the money is in the account.

To find the interest we use the simple interest formula, I=Prt.

SIMPLE INTEREST

If an amount of money, *P*, called the principal, is invested for a period of *t* years at an annual interest rate *r*, the amount of interest, *I*, earned is

$$I = Prt \quad \text{where} \quad \begin{aligned} I &= \text{interest} \\ P &= \text{principle} \\ r &= \text{rate} \\ t &= \text{time} \end{aligned} \tag{3.3.1}$$

Interest earned according to this formula is called **simple interest**.

Interest may also be calculated another way, called compound interest. This type of interest will be covered in later math classes.

The formula we use to calculate simple interest is I=Prt. To use the formula, we substitute in the values the problem gives us for the variables, and then solve for the unknown variable. It may be helpful to organize the information in a chart.

Example 3.3.25

Nathaly deposited $12,500 in her bank account where it will earn 4% interest. How much interest will Nathaly earn in 5 years?

$$\begin{aligned} I &= ? \\ P &= \$12,500 \\ r &= 4\% \\ t &= 5 \text{ years} \end{aligned}$$

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for. — the amount of interest earned

Step 3. Name what we are looking for. Choose a variable to represent that quantity — Let I = the amount of interest.

Step 4. Translate into an equation.

Write the formula. — $I = Prt$

Substitute in the given information. — $I = (12500)(.04)(5)$

Step 5. Solve the equation. — $I = 2500$

Step 6. Check: Does this make sense?

Is $2,500 is a reasonable interest on $12,500? Yes.

Step 7. Answer the question with a complete sentence. — The interest is $2500

Try It 3.3.26

Areli invested a principal of $950 in her bank account with interest rate 3%. How much interest did she earn in 5 years?

Answer

$142.50

Try It 3.3.27

Susana invested a principal of $36,000 in her bank account with interest rate 6.5%. How much interest did she earn in 3 years?

Answer

$7020

There may be times when we know the amount of interest earned on a given principal over a certain length of time, but we don't know the rate. To find the rate, we use the simple interest formula, substitute in the given values for the principal and time, and then solve for the rate.

Example 3.3.28

Loren loaned his brother $3,000 to help him buy a car. In 4 years his brother paid him back the $3,000 plus $660 in interest. What was the rate of interest?

$$\begin{aligned} I &= \$660 \\ P &= \$3000 \\ r &= ? \\ t &= 4 \text{ years} \end{aligned} \tag{3.3.2}$$

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for.	the rate of interest
Step 3. Name what we are looking for. Choose a variable to represent that quantity	Let r = the rate of interest.
Step 4. Translate into an equation.	
Write the formula.	$I = Prt$
Substitute in the given information.	$660 = (3000)r(4)$
Step 5. Solve the equation.	
	$660 = (12000)r$
Divide.	$0.055 = r$
Change to percent form.	$5.5\% = r$

Step 6. Check: Does this make sense?

$I = Prt$

$660 \stackrel{?}{=} (3000)(0.055)(4)$

$660 = 660$✓

Step 7. Answer the question with a complete sentence.	The rate of interest was 5.5%

Notice that in this example, Loren's brother paid Loren interest, just like a bank would have paid interest if Loren invested his money there.

? Try It 3.3.29

Jim loaned his sister $5,000 to help her buy a house. In 3 years, she paid him the $5,000, plus $900 interest. What was the rate of interest?

Answer

6%

? Try It 3.3.30

Hang borrowed $7,500 from her parents to pay her tuition. In 5 years, she paid them $1,500 interest in addition to the $7,500 she borrowed. What was the rate of interest?

Answer

4%

✓ Example 3.3.31

Eduardo noticed that his new car loan papers stated that with a 7.5% interest rate, he would pay $6,596.25 in interest over 5 years. How much did he borrow to pay for his car?

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for. Choose a variable to represent that quantity.	the amount borrowed (the principal)
Step 3. Name what we are looking for. Choose a variable to represent that quantity	Let P = principal borrowed.
Step 4. Translate into an equation.	
Write the formula.	$I = Prt$
Substitute in the given information.	$6596.25 = P(0.075)(5)$
Step 5. Solve the equation.	
	$6596.25 = 0.375P$
Divide.	$17590 = P$

Step 6. Check: Does this make sense?

$I = Prt$

$6596.25 \stackrel{?}{=} (17590)(0.075)(5)$

$6596.25 = 6596.25$✓

Step 7. Answer the question with a complete sentence. The principal is $17590

? Try It 3.3.32

Sean's new car loan statement said he would pay $4,866.25 in interest from an interest rate of 8.5% over 5 years. How much did he borrow to buy his new car?

Answer

$11,450

? Try It 3.3.33

In 5 years, Gloria's bank account earned $2,400 interest at 5%. How much had she deposited in the account?

Answer

$9,600

Solve Applications with Discount or Mark-up

Applications of discount are very common in retail settings. When you buy an item on sale, the original price has been discounted by some dollar amount. The **discount rate**, usually given as a percent, is used to determine the amount of the discount. To determine the **amount of discount**, we multiply the discount rate by the original price.

We summarize the discount model in the box below.

DISCOUNT

$$\begin{aligned} \text{amount of discount} &= \text{discount rate} \times \text{original price} \\ \text{sale price} &= \text{original price - amount of discount} \end{aligned} \tag{3.3.3}$$

Keep in mind that the sale price should always be less than the original price.

Example 3.3.34

Elise bought a dress that was discounted 35% off of the original price of $140. What was (a) the amount of discount and (b) the sale price of the dress?

Solution

(a) Original price = $140
Discount rate = 35%
Discount = ?

Step 1. Read the problem.

Step 2. Identify what we are looking for. — the amount of discount

Step 3. Name what we are looking for.
Choose a variable to represent that quantity. — Let d = the amount of discount.

Step 4. Translate into an equation. Write a sentence that gives the information to find it.
Translate into an equation — $d = 0.35(140)$

Step 5. Solve the equation. — $d = 49$

Step 6. Check: Does this make sense?

Is a $49 discount reasonable for a $140 dress?
Yes.

Step 7. Write a complete sentence to answer the question. — The amount of discount was $49

(b) Read the problem again.

Step 1. Identify what we are looking for.	the sale price of the dress
Step 2. Name what we are looking for.	
Choose a variable to represent that quantity.	Let s= the sale price.
Step 3. Translate into an equation.	
Write a sentence that gives the information to find it.	The sale price is the $140 minus the $49 discount
Translate into an equation.	$s = 140 - 49$
Step 4. Solve the equation.	$s = 91$
Step 5. Check. Does this make sense?	
Is the sale price less than the original price?	

Yes, $91 is less than $140.	
Step 6. Answer the question with a complete sentence.	The sale price of the dress was $91.

Try It 3.3.35

Find (a) the amount of discount and (b) the sale price:

Sergio bought a belt that was discounted 40% from an original price of $29.

Answer

(a) $11.60, (b) $17.40

Try It 3.3.36

Find (a) the amount of discount and (b) the sale price:

Oscar bought a barbecue that was discounted 65% from an original price of $395.

Answer

(a) $256.75, (b) $138.25

There may be times when we know the original price and the sale price, and we want to know the **discount rate**. To find the discount rate, first we will find the amount of discount and then use it to compute the rate as a percent of the original price. Example 3.3.37 will show this case.

Example 3.3.37

Jeannette bought a swimsuit at a sale price of $13.95. The original price of the swimsuit was $31. Find the (a) amount of discount and (b) discount rate.

Solution

(a) Original price = $31
Discount rate = ?
Discount = $13.95

Step 1. Read the problem.

Step 2. Identify what we are looking for. — the amount of discount

Step 3. Name what we are looking for.
Choose a variable to represent that quantity. — Let d = the amount of discount.

Step 4. Translate into an equation. Write a sentence that gives the information to find it. — The discount is the difference between the original price and the sale price.

Translate into an equation — $d = 31 - 13.95$

Step 5. Solve the equation. — $d = 17.05$

Step 6. Check: Does this make sense?

Is 17.05 less than 31? Yes.

Step 7. Answer the question with a complete sentence. — The amount of discount was $17.05

(b) Read the problem again.

Step 1. Identify what we are looking for.	the discount rate
Step 2. Name what we are looking for.	
Choose a variable to represent it.	Let r = the discount rate.
Step 3. Translate into an equation.	
Write a sentence that gives the information to find it.	The discount of $17.05 is what percent of $31 ?
Translate into an equation.	$17.05 = r \cdot 31$
Step 4. Solve the equation.	$17.05 = 31r$
Divide both sides by 31.	$0.55 = r$
Change to percent form.	$r = 55\%$
Step 5. Check. Does this make sense?	
Is $17.05 equal to 55% of $31?	
$17.05 \stackrel{?}{=} 0.55(31)$	
$17.05 = 17.05$✓	
Step 6. Answer the question with a complete sentence.	The rate of discount was 55%.

Try It 3.3.38

Find

a. the amount of discount and
b. the discount rate.

Lena bought a kitchen table at the sale price of $375.20. The original price of the table was $560.

Answer

a. $184.80
b. 33%

Try It 3.3.39

Find

a. the amount of discount and
b. the discount rate.

Nick bought a multi-room air conditioner at a sale price of $340. The original price of the air conditioner was $400.

Answer

a. $60
b. 15%

Applications of mark-up are very common in retail settings. The price a retailer pays for an item is called the **original cost**. The retailer then adds a **mark-up** to the original cost to get the **list price**, the price he sells the item for. The mark-up is usually calculated as a percent of the original cost. To determine the amount of mark-up, multiply the mark-up rate by the original cost.

We summarize the mark-up model in the box below.

MARK-UP

$$\begin{aligned} \text{amount of mark-up} &= \text{mark-up rate} \times \text{original cost} \\ \text{list price} &= \text{original cost} + \text{amount of mark up} \end{aligned} \tag{3.3.4}$$

Keep in mind that the list price should always be more than the original cost.

Example 3.3.40

Adam's art gallery bought a photograph at original cost $250. Adam marked the price up 40%. Find the

a. amount of mark-up and
b. the list price of the photograph.

Solution

a.

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the amount of mark-up
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let m = the amount of markup.
Step 4. Translate into an equation.	
Write a sentence that gives the information to find it.	The mark-up is 40% of the $250 original cost

Translate into an equation.	$m = 0.40 \cdot 250$
Step 5. Solve the equation.	$m = 100$
Step 6. Check. Does this make sense?	
Yes, 40% is less than one-half and 100 is less than half of 250.	
Step 7. Answer the question with a complete sentence.	The mark-up on the photograph was $100.

b.

Step 1. Read the problem again.	
Step 2. Identify what we are looking for.	the list price
Step 3. Name what we are looking for.	
Choose a variable to represent it.	Let p= the list price.
Step 4. Translate into an equation.	
Write a sentence that gives the information to find it.	The list price is original cost plus the mark-up
Translate into an equation.	$p = 250 + 100$
Step 5. Solve the equation.	$p = 350$
Step 6. Check. Does this make sense?	
Is the list price more than the net price? Is $350 more than $250? Yes.	
Step 7. Answer the question with a complete sentence.	The list price of the photograph was $350.

Try It 3.3.41

Find

a. the amount of mark-up and
b. the list price.

Jim's music store bought a guitar at original cost $1,200. Jim marked the price up 50%.

Answer

a. $600
b. $1,800

Try It 3.3.42

Find

a. the amount of mark-up and
b. the list price.

The Auto Resale Store bought Pablo's Toyota for $8,500. They marked the price up 35%.

Answer

a. $2,975
b. $11,475

Key Concepts

- **Percent Increase** To find the percent increase:
  1. Find the amount of increase. increase=new amount−originalamount
  2. Find the percent increase. Increase is what percent of the original amount?
- **Percent Decrease** To find the percent decrease:
  1. Find the amount of decrease. decrease=original amount−newamount
  2. Find the percent decrease. Decrease is what percent of the original amount?
- **Simple Interest** If an amount of money, *P*, called the principal, is invested for a period of *t* years at an annual interest rate *r*, the amount of interest, *I*, earned is

$$I = Prt \quad \text{where} \quad \begin{aligned} I &= \text{interest} \\ P &= \text{principle} \\ r &= \text{rate} \\ t &= \text{time} \end{aligned} \tag{3.3.5}$$

- **Discount**
  - amount of discount is discount rate × original price
  - sale price is original price – discount
- **Mark-up**
  - amount of mark-up is mark-up rate × original cost
  - list price is original cost + mark up

Glossary

amount of discount

The amount of discount is the amount resulting when a discount rate is multiplied by the original price of an item.

discount rate

The discount rate is the percent used to determine the amount of a discount, common in retail settings.

interest

Interest is the money that a bank pays its customers for keeping their money in the bank.

list price

The list price is the price a retailer sells an item for.

mark-up

A mark-up is a percentage of the original cost used to increase the price of an item.

original cost

The original cost in a retail setting, is the price that a retailer pays for an item.

principal

The principal is the original amount of money invested or borrowed for a period of time at a specific interest rate.

rate of interest

The rate of interest is a percent of the principal, usually expressed as a percent per year.

simple interest

Simple interest is the interest earned according to the formula I=Prt.

This page titled 3.3: Solve Percent Applications is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.2: Solve Percent Applications** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.3E: Exercises

Practice Makes Perfect

Translate and Solve Basic Percent Equations

In the following exercises, translate and solve.

Exercise 3.3E.1

What number is 45% of 120?

Answer

54

Exercise 3.3E.2

What number is 65% of 100?

Exercise 3.3E.3

What number is 24% of 112?

Answer

26.88

Exercise 3.3E.4

What number is 36% of 124?

Exercise 3.3E.5

250% of 65 is what number?

Answer

162.5

Exercise 3.3E.6

150% of 90 is what number?

Exercise 3.3E.7

800% of 2250 is what number?

Answer

18,000

Exercise 3.3E.8

600% of 1740 is what number?

Exercise 3.3E.9

28 is 25% of what number?

Answer

112

Exercise 3.3E.10

36 is 25% of what number?

Exercise 3.3E.11

81 is 75% of what number?

Answer

108

Exercise 3.3E.12

93 is 75% of what number?

Exercise 3.3E.13

8.2% of what number is $2.87?

Answer

$35

Exercise 3.3E.14

6.4% of what number is $2.88?

Exercise 3.3E.15

11.5% of what number is $108.10?

Answer

$940

Exercise 3.3E.16

12.3% of what number is $92.25?

Exercise 3.3E.17

What percent of 260 is 78?

Answer

30%

Exercise 3.3E.18

What percent of 215 is 86?

Exercise 3.3E.19

What percent of 1500 is 540?

Answer

36%

Exercise 3.3E.20

What percent of 1800 is 846?

Exercise 3.3E.21

30 is what percent of 20?

Answer

150%

Exercise 3.3E.22

50 is what percent of 40?

Exercise 3.3E.23

840 is what percent of 480?

Answer

175%

Exercise 3.3E.24

790 is what percent of 395?

Solve Percent Applications

In the following exercises, solve.

Exercise 3.3E.25

Geneva treated her parents to dinner at their favorite restaurant. The bill was $74.25. Geneva wants to leave 16% of the total bill as a tip. How much should the tip be?

Answer

$11.88

Exercise 3.3E. 26

When Hiro and his co-workers had lunch at a restaurant near their work, the bill was $90.50. They want to leave 18% of the total bill as a tip. How much should the tip be?

Exercise 3.3E. 27

Trong has 12% of each paycheck automatically deposited to his savings account. His last paycheck was $2165. How much money was deposited to Trong's savings account?

Answer

$259.80

Exercise 3.3E. 28

Cherise deposits 8% of each paycheck into her retirement account. Her last paycheck was $1,485. How much did Cherise deposit into her retirement account?

Exercise 3.3E. 29

One serving of oatmeal has eight grams of fiber, which is 33% of the recommended daily amount. What is the total recommended daily amount of fiber?

Answer

24.2 g

Exercise 3.3E. 30

One serving of trail mix has 67 grams of carbohydrates, which is 22% of the recommended daily amount. What is the total recommended daily amount of carbohydrates?

Exercise 3.3E. 31

A bacon cheeseburger at a popular fast food restaurant contains 2070 milligrams (mg) of sodium, which is 86% of the recommended daily amount. What is the total recommended daily amount of sodium?

Answer

2407 mg

Exercise 3.3E. 32

A grilled chicken salad at a popular fast food restaurant contains 650 milligrams (mg) of sodium, which is 27% of the recommended daily amount. What is the total recommended daily amount of sodium?

Exercise 3.3E. 33

After 3 months on a diet, Lisa had lost 12% of her original weight. She lost 21 pounds. What was Lisa's original weight?

Answer

175 lb.

Exercise 3.3E.34

Tricia got a 6% raise on her weekly salary. The raise was $30 per week. What was her original salary?

Exercise 3.3E.35

Yuki bought a dress on sale for $72. The sale price was 60% of the original price. What was the original price of the dress?

Answer

$120

Exercise 3.3E.36

Kim bought a pair of shoes on sale for $40.50. The sale price was 45% of the original price. What was the original price of the shoes?

Exercise 3.3E.37

Tim left a $9 tip for a $50 restaurant bill. What percent tip did he leave?

Answer

18%

Exercise 3.3E.38

Rashid left a $15 tip for a $75 restaurant bill. What percent tip did he leave?

Exercise 3.3E.39

The nutrition fact sheet at a fast food restaurant says the fish sandwich has 380 calories, and 171 calories are from fat. What percent of the total calories is from fat?

Answer

45%

Exercise 3.3E.40

The nutrition fact sheet at a fast food restaurant says a small portion of chicken nuggets has 190 calories, and 114 calories are from fat. What percent of the total calories is from fat?

Exercise 3.3E.41

Emma gets paid $3,000 per month. She pays $750 a month for rent. What percent of her monthly pay goes to rent?

Answer

25%

Exercise 3.3E.42

Dimple gets paid $3,200 per month. She pays $960 a month for rent. What percent of her monthly pay goes to rent?

Find Percent Increase and Percent Decrease

In the following exercises, solve.

Exercise 3.3E.43

Tamanika got a raise in her hourly pay, from $15.50 to $17.36. Find the percent increase.

Answer

12%

Exercise 3.3E.44

Ayodele got a raise in her hourly pay, from $24.50 to $25.48. Find the percent increase.

Exercise 3.3E.45

Annual student fees at the University of California rose from about $4,000 in 2000 to about $12,000 in 2010. Find the percent increase.

Answer

200%

Exercise 3.3E.46

The price of a share of one stock rose from $12.50 to $50. Find the percent increase.

Exercise 3.3E.47

According to *Time* magazine annual global seafood consumption rose from 22 pounds per person in the 1960s to 38 pounds per person in 2011. Find the percent increase. (Round to the nearest tenth of a percent.)

Answer

72.7%

Exercise 3.3E.48

In one month, the median home price in the Northeast rose from $225,400 to $241,500. Find the percent increase. (Round to the nearest tenth of a percent.)

Exercise 3.3E.49

A grocery store reduced the price of a loaf of bread from $2.80 to $2.73. Find the percent decrease.

Answer

2.5%

Exercise 3.3E.50

The price of a share of one stock fell from $8.75 to $8.54. Find the percent decrease.

Exercise 3.3E.51

Hernando's salary was $49,500 last year. This year his salary was cut to $44,055. Find the percent decrease.

Answer

11%

Exercise 3.3E.52

In 10 years, the population of Detroit fell from 950,000 to about 712,500. Find the percent decrease.

Exercise 3.3E.53

In 1 month, the median home price in the West fell from $203,400 to $192,300. Find the percent decrease. (Round to the nearest tenth of a percent.)

Answer

5.5%

Exercise 3.3E.54

Sales of video games and consoles fell from $1,150 million to $1,030 million in 1 year. Find the percent decrease. (Round to the nearest tenth of a percent.)

Solve Simple Interest Applications

In the following exercises, solve.

Exercise 3.3E.55

Casey deposited $1,450 in a bank account with interest rate 4%. How much interest was earned in two years?

Answer

$116

Exercise 3.3E.56

Terrence deposited $5,720 in a bank account with interest rate 6%. How much interest was earned in 4 years?

Exercise 3.3E.57

Robin deposited $31,000 in a bank account with interest rate 5.2%. How much interest was earned in 3 years?

Answer

$4,836

Exercise 3.3E.58

Carleen deposited $16,400 in a bank account with interest rate 3.9%. How much interest was earned in 8 years?

Exercise 3.3E.59

Hilaria borrowed $8,000 from her grandfather to pay for college. Five years later, she paid him back the $8,000, plus $1,200 interest. What was the rate of interest?

Answer

3%

Exercise 3.3E.60

Kenneth loaned his niece $1,200 to buy a computer. Two years later, she paid him back the $1,200, plus $96 interest. What was the rate of interest?

Exercise 3.3E.61

Lebron loaned his daughter $20,000 to help her buy a condominium. When she sold the condominium four years later, she paid him the $20,000, plus $3,000 interest. What was the rate of interest?

Answer

3.75%

Exercise 3.3E.62

Pablo borrowed $50,000 to start a business. Three years later, he repaid the $50,000, plus $9,375 interest. What was the rate of interest?

Exercise 3.3E.63

In 10 years, a bank account that paid 5.25% earned $18,375 interest. What was the principal of the account?

Answer

$35,000

Exercise 3.3E.64

In 25 years, a bond that paid 4.75% earned $2,375 interest. What was the principal of the bond?

Exercise 3.3E.65

Joshua's computer loan statement said he would pay $1,244.34 in interest for a 3-year loan at 12.4%. How much did Joshua borrow to buy the computer?

Answer

$3,345

Exercise 3.3E.66

Margaret's car loan statement said she would pay $7,683.20 in interest for a 5-year loan at 9.8%. How much did Margaret borrow to buy the car?

Solve Applications with Discount or Mark-up

In the following exercises, find the sale price.

Exercise 3.3E.67

Perla bought a cell phone that was on sale for $50 off. The original price of the cell phone was $189.

Answer

$139

Exercise 3.3E.68

Sophie saw a dress she liked on sale for $15 off. The original price of the dress was $96.

Exercise 3.3E.69

Rick wants to buy a tool set with original price $165. Next week the tool set will be on sale for $40 off.

Answer

$125

Exercise 3.3E.70

Angelo's store is having a sale on televisions. One television, with original price $859, is selling for $125 off.

In the following exercises, find

1. the amount of discount and
2. the sale price.

Exercise 3.3E.71

Janelle bought a beach chair on sale at 60% off. The original price was $44.95.

Answer

1. $26.97
2. $17.98

Exercise 3.3E.72

Errol bought a skateboard helmet on sale at 40% off. The original price was $49.95.

Exercise 3.3E.73

Kathy wants to buy a camera that lists for $389. The camera is on sale with a 33% discount.

Answer

1. $128.37
2. $260.63

Exercise 3.3E.74

Colleen bought a suit that was discounted 25% from an original price of $245.

Exercise 3.3E.75

Erys bought a treadmill on sale at 35% off. The original price was $949.95 (round to the nearest cent.)

Answer

1. $332.48
2. $617.47

Exercise 3.3E.76

Jay bought a guitar on sale at 45% off. The original price was $514.75 (round to the nearest cent.)

In the following exercises, find

1. the amount of discount and
2. the discount rate. (Round to the nearest tenth of a percent if needed.)

Exercise 3.3E.77

Larry and Donna bought a sofa at the sale price of $1,344. The original price of the sofa was $1,920.

Answer

1. $576
2. 30%

Exercise 3.3E.78

Hiroshi bought a lawnmower at the sale price of $240. The original price of the lawnmower is $300.

Exercise 3.3E.79

Patty bought a baby stroller on sale for $301.75. The original price of the stroller was $355.

Answer

1. $53.25
2. 15%

Exercise 3.3E.80

Bill found a book he wanted on sale for $20.80. The original price of the book was $32.

Exercise 3.3E.81

Nikki bought a patio set on sale for $480. The original price was $850. To the nearest tenth of a percent, what was the rate of discount?

Answer

1. $370
2. 43.5%

Exercise 3.3E.82

Stella bought a dinette set on sale for $725. The original price was $1,299. To the nearest tenth of a percent, what was the rate of discount?

In the following exercises, find ⓐ the amount of the mark-up and ⓑ the list price.

Exercise 3.3E.83

Daria bought a bracelet at original cost $16 to sell in her handicraft store. She marked the price up 45%.

Answer

1. $7.20
2. $23.20

Exercise 3.3E.84

Regina bought a handmade quilt at original cost $120 to sell in her quilt store. She marked the price up 55%.

Exercise 3.3E.85

Tom paid $0.60 a pound for tomatoes to sell at his produce store. He added a 33% mark-up.

Answer

1. $0.20
2. $0.80

Exercise 3.3E.86

Flora paid her supplier $0.74 a stem for roses to sell at her flower shop. She added an 85% mark-up.

Exercise 3.3E.87

Alan bought a used bicycle for $115. After re-conditioning it, he added 225% mark-up and then advertised it for sale.

Answer

1. $258.75
2. $373.75

Exercise 3.3E.88

Michael bought a classic car for $8,500. He restored it, then added 150% mark-up before advertising it for sale.

Everyday Math

Exercise 3.3E.89

Leaving a Tip At the campus coffee cart, a medium coffee costs $1.65. MaryAnne brings $2.00 with her when she buys a cup of coffee and leaves the change as a tip. What percent tip does she leave?

Answer

21.2%

Exercise 3.3E.90

Splitting a Bill Four friends went out to lunch and the bill came to $53.75. They decided to add enough tip to make a total of $64, so that they could easily split the bill evenly among themselves. What percent tip did they leave?

Writing Exercises

Exercise 3.3E.91

Without solving the problem "44 is 80% of what number" think about what the solution might be. Should it be a number that is greater than 44 or less than 44? Explain your reasoning.

Answer

The number should be greater than 44. Since 80% equals 0.8 in decimal form, 0.8 is less than one, and we must multiply the number by 0.8 to get 44, the number must be greater than 44.

Exercise 3.3E.92

Without solving the problem "What is 20% of 300?" think about what the solution might be. Should it be a number that is greater than 300 or less than 300? Explain your reasoning.

Exercise 3.3E.93

After returning from vacation, Alex said he should have packed 50% fewer shorts and 200% more shirts. Explain what Alex meant.

Answer

He meant that he should have packed half the shorts and twice the shirts.

Exercise 3.3E.94

Because of road construction in one city, commuters were advised to plan that their Monday morning commute would take 150% of their usual commuting time. Explain what this means.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
translate and solve basic percent equations.			
solve percent applications.			
find percent increase and percent decrease.			
solve simple interest applications.			
solve applications with discount or mark-up.			

ⓑ After reviewing this checklist, what will you do to become confident for all goals?

This page titled 3.3E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.2E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.4: Solve Mixture Applications

Learning Objectives

By the end of this section, you will be able to:

- Solve coin word problems
- Solve ticket and stamp word problems
- Solve mixture word problems
- Use the mixture model to solve investment problems using simple interest

Note

Before you get started, take this readiness quiz.

1. Multiply: $14(0.25)$.
   If you missed this problem, review Exercise 1.8.19.
2. Solve: $0.25x + 0.10(x + 4) = 2.5$.
   If you missed this problem, review Exercise 2.4.22.
3. The number of dimes is three more than the number of quarters. Let q represent the number of quarters. Write an expression for the number of dimes.
   If you missed this problem, review Exercise 1.3.43.

Solve Coin Word Problems

In **mixture problems**, we will have two or more items with different values to combine together. The mixture model is used by grocers and bartenders to make sure they set fair prices for the products they sell. Many other professionals, like chemists, investment bankers, and landscapers also use the mixture model.

Note

Doing the Manipulative Mathematics activity *Coin Lab* will help you develop a better understanding of mixture word problems.

We will start by looking at an application everyone is familiar with—money!

Imagine that we take a handful of coins from a pocket or purse and place them on a desk. How would we determine the value of that pile of coins? If we can form a step-by-step plan for finding the total value of the coins, it will help us as we begin solving coin word problems.

So what would we do? To get some order to the mess of coins, we could separate the coins into piles according to their value. Quarters would go with quarters, dimes with dimes, nickels with nickels, and so on. To get the total value of all the coins, we would add the total value of each pile.



Figure 3.4.1

How would we determine the value of each pile? Think about the dime pile—how much is it worth? If we count the number of dimes, we'll know how many we have—the *number* of dimes.

But this does not tell us the *value* of all the dimes. Say we counted 17 dimes, how much are they worth? Each dime is worth \$0.10—that is the *value* of one dime. To find the total value of the pile of 17 dimes, multiply 17 by \$0.10 to get \$1.70. This is the total

value of all 17 dimes. This method leads to the following model.

TOTAL VALUE OF COINS

For the same type of coin, the total value of a number of coins is found by using the model

$$number \cdot value = total\ value \tag{3.4.1}$$

where

number is the number of coins

value is the value of each coin

total value is the total value of all the coins

The number of dimes times the value of each dime equals the total value of the dimes.

$$\begin{aligned} \text{number.} \cdot \text{ value } &= \text{ total value} \\ 17 \cdot \$0.10 &= \$1.70 \end{aligned}$$

We could continue this process for each type of coin, and then we would know the total value of each type of coin. To get the total value of *all* the coins, add the total value of each type of coin.

Let's look at a specific case. Suppose there are 14 quarters, 17 dimes, 21 nickels, and 39 pennies.

Type	Number	• Value($)	= Total Value($)
Quarters	14	0.25	3.50
Dimes	17	0.10	1.70
Nickels	21	0.05	1.05
Pennies	39	0.01	0.39
			6.64

Table 3.4.1

The total value of all the coins is $6.64.

Notice how the chart helps organize all the information! Let's see how we use this method to solve a coin word problem.

Example 3.4.1

Adalberto has $2.25 in dimes and nickels in his pocket. He has nine more nickels than dimes. How many of each type of coin does he have?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.

Determine the types of coins involved.
Think about the strategy we used to find the value of the handful of coins. The first thing we need is to notice what types of coins are involved. Adalberto has dimes and nickels. **Create a table** to organize the information. See chart below.

- Label the columns "type," "number," "value," "total value."
- List the types of coins.
- Write in the value of each type of coin.
- Write in the total value of all the coins.

We can work this problem all in cents or in dollars. Here we will do it in dollars and put in the dollar sign ($) in the table as a reminder.
The value of a dime is $0.10 and the value of a nickel is $0.05. The total value of all the coins is $2.25. The table below shows

this information.

Type	Number	•	Value($)	=	Total Value($)
Dimes			0.10		
Nickels			0.05		
					2.25

Step 2. Identify what we are looking for.

We are asked to find the number of dimes and nickels Adalberto has.

Step 3. Name what we are looking for. Choose a variable to represent that quantity.

Use variable expressions to represent the number of each type of coin and write them in the table.

Multiply the number times the value to get the total value of each type of coin.

Next we counted the number of each type of coin. In this problem we cannot count each type of coin—that is what you are looking for—but we have a clue. There are nine more nickels than dimes. The number of nickels is nine more than the number of dimes.

$$\begin{aligned} \text{Let } d &= \text{ number of dimes.} \\ d+9 &= \text{ number of nickels} \end{aligned}$$

Fill in the "number" column in the table to help get everything organized.

Type	Number	•	Value($)	=	Total Value($)
Dimes	d		0.10		
Nickels	$d+9$		0.05		
					2.25

Now we have all the information we need from the problem!

We multiply the number times the value to get the total value of each type of coin. While we do not know the actual number, we do have an expression to represent it.

And so now multiply $\text{number} \cdot \text{value} = \text{total value}$. See how this is done in the table below.

Type	Number	•	Value($)	=	Total Value($)
Dimes	d		0.10		$0.10d$
Nickels	$d+9$		0.05		$0.05(d+9)$
					2.25

Notice that we made the heading of the table show the model.

Step 4. Translate into an equation. It may be helpful to restate the problem in one sentence. Translate the English sentence into an algebraic equation.

Write the equation by adding the total values of all the types of coins.

	Value of dimes	+	value of nickels	=	total value of coins
Translate to an equation.	$0.10d$	+	$0.05(d+9)$	=	2.25

Step 5. Solve the equation using good algebra techniques.

Now solve this equation.	$0.10d + 0.05(d+9) = 2.25$
Distribute.	$0.10d + 0.05d + 0.45 = 2.25$
Combine like terms.	$0.15d + 0.45 = 2.25$
Subtract 0.45 from each side.	$0.15d = 1.80$
Divide.	$d = 12$

So there are 12 dimes.	
The number of nickels is d+9d+9.	$d + 9$
	$12 + 9$
	21

Step 6. Check the answer in the problem and make sure it makes sense.

Does this check?

$$\begin{array}{lrcl} 12 \text{ dimes} & 12(0.10) & = & 1.20 \\ 21 \text{ nickels} & 21(0.05) & = & \underline{1.05} \\ & & & \$2.25 \checkmark \end{array} \qquad (3.4.2)$$

Step 7. Answer the question with a complete sentence.

Adalberto has twelve dimes and twenty-one nickels.

If this were a homework exercise, our work might look like the following.

Adalberto has $2.25 in dimes and nickels in his pocket. He has nine more nickels than dimes.

How many of each type does he have?

Type	Number • Value($)		= Total Value ($)
Dimes	d	0.10	0.10d
Nickels	d + 9	0.05	0.05(d + 9)
			2.25 ✓

12 dimes 12(0.10) = 1.20

21 nickels 21(0.05) = 1.05

$2.25 ✓

Adalberto has twelve dimes and twenty-one nickels.

$0.10d + 0.05d + 0.45 = 2.25$

$0.15d + 0.45 = 2.25$

$0.15d = 1.80$

$d = 12$ dimes

$d + 9$

$12 + 9$

21 nickels

? Try It 3.4.1

Michaela has $2.05 in dimes and nickels in her change purse. She has seven more dimes than nickels. How many coins of each type does she have?

Answer

9 nickels, 16 dimes

? Try It 3.4.2

Liliana has $2.10 in nickels and quarters in her backpack. She has 12 more nickels than quarters. How many coins of each type does she have?

Answer

17 nickels, 5 quarters

SOLVE COIN WORD PROBLEMS.

1. **Read** the problem. Make sure all the words and ideas are understood.
   - Determine the types of coins involved.
   - Create a table to organize the information.
   - Label the columns “type,” “number,” “value,” “total value.”
   - List the types of coins.
   - Write in the value of each type of coin.
   - Write in the total value of all the coins.

Type	Number •	Value($) =	Total Value($)

2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
   - Use variable expressions to represent the number of each type of coin and write them in the table.
   - Multiply the number times the value to get the total value of each type of coin.
4. **Translate** into an equation.
   It may be helpful to restate the problem in one sentence with all the important information. Then, translate the sentence into an equation.
   Write the equation by adding the total values of all the types of coins.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Example 3.4.2

Maria has $2.43 in quarters and pennies in her wallet. She has twice as many pennies as quarters. How many coins of each type does she have?

Solution

Step 1. Read the problem.

Determine the types of coins involved.

We know that Maria has quarters and pennies.

Create a table to organize the information.

- **Step 2. Identify** what you are looking for.
  - We are looking for the number of quarters and pennies.

  Step 3. Name. Represent the number of quarters and pennies using variables.
  - Multiply the ‘number’ and the ‘value’ to get the ‘total value’ of each type of coin.

Type	Number •	Value($) =	Total Value($)
Quarters	q	0.25	$0.25q$
Pennies	$2q$	0.01	$0.01(2q)$
			2.43

  Step 4. Translate. Write the equation by adding the ‘total value’ of all the types of coins.

Step5. Solve the equation.	$0.25q+0.01(2q)=2.43$
Multiply.	$0.25q+0.02q=2.43$
Combine like terms.	$0.27q=2.43$
Divide by 0.27	$q=9$ quarters
The number of pennies is 2q.	$2q$
	$2\cdot 9$
	18 pennies

Step 6. Check the answer in the problem.

Maria has 9 quarters and 18 pennies. Dies this make $2.43?

9 quarters	$9(0.25)$	$=$	2.25
18 pennies	$18(0.01)$	$=$	0.18
Total			$2.43✓

Step 7. Answer the question. Maria has nine quarters and eighteen pennies.

Try It 3.4.3

Sumanta has $4.20 in nickels and dimes in her piggy bank. She has twice as many nickels as dimes. How many coins of each type does she have?

Answer

42 nickels, 21 dimes

Try It 3.4.4

Alison has three times as many dimes as quarters in her purse. She has $9.35 altogether. How many coins of each type does she have?

Answer

51 dimes, 17 quarters

In the next example, we'll show only the completed table—remember the steps we take to fill in the table.

Example 3.4.3

Danny has $2.14 worth of pennies and nickels in his piggy bank. The number of nickels is two more than ten times the number of pennies. How many nickels and how many pennies does Danny have?

Solution

Step 1. Read the problem.	
Determine the types of coins involved.	pennies and nickels
Create a table.	
Write in the value of each type of coin.	Pennies are worth $0.01. Nickels are worth $0.05.
Step 2. Identify what we are looking for.	the number of pennies and nickels

Step 3. Name. Represent the number of each type of coin using variables.	
The number of nickels is defined in terms of the number of pennies, so start with pennies.	Let $p =$ number of pennies.
The number of nickels is two more than ten times the number of pennies.	And let $10p + 2 =$ number of nickels.
Multiply the number and the value to get the total value of each type of coin.	

Type	Number • Value ($) = Total Value ($)		
pennies	p	0.01	$0.01p$
nickels	$10p + 2$	0.05	$0.05(10p + 2)$
			$2.14

Step 4. Translate. Write the equation by adding the total value of all the types of coins.	$0.01p + 0.05(10p + 2) = 2.14$
Step 5. Solve the equation.	$0.01p + 0.50p + 0.10 = 2.14$ $0.51p + 0.10 = 2.14$ $0.51p = 2.04$ $p = 4$ pennies
How many nickels?	$10p + 2$ $10(4) + 2$ 42 nickels
Step 6. Check the answer in the problem and make sure it makes sense Danny has four pennies and 42 nickels. Is the total value $2.14? $4(0.01) + 42(0.05) \stackrel{?}{=} 2.14$ $2.14 = 2.14$✓	
Step 7. Answer the question.	Danny has four pennies and 42 nickels.

Try It 3.4.5

Jesse has $6.55 worth of quarters and nickels in his pocket. The number of nickels is five more than two times the number of quarters. How many nickels and how many quarters does Jesse have?

Answer

41 nickels, 18 quarters

Try It 3.4.6

Elane has $7.00 total in dimes and nickels in her coin jar. The number of dimes that Elane has is seven less than three times the number of nickels. How many of each coin does Elane have?

Answer

22 nickels, 59 dimes

Solve Ticket and Stamp Word Problems

Problems involving tickets or stamps are very much like coin problems. Each type of ticket and stamp has a value, just like each type of coin does. So to solve these problems, we will follow the same steps we used to solve coin problems.

Example 3.4.4

At a school concert, the total value of tickets sold was $1,506. Student tickets sold for $6 each and adult tickets sold for $9 each. The number of adult tickets sold was five less than three times the number of student tickets sold. How many student tickets and how many adult tickets were sold?

Solution

Step 1. Read the problem.

- Determine the types of tickets involved. There are student tickets and adult tickets.
- Create a table to organize the information.

Type	Number •	Value($) =	Total Value($)
Student		6	
Adult		9	
			1506

Step 2. Identify what we are looking for.

- We are looking for the number of student and adult tickets.

Step 3. Name. Represent the number of each type of ticket using variables.

We know the number of adult tickets sold was five less than three times the number of student tickets sold.

- Multiply the number times the value to get the total value of each type of ticket.

Type	Number •	Value($) =	Total Value($)
Student	s	6	$6s$
Adult	$3s-5$	9	$9(3s-5)$
			1506

Step 4. Translate. Write the equation by adding the total values of each type of ticket.

$$6s+9(3s-5)=1506$$

Step 5. Solve the equation.

$$\begin{aligned} 6s+27s-45 &= 1506 \\ 33s-45 &= 1506 \\ 33s &= 1551 \\ s &= 47 \text{ student tickets} \\ \text{Number of adult tickets} &= 3s-5 \\ &= 3(47)-5 \\ \text{So there were} \quad 136 &\quad \text{adult tickets} \end{aligned}$$

Step 6. Check the answer.

There were 47 student tickets at $6 each and 136 adult tickets at $9 each. Is the total value $1,506? We find the total value of each type of ticket by multiplying the number of tickets times its value then add to get the total value of all the tickets sold.

$$\begin{aligned} 47\cdot 6 &= 282 \\ 136\cdot 9 &= \underline{1224} \\ & \quad 1506\checkmark \end{aligned}$$

Step 7. Answer the question. They sold 47 student tickets and 136 adult tickets.

Try It 3.4.7

The first day of a water polo tournament the total value of tickets sold was $17,610. One-day passes sold for $20 and tournament passes sold for $30. The number of tournament passes sold was 37 more than the number of day passes sold. How many day passes and how many tournament passes were sold?

Answer

330 day passes, 367 tournament passes

Try It 3.4.8

At the movie theater, the total value of tickets sold was $2,612.50. Adult tickets sold for $10 each and senior/child tickets sold for $7.50 each. The number of senior/child tickets sold was 25 less than twice the number of adult tickets sold. How many senior/child tickets and how many adult tickets were sold?

Answer

112 adult tickets, 199 senior/child tickets

We have learned how to find the total number of tickets when the number of one type of ticket is based on the number of the other type. Next, we'll look at an example where we know the total number of tickets and have to figure out how the two types of tickets relate.

Suppose Bianca sold a total of 100 tickets. Each ticket was either an adult ticket or a child ticket. If she sold 20 child tickets, how many adult tickets did she sell?

- *Did you say '80'? How did you figure that out? Did you subtract 20 from 100?*

If she sold 45 child tickets, how many adult tickets did she sell?

- *Did you say '55'? How did you find it? By subtracting 45 from 100?*

What if she sold 75 child tickets? How many adult tickets did she sell?

- *The number of adult tickets must be 100−75. She sold 25 adult tickets.*

Now, suppose Bianca sold x child tickets. Then how many adult tickets did she sell? To find out, we would follow the same logic we used above. In each case, we subtracted the number of child tickets from 100 to get the number of adult tickets. We now do the same with x.

We have summarized this below.

Child tickets	Adult tickets
20	80
45	55
75	25
x	$100-x$

Table 3.4.2

We can apply these techniques to other examples

Example 3.4.5

Galen sold 810 tickets for his church's carnival for a total of $2,820. Children's tickets cost $3 each and adult tickets cost $5 each. How many children's tickets and how many adult tickets did he sell?

Solution

Step 1. Read the problem.

- Determine the types of tickets involved. There are children tickets and adult tickets.
- Create a table to organize the information.

Type	Number •	Value($) =	Total Value($)
Children		3	
Adult		5	
			2820

Step 2. Identify what we are looking for.

- We are looking for the number of children and adult tickets.

Step 3. Name. Represent the number of each type of ticket using variables.

- We know the total number of tickets sold was 810.
- This means the number of children's tickets plus the number of adult tickets must add up to 810.
- Let c be the number of children tickets.

Type	Number •	Value($) =	Total Value($)
Children	c	3	$3c$
Adult	$810-c$	5	$5(810-c)$
			2820

- Then $810-c$ is the number of adult tickets.
- Multiply the number times the value to get the total value of each type of ticket.

Step 4. Translate.

Write the equation by adding the total values of each type of ticket.

Step 5. Solve the equation.

$$\begin{aligned} 3c+5(810-c) &= 2,820 \\ 3c+4,050-5c &= 2,820 \\ -2c &= -1,230 \\ c &= 615 \text{ children tickets} \end{aligned}$$

How many adults?

$$\begin{aligned} &810-c \\ &810-615 \\ &195 \text{ adult tickets} \end{aligned}$$

Step 6. Check the answer. There were 615 children's tickets at $3 each and 195 adult tickets at $5 each. Is the total value $2,820?

$$\begin{aligned} 615\cdot 3 &= 1845 \\ 195\cdot 5 &= \underline{975} \\ & 2,820\checkmark \end{aligned}$$

Step 7. Answer the question. Galen sold 615 children's tickets and 195 adult tickets.

Try It 3.4.9

During her shift at the museum ticket booth, Leah sold 115 tickets for a total of $1,163. Adult tickets cost $12 and student tickets cost $5. How many adult tickets and how many student tickets did Leah sell?

Answer

84 adult tickets, 31 student tickets

Try It 3.4.10

A whale-watching ship had 40 paying passengers on board. The total collected from tickets was $1,196. Full-fare passengers paid $32 each and reduced-fare passengers paid $26 each. How many full-fare passengers and how many reduced-fare passengers were on the ship?

Answer

26 full-fare, 14 reduced fare

Now, we'll do one where we fill in the table all at once.

Example 3.4.6

Monica paid $8.36 for stamps. The number of 41-cent stamps was four more than twice the number of two-cent stamps. How many 41-cent stamps and how many two-cent stamps did Monica buy?

Solution

The types of stamps are 41-cent stamps and two-cent stamps. Their names also give the value!

"The number of 41-cent stamps was four more than twice the number of two-cent stamps."

$$\text{Let } x = \text{ number of 2 -cent stamps.}$$
$$2x+4 = \text{ number of 41 - cent stamps}$$

Type	Number •	Value($) =	Total Value($)
41 cent stamps	$2x+4$	0.41	$0.41(2x+4)$
2 cent stamps	x	0.02	$0.02x$
			8.36

Write the equation from the total values.

$$0.41(2x+4)+0.02x = 8.36$$
$$0.82x+1.64+0.02x = 8.36$$
$$0.84x+1.64 = 8.36$$

Solve the equation.

$$0.84x = 6.72$$
$$x = 8$$

Monica bought eight two-cent stamps.

Find the number of 41-cent stamps she bought by evaluating $2x+4$ for $x=8$

$$2x+4$$
$$2(8)+4$$
$$20$$

Check.

$$\begin{aligned} 8(0.02)+20(0.41) &\stackrel{?}{=} 8.36 \\ 0.16+8.20 &\stackrel{?}{=} 8.36 \\ 8.36 &= 8.36\checkmark \end{aligned} \tag{3.4.3}$$

Monica bought eight two-cent stamps and 20 41-cent stamps

Try It 3.4.11

Eric paid $13.36 for stamps. The number of 41-cent stamps was eight more than twice the number of two-cent stamps. How many 41-cent stamps and how many two-cent stamps did Eric buy?

Answer

32 at $0.41, 12 at $0.02

Try It 3.4.12

Kailee paid $12.66 for stamps. The number of 41-cent stamps was four less than three times the number of 20-cent stamps. How many 41-cent stamps and how many 20-cent stamps did Kailee buy?

Answer

26 at $0.41, 10 at $0.20

Solve Mixture Word Problems

Now we'll solve some more general applications of the mixture model. Grocers and bartenders use the mixture model to set a fair price for a product made from mixing two or more ingredients. Financial planners use the mixture model when they invest money in a variety of accounts and want to find the overall interest rate. Landscape designers use the mixture model when they have an assortment of plants and a fixed budget, and event coordinators do the same when choosing appetizers and entrees for a banquet.

Our first mixture word problem will be making trail mix from raisins and nuts.

Example 3.4.7

Henning is mixing raisins and nuts to make 10 pounds of trail mix. Raisins cost $2 a pound and nuts cost $6 a pound. If Henning wants his cost for the trail mix to be $5.20 a pound, how many pounds of raisins and how many pounds of nuts should he use?

Solution

As before, we fill in a chart to organize our information.

The 10 pounds of trail mix will come from mixing raisins and nuts.

$$\text{Let } x = \text{ number of pounds of raisins.}$$
$$10 - x = \text{ number of pounds of nuts}$$

We enter the price per pound for each item.

We multiply the number times the value to get the total value.

Type	Number of pounds	• Price per pound ($)	= Total Value ($)
Raisins	x	2	$2x$
Nuts	$10 - x$	6	$6(10 - x)$
Trail mix	10	5.20	10(5.20)

Notice that the last line in the table gives the information for the total amount of the mixture.

We know the value of the raisins plus the value of the nuts will be the value of the trail mix.

Write the equation from the total values.	$2x + 6(10 - x) = 10(5.20)$
Solve the equation.	$2x + 60 - 6x = 52$
	$-4x = -8$

	$x = 2$ pounds of raisins
Find the number of pounds of nuts.	$10 - x$
	$10 - 2$
	8 pounds of nuts
Check. $2(\$2) + 8(\$6) \stackrel{?}{=} 10(\$5.20)$ $\$4 + \$48 \stackrel{?}{=} \$52$ $\$52 = \$52\checkmark$	
	Henning mixed two pounds of raisins with eight pounds of nuts.

Try It 3.4.13

Orlando is mixing nuts and cereal squares to make a party mix. Nuts sell for \$7 a pound and cereal squares sell for \$4 a pound. Orlando wants to make 30 pounds of party mix at a cost of \$6.50 a pound, how many pounds of nuts and how many pounds of cereal squares should he use?

Answer

5 pounds cereal squares, 25 pounds nuts

Try It 3.4.14

Becca wants to mix fruit juice and soda to make a punch. She can buy fruit juice for \$3 a gallon and soda for \$4 a gallon. If she wants to make 28 gallons of punch at a cost of \$3.25 a gallon, how many gallons of fruit juice and how many gallons of soda should she buy?

Answer

21 gallons of fruit punch, 7 gallons of soda

We can also use the mixture model to solve investment problems using **simple interest**. We have used the simple interest formula, $I = Prt$, where t represented the number of years. When we just need to find the interest for one year, $t = 1$, so then $I = Pr$.

Example 3.4.8

Stacey has \$20,000 to invest in two different bank accounts. One account pays interest at 3% per year and the other account pays interest at 5% per year. How much should she invest in each account if she wants to earn 4.5% interest per year on the total amount?

Solution

We will fill in a chart to organize our information. We will use the simple interest formula to find the interest earned in the different accounts.

The interest on the mixed investment will come from adding the interest from the account earning 3% and the interest from the account earning 5% to get the total interest on the \$20,000.

$$\begin{aligned} \text{Let } x &= \text{amount invested at } 3\% \\ 20,000 - x &= \text{amount invested at } 5\% \end{aligned}$$

The amount invested is the *principal* for each account.

We enter the interest rate for each account.

We multiply the amount invested times the rate to get the interest.

Type	Amount Invested	• Rate	= Interest
3%	x	0.03	$0.03x$
5%	$20{,}000 - x$	0.05	$0.05(20{,}000 - x)$
4.5%	20,000	0.045	$0.045(20{,}000)$

Notice that the total amount invested, 20,000, is the sum of the amount invested at 3% and the amount invested at 5%. And the total interest, $0.045(20,000)$ is the sum of the interest earned in the 3% account and the interest earned in the 5% account.

As with the other mixture applications, the last column in the table gives us the equation to solve.

Write the equation from the interest earned. Solve the equation.	$0.03x + 0.05(20000 - x) = 0.045(20000)$ $0.03x + 1000 - 0.05x = 900$ $-0.02x = -100$ $x = 5000$ $\text{amount invested at } 3\%$
Find the amount invested at 5%.	$20{,}000 - x$ $20{,}000 - 5{,}000$ 15,000 = amount invested at 5%
Check. $0.03x + 0.05(15000 + x) \stackrel{?}{=} 0.045(20000)$ $150 + 750 \stackrel{?}{=} 900$ $900 = 900\checkmark$	
	Stacey should invest \$5,000 in the account that earns 3% and \$15,000 in the account that earns 5%.

Try It 3.4.15

Remy has \$14,000 to invest in two mutual funds. One fund pays interest at 4% per year and the other fund pays interest at 7% per year. How much should she invest in each fund if she wants to earn 6.1% interest on the total amount?

Answer

\$4,200 at 4%, \$9,800 at 7%

Try It 3.4.16

Marco has \$8,000 to save for his daughter's college education. He wants to divide it between one account that pays 3.2% interest per year and another account that pays 8% interest per year. How much should he invest in each account if he wants the interest on the total investment to be 6.5%?

Answer

\$2,500 at 3.2%, \$5,500 at 8%

Key Concepts

- **Total Value of Coins** For the same type of coin, the total value of a number of coins is found by using the model. number·value=total value where *number* is the number of coins and *value* is the value of each coin; *total value* is the total value of all the coins
- **Problem-Solving Strategy—Coin Word Problems**

1. **Read** the problem. Make all the words and ideas are understood. Determine the types of coins involved.
   - Create a table to organize the information.
   - Label the columns type, number, value, total value.
   - List the types of coins.
   - Write in the value of each type of coin.
   - Write in the total value of all the coins.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
   Use variable expressions to represent the number of each type of coin and write them in the table.
   Multiply the number times the value to get the total value of each type of coin.
4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information.
   Then, translate the sentence into an equation.
   Write the equation by adding the total values of all the types of coins.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Glossary

mixture problems

Mixture problems combine two or more items with different values together.

This page titled 3.4: Solve Mixture Applications is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.3: Solve Mixture Applications** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.4E: Exercises

Practice Makes Perfect

Solve Coin Word Problems

In the following exercises, solve each coin word problem.

Exercise 3.4E.1

Jaime has $2.60 in dimes and nickels. The number of dimes is 14 more than the number of nickels. How many of each coin does he have?

Answer

8 nickels, 22 dimes

Exercise 3.4E.2

Lee has $1.75 in dimes and nickels. The number of nickels is 11 more than the number of dimes. How many of each coin does he have?

Exercise 3.4E.3

Ngo has a collection of dimes and quarters with a total value of $3.50. The number of dimes is seven more than the number of quarters. How many of each coin does he have?

Answer

15 dimes, 8 quarters

Exercise 3.4E.4

Connor has a collection of dimes and quarters with a total value of $6.30. The number of dimes is 14 more than the number of quarters. How many of each coin does he have?

Exercise 3.4E.5

A cash box of $1 and $5 bills is worth $45. The number of $1 bills is three more than the number of $5 bills. How many of each bill does it contain?

Answer

10 at $1, 7 at $5

Exercise 3.4E.6

Joe’s wallet contains $1 and $5 bills worth $47. The number of $1 bills is five more than the number of $5 bills. How many of each bill does he have?

Exercise 3.4E.7

Rachelle has $6.30 in nickels and quarters in her coin purse. The number of nickels is twice the number of quarters. How many coins of each type does she have?

Answer

18 quarters, 36 nickels

Exercise 3.4E.8

Deloise has $1.20 in pennies and nickels in a jar on her desk. The number of pennies is three times the number of nickels. How many coins of each type does she have?

Exercise 3.4E.9

Harrison has $9.30 in his coin collection, all in pennies and dimes. The number of dimes is three times the number of pennies. How many coins of each type does he have?

Answer

30 pennies, 90 dimes

Exercise 3.4E.10

Ivan has $8.75 in nickels and quarters in his desk drawer. The number of nickels is twice the number of quarters. How many coins of each type does he have?

Exercise 3.4E.11

In a cash drawer there is $125 in $5 and $10 bills. The number of $10 bills is twice the number of $5 bills. How many of each are in the drawer?

Answer

10 at $10, 5 at $5

Exercise 3.4E.12

John has $175 in $5 and $10 bills in his drawer. The number of $5 bills is three times the number of $10 bills. How many of each are in the drawer?

Exercise 3.4E.13

Carolyn has $2.55 in her purse in nickels and dimes. The number of nickels is nine less than three times the number of dimes. Find the number of each type of coin.

Answer

12 dimes and 27 nickels

Exercise 3.4E.14

Julio has $2.75 in his pocket in nickels and dimes. The number of dimes is 10 less than twice the number of nickels. Find the number of each type of coin.

Exercise 3.4E.15

Chi has $11.30 in dimes and quarters. The number of dimes is three more than three times the number of quarters. How many of each are there?

Answer

63 dimes, 20 quarters

Exercise 3.4E.16

Tyler has $9.70 in dimes and quarters. The number of quarters is eight more than four times the number of dimes. How many of each coin does he have?

Exercise 3.4E.17

Mukul has $3.75 in quarters, dimes and nickels in his pocket. He has five more dimes than quarters and nine more nickels than quarters. How many of each coin are in his pocket?

Answer

16 nickels, 12 dimes, 7 quarters

Exercise 3.4E.18

Vina has $4.70 in quarters, dimes and nickels in her purse. She has eight more dimes than quarters and six more nickels than quarters. How many of each coin are in her purse?

Solve Ticket and Stamp Word Problems

In the following exercises, solve each ticket or stamp word problem.

Exercise 3.4E.19

The school play sold $550 in tickets one night. The number of $8 adult tickets was 10 less than twice the number of $5 child tickets. How many of each ticket were sold?

Answer

30 child tickets, 50 adult tickets

Exercise 3.4E.20

If the number of $8 child tickets is seventeen less than three times the number of $12 adult tickets and the theater took in $584, how many of each ticket were sold?

Exercise 3.4E.21

The movie theater took in $1,220 one Monday night. The number of $7 child tickets was ten more than twice the number of $9 adult tickets. How many of each were sold?

Answer

110 child tickets, 50 adult tickets

Exercise 3.4E.22

The ball game sold $1,340 in tickets one Saturday. The number of $12 adult tickets was 15 more than twice the number of $5 child tickets. How many of each were sold?

Exercise 3.4E. 23

The ice rink sold 95 tickets for the afternoon skating session, for a total of $828. General admission tickets cost $10 each and youth tickets cost $8 each. How many general admission tickets and how many youth tickets were sold?

Answer

34 general, 61 youth

Exercise 3.4E. 24

For the 7:30 show time, 140 movie tickets were sold. Receipts from the $13 adult tickets and the $10 senior tickets totaled $1,664. How many adult tickets and how many senior tickets were sold?

Exercise 3.4E. 25

The box office sold 360 tickets to a concert at the college. The total receipts were $4170. General admission tickets cost $15 and student tickets cost $10. How many of each kind of ticket was sold?

Answer

114 general, 246 student

Exercise 3.4E. 26

Last Saturday, the museum box office sold 281 tickets for a total of $3954. Adult tickets cost $15 and student tickets cost $12. How many of each kind of ticket was sold?

Exercise 3.4E. 27

Julie went to the post office and bought both $0.41 stamps and $0.26 postcards. She spent $51.40. The number of stamps was 20 more than twice the number of postcards. How many of each did she buy?

Answer

40 postcards, 100 stamps

Exercise 3.4E. 28

Jason went to the post office and bought both $0.41 stamps and $0.26 postcards and spent $10.28. The number of stamps was four more than twice the number of postcards. How many of each did he buy?

Exercise 3.4E. 29

Maria spent $12.50 at the post office. She bought three times as many $0.41 stamps as $0.02 stamps. How many of each did she buy?

Answer

30 at $0.41, 10 at $0.02

Exercise 3.4E. 30

Hector spent $33.20 at the post office. He bought four times as many $0.41 stamps as $0.02 stamps. How many of each did he buy?

Exercise 3.4E.31

Hilda has $210 worth of $10 and $12 stock shares. The numbers of $10 shares is five more than twice the number of $12 shares. How many of each does she have?

Answer

15 $10 shares, 5 $12 shares

Exercise 3.4E.32

Mario invested $475 in $45 and $25 stock shares. The number of $25 shares was five less than three times the number of $45 shares. How many of each type of share did he buy?

Solve Mixture Word Problems

In the following exercises, solve each mixture word problem.

Exercise 3.4E.33

Lauren in making 15 liters of mimosas for a brunch banquet. Orange juice costs her $1.50 per liter and champagne costs her $12 per liter. How many liters of orange juice and how many liters of champagne should she use for the mimosas to cost Lauren $5 per liter?

Answer

5 liters champagne, 10 liters orange juice

Exercise 3.4E.34

Macario is making 12 pounds of nut mixture with macadamia nuts and almonds. Macadamia nuts cost $9 per pound and almonds cost $5.25 per pound. How many pounds of macadamia nuts and how many pounds of almonds should Macario use for the mixture to cost $6.50 per pound to make?

Exercise 3.4E.35

Kaapo is mixing Kona beans and Maui beans to make 25 pounds of coffee blend. Kona beans cost Kaapo $15 per pound and Maui beans cost $24 per pound. How many pounds of each coffee bean should Kaapo use for his blend to cost him $17.70 per pound?

Answer

7.5 lbs Maui beans, 17.5 Kona beans

Exercise 3.4E.36

Estelle is making 30 pounds of fruit salad from strawberries and blueberries. Strawberries cost $1.80 per pound and blueberries cost $4.50 per pound. If Estelle wants the fruit salad to cost her $2.52 per pound, how many pounds of each berry should she use?

Exercise 3.4E.37

Carmen wants to tile the floor of his house. He will need 1000 square feet of tile. He will do most of the floor with a tile that costs $1.50 per square foot, but also wants to use an accent tile that costs $9.00 per square foot. How many square feet of each tile should he plan to use if he wants the overall cost to be $3 per square foot?

Answer

800 at $1.50, 200 at $9.00

Exercise 3.4E. 38

Riley is planning to plant a lawn in his yard. He will need nine pounds of grass seed. He wants to mix Bermuda seed that costs $4.80 per pound with Fescue seed that costs $3.50 per pound. How much of each seed should he buy so that the overall cost will be $4.02 per pound?

Exercise 3.4E. 39

Vartan was paid $25,000 for a cell phone app that he wrote and wants to invest it to save for his son's education. He wants to put some of the money into a bond that pays 4% annual interest and the rest into stocks that pay 9% annual interest. If he wants to earn 7.4% annual interest on the total amount, how much money should he invest in each account?

Answer

$8000 at 4%, $17,000 at 9%

Exercise 3.4E. 40

Vern sold his 1964 Ford Mustang for $55,000 and wants to invest the money to earn him 5.8% interest per year. He will put some of the money into Fund A that earns 3% per year and the rest in Fund B that earns 10% per year. How much should he invest into each fund if he wants to earn 5.8% interest per year on the total amount?

Exercise 3.4E. 41

Stephanie inherited $40,000. She wants to put some of the money in a certificate of deposit that pays 2.1% interest per year and the rest in a mutual fund account that pays 6.5% per year. How much should she invest in each account if she wants to earn 5.4% interest per year on the total amount?

Answer

$10,000 in CD, $30,000 in mutual fund

Exercise 3.4E. 42

Avery and Caden have saved $27,000 towards a down payment on a house. They want to keep some of the money in a bank account that pays 2.4% annual interest and the rest in a stock fund that pays 7.2% annual interest. How much should they put into each account so that they earn 6% interest per year?

Exercise 3.4E. 43

Dominic pays 7% interest on his $15,000 college loan and 12% interest on his $11,000 car loan. What average interest rate does he pay on the total $26,000 he owes? (Round your answer to the nearest tenth of a percent.)

Answer

9.1%

Exercise 3.4E.44

Liam borrowed a total of $35,000 to pay for college. He pays his parents 3% interest on the $8,000 he borrowed from them and pays the bank 6.8% on the rest. What average interest rate does he pay on the total $35,000? (Round your answer to the nearest tenth of a percent.)

Everyday Math

Exercise 3.4E.45

As the treasurer of her daughter's Girl Scout troop, Laney collected money for some girls and adults to go to a 3-day camp. Each girl paid $75 and each adult paid $30. The total amount of money collected for camp was $765. If the number of girls is three times the number of adults, how many girls and how many adults paid for camp?

Answer

9 girls, 3 adults

Exercise 3.4E.46

Laurie was completing the treasurer's report for her son's Boy Scout troop at the end of the school year. She didn't remember how many boys had paid the $15 full-year registration fee and how many had paid the $10 partial-year fee. She knew that the number of boys who paid for a full-year was ten more than the number who paid for a partial-year. If $250 was collected for all the registrations, how many boys had paid the full-year fee and how many had paid the partial-year fee?

Writing Exercises

Exercise 3.4E.47

Suppose you have six quarters, nine dimes, and four pennies. Explain how you find the total value of all the coins.

Answer

Answers will vary.

Exercise 3.4E.48

Do you find it helpful to use a table when solving coin problems? Why or why not?

Exercise 3.4E.49

In the table used to solve coin problems, one column is labeled "number" and another column is labeled "value." What is the difference between the "number" and the "value?"

Answer

Answers will vary.

Exercise 3.4E.50

What similarities and differences did you see between solving the coin problems and the ticket and stamp problems?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve coin word problems.			
solve ticket and stamp word problems.			
solve mixture word problems.			

ⓑ After reviewing this checklist, what will you do to become confident for all objectives?

This page titled 3.4E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.3E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.5: Triangles, Rectangles, and the Pythagorean Theorem

Learning Objectives

By the end of this section, you will be able to:

- Solve applications using properties of triangles
- Use the Pythagorean Theorem
- Solve applications using rectangle properties

Be prepared

Before you get started, take this readiness quiz.

1. Simplify: $12(6h)$.
   If you missed this problem, review Exercise 1.10.1.
2. The length of a rectangle is three less than the width. Let w represent the width. Write an expression for the length of the rectangle.
   If you missed this problem, review Exercise 1.3.43.
3. Solve: $A = \frac{1}{2}bh$ for b when A=260 and h=52.
   If you missed this problem, review Exercise 2.6.10.
4. Simplify: $\sqrt{144}$.
   If you missed this problem, review Exercise 1.9.10.

Solve Applications Using Properties of Triangles

In this section we will use some common geometry formulas. We will adapt our problem-solving strategy so that we can solve geometry applications. The geometry formula will name the variables and give us the equation to solve. In addition, since these applications will all involve shapes of some sort, most people find it helpful to draw a figure and label it with the given information. We will include this in the first step of the problem solving strategy for geometry applications.

SOLVE GEOMETRY APPLICATIONS

1. **Read** the problem and make sure all the words and ideas are understood. Draw the figure and label it with the given information.
2. **Identify** what we are looking for.
3. **Label** what we are looking for by choosing a variable to represent it.
4. **Translate** into an equation by writing the appropriate formula or model for the situation. Substitute in the given information.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer by substituting it back into the equation solved in step 5 and by making sure it makes sense in the context of the problem.
7. **Answer** the question with a complete sentence.

We will start geometry applications by looking at the properties of triangles. Let's review some basic facts about triangles. Triangles have three sides and three interior angles. Usually each side is labeled with a lowercase letter to match the uppercase letter of the opposite vertex.

The plural of the word *vertex* is *vertices*. All triangles have three **vertices**. Triangles are named by their vertices: The triangle in Figure 3.5.1 is called $\triangle ABC$.




Figure 3.5.1: Triangle ABC has vertices A, B, and C. The lengths of the sides are a, b, and c.

The three angles of a triangle are related in a special way. The sum of their measures is 180°. Note that we read $m\angle A$ as "the measure of angle A." So in $\triangle ABC$ in Figure 3.5.1.

$$m\angle A + m\angle B + m\angle C = 180^\circ$$

Because the **perimeter** of a figure is the length of its boundary, the perimeter of $\triangle ABC$ is the sum of the lengths of its three sides.

$$P = a + b + c$$

To find the **area** of a triangle, we need to know its base and height. The height is a line that connects the base to the opposite vertex and makes a 90° angle with the base. We will draw $\triangle ABC$ again, and now show the height, h. See Figure 3.5.2.





Figure 3.5.2: The formula for the area of $\triangle ABC$ is $A = \frac{1}{2}bh$ where b is the base and h is the height.

TRIANGLE PROPERTIES



For $\triangle ABC$

Angle measures:

$$m\angle A + m\angle B + m\angle C = 180^\circ \tag{3.5.1}$$

- The sum of the measures of the angles of a triangle is 180°.

Perimeter:

$$P = a + b + c \tag{3.5.2}$$

- The perimeter is the sum of the lengths of the sides of the triangle.

Area:


$A = \frac{1}{2}bh, b =$ base , $h =$ height

- The area of a triangle is one-half the base times the height.

Example 3.5.1

The measures of two angles of a triangle are 55 and 82 degrees. Find the measure of the third angle.

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	
Step 2. Identify what you are looking for.	the measure of the third angle in a triangle
Step 3. Name. Choose a variable to represent it.	Let $x =$ the measure of the angle.
Step 4. Translate.	
Write the appropriate formula and substitute.	$m\angle A + m\angle B + m\angle C = 180^\circ$
Step 5. Solve the equation.	$\begin{aligned} 55 + 82 + x &= 180 \\ 137 + x &= 180 \\ x &= 43 \end{aligned}$
Step 6. Check. $\begin{aligned} 55 + 82 + 43 &\stackrel{?}{=} 180 \\ 180 &= 180\checkmark \end{aligned}$	
Step 7. Answer the question.	The measure of the third angle is 43 degrees.

Try It 3.5.1

The measures of two angles of a triangle are 31 and 128 degrees. Find the measure of the third angle.

Answer

21 degrees

Try It 3.5.2

The measures of two angles of a triangle are 49 and 75 degrees. Find the measure of the third angle.


Answer

56 degrees

Example 3.5.2

The perimeter of a triangular garden is 24 feet. The lengths of two sides are four feet and nine feet. How long is the third side?

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	 P = 24 ft
Step 2. Identify what you are looking for.	length of the third side of a triangle
Step 3. Name. Choose a variable to represent it.	Let $c =$ the third side.
Step 4. Translate.	
Write the appropriate formula and substitute.	$P = a + b + c$
Substitute in the given information.	$24 \text{ ft} = 4 \text{ ft} + 9 \text{ ft} + c$
Step 5. Solve the equation.	$24 = 13 + c$ $11 = c$
Step 6. Check. $P = a+b+c$ $24 \stackrel{?}{=} 4+9+11$ $24 = 24\checkmark$	
Step 7. Answer the question.	The third side is 11 feet long.

Try It 3.5.3

The perimeter of a triangular garden is 48 feet. The lengths of two sides are 18 feet and 22 feet. How long is the third side?

Answer

8 feet

Try It 3.5.4

The lengths of two sides of a triangular window are seven feet and five feet. The perimeter is 18 feet. How long is the third side?


Answer

6 feet

Example 3.5.3

The area of a triangular church window is 90 square meters. The base of the window is 15 meters. What is the window's height?

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	 Area $= 90m^2$
Step 2. Identify what you are looking for.	height of a triangle
Step 3. Name. Choose a variable to represent it.	Let $h =$ the height.
Step 4. Translate.	
Write the appropriate formula.	$A = \frac{1}{2} \cdot b \cdot h$
Substitute in the given information.	$90\ \text{m}^2 = \frac{1}{2} \cdot 15\ \text{m} \cdot h$
Step 5. Solve the equation.	$90 = \frac{15}{2}h$ $12 = h$
Step 6. Check.	$A = \frac{1}{2}bh$ $90 \stackrel{?}{=} \frac{1}{2} \cdot 15 \cdot 12$ $90 = 90\checkmark$
Step 7. Answer the question.	The height of the triangle is 12 meters.

? Try It 3.5.5

The area of a triangular painting is 126 square inches. The base is 18 inches. What is the height?

Answer

14 inches

? Try It 3.5.6

A triangular tent door has area 15 square feet. The height is five feet. What is the base?

Answer

6 feet

The triangle properties we used so far apply to all triangles. Now we will look at one specific type of triangle—a right triangle. A **right triangle** has one 90° angle, which we usually mark with a small square in the corner.




Figure 3.5.3


Definition: RIGHT TRIANGLE

A **right triangle** has one 90° angle, which is often marked with a square at the vertex.

Example 3.5.4

One angle of a right triangle measures 28°. What is the measure of the third angle?

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	
Step 2. Identify what you are looking for.	the measure of an angle
Step 3. Name. Choose a variable to represent it.	Let $x =$ the measure of an angle.
Step 4. Translate.	$m\angle A + m\angle B + m\angle C = 180$
Write the appropriate formula and substitute.	$x + 90 + 28 = 180$
Step 5. Solve the equation.	$x = 62$
Step 6. Check. $180 \stackrel{?}{=} 90 + 28 + 62$ $180 = 180\checkmark$	
Step 7. Answer the question.	The measure of the third angle is 62°.

Try It 3.5.7

One angle of a right triangle measures 56°. What is the measure of the other small angle?

Answer

34°

Try It 3.5.8

One angle of a right triangle measures 45°. What is the measure of the other small angle?

Answer


45°

In the examples we have seen so far, we could draw a figure and label it directly after reading the problem. In the next example, we will have to define one angle in terms of another. We will wait to draw the figure until we write expressions for all the angles we are looking for.

Example 3.5.5

The measure of one angle of a right triangle is 20 degrees more than the measure of the smallest angle. Find the measures of all three angles.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the measures of all three angles
Step 3. Name. Choose a variable to represent it.	Let $a = 1^{st}$ angle. $a + 20 = 2^{nd}$ angle $90 = 3^{rd}$ angle (the right angle)
Draw the figure and label it with the given information	
Step 4. Translate	$m\angle A + m\angle B + m\angle C = 180$
Write the appropriate formula. Substitute into the formula.	$a + (a + 20) + 90 = 180$
Step 5. Solve the equation.	$2a + 110 = 180$ $2a = 70$ $a = 35$ first angle $a + 20$ second angle $35 + 20 = 55$ And the third angle is 90.
Step 6. Check. $35 + 55 + 90 \stackrel{?}{=} 180$ $180 = 180\checkmark$	
Step 7. Answer the question.	The three angles measure 35°, 55°, and 90°.

? Try It 3.5.9

The measure of one angle of a right triangle is 50° more than the measure of the smallest angle. Find the measures of all three angles.

Answer

20°,70°,90°

? Try It 3.5.10

The measure of one angle of a right triangle is 30° more than the measure of the smallest angle. Find the measures of all three angles.

Answer

30°,60°,90°

Use the Pythagorean Theorem

We have learned how the measures of the angles of a triangle relate to each other. Now, we will learn how the lengths of the sides relate to each other. An important property that describes the relationship among the lengths of the three sides of a right triangle is

called the **Pythagorean Theorem**. This theorem has been used around the world since ancient times. It is named after the Greek philosopher and mathematician, Pythagoras, who lived around 500 BC.

Before we state the Pythagorean Theorem, we need to introduce some terms for the sides of a triangle. Remember that a right triangle has a 90° angle, marked with a small square in the corner. The side of the triangle opposite the 90°90° angle is called the *hypotenuse* and each of the other sides are called *legs*.





Figure 3.5.4

The Pythagorean Theorem tells how the lengths of the three sides of a right triangle relate to each other. It states that in any right triangle, the sum of the squares of the lengths of the two legs equals the square of the length of the hypotenuse. In symbols we say: in any right triangle, $a^2+b^2=c^2$, where a and b are the lengths of the legs and cc is the length of the hypotenuse.

Writing the formula in every exercise and saying it aloud as you write it, may help you remember the Pythagorean Theorem.

THE PYTHAGOREAN THEOREM

In any right triangle, where a and b are the lengths of the legs, c is the length of the hypotenuse.



Then

$$a^2+b^2=c^2 \tag{3.5.3}$$

To solve exercises that use the Pythagorean Theorem (Equation 3.5.3), we will need to find square roots. We have used the notation $\sqrt{m}$ and the definition:


If $m=n^2$, then $\sqrt{m}=n$, for $n\geq 0$.

For example, we found that $\sqrt{25}$ is 5 because $25=5^2$.

Because the Pythagorean Theorem contains variables that are squared, to solve for the length of a side in a right triangle, we will have to use square roots.


Example 3.5.6

Use the Pythagorean Theorem to find the length of the hypotenuse shown below.




Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the length of the hypotenuse of the triangle

Step 3. Name. Choose a variable to represent it. Label side c on the figure.	Let c = the length of the hypotenuse. 
Step 4. Translate.	
Write the appropriate formula.	$a^2+b^2=c^2$
Substitute.	$3^2+4^2=c^2$
Step 5. Solve the equation.	$9+16=c^2$
Simplify.	$25=c^2$
Use the definition of square root.	$\sqrt{25}=c$
Simplify.	$5=c$
Step 6. Check. $3^2+4^2 \stackrel{?}{=} 5^2$ $9+16 \stackrel{?}{=} 25$ $25=25$ ✓	
Step 7. Answer the question.	The length of the hypotenuse is 5.

? Try It 3.5.11

Use the Pythagorean Theorem to find the length of the hypotenuse in the triangle shown below.




Answer

c=10

? Try It 3.5.12

Use the Pythagorean Theorem to find the length of the hypotenuse in the triangle shown below.




Answer


c=13

Example 3.5.7

Use the Pythagorean Theorem to find the length of the leg shown below.




Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the length of the leg of the triangle
Step 3. Name. Choose a variable to represent it.	Let $b =$ the leg of the triangle.
Label side b.	
Step 4. Translate	
Write the appropriate formula.	$a^2 + b^2 = c^2$
Substitute.	$5^2 + b^2 = 13^2$
Step 5. Solve the equation.	$25 + b^2 = 169$
Isolate the variable term.	$b^2 = 144$
Use the definition of square root.	$b = \sqrt{144}$
Simplify.	$b = 12$
Step 6. Check. $5^2 + 12^2 \stackrel{?}{=} 13^2$ $25 + 144 \stackrel{?}{=} 169$ $169 = 169$ ✓	
Step 7. Answer the question.	The length of the leg is 12.

Try It 3.5.13

Use the Pythagorean Theorem to find the length of the leg in the triangle shown below.




Answer

8

Try It 3.5.14


Use the Pythagorean Theorem to find the length of the leg in the triangle shown below.



Answer

12

Example 3.5.8



Kelvin is building a gazebo and wants to brace each corner by placing a 10″ piece of wood diagonally as shown above.

If he fastens the wood so that the ends of the brace are the same distance from the corner, what is the length of the legs of the right triangle formed? Approximate to the nearest tenth of an inch.

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for.	the distance from the corner that the bracket should be attached
Step 3. Name. Choose a variable to represent it.	Let x = distance from the corner.
Step 4. Translate Write the appropriate formula and substitute.	$a^2+b^2=c^2$ $x^2+x^2=10^2$
Step 5. Solve the equation.	
	$2x^2=100$
Isolate the variable.	$x^2=50$
Simplify. Approximate to the nearest tenth.	$x\approx 7.1$

Step 6. Check.
$a^2+b^2=c^2$
$(7.1)^2+(7.1)^2\approx 10^2$ Yes.

Step 7. Answer the question.	Kelven should fasten each piece of wood approximately 7.1'' from the corner.

? Try It 3.5.15

John puts the base of a 13-foot ladder five feet from the wall of his house as shown below. How far up the wall does the ladder reach?




Answer

12 feet

? Try It 3.5.16

Randy wants to attach a 17 foot string of lights to the top of the 15 foot mast of his sailboat, as shown below. How far from the base of the mast should he attach the end of the light string?




Answer

8 feet

Solve Applications Using Rectangle Properties


You may already be familiar with the properties of rectangles. Rectangles have four sides and four right (90°) angles. The opposite sides of a **rectangle** are the same length. We refer to one side of the rectangle as the length, L, and its adjacent side as the width, W.



The distance around this rectangle is $L+W+L+W$, or $2L+2W$. This is the **perimeter**, P, of the rectangle.

$$P=2L+2W \tag{3.5.4}$$

What about the **area** of a rectangle? Imagine a rectangular rug that is 2-feet long by 3-feet wide. Its area is 6 square feet. There are six squares in the figure.



$$\begin{aligned} A&=6\\ A&=2\cdot 3\\ A&=L\cdot W \end{aligned} \tag{3.5.5}$$

The area is the length times the width. The formula for the area of a rectangle is

$$A=LW. \tag{3.5.6}$$

PROPERTIES OF RECTANGLES

Rectangles have four sides and four right (90°) angles.

The lengths of opposite sides are equal.

The perimeter of a rectangle is the sum of twice the length and twice the width.

$$P = 2L + 2W \tag{3.5.7}$$


The area of a rectangle is the product of the length and the width.

$$A = L \cdot W \tag{3.5.8}$$

Example 3.5.9

The length of a rectangle is 32 meters and the width is 20 meters. What is the perimeter?

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	
Step 2. Identify what you are looking for.	the perimeter of a rectangle
Step 3. Name. Choose a variable to represent it.	Let $P =$ the perimeter.
Step 4. Translate.	
Write the appropriate formula.	$P = 2L + 2W$
Substitute.	$P = 2(32\text{ m}) + 2(20\text{ m})$
Step 5. Solve the equation.	$P = 64 + 40$ $P = 104$
Step 6. Check. $P \stackrel{?}{=} 104$ $20 + 32 + 20 + 32 \stackrel{?}{=} 104$ $104 = 104\checkmark$	
Step 7. Answer the question.	The perimeter of the rectangle is 104 meters.

Try It 3.5.17

The length of a rectangle is 120 yards and the width is 50 yards. What is the perimeter?

Answer

340 yards

Try It 3.5.18

The length of a rectangle is 62 feet and the width is 48 feet. What is the perimeter?



Answer

220 feet

Example 3.5.10

The area of a rectangular room is 168 square feet. The length is 14 feet. What is the width?

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	
Step 2. Identify what you are looking for.	the width of a rectangular room
Step 3. Name. Choose a variable to represent it.	Let $W =$ the width.
Step 4. Translate.	
Write the appropriate formula.	$A = LW$
Substitute.	$168 = 14W$
Step 5. Solve the equation.	$\frac{168}{14} = \frac{14W}{14}$ $12 = W$
Step 6. Check.  $A = LW$ $168 \stackrel{?}{=} 14 \cdot 12$ $168 = 168$✓	
Step 7. Answer the question.	The width of the room is 12 feet.

Try It 3.5.19

The area of a rectangle is 598 square feet. The length is 23 feet. What is the width?

Answer

26 feet

Try It 3.5.20

The width of a rectangle is 21 meters. The area is 609 square meters. What is the length?



Answer

29 meters

Example 3.5.11

Find the length of a rectangle with perimeter 50 inches and width 10 inches.

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	
Step 2. Identify what you are looking for.	the length of the rectangle
Step 3. Name. Choose a variable to represent it.	Let $L =$ the length.
Step 4. Translate.	
Write the appropriate formula.	$P = 2L + 2W$
Substitute.	$50 = 2L + 2(10)$
Step 5. Solve the equation.	$50 - 20 = 2L + 20 - 20$ $30 = 2L$ $\frac{30}{2} = \frac{2L}{2}$ $15 = L$
Step 6. Check.  $P = 50$ $15 + 10 + 15 + 10 \stackrel{?}{=} 50$ $50 = 50\checkmark$	
Step 7. Answer the question.	The length is 15 inches.

Try It 3.5.21

Find the length of a rectangle with: perimeter 80 and width 25.

Answer

15

Try It 3.5.22

Find the length of a rectangle with: perimeter 30 and width 6.

Answer



9

We have solved problems where either the length or width was given, along with the perimeter or area; now we will learn how to solve problems in which the width is defined in terms of the length. We will wait to draw the figure until we write an expression for the width so that we can label one side with that expression.

Example 3.5.12

The width of a rectangle is two feet less than the length. The perimeter is 52 feet. Find the length and width.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the length and width of a rectangle
Step 3. Name. Choose a variable to represent it. Since the width is defined in terms of the length, we let $L =$ length. The width is two feet less than the length, so we let $L-2$ width.	 $P = 52$ ft
Step 4. Translate.	
Write the appropriate formula. The formula for the perimeter of a rectangle relates all the information.	$P = 2L + 2W$
Substitute in the given information.	$52 = 2L + 2(L-2)$
Step 5. Solve the equation.	$52 = 2L + 2L - 4$
Combine like terms.	$52 = 4L - 4$
Add 4 to each side.	$56 = 4L$
Divide by 4.	$\frac{56}{4} = \frac{4L}{4}$ $14 = L$ The length is 14 feet.
Now we need to find the width.	The width is $L-2$. $L-2$ $14-2$ 12 The width is 12 feet.
Step 6. Check. Since $14+12+14+12=52$, this works! 	
Step 7. Answer the question.	The length is 14 feet and the width is 12 feet.

Try It 3.5.23

The width of a rectangle is seven meters less than the length. The perimeter is 58 meters. Find the length and width.

Answer

18 meters, 11 meters

Try It 3.5.24

The length of a rectangle is eight feet more than the width. The perimeter is 60 feet. Find the length and width.


Answer


19 feet, 11 feet

Example 3.5.13

The length of a rectangle is four centimeters more than twice the width. The perimeter is 32 centimeters. Find the length and width.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	the length and the width
Step 3. Name. Choose a variable to represent the width.	Let W = width
The length is four more than twice the width.	$2W + 4$ = length  $P = 32$ cm
Step 4. Translate	
Write the appropriate formula.	$P = 2L + 2W$
Substitute in the given information.	$32 = 2(2W + 4) + 2W$
Step 5. Solve the equation.	$32 = 4W + 8 + 2W$ $32 = 6W + 8$ $24 = 6W$ $4 = W$ (width) $2W + 4$ (length) $2(4) + 4$ 12 The length is 12 cm.

Step 6. Check.  $P = 2L + 2W$ $32 \stackrel{?}{=} 2 \cdot 12 + 2 \cdot 4$ $32 = 32\checkmark$	
Step 7. Answer the question.	The length is 12 cm and the width is 4 cm.

Try It 3.5.25

The length of a rectangle is eight more than twice the width. The perimeter is 64. Find the length and width.

Answer

24, 8

Try It 3.5.26

The width of a rectangle is six less than twice the length. The perimeter is 18. Find the length and width.


Answer

5, 4

Example 3.5.14

The perimeter of a rectangular swimming pool is 150 feet. The length is 15 feet more than the width. Find the length and width.

Solution

Step 1. Read the problem. Draw the figure and label it with the given information.	 $P = 150$ ft
Step 2. Identify what you are looking for.	the length and the width of the pool
Step 3. Name. Choose a variable to represent the width. The length is 15 feet more than the width.	Let W = width $W + 15$ = length
Step 4. Translate	
Write the appropriate formula.	$P = 2L + 2W$
Substitute.	$150 = 2(W + 15) + 2W$

Step 5. Solve the equation.	$150 = 2W + 30 + 2W$ $150 = 4W + 30$ $120 = 4W$ $30 = W$ (the width of the pool) $W + 15$ (the length of the pool) $30 + 15$ 45
Step 6. Check. $P = 2L + 2W$ $150 \stackrel{?}{=} 2(45) + 2(30)$ $150 = 150\checkmark$	
Step 7. Answer the question.	The length of the pool is 45 feet and the width is 30 feet.

? Try It 3.5.27

The perimeter of a rectangular swimming pool is 200 feet. The length is 40 feet more than the width. Find the length and width.

Answer

70 feet, 30 feet

? Try It 3.5.28

The length of a rectangular garden is 30 yards more than the width. The perimeter is 300 yards. Find the length and width.

Answer

90 yards, 60 yards

Key Concepts

- **Problem-Solving Strategy for Geometry Applications**
  1. **Read** the problem and make all the words and ideas are understood. Draw the figure and label it with the given information.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for by choosing a variable to represent it.
  4. **Translate** into an equation by writing the appropriate formula or model for the situation. Substitute in the given information.
  5. **Solve** the equation using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.
- **Triangle Properties For △ABC**
  Angle measures:
  - $m\angle A + m\angle B + m\angle C = 180$

  Perimeter:
  - $P = a + b + c$

  Area:
  - $A = \frac{1}{2}bh$, b=base,h=height

  A right triangle has one 90° angle.
- **The Pythagorean Theorem** In any right triangle, $a^2 + b^2 = c^2$ where c is the length of the hypotenuse and a and b are the lengths of the legs.

- **Properties of Rectangles**
  - Rectangles have four sides and four right (90°) angles.
  - The lengths of opposite sides are equal.
  - The perimeter of a rectangle is the sum of twice the length and twice the width: $P = 2L + 2W$.
  - The area of a rectangle is the length times the width: $A = LW$.

This page titled 3.5: Triangles, Rectangles, and the Pythagorean Theorem is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.4: Triangles, Rectangles, and the Pythagorean Theorem** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.5E: Exercises

Practice Makes Perfect

Solving Applications Using Triangle Properties

In the following exercises, solve using triangle properties.

Exercise 3.5E.1

The measures of two angles of a triangle are 26 and 98 degrees. Find the measure of the third angle.

Answer

56 degrees

Exercise 3.5E.2

The measures of two angles of a triangle are 61 and 84 degrees. Find the measure of the third angle.

Exercise 3.5E.3

The measures of two angles of a triangle are 105 and 31 degrees. Find the measure of the third angle.

Answer

44 degrees

Exercise 3.5E.4

The measures of two angles of a triangle are 47 and 72 degrees. Find the measure of the third angle.

Exercise 3.5E.5

The perimeter of a triangular pool is 36 yards. The lengths of two sides are 10 yards and 15 yards. How long is the third side?

Answer

11 feet

Exercise 3.5E.6

A triangular courtyard has perimeter 120 meters. The lengths of two sides are 30 meters and 50 meters. How long is the third side?

Exercise 3.5E.7

If a triangle has sides 6 feet and 9 feet and the perimeter is 23 feet, how long is the third side?

Answer

8 feet

Exercise 3.5E.8

If a triangle has sides 14 centimeters and 18 centimeters and the perimeter is 49 centimeters, how long is the third side?

Exercise 3.5E.9

A triangular flag has base one foot and height 1.5 foot. What is its area?

Answer

0.75 sq. ft.

Exercise 3.5E.10

A triangular window has base eight feet and height six feet. What is its area?

Exercise 3.5E.11

What is the base of a triangle with area 207 square inches and height 18 inches?

Answer

23 inches

Exercise 3.5E.12

What is the height of a triangle with area 893 square inches and base 38 inches?

Exercise 3.5E.13

One angle of a right triangle measures 33 degrees. What is the measure of the other small angle?

Answer

57

Exercise 3.5E.14

One angle of a right triangle measures 51 degrees. What is the measure of the other small angle?

Exercise 3.5E.15

One angle of a right triangle measures 22.5 degrees. What is the measure of the other small angle?

Answer

67.5

Exercise 3.5E.16

One angle of a right triangle measures 36.5 degrees. What is the measure of the other small angle?

Exercise 3.5E.17

The perimeter of a triangle is 39 feet. One side of the triangle is one foot longer than the second side. The third side is two feet longer than the second side. Find the length of each side.

Answer

13 ft., 12 ft., 14 ft.

Exercise 3.5E.18

The perimeter of a triangle is 35 feet. One side of the triangle is five feet longer than the second side. The third side is three feet longer than the second side. Find the length of each side.

Exercise 3.5E.19

One side of a triangle is twice the shortest side. The third side is five feet more than the shortest side. The perimeter is 17 feet. Find the lengths of all three sides.

Answer

3 ft., 6 ft., 8 ft.

Exercise 3.5E.20

One side of a triangle is three times the shortest side. The third side is three feet more than the shortest side. The perimeter is 13 feet. Find the lengths of all three sides.

Exercise 3.5E.21

The two smaller angles of a right triangle have equal measures. Find the measures of all three angles.

Answer

$45^\circ, 45^\circ, 90^\circ$

Exercise 3.5E.22

The measure of the smallest angle of a right triangle is 20° less than the measure of the next larger angle. Find the measures of all three angles.

Exercise 3.5E.23

The angles in a triangle are such that one angle is twice the smallest angle, while the third angle is three times as large as the smallest angle. Find the measures of all three angles.

Answer

$30^\circ, 60^\circ, 90^\circ$


Exercise 3.5E.24

The angles in a triangle are such that one angle is 20° more than the smallest angle, while the third angle is three times as large as the smallest angle. Find the measures of all three angles.

Use the Pythagorean Theorem

In the following exercises, use the Pythagorean Theorem to find the length of the hypotenuse.


Exercise 3.5E. 25




Answer

15

Exercise 3.5E. 26




Exercise 3.5E. 27



Answer


25

Exercise 3.5E. 28



In the following exercises, use the Pythagorean Theorem to find the length of the leg. Round to the nearest tenth, if necessary.


Exercise 3.5E. 29




Answer

8

Exercise 3.5E.30




Exercise 3.5E.31




Answer

12

Exercise 3.5E.32




Exercise 3.5E.33




Answer

10.2

Exercise 3.5E.34




Exercise 3.5E.35



Answer

9.8


Exercise 3.5E.36



In the following exercises, solve using the Pythagorean Theorem. Approximate to the nearest tenth, if necessary.

Exercise 3.5E.37

A 13-foot string of lights will be attached to the top of a 12-foot pole for a holiday display, as shown below. How far from the base of the pole should the end of the string of lights be anchored?



Answer

5 feet

Exercise 3.5E.38

Pam wants to put a banner across her garage door, as shown below, to congratulate her son for his college graduation. The garage door is 12 feet high and 16 feet wide. How long should the banner be to fit the garage door?



Exercise 3.5E. 39

Chi is planning to put a path of paving stones through her flower garden, as shown below. The flower garden is a square with side 10 feet. What will the length of the path be?



Answer

14.1 feet

Exercise 3.5E. 40

Brian borrowed a 20 foot extension ladder to use when he paints his house. If he sets the base of the ladder 6 feet from the house, as shown below, how far up will the top of the ladder reach?



Solve Applications Using Rectangle Properties

In the following exercises, solve using rectangle properties.

Exercise 3.5E. 41

The length of a rectangle is 85 feet and the width is 45 feet. What is the perimeter?

Answer

260 feet

Exercise 3.5E. 42

The length of a rectangle is 26 inches and the width is 58 inches. What is the perimeter?

Exercise 3.5E. 43

A rectangular room is 15 feet wide by 14 feet long. What is its perimeter?

Answer

58 feet

Exercise 3.5E.44

A driveway is in the shape of a rectangle 20 feet wide by 35 feet long. What is its perimeter?

Exercise 3.5E.45

The area of a rectangle is 414 square meters. The length is 18 meters. What is the width?

Answer

23 meters

Exercise 3.5E.46

The area of a rectangle is 782 square centimeters. The width is 17 centimeters. What is the length?

Exercise 3.5E.47

The width of a rectangular window is 24 inches. The area is 624 square inches. What is the length?

Answer

26 inches

Exercise 3.5E.48

The length of a rectangular poster is 28 inches. The area is 1316 square inches. What is the width?

Exercise 3.5E.49

Find the length of a rectangle with perimeter 124 and width 38.

Answer

24

Exercise 3.5E.50

Find the width of a rectangle with perimeter 92 and length 19.

Exercise 3.5E.51

Find the width of a rectangle with perimeter 16.2 and length 3.2.

Answer

4.9

Exercise 3.5E.52

Find the length of a rectangle with perimeter 20.2 and width 7.8.

Exercise 3.5E.53

The length of a rectangle is nine inches more than the width. The perimeter is 46 inches. Find the length and the width.

Answer

16 in., 7 in.

Exercise 3.5E.54

The width of a rectangle is eight inches more than the length. The perimeter is 52 inches. Find the length and the width.

Exercise 3.5E.55

The perimeter of a rectangle is 58 meters. The width of the rectangle is five meters less than the length. Find the length and the width of the rectangle.

Answer

17 m, 12 m

Exercise 3.5E.56

The perimeter of a rectangle is 62 feet. The width is seven feet less than the length. Find the length and the width.

Exercise 3.5E.57

The width of the rectangle is 0.7 meters less than the length. The perimeter of a rectangle is 52.6 meters. Find the dimensions of the rectangle.

Answer

13.5 m length, 12.8 m width

Exercise 3.5E.58

The length of the rectangle is 1.1 meters less than the width. The perimeter of a rectangle is 49.4 meters. Find the dimensions of the rectangle.

Exercise 3.5E.59

The perimeter of a rectangle is 150 feet. The length of the rectangle is twice the width. Find the length and width of the rectangle.

Answer

50 ft., 25 ft.

Exercise 3.5E.60

The length of a rectangle is three times the width. The perimeter of the rectangle is 72 feet. Find the length and width of the rectangle.

Exercise 3.5E.61

The length of a rectangle is three meters less than twice the width. The perimeter of the rectangle is 36 meters. Find the dimensions of the rectangle.

Answer

7 m width, 11 m length

Exercise 3.5E.62

The length of a rectangle is five inches more than twice the width. The perimeter is 34 inches. Find the length and width.

Exercise 3.5E.63

The perimeter of a rectangular field is 560 yards. The length is 40 yards more than the width. Find the length and width of the field.

Answer

160 yd., 120 yd.

Exercise 3.5E.64

The perimeter of a rectangular atrium is 160 feet. The length is 16 feet more than the width. Find the length and width of the atrium.

Exercise 3.5E.65

A rectangular parking lot has perimeter 250 feet. The length is five feet more than twice the width. Find the length and width of the parking lot.

Answer

85 ft., 40 ft.

Exercise 3.5E.66

A rectangular rug has perimeter 240 inches. The length is 12 inches more than twice the width. Find the length and width of the rug.

Everyday Math

Exercise 3.5E.67

Christa wants to put a fence around her triangular flowerbed. The sides of the flowerbed are six feet, eight feet and 10 feet. How many feet of fencing will she need to enclose her flowerbed?

Answer

24 feet

Exercise 3.5E.68

Jose just removed the children's playset from his back yard to make room for a rectangular garden. He wants to put a fence around the garden to keep out the dog. He has a 50 foot roll of fence in his garage that he plans to use. To fit in the backyard, the width of the garden must be 10 feet. How long can he make the other length?

Writing Exercises

Exercise 3.5E.69

If you need to put tile on your kitchen floor, do you need to know the perimeter or the area of the kitchen? Explain your reasoning.

Answer


area; answers will vary

Exercise 3.5E.70

If you need to put a fence around your backyard, do you need to know the perimeter or the area of the backyard? Explain your reasoning.

Exercise 3.5E.71

Look at the two figures below.



1. Which figure looks like it has the larger area?
2. Which looks like it has the larger perimeter?
3. Now calculate the area and perimeter of each figure.
4. Which has the larger area?
5. Which has the larger perimeter?

Answer

1. Answers will vary.
2. Answers will vary.
3. Answers will vary.
4. The areas are the same.
5. The 2x8 rectangle has a larger perimeter than the 4x4 square.

Exercise 3.5E.72

Write a geometry word problem that relates to your life experience, then solve it and explain all your steps.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve applications using triangle properties.			
use the Pythagorean Theorem.			
solve applications using rectangle properties.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 3.5E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.4E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.6: Solve Uniform Motion Applications

Learning Objectives

By the end of this section, you will be able to:

- Solve uniform motion applications

Note

Before you get started, take this readiness quiz.

1. Find the distance traveled by a car going 70 miles per hour for 3 hours.
   If you missed this problem, review Exercise 2.6.1.
2. Solve $x+1.2(x-10)=98$.
   If you missed this problem, review Exercise 2.4.7.
3. Convert 90 minutes to hours.
   If you missed this problem, review Exercise 1.11.1.

Solve Uniform Motion Applications

When planning a road trip, it often helps to know how long it will take to reach the destination or how far to travel each day. We would use the distance, rate, and time formula, D=rt, which we have already seen.

In this section, we will use this formula in situations that require a little more algebra to solve than the ones we saw earlier. Generally, we will be looking at comparing two scenarios, such as two vehicles traveling at different rates or in opposite directions. When the speed of each vehicle is constant, we call applications like this *uniform motion problems*.

Our problem-solving strategies will still apply here, but we will add to the first step. The first step will include drawing a diagram that shows what is happening in the example. Drawing the diagram helps us understand what is happening so that we will write an appropriate equation. Then we will make a table to organize the information, like we did for the money applications.

The steps are listed here for easy reference:

USE A PROBLEM-SOLVING STRATEGY IN DISTANCE, RATE, AND TIME APPLICATIONS.

1. **Read** the problem. Make sure all the words and ideas are understood.
   - Draw a diagram to illustrate what it happening.
   - Create a table to organize the information.
   - Label the columns rate, time, distance.
   - List the two scenarios.
   - Write in the information you know.

	Rate	• Time	= Distance

2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
   - Complete the chart.
   - Use variable expressions to represent that quantity in each row.
   - Multiply the rate times the time to get the distance.
4. **Translate** into an equation.
   - Restate the problem in one sentence with all the important information.
   - Then, translate the sentence into an equation.
5. **Solve** the equation using good algebra techniques.

6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.


Example 3.6.1

An express train and a local train leave Pittsburgh to travel to Washington, D.C. The express train can make the trip in 4 hours and the local train takes 5 hours for the trip. The speed of the express train is 12 miles per hour faster than the speed of the local train. Find the speed of both trains.

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.

Draw a diagram to illustrate what it happening. Shown below is a sketch of what is happening in the example.



	Rate (mph)	•	Time (hrs)	=	Distance (miles)
Express			4		
Local			5		

Create a table to organize the information. Label the columns “Rate,” “Time,” and “Distance.” List the two scenarios. Write in the information you know.

Step 2. Identify what we are looking for.

We are asked to find the speed of both trains. Notice that the distance formula uses the word “rate,” but it is more common to use “speed” when we talk about vehicles in everyday English.

Step 3. Name what we are looking for. Choose a variable to represent that quantity.

Complete the chart Use variable expressions to represent that quantity in each row. We are looking for the speed of the trains. Let’s let r represent the speed of the local train. Since the speed of the express train is 12 mph faster, we represent that as r+12.

$$\begin{aligned} r &= \text{speed of the local train} \\ r+12 &= \text{speed of the express train} \end{aligned}$$

Fill in the speeds into the chart.

	Rate (mph)	•	Time (hrs)	=	Distance (miles)
Express	$r+12$		4		
Local	r		5		

Multiply the rate times the time to get the distance.

	Rate (mph)	•	Time (hrs)	=	Distance (miles)
Express	$r+12$		4		$4(r+12)$
Local	r		5		$5r$

Step 4. Translate into an equation.

Restate the problem in one sentence with all the important information. Then, translate the sentence into an equation.

	distance traveled by express train	=	distance traveled by local train
Translate to an equation.	$4(r+12)$	=	$5r$

- The equation to model this situation will come from the relation between the distances. Look at the diagram we drew above. How is the distance traveled by the express train related to the distance traveled by the local train?
- Since both trains leave from Pittsburgh and travel to Washington, D.C. they travel the same distance. So we write:

Step 5. Solve the equation using good algebra techniques.

Now solve this equation.	$4(r+12)=5r$ $4r+48=5r$ $48=r$ So the speed of the local train is 48 mph.
Find the speed of the express train.	$r+12$ $48+12$ 60 he speed of the express train is 60 mph.

Step 6. Check the answer in the problem and make sure it makes sense. \[\begin{array}{ll}{\text { express train }} & {60 \mathrm{mph}(4 \text { hours })=240 \mathrm{miles}} \\ {\text { local train }} & {48 \mathrm{mph}(5 \text { hours })=240 \mathrm{miles} \checkmark \end{array}\]

Step 7. Answer the question with a complete sentence.

The speed of the local train is 48 mph and the speed of the express train is 60 mph.

Try It 3.6.2

Wayne and Dennis like to ride the bike path from Riverside Park to the beach. Dennis's speed is seven miles per hour faster than Wayne's speed, so it takes Wayne 2 hours to ride to the beach while it takes Dennis 1.5 hours for the ride. Find the speed of both bikers.

Answer

Wayne 21 mph, Dennis 28 mph

Try It 3.6.3

Jeromy can drive from his house in Cleveland to his college in Chicago in 4.5 hours. It takes his mother 6 hours to make the same drive. Jeromy drives 20 miles per hour faster than his mother. Find Jeromy's speed and his mother's speed.

Answer

Jeromy 80 mph, mother 60 mph

In Exercise 3.6.4, the last example, we had two trains traveling the same distance. The diagram and the chart helped us write the equation we solved. Let's see how this works in another case.

Example 3.6.4

Christopher and his parents live 115 miles apart. They met at a restaurant between their homes to celebrate his mother's birthday. Christopher drove 1.5 hours while his parents drove 1 hour to get to the restaurant. Christopher's average speed was 10 miles per hour faster than his parents' average speed. What were the average speeds of Christopher and of his parents as they drove to the restaurant?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.

Draw a diagram to illustrate what it happening. Below shows a sketch of what is happening in the example.



Create a table to organize the information.

Label the columns rate, time, distance.

List the two scenarios.

Write in the information you know.

	Rate (mph)	•	Time (hrs)	=	Distance (miles)
Christopher			1.5		
Parents			1		
					115

Step 2. Identify what we are looking for.

We are asked to find the average speeds of Christopher and his parents.

Step 3. Name what we are looking for. Choose a variable to represent that quantity.

Complete the chart.

Use variable expressions to represent that quantity in each row.

We are looking for their average speeds. Let's let r represent the average speed of the parents. Since the Christopher's speed is 10 mph faster, we represent that as r+10.

Fill in the speeds into the chart.

	Rate (mph)	•	Time (hrs)	=	Distance (miles)
Christopher	$r+10$		1.5		$1.5(r+10)$
Parents	r		1		r
					115

Multiply the rate times the time to get the distance.

Step 4. Translate into an equation.

Restate the problem in one sentence with all the important information. Then, translate the sentence into an equation. Again, we need to identify a relationship between the distances in order to write an equation. Look at the diagram we created above and notice the relationship between the distance Christopher traveled and the distance his parents traveled.

The distance Christopher travelled plus the distance his parents travel must add up to 115 miles. So we write:

	distance traveled by Christopher	+	distance traveled by his parents	= 115
Translate to an equation.	$1.5(r+10)$	+	r	$=115$

Step 5. Solve the equation using good algebra techniques.

	$1.5(r+10)+r=115$
	$1.5r+15+r=115$
Now solve this equation.	$2.5r+15=115$
	$2.5r=100$
	$r=40$
	so the parents' speed was 40 mph.
	$r+10$
Christopher's speed is r + 10	$40+10$
	50
	Christopher's speed was 50 mph.

Step 6. Check the answer in the problem and make sure it makes sense.

Christopher drove	50 mph (1.5 hours)	=	75 miles
His parents drove	40 mph (1 hour)	=	40 miles
			115 miles

Step 7. Answer the question with a complete sentence.

Christopher's speed was 50 mph.
His parents' speed was 40 mph.

? Try It 3.6.5

Carina is driving from her home in Anaheim to Berkeley on the same day her brother is driving from Berkeley to Anaheim, so they decide to meet for lunch along the way in Buttonwillow. The distance from Anaheim to Berkeley is 410 miles. It takes Carina 3 hours to get to Buttonwillow, while her brother drives 4 hours to get there. The average speed Carina's brother drove was 15 miles per hour faster than Carina's average speed. Find Carina's and her brother's average speeds.

Answer

Carina 50 mph, brother 65 mph

? Try It 3.6.6

Ashley goes to college in Minneapolis, 234 miles from her home in Sioux Falls. She wants her parents to bring her more winter clothes, so they decide to meet at a restaurant on the road between Minneapolis and Sioux Falls. Ashley and her parents both drove 2 hours to the restaurant. Ashley's average speed was seven miles per hour faster than her parents' average speed. Find Ashley's and her parents' average speed.

Answer

parents 55 mph, Ashley 62 mph

As you read the next example, think about the relationship of the distances traveled. Which of the previous two examples is more similar to this situation?

✓ Example 3.6.7

Two truck drivers leave a rest area on the interstate at the same time. One truck travels east and the other one travels west. The truck traveling west travels at 70 mph and the truck traveling east has an average speed of 60 mph. How long will they travel before they are 325 miles apart?

Solution

Step 1. Read the problem. Make sure all the words and ideas are understood.

Draw a diagram to illustrate what it happening.



Create a table to organize the information.

	Rate (mph)	• Time (hrs)	= Distance (miles)
West	70		
East	60		
			325

Step 2. Identify what we are looking for.

We are asked to find the amount of time the trucks will travel until they are 325 miles apart.

Step 3. Name what we are looking for. Choose a variable to represent that quantity.

We are looking for the time traveled. Both trucks will travel the same amount of time. Let's call the time t. Since their speeds are different, they will travel different distances. Complete the chart.

	Rate (mph)	• Time (hrs)	= Distance (miles)
West	70	t	$70t$
East	60	t	$60t$
			325

Step 4. Translate into an equation.

We need to find a relation between the distances in order to write an equation. Looking at the diagram, what is the relationship between the distance each of the trucks will travel? The distance traveled by the truck going west plus the distance traveled by the truck going east must add up to 325 miles. So we write:

	distance traveled by westbound truck	+	distance traveled by eastbound truck	= 32
Translate to an equation.	70t	+	60t	= 32

Step 5. Solve the equation using good algebra techniques.

$$\begin{aligned} \text{Now solve this equation.} \quad 70t+60t &= 325 \\ 130t &= 325 \\ t &= 2.5 \end{aligned} \tag{3.6.1}$$

Step 6. Check the answer in the problem and make sure it makes sense.

$$\begin{aligned} \text{Truck going West} \quad 70 \text{ mph } (2.5 \text{ hours}) &= 175 \text{ miles} \\ \text{Truck going East} \quad 60 \text{ mph } (2.5 \text{ hour}) &= \underline{150 \text{ miles}} \\ & \quad 325 \text{ miles} \end{aligned}$$

Step 7. Answer the question with a complete sentence. It will take the truck 2.5 hours to be 325 miles apart.

? Try It 3.6.8

Pierre and Monique leave their home in Portland at the same time. Pierre drives north on the turnpike at a speed of 75 miles per hour while Monique drives south at a speed of 68 miles per hour. How long will it take them to be 429 miles apart?

Answer

3 hours

Try It 3.6.9

Thanh and Nhat leave their office in Sacramento at the same time. Thanh drives north on I-5 at a speed of 72 miles per hour. Nhat drives south on I-5 at a speed of 76 miles per hour. How long will it take them to be 330 miles apart?

Answer

2.2 hours

MATCHING UNITS IN PROBLEMS


It is important to make sure the units match when we use the distance rate and time formula. For instance, if the rate is in miles per hour, then the time must be in hours.

Example 3.6.10

When Katie Mae walks to school, it takes her 30 minutes. If she rides her bike, it takes her 15 minutes. Her speed is three miles per hour faster when she rides her bike than when she walks. What are her walking speed and her speed riding her bike?

Solution

First, we draw a diagram that represents the situation to help us see what is happening.



We are asked to find her speed walking and riding her bike. Let's call her walking speed r. Since her biking speed is three miles per hour faster, we will call that speed r+3. We write the speeds in the chart.

The speed is in miles per hour, so we need to express the times in hours, too, in order for the units to be the same. Remember, one hour is 60 minutes. So:

$$\begin{aligned} 30 \text{ minutes is } \tfrac{30}{60} \text{ or } \tfrac{1}{2} \text{ hour} \\ 15 \text{ minutes is } \tfrac{15}{60} \text{ or } \tfrac{1}{4} \text{ hour} \end{aligned} \qquad (3.6.2)$$

Next, we multiply rate times time to fill in the distance column.

	Rate (mph)	· Time (hrs)	= Distance (miles)
Walk	r	$\frac{1}{2}$	$\frac{1}{2}r$
Bike	$r+3$	$\frac{1}{4}$	$\frac{1}{4}(r+3)$

The equation will come from the fact that the distance from Katie Mae's home to her school is the same whether she is walking or riding her bike.

So we say:

	distance walked = distance covered by bike
Translate into an equation.	$\frac{1}{2}r = \frac{1}{4}(r+3)$
Solve this equation.	$\frac{1}{2}r = \frac{1}{4}(r+3)$
Clear the fractions by multiplying by the LCD of all the fractions in the equation.	$8 \cdot \frac{1}{2}r = 8 \cdot \frac{1}{4}(r+3)$
Simplify.	$4r = 2(r+3)$ $4r = 2r + 6$ $2r = 6$ $r = 3$ mph (Katie Mae's walking speed) $r + 3$ biking speed $3 + 3$ 6 mph (Katie Mae's biking speed)
Let's check if this works. Walk 3 mph (0.5 hour) = 1.5 miles Bike 6 mph (0.25 hour) = 1.5 miles	
Yes, either way Katie Mae travels 1.5 miles to school.	Katie Mae's walking speed is 3 mph. Her speed riding her bike is 6 mph.

Try It 3.6.11

Suzy takes 50 minutes to hike uphill from the parking lot to the lookout tower. It takes her 30 minutes to hike back down to the parking lot. Her speed going downhill is 1.2 miles per hour faster than her speed going uphill. Find Suzy's uphill and downhill speeds.

Answer

uphill 1.8 mph, downhill three mph

Try It 3.6.12

Llewyn takes 45 minutes to drive his boat upstream from the dock to his favorite fishing spot. It takes him 30 minutes to drive the boat back downstream to the dock. The boat's speed going downstream is four miles per hour faster than its speed going upstream. Find the boat's upstream and downstream speeds.

Answer

upstream 8 mph, downstream 12 mph

In the distance, rate, and time formula, time represents the actual amount of **elapsed time** (in hours, minutes, etc.). If a problem gives us starting and ending times as clock times, we must find the elapsed time in order to use the formula.

Example 3.6.13

Hamilton loves to travel to Las Vegas, 255 miles from his home in Orange County. On his last trip, he left his house at 2:00 pm. The first part of his trip was on congested city freeways. At 4:00 pm, the traffic cleared and he was able to drive through the desert at a speed 1.75 times as fast as when he drove in the congested area. He arrived in Las Vegas at 6:30 pm. How fast was he driving during each part of his trip?

Solution

A diagram will help us model this trip.



Next, we create a table to organize the information.

We know the total distance is 255 miles. We are looking for the rate of speed for each part of the trip. The rate in the desert is 1.75 times the rate in the city. If we let r= the rate in the city, then the rate in the desert is 1.75r.

The times here are given as clock times. Hamilton started from home at 2:00 pm and entered the desert at 4:30 pm. So he spent two hours driving the congested freeways in the city. Then he drove faster from 4:00 pm until 6:30 pm in the desert. So he drove 2.5 hours in the desert.

Now, we multiply the rates by the times.

	Rate (mph)	• Time (hrs)	= Distance (miles)
City	r	2	$2r$
Desert	$1.75r$	2.5	$2.5(1.75r)$
			255

By looking at the diagram below, we can see that the sum of the distance driven in the city and the distance driven in desert is 255 miles.

	distance driven in the city + distance driven in desert = 255
Translate into an equation.	$2r + 2.5(1.75r) = 255$
Solve this equation.	$2r + 2.5(1.75r) = 255$ $2r + 4.375r = 255$ $6.375r = 255$ $r = 40$ mph city $1.75r$ desert speed $1.75(40)$ 70 mph
Check. City 40 mph (2 hours) = 80 miles Desert 70 mph (2.5 hours) = 175 miles 255 miles	
	Hamilton drove 40 mph in the city and 70 mph in the desert.

Try It 3.6.14

Cruz is training to compete in a triathlon. He left his house at 6:00 and ran until 7:30. Then he rode his bike until 9:45. He covered a total distance of 51 miles. His speed when biking was 1.6 times his speed when running. Find Cruz's biking and running speeds.

Answer

biking 16 mph, running 10 mph

Try It 3.6.15

Phuong left home on his bicycle at 10:00. He rode on the flat street until 11:15, then rode uphill until 11:45. He rode a total of 31 miles. His speed riding uphill was 0.6 times his speed on the flat street. Find his speed biking uphill and on the flat street.

Answer

uphill 12 mph, flat street 20 mph

Key Concepts

- **Distance, Rate, and Time**
  - $D = rt$ where D = distance, r = rate, t = time
- **Problem-Solving Strategy—Distance, Rate, and Time Applications**
  1. **Read** the problem. Make sure all the words and ideas are understood.
     Draw a diagram to illustrate what it happening.
     Create a table to organize the information: Label the columns rate, time, distance. List the two scenarios. Write in the information you know.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose a variable to represent that quantity.
     Complete the chart.
     Use variable expressions to represent that quantity in each row.
     Multiply the rate times the time to get the distance.
  4. **Translate** into an equation.
     Restate the problem in one sentence with all the important information.
     Then, translate the sentence into an equation.
  5. **Solve** the equation using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.

This page titled 3.6: Solve Uniform Motion Applications is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.5: Solve Uniform Motion Applications** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.6E: Exercises

Practice Makes Perfect

Solve Uniform Motion Applications

In the following exercises, solve.

Exercise 3.6E.1

Lilah is moving from Portland to Seattle. It takes her three hours to go by train. Mason leaves the train station in Portland and drives to the train station in Seattle with all Lilah's boxes in his car. It takes him 2.4 hours to get to Seattle, driving at 15 miles per hour faster than the speed of the train. Find Mason's speed and the speed of the train.

Answer

Mason 75 mph, train 60 mph

Exercise 3.6E.2

Kathy and Cheryl are walking in a fundraiser. Kathy completes the course in 4.8 hours and Cheryl completes the course in 8 hours. Kathy walks two miles per hour faster than Cheryl. Find Kathy's speed and Cheryl's speed.

Exercise 3.6E.3

Two busses go from Sacramento for San Diego. The express bus makes the trip in 6.8 hours and the local bus takes 10.2 hours for the trip. The speed of the express bus is 25 mph faster than the speed of the local bus. Find the speed of both busses.

Answer

express bus 75mph, local 50 mph

Exercise 3.6E.4

A commercial jet and a private airplane fly from Denver to Phoenix. It takes the commercial jet 1.1 hours for the flight, and it takes the private airplane 1.8 hours. The speed of the commercial jet is 210 miles per hour faster than the speed of the private airplane. Find the speed of both airplanes.

Exercise 3.6E.5

Saul drove his truck 3 hours from Dallas towards Kansas City and stopped at a truck stop to get dinner. At the truck stop he met Erwin, who had driven 4 hours from Kansas City towards Dallas. The distance between Dallas and Kansas City is 542 miles, and Erwin's speed was eight miles per hour slower than Saul's speed. Find the speed of the two truckers.

Answer

Saul 82 mph, Erwin 74 mph

Exercise 3.6E.6

Charlie and Violet met for lunch at a restaurant between Memphis and New Orleans. Charlie had left Memphis and drove 4.8 hours towards New Orleans. Violet had left New Orleans and drove 2 hours towards Memphis, at a speed 10 miles per hour faster than Charlie's speed. The distance between Memphis and New Orleans is 394 miles. Find the speed of the two drivers.

Exercise 3.6E.7

Sisters Helen and Anne live 332 miles apart. For Thanksgiving, they met at their other sister's house partway between their homes. Helen drove 3.2 hours and Anne drove 2.8 hours. Helen's average speed was four miles per hour faster than Anne's. Find Helen's average speed and Anne's average speed.

Answer

Helen 60 mph, Anne 56 mph

Exercise 3.6E.8

Ethan and Leo start riding their bikes at the opposite ends of a 65-mile bike path. After Ethan has ridden 1.5 hours and Leo has ridden 2 hours, they meet on the path. Ethan's speed is six miles per hour faster than Leo's speed. Find the speed of the two bikers.

Exercise 3.6E.9

Elvira and Aletheia live 3.1 miles apart on the same street. They are in a study group that meets at a coffee shop between their houses. It took Elvira half an hour and Aletheia two-thirds of an hour to walk to the coffee shop. Aletheia's speed is 0.6 miles per hour slower than Elvira's speed. Find both women's walking speeds.

Answer

Aletheia 2.4 mph, Elvira 3 mph

Exercise 3.6E.10

DaMarcus and Fabian live 23 miles apart and play soccer at a park between their homes. DaMarcus rode his bike for three-quarters of an hour and Fabian rode his bike for half an hour to get to the park. Fabian's speed was six miles per hour faster than DaMarcus' speed. Find the speed of both soccer players.

Exercise 3.6E.11

Cindy and Richard leave their dorm in Charleston at the same time. Cindy rides her bicycle north at a speed of 18 miles per hour. Richard rides his bicycle south at a speed of 14 miles per hour. How long will it take them to be 96 miles apart?

Answer

3 hours

Exercise 3.6E.12

Matt and Chris leave their uncle's house in Phoenix at the same time. Matt drives west on I-60 at a speed of 76 miles per hour. Chris drives east on I-60 at a speed of 82 miles per hour. How many hours will it take them to be 632 miles apart?

Exercise 3.6E.13

Two busses leave Billings at the same time. The Seattle bus heads west on I-90 at a speed of 73 miles per hour while the Chicago bus heads east at a speed of 79 miles an hour. How many hours will it take them to be 532 miles apart?

Answer

3.5 hours

Exercise 3.6E.14

Two boats leave the same dock in Cairo at the same time. One heads north on the Mississippi River while the other heads south. The northbound boat travels four miles per hour. The southbound boat goes eight miles per hour. How long will it take them to be 54 miles apart?

Exercise 3.6E.15

Lorena walks the path around the park in 30 minutes. If she jogs, it takes her 20 minutes. Her jogging speed is 1.5 miles per hour faster than her walking speed. Find Lorena's walking speed and jogging speed.

Answer

walking 3 mph, jogging 4.5 mph

Exercise 3.6E.16

Julian rides his bike uphill for 45 minutes, then turns around and rides back downhill. It takes him 15 minutes to get back to where he started. His uphill speed is 3.2 miles per hour slower than his downhill speed. Find Julian's uphill and downhill speed.

Exercise 3.6E.17

Cassius drives his boat upstream for 45 minutes. It takes him 30 minutes to return downstream. His speed going upstream is three miles per hour slower than his speed going downstream. Find his upstream and downstream speeds.

Answer

upstream 6 mph, downstream 9 mph

Exercise 3.6E.18

It takes Darline 20 minutes to drive to work in light traffic. To come home, when there is heavy traffic, it takes her 36 minutes. Her speed in light traffic is 24 miles per hour faster than her speed in heavy traffic. Find her speed in light traffic and in heavy traffic.

Exercise 3.6E.19

At 1:30 Marlon left his house to go to the beach, a distance of 7.6 miles. He rode his skateboard until 2:15, then walked the rest of the way. He arrived at the beach at 3:00. Marlon's speed on his skateboard is 2.5 times his walking speed. Find his speed when skateboarding and when walking.

Answer

skateboarding 8 mph, walking 3.2 mph

Exercise 3.6E.20

Aaron left at 9:15 to drive to his mountain cabin 108 miles away. He drove on the freeway until 10:45, and then he drove on the mountain road. He arrived at 11:05. His speed on the freeway was three times his speed on the mountain road. Find Aaron's speed on the freeway and on the mountain road.

Exercise 3.6E.21

Marisol left Los Angeles at 2:30 to drive to Santa Barbara, a distance of 95 miles. The traffic was heavy until 3:20. She drove the rest of the way in very light traffic and arrived at 4:20. Her speed in heavy traffic was 40 miles per hour slower than her speed in light traffic. Find her speed in heavy traffic and in light traffic.

Answer

heavy traffic 30 mph, light traffic 70 mph

Exercise 3.6E.22

Lizette is training for a marathon. At 7:00 she left her house and ran until 8:15, then she walked until 11:15. She covered a total distance of 19 miles. Her running speed was five miles per hour faster than her walking speed. Find her running and walking speeds.

Everyday Math

Exercise 3.6E.23

John left his house in Irvine at 8:35 am to drive to a meeting in Los Angeles, 45 miles away. He arrived at the meeting at 9:50. At 3:30 pm, he left the meeting and drove home. He arrived home at 5:18.

1. What was his average speed on the drive from Irvine to Los Angeles?
2. What was his average speed on the drive from Los Angeles to Irvine?
3. What was the total time he spent driving to and from this meeting?
4. John drove a total of 90 miles roundtrip. Find his average speed. (Round to the nearest tenth.)

Answer

1. 36 mph
2. 25 mph
3. 3.05 hours
4. 29.5 mph

Exercise 3.6E.24

Sarah wants to arrive at her friend's wedding at 3:00. The distance from Sarah's house to the wedding is 95 miles. Based on usual traffic patterns, Sarah predicts she can drive the first 15 miles at 60 miles per hour, the next 10 miles at 30 miles per hour, and the remainder of the drive at 70 miles per hour.

1. How long will it take Sarah to drive the first 15 miles?
2. How long will it take Sarah to drive the next 10 miles?
3. How long will it take Sarah to drive the rest of the trip?
4. What time should Sarah leave her house?

Writing Exercises

Exercise 3.6E.25

When solving a uniform motion problem, how does drawing a diagram of the situation help you?

Answer

Answers will vary.

Exercise 3.6E. 26

When solving a uniform motion problem, how does creating a table help you?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve uniform motion applications.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 3.6E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.5E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.7: Solve Applications with Linear Inequalities

Learning Objectives

By the end of this section, you will be able to:

- Solve applications with linear inequalities

Note

Before you get started, take this readiness quiz.

1. Write as an inequality: x is at least 30.
   If you missed this problem, review Exercise 2.7.34.
2. Solve $8-3y<41$.
   If you missed this problem, review Exercise 2.7.22.

Solve Applications with Linear Inequalities

Many real-life situations require us to solve inequalities. In fact, inequality applications are so common that we often do not even realize we are doing algebra. For example, how many gallons of gas can be put in the car for $20? Is the rent on an apartment affordable? Is there enough time before class to go get lunch, eat it, and return? How much money should each family member's holiday gift cost without going over budget?

The method we will use to solve applications with linear inequalities is very much like the one we used when we solved applications with equations. We will read the problem and make sure all the words are understood. Next, we will identify what we are looking for and assign a variable to represent it. We will restate the problem in one sentence to make it easy to translate into an inequality. Then, we will solve the inequality.

Example 3.7.1

Emma got a new job and will have to move. Her monthly income will be $5,265. To qualify to rent an apartment, Emma's monthly income must be at least three times as much as the rent. What is the highest rent Emma will qualify for?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the highest rent Emma will qualify for
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let r = rent
Step 4. Translate into an inequality. First write a sentence that gives the information to find it.	Emma's monthly income must be at least three times the rent.
Step 5. Solve the inequality. Remember, $a>x$ has the same meaning as $x<a$	$5265 \geq 3r$ $1755 \geq r$ $r \leq 1755$
Step 6. Check the answer in the problem and make sure it makes sense. A maximum rent of $1,755 seems reasonable for an income of $5,265.	
Step 7. Answer the answer in the problem complete sentence.	the question with a The maximum rent is $1,755.

Try It 3.7.2

Alan is loading a pallet with boxes that each weighs 45 pounds. The pallet can safely support no more than 900 pounds. How many boxes can he safely load onto the pallet?

Answer

There can be no more than 20 boxes.

Try It 3.7.3

The elevator in Yehire's apartment building has a sign that says the maximum weight is 2,100 pounds. If the average weight of one person is 150 pounds, how many people can safely ride the elevator?

Answer

A maximum of 14 people can safely ride in the elevator.

Sometimes an application requires the solution to be a whole number, but the algebraic solution to the inequality is not a whole number. In that case, we must round the algebraic solution to a whole number. The context of the application will determine whether we round up or down. To check applications like this, we will round our answer to a number that is easy to compute with and make sure that number makes the inequality true.

Example 3.7.4

Dawn won a mini-grant of $4,000 to buy tablet computers for her classroom. The tablets she would like to buy cost $254.12 each, including tax and delivery. What is the maximum number of tablets Dawn can buy?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the maximum number of tablets Dawn can buy
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let n = the number of tablets.
Step 4. Translate. write a sentence that gives the information to find it.	$254.12 times the number of tablets is no more than $4000.
Translate into an inequality.	$254.12n \leq 4000$
Step 5. Solve the inequality.	$n \leq 15.74$
But n must be a whole number of tablets, so round to 15.	$n \leq 15$
Step 6. Check the answer in the problem and make sure it makes sense. Rounding down the price to $250, 15 tablets would cost $3750, while 16 tablets would be $4000. So a maximum of 15 tablets at $254.12 seems reasonable.	
Step 7. Answer the answer in the problem complete sentence.	the question with a Dawn can buy a maximum of 15 tablets.

Try It 3.7.5

Angie has $20 to spend on juice boxes for her son's preschool picnic. Each pack of juice boxes costs $2.63. What is the maximum number of packs she can buy?

Answer

seven packs

Try It 3.7.6

Daniel wants to surprise his girlfriend with a birthday party at her favorite restaurant. It will cost $42.75 per person for dinner, including tip and tax. His budget for the party is $500. What is the maximum number of people Daniel can have at the party?

Answer

11 people

Example 3.7.7

Pete works at a computer store. His weekly pay will be either a fixed amount, $925, or $500 plus 12% of his total sales. How much should his total sales be for his variable pay option to exceed the fixed amount of $925?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the total sales needed for his variable pay option to exceed the fixed amount of $925
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let s = the total sales.
Step 4. Translate. write a sentence that gives the information to find it. Translate into an inequality. Remember to convert the percent to a decimal.	$500 plus 12% of total sales is more than $925. $500 + 0.12s > 925$
Step 5. Solve the inequality.	$0.12s > 425$ $s > 3541.\overline{66}$
Step 6. Check the answer in the problem and make sure it makes sense. Rounding down the price to $250, 15 tablets would cost $3750, while If we round the total sales up to $4000, we see that $500+0.12(4000) = 980$, which is more than $925.	
Step 7. Answer the the question with a complete sentence.	The total sales must be more than $3541.67

Try It 3.7.8

Tiffany just graduated from college and her new job will pay her $20000 per year plus 2% of all sales. She wants to earn at least $100000 per year. For what total sales will she be able to achieve her goal?

Answer

at least $4000000

Try It 3.7.9

Christian has been offered a new job that pays $24000 a year plus 3% of sales. For what total sales would this new job pay more than his current job which pays $60000?

Answer

at least $1200000

Example 3.7.10

Sergio and Lizeth have a very tight vacation budget. They plan to rent a car from a company that charges $75 a week plus $0.25 a mile. How many miles can they travel and still keep within their $200 budget?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the number of miles Sergio and Lizeth can travel
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let m = the number of miles.
Step 4. Translate. write a sentence that gives the information to find it. Translate into an inequality.	$75 plus 0.25 times the number of miles is less than or equal to $200. $75 + 25m \leq 200$
Step 5. Solve the inequality.	$0.25m \leq 125$ $m \leq 500$ miles
Step 6. Check the answer in the problem and make sure it makes sense. Yes, $75 + 0.25(500) = 200$.	
Step 7. Answer the the question with a complete sentence.	Sergio and Lizeth can travel 500 miles and still stay on budget.

Try It 3.7.11

Taleisha's phone plan costs her $28.80 a month plus $0.20 per text message. How many text messages can she use and keep her monthly phone bill no more than $50?

Answer

no more than 106 text messages

Try It 3.7.12

Rameen's heating bill is $5.42 per month plus $1.08 per therm. How many therms can Rameen use if he wants his heating bill to be a maximum of $87.50?

Answer

no more than 76 therms

A common goal of most businesses is to make a profit. *Profit* is the money that remains when the expenses have been subtracted from the money earned. In the next example, we will find the number of jobs a small businessman needs to do every month in order to make a certain amount of **profit**.

Example 3.7.13

Elliot has a landscape maintenance business. His monthly expenses are $1,100. If he charges $60 per job, how many jobs must he do to earn a profit of at least $4,000 a month?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the number of jobs Elliot needs
Step 3. Name what we are looking for.	
Choose a variable to represent it	Let j = the number of jobs.
Step 4. Translate. write a sentence that	$60 times the number of jobs minus $1,100 is at least $4,000.
gives the information to find it.	less than or equal to $200.
Translate into an inequality.	$60j - 1100 \geq 4000$
Step 5. Solve the inequality.	$60j \geq 5100$
	$j \geq 85$ jobs
Step 6. Check the answer in the problem and make sure it makes sense.	If Elliot did 90 jobs, his profit would be $60(90) - 1{,}100$, or $4,300. This is more than $4,000.
Step 7. Answer the the question with a complete sentence.	Elliot must work at least 85 jobs.

Try It 3.7.14

Caleb has a pet sitting business. He charges $32 per hour. His monthly expenses are $2272. How many hours must he work in order to earn a profit of at least $800 per month?

Answer

at least 96 hours

Try It 3.7.15

Felicity has a calligraphy business. She charges $2.50 per wedding invitation. Her monthly expenses are $650. How many invitations must she write to earn a profit of at least $2800 per month?

Answer

at least 1380 invitations

Sometimes life gets complicated! There are many situations in which several quantities contribute to the total expense. We must make sure to account for all the individual expenses when we solve problems like this.

Example 3.7.16

Brenda's best friend is having a destination wedding and the event will last 3 days. Brenda has $500 in savings and can earn $15 an hour babysitting. She expects to pay $350 airfare, $375 for food and entertainment and $60 a night for her share of a hotel room. How many hours must she babysit to have enough money to pay for the trip?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	the number of hours Brenda must babysit
Step 3. Name what we are looking for. Choose a variable to represent that quantity.	Let h = the number of hours.
Step 4. Translate. write a sentence that gives the information to find it.	The expenses must be less than or equal to the income. The cost of airfare plus the cost of food and entertainment and the hotel bill must be less than or equal to the savings plus the amount earned babysitting.
Translate into an inequality.	$\$350 + \$375 + \$60(3) \leq \$500 + \$15h$
Step 5. Solve the inequality.	$905 \leq 500 + 15h$ $405 \leq 15h$ $27 \leq h$ $h \geq 27$
Step 6. Check the answer in the problem and make sure it makes sense.	
We substitute 27 into the inequality.	
$905 \leq 500 + 15h$	
$905 \leq 500 + 15(27)$	
$905 \leq 905$	
Step 7. Answer the the question with a complete sentence.	Brenda must babysit at least 27 hours.

? Try It 3.7.17

Malik is planning a 6-day summer vacation trip. He has $840 in savings, and he earns $45 per hour for tutoring. The trip will cost him $525 for airfare, $780 for food and sightseeing, and $95 per night for the hotel. How many hours must he tutor to have enough money to pay for the trip?

Answer

at least 23 hours

? Try It 3.7.18

Josue wants to go on a 10-day road trip next spring. It will cost him $180 for gas, $450 for food, and $49 per night for a motel. He has $520 in savings and can earn $30 per driveway shoveling snow. How many driveways must he shovel to have enough money to pay for the trip?

Answer

at least 20 driveways

Key Concepts

- **Solving inequalities**
  1. **Read** the problem.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose a variable to represent that quantity.
  4. **Translate.** Write a sentence that gives the information to find it. Translate into an inequality.

5. **Solve** the inequality.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

This page titled 3.7: Solve Applications with Linear Inequalities is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **3.6: Solve Applications with Linear Inequalities** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

3.7E: Exercises

Practice Makes Perfect

Solve Applications with Linear Inequalities

In the following exercises, solve.

Exercise 3.7E.1

Mona is planning her son's birthday party and has a budget of $285. The Fun Zone charges $19 per child. How many children can she have at the party and stay within her budget?

Answer

15 children

Exercise 3.7E.2

Carlos is looking at apartments with three of his friends. They want the monthly rent to be no more than $2360. If the roommates split the rent evenly among the four of them, what is the maximum rent each will pay?

Exercise 3.7E.3

A water taxi has a maximum load of 1,800 pounds. If the average weight of one person is 150 pounds, how many people can safely ride in the water taxi?

Answer

12 people

Exercise 3.7E.4

Marcela is registering for her college classes, which cost $105 per unit. How many units can she take to have a maximum cost of $1,365?

Exercise 3.7E.5

Arleen got a $20 gift card for the coffee shop. Her favorite iced drink costs $3.79. What is the maximum number of drinks she can buy with the gift card?

Answer

five drinks

Exercise 3.7E.6

Teegan likes to play golf. He has budgeted $60 next month for the driving range. It costs him $10.55 for a bucket of balls each time he goes. What is the maximum number of times he can go to the driving range next month?

Exercise 3.7E.7

Joni sells kitchen aprons online for $32.50 each. How many aprons must she sell next month if she wants to earn at least $1,000?

Answer

31 aprons

Exercise 3.7E.8

Ryan charges his neighbors $17.50 to wash their car. How many cars must he wash next summer if his goal is to earn at least $1,500?

Exercise 3.7E.9

Keshad gets paid $2,400 per month plus 6% of his sales. His brother earns $3,300 per month. For what amount of total sales will Keshad's monthly pay be higher than his brother's monthly pay?

Answer

$15,000

Exercise 3.7E.10

Kimuyen needs to earn $4,150 per month in order to pay all her expenses. Her job pays her $3,475 per month plus 4% of her total sales. What is the minimum Kimuyen's total sales must be in order for her to pay all her expenses?

Exercise 3.7E.11

Andre has been offered an entry-level job. The company offered him $48,000 per year plus 3.5% of his total sales. Andre knows that the average pay for this job is $62,000. What would Andre's total sales need to be for his pay to be at least as high as the average pay for this job?

Answer

$400,000

Exercise 3.7E.12

Nataly is considering two job offers. The first job would pay her $83,000 per year. The second would pay her $66,500 plus 15% of her total sales. What would her total sales need to be for her salary on the second offer be higher than the first?

Exercise 3.7E.13

Jake's water bill is $24.80 per month plus $2.20 per ccf (hundred cubic feet) of water. What is the maximum number of ccf Jake can use if he wants his bill to be no more than $60?

Answer

16 ccf

Exercise 3.7E.14

Kiyoshi's phone plan costs $17.50 per month plus $0.15 per text message. What is the maximum number of text messages Kiyoshi can use so the phone bill is no more than $56.50?

Exercise 3.7E.15

Marlon's TV plan costs $49.99 per month plus $5.49 per first-run movie. How many first-run movies can he watch if he wants to keep his monthly bill to be a maximum of $100?

Answer

nine movies

Exercise 3.7E.16

Kellen wants to rent a banquet room in a restaurant for her cousin's baby shower. The restaurant charges $350 for the banquet room plus $32.50 per person for lunch. How many people can Kellen have at the shower if she wants the maximum cost to be $1,500?

Exercise 3.7E.17

Moshde runs a hairstyling business from her house. She charges $45 for a haircut and style. Her monthly expenses are $960. She wants to be able to put at least $1,200 per month into her savings account order to open her own salon. How many "cut & styles" must she do to save at least $1,200 per month?

Answer

48 cut & styles

Exercise 3.7E.18

Noe installs and configures software on home computers. He charges $125 per job. His monthly expenses are $1,600. How many jobs must he work in order to make a profit of at least $2,400?

Exercise 3.7E.19

Katherine is a personal chef. She charges $115 per four-person meal. Her monthly expenses are $3,150. How many four-person meals must she sell in order to make a profit of at least $1,900?

Answer

44 meals

Exercise 3.7E.20

Melissa makes necklaces and sells them online. She charges $88 per necklace. Her monthly expenses are $3745. How many necklaces must she sell if she wants to make a profit of at least $1,650?

Exercise 3.7E.21

Five student government officers want to go to the state convention. It will cost them $110 for registration, $375 for transportation and food, and $42 per person for the hotel. There is $450 budgeted for the convention in the student government savings account. They can earn the rest of the money they need by having a car wash. If they charge $5 per car, how many cars must they wash in order to have enough money to pay for the trip?

Answer

49 cars

Exercise 3.7E.22

Cesar is planning a 4-day trip to visit his friend at a college in another state. It will cost him $198 for airfare, $56 for local transportation, and $45 per day for food. He has $189 in savings and can earn $35 for each lawn he mows. How many lawns must he mow to have enough money to pay for the trip?

Exercise 3.7E.23

Alonzo works as a car detailer. He charges $175 per car. He is planning to move out of his parents' house and rent his first apartment. He will need to pay $120 for application fees, $950 for security deposit, and first and last months' rent at $1,140 per month. He has $1,810 in savings. How many cars must he detail to have enough money to rent the apartment?

Answer

9 cars

Exercise 3.7E.24

Eun-Kyung works as a tutor and earns $60 per hour. She has $792 in savings. She is planning an anniversary party for her parents. She would like to invite 40 guests. The party will cost her $1,520 for food and drinks and $150 for the photographer. She will also have a favor for each of the guests, and each favor will cost $7.50. How many hours must she tutor to have enough money for the party?

Everyday Math

Exercise 3.7E.25

Maximum Load on a Stage In 2014, a high school stage collapsed in Fullerton, California, when 250 students got on stage for the finale of a musical production. Two dozen students were injured. The stage could support a maximum of 12,750 pounds. If the average weight of a student is assumed to be 140 pounds, what is the maximum number of students who could safely be on the stage?

Answer

91 students

Exercise 3.7E.26

Maximum Weight on a Boat In 2004, a water taxi sank in Baltimore harbor and five people drowned. The water taxi had a maximum capacity of 3,500 pounds (25 people with average weight 140 pounds). The average weight of the 25 people on the water taxi when it sank was 168 pounds per person. What should the maximum number of people of this weight have been?

Exercise 3.7E.27

Wedding Budget Adele and Walter found the perfect venue for their wedding reception. The cost is $9,850 for up to 100 guests, plus $38 for each additional guest. How many guests can attend if Adele and Walter want the total cost to be no more than $12,500?

Answer

169 guests

Exercise 3.7E.28

Shower Budget Penny is planning a baby shower for her daughter-in-law. The restaurant charges $950 for up to 25 guests, plus $31.95 for each additional guest. How many guests can attend if Penny wants the total cost to be no more than $1,500?

Writing Exercises

Exercise 3.7E.29

Find your last month's phone bill and the hourly salary you are paid at your job. (If you do not have a job, use the hourly salary you would realistically be paid if you had a job.) Calculate the number of hours of work it would take you to earn at least enough money to pay your phone bill by writing an appropriate inequality and then solving it.

Answer

Answers will vary.

Exercise 3.7E.30

Find out how many units you have left, after this term, to achieve your college goal and estimate the number of units you can take each term in college. Calculate the number of terms it will take you to achieve your college goal by writing an appropriate inequality and then solving it.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve applications with linear inequalities.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 3.7E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **3.6E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 3 Review Exercises

Chapter 3 Review Exercises

3.1 Using a Problem Solving Strategy

Approach Word Problems with a Positive Attitude

In the following exercises, reflect on your approach to word problems.

Exercise 1

How has your attitude towards solving word problems changed as a result of working through this chapter? Explain.

Answer

answers will vary

Exercise 2

Did the problem-solving strategy help you solve word problems in this chapter? Explain.

Use a Problem-Solving Strategy for Word Problems

In the following exercises, solve using the problem-solving strategy for word problems. Remember to write a complete sentence to answer each question.

Exercise 3

Three-fourths of the people at a concert are children. If there are 87 children, what is the total number of people at the concert?

Answer

116

Exercise 4

There are nine saxophone players in the band. The number of saxophone players is one less than twice the number of tuba players. Find the number of tuba players.

Solve Number Problems

In the following exercises, solve each number word problem.

Exercise 5

The sum of a number and three is forty-one. Find the number.

Answer

38

Exercise 6

Twice the difference of a number and ten is fifty-four. Find the number.

Exercise 7

One number is nine less than another. Their sum is negative twenty-seven. Find the numbers.

Answer

$-18,-9$

Exercise 8

One number is eleven more than another. If their sum is increased by seventeen, the result is 90. Find the numbers.

Exercise 9

One number is two more than four times another. Their sum is -13. Find the numbers.

Answer

$-3,-10$

Exercise 10

The sum of two consecutive integers is -135. Find the numbers.

Exercise 11

Find three consecutive integers whose sum is -141.

Answer

$-48,-47,-46$

Exercise 12

Find three consecutive even integers whose sum is 234.

Exercise 13

Find three consecutive odd integers whose sum is 51.

Answer

15, 17, 19

Exercise 14

Koji has $5,502 in his savings account. This is $30 less than six times the amount in his checking account. How much money does Koji have in his checking account?

3.2 Solve Percent Applications

Translate and Solve Basic Percent Equations

In the following exercises, translate and solve.

Exercise 15

What number is 67% of 250?

Answer

167.5

Exercise 16

300% of 82 is what number?

Exercise 17

12.5% of what number is 20?

Answer

160

Exercise 18

72 is 30% of what number?

Exercise 19

What percent of 125 is 150?

Answer

120%

Exercise 20

127.5 is what percent of 850?

Solve Percent Applications

In the following exercises, solve.

Exercise 21

The bill for Dino's lunch was $19.45. He wanted to leave 20% of the total bill as a tip. How much should the tip be?

Answer

$3.89

Exercise 22

Reza was very sick and lost 15% of his original weight. He lost 27 pounds. What was his original weight?

Exercise 23

Dolores bought a crib on sale for $350. The sale price was 40% of the original price. What was the original price of the crib?

Answer

$875

Exercise 24

Jaden earns $2,680 per month. He pays $938 a month for rent. What percent of his monthly pay goes to rent?

Find Percent Increase and Percent Decrease

In the following exercises, solve.

Exercise 25

Angel's got a raise in his annual salary from $55,400 to $56,785. Find the percent increase.

Answer

2.5%

Exercise 26

Rowena's monthly gasoline bill dropped from $83.75 last month to $56.95 this month. Find the percent decrease.

Solve Simple Interest Applications

In the following exercises, solve.

Exercise 27

Winston deposited $3,294 in a bank account with interest rate 2.6%. How much interest was earned in 5 years?

Answer

$428.22

Exercise 28

Moira borrowed $4,500 from her grandfather to pay for her first year of college. Three years later, she repaid the $4,500 plus $243 interest. What was the rate of interest?

Exercise 29

Jaime's refrigerator loan statement said he would pay $1,026 in interest for a 4-year loan at 13.5%. How much did Jaime borrow to buy the refrigerator?

Answer

$1,900

Exercise 30

In 12 years, a bond that paid 6.35% interest earned $7,620 interest. What was the principal of the bond?

Solve Applications with Discount or Mark-up

In the following exercises, find the sale price.

Exercise 31

The original price of a handbag was $84. Carole bought it on sale for $21 off.

Answer

$63

Exercise 32

Marian wants to buy a coffee table that costs $495. Next week the coffee table will be on sale for $149 off.

In the following exercises, find

1. the amount of discount and
2. the sale price.

Exercise 33

Emmett bought a pair of shoes on sale at 40% off from an original price of $138.

Answer

1. $55.20
2. $82.80

Exercise 34

Anastasia bought a dress on sale at 75% off from an original price of $280.

In the following exercises, find ⓐ the amount of discount and ⓑ the discount rate. (Round to the nearest tenth of a percent, if needed.)

Exercise 35

Zack bought a printer for his office that was on sale for $380. The original price of the printer was $450.

Answer

1. $70
2. 15.6%

Exercise 36

Lacey bought a pair of boots on sale for $95. The original price of the boots was $200.

In the following exercises, find

1. the amount of the mark-up and
2. the list price.

Exercise 37

Nga and Lauren bought a chest at a flea market for $50. They re-finished it and then added a 350% mark-up.

Answer

1. $175
2. $225

Exercise 38

Carly bought bottled water for $0.24 per bottle at the discount store. She added a 75% mark-up before selling them at the football game.

3.3 Solve Mixture Applications

Solve Coin Word Problems

In the following exercises, solve each coin word problem.

Exercise 39

Francie has $4.35 in dimes and quarters. The number of dimes is five more than the number of quarters. How many of each coin does she have?

Answer

16 dimes, 11 quarters

Exercise 40

Scott has $0.39 in pennies and nickels. The number of pennies is eight times the number of nickels. How many of each coin does he have?

Exercise 41

Paulette has $140 in $5 and $10 bills. The number of $10 bills is one less than twice the number of $5 bills. How many of each does she have?

Answer

six $5 bills, 11 $10 bills

Exercise 42

Lenny has $3.69 in pennies, dimes, and quarters. The number of pennies is three more than the number of dimes. The number of quarters is twice the number of dimes. How many of each coin does he have?

Solve Ticket and Stamp Word Problems

In the following exercises, solve each ticket or stamp word problem.

Exercise 43

A church luncheon made $842. Adult tickets cost $10 each and children's tickets cost $6 each. The number of children was 12 more than twice the number of adults. How many of each ticket were sold?

Answer

35 adults, 82 children

Exercise 44

Tickets for a basketball game cost $2 for students and $5 for adults. The number of students was three less than 10 times the number of adults. The total amount of money from ticket sales was $619. How many of each ticket were sold?

Exercise 45

125 tickets were sold for the jazz band concert for a total of $1,022. Student tickets cost $6 each and general admission tickets cost $10 each. How many of each kind of ticket were sold?

Answer

57 students, 68 adults

Exercise 46

One afternoon the water park sold 525 tickets for a total of $13,545. Child tickets cost $19 each and adult tickets cost $40 each. How many of each kind of ticket were sold?

Exercise 47

Ana spent $4.06 buying stamps. The number of $0.41 stamps she bought was five more than the number of $0.26 stamps. How many of each did she buy?

Answer

three $0.26 stamps, eight $0.41 stamps

Exercise 48

Yumi spent $34.15 buying stamps. The number of $0.56 stamps she bought was 10 less than four times the number of $0.41 stamps. How many of each did she buy?

Solve Mixture Word Problems

In the following exercises, solve each mixture word problem.

Exercise 49

Marquese is making 10 pounds of trail mix from raisins and nuts. Raisins cost $3.45 per pound and nuts cost $7.95 per pound. How many pounds of raisins and how many pounds of nuts should Marquese use for the trail mix to cost him $6.96 per pound?

Answer

2.2 lb. of raisins, 7.8 lb. of nuts

Exercise 50

Amber wants to put tiles on the backsplash of her kitchen counters. She will need 36 square feet of tile. She will use basic tiles that cost $8 per square foot and decorator tiles that cost $20 per square foot. How many square feet of each tile should she use so that the overall cost of the backsplash will be $10 per square foot?

Exercise 51

Shawn has $15,000 to invest. She will put some of it into a fund that pays 4.5% annual interest and the rest in a certificate of deposit that pays 1.8% annual interest. How much should she invest in each account if she wants to earn 4.05% annual interest on the total amount?

Answer

$12,500 at 4.5%, $2,500 at 1.8%

Exercise 52

Enrique borrowed $23,500 to buy a car. He pays his uncle 2% interest on the $4,500 he borrowed from him, and he pays the bank 11.5% interest on the rest. What average interest rate does he pay on the total $23,500? (Round your answer to the nearest tenth of a percent.)

3.4 Solve Geometry Applications: Triangles, Rectangles and the Pythagorean Theorem

Solve Applications Using Triangle Properties

In the following exercises, solve using triangle properties.

Exercise 53

The measures of two angles of a triangle are 22 and 85 degrees. Find the measure of the third angle.

Answer

73°

Exercise 54

The playground at a shopping mall is a triangle with perimeter 48 feet. The lengths of two sides are 19 feet and 14 feet. How long is the third side?

Exercise 55

A triangular road sign has base 30 inches and height 40 inches. What is its area?

Answer

600 square inches

Exercise 56

What is the height of a triangle with area 67.5 square meters and base 9 meters?

Exercise 57

One angle of a triangle is 30° more than the smallest angle. The largest angle is the sum of the other angles. Find the measures of all three angles.

Answer

30°,60°,90°

Exercise 58

One angle of a right triangle measures 58°. What is the measure of the other angles of the triangle?

Exercise 59

The measure of the smallest angle in a right triangle is 45° less than the measure of the next larger angle. Find the measures of all three angles.

Answer

22.5°,67.5°,90°


Exercise 60

The perimeter of a triangle is 97 feet. One side of the triangle is eleven feet more than the smallest side. The third side is six feet more than twice the smallest side. Find the lengths of all sides.

Use the Pythagorean Theorem

In the following exercises, use the Pythagorean Theorem to find the length of the hypotenuse.


Exercise 61



Answer


26

Exercise 62



In the following exercises, use the Pythagorean Theorem to find the length of the missing side. Round to the nearest tenth, if necessary.


Exercise 63




Answer

8

Exercise 64




Exercise 65



Answer

8.1


Exercise 66



In the following exercises, solve. Approximate to the nearest tenth, if necessary.

Exercise 67

Sergio needs to attach a wire to hold the antenna to the roof of his house, as shown in the figure. The antenna is 8 feet tall and Sergio has 10 feet of wire. How far from the base of the antenna can he attach the wire?




Answer

$6'$

Exercise 68

Seong is building shelving in his garage. The shelves are 36 inches wide and 15 inches tall. He wants to put a diagonal brace across the back to stabilize the shelves, as shown. How long should the brace be?



Solve Applications Using Rectangle Properties

In the following exercises, solve using rectangle properties.

Exercise 69

The length of a rectangle is 36 feet and the width is 19 feet. Find the

1. perimeter
2. area.

Answer

1. 110 ft.
2. 684 sq. ft.

Exercise 70

A sidewalk in front of Kathy's house is in the shape of a rectangle four feet wide by 45 feet long. Find the

1. perimeter
2. area.

Exercise 71

The area of a rectangle is 2356 square meters. The length is 38 meters. What is the width?

Answer

62 m

Exercise 72

The width of a rectangle is 45 centimeters. The area is 2,700 square centimeters. What is the length?

Exercise 73

The length of a rectangle is 12 cm more than the width. The perimeter is 74 cm. Find the length and the width.

Answer

24.5 cm, 12.5 cm

Exercise 74

The width of a rectangle is three more than twice the length. The perimeter is 96 inches. Find the length and the width.

3.5 Solve Uniform Motion Applications

Solve Uniform Motion Applications

In the following exercises, solve.

Exercise 75

When Gabe drives from Sacramento to Redding it takes him 2.2 hours. It takes Elsa 2 hours to drive the same distance. Elsa's speed is seven miles per hour faster than Gabe's speed. Find Gabe's speed and Elsa's speed.

Answer

Gabe 70 mph, Elsa 77 mph

Exercise 76

Louellen and Tracy met at a restaurant on the road between Chicago and Nashville. Louellen had left Chicago and drove 3.2 hours towards Nashville. Tracy had left Nashville and drove 4 hours towards Chicago, at a speed one mile per hour faster than Louellen's speed. The distance between Chicago and Nashville is 472 miles. Find Louellen's speed and Tracy's speed.

Exercise 77

Two busses leave Amarillo at the same time. The Albuquerque bus heads west on the I-40 at a speed of 72 miles per hour, and the Oklahoma City bus heads east on the I-40 at a speed of 78 miles per hour. How many hours will it take them to be 375 miles apart?

Answer

2.5 hours

Exercise 78

Kyle rowed his boat upstream for 50 minutes. It took him 30 minutes to row back downstream. His speed going upstream is two miles per hour slower than his speed going downstream. Find Kyle's upstream and downstream speeds.

Exercise 79

At 6:30, Devon left her house and rode her bike on the flat road until 7:30. Then she started riding uphill and rode until 8:00. She rode a total of 15 miles. Her speed on the flat road was three miles per hour faster than her speed going uphill. Find Devon's speed on the flat road and riding uphill.

Answer

flat road 11 mph, uphill 8 mph

Exercise 80

Anthony drove from New York City to Baltimore, a distance of 192 miles. He left at 3:45 and had heavy traffic until 5:30. Traffic was light for the rest of the drive, and he arrived at 7:30. His speed in light traffic was four miles per hour more than twice his speed in heavy traffic. Find Anthony's driving speed in heavy traffic and light traffic.

3.6 Solve Applications with Linear Inequalities

Solve Applications with Linear Inequalities

In the following exercises, solve.

Exercise 81

Julianne has a weekly food budget of $231 for her family. If she plans to budget the same amount for each of the seven days of the week, what is the maximum amount she can spend on food each day?

Answer

$33 per day

Exercise 82

Rogelio paints watercolors. He got a $100 gift card to the art supply store and wants to use it to buy $12'' \times 16''$ canvases. Each canvas costs $10.99. What is the maximum number of canvases he can buy with his gift card?

Exercise 83

Briana has been offered a sales job in another city. The offer was for $42,500 plus 8% of her total sales. In order to make it worth the move, Briana needs to have an annual salary of at least $66,500. What would her total sales need to be for her to move?

Answer

at least $300,000

Exercise 84

Renee's car costs her $195 per month plus $0.09 per mile. How many miles can Renee drive so that her monthly car expenses are no more than $250?

Exercise 85

Costa is an accountant. During tax season, he charges $125 to do a simple tax return. His expenses for buying software, renting an office, and advertising are $6,000. How many tax returns must he do if he wants to make a profit of at least $8,000?

Answer

at least 112 jobs

Exercise 86

Jenna is planning a 5-day resort vacation with three of her friends. It will cost her $279 for airfare, $300 for food and entertainment, and $65 per day for her share of the hotel. She has $550 saved towards her vacation and can earn $25 per hour as an assistant in her uncle's photography studio. How many hours must she work in order to have enough money for her vacation?

Practice Test

Exercise 87

Four-fifths of the people on a hike are children. If there are 12 children, what is the total number of people on the hike?

Answer

15

Exercise 88

One number is three more than twice another. Their sum is −63. Find the numbers.

Exercise 89

The sum of two consecutive odd integers is −96. Find the numbers.

Answer

−49,−47

Exercise 90

Marla's breakfast was 525 calories. This was 35% of her total calories for the day. How many calories did she have that day?

Exercise 91

Humberto's hourly pay increased from $16.25 to $17.55. Find the percent increase.

Answer

8%

Exercise 92

Melinda deposited $5,985 in a bank account with an interest rate of 1.9%. How much interest was earned in 2 years?

Exercise 93

Dotty bought a freezer on sale for $486.50. The original price of the freezer was $695. Find

1. the amount of discount and
2. the discount rate.

Answer

1. $208.50
2. 30%

Exercise 94

Bonita has $2.95 in dimes and quarters in her pocket. If she has five more dimes than quarters, how many of each coin does she have?

Exercise 95

At a concert, $1,600 in tickets were sold. Adult tickets were $9 each and children's tickets were $4 each. If the number of adult tickets was 30 less than twice the number of children's tickets, how many of each kind were sold?

Answer

140 adult, 85 children

Exercise 96

Kim is making eight gallons of punch from fruit juice and soda. The fruit juice costs $6.04 per gallon and the soda costs $4.28 per gallon. How much fruit juice and how much soda should she use so that the punch costs $5.71 per gallon?

Exercise 97

The measure of one angle of a triangle is twice the measure of the smallest angle. The measure of the third angle is 14 more than the measure of the smallest angle. Find the measures of all three angles.

Answer


41.5°,55.5°,83°

Exercise 98

What is the height of a triangle with area 277.2 square inches and base 44 inches?

In the following exercises, use the Pythagorean Theorem to find the length of the missing side. Round to the nearest tenth, if necessary.


Exercise 99



Answer


10

Exercise 100



Exercise 101

A baseball diamond is really a square with sides of 90 feet. How far is it from home plate to second base, as shown?



Answer

127.3 ft.

Exercise 102

The length of a rectangle is two feet more than five times the width. The perimeter is 40 feet. Find the dimensions of the rectangle.

Exercise 103

Two planes leave Dallas at the same time. One heads east at a speed of 428 miles per hour. The other plane heads west at a speed of 382 miles per hour. How many hours will it take them to be 2,025 miles apart?

Answer

2.5 hours

Exercise 104

Leon drove from his house in Cincinnati to his sister's house in Cleveland, a distance of 252 miles. It took him 412412 hours. For the first half hour he had heavy traffic, and the rest of the time his speed was five miles per hour less than twice his speed in heavy traffic. What was his speed in heavy traffic?

Exercise 105

Chloe has a budget of $800 for costumes for the 18 members of her musical theater group. What is the maximum she can spend for each costume?

Answer

at most $44.44 per costume

Exercise 106

Frank found a rental car deal online for $49 per week plus $0.24 per mile. How many miles could he drive if he wants the total cost for one week to be no more than $150?

Chapter 3 Review Exercises is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **Chapter 3 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

CHAPTER OVERVIEW

4: Graphs

4.1: Use the Rectangular Coordinate System

4.1E: Exercises

4.2: Graph Linear Equations in Two Variables

4.2E: Exercises

4.3: Graph with Intercepts

4.3E: Exercises

4.4: Understanding the Slope of a Line

4.4E: Exercises

4.5: Use the Slope–Intercept Form of an Equation of a Line

4.5E: Exercises

4.6: Find the Equation of a Line

4.6E: Exercises

4.7: Graphs of Linear Inequalities

4.7E: Exercises

Chapter 4 Review Exercises

This page titled 4: Graphs is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

4.1: Use the Rectangular Coordinate System

Learning Objectives

By the end of this section, you will be able to:

- Plot points in a rectangular coordinate system
- Verify solutions to an equation in two variables
- Complete a table of solutions to a linear equation
- Find solutions to a linear equation in two variables

Note

Before you get started, take this readiness quiz.

1. Evaluate $x+3$ when $x=-1$.
   If you missed this problem, review Example 1.5.25.
2. Evaluate $2x-5y$ when $x=3$ and y=−2.
   If you missed this problem, review Example 1.5.28.
3. Solve for y: $40-4y=20$
   If you missed this problem, review Example 2.3.1.

Plot Points on a Rectangular Coordinate System

Just like maps use a grid system to identify locations, a grid system is used in algebra to show a relationship between two variables in a **rectangular coordinate system**. The rectangular coordinate system is also called the *xy*-plane or the 'coordinate plane'.

The horizontal number line is called the *x-axis*. The vertical number line is called the *y-axis*. The *x*-axis and the *y*-axis together form the rectangular coordinate system. These axes divide a plane into four regions, called **quadrants**. The quadrants are identified by Roman numerals, beginning on the upper right and proceeding counterclockwise. See Figure \(\PageIndex{1}\).




Figure 4.1.1: 'Quadrant' has the root 'quad,' which means 'four.'

In the **rectangular coordinate system**, every point is represented by an *ordered pair*. The first number in the ordered pair is the ***x*-coordinate** of the point, and the second number is the ***y*-coordinate** of the point.

ORDERED PAIR

An **ordered pair**, (x,y)(x,y), gives the coordinates of a point in a rectangular coordinate system.




Figure 4.1.2

The first number is the *x*-coordinate.

The second number is the *y*-coordinate.

The phrase 'ordered pair' means the order is important. What is the ordered pair of the point where the axes cross? At that point both coordinates are zero, so its ordered pair is $(0, 0)$. The point $(0, 0)$ has a special name. It is called the **origin**.

THE ORIGIN

The point $(0, 0)$ is called the **origin**. It is the point where the *x*-axis and *y*-axis intersect.

We use the coordinates to locate a point on the *xy*-plane. Let's plot the point $(1, 3)$ as an example. First, locate 1 on the *x*-axis and lightly sketch a vertical line through $x = 1$. Then, locate 3 on the *y*-axis and sketch a horizontal line through $y = 3$. Now, find the point where these two lines meet — that is the point with coordinates $(1, 3)$.




Figure 4.1.3

Notice that the vertical line through $x = 1$ and the horizontal line through $y = 3$ are not part of the graph. We just used them to help us locate the point $(1, 3)$.

Example 4.1.1

Plot each point in the rectangular coordinate system and identify the **quadrant** in which the point is located:

a. (−5,4)
b. (−3,−4)
c. (2,−3)
d. (−2,3)
e. $(3, \frac{5}{2})$

Solution

The first number of the coordinate pair is the *x*-coordinate, and the second number is the *y*-coordinate.

a. Since x=−5, the point is to the left of the *y*-axis. Also, since y=4, the point is above the *x*-axis. The point (−5,4) is in Quadrant II.
b. Since x=−3, the point is to the left of the *y*-axis. Also, since y=−4, the point is below the *x*-axis. The point (−3,−4) is in Quadrant III.
c. Since x=2, the point is to the right of the *y*-axis. Since y=−3, the point is below the *x*-axis. The point (2,−3) is in Quadrant IV.
d. Since x=−2, the point is to the left of the *y*-axis. Since y=3, the point is above the *x*-axis. The point (−2,3) is in Quadrant II.
e. Since x=3, the point is to the right of the *y*-axis. Since $y = \frac{5}{2}$, the point is above the *x*-axis. (It may be helpful to write $\frac{5}{2}$ as a mixed number or decimal.) The point $(3, \frac{5}{2})$ is in Quadrant I.




Figure 4.1.4


Try It 4.1.2

Plot each point in a rectangular coordinate system and identify the quadrant in which the point is located:

a. $(-2,1)$
b. $(-3,-1)$
c. $(4,-4)$
d. $(-4,4)$
e. $(-4, \frac{3}{2})$

Answer

The point a (-2,1) is in Quadrant II.
The point b (-3,-1) is in Quadrant III.
The point c (4,-4) is in Quadrant IV.
The point d (-4,4) is in Quadrant II.
The point e $(-4, \frac{3}{2})$ is in Quadrant II.




Try It 4.1.3

Plot each point in a rectangular coordinate system and identify the quadrant in which the point is located:

a. $(-4,1)$
b. $(-2,3)$
c. $(2,-5)$
d. $(-2,5)$
e. $(-3, \frac{5}{2})$

Answer

The point a (-4, 1) is in Quadrant II.
The point b (-2, 3) is in Quadrant II.
The point c (2, -5) is in Quadrant IV.
The point d (-2, 5) is in Quadrant II.
The point e $(-3, \frac{5}{2})$ is in Quadrant II.



How do the signs affect the location of the points? You may have noticed some patterns as you graphed the points in the previous example.

For the point in Figure 4.1.4 in Quadrant IV, what do you notice about the signs of the coordinates? What about the signs of the coordinates of points in the third quadrant? The second quadrant? The first quadrant?

Can you tell just by looking at the coordinates in which quadrant the point (−2,5) is located? In which quadrant is (2,−5) located?

QUADRANTS

We can summarize sign patterns of the quadrants in this way.

Quadrant I	Quadrant II	Quadrant III	Quadrant IV	
(x, y)	(x, y)	(x, y)	(x, y)	(4.1.1)
$(+, +)$	$(-, +)$	$(-, -)$	$(+, -)$	




Figure 4.1.5

What if one coordinate is zero as shown in Figure 4.1.6? Where is the point (0,4) located? Where is the point (−2,0) located?




Figure 4.1.6

The point (0,4) is on the *y*-axis and the point (−2,0) is on the *x*-axis.

POINTS ON THE AXES

Points with a ***y*-coordinate** equal to 0 are on the *x*-axis, and have coordinates (a,0).

Points with an ***x*-coordinate** equal to 0 are on the *y*-axis, and have coordinates (0,b).

Example 4.1.4

Plot each point:

1. (0,5)
2. (4,0)
3. (−3,0)
4. (0,0)
5. (0,−1)

Solution

1. Since x=0, the point whose coordinates are (0,5) is on the *y*-axis.
2. Since y=0, the point whose coordinates are (4,0) is on the *x*-axis.
3. Since y=0, the point whose coordinates are (−3,0) is on the *x*-axis.
4. Since x=0 and y=0, the point whose coordinates are (0,0) is the origin.
5. Since x=0, the point whose coordinates are (0,−1) is on the *y*-axis.





Figure 4.1.7

Try It 4.1.5

Plot each point:

1. (4,0)
2. (−2,0)
3. (0,0)
4. (0,2)
5. (0,−3).

Answer




https://math.libretexts.org/@go/page/30503

Try It 4.1.6

Plot each point:

1. (−5,0)
2. (3,0)
3. (0,0)
4. (0,−1)
5. (0,4).

Answer



In algebra, being able to identify the coordinates of a point shown on a graph is just as important as being able to plot points. To identify the *x*-coordinate of a point on a graph, read the number on the *x*-axis directly above or below the point. To identify the *y*-coordinate of a point, read the number on the *y*-axis directly to the left or right of the point. Remember, when you write the **ordered pair** use the correct order, (x,y).

Example 4.1.7

Name the ordered pair of each point shown in the rectangular coordinate system.




Figure 4.1.8

Solution

Point A is above −3 on the *x*-axis, so the *x*-coordinate of the point is −3.

The point is to the left of 3 on the *y*-axis, so the *y*-coordinate of the point is 3.

The coordinates of the point are (−3,3).

Point B is below −1 on the *x*-axis, so the *x*-coordinate of the point is −1.

The point is to the left of −3 on the *y*-axis, so the *y*-coordinate of the point is −3.

The coordinates of the point are (−1,−3).

Point C is above 2 on the *x*-axis, so the *x*-coordinate of the point is 2.

The point is to the right of 4 on the *y*-axis, so the *y*-coordinate of the point is 4.

The coordinates of the point are (2,4).

Point D is below 4 on the *x*-axis, so the *x*-coordinate of the point is 4.

The point is to the right of −4 on the *y*-axis, so the *y*-coordinate of the point is −4.

The coordinates of the point are (4,−4).

Point E is on the *y*-axis at y=−2. The coordinates of point E are (0,−2).

Point F is on the *x*-axis at x=3. The coordinates of point F are (3,0).

Try It 4.1.8

Name the ordered pair of each point shown in the rectangular coordinate system.




Figure 4.1.9

Answer

A: (5,1) B: (−2,4) C: (−5,−1) D: (3,−2) E: (0,−5) F: (4,0)

Try It 4.1.9

Name the ordered pair of each point shown in the rectangular coordinate system.




Figure 4.1.10

Answer

A: (4,2) B: (−2,3) C: (−4,−4) D: (3,−5) E: (−3,0) F: (0,2)

Verify Solutions to an Equation in Two Variables

Up to now, all the equations you have solved were equations with just one variable. In almost every case, when you solved the equation you got exactly one solution. The process of solving an equation ended with a statement like x=4. (Then, you checked the solution by substituting back into the equation.)

Here's an example of an equation in one variable, and its one solution.

$$\begin{aligned} 3x+5 &= 17 \\ 3x &= 12 \\ x &= 4 \end{aligned}$$

But equations can have more than one variable. Equations with two variables may be of the form Ax+By=C. Equations of this form are called **linear equations in two variables**.

LINEAR EQUATION

An equation of the form Ax+By=C, where A and B are not both zero, is called a **linear equation in two variables**.

Notice the word *line* in **linear**. Here is an example of a linear equation in two variables, x and y.

$$\begin{aligned} Ax+By&=C \\ x+4y&=8 \\ A=1,\ B&=4,\ C=8 \end{aligned}$$

Figure 4.1.11

The equation y=−3x+5 is also a **linear equation**. But it does not appear to be in the form Ax+By=C. We can use the Addition Property of Equality and rewrite it in Ax+By=C form.

	y	$=$	$-3x+5$
Add to both sides.	$y+3x$	$=$	$-3x+5+3x$
Simplify.	$y+3x$	$=$	5
Use the Commutative Property to put it in $Ax+By=C$ form.	$3x+y$	$=$	5

By rewriting y=−3x+5 as 3x+y=5, we can easily see that it is a linear equation in two variables because it is of the form Ax+By=C. When an equation is in the form Ax+By=C, we say it is in *standard form*.

STANDARD FORM OF LINEAR EQUATION

A linear equation is in standard form when it is written Ax+By=C.

Most people prefer to have A, B, and C be integers and $A \geq 0$ when writing a linear equation in standard form, although it is not strictly necessary.

Linear equations have infinitely many solutions. For every number that is substituted for x there is a corresponding y value. This pair of values is a *solution* to the linear equation and is represented by the ordered pair (x,y). When we substitute these values of x and y into the equation, the result is a true statement, because the value on the left side is equal to the value on the right side.

Solution OF A LINEAR EQUATION IN TWO VARIABLES

An **ordered pair** (x,y) is a **solution** of the linear equation Ax+By=C, if the equation is a true statement when the *x*- and *y*-values of the ordered pair are substituted into the equation.

Example 4.1.10

Determine which ordered pairs are solutions to the equation x+4y=8.

(a) (0,2)

(b) (2,−4)

(c) (−4,3)

Solution

Substitute the *x*- and *y*-values from each ordered pair into the equation and determine if the result is a true statement.

(a)	(b)	(c)
(0, 2)	(2, –4)	(–4, 3)
$x = 0, y = 2$	$x = 2, y = -4$	$x = -4, y = 3$
$x + 4y = 8$	$x + 4y = 8$	$x + 4y = 8$
$0 + 4 \cdot 2 \stackrel{?}{=} 8$	$2 + 4(-4) \stackrel{?}{=} 8$	$-4 + 4 \cdot 3 \stackrel{?}{=} 8$
$0 + 8 \stackrel{?}{=} 8$	$2 + (-16) \stackrel{?}{=} 8$	$-4 + 12 \stackrel{?}{=} 8$
$8 = 8$ ✓	$-14 \neq 8$	$8 = 8$ ✓
(0, 2) is a solution.	(2, –4) is not a solution.	(–4, 3) is a solution.

Try It 4.1.11

Which of the following ordered pairs are solutions to 2x+3y=6?

1. (3,0)
2. (2,0)
3. (6,−2)

Answer

1, 3

Try It 4.1.12

Which of the following ordered pairs are solutions to the equation 4x−y=8?

1. (0,8)
2. (2,0)
3. (1,−4)

Answer

2, 3

Example 4.1.13

Which of the following ordered pairs are solutions to the equation y=5x−1?

(a) (0,−1)

(b) (1,4)

(c) (−2,−7)

Solution

Substitute the *x*- and *y*-values from each **ordered pair** into the equation and determine if it results in a true statement.

(a)	(b)	(c)
(0, –1)	(1, 4)	(–2, –7)
$x = 0, y = -1$	$x = 1, y = 4$	$x = -2, y = -7$
$y = 5x - 1$	$y = 5x - 1$	$y = 5x - 1$
$-1 \stackrel{?}{=} 5(0) - 1$	$4 \stackrel{?}{=} 5(1) - 1$	$-7 \stackrel{?}{=} 5(-2) - 1$
$-1 \stackrel{?}{=} 0 - 1$	$4 \stackrel{?}{=} 5 - 1$	$-7 \stackrel{?}{=} -10 - 1$
$-1 = -1$ ✓	$4 = 4$ ✓	$-7 \neq -11$
(0, –1) is a solution.	(1, 4) is a solution.	(–2, –7) is not a solution.

Try It 4.1.14

Which of the following ordered pairs are solutions to the equation y=4x−3?

1. (0,3)
2. (1,1)
3. (−1,−1)

Answer

2

Try It 4.1.15

Which of the following ordered pairs are solutions to the equation y=−2x+6?

1. (0,6)
2. (1,4)
3. (−2,−2)

Answer

1, 2

Complete a Table of Solutions to a Linear Equation in Two Variables

In the examples above, we substituted the *x*- and *y*-values of a given ordered pair to determine whether or not it was a solution to a linear equation. But how do you find the ordered pairs if they are not given? It's easier than you might think—you can just pick a value for x and then solve the equation for y. Or, pick a value for y and then solve for x.

We'll start by looking at the solutions to the equation y = 5x−1 that we found in Example 4.1.13. We can summarize this information in a table of solutions, as shown in Table 4.1.1.

Table 4.1.1

y=5x−1		
x	**y**	**(x,y)**
0	−1	(0,−1)
1	4	(1,4)

To find a third solution, we'll let x=2 and solve for y.

	$y = 5x - 1$
Substitute $x = 2$.	$y = 5(2) - 1$
Multiply.	$y = 10 - 1$
Simplify.	$y = 9$

Figure 4.1.12

The ordered pair (2,9) is a solution to y=5x−1. We will add it to Table 4.1.2.

Table 4.1.2

y=5x−1		
x	**y**	**(x,y)**
0	−1	(0,−1)
1	4	(1,4)
2	9	(2,9)

We can find more solutions to the equation by substituting in any value of x or any value of y and solving the resulting equation to get another ordered pair that is a solution. There are infinitely many solutions of this equation.

Example 4.1.16

Complete the table to find three solutions to the equation $y=4x-2$.

Table 4.1.3

y=4x−2		
x	**y**	**(x,y)**
0		
−1		
2		

Solution

Substitute $x=0$, $x=-1$, and $x=2$ into $y=4x-2$.

$x=0$	$x=-1$	$x=2$
$y=4x-2$	$y=4x-2$	$y=4x-2$
$y=4\cdot 0-2$	$y=4(-1)-2$	$y=4\cdot 2-2$
$y=0-2$	$y=-4-2$	$y=8-2$
$y=-2$	$y=-6$	$y=6$
$(0,-2)$	$(-1,-6)$	$(2,6)$

The results are summarized in Table 4.1.4

Table 4.1.4

y=4x−2		
x	**y**	**(x,y)**
0	−2	(0,−2)
−1	−6	(−1,−6)
2	6	(2,6)

Try It 4.1.17

Complete the table to find three solutions to this equation: $y=3x-1$.

Table 4.1.5

y=3x−1		
x	**y**	**(x,y)**
0		
−1		
2		

Answer

Table 4.1.6

y=3x−1

x	y	(x,y)
0	-1	(0, -1)
−1	-4	(-1, -4)
2	5	(2, 5)

Try It 4.1.18

Complete the table to find three solutions to this equation: y=6x+1.

Table 4.1.7

y=6x+1		
x	**y**	**(x,y)**
-2		

Answer

Table 4.1.8

y=6x+1		
x	**y**	**(x,y)**
0	1	(0,1)
1	7	(1,7)
−2	−11	(−2,−11)

Example 4.1.19

Complete Table 4.1.9 to find three solutions to the equation 5x−4y=20.

Table 4.1.9

5x−4y=20		
x	**y**	**(x,y)**
	0	
	5	

Solution

Substitute the given value into the equation 5x−4y=20 and solve for the other variable. Then, fill in the values in the table.

https://math.libretexts.org/@go/page/30503

$x = 0$	$y = 0$	$y = 5$
$5x - 4y = 20$	$5x - 4y = 20$	$5x - 4y = 20$
$5 \cdot 0 - 4y = 20$	$5x - 4 \cdot 0 = 20$	$5x - 4 \cdot 5 = 20$
$0 - 4y = 20$	$5x - 0 = 20$	$5x - 20 = 20$
$-4y = 20$	$5x = 20$	$5x = 40$
$y = -5$	$x = 4$	$x = 8$
$(0, -5)$	$(4, 0)$	$(8, 5)$

The results are summarized in Table 4.1.10

Table 4.1.10

5x−4y=20		
x	**y**	**(x,y)**
0	−5	(0,−5)
4	0	(4,0)
8	5	(8,5)

Try It 4.1.20

Complete the table to find three solutions to this equation: 2x−5y=20.

Table 4.1.11

2x−5y=20		
x	**y**	**(x,y)**
-5		

Answer

Table 4.1.12

2x−5y=20		
x	**y**	**(x,y)**
0	−4	(0,−4)
10	0	(10,0)
−5	−6	(−5,−6)

Try It 4.1.21

Complete the table to find three solutions to this equation: 3x−4y=12.

Table 4.1.13

3x−4y=12		
x	**y**	**(x,y)**

-4		

Answer

Table 4.1.14

3x−4y=12		
x	**y**	**(x,y)**
0	−3	(0,−3)
4	0	(4,0)
−4	−6	(−4,−6)

Find Solutions to a Linear Equation

To find a solution to a linear equation, you really can pick *any* number you want to substitute into the equation for x or y. But since you'll need to use that number to solve for the other variable it's a good idea to choose a number that's easy to work with.

When the equation is in *y*-form, with the *y* by itself on one side of the equation, it is usually easier to choose values of x and then solve for y.

Example 4.1.22

Find three solutions to the equation y=−3x+2.

Solution

We can substitute any value we want for x or any value for y. Since the equation is in *y*-form, it will be easier to substitute in values of x. Let's pick x=0, x=1, and x=−1.

			$x=0$	$x=1$	$x=-1$
			$y=-3x+2$	$y=-3x+2$	$y=-3x+2$
Substitute the value into the equation.			$y=-3\cdot 0+2$	$y=-3\cdot 1+2$	$y=-3(-1)+2$
Simplify.			$y=0+2$	$y=-3+2$	$y=3+2$
Simplify.			$y=2$	$y=-1$	$y=5$
Write the ordered pair.			(0, 2)	(1, -1)	(-1, 5)
Check.					
y=−3x+2	y=−3x+2	y=−3x+2			
$2\stackrel{?}{=}-3\cdot 0+2$	$-1\stackrel{?}{=}-3\cdot 1+2$	$5\stackrel{?}{=}-3(-1)+2$			
$2\stackrel{?}{=}0+2$	$-1\stackrel{?}{=}-3+2$	$5\stackrel{?}{=}-3+2$			
$2=2\checkmark$	$-1=-1\checkmark$	$5=5\checkmark$			

Table 4.1.15

So, (0,2), (1,−1) and (−1,5) are all solutions to y=−3x+2. We show them in Table 4.1.16

Table 4.1.16

y=−3x+2

x	y	(x,y)
0	2	(0,2)
1	−1	(1,−1)
−1	5	(−1,5)

Try It 4.1.23

Find three solutions to this equation: y=−2x+3.

Answer

Answers will vary. Answers might include (0,3), (1,1) and (−1,5).

Try It 4.1.24

Find three solutions to this equation: y=−4x+1.

Answer

Answers will vary. Answers might include (0,1), (1,−3) and (−1,5).

We have seen how using zero as one value of x makes finding the value of y easy. When an equation is in standard form, with both the x and y on the same side of the equation, it is usually easier to first find one solution when x=0 find a second solution when y=0, and then find a third solution.

Example 4.1.25

Find three solutions to the equation 3x+2y=6.

Solution

We can substitute any value we want for x or any value for y. Since the equation is in standard form, let's pick first x=0, then y=0, and then find a third point.

Table 4.1.17

$x=0$	$y=0$	$x=1$			
			$3x+2y=6$	$3x+2y=6$	$3x+2y=6$
Substitute the value into the equation.			$3(0)+2y=6$	$3x+2(0)=6$	$3(1)+2y=6$
Simplify.			$0+2y=6$	$3x+0=6$	$3+2y=6$
Solve.			$2y=6$	$3x=6$	$2y=3$
			$y=3$	$x=2$	$y=\frac{3}{2}$
Write the ordered pair.			(0, 3)	(2, 0)	$(1, \frac{3}{2})$
Check.					
3x+2y=6	3x+2y=6	3x+2y=6			
$3\cdot 0+2\cdot 3\stackrel{?}{=}6$	$3\cdot 2+2\cdot 0\stackrel{?}{=}6$	$3\cdot 1+2\cdot\frac{3}{2}\stackrel{?}{=}6$			
$0+6\stackrel{?}{=}6$	$6+0\stackrel{?}{=}6$	$3+3\stackrel{?}{=}6$			
$6=6$✓	$6=6$✓	$6=6$✓			

So (0,3), (2,0), and $(1, \frac{3}{2})$ are all solutions to the equation 3x+2y=6. We can list these three solutions in Table 4.1.18

Table 4.1.18

3x+2y=6		
x	**y**	**(x,y)**
0	3	(0,3)
2	0	(2,0)
1	$\frac{3}{2}$	$(1, \frac{3}{2})$

? Try It 4.1.26

Find three solutions to the equation 2x+3y=6.

Answer

Answers will vary. Answers might include (0,2), (3,0) and (6, −2).

? Try It 4.1.27

Find three solutions to the equation 4x+2y=8.

Answer

Answers will vary. Answers might include (0,4), (2,0) and (−1,6).

Key Concepts

- **Sign Patterns of the Quadrants**

Quadrant I	Quadrant II	Quadrant III	Quadrant IV
(x, y)	(x, y)	(x, y)	(x, y)
$(+, +)$	$(-, +)$	$(-, -)$	$(+, -)$

- **Points on the Axes**
  - On the *x*-axis, y=0. Points with a *y*-coordinate equal to 0 are on the *x*-axis, and have coordinates (a,0).
  - On the *y*-axis, x=0. Points with an *x*-coordinate equal to 0 are on the *y*-axis, and have coordinates (0,b).
- **Solution of a Linear Equation**
  - An ordered pair (x,y) is a solution of the linear equation Ax+By=C, if the equation is a true statement when the *x*- and *y*-values of the ordered pair are substituted into the equation.

Glossary

linear equation

A linear equation is of the form Ax+By=C, where A and B are not both zero, is called a linear equation in two variables.

ordered pair

An ordered pair (x,y) gives the coordinates of a point in a rectangular coordinate system.

origin

The point (0,0) is called the origin. It is the point where the *x*-axis and *y*-axis intersect.

quadrant

The *x*-axis and the *y*-axis divide a plane into four regions, called quadrants.

rectangular coordinate system

A grid system is used in algebra to show a relationship between two variables; also called the *xy*-plane or the ‘coordinate plane’.

***x*-coordinate**

The first number in an ordered pair (x,y).

***y*-coordinate**

The second number in an ordered pair (x,y).

This page titled 4.1: Use the Rectangular Coordinate System is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.1: Use the Rectangular Coordinate System** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.1E: Exercises

Practice Makes Perfect


Plot Points in a Rectangular Coordinate System

In the following exercises, plot each point in a rectangular coordinate system and identify the quadrant in which the point is located.

Exercise 4.1E.1

1. (−4,2)
2. (−1,−2)
3. (3,−5)
4. (−3,5)
5. $(\frac{5}{3}, 2)$

Answer




Exercise 4.1E.2

1. (−2,−3)
2. (3,−3)
3. (−4,1)
4. (4,−1)
5. $(\frac{3}{2}, 1)$

Exercise 4.1E.3

1. (3,−1)
2. (−3,1)
3. (−2,2)
4. (−4,−3)
5. $\left(1, \frac{14}{5}\right)$

Answer



Exercise 4.1E.4


1. (−1,1)
2. (−2,−1)
3. (2,1)
4. (1,−4)
5. $\left(3, \frac{7}{2}\right)$

In the following exercises, plot each point in a rectangular coordinate system.

Exercise 4.1E.5

1. (−2,0)
2. (−3,0)
3. (0,0)
4. (0,4)
5. (0,2)

Answer




Exercise 4.1E.6

1. (0,1)
2. (0,−4)
3. (−1,0)
4. (0,0)
5. (5,0)

Exercise 4.1E.7

1. (0,0)
2. (0,−3)
3. (−4,0)
4. (1,0)
5. (0,−2)

Answer




Exercise 4.1E.8

1. (−3,0)
2. (0,5)
3. (0,−2)
4. (2,0)
5. (0,0)

In the following exercises, name the ordered pair of each point shown in the rectangular coordinate system.


Exercise 4.1E.9




Answer

$A:(-4,1)\quad B:(-3,-4)\quad C:(1,-3)\quad D:(4,3)$

Exercise 4.1E.10




Exercise 4.1E.11



Answer

$A:(0,-2)\quad B:(-2,0)\quad C:(0,5)\quad D:(5,0)$

Exercise 4.1E.12



Verify Solutions to an Equation in Two Variables

In the following exercises, which ordered pairs are solutions to the given equations?

Exercise 4.1E.13

2x+y=6

1. (1,4)
2. (3,0)
3. (2,3)

Answer

1, 2

Exercise 4.1E.14

1. x+3y=9
2. (0,3)
3. (6,1)
4. (−3,−3)

Exercise 4.1E.15

4x−2y=8

1. (3,2)
2. (1,4)
3. (0,−4)

Answer

1, 3

Exercise 4.1E.16

3x−2y=12

1. (4,0)
2. (2,−3)
3. (1,6)

Exercise 4.1E.17

y=4x+3

1. (4,3)
2. (−1,−1)
3. $\left(\frac{1}{2},5\right)$

Answer

2, 3

Exercise 4.1E.18

y=2x−5

1. (0,−5)
2. (2,1)
3. $\left(\frac{1}{2}, -4\right)$

Exercise 4.1E.19

$y = \frac{1}{2}x - 1$

1. (2,0)
2. (−6,−4)
3. (−4,−1)

Answer

1, 2

Exercise 4.1E.20

$y = \frac{1}{3}x + 1$

1. (−3,0)
2. (9,4)
3. (−6,−1)

Complete a Table of Solutions to a Linear Equation

In the following exercises, complete the table to find solutions to each linear equation.

Exercise 4.1E.21

$y = 2x - 4$

x	y	(x,y)
0		
2		
-1		

Answer

x	y	(x,y)
0	−4	(0,−4)
2	0	(2,0)
−1	−6	(−1,−6)

Exercise 4.1E.22

$y = 3x - 1$

x	y	(x,y)
0		
2		
-1		

Exercise 4.1E.23

$y = -x + 5$

x	y	(x,y)
0		
3		
-2		

Answer

x	y	(x,y)
0	5	(0,5)
3	2	(3,2)
−2	7	(−2,7)

Exercise 4.1E.24

$y = -x + 2$

x	y	(x,y)
0		
3		
-2		

Exercise 4.1E.25

$y = \frac{1}{3}x + 1$

x	y	(x,y)
0		
3		
6		

Answer

x	y	(x,y)
0	1	(0,1)
3	2	(3,2)
6	3	(6,3)

Exercise 4.1E.26

$y = \frac{1}{2}x + 4$

x	y	(x,
0		
2		
4		

Exercise 4.1E.27

$y = -\frac{3}{2}x - 2$

x	y	(x,y)
0		
2		
-2		

Answer

x	y	(x,y)
0	−2	(0,−2)
2	−5	(2,−5)
−2	1	(−2,1)

Exercise 4.1E.28

$y = -\frac{2}{3}x - 1$

x	y	(x,y)
0		
3		
-3		

Exercise 4.1E.29

$x + 3y = 6$

x	y	(x,y)
0		
3		

	0	

Answer

x	y	(x,y)
0	2	(0,2)
3	4	(3,1)
6	0	(6,0)

Exercise 4.1E.30

x+2y=8

x	y	(x,y)
0		
4		
	0	

Exercise 4.1E.31

$2x-5y=10$

x	y	(x,y)
0		
10		
	0	

Answer

x	y	(x,y)
0	−2	(0,−2)
10	2	(10,2)
5	0	(5,0)

Exercise 4.1E.32

x	y	(x,y)
0		
8		
	0	

Find Solutions to a Linear Equation

In the following exercises, find three solutions to each linear equation.

Exercise 4.1E.33

$y=5x-8$

Answer

Answers will vary.

Exercise 4.1E.34

$y=3x-9$

Exercise 4.1E.35

$y=-4x+5$

Answer

Answers will vary.

Exercise 4.1E.36

$y=-2x+7$

Exercise 4.1E.37

$x+y=8$

Answer

Answers will vary.

Exercise 4.1E.38

$x+y=6$

Exercise 4.1E.39

$x+y=-2$

Answer

Answers will vary.

Exercise 4.1E.40

$x+y=-1$

Exercise 4.1E.41

$3x+y=5$

Answer

Answers will vary.

Exercise 4.1E.42

$2x+y=3$

Exercise 4.1E.43

$4x-y=8$

Answer

Answers will vary.

Exercise 4.1E.44

$5x-y=10$

Exercise 4.1E.45

$2x+4y=8$

Answer

Answers will vary.

Exercise 4.1E.46

$3x+2y=6$

Exercise 4.1E.47

$5x-2y=10$

Answer

Answers will vary.

Exercise 4.1E.48


$4x-3y=12$

Everyday Math

Exercise 4.1E.49

Weight of a baby. Mackenzie recorded her baby's weight every two months. The baby's age, in months, and weight, in pounds, are listed in the table below, and shown as an ordered pair in the third column.

1. Plot the points on a coordinate plane.




2. Why is only Quadrant I needed?

Age x	Weight y	(x,y)
0	7	(0, 7)
2	11	(2, 11)
4	15	(4, 15)
6	16	(6, 16)
8	19	(8, 19)
10	20	(10, 20)
12	21	(12, 21)

Answer

1.




2. Age and weight are only positive.

Exercise 4.1E.50

Weight of a child. Latresha recorded her son's height and weight every year. His height, in inches, and weight, in pounds, are listed in the table below, and shown as an ordered pair in the third column.

1. Plot the points on a coordinate plane.



2. Why is only Quadrant I needed?

Height x	Weight y	(x,y)
28	22	(28, 22)
31	27	(31, 27)
33	33	(33, 33)
37	35	(37, 35)
40	41	(40, 41)
42	45	(42, 45)

Writing Exercises

Exercise 4.1E.51

Explain in words how you plot the point (4,−2) in a rectangular coordinate system?

Answer

Answers will vary.

Exercise 4.1E.52

How do you determine if an ordered pair is a solution to a given equation?

Exercise 4.1E.53

Is the point (−3,0) on the *x*-axis or *y*-axis? How do you know?

Answer

Answers will vary.

? Exercise $4.1E.54$

Is the point (0,8) on the *x*-axis or *y*-axis? How do you know?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
plot points in a rectangular coordinate system.			
identify points on a graph.			
verify solutions to an equation in two variables.			
complete a table of solutions to a linear equation.			
find solutions to a linear equation.			

ⓑ If most of your checks were:

…confidently. Congratulations! You have achieved the objectives in this section. Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific.

…with some help. This must be addressed quickly because topics you do not master become potholes in your road to success. In math every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no, I don't get it. This is a warning sign and you must not ignore it. You should get help right away or you will quickly be overwhelmed. See your instructor as soon as you can to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 4.1E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.1E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.2: Graph Linear Equations in Two Variables

Learning Objectives

By the end of this section, you will be able to:

- Recognize the relationship between the solutions of an equation and its graph.
- Graph a linear equation by plotting points.
- Graph vertical and horizontal lines.

Note

Before you get started, take this readiness quiz.

1. Evaluate $3x+2$ when $x=-1$.
   If you missed this problem, review Exercise 1.5.34.
2. Solve $3x+2y=12$ for y in general.
   If you missed this problem, review Exercise 2.6.16.

Recognize the Relationship Between the Solutions of an Equation and its Graph

In the previous section, we found several solutions to the equation $3x+2y=6$. They are listed in Table 4.2.1. So, the ordered pairs (0,3), (2,0), and $(1, \frac{3}{2})$ are some solutions to the equation $3x+2y=6$. We can plot these solutions in the rectangular coordinate system as shown in Figure 4.2.1.

Table 4.2.1

3x+2y=6		
x	**y**	**(x,y)**
0	3	(0,3)
2	0	(2,0)
1	$\frac{3}{2}$	$(1, \frac{3}{2})$




Figure 4.2.1

Notice how the points line up perfectly? We connect the points with a line to get the graph of the equation 3x+2y=6. See Figure 4.2.2. Notice the arrows on the ends of each side of the line. These arrows indicate the line continues.




Figure 4.2.2

Every point on the line is a solution of the equation. Also, every solution of this equation is a point on this line. Points *not* on the line are not solutions.

Notice that the point whose coordinates are (−2,6) is on the line shown in Figure 4.2.3. If you substitute x=−2 and y=6 into the equation, you find that it is a solution to the equation.




Figure 4.2.3

Test (−2, 6)

$$3x + 2y = 6$$

$$3(-2) + 2(6) = 6$$

$$-6 + 12 = 6$$

$$6 = 6 \checkmark$$

So the point (−2,6) is a solution to the equation $3x + 2y = 6$. (The phrase "the point whose coordinates are (−2,6)" is often shortened to "the point (−2,6).")

What about (4, 1) ?

$$3x + 2y = 6$$

$$3 \cdot 4 + 2 \cdot 1 = 6$$

$$12 + 2 \stackrel{?}{=} 6$$

$$14 \neq 6$$

Figure 4.2.3. This is an example of the saying, "A picture is worth a thousand words." The line shows you *all* the solutions to the equation. Every point on the line is a solution of the equation. And, every solution of this equation is on this line. This line is called the *graph* of the equation $3x + 2y = 6$.


GRAPH OF A LINEAR EQUATION

The **graph of a linear equation** Ax+By=C is a line.

- Every point on the line is a solution of the equation.
- Every solution of this equation is a point on this line.

Example 4.2.1

The graph of y=2x−3 is shown.



For each ordered pair, decide:

1. Is the ordered pair a solution to the equation?
2. Is the point on the line?

A (0,−3) B (3,3) C (2,−3) D (−1,−5)


Solution

Substitute the *x*- and *y*- values into the equation to check if the ordered pair is a solution to the equation.

1.

A: (0, –3)	B: (3, 3)	C: (2, –3)	D: (–1, –5)
$y = 2x - 3$	$y = 2x - 3$	$y = 2x - 3$	$y = 2x - 3$
$-3 \stackrel{?}{=} 2(0) - 3$	$3 \stackrel{?}{=} 2(3) - 3$	$-3 \stackrel{?}{=} 2(2) - 3$	$-5 \stackrel{?}{=} 2(-1) - 3$
$-3 = -3$ ✓	$3 = 3$ ✓	$-3 \neq 1$	$-5 = -5$ ✓
(0, –3) is a solution.	(3, 3) is a solution.	(2, –3) is not a solution.	(–1, –5) is a solution.

2. Plot the points A (0,−3), B (3,3), C (2,−3), and D (−1,−5).



The points that are solutions to y=2x−3 are on the line, but the point that is not a solution is not on the line.

The points (0,−3), (3,3), and (−1,−5) are on the line y=2x−3, and the point (2,−3) is not on the line.

Try It 4.2.2


Use the graph of y=3x−1 to decide whether each ordered pair is:

- a solution to the equation.

https://math.libretexts.org/@go/page/30505

- on the line.

1. (0,−1)
2. (2,5)



Answer


1. yes, yes
2. yes, yes

? Try It 4.2.3

Use graph of y=3x−1 to decide whether each ordered pair is:

- a solution to the equation
- on the line

1. (3,−1)
2. (−1,−4)



Answer

1. no, no
2. yes, yes


Graph a Linear Equation by Plotting Points


There are several methods that can be used to graph a linear equation. The method we used to graph 3x+2y=6 is called plotting points, or the Point–Plotting Method.

Example 4.2.4: How To Graph an Equation By Plotting Points

Graph the equation y=2x+1 by plotting points.

Solution


Step 1. Find three points whose coordinates are solutions to the equation.	You can choose any values for x or y. In this case, since y is isolated on the left side of the equation, it is easier to choose values for x.	$y = 2x + 1$ $x = 0$ $y = 2x + 1$ $y = 2 \cdot 0 + 1$ $y = 0 + 1$ $y = 1$ $x = 1$ $y = 2x + 1$ $y = 2 \cdot 1 + 1$ $y = 2 + 1$ $y = 3$ $x = -2$ $y = 2x + 1$ $y = 2(-2) + 1$ $y = -4 + 1$ $y = -3$
Organize the solutions in a table.	Put the three solutions in a table.	$y = 2x + 1$ x | y | (x, y) 0 | 1 | (0, 1) 1 | 3 | (1, 3) –2 | –3 | (–2, –3)
Step 2. Plot the points in a rectangular coordinate system. Check that the points line up. If they do not, carefully check your work!	Plot: (0, 1), (1, 3), (–2, –3). Do the points line up? Yes, the points line up.	

Step 3. Draw the line through the three points. Extend the line to fill the grid and put arrows on both ends of the line.	This line is the graph of $y = 2x + 1$.	

Try It 4.2.5

Graph the equation by plotting points: y=2x−3.


Answer



Try It 4.2.6

Graph the equation by plotting points: y=−2x+4.

Answer



The steps to take when graphing a linear equation by plotting points are summarized below.

GRAPH A LINEAR EQUATION BY PLOTTING POINTS

1. Find three points whose coordinates are solutions to the equation. Organize them in a table.
2. Plot the points in a rectangular coordinate system. Check that the points line up. If they do not, carefully check your work.
3. Draw the line through the three points. Extend the line to fill the grid and put arrows on both ends of the line.

It is true that it only takes two points to determine a line, but it is a good habit to use three points. If you only plot two points and one of them is incorrect, you can still draw a line but it will not represent the solutions to the equation. It will be the wrong line.

If you use three points, and one is incorrect, the points will not line up. This tells you something is wrong and you need to check your work. Look at the difference between part (a) and part (b) in Figure 4.2.4.




Figure 4.2.4

Let's do another example. This time, we'll show the last two steps all on one grid.

Example 4.2.7

Graph the equation y=−3x.

Solution

Find three points that are solutions to the equation. Here, again, it's easier to choose values for x. Do you see why?


$x = 0$	$x = 1$	$x = -2$
$y = -3x$	$y = -3x$	$y = -3x$
$y = -3 \cdot 0$	$y = -3 \cdot 1$	$y = -3(-2)$
$y = 0$	$y = -3$	$y = 6$

We list the points in Table 4.2.2.

Table 4.2.2

y=−3x		
x	**y**	**(x,y)**
0	0	(0,0)
1	−3	(1,−3)
−2	6	(−2,6)


Plot the points, check that they line up, and draw the line.



Try It 4.2.8

Graph the equation by plotting points: y=−4x.


Answer



Try It 4.2.9

Graph the equation by plotting points: y=x.

Answer



When an equation includes a fraction as the coefficient of x, we can still substitute any numbers for x. But the math is easier if we make ‘good’ choices for the values of x. This way we will avoid fraction answers, which are hard to graph precisely.

Example 4.2.10

Graph the equation $y = \frac{1}{2}x + 3$.

Solution

Find three points that are solutions to the equation. Since this equation has the fraction $\frac{1}{2}$ as a coefficient of x, we will choose values of x carefully. We will use zero as one choice and multiples of 2 for the other choices. Why are multiples of 2 a good choice for values of x?

$x = 0$	$x = 2$	$x = 4$
$y = \frac{1}{2}x + 3$	$y = \frac{1}{2}x + 3$	$y = \frac{1}{2}x + 3$
$y = \frac{1}{2}(0) + 3$	$y = \frac{1}{2}(2) + 3$	$y = \frac{1}{2}(4) + 3$
$y = 0 + 3$	$y = 1 + 3$	$y = 2 + 3$
$y = 3$	$y = 4$	$y = 5$


The points are shown in Table 4.2.3.

Table 4.2.3

y=12x+3

x	y	(x,y)
0	3	(0,3)
2	4	(2,4)
4	5	(4,5)


Plot the points, check that they line up, and draw the line.



Try It 4.2.11

Graph the equation $y = \frac{1}{3}x - 1$.


Answer



Try It 4.2.12

Graph the equation $y = \frac{1}{5}x + 2$.

Answer



So far, all the equations we graphed had y given in terms of x. Now we'll graph an equation with x and y on the same side. Let's see what happens in the equation 2x+y=3. If y=0 what is the value of x?

$$\begin{aligned} y &= 0 \\ 2x + y &= 3 \\ 2x + 0 &= 3 \\ 2x &= 3 \\ x &= \frac{3}{2} \\ \left(\frac{3}{2}, 0\right) & \end{aligned}$$

This point has a fraction for the *x*- coordinate and, while we could graph this point, it is hard to be precise graphing fractions. Remember in the example y=12x+3, we carefully chose values for x so as not to graph fractions at all. If we solve the equation 2x+y=3 for y, it will be easier to find three solutions to the equation.

$$\begin{aligned} 2x + y &= 3 \\ y &= -2x + 3 \end{aligned}$$

The solutions for x=0, x=1, and x=−1 are shown in the Table 4.2.4. The graph is shown in Figure 4.2.5.

Table 4.2.4

2x+y=3		
x	**y**	**(x,y)**
0	3	(0,3)
1	1	(1,1)
−1	5	(−1,5)




Figure 4.2.5

Can you locate the point $\left(\frac{3}{2}, 0\right)$ which we found by letting y=0, on the line?

Example 4.2.13

Graph the equation 3x+y=−1.

Solution

Find three points that are solutions to the equation.	$3x + y$	$=$	-1
First solve the equation for y.	y	$=$	$-3x - 1$

We'll let x be 0, 1, and −1 to find 3 points. The ordered pairs are shown in Table 4.2.5. Plot the points, check that they line up, and draw the line. See Figure 4.2.6.

Table 4.2.5

3x+y=−1		
x	**y**	**(x,y)**
0	−1	(0,−1)
1	−4	(1,−4)
−1	2	(−1,2)





Figure 4.2.6

Try It 4.2.14

Graph the equation 2x+y=2.


Answer



Try It 4.2.15

Graph the equation 4x+y=−3.

Answer



If you can choose any three points to graph a line, how will you know if your graph matches the one shown in the answers in the book? If the points where the graphs cross the *x*- and *y*-axis are the same, the graphs match!

The equation in Example 4.2.13 was written in standard form, with both x and y on the same side. We solved that equation for y in just one step. But for other equations in standard form it is not that easy to solve for y, so we will leave them in standard form. We can still find a first point to plot by letting x=0 and solving for y. We can plot a second point by letting y=0 and then solving for x. Then we will plot a third point by using some other value for x or y.

Example 4.2.16

Graph the equation $2x-3y=6$.

Solution

Find three points that are solutions to the equation.	$2x-3y = 6$
First let $x=0$.	$2x-3y = 6$
Solve for y.	$2(0)-3y = 6$
	$-3y = 6$
	$y = -2$
Now let $y=0$.	$2x-3(0) = 6$
Solve for x.	$2x = 6$
	$x = 3$
We need a third point. Remember, we can choose any value for x or y. We'll let x = 6.	$2(6)-3y = 6$
	$12-3y = 6$
Solve for y.	$-3y = -6$
	$y = 2$

We list the ordered pairs in Table 4.2.6. Plot the points, check that they line up, and draw the line. See Figure 4.2.7.

Table 4.2.6

2x−3y=6		
x	**yT**	**(x,y)**
0	−2	(0,−2)
3	0	(3,0)
6	2	(6,2)





Figure 4.2.7

Try It 4.2.17

Graph the equation $4x + 2y = 8$.


Answer



Try It 4.2.18

Graph the equation $2x - 4y = 8$.

Answer



Graph Vertical and Horizontal Lines

Can we graph an equation with only one variable? Just x and no y, or just y without an x? How will we make a table of values to get the points to plot?

Let's consider the equation x=−3. This equation has only one variable, x. The equation says that x is *always* equal to −3, so its value does not depend on y. No matter what y is, the value of x is always −3.

So to make a table of values, write −3 in for all the x values. Then choose any values for y. Since x does not depend on y, you can choose any numbers you like. But to fit the points on our coordinate graph, we'll use 1, 2, and 3 for the *y*-coordinates. See Table 4.2.7

Table 4.2.7

x=−3		
x	**y**	**(x,y)**
−3	1	(−3,1)
−3	2	(−3,2)
−3	3	(−3,3)

Plot the points from Table 4.2.7 and connect them with a straight line. Notice in Figure 4.2.8 that we have graphed a *vertical line*.




Figure 4.2.8

VERTICAL LINE

A **vertical line** is the graph of an equation of the form x=a.

The line passes through the *x*-axis at (a,0).

Example 4.2.19

Graph the equation x=2.

Solution

The equation has only one variable, x, and x is always equal to 2. We create Table 4.2.8 where x is always 2 and then put in any values for y. The graph is a vertical line passing through the *x*-axis at 2. See Figure 4.2.9.

Table 4.2.8

x=2		
x	**y**	**(x,y)**
2	1	(2,1)
2	2	(2,2)
2	3	(2,3)





Figure 4.2.9

Try It 4.2.20

Graph the equation x=5.


Answer



Try It 4.2.21

Graph the equation x=−2.

Answer



What if the equation has y but no x? Let's graph the equation y=4. This time the *y*- value is a constant, so in this equation, y does not depend on x. Fill in 4 for all the y's in Table 4.2.9 and then choose any values for x. We'll use 0, 2, and 4 for the *x*-coordinates.

Table 4.2.9

y=4		
x	**y**	**(x,y)**

0	4	(0,4)
2	4	(2,4)
4	4	(4,4)

The graph is a horizontal line passing through the *y*-axis at 4. See Figure 4.2.10.




Figure 4.2.10

HORIZONTAL LINE

A **horizontal line** is the graph of an equation of the form y=b.

The line passes through the *y*-axis at (0,b).

Example 4.2.22

Graph the equation y=−1.

Solution

The equation y=−1 has only one variable, y. The value of y is constant. All the ordered pairs in Table 4.2.10 have the same *y*-coordinate. The graph is a horizontal line passing through the *y*-axis at −1, as shown in Figure 4.2.11.

Table 4.2.10

y=−1		
x	**y**	**(x,y)**
	−1	(0,−1)
	−1	(3,−1)
−3	−1	(−3,−1)





Figure 4.2.11

Try It 4.2.23

Graph the equation y=−4.


Answer



Try It 4.2.24

Graph the equation y=3.

Answer



The equations for vertical and horizontal lines look very similar to equations like y=4x. What is the difference between the equations y=4x and y=4?

The equation y=4x has both x and y. The value of y depends on the value of x. The *y*-coordinate changes according to the value of x. The equation y=4 has only one variable. The value of y is constant. The *y*-coordinate is always 4. It does not depend on the value of x. See Table 4.2.11.

Table 4.2.11

y=4x				y=4		
x	**y**	**(x,y)**		**x**	**y**	**(x,y)**
0	0	(0,0)		0	4	(0,4)
1	4	(1,4)		1	4	(1,4)
2	8	(2,8)		2	4	(2,4)




Figure 4.2.12

Notice, in Figure 4.2.12, the equation y=4x gives a slanted line, while y=4 gives a horizontal line.

Example 4.2.25

Graph y=−3x and y=−3 in the same rectangular coordinate system.

Solution

Notice that the first equation has the variable x, while the second does not. See Table 4.2.12 The two graphs are shown in Figure 4.2.13

Table 4.2.12

y=−3x				y=−3		
x	**y**	**(x,y)**		**x**	**y**	**(x,y)**
		(0,0)			−3	(0,−3)
	−3	(1,−3)			−3	(1,−3)
	−6	(2,−6)			−3	(2,−3)





Figure 4.2.13

Try It 4.2.26

Graph y=−4x and y=−4 in the same rectangular coordinate system.


Answer



Try It 4.2.27

Graph y=3 and y=3x in the same rectangular coordinate system.

Answer



Key Concepts

- **Graph a Linear Equation by Plotting Points**
  1. Find three points whose coordinates are solutions to the equation. Organize them in a table.
  2. Plot the points in a rectangular coordinate system. Check that the points line up. If they do not, carefully check your work!
  3. Draw the line through the three points. Extend the line to fill the grid and put arrows on both ends of the line.

Glossary

graph of a linear equation

The graph of a linear equation Ax+By=C is a straight line. Every point on the line is a solution of the equation. Every solution of this equation is a point on this line.

horizontal line

A horizontal line is the graph of an equation of the form y=b. The line passes through the *y*-axis at (0,b).

vertical line

A vertical line is the graph of an equation of the form x=a. The line passes through the *x*-axis at (a,0).

This page titled 4.2: Graph Linear Equations in Two Variables is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.2: Graph Linear Equations in Two Variables** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.2E: Exercises

Practice Makes Perfect

Recognize the Relationship Between the Solutions of an Equation and its Graph


In the following exercises, for each ordered pair, decide:

1. Is the ordered pair a solution to the equation?
2. Is the point on the line?

Exercise 4.2E.1

y=x+2

1. (0,2)
2. (1,2)
3. (−1,1)
4. (−3,−1)




Answer

1. yes; no
2. no; no
3. yes; yes
4. yes; yes

Exercise 4.2E.2

y=x−4


1. (0,−4)
2. (3,−1)
3. (2,2)
4. (1,−5)



Exercise $4.2E.3$

$y = \frac{1}{2}x - 3$

1. (0,−3)
2. (2,−2)
3. (−2,−4)
4. (4,1)




Answer

1. yes; yes
2. yes; yes
3. yes; yes
4. no; no

Exercise $4.2E.4$

$y = \frac{1}{3}x + 2$

1. (0,2)
2. (3,3)
3. (−3,2)
4. (−6,0)




Graph a Linear Equation by Plotting Points

In the following exercises, graph by plotting points.

Exercise 4.2E.5

$y = 3x - 1$

Answer




Exercise 4.2E.6

$y = 2x + 3$

Exercise 4.2E.7

$y = -2x + 2$

Answer




Exercise 4.2E.8

$y = -3x + 1$

Exercise 4.2E.9

$y = x + 2$

Answer




Exercise 4.2E.10

$y = x - 3$

Exercise 4.2E.11

$y = -x - 3$

Answer




Exercise 4.2E.12

$y = -x - 2$

Exercise 4.2E.13

$y = 2x$

Answer




Exercise 4.2E.14

$y = 3x$

Exercise 4.2E.15

$y = -4x$

Answer




Exercise 4.2E.16

$y = -2x$

Exercise 4.2E.17

$y = \frac{1}{2}x + 2$

Answer




Exercise 4.2E.18

$y = \frac{1}{3}x - 1$

Exercise 4.2E.19

$y = \frac{4}{3}x - 5$

Answer




Exercise 4.2E.20

$y = \frac{3}{2}x - 3$

Exercise 4.2E.21

$y = -\frac{2}{5}x + 1$

Answer




Exercise 4.2E.22

$y = -\frac{4}{5}x - 1$

Exercise 4.2E.23

$y = -\frac{3}{2}x + 2$

Answer




Exercise 4.2E.24

$y = -\frac{5}{3}x + 4$

Exercise 4.2E.25

$x + y = 6$

Answer




Exercise 4.2E.26

$x + y = 4$

Exercise 4.2E.27

$x + y = -3$

Answer




Exercise 4.2E.28

$x + y = -2$

Exercise 4.2E.29

$x - y = 2$

Answer




Exercise 4.2E.30

$x - y = 1$

Exercise 4.2E.31

$x - y = -1$

Answer




Exercise 4.2E.32

$x - y = -3$

Exercise 4.2E.33

$3x + y = 7$

Answer




Exercise 4.2E.34

$5x + y = 6$

Exercise 4.2E.35

2x+y=−3

Answer




Exercise 4.2E.36

$4x + y = -5$

Exercise 4.2E.37

$\frac{1}{3}x + y = 2$

Answer




Exercise 4.2E.38

$\frac{1}{2}x + y = 3$

Exercise 4.2E.39

$\frac{2}{5}x - y = 4$

Answer




Exercise 4.2E.40

$\frac{3}{4}x - y = 6$

Exercise 4.2E.41

$2x + 3y = 12$

Answer




Exercise 4.2E.42

4x+2y=12

Exercise 4.2E.43

3x−4y=12

Answer




Exercise 4.2E.44

2x−5y=10

Exercise 4.2E.45

x−6y=3

Answer




Exercise 4.2E.46

x−4y=2

Exercise 4.2E.47

5x+2y=4

Answer



Exercise 4.2E.48

3x+5y=5


Graph Vertical and Horizontal Lines

In the following exercises, graph each equation.

Exercise 4.2E.49

x=4

Answer




Exercise 4.2E.50

x=3

Exercise 4.2E.51

x=−2

Answer




Exercise 4.2E.52

x=−5

Exercise 4.2E.53

y=3

Answer




Exercise 4.2E.54

y=1

Exercise 4.2E.55

y=−5

Answer




Exercise 4.2E.56

y=−2

Exercise 4.2E.57

$x = \frac{7}{3}$

Answer




Exercise 4.2E.58

$x = \frac{5}{4}$

Exercise 4.2E.59

$y = -\frac{15}{4}$

Answer



Exercise 4.2E.60


$y = -\frac{5}{3}$

In the following exercises, graph each pair of equations in the same rectangular coordinate system.

Exercise 4.2E.61

y=2x and y=2

Answer




Exercise 4.2E.62

y=5x and y=5

Exercise 4.2E.63

$y = -\frac{1}{2}x$ and $y = -\frac{1}{2}$

Answer



Exercise 4.2E.64

$y = -\frac{1}{3}x$ and $y = -\frac{1}{3}$


Mixed Practice

In the following exercises, graph each equation.

Exercise 4.2E.65

y=4x

Answer




Exercise 4.2E.66

y=2x

Exercise 4.2E.67

$y = -\frac{1}{2}x + 3$

Answer




Exercise 4.2E.68

$y = \frac{1}{4}x - 2$

Exercise 4.2E.69

y=−x

Answer




Exercise 4.2E.70

y=x

Exercise 4.2E.71

x−y=3

Answer




Exercise 4.2E.72

x+y=−5

Exercise 4.2E.73

4x+y=2

Answer




Exercise 4.2E.74

2x+y=6

Exercise 4.2E.75

y=−1

Answer




Exercise 4.2E.76

y=5

Exercise 4.2E.77

2x+6y=12

Answer




Exercise 4.2E.78

5x+2y=10

Exercise 4.2E.79

x=3

Answer



Exercise 4.2E.80

x=−4


Everyday Math

Exercise 4.2E.81

Motor home cost. The Robinsons rented a motor home for one week to go on vacation. It cost them $594 plus $0.32 per mile to rent the motor home, so the linear equation y=594+0.32x gives the cost, yy, for driving xx miles. Calculate the rental cost for driving 400, 800, and 1200 miles, and then graph the line.

Answer

$722, $850, $978



Exercise 4.2E.82

Weekly earnings. At the art gallery where he works, Salvador gets paid $200 per week plus 15% of the sales he makes, so the equation y=200+0.15x gives the amount, yy, he earns for selling x dollars of artwork. Calculate the amount Salvador earns for selling $900, $1600, and $2000, and then graph the line.

Writing Exercises

Exercise 4.2E.83

Explain how you would choose three x - values to make a table to graph the line $y = \frac{1}{5}x - 2$

Answer

Answers will vary.

Exercise $4.2E.84$

What is the difference between the equations of a vertical and a horizontal line?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
recognize the relation between the solutions of an equation and its graph.			
graph a linear equation by plotting points.			
graph vertical and horizontal lines.			

ⓑ After reviewing this checklist, what will you do to become confident for all goals?

This page titled 4.2E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.2E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.3: Graph with Intercepts

Learning Objectives

By the end of this section, you will be able to:

- Identify the x- and y- intercepts on a graph
- Find the x- and y- intercepts from an equation of a line
- Graph a line using the intercepts

Note

Before you get started, take this readiness quiz.

1. Solve: $3 \cdot 0 + 4y = -2$.
   If you missed this problem, review Exercise 2.2.13.

Identify the *x*- and *y*- Intercepts on a Graph


Every linear equation can be represented by a unique line that shows all the solutions of the equation. We have seen that when graphing a line by plotting points, you can use any three solutions to graph. This means that two people graphing the line might use different sets of three points.

At first glance, their two lines might not appear to be the same, since they would have different points labeled. But if all the work was done correctly, the lines should be exactly the same. One way to recognize that they are indeed the same line is to look at where the line crosses the *x*- axis and the *y*- axis. These points are called the *intercepts* of the line.

INTERCEPTS OF A LINE

The points where a line crosses the *x*- axis and the *y*- axis are called the **intercepts of a line**.

Let's look at the graphs of the lines in Figure 4.3.1.



(a) $2x + y = 6$ (b) $3x - 4y = 12$

(c) $x - y = 5$ (d) $y = -2x$

Figure 4.3.1: Examples of graphs crossing the x-negative axis.

First, notice where each of these lines crosses the x axis. See Figure 4.3.1.

Table 4.3.1

Figure	The line crosses the x-axis at:	Ordered pair of this point
Figure (a)	3	(3,0)
Figure (b)	4	(4,0)
Figure (c)	5	(5,0)
Figure (d)	0	(0,0)

Do you see a pattern?

For each row, the *y*- coordinate of the point where the line crosses the *x*- axis is zero. The point where the line crosses the *x*- axis has the form (a,0) and is called the ***x*- intercept of a line**. The *x*- intercept occurs when y is zero. Now, let's look at the points where these lines cross the *y*- axis. See Table 4.3.2.

Table 4.3.2

Figure	The line crosses the x-axis at:	Ordered pair of this point
Figure (a)	6	(0,6)
Figure (b)	−3	(0,−3)
Figure (c)	−5	(0,5)
Figure (d)	0	(0,0)

What is the pattern here?

In each row, the *x*- coordinate of the point where the line crosses the *y*- axis is zero. The point where the line crosses the *y*- axis has the form (0,b) and is called the *y*- *intercept* of the line. The *y*- intercept occurs when x is zero.

X- INTERCEPT AND Y- INTERCEPT OF A LINE

The *x*- intercept is the point (a,0) where the line crosses the *x*- axis.

The *y*- intercept is the point (0,b) where the line crosses the *y*- axis.

	x	*y*
• The *x*-intercept occurs when *y* is zero.	*a*	0
• The *y*-intercept occurs when *x* is zero.	0	*b*

Figure 4.3.2

Example 4.3.1

Find the *x*- and *y*- intercepts on each graph.




Figure 4.3.3

Solution


(a) The graph crosses the *x*- axis at the point (4,0). The *x*- intercept is (4,0).
The graph crosses the *y*- axis at the point (0,2). The *y*- intercept is (0,2).

(b) The graph crosses the x- axis at the point (2,0). The x- intercept is (2,0)
The graph crosses the y- axis at the point (0,−6). The y- intercept is (0,−6).

(c) The graph crosses the x- axis at the point (−5,0). The x- intercept is (−5,0).
The graph crosses the y- axis at the point (0,−5). The y- intercept is (0,−5).

Try It 4.3.2

Find the x- and y- intercepts on the graph.




Answer

x- intercept: (2,0); y- intercept: (0,−2)

Try It 4.3.3

Find the x- and y- intercepts on the graph.



Answer

x- intercept: (3,0), y- intercept: (0,2)

Find the x- and y- Intercepts from an Equation of a Line

Recognizing that the ***x*- intercept** occurs when y is zero and that the y- intercept occurs when x is zero, gives us a method to find the intercepts of a line from its equation. To find the x- intercept, let y=0 and solve for x. To find the ***y*- intercept**, let x=0 and solve for y.

X- AND Y- INTERCEPTS FROM THE EQUATION OF A LINE

Use the equation of the line. To find:

- the *x*- intercept of the line, let y=0 and solve for x.
- the *y*- intercept of the line, let x=0 and solve for y.

Example 4.3.4

Find the intercepts of 2x+y=6.

Solution

We will let y=0 to find the *x*- intercept, and let x=0 to find the *y*- intercept. We will fill in the table, which reminds us of what we need to find.

$2x+y=6$	
x	y
	0
0	

x-intercept

y-intercept

To find the *x*- intercept, let y=0.

Table 4.3.3

	$2x+y=6$
Let $y = 0$.	$2x+0=6$
Simplify.	$2x=6$
	$x=3$
The *x*-intercept is	(3, 0)
To find the *y*-intercept, let $x = 0$.	
	$2x+y=6$
Let $x = 0$.	$2\cdot 0+y=6$
Simplify.	$0+y=6$
	$y=6$
The *y*-intercept is	(0, 6)

The intercepts are the points (3,0) and (0,6) as shown in Table 4.3.4

Table 4.3.4

2x+y=6	
x	**y**
3	0
0	6

Try It 4.3.5

Find the intercepts of 3x+y=12.

Answer

x- intercept: (4,0), *y*- intercept: (0,12)

Try It 4.3.6

Find the intercepts of x+4y=8.

Answer

x- intercept: (8,0), *y*- intercept: (0,2)

Example 4.3.7

Find the intercepts of 4x–3y=12.

Solution

To find the *x*-intercept, let $y = 0$.	
	$4x - 3y = 12$
Let $y = 0$.	$4x - 3 \cdot 0 = 12$
Simplify.	$4x - 0 = 12$
	$4x = 12$
	$x = 3$
The *x*-intercept is	(3, 0)
To find the *y*-intercept, let $x = 0$.	
	$4x - 3y = 12$
Let $x = 0$.	$4 \cdot 0 - 3y = 12$
Simplify.	$0 - 3y = 12$
	$-3y = 12$
	$y = -4$
The *y*-intercept is	(0, −4)

Table 4.3.5

The intercepts are the points (3, 0) and (0, −4) as shown in the following table.

Table 4.3.6

4x−3y=12	
x	**y**
3	0
0	−4

Try It 4.3.8

Find the intercepts of 3x–4y=12.

Answer

x- intercept: (4,0), *y*- intercept: (0,−3)

Try It 4.3.9

Find the intercepts of 2x–4y=8.

Answer

x- intercept: (4,0), *y*- intercept: (0,−2)

Graph a Line Using the Intercepts

To graph a linear equation by plotting points, you need to find three points whose coordinates are solutions to the equation. You can use the *x*- and *y*- intercepts as two of your three points. Find the intercepts, and then find a third point to ensure accuracy. Make sure the points line up—then draw the line. This method is often the quickest way to graph a line.

Example 4.3.10: How to Graph a Line Using Intercepts


Graph –x+2y=6 using the intercepts.

Solution

Step 1. Find the *x*- and *y*-intercepts of the line. Let $y = 0$ and solve for x. Let $x = 0$ and solve for y.	Find the *x*-intercept.	Let $y = 0$ $-x + 2y = 6$ $-x + 2(0) = 6$ $-x = 6$ $x = -6$ The *x*-intercept is (–6, 0).
	Find the *y*-intercept.	Let $x = 0$ $-x + 2y = 6$ $-0 + 2y = 6$ $2y = 6$ $y = 3$ The *y*-intercept is (0, 3).
Step 2. Find another solution to the equation.	We'll use $x = 2$.	Let $x = 2$ $-x + 2y = 6$ $-2 + 2y = 6$ $2y = 8$ $y = 4$ A third point is (2, 4).


Step 3. Plot the three points. Check that the points line up.

x	y	(x, y)
–6	0	(–6, 0)
0	3	(0, 3)
2	4	(2, 4)



Step 4. Draw the line.


See the graph.



Try It 4.3.11

Graph x–2y=4 using the intercepts.


Answer



Try It 4.3.12

Graph –x+3y=6 using the intercepts.

Answer



The steps to graph a linear equation using the intercepts are summarized below.

GRAPH A LINEAR EQUATION USING THE INTERCEPTS.

1. Find the *x*- and *y*- intercepts of the line.
   - Let y=0 and solve for x
   - Let x=0 and solve for y.
2. Find a third solution to the equation.
3. Plot the three points and check that they line up.
4. Draw the line.

Example 4.3.13

Graph 4x–3y=12 using the intercepts.

Solution


Find the intercepts and a third point.

x-intercept, let $y = 0$	*y*-intercept, let $x = 0$	third point, let $y = 4$
$4x - 3y = 12$	$4x - 3y = 12$	$4x - 3y = 12$
$4x - 3(0) = 12$	$4(0) - 3y = 12$	$4x - 3(4) = 12$
$4x = 12$	$-3y = 12$	$4x - 12 = 12$
$x = 3$	$y = -4$	$4x = 24$
		$x = 6$

We list the points in Table 4.3.7 and show the graph below.

4x−3y=12		
x	**y**	**(x,y)**
3	0	(3,0)
0	−4	(0,−4)
6	4	(6,4)


Table 4.3.7



Try It 4.3.14

Graph 5x–2y=10 using the intercepts.


Answer



Try It 4.3.15

Graph 3x–4y=12 using the intercepts.

Answer



Example 4.3.16

Graph y=5x using the intercepts.

Solution

x-intercept	*y*-intercept
Let $y = 0$.	Let $x = 0$.
$y = 5x$	$y = 5x$
$0 = 5x$	$y = 5 \cdot 0$
$0 = x$	$y = 0$
(0, 0)	(0, 0)

This line has only one intercept. It is the point (0,0).

To ensure accuracy we need to plot three points. Since the x- and y- intercepts are the same point, we need *two* more points to graph the line.


Let $x = 1$.	Let $x = -1$.
$y = 5x$	$y = 5x$
$y = 5 \cdot 1$	$y = 5(-1)$
$y = 5$	$y = -5$

See Table 4.3.8.

y=5x		
x	**y**	**(x,y)**
0	0	(0,0)
1	5	(1,5)
−1	−5	(−1,−5)

Table 4.3.8


Plot the three points, check that they line up, and draw the line.



Try It 4.3.17

Graph y=4x using the intercepts.


Answer



Try It 4.3.18

Graph y=−x the intercepts.

Answer



Key Concepts

- **Find the *x*- and *y*- Intercepts from the Equation of a Line**
  - Use the equation of the line to find the *x*- intercept of the line, let y=0 and solve for *x*.
  - Use the equation of the line to find the *y*- intercept of the line, let x=0 and solve for *y*.
- **Graph a Linear Equation using the Intercepts**
  1. Find the *x*- and *y*- intercepts of the line.
     Let y=0 and solve for *x*.
     Let x=0 and solve for *y*.
  2. Find a third solution to the equation.
  3. Plot the three points and then check that they line up.
  4. Draw the line.
- **Strategy for Choosing the Most Convenient Method to Graph a Line:**
  - Consider the form of the equation.
  - If it only has one variable, it is a vertical or horizontal line.
    x=a is a vertical line passing through the *x*- axis at a
    y=b is a horizontal line passing through the *y*- axis at b.
  - If *y* is isolated on one side of the equation, graph by plotting points.
  - Choose any three values for *x* and then solve for the corresponding *y*- values.
  - If the equation is of the form ax+by=c, find the intercepts. Find the *x*- and *y*- intercepts and then a third point.

Glossary

intercepts of a line

The points where a line crosses the *x*- axis and the *y*- axis are called the intercepts of the line.

***x*- intercept**

The point (a,0) where the line crosses the *x*- axis; the *x*- intercept occurs when y is zero.

***y*-intercept**

The point (0,b) where the line crosses the *y*- axis; the *y*- intercept occurs when x is zero.

This page titled 4.3: Graph with Intercepts is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.3: Graph with Intercepts** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.


4.3E: Exercises

Practice Makes Perfect

Identify the *x*- and *y*- Intercepts on a Graph

In the following exercises, find the *x*- and *y*- intercepts on each graph.


Exercise 4.3E.1




Answer

(3,0),(0,3)

Exercise 4.3E.2




Exercise 4.3E.3




Answer

(5,0),(0,−5)

Exercise 4.3E.4




Exercise 4.3E.5




Answer

(−2,0),(0,−2)

Exercise 4.3E.6




Exercise 4.3E.7




Answer

(−1,0),(0,1)

Exercise 4.3E.8




Exercise 4.3E.9




Answer

(6,0),(0,3)

Exercise 4.3E.10




Exercise 4.3E.11



Answer

(0,0)

Exercise 4.3E.12



Find the *x*- and *y*- Intercepts from an Equation of a Line

In the following exercises, find the intercepts for each equation.

Exercise 4.3E.13

x+y=4

Answer

(4,0),(0,4)

Exercise 4.3E.14

x+y=3

Exercise 4.3E.15

x+y=−2

Answer

(−2,0),(0,−2)

Exercise 4.3E.16

x+y=−5

Exercise 4.3E.17

x–y=5

Answer

(5,0),(0,−5)

Exercise 4.3E.18

x–y=1

Exercise 4.3E.19

x–y=−3

Answer

(−3,0),(0,3)

Exercise 4.3E.20

x–y=−4

Exercise 4.3E.21

x+2y=8

Answer

(8,0),(0,4)

Exercise 4.3E.22

x+2y=10

Exercise 4.3E.23

3x+y=6

Answer

(2,0),(0,6)

Exercise 4.3E.24

3x+y=9

Exercise 4.3E.25

x–3y=12

Answer

(12,0),(0,−4)

Exercise 4.3E.26

x–2y=8

Exercise 4.3E.27

4x–y=8

Answer

(2,0),(0,−8)

Exercise 4.3E.28

5x–y=5

Exercise 4.3E.29

2x+5y=10

Answer

(5,0),(0,2)

Exercise 4.3E.30

2x+3y=6

Exercise 4.3E.31

3x–2y=12

Answer

(4,0),(0,−6)

Exercise 4.3E.32

3x–5y=30

Exercise 4.3E.33

$y = \frac{1}{3}x + 1$

Answer

(-3,0),(0,1)

Exercise 4.3E.34

$y = \frac{1}{4}x - 1$

Exercise 4.3E.35

$y = \frac{1}{5}x + 2$

Answer

(−10,0),(0,2)

Exercise 4.3E.36

$y = \frac{1}{3}x + 4$

Exercise 4.3E.37

y=3x

Answer

(0,0)

Exercise 4.3E.38

y=-2x

Exercise 4.3E.39

y=-4x

Answer

(0,0)

Exercise 4.3E. 40

y=5x


Graph a Line Using the Intercepts

In the following exercises, graph using the intercepts.

Exercise 4.3E. 41

$-x+5y=10$

Answer




Exercise 4.3E. 42

$-x+4y=8$

Exercise 4.3E. 43

$x+2y=4$

Answer




Exercise 4.3E. 44

$x+2y=6$

Exercise 4.3E. 45

$x+y=2$

Answer




Exercise 4.3E. 46

$x + y = 5$

Exercise 4.3E. 47

$x + y = -3$

Answer




Exercise 4.3E. 48

$x + y = -1$

Exercise 4.3E. 49

$x - y = 1$

Answer




Exercise 4.3E.50

$x - y = 2$

Exercise 4.3E.51

$x - y = -4$

Answer




Exercise 4.3E.52

$x - y = -3$

Exercise 4.3E.53

$4x + y = 4$

Answer




Exercise 4.3E.54

$3x + y = 3$

Exercise 4.3E.55

$2x + 4y = 12$

Answer




Exercise 4.3E.56

$3x + 2y = 12$

Exercise 4.3E.57

$3x - 2y = 6$

Answer




Exercise 4.3E.58

$5x - 2y = 10$

Exercise 4.3E.59

$2x - 5y = -20$

Answer




Exercise 4.3E. 60

$3x - 4y = -12$

Exercise 4.3E. 61

$3x - y = -6$

Answer




Exercise 4.3E. 62

$2x - y = -8$

Exercise 4.3E. 63

$y = -2x$

Answer




Exercise 4.3E. 64

$y = -4x$

Exercise 4.3E. 65

$y = x$

Answer




Exercise 4.3E.66

y=3x

Everyday Math

Exercise 4.3E.67

Road trip. Damien is driving from Chicago to Denver, a distance of 1000 miles. The *x*- axis on the graph below shows the time in hours since Damien left Chicago. The *y*- axis represents the distance he has left to drive.




1. Find the *x*- and *y*- intercepts.
2. Explain what the *x*- and *y*- intercepts mean for Damien.

Answer

1. (0,1000),(15,0)
2. At (0,1000), he has been gone 0 hours and has 1000 miles left. At (15,0), he has been gone 15 hours and has 0 miles left to go.

Exercise 4.3E.68

Road trip. Ozzie filled up the gas tank of his truck and headed out on a road trip. The *x*- axis on the graph below shows the number of miles Ozzie drove since filling up. The *y*- axis represents the number of gallons of gas in the truck's gas tank.



1. Find the *x*- and *y*- intercepts.
2. Explain what the *x*- and *y*- intercepts mean for Ozzie.

Writing Exercises

Exercise $4.3E.69$

How do you find the x -intercept of the graph of $3x-2y=6$?

Answer

Answers will vary.

Exercise $4.3E.70$

Do you prefer to use the method of plotting points or the method using the intercepts to graph the equation 4x+y=−4? Why?

Exercise $4.3E.71$

Do you prefer to use the method of plotting points or the method using the intercepts to graph the equation $y=\frac{2}{3}x-2$? Why?

Answer

Answers will vary.

Exercise $4.3E.72$

Do you prefer to use the method of plotting points or the method using the intercepts to graph the equation y=6? Why?

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
identify the *x* and *y* intercepts on a graph.			
find the *x* and *y* intercepts from an equation of a line.			
graph a line using the intercepts.			

ⓑ What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 4.3E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.3E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.4: Understanding the Slope of a Line

Learning Objectives

- By the end of this section, you will be able to:
- Use geoboards to model slope
- Use $m = \frac{\text{rise}}{\text{run}}$ to find the slope of a line from its graph
- Find the slope of horizontal and vertical lines
- Use the slope formula to find the slope of a line between two points
- Graph a line given a point and the slope
- Solve slope applications

Note

Before you get started, take this readiness quiz.

1. Simplify: $\frac{1-4}{8-2}$.
   If you missed this problem, review Exercise 1.6.31
2. Divide: $\frac{0}{4}, \frac{4}{0}$.
   If you missed this problem, review Exercise 1.10.16.
3. Simplify: $\frac{15}{-3}, \frac{-15}{3}, \frac{-15}{-3}$.
   If you missed this problem, review Exercise 1.6.4.

When you graph linear equations, you may notice that some lines tilt up as they go from left to right and some lines tilt down. Some lines are very steep and some lines are flatter. What determines whether a line tilts up or down or if it is steep or flat?

In mathematics, the 'tilt' of a line is called the *slope* of the line. The concept of slope has many applications in the real world. The pitch of a roof, grade of a highway, and a ramp for a wheelchair are some examples where you literally see slopes. And when you ride a bicycle, you feel the slope as you pump uphill or coast downhill.

In this section, we will explore the concept of slope.

Use Geoboards to Model Slope

A **geoboard** is a board with a grid of pegs on it. Using rubber bands on a geoboard gives us a concrete way to model lines on a coordinate grid. By stretching a rubber band between two pegs on a geoboard, we can discover how to find the slope of a line.

Doing the Manipulative Mathematics activity "Exploring Slope" will help you develop a better understanding of the slope of a line. (Graph paper can be used instead of a geoboard, if needed.)

We'll start by stretching a rubber band between two pegs as shown in Figure 4.4.1.




Figure 4.4.1

Doesn't it look like a line?

Now we stretch one part of the rubber band straight up from the left peg and around a third peg to make the sides of a right triangle, as shown in Figure 4.4.2




Figure 4.4.2

We carefully make a 90º angle around the third peg, so one of the newly formed lines is vertical and the other is horizontal.

To find the slope of the line, we measure the distance along the vertical and horizontal sides of the triangle. The vertical distance is called the **rise** and the horizontal distance is called the **run**, as shown in Figure 4.4.3.




Figure 4.4.3

If our **geoboard** and rubber band look just like the one shown in Figure 4.4.4, the rise is 2. The rubber band goes up 2 units. (Each space is one unit.)




Figure 4.4.4

The rise on this geoboard is 2, as the rubber band goes up two units.

What is the run?

The rubber band goes across 3 units. The run is 3 (see Figure 4.4.4).

The slope of a line is the ratio of the rise to the run. In mathematics, it is always referred to with the letter m.

SLOPE OF A LINE

The **slope of a line** of a line is $m = \frac{\text{rise}}{\text{run}}$.

The **rise** measures the vertical change and the **run** measures the horizontal change between two points on the line.

What is the slope of the line on the geoboard in Figure 4.4.4?


$$m = \frac{\text{rise}}{\text{run}}$$
$$m = \frac{2}{3}$$

The line has slope $\frac{2}{3}$. This means that the line rises 2 units for every 3 units of run.

When we work with geoboards, it is a good idea to get in the habit of starting at a peg on the left and connecting to a peg to the right. If the rise goes up it is positive and if it goes down it is negative. The run will go from left to right and be positive.

Example 4.4.1


What is the slope of the line on the geoboard shown?



Solution

Use the definition of slope: $m = \frac{\text{rise}}{\text{run}}$.

Start at the left peg and count the spaces up and to the right to reach the second peg.




$$\begin{aligned} &\text{The rise is } 3. && m = \tfrac{3}{\text{run}} \\ &\text{The run is } 4. && m = \tfrac{3}{4} \\ & && \text{The slope is } \tfrac{3}{4}. \end{aligned} \tag{4.4.1}$$

This means that the line rises 3 units for every 4 units of run.

Try It 4.4.2

What is the slope of the line on the geoboard shown?




Answer

$\frac{4}{3}$

Try It 4.4.3

What is the slope of the line on the geoboard shown?




Answer

$\frac{1}{4}$

Example 4.4.4


What is the slope of the line on the geoboard shown?



Solution

Use the definition of slope: $m = \frac{\text{rise}}{\text{run}}$.

Start at the left peg and count the units down and to the right to reach the second peg.




$$\begin{aligned} &\text{The rise is } -1. && m = \frac{-1}{\text{run}} \\ &\text{The run is } 3. && m = \frac{-1}{3} \\ & && m = -\frac{1}{3} \\ & && \text{The slope is } -\frac{1}{3} \end{aligned} \tag{4.4.2}$$

This means that the line drops 1 unit for every 3 units of run.

Try It 4.4.5

What is the slope of the line on the geoboard?




Answer

$-\frac{2}{3}$

Try It 4.4.6

What is the slope of the line on the geoboard?



Answer

$-\frac{4}{3}$

Notice that in Exercise 4.4.1 the slope is positive and in Exercise 4.4.4 the slope is negative. Do you notice any difference in the two lines shown in Figure(a) and Figure(b)?




Figure 4.4.5

We 'read' a line from left to right just like we read words in English. As you read from left to right, the line in Figure(a) is going up; it has **positive slope**. The line in Figure(b) is going down; it has **negative slope**.

POSITIVE AND NEGATIVE SLOPES




Figure 4.4.6

Example 4.4.7


Use a geoboard to model a line with slope $\frac{1}{2}$.

Solution

To model a line on a geoboard, we need the rise and the run.

Use the slope formula.	$m = \frac{\text{rise}}{\text{run}}$
Replace m with $\frac{1}{2}$.	$\frac{1}{2} = \frac{\text{rise}}{\text{run}}$

So, the rise is 1 and the run is 2 .
Start at a peg in the lower left of the geoboard.
Stretch the rubber band up 1 unit, and then right 2 units.




The hypotenuse of the right triangle formed by the rubber band represents a line whose slope is $\frac{1}{2}$.

Try It 4.4.8

Model the slope $m = \frac{1}{3}$. Draw a picture to show your results.


Answer



Try It 4.4.9

Model the slope $m = \frac{3}{2}$. Draw a picture to show your results.

Answer




Example 4.4.10

Use a geoboard to model a line with slope $\frac{-1}{4}$.

Solution

Use the slope formula. m = $\frac{\text{rise}}{\text{run}}$.
Replace m with $\frac{-1}{4}$.
So, the rise is -1 and the run is 4.
Since the rise is negative, we choose a starting peg on the upper left that will give us room to count down.
We stretch the rubber band down 1 unit, and then right 4 units.




The hypotenuse of the right triangle formed by the rubber band represents a line whose slope is $\frac{-1}{4}$.

Try It 4.4.11

Model the slope $m = -2 = \frac{-2}{1}$. Draw a picture to show your results.


Answer



Try It 4.4.12

Model the slope $m = \frac{-1}{3}$. Draw a picture to show your results.

Answer



Use $m = \frac{\text{rise}}{\text{run}}$ to Find the Slope of a Line from its Graph


Now, we'll look at some graphs on the xy-coordinate plane and see how to find their slopes. The method will be very similar to what we just modeled on our geoboards.

To find the slope, we must count out the rise and the run. But where do we start?




We locate two points on the line whose coordinates are integers. We then start with the point on the left and sketch a right triangle, so we can count the rise and run.

Example 4.4.13:

Find the slope of the line shown.




Solution

Step 1. Locate two points on the graph whose coordinates are integers.	Mark (0, –3) and (5, 1).	
Step 2. Starting with the point on the left, sketch a right triangle, going from the first point to the second point.	Starting at (0, –3), sketch a right triangle to (5, 1).	
Step 3. Count the rise and the run on the legs of the triangle.	Count the rise. Count the run.	 The rise is 4. The run is 5.
Step 4. Take the ratio of rise to run to find the slope. $m = \frac{\text{rise}}{\text{run}}$	Use the slope formula. Substitute the values of the rise and run.	$m = \frac{\text{rise}}{\text{run}}$ $m = \frac{4}{5}$ The slope of the line is $\frac{4}{5}$. This means that y increases 4 units as x increases 5 units.

Try It 4.4.14

Find the slope of the line shown.




Answer

$\frac{2}{5}$

https://math.libretexts.org/@go/page/30509

Try It 4.4.15

Find the slope of the line shown.



Answer


$\frac{3}{4}$

FIND THE SLOPE OF A LINE FROM ITS GRAPH


1. Locate two points on the line whose coordinates are integers.
2. Starting with the point on the left, sketch a right triangle, going from the first point to the second point.
3. Count the rise and the run on the legs of the triangle.
4. Take the ratio of rise to run to find the slope, $m = \frac{\text{rise}}{\text{run}}$.

Example 4.4.16

Find the slope of the line shown.




Solution

Locate two points on the graph whose coordinates are integers.	(0,5) and (3,3)
Which point is on the left?	(0,5)
Starting at (0,5), sketch a right triangle to (3,3).	
Count the rise—it is negative.	The rise is −2.
Count the run.	The run is 3.

Use the slope formula.	$m = \frac{\text{rise}}{\text{run}}$
Substitute the values of the rise and run.	$m = \frac{-2}{3}$
Simplify.	$m = -\frac{2}{3}$
	The slope of the line is $-\frac{2}{3}$.

So y increases by 3 units as x decreases by 2 units.


What if we used the points (−3,7) and (6,1) to find the slope of the line?



The rise would be −6 and the run would be 9. Then $m = \frac{-6}{9}$, and that simplifies to $m = -\frac{2}{3}$. Remember, it does not matter which points you use—the slope of the line is always the same.

Try It 4.4.17

Find the slope of the line shown.




Answer

$-\frac{4}{3}$

Try It 4.4.18

Find the slope of the line shown.




Answer

$-\frac{3}{5}$


In the last two examples, the lines had *y*-intercepts with integer values, so it was convenient to use the *y*-intercept as one of the points to find the slope. In the next example, the *y*-intercept is a fraction. Instead of using that point, we'll look for two other points whose coordinates are integers. This will make the slope calculations easier.

Example 4.4.19

Find the slope of the line shown.



Solution


Locate two points on the graph whose coordinates are integers.	(2,3) and (7,6)
Which point is on the left?	(2,3)
Starting at (2,3), sketch a right triangle to (7,6).	
Count the rise.	The rise is 3.
Count the run.	The run is 5.
Use the slope formula.	$m = \frac{\text{rise}}{\text{run}}$
Substitute the values of the rise and run.	$m = \frac{3}{5}$
	The slope of the line is $\frac{3}{5}$.

This means that y increases 5 units as x increases 3 units.

When we used geoboards to introduce the concept of slope, we said that we would always start with the point on the left and count the rise and the run to get to the point on the right. That way the run was always positive and the rise determined whether the slope was positive or negative.

What would happen if we started with the point on the right?

Let's use the points (2,3) and (7,6) again, but now we'll start at (7,6).




Count the rise.	The rise is -3.
Count the run. It goes from right to left, so it is negative.	The run is-5.
Use the slope formula.	$m = \frac{\text{rise}}{\text{run}}$
Substitute the values of the rise and run.	$m = \frac{-3}{-5}$
	The slope of the line is $\frac{3}{5}$

It does not matter where you start—the slope of the line is always the same.

? Try It 4.4.20

Find the slope of the line shown.




Answer

$\frac{5}{4}$

? Try It 4.4.21

Find the slope of the line shown.



Answer

$\frac{3}{2}$

Find the Slope of Horizontal and Vertical Lines

Do you remember what was special about horizontal and vertical lines? Their equations had just one variable.

Horizontal line $y = b$	**Vertical line** $x = a$
y -coordinates are the same.	x -coordinates are the same.

(4.4.3)

So how do we find the slope of the horizontal line y=4? One approach would be to graph the horizontal line, find two points on it, and count the rise and the run. Let's see what happens when we do this.




Figure 4.4.7

What is the rise?	The rise is 0.
What is the run?	The run is 3.
	$m = \frac{\text{rise}}{\text{run}}$
	$m = \frac{0}{3}$
What is the slope?	$m = 0$
	The slope of the horizontal line y = 4 is 0.

All horizontal lines have slope 0. When the *y*-coordinates are the same, the rise is 0.

SLOPE OF A HORIZONTAL LINE

The slope of a horizontal line, y=b, is 0.

The floor of your room is horizontal. Its slope is 0. If you carefully placed a ball on the floor, it would not roll away.

Now, we'll consider a vertical line, the line.




Figure 4.4.8

What is the rise?	The rise is 2.
What is the run?	The run is 0.
	$m = \frac{\text{rise}}{\text{run}}$
What is the slope?	$m = \frac{2}{0}$

But we can't divide by 0. Division by 0 is not defined. So we say that the slope of the vertical line x=3 is undefined.

The slope of any vertical line is undefined. When the *x*-coordinates of a line are all the same, the run is 0.

SLOPE OF A VERTICAL LINE

The slope of a vertical line, x=a, is undefined.

Example 4.4.22

Find the slope of each line:

(a) x=8 (b) y=−5.

Solution

(a) x=8
This is a vertical line.
Its slope is undefined.

(b) y=−5
This is a horizontal line.
It has slope 0.

Try It 4.4.23

Find the slope of the line: x=−4.

Answer

undefined

Try It 4.4.24

Find the slope of the line: y=7.

Answer

0

QUICK GUIDE TO THE SLOPES OF LINES




Figure 4.4.9

Remember, we ‘read’ a line from left to right, just like we read written words in English.

Use the Slope Formula to find the Slope of a Line Between Two Points

Doing the Manipulative Mathematics activity “Slope of Lines Between Two Points” will help you develop a better understanding of how to find the slope of a line between two points.

Sometimes we’ll need to find the slope of a line between two points when we don’t have a graph to count out the rise and the run. We could plot the points on grid paper, then count out the rise and the run, but as we’ll see, there is a way to find the slope without graphing. Before we get to it, we need to introduce some algebraic notation.

We have seen that an ordered pair (x,y) gives the coordinates of a point. But when we work with slopes, we use two points. How can the same symbol (x,y) be used to represent two different points? Mathematicians use subscripts to distinguish the points.

$$\begin{aligned}(x_1, y_1) &\quad \text{read } `x \text{ sub } 1, y \text{ sub } 1'\\ (x_2, y_2) &\quad \text{read } `x \text{ sub } 2, y \text{ sub } 2'\end{aligned} \tag{4.4.4}$$

The use of subscripts in math is very much like the use of last name initials in elementary school. Maybe you remember Laura C. and Laura M. in your third grade class?

We will use (x_1, y_1) to identify the first point and (x_2, y_2) to identify the second point.

If we had more than two points, we could use (x_3, y_3), (x_4, y_4), and so on.

Let's see how the rise and run relate to the coordinates of the two points by taking another look at the slope of the line between the points (2,3) and (7,6).




Figure 4.4.10

Since we have two points, we will use subscript notation, $\begin{pmatrix} x_1, y_1 \\ 2, 3 \end{pmatrix} \begin{pmatrix} x_2, y_2 \\ 7, 6 \end{pmatrix}$.

On the graph, we counted the rise of 3 and the run of 5.

Notice that the rise of 3 can be found by subtracting the *y*-coordinates 6 and 3.

$$3 = 6 - 3 \tag{4.4.5}$$

And the run of 5 can be found by subtracting the *x*-coordinates 7 and 2.

$$5 = 7 - 2 \tag{4.4.6}$$

We know $m = \frac{\text{rise}}{\text{run}}$. So $m = \frac{3}{5}$.

We rewrite the rise and run by putting in the coordinates $m = \frac{6-3}{7-2}$

But 6 is y_2, the *y*-coordinate of the second point and 3 is y_1, the *y*-coordinate of the first point.

So we can rewrite the slope using subscript notation. $m = \frac{y2-y1}{7-2}$

Also, 7 is x_2, the *x*-coordinate of the second point and 2 is x_1, the *x*-coordinate of the first point.

So, again, we rewrite the slope using subscript notation. $m = \frac{y2-y1}{x2-x1}$

We've shown that $m = \frac{y2-y1}{x2-x1}$ is really another version of $m = \frac{\text{rise}}{\text{run}}$. We can use this formula to find the slope of a line when we have two points on the line.

SLOPE FORMULA

The slope of the line between two points (x_1, y_1) and (x_2, y_2) is

$$m = \frac{y_2 - y_1}{x_2 - x_1} \tag{4.4.7}$$

This is the **slope formula**.

The slope is:

y of the second point minus y of the first point
over
x of the second point minus x of the first point. (4.4.8)


Example 4.4.25

Use the **slope formula** to find the slope of the line between the points (1,2) and (4,5).

Solution

We'll call (1,2) point #1 and (4,5) point #2.	$\begin{pmatrix} x_1, y_1 \\ 1, 2 \end{pmatrix} \begin{pmatrix} x_2, y_2 \\ 4, 5 \end{pmatrix}$
Use the slope formula.	$m = \frac{y_2 - y_1}{x_2 - x_1}$
Substitute the values.	
y of the second point minus y of the first point	$m = \frac{5-2}{x_2 - x_1}$
x of the second point minus x of the first point	$m = \frac{5-2}{4-1}$
Simplify the numerator and the denominator.	$m = \frac{3}{3}$
Simplify.	$m = 1$

Let's confirm this by counting out the slope on a graph using $m = \frac{\text{rise}}{\text{run}}$.



It doesn't matter which point you call point #1 and which one you call point #2. The slope will be the same. Try the calculation yourself.

Try It 4.4.26

Use the slope formula to find the slope of the line through the points: (8,5) and (6,3).

Answer

1

Try It 4.4.27

Use the slope formula to find the slope of the line through the points: (1,5) and (5,9).

Answer

1

Example 4.4.28

Use the slope formula to find the slope of the line through the points (−2,−3) and (−7,4).

Solution

We'll call (-2, -3) point #1 and (-7,4) point #2. $\begin{pmatrix} x_1, y_1 \\ -2, -3 \end{pmatrix} \begin{pmatrix} x_2, y_2 \\ -7, 4 \end{pmatrix}$

Use the slope formula. $m = \frac{y_2 - y_1}{x_2 - x_1}$

Substitute the values.


y of the second point minus y of the first point $m = \frac{4-(-3)}{x_2 - x_1}$

x of the second point minus x of the first point $m = \frac{4-(-3)}{-7-(-2)}$

Simplify the numerator and the denominator. $m = \frac{7}{-5}$

Simplify. $m = -\frac{7}{5}$

Let's verify this slope on the graph shown.



$$m = \frac{\text{rise}}{\text{run}}$$
$$m = \frac{-7}{5}$$
$$m = -\frac{7}{5}$$

Try It 4.4.29

Use the slope formula to find the slope of the line through the points: (−3,4) and (2,−1).

Answer

-1

Try It 4.4.30

Use the slope formula to find the slope of the line through the pair of points: (−2,6) and (−3,−4).

Answer

10

Graph a Line Given a Point and the Slope




Up to now, in this chapter, we have graphed lines by plotting points, by using intercepts, and by recognizing horizontal and vertical lines.

One other method we can use to graph lines is called the **point–slope method**. We will use this method when we know one point and the slope of the line. We will start by plotting the point and then use the definition of slope to draw the graph of the line.

Example 4.4.31

Graph the line passing through the point $(1,-1)$ whose slope is $m = \frac{3}{4}$.


Solution

Step 1. Plot the given point.	Plot (1, –1).	
Step 2. Use the slope formula $m = \frac{\text{rise}}{\text{run}}$ to identify the rise and the run.	Identify the rise and the run.	$m = \frac{3}{4}$ $\frac{\text{rise}}{\text{run}} = \frac{3}{4}$ rise = 3 run = 4
Step 3. Starting at the given point, count out the rise and run to mark the second point.	Start at (1, –1) and count the rise and the run. Up 3 units, right 4 units.	
Step 4. Connect the points with a line.	Connect the two points with a line.	

Try It 4.4.32

Graph the line passing through the point $(2,-2)$ with the slope $m = \frac{4}{3}$.


Answer



Try It 4.4.33

Graph the line passing through the point (−2,3) with the slope $m = \frac{1}{4}$.

Answer



GRAPH A LINE GIVEN A POINT AND THE SLOPE.


1. Plot the given point.
2. Use the slope formula $m = \frac{\text{rise}}{\text{rise}}$ to identify the rise and the run.
3. Starting at the given point, count out the rise and run to mark the second point.
4. Connect the points with a line.

Example 4.4.34

Graph the line with *y*-intercept 2 whose slope is $m = -\frac{2}{3}$.

Solution

Plot the given point, the *y*-intercept, (0,2).




https://math.libretexts.org/@go/page/30509


Identify the rise and the run. $m = -\frac{2}{3}$

$$\frac{\text{rise}}{\text{run}} = \frac{-2}{3}$$
$$\text{rise} = -2$$
$$\text{run} = 3$$

Count the rise and the run. Mark the second point.



Connect the two points with a line.




You can check your work by finding a third point. Since the slope is $m = -\frac{2}{3}$, it can be written as $m = \frac{2}{-3}$. Go back to (0,2) and count out the rise, 2, and the run, −3.

? Try It 4.4.35

Graph the line with the *y*-intercept 4 and slope $m = -\frac{5}{2}$.


Answer



? Try It 4.4.36

Graph the line with the *x*-intercept −3 and slope $m = -\frac{3}{4}$.

Answer




Example 4.4.37

Graph the line passing through the point $(-1,-3)$ whose slope is $m=4$.

Solution

Plot the given point.




Identify the rise and the run. $m = 4$

Write 4 as a fraction. $\frac{\text{rise}}{\text{run}} = \frac{4}{1}$


$\text{rise} = 4 \quad \text{run} = 1$

Count the rise and run and mark the second point.



Connect the two points with a line.

https://math.libretexts.org/@go/page/30509




You can check your work by finding a third point. Since the slope is m=4, it can be written as $m = \frac{-4}{-1}$. Go back to (−1,−3) and count out the rise, −4, and the run, −1.

Try It 4.4.38

Graph the line with the point (−2,1) and slope m=3.


Answer



Try It 4.4.39

Graph the line with the point (4,−2) and slope m=−2.

Answer




Solve Slope Applications

At the beginning of this section, we said there are many applications of slope in the real world. Let's look at a few now.

Example 4.4.40

The 'pitch' of a building's roof is the slope of the roof. Knowing the pitch is important in climates where there is heavy snowfall. If the roof is too flat, the weight of the snow may cause it to collapse. What is the slope of the roof shown?



Solution

Use the slope formula.	$m = \frac{\text{rise}}{\text{rise}}$
Substitute the values for rise and run.	$m = \frac{9}{18}$
Simplify.	$m = \frac{1}{2}$

The slope of the roof is $\frac{1}{2}$.

The roof rises 1 foot for every 2 feet of horizontal run.

Try It 4.4.41

Use Exercise 4.4.40, substituting the rise = 14 and run = 24.

Answer

$\frac{7}{12}$

Try It 4.4.42

Use Exercise 4.4.40, substituting rise = 15 and run = 36.

Answer

$\frac{5}{12}$

Example 4.4.43

Have you ever thought about the sewage pipes going from your house to the street? They must slope down $\frac{1}{4}$ inch per foot in order to drain properly. What is the required slope?



Solution

Use the slope formula.	$m = \frac{\text{rise}}{\text{run}}$
	$m = \frac{-\frac{1}{4}\text{inch}}{1\text{ foot}}$
	$m = \frac{-\frac{1}{4}\text{ inch}}{12\text{ inches}}$
Simplify.	$m = -\frac{1}{48}$
	The slope of the pipe is $-\frac{1}{48}$

The pipe drops 1 inch for every 48 inches of horizontal run.

https://math.libretexts.org/@go/page/30509

Try It 4.4.44

Find the slope of a pipe that slopes down $\frac{1}{3}$ inch per foot.

Answer

$-\frac{1}{36}$

Try It 4.4.45

Find the slope of a pipe that slopes down $\frac{3}{4}$ inch per yard.

Answer

$-\frac{1}{48}$

Access these online resources for additional instruction and practice with understanding slope of a line.

- Practice Slope with a Virtual Geoboard
- Small, Medium, and Large Virtual Geoboards
- Explore Area and Perimeter with a Geoboard

Key Concepts

- **Find the Slope of a Line from its Graph using $m = \frac{\text{rise}}{\text{run}}$**
  1. Locate two points on the line whose coordinates are integers.
  2. Starting with the point on the left, sketch a right triangle, going from the first point to the second point.
  3. Count the rise and the run on the legs of the triangle.
  4. Take the ratio of rise to run to find the slope.
- **Graph a Line Given a Point and the Slope**
  1. Plot the given point.
  2. Use the slope formula $m = \frac{\text{rise}}{\text{run}}$ to identify the rise and the run.
  3. Starting at the given point, count out the rise and run to mark the second point.
  4. Connect the points with a line.
- **Slope of a Horizontal Line**
  - The slope of a horizontal line, y=b, is 0.
- **Slope of a vertical line**
  - The slope of a vertical line, x=a, is undefined

This page titled 4.4: Understanding the Slope of a Line is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.4: Understanding the Slope of a Line** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.


4.4E: Exercises

Practice Makes Perfect

Use Geoboards to Model Slope

In the following exercises, find the slope modeled on each geoboard.


Exercise 4.4E.1




Answer

$\frac{1}{4}$

Exercise 4.4E.2




Exercise 4.4E.3




Answer

$\frac{2}{3}$

Exercise 4.4E.4




Exercise 4.4E.5




Answer

$\frac{-3}{2} = -\frac{3}{2}$

Exercise 4.4E.6




Exercise 4.4E.7



Answer

$-\frac{2}{3}$

Exercise 4.4E.8




In the following exercises, model each slope. Draw a picture to show your results.

Exercise 4.4E.9

$\frac{2}{3}$

Answer




Exercise 4.4E.10

$\frac{3}{4}$

Exercise 4.4E.11

$\frac{1}{4}$

Answer




Exercise 4.4E.12

$\frac{4}{3}$

Exercise 4.4E.13

$-\frac{1}{2}$

Answer




Exercise 4.4E.14

$-\frac{3}{4}$

Exercise 4.4E.15

$-\frac{2}{3}$

Answer




Exercise 4.4E.16

$-\frac{3}{2}$

Use $m = \frac{rise}{run}$ to find the Slope of a Line from its Graph

In the following exercises, find the slope of each line shown.


Exercise 4.4E.17




Answer

$\frac{2}{5}$

Exercise 4.4E.18




Exercise 4.4E.19




Answer

$\frac{5}{4}$

Exercise 4.4E.20




Exercise 4.4E.21




Answer

$-\frac{1}{3}$

Exercise 4.4E.22




Exercise $4.4E.23$




Answer

$-\frac{3}{4}$

Exercise $4.4E.24$




Exercise $4.4E.25$




Answer

$\frac{3}{4}$

Exercise $4.4E.26$




Exercise $4.4E.27$




Answer

$-\frac{5}{2}$

Exercise $4.4E.28$




Exercise $4.4E.29$




Answer

$-\frac{2}{3}$

Exercise $4.4E.30$




Exercise $4.4E.31$



Answer

$\frac{1}{4}$

Exercise 4.4E.32



Find the Slope of Horizontal and Vertical Lines

In the following exercises, find the slope of each line.

Exercise 4.4E.33

y=3

Answer

0

Exercise 4.4E.34

y=1

Exercise 4.4E.35

x=4

Answer

undefined

Exercise 4.4E.36

x=2

Exercise 4.4E.37

y=−2

Answer

0

Exercise 4.4E.38

y=−3

Exercise 4.4E. 39

x=−5

Answer

undefined

Exercise 4.4E. 40

x=−4

Use the Slope Formula to find the Slope of a Line between Two Points

In the following exercises, use the slope formula to find the slope of the line between each pair of points.

Exercise 4.4E. 41

(1,4),(3,9)

Answer

$\frac{5}{2}$

Exercise 4.4E. 42

(2,3),(5,7)

Exercise 4.4E. 43

(0,3),(4,6)

Answer

$\frac{3}{4}$

Exercise 4.4E. 44

(0,1),(5,4)

Exercise 4.4E. 45

(2,5),(4,0)

Answer

$-\frac{5}{2}$

Exercise 4.4E. 46

(3,6),(8,0)

Exercise 4.4E. 47

(−3,3),(4,−5)

Answer

$-\frac{8}{7}$

Exercise 4.4E.48

$(-2,4),(3,-1)$

Exercise 4.4E.49

$(-1,-2),(2,5)$

Answer

$\frac{7}{3}$

Exercise 4.4E.50

$(-2,-1),(6,5)$

Exercise 4.4E.51

$(4,-5),(1,-2)$

Answer

-1

Exercise 4.4E.52

$(3,-6),(2,-2)$


Graph a Line Given a Point and the Slope

In the following exercises, graph each line with the given point and slope.

Exercise 4.4E.53

$(1,-2); m=\frac{3}{4}$

Answer




Exercise 4.4E.54

$(1,-1); m=\frac{2}{3}$

Exercise 4.4E.55

$(2,5); m = -\frac{1}{3}$

Answer




Exercise 4.4E.56

$(1,4); m = -\frac{1}{2}$

Exercise 4.4E.57

$(-3,4); m = -\frac{3}{2}$

Answer




Exercise 4.4E.58

$(-2,5); m = -\frac{5}{4}$

Exercise 4.4E.59

$(-1,-4); m = \frac{4}{3}$

Answer




Exercise 4.4E.60

$(-3, -5); m = \frac{3}{2}$

Exercise 4.4E.61

y -intercept $3; m = -\frac{2}{5}$

Answer




Exercise 4.4E.62

y -intercept $5; m = -\frac{4}{3}$

Exercise 4.4E.63

x -intercept $-2; m = \frac{3}{4}$

Answer




Exercise 4.4E.64

x -intercept $-1; m = \frac{1}{5}$

Exercise 4.4E.65

$(-3, 3); m = 2$

Answer




Exercise 4.4E.66

$(-4, 2); m = 4$

Exercise 4.4E.67

$(1, 5); m = -3$

Answer



Exercise 4.4E.68


$(2, 3); m = -1$

Everyday Math

Exercise 4.4E.69

Slope of a roof. An easy way to determine the slope of a roof is to set one end of a 12 inch level on the roof surface and hold it level. Then take a tape measure or ruler and measure from the other end of the level down to the roof surface. This will give you the slope of the roof. Builders, sometimes, refer to this as pitch and state it as an "x 12 pitch" meaning $\frac{x}{12}$, where x is the measurement from the roof to the level—the rise. It is also sometimes stated as an "x-in-12 pitch".

1. What is the slope of the roof in this picture?
2. What is the pitch in construction terms?




Answer

1. $\frac{1}{3}$
2. 4 12 pitch or 4-in-12 pitch

Exercise 4.4E.70

The slope of the roof shown here is measured with a 12” level and a ruler. What is the slope of this roof?



Exercise 4.4E.71

Road grade. A local road has a grade of 6%. The grade of a road is its slope expressed as a percent. Find the slope of the road as a fraction and then simplify. What rise and run would reflect this slope or grade?

Answer

$\frac{3}{50}$; rise $= 3$, run $= 50$

Exercise 4.4E.72

Highway grade. A local road rises 2 feet for every 50 feet of highway.

1. What is the slope of the highway?
2. The grade of a highway is its slope expressed as a percent. What is the grade of this highway?

Exercise 4.4E.73

Wheelchair ramp. The rules for wheelchair ramps require a maximum 1-inch rise for a 12-inch run.

1. How long must the ramp be to accommodate a 24-inch rise to the door?
2. Create a model of this ramp.

Answer

1. 288 inches (24 feet)
2. Models will vary.

Exercise 4.4E.74

Wheelchair ramp. A 1-inch rise for a 16-inch run makes it easier for the wheelchair rider to ascend a ramp.

1. How long must a ramp be to easily accommodate a 24-inch rise to the door?
2. Create a model of this ramp.

Writing Exercises

Exercise 4.4E.75

What does the sign of the slope tell you about a line?

Answer

When the slope is a positive number the line goes up from left to right. When the slope is a negative number the line goes down from left to right.

Exercise 4.4E.76

How does the graph of a line with slope $m = \frac{1}{2}$ differ from the graph of a line with slope $m = 2$?

Exercise 4.4E.77

Why is the slope of a vertical line "undefined"?

Answer

A vertical line has 0 run and since division by 0 is undefined the slope is undefined.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it
use geoboards to model slope.			
use $m = \frac{rise}{run}$ to find the slope of a line from its graph.			
find the slope of horizontal and vertical lines.			
use the slope formula to find the slope of a line between two points.			
graph a line given a point and the slope.			
solve slope applications.			

ⓑ On a scale of 1–10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

Glossary

geoboard

A geoboard is a board with a grid of pegs on it.

negative slope

A negative slope of a line goes down as you read from left to right.

positive slope

A positive slope of a line goes up as you read from left to right.

rise

The rise of a line is its vertical change.

run

The run of a line is its horizontal change.

slope formula

The slope of the line between two points (x_1, y_1) and (x_2, y_2) is $m = \frac{y_2 - y_1}{x_2 - x_1}$.

slope of a line

The slope of a line is $m = \frac{\text{rise}}{\text{run}}$. The rise measures the vertical change and the run measures the horizontal change.

This page titled 4.4E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.4E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.5: Use the Slope–Intercept Form of an Equation of a Line

Learning Objectives

By the end of this section, you will be able to:

- Recognize the relation between the graph and the slope–intercept form of an equation of a line
- Identify the slope and y-intercept form of an equation of a line
- Graph a line using its slope and intercept
- Choose the most convenient method to graph a line
- Graph and interpret applications of slope–intercept
- Use slopes to identify parallel lines
- Use slopes to identify perpendicular lines

Note

Before you get started, take this readiness quiz.

1. Add: $\frac{x}{4}+\frac{1}{4}$.
   If you missed this problem, review Exercise 1.7.1.
2. Find the reciprocal of $\frac{3}{7}$.
   If you missed this problem, review Exercise 1.6.19.
3. Solve $2x-3y=12$ for y.
   If you missed this problem, review Exercise 2.6.16.

Recognize the Relation Between the Graph and the Slope–Intercept Form of an Equation of a Line

We have graphed linear equations by plotting points, using intercepts, recognizing horizontal and vertical lines, and using the point–slope method. Once we see how an equation in slope–intercept form and its graph are related, we'll have one more method we can use to graph lines.

In **Graph Linear Equations in Two Variables**, we graphed the line of the equation $y=12x+3$ by plotting points. See Figure 4.5.1. Let's find the slope of this line.




Figure 4.5.1

The red lines show us the rise is 1 and the run is 2. Substituting into the slope formula:

$$m=\frac{\text{rise}}{\text{run}}$$
$$m=\frac{1}{2}$$

What is the y-intercept of the line? The y-intercept is where the line crosses the y-axis, so y-intercept is $(0,3)$. The equation of this line is:

$y = \frac{1}{2}x + 3$

Notice, the line has:

slope $m = \frac{1}{2}$ and y-intercept $(0, 3)$

When a linear equation is solved for y, the coefficient of the x-term is the slope and the constant term is the y-coordinate of the y-intercept. We say that the equation $y = \frac{1}{2}x + 3$ is in slope–intercept form.

$m = \frac{1}{2}$; y-intercept is $(0, 3)$

$y = \frac{1}{2}x + 3$

$y = mx + b$

SLOPE-INTERCEPT FORM OF AN EQUATION OF A LINE

The **slope–intercept form** of an equation of a line with slope m and y-intercept, $(0, b)$ is,

$$y = mx + b \tag{4.5.1}$$

Sometimes the slope–intercept form is called the "y-form."


Example 4.5.1

Use the graph to find the slope and y-intercept of the line, $y = 2x + 1$.

Compare these values to the equation $y = mx + b$.

Solution


To find the slope of the line, we need to choose two points on the line. We'll use the points $(0, 1)$ and $(1, 3)$.

	
Find the rise and run.	$m = \frac{\text{rise}}{\text{run}}$
	$m = \frac{2}{1}$
	$m = 2$
Find the y-intercept of the line.	The y-intercept is the point $(0, 1)$.
We found slope $m = 2$ and y-intercept $(0, 1)$.	$y = 2x + 1$ $y = mx + b$

The slope is the same as the coefficient of x and the y-coordinate of the y-intercept is the same as the constant term.

Try It 4.5.2

Use the graph to find the slope and y-intercept of the line $y = \frac{2}{3}x - 1$. Compare these values to the equation $y = mx + b$.




Answer

slope $m = \frac{2}{3}$ and y-intercept $(0, -1)$

Try It 4.5.3

Use the graph to find the slope and y-intercept of the line $y = \frac{1}{2}x + 3$. Compare these values to the equation $y = mx + b$.



Answer

slope $m = \frac{1}{2}$ and y-intercept $(0, 3)$

Identify the Slope and y-Intercept From an Equation of a Line

In **Understand Slope of a Line**, we graphed a line using the slope and a point. When we are given an equation in slope–intercept form, we can use the y-intercept as the point, and then count out the slope from there. Let's practice finding the values of the slope and y-intercept from the equation of a line.

Example 4.5.4

Identify the slope and y-intercept of the line with equation $y = -3x + 5$.

Solution

We compare our equation to the slope–intercept form of the equation.

	$y = mx + b$
Write the equation of the line.	$y = -3x + 5$
Identify the slope.	$m = -3$
Identify the y-intercept.	y-intercept is $(0, 5)$

Try It 4.5.5

Identify the slope and y-intercept of the line $y = \frac{2}{5}x - 1$.

Answer

slope $\frac{2}{5}$; y-intercept $(0,-1)$

Try It 4.5.6

Identify the slope and y-intercept of the line $y = -\frac{4}{3}x + 1$.

Answer

slope $-\frac{4}{3}$; y-intercept $(0,1)$

When an equation of a line is not given in slope–intercept form, our first step will be to solve the equation for y.

Example 4.5.7

Identify the slope and y-intercept of the line with equation $x + 2y = 6$.

Solution

This equation is not in slope–intercept form. In order to compare it to the slope–intercept form we must first solve the equation for y.

Solve for y.	$x + 2y = 6$
Subtract x from each side.	$2y = -x + 6$
Divide both sides by 2.	$\frac{2y}{2} = \frac{-x+6}{2}$
Simplify.	$\frac{2y}{2} = \frac{-x}{2} + \frac{6}{2}$
(Remember: $\frac{a+b}{c} = \frac{a}{c} + \frac{b}{c}$)	
Simplify.	$y = -\frac{1}{2}x + 3$
Write the slope–intercept form of the equation of the line.	$y = mx + b$
Write the equation of the line.	$y = -\frac{1}{2}x + 3$
Identify the slope.	$m = -\frac{1}{2}$
Identify the y-intercept.	y-intercept is $(0, 3)$

Try It 4.5.8

Identify the slope and y-intercept of the line $x + 4y = 8$.

Answer

slope $-\frac{1}{4}$; y-intercept $(0,2)$

Try It 4.5.9

Identify the slope and y-intercept of the line $3x + 2y = 12$.

Answer

slope $-\frac{3}{2}$; y-intercept (0,6)


Graph a Line Using its Slope and Intercept



Now that we know how to find the slope and y-intercept of a line from its equation, we can graph the line by plotting the y-intercept and then using the slope to find another point.

Example 4.5.10: How to Graph a Line Using its Slope and Intercept

Graph the line of the equation $y = 4x - 2$ using its slope and y-intercept.

Solution


Step 1. Find the slope–intercept form of the equation.	This equation is in slope–intercept form.	$y = 4x - 2$
Step 2. Identify the slope and y-intercept.	Use $y = mx + b$ Find the slope. Find the y-intercept.	$y = mx + b$ $y = 4x + (-2)$ $m = 4$ $b = -2$, $(0, -2)$
Step 3. Plot the y-intercept.	Plot $(0, -2)$.	
Step 4. Use the slope formula $m = \frac{\text{rise}}{\text{run}}$ to identify the rise and the run.	Identify the rise and the run.	$m = 4$ $\frac{\text{rise}}{\text{run}} = \frac{4}{1}$ rise $= 4$ run $= 1$

Step 5. Starting at the *y*-intercept, count out the rise and run to mark the second point.	Start at (0, –2) and count the rise and the run. Up 4, right 1.	
Step 6. Connect the points with a line.	Connect the two points with a line.	

Try It 4.5.11

Graph the line of the equation $y = 4x + 1$ using its slope and y-intercept.


Answer



Try It 4.5.12

Graph the line of the equation $y = 2x - 3$ using its slope and y-intercept.

Answer




GRAPH A LINE USING ITS SLOPE AND y-INTERCEPT.

1. Find the slope-intercept form of the equation of the line.
2. Identify the slope and y-intercept.
3. Plot the y-intercept.
4. Use the slope formula $\frac{\text{rise}}{\text{run}}$ to identify the rise and the run.
5. Starting at the y-intercept, count out the rise and run to mark the second point.
6. Connect the points with a line.

Example 4.5.13

Graph the line of the equation $y = -x + 4$ using its slope and y-intercept.

Solution

	$y = mx + b$
The equation is in slope–intercept form.	$y = -x + 4$
Identify the slope and y-intercept.	$m = -1$
	y-intercept is $(0, 4)$
Plot the y-intercept.	See graph below.
Identify the rise and the run.	$m = \frac{-1}{1}$
Count out the rise and run to mark the second point.	rise -1, run 1
Draw the line.	
To check your work, you can find another point on the line and make sure it is a solution of the equation. In the graph we see the line goes through $(4, 0)$.	

check.

$y = -x + 4$


$0 \stackrel{?}{=} -4 + 4$

$0 = 0$✓

Try It 4.5.14

Graph the line of the equation $y = -x - 3$ using its slope and y-intercept.


Answer



Try It 4.5.15

Graph the line of the equation $y = -x - 1$ using its slope and y-intercept.

Answer




Example 4.5.16

Graph the line of the equation $y = -\frac{2}{3}x - 3$ using its slope and y-intercept.

Solution


	$y = mx + b$
The equation is in slope–intercept form.	$y = -\frac{2}{3}x - 3$
Identify the slope and y-intercept.	$m = -\frac{2}{3}$; y-intercept is $(0, -3)$
Plot the y-intercept.	See graph below.
Identify the rise and the run.	
Count out the rise and run to mark the second point.	

Draw the line.	

Try It 4.5.17

Graph the line of the equation $y = -\frac{5}{2}x + 1$ using its slope and y-intercept.


Answer



Try It 4.5.18

Graph the line of the equation $y = -\frac{3}{4}x - 2$ using its slope and y-intercept.

Answer




Example 4.5.19

Graph the line of the equation $4x - 3y = 12$ using its slope and y-intercept.

Solution


	$4x - 3y = 12$
Find the slope–intercept form of the equation.	$-3y = -4x + 12$
	$-\frac{3y}{3} = \frac{-4x+12}{-3}$

The equation is now in slope–intercept form.	$y = \frac{4}{3}x - 4$
Identify the slope and y-intercept.	$m = \frac{4}{3}$
	y-ntercept is $(0, -4)$
Plot the y-intercept.	See graph below.
Identify the rise and the run; count out the rise and run to mark the second point.	
Draw the line.	

Try It 4.5.20

Graph the line of the equation $2x - y = 6$ using its slope and y-intercept.


Answer



Try It 4.5.21

Graph the line of the equation $3x - 2y = 8$ using its slope and y-intercept.

Answer




We have used a grid with x and y both going from about -10 to 10 for all the equations we've graphed so far. Not all linear equations can be graphed on this small grid. Often, especially in applications with real-world data, we'll need to extend the axes to bigger positive or smaller negative numbers.

Example 4.5.22

Graph the line of the equation $y = 0.2x + 45$ using its slope and y-intercept.

Solution


We'll use a grid with the axes going from about -80 to 80.

	$y = mx + b$
The equation is in slope–intercept form.	$y = 0.2x + 45$
Identify the slope and y-intercept.	$m = 0.2$
	The y-intercept is $(0, 45)$
Plot the y-intercept.	See graph below.
Count out the rise and run to mark the second point. The slope is $m = 0.2$; in fraction form this means $m = \frac{2}{10}$. Given the scale of our graph, it would be easier to use the equivalent fraction $m = \frac{10}{50}$.	
Draw the line.	

Try It 4.5.23

Graph the line of the equation $y = 0.5x + 25$ using its slope and y-intercept.

Answer




https://math.libretexts.org/@go/page/30511

? Try It 4.5.24

Graph the line of the equation $y = 0.1x - 30$ using its slope and y-intercept.

Answer



Now that we have graphed lines by using the slope and y-intercept, let's summarize all the methods we have used to graph lines. See Figure 4.5.2.

Methods to Graph Lines

Point Plotting	Slope–Intercept	Intercepts	Recognize Vertical and Horizontal Lines
x | y	$y = mx + b$	x | y 0 | | 0	
Find three points. Plot the points, make sure they line up, then draw the line.	Find the slope and *y*-intercept. Start at the *y*-intercept, then count the slope to get a second point.	Find the intercepts and a third point. Plot the points, make sure they line up, then draw the line.	The equation has only one variable. $x = a$ vertical $y = b$ horizontal

Figure 4.5.2

Choose the Most Convenient Method to Graph a Line

Now that we have seen several methods we can use to graph lines, how do we know which method to use for a given equation?

While we could plot points, use the slope–intercept form, or find the intercepts for *any* equation, if we recognize the most convenient way to graph a certain type of equation, our work will be easier. Generally, plotting points is not the most efficient way to graph a line. We saw better methods in sections 4.3, 4.4, and earlier in this section. Let's look for some patterns to help determine the most convenient method to graph a line.

Here are six equations we graphed in this chapter, and the method we used to graph each of them.

$$\begin{array}{lll} \#1 & \text{Equation} & \text{Method} \\ \#2 & x = 2 & \text{Vertical line} \\ \#3 & y = 4 & \text{Horizontal line} \\ \#4 & -x + 2y = 6 & \text{Intercepts} \\ \#5 & 4x - 3y = 12 & \text{Intercepts} \\ \#6 & y = 4x - 2 & \text{Slope-intercept} \\ \#7 & y = -x + 4 & \text{Slope-intercept} \end{array} \tag{4.5.2}$$

Equations #1 and #2 each have just one variable. Remember, in equations of this form the value of that one variable is constant; it does not depend on the value of the other variable. Equations of this form have graphs that are vertical or horizontal lines.

In equations #3 and #4, both x and y are on the same side of the equation. These two equations are of the form $Ax + By = C$. We substituted $y = 0$ to find the x-intercept and $x = 0$ to find the y-intercept, and then found a third point by choosing another value for x or y.

Equations #5 and #6 are written in slope–intercept form. After identifying the slope and y-intercept from the equation we used them to graph the line.

This leads to the following strategy.

STRATEGY FOR CHOOSING THE MOST CONVENIENT METHOD TO GRAPH A LINE

Consider the form of the equation.

- If it only has one variable, it is a vertical or horizontal line.
  - $x = a$ is a vertical line passing through the x-axis at a.
  - $y = b$ is a horizontal line passing through the y-axis at b.
- If y is isolated on one side of the equation, in the form $y = mx + b$, graph by using the slope and y-intercept.
  - Identify the slope and y-intercept and then graph.
- If the equation is of the form $Ax + By = C$, find the intercepts.
  - Find the x- and y-intercepts, a third point, and then graph.

Example 4.5.25

Determine the most convenient method to graph each line.

1. $y = -6$
2. $5x - 3y = 15$
3. $x = 7$
4. $y = \frac{2}{5}x - 1$.

Solution

1. $y = -6$
This equation has only one variable, y. Its graph is a horizontal line crossing the y-axis at -6.

2. $5x - 3y = 15$
This equation is of the form $Ax + By = C$. The easiest way to graph it will be to find the intercepts and one more point.

3. $x = 7$
There is only one variable, x. The graph is a vertical line crossing the x-axis at 7.

4. $y = \frac{2}{5}x - 1$
Since this equation is in $y = mx + b$ form, it will be easiest to graph this line by using the slope and y-intercept.

Try It 4.5.26

Determine the most convenient method to graph each line:

1. $3x + 2y = 12$
2. $y = 4$
3. $y = \frac{1}{5}x - 4$
4. $x = -7$

Answer

1. intercepts
2. horizontal line
3. slope–intercept
4. vertical line

Try It 4.5.27

Determine the most convenient method to graph each line:

1. $x = 6$
2. $y = -\frac{3}{4}x + 1$
3. $y = -8$
4. $4x - 3y = -1$

Answer

1. vertical line
2. slope–intercept
3. horizontal line
4. intercepts

Graph and Interpret Applications of Slope–Intercept

Many real-world applications are modeled by linear equations. We will take a look at a few applications here so you can see how equations written in slope–intercept form relate to real-world situations.

Usually when a linear equation models a real-world situation, different letters are used for the variables, instead of x and y. The variable names remind us of what quantities are being measured.

Example 4.5.28

The equation $F = \frac{9}{5}C + 32$ is used to convert temperatures, C, on the Celsius scale to temperatures, F, on the Fahrenheit scale.

1. Find the Fahrenheit temperature for a Celsius temperature of 0.
2. Find the Fahrenheit temperature for a Celsius temperature of 20.
3. Interpret the slope and F-intercept of the equation.
4. Graph the equation.

Solution

1. Find the Fahrenheit temperature for a Celsius temperature of 0. $F = \frac{9}{5}C + 32$
   Find F when $C = 0$. $F = \frac{9}{5}(0) + 32$
   Simplify. $F = 32$

2.

$$\begin{aligned} &\text{Find the Fahrenheit temperature for a Celsius temperature of } 20. && F = \tfrac{9}{5}C + 32 \\ &\text{Find } F \text{ when } C = 20. && F = \tfrac{9}{5}(20) + 32 \\ &\text{Simplify.} && F = 36 + 32 \\ &\text{Simplify.} && F = 68 \end{aligned} \tag{4.5.3}$$

3. Interpret the slope and F-intercept of the equation.

Even though this equation uses F and C, it is still in slope–intercept form.

$y = mx + b$

$F = mC + b$

$F = \frac{9}{5}C + 32$

The slope, $\frac{9}{5}$, means that the temperature Fahrenheit (F) increases 9 degrees when the temperature Celsius (C) increases 5 degrees.

The F-intercept means that when the temperature is 0° on the Celsius scale, it is 32° on the Fahrenheit scale.

4. Graph the equation.

We'll need to use a larger scale than our usual. Start at the F-intercept $(0, 32)$ then count out the rise of 9 and the run of 5 to get a second point. See Figure $4.5.3$.





Figure 4.5.3

? Try It 4.5.29

The equation $h = 2s + 50$ is used to estimate a woman's height in inches, h, based on her shoe size, s.

1. Estimate the height of a child who wears women's shoe size 0.
2. Estimate the height of a woman with shoe size 8.
3. Interpret the slope and h-intercept of the equation.
4. Graph the equation.

Answer


1. 50 inches
2. 66 inches
3. The slope, 2, means that the height, h, increases by 2 inches when the shoe size, s, increases by 1. The h-intercept means that when the shoe size is 0, the height is 50 inches.
4. 

? Try It 4.5.30

The equation $T = \frac{1}{4}n + 40$ is used to estimate the temperature in degrees Fahrenheit, T, based on the number of cricket chirps, n, in one minute.

1. Estimate the temperature when there are no chirps.
2. Estimate the temperature when the number of chirps in one minute is 100.
3. Interpret the slope and T-intercept of the equation.
4. Graph the equation.

Answer

1. 40 degrees
2. 65 degrees
3. The slope, $\frac{1}{4}$, means that the temperature Fahrenheit (F) increases 1 degree when the number of chirps, n, increases by 4. The T-intercept means that when the number of chirps is 0, the temperature is 40°.
4. 

The cost of running some types business has two components—a *fixed cost* and a *variable cost*. The fixed cost is always the same regardless of how many units are produced. This is the cost of rent, insurance, equipment, advertising, and other items that must be paid regularly. The variable cost depends on the number of units produced. It is for the material and labor needed to produce each item.

Example 4.5.31


Stella has a home business selling gourmet pizzas. The equation $C = 4p + 25$ models the relation between her weekly cost, C, in dollars and the number of pizzas, p, that she sells.

1. Find Stella's cost for a week when she sells no pizzas.
2. Find the cost for a week when she sells 15 pizzas.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.

Solution

1. Find Stella's cost for a week when she sells no pizzas.	$C = 4p + 25$
Find C when $p = 0$.	$C = 4(0) + 25$
Simplify.	$C = 25$
	Stella's fixed cost is \$25 when she sells no pizzas.
2. Find the cost for a week when she sells 15 pizzas.	$C = 4p + 25$
Find C when $p = 15$.	$C = 4(15) + 25$
Simplify.	$C = 60 + 25$
	$C = 85$
	Stella's costs are \$85 when she sells 15 pizzas.
3. Interpret the slope and C-intercept of the equation.	$y = mx + b$ $C = 4p + 25$
	The slope, 4, means that the cost increases by \$4 for each pizza Stella sells. The C-intercept means that even when Stella sells no pizzas, her costs for the week are \$25.

4. Graph the equation. We'll need to use a larger scale than our usual. Start at the C-intercept $(0, 25)$ then count out the rise of 4 and the run of 1 to get a second point.




Try It 4.5.32

Sam drives a delivery van. The equation $C = 0.5m + 60$ models the relation between his weekly cost, C, in dollars and the number of miles, m, that he drives.

1. Find Sam's cost for a week when he drives 0 miles.
2. Find the cost for a week when he drives 250 miles.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.

Answer

1. $\$60$
2. $\$185$
3. The slope, 0.5, means that the weekly cost, C, increases by $\$0.50$ when the number of miles driven, n, increases by 1. The C-intercept means that when the number of miles driven is 0, the weekly cost is $\$60$.
4.



Try It 4.5.33


Loreen has a calligraphy business. The equation $C = 1.8n + 35$ models the relation between her weekly cost, C, in dollars and the number of wedding invitations, n, that she writes.

1. Find Loreen's cost for a week when she writes no invitations.
2. Find the cost for a week when she writes 75 invitations.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.

Answer

1. $\$35$
2. $\$170$
3. The slope, 1.8, means that the weekly cost, C, increases by $\$1.80$ when the number of invitations, n, increases by 1. The C-intercept means that when the number of invitations is 0, the weekly cost is $\$35$.

4.



Use Slopes to Identify Parallel Lines

The slope of a line indicates how steep the line is and whether it rises or falls as we read it from left to right. Two lines that have the same slope are called parallel lines. Parallel lines never intersect.




Figure 4.5.4.




Figure 4.5.4: Verify that both lines have the same slope, $m = \frac{2}{5}$, and different y-intercepts.

What about vertical lines? The slope of a vertical line is undefined, so vertical lines don't fit in the definition above. We say that vertical lines that have different x-intercepts are parallel. See Figure 4.5.5.




Figure 4.5.5: Vertical lines with different x-intercepts are parallel.

PARALLEL LINES


Parallel lines are lines in the same plane that do not intersect.

- Parallel lines have the same slope and different y-intercepts.
- If m_1 and m_2 are the slopes of two parallel lines then $m_1 = m_2$.
- Parallel vertical lines have different x-intercepts.

Let's graph the equations $y = -2x + 3$ and $2x + y = -1$ on the same grid. The first equation is already in slope–intercept form: $y = -2x + 3$. We solve the second equation for y:

$$\begin{aligned} 2x + y &= -1 \\ y &= -2x - 1 \end{aligned}$$

Graph the lines.



Notice the lines look parallel. What is the slope of each line? What is the y-intercept of each line?

$$\begin{array}{lll} y & = mx + b & y = mx + b \\ y & = -2x + 3 & y = -2x - 1 \\ m & = -2 & m = -2 \\ b & = 3, (0, 3) & b = -1, (0, -1) \end{array} \tag{4.5.4}$$

The slopes of the lines are the same and the y-intercept of each line is different. So we know these lines are parallel.

Since parallel lines have the same slope and different y-intercepts, we can now just look at the slope–intercept form of the equations of lines and decide if the lines are parallel.

Example 4.5.34

Use slopes and y-intercepts to determine if the lines $3x-2y=6$ and $y=\frac{3}{2}x+1$ are parallel.

Solution

Solve the first equation for y.

$$\begin{aligned} 3x-2y &= 6 \\ -2y &= -3x+6 \\ \frac{-2y}{-2} &= \frac{-3x+6}{-2} \\ y &= \frac{3}{2}x-3 \end{aligned}$$

The equation is now in slope–intercept form.

The equation of the second line is already in slope–intercept form.

Identify the slope and y-intercept of both lines.

$y=\frac{3}{2}x+1$	$y=\frac{3}{2}x-3$
$y=mx+b$	$y=mx+b$
$m=\frac{3}{2}$	$m=\frac{3}{2}$
y-intercept is $(0,1)$	y-intercept is $(0,-3)$

The lines have the same slope and different y-intercepts and so they are parallel. You may want to graph the lines to confirm whether they are parallel.

Try It 4.5.35

Use slopes and y-intercepts to determine if the lines $2x+5y=5$ and $y=-\frac{2}{5}x-4$ are parallel.

Answer

parallel

Try It 4.5.36

Use slopes and y-intercepts to determine if the lines $4x-3y=6$ and $y=\frac{4}{3}x-1$ are parallel.

Answer

parallel

Example 4.5.37

Use slopes and y-intercepts to determine if the lines $y=-4$ and $y=3$ are parallel.

Solution

Write each equation in slope-intercept form.	$y=-4$	and	$y=3$
Since there is no x term we write $0x$.	$y=0x-4$		$y=0x+3$
Identify the slope and y-intercept of both lines.	$y=mx+b$		$y=mx+b$
	$m=0$		$m=0$
	y-intercept is $(0,4)$		y-intercept is $(0,3)$

The lines have the same slope and different y-intercepts and so they are parallel.

There is another way you can look at this example. If you recognize right away from the equations that these are horizontal lines, you know their slopes are both 0. Since the horizontal lines cross the y-axis at $y=-4$ and at $y=3$, we know the y-intercepts are $(0,-4)$ and $(0,3)$. The lines have the same slope and different y-intercepts and so they are parallel.

? Try It 4.5.38

Use slopes and y-intercepts to determine if the lines $y=8$ and $y=-6$ are parallel.

Answer

parallel

? Try It 4.5.39

Use slopes and y-intercepts to determine if the lines $y=1$ and $y=-5$ are parallel.

Answer

parallel

✓ Example 4.5.40

Use slopes and y-intercepts to determine if the lines $x=-2$ and $x=-5$ are parallel.

Solution

$$x=-2 \text{ and } x=-5 \tag{4.5.5}$$

Since there is no y, the equations cannot be put in slope–intercept form. But we recognize them as equations of vertical lines. Their x-intercepts are -2 and -5. Since their x-intercepts are different, the vertical lines are parallel.

? Try It 4.5.41

Use slopes and y-intercepts to determine if the lines $x=1$ and $x=-5$ are parallel.

Answer

parallel

? Try It 4.5.42

Use slopes and y-intercepts to determine if the lines $x=8$ and $x=-6$ are parallel.

Answer

parallel

✓ Example 4.5.43

Use slopes and y-intercepts to determine if the lines $y=2x-3$ and $-6x+3y=-9$ are parallel. You may want to graph these lines, too, to see what they look like.

Solution

The first equation is already in slope-intercept form. $y = 2x - 3$

Solve the second equation for y

$$\begin{aligned} -6x + 3y &= -9 \\ 3y &= 6x - 9 \\ \tfrac{3y}{3} &= \tfrac{6x-9}{3} \\ y &= 2x - 3 \end{aligned}$$

The second equation is now in slope–intercept form as well.

Identify the slope and y-intercept of both lines.

$y = 2x - 3$	$y = 2x - 3$
$y = mx + b$	$y = mx + b$
$m = 2$	$m = 2$
The y-intercept is $(0, -3)$	The y-intercept is $(0, -3)$

The lines have the same slope, but they also have the same y-intercepts. Their equations represent the same line. They are not parallel; they are the same line.

Try It 4.5.44

Use slopes and y-intercepts to determine if the lines $y = -\frac{1}{2}x + 1$ and $x + 2y = 2$ are parallel.

Answer

not parallel; same line

Try It 4.5.45

Use slopes and y-intercepts to determine if the lines $y = \frac{3}{4}x - 3$ and $3x - 4y = 12$ are parallel.

Answer

not parallel; same line

Use Slopes to Identify Perpendicular Lines

Let's look at the lines whose equations are $y = \frac{1}{4}x - 1$ and $y = -4x + 2$, shown in Figure 4.5.5.




Figure 4.5.5

These lines lie in the same plane and intersect in right angles. We call these lines **perpendicular**.

What do you notice about the slopes of these two lines? As we read from left to right, the line $y = \frac{1}{4}x - 1$ rises, so its slope is positive. The line $y = -4x + 2$ drops from left to right, so it has a negative slope. Does it make sense to you that the slopes of two perpendicular lines will have opposite signs?

If we look at the slope of the first line, $m_1 = \frac{1}{4}$, and the slope of the second line, $m_2 = -4$, we can see that they are *negative reciprocals* of each other. If we multiply them, their product is -1.

$$\begin{array}{c} m_1 \cdot m_2 \\ \frac{1}{4}(-4) \\ -1 \end{array} \tag{4.5.6}$$

This is always true for **perpendicular lines** and leads us to this definition.

PERPENDICULAR LINES

Perpendicular lines are lines in the same plane that form a right angle.

If m_1 and m_2 are the slopes of two perpendicular lines, then:

$$m_1 \cdot m_2 = -1 \text{ and } m_1 = \frac{-1}{m_2} \tag{4.5.7}$$

Vertical lines and horizontal lines are always perpendicular to each other.

We were able to look at the slope–intercept form of linear equations and determine whether or not the lines were parallel. We can do the same thing for perpendicular lines.

We find the slope–intercept form of the equation, and then see if the slopes are negative reciprocals. If the product of the slopes is -1, the lines are perpendicular. Perpendicular lines may have the same y-intercepts.

Example 4.5.46

Use slopes to determine if the lines, $y = -5x - 4$ and $x - 5y = 5$ are perpendicular.

Solution

The first equation is already in slope–intercept form: $y = -5x - 4$

Solve the second equation for y.

$$\begin{array}{rcl} x - 5y & = & 5 \\ -5y & = & -x + 5 \\ \frac{-5y}{-5} & = & \frac{-x+5}{-5} \\ y & = & \frac{1}{5}x - 1 \end{array}$$

The second equation is now in slope-intercept form as well.

Identify the slope of each line.

$$\begin{array}{rclrcl} y & = & -5x - 4 & y & = & \frac{1}{5}x - 1 \\ y & = & mx + b & y & = & mx + b \\ m_1 & = & -5 & m_2 & = & \frac{1}{5} \end{array}$$

The slopes are negative reciprocals of each other, so the lines are perpendicular. We check by multiplying the slopes,

$$\begin{array}{c} m_1 \cdot m_2 \\ -5\left(\frac{1}{5}\right) \\ -1\checkmark \end{array} \tag{4.5.8}$$

Try It 4.5.47

Use slopes to determine if the lines $y = -3x + 2$ and $x - 3y = 4$ are perpendicular.

Answer

perpendicular

Try It 4.5.48

Use slopes to determine if the lines $y = 2x - 5$ and $x + 2y = -6$ are perpendicular.

Answer

perpendicular

Example 4.5.49

Use slopes to determine if the lines, $7x + 2y = 3$ and $2x + 7y = 5$ are perpendicular.

Solution

Solve the equations for y.

$$\begin{aligned} 7x + 2y &= 3 \\ 2y &= -7x + 3 \\ \frac{2y}{2} &= \frac{-7x+3}{2} \\ y &= -\frac{7}{2}x + \frac{3}{2} \end{aligned} \qquad \begin{aligned} 2x + 7y &= 5 \\ 7y &= -2x + 5 \\ \frac{7y}{7} &= \frac{-2x+5}{7} \\ y &= \frac{-2}{7}x + \frac{5}{7} \end{aligned}$$

Identify the slope of each line.

$$\begin{aligned} y &= mx + b \\ m_1 &= -\frac{7}{2} \end{aligned} \qquad \begin{aligned} y &= mx + b \\ m_2 &= -\frac{2}{7} \end{aligned}$$

The slopes are reciprocals of each other, but they have the same sign. Since they are not negative reciprocals, the lines are not perpendicular.

Try It 4.5.50

Use slopes to determine if the lines $5x + 4y = 1$ and $4x + 5y = 3$ are perpendicular.

Answer

not perpendicular

Try It 4.5.51

Use slopes to determine if the lines $2x - 9y = 3$ and $9x - 2y = 1$ are perpendicular.

Answer

not perpendicular

Note

Access this online resource for additional instruction and practice with graphs.

- Explore the Relation Between a Graph and the Slope–Intercept Form of an Equation of a Line

Key Concepts

- The slope–intercept form of an equation of a line with slope mm and y-intercept, $(0, b)$ is, $y = mx + b$.
- **Graph a Line Using its Slope and y-Intercept**
  1. Find the slope-intercept form of the equation of the line.
  2. Identify the slope and y-intercept.
  3. Plot the y-intercept.
  4. Use the slope formula $m = \frac{\text{rise}}{\text{run}}$ to identify the rise and the run.
  5. Starting at the y-intercept, count out the rise and run to mark the second point.
  6. Connect the points with a line.
- **Strategy for Choosing the Most Convenient Method to Graph a Line:** Consider the form of the equation.
  - If it only has one variable, it is a vertical or horizontal line.
    $x = a$ is a vertical line passing through the x-axis at a.
    $y = b$ is a horizontal line passing through the y-axis at b.
  - If y is isolated on one side of the equation, in the form $y = mx + b$, graph by using the slope and y-intercept.
    Identify the slope and y-intercept and then graph.
  - If the equation is of the form $Ax + By = C$, find the intercepts.
    Find the x- and y-intercepts, a third point, and then graph.
- Parallel lines are lines in the same plane that do not intersect.
  - Parallel lines have the same slope and different y-intercepts.
  - If m_1 and m_2 are the slopes of two parallel lines then $m_1 = m_2$.
  - Parallel vertical lines have different x-intercepts.
- Perpendicular lines are lines in the same plane that form a right angle.
  - If m_1 and m_2 are the slopes of two perpendicular lines, then $m_1 \cdot m_2 = -1$ and $m_1 = \frac{-1}{m_2}$.
  - Vertical lines and horizontal lines are always perpendicular to each other.

Glossary

parallel lines

Lines in the same plane that do not intersect.

perpendicular lines

Lines in the same plane that form a right angle.

slope-intercept form of an equation of a line

The slope–intercept form of an equation of a line with slope mm and y-intercept, $(0, b)$ is, $y = mx + b$.

This page titled 4.5: Use the Slope–Intercept Form of an Equation of a Line is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.5: Use the Slope–Intercept Form of an Equation of a Line** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.


4.5E: Exercises

Practice Makes Perfect

Recognize the Relation Between the Graph and the Slope–Intercept Form of an Equation of a Line


In the following exercises, use the graph to find the slope and y-intercept of each line. Compare the values to the equation $y = mx + b$.

Exercise $4.5E.1$



$y = 3x - 5$

Exercise $4.5E.2$




$y = 4x - 2$

Answer


slope $m = 4$ and y-intercept $(0, -2)$

Exercise $4.5E.3$



$y = -x + 4$

Exercise $4.5E.4$




$y = -3x + 1$

Answer


slope $m = -3$ and y-intercept $(0, 1)$

Exercise $4.5E.5$



$y = -\frac{4}{3}x + 1$

Exercise $4.5E.6$



$y = -\frac{2}{5}x + 3$

Answer

slope $m = -\frac{2}{5}$ and y -intercept $(0, 3)$

Identify the Slope and y-Intercept From an Equation of a Line

In the following exercises, identify the slope and y-intercept of each line.

Exercise $4.5E.7$

$y = -7x + 3$

Exercise $4.5E.8$

$y = -9x + 7$

Answer

$m = -9$; y-intercept: $(0, 7)$

Exercise $4.5E.9$

$y = 6x - 8$

Exercise $4.5E.10$

$y = 4x - 10$

Answer

$m = 4$; y-intercept: $(0, -10)$

Exercise $4.5E.11$

$3x + y = 5$

Exercise 4.5E.12

$4x + y = 8$

Answer

$m = -4 \backslash 0$; \(y-intercept: $(0, 8)$

Exercise 4.5E.13

$6x + 4y = 12$

Exercise 4.5E.14

$8x + 3y = 12$

Answer

$m = -\frac{8}{3}$; y-intercept: $(0, 4)$

Exercise 4.5E.15

$5x - 2y = 6$

Exercise 4.5E.16

$7x - 3y = 9$

Answer

$m = \frac{7}{3}$; y-intercept: $(0, -3)$

Graph a Line Using Its Slope and Intercept

In the following exercises, graph the line of each equation using its slope and y-intercept.


Exercise 4.5E.17

$y = x + 3$

Exercise 4.5E.18

$y = x + 4$

Answer




Exercise 4.5E.19

$y = 3x - 1$

Exercise 4.5E.20

$y = 2x - 3$

Answer




Exercise 4.5E.21

$y = -x + 2$

Exercise 4.5E.22

$y = -x + 3$

Answer




Exercise 4.5E.23

$y = -x - 4$

Exercise 4.5E.24

$y = -x - 2$

Answer




Exercise 4.5E.25

$y = -\frac{3}{4}x - 1$

Exercise 4.5E.26

$y = -\frac{2}{5}x - 3$

Answer




Exercise 4.5E.27

$y = -\frac{3}{5}x + 2$

Exercise 4.5E.28

$y = -\frac{2}{3}x + 1$

Answer




Exercise 4.5E.29

$3x - 4y = 8$

Exercise 4.5E.30

$4x - 3y = 6$

Answer




Exercise 4.5E.31

$y = 0.1x + 15$

Exercise 4.5E.32

$y = 0.3x + 25$

Answer



Choose the Most Convenient Method to Graph a Line

In the following exercises, determine the most convenient method to graph each line.

Exercise 4.5E.33

$x = 2$

Exercise 4.5E.34

$y = 4$

Answer

horizontal line

Exercise 4.5E.35

$y = 5$

Exercise 4.5E.36

$x = -3$

Answer

vertical line

Exercise 4.5E.37

$y = -3x + 4$

Exercise 4.5E.38

$y = -5x + 2$

Answer

slope–intercept

Exercise 4.5E.39

$x - y = 5$

Exercise 4.5E.40

$x - y = 1$

Answer

intercepts

Exercise 4.5E.41

$y = \frac{2}{3}x - 1$

Exercise 4.5E.42

$y = \frac{4}{5}x - 3$

Answer

slope–intercept

Exercise 4.5E.43

$y = -3$

Exercise 4.5E.44

$y = -1$

Answer

horizontal line

Exercise 4.5E.45

$3x - 2y = -12$

Exercise 4.5E.46

$2x - 5y = -10$

Answer

intercepts

Exercise 4.5E.47

$y = -\frac{1}{4}x + 3$

Exercise 4.5E.48

$y = -\frac{1}{3}x + 5$

Answer

slope–intercept

Graph and Interpret Applications of Slope–Intercept

Exercise 4.5E.49

The equation $P = 31 + 1.75w$ models the relation between the amount of Tuyet's monthly water bill payment, P, in dollars, and the number of units of water, w, used.

1. Find Tuyet's payment for a month when 0 units of water are used.
2. Find Tuyet's payment for a month when 12 units of water are used.
3. Interpret the slope and P-intercept of the equation.
4. Graph the equation.


Exercise 4.5E.50

The equation $P = 28 + 2.54w$ models the relation between the amount of Randy's monthly water bill payment, P, in dollars, and the number of units of water, w, used.

1. Find the payment for a month when Randy used 0 units of water.
2. Find the payment for a month when Randy used 15 units of water.
3. Interpret the slope and P-intercept of the equation.
4. Graph the equation.

Answer

1. $28
2. $66.10
3. The slope, 2.54, means that Randy's payment, P, increases by $2.54 when the number of units of water he used, w, ncreases by 1. The P-intercept means that if the number units of water Randy used was 0, the payment would be $28
4.



Exercise $4.5E.51$

Bruce drives his car for his job. The equation $R = 0.575m + 42$ models the relation between the amount in dollars, R, that he is reimbursed and the number of miles, m, he drives in one day.

1. Find the amount Bruce is reimbursed on a day when he drives 0 miles.
2. Find the amount Bruce is reimbursed on a day when he drives 220 miles.
3. Interpret the slope and R-intercept of the equation.
4. Graph the equation.


Exercise $4.5E.52$

Janelle is planning to rent a car while on vacation. The equation $C = 0.32m + 15$ models the relation between the cost in dollars, C, per day and the number of miles, m, she drives in one day.

1. Find the cost if Janelle drives the car 0 miles one day.
2. Find the cost on a day when Janelle drives the car 400 miles.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.

Answer

1. $\$15$
2. $\$143$
3. The slope, 0.32, means that the cost, C, increases by $\$0.32$when the number of miles driven, m, increases by 1. The C-intercept means that if Janelle drives 0 miles one day, the cost would be $\$15$.
4.



Exercise $4.5E.53$

Cherie works in retail and her weekly salary includes commission for the amount she sells. The equation $S = 400 + 0.15c$ models the relation between her weekly salary, S, in dollars and the amount of her sales, c, in dollars.


1. Find Cherie's salary for a week when her sales were 0.
2. Find Cherie's salary for a week when her sales were 3600.
3. Interpret the slope and S-intercept of the equation.
4. Graph the equation.

Exercise $4.5E.54$

Patel's weekly salary includes a base pay plus commission on his sales. The equation $S = 750 + 0.09c$ models the relation between his weekly salary, S, in dollars and the amount of his sales, c, in dollars.

1. Find Patel's salary for a week when his sales were 0.
2. Find Patel's salary for a week when his sales were $18,540$.
3. Interpret the slope and S-intercept of the equation.
4. Graph the equation.

Answer

1. $\$750$
2. $\$2418.60$
3. The slope, 0.09, means that Patel's salary, S, increases by $\$0.09$ for every $\$1$ increase in his sales. The S-intercept means that when his sales are $\$0$, his salary is $\$750$
4. 

Exercise $4.5E.55$

Costa is planning a lunch banquet. The equation $C = 450 + 28g$ models the relation between the cost in dollars, C, of the banquet and the number of guests, g.

1. Find the cost if the number of guests is 40.
2. Find the cost if the number of guests is 80.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.


Exercise $4.5E.56$

Margie is planning a dinner banquet. The equation $C = 750 + 42g$ models the relation between the cost in dollars, C, of the banquet and the number of guests, g.

1. Find the cost if the number of guests is 50.
2. Find the cost if the number of guests is 100.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.

Answer

1. $\$2850$
2. $\$4950$

3. The slope, 42, means that the cost, C, increases by $\$42$ for when the number of guests increases by 1. The C-intercept means that when the number of guests is 0, the cost would be $\$750$
4. 

Use Slopes to Identify Parallel Lines

In the following exercises, use slopes and y-intercepts to determine if the lines are parallel.

Exercise $4.5E.57$

$y = \frac{3}{4}x - 3; \quad 3x - 4y = -2$

Exercise $4.5E.58$

$y = \frac{2}{3}x - 1; \quad 2x - 3y = -2$

Answer

parallel

Exercise $4.5E.59$

$2x - 5y = -3; \quad y = \frac{2}{5}x + 1$

Exercise $4.5E.60$

$3x - 4y = -2; \quad y = \frac{3}{4}x - 3$

Answer

parallel

Exercise $4.5E.61$

$2x - 4y = 6; \quad x - 2y = 3$

Exercise $4.5E.62$

$6x - 3y = 9; \quad 2x - y = 3$

Answer

not parallel

Exercise 4.5E.63

$4x+2y=6; \quad 6x+3y=3$

Exercise 4.5E.64

$8x+6y=6; \quad 12x+9y=12$

Answer

parallel

Exercise 4.5E.65

$x=5; \quad x=-6$

Exercise 4.5E.66

$x=7; \quad x=-8$

Answer

parallel

Exercise 4.5E.67

$x=-4; \quad x=-1$

Exercise 4.5E.68

$x=-3; \quad x=-2$

Answer

parallel

Exercise 4.5E.69

$y=2; \quad y=6$

Exercise 4.5E.70

$y=5; \quad y=1$

Answer

parallel

Exercise 4.5E.71

$y=-4; \quad y=3$

Exercise 4.5E.72

$y = -1; \quad y = 2$

Answer

parallel

Exercise 4.5E.73

$x - y = 2; \quad 2x - 2y = 4$

Exercise 4.5E.74

$4x + 4y = 8; \quad x + y = 2$

Answer

not parallel

Exercise 4.5E.75

$x - 3y = 6; \quad 2x - 6y = 12$

Exercise 4.5E.76

$5x - 2y = 11; \quad 5x - y = 7$

Answer

not parallel

Exercise 4.5E.77

$3x - 6y = 12; \quad 6x - 3y = 3$

Exercise 4.5E.78

$4x - 8y = 16; \quad x - 2y = 4$

Answer

not parallel

Exercise 4.5E.79

$9x - 3y = 6; \quad 3x - y = 2$

Exercise 4.5E.80

$x - 5y = 10; \quad 5x - y = -10$

Answer

not parallel

Exercise 4.5E.81

$7x-4y=8; \quad 4x+7y=14$

Exercise 4.5E.82

$9x-5y=4; \quad 5x+9y=-1$

Answer

not parallel

Use Slopes to Identify Perpendicular Lines

In the following exercises, use slopes and y-intercepts to determine if the lines are perpendicular.

Exercise 4.5E.83

$3x-2y=8; \quad 2x+3y=6$

Exercise 4.5E.84

$x-4y=8; \quad 4x+y=2$

Answer

perpendicular

Exercise 4.5E.85

$2x+5y=3; \quad 5x-2y=6$

Exercise 4.5E.86

$2x+3y=5; \quad 3x-2y=7$

Answer

perpendicular

Exercise 4.5E.87

$3x-2y=1; \quad 2x-3y=2$

Exercise 4.5E.88

$3x-4y=8; \quad 4x-3y=6$

Answer

not perpendicular

Exercise 4.5E.89

$5x+2y=6; \quad 2x+5y=8$

Exercise 4.5E.90

$2x+4y=3; \quad 6x+3y=2$

Answer

not perpendicular

Exercise 4.5E.91

$4x-2y=5; \quad 3x+6y=8$

Exercise 4.5E.92

$2x-6y=4; \quad 12x+4y=9$

Answer

perpendicular

Exercise 4.5E.93

$6x-4y=5; \quad 8x+12y=3$

Exercise 4.5E.94

$8x-2y=7; \quad 3x+12y=9$

Answer

perpendicular

Everyday Math

Exercise 4.5E.95

The equation $C=\frac{5}{9}F-17.8$ can be used to convert temperatures, F, on the Fahrenheit scale to temperatures, C, on the Celsius scale.

1. Explain what the slope of the equation means.
2. Explain what the C-intercept of the equation means.

Exercise 4.5E.96

The equation $n=4T-160$ is used to estimate the number of cricket chirps, n, in one minute based on the temperature in degrees Fahrenheit, T.

1. Explain what the slope of the equation means.
2. Explain what the n-intercept of the equation means. Is this a realistic situation?

Answer

1. For every increase of one degree Fahrenheit, the number of chirps increases by four.
2. There would be -160 chirps when the Fahrenheit temperature is 0°. (Notice that this does not make sense; this model cannot be used for all possible temperatures.)

Writing Exercises

Exercise $4.5E.97$

Explain in your own words how to decide which method to use to graph a line.

Exercise $4.5E.98$

Why are all horizontal lines parallel?

Answer

Answers will vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
recognize the relation between the graph and the slope-intercept form of an equation of a line.			
identify the slope and *y*-intercept from an equation of a line.			
graph a line using its slope and intercept.			
choose the most convenient method to graph a line.			
graph and interpret applications of slope-intercept.			
use slopes to identify parallel lines.			

b. After looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

This page titled 4.5E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.5E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.6: Find the Equation of a Line

Learning Objectives

By the end of this section, you will be able to:

- Find an equation of the line given the slope and y-intercept
- Find an equation of the line given the slope and a point
- Find an equation of the line given two points
- Find an equation of a line parallel to a given line
- Find an equation of a line perpendicular to a given line

Note

Before you get started, take this readiness quiz.

1. Solve: $\frac{2}{3} = \frac{x}{5}$.
   If you missed this problem, review Exercise 2.2.4.
2. Simplify: $-\frac{2}{5}(x - 15)$.
   If you missed this problem, review Exercise 1.10.34.

How do online retailers know that 'you may also like' a particular item based on something you just ordered? How can economists know how a rise in the minimum wage will affect the unemployment rate? How do medical researchers create drugs to target cancer cells? How can traffic engineers predict the effect on your commuting time of an increase or decrease in gas prices? It's all mathematics.

You are at an exciting point in your mathematical journey as the mathematics you are studying has interesting applications in the real world.

The physical sciences, social sciences, and the business world are full of situations that can be modeled with linear equations relating two variables. Data is collected and graphed. If the data points appear to form a straight line, an equation of that line can be used to predict the value of one variable based on the value of the other variable.

To create a mathematical model of a linear relation between two variables, we must be able to find the equation of the line. In this section we will look at several ways to write the equation of a line. The specific method we use will be determined by what information we are given.

Find an Equation of the Line Given the Slope and *y*-Intercept

We can easily determine the slope and intercept of a line if the equation was written in slope–intercept form, y=mx+b. Now, we will do the reverse—we will start with the slope and *y*-intercept and use them to find the equation of the line.

Example 4.6.1

Find an equation of a line with slope −7 and *y*-intercept (0,−1).

Solution

Since we are given the slope and *y*-intercept of the line, we can substitute the needed values into the slope–intercept form, y=mx+b.

Name the slope.	$m = -7$
Name the *y*-intercept.	*y*-intercept $(0, -1)$
Substitute the values into y=mx+b.	$y = mx + b$
	$y = -7x + (-1)$
	$y = -7x + -1$

Try It 4.6.2

Find an equation of a line with slope $\frac{2}{5}$ and y-intercept (0,4).

Answer

$y = \frac{2}{5}x + 4$

Try It 4.6.3

Find an equation of a line with slope −1 and y-intercept (0,−3).


Answer

$y = -x - 3$

Sometimes, the slope and intercept need to be determined from the graph.

Example 4.6.4

Find the equation of the line shown.



Solution

We need to find the slope and y-intercept of the line from the graph so we can substitute the needed values into the slope–intercept form, $y = mx + b$.


To find the slope, we choose two points on the graph.

The y-intercept is (0,−4) and the graph passes through (3,−2).

Find the slope by counting the rise and run.	$m = \frac{\text{rise}}{\text{run}}$
	$m = \frac{2}{3}$
Find the y-intercept.	y-intercept (0, −4)
Substitute the values into y=mx+b.	$y = mx + b$
	$y = \frac{2}{3}x - 4$

Try It 4.6.5

Find the equation of the line shown in the graph.




Answer

$y = \frac{3}{5}x + 1$

Try It 4.6.6

Find the equation of the line shown in the graph.



Answer

$y = \frac{4}{3}x - 5$

Find an Equation of the Line Given the Slope and a Point

Finding an equation of a line using the slope–intercept form of the equation works well when you are given the slope and *y*-intercept or when you read them off a graph. But what happens when you have another point instead of the *y*-intercept?

We are going to use the slope formula to derive another form of an equation of the line. Suppose we have a line that has slope mm and that contains some specific point (x_1, y_1) and some other point, which we will just call (x,y). We can write the slope of this line and then change it to a different form.

	m	$= \frac{y-y_1}{x-x_1}$
Multiply both sides of the equation by $x - x_1$.	$m(x-x_1)$	$= \left(\frac{y-y_1}{x-x_1}\right)(x-x_1)$
Simplify.	$m(x-x_1)$	$= y - y_1$
Rewrite the equation with the y terms on the left.	$y - y_1$	$= m(x-x_1)$

This format is called the point–slope form of an equation of a line.

POINT–SLOPE FORM OF AN EQUATION OF A LINE

The **point–slope form** of an equation of a line with slope m and containing the point (x_1, y_1) is

$$y - y_1 = m(x - x_1)$$

We can use the point–slope form of an equation to find an equation of a line when we are given the slope and one point. Then we will rewrite the equation in slope–intercept form. Most applications of linear equations use the the slope–intercept form.

Example 4.6.7: Find an Equation of a Line Given the Slope and a Point

Find an equation of a line with slope $m = \frac{2}{5}$ that contains the point (10,3). Write the equation in slope–intercept form.

Solution

Step 1. Identify the slope.	The slope is given.	$m = \frac{2}{5}$
Step 2. Identify the point.	The point is given.	(x_1, y_1) $(10, 3)$
Step 3. Substitute the values into the point–slope form, $y - y_1 = m(x - x_1)$.		$y - y_1 = m(x - x_1)$
		$y - 3 = \frac{2}{5}(x - 10)$
	Simplify.	$y - 3 = \frac{2}{5}x - 4$
Step 4. Write the equation in slope–intercept form.		$y = \frac{2}{5}x - 1$

Try It 4.6.8

Find an equation of a line with slope $m = \frac{5}{6}$ and containing the point (6,3).

Answer

$y = \frac{5}{6}x - 2$

Try It 4.6.9

Find an equation of a line with slope $m = \frac{2}{3}$ and containing the point (9,2).

Answer

$y = \frac{2}{3}x - 4$

FIND AN EQUATION OF A LINE GIVEN THE SLOPE AND A POINT.

1. Identify the slope.
2. Identify the point.
3. Substitute the values into the point-slope form, $y - y_1 = m(x - x_1)$.
4. Write the equation in slope–intercept form.

Example 4.6.10

Find an equation of a line with slope $m = -\frac{1}{3}$ that contains the point (6,−4). Write the equation in slope–intercept form.

Solution

Since we are given a point and the slope of the line, we can substitute the needed values into the point–slope form, $y - y_1 = m(x - x_1)$.

Identify the slope.	$m = -\frac{1}{3}$
Identify the point.	$\begin{matrix} x_1, & y_1 \\ 6, & -4 \end{matrix}$
Substitute the values into $y - y_1 = m(x - x_1)$.	$y - y_1 = m(x - x_1)$
	$y - (-4) = -\frac{1}{3}(x - 6)$
Simplify.	$y + 4 = -\frac{1}{3}x + 2$
Write in slope–intercept form.	$y = -\frac{1}{3}x - 2$

Try It 4.6.11

Find an equation of a line with slope $m = -\frac{2}{5}$ and containing the point (10,−5).

Answer

$y = -\frac{2}{5}x - 1$

Try It 4.6.12

Find an equation of a line with slope $m = -\frac{3}{4}$, and containing the point (4,−7).

Answer

$y = -\frac{3}{4}x - 4$

Example 4.6.13

Find an equation of a horizontal line that contains the point (−1,2). Write the equation in slope–intercept form.

Solution

Every horizontal line has slope 0. We can substitute the slope and points into the point–slope form, $y - y_1 = m(x - x_1)$.

Identify the slope.	$m = 0$
Identify the point.	$\begin{matrix} x_1, & y_1 \\ -1, & 2 \end{matrix}$
Substitute the values into $y - y_1 = m(x - x_1)$.	$y - y_1 = m(x - x_1)$
	$y - 2 = 0(x - (-1))$
Simplify.	$y - 2 = 0(x + 1)$
	$y - 2 = 0$
	$y = 2$

Write in slope–intercept form.	It is in *y*-form, but could be written y=0x+2.

Did we end up with the form of a horizontal line, y=a?

Try It 4.6.14

Find an equation of a horizontal line containing the point (−3,8).

Answer

y = 8

Try It 4.6.15

Find an equation of a horizontal line containing the point (−1,4).

Answer

y = 4

Find an Equation of the Line Given Two Points

When real-world data is collected, a linear model can be created from two data points. In the next example we'll see how to find an equation of a line when just two points are given.

We have two options so far for finding an equation of a line: slope–intercept or point–slope. Since we will know two points, it will make more sense to use the point–slope form.

But then we need the slope. Can we find the slope with just two points? Yes. Then, once we have the slope, we can use it and one of the given points to find the equation.

Example 4.6.16: Find an Equation of a Line Given Two Points

Find an equation of a line that contains the points (5,4) and (3,6). Write the equation in slope–intercept form.

Solution

Step 1. Find the slope using the given points.	To use the point-slope form, we first find the slope.	$m=\frac{y_2-y_1}{x_2-x_1}$ $m=\frac{6-4}{3-5}$ $m=\frac{2}{-2}$ $m=-1$
Step 2. Choose one point.	Choose either point.	$(\overset{x_1}{5},\ \overset{y_1}{4})$
Step 3. Substitute the values into the point-slope form, $y-y_1=m(x-x_1)$.	Simplify.	$y-y_1=m(x-x_1)$ $y-4=-1(x-5)$ $y-4=-1x+5$
Step 4. Write the equation in slope–intercept form.		$y=-1x+9$

Use the point (3,6) and see that you get the same equation.

Try It 4.6.17

Find an equation of a line containing the points (3,1) and (5,6).

Answer

$y = \frac{5}{2}x - \frac{13}{2}$

Try It 4.6.18

Find an equation of a line containing the points (1,4) and (6,2).

Answer

$y = -\frac{2}{5}x + \frac{22}{5}$

FIND AN EQUATION OF A LINE GIVEN TWO POINTS.

1. Find the slope using the given points.
2. Choose one point.
3. Substitute the values into the point-slope form, $y - y_1 = m(x - x_1)$.
4. Write the equation in slope–intercept form.

Example 4.6.19

Find an equation of a line that contains the points (−3,−1) and (2,−2). Write the equation in slope–intercept form.

Solution

Since we have two points, we will find an equation of the line using the point–slope form. The first step will be to find the slope.

Find the slope of the line through (−3, −1) and (2, −2).	$m = \frac{y_2 - y_1}{x_2 - x_1}$
	$m = \frac{-2-(-1)}{2-(-3)}$
	$m = \frac{-1}{5}$
	$m = -\frac{1}{5}$
Choose either point.	$\begin{matrix} x_1 & y_1 \\ 2, & -2 \end{matrix}$
Substitute the values into $y - y_1 = m(x - x_1)$.	$y - y_1 = m(x - x_1)$
	$y-(-2) = -\frac{1}{5}(x-2)$
	$y + 2 = -\frac{1}{5}x + \frac{2}{5}$
Write in slope–intercept form.	$y = -\frac{1}{5}x - \frac{8}{5}$

Try It 4.6.20

Find an equation of a line containing the points (−2,−4) and (1,−3).

Answer

$y = \frac{1}{3}x - \frac{10}{3}$

Try It 4.6.21

Find an equation of a line containing the points (−4,−3) and (1,−5).

Answer

$y = -\frac{2}{5}x - \frac{23}{5}$

Example 4.6.22

Find an equation of a line that contains the points (−2,4) and (−2,−3). Write the equation in slope–intercept form.

Solution

Again, the first step will be to find the slope.

Find the slope of the line through $(-2, 4)$ and $(-2, -3)$

$$\begin{aligned} m &= \frac{y_2 - x_1}{x_2 - x_1} \\ m &= \frac{-3-4}{-2-(-2)} \\ m &= \frac{-7}{0} \end{aligned}$$

The slope is undefined.

This tells us it is a vertical line. Both of our points have an *x*-coordinate of −2. So our equation of the line is x=−2. Since there is no yy, we cannot write it in slope–intercept form.

You may want to sketch a graph using the two given points. Does the graph agree with our conclusion that this is a vertical line?

Try It 4.6.23

Find an equation of a line containing the points (5,1) and (5,−4).

Answer

x = 5

Try It 4.6.24

Find an equation of a line containing the points (−4,4) and (−4,3).

Answer

x=−4

We have seen that we can use either the slope–intercept form or the point–slope form to find an equation of a line. Which form we use will depend on the information we are given. This is summarized in Table 4.6.1.

Table 4.6.1

To Write an Equation of a Line		
If given:	Use:	Form:
Slope and *y*-intercept	slope–intercept	y=mx+b

https://math.libretexts.org/@go/page/30513

Slope and a point	point–slope	$y - y_1 = m(x - x_1)$
Two points	point–slope	$y - y_1 = m(x - x_1)$

Find an Equation of a Line Parallel to a Given Line

Suppose we need to find an equation of a line that passes through a specific point and is parallel to a given line. We can use the fact that parallel lines have the same slope. So we will have a point and the slope—just what we need to use the point–slope equation.

First let's look at this graphically.

The graph shows the graph of y=2x−3. We want to graph a line parallel to this line and passing through the point (−2,1).




Figure 4.6.1

We know that parallel lines have the same slope. So the second line will have the same slope as y=2x−3. That slope is $m_{\parallel} = 2$. We'll use the notation $m_{\parallel}$ to represent the slope of a line parallel to a line with slope m. (Notice that the subscript ∥ looks like two parallel lines.)

The second line will pass through (−2,1) and have m=2. To graph the line, we start at (−2,1) and count out the rise and run. With m=2 (or $m = \frac{2}{1}$), we count out the rise 2 and the run 1. We draw the line.




Figure 4.6.2

Do the lines appear parallel? Does the second line pass through (−2,1)?

Now, let's see how to do this algebraically.

We can use either the slope–intercept form or the point–slope form to find an equation of a line. Here we know one point and can find the slope. So we will use the point–slope form.

Example 4.6.25: How to Find an Equation of a Line Parallel to a Given Line

Find an equation of a line parallel to y=2x−3 that contains the point (−2,1). Write the equation in slope–intercept form.

Solution

Step 1. Find the slope of the given line.	The line is in slope–intercept form, $y = 2x - 3$.	$m = 2$
Step 2. Find the slope of the parallel line.	Parallel lines have the same slope.	$m_{\parallel} = 2$
Step 3. Identify the point.	The given point is, (–2, 1).	$\begin{pmatrix} x_1 & y_1 \\ -2, & 1 \end{pmatrix}$
Step 4. Substitute the values into the point–slope form, $y - y_1 = m(x - x_1)$.	Simplify.	$y - y_1 = m(x - x_1)$ $y - 1 = 2(x - (-2))$ $y - 1 = 2(x + 2)$ $y - 1 = 2x + 4$
Step 5. Write the equation in slope–intercept form.		$y = 2x + 5$

Does this equation make sense? What is the *y*-intercept of the line? What is the slope?

? Try It 4.6.26

Find an equation of a line parallel to the line y=3x+1 that contains the point (4,2). Write the equation in slope–intercept form.

Answer

y=3x−10

? Try It 4.6.27

Find an equation of a line parallel to the line $y = \frac{1}{2}x - 3$ that contains the point (6,4).

Answer

$y = \frac{1}{2}x + 1$

FIND AN EQUATION OF A LINE PARALLEL TO A GIVEN LINE.

1. Find the slope of the given line.
2. Find the slope of the parallel line.
3. Identify the point.
4. Substitute the values into the point–slope form, $y - y_1 = m(x - x_1)$.
5. Write the equation in slope–intercept form.

Find an Equation of a Line Perpendicular to a Given Line

Now, let's consider perpendicular lines. Suppose we need to find a line passing through a specific point and which is perpendicular to a given line. We can use the fact that perpendicular lines have slopes that are negative reciprocals. We will again use the point–slope equation, like we did with parallel lines.

The graph shows the graph of y=2x−3. Now, we want to graph a line perpendicular to this line and passing through (−2,1).




Figure 4.6.3

We know that perpendicular lines have slopes that are negative reciprocals. We'll use the notation $m_{\perp}$ to represent the slope of a line perpendicular to a line with slope m. (Notice that the subscript $_{\perp}$ looks like the right angles made by two perpendicular lines.)

$$\begin{aligned} y &= 2x - 3 \quad \text{perpendicular line} \\ m &= 2 \qquad m_{\perp} = -\tfrac{1}{2} \end{aligned} \tag{4.6.1}$$

We now know the perpendicular line will pass through (−2,1) with $m_{\perp} = -\frac{1}{2}$.

To graph the line, we will start at (−2,1) and count out the rise −1 and the run 2. Then we draw the line.




Figure 4.6.4

Do the lines appear perpendicular? Does the second line pass through (−2,1)?

Now, let's see how to do this algebraically. We can use either the slope–intercept form or the point–slope form to find an equation of a line. In this example we know one point, and can find the slope, so we will use the point–slope form.

Example 4.6.28

Find an equation of a line perpendicular to y=2x−3 that contains the point (−2,1). Write the equation in slope–intercept form.

Solution

Step 1. Find the slope of the given line.	The line is in slope–intercept form, $y = 2x - 3$.	$m = 2$
Step 2. Find the slope of the perpendicular line.	The slopes of perpendicular lines are negative reciprocals.	$m_{\perp} = -\frac{1}{2}$
Step 3. Identify the point.	The given point is, (–2, 1)	(x_1, y_1) $(-2, 1)$

Step 4. Substitute the values into the point–slope form, $y-y_1=m(x-x_1)$.	Simplify.	$y-y_1=m(x-x_1)$ $y-1=-\frac{1}{2}(x-(-2))$ $y-1=-\frac{1}{2}(x+2)$ $y-1=-\frac{1}{2}x-1$
Step 5. Write the equation in slope–intercept form.		$y=-\frac{1}{2}x$

Try It 4.6.29

Find an equation of a line perpendicular to the line y=3x+1 that contains the point (4,2). Write the equation in slope–intercept form.

Answer

$y=-\frac{1}{3}x+\frac{10}{3}$

Try It 4.6.30

Find an equation of a line perpendicular to the line $y=\frac{1}{2}x-3$ that contains the point (6,4).

Answer

y=−2x+16

FIND AN EQUATION OF A LINE PERPENDICULAR TO A GIVEN LINE.

1. Find the slope of the given line.
2. Find the slope of the perpendicular line.
3. Identify the point.
4. Substitute the values into the point–slope form, $y-y_1=m(x-x_1)$.
5. Write the equation in slope–intercept form.

Example 4.6.31

Find an equation of a line perpendicular to x=5 that contains the point (3,−2). Write the equation in slope–intercept form.

Solution

Again, since we know one point, the point–slope option seems more promising than the slope–intercept option. We need the slope to use this form, and we know the new line will be perpendicular to x=5. This line is vertical, so its perpendicular will be horizontal. This tells us the $m_\perp=0$.

Identify the point.	$(3$	,	$-2)$
Identify the slope of the perpendicular line.	$m_\perp$	$=$	0
Substitute the values into $y-y_1=m\,(x-x_1)$.	$y-y_1$	$=$	$m\,(x-x_1)$
	$y-(-2)$	$=$	$0(x-3)$
Simplify.	$y+2$	$=$	0
	y	$=$	-2

Sketch the graph of both lines. Do they appear to be perpendicular?

Try It 4.6.32

Find an equation of a line that is perpendicular to the line $x=4$ that contains the point $(4,-5)$. Write the equation in slope–intercept form.

Answer

$y=-5$

Try It 4.6.33

Find an equation of a line that is perpendicular to the line $x=2$ that contains the point $(2,-1)$. Write the equation in slope–intercept form.

Answer

$y=-1$

In Example 4.6.31, we used the point–slope form to find the equation. We could have looked at this in a different way.

We want to find a line that is perpendicular to $x=5$ that contains the point $(3,-2)$. The graph shows us the line $x=5$ and the point $(3,-2)$.




Figure 4.6.5

We know every line perpendicular to a vertical line is horizontal, so we will sketch the horizontal line through $(3,-2)$.




Figure 4.6.6

Do the lines appear perpendicular?

If we look at a few points on this horizontal line, we notice they all have *y*-coordinates of -2. So, the equation of the line perpendicular to the vertical line $x=5$ is $y=-2$.

Example 4.6.34

Find an equation of a line that is perpendicular to y=−4 that contains the point (−4,2). Write the equation in slope–intercept form.

Solution

The line y=−4 is a horizontal line. Any line perpendicular to it must be vertical, in the form x=a. Since the perpendicular line is vertical and passes through (−4,2), every point on it has an *x*-coordinate of −4. The equation of the perpendicular line is x=−4. You may want to sketch the lines. Do they appear perpendicular?

Try It 4.6.35

Find an equation of a line that is perpendicular to the line y=1 that contains the point (−5,1). Write the equation in slope–intercept form.

Answer

x=−5

Try It 4.6.36

Find an equation of a line that is perpendicular to the line y=−5 that contains the point (−4,−5).

Answer

x=−4

Note

Access this online resource for additional instruction and practice with finding the equation of a line.

- Use the Point-Slope Form of an Equation of a Line

Key Concepts

- **To Find an Equation of a Line Given the Slope and a Point**
  1. Identify the slope.
  2. Identify the point.
  3. Substitute the values into the point-slope form, $y-y_1=m(x-x_1)$.
  4. Write the equation in slope-intercept form.
- **To Find an Equation of a Line Given Two Points**
  1. Find the slope using the given points.
  2. Choose one point.
  3. Substitute the values into the point-slope form, $y-y_1=m(x-x_1)$.
  4. Write the equation in slope-intercept form.
- **To Write an Equation of a Line**
  - If given slope and y-intercept, use slope–intercept form $y=mx+b$.
  - If given slope and a point, use point–slope form $y-y_1=m(x-x_1)$.
  - If given two points, use point–slope form $y-y_1=m(x-x_1)$.
- **To Find an Equation of a Line Parallel to a Given Line**
  1. Find the slope of the given line.
  2. Find the slope of the parallel line.
  3. Identify the point.
  4. Substitute the values into the point-slope form, $y-y_1=m(x-x_1)$.

5. Write the equation in slope-intercept form.

- **To Find an Equation of a Line Perpendicular to a Given Line**
  1. Find the slope of the given line.
  2. Find the slope of the perpendicular line.
  3. Identify the point.
  4. Substitute the values into the point-slope form, $y - y_1 = m(x - x_1)$.
  5. Write the equation in slope-intercept form.

Glossary

point–slope form

The point–slope form of an equation of a line with slope mm and containing the point (x_1, y_1) is $y - y_1 = m\,(x - x_1)$.

This page titled 4.6: Find the Equation of a Line is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.6: Find the Equation of a Line** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.6E: Exercises

Practice Makes Perfect

Find an Equation of the Line Given the Slope and y-Intercept

In the following exercises, find the equation of a line with given slope and y-intercept. Write the equation in slope–intercept form.

Exercise $4.6E.1$

slope 3 and y-intercept $(0, 5)$

Exercise $4.6E.2$

slope 4 and y-intercept $(0, 1)$

Answer

$y = 4x + 1$

Exercise $4.6E.3$

slope 6 and y-intercept $(0, -4)$

Exercise $4.6E.4$

slope 8 and y-intercept $(0, -6)$

Answer

$y = 8x - 6$

Exercise $4.6E.5$

slope -1 and y-intercept $(0, 3)$

Exercise $4.6E.6$

slope -1 and y-intercept $(0, 7)$

Answer

$y = -x + 7$

Exercise $4.6E.7$

slope -2 and y-intercept $(0, -3)$

Exercise $4.6E.8$

slope -3 and y-intercept $(0, -1)$

Answer

$y = -3x - 1$

Exercise 4.6E.9

slope $\frac{3}{5}$ and y-intercept $(0, -1)$

Exercise 4.6E.10

slope $\frac{1}{5}$ and y-intercept $(0, -5)$

Answer

$y = \frac{1}{5}x - 5$

Exercise 4.6E.11

slope $-\frac{3}{4}$ and y-intercept $(0, -2)$

Exercise 4.6E.12

slope $-\frac{2}{3}$ and y-intercept $(0, -3)$

Answer

$y = -\frac{2}{3}x - 3$

Exercise 4.6E.13

slope 0 and y-intercept $(0, -1)$

Exercise 4.6E.14

slope 0 and y-intercept $(0, 2)$

Answer

$y = 2$

Exercise 4.6E.15

slope -3 and y-intercept $(0, 0)$

Exercise 4.6E.16


slope -4 and y-intercept $(0, 0)$

Answer


$y = -4x$

In the following exercises, find the equation of the line shown in each graph. Write the equation in slope–intercept form.

Exercise 4.6E.17




Exercise $4.6E.18$




Answer

$y = -2x + 4$

Exercise $4.6E.19$




Exercise $4.6E.20$




Answer

$y = \frac{3}{4}x + 2$

Exercise $4.6E.21$




Exercise $4.6E.22$




Answer

$y = -\frac{3}{2}x - 1$

Exercise $4.6E.23$



? Exercise $4.6E.24$



Answer

$y = 6$

Find an Equation of the Line Given the Slope and a Point

In the following exercises, find the equation of a line with given slope and containing the given point. Write the equation in slope–intercept form.

? Exercise $4.6E.25$

$m = \frac{5}{8}$, point $(8, 3)$

? Exercise $4.6E.26$

$m = \frac{3}{8}$, point $(8, 2)$

Answer

$y = \frac{3}{8}x - 1$

Exercise 4.6E.27

$m = \frac{1}{6}$, point $(6, 1)$

Exercise 4.6E.28

$m = \frac{5}{6}$, point $(6, 7)$

Answer

$y = \frac{5}{6}x + 2$

Exercise 4.6E.29

$m = -\frac{3}{4}$, point $(8, -5)$

Exercise 4.6E.30

$m = -\frac{3}{5}$, point $(10, -5)$

Answer

$y = -\frac{3}{5}x + 1$

Exercise 4.6E.31

$m = -\frac{1}{4}$, point $(-12, -6)$

Exercise 4.6E.32

$m = -\frac{1}{3}$, point $(-9, -8)$

Answer

$y = -\frac{1}{3}x - 11$

Exercise 4.6E.33

Horizontal line containing $(-2, 5)$

Exercise 4.6E.34

Horizontal line containing $(-1, 4)$

Answer

$y = 4$

Exercise 4.6E.35

Horizontal line containing $(-2, -3)$

Exercise 4.6E.36

Horizontal line containing $(-1, -7)$

Answer

$y = -7$

Exercise 4.6E.37

$m = -\frac{3}{2}$, point $(-4, -3)$

Exercise 4.6E.38

$m = -\frac{5}{2}$, point $(-8, -2)$

Answer

$y = -\frac{5}{2}x - 22$

Exercise 4.6E.39

$m = -7$, point $(-1, -3)$

Exercise 4.6E.40

$m = -4$, point $(-2, -3)$

Answer

$y = -4x - 11$

Exercise 4.6E.41

Horizontal line containing $(2, -3)$

Exercise 4.6E.42

Horizontal line containing $(4, -8)$

Answer

$y = -8$

Find an Equation of the Line Given Two Points

In the following exercises, find the equation of a line containing the given points. Write the equation in slope–intercept form.

Exercise 4.6E.43

$(2, 6)$ and $(5, 3)$

Exercise 4.6E.44

$(3, 1)$ and $(2, 5)$

Answer

$y = -4x + 13$

Exercise 4.6E.45

$(4, 3)$ and $(8, 1)$

Exercise 4.6E.46

$(2, 7)$ and $(3, 8)$

Answer

$y = x + 5$

Exercise 4.6E.47

$(-3, -4)$ and $(5, -2)$

Exercise 4.6E.48

$(-5, -3)$ and $(4, -6)$

Answer

$y = -\frac{1}{3}x - \frac{14}{3}$

Exercise 4.6E.49

$(-1, 3)$ and $(-6, -7)$

Exercise 4.6E.50

$(-2, 8)$ and $(-4, -6)$

Answer

$y = 7x + 22$

Exercise 4.6E.51

$(6, -4)$ and $(-2, 5)$

Exercise 4.6E.52

$(3, -2)$ and $(-4, 4)$

Answer

$y = -\frac{6}{7}x + \frac{4}{7}$

Exercise 4.6E.53

$(0, 4)$ and $(2, -3)$

Exercise 4.6E.54

$(0, -2)$ and $(-5, -3)$

Answer

$y = \frac{1}{5}x - 2$

Exercise 4.6E.55

$(7, 2)$ and $(7, -2)$

Exercise 4.6E.56

$(4, 2)$ and $(4, -3)$

Answer

$x = 4$

Exercise 4.6E.57

$(-7, -1)$ and $(-7, -4)$

Exercise 4.6E.58

$(-2, 1)$ and $(-2, -4)$

Answer

$x = -2$

Exercise 4.6E.59

$(6, 1)$ and $(0, 1)$

Exercise 4.6E.60

$(6, 2)$ and $(-3, 2)$

Answer

$y = 2$

Exercise 4.6E.61

$(3, -4)$ and $(5, -4)$

Exercise 4.6E.62

$(-6, -3)$ and $(-1, -3)$

Answer

$y = -3$

Exercise 4.6E.63

$(4, 3)$ and $(8, 0)$

Exercise 4.6E.64

$(0, 0)$ and $(1, 4)$

Answer

$y = 4x$

Exercise 4.6E.65

$(-2, -3)$ and $(-5, -6)$

Exercise 4.6E.66

$(-3, 0)$ and $(-7, -2)$

Answer

$y = \frac{1}{2}x + \frac{3}{2}$

Exercise 4.6E.67

$(8, -1)$ and $(8, -5)$

Exercise 4.6E.68

$(3, 5)$ and $(-7, 5)$

Answer

$y = 5$

Find an Equation of a Line Parallel to a Given Line

In the following exercises, find an equation of a line parallel to the given line and contains the given point. Write the equation in slope–intercept form.

Exercise 4.6E.69

line $y = 4x + 2$, point $(1, 2)$

Exercise 4.6E.70

line $y = 3x + 4$, point $(2, 5)$

Answer

$y = 3x - 1$

Exercise 4.6E.71

line $y = -2x - 3$, point $(-1, 3)$

Exercise 4.6E.72

line $y = -3x - 1$, point $(2, -3)$

Answer

$y = -3x + 3$

Exercise 4.6E.73

line $3x - y = 4$, point $(3, 1)$

Exercise 4.6E.74

line $2x - y = 6$, point $(3, 0)$

Answer

$y = 2x - 6$

Exercise 4.6E.75

line $4x + 3y = 6$, point $(0, -3)$

Exercise 4.6E.76

line $2x + 3y = 6$, point $(0, 5)$

Answer

$y = -\frac{2}{3}x + 5$

Exercise 4.6E.77

line $x = -3$, point $(-2, -1)$

Exercise 4.6E.78

line $x = -4$, point $(-3, -5)$

Answer

$x = -3$

Exercise 4.6E.79

line $x - 2 = 0$, point $(1, -2)$

Exercise 4.6E.80

line $x-6=0$, point $(4,-3)$

Answer

$x=4$

Exercise 4.6E.81

line $y=5$, point $(2,-2)$

Exercise 4.6E.82

line $y=1$, point $(3,-4)$

Answer

$y=-4$

Exercise 4.6E.83

line $y+2=0$, point $(3,-3)$

Exercise 4.6E.84

line $y+7=0$, point $(1,-1)$

Answer

$y=-1$

Find an Equation of a Line Perpendicular to a Given Line

In the following exercises, find an equation of a line perpendicular to the given line and contains the given point. Write the equation in slope–intercept form.

Exercise 4.6E.85

line $y=-2x+3$, point $(2,2)$

Exercise 4.6E.86

line $y=-x+5$, point $(3,3)$

Answer

$y=x$

Exercise 4.6E.87

line $y=\frac{3}{4}x-2$, point $(-3,4)$

Exercise 4.6E.88

line $y = \frac{2}{3}x - 4$, point $(2, -4)$

Answer

$y = -\frac{3}{2}x - 1$

Exercise 4.6E.89

line $2x - 3y = 8$, point $(4, -1)$

Exercise 4.6E.90

line $4x - 3y = 5$, point $(-3, 2)$

Answer

$y = -\frac{3}{4}x - \frac{1}{4}$

Exercise 4.6E.91

line $2x + 5y = 6$, point $(0, 0)$

Exercise 4.6E.92

line $4x + 5y = -3$, point $(0, 0)$

Answer

$y = \frac{5}{4}x$

Exercise 4.6E.93

line $y - 3 = 0$, point $(-2, -4)$

Exercise 4.6E.94

line $y - 6 = 0$, point $(-5, -3)$

Answer

$x = -5$

Exercise 4.6E.95

line y-axis, point $(3, 4)$

Exercise 4.6E.96

line y-axis, point $(2, 1)$

Answer

$y = 1$

Mixed Practice

In the following exercises, find the equation of each line. Write the equation in slope–intercept form.

Exercise 4.6E.97

Containing the points $(4, 3)$ and $(8, 1)$

Exercise 4.6E.98

Containing the points $(2, 7)$ and $(3, 8)$

Answer

$y = x + 5$

Exercise 4.6E.99

$m = \frac{1}{6}$, containing point $(6, 1)$

Exercise 4.6E.100

$m = \frac{5}{6}$, containing point $(6, 7)$

Answer

$y = \frac{5}{6}x + 2$

Exercise 4.6E.101

Parallel to the line $4x + 3y = 6$, containing point $(0, -3)$

Exercise 4.6E.102

Parallel to the line $2x + 3y = 6$, containing point $(0, 5)$

Answer

$y = -\frac{2}{3}x + 5$

Exercise 4.6E.103

$m = -\frac{3}{4}$, containing point $(8, -5)$

Exercise 4.6E.104

$m = -\frac{3}{5}$, containing point $(10, -5)$

Answer

$y = -\frac{3}{5}x + 1$

Exercise 4.6E.105

Perpendicular to the line $y-1=0,$ point $(-2,6)$

Exercise 4.6E.106

Perpendicular to the line y-axis, point $(-6,2)$

Answer

$y=2$

Exercise 4.6E.107

Containing the points $(4,3)$ and $(8,1)$

Exercise 4.6E.108

Containing the points $(-2,0)$ and $(-3,-2)$

Answer

$y=x+2$

Exercise 4.6E.109

Parallel to the line $x=-3$, containing point $(-2,-1)$

Exercise 4.6E.110

Parallel to the line $x=-4$, containing point $(-3,-5)$

Answer

$x=-3$

Exercise 4.6E.111

Containing the points $(-3,-4)$ and $(2,-5)$

Exercise 4.6E.112

Containing the points $(-5,-3)$ and $(4,-6)$

Answer

$y=-\frac{1}{3}x-\frac{14}{3}$

Exercise 4.6E.113

Perpendicular to the line $x-2y=5,$ containing point $(-2,2)$

Exercise 4.6E.114

Perpendicular to the line $4x+3y=1$, containing point $(0,0)$

Answer

$y=\frac{3}{4}x$

Everyday Math

Exercise 4.6E.115

Cholesterol. The age, x, and LDL cholesterol evel, y, of two men are given by the points $(18,68)$ and $(27,122)$. Find a linear equation that models the relationship between age and LDL cholesterol level.

Exercise 4.6E.116

Fuel consumption. The city mpg, x, and highway mpg, y, of two cars are given by the points $(29,40)$ and $(19,28)$. Find a linear equation that models the relationship between city mpg and highway mpg.

Answer

$y=1.2x+5.2$

Writing Exercises

Exercise 4.6E.117

Why are all horizontal lines parallel?

Exercise 4.6E.118

Explain in your own words why the slopes of two perpendicular lines must have opposite signs.

Answer

Answers will vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
find the equation of the line given the slope and *y*-intercept.			
find an equation of the line given the slope and a point.			
find an equation of the line given two points.			
find an equation of a line parallel to a given line.			
find an equation of a line perpendicular to a given line.			

b. On a scale of 1-10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 4.6E: Exercises is shared under a not declared license and was authored, remixed, and/or curated by OpenStax.

- **4.6E: Exercises** by OpenStax has no license indicated. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.7: Graphs of Linear Inequalities

Learning Objectives

By the end of this section, you will be able to:

- Verify solutions to an inequality in two variables
- Recognize the relation between the solutions of an inequality and its graph
- Graph linear inequalities

Note

Before you get started, take this readiness quiz.

1. Solve: $4x+3>23$.
   If you missed this problem, review Exercise 2.7.22.
2. Translate from algebra to English: $x<5$.
   If you missed this problem, review Exercise 1.3.1.
3. Evaluate $3x-2y$ when $x=1,\ y=-2$.
   If you missed this problem, review Exercise 1.5.28.

Verify Solutions to an Inequality in Two Variables

We have learned how to solve inequalities in one variable. Now, we will look at inequalities in two variables. Inequalities in two variables have many applications. If you ran a business, for example, you would want your revenue to be greater than your costs—so that your business would make a profit.

LINEAR INEQUALITY

A **linear inequality** is an inequality that can be written in one of the following forms:

$$Ax+By>C \quad Ax+By\geq C \quad Ax+By<C \quad Ax+By\leq C$$

where A and B are not both zero.

Do you remember that an inequality with one variable had many solutions? The solution to the inequality $x>3$ is any number greater than 3. We showed this on the number line by shading in the number line to the right of 3, and putting an open parenthesis at 3. See Figure 4.7.1.




Figure 4.7.1

Similarly, inequalities in two variables have many solutions. Any ordered pair (x,y) that makes the inequality true when we substitute in the values is a solution of the inequality.

Solution OF A LINEAR INEQUALITY

An ordered pair (x,y) is a **solution of a linear inequality** if the inequality is true when we substitute the values of x and y.

Example 4.7.1

Determine whether each ordered pair is a solution to the inequality $y>x+4$:

1. $(0,0)$
2. $(1,6)$
3. $(2,6)$
4. $(-5,-15)$

5. $(-8, 12)$

Solution

1.

$(0, 0)$	$y > x + 4$
Substitute 0 for x and 0 for y.	$0 \stackrel{?}{>} 0 + 4$
Simplify.	$0 \not> 4$ So, $(0, 0)$ is not a solution to $y > x + 4$.

2.

$(1, 6)$	$y > x + 4$
Substitute 1 for x and 6 for y.	$6 \stackrel{?}{>} 1 + 4$
Simplify.	$6 > 5$ So, $(1, 6)$ is a solution to $y > x + 4$.

3.

$(2, 6)$	$y > x + 4$
Substitute 2 for x and 6 for y.	$6 \stackrel{?}{>} 2 + 4$
Simplify.	$6 \not> 6$ So, $(2, 6)$ is not a solution to $y > x + 4$.

4.

$(-5, -15)$	$y > x + 4$
Substitute –5 for x and –15 for y.	$-15 \stackrel{?}{>} -5 + 4$
Simplify.	$-15 \not> -1$ So, $(-5, -15)$ is not a solution to $y > x + 4$.

5.

$(-8,12)$	$y > x + 4$
Substitute –8 for x and 12 for y.	$12 \stackrel{?}{>} -8 + 4$
Simplify.	$12 > -4$ So, $(-8, 12)$ is a solution to $y > x + 4$.

? Try It 4.7.2

Determine whether each ordered pair is a solution to the inequality $y > x - 3$:

1. $(0, 0)$
2. $(4, 9)$
3. $(-2, 1)$
4. $(-5, -3)$
5. $(5, 1)$

Answer

1. yes

2. yes
3. yes
4. yes
5. no

Try It 4.7.3

Determine whether each ordered pair is a solution to the inequality $y < x + 1$:

1. $(0, 0)$
2. $(8, 6)$
3. $(-2, -1)$
4. $(3, 4)$
5. $(-1, -4)$

Answer

1. yes
2. yes
3. no
4. no
5. yes

Recognize the Relation Between the Solutions of an Inequality and its Graph

Now, we will look at how the solutions of an inequality relate to its graph.

Let's think about the number line in Figure 4.7.1 again. The point $x = 3$ separated that number line into two parts. On one side of 3 are all the numbers less than 3. On the other side of 3 all the numbers are greater than 3. See Figure 4.7.2.




Figure 4.7.2

The solution to $x > 3$ is the shaded part of the number line to the right of $x = 3$.

Similarly, the line $y = x + 4$ separates the plane into two regions. On one side of the line are points with $y < x + 4$. On the other side of the line are the points with $y > x + 4$. We call the line $y = x + 4$ a boundary line.

BOUNDARY LINE

The line with equation $Ax + By = C$ is the **boundary line** that separates the region where $Ax + By > C$ from the region where $Ax + By < C$.

For an inequality in one variable, the endpoint is shown with a parenthesis or a bracket depending on whether or not aa is included in the solution:



Similarly, for an inequality in two variables, the boundary line is shown with a solid or dashed line to indicate whether or not it the line is included in the solution. This is summarized in Table 4.7.1.

Table 4.7.1

$Ax + By < C$	$Ax + By \leq C$
$Ax + By > C$	$Ax + By \geq C$
Boundary line is not included in solution.	Boundary line is included in solution.

Boundary line is dashed.	Boundary line is solid.

Now, let's take a look at what we found in Example 4.7.1. We'll start by graphing the line $y = x + 4$, and then we'll plot the five points we tested. See Figure 4.7.3.




Figure 4.7.3

In Example 4.7.1 we found that some of the points were solutions to the inequality $y > x + 4$ and some were not.

Which of the points we plotted are solutions to the inequality $y > x + 4$? The points $(1, 6)$ and $(-8, 12)$ are solutions to the inequality $y > x + 4$. Notice that they are both on the same side of the boundary line $y = x + 4$.

The two points $(0, 0)$ and $(-5, -15)$ are on the other side of the **boundary line** $y = x + 4$, and they are not solutions to the inequality $y > x + 4$. For those two points, $y < x + 4$.

What about the point $(2, 6)$? Because $6 = 2 + 4$, the point is a solution to the equation $y = x + 4$. So the point $(2, 6)$ is on the boundary line.

Let's take another point on the left side of the boundary line and test whether or not it is a solution to the inequality $y > x + 4$. The point $(0, 10)$ clearly looks to be to the left of the boundary line, doesn't it? Is it a solution to the inequality?

$$\begin{aligned} &y > x + 4 \\ &10 \overset{?}{>} 0 + 4 \\ &10 > 4 \qquad \text{So, } (0, 10) \text{ is a solution to } y > x + 4. \end{aligned} \tag{4.7.1}$$

Any point you choose on the left side of the boundary line is a solution to the inequality $y > x + 4$. All points on the left are solutions.

Similarly, all points on the right side of the boundary line, the side with $(0, 0)$ and $(-5, -15)$, are not solutions to $y > x + 4$. See Figure 4.7.4.




Figure 4.7.4

The graph of the inequality $y > x + 4$ is shown in Figure $4.7.5$ below. The line $y = x + 4$ divides the plane into two regions. The shaded side shows the solutions to the inequality $y > x + 4$.

The points on the boundary line, those where $y = x + 4$, are not solutions to the inequality $y > x + 4$, so the line itself is not part of the solution. We show that by making the line dashed, not solid.





Figure 4.7.5: The graph of the inequality y>x+4.

Example 4.7.4

The boundary line shown is $y = 2x - 1$. Write the inequality shown by the graph.



Solution

The line $y = 2x - 1$ is the boundary line. On one side of the line are the points with $y > 2x - 1$ and on the other side of the line are the points with $y < 2x - 1$.

Let's test the point $(0, 0)$ and see which inequality describes its side of the boundary line.

At $(0, 0)$, which inequality is true:

$$\begin{array}{lll} y > 2x - 1 & \text{or} & y < 2x - 1? \\ y > 2x - 1 & & y < 2x - 1 \\ 0 > 2 \cdot 0 - 1 & & 0 < 2 \cdot 0 - 1 \\ 0 > -1 \text{ True} & & 0 < -1 \text{ False} \end{array} \tag{4.7.2}$$

Since $y > 2x - 1$ is true, the side of the line with $(0, 0)$, is the solution. The shaded region shows the solution of the inequality $y > 2x - 1$.


Since the boundary line is graphed with a solid line, the inequality includes the equal sign.

The graph shows the inequality $y \geq 2x - 1$.

We could use any point as a test point, provided it is not on the line. Why did we choose $(0, 0)$? Because it's the easiest to evaluate. You may want to pick a point on the other side of the boundary line and check that $y < 2x - 1$.

? Try It 4.7.5

Write the inequality shown by the graph with the boundary line $y = -2x + 3$.




Answer

$y \geq -2x + 3$

Try It 4.7.6

Write the inequality shown by the graph with the boundary line $y = \frac{1}{2}x - 4$.




Answer

$y \leq \frac{1}{2}x - 4$

Example 4.7.7

The boundary line shown is $2x + 3y = 6$. Write the inequality shown by the graph.



Solution

The line $2x + 3y = 6$ is the boundary line. On one side of the line are the points with $2x + 3y > 6$ and on the other side of the line are the points with $2x + 3y < 6$.

Let's test the point $(0, 0)$ and see which inequality describes its side of the boundary line.

At $(0, 0)$, which inequality is true:

$$\begin{aligned} 2x + 3y &> 6 \\ 2x + 3y &> 6 \\ 2(0) + 3(0) &> 6 \\ 0 &> 6 \quad \text{False} \end{aligned} \qquad \text{or} \qquad \begin{aligned} 2x + 3y &< 6? \\ 2x + 3y &< 6 \\ 2(0) + 3(0) &< 6 \\ 0 &< 6 \quad \text{True} \end{aligned} \tag{4.7.3}$$

So the side with $(0, 0)$ is the side where $2x + 3y < 6$.


(You may want to pick a point on the other side of the boundary line and check that $2x + 3y > 6$.)

Since the boundary line is graphed as a dashed line, the inequality does not include an equal sign.

The graph shows the solution to the inequality $2x + 3y < 6$.

Try It 4.7.8

Write the inequality shown by the shaded region in the graph with the boundary line $x - 4y = 8$.




Answer

$x - 4y \leq 8$

Try It 4.7.9

Write the inequality shown by the shaded region in the graph with the boundary line $3x - y = 6$.



Answer

$3x - y \leq 6$



Graph Linear Inequalities

Now, we're ready to put all this together to graph linear inequalities.

Example 4.7.10: How to Graph Linear Inequalities

Graph the **linear inequality** $y \geq \frac{3}{4}x - 2$.


Solution

Step 1. Identify and graph the boundary line. • If the inequality is ≤ or ≥, the boundary line is solid. • If the inequality is < or >, the boundary line is dashed.	Replace the inequality sign with an equal sign to find the boundary line. Graph the boundary line $y = \frac{3}{4}x - 2$. The inequality sign is ≥, so we draw a solid line.	
Step 2. Test a point that is not on the boundary line. Is it a solution of the inequality?	We'll test (0, 0). Is it a solution of the inequality?	At (0, 0), is $y \geq \frac{3}{4}x - 2$? $0 \stackrel{?}{\geq} \frac{3}{4}(0) - 2$ $0 \geq -2$ So, (0, 0) is a solution.
Step 3. Shade in one side of the boundary line. • If the test point is a solution, shade in the side that includes the point. • If the test point is not a solution, shade in the opposite side.	The test point (0, 0), is a solution to $y \geq \frac{3}{4}x - 2$. So we shade in that side.	

Try It 4.7.11

Graph the linear inequality $y \geq \frac{5}{2}x - 4$.

Answer




https://math.libretexts.org/@go/page/30515

Try It 4.7.12

Graph the linear inequality $y < \frac{2}{3}x - 5$.

Answer



The steps we take to graph a linear inequality are summarized here.

GRAPH A LINEAR INEQUALITY.


1. Identify and graph the boundary line.
   - If the inequality is $\leq$ or $\geq$, the boundary line is solid.
   - If the inequality is $<$ or $>$, the boundary line is dashed.
2. Test a point that is not on the boundary line. Is it a solution of the inequality?
3. Shade in one side of the boundary line.
   - If the test point is a solution, shade in the side that includes the point.
   - If the test point is not a solution, shade in the opposite side.

Example 4.7.13

Graph the linear inequality $x - 2y < 5$.

Solution

First we graph the boundary line $x - 2y = 5$. The inequality is $<$ so we draw a dashed line.



Then we test a point. We'll use $(0, 0)$ again because it is easy to evaluate and it is not on the boundary line.

Is $(0, 0)$ a solution of $x - 2y < 5$?


$$0-2(0) \stackrel{?}{<} 5$$
$$0-0 \stackrel{?}{<} 5$$
$$0<5$$

The point $(0,0)$ is a solution of $x-2y<5$, so we shade in that side of the boundary line.

Try It 4.7.14

Graph the linear inequality $2x-3y \leq 6$.


Answer



Try It 4.7.15

Graph the linear inequality $2x-y>3$.

Answer




What if the boundary line goes through the origin? Then we won't be able to use $(0,0)$ as a test point. No problem—we'll just choose some other point that is not on the boundary line.

Example 4.7.16

Graph the linear inequality $y \leq -4x$.

Solution

First we graph the boundary line $y=-4x$. It is in slope–intercept form, with $m=-4$ and $b=0$. The inequality is $\leq$ so we draw a solid line.



Now, we need a test point. We can see that the point $(1, 0)$ is not on the boundary line.

Is $(1, 0)$ a solution of $y \leq -4x$?

$0 \stackrel{?}{\leq} -4(1)$

$0 \not\leq -4$

The point $(1, 0)$ is not a solution to $y \leq -4x$, so we shade in the opposite side of the boundary line. See Figure 4.7.6.





Figure 4.7.6

? Try It 4.7.17

Graph the linear inequality $y > -3x$.


Answer



Try It 4.7.18

Graph the linear inequality $y \geq -2x$.

Answer



Some linear inequalities have only one variable. They may have an x but no y, or a y but no x. In these cases, the boundary line will be either a vertical or a horizontal line. Do you remember?

$x = a$ vertical line
$y = b$ horizontal line

Example 4.7.19

Graph the linear inequality $y > 3$.

Solution


First we graph the boundary line $y = 3$. It is a horizontal line. The inequality is $>$ so we draw a dashed line.

We test the point $(0, 0)$.

$$\begin{array}{l} y > 3 \\ 0 \not> 3 \end{array} \tag{4.7.4}$$

$(0, 0)$ is not a solution to $y > 3$.


So we shade the side that does not include $(0, 0)$.



Try It 4.7.20

Graph the linear inequality $y < 5$.


Answer



Try It 4.7.21

Graph the linear inequality $y \leq -1$.

Answer



Key Concepts

- **To Graph a Linear Inequality**
  1. Identify and graph the boundary line.
     If the inequality is $\leq$ or $\geq$, the boundary line is solid.
     If the inequality is $<$ or $>$, the boundary line is dashed.
  2. Test a point that is not on the boundary line. Is it a solution of the inequality?
  3. Shade in one side of the boundary line.
     If the test point is a solution, shade in the side that includes the point.
     If the test point is not a solution, shade in the opposite side.

Glossary

boundary line

The line with equation $Ax + By = C$ that separates the region where $Ax + By > C$ from the region where $Ax + By < C$.

linear inequality

An inequality that can be written in one of the following forms:

$$Ax + By > C \quad Ax + By \geq C \quad Ax + By < C \quad Ax + By \leq C \tag{4.7.5}$$

where A and B are not both zero.

solution of a linear inequality

An ordered pair $(x,\ y)$ is a solution to a linear inequality the inequality is true when we substitute the values of x and y.

This page titled 4.7: Graphs of Linear Inequalities is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.7: Graphs of Linear Inequalities** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

4.7E: Exercises

Practice Makes Perfect

Verify Solutions to an Inequality in Two Variables

In the following exercises, determine whether each ordered pair is a solution to the given inequality.

Exercise 4.7E.1

Determine whether each ordered pair is a solution to the inequality $y > x - 1$:

1. $(0, 1)$
2. $(-4, -1)$
3. $(4, 2)$
4. $(3, 0)$
5. $(-2, -3)$

Exercise 4.7E.2

Determine whether each ordered pair is a solution to the inequality $y > x - 3$:

1. $(0, 0)$
2. $(2, 1)$
3. $(-1, -5)$
4. $(-6, -3)$
5. $(1, 0)$

Answer

1. yes
2. no
3. no
4. yes
5. no

Exercise 4.7E.3

Determine whether each ordered pair is a solution to the inequality $y < x + 2$:

1. $(0, 3)$
2. $(-3, -2)$
3. $(-2, 0)$
4. $(0, 0)$
5. $(-1, 4)$

Exercise 4.7E.4

Determine whether each ordered pair is a solution to the inequality $y < x + 5$:

1. $(-3, 0)$
2. $(1, 6)$
3. $(-6, -2)$
4. $(0, 1)$
5. $(5, -4)$

Answer

1. yes
2. no
3. no
4. yes
5. yes

Exercise $4.7E.5$

Determine whether each ordered pair is a solution to the inequality $x+y>4$:

1. $(5,1)$
2. $(-2,6)$
3. $(3,2)$
4. $(10,-5)$
5. $(0,0)$

Exercise $4.7E.6$

Determine whether each ordered pair is a solution to the inequality $x+y>2$:

1. $(1,1)$
2. $(4,-3)$
3. $(0,0)$
4. $(-8,12)$
5. $(3,0)$

Answer


1. no
2. no
3. no
4. yes
5. yes

Recognize the Relation Between the Solutions of an Inequality and its Graph

In the following exercises, write the inequality shown by the shaded region.

Exercise $4.7E.7$


Write the inequality shown by the graph with the boundary line $y=3x-4$.



https://math.libretexts.org/@go/page/30516

Exercise 4.7E.8

Write the inequality shown by the graph with the boundary line $y = 2x - 4$.




Answer

$y < 2x - 4$


Exercise 4.7E.9

Write the inequality shown by the graph with the boundary line $y = \frac{1}{2}x + 1$



Exercise 4.7E.10

Write the inequality shown by the graph with the boundary line $y = -\frac{1}{3}x - 2$




Answer

$y \leq -\frac{1}{3}x - 2$

Exercise 4.7E.11


Write the inequality shown by the shaded region in the graph with the boundary line $x + y = 5$.

https://math.libretexts.org/@go/page/30516



Exercise 4.7E.12

Write the inequality shown by the shaded region in the graph with the boundary line $x + y = 3$.




Answer

$x + y \geq 3$


Exercise 4.7E.13

Write the inequality shown by the shaded region in the graph with the boundary line $2x + y = -4$.



Exercise 4.7E.14

Write the inequality shown by the shaded region in the graph with the boundary line $x + 2y = -2$.




https://math.libretexts.org/@go/page/30516

Answer

$x + 2y \geq -2$


Exercise 4.7E.15

Write the inequality shown by the shaded region in the graph with the boundary line $3x - y = 6$.



Exercise 4.7E.16

Write the inequality shown by the shaded region in the graph with the boundary line $2x - y = 4$.




Answer

$2x - y < 4$

Exercise 4.7E.17


Write the inequality shown by the shaded region in the graph with the boundary line $2x - 5y = 10$.



Exercise 4.7E.18

Write the inequality shown by the shaded region in the graph with the boundary line $4x - 3y = 12$.

https://math.libretexts.org/@go/page/30516



Answer

$4x - 3y > 12$

Graph Linear Inequalities

In the following exercises, graph each linear inequality.


Exercise 4.7E.19

Graph the linear inequality $y > \frac{2}{3}x - 1$

Exercise 4.7E.20

Graph the linear inequality $y < \frac{3}{5}x + 2$

Answer




Exercise 4.7E.21

Graph the linear inequality $y \leq -\frac{1}{2}x + 4$

Exercise 4.7E.22

Graph the linear inequality $y \geq -\frac{1}{3}x - 2$

Answer



https://math.libretexts.org/@go/page/30516


Exercise 4.7E.23

Graph the linear inequality $x - y \leq 3$

Exercise 4.7E.24

Graph the linear inequality $x - y \geq -2$

Answer




Exercise 4.7E.25

Graph the linear inequality $4x + y > -4$

Exercise 4.7E.26

Graph the linear inequality $x + 5y < -5$

Answer




Exercise 4.7E.27

Graph the linear inequality $3x + 2y \geq -6$

Exercise 4.7E.28

Graph the linear inequality $4x + 2y \geq -8$

Answer




Exercise $4.7E.29$

Graph the linear inequality $y > 4x$

Exercise $4.7E.30$

Graph the linear inequality $y > x$

Answer




Exercise $4.7E.31$

Graph the linear inequality $y \leq -x$

Exercise $4.7E.32$

Graph the linear inequality $y \leq -3x$

Answer



Exercise $4.7E.33$


Graph the linear inequality $y \geq -2$

https://math.libretexts.org/@go/page/30516

Exercise 4.7E.34

Graph the linear inequality $y < -1$

Answer




Exercise 4.7E.35

Graph the linear inequality $y < 4$

Exercise 4.7E.36

Graph the linear inequality $y \geq 2$

Answer




Exercise 4.7E.37

Graph the linear inequality $x \leq 5$

Exercise 4.7E.38

Graph the linear inequality $x > -2$

Answer



https://math.libretexts.org/@go/page/30516


Exercise 4.7E. 39

Graph the linear inequality $x > -3$

Exercise 4.7E. 40

Graph the linear inequality $x \leq 4$

Answer




Exercise 4.7E. 41

Graph the linear inequality $x - y < 4$

Exercise 4.7E. 42

Graph the linear inequality $x - y < -3$

Answer




Exercise 4.7E. 43

Graph the linear inequality $y \geq \frac{3}{2}x$

Exercise 4.7E. 44

Graph the linear inequality $y \leq \frac{5}{4}x$

Answer




Exercise 4.7E.45

Graph the linear inequality $y > -2x + 1$

Exercise 4.7E.46

Graph the linear inequality $y < -3x - 4$

Answer




Exercise 4.7E.47

Graph the linear inequality $x \leq -1$

Exercise 4.7E.48

Graph the linear inequality $x \geq 0$

Answer



Everyday Math

https://math.libretexts.org/@go/page/30516

Exercise $4.7E.49$

Money. Gerry wants to have a maximum of \$100 cash at the ticket booth when his church carnival opens. He will have \$1 bills and \$5 bills. If x is the number of \$1 bills and y is the number of \$5 bills, the inequality $x+5y \leq 100$ models the situation.

1. Graph the inequality.
2. List three solutions to the inequality $x+5y \leq 100$ where both x and y are integers.


Exercise $4.7E.50$

Shopping. Tula has \$20 to spend at the used book sale. Hardcover books cost \$2 each and paperback books cost \$0.50 each. If x is the number of hardcover books Tula can buy and y is the number of paperback books she can buy, the inequality $2x+\frac{1}{2}y \leq 20$ models the situation.

1. Graph the inequality.
2. List three solutions to the inequality $2x+\frac{1}{2}y \leq 20$ where both x and y are whole numbers.

Answer

1.



2. Answers will vary.

Writing Exercises

Exercise $4.7E.51$

Lester thinks that the solution of any inequality with a $>$ sign is the region above the line and the solution of any inequality with a $<$ sign is the region below the line. Is Lester correct? Explain why or why not.

Exercise $4.7E.52$

Explain why in some graphs of linear inequalities the boundary line is solid but in other graphs it is dashed.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
verify solutions to an inequality in two variables.			
recognize the relation between the solutions of an inequality and its graph.			
graph linear inequalities.			

b. What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 4.7E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **4.7E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 4 Review Exercises

Chapter 4 Review Exercises

Rectangular Coordinate System

Plot Points in a Rectangular Coordinate System

In the following exercises, plot each point in a rectangular coordinate system.


? Exercise 1

1. $(-1,-5)$
2. $(-3,4)$
3. $(2,-3)$
4. $\left(1,\frac{5}{2}\right)$

? Exercise 2

1. $(4,3)$
2. $(-4,3)$
3. $(-4,-3)$
4. $(4,-3)$

Answer




? Exercise 3

1. $(-2,0)$
2. $(0,-4)$
3. $(0,5)$
4. $(3,0)$

? Exercise 4

1. $\left(2,\frac{3}{2}\right)$
2. $\left(3,\frac{4}{3}\right)$
3. $\left(\frac{1}{3},-4\right)$
4. $\left(\frac{1}{2},-5\right)$


Answer




Identify Points on a Graph

In the following exercises, name the ordered pair of each point shown in the rectangular coordinate system.

Exercise 5



Exercise 6



Answer

a. (2,0)

b (0,−5)

c (−4.0)

d (0,3)

Verify Solutions to an Equation in Two Variables

In the following exercises, which ordered pairs are solutions to the given equations?

Exercise 7

$5x + y = 10$

1. (5,1)
2. (2,0)
3. (4,−10)

Exercise 8

$y = 6x - 2$

1. (1,4)
2. $\left(\frac{1}{3}, 0\right)$
3. (6,−2)

Answer

1, 2

Complete a Table of Solutions to a Linear Equation in Two Variables

In the following exercises, complete the table to find solutions to each linear equation.

Exercise 9

$y = 4x - 1$

x	y	(x,y)
0		
1		
-2		

Exercise 10

$y = -\frac{1}{2}x + 3$

x	y	(x,y)
0		
4		
-2		

Answer

x	y	(x,y)
0	3	(0,3)
4	1	(4, 1)
−2	4	(−2,4)

Exercise 11

$x + 2y = 5$

x	**y**	**(x,y)**
	0	
1		
-1		

Exercise 12

$3x + 2y = 6$

x	y	(x,y)
0		
	0	
-2		

Answer

x	y	(x,y)
0	−3	(0,−3)
2	0	(2,0)
−2	−6	(−2,−6)

Find Solutions to a Linear Equation in Two Variables

In the following exercises, find three solutions to each linear equation.

Exercise 13

$x + y = 3$

Exercise 14

$x + y = -4$

Answer

Answers will vary.

Exercise 15

$y = 3x + 1$

Exercise 16

$y = -x - 1$

Answer

Answers will vary.

Graphing Linear Equations

Recognize the Relation Between the Solutions of an Equation and its Graph

In the following exercises, for each ordered pair, decide:


1. Is the ordered pair a solution to the equation?
2. Is the point on the line?

Exercise 17

$y = -x + 4$

(0,4) (−1,3)

(2,2) (−2,6)




Exercise 18

$y = \frac{2}{3}x - 1$

$(0, -1)(3, 1)$

$(-3, -3)(6, 4)$



Answer

1. yes; yes
2. yes; no

Graph a Linear Equation by Plotting Points

In the following exercises, graph by plotting points.


Exercise 19

$y = 4x - 3$

Exercise 20

$y = -3x$

Answer




Exercise 21

$y = \frac{1}{2}x + 3$

Exercise 22

$x - y = 6$

Answer



Exercise 23


$2x + y = 7$

Exercise 24

$3x - 2y = 6$

Answer

https://math.libretexts.org/@go/page/30517



Graph Vertical and Horizontal lines

In the following exercises, graph each equation.


Exercise 25

$y = -2$

Exercise 26

$x = 3$

Answer



In the following exercises, graph each pair of equations in the same rectangular coordinate system.


Exercise 27

$y = -2x$ and $y = -2$

Exercise 28

$y = \frac{4}{3}x$ and $y = \frac{4}{3}$


Answer




Identify the x- and y-Intercepts on a Graph

In the following exercises, find the x- and y-intercepts.

Exercise 29



Exercise 30



Answer

$(3, 0)$ and $(0, 3)$

Find the$x$- and y-Intercepts from an Equation of a Line

In the following exercises, find the intercepts of each equation.

Exercise 31

$x + y = 5$

Exercise 32

$x - y = -1$

Answer

$(-1, 0), (0, 1)$

Exercise 33

$x+2y=6$

Exercise 34

$2x+3y=12$

Answer

$(6,0),(0,4)$

Exercise 35

$y=\frac{3}{4}x-12$

Exercise 36

$y=3x$

Answer

$(0,0)$

Graph a Line Using the Intercepts

In the following exercises, graph using the intercepts.


Exercise 37

$-x+3y=3$

Exercise 38

$x+y=-2$

Answer




Exercise 39

$x-y=4$

Exercise 40

$2x-y=5$

Answer




Exercise 41

$2x - 4y = 8$

Exercise 42

$y = 2x$

Answer




Slope of a Line


Use Geoboards to Model Slope

In the following exercises, find the slope modeled on each geoboard.

Exercise 43




Exercise 44




Answer

$\frac{4}{3}$

Exercise 45



Exercise 46



Answer

$-\frac{2}{3}$

Exercise 47


$\frac{1}{3}$

Exercise 48

$\frac{3}{2}$

Answer

https://math.libretexts.org/@go/page/30517




Exercise 49

$-\frac{2}{3}$

Exercise 50

$-\frac{1}{2}$


Answer




Use $m = \frac{\text{rise}}{\text{run}}$ to find the Slope of a Line from its Graph

In the following exercises, find the slope of each line shown.

Exercise 51




Exercise 52



Answer


1

Exercise 53



Exercise 54

https://math.libretexts.org/@go/page/30517



Answer

$-\frac{1}{2}$

Find the Slope of Horizontal and Vertical Lines

In the following exercises, find the slope of each line.

? Exercise 55

$y = 2$

? Exercise 56

$x = 5$

Answer

undefined

? Exercise 57

$x = -3$

? Exercise 58

$y = -1$

Answer

0

Use the Slope Formula to find the Slope of a Line between Two Points

In the following exercises, use the slope formula to find the slope of the line between each pair of points.

Exercise 59

$(-1,-1),(0,5)$

Exercise 60

$(3,5),(4,-1)$

Answer

-6

Exercise 61

$(-5,-2),(3,2)$

Exercise 62

$(2,1),(4,6)$

Answer

$\frac{5}{2}$

Graph a Line Given a Point and the Slope

In the following exercises, graph each line with the given point and slope.


Exercise 63

$(2,-2);\quad m=\frac{5}{2}$

Exercise 64

$(-3,4);\quad m=-\frac{1}{3}$

Answer



Exercise 65


x -intercept $-4;\quad m=3$

https://math.libretexts.org/@go/page/30517

Exercise 66

y -intercept 1; $m = -\frac{3}{4}$

Answer




Solve Slope Applications

In the following exercises, solve these slope applications.

Exercise 67

The roof pictured below has a rise of 10 feet and a run of 15 feet. What is its slope?



Exercise 68

A mountain road rises 50 feet for a 500-foot run. What is its slope?

Answer


$\frac{1}{10}$

Intercept Form of an Equation of a Line

Recognize the Relation Between the Graph and the Slope–Intercept Form of an Equation of a Line


In the following exercises, use the graph to find the slope and y-intercept of each line. Compare the values to the equation $y = mx + b$.

Exercise 69



$y = 4x - 1$

Exercise 70



$y = -\frac{2}{3}x + 4$

Answer

slope $m = -\frac{2}{3}$ and y-intercept $(0, 4)$

Identify the Slope and y-Intercept from an Equation of a Line

In the following exercises, identify the slope and y-intercept of each line.

Exercise 71

$y = -4x + 9$

Exercise 72

$y = \frac{5}{3}x - 6$

Answer

$\frac{5}{3}; (0, -6)$

Exercise 73

$5x + y = 10$

Exercise 74

$4x - 5y = 8$

Answer

$\frac{4}{5}; \quad \left(0, -\frac{8}{5}\right)$

Graph a Line Using Its Slope and Intercept

In the following exercises, graph the line of each equation using its slope and y-intercept.


Exercise 75

$y = 2x + 3$

Exercise 76

$y = -x - 1$

Answer




Exercise 77

$y = -\frac{2}{5}x + 3$

Exercise 78

$4x - 3y = 12$

Answer



In the following exercises, determine the most convenient method to graph each line.

Exercise 79

$x = 5$

Exercise 80

$y = -3$

Answer

horizontal line

Exercise 81

$2x + y = 5$

Exercise 82

$x - y = 2$

Answer

intercepts

Exercise 83

$y = x + 2$

Exercise 84

$y = \frac{3}{4}x - 1$

Answer

plotting points

Graph and Interpret Applications of Slope–Intercept

Exercise 85

Katherine is a private chef. The equation $C = 6.5m + 42$ models the relation between her weekly cost, C, in dollars and the number of meals, m, that she serves.

1. Find Katherine's cost for a week when she serves no meals.
2. Find the cost for a week when she serves 14 meals.
3. Interpret the slope and C-intercept of the equation.
4. Graph the equation.


Exercise 86

Marjorie teaches piano. The equation $P = 35h - 250$ models the relation between her weekly profit, P, in dollars and the number of student lessons, s, that she teaches.

1. Find Marjorie's profit for a week when she teaches no student lessons.
2. Find the profit for a week when she teaches 20 student lessons.

3. Interpret the slope and P-intercept of the equation.
4. Graph the equation.

Answer

1. $-\$250$
2. $\$450$
3. The slope, 35, means that Marjorie's weekly profit, P, increases by $\$35$ for each additional student lesson she teaches. The P-intercept means that when the number of lessons is 0, Marjorie loses $\$250$
4. 

Use Slopes to Identify Parallel Lines

In the following exercises, use slopes and y-intercepts to determine if the lines are parallel.

? Exercise 87

$4x-3y=-1; \quad y=\frac{4}{3}x-3$

? Exercise 88

$2x-y=8; \quad x-2y=4$

Answer

not parallel

Use Slopes to Identify Perpendicular Lines

In the following exercises, use slopes and y-intercepts to determine if the lines are perpendicular.

? Exercise 89

$y=5x-1; \quad 10x+2y=0$

? Exercise 90

$3x-2y=5; \quad 2x+3y=6$

Answer

perpendicular

Find an Equation of the Line Given the Slope and *y*-Intercept

In the following exercises, find the equation of a line with given slope and y-intercept. Write the equation in slope–intercept form.

Exercise 91

slope $\frac{1}{3}$ and y-intercept $(0, -6)$

Exercise 92

slope -5 and y-intercept $(0, -3)$

Answer

$y = -5x - 3$

Exercise 93

slope 0 and y-intercept $(0, 4)$

Exercise 94


slope -2 and y-intercept $(0, 0)$

Answer


$y = -2x$

In the following exercises, find the equation of the line shown in each graph. Write the equation in slope–intercept form.

Exercise 95




Exercise 96




Answer

$y = -3x + 5$

Exercise 97



Exercise 98



Answer

$y = -4$

Find an Equation of the Line Given the Slope and a Point

In the following exercises, find the equation of a line with given slope and containing the given point. Write the equation in slope–intercept form.

Exercise 99

$m = -\frac{1}{4}$, point $(-8, 3)$

Exercise 100

$m = \frac{3}{5}$, point $(10, 6)$

Answer

$y = \frac{3}{5}x$

Exercise 101

Horizontal line containing $(-2, 7)$

Exercise 102

$m = -2$, point $(-1, -3)$

Answer

$y = -2x - 5$

Find an Equation of the Line Given Two Points

In the following exercises, find the equation of a line containing the given points. Write the equation in slope–intercept form.

Exercise 103

$(2, 10)$ and $(-2, -2)$

Exercise 104

$(7, 1)$ and $(5, 0)$

Answer

$y = \frac{1}{2}x - \frac{5}{2}$

Exercise 105

$(3, 8)$ and $(3, -4)$

Exercise 106

$(5, 2)$ and $(-1, 2)$

Answer

$y = 2$

Find an Equation of a Line Parallel to a Given Line

In the following exercises, find an equation of a line parallel to the given line and contains the given point. Write the equation in slope–intercept form.

? Exercise 107

line $y = -3x + 6$, point $(1, -5)$

? Exercise 108

line $2x + 5y = -10$, point $(10, 4)$

Answer

$y = -\frac{2}{5}x + 8$

? Exercise 109

line $x = 4$, point $(-2, -1)$

? Exercise 110

line $y = -5$, point $(-4, 3)$

Answer

$y = 3$

Find an Equation of a Line Perpendicular to a Given Line

In the following exercises, find an equation of a line perpendicular to the given line and contains the given point. Write the equation in slope–intercept form.

? Exercise 111

line $y = -\frac{4}{5}x + 2$, point $(8, 9)$

? Exercise 112

line $2x - 3y = 9$, point $(-4, 0)$

Answer

$y = -\frac{3}{2}x - 6$

? Exercise 113

line $y = 3$, point $(-1, -3)$

? Exercise 114

line $x = -5$ point $(2, 1)$

Answer

$y = 1$

Graph Linear Inequalities

Verify Solutions to an Inequality in Two Variables

In the following exercises, determine whether each ordered pair is a solution to the given inequality.

Exercise 115

Determine whether each ordered pair is a solution to the inequality $y < x - 3$:

1. $(0, 1)$
2. $(-2, -4)$
3. $(5, 2)$
4. $(3, -1)$
5. $(-1, -5)$

Exercise 116

Determine whether each ordered pair is a solution to the inequality $x + y > 4$:

1. $(6, 1)$
2. $(-3, 6)$
3. $(3, 2)$
4. $(-5, 10)$
5. $(0, 0)$

Answer


1. yes
2. no
3. yes
4. yes
5. no

Recognize the Relation Between the Solutions of an Inequality and its Graph

In the following exercises, write the inequality shown by the shaded region.


Exercise 117

Write the inequality shown by the graph with the boundary line $y = -x + 2$.



Exercise 118

Write the inequality shown by the graph with the boundary line $y = \frac{2}{3}x - 3$




Answer

$y > \frac{2}{3}x - 3$


Exercise 119

Write the inequality shown by the shaded region in the graph with the boundary line $x + y = -4$.



Exercise 120

Write the inequality shown by the shaded region in the graph with the boundary line $x - 2y = 6$.



Answer

$x - 2y \geq 6$

Graph Linear Inequalities

In the following exercises, graph each linear inequality.


Exercise 121

Graph the linear inequality $y > \frac{2}{5}x - 4$

Exercise 122

Graph the linear inequality $y \leq -\frac{1}{4}x + 3$

Answer




Exercise 123

Graph the linear inequality $x - y \leq 5$

Exercise 124

Graph the linear inequality $3x + 2y > 10$

Answer




Exercise 125

Graph the linear inequality $y \leq -3x$

Exercise 126

Graph the linear inequality $y < 6$

Answer



https://math.libretexts.org/@go/page/30517

Practice Test

Exercise 1

Plot each point in a rectangular coordinate system.

1. $(2, 5)$
2. $(-1, -3)$
3. $(0, 2)$
4. $\left(-4, \frac{3}{2}\right)$
5. $(5, 0)$

Exercise 2

Which of the given ordered pairs are solutions to the equation $3x - y = 6$?

1. $(3, 3)$
2. $(2, 0)$
3. $(4, -6)$

Answer

1. yes
2. yes
3. no

Exercise 3

Find three solutions to the linear equation $y = -2x - 4$

Exercise 4


Find the x- and y-intercepts of the equation $4x - 3y = 12$

Answer

$(3, 0), (0, -4)$


Find the slope of each line shown.

Exercise 5



https://math.libretexts.org/@go/page/30517


Exercise 6



Answer

undefined

Exercise 7



Exercise 8

Find the slope of the line between the points $(5, 2)$ and $(-1, -4)$

Answer

1

Exercise 9


Graph the line with slope $\frac{1}{2}$ containing the point $(-3, -4)$

Graph the line for each of the following equations.

Exercise 10

$y = \frac{5}{3}x - 1$

Answer




Exercise 11

$y = -x$

Exercise 12

$x - y = 2$

Answer




Exercise 13

$4x + 2y = -8$

Exercise 14

$y = 2$

Answer



Exercise 15

$x = -3$

Find the equation of each line. Write the equation in slope–intercept form.

Exercise 16

slope $-\frac{3}{4}$ and y-intercept $(0, -2)$

Answer

$y = -\frac{3}{4}x - 2$

Exercise 17

$m = 2$, point $(-3, -1)$

Exercise 18

containing $(10, 1)$ and $(6, -1)$

Answer

$y = \frac{1}{2}x - 4$

Exercise 19

parallel to the line $y = -\frac{2}{3}x - 1$, containing the point $(-3, 8)$

Exercise 20


perpendicular to the line $y = \frac{5}{4}x + 2$, containing the point $(-10, 3)$

Answer

$y = -\frac{4}{5}x - 5$

Exercise 21

Write the inequality shown by the graph with the boundary line $y = -x - 3$.




Graph each linear inequality.

https://math.libretexts.org/@go/page/30517

Exercise 22

$y > \frac{3}{2}x + 5$

Answer




Exercise 23

$x - y \geq -4$

Exercise 24

$y \leq -5x$

Answer



Exercise 1

$y < 3$

This page titled Chapter 4 Review Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **Chapter 4 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

CHAPTER OVERVIEW

5: Systems of Linear Equations

- 5.1: Solve Systems of Equations by Graphing
  - 5.1E: Exercises
- 5.2: Solve Systems of Equations by Substitution
  - 5.2E: Exercises
- 5.3: Solve Systems of Equations by Elimination
  - 5.3E: Exercises
- 5.4: Solve Applications with Systems of Equations
  - 5.4E: Exercises
- 5.5: Solve Mixture Applications with Systems of Equations
  - 5.5E: Exercises
- 5.6: Graphing Systems of Linear Inequalities
  - 5.6E: Exercises
- Chapter 5 Review Exercises

This page titled 5: Systems of Linear Equations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

5.1: Solve Systems of Equations by Graphing

Learning Objectives

By the end of this section, you will be able to:

- Determine whether an ordered pair is a solution of a system of equations
- Solve a system of linear equations by graphing
- Determine the number of solutions of linear system
- Solve applications of systems of equations by graphing

Note

Before you get started, take this readiness quiz.

1. For the equation $y = \frac{2}{3}x - 4$
   ⓐ is (6,0) a solution? ⓑ is (−3,−2) a solution?
   If you missed this problem, review Example 2.1.1.
2. Find the slope and y-intercept of the line 3x−y=12.
   If you missed this problem, review Example 4.5.7.
3. Find the x- and y-intercepts of the line 2x−3y=12.
   If you missed this problem, review Example 4.3.7.

Determine Whether an Ordered Pair is a Solution of a System of Equations

In the section on Solving Linear Equations and Inequalities we learned how to solve linear equations with one variable. Remember that the solution of an equation is a value of the variable that makes a true statement when substituted into the equation. Now we will work with **systems of linear equations**, two or more linear equations grouped together.

Definition: SYSTEM OF LINEAR EQUATIONS

When two or more linear equations are grouped together, they form a system of linear equations.

We will focus our work here on systems of two linear equations in two unknowns. Later, you may solve larger systems of equations.

An example of a system of two linear equations is shown below. We use a brace to show the two equations are grouped together to form a system of equations.

$$\begin{cases} 2x + y = 7 \\ x - 2y = 6 \end{cases} \tag{5.1.1}$$

A linear equation in two variables, like $2x + y = 7$, has an infinite number of solutions. Its graph is a line. Remember, every point on the line is a solution to the equation and every solution to the equation is a point on the line.

To solve a system of two linear equations, we want to find the values of the variables that are solutions to both equations. In other words, we are looking for the ordered pairs (x, y) that make both equations true. These are called the **solutions to a system of equations**.

Definition: SolutionS OF A SYSTEM OF EQUATIONS

Solutions of a system of equations are the values of the variables that make all the equations true. A solution of a system of two linear equations is represented by an ordered pair (x, y).

To determine if an ordered pair is a solution to a system of two equations, we substitute the values of the variables into each equation. If the ordered pair makes both equations true, it is a solution to the system.

Let's consider the system below:

$$\begin{cases} 3x-y=7 \\ x-2y=4 \end{cases} \tag{5.1.2}$$

Is the ordered pair (2,−1) a solution?

We substitute $x=2$ and $y=-1$ into both equations.

$$\begin{aligned} 3x-y &= 7 & \qquad x-2y &= 4 \\ 3(2)-(-1) &\stackrel{?}{=} 7 & \qquad 2-2(-1) &\stackrel{?}{=} 4 \\ 7 &= 7 \text{ true} & \qquad 4 &= 4 \text{ true} \end{aligned}$$

The ordered pair (2, −1) made both equations true. Therefore (2, −1) is a solution to this system.

Let's try another ordered pair. Is the ordered pair (3, 2) a solution?

We substitute $x=3$ and $y=2$ into both equations.

$$\begin{aligned} 3x-y &= 7 & \qquad x-2y &= 4 \\ 3(3)-2 &\stackrel{?}{=} 7 & \qquad 2-2(2) &\stackrel{?}{=} 4 \\ 7 &= 7 \text{ true} & \qquad -2 &= 4 \text{ false} \end{aligned}$$

The ordered pair (3, 2) made one equation true, but it made the other equation false. Since it is not a solution to **both** equations, it is not a solution to this system.

Example 5.1.1

Determine whether the ordered pair is a solution to the system: $\begin{cases} x-y=-1 \\ 2x-y=-5 \end{cases}$

1. (−2,−1)
2. (−4,−3)

Solution

1.

$$\begin{cases} x-y=-1 \\ 2x-y=-5 \end{cases}$$

We substitute $x=-2$ and $y=-1$ into both equations

$$\begin{aligned} x-y &= 1 & \qquad 2x-y &= -5 \\ -2-(-1) &= -1 & \qquad 2(-2)-(-1) &\stackrel{?}{=} -5 \\ -1 &= -1 \checkmark & \qquad -3 &= -5 \times \end{aligned}$$

(−2, −1) does not make both equations true. (−2, −1) is not a solution.

2.

We substitute $x=-4$ and $y=-3$ into both equations.

$$\begin{aligned} x-y &= -1 & \qquad 2x-y &= -5 \\ -4-(-3) &\stackrel{?}{=} -1 & \qquad 2(-4)-(-3) &\stackrel{?}{=} -5 \\ -1 &= -1 \checkmark & \qquad -5 &= -5 \checkmark \end{aligned}$$

(−4, −3) makes both equations true. (−4, −3) is a solution.

Try It 5.1.2

Determine whether the ordered pair is a solution to the system: $\begin{cases} 3x+y=0 \\ x+2y=-5 \end{cases}$

1. (1,−3)

2. (0,0)

Answer

1. yes
2. no

? Try It 5.1.3

Determine whether the ordered pair is a solution to the system: $\begin{cases} x-3y=-8 \\ -3x-y=4 \end{cases}$

1. (2,−2)
2. (−2,2)


Answer

1. no
2. yes


Solve a System of Linear Equations by Graphing

In this chapter we will use three methods to solve a system of linear equations. The first method we'll use is graphing. The graph of a linear equation is a line. Each point on the line is a solution to the equation. For a system of two equations, we will graph two lines. Then we can see all the points that are solutions to each equation. And, by finding what the lines have in common, we'll find the solution to the system.


Most linear equations in one variable have one solution, but we saw that some equations, called **contradictions**, have no solutions and for other equations, called **identities**, all numbers are solutions. Similarly, when we solve a system of two linear equations represented by a graph of two lines in the same plane, there are three possible cases, as shown in Figure 5.1.1:



The lines intersect.
Intersecting lines have one point in common.
There is one solution to this system.



The lines are parallel.
Parallel lines have no points in common.
There is no solution to this system.



Both equations give the same line.
Because we have just one line, there are infinitely many solutions.



Figure 5.1.1

For the first example of solving a system of linear equations in this section and in the next two sections, we will solve the same system of two linear equations. But we'll use a different method in each section. After seeing the third method, you'll decide which method was the most convenient way to solve this system.

✔ Example 5.1.4: How to Solve a System of Linear Equations by Graphing

Solve the system by graphing: $\begin{cases} 2x+y=7 \\ x-2y=6 \end{cases}$

Solution

Step 1. Graph the first equation.	To graph the first line, write the equation in slope–intercept form. $2x + y = 7$ $y = -2x + 7$ $m = -2$ $b = 7$	$\begin{cases} 2x + y = 7 \\ x - 2y = 6 \end{cases}$ 
Step 2. Graph the second equation on the same rectangular coordinate system.	To graph the second line, use intercepts. $x - 2y = 6$ $(0, -3)$ $(6, 0)$	
Step 3. Determine whether the lines intersect, are parallel, or are the same line.	Look at the graph of the lines.	The lines intersect.
Step 4. Identify the solution to the system. If the lines intersect, identify the point of intersection. Check to make sure it is a solution to both equations. This is the solution to the system. If the lines are parallel, the system has no solution If the lines are the same, the system has an infinite number of solutions.	Since the lines intersect, find the point of intersection. Check the point in both equations.	The lines intersect at $(4, -1)$. $2x + y = 7$ $2(4) + (-1) \stackrel{?}{=} 7$ $8 - 1 \stackrel{?}{=} 7$ $7 = 7$ ✓ $x - 2y = 6$ $4 - 2(-1) \stackrel{?}{=} 6$ $6 = 6$ ✓ The solution is $(4, -1)$.

? Try It 5.1.5

Solve each system by graphing: $\begin{cases} x - 3y = -3 \\ x + y = 5 \end{cases}$

Answer

(3,2)

Try It 5.1.6

Solve each system by graphing: $\begin{cases} -x+y=1 \\ 3x+2y=12 \end{cases}$

Answer

(2,3)

The steps to use to solve a system of linear equations by graphing are shown below.

TO SOLVE A SYSTEM OF LINEAR EQUATIONS BY GRAPHING.

1. Graph the first equation.
2. Graph the second equation on the same rectangular coordinate system.
3. Determine whether the lines intersect, are parallel, or are the same line.
4. Identify the solution to the system.
   - If the lines intersect, identify the point of intersection. Check to make sure it is a solution to both equations. This is the solution to the system.
   - If the lines are parallel, the system has no solution.
   - If the lines are the same, the system has an infinite number of solutions.

Example 5.1.7


Solve the system by graphing: $\begin{cases} y=2x+1 \\ y=4x-1 \end{cases}$

Solution

Both of the equations in this system are in slope-intercept form, so we will use their slopes and y-intercepts to graph them.

$\begin{cases} y=2x+1 \\ y=4x-1 \end{cases}$

Find the slope and y-intercept of the first equation.	$y=2x+1$ $m=2$ $b=1$
Find the slope and y-intercept of the first equation.	$y=4x-1$ $m=4$ $b=-1$
Graph the two lines.	
Determine the point of intersection.	The lines intersect at (1, 3).

	
Check the solution in both equations.	$y = 2x + 1$ $\quad$ $y = 4x - 1$ $3 \stackrel{?}{=} 2 \cdot 1 + 1$ $\quad$ $3 \stackrel{?}{=} 4 \cdot 1 - 1$ $3 = 3\checkmark$ $\quad$ $3 = 3\checkmark$
	The solution is (1, 3).

? Try It 5.1.8

Solve the system by graphing: $\begin{cases} y = 2x + 2 \\ y = -x - 4 \end{cases}$

Answer

(−2,−2)

? Try It 5.1.9

Solve the system by graphing: $\begin{cases} y = 3x + 3 \\ y = -x + 7 \end{cases}$

Answer

(1,6)


Both equations in Example 5.1.7 were given in slope–intercept form. This made it easy for us to quickly graph the lines. In the next example, we'll first re-write the equations into slope–intercept form.

✓ Example 5.1.10

Solve the system by graphing: $\begin{cases} 3x + y = -1 \\ 2x + y = 0 \end{cases}$

Solution

We'll solve both of these equations for y so that we can easily graph them using their slopes and y-intercepts. $\begin{cases} 3x + y = -1 \\ 2x + y = 0 \end{cases}$

Solve the first equation for y.	
Find the slope and y-intercept.	$3x+y=-1$ $y=-3x-1$ $m=-3$ $b=-1$
Solve the second equation for y.	$2x+y=0$ $y=-2x$
Find the slope and y-intercept.	$b=0$
Graph the lines.	
Determine the point of intersection.	The lines intersect at $(-1, 2)$.
Check the solution in both equations.	$3x+y=-1$ $\quad 3(-1)+2 \stackrel{?}{=} -1 \quad -1=-1\checkmark$ $2x+y=0$ $\quad 2(-1)+2 \stackrel{?}{=} 0 \quad 0=0\checkmark$
	The solution is $(-1, 2)$.

? Try It 5.1.11

Solve each system by graphing: $\begin{cases} -x+y=1 \\ 2x+y=10 \end{cases}$

Answer

(3,4)

? Try It 5.1.12

Solve each system by graphing: $\begin{cases} 2x+y=6 \\ x+y=1 \end{cases}$

Answer

(5,−4)


Usually when equations are given in standard form, the most convenient way to graph them is by using the intercepts. We'll do this in Example 5.1.13.

Example 5.1.13

Solve the system by graphing: $\begin{cases} x+y=2 \\ x-y=4 \end{cases}$

Solution

We will find the *x*- and *y*-intercepts of both equations and use them to graph the lines.

	$x+y=2$	
To find the intercepts, let $x = 0$ and solve for y, then let $y = 0$ and solve for x.	$x+y=2 \quad x+y=2$ $0+y=2 \quad x+0=2$ $y=2 \quad x=2$	x | y 0 | 2 2 | 0
	$x-y=4$	
To find the intercepts, let $x = 0$ then let $y = 0$.	$x-y=4 \quad x-y=4$ $0-y=4 \quad x-0=4$ $-y=4 \quad x=4$ $y=-4$	x | y 0 | –4 4 | 0
Graph the line.		
Determine the point of intersection.	The lines intersect at (3, −1).	
Check the solution in both equations.	$x+y=2 \quad x-y=4$ $3+(-1) \stackrel{?}{=} 2 \quad 3-(-1) \stackrel{?}{=} 4$ $2=2\checkmark \quad 4=4\checkmark$ The solution is (3, −1).	

(5.1.3)

Try It 5.1.14

Solve each system by graphing: $\begin{cases} x+y=6 \\ x-y=2 \end{cases}$

Answer

(4,2)

Try It 5.1.15

Solve each system by graphing: $\begin{cases} x+y=2 \\ x-y=8 \end{cases}$

Answer


(5,−3)

Do you remember how to graph a linear equation with just one variable? It will be either a vertical or a horizontal line.

Example 5.1.16

Solve the system by graphing: $\begin{cases} y=6 \\ 2x+3y=12 \end{cases}$

Solution

	$y=6$ $2x+3y=12$
We know the first equation represents a horizontal line whose y-intercept is 6.	$y=6$
The second equation is most conveniently graphed using intercepts.	$2x+3y=12$
To find the intercepts, let $x=0$ and then $y=0$.	x | y 0 | 4 6 | 0
Graph the lines.	
Determine the point of intersection.	The lines intersect at (−3, 6).
Check the solution to both equations.	$y = 6$, $6 \stackrel{?}{=} 6$, $6 = 6\checkmark$ $2x+3y = 12$, $2(-3)+3(6) \stackrel{?}{=} 12$, $-6+18 \stackrel{?}{=} 12$, $12 = 12\checkmark$
	The solution is (−3, 6).

Try It 5.1.17

Solve each system by graphing: $\begin{cases} y = -1 \\ x + 3y = 6 \end{cases}$

Answer

$(9,-1)$

Try It 5.1.18

Solve each system by graphing: $\begin{cases} x = 4 \\ 3x - 2y = 24 \end{cases}$

Answer

$(4,-6)$


In all the systems of linear equations so far, the lines intersected and the solution was one point. In the next two examples, we'll look at a system of equations that has no solution and at a system of equations that has an infinite number of solutions.

Example 5.1.19

Solve the system by graphing: $\begin{cases} y = \frac{1}{2}x - 3 \\ x - 2y = 4 \end{cases}$

Solution

	$y = \frac{1}{2}x - 3$ $x - 2y = 4$
To graph the first equation, we will use its slope and y-intercept.	$y = \frac{1}{2}x - 3$
	$m = \frac{1}{2}$
	$b = -3$
To graph the second equation, we will use the intercepts.	$x - 2y = 4$
	x | y 0 | −2 4 | 0

Graph the lines.	
Determine the point of intersection.	The lines are parallel.
	Since no point is on both lines, there is no ordered pair that makes both equations true. There is no solution to this system.

Try It 5.1.20

Solve each system by graphing: $\begin{cases} y = -\frac{1}{4}x + 2 \\ x + 4y = -8 \end{cases}$

Answer

no solution

Try It 5.1.21

Solve each system by graphing: $\begin{cases} y = 3x - 1 \\ 6x - 2y = 6 \end{cases}$

Answer


no solution

Example 5.1.22

Solve the system by graphing: $\begin{cases} y = 2x - 3 \\ -6x + 3y = -9 \end{cases}$

Solution

	$y = 2x - 3$ $-6x + 3y = -9$
Find the slope and y-intercept of the first equation.	$y = 2x - 3$ $m = 2$ $b = -3$
Find the intercepts of the second equation.	$-6x + 3y = -9$

	$\begin{array}{\|c\|c\|}\hline x & y \\ \hline 0 & -3 \\ \hline \frac{3}{2} & 0 \\ \hline \end{array}$
Graph the lines.	
Determine the point of intersection.	The lines are the same!
	Since every point on the line makes both equations true, there are infinitely many ordered pairs that make both equations true.
	There are infinitely many solutions to this system.

Try It 5.1.23

Solve each system by graphing: $\begin{cases} y = -3x - 6 \\ 6x + 2y = -12 \end{cases}$

Answer

infinitely many solutions

Try It 5.1.24

Solve each system by graphing: $\begin{cases} y = \frac{1}{2}x - 4 \\ 2x - 4y = 16 \end{cases}$

Answer

infinitely many solutions

If you write the second equation in Example 5.1.22 in slope-intercept form, you may recognize that the equations have the same slope and same *y*-intercept.

When we graphed the second line in the last example, we drew it right over the first line. We say the two lines are coincident. Coincident lines have the same slope and same *y*-intercept.

COINCIDENT LINES

Coincident lines have the same slope and same *y*-intercept.

Determine the Number of Solutions of a Linear System

There will be times when we will want to know how many solutions there will be to a system of linear equations, but we might not actually have to find the solution. It will be helpful to determine this without graphing.

We have seen that two lines in the same plane must either intersect or are parallel. The systems of equations in Example 5.1.4 through Example 5.1.18 all had two intersecting lines. Each system had one solution.

A system with parallel lines, like Example 5.1.19, has no solution. What happened in Example 5.1.22? The equations have **coincident lines**, and so the system had infinitely many solutions.

We'll organize these results in Figure 5.1.2 below:

Graph	Number of solutions
2 intersecting lines	1
Parallel lines	None
Same line	Infinitely many

Figure 5.1.2

Parallel lines have the same slope but different *y*-intercepts. So, if we write both equations in a system of linear equations in slope–intercept form, we can see how many solutions there will be without graphing! Look at the system we solved in Example 5.1.19.

$$\begin{cases} y = \frac{1}{2}x - 3 \\ x - 2y = 4 \end{cases}$$

The first line is in slope–intercept form. If we solve the second equation for y, we get

$$y = \frac{1}{2}x - 3 \qquad x - 2y = 4$$
$$x - 2y = -x + 4$$
$$y = \frac{1}{2}x - 2$$
$$m = \frac{1}{2}, b = -3 \qquad m = \frac{1}{2}, b = -2$$

The two lines have the same slope but different *y*-intercepts. They are parallel lines.

Figure 5.1.3 shows how to determine the number of solutions of a linear system by looking at the slopes and intercepts.

Number of Solutions of a Linear System of Equations			
Slopes	**Intercepts**	**Type of Lines**	**Number of Solutions**
Different		Intersecting	1 point
Same	Different	Parallel	No solution
Same	Same	Coincident	Infinitely many solutions

Figure 5.1.3

Let's take one more look at our equations in Example 5.1.19 that gave us parallel lines.

$$\begin{cases} y = \frac{1}{2}x - 3 \\ x - 2y = 4 \end{cases} \tag{5.1.4}$$

When both lines were in slope-intercept form we had:

$$y = \frac{1}{2}x - 3 \quad y = \frac{1}{2}x - 2 \tag{5.1.5}$$

Do you recognize that it is impossible to have a single ordered pair (x,y) that is a solution to both of those equations?

We call a system of equations like this **an inconsistent system**. It has no solution.

A system of equations that has at least one solution is called a **consistent system**.

CONSISTENT AND INCONSISTENT SYSTEMS

A **consistent system** of equations is a system of equations with at least one solution.

An **inconsistent system** of equations is a system of equations with no solution.

We also categorize the equations in a system of equations by calling the equations *independent* or *dependent*. If two equations are **independent equations**, they each have their own set of solutions. Intersecting lines and parallel lines are independent.

If two equations are dependent, all the solutions of one equation are also solutions of the other equation. When we graph two **dependent equations**, we get coincident lines.

INDEPENDENT AND DEPENDENT EQUATIONS

Two equations are **independent** if they have different solutions.

Two equations are **dependent** if all the solutions of one equation are also solutions of the other equation.

Let's sum this up by looking at the graphs of the three types of systems. See Figure 5.1.4 and Figure 5.1.5.




Figure 5.1.4

Lines	Intersecting	Parallel	Coincident
Number of solutions	1 point	No solution	Infinitely many
Consistent/inconsistent	Consistent	Inconsistent	Consistent
Dependent/independent	Independent	Independent	Dependent

Figure 5.1.5

Example 5.1.25

Without graphing, determine the number of solutions and then classify the system of equations: $\begin{cases} y = 3x - 1 \\ 6x - 2y = 12 \end{cases}$

Solution

We will compare the slopes and intercepts of the two lines.

$$\begin{cases} y = 3x - 1 \\ 6x - 2y = 12 \end{cases}$$

The first equation is already in slope-intercept form.

$$y = 3x - 1$$

Write the second equation inslope–intercept form.

$$\begin{aligned} 6x - 2y &= 12 \\ -2y &= -6x - 12 \\ \tfrac{-2y}{-2} &= \tfrac{-6x+12}{-2} \\ y &= 3x - 6 \end{aligned}$$

Find the slope and intercept of each line.

$$\begin{aligned} y &= 3x - 1 & y &= 3x - 6 \\ m &= 3 & m &= 3 \\ b &= -1 & b &= -6 \end{aligned}$$

Since the slopes are the same and y-intercepts are different, the lines are parallel.

A system of equations whose graphs are parallel lines has no solution and is inconsistent and independent.

Try It 5.1.26

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} y = -2x - 4 \\ 4x + 2y = 9 \end{cases}$$

Answer

no solution, inconsistent, independent

Try It 5.1.27

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} y = \frac{1}{3}x - 5 \\ x - 3y = 6 \end{cases}$$

Answer

no solution, inconsistent, independent

Example 5.1.28

Without graphing, determine the number of solutions and then classify the system of equations: $\begin{cases} 2x + y = -3 \\ x - 5y = 5 \end{cases}$

Solution

We will compare the slopes and intercepts of the two lines.

$$\begin{cases} 2x + y = -3 \\ x - 5y = 5 \end{cases}$$

Write the second equation in slope–intercept form.

$$\begin{aligned} 2x + y &= -3 \\ y &= -2x - 3 \end{aligned} \qquad \begin{aligned} x - 5y &= 5 \\ -5y &= -x + 5 \\ \tfrac{-5y}{-5} &= \tfrac{-x+5}{-5} \\ y &= \tfrac{1}{5}x - 1 \end{aligned}$$

Find the slope and intercept of each line.

$$\begin{aligned} y &= -2x - 3 \\ m &= -2 \\ b &= -3 \end{aligned} \qquad \begin{aligned} y &= \tfrac{1}{5}x - 1 \\ m &= \tfrac{1}{5} \\ b &= -1 \end{aligned}$$

Since the slopes are different, the lines intersect.

A system of equations whose graphs are intersect has 1 solution and is consistent and independent.

Try It 5.1.29

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} 3x + 2y = 2 \\ 2x + y = 1 \end{cases}$$

Answer

one solution, consistent, independent

Try It 5.1.30

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} x+4y=12 \\ -x+y=3 \end{cases}$$

Answer

one solution, consistent, independent

Example 5.1.31

Without graphing, determine the number of solutions and then classify the system of equations. $\begin{cases} 3x-2y=4 \\ y=\frac{3}{2}x-2 \end{cases}$

Solution

We will compare the slopes and intercepts of the two lines. $\begin{cases} 3x-2y &= 4 \\ y &= \frac{3}{2}x-2 \end{cases}$

Write the second equation in slope–intercept form.

$$\begin{aligned} 3x-2y &= 4 \\ -2y &= -3x+4 \\ \frac{-2y}{-2} &= \frac{-3x+4}{-2} \\ y &= \frac{3}{2}x-2 \end{aligned}$$

Find the slope and intercept of each line. $y = \frac{3}{2}x-2$

Since the equations are the same, they have the same slope and same y-intercept and so the lines are coincident.

A system of equations whose graphs are coincident lines has infinitely many solutions and is consistent and dependent.

Try It 5.1.32

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} 4x-5y=20 \\ y=\frac{4}{5}x-4 \end{cases}$$

Answer

infinitely many solutions, consistent, dependent

Try It 5.1.33

Without graphing, determine the number of solutions and then classify the system of equations.

$$\begin{cases} -2x-4y=8 \\ y=-\frac{1}{2}x-2 \end{cases}$$

Answer

infinitely many solutions, consistent, dependent

Solve Applications of Systems of Equations by Graphing

We will use the same problem solving strategy we used in Math Models to set up and solve applications of systems of linear equations. We'll modify the strategy slightly here to make it appropriate for systems of equations.

USE A PROBLEM SOLVING STRATEGY FOR SYSTEMS OF LINEAR EQUATIONS.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose variables to represent those quantities.
4. **Translate** into a system of equations.
5. **Solve** the system of equations using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Step 5 is where we will use the method introduced in this section. We will graph the equations and find the solution.

Example 5.1.34

Sondra is making 10 quarts of punch from fruit juice and club soda. The number of quarts of fruit juice is 4 times the number of quarts of club soda. How many quarts of fruit juice and how many quarts of club soda does Sondra need?

Solution

Step 1. Read the problem.

Step 2. Identify what we are looking for.

We are looking for the number of quarts of fruit juice and the number of quarts of club soda that Sondra will need.

Step 3. Name what we are looking for. Choose variables to represent those quantities.

Let f= number of quarts of fruit juice.
c= number of quarts of club soda

Step 4. Translate into a system of equations.

The number of quarts of fruit juice and the number of quarts of club soda is 10

$$f + c = 10$$

The number of quarts of fruit juice is four times the number of quarts of club soda

$$f = 4c$$

We now have the system. $\begin{cases} f + c = 10 \\ f = 4c \end{cases}$


Step 5. Solve the system of equations using good algebra techniques.

$f = 4c$ $\quad$ $f + c = 10$

$m = 4$ $\quad$ $f = -c + 10$

$b = 0$ $\quad$ $m = -1$

$b = 10$



The point of intersection (2, 8) is the solution. This means Sondra needs 2 quarts of club soda and 8 quarts of fruit juice.

Step 6. Check the answer in the problem and make sure it makes sense.

Does this make sense in the problem?

Yes, the number of quarts of fruit juice, 8 is 4 times the number of quarts of club soda, 2.

Yes, 10 quarts of punch is 8 quarts of fruit juice plus 2 quarts of club soda.

Step 7. Answer the question with a complete sentence.

Sondra needs 8 quarts of fruit juice and 2 quarts of soda.

Try It 5.1.35

Manny is making 12 quarts of orange juice from concentrate and water. The number of quarts of water is 3 times the number of quarts of concentrate. How many quarts of concentrate and how many quarts of water does Manny need?

Answer

Manny needs 3 quarts juice concentrate and 9 quarts water.

Try It 5.1.36

Alisha is making an 18 ounce coffee beverage that is made from brewed coffee and milk. The number of ounces of brewed coffee is 5 times greater than the number of ounces of milk. How many ounces of coffee and how many ounces of milk does Alisha need?

Answer

Alisha needs 15 ounces of coffee and 3 ounces of milk.

Note

Access these online resources for additional instruction and practice with solving systems of equations by graphing.

- Instructional Video Solving Linear Systems by Graphing
- Instructional Video Solve by Graphing

Key Concepts

- **To solve a system of linear equations by graphing**
  1. Graph the first equation.
  2. Graph the second equation on the same rectangular coordinate system.
  3. Determine whether the lines intersect, are parallel, or are the same line.
  4. Identify the solution to the system.
     If the lines intersect, identify the point of intersection. Check to make sure it is a solution to both equations. This is the solution to the system.
     If the lines are parallel, the system has no solution.
     If the lines are the same, the system has an infinite number of solutions.
  5. Check the solution in both equations.
- Determine the number of solutions from the graph of a linear system

Graph	Number of solutions
2 intersecting lines	1
Parallel lines	None
Same line	Infinitely many

- Determine the number of solutions of a linear system by looking at the slopes and intercepts

Number of Solutions of a Linear System of Equations			
Slopes	**Intercepts**	**Type of Lines**	**Number of Solutions**
Different		Intersecting	1 point
Same	Different	Parallel	No solution
Same	Same	Coincident	Infinitely many solutions

- Determine the number of solutions and how to classify a system of equations

Lines	**Intersecting**	**Parallel**	**Coincident**
Number of solutions	1 point	No solution	Infinitely many
Consistent/inconsistent	Consistent	Inconsistent	Consistent
Dependent/independent	Independent	Independent	Dependent

- **Problem Solving Strategy for Systems of Linear Equations**
  1. **Read** the problem. Make sure all the words and ideas are understood.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose variables to represent those quantities.
  4. **Translate** into a system of equations.
  5. **Solve** the system of equations using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.

Glossary

coincident lines

Coincident lines are lines that have the same slope and same y-intercept.

consistent system

A consistent system of equations is a system of equations with at least one solution.

dependent equations

Two equations are dependent if all the solutions of one equation are also solutions of the other equation.

inconsistent system

An inconsistent system of equations is a system of equations with no solution.

independent equations

Two equations are independent if they have different solutions.

solutions of a system of equations

Solutions of a system of equations are the values of the variables that make all the equations true. A solution of a system of two linear equations is represented by an ordered pair (x, y).

system of linear equations

When two or more linear equations are grouped together, they form a system of linear equations.

This page titled 5.1: Solve Systems of Equations by Graphing is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.1: Solve Systems of Equations by Graphing** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.1E: Exercises

Practice Makes Perfect

Determine Whether an Ordered Pair is a Solution of a System of Equations. In the following exercises, determine if the following points are solutions to the given system of equations.

Exercise 5.1E.1

$$\begin{cases} 2x-6y=0 \\ 3x-4y=5 \end{cases}$$

1. (3,1)
2. (−3,4)

Answer

1. yes
2. no

Exercise 5.1E.2

$$\begin{cases} 7x-4y=-1 \\ -3x-2y=1 \end{cases}$$

1. $(-\frac{3}{13}, -\frac{2}{13})$
2. (1,-2)

Answer

1. yes
2. no

Exercise 5.1E.3

$$\begin{cases} 2x+y=5 \\ x+y=1 \end{cases}$$

1. (4,−3)
2. (2,0)

Answer

1. yes
2. no

Exercise 5.1E.4

$$\begin{cases} -3x+y=8 \\ -x+2y=-9 \end{cases}$$

1. (−5,−7)
2. (−5,7)

Exercise 5.1E.5

$$\begin{cases} x+y=2 \\ y=\frac{3}{4}x \end{cases}$$

1. $\left(\frac{8}{7},\frac{6}{7}\right)$
2. $\left(1,\frac{3}{4}\right)$

Answer

1. yes
2. no

Exercise 5.1E.6

$$\begin{cases} x+y=1 \\ y=\frac{2}{5}x \end{cases}$$

1. $\left(\frac{5}{7},\frac{2}{7}\right)$
2. (5,2)

Exercise 5.1E.7

$$\begin{cases} x+5y=10 \\ y=\frac{3}{5}x+1 \end{cases}$$

1. (−10,4)
2. $\left(\frac{5}{4},\frac{7}{4}\right)$

Answer

1. no
2. yes

Exercise 5.1E.8

$$\begin{cases} x+3y=9 \\ y=\frac{2}{3}x-2 \end{cases}$$


1. (−6,5)
2. $\left(5,\frac{4}{3}\right)$

Solve a System of Linear Equations by Graphing In the following exercises, solve the following systems of equations by graphing.

Exercise 5.1E.9

$$\begin{cases} 3x+y=-3 \\ 2x+3y=5 \end{cases}$$

Answer



Solution Point: $(-2, 3)$


Exercise 5.1E. 10

$$\begin{cases} -x + y = 2 \\ 2x + y = -4 \end{cases}$$

Exercise 5.1E. 11

$$\begin{cases} -3x + y = -1 \\ 2x + y = 4 \end{cases}$$

Answer



Solution Point: $(1, 2)$

Exercise 5.1E. 12

$$\begin{cases} -2x + 3y = -3 \\ x + y = 4 \end{cases}$$

https://math.libretexts.org/@go/page/30520

Exercise 5.1E.13

$$\begin{cases} y = x + 2 \\ y = -2x + 2 \end{cases}$$

Answer

Solution Point: $(0, 2)$

Exercise 5.1E.14

$$\begin{cases} y = x - 2 \\ y = -3x + 2 \end{cases}$$

Exercise 5.1E.15

$$\begin{cases} y = \frac{3}{2}x + 1 \\ y = -\frac{1}{2}x + 5 \end{cases}$$

Answer

Solution Point: $(2, 4)$

Exercise 5.1E.16

$$\begin{cases} y = \frac{2}{3}x - 2 \\ y = -\frac{1}{3}x - 5 \end{cases}$$

Exercise 5.1E.17

$$\begin{cases} -x + y = -3 \\ 4x + 4y = 4 \end{cases}$$

Answer

Solution Point: $(2, -1)$

Exercise 5.1E.18

$$\begin{cases} x - y = 3 \\ 2x - y = 4 \end{cases}$$

Exercise 5.1E.19

$$\begin{cases} -3x + y = -1 \\ 2x + y = 4 \end{cases}$$

Answer

Solution Point: $(1, 2)$

https://math.libretexts.org/@go/page/30520

Exercise 5.1E.20

$$\begin{cases} -3x + y = -2 \\ 4x - 2y = 6 \end{cases}$$

Exercise 5.1E.21

$$\begin{cases} x + y = 5 \\ 2x - y = 4 \end{cases}$$

Answer

Solution Point: $(3, 2)$

Exercise 5.1E.22

$$\begin{cases} x - y = 2 \\ 2x - y = 6 \end{cases}$$

Exercise 5.1E.23

$$\begin{cases} x + y = 2 \\ x - y = 0 \end{cases}$$

Answer

Solution Point: $(1, 1)$

Exercise 5.1E.24

$$\begin{cases} x + y = 6 \\ x - y = -8 \end{cases}$$

Exercise 5.1E.25

$$\begin{cases} x + y = -5 \\ x - y = 3 \end{cases}$$

Answer

Solution Point: $(-1, -4)$

Exercise 5.1E.26

$$\begin{cases} x + y = 4 \\ x - y = 0 \end{cases}$$

Exercise 5.1E.27

$$\begin{cases} x + y = -4 \\ -x + 2y = -2 \end{cases}$$

Answer

Solution Point: $(3, 3)$

Exercise 5.1E.28

$$\begin{cases} -x+3y=3 \\ x+3y=3 \end{cases}$$

Exercise 5.1E.29

$$\begin{cases} -2x+3y=3 \\ x+3y=12 \end{cases}$$

Answer

Solution Point: $(-5, 6)$


Exercise 5.1E.30

$$\begin{cases} 2x-y=4 \\ 2x+3y=12 \end{cases}$$

Exercise 5.1E.31

$$\begin{cases} 2x+3y=6 \\ y=-2 \end{cases}$$

Answer



Solution Point: $(6, -2)$

Exercise 5.1E.32

$$\begin{cases} -2x+y=2 \\ y=4 \end{cases}$$

Exercise 5.1E.33

$$\begin{cases} x-3y=-3 \\ y=2 \end{cases}$$

Answer

Solution Point: $(3, 2)$

https://math.libretexts.org/@go/page/30520


Exercise $5.1E.34$

$$\begin{cases} 2x-2y=8 \\ y=-3 \end{cases}$$

Exercise $5.1E.35$

$$\begin{cases} 2x-y=-1 \\ x=1 \end{cases}$$

Answer



Solution Point: $(1,3)$

Exercise $5.1E.36$


$$\begin{cases} x+2y=2 \\ x=-2 \end{cases}$$

Exercise $5.1E.37$

$$\begin{cases} x-3y=-6 \\ x=-3 \end{cases}$$

Answer

https://math.libretexts.org/@go/page/30520



Solution Point: $(-3, 1)$


Exercise 5.1E. 38

$$\begin{cases} x + y = 4 \\ x = 1 \end{cases}$$

Exercise 5.1E. 39

$$\begin{cases} 4x - 3y = 8 \\ 8x - 6y = 14 \end{cases}$$

Answer



no solution

Exercise 5.1E. 40


$$\begin{cases} x + 3y = 4 \\ -2x - 6y = 3 \end{cases}$$

https://math.libretexts.org/@go/page/30520

Exercise 5.1E. 41

$$\begin{cases} -2x+4y=4 \\ y=\frac{1}{2}x \end{cases}$$

Answer



no solution

Exercise 5.1E. 42

$$\begin{cases} 3x+5y=10 \\ y=-\frac{3}{5}x+1 \end{cases}$$

Exercise 5.1E. 43

$$\begin{cases} x=-3y+4 \\ 2x+6y=8 \end{cases}$$

Answer

no solution

Exercise 5.1E. 44


$$\begin{cases} 4x=3y+7 \\ 8x-6y=14 \end{cases}$$

Exercise 5.1E. 45

$$\begin{cases} 2x+y=6 \\ -8x-4y=-24 \end{cases}$$

Answer

https://math.libretexts.org/@go/page/30520



infinitely many solutions with solution set: $\{(x, y)|2x + y = 6\}$

? Exercise 5.1E. 46

$$\begin{cases} 5x + 2y = 7 \\ -10x - 4y = -14 \end{cases}$$

? Exercise 5.1E. 47

$$\begin{cases} x + 3y = -6 \\ 4y = -\frac{4}{3}x - 8 \end{cases}$$

Answer

infinitely many solutions with solution set: $\{(x, y)|x + 3y = -6\}$

? Exercise 5.1E. 48

$$\begin{cases} -x + 2y = -6 \\ y = -\frac{1}{2}x - 1 \end{cases}$$

? Exercise 5.1E. 49

$$\begin{cases} -3x + 2y = -2 \\ y = -x + 4 \end{cases}$$

Answer

Solution Point: $(2, 2)$

? Exercise 5.1E. 50

$$\begin{cases} -x + 2y = -2 \\ y = -x - 1 \end{cases}$$

Determine the Number of Solutions of a Linear System Without graphing the following systems of equations, determine the number of solutions and then classify the system of equations.

Exercise 5.1E.51

$$\begin{cases} y = \frac{2}{3}x + 1 \\ -2x + 3y = 5 \end{cases}$$

Answer

0 solutions

Exercise 5.1E.52

$$\begin{cases} y = \frac{1}{3}x + 2 \\ x - 3y = 9 \end{cases}$$

Exercise 5.1E.53

$$\begin{cases} y = -2x + 1 \\ 4x + 2y = 8 \end{cases}$$

Answer

0 solutions

Exercise 5.1E.54

$$\begin{cases} y = 3x + 4 \\ 9x - 3y = 18 \end{cases}$$

Exercise 5.1E.55

$$\begin{cases} y = \frac{2}{3}x + 1 \\ 2x - 3y = 7 \end{cases}$$

Answer

0 solutions

Exercise 5.1E.56

$$\begin{cases} 3x + 4y = 12 \\ y = -3x - 1 \end{cases}$$

Exercise 5.1E.57

$$\begin{cases} 4x + 2y = 10 \\ 4x - 2y = -6 \end{cases}$$

Answer

consistent, 1 solution

Exercise 5.1E.58

$$\begin{cases} 5x+3y=4 \\ 2x-3y=5 \end{cases}$$

Exercise 5.1E.59

$$\begin{cases} y=-\frac{1}{2}x+5 \\ x+2y=10 \end{cases}$$

Answer

infinitely many solutions

Exercise 5.1E.60

$$\begin{cases} y=x+1 \\ -x+y=1 \end{cases}$$

Exercise 5.1E.61

$$\begin{cases} y=2x+3 \\ 2x-y=-3 \end{cases}$$

Answer

infinitely many solutions

Exercise 5.1E.62

$$\begin{cases} 5x-2y=10 \\ y=\frac{5}{2}x-5 \end{cases}$$

Solve Applications of Systems of Equations by Graphing In the following exercises, solve.

Exercise 5.1E.63

Molly is making strawberry infused water. For each ounce of strawberry juice, she uses three times as many ounces of water. How many ounces of strawberry juice and how many ounces of water does she need to make 64 ounces of strawberry infused water?

Answer

Molly needs 16 ounces of strawberry juice and 48 ounces of water.

Exercise 5.1E.64

Jamal is making a snack mix that contains only pretzels and nuts. For every ounce of nuts, he will use 2 ounces of pretzels. How many ounces of pretzels and how many ounces of nuts does he need to make 45 ounces of snack mix?

Exercise 5.1E.65

Enrique is making a party mix that contains raisins and nuts. For each ounce of nuts, he uses twice the amount of raisins. How many ounces of nuts and how many ounces of raisins does he need to make 24 ounces of party mix?

Answer

Enrique needs 8 ounces of nuts and 16 ounces of water.

Exercise 5.1E.66

Owen is making lemonade from concentrate. The number of quarts of water he needs is 4 times the number of quarts of concentrate. How many quarts of water and how many quarts of concentrate does Owen need to make 100 quarts of lemonade?

Everyday Math

Exercise 5.1E.67

Leo is planning his spring flower garden. He wants to plant tulip and daffodil bulbs. He will plant 6 times as many daffodil bulbs as tulip bulbs. If he wants to plant 350 bulbs, how many tulip bulbs and how many daffodil bulbs should he plant?

Answer

Leo should plant 50 tulips and 300 daffodils.

Exercise 5.1E.68

A marketing company surveys 1,200 people. They surveyed twice as many females as males. How many males and females did they survey?

Writing Exercises

Exercise 5.1E.69

In a system of linear equations, the two equations have the same slope. Describe the possible solutions to the system.

Answer

Given that it is only known that the slopes of both linear equations are the same, there are either no solutions (the graphs of the equations are parallel) or infinitely many.

Exercise 5.1E.70

In a system of linear equations, the two equations have the same intercepts. Describe the possible solutions to the system.

Self Check

After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
determine whether an ordered pair is a solution of a system of equations.			
solve a system of linear equations by graphing.			
determine the number of solutions of a linear system.			
solve applications of systems of equations by graphing.			

If most of your checks were:

…confidently. Congratulations! You have achieved the objectives in this section. Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific.

…with some help. This must be addressed quickly because topics you do not master become potholes in your road to success. In math every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no - I don't get it! This is a warning sign and you must not ignore it. You should get help right away or you will quickly be overwhelmed. See your instructor as soon as you can to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 5.1E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.1E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.2: Solve Systems of Equations by Substitution

Learning Objectives

By the end of this section, you will be able to:

- Solve a system of equations by substitution
- Solve applications of systems of equations by substitution

Note

Before you get started, take this readiness quiz.

1. Simplify −5(3−x).
   If you missed this problem, review Example 1.10.43.
2. Simplify 4−2(n+5).
   If you missed this problem, review Example 1.10.49.
3. Solve for y. 8y−8=32−2y
   If you missed this problem, review Example 2.3.22.
4. Solve for x. 3x−9y=−3
   If you missed this problem, review Example 2.6.22.

Solving systems of linear equations by graphing is a good way to visualize the types of solutions that may result. However, there are many cases where solving a system by graphing is inconvenient or imprecise. If the graphs extend beyond the small grid with x and y both between −10 and 10, graphing the lines may be cumbersome. And if the solutions to the system are not integers, it can be hard to read their values precisely from a graph.

In this section, we will solve systems of linear equations by the substitution method.

Solve a System of Equations by Substitution

We will use the same system we used first for graphing.

$$\begin{cases} 2x+y=7 \\ x-2y=6 \end{cases}$$

We will first solve one of the equations for either x or y. We can choose either equation and solve for either variable—but we'll try to make a choice that will keep the work easy.

Then we substitute that expression into the other equation. The result is an equation with just one variable—and we know how to solve those!

After we find the value of one variable, we will substitute that value into one of the original equations and solve for the other variable. Finally, we check our solution and make sure it makes both equations true.

We'll fill in all these steps now in Example 5.2.1.

Example 5.2.1: How to Solve a System of Equations by Substitution

Solve the system by substitution. $\begin{cases} 2x+y=7 \\ x-2y=6 \end{cases}$

Solution

Step 1. Solve one of the equations for either variable.	We'll solve the first equation for y.	$2x+y=7$ $y=7-2x$
Step 2. Substitute the expression from Step 1 into the other equation.	We replace y in the second equation with the expression $7-2x$.	$x-2y=6$ $x-2(7-2x)=6$
Step 3. Solve the resulting equation.	Now we have an equation with just 1 variable. We know how to solve this!	$x-2(7-2x)=6$ $x-14+4x=6$ $5x=20$ $x=4$
Step 4. Substitute the solution in Step 3 into one of the original equations to find the other variable.	We'll use the first equation and replace x with 4.	$2x+y=7$ $2(4)+y=7$ $8+y=7$ $y=-1$
Step 5. Write the solution as an ordered pair.	The ordered pair is (x, y).	$(4,-1)$
Step 6. Check that the ordered pair is a solution to **both** original equations.	Substitute $(4,-1)$ into both equations and make sure they are both true.	$2x+y=7$ $2(4)+(-1)\stackrel{?}{=}7$ $7=7$ ✓ $x-2y=6$ $4-2(-1)\stackrel{?}{=}6$ $6=6$ ✓ Both equations are true. $(4,-1)$ is the solution to the system.

Try It 5.2.2

Solve the system by substitution. $\begin{cases} -2x+y=-11 \\ x+3y=9 \end{cases}$

Answer

(6,1)

Try It 5.2.3

Solve the system by substitution. $\begin{cases} x+3y=10 \\ 4x+y=18 \end{cases}$

Answer

(4,2)

SOLVE A SYSTEM OF EQUATIONS BY SUBSTITUTION.

1. Solve one of the equations for either variable.
2. Substitute the expression from Step 1 into the other equation.
3. Solve the resulting equation.
4. Substitute the solution in Step 3 into one of the original equations to find the other variable.

5. Write the solution as an ordered pair.
6. Check that the ordered pair is a solution to **both** original equations.

If one of the equations in the system is given in slope–intercept form, Step 1 is already done! We'll see this in Example 5.2.4.

Example 5.2.4

Solve the system by substitution. $\begin{cases} x+y=-1 \\ y=x+5 \end{cases}$

Solution

The second equation is already solved for y. We will substitute the expression in place of y in the first equation.

	$x+y=-1$ $y=x+5$
The second equation is already solved for y. We will substitute into the first equation.	
Replace the y with $x+5$.	$y=x+5$ $x+y=-1$
Solve the resulting equation for x.	$x+x+5=-1$
	$2x+5=-1$
	$2x=-6$
Substitute $x=-3$ into $y=x+5$ to find y.	$x=-3$ $y=x+5$
	$y=-3+5$
The ordered pair is $(-3, 2)$.	$y=2$
Check the ordered pair in both equations: $x+y=-1 \quad y=x+5$ $-3+2 \stackrel{?}{=} -1 \quad 2 \stackrel{?}{=} -3+5$ $-1=-1\checkmark \quad 2=2\checkmark$	
	The solution is $(-3, 2)$.

Try It 5.2.5

Solve the system by substitution. $\begin{cases} x+y=6 \\ y=3x-2 \end{cases}$

Answer

(2,4)

Try It 5.2.6

Solve the system by substitution. $\begin{cases} 2x-y=1 \\ y=-3x-6 \end{cases}$

Answer

(−1,−3)

If the equations are given in standard form, we'll need to start by solving for one of the variables. In this next example, we'll solve the first equation for y.

Example 5.2.7

Solve the system by substitution. $\begin{cases} 3x+y=5 \\ 2x+4y=-10 \end{cases}$

Solution

We need to solve one equation for one variable. Then we will substitute that expression into the other equation.

Solve for y. Substitute into the other equation.	$3x+y=5$ $y=-3x+5$ $2x+4y=-10$
Replace the y with $-3x+5$.	$2x+4(-3x+5)=-10$
Solve the resulting equation for x.	$2x-12x+20=-10$
	$-10x+20=-10$ $-10x=-30$
Substitute $x=3$ into $3x+y=5$ to find y.	$x=3$ $3x+y=5$
	$3(3)+y=5$ $9+y=5$
The ordered pair is $(3,-4)$.	$y=-4$
Check the ordered pair in both equations: $3x+y=5$ $\quad$ $2x+4y=-10$ $3\cdot 3+(-4)\stackrel{?}{=}5$ $\quad$ $2\cdot 3+4(-4)\stackrel{?}{=}-10$ $9-4\stackrel{?}{=}5$ $\quad$ $6-16\stackrel{?}{=}-10$ $5=5\checkmark$ $\quad$ $-10=-10\checkmark$	
	The solution is $(3,-4)$.

Try It 5.2.8

Solve the system by substitution. $\begin{cases} 4x+y=2 \\ 3x+2y=-1 \end{cases}$

Answer

$(1,-2)$

Try It 5.2.9

Solve the system by substitution. $\begin{cases} -x+y=4 \\ 4x-y=2 \end{cases}$

Answer

$(2,6)$

In Example 5.2.7 it was easiest to solve for y in the first equation because it had a coefficient of 1. In Example 5.2.10 it will be easier to solve for x.

Example 5.2.10

Solve the system by substitution. $\begin{cases} x-2y=-2 \\ 3x+2y=34 \end{cases}$

Solution

We will solve the first equation for x and then substitute the expression into the second equation.

	$x-2y=-2$
Solve for x. Substitute into the other equation.	$x=2y-2$ $3x+2y=34$
Replace the x with $2y-2$.	$3(2y-2)+2y=34$
Solve the resulting equation for y.	$6y-6+2y=34$
Substitute $y=5$ into $x-2y=-2$ to find x.	$8y-6=34$ $8y=40$ $y=5$ $x-2y=-2$ $x-2\cdot 5=-2$ $x-10=-2$ $x=8$
The ordered pair is (8, 5).	
Check the ordered pair in both equations: $x-2y=-2$: $8-2\cdot 5 \stackrel{?}{=} -2$, $8-10 \stackrel{?}{=} -2$, $-2=-2$ ✓ $3x+2y=34$: $3\cdot 8+2\cdot 5 \stackrel{?}{=} 34$, $24+10 \stackrel{?}{=} 34$, $34=34$ ✓	
	The solution is (8, 5).

Try It 5.2.11

Solve the system by substitution. $\begin{cases} x-5y=13 \\ 4x-3y=1 \end{cases}$

Answer

(−2,−3)

Try It 5.2.12

Solve the system by substitution. $\begin{cases} x-6y=-6 \\ 2x-4y=4 \end{cases}$

Answer

(6,2)

When both equations are already solved for the same variable, it is easy to substitute!

Example 5.2.13

Solve the system by substitution. $\begin{cases} y = -2x + 5 \\ y = \frac{1}{2}x \end{cases}$

Solution

Since both equations are solved for *y*, we can substitute one into the other.

Substitute $\frac{1}{2}x$ for *y* in the first equation.	$y = \frac{1}{2}x$ $y = -2x + 5$
Replace the *y* with $\frac{1}{2}x$	$\frac{1}{2}x = -2x + 5$
Solve the resulting equation. Start by clearing the fraction.	$2\left(\frac{1}{2}x\right) = 2(-2x + 5)$
Solve for *x*.	$x = -4x + 10$
	$5x = 10$
Substitute $x = 2$ into $y = \frac{1}{2}x$ to find *y*.	$x = 2$ $y = \frac{1}{2}x$ $y = \frac{1}{2} \cdot 2$ $y = 1$
The ordered pair is (2,1).	
Check the ordered pair in both equations: $\begin{array}{rcl} y & = & \frac{1}{2}x \\ 1 & \stackrel{?}{=} & \frac{1}{2} \cdot 2 \\ 1 & = & 1\checkmark \end{array}$ $\begin{array}{rcl} y & = & -2x+5 \\ 1 & \stackrel{?}{=} & -2 \cdot 2+5 \\ 1 & = & -4+5 \\ 1 & = & 1\checkmark \end{array}$	
	The solution is (2,1).

Try It 5.2.14

Solve the system by substitution. $\begin{cases} y = 3x - 16 \\ y = \frac{1}{3}x \end{cases}$

Answer

(6,2)

Try It 5.2.15

Solve the system by substitution. $\begin{cases} y = -x + 10 \\ y = \frac{1}{4}x \end{cases}$

Answer

(8,2)

Be very careful with the signs in the next example.

Example 5.2.16

Solve the system by substitution. $\begin{cases} 4x+2y=4 \\ 6x-y=8 \end{cases}$

Solution

We need to solve one equation for one variable. We will solve the first equation for y.

	$4x+2y=4$
Solve the first equation for y.	$2y=-4x+4$
Substitute $-2x+2$ for y in the second equation.	$y=-2x+2$ $6x-y=8$
Replace the y with $-2x+2$.	$6x-(-2x+2)=8$
Solve the equation for x.	$6x+2x-2=8$
	$8x-2=8$ $8x=10$
Substitute $x=\frac{5}{4}$ into $4x+2y=4$ to find y.	$x=\frac{5}{4}$ $4x+2y=4$ $4\left(\frac{5}{4}\right)+2y=4$ $5+2y=4$ $2y=-1$ $y=-\frac{1}{2}$
The ordered pair is $(\frac{5}{4},-\frac{1}{2})$.	
Check the ordered pair in both equations.	$\begin{aligned} 4x+2y &= 4 \\ 4(\tfrac{5}{4})+2(-\tfrac{1}{2}) &\stackrel{?}{=} 4 \\ 5-1 &\stackrel{?}{=} 4 \\ 4 &= 4\checkmark \end{aligned}$ $\begin{aligned} 6x-y &= 8 \\ 6(\tfrac{5}{4})-(-\tfrac{1}{2}) &\stackrel{?}{=} 8 \\ \tfrac{15}{4}-(-\tfrac{1}{2}) &\stackrel{?}{=} 8 \\ \tfrac{16}{2} &\stackrel{?}{=} 8 \\ 8 &= 8\checkmark \end{aligned}$
	The solution is $\left(\frac{5}{4},-\frac{1}{2}\right)$.

Try It 5.2.17

Solve the system by substitution. $\begin{cases} x-4y=-4 \\ -3x+4y=0 \end{cases}$

Answer

$(2,\frac{3}{2})$

Try It 5.2.18

Solve the system by substitution. $\begin{cases} 4x-y=0 \\ 2x-3y=5 \end{cases}$

Answer

$(-\frac{1}{2}, -2)$

In Example, it will take a little more work to solve one equation for x or y.

Example 5.2.19

Solve the system by substitution. $\begin{cases} 4x-3y=6 \\ 15y-20x=-30 \end{cases}$

Solution

We need to solve one equation for one variable. We will solve the first equation for x.

	$4x - 3y = 6$
Solve the first equation for x.	$4x = 3y + 6$
Substitute $\frac{3}{4}y+\frac{3}{2}$ for x in the second equation.	$x = \frac{3}{4}y + \frac{3}{2}$ $15y - 20x = -30$
Replace the x with $\frac{3}{4}y+\frac{3}{2}$	$15y - 20\left(\frac{3}{4}y + \frac{3}{2}\right) = -30$
Solve for y.	$15y - 15y - 30 = -30$
	$0 - 30 = -30$
	$0 = 0$

Since 0 = 0 is a true statement, the system is consistent. The equations are dependent. The graphs of these two equations would give the same line. The system has infinitely many solutions.

Try It 5.2.20

Solve the system by substitution. $\begin{cases} 2x-3y=12 \\ -12y+8x=48 \end{cases}$

Answer

infinitely many solutions

Try It 5.2.21

Solve the system by substitution. $\begin{cases} 5x+2y=12 \\ -4y-10x=-24 \end{cases}$

Answer

infinitely many solutions

Look back at the equations in Example 5.2.22 Is there any way to recognize that they are the same line?

Let's see what happens in the next example.

Example 5.2.22

Solve the system by substitution. $\begin{cases} 5x-2y=-10 \\ y=\frac{5}{2}x \end{cases}$

Solution

The second equation is already solved for y, so we can substitute for y in the first equation.

Substitute x for y in the first equation.	$y=\left(\frac{5}{2}x\right)$ $5x-2y=-10$
Replace the y with $\frac{5}{2}x$.	$5x-2\left(\frac{5}{2}x\right)=-10$
Solve for x.	$5x-5x=-10$
	$0\neq-10$

Since $0 = -10$ is a false statement the equations are inconsistent. The graphs of the two equation would be parallel lines. The system has no solutions.

Try It 5.2.23

Solve the system by substitution. $\begin{cases} 3x+2y=9 \\ y=-\frac{3}{2}x+1 \end{cases}$

Answer

no solution

Try It 5.2.24

Solve the system by substitution. $\begin{cases} 5x-3y=2 \\ y=\frac{5}{3}x-4 \end{cases}$

Answer

no solution

Solve Applications of Systems of Equations by Substitution

We'll copy here the problem solving strategy we used in the Solving Systems of Equations by Graphing section for solving systems of equations. Now that we know how to solve systems by substitution, that's what we'll do in Step 5.

HOW TO USE A PROBLEM SOLVING STRATEGY FOR SYSTEMS OF LINEAR EQUATIONS.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose variables to represent those quantities.
4. **Translate** into a system of equations.
5. **Solve** the system of equations using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Some people find setting up word problems with two variables easier than setting them up with just one variable. Choosing the variable names is easier when all you need to do is write down two letters. Think about this in the next example—how would you have done it with just one variable?

Example 5.2.25

The sum of two numbers is zero. One number is nine less than the other. Find the numbers.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for two numbers.
Step 3. Name what we are looking for.	Let n= the first number Let m= the second number
Step 4. Translate into a system of equations.	The sum of two numbers is zero.
	$n+m=0$
	One number is nine less than the other.
	$n=m-9$
The system is:	$\begin{aligned} n+m&=0\\ n&=m-9\end{aligned}$
Step 5. Solve the system of equations. We will use substitution since the second equation is solved for n.	
Substitute $m-9$ for n in the first equation.	$n=m-9$ $n+m=0$
Solve for m.	$m-9+m=0$
	$2m-9=0$
	$2m=9$
Substitute $m=\frac{9}{2}$ into the second equation and then solve for n.	$m=\frac{9}{2}$ $n=m-9$
	$m=\frac{9}{2}-9$
	$m=\frac{9}{2}-\frac{18}{2}$
	$n=-\frac{9}{2}$
Step 6. Check the answer in the problem.	Do these numbers make sense in the problem? We will leave this to you!
Step 7. Answer the question.	The numbers are $\frac{9}{2}$ and $-\frac{9}{2}$.

Try It 5.2.26

The sum of two numbers is 10. One number is 4 less than the other. Find the numbers.

Answer

The numbers are 3 and 7.

Try It 5.2.27

The sum of two number is −6. One number is 10 less than the other. Find the numbers.

Answer


The numbers are 2 and −8.

In the Example 5.2.28, we'll use the formula for the perimeter of a rectangle, $P = 2L + 2W$.

Example 5.2.28

The perimeter of a rectangle is 88. The length is five more than twice the width. Find the length and width of the rectangle.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	We are looking for the length and width.
Step 3. Name what we are looking for.	Let L= the length W= the width
Step 4. Translate into a system of equations.	The perimeter of a rectangle is 88.
	$2L + 2W = P$ $2L + 2W = 88$
	The length is five more than twice the width.
	$L = 2W + 5$
The system is:	$2L + 2W = 88$ $L = 2W + 5$
Step 5. Solve the system of equations. We will use substitution since the second equation is solved for L. Substitute $2W + 5$ for L in the first equation.	$L = 2W + 5$ $2L + 2W = 88$
Solve for W.	$2(2W + 5) + 2W = 88$
	$4W + 10 + 2W = 88$
	$6W + 10 = 88$
	$6W = 78$
Substitute $W = 13$ into the second equation and then solve for L.	$W = 13$ $L = 2W + 5$
	$L = 2 \cdot 13 + 5$
	$L = 31$
Step 6. Check the answer in the problem.	Does a rectangle with length 31 and width 13 have perimeter 88? Yes.
Step 7. Answer the equation.	The length is 31 and the width is 13.

Try It 5.2.29

The perimeter of a rectangle is 40. The length is 4 more than the width. Find the length and width of the rectangle.

Answer

The length is 12 and the width is 8.

Try It 5.2.30

The perimeter of a rectangle is 58. The length is 5 more than three times the width. Find the length and width of the rectangle.

Answer

The length is 23 and the width is 6.


For Example 5.2.31 we need to remember that the sum of the measures of the angles of a triangle is 180 degrees and that a right triangle has one 90 degree angle.

Example 5.2.31

The measure of one of the small angles of a right triangle is ten more than three times the measure of the other small angle. Find the measures of both angles.

Solution

We will draw and label a figure.

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	We are looking for the measures of the angles.
Step 3. Name what we are looking for.	Let a= the measure of the 1st angle b= the measure of the 2nd angle
Step 4. Translate into a system of equations.	The measure of one of the small angles of a right triangle is ten more than three times the measure of the other small angle.
	$a = 3b + 10$
	The sum of the measures of the angles of a triangle is 180.
	$a + b + 90 = 180$
The system is:	$a = 3b + 10$ $a + b + 90 = 180$
Step 5. Solve the system of equations. We will use substitution since the first equation is solved for a.	$a = 3b + 10$ $a + b + 90 = 180$
Substitute $3b + 10$ for a in the second equation.	$(3b + 10) + b + 90 = 180$
Solve for b.	$4b + 100 = 180$

	$4b = 80$
	$b = 20$ $a = 3b + 10$
Substitute $b = 20$ into the first equation and then solve for a.	$a = 3 \cdot 20 + 10$ $a = 70$
Step 6. Check the answer in the problem.	We will leave this to you!
Step 7. Answer the question.	The measures of the small angles are 20 and 70.

Try It 5.2.32

The measure of one of the small angles of a right triangle is 2 more than 3 times the measure of the other small angle. Find the measure of both angles.

Answer

The measure of the angles are 22 degrees and 68 degrees.

Try It 5.2.33

The measure of one of the small angles of a right triangle is 18 less than twice the measure of the other small angle. Find the measure of both angles.

Answer

The measure of the angles are 36 degrees and 54 degrees.

Example 5.2.34

Heather has been offered two options for her salary as a trainer at the gym. Option A would pay her $25,000 plus $15 for each training session. Option B would pay her $10,000 + $40 for each training session. How many training sessions would make the salary options equal?

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	We are looking for the number of training sessions that would make the pay equal.
Step 3. Name what we are looking for.	Let s= Heather's salary. n= the number of training sessions
Step 4. Translate into a system of equations.	Option A would pay her $25,000 plus $15 for each training session.
	$s = 25{,}000 + 15n$
	Option B would pay her $10,000 + $40 for each training session
	$s = 10{,}000 + 40n$
The system is:	$s = 25{,}000 + 15n$ $s = 10{,}000 + 40n$

Step 5. Solve the system of equations. We will use substitution.	$s = 25{,}000 + 15n$ $s = 10{,}000 + 40n$
Substitute $25{,}000 + 15n$ for s in the second equation.	$25{,}000 + 15n = 10{,}000 + 40n$
Solve for n.	$25{,}000 = 10{,}000 + 25n$
	$15{,}000 = 25n$
	$600 = n$
Step 6. Check the answer.	Are 600 training sessions a year reasonable? Are the two options equal when $n = 600$?
Step 7. Answer the question.	The salary options would be equal for 600 training sessions.

Try It 5.2.35

Geraldine has been offered positions by two insurance companies. The first company pays a salary of $12,000 plus a commission of $100 for each policy sold. The second pays a salary of $20,000 plus a commission of $50 for each policy sold. How many policies would need to be sold to make the total pay the same?

Answer

There would need to be 160 policies sold to make the total pay the same.

Try It 5.2.36

Kenneth currently sells suits for company A at a salary of $22,000 plus a $10 commission for each suit sold. Company B offers him a position with a salary of $28,000 plus a $4 commission for each suit sold. How many suits would Kenneth need to sell for the options to be equal?

Answer

Kenneth would need to sell 1,000 suits.

Note

Access these online resources for additional instruction and practice with solving systems of equations by substitution.

- Instructional Video-Solve Linear Systems by Substitution
- Instructional Video-Solve by Substitution

Key Concepts

- **Solve a system of equations by substitution**
  1. Solve one of the equations for either variable.
  2. Substitute the expression from Step 1 into the other equation.
  3. Solve the resulting equation.
  4. Substitute the solution in Step 3 into one of the original equations to find the other variable.
  5. Write the solution as an ordered pair.
  6. Check that the ordered pair is a solution to both original equations.

This page titled 5.2: Solve Systems of Equations by Substitution is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.2: Solve Systems of Equations by Substitution** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.2E: Exercises

Practice Makes Perfect

Solve a System of Equations by Substitution

In the following exercises, solve the systems of equations by substitution.

Exercise 5.2E.1

$$\begin{cases} 2x+y=-4 \\ 3x-2y=-6 \end{cases}$$

Answer

$(-2,0)$

Exercise 5.2E.2

$$\begin{cases} 2x+y=-2 \\ 3x-y=7 \end{cases}$$

Exercise 5.2E.3

$$\begin{cases} x-2y=-5 \\ 2x-3y=-4 \end{cases}$$

Answer

$(7,6)$

Exercise 5.2E.4

$$\begin{cases} x-3y=-9 \\ 2x+5y=4 \end{cases}$$

Exercise 5.2E.5

$$\begin{cases} 5x-2y=-6 \\ y=3x+3 \end{cases}$$

Answer

$(0,3)$

Exercise 5.2E.6

$$\begin{cases} -2x+2y=6 \\ y=-3x+1 \end{cases}$$

Exercise 5.2E.7

$$\begin{cases} 2x+3y=3 \\ y=-x+3 \end{cases}$$

Answer

$(6, -3)$

Exercise 5.2E.8

$$\begin{cases} 2x + 5y = -14 \\ y = -2x + 2 \end{cases}$$

Exercise 5.2E.9

$$\begin{cases} 2x + 5y = 1 \\ y = \frac{1}{3}x - 2 \end{cases}$$

Answer

$(3, -1)$

Exercise 5.2E.10

$$\begin{cases} 3x + 4y = 1 \\ y = -\frac{2}{5}x + 2 \end{cases}$$

Exercise 5.2E.11

$$\begin{cases} 3x - 2y = 6 \\ y = \frac{2}{3}x + 2 \end{cases}$$

Answer

$(6, 6)$

Exercise 5.2E.12

$$\begin{cases} -3x - 5y = 3 \\ y = \frac{1}{2}x - 5 \end{cases}$$

Exercise 5.2E.13

$$\begin{cases} 2x + y = 10 \\ -x + y = -5 \end{cases}$$

Answer

$(5, 0)$

Exercise 5.2E.14

$$\begin{cases} -2x + y = 10 \\ -x + 2y = 16 \end{cases}$$

Exercise 5.2E.15

$$\begin{cases} 3x+y=1 \\ -4x+y=15 \end{cases}$$

Answer

$(-2,7)$

Exercise 5.2E.16

$$\begin{cases} x+y=0 \\ 2x+3y=-4 \end{cases}$$

Exercise 5.2E.17

$$\begin{cases} x+3y=1 \\ 3x+5y=-5 \end{cases}$$

Answer

$(-5,2)$

Exercise 5.2E.18

$$\begin{cases} x+2y=-1 \\ 2x+3y=1 \end{cases}$$

Exercise 5.2E.19

$$\begin{cases} 2x+y=5 \\ x-2y=-15 \end{cases}$$

Answer

$(-1,7)$

Exercise 5.2E.20

$$\begin{cases} 4x+y=10 \\ x-2y=-20 \end{cases}$$

Exercise 5.2E.21

$$\begin{cases} y=-2x-1 \\ y=-\frac{1}{3}x+4 \end{cases}$$

Answer

$(-3,5)$

Exercise 5.2E.22

$$\begin{cases} y=x-6 \\ y=-\frac{3}{2}x+4 \end{cases}$$

Exercise 5.2E.23

$$\begin{cases} y = 2x - 8 \\ y = \frac{3}{5}x + 6 \end{cases}$$

Answer

$(10, 12)$

Exercise 5.2E.24

$$\begin{cases} y = -x - 1 \\ y = x + 7 \end{cases}$$

Exercise 5.2E.25

$$\begin{cases} 4x + 2y = 8 \\ 8x - y = 1 \end{cases}$$

Answer

$\left(\frac{1}{2}, 3\right)$

Exercise 5.2E.26

$$\begin{cases} -x - 12y = -1 \\ 2x - 8y = -6 \end{cases}$$

Exercise 5.2E.27

$$\begin{cases} 15x + 2y = 6 \\ -5x + 2y = -4 \end{cases}$$

Answer

$\left(\frac{1}{2}, -\frac{3}{4}\right)$

Exercise 5.2E.28

$$\begin{cases} 2x - 15y = 7 \\ 12x + 2y = -4 \end{cases}$$

Exercise 5.2E.29

$$\begin{cases} y = 3x \\ 6x - 2y = 0 \end{cases}$$

Answer

Infinitely many solutions. The two equations represent the same line.
The solution set is: $\{(x, y) \mid y = 3x\}$
All points that are solutions of the equation $y = 3x$ are solutions of this system.

https://math.libretexts.org/@go/page/30522

Exercise 5.2E.30

$$\begin{cases} x = 2y \\ 4x - 8y = 0 \end{cases}$$

Exercise 5.2E.31

$$\begin{cases} 2x + 16y = 8 \\ -x - 8y = -4 \end{cases}$$

Answer

Infinitely many solutions. The two equations represent the same line.
The solution set is: $\{(x, y) \mid 2x + 16y = 8\}$
All points that are solutions of the equation $2x + 16y = 8$ are solutions of this system.

Exercise 5.2E.32

$$\begin{cases} 15x + 4y = 6 \\ -30x - 8y = -12 \end{cases}$$

Exercise 5.2E.33

$$\begin{cases} y = -4x \\ 4x + y = 1 \end{cases}$$

Answer

No solution

Exercise 5.2E.34

$$\begin{cases} y = -\frac{1}{4}x \\ x + 4y = 8 \end{cases}$$

Exercise 5.2E.35

$$\begin{cases} y = \frac{7}{8}x + 4 \\ -7x + 8y = 6 \end{cases}$$

Answer

No solution

Exercise 5.2E.36

$$\begin{cases} y = -\frac{2}{3}x + 5 \\ 2x + 3y = 11 \end{cases}$$

Solve Applications of Systems of Equations by Substitution

In the following exercises, translate to a system of equations and solve.

Exercise 5.2E.37

The sum of two numbers is 15. One number is 3 less than the other. Find the numbers.

Answer

The numbers are 6 and 9.

Exercise 5.2E.38

The sum of two numbers is 30. One number is 4 less than the other. Find the numbers.

Exercise 5.2E.39

The sum of two numbers is −26. One number is 12 less than the other. Find the numbers.

Answer

The numbers are −7 and −19.

Exercise 5.2E.40

The perimeter of a rectangle is 50. The length is 5 more than the width. Find the length and width.

Exercise 5.2E.41

The perimeter of a rectangle is 60. The length is 10 more than the width. Find the length and width.

Answer

The length is 20 and the width is 10.

Exercise 5.2E.42

The perimeter of a rectangle is 58. The length is 5 more than three times the width. Find the length and width.

Exercise 5.2E.43

The perimeter of a rectangle is 84. The length is 10 more than three times the width. Find the length and width.

Answer

The length is 34 and the width is 8.

Exercise 5.2E.44

The measure of one of the small angles of a right triangle is 14 more than 3 times the measure of the other small angle. Find the measure of both angles.

Exercise 5.2E.45

The measure of one of the small angles of a right triangle is 26 more than 3 times the measure of the other small angle. Find the measure of both angles.

Answer

The measures are 16° and 74°

Exercise 5.2E.46

The measure of one of the small angles of a right triangle is 15 less than twice the measure of the other small angle. Find the measure of both angles.

Exercise 5.2E.47

The measure of one of the small angles of a right triangle is 45 less than twice the measure of the other small angle. Find the measure of both angles.

Answer

The measures are 45° and 45°.

Exercise 5.2E.48

Maxim has been offered positions by two car dealers. The first company pays a salary of $10,000 plus a commission of $1,000 for each car sold. The second pays a salary of $20,000 plus a commission of $500 for each car sold. How many cars would need to be sold to make the total pay the same?

Exercise 5.2E.49

Jackie has been offered positions by two cable companies. The first company pays a salary of $ 14,000 plus a commission of $100 for each cable package sold. The second pays a salary of $20,000 plus a commission of $25 for each cable package sold. How many cable packages would need to be sold to make the total pay the same?

Answer

80 cable packages would need to be sold.

Exercise 5.2E.50

Amara currently sells televisions for company A at a salary of $17,000 plus a $100 commission for each television she sells. Company B offers her a position with a salary of $29,000 plus a $20 commission for each television she sells. How televisions would Amara need to sell for the options to be equal?

Exercise 5.2E.51

Mitchell currently sells stoves for company A at a salary of $12,000 plus a $150 commission for each stove he sells. Company B offers him a position with a salary of $24,000 plus a $50 commission for each stove he sells. How many stoves would Mitchell need to sell for the options to be equal?

Answer

Mitchell would need to sell 120 stoves.

Everyday Math

Exercise 5.2E.52

When Gloria spent 15 minutes on the elliptical trainer and then did circuit training for 30 minutes, her fitness app says she burned 435 calories. When she spent 30 minutes on the elliptical trainer and 40 minutes circuit training she burned 690

calories. Solve the system $\begin{cases} 15e+30c=435 \\ 30e+40c=690 \end{cases}$ for e, the number of calories she burns for each minute on the elliptical trainer, and cc, the number of calories she burns for each minute of circuit training.

? Exercise 5.2E.53

Stephanie left Riverside, California, driving her motorhome north on Interstate 15 towards Salt Lake City at a speed of 56 miles per hour. Half an hour later, Tina left Riverside in her car on the same route as Stephanie, driving 70 miles per hour. Solve the system $\begin{cases} 56s=70t \\ s=t+\frac{1}{2} \end{cases}$

1. for t to find out how long it will take Tina to catch up to Stephanie.
2. what is the value of ss, the number of hours Stephanie will have driven before Tina catches up to her?

Answer

1. $t=2$ hours
2. $s=2\frac{1}{2}$ hours

Writing Exercises

? Exercise 5.2E.54

Solve the system of equations
$\begin{cases} x+y=10 \\ x-y=6 \end{cases}$

1. by graphing.
2. by substitution.
3. Which method do you prefer? Why?

? Exercise 5.2E.55

Solve the system of equations
$\begin{cases} 3x+y=12 \\ x=y-8 \end{cases}$ by substitution and explain all your steps in words.

Answer

Answers will vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve a system of equations by substitution.			
solve applications of systems of equations by substitution.			

b. After reviewing this checklist, what will you do to become confident for all objectives?

This page titled 5.2E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.2E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.3: Solve Systems of Equations by Elimination

Learning Objectives

By the end of this section, you will be able to:

- Solve a system of equations by elimination
- Solve applications of systems of equations by elimination
- Choose the most convenient method to solve a system of linear equations

Note

Before you get started, take this readiness quiz.

1. Simplify −5(6−3a).
   If you missed this problem, review Example 1.10.43.
2. Solve the equation $\frac{1}{3}x+\frac{5}{8}=\frac{31}{24}$.
   If you missed this problem, review Example 2.5.1.

We have solved systems of linear equations by graphing and by substitution. Graphing works well when the variable coefficients are small and the solution has integer values. Substitution works well when we can easily solve one equation for one of the variables and not have too many fractions in the resulting expression.

The third method of solving systems of linear equations is called the Elimination Method. When we solved a system by substitution, we started with two equations and two variables and reduced it to one equation with one variable. This is what we'll do with the elimination method, too, but we'll have a different way to get there.

Solve a System of Equations by Elimination

The **Elimination Method** is based on the Addition Property of Equality. The Addition Property of Equality says that when you add the same quantity to both sides of an equation, you still have equality. We will extend the Addition Property of Equality to say that when you add equal quantities to both sides of an equation, the results are equal.

For any expressions *a*, *b*, *c*, and *d*,

$$\begin{array}{ll} \text{if} & a=b \\ \text{and} & c=d \\ \text{then} & a+c=b+d \end{array} \tag{5.3.1}$$

To solve a system of equations by elimination, we start with both equations in standard form. Then we decide which variable will be easiest to eliminate. How do we decide? We want to have the coefficients of one variable be opposites, so that we can add the equations together and eliminate that variable.

Notice how that works when we add these two equations together:

$$\begin{array}{r} 3x+y=5 \\ \underline{2x-y=0} \\ 5x \quad\quad =5 \end{array} \tag{5.3.2}$$

The *y*'s add to zero and we have one equation with one variable.

Let's try another one:

$$\begin{cases} x+4y=2 \\ 2x+5y=-2 \end{cases} \tag{5.3.3}$$

This time we don't see a variable that can be immediately eliminated if we add the equations.

But if we multiply the first equation by −2, we will make the coefficients of *x* opposites. We must multiply every term on both sides of the equation by −2.

$$\begin{cases} -2(x+4y) = -2(2) \\ 2x+5y = -2 \end{cases}$$

$$\begin{cases} -2x-8y = -4 \\ 2x+5y = -2 \end{cases}$$

Now we see that the coefficients of the x terms are opposites, so x will be eliminated when we add these two equations.

Add the equations yourself—the result should be $-3y = -6$. And that looks easy to solve, doesn't it? Here is what it would look like.

$$\begin{array}{r} -2x-8y=-4 \\ 2x+5y=-2 \\ \hline -3y=-6 \end{array}$$

We'll do one more:

$$\begin{cases} 4x-3y=10 \\ 3x+5y=-7 \end{cases} \tag{5.3.4}$$

It doesn't appear that we can get the coefficients of one variable to be opposites by multiplying one of the equations by a constant, unless we use fractions. So instead, we'll have to multiply both equations by a constant.

We can make the coefficients of x be opposites if we multiply the first equation by 3 and the second by -4, so we get $12x$ and $-12x$.

$$3(4x-3y) = 3(10)$$
$$-4(3x+5y) = -4(-7)$$

This gives us these two new equations:

$$\begin{cases} 12x-9y=30 \\ -12x-20y=28 \end{cases}$$

When we add these equations,

$$\begin{array}{r} 12x-9y=30 \\ -12x-20y=28 \\ \hline -29y=58 \end{array} \tag{5.3.5}$$

the x's are eliminated and we just have $-29y = 58$.

Once we get an equation with just one variable, we solve it. Then we substitute that value into one of the original equations to solve for the remaining variable. And, as always, we check our answer to make sure it is a solution to both of the original equations.

Now we'll see how to use elimination to solve the same system of equations we solved by graphing and by substitution.

Example 5.3.1: How to Solve a System of Equations by Elimination

Solve the system by elimination. $\begin{cases} 2x+y=7 \\ x-2y=6 \end{cases}$

Solution

Step 1. Write both equations in standard form. If any coefficients are fractions, clear them.	Both equations are in standard form, $Ax + By = C$. There are no fractions.	$\begin{cases} 2x + y = 7 \\ x - 2y = 6 \end{cases}$
Step 2. Make the coefficients of one variable opposites. Decide which variable you will eliminate. Multiply one or both equations so that the coefficients of that variable are opposites.	We can eliminate the y's by multiplying the first equation by 2. Multiply both sides of $2x + y = 7$ by 2.	$\begin{cases} 2x + y = 7 \\ x - 2y = 6 \end{cases}$ $\begin{cases} 2(2x + y) = 2(7) \\ x - 2y = 6 \end{cases}$
Step 3. Add the equations resulting from Step 2 to eliminate one variable.	We add the x's, y's, and constants.	$\begin{array}{r} 4x + 2y = 14 \\ x - 2y = 6 \\ \hline 5x \quad\quad = 20 \end{array}$
Step 4. Solve for the remaining variable.	Solve for x.	$x = 4$
Step 5. Substitute the solution from Step 4 into one of the original equations. Then solve for the other variable.	Substitute $x = 4$ into the second equation, $x - 2y = 6$. Then solve for y.	$x - 2y = 6$ $4 - 2y = 6$ $-2y = 2$ $y = -1$
Step 6. Write the solution as an ordered pair.	Write it as (x, y).	$(4, -1)$
Step 7. Check that the ordered pair is a solution to **both** original equations.	Substitute $(4, -1)$ into $2x + y = 7$ and $x - 2y = 6$ Do they make both equations true? Yes!	$2x + y = 7$: $2(4) + (-1) \stackrel{?}{=} 7$: $7 = 7$ ✓ $x - 2y = 6$: $4 - 2(-1) \stackrel{?}{=} 6$: $6 = 6$ ✓ The solution is $(4, -1)$.

Try It 5.3.2

Solve the system by elimination. $\begin{cases} 3x + y = 5 \\ 2x - 3y = 7 \end{cases}$

Answer

$(2, -1)$

Try It 5.3.3

Solve the system by elimination. $\begin{cases} 4x + y = -5 \\ -2x - 2y = -2 \end{cases}$

Answer

$(-2, 3)$

The steps are listed below for easy reference.

HOW TO SOLVE A SYSTEM OF EQUATIONS BY ELIMINATION.

1. Write both equations in standard form. If any coefficients are fractions, clear them.
2. Make the coefficients of one variable opposites.
   - Decide which variable you will eliminate.
   - Multiply one or both equations so that the coefficients of that variable are opposites.
3. Add the equations resulting from Step 2 to eliminate one variable.
4. Solve for the remaining variable.
5. Substitute the solution from Step 4 into one of the original equations. Then solve for the other variable.
6. Write the solution as an ordered pair.
7. Check that the ordered pair is a solution to **both** original equations.

First we'll do an example where we can eliminate one variable right away.

Example 5.3.4

Solve the system by elimination. $\begin{cases} x+y=10 \\ x-y=12 \end{cases}$

Solution

	$\begin{array}{r} x+y=10 \\ x-y=12 \end{array}$
Both equations are in standard form.	
The coefficients of y are already opposites.	
Add the two equations to eliminate y. The resulting equation has only 1 variable, x.	$\begin{array}{r} x+y=10 \\ x-y=12 \\ \hline 2x \quad =22 \end{array}$
Solve for x, the remaining variable. Substitute $x = 11$ into one of the original equations.	$x=11$ $x+y=10$
	$11+y=10$
Solve for the other variable, y.	$y=-1$
Write the solution as an ordered pair.	The ordered pair is (11, −1).
Check that the ordered pair is a solution to **both** original equations. $\begin{array}{rcl} x+y & = & 10 \\ 11+(-1) & \stackrel{?}{=} & 10 \\ 10 & = & 10\checkmark \end{array}$ $\quad$ $\begin{array}{rcl} x-y & = & 12 \\ 11-(-1) & \stackrel{?}{=} & 12 \\ 12 & = & 12\checkmark \end{array}$	
	The solution is (11, −1).

Try It 5.3.5

Solve the system by elimination. $\begin{cases} 2x+y=5 \\ x-y=4 \end{cases}$

Answer

(3,−1)

Try It 5.3.6

Solve the system by elimination. $\begin{cases} x+y=3 \\ -2x-y=-1 \end{cases}$

Answer

$(-2,5)$

In Example 5.3.7, we will be able to make the coefficients of one variable opposites by multiplying one equation by a constant.

Example 5.3.7

Solve the system by elimination. $\begin{cases} 3x-2y=-2 \\ 5x-6y=10 \end{cases}$

Solution

	$\begin{aligned} 3x-2y&=-2\\ 5x-6y&=10 \end{aligned}$
Both equations are in standard form.	
None of the coefficients are opposites.	
We can make the coefficients of y opposites by multiplying the first equation by -3.	$\begin{aligned} -3(3x-2y)&=-3(-2)\\ 5x-6y&=10 \end{aligned}$
Simplify.	$\begin{aligned} -9x+6y&=6\\ 5x-6y&=10 \end{aligned}$
Add the two equations to eliminate y.	$\begin{array}{rl} -9x+6y&=6\\ 5x-6y&=10\\ \hline -4x&=16 \end{array}$
Solve for the remaining variable, x. Substitute $x=-4$ into one of the original equations.	$\begin{aligned} x&=-4\\ 3x-2y&=-2 \end{aligned}$
	$3(-4)-2y=-2$
Solve for y.	$\begin{aligned} -12-2y&=-2\\ -2y&=10\\ y&=-5 \end{aligned}$
Write the solution as an ordered pair.	The ordered pair is $(-4,-5)$.
Check that the ordered pair is a solution to both original equations.	$\begin{array}{rcl\|rcl} 3x-2y&=&-2 & 5x-6y&=&10\\ 3(-4)-2(-5)&\stackrel{?}{=}&-2 & 5(-4)-6(-5)&\stackrel{?}{=}&10\\ -12+10&\stackrel{?}{=}&-2 & -20+30&\stackrel{?}{=}&10\\ -2&=&-2\checkmark & 10&=&10\checkmark \end{array}$
	The solution is $(-4,-5)$.

Try It 5.3.8

Solve the system by elimination. $\begin{cases} 4x-3y=1 \\ 5x-9y=-4 \end{cases}$

Answer

(1,1)

Try It 5.3.9

Solve the system by elimination. $\begin{cases} 3x+2y=2 \\ 6x+5y=8 \end{cases}$

Answer

(−2,4)

Now we'll do an example where we need to multiply both equations by constants in order to make the coefficients of one variable opposites.

Example 5.3.10

Solve the system by elimination. $\begin{cases} 4x-3y=9 \\ 7x+2y=-6 \end{cases}$

Solution

In this example, we cannot multiply just one equation by any constant to get opposite coefficients. So we will strategically multiply both equations by a constant to get the opposites.

	$\begin{aligned} 4x-3y&=9 \\ 7x+2y&=-6 \end{aligned}$
Both equations are in standard form. To get opposite coefficients of y, we will multiply the first equation by 2 and the second equation by 3.	$\begin{aligned} 2(4x-3y)&=2(9) \\ 3(7x+2y)&=3(-6) \end{aligned}$
Simplify.	$\begin{aligned} 8x-6y&=18 \\ 21x+6y&=-18 \end{aligned}$
Add the two equations to eliminate y.	$\begin{aligned} 8x-6y&=18 \\ 21x+6y&=-18 \\ \hline 39x&=0 \end{aligned}$
Solve for x. Substitute $x = 0$ into one of the original equations.	$x=0$ $7x+2y=-6$
	$7\cdot 0+2y=-6$
Solve for y.	$2y=-6$
	$y=-3$
Write the solution as an ordered pair.	The ordered pair is (0, −3).
Check that the ordered pair is a solution to **both** original equations. $\begin{aligned} 4x-3y &= 9 \\ 4(0)-3(-3) &\stackrel{?}{=} 9 \\ 9 &= 9\checkmark \end{aligned}$ $\quad$ $\begin{aligned} 7x+2y &= -6 \\ 7(0)+2(-3) &\stackrel{?}{=} -6 \\ -6 &= -6\checkmark \end{aligned}$	
	The solution is (0, −3).

Try It 5.3.11

Solve the system by elimination. $\begin{cases} 3x-4y=-9 \\ 5x+3y=14 \end{cases}$

Answer

(1,3)

Try It 5.3.12

Solve the system by elimination. $\begin{cases} 7x+8y=4 \\ 3x-5y=27 \end{cases}$

Answer

(4,−3)

When the system of equations contains fractions, we will first clear the fractions by multiplying each equation by its LCD.

Example 5.3.13

Solve the system by elimination. $\begin{cases} x+\frac{1}{2}y=6 \\ \frac{3}{2}x+\frac{2}{3}y=\frac{17}{2} \end{cases}$

Solution

In this example, both equations have fractions. Our first step will be to multiply each equation by its LCD to clear the fractions.

	$x+\frac{1}{2}y=6$ $\frac{3}{2}x+\frac{2}{3}y=\frac{17}{2}$
To clear the fractions, multiply each equation by its LCD.	$2\left(x+\frac{1}{2}y\right)=2(6)$ $6\left(\frac{3}{2}x+\frac{2}{3}y\right)=6\left(\frac{17}{2}\right)$
Simplify.	$2x+y=12$ $9x+4y=51$
Now we are ready to eliminate one of the variables. Notice that both equations are in standard form.	
We can eliminate y multiplying the top equation by −4.	$-4(2x+y)=-4(12)$ $9x+4y=51$
Simplify and add.	$-8x-4y=-48$ $9x+4y=51$ $x=3$
Substitute $x = 3$ into one of the original equations.	$x+\frac{1}{2}y=6$
Solve for y.	$3+\frac{1}{2}y=6$
	$\frac{1}{2}y=3$
	$y=6$
Write the solution as an ordered pair.	The ordered pair is (3, 6).

Check that the ordered pair is a solution to **both** original equations.

$$\begin{aligned} x+\tfrac{1}{2}y &= 6 \\ 3+\tfrac{1}{2}(6) &\stackrel{?}{=} 6 \\ 3+3 &\stackrel{?}{=} 6 \\ 6 &= 6\checkmark \end{aligned} \qquad \begin{aligned} \tfrac{3}{2}x+\tfrac{2}{3}y &= \tfrac{17}{2} \\ \tfrac{3}{2}(3)+\tfrac{2}{3}(6) &\stackrel{?}{=} \tfrac{17}{2} \\ \tfrac{9}{2}+4 &\stackrel{?}{=} \tfrac{17}{2} \\ \tfrac{9}{2}+\tfrac{8}{2} &\stackrel{?}{=} \tfrac{17}{2} \\ \tfrac{17}{2} &= \tfrac{17}{2}\checkmark \end{aligned}$$

The solution is (3, 6).

Try It 5.3.14

Solve the system by elimination. $\begin{cases} \frac{1}{3}x-\frac{1}{2}y=1 \\ \frac{3}{4}x-y=\frac{5}{2} \end{cases}$

Answer

(6,2)

Try It 5.3.15

Solve the system by elimination. $\begin{cases} x+\frac{3}{5}y=-\frac{1}{5} \\ -\frac{1}{2}x-\frac{2}{3}y=\frac{5}{6} \end{cases}$

Answer

(1,−2)

In the Solving Systems of Equations by Graphing we saw that not all systems of linear equations have a single ordered pair as a solution. When the two equations were really the same line, there were infinitely many solutions. We called that a consistent system. When the two equations described parallel lines, there was no solution. We called that an inconsistent system.

Example 5.3.16

Solve the system by elimination. $\begin{cases} 3x+4y=12 \\ y=3-\frac{3}{4}x \end{cases}$

Solution

$$\begin{cases} 3x+4y=12 \\ y=3-\frac{3}{4}x \end{cases}$$

Write the second equation in standard form.

$$\begin{cases} 3x+4y=12 \\ \frac{3}{4}x+y=3 \end{cases}$$

Clear the fractions by multiplying thesecond equation by 4.

$$\begin{cases} 3x+4y=12 \\ 4\left(\frac{3}{4}x+y\right)=4(3) \end{cases}$$

Simplify.

$$\begin{cases} 3x+4y=12 \\ 3x+4y=12 \end{cases}$$

To eliminate a variable, we multiply thesecond equation by -1.

$$\begin{cases} 3x+4y=12 \\ \underline{-3x-4y=-12} \end{cases}$$

Simplify and add.

$$0=0$$

This is a true statement. The equations are consistent but dependent. Their graphs would be the same line. The system has infinitely many solutions.

After we cleared the fractions in the second equation, did you notice that the two equations were the same? That means we have coincident lines.

Try It 5.3.17

Solve the system by elimination. $\begin{cases} 5x-3y=15 \\ y=-5+\frac{5}{3}x \end{cases}$

Answer

infinitely many solutions

Try It 5.3.18

Solve the system by elimination. $\begin{cases} x+2y=6 \\ y=-\frac{1}{2}x+3 \end{cases}$

Answer

infinitely many solutions

Example 5.3.19

Solve the system by elimination. $\begin{cases} -6x+15y=10 \\ 2x-5y=-5 \end{cases}$

Solution

The equations are in standard form. $\begin{cases} -6x+15y=10 \\ 2x-5y=-5 \end{cases}$

Multiply the second equation by 3 to eliminate a variable. $\begin{cases} -6x+15y=10 \\ 3(2x-5y)=3(-5) \end{cases}$

Simplify and add. $\begin{cases} -6x+15y=10 \\ \underline{6x-15y=-15} \\ 0 \neq 5 \end{cases}$

This statement is false. The equations are inconsistent and so their graphs would be parallel lines.

The system does not have a solution.

Try It 5.3.20

Solve the system by elimination. $\begin{cases} -3x+2y=8 \\ 9x-6y=13 \end{cases}$

Answer

no solution

Try It 5.3.21

Solve the system by elimination. $\begin{cases} 7x-3y=-2 \\ -14x+6y=8 \end{cases}$

Answer

no solution

Solve Applications of Systems of Equations by Elimination

Some applications problems translate directly into equations in standard form, so we will use the elimination method to solve them. As before, we use our Problem Solving Strategy to help us stay focused and organized.

Example 5.3.22

The sum of two numbers is 39. Their difference is 9. Find the numbers.

Solution

Step 1. Read the problem	
Step 2. Identify what we are looking for.	We are looking for two numbers.
Step 3. Name what we are looking for.	Let n = the first number. m = the second number
Step 4. Translate into a system of equations.	
	The sum of two numbers is 39. $n+m=39$ Their difference is 9. $n-m=9$
The system is:	$\begin{cases} n+m=39 \\ n-m=9 \end{cases}$
Step 5. Solve the system of equations. To solve the system of equations, use elimination. The equations are in standard form and the coefficients of m are opposites. Add.	$\begin{cases} n+m=39 \\ \underline{n-m=9} \\ 2n \quad\quad =48 \end{cases}$
Solve for n.	$n=24$
Substitute n=24 into one of the original equations and solve form.	$n+m=39$ $24+m=39$ $m=15$
Step 6. Check the answer.	Since 24+15=39 and 24−15=9, the answers check.
Step 7. Answer the question.	The numbers are 24 and 15.

? Try It 5.3.23

The sum of two numbers is 42. Their difference is 8. Find the numbers.

Answer

The numbers are 25 and 17.

? Try It 5.3.24

The sum of two numbers is −15. Their difference is −35. Find the numbers.

Answer

The numbers are −25 and 10.

✓ Example 5.3.25

Joe stops at a burger restaurant every day on his way to work. Monday he had one order of medium fries and two small sodas, which had a total of 620 calories. Tuesday he had two orders of medium fries and one small soda, for a total of 820 calories. How many calories are there in one order of medium fries? How many calories in one small soda?

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for the number of calories in one order of medium fries and in one small soda.
Step 3. Name what we are looking for.	Let f = the number of calories in 1 order of medium fries. s = the number of calories in 1 small soda.
Step 4. Translate into a system of equations:	one medium fries and two small sodas had a total of 620 calories
	$f+2s=620$
	two medium fries and one small soda had a total of 820 calories.
	$2f+s=820$
Our system is:	$\begin{cases} f+2s=620 \\ 2f+s=820 \end{cases}$
Step 5. Solve the system of equations. To solve the system of equations, use elimination. The equations are in standard form. To get opposite coefficients of f, multiply the top equation by -2.	$\begin{cases} -2(f+2s)=-2(620) \\ 2f+s=820 \end{cases}$
Simplify and add.	$\begin{array}{r} -2f-4s=-1240 \\ 2f+s=820 \\ \hline -3s=-420 \end{array}$
Solve for s.	$s=140$
Substitute $s=140$ into one of the original equations and then solve for f.	$f+2s=620$
	$f+2\cdot 140=620$
	$f+280=620$
	$f=340$
Step 6. Check the answer.	Verify that these numbers make sense in the problem and that they are solutions to both equations. We leave this to you!
Step 7. Answer the question.	The small soda has 140 calories and the fries have 340 calories.

? Try It 5.3.26

Malik stops at the grocery store to buy a bag of diapers and 2 cans of formula. He spends a total of $37. The next week he stops and buys 2 bags of diapers and 5 cans of formula for a total of $87. How much does a bag of diapers cost? How much is one can of formula?

Answer

The bag of diapers costs $11 and the can of formula costs $13.

? Try It 5.3.27

To get her daily intake of fruit for the day, Sasha eats a banana and 8 strawberries on Wednesday for a calorie count of 145. On the following Wednesday, she eats two bananas and 5 strawberries for a total of 235 calories for the fruit. How many calories are there in a banana? How many calories are in a strawberry?

Answer

There are 105 calories in a banana and 5 calories in a strawberry.

Choose the Most Convenient Method to Solve a System of Linear Equations

When you will have to solve a system of linear equations in a later math class, you will usually not be told which method to use. You will need to make that decision yourself. So you'll want to choose the method that is easiest to do and minimizes your chance of making mistakes.

Graphing	Substitution	Elimination
Use when you need a picture of the situation.	Use when one equation is already solved for one variable.	Use when the equations are in standard form.

✓ Example 5.3.28

For each system of linear equations decide whether it would be more convenient to solve it by substitution or elimination. Explain your answer.

1. $\begin{cases} 3x+8y=40 \\ 7x-4y=-32 \end{cases}$
2. $\begin{cases} 5x+6y=12 \\ y=\frac{2}{3}x-1 \end{cases}$

Solution

1. $\begin{cases} 3x+8y=40 \\ 7x-4y=-32 \end{cases}$

Since both equations are in standard form, using elimination will be most convenient.

2. $\begin{cases} 5x+6y=12 \\ y=\frac{2}{3}x-1 \end{cases}$

Since one equation is already solved for y, using substitution will be most convenient.

? Try It 5.3.29

For each system of linear equations, decide whether it would be more convenient to solve it by substitution or elimination. Explain your answer.

1. $\begin{cases} 4x-5y=-32 \\ 3x+2y=-1 \end{cases}$
2. $\begin{cases} x=2y-1 \\ 3x-5y=-7 \end{cases}$

Answer

1. Since both equations are in standard form, using elimination will be most convenient.
2. Since one equation is already solved for xx, using substitution will be most convenient.

Try It 5.3.30

For each system of linear equations, decide whether it would be more convenient to solve it by substitution or elimination. Explain your answer.

1. $\begin{cases} y = 2x - 1 \\ 3x - 4y = -6 \end{cases}$
2. $\begin{cases} 6x - 2y = 12 \\ 3x + 7y = -13 \end{cases}$

Answer

1. Since one equation is already solved for y, using substitution will be most convenient;
2. Since both equations are in standard form, using elimination will be most convenient.

Note

Access these online resources for additional instruction and practice with solving systems of linear equations by elimination.

- Instructional Video-Solving Systems of Equations by Elimination
- Instructional Video-Solving by Elimination
- Instructional Video-Solving Systems by Elimination

Key Concepts

- **To Solve a System of Equations by Elimination**
  1. Write both equations in standard form. If any coefficients are fractions, clear them.
  2. Make the coefficients of one variable opposites.
     - Decide which variable you will eliminate.
     - Multiply one or both equations so that the coefficients of that variable are opposites.
  3. Add the equations resulting from Step 2 to eliminate one variable.
  4. Solve for the remaining variable.
  5. Substitute the solution from Step 4 into one of the original equations. Then solve for the other variable.
  6. Write the solution as an ordered pair.
  7. Check that the ordered pair is a solution to **both** original equations.

This page titled 5.3: Solve Systems of Equations by Elimination is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.3: Solve Systems of Equations by Elimination** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.3E: Exercises

Practice Makes Perfect

Solve a System of Equations by Elimination

In the following exercises, solve the systems of equations by elimination.

Exercise 5.3E.1

$$\begin{cases} 5x+2y=2 \\ -3x-y=0 \end{cases}$$

Exercise 5.3E.2

$$\begin{cases} -3x+y=-9 \\ x-2y=-12 \end{cases}$$

Answer

$(6,9)$

Exercise 5.3E.3

$$\begin{cases} 6x-5y=-1 \\ 2x+y=13 \end{cases}$$

Exercise 5.3E.4

$$\begin{cases} 3x-y=-7 \\ 4x+2y=-6 \end{cases}$$

Answer

$(-2,1)$

Exercise 5.3E.5

$$\begin{cases} x+y=-1 \\ x-y=-5 \end{cases}$$

Exercise 5.3E.6

$$\begin{cases} x+y=-8 \\ x-y=-6 \end{cases}$$

Answer

$(-7,-1)$

Exercise 5.3E.7

$$\begin{cases} 3x-2y=1 \\ -x+2y=9 \end{cases}$$

Exercise 5.3E.8

$$\begin{cases} -7x+6y=-10 \\ x-6y=22 \end{cases}$$

Answer

$(-2,-4)$

Exercise 5.3E.9

$$\begin{cases} 3x+2y=-3 \\ -x-2y=-19 \end{cases}$$

Exercise 5.3E.10

$$\begin{cases} 5x+2y=1 \\ -5x-4y=-7 \end{cases}$$

Answer

$(-1,3)$

Exercise 5.3E.11

$$\begin{cases} 6x+4y=-4 \\ -6x-5y=8 \end{cases}$$

Exercise 5.3E.12

$$\begin{cases} 3x-4y=-11 \\ x-2y=-5 \end{cases}$$

Answer

$(-1,2)$

Exercise 5.3E.13

$$\begin{cases} 5x-7y=29 \\ x+3y=-3 \end{cases}$$

Exercise 5.3E.14

$$\begin{cases} 6x-5y=-75 \\ -x-2y=-13 \end{cases}$$

Answer

$(-5,9)$

Exercise 5.3E.15

$$\begin{cases} -x+4y=8 \\ 3x+5y=10 \end{cases}$$

Exercise 5.3E.16

$$\begin{cases} 2x-5y=7 \\ 3x-y=17 \end{cases}$$

Answer

$(6,1)$

Exercise 5.3E.17

$$\begin{cases} 5x-3y=-1 \\ 2x-y=2 \end{cases}$$

Exercise 5.3E.18

$$\begin{cases} 7x+y=-4 \\ 13x+3y=4 \end{cases}$$

Answer

$(-2,10)$

Exercise 5.3E.19

$$\begin{cases} -3x+5y=-13 \\ 2x+y=-26 \end{cases}$$

Exercise 5.3E.20

$$\begin{cases} 3x-5y=-9 \\ 5x+2y=16 \end{cases}$$

Answer

$(2,3)$

Exercise 5.3E.21

$$\begin{cases} 4x-3y=3 \\ 2x+5y=-31 \end{cases}$$

Exercise 5.3E.22

$$\begin{cases} 4x+7y=14 \\ -2x+3y=32 \end{cases}$$

Answer

$(-7,6)$

Exercise 5.3E.23

$$\begin{cases} 5x+2y=21 \\ 7x-4y=9 \end{cases}$$

Exercise 5.3E.24

$$\begin{cases} 3x+8y=-3 \\ 2x+5y=-3 \end{cases}$$

Answer

$(-9,3)$

Exercise 5.3E.25

$$\begin{cases} 11x+9y=-5 \\ 7x+5y=-1 \end{cases}$$

Exercise 5.3E.26

$$\begin{cases} 3x+8y=67 \\ 5x+3y=60 \end{cases}$$

Answer

$(9,5)$

Exercise 5.3E.27

$$\begin{cases} 2x+9y=-4 \\ 3x+13y=-7 \end{cases}$$

Exercise 5.3E.28

$$\begin{cases} \frac{1}{3}x-y=-3 \\ x+\frac{5}{2}y=2 \end{cases}$$

Answer

$(-3,2)$

Exercise 5.3E.29

$$\begin{cases} x+\frac{1}{2}y=\frac{3}{2} \\ \frac{1}{5}x-\frac{1}{5}y=3 \end{cases}$$

Exercise 5.3E.30

$$\begin{cases} x+\frac{1}{3}y=-1 \\ \frac{1}{2}x-\frac{1}{3}y=-2 \end{cases}$$

Answer

$(-2,3)$

Exercise 5.3E.31

$$\begin{cases} \frac{1}{3}x - y = -3 \\ \frac{2}{3}x + \frac{5}{2}y = 3 \end{cases}$$

Exercise 5.3E.32

$$\begin{cases} 2x + y = 3 \\ 6x + 3y = 9 \end{cases}$$

Answer

infinitely many solutions

Exercise 5.3E.33

$$\begin{cases} x - 4y = -1 \\ -3x + 12y = 3 \end{cases}$$

Exercise 5.3E.34

$$\begin{cases} -3x - y = 8 \\ 6x + 2y = -16 \end{cases}$$

Answer

infinitely many solutions

Exercise 5.3E.35

$$\begin{cases} 4x + 3y = 2 \\ 20x + 15y = 10 \end{cases}$$

Exercise 5.3E.36

$$\begin{cases} 3x + 2y = 6 \\ -6x - 4y = -12 \end{cases}$$

Answer

infinitely many solutions

Exercise 5.3E.37

$$\begin{cases} 5x - 8y = 12 \\ 10x - 16y = 20 \end{cases}$$

Exercise 5.3E.38

$$\begin{cases} -11x + 12y = 60 \\ -22x + 24y = 90 \end{cases}$$

Answer

inconsistent, no solution

Exercise 5.3E.39

$$\begin{cases} 7x-9y=16 \\ -21x+27y=-24 \end{cases}$$

Exercise 5.3E.40

$$\begin{cases} 5x-3y=15 \\ y=\frac{5}{3}x-2 \end{cases}$$

Answer

inconsistent, no solution

Exercise 5.3E.41

$$\begin{cases} 2x+4y=7 \\ y=-\frac{1}{2}x-4 \end{cases}$$

Solve Applications of Systems of Equations by Elimination

In the following exercises, translate to a system of equations and solve.

Exercise 5.3E.42

The sum of two numbers is 65. Their difference is 25. Find the numbers.

Answer

The numbers are 20 and 45.

Exercise 5.3E.43

The sum of two numbers is 37. Their difference is 9. Find the numbers.

Exercise 5.3E.44

The sum of two numbers is −27. Their difference is −59. Find the numbers.

Answer

The numbers are 16 and −43.

Exercise 5.3E.45

The sum of two numbers is −45. Their difference is −89. Find the numbers.

Exercise 5.3E.46

Andrea is buying some new shirts and sweaters. She is able to buy 3 shirts and 2 sweaters for $114 or she is able to buy 2 shirts and 4 sweaters for $164. How much does a shirt cost? How much does a sweater cost?

Answer

A shirt costs $16 and a sweater costs $33.

Exercise 5.3E.47

Peter is buying office supplies. He is able to buy 3 packages of paper and 4 staplers for $40 or he is able to buy 5 packages of paper and 6 staplers for $62. How much does a package of paper cost? How much does a stapler cost?

Exercise 5.3E.48

The total amount of sodium in 2 hot dogs and 3 cups of cottage cheese is 4720 mg. The total amount of sodium in 5 hot dogs and 2 cups of cottage cheese is 6300 mg. How much sodium is in a hot dog? How much sodium is in a cup of cottage cheese?

Answer

There are 860 mg in a hot dog. There are 1,000 mg in a cup of cottage cheese.

Exercise 5.3E.49

The total number of calories in 2 hot dogs and 3 cups of cottage cheese is 960 calories. The total number of calories in 5 hot dogs and 2 cups of cottage cheese is 1190 calories. How many calories are in a hot dog? How many calories are in a cup of cottage cheese?

Choose the Most Convenient Method to Solve a System of Linear Equations

In the following exercises, decide whether it would be more convenient to solve the system of equations by substitution or elimination.

Exercise 5.3E.50

1. $\begin{cases} 8x-15y=-32 \\ 6x+3y=-5 \end{cases}$
2. $\begin{cases} x=4y-3 \\ 4x-2y=-6 \end{cases}$

Answer

1. elimination
2. substitution

Exercise 5.3E.51

1. $\begin{cases} y=7x-5 \\ 3x-2y=16 \end{cases}$
2. $\begin{cases} 12x-5y=-42 \\ 3x+7y=-15 \end{cases}$

Exercise 5.3E.52

1. $\begin{cases} y=4x+9 \\ 5x-2y=-21 \end{cases}$
2. $\begin{cases} 9x-4y=24 \\ 3x+5y=-14 \end{cases}$

Answer

1. substitution
2. elimination

Exercise 5.3E.53

1. $\begin{cases} 14x-15y=-30 \\ 7x+2y=10 \end{cases}$
2. $\begin{cases} x=9y-11 \\ 2x-7y=-27 \end{cases}$

Everyday Math

Exercise 5.3E.54

In one hour Norris can row 3 miles upstream against the current. In the same amount of time he can row 5 miles downstream, with the current. Solve the system. $\begin{cases} r-c=3 \\ r+c=5 \end{cases}$

1. for r, his rowing speed in still water.
2. Then solve for c, the speed of the river current.

Answer

1. r=4
2. c=1

Exercise 5.3E.55

Josie wants to make 10 pounds of trail mix using nuts and raisins, and she wants the total cost of the trail mix to be $54. Nuts cost $6 per pound and raisins cost $3 per pound. Solve the system $\begin{cases} n+r=10 \\ 6n+3r=54 \end{cases}$ to find n, the number of pounds of nuts, and rr, the number of pounds of raisins she should use.


Writing Exercises

Exercise 5.3E.56

Solve the system
$\begin{cases} x+y=10 \\ 5x+8y=56 \end{cases}$

1. by substitution
2. by graphing
3. Which method do you prefer? Why?

Answer

1. (8, 2)
2. 

3. Answers will vary.

https://math.libretexts.org/@go/page/30524

Exercise $5.3E.57$

Solve the system $\begin{cases} x+y=-12 \\ y=4-\frac{1}{2}x \end{cases}$

1. by substitution
2. by graphing
3. Which method do you prefer? Why?

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve a system of equations by elimination.			
solve applications of systems of equations by elimination.			
choose the most convenient method to solve a system of linear equations.			

b. What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 5.3E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.3E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.4: Solve Applications with Systems of Equations

Learning Objectives

By the end of this section, you will be able to:

- Translate to a system of equations
- Solve direct translation applications
- Solve geometry applications
- Solve uniform motion applications

Note

Before you get started, take this readiness quiz.

1. The sum of twice a number and nine is 31. Find the number.
   If you missed this problem, review Example 3.1.10.
2. Twins Jon and Ron together earned $96,000 last year. Ron earned $8,000 more than three times what Jon earned. How much did each of the twins earn?
   If you missed this problem, review Example 3.1.31.
3. Alessio rides his bike $3\frac{1}{2}$ hours at a rate of 10 miles per hour. How far did he ride?
   If you missed this problem, review Example 2.6.1.

Previously in this chapter we solved several applications with systems of linear equations. In this section, we'll look at some specific types of applications that relate two quantities. We'll translate the words into linear equations, decide which is the most convenient method to use, and then solve them.

We will use our Problem Solving Strategy for Systems of Linear Equations.

USE A PROBLEM SOLVING STRATEGY FOR SYSTEMS OF LINEAR EQUATIONS.

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose variables to represent those quantities.
4. **Translate** into a system of equations.
5. **Solve** the system of equations using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

Translate to a System of Equations

Many of the problems we solved in earlier applications related two quantities. Here are two of the examples from the chapter on Math Models.

- The sum of two numbers is negative fourteen. One number is four less than the other. Find the numbers.
- A married couple together earns $110,000 a year. The wife earns $16,000 less than twice what her husband earns. What does the husband earn?

In that chapter we translated each situation into one equation using only one variable. Sometimes it was a bit of a challenge figuring out how to name the two quantities, wasn't it?

Let's see how we can translate these two problems into a system of equations with two variables. We'll focus on Steps 1 through 4 of our Problem Solving Strategy.

Example 5.4.1: How to Translate to a System of Equations

Translate to a system of equations:

The sum of two numbers is negative fourteen. One number is four less than the other. Find the numbers.

Solution

Step 1. Read the problem. Make sure you understand all the words and ideas.	This is a number problem.	The sum of two numbers is negative fourteen. One number is four less than the other. Find the numbers.
Step 2. Identify what you are looking for.	"Find the numbers."	We are looking for 2 numbers.
Step 3. Name what you are looking for. Choose variables to represent those quantities.	We will use two variables, m and n.	Let m = one number n = second number
Step 4. Translate into a system of equations.	We will write one equation for each sentence.	The sum of the numbers ($m+n$) is ($=$) -14 (-14) One number (m) is ($=$) four less than the other ($n-4$) The system is: $\begin{cases} m+n=-14 \\ m=n-4 \end{cases}$

Try It 5.4.2

Translate to a system of equations:

The sum of two numbers is negative twenty-three. One number is 7 less than the other. Find the numbers.

Answer

$$\begin{cases} m+n=-23 \\ m=n-7 \end{cases}$$

Try It 5.4.3

Translate to a system of equations:

The sum of two numbers is negative eighteen. One number is 40 more than the other. Find the numbers.

Answer

$$\begin{cases} m+n=-18 \\ m=n+40 \end{cases}$$

We'll do another example where we stop after we write the system of equations.

Example 5.4.4

Translate to a system of equations:

A married couple together earns $110,000 a year. The wife earns $16,000 less than twice what her husband earns. What does the husband earn?

Solution

We are looking for the amount that the husband and wife each earn.	Let $h =$ the amount the husband earns.
	$w =$ the amount the wife earns
Translate.	A married couple together earns $110,000.
	$w + h = 110000$
	The wife earns $16,000 less than twice what husband earns.
	$w = 2h - 16,000$
The system of equations is:	$\begin{cases} w + h = 110,000 \\ w = 2h - 16,000 \end{cases}$

Try It 5.4.5

Translate to a system of equations:

A couple has a total household income of $84,000. The husband earns $18,000 less than twice what the wife earns. How much does the wife earn?

Answer

$$\begin{cases} w + h = 84,000 \\ h = 2w - 18,000 \end{cases}$$

Try It 5.4.6

Translate to a system of equations:

A senior employee makes $5 less than twice what a new employee makes per hour. Together they make $43 per hour. How much does each employee make per hour?

Answer

$$\begin{cases} s = 2n - 5 \\ s + n = 43 \end{cases}$$

Solve Direct Translation Applications

We set up, but did not solve, the systems of equations in Example 5.4.1 and Example 5.4.4 Now we'll translate a situation to a system of equations and then solve it.

Example 5.4.7

Translate to a system of equations and then solve:

Devon is 26 years older than his son Cooper. The sum of their ages is 50. Find their ages.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for the ages of Devon and Cooper.
Step 3. Name what we are looking for.	Let d= Devon's age. c= Cooper's age
Step 4. Translate into a system of equations.	Devon is 26 years older than Cooper.
	$d = c + 26$
	The sum of their ages is 50.

	$d + c = 50$
The system is:	$d = c + 26$ $d + c = 50$
Step 5. Solve the system of equations. Solve by substitution.	$d = c + 26$ $d + c = 50$ $d + c = 50$
Substitute $c + 26$ into the second equation.	$c + 26 + c = 50$
Solve for c.	$2c + 26 = 50$
	$2c = 24$
	$c = 12$ $d = c + 26$
Substitute $c = 12$ into the first equation and then solve for d.	$d = 12 + 26$
	$d = 38$
Step 6. Check the answer in the problem.	Is Devon's age 26 more than Cooper's? Yes, 38 is 26 more than 12. Is the sum of their ages 50? Yes, 38 plus 12 is 50.
Step 7. Answer the question.	Devon is 38 and Cooper is 12 years old.

Try It 5.4.8

Translate to a system of equations and then solve:

Ali is 12 years older than his youngest sister, Jameela. The sum of their ages is 40. Find their ages.

Answer

Ali is 26 and Jameela is 14.

Try It 5.4.9

Translate to a system of equations and then solve:

Jake's dad is 6 more than 3 times Jake's age. The sum of their ages is 42. Find their ages.

Answer

Jake is 9 and his dad is 33.

Example 5.4.10

Translate to a system of equations and then solve:

When Jenna spent 10 minutes on the elliptical trainer and then did circuit training for 20 minutes, her fitness app says she burned 278 calories. When she spent 20 minutes on the elliptical trainer and 30 minutes circuit training she burned 473 calories. How many calories does she burn for each minute on the elliptical trainer? How many calories does she burn for each minute of circuit training?

Solution

Step 1. Read the problem.	

Step 2. Identify what we are looking for.	We are looking for the number of calories burned each minute on the elliptical trainer and each minute of circuit training.
Step 3. Name what we are looking for.	Let e= number of calories burned per minute on the elliptical trainer. c= number of calories burned per minute while circuit training
Step 4. Translate into a system of equations.	10 minutes on the elliptical and circuit training for 20 minutes, burned 278 calories
	$10e + 20c = 278$
	20 minutes on the elliptical and 30 minutes of circuit training burned 473 calories
	$20e + 30c = 473$
The system is:	$\begin{aligned} 10e + 20c &= 278 \\ 20e + 30c &= 473 \end{aligned}$
Step 5. Solve the system of equations.	
Multiply the first equation by −2 to get opposite coefficients of e.	$\begin{aligned} -2(10e + 20c) &= -2(278) \\ 20e + 30c &= 473 \end{aligned}$
Simplify and add the equations. Solve for c.	$\begin{aligned} -20e - 40c &= -556 \\ 20e + 30c &= 473 \\ \hline -10c &= -83 \\ c &= 8.3 \end{aligned}$
Substitute $c = 8.3$ into one of the original equations to solve for e.	$10e + 20c = 278$
	$10e + 20(8.3) = 278$
	$10e + 166 = 278$
	$10e = 112$
	$e = 11.2$
Step 6. Check the answer in the problem.	Check the math on your own.
$\begin{aligned} 10(11.2) + 20(8.3) &\stackrel{?}{=} 278 \\ 20(11.2) + 30(8.3) &\stackrel{?}{=} 473 \end{aligned}$	
Step 7. Answer the question.	Jenna burns 8.3 calories per minute circuit training and 11.2 calories per minute while on the elliptical trainer.

Try It 5.4.11

Translate to a system of equations and then solve:

Mark went to the gym and did 40 minutes of Bikram hot yoga and 10 minutes of jumping jacks. He burned 510 calories. The next time he went to the gym, he did 30 minutes of Bikram hot yoga and 20 minutes of jumping jacks burning 470 calories. How many calories were burned for each minute of yoga? How many calories were burned for each minute of jumping jacks?

Answer

Mark burned 11 calories for each minute of yoga and 7 calories for each minute of jumping jacks.

Try It 5.4.12

Translate to a system of equations and then solve:

Erin spent 30 minutes on the rowing machine and 20 minutes lifting weights at the gym and burned 430 calories. During her next visit to the gym she spent 50 minutes on the rowing machine and 10 minutes lifting weights and burned 600 calories. How many calories did she burn for each minutes on the rowing machine? How many calories did she burn for each minute of weight lifting?

Answer

Erin burned 11 calories for each minute on the rowing machine and 5 calories for each minute of weight lifting.

Solve Geometry Applications

When we learned about Math Models, we solved geometry applications using properties of triangles and rectangles. Now we'll add to our list some properties of angles.

The measures of two **complementary angles** add to 90 degrees. The measures of two **supplementary angles** add to 180 degrees.

COMPLEMENTARY AND SUPPLEMENTARY ANGLES

Two angles are **complementary** if the sum of the measures of their angles is 90 degrees.

Two angles are **supplementary** if the sum of the measures of their angles is 180 degrees.

If two angles are complementary, we say that *one angle is the complement of the other.*

If two angles are supplementary, we say that *one angle is the supplement of the other.*

Example 5.4.13

Translate to a system of equations and then solve:

The difference of two complementary angles is 26 degrees. Find the measures of the angles.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for the measure of each angle.
Step 3. Name what we are looking for.	Let x = the measure of the first angle. y = the measure of the second angle
Step 4. Translate into a system of equations.	The angles are complementary. $x+y=90$ The difference of the two angles is 26 degrees. $x-y=26$
The system is	$\begin{cases} x+y=90 \\ x-y=26 \end{cases}$
Step 5. Solve the system of equations by elimination.	$\begin{cases} x+y=90 \\ x-y=26 \end{cases}$ $2x = 116$ $x = 58$
Substitute $x=58$ into the first equation.	$x+y = 90$ $58+y = 90$ $y = 32$

Step 6. Check the answer in the problem. $58+32=90$✓ $58-32=36$✓	
Step 7. Answer the question.	The angle measures are 58 degrees and 32 degrees.

? Try It 5.4.14

Translate to a system of equations and then solve:

The difference of two complementary angles is 20 degrees. Find the measures of the angles.

Answer

The angle measures are 55 degrees and 35 degrees.

? Try It 5.4.15

Translate to a system of equations and then solve:

The difference of two complementary angles is 80 degrees. Find the measures of the angles.

Answer

The angle measures are 5 degrees and 85 degrees.

✓ Example 5.4.16

Translate to a system of equations and then solve:

Two angles are supplementary. The measure of the larger angle is twelve degrees less than five times the measure of the smaller angle. Find the measures of both angles.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for the measure of each angle.
Step 3. Name what we are looking for.	Let x= the measure of the first angle. y= the measure of the second angle
Step 4. Translate into a system of equations.	The angles are supplementary.
	$x+y=180$
	The larger angle is twelve less than five times the smaller angle
	$y=5x-12$
The system is: **Step 5. Solve** the system of equations substitution.	$x+y=180$ $y=5x-12$ $x+y=180$
Substitute $5x-12$ for y in the first equation.	$x+5x-12=180$
Solve for x.	$6x-12=180$
	$6x=192$

	$x = 32$ $y = 5x - 12$
Substitute 32 for in the second equation, then solve for y.	$y = 5 \cdot 32 - 12$
	$y = 160 - 12$
	$y = 148$
Step 6. Check the answer in the problem. $32 + 158 = 180$✓ $5 \cdot 32 - 12 = 148$✓	
Step 7. Answer the question.	The angle measures are 148 and 32.

Try It 5.4.17

Translate to a system of equations and then solve:

Two angles are supplementary. The measure of the larger angle is 12 degrees more than three times the smaller angle. Find the measures of the angles.

Answer

The angle measures are 42 degrees and 138 degrees.

Try It 5.4.18

Translate to a system of equations and then solve:

Two angles are supplementary. The measure of the larger angle is 18 less than twice the measure of the smaller angle. Find the measures of the angles.

Answer


The angle measures are 66 degrees and 114 degrees.

Example 5.4.19

Translate to a system of equations and then solve:

Randall has 125 feet of fencing to enclose the rectangular part of his backyard adjacent to his house. He will only need to fence around three sides, because the fourth side will be the wall of the house. He wants the length of the fenced yard (parallel to the house wall) to be 5 feet more than four times as long as the width. Find the length and the width.

Solution

Step 1. Read the problem.	
Step 2. Identify what you are looking for.	We are looking for the length and width.
	

Step 3. Name what we are looking for.	Let L= the length of the fenced yard. W= the width of the fenced yard
Step 4. Translate into a system of equations.	One length and two widths equal 125.
	$L+2W=125$
	The length will be 5 feet more than four times the width.
	$L=4W+5$
The system is: **Step 5. Solve** the system of equations by substitution.	$L+2W=125$ $L=4W+5$ $L+2W=125$
Substitute $L = 4W + 5$ into the first equation, then solve for W.	$4W+5+2W=125$
	$6W+5=125$
	$6W=120$
Substitute 20 for W in the second equation, then solve for L.	$W=20$ $L=4W+5$
	$L=4\cdot 20+5$
	$L=80+5$
	$L=85$
Step 6. Check the answer in the problem. $20+85+20 = 125\checkmark$ $85 = 4\cdot 20+5\checkmark$	
Step 7. Answer the equation.	The length is 85 feet and the width is 20 feet.

Try It 5.4.20

Translate to a system of equations and then solve:

Mario wants to put a rectangular fence around the pool in his backyard. Since one side is adjacent to the house, he will only need to fence three sides. There are two long sides and the one shorter side is parallel to the house. He needs 155 feet of fencing to enclose the pool. The length of the long side is 10 feet less than twice the width. Find the length and width of the pool area to be enclosed.

Answer

The length is 60 feet and the width is 35 feet.

Try It 5.4.21

Translate to a system of equations and then solve:

Alexis wants to build a rectangular dog run in her yard adjacent to her neighbor's fence. She will use 136 feet of fencing to completely enclose the rectangular dog run. The length of the dog run along the neighbor's fence will be 16 feet less than twice the width. Find the length and width of the dog run.

Answer

The length is 60 feet and the width is 38 feet.

Solve Uniform Motion Applications

We used a table to organize the information in uniform motion problems when we introduced them earlier. We'll continue using the table here. The basic equation was $D = rt$ where D is the distance traveled, r is the rate, and t is the time.

Our first example of a uniform motion application will be for a situation similar to some we have already seen, but now we can use two variables and two equations.


Example 5.4.22

Translate to a system of equations and then solve:

Joni left St. Louis on the interstate, driving west towards Denver at a speed of 65 miles per hour. Half an hour later, Kelly left St. Louis on the same route as Joni, driving 78 miles per hour. How long will it take Kelly to catch up to Joni?

Solution

A diagram is useful in helping us visualize the situation.



Identify and name what we are looking for. A chart will help us organize the data. We know the rates of both Joni and Kelly, and so we enter them in the chart.	
We are looking for the length of time Kelly, k, and Joni, j, will each drive. Since D=r·t we can fill in the Distance column.	

Type	Rate	• Time	= Distance
Joni	65	j	$65j$
Kelly	78	k	$78k$

Translate into a system of equations.

To make the system of equations, we must recognize that Kelly and Joni will drive the same distance. So, 65j=78k.

Also, since Kelly left later, her time will be $\frac{1}{2}$ hour less than Joni's time.

So, $k = j - \frac{1}{2}$.

Now we have the system.	$\begin{aligned} k &= j - \frac{1}{2} \\ 65j &= 78k \end{aligned}$
Solve the system of equations by substitution.	$65j = 78k$
Substitute k=j−12 into the second equation, then solve for j.	$65j = 78\left(j - \frac{1}{2}\right)$
	$65j = 78j - 39$
	$-13j = -39$
	$j = 3$
To find Kelly's time, substitute j = 3 into the first equation, then solve for k.	$k = j - \frac{1}{2}$
	$k = 3 - \frac{1}{2}$

	$k=\frac{5}{2}$ or $k=2\frac{1}{2}$
Check the answer in the problem. Joni 3 hours (65 mph) = 195 miles. Kelly $2\frac{1}{2}$ hours (78 mph) = 195 miles. Yes, they will have traveled the same distance when they meet.	
Answer the question.	Kelly will catch up to Joni in $2\frac{1}{2}$ hours. By then, Joni will have traveled 3 hours.

Try It 5.4.23

Translate to a system of equations and then solve: Mitchell left Detroit on the interstate driving south towards Orlando at a speed of 60 miles per hour. Clark left Detroit 1 hour later traveling at a speed of 75 miles per hour, following the same route as Mitchell. How long will it take Clark to catch Mitchell?

Answer

It will take Clark 4 hours to catch Mitchell.

Try It 5.4.24

Translate to a system of equations and then solve: Charlie left his mother's house traveling at an average speed of 36 miles per hour. His sister Sally left 15 minutes (1/4 hour) later traveling the same route at an average speed of 42 miles per hour. How long before Sally catches up to Charlie?

Answer

It will take Sally $1\frac{1}{2}$ hours to catch up to Charlie.

Many real-world applications of uniform motion arise because of the effects of currents—of water or air—on the actual speed of a vehicle. Cross-country airplane flights in the United States generally take longer going west than going east because of the prevailing wind currents.

Let's take a look at a boat traveling on a river. Depending on which way the boat is going, the current of the water is either slowing it down or speeding it up.

Figure 5.4.1 and Figure 5.4.2 show how a river current affects the speed at which a boat is actually traveling. We'll call the speed of the boat in still water b and the speed of the river current c.

In Figure 5.4.1 the boat is going downstream, in the same direction as the river current. The current helps push the boat, so the boat's actual speed is faster than its speed in still water. The actual speed at which the boat is moving is $b + c$.




Figure 5.4.1

In Figure 5.4.2 the boat is going upstream, opposite to the river current. The current is going against the boat, so the boat's actual speed is slower than its speed in still water. The actual speed of the boat is b−c.




Figure 5.4.2

We'll put some numbers to this situation in Example 5.4.25

Example 5.4.25


Translate to a system of equations and then solve:

A river cruise ship sailed 60 miles downstream for 4 hours and then took 5 hours sailing upstream to return to the dock. Find the speed of the ship in still water and the speed of the river current.

Solution

Read the problem.

This is a uniform motion problem and a picture will help us visualize the situation.



Identify what we are looking for.	We are looking for the speed of the ship in still water and the speed of the current.
Name what we are looking for.	Let s=s= the rate of the ship in still water. c=c= the rate of the current

A chart will help us organize the information. The ship goes downstream and then upstream. Going downstream, the current helps the ship; therefore, the ship's actual rate is $s + c$. Going upstream, the current slows the ship; therefore, the actual rate is $s - c$.	
Downstream it takes 4 hours. Upstream it takes 5 hours. Each way the distance is 60 miles.	
Translate into a system of equations. Since rate times time is distance, we can write the system of equations.	$\begin{cases} 4(s+c)=60 \\ 5(s-c)=60 \end{cases}$
Solve the system of equations. Distribute to put both equations in standard form, then solve by elimination.	$\begin{cases} 4s+4c=60 \\ 5s-5c=60 \end{cases}$
Multiply the top equation by 5 and the bottom equation by 4. Add the equations, then solve for s.	$\begin{array}{r} 20s+20c=300 \\ 20s-20c=240 \\ \hline 40s \quad\quad =540 \end{array}$
Substitute $s = 13.5$ into one of the original equations.	$s = 13.5$ $4(s+c)=60$
	$4(13.5+c)=60$
	$54+4c=60$
	$4c=6$
	$c=1.5$
Check the answer in the problem. The downstream rate would be 13.5 + 1.5 = 15 mph. In 4 hours the ship would travel 15 · 4 = 60 miles. The upstream rate would be 13.5 − 1.5 = 12 mph. In 5 hours the ship would travel 12 · 5 = 60 miles.	
Answer the question.	The rate of the ship is 13.5 mph and the rate of the current is 1.5 mph.

	Rate	• Time =	Distance
downstream	$s+c$	4	60
upstream	$s-c$	5	60

? Try It 5.4.26

Translate to a system of equations and then solve: A Mississippi river boat cruise sailed 120 miles upstream for 12 hours and then took 10 hours to return to the dock. Find the speed of the river boat in still water and the speed of the river current.

Answer

The rate of the boat is 11 mph and the rate of the current is 1 mph.

Try It 5.4.27

Translate to a system of equations and then solve: Jason paddled his canoe 24 miles upstream for 4 hours. It took him 3 hours to paddle back. Find the speed of the canoe in still water and the speed of the river current.

Answer

The speed of the canoe is 7 mph and the speed of the current is 1 mph.

Wind currents affect airplane speeds in the same way as water currents affect boat speeds. We'll see this in Example 5.4.28 A wind current in the same direction as the plane is flying is called a *tailwind*. A wind current blowing against the direction of the plane is called a *headwind*.

Example 5.4.28


Translate to a system of equations and then solve:

A private jet can fly 1095 miles in three hours with a tailwind but only 987 miles in three hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Solution

Read the problem.

This is a uniform motion problem and a picture will help us visualize.



Identify what we are looking for.	We are looking for the speed of the jet in still air and the speed of the wind.
Name what we are looking for.	Let j= the speed of the jet in still air. w= the speed of the wind
A chart will help us organize the information. The jet makes two trips-one in a tailwind and one in a headwind. In a tailwind, the wind helps the jet and so the rate is $j + w$. In a headwind, the wind slows the jet and so the rate is $j - w$.	

	Rate • Time = Distance		
tailwind	$j + w$	3	1095
headwind	$j - w$	3	987

Each trip takes 3 hours. In a tailwind the jet flies 1095 miles. In a headwind the jet flies 987 miles.	
Translate into a system of equations. Since rate times time is distance, we get the system of equations.	$\begin{aligned} 3(j+w) &= 1095 \\ 3(j-w) &= 987 \end{aligned}$
Solve the system of equations. Distribute, then solve by elimination.	$\begin{aligned} 3j + 3w &= 1095 \\ 3j - 3w &= 987 \\ \hline 6j &= 2082 \end{aligned}$

Add, and solve for j. Substitute $j = 347$ into one of the original equations, then solve for w.	$j = 347$ $3(j + w) = 1095$
	$3(347 + w) = 1095$
	$1041 + 3w = 1095$
	$3w = 54$
	$w = 18$
Check the answer in the problem. With the tailwind, the actual rate of the jet would be $347 + 18 = 365$ mph. In 3 hours the jet would travel $365 \cdot 3 = 1095$ miles. Going into the headwind, the jet's actual rate would be $347 - 18 = 329$ mph. In 3 hours the jet would travel $329 \cdot 3 = 987$ miles.	
Answer the question.	The rate of the jet is 347 mph and the rate of the wind is 18 mph.

? Try It 5.4.29

Translate to a system of equations and then solve: A small jet can fly 1,325 miles in 5 hours with a tailwind but only 1025 miles in 5 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Answer

The speed of the jet is 235 mph and the speed of the wind is 30 mph.

? Try It 5.4.30

Translate to a system of equations and then solve: A commercial jet can fly 1728 miles in 4 hours with a tailwind but only 1536 miles in 4 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Answer

The speed of the jet is 408 mph and the speed of the wind is 24 mph.

Glossary

complementary angles

Two angles are complementary if the sum of the measures of their angles is 90 degrees.

supplementary angles

Two angles are supplementary if the sum of the measures of their angles is 180 degrees.

This page titled 5.4: Solve Applications with Systems of Equations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.4: Solve Applications with Systems of Equations** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.4E: Exercises

Practice Makes Perfect

Translate to a System of Equations

In the following exercises, translate to a system of equations and solve the system.

Exercise 5.4E.1

The sum of two numbers is fifteen. One number is three less than the other. Find the numbers.

Answer

The numbers are 6 and 9.

Exercise 5.4E.2

The sum of two numbers is twenty-five. One number is five less than the other. Find the numbers.

Exercise 5.4E.3

The sum of two numbers is negative thirty. One number is five times the other. Find the numbers.

Answer

The numbers are −5 and −25.

Exercise 5.4E.4

The sum of two numbers is negative sixteen. One number is seven times the other. Find the numbers.

Exercise 5.4E.5

Twice a number plus three times a second number is twenty-two. Three times the first number plus four times the second is thirty-one. Find the numbers.

Answer

The numbers are 5 and 4.

Exercise 5.4E.6

Six times a number plus twice a second number is four. Twice the first number plus four times the second number is eighteen. Find the numbers.

Exercise 5.4E.7

Three times a number plus three times a second number is fifteen. Four times the first plus twice the second number is fourteen. Find the numbers.

Answer

The numbers are 2 and 3.

Exercise 5.4E.8

Twice a number plus three times a second number is negative one. The first number plus four times the second number is two. Find the numbers.

Exercise 5.4E.9

A married couple together earn $75,000. The husband earns $15,000 more than five times what his wife earns. What does the wife earn?

Answer

$10,000

Exercise 5.4E.10

During two years in college, a student earned $9,500. The second year she earned $500 more than twice the amount she earned the first year. How much did she earn the first year?

Exercise 5.4E.11

Daniela invested a total of $50,000, some in a certificate of deposit (CD) and the remainder in bonds. The amount invested in bonds was $5000 more than twice the amount she put into the CD. How much did she invest in each account?

Answer

She put $15,000 into a CD and $35,000 in bonds.

Exercise 5.4E.12

Jorge invested $28,000 into two accounts. The amount he put in his money market account was $2,000 less than twice what he put into a CD. How much did he invest in each account?

Exercise 5.4E.13

In her last two years in college, Marlene received $42,000 in loans. The first year she received a loan that was $6,000 less than three times the amount of the second year's loan. What was the amount of her loan for each year?

Answer

The amount of the first year's loan was $30,000 and the amount of the second year's loan was $12,000.

Exercise 5.4E.14

Jen and David owe $22,000 in loans for their two cars. The amount of the loan for Jen's car is $2000 less than twice the amount of the loan for David's car. How much is each car loan?

Solve Direct Translation Applications

In the following exercises, translate to a system of equations and solve.

Exercise 5.4E.15

Alyssa is twelve years older than her sister, Bethany. The sum of their ages is forty-four. Find their ages.

Answer

Bethany is 16 years old and Alyssa is 28 years old.

Exercise 5.4E.16

Robert is 15 years older than his sister, Helen. The sum of their ages is sixty-three. Find their ages.

Exercise 5.4E.17

The age of Noelle's dad is six less than three times Noelle's age. The sum of their ages is seventy-four. Find their ages.

Answer

Noelle is 20 years old and her dad is 54 years old.

Exercise 5.4E.18

The age of Mark's dad is 4 less than twice Marks's age. The sum of their ages is ninety-five. Find their ages.

Exercise 5.4E.19

Two containers of gasoline hold a total of fifty gallons. The big container can hold ten gallons less than twice the small container. How many gallons does each container hold?

Answer

The small container holds 20 gallons and the large container holds 30 gallons.

Exercise 5.4E.20

June needs 48 gallons of punch for a party and has two different coolers to carry it in. The bigger cooler is five times as large as the smaller cooler. How many gallons can each cooler hold?

Exercise 5.4E.21

Shelly spent 10 minutes jogging and 20 minutes cycling and burned 300 calories. The next day, Shelly swapped times, doing 20 minutes of jogging and 10 minutes of cycling and burned the same number of calories. How many calories were burned for each minute of jogging and how many for each minute of cycling?

Answer

There were 10 calories burned jogging and 10 calories burned cycling.

Exercise 5.4E.22

Drew burned 1800 calories Friday playing one hour of basketball and canoeing for two hours. Saturday he spent two hours playing basketball and three hours canoeing and burned 3200 calories. How many calories did he burn per hour when playing basketball?

Exercise 5.4E.23

Troy and Lisa were shopping for school supplies. Each purchased different quantities of the same notebook and thumb drive. Troy bought four notebooks and five thumb drives for $116. Lisa bought two notebooks and three thumb dives for $68. Find the cost of each notebook and each thumb drive.

Answer

Notebooks are $4 and thumb drives are $20.

Exercise 5.4E.24

Nancy bought seven pounds of oranges and three pounds of bananas for $17. Her husband later bought three pounds of oranges and six pounds of bananas for $12. What was the cost per pound of the oranges and the bananas?

Solve Geometry Applications In the following exercises, translate to a system of equations and solve.

Exercise 5.4E.25

The difference of two complementary angles is 30 degrees. Find the measures of the angles.

Answer

The measures are 60 degrees and 30 degrees.

Exercise 5.4E.26

The difference of two complementary angles is 68 degrees. Find the measures of the angles.

Exercise 5.4E.27

The difference of two supplementary angles is 70 degrees. Find the measures of the angles.

Answer

The measures are 125 degrees and 55 degrees.

Exercise 5.4E.28

The difference of two supplementary angles is 24 degrees. Find the measure of the angles.

Exercise 5.4E.29

The difference of two supplementary angles is 8 degrees. Find the measures of the angles.

Answer

94 degrees and 86 degrees

Exercise 5.4E.30

The difference of two supplementary angles is 88 degrees. Find the measures of the angles.

Exercise 5.4E.31

The difference of two complementary angles is 55 degrees. Find the measures of the angles.

Answer

72.5 degrees and 17.5 degrees

Exercise 5.4E.32

The difference of two complementary angles is 17 degrees. Find the measures of the angles.

Exercise 5.4E.33

Two angles are supplementary. The measure of the larger angle is four more than three times the measure of the smaller angle. Find the measures of both angles.

Answer

The measures are 44 degrees and 136 degrees.

Exercise 5.4E.34

Two angles are supplementary. The measure of the larger angle is five less than four times the measure of the smaller angle. Find the measures of both angles.

Exercise 5.4E.35

Two angles are complementary. The measure of the larger angle is twelve less than twice the measure of the smaller angle. Find the measures of both angles.

Answer

The measures are 34 degrees and 56 degrees.

Exercise 5.4E.36

Two angles are complementary. The measure of the larger angle is ten more than four times the measure of the smaller angle. Find the measures of both angles.

Exercise 5.4E.37

Wayne is hanging a string of lights 45 feet long around the three sides of his rectangular patio, which is adjacent to his house. The length of his patio, the side along the house, is five feet longer than twice its width. Find the length and width of the patio.

Answer

The width is 10 feet and the length is 25 feet.

Exercise 5.4E.38

Darrin is hanging 200 feet of Christmas garland on the three sides of fencing that enclose his rectangular front yard. The length, the side along the house, is five feet less than three times the width. Find the length and width of the fencing.

Exercise 5.4E.39

A frame around a rectangular family portrait has a perimeter of 60 inches. The length is fifteen less than twice the width. Find the length and width of the frame.

Answer

The width is 15 feet and the length is 15 feet.

Exercise 5.4E.40

The perimeter of a rectangular toddler play area is 100 feet. The length is ten more than three times the width. Find the length and width of the play area.

Solve Uniform Motion Applications In the following exercises, translate to a system of equations and solve.

Exercise 5.4E.41

Sarah left Minneapolis heading east on the interstate at a speed of 60 mph. Her sister followed her on the same route, leaving two hours later and driving at a rate of 70 mph. How long will it take for Sarah's sister to catch up to Sarah?

Answer

It took Sarah's sister 12 hours.

Exercise 5.4E.42

College roommates John and David were driving home to the same town for the holidays. John drove 55 mph, and David, who left an hour later, drove 60 mph. How long will it take for David to catch up to John?

Exercise 5.4E.43

At the end of spring break, Lucy left the beach and drove back towards home, driving at a rate of 40 mph. Lucy's friend left the beach for home 30 minutes (half an hour) later, and drove 50 mph. How long did it take Lucy's friend to catch up to Lucy?

Answer

It took Lucy's friend 2 hours.

Exercise 5.4E.44

Felecia left her home to visit her daughter driving 45 mph. Her husband waited for the dog sitter to arrive and left home twenty minutes (1/3 hour) later. He drove 55 mph to catch up to Felecia. How long before he reaches her?

Exercise 5.4E.45

The Jones family took a 12 mile canoe ride down the Indian River in two hours. After lunch, the return trip back up the river took three hours. Find the rate of the canoe in still water and the rate of the current.

Answer

The canoe rate is 5 mph and the current rate is 1 mph.

Exercise 5.4E.46

A motor boat travels 60 miles down a river in three hours but takes five hours to return upstream. Find the rate of the boat in still water and the rate of the current.

Exercise 5.4E.47

A motor boat traveled 18 miles down a river in two hours but going back upstream, it took 4.5 hours due to the current. Find the rate of the motor boat in still water and the rate of the current.

Answer

The boat rate is 6.5 mph and the current rate is 2.5 mph.

Exercise 5.4E.48

A river cruise boat sailed 80 miles down the Mississippi River for four hours. It took five hours to return. Find the rate of the cruise boat in still water and the rate of the current.

Exercise 5.4E.49

A small jet can fly 1,072 miles in 4 hours with a tailwind but only 848 miles in 4 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Answer

The jet rate is 240 mph and the wind speed is 28 mph.

Exercise 5.4E.50

A small jet can fly 1,435 miles in 5 hours with a tailwind but only 1215 miles in 5 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Exercise 5.4E.51

A commercial jet can fly 868 miles in 2 hours with a tailwind but only 792 miles in 2 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Answer

The jet rate is 415 mph and the wind speed is 19 mph.

Exercise 5.4E.52

A commercial jet can fly 1,320 miles in 3 hours with a tailwind but only 1,170 miles in 3 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Everyday Math

Exercise 5.4E.53

At a school concert, 425 tickets were sold. Student tickets cost $5 each and adult tickets cost $8 each. The total receipts for the concert were $2,851. Solve the system

$$\begin{cases} s+a=425 \\ 5s+8a=2,851 \end{cases}$$

to find s, the number of student tickets and aa, the number of adult tickets.

Answer

s=183,a=242

Exercise 5.4E.54

The first graders at one school went on a field trip to the zoo. The total number of children and adults who went on the field trip was 115. The number of adults was $\frac{1}{4}$ the number of children. Solve the system

$$\begin{cases} c+a=115 \\ a=\frac{1}{4}c \end{cases}$$

to find c, the number of children and aa, the number of adults.

Writing Exercises

? Exercise $5.4E.55$

Write an application problem similar to Example using the ages of two of your friends or family members. Then translate to a system of equations and solve it.

Answer

Answers will vary.

? Exercise $5.4E.56$

Write a uniform motion problem similar to Example that relates to where you live with your friends or family members. Then translate to a system of equations and solve it.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
translate to a system of equations.			
solve direct translation applications.			
solve geometry applications.			
solve uniform motion applications.			

b. On a scale of 1-10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

More Practice: Number Problems

? Exercise $5.4E.57$

Set up a linear system and solve.

1. The sum of two integers is 45. The larger integer is 3 less than twice the smaller. Find the two integers.

2. The sum of two integers is 126. The larger is 18 less than 5 times the smaller. Find the two integers.

3. The sum of two integers is 41. When 3 times the smaller is subtracted from the larger the result is 17. Find the two integers.

4. The sum of two integers is 46. When the larger is subtracted from twice the smaller the result is 2. Find the two integers.

5. The difference of two integers is 11. When twice the larger is subtracted from 3 times the smaller, the result is 3. Find the integers.

6. The difference of two integers is 6. The sum of twice the smaller and the larger is 72. Find the integers.

7. The sum of 3 times a larger integer and 2 times a smaller is 15. When 3 times the smaller integer is subtracted from twice the larger, the result is 23. Find the integers.

8. The sum of twice a larger integer and 3 times a smaller is 10. When the 4 times the smaller integer is added to the larger, the result is 0. Find the integers.

9. The difference of twice a smaller integer and 7 times a larger is 4. When 5 times the larger integer is subtracted from 3 times the smaller, the result is -5. Find the integers.

10. The difference of a smaller integer and twice a larger is 0. When 3 times the larger integer is subtracted from 2 times the smaller, the result is -5. Find the integers.

Answer

1. The integers are 16 and 29.
3. The integers are 6 and 35.
5. The integers are 25 and 36.
7. The integers are -3 and 7.
9. The integers are -5 and -2.


More Practice: Geometry Problems

? Exercise 5.4E. 58


1. The length of a rectangle is 5 more than twice its width. If the perimeter measures 46 meters, then find the dimensions of the rectangle.

2. The width of a rectangle is 2 centimeters less than one-half its length. If the perimeter measures 62 centimeters, then find the dimensions of the rectangle.

3. A partitioned rectangular pen next to a river is constructed with a total 136 feet of fencing (see illustration). If the outer fencing measures 114 feet, then find the dimensions of the pen.



4. A partitioned rectangular pen is constructed with a total 168 feet of fencing (see illustration). the perimeter measures 138 feet, then find the dimensions of the pen.



Add exercises text here.

Answer

1. Length: 17 meters; width: 6 meters
3. Width: 22 feet; length: 70 feet.

More Practice: Distance Problems

? Exercise 5.4E. 59

Set up a linear system and solve.

1. The two legs of a 432-mile trip took 8 hours. The average speed for the first leg of the trip was 52 miles per hour and the average speed for the second leg of the trip was 60 miles per hour. How long did each leg of the trip take?
2. Jerry took two buses on the 265-mile trip from Los Angeles to Las Vegas. The first bus averaged 55 miles per hour and the second bus was able to average 50 miles per hour. If the total trip took 5 hours, then how long was spent in each bus?
3. An executive was able to average 48 miles per hour to the airport in her car and then board an airplane that averaged 210 miles per hour. The 549-mile business trip took 3 hours. How long did it take her to drive to the airport?
4. Joe spends 1 hour each morning exercising by jogging and then cycling for a total of 15 miles. He is able to average 6 miles per hour jogging and 18 miles per hour cycling. How long does he spend jogging each morning?
5. Swimming with the current Jack can swim 2.5 miles in $\frac{1}{2}$ hour. Swimming back, against the same current, he can only swim 2 miles in the same amount of time. How fast is the current?
6. A light aircraft flying with the wind can travel 180 miles in $1\frac{1}{2}$ hours. The aircraft can fly the same distance against the wind in 2 hours. Find the speed of the wind.

https://math.libretexts.org/@go/page/30526

7. A light airplane flying with the wind can travel 600 miles in 4 hours. On the return trip, against the wind, it will take 5 hours. What are the speeds of the airplane and of the wind?
8. A boat can travel 15 miles with the current downstream in $1\frac{1}{4}$ hours. Returning upstream against the current, the boat can only travel $8\frac{3}{4}$ miles in the same amount of time. Find the speed of the current.
9. Mary jogged the trail from her car to the cabin at the rate of 6 miles per hour. She then walked back to her car at a rate of 4 miles per hour. If the entire trip took 1 hour, then how long did it take her to walk back to her car?
10. A jogger can sustain an average running rate of 8 miles per hour to his destination and 6 miles an hour on the return trip. Find the total distance the jogger ran if the total time running was $1\frac{3}{4}$ hour.
11. Two trains leave the station traveling in opposite directions. One train is 12 miles per hour faster than the other and in 3 hours they are 300 miles apart. Determine the average speed of each train.
12. Two trains leave the station traveling in opposite directions. One train is 8 miles per hour faster than the other and in $2\frac{1}{2}$ hours they are 230 miles apart. Determine the average speed of each train.

Answer

1. The first leg of the trip took 6 hours and the second leg took 2 hours.

3. It took her $\frac{1}{2}$ hour to drive to the airport.

5. 0.5 miles per hour.

7. Airplane: 135 miles per hour; wind: 15 miles per hour

9. $\frac{3}{5}$ hour

11. One train averaged 44 miles per hour and the other averaged 56 miles per hour.

This page titled 5.4E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.4E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **3.3: Applications of Linear Systems with Two Variables** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/advanced-algebra/index.html.

5.5: Solve Mixture Applications with Systems of Equations

Learning Objectives

By the end of this section, you will be able to:

- Solve mixture applications
- Solve interest applications

Note

Before you get started, take this readiness quiz.

1. Multiply 4.025(1,562).
   If you missed this problem, review Example 1.8.22.
2. Write 8.2% as a decimal.
   If you missed this problem, review Example 1.8.46.
3. Earl's dinner bill came to $32.50 and he wanted to leave an 18% tip. How much should the tip be?
   If you missed this problem, review Example 3.2.10.

Solve Mixture Applications

When we solved mixture applications with coins and tickets earlier, we started by creating a table so we could organize the information. For a coin example with nickels and dimes, the table looked like this:

Type	Number	• Value($)	= Total Value($)
nickels		0.05	
dimes		0.10	

Using one variable meant that we had to relate the number of nickels and the number of dimes. We had to decide if we were going to let n be the number of nickels and then write the number of dimes in terms of n, or if we would let d be the number of dimes and write the number of nickels in terms of d.

Now that we know how to solve systems of equations with two variables, we'll just let n be the number of nickels and d be the number of dimes. We'll write one equation based on the total value column, like we did before, and the other equation will come from the number column.

For the first example, we'll do a ticket problem where the ticket prices are in whole dollars, so we won't need to use decimals just yet.

Example 5.5.1

Translate to a system of equations and solve:

The box office at a movie theater sold 147 tickets for the evening show, and receipts totaled $1,302. How many $11 adult and how many $8 child tickets were sold?

Solution

Step 1. Read the problem.	We will create a table to organize the information.
Step 2. Identify what we are looking for.	We are looking for the number of adult tickets and the number of child tickets sold.
Step 3. Name what we are looking for.	Let a= the number of adult tickets. c= the number of child tickets
A table will help us organize the data. We have two types of tickets: adult and child.	Write a and c for the number of tickets.

Write the total number of tickets sold at the bottom of the Number column.	Altogether 147 were sold.
Write the value of each type of ticket in the Value column.	The value of each adult ticket is $11. The value of each child tickets is $8.
The number times the value gives the total value, so the total value of adult tickets is $a \cdot 11 = 11a$, and the total value of child tickets is $c \cdot 8 = 8c$.	<table><tr><th>Type</th><th>Number</th><th>•</th><th>Value ($)</th><th>=</th><th>Total Value ($)</th></tr><tr><td>adult</td><td>a</td><td></td><td>11</td><td></td><td>$11a$</td></tr><tr><td>child</td><td>c</td><td></td><td>8</td><td></td><td>$8c$</td></tr><tr><td></td><td>147</td><td></td><td></td><td></td><td>1302</td></tr></table>
Altogether the total value of the tickets was $1,302.	Fill in the Total Value column.
Step 4. Translate into a system of equations.	
The Number column and the Total Value column give us the system of equations. We will use the elimination method to solve this system.	$a + c = 147$ $11a + 8c = 1302$
Multiply the first equation by −8.	$-8(a + c) = -8(147)$ $11a + 8c = 1302$
Simplify and add, then solve for a.	$-8a - 8c = -1176$ $11a + 8c = 1302$ $3a = 126$
	$a = 42$
Substitute $a = 42$ into the first equation, then solve for c.	$42 + c = 147$
	$c = 105$
Step 5. Check the answer in the problem. 42 adult tickets at $11 per ticket makes $462 105 child tickets at $8 per ticket makes $840. The total receipts are $1,302.✓	
Step 6. Answer the question.	The movie theater sold 42 adult tickets and 105 child tickets.

Try It 5.5.2

Translate to a system of equations and solve:

The ticket office at the zoo sold 553 tickets one day. The receipts totaled $3,936. How many $9 adult tickets and how many $6 child tickets were sold?

Answer

There were 206 adult tickets sold and 347 children tickets sold.

Try It 5.5.3

Translate to a system of equations and solve:

A science center sold 1,363 tickets on a busy weekend. The receipts totaled $12,146. How many $12 adult tickets and how many $7 child tickets were sold?

Answer

There were 521 adult tickets sold and 842 children tickets sold.


In Example 5.5.4 we'll solve a coin problem. Now that we know how to work with systems of two variables, naming the variables in the 'number' column will be easy.

Example 5.5.4

Translate to a system of equations and solve:

Priam has a collection of nickels and quarters, with a total value of $7.30. The number of nickels is six less than three times the number of quarters. How many nickels and how many quarters does he have?

Solution

Step 1. Read the problem.	We will create a table to organize the information.
Step 2. Identify what we are looking for.	We are looking for the number of nickels and the number of quarters.
Step 3. Name what we are looking for.	Let n= the number of nickels. q= the number of quarters
A table will help us organize the data. We have two types of coins, nickels and quarters.	Write n and q for the number of each type of coin.
Fill in the Value column with the value of each type of coin.	The value of each nickel is $0.05. The value of each quarter is $0.25.
The number times the value gives the total value, so, the total value of the nickels is n (0.05) = 0.05n and the total value of quarters is q(0.25) = 0.25q. Altogether the total value of the coins is $7.30.	
Step 4. Translate into a system of equations.	
The Total value column gives one equation.	$0.05n + 0.25q = 7.30$
We also know the number of nickels is six less than three times the number of quarters. Translate to get the second equation.	$n = 3q - 6$
Now we have the system to solve.	$0.05n + 0.25q = 7.30$ $n = 3q - 6$
Step 5. Solve the system of equations We will use the substitution method. Substitute $n = 3q - 6$ into the first equation. Simplify and solve for q.	$0.05n + 0.25q = 7.30$
	$0.05(3q - 6) + 0.25q = 7.3$

	$0.15q - 0.3 + 0.25q = 7.3$
	$0.4q - 0.3 = 7.3$
	$0.4q = 7.6$
	$q = 19$
To find the number of nickels, substitute $q = 19$ into the second equation.	$n = 3q - 6$
	$n = 3 \cdot 19 - 6$
	$n = 51$
Step 6. Check the answer in the problem. $19 \text{ quarters at } \$0.25 = \$4.75$ $51 \text{ ickels at } \$0.05 = \$2.55$ $\text{Total } = \$7.30\checkmark$ $3 \cdot 19 - 16 = 51\checkmark$	
Step 7. Answer the question.	Priam has 19 quarters and 51 nickels.

Try It 5.5.5

Translate to a system of equations and solve:

Matilda has a handful of quarters and dimes, with a total value of $8.55. The number of quarters is 3 more than twice the number of dimes. How many dimes and how many quarters does she have?

Answer

Matilda has 13 dimes and 29 quarters.

Try It 5.5.6

Translate to a system of equations and solve:

Juan has a pocketful of nickels and dimes. The total value of the coins is $8.10. The number of dimes is 9 less than twice the number of nickels. How many nickels and how many dimes does Juan have?

Answer

Juan has 36 nickels and 63 dimes.

Some mixture applications involve combining foods or drinks. Example situations might include combining raisins and nuts to make a trail mix or using two types of coffee beans to make a blend.

Example 5.5.7

Translate to a system of equations and solve:

Carson wants to make 20 pounds of trail mix using nuts and chocolate chips. His budget requires that the trail mix costs him $7.60 per pound. Nuts cost $9.00 per pound and chocolate chips cost $2.00 per pound. How many pounds of nuts and how many pounds of chocolate chips should he use?

Solution

Step 1. Read the problem.	We will create a table to organize the information.

Step 2. Identify what we are looking for.	We are looking for the number of pounds of nuts and the number of pounds of chocolate chips.
Step 3. Name what we are looking for.	Let n= the number of pound of nuts. c= the number of pounds of chips
Carson will mix nuts and chocolate chips to get trail mix. Write in n and c for the number of pounds of nuts and chocolate chips. There will be 20 pounds of trail mix. Put the price per pound of each item in the Value column. Fill in the last column using	

Type	Number of pounds	• Value ($)	= Total Value ($)
nuts	n	9.00	$9n$
chocolate chips	c	2.00	$2c$
trail mix	20	7.60	$7.60(20) = 152$

Number · Value = Total Value	
Step 4. Translate into a system of equations. We get the equations from the Number and Total Value columns.	$\begin{aligned} n + c &= 20 \\ 9n + 2c &= 152 \end{aligned}$
Step 5. Solve the system of equations We will use elimination to solve the system.	
Multiply the first equation by −2 to eliminate c.	$\begin{aligned} -2(n + c) &= -2(20) \\ 9n + 2c &= 152 \end{aligned}$
Simplify and add. Solve for n.	$\begin{aligned} -2n - 2c &= -40 \\ 9n + 2c &= 152 \\ \hline 7n \quad &= 112 \end{aligned}$
	$n = 16$
To find the number of pounds of chocolate chips, substitute n = 16 into the first equation, then solve for c.	$\begin{aligned} n + c &= 20 \\ 16 + c &= 20 \end{aligned}$
	c=4
Step 6. Check the answer in the problem. $16 + 4 = 20$✓ $9 \cdot 16 + 2 \cdot 4 = 152$✓	
Step 7. Answer the question.	Carson should mix 16 pounds of nuts with 4 pounds of chocolate chips to create the trail mix.

? Try It 5.5.8

Translate to a system of equations and solve:

Greta wants to make 5 pounds of a nut mix using peanuts and cashews. Her budget requires the mixture to cost her $6 per pound. Peanuts are $4 per pound and cashews are $9 per pound. How many pounds of peanuts and how many pounds of cashews should she use?

Answer

Greta should use 3 pounds of peanuts and 2 pounds of cashews.

Try It 5.5.9

Translate to a system of equations and solve:

Sammy has most of the ingredients he needs to make a large batch of chili. The only items he lacks are beans and ground beef. He needs a total of 20 pounds combined of beans and ground beef and has a budget of $3 per pound. The price of beans is $1 per pound and the price of ground beef is $5 per pound. How many pounds of beans and how many pounds of ground beef should he purchase?

Answer

Sammy should purchase 10 pounds of beans and 10 pounds of ground beef.

Another application of mixture problems relates to concentrated cleaning supplies, other chemicals, and mixed drinks. The concentration is given as a percent. For example, a 20% concentrated household cleanser means that 20% of the total amount is cleanser, and the rest is water. To make 35 ounces of a 20% concentration, you mix 7 ounces (20% of 35) of the cleanser with 28 ounces of water.


For these kinds of mixture problems, we'll use percent instead of value for one of the columns in our table.

Example 5.5.10

Translate to a system of equations and solve:

Sasheena is a lab assistant at her community college. She needs to make 200 milliliters of a 40% solution of sulfuric acid for a lab experiment. The lab has only 25% and 50% solutions in the storeroom. How much should she mix of the 25% and the 50% solutions to make the 40% solution?

Solution

Step 1. Read the problem.	A figure may help us visualize the situation, then we will create a table to organize the information.
Sasheena must mix some of the 25% solution and some of the 50% solution together to get 200 ml of the 40% solution.	
Step 2. Identify what we are looking for.	We are looking for how much of each solution she needs.
Step 3. Name what we are looking for.	Let x= number of ml of 25% solution. y= number of ml of 50% solution
A table will help us organize the data. She will mix x ml of 25% with y ml of 50% to get 200 ml of 40% solution. We write the percents as decimals in the chart. We multiply the number of units times the concentration to get the total amount of sulfuric acid in each solution.	

Type	Number of units	• Concentration %	= Amount
25%	x	0.25	$0.25x$
50%	y	0.50	$0.50y$
40%	200	0.40	0.40(200)

Step 4. Translate into a system of equations. We get the equations from the Number column and the Amount column.	
Now we have the system.	$\begin{cases} x+y=200 \\ 0.25x+0.50y=0.40(200) \end{cases}$
Step 5. Solve the system of equations. We will solve the system by elimination. Multiply the first equation by −0.5 to eliminate y.	$\begin{cases} -0.5(x+y)=-0.5(200) \\ 0.25x+0.50y=80 \end{cases}$
Simplify and add to solve for x.	$\begin{cases} -0.5x-0.5y=-100 \\ 0.25x+0.5y=80 \end{cases}$ $-0.25x = -20$ $x = 80$
To solve for y, substitute $x = 80$ into the first equation.	$x+y=200$
	$80+y=200$
	$y=120$
Step 6. Check the answer in the problem. $80+120 = 120$✓ $0.25(80)+0.50(120) = 80$✓ Yes!	
Step 7. Answer the question.	Sasheena should mix 80 ml of the 25% solution with 120 ml of the 50% solution to get the 200 ml of the 40% solution.

Try It 5.5.11

Translate to a system of equations and solve:

LeBron needs 150 milliliters of a 30% solution of sulfuric acid for a lab experiment but only has access to a 25% and a 50% solution. How much of the 25% and how much of the 50% solution should he mix to make the 30% solution?

Answer

LeBron needs 120 ml of the 25% solution and 30 ml of the 50% solution.

Try It 5.5.12

Translate to a system of equations and solve:

Anatole needs to make 250 milliliters of a 25% solution of hydrochloric acid for a lab experiment. The lab only has a 10% solution and a 40% solution in the storeroom. How much of the 10% and how much of the 40% solutions should he mix to make the 25% solution?

Answer

Anatole should mix 125 ml of the 10% solution and 125 ml of the 40% solution.

Solve Interest Applications

The formula to model interest applications is $I = Prt$. Interest, I, is the product of the principal, P, the rate, r, and the time, t. In our work here, we will calculate the interest earned in one year, so t will be 1.

We modify the column titles in the mixture table to show the formula for interest, as you'll see in Example 5.5.13.

Example 5.5.13

Translate to a system of equations and solve:

Adnan has $40,000 to invest and hopes to earn 7.1% interest per year. He will put some of the money into a stock fund that earns 8% per year and the rest into bonds that earns 3% per year. How much money should he put into each fund?

Solution

Step 1. Read the problem.	A chart will help us organize the information.
Step 2. Identify what we are looking for.	We are looking for the amount to invest in each fund.
Step 3. Name what we are looking for.	Let s= the amount invested in stocks. b= the amount invested in bonds.
Write the interest rate as a decimal for each fund. Multiply: Principal · Rate · Time to get the Interest.	<table><tr><th>Account</th><th>Principal</th><th>Rate</th><th>Time</th><th>Interest</th></tr><tr><td>stock fund</td><td>s</td><td>0.08</td><td>1</td><td>0.08s</td></tr><tr><td>bonds</td><td>b</td><td>0.03</td><td>1</td><td>0.03b</td></tr><tr><td>Total</td><td>40,000</td><td>0.071</td><td></td><td>0.071(40,000)</td></tr></table>
Step 4. Translate into a system of equations. We get our system of equations from the Principal column and the Interest column.	$\begin{cases} s + b = 40{,}000 \\ 0.08s + 0.03b = 0.071(40{,}000) \end{cases}$
Step 5. Solve the system of equations Solve by elimination. Multiply the top equation by −0.03.	$\begin{cases} -0.03(s + b) = -0.03(40{,}000) \\ 0.08s + 0.03b = 2{,}840 \end{cases}$
Simplify and add to solve for s.	$\begin{array}{r} -0.03s - 0.03b = -1{,}200 \\ 0.08s + 0.03b = 2{,}840 \\ \hline 0.05s = 1{,}640 \end{array}$
	$s = 32{,}800$
To find b, substitute s = 32,800 into the first equation.	$\begin{array}{r} s + b = 40{,}000 \\ 32{,}800 + b = 40{,}000 \end{array}$
	$b = 7{,}200$
Step 6. Check the answer in the problem.	We leave the check to you.
Step 7. Answer the question.	Adnan should invest $32,800 in stock and $7,200 in bonds.

Did you notice that the Principal column represents the total amount of money invested while the Interest column represents only the interest earned? Likewise, the first equation in our system, $s + b = 40{,}000$, represents the total amount of money invested and the second equation, $0.08s + 0.03b = 0.071(40{,}000)$, represents the interest earned.

Try It 5.5.14

Translate to a system of equations and solve:

Leon had $50,000 to invest and hopes to earn 6.2 % interest per year. He will put some of the money into a stock fund that earns 7% per year and the rest in to a savings account that earns 2% per year. How much money should he put into each fund?

Answer

Leon should put $42,000 in the stock fund and $8000 in the savings account.

Try It 5.5.15

Translate to a system of equations and solve:

Julius invested $7,000 into two stock investments. One stock paid 11% interest and the other stock paid 13% interest. He earned 12.5% interest on the total investment. How much money did he put in each stock?

Answer

Julius invested $1,750 at 11% and $5,250 at 13%.

Example 5.5.16

Translate to a system of equations and solve:

Rosie owes $21,540 on her two student loans. The interest rate on her bank loan is 10.5% and the interest rate on the federal loan is 5.9%. The total amount of interest she paid last year was $1,669.68. What was the principal for each loan?

Solution

Step 1. Read the problem.	A chart will help us organize the information.
Step 2. Identify what we are looking for.	We are looking for the principal of each loan.
Step 3. Name what we are looking for.	Let b= the principal for the bank loan. f= the principal on the federal loan
The total loans are $21,540.	
Record the interest rates as decimals in the chart.	

Account	Principal •	Rate •	Time =	Interest
bank	b	0.105	1	$0.105b$
federal	f	0.059	1	$0.059f$
Total	21,540			1669.68

Multiply using the formula $I = Prt$ to get the Interest.	
Step 4. Translate into a system of equations. The system of equations comes from the Principal column and the Interest column.	$b + f = 21{,}540$ $0.105b + 0.059f = 1669.68$
Step 5. Solve the system of equations We will use substitution to solve. Solve the first equation for b.	$b + f = 21{,}540$ $b = -f + 21{,}540$
Substitute $b = -f + 21{,}540$ into the second equation.	$0.105b + 0.059f = 1669.68$ $0.105(-f + 21{,}540) + 0.059f = 1669.68$

Simplify and solve for f.	$-0.105f + 2261.70 + 0.059f = 1669.68$
	$-0.046f + 2261.70 = 1669.68$
	$-0.046f = -592.02$
	$f = 12{,}870$
To find b, substitute $f = 12{,}870$ into the first equation.	$b + f = 21{,}540$ $12{,}870 + f = 21{,}540$
	$f = 8{,}670$
Step 6. Check the answer in the problem.	We leave the check to you.
Step 7. Answer the question.	The principal of the bank loan is \$12,870 and the principal for the federal loan is \$8,670.

Try It 5.5.17

Translate to a system of equations and solve:

Laura owes \$18,000 on her student loans. The interest rate on the bank loan is 2.5% and the interest rate on the federal loan is 6.9 %. The total amount of interest she paid last year was \$1,066. What was the principal for each loan?

Answer

The principal amount for the bank loan was \$4,000. The principal amount for the federal loan was \$14,000.

Try It 5.5.18

Translate to a system of equations and solve:

Jill's Sandwich Shoppe owes \$65,200 on two business loans, one at 4.5% interest and the other at 7.2% interest. The total amount of interest owed last year was \$3,582. What was the principal for each loan?

Answer

The principal amount for was \$41,200 at 4.5%. The principal amount was, \$24,000 at 7.2%.

Note

Access these online resources for additional instruction and practice with solving application problems with systems of linear equations.

- Cost and Mixture Word Problems
- Mixture Problems

Key Concepts

- **Table for coin and mixture applications**

Type	Number	• Value(\$)	= Total Value(\$)
Total			

- **Table for concentration applications**

Type	Number of units	•	Concentration %	=	Amount
Total					

- **Table for interest applications**

Account	Principal	•	Rate	•	Time	=	Interest
					1		
					1		
Total							

This page titled 5.5: Solve Mixture Applications with Systems of Equations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.5: Solve Mixture Applications with Systems of Equations** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.5E: Exercises

Practice Makes Perfect

Solve Mixture Applications

In the following exercises, translate to a system of equations and solve.

Exercise 5.5E.1

Tickets to a Broadway show cost $35 for adults and $15 for children. The total receipts for 1650 tickets at one performance were $47,150. How many adult and how many child tickets were sold?

Answer

There 1120 adult tickets and 530 child tickets sold.

Exercise 5.5E.2

Tickets for a show are $70 for adults and $50 for children. One evening performance had a total of 300 tickets sold and the receipts totaled $17,200. How many adult and how many child tickets were sold?

Exercise 5.5E.3

Tickets for a train cost $10 for children and $22 for adults. Josie paid $1,200 for a total of 72 tickets. How many children's tickets and how many adult tickets did Josie buy?

Answer

Josie bought 40 adult tickets and 32 children tickets.

Exercise 5.5E.4

Tickets for a baseball game are $69 for Main Level seats and $39 for Terrace Level seats. A group of sixteen friends went to the game and spent a total of $804 for the tickets. How many of Main Level and how many Terrace Level tickets did they buy?

Exercise 5.5E.5

Tickets for a dance recital cost $15 for adults and $7 for children. The dance company sold 253 tickets and the total receipts were $2,771. How many adult tickets and how many child tickets were sold?

Answer

There were 125 adult tickets and 128 children tickets sold.

Exercise 5.5E.6

Tickets for the community fair cost $12 for adults and $5 dollars for children. On the first day of the fair, 312 tickets were sold for a total of $2,204. How many adult tickets and how many child tickets were sold?

Exercise 5.5E.7

Brandon has a cup of quarters and dimes with a total value of $3.80. The number of quarters is four less than twice the number of dimes. How many quarters and how many dimes does Brandon have?

Answer

Brandon has 12 quarters and 8 dimes.

Exercise 5.5E.8

Sherri saves nickels and dimes in a coin purse for her daughter. The total value of the coins in the purse is $0.95. The number of nickels is two less than five times the number of dimes. How many nickels and how many dimes are in the coin purse?

Exercise 5.5E.9

Peter has been saving his loose change for several days. When he counted his quarters and dimes, he found they had a total value $13.10. The number of quarters was fifteen more than three times the number of dimes. How many quarters and how many dimes did Peter have?

Answer

Peter had 11 dimes and 48 quarters.

Exercise 5.5E.10

Lucinda had a pocketful of dimes and quarters with a value of $ $6.20. The number of dimes is eighteen more than three times the number of quarters. How many dimes and how many quarters does Lucinda have?

Exercise 5.5E.11

A cashier has 30 bills, all of which are $10 or $20 bills. The total value of the money is $460. How many of each type of bill does the cashier have?

Answer

The cashier has fourteen $10 bills and sixteen $20 bills.

Exercise 5.5E.12

A cashier has 54 bills, all of which are $10 or $20 bills. The total value of the money is $910. How many of each type of bill does the cashier have?

Exercise 5.5E.13

Marissa wants to blend candy selling for $1.80 per pound with candy costing $1.20 per pound to get a mixture that costs her $1.40 per pound to make. She wants to make 90 pounds of the candy blend. How many pounds of each type of candy should she use?

Answer

Marissa should use 60 pounds of the $1.20/lb candy and 30 pounds of the $1.80/lb candy.

Exercise 5.5E.14

How many pounds of nuts selling for $6 per pound and raisins selling for $3 per pound should Kurt combine to obtain 120 pounds of trail mix that cost him $5 per pound?

Exercise 5.5E.15

Hannah has to make twenty-five gallons of punch for a potluck. The punch is made of soda and fruit drink. The cost of the soda is $1.79 per gallon and the cost of the fruit drink is $2.49 per gallon. Hannah's budget requires that the punch cost $2.21 per gallon. How many gallons of soda and how many gallons of fruit drink does she need?

Answer

Hannah needs 10 gallons of soda and 15 gallons of fruit drink.

Exercise 5.5E.16

Joseph would like to make 12 pounds of a coffee blend at a cost of $6.25 per pound. He blends Ground Chicory at $4.40 a pound with Jamaican Blue Mountain at $8.84 per pound. How much of each type of coffee should he use?

Exercise 5.5E.17

Julia and her husband own a coffee shop. They experimented with mixing a City Roast Columbian coffee that cost $7.80 per pound with French Roast Columbian coffee that cost $8.10 per pound to make a 20 pound blend. Their blend should cost them $7.92 per pound. How much of each type of coffee should they buy?

Answer

Julia and her husband should buy 12 pounds of City Roast Columbian coffee and 8 pounds of French Roast Columbian coffee.

Exercise 5.5E.18

Melody wants to sell bags of mixed candy at her lemonade stand. She will mix chocolate pieces that cost $4.89 per bag with peanut butter pieces that cost $3.79 per bag to get a total of twenty-five bags of mixed candy. Melody wants the bags of mixed candy to cost her $4.23 a bag to make. How many bags of chocolate pieces and how many bags of peanut butter pieces should she use?

Exercise 5.5E.19

Jotham needs 70 liters of a 50% alcohol solution. He has a 30% and an 80% solution available. How many liters of the 30% and how many liters of the 80% solutions should he mix to make the 50% solution?

Answer

Jotham should mix 42 liters of the 30% solution and 28 liters of the 80% solution.

Exercise 5.5E.20

Joy is preparing 15 liters of a 25% saline solution. She only has 40% and 10% solution in her lab. How many liters of the 40% and how many liters of the 10% should she mix to make the 25% solution?

Exercise 5.5E.21

A scientist needs 65 liters of a 15% alcohol solution. She has available a 25% and a 12% solution. How many liters of the 25% and how many liters of the 12% solutions should she mix to make the 15% solution?

Answer

The scientist should mix 15 liters of the 25% solution and 50 liters of the 12% solution.

Exercise 5.5E.22

A scientist needs 120 liters of a 20% acid solution for an experiment. The lab has available a 25% and a 10% solution. How many liters of the 25% and how many liters of the 10% solutions should the scientist mix to make the 20% solution?

Exercise 5.5E.23

A 40% antifreeze solution is to be mixed with a 70% antifreeze solution to get 240 liters of a 50% solution. How many liters of the 40% and how many liters of the 70% solutions will be used?

Answer

160 liters of the 40% solution and 80 liters of the 70% solution will be used.

Exercise 5.5E.24

A 90% antifreeze solution is to be mixed with a 75% antifreeze solution to get 360 liters of a 85% solution. How many liters of the 90% and how many liters of the 75% solutions will be used?

Solve Interest Applications

In the following exercises, translate to a system of equations and solve.

Exercise 5.5E.25

Hattie had $3,000 to invest and wants to earn 10.6% interest per year. She will put some of the money into an account that earns 12% per year and the rest into an account that earns 10% per year. How much money should she put into each account?

Answer

Hattie should invest $900 at 12% and $2,100 at 10%.

Exercise 5.5E.26

Carol invested $2,560 into two accounts. One account paid 8% interest and the other paid 6% interest. She earned 7.25% interest on the total investment. How much money did she put in each account?

Exercise 5.5E.27

Sam invested $48,000, some at 6% interest and the rest at 10%. How much did he invest at each rate if he received $4,000 in interest in one year?

Answer

Sam invested $28,000 at 10% and $20,000 at 6%.

Exercise 5.5E.28

Arnold invested $64,000, some at 5.5% interest and the rest at 9%. How much did he invest at each rate if he received $4,500 in interest in one year?

Exercise 5.5E.29

After four years in college, Josie owes $65,800 in student loans. The interest rate on the federal loans is 4.5% and the rate on the private bank loans is 2%. The total interest she owed for one year was $2,878.50. What is the amount of each loan?

Answer

The federal loan is $62,500 and the bank loan is $3,300.

Exercise 5.5E.30

Mark wants to invest $10,000 to pay for his daughter's wedding next year. He will invest some of the money in a short term CD that pays 12% interest and the rest in a money market savings account that pays 5% interest. How much should he invest at each rate if he wants to earn $1,095 in interest in one year?

Exercise 5.5E.31

A trust fund worth $25,000 is invested in two different portfolios. This year, one portfolio is expected to earn 5.25% interest and the other is expected to earn 4%. Plans are for the total interest on the fund to be $1150 in one year. How much money should be invested at each rate?

Answer

$12,000 should be invested at 5.25% and $13,000 should be invested at 4%.

Exercise 5.5E.32

A business has two loans totaling $85,000. One loan has a rate of 6% and the other has a rate of 4.5%. This year, the business expects to pay $4650 in interest on the two loans. How much is each loan?

Everyday Math

In the following exercises, translate to a system of equations and solve.

Exercise 5.5E.33

Laurie was completing the treasurer's report for her son's Boy Scout troop at the end of the school year. She didn't remember how many boys had paid the $15 full-year registration fee and how many had paid the $10 partial-year fee. She knew that the number of boys who paid for a full-year was ten more than the number who paid for a partial-year. If $250 was collected for all the registrations, how many boys had paid the full-year fee and how many had paid the partial-year fee?

Answer

14 boys paid the full-year fee. 4 boys paid the partial-year fee

Exercise 5.5E.34

As the treasurer of her daughter's Girl Scout troop, Laney collected money for some girls and adults to go to a three-day camp. Each girl paid $75 and each adult paid $30. The total amount of money collected for camp was $765. If the number of girls is three times the number of adults, how many girls and how many adults paid for camp?

Writing Exercises

Exercise 5.5E.35

Take a handful of two types of coins, and write a problem similar to Example relating the total number of coins and their total value. Set up a system of equations to describe your situation and then solve it.

Answer

Answers will vary.

Exercise 5.5E.36

In Example we solved the system of equations $\begin{cases} b+f=21,540 \\ 0.105b+0.059f=1669.68 \end{cases}$ by substitution. Would you have used substitution or elimination to solve this system? Why?

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve mixture applications.			
solve interest applications.			

b. After looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

More Practice: Money Problems

Exercise 5.5E.37

1. Two families bought tickets for the home basketball game. One family ordered 2 adult tickets and 4 children's tickets for a total of $36.00. Another family ordered 3 adult tickets and 2 children's tickets for a total of $32.00. How much did each ticket cost?
2. Two friends found shirts and shorts on sale at a flea market. One bought 4 shirts and 2 shorts for a total of $28.00. The other bought 3 shirts and 3 shorts for a total of $30.75. How much was each shirt and each pair of shorts?
3. A community theater sold 140 tickets to the evening musical for a total of $1,540. Each adult ticket was sold for $12 and each child ticket was sold for $8. How many adult tickets were sold?
4. The campus bookstore sells graphing calculators for $110 and scientific calculators for $16. On the first day of classes 50 calculators were sold for a total of $1,646. How many of each were sold?
5. Jim was able to purchase a pizza for $12.35 with quarters and dimes. If he uses 71 coins to buy the pizza, then how many of each did he have?
6. A cash register contains $5 bills and $10 bills with a total value of $350. If there are 46 bills total, then how many of each does the register contain?
7. A jar consisting of only nickels and quarters contains 70 coins. If the total value is $9.10, how many of each coin are in the jar?
8. Jill has $9.20 worth of dimes and quarters. If there are 68 coins in total, how many of each does she have?

Answer

1. Adults $7.00 each and children $5.50 each.
3. 105 adult tickets were sold.
5. 35 quarters and 36 dimes
7. The jar contains 42 nickels and 28 quarters

More Practice: Combined fixed and variable costs

A taxi fare includes a fixed flat fee, plus a variable cost that depends on the number of miles driven.
Total manufacturing costs include a fixed overhead cost, plus a variable cost that depends on the number of items manufactured.

Exercise 5.5E.38

1. In a moderate-size city, taxi fare is $51 for 12 miles and is $31 for 7 miles. What is the flat fee and what is the charge per mile?
2. If it costs $90 to rent a car driven 100 miles and $140 for one driven 200 miles, find the daily rental fee and the mileage fee.
3. To manufacture 30 items, it costs $2700, and to manufacture 50 items, it costs $3200. Find the overhead cost and the cost to manufacture one item

4. To manufacture 100 items, it costs $32,000, and to manufacture 200 items, it costs $40,000. Find the overhead cost and the cost to manufacture one item
5. If it costs a total of $750 to produce 20 items and the cost to manufacture an item is $25, what is the overhead cost?
6. If it costs $1900 to manufacture 60 items, and the overhead is $700, what is the cost to produce one item?
7. Sally earns $\$1,000$ per month plus a commission of 2% of sales. Jane earns $\$200$ per month plus 6% of her sales. At what monthly sales figure will both Sally and Jane earn the same amount of pay?
8. The cost of producing specialty book shelves includes a daily production cost plus an additional cost per book shelf. The cost to produce 7 book shelves in 2 days was $\$590$. The cost to produce 11 book shelves in 3 days was $\$910$. Find the daily cost and the cost for one bookshelf.

Answers

1. $\$4$ per mile 3. $\$25$ per item; $\$1950$ overhead costs. 5. $\$250$ overhead 7. Sales of $\$20,000$

More Practice: Percent Solution Problems

? Exercise $5.5E.39$

Set up a linear system and solve.

1. A 17% acid solution is to be mixed with a 9% acid solution to produce 8 gallons of a 10% acid solution. How much of each is needed?

2. A nurse wishes to obtain 28 ounces of a 1.5% saline solution. How much of a 1% saline solution must she mix with a 4.5% saline solution to achieve the desired mixture?

3. A customer ordered 4 pounds of a mixed peanut product containing 12% cashews. The inventory consists of only two mixes containing 10% and 26% cashews. How much of each type must be mixed to fill the order?

4. One alcohol solution contains 10% alcohol and another contains 25% alcohol. How much of each should be mixed together to obtain 2 gallons of a 13.75% alcohol solution?

5. A 50% fruit juice concentrate can be purchased wholesale. Best taste is achieved when water is mixed with the concentrate in such a way as to obtain a 15% fruit juice mixture. How much water and concentrate is needed to make a 60-ounce fruit juice drink?

6. How much cleaning fluid concentrate, with 60% alcohol content, must be mixed with water to obtain a 24-ounce mixture with 15% alcohol content?

7. How many pounds of pure peanuts must be combined with a 20% peanut mix to produce 2 pounds of a 50% peanut mix?

8. Pure sugar is to be mixed with a fruit salad containing 10% sugar to produce 63 ounces of a salad containing 18% sugar. How much pure sugar is required?

9. A custom aluminum alloy is created by mixing 150 grams of a 15% aluminum alloy and 350 grams of a 55% aluminum alloy. What percentage of aluminum is in the resulting mixture?

10. A research assistant mixed 500 milliliters of a solution that contained a 12% acid with 300 milliliters of water. What percentage of acid is in the resulting solution?

Answer

1. 7 gallons of the 9% acid solution and 1 gallon of the 17% acid solution

3. 3.5 pounds of the 10% cashew mix and 0.5 pounds of the 26% cashew mix

5. 18 ounces of fruit juice concentrate and 42 ounces of water

7. .75 pounds of pure peanuts

9. 43%.

More Practice: Per Cent Interest

Exercise 5.5E. 40

1. A $\$5,200$ principal is invested in two accounts, one earning 3% interest and another earning 6% interest. If the total interest for the year is $\$210$, then how much is invested in each account?
2. Harry's $\$2,200$ savings is in two accounts. One account earns 2% annual interest and the other earns 4%. His total interest for the year is $\$69$. How much does he have in each account?
3. Janine has two savings accounts totaling $\$6,500$. One account earns $2\frac{3}{4}\%$ annual interest and the other earns $3\frac{1}{2}\%$. If her total interest for the year is $\$211$, then how much is in each account?
4. Margaret has her total savings of $\$24,200$ in two different CD accounts. One CD earns 4.6% interest and another earns 3.4% interest. If her total interest for the year is $\$1,007.60$, then how much does she have in each CD account?
5. Last year Mandy earned twice as much interest in her Money Market fund as she did in her regular savings account. The total interest from the two accounts was $\$246$. How much interest did she earn in each account?
6. A small business invested $\$120,000$ in two accounts. The account earning 4% annual interest yielded twice as much interest as the account earning 3% annual interest. How much was invested in each account?

Answer

1. $\$3,400$ at 3% and $\$1,800$ at 6%
3. $\$2,200$ at $2\frac{3}{4}\%$ and $\$4,300$ at $3\frac{1}{2}\%$
5. Savings: $\$82$; Money Market: $\$164$

This page titled 5.5E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.5E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.
- **3.3: Applications of Linear Systems with Two Variables** by Anonymous is licensed CC BY-NC-SA 3.0. Original source: https://2012books.lardbucket.org/books/advanced-algebra/index.html.
- **1.4.1: Applications (Exercises)** by Rupinder Sekhon and Roberta Bloom is licensed CC BY 4.0. Original source: https://www.deanza.edu/faculty/bloomroberta/math11/afm3files.html.html.

5.6: Graphing Systems of Linear Inequalities

Learning Objectives

By the end of this section, you will be able to:

- Determine whether an ordered pair is a solution of a system of linear inequalities
- Solve a system of linear inequalities by graphing
- Solve applications of systems of inequalities

Note

Before you get started, take this readiness quiz.

1. Graph x>2 on a number line.
   If you missed this problem, review Example 2.7.1.
2. Solve the inequality 2a<5a+12.
   If you missed this problem, review Example 2.7.22.
3. Determine whether the ordered pair $(3, \frac{1}{2})$ is a solution to the system $\begin{cases} x+2y=4 \\ y=6x \end{cases}$.
   If you missed this problem, review Example 5.1.1.

Determine Whether an Ordered Pair is a Solution of a System of Linear Inequalities

The definition of a system of linear inequalities is very similar to the definition of a system of linear equations.

SYSTEM OF LINEAR INEQUALITIES

Two or more linear inequalities grouped together form a **system of linear inequalities**.

A system of linear inequalities looks like a system of linear equations, but it has inequalities instead of equations. A system of two linear inequalities is shown below.

$$\begin{cases} x+4y \geq 10 \\ 3x-2y < 12 \end{cases} \tag{5.6.1}$$

To solve a system of linear inequalities, we will find values of the variables that are solutions to both inequalities. We solve the system by using the graphs of each inequality and show the solution as a graph. We will find the region on the plane that contains all ordered pairs (x,y)(x,y) that make both inequalities true.

SolutionS OF A SYSTEM OF LINEAR INEQUALITIES

Solutions of a system of linear inequalities are the values of the variables that make all the inequalities true.

The solution of a system of linear inequalities is shown as a shaded region in the *x*-*y* coordinate system that includes all the points whose ordered pairs make the inequalities true.

To determine if an ordered pair is a solution to a system of two inequalities, we substitute the values of the variables into each inequality. If the ordered pair makes both inequalities true, it is a solution to the system.

Example 5.6.1

Determine whether the ordered pair is a solution to the system. $\begin{cases} x+4y \geq 10 \\ 3x-2y < 12 \end{cases}$

1. (−2, 4)
2. (3,1)

Solution

1. Is the ordered pair (−2, 4) a solution?
We substitute $x = -2$ and $y = 4$ into both inequalities.

$$x + 4y \geq 10 \qquad 3x - 2y < 12$$
$$-2 + 4(4) \stackrel{?}{\geq} 10 \qquad 3(-2) - 2(4) \stackrel{?}{<} 12$$
$$14 \geq 10 \text{ true} \qquad -14 < 12 \text{ true}$$

The ordered pair (−2, 4) made both inequalities true. Therefore (−2, 4) is a solution to this system.

2. Is the ordered pair (3,1) a solution?
We substitute $x = 3$ and $y = 1$ into both inequalities.

$$x + 4y \geq 10 \qquad 3x - 2y < 12$$
$$3 + 4(1) \stackrel{?}{\geq} 10 \qquad 3(3) - 2(1) \stackrel{?}{<} 12$$
$$7 \geq 10 \text{ false} \qquad 7 < 12 \text{ true}$$

The ordered pair (3,1) made one inequality true, but the other one false. Therefore (3,1) is not a solution to this system.

Try It 5.6.2

Determine whether the ordered pair is a solution to the system. $\begin{cases} x - 5y > 10 \\ 2x + 3y > -2 \end{cases}$

1. (3,−1)
2. (6,−3)

Answer

1. no
2. yes

Try It 5.6.3

Determine whether the ordered pair is a solution to the system. $\begin{cases} y > 4x - 2 \\ 4x - y < 20 \end{cases}$

1. (2,1)
2. (4,−1)

Answer

1. no
2. no



Solve a System of Linear Inequalities by Graphing

The solution to a single linear inequality is the region on one side of the boundary line that contains all the points that make the inequality true. The solution to a system of two linear inequalities is a region that contains the solutions to both inequalities. To find this region, we will graph each inequality separately and then locate the region where they are both true. The solution is always shown as a graph.

Example 5.6.4: How to Solve a System of Linear inequalities

Solve the system by graphing. $\begin{cases} y \geq 2x - 1 \\ y < x + 1 \end{cases}$


Solution

Step 1. Graph the first inequality. Graph the boundary line. Shade in the side of the boundary line where the inequality is true.	We will graph $y \geq 2x - 1$. We graph the line $y = 2x - 1$. It is a solid line because the inequality sign is $\geq$. We choose (0,0) as a test point. It is a solution to $y \geq 2x - 1$, so we shade in the left side of the boundary line.	$\begin{cases} y \geq 2x - 1 \\ y < x + 1 \end{cases}$ 
Step 2. On the same grid, graph the second inequality. Graph the boundary line. Shade in the side of that boundary line where the inequality is true.	We will graph $y < x + 1$ on the same grid. We graph the line $y = x + 1$. It is a dashed line because the inequality sign is $<$. Again, we use (0,0) as a test point. It is a solution so we shade in that side of the line $y = x + 1$.	
Step 3. The solution is the region where the shading overlaps.	The point where the boundary lines intersect is not a solution because it is not a solution to $y < x + 1$.	The solution is all points in the darker shaded region.
Step 4. Check by choosing a test point.	We'll use (–1, –1) as a test point.	Is (–1, –1) a solution to $y \geq 2x - 1$? $-1 \stackrel{?}{\geq} 2(-1) - 1$ $-1 \geq -3$ true Is (–1, –1) a solution to $y < x + 1$? $-1 \stackrel{?}{<} -1 + 1$ $-1 < 0$ true The region containing (–1, –1) is the solution to this system.

? Try It 5.6.5

Solve the system by graphing. $\begin{cases} y < 3x + 2 \\ y > -x - 1 \end{cases}$


Answer



Try It 5.6.6

Solve the system by graphing. $\begin{cases} y < -\frac{1}{2}x + 3 \\ y < 3x - 4 \end{cases}$

Answer



SOLVE A SYSTEM OF LINEAR INEQUALITIES BY GRAPHING.

1. Graph the first inequality.
   - Graph the boundary line.
   - Shade in the side of the boundary line where the inequality is true.
2. On the same grid, graph the second inequality.
   - Graph the boundary line.
   - Shade in the side of that boundary line where the inequality is true.
3. The solution is the region where the shading overlaps.
4. Check by choosing a test point.

Example 5.6.7


Solve the system by graphing. $\begin{cases} x - y > 3 \\ y < -\frac{1}{5}x + 4 \end{cases}$

Solution

Graph $x - y > 3$, by graphing $x - y = 3$ and testing a point.

The intercepts are $x = 3$ and $y = -3$ and the boundary line will be dashed.


Test (0, 0). It makes the inequality false. So, shade the side that does not contain (0, 0) red.



Graph $y<-15x+4$ by graphing $y=-15x+4$ using the slope $m=-15$ and y−intercept $b = 4$. The boundary line will be dashed.

Test (0, 0). It makes the inequality true, so shade the side that contains (0, 0) blue.

Choose a test point in the solution and verify that it is a solution to both inequalities.




The point of intersection of the two lines is not included as both boundary lines were dashed. The solution is the area shaded twice which is the darker-shaded region.

Try It 5.6.8

Solve the system by graphing. $\begin{cases} x + y \leq 2 \\ y \geq \frac{2}{3}x - 1 \end{cases}$


Answer



Try It 5.6.9

Solve the system by graphing. $\begin{cases} 3x - 2y \leq 6 \\ y > -\frac{1}{4}x + 5 \end{cases}$

Answer




Example 5.6.10

Solve the system by graphing. $\begin{cases} x - 2y < 5 \\ y > -4 \end{cases}$

Solution


Graph x−2y<5, by graphing x−2y=5 and testing a point.
The intercepts are $x = 5$ and $y = -2.5$ and the boundary line will be dashed.

Test (0, 0). It makes the inequality true. So, shade the side that contains (0, 0) red.



Graph $y > -4$, by graphing $y = -4$ and recognizing that it is a horizontal line through $y = -4$. The boundary line will be dashed.

Test (0, 0). It makes the inequality true. So, shade (blue) the side that contains (0, 0) blue.




The point (0, 0) is in the solution and we have already found it to be a solution of each inequality. The point of intersection of the two lines is not included as both boundary lines were dashed.

The solution is the area shaded twice which is the darker-shaded region.

Try It 5.6.11

Solve the system by graphing. $\begin{cases} y \geq 3x - 2 \\ y < -1 \end{cases}$


Answer



Try It 5.6.12

Solve the system by graphing. $\begin{cases} x > -4 \\ x - 2y \leq -4 \end{cases}$

Answer



Systems of linear inequalities where the boundary lines are parallel might have no solution. We'll see this in Example 5.6.13


Example 5.6.13

Solve the system by graphing. $\begin{cases} 4x + 3y \geq 12 \\ y < -\frac{4}{3}x + 1 \end{cases}$

Solution


Graph $4x + 3y \geq 12$, by graphing 4x+3y=12 and testing a point.
The intercepts are $x = 3$ and $y = 4$ and the boundary line will be solid.

Test (0, 0). It makes the inequality false. So,
shade the side that does not contain (0, 0) red.



Graph $y < -\frac{4}{3}x + 1$ by graphing $y = -\frac{4}{3}x + 1$ using the slope $m = \frac{4}{3}$ and the y-intercept $b = 1$. The boundary line will be dashed.

Test (0, 0). It makes the inequality true. So,
shade the side that contains (0, 0) blue.




There is no point in both shaded regions, so the system has no solution. This system has no solution.

? Try It 5.6.14

Solve the system by graphing. $\begin{cases} 3x - 2y \leq 12 \\ y \geq \frac{3}{2}x + 1 \end{cases}$

Answer

no solution




Try It 5.6.15

Solve the system by graphing. $\begin{cases} x+3y>8 \\ y<-\frac{1}{3}x-2 \end{cases}$

Answer

no solution




Example 5.6.16

Solve the system by graphing. $\begin{cases} y>\frac{1}{2}x-4 \\ x-2y<-4 \end{cases}$

Solution


Graph $y>\frac{1}{2}x-4$ by graphing $y=\frac{1}{2}x-4$ using the slope $m=\frac{1}{2}$ and the intercept $b=-4$. The boundary line will be dashed. Test (0, 0). It makes the inequality true. So, shade the side that contains (0, 0) red.



https://math.libretexts.org/@go/page/30529

Graph x−2y<−4x−2y<−4 by graphing x−2y=−4x−2y=−4 and testing a point.
The intercepts are $x = -4$ and $y = 2$ and the boundary line will be dashed.

Choose a test point in the solution and verify that it is a solution to both inequalities.



No point on the boundary lines is included in the solution as both lines are dashed.


The solution is the region that is shaded twice, which is also the solution to x−2y<−4.

Try It 5.6.17

Solve the system by graphing. $\begin{cases} y \geq 3x+1 \\ -3x+y \geq -4 \end{cases}$

Answer

$y \geq 3x+1$




Try It 5.6.18

Solve the system by graphing. $\begin{cases} y \leq -\frac{1}{4}x+2 \\ x+4y \leq 4 \end{cases}$

Answer

$x+4y \leq 4$



Solve Applications of Systems of Inequalities

The first thing we'll need to do to solve applications of systems of inequalities is to translate each condition into an inequality. Then we graph the system as we did above to see the region that contains the solutions. Many situations will be realistic only if both variables are positive, so their graphs will only show Quadrant I.

Example 5.6.19

Christy sells her photographs at a booth at a street fair. At the start of the day, she wants to have at least 25 photos to display at her booth. Each small photo she displays costs her $4 and each large photo costs her $10. She doesn't want to spend more than $200 on photos to display.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could she display 15 small and 5 large photos?
4. Could she display 3 large and 22 small photos?

Solution

1.Let x= the number of small photos.

y= the number of large photos

To find the system of inequalities, translate the information.

$$\text{She wants to have at least 25 photos.}$$

$$\text{The number of small plus the number of large should be at least 25.}$$

$$x+y \geq 25$$

$$\$4 \text{ for each small and } \$10 \text{ for each large must be no more than } \$200$$


$$4x+10y \leq 200$$

We have our system of inequalities. $\begin{cases} x+y \geq 25 \\ 4x+10y \leq 200 \end{cases}$

2.

To graph $x+y \geq 25$, graph $x + y = 25$ as a solid line.
Choose (0, 0) as a test point. Since it does not make the inequality
true, shade the side that does not include the point (0, 0) red.

To graph $4x+10y \leq 200$, graph $4x + 10y = 200$ as a solid line.
Choose (0, 0) as a test point. Since it does not make the inequality
true, shade the side that includes the point (0, 0) blue.



The solution of the system is the region of the graph that is double shaded and so is shaded darker.

3. To determine if 10 small and 20 large photos would work, we see if the point (10, 20) is in the solution region. It is not. Christy would not display 10 small and 20 large photos.

4. To determine if 20 small and 10 large photos would work, we see if the point (20, 10) is in the solution region. It is. Christy could choose to display 20 small and 10 large photos.


Notice that we could also test the possible solutions by substituting the values into each inequality.

Try It 5.6.20

A trailer can carry a maximum weight of 160 pounds and a maximum volume of 15 cubic feet. A microwave oven weighs 30 pounds and has 2 cubic feet of volume, while a printer weighs 20 pounds and has 3 cubic feet of space.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could 4 microwaves and 2 printers be carried on this trailer?
4. Could 7 microwaves and 3 printers be carried on this trailer?

Answer

1. $\begin{cases} 30m+20p \le 160 \\ 2m+3p \le 15 \end{cases}$
2. 

3. yes
4. no


Try It 5.6.21

Mary needs to purchase supplies of answer sheets and pencils for a standardized test to be given to the juniors at her high school. The number of the answer sheets needed is at least 5 more than the number of pencils. The pencils cost $2 and the answer sheets cost $1. Mary's budget for these supplies allows for a maximum cost of $400.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could Mary purchase 100 pencils and 100 answer sheets?
4. Could Mary purchase 150 pencils and 150 answer sheets?

Answer

1. $\begin{cases} a \ge p+5 \\ a+2p \le 400 \end{cases}$
2.



3. no

4. no

Example 5.6.22

Omar needs to eat at least 800 calories before going to his team practice. All he wants is hamburgers and cookies, and he doesn't want to spend more than $5. At the hamburger restaurant near his college, each hamburger has 240 calories and costs $1.40. Each cookie has 160 calories and costs $0.50.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could he eat 1 cookie and 3 hamburgers?
4. Could he eat 4 cookies and 2 hamburgers?

Solution

1.

Let h= the number of hamburgers.
c= the number of cookies
To find the system of inequalities, translate the information.
The calories from hamburgers at 240 calories each, plus the calories from cookies at 160 calories each must be more that 800.

$$240h + 160c \geq 800 \tag{5.6.2}$$

The amount spent on hamburgers at $1.40 each, plus the amount spent on cookies at $0.50 each must be no more than $5.00.


$$1.40h + 0.50c \leq 5 \tag{5.6.3}$$

We have our system of inequalities. $\begin{cases} 240h + 160c \geq 800 \\ 1.40h + 0.50c \leq 5 \end{cases}$

2.

To graph $240h + 160c \geq 800$ graph 240h+160c=800 as a solid line.
Choose (0, 0) as a test point. it does not make the inequality true.
So, shade (red) the side that does not include the point (0, 0).

To graph $1.40h + 0.50c \leq 5$, graph 1.40h+0.50c=5 as a solid line.
Choose (0,0) as a test point. It makes the inequality true. So, shade
(blue) the side that includes the point.



The solution of the system is the region of the graph that is double shaded and so is shaded darker.

3. To determine if 1 cookie and 3 hamburgers would meet Omar's criteria, we see if the point (1, 3) is in the solution region. It is not.
4. To determine if 4 cookies and 2 hamburgers would meet Omar's criteria, we see if the point (4, 2) is in the solution region. It is. He might choose to eat 4 cookies and 2 hamburgers.


We could also test the possible solutions by substituting the values into each inequality.

Try It 5.6.23

Tension needs to eat at least an extra 1,000 calories a day to prepare for running a marathon. He has only $25 to spend on the extra food he needs and will spend it on $0.75 donuts which have 360 calories each and $2 energy drinks which have 110 calories.

1. Write a system of inequalities that models this situation.
2. Graph the system.
3. Can he buy 8 donuts and 4 energy drinks?
4. Can he buy 1 donut and 3 energy drinks?

Answer

1. $\begin{cases} 0.75d+2e \le 25 \\ 360d+110e \ge 1000 \end{cases}$
2. 

3. yes
4. no


Try It 5.6.24

Philip's doctor tells him he should add at least 1000 more calories per day to his usual diet. Philip wants to buy protein bars that cost $1.80 each and have 140 calories and juice that costs $1.25 per bottle and have 125 calories. He doesn't want to spend more than $12.

1. Write a system of inequalities that models this situation.
2. Graph the system.
3. Can he buy 3 protein bars and 5 bottles of juice?
4. Can he buy 5 protein bars and 3 bottles of juice?

Answer

1. $\begin{cases} 140p+125j \ge 1000 \\ 1.80p+1.25j \le 12 \end{cases}$
2.



3. yes

4. no

Note

Access these online resources for additional instruction and practice with graphing systems of linear inequalities.

- Graphical System of Inequalities
- Systems of Inequalities
- Solving Systems of Linear Inequalities by Graphing

Key Concepts

- **To Solve a System of Linear Inequalities by Graphing**
  1. Graph the first inequality.
     - Graph the boundary line.
     - Shade in the side of the boundary line where the inequality is true.
  2. On the same grid, graph the second inequality.
     - Graph the boundary line.
     - Shade in the side of that boundary line where the inequality is true.
  3. The solution is the region where the shading overlaps.
  4. Check by choosing a test point.

Glossary

system of linear inequalities

Two or more linear inequalities grouped together form a system of linear inequalities.

This page titled 5.6: Graphing Systems of Linear Inequalities is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.6: Graphing Systems of Linear Inequalities** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

5.6E: Exercises

Practice Makes Perfect

Determine Whether an Ordered Pair is a Solution of a System of Linear Inequalities

In the following exercises, determine whether each ordered pair is a solution to the system.

Exercise 5.6E.1

$$\begin{cases} 3x+y>5 \\ 2x-y\leq 10 \end{cases}$$

1. (3,−3)
2. (7,1)

Answer

1. true
2. false

Exercise 5.6E.2

$$\begin{cases} 4x-y<10 \\ -2x+2y>-8 \end{cases}$$

1. (5,−2)
2. (−1,3)

Exercise 5.6E.3

$$\begin{cases} y>\frac{2}{3}x-5 \\ x+\frac{1}{2}y\leq 4 \end{cases}$$

1. (6,−4)
2. (3,0)

Answer

1. false
2. true

Exercise 5.6E.4

$$\begin{cases} y<\frac{3}{2}x+3 \\ \frac{3}{4}x-2y<5 \end{cases}$$

1. (−4,−1)
2. (8,3)

Exercise 5.6E.5

$$\begin{cases} 7x+2y>14 \\ 5x-y\leq 8 \end{cases}$$

1. (2,3)
2. (7,−1)

Answer

1. true

false

Exercise 5.6E.6

$$\begin{cases} 6x-5y<20 \\ -2x+7y>-8 \end{cases}$$

1. (1,−3)
2. (−4,4)

Exercise 5.6E.7

$$\begin{cases} 2x+3y\geq 2 \\ 4x-6y<-1 \end{cases}$$

1. $\left(\frac{3}{2},\frac{4}{3}\right)$
2. $\left(\frac{1}{4},\frac{7}{6}\right)$

Answer

1. true
2. true

Exercise 5.6E.8

$$\begin{cases} 5x-3y<-2 \\ 10x+6y>4 \end{cases}$$

1. $\left(\frac{1}{5},\frac{2}{3}\right)$
2. $\left(-\frac{3}{10},\frac{7}{6}\right)$


Solve a System of Linear Inequalities by Graphing

In the following exercises, solve each system by graphing.

Exercise 5.6E.9

$$\begin{cases} y\leq 3x+2 \\ y>x-1 \end{cases}$$

Answer



https://math.libretexts.org/@go/page/30530


Exercise 5.6E.10

$$\begin{cases} y < -2x + 2 \\ y \geq -x - 1 \end{cases}$$

Exercise 5.6E.11

$$\begin{cases} y < 2x - 1 \\ y \leq -\frac{1}{2}x + 4 \end{cases}$$

Answer




Exercise 5.6E.12

$$\begin{cases} y \geq -\frac{2}{3}x + 2 \\ y > 2x - 3 \end{cases}$$

Exercise 5.6E.13

$$\begin{cases} x - y > 1 \\ y < -\frac{1}{4}x + 3 \end{cases}$$

Answer



Exercise 5.6E.14


$$\begin{cases} x + 2y < 4 \\ y < x - 2 \end{cases}$$

Exercise 5.6E.15

$$\begin{cases} 3x - y \leq 6 \\ y \geq -\frac{1}{2}x \end{cases}$$

https://math.libretexts.org/@go/page/30530

Answer




Exercise 5.6E.16

$$\begin{cases} 2x+4y \geq 8 \\ y \leq \frac{3}{4}x \end{cases}$$

Exercise 5.6E.17

$$\begin{cases} 2x-5y < 10 \\ 3x+4y \geq 12 \end{cases}$$

Answer




Exercise 5.6E.18

$$\begin{cases} 3x-2y \leq 6 \\ -4x-2y > 8 \end{cases}$$

Exercise 5.6E.19

$$\begin{cases} 2x+2y > -4 \\ -x+3y \geq 9 \end{cases}$$

Answer



https://math.libretexts.org/@go/page/30530


Exercise 5.6E. 20

$$\begin{cases} 2x+y>-6 \\ -x+2y \geq -4 \end{cases}$$

Exercise 5.6E. 21

$$\begin{cases} x-2y<3 \\ y \leq 1 \end{cases}$$

Answer




Exercise 5.6E. 22

$$\begin{cases} x-3y>4 \\ y \leq -1 \end{cases}$$

Exercise 5.6E. 23

$$\begin{cases} y \geq -\frac{1}{2}x-3 \\ x \leq 2 \end{cases}$$

Answer



Exercise 5.6E. 24


$$\begin{cases} y \leq -\frac{2}{3}x+5 \\ x \geq 3 \end{cases}$$

Exercise 5.6E. 25

$$\begin{cases} y \geq \frac{3}{4}x-2 \\ y<2 \end{cases}$$

Answer

https://math.libretexts.org/@go/page/30530




Exercise 5.6E.26

$$\begin{cases} y \leq -\frac{1}{2}x + 3 \\ y < 1 \end{cases}$$

Exercise 5.6E.27

$$\begin{cases} 3x - 4y < 8 \\ x < 1 \end{cases}$$

Answer




Exercise 5.6E.28

$$\begin{cases} -3x + 5y > 10 \\ x > -1 \end{cases}$$

Exercise 5.6E.29

$$\begin{cases} x \geq 3 \\ y \leq 2 \end{cases}$$

Answer



https://math.libretexts.org/@go/page/30530


Exercise 5.6E.30

$$\begin{cases} x \leq -1 \\ y \geq 3 \end{cases}$$

Exercise 5.6E.31

$$\begin{cases} 2x + 4y > 4 \\ y \leq -\frac{1}{2}x - 2 \end{cases}$$

Answer



Exercise 5.6E.32


$$\begin{cases} x - 3y \geq 6 \\ y > \frac{1}{3}x + 1 \end{cases}$$

Exercise 5.6E.33

$$\begin{cases} -2x + 6y < 0 \\ 6y > 2x + 4 \end{cases}$$

Answer

No solution




Exercise 5.6E.34

$$\begin{cases} -3x + 6y > 12 \\ 4y \leq 2x - 4 \end{cases}$$

Exercise 5.6E.35

$$\begin{cases} y \geq -3x + 2 \\ 3x + y > 5 \end{cases}$$

Answer



Exercise 5.6E.36


$$\begin{cases} y \geq \frac{1}{2}x - 1 \\ -2x + 4y \geq 4 \end{cases}$$

Exercise 5.6E.37

$$\begin{cases} y \leq -\frac{1}{4}x - 2 \\ x + 4y < 6 \end{cases}$$

Answer

x+4y<6



Exercise 5.6E.38

$$\begin{cases} y \geq 3x - 1 \\ -3x + y > -4 \end{cases}$$


Exercise 5.6E.39

$$\begin{cases} 3y > x + 2 \\ -2x + 6y > 8 \end{cases}$$

Answer

$-2x + 6y > 8$

https://math.libretexts.org/@go/page/30530



Exercise 5.6E. 40

$$\begin{cases} y < \frac{3}{4}x - 2 \\ -3x + 4y < 7 \end{cases}$$

Solve Applications of Systems of Inequalities


In the following exercises, translate to a system of inequalities and solve.

Exercise 5.6E. 41

Caitlyn sells her drawings at the county fair. She wants to sell at least 60 drawings and has portraits and landscapes. She sells the portraits for $15 and the landscapes for $10. She needs to sell at least $800 worth of drawings in order to earn a profit.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Will she make a profit if she sells 20 portraits and 35 landscapes?
4. Will she make a profit if she sells 50 portraits and 20 landscapes?

Answer

1. $\begin{cases} p + l \geq 60 \\ 15p + 10l \geq 800 \end{cases}$
2. 

3. No
4. Yes

Exercise 5.6E. 42

Jake does not want to spend more than $50 on bags of fertilizer and peat moss for his garden. Fertilizer costs $2 a bag and peat moss costs $5 a bag. Jake's van can hold at most 20 bags.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Can he buy 15 bags of fertilizer and 4 bags of peat moss?

https://math.libretexts.org/@go/page/30530


4. Can he buy 10 bags of fertilizer and 10 bags of peat moss?

Exercise $5.6E.43$

Reiko needs to mail her Christmas cards and packages and wants to keep her mailing costs to no more than $500. The number of cards is at least 4 more than twice the number of packages. The cost of mailing a card (with pictures enclosed) is $3 and for a package the cost is $7.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Can she mail 60 cards and 26 packages?
4. Can she mail 90 cards and 40 packages?

Answer

1. $\begin{cases} 7p+3c \leq 500 \\ p \geq 2c+4 \end{cases}$
2. 

3. Yes
4. No

Exercise $5.6E.44$

Juan is studying for his final exams in Chemistry and Algebra. He knows he only has 24 hours to study, and it will take him at least three times as long to study for Algebra than Chemistry.


1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Can he spend 4 hours on Chemistry and 20 hours on Algebra?
4. Can he spend 6 hours on Chemistry and 18 hours on Algebra?

Exercise $5.6E.45$

Jocelyn is pregnant and needs to eat at least 500 more calories a day than usual. When buying groceries one day with a budget of $15 for the extra food, she buys bananas that have 90 calories each and chocolate granola bars that have 150 calories each. The bananas cost $0.35 each and the granola bars cost $2.50 each.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could she buy 5 bananas and 6 granola bars?
4. Could she buy 3 bananas and 4 granola bars?

Answer

1. $\begin{cases} 90b+150g \geq 500 \\ 0.35b+2.50g \leq 15 \end{cases}$
2. 

3. No
4. Yes

? Exercise $5.6E.46$

Mark is attempting to build muscle mass and so he needs to eat at least an additional 80 grams of protein a day. A bottle of protein water costs $3.20 and a protein bar costs $1.75. The protein water supplies 27 grams of protein and the bar supplies 16 gram. If he has $ 10 dollars to spend


1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could he buy 3 bottles of protein water and 1 protein bar?
4. Could he buy no bottles of protein water and 5 protein bars?

? Exercise $5.6E.47$

Jocelyn desires to increase both her protein consumption and caloric intake. She desires to have at least 35 more grams of protein each day and no more than an additional 200 calories daily. An ounce of cheddar cheese has 7 grams of protein and 110 calories. An ounce of parmesan cheese has 11 grams of protein and 22 calories.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could she eat 1 ounce of cheddar cheese and 3 ounces of parmesan cheese?
4. Could she eat 2 ounces of cheddar cheese and 1 ounce of parmesan cheese?

Answer

1. $\begin{cases} 7c+11p \geq 35 \\ 110c+22p \leq 200 \end{cases}$
2. 

3. Yes
4. No

Exercise 5.6E.48

Mark is increasing his exercise routine by running and walking at least 4 miles each day. His goal is to burn a minimum of 1,500 calories from this exercise. Walking burns 270 calories/mile and running burns 650 calories.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could he meet his goal by walking 3 miles and running 1 mile?
4. Could he meet his goal by walking 2 miles and running 2 mile?

Everyday Math

Exercise 5.6E.49

Tickets for an American Baseball League game for 3 adults and 3 children cost less than $75, while tickets for 2 adults and 4 children cost less than $62.


1. Write a system of inequalities to model this problem.
2. Graph the system.
3. Could the tickets cost $20 for adults and $8 for children?
4. Could the tickets cost $15 for adults and $5 for children?

Answer

1. \(\left\{

$$\begin{aligned} 3a+3c &< 75 \\ 2a+4c &< 62 \end{aligned} \tag{5.6E.1}$$

\right.\

2. 

3. No
4. Yes

Exercise 5.6E.50

Grandpa and Grandma are treating their family to the movies. Matinee tickets cost $4 per child and $4 per adult. Evening tickets cost $6 per child and $8 per adult. They plan on spending no more than $80 on the matinee tickets and no more than $100 on the evening tickets.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Could they take 9 children and 4 adults to both shows?
4. Could they take 8 children and 5 adults to both shows?

Writing Exercises

Exercise $5.6E.51$

Graph the inequality $x - y \geq 3$. How do you know which side of the line $x - y = 3$ should be shaded?

Answer

Answers will vary.

Exercise $5.6E.52$

Graph the system $\begin{cases} x + 2y \leq 6 \\ y \geq -\frac{1}{2}x - 4 \end{cases}$. What does the solution mean?

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
determine whether an ordered pair is a solution of a system of linear inequalities.			
solve a system of linear inequalities by graphing.			
solve applications of systems of inequalities.			

b. After reviewing this checklist, what will you do to become confident for all objectives?

This page titled 5.6E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **5.6E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 5 Review Exercises

Chapter 5 Review Exercises

Solve Systems of Equations by Graphing

Determine Whether an Ordered Pair is a Solution of a System of Equations.

In the following exercises, determine if the following points are solutions to the given system of equations.

Exercise 1

$$\begin{cases} x+3y=-9 \\ 2x-4y=12 \end{cases}$$

1. $(-3,-2)$
2. $(0,-3)$

Answer

1. no
2. yes

Exercise 2

$$\begin{cases} x+y=8 \\ y=x-4 \end{cases}$$

1. $(6,2)$
2. $(9,-1)$

Solve a System of Linear Equations by Graphing


In the following exercises, solve the following systems of equations by graphing.

Exercise 3

$$\begin{cases} 3x+y=6 \\ x+3y=-6 \end{cases}$$

Answer

$(3,-3)$



Exercise 4


$$\begin{cases} y=x-2 \\ y=-2x-2 \end{cases}$$

Exercise 5

$$\begin{cases} 2x - y = 6 \\ y = 4 \end{cases}$$

Answer

(5,4)



Exercise 6


$$\begin{cases} x + 4y = -1 \\ x = 3 \end{cases}$$

Exercise 7

$$\begin{cases} 2x - y = 5 \\ 4x - 2y = 10 \end{cases}$$

Answer

coincident lines



Exercise 8

$$\begin{cases} -x + 2y = 4 \\ y = \frac{1}{2}x - 3 \end{cases}$$

Determine the Number of Solutions of a Linear System

In the following exercises, without graphing determine the number of solutions and then classify the system of equations.

Exercise 9

$$\begin{cases} y = \frac{2}{5}x + 2 \\ -2x + 5y = 10 \end{cases}$$

Answer

infinitely many solutions, consistent system, dependent equations

Exercise 10

$$\begin{cases} 3x+2y=6 \\ y=-3x+4 \end{cases}$$

Exercise 11

$$\begin{cases} 5x-4y=0 \\ y=\frac{5}{4}x-5 \end{cases}$$

Answer

no solutions, inconsistent system, independent equations

Exercise 12

$$\begin{cases} y=-\frac{3}{4}x+1 \\ 6x+8y=8 \end{cases}$$

Solve Applications of Systems of Equations by Graphing

Exercise 13

LaVelle is making a pitcher of caffe mocha. For each ounce of chocolate syrup, she uses five ounces of coffee. How many ounces of chocolate syrup and how many ounces of coffee does she need to make 48 ounces of caffe mocha?

Answer

LaVelle needs 8 ounces of chocolate syrup and 40 ounces of coffee.

Exercise 14

Eli is making a party mix that contains pretzels and chex. For each cup of pretzels, he uses three cups of chex. How many cups of pretzels and how many cups of chex does he need to make 12 cups of party mix?

Solve Systems of Equations by Substitution

Solve a System of Equations by Substitution

In the following exercises, solve the systems of equations by substitution.

Exercise 15

$$\begin{cases} 3x-y=-5 \\ y=2x+4 \end{cases}$$

Answer

$(-1,2)$

Exercise 16

$$\begin{cases} 3x-2y=2 \\ y=\frac{1}{2}x+3 \end{cases}$$

Exercise 17

$$\begin{cases} x-y=0 \\ 2x+5y=-14 \end{cases}$$

Answer

$(-2,-2)$

Exercise 18

$$\begin{cases} y=-2x+7 \\ y=\frac{2}{3}x-1 \end{cases}$$

Exercise 19

$$\begin{cases} y=-5x \\ 5x+y=6 \end{cases}$$

Answer

no solution

Exercise 20

$$\begin{cases} y=-\frac{1}{3}x+2 \\ x+3y=6 \end{cases}$$

Solve Applications of Systems of Equations by Substitution

In the following exercises, translate to a system of equations and solve.

Exercise 21

The sum of two number is 55. One number is 11 less than the other. Find the numbers.

Answer

The numbers are 22 and 33.

Exercise 22

The perimeter of a rectangle is 128. The length is 16 more than the width. Find the length and width.

Exercise 23

The measure of one of the small angles of a right triangle is 2 less than 3 times the measure of the other small angle. Find the measure of both angles.

Answer

The measures are 23 degrees and 67 degrees.

Exercise 24

Gabriela works for an insurance company that pays her a salary of $32,000 plus a commission of $100 for each policy she sells. She is considering changing jobs to a company that would pay a salary of $40,000 plus a commission of $80 for each policy sold. How many policies would Gabriela need to sell to make the total pay the same?

Solve Systems of Equations by Elimination

Solve a System of Equations by Elimination In the following exercises, solve the systems of equations by elimination.

Exercise 25

$$\begin{cases} x+y=12 \\ x-y=-10 \end{cases}$$

Answer

(1,11)

Exercise 26

$$\begin{cases} 4x+2y=2 \\ -4x-3y=-9 \end{cases}$$

Exercise 27

$$\begin{cases} 3x-8y=20 \\ x+3y=1 \end{cases}$$

Answer

(4,−1)

Exercise 28

$$\begin{cases} 3x-2y=6 \\ 4x+3y=8 \end{cases}$$

Exercise 29

$$\begin{cases} 9x+4y=2 \\ 5x+3y=5 \end{cases}$$

Answer

(−2,5)

Exercise 30

$$\begin{cases} -x+3y=8 \\ 2x-6y=-20 \end{cases}$$

Solve Applications of Systems of Equations by Elimination

In the following exercises, translate to a system of equations and solve.

Exercise 31

The sum of two numbers is −90. Their difference is 16. Find the numbers.

Answer

The numbers are −37 and −53.

Exercise 32

Omar stops at a donut shop every day on his way to work. Last week he had 8 donuts and 5 cappuccinos, which gave him a total of 3,000 calories. This week he had 6 donuts and 3 cappuccinos, which was a total of 2,160 calories. How many calories are in one donut? How many calories are in one cappuccino?

Choose the Most Convenient Method to Solve a System of Linear Equations

In the following exercises, decide whether it would be more convenient to solve the system of equations by substitution or elimination.

Exercise 33

$$\begin{cases} 6x-5y=27 \\ 3x+10y=-24 \end{cases}$$

Answer

elimination

Exercise 34

$$\begin{cases} y=3x-9 \\ 4x-5y=23 \end{cases}$$

Solve Applications with Systems of Equations

Translate to a System of Equations

In the following exercises, translate to a system of equations. Do not solve the system.

Exercise 35

The sum of two numbers is −32. One number is two less than twice the other. Find the numbers.

Answer

$$\begin{cases} x+y=-32 \\ x=2y-2 \end{cases}$$

Exercise 36

Four times a number plus three times a second number is −9. Twice the first number plus the second number is three. Find the numbers.

Exercise 37

Last month Jim and Debbie earned $7,200. Debbie earned $1,600 more than Jim earned. How much did they each earn?

Answer

$$\begin{cases} j+d=7200 \\ d=j+1600 \end{cases}$$

Exercise 38

Henri has $24,000 invested in stocks and bonds. The amount in stocks is $6,000 more than three times the amount in bonds. How much is each investment?

Solve Direct Translation Applications

In the following exercises, translate to a system of equations and solve.

Exercise 39

Pam is 3 years older than her sister, Jan. The sum of their ages is 99. Find their ages.

Answer

Pam is 51 and Jan is 48.

Exercise 40

Mollie wants to plant 200 bulbs in her garden. She wantsall irises and tulips. She wants to plant three times as many tulips as irises. How many irises and how many tulips should she plant?

Solve Geometry Applications

In the following exercises, translate to a system of equations and solve.

Exercise 41

The difference of two supplementary angles is 58 degrees. Find the measures of the angles.

Answer

The measures are 119 degrees and 61 degrees.

Exercise 42

Two angles are complementary. The measure of the larger angle is five more than four times the measure of the smaller angle. Find the measures of both angles.

Exercise 43

Becca is hanging a 28 foot floral garland on the two sides and top of a pergola to prepare for a wedding. The height is four feet less than the width. Find the height and width of the pergola.

Answer

The pergola is 8 feet high and 12 feet wide.

Exercise 44

The perimeter of a city rectangular park is 1428 feet. The length is 78 feet more than twice the width. Find the length and width of the park.

Solve Uniform Motion Applications

In the following exercises, translate to a system of equations and solve.

Exercise 45

Sheila and Lenore were driving to their grandmother's house. Lenore left one hour after Sheila. Sheila drove at a rate of 45 mph, and Lenore drove at a rate of 60 mph. How long will it take for Lenore to catch up to Sheila?

Answer

It will take Lenore 3 hours.

Exercise 46

Bob left home, riding his bike at a rate of 10 miles per hour to go to the lake. Cheryl, his wife, left 45 minutes ($\frac{3}{4}$ hour) later, driving her car at a rate of 25 miles per hour. How long will it take Cheryl to catch up to Bob?

Exercise 47

Marcus can drive his boat 36 miles down the river in three hours but takes four hours to return upstream. Find the rate of the boat in still water and the rate of the current.

Answer

The rate of the boat is 10.5 mph. The rate of the current is 1.5 mph.

Exercise 48

A passenger jet can fly 804 miles in 2 hours with a tailwind but only 776 miles in 2 hours into a headwind. Find the speed of the jet in still air and the speed of the wind.

Solve Mixture Applications with Systems of Equations

Solve Mixture Applications

In the following exercises, translate to a system of equations and solve.

Exercise 49

Lynn paid a total of $2,780 for 261 tickets to the theater. Student tickets cost $10 and adult tickets cost $15. How many student tickets and how many adult tickets did Lynn buy?

Answer

Lynn bought 227 student tickets and 34 adult tickets.

Exercise 50

Priam has dimes and pennies in a cup holder in his car. The total value of the coins is $4.21. The number of dimes is three less than four times the number of pennies. How many dimes and how many pennies are in the cup?

Exercise 51

Yumi wants to make 12 cups of party mix using candies and nuts. Her budget requires the party mix to cost her $1.29 per cup. The candies are $2.49 per cup and the nuts are $0.69 per cup. How many cups of candies and how many cups of nuts should she use?

Answer

Yumi should use 4 cups of candies and 8 cups of nuts.

Exercise 52

A scientist needs 70 liters of a 40% solution of alcohol. He has a 30% and a 60% solution available. How many liters of the 30% and how many liters of the 60% solutions should he mix to make the 40% solution?

Solve Interest Applications

In the following exercises, translate to a system of equations and solve.

Exercise 53

Jack has $12,000 to invest and wants to earn 7.5% interest per year. He will put some of the money into a savings account that earns 4% per year and the rest into CD account that earns 9% per year. How much money should he put into each account?

Answer

Jack should put $3600 into savings and $8400 into the CD.

Exercise 54

When she graduates college, Linda will owe $43,000 in student loans. The interest rate on the federal loans is 4.5% and the rate on the private bank loans is 2%. The total interest she owes for one year was $1585. What is the amount of each loan?

Graphing Systems of Linear Inequalities

Determine Whether an Ordered Pair is a Solution of a System of Linear Inequalities

In the following exercises, determine whether each ordered pair is a solution to the system.

Exercise 55

$$\begin{cases} 4x+y>6 \\ 3x-y\leq 12 \end{cases}$$

1. (2,−1)
2. (3,−2)

Answer

1. yes
2. no

Exercise 56

$$\begin{cases} y>\frac{1}{3}x+2 \\ x-\frac{1}{4}y\leq 10 \end{cases}$$

1. (6,5)
2. (15,8)


Solve a System of Linear Inequalities by Graphing

In the following exercises, solve each system by graphing.

Exercise 57

$$\begin{cases} y < 3x + 1 \\ y \geq -x - 2 \end{cases}$$

Answer




Exercise 58

$$\begin{cases} x - y > -1 \\ y < \frac{1}{3}x - 2 \end{cases}$$

Exercise 59

$$\begin{cases} 2x - 3y < 6 \\ 3x + 4y \geq 12 \end{cases}$$

Answer



Exercise 60


$$\begin{cases} y \leq -\frac{3}{4}x + 1 \\ x \geq -5 \end{cases}$$

Exercise 61

$$\begin{cases} x + 3y < 5 \\ y \geq -\frac{1}{3}x + 6 \end{cases}$$

Answer

No solution



Exercise 62

$$\begin{cases} y \geq 2x - 5 \\ -6x + 3y > -4 \end{cases}$$

Solve Applications of Systems of Inequalities

In the following exercises, translate to a system of inequalities and solve.


Exercise 63

Roxana makes bracelets and necklaces and sells them at the farmers' market. She sells the bracelets for $12 each and the necklaces for $18 each. At the market next weekend she will have room to display no more than 40 pieces, and she needs to sell at least $500 worth in order to earn a profit.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Should she display 26 bracelets and 14 necklaces?
4. Should she display 39 bracelets and 1 necklace?

Answer

1. $\begin{cases} b + n \leq 40 \\ 12b + 18n \geq 500 \end{cases}$
2.



3. yes
4. no

Exercise 64

Annie has a budget of $600 to purchase paperback books and hardcover books for her classroom. She wants the number of hardcover to be at least 5 more than three times the number of paperback books. Paperback books cost $4 each and hardcover books cost $15 each.

1. Write a system of inequalities to model this situation.
2. Graph the system.

3. Can she buy 8 paperback books and 40 hardcover books?
4. Can she buy 10 paperback books and 37 hardcover books?

Practice Test

? Exercise 1

$$\begin{cases} x-4y=-8 \\ 2x+5y=10 \end{cases}$$

1. (0,2)
2. (4,3)

Answer

1. yes
2. no

In the following exercises, solve the following systems by graphing.


? Exercise 2

$$\begin{cases} x-y=5 \\ x+2y=-4 \end{cases}$$

? Exercise 3

$$\begin{cases} x-y>-2 \\ y\leq 3x+1 \end{cases}$$

Answer



In the following exercises, solve each system of equations. Use either substitution or elimination.

? Exercise 4

$$\begin{cases} 3x-2y=3 \\ y=2x-1 \end{cases}$$

? Exercise 5

$$\begin{cases} x+y=-3 \\ x-y=11 \end{cases}$$

Answer

(4,−7)

Exercise 6

$$\begin{cases} 4x-3y=7 \\ 5x-2y=0 \end{cases}$$

Exercise 7

$$\begin{cases} y=-\frac{4}{5}x+1 \\ 8x+10y=10 \end{cases}$$

Answer

infinitely many solutions

Exercise 8

$$\begin{cases} 2x+3y=12 \\ -4x+6y=-16 \end{cases}$$

In the following exercises, translate to a system of equations and solve.

Exercise 9

The sum of two numbers is −24. One number is 104 less than the other. Find the numbers.

Answer

The numbers are 40 and 64

Exercise 10

Ramon wants to plant cucumbers and tomatoes in his garden. He has room for 16 plants, and he wants to plant three times as many cucumbers as tomatoes. How many cucumbers and how many tomatoes should he plant?

Exercise 11

Two angles are complementary. The measure of the larger angle is six more than twice the measure of the smaller angle. Find the measures of both angles.

Answer

The measures of the angles are 28 degrees and 62 degrees.

Exercise 12

On Monday, Lance ran for 30 minutes and swam for 20 minutes. His fitness app told him he had burned 610 calories. On Wednesday, the fitness app told him he burned 695 calories when he ran for 25 minutes and swam for 40 minutes. How many calories did he burn for one minute of running? How many calories did he burn for one minute of swimming?

Exercise 13

Kathy left home to walk to the mall, walking quickly at a rate of 4 miles per hour. Her sister Abby left home 15 minutes later and rode her bike to the mall at a rate of 10 miles per hour. How long will it take Abby to catch up to Kathy?

Answer

It will take Kathy $\frac{1}{6}$ of an hour (or 10 minutes)

Exercise 14

It takes $5\frac{1}{2}$ hours for a jet to fly 2,475 miles with a headwind from San Jose, California to Lihue, Hawaii. The return flight from Lihue to San Jose with a tailwind, takes 5 hours. Find the speed of the jet in still air and the speed of the wind.

Exercise 15

Liz paid $160 for 28 tickets to take the Brownie troop to the science museum. Children's tickets cost $5 and adult tickets cost $9. How many children's tickets and how many adult tickets did Liz buy?

Answer

Liz bought 23 children's tickets and 5 adult tickets.

Exercise 16

A pharmacist needs 20 liters of a 2% saline solution. He has a 1% and a 5% solution available. How many liters of the 1% and how many liters of the 5% solutions should she mix to make the 2% solution?


Exercise 1

Translate to a system of inequalities and solve.

Andi wants to spend no more than $50 on Halloween treats. She wants to buy candy bars that cost $1 each and lollipops that cost $0.50 each, and she wants the number of lollipops to be at least three times the number of candy bars.

1. Write a system of inequalities to model this situation.
2. Graph the system.
3. Can she buy 20 candy bars and 70 lollipops?
4. Can she buy 15 candy bars and 65 lollipops?

Answer

1. $\begin{cases} C+0.5L \leq 50 \\ L \geq 3C \end{cases}$
2. 

3. No
4. Yes

Chapter 5 Review Exercises is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **Chapter 5 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

CHAPTER OVERVIEW

6: Polynomials

- 6.1: Add and Subtract Polynomials
  - 6.1E: Exercises
- 6.2: Use Multiplication Properties of Exponents
  - 6.2E: Exercises
- 6.3: Multiply Polynomials
  - 6.3E: Exercises
- 6.4: Special Products
  - 6.4E: Exercises
- 6.5: Divide Monomials
  - 6.5E: Exercises
- 6.6: Divide Polynomials
  - 6.6E: Exercises
- 6.7: Integer Exponents and Scientific Notation
  - 6.7E: Exercises
- Chapter 6 Review Exercises

This page titled 6: Polynomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

6.1: Add and Subtract Polynomials

Learning Objectives

By the end of this section, you will be able to:

- Identify polynomials, monomials, binomials, and trinomials
- Determine the degree of polynomials
- Add and subtract monomials
- Add and subtract polynomials
- Evaluate a polynomial for a given value

Quiz

Before you get started, take this readiness quiz.

1. Simplify: $8x+3x$.
   If you missed this problem, review Example 1.3.37.
2. Subtract: $(5n+8)-(2n-1)$.
   If you missed this problem, review Example 1.10.52.
3. Write in expanded form: a^5.
   If you missed this problem, review Example 1.3.7.

Identify Polynomials, Monomials, Binomials and Trinomials

You have learned that a *term* is a constant or the product of a constant and one or more variables. When it is of the form ax^m, where a is a constant and m is a whole number, it is called a *monomial*. Some examples of monomial are 8, $-2x^2$, $4y^3$, and $11z^7$.

Definition: Monomials

A **monomial** is a term of the form ax^m, where a is a constant and m is a positive whole number.

A monomial, or two or more monomials combined by addition or subtraction, is a polynomial. Some polynomials have special names, based on the number of terms. A monomial is a polynomial with exactly one term. A binomial has exactly two terms, and a trinomial has exactly three terms. There are no special names for polynomials with more than three terms.

Definitions: Polynomials

- **polynomial**—A monomial, or two or more monomials combined by addition or subtraction, is a *polynomial*.
- **monomial**—A polynomial with exactly one term is called a *monomial*.
- **binomial**—A polynomial with exactly two terms is called a *binomial*.
- **trinomial**—A polynomial with exactly three terms is called a *trinomial*.

Here are some examples of polynomials.

Polynomial	$b+1$	$4y^2-7y+2$	$4x^4+x^3+8x^2-9x+1$	
Monomial	14	$8y^2$	$-9x^3y^5$	-13
Binomial	$a+7$	$4b-5$	y^2-16	$3x^3-9x^2$
Trinomial	$x^2-7x+12$	$9y^2+2y-8$	$6m^4-m^3+8m$	z^4+3z^2-1

Notice that every monomial, binomial, and trinomial is also a polynomial. They are just special members of the “family” of polynomials and so they have special names. We use the words *monomial*, *binomial*, and *trinomial* when referring to these special polynomials and just call all the rest *polynomials*.

Example 6.1.1

Determine whether each polynomial is a monomial, binomial, trinomial, or other polynomial.

a. $4y^2 - 8y - 6$
b. $-5a^4b^2$
c. $2x^5 - 5x^3 - 3x + 4$
d. $13 - 5m^3$
e. q

Solution

	Polynomial	Number of terms	Type
(a)	$4y^2 - 8y - 6$	3	Trinomial
(b)	$-5a^4b^2$	1	Monomial
(c)	$2x^5 - 5x^3 - 9x^2 + 3x + 4$	5	Ponomial
(d)	$13 - 5m^3$	2	Binomial
(e)	q	1	Monomial

Try It 6.1.2

Determine whether each polynomial is a monomial, binomial, trinomial, or other polynomial:

a. 5b
b. $8y^3 - 7y^2 - y - 3$
c. $-3x^2 - 5x + 9$
d. $81 - 4a^2$
e. $-5x^6$

Answer

1. monomial
2. polynomial
3. trinomial
4. binomial
5. monomial

Try It 6.1.3

Determine whether each polynomial is a monomial, binomial, trinomial, or other polynomial:

a. $27z^3 - 8$
b. $12m^3 - 5m^2 - 2m$
c. $\frac{5}{6}$
d. $8x^4 - 7x^2 - 6x - 5$
e. $-n^4$

Answer

a. binomial
b. trinomial
c. monomial
d. polynomial
e. monomial

Determine the Degree of Polynomials

The degree of a polynomial and the degree of its terms are determined by the exponents of the variable. A **monomial** that has no variable, just a constant, is a special case. The degree of a constant is 0, i.e., it has no variable.

Definition: Degree of a Polynomial

- The **degree of a term** is the sum of the exponents of its variables.
- The **degree of a constant** is 0.
- The **degree of a polynomial** is the highest degree of all its terms.

Let's see how this works by looking at several polynomials. We'll take it step by step, starting with monomials, and then progressing to polynomials with more terms.

Monomial	14	$8y^2$	$-9x^3y^6$	$-13a$
Degree	0	2	8	1
Binomial	$a+7$	$4b^2-5b$	x^2y^2-16	$3n^3-9n^2$
Degree of each term	1 0	2 1	4 0	3 2
Degree of polynomial	1	2	4	3
Trinomial	$x^2-7x+12$	$9a^2+6ab+b^2$	$6m^4-m^3n^2+8mn^5$	z^4+3z^2-1
Degree of each term	2 1 0	2 2 2	4 5 6	4 2 0
Degree of polynomial	2	2	6	4
Polynomial	$b+1$	$4y^2-7y+2$	$4x^4+x^3+8x^2-9x+1$	
Degree of each term	1 0	2 1 0	4 3 2 1 0	
Degree of polynomial	1	2	4	

A polynomial is in **standard form** when the terms of a polynomial are written in descending order of degrees. Get in the habit of writing the term with the highest degree first.

Example 6.1.4

Find the degree of the following polynomials.

a. 10y
b. $4x^3-7x+5$
c. -15
d. $-8b^2+9b-2$
e. $8xy^2+2y$

Solution

a. The exponent of y is one. $y=y^1$ $10y$ The degree is 1.
b. The highest degree of all the terms is 3. $4x^3-7x+5$ The degree is 3.
c. The degree of a constant is 0. -15 The degree is 0.
d. The highest degree of all the terms is 2. $-8b^2+9b-2$ The degree is 2.

e.	$8xy^2+2y$
The highest degree of all the terms is 3.	The degree is 3.

? Try It 6.1.5

Find the degree of the following polynomials:

a. $-15b$
b. $10z^4+4z^2-5$
c. $12c^5d^4+9c^3d^9-7$
d. $3x^2y-4x$
e. -9

Answer

a. 1
b. 4
c. 12
d. 3
e. 0

? Try It 6.1.6

Find the degree of the following polynomials:

a. 52
b. a^4b-17a^4
c. $5x+6y+2z$
d. $3x^2-5x+7$
e. $-a^3$

Answer

a. 0
b. 5
c. 1
d. 2
e. 3

Add and Subtract Monomials

You have learned how to simplify expressions by combining like terms. Remember, like terms must have the same variables with the same exponent. Since monomials are terms, adding and subtracting monomials is the same as combining like terms. If the monomials are like terms, we just combine them by adding or subtracting the coefficient.

✔ Example 6.1.7

Add: $25y^2+15y^2$

Solution

	$25y^2+15y^2$
Combine like terms.	$40y^2$

Try It 6.1.8

Add: $12q^2 + 9q^2$

Answer

$21q^2$

Try It 6.1.9

Add: $-15c^2 + 8c^2$

Answer

$-7c^2$

Example 6.1.10

Subtract: 16p−(−7p)

Solution

	$16p - (-7p)$
Combine like terms.	$23p$

Try It 6.1.11

Subtract: 8m−(−5m).

Answer

13m

Try It 6.1.12

Subtract: $-15z^3 - \left(-5z^3\right)$

Answer

$-10z^3$

Remember that like terms must have the same variables with the same exponents.

Example 6.1.13

Simplify: $c^2 + 7d^2 - 6c^2$

Solution

	$c^2 + 7d^2 - 6c^2$
Combine like terms.	$-5c^2 + 7d^2$

Try It 6.1.14

Add: $8y^2 + 3z^2 - 3y^2$

Answer

$5y^2 + 3z^2$

Try It 6.1.15

Add: $3m^2+n^2-7m^2$

Answer

$-4m^2+n^2$

Example 6.1.16

Simplify: $u^2v+5u^2-3v^2$

Solution

	$u^2v+5u^2-3v^2$
There are no like terms to combine.	$u^2v+5u^2-3v^2$

Try It 6.1.17

Simplify: $m^2n^2-8m^2+4n^2$

Answer

There are no like terms to combine.

Try It 6.1.18

Simplify: $pq^2-6p-5q^2$

Answer

There are no like terms to combine.

Add and Subtract Polynomials

We can think of adding and subtracting polynomials as just adding and subtracting a series of monomials. Look for the like terms—those with the same variables and the same exponent. The Commutative Property allows us to rearrange the terms to put like terms together.

Example 6.1.19

Find the sum: $\left(5y^2-3y+15\right)+\left(3y^2-4y-11\right)$

Solution

Identify like terms.	$(5y^2-3y+15)+(3y^2-4y-11)$
Rearrange to get the like terms together.	$5y^2+3y^2-3y-4y+15-11$
Combine like terms.	$8y^2-7y+4$

Try It 6.1.20

Find the sum: $\left(7x^2-4x+5\right)+\left(x^2-7x+3\right)$

Answer

$8x^2-11x+1$

Try It 6.1.21

Find the sum: $\left(14y^2+6y-4\right)+\left(3y^2+8y+5\right)$

Answer

$17y^2+14y+1$

Example 6.1.22

Find the difference: $\left(9w^2-7w+5\right)-\left(2w^2-4\right)$

Solution

	$(9w^2-7w+5)-(2w^2-4)$
Distribute and identify like terms.	$9w^2-7w+5-2w^2+4$
Rearrange the terms.	$9w^2-2w^2-7w+5+4$
Combine like terms.	$7w^2-7w+9$

Try It 6.1.23

Find the difference: $\left(8x^2+3x-19\right)-\left(7x^2-14\right)$

Answer

$15x^2+3x-5$

Try It 6.1.24

Find the difference: $\left(9b^2-5b-4\right)-\left(3b^2-5b-7\right)$

Answer

$6b^2+3$

Example 6.1.25

Subtract: $\left(c^2-4c+7\right)$ from $\left(7c^2-5c+3\right)$

Solution

	Subtract (c^2-4c+7) from $(7c^2-5c+3)$.
	$(7c^2-5c+3)-(c^2-4c+7)$
Distribute and identify like terms.	$7c^2-5c+3-c^2+4c-7$
Rearrange the terms.	$7c^2-c^2-5c+4c+3-7$
Combine like terms.	$6c^2-c-4$

Try It 6.1.26

Subtract: $\left(5z^2-6z-2\right)$ from $\left(7z^2+6z-4\right)$

Answer

$2z^2+12z-2$

Try It 6.1.27

Subtract: $\left(x^2-5x-8\right)$ from $\left(6x^2+9x-1\right)$

Answer

$5x^2+14x+7$

Example 6.1.28

Find the sum: $\left(u^2-6uv+5v^2\right)+\left(3u^2+2uv\right)$

Solution

	$\left(u^2-6uv+5v^2\right)+\left(3u^2+2uv\right)$
Distribute.	$u^2-6uv+5v^2+3u^2+2uv$
Rearrange the terms, to put like terms together	$u^2+3u^2-6uv+2uv+5v^2$
Combine like terms.	$4u^2-4uv+5v^2$

Try It 6.1.29

Find the sum: $\left(3x^2-4xy+5y^2\right)+\left(2x^2-xy\right)$

Answer

$5x^2-5xy+5y^2$

Try It 6.1.30

Find the sum: $\left(2x^2-3xy-2y^2\right)+\left(5x^2-3xy\right)$

Answer

$7x^2-6xy-2y^2$

Example 6.1.31

Find the difference: $\left(p^2+q^2\right)-\left(p^2+10pq-2q^2\right)$

Solution

	$\left(p^2+q^2\right)-\left(p^2+10pq-2q^2\right)$
Distribute.	$p^2+q^2-p^2-10pq+2q^2$
Rearrange the terms, to put like terms together	$p^2-p^2-10pq+q^2+2q^2$
Combine like terms.	$-10pq+3q^2$

Try It 6.1.32

Find the difference: $\left(a^2+b^2\right)-\left(a^2+5ab-6b^2\right)$

Answer

$-5ab-5b^2$

Try It 6.1.33

Find the difference: $\left(m^2+n^2\right)-\left(m^2-7mn-3n^2\right)$

Answer

$4n^2+7mn$

Example 6.1.34

Simplify: $\left(a^3-a^2b\right)-\left(ab^2+b^3\right)+\left(a^2b+ab^2\right)$

Solution

	$\left(a^3-a^2b\right)-\left(ab^2+b^3\right)+\left(a^2b+ab^2\right)$
Distribute.	$a^3-a^2b-ab^2-b^3+a^2b+ab^2$
Rearrange the terms, to put like terms together	$a^3-a^2b+a^2b-ab^2+ab^2-b^3$
Combine like terms.	a^3-b^3

Try It 6.1.35

Simplify: $\left(x^3-x^2y\right)-\left(xy^2+y^3\right)+\left(x^2y+xy^2\right)$

Answer

x^3-y^3

Try It 6.1.36

Simplify: $\left(p^3-p^2q\right)+\left(pq^2+q^3\right)-\left(p^2q+pq^2\right)$

Answer

$p^3-2p^2q+q^3$

Evaluate a Polynomial for a Given Value

We have already learned how to evaluate expressions. Since polynomials are expressions, we'll follow the same procedures to evaluate a **polynomial**. We will substitute the given value for the variable and then simplify using the order of operations.

Example 6.1.37

Evaluate $5x^2-8x+4$ when

1. x=4
2. x=−2
3. x=0

Solution

1. x=4	
	$5x^2-8x+4$
Substitute 4 for x.	$5(4)^2-8(4)+4$
Simplify the exponents.	$5\cdot16-8(4)+4$
Multiply.	$80-32+4$

Simplify.	52

2. x=−2	
	$5x^2 - 8x + 4$
Substitute –2 for *x*.	$5(-2)^2 - 8(-2) + 4$
Simplify the exponents.	$5 \cdot 4 - 8(-2) + 4$
Multiply.	$20 + 16 + 4$
Simplify.	40

3. x=0	
	$5x^2 - 8x + 4$
Substitute 0 for *x*.	$5(0)^2 - 8(0) + 4$
Simplify the exponents.	$5 \cdot 0 - 8(0) + 4$
Multiply.	$0 + 0 + 4$
Simplify.	4

? Try It 6.1.38

Evaluate: $3x^2 + 2x - 15$ when

1. x=3
2. x=−5
3. x=0

Answer

1. 18
2. 50
3. −15

? Try It 6.1.39

Evaluate: $5z^2 - z - 4$ when

1. z=−2
2. z=0
3. z=2

Answer

1. 18
2. −4
3. 14

✓ Example 6.1.40

The polynomial $-16t^2 + 250$ gives the height of a ball tt seconds after it is dropped from a 250 foot tall building. Find the height after t=2 seconds.

Solution

	$-16t^2+250$
Substitute t = 2.	$-16(2)^2+250$
Simplify	$-16\cdot 4+250$
Simplify	$-64+250$
Simplify	186
	After 2 seconds the height of the ball is 186 feet.

Try It 6.1.41

The polynomial $-16t^2+250$ gives the height of a ball tt seconds after it is dropped from a 250 foot tall building. Find the height after t=0 seconds.

Answer

250

Try It 6.1.42

The polynomial $-16t^2+250$ gives the height of a ball tt seconds after it is dropped from a 250 foot tall building. Find the height after t=3 seconds.

Answer

106

Example 6.1.43

The polynomial $6x^2+15xy$ gives the cost, in dollars, of producing a rectangular container whose top and bottom are squares with side x feet and sides of height y feet. Find the cost of producing a box with x=4 feet and y=6y=6 feet.

Solution

	$6x^2+15xy$
Substitute $x=4, y=6$.	$6(4)^2+15(4)(6)$
Simplify.	$6\cdot 16+15(4)(6)$
Simplify.	$96+360$
Simplify.	456
	The cost of producing the box is $456.

Try It 6.1.43

The polynomial $6x^2+15xy$ gives the cost, in dollars, of producing a rectangular container whose top and bottom are squares with side x feet and sides of height y feet. Find the cost of producing a box with x=6 feet and y=4 feet.

Answer

$576

Try It 6.1.44

The polynomial $6x^2 + 15xy$ gives the cost, in dollars, of producing a rectangular container whose top and bottom are squares with side *x* feet and sides of height *y* feet. Find the cost of producing a box with x=5 feet and y=8 feet.

Answer

$750

Key Concepts

- **Monomials**
  - A monomial is a term of the form ax^m, where aa is a constant and mm is a whole number
- **Polynomials**
  - **polynomial**—A monomial, or two or more monomials combined by addition or subtraction is a polynomial.
  - **monomial**—A polynomial with exactly one term is called a monomial.
  - **binomial**—A polynomial with exactly two terms is called a binomial.
  - **trinomial**—A polynomial with exactly three terms is called a trinomial.
- **Degree of a Polynomial**
  - The **degree of a term** is the sum of the exponents of its variables.
  - The **degree of a constant** is 0.
  - The **degree of a polynomial** is the highest degree of all its terms.

Glossary

binomial

A binomial is a polynomial with exactly two terms.

degree of a constant

The degree of any constant is 0.

degree of a polynomial

The degree of a polynomial is the highest degree of all its terms.

degree of a term

The degree of a term is the exponent of its variable.

monomial

A monomial is a term of the form ax^m, where a is a constant and m is a whole number; a monomial has exactly one term.

polynomial

A polynomial is a monomial, or two or more monomials combined by addition or subtraction.

standard form

A polynomial is in standard form when the terms of a polynomial are written in descending order of degrees.

trinomial

A trinomial is a polynomial with exactly three terms.

This page titled 6.1: Add and Subtract Polynomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.1: Add and Subtract Polynomials** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.1E: Exercises

Identify Polynomials, Monomials, Binomials, and Trinomials

In the following exercises, determine if each of the following polynomials is a monomial, binomial, trinomial, or other polynomial.

Exercise 1

a. $81b^5 - 24b^3 + 1$
b. $5c^3 + 11c^2 - c - 8$
c. $\frac{14}{15}y + \frac{1}{7}$
d. 5
e. $4y + 17$

Answer

a. trinomial
b. polynomial
c. binomial
d. monomial
e. binomial

Exercise 2

a. $x^2 - y^2$
b. $-13c^4$
c. $x^2 + 5x - 7$
d. $x^2y^2 - 2xy + 8$
e. 19

Exercise 3

a. $8 - 3x$
b. $z^2 - 5z - 6$
c. $y^3 - 8y^2 + 2y - 16$
d. $81b^5 - 24b^3 + 1$
e. -18

Answer

a. binomial
b. trinomial
c. polynomial
d. trinomial
e. monomial

Exercise 4

a. $11y^2$
b. -73
c. $6x^2 - 3xy + 4x - 2y + y^2$
d. $4y + 17$
e. $5c^3 + 11c^2 - c - 8$

Determine the Degree of Polynomials

In the following exercises, determine the degree of each polynomial.

Exercise 5

a. $6a^2+12a+14$
b. $18xy^2z$
c. $5x+2$
d. $y^3-8y^2+2y-16$
e. -24

Answer

a. 2
b. 4
c. 1
d. 3
e. 0

Exercise 6

a. $9y^3-10y^2+2y-6$
b. $-12p^4$
c. $a^2+9a+18$
d. $20x^2y^2-10a^2b^2+30$
e. 17

Exercise 7

a. $14-29x$
b. z^2-5z-6
c. $y^3-8y^2+2y-16$
d. $23ab^2-14$
e. -3

Answer

a. 1
b. 2
c. 3
d. 3
e. 0

Exercise 8

a. $62y^2$
b. 15
c. $6x^2-3xy+4x-2y+y^2$
d. $10-9x$
e. $m^4+4m^3+6m^2+4m+1$

Add and Subtract Monomials

In the following exercises, add or subtract the monomials.

Exercise 9

$7x^2 + 5x^2$

Answer

$12x^2$

Exercise 10

$4y^3 + 6y^3$

Exercise 11

$-12w + 18w$

Answer

$6w$

Exercise 12

$-3m + 9m$

Exercise 13

$4a - 9a$

Answer

$-5a$

Exercise 14

$-y - 5y$

Exercise 15

$28x - (-12x)$

Answer

$40x$

Exercise 16

$13z - (-4z)$

Exercise 17

$-5b - 17b$

Answer

$-22b$

Exercise 18

$-10x - 35x$

Exercise 19

$12a + 5b - 22a$

Answer

$-10a + 5b$

Exercise 20

$14x - 3y - 13x$

Exercise 21

$2a^2 + b^2 - 6a^2$

Answer

$-4a^2 + b^2$

Exercise 22

$5u^2 + 4v^2 - 6u^2$

Exercise 23

$xy^2 - 5x - 5y^2$

Answer

$xy^2 - 5x - 5y^2$

Exercise 24

$pq^2 - 4p - 3q^2$

Exercise 25

$a^2b - 4a - 5ab^2$

Answer

$a^2b - 4a - 5ab^2$

Exercise 26

$x^2y - 3x + 7xy^2$

Exercise 27

$12a+8b$

Answer

$12a+8b$

Exercise 28

$19y+5z$

Exercise 29

Add: $4a,\ -3b,\ -8a$

Answer

$-4a-3b$

Exercise 30

Add: $4x,\ 3y,\ -3x$

Exercise 31

Subtract $5x^6$ from $-12x^6$

Answer

$-17x^6$

Exercise 32

Subtract $2p^4$ from $-7p^4$

Add and Subtract Polynomials

In the following exercises, add or subtract the polynomials.

Exercise 33

$(5y^2+12y+4)+(6y^2-8y+7)$

Answer

$11y^2+4y+11$

Exercise 34

$(4y^2+10y+3)+(8y^2-6y+5)$

Exercise 35

$(x^2+6x+8)+(-4x^2+11x-9)$

Answer

$-3x^2+17x-1$

Exercise 36

$(y^2+9y+4)+(-2y^2-5y-1)$

Exercise 37

$(8x^2-5x+2)+(3x^2+3)$

Answer

$11x^2-5x+5$

Exercise 38

$(7x^2-9x+2)+(6x^2-4)$

Exercise 39

$(5a^2+8)+(a^2-4a-9)$

Answer

$6a^2-4a-1$

Exercise 40

$(p^2-6p-18)+(2p^2+11)$

Exercise 41

$(4m^2-6m-3)-(2m^2+m-7)$

Answer

$2m^2-7m+4$

Exercise 42

$(3b^2-4b+1)-(5b^2-b-2)$

Exercise 43

$(a^2+8a+5)-(a^2-3a+2)$

Answer

$11a+3$

Exercise 44

$(b^2-7b+5)-(b^2-2b+9)$

Exercise 45

$(12s^2-15s)-(s-9)$

Answer

$12s^2-16s+9$

Exercise 46

$(10r^2-20r)-(r-8)$

Exercise 47

Subtract $(9x^2+2)$ from $(12x^2-x+6)$

Answer

$3x^2-x+4$

Exercise 48

Subtract $(5y^2-y+12)$ from $(10y^2-8y-20)$

Exercise 49

Subtract $(7w^2-4w+2)$ from $(8w^2-w+6)$

Answer

w^2+3w+4

Exercise 50

Subtract $(5x^2-x+12)$ from $(9x^2-6x-20)$

Exercise 51

Find the sum of $(2p^3-8)$ and $(p^2+9p+18)$

Answer

$2p^3+p^2+9p+10$

Exercise 52

Find the sum of
$(q^2+4q+13)$ and $(7q^3-3)$

Exercise 53

Find the sum of $(8a^3 - 8a)$ and $(a^2 + 6a + 12)$

Answer

$8a^3 + a^2 - 2a + 12$

Exercise 54

Find the sum of
$(b^2 + 5b + 13)$ and $(4b^3 - 6)$

Exercise 55

Find the difference of

$(w^2 + w - 42)$ and
$(w^2 - 10w + 24)$.

Answer

$11w - 66$

Exercise 56

Find the difference of
$(z^2 - 3z - 18)$ and
$(z^2 + 5z - 20)$

Exercise 57

Find the difference of
$(c^2 + 4c - 33)$ and
$(c^2 - 8c + 12)$

Answer

$12c - 45$

Exercise 58

Find the difference of
$(t^2 - 5t - 15)$ and
$(t^2 + 4t - 17)$

Exercise 59

$(7x^2 - 2xy + 6y^2) + (3x^2 - 5xy)$

Answer

$10x^2 - 7xy + 6y^2$

Exercise 60

$(-5x^2-4xy-3y^2)+(2x^2-7xy)$

Exercise 61

$(7m^2+mn-8n^2)+(3m^2+2mn)$

Answer

$10m^2+3mn-8n^2$

Exercise 62

$(2r^2-3rs-2s^2)+(5r^2-3rs)$

Exercise 63

$(a^2-b^2)-(a^2+3ab-4b^2)$

Answer

$-3ab+3b^2$

Exercise 64

$(m^2+2n^2)-(m^2-8mn-n^2)$

Exercise 65

$(u^2-v^2)-(u^2-4uv-3v^2)$

Answer

$4uv+2v^2$

Exercise 66

$(j^2-k^2)-(j^2-8jk-5k^2)$

Exercise 67

$(p^3-3p^2q)+(2pq^2+4q^3)-(3p^2q+pq^2)$

Answer

$p^3-6p^2q+pq^2+4q^3$

Exercise 68

$(a^3-2a^2b)+(ab^2+b^3)-(3a^2b+4ab^2)$

Exercise 69

$(x^3 - x^2y) - (4xy^2 - y^3) + (3x^2y - xy^2)$

Answer

$x^3 + 2x^2y - 5xy^2 + y^3$

Exercise 70

$(x^3 - 2x^2y) - (xy^2 - 3y^3) - (x^2y - 4xy^2)$

Evaluate a Polynomial for a Given Value

In the following exercises, evaluate each polynomial for the given value.

Exercise 71

Evaluate $8y^2 - 3y + 2$ when:

a. $y = 5$
b. $y = -2$
c. $y = 0$

Answer

a. 187
b. 46
c. 2

Exercise 72

Evaluate $5y^2 - y - 7$ when:

a. $y = -4$
b. $y = 1$
c. $y = 0$

Exercise 73

Evaluate $4 - 36x$ when:

a. $x = 3$
b. $x = 0$
c. $x = -1$

Answer

a. -104
b. 4
c. 40

Exercise 74

Evaluate $16 - 36x^2$ when:

a. $x = -1$
b. $x = 0$
c. $x = 2$

Exercise 75

A painter drops a brush from a platform 75 feet high. The polynomial $-16t^2+75$ gives the height of the brush t seconds after it was dropped. Find the height after $t=2$ seconds.

Answer

11

Exercise 76

A girl drops a ball off a cliff into the ocean. The polynomial $-16t^2+250$ gives the height of a ball t seconds after it is dropped from a 250-foot tall cliff. Find the height after $t=2$ seconds.

Exercise 77

A manufacturer of stereo sound speakers has found that the revenue received from selling the speakers at a cost of p dollars each is given by the polynomial $-4p^2+420p$. Find the revenue received when $p=60$ dollars.

Answer

$\$10,800$

Exercise 78

A manufacturer of the latest basketball shoes has found that the revenue received from selling the shoes at a cost of p dollars each is given by the polynomial $-4p^2+420p$. Find the revenue received when $p=90$ dollars.

Everyday Math

Exercise 79

Fuel Efficiency The fuel efficiency (in miles per gallon) of a car going at a speed of x miles per hour is given by the polynomial $-\frac{1}{150}x^2+\frac{1}{3}x$, where $x=30$ mph.

Answer

4

Exercise 80

Stopping Distance The number of feet it takes for a car traveling at x miles per hour to stop on dry, level concrete is given by the polynomial $0.06x^2+1.1x$, where $x=40$ mph.

Exercise 81

Rental Cost The cost to rent a rug cleaner for d days is given by the polynomial $5.50d+25$. Find the cost to rent the cleaner for 6 days.

Answer

$\$58$

Exercise 82

Height of Projectile The height (in feet) of an object projected upward is given by the polynomial $-16t^2+60t+90$ where t represents time in seconds. Find the height after $t=2.5$ seconds.

Exercise 83

Temperature Conversion The temperature in degrees Fahrenheit is given by the polynomial $\frac{9}{5}c+32$ where c represents the temperature in degrees Celsius. Find the temperature in degrees Fahrenheit when $c=65^\circ$.

Answer

149°F

Writing Exercises

Exercise 84

Using your own words, explain the difference between a monomial, a binomial, and a trinomial.

Exercise 85

Using your own words, explain the difference between a polynomial with five terms and a polynomial with a degree of 5.

Answer

Answers will vary.

Exercise 86

Ariana thinks the sum $6y^2+5y^4$ is $11y^6$

Exercise 87

Jonathan thinks that $\frac{1}{3}$ and $\frac{1}{x}$ are both monomials. What is wrong with his reasoning?

Answer

Answers will vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
identify polynomials, monomials, binomials, and trinomials.			
determine the degree of polynomials.			
add and subtract monomials.			
add and subtract polynomials.			
evaluate a polynomial for a given value.			

b. If most of your checks were:

…confidently. Congratulations! You have achieved the objectives in this section. Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific.

…with some help. This must be addressed quickly because topics you do not master become potholes in your road to success. In math every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no - I don't get it! This is a warning sign and you must not ignore it. You should get help right away or you will quickly be overwhelmed. See your instructor as soon as you can to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 6.1E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.1E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.2: Use Multiplication Properties of Exponents

Learning Objectives

By the end of this section, you will be able to:

- Simplify expressions with exponents
- Simplify expressions using the Product Property for Exponents
- Simplify expressions using the Power Property for Exponents
- Simplify expressions using the Product to a Power Property
- Simplify expressions by applying several properties
- Multiply monomials

Note

Before you get started, take this readiness quiz.


1. Simplify: $\frac{3}{4} \cdot \frac{3}{4}$
   If you missed this problem, review Example 1.6.13.
2. Simplify: $(-2)(-2)(-2)$.
   If you missed this problem, review Example 1.5.13.

Simplify Expressions with Exponents

Remember that an exponent indicates repeated multiplication of the same quantity. For example, 2^4 means the product of 4 factors of 2, so 2^4 means $2 \cdot 2 \cdot 2 \cdot 2$.

Let's review the vocabulary for expressions with exponents.

EXPONENTIAL NOTATION



This is read a to the mth power.

In the expression a^m, the *exponent* m tells us how many times we use the *base* a as a factor.



Before we begin working with variable expressions containing exponents, let's simplify a few expressions involving only numbers.

Example 6.2.1

Simplify:

a. 4^3
b. 7^1
c. $\left(\frac{5}{6}\right)^2$
d. $(0.63)^2$

Solution

a. 4^3

Multiply three factors of 4. $4 \cdot 4 \cdot 4$

Simplify. 64

b. 7^1

Multiply one factor of 7. 7

c. $\left(\frac{5}{6}\right)^2$

Multiply two factors. $\left(\frac{5}{6}\right)\left(\frac{5}{6}\right)$

Simplify. $\frac{25}{36}$

d. $(0.63)^2$

Multiply two factors. $(0.63)(0.63)$

Simplify. 0.3969

Try It 6.2.2

Simplify:

a. 6^3
b. 15^1
c. $\left(\frac{3}{7}\right)^2$
d. $(0.43)^2$

Answer

a. 216
b. 15
c. $\frac{9}{49}$
d. 0.1849

Try It 6.2.3

Simplify:

a. 2^5
b. 21^1
c. $\left(\frac{2}{5}\right)^3$
d. $(0.218)^2$

Answer

a. 32
b. 21
c. $\frac{8}{125}$
d. 0.047524

Example 6.2.4

Simplify:

a. $(-5)^4$
b. -5^4

Solution

		$(-5)^4$
a.	Multiply four factors of -5	$(-5)(-5)(-5)(-5)$
	Simplify.	625
		-5^4
b.	Multiply four factors of 5.	$-(5 \cdot 5 \cdot 5 \cdot 5)$
	Simplify.	-625

Notice the similarities and differences in Example $6.2.4$ part 1 and Example $6.2.4$ part 2! Why are the answers different? As we follow the order of operations in part 1 the parentheses tell us to raise the (-5) to the 4th power. In part 2 we raise just the 5 to the 4th power and then take the opposite.

Try It 6.2.5

Simplify:

a. $(-3)^4$
b. -3^4

Answer

a. 81
b. −81

Try It 6.2.6

Simplify:

a. $(-13)^4$
b. -13^4

Answer

a. 169
b. −169

Simplify Expressions Using the Product Property for Exponents

You have seen that when you combine like terms by adding and subtracting, you need to have the same base with the same exponent. But when you multiply and divide, the exponents may be different, and sometimes the bases may be different, too.

We'll derive the properties of exponents by looking for patterns in several examples.

First, we will look at an example that leads to the Product Property.

	$x^2 \cdot x^3$
What does this mean? How many factors altogether?	$\underbrace{x \cdot x}_{2 \text{ factors}} \cdot \underbrace{x \cdot x \cdot x}_{3 \text{ factors}}$ 5 factors
So, we have	x^5
Notice that 5 is the sum of the exponents, 2 and 3.	$x^2 \cdot x^3$ is x^{2+3} or x^5

We write:

$$\begin{aligned} & x^2 \cdot x^3 \\ & x^{2+3} \\ & x^5 \end{aligned} \tag{6.2.1}$$

The base stayed the same and we added the exponents. This leads to the **Product Property for Exponents**.

PRODUCT PROPERTY FOR EXPONENTS

If a is a real number, and m and n are counting numbers, then

$$a^m \cdot a^n = a^{m+n} \tag{6.2.2}$$

To multiply with like bases, add the exponents.

An example with numbers helps to verify this property.

$$\begin{aligned} 2^3 \cdot 2^2 &\stackrel{?}{=} 2^{2+3} \\ 4 \cdot 8 &\stackrel{?}{=} 2^5 \\ 32 &= 32\checkmark \end{aligned} \tag{6.2.3}$$

Example 6.2.7

Simplify: $y^5 \cdot y^6$

Solution

	$y^5 \cdot y^6$
Use the product property, $a^m \cdot a^n = a^{m+n}$.	y^{5+6}
Simplify.	y^{11}

Try It 6.2.8

Simplify: $b^9 \cdot b^8$

Answer

b^{17}

Try It 6.2.9

Simplify: $x^{12} \cdot x^4$

Answer

x^{16}

Example 6.2.10

Simplify:

a. $2^5 \cdot 2^9$
b. $3 \cdot 3^4$

Solution

a.

	$2^5 \cdot 2^9$
Use the product property, $a^m \cdot a^n = a^{m+n}$.	2^{5+9}
Simplify.	2^{14}

b.

	$3 \cdot 3^4$
Use the product property, $a^m \cdot a^n = a^{m+n}$.	3^{1+4}
Simplify.	3^5

Try It 6.2.11

Simplify:

a. $5 \cdot 5^5$
b. $4^9 \cdot 4^9$

Answer

a. 5^6
b. 4^{18}

Try It 6.2.12

Simplify:

a. $7^6 \cdot 7^8$
b. $10 \cdot 10^{10}$

Answer

a. 7^{14}
b. 10^{11}

Example 6.2.13

Simplify:

a. $a^7 \cdot a$
b. $x^{27} \cdot x^{13}$

Solution

a.

	$a^7 \cdot a$
Rewrite, $a = a^1$	$a^7 \cdot a^1$
Use the product property, $a^m \cdot a^n = a^{m+n}$.	a^{7+1}
Simplify.	a^8

b.

	$x^{27} \cdot x^{13}$
Notice, the bases are the same, so add the exponents.	x^{27+13}
Simplify.	x^{40}

Try It 6.2.14

Simplify:

a. $p^5 \cdot p$
b. $y^{14} \cdot y^{29}$

Answer

a. p^6
b. y^{43}

Try It 6.2.15

Simplify:

a. $z \cdot z^7$
b. $b^{15} \cdot b^{34}$

Answer

a. z^8
b. b^{49}

We can extend the Product Property for Exponents to more than two factors.

Example 6.2.16

Simplify: $d^4 \cdot d^5 \cdot d^2$

Solution

	$d^4 \cdot d^5 \cdot d^2$
Add the exponents, since bases are the same.	d^{4+5+2}
Simplify.	d^{11}

Try It 6.2.17

Simplify: $x^6 \cdot x^4 \cdot x^8$

Answer

x^{18}

Try It 6.2.18

Simplify: $b^5 \cdot b^9 \cdot b^5$

Answer

b^{19}

Simplify Expressions Using the Power Property for Exponents

Now let's look at an exponential expression that contains a power raised to a power. See if you can discover a general property.

	$(x^2)^3$

What does this mean? How many factors altogether?	$\underbrace{x^2 \cdot x^2 \cdot x^2}_{6 \text{ factors}}$ $\underbrace{x \cdot x}_{2 \text{ factors}} \cdot \underbrace{x \cdot x}_{2 \text{ factors}} \cdot \underbrace{x \cdot x}_{2 \text{ factors}}$
So we have	x^6
Notice that 6 is the product of the exponents, 2 and 3.	$\left(x^2\right)^3$ is $x^{2 \cdot 3}$ or x^6

We write:

$$\begin{aligned} &\left(x^2\right)^3 \\ &x^{2 \cdot 3} \\ &x^6 \end{aligned} \tag{6.2.4}$$

We multiplied the exponents. This leads to the **Power Property for Exponents.**

POWER PROPERTY FOR EXPONENTS

If a is a real number, and m and n are whole numbers, then

$$(a^m)^n = a^{m \cdot n} \tag{6.2.5}$$

To raise a power to a power, multiply the exponents.

An example with numbers helps to verify this property.

$$\begin{aligned} \left(3^2\right)^3 &\stackrel{?}{=} 3^{2 \cdot 3} \\ (9)^3 &\stackrel{?}{=} 3^6 \\ 729 &= 729 \checkmark \end{aligned} \tag{6.2.6}$$

Example 6.2.19

Simplify:

a. $\left(y^5\right)^9$

b. $\left(4^4\right)^7$

Solution

a.

	$\left(y^5\right)^9$
Use the power property, $\left(a^m\right)^n = a^{m \cdot n}$.	$y^{5 \cdot 9}$
Simplify.	y^{45}

b.

	$\left(4^4\right)^7$
Use the power property.	$y^{4 \cdot 7}$
Simplify.	4^{28}

Try It 6.2.20

Simplify:

a. $\left(b^7\right)^5$
b. $\left(5^4\right)^3$

Answer

a. b^{35}
b. 5^{12}

Try It 6.2.21

Simplify:

a. $\left(z^6\right)^9$
b. $\left(3^7\right)^7$

Answer

a. z^{54}
b. 3^{49}

Simplify Expressions Using the Product to a Power Property

We will now look at an expression containing a product that is raised to a power. Can you find this pattern?

What does this mean?	$(2x)^3$
We group the like factors together.	$2x \cdot 2x \cdot 2x$
How many factors of 2 and of x?	$2 \cdot 2 \cdot 2 \cdot x^3$

Notice that each factor was raised to the power and $(2x)^3$ is $2^3 \cdot x^3$

We write: $(2x)^3$
$2^3 \cdot x^3$

The exponent applies to each of the factors! This leads to the **Product to a Power Property for Exponents.**

PRODUCT TO A POWER PROPERTY FOR EXPONENTS

If a and b are real numbers and m is a whole number, then

$$(ab)^m = a^m b^m \tag{6.2.7}$$

To raise a product to a power, raise each factor to that power.

An example with numbers helps to verify this property:

$$\begin{aligned} (2 \cdot 3)^2 &\stackrel{?}{=} 2^2 \cdot 3^2 \\ 6^2 &\stackrel{?}{=} 4 \cdot 9 \\ 36 &= 36\checkmark \end{aligned} \tag{6.2.8}$$

Example 6.2.22

Simplify:

a. $(-9d)^2$
b. $(3mn)^3$.

Solution

a.

	$(-9d)^2$
Use Power of a Product Property, $(ab)^m = a^m b^m$.	$(-9)^2 d^2$
Simplify.	$81d^2$

b.

	$(3mn)^3$.
Use Power of a Product Property, $(ab)^m = a^m b^m$.	$(3)^3 m^3 n^3$
Simplify.	$27m^3 n^3$

Try It 6.2.23

Simplify:

a. $(-12y)^2$
b. $(2wx)^5$

Answer

a. $144y^2$
b. $32w^5x^5$

Try It 6.2.24

Simplify:

a. $(5wx)^3$
b. $(-3y)^3$

Answer

a. $125w^3x^3$
b. $-27y^3$

Simplify Expressions by Applying Several Properties

We now have three properties for multiplying expressions with exponents. Let's summarize them and then we'll do some examples that use more than one of the properties.

PROPERTIES OF EXPONENTS

If a and b are real numbers, and m and n are whole numbers, then

$$\begin{aligned} &\textbf{Product Property} && a^m \cdot a^n &&= a^{m+n} \\ &\textbf{Power Property} && (a^m)^n &&= a^{mn} \\ &\textbf{Product to a Power} && (ab)^m &&= a^m b^m \end{aligned} \tag{6.2.9}$$

All exponent properties hold true for any real numbers m and n. Right now, we only use whole number exponents.

Example 6.2.25

Simplify:

a. $\left(y^3\right)^6\left(y^5\right)^4$
b. $\left(-6x^4y^5\right)^2$

Solution

a.

	$\left(y^3\right)^6\left(y^5\right)^4$
Use the Power Property.	$y^{18}\cdot y^{20}$
Add the exponents.	y^{38}

b.

	$\left(-6x^4y^5\right)^2$
Use the Product to a Power Property.	$(-6)^2\left(x^4\right)^2\left(y^5\right)^2$
Use the Power Property.	$(-6)^2\left(x^8\right)\left(y^{10}\right)^2$
Simplify.	$36x^8y^{10}$

Try It 6.2.26

Simplify:

a. $\left(a^4\right)^5\left(a^7\right)^4$
b. $\left(-2c^4d^2\right)^3$

Answer

a. a^{48}
b. $-8c^{12}d^6$

Try It 6.2.27

Simplify:

a. $\left(-3x^6y^7\right)^4$
b. $\left(q^4\right)^5\left(q^3\right)^3$

Answer

a. $81x^{24}y^{28}$
b. q^{29}

Example 6.2.28

Simplify:

a. $(5m)^2\left(3m^3\right)$
b. $\left(3x^2y\right)^4\left(2xy^2\right)^3$

Solution

a.

	$(5m)^2\left(3m^3\right)$
Raise $5m$ to the second power.	$5^2m^2\cdot 3m^3$
Simplify.	$25m^2\cdot 3m^3$
Use the Commutative Property.	$25\cdot 3\cdot m^2\cdot m^3$
Multiply the constants and add the exponents.	$75m^5$

	$\left(3x^2y\right)^4\left(2xy^2\right)^3$
Use the Product to a Power Property.	$\left(3^4x^8y^4\right)\left(2^3x^3y^6\right)$
b. Simplify.	$\left(81x^8y^4\right)\left(8x^3y^6\right)$
Use the Commutative Property.	$81\cdot 8\cdot x^8\cdot x^3\cdot y^4\cdot y^6$
Multiply the constants and add the exponents.	$648x^{11}y^{10}$

Try It 6.2.29

Simplify:

a. $(5n)^2\left(3n^{10}\right)$
b. $\left(c^4d^2\right)^5\left(3cd^5\right)^4$

Answer

a. $75n^{12}$
b. $81c^{24}d^{30}$

Try It 6.2.30

Simplify:

a. $\left(a^3b^2\right)^6\left(4ab^3\right)^4$
b. $(2x)^3\left(5x^7\right)$

Answer

a. $256a^{22}b^{24}$
b. $40x^{10}$

Multiply Monomials

Since a **monomial** is an algebraic expression, we can use the properties of exponents to multiply monomials.

Example 6.2.31

Multiply: $\left(3x^2\right)\left(-4x^3\right)$

Solution

	$\left(3x^2\right)\left(-4x^3\right)$
Use the Commutative Property to rearrange the terms.	$3\cdot(-4)\cdot x^2\cdot x^3$
Multiply.	$-12x^5$

Try It 6.2.32

Multiply: $\left(5y^7\right)\left(-7y^4\right)$

Answer

$-35y^{11}$

Try It 6.2.33

Multiply: $\left(-6b^4\right)\left(-9b^5\right)$

Answer

$54b^9$

Example 6.2.34

Multiply: $\left(\frac{5}{6}x^3y\right)\left(12xy^2\right)$

Solution

	$\left(\frac{5}{6}x^3y\right)\left(12xy^2\right)$
Use the Commutative Property to rearrange the terms.	$\frac{5}{6}\cdot 12\cdot x^3\cdot x\cdot y\cdot y^2$
Multiply.	$10x^4y^3$

Try It 6.2.35

Multiply: $\left(\frac{2}{5}a^4b^3\right)\left(15ab^3\right)$

Answer

$6a^5b^6$

Try It 6.2.36

Multiply: $\left(\frac{2}{3}r^5s\right)\left(12r^6s^7\right)$

Answer

$8r^{11}s^8$


Note

Access these online resources for additional instruction and practice with using multiplication properties of exponents:

- Multiplication Properties of Exponents

Key Concepts

- **Exponential Notation**



- **Properties of Exponents**
  - If a and b are real numbers and m and n are whole numbers, then

$$\begin{array}{lrcll} \textbf{Product Property} & a^m\cdot a^n & = & a^{m+n} & \\ \textbf{Power Property} & (a^m)^n & = & a^{mn} & (6.2.10) \\ \textbf{Product to a Power} & (ab)^m & = & a^mb^m & \end{array}$$

This page titled 6.2: Use Multiplication Properties of Exponents is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.2: Use Multiplication Properties of Exponents** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.2E: Exercises

Practice Makes Perfect

Simplify Expressions with Exponents

In the following exercises, simplify each expression with exponents.

Exercise 1

a. 3^5
b. 9^1
c. $(\frac{1}{3})^2$
d. $(0.2)^4$

Exercise 2

a. 10^4
b. 17^1
c. $(\frac{2}{9})^2$
d. $(0.5)^3$

Answer

a. 10,000
b. 17
c. $\frac{4}{81}$
d. 0.125

Exercise 3

a. 2^6
b. 14^1
c. $(\frac{2}{5})^3$
d. $(0.7)^2$

Exercise 4

a. 8^3
b. 8^1
c. $(\frac{3}{4})^3$
d. $(0.4)^3$

Answer

a. 512
b. 8
c. $\frac{27}{64}$
d. 0.064

Exercise 5

a. $(-6)^4$
b. -6^4

Exercise 6

a. $(-2)^6$
b. -2^6

Answer

a. 64
b. −64

Exercise 7

a. $-(\frac{1}{4})^4$
b. $(-\frac{1}{4})^4$

Exercise 8

a. $-(\frac{2}{3})^2$
b. $(-\frac{2}{3})^2$

Answer

a. $-\frac{4}{9}$
b. $\frac{4}{9}$

Exercise 9

a. -0.5^2
b. $(-0.5)^2$

Exercise 10

a. -0.1^4
b. $(-0.1)^4$

Answer

a. −0.0001
b. 0.0001

Simplify Expressions Using the Product Property for Exponents

In the following exercises, simplify each expression using the Product Property for Exponents.

Exercise 11

$d^3 \cdot d^6$

Exercise 12

$x^4 \cdot x^2$

Answer

x^6

Exercise 13

$n^{19} \cdot n^{12}$

Exercise 14

$q^{27} \cdot q^{15}$

Answer

q^{42}

Exercise 15

a. $4^5 \cdot 4^9$
b. $8^9 \cdot 8$

Exercise 16

a. $3^{10} \cdot 3^6$
b. $5 \cdot 5^4$

Answer

a. 3^{16}
b. 5^5

Exercise 17

a. $y \cdot y^3$
b. $z^{25} \cdot z^8$

Exercise 18

a. $w^5 \cdot w$
b. $u^{41} \cdot u^{53}$

Answer

a. w^6
b. u^{94}

Exercise 19

$w \cdot w^2 \cdot w^3$

Exercise 20

$y \cdot y^3 \cdot y^5$

Answer

y^9

https://math.libretexts.org/@go/page/30536

Exercise 21

$a^4 \cdot a^3 \cdot a^9$

Exercise 22

$c^5 \cdot c^{11} \cdot c^2$

Answer

c^{18}

Exercise 23

$m^x \cdot m^3$

Exercise 24

$n^y \cdot n^2$

Answer

n^{y+2}

Exercise 25

$y^a \cdot y^b$

Exercise 26

$x^p \cdot x^q$

Answer

x^{p+q}

Simplify Expressions Using the Power Property for Exponents

In the following exercises, simplify each expression using the Power Property for Exponents.

Exercise 27

a. $(m^4)^2$
b. $(10^3)^6$

Exercise 28

a. $(b^2)^7$
b. $(3^8)^2$

Answer

a. b^{14}
b. 3^{16}

Exercise 29

a. $(y^3)^x$
b. $(5^x)^y$

Exercise 30

a. $(x^2)^y$
b. $(7^a)^b$

Answer

a. x^{2y}
b. 7^{ab}

Simplify Expressions Using the Product to a Power Property

In the following exercises, simplify each expression using the Product to a Power Property.

Exercise 31

a. $(6a)^2$
b. $(3xy)^2$

Exercise 32

a. $(5x)^2$
b. $(4ab)^2$

Answer

a. $25x^2$
b. $16a^2b^2$

Exercise 33

a. $(-4m)^3$
b. $(5ab)^3$

Exercise 34

a. $(-7n)^3$
b. $(3xyz)^4$

Answer

a. $-343n^3$
b. $81x^4y^4z^4$

Simplify Expressions by Applying Several Properties

In the following exercises, simplify each expression.

Exercise 35

a. $(y^2)^4 \cdot (y^3)^2$
b. $(10a^2b)^3$

Exercise 36

a. $(w^4)^3 \cdot (w^5)^2$
b. $(2xy^4)^5$

Answer

a. w^{22}
b. $32x^5y^{20}$

Exercise 37

a. $(-2r^3s^2)^4$
b. $(m^5)^3 \cdot (m^9)^4$

Exercise 38

a. $(-10q^2p^4)^3$
b. $(n^3)^{10} \cdot (n^5)^2$

Answer

a. $-1000q^6p^{12}$
b. n^{40}

Exercise 39

a. $(3x)^2(5x)$
b. $(5t^2)^3(3t)^2$

Exercise 40

a. $(2y)^3(6y)$
b. $(10k^4)^3(5k^6)^2$

Answer

a. $48y^4$
b. $25,000k^{24}$

Exercise 41

a. $(5a)^2(2a)^3$
b. $(12y^2)^3(23y)^2$

Exercise 42

a. $(4b)^2(3b)^3$
b. $(12j^2)^5(25j^3)^2$

Answer

a. $432b^5$
b. $1200j^{16}$

https://math.libretexts.org/@go/page/30536

Exercise 43

a. $(25x^2y)^3$
b. $(89xy^4)^2$

Exercise 44

a. $(2r^2)^3(4r)^2$
b. $(3x^3)^3(x^5)^4$

Answer

a. $128r^8$
b. $27x^{29}$

Exercise 45

a. $(m^2n)^2(2mn^5)^4$
b. $(3pq^4)^2(6p^6q)^2$

Multiply Monomials

In the following exercises, multiply the monomials.

Exercise 46

$(6y^7)(-3y^4)$

Answer

$-18y^{11}$

Exercise 47

$(-10x^5)(-3x^3)$

Exercise 48

$(-8u^6)(-9u)$

Answer

$72u^7$

Exercise 49

$(-6c^4)(-12c)$

Exercise 50

$(\frac{1}{5}f^8)(20f^3)$

Answer

$4f^{11}$

Exercise 51

$(\frac{1}{4}d^5)(36d^2)$

Exercise 52

$(4a^3b)(9a^2b^6)$

Answer

$36a^5b^7$

Exercise 53

$(6m^4n^3)(7mn^5)$

Exercise 54

$(\frac{4}{7}rs^2)(14rs^3)$

Answer

$8r^2s^5$

Exercise 55

$(\frac{5}{8}x^3y)(24x^5y)$

Exercise 56

$(\frac{2}{3}x^2y)(\frac{3}{4}xy^2)$

Answer

$\frac{1}{2}x^3y^3$

Exercise 57

$(\frac{3}{5}m^3n^2)(\frac{5}{9}m^2n^3)$

Mixed Practice

In the following exercises, simplify each expression.

Exercise 58

$(x^2)^4 \cdot (x^3)^2$

Answer

x^{14}

Exercise 59

$(y^4)^3 \cdot (y^5)^2$

Exercise 60

$(a^2)^6 \cdot (a^3)^8$

Answer

a^{36}

Exercise 61

$(b^7)^5 \cdot (b^2)^6$

Exercise 62

$(2m^6)^3$

Answer

$8m^{18}$

Exercise 63

$(3y^2)^4$

Exercise 64

$(10x^2y)^3$

Answer

$1000x^6y^3$

Exercise 65

$(2mn^4)^5$

Exercise 66

$(-2a^3b^2)^4$

Answer

$16a^{12}b^8$

Exercise 67

$(-10u^2v^4)^3$

https://math.libretexts.org/@go/page/30536

Exercise 68

$(\frac{2}{3}x^2y)^3$

Answer

$\frac{8}{27}x^6y^3$

Exercise 69

$(\frac{7}{9}pq^4)^2$

Exercise 70

$(8a^3)^2(2a)^4$

Answer

$1024a^{10}$

Exercise 71

$(5r^2)^3(3r)^2$

Exercise 72

$(10p^4)^3(5p^6)^2$

Answer

$25000p^{24}$

Exercise 73

$(4x^3)^3(2x^5)^4$

Exercise 74

$(\frac{1}{2}x^2y^3)^4(4x^5y^3)^2$

Answer

$x^{18}y^{18}$

Exercise 75

$(\frac{1}{3}m^3n^2)^4(9m^8n^3)^2$

Exercise 76

$(3m^2n)^2(2mn^5)^4$

Answer

$144m^8n^{22}$

Exercise 77

$(2pq^4)^3(5p^6q)^2$

Everyday Math

Exercise 78

Email Kate emails a flyer to ten of her friends and tells them to forward it to ten of their friends, who forward it to ten of their friends, and so on. The number of people who receive the email on the second round is 10^2, on the third round is 10^3, as shown in the table below. How many people will receive the email on the sixth round? Simplify the expression to show the number of people who receive the email.

Round	Number of People
1	10
2	10^2
3	10^3
...	...
6	?

Answer

1,000,000

Exercise 79

Salary Jamal's boss gives him a 3% raise every year on his birthday. This means that each year, Jamal's salary is 1.03 times his last year's salary. If his original salary was \$35,000, his salary after 1 year was \$35,000(1.03), after 2 years was $\$35,000(1.03)^2$, after 3 years was $\$35,000(1.03)^3$, as shown in the table below. What will Jamal's salary be after 10 years? Simplify the expression, to show Jamal's salary in dollars.

Year	Salary
1	\$35,000(1.03)
2	$\$35,000(1.03)^2$
3	$\$35,000(1.03)^3$
...	...
10	?

Exercise 80

Clearance A department store is clearing out merchandise in order to make room for new inventory. The plan is to mark down items by 30% each week. This means that each week the cost of an item is 70% of the previous week's cost. If the original cost of a sofa was \$1,000, the cost for the first week would be \$1,000(0.70) and the cost of the item during the second week would be $\$1,000(0.70)^2$. Complete the table shown below. What will be the cost of the sofa during the fifth week? Simplify the expression, to show the cost in dollars.

Week	Cost

Week	Cost
1	$1,000(0.70)
2	$1,000(0.70)^2$
3	$1,000(0.70)^2$
4	...
5	?
...	...
n	?

Answer

$168.07

Exercise 81

Depreciation Once a new car is driven away from the dealer, it begins to lose value. Each year, a car loses 10% of its value. This means that each year the value of a car is 90% of the previous year's value. If a new car was purchased for $20,000, the value at the end of the first year would be $20,000(0.90) and the value of the car after the end of the second year would be $ $20,000(0.90)^2$. Complete the table shown below. What will be the value of the car at the end of the eighth year? Simplify the expression, to show the value in dollars.

Year	Cost
1	$20,000(0.90)
2	$20,000(0.90)^2$
3	
...	...
8	?

Writing Exercises

Exercise 82

Use the Product Property for Exponents to explain why $x \cdot x = x^2$

Answer

Answers will vary.

Exercise 83

Explain why $-5^3 = (-5)^3$, but $-5^4 \neq (-5)^4$.

Exercise 84

Jorge thinks $(\frac{1}{2})^2$ is 1. What is wrong with his reasoning?

Answer

Answers will vary.

Exercise 85

Explain why $x^3 \cdot x^5$ is x^8, and not x^{15}.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
simplify expressions with exponents.			
simplify expressions using the Product Property for Exponents.			
simplify expressions using the Power Property for Exponents.			
simplify expressions using the Product to a Power Property.			
simplify expressions by applying several properties.			
multiply monomials.			

b. After reviewing this checklist, what will you do to become confident for all goals?

This page titled 6.2E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.2E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.3: Multiply Polynomials

Learning Objectives

By the end of this section, you will be able to:

- Multiply a polynomial by a monomial
- Multiply a binomial by a binomial
- Multiply a trinomial by a binomial

Note

Before you get started, take this readiness quiz.

1. Distribute: $2(x+3)$.
   If you missed this problem, review Example 1.10.31.
2. Combine like terms: $x^2+9x+7x+63$.
   If you missed this problem, review Example 1.3.37.

Multiply a Polynomial by a Monomial

We have used the Distributive Property to simplify expressions like $2(x-3)$. You multiplied both terms in the parentheses, x and 3, by 2, to get $2x-6$. With this chapter's new vocabulary, you can say you were multiplying a binomial, $x-3$, by a monomial, 2.

Multiplying a **binomial** by a **monomial** is nothing new for you! Here's an example:

Example 6.3.1

Multiply: $4(x+3)$.

Solution

	$4(x+3)$
Distribute.	$4\cdot x+4\cdot 3$
Simplify.	$4x+12$

Try It 6.3.2

Multiply: $5(x+7)$.

Answer

5x+35

Try It 6.3.3

Multiply: $3(y+13)$.

Answer

3y+39

Example 6.3.4

Multiply: y(y−2).

Solution

	$y(y-2)$
Distribute.	$y \cdot y - y \cdot 2$
Simplify.	$y^2 - 2y$

Try It 6.3.5

Multiply: $x(x-7)$.

Answer

$x^2 - 7x$

Try It 6.3.6

Multiply: $d(d-11)$.

Answer

$d^2 - 11d$

Example 6.3.7

Multiply: $7x(2x+y)$

Solution

	$7x(2x+y)$
Distribute.	$7x \cdot 2x + 7x \cdot y$
Simplify.	$14x^2 + 7xy$

Try It 6.3.8

Multiply: $5x(x+4y)$

Answer

$5x^2 + 20xy$

Try It 6.3.9

Multiply: $2p(6p+r)$

Answer

$12p^2 + 2pr$

Example 6.3.10

Multiply: $-2y\left(4y^2+3y-5\right)$

Solution

	
Distribute.	$-2y \cdot 4y^2+(-2y) \cdot 3y-(-2y) \cdot 5$
Simplify.	$-8y^3-6y^2+10y$

Try It 6.3.11

Multiply: $-3y\left(5y^2+8y-7\right)$

Answer

$-15y^3-24y^2+21y$

Try It 6.3.12

Multiply: $4x^2\left(2x^2-3x+5\right)$

Answer

$8x^4-24x^3+20x^2$

Example 6.3.13

Multiply: $2x^3\left(x^2-8x+1\right)$

Solution

	
Distribute.	$2x^3 \cdot x^2+\left(2x^3\right) \cdot(-8x)+\left(2x^3\right) \cdot 1$
Simplify.	$2x^5-16x^4+2x^3$

Try It 6.3.14

Multiply: $4x\left(3x^2-5x+3\right)$

Answer

$12x^3-20x^2+12x$

Try It 6.3.15

Multiply: $-6a^3\left(3a^2-2a+6\right)$

Answer

$-18a^5+12a^4-36a^3$

Example 6.3.16

Multiply: $(x+3)p$

Solution

The monomial is the second factor.	
Distribute.	$x \cdot p + 3 \cdot p$
Simplify.	$xp + 3p$

Try It 6.3.17

Multiply: $(x+8)p$

Answer

$xp + 8p$

Try It 6.3.18

Multiply: $(a+4)p$

Answer


$ap + 4p$

Multiply a Binomial by a Binomial

Just like there are different ways to represent multiplication of numbers, there are several methods that can be used to multiply a **binomial** times a binomial. We will start by using the Distributive Property.

Multiply a Binomial by a Binomial Using the Distributive Property

Look at Example 6.3.16, where we multiplied a binomial by a **monomial**.

Instructions	Expression
Starting Expressions	
We distributed the p to get:	
What if we have $(x+7)$ instead of p?	
Distribute $(x+7)$.	
Distribute again.	$x^2 + 7x + 3x + 21$
Combine like terms.	$x^2 + 10x + 21$

Notice that before combining like terms, you had four terms. You multiplied the two terms of the first binomial by the two terms of the second binomial—four multiplications.

Example 6.3.19

Multiply: $(y+5)(y+8)$

Solution

	$(y+5)(y+8)$
Distribute $(y+8)$.	$y(y+8)+5(y+8)$
Distribute again	$y^2+8y+5y+40$
Combine like terms.	$y^2+13y+40$

Try It 6.3.20

Multiply: $(x+8)(x+9)$

Answer

$x^2+17x+72$

Try It 6.3.21

Multiply: $(5x+9)(4x+3)$

Answer

$20x^2+51x+27$

Example 6.3.22

Multiply: $(2y+5)(3y+4)$

Solution

	$(2y+5)(3y+4)$
Distribute $(3y+4)$.	$2y(3y+4)+5(3y+4)$
Distribute again	$6y^2+8y+15y+20$
Combine like terms.	$6y^2+23y+20$

Try It 6.3.23

Multiply: $(3b+5)(4b+6)$

Answer

$12b^2+38b+30$

Try It 6.3.24

Multiply: $(a+10)(a+7)$

Answer

$a^2+17a+70$

Example 6.3.25

Multiply: $(4y+3)(2y-5)$

Solution

	$(4y+3)(2y-5)$
Distribute.	$4y(2y-5)+3(2y-5)$
Distribute again.	$8y^2-20y+6y-15$
Combine like terms.	$8y^2-14y-15$

Try It 6.3.26

Multiply: $(5y+2)(6y-3)$

Answer

$30y^2-3y-6$

Try It 6.3.27

Multiply: $(3c+4)(5c-2)$

Answer

$15c^2+14c-8$

Example 6.3.28

Multiply: $(x-2)(x-y)$

Solution

	$(x-2)(x-y)$
Distribute.	$x(x-y)-2(x-y)$
Distribute again.	$x^2-xy-2x+2y$
There are no like terms to combine.	

Try It 6.3.29

Multiply: $(a+7)(a-b)$

Answer

$a^2-ab+7a-7b$

Try It 6.3.30

Multiply: $(x+5)(x-y)$

Answer

$x^2 - xy + 5x - 5y$

Multiply a Binomial by a Binomial Using the FOIL Method

Remember that when you multiply a binomial by a binomial you get four terms. Sometimes you can combine like terms to get a **trinomial**, but sometimes, like in Example 6.3.28, there are no like terms to combine.

Let's look at the last example again and pay particular attention to how we got the four terms.


$$(x-2)(x-y)$$
$$x^2 - xy - 2x + 2y$$

Where did the first term, x^2, come from?


It is the product of *x* and *x*, the *first* terms in (*x* – 2) and (*x* – *y*).




The next term, –*xy*, is the product of *x* and –*y*, the two *outer* terms.



The third term, –2*x*, is the product of –2 and *x*, the two *inner* terms.



And the last term, +2*y*, came from multiplying the two *last* terms, –2 and –*y*.



We abbreviate "First, Outer, Inner, Last" as FOIL. The letters stand for '**F**irst, **O**uter, **I**nner, **L**ast'. The word FOIL is easy to remember and ensures we find all four products.

$$\begin{array}{c} (x-2)(x-y) \\ x^2 - xy - 2x + 2y \\ F \quad O \quad I \quad L \end{array} \tag{6.3.1}$$

Let's look at (x+3)(x+7).

Distributive Property	FOIL
$(x+3)(x+7)$	$(x+3)(x+7)$
$x(x+7)+3(x+7)$	
$x^2+7x+3x+21$ F O I L	$x^2+7x+3x+21$ F O I L
$x^2+10x+21$	$x^2+10x+21$

Notice how the terms in third line fit the FOIL pattern.

Now we will do an example where we use the FOIL pattern to multiply two binomials.

Example 6.3.31: How to Multiply a Binomial by a Binomial using the FOIL Method

Multiply using the FOIL method: $(x+5)(x+9)$

Solution

Step 1. Multiply the *First* terms.	$(x+5)(x+9)$	$x^2+\_+\_+\_$ F O I L
Step 2. Multiply the *Outer* terms.	$(x+5)(x+9)$	$x^2+9x+\_+\_$ F O I L
Step 3. Multiply the *Inner* terms.	$(x+5)(x+9)$	$x^2+9x+5x+\_$ F O I L
Step 4. Multiply the *Last* terms.	$(x+5)(x+9)$	$x^2+9x+5x+45$ F O I L
Step 5. Combine like terms, when possible.		$x^2+14x+45$

Try It 6.3.32

Multiply using the FOIL method: $(x+6)(x+8)$

Answer

$x^2+14x+48$

Try It 6.3.33

Multiply using the FOIL method: $(y+17)(y+3)$

Answer

$y^2+20y+51$

We summarize the steps of the FOIL method below. The FOIL method only applies to multiplying binomials, not other polynomials!

MULTIPLY TWO BINOMIALS USING THE FOIL METHOD

Step 1. Multiply the *First* terms.

Step 2. Multiply the *Outer* terms.

Step 3. Multiply the *Inner* terms.

Step 4. Multiply the *Last* terms.

Step 5. Combine like terms, when possible.

$$\underset{\text{first}}{(a}+\underbrace{\underset{\text{last}}{b})(\underset{\text{first}}{c}}_{\text{inner}}+\underbrace{\underset{\text{last}}{d}}_{\text{outer}})$$

Say it as you multiply!
FOIL
First
Outer
Inner
Last

When you multiply by the FOIL method, drawing the lines will help your brain focus on the pattern and make it easier to apply.

Example 6.3.34

Multiply: $(y-7)(y+4)$.

Solution

		$(y-7)(y+4)$
Multiply the *First* terms.	$(y-7)(y+4)$	$y^2+\_+\_+\_$ F O I L
Multiply the *Outer* terms.	$(y-7)(y+4)$	$y^2+4y+\_+\_$ F O I L
Multiply the *Inner* terms.	$(y-7)(y+4)$	$y^2+4y-7y+\_$ F O I L
Multiply the *Last* terms.	$(y-7)(y+4)$	$y^2+4y-7y-28$ F O I L
Combine like terms.		$y^2-3y-28$

Try It 6.3.35

Multiply: $(x-7)(x+5)$.

Answer

$x^2-2x-35$

Try It 6.3.36

Multiply: (b−3)(b+6).

Answer

$b^2+3b-18$

Example 6.3.37

Multiply: $(4x+3)(2x-5)$.

Solution

	$(4x-3)(2x-5)$ $(4x+3)(2x-5)$
Multiply the *First* terms, $4x \cdot 2x$.	$8x^2+\_+\_+\_$ F O I L
Multiply the *Outer* terms, $4x \cdot (-5)$.	$8x^2-20x+\_+\_$ F O I L
Multiply the *Inner* terms, $3 \cdot 2x$.	$8x^2-2x+6x+\_$ F O I L
Multiply the *Last* terms, $3 \cdot (-5)$.	$8x^2-20x+6x-15$ F O I L
Combine like terms.	$8x^2-14x-15$

Try It 6.3.38

Multiply: $(3x+7)(5x-2)$.

Answer

$15x^2+29x-14$

Try It 6.3.39

Multiply: $(4y+5)(4y-10)$.

Answer

$16y^2-20y-50$

The final products in the last four examples were trinomials because we could combine the two middle terms. This is not always the case.

Example 6.3.40

Multiply: $(3x-y)(2x-5)$.

Solution

	$(3x-y)(2x-5)$
	
Multiply the *First*.	$6x^2+\_+\_+\_$ F O I L
Multiply the *Outer*.	$6x^2-15x+\_+\_$ F O I L
Multiply the *Inner*.	$6x^2-15x-2xy+\_$ F O I L
Multiply the *Last*.	$6x^2-15x-2xy+5y$ F O I L
Combine like terms—there are none.	$6x^2-15x-2xy+5y$

Try It 6.3.41

Multiply: (10c−d)(c−6).

Answer

$10c^2-60c-cd+6d$

Try It 6.3.42

Multiply: (7x−y)(2x−5).

Answer


$14x^2-35x-2xy+10y$

Be careful of the exponents in the next example.

Example 6.3.43

Multiply: $\left(n^2+4\right)(n-1)$

Solution

	$(n^2+4)(n-1)$
	
Multiply the *First*.	$n^3+\_+\_+\_$
Multiply the *Outer*.	$n^3-n^2+\_+\_$
Multiply the *Inner*.	$n^3-n^2+4n+\_$
Multiply the *Last*.	n^3-n^2+4n-4
Combine like terms—there are none.	n^3-n^2+4n-4

Try It 6.3.44

Multiply: $(x^2+6)(x-8)$

Answer

$x^3-8x^2+6x-48$

Try It 6.3.45

Multiply: $(y^2+7)(y-9)$


Answer

$y^3-9y^2+7y-63$

Example 6.3.46

Multiply: $(3pq+5)(6pq-11)$

Solution

	$(3pq+5)(6pq-11)$	
Multiply the *First*.	$18p^2q^2+\_+\_+\_$	
Multiply the *Outer*.	$18p^2q^2-33pq+\_+\_$	
Multiply the *Inner*.	$18p^2q^2-33pq+30pq+\_$	
Multiply the *Last*.	$18p^2q^2-33pq+30pq-55$	
Combine like terms—there are none.	$18p^2q^2-3pq-55$	

Try It 6.3.47

Multiply: $(2ab+5)(4ab-4)$

Answer

$8a^2b^2+12ab-20$

Try It 6.3.48

Multiply: $(2xy+3)(4xy-5)$

Answer

$8x^2y^2+2xy-15$

Multiply a Binomial by a Binomial Using the Vertical Method

The FOIL method is usually the quickest method for multiplying two binomials, but it *only* works for binomials. You can use the Distributive Property to find the product of any two polynomials. Another method that works for all polynomials is the Vertical Method. It is very much like the method you use to multiply whole numbers. Look carefully at this example of multiplying two-digit numbers.

23	
×46	
138 partial product	Start by multiplying 23 by 6 to get 138.
92 partial product	Next, multiply 23 by 4, lining up the partial product in the correct columns.
1058 product	Last you add the partial products.

Now we'll apply this same method to multiply two binomials.

Example 6.3.49

Multiply using the Vertical Method: $(3y-1)(2y-6)$

Solution

It does not matter which binomial goes on the top.

Multiply $3y-1$ by -6.
Multiply $3y-1$ by $2y$.

$$\begin{array}{rl} 3y-1 & \\ \times\ 2y-6 & \\ \hline -18y+6 & \text{partial product} \\ 6y^2-2y & \text{partial product} \\ \hline 6y^2-20y+6 & \text{product} \end{array} \qquad (6.3.2)$$

Add like terms.

Notice the partial products are the same as the terms in the FOIL method.

$$\begin{array}{rl} (3y-1)(2y-6) & \\ 6y^2-2y-18y+6 & \\ 6y^2-20y+6 & \end{array} \qquad \begin{array}{r} 3y-1 \\ \times\ 2y-6 \\ \hline -18y+6 \\ 6y^2-2y \\ \hline 6y^2-20x+6 \end{array}$$

Try It 6.3.50

Multiply using the Vertical Method: $(5m-7)(3m-6)$

Answer

$15m^2-51m+42$

Try It 6.3.51

Multiply using the Vertical Method: $(6b-5)(7b-3)$

Answer

$42b^2-53b+15$

We have now used three methods for multiplying binomials. Be sure to practice each method, and try to decide which one you prefer. The methods are listed here all together, to help you remember them.

MULTIPLYING TWO BINOMIALS

To multiply binomials, use the:

- Distributive Property
- FOIL Method
- Vertical Method

Remember, FOIL only works when multiplying two binomials.

Multiply a Trinomial by a Binomial

We have multiplied monomials by monomials, monomials by polynomials, and binomials by binomials. Now we're ready to multiply a **trinomial** by a **binomial**. Remember, FOIL will not work in this case, but we can use either the Distributive Property or the Vertical Method. We first look at an example using the Distributive Property.

Example 6.3.52

Multiply using the Distributive Property: $(b+3)\left(2b^2-5b+8\right)$

Solution

	
Distribute.	
Multiply.	$2b^3-5b^2+8b+6b^2-15b+24$
Combine like terms.	$2b^3+b^2-7b+24$

Try It 6.3.53

Multiply using the Distributive Property: $(y-3)\left(y^2-5y+2\right)$

Answer

$y^3-8y^2+17y-6$

Try It 6.3.54

Multiply using the Distributive Property: $(x+4)\left(2x^2-3x+5\right)$

Answer

$2x^3+5x^2-7x+20$

Now let's do this same multiplication using the Vertical Method.

Example 6.3.55

Multiply using the Vertical Method: $(b+3)\left(2b^2-5b+8\right)$

Solution

It is easier to put the polynomial with fewer terms on the bottom because we get fewer partial products this way.

Multiply $(2b^2-5b+8)$ by 3.	$2b^2-5b+8$ $\times \quad b+3$ $6b^2-15b+24$
	$2b^3-5b^2+8b$
Multiply $(2b^2-5b+8)$ by b.	$2b^3+b^2-7b+24$
Add like terms.	

Try It 6.3.56

Multiply using the Vertical Method: $(y-3)\left(y^2-5y+2\right)$

Answer

$y^3-8y^2+17y-6$

Try It 6.3.57

Multiply using the Vertical Method: $(x+4)\left(2x^2-3x+5\right)$

Answer

$2x^3+5x^2-7x+20$

We have now seen two methods you can use to multiply a **trinomial** by a binomial. After you practice each method, you'll probably find you prefer one way over the other. We list both methods are listed here, for easy reference.

MULTIPLYING A TRINOMIAL BY A BINOMIAL

To multiply a trinomial by a binomial, use the:

- Distributive Property
- Vertical Method

Note

Access these online resources for additional instruction and practice with multiplying polynomials:

- Multiplying Exponents 1
- Multiplying Exponents 2
- Multiplying Exponents 3

Key Concepts

- **FOIL Method for Multiplying Two Binomials**—To multiply two binomials:
  1. Multiply the **First** terms.
  2. Multiply the **Outer** terms.

3. Multiply the **Inner** terms.
4. Multiply the **Last** terms.

- **Multiplying Two Binomials**—To multiply binomials, use the:
  - Distributive Property (Example)
  - FOIL Method (Example)
  - Vertical Method (Example)
- **Multiplying a Trinomial by a Binomial**—To multiply a trinomial by a binomial, use the:
  - Distributive Property (Example)
  - Vertical Method (Example)

This page titled 6.3: Multiply Polynomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.3: Multiply Polynomials** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.3E: Exercises

Practice Makes Perfect

Multiply a Polynomial by a Monomial

In the following exercises, multiply.

Exercise 6.3E.1

$4(w+10)$

Answer

4w+40

Exercise 6.3E.2

6(b+8)

Exercise 6.3E.3

−3(a+7)

Answer

−3a−21

Exercise 6.3E.4

−5(p+9)

Exercise 6.3E.5

2(x−7)

Answer

2x−14

Exercise 6.3E.6

7(y−4)

Exercise 6.3E.7

−3(k−4)

Answer

−3k+12

Exercise 6.3E.8

−8(j−5)

Exercise 6.3E.9

q(q+5)

Answer

q^2+5q

Exercise 6.3E.10

k(k+7)

Exercise 6.3E.11

−b(b+9)

Answer

$-b^2-9b$

Exercise 6.3E.12

−y(y+3)

Exercise 6.3E.13

−x(x−10)

Answer

$-x^2+10x$

Exercise 6.3E.14

−p(p−15)

Exercise 6.3E.15

6r(4r+s)

Answer

$24r^2+6rs$

Exercise 6.3E.16

5c(9c+d)

Exercise 6.3E.17

12x(x−10)

Answer

$12x^2-120x$

Exercise 6.3E.18

9m(m−11)

Exercise 6.3E.19

−9a(3a+5)

Answer

$-27a^2-45a$

Exercise 6.3E.20

−4p(2p+7)

Exercise 6.3E.21

$3\left(p^2+10p+25\right)$

Answer

$3p^2+30p+75$

Exercise 6.3E.22

$6\left(y^2+8y+16\right)$

Exercise 6.3E.23

$-8x\left(x^2+2x-15\right)$

Answer

$-8x^3-16x^2+120x$

Exercise 6.3E.24

$-5t\left(t^2+3t-18\right)$

Exercise 6.3E.25

$5q^3\left(q^3-2q+6\right)$

Answer

$5q^6-10q^4+30q^3$

Exercise 6.3E.26

$4x^3\left(x^4-3x+7\right)$

Exercise 6.3E.27

$-8y\left(y^2+2y-15\right)$

Answer

$-8y^3-16y^2+120y$

Exercise 6.3E.28

$-5m\left(m^2+3m-18\right)$

Exercise 6.3E.29

$5q^3\left(q^2-2q+6\right)$

Answer

$5q^5-10q^4+30q^3$

Exercise 6.3E.30

$9r^3\left(r^2-3r+5\right)$

Exercise 6.3E.31

$-4z^2\left(3z^2+12z-1\right)$

Answer

$-12z^4-48z^3+4z^2$

Exercise 6.3E.32

$-3x^2\left(7x^2+10x-1\right)$

Exercise 6.3E.33

$(2m-9)m$

Answer

$2m^2-9m$

Exercise 6.3E.34

$(8j-1)j$

Exercise 6.3E.35

$(w-6)\cdot 8$

Answer

$8w-48$

Exercise 6.3E.36

$(k-4)\cdot 5$

Exercise 6.3E.37

$4(x+10)$

Answer

4x+40

Exercise 6.3E.38

6(y+8)

Exercise 6.3E.39

15(r−24)

Answer

15r−360

Exercise 6.3E.40

12(v−30)

Exercise 6.3E.41

−3(m+11)

Answer

−3m−33

Exercise 6.3E.42

−4(p+15)

Exercise 6.3E.43

−8(z−5)

Answer

−8z+40

Exercise 6.3E.44

−3(x−9)

Exercise 6.3E.45

u(u+5)

Answer

u^2+5u

Exercise 6.3E.46

$q(q+7)$

Exercise 6.3E.47

$n\left(n^2-3n\right)$

Answer

n^3-3n^2

Exercise 6.3E.48

$s\left(s^2-6s\right)$

Exercise 6.3E.49

$6x(4x+y)$

Answer

$24x^2+6xy$

Exercise 6.3E.50

5a(9a+b)

Exercise 6.3E.51

5p(11p−5q)

Answer

$55p^2-25pq$

Exercise 6.3E.52

$12u(3u-4v)$

Exercise 6.3E.53

$3\left(v^2+10v+25\right)$

Answer

$3v^2+30v+75$

Exercise 6.3E.54

$6\left(x^2+8x+16\right)$

Exercise 6.3E.55

$2n\left(4n^2-4n+1\right)$

Answer

$8n^3-8n^2+2n$

Exercise 6.3E.56

$3r\left(2r^2-6r+2\right)$

Exercise 6.3E.57

$-8y\left(y^2+2y-15\right)$

Answer

$-8y^3-16y^2+120y$

Exercise 6.3E.58

$-5m\left(m^2+3m-18\right)$

Exercise 6.3E.59

$5q^3\left(q^2-2q+6\right)$

Answer

$5q^5-10q^4+30q^3$

Exercise 6.3E.60

$9r^3\left(r^2-3r+5\right)$

Exercise 6.3E.61

$-4z^2\left(3z^2+12z-1\right)$

Answer

$-12z^4-48z^3+4z^2$

Exercise 6.3E.62

$-3x^2\left(7x^2+10x-1\right)$

Exercise 6.3E.63

$(2y-9)y$

Answer

$18y^2-9y$

Exercise 6.3E.64

$(8b-1)b$

Multiply a Binomial by a Binomial

In the following exercises, multiply the following binomials using: ⓐ the Distributive Property ⓑ the FOIL method ⓒ the Vertical Method.

Exercise 6.3E.65

(w+5)(w+7)

Answer

$w^2+12w+35$

Exercise 6.3E.66

(y+9)(y+3)

Exercise 6.3E.67

(p+11)(p−4)

Answer

$p^2+7p-44$

Exercise 6.3E.68

(q+4)(q−8)

In the following exercises, multiply the binomials. Use any method.

Exercise 6.3E.69

(x+8)(x+3)

Answer

$x^2+11x+24$

Exercise 6.3E.70

(y+7)(y+4)

Exercise 6.3E.71

(y−6)(y−2)

Answer

$y^2 - 8y + 12$

Exercise 6.3E.72

(x−7)(x−2)

Exercise 6.3E.73

(w−4)(w+7)

Answer

$w^2 + 3w - 28$

Exercise 6.3E.74

$(q-5)(q+8)$

Exercise 6.3E.75

(p+12)(p−5)

Answer

$p^2 + 7p - 60$

Exercise 6.3E.76

(m+11)(m−4)

Exercise 6.3E.77

(6p+5)(p+1)

Answer

$6p^2 + 11p + 5$

Exercise 6.3E.78

$(7m+1)(m+3)$

Exercise 6.3E.79

(2t−9)(10t+1)

Answer

$20t^2 - 88t - 9$

Exercise 6.3E.80

(3r−8)(11r+1)

Exercise 6.3E.81

(5x−y)(3x−6)

Answer

$15x^2 - 3xy - 30x + 6y$

Exercise 6.3E.82

(10a−b)(3a−4)

Exercise 6.3E.83

(a+b)(2a+3b)

Answer

$2a^2 + 5ab + 3b^2$

Exercise 6.3E.84

(r+s)(3r+2s)

Exercise 6.3E.85

(4z−y)(z−6)

Answer

$4z^2 - 24z - zy + 6y$

Exercise 6.3E.86

(5x−y)(x−4)

Exercise 6.3E.87

$\left(x^2 + 3\right)(x + 2)$

Answer

$x^3 + 2x^2 + 3x + 6$

Exercise 6.3E.88

$\left(y^2 - 4\right)(y + 3)$

Exercise 6.3E.89

$\left(x^2+8\right)\left(x^2-5\right)$

Answer

x^4+3x^2-40

Exercise 6.3E.90

$\left(y^2-7\right)\left(y^2-4\right)$

Exercise 6.3E.91

(5ab−1)(2ab+3)

Answer

$10a^2b^2+13ab-3$

Exercise 6.3E.92

(2xy+3)(3xy+2)

Exercise 6.3E.93

(6pq−3)(4pq−5)

Answer

$24p^2q^2-42pq+15$

Exercise 6.3E.94

(3rs−7)(3rs−4)

Multiply a Trinomial by a Binomial

In the following exercises, multiply using ⓐ the Distributive Property ⓑ the Vertical Method.

Exercise 6.3E.95

$(x+5)\left(x^2+4x+3\right)$

Answer

$x^3+9x^2+23x+15$

Exercise 6.3E.96

$(u+4)\left(u^2+3u+2\right)$

Exercise 6.3E.97

$(y+8)\left(4y^2+y-7\right)$

Answer

https://math.libretexts.org/@go/page/30538

$4y^3+33y^2+y-56$

Exercise 6.3E.98

$(a+10)\left(3a^2+a-5\right)$

In the following exercises, multiply. Use either method.

Exercise 6.3E.99

$(w-7)\left(w^2-9w+10\right)$

Answer

$w^3-16w^2+73w-70$

Exercise 6.3E.100

$(p-4)\left(p^2-6p+9\right)$

Exercise 6.3E.101

$(3q+1)\left(q^2-4q-5\right)$

Answer

$3q^3-11q^2-19q-5$

Exercise 6.3E.102

$(6r+1)\left(r^2-7r-9\right)$

Mixed Practice

Exercise 6.3E.103

(10y−6)+(4y−7)

Answer

14y−13

Exercise 6.3E.104

(15p−4)+(3p−5)

Exercise 6.3E.105

$\left(x^2-4x-34\right)-\left(x^2+7x-6\right)$

Answer

−11x−28

Exercise 6.3E.106

$\left(j^2-8j-27\right)-\left(j^2+2j-12\right)$

Exercise 6.3E.107

$5q\left(3q^2-6q+11\right)$

Answer

$15q^3-30q^2+55q$

Exercise 6.3E.108

$8t\left(2t^2-5t+6\right)$

Exercise 6.3E.109

(s−7)(s+9)

Answer

$s^2+2s-63$

Exercise 6.3E.110

(x−5)(x+13)

Exercise 6.3E.111

$\left(y^2-2y\right)(y+1)$

Answer

y^3-y^2-2y

Exercise 6.3E.112

$\left(a^2-3a\right)(4a+5)$

Exercise 6.3E.113

$(3n-4)\left(n^2+n-7\right)$

Answer

$3n^3-n^2-25n+28$

Exercise 6.3E.114

$(6k-1)\left(k^2+2k-4\right)$

https://math.libretexts.org/@go/page/30538

Exercise 6.3E.115

$(7p+10)(7p-10)$

Answer

$49p^2-100$

Exercise 6.3E.116

(3y+8)(3y−8)

Exercise 6.3E.117

$\left(4m^2-3m-7\right)m^2$

Answer

$4m^4-3m^3-7m^2$

Exercise 6.3E.118

$\left(15c^2-4c+5\right)c^4$

Exercise 6.3E.119

$(5a+7b)(5a+7b)$

Answer

$25a^2+70ab+49b^2$

Exercise 6.3E.120

(3x−11y)(3x−11y)

Exercise 6.3E.121

(4y+12z)(4y−12z)

Answer

$16y^2-144z^2$

Everyday Math

Exercise 6.3E.122

Mental math You can use binomial multiplication to multiply numbers without a calculator. Say you need to multiply 13 times 15. Think of 13 as 10+3 and 15 as 10+5.

1. Multiply (10+3)(10+5) by the FOIL method.
2. Multiply 13·15 without using a calculator.
3. Which way is easier for you? Why?

Exercise 6.3E.123

Mental math You can use binomial multiplication to multiply numbers without a calculator. Say you need to multiply 18 times 17. Think of 18 as 20−2 and 17 as 20−3.

1. Multiply (20−2)(20−3) by the FOIL method.
2. Multiply 18·17 without using a calculator.
3. Which way is easier for you? Why?

Answer

1. 306
2. 306
3. Answers will vary.

Writing Exercises

Exercise 6.3E.124

Which method do you prefer to use when multiplying two binomials: the Distributive Property, the FOIL method, or the Vertical Method? Why?

Exercise 6.3E.125

Which method do you prefer to use when multiplying a trinomial by a binomial: the Distributive Property or the Vertical Method? Why?

Answer

Answers will vary.

Exercise 6.3E.126

Multiply the following:

$(x+2)(x-2)$
$(y+7)(y-7)$
$(w+5)(w-5)$

Explain the pattern that you see in your answers.

Exercise 6.3E.127

Multiply the following:

$(m-3)(m+3)$
$(n-10)(n+10)$
$(p-8)(p+8)$

Explain the pattern that you see in your answers.

Answer

Answers may vary.

Exercise 6.3E.128

Multiply the following:

$(p+3)(p+3)$
$(q+6)(q+6)$
$(r+1)(r+1)$

Explain the pattern that you see in your answers.

Exercise 6.3E.129

Multiply the following:

$(x-4)(x-4)$
$(y-1)(y-1)$
$(z-7)(z-7)$

Explain the pattern that you see in your answers.

Answer

Answers may vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
multiply a polynomial by a monomial.			
multiply a binomial by a binomial.			
multiply a trinomial by a binomial.			

b. What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 6.3E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.3E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.4: Special Products

Learning Objectives

By the end of this section, you will be able to:

- Square a binomial using the Binomial Squares Pattern
- Multiply conjugates using the Product of Conjugates Pattern
- Recognize and use the appropriate special product pattern

Note

Before you get started, take this readiness quiz.

Simplify: a. 9^2 b. $(-9)^2$ c. -9^2.

If you missed this problem, review Exercise 1.5.13.

Square a Binomial Using the Binomial Squares Pattern

Mathematicians like to look for patterns that will make their work easier. A good example of this is squaring binomials. While you can always get the product by writing the **binomial** twice and using the methods of the last section, there is less work to do if you learn to use a pattern.

Let's start by looking at $(x+9)^2$.	
What does this mean?	$(x+9)^2$
It means to multiply $(x+9)$ by itself.	$(x+9)(x+9)$
Then, using FOIL, we get:	$x^2+9x+9x+81$
Combining like terms gives:	$x^2+18x+81$
Here's another one:	$(y-7)^2$
Multiply $(y-7)$ by itself.	$(y-7)(y-7)$
Using FOIL, we get:	$y^2-7y-7y+49$
And combining like terms:	$y^2-14y+49$
And one more:	$(2x+3)^2$
Multiply.	$(2x+3)(2x+3)$
Use FOIL:	$4x+6x+6x+9$
Combine like terms.	$4x^2+12x+9$

Look at these results. Do you see any patterns?

What about the number of terms? In each example we squared a binomial and the result was a **trinomial**.

$$(a+b)^2 = \_\_\_\_ + \_\_\_\_ + \_\_\_\_$$

Now look at the ***first term*** in each result. Where did it come from?

$(x+9)^2$	$(y-7)^2$	$(2x+3)^2$
$(x+9)(x+9)$	$(y-7)(y-7)$	$(2x+3)(2x+3)$
$x^2+9x+9x+81$	$y^2-7y-7y+49$	$4x^2+6x+6x+9$
$x^2+18x+81$	$y^2-14y+49$	$4x^2+12x+9$

The first term is the product of the first terms of each binomial. Since the binomials are identical, it is just the square of the first term!

$$(a+b)^2 = a^2 + \_\_\_\_ + \_\_\_\_$$

To get the ***first term*** of the product, ***square the first term***.

Where did the ***last term*** come from? Look at the examples and find the pattern.

The last term is the product of the last terms, which is the square of the last term.

$$(a+b)^2 = \_\_\_\_ + \_\_\_\_ + b^2$$

*To get the **last term** of the product, **square the last term**.*

Finally, look at the ***middle term***. Notice it came from adding the "outer" and the "inner" terms—which are both the same! So the middle term is double the product of the two terms of the binomial.

$$(a+b)^2 = \_\_\_\_ + 2ab + \_\_\_\_$$

$$(a+b)^2 = \_\_\_\_ - 2ab + \_\_\_\_$$

*To get the **middle term** of the product, **multiply the terms and double their product**.*

Putting it all together:

Binomial Square Pattern

If a and b are real numbers, the binomial square pattern is

$$\underbrace{(a+b)^2}_{(\text{binomial})^2} = \underbrace{a^2}_{(\text{first term})^2} + \underbrace{2ab}_{2\times(\text{product of terms})} + \underbrace{a^2}_{(\text{last term})^2}$$

Applying this to two binomial types:

$$\begin{aligned}(a+b)^2 &= a^2+2ab+b^2\\ (a-b)^2 &= a^2-2ab+b^2\end{aligned}$$

To square a binomial:

1. square the first term
2. square the last term
3. double their product

A number example helps verify the pattern.

	$(10+4)^2$
Square the first term.	$10^2 + \_\_\_\_ + \_\_\_\_$
Square the last term.	$10^2 + \_\_\_\_ + \frac{1}{4^2}$
Double their product.	$10^2 + 2\cdot 10\cdot 4 + 4^2$
Simplify.	$100 + 80 + 16$
Simplify.	196

To multiply $(10+4)^2$ usually you'd follow the Order of Operations.

$$\begin{gathered}(10+4)^2\\ (14)^2\\ 196\end{gathered}$$

The pattern works!

Example 6.4.1

Multiply: $(x+5)^2$

Solution:

	$(x+5)^2$
Square the first term.	$x^2+\_\_\_+\_\_\_$
Square the last term.	$x^2+\_\_\_+5^2$
Double the product.	$x^2+2\cdot x\cdot 5+5^2$
Simplify.	$x^2+10x+25$

Try It! 6.4.1

Multiply: $(x+9)^2$

Answer

$x^2+18x+81$

Try It! 6.4.2

Multiply: $(y+11)^2$

Answer

$y^2+22y+121$

Example 6.4.2

Multiply: $(y-3)^2$

Solution:

	$(y-3)^2$
Square the first term.	$y^2-\_\_\_+\_\_\_$
Square the last term.	$y^2-\_\_\_+3^2$
Double the product.	$y^2-2\cdot y\cdot 3+3^2$
Simplify.	y^2-6y+9

Try It! 6.4.3

Multiply: $(x-9)^2$

Answer

$x^2-18x+81$

Try It! 6.4.4

Multiply: $(p-13)^2$

Answer

$p^2-26p+169$

Example 6.4.3

Multiply: $(4x+6)^2$

Solution:

	$(a+b)^2$ $(4x+6)^2$
Use the pattern.	$a^2+2\cdot a\cdot b+b^2$ $(4x)^2+2\cdot 4x\cdot 6+6^2$
Simplify.	$16x^2+48x+36$

Try It! 6.4.5

Multiply: $(6x+3)^2$

Answer

$36x^2+36x+9$

Try It! 6.4.6

Multiply: $(4x+9)^2$

Answer

$16x^2+72x+81$

Example 6.4.4

Multiply:$(2x-3y)^2$

Solution:

	$(a-b)^2$ $(2x-3y)^2$
Use the pattern.	$a^2-2\cdot a\cdot b+b^2$ $(2x)^2-2\cdot 2x\cdot 3y+(3y)^2$
Simplify.	$4x^2-12xy+9y^2$

Try It! 6.4.7

Multiply: $(2c-d)^2$

Answer

$4c^2-4cd+d^2$

Try It! 6.4.8

Multiply: $(4x-5y)^2$

Answer

$16x^2-40xy+25y^2$

Example 6.4.5

Multiply: $\left(4u^3+1\right)^2$

Solution:

	$\left(4u^3+1\right)^2$
Use the pattern.	$(4u^3)^2+2\cdot 4u^3\cdot 1+(1)^2$
Simplify.	$16u^6+8u^3+1$

Try It! 6.4.9

Multiply: $\left(2x^2+1\right)^2$

Answer

$4x^4+4x^2+1$

Try It! 6.4.10

Multiply: $\left(3y^3+2\right)^2$

Answer

$9y^6+12y^3+4$

Multiply Conjugates Using the Product of Conjugates Pattern

We just saw a pattern for squaring binomials that we can use to make multiplying some binomials easier. Similarly, there is a pattern for another product of binomials. But before we get to it, we need to introduce some vocabulary.

What do you notice about these pairs of binomials?

$$(x-9)(x+9) \qquad (y-8)(y+8) \qquad (2x-5)(2x+5)$$

Look at the first term of each **binomial** in each pair.

$$(x-9)(x+9) \qquad (y-8)(y+8) \qquad (2x-5)(2x+5)$$

Notice the first terms are the same in each pair.

Look at the last terms of each binomial in each pair.

$$(x-9)(x+9) \qquad (y-8)(y+8) \qquad (2x-5)(2x+5)$$

Notice the last terms are the same in each pair.

Notice how each pair has one sum and one difference.

$(x-9)(x+9)$ (Difference, Sum) $(y-8)(y+8)$ (Difference, Sum) $(2x-5)(2x+5)$ (Difference, Sum)

A pair of binomials that each have the same first term and the same last term, but one is a sum and one is a difference has a special name. It is called a *conjugate pair* and is of the form (a−b),(a+b).

Definition: Conjugate Pair

A **conjugate pair** is two binomials of the form

$$(a-b),(a+b)$$

The pair of binomials each have the same first term and the same last term, but one binomial is a sum and the other is a difference.

There is a nice pattern for finding the product of conjugates. You could, of course, simply FOIL to get the product, but using the pattern makes your work easier.

Let's look for the pattern by using FOIL to multiply some conjugate pairs.

$(x-9)(x+9)$	$(y-8)(y+8)$	$(2x-5)(2x+5)$
$x^2+9x-9x-81$	$y^2+8y-8y-64$	$4x^2+10x-10x-25$
x^2-81	y^2-64	$4x^2-25$

Each **first term** is the product of the first terms of the binomials, and since they are identical it is the square of the first term.

$$(a+b)(a-b)=a^2-\_\_\_\_$$

To get the **first term, square the first term.**

The **last term** came from multiplying the last terms, the square of the last term.

$$(a+b)(a-b)=a^2-b^2$$

To get the **last term, square the last term.**

What do you observe about the products?

The product of the two binomials is also a binomial! Most of the products resulting from FOIL have been trinomials.


Why is there no middle term? Notice the two middle terms you get from FOIL combine to 0 in every case, the result of one addition and one subtraction.

The product of conjugates is always of the form a^2-b^2. This is called a difference of squares.

This leads to the pattern:

PRODUCT OF CONJUGATES PATTERN

If a and b are real numbers,



The product is called a **difference of squares**.

To multiply conjugates, square the first term, square the last term, and write the product as a difference of squares.

Let's test this pattern with a numerical example.

	$(10-2)(10+2)$
It is the product of conjugates, so the result will be the difference of two squares.	
	$\_\_\_\_ - \_\_\_\_$
Square the first term.	$10^2 - \_\_\_\_$
Square the last term.	$10^2 - 2^2$
Simplify.	$100 - 4$
Simplify.	96
What do you get using the Order of Operations?	
	$(10-2)(10+2)$
	$(8)(12)$
	96

Notice, the result is the same!

Example 6.4.6

Multiply: $(x-8)(x+8)$

Solution:

First, recognize this as a product of conjugates. The binomials have the same first terms, and the same last terms, and one binomial is a sum and the other is a difference.

It fits the pattern.	$(a-b)(a+b)$ $(x-8)(x+8)$
Square the first term, x.	$a^2 - b^2$ $x^2 - \_\_$
Square the last term, 8.	$a^2 - b^2$ $x^2 - 8^2$
The product is a difference of squares.	$a^2 - b^2$ $x^2 - 64$

Try It! 6.4.11

Multiply: $(x-5)(x+5)$

Answer

$x^2 - 25$

Try It! 6.4.12

Multiply: $(w-3)(w+3)$

Answer

$w^2 - 9$

Example 6.4.7

Multiply: $(2x+5)(2x-5)$

Solution:

Are the binomials conjugates?

It is the product of conjugates.	$(a+b)(a-b)$ $(2x+5)(2x-5)$
Square the first term, $2x$.	a^2-b^2 $(2x)^2-\_\_$
Square the last term, 5.	a^2-b^2 $(2x)^2-5^2$
Simplify. The product is a difference of squares.	a^2-b^2 $4x^2-25$

Try It! 6.4.13

Multiply: $(6x+5)(6x-5)$

Answer

$36x^2-25$

Try It! 6.4.14

Multiply: $(2x+7)(2x-7)$

Answer

$4x^2-49$

The binomials in the next example may look backwards – the variable is in the second term. But the two binomials are still conjugates, so we use the same pattern to multiply them.

Example 6.4.8

Find the product:$(3+5x)(3-5x)$

Solution:

It is the product of conjugates.	$(a-b)(a+b)$ $(3+5x)(3-5x)$
Use the pattern.	a^2-b^2 $3^2-(5x)^2$
Simplify.	$9-25x^2$

Try It! 6.4.15

Multiply: $(7+4x)(7-4x)$

Answer

$49-16x^2$

Try It! 6.4.16

Multiply: $(9-2y)(9+2y)$

Answer

$81-4y^2$

Now we'll multiply conjugates that have two variables.

Example 6.4.9

Find the product: $(5m-9n)(5m+9n)$

Solution:

This fits the pattern.	$(\underset{5m}{a}-\underset{9n}{b})(\underset{5m}{a}+\underset{9n}{b})$
Use the pattern.	$\underset{(5m)^2}{a^2}-\underset{(9n)^2}{b^2}$
Simplify.	$25m^2-81n^2$

Try It! 6.4.17

Find the product: $(4p-7q)(4p+7q)$

Answer

$16p^2-49q^2$

Try It! 6.4.18

Find the product: $(3x-y)(3x+y)$

Answer

$9x^2-y^2$

Example 6.4.10

Find the product: $(cd-8)(cd+8)$

Solution:

This fits the pattern.	$(\underset{cd}{a}-\underset{8}{b})(\underset{cd}{a}+\underset{8}{b})$
Use the pattern.	$\underset{(cd)^2}{a^2}-\underset{(8)^2}{b^2}$
Simplify.	c^2d^2-64

Try It! 6.4.19

Find the product: $(xy-6)(xy+6)$

Answer

x^2y^2-36

Try It! 6.4.20

Find the product: $(ab-9)(ab+9)$

Answer

a^2b^2-81

Example 6.4.11

Find the product: $\left(6u^2-11v^5\right)\left(6u^2+11v^5\right)$

Solution:

This fits the pattern.	$\left(\underset{6u^2}{a}-\underset{11v^5}{b}\right)\left(\underset{6u^2}{a}+\underset{11v^5}{b}\right)$
Use the pattern.	$\underset{(6u^2)^2}{a^2}-\underset{(11v^5)^2}{b^2}$
Simplify.	$36u^4-121v^{10}$

Try It! 6.4.21

Find the product: $\left(3x^2-4y^3\right)\left(3x^2+4y^3\right)$

Answer

$9x^4-16y^6$

Try It! 6.4.22

Find the product: $\left(2m^2-5n^3\right)\left(2m^2+5n^3\right)$

Answer

$4m^4-25n^6$

Recognize and Use the Appropriate Special Product Pattern

We just developed special product patterns for Binomial Squares and for the Product of Conjugates. The products look similar, so it is important to recognize when it is appropriate to use each of these patterns and to notice how they differ. Look at the two patterns together and note their similarities and differences.

COMPARING THE SPECIAL PRODUCT PATTERNS

Binomial Squares	**Product of Conjugates**
$(a+b)^2=a^2+2ab+b^2$	$(a-b)(a+b)=a^2-b^2$
$(a-b)^2=a^2+2ab+b^2$	
- Squaring a binomial	- Multiplying conjugates
- Product is a trinomial	- Product is a binomial
- Inner and outer terms with FOIL are the same.	- Inner and outer terms with FOIL are opposites.
- Middle term is double the product of the terms.	- There is no middle term.

Example 6.4.12

Choose the appropriate pattern and use it to find the product:

1. $(2x-3)(2x+3)$
2. $(8x-5)^2$
3. $(6m+7)^2$
4. $(5x-6)(6x+5)$

Solution:

1. $(2x-3)(2x+3)$ These are conjugates. They have the same first numbers, and the same last numbers, and one binomial is a sum and the other is a difference. It fits the Product of Conjugates pattern.

This fits the pattern.	$(\underset{2x}{a}-\underset{3}{b})(\underset{2x}{a}+\underset{3}{b})$
Use the pattern.	$\underset{(2x)^2}{a^2}-\underset{3^2}{b^2}$
Simplify.	$4x^2-9$

2. $(8x-5)^2$ We are asked to square a binomial. It fits the **binomial squares** pattern.

	$(\underset{8x}{a}-\underset{5}{b})^2$
Use the pattern.	$\underset{(8x)^2}{a^2}-\underset{2\cdot 8x\cdot 5}{2ab}+\underset{5^2}{b^2}$
Simplify.	$64x^2-80x+25$

3. $(6m+7)^2$ Again, we will square a binomial so we use the **binomial squares** pattern.

	$(\underset{6m}{a}+\underset{7}{b})^2$
Use the pattern.	$\underset{(6m)^2}{a^2}+\underset{2\cdot 6m\cdot 7}{2ab}+\underset{7^2}{b^2}$
Simplify.	$36m^2+84m+49$

4. $(5x-6)(6x+5)$ This product does not fit the patterns, so we will use FOIL.

	$(5x-6)(6x+5)$
Use FOIL.	$30x^2+25x-36x-30$
Simplify.	$30x^2-11x-30$

? Try It! 6.4.23

Choose the appropriate pattern and use it to find the product:

1. $(9b-2)(2b+9)$
2. $(9p-4)2$
3. $(7y+1)^2$
4. $(4r-3)(4r+3)$

Answer

1. FOIL; $18b^2+77b-18$
2. Binomial Squares; $81p^2-72p+16$
3. Binomial Squares; $49y^2+14y+1$
4. Product of Conjugates; $16r^2-9$

? Try It! 6.4.24

Choose the appropriate pattern and use it to find the product:

1. $(6x+7)^2$
2. $(3x-4)(3x+4)$
3. $(2x-5)(5x-2)$
4. $(6n-1)^2$

Answer

1. Binomial Squares; $36x^2+84x+49$
2. Product of Conjugates; $9x^2-16$
3. FOIL; $10x^2-29x+10$
4. Binomial Squares; $36n^2-12n+1$

Note

Access these online resources for additional instruction and practice with special products:

- Special Products

Key Concepts

- **Binomial Squares Pattern**
  - If a,b are real numbers,

$$\underbrace{(a+b)^2}_{(\text{binomial})^2} = \underbrace{a^2}_{(\text{first term})^2} + \underbrace{2ab}_{2(\text{product of terms})} + \underbrace{b^2}_{(\text{last term})^2}$$

  - $(a+b)^2=a^2+2ab+b^2$
  - $(a-b)^2=a^2-2ab+b^2$
  - To square a binomial: square the first term, square the last term, double their product.
- **Product of Conjugates Pattern**
  - If a,ba,b are real numbers,

$$\underbrace{(a-b)(a+b)}_{\text{conjugates}} = \underbrace{a^2}_{\text{squares}} \overbrace{-}^{\text{difference}} \underbrace{b^2}_{\text{squares}}$$

  - $(a-b)(a+b)=a^2-b^2$
  - The product is called a difference of squares.
- **To multiply conjugates:**
  - **square the first term square the last term** write it as a difference of squares

Glossary

conjugate pair

A conjugate pair is two binomials of the form $(a-b)$ and $(a+b)$; the pair of binomials each have the same first term and the same last term, but one binomial is a sum and the other is a difference.

This page titled 6.4: Special Products is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.4: Special Products** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.4E: Exercises

Practice Makes Perfect

Square a Binomial Using the Binomial Squares Pattern

In the following exercises, square each binomial using the Binomial Squares Pattern.

Exercise 1

$(w+4)^2$

Exercise 2

$(q+12)^2$

Answer

$q^2+24q+144$

Exercise 3

$(y+14)^2$

Exercise 4

$(x+\frac{2}{3})^2$

Answer

$x^2+\frac{4}{3}x+\frac{4}{9}$

Exercise 5

$(b-7)^2$

Exercise 6

$(y-6)^2$

Answer

$y^2-12y+36$

Exercise 7

$(m-15)^2$

Exercise 8

$(p-13)^2$

Answer

$p^2-26p+169$

Exercise 9

$(3d+1)^2$

Exercise 10

$(4a+10)^2$

Answer

$16a^2+80a+100$

Exercise 11

$(2q+13)^2$

Exercise 12

$(3z+15)^2$

Answer

$9z^2+65z+125$

Exercise 13

$(3x-y)^2$

Exercise 14

$(2y-3z)^2$

Answer

$4y^2-12yz+9z^2$

Exercise 15

$(15x-17y)^2$

Exercise 16

$(18x-19y)^2$

Answer

$164x^2-136xy+181y^2$

Exercise 17

$(3x2+2)^2$

Exercise 18

$(5u^2+9)^2$

Answer

$25u^4+90u^2+81$

Exercise 19

$(4y^3-2)^2$

Exercise 20

$(8p^3-3)^2$

Answer

$64p^6-48p^3+9$

Multiply Conjugates Using the Product of Conjugates Pattern

In the following exercises, multiply each pair of conjugates using the Product of Conjugates Pattern.

Exercise 21

$(m-7)(m+7)$

Exercise 22

$(c-5)(c+5)$

Answer

c^2-25

Exercise 23

$(x+34)(x-34)$

Exercise 24

$(b+\frac{6}{7})(b-\frac{6}{7})$

Answer

$b^2-\frac{36}{49}$

Exercise 25

$(5k+6)(5k-6)$

Exercise 26

$(8j+4)(8j-4)$

Answer

$64j^2 - 16$

Exercise 27

$(11k + 4)(11k - 4)$

Exercise 28

$(9c + 5)(9c - 5)$

Answer

$81c^2 - 25$

Exercise 29

$(11 - b)(11 + b)$

Exercise 30

$(13 - q)(13 + q)$

Answer

$169 - q^2$

Exercise 31

$(5 - 3x)(5 + 3x)$

Exercise 32

$(4 - 6y)(4 + 6y)$

Answer

$16 - 36y^2$

Exercise 33

$(9c - 2d)(9c + 2d)$

Exercise 34

$(7w + 10x)(7w - 10x)$

Answer

$49w^2 - 100x^2$

Exercise 35

$(m + \frac{2}{3}n)(m - \frac{2}{3}n)$

Exercise 36

$(p+\frac{4}{5}q)(p-\frac{4}{5}q)$

Answer

$p^2-\frac{16}{25}q^2$

Exercise 37

$(ab-4)(ab+4)$

Exercise 38

$(xy-9)(xy+9)$

Answer

x^2y^2-81

Exercise 39

$(uv-\frac{3}{5})(uv+\frac{3}{5})$

Exercise 40

$(rs-\frac{2}{7})(rs+\frac{2}{7})$

Answer

$r^2s^2-\frac{4}{49}$

Exercise 41

$(2x^2-3y^4)(2x^2+3y^4)$

Exercise 42

$(6m^3-4n^5)(6m^3+4n^5)$

Answer

$36m^6-16n^{10}$

Exercise 43

$(12p^3-11q^2)(12p^3+11q^2)$

Exercise 44

$(15m^2-8n^4)(15m^2+8n^4)$

Answer

$225m^4-64n^8$

Recognize and Use the Appropriate Special Product Pattern

In the following exercises, find each product.

Exercise 45

a. $(p-3)(p+3)$

b. $(t-9)^2$

c. $(m+n)^2$

d. $(2x+y)(x-2y)$

Exercise 46

a. $(2r+12)^2$

b. $(3p+8)(3p-8)$

c. $(7a+b)(a-7b)$

d. $(k-6)^2$

Answer

a. $4r^2+48r+144$

b. $9p^2-64$

c. $7a^2-48ab-7b^2$

d. $k^2-12k+36$

Exercise 47

a. $(a^5-7b)^2$

b. $(x^2+8y)(8x-y^2)$

c. $(r^6+s^6)(r^6-s^6)$

d. $(y^4+2z)^2$

Exercise 48

a. $(x^5+y^5)(x^5-y^5)$

b. $(m^3-8n)^2$

c. $(9p+8q)^2$

d. $(r^2-s^3)(r^3+s^2)$

Answer

a. $x^{10}-y^{10}$

b. $m^6-16m^3n+64n^2$

c. $81p^2+144pq+64q^2$

d. $r^5+r^2s^2-r^3s^3-s^5$

Everyday Math

Exercise 49

Mental math You can use the product of conjugates pattern to multiply numbers without a calculator. Say you need to multiply 47 times 53. Think of 47 as 50−3 and 53 as 50+3

1. Multiply (50−3)(50+3) by using the product of conjugates pattern, $(a-b)(a+b)=a^2-b^2$
2. Multiply 47·53 without using a calculator.
3. Which way is easier for you? Why?

Exercise 50

Mental math You can use the binomial squares pattern to multiply numbers without a calculator. Say you need to square 65. Think of 65 as 60+5.

1. Multiply $(60+5)^2$ by using the binomial squares pattern, $(a+b)^2=a^2+2ab+b^2$
2. Square 65 without using a calculator.
3. Which way is easier for you? Why?

Answer

1. 4,225
2. 4,225
3. Answers will vary.

Writing Exercises

Exercise 51

How do you decide which pattern to use?

Exercise 52

Why does $(a+b)^2$ result in a trinomial, but (a−b)(a+b) result in a binomial?

Answer

Answers will vary.

Exercise 53

Marta did the following work on her homework paper:

$$(3-y)^2$$
$$3^2-y^2$$
$$9-y^2$$

Explain what is wrong with Marta's work.

Exercise 54

Use the order of operations to show that $(3+5)^2$ is 64, and then use that numerical example to explain why $(a+b)^2\neq a^2+b^2$

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
square a binomial using the binomial squares pattern.			
multiply conjugates using the product of conjugates pattern.			
recognize and use the appropriate special product pattern.			

ⓑ On a scale of 1-10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 6.4E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.4E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.5: Divide Monomials

Learning Objectives

By the end of this section, you will be able to:

- Simplify expressions using the Quotient Property for Exponents
- Simplify expressions with zero exponents
- Simplify expressions using the quotient to a Power Property
- Simplify expressions by applying several properties
- Divide monomials

Note

Before you get started, take this readiness quiz.

1. Simplify: $\frac{8}{24}$.
   If you missed this problem, review Example 1.6.4.
2. Simplify: $(2m^3)^5$.
   If you missed this problem, review Example 6.2.22.
3. Simplify: $\frac{12x}{12y}$
   If you missed this problem, review Example 1.6.10.

Simplify Expressions Using the Quotient Property for Exponents

Earlier in this chapter, we developed the properties of exponents for multiplication. We summarize these properties below.

SUMMARY OF EXPONENT PROPERTIES FOR MULTIPLICATION

If a and b are real numbers, and m and n are whole numbers, then

$$\begin{aligned} &\textbf{Product Property} && a^m \cdot a^n = a^{m+n} \\ &\textbf{Power Property} && (a^m)^n = a^{mn} \\ &\textbf{Product to a Power} && (ab)^m = a^m b^m \end{aligned} \tag{6.5.1}$$

Now we will look at the exponent properties for division. A quick memory refresher may help before we get started. You have learned to simplify fractions by dividing out common factors from the numerator and denominator using the Equivalent Fractions Property. This property will also help you work with algebraic fractions—which are also quotients.

EQUIVALENT FRACTIONS PROPERTY

If a, b, and c are whole numbers where $b \neq 0, c \neq 0$.

$$\text{then} \quad \frac{a}{b} = \frac{a \cdot c}{b \cdot c} \quad \text{and} \quad \frac{a \cdot c}{b \cdot c} = \frac{a}{b} \tag{6.5.2}$$

As before, we'll try to discover a property by looking at some examples.

Consider	$\frac{x^5}{x^2}$	and	$\frac{x^2}{x^3}$
What do they mean?	$\frac{x \cdot x \cdot x \cdot x \cdot x}{x \cdot x}$		$\frac{x \cdot x}{x \cdot x \cdot x}$
Use the Equivalent Fractions Property.	$\frac{\cancel{x} \cdot \cancel{x} \cdot x \cdot x \cdot x}{\cancel{x} \cdot \cancel{x}}$		$\frac{\cancel{x} \cdot \cancel{x} \cdot 1}{\cancel{x} \cdot \cancel{x} \cdot x}$
Simplify.	x^3		$\frac{1}{x}$

(6.5.3)

Notice, in each case the bases were the same and we subtracted exponents.

When the larger exponent was in the numerator, we were left with factors in the numerator.

When the larger exponent was in the denominator—notice the numerator of 1.

We write:

$\frac{x^5}{x^2}$	$\frac{x^2}{x^3}$
x^{5-2}	$\frac{1}{x^{3-2}}$
x^3	$\frac{1}{x}$

(6.5.4)

This leads to the *Quotient Property for Exponents*.

QUOTIENT PROPERTY FOR EXPONENTS

If a is a real number, $a \neq 0$, and m and n are whole numbers, then

$$\frac{a^m}{a^n} = a^{m-n}, m > n \text{ and } \frac{a^m}{a^n} = \frac{1}{a^{n-m}}, n > m \tag{6.5.5}$$

A couple of examples with numbers may help to verify this property.

$\frac{3^4}{3^2}$	$=$	3^{4-2}	$\frac{5^2}{5^3}$	$=$	$\frac{1}{5^{3-2}}$
$\frac{81}{9}$	$=$	3^2	$\frac{25}{125}$	$=$	$\frac{1}{5^1}$
9	$=$	$9\checkmark$	$\frac{1}{5}$	$=$	$\frac{1}{5}\checkmark$

(6.5.6)

Example 6.5.1

Simplify:

1. $\frac{x^9}{x^7}$
2. $\frac{3^{10}}{3^2}$

Solution

To simplify an expression with a quotient, we need to first compare the exponents in the numerator and denominator.

1.

Since 9 > 7, there are more factors of *x* in the numerator.	$\frac{x^9}{x^7}$
Use the Quotient Property, $\frac{a^m}{a^n} = a^{m-n}$	x^{9-7}

Simplify.	x^2

2.

Since 10 > 2, there are more factors of x in the numerator.	$\frac{3^{10}}{3^2}$
Use the Quotient Property, $\frac{a^m}{a^n} = a^{m-n}$	3^{10-2}
Simplify.	3^8

Notice that when the larger exponent is in the numerator, we are left with factors in the numerator.

Try It 6.5.2

Simplify:

a. $\frac{x^{15}}{x^{10}}$
b. $\frac{6^{14}}{6^5}$

Answer

1. x^5
2. 6^9

Try It 6.5.3

Simplify:

1. $\frac{y^{43}}{y^{37}}$
2. $\frac{10^{15}}{10^7}$

Answer

1. y^6
2. 10^8

Example 6.5.4

Simplify:

1. $\frac{b^8}{b^{12}}$
2. $\frac{7^3}{7^5}$

Solution

To simplify an expression with a quotient, we need to first compare the exponents in the numerator and denominator.

1.

Since 12 > 8, there are more factors of b in the denominator.	$\frac{b^8}{b^{12}}$

Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	$\frac{1}{b^{12-8}}$
Simplify.	$\frac{1}{b^4}$

2.

Since 5 > 3, there are more factors of 3 in the denominator.	$\frac{7^3}{7^5}$
Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	$\frac{1}{7^{5-3}}$
Simplify.	$\frac{1}{7^2}$
Simplify.	$\frac{1}{49}$

Notice that when the larger exponent is in the denominator, we are left with factors in the denominator.

Try It 6.5.5

Simplify:

1. $\frac{x^{18}}{x^{22}}$
2. $\frac{12^{15}}{12^{30}}$

Answer

1. $\frac{1}{x^4}$
2. $\frac{1}{12^{15}}$

Try It 6.5.6

Simplify:

1. $\frac{m^7}{m^{15}}$
2. $\frac{9^8}{9^{19}}$

Answer

1. $\frac{1}{m^8}$
2. $\frac{1}{9^{11}}$

Notice the difference in the two previous examples:

- If we start with more factors in the numerator, we will end up with factors in the numerator.
- If we start with more factors in the denominator, we will end up with factors in the denominator.

The first step in simplifying an expression using the Quotient Property for Exponents is to determine whether the exponent is larger in the numerator or the denominator.

Example 6.5.7

Simplify:

1. $\frac{a^5}{a^9}$
2. $\frac{x^{11}}{x^7}$

Solution

1. Is the exponent of a larger in the numerator or denominator? Since 9 > 5, there are more a's in the denominator and so we will end up with factors in the denominator.

	$\frac{a^5}{a^9}$
Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	$\frac{1}{a^{9-5}}$
Simplify.	$\frac{1}{a^4}$

2. Notice there are more factors of xx in the numerator, since 11 > 7. So we will end up with factors in the numerator.

	$\frac{x^{11}}{x^7}$
Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	x^{11-7}
Simplify.	x^4

Try It 6.5.8

Simplify:

1. $\frac{b^{19}}{b^{11}}$
2. $\frac{z^5}{z^{11}}$

Answer

1. b^8
2. $\frac{1}{z^6}$

Try It 6.5.9

Simplify:

1. $\frac{p^9}{p^{17}}$
2. $\frac{w^{13}}{w^9}$

Answer

1. $\frac{1}{p^8}$
2. w^4

Simplify Expressions with an Exponent of Zero

A special case of the Quotient Property is when the exponents of the numerator and denominator are equal, such as an expression like $\frac{a^m}{a^m}$. From your earlier work with fractions, you know that:

$$\frac{2}{2}=1 \quad \frac{17}{17}=1 \quad \frac{-43}{-43}=1 \tag{6.5.7}$$

In words, a number divided by itself is 1. So, $\frac{x}{x}=1$, for any $x(x\neq 0)$, since any number divided by itself is 1.

The Quotient Property for Exponents shows us how to simplify $\frac{a^m}{a^n}$ when $m>n$ and when $n<m$ by subtracting exponents. What if $m=n$?

Consider $\frac{8}{8}$, which we know is 1.

	$\frac{8}{8}$	$=$	1
Write 8 as 2^3.	$\frac{2^3}{2^3}$	$=$	1
Subtract exponents.	2^{3-3}	$=$	1
Simplify.	2^0	$=$	1

Now we will simplify $\frac{a^m}{a^m}$ in two ways to lead us to the definition of the zero exponent. In general, for $a\neq 0$:

$\frac{a^m}{a^m}$	$\frac{a^m}{a^m}$
a^{m-m}	$\frac{\overbrace{y\cdot y\cdot\cdots\cdot y}^{m \text{ factors}}}{\underbrace{y\cdot y\cdot\cdots\cdot y}_{m \text{ factors}}}$
a^0	1

We see $\frac{a^m}{a^m}$ simplifies to a^0 and to 1. So $a^0=1$.

ZERO EXPONENT

If a is a non-zero number, then $a^0=1$.

Any nonzero number raised to the zero power is 1.

In this text, we assume any variable that we raise to the zero power is not zero.

Example 6.5.10

Simplify:

a. 9^0
b. n^0

Solution

The definition says any non-zero number raised to the zero power is 1.

a.	Use the definition of the zero exponent.	9^0 1
b.	Use the definition of the zero exponent.	n^0 1

Try It 6.5.11

Simplify:

1. 15^0
2. m^0

Answer

1. 1
2. 1

Try It 6.5.12

Simplify:

a. k^0
b. 29^0

Answer

a. 1
b. 1

Now that we have defined the zero exponent, we can expand all the Properties of Exponents to include whole number exponents.

What about raising an expression to the zero power? Let's look at $(2x)^0$. We can use the product to a power rule to rewrite this expression.

$$\begin{array}{ll} & (2x)^0 \\ \text{Use the product to a power rule.} & 2^0x^0 \\ \text{Use the zero exponent property.} & 1 \cdot 1 \\ \text{Simplify.} & 1 \end{array} \tag{6.5.8}$$

This tells us that any nonzero expression raised to the zero power is one.

Example 6.5.13

Simplify:

a. $(5b)^0$
b. $(-4a^2b)^0$.

Solution

a.	Use the definition of the zero exponent.	$(5b)^0$ 1
b.	Use the definition of the zero exponent.	$(-4a^2b)^0$ 1

Try It 6.5.14

Simplify:

a. $(11z)^0$
b. $(-11pq^3)^0$.

Answer

a. 1
b. 1

Try It 6.5.15

Simplify:

a. $(-6d)^0$
b. $(-8m^2n^3)^0$.

Answer

a. 1
b. 1

Simplify Expressions Using the Quotient to a Power Property

Now we will look at an example that will lead us to the Quotient to a Power Property.

$$\begin{array}{ll} & \left(\frac{x}{y}\right)^3 \\ \text{This means:} & \frac{x}{y}\cdot\frac{x}{y}\cdot\frac{x}{y} \\ \text{Multiply the fractions.} & \frac{x\cdot x\cdot x}{y\cdot y\cdot y} \\ \text{Write with exponents.} & \frac{x^3}{y^3} \end{array} \tag{6.5.9}$$

Notice that the exponent applies to both the numerator and the denominator.

$$\begin{array}{ll} \text{We see that } \left(\frac{x}{y}\right)^3 \text{ is } \frac{x^3}{y^3} & \\ \text{We write:} & \left(\frac{x}{y}\right)^3 \\ & \frac{x^3}{y^3} \end{array} \tag{6.5.10}$$

This leads to the *Quotient to a Power Property for Exponents*.

QUOTIENT TO A POWER PROPERTY FOR EXPONENTS

If a and b are real numbers, $b\neq 0$, and m is a counting number, then

$$\left(\frac{a}{b}\right)^m=\frac{a^m}{b^m} \tag{6.5.11}$$

To raise a fraction to a power, raise the numerator and denominator to that power.

An example with numbers may help you understand this property:

$$\left(\frac{2}{3}\right)^3 = \frac{2^3}{3^3}$$
$$\frac{2}{3}\cdot\frac{2}{3}\cdot\frac{2}{3} = \frac{8}{27}$$
$$\frac{8}{27} = \frac{8}{27}\checkmark$$

Example 6.5.16

Simplify:

a. $\left(\frac{3}{7}\right)^2$

b. $\left(\frac{b}{3}\right)^4$

c. $\left(\frac{k}{j}\right)^3$

Solution

1.

	$\left(\frac{3}{7}\right)^2$
Use the Quotient Property, $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$	$\frac{3^2}{7^2}$
Simplify.	$\frac{9}{49}$

2.

	$\left(\frac{b}{3}\right)^4$
Use the Quotient Property, $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$	$\frac{b^4}{3^4}$
Simplify.	$\frac{b^4}{81}$

3.

	$\left(\frac{k}{j}\right)^3$
Raise the numerator and denominator to the third power.	$\frac{k^3}{j^3}$

Try It 6.5.17

Simplify:

a. $\left(\frac{5}{8}\right)^2$

b. $\left(\frac{p}{10}\right)^4$

c. $\left(\frac{m}{n}\right)^7$

Answer

a. $\frac{25}{64}$
b. $\frac{p^4}{10,000}$
c. $\frac{m^7}{n^7}$

Try It 6.5.18

Simplify:

a. $\left(\frac{1}{3}\right)^3$
b. $\left(\frac{-2}{q}\right)^3$
c. $\left(\frac{w}{x}\right)^4$

Answer

a. $\frac{1}{27}$
b. $\frac{-8}{q^3}$
c. $\frac{w^4}{x^4}$

Simplify Expressions by Applying Several Properties

We'll now summarize all the properties of exponents so they are all together to refer to as we simplify expressions using several properties. Notice that they are now defined for whole number exponents.

SUMMARY OF EXPONENT PROPERTIES

If a and b are real numbers, and m and n are whole numbers, then

$$\begin{array}{lrcll} \textbf{Product Property} & a^m \cdot a^n & = & a^{m+n} & \\ \textbf{Power Property} & (a^m)^n & = & a^{m \cdot n} & \\ \textbf{Product to a Power} & (ab)^m & = & a^m b^m & \\ \textbf{Quotient Property} & \frac{a^m}{a^n} & = & a^{m-n}, & a \neq 0, m > n \\ & \frac{a^n}{a^n} & = & 1, & a \neq 0, n > m \\ \textbf{Zero Exponent Definition} & a^0 & = & 1, & a \neq 0 \\ \textbf{Quotient to a Power Property} & \left(\frac{a}{b}\right)^m & = & \frac{a^m}{b^m}, & b \neq 0 \end{array} \tag{6.5.12}$$

Example 6.5.19

Simplify: $\frac{\left(y^4\right)^2}{y^6}$

Solution

	$\frac{\left(y^4\right)^2}{y^6}$
Multiply the exponents in the numerator.	$\frac{y^8}{y^6}$
Subtract the exponents.	y^2

Try It 6.5.20

Simplify: $\frac{\left(m^5\right)^4}{m^7}$

Answer

m^{13}

Try It 6.5.21

Simplify: $\frac{\left(k^2\right)^6}{k^7}$

Answer

k^5

Example 6.5.22

Simplify: $\frac{b^{12}}{\left(b^2\right)^6}$

Solution

	$\frac{b^{12}}{\left(b^2\right)^6}$
Multiply the exponents in the numerator.	$\frac{b^{12}}{b^{12}}$
Subtract the exponents.	b^0
Simplify	1

Notice that after we simplified the denominator in the first step, the numerator and the denominator were equal. So the final value is equal to 1.

Try It 6.5.23

Simplify $\frac{n^{12}}{\left(n^3\right)^4}$.

Answer

1

Try It 6.5.24

Simplify $\frac{x^{15}}{\left(x^3\right)^5}$.

Answer

1

Example 6.5.25

Simplify: $\left(\frac{y^9}{y^4}\right)^2$

Solution

	$\left(\frac{y^9}{y^4}\right)^2$
Remember parentheses come before exponents.	
Notice the bases are the same, so we can simplify inside the parentheses. Subtract the exponents.	$\left(y^5\right)^2$
Multiply the exponents.	y^{10}

Try It 6.5.26

Simplify: $\left(\frac{r^5}{r^3}\right)^4$

Answer

r^8

Try It 6.5.27

Simplify: $\left(\frac{v^6}{v^4}\right)^3$

Answer

v^6

Example 6.5.28

Simplify: $\left(\frac{j^2}{k^3}\right)^4$

Solution

Here we cannot simplify inside the parentheses first because the bases are not the same.	$\left(\frac{j^2}{k^3}\right)^4$
Raise the numerator and denominator to the 4th power using the Quotient to a Power Property, $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$	$\frac{\left(j^2\right)^4}{\left(k^3\right)^4}$
Use the Power Property and simplify.	$\frac{j^8}{k^{12}}$

Try It 6.5.29

Simplify: $\left(\frac{a^3}{b^2}\right)^4$

Answer

$\frac{a^{12}}{b^8}$

Try It 6.5.30

Simplify: $\left(\frac{q^7}{r^5}\right)^3$

Answer

$\frac{q^{21}}{r^{15}}$

Example 6.5.31

Simplify: $\left(\frac{2m^2}{5n}\right)^4$

Solution

	$\left(\frac{2m^2}{5n}\right)^4$
Raise the numerator and denominator to the 4th power using the Quotient to a Power Property, $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$	$\frac{(2m^2)^4}{(5n)^4}$
Use the Product to a Power Property, $(ab)^m = a^m b^m$	$\frac{2^4(m^2)^4}{5^4 n^4}$
Use the Power Property and simplify.	$\frac{16m^8}{625n^4}$

Try It 6.5.32

Simplify: $\left(\frac{7x^3}{9y}\right)^2$

Answer

$\frac{49x^6}{81y^2}$

Try It 6.5.33

Simplify: $\left(\frac{3x^4}{7y}\right)^2$

Answer

$\frac{9x^8}{49y^2}$

Example 6.5.34

Simplify: $\frac{\left(x^3\right)^4\left(x^2\right)^5}{\left(x^6\right)^5}$

Solution

	$\frac{\left(x^3\right)^4\left(x^2\right)^5}{\left(x^6\right)^5}$
Use the Power Property, $(a^m)^n = a^{m \cdot n}$	$\frac{\left(x^{12}\right)\left(x^{10}\right)}{\left(x^{30}\right)}$
Add the exponents in the numerator.	$\frac{x^{22}}{x^{30}}$
Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	$\frac{1}{x^8}$

Try It 6.5.35

Simplify: $\frac{\left(a^2\right)^3\left(a^2\right)^4}{\left(a^4\right)^5}$

Answer

$\frac{1}{a^6}$

Try It 6.5.36

Simplify: $\frac{\left(p^3\right)^4\left(p^5\right)^3}{\left(p^7\right)^6}$

Answer

$\frac{1}{p^{15}}$

Example 6.5.37

Simplify: $\frac{\left(10p^3\right)^2}{(5p)^3\left(2p^5\right)^4}$

Solution

	$\frac{(10p^3)^2}{(5p)^3(2p^5)^4}$
Use the Product to a Power Property, $(ab)^m = a^m b^m$	$\frac{(10)^2(p^3)^2}{(5)^3(p)^3(2)^4(p^5)^4}$
Use the Power Property, $(a^m)^n = a^{m \cdot n}$	$\frac{100p^6}{125p^3 \cdot 16p^{20}}$
Add the exponents in the denominator.	$\frac{100p^6}{125 \cdot 16p^{23}}$
Use the Quotient Property, $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$	$\frac{100}{125 \cdot 16p^{17}}$
Simplify.	$\frac{1}{20p^{17}}$

? Try It 6.5.38

Simplify: $\frac{(3r^3)^2(r^3)^7}{(r^3)^3}$

Answer

$9r^{18}$

? Try It 6.5.39

Simplify: $\frac{(2x^4)^5}{(4x^3)^2(x^3)^5}$

Answer

$\frac{2}{x}$

Divide Monomials

You have now been introduced to all the properties of exponents and used them to simplify expressions. Next, you'll see how to use these properties to divide monomials. Later, you'll use them to divide polynomials.

✓ Example 6.5.40

Find the quotient: $56x^7 \div 8x^3$

Solution

	$56x^7 \div 8x^3$
Rewrite as a fraction.	$\frac{56x^7}{8x^3}$
Use fraction multiplication.	$\frac{56}{8} \cdot \frac{x^7}{x^3}$
Simplify and use the Quotient Property.	$7x^4$

Try It 6.5.41

Find the quotient: $42y^9 \div 6y^3$

Answer

$7y^6$

Try It 6.5.42

Find the quotient: $48z^8 \div 8z^2$

Answer

$6z^6$

Example 6.5.43

Find the quotient: $\dfrac{45a^2b^3}{-5ab^5}$

Solution

When we divide monomials with more than one variable, we write one fraction for each variable.

	$\dfrac{45a^2b^3}{-5ab^5}$
Use fraction multiplication.	$\dfrac{45}{-5} \cdot \dfrac{a^2}{a} \cdot \dfrac{b^3}{b^5}$
Simplify and use the Quotient Property.	$-9 \cdot a \cdot \dfrac{1}{b^2}$
Multiply.	$-\dfrac{9a}{b^2}$

Try It 6.5.44

Find the quotient: $\dfrac{-72a^7b^3}{8a^{12}b^4}$

Answer

$-\dfrac{9}{a^5b}$

Try It 6.5.45

Find the quotient: $\dfrac{-63c^8d^3}{7c^{12}d^2}$

Answer

$\dfrac{-9d}{c^4}$

Example 6.5.46

Find the quotient: $\frac{24a^5b^3}{48ab^4}$

Solution

	$\frac{24a^5b^3}{48ab^4}$
Use fraction multiplication.	$\frac{24}{48}\cdot\frac{a^5}{a}\cdot\frac{b^3}{b^4}$
Simplify and use the Quotient Property.	$\frac{1}{2}\cdot a^4\cdot\frac{1}{b}$
Multiply.	$\frac{a^4}{2b}$

Try It 6.5.47

Find the quotient: $\frac{16a^7b^6}{24ab^8}$

Answer

$\frac{2a^6}{3b^2}$

Try It 6.5.48

Find the quotient: $\frac{27p^4q^7}{-45p^{12}q}$

Answer

$-\frac{3q^6}{5p^8}$

Once you become familiar with the process and have practiced it step by step several times, you may be able to simplify a fraction in one step.

Example 6.5.49

Find the quotient: $\frac{14x^7y^{12}}{21x^{11}y^6}$

Solution

Be very careful to simplify $\frac{14}{21}$ by dividing out a common factor, and to simplify the variables by subtracting their exponents.

	$\frac{14x^7y^{12}}{21x^{11}y^6}$
Simplify and use the Quotient Property.	$\frac{2y^6}{3x^4}$

Try It 6.5.50

Find the quotient: $\frac{28x^5y^{14}}{49x^9y^{12}}$

Answer

$\frac{4y^2}{7x^4}$

Try It 6.5.51

Find the quotient: $\frac{30m^5n^{11}}{48m^{10}n^{14}}$

Answer

$\frac{5}{8m^5n^3}$

In all examples so far, there was no work to do in the numerator or denominator before simplifying the fraction. In the next example, we'll first find the product of two monomials in the numerator before we simplify the fraction. This follows the order of operations. Remember, a fraction bar is a grouping symbol.

Example 6.5.52

Find the quotient: $\frac{\left(6x^2y^3\right)\left(5x^3y^2\right)}{\left(3x^4y^5\right)}$

Solution

Answer

	$\frac{\left(6x^2y^3\right)\left(5x^3y^2\right)}{\left(3x^4y^5\right)}$
Simplify the numerator.	$\frac{30x^5y^5}{3x^4y^5}$
Simplify.	$10x$

Try It 6.5.53

Find the quotient: $\frac{\left(6a^4b^5\right)\left(4a^2b^5\right)}{12a^5b^8}$

Answer

$2ab^2$

Try It 6.5.54

Find the quotient: $\frac{\left(-12x^6y^9\right)\left(-4x^5y^8\right)}{-12x^{10}y^{12}}$

Answer

$-4xy^5$

Note

Access these online resources for additional instruction and practice with dividing monomials:

- Rational Expressions
- Dividing Monomials
- Dividing Monomials 2

Key Concepts

- **Quotient Property for Exponents:**
  - If a is a real number, $a \neq 0$, and m,n are whole numbers, then: $\frac{a^m}{a^n} = a^{m-n}, m > n$ and $\frac{a^m}{a^n} = \frac{1}{a^{m-n}}, n > m$
- **Zero Exponent**
  - If a is a non-zero number, then $a^0 = 1$.
- **Quotient to a Power Property for Exponents**:
  - If a and b are real numbers, $b \neq 0$, and mm is a counting number, then: $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$
  - To raise a fraction to a power, raise the numerator and denominator to that power.
- **Summary of Exponent Properties**
  - If a,b are real numbers and m,nm,n are whole numbers, then

Product Property	$a^m \cdot a^n$	$=$	a^{m+n}
Power Property	$(a^m)^n$	$=$	$a^{m \cdot n}$
Product to a Power	$(ab)^m$	$=$	$a^m b^m$
Quotient Property	$\frac{a^m}{a^n}$	$=$	$a^{m-n}, a \neq 0, m > n$
	$\frac{a^n}{a^n}$	$=$	$1, a \neq 0, n > m$
Zero Exponent Definition	a^0	$=$	$1, a \neq 0$
Quotient to a Power Property	$\left(\frac{a}{b}\right)^m$	$=$	$\frac{a^m}{b^m}, b \neq 0$

This page titled 6.5: Divide Monomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.5: Divide Monomials** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.5E: Exercises

Practice Makes Perfect

Simplify Expressions Using the Quotient Property for Exponents

In the following exercises, simplify.

Exercise 6.5E.1

a. $\frac{x^{18}}{x^3}$

b. $\frac{5^{12}}{5^3}$

Exercise 6.5E.2

a. $\frac{y^{20}}{y^{10}}$

b. $\frac{7^{16}}{7^2}$

Answer

a. y^{10}

b. 7^{14}

Exercise 6.5E.3

a. $\frac{p^{21}}{p^7}$

b. $\frac{4^{16}}{4^4}$

Exercise 6.5E.4

a. $\frac{u^{24}}{u^3}$

b. $\frac{9^{15}}{9^5}$

Answer

a. u^{21}

b. 9^{10}

Exercise 6.5E.5

a. $\frac{q^{18}}{q^{36}}$

b. $\frac{10^2}{10^3}$

Exercise 6.5E.6

a. $\frac{t^{10}}{t^{40}}$
b. $\frac{8^3}{8^5}$

Answer

a. $\frac{1}{t^{30}}$
b. $\frac{1}{64}$

Exercise 6.5E.7

a. $\frac{b}{b^9}$
b. $\frac{4}{4^6}$

Exercise 6.5E.8

a. $\frac{x}{x^7}$
b. $\frac{10}{10^3}$

Answer

a. $\frac{1}{x^6}$
b. $\frac{1}{100}$

Simplify Expressions with Zero Exponents

In the following exercises, simplify.

Exercise 6.5E.9

a. 20^0
b. b^0

Exercise 6.5E.10

a. 13^0
b. k^0

Answer

a. 1
b. 1

Exercise 6.5E.11

a. -27^0
b. $-\left(27^0\right)$

Exercise 6.5E.12

a. -15^0
b. $-\left(15^0\right)$

Answer

a. -1
b. -1

Exercise 6.5E.13

a. $(25x)^0$
b. $25x^0$

Exercise 6.5E.14

a. $(6y)^0$
b. $6y^0$

Answer

a. 1
b. 6

Exercise 6.5E.15

a. $(12x)^0$
b. $\left(-56p^4q^3\right)^0$

Exercise 6.5E.16

a. $7y^0(17y)^0$
b. $\left(-93c^7d^{15}\right)^0$

Answer

a. 7
b. 1

Exercise 6.5E.17

a. $12n^0-18m^0$
b. $(12n)^0-(18m)^0$

Exercise 6.5E.18

a. $15r^0-22s^0$
b. $(15r)^0-(22s)^0$

Answer

a. -7
b. 0

Simplify Expressions Using the Quotient to a Power Property

In the following exercises, simplify.

Exercise 6.5E.19

a. $\left(\frac{3}{4}\right)^3$

b. $\left(\frac{p}{2}\right)^5$

c. $\left(\frac{x}{y}\right)^6$

Exercise 6.5E.20

a. $\left(\frac{2}{5}\right)^2$

b. $\left(\frac{x}{3}\right)^4$

c. $\left(\frac{a}{b}\right)^5$

Answer

a. $\frac{4}{25}$

b. $\frac{x^4}{81}$

c. $\left(\frac{a}{b}\right)^5$

Exercise 6.5E.21

a. $\left(\frac{a}{3b}\right)^4$

b. $\left(\frac{5}{4m}\right)^2$

Exercise 6.5E.22

a. $\left(\frac{a}{3b}\right)^4$

b. $\left(\frac{10}{3q}\right)^4$

Answer

a. $\frac{x^3}{8y^3}$

b. $\frac{10,000}{81q^4}$

Simplify Expressions by Applying Several Properties

In the following exercises, simplify.

https://math.libretexts.org/@go/page/30542

Exercise 6.5E.23

$$\frac{\left(a^2\right)^3}{a^4}$$

Exercise 6.5E.24

$$\frac{\left(p^3\right)^4}{p^5}$$

Answer

p^7

Exercise 6.5E.25

$$\frac{\left(y^3\right)^4}{y^{10}}$$

Exercise 6.5E.26

$$\frac{\left(x^4\right)^5}{x^{15}}$$

Answer

x^5

Exercise 6.5E.27

$$\frac{u^6}{\left(u^3\right)^2}$$

Exercise 6.5E.28

$$\frac{v^{20}}{\left(v^4\right)^5}$$

Answer

1

Exercise 6.5E.29

$$\frac{m^{12}}{\left(m^8\right)^3}$$

Exercise 6.5E.30

$$\frac{n^8}{\left(n^6\right)^4}$$

Answer

https://math.libretexts.org/@go/page/30542

$$\frac{1}{n^{16}}$$

Exercise 6.5E.31

$$\left(\frac{p^9}{p^3}\right)^5$$

Exercise 6.5E.32

$$\left(\frac{q^8}{q^2}\right)^3$$

Answer

$$q^{18}$$

Exercise 6.5E.33

$$\left(\frac{r^2}{r^6}\right)^3$$

Exercise 6.5E.34

$$\left(\frac{m^4}{m^7}\right)^4$$

Answer

$$\frac{1}{m^{12}}$$

Exercise 6.5E.35

$$\left(\frac{p}{r^{11}}\right)^2$$

Exercise 6.5E.36

$$\left(\frac{a}{b^6}\right)^3$$

Answer

$$\frac{a^3}{b^{18}}$$

Exercise 6.5E.37

$$\left(\frac{w^5}{x^3}\right)^8$$

Exercise 6.5E.38

$$\left(\frac{y^4}{z^{10}}\right)^5$$

Answer

$$\frac{y^{20}}{z^{50}}$$

Exercise 6.5E.39

$$\left(\frac{2j^3}{3k}\right)^4$$

Exercise 6.5E.40

$$\left(\frac{3m^5}{5n}\right)^3$$

Answer

$$\frac{27m^{15}}{125n^3}$$

Exercise 6.5E.41

$$\left(\frac{3c^2}{4d^6}\right)^3$$

Exercise 6.5E.42

$$\left(\frac{5u^7}{2v^3}\right)^4$$

Answer

$$\frac{625u^{28}}{16v^{12}}$$

Exercise 6.5E.43

$$\left(\frac{k^2k^8}{k^3}\right)^2$$

Exercise 6.5E.44

$$\left(\frac{j^2j^5}{j^4}\right)^3$$

Answer

$$j^9$$

https://math.libretexts.org/@go/page/30542

Exercise 6.5E.45

$$\frac{\left(t^2\right)^5\left(t^4\right)^2}{\left(t^3\right)^7}$$

Exercise 6.5E.46

$$\frac{\left(q^3\right)^6\left(q^2\right)^3}{\left(q^4\right)^8}$$

Answer

$$\frac{1}{q^8}$$

Exercise 6.5E.47

$$\frac{\left(-2p^2\right)^4\left(3p^4\right)^2}{\left(-6p^3\right)^2}$$

Exercise 6.5E.48

$$\frac{\left(-2k^3\right)^2\left(6k^2\right)^4}{\left(9k^4\right)^2}$$

Answer

$64k^6$

Exercise 6.5E.49

$$\frac{\left(-4m^3\right)^2\left(5m^4\right)^3}{\left(-10m^6\right)^3}$$

Exercise 6.5E.50

$$\frac{\left(-10n^2\right)^3\left(4n^5\right)^2}{\left(2n^8\right)^2}$$

Answer

−4,000

Divide Monomials

In the following exercises, divide the monomials.

Exercise 6.5E.51

$56b^8 \div 7b^2$

https://math.libretexts.org/@go/page/30542

Exercise 6.5E.52

$63v^{10} \div 9v^2$

Answer

$7v^8$

Exercise 6.5E.53

$-88y^{15} \div 8y^3$

Exercise 6.5E.54

$-72u^{12} \div 12u^4$

Answer

$-6u^8$

Exercise 6.5E.55

$$\frac{45a^6b^8}{-15a^{10}b^2}$$

Exercise 6.5E.56

$$\frac{54x^9y^3}{-18x^6y^{15}}$$

Answer

$$-\frac{3x^3}{y^{12}}$$

Exercise 6.5E.57

$$\frac{15r^4s^9}{18r^9s^2}$$

Exercise 6.5E.58

$$\frac{20m^8n^4}{30m^5n^9}$$

Answer

$$\frac{2m^3}{3n^5}$$

Exercise 6.5E.59

$$\frac{18a^4b^8}{-27a^9b^5}$$

https://math.libretexts.org/@go/page/30542

Exercise 6.5E.60

$$\frac{45x^5y^9}{-60x^8y^6}$$

Answer

$$\frac{-3y^3}{4x^3}$$

Exercise 6.5E.61

$$\frac{64q^{11}r^9s^3}{48q^6r^8s^5}$$

Exercise 6.5E.62

$$\frac{65a^{10}b^8c^5}{42a^7b^6c^8}$$

Answer

$$\frac{65a^3b^2}{42c^3}$$

Exercise 6.5E.63

$$\frac{\left(10m^5n^4\right)\left(5m^3n^6\right)}{25m^7n^5}$$

Exercise 6.5E.64

$$\frac{\left(-18p^4q^7\right)\left(-6p^3q^8\right)}{-36p^{12}q^{10}}$$

Answer

$$\frac{-3q^5}{p^5}$$

Exercise 6.5E.65

$$\frac{\left(6a^4b^3\right)\left(4ab^5\right)}{\left(12a^2b\right)\left(a^3b\right)}$$

Exercise 6.5E.66

$$\frac{\left(4u^2v^5\right)\left(15u^3v\right)}{\left(12u^3v\right)\left(u^4v\right)}$$

Answer

$$\frac{5v^4}{u^2}$$

Mixed Practice

Exercise 6.5E.67

a. $24a^5 + 2a^5$
b. $24a^5 - 2a^5$
c. $24a^5 \cdot 2a^5$
d. $24a^5 \div 2a^5$

Exercise 6.5E.68

a. $15n^{10} + 3n^{10}$
b. $15n^{10} - 3n^{10}$
c. $15n^{10} \cdot 3n^{10}$
d. $15n^{10} \div 3n^{10}$

Answer

a. $18n^{10}$
b. $12n^{10}$
c. $45n^{20}$
d. 5

Exercise 6.5E.69

a. $p^4 \cdot p^6$
b. $\left(p^4\right)^6$

Exercise 6.5E.70

a. $q^5 \cdot q^3$
b. $\left(q^5\right)^3$

Answer

a. q^8
b. q^{15}

Exercise 6.5E.71

a. $\dfrac{y^3}{y}$
b. $\dfrac{y}{y^3}$

Exercise 6.5E.72

a. $\dfrac{z^6}{z^5}$
b. $\dfrac{z^5}{z^6}$

Answer

a. z
b. $\dfrac{1}{z}$

https://math.libretexts.org/@go/page/30542

Exercise 6.5E.73

$\left(8x^5\right)(9x) \div 6x^3$

Exercise 6.5E.74

$(4y)\left(12y^7\right) \div 8y^2$

Answer

$6y^6$

Exercise 6.5E.75

$\frac{27a^7}{3a^3} + \frac{54a^9}{9a^5}$

Exercise 6.5E.76

$\frac{32c^{11}}{4c^5} + \frac{42c^9}{6c^3}$

Answer

$15c^6$

Exercise 6.5E.77

$\frac{32y^5}{8y^2} - \frac{60y^{10}}{5y^7}$

Exercise 6.5E.78

$\frac{48x^6}{6x^4} - \frac{35x^9}{7x^7}$

Answer

$3x^2$

Exercise 6.5E.79

$\frac{63r^6s^3}{9r^4s^2} - \frac{72r^2s^2}{6s}$

Exercise 6.5E.80

$\frac{56y^4z^5}{7y^3z^3} - \frac{45y^2z^2}{5y}$

Answer

$-yz^2$

Everyday Math

Exercise 6.5E.81

Memory One megabyte is approximately 10^6 bytes. One gigabyte is approximately 10^9 bytes. How many megabytes are in one gigabyte?

Exercise 6.5E.82

Memory One gigabyte is approximately 10^9 bytes. One terabyte is approximately 10^12 bytes. How many gigabytes are in one terabyte?

Answer

10^3

Writing Exercises

Exercise 6.5E.83

Jennifer thinks the quotient $\frac{a^{24}}{a^6}$ simplifies to a^4. What is wrong with her reasoning?

Exercise 6.5E.84

Maurice simplifies the quotient $\frac{d^7}{d}$ by writing $\frac{\not{d}^7}{\not{d}} = 7$. What is wrong with his reasoning?

Answer

Answers will vary.

Exercise 6.5E.85

When Drake simplified -3^0 and $(-3)^0$ he got the same answer. Explain how using the Order of Operations correctly gives different answers.

Exercise 6.5E.86

Robert thinks x^0 simplifies to 0. What would you say to convince Robert he is wrong?

Answer

Answers will vary.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
simplify expressions using the Quotient Property for Exponents.			
simplify expressions with zero exponents.			
simplify expressions using the Quotient to a Power Property.			
simplify expressions by applying several properties.			
divide monomials.			

b. On a scale of 1-10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 6.5E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.5E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.6: Divide Polynomials

Learning Objectives

By the end of this section, you will be able to:

- Divide a polynomial by a monomial
- Divide a polynomial by a binomial

Note

Before you get started, take this readiness quiz.

1. Add: $\frac{3}{d}+\frac{x}{d}$
   If you missed this problem, review Example 1.7.1.
2. Simplify: $\frac{30xy^3}{5xy}$
   If you missed this problem, review Example 6.5.37.
3. Combine like terms: $8a^2+12a+1+3a^2-5a+4$
   If you missed this problem, review Example 1.3.37.

Divide a Polynomial by a Monomial

In the last section, you learned how to divide a monomial by a monomial. As you continue to build up your knowledge of polynomials the next procedure is to divide a polynomial of two or more terms by a **monomial**.

The method we'll use to divide a polynomial by a monomial is based on the properties of fraction addition. So we'll start with an example to review fraction addition.

The sum, $\frac{y}{5}+\frac{2}{5}$

simplifies to $\frac{y+2}{5}$

Now we will do this in reverse to split a single fraction into separate fractions.

We'll state the fraction addition property here just as you learned it and in reverse.

FRACTION ADDITION

If a,b, and c are numbers where $c\neq 0$, then

$$\frac{a}{c}+\frac{b}{c}=\frac{a+b}{c} \quad \text{and} \quad \frac{a+b}{c}=\frac{a}{c}+\frac{b}{c} \tag{6.6.1}$$

We use the form on the left to add fractions and we use the form on the right to divide a polynomial by a monomial.

We use this form of fraction addition to divide polynomials by monomials.

DIVISION OF A POLYNOMIAL BY A MONOMIAL

To divide a polynomial by a monomial, divide each term of the polynomial by the **monomial**.

Example 6.6.1

Find the quotient: $\frac{7y^2+21}{7}$

Solution

	$\frac{7y^2+21}{7}$
Divide each term of the numerator by the denominator.	$\frac{7y^2}{7}+\frac{21}{7}$
Simplify each fraction.	y^2+3

Try It 6.6.2

Find the quotient: $\frac{8z^2+24}{4}$

Answer

$2z^2+6$

Try It 6.6.3

Find the quotient: $\frac{18z^2-27}{9}$

Answer

$2z^2-3$

Remember that division can be represented as a fraction. When you are asked to divide a polynomial by a monomial and it is not already in fraction form, write a fraction with the polynomial in the numerator and the monomial in the denominator.

Example 6.6.4

Find the quotient: $\left(18x^3-36x^2\right)\div 6x$

Solution

	$\left(18x^3-36x^2\right)\div 6x$
Rewrite as a fraction.	$\frac{18x^3-36x^2}{6x}$
Divide each term of the numerator by the denominator.	$\frac{18x^3}{6x}-\frac{36x^2}{6x}$
Simplify.	$3x^2-6x$

Try It 6.6.5

Find the quotient: $\left(27b^3-33b^2\right)\div 3b$

Answer

$9b^2-11b$

Try It 6.6.6

Find the quotient: $\left(25y^3-55y^2\right)\div 5y$

Answer

$5y^2-11y$

When we divide by a negative, we must be extra careful with the signs.

Example 6.6.7

Find the quotient: $\frac{12d^2-16d}{-4}$

Solution

	$\frac{12d^2-16d}{-4}$
Divide each term of the numerator by the denominator.	$\frac{18x^3-36x^2}{6x}$
Simplify. Remember, subtracting a negative is like adding a positive!	$-3d^2+4d$

Try It 6.6.8

Find the quotient: $\frac{25y^2-15y}{-5}$

Answer

$-5y^2+3y$

Try It 6.6.9

Find the quotient: $\frac{42b^2-18b}{-6}$

Answer

$-7b^2+3b$

Example 6.6.10

Find the quotient: $\frac{105y^5+75y^3}{5y^2}$

Solution

	$\frac{105y^5+75y^3}{5y^2}$
Separate the terms.	$\frac{105y^5}{5y^2}+\frac{75y^3}{5y^2}$
Simplify.	$21y^3+15y$

Try It 6.6.11

Find the quotient: $\frac{60d^7+24d^5}{4d^3}$

Answer

$15d^4+6d^2$

Try It 6.6.12

Find the quotient: $\frac{216p^7-48p^5}{6p^3}$

Answer

$36p^4 - 8p^2$

Example 6.6.13

Find the quotient: $\left(15x^3y - 35xy^2\right) \div (-5xy)$

Solution

	$\left(15x^3y - 35xy^2\right) \div (-5xy)$
Rewrite as a fraction.	$\dfrac{15x^3y - 35xy^2}{-5xy}$
Separate the terms. Be careful with the signs!	$\dfrac{15x^3y}{-5xy} - \dfrac{35xy^2}{-5xy}$
Simplify.	$-3x^2 + 7y$

Try It 6.6.14

Find the quotient: $\left(32a^2b - 16ab^2\right) \div (-8ab)$

Answer

$-4a + 2b$

Try It 6.6.15

Find the quotient: $\left(-48a^8b^4 - 36a^6b^5\right) \div \left(-6a^3b^3\right)$

Answer

$8a^5b + 6a^3b^2$

Example 6.6.16

Find the quotient: $\dfrac{36x^3y^2 + 27x^2y^2 - 9x^2y^3}{9x^2y}$

Solution

	$\dfrac{36x^3y^2 + 27x^2y^2 - 9x^2y^3}{9x^2y}$
Separate the terms.	$\dfrac{36x^3y^2}{9x^2y} + \dfrac{27x^2y^2}{9x^2y} - \dfrac{9x^2y^3}{9x^2y}$
Simplify.	$4xy + 3y - y^2$

Try It 6.6.17

Find the quotient: $\dfrac{40x^3y^2 + 24x^2y^2 - 16x^2y^3}{8x^2y}$

Answer

$5xy + 3y - 2y^2$

Try It 6.6.18

Find the quotient: $\frac{35a^4b^2+14a^4b^3-42a^2b^4}{7a^2b^2}$

Answer

$5a^2+2a^2b-6b^2$

Example 6.6.19

Find the quotient: $\frac{10x^2+5x-20}{5x}$

Solution

	$\frac{10x^2+5x-20}{5x}$
Separate the terms.	$\frac{10x^2}{5x}+\frac{5x}{5x}-\frac{20}{5x}$
Simplify.	$2x+1-\frac{4}{x}$

Try It 6.6.20

Find the quotient: $\frac{18c^2+6c-9}{6c}$

Answer

$3c+1-\frac{3}{2c}$

Try It 6.6.21

Find the quotient: $\frac{10d^2-5d-2}{5d}$

Solution

$2d-1-\frac{2}{5d}$

Divide a Polynomial by a Binomial

To divide a polynomial by a **binomial**, we follow a procedure very similar to long division of numbers. So let's look carefully the steps we take when we divide a 3-digit number, 875, by a 2-digit number, 25.

We write the long division	$25\overline{)875}$
We divide the first two digits, 87, by 25.	$\begin{array}{r} 3 \\ 25\overline{)875} \end{array}$
We multiply 3 times 25 and write the product under the 87.	$\begin{array}{r} 3 \\ 25\overline{)875} \\ \underline{75} \end{array}$

Now we subtract 75 from 87.	$$\begin{array}{r} 3 \\ 25\overline{)875} \\ \underline{-75} \\ 12 \end{array}$$ $$\begin{array}{r} 3 \\ 25\longdiv{875} \\ \underline{-75} \\ 125 \end{array} \quad (6.6.2)$$
Then we bring down the third digit of the dividend, 5.	$$\begin{array}{r} 3 \\ 25\overline{)875} \\ \underline{-75} \\ 125 \end{array}$$
Repeat the process, dividing 25 into 125.	$$\begin{array}{r} 35 \\ 25\overline{)875} \\ \underline{-75} \\ 125 \\ \underline{-125} \end{array}$$

We check division by multiplying the quotient by the divisor.

If we did the division correctly, the product should equal the dividend.

$$\begin{array}{l} 35 \cdot 25 \\ 875\checkmark \end{array} \quad (6.6.3)$$

Now we will divide a **trinomial** by a binomial. As you read through the example, notice how similar the steps are to the numerical example above.

Example 6.6.22

Find the quotient: $\left(x^2+9x+20\right) \div (x+5)$

Solution

	$(x^2 + 9x + 20) \div (x + 5)$
Write it as a long division problem.	
Be sure the dividend is in standard form.	$x + 5 \overline{)\, x^2 + 9x + 20}$
Divide x^2 by x. It may help to ask yourself, "What do I need to multiply x by to get x^2?"	
Put the answer, x, in the quotient over the x term.	$\begin{array}{r} x \\ x + 5 \overline{)\, x^2 + 9x + 20} \end{array}$
Multiply x times $x + 5$. Line up the like terms under the dividend.	$\begin{array}{r} x \\ x + 5 \overline{)\, x^2 + 9x + 20} \\ \underline{x^2 + 5x} \end{array}$
Subtract $x^2 + 5x$ from $x^2 + 9x$.	
You may find it easier to change the signs and then add. Then bring down the last term, 20.	$\begin{array}{r} x \\ x + 5 \overline{)\, x^2 + 9x + 20} \\ \underline{-x^2 + (-5x)} \\ 4x + 20 \end{array}$
Divide $4x$ by x. It may help to ask yourself, "What do I need to multiply x by to get $4x$?"	

Put the answer, 4, in the quotient over the constant term.	$x+4$ $x+5 \overline{)\, x^2+9x+20}$ $-x^2+(-5x)$ $4x+20$
Multiply 4 times x + 5.	$x+4$ $x+5 \overline{)\, x^2+9x+20}$ $-x^2+(-5x)$ $4x+20$ $4x+20$
Subtract 4x + 20 from 4x + 20.	$x+4$ $x+5 \overline{)\, x^2+9x+20}$ $-x^2+(-5x)$ $4x+20$ $-4x+(-20)$ 0
Check:	
Multiply the quotient by the divisor.	
(x + 4)(x + 5)	
You should get the dividend.	
x^2 + 9x + 20✓	

Try It 6.6.23

Find the quotient: $\left(y^2+10y+21\right) \div (y+3)$

Answer

y+7

Try It 6.6.24

Find the quotient: $\left(m^2+9m+20\right) \div (m+4)$

Answer

m+5

When the divisor has subtraction sign, we must be extra careful when we multiply the partial quotient and then subtract. It may be safer to show that we change the signs and then add.

Example 6.6.25

Find the quotient: $\left(2x^2-5x-3\right) \div (x-3)$

Solution

	$(2x^2-5x-3) \div (x-3)$
Write it as a long division problem.	
Be sure the dividend is in standard form.	$x-3 \overline{)\, 2x^2-5x-3}$
Divide 2x^2 by x. Put the answer, 2x, in the quotient over the x term.	$2x$ $x-3 \overline{)\, 2x^2-5x-3}$

Multiply $2x$ times $x - 3$. Line up the like terms under the dividend.	$$\begin{array}{r} 2x \\ x-3\overline{)\,2x^2-5x-3} \\ \underline{2x^2-6x} \end{array}$$
Subtract $2x^2 - 6x$ from $2x^2 - 5x$. Change the signs and then add. Then bring down the last term.	$$\begin{array}{r} 2x \\ x-3\overline{)\,2x^2-5x-3} \\ \underline{-2x^2+6x} \\ x-3 \end{array}$$
Divide x by x. Put the answer, 1, in the quotient over the constant term.	$$\begin{array}{r} 2x+1 \\ x-3\overline{)\,2x^2-5x-3} \\ \underline{-2x^2-(-6x)} \\ x-3 \end{array}$$
Multiply 1 times $x - 3$.	$$\begin{array}{r} 2x+1 \\ x-3\overline{)\,2x^2-5x-3} \\ \underline{-2x^2+6x} \\ x-3 \\ \underline{x-3} \end{array}$$
Subtract $x - 3$ from $x - 3$ by changing the signs and adding.	$$\begin{array}{r} 2x+1 \\ x-3\overline{)\,2x^2-5x-3} \\ \underline{-2x^2+6x} \\ x-3 \\ \underline{-x+3} \\ 0 \end{array}$$
To check, multiply $(x - 3)(2x + 1)$.	
The result should be $2x^2 - 5x - 3$.	

? Try It 6.6.26

Find the quotient: $\left(2x^2-3x-20\right) \div (x-4)$

Answer

2x+5

? Try It 6.6.27

Find the quotient: $\left(3x^2-16x-12\right) \div (x-6)$

Answer

3x+2

When we divided 875 by 25, we had no remainder. But sometimes division of numbers does leave a remainder. The same is true when we divide polynomials. In Example 6.6.25, we'll have a division that leaves a remainder. We write the remainder as a fraction with the divisor as the denominator.

✔ Example 6.6.28

Find the quotient: $\left(x^3-x^2+x+4\right) \div (x+1)$

Solution

	$(x^3 - x^2 + x + 4) \div (x + 1)$
Write it as a long division problem.	

Be sure the dividend is in standard form.	$x+1\overline{)\,x^3-x^2+x+4}$
Divide x^3 by x. Put the answer, x^2, in the quotient over the x^2 term. Multiply x^2 times $x + 1$. Line up the like terms under the dividend.	$\begin{array}{r} x^2 \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{x^3+x^2} \end{array}$
Subtract $x^3 + x^2$ from $x^3 - x^2$ by changing the signs and adding. Then bring down the next term.	$\begin{array}{r} x^2 \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{-x^3+(-x^2)} \\ -2x^2+x \end{array}$
Divide $-2x^2$ by x. Put the answer, $-2x$, in the quotient over the x term. Multiply $-2x$ times $x + 1$. Line up the like terms under the dividend.	$\begin{array}{r} x^2-2x \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{-x^3+(-x^2)} \\ -2x^2+x \\ \underline{-2x^2-2x} \end{array}$
Subtract $-2x^2 - 2x$ from $-2x^2 + x$ by changing the signs and adding. Then bring down the last term.	$\begin{array}{r} x^2-2x \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{-x^3+(-x^2)} \\ -2x^2+x \\ \underline{+2x^2+2x} \\ 3x+4 \end{array}$
Divide $3x$ by x. Put the answer, 3, in the quotient over the constant term. Multiply 3 times $x + 1$. Line up the like terms under the dividend.	$\begin{array}{r} x^2-2x+3 \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{-x^3+(-x^2)} \\ -2x^2+x \\ \underline{+2x^2+2x} \\ 3x+4 \\ \underline{3x+3} \end{array}$
Subtract $3x + 3$ from $3x + 4$ by changing the signs and adding. Write the remainder as a fraction with the divisor as the denominator.	$\begin{array}{r} x^2-2x+3+\dfrac{1}{x+1} \\ x+1\overline{)\,x^3-x^2+x+4} \\ \underline{-x^3+(-x^2)} \\ -2x^2+x \\ \underline{+2x^2+2x} \\ 3x+4 \\ \underline{-3x+(-3)} \\ 1 \end{array}$
To check, multiply $(x+1)\left(x^2-2x+3+\dfrac{1}{x+1}\right)$ The result should be x^3-x^2+x+4	

? Try It 6.6.29

Find the quotient: $\left(x^3+5x^2+8x+6\right)\div(x+2)$

Answer

$$x^2+3x+2+\frac{2}{x+2}$$

Try It 6.6.30

Find the quotient: $\left(2x^3+8x^2+x-8\right)\div(x+1)$

Answer

$$2x^2+6x-5-\frac{3}{x+1}$$

Look back at the dividends in Example, Example, and Example. The terms were written in descending order of degrees, and there were no missing degrees. The dividend in Example will be x^4-x^2+5x-2 . It is missing an x^3 term. We will add in $0x^3$ as a placeholder.

Example 6.6.31

Find the quotient: $\left(x^4-x^2+5x-2\right)\div(x+2)$

Solution

Notice that there is no x^3 term in the dividend. We will add $0x^3$ as a placeholder.

	$(x^4 - x^2 + 5x - 2) \div (x + 2)$
Write it as a long division problem. Be sure the dividend is in standard form with placeholders for missing terms.	$x+2\overline{)\,x^4-0x^3-x^2+5x-2}$
Divide x^4 by x. Put the answer, x^3, in the quotient over the x^3 term. Multiply x^3 times $x + 2$. Line up the like terms. Subtract and then bring down the next term.	x^3 $x+2\overline{)\,x^4+0x^3-x^2+5x-2}$ $-(x^4+2x^3)$ $-2x^3-x^2$ It may be helpful to change the signs and add.
Divide $-2x^3$ by x. Put the answer, $-2x^2$, in the quotient over the x^2 term. Multiply $-2x^2$ times $x + 1$. Line up the like terms. Subtract and bring down the next term.	x^3-2x^2 $x+2\overline{)\,x^4+0x^3-x^2+5x-2}$ $-(x^4+2x^3)$ $-2x^3-x^2$ $-(-2x^3-4x^2)$ $3x^2+5x$ It may be helpful to change the signs and add.
Divide $3x^2$ by x. Put the answer, $3x$, in the quotient over the x term. Multiply $3x$ times $x + 1$. Line up the like terms. Subtract and bring down the next term.	x^3-2x^2+3x $x+2\overline{)\,x^4+0x^3-x^2+5x-2}$ $-(x^4+2x^3)$ $-2x^3-x^2$ $-(-2x^3-4x^2)$ $3x^2+5x$ $-(3x^2+6x)$ $-x-2$ It may be helpful to change the signs and add.
Divide $-x$ by x. Put the answer, -1, in the quotient over the constant term. Multiply -1 times $x + 1$. Line up the like terms. Change the signs, add.	x^3-2x^2+3x-1 $x+2\overline{)\,x^4+0x^3-x^2+5x-2}$ $-(x^4+2x^3)$ $-2x^3-x^2$ $-(-2x^3-4x^2)$ $3x^2+5x$ $-(3x^2+6x)$ $-x-2$ $-(-x-2)$ 0 It may be helpful to change the signs and add.
To check, multiply $(x+2)\left(x^3-2x^2+3x-1\right)$	
The result should be x^4-x^2+5x-2	

Try It 6.6.32

Find the quotient: $\left(x^3+3x+14\right) \div (x+2)$

Answer

x^2-2x+7

Try It 6.6.33

Find the quotient: $\left(x^4-3x^3-1000\right) \div (x+5)$

Answer

$x^3-8x^2+40x-200$


In Example 6.6.31, we will divide by $2a-3$. As we divide we will have to consider the constants as well as the variables.

Example 6.6.34

Find the quotient: $\left(8a^3+27\right) \div (2a+3)$

Solution

This time we will show the division all in one step. We need to add two placeholders in order to divide.



To check, multiply $(2a+3)\left(4a^2-6a+9\right)$

The result should be $8a^3+27$

Try It 6.6.35

Find the quotient: $\left(x^3-64\right) \div (x-4)$

Answer

$x^2+4x+16$

Try It 6.6.36

Find the quotient: $\left(125x^3-8\right) \div (5x-2)$

Answer

$25x^2+10x+4$

Note

Access these online resources for additional instruction and practice with dividing polynomials:

- Divide a Polynomial by a Monomial
- Divide a Polynomial by a Monomial 2
- Divide Polynomial by Binomial

Key Concepts

- **Fraction Addition**
  - If a,b, and c are numbers where $c \neq 0$, then $\frac{a}{c}+\frac{b}{c}=\frac{a+b}{c}$ and $\frac{a+b}{c}=\frac{a}{c}+\frac{b}{c}$
- **Division of a Polynomial by a Monomial**
  - To divide a polynomial by a monomial, divide each term of the polynomial by the monomial.

This page titled 6.6: Divide Polynomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.6: Divide Polynomials** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.6E: Exercises

Practice Makes Perfect

In the following exercises, divide each polynomial by the monomial.

Exercise 1

$\frac{45y+36}{9}$

Exercise 2

$\frac{30b+75}{5}$

Answer

$6b+15$

Exercise 3

$\frac{8d^2-4d}{2}$

Exercise 4

$\frac{42x^2-14x}{7}$

Answer

$6x^2-2x$

Exercise 5

$(16y^2-20y)\div 4y$

Exercise 6

$(55w^2-10w)\div 5w$

Answer

$11w-2$

Exercise 7

$(9n^4+6n^3)\div 3n$

Exercise 8

$(8x^3+6x^2)\div 2x$

Answer

$4x^2+3x$

Exercise 9

$$\frac{18y^2 - 12y}{-6}$$

Exercise 10

$$\frac{20b^2 - 12b}{-4}$$

Answer

$$-5b^2 + 3b$$

Exercise 11

$$\frac{35a^4 + 65a^2}{-5}$$

Exercise 12

$$\frac{51m^4 + 72m^3}{-3}$$

Answer

$$-17m^4 - 24m^3$$

Exercise 13

$$\frac{310y^4 - 200y^3}{5y^2}$$

Exercise 14

$$\frac{412z^8 - 48z^5}{4z^3}$$

Answer

$$103z^5 - 12z^2$$

Exercise 15

$$\frac{46x^3 + 38x^2}{2x^2}$$

Exercise 16

$$\frac{51y^4 + 42y^2}{3y^2}$$

Answer

$$17y^2 + 14$$

Exercise 17

$(24p^2 - 33p) \div (-3p)$

Exercise 18

$(35x^4 - 21x) \div (-7x)$

Answer

$-5x^3 + 3$

Exercise 19

$(63m^4 - 42m^3) \div (-7m^2)$

Exercise 20

$(48y^4 - 24y^3) \div (-8y^2)$

Answer

$-6y^2 + 3y$

Exercise 21

$(63a^2b^3 + 72ab^4) \div (9ab)$

Exercise 22

$(45x^3y^4 + 60xy^2) \div (5xy)$

Answer

$9x^2y^3 + 12y$

Exercise 23

$$\frac{52p^5q^4 + 36p^4q^3 - 64p^3q^2}{4p^2q}$$

Exercise 24

$$\frac{49c^2d^2 - 70c^3d^3 - 35c^2d}{7cd^2}$$

Answer

$$7c - 10c^2d - \frac{5c}{d}$$

Exercise 25

$$\frac{66x^3y^2 - 110x^2y^3 - 44x^4y^3}{11x^2y^2}$$

Exercise 26

$$\frac{72r^5s^2+132r^4s^3-96r^3s^5}{12r^2s^2}$$

Answer

$$6r^3+11r^2s-8rs^3$$

Exercise 27

$$\frac{4w^2+2w-5}{2w}$$

Exercise 28

$$\frac{12q^2+3q-1}{3q}$$

Answer

$$4q+1-\frac{1}{3q}$$

Exercise 29

$$\frac{10x^2+5x-4}{-5x}$$

Exercise 30

$$\frac{20y^2+12y-1}{-4y}$$

Answer

$$-5y-3+\frac{1}{4y}$$

Exercise 31

$$\frac{36p^3+18p^2-12p}{6p^2}$$

Exercise 32

$$\frac{63a^3-108a^2+99a}{9a^2}$$

Answer

$$7a-12+\frac{11}{a}$$

Divide a Polynomial by a Binomial

In the following exercises, divide each polynomial by the binomial.

Exercise 33

$(y^2+7y+12)\div(y+3)$

Exercise 34

$(d^2+8d+12)\div(d+2)$

Answer

$d+6$

Exercise 35

$(x^2-3x-10)\div(x+2)$

Exercise 36

$(a^2-2a-35)\div(a+5)$

Answer

$a-7$

Exercise 37

$(t^2-12t+36)\div(t-6)$

Exercise 38

$(x^2-14x+49)\div(x-7)$

Answer

$x-7$

Exercise 39

$(6m^2-19m-20)\div(m-4)$

Exercise 40

$(4x^2-17x-15)\div(x-5)$

Answer

$4x+3$

Exercise 41

$(q^2+2q+20)\div(q+6)$

Exercise 42

$(p^2+11p+16)\div(p+8)$

Answer

$$p+3-\frac{8}{p+8}$$

Exercise 43

$(y^2-3y-15)\div(y-8)$

Exercise 44

$(x^2+2x-30)\div(x-5)$

Answer

$$x+7+\frac{5}{x-5}$$

Exercise 45

$(3b^3+b^2+2)\div(b+1)$

Exercise 46

$(2n^3-10n+28)\div(n+3)$

Answer

$2n^2-6n+8+\frac{4}{n+3}$

Exercise 47

$(2y^3-6y-36)\div(y-3)$

Exercise 48

$(7q^3-5q-2)\div(q-1)$

Answer

$7q^2+7q+2$

Exercise 49

$(z^3+1)\div(z+1)$

Exercise 50

$(m^3+1000)\div(m+10)$

Answer

$m^2-10m+100$

Exercise 51

$(a^3 - 125) \div (a - 5)$

Exercise 52

$(x^3 - 216) \div (x - 6)$

Answer

$x^2 + 6x + 36$

Exercise 53

$(64x^3 - 27) \div (4x - 3)$

Exercise 54

$(125y^3 - 64) \div (5y - 4)$

Answer

$25y^2 + 20x + 16$

Everyday Math

Exercise 55

Average cost Pictures Plus produces digital albums. The company's average cost (in dollars) to make x albums is given by the expression $\frac{7x + 500}{x}$

1. Find the quotient by dividing the numerator by the denominator.
2. What will the average cost (in dollars) be to produce 20 albums?

Exercise 56

Handshakes At a company meeting, every employee shakes hands with every other employee. The number of handshakes is given by the expression $\frac{n^2 - n}{2}$ nn represents the number of employees. How many handshakes will there be if there are 10 employees at the meeting?

Answer

45

Writing Exercises

Exercise 57

James divides $48y + 6$ by 6 this way: $\frac{48y + 6}{6} = 48y$

Exercise 58

Divide $\frac{10x^2+x-12}{2x}$ and explain with words how you get each term of the quotient.

Answer

Answers will vary.

Self Check

ⓐ After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
divide a polynomial by a monomial.			
divide a polynomial by a binomial.			

ⓑ After reviewing this checklist, what will you do to become confident for all goals?

This page titled 6.6E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.6E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.7: Integer Exponents and Scientific Notation

Learning Objectives

By the end of this section, you will be able to:

- Use the definition of a negative exponent
- Simplify expressions with integer exponents
- Convert from decimal notation to scientific notation
- Convert scientific notation to decimal form
- Multiply and divide using scientific notation

Note

Before you get started, take this readiness quiz.

1. What is the place value of the 6 in the number 64891?
   If you missed this problem, review Example 1.2.1.
2. Name the decimal: 0.0012.
   If you missed this problem, review Example 1.8.1.
3. Subtract: 5−(−3).
   If you missed this problem, review Example 1.4.33.

Use the Definition of a Negative Exponent

We saw that the Quotient Property for Exponents introduced earlier in this chapter, has two forms depending on whether the exponent is larger in the numerator or the denominator.

QUOTIENT PROPERTY FOR EXPONENTS

If a is a real number, $a \neq 0$, and m and n are whole numbers, then

$$\frac{a^m}{a^n} = a^{m-n}, m > n \tag{6.7.1}$$

and

$$\frac{a^m}{a^n} = \frac{1}{a^{n-m}}, n > m \tag{6.7.2}$$

What if we just subtract exponents regardless of which is larger?

Let's consider $\frac{x^2}{x^5}$.

We subtract the exponent in the denominator from the exponent in the numerator.

$$\begin{gathered} \frac{x^2}{x^5} \\ x^{2-5} \\ x^{-3} \end{gathered} \tag{6.7.3}$$

We can also simplify $\frac{x^2}{x^5}$ by dividing out common factors:

$$\begin{gathered} \frac{x \cdot x}{x \cdot x \cdot x \cdot x \cdot x} \\ \frac{1}{x^3} \end{gathered}$$

This implies that $x^{-3} = \frac{1}{x^3}$ and it leads us to the definition of a *negative exponent*.

Definition: NEGATIVE EXPONENT

If n is an integer and $a \neq 0$, then $a^{-n} = \frac{1}{a^n}$

The **negative exponent** tells us we can re-write the expression by taking the reciprocal of the base and then changing the sign of the exponent.

Any expression that has negative exponents is not considered to be in simplest form. We will use the definition of a **negative exponent** and other properties of exponents to write the expression with only positive exponents.

For example, if after simplifying an expression we end up with the expression x^{-3}, we will take one more step and write $\frac{1}{x^3}$. The answer is considered to be in simplest form when it has only positive exponents.

Example 6.7.1

Simplify:

1. 4^{-2}
2. 10^{-3}

Solution

1.

	4^{-2}
Use the definition of a negative exponent, $a^{-n} = \frac{1}{a^n}$,	$\frac{1}{4^2}$
Simplify.	$\frac{1}{16}$

2.

	10^{-3}
Use the definition of a negative exponent, $a^{-n} = \frac{1}{a^n}$,	$\frac{1}{10^3}$
Simplify.	$\frac{1}{1000}$

Try It 6.7.2

Simplify:

1. 2^{-3}
2. 10^{-7}

Answer

1. $\frac{1}{8}$
2. $\frac{1}{10^7}$

Try It 6.7.3

Simplify:

1. 3^{-2}
2. 10^{-4}

Answer

1. $\frac{1}{9}$
2. $\frac{1}{10,000}$

In Example 6.7.1 we raised an integer to a negative exponent. What happens when we raise a fraction to a negative exponent? We'll start by looking at what happens to a fraction whose numerator is one and whose denominator is an integer raised to a negative exponent.

	$\frac{1}{a^{-n}}$
Use the definition of a negative exponent, $a^{-n}=\frac{1}{a^n}$	$\frac{1}{\frac{1}{a^n}}$
Simplify the complex fraction.	$1\cdot\frac{a^n}{1}$
Multiply.	a^n

This leads to the Property of Negative Exponents.

PROPERTY OF NEGATIVE EXPONENTS

If n is an integer and $a\neq 0$, then $\frac{1}{a^{-n}}=a^n$.

Example 6.7.4

Simplify:

1. $\frac{1}{y^{-4}}$
2. $\frac{1}{3^{-2}}$

Solution

1.

	$\frac{1}{y^{-4}}$
Use the property of a negative exponent, $\frac{1}{a^{-n}}=a^n$.	y^4

2.

	$\frac{1}{3^{-2}}$
Use the property of a negative exponent, $\frac{1}{a^{-n}}=a^n$.	3^2
Simplify.	9

Try It 6.7.5

Simplify:

1. $\frac{1}{p^{-8}}$
2. $\frac{1}{4^{-3}}$

Answer

1. p^8
2. 64

Try It 6.7.6

Simplify:

1. $\dfrac{1}{q^{-7}}$
2. $\dfrac{1}{2^{-4}}$

Answer

1. q^7
2. 16

Suppose now we have a fraction raised to a negative exponent. Let's use our definition of negative exponents to lead us to a new property.

	$\left(\frac{3}{4}\right)^{-2}$
Use the definition of a negative exponent, $a^{-n}=\frac{1}{a^n}$	$\frac{1}{\left(\frac{3}{4}\right)^{2}}$
Simplify the denominator.	$\frac{1}{\frac{9}{16}}$
Simplify the complex fraction.	$\frac{16}{9}$
But we know that $\frac{16}{9}$ is $\left(\frac{4}{3}\right)^{2}$	
This tells us that:	$\left(\frac{3}{4}\right)^{-2}=\left(\frac{4}{3}\right)^{2}$

To get from the original fraction raised to a negative exponent to the final result, we took the reciprocal of the base—the fraction—and changed the sign of the exponent.

This leads us to the *Quotient to a Negative Power Property*.

QUOTIENT TO A NEGATIVE EXPONENT PROPERTY

If a and b are real numbers, $a\neq 0, b\neq 0,$ and n is an integer, then $\left(\frac{a}{b}\right)^{-n}=\left(\frac{b}{a}\right)^{n}$

Example 6.7.7

Simplify:

1. $\left(\frac{5}{7}\right)^{-2}$
2. $\left(-\frac{2x}{y}\right)^{-3}$

Solution

1. $\left(\frac{5}{7}\right)^{-2}$

   Use the Quotient to a Negative Exponent Property, $\left(\frac{a}{b}\right)^{-n}=\left(\frac{b}{a}\right)^{n}$. Take the reciprocal of the fraction and change the sign of the exponent. . $\left(\frac{7}{5}\right)^{2}$

   Simplify. $\frac{49}{25}$

2. $\left(-\frac{2x}{y}\right)^{-3}$

   Use the Quotient to a Negative Exponent Property, $\left(\frac{a}{b}\right)^{-n}=\left(\frac{b}{a}\right)^{n}$. Take the reciprocal of the fraction and change the sign of the exponent. $\left(-\frac{y}{2x}\right)^{3}$

   Simplify. $-\frac{y^3}{8x^3}$

? Try It 6.7.8

Simplify:

1. $\left(\frac{2}{3}\right)^{-4}$
2. $\left(-\frac{6m}{n}\right)^{-2}$

Answer

1. $\frac{81}{16}$
2. $\frac{n^2}{36m^2}$

? Try It 6.7.9

Simplify:

1. $\left(\frac{3}{5}\right)^{-3}$
2. $\left(-\frac{a}{2b}\right)^{-4}$

Answer

1. $\frac{125}{27}$
2. $\frac{16b^4}{a^4}$

When simplifying an expression with exponents, we must be careful to correctly identify the base.

Example 6.7.10

Simplify:

1. $(-3)^{-2}$
2. -3^{-2}
3. $\left(-\frac{1}{3}\right)^{-2}$
4. $-\left(\frac{1}{3}\right)^{-2}$

Solution

1. Here the exponent applies to the base −3.

	$(-3)^{-2}$
Take the reciprocal of the base and change the sign of the exponent.	$\frac{1}{(-3)^2}$
Simplify.	$\frac{1}{9}$

2. The expression -3^{-2} means "find the opposite of 3^{-2}". Here the exponent applies to the base 3.

	-3^{-2}
Rewrite as a product with -1	$-1 \cdot 3^{-2}$
Take the reciprocal of the base and change the sign of the exponent.	$-1 \cdot \frac{1}{3^2}$
Simplify.	$-\frac{1}{9}$

3. Here the exponent applies to the base $\left(-\frac{1}{3}\right)$.

	$\left(-\frac{1}{3}\right)^{-2}$
Take the reciprocal of the base and change the sign of the exponent.	$\left(-\frac{3}{1}\right)^{2}$
Simplify.	9

4. The expression $-\left(\frac{1}{3}\right)^{-2}$ means "find the opposite of $\left(\frac{1}{3}\right)^{-2}$". Here the exponent applies to the base $\left(\frac{1}{3}\right)$.

	$-\left(\frac{1}{3}\right)^{-2}$
Rewrite as a product with -1	$-1 \cdot \left(\frac{1}{3}\right)^{-2}$
Take the reciprocal of the base and change the sign of the exponent.	$-1 \cdot \left(\frac{3}{1}\right)^{2}$
Simplify.	-9

Try It 6.7.11

Simplify:

1. $(-5)^{-2}$
2. -5^{-2}
3. $\left(-\frac{1}{5}\right)^{-2}$

4. $-\left(\frac{1}{5}\right)^{-2}$

Answer

1. $\frac{1}{25}$
2. $-\frac{1}{25}$
3. 25
4. -25

Try It 6.7.12

Simplify:

1. $(-7)^{-2}$
2. -7^{-2}
3. $\left(-\frac{1}{7}\right)^{-2}$
4. $-\left(\frac{1}{7}\right)^{-2}$

Answer

1. $\frac{1}{49}$
2. $-\frac{1}{49}$
3. 49
4. -49

We must be careful to follow the Order of Operations. In the next example, parts (a) and (b) look similar, but the results are different.

Example 6.7.13

Simplify:

1. $4 \cdot 2^{-1}$
2. $(4 \cdot 2)^{-1}$

Solution

1.

Do exponents before multiplication.	$4 \cdot 2^{-1}$
Use $a^{-n} = \frac{1}{a^n}$	$4 \cdot \frac{1}{2^1}$
Simplify.	2

2.

	$(4 \cdot 2)^{-1}$
Simplify inside the parentheses first.	$(8)^{-1}$
Use $a^{-n} = \frac{1}{a^n}$	$\frac{1}{8^1}$
Simplify.	$\frac{1}{8}$

Try It 6.7.14

Simplify:

1. $6 \cdot 3^{-1}$
2. $(6 \cdot 3)^{-1}$

Answer

1. 2
2. $\frac{1}{18}$

Try It 6.7.15

Simplify:

1. $8 \cdot 2^{-2}$
2. $(8 \cdot 2)^{-2}$

Answer

1. 2
2. $\frac{1}{256}$

When a variable is raised to a negative exponent, we apply the definition the same way we did with numbers. We will assume all variables are non-zero.

Example 6.7.16

Simplify:

1. x^{-6}
2. $\left(u^4\right)^{-3}$

Solution

1. x^{-6}

   Use the definition of a negative exponent, $a^{-n} = \frac{1}{a^n}$ $\frac{1}{x^6}$

2. $\left(u^4\right)^{-3}$

   Use the definition of a negative exponent, $a^{-n} = \frac{1}{a^n}$ $\frac{1}{\left(u^4\right)^3}$

   Simplify. $\frac{1}{u^{12}}$

Try It 6.7.17

Simplify:

1. y^{-7}
2. $\left(z^3\right)^{-5}$

Answer

1. $\frac{1}{y^7}$

2. $\dfrac{1}{z^{15}}$

Try It 6.7.18

Simplify:

1. p^{-9}
2. $\left(q^4\right)^{-6}$

Answer

1. $\dfrac{1}{p^9}$
2. $\dfrac{1}{q^{24}}$

When there is a product and an exponent we have to be careful to apply the exponent to the correct quantity. According to the Order of Operations, we simplify expressions in parentheses before applying exponents. We'll see how this works in the next example.

Example 6.7.19

Simplify:

1. $5y^{-1}$
2. $(5y)^{-1}$
3. $(-5y)^{-1}$

Solution

1. Notice the exponent applies to just the base y.

	$5y^{-1}$
Take the reciprocal of the base y, and change the sign of the exponent.	$5 \cdot \dfrac{1}{y^1}$
Simplify.	$\dfrac{5}{y}$

2. Here the parentheses make the exponent apply to the base $5y$

	$(5y)^{-1}$
Take the reciprocal of the base $5y$, and change the sign of the exponent.	$\dfrac{1}{(5y)^1}$
Simplify.	$\dfrac{1}{5y}$

3. The base here is $-5y$

	$(-5y)^{-1}$
Take the reciprocal of the base $-5y$ and change the sign of the exponent.	$\dfrac{1}{(-5y)^1}$
Simplify.	$\dfrac{1}{-5y}$
Use $\dfrac{a}{-b} = -\dfrac{a}{b}$	$-\dfrac{1}{5y}$

Try It 6.7.20

Simplify:

1. $8p^{-1}$
2. $(8p)^{-1}$
3. $(-8p)^{-1}$

Answer

1. $\frac{8}{p}$
2. $\frac{1}{8p}$
3. $-\frac{1}{8p}$

Try It 6.7.21

Simplify:

1. $11q^{-1}$
2. $(11q)^{-1}$
3. $(-11q)^{-1}$

Answer

1. $\frac{11}{q}$
2. $\frac{1}{11q}$
3. $-\frac{1}{11q}$

With negative exponents, the Quotient Rule needs only one form $\frac{a^m}{a^n} = a^{m-n}$, for $a \neq 0$. When the exponent in the denominator is larger than the exponent in the numerator, the exponent of the quotient will be negative.

Simplify Expressions with Integer Exponents

All of the exponent properties we developed earlier in the chapter with whole number exponents apply to integer exponents, too. We restate them here for reference.

SUMMARY OF EXPONENT PROPERTIES

If a and b are real numbers, and m and n are integers, then

Product Property	$a^m \cdot a^n$	$=$	a^{m+n}
Power Property	$(a^m)^n$	$=$	$a^{m \cdot n}$
Product to a Power	$(ab)^m$	$=$	$a^m b^m$
Quotient Property	$\frac{a^m}{a^n}$	$=$	$a^{m-n}, \quad a \neq 0$
Zero Exponent Property	a^0	$=$	$1, \quad a \neq 0$
Quotient to a Power Property	$\left(\frac{a}{b}\right)^m$	$=$	$\frac{a^m}{b^m}, \quad b \neq 0$
Properties of Negative Exponents	a^{-n}	$=$	$\frac{1}{a^n}$ and $\frac{1}{a^{-n}} = a^n$
Quotient to a Negative Exponents	$\left(\frac{a}{b}\right)^{-n}$	$=$	$\left(\frac{b}{a}\right)^n$

Example 6.7.22

Simplify:

1. $x^{-4} \cdot x^6$
2. $y^{-6} \cdot y^4$
3. $z^{-5} \cdot z^{-3}$

Solution

1.

	$x^{-4} \cdot x^6$
Use the Product Property, $a^m \cdot a^n = a^{m+n}$	x^{-4+6}
Simplify.	x^2

2.

	$y^{-6} \cdot y^4$
Notice the same bases, so add the exponents.	y^{-6+4}
Simplify.	y^{-2}
Use the definition of a negative exponent, $a^{-n} = \frac{1}{a^n}$	$\frac{1}{y^2}$

3.

	$z^{-5} \cdot z^{-3}$
Add the exponents, since the bases are the same.	z^{-5-3}
Simplify.	z^{-8}
Use the definition of a negative exponent to take the reciprocal of the base and change the sign of the exponent.	$\frac{1}{z^8}$

Try It 6.7.23

Simplify:

1. $x^{-3} \cdot x^7$
2. $y^{-7} \cdot y^2$
3. $z^{-4} \cdot z^{-5}$

Answer

1. x^4
2. $\frac{1}{y^5}$

3. $\frac{1}{z^9}$

Try It 6.7.24

Simplify:

1. $a^{-1} \cdot a^6$
2. $b^{-8} \cdot b^4$
3. $c^{-8} \cdot c^{-7}$

Answer

1. a^5
2. $\frac{1}{b^4}$
3. $\frac{1}{c^{15}}$

In the next two examples, we'll start by using the Commutative Property to group the same variables together. This makes it easier to identify the like bases before using the Product Property.

Example 6.7.25

Simplify: $\left(m^4 n^{-3}\right)\left(m^{-5} n^{-2}\right)$

Solution

	$\left(m^4 n^{-3}\right)\left(m^{-5} n^{-2}\right)$
Use the Commutative Property to get like bases together.	$m^4 m^{-5} \cdot n^{-2} n^{-3}$
Add the exponents for each base.	$m^{-1} \cdot n^{-5}$
Take reciprocals and change the signs of the exponents.	$\frac{1}{m^1} \cdot \frac{1}{n^5}$
Simplify.	$\frac{1}{mn^5}$

Try It 6.7.26

Simplify: $\left(p^6 q^{-2}\right)\left(p^{-9} q^{-1}\right)$

Answer

$\frac{1}{p^3 q^3}$

Try It 6.7.27

Simplify: $\left(r^5 s^{-3}\right)\left(r^{-7} s^{-5}\right)$

Answer

$\frac{1}{r^2 s^8}$

If the monomials have numerical coefficients, we multiply the coefficients, just like we did earlier.

Example 6.7.28

Simplify: $\left(2x^{-6}y^{8}\right)\left(-5x^{5}y^{-3}\right)$

Solution

	$\left(2x^{-6}y^{8}\right)\left(-5x^{5}y^{-3}\right)$
Rewrite with the like bases together.	$2(-5)\cdot\left(x^{-6}x^{5}\right)\cdot\left(y^{8}y^{-3}\right)$
Multiply the coefficients and add the exponents of each variable.	$-10\cdot x^{-1}\cdot y^{5}$
Use the definition of a negative exponent, $a^{-n}=\frac{1}{a^{n}}$	$-10\cdot\frac{1}{x^{1}}\cdot y^{5}$
Simplify.	$\frac{-10y^{5}}{x}$

Try It 6.7.29

Simplify: $\left(3u^{-5}v^{7}\right)\left(-4u^{4}v^{-2}\right)$

Answer

$-\frac{12v^{5}}{u}$

Try It 6.7.30

Simplify: $\left(-6c^{-6}d^{4}\right)\left(-5c^{-2}d^{-1}\right)$

Answer

$\frac{30d^{3}}{c^{8}}$

In the next two examples, we'll use the Power Property and the Product to a Power Property.

Example 6.7.31

Simplify: $\left(6k^{3}\right)^{-2}$

Solution

	$\left(6k^{3}\right)^{-2}$
Use the Product to a Power Property, $(ab)^{m}=a^{n}b^{m}$	$(6)^{-2}\left(k^{3}\right)^{-2}$
Use the Power Property, $(a^{m})^{n}=a^{m\cdot n}$	$6^{-2}k^{-6}$
Use the definition of a negative exponent, $a^{-n}=\frac{1}{a^{n}}$	$\frac{1}{6^{2}}\cdot\frac{1}{k^{6}}$
Simplify.	$\frac{1}{36k^{6}}$

Try It 6.7.32

Simplify: $\left(-4x^{4}\right)^{-2}$

Answer

$\frac{1}{16x^{8}}$

Try It 6.7.33

Simplify: $\left(2b^3\right)^{-4}$

Answer

$$\frac{1}{16b^{12}}$$

Example 6.7.34

Simplify: $\left(5x^{-3}\right)^2$

Solution

	$\left(5x^{-3}\right)^2$
Use the Product to a Power Property, $(ab)^m = a^n b^m$	$5^2\left(x^{-3}\right)^2$
Simplify 5^2 and multiply the exponents of x using the Power Property, $(a^m)^n = a^{m \cdot n}$.	$25 \cdot x^{-6}$
Rewrite x^{-6} by using the Definition of a Negative Exponent, $a^{-n} = \frac{1}{a^n}$	$25 \cdot \frac{1}{x^6}$
Simplify.	$\frac{25}{x^6}$

Try It 6.7.35

Simplify: $\left(8a^{-4}\right)^2$

Answer

$$\frac{64}{a^8}$$

Try It 6.7.36

Simplify: $\left(2c^{-4}\right)^3$

Answer

$$\frac{8}{c^{12}}$$

To simplify a fraction, we use the Quotient Property and subtract the exponents.

Example 6.7.37

Simplify: $\frac{r^5}{r^{-4}}$

Solution

	$\frac{r^5}{r^{-4}}$
Use the Quotient Property, $\frac{a^n}{a^n} = a^{m-n}$	$r^{5-(-4)}$
Simplify.	r^9

Try It 6.7.38

Simplify: $\frac{x^8}{x^{-3}}$

Answer

x^{11}

Try It 6.7.39

Simplify: $\frac{y^8}{y^{-6}}$

Answer

y^{14}

Convert from Decimal Notation to Scientific Notation

Remember working with place value for whole numbers and decimals? Our number system is based on powers of 10. We use tens, hundreds, thousands, and so on. Our decimal numbers are also based on powers of tens—tenths, hundredths, thousandths, and so on. Consider the numbers 4,000 and 0.004. We know that 4,000 means $4 \times 1,000$ and 0.004 means $4 \times \frac{1}{1,000}$.

If we write the 1000 as a power of ten in exponential form, we can rewrite these numbers in this way:

$$\begin{array}{ll} 4,000 & 0.004 \\ 4 \times 1,000 & 4 \times \frac{1}{1,000} \\ 4 \times 10^3 & 4 \times \frac{1}{10^3} \\ & 4 \times 10^{-3} \end{array} \tag{6.7.4}$$

When a number is written as a product of two numbers, where the first factor is a number greater than or equal to one but less than 10, and the second factor is a power of 10 written in exponential form, it is said to be in *scientific notation.*

SCIENTIFIC NOTATION

A number is expressed in **scientific notation** when it is of the form

$$a \times 10^n \text{ where } 1 \leq a < 10 \text{ and } n \text{ is an integer} \tag{6.7.5}$$

It is customary in scientific notation to use as the $\times$ multiplication sign, even though we avoid using this sign elsewhere in algebra.

If we look at what happened to the decimal point, we can see a method to easily convert from decimal notation to scientific notation.

$4000. = 4 \times 10^3$	$0.004 = 4 \times 10^{-3}$
$4000. = 4 \times 10^3$	$0.004 = 4 \times 10^{-3}$
Moved the decimal point 3 places to the left.	Moved the decimal point 3 places to the right.

In both cases, the decimal was moved 3 places to get the first factor between 1 and 10.

The power of 10 is positive when the number is larger than 1 : $4,000 = 4 \times 10^3$

The power of 10 is negative when the number is between 0 and 1 : $0.004 = 4 \times 10^{-3}$

Example 6.7.40: HOW TO CONVERT FROM DECIMAL NOTATION TO SCIENTIFIC NOTATION

Write in scientific notation: 37000.

Solution

Step 1. Move the decimal point so that the first factor is greater than or equal to 1 but less than 10.	Remember, there is a decimal at the end of 37,000. Move the decimal after the 3. 3.700 is between 1 and 10.	37,000.
Step 2. Count the number of decimal places, *n*, that the decimal point was moved.	The decimal point was moved 4 places to the left.	37000.
Step 3. Write the number as a product with a power of 10. If the original number is: Greater than 1, the power of 10 will be 10^n. Between 0 and 1, the power of 10 will be 10^{-n}.	37,000 is greater than 1 so the power of 10 will have exponent 4.	3.7×10^4
Step 4. Check.	Check to see if your answer makes sense.	10^4 is 10,000 and 10,000 times 3.7 will be 37,000. $37{,}000 = 3.7 \times 10^4$

Try It 6.7.41

Write in scientific notation: 96000.

Answer

9.6×10^4

Try It 6.7.42

Write in scientific notation: 48300.

Answer

4.83×10^4

HOW TO: Convert from decimal notation to scientific notation

1. Step 1. Move the decimal point so that the first factor is greater than or equal to 1 but less than 10.
2. Step 2. Count the number of decimal places, *n*, that the decimal point was moved.
3. Step 3. Write the number as a product with a power of 10.
   If the original number is:
   - greater than 1, the power of 10 will be 10^n.
   - between 0 and 1, the power of 10 will be 10^{-n}.
4. Step 4. Check.

https://math.libretexts.org/@go/page/30545

Example 6.7.43

Write in scientific notation: 0.0052.

Solution

The original number, 0.0052, is between 0 and 1 so we will have a negative power of 10.

	0.0052
Move the decimal point to get 5.2, a number between 1 and 10.	0.0052
Count the number of decimal places the point was moved.	3 places
Write as a product with a power of 10.	5.2×10^{-3}
Check.	
5.2×10^{-3} $5.2 \times \frac{1}{10^3}$ $5.2 \times \frac{1}{1000}$ 5.2×0.001	
0.0052	$0.0052 = 5.2 \times 10^{-3}$

Try It 6.7.44

Write in scientific notation: 0.0078

Answer

7.8×10^{-3}

Try It 6.7.45

Write in scientific notation: 0.0129

Answer

1.29×10^{-2}

Convert Scientific Notation to Decimal Form

How can we convert from scientific notation to decimal form? Let's look at two numbers written in scientific notation and see.

$$\begin{array}{cc} 9.12 \times 10^{4} & 9.12 \times 10^{-4} \\ 9.12 \times 10,000 & 9.12 \times 0.0001 \\ 91,200 & 0.000912 \end{array} \tag{6.7.6}$$

If we look at the location of the decimal point, we can see an easy method to convert a number from scientific notation to decimal form.

$$9.12 \times 10^{4} = 91,200 \quad 9.12 \times 10^{-4} = 0.000912 \tag{6.7.7}$$

$9.12 \times 10^4 = 91{,}200$	$9.12 \times 10^{-4} = 0.000912$
$9.12\_\_\_\times 10^4 = 91{,}200$	$\_\_\_9.12 \times 10^{-4} = 0.000912$
Move the decimal point 4 places to the right.	Move the decimal point 4 places to the left.

In both cases the decimal point moved 4 places. When the exponent was positive, the decimal moved to the right. When the exponent was negative, the decimal point moved to the left.

Example 6.7.46

Convert to decimal form: 6.2×10^3

Solution

Step 1. Determine the exponent, *n*, on the factor 10.	The exponent is 3.	6.2×10^3
Step 2. Move the decimal *n* places, adding zeros if needed. If the exponent is positive, move the decimal point *n* places to the right. If the exponent is negative, move the decimal point $\lvert n \rvert$ places to the left.	The exponent is positive, so move the decimal point 3 places to the right. We need to add 2 zeros as placeholders.	6.200. 6,200
Step 3. Check to see if your answer makes sense.		10^3 is 1000 and 1000 times 6.2 will be 6,200. $6.2 \times 10^3 = 6{,}200$

Try It 6.7.47

Convert to decimal form: 1.3×10^3

Answer

$1,300$

Try It 6.7.48

Convert to decimal form: 9.25×10^4

Answer

$92,500$

The steps are summarized below.

HOW TO

Convert scientific notation to decimal form.

To convert scientific notation to decimal form:

1. Step 1. Determine the exponent, n, on the factor 10.
2. Step 2. Move the decimal n places, adding zeros if needed.

- If the exponent is positive, move the decimal point n places to the right.
- If the exponent is negative, move the decimal point $|n|$ places to the left.

3. Step 3. Check.

Example 6.7.49

Convert to decimal form: 8.9×10^{-2}

Solution

	8.9×10^{-2}
Determine the exponent, n, on the factor 10.	The exponent is –2.
Since the exponent is negative, move the decimal point 2 places to the left.	8.9
Add zeros as needed for placeholders.	$8.9 \times 10^{-2} = 0.089$

Try It 6.7.50

Convert to decimal form: 1.2×10^{-4}

Answer

0.00012

Try It 6.7.51

Convert to decimal form: 7.5×10^{-2}

Answer

0.075

Multiply and Divide Using Scientific Notation

Astronomers use very large numbers to describe distances in the universe and ages of stars and planets. Chemists use very small numbers to describe the size of an atom or the charge on an electron. When scientists perform calculations with very large or very small numbers, they use scientific notation. Scientific notation provides a way for the calculations to be done without writing a lot of zeros. We will see how the Properties of Exponents are used to multiply and divide numbers in scientific notation.

Example 6.7.52

Multiply. Write answers in decimal form: $\left(4 \times 10^{5}\right)\left(2 \times 10^{-7}\right)$

Solution

	$\left(4 \times 10^{5}\right)\left(2 \times 10^{-7}\right)$
Use the Commutative Property to rearrange the factors.	$4 \cdot 2 \cdot 10^{5} \cdot 10^{-7}$
Multiply.	8×10^{-2}
Change to decimal form by moving the decimal two places left.	0.08

Try It 6.7.53

Multiply $(3 \times 10^{6})(2 \times 10^{-8})$. Write answers in decimal form.

Answer

0.06

Try It 6.7.54

Multiply $\left(3 \times 10^{-2}\right)\left(3 \times 10^{-1}\right)$. Write answers in decimal form.

Answer

0.009

Example 6.7.55

Divide. Write answers in decimal form: $\dfrac{9 \times 10^{3}}{3 \times 10^{-2}}$

Solution

	$\dfrac{9 \times 10^{3}}{3 \times 10^{-2}}$
Separate the factors, rewriting as the product of two fractions.	$\dfrac{9}{3} \times \dfrac{10^{3}}{10^{-2}}$
Divide.	3×10^{5}
Change to decimal form by moving the decimal five places right.	300000

Try It 6.7.56

Divide $\dfrac{8 \times 10^{4}}{2 \times 10^{-1}}$. Write answers in decimal form.

Answer

$400,000$

Try It 6.7.57

Divide $\dfrac{8 \times 10^{2}}{4 \times 10^{-2}}$. Write answers in decimal form.

Answer

$20,000$

MEDIA ACCESS ADDITIONAL ONLINE RESOURCES

Access these online resources for additional instruction and practice with integer exponents and scientific notation:

- Negative Exponents
- Scientific Notation
- Scientific Notation 2

Key Concepts

- **Property of Negative Exponents**
  - If n is a positive integer and $a \neq 0$, then $\frac{1}{a^{-n}} = a^n$
- **Quotient to a Negative Exponent**
  - If a and b are real numbers, $b \neq 0$ and n is an integer , then $\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n$
- **To convert scientific notation to decimal form:**
  1. Determine the exponent, n on the factor 10.
  2. Move the decimal n places, adding zeros if needed.
     - If the exponent is positive, move the decimal point n places to the right.
     - If the exponent is negative, move the decimal point $|n|$ places to the left.
  3. Check.
- **To convert a decimal to scientific notation:**
  1. Move the decimal point so that the first factor is greater than or equal to 1 but less than 10.
  2. Count the number of decimal places, n that the decimal point was moved.
  3. Write the number as a product with a power of 10. If the original number is:
     - greater than 1, the power of 10 will be 10^n
     - between 0 and 1, the power of 10 will be 10^{-n}
  4. Check.

Glossary

negative exponent

If n is a positive integer and $a \neq 0$, then $a^{-n} = \frac{1}{a^n}$.

scientific notation

A number is expressed in scientific notation when it is of the form $a \times 10^n$ where $a \geq 1$ and a<10 and n is an integer.

This page titled 6.7: Integer Exponents and Scientific Notation is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.7: Integer Exponents and Scientific Notation** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

6.7E: Exercises

Practice Makes Perfect

Use the Definition of a Negative Exponent

In the following exercises, simplify.

Exercise 1

1. 4^{-2}
2. 10^{-3}

Exercise 2

1. 3^{-4}
2. 10^{-2}

Answer

1. $\frac{1}{81}$
2. $\frac{1}{100}$

Exercise 3

1. 5^{-3}
2. 10^{-5}

Exercise 4

1. 2^{-8}
2. 10^{-2}

Answer

1. $\frac{1}{256}$
2. $\frac{1}{100}$

Exercise 5

1. $\frac{1}{c^{-5}}$
2. $\frac{1}{3^{-2}}$

Exercise 6

1. $\frac{1}{c^{-5}}$
2. $\frac{1}{5^{-2}}$

Answer

1. c^5
2. 25

Exercise 7

1. $\frac{1}{q^{-10}}$
2. $\frac{1}{10^{-3}}$

Exercise 8

1. $\frac{1}{t^{-9}}$
2. $\frac{1}{10^{-4}}$

Answer

1. t^9
2. 10000

Exercise 9

1. $\left(\frac{5}{8}\right)^{-2}$
2. $\left(-\frac{3m}{n}\right)^{-2}$

Exercise 10

1. $\left(\frac{3}{10}\right)^{-2}$
2. $\left(-\frac{2c}{d}\right)^{-3}$

Answer

1. $\frac{100}{9}$
2. $-\frac{d^3}{8c^3}$

Exercise 11

1. $\left(\frac{4}{9}\right)^{-3}$
2. $\left(-\frac{u^2}{2v}\right)^{-5}$

Exercise 12

1. $\left(\frac{7}{2}\right)^{-3}$
2. $\left(-\frac{3}{xy^2}\right)^{-3}$

Answer

1. $\frac{8}{343}$
2. $-\frac{x^3y^6}{27}$

Exercise 13

1. $(-5)^{-2}$
2. -5^{-2}
3. $\left(-\frac{1}{5}\right)^{-2}$
4. $-\left(\frac{1}{5}\right)^{-2}$

Exercise 14

1. $(-7)^{-2}$
2. -7^{-2}
3. $\left(-\frac{1}{7}\right)^{-2}$
4. $-\left(\frac{1}{7}\right)^{-2}$

Answer

1. $\frac{1}{49}$
2. $-\frac{1}{49}$
3. 49
4. −49

Exercise 15

1. -3^{-3}
2. $\left(-\frac{1}{3}\right)^{-3}$
3. $-\left(\frac{1}{3}\right)^{-3}$
4. $(-3)^{-3}$

Exercise 16

1. -5^{-3}
2. $\left(-\frac{1}{5}\right)^{-3}$
3. $-\left(\frac{1}{5}\right)^{-3}$
4. $(-5)^{-3}$

Answer

1. $-\frac{1}{125}$
2. −125
3. −125
4. $-\frac{1}{125}$

Exercise 17

1. $3 \cdot 5^{-1}$
2. $(3 \cdot 5)^{-1}$

Exercise 18

1. $2 \cdot 5^{-1}$
2. $(2 \cdot 5)^{-1}$

Answer

1. $\frac{2}{5}$
2. $\frac{1}{10}$

Exercise 19

1. $4 \cdot 5^{-2}$
2. $(4 \cdot 5)^{-2}$

Exercise 20

1. $3 \cdot 4^{-2}$
2. $(3 \cdot 4)^{-2}$

Answer

1. $\frac{3}{16}$
2. $\frac{1}{144}$

Exercise 21

1. m^{-4}
2. $(x^3)^{-4}$

Exercise 22

1. b^{-5}
2. $(k^2)^{-5}$

Answer

1. $\frac{1}{b^5}$
2. $\frac{1}{k^{10}}$

Exercise 23

1. p^{-10}
2. $(q^6)^{-8}$

Exercise 24

1. s^{-8}
2. $(a^9)^{-10}$

Answer

1. $\frac{1}{s^8}$
2. $\frac{1}{a^{90}}$

Exercise 25

1. $7n^{-1}$
2. $(7n)^{-1}$
3. $(-7n)^{-1}$

Exercise 26

1. $6r^{-1}$
2. $(6r)^{-1}$
3. $(-6r)^{-1}$

Answer

1. $\frac{6}{r}$
2. $\frac{1}{6r}$
3. $-\frac{1}{6r}$

Exercise 27

1. $(3p)^{-2}$
2. $3p^{-2}$
3. $-3p^{-2}$

Exercise 28

1. $(2q)^{-4}$
2. $2q^{-4}$
3. $-2q^{-4}$

Answer

1. $\frac{1}{16q^4}$
2. $\frac{2}{q^4}$
3. $-\frac{2}{q^4}$

Simplify Expressions with Integer Exponents

In the following exercises, simplify.

Exercise 29

1. b^4b^{-8}
2. $r^{-2}r^5$
3. $x^{-7}x^{-3}$

Exercise 30

1. $s^3 \cdot s^{-7}$
2. $q^{-8} \cdot q^3$
3. $y^{-2} \cdot y^{-5}$

Answer

1. $\frac{1}{s^4}$
2. $\frac{1}{q^5}$
3. $\frac{1}{y^7}$

Exercise 31

1. $a^3 \cdot a^{-3}$
2. $a \cdot a^3$
3. $a \cdot a^{-3}$

Exercise 32

1. $y^5 \cdot y^{-5}$
2. $y \cdot y^5$
3. $y \cdot y^{-5}$

Answer

1. 1
2. y^6
3. $\frac{1}{y^4}$

Exercise 33

$p^5 \cdot p^{-2} \cdot p^{-4}$

Exercise 34

$x^4 \cdot x^{-2} \cdot x^{-3}$

Answer

$\frac{1}{x}$

Exercise 35

$(w^4x^{-5})(w^{-2}x^{-4})$

Exercise 36

$(m^3n^{-3})(m^{-5}n^{-1})$

Answer

$\frac{1}{m^2n^4}$

Exercise 37

$(uv^{-2})(u^{-5}v^{-3})$

Exercise 38

$(pq^{-4})(p^{-6}q^{-3})$

Answer

$$\frac{1}{p^5q^7}$$

Exercise 39

$(-6c^{-3}d^9)(2c^4d^{-5})$

Exercise 40

$(-2j^{-5}k^8)(7j^2k^{-3})$

Answer

$$-\frac{14k^5}{j^3}$$

Exercise 41

$(-4r^{-2}s^{-8})(9r^4s^3)$

Exercise 42

$(-5m^4n^6)(8m^{-5}n^{-3})$

Answer

$$-\frac{40n^3}{m}$$

Exercise 43

$(5x^2)^{-2}$

Exercise 44

$(4y^3)^{-3}$

Answer

$$\frac{1}{64y^9}$$

https://math.libretexts.org/@go/page/30546

Exercise 45

$(3z^{-3})^2$

Exercise 46

$(2p^{-5})^2$

Answer

$$\frac{4}{p^{10}}$$

Exercise 47

$$\frac{t^9}{t^{-3}}$$

Exercise 48

$$\frac{n^5}{n^{-2}}$$

Answer

n^7

Exercise 49

$$\frac{x^{-7}}{x^{-3}}$$

Exercise 50

$$\frac{y^{-5}}{y^{-10}}$$

Answer

y^5

Convert from Decimal Notation to Scientific Notation

In the following exercises, write each number in scientific notation.

Exercise 51

57,000

Exercise 52

340,000

Answer

3.4×10^5

Exercise 53

8,750,000

Exercise 54

1,290,000

Answer

1.29×10^6

Exercise 55

0.026

Exercise 56

0.041

Answer

4.1×10^{-2}

Exercise 57

0.00000871

Exercise 58

0.00000103

Answer

1.03×10^{-6}

Convert Scientific Notation to Decimal Form

In the following exercises, convert each number to decimal form.

Exercise 59

5.2×10^2

Exercise 60

8.3×10^2

Answer

830

Exercise 61

7.5×10^6

Exercise 62

1.6×10^{10}

Answer

16,000,000,000

Exercise 63

2.5×10^{-2}

Exercise 64

3.8×10^{-2}

Answer

0.038

Exercise 65

4.13×10^{-5}

Exercise 66

1.93×10^{-5}

Answer

0.0000193

Multiply and Divide Using Scientific Notation

In the following exercises, multiply. Write your answer in decimal form.

Exercise 67

$\left(3 \times 10^{-5}\right)\left(3 \times 10^{9}\right)$

Exercise 68

$\left(2 \times 10^{2}\right)\left(1 \times 10^{-4}\right)$

Answer

0.02

Exercise 69

$\left(7.1 \times 10^{-2}\right)\left(2.4 \times 10^{-4}\right)$

Exercise 70

$\left(3.5 \times 10^{-4}\right)\left(1.6 \times 10^{-2}\right)$

Answer

5.6×10^{-6}

In the following exercises, divide. Write your answer in decimal form.

Exercise 71

$$\frac{7 \times 10^{-3}}{1 \times 10^{-7}}$$

Exercise 72

$$\frac{5 \times 10^{-2}}{1 \times 10^{-10}}$$

Answer

500,000,000

Exercise 73

$$\frac{6 \times 10^{4}}{3 \times 10^{-2}}$$

Exercise 74

$$\frac{8 \times 10^{6}}{4 \times 10^{-1}}$$

Answer

20,000,000

Everyday Math

Exercise 75

The population of the United States on July 4, 2010 was almost 310,000,000. Write the number in scientific notation.

Exercise 76

The population of the world on July 4, 2010 was more than 6,850,000,000. Write the number in scientific notation

Answer

6.85×10^{9}

Exercise 77

The average width of a human hair is 0.0018 centimeters. Write the number in scientific notation.

Exercise 78

The probability of winning the 2010 Megamillions lottery was about 0.0000000057. Write the number in scientific notation.

Answer

5.7×10^{-9}

Exercise 79

In 2010, the number of Facebook users each day who changed their status to 'engaged" was 2×10^4. Convert this number to decimal form.

Exercise 80

At the start of 2012, the US federal budget had a deficit of more than $\$1.5 \times 10^{13}$. Convert this number to decimal form.

Answer

15,000,000,000,000

Exercise 81

The concentration of carbon dioxide in the atmosphere is 3.9×10^{-4}. Convert this number to decimal form.

Exercise 82

The width of a proton is 1×10^{-5} of the width of an atom. Convert this number to decimal form.

Answer

0.00001

Exercise 83

Health care costs The Centers for Medicare and Medicaid projects that consumers will spend more than $4 trillion on health care by 2017.

1. Write 4 trillion in decimal notation.
2. Write 4 trillion in scientific notation.

Exercise 84

Coin production In 1942, the U.S. Mint produced 154,500,000 nickels. Write 154,500,000 in scientific notation.

Answer

1.545×10^8

Exercise 85

Distance The distance between Earth and one of the brightest stars in the night star is 33.7 light years. One light year is about 6,000,000,000,000 (6 trillion), miles.

1. Write the number of miles in one light year in scientific notation.
2. Use scientific notation to find the distance between Earth and the star in miles. Write the answer in scientific notation.

Exercise 86

Debt At the end of fiscal year 2015 the gross United States federal government debt was estimated to be approximately $18,600,000,000,000 ($18.6 trillion), according to the Federal Budget. The population of the United States was approximately 300,000,000 people at the end of fiscal year 2015.

1. Write the debt in scientific notation.
2. Write the population in scientific notation.
3. Find the amount of debt per person by using scientific notation to divide the debt by the population. Write the answer in scientific notation.

Answer

1. 1.86×10^{13}
2. 3×10^{8}
3. 6.2×10^{4}

Writing Exercises

Exercise 87

1. Explain the meaning of the exponent in the expression 2^3.
2. Explain the meaning of the exponent in the expression 2^{-3}.

Exercise 88

When you convert a number from decimal notation to scientific notation, how do you know if the exponent will be positive or negative?

Answer

answers will vary

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
use the definition of a negative exponent.			
simplify expressions with integer exponents.			
convert from decimal notation to scientific notation.			
convert scientific notation to decimal form.			
multiply and divide using scientific notation.			

b. Overall, after looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

More Practice with Exponents in Quotients

Quotients

Exercise $6.7E.89$

★ **Simplify. (Assume all variables represent nonzero numbers.)**

1. $\dfrac{10^2 \cdot 10^4}{10^5}$
2. $\dfrac{7^5 \cdot 7^9}{7^2}$
3. $\dfrac{a^8 \cdot a^6}{a^5}$
4. $\dfrac{b^4 \cdot b^{10}}{b^8}$
5. $\dfrac{a^8 \cdot a^{-3}}{a^{-6}}$
6. $\dfrac{b^{-10} \cdot b^4}{b^{-2}}$
7. $\dfrac{25x^{-3}y^2}{5x^{-1}y^{-3}}$
8. $\dfrac{-9x^{-1}y^3z^{-5}}{3x^{-2}y^2z^{-1}}$
9. $\dfrac{40x^5y^3z}{4x^2y^2z}$
10. $\dfrac{8x^2y^5z^3}{16x^2yz}$
11. $\dfrac{24a^8b^3(a-5b)^{10}}{8a^5b^3(a-5b)^2}$
12. $\dfrac{175m^9n^5(m+n)^7}{25m^8n(m+n)^3}$

Answers to odd exercises.

1. 10
3. a^9
5. a^{11}
7. $\dfrac{5y^5}{x^2}$
9. $10x^3y$
11. $3a^3(a-5b)^8$

Powers of Quotients

Exercise $6.7E.90$

★ **Simplify. (Assume all variables represent nonzero numbers.)**

1. $\left(\dfrac{-3ab^2}{2c^3}\right)^3$
2. $\left(\dfrac{-10a^3b}{3c^2}\right)^2$
3. $\left(\dfrac{-2xy^4}{z^3}\right)^4$
4. $\left(\dfrac{-7x^9y}{z^4}\right)^3$
5. $\left(\dfrac{12x^3y^2z}{2x^7yz^8}\right)^3$
6. $\left(\dfrac{150xy^8z^2}{90x^7y^2z}\right)^2$
7. $\left(\dfrac{2x^{-3}z}{y^2}\right)^{-5}$
8. $\left(\dfrac{5x^5z^{-2}}{2y^{-3}}\right)^{-3}$
9. $\left(\dfrac{xy^2}{z^3}\right)^3$
10. $\left(\dfrac{2x^2y^3}{z}\right)^4$
11. $\left(\dfrac{-9a^{-3}b^4c^{-2}}{3a^3b^5c^{-7}}\right)^{-4}$
12. $\left(\dfrac{-15a^7b^5c^{-8}}{3a^{-6}b^2c^3}\right)^{-3}$

Answers to odd exercises:

1. $-\dfrac{27a^3b^6}{8c^9}$
3. $\dfrac{16x^4y^{16}}{z^{12}}$
5. $\dfrac{216y^3}{x^{12}z^{21}}$
7. $\dfrac{x^{15}y^{10}}{32z^5}$
9. $\dfrac{x^3y^6}{z^9}$
11. $\dfrac{a^{24}b^4}{81c^{20}}$

This page titled 6.7E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **6.7E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 6 Review Exercises

Chapter 6 Review Exercises

Add and Subtract Polynomials

Identify Polynomials, Monomials, Binomials and Trinomials

In the following exercises, determine if each of the following polynomials is a monomial, binomial, trinomial, or other polynomial.

? Exercise 1

1. $11c^4 - 23c^2 + 1$
2. $9p^3 + 6p^2 - p - 5$
3. $\frac{3}{7}x + \frac{5}{14}$
4. 10
5. 2y−12

? Exercise 2

1. $a^2 - b^2$
2. $24d^3$
3. $x^2 + 8x - 10$
4. $m^2n^2 - 2mn + 6$
5. $7y^3 + y^2 - 2y - 4$

Answer

1. binomial
2. monomial
3. trinomial
4. trinomial
5. other polynomial

Determine the Degree of Polynomials

In the following exercises, determine the degree of each polynomial.

? Exercise 3

1. $3x^2 + 9x + 10$
2. $14a^2bc$
3. 6y+1
4. $n^3 - 4n^2 + 2n - 8$
5. −19

? Exercise 4

1. $5p^3 - 8p^2 + 10p - 4$
2. $-20q^4$
3. $x^2 + 6x + 12$
4. $23r^2s^2 - 4rs + 5$
5. 100

Answer

1. 3
2. 4

3. 2
4. 4
5. 0

Add and Subtract Monomials

In the following exercises, add or subtract the monomials.

Exercise 5

$5y^3 + 8y^3$

Exercise 6

$-14k + 19k$

Answer

5k

Exercise 7

12q−(−6q)

Exercise 8

−9c−18c

Answer

−27c

Exercise 9

12x−4y−9x

Exercise 2

$3m^2 + 7n^2 - 3m^2$

Answer

$7n^2$

Exercise 3

$6x^2y - 4x + 8xy^2$

Exercise 4

13a+b

Answer

13a+b

Add and Subtract Polynomials

In the following exercises, add or subtract the polynomials.

Exercise 5

$(5x^2+12x+1)+(6x^2-8x+3)$

Exercise 6

$(9p^2-5p+3)+(4p^2-4)$

Answer

$13p^2-5p-1$

Exercise 7

$(10m^2-8m-1)-(5m^2+m-2)$

Exercise 8

$(7y^2-8y)-(y-4)$

Answer

$7y^2-9y+4$

Exercise 9

Subtract
$(3s^2+10)$ from $(15s^2-2s+8)$

Exercise 10

Find the sum of (a^2+6a+9) and $(5a^3-7)$

Answer

$5a^3+a^2+6a+2$

Evaluate a Polynomial for a Given Value of the Variable

In the following exercises, evaluate each polynomial for the given value.

Exercise 11

Evaluate $3y^2-y+1$ when:

1. y=5
2. y=−1
3. y=0

Exercise 12

Evaluate 10−12x when:

1. x=3

2. x=0
3. x=−1

Answer

1. −26
2. 10
3. 22

Exercise 13

Randee drops a stone off the 200 foot high cliff into the ocean. The polynomial $-16t^2+200$ gives the height of a stone t seconds after it is dropped from the cliff. Find the height after t=3 seconds.

Exercise 14

A manufacturer of stereo sound speakers has found that the revenue received from selling the speakers at a cost of p dollars each is given by the polynomial $-4p^2+460p$. Find the revenue received when p=75 dollars.

Answer

12,000

Use Multiplication Properties of Exponents

Simplify Expressions with Exponents

In the following exercises, simplify.

Exercise 15

10^4

Exercise 16

17^1

Answer

17

Exercise 17

$\left(\frac{2}{9}\right)^2$

Exercise 18

$(0.5)^3$

Answer

0.125

Exercise 19

$(-2)^6$

Exercise 20

-2^6

Answer

-64

Simplify Expressions Using the Product Property for Exponents

In the following exercises, simplify each expression.

Exercise 21

$x^4 \cdot x^3$

Exercise 22

$p^{15} \cdot p^{16}$

Answer

p^{31}

Exercise 23

$4^{10} \cdot 4^6$

Exercise 24

$8 \cdot 8^5$

Answer

8^6

Exercise 25

$n \cdot n^2 \cdot n^4$

Exercise 26

$y^c \cdot y^3$

Answer

y^{c+3}

Simplify Expressions Using the Power Property for Exponents

In the following exercises, simplify each expression.

Exercise 27

$(m^3)^5$

Exercise 28

$(5^3)^2$

Answer

5^6

Exercise 29

$(y^4)^x$

Exercise 30

$(3^r)^s$

Answer

3^{rs}

Simplify Expressions Using the Product to a Power Property

In the following exercises, simplify each expression.

Exercise 31

$(4a)^2$

Exercise 32

$(-5y)^3$

Answer

$-125y^3$

Exercise 33

$(2mn)^5$

Exercise 34

$(10xyz)^3$

Answer

$1000x^3y^3z^3$

Simplify Expressions by Applying Several Properties

In the following exercises, simplify each expression.

? Exercise 35

$\left(p^2\right)^5 \cdot \left(p^3\right)^6$

? Exercise 36

$\left(4a^3b^2\right)^3$

Answer

$64a^9b^6$

? Exercise 37

$(5x)^2(7x)$

? Exercise 38

$\left(2q^3\right)^4(3q)^2$

Answer

$48q^{14}$

? Exercise 39

$\left(\frac{1}{3}x^2\right)^2\left(\frac{1}{2}x\right)^3$

? Exercise 40

$\left(\frac{2}{5}m^2n\right)^3$

Answer

$\frac{8}{125}m^6n^3$

Multiply Monomials

In the following exercises 8, multiply the monomials.

? Exercise 41

$\left(-15x^2\right)\left(6x^4\right)$

? Exercise 42

$\left(-9n^7\right)(-16n)$

Answer

$144n^8$

Exercise 43

$\left(7p^5q^3\right)\left(8pq^9\right)$

Exercise 44

$\left(\frac{5}{9}ab^2\right)\left(27ab^3\right)$

Answer

$15a^2b^5$

Multiply Polynomials

Multiply a Polynomial by a Monomial

In the following exercises, multiply.

Exercise 45

7(a+9)

Exercise 46

−4(y+13)

Answer

−4y−52

Exercise 47

−5(r−2)

Exercise 48

p(p+3)

Answer

p^2+3p

Exercise 49

−m(m+15)

Exercise 50

−6u(2u+7)

Answer

$-12u^2-42u$

Exercise 51

$9\left(b^2+6b+8\right)$

Exercise 52

$3q^2\left(q^2-7q+6\right)3$

Answer

$3q^4-21q^3+18q^2$

Exercise 53

$(5z-1)z$

Exercise 54

$(b-4)\cdot 11$

Answer

11b−44

Multiply a Binomial by a Binomial

In the following exercises, multiply the binomials using:

1. the Distributive Property,
2. the FOIL method,
3. the Vertical Method.

Exercise 55

(x−4)(x+10)

Exercise 56

(6y−7)(2y−5)

Answer

1. $12y^2-44y+35$
2. $12y^2-44y+35$
3. $12y^2-44y+35$

In the following exercises, multiply the binomials. Use any method.

Exercise 57

(x+3)(x+9)

Exercise 58

(y−4)(y−8)

Answer

$y^2 - 12y + 32$

Exercise 59

(p−7)(p+4)

Exercise 60

(q+16)(q−3)

Answer

$q^2 + 13q - 48$

Exercise 61

(5m−8)(12m+1)

Exercise 62

$\left(u^2 + 6\right)\left(u^2 - 5\right)$

Answer

$u^4 + u^2 - 30$

Exercise 63

(9x−y)(6x−5)

Exercise 64

(8mn+3)(2mn−1)

Answer

$16m^2n^2 - 2mn - 3$

Multiply a Trinomial by a Binomial

In the following exercises, multiply using

1. the Distributive Property,
2. the Vertical Method.

Exercise 65

$(n + 1)\left(n^2 + 5n - 2\right)$

Exercise 66

$(3x - 4)\left(6x^2 + x - 10\right)$

Answer

1. $18x^3 - 21x^2 - 34x + 40$
2. $18x^3 - 21x^2 - 34x + 40$

In the following exercises, multiply. Use either method.

Exercise 67

$(y-2)\left(y^2-8y+9\right)$

Exercise 68

$(7m+1)\left(m^2-10m-3\right)$

Answer

$7m^3-69m^2-31m-3$

Special Products

Square a Binomial Using the Binomial Squares Pattern

In the following exercises, square each binomial using the Binomial Squares Pattern.

Exercise 69

$(c+11)^2$

Exercise 70

$(q-15)^2$

Answer

$q^2-30q+225$

Exercise 71

$\left(x+\frac{1}{3}\right)^2$

Exercise 72

$(8u+1)^2$

Answer

$64u^2+16u+1$

Exercise 73

$\left(3n^3-2\right)^2$

Exercise 74

$(4a-3b)^2$

Answer

$16a^2-24ab+9b^2$

Multiply Conjugates Using the Product of Conjugates Pattern

In the following exercises, multiply each pair of conjugates using the Product of Conjugates Pattern.

Exercise 75

(s−7)(s+7)

Exercise 76

$\left(y+\frac{2}{5}\right)\left(y-\frac{2}{5}\right)$

Answer

$y^2-\frac{4}{25}$

Exercise 77

$(12c+13)(12c-13)$

Exercise 78

(6−r)(6+r)

Answer

$36-r^2$

Exercise 79

$\left(u+\frac{3}{4}v\right)\left(u-\frac{3}{4}v\right)$

Exercise 80

$\left(5p^4-4q^3\right)\left(5p^4+4q^3\right)$

Answer

$25p^8-16q^6$

Recognize and Use the Appropriate Special Product Pattern

In the following exercises, find each product.

Exercise 81

$(3m+10)^2$

Exercise 82

(6a+11)(6a−11)

Answer

$36a^2-121$

Exercise 83

(5x+y)(x−5y)

Exercise 84

$\left(c^4+9d\right)^2$

Answer

$c^8+18c^4d+81d^2$

Exercise 85

$\left(p^5+q^5\right)\left(p^5-q^5\right)$

Exercise 86

$\left(a^2+4b\right)\left(4a-b^2\right)$

Answer

$4a^3+3a^2b-4b^3$

Divide Monomials

Simplify Expressions Using the Quotient Property for Exponents

In the following exercises, simplify.

Exercise 87

$\frac{u^{24}}{u^6}$

Exercise 88

$\frac{10^{25}}{10^5}$

Answer

10^{20}

Exercise 89

$\frac{3^4}{3^6}$

Exercise 90

$\frac{v^{12}}{v^{48}}$

Answer

$\frac{1}{v^{36}}$

Exercise 91

$\frac{x}{x^5}$

Exercise 92

$\frac{5}{5^8}$

Answer

$\frac{1}{5^7}$

Simplify Expressions with Zero Exponents

In the following exercises, simplify.

Exercise 93

75^0

Exercise 94

x^0

Answer

1

Exercise 95

-12^0

Exercise 96

$\left(-12^0\right)(-12)^0$

Answer

1

Exercise 97

$25x^0$

Exercise 98

$(25x)^0$

Answer

1

Exercise 99

$19n^0 - 25m^0$

Exercise 100

$(19n)^0 - (25m)^0$

Answer

0

Simplify Expressions Using the Quotient to a Power Property

In the following exercises, simplify.

Exercise 101

$\left(\frac{2}{5}\right)^3$

Exercise 102

$\left(\frac{m}{3}\right)^4$

Answer

$\frac{m^4}{81}$

Exercise 103

$\left(\frac{r}{s}\right)^8$

Exercise 104

$\left(\frac{x}{2y}\right)^6$

Answer

$\frac{x^6}{64y^6}$

Simplify Expressions by Applying Several Properties

In the following exercises, simplify.

Exercise 105

$\frac{\left(x^3\right)^5}{x^9}$

Exercise 106

$\frac{n^{10}}{\left(n^5\right)^2}$

Answer

1

Exercise 107

$\left(\frac{q^6}{q^8}\right)^3$

Exercise 108

$\left(\frac{r^8}{r^3}\right)^4$

Answer

r^{20}

Exercise 109

$\left(\frac{c^2}{d^5}\right)^9$

Exercise 110

$\left(\frac{3x^4}{2y^2}\right)^5$

Answer

$\frac{343x^{20}}{32y^{10}}$

Exercise 111

$\left(\frac{v^3v^9}{v^6}\right)^4$

Exercise 112

$\frac{(3n^2)^4(-5n^4)^3}{(-2n^5)^2}$

Answer

$-\frac{10{,}125n^{10}}{4}$

Divide Monomials

In the following exercises, divide the monomials.

Exercise 113

$-65y^{14} \div 5y^2$

Exercise 114

$\frac{64a^5b^9}{-16a^{10}b^3}$

Answer

$-\frac{4b^6}{a^5}$

? Exercise 115

$\frac{144x^{15}y^8z^3}{18x^{10}y^2z^{12}}$

? Exercise 116

$\frac{(8p^6q^2)(9p^3q^5)}{16p^8q^7}$

Answer

$\frac{9p}{2}$

Divide Polynomials

Divide a Polynomial by a Monomial

In the following exercises, divide each polynomial by the monomial.

? Exercise 117

$\frac{42z^2-18z}{6}$

? Exercise 118

$(35x^2-75x)\div 5x$

Answer

7x−15

? Exercise 119

$\frac{81n^4+105n^2}{-3}$

? Exercise 120

$\frac{550p^6-300p^4}{10p^3}$

Answer

$55p^3-30p$

? Exercise 121

$(63xy^3+56x^2y^4)\div(7xy)$

? Exercise 122

$\frac{96a^5b^2-48a^4b^3-56a^2b^4}{8ab^2}$

Answer

$12a^4-6a^3b-7ab^2$

Exercise 123

$\frac{57m^2-12m+1}{-3m}$

Exercise 124

$\frac{105y^5+50y^3-5y}{5y^3}$

Answer

$21y^2+10-\frac{1}{y^2}$

Divide a Polynomial by a Binomial

In the following exercises, divide each polynomial by the binomial.

Exercise 125

$\left(k^2-2k-99\right)\div(k+9)$

Exercise 126

$\left(v^2-16v+64\right)\div(v-8)$

Answer

v−8

Exercise 127

$\left(3x^2-8x-35\right)\div(x-5)$

Exercise 128

$\left(n^2-3n-14\right)\div(n+3)$

Answer

$n-6+\frac{4}{n+3}$

Exercise 129

$\left(4m^3+m-5\right)\div(m-1)$

Exercise 130

$\left(u^3-8\right)\div(u-2)$

Answer

u^2+2u+4

Integer Exponents and Scientific Notation

Use the Definition of a Negative Exponent

In the following exercises, simplify.

Exercise 131

9^{-2}

Exercise 132

$(-5)^{-3}$

Answer

$-\frac{1}{125}$

Exercise 133

$3 \cdot 4^{-3}$

Exercise 134

$(6u)^{-3}$

Answer

$\frac{1}{216u^3}$

Exercise 135

$\left(\frac{2}{5}\right)^{-1}$

Exercise 136

$\left(\frac{3}{4}\right)^{-2}$

Answer

$\frac{16}{9}$

Simplify Expressions with Integer Exponents

In the following exercises, simplify.

Exercise 137

$p^{-2} \cdot p^{8}$

Exercise 138

$q^{-6} \cdot q^{-5}$

Answer

$\frac{1}{q^{11}}$

Exercise 139

$\left(c^{-2}d\right)\left(c^{-3}d^{-2}\right)$

Exercise 140

$\left(y^{8}\right)^{-1}$

Answer

$\frac{1}{y^{8}}$

Exercise 141

$\left(q^{-4}\right)^{-3}$

Exercise 142

$\frac{a^{8}}{a^{12}}$

Answer

$\frac{1}{a^{4}}$

Exercise 143

$\frac{n^{5}}{n^{-4}}$

Exercise 144

$\frac{r^{-2}}{r^{-3}}$

Answer

r

Convert from Decimal Notation to Scientific Notation

In the following exercises, write each number in scientific notation.

Exercise 145

8,500,000

Exercise 146

0.00429

Answer

4.29×10^{-3}

Exercise 147

The thickness of a dime is about 0.053 inches.

Exercise 148

In 2015, the population of the world was about 7,200,000,000 people.

Answer

7.2×10^9

Convert Scientific Notation to Decimal Form

In the following exercises, convert each number to decimal form.

Exercise 149

3.8×10^5

Exercise 150

1.5×10^{10}

Answer

15,000,000,000

Exercise 151

9.1×10^{-7}

Exercise 152

5.5×10^{-1}

Answer

0.55

Multiply and Divide Using Scientific Notation

In the following exercises, multiply and write your answer in decimal form.

Exercise 153

$\left(2 \times 10^5\right)\left(4 \times 10^{-3}\right)$

Exercise 154

$\left(3.5 \times 10^{-2}\right)\left(6.2 \times 10^{-1}\right)$

Answer

0.0217

In the following exercises, divide and write your answer in decimal form.

Exercise 155

$\frac{8\times10^{5}}{4\times10^{-1}}$

Exercise 156

$\frac{9\times10^{-5}}{3\times10^{2}}$

Answer

0.0000003

Chapter Practice Test

Exercise 1

For the polynomial $10x^4+9y^2-1$
ⓐ Is it a monomial, binomial, or trinomial?
ⓑ What is its degree?

In the following exercises, simplify each expression.

Exercise 2

$\left(12a^2-7a+4\right)+\left(3a^2+8a-10\right)$

Answer

$15a^2+a-6$

Exercise 3

$\left(9p^2-5p+1\right)-\left(2p^2-6\right)$

Exercise 4

$\left(-\frac{2}{5}\right)^3$

Answer

$-\frac{8}{125}$

Exercise 5

$u\cdot u^4$

Exercise 6

$\left(4a^3b^5\right)^2$

Answer

$16a^6b^{10}$

Exercise 7

$\left(-9r^4s^5\right)\left(4rs^7\right)$

Exercise 8

$3k\left(k^2-7k+13\right)$

Answer

$3k^3-21k^2+39k$

Exercise 9

$(m+6)(m+12)$

Exercise 10

(v−9)(9v−5)

Answer

$9v^2-86v+45$

Exercise 11

(4c−11)(3c−8)

Exercise 12

$(n-6)\left(n^2-5n+4\right)$

Answer

$n^3-11n^2+34n-24$

Exercise 13

$(2x-15y)(5x+7y)$

Exercise 14

$(7p-5)(7p+5)$

Answer

$49p^2-25$

Exercise 15

$(9v-2)^2$

Exercise 16

$\frac{3^8}{3^{10}}$

Answer

$\frac{1}{9}$

Exercise 17

$\left(\frac{m^4 \cdot m}{m^3}\right)^6$

Exercise 18

$\left(87x^{15}y^3z^{22}\right)^0$

Answer

1

Exercise 19

$\frac{80c^8d^2}{16cd^{10}}$

Exercise 20

$\frac{12x^2+42x-6}{2x}$

Answer

$6x+21-\frac{3}{x}$

Exercise 21

$\left(70xy^4+95x^3y\right) \div 5xy$

Exercise 22

$\frac{64x^3-1}{4x-1}$

Answer

$16x^2+4x+1$

Exercise 23

$\left(y^2-5y-18\right) \div (y+3)$

Exercise 24

5^{-2}

Answer

$\frac{1}{25}$

Exercise 25

$(4m)^{-3}$

Exercise 26

$q^{-4} \cdot q^{-5}$

Answer

$\frac{1}{q^9}$

Exercise 27

$\frac{n^{-2}}{n^{-10}}$

Exercise 28

Convert 83,000,000 to scientific notation.

Answer

8.3×10^7

Exercise 29

Convert 6.91×10^{-5} to decimal form.

In the following exercises, simplify, and write your answer in decimal form.

Exercise 30

$\left(3.4 \times 10^9\right)\left(2.2 \times 10^{-5}\right)$

Answer

74,800

Exercise 31

$\frac{8.4 \times 10^{-3}}{4 \times 10^3}$

Exercise 32

A helicopter flying at an altitude of 1000 feet drops a rescue package. The polynomial $-16t^2 + 1000$ gives the height of the package t seconds a after it was dropped. Find the height when t=6 seconds.

Answer

424 feet

Chapter 6 Review Exercises is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **Chapter 6 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

CHAPTER OVERVIEW

7: Factoring

7.1: Greatest Common Factor and Factor by Grouping

7.1E: Exercises

7.2: Factor Quadratic Trinomials with Leading Coefficient 1

7.2E: Exercises

7.3: Factor Quadratic Trinomials with Leading Coefficient Other than 1

7.3E: Exercises

7.4: Factor Special Products

7.4E: Exercises

7.5: General Strategy for Factoring Polynomials

7.5E: Exercises

7.6: Quadratic Equations

7.6E: Exercises

Chapter 7 Review Exercises

This page titled 7: Factoring is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

7.1: Greatest Common Factor and Factor by Grouping

Learning Objectives

By the end of this section, you will be able to:

- Find the greatest common factor of two or more expressions
- Factor the greatest common factor from a polynomial
- Factor by grouping


BE PREPARED

Before you get started, take this readiness quiz.

1. Factor 56 into primes.
   If you missed this problem, review Example 1.2.19.
2. Find the least common multiple of 18 and 24.
   If you missed this problem, review Example 1.2.28.
3. Simplify $-3(6a+11)$.
   If you missed this problem, review Example 1.10.40.

Find the Greatest Common Factor of Two or More Expressions

Earlier we multiplied factors together to get a product. Now, we will be reversing this process; we will start with a product and then break it down into its factors. Splitting a product into factors is called **factoring**.



We have learned how to factor numbers to find the least common multiple (LCM) of two or more numbers. Now we will factor expressions and find the **greatest common factor** of two or more expressions. The method we use is similar to what we used to find the LCM.

GREATEST COMMON FACTOR


The greatest common factor (GCF) of two or more expressions is the largest expression that is a factor of all the expressions.

First we'll find the GCF of two numbers.

Example 7.1.1: HOW TO FIND THE GREATEST COMMON FACTOR OF TWO OR MORE EXPRESSIONS

Find the GCF of 54 and 36.

Solution

Step 1. Factor each coefficient into primes. Write all variables with exponents in expanded form.	Factor 54 and 36.	
Step 2. In each column, circle the common factors.	Circle the 2, 3, and 3 that are shared by both numbers.	$36 = 2 \cdot 2 \cdot 3 \cdot 3$ $18 = 2 \cdot 3 \cdot 3 \cdot 3$
Step 3. Bring down the common factors that all expressions share.	Bring down the 2, 3, and 3, and then multiply.	$GCF = 2 \cdot 3 \cdot 3$
Step 4. Multiply the factors.		$GCF = 18$ The GCF of 54 and 36 is 18.

Notice that, because the GCF is a factor of both numbers, 54 and 36 can be written as multiples of 18.

$$\begin{aligned} 54 &= 18 \cdot 3 \\ 36 &= 18 \cdot 2 \end{aligned} \tag{7.1.1}$$

? Try It 7.1.2

Find the GCF of 48 and 80.

Answer

16

? Try It 7.1.3

Find the GCF of 18 and 40.

Answer

2

We summarize the steps we use to find the GCF below.

HOW TO

Find the Greatest Common Factor (GCF) of two expressions.

1. Step 1. Factor each coefficient into primes. Write all variables with exponents in expanded form.
2. Step 2. List all factors—matching common factors in a column. In each column, circle the common factors.
3. Step 3. Bring down the common factors that all expressions share.
4. Step 4. Multiply the factors.

In the first example, the GCF was a constant. In the next two examples, we will get variables in the greatest common factor.

Example 7.1.4

Find the greatest common factor of $27x^3$ and $18x^4$.

Solution

Factor each coefficient into primes and write the variables with exponents in expanded form. Circle the common factors in each column.	$27x^3 = 3 \cdot 3 \cdot 3 \cdot x \cdot x \cdot x$ $18x^4 = 2 \cdot 3 \cdot 3 \cdot x \cdot x \cdot x \cdot x$
Bring down the common factors.	$\text{GCF} = 3 \cdot 3 \cdot x \cdot x \cdot x$
Multiply the factors.	$\text{GCF} = 9x^3$
	The GCF of $27x^3$ and $18x^4$ is $9x^3$.

Try It 7.1.5

Find the GCF: $12x^2, 18x^3$

Answer

$6x^2$

Try It 7.1.6

Find the GCF: $16y^2, 24y^3$

Answer

$8y^2$

Example 7.1.7

Find the GCF of $4x^2y, 6xy^3$

Solution

Factor each coefficient into primes and write the variables with exponents in expanded form. Circle the common factors in each column.	$4x^2y = 2 \cdot 2 \cdot x \cdot x \cdot y$ $6xy^3 = 2 \cdot 3 \cdot x \cdot y \cdot y \cdot y$
Bring down the common factors.	$\text{GCF} = 2 \cdot x \cdot y$
Multiply the factors.	$\text{GCF} = 2xy$
	The GCF of $4x^2y$ and $6xy^3$ is $2xy$.

Try It 7.1.8

Find the GCF: $6ab^4, 8a^2b$

Answer

$2ab$

Try It 7.1.9

Find the GCF: $9m^5n^2, 12m^3n$

Answer

$3m^3n$

Example 7.1.10

Find the GCF of: $21x^3, 9x^2, 15x$

Solution

Factor each coefficient into primes and write the variables with exponents in expanded form. Circle the common factors in each column.	$21x^3 = 3 \cdot 7 \cdot x \cdot x \cdot x$ $9x^2 = 3 \cdot 3 \cdot x \cdot x$ $15x = 3 \cdot 5 \cdot x$
Bring down the common factors.	GCF = $3 \cdot x$
Multiply the factors.	GCF = $3x$
	The GCF of $21x^3, 9x^2$ and $15x$ is $3x$

Try It 7.1.11

Find the greatest common factor: $25m^4, 35m^3, 20m^2$

Answer

$5m^2$

Try It 7.1.12

Find the greatest common factor: $14x^3, 70x^2, 105x$

Answer

$7x$

Factor the Greatest Common Factor from a Polynomial

Just like in arithmetic, where it is sometimes useful to represent a number in factored form (for example, 12 as 2·6or3·4),2·6or3·4), in algebra, it can be useful to represent a polynomial in factored form. One way to do this is by finding the GCF of all the terms. Remember, we multiply a polynomial by a monomial as follows:

$$\begin{aligned} &2(x+7) && \text{factors} \\ &2\cdot x+2\cdot 7 && \\ &2x+14 && \text{product} \end{aligned} \tag{7.1.2}$$

Now we will start with a product, like $2x+14$, and end with its factors, $2(x+7)$. To do this we apply the Distributive Property "in reverse."

We state the Distributive Property here just as you saw it in earlier chapters and "in reverse."

DISTRIBUTIVE PROPERTY

If a, b, c are real numbers, then

$$a(b+c) = ab+ac \quad \text{and} \quad ab+ac = a(b+c) \tag{7.1.3}$$

The form on the left is used to multiply. The form on the right is used to factor.

So how do you use the Distributive Property to factor a polynomial? You just find the GCF of all the terms and write the polynomial as a product!

Example 7.1.13: HOW TO FACTOR THE GREATEST COMMON FACTOR FROM A POLYNOMIAL

Factor: $4x+12$

Solution

Step 1. Find the GCF of all the terms of the polynomial.	Find the GCF of $4x$ and 12.	$4x = 2 \cdot 2 \cdot \cdot x$ $12 = 2 \cdot 2 \cdot 3$ $\text{GCF} = 2 \cdot 2$ $\text{GCF} = 4$
Step 2. Rewrite each term as a product using the GCF.	Rewrite $4x$ and 12 as products of their GCF, 4. $4x = 4 \cdot x$ $12 = 4 \cdot 3$	$4x + 12$ $4 \cdot x + 4 \cdot 3$
Step 3. Use the "reverse" Distributive Property to factor the expression.		$4(x + 3)$
Step 4. Check by multiplying the factors.		$4(x + 3)$ $4 \cdot x + 4 \cdot 3$ $4x + 12$ ✓

Try It 7.1.14

Factor: $6a+24$

Answer

$6(a+4)$

Try It 7.1.15

Factor: $2b+14$

Answer

$2(b+7)$

HOW TO

Factor the greatest common factor from a polynomial.

Step 1. Find the GCF of all the terms of the polynomial.

Step 2. Rewrite each term as a product using the GCF.

Step 3. Use the "reverse" Distributive Property to factor the expression.

Step 4. Check by multiplying the factors.

FACTOR AS A NOUN AND A VERB

We use “factor” as both a noun and a verb.

Noun	7 is a factor of 14
Verb	factor 3 from $3a+3$

Example 7.1.16

Factor: $5a+5$

Solution

Find the GCF of $5a$ and 5.	$5a = 5 \cdot a$ $5 = 5$ $\text{GCF} = 5$
	$5a+5$
Rewrite each term as a product using the GCF.	$5 \cdot a + 5 \cdot 1$
Use the Distributive Property "in reverse" to factor the GCF.	$5(a+1)$
Check by multiplying the factors to get the original polynomial.	
$5(a+1)$	
$5 \cdot a + 5 \cdot 1$	
$5a+5$✓	

Try It 7.1.17

Factor: $14x+14$

Answer

$14(x+1)$

Try It 7.1.18

Factor: $12p+12$

Answer

$12(p+1)$

The expressions in the next example have several factors in common. Remember to write the GCF as the product of all the common factors.

Example 7.1.19

Factor: $12x-60$

Solution

Find the GCF of $12x$ and 60.	$12x = 2 \cdot 2 \cdot 3 \cdot x$ $60 = 2 \cdot 2 \cdot 3 \cdot 5$ $GCF = 2 \cdot 2 \cdot 3$ $GCF = 12$
	$12x - 60$
Rewrite each term as a product using the GCF.	$12 \cdot x - 12 \cdot 5$
Factor the GCF.	$12(x - 5)$
Check by multiplying the factors.	
$12(x-5)$	
$12 \cdot x - 12 \cdot 5$	
$12x - 60$✓	

Try It 7.1.20

Factor: $18u - 36$

Answer

$8(u - 2)$

Try It 7.1.21

Factor: $30y - 60$

Answer

$30(y - 2)$

Now we'll factor the greatest common factor from a trinomial. We start by finding the GCF of all three terms.

Example 7.1.22

Factor: $4y^2 + 24y + 28$

Solution

We start by finding the GCF of all three terms.

Find the GCF of $4y^2$, $24y$ and 28	$4y^2 = 2 \cdot 2 \cdot y \cdot y$ $24y = 2 \cdot 2 \cdot 2 \cdot 3 \cdot y$ $28 = 2 \cdot 2 \cdot 7$ $GCF = 2 \cdot 2$ $GCF = 4$
	$4y^2 + 24y + 28$
Rewrite each term as a product using the GCF.	$4 \cdot y^2 + 4 \cdot 6y + 4 \cdot 7$
Factor the GCF.	$4(y^2 + 6y + 7)$
Check by multiplying.	
$4\left(y^2 + 6y + 7\right)$	
$4 \cdot y^2 + 4 \cdot 6y + 4 \cdot 7$	

$4y^2+24y+28$✓

Try It 7.1.23

Factor: $5x^2-25x+15$

Answer

$5\left(x^2-5x+3\right)$

Try It 7.1.24

Factor: $3y^2-12y+27$

Answer

$3\left(y^2-4y+9\right)$

Example 7.1.25

Factor: $5x^3-25x^2$

Solution

Find the GCF of $5x^3$ and $25x^2$	$5x^3 = 5 \cdot x \cdot x \cdot x$ $25x^2 = 5 \cdot 5 \cdot x \cdot x$ $GCF = 5 \cdot x \cdot x$ $GCF = 5x^2$
	$5x^3-25x^2$
Rewrite each term.	$5x^2 \cdot x - 5x^2 \cdot 5$
Factor the GCF.	$5x^2(x-5)$
Check.	
$5x^2(x-5)$	
$5x^2 \cdot x - 5x^2 \cdot 5$	
$5x^3-25x^2$✓	

Try It 7.1.26

Factor: $2x^3+12x^2$

Answer

$2x^2(x+6)$

Try It 7.1.27

Factor: $6y^3-15y^2$

Answer

$3y^2(2y-5)$

Example 7.1.28

Factor: $21x^3 - 9x^2 + 15x$

Solution

In a previous example we found the GCF of $21x^3, 9x^2, 15x$ to be $3x$.

	$21x^3 - 9x^2 + 15x$
Rewrite each term using the GCF, $3x$.	$3x \cdot 7x^2 - 3x \cdot 3x + 3x \cdot 5$
Factor the GCF.	$3x(7x^2 - 3x + 5)$
Check.	
$3x\left(7x^2 - 3x + 5\right)$	
$3x \cdot 7x^2 - 3x \cdot 3x + 3x \cdot 5$	
$21x^3 - 9x^2 + 15x$✓	

Try It 7.1.29

Factor: $20x^3 - 10x^2 + 14x$

Answer

$2x(10x^2 - 5x + 7)$

Try It 7.1.30

Factor: $24y^3 - 12y^2 - 20y$

Answer

$4y(6y^2 - 3y - 5)$

Example 7.1.31

Factor: $8m^3 - 12m^2n + 20mn^2$

Solution

Find the GCF of $8m^3, 12m^2n, 20mn^2$	$8m^3 = 2 \cdot 2 \cdot 2 \cdot m \cdot m \cdot m$ $12m^2n = 2 \cdot 2 \cdot 3 \cdot m \cdot m \cdot n$ $20mn^2 = 2 \cdot 2 \cdot 5 \cdot m \cdot n \cdot n$ $GCF = 2 \cdot 2 \cdot m$ $GCF = 4m$
	$8m^3 - 12m^2n + 20mn^2$
Rewrite each term.	$4m \cdot 2m^2 - 4m \cdot 3m\,n + 4m \cdot 5n^2$
Factor the GCF.	$4m(2m^2 - 3m\,n + 5n^2)$
Check.	
$4m\left(2m^2 - 3mn + 5n^2\right)$	
$4m \cdot 2m^2 - 4m \cdot 3mn + 4m \cdot 5n^2$	

$8m^3 - 12m^2n + 20mn^2$ ✓

Try It 7.1.32

Factor: $9xy^2 + 6x^2y^2 + 21y^3$

Answer

$3y^2(3x + 2x^2 + 7y)$

Try It 7.1.33

Factor: $3p^3 - 6p^2q + 9pq^3$

Answer

$3p(p^2 - 2pq + 3q^2$

When the leading coefficient is negative, we factor the negative out as part of the GCF.

Example 7.1.34

Factor: $-8y - 24$

Solution

When the leading coefficient is negative, the GCF will be negative.

Ignoring the signs of the terms, we first find the GCF of 8*y* and 24 is 8. Since the expression −8*y* − 24 has a negative leading coefficient, we use −8 as the GCF.	$8y = 2 \cdot 2 \cdot 2 \cdot y$ $24 = 2 \cdot 2 \cdot 2 \cdot 3$ $GCF = 2 \cdot 2 \cdot 2$ $GCF = 8$
Rewrite each term using the GCF.	$-8y - 24$ $-8 \cdot y + (-8) \cdot 3$
Factor the GCF.	$-8(y + 3)$
Check.	
$-8(y + 3)$	
$-8 \cdot y + (-8) \cdot 3$	
$-8y - 24$ ✓	

Try It 7.1.35

Factor: $-16z - 64$

Answer

$-16(z + 4)$

Try It 7.1.36

Factor: $-9y-27$

Answer

$-9(y+3)$

Example 7.1.37

Factor: $-6a^2+36a$

Solution

The leading coefficient is negative, so the GCF will be negative.?

Since the leading coefficient is negative, the GCF is negative, $-6a$.	$6a^2 = 2 \cdot 3 \cdot a \cdot a$ $36a = 2 \cdot 2 \cdot 3 \cdot 3 \cdot a$ $GCF = 2 \cdot 3 \cdot a$ $GCF = 6a$ $-6a^2 + 36a$
Rewrite each term using the GCF.	$-6a \cdot a - (-6a) \cdot 6$
Factor the GCF.	$-6a(a - 6)$
Check.	
$-6a(a-6)$	
$-6a \cdot a + (-6a)(-6)$	
$-6a^2+36av$	

Try It 7.1.38

Factor: $-4b^2+16b$

Answer

$-4b(b-4)$

Try It 7.1.39

Factor: $-7a^2+21a$

Answer

$-7a(a-3)$

Example 7.1.40

Factor: $5q(q+7)-6(q+7)$

Solution

The GCF is the binomial q+7.

	$5q(q + 7) - 6(q + 7)$

Factor the GCF, $(q + 7)$.	$(q + 7)(5q - 6)$
Check on your own by multiplying.	

Try It 7.1.41

Factor: $4m(m+3)-7(m+3)$

Answer

$(m+3)(4m-7)$

Try It 7.1.42

Factor: $8n(n-4)+5(n-4)$

Answer

$(n-4)(8n+5)$

Factor by Grouping

When there is no common factor of all the terms of a polynomial, look for a common factor in just some of the terms. When there are four terms, a good way to start is by separating the polynomial into two parts with two terms in each part. Then look for the GCF in each part. If the polynomial can be factored, you will find a common factor emerges from both parts.

(Not all polynomials can be factored. Just like some numbers are prime, some polynomials are prime.)

Example 7.1.43

Factor: $xy+3y+2x+6$

Solution

Step 1. Group terms with common factors.	Is there a greatest common factor of all four terms? No, so let's separate the first two terms from the second two.	$xy + 3y + 2x + 6$ $\underline{xy + 3y}\underline{+ 2x + 6}$
Step 2. Factor out the common factor in each group.	Factor the GCF from the first two terms. Factor the GCF from the second two terms.	$y(x + 3)\underline{+ 2x + 6}$ $y(x + 3) + 2(x + 3)$
Step 3. Factor the common factor from the expression.	Notice that each term has a common factor of $(x + 3)$. Factor out the common factor.	$y(x + 3) + 2(x + 3)$ $(x + 3)\,(y + 2)$
Step 4. Check.	Multiply $(x + 3)(y + 2)$. Is the product the original expression?	$(x + 3)\,(y + 2)$ $xy + 2x + 3y + 6$ $xy + 3y + 2x + 6$ ✓

Try It 7.1.44

Factor: $xy+8y+3x+24$

Answer

$(x+8)(y+3)$

Try It 7.1.45

Factor: $ab+7b+8a+56$

Answer

$(a+7)(b+8)$

HOW TO

Factor by grouping.

Step 1. Group terms with common factors.

Step 2. Factor out the common factor in each group.

Step 3. Factor the common factor from the expression.

Step 4. Check by multiplying the factors.

Example 7.1.46

Factor: $x^2+3x-2x-6$

Solution

There is no GCF in all four terms.	$x^2+3x-2x-6$
Separate into two parts.	$\underbrace{x^2+3x}\ \underbrace{-2x-6}$
Factor the GCF from both parts. Be careful with the signs when factoring the GCF from the last two terms.	$x(x+3)-2(x+3)$
	$(x+3)(x-2)$

Check on your own by multinlying.

Try It 7.1.47

Factor: $x^2+2x-5x-10$

Answer

$(x-5)(x+2)$

Try It 7.1.48

Factor: $y^2+4y-7y-28$

Answer

$(y+4)(y-7)$

MEDIA ACCESS ADDITIONAL ONLINE RESOURCES

Access these online resources for additional instruction and practice with greatest common factors (GFCs) and factoring by grouping.

- Greatest Common Factor (GCF)
- Factoring Out the GCF of a Binomial
- Greatest Common Factor (GCF) of Polynomials

Glossary

factoring

Factoring is splitting a product into factors; in other words, it is the reverse process of multiplying.

greatest common factor

The greatest common factor is the largest expression that is a factor of two or more expressions is the greatest common factor (GCF).

This page titled 7.1: Greatest Common Factor and Factor by Grouping is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.1: Greatest Common Factor and Factor by Grouping** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.1E: Exercises

Practice Makes Perfect

Find the Greatest Common Factor of Two or More Expressions

In the following exercises, find the greatest common factor.

Exercise 1

$8,\ 18$

Answer

2

Exercise 2

$24,\ 40$

Exercise 3

$72,\ 162$

Answer

18

Exercise 4

$150,\ 275$

Exercise 5

$10a,\ 50$

Answer

10

Exercise 6

$5b,\ 30$

Exercise 7

$3x,\ 10x^2$

Answer

x

Exercise 8

$21b^2,\ 14b$

Exercise 9

$8w^2, 24w^3$

Answer

$8w^2$

Exercise 10

$30x^2, 18x^3$

Exercise 11

$10p^3q, 12pq^2$

Answer

$2pq$

Exercise 12

$8a^2b^3, 10ab^2$

Exercise 13

$12m^2n^3, 30m^5n^3$

Answer

$6m^2n^3$

Exercise 14

$28x^2y^4, 42x^4y^4$

Exercise 15

$10a^3, 12a^2$, 14a

Answer

$2a$

Exercise 16

$20y^3, 28y^2$, 40y

Exercise 17

$35x^3, 10x^4, 5x^5$

Answer

$5x^3$

Exercise 18

$27p^2$, $45p^3$, $9p^4$

Factor the Greatest Common Factor from a Polynomial

In the following exercises, factor the greatest common factor from each polynomial.

Exercise 19

$4x+20$

Answer

4(x+5)

Exercise 20

$8y+16$

Exercise 21

$6m+9$

Answer

$3(2m+3)$

Exercise 22

$14p+35$

Exercise 23

$9q+9$

Answer

$9(q+1)$

Exercise 24

$7r+7$

Exercise 25

$8m-8$

Answer

$8(m-1)$

Exercise 26

$4n-4$

Exercise 27

$9n - 63$

Answer

$9(n - 7)$

Exercise 28

$45b - 18$

Exercise 29

$3x^2 + 6x - 9$

Answer

$3(x^2 + 2x - 3)$

Exercise 30

$4y^2 + 8y - 4$

Exercise 31

$8p^2 + 4p + 2$

Answer

$2(4p^2 + 2p + 1)$

Exercise 32

$10q^2 + 14q + 20$

Exercise 33

$8y^3 + 16y^2$

Answer

$8y^2(y + 2)$

Exercise 34

$12x^3 - 10x$

Exercise 35

$5x^3 - 15x^2 + 20x$

Answer

$5x(x^2 - 3x + 4)$

https://math.libretexts.org/@go/page/30550

Exercise 36

$8m^2 - 40m + 16$

Exercise 37

$12xy^2 + 18x^2y^2 - 30y^3$

Answer

$6y^2(2x + 3x^2 - 5y)$

Exercise 38

$21pq^2 + 35p^2q^2 - 28q^3$

Exercise 39

$-2x - 4$

Answer

$-2(x + 2)$

Exercise 40

$-3b + 12$

Exercise 41

$5x(x + 1) + 3(x + 1)$

Answer

$(x + 1)(5x + 3)$

Exercise 42

$2x(x - 1) + 9(x - 1)$

Exercise 43

$3b(b - 2) - 13(b - 2)$

Answer

$(b - 2)(3b - 13)$

Exercise 44

$6m(m - 5) - 7(m - 5)$

Factor by Grouping

In the following exercises, factor by grouping.

https://math.libretexts.org/@go/page/30550

Exercise 45

$xy+2y+3x+6$

Answer

$(y+3)(x+2)$

Exercise 46

$mn+4n+6m+24$

Exercise 47

$uv-9u+2v-18$

Answer

$(u+2)(v-9)$

Exercise 48

$pq-10p+8q-80$

Exercise 49

$b^2+5b-4b-20$

Answer

$(b-4)(b+5)$

Exercise 50

$m^2+6m-12m-72$

Exercise 51

$p^2+4p-9p-36$

Answer

$(p-9)(p+4)$

Exercise 52

$x^2+5x-3x-15$

Mixed Practice

In the following exercises, factor.

Exercise 53

$-20x-10$

Answer

https://math.libretexts.org/@go/page/30550

$-10(2x+1)$

Exercise 54

$5x^3-x^2+x$

Exercise 55

$3x^3-7x^2+6x-14$

Answer

$(x^2+2)(3x-7)$

Exercise 56

x^3+x^2-x-1

Exercise 57

$x^2+xy+5x+5y$

Answer

$(x+y)(x+5)$

Exercise 58

$5x^3-3x^2-5x-3$

Everyday Math

Exercise 59

Area of a rectangle The area of a rectangle with length 6 less than the width is given by the expression w^2-6w, where $w=$ width. Factor the greatest common factor from the polynomial.

Answer

$w(w-6)$

Exercise 60

Height of a baseball The height of a baseball t seconds after it is hit is given by the expression $-16t^2+80t+4$

Writing Exercises

Exercise 61

The greatest common factor of 36 and 60 is 12. Explain what this means.

Answer

Answers will vary.

Exercise 62

What is the GCF of y^4, y^5, and y^{10}? Write a general rule that tells you how to find the GCF of y^a, y^b, and y^c.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
find the greatest common factor of two or more expressions.			
factor the greatest common factor from a polynomial.			
factor by grouping.			

b. If most of your checks were:

…confidently. Congratulations! You have achieved your goals in this section! Reflect on the study skills you used so that you can continue to use them. What did you do to become confident of your ability to do these things? Be specific!

…with some help. This must be addressed quickly as topics you do not master become potholes in your road to success. Math is sequential—every topic builds upon previous work. It is important to make sure you have a strong foundation before you move on. Who can you ask for help? Your fellow classmates and instructor are good resources. Is there a place on campus where math tutors are available? Can your study skills be improved?

…no - I don't get it! This is critical and you must not ignore it. You need to get help immediately or you will quickly be overwhelmed. See your instructor as soon as possible to discuss your situation. Together you can come up with a plan to get you the help you need.

This page titled 7.1E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.1E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.2: Factor Quadratic Trinomials with Leading Coefficient 1

Learning Objectives

By the end of this section, you will be able to:

- Factor trinomials of the form x^2+bx+c
- Factor trinomials of the form $x^2+bxy+cy^2$

BE PREPARED

Before you get started, take this readiness quiz.

1. Multiply: (x+4)(x+5).
   If you missed this problem, review Example 6.3.31.
2. Simplify: (a) $-9+(-6)$ (b) $-9+6$.
   If you missed this problem, review Example 1.4.18.
3. Simplify: (a) $-9(6)$ (b) $-9(-6)$.
   If you missed this problem, review Example 1.5.1.
4. Simplify: (a) $|-5|$ (b) $|3|$.
   If you missed this problem, review Example 1.4.2.

Factor Trinomials of the Form x^2+bx+c

You have already learned how to multiply binomials using FOIL. Now you'll need to "undo" this multiplication—to start with the product and end up with the factors. Let's look at an example of multiplying binomials to refresh your memory.

$(x+2)(x+3)$ factors
F O I L
$x^2+3x+2x+6$
x^2+5x+6 product

To factor the trinomial means to start with the product, x^2+5x+6, and end with the factors, $(x+2)(x+3)$. You need to think about where each of the terms in the trinomial came from.

The *first term* came from multiplying the first term in each binomial. So to get x^2 in the product, each binomial must start with an *x*.

$$\begin{array}{l} x^2+5x+6 \\ (x\quad)(x\quad) \end{array} \tag{7.2.1}$$

The *last term* in the trinomial came from multiplying the last term in each binomial. So the last terms must multiply to 6.

What two numbers multiply to 6?

The factors of 6 could be 1 and 6, or 2 and 3. How do you know which pair to use?

Consider the *middle term*. It came from adding the outer and inner terms.

So the numbers that must have a product of 6 will need a sum of 5. We'll test both possibilities and summarize the results in Table 7.2.1—the table will be very helpful when you work with numbers that can be factored in many different ways.

Factors of 6	Sum of factors
1,6	$1+6=7$
2,3	$2+3=5$

Table 7.2.1

We see that 2 and 3 are the numbers that multiply to 6 and add to 5. So we have the factors of x^2+5x+6. They are $(x+2)(x+3)$.

$$\begin{array}{ll} x^2+5x+6 & \text{product} \\ (x+2)(x+3) & \text{factors} \end{array} \tag{7.2.2}$$

You should check this by multiplying.

Looking back, we started with x^2+5x+6, which is of the form x^2+bx+c, where b=5 and c=6. We factored it into two binomials of the form (x+m) and (x+n).

$$\begin{array}{ll} x^2+5x+6 & x^2+bx+c \\ (x+2)(x+3) & (x+m)(x+n) \end{array} \tag{7.2.3}$$

To get the correct factors, we found two numbers *m* and *n* whose product is *c* and sum is *b*.

Example 7.2.1: HOW TO FACTOR TRINOMIALS OF THE FORM x^2+bx+c

Factor: $x^2+7x+12$

Solution

Step 1. Write the factors as two binomials with first terms *x*.	Write two sets of parentheses and put *x* as the first term.	$x^2+7x+12$ $(x\quad)(x\quad)$
Step 2. Find two numbers *m and n* that multiply to *c*, $m \cdot n = c$ add to *b*, $m+n=b$	Find two numbers that multiply to 12 and add to 7. **Factors of 12** / **Sum of factors** 1, 12 / 1 + 12 = 13 2, 6 / 2 + 6 = 8 3, 4 / 3 + 4 = 7*	
Step 3. Use *m and n* as the last terms of the factors.	Use 3 and 4 as the last terms of the binomials.	$(x+3)(x+4)$
Step 4. Check by multiplying the factors.		$(x+3)(x+4)$ $x^2+4x+3x+12$ $x^2+7x+12$ ✓

Try It 7.2.2

Factor: x^2+6x+8

Try It 7.2.3

Factor: $y^2+8y+15$

Let's summarize the steps we used to find the factors.

HOW TO

Factor trinomials of the form x^2+bx+c.

Step 1. Write the factors as two binomials with first terms *x*: $(x\quad)(x\quad)$

Step 2. Find two numbers *m* and *n* that
Multiply to *c*, $m \cdot n = c$
Add to *b*, $m+n=b$

Step 3. Use *m* and *n* as the last terms of the factors: $(x+m)(x+n)$

Step 4. Check by multiplying the factors.

Example 7.2.4

Factor: $u^2+11u+24$

Solution

Notice that the variable is *u*, so the factors will have first terms *u*.

	$u^2+11u+24$
Write the factors as two binomials with first terms u .	$(u\quad)(u\quad)$
Find two numbers that: multiply to 24 and add to 11.	

Factors of 24	Sum of factors
1,24	1+24=25
2,12	2+12=14
3,8	3+8=11*
4,6	4+6=10

Use 3 and 8 as the last terms of the binomials. $(u+3)(u+8)$

Check.

$(u+3)(u+8)$
$u^2+3u+8u+24$
$u^2+11u+24v$✓

Try It 7.2.5

Factor: $q^2+10q+24$

Try It 7.2.6

Factor: $t^2+14t+24$

Example 7.2.7

Factor: $y^2+17y+60$

Solution

	$y^2+17y+60$
Write the factors as two binomials with first terms y.	$(y\quad)(y\quad)$
Find two numbers that multiply to 60 and add to 17.	

Factors of 60	Sum of factors
1,60	1+60=61
2,30	2+30=32
3,20	3+20=23
4,15	4+15=19

Factors of 60	Sum of factors
5,12	5+12=17*
6,10	

Use 5 and 12 as the last terms. $(y+5)(y+12)$
Check.

$(y+5)(y+12)$
$\left(y^2+12y+5y+60\right)$
$\left(y^2+17y+60\right)\checkmark$

Try It 7.2.8

Factor: $x^2+19x+60$

Try It 7.2.9

Factor: $v^2+23v+60$

Factor Trinomials of the Form $x^2 + bx + c$ with b Negative, c Positive

In the examples so far, all terms in the trinomial were positive. What happens when there are negative terms? Well, it depends which term is negative. Let's look first at trinomials with only the middle term negative.

Remember: To get a negative sum and a positive product, the numbers must both be negative.

Again, think about FOIL and where each term in the trinomial came from. Just as before,

- the first term, x^2, comes from the product of the two first terms in each binomial factor, *x* and *y*;
- the positive last term is the product of the two last terms
- the negative middle term is the sum of the outer and inner terms.

How do you get a *positive product* and a *negative sum*? With two negative numbers.

Example 7.2.10

Factor: $t^2-11t+28$

Solution

Again, with the positive last term, 28, and the negative middle term, −11t, we need two negative factors. Find two numbers that multiply 28 and add to −11.

Write the factors as two binomials with first terms t $t^2-11t+28$ $(t\quad)(t\quad)$

Find two numbers that: multiply to 28 and add to −11.

Factors of 28	Sum of factors
−1,−28	−1+(−28)=−29
−2,−14	−2+(−14)=−16
−4,−7	$-4+(-7)=-11^*$

Use $-4, -7$ as the last terms of the binomials. $(t-4)(t-7)$
Check.

$(t-4)(t-7)$
$t^2-7t-4t+28$
$t^2-11t+28$✓

Try It 7.2.11

Factor: $u^2-9u+18$

Try It 7.2.12

Factor: $y^2-16y+63$

Factor Trinomials of the Form x2+bx+c with *c* Negative

Now, what if the last term in the trinomial is negative? Think about FOIL. The last term is the product of the last terms in the two binomials. A negative product results from multiplying two numbers with opposite signs. You have to be very careful to choose factors to make sure you get the correct sign for the middle term, too.

Remember: To get a negative product, the numbers must have different signs.

Example 7.2.13

Factor: z^2+4z-5

Solution

To get a negative last term, multiply one positive and one negative. We need factors of −5 that add to positive 4.

Factors of −5	Sum of factors
1,−5	1+(−5)=−4
−1,5	−1+5=4*

Notice: We listed both 1,−5 and −1,5 to make sure we got the sign of the middle term correct.

	z^2+4z-5
Factors will be two binomials with first terms z.	$(z\quad)(z\quad)$
Use $-1, 5$ as the last terms of the binomials.	$(z-1)(z+5)$
Check.	

$(z-1)(z+5)$
$z^2+5z-1z-5$
z^2+4z-5✓

Try It 7.2.14

Factor: $h^2+4h-12$

Answer

$(h-2)(h+6)$

Try It 7.2.15

Factor: : 2^2+k-20

Answer

$(k-4)(k+5)$

Let's make a minor change to the last trinomial and see what effect it has on the factors.

Example 7.2.16

Factor: z^2-4z-5

Solution

Solution

This time, we need factors of −5 that add to −4.

Factors of −5	Sum of factors
1,−5	1+(−5)=−4*
−1,5	−1+5=4

	z^2-4z-5
Factors will be two binomials with first terms z.	$(z\quad)(z\quad)$
Use $1,-5$ as the last terms of the binomials.	$(z+1)(z-5)$
Check.	

$(z+1)(z-5)$
$z^2-5z+1z-5$
z^2-4z-5✓

Try It 7.2.17

Factor: $x^2-4x-12$

Answer

$(x+2)(x-6)$

Try It 7.2.18

Factor: y^2-y-20

Answer

$(y+4)(y-5)$

Example 7.2.19

Factor: $q^2-2q-15$

Solution

Factors will be two binomials with first terms q.	$q^2-2q-15$ $(q\quad)(q\quad)$
You can use $3,-5$ as the last terms of the binomials.	$(q+3)(q-5)$

Factors of −15	Sum of factors
1,−15	1+(−15)=−14
−1,15	−1+15=14
3,−5	3+(−5)=−2*
−3,5	

Check.

$(q+3)(q-5)$
$q^2-5q+3z-15$
$q^2-2q-15$✓

Try It 7.2.20

Factor: $r^2-3r-40$

Answer

$(r+5)(r-8)$

Try It 7.2.21

Factor: $s^2-3s-10$

Answer

$(s+2)(s-5)$

Some trinomials are prime. The only way to be certain a trinomial is prime is to list all the possibilities and show that none of them work.

Example 7.2.22

Factor: $y^2-6y+15$

Solution

Factors will be two binomials with first terms y.	$y^2-6y+15$ $(y\quad)(y\quad)$

Factors of 15	Sum of factors
−1,−15	−1+(−15)=−16
−3,−5	−3+(−5)=−8

As shown in the table, none of the factors add to −6; therefore, the expression is prime.

Try It 7.2.23

Factor: $m^2+4m+18$

Answer

prime

Try It 7.2.24

Factor: $n^2-10n+12$

Answer

prime

Example 7.2.25

Factor: $2x+x^2-48$

Solution

	$2x+x^2-48$
First we put the terms in decreasing degree order.	$x^2+2x-48$
Factors will be two binomials with first terms x .	$(x\quad)(x\quad)$

As shown in the table, you can use $-6,8$ as the last terms of the binomials.

$$(x-6)(x+8) \tag{7.2.4}$$

Factors of -48	Sum of factors
$-1,48$	$-1+48=47$
$-2,24$ $-3,16$ $-4,12$ $-6,8$	$-2+24=22$ $-3+16=13$ $-4+12=8$ $-6+8=2$

Check.

$(x-6)(x+8)$

$x^2-6q+8q-48$

$x^2+2x-48$✓

Try It 7.2.26

Factor: $9m+m^2+18$

Answer

$(m+3)(m+6)$

Try It 7.2.27

Factor: $-7n+12+n^2$

Answer

$(n-3)(n-4)$

Let's summarize the method we just developed to factor trinomials of the form x^2+bx+c

Note

When we factor a trinomial, we look at the signs of its terms first to determine the signs of the binomial factors.

$$\begin{array}{c} x^2+bx+c \\ (x+m)(x+n) \end{array} \tag{7.2.5}$$

When *c* is positive, *m* and *n* have the same sign.

$$\begin{array}{cc} \text{b positive} & \text{b negative} \\ m, n \text{ positive} & m, n \text{ negative} \\ x^2+5x+6 & x^2-6x+8 \\ (x+2)(x+3) & (x-4)(x-2) \\ \text{same signs} & \text{same signs} \end{array} \tag{7.2.6}$$

When *c* is negative, *m* and *n* have opposite signs.

$$\begin{array}{cc} x^2+x-12 & x^2-2x-15 \\ (x+4)(x-3) & (x-5)(x+3) \\ \text{opposite signs} & \text{opposite signs} \end{array} \tag{7.2.7}$$

Notice that, in the case when *m* and *n* have opposite signs, the sign of the one with the larger absolute value matches the sign of *b*.

Factor Trinomials of the Form $x^2 + bxy + cy^2$

Sometimes you'll need to factor trinomials of the form $x^2+bxy+cy^2$ with two variables, such as $x^2+12xy+36y^2$. The first term, x^2, is the product of the first terms of the binomial factors, $x \cdot x$. The y^2 in the last term means that the second terms of the binomial factors must each contain *y*. To get the coefficients *b* and *c*, you use the same process summarized in the previous objective.

Example 7.2.28

Factor: $x^2+12xy+36y^2$

Solution

	$x^2+12xy+36y^2$
Note that the first terms are x, last terms contain y	$(x\_y)\,(x\_y)$

Find the numbers that multiply to 36 and add to 12.

Factors of 36	Sum of factors
1, 36	1+36=37
2, 18	2+18=20
3, 12	3+12=15
4, 9	4+9=13
6, 6	6+6=12*

Use 6 and 6 as the coefficients of the last terms.	$(x+6y)(x+6y)$
Check your answer.	

$(x+6y)(x+6y)$
$x^2+6xy+6xy+36y^2$
$x^2+12xy+36y^2$ ✓

Try It 7.2.29

Factor: $u^2+11uv+28v^2$

Answer

$(u+4v)(u+7v)$

Try It 7.2.30

Factor: $x^2+13xy+42y^2$

Answer

$(x+6y)(x+7y)$

Example 7.2.31

Factor: $r^2-8rs-9s^2$

Solution

We need r in the first term of each binomial and s in the second term. The last term of the trinomial is negative, so the factors must have opposite signs.

	$r^2-8rs-9s^2$
Note that the first terms are r, last terms contain s	$(r\_s)\,(r\_s)$

Factors of −9	Sum of factors
1,−9	1+(−9)=−8*
−1,9	−1+9=8
3,−3	3+(−3)=0

Use $1,-9$ as coefficients of the last terms.	$(r+s)(r-9s)$

Check your answer.

$(r-9s)(r+s)$
$r^2+rs-9rs-9s^2$
$r^2-8rs-9s^2$ ✓

Try It 7.2.32

Factor: $a^2-11ab+10b^2$

Answer

$(a-b)(a-10b)$

Try It 7.2.33

Factor: $m^2 - 13mn + 12n^2$

Answer

$(m - n)(m - 12n)$

Example 7.2.34

Factor: $u^2 - 9uv - 12v^2$

Solution

We need *u* in the first term of each binomial and v in the second term. The last term of the trinomial is negative, so the factors must have opposite signs.

$$u^2 - 9uv - 12v^2$$

Note that the first terms are u, last terms contain v $\quad (u\_v)\,(u\_v)$

Find the numbers that multiply to −12 and add to −9.

Factors of −12	Sum of factors
1,−12	1+(−12)=−11
−1,12	−1+12=11
2,−6	2+(−6)=−4
−2,6	−2+6=4
3,−4	3+(−4)=−1
−3,4	−3+4=1

Note there are no factor pairs that give us −9 as a sum. The trinomial is prime.

Try It 7.2.35

Factor: $x^2 - 7xy - 10y^2$

Answer

prime

Try It 7.2.36

Factor: $p^2 + 15pq + 20q^2$

Answer

prime

Key Concepts

- **Factor trinomials of the form $x^2 + bx + c$**
  1. Write the factors as two binomials with first terms x: $(x \quad)(x \quad)$
  2. Find two numbers m and n that
     Multiply to c, $m \cdot n = c$
     Add to b, $m + n = b$

3. Use m and n as the last terms of the factors: $(x+m)(x+n)$.
4. Check by multiplying the factors.

This page titled 7.2: Factor Quadratic Trinomials with Leading Coefficient 1 is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.2: Factor Quadratic Trinomials with Leading Coefficient 1** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.2E: Exercises

Practice Makes Perfect

Factor Trinomials of the Form x^2+bx+c

In the following exercises, factor each trinomial of the form x^2+bx+c

Exercise 1

x^2+4x+3

Answer

$(x+1)(x+3)$

Exercise 2

y^2+8y+7

Exercise 3

$m^2+12m+11$

Answer

$(m+1)(m+11)$

Exercise 4

$b^2+14b+13$

Exercise 5

$a^2+9a+20$

Answer

$(a+4)(a+5)$

Exercise 6

$m^2+7m+12$

Exercise 7

$p^2+11p+30$

Answer

$(p+5)(p+6)$

Exercise 8

$w^2+10w+21$

Exercise 9

$n^2+19n+48$

Answer

$(n+3)(n+16)$

Exercise 10

$b^2+14b+48$

Exercise 11

$a^2+25a+100$

Answer

$(a+5)(a+20)$

Exercise 12

$u^2+101u+100$

Exercise 13

$x^2-8x+12$

Answer

$(x-2)(x-6)$

Exercise 14

$q^2-13q+36$

Exercise 15

$y^2-18y+45$

Answer

$(y-3)(y-15)$

Exercise 16

$m^2-13m+30$

Exercise 17

x^2-8x+7

Answer

$(x-1)(x-7)$

Exercise 18

$y^2 - 5y + 6$

Exercise 19

$p^2 + 5p - 6$

Answer

$(p-1)(p+6)$

Exercise 20

$n^2 + 6n - 7$

Exercise 21

$y^2 - 6y - 7$

Answer

$(y+1)(y-7)$

Exercise 22

$v^2 - 2v - 3$

Exercise 23

$x^2 - x - 12$

Answer

$(x-4)(x+3)$

Exercise 24

$r^2 - 2r - 8$

Exercise 25

$a^2 - 3a - 28$

Answer

$(a-7)(a+4)$

Exercise 26

$b^2 - 13b - 30$

https://math.libretexts.org/@go/page/30552

Exercise 27

$w^2 - 5w - 36$

Answer

$(w-9)(w+4)$

Exercise 28

$t^2 - 3t - 54$

Exercise 29

$x^2 + x + 5$

Answer

prime

Exercise 30

$x^2 - 3x - 9$

Exercise 31

$8 - 6x + x^2$

Answer

$(x-4)(x-2)$

Exercise 32

$7x + x^2 + 6$

Exercise 33

$x^2 - 12 - 11x$

Answer

$(x-12)(x+1)$

Exercise 34

$-11 - 10x + x^2$

Factor Trinomials of the Form $x^2 + bxy + cy^2$

In the following exercises, factor each trinomial of the form $x^2 + bxy + cy^2$

Exercise 33.2

$p^2 + 3pq + 2q^2$

Answer

$(p+q)(p+2q)$

Exercise 34.2

$m^2+6mn+5n^2$

Exercise 35

$r^2+15rs+36s^2$

Answer

$(r+3s)(r+12s)$

Exercise 36

$u^2+10uv+24v^2$

Exercise 37

$m^2-12mn+20n^2$

Answer

$(m-2n)(m-10n)$

Exercise 38

$p^2-16pq+63q^2$

Exercise 39

$x^2-2xy-80y^2$

Answer

$(x+8y)(x-10y)$

Exercise 40

$p^2-8pq-65q^2$

Exercise 41

$m^2-64mn-65n^2$

Answer

$(m+n)(m-65n)$

Exercise 42

$p^2-2pq-35q^2$

https://math.libretexts.org/@go/page/30552

Exercise 43

$a^2+5ab-24b^2$

Answer

$(a+8b)(a-3b)$

Exercise 44

$r^2+3rs-28s^2$

Exercise 45

$x^2-3xy-14y^2$

Answer

prime

Exercise 46

$u^2-8uv-24v^2$

Exercise 47

$m^2-5mn+30n^2$

Answer

prime

Exercise 48

$c^2-7cd+18d^2$

Mixed Practice

In the following exercises, factor each expression.

Exercise 49

$u^2-12u+36$

Answer

$(u-6)(u-6)$

Exercise 50

$w^2+4w-32$

Exercise 51

$x^2-14x-32$

Answer

$(x+2)(x-16)$

Exercise 52

$y^2+41y+40$

Exercise 53

$r^2-20rs+64s^2$

Answer

$(r-4s)(r-16s)$

Exercise 54

$x^2-16xy+64y^2$

Exercise 55

$k^2+34k+120$

Answer

$(k+4)(k+30)$

Exercise 56

$m^2+29m+120$

Exercise 57

$y^2+10y+15$

Answer

prime

Exercise 58

$z^2-3z+28$

Exercise 59

$m^2+mn-56n^2$

Answer

$(m+8n)(m-7n)$

Exercise 60

$q^2-29qr-96r^2$

Exercise 61

$u^2 - 17uv + 30v^2$

Answer

$(u - 15v)(u - 2v)$

Exercise 62

$m^2 - 31mn + 30n^2$

Exercise 63

$c^2 - 8cd + 26d^2$

Answer

prime

Exercise 64

$r^2 + 11rs + 36s^2$

Everyday Math

Exercise 65

Consecutive integers Deirdre is thinking of two consecutive integers whose product is 56. The trinomial $x^2 + x - 56$ describes how these numbers are related. Factor the trinomial.

Answer

$(x + 8)(x - 7)$

Exercise 66

Consecutive integers Deshawn is thinking of two consecutive integers whose product is 182. The trinomial $x^2 + x - 182$ describes how these numbers are related. Factor the trinomial describes how these numbers are related. Factor the trinomial.

Writing Exercises

Exercise 67

Many trinomials of the form $x^2 + bx + c$ factor into the product of two binomials $(x + m)(x + n)$. Explain how you find the values of m and n.

Answer

Answers may vary

Exercise 68

How do you determine whether to use plus or minus signs in the binomial factors of a trinomial of the form x^2+bx+c where b and c may be positive or negative numbers?

Exercise 69

Will factored x^2-x-20 as $(x+5)(x-4)$. Bill factored it as $(x+4)(x-5)$. Phil factored it as $(x-5)(x-4)$. Who is correct? Explain why the other two are wrong.

Answer

Answers may vary

Exercise 70

Look at Example, where we factored $y^2+17y+60$. We made a table listing all pairs of factors of 60 and their sums. Do you find this kind of table helpful? Why or why not?

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
factor trinomials of the form x^2+bx+c.			
factor trinomials of the form $x^2+bxy+cy^2$.			

b. After reviewing this checklist, what will you do to become confident for all goals?

This page titled 7.2E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.2E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.3: Factor Quadratic Trinomials with Leading Coefficient Other than 1

Learning Objectives

By the end of this section, you will be able to:

- Recognize a preliminary strategy to factor polynomials completely
- Factor trinomials of the form ax^2+bx+c with a GCF
- Factor trinomials using trial and error
- Factor trinomials using the 'ac' method

Note

Before you get started, take this readiness quiz.

1. Find the GCF of $45p^2$ and $30p^6$
   If you missed this problem, review Example 7.1.4.
2. Multiply $(3y+4)(2y+5)$
   If you missed this problem, review Example 6.3.37.
3. Combine like terms $12x^2+3x+5x+9$
   If you missed this problem, review Example 1.3.37.

Recognize a Preliminary Strategy for Factoring

Let's summarize where we are so far with factoring polynomials. In the first two sections of this chapter, we used three methods of factoring: factoring the GCF, factoring by grouping, and factoring a trinomial by "undoing" FOIL. More methods will follow as you continue in this chapter, as well as later in your studies of algebra.


How will you know when to use each factoring method? As you learn more methods of factoring, how will you know when to apply each method and not get them confused? It will help to organize the factoring methods into a strategy that can guide you to use the correct method.

As you start to factor a polynomial, always ask first, "Is there a greatest common factor?" If there is, factor it first.

The next thing to consider is the type of polynomial. How many terms does it have? Is it a binomial? A trinomial? Or does it have more than three terms?

- If it is a trinomial where the leading coefficient is one, x^2+bx+c, use the "undo FOIL" method.
- If it has more than three terms, try the grouping method. This is the only method to use for polynomials of more than three terms.

Some polynomials cannot be factored. They are called "prime." Below we summarize the methods we have so far.



CHOOSE A STRATEGY TO FACTOR POLYNOMIALS COMPLETELY.

1. Is there a greatest common factor?
   - Factor it out.
2. Is the polynomial a binomial, trinomial, or are there more than three terms?
   - If it is a binomial, right now we have no method to factor it.
   - If it is a trinomial of the form x^2+bx+c: Undo FOIL $(x\quad)(x\quad)$
   - If it has more than three terms: Use the grouping method.

3. Check by multiplying the factors.

Use the preliminary strategy to completely factor a polynomial. A polynomial is factored completely if, other than monomials, all of its factors are prime.

Example 7.3.1

Identify the best method to use to factor each polynomial.

a. $6y^2-72$
b. $r^2-10r-24$
c. $p^2+5p+pq+5q$

Answer a

	$6y^2-72$
Is there a greatest common factor?	Yes, 6.
Factor out the 6	$6\left(y^2-12\right)$
Is it a binomial, trinomial, or are there more than 3 terms?	Binomial, we have no method to factor binomials yet.

Answer b

	$r^2-10r-24$
Is there a greatest common factor?	No, there is no common factor.
Is it a binomial, trinomial, or are there more than three terms?	Trinomial, with leading coefficient 1, so "undo" FOIL.

Answer c

	$p^2+5p+pq+5q$
Is there a greatest common factor?	No, there is no common factor.
Is it a binomial, trinomial, or are there more than three terms?	More than three terms, so factor using grouping.

Try It 7.3.2

Identify the best method to use to factor each polynomial:

a. $4y^2+32$
b. $y^2+10y+21$
c. $yz+2y+3z+6$

Answer a

no method

Answer b

undo using FOIL

Answer c

factor with grouping

Try It 7.3.3

Identify the best method to use to factor each polynomial:

a. $ab+a+4b+4$

b. $3k^2+15$
c. p^2+9p+8

Answer a

factor using grouping

Answer b

no method

Answer c

undo using FOIL

Factor Trinomials of the form $ax^2 + bx + c$ with a GCF

Now that we have organized what we've covered so far, we are ready to factor trinomials whose leading coefficient is not 1, trinomials of the form ax^2+bx+c . Remember to always check for a GCF first! Sometimes, after you factor the GCF, the leading coefficient of the trinomial becomes 1 and you can factor it by the methods in the last section. Let's do a few examples to see how this works. Watch out for the signs in the next two examples.

Example 7.3.4

Factor completely: $2n^2-8n-42$.

Solution

Use the preliminary strategy.

Is there a greatest common factor?	$2n^2-8n-42$
Yes, GCF $=2$. Factor it out.	$2\left(n^2-4n-21\right)$
Inside the parentheses, is it a binomial, trinomial, or are there more than three terms?	
It is a trinomial whose coefficient is 1, so undo FOIL.	$2(n\quad)(n\quad)$
Use 3 and -7 as the last terms of the binomials.	$2(n+3)(n-7)$

Factors of −21	Sum of factors
1,−21	1+(−21)=−20
3,−7	3+(−7)=−4*

Check.
$2(n+3)(n-7)$
$2\left(n^2-7n+3n-21\right)$
$2\left(n^2-4n-21\right)$
$2n^2-8n-42$✓

Try It 7.3.5

Factor completely: $4m^2-4m-8$

Answer

$4(m+1)(m-2)$

Try It 7.3.6

Factor completely: $5k^2-15k-50$

Answer

$5(k+2)(k-5)$

Example 7.3.7

Factor completely: $4y^2-36y+56$

Solution

Use the preliminary strategy.

Is there a greatest common factor?	$4y^2-36y+56$
Yes, GCF $=4$. Factor it out.	$4\left(y^2-9y+14\right)$
Inside the parentheses, is it a binomial, trinomial, or are there more than three terms?	
It is a trinomial whose coefficient is 1, so undo FOIL.	$4(y\quad)(y\quad)$
Use a table like the one below to find two numbers that multiply to 14 and add to -9	
Both factors of 14 must be negative.	$4(y-2)(y-7)$

Factors of 14	Sum of factors
−1,−14	−1+(−14)=−15
−2,−7	−2+(−7)=−9*

Check.

$4(y-2)(y-7)$

$4\left(y^2-7y-2y+14\right)$

$4\left(y^2-9y+14\right)$

$4y^2-36y+42$✓

Try It 7.3.8

Factor completely: $3r^2-9r+6$

Answer

$3(r-1)(r-2)$

Try It 7.3.9

Factor completely: $2t^2-10t+12$

Answer

$2(t-2)(t-3)$

In the next example the GCF will include a variable.

Example 7.3.10

Factor completely: $4u^3+16u^2-20u$

Solution

Use the preliminary strategy.

Is there a greatest common factor?	$4u^3+16u^2-20u$
Yes, GCF $=4u$. Factor it.	$4u\left(u^2+4u-5\right)$
Binomial, trinomial, or more than three terms? more than three terms?	
It is a trinomial. So "undo FOIL."	$4u(u\quad)(u\quad)$
Use a table like the table below to find two numbers that multiply to -5 and add to 4	$4u(u-1)(u+5)$

Factors of −5	Sum of factors
−1,5	−1+5=4*
1,−5	1+(−5)=−4

Check.

$4u(u-1)(u+5)$

$4u\left(u^2+5u-u-5\right)$

$4u\left(u^2+4u-5\right)$

$4u^3+16u^2-20u$✓

Try It 7.3.11

Factor completely: $5x^3+15x^2-20x$

Answer

$5x(x-1)(x+4)$

Try It 7.3.12

Factor completely: $6y^3+18y^2-60y$

Answer

$6y(y-2)(y+5)$

Factor Trinomials using Trial and Error

What happens when the leading coefficient is not 1 and there is no GCF? There are several methods that can be used to factor these trinomials. First we will use the Trial and Error method.

Let's factor the trinomial $3x^2+5x+2$

From our earlier work we expect this will factor into two binomials.

$$\begin{array}{c} 3x^2+5x+2 \\ (\quad)(\quad) \end{array} \tag{7.3.1}$$

We know the first terms of the binomial factors will multiply to give us $3x^2$. The only factors of $3x^2$ are $1x, 3x$. We can place them in the binomials.

$3x^2 + 5x + 2$

$1x, 3x$

$(x \quad)(3x \quad)$

Check. Does $1x \cdot 3x = 3x^2$?

We know the last terms of the binomials will multiply to 2. Since this trinomial has all positive terms, we only need to consider positive factors. The only factors of 2 are 1 and 2. But we now have two cases to consider as it will make a difference if we write 1, 2, or 2, 1.

$3x^2 + 5x + 2$ $\quad$ $3x^2 + 5x + 2$

$1x, 3x \quad 1, 2$ $\quad$ $1x, 3x \quad 1, 2$

$(x + 1)(3x + 2)$ or $(x + 2)(3x + 1)$

Which factors are correct? To decide that, we multiply the inner and outer terms.

$3x^2 + 5x + 2$ $\quad$ $3x^2 + 5x + 2$

$1x, 3x \quad 1, 2$ $\quad$ $1x, 3x \quad 1, 2$

$(x + 1)(3x + 2)$ or $(x + 2)(3x + 1)$

$3x$, $2x$, $5x$ $\quad$ $6x$, $1x$, $7x$

Since the middle term of the trinomial is $5x$, the factors in the first case will work. Let's FOIL to check.

$$\begin{aligned} &(x+1)(3x+2) \\ &3x^2+2x+3x+2 \\ &3x^2+5x+2\checkmark \end{aligned} \tag{7.3.2}$$

Our result of the factoring is:

$$\begin{aligned} &3x^2+5x+2 \\ &(x+1)(3x+2) \end{aligned} \tag{7.3.3}$$

Example 7.3.13: How to Factor Trinomials of the Form $ax^2 + bx + c$ Using Trial and Error

Factor completely: $3y^2+22y+7$

Solution

Step 1. Write the trinomial in descending order.	The trinomial is already in descending order.	$3y^2 + 22y + 7$
Step 2. Find all the factor pairs of the first term.	The only factors of $3y^2$ are $1y$, $3y$ Since there is only one pair, we can put them in the parentheses.	$3y^2 + 22y + 7$ $1y, 3y$ $3y^2 + 22y + 7$ $1y, 3y$ $(y \quad)(3y \quad)$

Step 3. Find all the factor pairs of the third term.	The only factors of 7 are 1, 7.	$3y^2 + 22y + 7$ 1y, 3y 1, 7 $(y \quad)(3y \quad)$
Step 4. Test all the possible combinations of the factors until the correct product is found.	$3y^2 + 22y + 7$ 1y, 3y 1, 7 $(y + 1)(3y + 7)$ $3y$ $7y$ $10y$ No. We need $22y$ $3y^2 + 22y + 7$ 1y, 3y 1, 7 $(y + 7)(3y + 1)$ $21y$ y $22y$	$3y^2 + 22y + 7$ Possible factors: $(y + 1)(3y + 7)$ — Product: $3y^2 + 10y + 7$ $(y + 7)(3y + 1)$ — $3y^2 + 22y + 7$ $(y + 7)(3y + 1)$
Step 5. Check by multiplying.		$(y + 7)(3y + 1)$ $3y^2 + 22y + 7$ ✓

Try It 7.3.14

Factor completely: $2a^2 + 5a + 3$

Answer

$(a + 1)(2a + 3)$

Try It 7.3.15

Factor completely: $4b^2 + 5b + 1$

Answer

$(b + 1)(4b + 1)$

FACTOR TRINOMIALS OF THE FORM $ax^2 + bx + c$ USING TRIAL AND ERROR.

1. Write the trinomial in descending order of degrees.
2. Find all the factor pairs of the first term.
3. Find all the factor pairs of the third term.
4. Test all the possible combinations of the factors until the correct product is found.
5. Check by multiplying.

When the middle term is negative and the last term is positive, the signs in the binomials must both be negative.

Example 7.3.16

Factor completely: $6b^2 - 13b + 5$

Solution

The trinomial is already in descending order.	$6b^2 - 13b + 5$
Find the factors of the first term.	$6b^2 - 13b + 5$ $1b \cdot 6b$ $2b \cdot 3b$

Find the factors of the last term. Consider the signs. Since the last term, 5 is positive its factors must both be positive or both be negative. The coefficient of the middle term is negative, so we use the negative factors.	$6b^2 - 13b + 5$ $1b \cdot 6b$ $\quad -1, -5$ $2b \cdot 3b$

Consider all the combinations of factors.

$6b^2 - 13b + 5$

Possible factors	Product
(b−1)(6b−5)	$6b^2 - 11b + 5$
(b−5)(6b−1)	$6b^2 - 31b + 5$
(2b−1)(3b−5)	$6b^2 - 13b + 5$ *
(2b−5)(3b−1)	$6b^2 - 17b + 5$

The correct factors are those whose product is the original trinomial. $(2b - 1)(3b - 5)$

Check by multiplying.

$(2b - 1)(3b - 5)$
$6b^2 - 10b - 3b + 5$
$6b^2 - 13b + 5v$✓

? Try It 7.3.17

Factor completely: $8x^2 - 14x + 3$

Answer

$(2x - 3)(4x - 1)$

? Try It 7.3.18

Factor completely: $10y^2 - 37y + 7$

Answer

$(2y - 7)(5y - 1)$

When we factor an expression, we always look for a greatest common factor first. If the expression does not have a greatest common factor, there cannot be one in its factors either. This may help us eliminate some of the possible factor combinations.

✔ Example 7.3.19

Factor completely: $14x^2 - 47x - 7$

Solution

The trinomial is already in descending order.	$14x^2 - 47x - 7$
Find the factors of the first term.	$14x^2 - 47x - 7$ $1x \cdot 14x$ $2x \cdot 7x$

Find the factors of the last term. Consider the signs. Since it is negative, one factor must be positive and one negative.	$14x^2 - 47x - 7$ $1x \cdot 14x$ $\quad$ $1, -7$ $2x \cdot 7x$ $\quad$ $-1, 7$

Consider all the combinations of factors. We use each pair of the factors of $14x^2$ with each pair of factors of -7.

Factors of $14x^2$	Pair with	Factors of -7
$x, 14x$		$11, -7$ $-7, 11$ (reverse order)
$x, 14x$		$-1, 77$ $77, -1$ (reverse order)
$2x, 7x$		$11, -7$ $-7, 11$ (reverse order)
$2x, 7x$		$-1, 77$ $77, -1$ (reverse order)

These pairings lead to the following eight combinations.

$14x^2 - 47x - 7$	
Possible factors	**Product**
$(x + 1)(14x - 7)$	Not an option
$(x - 7)(14x + 1)$	$14x^2 - 97x - 7$
$(x - 1)(14x + 7)$	Not an option
$(x + 7)(14x - 1)$	$14x^2 + 97x - 7$
$(2x + 1)(7x - 7)$	Not an option
$(2x - 7)(7x + 1)$	$14x^2 - 47x - 7$*
$(2x - 1)(7x + 7)$	Not an option
$(2x + 7)(7x - 1)$	$14x^2 + 47x - 7$

If the trinomial has no common factors, then neither factor can contain a common factor. That means each of these combinations is not an option.

The correct factors are those whose product is the original trinomial. $(2x - 7)(7x + 1)$

Check by multiplying.

$(2b - 1)(3b - 5)$
$6b^2 - 10b - 3b + 5$
$6b^2 - 13b + 5$✓

? Try It 7.3.20

Factor completely: $8a^2 - 3a - 5$

Answer

$(a - 1)(8a + 5)$

? Try It 7.3.21

Factor completely: $6b^2 - b - 15$

Answer

$(2b + 3)(3b - 5)$

Example 7.3.22

Factor completely: $18n^2 - 37n + 15$

Solution

The trinomial is already in descending order.	$18n^2 - 37n + 15$
Find the factors of the first term.	$18n^2 - 37n + 15$ $1n \cdot 18n$ $2n \cdot 9n$ $3n \cdot 6n$
Find the factors of the last term. Consider the signs. Since 15 is positive and the coefficient of the middle term is negative, we use the negative factors.	$18n^2 - 37n + 15$ $1n \cdot 18n$ $\quad -1(-15)$ $2n \cdot 9n$ $\quad -3(-5)$ $3n \cdot 6n$

Consider all the combinations of factors.

$18n^2 - 37n + 15$	
Possible factors	**Product**
$(n-1)(18n-15)$	Not an option
$(n-15)(18n-1)$	$18n^2 - 271n + 15$
$(n-3)(18n-5)$	$18n^2 - 59n + 15$
$(n-5)(18n-3)$	Not an option
$(2n-1)(9n-15)$	Not an option
$(2n-15)(9n-1)$	$18n^2 - 137n + 15$
$(2n-3)(9n-5)$	$18n^2 - 37n + 15$*
$(2n-5)(9n-3)$	Not an option
$(3n-1)(6n-15)$	Not an option
$(3n-15)(6n-1)$	Not an option
$(3n-3)(6n-5)$	Not an option
$(3n-5)(6n-3)$	Not an option

If the trinomial has no common factors, then neither factor can contain a common factor. That means this combination is not an option.

The correct factors are those whose product is the original trinomial. $(2n-3)(9n-5)$

Check by multiplying.

$(2n-3)(9n-5)$
$18n^2 - 10n - 27n + 15$
$18n^2 - 37n + 15$✓

Try It 7.3.23

Factor completely: $18x^2 - 3x - 10$

Answer

$(3x+2)(6x-5)$

Try It 7.3.24

Factor completely: $30y^2 - 53y - 21$

Answer

$(3y+1)(10y-21)$

Don’t forget to look for a GCF first.

Example 7.3.25

Factor completely: $10y^4+55y^3+60y^2$

Solution

	$10y^4+55y^3+60y^2$
Notice the greatest common factor, and factor it first.	$5y^2\left(2y^2+11y+12\right)$
Factor the trinomial.	$5y^2(2y^2+11y+12)$ $y\cdot 2y$ $1\cdot 12$ $2\cdot 6$ $3\cdot 4$

Consider all the combinations.

$2y^2+11y+12$	
Possible factors	**Product**
$(y+1)(2y+12)$	Not an option
$(y+12)(2y+1)$	$2y^2+25y+12$
$(y+2)(2y+6)$	Not an option
$(y+6)(2y+2)$	Not an option
$(y+3)(2y+4)$	Not an option
$(y+4)(2y+3)$	$2y^2+11y+12$*

If the trinomial has no common factors, then neither factor can contain a common factor. That means this combination is not an option.

The correct factors are those whose product is the original trinomial. Remember to include the factor $5y^2$

$5y^2(y+4)(2y+3)$

Check by multiplying.

$5y^2(y+4)(2y+3)$
$5y^2\left(2y^2+8y+3y+12\right)$
$10y^4+55y^3+60y^2$✓

Try It 7.3.26

Factor completely: $15n^3-85n^2+100n$

Answer

$5n(n-4)(3n-5)$

Try It 7.3.27

Factor completely: $56q^3+320q^2-96q$

Answer

$8q(q+6)(7q-2)$

Factor Trinomials using the “ac” Method

Another way to factor trinomials of the form ax^2+bx+c is the “ac” method. (The “ac” method is sometimes called the grouping method.) The “ac” method is actually an extension of the methods you used in the last section to factor trinomials with leading coefficient one. This method is very structured (that is step-by-step), and it always works!

Example 7.3.28: How to Factor Trinomials Using the “ac” Method

Factor: $6x^2+7x+2$

Solution

Step 1. Factor any GCF.	Is there a greatest common factor? No.	$6x^2+7x+2$
Step 2. Find the product ac.	$a \cdot c$ $6 \cdot 2$ 12	ax^2+bx+c $6x^2+7x+2$
Step 3. Find two numbers m and n that: Multiply to ac $m \cdot n = a \cdot c$ Add to b $m+n=b$	Find two numbers that multiply to 12 and add to 7. Both factors must be positive. $3 \cdot 4 = 12$ $3+4=7$	
Step 4. Split the middle term using m, and n ax^2+bx+c $ax^2+\underbrace{mx+nx}_{bx}+c$	Rewrite $7x$ as $3x+4x$. Notice that $6x^2+3x+4x+2$ is equal to $6x^2+7x+2$. We just split the middle term to get a more useful form.	$6x^2+7x+2$ $6x^2+\underbrace{3x+4x}_{7x}+2$
Step 5. Factor by grouping.		$3x(2x+1)+2(2x+1)$ $(2x+1)(3x+2)$
Step 6. Check by multiplying.		$(2x+1)(3x+2)$ $6x^2+4x+3x+2$ $6x^2+7x+2$ ✓

Try It 7.3.29

Factor: $6x^2+13x+2$

Answer

$(x+2)(6x+1)$

Try It 7.3.30

Factor: $4y^2+8y+3$

Answer

$(2y+1)(2y+3)$

FACTOR TRINOMIALS OF THE FORM USING THE “AC” METHOD.

1. Factor any GCF.
2. Find the product ac.
3. Find two numbers m and n that:
   Multiply to ac $m \cdot n = a \cdot c$
   Add to b $m+n=b$

4. Split the middle term using m and n: $ax^2+bx+c \quad bx \quad ax^2+mx+nx+c$
5. Factor by grouping.
6. Check by multiplying the factors.

When the third term of the trinomial is negative, the factors of the third term will have opposite signs.

Example 7.3.31

Factor: $8u^2-17u-21$

Solution

		ax^2+bx+c $8u^2-17u-21$
Is there a greatest common factor? No.		
Find $a \cdot c$	$a \cdot c$	
	$8(-21)$	
	-168	

Find two numbers that multiply to -168 and add to -17. The larger factor must be negative.

Factors of -168	Sum of factors
$1,-168$	$1+(-168)=-167$
$2,-84$	$2+(-84)=-82$
$3,-56$	$3+(-56)=-53$
$4,-42$	$4+(-42)=-38$
$6,-28$	$6+(-28)=-22$
$7,-24$	$7+(-24)=-17$*
$8,-21$	$8+(-21)=-13$

Split the middle term using $7u$ and $-24u$	$8u^2-17u-21$ $\underbrace{8u^2+7u}\ \underbrace{-24u-21}$
Factor by grouping.	$u(8u+7)-3(8u+7)$ $(8u+7)(u-3)$
Check by multiplying.	$(8u+7)(u-3)$ $8u^2-24u+7u-21$ $8u^2-17u-21$✓

Try It 7.3.32

Factor: $20h^2+13h-15$

Answer

$(4h-5)(5h+3)$

Try It 7.3.33

Factor: $6g^2+19g-20$

Answer

$(q+4)(6q-5)$

Example 7.3.34

Factor: $2x^2+6x+5$

Solution

Is there a greatest common factor? No.	ax^2+bx+c $2x^2+6x+5$
Find a·c	a·c
	2(5)
	10

Find two numbers that multiply to 10 and add to 6.

Factors of 10	Sum of factors
1,10	1+10=11
2, 5	2+5=7

There are no factors that multiply to 10 and add to 6. The polynomial is prime.

Try It 7.3.35

Factor: $10t^2+19t-15$

Answer

$(2t+5)(5t-3)$

Try It 7.3.36

Factor: $3u^2+8u+5$

Answer

$(u+1)(3u+5)$

Don’t forget to look for a common factor!

Example 7.3.37

Factor: $10y^2-55y+70$

Solution

Is there a greatest common factor? Yes. The GCF is 5.	$10y^2-55y+70$
Factor it. Be careful to keep the factor of 5 all the way through the solution!	$5(2y^2-11y+14)$

The trinomial inside the parentheses has a leading coefficient that is not 1.	$ax^2 + bx + c$ $5(2y^2 - 11y + 14)$
Factor the trinomial.	$5(y-2)(2y-7)$
Check by multiplying all three factors.	
$5(2y^2 - 2y - 4y + 14)$	
$5(2y^2 - 11y + 14)$	
$10y^2 - 55y + 70$ ✓	

? Try It 7.3.38

Factor: $16x^2 - 32x + 12$

Answer

$4(2x-3)(2x-1)$

? Try It 7.3.39

Factor: $18w^2 - 39w + 18$

Answer

$3(3w-2)(2w-3)$

We can now update the Preliminary Factoring Strategy, as shown in Figure 7.3.1 and detailed in Choose a strategy to factor polynomials completely (updated), to include trinomials of the form $ax^2 + bx + c$. Remember, some polynomials are prime and so they cannot be factored.




Figure 7.3.1

CHOOSE A STRATEGY TO FACTOR POLYNOMIALS COMPLETELY (UPDATED).

1. Is there a greatest common factor?
   - Factor it.
2. Is the polynomial a binomial, trinomial, or are there more than three terms?
   - If it is a binomial, right now we have no method to factor it.
   - If it is a trinomial of the form $x^2 + bx + c$
     Undo FOIL \(x)(x).
   - If it is a trinomial of the form $ax^2 + bx + c$
     Use Trial and Error or the "ac" method.
   - If it has more than three terms
     Use the grouping method.
3. Check by multiplying the factors.

Note

Access these online resources for additional instruction and practice with factoring trinomials of the form ax^2+bx+c

- Factoring Trinomials, a is not 1

Key Concepts

- **Factor Trinomials of the Form ax^2+bx+c using Trial and Error:** See Example.
  1. Write the trinomial in descending order of degrees.
  2. Find all the factor pairs of the first term.
  3. Find all the factor pairs of the third term.
  4. Test all the possible combinations of the factors until the correct product is found.
  5. Check by multiplying.
- **Factor Trinomials of the Form ax^2+bx+c Using the "ac" Method:** See Example.
  1. Factor any GCF.
  2. Find the product ac.
  3. Find two numbers *m* and *n* that: $\begin{array}{ll}\text{Multiply to } ac & m\cdot n=a\cdot c\\ \text{Add to } b & m+n=b\end{array}$
  4. Split the middle term using *m* and *n*:
     ax^2+bx+c
     $ax^2+\underset{bx}{mx+nx}+c$
  5. Factor by grouping.
  6. Check by multiplying the factors.
- **Choose a strategy to factor polynomials completely (updated):**
  1. Is there a greatest common factor? Factor it.
  2. Is the polynomial a binomial, trinomial, or are there more than three terms?
     If it is a binomial, right now we have no method to factor it.
     If it is a trinomial of the form x^2+bx+c
     Undo FOIL $(x\quad)(x\quad)$.
     If it is a trinomial of the form ax^2+bx+c
     Use Trial and Error or the "ac" method.
     If it has more than three terms
     Use the grouping method.
  3. Check by multiplying the factors.

Glossary

prime polynomials

Polynomials that cannot be factored are prime polynomials.

This page titled 7.3: Factor Quadratic Trinomials with Leading Coefficient Other than 1 is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.3: Factor Quadratic Trinomials with Leading Coefficient Other than 1** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.3E: Exercises

Practice Makes Perfect

Recognize a Preliminary Strategy to Factor Polynomials Completely

In the following exercises, identify the best method to use to factor each polynomial.

Exercise 1

a. $10q^2+50$
b. $a^2-5a-14$
c. $uv+2u+3v+6$

Answer

a. factor the GCF, binomial
b. Undo FOIL
c. factor by grouping

Exercise 2

a. $n^2+10n+24$
b. $8u^2+16$
c. $pq+5p+2q+10$

Exercise 3

a. $x^2+4x-21$
b. $ab+10b+4a+40$
c. $6c^2+24$

Answer

a. undo FOIL
b. factor by grouping
c. factor the GCF, binomial

Exercise 4

a. $20x^2+100$
b. $uv+6u+4v+24$
c. $y^2-8y+15$

Factor Trinomials of the form ax^2+bx+c with a GCF

In the following exercises, factor completely.

Exercise 5

$5x^2+35x+30$

Answer

$5(x+1)(x+6)$

Exercise 6

$12s^2+24s+12$

Exercise 7

$2z^2-2z-24$

Answer

$2(z-4)(z+3)$

Exercise 8

$3u^2-12u-36$

Exercise 9

$7v^2-63v+56$

Answer

$7(v-1)(v-8)$

Exercise 10

$5w^2-30w+45$

Exercise 11

p^3-8p^2-20p

Answer

$p(p-10)(p+2)$

Exercise 12

q^3-5q^2-24q

Exercise 13

$3m^3-21m^2+30m$

Answer

$3m(m-5)(m-2)$

Exercise 14

$11n^3-55n^2+44n$

Exercise 15

$5x^4+10x^3-75x^2$

Answer

$5x^2(x-3)(x+5)$

Exercise 16

$6y^4+12y^3-48y^2$

Factor Trinomials Using Trial and Error

In the following exercises, factor.

Exercise 17

$2t^2+7t+5$

Answer

$(2t+5)(t+1)$

Exercise 18

$5y^2+16y+11$

Exercise 19

$11x^2+34x+3$

Answer

$(11x+1)(x+3)$

Exercise 20

$7b^2+50b+7$

Exercise 21

$4w^2-5w+1$

Answer

$(4w-1)(w-1)$

Exercise 22

$5x^2-17x+6$

Exercise 23

$6p^2-19p+10$

Answer

$(3p-2)(2p-5)$

Exercise 24

$21m^2-29m+10$

Exercise 25

$4q^2-7q-2$

Answer

$(4q+1)(q-2)$

Exercise 26

$10y^2-53y-11$

Exercise 27

$4p^2+17p-15$

Answer

$(4p-3)(p+5)$

Exercise 28

$6u^2+5u-14$

Exercise 29

$16x^2-32x+16$

Answer

$16(x-1)(x-1)$

Exercise 30

$81a^2+153a-18$

Exercise 31

$30q^3+140q^2+80q$

Answer

$10q(3q+2)(q+4)$

Exercise 32

$5y^3+30y^2-35y$

Factor Trinomials using the 'ac' Method

In the following exercises, factor.

Exercise 33

$5n^2+21n+4$

Answer

$(5n+1)(n+4)$

Exercise 34

$8w^2+25w+3$

Exercise 35

$9z^2+15z+4$

Answer

$(3z+1)(3z+4)$

Exercise 36

$3m^2+26m+48$

Exercise 37

$4k^2-16k+15$

Answer

$(2k-3)(2k-5)$

Exercise 38

$4q^2-9q+5$

Exercise 39

$5s^2-9s+4$

Answer

$(5s-4)(s-1)$

Exercise 40

$4r^2-20r+25$

Exercise 41

$6y^2+y-15$

Answer

$(3y+5)(2y-3)$

Exercise 42

$6p^2+p-22$

Exercise 43

$2n^2-27n-45$

Answer

$(2n+3)(n-15)$

Exercise 44

$12z^2-41z-11$

Exercise 45

$3x^2+5x+4$

Answer

prime

Exercise 46

$4y^2+15y+6$

Exercise 47

$60y^2+290y-50$

Answer

$10(6y-1)(y+5)$

Exercise 48

$6u^2-46u-16$

Exercise 49

$48z^3-102z^2-45z$

Answer

$3z(8z+3)(2z-5)$

Exercise 50

$90n^3+42n^2-216n$

Exercise 51

$16s^2+40s+24$

Answer

$8(2s+3)(s+1)$

Exercise 52

$24p^2+160p+96$

Exercise 53

$48y^2+12y-36$

Answer

$12(4y-3)(y+1)$

Exercise 54

$30x^2+105x-60$

Mixed Practice

In the following exercises, factor.

Exercise 55

$12y^2-29y+14$

Answer

$(4y-7)(3y-2)$

Exercise 56

$12x^2+36y-24z$

Exercise 57

a^2-a-20

Answer

$(a-5)(a+4)$

Exercise 58

m^2-m-12

Exercise 59

$6n^2+5n-4$

Answer

$(2n-1)(3n+4)$

Exercise 60

$12y^2-37y+21$

Exercise 61

$2p^2+4p+3$

Answer

prime

Exercise 62

$3q^2+6q+2$

Exercise 63

$13z^2+39z-26$

Answer

$13(z^2+3z-2)$

Exercise 64

$5r^2+25r+30$

Exercise 65

$x^2+3x-28$

Answer

$(x+7)(x-4)$

Exercise 66

$6u^2+7u-5$

Exercise 67

$3p^2+21p$

Answer

$3p(p+7)$

Exercise 68

$7x^2-21x$

Exercise 69

$6r^2+30r+36$

Answer

$6(r+2)(r+3)$

Exercise 70

$18m^2+15m+3$

Exercise 71

$24n^2+20n+4$

Answer

$4(2n+1)(3n+1)$

Exercise 72

$4a^2+5a+2$

Exercise 73

$x^2+2x-24$

Answer

$(x+6)(x-4)$

Exercise 74

$2b^2-7b+4$

Everyday Math

Exercise 75

Height of a toy rocket The height of a toy rocket launched with an initial speed of 80 feet per second from the balcony of an apartment building is related to the number of seconds, t, since it is launched by the trinomial $-16t^2+80t+96$. Factor this trinomial.

Answer

$-16(t-6)(t+1)$

Exercise 76

Height of a beach ball The height of a beach ball tossed up with an initial speed of 12 feet per second from a height of 4 feet is related to the number of seconds, t, since it is tossed by the trinomial $-16t^2+12t+4$. Factor this trinomial.

Writing Exercises

Exercise 77

List, in order, all the steps you take when using the "ac" method to factor a trinomial of the form $ax^2 + bx + c$.

Answer

Answers may vary.

Exercise 78

How is the "ac" method similar to the "undo FOIL" method? How is it different?

Exercise 79

What are the questions, in order, that you ask yourself as you start to factor a polynomial? What do you need to do as a result of the answer to each question?

Answer

Answers may vary.

Exercise 80

On your paper draw the chart that summarizes the factoring strategy. Try to do it without looking at the book. When you are done, look back at the book to finish it or verify it.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
recognize a preliminary strategy to factor polynomials completely.			
factor trinomials of the form $ax^2 + bx + c$ with a GCF.			
factor trinomials using trial and error.			
factor trinomials using the "ac" method.			

b. What does this checklist tell you about your mastery of this section? What steps will you take to improve?

This page titled 7.3E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.3E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.4: Factor Special Products

Learning Objectives

By the end of this section, you will be able to:

- Factor perfect square trinomials
- Factor differences of squares
- Factor sums and differences of cubes
- Choose method to factor a polynomial completely

Note

Before you get started, take this readiness quiz.

1. Simplify: $(12x)^2$
   If you missed this problem, review Example 6.2.22.
2. Multiply: $(m+4)^2$
   If you missed this problem, review Example 6.4.1.
3. Multiply: $(p-9)^2$
   If you missed this problem, review Example 6.4.4.
4. Multiply: $(k+3)(k-3)$
   If you missed this problem, review Example 6.4.16.

The strategy for factoring we developed in the last section will guide you as you factor most binomials, trinomials, and polynomials with more than three terms. We have seen that some binomials and trinomials result from special products—squaring binomials and multiplying conjugates. If you learn to recognize these kinds of polynomials, you can use the special products patterns to factor them much more quickly.

Factor Perfect Square Trinomials

Some trinomials are perfect squares. They result from multiplying a binomial times itself. You can square a binomial by using FOIL, but using the Binomial Squares pattern you saw in a previous chapter saves you a step. Let's review the Binomial Squares pattern by squaring a binomial using FOIL.

$$(3x+4)^2$$
$$(3x+4)(3x+4)$$
$$\underset{9x^2}{F}+\underset{12x}{O}+\underset{12x}{I}+\underset{16}{L}$$
$$9x^2+24x+16$$

The first term is the square of the first term of the binomial and the last term is the square of the last. The middle term is twice the product of the two terms of the binomial.

$$\begin{aligned}(3x)^2+2(3x\cdot 4)+4^2\\ 9x^2+24x+16\end{aligned} \qquad (7.4.1)$$

The trinomial $9x^2+24+16$ is called a perfect square trinomial. It is the square of the binomial $3x+4$.

We'll repeat the Binomial Squares Pattern here to use as a reference in factoring.

BINOMIAL SQUARES PATTERN

If a and b are real numbers,

$$(a+b)^2=a^2+2ab+b^2 \qquad (a-b)^2=a^2-2ab+b^2 \qquad (7.4.2)$$

When you square a binomial, the product is a perfect square trinomial. In this chapter, you are learning to factor—now, you will start with a perfect square trinomial and factor it into its prime factors.

You could factor this trinomial using the methods described in the last section, since it is of the form ax^2+bx+c. But if you recognize that the first and last terms are squares and the trinomial fits the **perfect square trinomials pattern**, you will save yourself a lot of work.

Here is the pattern—the reverse of the binomial squares pattern.

PERFECT SQUARE TRINOMIALS PATTERN

If a and b are real numbers,

$$a^2+2ab+b^2=(a+b)^2 \qquad a^2-2ab+b^2=(a-b)^2 \tag{7.4.3}$$

To make use of this pattern, you have to recognize that a given trinomial fits it. Check first to see if the leading coefficient is a perfect square, a^2. Next check that the last term is a perfect square, b^2. Then check the middle term—is it twice the product, $2ab$? If everything checks, you can easily write the factors.

Example 7.4.1: How to Factor Perfect Square Trinomials

Factor: $9x^2+12x+4$

Solution

Step 1. Does the trinomial fit the perfect square trinomials pattern, $a^2+2ab+b^2$?		
• Is the first term a perfect square? Write it as a square, a^2.	Is $9x^2$ a perfect square? Yes—write it as $(3x)^2$.	$9x^2+12x+4$ $(3x)^2$
• Is the last term a perfect square? Write it as a square, b^2.	Is 4 a perfect square? Yes—write it as $(2)^2$.	$(3x)^2 \qquad (2)^2$
• Check the middle term. Is it $2ab$?	Is $12x$ twice the product of $3x$ and 2? Does it match? Yes, so we have a perfect square trinomial!	$(3x)^2 \qquad (2)^2$ $2(3x)(2)$ $12x$
Step 2. Write the square of the binomial.	Write it as the square of a binomial.	$9x^2+12x+4$ $a^2 + 2 \cdot a \cdot b + b^2$ $(3x)^2+2\cdot 3x\cdot 2+2^2$ $(a+b)^2$ $(3x+2)^2$
Step 3. Check. $(3x+2)^2$ $(3x)^2+2\cdot 3x\cdot 2+2^2$ $9x^2+12x+4$ ✓		

Try It 7.4.2

Factor: $4x^2+12x+9$

Answer

$(2x+3)^2$

Try It 7.4.3

Factor: $9y^2+24y+16$

Answer

$(3y+4)^2$

The sign of the middle term determines which pattern we will use. When the middle term is negative, we use the pattern $a^2-2ab+b^2$, which factors to $(a-b)^2$.

The steps are summarized here.

FACTOR PERFECT SQUARE TRINOMIALS.

Step 1. Does the trinomial fit the pattern? $a^2+2ab+b^2$ $a^2-2ab+b^2$

- Is the first term a perfect square? $(a)^2$ $(a)^2$
  Write it as a square.
- Is the last term a perfect square? $(a)^2$ $(b)^2$ $(a)^2$ $(b)^2$
  Write it as a square.
- Check the middle term. Is it $2ab$? $(a)^2$ $\searrow 2\cdot a\cdot b \swarrow$ $(b)^2$ $(a)^2$ $\searrow 2\cdot a\cdot b \swarrow$ $(b)^2$

Step 2. Write the square of the binomial. $(a+b)^2$ $(a-b)^2$

Step 3. Check by multiplying.

We'll work one now where the middle term is negative.

Example 7.4.4

Factor: $81y^2-72y+16$

Solution

The first and last terms are squares. See if the middle term fits the pattern of a perfect square trinomial. The middle term is negative, so the binomial square would be $(a-b)^2$.

	$81y^2-72y+16$
Are the first and last terms perfect squares?	$(9y)^2$ $(4)^2$
Check the middle term.	$(9y)^2$ $(4)^2$ $2(9y)(4)$ $72y$
Does is match $(a-b)^2$? Yes.	a^2 -2 a $b+b^2$ $(9y)^2-2\cdot 9y\cdot 4+4^2$
Write the square of a binomial.	$(9y-4)^2$
Check by multiplying.	
$(9y-4)^2$	
$(9y)^2-2\cdot 9y\cdot 4+4^2$	
$81y^2-72y+16$✓	

Try It 7.4.5

Factor: $64y^2 - 80y + 25$

Answer

$(8y - 5)^2$

Try It 7.4.6

Factor: $16z^2 - 72z + 81$

Answer

$(4z - 9)^2$

The next example will be a perfect square trinomial with two variables.

Example 7.4.7

Factor: $36x^2 + 84xy + 49y^2$

Solution

	$36x^2 + 84xy + 49y^2$
Test each term to verify the pattern.	$\overset{a^2}{(6x)^2} \overset{+\,2}{+\,2} \overset{a}{\cdot\, 6x} \overset{b}{\cdot\, 7y} \overset{+}{+} \overset{b^2}{(7y)^2}$
Factor.	$(6x + 7y)^2$
Check by multiplying.	
$(6x + 7y)^2$	
$(6x)^2 + 2 \cdot 6x \cdot 7y + (7y)^2$	
$36x^2 + 84xy + 49y^2$ ✓	

Try It 7.4.8

Factor: $49x^2 + 84xy + 36y^2$

Answer

$(7x + 6y)^2$

Try It 7.4.9

Factor: $64m^2 + 112mn + 49n^2$

Answer

$(8m + 7n)^2$

Example 7.4.10

Factor: $9x^2 + 50x + 25$

Solution

	$9x^2+50x+25$
Are the first and last terms perfect squares?	$(3x)^2 \qquad (5)^2$
Check the middle term-is it 2ab?	$(3x)^2 \searrow_{2(3x)(5)} \swarrow (5)^2.$
	$30x$
No! $30x \neq 50x$	This does not fit the pattern!
Factor using the "ac" method.	$9x^2+50x+25$
ac	
Notice: $9 \cdot 25$ and $5 \cdot 45 = 225$	
225	
	$9x^2+5x+45x+25$
Split the middle term.	$x(9x+5)+5(9x+5)$
	$(9x+5)(x+5)$
Factor by grouping.	
Check.	
$(9x+5)(x+5)$	
$9x^2+45x+5x+25$	
$9x^2+50x+25$✓	

? Try It 7.4.11

Factor: $16r^2+30rs+9s^2$

Answer

$(8r+3s)(2r+3s)$

? Try It 7.4.12

Factor: $9u^2+87u+100$

Answer

$(3u+4)(3u+25)$

Remember the very first step in our Strategy for Factoring Polynomials? It was to ask "is there a greatest common factor?" and, if there was, you factor the GCF before going any further. Perfect square trinomials may have a GCF in all three terms and it should be factored out first. And, sometimes, once the GCF has been factored, you will recognize a perfect square trinomial.

✓ Example 7.4.13

Factor: $36x^2y-48xy+16y$

Solution

	$36x^2y-48xy+16y$
Is there a GCF? Yes, $4y$, so factor it out.	$4y\left(9x^2-12x+4\right)$
Is this a perfect square trinomial?	
Verify the pattern.	$4y[(3x)^2-2 \cdot 3x \cdot 2+2^2]$ (a^2 -2 a b + b^2)
Factor.	$4y(3x-2)^2$
Remember: Keep the factor $4y$ in the final product.	

Check.
$4y(3x-2)^2$
$4y[(3x)^2-2\cdot 3x\cdot 2+2^2]$
$4y(9x)^2-12x+4$
$36x^2y-48xy+16y$✓

? Try It 7.4.14

Factor: $8x^2y-24xy+18y$

Answer

$2y(2x-3)^2$

? Try It 7.4.15

Factor: $27p^2q+90pq+75q$

Answer

$3q(3p+5)^2$

Factor Differences of Squares

The other special product you saw in the previous was the Product of Conjugates pattern. You used this to multiply two binomials that were conjugates. Here's an example:

$$\begin{array}{c}(3x-4)(3x+4)\\ 9x^2-16\end{array} \tag{7.4.4}$$

Remember, when you multiply conjugate binomials, the middle terms of the product add to 0. All you have left is a binomial, the difference of squares.

Multiplying conjugates is the only way to get a binomial from the product of two binomials.

PRODUCT OF CONJUGATES PATTERN

If a and b are real numbers


$$(a-b)(a+b)=a^2-b^2 \tag{7.4.5}$$

The product is called a difference of squares.

To factor, we will use the product pattern "in reverse" to factor the difference of squares. A **difference of squares** factors to a product of conjugates.

DIFFERENCE OF SQUARES PATTERN

If a and b are real numbers,



https://math.libretexts.org/@go/page/30555

Remember, “difference” refers to subtraction. So, to use this pattern you must make sure you have a binomial in which two squares are being subtracted.

Example 7.4.16: How to Factor Differences of Squares

Factor: $x^2 - 4$

Solution

Step 1. Does the binomial fit the pattern?		$x^2 - 4$
• Is this a difference?	Yes	$x^2 - 4$
• Are the first and last terms perfect squares?	Yes	
Step 2. Write them as squares.	Write them as x^2 and 2^2.	$a^2 - b^2$ $(x)^2 - 2^2$
Step 3. Write the product of conjugates.		$(a - b)(a + b)$ $(x - 2)(x + 2)$
Step 4. Check.		$(x - 2)(x + 2)$ $x^2 - 4$ ✓

Try It 7.4.17

Factor: $h^2 - 81$

Answer

$(h-9)(h+9)$

Try It 7.4.18

Factor: $k^2 - 121$

Answer

$(k-11)(k+11)$

FACTOR DIFFERENCES OF SQUARES.

Step 1. Does the binomial fit the pattern? $a^2 - b^2$
- Is this a difference? $\_\_ - \_\_$
- Are the first and last terms perfect squares?

Step 2. Write them as squares. $(a)^2 - (b)^2$

Step 3. Write the product of conjugates. $(a-b)(a+b)$

Step 4. Check by multiplying.

It is important to remember that *sums of squares do not factor into a product of binomials*. There are no binomial factors that multiply together to get a sum of squares. After removing any GCF, the expression $a^2 + b^2$ is prime!

Don’t forget that 1 is a perfect square. We’ll need to use that fact in the next example.

Example 7.4.19

Factor: $64y^2 - 1$

Solution

	$64y^2 - 1$
Is this a difference? Yes.	$64y^2 - 1$
Are the first and last terms perfect squares?	
Yes - write them as squares.	$a^2 - b^2$ $(8y)^2 - 1^2$
Factor as the product of conjugates.	$(a - b)(a + b)$ $(8y - 1)(8y + 1)$
Check by multiplying.	
$(8y - 1)(8y + 1)$	
$64y^2 - 1$✓	

Try It 7.4.20

Factor: $m^2 - 1$

Answer

$(m - 1)(m + 1)$

Try It 7.4.21

Factor: $81y^2 - 1$

Answer

$(9y - 1)(9y + 1)$

Example 7.4.22

Factor: $121x^2 - 49y^2$

Solution

	$121x^2 - 49y^2$
Is this a difference of squares? Yes.	$(11x)^2 - (7y)^2$
Factor as the product of conjugates.	$(11x - 7y)(11x + 7y)$
Check by multiplying.	
$(11x - 7y)(11x + 7y)$	
$121x^2 - 49y^2$✓	

Try It 7.4.23

Factor: $196m^2 - 25n^2$

Answer

$(16m - 5n)(16m + 5n)$

Try It 7.4.24

Factor: $144p^2 - 9q^2$

Answer

$(12p - 3q)(12p + 3q)$

The binomial in the next example may look "backwards," but it's still the difference of squares.

Example 7.4.25

Factor: $100 - h^2$

Solution

	$100 - h^2$
Is this a difference of squares? Yes.	$(10)^2 - (h)^2$
Factor as the product of conjugates.	$(10 - h)(10 + h)$

Check by multiplying.

$(10 - h)(10 + h)$

$100 - h^2$ ✓

Be careful not to rewrite the original expression as $h^2 - 100$.

Factor $h^2 - 100$ on your own and then notice how the result differs from $(10 - h)(10 + h)$.

Try It 7.4.26

Factor: $144 - x^2$

Answer

$(12 - x)(12 + x)$

Try It 7.4.27

Factor: $169 - p^2$

Answer

$(13 - p)(13 + p)$

To completely factor the binomial in the next example, we'll factor a difference of squares twice!

Example 7.4.28

Factor: $x^4 - y^4$

Solution

Is this a difference of squares? Yes.	x^4-y^4
Factor it as the product of conjugates.	$\left(x^2\right)^2-\left(y^2\right)^2$
Notice the first binomial is also a difference of squares!	$\left(x^2-y^2\right)\left(x^2+y^2\right)$
Factor it as the product of conjugates. The last factor, the sum of squares, cannot be factored.	$(x-y)(x+y)\left(x^2+y^2\right)$

Check by multiplying.

$(x-y)(x+y)\left(x^2+y^2\right)$

$[(x-y)(x+y)]\left(x^2+y^2\right)$

$\left(x^2-y^2\right)\left(x^2+y^2\right)$

x^4-y^4✓

Try It 7.4.29

Factor: a^4-b^4

Answer

$\left(a^2+b^2\right)(a+b)(a-b)$

Try It 7.4.30

Factor: x^4-16

Answer

$\left(x^2+4\right)(x+2)(x-2)$

As always, you should look for a common factor first whenever you have an expression to factor. Sometimes a common factor may "disguise" the difference of squares and you won't recognize the perfect squares until you factor the GCF.

Example 7.4.31

Factor: $8x^2y-98y$

Solution

	$8x^2y-98y$
Is there a GCF? Yes, $2y-$ factor it out!	$2y\left(4x^2-49\right)$
Is the binomial a difference of squares? Yes.	$2y\left((2x)^2-(7)^2\right)$
Factor as a product of conjugates.	$2y(2x-7)(2x+7)$
Check by multiplying.	

$2y(2x-7)(2x+7)$

$2y[(2x-7)(2x+7)]$

$2y\left(4x^2-49\right)$

$8x^2y-98y$✓

Try It 7.4.32

Factor: $7xy^2 - 175x$

Answer

$7x(y-5)(y+5)$

Try It 7.4.33

Factor: $45a^2b - 80b$

Answer

$5b(3a-4)(3a+4)$

Example 7.4.34

Factor: $6x^2 + 96$

Solution

	$6x^2 + 96$
Is there a GCF? Yes, 6 — factor it out!	$6\left(x^2 + 16\right)$
Is the binomial a difference of squares? No, it is a sum of squares. Sums of squares do not factor!	
Check by multiplying.	

$6\left(x^2 + 16\right)$

$6x^2 + 96$✓

Try It 7.4.35

Factor: $8a^2 + 200$

Answer

$8\left(a^2 + 25\right)$

Try It 7.4.36

Factor: $36y^2 + 81$

Answer

$9\left(4y^2 + 9\right)$

Factor Sums and Differences of Cubes

There is another special pattern for factoring, one that we did not use when we multiplied polynomials. This is the pattern for the sum and difference of cubes. We will write these formulas first and then check them by multiplication.

$$\begin{aligned} a^3 + b^3 &= (a+b)\left(a^2 - ab + b^2\right) \\ a^3 - b^3 &= (a-b)\left(a^2 + ab + b^2\right) \end{aligned}$$

We'll check the first pattern and leave the second to you.

	$(a+b)\left(a^2-ab+b^2\right)$
Distribute.	$a\left(a^2-ab+b^2\right)+b\left(a^2-ab+b^2\right)$
Multiply.	$a^3-a^2b+ab^2+a^2b-ab^2+b^3$
Combine like terms.	a^3+b^3

SUM AND DIFFERENCE OF CUBES PATTERN

$$\begin{aligned} a^3+b^3&=(a+b)\left(a^2-ab+b^2\right)\\ a^3-b^3&=(a-b)\left(a^2+ab+b^2\right)\end{aligned} \tag{7.4.6}$$

The two patterns look very similar, don't they? But notice the signs in the factors. The sign of the binomial factor matches the sign in the original binomial. And the sign of the middle term of the trinomial factor is the opposite of the sign in the original binomial. If you recognize the pattern of the signs, it may help you memorize the patterns.

$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$ — same sign; opposite signs

$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$ — same sign; opposite signs

The trinomial factor in the **sum and difference of cubes pattern** cannot be factored.

It can be very helpful if you learn to recognize the cubes of the integers from 1 to 10, just like you have learned to recognize squares. We have listed the cubes of the integers from 1 to 10 in Figure 7.4.1.




Figure 7.4.1

Example 7.4.37: How to Factor the Sum or Difference of Cubes

Factor: x^3+64

Solution

Step 1. Does the binomial fit the sum or difference of cubes pattern?		$x^3 + 64$
• Is it a sum or difference?	This is a sum.	$x^3 + 64$
• Are the first and last terms perfect cubes?	Yes	
Step 2. Write the terms as cubes.	Write them as x^3 and 4^3	$a^3 + b^3$ $x^3 + 4^3$
Step 3. Use either the sum or difference of cubes pattern.	This is a sum of cubes.	$(a + b)(a^2 - ab + b^2)$ $(x + 4)(x^2 - 4x + 4^2)$

Step 4. Simplify inside the parentheses.	Simplify 4^2.	$(x+4)(x^2-4x+16)$
Step 5. Check by multiplying the factors.		$\begin{array}{r} x^2-4x+16 \\ x+4 \\ \hline 4x^2-16x+64 \\ x^3-4x^2+16x \\ \hline x^3 \qquad +64 \checkmark \end{array}$

? Try It 7.4.38

Factor: x^3+27

Answer

$(x+3)\left(x^2-3x+9\right)$

? Try It 7.4.39

Factor: y^3+8

Answer

$(y+2)\left(y^2-2y+4\right)$

FACTOR THE SUM OR DIFFERENCE OF CUBES.

To factor the sum or difference of cubes:

1. Does the binomial fit the sum or difference of cubes pattern?
   - Is it a sum or difference?
   - Are the first and last terms perfect cubes?
2. Write them as cubes.
3. Use either the sum or difference of cubes pattern.
4. Simplify inside the parentheses
5. Check by multiplying the factors.

✓ Example 7.4.40

Factor: : x^3-1000

Solution

	x^3-1000
This binomial is a difference. The first and last terms are perfect cubes.	
Write the terms as cubes.	$\begin{matrix} a^3 & - & b^3 \\ x^3 & - & 10^3 \end{matrix}$
Use the difference of cubes pattern.	$\begin{matrix} (a & - & b) & (a^2 & + & ab & + & b^2) \\ (x & - & 10) & (x^2 & + & 10\cdot x & + & 10^2) \end{matrix}$
Simplify.	$\begin{matrix} (a & - & b) & (a^2 & + & ab & + & b^2) \\ (x & - & 10) & (x^2 & + & 10x & + & 100) \end{matrix}$
Check by multiplying.	

$(x-10)\ \ (x^2+10x+100)$

$x^2+10x+100$

$x-100$

x^3+10x^2+100x

$-10x^2-100x-1000$ ✓

$x^3 \qquad -1000$

Try It 7.4.41

Factor: u^3-125

Answer

$(u-5)\left(u^2+5u+25\right)$

Try It 7.4.42

Factor: v^3-343

Answer

$(v-7)\left(v^2+7v+49\right)$

Be careful to use the correct signs in the factors of the sum and difference of cubes.

Example 7.4.43

Factor: $512-125p^3$

Solution

	$512-125p^3$
This binomial is a difference. The first and last terms are perfect cubes.	
Write the terms as cubes.	$8^3-(5p)^3$
Use the difference of cubes pattern.	$(8-5p)\left(8^2+8\cdot 5p+(5p)^2\right)$
Simplify.	$(8-5p)\left(64+40p+25p^2\right)$
Check by multiplying.	We'll leave the check to you.

Try It 7.4.44

Factor: $64-27x^3$

Answer

$(4-3x)\left(16+12x+9x^2\right)$

Try It 7.4.45

Factor: $27-8y^3$

Answer

$(3-2y)\left(9+6y+4y^2\right)$

Example 7.4.46

Factor: $27u^3-125v^3$

Solution

	$27u^3-125v^3$
This binomial is a difference. The first and last terms are perfect cubes.	
Write the terms as cubes.	$(3u)^3-(5v)^3$
Use the difference of cubes pattern.	$(3u-5v)\left((3u)^2+3u\cdot 5v+(5v)^2\right)$
Simplify.	$(3u-5v)\left(9u^2+15uv+25v^2\right)$
Check by multiplying.	We'll leave the check to you.

Try It 7.4.47

Factor: $8x^3-27y^3$

Answer

$(2x-3y)\left(4x^2+6xy+9y^2\right)$

Try It 7.4.48

Factor: $1000m^3-125n^3$

Answer

$(10m-5n)\left(100m^2+50mn+25n^2\right)$

In the next example, we first factor out the GCF. Then we can recognize the sum of cubes.

Example 7.4.49

Factor: $5m^3+40n^3$

Solution

	$5m^3+40n^3$
Factor the common factor.	$5(m^3+8n^3)$
This binomial is a sum. The first and last terms are perfect cubes.	

Write the terms as cubes.	$5\left(m^3+(2n)^3\right)$
Use the sum of cubes pattern.	$5(m+2n)\left(m^2-m\cdot 2n+(2n)^2\right)$
Simplify.	$5(m+2n)\left(m^2-2mn+4n^2\right)$

Check. To check, you may find it easier to multiply the sum of cubes factors first, then multiply that product by 5. We'll leave the multiplication for you.

$5(m+2n)\left(m^2-2mn+4n^2\right)$

Try It 7.4.50

Factor: $500p^3+4q^3$

Answer

$4(5p+q)\left(25p^2-5pq+q^2\right)$

Try It 7.4.51

Factor: $432c^3+686d^3$

Answer

$2(6c+7d)\left(36c^2-42cd+49d^2\right)$

Note

Access these online resources for additional instruction and practice with factoring special products.

- Sum of Difference of Cubes
- Difference of Cubes Factoring

Key Concepts

- **Factor perfect square trinomials** See Example.

  Step 1 . Does the trinomial fit the pattern? $a^2+2ab+b^2$ $a^2-2ab+b^2$

  - Is the first term a perfect square? $(a)^2$ $(a)^2$

    Write it as a square.

  - Is the last term a perfect square? $(a)^2$ $(b)^2$ $(a)^2$ $(b)^2$

    Write it as a square.

  - Check the middle term. Is it $2ab$? $(a)^2$ $\searrow 2\cdot a\cdot b \swarrow$ $(b)^2$ $(a)^2$ $\searrow 2\cdot a\cdot b \swarrow$ $(b)^2$

  Step 2. Write the square of the binomial. $(a+b)^2$ $(a-b)^2$

  Step 3. Check by multiplying.

- **Factor differences of squares** See Example.

  Step 1. Does the binomial fit the pattern? a^2-b^2

  - Is this a difference? $\_\_-\_\_$
  - Are the first and last terms perfect squares?

  Step 2. Write them as squares. $(a)^2-(b)^2$

  Step 3. Write the product of conjugates. $(a-b)(a+b)$

  Step 4. Check by multiplying.

- **Factor sum and difference of cubes** To factor the sum or difference of cubes: See Example.

  1. Does the binomial fit the sum or difference of cubes pattern? Is it a sum or difference? Are the first and last terms perfect cubes?

2. Write them as cubes.
3. Use either the sum or difference of cubes pattern.
4. Simplify inside the parentheses
5. Check by multiplying the factors.

Glossary


perfect square trinomials pattern

If a and b are real numbers,

$$a^2+2ab+b^2=(a+b)^2 \qquad a^2-2ab+b^2=(a-b)^2$$

difference of squares pattern

If a and b are real numbers,



sum and difference of cubes pattern

$$a^3+b^3=(a+b)(a^2-ab+b^2) \qquad a^3-b^3=(a-b)(a^2+ab+b^2)$$

This page titled 7.4: Factor Special Products is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.4: Factor Special Products** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.4E: Exercises

Practice Makes Perfect

Factor Perfect Square Trinomials

In the following exercises, factor.

Exercise 1

$16y^2+24y+9$

Answer

$(4y+3)^2$

Exercise 2

$25v^2+20v+4$

Exercise 3

$36s^2+84s+49$

Answer

$(6s+7)^2$

Exercise 4

$49s^2+154s+121$

Exercise 5

$100x^2-20x+1$

Answer

$(10x-1)^2$

Exercise 6

$64z^2-16z+1$

Exercise 7

$25n^2-120n+144$

Answer

$(5n-12)^2$

Exercise 8

$4p^2-52p+169$

Exercise 9

$49x^2 - 28xy + 4y^2$

Answer

$(7x - 2y)^2$

Exercise 10

$25r^2 - 60rs + 36s^2$

Exercise 11

$25n^2 + 25n + 4$

Answer

$(5n + 4)(5n + 1)$

Exercise 12

$100y^2 - 20y + 1$

Exercise 13

$64m^2 - 16m + 1$

Answer

$(8m - 1)^2$

Exercise 14

$100x^2 - 25x + 1$

Exercise 15

$10k^2 + 80k + 160$

Answer

$10(k + 4)^2$

Exercise 16

$64x^2 - 96x + 36$

Exercise 17

$75u^3 - 30u^2v + 3uv^2$

Answer

$3u(5u - v)^2$

Exercise 18

$90p^3 + 300p^2q + 250pq^2$

Factor Differences of Squares

In the following exercises, factor.

Exercise 19

$x^2 - 16$

Answer

$(x-4)(x+4)$

Exercise 20

$n^2 - 9$

Exercise 21

$25v^2 - 1$

Answer

$(5v-1)(5v+1)$

Exercise 22

$169q^2 - 1$

Exercise 23

$121x^2 - 144y^2$

Answer

$(11x-12y)(11x+12y)$

Exercise 24

$49x^2 - 81y^2$

Exercise 25

$169c^2 - 36d^2$

Answer

$(13c-6d)(13c+6d)$

Exercise 26

$36p^2 - 49q^2$

Exercise 27

$4-49x^2$

Answer

$(2-7x)(2+7x)$

Exercise 28

$121-25s^2$

Exercise 29

$16z^4-1$

Answer

$(2z-1)(2z+1)(4z^2+1)$

Exercise 30

m^4-n^4

Exercise 31

$5q^2-45$

Answer

$5(q-3)(q+3)$

Exercise 32

$98r^3-72r$

Exercise 33

$24p^2+54$

Answer

$6(4p^2+9)$

Exercise 34

$20b^2+140$

Factor Sums and Differences of Cubes

In the following exercises, factor.

Exercise 35

x^3+125

Answer

$(x+5)(x^2-5x+25)$

Exercise 36

n^3+512

Exercise 37

z^3-27

Answer

$(z-3)(z^2+3z+9)$

Exercise 38

v^3-216

Exercise 39

$8-343t^3$

Answer

$(2-7t)(4+14t+49t^2)$

Exercise 40

$125-27w^3$

Exercise 41

$8y^3-125z^3$

Answer

$(2y-5z)(4y^2+10yz+25z^2)$

Exercise 42

$27x^3-64y^3$

Exercise 43

$7k^3+56$

Answer

$7(k+2)(k^2-2k+4)$

Exercise 44

$6x^3-48y^3$

https://math.libretexts.org/@go/page/30556

Exercise 45

$2-16y^3$

Answer

$2(1-2y)(1+2y+4y^2)$

Exercise 46

$-2x^3-16y^3$

Mixed Practice

In the following exercises, factor.

Exercise 47

$64a^2-25$

Answer

$(8a-5)(8a+5)$

Exercise 48

$121x^2-144$

Exercise 49

$27q^2-3$

Answer

$3(3q-1)(3q+1)$

Exercise 50

$4p^2-100$

Exercise 51

$16x^2-72x+81$

Answer

$(4x-9)^2$

Exercise 52

$36y^2+12y+1$

Exercise 53

$8p^2+2$

Answer

$2(4p^2+1)$

Exercise 54

$81x^2+169$

Exercise 55

$125-8y^3$

Answer

$(5-2y)(25+10y+4y^2)$

Exercise 56

$27u^3+1000$

Exercise 57

$45n^2+60n+20$

Answer

$5(3n+2)^2$

Exercise 58

$48q^3-24q^2+3q$

Everyday Math

Exercise 59

Landscaping Sue and Alan are planning to put a 15 foot square swimming pool in their backyard. They will surround the pool with a tiled deck, the same width on all sides. If the width of the deck is w, the total area of the pool and deck is given by the trinomial $4w^2+60w+225$.

Answer

$(2w+15)^2$

Exercise 60

Home repair The height a twelve foot ladder can reach up the side of a building if the ladder's base is b feet from the building is the square root of the binomial $144-b^2$.

Writing Exercises

Exercise 61

Why was it important to practice using the binomial squares pattern in the chapter on multiplying polynomials?

Answer

Answers may vary.

Exercise 62

How do you recognize the binomial squares pattern?

Exercise 63

Explain why $n^2 + 25 \neq (n+5)^2$.

Answer

Answers may vary.

Exercise 64

Maribel factored $y^2 - 30y + 81$ as (y−9)^2. How do you know that this is incorrect?

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
factor perfect square trinomials.			
factor differences of squares.			
factor sums and differences of cubes.			

b. On a scale of 1–10, how would you rate your mastery of this section in light of your responses on the checklist? How can you improve this?

This page titled 7.4E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.4E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.5: General Strategy for Factoring Polynomials

Learning Objectives

By the end of this section, you will be able to:

- Recognize and use the appropriate method to factor a polynomial completely

Note

Before you get started, take this readiness quiz.

1. Factor $y^2 - 2y - 24$.
   If you missed this problem, review Example 7.2.19.
2. Factor $3t^2 + 17t + 10$.
   If you missed this problem, review Example 7.3.28.
3. Factor $36p^2 - 60p + 25$.
   If you missed this problem, review Example 7.4.1.
4. Factor $5x^2 - 80$.
   If you missed this problem, review Example 7.4.31.

Recognize and Use the Appropriate Method to Factor a Polynomial Completely

You have now become acquainted with all the methods of factoring that you will need in this course. (In your next algebra course, more methods will be added to your repertoire.) The figure below summarizes all the factoring methods we have covered. Figure 7.5.1 outlines a strategy you should use when factoring polynomials.




Figure 7.5.1

FACTOR POLYNOMIALS.

1. Is there a greatest common factor?
   - Factor it out.
2. Is the polynomial a binomial, trinomial, or are there more than three terms?
   - If it is a binomial:
     Is it a sum?
     - Of squares? Sums of squares do not factor.
     - Of cubes? Use the sum of cubes pattern.

     Is it a difference?
     - Of squares? Factor as the product of conjugates.
     - Of cubes? Use the difference of cubes pattern.

- If it is a trinomial:
  Is it of the form x^2+bx+c? ? Undo FOIL.
  Is it of the form ax^2+bx+c ?
  - If aa and cc are squares, check if it fits the trinomial square pattern.
  - Use the trial and error or "ac" method.
- If it has more than three terms:
  Use the grouping method.

3. Check.
   - Is it factored completely?
   - Do the factors multiply back to the original polynomial?

Remember, a polynomial is completely factored if, other than monomials, its factors are prime!

Example 7.5.1

Factor completely: $4x^5+12x^4$

Solution

Is there a GCF?	Yes, $4x^4$	$4x^5+12x^4$
Factor out the GCF.		$4x^4(x+3)$
In the parentheses, is it a binomial, a trinomial, or are there more than three terms?	Binomial.	
Is it a sum?		Yes.
Of squares? Of cubes?		No.
Check.		
Is the expression factored completely?		Yes.
Multiply.		

$4x^4(x+3)$
$4x^4\cdot x+4x^4\cdot 3$
$4x^5+12x^4$✓

Try It 7.5.2

Factor completely: $3a^4+18a^3$

Answer

$3a^3(a+6)$

Try It 7.5.3

Factor completely: $45b^6+27b^5$

Answer

$9b^5(5b+3)$

Example 7.5.4

Factor completely: $12x^2-11x+2$

Solution

		$12x^2 - 11x + 2$
Is there a GCF?	No.	
Is it a binomial, trinomial, or are there more than three terms?	Trinomial.	
Are a and c perfect squares?	No, $a = 12$, not a perfect square.	
Use trial and error or the "ac" method. We will use trial and error here.		$12x^2 - 11x + 2$ $1x, 12x$ $\quad -1, -2$ $2x, 6x$ $3x, 4x$

$12x^2 - 11x + 2$	
Possible factors	**Product**
$(x - 1)(12x - 2)$	Not an option
$(x - 2)(12x - 1)$	$12x^2 - 25x + 2$
$(2x - 1)(6x - 2)$	Not an option
$(2x - 2)(6x - 1)$	Not an option
$(3x - 1)(4x - 2)$	Not an option
$(3x - 2)(4x - 1)$	$12x^2 - 11x + 2$

If the trinomial has no common factors, then neither factor can contain a common factor. That means each of these combinations is not an option.

$(3x - 2)(4x - 1)$

Check. $12x^2 - 3x - 8x + 2$

$12x^2 - 11x + 2$ ✓

? Try It 7.5.5

Factor completely: $10a^2 - 17a + 6$

Answer

$(5a - 6)(2a - 1)$

? Try It 7.5.6

Factor completely: $8x^2 - 18x + 9$

Answer

$(2x - 3)(4x - 3)$

✓ Example 7.5.7

Factor completely: $g^3 + 25g$

Solution

Is there a GCF?	Yes, g.	g^3+25g
Factor out the GCF.		$g\left(g^2+25\right)$
In the parentheses, is it a binomial, trinomial, or are there more than three terms?	Binomial.	
Is it a sum? Of squares?	Yes.	Sums of squares are prime.
Check.		
Is the expression factored completely?	Yes.	
Multiply.		
$g\left(g^2+25\right)$		
g^3+25g✓		

Try It 7.5.8

Factor completely: x^3+36x

Answer

$x\left(x^2+36\right)$

Try It 7.5.9

Factor completely: $27y^2+48$

Answer

$3\left(9y^2+16\right)$

Example 7.5.10

Factor completely: $12y^2-75$

Solution

Is there a GCF?	Yes, 3.	$12y^2-75$
Factor out the GCF.		$3\left(4y^2-25\right)$
In the parentheses, is it a binomial, trinomial, or are there more than three terms?	Binomial.	
Is it a sum?	No.	
Is it a difference? Of squares or cubes?	Yes, squares.	$3\left((2y)^2-(5)^2\right)$
Write as a product of conjugates.		$3(2y-5)(2y+5)$
Check.		
Is the expression factored completely?	Yes.	
Neither binomial is a difference of squares.		
Multiply.		
$3(2y-5)(2y+5)$		
$3\left(4y^2-25\right)$		
$12y^2-75$✓		

Try It 7.5.11

Factor completely: $16x^3 - 36x$

Answer

$4x(2x-3)(2x+3)$

Try It 7.5.12

Factor completely: $27y^2 - 48$

Answer

$3(3y-4)(3y+4)$

Example 7.5.13

Factor completely: $4a^2 - 12ab + 9b^2$

Solution

Is there a GCF?	No.	$4a^2 - 12ab + 9b^2$
Is it a binomial, trinomial, or are there more terms?		
Trinomial with $a \neq 1$. But the first term is a perfect square.		
Is the last term a perfect square?	Yes.	$(2a)^2 - 12ab + (3b)^2$
Does it fit the pattern, $a^2 - 2ab + b^2$?	Yes.	$(2a)^2 - 12ab + (3b)^2$ $-2(2a)(3b)$ $12ab$
Write it as a square.		$(2a - 3b)^2$
Check your answer.		
Is the expression factored completely?		
Yes.		
The binomial is not a difference of squares.		
Multiply.		
$(2a - 3b)^2$		
$(2a)^2 - 2 \cdot 2a \cdot 3b + (3b)^2$		
$4a^2 - 12ab + 9b^2$ ✓		

Try It 7.5.14

Factor completely: $4x^2 + 20xy + 25y^2$

Answer

$(2x+5y)^2$

Try It 7.5.15

Factor completely: $9m^2+42mn+49n^2$

Answer

$(3m+7n)^2$

Example 7.5.16

Factor completely: $6y^2-18y-60$

Solution

Is there a GCF?	Yes, 6.	$6y^2-18y-60$
Factor out the GCF.	Trinomial with leading coefficient 1	$6\left(y^2-3y-10\right)$
In the parentheses, is it a binomial, trinomial, or are there more terms?		
"Undo' FOIL.	$6(y\quad)(y\quad)$	$6(y+2)(y-5)$
Check your answer.		
Is the expression factored completely?		Yes.
Neither binomial is a difference of squares.		
Multiply.		

$$6(y+2)(y-5)$$
$$6\left(y^2-5y+2y-10\right)$$
$$6\left(y^2-3y-10\right)$$
$$6y^2-18y-60\checkmark$$

Try It 7.5.17

Factor completely: $8y^2+16y-24$

Answer

$8(y-1)(y+3)$

Try It 7.5.18

Factor completely: $5u^2-15u-270$

Answer

$5(u-9)(u+6)$

Example 7.5.19

Factor completely: $24x^3+81$

Solution

Is there a GCF?	Yes, 3.	$24x^3+81$
Factor it out.		$3\left(8x^3+27\right)$

In the parentheses, is it a binomial, trinomial, or are there more than three terms?	Binomial.	
Is it a sum or difference?	Sum.	
Of squares or cubes?	Sum of cubes.	$3\left(\overset{a^3}{(2x)^3}+\overset{b^3}{(3)^3}\right)$
Write it using the sum of cubes pattern.		$3\left(\overset{a}{2x}+\overset{b}{3}\right)\left(\overset{a^2}{(2x)^2}-\overset{ab}{2x\cdot 3}+\overset{b^2}{3^2}\right)$
Is the expression factored completely?	Yes.	$3(2x+3)\left(4x^2-6x+9\right)$
Check by multiplying.		We leave the check to you.

Try It 7.5.20

Factor completely: $250m^3+432$

Answer

$2(5m+6)\left(25m^2-30m+36\right)$

Try It 7.5.21

Factor completely: $81q^3+192$

Answer

$3(3q+4)\left(9q^2-12q+16\right)$

Example 7.5.22

Factor completely: $2x^4-32$

Solution

Is there a GCF?	Yes, 2.	$2x^4-32$
Factor out the GCF.		$2\left(x^4-16\right)$
In the parentheses, is it a binomial, trinomial, or are there more than three terms?	Binomial.	
Is it a sum or difference?	Yes.	
Of squares or cubes?	Difference of squares.	$2\left(\left(x^2\right)^2-(4)^2\right)$
Write it as a product of conjugates.		$2\left(x^2-4\right)\left(x^2+4\right)$
The first binomial is again a difference of squares.		$2\left((x)^2-(2)^2\right)\left(x^2+4\right)$
Write it as a product of conjugates.		$2(x-2)(x+2)\left(x^2+4\right)$
Is the expression factored completely?	Yes.	

None of these binomials is a difference of squares.
Check your answer.
Multiply.

$$2(x-2)(x+2)\left(x^2+4\right)$$
$$2(x-2)(x+2)\left(x^2+4\right)$$
$$2(x-10)$$
$$2x^4-32\checkmark$$

Try It 7.5.23

Factor completely: $4a^4-64$

Answer

$4\left(a^2+4\right)(a-2)(a+2)$

Try It 7.5.24

Factor completely: $7y^4-7$

Answer

$7\left(y^2+1\right)(y-1)(y+1)$

Example 7.5.25

Factor completely: $3x^2+6bx-3ax-6ab$

Solution

Is there a GCF?	Yes, 3.	$3x^2+6bx-3ax-6ab$
Factor out the GCF.		$3\left(x^2+2bx-ax-2ab\right)$
In the parentheses, is it a binomial, trinomial, or are there more terms?	More than 3 terms.	
Use grouping.		$3[x(x+2b)-a(x+2b)]$
		$3(x+2b)(x-a)$
Check your answer.		

Is the expression factored completely? Yes.

Multiply.

$3(x+2b)(x-a)$

$3\left(x^2-ax+2bx-2ab\right)$

$3x^2-3ax+6bx-6ab$✓

? Try It 7.5.26

Factor completely: $6x^2-12xc+6bx-12bc$

Answer

$6(x+b)(x-2c)$

? Try It 7.5.27

Factor completely: $16x^2+24xy-4x-6y$

Answer

$2(4x-1)(x+3y)$

✓ Example 7.5.28

Factor completely: $10x^2-34x-24$

Solution

Is there a GCF?	Yes, 2.	$10x^2-34x-24$
Factor out the GCF.		$2\left(5x^2-17x-12\right)$
In the parentheses, is it a binomial, trinomial, or are there more than three terms?	Trinomial with $a\neq 1$	
Use trial and error or the "ac" method.		$2\left(5x^2-17x-12\right)$
		$2(5x+3)(x-4)$
Check your answer. Is the expression factored completely? Yes.		

Multiply.

$2(5x+3)(x-4)$

$2\left(5x^2-20x+3x-12\right)$

$2\left(5x^2-17x-12\right)$

$10x^2-34x-24$✓

? Try It 7.5.29

Factor completely: $4p^2 - 16p + 12$

Answer

$4(p-1)(p-3)$

? Try It 7.5.30

Factor completely: $6q^2 - 9q - 6$

Answer

$3(q-2)(2q+1)$

Key Concepts

- **General Strategy for Factoring Polynomials See Figure 7.5.1.**
- **How to Factor Polynomials**
  1. Is there a greatest common factor? Factor it out.
  2. Is the polynomial a binomial, trinomial, or are there more than three terms?
     - If it is a binomial:
       Is it a sum?
       - Of squares? Sums of squares do not factor.
       - Of cubes? Use the sum of cubes pattern.

       Is it a difference?
       - Of squares? Factor as the product of conjugates.
       - Of cubes? Use the difference of cubes pattern.
     - If it is a trinomial:
       Is it of the form $x^2 + bx + c$? Undo FOIL.
       Is it of the form $ax^2 + bx + c$?
       - If 'a' and 'c' are squares, check if it fits the trinomial square pattern.
       - Use the trial and error or 'ac' method.
     - If it has more than three terms:
       Use the grouping method.
  3. Check. Is it factored completely? Do the factors multiply back to the original polynomial?

This page titled 7.5: General Strategy for Factoring Polynomials is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.5: General Strategy for Factoring Polynomials** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.5E: Exercises

Practice Makes Perfect

Recognize and Use the Appropriate Method to Factor a Polynomial Completely

In the following exercises, factor completely.

Exercise 7.5E.31

$10x^4+35x^3$

Answer

$5x^3(2x+7)$

Exercise 7.5E.32

$18p^6+24p^3$

Exercise 7.5E.33

$y^2+10y-39$

Answer

$(y-3)(y+13)$

Exercise 7.5E.34

$b^2-17b+60$

Exercise 7.5E.35

$2n^2+13n-7$

Answer

$(2n-1)(n+7)$

Exercise 7.5E.36

$8x^2-9x-3$

Exercise 7.5E.37

a^5+9a^3

Answer

$a^3(a^2+9)$

Exercise 7.5E.38

$75m^3+12m$

Exercise 7.5E.39

$121r^2 - s^2$

Answer

$(11r - s)(11r + s)$

Exercise 7.5E.40

$49b^2 - 36a^2$

Exercise 7.5E.41

$8m^2 - 32$

Answer

$8(m - 2)(m + 2)$

Exercise 7.5E.42

$36q^2 - 100$

Exercise 7.5E.43

$25w^2 - 60w + 36$

Answer

$(5w - 6)^2$

Exercise 7.5E.44

$49b^2 - 112b + 64$

Exercise 7.5E.45

$m^2 + 14mn + 49n^2$

Answer

$(m + 7n)^2$

Exercise 7.5E.46

$64x^2 + 16xy + y^2$

Exercise 7.5E.47

$7b^2 + 7b - 42$

Answer

$7(b + 3)(b - 2)$

Exercise 7.5E.48

$3n^2+30n+72$

Exercise 7.5E.49

$3x^3-81$

Answer

$3(x-3)(x^2+3x+9)$

Exercise 7.5E.50

$5t^3-40$

Exercise 7.5E.51

k^4-16

Answer

$(k-2)(k+2)(k^2+4)$

Exercise 7.5E.52

m^4-81

Exercise 7.5E.53

$15pq-15p+12q-12$

Answer

$3(5p+4)(q-1)$

Exercise 7.5E.54

$12ab-6a+10b-5$

Exercise 7.5E.55

$4x^2+40x+84$

Answer

$4(x+3)(x+7)$

Exercise 7.5E.56

$5q^2-15q-90$

Exercise 7.5E.57

$u^5 + u^2$

Answer

$u^2(u+1)(u^2-u+1)$

Exercise 7.5E.58

$5n^3 + 320$

Exercise 7.5E.59

$4c^2 + 20cd + 81d^2$

Answer

prime

Exercise 7.5E.60

$25x^2 + 35xy + 49y^2$

Exercise 7.5E.61

$10m^4 - 6250$

Answer

$10(m-5)(m+5)(m^2+25)$

Exercise 7.5E.62

$3v^4 - 768$

Everyday Math

Exercise 7.5E.63

Watermelon drop A springtime tradition at the University of California San Diego is the Watermelon Drop, where a watermelon is dropped from the seventh story of Urey Hall.

1. The binomial $-16t^2 + 80$ gives the height of the watermelon t seconds after it is dropped. Factor the greatest common factor from this binomial.
2. If the watermelon is thrown down with initial velocity 8 feet per second, its height after t seconds is given by the trinomial $-16t2 - 8t + 80$

Answer

1. $-16(t^2 - 5)$
2. −8(2t+5)(t−2)

Exercise 7.5E.64

Pumpkin drop A fall tradition at the University of California San Diego is the Pumpkin Drop, where a pumpkin is dropped from the eleventh story of Tioga Hall.

1. The binomial $-16t^2+128$ gives the height of the pumpkin *t* seconds after it is dropped. Factor the greatest common factor from this binomial.
2. If the pumpkin is thrown down with initial velocity 32 feet per second, its height after t seconds is given by the trinomial $-16t^2-32t+128$

Writing Exercises

Exercise 7.5E.65

The difference of squares y^4-625 can be factored as $(y^2-25)(y^2+25)$ *completely* factored. What more must be done to completely factor it?

Exercise 7.5E.66

Of all the factoring methods covered in this chapter (GCF, grouping, undo FOIL, 'ac' method, special products) which is the easiest for you? Which is the hardest? Explain your answers.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
recognize and use the appropriate method to factor a polynomial completely.			

b. Overall, after looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

This page titled 7.5E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.5E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.6: Quadratic Equations

Learning Objectives

By the end of this section, you will be able to:

- Solve quadratic equations by using the Zero Product Property
- Solve quadratic equations factoring
- Solve applications modeled by quadratic equations

Note

Before you get started, take this readiness quiz.

1. Solve: $5y-3=0$.
   If you missed this problem, review Example 2.3.1.
2. Solve: $10a=0$.
   If you missed this problem, review Example 2.2.1.
3. Combine like terms: $12x^2-6x+4x$.
   If you missed this problem, review Example 1.3.37.
4. Factor n^3-9n^2-22n completely.
   If you missed this problem, review Example 7.3.10.

We have already solved linear equations, equations of the form $ax+by=c$. In linear equations, the variables have no exponents. Quadratic equations are equations in which the variable is squared. Listed below are some examples of quadratic equations:

$$x^2+5x+6=0 \quad 3y^2+4y=10 \quad 64u^2-81=0 \quad n(n+1)=42 \tag{7.6.1}$$

The last equation doesn't appear to have the variable squared, but when we simplify the expression on the left we will get n^2+n.

The general form of a quadratic equation is $ax^2+bx+c=0$,with $a\neq 0$.

QUADRATIC EQUATION

An equation of the form $ax^2+bx+c=0$ is called a quadratic equation.

$$a, b, \text{ and } c \text{ are real numbers and } a\neq 0 \tag{7.6.2}$$

To solve **quadratic equations** we need methods different than the ones we used in solving linear equations. We will look at one method here and then several others in a later chapter.

Solve Quadratic Equations Using the Zero Product Property

We will first solve some quadratic equations by using the Zero Product Property. The **Zero Product Property** says that if the product of two quantities is zero, it must be that at least one of the quantities is zero. The only way to get a product equal to zero is to multiply by zero itself.

ZERO PRODUCT PROPERTY

If $a\cdot b=0$, then either $a=0$ or $b=0$ or both.

We will now use the **Zero Product Property**, to solve a quadratic equation.

Example 7.6.1: How to Use the Zero Product Property to Solve a Quadratic Equation

Solve: $(x+1)(x-4)=0$

Solution

Step 1. Set each factor equal to zero.	The product equals zero, so at least one factor must equal zero.	$(x+1)(x-4)=0$ $x+1=0$ or $x-4=0$
Step 2. Solve the linear equations.	Solve each equation.	$x=-1$ or $x=4$

Step 3. Check.	Substitute each solution separately into the original equation.	$x=-1$ $(x+1)(x-4)=0$ $(-1+1)(-1-4)\stackrel{?}{=}0$ $(0)(-5)\stackrel{?}{=}0$ $0=0$ ✓ $x=4$ $(x+1)(x-4)=0$ $(4+1)(4-4)\stackrel{?}{=}0$ $(5)(0)\stackrel{?}{=}0$ $0=0$ ✓

Try It 7.6.2

Solve: $(x-3)(x+5)=0$

Answer

$x=3, x=-5$

Try It 7.6.3

Solve: $(y-6)(y+9)=0$

Answer

$y=6, y=-9$

We usually will do a little more work than we did in this last example to solve the linear equations that result from using the Zero Product Property.

Example 7.6.4

Solve: $(5n-2)(6n-1)=0$

Solution

	$(5n-2)(6n-1)=0$	
Use the Zero Product Property to set each factor to 0.	$5n-2=0$	$6n-1=0$
Solve the equations.	$n=\frac{2}{5}$	$n=\frac{1}{6}$
Check your answers.		
$n=\frac{2}{5}$ $(5n-2)(6n-1)=0$ $\left(5\cdot\frac{2}{5}-2\right)\left(6\cdot\frac{2}{5}-1\right)\stackrel{?}{=}0$ $(2-2)\left(\frac{12}{5}-\frac{5}{5}\right)\stackrel{?}{=}0$ $(0)\left(\frac{7}{5}\right)\stackrel{?}{=}0$ $0=0$ ✓	$n=\frac{1}{6}$ $(5n-2)(6n-1)=0$ $\left(5\cdot\frac{1}{6}-2\right)\left(6\cdot\frac{1}{6}-1\right)\stackrel{?}{=}0$ $\left(\frac{5}{6}-\frac{12}{6}\right)(1-1)\stackrel{?}{=}0$ $\left(-\frac{7}{6}\right)(0)\stackrel{?}{=}0$ $0=0$ ✓	

Try It 7.6.5

Solve: $(3m-2)(2m+1)=0$

Answer

$m=\frac{2}{3}, m=-\frac{1}{2}$

Try It 7.6.6

Solve: $(4p+3)(4p-3)=0$

Answer

$p=-\frac{3}{4}, p=\frac{3}{4}$

Notice when we checked the solutions that each of them made just one factor equal to zero. But the product was zero for both solutions.

Example 7.6.7

Solve: $3p(10p+7)=0$

Solution

	$3p(10p+7)=0$	
Use the Zero Product Property to set each factor to 0.	3p=0	10p+7=0
Solve the equations.	p=0	10p=−7
		$p=-\frac{7}{10}$
Check your answers.		

$p=0$	$p=-\frac{7}{10}$
$3p(10p+7)=0$	$3p(10p+7)=0$
$3\cdot 0(10\cdot 0+7)\stackrel{?}{=}0$	$3\left(-\frac{7}{10}\right)10\left(-\frac{7}{10}\right)+7\stackrel{?}{=}0$
$0(0+7)\stackrel{?}{=}0$	$\left(-\frac{21}{10}\right)(-7+7)\stackrel{?}{=}0$
$0(7)\stackrel{?}{=}0$	$\left(-\frac{21}{10}\right)(0)\stackrel{?}{=}0$
$0=0$ ✓	$0=0$ ✓

Try It 7.6.8

Solve: $2u(5u-1)=0$

Answer

$u=0, u=\frac{1}{5}$

Try It 7.6.9

Solve: $w(2w+3)=0$

Answer

$w=0, w=-\frac{3}{2}$

It may appear that there is only one factor in the next example. Remember, however, that $(y-8)^2$ means $(y-8)(y-8)$.

Example 7.6.10

Solve: $(y-8)^2=0$

Solution

	$(y-8)^2=0$	
Rewrite the left side as a product.	(y−8)(y−8)=0	
Use the Zero Product Property and set each factor to 0.	y−8=0	y−8=0
Solve the equations.	y=8	y=8

When a solution repeats, we call it a double root.		
Check your answer.		
$y = 8$ $(y-8)^2 = 0$ $(8-8)^2 \stackrel{?}{=} 0$ $(0)^2 \stackrel{?}{=} 0$ $0 = 0$ ✓		

? Try It 7.6.11

Solve: $(x+1)^2 = 0$

Answer

$x = 1$

? Try It 7.6.12

Solve: $(v-2)^2 = 0$

Answer

$v = 2$

Solve Quadratic Equations by Factoring

Each of the equations we have solved in this section so far had one side in factored form. In order to use the Zero Product Property, the quadratic equation must be factored, with zero on one side. So we be sure to start with the quadratic equation in standard form, $ax^2 + bx + c = 0$. Then we factor the expression on the left.

✓ Example 7.6.13

Solve: $x^2 + 2x - 8 = 0$

Solution

Step 1. Write the quadratic equation in standard form, $ax^2 + bx + c = 0$.	The equation is already in standard form.	$x^2 + 2x - 8 = 0$
Step 2. Factor the quadratic expression.	Factor $x^2 + 2x - 8$ $(x+4)(x-2)$	$(x+4)(x-2) = 0$
Step 3. Use the Zero Product Property.	Set each factor equal to zero.	$x + 4 = 0$ or $x - 2 = 0$
Step 4. Solve the linear equations.	We have two linear equations.	$x = -4$ or $x = 2$
Step 5. Check.	Substitute each solution separately into the original equation.	$x^2 + 2x - 8 = 0$ $x = -4$ $(-4)^2 - 2(-4) - 8 \stackrel{?}{=} 0$ $16 + (-8) - 8 \stackrel{?}{=} 0$ $0 = 0$ ✓ $x^2 + 2x - 8 = 0$ $x = 2$ $2^2 - 2(2) - 8 \stackrel{?}{=} 0$ $4 + 4 - 8 \stackrel{?}{=} 0$ $0 = 0$ ✓

Try It 7.6.14

Solve: $x^2 - x - 12 = 0$

Answer

$x = 4, x = -3$

Try It 7.6.15

Solve: $b^2 + 9b + 14 = 0$

Answer

$b = -2, b = -7$

SOLVE A QUADRATIC EQUATION BY FACTORING.

1. Write the quadratic equation in standard form, $ax^2 + bx + c = 0$.
2. Factor the quadratic expression.
3. Use the Zero Product Property.
4. Solve the linear equations.
5. Check.

Before we factor, we must make sure the quadratic equation is in standard form.

Example 7.6.16

Solve: $2y^2 = 13y + 45$

Solution

	$2y^2 = 13y + 45$	
Write the quadratic equation in standard form.	$2y^2 - 13y - 45 = 0$	
Factor the quadratic expression.	$(2y + 5)(y - 9) = 0$	
Use the Zero Product Property to set each factor to 0.	$2y + 5 = 0$	$y - 9 = 0$
Solve each equation.	$y = -\frac{5}{2}$	$y = 9$
Check your answers.		
$y = -\frac{5}{2}$ $2y^2 = 13y + 45$ $2\left(-\frac{5}{2}\right)^2 \stackrel{?}{=} 13\left(-\frac{5}{2}\right) + 45$ $2\left(\frac{25}{4}\right) \stackrel{?}{=} \left(-\frac{65}{2}\right) + \frac{90}{2}$ $\frac{25}{2} = \frac{25}{2}$ ✓	$y = 9$ $2y^2 = 13y + 45$ $2(9)^2 \stackrel{?}{=} 13(9) + 45$ $2(81) \stackrel{?}{=} 117 + 45$ $162 = 162$ ✓	

Try It 7.6.17

Solve: $3c^2 = 10c - 8$

Answer

$c = 0, c = \frac{4}{3}$

Try It 7.6.18

Solve: $2d^2 - 5d = 3$

Answer

$d = 3, d = -\frac{1}{2}$

Example 7.6.19

Solve: $5x^2 - 13x = 7x$

Solution

	$5x^2 - 13x = 7x$	
Write the quadratic equation in standard form.	$5x^2 - 20x = 0$	
Factor the left side of the equation.	$5x(x-4) = 0$	
Use the Zero Product Property to set each factor to 0.	$5x = 0$	$x - 4 = 0$
Solve each equation.	$x = 0$	$x = 4$
Check your answers.		

$x = 0$

$5x^2 - 13x = 7x$

$5(0)^2 - 13(0) \stackrel{?}{=} 7(0)$

$0 - 0 \stackrel{?}{=} 0$

$0 = 0$ ✓

$x = 4$

$5x^2 - 13x = 7x$

$5(4)^2 - 13(4) \stackrel{?}{=} 7(4)$

$5(16) - 52 \stackrel{?}{=} 28$

$28 = 28$ ✓

Try It 7.6.20

Solve: $6a^2 + 9a = 3a$

Answer

$a = 0, a = -1$

Try It 7.6.21

Solve: $45b^2 - 2b = -17b$

Answer

$b = 0, b = -\frac{1}{3}$

Solving quadratic equations by factoring will make use of all the factoring techniques you have learned in this chapter! Do you recognize the special product pattern in the next example?

Example 7.6.22

Solve: $144q^2 = 25$

Solution

	$144q^2$	$=$	25			
Write the quadratic equation in standard form.	$144q^2 - 25$	$=$	0			
Factor. It is a difference of squares.	$(12q-5)(12q+5)$	$=$	0			
Use the Zero Product Property to set each factor to 0.	$12q-5$	$=$	0	$12q+5$	$=$	0
Solve each equation.	$12q$	$=$	5	$12q$	$=$	-5
	q	$=$	$\frac{5}{12}$	q	$=$	$-\frac{5}{12}$

Check your answers.

Try It 7.6.23

Solve: $25p^2 = 49$

Answer

$p = \frac{7}{5}, p = -\frac{7}{5}$

Try It 7.6.24

Solve: $36x^2 = 121$

Answer

$x = \frac{11}{6}, x = -\frac{11}{6}$

The left side in the next example is factored, but the right side is not zero. In order to use the Zero Product Property, one side of the equation must be zero. We'll multiply the factors and then write the equation in standard form.

Example 7.6.25

Solve: $(3x-8)(x-1) = 3x$

Solution

	$(3x-8)(x-1) = 3x$
Multiply the binomials.	$3x^2 - 11x + 8 = 3x$
Write the quadratic equation in standard form.	$3x^2 - 14x + 8 = 0$
Factor the trinomial.	$(3x-2)(x-4) = 0$
Use the Zero Product Property to set each factor to 0.	$3x-2=0 \quad x-4=0$
Solve each equation.	$3x = 2 \quad x = 4$
	$x = \frac{2}{3}$
Check your answers.	The check is left to you!

Try It 7.6.26

Solve: $(2m+1)(m+3) = 12m$

Answer

$m = 1, m = \frac{3}{2}$

Try It 7.6.27

Solve: $(k+1)(k-1) = 8$

Answer

$k = 3, k = -3$

The Zero Product Property also applies to the product of three or more factors. If the product is zero, at least one of the factors must be zero. We can solve some equations of degree more than two by using the Zero Product Property, just like we solved quadratic equations.

Example 7.6.28

Solve: $9m^3 + 100m = 60m^2$

Solution

	$9m^3 + 100m$	$=$	$60m^2$					
Bring all the terms to one side so that the other side is zero.	$9m^3 - 60m^2 + 100m$	$=$	0					
Factor the greatest common factor first.	$m\left(9m^2 - 60m + 100\right)$	$=$	0					
Factor the trinomial.	$m(3m-10)(3m-10)$	$=$	0					
Use the Zero Product Property to set each factor to 0.	m	$=$	0	$3m-10$	$=$	0	$3m-10$	$=$ 0
Solve each equation.	m	$=$	0	m	$=$	$\frac{10}{3}$	m	$=$ $\frac{10}{3}$
Check your answers.	The check is left to you.							

Try It 7.6.29

Solve: $8x^3 = 24x^2 - 18x$

Answer

$x = 0, x = \frac{3}{2}$

Try It 7.6.30

Solve: $16y^2 = 32y^3 + 2y$

Answer

$y = 0, y = \frac{1}{4}$

When we factor the quadratic equation in the next example we will get three factors. However the first factor is a constant. We know that factor cannot equal 0.

Example 7.6.31

Solve: $4x^2 = 16x + 84$

Solution

	$4x^2$	$=$	$16x+84$					
Write the quadratic equation in standard form.	$4x^2-16x-84$	$=$	0					
Factor the greatest common factor first.	$4\left(x^2-4x-21\right)$	$=$	0					
Factor the trinomial.	$4(x-7)(x+3)$	$=$	0					
Use the Zero Product Property to set each factor to 0.	4	$\neq$	0	$x-7$	$=$	0	$x+3$	$=$ 0
Solve each equation.	4	$\neq$	0	x	$=$	7	x	$=$ -3
Check your answers.	The check is left to you.							

Try It 7.6.32

Solve: $18a^2 - 30 = -33a$

Answer

$a = -\frac{5}{2}, a = \frac{2}{3}$

Try It 7.6.33

Solve: $123b = -6 - 60b^2$

Answer

$b = 2, b = \frac{1}{20}$

Solve Applications Modeled by Quadratic Equations

The problem solving strategy we used earlier for applications that translate to linear equations will work just as well for applications that translate to quadratic equations. We will copy the problem solving strategy here so we can use it for reference.

USE A PROBLEM-SOLVING STRATEGY TO SOLVE WORD PROBLEMS

1. **Read** the problem. Make sure all the words and ideas are understood.
2. **Identify** what we are looking for.
3. **Name** what we are looking for. Choose a variable to represent that quantity.
4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information. Then, translate the English sentence into an algebra equation.
5. **Solve** the equation using good algebra techniques.
6. **Check** the answer in the problem and make sure it makes sense.
7. **Answer** the question with a complete sentence.

We will start with a number problem to get practice translating words into a quadratic equation.

Example 7.6.34

The product of two consecutive integers is 132. Find the integers.

Solution

Step 1. Read the problem.	
Step 2. Identify what we are looking for.	We are looking for two consecutive integers.
Step 3. Name what we are looking for.	Let $n =$ the first integer $n+1 =$ the next consecutive integer
Step 4. Translate into an equation. Restate the problem in a sentence.	The product of the two consecutive integers is 132.
Translate to an equation.	The first integer times the next integer is 132. $n(n+1) = 132$
Step 5. Solve the equation.	$n^2 + n = 132$
Bring all the terms to one side.	$n^2 + n - 132 = 0$
Factor the trinomial.	$(n-11)(n+12) = 0$
Use the zero product property.	$n-11=0 \quad n+12=0$
Solve the equations.	$n = 11 \quad n = -12$

Try It 7.6.35

The product of two consecutive integers is 240. Find the integers.

Answer

$-15, -16$ and $15, 16$

Try It 7.6.36

The product of two consecutive integers is 420. Find the integers.

Answer

$-21, -20$ and $20, 21$


Were you surprised by the pair of negative integers that is one of the solutions to the previous example? The product of the two positive integers and the product of the two negative integers both give 132.


In some applications, negative solutions will result from the algebra, but will not be realistic for the situation.

Example 7.6.37

A rectangular garden has an area of 15 square feet. The length of the garden is two feet more than the width. Find the length and width of the garden.

Solution

Step 1. Read the problem. In problems involving geometric figures, a sketch can help you visualize the situation.		
Step 2. Identify what you are looking for.	We are looking for the length and width.	
Step 3. Name what you are looking for. The length is two feet more than width.	Let W = the width of the garden. $W + 2$ = the length of the garden	
Step 4. Translate into an equation. Restate the important information in a sentence.	The area of the rectangular garden is 15 square feet.	
Use the formula for the area of a rectangle.	$A = L \cdot W$	
Substitute in the variables.	$15 = (W+2)W$	
Step 5. Solve the equation. Distribute first.	$15 = W^2 + 2W$	
Get zero on one side.	$0 = W^2 + 2W - 15$	
Factor the trinomial.	$0 = (W+5)(W-3)$	
Use the Zero Product Property.	$0 = W + 5$	$0 = W - 3$
Solve each equation.	$-5 = W$	$3 = W$

Since W is the width of the garden, it does not make sense for it to be negative. We eliminate that value for W.	$W = -5$ cannot be the width, since it's negative. $W = 3$	$3 = W$ Width is 3 feet.
Find the value of the length.	$\text{length} = W + 2$	
	$\text{length} = 3 + 2$	
	$\text{length} = 5$	Length is 5 feet.
Step 6. Check the answer. Does the answer make sense?		
 $A = L \cdot W$ $A = 3 \cdot 5$ $A = 15$		
	Yes, this makes sense.	
Step 7. Answer the question.	The width of the garden is 3 feet and the length is 5 feet.	

? Try It 7.6.38

A rectangular sign has an area of 30 square feet. The length of the sign is one foot more than the width. Find the length and width of the sign.

Answer

55 feet and 66 feet


? Try It 7.6.39

A rectangular patio has an area of 180 square feet. The width of the patio is three feet less than the length. Find the length and width of the patio.

Answer

12 feet and 15 feet


In an earlier chapter, we used the Pythagorean Theorem $\left(a^2 + b^2 = c^2\right)$. It gave the relation between the legs and the hypotenuse of a right triangle.



We will use this formula to in the next example.

✓ Example 7.6.40


Justine wants to put a deck in the corner of her backyard in the shape of a right triangle, as shown below. The hypotenuse will be 17 feet long. The length of one side will be 7 feet less than the length of the other side. Find the lengths of the sides of the deck.



Solution

Step 1. Read the problem.			
Step 2. Identify what you are looking for.	We are looking for the lengths of the sides of the deck.		
Step 3. Name what you are looking for. One side is 7 less than the other.	Let x = length of a side of the deck $x - 7$ = length of other side		

https://math.libretexts.org/@go/page/30559

Step 4. Translate into an equation. Since this is a right triangle we can use the Pythagorean Theorem.	$a^2+b^2=c^2$		
Substitute in the variables.	$x^2+(x-7)^2=17^2$		
Step 5. Solve the equation.	$x^2+x^2-14x+49=289$		
Simplify.	$2x^2-14x+49=289$		
It is a quadratic equation, so get zero on one side.	$2x^2-14x-240=0$		
Factor the greatest common factor.	$2\left(x^2-7x-120\right)=0$		
Factor the trinomial.	$2(x-15)(x+8)=0$		
Use the Zero Product Property.	$2\neq 0$	$x-15=0$	$x+8=0$
Solve.	$2\neq 0$	$x=15$	$x=-8$
Since x is a side of the triangle, $x=-8$ does not make sense.	$2\neq 0$	$x=15$	$x=-8$
Find the length of the other side.			
If the length of one side is		$x=15$	
then the length of the other side is		$x-7$	
		$15-7=8$	
	8 is the length of the other side.		
Step 6. Check the answer. Do these numbers make sense?			
			
Step 7. Answer the question.	The sides of the deck are 8, 15, and 17 feet.		

? Try It 7.6.41

A boat's sail is a right triangle. The length of one side of the sail is 7 feet more than the other side. The hypotenuse is 13. Find the lengths of the two sides of the sail.

Answer

5 feet and 12 feet

? Try It 7.6.42

A meditation garden is in the shape of a right triangle, with one leg 7 feet. The length of the hypotenuse is one more than the length of one of the other legs. Find the lengths of the hypotenuse and the other leg.

Answer

24 feet and 25 feet

Key Concepts

- **Zero Product Property** If $a\cdot b=0$, then either a=0 or b=0 or both. See Example.
- **Solve a quadratic equation by factoring** To solve a quadratic equation by factoring: See Example.
  1. Write the quadratic equation in standard form, $ax^2+bx+c=0$.
  2. Factor the quadratic expression.
  3. Use the Zero Product Property.

4. Solve the linear equations.
5. Check.

- **Use a problem solving strategy to solve word problems** See Example.
  1. **Read** the problem. Make sure all the words and ideas are understood.
  2. **Identify** what we are looking for.
  3. **Name** what we are looking for. Choose a variable to represent that quantity.
  4. **Translate** into an equation. It may be helpful to restate the problem in one sentence with all the important information. Then, translate the English sentence into an algebra equation.
  5. **Solve** the equation using good algebra techniques.
  6. **Check** the answer in the problem and make sure it makes sense.
  7. **Answer** the question with a complete sentence.

Glossary

quadratic equations

are equations in which the variable is squared.

Zero Product Property

The Zero Product Property states that, if the product of two quantities is zero, at least one of the quantities is zero.

This page titled 7.6: Quadratic Equations is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.6: Quadratic Equations** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

7.6E: Exercises

Practice Makes Perfect

Use the Zero Product Property

In the following exercises, solve.

Exercise 1

$(x-3)(x+7)=0$

Answer

$x=3$, $x=-7$ so the solution set is: $\{3,-7\}$

Exercise 2

$(y-11)(y+1)=0$

Exercise 3

$(3a-10)(2a-7)=0$

Answer

$a=\frac{10}{3}$, $a=\frac{7}{2}$ so the solution set is: $\left\{\frac{10}{3},\frac{7}{2}\right\}$

Exercise 4

$(5b+1)(6b+1)=0$

Exercise 5

$6m(12m-5)=0$

Answer

$m=0$, $m=\frac{5}{12}$ so the solution set is: $\left\{0,\frac{5}{12}\right\}$

Exercise 6

$2x(6x-3)=0$

Exercise 7

$(y-3)^2=0$

Answer

$y=3$ so the solution set is: $\{3\}$

Exercise 8

$(b+10)^2=0$

Exercise 9

$(2x-1)^2=0$

Answer

$x=\frac{1}{2}$ so the solution set is: $\left\{\frac{1}{2}\right\}$

Exercise 10

$(3y+5)^2=0$

Solve Quadratic Equations by Factoring

In the following exercises, solve.

Exercise 11

$x^2+7x+12=0$

Answer

$x=-3$, $x=-4$ so the solution set is: $\{-3,-4\}$

Exercise 12

$y^2-8y+15=0$

Exercise 13

$5a^2-26a=24$

Answer

$a=-\frac{4}{5}$, $a=6$ so the solution set is: $\left\{-\frac{4}{5},6\right\}$

Exercise 14

$4b^2+7b=-3$

Exercise 15

$4m^2=17m-15$

Answer

$m=\frac{5}{4}$, $m=3$ so the solution set is: $\left\{\frac{5}{4},3\right\}$

Exercise 16

$n^2 = 5 - 6n$

Exercise 17

$7a^2 + 14a = 7a$

Answer

$a = -1$, $a = 0$ so the solution set is: $\{-1, 0\}$

Exercise 18

$12b^2 - 15b = -9b$

Exercise 19

$49m^2 = 144$

Answer

$m = \frac{12}{7}$, $m = -\frac{12}{7}$ so the solution set is: $\left\{-\frac{12}{7}, \frac{12}{7}\right\}$

Exercise 20

$625 = x^2$

Exercise 21

$(y-3)(y+2) = 4y$

Answer

$y = -1$, $y = 6$ so the solution set is: $\{-1, 6\}$

Exercise 22

$(p-5)(p+3) = -7$

Exercise 23

$(2x+1)(x-3) = -4x$

Answer

$x = \frac{3}{2}$, $x = -1$ so the solution set is: $\left\{-1, \frac{3}{2}\right\}$

Exercise 24

$(x+6)(x-3) = -8$

Exercise 25

$16p^3 = 24p^2 - 9p$

Answer

$p = 0$, $p = \frac{3}{4}$ so the solution set is: $\left\{0, \frac{3}{4}\right\}$

Exercise 26

$m^3 - 2m^2 = -m$

Exercise 27

$20x^2 - 60x = -45$

Answer

$x = \frac{3}{2}$ so the solution set is: $\left\{\frac{3}{2}\right\}$

Exercise 28

$3y^2 - 18y = -27$

Solve Applications Modeled by Quadratic Equations

In the following exercises, solve.

Exercise 29

The product of two consecutive integers is 56. Find the integers.

Answer

7 and 8; −8 and −7

Exercise 30

The product of two consecutive integers is 42. Find the integers.

Exercise 31

The area of a rectangular carpet is 28 square feet. The length is three feet more than the width. Find the length and the width of the carpet.

Answer

4 feet and 7 feet

Exercise 32

A rectangular retaining wall has area 15 square feet. The height of the wall is two feet less than its length. Find the height and the length of the wall.

Exercise 33

A pennant is shaped like a right triangle, with hypotenuse 10 feet. The length of one side of the pennant is two feet longer than the length of the other side. Find the length of the two sides of the pennant.

Answer

6 feet and 8 feet

Exercise 34

A reflecting pool is shaped like a right triangle, with one leg along the wall of a building. The hypotenuse is 9 feet longer than the side along the building. The third side is 7 feet longer than the side along the building. Find the lengths of all three sides of the reflecting pool.

Mixed Practice

In the following exercises, solve.

Exercise 35

(x+8)(x−3)=0

Answer

$x = -8,\ x = 3$ so the solution set is: $\{-8, 3\}$

Exercise 36

(3y−5)(y+7)=0

Exercise 37

$p^2 + 12p + 11 = 0$

Answer

$p = -1,\ p = -11$ so the solution set is: $\{-11, -1\}$

Exercise 38

$q^2 - 12q - 13 = 0$

Exercise 39

$m^2 = 6m + 16$

Answer

$m = -2,\ m = 8$ so the solution set is: $\{-2, 8\}$

Exercise 40

$4n^2 + 19n = 5$

Exercise 41

$a^3 - a^2 - 42a = 0$

Answer

$a = 0,\ a = -6,\ a = 7$ so the solution set is: $\{-6, 0, 7\}$

Exercise 42

$4b^2 - 60b + 224 = 0$

Exercise 43

The product of two consecutive integers is 110. Find the integers.

Answer

10 and 11; −11 and −10

Exercise 44

The length of one leg of a right triangle is three more than the other leg. If the hypotenuse is 15, find the lengths of the two legs.

Everyday Math

Exercise 45

Area of a patio If each side of a square patio is increased by 4 feet, the area of the patio would be 196 square feet. Solve the equation (s+4)2=196(s+4)2=196 for *s* to find the length of a side of the patio.

Answer

10 feet

Exercise 46

Watermelon drop A watermelon is dropped from the tenth story of a building. Solve the equation −16t2+144=0−16t2+144=0 for tt to find the number of seconds it takes the watermelon to reach the ground.

Writing Exercises

Exercise 47

Explain how you solve a quadratic equation. How many answers do you expect to get for a quadratic equation?

Answer

Answers may vary for the explanation. You should expect no more than 2 solutions for a quadratic equation. Often it has two solutions, but sometimes, it can have one repeated solution or even no solution.

? Exercise 48

Give an example of a quadratic equation that has a GCF and none of the solutions to the equation is zero.

Self Check

a. After completing the exercises, use this checklist to evaluate your mastery of the objectives of this section.

I can...	Confidently	With some help	No-I don't get it!
solve quadratic equations by using the Zero Product Property.			
solve quadratic equations by factoring.			
solve applications modeled by quadratic equations.			

b. Overall, after looking at the checklist, do you think you are well-prepared for the next section? Why or why not?

This page titled 7.6E: Exercises is shared under a CC BY license and was authored, remixed, and/or curated by OpenStax.

- **7.6E: Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.

Chapter 7 Review Exercises

Chapter 7 Review Exercises

7.1 Greatest Common Factor and Factor by Grouping

Find the Greatest Common Factor of Two or More Expressions

In the following exercises, find the greatest common factor.

Exercise 1

42, 60

Answer

6

Exercise 2

450, 420

Exercise 3

90, 150, 105

Answer

15

Exercise 4

60, 294, 630

Factor the Greatest Common Factor from a Polynomial

In the following exercises, factor the greatest common factor from each polynomial.

Exercise 5

$24x - 42$

Answer

$6(4x - 7)$

Exercise 6

$35y + 84$

Exercise 7

$15m^4 + 6m^2n$

Answer

$3m^2(5m2 + 2n)$

? Exercise 8

$24pt^4 + 16t^7$

Factor by Grouping

In the following exercises, factor by grouping.

? Exercise 9

$ax - ay + bx - by$

Answer

$(a+b)(x-y)$

? Exercise 10

$x^2y - xy^2 + 2x - 2y$

? Exercise 11

$x^2 + 7x - 3x - 21$

Answer

$(x-3)(x+7)$

? Exercise 12

$4x^2 - 16x + 3x - 12$

? Exercise 13

$m^3 + m^2 + m + 1$

Answer

$(m^2+1)(m+1)$

? Exercise 14

$5x - 5y - y + x$

7.2 Factor Trinomials of the form $x^2 + bx + c$

Factor Trinomials of the Form $x^2 + bx + c$

In the following exercises, factor each trinomial of the form $x^2 + bx + c$

? Exercise 15

$u^2 + 17u + 72$

Answer

$(u+8)(u+9)$

Exercise 16

$a^2+14a+33$

Exercise 17

$k^2-16k+60$

Answer

$(k-6)(k-10)$

Exercise 18

$r^2-11r+28$

Exercise 19

y^2+6y-7

Answer

$(y+7)(y-1)$

Exercise 20

$m^2+3m-54$

Exercise 21

s^2-2s-8

Answer

$(s-4)(s+2)$

Exercise 22

$x^2-3x-10$

Factor Trinomials of the Form $x^2+bxy+cy^2$

In the following examples, factor each trinomial of the form $x^2+bxy+cy^2$

Exercise 23

$x^2+12xy+35y^2$

Answer

$(x+5y)(x+7y)$

Exercise 24

$u^2+14uv+48v^2$

Exercise 25

$a^2+4ab-21b^2$

Answer

$(a+7b)(a-3b)$

Exercise 26

$p^2-5pq-36q^2$

7.3 Factoring Trinomials of the form ax^2+bx+c

Recognize a Preliminary Strategy to Factor Polynomials Completely

In the following exercises, identify the best method to use to factor each polynomial.

Exercise 27

$y^2-17y+42$

Answer

Undo FOIL

Exercise 28

$12r^2+32r+5$

Exercise 29

$8a^3+72a$

Answer

Factor the GCF

Exercise 30

$4m-mn-3n+12$

Factor Trinomials of the Form ax^2+bx+c with a GCF

In the following exercises, factor completely.

Exercise 31

$6x^2+42x+60$

Answer

$6(x+2)(x+5)$

Exercise 32

$8a^2+32a+24$

Exercise 33

$3n^4 - 12n^3 - 96n^2$

Answer

$3n^2(n-8)(n+4)$

Exercise 34

$5y^4 + 25y^2 - 70y$

Factor Trinomials Using the "ac" Method

In the following exercises, factor.

Exercise 35

$2x^2 + 9x + 4$

Answer

$(x+4)(2x+1)$

Exercise 36

$3y^2 + 17y + 10$

Exercise 37

$18a^2 - 9a + 1$

Answer

$(3a-1)(6a-1)$

Exercise 38

$8u^2 - 14u + 3$

Exercise 39

$15p^2 + 2p - 8$

Answer

$(5p+4)(3p-2)$

Exercise 40

$15x^2 + 6x - 2$

Exercise 41

$40s^2 - s - 6$

Answer

$(5s-2)(8s+3)$

Exercise 42

$20n^2-7n-3$

Factor Trinomials with a GCF Using the “ac” Method

In the following exercises, factor.

Exercise 43

$3x^2+3x-36$

Answer

$3(x+4)(x-3)$

Exercise 44

$4x^2+4x-8$

Exercise 45

$60y^2-85y-25$

Answer

$5(4y+1)(3y-5)$

Exercise 46

$18a^2-57a-21$

7.4 Factoring Special Products

Factor Perfect Square Trinomials

In the following exercises, factor.

Exercise 47

$25x^2+30x+9$

Answer

$(5x+3)^2$

Exercise 48

$16y^2+72y+81$

Exercise 49

$36a^2-84ab+49b^2$

Answer

$(6a-7b)^2$

Exercise 50

$64r^2-176rs+121s^2$

Exercise 51

$40x^2+360x+810$

Answer

$10(2x+9)^2$

Exercise 52

$75u^2+180u+108$

Exercise 53

$2y^3-16y^2+32y$

Answer

$2y(y-4)^2$

Exercise 54

$5k^3-70k^2+245k$

Factor Differences of Squares

In the following exercises, factor.

Exercise 55

$81r^2-25$

Answer

$(9r-5)(9r+5)$

Exercise 56

$49a^2-144$

Exercise 57

$169m^2-n^2$

Answer

$(13m+n)(13m-n)$

Exercise 58

$64x^2 - y^2$

Exercise 59

$25p^2 - 1$

Answer

$(5p-1)(5p+1)$

Exercise 60

$1 - 16s^2$

Exercise 61

$9 - 121y^2$

Answer

$(3+11y)(3-11y)$

Exercise 62

$100k^2 - 81$

Exercise 64

$20x^2 - 125$

Answer

$5(2x-5)(2x+5)$

Exercise 64

$18y^2 - 98$

Exercise 65

$49u^3 - 9u$

Answer

$u(7u+3)(7u-3)$

Exercise 66

$169n^3 - n$

Factor Sums and Differences of Cubes

In the following exercises, factor.

Exercise 67

$a^3 - 125$

Answer

$(a-5)(a^2+5a+25)$

Exercise 68

$b^3 - 216$

Exercise 69

$2m^3 + 54$

Answer

$2(m+3)(m^2-3m+9)$

Exercise 70

$81x^3 + 3$

7.5 General Strategy for Factoring Polynomials

Recognize and Use the Appropriate Method to Factor a Polynomial Completely

In the following exercises, factor completely.

Exercise 71

$24x^3 + 44x^2$

Answer

$4x^2(6x+11)$

Exercise 72

$24a^4 - 9a^3$

Exercise 73

$16n^2 - 56mn + 49m^2$

Answer

$(4n-7m)^2$

Exercise 74

$6a^2 - 25a - 9$

Exercise 75

$5r^2 + 22r - 48$

Answer

(r+6)(5r−8)

Exercise 76

$5u^4 - 45u^2$

Exercise 77

$n^4 - 81$

Answer

$(n^2 + 9)(n + 3)(n - 3)$

Exercise 78

$64j^2 + 225$

Exercise 79

$5x^2 + 5x - 60$

Answer

$5(x - 3)(x + 4)$

Exercise 80

$b^3 - 64$

Exercise 81

$m^3 + 125$

Answer

$(m + 5)(m^2 - 5m + 25)$

Exercise 82

$2b^2 - 2bc + 5cb - 5c^2$

7.6 Quadratic Equations

Use the Zero Product Property

In the following exercises, solve.

Exercise 83

$(a-3)(a+7)=0$

Answer

$a=3, a=-7$

Exercise 84

$(b-3)(b+10)=0$

Exercise 85

$3m(2m-5)(m+6)=0$

Answer

$m=0, m=-6, m=\frac{5}{2}$

Exercise 86

$7n(3n+8)(n-5)=0$

Solve Quadratic Equations by Factoring

In the following exercises, solve.

Exercise 87

$x^2+9x+20=0$

Answer

$x=-4, x=-5$

Exercise 88

$y^2-y-72=0$

Exercise 89

$2p^2-11p=40$

Answer

$p=-\frac{5}{2}$, p=8

Exercise 90

$q^3+3q^2+2q=0$

Exercise 91

$144m^2-25=0$

Answer

$m = \frac{5}{12}, m = -\frac{5}{12}$

Exercise 92

$4n^2 = 36$

Solve Applications Modeled by Quadratic Equations

In the following exercises, solve.

Exercise 93

The product of two consecutive numbers is 462.

Answer

$-21, -22$

21, 22

Exercise 94

The area of a rectangular shaped patio 400 square feet. The length of the patio is 99 feet more than its width. Find the length and width.

Practice Test

In the following exercises, find the Greatest Common Factor in each expression.

Exercise 95

$14y - 42$

Answer

$7(y - 6)$

Exercise 96

$-6x^2 - 30x$

Exercise 97

$80a^2 + 120a^3$

Answer

$40a^2(2 + 3a)$

Exercise 98

$5m(m - 1) + 3(m - 1)$

In the following exercises, factor completely.

Exercise 99

$x^2+13x+36$

Answer

$(x+7)(x+6)$

Exercise 100

$p^2+pq-12q^2$

Exercise 101

$3a^3-6a^2-72a$

Answer

$3a(a+4)(a-6)$

Exercise 102

$s^2-25s+84$

Exercise 103

$5n^2+30n+45$

Answer

$5(n+3)^2$

Exercise 104

$64y^2-49$

Exercise 105

$xy-8y+7x-56$

Answer

$(x-8)(y+7)$

Exercise 106

$40r^2+810$

Exercise 107

$9s^2-12s+4$

Answer

$(3s-2)^2$

Exercise 1008

$n^2+12n+36$

Exercise 109

$100-a^2$

Answer

$(10-a)(10+a)$

Exercise 110

$6x^2-11x-10$

Exercise 111

$3x^2-75y^2$

Answer

$3(x+5y)(x-5y)$

Exercise 112

$c^3-1000d^3$

Exercise 113

$ab-3b-2a+6$

Answer

$(a-3)(b-2)$

Exercise 114

$6u^2+3u-18$

Exercise 115

$8m^2+22m+5$

Answer

$(4m+1)(2m+5)$

In the following exercises, solve.

Exercise 116

$x^2+9x+20=0$

Exercise 117

$y^2 = y + 132$

Answer

$y = -11, y = 12$

Exercise 118

$5a^2 + 26a = 24$

Exercise 119

$9b^2 - 9 = 0$

Answer

$b = 1, b = -1$

Exercise 120

$16 - m^2 = 0$

Exercise 121

$4n^2 + 19n + 21 = 0$

Answer

$n = -\frac{7}{4}$, n=−3

Exercise 122

$(x - 3)(x + 2) = 6$

Exercise 123

The product of two consecutive integers is 156.

Answer

12 and 13; −13 and −12

Exercise 124

The area of a rectangular place mat is 168 square inches. Its length is two inches longer than the width. Find the length and width of the placemat.

Chapter 7 Review Exercises is shared under a not declared license and was authored, remixed, and/or curated by LibreTexts.

- **Chapter 7 Review Exercises** by OpenStax is licensed CC BY 4.0. Original source: https://openstax.org/details/books/elementary-algebra-2e.