

FIRST-SEMESTER ABSTRACT ALGEBRA: A STRUCTURAL APPROACH



Jessica K. Sklar
Pacific Lutheran University

Pacific Lutheran University

First-Semester Abstract Algebra: A Structural
Approach (Sklar)

Jessica K. Sklar

This text is disseminated via the Open Education Resource (OER) LibreTexts Project (<https://LibreTexts.org>) and like the thousands of other texts available within this powerful platform, it is freely available for reading, printing, and "consuming."

The LibreTexts mission is to bring together students, faculty, and scholars in a collaborative effort to provide an accessible, and comprehensive platform that empowers our community to develop, curate, adapt, and adopt openly licensed resources and technologies; through these efforts we can reduce the financial burden borne from traditional educational resource costs, ensuring education is more accessible for students and communities worldwide.

Most, but not all, pages in the library have licenses that may allow individuals to make changes, save, and print this book. Carefully consult the applicable license(s) before pursuing such effects. Instructors can adopt existing LibreTexts texts or Remix them to quickly build course-specific resources to meet the needs of their students. Unlike traditional textbooks, LibreTexts' web based origins allow powerful integration of advanced features and new technologies to support learning.



LibreTexts is the adaptable, user-friendly non-profit open education resource platform that educators trust for creating, customizing, and sharing accessible, interactive textbooks, adaptive homework, and ancillary materials. We collaborate with individuals and organizations to champion open education initiatives, support institutional publishing programs, drive curriculum development projects, and more.

The LibreTexts libraries are Powered by [NICE CXone Expert](#) and were supported by the Department of Education Open Textbook Pilot Project, the California Education Learning Lab, the UC Davis Office of the Provost, the UC Davis Library, the California State University Affordable Learning Solutions Program, and Merlot. This material is based upon work supported by the National Science Foundation under Grant No. 1246120, 1525057, and 1413739.

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation nor the US Department of Education.

Have questions or comments? For information about adoptions or adaptations contact info@LibreTexts.org or visit our main website at <https://LibreTexts.org>.

The LibreTexts name, logo, and book covers are trademarked by LibreTexts, Inc. (a 501(c)(3) not-for-profit organization) and protected against unauthorized use.

This text was compiled on 09/20/2026

TABLE OF CONTENTS

Licensing

1: Preliminaries

- 1.1: Sets
- 1.2: Functions
- 1.3: Cardinality
- 1.4: Exercises

2: Groups

- 2.1: Binary Operations and Structures
- 2.2: Exercises, Part I
- 2.3: The Definition of a Group
- 2.4: Examples of Groups/Nongroups, Part I
- 2.5: Group Conventions and Properties
- 2.6: Examples of Groups/Nongroups, Part II
- 2.7: Summaries of Groups We've Seen
- 2.8: Exercises, Part II

3: Homomorphisms and Isomorphisms

- 3.1: Groups of Small Order
- 3.2: Definitions of Homomorphisms and Isomorphisms
- 3.3: Isomorphic Groups
- 3.4: Exercises

4: Subgroups

- 4.1: Introduction to Subgroups
- 4.2: Subgroup Proofs and Lattices
- 4.3: Exercises

5: Cyclic Groups

- 5.1: Introduction to Cyclic Groups
- 5.2: The Subgroup Lattices of Cyclic Groups
- 5.3: Exercises

6: Permutation and Dihedral Groups

- 6.1: Introduction to Permutation Groups
- 6.2: Symmetric Groups
- 6.3: Alternating Groups
- 6.4: Cayley's Theorem
- 6.5: Dihedral Groups
- 6.6: Exercises

7: The Wonderful World of Cosets

- 7.1: Partitions of and Equivalence Relations on Sets
- 7.2: Introduction to Cosets and Normal Subgroups
- 7.3: The Index of a Subgroup and Lagrange's Theorem
- 7.4: Exercises

8: Factor Groups

- 8.1: Motivation
- 8.2: Focusing on Normal Subgroups
- 8.3: Introduction to Factor Groups
- 8.4: Exercises

9: The Isomorphism Theorem

- 9.1: The First Isomorphism Theorem
- 9.2: The Second and Third Isomorphism Theorems
- 9.3: Exercises

Index

Glossary

Bibliography

Detailed Licensing

Notation

Solutions to Exercises

Licensing

A detailed breakdown of this resource's licensing can be found in **Back Matter/Detailed Licensing**.

1: PRELIMINARIES

1.1: Sets

1.2: Functions

1.3: Cardinality

1.4: Exercises

This page titled [1: Preliminaries](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

1.1: Sets

We start off by providing a “definition” of a basic mathematical structure to which we will soon add bells and whistles. We use quotes here because what follows doesn’t have the precision we usually require when defining a mathematical object.

Definition: Set

A **set** is an (unordered) collection of objects.

This is only sort of a “definition” because it is not a rigorous definition of a set. For instance, what do we mean by a “collection” of objects? This “definition” will be sufficient for our course, but be warned that defining a set in this vague way can lead to some serious mathematical issues, such as *Russell’s paradox*. A mathematician whose expertise is in set theory may scowl disagreeably if you try to define a set as we have above.

Note

Let S be the set of all sets that aren’t members of themselves. Is S a member of itself? If you think carefully about this, you’ll see that S can be neither a member of itself, nor *not* a member of itself. Uh oh! This contradiction is known as “Russell’s paradox” (named for the British philosopher, mathematician, and all-round academic Bertrand Russell). Mathematicians deal with this by declaring that some object collections, called **classes**, are not in fact sets.

Definition: Elements and Empty Set

The members of a set are called its **elements**. If S is a set, we write $x \in S$ to indicate “ x is an element of S ,” and $x \notin S$ to indicate “ x is not an element of S .” There is a unique set containing no elements; it is called the **empty set**, and denoted by $\emptyset$.

Sets must be **well-defined**: that is, it must be clear exactly which objects are in a set, and which objects aren’t. For instance, the set of all integers is well-defined, but the set of all big integers is not well-defined, since it is unclear what “big” means in this context.

We refer to some sets so frequently in mathematics that we have special notation for them.

Example 1.1.1

Some common sets are:

Illegal pream-token (&)

We can further write denote by $\mathbb{Z}^+$ the set of all positive integers, and by $\mathbb{Z}^*$ the set of all nonzero integers. Can you guess what the penultimate notation represents if we replace $\mathbb{Z}$ with $\mathbb{Q}$ or $\mathbb{R}$, and/or $+$ with $-$? What about what the last notation, if we replace $\mathbb{Z}$ with $\mathbb{Q}$, $\mathbb{R}$, or $\mathbb{C}$?

We also provide notation for commonly considered sets of matrices:

Definition

Given $m, n \in \mathbb{Z}^+$ and a set S , we define $\mathbb{M}_{m \times n}(S)$ to be the set of all $m \times n$ matrices over S (that is, of all $m \times n$ matrices with entries in S). We use the shorthand notation $\mathbb{M}_n(S)$ for the set $\mathbb{M}_{n \times n}(S)$.

One common way of describing a set is to list its elements in curly braces, separated by commas; you can use ellipses to indicate a repeated pattern of elements. A few examples are $\{1, 4, \pi\}$, $\{3, 4, 5, \dots\}$, and $\{\dots, -4, -2, 0, 2, 4, \dots\}$; the last of these can be written more concisely as $\{0, \pm 2, \pm 4, \dots\}$. Note that since elements of a set are unordered, the sets $\{1, 4, \pi\}$ and $\{4, \pi, 1\}$, for instance, are identical.

Another method is using **set-builder notation**. This consists of an element name (or names), followed by a colon (meaning “such that”), followed by a Boolean expression involving the element name(s), all surrounded by curly braces.

Note

The use of a colon to denote “such that” is *only* valid in the above set-builder notation context. Outside of this context, you should never use a colon to denote “such that”; instead, use the abbreviation “s.t.” or write out the actual words. Conversely, never use one of those ways of indicating “such that” within set-builder notation; always use a colon there. Why? Convention.

For example,

$$\{x \in \mathbb{Z} : x > 4\}$$

is the set $\{5, 6, 7, \dots\}$, while

$$\{z \in \mathbb{C} : |z| = 1\}$$

is the set of all complex numbers at distance 1 from the origin in the complex plane.

Note

If one simply writes $\{x : x > 4\}$, it is unclear what this set is; it could be the set of all integers greater than 4, or the set of all real numbers greater than 4, or something else. When one can, it is better to identify the named element(s) as a member (members) of a known set, such as $\mathbb{R}$ or $\mathbb{Z}$, whenever possible.

Definition: Subset and Superset

Set A is a **subset** of B (and set B is a **superset** of A) if every element in A is also in B . We denote “ A is a subset of B ” by $A \subseteq B$. Sets A and B are said to be **equal**, and we write $A = B$, if they contain exactly the same elements; equivalently, $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$. Set A is a **proper** subset of set B if $A \subseteq B$ but $A \neq B$; we write this $A \subsetneq B$.

Remark

Sometimes the notation $A \subset B$ is used to indicate that A is a proper subset of B , and sometimes it is simply used to mean that A is a subset—proper or improper—of B . We will not use the notation $A \subset B$ in this text.

Note

One often proves that two sets A and B are equal by proving that $A \subseteq B$ and $B \subseteq A$.

Example 1.1.2

We have the following: $\mathbb{Z}^+ \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.

Definition: Power Set

The **power set** of A , denoted $P(A)$, is the set of all subsets of A . (Note that the power set of any set contains the empty set as an element.)

Note

Be careful to use your curly braces correctly when writing power sets! Remember, the power set of a set is a *set of sets*.

The following provides a good example of using braces correctly.

Example 1.1.3

If $A = \{a, b\}$, then $P(A) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$. Note that the element $\{a, b\}$ of $P(A)$ could also be written simply as A .

Definition: Union, Intersection, Difference, and Disjoint

1. If A and B are subsets of a set U , then the **union** of A and B , denoted $A \cup B$, is the set $A \cup B = \{x^{\text{int}}U : x^{\text{int}}A \text{ or } x^{\text{int}}B\}$; the **intersection** of A and B , denoted $A \cap B$, is the set $A \cap B = \{x^{\text{int}}U : x^{\text{int}}A \text{ and } x^{\text{int}}B\}$; and the **difference** of A and B , denoted $A - B$ (or $A \setminus B$), is the set $A - B = \{x : x^{\text{int}}A \text{ and } x \notin B\}$.
2. More generally, given any collection of subsets A_i (i in some index set I) of a set U , the **union** of the A_i is

$$\bigcup_{i^{\text{int}}I} A_i = \{x^{\text{int}}U : x^{\text{int}}A_i \text{ for some } i^{\text{int}}I\},$$

and the **intersection** of the A_i is

$$\bigcap_{i^{\text{int}}I} A_i = \{x^{\text{int}}U : x^{\text{int}}A_i \text{ for every } i^{\text{int}}I\}.$$

3. Sets A and B are **disjoint** if $A \cap B = \emptyset$. More generally, sets A_i (i in some index set I) are *disjoint* if

$$\bigcap_{i^{\text{int}}I} A_i = \emptyset$$

and are **mutually disjoint** if

$$A_i \cap A_j = \emptyset \text{ for all } i \neq j^{\text{int}}I.$$

Notice that for any subsets A and B of a set U , $A \cap B \subseteq A \subseteq A \cup B$ and $A \cap B \subseteq B \subseteq A \cup B$. Also note that if sets A_i ($i^{\text{int}}I$) are mutually disjoint then they are also disjoint, but they may be disjoint without being mutually disjoint. For example, the sets $\{i, i + 1\}$ for $i^{\text{int}}\mathbb{Z}$ are disjoint but not mutually disjoint. (Do you see why?)

We define one more way of “combining” sets.

Definition: Direct Product and Ordered Pair

Let A and B be sets. Then the **direct product** (or **Cartesian product**) $A \times B$ of A and B is the set

$$A \times B = \{(a, b) : a \in A, b \in B\}.$$

An element (a, b) of $A \times B$ is called an **ordered pair**. More generally, if $A_1, A_2, \dots, A_n$ are sets for some $n \in \mathbb{Z}^+$, then the **product** of the A_i is

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for } i = 1, 2, \dots, n\};$$

the elements $a_1, a_2, \dots, a_n$ of this product are called n -tuples (or triples, if $n = 3$). (You can also have products of infinitely many sets, but we will not discuss that in this course.) Finally, if each set A_i is the same set A , we can use the notation A^n to denote the product

$$A \times A \times \dots \times A = \{(a_1, a_2, \dots, a_n) : a_i \in A \text{ for } i = 1, 2, \dots, n\}.$$

of n copies of A .

Example 1.1.4

For example, the Cartesian plane is the set $\mathbb{R}^2$, and the set $\mathbb{Z} \times \mathbb{R}$ consists of exactly the points in the plane with integer x -coordinates (that is, the points that lie on vertical lines intersecting the x -axis at integer values).

1.2: Functions

You have probably encountered functions before. In introductory calculus, for instance, you typically deal with functions from $\mathbb{R}$ to $\mathbb{R}$ (e.g., the function $f(x) = x^2$). More generally, functions “send” elements of one set to elements of another set; these sets may or may not be sets of real numbers. We provide below a “good enough for government work” definition of a function.

Definition: Function

A **function** f from a set S to a set T is a “rule” that assigns to each element s in S a unique element $f(s)$ in T ; the element $f(s)$ is called the **image** of s under f . If f is a function from S to T , we write $f : S \rightarrow T$, and call S the **domain** of f and T the **codomain** of f . The **range** of f is

$$f(S) = \{f(s) \mid s \in S\} \subseteq T.$$

More generally, if $U \subseteq S$, the image of U in T under f is

$$f(U) = \{f(u) \mid u \in U\}.$$

If $V \subseteq T$, the **preimage** of V in S under f is the set

$$f^{-1}(V) = \{s \in S \mid f(s) \in V\}.$$

Example 1.2.1

Consider the function $f : \mathbb{Z} \rightarrow \mathbb{R}$ defined by $f(x) = x^2$. The domain of f is $\mathbb{Z}$ and the codomain of f is $\mathbb{R}$; the range of f is $\{x^2 \mid x \in \mathbb{Z}\} = \{0, 1, 4, 9, \dots\}$. The image of $\{-2, -1, 1, 2\}$ under f is the two-element set $\{1, 4\} \subseteq \mathbb{R}$, and the preimage of $\{4, 25, \pi\}$ under f is the set $\{\pm 2, \pm 5\}$. (Do you see why $\pm\sqrt{\pi}$ are not in this preimage?) What is the preimage of just $\{\pi\}$ under f ?

The following definitions will be very important in our future work.

Definition: One-to-one, Onto, and Bijection

Let S and T be sets, and $f : S \rightarrow T$.

1. Function f is **one-to-one** (1-1) if whenever $s_1, s_2 \in S$ with $f(s_1) = f(s_2)$, we have $s_1 = s_2$. Equivalently, f is one-to-one if whenever $s_1 \neq s_2 \in S$, then $f(s_1) \neq f(s_2)$.
2. Function f is **onto** if for every $t \in T$, there exists an element $s \in S$ such that $f(s) = t$. Equivalently, f is onto if $f(S) = T$.
3. Function f is a **bijection** if it is both one-to-one and onto.

Remark

We will often have to show functions are one-to-one or onto. Given a function $f : S \rightarrow T$, the following methods are recommended.

- To prove that f is one-to-one: Let $s_1, s_2 \in S$ with $f(s_1) = f(s_2)$ and prove that then $s_1 = s_2$.

Note: It is not sufficient to prove that if $s_1 = s_2$ in S then $f(s_1) = f(s_2)$; that holds true for ANY function from S to T ! Be careful to assume and prove the correct facts.

- To prove that f is not one-to-one: Identify two elements $s_1 \neq s_2$ of S such that $f(s_1) = f(s_2)$.
- To prove that f is onto: Let $t \in T$ and prove that there exists an element $s \in S$ with $f(s) = t$.

Note: It is not sufficient to prove that if $s \in S$ then $f(s)$ is in T ; that holds true for ANY function from S to T ! Again, be careful to assume and prove the correct facts.

- To prove that f is *not* onto: Identify an element $t \in T$ for which there is no $s \in S$ with $f(s) = t$.

Example 1.2.2

Consider the function $f: \mathbb{R}^* \rightarrow \mathbb{R}^+$ defined by $f(x) = x^2$. Function f is *not* one-to-one: indeed, -1 and 1 are in $\mathbb{R}^*$ with $f(-1) = 1 = f(1)$ in $\mathbb{R}^+$. However, f is onto: indeed, let $t \in \mathbb{R}^+$. Then $\sqrt{t} \in \mathbb{R}^*$ with $t = f(\sqrt{t})$, so we're done.

Example 1.2.3

Consider the function $f: \mathbb{Z}^+ \rightarrow \mathbb{R}$ defined by $f(x) = \frac{x}{2}$. Function f is one-to-one: indeed, let $s_1, s_2 \in \mathbb{Z}^+$ with $f(s_1) = f(s_2)$. Then $\frac{s_1}{2} = f(s_1) = f(s_2) = \frac{s_2}{2}$; multiplying both sides of the equation $\frac{s_1}{2} = \frac{s_2}{2}$ by 2, we obtain $s_1 = s_2$. However, f is *not* onto: for example, $\pi \in \mathbb{R}$ but there is no positive integer s for which $f(s) = \frac{s}{2} = \pi$.

Recall that we can combine certain functions using **composition**:

Definition: Composition and Identity Function

If $f: S \rightarrow T$ and $g: T \rightarrow U$, then the **composition of** f and g is the function $g \circ f: S \rightarrow U$ defined by

$$(g \circ f)(s) = g(f(s))$$

for all $s \in S$. (More generally, you can **compose** functions $f: S \rightarrow T$ and $g: T \rightarrow U$ to form $g \circ f: S \rightarrow U$, as long as $f(S) \subseteq T$.) Also recall that given any set S , the **identity function** on S is the function $1_S: S \rightarrow S$ defined by $1_S(s) = s$ for every $s \in S$.

Definition: Inverse

Let f be a function from S to T . A function g from T to S is an **inverse** of f if $g \circ f$ and $f \circ g$ are the identity functions on S and T , respectively; that is, if for all $s \in S$ and $t \in T$, $g(f(s)) = s$ and $f(g(t)) = t$.

We say a function is **invertible** if it has an inverse.

We have the following useful theorems.

Theorem 1.2.1

Function $f: S \rightarrow T$ is invertible if and only if f is a bijection.

Proof available at the end of this section.

Theorem 1.2.2

Let $f: S \rightarrow T$ be invertible. Then the inverse of f is unique; we denote the unique inverse of f by f^{-1} .

Proof available at the end of this section.

Theorem 1.2.3

Let $f : S \rightarrow T$ and $g : T \rightarrow U$ be functions. If f and g are both 1-1 [resp., onto], then $g \circ f : S \rightarrow U$ is 1-1 [resp., onto]. It follows that if f and g are both bijections, then $g \circ f : S \rightarrow U$ is a bijection.

Proof available at the end of this section.

Answers

Theorem 1.2.1

Suppose f has inverse g . Let $t \in T$. Then $f(g(t)) = t$, so f is onto. Next, suppose $s_1, s_2 \in S$ with $f(s_1) = f(s_2)$. Then

$$s_1 = g(f(s_1)) = g(f(s_2)) = s_2,$$

so f is one-to-one.

Conversely, suppose f is a bijection. Then for every $t \in T$, there exists a unique element $s_t \in S$ such that $f(s_t) = t$. It is then straightforward to show that the function $g : T \rightarrow S$ defined by

$$g(t) = s_t \text{ for every } t \in T$$

is an inverse of f .

Theorem 1.2.2

Suppose f has inverses g and h . Then for every $t \in T$, $f(g(t)) = t = f(h(t))$. But since f is invertible, it's one-to-one, so we must have $g(t) = h(t)$ for every $t \in T$. Thus, $g = h$, and so f cannot have two or more distinct inverses.

Theorem 1.2.3

Assume f and g are 1-1. Let $s_1, s_2 \in S$ with $(g \circ f)(s_1) = (g \circ f)(s_2)$. Then $g(f(s_1)) = g(f(s_2))$; since g is one-to-one, this shows that $f(s_1) = f(s_2)$. Then since f is one-to-one, we must have $s_1 = s_2$. Thus, $g \circ f$ is one-to-one.

The proof that $g \circ f$ is onto if f and g are onto is left as an exercise for the reader.

1.3: Cardinality

One of the set traits that will be useful to us in distinguishing between algebraic structures is **cardinality**.

Definition: Cardinality, Finite, and Infinite

A set is **finite** if it contains a finite number of elements; otherwise, it's **infinite**. The **cardinality** of a finite set S is the number of elements in S ; we denote the cardinality of S by $|S|$. When S is infinite, we may write $|S| = \infty$.

Note

Of course, vertical bars are used to denote other mathematical concepts; for instance, if x is a real number, $|x|$ usually denotes the absolute value of x . You must determine from context, and from the nature of the expression within the bars, what vertical bars mean in a particular context.

Note that in the above definition, we omitted the definition of the cardinality of an infinite set. This is because defining the cardinality of an infinite set is a more complicated endeavor, and one which is, in the most general context, beyond the scope of this class. For us, it will suffice to distinguish between two types of infinite sets: **countably infinite** sets and **uncountable** (or **uncountably infinite**) sets.

Definition: Countable and Uncountable

A set S is said to be **countably infinite** if there exists a bijection from S to $\mathbb{Z}$ (equivalently, if there exists a bijection from $\mathbb{Z}$ to S). It is said to be **countable** if it is finite or countably infinite, and **uncountable** (or **uncountably infinite**) if it is not countable.

Here are some straightforward examples to get us started:

Example 1.3.1

1. $|\{a, b\}| = 2$.
2. $|\emptyset| = 0$.
3. Clearly, $\mathbb{Z}$ itself is countably infinite.

Remark

It turns out that countably infinite sets have the same cardinality as one another, while a countably infinite set and an uncountable set have different cardinalities. Intuitively, you can think of two countably infinite sets as having the same “size,” and a countable set and an uncountable set as having different “sizes”; however, this is a risky way of framing things, since it can make some results seem counterintuitive when you're used to dealing only with finite sets (see, for instance, Example 1.3.2). Luckily, exploring the cardinality of infinite sets isn't the focus of this class.

Perhaps surprisingly, a proper subset of a set can have the same cardinality as its superset, as the following example shows.

Example 1.3.2

We claim that $\mathbb{Z}^+$ is countably infinite. Indeed, consider the function $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ defined by $f(n) = (-1)^n \lfloor \frac{n}{2} \rfloor$, where $\lfloor x \rfloor$ denotes the greatest integer less than or equal to x , for each $x \in \mathbb{R}$. The fact that f is a bijection is demonstrated (though not proven) by the following visual representation, where each number maps via f to the value directly to the right of it:

Illegal pream-token (&)

Note that this means that $\mathbb{Z}$ and its proper subset $\mathbb{Z}^+$ have the same cardinality, that is, the same “size”!

We summarize here examples of countably and uncountably infinite sets. (On pp. 5–6 of [1], Fraleigh sketches proofs of the facts that $\mathbb{Q}$ is countable and that the interval $(0, 1)$ in $\mathbb{R}$ is uncountable. The proof then that $\mathbb{R}$ is uncountable follows from Theorem 1.3.1, below.)

Countably infinite sets: $\mathbb{Z}, \mathbb{Z}^+, \mathbb{Z}^-, \mathbb{Z}^*, \mathbb{Q}, \mathbb{Q}^+, \mathbb{Q}^-, \mathbb{Q}^*, \mathbb{N}$

Uncountably infinite sets: $\mathbb{R}, \mathbb{R}^+, \mathbb{R}^-, \mathbb{R}^*, \mathbb{C}, \mathbb{C}^*$, the interval $(0, 1)$ in $\mathbb{R}$

The key idea here for us is that if two sets are essentially “the same,” then they must have the same “size.” Thus, we see that there is some fundamental difference between the sets $\mathbb{Z}$ and $\mathbb{R}$ (in fact, there are many such differences). On the other hand, cardinality alone won’t allow us to distinguish structurally between the sets $\mathbb{Z}$ and $\mathbb{Z}^+$.

We end our preliminary chapter with the following theorem and a corollary of it (which can be proved using induction on n). We omit the proofs of the first two statements.

Theorem 1.3.1

Let A and B be sets.

1. If $A \subseteq B$ and A is infinite [uncountable] then so is B .
2. If $A \subseteq B$ and B is finite [countable] then so is A .
3. If $|A| = m < \infty$ and $|B| = n < \infty$, then $|A \times B| = mn$. More generally, if $n > 1$ is an integer and $A_1, A_2, \dots, A_n$ are sets with $|A_i| = k_i < \infty$ for each $i = 1, 2, \dots, n$, then $|A_1 \times A_2 \times \dots \times A_n| = k_1 k_2 \dots k_n$.
4. Let A and B be countable sets. Then $A \times B$ is countable. More generally, if $n > 1$ be an integer and $A_1, A_2, \dots, A_n$ are countable sets, then $A_1 \times A_2 \times \dots \times A_n$ is countable.

Proof available at the end of this section.

Remark

It is *not* true that any countable product of countable (or even finite) sets is countable. Indeed, even the set $\{0, 1\} \times \{0, 1\} \times \dots$ is uncountable. (If you want to get into the gory details of this, the key is that there is a bijection from this set to the power set of the natural numbers, which Cantor’s Theorem tells us is uncountable. You are welcome to jump down this rabbit hole by googling “Cantor’s Theorem,” if you desire, but know that you will not be responsible for that material in class.)

Answers

Theorem 1.3.1

You prove the first statement in Part 3 in your homework. The second statement in Part 3 follows using induction on the number of sets.

To prove the first statement in Part 4: Assume that A and B are both countably infinite. Since $\mathbb{Z}^+$ is countably infinite, we can index the elements of A and of B by $\mathbb{Z}^+$, writing

$$A = \{a_1, a_2, \dots\} \text{ and } B = \{b_1, b_2, \dots\}.$$

Notice that the table

Illegal pream-token (&)

contains every element of $A \times B$. We can then list the elements of $A \times B$ by listing the elements in progressive upper-right to lower-left diagonals, starting with (a_1, b_1) and moving to the right along the top row: that is, we can write

$$A \times B = \{(a_1, b_1), (a_1, b_2), (a_2, b_1), (a_2, b_3), (a_3, b_2), (a_3, b_1), \dots\}.$$

This implicitly yields a bijection from $\mathbb{Z}^+$ to $A \times B$; thus, $A \times B$ is countably infinite, and hence countable.

The proof in the case that one or both A and B are finite is similar; the corresponding table in that case will simply have either only finitely many rows or finitely many columns, or both.

The second statement in Part 4 follows using induction on the number of sets.

1.4: Exercises

1. Yes/No. For each of the following, write Y if the object described is a well-defined set; otherwise, write N. You do NOT need to provide explanations or show work for this problem.

- a. $\{z^{\text{int}}\mathbb{C} : |z| = 1\}$
- b. $\{\epsilon^{\text{int}}\mathbb{R}^+ : \epsilon \text{ is sufficiently small}\}$
- c. $\{q^{\text{int}}\mathbb{Q} : q \text{ can be written with denominator 4}\}$
- d. $\{n^{\text{int}}\mathbb{Z} : n^2 < 0\}$

2. List the elements in the following sets, writing your answers as sets.

Example: $\{z^{\text{int}}\mathbb{C} : z^4 = 1\}$ *Solution:* $\{\pm 1, \pm i\}$

- a. $\{z^{\text{int}}\mathbb{R} : z^2 = 5\}$
- b. $\{m^{\text{int}}\mathbb{Z} : mn = 50 \text{ for some } n^{\text{int}}\mathbb{Z}\}$
- c. $\{a, b, c\} \times \{1, d\}$
- d. $P(\{a, b, c\})$

3. Let S be a set with cardinality $n^{\text{int}}\mathbb{N}$. Use the cardinalities of $P(\{a, b\})$ and $P(\{a, b, c\})$ to make a conjecture about the cardinality of $P(S)$. You do not need to prove that your conjecture is correct (but you should try to ensure it is correct).

4. Let $f : \mathbb{Z}^2 \rightarrow \mathbb{R}$ be defined by $f(a, b) = ab$. (Note: technically, we should write $f((a, b))$, not $f(a, b)$, since f is being applied to an ordered pair, but this is one of those cases in which mathematicians abuse notation in the interest of concision.)

- a. What are f 's domain, codomain, and range?
- b. Prove or disprove each of the following statements. (Your proofs do not need to be long to be correct!)
 - i. f is onto;
 - ii. f is 1-1;
 - iii. f is a bijection. (You may refer to parts (i) and (ii) for this part.)
- c. Find the images of the element $(6, -2)$ and of the set $\mathbb{Z}^- \times \mathbb{Z}^-$ under f . (Remember that the image of an element is an element, and the image of a set is a set.)
- d. Find the preimage of $\{2, 3\}$ under f . (Remember that the preimage of a set is a set.)

5. Let S, T , and U be sets, and let $f : S \rightarrow T$ and $g : T \rightarrow U$ be onto. Prove that $g \circ f$ is onto.

6. Let A and B be sets with $|A| = m < \infty$ and $|B| = n < \infty$. Prove that $|A \times B| = mn$.

2: GROUPS

2.1: Binary Operations and Structures

2.2: Exercises, Part I

2.3: The Definition of a Group

2.4: Examples of Groups/Nongroups, Part I

2.5: Group Conventions and Properties

2.6: Examples of Groups/Nongroups, Part II

2.7: Summaries of Groups We've Seen

2.8: Exercises, Part II

This page titled [2: Groups](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

2.1: Binary Operations and Structures

So far we have been discussing sets. These are extremely simple objects, essentially mathematical “bags of stuff.” Without any added structure, their usefulness is very limited. Imagine, for instance, living with friends in a two-story house without rooms, stairs, closets, or hallways. You have no privacy, cannot access the second floor, etc. A set with no added structure will not help us, say, solve a linear equation. What *will* help us with such things are objects such as groups, rings, fields, and vector spaces. These are sets equipped with *binary operations* which allow us to combine set elements in various ways. We first rigorously define a binary operation.

Definition: Binary Operation

A **binary operation** on a set S is a function from $S \times S$ to S . Given a binary operation $*$ on S , for each $(a, b) \in S \times S$ we denote $*((a, b))$ in S more simply by $a * b$. (Intuitively, a binary operation $*$ on S assigns to each pair of elements $a, b \in S$ a unique element $a * b$ of S .)

A set S equipped with a binary operation $*$ is called a **binary (algebraic) structure**, and is denoted by $\langle S, * \rangle$, or just by S , if $*$ is understood from context.

Remark

1. For $*$ to be a binary operation on S , $a * b$ must be *well-defined* and *in* S for each $a, b \in S$. For instance, we cannot define a binary operation on $\mathbb{R}$ by

$$a * b = \text{the greatest number less than } a + b$$

since there is no such “greatest number.” Nor can we define a binary operation on $\mathbb{Z}$ by $a * b = \frac{ab}{2}$, since for, say, $a = b = 1 \in \mathbb{Z}$, $\frac{ab}{2} = \frac{1}{2} \notin \mathbb{Z}$.

2. Not every binary operation is denoted by $*$. In fact, many already have common notations: for instance, $+$ on $\mathbb{Z}$ or $\circ$ on the set of functions from $\mathbb{R}$ to $\mathbb{R}$. We will assume these common notations represent the “usual” binary operations we know them to mean, unless otherwise noted.
3. Do not mix up the $*$ that indicates a binary operation and the superscript $*$ that indicates that we are only considering the nonzero elements of a given set (e.g., $\mathbb{R}^*$). You should be able to tell which type of $*$ we are using from context and placement. Also, make sure you correctly place these symbols!

Definition: Associative

A binary operation $*$ on a set S is **associative** if $(a * b) * c = a * (b * c)$ for all $a, b, c \in S$.

Remark

When a binary operation is associative, we can omit parentheses when operating on set elements. For example, $+$ is associative on $\mathbb{Z}$, so we can unambiguously write the (equal) expressions $1 + (2 + 3)$ and $(1 + 2) + 3$ as $1 + 2 + 3$.

Definition: Commutative

A binary operation $*$ on set S is **commutative** if

$$a * b = b * a$$

for all $a, b \in S$. We say that specific elements a and b of S **commute** if $a * b = b * a$.

Definition: Identity Element

Let $\langle S, * \rangle$ be a binary structure. An element e in S is an **identity element** of $\langle S, * \rangle$ if $x * e = e * x = x$ for all $x \in S$.

Note

Sometimes an identity element of $\langle S, * \rangle$ is referred to as an *identity element of S under $*$* , or, when $*$ is clear from context, simply as an **identity element** of S .

Note

Not every binary structure contains an identity element! (Ex: $\langle \mathbb{Z}, - \rangle$).

A natural question to ask is if a binary structure can have more than one identity element? The answer is no!

Theorem 2.1.1

A binary structure $\langle S, * \rangle$ has *at most one* identity element. That is, identity elements in binary structures, when they exist, are unique.

Proof available at the end of this section.

Definition: Two-sided Inverse

Let $\langle S, * \rangle$ be a binary structure with identity element e . Then for $a \in S$, b is a *(two-sided) inverse of a in $\langle S, * \rangle$* if $a * b = b * a = e$.

Note

We can also refer to such an element b as an *inverse for a in S under $*$* , or, when $*$ is clear from context, simply as an *inverse of a* .

Note

- Not every element in a binary structure with an identity element has an inverse!
- If a binary structure *does not have* an identity element, it *doesn't even make sense* to say an element in the structure does or does not have an inverse!

Theorem 2.1.2

Let $\langle S, * \rangle$ be a binary structure with an identity element, where $*$ is associative. Let $a \in S$. If a has an inverse, then its inverse is unique.

Proof available at the end of this section.

We end by discussing the idea of **closure** under a binary operation.

Definition: Closed

Let $\langle S, * \rangle$ be a binary structure and let $T \subseteq S$. Then T is said to be **closed** under $*$ if $t_1 * t_2 \in T$ whenever $t_1, t_2 \in T$.

Example 2.1.1

Consider the binary structure $\langle M_2(\mathbb{R}), + \rangle$, where $+$ denotes matrix addition. Let

$$T = \{A \in M_2(\mathbb{R}) : A \text{ is invertible}\}. \quad (2.1.1)$$

We claim that T is *not* closed under $+$. Indeed, if we denote the identity matrix in $M_2(\mathbb{R})$ by I_2 , then we observe that $I_2, -I_2 \in T$, but $I_2 + (-I_2) = 0 \notin T$, since the zero matrix isn't invertible.

Example 2.1.2

Consider the binary structure $\langle M_2(\mathbb{R}), \cdot \rangle$, where $\cdot$ denotes matrix multiplication. Again, let T be the set of all invertible matrices in $M_2(\mathbb{R})$. We claim that T is closed under $\cdot$. Indeed, let $A, B \in T$. Then A and B are invertible, so their determinants are nonzero. Thus, $\det(AB) = (\det A)(\det B) \neq 0$, so AB is invertible, and hence $AB \in T$.

Answers

Theorem 2.1.1

Suppose that e and f are identity elements of S . Then since e is an identity element, $e * f = f$ and since f is an identity element, $e * f = e$. Thus, $e = f$.

Theorem 2.1.2

Let e be the identity element of S . Suppose a has inverses b and c . Then $a * b = e$ so, multiplying both sides of the equation by c on the left, we have $c * (a * b) = c * e = c$. But since $*$ is associative, we have $c * (a * b) = (c * a) * b = e * b = b$. But $b = c$. Thus, a 's inverse is unique.

2.2: Exercises, Part I

- For each of the following, write Y if the given “operation” is a well-defined binary operation on the given set; otherwise, write N. In each case in which it *isn't* a well-defined binary operation on the set, provide a *brief* explanation. You do not need to prove or explain anything in the cases in which it *is* a binary operation.
 - $+$ on $\mathbb{C}^*$
 - $*$ on $\mathbb{R}^+$ defined by $x * y = \log_x y$
 - $*$ on $M_2(\mathbb{R})$ defined by $A * B = AB^{-1}$
 - $*$ on $\mathbb{Q}^*$ defined by $z * w = \frac{z}{w}$
- Define $*$ on $\mathbb{Q}$ by $p * q = pq + 1$. Prove or disprove that $*$ is (a) commutative; (b) associative.
- Prove that matrix multiplication is not commutative on $M_2(\mathbb{R})$.
- Prove or disprove each of the following statements.
 - The set $2\mathbb{Z} = \{2x : x \in \mathbb{Z}\}$ is closed under addition in $\mathbb{Z}$.
 - The set $S = \{1, 2, 3\}$ is closed under multiplication in $\mathbb{R}$.
 - The set

$$U = \left\{ \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}$$

is closed under multiplication in $M_2(\mathbb{R})$. (Recall that U is the set of *upper-triangular matrices* in $M_2(\mathbb{R})$.)

- Let $*$ be an associative and commutative binary operation on a set S . An element $u \in S$ is said to be an **idempotent** in S if $u * u = u$. Let H be the set of all idempotents in S . Prove that H is closed under $*$.




Figure 2.2.1: (Copyright; Bill Griffith. Reprinted with permission)

This page titled [2.2: Exercises, Part I](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

2.3: The Definition of a Group

At this point you may be asking yourself, why do we care? We've covered a lot of definitions and proved some theorems, but what is the goal of all this? Well, there are actually many goals that we can achieve using such material. Consider the following as an example. Suppose we want to solve the equation $5 + x = 2$. We can probably solve this quite easily almost just by looking at it ($x = -3$), but what facts are we actually using there? If we break down the reasoning leading to this answer, we may obtain something like the following, where each line is equivalent to the one preceding it.

Illegal pream-token (&)

In the second line we add the inverse of 5 in $\langle \mathbb{Z}, + \rangle$ to each side of the equation; in the third line we use associativity of $+$ in $\mathbb{Z}$ (along with computation); and in the fourth line we use the fact that -5 is the additive inverse of 5 (that is, the inverse of 5 in $\mathbb{Z}$ under $+$). Finally, we use the fact that 0 is an additive identity element in $\mathbb{Z}$ (that is, the identity element in $\mathbb{Z}$ under $+$).

In summary, we used **associativity**, **identity elements**, and **inverses** in $\mathbb{Z}$ to solve the given equation. This perhaps suggests that these would be useful traits for a binary structure and/or its operation to have. They are in fact so useful that a binary structure displaying these characteristics is given a special name. We note that these axioms are rather strong; “most” binary structures aren't groups.

Definition: Group Axioms

A **group** is a set G , equipped with a binary operation $*$, that satisfies the following three **group axioms**:

- I. $*$ is associative on G . (Axiom $\mathcal{G}_1$)
- II. There exists an identity element for $*$ in G . (Axiom $\mathcal{G}_2$)
- III. Every element $a \in G$ has an inverse in G . (Axiom $\mathcal{G}_3$)

We denote group G under $*$ by the binary structure notation $\langle G, * \rangle$, or simply by G if the operation $*$ is known from context (or need not be known in the current situation).

Remark

When proving/disproving that a set G is/is not a group under an (apparent) operation $*$:

- The first thing you should do is check to make sure that $\langle G, * \rangle$ is a binary structure: that is, you should make sure that $a * b$ is well-defined and in G for every $a, b \in G$. If this is not the case, it doesn't make sense to check to see if axioms $\mathcal{G}_1$ – $\mathcal{G}_3$ hold). (For instance: $\mathbb{Z}^*$ has no chance of being a group under $\div$, since, e.g., $3, 4 \in \mathbb{Z}^*$ but $3 \div 4 \notin \mathbb{Z}^*$.)
- You should never check $\mathcal{G}_3$ before confirming $\mathcal{G}_2$ holds, because it makes no sense to look for inverses if you haven't confirmed that G contains an identity element under $*$.

2.4: Examples of Groups/Nongroups, Part I

Let's look at some examples of groups/nongroups.

Example 2.4.1

We claim that $\mathbb{Z}$ is a group under addition. Indeed, $\langle \mathbb{Z}, + \rangle$ is a binary structure and that addition is associative on the integers. The integer 0 acts as an identity element of $\mathbb{Z}$ under addition (since $a + 0 = 0 + a = a$ for each $a \in \mathbb{Z}$), and each element a in G has inverse $-a$ since $a + (-a) = -a + a = 0$.

Example 2.4.2

For each following binary structure $\langle G, * \rangle$, determine whether or not G is a group. For those that are *not* groups, determine the first group axiom that fails, and provide a proof that it fails.

1. $\langle \mathbb{Q}, + \rangle$
2. $\langle \mathbb{Z}, - \rangle$
3. $\langle \mathbb{R}, \cdot \rangle$
4. $\langle \mathbb{C}^*, \cdot \rangle$
5. $\langle \mathbb{R}, + \rangle$
6. $\langle \mathbb{Z}^+, + \rangle$
7. $\langle \mathbb{Z}^*, \cdot \rangle$
8. $\langle \mathbb{M}_n(\mathbb{R}), + \rangle$
9. $\langle \mathbb{C}, + \rangle$
10. $\langle \mathbb{Z}, \cdot \rangle$
11. $\langle \mathbb{R}^*, \cdot \rangle$
12. $\langle \mathbb{M}_n(\mathbb{R}), \cdot \rangle$

If you have taken linear algebra, you have also probably seen a collection of matrices that is a group under matrix multiplication.

Definition: General and Special Linear Groups

Recall that given a square matrix A , the notation $\det A$ denotes the determinant of A . Let

$$GL(n, \mathbb{R}) = \{M \in \mathbb{M}_n(\mathbb{R}) : \det M \neq 0\}$$

(that is, let $GL(n, \mathbb{R})$ be the set of all invertible $n \times n$ matrices over $\mathbb{R}$), and let

$$SL(n, \mathbb{R}) = \{M \in \mathbb{M}_n(\mathbb{R}) : \det M = 1\}.$$

These subsets of $\mathbb{M}_n(\mathbb{R})$ are called, respectively, the **general and special linear groups** of degree n over $\mathbb{R}$.

Note that these definitions imply that these subsets of $\mathbb{M}_n(\mathbb{R})$ are groups (under some operation). Sure enough, they are!

Theorem 2.4.1

$GL(n, \mathbb{R})$ is a group under matrix multiplication.

Proof available at the end of this section.

Theorem 2.4.2

$SL(n, \mathbb{R})$, is a group under matrix multiplication.

Proof available at the end of this section.

Remark

Throughout this course, if we are discussing a set $GL(n, \mathbb{R})$ or $SL(n, \mathbb{R})$, you should assume $n \in \mathbb{Z}^+$, unless otherwise noted.

We end this section with a final example.

Example 2.4.3

Define $*$ on $\mathbb{Q}^*$ by $a * b = (ab)/2$ for all $a, b \in \mathbb{Q}^*$. Prove that $\langle \mathbb{Q}^*, * \rangle$ is a group.

Solution

First, $\langle \mathbb{Q}^*, * \rangle$ is a binary structure, since $(ab)/2$ is rational and nonzero whenever a, b are rational and nonzero.

Next, we check that $\mathbb{Q}^*$ under $*$ satisfies the group axioms. Since multiplication is commutative on $\mathbb{Q}$, $*$ is clearly commutative on $\mathbb{Q}^*$, and so our work to show $\mathcal{G}_2$ and $\mathcal{G}_3$ is marginally reduced.

First, associativity of $*$ on $\mathbb{Q}^*$ is inherited from associativity of multiplication on $\mathbb{Q}^*$.

Notice that the perhaps “obvious” choice, 1, is *not* an identity element for $\mathbb{Q}^*$ under $*$: for instance, $1 * 3 = 3/2 \neq 3$. Rather, e is such an identity element if and only if for all $a \in \mathbb{Q}$ we have $a = e * a = (ea)/2$. We clearly have $a = (2a)/2$ for all $a \in \mathbb{Q}^*$; so 2 acts as an identity element for $\mathbb{Q}^*$ under $*$.

Finally, let $a \in \mathbb{Q}^*$. Since $a \neq 0$, it makes sense to divide by a ; then $4/a \in \mathbb{Q}^*$, with $a * (4/a) = (a(4/a))/2 = 2$.

Thus, $\langle \mathbb{Q}^*, * \rangle$ is a group.

Answers

Theorem 2.4.1

Let $A, B \in GL(n, \mathbb{R})$. Then $\det(AB) = (\det A)(\det B) \neq 0$ (since $\det A, \det B \neq 0$), so $AB \in GL(n, \mathbb{R})$. Thus, $\langle GL(n, \mathbb{R}), \cdot \rangle$ is a binary structure.

We know that matrix multiplication is always associative, so $\mathcal{G}_2$ holds. Next,

$$\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}$$

the is in $GL(n, \mathbb{R})$ since $\det I_n = 1 \neq 0$, and it acts as an identity element for $\langle GL(n, \mathbb{R}), \cdot \rangle$ since

$$AI_n = I_n A = A$$

for all $A \in GL(n, \mathbb{R})$.

Finally, let $A \in GL(n, \mathbb{R})$. Since $\det A \neq 0$, A has (matrix multiplicative) inverse A^{-1} in $M_2(\mathbb{R})$. But we need to verify that A^{-1} is in G . This is in fact the case, however, since A^{-1} is invertible (it has inverse A), hence $\det A^{-1} \neq 0$. Thus, A^{-1} is also in $GL(n, \mathbb{R})$.

So $GL(n, \mathbb{R})$ is a group under multiplication.

Theorem 2.4.2

Let $A, B \in SL(n, \mathbb{R})$. Then $\det(AB) = (\det A)(\det B) = 1(1) = 1$, so $AB \in SL(n, \mathbb{R})$. Thus, $\langle SL(n, \mathbb{R}), \cdot \rangle$ is a binary structure.

The rest of the proof is left as an exercise for the reader.

This page titled [2.4: Examples of Groups/Nongroups, Part I](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

2.5: Group Conventions and Properties

Before we discuss more examples, we present a theorem and look at some conventions we follow and notation we use when discussing groups in general; we also discuss some properties of groups.

2.5.1: Some Group Conventions

Theorem 2.5.1

The identity element of a group is unique (by Theorem 2.1.9), and given any element a of a group G , the inverse of a in G is unique (by Theorem 2.1.2).

Note

We will generally use e or e_G as our default notation for an identity element of group, but be aware that many mathematicians denote a group's identity element by 1. We denote the inverse of element a in G by a^{-1} .

Note

Although it is written in what we call *multiplicative notation*, do *not* assume a^{-1} is what we usually think of as a multiplicative inverse for a ; remember, we don't even know if elements of a group are numbers! The type of inverse that a^{-1} is (a multiplicative inverse for a real number? an additive inverse for a real number? a multiplicative inverse for a matrix? an inverse function for a function from $\mathbb{R}$ to $\mathbb{R}$?) depends on both G 's elements and its operation.

Note

- We usually *don't* use the notation $*$ when describing group operations. Instead, we use the multiplication symbol $\cdot$ for the operation in an arbitrary group, and call applying the operation “multiplying”—even though the operation may not be “multiplication” in the non-abstract, traditional sense! It may actually be addition of real numbers, composition of functions, etc. Moreover, when actually operating in a group $\langle G, \cdot \rangle$, we typically omit the $\cdot$. That is, for $a, b \in G$, we write the product $a \cdot b$ as ab . We call this the “product” of a and b .

For every element a in a group $\langle G, \cdot \rangle$ and $n \in \mathbb{Z}^+$, we use the expression a^n to denote the product

$$a \cdot a \cdot \cdots \cdot a$$

of n copies of a , and a^{-n} to denote $(a^{-1})^n$ (that is, the product of n copies of a^{-1}). Finally, we define a^0 to be e . Note that our “usual” rules for exponents then hold in an arbitrary group: that is, if a is in group $\langle G, \cdot \rangle$ and $m, n \in \mathbb{Z}$, then $a^m a^n = a^{m+n}$ and $(a^m)^n = a^{mn} = (a^n)^m$.

However:

- When we know our operation is commutative, we typically use additive notation, denoting the group operation by $+$, calling the group operation “addition,” and denoting the inverse of an element a by $-a$. When we use additive notation, we do *not* omit the $+$ when operating in a group $\langle G, + \rangle$, and we call $a + b$ a *sum* rather than a product. Also, when working with an operation that is known to be commutative, the identity element may be denoted by 0 rather than by e , e_G , or 1 , and for $n \in \mathbb{Z}$, we write na instead of a^n .

Finally, note that $(-n)a = n(-a) = -(na)$ (where $-a$ and $-(na)$ indicate the additive inverses of a and na , respectively); we can therefore unambiguously use the notation $-na$ for this element. Using this notation, note that for $m^{\text{int}}\mathbb{Z}$, $na + ma = (n + m)a$ and $n(ma) = (nm)a$.

Note

Be careful to always know where an element you are working with lives! For instance, if, as above, $n^{\text{int}}\mathbb{Z}^+$ and a is a group element, $-n$ and $-a$ look similar but may mean very different things. While $-n$ is a negative integer, $-a$ may be the additive inverse of a matrix in $M_2(\mathbb{R})$, the additive inverse 2 of the number 4 in $\mathbb{Z}_6$, or even something completely unrelated to numbers.

Remark

Multiplicative notation can be used when working with any arbitrary group, while additive notation should be used only when working with a group whose binary operation is commutative.

We summarize multiplicative versus addition notation in the following table, where a, b are elements of a group G .

Table 2.5.1: Summary of Multiplicative and Addition Notation in Groups.

	Multiplicative Notation	Additive Notation
Operation Notation	$\cdot$	$+$
a operated with b	ab	$a + b$
Identity Element	e or e_G (or 1)	e or e_G (or 0)
Inverse of a	a^{-1}	$-a$

Note

We *do* use the notation $*$ when using multiplicative or additive notation would lead to confusion. For instance, if we want to define an operation on $\mathbb{Q}^*$ that assigns to pair (a, b) the quantity $ab/2$, it would be unwise to use multiplicative or additive notation for this operation since we already have conventional meanings of ab and $a + b$. Similarly, we would not denote the identity element of $\mathbb{Q}^*$ under this operation by 0 or 1, since the identity element in this group is the rational number 2, and writing $0 = 2$ or $1 = 2$ would look weird.

Finally:

Note

If there is a default notation for a particular operation (say, $\circ$ for composition of functions) or identity element (say, I_n in $GL(n, \mathbb{R})$) we usually use that notation instead.

2.5.2: Some Group Properties

While we don't need to worry about "order" when multiplying a group element a by itself, we *do* need to worry about it in general.

Note

Group operations need *not* be commutative!

Definition: Abelian and Nonabelian

A group $\langle G, \cdot \rangle$ is said to be **abelian** if $ab = ba$ for all $a, b \in G$. Otherwise, G is **nonabelian**. (The word “abelian” derives from the surname of mathematician Niels Henrik Abel.)

Remark

If we know that a binary operation $\cdot$ on a set G is commutative, then in checking to see if axioms $\mathcal{G}_2$ and $\mathcal{G}_3$ hold we need only verify that there exists $e \in G$ such that $ae = a$ (we don't need to check that $ea = a$) for all $a \in G$ and that for each $a \in G$ there exists $b \in G$ such that $ab = e$ (we don't need to check that $ba = e$).

Remark

If G is not known to be abelian, we must be careful when multiplying elements of G by one another: multiplying on the left is, in general, not the same as multiplying on the right!

Definition: Order, Finite Group, and Infinite Group

If G is a group, then the cardinality $|G|$ of G is called the **order** of G . If $|G|$ is finite, then G is said to be a **finite group**; otherwise, it's an **infinite group**.

Example 2.5.1

Of the groups we've discussed, which are abelian? Which are infinite/finite?

We have already seen that identity elements of groups are unique, and that each element a of a group G has a unique inverse. Double exponent: use braces to clarify Here are some other basic properties of groups.

Theorem 2.5.2

If $\langle G, \cdot \rangle$ is a group, then **left and right cancellation laws** hold in G . That is, if $a, b, c \in G$, then

1. If $ab = ac$, we have $b = c$ (the left cancellation law); and
2. If $ba = ca$, we have $b = c$ (the right cancellation law).

Proof available at the end of this section.

Theorem 2.5.3

Let $\langle G, \cdot \rangle$ be a group and let $a, b \in G$. Then there exist *unique* elements $x, y \in G$ such that $ax = b$ and $ya = b$.

Proof available at the end of this section.

Note

We only of necessity have $(ab)^{-1} = a^{-1}b^{-1}$ if G is known to be abelian!

However, we do have the following:

Theorem 2.5.4

If a and b are elements of a group $\langle G, \cdot \rangle$, then

$$(ab)^{-1} = b^{-1}a^{-1}.$$

Proof

We have that

$$(ab)(b^{-1}a^{-1}) = a(bb^{-1})a^{-1} = aea^{-1} = aa^{-1} = e$$

Similarly, $(b^{-1}a^{-1})(ab) = e$.

Answers

Theorem 2.5.2

Let $a, b, c \in G$ and assume that $ab = ac$. Multiplying both equation sides on the left by a^{-1} , we obtain

Illegal pream-token (&)

This proves that the left cancellation law holds. A similar proof shows that the right cancellation law holds.

Theorem 2.5.3

If $x = a^{-1}b$ and $y = ba^{-1}$, then $ax = a(a^{-1}b) = (aa^{-1})b = eb = b$ and $ya = (ba^{-1})a = b(a^{-1}a) = be = b$. So such elements x and y exist. The fact that they are unique follows from the cancellation laws: if $ax = b$ and $ax' = b$ then $x = x'$ by left cancellation, and if $ya = b$ and $y'a = b$ then $y = y'$ by right cancellation.

2.6: Examples of Groups/Nongroups, Part II

Example 2.6.1

Let $n^{\text{int}}\mathbb{Z}^+$. We define $n\mathbb{Z}$ by

$$n\mathbb{Z} = \{nx : x^{\text{int}}\mathbb{Z}\} :$$

that is, $n\mathbb{Z}$ is the set of all (integer) multiples of n .

Theorem 2.6.1

$n\mathbb{Z}$ is a group under $+$ (the usual addition of integers).

Proof available at the end of this section.

Remark

When we are discussing a group $n\mathbb{Z}$, assume that $n^{\text{int}}\mathbb{Z}^+$, unless otherwise noted.

We use an example from our next class of groups all the time; in fact, most six-year-olds do as well, since it is used when telling time! Before we get to the example, we need some more definitions and some notation. Throughout the following discussion, assume n is a fixed positive integer.

Definition: Congruent modulo (mod)

We say integers a and b are **congruent modulo** [or **mod**] n if n divides $a - b$. If a and b are congruent mod n , we write $a \equiv_n b$.

Example 2.6.2

1, 7, 13, and -5 are all congruent mod 6.

The following is a profoundly useful theorem; it's so important, it has a special name. We omit the proof of this theorem, but direct interested readers to for, instance, p. 5 in [3].

Theorem 2.6.2: Division Algorithm

Let $n^{\text{int}}\mathbb{Z}^+$ and let a be any integer. Then there exist unique integers q and r , with $0 \leq r < n$, such that $a = qn + r$.

(This is actually a special case of a more general theorem, which states that given any integers n and a , there exist unique integers q and r , with $0 \leq r < |n|$, such that $a = qn + r$.)

It follows that for each positive integer n and integer a , there exists a unique element $R_n(a)$ (the r in the above theorem) of the set $\{0, 1, 2, \dots, n-1\}$ such that a is congruent to $R_n(a)$ modulo n . For example, $R_3(4) = 1$, $R_3(0) = 0$, $R_3(17) = 2$, and $R_3(-5) = 1$.

Definition: Remainder

$R_n(a)$ is the **remainder** when we divide a by n .

(Note: You were probably already familiar with the remainder when you divide a positive integer by n .)

Definition: Addition modulo

We define **addition modulo** n , $+_n$, on $\mathbb{Z}$ by, for all $a, b \in \mathbb{Z}$,

$$a +_n b = R_n(a + b),$$

that is, the unique element of $\{0, 1, \dots, n-1\}$ that's congruent to the integer $a + b$ modulo n .

Remark

Addition mod 24 is what we use to tell time!

The set $\{0, 1, 2, \dots, n-1\}$ of remainders when dividing by n is so important we give it a special notation.

Definition

We define $\mathbb{Z}_n$ to be the set $\{0, 1, 2, \dots, n-1\}$.

Note

Note that by our definition of $\mathbb{Z}_n$, the integer n itself is *not* in $\mathbb{Z}_n$!

We are now ready to consider our next type of group.

Example 2.6.3

For each $n \in \mathbb{Z}^+$, $\langle \mathbb{Z}_n, +_n \rangle$ is a group, called the **cyclic group of order n** (we will see later why we use the word “cyclic” here). This group is abelian and of order n .

Proof

We first check that $\langle \mathbb{Z}_n, +_n \rangle$ is a binary structure. Note that by the definition of $+_n$, $a +_n b \in \mathbb{Z}_n$ for each $a, b \in \mathbb{Z}$. Thus, $a +_n b \in \mathbb{Z}_n$ for each $a, b \in \mathbb{Z}_n$.

We next check that $\mathbb{Z}_n$ under $+_n$ satisfies the three group axioms. Note that since addition is commutative on $\mathbb{Z}$,

$$a +_n b = R_n(a + b) = R_n(b + a) = b +_n a$$

for all $a, b \in \mathbb{Z}_n$. Again, a simpler way of stating this is that commutativity of $+_n$ on $\mathbb{Z}_n$ is inherited from the commutativity of addition on $\mathbb{Z}$. One nice result of this is that since $+_n$ is commutative on $\mathbb{Z}_n$, we have less to check when verifying group axioms $\mathcal{G}_2$ and $\mathcal{G}_3$.

Now, let $a, b, c \in \mathbb{Z}_n$. We want to show that $(a +_n b) +_n c = a +_n (b +_n c)$. Now,

Illegal pream-token (&)

So $(a +_n b) +_n c$ and $a +_n (b +_n c)$ are congruent mod n . Since both of these values are in $\{0, 1, \dots, n-1\}$, this implies that they are equal, as desired.

Next, clearly, $0 \in \mathbb{Z}_n$ acts as an identity element under $+_n$, since

$$0 +_n a = R_n(0 + a) = R_n(a)$$

for each $a \in \mathbb{Z}_n$.

Finally, let $a \in \mathbb{Z}_n$. If $a = 0$, then clearly a has inverse $0 \in \mathbb{Z}_n$ since $0 +_n 0 = 0$. If $a \neq 0$, then the element $n - a \in \mathbb{Z}_n$ is an inverse for a since

$$a +_n (n - a) = R_n(a + (n - a)) = R_n(n) = 0.$$

Since $+_n$ is commutative on $\mathbb{Z}_n$, $\mathbb{Z}_n$ is an abelian group under $+_n$.

Finally, we already know that $|\mathbb{Z}_n| = |\{0, 1, 2, \dots, n-1\}| = n$.

Remark

In practice, we often omit the subscript n and just write $+$ when discussing addition modulo n on $\mathbb{Z}_n$.

Note

Do not confuse $n\mathbb{Z}$ and $\mathbb{Z}_n$! They are very different as sets and as groups.

Example 2.6.4

In the group $\langle \mathbb{Z}_8, + \rangle$ (where, as indicated by our above remark, $+$ means addition modulo 8), we have, for instance, $3 + 7 = 2$ and $7 + 7 = 6$. The numbers 2 and 6 are each other's inverse, and $7^{-1} = 1$. The number 0 has inverse 0 (it can't be 8, since $8 \notin \mathbb{Z}_8$!).

Definition: Multiplication modulo

For $n \in \mathbb{Z}^+$, we define **multiplication modulo n** , denoted $\cdot_n$, on $\mathbb{Z}_n$ by $a \cdot_n b = R_n(ab)$, the remainder when ab is divided by n .

Remark

$\mathbb{Z}_n$ is never a group under $\cdot_n$ (do you see why?).

But we can consider the following

Definition

For $n \in \mathbb{Z}^+$, we define $\mathbb{Z}_n^\times$ to be the set

$$\{a \in \mathbb{Z}_n : \gcd(a, n) = 1\}.$$

Example 2.6.5

$\langle \mathbb{Z}_n^\times, \cdot_n \rangle$ is a group under multiplication. We omit the proof.

We end this section by considering a few more examples.

Example 2.6.6

Let F be the set of all functions from $\mathbb{R}$ to $\mathbb{R}$, and define **pointwise addition** $+$ on F by

$$(f + g)(x) = f(x) + g(x)$$

for all $f, g \in F$ and $x \in \mathbb{R}$. We claim that F is a group under pointwise addition. (For variety, in this proof we don't explicitly refer to $\mathcal{G}_1$ – $\mathcal{G}_3$, though we certainly do verify they hold.)

Proof

If $f, g \in F$ then clearly $f + g$ is also a function from $\mathbb{R}$ to $\mathbb{R}$, so $\langle F, + \rangle$ is a binary structure.

Next, let $f, g, h \in F$. Then for all $x \in \mathbb{R}$,

Illegal pream-token (&)

Note that the key fact used in this argument is that $f(x)$, $g(x)$ and $h(x)$ all lie in $\mathbb{R}$, and addition is associative on $\mathbb{R}$. When you get used to such arguments, it is sufficient to say that associativity of $+$ on F is *inherited* from the associativity of addition on $\mathbb{R}$.

Next, let $z : \mathbb{R} \rightarrow \mathbb{R}$ be the function $z(x) = 0$ for all x . Then for all $f \in F$ and $x \in \mathbb{R}$,

Illegal pream-token (&)

So z is an identity element of $\langle F, + \rangle$.

Finally, let $f \in F$, and define $g \in F$ by $g(x) = -f(x)$ for all $x \in \mathbb{R}$. It is easy then to see that g is an inverse for f in F .

Hence, F is a group under pointwise addition. Note that it is uncountably infinite and abelian.

Example 2.6.7

The set F is *not* a group under function composition (do you see why?). But if we define B to be the set of all **bijections** from $\mathbb{R}$ to $\mathbb{R}$, then B is a group under function composition. (Prove it!) B is uncountably infinite and nonabelian.

Example 2.6.8

Let $\langle G_1, *_1 \rangle, \langle G_2, *_2 \rangle, \dots, \langle G_n, *_n \rangle$ be groups $(n \in \mathbb{Z}^+)$. Then the *group product*

$$G = G_1 \times G_2 \times \cdots \times G_n$$

is a group under the *componentwise* operation $*$ defined by

$$(g_1, g_2, \dots, g_n) * (h_1, h_2, \dots, h_n) = (g_1 *_1 h_1, g_2 *_2 h_2, \dots, g_n *_n h_n)$$

for all $(g_1, g_2, \dots, g_n), (h_1, h_2, \dots, h_n) \in G$.

For instance, considering multiplication on $\mathbb{R}^*$, matrix multiplication on $GL(2, \mathbb{R})$, and addition modulo 6 on $\mathbb{Z}_6$, we have that $\langle \mathbb{R}^* \times GL(2, \mathbb{R}) \times \mathbb{Z}_6, * \rangle$ is a group in which, for instance,

$$\left(-1, \begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix}, 3\right) * \left(\pi, \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, 4\right) = \left(-\pi, \begin{bmatrix} 5 & 4 \\ -1 & -1 \end{bmatrix}, 1\right).$$

Example 2.6.9

A common example of a group product is the group $\mathbb{Z}_2^2$, equipped with componentwise addition modulo 2.

Definition: Klein-4-group

The group $\mathbb{Z}_2^2$ is known as the **Klein 4-group**. (Felix Klein was a German mathematician; you may have heard of him in relation to the Klein Bottle.) The group $\mathbb{Z}_2^2$ is sometimes denoted by V , which stands for “Vierergruppe,” the German word for “four-group”.

Answers

Theorem 2.6.1

Let $x, y \in n\mathbb{Z}$. Then there exist $a, b \in \mathbb{Z}$ such that $x = na$ and $y = nb$. Then $x + y = na + nb = n(a + b) \in n\mathbb{Z}$. So $\langle n\mathbb{Z}, + \rangle$ is a binary structure. The remainder of the proof is left as an exercise for the reader.

2.7: Summaries of Groups We've Seen

When you see the following groups in the wild, you should assume they are equipped with the following default operations, unless otherwise noted. You should know:

- What elements the groups contain;
- What their default operations are;
- Their orders (and, if they're infinite, whether they're countably infinite or uncountable);
- Whether or not they are abelian.

Group(s)	Operation
$\mathbb{Z}, \mathbb{Q}$	Addition of Numbers
$n\mathbb{Z}$	Addition of Numbers
$\mathbb{R}, \mathbb{C}$	Addition of Numbers
$\mathbb{Q}^*, \mathbb{Q}^+$	Multiplication of Numbers
$\mathbb{R}^*, \mathbb{R}^+, \mathbb{C}^*$	Multiplication of Numbers
$M_{m \times n}(\mathbb{R}), M_n(\mathbb{R})$	Matrix Addition
$GL(n, \mathbb{R}), SL(n, \mathbb{R})$	Matrix Multiplication
$\mathbb{Z}_n$	Addition mod n
$\mathbb{Z}_2^2$	Componentwise Addition mod 2
F (see Example 2.6.6)	Pointwise Addition
B (see Example 2.6.7)	Composition

This page titled [2.7: Summaries of Groups We've Seen](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

2.8: Exercises, Part II

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem.
 - a. For every positive integer n , there exists a group of order n .
 - b. For every integer $n \geq 2$, $\mathbb{Z}_n$ is abelian.
 - c. Every abelian group is finite.
 - d. For every integer m and integer $n \geq 2$, there exist infinitely many integers a such that a is congruent to m modulo n .
 - e. A binary operation $*$ on a set S is commutative if and only if there exist $a, b \in S$ such that $a * b = b * a$.
 - f. If $\langle S, * \rangle$ is a binary structure, then the elements of S must be numbers.
 - g. If e is an identity element of a binary structure (not necessarily a group) $\langle S, * \rangle$, then e is an idempotent in S (that is, $e * e = e$).
 - h. If s is an idempotent in a binary structure (not necessarily a group) $\langle S, * \rangle$, then s must be an identity element of S .
2. Let G be the set of all functions from $\mathbb{Z}$ to $\mathbb{R}$. Prove that pointwise multiplication on G (that is, the operation defined by $(fg)(x) = f(x)g(x)$ for all $f, g \in G$) is commutative. (Note. To prove that two functions, h and j , sharing the same domain D are equal, you need to show that $h(x) = j(x)$ for every $x \in D$.)
3. Decide which of the following binary structures are groups. For each, if the binary structure *isn't* a group, prove that. (Remember, you should *not* state that inverses do or do not exist for elements until you have made sure that the structure contains an identity element!) If the binary structure *is* a group, prove that.
 - a. $\mathbb{Q}$ under multiplication
 - b. $M_2(\mathbb{R})$ under addition
 - c. $M_2(\mathbb{R})$ under multiplication
 - d. $\mathbb{R}^+$ under $*$, defined by $a * b = \sqrt{ab}$ for all $a, b \in \mathbb{R}^+$
4. Give an example of an abelian group containing 711 elements.
5. Let $n\mathbb{Z}$. Prove that $n\mathbb{Z}$ is a group under the usual addition of integers. Note: You may use the fact that $\langle n\mathbb{Z}, + \rangle$ is a binary structure if you provide a reference for this fact.
6. Let $n\mathbb{Z}^+$. Prove that $SL(n, \mathbb{R})$ is a group under matrix multiplication. Note: You may use the fact that $\langle SL(n, \mathbb{R}), \cdot \rangle$ is a binary structure if you provide a reference for this fact.
7.
 - a. List three distinct integers that are congruent to 6 modulo 5.
 - b. List the elements of $\mathbb{Z}_5$.
 - c. Compute:
 - i. $4 + 5$ in $\mathbb{Z}$;
 - ii. $4 + 5$ in $\mathbb{Q}$;
 - iii. $4 +_6 5$ in $\mathbb{Z}_6$;
 - iv. the inverse of 4 in $\mathbb{Z}$;
 - v. the inverse of 4 in $\mathbb{Z}_6$.
 - d. Why does it not make sense for me to ask you to compute $4 +_3 2$ in $\mathbb{Z}_3$? Please answer this using a complete, grammatically correct sentence.
8. Let G be a group with identity element e . Prove that if every element of G is its own inverse, then G is abelian.
9. Let G be a group. The subset

$$Z(G) := \{z \in G : zg = gz \text{ for all } g \in G\}$$

of G is called the *center* of G . In other words, $Z(G)$ is the set of all elements of G that commute with every element of G . Prove that $Z(G)$ is closed in G .

3: HOMOMORPHISMS AND ISOMORPHISMS

3.1: Groups of Small Order

3.2: Definitions of Homomorphisms and Isomorphisms

3.3: Isomorphic Groups

3.4: Exercises

This page titled [3: Homomorphisms and Isomorphisms](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

3.1: Groups of Small Order

Let's start exploring groups in order of increasing, um, order. Before we do this, it will be helpful to introduce the notion of a **group table** (also known as a **Cayley table**¹) for a finite group. Given a finite group G , list its elements in some fixed order, say, $a_1, a_2, \dots, a_n$, and then construct its group table by creating an array with exactly one row and exactly one column corresponding to each group element. We then put in the row i and column j the element $a_i a_j$ of G . Note that a single group can have group tables that look different from one another, since reordering a group's elements will change its table.

Example 3.1.1

Consider the group $\mathbb{Z}_4$ under addition modulo 4. Ordering the elements of $\mathbb{Z}_4$ as 1, 2, 3, 4, we have the following group table for $\langle \mathbb{Z}_4, + \rangle$:

+	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

Now, clearly, there is no group of order 0 (do you see why?). Is there a group of order 1? Well, suppose $\langle G, * \rangle$ is such a group. Since G must contain an identity element e , we must have $G = \{e\}$, and since e is G 's identity element, we must have $e * e = e$. Clearly, in this case, the three group axioms hold. So G is a valid group, without much going on in it.

Definition: Trivial Group

If G is a group with $|G| = 1$, then G is called the **trivial group**.

Remark

A good question to ask here is why it's called "the" trivial group, rather than "a" trivial group. Indeed, there are infinitely many groups of order 1! (Do you see why?) But it turns out that all of these groups are *structurally* the same. Hence mathematicians end up thinking of them as various instantiations of one group, rather than separate groups. We will discuss this in more depth shortly, when we introduce the idea of *isomorphism*.

Next suppose that group $\langle G, \cdot \rangle$ has order 2. Then G must contain an identity element, e , and a non-identity element, a . Since e is its own inverse, and inverses are unique, a must be its own inverse as well. So G must have the following table.

*	e	a
e	e	a
a	a	e

It is straightforward to show that such a structure does satisfy all the group axioms (the only one we really need to check is associativity).

Now, what if group $\langle G, \cdot \rangle$ has order 3? Note that *you can't have any entry appear more than once in the same row or same column* (excluding of course the labels outside the grid we're filling in), given Theorem 2.5.3. Is there only one way of filling in the table for a group of order 3? (Hint: consider what element must be the second row, third column entry.)

*	e	a	b
e			
a			
b			

Finally, what if group $\langle G, \cdot \rangle$ has order 4? It turns out in this case there are *two* valid ways of filling in a group table!

What we have been doing here is really getting into the idea of the *structure* of groups, and when we can consider groups to be essentially “the same” or fundamentally “different.” We approach this more formally via the concepts of homomorphism and isomorphism.

Notes:

1. Named after the British mathematician Arthur Cayley.

3.2: Definitions of Homomorphisms and Isomorphisms

Definition: Homomorphism and Isomorphism

Let $\langle S, * \rangle$ and $\langle S', *' \rangle$ be binary structures. A function ϕ from S to S' is a **homomorphism** if

$$\phi(a * b) = \phi(a) *' \phi(b)$$

for all $a, b \in S$. An **isomorphism** is a homomorphism that is also a bijection.

Intuitively, you can think of a homomorphism ϕ as a “structure-preserving” map: if you multiply and then apply ϕ , you get the same result as when you first apply ϕ and then multiply. Isomorphisms, then, are both structure-preserving and cardinality-preserving.

Note

We may omit the $*$ and $*'$, as per our group conventions, but we include them here to emphasize that the operations in the structures may be distinct from one another. When we omit them and write $\phi(st) = \phi(s)\phi(t)$, then it is the writers' and readers' responsibility to keep in mind that s and t are being operated together using the operation in S , while $\phi(s)$ and $\phi(t)$ are being operated together using the operation in S' .

Remark

There may be more than one homomorphism [isomorphism] from one binary structure to another (see Example 3.2.1).

Example 3.2.1

For each of the following, decide whether or not the given function ϕ from one binary structure to another is a homomorphism, and, if so, if it is an isomorphism. Prove or disprove your answers! For Parts 6 and 7, C^0 is the set of all continuous functions from $\mathbb{R}$ to $\mathbb{R}$; C^1 is the set of all differentiable functions from $\mathbb{R}$ to $\mathbb{R}$ whose derivatives are continuous; and each $+$ indicates pointwise addition on C^0 and C^1 .

1. $\phi : \langle \mathbb{Z}, + \rangle \rightarrow \langle \mathbb{Z}, + \rangle$ defined by $\phi(x) = x$;
2. $\phi : \langle \mathbb{Z}, + \rangle \rightarrow \langle \mathbb{Z}, + \rangle$ defined by $\phi(x) = -x$;
3. $\phi : \langle \mathbb{Z}, + \rangle \rightarrow \langle \mathbb{Z}, + \rangle$ defined by $\phi(x) = 2x$;
4. $\phi : \langle \mathbb{R}, + \rangle \rightarrow \langle \mathbb{R}^+, \cdot \rangle$ defined by $\phi(x) = e^x$;
5. $\phi : \langle \mathbb{R}, + \rangle \rightarrow \langle \mathbb{R}^*, \cdot \rangle$ defined by $\phi(x) = e^x$;
6. $\phi : \langle C^1, + \rangle \rightarrow \langle C^0, + \rangle$ defined by $\phi(f) = f'$ (the derivative of f);
7. $\phi : \langle C^0, + \rangle \rightarrow \langle \mathbb{R}, + \rangle$ defined by $\phi(f) = \int_0^1 f(x) dx$.

Example 3.2.2

Let $\langle G, \cdot \rangle$ be a group and let $a \in G$. Then the function c_a from G to G defined by $c_a(x) = axa^{-1}$ (for all $x \in G$) is a homomorphism. Indeed, let $x, y \in G$. Then

Illegal pream-token (&)

The homomorphism c_a is called **conjugation** by a .

Definition: Endomorphism and Automorphism

Homomorphisms from a group G to itself are called **endomorphisms**, and isomorphisms from a group to itself are called **automorphisms**.

It can be shown that conjugation by any element a of a group G is a bijection from G to itself (can you prove this?), so such conjugation is an automorphism of G . (Beware: Some texts use “conjugation by a ” to refer to the function $x \mapsto a^{-1}xa$.) Both versions of conjugation by a in group G are automorphisms of G .)

We end with a theorem stating basic facts about homomorphisms from one group to another. (*Note.* This doesn't apply to arbitrary binary structures, which may or may not even have identity elements.)

Theorem 3.2.1

Let $\langle G, \cdot \rangle$ and $\langle G', \cdot' \rangle$ be groups with identity elements e and e' , respectively, and let ϕ be a homomorphism from G to G' . Then:

1. $\phi(e) = e'$; and
2. For every $a \in G$, $\phi(a)^{-1} = \phi(a^{-1})$.

Proof available at the end of this section.

Answers

Theorem 3.2.1

For Part 1, note that

Illegal pream-token (&)

Thus, by left cancellation, $e' = \phi(e)$. The proof of Part 2 is left as an exercise for the reader.

3.3: Isomorphic Groups

One of the key ideas we've discussed in determining whether binary structures are essentially "the same" or "different." We approach this rigorously using the concept of *isomorphic* groups.

Definition: Isomorphic and Nonisomorphic

We say that two groups G and G' (or binary structures S and S') are **isomorphic** (or G is isomorphic to G'), and write $G \simeq G'$, if there exists an isomorphism from G to G' . We say that G is **isomorphic to G' via ϕ** if ϕ is an isomorphism from G to G' . If there exists no isomorphism from G and G' , then we say that G and G' are **nonisomorphic** (or G is not isomorphic to G'), and write $G \not\simeq G'$.

Note

Just because a particular map (even an "obvious" one) from group G to group G' is not an isomorphism, we do not know that G and G' are not isomorphic! For instance, the map $\phi : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $\phi(x) = 2x$ for all x is *not* an isomorphism (since it's not onto), but $\mathbb{Z}$ is isomorphic to itself, as we will see in Part 1 of Theorem 3.3.1.

Isomorphic groups have *the same structure* as far as algebraists are concerned. Again, picture two houses that are identical except for the colors they are painted. Though they differ in some ways (one house is red while the other is green), they are structurally identical. Isomorphic groups have identical structures, though the elements of one group may differ greatly from those of the other. Returning to the house analogy: if two houses are structurally identical, we can learn many things about one house by looking at the other (e.g., how many bathrooms it has, whether it has a basement, etc.). Similarly, suppose we know a great deal about group G and are given a new group, G' . If we prove that G' is isomorphic to G , then we can likely deduce information about G' from the information we know about G .

Theorem 3.3.1

Let $\langle G, \cdot \rangle$, $\langle G', \cdot' \rangle$, and $\langle G'', \cdot'' \rangle$ be groups.

1. Group G is isomorphic to itself.
2. If ϕ is an isomorphism from G to group G' , then there exists an isomorphism from G' to G . Hence, $G \simeq G'$ if and only if $G' \simeq G$.
3. If $G \simeq G'$ and $G' \simeq G''$, then $G \simeq G''$.

Proof available at the end of this section.

Remark

To show that given groups G and G' are isomorphic, we must do three things:

1. *Define* a function ϕ from G to G' (or from G' to G , as we have Theorem 3.3.1).
2. Prove that ϕ is a homomorphism.
3. Prove that ϕ is a bijection.

Note

Remember, you can show that ϕ is a bijection by proving that it's one-to-one and onto, *or* by showing that it has an inverse.

Note

Do **NOT** try to prove that a function ϕ is an isomorphism WITHOUT DEFINING ϕ !

We now provide some terminology that will be helpful for our study of the structures of groups.

Definition: Unique Group

Given a certain property (or properties), we say there is a **unique group** with that property (or properties) **up to isomorphism** if any two groups sharing that property (or properties) are isomorphic to one another.

This may seem a little abstruse at the moment, but seeing examples will help illuminate the concept.

Example 3.3.1

1. If G and G' are groups with $|G| = |G'| = 1$, then $G \simeq G'$, since the map from G to G' sending G 's identity (and sole) element to G' 's identity (and sole) element is clearly an isomorphism. This is why we can mildly abuse terminology and call any group of order 1 *the* trivial group instead of *a* trivial group: G and G' may technically be different groups, but structurally they are identical, so we can consider them to be more or less "the same." Thus, there is a unique group of order 1, up to isomorphism.
2. Let G be a group with $|G| = 2$. Then G must consist of an identity element e and a nonidentity element a , and have the following group table. Compare the respective group tables for G and for the specific two-element group $\mathbb{Z}_2$.

*	e	a
e	e	a
a	a	e

+	0	1
0	0	1
1	1	0

Note that the first table looks exactly like the second table if we replace $*$ with $+$, each e with 0, and each a with 1. This shows that groups G and $\mathbb{Z}_2$ have identical structures; more precisely, it shows that the function ϕ from G to $\mathbb{Z}_2$ defined by $\phi(e) = 0$ and $\phi(a) = 1$ is an isomorphism. Since any group of order 2 is isomorphic to $\mathbb{Z}_2$, using Theorem 3.3.1 we see that there is a unique group of order 2, up to isomorphism.

3. A similar argument shows that there is a unique group of order 3 up to isomorphism: specifically, any group of order 3 is isomorphic to $\mathbb{Z}_3$.
4. We will see later, in Example 3.3.6, that there is *not* a unique group of order 4 up to isomorphism: that is, there are two nonisomorphic groups of order 4.

Theorem 3.3.2

The groups $\langle \mathbb{R}, + \rangle$ and $\langle \mathbb{R}^+, \cdot \rangle$ are isomorphic.

Proof available at the end of this section.

Example 3.3.2

Let $n\mathbb{Z}^+$. Then the groups $\langle n\mathbb{Z}, + \rangle$ and $\langle \mathbb{Z}, + \rangle$ are isomorphic. The proof of this is left as an exercise for the reader.

We have now seen examples in which we have proved that two groups are isomorphic. How, though, do we prove that two groups are *not* isomorphic? It is usually wildly impractical, if not impossible, to check that no function from one group to the other is an isomorphism. For instance, it turns out that $\mathbb{R}^*$ is not isomorphic to $GL(2, \mathbb{R})$ (see Example 3.3.3), but there are infinitely many bijections from $\mathbb{R}^*$ to $GL(2, \mathbb{R})$ —it is impossible to check that each one is not an isomorphism. Instead, we use **group invariants**.

Definition: Group Invariant

A group property is called a **group invariant** if it is preserved under isomorphism.

Group invariants are **structural properties**. Some examples of group invariants are:

1. Cardinality (since any isomorphism between groups is a bijection);
2. Abelianness (the proof that this is a group invariant is left as an exercise for the reader);
3. Number of elements which are their own inverses (proven by an argument similar to that in Example 3.3.6).

A **nonexample** of a group invariant is the property of being a subset of $\mathbb{R}$.

Example 3.3.3

The group $\mathbb{R}$ cannot be isomorphic to the group $GL(2, \mathbb{R})$ since the former group is abelian and the latter nonabelian.

Example 3.3.4

The groups $\mathbb{R}$ and $\mathbb{Q}$ cannot be isomorphic since the former group is uncountable and the latter countable.

Sometimes we must resort to trickier methods in order to decide whether or not two groups are isomorphic.

Example 3.3.5

The groups $\mathbb{Z}$ and $\mathbb{Q}$ are not isomorphic. We use contradiction to prove this. Suppose that $\mathbb{Z}$ and $\mathbb{Q}$ are isomorphic via isomorphism $\phi: \mathbb{Q} \rightarrow \mathbb{Z}$. Let $a \in \mathbb{Q}$. Then $\frac{a}{2} \in \mathbb{Q}$ with $\frac{a}{2} + \frac{a}{2} = a$. Then

$$\phi\left(\frac{a}{2}\right) + \phi\left(\frac{a}{2}\right) = \phi\left(\frac{a}{2} + \frac{a}{2}\right) = \phi(a);$$

since $\phi\left(\frac{a}{2}\right)$ is in $\mathbb{Z}$, $\phi(a)$ must be evenly divisible by 2. But a was arbitrary in $\mathbb{Q}$ and ϕ is onto, so this means every element of $\mathbb{Z}$ must be evenly divisible by 2, which is clearly false. Thus, $\mathbb{Z} \not\cong \mathbb{Q}$.

Example 3.3.6

The groups $\mathbb{Z}_4$ and $V = \mathbb{Z}_2^2$ are not isomorphic. Why? Well, they are both abelian of order 4, so we cannot use cardinality or commutativity to prove they are nonisomorphic. The gist of our argument will be to note that every element in V is its own inverse; so if V and $\mathbb{Z}_4$ are isomorphic (hence structurally identical) we must have that every element of $\mathbb{Z}_4$ is also its own inverse, which doesn't hold (e.g., in $\mathbb{Z}_4$, $3 + 3 = 2$, not 0).

A rigorous proof of the fact that V and $\mathbb{Z}_4$ are not isomorphic is as follows: For now, denote the usual operation on V by $*$. Suppose that V and $\mathbb{Z}_4$ are isomorphic, via isomorphism ϕ from V to $\mathbb{Z}_4$. Then since ϕ is onto, there exists an element $a \in V$ such that $\phi(a) = 3$. Then

Illegal pream-token (&)

since ϕ is a homomorphism, so sends identity element to identity element. But this is a contradiction, since $3 + 3 = 2 \neq 0$ in $\mathbb{Z}_4$. Thus, $V \not\cong \mathbb{Z}_4$.

Answers

Theorem 3.3.1

For Part 1: The identity map $1_G : G \rightarrow G$ defined by $1_G(a) = a$ for all $a \in G$ is clearly an isomorphism.

For Part 2: Since ϕ is an isomorphism, it's a bijection, hence has inverse ϕ^{-1} . From Theorem 1.2.1, we know that ϕ^{-1} must also be a bijection (in this case, from G' to G). So it suffices to show that ϕ^{-1} is a homomorphism. Let $a, b \in G'$. We want to show that $\phi^{-1}(a \cdot' b) = \phi^{-1}(a) \cdot \phi^{-1}(b)$. Since ϕ is 1-1, it suffices to show that

$$\phi(\phi^{-1}(a \cdot' b)) = \phi(\phi^{-1}(a) \cdot \phi^{-1}(b)).$$

Notice, we have $\phi(\phi^{-1}(a \cdot' b)) = a \cdot' b$; further, we have

$$\phi(\phi^{-1}(a) \cdot \phi^{-1}(b)) = \phi(\phi^{-1}(a)) \cdot' \phi(\phi^{-1}(b)) = a \cdot' b.$$

This shows that $\phi^{-1}(a \cdot' b) = \phi^{-1}(a) \cdot \phi^{-1}(b)$, as desired.

For Part 3: Since $G \simeq G'$ and $G' \simeq G''$, there exist isomorphisms $\phi : G \rightarrow G'$ and $\psi : G' \rightarrow G''$. Define $\theta : G \rightarrow G''$ by $\theta = \psi \circ \phi$. Since ϕ and ψ are both bijections, θ is a bijection (Theorem 1.2.3). Next, let $a, b \in G$. The

Illegal pream-token (&)

Thus, θ is a homomorphism, and hence, since it is also a bijection, an isomorphism.

Theorem 3.3.2

Define $\phi : \mathbb{R} \rightarrow \mathbb{R}^+$ by $\phi(x) = e^x$. Our map ϕ is a homomorphism since for every $x, y \in \mathbb{R}$, we have

$$\phi(x + y) = e^{x+y} = e^x e^y = \phi(x) \phi(y).$$

Moreover, ϕ is a bijection, since it has inverse function $\phi^{-1} : \mathbb{R}^+ \rightarrow \mathbb{R}$ defined by $\phi^{-1}(x) = \ln x$. Hence, $\mathbb{R} \simeq \mathbb{R}^+$ via isomorphism ϕ .

3.4: Exercises

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G and G' be groups.
 - a. If there exists a homomorphism $\phi : G \rightarrow G'$, then G and G' must be isomorphic groups.
 - b. There is an integer $n \geq 2$ such that $\mathbb{Z} \simeq \mathbb{Z}_n$.
 - c. If $|G| = |G'| = 3$, then we must have $G \simeq G'$.
 - d. If $|G| = |G'| = 4$, then we must have $G \simeq G'$.
2. For each of the following functions, prove or disprove that the function is (i) a homomorphism; (ii) an isomorphism. (Remember to work with the default operation on each of these groups!)
 - a. The function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(n) = 2n$.
 - b. The function $g : \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) = x^2$.
 - c. The function $h : \mathbb{Q}^* \rightarrow \mathbb{Q}^*$ defined by $h(x) = x^2$.
3. Define $d : GL(2, \mathbb{R}) \rightarrow \mathbb{R}^*$ by $d(A) = \det A$. Prove/disprove that d is:
 - a. a homomorphism
 - b. 1-1
 - c. onto
 - d. an isomorphism.
4. Complete the group tables for $\mathbb{Z}_4$ and $\mathbb{Z}_8^\times$. Use the group tables to decide whether or not $\mathbb{Z}_4$ and $\mathbb{Z}_8^\times$ are isomorphic to one another. (You do not need to provide a proof.)
5. Let $n^{\text{int}}\mathbb{Z}^+$. Prove that $\langle n\mathbb{Z}, + \rangle \simeq \langle \mathbb{Z}, + \rangle$.
6.
 - a. Let G and G' be groups, where G is abelian and $G \simeq G'$. Prove that G' is abelian.
 - b. Give an example of groups G and G' , where G is abelian and there exists a homomorphism from G to G' , but G' is NOT abelian.
7. Let $\langle G, \cdot \rangle$ and $\langle G', \cdot' \rangle$ be groups with identity elements e and e' , respectively, and let ϕ be a homomorphism from G to G' . Let $a^{\text{int}}G$. Prove that $\phi(a)^{-1} = \phi(a^{-1})$.

4: SUBGROUPS

4.1: Introduction to Subgroups

4.2: Subgroup Proofs and Lattices

4.3: Exercises

This page titled [4: Subgroups](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

4.1: Introduction to Subgroups

Sometimes groups are too complicated to understand directly. One method that can be used to identify a group's structure is to study its **subgroups**.

Definition: Subgroup

A **subgroup** of a group G is a subset of G that is also a group under G 's operation. If H is a subgroup of G , we write $H \leq G$; if $H \subseteq G$ is not a subgroup of G , we write $H \not\leq G$.

Note

Do not confuse *subgroups* with *subsets*! All subgroups of a group G are, by definition, subsets of G , but not all subsets of G are subgroups of G (see Example 4.1.1, Parts 2–5, below). Whether or not a subset of G is a subgroup of G depends on the operation of G .

Example 4.1.1

1. Consider the subset $\mathbb{Z}$ of the group $\mathbb{Q}$, assuming that $\mathbb{Q}$ is equipped with the usual addition of real numbers (as we indicated above that we would assume, by default). Since we already know that $\mathbb{Z}$ is a group under this operation, $\mathbb{Z}$ is not just a subset but in fact a subgroup of $\mathbb{Q}$ (under addition).
2. Instead, consider the subset $\mathbb{Q}^+$ of the group $\mathbb{Q}$. This subset is *not* a group under $\mathbb{Q}$'s operation $+$, since it does not contain an identity element for $+$. Therefore, $\mathbb{Q}^+$ is a subset but not a subgroup of $\mathbb{Q}$.
3. Let I be the subset

$$I = \mathbb{R} - \mathbb{Q} = \{x^{\text{int}}\mathbb{R} : x \text{ is irrational}\}$$

of the group $\mathbb{R}$. The set I is *not* a group under $\mathbb{R}$'s operation $+$ since it is not closed under addition: for instance, $\pi, -\pi^{\text{int}}I$, but $\pi + (-\pi) = 0 \notin I$. So I is a subset but not a subgroup of $\mathbb{R}$.

4. Consider the subset $\mathbb{Z}^+$ of the group $\mathbb{R}^+$. The set $\mathbb{Z}^+$ is closed under multiplication, multiplication is associative on $\mathbb{Z}^+$, and $\mathbb{Z}^+$ does contain an identity element (namely, 1). However, most elements of $\mathbb{Z}^+$ do not have inverses in $\mathbb{Z}^+$ under multiplication: for instance, the inverse of 3 would have to be $\frac{1}{3}$, but $\frac{1}{3} \notin \mathbb{Z}^+$. Therefore, $\mathbb{Z}^+$ is a subset but not a subgroup of $\mathbb{R}^+$.
5. Consider the subset $GL(n, \mathbb{R})$ of $M_n(\mathbb{R})$. We know that $GL(n, \mathbb{R})$ is a group, so it might be tempting to say that it is a subgroup of $M_n(\mathbb{R})$; to be a subgroup of $M_n(\mathbb{R})$, $GL(n, \mathbb{R})$ must be a group under $M_n(\mathbb{R})$'s operation, which is matrix addition. While $GL(n, \mathbb{R})$ is a group under matrix multiplication, it is not a group under matrix addition: for instance, it is not closed under matrix addition, since $I_n, -I_n^{\text{int}}GL(n, \mathbb{R})$ but $I_n + (-I_n)$ is the matrix consisting of all zeros, which is not in $GL(n, \mathbb{R})$. So $GL(n, \mathbb{R})$ is a subset but not a subgroup of $M_n(\mathbb{R})$.
6. Consider the subset $H = \{0, 2\}$ of $\mathbb{Z}_4$. The subset H is closed under addition modulo 4 ($0 + 0 = 0, 0 + 2 = 2 + 0 = 2, 2 + 2 = 0$), addition modulo 4 is always associative, H contains an identity element (namely, 0) under addition modulo 4, and both 0 and 2 have inverses in $\mathbb{Z}_4$ under this operation (0 and 2 are each their own inverses). Thus, H is a subgroup of $\mathbb{Z}_4$.
7. Let G be a group. Then $\{e_G\}$ and G are clearly both subgroups of G .

Definition: Trivial, Nontrivial, Proper, and Improper Subgroup

Let G be a group. The subgroups $\{e_G\}$ and G of G are called the **trivial subgroup** and the **improper subgroup** of G , respectively. Not surprisingly, if $H \leq G$ and $H \neq \{e_G\}$, H is called a **nontrivial subgroup** of G , and if $H \leq G$ and $H \neq G$, H is called a **proper subgroup** of G .

Remark

Sometimes the notation $H < G$ is used to indicate that H is a proper subgroup of G , but sometimes it is simply used to mean that H is a subgroup—proper or improper—of G . We will not use the notation $H < G$ in this text.

Notice that in the cases above, we saw subsets of groups fail to be subgroups because they were not closed under the groups' operations; because they did not contain identity elements; or because they didn't contain an inverse for each of their elements. None, however, failed because the relevant group's operation was not associative on them. This is not a coincidence: rather, since any element of a subset of a group G also lives in G , any associative operation on G is of necessity associative on any closed subset of G . Therefore, when we are checking to see if $H \subseteq G$ is a subgroup of group G , we need only check for closure, an identity element, and inverses.

Lemma 4.1.1

Let G be a group.

1. If H is a subgroup of G then the identity element e_H of H is e_G , the identity element of G .
2. If H is a subgroup of G and $a^{\text{int}} H$ has inverse a^{-1} in G , then a 's inverse in H is also a^{-1} .

Proof available at the end of this section.

Corollary 4.1.1

Let $H \subseteq G$. If the identity element of G is not in H , then $H \not\leq G$.

Answers

Lemma 4.1.1

For Part 1: Since e_H is in both H and G , by the definition of e_H , we have $e_H e_H = e_H$, and by the definition of e_G we have $e_G e_H = e_H$. So $e_H e_H = e_G e_H$, and thus by right cancellation, $e_H = e_G$.

Next, for Part 2, let b be the inverse of a in H . Then using Part 1 of this lemma and the definition of an inverse, $ab = e_H = e_G = aa^{-1}$. By left cancellation, then, we have that $b = a^{-1}$.

4.2: Subgroup Proofs and Lattices

Using Lemma 4.1.1 and the argument preceding it, we have the following.

Theorem 4.2.1

A subset H of a group G is a subgroup of G if and only if

1. H is closed under G 's operation;
2. The identity element of G is in H ; and
3. For each $a \in H$, a 's inverse in G is contained in H .

Example 4.2.1

For each of the following, prove that the given subset H of group G is or is not a subgroup of G .

1. $H = 3\mathbb{Z}$, $G = \mathbb{Z}$.
2. $H = \{0, 1, 2, 3\}$, $G = \mathbb{Z}_6$;
3. $H = \mathbb{R}^*$, $G = \mathbb{R}$;
4. $H = \{(0, x, y, z) : x, y, z \in \mathbb{R}\}$, $G = \mathbb{R}^4$.

Example 4.2.2

Generalizing Part 1 of the above theorem, we have $n\mathbb{Z} \leq \mathbb{Z}$ for every $n \in \mathbb{Z}^+$.

The proof of this is left as an exercise for the reader.

Example 4.2.3

Consider the group $\langle F, + \rangle$, where F is the set of all functions from $\mathbb{R}$ to $\mathbb{R}$ and $+$ is pointwise addition. Which of the following are subgroups of F ?

1. $H = \{f \in F : f(5) = 0\}$;
2. $K = \{f \in F : f \text{ is continuous}\}$;
3. $L = \{f \in F : f \text{ is differentiable}\}$.

Are any of H , K , and L subgroups of one another?

In fact, we can narrow down the number of facts we need to check to prove a subset $H \subseteq G$ is a subgroup of G to only two.

Theorem 4.2.2: Two-Step Subgroup Test

Let G be a group and $H \subseteq G$. Then H is a subgroup of G if

1. $H \neq \emptyset$; and
2. For each $a, b \in H$, $ab^{-1} \in H$. Double exponent: use braces to clarify

Proof available at the end of this section.

Example 4.2.4

1. Use the Two-Step Subgroup Test to prove that $3\mathbb{Z}$ is a subgroup of $\mathbb{Z}$.
2. Use the Two-Step Subgroup Test to prove that $SL(n, \mathbb{R})$ is a subgroup of $GL(n, \mathbb{R})$.

It is straightforward to prove the following theorem.


Theorem 4.2.3

If H is a subgroup of a group G and K is a subset of H , then K is a subgroup of H if and only if it's a subgroup of G .

It can be useful to look at how subgroups of a group relate to one another. One way of doing this is to consider **subgroup lattices** (also known as **subgroup diagrams**). To draw a subgroup lattice for a group G , we list all the subgroups of G , writing a subgroup K below a subgroup H , and connecting them with a line, if K is a subgroup of H and there is no proper subgroup L of H that contains K as a proper subgroup.

Example 4.2.5

Consider the group $\mathbb{Z}_8$. We will see later that the subgroups of $\mathbb{Z}_8$ are $\{0\}$, $\{0, 2, 4, 6\}$, $\{0, 4\}$ and $\mathbb{Z}_8$ itself. So $\mathbb{Z}_8$ has the following subgroup lattice.



Example 4.2.6

Referring to Example 4.2.3, draw the portion of the subgroup lattice for F that shows the relationships between itself and its proper subgroups H , K , and L .

Example 4.2.7

Indicate the subgroup relationships between the following groups: $\mathbb{Z}$, $12\mathbb{Z}$, $\mathbb{Q}^+$, $\mathbb{R}$, $6\mathbb{Z}$, $\mathbb{R}^+$, $3\mathbb{Z}$, $G = \langle \{\pi^n : n \in \mathbb{Z}\}, \cdot \rangle$ and $J = \langle \{6^n : n \in \mathbb{Z}\}, \cdot \rangle$.

We end with a theorem about homomorphisms and subgroups that leads us to another group invariant.

Theorem 4.2.4

Let G and G' be groups, let ϕ a homomorphism from G to G' , and let H a subgroup of G . Then $\phi(H)$ is a subgroup of G' .

Proof available at the end of this section.

Corollary 4.2.1


The probabilities assigned to events by a distribution function on a sample space are given by.

Proof available at the end of this section.

This is another way of, for instance, distinguishing between the groups $\mathbb{Z}_4$ and the Klein 4-group $\mathbb{Z}_2^2$.

Example 4.2.8

By inspection, $\mathbb{Z}_4$ and $\mathbb{Z}_2^2$ have, respectively, the following subgroup lattices.



Since $\mathbb{Z}_4$ contains exactly 3 subgroups and $\mathbb{Z}_2^2$ exactly 5, we have that $\mathbb{Z}_4 \not\cong \mathbb{Z}_2^2$.

Answers

Theorem 4.2.2: Two-Step Subgroup Test

Assume that the above two properties hold. Since $H \neq \emptyset$, there exists an $x \in H$ such that $x \in H$. Then $e_G = xx^{-1}$ is in H , by the second property. Next, for every $a \in H$ we have $a^{-1} = e_G a^{-1} \in H$ (again by the second property). Finally, if $a, b \in H$ then we've already shown $b^{-1} \in H$; so $ab = a(b^{-1})^{-1} \in H$, yet again by the second property. Thus, $H \leq G$.

Theorem 4.2.4

The proof is left as an exercise for the reader.

Corollary 4.2.1

If $G \simeq G'$ and G contains exactly n subgroups ($n^{\text{int}} \mathbb{Z}^+$), then so does G' .

4.3: Exercises

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G and G' be groups.
 - a. Every group contains at least two distinct subgroups.
 - b. If H is a proper subgroup of group G and G is finite, then we must have $|H| < |G|$.
 - c. $7\mathbb{Z}$ is a subgroup of $14\mathbb{Z}$.
 - d. A group G may have two distinct proper subgroups which are isomorphic (to one another).
2. Give specific, precise examples of the following groups G with subgroups H :
 - a. A group G with a proper subgroup H of G such that $|H| = |G|$.
 - b. A group G of order 12 containing a subgroup H with $|H| = 3$.
 - c. A nonabelian group G containing a nontrivial abelian subgroup H .
 - d. A finite subgroup H of an infinite group G .
3. Let $n^{\text{int}}\mathbb{Z}^+$.
 - a. Prove that $n\mathbb{Z} \leq \mathbb{Z}$.
 - b. Prove that the set $H = \{A^{\text{int}}\mathbb{M}_n(\mathbb{R}) : \det A = \pm 1\}$ is a subgroup of $GL(n, \mathbb{R})$.

(Note: Your proofs do not need to be long to be correct!)
4. Let $n^{\text{int}}\mathbb{Z}^+$. For each group G and subset H , decide whether or not H is a subgroup of G . In the cases in which H is *not* a subgroup of G , provide a proof. (Note. Your proofs do not need to be long to be correct!)
 - a. $G = \mathbb{R}, H = \mathbb{Z}$
 - b. $G = \mathbb{Z}_{15}, H = \{0, 5, 10\}$
 - c. $G = \mathbb{Z}_{15}, H = \{0, 4, 8, 12\}$
 - d. $G = \mathbb{C}, H = \mathbb{R}^*$
 - e. $G = \mathbb{C}^*, H = \{1, i, -1, -i\}$
 - f. $G = \mathbb{M}_n(\mathbb{R}), H = GL(n, \mathbb{R})$
 - g. $G = GL(n, \mathbb{R}), H = \{A^{\text{int}}\mathbb{M}_n(\mathbb{R}) : \det A = -1\}$
5. Let G and G' be groups, let ϕ be a homomorphism from G to G' , and let H be a subgroup of G . Prove that $\phi(H)$ is a subgroup of G' .
6. Let G be an abelian group, and let $U = \{g^{\text{int}}G : g^{-1} = g\}$. Prove that U is a subgroup of G .

5: CYCLIC GROUPS

5.1: Introduction to Cyclic Groups

5.2: The Subgroup Lattices of Cyclic Groups

5.3: Exercises

This page titled [5: Cyclic Groups](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

5.1: Introduction to Cyclic Groups

Certain groups and subgroups of groups have particularly nice structures.

Definition: Cyclic

A group is **cyclic** if it is isomorphic to $\mathbb{Z}_n$ for some $n \geq 1$, or if it is isomorphic to $\mathbb{Z}$.

Example 5.1.1

Examples/nonexamples of cyclic groups.

1. $n\mathbb{Z}$ and $\mathbb{Z}_n$ are cyclic for every $n \in \mathbb{Z}^+$.
2. $\mathbb{R}$, $\mathbb{R}^*$, $M_2(\mathbb{R})$, and $GL(2, \mathbb{R})$ are uncountable and hence can't be cyclic.
3. $\mathbb{Z}_2^2$ is *not* cyclic since it would have to be isomorphic to $\mathbb{Z}_4$ if it were (since it has order 4).
4. $\mathbb{Q}$ isn't cyclic. If it were cyclic it would have to be isomorphic to $\mathbb{Z}$, since $\mathbb{Q}$ is an infinite group (so can't be isomorphic to $\mathbb{Z}_n$ for any n). But we showed in Example 3.3.5 that $\mathbb{Q} \neq \mathbb{Z}$.

Remark

Clearly, cyclicity is a group invariant.

Theorem 5.1.1

If G is cyclic, then G is abelian; however, G can be abelian but not cyclic.

Proof available at the end of this section.

Definition: Cyclic Subgroup of G Generated by a

Let a be an element of a group G . Then

$$\langle a \rangle = \{a^n : n \in \mathbb{Z}\}$$

is called the **(cyclic) subgroup of G generated by a** . (We will later see why the words “cyclic” and “generated” are used here.)

Remark

Note that in the above definition, we were using multiplicative notation. Using additive notation, we have

$$\langle a \rangle = \{na : n \in \mathbb{Z}\} = \{\dots, -2a, -a, 0a, 1a, 2a, \dots\}.$$

Theorem 5.1.2

Let a be an element of a group G . Then $\langle a \rangle \leq G$.

Proof available at the end of this section.

We next prove a lemma to which we will refer when proving the theorem which succeeds it.

Lemma 5.1.1: Clock Lemma

Let G be a group and let $a \in G$. Suppose there is a positive power of a which equals the identity element e of G . Let n be the least such positive integer. Then for all $s, t \in \mathbb{Z}$, $a^s = a^t$ if and only if s is congruent to t modulo n . In particular, if $a^s = e$, then n divides s .

Proof available at the end of this section.

Theorem 5.1.3

Let G be a group with identity element e , and let $a \in G$. Then the group $\langle a \rangle$ is cyclic.

Proof available at the end of this section.

Definition: Order

Let G be a group and let $a \in G$. We define the **order** of a , denoted $o(a)$, to be $|\langle a \rangle|$. (Note: If there exists a positive integer k such that $a^k = e$, then the least such integer is the order of a ; otherwise, $o(a) = \infty$.)

Remark

Do not confuse the order of a group with the order of an *element of a group*. These are related concepts, but they are distinct, and have distinct notations: as we've seen, the order of a group, G , is denoted by $|G|$, while the order of an element a of a group is denoted by $o(a)$.

Theorem 5.1.4

Let element a in group G have $o(a) = k < \infty$. Then for $m \in \mathbb{Z}^+$, $a^m = e$ if and only if k divides m .

Proof available at the end of this section.

We have the following handy theorem.

Theorem 5.1.5

Let $a \in G$. Then $o(a) = o(a^{-1})$.

Proof available at the end of this section.

Unfortunately, there's no formula one can simply use to compute the order of an element in an arbitrary group. However, in the special case that the group is cyclic of order n , we do have such a formula. We present the following result without proof.

Theorem 5.1.6

For each $a \in \mathbb{Z}_n$, $o(a) = n / \gcd(n, a)$.

Theorem 5.1.7

Let $m, n \in \mathbb{Z}^+$ and let $(a, b) \in \mathbb{Z}_m \times \mathbb{Z}_n$. Let c, d be the respective orders of a in $\mathbb{Z}_m$ and b in $\mathbb{Z}_n$. Then $o((a, b)) = \text{lcm}(c, d)$.

Proof available at the end of this section.

Corollary 5.1.1

Suppose $m, n \in \mathbb{Z}^+$ with $\gcd(m, n) = 1$. Then $\mathbb{Z}_m \times \mathbb{Z}_n$ is cyclic.

Proof available at the end of this section.

Here are some examples of cyclic subgroups of groups, and orders of group elements.

Example 5.1.2

1. In $\mathbb{Z}$, $\langle 2 \rangle = \{\dots, -4, -2, 0, 2, 4, \dots\} = 2\mathbb{Z}$. More generally, given any $n \in \mathbb{Z}$, in $\mathbb{Z}$ we have $\langle n \rangle = n\mathbb{Z}$. For $a \in \mathbb{Z}$, $o(a) = \infty$ if $a \neq 0$; $o(0) = 1$.
2. In $\mathbb{Z}_8$, we have $\langle 0 \rangle = \{0\}$, $\langle 1 \rangle = \langle 3 \rangle = \langle 5 \rangle = \langle 7 \rangle = \mathbb{Z}_8$, $\langle 2 \rangle = \langle 6 \rangle = \{0, 2, 4, 6\}$, and $\langle 4 \rangle = \{0, 4\}$.
3. In $\mathbb{R}$, $\langle \pi \rangle = \pi\mathbb{Z}$, so $o(\pi) = \infty$.
4. In $\mathbb{R}^*$, $\langle \pi \rangle = \{\pi^n : n \in \mathbb{Z}\}$. Again, $o(\pi) = \infty$.
5. In $M_2(\mathbb{R})$,

$$\left\langle \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \right\rangle = \left\{ \begin{bmatrix} c & c \\ 0 & c \end{bmatrix} : c \in \mathbb{Z} \right\} = \left\{ c \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} : c \in \mathbb{Z} \right\}.$$

The matrix therefore has infinite order.

6. In $M_2(\mathbb{Z}_2)$, if $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, then $\langle A \rangle = \left\{ A, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \right\}$, so A has order 2.
7. In $GL(2, \mathbb{Z}_2)$, $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ has order 2. (Why?)
8. In $GL(2, \mathbb{R})$, $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ has order 4. (Why?)
9. In $GL(2, \mathbb{R})$, $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ has infinite order. (Why?)
10. In $GL(4, \mathbb{Z}_2)$, $A = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$ has order 2. (Why?)

Theorem 5.1.8

Every cyclic group G is of the form $\langle a \rangle$ for some $a \in G$.

Proof available at the end of this section.

Definition: Generator of G

Let G be a group. An element $a \in G$ is a **generator of G** (equivalently, a **generates G**) if $\langle a \rangle = G$.

Remark

1. Note that if G has a generator, then it is necessarily a cyclic group.
2. Note that an element a of a group G generates G if and only if every element of G is of the form a^n for some $n \in \mathbb{Z}$.
3. Generators of groups *need not* be unique. For instance, we saw in Example 5.1.2 that each of the elements 1, 3, 5 and 7 of $\mathbb{Z}_8$ is a generator for $\mathbb{Z}_8$.

Example 5.1.3

1. $\mathbb{Z}$ has generator 1 and generator -1 .
2. Given $n \in \mathbb{Z}$, $n\mathbb{Z}$ has generator n and generator $-n$.
3. Given $n \geq 2$ in $\mathbb{Z}$, the generators of $\mathbb{Z}_n$ are exactly the elements $a \in \mathbb{Z}_n$ such that $\gcd(n, a) = 1$. (This follows from Theorem 5.1.6.)

Remark

Order of elements provides another group invariant.

Theorem 5.1.9

Let $\phi : G \rightarrow G'$ be a group isomorphism and let $a \in G$. Then $o(\phi(a)) = o(a)$.

Proof available at the end of this section.

Corollary 5.1.2

If groups G and G' are isomorphic then for any $n \in \mathbb{Z}^+$, the number of elements of G of order n is the same as the number of elements of G' of order n .

Example 5.1.4

Since 1 in $\mathbb{Z}_4$ has order 4 but every element in $\mathbb{Z}_2 \times \mathbb{Z}_2$ has order less than or equal to 2, these groups cannot be isomorphic.

Note

Note, however, that just because the orders of elements of two groups “match up” the groups need not be isomorphic. For example, every element of $\mathbb{Z}$ has infinite order, except for its identity element, which has order 1; the same is true for the group $\mathbb{Q}$. However, we have previously proven that these groups are not isomorphic.

Answers

Theorem 5.1.1

Since $\mathbb{Z}$ and each $\mathbb{Z}_n$ are abelian, every cyclic group is abelian. But, for instance, $\mathbb{R}$ is an abelian noncyclic group (it can't be cyclic because it is uncountable).

Theorem 5.1.2

Let $x, y \in \langle a \rangle$. Then $x = a^i, y = a^j$ for some $i, j \in \mathbb{Z}$. So $xy = a^i a^j = a^{i+j} \in \langle a \rangle$. Next, $e_G = a^0 \in \langle a \rangle$. Finally, $(a^i)^{-1} = a^{-i} \in \langle a \rangle$. So $\langle a \rangle$ is a subgroup of G .

Lemma 5.1.1: Clock Lemma

Let $s, t \in \mathbb{Z}$. By the Division Algorithm, there exist $q \in \mathbb{Z}$ and $r \in \mathbb{Z}_n$ with $s - t = qn + r$. Then $a^{s-t} = a^{qn+r} = (a^n)^q a^r = e^q a^r = a^r$. Since $\forall 0 \leq r < n$

The last statement follows from the fact that n divides any positive integer to which it is congruent modulo itself.

Theorem 5.1.3

Case 1. There is no positive integer k with $a^k = e$. In this case, we claim $\langle a \rangle \simeq \mathbb{Z}$. Indeed: Let $\phi : \mathbb{Z} \rightarrow \langle a \rangle$ be defined by $\phi(i) = a^i$. Clearly, ϕ is a homomorphism that is onto. So to show that ϕ is an isomorphism, it suffices to show that ϕ is one-to-one. Let $i, j \in \mathbb{Z}$ with $\phi(i) = \phi(j)$. Then $a^i = a^j$. Without loss of generality, we may assume $i \geq j$. Then, multiplying both sides of the equation by a^{-j} (on the right or on the left) we obtain $a^{i-j} = e$. Since $i - j \in \mathbb{N}$ and there is no positive integer k with $a^k = e$, we must have $i - j = 0$, so $i = j$. Thus, ϕ is an isomorphism from $\mathbb{Z}$ to $\langle a \rangle$.

Case 2. There is a positive integer k such that $a^k = e$. In this case, let n be the least such positive integer. Define $\phi : \mathbb{Z}_n \rightarrow \langle a \rangle$ be defined by $\phi(i) = a^i$. We claim that ϕ is an isomorphism.

First, let $i, j \in \mathbb{Z}$. We want to show that $\phi(i +_n j) = \phi(i)\phi(j)$, that is, that $a^{i+_n j} = a^i a^j$. Since $i + j$ is congruent to $i +_n j$ modulo n , $a^{i+j} = a^{i+_n j} = a^{i+_n j}$, by the Clock Lemma. So ϕ is a homomorphism. Next, we show that ϕ is one-to-one. Let $i, j \in \mathbb{Z}_n$ with $\phi(i) = \phi(j)$, so $a^i = a^j$. Then, by the Clock Lemma, i and j are congruent modulo n . Since they're both in $\mathbb{Z}_n$, they must be equal, so ϕ is one-to-one. Finally, we show that ϕ is onto. Let $x \in \langle a \rangle$. Then $x = a^i$ for some $i \in \mathbb{Z}$. Letting r be the remainder when we divide i by n , we have $r \in \mathbb{Z}_n$ with i congruent to r modulo n ; so, again using the Clock Lemma, $x = a^i = a^r$. Since $r \in \mathbb{Z}_n$, $x = \phi(r)$. So ϕ is onto.

Thus, ϕ is an isomorphism, and so in this case we have $\mathbb{Z}_n \simeq \langle a \rangle$.

Theorem 5.1.4

This follows from the last statement of the Clock Lemma.

Theorem 5.1.5

First, assume $o(a) = n < \infty$. Then

$$(a^{-1})^n = (a^n)^{-1} = e^{-1} = e,$$

so $o(a^{-1}) \leq n = o(a)$. Using the same argument, we have $n = o(a) \leq o(a^{-1})$. Since $o(a^{-1}) \leq n$ and $n \leq o(a^{-1})$, $o(a^{-1}) = n = o(a)$.

On the other hand, assume $o(a) = \infty$. Then a^{-1} must also have infinite order, since if it had finite order m , a would, by the above argument, have order less than or equal to m .

Theorem 5.1.7

Let $k \in \mathbb{Z}^+$. By definition of c and d and by Theorem 5.1.4, we have $k(a, b) = (ka, kb) = (0, 0)$ if and only if c and d divide k . Since $\gcd(c, d) = 1$, the least positive integer divisible by both c and d is $\text{lcm}(c, d)$. Thus, $o((a, b)) = \text{lcm}(c, d)$, as desired.

Corollary 5.1.1

Since $\gcd(m, n) = 1$, $\text{lcm}(m, n) = mn$. So

$$o((1, 1)) = \text{lcm}(m, n) = mn = |\mathbb{Z}_m \times \mathbb{Z}_n|,$$

and we see that $\langle (1, 1) \rangle$ generates $\mathbb{Z}_m \times \mathbb{Z}_n$.

Theorem 5.1.8

Let G be cyclic. Suppose $|G| = \infty$. Then there is an isomorphism $\phi : \mathbb{Z} \rightarrow G$. Note that $\mathbb{Z} = \langle 1 \rangle$. So

Illegal pream-token (&)

A similar argument shows that there exists $a \in G$ such that $G = \langle a \rangle$ when $|G| < \infty$.

Theorem 5.1.9

This follows from the fact that $\phi(\langle a \rangle) = \langle \phi(a) \rangle$.

5.2: The Subgroup Lattices of Cyclic Groups

We now explore the subgroups of cyclic groups. A complete proof of the following theorem is provided on p. 61 of [1].

Theorem 5.2.1

Every subgroup of a cyclic group is cyclic.

Sketch of proof: Let $G = \langle a \rangle$ and $H \leq G$. If $H = \{e\}$, then clearly H is cyclic. Else, there exists an element a^i in H with $i > 0$; let d be the least positive integer such that $a^d \in H$. It turns out that $H = \langle a^d \rangle$.

Corollary 5.2.1

Every subgroup of $\mathbb{Z}$ is of the form $n\mathbb{Z}$ for $n \in \mathbb{Z}$. (Note that $n\mathbb{Z} \simeq \mathbb{Z}$ unless $n = 0$.)

Really, it suffices to study the subgroups of $\mathbb{Z}$ and $\mathbb{Z}_n$ to understand the subgroup lattice of every cyclic group.

We provide the following theorems without proof.

Theorem 5.2.2

The nontrivial subgroups of $\mathbb{Z}_n$ are exactly those of the form $\langle d \rangle$, where d is a positive divisor of n . Note that

$$|\langle d \rangle| = n/d$$

for each such d .

Theorem 5.2.3

For each $0 \neq a \in \mathbb{Z}_n$,

$$\langle a \rangle = \left\langle \frac{n}{\gcd(n, a)} \right\rangle.$$

It follows that for each $0 \neq a \in \mathbb{Z}_n$,

$$|\langle a \rangle| = \frac{n}{\gcd(n, a)}.$$

In fact:

Theorem 5.2.4

$\mathbb{Z}_n$ has a unique subgroup of order k , for each positive divisor k of n .

Example 5.2.1

How many subgroups does $\mathbb{Z}_{18}$ have? What are the generators of $\mathbb{Z}_{15}$?

Well, the positive divisors of 18 are 1, 2, 3, 6, 9, and 18, so $\mathbb{Z}_{18}$ has exactly six subgroups (namely, $\langle 1 \rangle$, $\langle 2 \rangle$, etc.). The generators of $\mathbb{Z}_{15}$ are the elements of $\mathbb{Z}_{15}$ that are relatively prime to 15, namely 1, 2, 4, 7, 8, 11, 13, and 14.

Example 5.2.2

Draw a subgroup lattice for $\mathbb{Z}_{12}$.

The positive divisors of 12 are 1, 2, 3, 4, 6, and 12; so $\mathbb{Z}_{12}$'s subgroups are of the form $\langle 1 \rangle$, $\langle 2 \rangle$, etc. So $\mathbb{Z}_{12}$ has the following subgroup lattice.

5.3: Exercises

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G be a group with identity element e .

- If G is infinite and cyclic, then G must have infinitely many generators.
- There may be two distinct elements a and b of a group G with $\langle a \rangle = \langle b \rangle$.
- If $a, b \in G$ and $a \in \langle b \rangle$ then we must have $b \in \langle a \rangle$.
- If $a \in G$ with $a^4 = e$, then $o(a)$ must equal 4.
- If G is countable then G must be cyclic.

2. Give examples of the following.

- An infinite noncyclic group G containing an infinite cyclic subgroup H .
- An infinite noncyclic group G containing a finite nontrivial cyclic subgroup H .
- A cyclic group G containing exactly 20 elements.
- A nontrivial cyclic group G whose elements are all matrices.
- A noncyclic group G such that every proper subgroup of G is cyclic.

3. Find the orders of the following elements in the given groups.

- $2 \in \mathbb{Z}$
- $-i \in \mathbb{C}^*$
- $-I_2 \in GL(2, \mathbb{R})$
- $-I_2 \in M_2(\mathbb{R})$
- $(6, 8) \in \mathbb{Z}_{10} \times \mathbb{Z}_{10}$

4. For each of the following, if the group is cyclic, list *all* of its generators. If the group is *not* cyclic, write NC.

- $5\mathbb{Z}$
- $\mathbb{Z}_{18}$
- $\mathbb{R}$
- $\langle \pi \rangle$ in $\mathbb{R}$
- $\mathbb{Z}_2^2$
- $\langle 8 \rangle$ in $\mathbb{Q}^*$

5. Explicitly identify the elements of the following subgroups of the given groups. You may use set-builder notation if the subgroup is infinite, or a conventional name for the subgroup if we have one.

- $\langle 3 \rangle$ in $\mathbb{Z}$
- $\langle i \rangle$ in $\mathbb{C}^*$
- $\langle A \rangle$, for $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \in M_2(\mathbb{R})$
- $\langle (2, 3) \rangle$ in $\mathbb{Z}_4 \times \mathbb{Z}_5$
- $\langle B \rangle$, for $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \in GL(2, \mathbb{R})$

6. Draw subgroup lattices for the following groups

- $\mathbb{Z}_6$
- $\mathbb{Z}_{13}$
- $\mathbb{Z}_{18}$

7. Let G be a group with no nontrivial proper subgroups. Prove that G is cyclic.

This page titled [5.3: Exercises](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

6: PERMUTATION AND DIHEDRAL GROUPS

We have already been introduced to two important classes of nonabelian groups: namely, the matrix groups $GL(n, \mathbb{R})$ and $SL(n, \mathbb{R})$ for $n \geq 2$. We now consider a more general class of (mostly) nonabelian groups: permutation groups.

6.1: Introduction to Permutation Groups

6.2: Symmetric Groups

6.3: Alternating Groups

6.4: Cayley's Theorem

6.5: Dihedral Groups

6.6: Exercises

6.1: Introduction to Permutation Groups

Definition: Permutation

A **permutation** on a set A is a bijection from A to A . We say a permutation σ on A **fixes** $a^{\text{int}} A$ if $\sigma(a) = a$.

Example 6.1.1

Let A be the set $A = \{\Delta, \star, 4\}$. Then the functions $\sigma : A \rightarrow A$ defined by

$$\sigma(\Delta) = \star, \sigma(\star) = \Delta, \text{ and } \sigma(4) = 4;$$

and $\tau : A \rightarrow A$ defined by

$$\tau(\Delta) = 4, \tau(\star) = \Delta, \text{ and } \tau(4) = \star$$

are both permutations on A .

Definition: Permutation Multiplication

Composition of permutations on a set A is often called **permutation multiplication**, and if σ and τ are permutations on a set A , we usually omit the composition symbol and write $\sigma \circ \tau$ simply as $\sigma\tau$.

Note

For us, if σ and τ are permutations on a set A , then applying $\sigma\tau$ to A means first applying τ and *then* applying σ . This is due to the conventional right-to-left reading of function compositions.

That is, if $a^{\text{int}} A$, by $\sigma\tau(a)$ we mean $\sigma(\tau(a))$. (Some other books/mathematicians do not use this convention, and read permutation multiplication from left-to-right. Make sure to always know what convention your particular author or colleague is using!)

Example 6.1.2

Let A , σ , and τ be as in Example 6.1.1. Then $\sigma\tau$ is the function from A to A defined by

$$\sigma\tau(\Delta) = 4, \sigma\tau(\star) = \star, \text{ and } \sigma\tau(4) = \Delta,$$

while $\tau\sigma$ is the function from A to A defined by

$$\tau\sigma(\Delta) = \Delta, \tau\sigma(\star) = 4, \text{ and } \tau\sigma(4) = \star$$

Definition

Given a set A , we define S_A to be the set of all permutations on A .

Theorem 6.1.1

Given a set A :

1. S_A is a group under permutation multiplication.
2. If A has finite cardinality n , then $|S_A| = n!$ (if $|A| = \infty$ then $|S_A|$ is also ∞).
3. S_A is abelian if $|A| = 1$ or 2 , and nonabelian otherwise.

Proof available at the end of this section.

We will in the future use language provided by the following definition:

Definition: Permutation Group

A group is said to be a **permutation group** if it is a subgroup of S_A for some set A .

Remark

Notice that if A and B are sets, then $|A| = |B|$ if and only if $S_A \simeq S_B$.

Thus, for any set B with $|B| = n^{\text{int}}\mathbb{Z}^+$, we have $S_B \simeq S_A$, where $A = \{1, 2, \dots, n\}$. Since we are concerned in this course primarily with group structures which are invariant under isomorphism, we may focus now on groups of permutations on the set $\{1, 2, \dots, n\}$ ($n^{\text{int}}\mathbb{Z}^+$).

Answers

Theorem 6.1.1

1. Let $\sigma, \tau \in S_A$. Since a composition of bijections is a bijection (see Theorem 1.2.3), $\sigma\tau$ is a bijection from A to A , hence is in S_A . So $\langle S_A, \circ \rangle$ is a binary structure. Next, function composition is always associative.

The identity function $1_A : A \rightarrow A$ defined by

$$1_A(a) = a \text{ for all } a \in A$$

clearly acts as an identity element in S_A . Henceforth, we will denote 1_A by e .

Finally, let $\sigma \in S_A$. Since σ is a bijection, σ has an inverse function σ^{-1} that is also a bijection from A to A (Theorem 1.2.1). Since $\sigma^{-1} \in S_A$ with $\sigma\sigma^{-1} = \sigma^{-1}\sigma = 1_A$, every element of S_A has an inverse element in S_A .

So S_A is a group.

2. Clearly $|S_A| = \infty$ when $|A| = \infty$, and a straightforward combinatorial argument yields that when $|A| = n \in \mathbb{Z}^+$, we have $|S_A| = n!$.
3. Finally, if $|A| = 1$ or 2 , then $|S_A| = 1! = 1$ or $|S_A| = 2! = 2$ so S_A must be abelian (as it's a group of order 1 or 2). On the other hand, suppose $|A| > 2$. Then A contains at least three distinct elements, say x, y , and z . Let σ be the permutation of A swapping x and y and fixing every other element of A , and let τ be the permutation of A swapping y and z and fixing every other element of A . Then $\sigma\tau(x) = y$ while $\tau\sigma(x) = z$, so $\sigma\tau \neq \tau\sigma$, and hence S_A is nonabelian.

6.2: Symmetric Groups

Definition: Symmetric Group

When $A = \{1, 2, \dots, n\}$ ($n \in \mathbb{Z}^+$), we call S_A the **symmetric group** on n letters and denote it by S_n .

(We can, in fact, define S_0 , the set of all permutations on the empty set. One can show, using the fact that a function is a relation on a Cartesian product of sets, that S_0 is the trivial group. However, this will not be relevant in this text.)

Remark

Throughout this course, if we are discussing a group S_n , you should assume $n \in \mathbb{Z}^+$.

It is important for us to be able to easily describe specific elements of S_n . It would be cumbersome to describe, for instance, an element of S_3 by saying it swaps 1 and 2 and fixes 3; imagine how much more cumbersome it could be to describe an element of, say, S_{100} ! One can somewhat concisely describe a permutation σ of S_n by listing out the elements $1, 2, \dots, n$ and writing the element $\sigma(i)$ below each i for $i = 1, 2, \dots, n$. For instance, if σ sends 1 to 2, we'd write the number 2 below the number 1. The convention is to enclose these two rows of numbers in a single set of parentheses, as in the following example.

Example 6.2.1

We can denote the element σ of S_3 that swaps 1 and 2 and fixes 3 by

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix},$$

and the element τ of S_3 that sends 1 to 3, 2 to 1, and 3 to 2 by

$$\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix}.$$

Then

$$\sigma\tau = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

and

$$\tau\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix}.$$

But even this notation is unnecessarily cumbersome. Instead, we use **cycle notation**.

Definition: Cycle, Cycle Notation, and Transposition

A permutation σ in S_n is called a **k -cycle** or a **cycle of length k** (or, less specifically, a **cycle**) if there exist distinct elements $a_1, a_2, \dots, a_k$ in $\{1, 2, \dots, n\}$ such that

- $\sigma(a_i) = a_{i+1}$ for each $i = 1, 2, \dots, k-1$;
- $\sigma(a_k) = a_1$;
- $\sigma(x) = x$ for every other element of $\{1, 2, \dots, n\}$.

We use the **cycle notation** $\sigma = (a_1 a_2 \cdots a_k)$ to describe such a cycle. A 2-cycle is often called a **transposition**.

Example 6.2.2

The permutation τ in S_3 that sends 1 to 3, 2 to 1, and 3 to 2 is a cycle. It can be denoted by $\tau = (132)$. Similarly, the cycle ρ in S_3 swapping 1 and 3 can be denoted by $\rho = (13)$. On the other hand, the permutation in S_4 that swaps 1 with 2 and 3 with 4 is *not* a cycle.

Remark

Given a k -cycle $\sigma = (a_1 a_2 \cdots a_k)$, there are k different expressions for σ . Indeed, we have

$$\sigma = (a_1 a_2 \cdots a_k) = (a_2 a_3 \cdots a_k a_1) = (a_3 a_4 \cdots a_k a_1 a_2) = \cdots = (a_k a_1 \cdots a_{k-1}).$$

Example 6.2.3

The permutation τ described in Example 6.2.2 can also be written as (321) and as (213) .

However, by convention, we usually “start” a cycle σ with the smallest of the numbers that σ doesn't fix: e.g., we'd write $\sigma = (213)$ as (132) .

Definition: Disjoint and Mutual Disjointness

Two cycles $\sigma = (a_1 a_2 \cdots a_k)$ and $\tau = (b_1 b_2 \cdots b_m)$ are said to be **disjoint** if $a_i \neq b_j$ for all i and j . Cycles $\sigma_1, \sigma_2, \dots, \sigma_m$ are **disjoint** if σ_i and σ_j are disjoint for each $i \neq j$. (Notice: this version of disjointness is what we usually refer to as **mutual disjointness**.)

Remark

Note that if cycles σ and τ are disjoint, then σ and τ commute; that is, $\sigma\tau = \tau\sigma$.

Note

If cycles σ and τ are *not* disjoint then they may not commute. For instance, see Example 6.2.3, where $\sigma\tau \neq \tau\sigma$.

Note that any permutation of S_n is a product of disjoint cycles (where by “product” we mean the permutation resulting from permutation multiplication).

Definition: (Disjoint) Cycle Notation

Writing a permutation in **(disjoint) cycle notation** means writing it as a product of disjoint cycles, where each cycle is written in cycle notation.

Remark

Note that if σ in S_n is written in cycle notation and the number $a^{\text{int}}\{1, 2, \dots, n\}$ appears nowhere in σ 's representation, this means that σ fixes a . The only permutation that we cannot really write in cycle notation is the identity element of S_n , which we henceforth denote by e .

Example 6.2.4

The permutation

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 2 & 6 & 5 & 4 \end{pmatrix}$$

is the product of disjoint cycles (132) and (46) , so in cycle notation we have

$$\sigma = (132)(46).$$

Note that we could also write σ as $(321)(46)$, $(213)(64)$, $(64)(132)$, etc.

While it is true that we also have $\sigma = (13)(23)(46)$, this is not a disjoint cycle representation of σ since both (13) and (23) “move” the element 3.

Example 6.2.5

In S_4 , let $\sigma = (243)$ and $\tau = (13)(24)$. Then $\sigma\tau = (123)$ and $\tau\sigma = (134)$.

Example 6.2.6

In S_9 , let $\sigma = (134)$, $\tau = (26)(17)$, and $\rho = (358)(12)$. Find the following, writing your answers using disjoint cycle notation.

1. σ^{-1}
2. $\sigma^{-1}\tau\sigma$
3. σ^2
4. σ^3
5. ρ^2
6. ρ^{-2}
7. $\sigma\tau$
8. $\sigma\rho$

Example 6.2.7

Explicitly express all the elements of S_4 in disjoint cycle notation.

Theorem 6.2.1

Any k -cycle has order k in S_n . More generally, if permutation σ can be written in disjoint cycle notation as $\sigma = \sigma_1\sigma_2\cdots\sigma_m$, then

$$\begin{aligned} o(\sigma) &= \text{lcm}(o(\sigma_1), o(\sigma_2), \dots, o(\sigma_m)) \\ &= \text{lcm}(\text{length}(\sigma_1), \text{length}(\sigma_2), \dots, \text{length}(\sigma_m)), \end{aligned}$$

where lcm denotes the least common multiple.

Note

Permutation σ must be in *disjoint* cycle notation for the above formula to hold. For instance, let $\sigma = (12)(23)$ in S_3 . The transpositions (12) and (23) both have order 2, but $o(\sigma) \neq \text{lcm}(2, 2) = 2$. Rather, $o(\sigma) = 3$, since in disjoint cycle notation σ can be written as $\sigma = (123)$. You must write a permutation using disjoint cycle notation before attempting to use this method to compute its order!

Example 6.2.8

1. Find the orders of each of the elements in Example 6.2.6, including σ , τ , and ρ themselves.
2. Explicitly list the elements of $\langle \sigma \rangle$, $\langle \tau \rangle$, and $\langle \rho \rangle$.

6.3: Alternating Groups

Note that every k -cycle $(a_1 a_2 \dots a_k)^{\text{int}} S_n$ can be written as a product of (not necessarily disjoint) transpositions:

$$(a_1 a_2 \dots a_k) = (a_1 a_k)(a_1 a_{k-1}) \cdots (a_1 a_3)(a_1 a_2).$$

We, therefore, have the following theorem.

Theorem 6.3.1

Every permutation in S_n can be written as a product of transpositions.

Definition: Even and Odd

We say that a permutation in S_n is **even** [resp., **odd**] if it can be written as a product of an even [resp., odd] number of transpositions.

Theorem 6.3.2

Every permutation in S_n is even or odd, but not both.

Proof available at the end of this section.

Lemma 6.3.1

For each $2 \leq k \leq n$, then a k -cycle is even if k is odd, and odd if k is even.

The proof of this is left as an exercise for the reader.

Example 6.3.1

In S_3 , the permutations e , $(123) = (13)(12)$, and $(132) = (12)(13)$ are even, while the permutations (12) , (13) , and (23) are odd.

Example 6.3.2

List all of the even [resp., odd] permutations in S_4 .

We have the following theorem, whose proof is left as an exercise for the reader.

Theorem 6.3.3

The set of all even permutations in S_n is a subgroup of S_n .

Definition: Alternating Group

The **alternating group** on n letters is the subgroup A_n of S_n consisting of all of the even permutations in S_n .

We end with this theorem, whose proof can be found on p. 93 of [1].

Theorem 6.3.4

$$|A_n| = \frac{(n!)}{2}.$$

Answers

Theorem 6.3.2

We already know that every permutation in S_n is a product of transpositions, so must be even or odd. For proof that no permutation is both even *and* odd, see, for instance, Proof 1 or 2 of Theorem 9.15 on p. 91 in [1].

6.4: Cayley's Theorem

One might wonder how “common” permutation groups are in math. They are, it turns out, ubiquitous in abstract algebra: in fact, *every group* can be thought of as a group of permutations! We will prove this, but we first need the following lemma. (We will not use the maps ρ_a or c_a , defined below, in our theorem, but define them here for potential future use.)

Lemma 6.4.1

Let G be a group and $a \in G$. Then the following functions are permutations on G , and hence are elements of S_G :

- $\lambda_a : G \rightarrow G$ defined by $\lambda_a(x) = ax$;
- $\rho_a : G \rightarrow G$ defined by $\rho_a(x) = xa$;
- $c_a : G \rightarrow G$ defined by $c_a(x) = axa^{-1}$.

Proof available at the end of this section.

Definition: Left Multiplication, Right Multiplication, and Conjugation

We say that λ_a , ρ_a , and c_a perform on G , respectively, **left multiplication** by a , **right multiplication** by a , and **conjugation** by a . (Note: Sometimes when people talk about conjugation by a they instead are referring to the permutation of G that sends each x to $a^{-1}xa$.)

Now we are ready for our theorem:

Theorem 6.4.1: Cayley's Theorem

Let G be a group. Then G is isomorphic to a subgroup of S_G . Thus, every group can be thought of as a group of permutations.

Proof available at the end of this section.

Remark

In general, $\phi(G) \neq S_G$, so we cannot conclude that G is isomorphic to S_G itself; rather, we may only conclude that it is isomorphic to some *subgroup* of S_G .

Remark

While we chose to use the maps λ_a to prove the above theorem, we could just as well have used the maps ρ_a or c_a , instead.

Answers

Lemma 6.4.1

To show that λ_a is a bijection, first assume $x_1, x_2 \in G$ with $\lambda_a(x_1) = \lambda_a(x_2)$. Then $ax_1 = ax_2$; so, by left cancellation, $x_1 = x_2$. Thus, λ_a is one-to-one. Further, each $y \in G$ equals $\lambda_a(a^{-1}y)$ for $a^{-1}y \in G$, so λ_a is onto. Thus, λ_a is a bijection from G to G ; that is, it's a permutation on G . The proofs that ρ_a and c_a are bijections are similar.

Theorem 6.4.1: Cayley's Theorem

For each $a \in G$, let λ_a be defined, as above, by $\lambda_a(x) = ax$ for each $x \in G$; recall that each λ_a is in S_G . Now define $\phi : G \rightarrow S_G$ by $\phi(a) = \lambda_a$, for each $a \in G$.

We claim that ϕ is both a homomorphism and one-to-one. Indeed, let $a, b \in G$. Now, $\phi(a)\phi(b)$ and $\phi(ab)$ are both functions with domain G , so we need to show $(\phi(a)\phi(b))(x) = (\phi(ab))(x)$ for each $x \in G$. Well, let $x \in G$. Then

Illegal pream-token (&)

So ϕ is a homomorphism. Further, if $a, b \in G$ with $\phi(a) = \phi(b)$, then $\lambda_a = \lambda_b$. In particular, $\lambda_a(e) = \lambda_b(e)$. But $\lambda_a(e) = ae = a$ and $\lambda_b(e) = be = b$, so $a = b$. Thus, ϕ is one-to-one.


Since by definition $\phi(G)$ we have that ϕ maps G onto $\phi(G)$, we conclude that ϕ provides an isomorphism from G to the subgroup $\phi(G)$ of S_G .

6.5: Dihedral Groups

Dihedral groups are groups of symmetries of regular n -gons. We start with an example.

Example 6.5.1

Consider a regular triangle T , with vertices labeled 1, 2, and 3. We show T below, also using dotted lines to indicate a vertical line of symmetry of T and a rotation of T .



Note that if we reflect T over the vertical dotted line (indicated in the picture by f), T maps onto itself, with 1 mapping to 1, and 2 and 3 mapping to each other. Similarly, if we rotate T clockwise by 120° (indicated in the picture by r), T again maps onto itself, this time with 1 mapping to 2, 2 mapping to 3, and 3 mapping to 1. Both of these maps are called **symmetries** of T ; f is a **reflection** or **flip** and r is a **rotation**.

Of course, these are not the only symmetries of T . If we compose two symmetries of T , we obtain a symmetry of T : for instance, if we apply the map $f \circ r$ to T (meaning first do r , then do f) we obtain reflection over the line connecting 2 to the midpoint of line segment $\overline{13}$. Similarly, if we apply the map $f \circ (r \circ r)$ to T (first do r twice, then do f) we obtain reflection over the line connecting 3 to the midpoint of line segment $\overline{12}$. In fact, every symmetry of T can be obtained by composing applications of f and applications of r .

For convenience of notation, we omit the composition symbols, writing, for instance, fr for $f \circ r$, $r \circ r$ as r^2 , etc. It turns out there are exactly six symmetries of T , namely:

1. the map e from T to T sending every element to itself;
2. f (that is, reflection over the line connecting 1 and the midpoint of $\overline{23}$);
3. r (that is, clockwise rotation by 120°);
4. r^2 (that is, clockwise rotation by 240°);
5. fr (that is, reflection over the line connecting 2 and the midpoint of $\overline{13}$); and
6. fr^2 (that is, reflection over the line connecting 3 and the midpoint of $\overline{12}$).

Declaring that $f^0 = r^0 = e$, the set

$$D_3 = \{e, f, r, r^2, fr, fr^2\} = \{f^i r^j : i = 0, 1, j = 0, 1, 2\}$$

is the collection of all symmetries of T .

Remark

Notice that $rf = fr^2$ and that $f^2 = r^3 = e$.

Theorem 6.5.1

D_3 is a group under composition.

Proof available at the end of this section.

Let us look at D_3 another way. Note that each map in D_3 can be uniquely described by how it permutes the vertices 1, 2, 3 of T : that is, each map in D_3 can be uniquely identified with a unique element of S_3 . For instance, f corresponds to the permutation (23) in S_3 , while fr corresponds to the permutation (13). It turns out that $D_3 \simeq S_3$, via the following correspondence.

$$e \mapsto e$$

$$f \mapsto (23)$$

$$r \mapsto (123)$$

$$r^2 \mapsto (132)$$

$$fr \mapsto (13)$$

$$fr^2 \mapsto (12)$$

The group D_3 is an example of class of groups called *dihedral groups*.

Definition: Dihedral Group

Let n be an integer greater than or equal to 3. We let D_n be the collection of symmetries of the regular n -gon. It turns out that D_n is a group (see below), called the **dihedral group of order $2n$** . (Note: Some books and mathematicians instead denote the group of symmetries of the regular n -gon by D_{2n} —so, for instance, our D_3 , above, would instead be called D_6 . Make sure you are aware of the convention your book or colleague is using.)

Theorem 6.5.2

Let n be an integer greater than or equal to 3. Then, again using the convention that $f^0 = r^0 = e$, D_n can be uniquely described as

$$D_n = \{f^i r^j : i = 0, 1, j = 0, 1, \dots, n-1\}$$

with the relations

$$rf = fr^{n-1} \text{ and } f^2 = r^n = e.$$

The dihedral group D_n is a nonabelian group of order $2n$.

Proof available at the end of this section.

Remark

Throughout this course, if we are discussing a group D_n you should assume $n \in \mathbb{Z}^+$, $n \geq 3$, unless otherwise noted.

Definition: Standard Form

We say that an element of D_n is written in **standard form** if it is written in the form $f^i r^j$ where $i \in \{0, 1\}$ and $j \in \{0, 1, \dots, n-1\}$.

Theorem 6.5.3

Each D_n is isomorphic to a subgroup of S_n .

Proof available at the end of this section.

Note

While D_3 is actually isomorphic to S_3 itself, for $n > 3$ we have that D_n is *not* isomorphic to S_n but is rather isomorphic to a *proper subgroup* of S_n . When $n > 3$ you can see that D_n cannot be isomorphic to S_n since $|D_n| = 2n < n! = |S_n|$ for $n > 3$.

It is important to be able to do computations with specific elements of dihedral groups. We have the following theorem.

Theorem 6.5.4

The following relations hold in D_n , for every n :

1. For every i , $r^i f = f r^{-i}$ (in particular, $r f = f r^{-1} = f r^{n-1}$);
2. $o(f r^i) = 2$ for every i (in particular, $f^2 = e$);
3. $o(r) = o(r^{-1}) = n$;
4. If n is even, then $r^{n/2}$ commutes with every element of D_n .

Proof available at the end of this section.

Example 6.5.2

1. Write $f r^2 f$ in D_3 in standard form. Do the same for $f r^2 f$ in D_4 .
2. What is the inverse of $f r^3$ in D_5 ? Write it in standard form.
3. Explicitly describe an isomorphism from D_4 to a subgroup of S_4 .

Example 6.5.3

Classify the following groups up to isomorphism. (*Hint*: You may want to look at the number of group elements that have a specific finite order.)

$$\mathbb{Z}, \mathbb{Z}_6, \mathbb{Z}_2, S_6, \mathbb{Z}_4, \mathbb{Q}, 3\mathbb{Z}, \mathbb{R}, S_2, \mathbb{R}^*, S_3, \mathbb{Q}^*, \mathbb{C}^*, \langle \pi \rangle \text{ in } \mathbb{R}^*,$$

$$D_6, \langle (134)(25) \rangle \text{ in } S_5, \mathbb{R}^+, D_3, \langle r \rangle \text{ in } D_4, 17\mathbb{Z}$$

Answers

Theorem 6.5.1

First, as noted above, $rf = fr^2$. So any map of the form $f^i r^j f^k r^l$ ($k = 0, 1, \quad j, l = 0, 1, 2$) can be written in the form $f^s r^t$ for some $s, t \in N$. Finally, let $R_2(s)$ and $R_3(t)$ be the remainders when you divide s by 2 and t by 3; then $f^s r^t = f^{R_2(s)} r^{R_3(t)} \in D_3$. So $\langle D_3, \circ \rangle$ is a binary structure.

Next, function composition is always associative, and the function ee clearly acts as identity element in D_3 . Finally, let $x = f^i r^j \in D_3$. Then $y = r^{3-j} f^{2-i}$ is in D_3 with $xy = yx = e$. So D_3 is a group.

Theorem 6.5.2

The proof that D_n is a group parallels the proof, above, that D_3 is a group. It is clear that D_n is nonabelian (e.g., $rf = fr^{n-1} \neq fr$) and has order $2n$.

Theorem 6.5.3

We provide here a sketch of a proof; the details are left as an exercise for the reader. We described above how D_3 is isomorphic to a subgroup (namely, the improper subgroup) of S_3 . One can show that each D_n is isomorphic to a subgroup of S_n by similarly labeling the vertices of the regular n -gon $1, 2, \dots, n$ and determining how these vertices are permuted by each element of D_n .

Theorem 6.5.4

1. We use induction on the exponent of r . We already know that $r^1 f = fr^{-1}$. Now suppose $r^{i-1} f = fr^{-(i-1)}$ for some $i \geq 2$. Then

$$r^i f = r(r^{i-1} f) = r(fr^{-(i-1)}) = (rf)r^{-i+1} = (fr^{-1})r^{-i+1} = fr^{-i}.$$

2. For every i , $fr^i \neq e$, but

$$(fr^i)^2 = (fr^i)(fr^i) = f(r^i f)r^i = f(fr^{-i})r^i = f^2 r^0 = e.$$

3. This follows from Theorem 5.1.5 and the fact that $o(r)=n, o(r)=n$.
4. The proof of this statement is left as an exercise for the reader.

6.6: Exercises

Throughout, write all permutations using *disjoint cycle notation*, and write all dihedral group elements in *standard form*.

1. Let $\sigma = (134)$, $\tau = (23)(145)$, $\rho = (56)(78)$, and $\alpha = (12)(145)$ in S_8 . Compute the following.
 - a. $\sigma\tau$
 - b. $\tau\sigma$
 - c. τ^2
 - d. τ^{-1}
 - e. $o(\tau)$
 - f. $o(\rho)$
 - g. $o(\alpha)$
 - h. $\langle \tau \rangle$
2. Prove Lemma 6.3.4.
3. Prove that A_n is a subgroup of S_n .
4. Prove or disprove: The set of all odd permutations in S_n is a subgroup of S_n .
5. Let n be an integer greater than 2. $m^{\text{int}}\{1, 2, \dots, n\}$, and let $H = \{\sigma^{\text{int}} S_n : \sigma(m) = m\}$ (in other words, H is the set of all permutations in S_n that fix m).
 - a. Prove that $H \leq S_n$.
 - b. Identify a familiar group to which H is isomorphic. (You do not need to show any work.)
6. Write $rf r^2 f r f r$ in D_5 in standard form.
7. Prove or disprove: $D_6 \simeq S_6$.
8. Which elements of D_4 (if any)
 - a. have order 2?
 - b. have order 3?
9. Let n be an even integer that's greater than or equal to 4. Prove that $r^{n/2}$ commutes with every element of D_n . (Do NOT simply refer to the last statement in Theorem 6.5.4; that is the statement you are proving here.)

7: THE WONDERFUL WORLD OF COSETS

We have already seen one way we can examine a complicated group G : namely, study its subgroups, whose group structures are in some cases much more directly understandable than the structure of G itself. But if H is a subgroup of a group G , if we only study H we lose all the information about G 's structure “outside” of H . We might hope that $G - H$ (that is, the set of elements of G that are not in H) is also a subgroup of G , but we immediately see that cannot be the case since the identity element of G must be in H , and $H \cap (G - H) = \emptyset$. Instead, let's ask how we can get at some understanding of G 's entire structure using a subgroup H ? It turns out we use what are called cosets of H ; but before we get to those, we need to cover some preliminary material.

[7.1: Partitions of and Equivalence Relations on Sets](#)

[7.2: Introduction to Cosets and Normal Subgroups](#)

[7.3: The Index of a Subgroup and Lagrange's Theorem](#)

[7.4: Exercises](#)

7.1: Partitions of and Equivalence Relations on Sets

Definition: Partition and Cells

Let S be a set. Then a collection of subsets $P = \{S_i\}_{i \in I}$ (where I is some index set) is a **partition** of S if each $S_i \neq \emptyset$ and each element of S is in exactly one S_i . In other words, $P = \{S_i\}_{i \in I}$ is a partition of S if and only if:

1. each $S_i \neq \emptyset$;
2. the S_i are mutually disjoint (that is, $S_i \cap S_j = \emptyset$ for $i \neq j \in I$); and
3. $S = \bigcup_{i \in I} S_i$.

The S_i are called the **cells** of the partition.

Example 7.1.1

1. The set $\{1, 2, 3\}$ has 5 partitions: namely,

$$\{\{1, 2, 3\}\}, \{\{1, 2\}, \{3\}\}, \{\{1, 3\}, \{2\}\}, \{\{2, 3\}, \{1\}\} \text{ and } \{\{1\}, \{2\}, \{3\}\}.$$

The first partition we've mentioned has one cell, the next three have two cells, and the last one has three cells.

2. The following is a 2-celled partition of $\mathbb{Z}$:

$$\{\{x \in \mathbb{Z} : x \text{ is even}\}, \{x \in \mathbb{Z} : x \text{ is odd}\}\}.$$

The number of partitions of a finite set of n elements gets large very quickly as n goes to infinity. Indeed, there are 52 partitions of a set containing just 5 elements! (The total number of partitions of an n -element set is the **Bell number**, B_n . There is no trivial way of computing B_n , in general, though the B_n do satisfy the relatively simple recurrence relation

$$B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k,$$

for each $n \geq 1$.) But our goal here is not to count the number of partitions of a given set, but rather to use particular partitions of a group G to help us study that group's structure. We turn now to our next definition.

Definition: Relation

Let S be a set. Then a **relation** on S is a subset R of $S \times S$.¹ If R is a relation on S and $x, y \in S$, then we say x is **related to** y , and write xRy , if $(x, y) \in R$; otherwise, we say that x is **not related to** y , and write $x \not R y$.

Remark

If there is a conventional notation used to denote a particular relation on a set, we will usually use that notation, rather than R , to denote the relation.

We are already familiar with some relations on $\mathbb{R}$. Common such relations are $=$, $\neq$, $<$, $\leq$, $>$, and $\geq$; they contain exactly the elements we'd expect them to contain.

Example 7.1.2

1. $<$ is a relation on $\mathbb{R}$ that contains, for instance, $(3, 4)$ but not $(3, 3)$ or $(4, 3)$; $\leq$ is a relation on $\mathbb{R}$ that contains $(3, 4)$ and $(3, 3)$ but not $(4, 3)$.
2. Given any $n \in \mathbb{Z}^+$, congruence modulo n , denoted $\equiv_n$, is a relation on $\mathbb{Z}$, where $\equiv_n$ is defined to be $\{(x, y) \in \mathbb{Z} \times \mathbb{Z} : n \text{ divides } x - y\}$.
3. We can define a relation R on C^1 (the set of all differentiable functions from $\mathbb{R}$ to $\mathbb{R}$ whose derivatives are continuous) by declaring that $(f, g) \in C^1 \times C^1$ is in R if and only if $f' = g'$.

Definition: Equivalence Relation

Let R be a relation on a set S . Then R is said to be:

- **reflexive** on S if xRx for every $x \in S$;
- **symmetric** on S if whenever $x, y \in S$ with xRy we also have yRx ; and
- **transitive** on S if whenever $x, y, z \in S$ with xRy and yRz we also have xRz .

A relation that is reflexive, symmetric, and transitive is called an **equivalence relation**.

If we know, or plan to prove, that a relation is an equivalence relation, by convention we may denote the relation by $\sim$, rather than by R .

Remark

You can think of an equivalence relation on a set S as being a “weak version” of equality on S . Indeed, $=$ is an equivalence relation on any set S , but it also has a very special property that most equivalence relations *don't* have: namely, no element of S is related to *any other element* of S under $=$.

Example 7.1.3

1. $<$ is *not* an equivalence relation on $\mathbb{R}$ because it is not reflexive: for instance, $3 \not< 3$. $\leq$ is also *not* an equivalence relation on $\mathbb{R}$, since even though it is reflexive, it's not symmetric: indeed, $3 \leq 4$ but $4 \not\leq 3$.
2. Given any $n \in \mathbb{Z}^+$, $\equiv_n$ is an equivalence relation on $\mathbb{Z}$. The proof of this is left as an exercise for the reader.
3. The relation R on C^1 discussed above (fRg iff $f' = g'$) is an equivalence relation on C^1 .
4. $\simeq$ is an equivalence relation on any set of groups. This follows from Theorem 3.3.1.
5. Define relation R on $\mathbb{Z}$ by xRy if and only if $xy \geq 0$. Is R an equivalence relation on $\mathbb{Z}$? Well, for every $x \in \mathbb{Z}$, $x^2 \geq 0$, so R is reflexive. Moreover, if $x, y \in \mathbb{Z}$ with $xy \geq 0$, then $yx \geq 0$; so R is symmetric. But R is *not* transitive: indeed, $3, 0, -4 \in \mathbb{Z}$ with $3(0) = 0 \geq 0$ and $0(-4) = 0 \geq 0$, so $3R0$ and $0R-4$; but $3(-4) = -12 \not\geq 0$. So R isn't transitive, and hence is *not* an equivalence relation.

Definition: Equivalence Class of x under $\sim$

Given an equivalence relation $\sim$ on a set S , for each $x \in S$ we define the **equivalence class of x under $\sim$** to be

$$[x] = \{y \in S : y \sim x\}.$$

These sets $[x]$ ($x \in S$) are called the **equivalence classes of S under $\sim$** .

Remark

Note that, by reflexivity of $\sim$, $x \in [x]$ for every $x \in S$.

We have now the following Very Important Lemma:

Lemma 7.1.1

Let $\sim$ be an equivalence relation on set S . Then for $x, y \in S$, the following are equivalent:

1. $y \in [x]$;
2. $[x] = [y]$;
3. $x \in [y]$.

Proof available at the end of this section.

Definition: Representative

In many cases there are many distinct elements $x, y \in S$ with $[x] = [y]$; thus, there are usually many different ways we could denote a particular equivalence class of S under $\sim$. Element $y \in [x]$ is called a **representative** of class X if $y \in X$.

Example 7.1.4

1. Consider the equivalence relation $\equiv_2$ on $\mathbb{Z}$. Under this relation, $[0] = \{0, \pm 2, \pm 4, \dots\}$ and $[1] = \{\dots, -3, -1, 1, 3, \dots\}$; in fact, if we let E be the set of all even integers and O the set of all odd integers, then for $x \in \mathbb{Z}$, $[x] = E$ if x is even and O if x is odd. Thus, the set of all equivalence classes of $\mathbb{Z}$ under $\equiv_2$ is the 2-element set $\{E, O\}$: every even integer is a representative of E , while every odd integer is a representative of O .
2. For $f \in C^1$, the equivalence class of f under R (defined above) is the set of all functions in C^1 of the form $g(x) = f(x) + c$, where $c \in \mathbb{R}$.
3. Let $A = \{a, b, c\}$, and define $\sim$ on the power set $P(A)$ of A by $X \sim Y$ if and only if $|X| = |Y|$. It is straightforward to show that $\sim$ is an equivalence relation on $P(A)$, under which $P(A)$ has exactly 4 distinct equivalence classes.

$$\begin{aligned} [\emptyset] &= \{\emptyset\}, \\ [\{a\}] &= [\{b\}] = [\{c\}] = \{\{a\}, \{b\}, \{c\}\}, \\ [\{a, b\}] &= [\{a, c\}] = [\{b, c\}] = \{\{a, b\}, \{a, c\}, \{b, c\}\}, \text{ and} \\ [A] &= \{A\}. \end{aligned}$$

Remark

Notice that the complete set $\{E, O\}$ of distinct equivalence classes of $\mathbb{Z}$ under $\equiv_2$ is a partition of $\mathbb{Z}$, and the complete set

$$\{[\emptyset], [\{a\}], [\{a, b\}], [A]\}$$

of distinct equivalence classes of $P(A)$ under $\sim$ is a partition of $P(A)$. This is not a coincidence! It turns out that equivalence relations and partitions go hand in hand.

Theorem 7.1.1

Let S be a set. Then:

1. If $\sim$ is an equivalence relation on S , then the set of all equivalence classes of S under $\sim$ is a partition of S ; and
2. If P is a partition of S , then the relation on S defined by $x \sim y$ if and only if x is in the same cell of P as y is an equivalence relation on S .

Notice that in each case, the cells of the partition are the equivalence classes of the set under the corresponding equivalence relation.

Proof available at the end of this section.

Example 7.1.5

1. For $n^{\text{int}}\mathbb{Z}^+$, the set of the equivalence classes of $\mathbb{Z}$ under $\equiv_n$ is the partition $\{[0], [1], \dots, [n-1]\}$ of $\mathbb{Z}$. (The partition $\{E, O\}$ of $\mathbb{Z}$ discussed above is this partition when $n = 2$.)

Notes

1. More generally, if S and T are sets, a **relation between S and T** is a subset of $S \times T$. We will not, however, in our class consider the cases in which S and T are different sets.

Answers

Lemma 7.1.1

To prove that the first and second statements are equivalent: Let $x, y \in S$. If $y \in [x]$, then $y \sim x$, so for every $z \in [y]$ (that is, for every z in S with $z \sim y$), we have, by transitivity, $z \sim x$, so $z \in [x]$. On the other hand, by the symmetry of $\sim$ we have $x \sim y$; so for every $z \in [x]$, we have, again by transitivity, that $z \in [y]$. Thus, $[x] = [y]$. Conversely, if $[x] = [y]$, then since $y \in [y] = [x]$.

The proof that the second and third statements are equivalent is nearly identical.

Theorem 7.1.1

To prove the first statement: Let $\sim$ be an equivalence relation on S . Clearly, the equivalence classes of S under $\sim$ are nonempty sets whose union is S . Thus, it suffices to show $X \cap Y = \emptyset$ for each pair of equivalence classes $X \neq Y$ of S under $\sim$. Let X, Y be equivalence classes of S under $\sim$ that are NOT disjoint. Then there exists an element $z \in X \cap Y$. Then by Lemma 7.1.1, $[z] = X$ and $[z] = Y$; so $X = Y$. Thus, if $X \neq Y$, then $X \cap Y = \emptyset$.

Finally, it is straightforward (almost silly!) to prove that $\sim$ is reflexive, symmetric, and transitive.

7.2: Introduction to Cosets and Normal Subgroups

Definition

Given a group G with subgroup H , we define $\sim_L$ on G by

$$a \sim_L b \text{ if and only if } a^{-1}b \in H$$

and $\sim_R$ on G by

Double exponent: use braces to clarify

Theorem 7.2.1

$\sim_L$ and $\sim_R$ are equivalence relations on G .

Proof available at the end of this section.

Remark

Of course, different subgroups H and K in a group G will give rise to different relations $\sim_L$ and $\sim_R$ on G ; that is, these relations are really defined with respect to a particular subgroup of G .

From now on, whenever we discuss $\sim_L$ or $\sim_R$ on a group, assume that it is with respect to a particular subgroup H of G .

Now, as equivalence relations on a group G , each of $\sim_L$ and $\sim_R$ gives rise to a partition of G . What are the cells of those partitions?

Definition: Left and Right Cosets

Given $a \in G$, we define

$$aH = \{ah : h \in H\}$$

and

$$Ha = \{ha : h \in H\}.$$

We call aH and Ha , respectively, the **left and right cosets of H containing a** .

If we know that G is abelian, with operation denoted by $+$, we may denote these left and right cosets by $a + H$ and $H + a$, respectively.

Note

In the following, we use the notation $\Leftrightarrow$ to denote the phrase “if and only if.”

Theorem 7.2.2

Let $a \in G$. Then under $\sim_L$, $[a] = aH$ while under $\sim_R$, $[a] = Ha$.

Proof available at the end of this section.

We next summarize some facts about the left and right cosets of a subgroup H of a group G :

Theorem 7.2.3

Let G be a group with $H \leq G$ and $a, b \in G$.

1. The left [right] cosets of H in G partition G .

$$2. \quad b^{-1}aH \Leftrightarrow aH = bH \Leftrightarrow a^{-1}bH$$

and

$$b^{-1}Ha \Leftrightarrow Ha = Hb \Leftrightarrow a^{-1}Hb.$$

In particular,

$$a^{-1}H \Leftrightarrow aH = H \Leftrightarrow Ha = H.$$

3. H is the only left or right coset of H that is a **subgroup** of G .

4. $|aH| = |H| = |Ha|$.

Proof available at the end of this section.

Remark

We can use Statements 2 and 3, above, to save some time when computing left and right cosets of a subgroup of a group.

Example 7.2.1

Find the left and right cosets of $H = \langle (12) \rangle$ in S_3 .

The left cosets are

$$\begin{aligned} eH &= H = (12)H, \\ (13)H &= \{(13), (123)\} = (123)H, \\ \text{and } (23)H &= \{(23), (132)\} = (132)H, \end{aligned}$$

and the right cosets are

$$\begin{aligned} He &= H = H(12), \\ H(13) &= \{(13), (132)\} = H(132), \\ \text{and } H(23) &= \{(23), (123)\} = H(123). \end{aligned}$$

Thus, $\sim_L$ partitions S_3 into $\{H, \{(13), (123)\}, \{(23), (132)\}\}$ and $\sim_R$ partitions S_3 into $\{H, \{(13), (132)\}, \{(23), (123)\}\}$.

Example 7.2.2

Find the left and right cosets of $H = \langle f \rangle$ in D_4 .

This example is left as an exercise for the reader.

We now draw attention to some very important facts:

Note

For $a, b \in G$:

1. In general, $aH \neq Ha$!
2. $aH = bH$ does not necessarily imply $a = b$ or that there exists an $h \in H$ with $ah = bh$; similarly, $Ha = Hb$ does not necessarily imply $a = b$ or that there exists an $h \in H$ with $ha = hb$.

Example 7.2.3

We saw above that in S_3 with $H = \langle (12) \rangle$,

$$(13)H = \{(13), (123)\} \neq \{(13), (132)\} = H(13).$$

Also, $(13)H = (123)H$ but $(13)e \neq (123)e$ and $(13)(12) \neq (123)(12)$.

It turns out that subgroups H for which $aH = Ha$ for all $a \in G$ will be very important to us.

Definition: Normal Subgroup

We say that subgroup H of G is **normal** in G (or is **normal subgroup** of G) if $aH = Ha$ for all $a \in G$. We denote that fact that H is normal in G by writing $H \trianglelefteq G$.

Remark

1. If H is normal in G , we may refer to the left and right cosets of G as simply **cosets**.
2. Of course, if G is abelian, every subgroup of G is normal in G . But there can also be normal subgroups of nonabelian groups: for instance, the trivial and improper subgroups of every group are normal in that group.

Example 7.2.4

Find the cosets of $5\mathbb{Z}$ in $\mathbb{Z}$.

Notice that in additive notation, the statement " $a^{-1}b \in H$ " becomes $-a + b \in H$. So for $a, b \in \mathbb{Z}$, $a \sim_L b$ if and only if $-a + b \in 5\mathbb{Z}$; that is, if and only if 5 divides $b - a$. In other words, $a \sim_L b$ if and only if $a \equiv_5 b$. So in this case, $\sim_L$ is just congruence modulo 5. Thus, the cosets of $5\mathbb{Z}$ in $\mathbb{Z}$ are

$$\begin{aligned} 5\mathbb{Z} &= \{\dots, -5, 0, 5, \dots\} \\ 1 + 5\mathbb{Z} &= \{\dots, -4, 1, 6, \dots\}, \\ 2 + 5\mathbb{Z} &= \{\dots, -3, 2, 7, \dots\}, \\ 3 + 5\mathbb{Z} &= \{\dots, -2, 3, 8, \dots\}, \\ 4 + 5\mathbb{Z} &= \{\dots, -1, 4, 9, \dots\}. \end{aligned}$$

Do you see how this example would generalize for $n\mathbb{Z}$ ($n^{\text{int}}\mathbb{Z}^+$) in $\mathbb{Z}$?

Example 7.2.5

Find the cosets of $H = \langle 12 \rangle$ in $4\mathbb{Z}$.

They are

$$\begin{aligned} H &= \{\dots, -12, 0, 12, \dots\}, \\ 4 + H &= \{\dots, -8, 4, 16, \dots\}, \\ 8 + H &= \{\dots, -4, 8, 20, \dots\}. \end{aligned}$$

Example 7.2.6

Find the cosets of $H = \langle 4 \rangle$ in $\mathbb{Z}_{12}$.

They are

$$\begin{aligned} H &= \{0, 4, 8\}, \\ 1 + H &= \{1, 5, 9\}, \\ 2 + H &= \{2, 6, 10\}, \\ 3 + H &= \{3, 7, 11\}. \end{aligned}$$

We now consider the set of all left cosets of a subgroup of a group.

Definition: $G \text{ mod } H$

We let G/H be the set of all left cosets of subgroup H in G . We read G/H as “ $G \text{ mod } H$.”

(We may denote the set of all right cosets of subgroup H in G by $H \backslash G$, but we will not use that notation in this class.)

Answers

Theorem 7.2.1

First, let $a \in G$. Then $a^{-1}a = e \in H$, so $a \sim_L a$. Thus, $\sim_L$ is reflexive.

Next, let $a, b \in G$ with $a \sim_L b$. Then $a^{-1}b \in H$, so, since H is a subgroup of G , $(a^{-1}b)^{-1} \in H$. But $(a^{-1}b)^{-1} = b^{-1}(a^{-1})^{-1} = b^{-1}a$; thus, $b \sim_L a$, and so $\sim_L$ is symmetric.

Finally, let $a, b, c \in G$ with $a \sim_L b$ and $b \sim_L c$. Then $a^{-1}b$ and $b^{-1}c$ are in H . Since H is a subgroup of G , we must then have $(a^{-1}b)(b^{-1}c) \in H$; but $(a^{-1}b)(b^{-1}c)$ equals $a^{-1}c$. Thus, $a \sim_L c$, and so $\sim_L$ is transitive.

Thus, $\sim_L$ is an equivalence relation on G . The proof that $\sim_R$ is an equivalence relation is left as an exercise for the reader.

Theorem 7.2.2

Let $b \in G$. Then $b \sim_L a \Leftrightarrow a \sim_L b \Leftrightarrow a^{-1}b \in H \Leftrightarrow a^{-1}b = h$ for some $h \in H \Leftrightarrow b = ah$ for some $h \in H \Leftrightarrow b \in aH$. So under $\sim_L$ we have $[a] = aH$. Similarly, under $\sim_R$ we have $[a] = Ha$.

Theorem 7.2.3

Statements 1 and 2 hold because the left and right cosets of H in G are equivalence classes. Statement 3 holds because no left or right coset of H other than H itself can contain e , since the left [right] cosets of H are mutually disjoint. For Statement 4: Define $f : H \rightarrow aH$ by $f(h) = ah$. It is straightforward to show that f is a bijection, so $|H| = |aH|$. Similarly, $|Ha| = |H|$.

This page titled [7.2: Introduction to Cosets and Normal Subgroups](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

Definition: Index

We define the **index** of H in G , denoted $(G : H)$, to be the cardinality of G/H .

Remark

If $H \leq G$, there are always have the same number of left cosets and right cosets of H in G . The proof of this is left as an exercise for the reader.

(Keep in mind your proof must apply even when there are infinitely many left and/or right cosets of H in G . Hint: Show there is a bijection between G/H and the set of all right cosets of H in G .)

In the cases in which G/H contains infinitely many elements, we won't worry about specific cardinality, and may simply say that the index of H in G is infinite, and write $(G : H) = \infty$. However, we can, of course, distinguish between countably infinite and uncountably infinite indices.

Note that if G is finite then $(G : H)$ must be finite; however, we see below that if G is infinite then $(G : H)$ could be finite or infinite.

Example 7.3.1

If $(\mathbb{R} : \mathbb{Z})$ were finite, then we'd be able to write $\mathbb{R}$ as a finite union of countable sets, implying that $\mathbb{R}$ is countable—which it isn't. Thus, $(\mathbb{R} : \mathbb{Z}) = \infty$.

Example 7.3.2

Since $\langle i \rangle = \{i, -1, -i, 1\}$ is a finite subgroup of C^* and C^* is infinite, we must have that $(C^* : \langle i \rangle)$ is infinite. However, $(C^* : C^*) = 1$.

Example 7.3.3

Referring to our previous examples, we have:

$$\begin{aligned}(S_3 : \langle (12) \rangle) &= 3, \\ (D_4 : \langle f \rangle) &= 4, \\ (\mathbb{Z} : 5\mathbb{Z}) &= 5, \\ (4\mathbb{Z} : 12\mathbb{Z}) &= 3, \\ \text{and } (\mathbb{Z}_{12} : \langle 4 \rangle) &= 4.\end{aligned}$$

Notice that in the cases in which G is finite, $(G : H) = |G|/|H|$. This makes sense, since the left cosets of H in G partition G , and each left coset has cardinality $|H|$.

Since the left cosets of a subgroup H of a group G partition G and all have the same cardinality, we have the following two theorems. The first is known as **Lagrange's Theorem** (named for the French mathematician Joseph-Louis Lagrange).

Theorem 7.3.1: Lagrange's Theorem

If G is a finite group and $H \leq G$, then $|H|$ divides $|G|$.

Theorem 7.3.2

If G is a finite group and $H \leq G$, then $(G : H) = |G|/|H|$.

Remark

The converse of Lagrange's Theorem does *not* hold. Indeed A_4 is a group of order 12 which contains no subgroup of order 6 even though 6 divides $|A_4| = 12$.

We end this chapter with two corollaries to Lagrange's Theorem.

Corollary 7.3.1

Let G be a finite group and let $a \in G$. Then $a^{|G|} = e_G$.

Proof available at the end of this section.

Finally, we have the following corollary, whose proof is left as an exercise for the reader.

Corollary 7.3.2

Let G be a group of prime order. Then G is cyclic. It follows that for every prime p , there exists a unique group of order p , up to isomorphism.

Answers

Corollary 7.3.1

Let $d = o(a)$. By Lagrange's Theorem, d divides $|G|$, so there exists $k \in \mathbb{Z}$ with $|G| = dk$. Then $a^{|G|} = a^{dk} = (a^d)^k = (e_G)^k = e_G$.

7.4: Exercises

1. How many distinct partitions of the set $S = \{a, b, c, d\}$ are there? You do not need to list them. (Yes, you can find this answer online. But I recommend doing the work yourself for practice working with partitions!)

2.

- Let $n^{\text{int}}\mathbb{Z}^+$. Prove that $\equiv_n$ is an equivalence relation on $\mathbb{Z}$.
- The cells of the induced partition of $\mathbb{Z}$ are called the **residue classes** (or **congruence classes**) of $\mathbb{Z}$ modulo n . Using set notation of the form $\{\dots, \#, \#, \#, \dots\}$ for each class, write down the residue classes of $\mathbb{Z}$ modulo 4.

3. Let G be a group with subgroup H . Prove that $\sim_R$ is an equivalence relation on G .

4. Find the indices of:

- $H = \langle (15)(24) \rangle$ in S_5
- $K = \langle (2354)(34) \rangle$ in S_6
- A_n in S_n

5. For each subgroup H of group G , (i) find the left and the right cosets of H in G , (ii) decide whether or not H is normal in G , and (iii) find $(G : H)$.

Write all permutations using **disjoint cycle notation**, and write all dihedral group elements using *standard form*.

- $H = 6\mathbb{Z}$ in $G = 2\mathbb{Z}$
- $H = \langle 4 \rangle$ in $\mathbb{Z}_{20}$
- $H = \langle (23) \rangle$ in $G = S_3$
- $H = \langle r \rangle$ in $G = D_4$
- $H = \langle f \rangle$ in $G = D_4$

6. For each of the following, give an example of a group G with a subgroup H that matches the given conditions. If no such example exists, prove that.

- A group G with subgroup H such that $|G/H| = 1$.
- A finite group G with subgroup H such that $|G/H| = |G|$.
- An abelian group G of order 8 containing a non-normal subgroup H of order 2.
- A group G of order 8 containing a normal subgroup of order 2.
- A nonabelian group G of order 8 containing a normal subgroup of index 2.
- A group G of order 8 containing a subgroup of order 3.
- An infinite group G containing a subgroup H of finite index.
- An infinite group G containing a finite nontrivial subgroup H .

7. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G be a group with subgroup H and elements $a, b^{\text{int}}G$.

- If $a^{\text{int}}bH$ then aH must equal bH .
- aH must equal Ha .
- If $aH = bH$ then Ha must equal Hb .
- If $a^{\text{int}}H$ then aH must equal Ha .
- H must be normal in G if there exists $a^{\text{int}}G$ such that $aH = Ha$.
- If $aH = bH$ then $ah = bh$ for every $h^{\text{int}}H$.
- If G is finite, then $|G/H|$ must be less than $|G|$.
- If G is finite, then $(G : H)$ must be less than or equal to $|G|$.

8. Let G be a group of order pq , where p and q are prime, and let H be a proper subgroup of G . Prove that H is cyclic.

9. Prove Corollary 7.3.2: that is, let G be a group of prime order, and prove that G is cyclic.

This page titled [7.4: Exercises](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

8: FACTOR GROUPS

8.1: Motivation

8.2: Focusing on Normal Subgroups

8.3: Introduction to Factor Groups

8.4: Exercises

This page titled [8: Factor Groups](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

8.1: Motivation

We mentioned previously that given a subgroup H of G , we'd like to use H to get at some understanding of G 's entire structure. Recall that we've defined G/H to be the set of all left cosets of H in G . What we'd like to do now is equip G/H with some operation under which G/H is a group! The natural way to do this would be to define multiplication on G/H by

$$(aH)(bH) = abH \text{ for all } a, b \in G.$$

Ok, so let's do that! But wait: we're defining this operation by referring to coset representatives, so we'd better make sure our operation is well-defined. Only it sadly turns out that in general this operation is *not* well-defined. For example:

Example 8.1.1

Let $H = \langle (12) \rangle$ in S_3 , and let $a = (13)$ and $b = (132)$. Let $x = aH$ and $y = bH$ in S_3/H . Then using the above-defined operation on G/H we'd have

$$xy = (aH)(bH) = abH = (13)(132)H = (23)H.$$

But also $x = (123)H$ and $y = (23)H$, so we'd also have

$$xy = ((123)H)((23)H) = (123)(23)H = (12)H = H \neq (23)H.$$

So this operation isn't well-defined.

The question for us now becomes: what conditions must hold for H in G in order for operation

$$(aH)(bH) = abH$$

on G/H to be well-defined? It turns out that this operation is well-defined exactly when H is normal in G ! We state this in the following theorem:

Theorem 8.1.1

Let $H \leq G$. Then the operation

$$(aH)(bH) = abH$$

on G/H is well-defined if and only if $H \trianglelefteq G$.

Proof available at the end of this section.

Definition: Left Coset Multiplication

When $H \trianglelefteq G$, the well-defined operation $(aH)(bH) = abH$ on G/H is called **left coset multiplication**.

It is clear that normal subgroups will be very important for us in studying group structures. We therefore spend some time looking at normal subgroups before returning to equipping G/H with a group structure.

Answers

Theorem 8.1.1

First, assume that the described operation is well-defined. Let $a \in G$; we want to show that $aH = Ha$.

Well, let $x \in aH$. Then

$$(xH)(a^{-1}H) = xa^{-1}H$$

and, since $xH = aH$

$$(xH)(a^{-1}H) = (aH)(a^{-1}H) = aa^{-1}H = H.$$

Since our operation on G/H is well-defined, this means that $xa^{-1}H = H$, so $xa^{-1} \in H$; thus, $x \in Ha$. We conclude that $aH \subseteq Ha$. The proof that $Ha \subseteq aH$ is similar. So $aH = Ha$, and thus, since $a \in G$ was arbitrary, H is normal in G .

Conversely, assume $H \trianglelefteq G$. Let $a_1, a_2, b_1, b_2 \in G$ with $a_1H = a_2H$ and $b_1H = b_2H$. We want to show that then $a_1b_1H = a_2b_2H$, that is, that $(a_2b_2)^{-1}a_1b_1 \in H$. Well, since $a_1H = a_2H$ we have $a_2^{-1}a_1 \in H$. So

$$(a_2b_2)^{-1}a_1b_1 = b_2^{-1}(a_2^{-1}a_1)b_1 \in b_2^{-1}Hb_1.$$

Next, since $H \trianglelefteq G$, we have $Hb_1 = b_1H$, so $b_2^{-1}Hb_1 = b_2^{-1}b_1H$; but since $b_1H = b_2H$, we have $b_2^{-1}b_1 \in H$, so $b_2^{-1}b_1H = H$. Thus, $(a_2b_2)^{-1}a_1b_1 \in H$, as desired.

8.2: Focusing on Normal Subgroups

We first provide a theorem that will help us in identifying when a subgroup of a group is normal. First, we provide a definition.

Definition

Let H be a subgroup of G and a, b in G . We define

$$aHb = \{ahb : h \in H\}.$$

Theorem 8.2.1

Let H be a subgroup of a group G . Then the following are equivalent:

1. H is normal in G ;
2. $aHa^{-1} = H$ for all $a \in G$;
3. $aHa^{-1} \subseteq H$ for all $a \in G$.

Proof available at the end of this section.

We now consider some examples and nonexamples of normal subgroups of groups.

Example 8.2.1

1. As previously mentioned, the trivial and improper subgroups of any group G are normal in G .
2. As previously mentioned, if group G is abelian then each of its subgroups is normal in G .
3. Suppose $H \leq G$ has $(G : H) = 2$. Then $H \trianglelefteq G$. The proof of this is left as an exercise for the reader.
4. Example 7.2.1 shows that subgroup $H = \langle (12) \rangle$ isn't normal in S_3 (for example, $(13)H \neq H(13)$). But $\langle (123) \rangle$ must be normal in S_3 since $(S_3 : \langle (123) \rangle) = 6/3 = 2$.
5. $\langle r \rangle$ is normal in D_n since $(D_n : \langle r \rangle) = 2$.
6. $\langle f \rangle$ isn't normal in D_4 : for instance,

$$r\langle f \rangle r^{-1} = \{e, rfr^3\} = \{e, fr^3r^3\} = \{e, fr^2\} \not\subseteq \langle f \rangle.$$

We consider two other very significant examples. First, we revisit the idea of the center of a group, first introduced in Example 2.8.9.

Definition: Center

Let G be a group. We let

$$Z(G) = \{z \in G : az = za \text{ for all } a \in G\}.$$

$Z(G)$ is called the **center** of G .

(The Z stands for “zentrum,” the German word for “center.”)

Theorem 8.2.2

Let G be a group. Then $Z(G)$ is a subgroup of G .

Proof available at the end of this section.

Theorem 8.2.3

Let G be a group and let H be a subgroup of G with $H \subseteq Z(G)$. Then $H \trianglelefteq G$. In particular, $Z(G)$ is itself a normal subgroup of G .

Proof available at the end of this section.

The next definition is profoundly important for us.

Definition: Kernel

Let G and G' be groups and let ϕ a homomorphism from G to G' . Letting e' be the identity element of G' , we define the **kernel** of ϕ , $\text{Ker } \phi$, by

$$\text{Ker } \phi = \{k^{\text{int}} G : \phi(k) = e'\}.$$

Theorem 8.2.4

Let G and G' be groups and let ϕ be a homomorphism from G to G' . Then $\text{Ker } \phi$ is a normal subgroup of G .

Proof available at the end of this section.

One slick way, therefore, of showing that a particular set is a normal subgroup of a group G is by showing it's the kernel of a homomorphism from G to another group.

Example 8.2.2

Let $n^{\text{int}}\mathbb{Z}^+$. Here is a rather elegant proof of the fact that $SL(n, \mathbb{R})$ is a normal subgroup of $GL(n, \mathbb{R})$: Define $\phi : GL(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ by $\phi(A) = \det A$. Clearly, ϕ is a homomorphism, and since the identity element of $\mathbb{R}^*$ is 1,

$$\text{Ker } \phi = \{A^{\text{int}} GL(n, \mathbb{R}) : \det A = 1\} = SL(n, \mathbb{R}).$$

So $SL(n, \mathbb{R}) \trianglelefteq GL(n, \mathbb{R})$.

Answers

Theorem 8.2.1

The equivalence of Statements 1 and 2 is clear, as is the fact that Statement 2 implies Statement 3. So it suffices to show that Statement 3 implies Statement 2. Well, assume that $aHa^{-1} \subseteq H$ for all $a \in G$, and let $b \in G$. We want to show that $bHb^{-1} = H$. Since Statement 3 holds, we clearly have $bHb^{-1} \subseteq H$. But, again using Statement 3, we also have $b^{-1}Hb \subseteq H$; multiplying both sides of this equation on the left by b and on the right by b^{-1} , we obtain $H \subseteq bHb^{-1}$. Hence, since $bHb^{-1} \subseteq H$ and $H \subseteq bHb^{-1}$, those two sets are equal. Thus, Statement 2 is proven.

Theorem 8.2.2

Clearly, $e \in Z(G)$. Next, let $z, w \in Z(G)$. Then for all $a \in G$,

$$a(zw) = (az)w = (za)w = z(aw) = z(wa) = (zw)a,$$

so $zw \in Z(G)$. Finally, $az = za$ so, multiplying both sides on the left and right by z^{-1} , we have $z^{-1}a = az^{-1}$; thus, $z^{-1} \in Z(G)$. Hence, $Z(G) \leq G$.

Theorem 8.2.3

Let $a \in G$. Then since every element of $Z(G)$ commutes with every element of G , every element of H commutes with every element of G ; so we must have $aH = Ha$. Thus, $H \trianglelefteq G$.

Theorem 8.2.4

Let $K = \text{Ker } \phi$. We first show that K is a subgroup of G . Clearly, the identity element of G is in K , so $K \neq \emptyset$. Next, let $k, m \in K$. Then, letting e' denote the identity element of G' , we have

$$\phi(km^{-1}) = \phi(k)\phi(m)^{-1} = e'(e')^{-1} = e',$$

so $km^{-1} \in K$. Thus, by the Two-Step Subgroup Test, we have that K is a subgroup of G .

Next, let $k \in K$ and let $a \in G$. Then

$$\phi(aka^{-1}) = \phi(a)\phi(k)\phi(a)^{-1} = \phi(a)e'e^{-1} = \phi(a)\phi(a)^{-1} = e'.$$

So $aka^{-1} \in K$. Thus, $K \trianglelefteq G$.

8.3: Introduction to Factor Groups

We now return to the notion of equipping G/H , when $H \trianglelefteq G$, with a group structure. We have already seen that left coset multiplication on G/H is well-defined when $H \trianglelefteq G$ (Theorem 8.1.1); it turns out that given this, it is very easy to prove that G/H under this operation is a group.

Before we prove this, we introduce a change in our notation: We initiate the convention of frequently using N , rather than H , to denote a subgroup of a group G when we know that that subgroup is **normal** in G .

Theorem 8.3.1

Let G be a group with identity element e , and let $N \trianglelefteq G$. Then G/N is a group under left coset multiplication (that is, under the operation $(aN)(bN) = abN$ for all $a, b \in G$), and $|G/N| = (G : N)$ (in particular, if $|G| < \infty$, then $|G/N| = |G|/|N|$).

Proof available at the end of this section.

Definition: Factor Group

When G is a group and $N \trianglelefteq G$, the above group $(G/N$ under left coset multiplication) is called a **factor group** or **quotient group**.

Example 8.3.1

Let $G = \mathbb{Z}$ and $N = 3\mathbb{Z}$. Then N is normal in G , since G is abelian, so the set $G/N = \{N, 1 + N, 2 + N\}$ is a group under left coset multiplication. Noting that $N = 0 + N$, it is straightforward to see that G/N (that is, $\mathbb{Z}/3\mathbb{Z}$) has the following group table:

+	$0 + N$	$1 + N$	$2 + N$
$0 + N$	$0 + N$	$1 + N$	$2 + N$
$1 + N$	$1 + N$	$2 + N$	$0 + N$
$2 + N$	$2 + N$	$0 + N$	$1 + N$

Clearly, if we ignore all the $+N$'s after the 0's, 1's, and 2's in the above table, and consider $+$ to be addition mod 3, rather than left coset addition in $\mathbb{Z}/3\mathbb{Z}$, we obtain the group table for $\mathbb{Z}_3$:

+	0	1	2
0	0	1	2
1	1	2	0
2	2	0	1

Thus, we see that $\mathbb{Z}/3\mathbb{Z}$ is isomorphic to $\mathbb{Z}_3$.

This is not a coincidence! In fact, we have the following:

Theorem 8.3.2

Let $n, d \in \mathbb{Z}^+$ with d dividing n . Then we have:

1. $n\mathbb{Z} \trianglelefteq d\mathbb{Z}$ and $\langle d \rangle \trianglelefteq \mathbb{Z}_n$;
2. $d\mathbb{Z}/n\mathbb{Z} \simeq \mathbb{Z}_{n/d}$ (special case: $\mathbb{Z}/n\mathbb{Z} \simeq \mathbb{Z}_n$); and
3. $\mathbb{Z}_n/\langle d \rangle \simeq \mathbb{Z}_d$.

Proof available at the end of this section.

Example 8.3.2

Let $N = \langle (123) \rangle \leq S_3$. Since $(S_3 : N) = 2$, N must be normal in S_3 , so S_3/N is a group under left coset multiplication. Since $|S_3/N| = 2$, S_3/N must be isomorphic to $\mathbb{Z}_2$.

In the above examples, we were able to identify G/N up to isomorphism relatively easily because we could determine that G/N was cyclic. In general, however, it can be quite difficult to identify the group structure of a factor group. We explore some powerful tools we can use in this identification in the next section, but first we note a couple properties of G that are “inherited” by any of its factor groups.

Theorem 8.3.3

Let G be a group and $N \trianglelefteq G$. Then:

1. If G is finite, then G/N is finite.
2. If G is abelian, then G/N is abelian; and
3. If G is cyclic, then G/N is cyclic.

Proof available at the end of this section.

Note

However, G can be nonabelian [noncyclic, nonfinite] and yet have a normal subgroup N such that G/N is abelian [cyclic, finite]. (See the examples below.) Intuitively, the idea is that modding out a group by a normal subgroup can *introduce* abelianness or cyclicity, or finiteness, but not *remove* one of those characteristics.

Example 8.3.3

S_3 is nonabelian (and therefore of course noncyclic), but we saw above that $N = \langle (123) \rangle$ is a normal subgroup of S_3 with $S_3/N \simeq \mathbb{Z}_2$, a cyclic (and therefore of course abelian) group.

Example 8.3.4

$\mathbb{Z}$ is an infinite group, but it has finite factor group $\mathbb{Z}/2\mathbb{Z}$.

What do the (normal) subgroups of a factor group G/N look like? Well, they come from the (normal) subgroups of G ! We have the following theorem, whose proof is tedious but straightforward.

Theorem 8.3.4: Correspondence Theorem

Let G be a group, and let $K \trianglelefteq G$. Then the subgroups of G/K are exactly those of the form H/K , where $H \leq G$ with $K \subseteq H$. Moreover, the normal subgroups of G/K are exactly those of the form N/K , where $N \trianglelefteq G$ with $K \subseteq N$.

Example 8.3.5

Let $N = 18\mathbb{Z}$ in $\mathbb{Z}$. Since the subgroups of $\mathbb{Z}$ containing N are the sets $d\mathbb{Z}$ where d is a positive divisor of 18, the subgroups of $\mathbb{Z}/N$ are the sets $d\mathbb{Z}/N$, where, again, d is a positive divisor of 18.

We end this chapter by noting that given any group G and factor group G/N of G , there is a homomorphism from G to G/N that is onto. Before we define this homomorphism, we provide some more terminology.

Definition: Epimorphism and Monomorphism

Let $\phi : G \rightarrow G'$ be a homomorphism of groups. Then ϕ can be called an **epimorphism** if ϕ is onto, and a **monomorphism** if ϕ is one-to-one. (Of course, we already know that an epimorphism that is also a monomorphism is called an “isomorphism.”)

We now present the following theorem, whose straightforward proof is left to the reader.

Theorem 8.3.5

Let G be a group and let $N \trianglelefteq G$. Then the map $\Psi : G \rightarrow G/N$ defined by $\Psi(g) = gN$ is an epimorphism.

Definition: Canonical Epimorphism

We call this map Ψ the **canonical epimorphism** from G to G/N .

Notice that given $N \trianglelefteq G$, the kernel of the canonical epimorphism from G to G/N is N . Thus, putting this fact together with the fact that every kernel of a homomorphism is a normal subgroup of the homomorphism's domain, we have the following:

Theorem 8.3.6

Let G be a group and N a subset of G . Then N is a normal subgroup of G if and only if N is the kernel of a homomorphism from G to some group G' .

Answers

Theorem 8.3.1

We already know that since $N \trianglelefteq G$, left coset multiplication on G/N is well-defined; further, by definition, that $(aN)(bN) = abN \in G/N$ for every $aN, bN \in G/N$. Associativity of this operation on G/N follows from the associativity of G 's operation: indeed, if $aN, bN, cN \in G/N$, then

Illegal pream-token (&)

Next, $N = eN \in G/N$ acts as an identity element since for all $a \in G$,

$$(aN)(N) = aeN = aN \text{ and } N(aN) = eaN = aN.$$

Finally, let $aN \in G/N$. Then $a^{-1}N \in G/N$ with $(aN)(a^{-1}N) = aa^{-1}N = N$ and $(a^{-1}N)(aN) = a^{-1}aN = N$.

So G/N is a group under left coset multiplication. Since $(G : N)$ is, by definition, the cardinality of G/N , we're done.

Theorem 8.3.2

- Since d is a positive divisor of n , $n\mathbb{Z}$ and $\langle d \rangle$ are clearly subgroups of, respectively, $d\mathbb{Z}$ and $\mathbb{Z}_n$. Moreover, since $d\mathbb{Z}$ and $\mathbb{Z}_n$ are abelian, all of their subgroups are normal.
- This follows from the facts that $d\mathbb{Z}/n\mathbb{Z} = \langle d + n\mathbb{Z} \rangle$, hence is cyclic, and that $|d\mathbb{Z}/n\mathbb{Z}| = (d\mathbb{Z} : n\mathbb{Z}) = n/d$ (see Example 7.3.3). (Unpacking the statement that $d\mathbb{Z}/n\mathbb{Z} = \langle d + n\mathbb{Z} \rangle$: Notice that $d\mathbb{Z} = \langle d \rangle$, so every element of $d\mathbb{Z}$ is of the form kd for some integer k . Thus, every element of $d\mathbb{Z}/n\mathbb{Z}$ is of the form $kd + n\mathbb{Z}$ for some integer k . But for each $k \in \mathbb{Z}$, using the definition of left coset multiplication we have that $kd + n\mathbb{Z} = k(d + n\mathbb{Z})$. So $d\mathbb{Z}/n\mathbb{Z} = \langle d + n\mathbb{Z} \rangle$.)
- Similarly, since $\mathbb{Z}_n = \langle 1 \rangle$, $\mathbb{Z}_n/\langle d \rangle = \langle 1 + \langle d \rangle \rangle$, so is cyclic. Since $(\mathbb{Z}_n : \langle d \rangle) = d$ (again, see Example 7.3.3), $\mathbb{Z}_n/\langle d \rangle$ is isomorphic to $\mathbb{Z}_d$, as desired.

Q.E.D.

Theorem 8.3.3

- Clearly this holds, since in this case $|G/N| = |G|/|N|$.
- The proof of this statement is left as an exercise for the reader.
- Let G be cyclic. Then there exists $a \in G$ with $G = \langle a \rangle$. We claim $G/N = \langle aN \rangle$. Indeed:

$$\langle aN \rangle = (aN)i : i \in \mathbb{Z} = aiN : i \in \mathbb{Z}.$$

But every element of G is an integer power of a , so this set equals $\{xN : x \in G\}$, that is, it equals G/N .

8.4: Exercises

1. Let G be a group and let $H \leq G$ have index 2. Prove that $H \trianglelefteq G$.
2. Let G be an abelian group with $N \trianglelefteq G$. Prove that G/N is abelian.
3. Find the following.
 - a. $|2\mathbb{Z}/6\mathbb{Z}|$
 - b. $|H|$, for $H = 2 + \langle 6 \rangle \subseteq \mathbb{Z}_{12}$
 - c. $o(2 + \langle 6 \rangle)$ in $\mathbb{Z}_{12}/\langle 6 \rangle$
 - d. $\langle fH \rangle$ in D_4/H , where $H = \{e, r^2\}$
 - e. $|(\mathbb{Z}_6 \times \mathbb{Z}_8)/(\langle 3 \rangle \times \langle 2 \rangle)|$
 - f. $|(\mathbb{Z}_{15} \times \mathbb{Z}_{24})/(\langle 5, 4 \rangle)|$
4. For each of the following, find a familiar group to which the given group is isomorphic. (Hint: Consider the group order, properties such as abelianness and cyclicity, group tables, orders of elements, etc.)
 - a. $\mathbb{Z}/14\mathbb{Z}$
 - b. $3\mathbb{Z}/12\mathbb{Z}$
 - c. S_8/A_8
 - d. $(\mathbb{Z}_4 \times \mathbb{Z}_{15})/(\langle 2 \rangle \times \langle 3 \rangle)$
 - e. $D_4/\langle r^2 \rangle$
5. Let $H \trianglelefteq G$ with index k , and let $a^{\text{int}} G$. Prove that Double exponent: use braces to clarify

9: THE ISOMORPHISM THEOREM

Recall that our goal here is to use a subgroup of a group G to study not just the structure of the subgroup, but the structure of G outside of that subgroup (the ultimate goal being to get a feeling for the structure of G as a whole). We've further seen that if we choose N to be a normal subgroup of G , we can do this by studying both N and the factor group G/N . Now, we've noticed that in some cases—in particular, when G is cyclic—it is not too hard to identify the structure of a factor group of G . But what about when G and N are more complicated? For instance, we have seen that $SL(5, \mathbb{R})$ is a normal subgroup of $GL(5, \mathbb{R})$. What is the structure of $GL(5, \mathbb{R})/SL(5, \mathbb{R})$? That is not so easy to figure out by looking directly at left coset multiplication in the factor group.

9.1: The First Isomorphism Theorem

9.2: The Second and Third Isomorphism Theorems

9.3: Exercises

9.1: The First Isomorphism Theorem

A very powerful theorem, called the *First Isomorphism Theorem*, lets us in many cases identify factor groups (up to isomorphism) in a very slick way. Kernels will play an extremely important role in this. We therefore first provide some theorems relating to kernels.

Theorem 9.1.1

Let G and G' be groups, let ϕ be a homomorphism from G to G' , and let $K = \text{Ker } \phi$. Then for $a, b \in G$, $aK = bK$ if and only if $\phi(a) = \phi(b)$.

Proof available at the end of this section.

Corollary 9.1.1

Let ϕ be a homomorphism from group G to group G' . Then ϕ is one-to-one (hence a monomorphism) if and only if $\text{Ker } \phi = \{e_G\}$.


Proof available at the end of this section.

We now prove a theorem that provides the meat and potatoes of the First Isomorphism Theorem.

Theorem 9.1.2: Factorization Theorem

Let G and G' be groups, let ϕ be a homomorphism from G to G' , let $K = \text{Ker } \phi$, let N be a normal subgroup of G with $N \subseteq K$, and let Ψ be the canonical epimorphism from G to G/N . Then the map $\bar{\phi} : G/N \rightarrow G'$ defined by $\bar{\phi}(aN) = \phi(a)$ is a well-defined homomorphism, with $\bar{\phi} \circ \Psi = \phi$.

We summarize this in the following picture:



Proof available at the end of this section.

Note that the above theorem does *not* state that $\overline{\phi}$ is a monomorphism or an epimorphism. This is because in general it may be neither! We do have the following theorem:

Theorem 9.1.3

Let G , G' , ϕ , N , and $\overline{\phi}$ be as defined in the Factorization Theorem. Then

1. $\overline{\phi}$ is onto (an epimorphism) if and only if ϕ is onto (an epimorphism); and
2. $\overline{\phi}$ is one-to-one (a monomorphism) if and only if $N = \text{Ker}\phi$.

Proof available at the end of this section.

We are now ready to state the all-important First Isomorphism Theorem, which follows directly from the Factorization Theorem and Theorem 9.1.3.

Theorem 9.1.4: First Isomorphism Theorem

Let G and G' be groups, with homomorphism $\phi : G \rightarrow G'$. Let $K = \text{Ker}\phi$. Then $G/K \simeq \phi(G)$. In particular, if ϕ is onto, then $G/K \simeq G'$.

So to prove that a factor group G/N is isomorphic to a group G' , it suffices to show there exists an epimorphism from G to G' that has N as its kernel.

Example 9.1.1

Letting $n^{\text{int}}\mathbb{Z}^+$, let's identify a familiar group to which $GL(n, \mathbb{R})/SL(n, \mathbb{R})$ is isomorphic. As in Example 8.2.3, the map $\phi : GL(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ defined by $\phi(A)$ is a homomorphism with kernel $SL(n, \mathbb{R})$. Moreover, ϕ clearly maps onto $\mathbb{R}^*$: indeed, given $\lambda \in \mathbb{R}^*$, the diagonal matrix having λ in the uppermost left position and 1's elsewhere down the diagonal gets sent to λ by ϕ . So by the First Isomorphism Theorem, we have $GL(n, \mathbb{R})/SL(n, \mathbb{R}) \simeq \mathbb{R}^*$.

Example 9.1.2

Let $G = S_3 \times \mathbb{Z}_{52}$ and let $N = S_3 \times \{0\} \subseteq G$. It is straightforward to show that N is normal in G . What is the structure of G/N ? Well, define $\phi : G \rightarrow \mathbb{Z}_{52}$ by $\phi((\sigma, a)) = a$. Then ϕ is clearly an epimorphism and $\text{Ker}\phi = \{(\sigma, a)^{\text{int}}G : a = 0\} = N$. So G/N is isomorphic to $\mathbb{Z}_{52}$.

Generalizing the above example, we have the following theorem, whose proof we leave to the reader.

Theorem 9.1.5

Let $G = G_1 \times G_2 \times \cdots \times G_k$ (where $k \in \mathbb{Z}^+$) and let N_i be a normal subgroup of G_i for each $i = 1, 2, \dots, k$. Then $N = N_1 \times N_2 \times \cdots \times N_k$ is a normal subgroup of G , with $G/N \simeq G_1/N_1 \times G_2/N_2 \times \cdots \times G_k/N_k$.

We provide one more cool example of using the First Isomorphism Theorem. Clearly, since $\mathbb{R}$ is abelian, $\mathbb{Z}$ is a normal subgroup of $\mathbb{R}$. What is the structure of $\mathbb{R}/\mathbb{Z}$? Well, in modding $\mathbb{R}$ out by $\mathbb{Z}$ we have essentially identified together all real numbers that are an integer distance apart. So we can think of the canonical epimorphism from $\mathbb{R}$ to $\mathbb{R}/\mathbb{Z}$ as wrapping up $\mathbb{R}$ like a garden hose! Thus, one might guess that $\mathbb{R}/\mathbb{Z}$ has some circle-like structure—but if we want to think of it as a group, we have to figure out what the group structure on such a “circle” would be!

We leave, for a moment, our group $\mathbb{R}/\mathbb{Z}$, and look at how we can consider a circle to be a group.

Recall that for every $\theta \in \mathbb{R}$, $e^{i\theta}$ is defined to be $\cos \theta + i \sin \theta$. It is clear then that the set $S^1 = \{e^{i\theta} : \theta \in \mathbb{R}\}$ is the unit circle in the complex plane.

Remark

Note that if $\theta_1, \theta_2 \in \mathbb{R}$, then $e^{i\theta_1} = e^{i\theta_2}$ if and only if $\theta_1 - \theta_2 \in 2\pi\mathbb{Z}$.

Theorem 9.1.6

S^1 is a group under the multiplication $e^{i\theta_1} e^{i\theta_2} = e^{i(\theta_1 + \theta_2)}$.

Proof available at the end of this section.

Beautifully, it turns out that our group $\mathbb{R}/\mathbb{Z}$ is isomorphic to S^1 .

Theorem 9.1.7

$\mathbb{R}/\mathbb{Z} \simeq S^1$.

Proof available at the end of this section.

Answers

Theorem 9.1.1

Let $a, b \in G$. Then

as desired.

Corollary 9.1.1

Clearly, if ϕ is one-to-one then $\text{Ker} \phi = \{e_G\}$. Conversely, assume $\text{Ker} \phi = \{e_G\}$.

If $a, b \in G$ with $\phi(a) = \phi(b)$, then by the above theorem, $a \text{Ker} \phi = b \text{Ker} \phi$. But $a \text{Ker} \phi = a\{e_G\} = \{a\}$ and $b \text{Ker} \phi = b\{e_G\} = \{b\}$. Thus, $a = b$, and we see that ϕ is one-to-one.

Theorem 9.1.2: Factorization Theorem

We define $\bar{\phi} : G/N \rightarrow G'$, as indicated above, by $\bar{\phi}(aN) = \phi(a)$ for every $aN \in G/N$. Since $\bar{\phi}$ is defined using coset representatives, we must first show that $\bar{\phi}$ is well-defined. So let $aN = bN \in G/N$. Then $\bar{\phi}(aN) = \phi(a)$ and $\bar{\phi}(bN) = \phi(b)$, so we must show that $\phi(a) = \phi(b)$. Since $aN = bN$ and $N \subseteq \text{Ker} \phi$, we have that $aK = bK$ by Statement 6 of Theorem 7.2.3; thus $\phi(a) = \phi(b)$ (using Theorem 9.1.1). So $\bar{\phi}$ is well-defined.

Next, we show that $\bar{\phi}$ is a homomorphism. Let $aN, bN \in G/N$. Then

$$\bar{\phi}((aN)(bN)) = \bar{\phi}(abN) = \phi(ab) = \phi(a)\phi(b) = \bar{\phi}(aN)\bar{\phi}(bN).$$

Finally, for every $a \in G$,

$$(\bar{\phi} \circ \Psi)(a) = \bar{\phi}(\Psi(a)) = \bar{\phi}(aN) = \phi(a);$$

so $\bar{\phi} \circ \Psi = \phi$, as desired.

Theorem 9.1.3

This is clear, since $\bar{\phi}(G/N) = \phi(G)$.

By Corollary 9.1.1, $\bar{\phi}$ is one-to-one if and only if $\text{Ker} \bar{\phi} = \{N\}$. But

So $\text{Ker} \bar{\phi} = \{N\}$ if and only if

$$\{aN \in G/N : a \in \text{Ker} \phi\} = \{N\},$$

in other words if and only if $aN = N$ for all $a \in \text{Ker} \phi$. But

since we are given that $N \subseteq \text{Ker} \phi$. Thus, $\bar{\phi}$ is one-to-one if and only if $N = \text{Ker} \phi$, as desired.

Theorem 9.1.6

The tricky part is showing that the operation is well-defined. Suppose θ_1, θ_2, t_1 , and t_2 are in $\mathbb{R}$, with $e^{i\theta_1} = e^{it_1}$ and $e^{i\theta_2} = e^{it_2}$. We want to show that

$$e^{i\theta_1} e^{i\theta_2} = e^{it_1} e^{it_2},$$

i.e., that

$$e^{i(\theta_1 + \theta_2)} = e^{i(t_1 + t_2)}.$$

Now, $e^{i\theta_1} = e^{it_1}$ and $e^{i\theta_2} = e^{it_2}$ imply that $\theta_1 - t_1 = 2\pi m$ and $\theta_2 - t_2 = 2\pi n$ for some m and n in $\mathbb{Z}$; hence,

$$\theta_1 + \theta_2 - (t_1 + t_2) = 2\pi(m + n) \in 2\pi\mathbb{Z}.$$

Thus, $e^{i(\theta_1 + \theta_2)} = e^{i(t_1 + t_2)}$, so our operation is well-defined.

Since products of elements in S^1 are in S^1 , we have that S^1 is a binary structure; moreover, associativity of the operation in S^1 follows from the associativity of addition in $\mathbb{R}$. Moreover, $e^{i0} = 1$ clearly acts as an identity element in S^1 , and if $e^{i\theta} \in S^1$, then $e^{i\theta}$ has inverse $e^{i(-\theta)} \in S^1$. So S^1 is a group under the described operation.

Theorem 9.1.7

We will define an epimorphism ϕ from $\mathbb{R}$ to S^1 with $\text{Ker}\phi = \mathbb{Z}$; then we'll have $\mathbb{R}/\mathbb{Z} \simeq S^1$, by the First Isomorphism Theorem.

Define $\phi : \mathbb{R} \rightarrow S^1$ by $\phi(r) = e^{i2\pi r}$. We have that ϕ is a homomorphism, since for every $r, s \in \mathbb{R}$, we have

$$\phi(r + s) = e^{i2\pi(r+s)} = e^{i2\pi r + i2\pi s} = e^{i2\pi r} e^{i2\pi s} = \phi(r)\phi(s).$$

Moreover, ϕ is clearly onto, since if $e^{i\theta} \in S^1$, then

$$e^{i\theta} = e^{i2\pi(\theta/2\pi)} = \phi(\theta/2\pi).$$

Finally, $\text{Ker}\phi = \mathbb{Z}$: indeed,

Illegal pream-token (&)

Thus, $\text{Ker}\phi = \mathbb{Z}$, and hence $\mathbb{R}/\mathbb{Z} \simeq S^1$, as desired.

9.2: The Second and Third Isomorphism Theorems

The following theorems can be proven using the First Isomorphism Theorem. They are very useful in special cases.

Theorem 9.2.1: Second Isomorphism Theorem

Let G be a group, let $H \leq G$, and let $N \trianglelefteq G$. Then the set

$$HN = \{hn : h \in H, n \in N\}$$

is a subgroup of G , $H \cap N \trianglelefteq H$, and

$$H/(H \cap N) \simeq HN/N.$$

Proof available at the end of this section.

Theorem 9.2.2: Third Isomorphism Theorem

Let G be a group, and let K and N be normal subgroups of G , with $K \subseteq N$. Then $N/K \trianglelefteq G/K$, and

$$(G/K)/(N/K) \simeq G/N.$$

Proof available at the end of this section.

Example 9.2.1

Using the Third Isomorphism Theorem, we see that the group

$$(\mathbb{Z}/12\mathbb{Z})/(\langle 6\mathbb{Z}/12\mathbb{Z} \rangle)$$

is isomorphic to the group $\mathbb{Z}/6\mathbb{Z}$, or $\mathbb{Z}_6$.

Answers

Theorem 9.2.1: Second Isomorphism Theorem

We first prove that HN is a subgroup of G . Since $eG \in HN$, $HN \neq \emptyset$. Next, let $x = h_1n_1, y = h_2n_2 \in HN$ (where $h_1, h_2 \in H$ and $n_1, n_2 \in N$). Then

$$xy^{-1} = h_1n_1(h_2n_2)^{-1} = h_1n_1n_2^{-1}h_2^{-1}.$$

Since $N \leq G$, $n_1n_2^{-1}$ is in N ; so $h_1n_1n_2^{-1}h_2^{-1} \in h_1Nh_2^{-1}$, which equals $h_1h_2^{-1}N$, since $N \trianglelefteq G$ implies $Nh_2^{-1} = h_2^{-1}N$. So $xy^{-1} \in h_1h_2^{-1}N$. But $H \leq G$ implies $h_1h_2^{-1} \in H$; thus, $xy^{-1} \in HN$, and so HN is a subgroup of G .

Since $N \trianglelefteq G$ and $N \subseteq HN$, N is normal in HN (do you see why?). So HN/N is a group under left coset multiplication. We define $\phi : H \rightarrow HN/N$ by $\phi(h) = hN$ (notice that when $h \in H$, $h \in HN$ since $h = he_G$). Clearly, ϕ is a homomorphism. Further, ϕ is onto: Indeed, let $y \in HN/N$. Then $y = h_nN$ for some $h \in H$ and $n \in N$. But $nN = N$, so $y = hN = \phi(h)$. Finally,

$$H/(H \cap N) \simeq HN/N.$$

Thus,

by the First Isomorphism Theorem.

Theorem 9.2.2: Third Isomorphism Theorem

Define $\phi : G/K \rightarrow G/N$ by $\phi(aK) = aN$. We have that ϕ is well-defined: indeed, let $aK = bK \in G/K$. Then by Statement 6 of Theorem 7.2.3, we have $aN = bN$, that is, $\phi(aK) = \phi(bK)$. So ϕ is well-defined. ϕ is clearly onto, and we have

$$(G/K)/(N/K) \simeq G/N.$$

So the desired results hold, by the First Isomorphism Theorem.

This page titled [9.2: The Second and Third Isomorphism Theorems](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by [Jessica K. Sklar](#) via [source content](#) that was edited to the style and standards of the LibreTexts platform.

9.3: Exercises

1. Let F be the group of all functions from $[0, 1]$ to $\mathbb{R}$, under pointwise addition. Let

$$N = \{f \in F : f(1/4) = 0\}.$$

Prove that F/N is a group that's isomorphic to $\mathbb{R}$.

2. Let $N = \{1, -1\} \subseteq \mathbb{R}^*$. Prove that $\mathbb{R}^*/N$ is a group that's isomorphic to $\mathbb{R}^+$.

3. Let $n \in \mathbb{Z}^+$ and let $H = \{A \in GL(n, \mathbb{R}) : \det A = \pm 1\}$. Identify a group familiar to us that is isomorphic to $GL(n, \mathbb{R})/H$.

4. Let G and G' be groups with respective normal subgroups N and N' . Prove or disprove: If $G/N \cong G'/N'$ then $G \cong G'$.

INDEX

A

- abelian
 - [2.5: Group Conventions and Properties](#)
- addition modulo
 - [2.6: Examples of Groups/Nongroups, Part II](#)
- alternating group
 - [6.3: Alternating Groups](#)
- associative
 - [2.1: Binary Operations and Structures](#)
- associativity
 - [2.3: The Definition of a Group](#)
- automorphism
 - [3.2: Definitions of Homomorphisms and Isomorphisms](#)

B

- Bell numbers
 - [7.1: Partitions of and Equivalence Relations on Sets](#)
- bijection
 - [1.2: Functions](#)
 - [2.6: Examples of Groups/Nongroups, Part II](#)
- binary (algebraic) structure
 - [2.1: Binary Operations and Structures](#)
- binary operation
 - [2.1: Binary Operations and Structures](#)

C

- canonical epimorphism
 - [8.3: Introduction to Factor Groups](#)
- cardinality
 - [1.3: Cardinality](#)
- Cartesian product
 - [1.1: Sets](#)
- Cayley table
 - [3.1: Groups of Small Order](#)
- closed
 - [2.1: Binary Operations and Structures](#)
- closure
 - [2.1: Binary Operations and Structures](#)
- codomain
 - [1.2: Functions](#)
- commutative
 - [2.1: Binary Operations and Structures](#)
- commute
 - [2.1: Binary Operations and Structures](#)
- compose
 - [1.2: Functions](#)
- composition
 - [1.2: Functions](#)
- conjugation
 - [3.2: Definitions of Homomorphisms and Isomorphisms](#)
- countable
 - [1.3: Cardinality](#)
- countably infinite
 - [1.3: Cardinality](#)
- cyclic
 - [5.1: Introduction to Cyclic Groups](#)

cyclic subgroup

- [5.1: Introduction to Cyclic Groups](#)

D

- difference
 - [1.1: Sets](#)
- direct product
 - [1.1: Sets](#)
- disjoint
 - [1.1: Sets](#)
- domain
 - [1.2: Functions](#)

E

- elements
 - [1.1: Sets](#)
- empty set
 - [1.1: Sets](#)
- endomorphism
 - [3.2: Definitions of Homomorphisms and Isomorphisms](#)
- epimorphism
 - [8.3: Introduction to Factor Groups](#)
- equivalence relation
 - [7.1: Partitions of and Equivalence Relations on Sets](#)
- even
 - [6.3: Alternating Groups](#)

F

- factor group
 - [8.3: Introduction to Factor Groups](#)
- factorization theorem
 - [9.1: The First Isomorphism Theorem](#)
- finite
 - [1.3: Cardinality](#)
- finite group
 - [2.5: Group Conventions and Properties](#)
- first isomorphism theorem
 - [9.1: The First Isomorphism Theorem](#)
- function
 - [1.2: Functions](#)

G

- generator
 - [5.1: Introduction to Cyclic Groups](#)
- group
 - [2.3: The Definition of a Group](#)
- group axiom
 - [2.3: The Definition of a Group](#)
- group invariant
 - [3.3: Isomorphic Groups](#)
- group table
 - [3.1: Groups of Small Order](#)

H

- homomorphism
 - [3.2: Definitions of Homomorphisms and Isomorphisms](#)

I

- idempotent
 - [2.2: Exercises, Part I](#)
- identity element
 - [2.1: Binary Operations and Structures](#)
- identity elements
 - [2.3: The Definition of a Group](#)
- identity function
 - [1.2: Functions](#)

image

- [1.2: Functions](#)

improper subgroup

- [4.1: Introduction to Subgroups](#)

infinite

- [1.3: Cardinality](#)

infinite group

- [2.5: Group Conventions and Properties](#)

intersection

- [1.1: Sets](#)

inverse

- [1.2: Functions](#)
- [2.3: The Definition of a Group](#)

invertible

- [1.2: Functions](#)

isomorphic

- [3.3: Isomorphic Groups](#)

isomorphism

- [3.2: Definitions of Homomorphisms and Isomorphisms](#)

L

left cancellation law

- [2.5: Group Conventions and Properties](#)

M

mod

- [2.6: Examples of Groups/Nongroups, Part II](#)

monomorphism

- [8.3: Introduction to Factor Groups](#)

multiplication modulo

- [2.6: Examples of Groups/Nongroups, Part II](#)

mutually disjoint

- [1.1: Sets](#)

N

nonabelian

- [2.5: Group Conventions and Properties](#)

nonexample

- [3.3: Isomorphic Groups](#)

nonisomorphic

- [3.3: Isomorphic Groups](#)

nontrivial subgroup

- [4.1: Introduction to Subgroups](#)

O

odd

[6.3: Alternating Groups](#)

onto

[1.2: Functions](#)

order

[2.5: Group Conventions and Properties](#)

[5.1: Introduction to Cyclic Groups](#)

ordered pair

[1.1: Sets](#)

P

partition

[7.1: Partitions of and Equivalence Relations on Sets](#)

permutation

[6.1: Introduction to Permutation Groups](#)

permutation group

[6.1: Introduction to Permutation Groups](#)

permutation multiplication

[6.1: Introduction to Permutation Groups](#)

pointwise addition

[2.6: Examples of Groups/Nongroups, Part II](#)

preimage

[1.2: Functions](#)

product

[1.1: Sets](#)

proper subgroup

[4.1: Introduction to Subgroups](#)

Q

quotient group

[8.3: Introduction to Factor Groups](#)

R

range

[1.2: Functions](#)

remainder

[2.6: Examples of Groups/Nongroups, Part II](#)

representative (of class)

[7.1: Partitions of and Equivalence Relations on Sets](#)

right cancellation law

[2.5: Group Conventions and Properties](#)

S

set

[1.1: Sets](#)

structural properties

[3.3: Isomorphic Groups](#)

subgroup

[4.1: Introduction to Subgroups](#)

subgroup diagram

[4.2: Subgroup Proofs and Lattices](#)

subgroup lattice

[4.2: Subgroup Proofs and Lattices](#)

subset

[1.1: Sets](#)

superset

[1.1: Sets](#)

T

trivial group

[3.1: Groups of Small Order](#)

trivial subgroup

[4.1: Introduction to Subgroups](#)

U

uncountable

[1.3: Cardinality](#)

uncountably infinite

[1.3: Cardinality](#)

union

[1.1: Sets](#)

unique group

[3.3: Isomorphic Groups](#)

GLOSSARY

S

Sample Word 1 | Sample Definition 1

Bibliography

[1] John B. Fraleigh, *A First Course in Abstract Algebra* (7th ed.), Addison Wesley, 2002.

[2] Thomas W. Judson, *Abstract Algebra: Theory and Applications*. Revised edition published under the GNU Free Documentation License, 1997 (revised 2016). <http://abstract.pugetsound.edu/aata>

[3] Ivan Niven, Herbert S. Zuckerman, and Hugh L. Montgomery. *An Introduction to the Theory of Numbers* (5th ed.), John Wiley and Sons, 1991.

DETAILED LICENSING

Overview

Title: First-Semester Abstract Algebra: A Structural Approach (Sklar)

Webpages: 61

All licenses found:

- [GNU Free Documentation License](#): 80.3% (49 pages)
- [Undeclared](#): 19.7% (12 pages)

By Page

- First-Semester Abstract Algebra: A Structural Approach (Sklar) - [GNU Free Documentation License](#)
 - Front Matter - [Undeclared](#)
 - TitlePage - [Undeclared](#)
 - InfoPage - [Undeclared](#)
 - Table of Contents - [Undeclared](#)
 - Licensing - [Undeclared](#)
 - 1: Preliminaries - [GNU Free Documentation License](#)
 - 1.1: Sets - [GNU Free Documentation License](#)
 - 1.2: Functions - [GNU Free Documentation License](#)
 - 1.3: Cardinality - [GNU Free Documentation License](#)
 - 1.4: Exercises - [GNU Free Documentation License](#)
 - 2: Groups - [GNU Free Documentation License](#)
 - 2.1: Binary Operations and Structures - [GNU Free Documentation License](#)
 - 2.2: Exercises, Part I - [GNU Free Documentation License](#)
 - 2.3: The Definition of a Group - [GNU Free Documentation License](#)
 - 2.4: Examples of Groups/Nongroups, Part I - [GNU Free Documentation License](#)
 - 2.5: Group Conventions and Properties - [GNU Free Documentation License](#)
 - 2.6: Examples of Groups/Nongroups, Part II - [GNU Free Documentation License](#)
 - 2.7: Summaries of Groups We've Seen - [GNU Free Documentation License](#)
 - 2.8: Exercises, Part II - [GNU Free Documentation License](#)
 - 3: Homomorphisms and Isomorphisms - [GNU Free Documentation License](#)
 - 3.1: Groups of Small Order - [GNU Free Documentation License](#)
 - 3.2: Definitions of Homomorphisms and Isomorphisms - [GNU Free Documentation License](#)
 - 3.3: Isomorphic Groups - [GNU Free Documentation License](#)
 - 3.4: Exercises - [GNU Free Documentation License](#)
 - 4: Subgroups - [GNU Free Documentation License](#)
 - 4.1: Introduction to Subgroups - [GNU Free Documentation License](#)
 - 4.2: Subgroup Proofs and Lattices - [GNU Free Documentation License](#)
 - 4.3: Exercises - [GNU Free Documentation License](#)
 - 5: Cyclic Groups - [GNU Free Documentation License](#)
 - 5.1: Introduction to Cyclic Groups - [GNU Free Documentation License](#)
 - 5.2: The Subgroup Lattices of Cyclic Groups - [GNU Free Documentation License](#)
 - 5.3: Exercises - [GNU Free Documentation License](#)
 - 6: Permutation and Dihedral Groups - [GNU Free Documentation License](#)
 - 6.1: Introduction to Permutation Groups - [GNU Free Documentation License](#)
 - 6.2: Symmetric Groups - [GNU Free Documentation License](#)
 - 6.3: Alternating Groups - [GNU Free Documentation License](#)
 - 6.4: Cayley's Theorem - [GNU Free Documentation License](#)
 - 6.5: Dihedral Groups - [GNU Free Documentation License](#)
 - 6.6: Exercises - [GNU Free Documentation License](#)
 - 7: The Wonderful World of Cosets - [GNU Free Documentation License](#)
 - 7.1: Partitions of and Equivalence Relations on Sets - [GNU Free Documentation License](#)
 - 7.2: Introduction to Cosets and Normal Subgroups - [GNU Free Documentation License](#)
 - 7.3: The Index of a Subgroup and Lagrange's Theorem - [GNU Free Documentation License](#)
 - 7.4: Exercises - [GNU Free Documentation License](#)

- License*
- 8: Factor Groups - *GNU Free Documentation License*
 - 8.1: Motivation - *GNU Free Documentation License*
 - 8.2: Focusing on Normal Subgroups - *GNU Free Documentation License*
 - 8.3: Introduction to Factor Groups - *GNU Free Documentation License*
 - 8.4: Exercises - *GNU Free Documentation License*
- 9: The Isomorphism Theorem - *GNU Free Documentation License*
 - 9.1: The First Isomorphism Theorem - *GNU Free Documentation License*
 - 9.2: The Second and Third Isomorphism Theorems - *GNU Free Documentation License*
 - 9.3: Exercises - *GNU Free Documentation License*
- Back Matter - *Undeclared*
 - Index - *Undeclared*
 - Glossary - *Undeclared*
 - Bibliography - *Undeclared*
 - Detailed Licensing - *Undeclared*
 - Notation - *Undeclared*
 - Solutions to Exercises - *Undeclared*

This page titled [First-Semester Abstract Algebra: A Structural Approach \(Sklar\)](#) is shared under a [GNU FDL](#) license and was authored, remixed, and/or curated by Jessica K. Sklar via [source content](#) that was edited to the style and standards of the LibreTexts platform.

Notation

The following table defines the notation used in this book. Page numbers or references refer to the first appearance of each symbol.

Symbol	Description	Location
$x^{\text{int}} S$	x is an element of S	Definition 1.1.2
$x \not\text{int} S$	x is not an element of S	Definition 1.1.2
$\emptyset$	the empty set, $\{\}$	Definition 1.1.2
$\mathbb{Z}$	the set of all integers	Example 1.1.1
$\mathbb{Q}$	the set of all rational numbers	Example 1.1.1
$\mathbb{R}$	the set of all real numbers	Example 1.1.1
$\mathbb{C}$	the set of all complex numbers	Example 1.1.1
$\mathbb{N}$	the set of all natural numbers, $\{0, 1, 2, \dots\}$	Example 1.1.1
$\mathbb{Z}^+, \mathbb{Q}^+, \mathbb{R}^+$	the set of all positive elements of $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$	Example 1.1.1
$\mathbb{Z}^-, \mathbb{Q}^-, \mathbb{R}^-$	the set of all negative elements of $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$	Example 1.1.1
$\mathbb{Z}^*, \mathbb{Q}^*, \mathbb{R}^*, \mathbb{C}^*$	the set of all nonzero elements of $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$	Example 1.1.1
$\mathbb{M}_{m \times n}(S)$	the set of all $m \times n$ matrices over S	Definition 1.1.3
$\mathbb{M}_n(S)$	the set of all $n \times n$ matrices over S	Definition 1.1.3
$A \subseteq B$	A is a subset of the B	Definition 1.1.4
$A \subsetneq B$	A is a proper subset of B	Definition 1.1.4
$P(A)$	the power set of A	Definition 1.1.5
$A \cap B$	the intersection of A and B	Definition 1.1.6
$A \cup B$	the union of A and B	Definition 1.1.6
$A - B$	the difference of A and B	Definition 1.1.6
$\bigcup_{i \text{int} I} A_i$	$\{x : x^{\text{int}} A_i \text{ for some } i^{\text{int}} I\}$	Definition 1.1.6
$\bigcap_{i \text{int} I} A_i$	$\{x : x^{\text{int}} A_i \text{ for every } i^{\text{int}} I\}$	Definition 1.1.6
$A \times B$	the direct product of A and B	Definition 1.1.7
$f : S \rightarrow T$	function f from S to T	Definition 1.2.1
$f(U)$	the image of a set U under f	Definition 1.2.1
$f^{\leftarrow}(V)$	the preimage of a set V under f	Definition 1.2.1
$f \circ g$	the composition of f with g	Definition 1.2.3
1_S	the identity function on S	Definition 1.2.3
f^{-1}	the inverse of f	Theorem 1.2.2
$ S $	the cardinality of S	Definition 1.3.1
$\langle S, * \rangle$	binary structure	Definition 2.1.1
e	the identity element in a binary structure/group	Definition 2.1.4
$\det A$	the determinant of A	Definition 2.4.1

Symbol	Description	Location
$GL(n, \mathbb{R})$	the general linear group of degree n over $\mathbb{R}$	Definition 2.4.1
I_n	the $n \times n$ identity matrix	Theorem 2.4.1
e_G	the identity element in a group G	Convention 2.5.1
a^{-1}	the inverse of a in a group	Convention 2.5.1
$-a$	the inverse of a in an abelian group	Item
$n\mathbb{Z}$	$\{nm : m \in \mathbb{Z}\}$	Example 2.6.1
$a \equiv_n b$	a is congruent to $b \bmod n$	Definition 2.6.1
$R_n(a)$	the remainder when a is divided by n	Definition 2.6.2
$+_n$	addition modulo n	Definition 2.6.3
$\mathbb{Z}_n$	the cyclic group of order n	Example 2.6.3
$\mathbb{Z}_n^\times$	$\{a \in \mathbb{Z}_n : \gcd(a, n) = 1\}$	Definition 2.6.7
F	the set of all functions from $\mathbb{R}$ to $\mathbb{R}$	Example 2.6.6
B	the set of all bijections from $\mathbb{R}$ to $\mathbb{R}$	Example 2.6.7
$Z(G)$	the center of a group G	Exercise 2.8.9
C^1	the set of all differentiable functions from $\mathbb{R}$ to $\mathbb{R}$ whose derivatives are continuous	Item 6
C^0	the set of all continuous functions from $\mathbb{R}$ to $\mathbb{R}$	Item 7
c_a	conjugation by a	Example 3.2.2
$G \simeq G'$	G is isomorphic to G'	Definition 3.3.1
$G \not\simeq G'$	G is not isomorphic to G'	Definition 3.3.1
$H \leq G$	H is a subgroup of G	Definition 4.1.1
$H \not\leq G$	H is not a subgroup of G	Definition 4.1.1
$\langle a \rangle$	the (cyclic) subgroup generated by a	Definition 5.1.2
$o(a)$	the order of element a	Definition 5.1.2
S_A	the set of all permutations on A	Definition 6.1.3
S_n	the symmetric group on n letters	Definition 6.2.1
A_n	the alternating group on n letters	Definition 6.3.2
λ_a	left multiplication by a	Definition 6.4.1
ρ_a	right multiplication by a	Definition 6.4.1
$\mapsto$	maps to	Paragraph
D_n	the dihedral group of order $2n$	Definition 6.5.1
xRy	x is related to y	Definition 7.1.2
$x \not R y$	x is not related to y	Definition 7.1.2
$[x]$	the equivalence class of x	Definition 7.1.4
$a \sim_L b$	$a^{-1}b \in H$, where $H \leq G$ is specified	Definition 7.2.1
$a \sim_R b$	Double exponent: use braces to clarify $a^{-1}b^{-1}ab \in H$ where $H \leq G$ is specified	Definition 7.2.1
$aH, a + H$	the left coset of H containing a	Definition 7.2.2
$Ha, H + a$	the right coset of H containing a	Definition 7.2.2

Symbol	Description	Location
$\Leftrightarrow$	if and only if	Note 7.2
$H \trianglelefteq G$	H is a normal subgroup of G	Definition 7.2.3
G/H	the set of all left cosets of H in G	Definition 7.2.4
$(G : H)$	$ G/H $	Definition 7.3.1
aHb	$\{ahb : h \in H\}$	Definition 8.2.1
$\text{Ker}\phi$	the kernel of ϕ	Definition 8.2.3
G/N	the factor group G/N , when $N \trianglelefteq G$	Definition 8.3.1
Ψ	the canonical epimorphism from G to G/N	Definition 8.3.3
S^1	the unit circle $\{e^{i\theta} : \theta \in \mathbb{R}\}$ in the complex plane	Paragraph

Solutions to Exercises

Exercises 1.4

1. Yes/No. For each of the following, write Y if the object described is a well-defined set; otherwise, write N. You do NOT need to provide explanations or show work for this problem.

- $\{z^{\text{int}}\mathbb{C} : |z| = 1\}$
- $\{\epsilon^{\text{int}}\mathbb{R}^+ : \epsilon \text{ is sufficiently small}\}$
- $\{q^{\text{int}}\mathbb{Q} : q \text{ can be written with denominator 4}\}$
- $\{n^{\text{int}}\mathbb{Z} : n^2 < 0\}$

Solution

- Y
- N
- Y (it's $\mathbb{Q}$)
- Y (it's $\emptyset$)

2. List the elements in the following sets, writing your answers as sets.

Example: $\{z^{\text{int}}\mathbb{C} : z^4 = 1\}$ *Solution:* $\{\pm 1, \pm i\}$

- $\{z^{\text{int}}\mathbb{R} : z^2 = 5\}$
- $\{m^{\text{int}}\mathbb{Z} : mn = 50 \text{ for some } n^{\text{int}}\mathbb{Z}\}$
- $\{a, b, c\} \times \{1, d\}$
- $P(\{a, b, c\})$

Solution

- $\{\pm \sqrt{5}\}$
- $\{\pm 50, \pm 25, \pm 10, \pm 5, \pm 2, \pm 1\}$
- $\{(a, 1), (a, d), (b, 1), (b, d), (c, 1), (c, d)\}$
- $\{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$

3. Let S be a set with cardinality $n^{\text{int}}\mathbb{N}$. Use the cardinalities of $P(\{a, b\})$ and $P(\{a, b, c\})$ to make a conjecture about the cardinality of $P(S)$. You do not need to prove that your conjecture is correct (but you should try to ensure it is correct).

Solution

$|P(\{a, b\})| = 4 = 2^2$ and $|P(\{a, b, c\})| = 8 = 2^3$; we may conjecture that when $|S| = n$, $|P(S)| = 2^n$.

4. Let $f : \mathbb{Z}^2 \rightarrow \mathbb{R}$ be defined by $f(a, b) = ab$. (Note: technically, we should write $f((a, b))$, not $f(a, b)$, since f is being applied to an ordered pair, but this is one of those cases in which mathematicians abuse notation in the interest of concision.)

- What are f 's domain, codomain, and range?
- Prove or disprove each of the following statements. (Your proofs do not need to be long to be correct!)
 - f is onto;
 - f is 1-1;
 - f is a bijection. (You may refer to parts (i) and (ii) for this part.)
- Find the images of the element $(6, -2)$ and of the set $\mathbb{Z}^- \times \mathbb{Z}^-$ under f . (Remember that the image of an element is an element, and the image of a set is a set.)
- Find the preimage of $\{2, 3\}$ under f . (Remember that the preimage of a set is a set.)

Solution

- a. f 's domain, codomain, and range are, respectively, $\mathbb{Z}^2$, $\mathbb{R}$, and $\mathbb{Z}$.
- b.
 - i. f is not onto, since, for instance, $1/2 \in \mathbb{R} - \mathbb{Z}$.
 - ii. f is not 1-1: for instance, $f(-2, 2) = -4 = f(2, -2)$.
 - iii. f is not a bijection since it's not 1-1. (It would be equally valid to answer that it's not a bijection since it's not onto, or that it's not a bijection since it's neither 1-1 nor onto.)
- c. $f(6, -2) = -12$ and $f(\mathbb{Z}^- \times \mathbb{Z}^-) = \mathbb{Z}^+$.
- d. Illegal pream-token (&)

which is the set

$$\{(1, 2), (2, 1), (-1, -2), (-2, -1), (1, 3), (3, 1), (-1, -3), (-3, -1)\}.$$

5. Let S , T , and U be sets, and let $f : S \rightarrow T$ and $g : T \rightarrow U$ be onto. Prove that $g \circ f$ is onto.

Solution

Let $u \in U$. We want to show there is an element of S that gets mapped to u by $g \circ f$.

Since $g : T \rightarrow U$ is onto, there is an element $t \in T$ such that $g(t) = u$; next, since $f : S \rightarrow T$ is onto, there is an element $s \in S$ such that $f(s) = t$.

Then $(g \circ f)(s) = g(f(s)) = g(t) = u$. Thus, $g \circ f$ is onto.

6. Let A and B be sets with $|A| = m < \infty$ and $|B| = n < \infty$. Prove that $|A \times B| = mn$.

Solution

We can list the elements of A and B as so:

$$A = \{a_1, a_2, \dots, a_m\} \text{ and } B = \{b_1, b_2, \dots, b_n\}.$$

Consider the table

Illegal pream-token (&)

Clearly this table contains mn elements, and contains each element of $A \times B$ exactly once. Therefore, $|A \times B| = mn$.

Exercise 2.2 (Part I)

1. For each of the following, write Y if the given "operation" is a well-defined binary operation on the given set; otherwise, write N. In each case in which it *isn't* a well-defined binary operation on the set, provide a *brief* explanation. You do not need to prove or explain anything in the cases in which it *is* a binary operation.

- a. $+$ on $\mathbb{C}^*$
- b. $*$ on $\mathbb{R}^+$ defined by $x * y = \log_x y$
- c. $*$ on $M_2(\mathbb{R})$ defined by $A * B = AB^{-1}$
- d. $*$ on $\mathbb{Q}^*$ defined by $z * w = z/w$

Solution

- a. N; for instance, $1, -1 \in \mathbb{C}^*$, but $1 + (-1) = 0 \notin \mathbb{C}^*$.
- b. N; for instance, $10 * 1 = \log_{10} 0.1 = -1 \notin \mathbb{R}^+$.
- c. N; for instance, the zero matrix, 0 , in $M_2(\mathbb{R})$ is not invertible, so $A * 0$ is undefined for every $A \in M_2(\mathbb{R})$.
- d. Y

2. Define $*$ on $\mathbb{Q}$ by $p * q = pq + 1$. Prove or disprove that $*$ is (a) commutative; (b) associative.

Solution

- Let $p, q \in \mathbb{Q}$. Then $p * q = pq + 1 = qp + 1$ (since multiplication is commutative on $\mathbb{Q}$), which equals $q * p$. So $*$ is commutative.
- $*$ is not associative: for instance, $1, 2, 3 \in \mathbb{Q}$ and $1 * (2 * 3) = 1 * 7 = 8$, while $(1 * 2) * 3 = 3 * 3 = 10 \neq 8$.

3. Prove that matrix multiplication is not commutative on $\mathbb{M}_2(\mathbb{R})$.

Solution

Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ in $\mathbb{M}^2(\mathbb{R})$. Then $AB = B$ while BA is the zero matrix. Since $AB \neq BA$, matrix multiplication isn't commutative on $\mathbb{M}^2(\mathbb{R})$.

4. Prove or disprove each of the following statements.

- The set $2\mathbb{Z} = \{2x : x \in \mathbb{Z}\}$ is closed under addition in $\mathbb{Z}$.
- The set $S = \{1, 2, 3\}$ is closed under multiplication in $\mathbb{R}$.
- The set

$$U = \left\{ \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}$$

is closed under multiplication in $\mathbb{M}_2(\mathbb{R})$. (Recall that U is the set of *upper-triangular matrices* in $\mathbb{M}_2(\mathbb{R})$.)

Solution

- Let $x, y \in 2\mathbb{Z}$. Then there exist $a, b \in \mathbb{Z}$ such that $x = 2a$ and $y = 2b$. Then $x + y = 2a + 2b = 2(a + b) \in 2\mathbb{Z}$.

So $2\mathbb{Z}$ is closed under addition.

- Since, for instance, $2, 3 \in S$ but $2(3) = 6 \notin S$, S isn't closed under multiplication.

- Let $A = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}$ and $B = \begin{bmatrix} \alpha & \beta \\ 0 & \gamma \end{bmatrix}$ in U . Then $AB = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \begin{bmatrix} \alpha & \beta \\ 0 & \gamma \end{bmatrix} = \begin{bmatrix} a\alpha & a\beta + b\gamma \\ 0 & c\gamma \end{bmatrix} \in U$

So U is closed under matrix multiplication.

5. Let $*$ be an associative and commutative binary operation on a set S . An element $u \in S$ is said to be an *idempotent* in S if $u * u = u$. Let H be the set of all idempotents in S . Prove that H is closed under $*$.




Figure 2.2.1: ©Bill Griffith. Reprinted with permission.

Solution

Let $u, v \in H$. To show $u * v \in H$, we need to show that $(u * v) * (u * v) = u * v$. Now,

Illegal pream-token (&)

So $u * v \in H$.

Exercise 2.8 (Part II)

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem.

- For every positive integer n , there exists a group of order n .
- For every integer $n \geq 2$, $\mathbb{Z}_n$ is abelian.
- Every abelian group is finite.
- For every integer m and integer $n \geq 2$, there exist infinitely many integers a such that a is congruent to m modulo n .
- A binary operation $*$ on a set S is commutative if and only if there exist $a, b \in S$ such that $a * b = b * a$.
- If $\langle S, * \rangle$ is a binary structure, then the elements of S must be numbers.
- If e is an identity element of a binary structure (not necessarily a group) $\langle S, * \rangle$, then e is an idempotent in S (that is, $e * e = e$).
- If s is an idempotent in a binary structure (not necessarily a group) $\langle S, * \rangle$, then s must be an identity element of S .

Solution

- T
- T
- F
- T
- F
- F
- T
- F

2. Let G be the set of all functions from $\mathbb{Z}$ to $\mathbb{R}$. Prove that pointwise multiplication on G (that is, the operation defined by $(fg)(x) = f(x)g(x)$ for all $f, g \in G$ and $x \in \mathbb{Z}$) is commutative. (Note. To prove that two functions, h and j , sharing the same domain D are equal, you need to show that $h(x) = j(x)$ for every $x \in D$.)

Solution

Let $f, g \in G$. Then for every $x \in \mathbb{Z}$,

$$(fg)(x) = f(x)g(x) = g(x)f(x),$$

since $f(x), g(x) \in \mathbb{R}$ and multiplication of real numbers is commutative. Since $g(x)f(x) = (gf)(x)$, $fg = gf$, and so pointwise multiplication on G is commutative.

3. Decide which of the following binary structures are groups. For each, if the binary structure *isn't* a group, prove that. (Remember, you should *not* state that inverses do or do not exist for elements until you have made sure that the structure contains an identity element!) If the binary structure *is* a group, prove that.

- $\mathbb{Q}$ under multiplication
- $M_2(\mathbb{R})$ under addition
- $M_2(\mathbb{R})$ under multiplication
- $\mathbb{R}^+$ under $*$, defined by $a * b = \sqrt{ab}$ for all $a, b \in \mathbb{R}^+$

Solution

- $\mathbb{Q}$ isn't a group under multiplication since $0 \in \mathbb{Q}$ has no inverse.
- Matrix multiplication is always associative, and the zero matrix in $M^2(\mathbb{R})$ acts as an additive identity element. Finally, let $A \in M^2(\mathbb{R})$. Then $-A \in M^2(\mathbb{R})$ is an inverse for A under addition. Thus, $M^2(\mathbb{R})$ is a group under addition.
- $M^2(\mathbb{R})$ isn't a group under multiplication since the zero matrix has no inverse.
- $1 * (4 * 9) = 1 * 6 = \sqrt{6}$ while $(1 * 4) * 9 = 2 * 9 = \sqrt{18}$, so $*$ isn't associative. Thus, $\mathbb{R}^+$ isn't a group under $*$.

4. Give an example of an abelian group containing 711 elements.

Solution

$\mathbb{Z}_{711}$. (Other answers are possible.)

5. Let $n \in \mathbb{Z}^+$. Prove that $n\mathbb{Z}$ is a group under the usual addition of integers. *Note:* You may use the fact that $\langle n\mathbb{Z}, + \rangle$ is a binary structure if you provide a reference for this fact.

Solution

We know from Theorem 2.6.1 that $\langle n\mathbb{Z}, + \rangle$ is a binary structure.

Next, $+$ is associative on n , since it's associative on $\mathbb{Z}$, and $n\mathbb{Z} \subseteq \mathbb{Z}$.

Notice that $0 \in n\mathbb{Z}$ (since $0 = n(0)$); 0 then clearly acts as an identity element for $+$ in $n\mathbb{Z}$.

Finally, let $x \in n\mathbb{Z}$, then $x = nm$ for some $m \in \mathbb{Z}$, so $-x = n(-m)$ for $m \in \mathbb{Z}$, implying $-x \in n\mathbb{Z}$; and clearly $-x$ acts as an inverse for x in $n\mathbb{Z}$.

Thus, $\langle n\mathbb{Z}, + \rangle$ is a group.

6. Let $n \in \mathbb{Z}^+$. Prove that $SL(n, \mathbb{R})$ is a group under matrix multiplication. *Note:* You may use the fact that $\langle SL(n, \mathbb{R}), \cdot \rangle$ is a binary structure if you provide a reference for this fact.

Solution

We know from Theorem 2.4.2 that $\langle SL(n, \mathbb{R}), \cdot \rangle$ is a binary structure, and we know that matrix multiplication is always associative. Next, since $\det I_n = 1$, I_n is in $SL(n, \mathbb{R})$, and clearly acts as an identity element in $SL(n, \mathbb{R})$. Finally, let $A \in SL(n, \mathbb{R})$. Since $\det A = 1 \neq 0$, A has an inverse matrix A^{-1} in $GL(n, \mathbb{R})$. A^{-1} is in $SL(n, \mathbb{R})$ since

$$\det(A^{-1}) = \frac{1}{\det A} = \frac{1}{1} = 1.$$

So A has an inverse in $SL(n, \mathbb{R})$.

Thus, our proof is complete.

7.

- a. List three distinct integers that are congruent to 6 modulo 5.
- b. List the elements of $\mathbb{Z}_5$.
- c. Compute:
 - i. $4 + 5$ in $\mathbb{Z}$;
 - ii. $4 + 5$ in $\mathbb{Q}$;
 - iii. $4 +_6 5$ in $\mathbb{Z}_6$;
 - iv. the inverse of 4 in $\mathbb{Z}$;
 - v. the inverse of 4 in $\mathbb{Z}_6$.
- d. Why does it not make sense for me to ask you to compute $4 +_3 2$ in $\mathbb{Z}_3$? *Please answer this using a complete, grammatically correct sentence.*

Solution

- a. 11, 1, -4 . (Other answers are possible.)
- b. $\mathbb{Z}_5 = \{0, 1, 2, 3, 4\}$
- c. Compute:
 - i. 9
 - ii. 9
 - iii. 3
 - iv. -4
 - v. 2
- d. It doesn't make sense because $4 \notin \mathbb{Z}_3$.

8. Let G be a group with identity element e . Prove that if every element of G is its own inverse, then G is abelian.

Solution

Let $a, b \in G$. Then

Illegal pream-token (&)

since a, b are their own inverses.

So G is abelian.

9. Let G be a group. The subset

$$Z(G) := \{z \in G : zg = gz \text{ for all } g \in G\}$$

of G is called the *center* of G . In other words, $Z(G)$ is the set of all elements of G that commute with every element of G . Prove that $Z(G)$ is closed in G .

Solution

Let $z_1, z_2 \in Z(G)$. Let $z = z_1 z_2$; we want to show that z is in Z : that is, we want to show that for every $g \in G$, $zg = gz$. But for every $g \in G$,

Illegal pream-token (&)

Thus, $Z(G)$ is closed in G .

Exercise 3.4

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G and G' be groups.

- If there exists a homomorphism $\phi : G \rightarrow G'$, then G and G' must be isomorphic groups.
- There is an integer $n \geq 2$ such that $\mathbb{Z} \simeq \mathbb{Z}_n$.
- If $|G| = |G'| = 3$, then we must have $G \simeq G'$.
- If $|G| = |G'| = 4$, then we must have $G \simeq G'$.

Solution

- F
- F
- T
- F

2. For each of the following functions, prove or disprove that the function is (i) a homomorphism; (ii) an isomorphism. (Remember to work with the default operation on each of these groups!)

- The function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(n) = 2n$.
- The function $g : \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) = x^2$.
- The function $h : \mathbb{Q}^* \rightarrow \mathbb{Q}^*$ defined by $h(x) = x^2$.

Solution

- The function f is a homomorphism, since for every $a, b \in \mathbb{Z}$, we have $f(a + b) = 2(a + b) = 2a + 2b = f(a) + f(b)$.
 - The function f isn't a bijection, since it isn't onto: for instance, there is no element a in $\mathbb{Z}$ such that $f(a) = 3$. Thus, f is not an isomorphism.
- The function g isn't a homomorphism: for instance, we have $g(2 + 3) = g(5) = 25 \neq 13 = 4 + 9 = g(2) + g(3)$.
 - Since it isn't a homomorphism, it isn't an isomorphism.
- The function h is a homomorphism, since for every $a, b \in \mathbb{Q}$, we have $h(ab) = (ab)^2 = a^2b^2 = h(a)h(b)$.
 - The function h isn't onto: its range is the set of nonnegative rational numbers, not all of $\mathbb{Q}$. Thus, h is not an isomorphism.

3. Define $d : GL(2, \mathbb{R}) \rightarrow \mathbb{R}^*$ by $d(A) = \det A$. Prove/disprove that d is:

- a homomorphism
- 1-1
- onto
- an isomorphism.

Solution

- The function d is a homomorphism, since for every $A, B \in GL(2, \mathbb{R})$, we have $d(AB) = \det(AB) = (\det A)(\det B) = d(A)d(B)$.
- d isn't 1-1: For instance, $d(I_2) = d(-I_2)$.
- d is onto: Let $a \in \mathbb{R}^*$. Then the matrix $A = \begin{bmatrix} a & 0 \\ 0 & 1 \end{bmatrix}$

is in $GL(2, \mathbb{R})$, with $d(A) = a$.

- Since d isn't 1-1, it isn't an isomorphism.

4. Complete the group tables for $\mathbb{Z}_4$ and $\mathbb{Z}_8^\times$. Use the group tables to decide whether or not $\mathbb{Z}_4$ and $\mathbb{Z}_8^\times$ are isomorphic to one another. (You do not need to provide a proof.)

Solution

The group tables of $\mathbb{Z}_4$ and $\mathbb{Z}_8^\times$ are, respectively,

+	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

and

$\cdot_8$	1	3	5	7
1	1	3	5	7
3	3	5	7	1
5	5	7	1	3
7	7	1	3	5

We can see from the group table for $\mathbb{Z}_8^\times$ that every element of that group is its own inverse; that is not the case in $\mathbb{Z}_4$. Thus, $\mathbb{Z}_4 \not\cong \mathbb{Z}_8^\times$.

5. Let $n^{\text{int}}\mathbb{Z}^+$. Prove that $\langle n\mathbb{Z}, + \rangle \simeq \langle \mathbb{Z}, + \rangle$.

Solution

Define $\phi : \mathbb{Z} \rightarrow n\mathbb{Z}$ by $\phi(x) = nx$, for all $x \in \mathbb{Z}$. Clearly, ϕ is a bijection. Moreover, for all $x, y \in \mathbb{Z}$,

$$\phi(x + y) = n(x + y) = nx + ny = \phi(x) + \phi(y);$$

so ϕ is a homomorphism. Thus, ϕ is an isomorphism, and hence

$$\langle \mathbb{Z}, + \rangle \simeq \langle n\mathbb{Z}, + \rangle.$$

6.

- Let G and G' be groups, where G is abelian and $G \simeq G'$. Prove that G' is abelian.
- Give an example of groups G and G' , where G is abelian and there exists a homomorphism from G to G' , but G' is NOT abelian.

Solution

- Since $G \simeq G'$, there exists some isomorphism $\phi : G \rightarrow G'$. Let $x, y \in G'$. Since ϕ is onto, there exist $a, b \in G$ such that $\phi(a) = x$ and $\phi(b) = y$. So $\phi(a + b) = \phi(a) + \phi(b) = x + y$. Thus, G' is abelian.

Thus, G' is abelian.

- Let $G = \{I_2\}$ (under multiplication) and let $G' = GL(2, \mathbb{R})$. Then the map $\phi : G \rightarrow G'$ defined by $\phi(I_2) = I_2$ is clearly a homomorphism, but note that G is abelian while G' is not.

7. Let $\langle G, \cdot \rangle$ and $\langle G', \cdot' \rangle$ be groups with identity elements e and e' , respectively, and let ϕ be a homomorphism from G to G' . Let $a^{\text{int}}G$. Prove that $\phi(a)^{-1} = \phi(a^{-1})$.

Solution

We omit the group operations in this solution in order to increase familiarity with the convention.

We want to show that $\phi(a)^{-1} = \phi(a^{-1})$. Notice that

Illegal pream-token (&)

Thus, $\phi(a^{-1}) = \phi(a)^{-1}$, by right cancellation.

Exercise 4.3

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G and G' be groups.

- Every group contains at least two distinct subgroups.
- If H is a proper subgroup of group G and G is finite, then we must have $|H| < |G|$.
- $7\mathbb{Z}$ is a subgroup of $14\mathbb{Z}$.
- A group G may have two distinct proper subgroups which are isomorphic (to one another).

Solution

- F
- T
- F
- T

2. Give specific, precise examples of the following groups G with subgroups H :

- A group G with a proper subgroup H of G such that $|H| = |G|$.
- A group G of order 12 containing a subgroup H with $|H| = 3$.
- A nonabelian group G containing a nontrivial abelian subgroup H .
- A finite subgroup H of an infinite group G .

Solution

(Other answers are possible.)

- $G = \mathbb{Z}$, $H = 2\mathbb{Z}$
- $G = \mathbb{Z}_{12}$, $H = \{0, 4, 8\}$
- $G = GL(2, \mathbb{R})$, $H = \{\pm I_2\}$
- $G = \mathbb{R}^*$, $H = \{\pm 1\}$.

3. Let $n^{\text{int}}\mathbb{Z}^+$.

- Prove that $n\mathbb{Z} \leq \mathbb{Z}$.
- Prove that the set $H = \{A^{\text{int}}\mathbb{M}_n(\mathbb{R}) : \det A = \pm 1\}$ is a subgroup of $GL(n, \mathbb{R})$.

(Note: Your proofs do not need to be long to be correct!)

Solution

- We know that $n\mathbb{Z} \subseteq \mathbb{Z}$ and that $n\mathbb{Z}$ is a group under addition, the group operation in $\mathbb{Z}$. Therefore, $n\mathbb{Z} \leq \mathbb{Z}$.
- Clearly, H is a nonempty subset of $GL(n, \mathbb{R})$ (for instance, $I_n \in H$). Next, let $A, B \in H$. Then $\det(AB^{-1}) = (\det A)(\det(B^{-1})) = (\pm 1)(\pm 1) = \pm 1$,

so $AB^{-1} \in H$. Thus, $H \leq GL(n, \mathbb{R})$ by the Two-Step Subgroup Test.

4. Let $n^{\text{int}}\mathbb{Z}^+$. For each group G and subset H , decide whether or not H is a subgroup of G . In the cases in which H is *not* a subgroup of G , provide a proof. (Note. Your proofs do not need to be long to be correct!)

- $G = \mathbb{R}, H = \mathbb{Z}$
- $G = \mathbb{Z}_{15}, H = \{0, 5, 10\}$
- $G = \mathbb{Z}_{15}, H = \{0, 4, 8, 12\}$
- $G = \mathbb{C}, H = \mathbb{R}^*$
- $G = \mathbb{C}^*, H = \{1, i, -1, -i\}$
- $G = M_n(\mathbb{R}), H = GL(n, \mathbb{R})$
- $G = GL(n, \mathbb{R}), H = \{A^{\text{int}} M_n(\mathbb{R}) : \det A = -1\}$

Solution

- $H \leq G$.
- $H \leq G$.
- H is a subset of G but isn't closed under $+$: for instance $12 +_{15} 4 = 1 \notin H$. So $H \not\leq G$.
- H is a subset of G , but the identity element, 0, of G isn't in H , so $H \not\leq G$.
- $H \leq G$.
- H is a subset of G , but the identity element, 0, of G isn't in H , so $H \not\leq G$.
- H is a subset of G , but the identity element, I_n , of G isn't in H , since $\det I_n = 1$. Thus, $H \not\leq G$.

5. Let G and G' be groups, let ϕ be a homomorphism from G to G' , and let H be a subgroup of G . Prove that $\phi(H)$ is a subgroup of G' .

Solution

Let $x, y \in \phi(H)$. Then $x = \phi(a)$ and $y = \phi(b)$ for some $a, b \in H$. So $xy = \phi(a)\phi(b) = \phi(ab)$ (since ϕ is a homomorphism). Since $H \leq G$, $ab \in H$. Thus, $xy \in \phi(H)$. Next, $e_G \in H$, so $e_G' = \phi(e_G) \in \phi(H)$. Finally, since $H \leq G$ and $a \in H$, $a^{-1} \in H$; so $x^{-1} = \phi(a)^{-1} = \phi(a^{-1}) \in \phi(H)$. Thus, $\phi(H) \leq G'$.

6. Let G be an abelian group, and let $U = \{g^{\text{int}}G : g^{-1} = g\}$. Prove that U is a subgroup of G .

Solution

Clearly, $U \subseteq G$ and $e \in U$. Next, let $u, v \in U$. Then

$$(uv)^2 = (uv)(uv) = u^2v^2 = uv,$$

so $uv \in U$. Finally, since $u \in U$, we have $u^{-1} = u$, so $(u^{-1})^2 = u^2 = e$; thus, $u^{-1} \in U$.

Exercise 5.3

1. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G be a group with identity element e .

- If G is infinite and cyclic, then G must have infinitely many generators.
- There may be two distinct elements a and b of a group G with $\langle a \rangle = \langle b \rangle$.
- If $a, b^{\text{int}}G$ and $a^{\text{int}}\langle b \rangle$ then we must have $b^{\text{int}}\langle a \rangle$.
- If $a^{\text{int}}G$ with $a^4 = e$, then $o(a)$ must equal 4.
- If G is countable then G must be cyclic.

Solution

- F

- b. T
- c. F
- d. F
- e. F

2. Give examples of the following.

- a. An infinite noncyclic group G containing an infinite cyclic subgroup H .
- b. An infinite noncyclic group G containing a finite nontrivial cyclic subgroup H .
- c. A cyclic group G containing exactly 20 elements.
- d. A nontrivial cyclic group G whose elements are all matrices.
- e. A noncyclic group G such that every proper subgroup of G is cyclic.

Solution

(Other answers are possible.)

- a. $G = \mathbb{R}, H = \mathbb{Z}$
- b. $G = GL(n, \mathbb{R}), H = \{\pm I_2\}$
- c. $G = \mathbb{Z}_{20}$
- d. $G = \{\pm I_2\}$, under matrix multiplication
- e. $G = \mathbb{Z}_2^2$

3. Find the orders of the following elements in the given groups.

- a. $2^{\text{int}} \mathbb{Z}$
- b. $-i^{\text{int}} \mathbb{C}^*$
- c. $-I_2^{\text{int}} GL(2, \mathbb{R})$
- d. $-I_2^{\text{int}} M_2(\mathbb{R})$
- e. $(6, 8)^{\text{int}} \mathbb{Z}_{10} \times \mathbb{Z}_{10}$

Solution

- a. $o(2) = \infty$, since $\langle 2 \rangle = 2\mathbb{Z}$.
- b. $o(-i) = 4$, since $\langle -i \rangle = \{-i, -1, i, 1\}$.
- c. $o(I_2) = 2$, since $\langle -I_2 \rangle = \{-I_2, I_2\}$.
- d. $o(I_2) = \infty$, since $\langle -I_2 \rangle = \{kI_2 : k \in \mathbb{Z}\}$.
- e. $o((6, 8)) = 5$, since $\langle (6, 8) \rangle = \{(6, 8), (2, 6), (8, 4), (4, 2), (0, 0)\}$.

4. For each of the following, if the group is cyclic, list *all* of its generators. If the group is *not* cyclic, write NC.

- a. $5\mathbb{Z}$
- b. $\mathbb{Z}_{18}$
- c. $\mathbb{R}$
- d. $\langle \pi \rangle$ in $\mathbb{R}$
- e. $\mathbb{Z}_2^2$
- f. $\langle 8 \rangle$ in $\mathbb{Q}^*$

Solution

- a. ± 5
- b. 1, 5, 7, 11, 13, 17
- c. NC
- d. $\pm \pi$
- e. NC
- f. $8, \frac{1}{8}$

5. Explicitly identify the elements of the following subgroups of the given groups. You may use set-builder notation if the subgroup is infinite, or a conventional name for the subgroup if we have one.

- $\langle 3 \rangle$ in $\mathbb{Z}$
- $\langle i \rangle$ in C^*
- $\langle A \rangle$, for $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}^{\text{int}} \mathbb{M}_2(\mathbb{R})$
- $\langle (2, 3) \rangle$ in $\mathbb{Z}_4 \times \mathbb{Z}_5$
- $\langle B \rangle$, for $B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{\text{int}} GL(2, \mathbb{R})$

Solution

- $3\mathbb{Z}$
- $\{i, -1, -i, 1\}$
- $\left\{ \begin{bmatrix} k & 0 \\ 0 & 0 \end{bmatrix} : k \in \mathbb{Z} \right\}$
- $\{(2, 3), (0, 1), (2, 4), (0, 2), (2, 0), (0, 3), (2, 1), (0, 4), (2, 2), (0, 0)\}$
- $\left\{ \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} : k \in \mathbb{Z} \right\}$


6. Draw subgroup lattices for the following groups.

- $\mathbb{Z}_6$
- $\mathbb{Z}_{13}$
- $\mathbb{Z}_{18}$

Solution

The respective solutions are:


a.



b.



c.



7. Let G be a group with no nontrivial proper subgroups. Prove that G is cyclic.

Solution

Let e be the identity element of G . If $G = \{e\}$, then G is clearly cyclic. Else, there exists $a \neq e$ in G . Then $\langle a \rangle$ is a subgroup of G . Since $e \neq a \in \langle a \rangle$, $\langle a \rangle$ is a nontrivial subgroup of G . Therefore, we must have $\langle a \rangle = G$. Hence, G is cyclic, as desired.

Exercise 6.6

1. Let $\sigma = (134)$, $\tau = (23)(145)$, $\rho = (56)(78)$, and $\alpha = (12)(145)$ in S_8 . Compute the following.

- $\sigma\tau$
- $\tau\sigma$
- τ^2
- τ^{-1}
- $o(\tau)$
- $o(\rho)$
- $o(\alpha)$
- $\langle \tau \rangle$

Solution

- $\sigma\tau = (134)(23)(145) = (2453)$

- b. $\tau\sigma = (23)(145)(134) = (1235)$
- c. $\tau^2 = (154)$
- d. $\tau - 1 = (23)(154)$
- e. $o(\tau) = \text{lcm}(2, 3) = 6$ (since (23) and (145) are disjoint)
- f. $o(\rho) = \text{lcm}(2, 2) = 2$
- g. In disjoint cycle notation, $\alpha = (1452)$, so $o(\alpha) = o(1452) = 4$.
- h. $\langle \tau \rangle = (23)(145), (154), (23), (145), e$

2. Prove Lemma 6.3.1.

Solution

Let $\sigma = (a_1 a_2 \cdots a_k) \in S_n$. Then $\sigma = (a_1 a_k)(a_1 a_{k-1}) \cdots (a_1 a_3)(a_1 a_2)$, so is a product of $k - 1$ transpositions. Thus, σ is even if $k - 1$ is even, that is, if k is odd, and odd if $k - 1$ is odd, that is, if k is even.

3. Prove that A_n is a subgroup of S_n .

Solution

Since $e \in A_n$, $A_n \neq \emptyset$. Next, let $\sigma, \tau \in A_n$. Then σ can be written as a product of k transpositions and τ can be written as $\tau = \tau_1 \tau_2 \cdots \tau_m$, where k and m are even and the τ_i are transpositions. Since each τ_i is a transposition, each τ_i is its own inverse; so

$$\tau^{-1} = (\tau_m)^{-1}(\tau_{m-1})^{-1} \cdots \tau_1^{-1} = \tau_m^{-1} \tau_{m-1}^{-1} \cdots \tau_1^{-1}.$$

Thus, $\sigma\tau^{-1}$ is a product of $k + m$ transpositions; since k and m are even, $k + m$ is even, and so $\sigma\tau^{-1} \in A_n$. Thus, $A_n \leq S_n$, by the Two-Step Subgroup Test.

4. Prove or disprove: The set of all odd permutations in S_n is a subgroup of S_n .

Solution

This set is *not* a subgroup of S_n because, for instance, it doesn't contain e .

5. Let n be an integer greater than 2. $m^{\text{int}}\{1, 2, \dots, n\}$, and let $H = \{\sigma^{\text{int}} S_n : \sigma(m) = m\}$ (in other words, H is the set of all permutations in S_n that fix m).

- a. Prove that $H \leq S_n$.
- b. Identify a familiar group to which H is isomorphic. (You do not need to show any work.)

Solution

- a. Since $e \in H$, $H \neq \emptyset$. Next, let $\sigma, \tau \in H$. Since $\tau(m) = m$ and τ is a bijection, $\tau^{-1}(m) = m$. So $\sigma\tau^{-1}(m) = \sigma(\tau^{-1}(m)) = \sigma(m) = m$. So $\sigma\tau^{-1} \in H$; therefore, $H \leq S_n$, by the Two-Step Subgroup Test.
- b. S_{n-1} .

6. Write $rf r^2 f r f r$ in D_5 in standard form.

Solution

In D_5 , $f r f r = (f r)^2 = e$ and $r f = f r^{-1}$, so

$$r f r^2 f r f r = r f r^2 = f r^{-1} r^2 = f r.$$

7. Prove or disprove: $D_6 \simeq S_6$.

Solution

$|D_6| = 2(6) = 12$ while $|S_6| = 720$, so $D_6 \not\cong S_6$.

8. Which elements of D_4 (if any)

- have order 2?
- have order 3?

Solution

We know $o(fr^i) = 2$ for every i , and $o(e) = 1$. Moreover, since 4 is even, $r^{4/2} = r^2$ has order 2. Finally, $o(r) = o(r^{-1}) = 4$. So (a) the elements of order 2 in D_4 are f, fr, fr^2, fr^3 , and r^2 , and (b) D_4 contains no elements of order 3.

9. Let n be an even integer that's greater than or equal to 4. Prove that **Double exponent: use braces to clarify** that is, prove that $r^{n/2}$ commutes with every element of D_n . (Do NOT simply refer to the last statement in Theorem 6.5.4; that is the statement you are proving here.)

Solution

Clearly, $r^{n/2}$ commutes with every r^k for $0 \leq k \leq n$. Moreover, by the first statement of Theorem 6.5.4,

$$r^{n/2}f = f(r^{n/2})^{-1} = fr^{n-n/2} = fr^{n/2}.$$

So $r^{n/2}$ commutes with f . It clearly follows that $r^{n/2} \in \mathbb{Z}(D_n)$.

Exercise 7.4

1. How many distinct partitions of the set $S = \{a, b, c, d\}$ are there? You do not need to list them. (Yes, you can find this answer online. But I recommend doing the work yourself for practice working with partitions!)

Solution

There are 15 distinct partitions of the set $\{a, b, c, d\}$:

$$\{\{a\}, \{b\}, \{c\}, \{d\}\}, \{\{a, b\}, \{c\}, \{d\}\}, \{\{a, c\}, \{b\}, \{d\}\}, \{\{a, d\}, \{b\}, \{c\}\}, \{\{b, c\}, \{a\}, \{d\}\}, \{\{b, d\}, \{a\}, \{c\}\}, \{\{c, d\}, \{a\}, \{b\}\}, \\ \{\{a, b\}, \{c, d\}\}, \{\{a, c\}, \{b, d\}\}, \{\{a, d\}, \{b, c\}\}, \{\{a, b, c\}, \{d\}\}, \{\{a, b, d\}, \{c\}\}, \{\{a, c, d\}, \{b\}\}, \{\{b, c, d\}, \{a\}\}, \{\{a, b, c, d\}\}.$$

2.

- Let $n^{\text{int}}\mathbb{Z}^+$. Prove that $\equiv_n$ is an equivalence relation on $\mathbb{Z}$.
- The cells of the induced partition of $\mathbb{Z}$ are called the residue classes (or congruence classes) of $\mathbb{Z}$ modulo n . Using set notation of the form $\{\dots, \#, \#, \#, \dots\}$ for each class, write down the residue classes of $\mathbb{Z}$ modulo 4.

Solution

- First, for every $x \in \mathbb{Z}$, n divides $0 = x - x$; so $x \equiv_n x$ for every $x \in \mathbb{Z}$. Thus, $\equiv_n$ is reflexive. Next, let $x, y \in \mathbb{Z}$ with $x \equiv_n y$. Then there exists some $q \in \mathbb{Z}$ with $x - y = nq$; so $y - x = n(-q)$. Since $-q \in \mathbb{Z}$, this shows us that $y \equiv_n x$. Hence, $\equiv_n$ is symmetric. Finally, let x, y and z be in $\mathbb{Z}$ with $x \equiv_n y$ and $y \equiv_n z$. Then there exist $q_1, q_2 \in \mathbb{Z}$ with $x - y = nq_1$ and $y - z = nq_2$. Thus, $x - z = (x - y) + (y - z) = nq_1 + nq_2 = n(q_1 + q_2)$;

since $q_1 + q_2 \in \mathbb{Z}$, this shows us that $x \equiv_n z$. Thus, $\equiv_n$ is transitive. Hence, $\equiv_n$ is an equivalence relation, as desired.

$$\text{b. } \{\dots, -8, -4, 0, 4, \dots\}, \{\dots, -7, -3, 1, 5, \dots\}, \{\dots, -6, -2, 2, 6, \dots\}, \{\dots, -5, -1, 3, 7, \dots\}$$

3. Let G be a group with subgroup H . Prove that $\sim_R$ is an equivalence relation on G .

Solution

First, let $a \in G$. Then $aa^{-1} = e \in H$, so $a \sim_R a$. Thus, $\sim_R$ is reflexive.

Next, let $a, b \in G$ with $asim_R b$. Then $ab^{-1} \in H$, so, since H is a subgroup of G , $(ab^{-1})^{-1} \in H$. But $(ab^{-1})^{-1} = ba^{-1}$; thus, $b \sim_R a$, and so $\sim_R$ is symmetric.

Finally, let $a, b, c \in G$ with $a \sim_R b$ and $b \sim_L c$. Then ab^{-1} and bc^{-1} are in H . Since H is a subgroup of G , we must then have $(ab^{-1})(bc^{-1}) \in H$; but $(ab^{-1})(bc^{-1})$ equals ac^{-1} . Thus, $a \sim_R c$, and so $\sim_R$ is transitive.

Hence, $\sim_R$ is an equivalence relation on G , as desired.

4. Find the indices of:

- $H = \langle (15)(24) \rangle$ in S_5
- $K = \langle (2354)(34) \rangle$ in S_6
- A_n in S_n

Solution

- $(S_5 : H) = \frac{|S_5|}{|H|} = \frac{120}{2} = 60$.
- In disjoint cycle notation, the permutation $(2354)(34)$ is written $(23)(45)$, so L has order 2. Thus, $(S_6 : K) = \frac{|S_6|}{|K|} = \frac{720}{2} = 360$.
- $(S_n : A_n) = 2$.

5. For each subgroup H of group G , (i) find the left and the right cosets of H in G , (ii) decide whether or not H is normal in G , and (iii) find $(G : H)$.

Write all permutations using *disjoint cycle notation*, and write all dihedral group elements using *standard form*.

- $H = 6\mathbb{Z}$ in $G = 2\mathbb{Z}$
- $H = \langle 4 \rangle$ in $\mathbb{Z}_{20}$
- $H = \langle (23) \rangle$ in $G = S_3$
- $H = \langle r \rangle$ in $G = D_4$
- $H = \langle f \rangle$ in $G = D_4$

Solution

- The left and the right cosets of $6\mathbb{Z}$ in $2\mathbb{Z}$ are $6\mathbb{Z}$, $2 + 6\mathbb{Z} = 6\mathbb{Z} + 2$, and $4 + 6\mathbb{Z} = 6\mathbb{Z} + 4$.
 - $6\mathbb{Z} \trianglelefteq 2\mathbb{Z}$.
 - $(2\mathbb{Z} : 6\mathbb{Z}) = 3$.
- The left and the right cosets of $H = \langle 4 \rangle = \{0, 4, 8, 12, 16\}$ in $\mathbb{Z}_{20}$ are H , $1 + H = \{1, 5, 9, 13, 17\} = H + 1$, $2 + H = \{2, 6, 10, 14, 18\} = H + 2$, and $3 + H = \{3, 7, 11, 15, 19\} = H + 3$.
 - $H \trianglelefteq \mathbb{Z}_{20}$.
 - $(\mathbb{Z}_{20} : H) = 4$.
- The left cosets of H in S_3 are $H = \{e, (23)\}$, $(12)H = \{(12), (123)\}$, and $(13)H = \{(13), (132)\}$. The right cosets of H in S_3 are H , $H(12) = \{(12), (132)\}$, and $H(13) = \{(13), (123)\}$.
 - H is not normal in S_3 .
 - $(S_3 : H) = 3$.
- The left cosets and the right cosets of H in D_4 are $H = \{e, r, r^2, r^3\}$ and $fH = \{f, fr, fr^2, fr^3\} = Hf$.
 - $H \trianglelefteq D_4$.
 - $(D_4 : H) = 2$.
- The left cosets of H in D_4 are $H = \{e, f\}$, $rH = \{r, fr^3\}$, $r^2H = \{r^2, fr^2\}$, and $r^3H = \{r^3, fr\}$. The right cosets of H in D_4 are H , $Hr = \{r, fr\}$, $Hr^2 = \{r^2, fr^2\}$, and $Hr^3 = \{r^3, fr^3\}$.
 - H is not normal in D_4 .
 - $(D_4 : H) = 4$.

6. For each of the following, give an example of a group G with a subgroup H that matches the given conditions. If no such example exists, prove that.

- A group G with subgroup H such that $|G/H| = 1$.
- A finite group G with subgroup H such that $|G/H| = |G|$.
- An abelian group G of order 8 containing a non-normal subgroup H of order 2.
- A group G of order 8 containing a normal subgroup of order 2.
- A nonabelian group G of order 8 containing a normal subgroup of index 2.
- A group G of order 8 containing a subgroup of order 3.
- An infinite group G containing a subgroup H of finite index.
- An infinite group G containing a finite nontrivial subgroup H .

Solution

(Other answers are possible.)

- $G = H = S_3$.
- $G = S_3, H = \{e\}$.
- No such example exists, since every subgroup of an abelian group is normal.
- $G = \mathbb{Z}_8, H = \{0, 4\}$.
- $G = D_4, H = \langle r \rangle$.
- No such example exists, since by Lagrange's Theorem $|H|$ must divide $|G|$, and 3 doesn't divide 8.
- $G = \mathbb{Z}, H = 2\mathbb{Z}$.
- $G = GL(2, \mathbb{R}), H = \{\pm I_2\}$.

7. True/False. For each of the following, write T if the statement is true; otherwise, write F. You do NOT need to provide explanations or show work for this problem. Throughout, let G be a group with subgroup H and elements $a, b \in G$.

- If $a \in H$ then aH must equal bH .
- aH must equal Ha .
- If $aH = bH$ then Ha must equal Hb .
- If $a \in H$ then aH must equal Ha .
- H must be normal in G if there exists $a \in G$ such that $aH = Ha$.
- If $aH = bH$ then $ah = bh$ for every $h \in H$.
- $|G/H|$ must be less than $|G|$.
- $(G : H)$ must be less than or equal to $|G|$.

Solution

- T
- F
- F
- T
- F
- F
- F
- T

8. Let G be a group of order pq , where p and q are prime, and let H be a proper subgroup of G . Prove that H is cyclic.

Solution

By Lagrange's Theorem, $|H|$ divides pq , and since H is proper, $|H| \neq pq$. Thus, $|H| = 1, p$, or q . Since the order of H is either 1 or a prime number, H must be cyclic.

9. Prove Corollary 7.3.2: that is, let G be a group of prime order, and prove that G is cyclic.

Solution

Since $|G|$ is prime, $|G| > 1$, so there exists $a \in G$ with $a \neq e$. Then $\langle a \rangle$ is a subgroup of G , so by Lagrange's Theorem, $|\langle a \rangle|$ divides $|G|$. But $|G|$ is prime and $a \neq e$, so we must have $|\langle a \rangle| = |G|$; thus, $G = \langle a \rangle$.

10. Let G be a group of finite order n , containing identity element e . Let $a^{\text{int}} G$. Prove that $a^n = e$.

Solution

By Lagrange's Theorem, $o(a) = |\langle a \rangle|$ divides n ; that is, $n = o(a)m$ for some $m \in \mathbb{Z}^+$. Thus,

$$a^n = a^{o(a)m} = (a^{o(a)})^m = e^m = e.$$

Exercise 8.4

1. Let G be a group and let $H \leq G$ have index 2. Prove that $H \trianglelefteq G$.

Solution

Let $a \in G$. If $a \in H$, then $aH = H = Ha$. On the other hand, if $a \notin H$, then $aH = G - H = Ha$, since $(G : H) = 2$. Thus, $H \trianglelefteq G$.

2. Let G be an abelian group with $N \trianglelefteq G$. Prove that G/N is abelian.

Solution

Let $aN, bN \in G/N$. Then

$$(aN)(bN) = abN = baN = (bN)(aN).$$

So G/N is abelian.

3. Find the following.

- $|2\mathbb{Z}/6\mathbb{Z}|$
- $|H|$, for $H = 2 + \langle 6 \rangle \subseteq \mathbb{Z}_{12}$
- $o(2 + \langle 6 \rangle)$ in $\mathbb{Z}_{12}/\langle 6 \rangle$
- $\langle f + H \rangle$ in D_4/H , where $H = \{e, r^2\}$
- $|(\mathbb{Z}_6 \times \mathbb{Z}_8)/(\langle 3 \rangle \times \langle 2 \rangle)|$
- $|(\mathbb{Z}_{15} \times \mathbb{Z}_{24})/\langle (5, 4) \rangle|$

Solution

- $|2\mathbb{Z}/6\mathbb{Z}| = (2\mathbb{Z} : 6\mathbb{Z}) = 3$
- $\langle 6 \rangle = \{6, 0\}$, so $|H| = |\{8, 2\}| = 2$
- $o(2 + \langle 6 \rangle) = 3$
- $\langle f + H \rangle = \{f + H, H\}$
- $|\mathbb{Z}_6 \times \mathbb{Z}_8| = 6(8) = 48$ and $|\langle 3 \rangle \times \langle 2 \rangle| = 2(4) = 8$, so $|\mathbb{Z}_6 \times \mathbb{Z}_8 / \langle 3 \rangle \times \langle 2 \rangle| = \frac{48}{8} = 6$.
- $\langle (5, 4) \rangle = \{(5, 4), (10, 8), (0, 12), (5, 16), (10, 20), (0, 0)\}$; so the order of $(\mathbb{Z}_{15} \times \mathbb{Z}_{24}) / \langle (5, 4) \rangle$ is $\frac{(15)(24)}{6} = 60$.

4. For each of the following, find a familiar group to which the given group is isomorphic. (Hint: Consider the group order, properties such as abelianness and cyclicity, group tables, orders of elements, etc.)

- $\mathbb{Z}/14\mathbb{Z}$
- $3\mathbb{Z}/12\mathbb{Z}$

- c. S_8/A_8
- d. $(4\mathbb{Z} \times 15\mathbb{Z})/(\langle 2 \rangle \times \langle 3 \rangle)$
- e. $D_4/\langle r^2 \rangle$

Solution

- a. $\mathbb{Z}_{14}$
- b. $3\mathbb{Z}/12\mathbb{Z}$ is cyclic (since $3\mathbb{Z}$ is cyclic) with order 4, so is isomorphic to $\mathbb{Z}_4$
- c. S_8/A_8 has order 2, so is isomorphic to $\mathbb{Z}_2$.
- d. Since $\gcd(4, 15) = 1$, the group $4\mathbb{Z} \times 15\mathbb{Z}$ is cyclic. So since $|4\mathbb{Z} \times 15\mathbb{Z}|/|\langle 2 \rangle \times \langle 3 \rangle| = \frac{60}{10} = 6$, we have $(4\mathbb{Z} \times 15\mathbb{Z})/(\langle 2 \rangle \times \langle 3 \rangle) \simeq \mathbb{Z}_6$.
- e. $|D_4/\langle r^2 \rangle| = \frac{8}{2} = 4$, and $f\langle r^2 \rangle$ and $r\langle r^2 \rangle$ in $D_4/\langle r^2 \rangle$ each have order 2, so $D_4/\langle r^2 \rangle \simeq \mathbb{Z}_2^2$.

5. Let $H \trianglelefteq G$ with index k , and let $a^{\text{int}} G$. Prove that **Double exponent: use braces to clarify**

Solution

Since $k = (G : H)$, $|G/H| = k$. Then $aH \in G/H$ has an order, call it d , which must divide $|G/H| = k$. So $k = dm$ for some $m \in \mathbb{Z}$. Then $(aH)^k = (aH)^{dm} = ((aH)^d)^m = H^m = H$. Since $(aH)^k = a^k H$, we thus have $a^k H = H$, implying that $a^k \in H$, as desired.

Exercise 9.3

1. Let F be the group of all functions from $[0, 1]$ to $\mathbb{R}$, under pointwise addition. Let

$$N = \{f^{\text{int}} F : f(1/4) = 0\}.$$

Prove that F/N is a group that's isomorphic to $\mathbb{R}$.

Solution

Define $\Phi : F \rightarrow \mathbb{R}$ by $\Phi(f) = f(1/4)$, for every $f \in F$. We have that Φ is a homomorphism, since for every $f, g \in F$,

$$\Phi(f + g) = (f + g)\left(\frac{1}{4}\right) = f\left(\frac{1}{4}\right) + g\left(\frac{1}{4}\right) = \Phi(f) + \Phi(g).$$

We also have that Φ is onto, since if $r \in \mathbb{R}$, then the constant function c_r defined by

$$c_r(x) = r \text{ for every } x \in [0, 1]$$

is sent to r by Φ . So $\Phi(F) = \mathbb{R}$. Finally, if $f \in F$, then

$$f \in \text{Ker}\Phi \Leftrightarrow \Phi(f) = 0 \Leftrightarrow f\left(\frac{1}{4}\right) = 0 \Leftrightarrow f \in N;$$

so $\text{Ker}\Phi = N$. Thus, $F/N \cong \mathbb{R}$, by the First Isomorphism Theorem.

2. Let $N = \{1, -1\} \subseteq \mathbb{R}^*$. Prove that $\mathbb{R}^*/N$ is a group that's isomorphic to $\mathbb{R}^+$.

Solution

Define $\Phi : \mathbb{R}^* \rightarrow \mathbb{R}^+$ by $\Phi(x) = |x|$. We know that Φ is a homomorphism, since $\Phi(xy) = |xy| = |x||y| = \Phi(x)\Phi(y)$, for every $x, y \in \mathbb{R}^*$. Moreover, Φ is clearly onto (so $\Phi(\mathbb{R}^*) = \mathbb{R}^+$), and has

$$\text{Ker}\Phi = \{x \in \mathbb{R}^* : \Phi(x) = 1\} = \{1, -1\} = N.$$

Thus, $\mathbb{R}^*/N \cong \mathbb{R}^+$, by the First Isomorphism Theorem.

3. Let $n^{\text{int}}\mathbb{Z}^+$ and let $H = \{A^{\text{int}}GL(n, \mathbb{R}) : \det A = \pm 1\}$. Identify a group familiar to us that is isomorphic to $GL(n, \mathbb{R})/H$.

Solution

Define $\Phi : GL(n, \mathbb{R}) \rightarrow \mathbb{R}^+$ by $\Phi(A) = |\det A|$. We have that Φ is an epimorphism since for all $A, B \in GL(n, \mathbb{R})$,

$$\Phi(AB) = |\det AB| = |\det A \det B| = |\det A| |\det B| = \Phi(A)\Phi(B),$$

and for all $\lambda \in \mathbb{R}^+$, the diagonal matrix having λ in the uppermost left position and 1's elsewhere down the diagonal gets sent to λ . Since $\text{Ker}\Phi = H$, we have that $GL(n, \mathbb{R})/H \simeq \mathbb{R}^+$, by the First Isomorphism Theorem.

4. Let G and G' be groups with respective normal subgroups N and N' . Prove or disprove: If $G/N \simeq G'/N'$ then $G \simeq G'$.

Solution

The statement is false. Indeed, using the preceding two problems, we see that $\mathbb{R}^*/\{1, -1\}$ and $GL(2, \mathbb{R})/H$, where $H = \{A \in GL(2, \mathbb{R}) : \det A = \pm 1\}$, are both isomorphic to $\mathbb{R}^+$, hence are isomorphic to each other. But we know that $\mathbb{R}^* \not\simeq GL(2, \mathbb{R})$.